United States
Environmental Protection
Agency

Environmental Monitoring and
Support Laboratory
Cincinnati OH 45268

EPA 600/3-82-026
April 1982

EPA

Freshwater Snails (Mollusca: Gastropoda) of North America

EPA-600/3-82-026
April 1982

FRESHWATER SNAILS (MOLLUSCA: GASTROPODA) OF NORTH AMERICA

by

J.B. Burch

Museum of Zoology

Department of Ecology and Evolutionary Biology
Division of Biological Sciences

and

School of Natural Resources

The University of Michigan
Ann Arbor, Michigan 48109

CONTRACT NO. 68-03-1280

Project Officer

Donald J. Klemm

Aquatic Biology Section
Biological Methods Branch
Environmental Monitoring and Support Laboratory
Cincinnati, Ohio 45268

ENVIRONMENTAL MONITORING AND SUPPORT LABORATORY
OFFICE OF RESEARCH AND DEVELOPMENT
U.S. ENVIRONMENTAL PROTECTION AGENCY
CINCINNATI, OHIO 45268

DISCLAIMER

This report has been reviewed by the Environmental Monitoring and Support Laboratory, U.S. Environmental Protection Agency, and approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

FOREWORD

Environmental measurements are required to determine the quality of ambient water, the character of effluents, and the effects of pollutants on aquatic life. The Environmental Monitoring and Support Laboratory - Cincinnati conducts an Agency-wide quality assurance program to assure standardization and quality control of systems for monitoring water and wastewater and carries out research to develop, evaluate, standardize, and promulgate analytical methods to:

* Measure the presence and concentration of physical, chemical, and radiological pollutants in water, wastewater, bottom sediments, and solid waste.
* Concentrate, recover, and identify enteric viruses, bacteria, and other microorganisms in water.
* Measure the effects of pollution on freshwater, estuarine, and marine organisms, including the phytoplankton, zooplankton, periphyton, macrophyton, macroinvertebrates, and fish.
* Automate the measurement of physical, chemical, and biological quality of water.

The effectiveness of measures taken to maintain and restore the biological integrity of the Nation's surface waters is dependent upon our knowledge of the changes in the taxonomic composition of the aquatic life caused by discharges of toxic substances and other pollutants and upon the level of our understanding of the complex relationships that prevail in aquatic ecosystems. Snails occur in almost all types of freshwater habitats and are important components of aquatic food-webs. They also may serve as carriers of parasitic diseases of man and other vertebrates. Snails respond to various chemical and physical characteristics of the environment and are sensitive to changes caused by urban and agricultural run off, industrial, thermal, and domestic pollution. Certain toxic substances (heavy metals, pesticides, and radioactive materials) can concentrate in the tissues and shells of gastropods even when present in trace amounts. The analysis of snails living in biotopes subjected to these materials can indicate those types of pollutants occurring there. This report fills a long standing need for a comprehensive identification manual to the species of freshwater gastropods in North America to assist biologists in analyzing samples and evaluating data collected during studies of the effects of toxic substances and other pollutants on the communities of aquatic organisms.

Robert L. Booth
Acting Director
Environmental Monitoring and Support
Laboratory - Cincinnati

ABSTRACT

Freshwater gastropod mollusks are represented in North America (north of Mexico) by 15 families, 78 genera and, as treated in this manual, 499 species. They are grouped into two large subclasses, the gill-breathing, operculated Prosobranchia and the lung-breathing, non-operculated Pulmonata. The prosobranch snails are represented by 49 genera and 349 species, and the pulmonate snails by 29 genera and 150 species. Systematics are not well worked out in many groups, which makes a definitive listing of species somewhat arbitrary at this time. The species listed in some groups reflect a conservative approach to the lower taxa; in other groups, some of the so-called species possibly may represent merely forms resulting from intraspecific variation.

The main feature of this publication is a series of illustrated taxonomic keys for the identification of the North American (north of Mexico) freshwater snails. Also presented are distributions for each of the species, a generic synonymy and an extensive bibliography.

CONTENTS

Foreword iii
Abstract iv
Acknowledgements vi

I. Introduction 1
II. Identification 7
III. Species List and Distribution Ranges 15
IV. Keys to the Freshwater Gastropods of North America 64
Families and Higher Taxa 64
Family Neritinidae 67
Family Valvatidae (by William H. Heard) 67
Family Viviparidae (by J.B. Burch and Virginia A. Vail) 75
Family Ampullariidae 81
Family Bithyniidae 81
Family Micromelaniidae 83
Family Hydrobiidae 83
Family Pomatiopsidae 115
Family Thiaridae 117
Family Pleuroceridae 118
Family Acroloxidae 147
Family Lymnaeidae 147
Family Physidae 160
Family Planorbidae 168
Family Ancylidae 188

Supplemental Notes 194
Generic Synonymy 210
Glossary 214
References 222
Index to Scientific Names 267

ACKNOWLEDGEMENTS

I wish to express my appreciation to Janet White-Rudolph, whose skills in editing and composition have been indispensable in producing this manual. I am also grateful to Margaret S. Burch for help with photographic work on the plates of figures, to C.M. Patterson for assembling the plates, and to Paul H. Rudolph for constructing the index. The majority of the illustrations are by John L. Tottenham, and once again the high quality of his work is clearly evident. Appreciation is also noted to William H. Heard and Fred G. Thompson for their helpful comments on the manuscript, and to the latter for use of illustrations from his publications on Hydrobiidae.

A note of special thanks is due Donald J. Klemm for his unfailing support and understanding throughout the preparation of this manual. It is hoped that the quality and usefulness of this publication will justify his faith in sponsoring it.

SECTION I

INTRODUCTION

The last manual on freshwater gastropods of North America was written over a hundred years ago by W.G. Binney and published by the Smithsonian Institution in 1865. Many species and genera have been named since that time, and many changes have been made in systematics and in nomenclature. So a new manual on North American freshwater snails is now timely.

Over the years since Binney's manual, a few regional handbooks or monographs have been published on, or which have included, freshwater snails. Some of these are listed chronologically below.

Augustus A. Gould. 1870. *Report on the Invertebrata of Massachusetts.* Edited by W.G. Binney. Wright and Potter, Boston.

R. Ellsworth Call. 1884. *On the Quaternary and Recent Mollusca of the Great Basin, with descriptions of new forms.* Bulletin of the U.S. Geological Survey, No. 11.

R. Ellsworth Call. 1900. *A descriptive illustrated catalogue of the Mollusca of Indiana.* 24th Annual Report of the Department of Geology and Natural Resources of Indiana. (Revised [without figures] by Calvin Goodrich & Henry van der Schalie, 1944.)

Frank Collins Baker. 1902. *The Mollusca of the Chicago area. Part 2, The Gastropoda.* Chicago Academy of Science.

William H. Dall. 1905. *Land and fresh water mollusks of Alaska and adjoining regions.* Smithsonian Institution Harriman Alaska Expedition.

Harold Hannibal. 1912. *A synopsis of the Recent and Tertiary freshwater Mollusca of the California province, based upon an ontogenetic classification.* Proceedings of the Malacological Society of London, Vol. 10.

Junius Henderson. 1924. *Mollusca of Colorado, Utah, Montana, Idaho and Wyoming.* University of Colorado Studies, Vol. 13. (A supplement was published in 1936.)

Frank Collins Baker. 1928. *The fresh water Mollusca of Wisconsin.* Wisconsin Geological and Natural History Survey, Bull. 70.

Ralph V. Chamberlin & David T. Jones. 1929. *A descriptive catalogue of the Mollusca of Utah.* Bulletin of the University of Utah, Vol. 19.

Junius Henderson. 1929. *Non-marine Mollusca of Oregon and Washington.* University of Colorado Studies, Vol. 17. (A supplement was published in 1936.)

Calvin Goodrich. 1932. *The Mollusca of Michigan.* Museum of Zoology, The University of Michigan.

Imogene C.S. Robertson & Clifford L. Blakeslee. 1948. *The Mollusca of the Niagara Frontier region.* Bulletin of the Buffalo Society of Natural Science, Vol. 19.

William J. Clench & Ruth D. Turner. 1956. *Freshwater mollusks of Alabama, Georgia, and Florida from the Escambia to the Suwannee River.* Bulletin of the Florida State Museum (Biological Science), Vol. 1.

A. Byron Leonard. 1959. *Handbook of gastropods in Kansas.* University of Kansas Museum of Natural History Miscellaneous Publication, No. 20.

Aurèle La Rocque. 1968. *Pleistocene Mollusca of Ohio.* Division of Geological Survey, Department of Natural Resources, State of Ohio, Bull. 62. (Includes Recent species.)

Willard N. Harman & Clifford O. Berg. 1971. *The freshwater snails of central New York, with illustrated keys to the genera and species.* Search, Agriculture, Entomology (Ithaca) 2, Vol. 1.

Arthur H. Clarke. 1973. *The freshwater mollusks of the Canadian Interior Basin.* Malacologia, Vol. 13.

Joseph C. Bequaert & Walter B. Miller. 1973. *The mollusks of the arid Southwest, with an Arizona check list.* The University of Arizona Press, Tucson.

William K. Emerson & Morris K. Jacobson. 1976. *Guide to shells, land, freshwater, and marine, from Nova Scotia to Florida.* Alfred A. Knopf, New York.

Many of these, particularly the older publications, are now badly out-of-date, especially in regard to zoological nomenclature.

Also since Binney's manual, various taxonomic groups of gastropods have been monographed, either regionally or totally:

Frank Collins Baker. 1911. *The Lymnaeidae of North and Middle America, Recent and fossil.* Chicago Academy of Science.

Calvin Goodrich. 1917-44. [Numerous papers revising the large family Pleuroceridae, mostly published in the *Occasional Papers of the Museum of Zoology, University of Michigan* and the *Miscellaneous Publications of the Museum of Zoology, University of Michigan.*]

Elmer G. Berry. 1943. *The Amnicolidae of Michigan: distribution, ecology, and taxonomy.* Miscellaneous Publications of the Museum of Zoology, University of Michigan, No. 57.

Frank Collins Baker. 1945. *The molluscan family Planorbidae.* University of Illinois Press, Urbana.

Bengt Hubendick. 1951. *Recent Lymnaeidae. Their variation, morphology, taxonomy, nomenclature, and distribution.* Kungliga Svenska Vetenskapsakademiens Handlingar, Vol. 3.

Paul F. Basch. 1963. *A review of the Recent freshwater limpet snails of North America (Mollusca: Pulmonata).* Bulletin of the Museum of Comparative Zoology at Harvard College, Vol. 129.

Fred G. Thompson. 1968. *The aquatic snails of the family Hydrobiidae of peninsular Florida.* University of Florida Press, Gainesville.

George A. Te. 1975. *Michigan Physidae, with systematic notes on* Physella *and* Physodon *(Basommatophora: Pulmonata).* Malacological Review, Vol. 8.

Fred G. Thompson. 1977. *The hydrobiid snail genus* Marstonia. Bulletin of the Florida State Museum (Biological Science), Vol. 21.

The taxonomic philosophies of the authors of the above have varied considerably, and the taxonomic treatments in the older publications especially need revision. Nevertheless, these, as well as the first list above, have been invaluable source material for the present manual, and may prove to be especially useful to current biologists who must deal with mollusks of the geographic areas or taxonomic groups covered.

In addition to the major publications listed above (and hundreds of more minor ones listed in the References, pp. 222-266), an additional publication should be mentioned: Bryant Walker's (1918) *A synopsis of the classification of the freshwater Mollusca of North America, north of Mexico, and a catalogue of the more recently described species, with notes*, Miscellaneous Publications of the Museum of Zoology, University of Michigan, No. 6. Although this publication is now out-of-date, it has been one of the landmark publications on the classification of North American mollusks for the past sixty years.

Below is a list of the higher taxa and genera of North American freshwater gastropods according to the taxonomic scheme used in this manual. The genera are arranged alphabetically under their tribes, subfamilies or families.

	Number of North American Species	Page Numbers
Subclass Prosobranchia		
Order Neritinacea		
Superfamily **Neritinoidea**		
Family NERITINIDAE		
Genus *Neritina*	1	15, 67
Order Mesogastropoda		
Superfamily **Valvatoidea**		
Family VALVATIDAE		
Genus *Valvata*	11	15, 68
Superfamily **Ampullarioidea**		
Family VIVIPARIDAE		
Subfamily Viviparinae		
Genus *Tulotoma*	1	16, 75
Genus *Viviparus*	3	16, 76
Subfamily Bellamyinae		
Genus *Cipangopaludina*	2	17, 75
Subfamily Lioplacinae		
Genus *Campeloma*	8	17, 78
Genus *Lioplax*	5	18, 75
Family AMPULLARIIDAE		
Genus *Marisa*	1	18, 81
Genus *Pomacea*	2	19, 81
Family BITHYNIIDAE		
Genus *Bithynia*	1	19, 81
Superfamily **Truncatelloidea**		
Family MICROMELANIIDAE		
Genus *Antroselates*	1	19, 83
Family HYDROBIIDAE		
Subfamily Hydrobiinae		
Genus *Aphaostracon*	9	19, 98
Genus *Hoyia*	1	20, 90
Genus *Hyalopyrgus*	2	20, 98
Genus *Littoridinops*	2	20, 90
Genus *Probythinella*	1	21, 88
Genus *Pyrgophorus*	2	21, 90
Genus *Tryonia*	5	21, 90
Subfamily Lithoglyphinae		
Genus *Antrobia*	1	21, 88
Genus *Clappia*	2	22, 88
Genus *Cochliopina*	1	22, 84
Genus *Fluminicola*	12	22, 84
Genus *Gillia*	1	23, 88
Genus *Lepyrium*	1	23, 84
Genus *Somatogyrus*		
Subgenus *Somatogyrus* s.s.	32	23, 88
Subgenus *Walkerilla*	3	24, 88

Subfamily Nymphophilinae		
Genus *Birgella*	1	25, 98
Genus *Cincinnatia*	14	25, 110
Genus *Fontelicella*		
Subgenus *Fontelicella* s.s.	6	26, 110
Subgenus *Microamnicola*	1	26, 110
Subgenus *Natricola*	3	26, 110
Genus *Marstonia*	8	27, 102
Genus *Notogillia*	2	27, 104
Genus *Orygoceras*	1	27, 98
Genus *Pyrgulopsis*	5	27, 110
Genus *Rhapinema*	1	28, 102
Genus *Spilochlamys*	3	28, 104
Genus *Stiobia*	1	28, 98
Subfamily Amnicolinae		
Genus *Amnicola*		
Subgenus *Amnicola* s.s.	11	28, 115
Subgenus *Lyogyrus*	7	19, 115
Genus *Hauffenia*	1	30, 115
Genus *Horatia*	1	30, 115
Subfamily Fontigentinae		
Genus *Fontigens*	8	30, 110
Incertae Sedis	3	31
Family POMATIOPSIDAE		
Genus *Pomatiopsis*	6	31, 115
Superfamily **Vermetoidea**		
Family THIARIDAE		
Genus *Melanoides*	1	31, 117
Genus *Thiara*	1	32, 118
Family PLEUROCERIDAE		
Genus *Elimia*	83	32, 123
Genus *Gyrotoma*	6	40, 118
Genus *Io*	1	41, 118
Genus *Juga*		
Subgenus *Juga* s.s.	3	41
Subgenus *Calibasis*	2	41
Subgenus *Oreobasis*	4	41
Genus *Leptoxis*		
Subgenus *Leptoxis* s.s.	16	42, 120
Subgenus *Athearnia*	1	44, 120
Subgenus *Mudalia*	6	43, 120
Genus *Lithasia*		
Subgenus *Lithasia* s.s.	3	44, 123
Subgenus *Angitrema*	7	44, 123
Genus *Pleurocera*		
Subgenus *Pleurocera* s.s.	18	45
Subgenus *Strephobasis*	3	47
Subclass Pulmonata		
Order Limnophila		
Superfamily **Acroloxoidea**		
Family ACROLOXIDAE		
Genus *Acroloxus*	1	48, 147

Superfamily **Lymnaeoidea**		
Family LYMNAEIDAE		
Subfamily Lymnaeinae		
Genus *Acella*	1	48, 148
Genus *Bulimnea*	1	48, 149
Genus *Fossaria*		
Subgenus *Fossaria* s.s.	11	48, 149
Subgenus *Bakerilymnaea*	11	49, 149
Genus *Lymnaea*	2	50, 148
Genus *Pseudosuccinea*	1	50, 148
Genus *Radix*	1	51, 148
Genus *Stagnicola*		
Subgenus *Stagnicola* s.s.	21	51
Subgenus *Hinkleyia*	3	52
Subfamily Lancinae		
Genus *Fisherola*	1	52, 147
Genus *Lanx*		
Subgenus *Lanx* s.s.	3	53, 147
Subgenus *Walkerola*	1	53, 147
Superfamily **Ancyloidea**		
Family PHYSIDAE		
Subfamily Physinae		
Genus *Physa*	2	53, 160
Genus *Physella*		
Subgenus *Physella* s.s.	16	53
Subgenus *Costatella*	14	55
Subgenus *Petrophysa*	1	56, 168
Subfamily Aplexinae		
Genus *Aplexa*	2	56, 168
Genus *Stenophysa*	2	57, 160
Family PLANORBIDAE		
Subfamily Planorbinae		
Tribe Planorbini		
Genus *Gyraulus*		
Subgenus *Gyraulus* s.s.	1	57, 172
Subgenus *Armiger*	1	57, 168
Subgenus *Torquis*	3	57, 172
Tribe Drepanotremini		
Genus *Drepanotrema*		
Subgenus *Antillorbis*	1	57, 171
Subgenus *Fossulorbis*	2	58, 171
Tribe Biomphalariini		
Genus *Biomphalaria*	2	58, 173
Tribe Helisomini		
Genus *Helisoma*		
Subgenus *Helisoma* s.s.	2	58, 174
Subgenus *Carinifex*	1	58, 174
Genus *Menetus*		
Subgenus *Menetus* s.s.	1	59, 173
Subgenus *Micromenetus*	3	59, 172
Genus *Planorbella*		
Subgenus *Planorbella* s.s.	2	59, 175

Subgenus *Pierosoma*	12	59, 175
Subgenus *Seminolina*	2	61, 175
Genus *Planorbula*	2	61, 171
Genus *Promenetus*	2	61, 173
Genus *Vorticifex*		
Subgenus *Parapholyx*	2	61, 174
Subfamily Neoplanorbinae		
Genus *Amphigyra*	1	62, 171
Genus *Neoplanorbis*	4	62, 171
Family ANCYLIDAE		
Subfamily Ancylinae		
Genus *Rhodacmea*	3	62, 188
Subfamily Ferrissinae		
Genus *Ferrissia*	5	62, 190
Subfamily Laevapecinae		
Genus *Hebetancylus*	1	63, 193
Genus *Laevapex*	2	63, 193

The Recent freshwater gastropod fauna of North America is a very diverse one, comprising two molluscan subclasses, three orders, eight superfamilies, 15 families, 78 genera and, as listed in this manual, 499 species (plus an additional 143 subspecies and forms). This is a large fauna, and, since systematists dealing with these mollusks over the years have not been many, systematics have not been well worked out in many groups. Because of this, a definitive list of species can be made only somewhat arbitrarily at this time.

While the treatment of various groups represents my own interpretation of species-level taxonomy, in other groups I have simply followed the latest reviser. Because of the latter procedure, the overall treatment of the various groups may be a bit uneven. For example, within the Pleuroceridae, the treatment of the genera *Elimia* and *Pleurocera* probably reflects a nomenclatural system partially imposed on clinal variation, i.e., there is a liberal supply of latinized binomials and trinomials, while in the Ancylidae, the species lists and keys may reflect an overly conservative approach to the lower taxa. However, the purpose of this manual is not to present a major revision of the systematics of North American freshwater gastropods – such an undertaking would require many years – but to present a reasonable, current classification, a means to identify the species of North American gastropods, and a generalized distribution of the included species.

SECTION II

IDENTIFICATION

Characters of the shells of freshwater gastropods are very important in species recognition and usually for generic and familial placement as well. Especially useful are the size and general form of the shell. Among the many species, the shell may take various shapes, yet, for any one species, the shell shape is usually quite constant (excepting, of course, minor clinal, populational and ecophenotypic variations exhibited by some species). The shells among the different species may vary from very elongate, to globose, depressed and discoidal. The shell may be longer than wide, or wider than long (the columella determining the antero-posterior (length/width) axis). Its coils (whorls) may turn either to the left or to the right, be round, angular, shouldered or flattened, and have shallow or impressed sutures. The shell may have few or many whorls, may lack an opening (umbilicus) at its base, or may have either a narrow or wide opening. The columella or central axial column of the shell may be either twisted or straight and may or may not end abruptly. The outer lip of the shell may be either straight or variously curved and is sometimes turned back or reflected. The surface of the shell may be marked in various ways, i.e., differentially colored or sculptured, or may be simply unicolored and smooth. The outline of the shell aperture ("mouth") may take many forms due to the shape and relation of the whorls to each other. The aperture may or may not be closed by an operculum, which itself has important recognition characters. The operculum may be round, oval or spindle-shaped, and concentric, paucispiral or multispiral, depending on the way in which it is formed.

In some groups, aspects of soft anatomy are essential for identification, because the various taxa in these groups have shells which are relatively uniform or have few distinctive characteristics. This is true especially of the Hydrobiidae, where not only are many of the genera distinguished on the basis of the characteristics of their terminal male genitalia, but the subfamilies are recognized in this way also. This, of course, makes identification very difficult for species of such a family when only empty shells are available. However, identification can be aided by careful inspection of the shell illustrations and by taking into account the known distributions of the various species.



FIG. 1. An operculated snail, i.e., one which carries an operculum attached to its dorsal posterior foot. a, Position of the operculum when the snail is active; b, position of the operculum when the snail has withdrawn into its shell.



FIG. 2. Direction of coiling of gastropod shells. a, Shell coiled to the left, i.e., *sinistral*; b, shell coiled to the right, i.e., *dextral*.



FIG. 3. a, A snail with *sinistral* organization of its body, i.e., respiratory, excretory and reproductive openings are on the *left* side; b, a snail with *dextral* organization of its body, i.e., respiratory, excretory and reproductive openings are on the *right* side.



FIG. 4. Orientation of the shell.



FIG. 5. Shell terminology.



FIG. 6. Shell surface sculpture.



FIG. 7. Shell sizes: up to 10 mm = small; 10-30 mm = medium; over 30 mm = large.



FIG. 8. Shell shapes. a, subglobose; b, oval; c, fusiform or spindle-shaped; d, turbiniform; e, cylindrical.



FIG. 9. Shell shapes. a, narrowly conic; b, elongately conic; c, broadly (ovately) conic; d, globosely conic; e, depressed conic.



FIG. 10. Method of counting whorls. This shell has 3¼ whorls. FIGS. 11-14. Shell terminology. Fig. 11. a, Shell with well-rounded whorls and indented sutures; b, shell with flattened whorls and shallow sutures; c, shell with shouldered whorls. Fig. 12. Planorbiform or discoidal shell. Fig. 13. Ancyliform or limpet-shaped shell. Fig. 14. Neritiniform shell. FIG. 15. Types of opercula. a, Multispiral; b, paucispiral; c, concentric; d, concentric with spiral nucleus. FIG. 16. A valvatid snail, showing bipectinate gill and pallial tentacle (from Harman & Berg, 1971, as modified from F.C. Baker, 1928c).

ARCHAEOGASTROPODA and NERITINACEA

RHIPIDOGLOSSA (=ZYGOBRANCHIA +ASPIDOBRANCHIA, =DIOTOCARDIA)

All the Archaeogastropoda [all marine] except the limpets (i.e., except the so-called Docoglossa: Acmaeidae, Patellidae and Lepetidae).

The Neritinacea [marine, freshwater and land].





DOCOGLOSSA (=PATELLOIDEA)

The archaeogastropod limpets [all marine].

Families Acmaeidae
Patellidae
Lepetidae



MESOGASTROPODA

TAENIOGLOSSA

Contains most of the mesogastropods [marine and freshwater], and in North American freshwaters all the Prosobranchia except the Neritinacea.

PTENOGLOSSA (=EPITONIOIDEA)

Contains the specialized Epitoniidae (=Scalidae) and Janthinidae [both marine].

NEOGASTROPODA = Stenoglossa

RACHIGLOSSA

Contains many of the neogastropods [all marine].



TOXOGLOSSA

Radula consists of only long teeth (marginals). The name refers to the poison gland associated with the radula of *Conus*.

Families Conidae
Turridae [all marine]
Cancellariidae

GYMNOGLOSSA

No radula; radula not needed because of parasitic existence.
Eulimidae - a mesogastropod [marine] family.
Pyramidellidae - an opisthobranch [marine] family.

FIG. 17. Prosobranch snail classification based on radulae. The Prosobranchia have been divided in the past into a number of groups which take their names from the prevalent type of radulae they possess. This classification generally separates assemblages that are also distinct in their soft anatomy, but in most groups there are species that show anomalies. North American freshwater prosobranchs possess only the rhipidoglossate (in the Neritinidae) and the taenioglossate (in the other prosobranch families) types of radulae.



FIG. 18. The radula of a freshwater pulmonate snail (*Biomphalaria*) and its relation to the alimentary system. a, The alimentary system, mainly dorsal view; b, longitudinal section through the buccal mass, showing the orientation of the radula as viewed from the right side; c, right side and dorsal views of the radula; d, four teeth from one transverse row of the radular ribbon, r = rachidian or central tooth, 1 = first tooth to the right of the central tooth (a lateral tooth), 10, 11 = 10th and 11th teeth to the right of the central tooth (marginal teeth). From Barbosa et al. (1968), after Demian.



FIG. 19. Central or rachidian tooth of a gastropod (a truncatelloid pomatiopsid snail) showing the arrangement of the cutting edges. a, Tooth from above; b, profile (side) view of tooth (from Pilsbry & Bequaert, 1927).

FIG. 20. Major drainages of North America (north of Mexico).

SECTION III

SPECIES LIST AND DISTRIBUTION RANGES

Family NERITINIDAE[1]

Genus *Neritina* Lamarck 1816

Neritina reclivata reclivata (Say 1822) [Figs. 21, 22][2]
Florida to Mississippi. Also Cuba, northern Mexico and Venezuela (H. B. Baker, 1923).

Neritina reclivata sphaera Pilsbry 1931
Drainage canal draining Lake Okechobee, a few miles from the Atlantic, Ojus, Florida (Pilsbry, 1931).

Neritina reclivata palmae Dall 1885
Brook near Palma Sola, Florida (Dall, 1885)

Family VALVATIDAE*

Genus *Valvata* Müller 1774

Valvata bicarinata bicarinata Lea 1841 [Fig. 23]
Of discontinuous distribution: New Jersey and Pennsylvania; and Iowa, Illinois, Tennessee, Alabama, Georgia and North Carolina.

Valvata bicarinata morph *normalis* Walker 1902 [Fig. 24]
Distribution nearly as for *V. bicarinata* s.s., but not in Georgia and North Carolina.

Valvata humeralis Say 1829 [Fig. 25]
Known from Montana south to Colorado, and west to British Columbia and California.

Valvata lewisi lewisi Currier 1868 [Fig. 26]
Southern Canada from Quebec west to British Columbia, and northern U.S.A. from New York west to Minnesota.

Valvata lewisi morph *ontarioensis* F.C. Baker 1931 [Fig. 27]
Northwestern Ontario in the region of Lake Superior drained by the headwaters of the Attawapiskat, Albany and Severn river systems (Clarke, 1973).

Valvata mergella Westerlund 1883 [Fig. 28]
Northwestern North America: Alaska to Washington.

---

[1] Superscript numbers throughout the text refer to corresponding comments under Supplemental Notes, which appear on pp. 194-209.

*The species list and distributions for the Valvatidae is by William H. Heard.

Valvata perdepressa Walker 1906 [Fig. 29]

V. perdepressa s.s. and *V. perdepressa* ?form *walkeri* F.C. Baker 1930: the Great Lakes (Lakes Michigan, Huron, Erie and Ontario).

Valvata piscinalis (Müller 1774) (?form *obtusa* Draparnaud 1801) [Fig. 30]

Introduced from Europe into the lower Great Lakes (Lake Ontario, and perhaps tributaries near the lake).

Valvata sincera sincera Say 1824 [Fig. 31]

Maine west to Alberta, and south to South Dakota and Indiana.

Valvata sincera ?form *danielsi* Walker 1906 [Fig. 32]

Of discontinuous distribution: New Brunswick in eastern Canada, and Illinois, Wisconsin and Minnesota in north central U.S.A.

Valvata sincera nylanderi Dall 1905

Quebec and Maine west to Ontario and Minnesota.

Valvata tricarinata (Say 1817) [Fig. 33] and its morphs: *bakeri* Fluck 1932, *basalis* Vanatta 1915, *infracarinata* Vanatta 1915, *mediocarinata* F.C. Baker 1932, *perconfusa* Walker 1917, *simplex* Gould 1841, *tricarinata* s.s., and *unicarinata* DeKay 1843

Quebec and New Brunswick west to Alberta, and south to Wyoming, Arkansas and Virginia.

Valvata utahensis Call 1884 [Fig. 34]

V. utahensis s.s. and *V. utahensis* morph *horati* Baily & Baily 1951 are known only from Idaho and Utah.

Valvata virens Tryon 1863 [Fig. 35]

California, Oregon and Nevada.

Valvata winnebagoensis F.C. Baker 1928 [Fig. 36]

Of sporadic occurrence in Michigan (Ottawa County), Wisconsin (Winnebago and Oshkosh counties) and Minnesota (Rice County).

Family VIVIPARIDAE

Subfamily Viviparinae

Genus *Tulotoma* Haldeman 1840

Tulotoma magnifica (Conrad 1834)[3] [Figs. 44, 45]

Coosa-Alabama river system in Alabama (Clench, 1962a).

Genus *Viviparus* Montfort 1810

Viviparus georgianus (Lea 1834) [Figs. 46, 47]

South central Florida, Georgia, Alabama and north, mainly in the Mississippi River system, to Illinois and northwestern Indiana; it has invaded Ohio, Michigan, Wisconsin, Virginia, Pennsylvania, New York, New Jersey, New England and Quebec since 1867 (Clench, 1962a; Clench & Fuller, 1965).

Viviparus intertextus (Say 1829) [Fig. 48]

The Houston ship channel system west of Houston, Harris County, and the San Jacinto, Liberty and Neches river systems, Texas; the Bayou Teche system in

Louisiana; the Mississippi River system in Louisiana, eastern Arkansas, northwestern Tennessee, Illinois, eastern Iowa, Minneapolis and White Bear Lake, Minnesota; Pearl River system, Mississippi; Coosa-Alabama river system, Alabama; Altamaha River system, Georgia; Edisto and Santee river systems, South Carolina; Rainy Lake, Koochiching County, Minnesota (Clench & Fuller, 1965).

Viviparus subpurpureus (Say 1829) [Figs. 49-51]

Mississippi River system north to southeastern Iowa, northwestern Illinois and northern Kentucky; Neches and Sabine river systems in eastern Texas and Sabine and Atchafalaya river systems in western Louisiana; Pascagoula River system in southeastern Mississippi (Clench & Fuller, 1965).

Subfamily Bellamyinae

Genus *Cipangopaludina* Hannibal 1912

Cipangopaludina chinensis malleata (Reeve 1863) [Fig. 52]

Widely introduced in the United States. Clench & Fuller (1965) list localities in Arizona, California, Colorado, Delaware, Florida, Indiana, Maine, Massachusetts, Michigan, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Rhode Island, Utah, Vermont and Washington. Originally from Asia.

Cipangopaludina japonica (Martens 1861) [Fig. 53]

Widely introduced in the United States (some reports may be confused with *C. chinensis malleata*). Clench & Fuller (1965) list localities in Massachusetts, Michigan and Oklahoma. Originally from Asia.

Subfamily Lioplacinae

Genus *Campeloma* Rafinesque 1819[4]

Campeloma crassula Rafinesque 1819 [Figs. 42, 54, 55]

Midwestern United States in the Great Lakes-St. Lawrence and Mississippi drainages as far west as Iowa and south to Tennessee.

Campeloma decampi ('Currier' Binney 1865) [Fig. 56]

Jackson, Limestone and Madison counties, northern Alabama (Clench & Turner, 1955).

Campeloma decisum (Say 1817) s.l. (includes forms such as *C. brevispirum* [Figs. 58, 59], *C. decisum* s.s. [Fig. 57], *C. exilis, C. gibbum* [Fig. 60], *C. integrum*[5] [Fig. 38], *C. leptum, C. lewisi* [Fig. 37], *C. milesi* [Fig. 39] and *C. tannum* [Fig. 61]

Eastern North America, from Nova Scotia, southern Ontario and southern Manitoba south to Texas, Louisiana, Mississippi, Alabama, northern Georgia and Virginia.

Campeloma floridense Call 1886 [Fig. 62]

Eastern Florida: the upper St. John's River and its tributaries; Lake Jessup;

Miami (?), Dade County.

Campeloma geniculum (Conrad 1834) [Figs. 63, 64]
Suwannee River, Florida, west to the Escambia River, Alabama (Clench & Turner, 1956).

Campeloma limum (Anthony 1860) [Fig. 41]
Atlantic drainage, from Georgia to North Carolina.

Campeloma parthenum Vail 1979 [Fig. 65]
Ochlockonee River drainage in Florida: Lake Talquin and the Little and Ochlockonee rivers (Vail, 1979a).

Campeloma regulare (Lea 1841) [Figs. 40, 66]
Alabama-Coosa river system, Alabama.

Genus *Lioplax* Troschel 1857

Lioplax cyclostomaformis (Lea 1841) [Fig. 43]
Coosa-Alabama-Tombigbee river system from northwestern Georgia, south to Selma, Dallas County, on the Alabama River, and Big Prairie Creek, Marengo County, on the Tombigbee River in Alabama (Clench & Turner, 1955); Tensas River, near Delhi, Madison County, Louisiana (Clench, 1962b).

Lioplax pilsbryi pilsbryi Walker 1905 [Fig. 67]
Chipola River, Florida.

Lioplax pilsbryi choctawhatchensis Vanatta 1935[6]
Choctawhatchee, Escambia, Flint and Suwannee river systems, Florida and Georgia.

Lioplax subcarinata (Say 1816)[7] [Fig. 68]
Atlantic drainage. Cedar Lake near Litchfield (upper Susquehanna drainage) and Albany (Hudson River drainage), New York, south to South Carolina (Clench & Turner, 1955; Clench, 1965c; Vail, 1979b).

Lioplax sulculosa (Menke 1828)[7]
Mississippi drainage. Northwestern Wisconsin and eastern Minnesota south to northwestern Arkansas and east to southwestern Ohio and northern Kentucky (Clench & Turner, 1955; Vail, 1979b); Paint Rock River of the Tennessee River system, near Paint Rock, Jackson County, Alabama (Clench, 1962b).

Lioplax talquinensis Vail 1979 [Fig. 69]
Ochlockonee River, Florida (Lake Talquin and upstream), and Yellow River (northwestern Florida and southern Alabama) drainages (Vail, 1979b).

Family AMPULLARIIDAE

Genus *Marisa* Gray 1824

Marisa cornuarietis (Linnaeus 1758) [Figs. 70, 71]
Northern South America and southern Central America (H.B. Baker, 1930); introduced into southern Florida (Hunt, 1958; Robins, 1970).

Genus *Pomacea* Perry 1810

Pomacea bridgesi (Reeve 1856)
Brazil; introduced into Florida (Clench, 1966).
Pomacea paludosa (Say 1829) [Figs. 72, 73]
Choctawhatchee, Econfina, St. Marks and Suwannee river systems, Florida, and the Apalachicola River system in Georgia and Florida (Clench & Turner, 1956); Gantt, Covington County, Alabama (Hubricht, 1962).

Family BITHYNIIDAE

Genus *Bithynia* Leach (in Abel) 1818

Bithynia tentaculata tentaculata (Linnaeus 1758), introduced, and *Bithynia tentaculata magnalacustris* F.C. Baker 1928, native(?) [Fig. 74]
Great Lakes region from Albany, New York, west to Winnebago Lake, Calumet and Winnebago counties, Wisconsin; recorded from New York, Pennsylvania, Ohio, Indiana, Illinois, Michigan and Wisconsin (F.C. Baker, 1928a,c).

Family MICROMELANIIDAE

Genus *Antroselates* Hubricht 1963

Antroselates spiralis Hubricht 1963 [Fig. 108]
Springs and streams in caves in Crawford County, Indiana, and Edmonson County, Kentucky (Hubricht, 1963b).

Family HYDROBIIDAE

Subfamily Hydrobiinae s.s.

Genus *Aphaostracon* Thompson 1968

Aphaostracon asthenes Thompson 1968 [Figs. 95, 109]
Blue Springs, three miles west of Orange City, Volusia County, Florida (Thompson, 1968).
Aphaostracon chalarogyrus Thompson 1968 [Figs. 96, 110]
Magnesia Springs, 3.7 miles west of Hawthorne, Alachua County, Florida (Thompson, 1968).
Aphaostracon hypohyalina Thompson 1968 [Figs. 97, 111, 112]
North central Florida in springs along the lower half of the Suwannee River and its tributary, the Santa Fe River, and in a nearby landlocked spring (Thompson, 1968).
Aphaostracon monas (Pilsbry 1899) [Figs. 98, 113, 114]
Wekiwa Springs, Florida, and the Wekiva River for about one mile below the springs (Thompson, 1968).

Aphaostracon pachynotus Thompson 1968 [Figs. 100, 115, 116]
Eastern Florida, from the upper half of the St. Johns River near Sanford, south to the Miami region (Thompson, 1968).

Aphaostracon pycnus Thompson 1968 [Figs. 101, 117, 118]
Alexander Springs Run, Lake County, Florida (Thompson, 1968).

Aphaostracon rhadinus Thompson 1968 [Figs. 75, 99]
In small streams and sloughs in northeastern Florida draining into the St. Johns River north of Palatka, and in an artificial lake near the coast in St. Johns County (Thompson, 1968).

Aphaostracon theiocrenetus Thompson 1968 [Figs. 102, 119, 120]
Clifton Springs Run, about two miles north of Oviedo, Seminole County, Florida (Thompson, 1968).

Aphaostracon xynoelictus Thompson 1968 [Figs. 103, 121, 122]
Fenney Springs, two miles east of Coleman, Sumter County, Florida (Thompson, 1968).

Genus *Hoyia* F.C. Baker 1926

Hoyia sheldoni (Pilsbry 1890) [Fig. 76]
Lake Michigan, off Racine, Wisconsin (Pilsbry 1890d; F.C. Baker, 1928c).

Genus *Hyalopyrgus* Thompson 1968

Hyalopyrgus aequicostatus (Pilsbry 1889) [Figs. 77, 78, 83, 84, 104]
Lower half of the St. Johns River system and the Withlacoochee River system, south to Tampa Bay and the Orlando area; also Lake Okeechobee (Thompson, 1968).

Hyalopyrgus brevissimus (Pilsbry 1890) [Figs. 123, 124]
Orange, Seminole, Sumter and Volusia counties in central Florida (Thompson, 1968).

Genus *Littoridinops* Pilsbry 1952

Littoridinops monroensis (Frauenfeld 1863) [Figs. 80, 85, 86, 105]
Florida and Bahama Islands; in Florida it is generally distributed along both coasts and the Florida Keys, primarily in brackish water, but it has invaded marginal fresh water, and occurs throughout the St. Johns drainage system (Thompson, 1968).

Littoridinops tenuipes Couper (in Haldeman) 1844 [Figs. 79, 87, 106, 125]
Streams draining into the Atlantic Ocean and the Inland waterway of east Florida and Georgia, from Dade County, Florida, north to at least McIntosh County, Georgia (Thompson, 1968).

Genus *Probythinella* Thiele 1928

Probythinella lacustris (F.C. Baker 1928)[8] [Figs. 107, 129-131]
Canada: throughout Ontario and Manitoba, northern Saskatchewan, and in the Northwest Territories south of the tree-line (Clarke, 1973); United States: New York west to Iowa, south to Kentucky and Arkansas (F.C. Baker, 1928c); also North Dakota, South Dakota, Nebraska, Missouri and Alabama (Hibbard & Taylor, 1960).

Genus *Pyrgophorus* Ancey 1888

Pyrgophorus platyrachis Thompson 1968 [Figs. 88, 132]
Throughout the southern part of peninsular Florida from Lake Okeechobee south, and along the coast as far north as southern Brevard County (Thompson, 1968).

Pyrgophorus spinosus (Call & Pilsbry 1886) [Fig. 126]
Guadalupe River and its tributary, Comal Creek, Comal County, Texas (Call & Pilsbry, 1886; Pilsbry, 1887b).

Genus *Tryonia* Stimpson 1865

Tryonia cheatumi (Pilsbry 1935) [Figs. 127, 128, 133]
Phantom Lake, near Toyahvale, Reeves County, Texas (Pilsbry, 1935a).

Tryonia clathrata Stimpson 1865 [Fig. 134]
Pahranagat Valley, southern Nevada (Taylor, 1966b).

Tryonia diaboli (Pilsbry & Ferriss 1906) [Fig. 135]
Devil's River, and Rio San Filipe, Val Verde County, Texas (Pilsbry & Ferriss, 1906).

Tryonia imitator (Pilsbry 1899)
San Francisco Bay to San Diego County, California, in brackish water (Taylor, 1966b); Quitobaquito Springs, Organ Pipe Cactus National Monument, Pima County, Arizona (Bequaert & Miller, 1973).

Tryonia protea (Gould 1855) [Figs. 136, 137]
Colorado Desert and Fish Springs, Imperial County, California (subfossil) (Gould, 1855a; Taylor, 1966b); Santa Cruz River, Tucson, Pima County, Arizona (Pilsbry & Ferriss, 1915); near Buckeye, Maricopa County, Arizona (Bequaert & Miller, 1973).

Subfamily Lithoglyphinae

Genus *Antrobia* Hubricht 1971

Antrobia culveri Hubricht 1972 [Fig. 138]
Stream in Tumbling Creek Cave, near Protem, Taney County, Missouri (Hubricht, 1971).

Genus *Clappia* Walker 1909[9]

Clappia cahabensis Clench 1965
Cahaba River, Bibb County, Alabama (Clench, 1965b).
Clappia umbilicata (Walker 1904) [Figs. 139, 143, 144]
Coosa River, Alabama (Walker, 1904a, 1909c).

Genus *Cochliopina* Morrison 1946

Cochliopina riograndensis (Pilsbry & Ferriss 1906) [Fig. 140]
Lower Pecos River and Rio Grande valleys, Texas; coastal plain in Tamaulipas, Mexico (Taylor, 1966b).

Genus *Fluminicola* Stimpson 1865[10]

Fluminicola columbiana Hemphill (in Pilsbry) 1899 [Fig. 145]
Middle portions of Columbia River, Washington, and lower Snake River, Washington and Idaho.
Fluminicola erythopoma Pilsbry 1899
Ash Meadows, Nye County, Nevada (Pilsbry, 1899a).
Fluminicola fusca (Haldeman (in Chenu) 1847) [Fig. 141]
"Oregon territory" (Haldeman, 1847).
Fluminicola hindsi (Baird 1863)
Upper Green River and tributaries, Wyoming; tributaries of Great Salt Lake, Utah; upper Snake River and tributaries and Salmon River, Idaho; Spokane, Little Spokane and Grande Ronde rivers, Washington (see Taylor, 1966a).
Fluminicola merriumi Pilsbry & Beecher (in Pilsbry) 1892 [Fig. 146]
Pahranagat Valley, Nevada (Pilsbry, 1892a).
Fluminicola minutissima Pilsbry 1907 [Fig. 147]
Price Valley, Weiser Canyon, Washington County, Idaho (Pilsbry, 1907).
Fluminicola modoci Hannibal 1912
California: Fletcher's Spring, south end of Goose Lake; Fritter's Spring, head of Willow Creek, Honey Lake basin; Troxel's Spring, Eagle Lake (Hannibal, 1912b).
Fluminicola nevadensis Walker 1916 [Fig. 148]
Cortez foothills, Humboldt Valley, Elko County, Nevada (Walker, 1916).
Fluminicola nuttalliana (Lea 1838) [Fig. 142]
Probably inhabits the entire Columbia Valley (Pilsbry, 1899a).
Fluminicola seminalis (Hinds 1842)
Sacramento, Pitt and Fall rivers and tributaries, and Surprise Valley, California; Klamath River, Oregon (Pilsbry, 1899a).
Fluminicola turbiniformis (Tryon 1865) [Fig 152]
Upper Sacramento and Pitt rivers and various tributaries in northeastern California, western Nevada and central southern Oregon (see Taylor, 1966a).
Fluminicola virens (Lea 1838)
Willamette, lower Columbia, upper Deschutes and Umpqua rivers, Oregon;

Olympia and San Juan County, Washington; Vancouver Island (Pilsbry, 1899a).

Genus *Gillia* Stimpson 1865

Gillia altilis (Lea 1841) [Fig. 191]
Atlantic drainage from New Jersey to South Carolina (Walker, 1918a).

Genus *Lepyrium* Dall 1896

Lepyrium showalteri (Lea 1861) [Figs. 192, 193]
Cahaba and Coosa rivers, Alabama.

Genus *Somatogyrus* Gill 1863

Somatogyrus alcoviensis Krieger 1972
Alcovy and Yellow rivers, Georgia (Krieger, 1972).
Somatogyrus amnicoloides Walker 1915 [Fig. 149]
Ouachita River, Arkadelphia, Arkansas (Walker, 1915c).
Somatogyrus aureus Tryon 1865 [Figs. 151, 194]
Tennessee River (Tryon, 1865i).
Somatogyrus biangulatus Walker 1906 [Fig. 153]
Tennessee River, Florence, Alabama (Walker, 1906a).
Somatogyrus constrictus Walker 1904 [Fig. 154]
Coosa River, Alabama (Walker, 1904a).
Somatogyrus crassilabris Walker 1915 [Fig. 155]
North Fork of the White River, Norfolk, Arkansas (Hinkley, 1915).
Somatogyrus crassus Walker 1904 [Figs. 156, 157]
Coosa River, Alabama (Walker, 1904a).
Somatogyrus currierianus (Lea 1863) [Fig. 158]
Huntsville, Alabama (Lea, 1863).
Somatogyrus decipiens Walker 1909 [Figs. 159, 160]
Coosa River, Alabama (Walker, 1909c).
Somatogyrus depressus (Tryon 1862) [Fig. 195]
Mississippi River, Davenport, Iowa (Tryon, 1862); Wisconsin, Iowa and Illinois (F.C. Baker, 1928c).
Somatogyrus excavatus Walker 1906 [Figs. 161, 162]
Shoal Creek, Florence, Alabama (Walker, 1906a).
Somatogyrus georgianus Walker 1904 [Fig. 163]
Chattanooga River, Georgia; Tennessee, Cahaba and Alabama rivers, Alabama (Walker, 1904a).
Somatogyrus hendersoni Walker 1909 [Fig. 164]
Coosa River, Alabama (Walker, 1909c).
Somatogyrus hinkleyi Walker 1904 [Figs. 165, 166]
Coosa and Tallapoosa rivers, Alabama (Walker, 1904a).
Somatogyrus humerosus Walker 1906 [Fig. 167]
Tennessee River, Florence, Alabama (Walker, 1906a).

Somatogyrus integra (Say 1829)[11]
Ohio River and tributaries in Ohio, Indiana, Kentucky and eastern Illinois.
Somatogyrus nanus Walker 1904 [Fig. 168]
Coosa River, Alabama (Walker, 1904a).
Somatogyrus obtusus Walker 1904 [Fig. 169]
Coosa River, Alabama (Walker, 1904a).
Somatogyrus parvulus Tryon 1865 [Figs. 170, 171]
Powells River, Tennessee (Tryon, 1865i).
Somatogyrus pennsylvanicus Walker 1904 [Figs. 172-174]
Columbia, Pennsylvania (Walker, 1904a); Potomac River, Harper's Ferry, West Virginia (Walker, 1906a).
Somatogyrus pilsbryanus Walker 1904 [Fig. 175]
Tallapoosa River, Tallassee, Alabama (Walker, 1904a).
Somatogyrus pumilus (Conrad 1834) [Fig. 176]
Black Warrior River and Cahatchee Creek, Shelby, Alabama (Walker, 1906a).
Somatogyrus pygmaeus Walker 1909 [Fig. 177]
Coosa River, Alabama (Walker, 1909c).
Somatogyrus quadratus Walker 1906 [Fig. 178]
Tennessee River and Shoal Creek, Florence, Alabama (Walker, 1906a).
Somatogyrus sargenti Pilsbry 1895 [Fig. 179]
Mud Creek and tributary, tributaries of the Tennessee River, Alabama (Pilsbry, 1895a; Sargent, 1895).
Somatogyrus strengi Pilsbry & Walker 1906 [Fig. 180]
Tennessee River and Shoal Creek, Florence, Alabama; Bridgeport, Alabama (Pilsbry & Walker, 1906a).
Somatogyrus substriatus Walker 1906 [Fig. 181]
Tennessee River, Florence, Alabama; Tombigbee River, Columbus, Mississippi (Walker, 1906a); Uchee Creek, Fort Mitchell, Russell County, and Choctawhatchee River, Dale County, Alabama (Clench & Turner, 1956).
Somatogyrus tennesseensis Walker 1906 [Fig. 182]
Shoal Creek, Florence, Tennessee (Walker, 1906a).
Somatogyrus trothis Doherty 1878
Ohio River, Campbell County, Kentucky (Doherty, 1878).
Somatogyrus tryoni Pilsbry & Baker 1927[9]
Ashippun, Bark and Crawfish rivers, and Lake Michigan drainage, Milwaukee, Wisconsin; Mukwonago River and Creek, Waukesha County, Illinois (Pilsbry & F.C. Baker, 1927).
Somatogyrus walkerianus Aldrich 1905 [Fig. 185]
Conecut River, Escambia County, Alabama (Aldrich, 1905).
Somatogyrus wheeleri Walker 1915 [Figs. 183, 184]
Ouachita River, Arkadelphia, Arkansas (Walker, 1915c).

Subgenus *Walkerilla* Thiele 1928

Somatogyrus (Walkerilla) coosaensis Walker 1904 [Figs. 150, 186, 196]
Coosa and Catawba rivers, Alabama (Walker, 1904a, 1906a).
Somatogyrus (Walkerilla) tenax Thompson 1969 [Figs. 89, 197, 201]
Broad River, Elbert County, Georgia (Thompson, 1969).

Somatogyrus (Walkerilla) virginicus Walker 1904[9, 12] [Fig. 187]
Barnard's Ford, Rapidan River, Virginia (Walker, 1904a).

Subfamily Nymphophilinae

Genus *Birgella* F.C. Baker 1926

Birgella subglobosa (Say 1825) [Figs. 188, 198, 202]
Great Lakes; the river and creek form (*isogona* Say 1829) ranges from Ohio west to Iowa, and from Michigan south to Alabama and Arkansas (F.C. Baker, 1928c).

Genus *Cincinnatia* Pilsbry 1891

Cincinnatia cincinnatiensis (Anthony 1840)[5] [Figs. 189, 199, 203]
New York and Pennsylvania west to southern Manitoba, southern Saskatchewan, North Dakota, Utah and Texas (Clarke, 1973).
Cincinnatia comalensis (Pilsbry & Ferriss 1906) [Fig. 190]
Comal Creek, near New Braunfels, and the Guadelupe River, about four miles above New Braunfels, Comal County, Texas (Pilsbry & Ferriss, 1906).
Cincinnatia floridana (Frauenfeld 1863) [Figs. 204, 235]
Confined to Florida: from the Suwannee River south to Orlando and Hillsborough County (Thompson, 1968).
Cincinnatia fraterna Thompson 1968 [Figs. 200, 205]
Creeks, small streams and sloughs along the lower third of the St. Johns River, Florida (Thompson, 1968).
Cincinnatia helicogyra Thompson 1968 [Figs. 206, 222]
Spring-fed lagoon on the south side of the head of the Crystal River, Citrus County, Florida (Thompson, 1968).
Cincinnatia integra (Say 1829)
Ohio River and tributaries in Ohio, Indiana, Kentucky and southeastern Illinois.
Cincinnatia mica Thompson 1968 [Figs. 207, 223]
Small spring along the west bank of the Ichetucknee River about one mile northeast of U.S. Highway 27, Suwannee County, Florida (Thompson, 1968).
Cincinnatia monroensis (Dall 1885) [Fig. 208]
Brook flowing from Benson's Mineral Spring, Enterprise, Volusia County, Florida (Dall, 1885; Thompson, 1968).
Cincinnatia parva Thompson 1968 [Figs. 209, 224]
Blue Springs, three miles west of Orange City, Volusia County, Florida (Thompson, 1968).
Cincinnatia peracuta Pilsbry & Walker (in Pilsbry) 1889 [Fig. 225]
Spivey's Lake, Navarro County, Texas (Pilsbry, 1889).
Cincinnatia petrifons Thompson 1968 [Figs. 210, 226]
Rock Springs, 6.5 miles north of Apopka, Orange County, Florida (Thompson, 1968).
Cincinnatia ponderosa Thompson 1968 [Figs. 211, 227]
Sanlando Springs, 3.1 miles west of Longwood, Seminole County, Florida

(Thompson, 1968).

Cincinnatia vanhyningi (Vanatta 1934) [Figs. 212, 236]

Seminole Springs, 3.4 miles northeast of Sorrento, Lake County, Florida (Vanatta, 1934; Thompson, 1968).

Cincinnatia wekiwae Thompson 1968 [Figs. 213, 228]

Wekiwa Springs, about five miles northeast of Apopka, Seminole County, Florida (Thompson, 1968).

Genus *Fontelicella* Gregg & Taylor 1965

Subgenus *Fontelicella* s.s.

Fontelicella californiensis Gregg & Taylor 1965 [Fig. 229]

Southern California and northwestern Baja California (Gregg & Taylor, 1965).

Fontelicella deserta (Pilsbry 1916) [Fig. 237]

Washington County, Utah.

Fontelicella intermedia (Tryon 1865) [Fig. 238]

Owyhee River, Malheur County, Oregon (Tryon, 1865i; Gregg & Taylor, 1965).

Fontelicella neomexicana (Pilsbry 1916) [Fig. 239]

In warm springs at Socorro, New Mexico (Pilsbry, 1916a).

Fontelicella pilsbryana (Baily & Baily 1952)

Bear Lake Valley, southeastern Idaho-northeastern Utah (Baily & Baily, 1951; Gregg & Taylor, 1965).

Fontelicella stearnsiana (Pilsbry 1899)

San Francisco Bay region eastward to the Sierra Nevada foothills, California (Gregg & Taylor, 1965).

Subgenus *Natricola* Gregg & Taylor 1965

Fontelicella (Natricola) hendersoni (Pilsbry 1933) [Fig. 240]

Harney Lake basin, Harney County, Oregon (Gregg & Taylor, 1965).

Fontelicella (Natricola) idahoensis (Pilsbry 1933) [Figs. 241, 242]

Snake River, southwestern Idaho (Gregg & Taylor, 1965).

Fontelicella (Natricola) robusta (Walker 1908) [Figs. 230, 243]

Snake River drainage of western Wyoming and southern Idaho; Harney Lake basin, eastern Oregon (Gregg & Taylor, 1965).

Subgenus *Microamnicola* Gregg & Taylor 1965

Fontelicella (Microamnicola) micrococcus Pilsbry (in Stearns) 1893 [Figs. 231, 244]

Amargosa River drainage, in southern Nye County, Nevada; eastern Inyo County and northern San Bernardino County, California (Gregg & Taylor, 1965).

Genus *Marstonia* F.C. Baker 1926[13]

Marstonia agarhecta Thompson 1969 [Figs. 214, 232]
Bluff Creek, Pulaski County, Georgia (Thompson, 1969, 1977).

Marstonia arga Thompson 1977 [Figs. 215, 233]
Tennessee River in the vicinity of the Guntersville Reservoir in northeastern Alabama and from Shoal Creek (Thompson, 1977).

Marstonia castor Thompson 1977 [Figs. 216, 234]
Cedar Creek, Crisp County, Georgia (Thompson, 1977).

Marstonia halcyon Thompson 1977 [Figs. 217, 249]
Lower half of the Ogeechee River system in eastern Georgia (Thompson, 1977).

Marstonia lustrica (Pilsbry 1890)[14] [Figs. 218, 219, 245, 246, 250, 251]
Canada: southern Quebec and Ontario; United States: Maine and New York west through northwestern Pennsylvania, Ohio, northern Indiana and northern Illinois to Iowa and Minnesota (Thompson, 1977).

Marstonia ogmorphaphe Thompson 1977 [Figs. 220, 252]
Owen Springs, Sequatchie, Marion County, Tennessee (Thompson, 1977).

Marstonia olivacea (Pilsbry 1895) [Fig. 247]
Big Spring Creek, Madison County, Alabama (Thompson, 1977).

Marstonia pachyta Thompson 1977 [Figs. 221, 253]
Limestone Creek and Piney Creek, Limestone County, Alabama (Thompson, 1977).

Genus *Notogillia* Pilsbry 1953

Notogillia sathon Thompson 1969 [Figs. 90, 254]
Small streams and springs draining into the Ocmulgee River in south central Georgia (Thompson, 1969).

Notogillia wetherbyi (Dall 1885) [Figs. 255, 260]
Alabama: Decatur and Seminole counties (Clench & Turner, 1956); Florida: northern half of the peninsula west of the St. Johns River and as far north as the Suwannee River system; also in Jackson and Calhoun counties (Thompson, 1968); Georgia: Barbour County (Hubricht, 1963c).

Genus *Orygoceras* Brusina 1882

Orygoceras sp. [Fig. 248]
Roaring Springs, Real County, Texas (Taylor, 1974).

Genus *Pyrgulopsis* Call & Pilsbry 1886

Pyrgulopsis archimedis S.S. Berry 1947 [Fig. 274]
Upper Klamath Lake, near Algoma, Oregon (S.S. Berry, 1947).

Pyrgulopsis letsoni (Walker 1901)[13] [Fig. 261]
Ontario, New York, Ohio and Michigan (F.C. Baker, 1928c; LaRocque, 1968).

Pyrgulopsis nevadensis nevadensis (Stearns 1883) [Figs. 256, 270-272]
Pyramid Lake and Walker's Lake, Nevada (Stearns, 1883b); Upper Klamath Lake, Oregon (Hanna, 1930).
Pyrgulopsis nevadensis paiutica Baily & Baily 1951
Pyramid Lake, Nevada (Baily & Baily, 1951).
Pyrgulopsis ozarkensis Hinkley 1915[13]
North Fork of the White River, above Norfolk, Arkansas (Hinkley, 1915).
Pyrgulopsis scalariformis (Wolf 1869)[13] [Fig. 273]
Shoal Creek, near Florence, Alabama; Illinois River, Tazewell County, and Rock River, Rock Island County, Illinois, as Pleistocene fossils (Wolf, 1869; F.C. Baker, 1928c).

Genus *Rhapinema* Thompson 1969

Rhapinema dacryon Thompson 1969 [Figs. 91, 257, 262]
Chipola River drainage in Jackson County, Florida (Thompson, 1969).

Genus *Spilochlamys* Thompson 1968

Spilochlamys conica Thompson 1968 [Figs. 92, 258, 263]
River systems draining into the Gulf of Mexico in north central Florida, from Levy County north and west to Jackson County (Thompson, 1968).
Spilochlamys gravis Thompson 1968 [Figs. 264, 275, 276]
North central Florida, in the St. Johns drainage system from Palatka south to the Wekiva River (Thompson, 1968).
Spilochlamys turgida Thompson 1969 [Fig. 259]
Small streams and springs draining into the Ocmulgee River in south central Georgia (Thompson, 1969).

Genus *Stiobia* Thompson 1978

Stiobia nana Thompson 1978 [Figs. 265, 297]
Coldwater Spring Run, west of Oxford, Calhoun County, Alabama (Thompson & McCaleb, 1978).

Subfamily Amnicolinae

Genus *Amnicola* Gould & Haldeman 1840[15]

Subgenus *Amnicola* s.s.

Amnicola aldrichi aldrichi (Call & Beecher 1886)[16] [Fig. 277]
Tributary of Black River, Reynolds County, Missouri (Call & Beecher, 1886).
Amnicola aldrichi antroecetes Hubricht 1940
Caves in southwestern Illinois and in eastern and southeastern Missouri (Hubricht, 1940a).

Amnicola aldrichi insolita Hubricht 1940
Springs in southeastern Missouri (Hubricht, 1940a).
Amnicola bakeriana Pilsbry 1917[16]
Oneida Lake, New York (Pilsbry, 1917c).
Amnicola clarkei Pilsbry 1917[16]
Oneida Lake, New York (Pilsbry, 1917c).
Amnicola cora Hubricht 1979 [Fig. 285]
Stream in Foushee Case, three miles west of Locust Grove, Independence County, Arkansas (Hubricht, 1979).
Amnicola dalli dalli (Pilsbry & Beecher 1892) [Figs. 266, 298]
Throughout the northern half of peninsular Florida and west into the Florida panhandle to Leon County (Thompson, 1968).
Amnicola dalli johnsoni Pilsbry 1899 [Figs. 93, 267, 278, 284]
Throughout the northern two-thirds of peninsular Florida, and near Tallahassee (Thompson, 1968).
Amnicola decisa Haldeman 1845[16] [Fig. 287]
Tributaries of the Susquehanna River and in the Schuylkill River (Haldeman, 1845).
Amnicola limosa limosa (Say 1817) [Fig. 268, 286, 288-290, 299]
From the Atlantic coast to as far west as Utah, and from Labrador to Florida (E.G. Berry, 1943).
Amnicola limosa parva Lea 1841
Atlantic and Middle States, including Ohio, Indiana, Illinois, Iowa and Missouri (F.C. Baker, 1928c).
Amnicola missouriensis Pilsbry 1898[16]
Carter County, Missouri (Pilsbry, 1898a).
Amnicola proserpina Hubricht 1940[16]
Spring in St. Louis County and caves in St. Genevieve and Jefferson counties, eastern Missouri (Hubricht, 1940a, 1942).
Amnicola rhombostoma Thompson 1968 [Figs. 269, 300]
In small sand-bottomed streams and rivers draining into the west side of the St. Johns River in Clay and Putnam counties, Florida (Thompson, 1968).
Amnicola stygia Hubricht 1971 [Fig. 291]
Caves in Perry County, Missouri (Hubricht, 1971).

Subgenus *Lyogyrus* Gill 1863

Amnicola (Lyogyrus) browni Carpenter 1872 [Fig. 301]
Massachusetts and Rhode Island (see E.G. Berry, 1943).
Amnicola (Lyogyrus) grana (Say 1822) [Figs. 279, 302]
Atlantic drainage in southeastern Pennsylvania and New Jersey (Walker, 1918b); headwaters of the Pearl River, Mississippi (Hubricht, 1963a).
Amnicola (Lyogyrus) greggi Pilsbry 1935 [Figs. 292, 303]
Cliff Creek canyon, a fork of Hoback canyon, about 29 miles south of Jackson, Wyoming, in the Snake River drainage (Pilsbry, 1935a); also in western Montana and southeastern Idaho (Taylor, 1966b).
Amnicola (Lyogyrus) pilsbryi Walker 1906 [Figs. 293-296, 304]
Wisconsin east to New Philadelphia, Ohio, and south to northern Illinois (F.C.

Baker, 1928c).

Amnicola (Lyogyrus) pupoidea (Gould 1841) [Figs. 280, 305]
Canada, Maine, Connecticut, Massachusetts and the District of Columbia (Binney, 1865d).

Amnicola (Lyogyrus) retromargo Thompson 1968 [Figs. 94, 281, 306]
Occurs in a narrow zone across the neck of the Florida peninsula from the west side of the St. Johns River in Clay and Putnam counties west to Dixie County (Thompson, 1968).

Amnicola (Lyogyrus) walkeri Pilsbry 1898 [Figs. 282, 307, 309]
St. Lawrence River and Great Lake drainages, upper Mississippi drainage, the Canadian Interior basin in the Albany and Winnipeg river systems and in Lake Winnipeg (Clarke, 1973).

Genus *Hauffenia* Pollonera 1898[17]

Hauffenia micra (Pilsbry & Ferriss 1906 [Fig. 308]
Found in drift debris of the Guadalupe River, near New Braunfels, Texas (Pilsbry & Ferriss, 1906).

Genus *Horatia* Bourguignat 1887[17]

Horatia nugax (Pilsbry & Ferriss 1906) [Fig. 316]
Found in drift debris of the Guadalupe River, near New Braunfels, Texas (Pilsbry & Ferriss, 1906).

Subfamily Fontigentinae

Genus *Fontigens* Pilsbry 1933

Fontigens binneyana (Hannibal 1912)[18]
"Ohio" (Lea, 1841, for *"Paludina" obtusa*, preoccupied; renamed *binneyana*).

Fontigens cryptica Hubricht 1963 [Fig. 315]
Small spring in Clarke County, Indiana (Hubricht, 1963b).

Fontigens holsingeri Hubricht 1976 [Fig. 311]
Streams in caves in Randolph and Pocahontas counties, West Virginia (Hubricht, 1976).

Fontigens nickliniana (Lea 1838) [Figs. 283, 319]
In cool, shallow springs from Pennsylvania to Wisconsin and from Ontario to Alabama (E.G. Berry, 1943).

Fontigens orolibas Hubricht 1957 [Figs. 312, 313]
Springs in the Shenandoah National Park and along the Blue Ridge Parkway, Virginia (Hubricht, 1957).

Fontigens tartarea Hubricht 1963 [Fig. 314]
Stream in Organ Cave, Greenbrier County, West Virginia (Hubricht, 1963b).

Fontigens turritella Hubricht 1976 [Fig. 310]
Caves in Greenbrier County, West Virginia (Hubricht, 1976).

Fontigens weberi Pilsbry 1950[19]
West Lake, Cape Sable, Florida (Pilsbry, 1950a).

Incertae Sedis

"Bythinella" hemphilli Pilsbry 1890 [Fig. 320]
Near Kentucky Ferry, Snake River, Washington (Pilsbry, 1890e)
"Cochliopa" texana Pilsbry 1935 [Fig. 317]
Phantom Lake, near Toyahvale, Reeves County, Texas (Pilsbry, 1935a).
"Paludestrina" bottimeri Walker 1925 [Fig. 318]
Glen Echo, Montgomery County, Maryland (Walker, 1925a).

Family POMATIOPSIDAE

Genus *Pomatiopsis* Tryon 1862

Pomatiopsis binneyi Tryon 1863 [Fig. 321]
Bolinas, Marin County, California (Tryon, 1863a); Mt. Tamalpais, Marin County (Davis, 1967).
Pomatiopsis californica Pilsbry 1899 [Fig. 322]
San Francisco and Oakland, California (Pilsbry, 1899); Bolinas Bay, Marin County, California (Davis, 1967).
Pomatiopsis chacei Pilsbry 1937
Near Klamath, Humboldt County, California (Pilsbry, 1937a); Crescent City, Del Norte County, and Wilson Creek, California (E.G. Berry, 1947b).
Pomatiopsis cincinnatiensis (Lea 1840) [Fig. 323]
Tennessee and southwest Virginia to southern Michigan, Illinois and Iowa (Burch & Van Devender, 1980).
Pomatiopsis hinkleyi Pilsbry 1896[20] [Fig. 324]
The original localities (near Florence, Alabama) are now covered by the water impounded by Wilson Dam; also found at a spring near Ashland City, Tennessee, and near Eberhardt, South Carolina (Hubricht, 1960).
Pomatiopsis lapidaria (Say 1817) [Fig. 325]
Widely distributed in the eastern United States, with occasional occurrences west to northern Texas and New Mexico (Burch & Van Devender, 1980).

Family THIARIDAE

Genus *Melanoides* Olivier 1904

Melanoides tuberculata (Müller 1774) [Fig. 327]
Much of Africa and the eastern Mediterranean countries, throughout India, Southeast Asia, Malaysia and southern China, north to the Ryukyu Islands of Japan, south and east through many of the Pacific islands to northern Australia and the New Hebrides (Pace, 1973); introduced into Florida, Texas and Arizona (see Dundee, 1974; Murray, 1964, 1976).

Genus *Thiara* Röding 1798

Thiara granifera (Lamarck 1822) [Fig. 326]
Madagascar and India eastward throughout Malaysia and the Philippines to the Society Islands and north to the Ryukyu Islands and Hawaii (Pace, 1973); introduced into Florida (Abbott, 1952) and Texas (Murray, 1964).

Family PLEUROCERIDAE[21, 22]

Genus *Elimia* H. & A. Adams 1854[23]

Elimia acuta group

Elimia acuta acuta (Lea 1831)
Tributaries of the Tennessee River in southern Tennessee and northern Alabama (Goodrich, 1930a, 1941b).
Elimia acuta clavula (Lea 1868)
Tributaries of the Tennessee River in Madison County, Tennessee, and Jackson County, Alabama (Goodrich, 1940d).
Elimia comma (Conrad 1834)
Springs and spring branches of the Black Warrior River in Jefferson and Blount counties, Alabama (Goodrich, 1941b).

Elimia boykiniana group

Elimia boykiniana boykiniana (Lea 1840) [Fig. 328]
Chattahoochee and Flint rivers, Georgia (Goodrich, 1942b).
Elimia boykiniana albanyensis (Lea 1864)
Flint River, Georgia, and tributaries; Uchee Creek, Russell County, Alabama (Goodrich, 1942b).
Elimia boykiniana viennaensis (Lea 1862) [Fig. 329]
Flint River and creeks of western Georgia; Uchee Creek, Russell County, Alabama (Goodrich, 1942b).
Elimia clenchi (Goodrich 1924) [Fig. 330]
Tributaries of Choctawhatchee and Chipola rivers, Alabama and Florida; branches of Conecuh River, Covington County, Alabama (Goodrich, 1942b).
Elimia ucheensis (Lea 1862) [Fig. 346]
Uchee and Little Uchee creeks, Russell County, Alabama (Goodrich, 1942b).

Elimia carinifera group

Elimia bellacrenata (Haldeman 1841) [Fig. 345]
Tributary springs, spring-fed brooks and creeks of the Cahaba River (Goodrich, 1941c).
Elimia carinifera (Lamarck 1822) [Fig. 331]
Springs, brooks, creeks and occasionally rivers of the Alabama River drainage

basin, from north Georgia to Monroe County, Alabama; parts of the Tennessee River system in the vicinity of Chattanooga, Hamilton County, Tennessee (Goodrich, 1941b,c).

Elimia catenaria group

Elimia arachnoidea arachnoidea (Anthony 1854) [Fig. 332]
Small streams of East Tennessee (Goodrich, 1940d).

Elimia arachnoidea spinella (Lea 1862) [Fig. 333]
Small streams of Lee and Scott counties, Virginia, and Claiborne County, Tennessee (Goodrich, 1940d).

Elimia athearni (Clench & Turner 1956)
Central part of the Chipola River system (Clench & Turner, 1956).

Elimia brevis (Reeve 1860) [Figs. 347, 348]
Middle and lower reaches of the Coosa River, Alabama (Goodrich, 1944d).

Elimia capillaris (Lea 1861) [Fig. 334]
Coosa River, Floyd County, Georgia, to shoals of Chilton and Coosa counties, Alabama; in the Etowah River, at Rome, Georgia, and creeks to Talladega County, Alabama (Goodrich, 1944d).

Elimia catenaria catenaria (Say 1822) [Fig. 335]
Springs of eastern South Carolina, possibly in streams southward to the Savannah River (Goodrich, 1942b).

Elimia catenaria dislocata (Reeve 1861) [Figs. 336, 349]
Streams of Durham, Burke, Franklin, Madison and Mecklenburg counties, North Carolina; headstreams in South Carolina, Greenville County, Virginia (Goodrich, 1942b).

Elimia catenaria inclinans (Lea 1862)
Flint River and tributaries, Georgia (Goodrich, 1942b).

Elimia catenaria postelli (Lea 1858) [Fig. 337]
Altamaha, Ogeechee and Canoochee rivers, and possibly Savannah River, Georgia (Goodrich, 1942b).

Elimia catenaria vanhyningiana (Goodrich 1921) [Fig. 338]
Lake, Marion and Orange counties, Florida (Goodrich, 1942b).

Elimia cochilaris (Lea 1868)
Found in springs and spring brooks of the Little Cahaba River in Bibb, Jefferson and Tuscaloosa counties, Alabama (Goodrich, 1941c).

Elimia comalensis comalensis (Pilsbry 1890) [Fig. 339]
Drainage of Guadeloupe River, Texas; ? basin of Brazos River (Goodrich, 1942b).

Elimia comalensis fontinalis (Pilsbry & Ferriss 1906) [Fig. 350]
Comal Creek and its springs, New Braunfels, Comal County, Texas (Goodrich, 1942b).

Elimia crenatella (Lea 1860) [Fig. 340]
Coosa River Basin: in the Coosa River from St. Clair to Chilton County, Alabama, and in creeks of St. Clair, Etowah and Talladega counties (Goodrich, 1944d).

Elimia edgariana (Lea 1841) [Fig. 341]
Streams of Cumberland, Duck and Elk rivers, Tennessee (Goodrich, 1940d).

Elimia floridensis (Reeve 1860)
Upper reaches of the St. Johns and Hillsborough rivers in central Florida, north and west to the Apalachicola River and the upper reaches of Holmes Creek (Clench & Turner, 1956).

Elimia fusiformis (Lea 1841) [Figs. 351, 352]
Coosa River: Weduska Shoals, Shelby County, to Wetumpka, Elmore County, Alabama (Goodrich, 1944d).

Elimia impressa (Lea 1841) [Fig. 342]
Coosa River: Leoto Shoals, St. Clair County, to rapids of Coosa County, Alabama (Goodrich, 1944d).

Elimia nassula (Conrad 1834) [Fig. 353]
Springs and spring branches of Madison and Colbert counties, Alabama (Goodrich, 1940d).

Elimia perstriata perstriata (Lea 1852)
Springs and small streams of north Alabama (Goodrich, 1940d).

Elimia perstriata crispa (Lea 1862)
Lawrence and Madison counties, Alabama (Goodrich, 1940d).

Elimia perstriata decampi (Lea 1863)[24] [Fig. 343]
Madison County, Alabama (Goodrich, 1940d).

Elimia plicatastriata (Wetherby 1876)
Small branches of the Cumberland River, Tennessee and Kentucky; Big Richland Creek of the Tennessee River, Humphreys County, Tennessee (Goodrich, 1940d).

Elimia porrecta (Lea 1863) [Figs. 368, 369]
Springs and streams of Claiborne County, Tennessee (Goodrich, 1940d).

Elimia pupaeformis (Lea 1864) [Fig. 354]
Coosa River, from the vicinity of Riverside, St. Clair County, to Wetumpka, Alabama (Goodrich, 1944d).

Elimia striatula (Lea 1842) [Fig. 344]
In the Tennessee River system at springs in Monroe County and in a reservoir near Cleveland, Bradley County, Tennessee (Goodrich, 1940d); in the Alabama River system at Coahulla Creek, Whitfield County, Georgia (Goodrich, 1941b).

Elimia strigosa (Lea 1841) [Fig. 355]
Small streams near Knoxville, Knox County, Tennessee (Goodrich, 1940d).

Elimia teres (Lea 1841) [Fig. 356]
Small streams of Walden Ridge, Tennessee, flowing eastward (Goodrich, 1940d).

Elimia troostiana (Lea 1838) [Fig. 357]
Mossy Creek, Jefferson County, Tennessee (Goodrich, 1940d).

Elimia carinocostata group

Elimia bentoniensis (Lea 1862)
Coosa River Basin, in small streams of Calhoun, St. Clair and Talladega counties, Alabama (Goodrich, 1941b, 1944d).

Elimia carinocostata (Lea 1845)
Tributaries of Black Warrior River (Goodrich, 1941b); upper Cahaba River to

Nunley Ford, Shelby County, and upper Little Cahaba River, Alabama (Goodrich, 1941c); Coosa River Basin, in creeks from Whitfield County, Georgia, to Elmore County, Alabama (Goodrich, 1944d).

Elimia curvicostata (Reeve 1861) [Fig. 358]

Streams of western Georgia and Florida; rivers and creeks of southeastern Alabama (Goodrich, 1942b).

Elimia dickinsoni (Clench & Turner 1956) [Fig. 359]

Upper tributaries of the Chipola River in Florida and Alabama, and the tributaries of the Choctawhatchee immediately to the west (Clench & Turner, 1956).

Elimia induta (Lea 1862)

Flint River basin, Crisp and Dooly counties, Georgia (Goodrich, 1942b).

Elimia ebenum group

Elimia ebenum ebenum (Lea 1841) [Fig. 370]

Cumberland River above the Falls; Smith's Shoals, Pulaski County, Kentucky; springs and small streams of the river downstream to Dickson County, Tennessee (Goodrich, 1940d).

Elimia ebenum emeryensis (Lea 1864)

In branches of the Cumberland River in eastern Kentucky and Tennessee (Goodrich, 1940d).

Elimia gerhardti group

Elimia annettae (Goodrich 1941) [Fig. 360]

Cahaba River, Lily Shoals, Bibb County, Alabama (Goodrich, 1941a).

Elimia cahawbensis cahawbensis (Lea 1861) [Fig. 371]

Headwaters and small streams and creeks of the Cahaba River; in a few tributaries of the Black Warrior River; Waxahatchee Creek and branches of the Coosa River basin (Goodrich, 1941b).

Elimia cahawbensis fraterna (Lea 1864)

A branch of the Cahaba River in Bibb County, and in the Black Warrior River basin at Murphy's Creek, Blount County, Alabama (Goodrich, 1941c).

Elimia flava (Lea 1862) [Fig. 372]

Confined to the Tallapoosa River and its tributaries (Goodrich, 1941b).

Elimia gerhardti (Lea 1862) [Figs. 361, 362, 373]

Coosa River basin, from north Georgia to the lower tributaries of the Coosa River (Goodrich, 1944d).

Elimia varians (Lea 1861)

Cahaba River, Bibb County, Alabama, from Pratt's Ferry to seven miles below Centerville (Goodrich, 1941c).

Elimia hartmaniana group

Elimia hartmaniana (Lea 1861) [Fig. 363]

Coosa River: St. Clair to Elmore County, Alabama (Goodrich, 1944d).

Elimia macglameriana (Goodrich 1936) [Fig. 374]
Coosa River: Yancy's Landing, Floyd County, Georgia, to Riddle's Bend, St. Clair County, Alabama (Goodrich, 1944d).
Elimia pygmaea (Smith (in Goodrich) 1936) [Fig. 375]
Three Island Shoals, Coosa River, Talladega County, Alabama (Goodrich, 1944d).

Elimia haysiana group

Elimia alabamensis (Lea 1861) [Figs. 376, 377]
In middle sections of the Coosa River, and in creeks of Talladega County, Alabama (Goodrich, 1944d).
Elimia clausa (Lea 1861) [Fig. 364]
Coosa River, in shoals of St. Clair County, Alabama (Goodrich, 1944d).
Elimia haysiana (Lea 1843) [Figs. 378, 379]
Lower part of the Coosa River (Goodrich, 1944d).
Elimia pupoidea (Anthony 1854) [Figs. 380, 381]
Cahaba and Black Warrior rivers, and in the vicinity of Selma, Alabama River, Alabama (Goodrich, 1941c).

Elimia hydei group

Elimia hydei (Conrad 1834) [Fig. 365]
Confined to the Black Warrior River, Alabama, and its branches (Goodrich, 1941b).

Elimia laqueata group

Elimia costifera (Reeve 1861)
Tributaries of the Ohio River in Kentucky and Illinois (Goodrich, 1940d).
Elimia curreyana (Lea 1841)
Green River, Kentucky, and tributaries; streams of Cumberland River, middle Tennessee (Goodrich, 1940d).
Elimia interveniens (Lea 1862)
Tributaries of the Tennessee River in north Alabama (Goodrich, 1940d).
Elimia laqueata laqueata (Say 1829) [Figs. 366, 367, 391]
Green River and tributaries, Kentucky; tributaries of middle parts of Cumberland River, Tennessee; Duck River and branches, Tennessee; tributaries of the Tennessee River, Tennessee and Alabama (Goodrich, 1940d).
Elimia laqueata castanea (Lea 1841) [Fig. 382]
Headwaters of the Duck River, Tennessee (Goodrich, 1940d).
Elimia laqueata costulata (Lea 1841) [Fig. 392]
Green River of Kentucky and branches; branches of the Duck River, Tennessee (Goodrich, 1940d).
Elimia laqueata tortum (Lea 1845)
Elk River drainage in Lynn Creek, Giles County, Tennessee, and Richland

Creek, Lawrence County, Tennessee; headwaters of Big Creek, Lawrence County, Tennessee (Goodrich, 1930a, 1940d).

Elimia paupercula (Lea 1862) [Fig. 383]
Creeks of northern Alabama (Goodrich, 1940d).

Elimia pybasi (Lea 1862)
Springs and streams of northern Alabama (Goodrich, 1941b).

Elimia livescens group

Elimia livescens livescens (Menke 1830) [Fig. 393]
St. Lawrence River drainage from the Great Lakes to Lake Champlain and Quebec; tributaries of the Ohio River, east of Scioto River in Ohio; Wabash River and branches, west to the Illinois River; through the Erie Canal it has invaded the Hudson River basin (Goodrich, 1940d, 1945).

Elimia livescens gracilior (Lea 1861)
Lakes of Summit and Stark counties, Ohio (Goodrich, 1939d).

Elimia livescens haldemani (Tryon 1865) [Fig. 384]
Lake Erie; ? Lake Champlain (Goodrich, 1939d).

Elimia mutabilis group

Elimia mutabilis mutabilis (Lea 1862) [Fig. 394]
Streams of western Georgia and Florida, and southern Alabama; also a few creeks and springs of Alabama within the Alabama River system (Goodrich, 1942b).

Elimia mutabilis timidus (Goodrich 1942)
Spring two miles northwest of Hawkinsville, Pulaski County, Georgia, in the basin of the Altamaha River (Goodrich, 1942b).

Elimia taitiana (Lea 1841) [Fig. 395]
Branch of Sepulga River, Conecuh County, Alabama; small streams of the Alabama River system, Sumpter, Marengo, Monroe and Wilcox counties, Alabama (Goodrich, 1942b).

Elimia olivula group

Elimia bellula (Lea 1861) [Fig. 396]
In the middle part of the Coosa River, and in Yellowleaf Creek, Shelby County, and Choccolocco Creek, Talladega County, Alabama (Goodrich, 1944d).

Elimia chiltonensis (Goodrich, 1941) [Fig. 397]
Waxahatchee Creek of Chilton and Shelby counties, Alabama, and three of its tributaries, and in Weguska Creek, Coosa County, Alabama (Goodrich, 1941a).

Elimia cylindracea (Conrad 1834) [Fig. 398]
Tombigbee River from Columbus, Mississippi, to near its mouth, and in the lower part of the Black Warrior River, Alabama (Goodrich, 1936).

Elimia gibbera (Smith 1936) [Fig. 399]
Coosa River, shoals of St. Clair County, Alabama (Goodrich, 1944d).

Elimia lachryma (Reeve 1861) [Fig. 400]
Coosa River: Gilbert's Ferry, Etowah County, to near Childersburg, Talladega County, Alabama (Goodrich, 1944d).

Elimia laeta (Jay 1839) [Fig. 401]
Coosa River, from Cherokee County to Elmore County, Alabama (Goodrich, 1944d).

Elimia olivula (Conrad 1834) [Fig. 402]
Alabama River and the lower Cahaba River (below the Falls Line), Alabama (Goodrich, 1941c).

Elimia pilsbryi (Goodrich 1927)[25] [Fig. 385]
Coosa River: Hall's Island, Talladega County, to mouth of Yellowleaf Creek of Chilton County, Alabama (Goodrich, 1944d).

Elimia showalteri (Lea 1860) [Fig. 386]
Cahaba River, from Lily Shoals to two miles east of Harrisburg, Bibb County, Alabama (Goodrich, 1941c).

Elimia variata (Lea 1861) [Fig. 387]
Cahaba River and tributaries in Shelby and Bibb counties, Alabama; Little Cahaba River, Jefferson County (Goodrich, 1941c).

Elimia semicarinata group

Elimia semicarinata (Say 1829)
Tributaries of Ohio River, Scioto River, Ohio, to Big Blue River, Indiana; Licking River to Salt River in Kentucky; two creeks of Green River of Kentucky (Goodrich, 1940d).

Elimia simplex group

Elimia aterina (Lea 1863) [Fig. 388]
Springs and small streams of Claiborne and Hancock counties, Tennessee (Goodrich, 1940d).

Elimia clavaeformis (Lea 1841) [Figs. 403-405]
Tributaries of the upper Tennessee River in Virginia, Tennessee and North Carolina (Goodrich, 1940d).

Elimia simplex (Say 1825) [Fig. 406]
Headwaters of the Tennessee River system in Virginia, Tennessee and North Carolina; Beaver Fork of the Bluestone River, which is a tributary of the Kanawha River in Mercer County, West Virginia (Goodrich, 1940d).

Elimia vanuxemiana group

Elimia bullula (Lea 1861) [Figs. 389, 390]
Coosa River, Cherokee County, Alabama, to near the Narrows, Coosa County, and in five tributaries between these points (Goodrich, 1944d).

Elimia caelatura caelatura (Reeve 1860) [Fig. 407]
Coosa River Basin, from Georgia headwaters to side streams of Talladega

County, Alabama (Goodrich, 1944d).

Elimia caelatura excellens (Goodrich, 1935) [Fig. 408]

Known from three streams in the Alabama River system of northwestern Georgia and northeastern Alabama (Goodrich, 1941b).

Elimia caelatura georgiana (Lea 1862) [Fig. 409]

Chattooga River, Georgia (Goodrich, 1941b).

Elimia caelatura infuscata (Lea 1862) [Fig. 415]

Small streams of the Alabama River system in Bartow, Floyd, Gordon and Murray counties, Georgia, and Cherokee, Etowah and St. Clair counties, Alabama (Goodrich, 1941b).

Elimia caelatura lecontiana (Lea 1841)

Creeks of the Alabama River system in northwestern Georgia to northeastern Alabama (Goodrich, 1941b).

Elimia caelatura luteocella (Lea 1868) [Fig. 410]

Small streams of the Alabama River system in northwestern Georgia, northeastern Alabama and Talladega County, Alabama (Goodrich, 1941b).

Elimia caelatura stearnsiana (Call 1886) [Figs. 411, 416]

Small streams of the Alabama River from north Georgia to Calhoun, Shelby and Talladega counties, Alabama (Goodrich, 1941b).

Elimia fascinans (Lea 1861) [Fig. 417]

Coosa River Basin, in creeks from Calhoun to Coosa County, and occasionally in the Coosa River (Goodrich, 1944d).

Elimia jonesi (Goodrich 1936) [Fig. 418]

Coosa River: Ten Island Shoals, St. Clair County, to the Bar, Chilton County, Alabama (Goodrich, 1944d).

Elimia vanuxemiana (Lea 1843) [Figs. 419-422]

Coosa River Basin: in the Coosa River at Etowah County and downstream, and in the mouths of a few tributaries of the same range (Goodrich, 1944d).

Elimia virginica group

Elimia proxima (Say 1825) [Fig. 412]

Highlands of North and South Carolina. *Elimia proxima* may be a composite group, those in the Atlantic drainage having been derived from *E. symmetrica* and those in the Tennessee drainage from *E. simplex* (Goodrich, 1942b, 1950). Say (1825) originally described *E. proxima* from specimens from three localities: a small brook which discharges into the Catawba River near Landsford, South Carolina [Atlantic drainage], and in the warm springs and in the French Broad River, both in Buncombe County, North Carolina [Tennessee drainage].

Elimia symmetrica (Haldeman 1841) [Figs. 423-425]

Southern Virginia; North Carolina (Goodrich, 1942b).

Elimia virginica (Say 1817) [Fig. 413]

Connecticut River, Massachusetts and Connecticut, to Virginia; ? also North Carolina; westward through the Erie Canal to Monroe County, New York, in the Great Lakes basin (Goodrich, 1942b).

Elimia clara (group ?)

Elimia clara (Anthony 1854) [Fig. 414]
Cahaba River, Alabama, and its tributaries (Goodrich, 1941c).
Elimia ampla (Anthony 1854)[26] [Fig. 426]
Cahaba River, Alabama, at Centerville and Lily Shoals (Goodrich, 1941c).

Elimia potosiensis (group ?)

Elimia potosiensis potosiensis (Lea 1841) [Fig. 427]
Upland streams of a few Missouri counties (Goodrich, 1939e).
Elimia potosiensis crandalli (Pilsbry 1890)
Known only from Mammoth Springs, Fulton County, Arkansas (Goodrich, 1939e).
Elimia potosiensis ozarkensis (Call 1886) [Fig. 458]
In springs of Shannon, Carter, Washington; Dent and Camden counties, Missouri (Goodrich, 1939e).
Elimia potosiensis plebius (Gould 1850) [Figs. 459, 460]
Common in rivers and creeks of the Ozarkian area of Missouri and Arkansas, and in Oklahoma counties bordering Missouri (Goodrich, 1939e).

group ?

Elimia interrupta (Haldeman 1840) [Fig. 428]
Hiwassee River and its streams, North Carolina and eastern Tennessee (Goodrich, 1940d).

(? hybrid)

Elimia ornata (Lea 1868)[27]
Coosa River Basin, confined to a few miles of the Connesauga River, Georgia, and nearby tributaries (Goodrich, 1944d).

Genus *Gyrotoma* Shuttleworth 1845[28,29]

Gyrotoma excisum (Lea 1843) [Figs. 431-440]
Coosa River, Alabama, in Chilton, Coosa, Elmore, Shelby, St. Clair and Talladega counties.
Gyrotoma lewisi (Lea 1869) [Fig. 441]
Coosa River, Alabama: Fort William Shoals, Talladega County, and Three Island Shoals, Wilsonville, Shelby County (Goodrich, 1924a, 1944d).
Gyrotoma pagodum (Lea 1845) [Figs. 442, 443]
Coosa River, Alabama: The Bar, Chilton County, to Wetumpka, Elmore County (Goodrich, 1944d).

Gyrotoma pumilum (Lea 1860) [Figs. 444, 445]
Coosa River, Alabama: Fort William Shoals, Talladega County, and Weduska Shoals, Shelby County, to Wetumpka (Goodrich, 1924a, 1944d).
Gyrotoma pyramidatum Shuttleworth 1845 [Fig. 446]
Coosa River, Alabama: Ten Island Shoals, St. Clair County, to the mouth of Yellowleaf Creek, Shelby County (Goodrich, 1944d).
Gyrotoma walkeri Smith (in Goodrich) 1924 [Fig. 447]
Coosa River, Alabama: Weduska Shoals, Shelby County, to Butting Ram Shoals, Coosa County (Goodrich, 1944d).

Genus *Io* Lea 1831

Io fluvialis (Say 1825)[30] [Figs. 429, 430, 461-465]
Tennessee River and several of its main tributaries in western Virginia and eastern Tennessee (Clinch, French Broad, Holston, Nolichucky and Powell rivers).

Genus *Juga* H. & A. Adams 1854

Subgenus *Juga* s.s.

Juga hemphilli hemphilli (Henderson 1935) [Fig. 454]
Near Portland, Oregon (Henderson, 1935a).
Juga hemphilli dallesensis (Henderson 1935) [Fig. 455]
Mill Creek, The Dalles, Oregon (Henderson, 1935a).
Juga plicifera (Lea 1838) [Fig. 448]
Larger streams of Oregon and Washington (Goodrich, 1942d).
Juga silicula (Gould 1847) [Fig. 449]
Streams of Oregon and Washington (Goodrich, 1942d).

Subgenus *Calibasis* Taylor 1966

Juga (Calibasis) acutifilosa acutifilosa (Stearns 1890) [Fig. 450]
Shasta and Lassen counties, California (Goodrich, 1942d).
Juga (Calibasis) acutifilosa pittensis (Henderson 1935)
Fall River, Shasta County, California (Henderson, 1935a).
Juga (Calibasis) acutifilosa siskiyouensis (Pilsbry 1899)
Siskiyou County, California (Goodrich, 1942d).
Juga (Calibasis) occata (Hinds 1844) [Fig. 451]
Sacramento and San Joaquin rivers, California (Goodrich, 1942d).

Subgenus *Oreobasis* Taylor 1966

Juga (Oreobasis) bulbosa (Gould 1847) [Fig. 452]
Lower Columbia River in Oregon and Washington and several of its tributaries

(Deschutes and Owyhee rivers) (Pilsbry, 1899f).

Juga (Oreobasis) interioris (Goodrich 1944) [Fig. 466]

Badger Creek, Bitner Ranch, Washoe County, and in the outlet of artesian wells, nine miles west of Gerlach, Washoe County, Nevada (Goodrich, 1944a).

Juga (Oreobasis) laurae (Goodrich 1944) [Fig. 467]

Found in a spring west of Home Camp and in Boulder Springs, both in Long Valley, Nevada, and in springs of Grasshopper Valley, Lassen County, California (Goodrich, 1944a).

Juga (Oreobasis) nigrina (Lea 1856) [Fig. 453]

Head streams and river tributaries of Oregon and northern California (Goodrich, 1942d).

Genus *Leptoxis* Rafinesque 1819

Subgenus *Leptoxis* s.s.[32]

Leptoxis ampla (Anthony 1855) [Figs. 456, 457]

Cahaba River, Alabama, and some of its tributaries (Goodrich, 1941b).

Leptoxis clipeata (Smith (in Goodrich) 1922) [Fig. 468]

Coosa River, Alabama, from near Riverside, St. Clair County, to Butting Ram Shoals (Goodrich, 1944d).

Leptoxis compacta (Anthony 1854) [Figs. 469, 470]

Mostly confined to the middle parts of the Cahaba River, but taken at two upstream localities and in Buck Creek, Shelby County, Alabama (Goodrich, 1941b).

Leptoxis foremani (Lea 1843) [Figs. 471, 472]

Coosa River, Alabama, from Three Island Shoals, Talladega County, to Butting Ram Shoals (Goodrich, 1944d).

Leptoxis formosa (Lea 1860)

Coosa River, from the head streams in northwestern Georgia to Coosa County, Alabama; Terrapin Creek, Cherokee County, Alabama (Goodrich, 1941b, 1944d).

Leptoxis ligata (Anthony 1860) [Fig. 473]

Coosa River, Alabama, from Weduska Shoals, Shelby County, to Wetumpka (Goodrich, 1944d).

Leptoxis lirata (Smith (in Goodrich) 1922)[33]

Three Island and Fort William shoals, Coosa River, Talladega County, Alabama (Goodrich, 1922).

Leptoxis melanoides (Conrad 1834) [Fig. 474]

Black Warrior River (Goodrich, 1922).

Leptoxis occultata (Smith (in Goodrich) 1922) [Fig. 475]

Coosa River, Alabama, confined to the shoals bordering Chilton and Coosa counties (Goodrich, 1944d).

Leptoxis picta (Conrad 1834) [Fig. 476]

Alabama River from the Coosa River to Clairborne, Monroe County, Alabama; in the Coosa River only as far up as the gravel bars below the last series of rapids below Wetumpka (Goodrich, 1922, 1944d).

Leptoxis plicata (Conrad 1834) [Fig. 477]
Headwaters of the Black Warrior River, and Valley Creek, Jefferson County, Alabama (Goodrich, 1922, 1941b).

Leptoxis praerosa (Say 1821) [Figs. 478-482]
Ohio River, below Cincinnati, Ohio, to Elizabethtown, Illinois, together with a few tributaries; Cumberland River and branches; Duck River, Coffee County, Tennessee, to its mouth; Tennessee River and the lower parts of its tributaries (Goodrich, 1940d).

Leptoxis showalteri (Lea 1860) [Fig. 483]
Coosa River, Alabama, from Ten Island Shoals, St. Clair County, to Fort William and Peckerwood Shoals, Talladega County (Goodrich, 1944d).

Leptoxis taeniata (Conrad 1834) [Figs. 484-486]
Alabama River and the Coosa River and its tributaries, Alabama, and into the Cahaba River for a short distance (Goodrich, 1922, 1944d).

Leptoxis umbilicata (Wetherby 1876) [Fig. 528]
Stone's River, Red River, and Ringgold Creek of the Cumberland River, all in Tennessee; Elk River, Franklin County, Tennessee (Goodrich, 1940d).

Leptoxis vittata (Lea 1860) [Fig. 487]
Coosa River, Alabama, from The Bar, Chilton County, to Wetumpka (Goodrich, 1922).

Subgenus *Mudalia* Haldeman 1840

Leptoxis (Mudalia) arkansensis (Hinkley 1915) [Fig. 488]
White River, Baxter County, Arkansas, and North Fork of the White River, east of Richville, Douglas County, Missouri (Goodrich, 1939e).

Leptoxis (Mudalia) carinata carinata (Bruguière 1792) [Figs. 489-492]
New York to North Carolina; ? also South Carolina (Goodrich, 1942b).

Leptoxis (Mudalia) carinata nickliniata (Lea 1841) [Fig. 493]
Hot Springs, Bath County, Virginia (Goodrich, 1942b).

Leptoxis (Mudalia) dilatata (Conrad 1835) [Fig. 494]
Kanawha River, West Virginia; its head streams and branches (Goodrich, 1940d).

Leptoxis (Mudalia) minor (Hinkley 1912) [Fig. 495]
Tennessee River at Muscle Shoals, Lauderdale County, Alabama (Goodrich, 1940d).

Leptoxis (Mudalia) trilineata (Say 1829) [Figs. 496, 497]
Ohio River, Cincinnati, Ohio, to Louisville, Kentucky; Little Miami River, Ohio, near its mouth; Five-mile Creek, Campbell County, Kentucky (Goodrich, 1940d).

Leptoxis (Mudalia) virgata (Lea 1841) [Figs. 498-500]
Holston River and its forks, Sullivan County to Knox County, Tennessee; Tennessee River, Knox County, Tennessee, to Jackson County, Alabama; Hiwassee River, North Carolina (Goodrich, 1940d).

Subgenus *Athearnia* Morrison 1971

Leptoxis (Athearnia) crassa crassa (Haldeman 1841)[34] [Fig. 501]
Eastern Tennessee: Powell River, near its mouth, and the Clinch River in Anderson, Knox and Roane counties (Goodrich, 1940d).

Leptoxis (Athearnia) crassa anthonyi (Redfield 1854) [Fig. 502]
Tennessee River, Knox County, Tennessee, to Lauderdale County, Alabama; lower French Broad and Clinch rivers, eastern Tennessee; Elk River, Alabama; smaller tributaries of the Tennessee River from the Little Tennessee River, Tennessee, to Limestone County, Alabama (Goodrich, 1940d).

Genus *Lithasia* Haldeman 1840

Subgenus *Lithasia* s.s.

Lithasia geniculata geniculata (Haldeman 1840) [Figs. 503, 504]
Cumberland River, above Burnside, Pulaski County, Kentucky, to points below Nashville, Davidson County, Tennessee; branches in Tennessee; Duck River, Maury County to its mouth (Goodrich, 1940d).

Lithasia geniculata fuliginosa (Lea 1841) [Fig. 505]
Tennessee: Duck River, Bedford County, to below Maury County; Buffalo River; Harpeth River; Red River, Robertson County (Goodrich, 1940d).

Lithasia geniculata pinguis (Lea 1852)[35] [Fig. 506]
Tennessee: Caney Fork and branches; Duck River, Coffee County (Goodrich, 1940d).

Lithasia obovata (Say 1829)[35] [Figs. 507-510)
Ohio River and tributaries, in Pennsylvania, Ohio, Indiana, Illinois, Kentucky and Tennessee.

Lithasia salebrosa salebrosa (Conrad 1834)[36] [Fig. 511]
Tennessee River and Cypress Creek, Lauderdale County, Alabama; lower Cumberland River, Montgomery County, Tennessee, to Trigg County, Kentucky (Goodrich, 1940d).

Lithasia salebrosa florentiana (Lea 1841) [Fig. 512]
Tennessee River, Muscle Shoals, Alabama, and a near-by tributary; Elk River, Tennessee and Alabama (Goodrich, 1940d).

Lithasia salebrosa subglobosa (Lea 1861) [Fig. 513]
Tennessee River, Muscle Shoals, Alabama (Goodrich, 1940d).

Subgenus *Angitrema* Haldeman 1841

Lithasia (Angitrema) armigera (Say 1821) [Fig. 514]
Lower Ohio River, lower Wabash River; Cumberland River from above Burnside, Pulaski County, Kentucky, to branches in Trigg County, Kentucky; Tennessee River in the vicinity of Florence, Lauderdale County, Alabama (Goodrich, 1940d).

Lithasia (Angitrema) curta Lea 1868 [Fig. 515]
Tennessee River, Muscle Shoals; Shoals Creek, Lauderdale County, Alabama

(Goodrich, 1940d).

Lithasia (Angitrema) duttoniana (Lea 1841) [Fig. 516]

Tennessee: Duck River, Bedford County to Humphreys County; two tributaries in Bedford County (Goodrich, 1940d).

Lithasia (Angitrema) hubrichti Clench 1956

Big Black River, Mississippi (Clench, 1965a).

Lithasia (Angitrema) jayana (Lea 1841) [Fig. 517]

Forks of Cumberland River; Caney Fork, Tennessee, near mouth.

Lithasia (Angitrema) lima (Conrad 1834) [Figs. 518, 519]

Elk River, Tennessee and Alabama; branch of Elk River in Franklin County, Tennessee; Tennessee River, Alabama, Muscle Shoals and three near-by creeks (Goodrich, 1940d).

Lithasia (Angitrema) verrucosa (Rafinesque 1820) [Fig. 520]

Branch of Ohio River near Cincinnati to lower part of river; lower Wabash River; lower parts of East Tennessee head streams of Tennessee River to Marshall County, Kentucky; Black and Spring rivers, Arkansas (Goodrich, 1940d).

Genus *Pleurocera* Rafinesque 1818[37]

Pleurocera acuta group

Pleurocera acuta acuta Rafinesque 1831 [Fig. 521]

Ohio River head streams and tributaries; Great Lakes and tributaries; Mississippi River and westward to Nebraska and Kansas; through the Erie Canal into the basin of the Hudson River; Cumberland and Duck rivers, Tennessee (Goodrich, 1940d).

Pleurocera acuta hinkleyi Goodrich 1921

Little Muddy River, Dubois, Washington County, Illinois (Goodrich, 1939e).

Pleurocera acuta lewisi (Lea 1862) [Fig. 522]

Illinois and Kankakee rivers, Illinois (Goodrich, 1939e, 1940d).

Pleurocera alveare group

Pleurocera alveare (Conrad 1834) [Figs. 523, 529]

Lower parts of Ohio, Wabash and Green rivers, together with a few tributaries; Cumberland River, above Burnside, Pulaski County, Kentucky, to tributaries of the river in Trigg County, Kentucky; Tennessee River, Muscle Shoals, and near-by creeks, Alabama; streams of northern Arkansas and southern Missouri (Goodrich, 1940d).

Pleurocera canaliculatum group

Pleurocera canaliculatum canaliculatum (Say 1821)

Ohio River from vicinity of Pittsburgh, Pennsylvania, to Illinois; Wabash River and its tributaries; aberrantly in the Tennessee River system; Omaha, Nebraska (Goodrich, 1940d).

Pleurocera canaliculatum alabamense (Lea 1862) [Fig. 524]
Tributaries of the Tennessee River in northern Alabama (Goodrich, 1940d).

Pleurocera canaliculatum excuratum (Conrad 1834) [Fig. 530]
Tennessee River at Muscle Shoals, Alabama, and lower parts of a few near-by tributaries; Cumberland River, Nashville, Tennessee, to parts of the river in Kentucky; aberrant in Clinch and Wabash rivers (Goodrich, 1940d).

Pleurocera canaliculatum filum (Lea 1845) [Figs. 525, 531]
Upper Cumberland River to a point above Nashville, Davidson County, Tennessee; Duck River, Coffee County, to near the mouth, Tennessee; aberrant in Tennessee River (Goodrich, 1940d).

Pleurocera canaliculatum moriforme (Lea 1862) [Fig. 532]
Muscle Shoals, Tennessee River, Alabama (Goodrich, 1940d).

Pleurocera canaliculatum undulatum (Say 1829) [Figs. 526, 527, 533, 534]
Kentucky River, Kentucky (typical form); Ohio River and tributaries and Cumberland and Tennessee rivers and branches (carinate or angled forms); Rock River, Illinois (Goodrich, 1940d).

Pleurocera gradatum (Anthony 1854) [Fig. 538]
Holston River, Washington County, southwestern Virginia (Tryon, 1873b).

Pleurocera nobile nobile (Lea 1845) [Fig. 550]
Tennessee River, Jackson County, to Marion County, Alabama; Sequatchie River, Tennessee, near mouth; Flint Creek, Morgan County, Alabama (Goodrich, 1940d).

Pleurocera nobile nodosa (Lea 1861) [Fig. 535]
Tennessee River above Chattanooga, Hamilton County, Tennessee (Goodrich, 1940d).

Pleurocera parvum (Lea 1862) [Figs. 536, 537]
Tributaries of the Tennessee River, East Tennessee; apparently extending into South Carolina (Goodrich, 1940d).

Pleurocera postelli (Lea 1862) [Fig. 539]
Small streams of northern Alabama in the vicinity of Muscle Shoals (Goodrich, 1940d).

Pleurocera pyrenellum group

Pleurocera brumbyi (Lea 1852) [Fig. 551]
Springs and streams of the Tennessee River in Madison, Limestone and Courtland counties, Alabama (Goodrich, 1940d).

Pleurocera currierianum (Lea 1863)[38] [Fig. 552]
Alabama: Florence, Lauderdale County; Swan Lake, near Decatur, Limestone County; discharge of a spring in Madison County (Goodrich, 1940d).

Pleurocera pyrenellum (Conrad 1834) [Figs. 540, 541, 553]
Tributaries of the Tennessee River in Morgan and Limestone counties, Alabama, and Walker County, Georgia (Goodrich, 1940d).

Pleurocera trochiformis (Conrad 1834) [Fig. 554]
Tennessee River, Bridgeport, Jackson County, to Florence, Lauderdale County, Alabama; tributaries in Walker County, Georgia, to those near Muscle Shoals, Alabama (Goodrich, 1940d).

Pleurocera viridulum (Anthony 1854)[39] [Fig. 556]
Chickamauga Creek, Walker County, Georgia (Goodrich, 1940d).

Pleurocera unciale group

Pleurocera unciale unciale (Reeve 1861) [Figs. 542, 543, 555]
Upper tributaries of the Tennessee River in Virginia and eastern Tennessee (Goodrich, 1940d).
Pleurocera unciale hastatum (Anthony 1854) [Fig. 557]
North and South Fork of the Holston River, Sullivan County, Tennessee (Goodrich, 1940d).

Pleurocera prasinatum group

Pleurocera annuliferum (Conrad 1834) [Figs. 544, 558]
Upper and middle parts of the Black Warrior River; also known from Village Creek, Jefferson County, Alabama (Goodrich, 1941b).
Pleurocera foremani (Lea 1843) [Fig. 545]
Cahaba River and Coosa River basin from the Etowah River of Georgia downstream, and at the mouths of a few side streams (Goodrich, 1944d).
Pleurocera prasinatum (Conrad 1834) [Figs. 546, 547]
In quiet stretches in the middle and lower Cahaba and Coosa rivers and in the Alabama River (Goodrich, 1941b,c, 1944d).
Pleurocera showalteri (Lea 1862) [Fig. 548]
Lower part of the main Coosa River headwaters and that part of the river which is in Georgia (Goodrich, 1944d).
Pleurocera vestitum (Conrad 1834) [Figs. 549, 559]
Headwaters, creeks and springs from northern Georgia and Alabama to small streams as far south as the first county above Mobile (Goodrich, 1941b).

Subgenus *Strephobasis* Lea 1861

Pleurocera (Strephobasis) corpulentum (Anthony 1854) [Fig. 560]
Tennessee River between Bridgeport and Florence, Alabama; Battle Creek at Ketchall, Marion County, Tennessee (Goodrich, 1928a).
Pleurocera (Strephobasis) curtum curtum (Haldeman 1841) [Fig. 561]
Holston and Tennessee rivers from McMillan, Knox County, Tennessee, to the Muscle Shoals area in Alabama, and probably below it; Cumberland River in the vicinity of Nashville, Tennessee, and Caney Fork near Carthage, Tennessee; Clinch, Little and Little Tennessee rivers a few miles above their mouths; Paint Rock and Flint rivers, Alabama (Goodrich, 1928a).
Pleurocera (Strephobasis) curtum roanense (Lea 1864) [Fig. 562]
Emory River, Roane and Morgan counties, and the Little River, Blount County, Tennessee (Goodrich, 1928a).
Pleurocera (Strephobasis) walkeri Goodrich 1928 [Fig. 563]
Sequatchie and Little Sequatchie rivers, Marion County, Tennessee; Cumber-

land River, Jackson County, the Tennessee River at Muscle Shoals and Shoals Creek, Lauderdale County, Alabama (Goodrich, 1928a).

Family ACROLOXIDAE

Genus *Acroloxus* Beck 1837

Acroloxus coloradensis (Henderson 1930) [Fig. 564]
Isolated lakes high in the Rocky Mountains in Colorado, Montana and Alberta, and a few pond and lake localities in northern Quebec and eastern Ontario (Clarke, 1973).

Family LYMNAEIDAE

Subfamily Lymnaeinae[40]

Genus *Acella* Haldeman 1841

Acella haldemani ('Deshayes' W.G. Binney 1867) [Fig. 565]
Vermont and eastern Ontario west to northern Minnesota, south to northern Illinois and Ohio (F.C. Baker, 1928c).

Genus *Bulimnea* Haldeman 1841

Bulimnea megasoma (Say 1824) [Fig. 566]
Great Lakes and St. Lawrence River drainage area, upper tributaries of the Mississippi drainage area, parts of the Albany, Winnipeg and Nelson river systems in the Canadian Interior Basin (see Clarke, 1973).

Genus *Fossaria* Westerlund 1885[41]

Subgenus *Fossaria* s.s.

Fossaria cyclostoma (Walker 1908) [Fig. 567]
New York to Michigan; a species of the Canadian region and of the Transition life zone (F.C. Baker, 1911a); Great Lakes region; Ontario (F.C. Baker, 1928c).

Fossaria galbana (Say 1825) [Fig. 568]
Great Lakes-St. Lawrence River basin northward in the region west of James Bay to the Attawapiskat and Severn river systems, and northwestward in the boreal forest region to the vicinity of Great Slave Lake (Clarke, 1973, as given for *F. decampi*, here considered a synonym of *F. galbana*).

Fossaria humilis (Say 1822) [Fig. 569]
Atlantic drainage area from southern New Jersey south to South Carolina (see F.C. Baker, 1911a).

Fossaria obrussa (Say 1825) group [Figs. 570, 573-577]

North America generally, except for the Atlantic drainage from southern Virginia south.

exigua Lea 1841 [Fig. 573]

Throughout the St. Lawrence River system, south to Alabama in the Mississippi-Missouri river basin, north to the Hudson Bay lowlands in northern Ontario, and west to the Red River and Lake Winnipeg region in Minnesota and Manitoba (Clarke, 1973).

modicella Say 1825 [Fig. 574]

Eastern Quebec, Nova Scotia and New Jersey west to Vancouver Island, Manitoba south to southern California, Arizona, Texas and Alabama (F.C. Baker, 1928c); also Saskatchewan, Alberta and Northwest Territories (Clarke, 1973).

obrussa Say 1825 [Figs. 570, 575]

From the Atlantic to the Pacific oceans, and from Mackenzie Territory, Canada, south to Arizona and northern Mexico (F.C. Baker, 1928c).

peninsulae Walker 1908 [Fig. 576]

Northern Maine west to Wisconsin; in Wisconsin and Michigan found in streams flowing into Lake Superior (F.C. Baker, 1928c).

rustica Lea 1841 [Fig. 577]

New York west to Utah, Nebraska south to New Mexico (F.C. Baker, 1928c); Ontario, Manitoba, Saskatchewan, Alberta and Northwest Territories (Clarke, 1973).

Fossaria parva (Lea 1841) [Fig. 571]

Connecticut west to Idaho, James Bay and Montana south to Maryland, Kentucky, Oklahoma, southern New Mexico and Arizona (F.C. Baker, 1928c); in Canada, from eastern James Bay drainage to Alberta and north to the region of Great Slave Lake (Clarke, 1973).

Fossaria tazewelliana (Wolf 1869) [Fig. 572]

Northeastern Illinois (Pleistocene); Iowa.

Fossaria truncatula (Müller 1774) [Fig. 583]

Europe, northern Asia and portions of Alaska and Yukon Territory (F.C. Baker, 1911a).

Subgenus *Bakerilymnaea* Weyrauch 1964

Fossaria (Bakerilymnaea) bulimoides (Lea 1841) group [Figs. 584-586]

United States west of the vicinity of the Mississippi River; also southern Saskatchewan and Alberta (chiefly in the prairie and Rocky foothill regions) and southern British Columbia (Clarke, 1973).

alberta F.C. Baker 1919

Brazean Lake, Alberta, Canada (F.C. Baker, 1919e).

bulimoides Lea 1841 [Fig. 584]

Pacific Coast, from Vancouver Island south to southern California (F.C. Baker, 1911a).

cockerelli Pilsbry & Ferriss 1906 [Fig. 585]

Sporadic over most of the United States west of the Mississippi River (Hibbard & Taylor, 1960).

hendersoni F.C. Baker 1909

West of Fort Collins, Larimer County, Colorado (F.C. Baker, 1909a).

perplexa F.C. Baker & Henderson 1929

Park Lake, Grand Coulee, Washington (F.C. Baker & Henderson, 1929).

techella Haldeman 1867 [Fig. 586]

Southwestern and south central United States to central Mexico; from southern California through Utah, Colorado, southernmost Nebraska and Kansas to Missouri and Alabama (Hibbard & Taylor, 1960).

vancouverensis F.C. Baker 1939

Southern part of Vancouver Island, British Columbia, Canada (F.C. Baker, 1939a).

Fossaria (Bakerilymnaea) cubensis (Pfeiffer 1839) [Fig. 587]

Southern United States, from Florida to southern Texas (see F.C. Baker, 1911a).

Fossaria (Bakerilymnaea) dalli (F.C. Baker 1907) [Fig. 588]

Ohio to northern Michigan and Montana, south to Kansas and Arizona (F.C. Baker, 1928c); in the Canadian Interior Basin from southern Manitoba to Alberta (Clarke, 1973).

Fossaria (Bakerilymnaea) perpolita (Dall 1905)

Nushagak, Bristol Bay, Alaska (Dall, 1905).

Fossaria (Bakerilymnaea) sonomaensis Hemphill (in Pilsbry & Ferriss) 1906 [Fig. 589]

Sonoma County, California (Pilsbry & Ferriss, 1906).

Genus *Lymnaea* Lamarck 1799[40]

Lymnaea atkaensis Dall 1884 [Fig. 590]

Throughout most of Alaska and the Yukon Territory, in northern British Columbia in the Liard and Yukon river systems, and along the Arctic Coast to Cape Perry, Northwest Territory (Clarke, 1973).

Lymnaea stagnalis appressa Say 1821 [Fig. 591]

Great Lakes-St. Lawrence River drainage area, northwest to the Mackenzie and Yukon river drainage areas, west to the Rocky Mountains, south in the Rocky Mountains to Colorado, and in Illinois and Ohio in the Mississippi drainage (Clarke, 1973).

Lymnaea stagnalis sanctaemariae Walker 1892 [Fig. 592]

Lake Superior drainage area and adjacent parts of the Lake Huron, Wisconsin River and Winnipeg River drainage areas (Clarke, 1973).

Genus *Pseudosuccinea* F.C. Baker 1908

Pseudosuccinea columella (Say 1817) [Fig. 593]

Eastern North America generally. Nova Scotia and Quebec west to Manitoba, Minnesota and eastern Kansas, south to central Texas and Florida (F.C. Baker, 1911a).

Genus *Radix* Montfort 1810

Radix auricularia (Linnaeus 1758) [Fig. 594]
Europe and northern Asia; widely introduced but of spotty occurrence in North America.

Genus *Stagnicola* Leach (in Jeffreys) 1830[42,43]

Subgenus *Stagnicola* s.s.

Stagnicola elodes group

Stagnicola elodes (Say 1821)[42] [Figs. 600-606, 611]
New England west to Oregon and California, south to New Mexico; widely distributed in the Canadian Interior Basin (see Clarke, 1973).
Stagnicola elrodiana F.C. Baker 1935 [Fig. 595]
Western Montana (Lakes Sin-yale-a-min and McDonald) (Elrod, 1902).
Stagnicola exilis (Lea 1834) [Figs. 596, 597]
Ohio to Kansas, northward to northern Minnesota and Michigan (F.C. Baker, 1928c).
Stagnicola neopalustris (F.C. Baker 1911) [Fig. 598]
Orange County, Virginia (F.C. Baker, 1911a).
Stagnicola traski (Tryon 1863) [Fig. 599]
California to Wyoming, north to southern Alberta (F.C. Baker, 1911a).

Stagnicola emarginata/catascopium group

Stagnicola apicina (Lea 1838)
Northern part of the lower peninsula of Michigan west to western Washington; Ontario south to southern Wyoming and South Dakota (F.C. Baker, 1911a).
Stagnicola arctica (Lea 1864) [Fig. 607]
Newfoundland to the vicinity of Hudson Bay, north and northwest in subarctic and arctic Canada to Ungava, southern Victoria Island, the Mackenzie River Delta and the vicinity of Point Barrow, Alaska (Clarke, 1973).
Stagnicola bonnevillensis (Call 1884) [Fig. 608]
Wyoming (D.W. Taylor!).
Stagnicola catascopium (Say 1817) [Figs. 609, 610]
Eastern Canada and Nova Scotia west to North Dakota, Great Slave Lake south to northern Iowa, northern Ohio and Maryland (F.C. Baker, 1928c).
Stagnicola contracta (Currier (in DeCamp) 1881) [Fig. 612]
Higgins Lake, Roscommon County, Michigan (DeCamp, 1881).
Stagnicola elrodi (F.C. Baker & Henderson 1933) [Fig. 613]
Flathead Lake, Montana (F.C. Baker & Henderson, 1933).
Stagnicola emarginata (Say 1821) [Figs. 614-617]
Maine west to Minnesota and Wisconsin, Canadian Interior Basin south to Michigan, Pennsylvania and New York (F.C. Baker, 1928c).

Stagnicola gabbi (Tryon 1865) [Fig. 618]
California (see F.C. Baker, 1911a).
Stagnicola hinkleyi (F.C. Baker 1906) [Fig. 620]
Columbia River drainage, Idaho and Oregon (F.C. Baker, 1911a).
Stagnicola idahoense (Henderson 1931) [Fig. 619]
Little Salmon River, Idaho (Henderson, 1931a).
Stagnicola mighelsi (Binney 1865) [Fig. 621]
Lakes in Maine (see F.C. Baker, 1911a).
Stagnicola oronoensis (F.C. Baker 1904) [Fig. 622]
Maine to eastern Ontario (F.C. Baker, 1911a).
Stagnicola petoskeyensis (Walker 1908) [Fig. 623]
Small spring-brook flowing into Little Traverse Bay, near Petoskey, Michigan (Walker, 1908e).
Stagnicola utahensis (Call 1884) [Fig. 624]
Lake Utah, Utah (Call, 1884).
Stagnicola walkeriana F.C. Baker 1926 [Fig. 625]
Great Lakes (Michigan and Superior) (F.C. Baker, 1928c).
Stagnicola woodruffi (F.C. Baker 1901) [Figs. 626, 627]
Great Lakes (Huron, Michigan, Ontario); Lake Geneva, Wisconsin; Rainy River system; Lake of the Woods (Clarke, 1973).

Subgenus *Hinkleyia* F.C. Baker 1928

Stagnicola (Hinkleyia) caperata (Say 1829) [Figs. 628, 629]
Quebec and Massachusetts west to California; Yukon Territory and James Bay south to Maryland, Indiana, Colorado and California (F.C. Baker, 1928c).
Stagnicola (Hinkleyia) montanensis (F.C. Baker 1913) [Fig. 630]
Hays Creek near Ward, Montana; upper Snake River drainage in Idaho and Wyoming; Beaver, Cache and Summit counties, Utah; Nye County, Nevada (Taylor, Walter & Burch, 1963).
Stagnicola (Hinkleyia) pilsbryi (Hemphill 1890) [Fig. 631]
Fish Springs, Juab County, Utah (Russell, 1971b).

Subfamily Lancinae[44]

Genus *Fisherola* Hannibal 1912

Fisherola nuttalli nuttalli (Haldeman 1841)[45]
Columbia River drainage (Pilsbry, 1925a); Snake River drainage, Idaho, and Deschutes River and The Dalles, Oregon (Henderson, 1936c).
Fisherola nuttalli kootaniensis (Baird 1863)
Spokane River [eastern Washington] and Kootenai River, British Columbia (Baird, 1863).
Fisherola nuttalli lancides Hannibal 1912 [Fig. 632]
Snake River basin (Hannibal, 1912b), = ? Spokane River (Henderson, 1936c).

Genus *Lanx* Clessin 1882

Subgenus *Lanx* s.s.

Lanx alta (Tryon 1865) [Fig. 578]
Klamath River, California (Tryon, 1865j).
Lanx patelloides (Lea 1856) [Figs. 580, 633]
Sacramento River, California (Lea, 1856).
Lanx subrotundata (Tryon 1865) [Fig. 579]
Umpqua River, Oregon (Tryon, 1865j); Umpqua River system, Oregon (Henderson, 1929c, 1936c).

Subgenus *Walkerola* Hannibal 1912

Lanx (Walkerola) klamathensis Hannibal 1912 [Fig. 634]
Klamath system in basin of Klamath River, Oregon (Hannibal, 1912b).

Family PHYSIDAE[46]

Subfamily Physinae

Genus *Physa* Draparnaud 1801

Physa jennessi Dall 1919 [Fig. 635]
Alaska, Northwest Territories and British Columbia.
Physa skinneri Taylor 1954 [Fig. 636]
Canada from Quebec to Northwest Territories and British Columbia; south to Washington, Montana, Wyoming, Nebraska, Iowa, Ohio, Pennsylvania and New England.
Physa skinneri, large unnamed morph [Fig. 637]
Canada from Ontario west to Saskatchewan; Massachusetts and Pennsylvania west to Michigan.

Genus *Physella* Haldeman 1843

Physella ancillaria (Say 1825) [Fig. 666]
New Brunswick to Ontario, Canada, and New York and Pennsylvania east into New England.
Physella boucardi (Crosse & Fischer 1881) [Figs. 581, 638]
Nevada and California south into Mexico.
Physella columbiana (Hemphill 1890) [Fig. 639]
Wyoming and Montana west to Washington.
Physella cooperi (Tryon 1865) [Fig. 640]
Wyoming west to California and east to Colorado.
Physella globosa (Haldeman 1841) [Fig. 667]
Kentucky, Ohio and Tennessee.

Physella gyrina gyrina (Say 1821) [Fig. 647]
In Canada, Quebec to Ontario; south to Nebraska and east to New York.
Physella gyrina gyrina morph *elliptica* (Lea 1834) [Fig. 648]
Ontario south to Iowa and Missouri and east to New York.
Physella gyrina gyrina morph *hildrethiana* (Lea 1841) [Fig. 649]
Ontario south to Iowa and Missouri and east to New York.
Physella gyrina alba (Crandall 1901) [Fig. 650]
Eastern Canada to Ontario and northeastern United States.
Physella gyrina ampullacea Gould 1855 [Fig. 651]
In Canada from Manitoba west to British Columbia; south to California, east to Arizona and north to Minnesota.
Physella gyrina athearni (Clarke 1973) [Fig. 652]
Alberta, Canada.
Physella gyrina aurea (Lea 1838) [Fig. 653]
New Jersey to Kansas, south to Arkansas and Florida.
Physella gyrina aurea morph *albofilata* ('Ancey' Sampson 1893) [Fig. 654]
Pennsylvania west to Kansas, south to Oklahoma and Alabama.
Physella gyrina bayfieldensis (Baker 1928) [Fig. 655]
Northwest Territories of Canada south to Kansas.
Physella gyrina cylindrica (Newcomb 1843) [Fig. 656]
Ontario and New York south to Virginia.
Physella gyrina gouldi (Clench 1935) [Fig. 657]
Northwest Territories south to Wisconsin and Colorado.
Physella gyrina hawni (Lea 1864) [Fig. 658]
Ohio west to Kansas and south to Texas and Alabama.
Physella gyrina microstoma (Haldeman 1840) [Fig. 659]
Virginia to Missouri and south to Arkansas and Alabama.
Physella gyrina sayi (Tappan 1838) [Fig. 660]
Quebec to Northwest Territories, south to Saskatchewan, the Dakotas and New York.
Physella gyrina smithiana (Baker 1920) [Fig. 661]
Kansas to Texas, Wyoming and California.
Physella hordacea (Lea 1864) [Fig. 641]
British Columbia, Washington and Oregon.
Physella lordi (Baird 1863) [Fig. 642]
British Columbia south to Montana, Nevada and California.
Physella magnalacustris (Walker 1901) [Fig. 668]
Ontario south to the Great Lakes states and Indiana, east to Vermont and Maine.
Physella microstriata (Chamberlain & E.G. Befry 1930) [Fig. 643]
Utah.
Physella parkeri parkeri (Currier (in DeCamp) 1881) [Fig. 669]
Michigan and Wisconsin.
Physella parkeri latchfordi (Baker 1928) [Fig. 670]
Quebec, Ontario, Wisconsin, Michigan and Maine.
Physella propinqua propinqua (Tryon 1865) [Fig. 662]
Montana and Washington south to Wyoming and California.

Physella propinqua nuttalli (Lea 1864) [Fig. 663]
British Columbia south to Montana, Wyoming and California.
Physella propinqua nuttalli morph *triticea* (Lea 1856) [Fig. 664]
Idaho and Washington, south to California and Nevada.
Physella propinqua nuttalli morph *venusta* (Lea 1864) [Fig. 665]
Montana to Washington, south to California and northeast to Utah and Wyoming.
Physella traski (Lea 1864) [Fig. 644]
Oregon and California.
Physella utahensis (Clench 1925) [Fig. 645]
Wyoming, Colorado and Utah.
Physella vinosa (Gould 1847) [Fig. 671]
Ontario, Canada, and the Great Lakes states.
Physella virginea (Gould 1847) [Fig. 646]
British Columbia south to California.

Subgenus *Costatella* Dall 1870

Physella (Costatella) acuta Draparnaud 1805) [Fig. 678]
Europe, Mediterranean regions, and Africa; introduced into Australia, Hawaii and perhaps parts of continental United States.
Physella (Costatella) bottimeri (Clench 1924) [Fig. 679]
New Mexico, Oklahoma and Texas.
Physella (Costatella) conoidea (Fischer & Crosse 1886) [Fig. 582]
Texas.
Physella (Costatella) costata (Newcomb 1861) [Fig. 674]
California.
Physella (Costatella) cubensis cubensis (Pfeiffer 1839) [Fig. 672]
Bahamas, Cuba, Jamaica, Puerto Rico, West Indies, Honduras and Florida.
Physella (Costatella) cubensis peninsulae (Pilsbry 1899) [Fig. 673]
Florida.
Physella (Costatella) hendersoni hendersoni (Clench 1925) [Fig. 675]
West Virginia, Tennessee and Missouri, south to the Carolinas, Mississippi, and Florida.
Physella (Costatella) hendersoni hendersoni morph *ariomus* (Clench 1925) [Fig. 676]
Virginia, South Carolina, Georgia, Florida and Alabama.
Physella (Costatella) hendersoni ssp. [Fig. 677]
Virginia, North Carolina, Florida and Alabama.
Physella (Costatella) heterostropha heterostropha (Say 1817) [Fig. 680]
Nova Scotia to Ontario; New England to Ohio, Tennessee and the Virginias; the Bahamas.
Physella (Costatella) heterostropha halei (Lea 1864) [Fig. 682]
Illinois, Missouri, Kansas, Arkansas and Texas.
Physella (Costatella) heterostropha pomila (Conrad 1834) [Fig. 681]
Eastern United States to the Ohio and Mississippi rivers and in Iowa.
Physella (Costatella) humerosa (Gould 1855) [Fig. 683]
California, Arizona and Colorado.

Physella (Costatella) integra integra (Haldeman 1841) [Fig. 695]
Quebec to Manitoba, Canada, and the Great Lakes states, Iowa, South Dakota, Tennessee, Kentucky and West Virginia.
Physella (Costatella) integra integra morph *walkeri* (Crandall 1901) [Fig. 696]
Quebec, Ontario and the Great Lakes states.
Physella (Costatella) integra brevispira (Lea 1864) [Fig. 697]
New York, Ohio, Wisconsin and Minnesota.
Physella (Costatella) johnsoni (Clench 1926) [Fig. 684]
Alberta, Canada, and Montana, Wyoming and Colorado.
Physella (Costatella) osculans (Haldeman 1841) [Fig. 685]
Colorado west to California and southeast to Arizona and into Mexico.
Physella (Costatella) spelunca (Turner & Clench 1974) [Fig. 686]
Wyoming.
Physella (Costatella) squalida (Morelet 1851) [Fig. 687]
Texas into Mexico, Central and South America, and in Costa Rica.
Physella (Costatella) virgata virgata (Gould 1855) [Fig. 688]
Nebraska west to California, east to Texas and into Mexico.
Physella (Costatella) virgata virgata morph *parva* (Lea 1864) [Fig. 689]
Iowa west to California, east to Texas and north to Kansas.
Physella (Costatella) virgata anatina (Lea 1864) [Fig. 690]
Wisconsin and South Dakota southwest to Colorado and Nevada; Texas and Arkansas north to Illinois and Nebraska.
Physella (Costatella) virgata berendti (Fischer & Crosse 1886) [Fig. 691]
Wyoming to California, southeast to Texas and Mexico and north to Kansas.
Physella (Costatella) virgata concolor (Haldeman 1841) [Fig. 692]
Manitoba and Wisconsin to Idaho.
Physella (Costatella) virgata concolor morph [Fig. 693]
Wyoming.
Physella (Costatella) virgata rhyssa (Pilsbry 1899) [Fig. 694]
California, New Mexico and Texas into Mexico.

Subgenus *Petrophysa* Pilsbry 1926

Physella (Petrophysa) zionis (Pilsbry 1926) [Fig. 698]
Utah.

Subfamily Aplexinae

Genus *Aplexa* Fleming 1820

Aplexa elongata (Say 1821) [Fig. 699]
Ontario to Saskatchewan, Canada, and Alaska; New England through the Great Lakes states to Washington; south to Idaho, Utah and Wyoming.
Aplexa elongata morph *tryoni* (Currier 1867) [Fig. 700]
Ontario and Alberta, Canada; Michigan and Minnesota west to Washington; also in Utah, Illinois and Indiana.

Genus *Stenophysa* Martens 1898[47]

Stenophysa marmorata (Guilding 1828) [Fig. 701]
Brazil, Guatemala, Uruguay, Venezuela and the West Indies. Introduced into Texas (*teste* Te, 1978).
Stenophysa maugeriae (Gray 1837) [Fig. 702]
Mexico. Introduced into Texas (*teste* Te, 1978).

Family PLANORBIDAE

Subfamily Planorbinae

Tribe Planorbini

Genus *Gyraulus* 'Agassiz' Charpentier 1837

Subgenus *Gyraulus* s.s.

Gyraulus deflectus (Say 1824) [Fig. 705]
Along the Atlantic Coast from Prince Edward Island south to Virginia, west to Ohio, Illinois, Alberta and Idaho (Miller, 1966); north to near the Arctic Coast in the Ungava, Coppermine River and Mackenzie River districts (Clarke, 1973).

Subgenus *Armiger* Hartmann 1840

Gyraulus (Armiger) crista (Linnaeus 1758) [Fig. 706]
Holarctic. In North America from Ontario and Maine to Minnesota, northwestern Northwest Territories and Alaska (Clarke, 1973).

Subgenus *Torquis* Dall 1905

Gyraulus (Torquis) circumstriatus (Tryon 1866) [Fig. 707]
Connecticut north to Quebec, west to Alberta and south in the Rocky Mountains to New Mexico (Clarke, 1973).
Gyraulus (Torquis) hornensis F.C. Baker 1934[48] [Fig. 708]
Mackenzie River region west of Great Slave Lake; western Ontario, Wisconsin and North Dakota (F.C. Baker, 1934).
Gyraulus (Torquis) parvus (Say 1817) [Fig. 709]
North America, from Alaska and northern Canada to Cuba and from the Atlantic to the Pacific Coast (Taylor, 1960).

Tribe Drepanotremini[49]

Genus *Drepanotrema* Fischer & Crosse 1880

Subgenus *Antillorbis* Harry & Hubendick 1964

Drepanotrema (Antillorbis) aeruginosum (Morelet 1851) [Fig. 710]
Southern Texas and southern Arizona, Mexico, Guatemala and Antilles (Bequaert & Miller, 1973).

Subgenus *Fossulorbis* Pilsbry 1934

Drepanotrema (Fossulorbis) cimex (Moricand 1839) [Fig. 715]
Southern Texas, Mexico, Central America, Venezuela, Brazil and the Greater Antilles (Harry & Hubendick, 1964).
Drepanotrema (Fossulorbis) kermatoides (d'Orbigny 1835) [Fig. 711]
Florida, Texas, Mexico, Central America, Venezuela, Peru, Brazil and the Lesser Antilles (Harry & Hubendick, 1964).

Tribe Biomphalariini

Genus *Biomphalaria* Preston 1910

Biomphalaria glabrata (Say 1818) [Fig. 712]
West Indies, Venezuela, Surinam, French Guiana and Brazil (Barbosa et al., 1968); introduced to Florida.
Biomphalaria havanensis (Pfeiffer 1839) [Fig. 713]
Florida, Louisiana and Texas (Malek, 1969); Arizona, Mexico, Central America (Bequaert & Miller, 1973); also Puerto Rico and Cuba.

Tribe Helisomini

Genus *Helisoma* Swainson 1840

Subgenus *Helisoma* s.s.

Helisoma anceps anceps (Menke 1830)[50] [Fig. 714]
Throughout North America from James and Hudson bays south to Georgia, Alabama, Texas and northwestern Mexico, west to southwestern Northwest Territories, Alberta and Oregon (see Walker, 1909e; Clarke, 1973).
Helisoma anceps royalense (Walker 1909)
Isle Royale, Lake Superior, and the adjacent portion of Ontario north and west of Lake Superior in parts of the Albany, Attawapiskat and Winnipeg river systems (Clarke, 1973).
Helisoma eucosmium (Bartsch 1908)[50]
Greenfield Pond near Wilmington, North Carolina; Burks Place, Louisiana (form or subspecies *vaughani*) (Bartsch, 1908).

Subgenus *Carinifex* W.G. Binney 1865[51]

Helisoma (Carinifex) newberryi newberryi (Lea 1858) [Figs. 720, 721]
Idaho, Utah, Nevada, Oregon and California.

Helisoma (Carinifex) newberryi jacksonense Henderson 1932 [Figs. 716, 717]
Jackson Lake, Wyoming (Henderson, 1932a).
Helisoma (Carinifex) newberryi occidentale Hanna 1924 [Figs. 718, 719]
Eagle Lake, Lassen County, California (Hanna, 1924).

Genus *Menetus* H. & A. Adams 1855

Subgenus *Menetus* s.s.[52]

Menetus opercularis (Gould 1847) [Figs. 722, 723]
Alaska south to Alberta and southern California.

Subgenus *Micromenetus* F.C. Baker 1945[53, 54, 55]

Menetus (Micromenetus) brogniartianus (Lea 1842) [Fig. 725]
Near Cincinnati, Ohio (Lea, 1842b); Woodville, Alabama (Pilsbry, 1895b; for *alabamensis*).
Menetus (Micromenetus) dilatatus (Gould 1841) [Figs. 724, 726]
Eastern United States, from Maine west to Iowa, south to Texas and Florida.
Menetus (Micromenetus) sampsoni 'Ancey' Sampson 1885 [Fig. 727]
Illinois, Missouri and Arkansas (Sampson, 1913).

Genus *Planorbella* Haldeman 1842

Subgenus *Planorbella* s.s.[56]

Planorbella campanulata campanulata (Say 1821) [Fig. 728]
Vermont west to North Dakota, south to Ohio and Illinois, northward to Great Slave Lake (F.C. Baker, 1928c).
Planorbella campanulata collinsi (F.C. Baker 1939)
Northwestern Ontario in the headwaters of the Albany, Winnipeg and Severn river systems (Clarke, 1973).
Planorbella multivolvis (Case 1847) [Fig. 729]
Howe Lake, Marquette County, Michigan (Walker, 1907d).

Subgenus *Pierosoma* Dall 1905

Planorbella (Pierosoma) ammon (Gould 1855) [Fig. 730]
Cienaga Grande, or Colorado Low Desert (Gould, 1855a; Henderson, 1936d); Sacramento and San Joaquin river drainages and near Watsonville, California (Henderson, 1934a).
Planorbella (Pierosoma) binneyi (Tryon 1867)
California to British Columbia in the Pacific drainage area and British Columbia and Alberta in the headwaters of the Peace and North Saskatchewan river systems (Clarke, 1973).

Planorbella (Pierosoma) columbiensis (F.C. Baker 1945)
Lac La Hache, Cariboo District, British Columbia (F.C. Baker, 1945).

Planorbella (Pierosoma) corpulenta corpulenta (Say 1824)
Western Ontario, eastern Manitoba and northern Minnesota in the Winnipeg River system; upper Mississippi River system in northern Minnesota (Clarke, 1973).

Planorbella (Pierosoma) corpulenta vermilionensis (F.C. Baker 1929)
Vermilion Lake, St. Louis County, Minnesota (F.C. Baker, 1929b).

Planorbella (Pierosoma) corpulenta whiteavesi (F.C. Baker 1932)
Greenwater Lake and Lac des Mille Lacs, Thunder Bay District, Ontario (Clarke, 1973).

Planorbella (Pierosoma) magnifica (Pilsbry 1903) [Fig. 732]
Greenfield Pond, near Wilmington, North Carolina (Bartsch, 1908).

Planorbella (Pierosoma) occidentalis (Cooper 1870) [Fig. 733]
Lakes, rivers, creeks, ditches, sloughs and swamps in California, Oregon and Washington (see Henderson, 1936c).

Planorbella (Pierosoma) oregonensis (Tryon 1865)
Pueblo Valley, Oregon (Tryon, 1865j); Tooele County, Utah (F.C. Baker, 1945).

Planorbella (Pierosoma) pilsbryi (F.C. Baker 1926) [Fig. 731]
Massachusetts west to Minnesota, northern New York and central Wisconsin northward (F.C. Baker, 1928c) [form *pilsbryi* s.s.]; St. Lawrence River drainage area in Georgian Bay and the St. Lawrence River and Rideau River; Canadian Interior Basin from eastern Ontario to central Saskatchewan (Clarke, 1973) [form *infracarinata*]; Vilas County, Wisconsin (F.C. Baker, 1928c) [form *winslowi*].

Planorbella (Pierosoma) tenuis (Dunker 1850) [Fig. 735]
Texas, Arizona, New Mexico, southern California and Mexico (Bequaert & Miller, 1973).

Planorbella (Pierosoma) traski (Lea 1856)
California: Kern Lake (Lea, 1856), Stockton (Henderson, 1934a), Bakersfield, Kern County, and Buena Vista Lake (F.C. Baker, 1945).

Planorbella (Pierosoma) trivolvis trivolvis (Say 1817) [Fig. 736]
Northern North America east of the Rocky Mountains, south to Nebraska, northern Illinois, Pennsylvania and New Jersey.

Planorbella (Pierosoma) trivolvis lenta (Say 1834)
Central United States from Kansas and central Illinois to (?) Texas and Louisiana.

Planorbella (Pierosoma) trivolvis subcrenata (Carpenter 1857) [Fig. 734]
California to British Columbia and Yukon Territory and east to Utah, Colorado, Minnesota and Manitoba (Clarke, 1973).

Planorbella (Pierosoma) trivolvis turgida (Jeffreys 1830)
From Long Pine Key, in the southern Everglades, throughout peninsular Florida and north along the coast to Lake Waccamaw, North Carolina (Pilsbry, 1934a) and Delaware and Maryland, west to Alabama, Arkansas and (?) Texas.

Planorbella (Pierosoma) truncata (Miles 1861) [Fig. 737]
Michigan, northern Illinois, and Wisconsin (F.C. Baker, 1928c).

Subgenus *Seminolina* Pilsbry 1934

Planorbella (Seminolina) duryi (Wetherby 1879) [Figs. 738, 739]
Northern to southern Florida (see Pilsbry, 1934a).
Planorbella (Seminolina) scalaris (Jay 1839) [Fig. 740]
Southern Florida (see Pilsbry, 1934a).

Genus *Planorbula* Haldeman 1840

Planorbula armigera armigera (Say 1821) [Figs. 741, 742]
New Brunswick west to southeastern Ontario, west to Saskatchewan, northwest to the Mackenzie River system (Clarke, 1973); south to Georgia and Louisiana and west to Nebraska (F.C. Baker, 1928c).
Planorbula armigera wheatleyi (Lea 1858)[57] [Figs. 743, 744]
Alabama and Florida.
Planorbula campestris (Dawson 1875) [Fig. 745]
Southern Manitoba and North Dakota, south to Utah and New Mexico, west to British Columbia, and north to the Mackenzie River system (Clarke, 1973).

Genus *Promenetus* F.C. Baker 1935[58]

Promenetus exacuous (Say 1821) [Fig. 746]
United States east of the Rocky Mountains, north to Alaska and the Mackenzie River, south to New Mexico (F.C. Baker, 1928); in Canada absent from Quebec, but widely distributed west of James and Hudson bays, mainly south of the tree-line (Clarke, 1973).
Promenetus umbilicatellus (Cockerell 1887)[59] [Fig. 747]
Alaska south to Oregon, northern Utah, Colorado, New Mexico and eastern Oklahoma, east to eastern Ohio, western New York; in Canada, in Alberta, Saskatchewan and Manitoba (Hibbard & Taylor, 1960); also Texas, if *P. carus* (Pilsbry & Ferriss) is a synonym of *P. umbilicatellus*.

Genus *Vorticifex* Meek (in Dall) 1870[60]

Subgenus *Parapholyx* Hanna 1922

Vorticifex (Parapholyx) effusa (Lea 1856) [Fig. 748]
Rivers and lakes in California and Oregon.
Vorticifex (Parapholyx) solida Dall 1870)[61] [Fig. 751]
Lakes in Nevada and California.

Subfamily Neoplanorbinae

Genus *Amphigyra* Pilsbry 1906

Amphigyra alabamensis Pilsbry 1906 [Fig. 749]
Coosa River, Alabama (Pilsbry, 1906b).

Genus *Neoplanorbis* Pilsbry 1906[62]

Neoplanorbis carinatus Walker 1908 [Fig. 752]
Coosa River, Alabama (Walker, 1908c).
Neoplanorbis smithi Walker 1908 [Fig. 753]
Coosa River, Alabama (Walker, 1908c).
Neoplanorbis tantillus Pilsbry 1906 [Fig. 750]
Coosa River, Alabama (Pilsbry, 1906b).
Neoplanorbis umbilicatus Walker 1908 [Fig. 754]
Coosa River, Alabama (Walker, 1908c).

Family ANCYLIDAE

Subfamily Ancylinae

Genus *Rhodacmea* Walker 1917

Rhodacmea elatior (Anthony 1855) [Fig. 756]
Tennessee and Cahaba river systems (Basch, 1963).
Rhodacmea filosa (Conrad 1834) [Figs. 757, 759]
Black Warrior and Coosa rivers, Alabama, and tributaries; ? also in Tennessee River system (Basch, 1963).
Rhodacmea hinkleyi (Walker 1908) [Figs. 758, 760]
Coosa River, Alabama, and the Tennessee River drainage, extending irregularly northward to the southern borders of Illinois and Indiana (Basch, 1963).

Subfamily Ferrissinae

Genus *Ferrissia* Walker 1903

Ferrissia fragilis (Tryon 1863) [Figs. 764, 765]
"Among the most widely distributed of North American freshwater snails" (Basch, 1963); New York to Michigan, California and Texas.
Ferrissia mcneili Walker 1925 [Fig. 766]
Mobile area, Alabama (see Basch, 1963).
Ferrissia parallelus (Haldeman 1841)
In Canada and the northern United States from the Atlantic coast westward (Basch, 1963); Nova Scotia and New England west to Manitoba, Minnesota

and Illinois in the Atlantic, St. Lawrence River, Hudson Bay and upper Mississippi River drainage areas (Clarke, 1973).

Ferrissia rivularis (Say 1817) [Figs. 761, 767]

Throughout most of North America; it extends northward into the Hudson Bay lowlands and northwestward to at least central Saskatchewan; south to North Carolina and New Mexico and west to California and Oregon (see Clarke, 1973).

Ferrissia walkeri (Pilsbry & Ferriss 1907) [Fig. 768]

Arkansas, Michigan and southern California (Basch, 1963).

Subfamily Laevapecinae

Genus *Hebetancylus* Pilsbry 1914

Hebetancylus excentricus (Morelet 1851) [Figs. 762, 769]

Central America; Georgia, Florida and Texas (Basch, 1963).

Genus *Laevapex* Walker 1903

Laevapex diaphanus (Haldeman 1841) [Fig. 770]

Delaware, Illinois, Ohio, Holston and Tennessee rivers (Walker, 1903b); Georgia and Alabama (Walker, 1908d, for *L. hemisphaericus*).

Laevapex fuscus (C.B. Adams 1841) [Figs. 763, 771]

United States and Canada, generally east of the Great Plains; Great Lakes area, Florida and southeastern states; generally absent from mountainous areas (Basch, 1963); west to Iowa, Kansas and Oklahoma (Clarke, 1973).

SECTION IV

KEYS TO THE FRESHWATER GASTROPODS OF NORTH AMERICA

* * *

FAMILIES AND HIGHER TAXA

1 Animal with an operculum (which seals the shell aperture when the snail's body is withdrawn into the shell) (Fig. 1); respiration by gills; mantle opening facing anteriorly. Subclass **Prosobranchia** 2

Animal without an operculum to seal its shell aperture when withdrawn; respiration by the vascularized lining of the mantle cavity (true gills are lacking) or by a pseudobranch (false gill) outside the mantle cavity (Fig. 3a); mantle opening directed to the side (to the right or left, depending on whether the animal is dextral [right coiled] or sinistral [left coiled] (Fig. 3a,b)). Subclass **Pulmonata**, Order **Lymnophila** 11

2(1) Shell globose, subspherical or hemispherical (Fig. 21), solid, with a very low spire; aperture semi-circular or half-moon shaped, with "teeth" or tubercles on the parietal columellar margin of the aperture; operculum calcareous, paucispiral, with a pair of projecting processes on the inner columellar side (Fig. 22); shell usually with a pattern of pale variegations on a greenish-olive background; adult shell of medium size, its height about 20 mm; shell with three to four whorls, the last one making up most of the shell; gill bipectinate or feather-like, i.e., with gill laminae on both sides of the gill axis; radula rhipidoglossate (Fig. 17), with many marginal teeth. Florida and southern Georgia . . . . . Family NERITINIDAE [Order **Neritinacea**, Superfamily **Neritinoidea**] (page 67)

Shell of various shapes and sizes, but if neritiniform (see above, Neritinidae; Fig. 14) the shell is small (no more than 5 mm in height); operculum without a projecting process on the inner side; shell color patterns variable, but not of the variegated kind (see above, Neritinidae); gill monopectinate (except in the Valvatidae), i.e., with gill laminae only on one side of the gill axis (which is adnate along its entire length to the pallial wall); radula taenioglossate (Fig. 17), with few (two) marginal teeth. Order **Mesogastropoda** 3

3(2) Shell small (8 mm or less in diameter), spire generally depressed, some species with carina; operculum multispiral (Fig. 15a); gill bipectinate or feather-like, protruding from the mantle cavity when the snail is active (Fig. 16); pallial tentacle (Fig. 16) present. Superfamily **Valvatoidea** . . . . Family VALVATIDAE (page 67)

Shell small to large, spire depressed to elongate; operculum multispiral (Fig. 15a), paucispiral (Fig. 15b) or concentric (Fig. 15c, d); gill monopectinate; pallial tentacle absent 4

4(3) Operculum multispiral or paucispiral (Fig. 15a, b), the distal margins not concentric 5

Operculum concentric (although the nucleus may be paucispiral) (Fig. 15c, d). Superfamily **Ampullarioidea** (Viviparoidea) 9

5(4) Adult shells usually less than 5 mm in length (but a few species reach this length or exceed it by 1 or 2 mm, and the shell of one hydrobiid species (*Fluminicola nuttalliana* Lea) reaches 10 mm in length); males possess a verge (see Figs. 83, 85-92). Superfamily **Truncatelloidea** (Rissooidea) 6

Adult shells of medium to large size (usually more than 15 mm in length, but some shells are smaller, to 10 mm in length, and in several species the adult shells are no longer than 6-9 mm); males lack a verge. Superfamily **Vermetoidea** (Cerithioidea) 8

6(5) Shell globose-conic, sculptured with numerous spiral epidermal ridges; central radular tooth lacks basal denticles (Fig. 81a). Inhabits streams in caves in Indiana and Kentucky . . . . . Family MICROMELANIIDAE (page 83)

Shell of various shapes, usually smooth, but if sculpturing is present it does not consist of spiral epidermal ridges; central radular tooth with one or more basal denticles or cusps on each side (Fig. 81b, c) 7

7(6) Shell high-spired, turriform; the head-foot region of the body is subdivided on each side by a longitudinal groove; central radular tooth with two or more basal cusps, which are situated on antero-posterior ridges (Fig. 81c); eyes in prominent swellings on the outer bases of the tentacles; amphibious or terrestrial in habit . . . . . Family POMATIOPSIDAE (page 115)

Shell high-spired to depressed; head-foot region not subdivided by a longitudinal groove; central radular tooth with 1-10 basal cusps attached to a thickened ridge along the lateral angle (Fig. 81b), not on antero-posterior ridges; eyes at the outer bases of the tentacles, but not on prominent swellings; totally aquatic in habit . . . . . Family HYDROBIIDAE (page 83)

8(5) Mantle edge smooth; males always present, reproduction dioecious; females lay eggs, having an egg-laying sinus on the right side of the foot . . . . . Family PLEUROCERIDAE[21] (page 118)

Mantle edge papillate; males generally absent (parthenogenetic reproduction common, often the rule); females brood their young in an adventitious ("subhaemocoelic"; not uterine) brood pouch in the postero-dorsal head-foot region. Introduced sporadically in the southernmost United States from Florida to Texas . . . . . Family THIARIDAE (page 117)

9(4) Shells of adults medium to large, more than 20 mm in shell length (in some species reaching more than 50 or 60 mm); operculum corneous 10

Shells of adults smaller, less than 15 mm in length; operculum calcareous. Great Lakes and St. Lawrence regions from Wisconsin to Pennsylvania and New York . . . . . Family BITHYNIIDAE (page 81)

10(9) Shell globose and large (height often up to or exceeding 60 mm), or shell planate (discoidal, with sunken spire), its width exceeding 40 mm; ends of labial palps whip-like; in males the penis arises from the right side of the mantle edge; females lay calcareous (*Pomacea*) or gelatinous (*Marisa*) eggs. Alabama, Florida and Georgia . . . . . Family AMPULLARIIDAE (page 81)

Shell subglobose to turreted, medium to large; ends of labial palps blunt, not whip-like; in males the right tentacle is modified as a penis sheath; females ovoviviparous. Found throughout the United States and Canada . . . . . Family VIVIPARIDAE (page 75)

11(1) Shell coiled 12

Shell an uncoiled, obtuse cone (limpet- or cap-shaped) (Fig. 13) 14

12(11) Animal and shell dextral (coiled to the right) (Figs. 2b, 3b). Superfamily **Lymnaeoidea**, in part . . . . . Family LYMNAEIDAE, in part (page 147)

Animal and shell sinistral (coiled to the left) (Figs. 2a, 3a). Superfamily **Ancyloidea**, in part 13

13(12) Shell with a raised spire; blood (haemolymph) nearly colorless (the respiratory pigment is haemocyanin); animal without pseudobranch (false gill); mantle margin digitate or lobed . . . . . Family PHYSIDAE (page 160)

Shell discoidal, with a sunken spire (Figs. 12, 704) (in some species the smaller (older) shell coils protrude on the umbilical side ("ultrasinistral" or pseudo-dextral shells)); blood (haemolymph) in nearly all species is red (contains haemoglobin); a pseudobranch (false gill) is situated near the pneumostome or anus (Fig. 3a); mantle margin simple . . . . . Family PLANORBIDAE (page 168)

14(11) Adult shell relatively large (up to 12 mm in length), apex nearly central, not distinctly to the right or left of the median line; animal dextral. Pacific drainage. Superfamily **Lymnaeoidea**, in part . . . . . Family LYMNAEIDAE, in part (page 147)

Adult shell smaller (7 mm or less in length), apex may be nearly central but often to the right or left of the median line; animal dextral or sinistral 15

15(14) Animal and shell dextral (Fig. 755a). Several lakes in the Rocky Mountains, northeastern Ontario and northcentral Quebec. Superfamily **Acroloxoidea** . . . . . Family ACROLOXIDAE (page 147)

Animal and shell sinistral (Fig. 755b). Generally distributed throughout North America. Superfamily **Ancyloidea**, in part . . . . . Family ANCYLIDAE (page 188)

FAMILY NERITINIDAE

The Neritinidae[1] are largely marine and are well represented throughout the world, especially in tropical and subtropical regions. There has been a tendency for various lineages of neritinids to invade estuarine habitats, and freshwater and terrestrial ones as well. Only one species occurs in the United States, *Neritina reclivata* (Say) (Figs. 21, 22). It is found from Florida to Mississippi. Dall (1885) named a subspecies (*palmae*) from near Palma Sola, Florida, and Pilsbry (1931) named a subspecies (*sphaera*) from Ojus, Florida. Both of these may be simply "forms" of *N. reclivata*.

The shells of neritinids are usually subglobose or hemispherical, have few whorls, very reduced spires and very large body whorls. These characteristics, together with the generally thickened shell with heavily calloused and expanded parietal apertural margin, produce a rather typical shape, referred to as *neritiform* (Burch, 1968) or *neritiniform*. The shell is generally smooth, often polished, and its columellar margin is toothed. The operculum (Fig. 22) is paucispiral, calcified, and contains a pair of projections, or apophyses, on the inner columellar side.

The shell of *Neritina reclivata palmae* Dall is "quite small [maximum length 1 cm], . . . black, with a cerous labrum, but the light zigzag lines, characteristic of some color varieties of *reclivata*, [are] beautifully clear by transmitted light" (Dall, 1885).

The shell of *Neritina reclivata sphaera* Pilsbry "is less elevated than *N. reclivata*, the spire *extremely short*, rising very little, the last whorl *strongly convex above the periphery*, not flattened and sloping as in *reclivata*. Color grape green, densely marked with fine black lines and with a black line following the suture, as in *reclivata*" (Pilsbry, 1931).

FAMILY VALVATIDAE

By William H. Heard

The Valvatidae comprise a total of about 20 extant species inhabiting permanent standing and flowing fresh waters in the Northern Hemisphere. Except for *Borysthenia naticina* (Menke) of the Danube River drainage in eastern Europe, the family is represented by species of the genus *Valvata* Müller. The animals of *Valvata* are oviparous hermaphrodites. A single bipectinate gill is directed to the left, and a pallial tentacle occurs on the right side of the mantle cavity (Fig. 16).

The shells of North American *Valvata* are comparatively small (diameter up to 5 mm), have up to 4½ whorls, are dextral, and vary in form from discoid to high-turbinate. The nuclear whorls possess both axial and spiral sculpture; the rest of the shell contains lamellate to obsolete axial sculpture and is either spirally angulated, carinated or smooth. Several species are polymorphic in shell form and sculpture. The operculum is corneous, thin, flattened but slightly concave, circular in outline and multispiral (Fig. 15a).

Shell features are used to identify North American species of *Valvata*, several of which are polymorphic. For example, the "kinds" of *V. tricarinata* s.lat. are characterized by differing numbers and locations of spiral carinae or angulations. A single population usually contains several of these variants, which have often been treated taxonomically as subspecies. However, these variants are neither geographical races nor environmental forms (ecophenotypes), and they are treated as morphs here. *V. lewisi* morph *ontarioensis* (Fig. 27), which often comprises monomorphic populations, does

have a distinctive range, but it is called a morph because of its peculiar shell form. The nature of the variation in some other species is not understood at this time, and several variants are thus treated as possible forms.

The extensive polymorphism in some species has not precluded the construction of a dichotomous key comprised of two alternative choices per couplet, but has in four places provided for a more convenient choice among three alternatives (see "couplets" 2, 3, 5 and 8, below). Extremely rare, atypical variations (e.g., disjunctly coiled *Valvata sincera* s.str. and *V. tricarinata* s.str., and also tetracarinate *V. tricarinata* s.str.) are not included here.

Identification Key for the Valvatidae

1 Shell with one to three postnuclear spiral carinae or angulations 2

Shell lacking postnuclear spiral carinae or angulations 9

2(1) Shell with one spiral carina or angulation 3

Shell with two spiral carinae or angulations 5

Shell with three spiral carinae or angulations 8

3(2) Carina or angulation in dorsal location on the body whorl 4

Carina or angulation in peripheral location on the body whorl. Quebec and New Brunswick west to Alberta, and south to Wyoming, Arkansas and Virginia ..... *Valvata tricarinata* morph *mediocarinata* F.C. Baker

Carina or angulation in ventral location on the body whorl. Quebec and New Brunswick west to Alberta, and south to Wyoming, Arkansas and Virginia ..... *Valvata tricarinata* morph *infracarinata* Vanatta

4(3) Angulation incomplete, becoming obsolete toward the outer lip of the aperture (Fig. 34). Idaho and Utah ..... *Valvata utahensis utahensis* Call

Carina or angulation complete, continuing to the outer lip of the aperture. Quebec and New Brunswick west to Alberta, and south to Wyoming, Arkansas and Virginia ..... *Valvata tricarinata* morph *unicarinata* DeKay

5(2) Carinae or angulations in dorsal and peripheral locations on the body whorl. Quebec and New Brunswick west to Alberta, and south to Wyoming, Arkansas and Virginia ..... *Valvata tricarinata* morph *basalis* Vanatta

Carinae or angulations in peripheral and ventral locations on the body whorl. Quebec and New Brunswick west to Alberta, and south to Wyoming, Arkansas and Virginia ..... *Valvata tricarinata* morph *bakeri* Fluck

Carinae or angulations in dorsal and ventral locations on the body whorl 6



FIGS. 21-25. Shells of Neritinidae (Figs. 21, 22) and Valvatidae (Figs. 23-25). FIG. 21. *Neritina reclivata reclivata*, shell. FIG. 22. *N. reclivata reclivata*, operculum; external view (on left) and internal view (on right). FIG. 23. *Valvata bicarinata*, spire, apertural and umbilical views (left to right). FIG. 24. *V. bicarinata* morph *normalis*. FIG. 25. *V. humeralis*. Measurement lines = 1 mm or are divided into millimeters. Figs. 21-25 are from Burch & Tottenham (1980).

6(5) Shoulder on the body whorl sloping upward from the dorsal carina or angulation to the suture 7

Shoulder on the body whorl sloping downward from the dorsal carina to the suture (Fig. 23). Discontinuously distributed in eastern United States from New Jersey south to Alabama and west to Iowa . . . . . *Valvata bicarinata bicarinata* Lea

7(6) Dorsal angulation incomplete, becoming obsolete on the body whorl. Idaho and Utah . . . . . *Valvata utahensis* morph *horati* Baily & Baily

Dorsal carina or angulation complete, continuing to the outer lip of the aperture. Quebec and New Brunswick west to Alberta, and south to Wyoming, Arkansas and Virginia . . . . . *Valvata tricarinata* morph *perconfusa* Walker

8(2) Shoulder of the body whorl sloping downward from the dorsal carina to the suture (Fig. 24). Discontinuously distributed in eastern United States from New Jersey south to Alabama and west to Iowa . . . . . *Valvata bicarinata* morph *normalis* Walker

Shoulder of the body whorl sloping upward from the dorsal carina or angulation to the suture (Fig. 33). Quebec and New Brunswick west to Alberta, and south to Wyoming, Arkansas and Virginia . . . . . *Valvata tricarinata tricarinata* (Say)

Shoulder of the body whorl sloping upward from the dorsal carina nearly to the suture, then turning downward (Fig. 36). Michigan, Minnesota and Wisconsin . . . . . *Valvata winnebagoensis* F.C. Baker

9(1) Shell partly uncoiled with the body whorl broadly separated from the penultimate whorl (Fig. 27). Ontario in the region of Lake Superior drained by the headwaters of the Attawapiskat, Albany and Severn river systems . . . . . *Valvata lewisi* morph *ontarioensis* F.C. Baker

Shell not disjunctly coiled 10

10(9) Shell of discoid shape. Lakes Erie, Huron, Michigan and Ontario . . . . . *Valvata perdepressa* ?form *walkeri* F.C. Baker

Shell with spire elevated above the body whorl 11

11(10) Shoulder of the body whorl flattened, sloping slightly upward toward the suture; often with a very faint angulation in dorsal location (and rarely also in peripheral locations). Quebec and New Brunswick west to Alberta, and south to Wyoming, Arkansas and Virginia . . . . . *Valvata tricarinata* morph *simplex* Gould

Body whorl evenly convex, not flattened above (or elsewhere) 12



Reproduced from best available copy.

FIGS. 26-30. Shells of Valvatidae. FIG. 26. *Valvata lewisi*, spire, apertural and umbilical views (left to right). FIG. 27. *V. lewisi* morph *ontarioensis*. FIG. 28. *V. mergella*. FIG. 29. *V. perdepressa*. FIG. 30. *V. piscinalis* ?form *obtusa*. Measurement lines = 1 mm. Figs. 26-30 are from Burch & Tottenham (1980).

12(11) Shell depressed-turbinate, spire but little elevated 13

Shell high-turbinate or subconical, spire markedly elevated 16

13(12) Shell diameter exceeding 5 mm 14

Shell diameter less than 5 mm 15

14(13) Axial striae lamellate; luster of shell dull (Fig. 26). Southern Canada from Quebec west to British Columbia, and northern United States from New York west to Minnesota . . . . . *Valvata lewisi lewisi* Currier

Axial striae obsolete; shell with a high gloss (Fig. 28). Alaska to Washington state . . . . . *Valvata mergella* Westerlund

15(13) Color of the apical whorls of the shell usually dull purple, or violet or pink; luster of shell dull (Fig. 29). Lakes Erie, Huron, Michigan and Ontario . . . . . *Valvata perdepressa perdepressa* Walker

Color of the apical whorls of the shell pale green to white; shell glossy (Fig. 25). Montana south to Colorado, west to British Columbia and California and south into Mexico . . . . . *Valvata humeralis* Say

16(12) Shell high-turbinate 17

Shell subconical 19

17(16) Apex of shell flattened, appearing truncated (Fig. 30). Lower Great Lakes . . . . . *Valvata piscinalis* ?form *obtusa* Draparnaud

Apex of shell acute 18

18(17) Shell color pale green; shell diameter greater than 5 mm (Fig. 32). Eastern Canada and north central United States . . . . . *Valvata sincera* ?form *danielsi* Walker

Shell color dark to often brilliant green; shell diameter less than 5 mm (Fig. 35). California, Nevada and Oregon . . . . . *Valvata virens* Tryon

19(16) Axial striae lamellate. Quebec and Maine west to Ontario and Minnesota . . . . . *Valvata sincera nylanderi* Dall

Axial striae fine (Fig. 31). Maine west to Alberta, and south to South Dakota and Illinois . . . . . *Valvata sincera sincera* Say



FIGS. 31-35. Shells of Valvatidae. FIG. 31. *Valvata sincera sincera*, spire, apertural and umbilical views (left to right). FIG. 32. *V. sincera* ? form *danielsi*. FIG. 33. *V. tricarinata*. FIG. 34. *V. utahensis*. FIG. 35. *V. virens*. Measurement lines = 1 mm or are divided into millimeters. Figs. 31-35 are from Burch & Tottenham (1980).

Reproduced from best available copy.



FIGS. 36-43. Shells of Valvatidae (Fig. 36) and Viviparidae (Lioplacinae) (Figs. 37-43). FIG. 36. *Valvata winnebagoensis*, spire, apertural and umbilical views (left to right). FIG. 37. *Campeloma lewisi* = *C. decisum*. FIG. 38. *C. integrum* = *C. decisum*. FIG. 39. *C. milesi* = *C. decisum*. FIG. 40. *C. regulare*. FIG. 41. *C. limum*. FIG. 42. *C. crassula*, operculum. FIG. 43. *Lioplax cyclostomaformis*. Measurement lines = 1 mm or are divided into millimeters. Figs. 36-43 are from Burch & Tottenham (1980).

FAMILY VIVIPARIDAE

By J.B. Burch and Virginia A. Vail

The Viviparidae are nearly world-wide in distribution and in North America occur throughout the eastern United States and Canada. *Campeloma, Lioplax* and *Tulotoma* are endemic to (i.e., restricted to) North America. *Viviparus* has a Holarctic distribution, and *Cipangopaludina* is an Asian genus. *Campeloma, Lioplax* and *Viviparus* are relatively common and have wide distributions. *Tulotoma* is confined to the Coosa-Alabama river system in Alabama and is rare, perhaps now nearly extinct. The two introduced species of the Asian *Cipangopaludina* have rather wide although sporadic distributions in the United States.

The Viviparidae are all "live-bearers", i.e., are ovoviviparous, giving birth to young crawling snails, rather than laying eggs that hatch in the external environment. It is this reproductive trait which has provided the family with its name.

The sexes are separate in the Viviparidae, the males being readily distinguishable by their modified right tentacle, which serves as a copulatory organ. This modified tentacle in the males is shorter and thicker than the left tentacle or either of the bilaterally symmetrical tentacles of the females. Some populations of *Campeloma* are parthenogenetic, consisting entirely of females.

Identification Key for the Viviparidae

1 Shell large, adults over 35 mm and up to 50 mm in length; shell relatively thin; whorls not shouldered. Genus *Cipangopaludina* 2

Shell medium to large, generally less than 35 mm in length, but if large, the shell is thick and ponderous, and the whorls are generally shouldered 3

2(1) Shell with acute spire and usually with spiral angulations or low carinae on the whorls; not malleated (Fig. 53). Sporadically but widely distributed in the United States ..... *Cipangopaludina japonica* (Martens)

Shell with obtuse spire and without spiral angulations or low carinae; generally with surface malleations (Fig. 52). Sporadically but widely distributed in the United States ..... *Cipangopaludina chinensis malleata* (Reeve)

3(1) Shell with or without one or two spiral rows of nodules; outer margin of shell aperture concave (when observed from an angle parallel to the plane of the aperture) and its oblique margin to the shell axis quite exaggerated (Fig. 772); columellar margin of operculum reflected inward; restricted to the Coosa-Alabama river system in Alabama ..... *Tulotoma magnifica* (Conrad)[3]

Shell without rows of spiral nodules; outer margin of shell aperture not concave (when observed from an angle parallel to the plane of the aperture) and its oblique angle to the shell axis not exaggerated (Fig. 772); columellar margin of operculum not reflected inward 4

4(3) Operculum concentric, but with spiral nucleus; whorls commonly with a median spiral angle or low ridge or a spiral subsutural sulcus. Genus *Lioplax* 5

Operculum entirely concentric, including its nucleus; whorls without spiral angles, ridges or sulci 10

5(4) Shell attenuate, compressed; whorls rarely angular (Fig. 43). Coosa-Alabama-Tombigbee river system in Georgia and Alabama, and Tensas River, Alabama . . . . . *Lioplax cyclostomaformis* (Lea)

Shell subglobose, not attenuate and compressed; at least some of the whorls are generally angular or with a spiral subsutural sulcus 6

6(5) Shell large for the genus, adults up to 30 mm in length, dark olive-green to nearly black (Fig. 67). Chipola River, Florida . . . . . *Lioplax pilsbryi pilsbryi* Walker

Shell smaller, adults less than 25 mm in length and seldom more than 20 mm, horn to pale or occasionally dark olive-green in color 7

7(6) Atlantic drainage and Gulf drainage 8

Mississippi drainage (Minnesota to Arkansas and Ohio)[7] . . . . . *Lioplax sulculosa* (Menke)

8(7) Atlantic drainage (New York to South Carolina)[7] (Fig. 68) . . . . . *Lioplax subcarinata* (Say)

Gulf drainage 9

9(8) Whorls generally with a spiral subsutural sulcus, which tends to constrict the posterior aperture (Fig. 69). Ochlockonee and Yellow river systems, Florida and Alabama . . . . . *Lioplax talquinensis* Vail

Whorls without a spiral subsutural sulcus; aperture rounded posteriorly. Choctawhatchee, Escambia, Flint and Suwannee river systems, Florida and Georgia . . . . . *Lioplax pilsbryi choctawhatchensis* Vanatta[6]

10(4) Shell with or without spiral color bands; width and length of aperture usually nearly equal, making it round, or nearly so; lateral and marginal radular teeth with prominent cusps. Genus *Viviparus* 11

Shell without spiral color bands; length of aperture noticeably greater than width; lateral and marginal teeth simple with very fine, difficult-to-distinguish cusps. Genus *Campeloma*[4] 13

11(10) Shell dark yellowish-green to (usually) dark olivaceous-green, without spiral color bands; shell broadly ovate, whorls globosely rounded, spire obtuse (Fig. 48). Minnesota south to Louisiana, mainly in the Mississippi river drainage; Gulf drainage from Texas to Alabama; Atlantic drainage in Georgia and South Carolina . . . . . *Viviparus intertextus* (Say)



FIGS. 44-53. Shells of Viviparidae (Viviparinae and Bellamyinae). FIG. 44. *Tulotoma magnifica*. FIG. 45. *T. angulata* ? = *T. magnifica*. FIG. 46. *Viviparus georgianus*. FIG. 47. *V. georgianus*. FIG. 48. *V. intertextus*. FIG. 49. *V. subpurpureus*. FIG. 50. *V. subpurpureus*. FIG. 51. *V. subpurpureus*. FIG. 52. *Cipangopaludina chinensis malleata*. FIG. 53. *C. japonica*. Measurement lines are divided into millimeters. Figs. 44-53 are from Burch & Tottenham (1980).

Reproduced from best available copy.

Shell pale olivaceous-green to olivaceous-brown, with or without spiral color bands, ovate but not broadly so, whorls flattened to well rounded but not globosely rounded, spire relatively acute 12

12(11) Shell yellowish-brown or olivaceous-brown; color bands, when present, three in number; shell rather heavy; whorls often flat-sided (Figs. 49-51). Mississippi river drainage from Iowa to Louisiana; Gulf drainage in Texas and Mississippi ..... *Viviparus subpurpureus* (Say)

Shell yellowish-green or olivaceous-green; color bands, when present, usually four in number; shell relatively thin, but sturdy; whorls usually well rounded (Figs. 46, 47). Alabama, Florida and Georgia north to Illinois and Indiana; northern states from Wisconsin to New England and Quebec ..... *Viviparus georgianus* (Lea)

13(10) Inside of shell aperture deep reddish-brown or brown (Fig. 62); shell of newborn young uniformly dark brown. Eastern Florida .... *Campeloma floridense* Call

Inside of shell aperture white, bluish or faintly pinkish; shell of newborn young opaque white or light translucent beige 14

14(13) Shell whorls generally with angled shoulders. Southern in distribution 15

Shell whorls unshouldered or with rounded shoulders 16

15(14) Shell broadly ovate (Figs. 63, 64). Northwestern Florida, southwestern Georgia and southeastern Alabama ..... *Campeloma geniculum* (Conrad)

Shell narrowly ovate (Fig. 41). Atlantic drainage from North Carolina to Georgia ..... *Campeloma limum* (Anthony)

16(14) Shell narrow, relatively thin, generally with prominent raised spiral lines (Fig. 56). Northern Alabama ..... *Campeloma decampi* Binney

Shell broader, relatively thin to thick and ponderous, spiral lines on adult shells when present are not prominent 17

17(16) Spire typically depressed and obtuse, body whorl large and often cylindrical (Figs. 40, 66). Alabama-Coosa drainage ..... *Campeloma regulare* (Lea)

Spire elongate, seldom depressed, body whorl rounded 18

18(17) Shell large, heavy and ponderous (Figs. 42, 54, 55). Midwestern United States in the Great Lakes-St. Lawrence and Mississippi drainages .... *Campeloma crassula* Rafinesque

Shell medium or a little larger, relatively thin to strong, but not very large or heavy and ponderous 19



FIGS. 54-63. Shells of Viviparidae (Lioplacinae). FIG. 54. *Campeloma subsolidum* = *C. crassula*. FIG. 55. *C. obesum* = *C. crassula*. FIG. 56. *C. decampi*. FIG. 57. *C. decisum*. FIGS. 58, 59. *C. brevispirum* = *C. decisum*. FIG. 60. *C. gibbum* = *C. decisum*. FIG. 61. *C. tannum* = *C. decisum*. FIG. 62. *C. floridense*. FIG. 63. *C. geniculum*. Measurement lines are divided into millimeters. Figs. 54-63 are from Burch & Tottenham (1980).

Reproduced from best available copy.



FIGS. 64-71. Shells of Viviparidae (Lioplacinae) (Figs. 64-69) and Ampullariidae (Figs. 70, 71). FIG. 64. *Campeloma geniculum*. FIG. 65. *C. parthenum*. FIG. 66. *C. regulare*. FIG. 67. *Lioplax pilsbryi* s.str. FIG. 68. *L. subcarinata*. FIG. 69. *L. talquinensis*. FIG. 70. *Marisa cornuarietis*, apertural (left figure), spire (right top figure) and umbilical (right bottom figure) views. FIG. 71. *M. cornuarietis*, operculum. Measurement lines are divided into millimeters. Figs. 64-71 are from Burch & Tottenham (1980).

Reproduced from best available copy.

19(18) Widely distributed, from southern Canada to Texas, Louisiana, Mississippi, Alabama, northern Georgia and Virginia. Figs. 37-39, 57-61 .... *Campeloma decisum* (Say)

Ochlockonee river drainage in southern Georgia and northern Florida. Fig. 65 ..... *Campeloma parthenum* Vail

FAMILY AMPULLARIIDAE

The family Ampullariidae contains the "apple snails", many of which are very large and globose or subglobose in shape. The family is represented world-wide in the tropics. They are mostly amphibious snails which can survive for long periods out of water, including during the dry season when they burrow into the mud. Their mantle cavity is divided into two compartments, the left one containing a gill for aquatic respiration and the right compartment serving as a lung for air-breathing. From the left side a long siphon extends, by which the snail can admit air to the pulmonary chamber when immersed.

Pomacea paludosa (Say) is the largest freshwater gastropod found in North America, its height and width commonly exceeding 60 mm. Its color is dark to light olive green with a dozen or more reddish or brownish spiral bands. The operculum is concentric, thin and corneous. Pilsbry (1899e) gave the name *miamiensis* to a small, reddish-brown population from the vicinity of Miami, Florida, but according to Clench & Turner (1956) this is a synonym of *paludosa* Say. A Brazilian species, *P. bridgesi* (Reeve), recently has been introduced to Florida (Clench, 1966).

Marisa cornuarietis is also a large snail, and its shell also has an olive color with spiral reddish or brown bands. However, the shell is peculiar in that its spire is sunken below the body whorl and the umbilicus is very wide.

Identification Key for the Ampullariidae

1 Shell subglobose in shape. Alabama, Florida and Georgia. Genus *Pomacea* 2

Shell discoidal or planispiral in shape (Figs. 70, 71). Southern Florida .... *Marisa cornuarietis* (Linnaeus)

2(1) Shell large, often up to 60 mm or more in length, whorls with only weak or without shoulders, body whorl very wide, spire depressed, aperture narrowly oval (Figs. 72, 73). Alabama, Florida and Georgia .... *Pomacea paludosa* (Say)

Shell smaller, less than 50 mm in length, whorls more strongly shouldered, body whorl narrower, spire projecting and turreted, aperture more broadly oval. Florida ..... *Pomacea bridgesi* (Reeve)

FAMILY BITHYNIIDAE

The Bithyniidae are found throughout Europe and Asia, and in Africa, Indonesia, the Philippines and Australia. The European *Bithynia tentaculata* (Linnaeus) was introduced long ago by man into



Reproduced from best available copy.

FIGS. 72-80. Shells of Ampullariidae (Figs. 72, 73) and Hydrobiidae (Hydrobiinae) (Figs. 74-80). FIG. 72. *Pomacea paludosa*, apertural (left figure), spire (right top figure) and umbilical (right bottom figure) views. FIG. 73. *P. paludosa*, operculum. FIG. 74. *Bithynia tentaculata magnalacustris*. FIG. 75. *Aphaostracon rhadinus*. FIG. 76. *Hoyia sheldoni*. FIG. 77. *Hyalopyrgus aequicostatus*, female. FIG. 78. *H. aequicostatus*, male. FIG. 79. *Littoridinops tenuipes*. FIG. 80. *L. monroensis*. Measurement lines = 1 mm or are divided into millimeters.

North America and has spread widely. However, *B. tentaculata* has been reported in Pleistocene deposits in Chicago, so it may already have been living in the Western Hemisphere when Europeans first arrived. F.C. Baker (1928c) gave the varietal name *magnalacustris* to the supposedly North American form, which he considered to have "more rounded whorls with deeper sutures and an apex that stands well above the second whorl."

The Bithyniidae traditionally have been included in the family Hydrobiidae. However, Taylor (1966b) has recently argued that the bithyniids should be separated from the hydrobiids and transferred to the Viviparoidea (Ampullarioidea). Viviparoid characters of *Bithynia* are its size (adult shells are more than 10 mm long), calcareous operculum with paucispiral nucleus and concentric edges, nuchal lobes of the head-foot, relatively long, flexible and acute tentacles, yellow and orange skin pigment granules, spirally constructed fecal pellets, use of the ctenidium in food gathering, pallial innervation of the penis, and dimorphic sperm.

Bithynia tentaculata (Linnaeus) has a broadly conic or narrowly ovate shell (Fig. 74). It is larger than any of the Hydrobiidae, the shells of many adults measuring more than 12 mm in length. The color of the shell ranges from yellowish to greenish, and is covered by a thin brownish periostracum. Surface sculpture consists of fine transverse growth lines and fine incised spiral lines. In contrast to most hydrobiids, the concentric operculum just fits the outer aperture, and does not go past the peritreme when the animal withdraws its head-foot into the shell.

Bithynia tentaculata occurs in the Great Lakes region from Albany, New York, to Winnebago Lake, Wisconsin.

FAMILY MICROMELANIIDAE

The Micromelaniidae are a family of hydrobiid-like snails which lack basal denticles on their central radular teeth. They are found mainly in the ancient lakes Baical (Siberia) and Ohrid (Macedonia and Albania), the Caspian Sea, southeastern Europe, Asia Minor and eastern India. *Emmericiella* occurs in Mexico, and the monotypic *Antroselates* occurs in caves in southern Indiana and west-central Kentucky. The latter was transferred to the Micromelaniidae by Taylor (1966b) because of its radular characters.

Antroselates spiralis Hubricht has a small, solid, globosely conic, turbinate, narrowly perforate or rimate shell (Fig. 108). Its sculpture consists of numerous spiral periostracal threads. The operculum is paucispiral and hyaline. The animal is white. Males have a simple, long, slender, tapering verge. The central and lateral teeth have many small cusps of uniform size (Hubricht, 1963b).

FAMILY HYDROBIIDAE

The Hydrobiidae are one of the most common and widely distributed gastropod families, occurring in temperate, subtropical and tropical regions throughout much of the world. The family is a large one, comprising some 103 genera (Taylor & Sohl, 1962). Most hydrobiid species live in fresh water, although some are associated with brackish water. Only the North American freshwater species are dealt with in this manual.

Shells of hydrobiids are small (many are minute), generally elongate, dextral (Fig. 2b), nearly always drab and unicolored, and generally have relatively few whorls. The shells of most species are plain, but some species have prominent surface sculpture, and one species in North America (north of Mexico), *Cochliopina riograndensis* (Pilsbry & Ferriss) has spiral color bands (Fig. 140). The shell aperture is closed by an operculum, which is generally paucispiral (Fig. 15b), but some species have

round, multispiral opercula (Fig. 15a). Like most North American freshwater prosobranch snails, the sexes are separate in the Hydrobiidae, and the shells of some genera exhibit sexual dimorphism.

Because of the similarity of the shells of many species occurring in different genera and subfamilies, reliance must be placed on anatomical characters, especially those of the verge (male copulatory organ), in making identifications and for assigning species to genera and genera to subfamilies (Fig. 82). Since the anatomical characteristics of some species (and even genera) are not known, their taxonomic placement in this manual is presumptive. Further studies may change their systematic status.

Since so few hydrobiids have been studied anatomically in any great detail, a subfamilial classification based entirely on the male verge may be proven eventually to be inadequate or inaccurate. However, from a standpoint of practicality for presenting a workable classification for this identification manual, the hydrobiid genera are grouped according to the major characters of the verges of their species and these groups assigned to previously named subfamilies. While this possibly may not represent the true systematic and phylogenetic relationships of the various genera, it is a useful system at present.

Identification Key for the Hydrobiidae

1 Males with single-ducted verges (Fig. 82a,b,c) 2

Males with two- or three-ducted verges (Fig. 82d,e) 52

2(1) Males with simple verges, lacking accessory lobes and glandular apical and subapical crests (Fig. 82a). Subfamily Lithoglyphinae 3

Males with verges bearing accessory lobes or glandular apical and subapical crests (Fig. 82b,c) 13

3(2) Shell neritiniform (Figs. 14, 192, 193). Cahaba and Coosa rivers, Alabama . . . . . *Lepyrium showalteri* (Lea)

Shell conical, subglobose or heliciform 4

4(3) Shell depressed, heliciform, with spiral brown bands (Fig. 140). Texas . . . . . *Cochliopina riograndensis* (Pilsbry & Ferriss)

Shell conical to subglobose, without spiral color bands 5

5(4) Shell imperforate or narrowly perforate 6

Shell umbilicate 11

6(5) Western in distribution, in the Pacific drainage (Figs. 141, 142, 145-148, 152) . . . . . Genus *Fluminicola*[63]

Eastern in distribution, in the Mississippi, Gulf and Atlantic drainages 7



FIG. 81. Central radular tooth of truncatelloid snails. a, A micromelaniid, without basal cusps; b, a hydrobiid, with basal cusps on a thickened ridge along the lateral angle of the tooth; c, a pomatiopsid, with basal cusps on the anterio-posterior ridges. b = basal cusp.



FIG. 82. Basic types of verges of North American hydrobiid snails. a, Simple verge with a single duct (Lithoglyphinae); b, verge with a single duct and accessory lobes (Hydrobiinae); c, verge with a single duct and glandular apical and subapical crests (Nymphophilinae); d, verge with two ducts (Amnicolinae); e, verge with three ducts (Fontigentinae).



FIGS. 83-86. Animals of hydrobiid snails (Hydrobiinae), with shells removed. FIG. 83. *Hyalopyrgus aequicostatus*, male, dorsal view. FIG. 84. *H. aequicostatus*, female, dorsal view. FIG. 85. *Littoridinops monroensis*, male, dorsal view with mantle and viscera removed. FIG. 86. *L. monroensis*, male, right lateral view. Measurement lines = 1 mm. Figs. 83-86 are from Thompson (1968).



FIGS. 87-94. Animals of hydrobiid snails (Hydrobiinae, Lithoglyphinae, Nymphophilinae and Amnicolinae), with shells removed. FIG. 87. *Littoridinops tenuipes*, male, right lateral view. FIG. 88. *Pyrgophorus platyrachis*, male, dorsal view. FIG. 89. *Somatogyrus (Walkerilla) tenax*, male. FIG. 90. *Notogillia sathon*, male. FIG. 91. *Rhapinema dacryon*, male. FIG. 92. *Spilochlamys conica*, male. FIG. 93. *Amnicola dalli johnsoni*, female. FIG. 94. *Amnicola (Lyogyrus) retromargo*, female. Measurement lines = 1 mm. Figs. 87-94 are from Thompson (1968, 1969).



FIGS. 95-107. Verges of hydrobiid snails (Hydrobiinae). FIG. 95. *Aphaostracon asthenes*. FIG. 96. *A. chalarogyrus*. FIG. 97. *A. hypohyalina*. FIG. 98. *A. monas*. FIG. 99. *A. rhadinus*. FIG. 100. *A. pachynotus*. FIG. 101. *A. pycnus*. FIG. 102. *A. theiocrenetus*. FIG. 103. *A. xynoelictus*. FIG. 104. *Hyalopyrgus aequicostatus*. FIG. 105. *Littoridinops monroensis*. FIG. 106. *L. tenuipes*. FIG. 107. *Probythinella lacustris*. Measurement lines = ½ mm. Figs. 95-106 are from Thompson (1968); Fig. 107 is from E.G. Berry (1943).

7(6) Shell generally thick and solid, columella thickened. Mississippi and Gulf of Mexico drainage (except for *S. pennsylvanicus* and *S. virginicus*). Genus *Somatogyrus* 8

Shell rather thin, columella not thickened (Fig. 191). Atlantic drainage from New Jersey to South Carolina . . . . . *Gillia altilis* (Lea)

8(7) Shell with spirally striate apical whorls. Subgenus *Walkerilla*[64] 9

Shell without spirally striate apical whorls (Figs. 149, 151, 153-185, 194, 195). Widely distributed in eastern North America in the Midwest and South . . . . . Subgenus *Somatogyrus* s.s.[65]

9(8) Spire very depressed (Figs. 150, 186, 196). Catawba and Coosa rivers, Alabama . . . . . *Somatogyrus (Walkerilla) coosaensis* Walker

Spire not depressed. Georgia and Virginia 10

10(9) Shell perforate (Figs. 89, 197, 201). Broad River, Georgia . . . . . *Somatogyrus (Walkerilla) tenax* Thompson

Shell imperforate (Fig. 187). Rapidan River, Virginia . . . . . *Somatogyrus (Walkerilla) virginicus* Walker

11(5) Shell small (less than 2.5 mm in length), aperture round, columella thin (Fig. 138). Missouri . . . . . *Antrobia culveri* Hubricht

Shell larger (3.0-3.5 mm in length), aperture ovate, columella thickened. Alabama. Genus *Clappia* 12

12(11) Shell aperture more elongate, spire less attenuate, umbilicus larger, animal black (Figs. 139, 143, 144). Coosa River, Alabama . . . . . *Clappia umbilicata* (Walker)

Shell aperture broader, less elongate, spire relatively attenuate, umbilicus smaller, animal white. Cahaba River, Alabama . . . . . *Clappia cahabensis* Clench

13(2) Males with verges bearing accessory lobes (Fig. 82b). Subfamily Hydrobiinae[66] 14

Males with verges bearing glandular apical crests (Fig. 82c). Subfamily Nymphophilinae 27

14(13) Top of shell spire truncated. The first several spire whorls coiled in the same plane (Figs. 107, 129-131). Widely distributed in eastern North America . . . . . *Probythinella lacustris* (F. C. Baker)

Top of shell spire not truncated, the first several spire whorls coiled in a descending spiral 15



FIGS. 108-118. Shells of Micromelaniidae (Fig. 108) and Hydrobiidae (Figs. 109-118). FIG. 108. *Antroselates spiralis*. FIG. 109. *Aphaostracon asthenes*. FIG. 110. *A. chalarogyrus*. FIG. 111. *A. hypohyalina*, female. FIG. 112. *A. hypohyalina*, male. FIG. 113. *A. monas*, female. FIG. 114. *A. monas*, male. FIG. 115. *A. pachynotus*, female. FIG. 116. *A. pachynotus*, male. FIG. 117. *A. pycnus*, female. FIG. 118. *A. pycnus*, male. Measurement lines = ½ mm. Fig. 108 is after Hubricht (1963b); Figs. 109-118 are from Thompson (1968).

15(14) Northern in distribution (Fig. 76). Lake Michigan, Wisconsin . . . . . *Hoyia sheldoni* (Pilsbry)[67]

Southern and western in distribution 16

16(15) Western in distribution. Texas, Arizona, Nevada and California. Genus *Tryonia*[68] 17

Southern in distribution. Georgia and Florida 22

17(16) Found in Texas 18

Further western in distribution, Arizona, Nevada and California 20

18(17) Shell minute, that of adults with four to five whorls less than 1.5 mm in shell length; umbilicus small but distinct (Fig. 135). Texas . . . . . *Tryonia diaboli* (Pilsbry & Ferriss)

Shell larger, that of adults with about five whorls more than 3 mm; imperforate 19

19(18) Shell surface smooth, except for fine transverse growth lines (Figs. 127, 128, 133). Texas . . . . . *Tryonia cheatumi* (Pilsbry)

Shell surface sculptured with revolving striae or carinae which are commonly modified into spines (Fig. 126). Texas . . . . . *Pyrgophorus spinosus* (Call & Pilsbry)

20(17) Shell surface smooth, except for fine transverse growth lines. California (in brackish water), Arizona . . . . . *Tryonia imitator* (Pilsbry)

Shell surface sculptured with transverse ribs and sometimes with spiral lirae also 21

21(20) Shell narrowly conic, ribbed, with or without lirae, ribs not angular except where crossed by lirae (Figs. 136, 137). California (subfossil), Arizona . . . . . *Tryonia protea* (Gould)

Shell elongately conic, ribbed, but without lirae, ribs angular (Fig. 134). Nevada . . . . . *Tryonia clathrata* Stimpson

22(16) Periphery of whorls flattened, sutures shallow; verge with 7-50 papillae along its right margin, 1-4 papillae along the distal third of the left margin and with or without papillae about the base. Genus *Littoridinops* 23

Periphery of whorls inflated, sutures impressed; verge with 1-7 papillae along the right margin and usually with one or two papillae on the left margin either at the base or distal end 24



FIGS. 119-128. Shells of Hydrobiidae (Hydrobiinae). FIG. 119. *Aphaostracon theiocrenetus*, female. FIG. 120. *A. theiocrenetus*, male. FIG. 121. *A. xynoelictus*, female. FIG. 122. *A. xynoelictus*, male. FIG. 123. *Hyalopyrgus brevissimus*, female. FIG. 124. *H. brevissimus*, male. FIG. 125. *Littoridinops tenuipes*. FIG. 126. *Pyrgophorus spinosus*. FIG. 127. *Tryonia cheatumi*. FIG. 128. *T. cheatumi*. Measurement lines = ½ mm. Figs. 119-124 are from Thompson (1968); Fig. 126 is from Call & Pilsbry (1886); Fig. 127 is from Pilsbry (1935a).



FIGS. 129-142. Shells of Hydrobiidae (Hydrobiinae and Lithoglyphinae). FIG. 129. *Probythinella lacustris*. FIG. 130. *Pr. lacustris*. FIG. 131. *Pr. lacustris*. FIG. 132. *Pyrgophorus platyrachis*. FIG. 133. *Tryonia cheatumi*. FIG. 134. *T. clathrata*. FIG. 135. *T. diaboli*. FIG. 136. *T. protea*. FIG. 137. *T. protea*. FIG. 138. *Antrobia culveri*. FIG. 139. *Clappia clappi* = *C. umbilicata*. FIG. 140. *Cochliopina riograndensis*. FIG. 141. *Fluminicola fusca*. FIG. 142. *F. nuttalliana*. Measurement lines = 1 mm or are divided into millimeters.

Reproduced from best available copy.



FIGS. 143-152. Shells of Hydrobiidae (Lithoglyphinae). FIG. 143. *Clappia clappi* = *C. umbilicata*. FIG. 144. *C. umbilicata*. FIG. 145. *Fluminicola columbiana*. FIG. 146. *F. merriami*. FIG. 147. *F. minutissima*. FIG. 148. *F. nevadensis*. FIG. 149. *Somatogyrus amnicoloides*. FIG. 150. *S. aldrichi* = *S. (Walkerilla) coosaensis*. FIG. 151. *S. aureus*. FIG. 152. *F. turbiniformis*. Measurement lines = 1 mm. Figs. 143, 144 and 148-150 are from Walker (1904a, 1906a, 1909c, 1915c, 1916); Figs. 145 and 146 are from Stearns (1901b); Fig. 147 is from Pilsbry (1908a); Figs. 151 and 152 are from Tryon (1865i).



FIGS. 153-161. Shells of Hydrobiidae (Lithoglyphinae). FIG. 153. *Somatogyrus biangulatus*. FIG. 154. *S. constrictus*. FIG. 155. *S. crassilabris*. FIG. 156. *S. crassus*. FIG. 157. *S. crassus*, immature. FIG. 158. *S. currierianus*. FIG. 159. *S. decipiens*, immature. FIG. 160. *S. decipiens*. FIG. 161. *S. excavatus*. Measurement lines = 1 mm. Figs. 153-161 are from Walker (1904a, 1906a, 1909c, 1915c).



FIGS. 162-171. Shells of Hydrobiidae (Lithoglyphinae). FIG. 162. *Somatogyrus excavatus*. FIG. 163. *S. georgianus*. FIG. 164. *S. hendersoni*. FIG. 165. *S. hinkleyi*. FIG. 166. *S. hinkleyi*, immature. FIG. 167. *S. humerosus*. FIG. 168. *S. nanus*. FIG. 169. *S. obtusus*. FIG. 170. *S. parvulus*. FIG. 171. *S. parvulus*. Measurement lines = 1 mm. Figs. 163-170 are from Walker (1904a, 1906a, 1909c); Fig. 171 is from Tryon (1865i).



FIGS. 172-180. Shells of Hydrobiidae (Lithoglyphinae). FIG. 172. *Somatogyrus pennsylvanicus*. FIG. 173. *S. pennsylvanicus*. FIG. 174. *S. pennsylvanicus*. FIG. 175. *S. pilsbryanus*. FIG. 176. *S. pumilus*. FIG. 177. *S. pygmaeus*. FIG. 178. *S. quadratus*. FIG. 179. *S. sargenti*. FIG. 180. *S. strengi*. Measurement lines = 1 mm. Figs. 172-180 are from Walker (1904a, 1906a, 1909c).



FIGS. 181-190. Shells of Hydrobiidae (Lithoglyphinae and Nymphophilinae). FIG. 181. *Somatogyrus substriatus*. FIG. 182. *S. tennesseensis*. FIG. 183. *S. wheeleri*, apex. FIG. 184. *S. wheeleri*. FIG. 185. *S. walkerianus*. FIG. 186. *S. (Walkerilla) coosaensis*. FIG. 187. *S. (W.) virginicus*. FIG. 188. *Birgella subglobosa*. FIG. 189. *Cincinnatia judayi* = *C. cincinnatiensis*. FIG. 190. *C. comalensis*. Measurement lines = 1 mm or are divided into millimeters. Figs. 181-184, 186 and 187 are from Walker (1904a, 1906a, 1915c); Fig. 185 is from Aldrich (1905); Fig. 190 is from Pilsbry & Ferriss (1906).

23(22) Verge with a single row of 7-15 papillae along the right margin and 3-10 papillae around the base (Figs. 79, 87, 106, 125). Atlantic drainage of Florida and Georgia . . . . . *Littoridinops tenuipes* Couper

Verge with 17-50 papillae arranged in three to five rows along the right margin, and no papillae at the base (Figs. 80, 85, 86, 105). Florida . . . . . *Littoridinops monroensis* (Frauenfeld)

24(22) Shell sculptured with fine spiral lines; verge with 1-7 papillae along the right margin and papillae along the left margin 25

Shell without fine spiral sculpturing; verge with 0-6 papillae along the right margin, no other papillae present (Figs. 75, 95-103, 109-122). Florida . . . . . Genus *Aphaostracon*[69]

25(24) Spiral sculpturing consisting of raised threads; verge with 3-7 papillae along the right margin, left margin usually with a papilla near the base and 1-4 papillae on a projection near the distal end (Figs. 88, 132). Southern Florida . . . . . *Pyrgophorus platyrachis* Thompson

Spiral sculpturing consisting of fine incised striations; verge with one large papilla on the right margin near the base, and one or two smaller papillae on the left margin near the distal end. Genus *Hyalopyrgus* 26

26(25) Shell elongated conical, rimate or imperforate; verge with two papillae and an apical protrusion on the left margin (Figs. 77, 78, 83, 84, 104). Florida . . . . . *Hyalopyrgus aequicostatus* (Pilsbry)

Shell ovate, openly umbilicate; verge with one papilla on the left margin (Figs. 123, 124). Central Florida . . . . . *Hyalopyrgus brevissimus* (Pilsbry)

27(13) Shell almost completely uncoiled (Fig. 248). Texas . . . . . Genus *Orygoceras*

Shell coiled 28

28(27) Shell relatively large (that of adults to nearly 10 mm in length), subglobose (Figs. 188, 198, 202). Widely distributed in central United States from the Great Lakes to Alabama and Arkansas . . . . . *Birgella subglobosa* (Say)

Shell smaller (that of adults generally less than 5 mm in length), globosely conic to broadly conic and rarely elongately conic, or subglobose, ovate or turbiniform 29

29(28) Shell turbiniform, minute (that of adults 1.2-1.4 mm long) (Figs. 265, 297). Alabama river system . . . . . *Stiobia nana* Thompson

Shell conic, subglobose or ovate 30



FIGS. 191-200. Shells of Hydrobiidae (Lithoglyphinae and Nymphophilinae). FIG. 191. *Gillia altilis*. FIG. 192. *Lepyrium showalteri*. FIG. 193. *L. showalteri*, operculum. FIG. 194. *Somatogyrus aureus*. FIG. 195. *S. depressus*. FIG. 196. *S. (Walkerilla) coosaensis*. FIG. 197. *S. (W.) tenax*. FIG. 198. *Birgella subglobosa*. FIG. 199. *Cincinnatia cincinnatiensis*. FIG. 200. *C. fraterna*. Measurement lines = 1 mm.

Reproduced from best available copy.



FIGS. 201-210. Verges of hydrobiid snails (Lithoglyphinae and Nymphophilinae). FIG. 201. *Somatogyrus (Walkerilla) tenax*. FIG. 202. *Birgella subglobosa*. FIG. 203. *Cincinnatia cincinnatiensis*. FIG. 204. *C. floridana*. FIG. 205. *C. fraterna*. FIG. 206. *C. helicogyra*. FIG. 207. *C. mica*. FIG. 208. *C. monroensis*. FIG. 209. *C. parva*. FIG. 210. *C. petrifons*. Measurement lines = 1 mm. Figs. 201 and 204-210 are from Thompson (1968); Figs. 202 and 203 are from E.G. Berry (1943).



FIGS. 211-221. Verges of hydrobiid snails (Nymphophilinae). FIG. 211. *Cincinnatia ponderosa*. FIG. 212. *C. vanhyningi*. FIG. 213. *C. wekiwae*. FIG. 214. *Marstonia agarhecta*. FIG. 215. *M. arga*. FIG. 216. *M. castor*. FIG. 217. *M. halcyon*. FIG. 218. *M. lustrica*. FIG. 219. *M. lustrica*. FIG. 220. *M. ogmorphaphe*. FIG. 221. *M. pachyta*. Measurement lines = ½ mm. Figs. 211-218, 220 and 221 are from Thompson (1968, 1969, 1977); Fig. 219 is from E.G. Berry (1943).

30(29) Verge with a relatively simple glandular pattern 31

Verge with elaborate patterns of many glands 39

31(30) Shell elongate (conic or ovate); verge with a short terminal penis. Widely distributed in eastern North America. Genus *Marstonia*[70] 32

Shell subglobose; verge with a long, slender flagellar penis (Figs. 91, 257, 262). Chipola river drainage, Florida[70] . . . . . *Rhapinema dacryon* Thompson

32(31) Shell minute, that of adults (with four or more whorls) less than 2.7 mm in length; verge with an elongate apical lobe, penis large and robust 33

Shell larger, that of adults (with 4½ or more whorls) 3.5 mm long; verge with a squarish apical lobe, penis short and slender 35

33(32) Shell thin, fragile, transparent, conical, with an incomplete peristome across the parietal margin; verge with a single gland on the apical lobe (Figs. 214, 232). Ocmulgee river system, Georgia . . . . . *Marstonia agarhecta* Thompson

Shell thick, solid, nearly opaque, ovate-conical; peristome complete across the parietal margin; verge with two glands, one near the base and one on the apical lobe 34

34(33) Shell broadly ovate, 0.70-0.80 times as wide as high, whorls strongly shouldered, flattened at the shell periphery, umbilicus wide, suture descending in lateral profile (Figs. 217, 249). Ogeechee river system, Georgia . . . . . *Marstonia halcyon* Thompson

Shell ovately conical, 0.66-0.73 times as wide as high, whorls rounded, not strongly shouldered, umbilicus narrow, suture not descending to the aperture in lateral profile (Figs. 216, 234). Flint river system, Georgia . . . . . *Marstonia castor* Thompson

35(32) Shell thick, solid, nearly opaque, umbilicus closed or narrowly rimate 36

Shell thinner, transparent or translucent, openly umbilicate 37

36(35) Shell ovately conical in shape, spire convex in outline, outer lip straight in lateral profile, sutures shallow, whorls not shouldered; verge with two small glands on the apical lobe and a small raised gland near the base of the verge (Figs. 221, 253). Creeks in Limestone County, Alabama . . . . . *Marstonia pachyta* Thompson

Shell nearly conical, spire straight-sided, outer lip strongly curved in lateral profile, whorls shouldered, suture deep; verge with a single large gland on the apical lobe (Figs. 215, 233). Tennessee River, Alabama . . . . . *Marstonia arga* Thompson



FIGS. 222-234. Shells of Hydrobiidae (Nymphophilinae). FIG. 222. *Cincinnatia helicogyra*. FIG. 223. *C. mica*. FIG. 224. *C. parva*. FIG. 225. *C. peracuta*. FIG. 226. *C. petrifons*. FIG. 227. *C. ponderosa*. FIG. 228. *C. wekiwae*. FIG. 229. *Fontelicella californiensis*. FIG. 230. *F. (Natricola) robusta*. FIG. 231. *F. (Microamnicola) micrococcus*. FIG. 232. *Marstonia agarhecta*. FIG. 233. *M. arga*. FIG. 234. *M. castor*. Measurement lines = 1 mm.

Reproduced from best available copy.

37(35) Shell sutures deep, whorls shouldered, outer lip arched slightly forward in lateral profile (Figs. 220, 252). Marion County, Tennessee . . . . . *Marstonia ogmorphaphe* Thompson

Shell sutures shallow, whorls not shouldered, outer lip straight in lateral profile 38

38(37) Northern in distribution: southern Canada, Maine west to Minnesota and Iowa (Figs. 218, 219, 245, 246, 250, 251) . . . . . *Marstonia lustrica* (Pilsbry)

Southern: Madison County, Alabama (Fig. 247) . . . . . *Marstonia olivacea* (Pilsbry)

39(30) Shell subglobose or broadly ovate, imperforate. Alabama, Florida and Georgia. Genus *Notogillia* 40

Shell conic or ovate, but if subglobose or broadly ovate then it is umbilicate 41

40(39) Shell subglobose, relatively small (that of adults is 4.0-4.5 mm in length), periostracum greyish white (Figs. 90, 254). Southcentral Georgia . . . . . *Notogillia sathon* Thompson

Shell broadly ovate, larger (that of adults is 4.5-7.5 mm in length), periostracum olivaceous-brown (Figs. 255, 260). Alabama, Florida and Georgia . . . . . *Notogillia wetherbyi* (Dall)

41(39) Distribution east of the Continental Divide 42

Distribution west of the Continental Divide 48

42(41) Penis relatively large, spatulate, and having a long narrow gland running along each margin from the base to near its tip. Georgia and Florida. Genus *Spilochlamys* 43

Penis small, slender, conical 45

43(42) Shell subglobose, spire depressed (Fig. 259). Tributaries of the Ocmulgee River, Georgia . . . . . *Spilochlamys turgida* Thompson

Shell ovate, spire prominent. Florida 44

44(43) Shell solid, thick (Figs. 275, 276); apex of the accessory lobe of the verge without a terminal glandular crest (Fig. 264). St. Johns river drainage, Florida . . . . . *Spilochlamys gravis* Thompson

Shell thin or only moderately thick (Fig. 258); apex of the accessory lobe of the verge with an apical glandular crest (Figs. 92, 263). Gulf of Mexico drainage in northcentral Florida . . . . . *Spilochlamys conica* Thompson



FIGS. 235-241. Shells of Hydrobiidae (Nymphophilinae). FIG. 235. *Cincinnatia augustina* = *C. floridana*. FIG. 236. *C. vanhyningi*. FIG. 237. *Fontelicella deserta*. FIG. 238. *F. intermedia*. FIG. 239. *F. neomexicana*. FIG. 240. *F. (Natricola) hendersoni*. FIG. 241. *F. (N.) idahoensis*. Measurement lines = 1 mm. Fig. 235 is from Walker (1906a); Fig. 236 is from Vanatta (1934); Figs. 237 and 239-241 are from Pilsbry (1916a); Fig. 238 is from Tryon (1865i).



FIGS. 242-248. Shells of Hydrobiidae (Nymphophilinae). FIG. 242. *Fontelicella (Natricola) idahoensis*. FIG. 243. *F. (N.) robusta*. FIG. 244. *F. (Microamnicola) micrococcus*. FIG. 245. *Marstonia winkleyi mozleyi* = *M. lustrica*, female. FIG. 246. *M. winkleyi mozleyi* = *M. lustrica*, male. FIG. 247. *M. olivacea*. FIG. 248. *Orygoceras* sp., dorsal and lateral views. Measurement lines = 1 mm. Figs. 242 and 243 are from Pilsbry (1933); Fig. 244 is from Stearns (1893); Figs. 245 and 246 are from Walker (1925a); Fig. 247 is from Thompson (1977); Fig. 248 is after Taylor (1974).



FIGS. 249-259. Shells of Hydrobiidae (Nymphophilinae). FIG. 249. *Marstonia halcyon*. FIG. 250. *M. lustrica*. FIG. 251. *M. lustrica*. FIG. 252. *M. ogmorphaphe*. FIG. 253. *M. pachyta*. FIG. 254. *Notogillia sathon*. FIG. 255. *N. wetherbyi*. FIG. 256. *Pyrgulopsis nevadensis*. FIG. 257. *Rhapinema dacryon*. FIG. 258. *Spilochlamys conica*. FIG. 259. *S. turgida*. Measurement lines = 1 mm.



FIGS. 260-269. Verges of hydrobiid snails (Nymphophilinae and Amnicolinae). FIG. 260. *Notogillia wetherbyi*. FIG. 261. *Pyrgulopsis letsoni*. FIG. 262. *Rhapinema dacryon*. FIG. 263. *Spilochlamys conica*. FIG. 264. *Sp. gravis*. FIG. 265. *Stiobia nana*. FIG. 266. *Amnicola dalli dalli*. FIG. 267. *A. dalli johnsoni*. FIG. 268. *A. limosa*. FIG. 269. *A. rhombostoma*. Measurement lines = 1 mm. Figs. 260, 262-267 and 269 are from Thompson (1968, 1969) and Thompson & McCaleb (1978); Figs. 261 and 268 are from E.G. Berry (1943).



FIGS. 270-278. Shells of Hydrobiidae (Nymphophilinae and Amnicolinae). FIG. 270. *Pyrgulopsis nevadensis nevadensis*. FIG. 271. *P. nevadensis nevadensis*. FIG. 272. *P. nevadensis nevadensis*. FIG. 273. *P. scalariformis*. FIG. 274. *P. archimedis*. FIG. 275. *Spilochlamys gravis*. FIG. 276. *S. gravis*. FIG. 277. *Amnicola aldrichi aldrichi*. FIG. 278. *A. dalli johnsoni*. Measurement lines = 1 mm. Fig. 270 is from Stearns (1883b); Fig. 271 is from Call & Pilsbry (1886); Figs. 272 and 274 are from S.S. Berry (1947); Fig. 273 is from Wolf (1869); Figs. 275, 276 and 278 are from Thompson (1968); Fig. 277 is from Call & Beecher (1886).

45(42) Shell elongately conical. Genus *Pyrgulopsis*, in part. Widely distributed 46

Shell broadly conical, globosely conical or ovate. Widely distributed (Figs. 189, 190, 199, 200, 203-213, 222-228, 235, 236) . . . . . Genus *Cincinnatia*[71]

46(45) Shell umbilicate (Fig. 261). Ontario and Michigan to New York . . . . . *Pyrgulopsis letsoni* (Walker)

Shell imperforate. Alabama and Arkansas 47

47(46) Whorls flat-sided, periphery angular or carinate (Fig. 273). Alabama . . . . . *Pyrgulopsis scalariformis* (Wolf)

Whorls rounded, periphery rounded. Arkansas . . . . . *Pyrgulopsis ozarkensis* Hinkley

48(41) Shell elongately conical, whorls wholly or nearly flat-sided, or concave, usually angulate or carinate. Genus *Pyrgulopsis*, in part 49

Shell conical, narrowly ovate to globosely conic, whorls rounded, not angulate or carinate. Genus *Fontelicella* 50

49(48) Periphery of body whorl concave (Fig. 274). Upper Klamath Lake, Oregon . . . . . *Pyrgulopsis archimedis* S. S. Berry

Periphery of body whorl flat-sided (Figs. 256, 270-272). Pyramid and Walker's lakes, Nevada . . . . . *Pyrgulopsis nevadensis* (Stearns)

50(48) Shell conical or narrowly ovate 51

Shell globosely conic, minute (that of adults is less than 2 mm in length) (Figs. 231, 244). Subgenus *Microamnicola*. California and Nevada . . . . . *Fontelicella (Microamnicola) micrococcus* Pilsbry (in Stearns) 1893

51(50) Shell relatively small (that of adults is 5 mm or less in length); the terminal lobe of the verge is usually a little longer than the penis (Figs. 229, 237-239). California, Idaho, New Mexico, Oregon and Utah . . . . . Subgenus *Fontelicella* s.s.[71]

Shell relatively large (that of adults is up to 8 mm in length); terminal lobe of the verge is about twice as long as the penis (Figs. 230, 240-243). Idaho, Oregon and Wyoming . . . . . Subgenus *Natricola*[71]

52(1) Males with two-ducted verges (Fig. 82d). Subfamily Amnicolinae 53

Males with three-ducted verges (Fig. 82e). Subfamily Fontigentinae (Figs. 283, 310-315, 319). Widely distributed in eastern North America . . . Genus *Fontigens*[71]



FIGS. 279-283. Verges of hydrobiid snails (Amnicolinae and Fontigentinae). FIG. 279. *Amnicola (Lyogyrus) grana*. FIG. 280. *A. (L.) pupoidea*. FIG. 281. *A. (L.) retromargo*. FIG. 282. *A. (L.) walkeri*. FIG. 283. *Fontigens nickliniana*. Measurement lines = 1 mm. Figs. 279-281 are from Thompson (1968); Figs. 282 and 283 are from E.G. Berry (1943).



FIGS. 284-287. Shells of Hydrobiidae (Amnicolinae). FIG. 284. *Amnicola dalli johnsoni*. FIG. 285. *A. cora*. FIG. 286. *A. limosa*. FIG. 287. *A. decisa*. Measurement lines = 1 mm. Fig. 284 is from Thompson (1968); Fig. 285 is from Hubricht (1979); Figs. 286 and 287 are from Haldeman (1845).



FIGS. 288-296. Shells of Hydrobiidae (Amnicolinae). FIG. 288. *Amnicola pallida* = *A. limosa*. FIG. 289. *A. porata* = *A. limosa*. FIG. 290. *A. porata* = *A. limosa*. FIG. 291. *A. stygia*. FIG. 292. *A. (Lyogyrus) greggi*. FIG. 293. *A. (L.) pilsbryi*. FIG. 294. *A. (L.) pilsbryi*. FIG. 295. *A. (L.) pilsbryi*. FIG. 296. *A. (L.) pilsbryi*. Measurement lines = 1 mm. Figs. 288 and 289 are from Haldeman (1845); Fig. 291 is from Hubricht (1971); Fig. 292 is from Pilsbry (1935a); Figs. 293 and 294 are from Walker (1906a).



FIGS. 297-308. Shells of Hydrobiidae (Nymphophilinae and Amnicolinae). FIG. 297. *Stiobia nana*. FIG. 298. *Amnicola dalli dalli*. FIG. 299. *A. limosa limosa*. FIG. 300. *A. rhombostoma*. FIG. 301. *A. (Lyogyrus) browni*. FIG. 302. *A. (L.) grana*. FIG. 303. *A. (L.) greggi*. FIG. 304. *A. (L.) pilsbryi*. FIG. 305. *A. (L.) pupoidea*. FIG. 306. *A. (L.) retromargo*. FIG. 307. *A. (L.) walkeri*. FIG. 308. *Hauffenia micra*. Measurement lines = 1 mm.

Reproduced from best available copy.



FIGS. 309-318. Shells of Hydrobiidae (Nymphophilinae and Amnicolinae). FIG. 309. *Amnicola (Lyogyrus) walkeri*. FIG. 310. *Fontigens turritella*. FIG. 311. *F. holsingeri*. FIG. 312. *F. orolibas*. FIG. 313. *F. orolibas*. FIG. 314. *F. tartarea*. FIG. 315. *F. cryptica*. FIG. 316. *Horatia nugax*. FIG. 317. *"Cochliopa" texana*. FIG. 318. *"Paludestrina" bottimeri*. Measurement lines = ½ mm. Fig. 309 is from Walker (1906a); Figs. 310-315 are from or after Hubricht (1957, 1963b, 1976); Fig. 316 is after Pilsbry & Ferriss (1906); Fig. 317 is from Pilsbry (1935a); Fig. 318 is from Walker (1925a).

53(52) Shell ovate or turbinate to globosely conic. Widely distributed. Genus *Amnicola* 54

Shell discoidal or subdiscoidal. Texas (? also Alabama) 55

54(53) Nuclear whorl of shell relatively large (0.38-0.48 mm in diameter); mantle heavily mottled with black; penis and flagellum relatively stout (Figs. 93, 266-269, 277, 278, 284-291, 298-300). Widely distributed in eastern North America . . . . . Subgenus *Amnicola* s.s.[72]

Nuclear whorl of shell small (0.29-0.36 mm in diameter); mantle diffusely shaded with pigment; penis and flagellum relatively slender and elongate (Figs. 94, 279-282, 292-296, 301-309). Widely distributed in North America . . . . . Subgenus *Lyogyrus*[72]

55(53) Shell discoidal, spire hardly raised above the body whorl (Fig. 308). Texas . . . . . *Hauffenia micra* (Pilsbry & Ferriss)[73]

Shell subdiscoidal, spire noticeably raised above the body whorl (Fig. 316). Texas . . . . . *Horatia nugax* (Pilsbry & Ferriss)[73]

FAMILY POMATIOPSIDAE

The Pomatiopsidae are represented in North America by six species, three in the east and three in California. Their general appearance is that of a hydrobioid, and in the past they frequently have been included in the Hydrobiidae as a subfamily (see Davis, 1967, for a review of familial classification). For the most recent diagnoses of the families Pomatiopsidae and Hydrobiidae, see Davis (1979).

Because of their obvious close systematic relationship to the medically important Oriental genus *Oncomelania*, North American *Pomatiopsis*, especially *P. cincinnatiensis* (Lea) and *P. lapidaria* (Say), have received considerable attention.

The genus *Pomatiopsis* comprises a group of amphibious species which inhabit river banks or moist areas near streams. In contrast, the hydrobiids live *in* the water of springs, streams, pools and lakes.

Identification Key for the Pomatiopsidae

1 Eastern in distribution 2

Restricted to California 4

2(1) Shell elongate, with relatively flattened whorls and oval aperture 3

Shell more depressed, broadly conical, with rounded whorls and aperture (Fig. 323). Tennessee and southwestern Virginia to southern Michigan, Illinois and Iowa . . . . . *Pomatiopsis cincinnatiensis* (Lea)



FIGS. 319-332. Shells of Hydrobiidae (Figs. 319, 320), Pomatiopsidae (Figs. 321-325), Thiaridae (Figs. 326, 327) and Pleuroceridae (Figs. 328-332). FIG. 319. *Fontigens nickliniana*. FIG. 320. *"Bythinella" hemphilli*. FIG. 321. *Pomatiopsis binneyi*. FIG. 322. *P. californica*. FIG. 323. *P. cincinnatiensis*. FIG. 324. *P. hinkleyi*. FIG. 325. *P. lapidaria*. FIG. 326. *Thiara granifera*. FIG. 327. *Melanoides tuberculata*. FIG. 328. *Elimia boykiniana boykiniana*. FIG. 329. *E. boykiniana viennaensis*. FIG. 330. *E. clenchi*. FIG. 331. *E. carinifera*. FIG. 332. *E. arachnoidea arachnoidea*. Measurement lines = 1 mm or are divided into millimeters.

3(2) Spire more acute, body whorl proportionately smaller, aperture broadly oval, umbilicus wider, more open (Fig. 325). Widely distributed in the eastern United States, with occasional occurrences west to northern Texas and New Mexico . . . . . *Pomatiopsis lapidaria* (Say)

Spire more obtuse, body whorl proportionately larger, aperture narrowly oval, umbilicus nearly closed (Fig. 324). Found in several localities in Alabama, South Carolina and Tennessee . . . . . *Pomatiopsis hinkleyi* Pilsbry[20]

4(1) Shell quite small, that of adults with four to five whorls about 3 mm in length, light horn in color, imperforate (Fig. 321). Marin County, California . . . . . *Pomatiopsis binneyi* Tryon

Shell larger, that of adults more than 4 mm in length, brownish-olive or chestnut brown in color 5

5(4) Shell chestnut brown in color (Fig. 322). San Francisco area . . . . . *Pomatiopsis californica* Pilsbry

Shell brownish-olive in color. Northeastern California . . . . . *Pomatiopsis chacei* Pilsbry

FAMILY THIARIDAE

The Thiaridae and the Pleuroceridae contain various genera with very similar shells, and because of this they were long considered to all belong to one and the same family, traditionally called the Melaniidae. The latter name is based on the genus *Melania* Lamarck 1799, a synonym of *Thiara* Röding 1798. Morrison (1954) used biological characters to separate the various melanoid/cerithioid families, and separated the Thiaridae and the Pleuroceridae as follows:

Thiaridae: Reproduction parthenogenetic, without males; brood pouch not uterine, but adventitious (subhaemocoelic) in the neck region, with opening on right side of neck.

Pleuroceridae: Reproduction dioecious, with males present; females with egg-laying sinus on right side of foot; lays numerous eggs of small size.

A feature distinguishing *Thiara* and *Melanoides* from the pleurocerids is their mantle edge, which in the thiarids has a number of fleshy protuberances or papillae. The mantle edge of the Pleuroceridae is smooth.

Identification Key for the Thiaridae

1 Shells with rounded whorls which are sculptured with spiral threads and grooves, and transverse lines which commonly develop into low costae; this type of sculpture sometimes produces a reticulate or nodular pattern where the spiral and transverse elements intersect (Fig. 327). Florida, Texas and Arizona . . . . . *Melanoides tuberculata* (Müller)

Shell with flattened whorls, especially those of the spire; sculpturing of spiral rows of beads and nodules which are generally aligned in transverse rows (Fig. 326). Florida and Texas ..... *Thiara granifera* (Lamarck)

FAMILY PLEUROCERIDAE

The Pleuroceridae are widely distributed, occurring not only widely in North America, but in Central and South America, Africa and Asia as well. But, it is in North America that the family has reached its greatest development. Morrison (1954) has characterized the family as being dioecious, with the females having an egg-laying sinus on the right side of the foot. The types of eggs vary between some of the species, and attempts have been made to use egg-mass characteristics in generic taxonomy (see Dazo, 1965, for review). Unfortunately, egg-mass characters have been described for very few species. The generic groups traditionally have been distinguished on shell characters, and the classification of these groups as based on shells is not entirely satisfactory. Nevertheless, shell characters are useful in recognizing the genera and are essential for species identification.

As presented in this manual, the Pleuroceridae comprise seven nominal generic groups, several of which have subgroups. Many of the species within these groups exhibit considerable variation in shell characters, and in some cases this variation seems to be clinal. In other cases it may be ecological. *Io* is the only genus in which geographic variation has been carefully investigated, in a remarkable study by C. C. Adams (1915), which did much to clarify systematics within the genus.

The shells of pleurocerids are thick and solid and vary in shape from elongately conical to subglobose. The aperture is frequently entire and in many species it is canaliculated anteriorly. The operculum is paucispiral and corneous.

Identification Key for the Pleuroceridae

1 Shell large, fusiform, periphery of whorls angulated or inflated, periphery commonly with elongated spines (although some forms are smooth); anterior end or "base" of aperture prolonged into a long canal (Figs. 429, 430, 461-465). Tennessee River and several of its main tributaries in western Virginia and eastern Tennessee ..... *Io fluvialis* (Say)[30]

Shell large to small, conical to subglobose*, surface smooth or sculptured, with or without short spines, nodules, lirae, carina and costae; anterior end or "base" of aperture without a long canal (a short canal may be present or the canal may be absent altogether) 2

2(1) Terminal whorl with a posterior slit along the sutural juncture. Coosa River, Alabama. Genus *Gyrotoma*[28, 29] 3

Terminal whorl without a posterior slit along the sutural juncture 8

---

*Shell shape refers to undecollated shells.

3(2) Shell sculptured with numerous and closely spaced lirae, nine or more on the body whorls of adults 4

Shell relatively smooth or sculptured with eight or less lirae on the body whorls of adults 5

4(3) Lirae fine and numerous, 20 or more on the body whorl; color bands 8-10 (Fig. 441). Coosa River in Shelby and Talladega counties, Alabama . . . . . *Gyrotoma lewisi* (Lea)

Lirae coarser and less numerous, 9-12 on the body whorl; color bands seven or less (Figs. 444, 445). Coosa River, from Fort William Shoals to Wetumpka, Alabama . . . . . *Gyrotoma pumilum* (Lea)

5(3) Spire with a single, very accentuated lira (sometimes a second lower lira is present) on the spire whorls, giving the shell a pagoda-like appearance (Figs. 442, 443). Coosa River, from The Bar to Wetumpka, Alabama . . . . . *Gyrotoma pagodum* (Lea)

Spire not pagoda-like 6

6(5) Whorls flattened, tapering and lumpy, giving the shell a pyramidal shape (Fig. 446). Coosa River in Shelby and St. Clair counties, Alabama . . . . . *Gyrotoma pyramidatum* Shuttleworth

Whorls not both flattened and tapering, or if so, not lumpy 7

7(6) Small, decollated adult shells rarely over 16 mm long; sutural fissure very shallow (Fig. 447). Coosa River in Coosa and Shelby counties, Alabama . . . . . *Gyrotoma walkeri* Smith

Larger, decollated adult shells usually more than 20 mm long; sutural fissure moderate to deep, not exceedingly shallow (Figs. 431-440). Coosa River in Chilton, Coosa, Elmore, Shelby, St. Clair and Talladega counties, Alabama . . . . . *Gyrotoma excisum* (Lea)

8(2) Lateral radular teeth with broad, bluntly rounded or cleaver-like median cusps; shell medium to small, subglobose, globosely or broadly conic, or ovate. Genus *Leptoxis* 9

Lateral radular teeth with narrow, pointed, spade-shaped or triangular median cusps; shell large to small, generally elongately or narrowly conic, but several species are broadly conic, ovate or cylindrical 34

9(8) Shell with an elongated or short spire, body whorl generally tapering and usually without prominent surface sculpture, although several species have spiral striae, carinae or small shoulder nodules; aperture broadly ovate, its anterior end nearly always rounded 10

Shell with a very short spire and a nearly cylindrical body whorl with relatively large bumps or nodules on the shoulders; aperture pyriform, its anterior end pointed (Figs. 501, 502). Tennessee River and tributaries in Alabama and Tennessee. Subgenus *Athearnia* . . . . . *Leptoxis (Athearnia) crassa* (Haldeman)[34]

10(9) Shell generally thick and solid. Ohio and Alabama river drainages. Subgenus *Leptoxis* s.s.[32, 74] 11

Shell commonly relatively thin. Ohio river and Atlantic drainages and White River, Arkansas. Subgenus *Mudalia*[75] 28

11(10) Ohio river drainage, including the Tennessee, Cumberland, Duck and Elk river drainages 12

Alabama river drainage 13

12(11) Base of adult shell without an umbilicus (Figs. 478-482). Cumberland, Duck, Ohio and Tennessee rivers and some of their drainages . . . . . *Leptoxis praerosa* (Say)

Base of adult shell with an umbilicus (Fig. 528). Elk, Red and Stone's rivers, Tennessee, and in Ringgold Creek of the Cumberland River . . . . . *Leptoxis umbilicata* (Wetherby)

13(11) Species inhabiting the Alabama River proper and very short distances up the Cahaba or Coosa rivers from their mouths 14

Species confined to tributaries of the Alabama River 15

14(13) Operculum ovate, loosely paucispiral (Fig. 476). Alabama and Coosa rivers, Alabama . . . . . *Leptoxis picta* (Conrad)

Operculum elongate, tightly paucispiral (Figs. 484-486). Alabama and Cahaba rivers and the Coosa River and tributaries . . . . . *Leptoxis taeniata* (Conrad)[76]

15(13) Species confined to the Coosa River and its tributaries 16

Species confined to the Cahaba and Black Warrior rivers and their tributaries 25

16(15) Shell strongly lirate 17

Shell smooth to spirally striate or weakly lirate, but not strongly lirate 19

17(16) Carinae may be well developed, but not highly accentuated (Figs. 484-486). Alabama and Cahaba rivers and the Coosa River and tributaries . . . . . *Leptoxis taeniata* (Conrad)[76]

Carinae high, accentuated 18

18(17) Shell relatively large (that of adults 15-22 mm in length), spire rather depressed, body whorl and aperture wide (Fig. 483). Coosa River, Alabama . . . . . *Leptoxis showalteri* (Lea)

Shell relatively small (that of adults 10-13 mm in length), high-spired, body whorl and aperture narrow. Coosa River, Alabama . . . . . *Leptoxis lirata* (Smith)[33]

19(16) Shell relatively large (that of adults more than 13 mm in length) 20

Shell relatively small (that of adults less than 12 mm in length) 24

20(19) Margin of operculum relatively smooth, without regular serrations 21

Margin of operculum serrated regularly either on the right or at the anterior ("base") 23

21(20) Operculum tightly paucispiral (Figs. 484-486). Alabama and Cahaba rivers and the Coosa River and tributaries . . . . . *Leptoxis taeniata* (Conrad)[76]

Operculum loosely paucispiral 22

22(21) Shell surface with widely spaced spiral striae (incised lines). Coosa River in Alabama and Georgia, and in Terrapin Creek, Cherokee County, Alabama . . . . . *Leptoxis formosa* (Lea)[77]

Shell surface smooth (Fig. 468). Coosa River, Alabama . . . . . *Leptoxis clipeata* (Smith)

23(20) Right margin of operculum serrated regularly, anterior or "basal" margin smooth (Figs. 471, 472). Coosa River, Alabama . . . . . *Leptoxis foremani* (Lea)

Right margin of operculum smooth, anterior or "basal" margin serrated regularly (Fig. 473). Coosa River, Alabama . . . . . *Leptoxis ligata* (Anthony)

24(19) Shells of adults 8 mm or less in length, with a noticeable spire (Fig. 487). Coosa River, Alabama . . . . . *Leptoxis vittata* (Lea)

Shells of adults 10 mm or more in length, spire greatly depressed (Fig. 475). Coosa River, Alabama . . . . . *Leptoxis occultata* (Smith)

25(15) Species confined to the Cahaba River 26

Species confined to the Black Warrior River 27

26(25) Shell with depressed spire and subglobose body whorl (Figs. 456, 457). Cahaba River, Alabama, and tributaries ..... *Leptoxis ampla* (Anthony)

Shell with elevated spire and elongated body whorl (Figs. 469, 470). Cahaba River and Buck Creek, Alabama ..... *Leptoxis compacta* (Anthony)

27(25) Shell ovate, relatively large (that of adults more than 13 mm in length) (Fig. 477). Black Warrior River and Valley Creek, Alabama ..... *Leptoxis plicata* (Conrad)

Shell broadly conic, relatively small (that of adults less than 13 mm in length) (Fig. 474). Black Warrior River, Alabama ..... *Leptoxis melanoides* (Conrad)

28(10) In streams of the Atlantic drainage 29

In streams of the Mississippi river drainage 30

29(28) Shell of adults medium, 13 or more mm in length, commonly with one or several carinae (Figs. 489-492). New York to North Carolina .... *Leptoxis (Mudalia) carinata carinata* (Bruguière)

Shells of adults small, about 10 mm in length, elongately conic, without carinae (Fig. 493). Hot Springs, Bath County, Virginia .... *Leptoxis (Mudalia) carinata nickliniata* (Lea)

30(28) In streams of the Ohio river drainage 31

In the White River, Arkansas, and its North Fork, in Missouri; shell typically covered with thick whitish calcium deposits (Fig. 488) .... *Leptoxis (Mudalia) arkansensis* (Hinkley)

31(30) Shell small (that of adults 8 mm or less in length), periphery with a single angulation or carina (Fig. 495). Tennessee River at Muscle Shoals, Alabama ..... *Leptoxis (Mudalia) minor* (Hinkley)

Shell medium in size (that of adults 10 mm or more in length), periphery smooth or with one, two or three angulations or carinae 32

32(31) Shell relatively large (that of adults 15 mm or more in length), high-spired, ovately conic, nearly always without color bands and carinae (Fig. 494). Kanawha River and tributaries, West Virginia ..... *Leptoxis (Mudalia) dilatata* (Conrad)

Shell smaller (that of adults 10-13 mm in length), ovately or globosely conic to subglobose, with or without color bands and carinae 33

33(32) Shell subglobose, generally with one to several carinae, usually without color bands (Figs. 496, 497). Ohio River in western Ohio and northern Kentucky and tributaries ..... *Leptoxis (Mudalia) trilineata* (Say)

Shell subglobose, without carinae, with spiral color bands (Figs. 498-500). Upper Tennessee River and tributaries . . . . . *Leptoxis (Mudalia) virgata* (Lea)

34(8) Shell medium (except for one large species, *Lithasia lima*), elongately conic, subglobose, ovate, or cylindrical, surface of most species sculptured with obtuse spines or prominent nodules (one species is smooth and several nodulate species have smooth forms); columellar margin of the aperture thickened, meeting the anterior or "basal" lip with a channel or strong angle (except for *L. obovata* and *L. geniculata pinguis*[35]); a calloused thickening usually occurs on the parietal wall at the posterior end of the aperture. Genus *Lithasia* 35

Shell large to small, narrowly or elongately conic, or cylindrical; surface smooth, carinate, lirate, costate, or occasionally with nodules; anterior or basal end of aperture either rounded and smooth or produced into a short canal; columellar margin of the aperture and posterior parietal wall without a thickening 36

35(34) The most prominent spiral row of nodules or tubercles is along the shoulder of the whorls (Figs. 503-513). Ohio and Tennessee rivers and their tributaries . . . . . Subgenus *Lithasia* s.s.[78]

The most prominent spiral row of nodules, tubercles or spines is along or near the median periphery of the whorls (Figs. 514-520). Ohio and Tennessee rivers and their tributaries; Black and Spring rivers, Arkansas; Big Black River, Mississippi . . . . . Subgenus *Angitrema*[78]

36(34) Anterior or "basal" end of aperture prolonged into a short canal, producing an auger-shaped base to the shell (Figs. 521-527, 529-563). Mississippi river and Great Lakes drainages, and through the Erie Canal into the basin of the Hudson River . . . . . Genus *Pleurocera*[78]

Anterior or "basal" end of aperture not channeled or auger-shaped 37

37(36) Eastern in distribution, east of the Continental Divide, occurring in drainages of the Mississippi River, the Gulf of Mexico, the Atlantic slope, the Great Lakes-St. Lawrence River or Hudson Bay (Figs. 323-428, 458-460) . . . . . Genus *Elimia*[78]

Western in distribution, west of the Continental Divide, occurring in the drainages of the Great Basin or the Pacific slope (Figs. 448-455, 466, 467) . . . . . Genus *Juga*[78]



FIGS. 333-344. Shells of Pleuroceridae. FIG. 333. *Elimia arachnoidea spinella*. FIG. 334. *E. capillaris*. FIG. 335. *E. catenaria catenaria*. FIG. 336. *E. catenaria dislocata*. FIG. 337. *E. catenaria postelli*. FIG. 338. *E. catenaria vanhyningiana*. FIG. 339. *E. comalensis comalensis*. FIG. 340. *E. crenatella*. FIG. 341. *E. edgariana*. FIG. 342. *E. impressa*. FIG. 343. *E. perstriata decampi*. FIG. 344. *E. striatula*. Measurement lines are divided into millimeters.



FIGS. 345-354. Shells of Pleuroceridae. FIG. 345. *Elimia bellacrenata*. FIG. 346. *E. ucheensis*. FIG. 347. *E. brevis*. FIG. 348. *E. brevis*. FIG. 349. *E. catenaria dislocata*. FIG. 350. *E. comalensis fontinalis*. FIG. 351. *E. fusiformis*. FIG. 352. *E. fusiformis*. FIG. 353. *E. nassula*. FIG. 354. *E. pupaeformis*. Measurement lines are divided into millimeters. Figs. 345-347, 349, 351 and 353 are from Tryon (1865-66, 1873b); Figs. 348, 352 and 354 are from Goodrich (1936); Fig. 350 is from Pilsbry & Ferriss (1906).



FIGS. 355-367. Shells of Pleuroceridae. FIG. 355. *Elimia strigosa*. FIG. 356. *E. teres*. FIG. 357. *E. troostiana*. FIG. 358. *E. curvicostata*. FIG. 359. *E. dickinsoni*. FIG. 360. *E. annettae*. FIG. 361. *E. gerhardti*. FIG. 362. *E. murrayensis* = *E. gerhardti*. FIG. 363. *E. hartmaniana*. FIG. 364. *E. clausa*. FIG. 365. *E. hydei*. FIG. 366. *E. laqueata laqueata*. FIG. 367. *E. laqueata laqueata*. Measurement lines are divided into millimeters.



FIGS. 368-378. Shells of Pleuroceridae. FIG. 368. *Elimia porrecta*. FIG. 369. *E. rubella* = ?*E. porrecta*. FIG. 370. *E. ebenum ebenum*. FIG. 371. *E. cahawbensis cahawbensis*. FIG. 372. *E. flava*. FIG. 373. *E. gerhardti*. FIG. 374. *E. macglameriana*. FIG. 375. *E. pygmaea*. FIG. 376. *E. alabamensis*. FIG. 377. *E. alabamensis*. FIG. 378. *E. haysiana*. Measurement lines are divided into millimeters. Figs. 368-373, 376 and 378 are from Tryon (1865-66); Figs. 374, 375 and 377 are from Goodrich (1936).



FIGS. 379-390. Shells of Pleuroceridae. FIG. 379. *Elimia haysiana*. FIG. 380. *E. pupoidea*. FIG. 381. *E. pupoidea*. FIG. 382. *E. laqueata castanea*. FIG. 383. *E. paupercula*. FIG. 384. *E. livescens haldemani*. FIG. 385. *E. pilsbryi*. FIG. 386. *E. showalteri*. FIG. 387. *E. variata*. FIG. 388. *E. aterina*. FIG. 389. *E. bullula*. FIG. 390. *E. bullula*. Measurement lines are divided into millimeters. Figs. 379, 380, 382-384, 386-389 are from Tryon (1865-66, 1873b); Figs. 381, 385 and 390 are from Goodrich (1936).



FIGS. 391-402. Shells of Pleuroceridae. FIG. 391. *Elimia mutata* = *E. laqueata laqueata*. FIG. 392. *E. laqueata costulata*. FIG. 393. *E. livescens livescens*. FIG. 394. *E. mutabilis mutabilis*. FIG. 395. *E. taitiana*. FIG. 396. *E. bellula*. FIG. 397. *E. chiltonensis*. FIG. 398. *E. cylindracea*. FIG. 399. *E. gibbera*. FIG. 400. *E. lachryma*. FIG. 401. *E. laeta*. FIG. 402. *E. olivula*. Measurement lines are divided into millimeters.



FIGS. 403-414. Shells of Pleuroceridae. FIG. 403. *Elimia clavaeformis*. FIG. 404. *E. acutocarinata* = ? *E. clavaeformis*. FIG. 405. *E. acutocarinata* = ? *E. clavaeformis*. FIG. 406. *E. simplex*. FIG. 407. *E. caelatura caelatura*. FIG. 408. *E. caelatura excellens*. FIG. 409. *E. caelatura georgiana*. FIG. 410. *E. caelatura luteocella*. FIG. 411. *E. caelatura stearnsiana*. FIG. 412. *E. proxima*. FIG. 413. *E. virginica*. FIG. 414. *E. clara*. Measurement lines are divided into millimeters.



FIGS. 415-425. Shells of Pleuroceridae. FIG. 415. *Elimia caelatura infuscata*. FIG. 416. *E. caelatura stearnsiana*. FIG. 417. *E. fascinans*. FIG. 418. *E. jonesi*. FIG. 419. *E. vanuxemiana*. FIG. 420. *E. vanuxemiana*. FIG. 421. *E. vanuxemiana*. FIG. 422. *E. vanuxemiana*. FIG. 423. *E. symmetrica*. FIG. 424. *E. symmetrica*. FIG. 425. *E. symmetrica*. Measurement lines are divided into millimeters. Figs. 415, 417, 419, 423-425 are from Tryon (1865-66, 1873b); Fig. 416 is from Call (1886c); Figs. 418 and 420-422 are from Goodrich (1936).



FIGS. 426-434. Shells of Pleuroceridae. FIG. 426. *Elimia ampla*. FIG. 427. *E. potosiensis potosiensis*. FIG. 428. *E. interrupta*. FIG. 429. *Io fluvialis*. FIG. 430. *I. fluvialis* form *angitremoides*. FIG. 431. *Gyrotoma excisum*, apertural and right lateral views. FIG. 432. *G. excisum*. FIG. 433. *G. excisum*, operculum. FIG. 434. *G. excisum*, right lateral view of the shell in Fig. 432. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 435-440. Shells of Pleuroceridae, apertural and right lateral views. FIG. 435. *Gyrotoma alabamensis* = ? *G. excisum*. FIG. 436. *G. amplum* = ? *G. excisum*. FIG. 437. *G. cariniferum* = ? *G. excisum*. FIG. 438. *G. incisum* = ? *G. excisum*. FIG. 439. *G. laciniatum* = ? *G. excisum*. FIG. 440. *G. spillmani* = ? *G. excisum*. Measurement lines are divided into millimeters.



FIGS. 441-446. Shells of Pleuroceridae, apertural and right lateral views. FIG. 441. *Gyrotoma lewisi*. FIG. 442. *G. pagodum*. FIG. 443. *G. pagodum*. FIG. 444. *G. pumilum*. FIG. 445. *G. hendersoni* = ? *G. pumilum*. FIG. 446. *G. pyramidatum*. Measurement lines are divided into millimeters.



FIGS. 447-457. Shells of Pleuroceridae. FIG. 447. *Gyrotoma walkeri*, apertural and right lateral views. FIG. 448. *Juga plicifera*. FIG. 449. *J. silicula*. FIG. 450. *J. (Calibasis) acutifilosa*. FIG. 451. *J. (C.) occata*. FIG. 452. *J. (Oreobasis) bulbosa*. FIG. 453. *J. (O.) nigrina*. FIG. 454. *J. hemphilli hemphilli*. FIG. 455. *J. hemphilli dallesensis*. FIG. 456. *Leptoxis ampla*. FIG. 457. *L. ampla*. Measurement lines are divided into millimeters.



FIGS. 458-467. Shells of Pleuroceridae. FIG. 458. *Elimia potosiensis ozarkensis*. FIG. 459. *E. potosiensis plebius*. FIG. 460. *E. potosiensis plebius*. FIG. 461. *Io fluvialis* form *turrita*. FIG. 462. *I. fluvialis* form *verrucosa*. FIG. 463. *I. fluvialis* form *recta*. FIG. 464. *I. fluvialis* form *brevis*. FIG. 465. *I. fluvialis* form *spinosa*. FIG. 466. *Juga (Oreobasis) interioris*. FIG. 467. *J. (O.) laurae*. Measurement lines are divided into millimeters. Fig. 458 is from Call (1886b); Figs. 459-465 are from Tryon (1873b); Figs. 466 and 467 are from Goodrich (1944a).



FIGS. 468-477. Shells of Pleuroceridae. FIG. 468. *Leptoxis clipeata*. FIG. 469. *L. compacta*. FIG. 470. *L. compacta*. FIG. 471. *L. foremani*. FIG. 472. *L. downiei* = *L. foremani*. FIG. 473. *L. ligata*. FIG. 474. *L. melanoides*. FIG. 475. *L. occultata*. FIG. 476. *L. picta*. FIG. 477. *L. plicata*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 478-487. Shells of Pleuroceridae. FIG. 478. *Leptoxis praerosa*. FIG. 479. *L. praerosa*. FIG. 480. *L. subglobosa* = *L. praerosa*. FIG. 481. *L. gibbosa* = *L. praerosa*. FIG. 482. *L. tintinabulum* = *L. praerosa*. FIG. 483. *L. showalteri*. FIG. 484. *L. coosaensis* = *L. taeniata*. FIG. 485. *L. brevispira* = *L. taeniata*. FIG. 486. *L. taeniata*. FIG. 487. *L. vittata*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 488-497. Shells of Pleuroceridae. FIG. 488. *Leptoxis (Mudalia) arkansensis*. FIG. 489. *L. (M.) carinata carinata*. FIG. 490. *L. (M.) carinata carinata*. FIG. 491. *L. (M.) carinata carinata*. FIG. 492. *L. (M.) corpulenta* = *L. (M.) carinata*. FIG. 493. *L. (M.) carinata nickliniata*. FIG. 494. *L. (M.) dilatata*. FIG. 495. *L. (M.) minor*. FIG. 496. *L. (M.) trilineata*. FIG. 497. *L. (M.) trilineata*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 498-506. Shells of Pleuroceridae. FIG. 498. *Leptoxis (Mudalia) virgata*. FIG. 499. *Le. (M.) virgata*. FIG. 500. *Le. (M.) virgata*. FIG. 501. *Le. (Athearnia) crassa crassa*. FIG. 502. *Le. (A.) crassa anthonyi*. FIG. 503. *Lithasia geniculata geniculata*. FIG. 504. *Li. geniculata geniculata*. FIG. 505. *Li. geniculata fuliginosa*. FIG. 506. *Li. geniculata pinguis*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 507-516. Shells of Pleuroceridae. FIG. 507. *Lithasia obovata*. FIG. 508. *L. obovata* form *depygis*. FIG. 509. *L. obovata* form *pennsylvanica*. FIG. 510. *L. obovata* form *sordida*. FIG. 511. *L. salebrosa salebrosa*. FIG. 512. *L. salebrosa florentiana*. FIG. 513. *L. salebrosa subglobosa*. FIG. 514. *L. (Angitrema) armigera*. FIG. 515. *L. (A.) curta*. FIG. 516. *L. (A.) duttoniana*. Measurement lines are divided into millimeters.



FIGS. 517-527. Shells of Pleuroceridae. FIG. 517. *Lithasia (Angitrema) jayana*. FIG. 518. *L. (A.) lima*. FIG. 519. *L. (A.) lima* form. FIG. 520. *L. (A.) verrucosa*. FIG. 521. *Pleurocera acuta acuta*. FIG. 522. *P. acuta lewisi*. FIG. 523. *P. alveare*. FIG. 524. *P. striatum* = *P. canaliculatum alabamense*. FIG. 525. *P. canaliculatum filum*. FIG. 526. *P. canaliculatum undulatum*. FIG. 527. *P. canaliculatum undulatum*. Measurement lines are divided into millimeters.



FIGS. 528-537. Shells of Pleuroceridae. FIG. 528. *Leptoxis umbilicata*. FIG. 529. *Pleurocera alveare*. FIG. 530. *P. canaliculatum excuratum*. FIG. 531. *P. canaliculatum filum*. FIG. 532. *P. canaliculatum moriforme*. FIG. 533. *P. canaliculatum undulatum*. FIG. 534. *P. ponderosum* = *P. canaliculatum undulatum*. FIG. 535. *P. moniliferum* = *P. nobile nodosa*. FIG. 536. *P. parvum*. FIG. 537. *P. modestum* = *P. parvum*. Measurement lines are divided into millimeters. Figs. 529-537 are from Tryon (1865-66).



FIGS. 538-549. Shells of Pleuroceridae. FIG. 538. *Pleurocera gradatum*. FIG. 539. *P. postelli*. FIG. 540. *P. pyrenellum*. FIG. 541. *P. planogyrum* = *P. pyrenellum*. FIG. 542. *P. unciale unciale*. FIG. 543. *P. esterbrooki* = *P. unciale*. FIG. 544. *P. annuliferum*. FIG. 545. *P. foremani*. FIG. 546. *P. prasinatum*. FIG. 547. *P. prasinatum*. FIG. 548. *P. showalteri*. FIG. 549. *P. vestitum*. Measurement lines are divided into millimeters. Figs. 538-549 are from Tryon (1865-66).



FIGS. 550-561. Shells of Pleuroceridae. FIG. 550. *Pleurocera nobile nobile*. FIG. 551. *P. brumbyi*. FIG. 552. *P. currierianum*. FIG. 553. *P. pyrenellum*. FIG. 554. *P. trochiformis*. FIG. 555. *P. unciale unciale*. FIG. 556. *P. viridulum*. FIG. 557. *P. unciale hastatum*. FIG. 558. *P. annuliferum*. FIG. 559. *P. vestitum*. FIG. 560. *P. (Strephobasis) corpulentum*. FIG. 561. *P. (S.) curtum curtum*. Measurement lines are divided into millimeters.



FIGS. 562-572. Shells of Pleuroceridae (Figs. 562, 563), Acroloxidae (Fig. 564) and Lymnaeidae (Lymnaeinae) (Figs. 565-572). FIG. 562. *Pleurocera (Strephobasis) curtum roanense*. FIG. 563. *P. (S.) walkeri*. FIG. 564. *Acroloxus coloradensis*. FIG. 565. *Acella haldemani*. FIG. 566. *Bulimnea megasoma*. FIG. 567. *Fossaria cyclostoma*. FIG. 568. *F. galbana*. FIG. 569. *F. humilis*. FIG. 570. *F. obrussa*. FIG. 571. *F. parva*. FIG. 572. *F. tazewelliana*. Measurement lines = 1 mm or are divided into millimeters.

FAMILY ACROLOXIDAE

The family Acroloxidae is mainly a Eurasian one of ancient lakes (Baikal and Ohrid), although one species, *Acroloxus lacustris* (Linnaeus), is the common, widespread pond and lake limpet of Europe. One species occurs in North America, *A. coloradensis* (Henderson), which has a spotty, probably relic, distribution. It is known from three localities in the Rocky Mountains, and from a few ponds and lakes in northern Quebec and eastern Ontario.

Acroloxus is peculiar for a freshwater limpet because its body has a *dextral* organization (Fig. 755a). The common freshwater limpets, members of the Ancylidae, are all sinistral (Fig. 755b). The consequences of this right- and left-handedness can be seen in the reduced and very simplified patelliform shells of the two families. In *Acroloxus* the apex is inclined to the *left*, and in the Ancylidae it is inclined to the *right*.

Acroloxus coloradensis has a small, depressed shell with a striate, projecting apex (Fig. 564). Shells which reach 5 mm in length are only about 1 mm high. The shell surface is covered with delicate radial striae and fine, regular growth lines.

FAMILY LYMNAEIDAE

The Lymnaeidae are world-wide in distribution, but their greatest diversity is found in the northern United States and central Canada. Their shells range in shape from the coiled, needle-like *Acella haldemani* (Binney) (Fig. 565) to the uncoiled, limpet-shaped *Lanx* (Figs. 578-580, 633, 634) and *Fisherola* (Fig. 632). Those with coiled shells are easily distinguished from the Physidae by their dextral shells (the lone exception in the Lymnaeidae is the sinistral *Pseudisidora producta* (Mighels), which is restricted to Hawaii). No lymnaeids have planispiral shells, which immediately distinguishes them from the North American Planorbidae. The patelliform Lancinae, which occur only in the Pacific drainage region, can be distinguished from the Ancylidae by their much larger size and by their anterior rather than posterior shell apex.

The tentacles of lymnaeids are broad, flat and triangular (see Fig. 4), rather than being long, thin and filamentous as in the Physidae, Planorbidae and Ancylidae. Also, in contrast to the three latter families, all Lymnaeidae lack a respiratory pseudobranch (see Fig. 4).

Identification Key for the Lymnaeidae

1 Shell cap-shaped (ancyliform, limpet-shaped), not coiled. Western North America, in stream systems draining into the Pacific Ocean. Subfamily Lancinae[44] 2

Shell coiled. Common throughout North America. Subfamily Lymnaeinae 4

2(1) Apex subcentral. Genus *Lanx* 3

Apex close to the anterior end (Fig. 632). Columbia river drainage . . . . . *Fisherola nuttalli* (Haldeman)

3(2) Entire shell or at least its apex elevated (Figs. 578-580, 633). Klamath and Sacramento rivers, California; Umpqua river system, Oregon . . . . . Subgenus *Lanx* s.s.[79]

Shell and apex depressed (Fig. 634). Subgenus *Walkerola*. Klamath system in basin of Klamath River, Oregon . . . . . *Lanx (Walkerola) klamathensis* Hannibal

4(1) Adult shell with large, globose body whorl, without spiral striations (Fig. 594). Widely distributed, but of spotty occurrence ..... *Radix auricularia* (Linnaeus)

Adult shell with narrow or globose body whorl, but if globose, the shell is well sculptured with microscopic spiral striations 5

5(4) Shell attenuate, very narrow, almost needle-like (Fig. 565). Southern Ontario; north central United States to Vermont ..... *Acella haldemani* (Binney)

Shell thicker, not especially narrow 6

6(5) Shell succiniform, i.e., thin and fragile, with a large, oval aperture and body whorl, and small spire; surface sculptured with microscopic, raised, spiral periostracal threads (Fig. 593). Eastern North America generally .... *Pseudosuccinea columella* (Say)

Shell not succiniform, aperture may or may not be large and oval, but if so, the shell is not thin and fragile and is not sculptured with microscopic, raised, spiral periostracal threads 7

7(6) Shell large, that of adults more than 35 mm in length 8

Shell smaller, that of adults less than 35 mm in length 13

8(7) Shell with a relatively narrow body whorl. Genus *Stagnicola*, in part[43, 79] 13

Shell with a wider, expanded, elongately oval to globose body whorl 9

9(8) Shell with a narrow, pointed spire. Genus *Lymnaea*[40] 10

Shell with a relatively wider spire 12

10(9) Shell rimate, i.e., with a narrowly open umbilicus partially covered by the flare of the columellar lip (Fig. 590). Alaska and northwestern Canada .... *Lymnaea atkaensis* Dall

Shell imperforate 11

11(10) Shell with a large, subglobose body whorl (Fig. 592). Lake Superior, northern Lake Huron, Wisconsin river and Winnipeg river drainages .... *Lymnaea stagnalis sanctaemariae* Walker

Shell with an ample but not broad and subglobose body whorl (Fig. 591). Throughout much of Canada; in the northern United States and south to Colorado in the Rocky Mountains ..... *Lymnaea stagnalis appressa* Say

12(9) Shell spire rather depressed, whorls shouldered (Fig. 621). Lakes in Maine .... *Stagnicola mighelsi* (Binney)

Shell spire more elongated, whorls not shouldered (Fig. 566). Great Lakes and St. Lawrence river drainage area and parts of the Canadian Interior Basin ..... *Bulimnea megasoma* (Say)

13(7, 8) Adult shell medium to large, generally more than 13 mm (but occasionally 13 mm or less) in length; surface sculptured with microscopic spiral striations; columella usually with a well-developed twist or plait (Figs. 595-631). Widely distributed in North America ..... Genus *Stagnicola*[43, 79]

Adult shell small, generally less than 13 mm (but occasionally up to 15 or 16 mm) in length; spiral sculpture usually absent, very weak when present; columella generally without a twist or plait. Genus *Fossaria*[41] 14

14(13) Lateral teeth of the radula tricuspid (i.e., with three prominent cusps)[80]. Subgenus *Fossaria* s.str. 15

[The genus *Fossaria* contains the small lymnaeids, very few specimens of which have shells more than 12 or 13 mm in length, most being smaller. The spiral striations of the shell, characteristic of most other members of the family, are absent or poorly developed. The columella is most commonly smooth, without a twist or plait.

The type species of *Fossaria* is the Holarctic (but mainly Eurasian) *F. truncatula* (Müller)[81]. *Galba* Schrank 1803 is another name sometimes used for the genus, especially in Europe, but the type species (*Galba pusilla* Schrank) on which the name is based is unidentifiable (Pilsbry & Bequaert, 1927). Other synonyms are *Simpsonia* F.C. Baker 1911, preoccupied by *Simpsonia* Rochebrune 1905, and *Pseudogalba* F.C. Baker 1913, a replacement name for *Simpsonia* Baker.

Some 40 species or subspecies of North American fossarias have been named, but the majority of these will prove to be synonyms. Hubendick (1951) recognized only three species ("*Lymnaea*" *bulimoides*, "*L.*" *cubensis* and "*L.*" *humilis*), but that amount of "lumping" seems excessive. A definitive determination of the *Fossaria* species must await careful and detailed biological/morphological/conchological studies.]

Lateral teeth of the radula bicuspid (i.e., with only two prominent cusps)[80]. Subgenus *Bakerilymnaea* 21

[The main distinguishing feature of the subgenus *Bakerilymnaea* is the bicuspid lateral teeth of the radula, in contrast to the tricuspid lateral teeth of *Fossaria* s.str. Also, the species of *Bakerilymnaea* are mostly more globose and larger, and frequently more glossy. Because of their bicuspid lateral radular teeth, F.C. Baker (1928c) grouped the bakerilymnaeas (as the subgenus *Nasonia*, preoccupied by *Nasonia* Ashmead 1904) with *Stagnicola*. However, they are more closely allied to *Fossaria*.]

15(14) Adult shell (with about five whorls) very small, less than 7 mm in length (Fig. 571). Widely distributed, absent from eastern Canada, most of New England, and the Gulf and South Atlantic states ..... *Fossaria parva* (Lea)

Adult shell larger, more than 8 mm in length 16

16(15) Shell thickened, commonly whitish; spire generally obtuse, but it may be elongated; whorls usually strongly shouldered, especially at the aperture lip; outer apertural lip flattened. Inhabitant of northern cold-water lakes and streams (Fig. 568) . . . . . *Fossaria galbana* (Say)

Shell generally relatively thin, but it may be solid; spire elongate; whorls not shouldered, or with only weak or moderate shoulders; outer apertural lip rounded, sometimes compressed, but not flattened 17

17(16) Shell spire elongate and generally narrow, its length noticeably larger than the aperture length. Northern, from New York to Michigan and Iowa; southwestern Yukon and southern Alaska 18

Shell spire broad to narrow, but in shells with narrow spires, the spire length is not much greater than the aperture length 20

18(17) Body whorl tumid, globular; aperture subcircular (cyclostomoid) (Fig. 567). New York to Michigan . . . . . *Fossaria cyclostoma* (Walker)

Body whorl elongate-oval; aperture oval 19

19(18) Eastern North America, from New York to Iowa (Fig. 572) . . . *Fossaria tazewelliana* (Wolf)

Southwestern Yukon and southern Alaska (Fig. 583) . . . . . *Fossaria truncatula* (Müller)[81]

20(17) Whorls regularly increasing in size, terminating in a tumid, ovate body whorl; whorls evenly convex; spire broad; aperture ovate. Eastern and southeastern United States in distribution (Fig. 569) . . . . . *Fossaria humilis* (Say)

Whorls regularly or irregularly increasing in size, terminating in an elongate-ovate, sometimes narrow body whorl; whorls convex to flattened; spire broad to narrow; aperture elongate-oval. Widely distributed in North America, but absent from the southeastern United States (Figs. 570, 573-577) . . . . . *Fossaria obrussa* (Say) group

[Shells of the *Fossaria obrussa* group are rather variable, and about 15 forms have been described as "new" species. However, there are probably only several species in this group, and these are not defined by constantly different shell characters. Names that are in common use, in addition to *obrussa*, are *exigua* Lea, *modicella* Say, *peninsulae* Walker and *rustica* Lea. F.C. Baker (1928c) characterized these forms as follows:

obrussa [Figs. 570, 575] – ". . . one of the most widely distributed . . . [and] . . . most variable, of the American Lymnaeas. . . . Typically, *obrussa* may be known by its pointed spire, compressed body whorl and elongated and shouldered aperture, which is also strongly effuse at the anterior end; the inner lip is appressed to the body whorl about the middle of the aperture. The shape of the shell, of the aperture and of the inner lip is quite different from *modicella*, the shell being larger and more elongated, the last whorl not so convex; the aperture is longer and narrower and much more effuse, besides forming a distinct shoulder at its junction with the body whorl; the inner lip is more compressed in the middle where it joins the parietal wall. In shells of the same size, *modicella* has five whorls, while *obrussa* has four whorls; in form the young *obrussa* somewhat approach *modicella*. The shell is, typically, much larger than *modicella, parva* and the other members of the *humilis* group."

exigua [Fig. 573] – "... appears quite separable from *obrussa*. The spire is usually long and the whorls flatly rounded, the body whorl more or less compressed; the most noteworthy feature appears to be the very deep suture, which is almost channelled in some specimens, causing the whorls to be turban-shaped. This feature is present in the majority of the specimens examined. The aperture is also more regularly ovate than in *obrussa*, and the inner lip is peculiarly flattened near the umbilical region, giving rise to a pseudoplait. Some specimens resemble *modicella rustica*, but in that race the spire is acutely conical, the whorls regularly increase in size, the body whorl is not compressed in the middle, and the aperture is roundly ovate, while in *exigua* the spire is broadly turreted, the whorls are more or less disproportionate in size and the body whorl is very cylindrical."

modicella [Fig. 574] – "... closely related to the *humilis* of the southeastern part of the United States, differing in its narrower shell and longer aperture, and more or less impressed inner lip where it joins the parietal wall. ... *Obrussa* is larger and more elongated and the inner lip is notably compressed and bent inward at its junction with the parietal wall."

peninsulae [Fig. 576] – "... differs from typical *obrussa* in being more slender, with a longer, more turreted spire, deeper sutures and a more oval aperture. The body whorl is more cylindrical than in the typical form [*obrussa*]."

rustica [Fig. 577] – "... appears to be a modification of the *modicella* type of shell, characterized principally by its long, very acute spire and ovate aperture. Its long, pointed spire will distinguish it from any form of *modicella*. It is liable to be confounded with forms of *exigua*, but in that species the aperture is longer and narrower and inclined to be squarish, while in *rustica* it is more acutely rounded at the extremities. The spire in *rustica* is longer and more acute than in *exigua*, the spire whorls being less inflated. Half-grown specimens of *obrussa* are similar in general form, but differ in the form of the aperture, which is longer and narrower and forms a distinct shoulder at the junction of the outer lip with the body whorl, while in *rustica* this part of the lip is gracefully curved. The aperture is sometimes almost round and the spire varies much in height. *Rustica* is evidently more nearly related to *modicella* than to *obrussa* and may be considered a variety of the former."]

21(14) Shell ovate, dark amber in color, very highly polished. Southwestern Alaska ..... *Fossaria (Bakerilymnaea) perpolita* (Dall)

Shell globose, subglobose, ovate or conic, horn, pale yellowish, light to dark brown or pearl gray in color, generally moderately glossy, but may be dull 22

22(21) Shell globose, thin and fragile, whorls rapidly expanding, producing a very small spire and an obese body whorl; umbilicus small to perforate (Fig. 589). Sonoma County, California .... *Fossaria (Bakerilymnaea) sonomaensis* (Hemphill (in Pilsbry & Ferriss) 1906)[82]

Shell ovate to conic, umbilicus relatively large to practically imperforate 23

23(22) Adult shell (with above five whorls) moderately small to very small, less than 10 mm in length 24

Adult shell larger, nearly always more than 10 mm in length, generally 11-13 mm (occasionally up to 15 or 16 mm). Alabama west to northern Mexico and southern California, north to southern Canada from British Columbia to Saskatchewan (Figs. 584-586) .... *Fossaria (Bakerilymnaea) bulimoides* group

[Shells of the *Fossaria (Bakerilymnaea) bulimoides* group are quite variable, and several forms have been recognized as species, subspecies or morphs. The best known of these are *cockerelli* Pilsbry & Ferriss and *techella* Haldeman. Hibbard & Taylor (1960) believed *cockerelli* to be specifically distinct from *bulimoides* s.str. and *bulimoides*' subspecies *techella*. *Cockerelli* and *techella*, as well as *alberta* and *perplexa*, were considered to be only "morphs" of *bulimoides* by Clarke (1973). Taylor (1975) lists *perplexa* with *Fossaria* s.str. All of these taxa must be studied much more thoroughly before their exact systematic status can be determined. Described characteristics of these forms, along with those of *hendersoni* and *vancouverensis*, are given below:

bulimoides [Fig. 584] – "*Bulimoides* may be distinguished from *techella* and other races by its more *regularly ovate shape, less globose body-whorl, more elongate-ovate aperture and by the different manner in which the inner lip is appressed to the columellar region.* There is considerable variation in the rotundity of the whorls and in the length and acuteness of the spire. The inner lip also varies greatly, in some specimens being rolled or folded over into the umbilical region while in others it is expanded, approaching the *techella* form. *Bulimoides* somewhat resembles *cubensis*, differing in its nearly closed umbilical chink, folded inner lip, shorter and broader spire and its ovate shell. The whorls of *cubensis* are also rounder and more distinctly shouldered than are those of *bulimoides*" (F. C. Baker, 1911a: 213).

alberta – ". . . may be . . . recognized by its elongate-ovate outline, strong spiral striation, and smooth, folded inner lip" (F. C. Baker, 1919e: 538)[83].

cockerelli [Fig. 585] – "Shell subglobose, pale yellowish-corneous. . . . *Spire very short,* last whorl and aperture very large. Aperture short-ovate, its length three-fifths to two-thirds that of the shell. Columella broadly expanded, not folded. Umbilicus large. . . . This form differs from *L. bulimoides* and *L. techella* by its more globose shape and shorter spire. . . . *L.* . . . *sonomaensis* Hemphill [Fig. 589], from Sonoma county, California, approaches *cockerelli*, but differs by the more rapidly expanding last whorl, narrower flat columella and *narrower umbilicus*, which is like that of typical *bulimoides*" (Pilsbry & Ferriss, 1906: 162-163).

hendersoni – "Globose, very thin and fragile; periostracum light yellowish or brownish horn; . . . spire very short, depressed. . . . The only Lymnaeid likely to be confounded with *hendersoni* is *sonomaensis*, which differs in the form of the spire [higher] and the inner lip [not rolled over as much]. . . . The outline of the shell is . . . more ovate than in *sonomaensis* and the aperture is not expanded" (F. C. Baker, 1911a: 223, 224). "*Lymnaea hendersoni* Baker is within the range of variation of *S.* ["*Stagnicola*"] *cockerelli* as considered here. Two paratypes (USNM 570386) are smaller than usual for *S. cockerelli*, but can be matched by lots from Colorado and elsewhere. They probably were exposed to acid water, for the first one or two whorls have been etched; hence, on the low spire of these shells the effect is that of a truncate shell. This is an environmental, adventitious effect; the whorls are not 'coiled in the same plane' as Baker thought" (Hibbard & Taylor, 1960: 92).

perplexa – ". . . resembles both *parva* and *dalli*. It appears to stand midway between these species, being larger than *dalli* and smaller than *parva*. Its brown color of shell and aperture, deep sutures, fine, regular lines of growth without spiral lines, and its flattened and wide inner lip will distinguish it from related species" (F.C. Baker & Henderson, 1929: 104)[84].

techella [Fig. 586] – "Shell obese, with *acutely conic spire*, of five or six convex whorls; pale yellowish or light brown, finely striate and *usually malleated* . . . Last whorl very ventricose, umbilicus large. Aperture short-ovate, about three-fifths the total length; basal lip expanded, *columellar lip broadly dilated*, without a fold. . . . *Cubensis* has a more triangular and less broadly developed columellar expansion" (Pilsbry & Ferriss, 1906: 163, 164).

vancouverensis – "Shell differing from typical *bulimoides* in its larger size, more ovate and widely expanded aperture, wider inner lip which is less triangular than in typical *bulimoides*, and coarser sculpture which is almost rib-striate in some specimens" (F. C. Baker, 1939a: 144).]

24(23) Adult shell (with about five whorls) very small, less than 6 mm in length 25



FIGS. 573-582. Shells of Lymnaeidae (Lymnaeinae and Lancinae) (Figs. 573-580) and Physidae (Figs. 581, 582). FIG. 573. *Fossaria exigua*. FIG. 574. *F. modicella*. FIG. 575. *F. obrussa*. FIG. 576. *F. peninsulae*. FIG. 577. *F. rustica*. FIG. 578. *Lanx alta*. FIG. 579. *L. subrotunda*. FIG. 580. *Lanx hanni* = ? *L. patelloides*. FIG. 581. *Physella boucardi*. FIG. 582. *P. (Costatella) conoidea*. Measurement lines = 1 mm. Figs. 578 and 579 are from Tryon (1865i). Fig. 580 is from Walker (1925b).



FIGS. 583-592. Shells of Lymnaeidae (Lymnaeinae). FIG. 583. *Fossaria truncatula*. FIG. 584. *F. (Bakerilymnaea) bulimoides*. FIG. 585. *F. (B.) cockerelli*. FIG. 586. *F. (B.) techella*. FIG. 587. *F. (B.) cubensis*. FIG. 588. *F. (B.) dalli*. FIG. 589. *F. (B.) sonomaensis*. FIG. 590. *Lymnaea atkaensis*. FIG. 591. *L. stagnalis appressa*. FIG. 592. *L. stagnalis sanctaemariae*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 593-602. Shells of Lymnaeidae (Lymnaeinae). FIG. 593. *Pseudosuccinea columella*. FIG. 594. *Radix auricularia*. FIG. 595. *Stagnicola elrodiana*. FIG. 596. *S. exilis*. FIG. 597. *S. exilis*. FIG. 598. *S. neopalustris*. FIG. 599. *S. traski*. FIG. 600. *S. elodes*. FIG. 601. *S. jolietensis* = *S. elodes*. FIG. 602. *S. alpenensis* = *S. elodes*. Measurement lines are divided into millimeters.



FIGS. 603-613. Shells of Lymnaeidae (Lymnaeinae). FIG. 603. *Stagnicola impedita* = ? *S. elodes*. FIG. 604. *S. newfoundlandensis* = ? *S. elodes*. FIG. 605. *S. elodes* form *reflexa*. FIG. 606. *S. wyomingensis* = ? *S. elodes*. FIG. 607. *S. arctica*. FIG. 608. *S. bonnevillensis*. FIG. 609. *S. catascopium*. FIG. 610. *S.* ? *catascopium*. FIG. 611. *S. laurentiana* = ? *S. elodes*. FIG. 612. *S. contracta*. FIG. 613. *S. elrodi*. Measurement lines are divided into millimeters.



FIGS. 614-623. Shells of Lymnaeidae (Lymnaeinae). FIG. 614. *Stagnicola emarginata*. FIG. 615. *S. emarginata* form *serrata*. FIG. 616. *S. emarginata* form *canadensis*. FIG. 617. *S. emarginata* form *nashotahensis*. FIG. 618. *S. gabbi*. FIG. 619. *S. idahoense*. FIG. 620. *S. hinkleyi*. FIG. 621. *S. mighelsi*. FIG. 622. *S. oronoensis*. FIG. 623. *S. petoskeyensis*. Measurement lines are divided into millimeters.



FIGS. 624-632. Shells of Lymnaeidae (Lymnaeinae and Lancinae). FIG. 624. *Stagnicola kingi* = *S. utahensis*. FIG. 625. *S. walkeriana*. FIG. 626. *S. woodruffi*. FIG. 627. *S. nasoni* = ?*S. woodruffi*. FIG. 628. *S. (Hinkleyia) caperata*. FIG. 629. *S. (H.) caperata*, periostracal ridges on body whorl. FIG. 630. *S. (H.) montanensis*. FIG. 631. *S. (H.) pilsbryi*. FIG. 632. *Fisherola nuttalli lancides*, top (left figure) and right lateral (right figure) views. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 633-640. Shells of Lymnaeidae (Lancinae) (Figs. 633, 634) and Physidae (Physinae) (Figs. 635-640). FIG. 633. *Lanx patelloides*. FIG. 634. *L. (Walkerola) klamathensis*. FIG. 635. *Physa jennessi*. FIG. 636. *Physa skinneri*. FIG. 637. *Physa skinneri*, large unnamed morph. FIG. 638. *Physella boucardi*. FIG. 639. *Physella columbiana*. FIG. 640. *Physella cooperi*. Measurement lines = 1 mm or are divided into millimeters.

Adult shell moderately small, 7 to 9 mm in length (Fig. 587). Southern United States from Florida to Texas ..... *Fossaria (Bakerilymnaea) cubensis* (Pfeiffer)

25(24) Shell pale brown. Southern Manitoba and southern Alberta, western region of the Great Lakes system, upper Mississippi drainage, and south in the Rocky Mountains to Arizona (Fig. 588) ..... *Fossaria (Bakerilymnaea) dalli* (F. C. Baker)

Shell dark brown. Found sporadically in Washington, California, Montana, Utah, Nevada and Arizona ..... *Fossaria (Bakerilymnaea) bulimoides* form *perplexa* (F. C. Baker)[84]

FAMILY PHYSIDAE

The Physidae are mainly a New World family, with only a few species occurring in Eurasia and Africa. In North America, the physids are readily recognized by a combination of several characters. Their lack of an operculum distinguishes them from all of the Prosobranchia. Their high-spired shell separates them from the Planorbidae and Ancylidae, and their sinistral (left coiled) shell marks them as being different from the Lymnaeidae.

In North America, the Physidae are the most abundant and wide-spread of the freshwater gastropods. They may be found in all types of habitats, and some species seem to be the most resistant to pollution of all the freshwater mollusks. In addition to being highly adaptable, the physids have undergone considerable diversification, much of which is not clearly exhibited in their shells. Many of the species are not easy to identify on shell characters alone.

Identification Key for the Physidae[85]

1 Mantle edge digitate (with finger-like projections) 2

Mantle edge without digitations; mantle edge may or may not be serrated 3

2(1) Digitations occur on both sides of the mantle; tip of shell spire rounded (Figs. 635-637). Canada and northern United States ..... Genus *Physa*[85]

Digitations occur only on the parietal side of the mantle (Figs. 581, 582, 638-698). Widely distributed and common throughout North America ..... Genus *Physella*[85]

3(1) Mantle edge smooth; mantle does not extend beyond the edge of the shell apertural lip 5

Mantle edge serrated and extending beyond the edge of the shell apertural lip, partly overlapping the shell. Texas. Genus *Stenophysa*[47] 4



FIGS. 641-648. Shells of Physidae (Physinae). FIG. 641. *Physella hordacea*. FIG. 642. *P. lordi*. FIG. 643. *P. microstriata*. FIG. 644. *P. traski*. FIG. 645. *P. utahensis*. FIG. 646. *P. virginea*. FIG. 647. *P. gyrina gyrina*. FIG. 648. *P. gyrina gyrina* morph *elliptica*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 649-658. Shells of Physidae (Physinae). FIG. 649. *Physella gyrina gyrina* morph *hildrethiana*. FIG. 650. *P. gyrina alba*. FIG. 651. *P. gyrina ampullacea*. FIG. 652. *P. gyrina athearni*. FIG. 653. *P. gyrina aurea*. FIG. 654. *P. gyrina aurea* morph *albofilata*. FIG. 655. *P. gyrina bayfieldensis*. FIG. 656. *P. gyrina cylindrica*. FIG. 657. *P. gyrina gouldi*. FIG. 658. *P. gyrina hawnii*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 659-667. Shells of Physidae (Physinae). FIG. 659. *Physella gyrina microstoma*. FIG. 660. *P. gyrina sayi*. FIG. 661. *P. gyrina smithiana*. FIG. 662. *P. propinqua propinqua*. FIG. 663. *P. propinqua nuttalli*. FIG. 664. *P. propinqua nuttalli* morph *triticea*. FIG. 665. *P. propinqua nuttalli* morph *venusta*. FIG. 666. *P. ancillaria*. FIG. 667. *P. globosa*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 668-676. Shells of Physidae (Physinae). FIG. 668. *Physella magnalacustris*. FIG. 669. *P. parkeri parkeri*. FIG. 670. *P. parkeri latchfordi*. FIG. 671. *P. vinosa*. FIG. 672. *P. (Costatella) cubensis cubensis*. FIG. 673. *P. (C.) cubensis peninsulae*. FIG. 674. *P. (C.) costata*. FIG. 675. *P. (C.) hendersoni hendersoni*. FIG. 676. *P. (C.) hendersoni hendersoni* morph *ariomus*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 677-685. Shells of Physidae (Physinae). FIG. 677. *Physella (Costatella) hendersoni* ssp. FIG. 678. *P. (C.) acuta*. FIG. 679. *P. (C.) bottimeri*. FIG. 680. *P. (C.) heterostropha heterostropha*. FIG. 681. *P. (C.) heterostropha pomila*. FIG. 682. *P. (C.) heterostropha halei*. FIG. 683. *P. (C.) humerosa*. FIG. 684. *P. (C.) johnsoni*. FIG. 685. *P. (C.) osculans*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 686-694. Shells of Physidae (Physinae). FIG. 686. *Physella (Costatella) spelunca*. FIG. 687. *P. (C.) squalida*. FIG. 688. *P. (C.) virgata virgata*. FIG. 689. *P. (C.) virgata virgata* morph *parva*. FIG. 690. *P. (C.) virgata anatina*. FIG. 691. *P. (C.) virgata berendti*. FIG. 692. *P. (C.) virgata concolor*. FIG. 693. *P. (C.) virgata concolor* morph. FIG. 694. *P. (C.) virgata rhyssa*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 695-702. Shells of Physidae (Physinae and Aplexinae). FIG. 695. *Physella (Costatella) integra integra*. FIG. 696. *P. (C.) integra integra* morph *walkeri*. FIG. 697. *P. (C.) integra brevispira*. FIG. 698. *P. (Petrophysa) zionis*. FIG. 699. *Aplexa elongata*. FIG. 700. *A. elongata* morph *tryoni*. FIG. 701. *Stenophysa marmorata*. FIG. 702. *S. maugeriae*. Measurement lines = 1 mm or are divided into millimeters.

4(3) Shell relatively small, less than 16 mm in length, horn to light or dark tan in color, usually translucent, seldom variegated (Fig. 701). Texas . . . . . *Stenophysa marmorata* (Guilding)

Shell relatively large, up to 30 mm or more in length, tan to chestnut brown in color, opaque, commonly variegated (Fig. 702). Texas . . . . . *Stenophysa maugeriae* (Gray)

5(3) Shell elongate, nearly spindle-shaped; shell surface glossy; spire long (Figs. 699, 700). Canada and northern United States . . . . . *Aplexa elongata* (Say)[86]

Shell subglobose, globular; shell surface dull; spire very short (Fig. 698). Utah . . . . . *Physella (Petrophysa) zionis* (Pilsbry)

FAMILY PLANORBIDAE

The Planorbidae in North America range in size from minute to relatively large (i.e., from about 1 mm in diameter to over 30 mm), but with few exceptions their shells are all discoidal, i.e., coiled in one plane. The animals are all sinistral, i.e., coiled to the left or in a counter-clockwise manner and having respiratory, excretory and reproductive systems terminating on the left side (Fig. 703). However, their shells do not always appear to be sinistral; those of many species seem to be dextral. This is because such shells tip to the left side in life and the type of apertural margin which develops in such cases is correspondingly slanted. In shells tipped to the left in such a fashion, the lower side (left side) is the spire side and the upper side (right side) is the umbilical side (Fig. 704). Such dextral-appearing shells on a sinistral animal are termed "pseudodextral" or "ultrasinistral".

A secondary gill (a pseudobranch) is situated on the left side of the animal, near the pneumostome and in close proximity to the anus (Fig. 703). The pseudobranch aids the mantle cavity in respiration.

A striking characteristic of nearly all planorbid snails is that the respiratory pigment of the blood or haemolymph is haemoglobin. This gives a reddish appearance to the animal, if the color is not masked by melanin pigments of the skin. Albino snails, and those with little pigment, appear bright red. (The genus *Drepanotrema* apparently lacks red haemolymph.)

The Planorbidae appear to be closely related to the Ancylidae, and some authors (e.g., Starobogatov, 1970) have combined the two as a single family.

Identification Key for the Planorbidae

1 Shell small, that of adults less than 8 mm in diameter 2

Shell larger, that of adults more than 8 mm and up to or more than 30 mm in diameter 23

2(1) Shell costate (Fig. 706). Canada and northern United States . . . . . *Gyraulus (Armiger) crista* (Linnaeus)

Shell not costate 3



FIG. 703. Planorbid (*Biomphalaria*) with shell removed to show aspects of anatomy (left side) (from Barbosa et al., 1968, after Demian).



FIG. 704. Terminology of a planorbid shell.

3(2) Shell minute, that of adults 2 mm or less in diameter. Coosa River, Alabama 4

Shell larger, that of adults more than 2 mm in diameter 8

4(3) Shell crepidulaform in shape, i.e., limpet-like with a small coil at the apex (Fig. 749). Coosa River, Alabama . . . . . *Amphigyra alabamensis* Pilsbry

Shell planorboid. Genus *Neoplanorbis*[62, 87] 5

5(4) Shell umbilicate, columella dentate 6

Shell perforate, columella smooth 7

6(5) Shell periphery carinate, umbilicus narrow (Fig. 752). Coosa River, Alabama . . . . . *Neoplanorbis carinatus* Walker

Shell periphery obtusely angled, umbilicus wider (Fig. 754). Coosa River, Alabama . . . . . *Neoplanorbis umbilicatus* Walker

7(5) Shell spirally striate, periphery carinate (Fig. 750). Coosa River, Alabama . . . . . *Neoplanorbis tantillus* Pilsbry

Shell without spiral striae, periphery rounded (Fig. 753). Coosa River, Alabama . . . . . *Neoplanorbis smithi* Walker

8(3) Shell very compressed, body whorl relatively flattened; aperture or body whorl without "teeth" or lamellae 9

Shell higher, body whorl moderately high; inside aperture or body whorl with "teeth" or lamellae. Genus *Planorbula*, in part 22

9(8) Shell either extremely flattened and multi-whorled or with numerous, low, close-set spiral ridges (lirae). Florida, Texas and southern Arizona. Genus *Drepanotrema* 10

Shell flattened, but not extremely so; not multi-whorled; without spiral ridges (lirae) 12

10(9) Shell extremely flattened; multi-whorled; without spiral ridges (lirae). Subgenus *Fossulorbis* 11

Shell not extremely flattened; with fewer, more rapidly enlarging whorls; sculptured with numerous, low lirae. Subgenus *Antillorbis*. (Fig. 710). Southern Arizona and southern Texas . . . . . *Drepanotrema (Antillorbis) aeruginosum* (Morelet)

11(10) Shell periphery strongly keeled (Fig. 711). Florida, Texas . . . . . *Drepanotrema (Fossulorbis) kermatoides* (d'Orbigny)

Shell periphery rounded or obtusely angular (Fig. 715). Southern Texas . . . . . *Drepanotrema (Fossulorbis) cimex* (Moricand)

12(9) Spire pit (on left side of shell) shallow and wide 13

Spire pit (on left side of shell) relatively deep and narrow 17

13(12) Height of body whorl relatively rapidly increasing toward the aperture (Fig. 727). Illinois, Missouri and Arkansas . . . . . *Menetus (Micromenetus) sampsoni* (Sampson)[53, 54, 55]

Height of body whorl nearly equal from one side to the other. Genus *Gyraulus* 14

14(13) Adult shells 4 to 7 mm in diameter, variable, with the body whorl not evenly rounded or with a peripheral keel or with a hirsute periostracum or a malleated surface or with any combination of these features.[88] Subgenus *Gyraulus* s.s. (Fig. 705). Canada and northern United States from Maine to Virginia and west to Idaho . . . . . *Gyraulus deflectus* (Say)

Adult shells 3 to 5 mm in diameter, variable, with the body whorl evenly rounded or with upper lateral surface slightly flattened; without a peripheral keel or a hirsute periostracum or malleated surface.[88] Subgenus *Torquis* 15

15(14) Shell relatively high (Fig. 708). Canada, North Dakota and Wisconsin . . . . . *Gyraulus (Torquis) hornensis* F.C. Baker[48]

Shell relatively flattened 16

16(15) Shell whitish or yellowish, semi-transparent, entirely or nearly planispiral, appearing almost the same from both sides. Characteristic of aquatic habitats that are subject to periodic drying[88] (Fig. 707). Canada and northern United States, south in the Rocky Mountains to New Mexico . . . . . *Gyraulus (Torquis) circumstriatus* (Tryon)

Shell brownish, translucent but not transparent, not planispiral but with apical and umbilical aspects clearly different. Characteristic of permanent and (occasionally) temporary aquatic habitats[88] (Fig. 709). Widely distributed throughout North America . . . . . *Gyraulus (Torquis) parvus* (Say)

17(12) Shell with carinate periphery 18

Shell with rounded, subangular or angular periphery 20

18(17) Western in distribution. Alaska south to Alberta and southern California (Figs. 722, 723) . . . . . *Menetus opercularis* (Gould)[52]

Found east of the Rocky Mountains 19

19(18) Relative height of body whorl rapidly increasing toward the aperture (Fig. 725). Ohio, Alabama . . . . . *Menetus (Micromenetus) brogniartianus* (Lea)[53, 54, 55]

Relative height of body whorl nearly equal from one side to the other (Fig. 746). Widely distributed in North America . . . . . *Promenetus exacuous* (Say)

20(17) Relative height of body whorl rapidly increasing toward the aperture (Figs. 724, 726). Widely distributed in the eastern United States . . . . . *Menetus (Micromenetus) dilatatus* (Gould)[53, 54, 55]

Relative height of body whorl nearly equal from one side to the other 21

21(20) Periphery of body whorl more or less angular or subangular (Figs. 722, 723). Alaska south to Alberta and southern California . . . . . *Menetus opercularis* (Gould)

Periphery of body whorl rounded (Fig. 747). Widely distributed in Canada, the western United States, and east to Oklahoma, Ohio and New York . . . . . *Promenetus umbilicatellus* (Cockerell)[58, 59]

22(8) Lamellae in last whorl prominent but not especially large; lower palatal lamella relatively short and straight or only slightly curved (Figs. 741, 742). Widely distributed in eastern North America . . . . . *Planorbula armigera armigera* (Say)

Lamellae in last whorl especially large; lower palatal lamella long, prominently curved (Figs. 743, 744). Alabama and Florida . . . . . *Planorbula armigera wheatleyi* (Lea)[57]

23(1) Shell thin, often rather fragile, body whorl relatively depressed 24

Shell thicker, usually rather solid, body whorl may or may not be relatively depressed, often high 26

24(23) Southern in distribution (Florida to Texas and Arizona). Genus *Biomphalaria* 25

Distribution northern and in the western mountains (Canada and North Dakota, south to New Mexico in the Rocky Mountains) (Fig. 745) . . . . . *Planorbula campestris* (Dawson)

25(24) Shell medium in size, that of adults with five or more whorls larger than 15 mm in diameter (Fig. 712). Florida . . . . . *Biomphalaria glabrata* (Say)

Shell small, that of adults with five or more whorls less than 10 mm in diameter (Fig. 713). Florida to Texas and Arizona . . . . . *Biomphalaria havanensis* (Pfeiffer)

26(23) Body whorl containing lamellae or "teeth" (Figs. 741, 742). Widely distributed in eastern North America . . . . . *Planorbula armigera armigera* (Say)

Body whorl without lamellae or "teeth" 27

27(26) Shell with few, rapidly increasing whorls; body whorl disproportionately large. Genus *Vorticifex*, subgenus *Parapholyx*.[60] Western in distribution 28

Shell with more than a few, often many whorls, that do not increase especially rapidly in size; body whorl not disproportionately large 29

28(27) Whorl angular or subangular around the concave columellar area (Fig. 751). Lakes in Nevada and California . . . . . *Vorticifex (Parapholyx) solida* (Dall)[61]

Whorl not angular or subangular around the basal columellar area (Fig. 748). Rivers and lakes in California and Oregon . . . . . *Vorticifex (Parapholyx) effusa* (Lea)

29(27) Shell spire (left side) strongly inverted, with a more or less deep conical depression; spire side of body whorl with or without a strong keel. Genus *Helisoma* 30

Shell spire (left side) not strongly inverted, with a shallow depression, no depression or exverted (raised above body whorl); spire side of body whorl rounded or angular. Genus *Planorbella* 35

30(29) Shell concave on both sides. Subgenus *Helisoma* s.s. 31

Shell concave on the left side, convex on the right side. Western in distribution. Subgenus *Carinifex* 33

31(30) Shell smaller, less than 7 mm in diameter, umbilical (basal, right) side with two chestnut-brown spiral bands. Isolated localities in North Carolina and Louisiana . . . . . *Helisoma eucosmium* (Bartsch)[50]

Shell larger, adults more than 7 mm in diameter, umbilical (basal, right) side without spiral color bands 32

32(31) Shell with basal (right) carina variously developed, but not close to the shoulder; transverse sculpture moderate to fine (Fig. 714). Widely distributed in most of North America . . . . . *Helisoma anceps anceps* (Menke)[50]

Shell with basal (right) carina very accentuated and at or close to the lower basal peripheral angle; transverse sculpture coarse. Lake Superior and Albany, Attawapiskat and Winnipeg river systems, Ontario . . . . . *Helisoma anceps royalense* (Walker)[50]

33(30) Widely distributed and quite variable (Figs. 720, 721). California, Idaho, Nevada, Oregon and Utah . . . . . *Helisoma (Carinifex) newberryi newberryi* (Lea)[51]

Restricted to either Jackson Lake, Wyoming, or Eagle Lake, California 34

34(33) Shell smaller (that of adults less than 12 mm in diameter), buff or tan in color (Figs. 716, 717). Jackson Lake, Wyoming . . . . . *Helisoma (Carinifex) newberryi jacksonense* Henderson

Shell larger (that of adults up to 13.5 mm in diameter), white or horn in color (Figs. 718, 719). Eagle Lake, California . . . . . *Helisoma (Carinifex) newberryi occidentale* Hanna

35(29) Body whorl at shell aperture campanulate (flared). Subgenus *Planorbella* s.s.[56, 89] 36

Body whorl at shell aperture straight, not campanulate 38

36(35) Shell spire (left side) conically raised above body whorl (Fig. 729). Howe Lake, Michigan . . . . . *Planorbella multivolvis* (Case)

Shell spire (left side) either slightly inverted, flat or obtusely raised above body whorl 37

37(36) Shell spire (left side) slightly inverted, flat or very slightly raised above the body whorl (Fig. 728). Widely distributed in northern United States and Canada . . . . . *Planorbella campanulata campanulata* (Say)

Shell spire (left side) obtusely raised above body whorl. Northwestern Ontario . . . . . *Planorbella campanulata collinsi* (F.C. Baker)

38(35) Shell surface usually dull, usually rough in texture, with raised transverse thread-like striae. Widely distributed in North America. Subgenus *Pierosoma*[90] 39

Shell surface usually glossy, relatively smooth, without raised transverse thread-like striae (Figs. 738-740). Florida. Subgenus *Seminolina*[91] 48

39(38) Species of western North America 40

Species of central and eastern North America 42

40(39) Shell small, specimens with four whorls about 10 mm in major diameter. Southeastern Oregon and northwestern Utah . . . . . *Planorbella (Pierosoma) oregonensis* (Tryon)

Shell larger, adults 15-30 mm in major diameter 41

41(40) Greatest height of adults exceeding 12 mm; greatest width of shell less than twice the greatest height (Figs. 730, 733). Widely distributed in western North America ..... *Planorbella (Pierosoma) ammon* (Gould) group[92]

Greatest height of adults 10-12 mm; greatest width of shell generally more than twice the greatest height (Fig. 734). Widely distributed in western North America ..... *Planorbella (Pierosoma) trivolvis subcrenata* (Carpenter)[93]

42(39) Carinae or strong angulations present on the outer edges of both the right (umbilical) and left (spire) side of the body whorl of the shell 43

Carinae absent, although a rather strong angulation may be present on the upper surface of the body whorl of the spire 46

43(42) Shells larger, those of adults more than 18 mm in greatest diameter; spire may be flat or sunken into a bowl-like depression 44

Shells smaller, those of adults less than 18 mm in greatest diameter; spire flat, not inverted or sunken into a bowl-like depression (Fig. 737). Michigan, northern Illinois and Wisconsin ..... *Planorbella (Pierosoma) truncata* (Miles)

44(43) Carinae cord-like, strong and acutely angled; body whorl flat or concave abaxially. Northern Minnesota .... *Planorbella (Pierosoma) corpulenta vermilionensis* (F.C. Baker)[94]

Carinae not cord-like 45

45(44) Upper surface of shell almost entirely flat; maximum height at aperture 14 mm or more; ratio of greater height to greater diameter more than 0.75 in many specimens. Headwaters of Rainy River system, western Ontario ..... *Planorbella (Pierosoma) corpulenta whiteavesi* (F.C. Baker)[94]

Body whorl higher than penultimate whorl, causing spire to be sunken; maximum height at aperture less than 14 mm; ratio of greater height to greater diameter less than 0.75. Western Ontario, Minnesota and Manitoba .... *Planorbella (Pierosoma) corpulenta corpulenta* (Say)[95]

46(42) Shell height up to 24 mm or more; surface glossy, growth lines fine (Fig. 732). Lower Cape Fear River, North Carolina .... *Planorbella (Pierosoma) magnifica* (Pilsbry)

Shell more compressed, less than 16 mm in height; surface dull, growth lines pronounced 47

47(46) Inverted portion of shell spire relatively wide, concavely smooth-sided and bowl-like (Fig. 731). Canadian Interior Basin and northern United States from Massachusetts west to Minnesota ... *Planorbella (Pierosoma) pilsbryi* (F.C. Baker)[96]



FIGS. 705-709. Shells of Planorbidae (Planorbinae, Planorbini). FIG. 705. *Gyraulus deflectus*, umbilical, apertural and spire views (left to right). FIG. 706. *G. (Armiger) crista*. FIG. 707. *G. (Torquis) circumstriatus*. FIG. 708. *G. (T.) hornensis*. FIG. 709. *G. (T.) parvus*. Measurement lines = 1 mm.



FIGS. 710-714. Shells of Planorbidae (Planorbinae, Drepanotremini, Biomphalariini and Helisomini). FIG. 710. *Drepanotrema (Antillorbis) aeruginosum*, umbilical, apertural and spire views (left to right). FIG. 711. *D. (Fossulorbis) kermatoides*. FIG. 712. *Biomphalaria glabrata*. FIG. 713. *B. havanensis*. FIG. 714. *Helisoma anceps anceps*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 715-719. Shells of Planorbidae (Planorbinae, Drepanotremini and Helisomini). FIG. 715. *Drepanotrema (Fossulorbis) cimex*, umbilical, apertural and spire views (left to right). FIG. 716. *Helisoma (Carinifex) newberryi jacksonense*. FIG. 717. *H. (C.) newberryi jacksonense*. FIG. 718. *H. (C.) newberryi occidentale*. FIG. 719. *H. (C.) newberryi occidentale*. Measurement lines = 1 mm or are divided into millimeters. FIG. 715 is from Barbosa et al. (1968).



FIGS. 720-724. Shells of Planorbidae (Planorbinae, Helisomini). FIG. 720. *Helisoma (Carinifex) newberryi newberryi*, umbilical, apertural and spire views (left to right). FIG. 721. *H. (C.) newberryi newberryi* form *ponsonbyi*. FIG. 722. *Menetus opercularis*. FIG. 723. *M. opercularis* ? form *callioglyptus*. FIG. 724. *M. (Micromenetus) dilatatus*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 725-727. Shells of Planorbidae (Planorbinae, Helisomini). FIG. 725. *Menetus (Micromenetus) brogniartianus*, umbilical, apertural and spire views (left to right). FIG. 726. *Me. (Mi.) dilatatus*. FIG. 727. *Me. (Mi.) sampsoni*. Measurement lines = 1 mm.



FIGS. 728-731. Shells of Planorbidae (Planorbinae, Helisomini). FIG. 728. *Planorbella campanulata campanulata*, umbilical apertural and spire views (left to right). FIG. 729. *P. multivolvis*. FIG. 730. *Pl. (Pierosoma) ammon*. FIG. 731. *Pl. (Pi.) pilsbryi infracarinata*. Measurement lines are divided into millimeters.



FIGS. 732-735. Shells of Planorbidae (Planorbinae, Helisomini). FIG. 732. *Planorbella (Pierosoma) magnifica*, umbilical, apertural and spire views (left to right). FIG. 733. *Pl. (Pi.) occidentalis*. FIG. 734. *Pl. (Pi.) trivolvis subcrenata*. FIG. 735. *Pl. (Pi.) tenuis*. Measurement lines are divided into millimeters.



FIGS. 736-740. Shells of Planorbidae (Planorbinae, Helisomini). FIG. 736. *Planorbella (Pierosoma) trivolvis trivolvis*, umbilical, apertural and spire views (left to right). FIG. 737. *Pl. (Pi.) truncata*. FIG. 738. *Pl. (Seminolina) duryi*. FIG. 739. *Pl. (S.) duryi* form *seminolis*. FIG. 740. *Pl. (S.) scalaris*. Measurement lines are divided into millimeters.



FIGS. 741-745. Shells of Planorbidae (Planorbinae, Helisomini). FIG. 741. *Planorbula armigera armigera*, umbilical, apertural and spire views (left to right). FIG. 742. *P. jenksii* = *P. armigera armigera*. FIG. 743. *P. armigera wheatleyi*. FIG. 744. *P. armigera wheatleyi*, showing lamellae in body whorl. FIG. 745. *P. campestris*. Measurement lines = 1 mm or are divided into millimeters.



FIGS. 746-750. Shells of Planorbidae (Planorbinae (Helisomini) and Neoplanorbinae). FIG. 746. *Promenetus exacuous*, umbilical, apertural and spire views (left to right). FIG. 747. *P. umbilicatellus*. FIG. 748. *Vorticifex (Parapholyx) effusa*. FIG. 749. *Amphigyra alabamensis*. FIG. 750. *Neoplanorbis tantillus*. Measurement lines = 1 mm or are divided into millimeters.



751



752



753



754

FIGS. 751-754. Shells of Planorbidae (Planorbinae (Helisomini) and Neoplanorbinae). FIG. 751. *Vorticifex (Parapholyx) solida* form *optima*. FIG. 752. *Neoplanorbis carinatus*. FIG. 753. *N. smithi*. FIG. 754. *N. umbilicatus*. Measurement lines = 1 mm. Figs. 752-754 are from Walker (1908c).

Inverted portion of shell spire narrower, generally not smooth-sided or bowl-like (Figs. 734, 736). Found throughout North America . . . . . *Planorbella (Pierosoma) trivolvis* (Say)[93, 97]

48(38) Shell either planate, with an inverted spire, or physoid, i.e., with an everted, raised spire; physoid individuals wider, usually more widely umbilicate and generally with the anterior aperture margin protruding more than the posterior shell margin (when viewed from the spire end) (Fig. 772). Northern to southern Florida . . . . . *Planorbella (Seminolina) duryi* (Wetherby)[98]

Shell physoid only, narrower, usually more narrowly umbilicate and generally with the posterior aperture margin protruding more than the anterior shell margin (when viewed from the spire end) (Fig. 774). Southern Florida . . . . . *Planorbella (Seminolina) scalaris* (Jay)

FAMILY ANCYLIDAE

The Ancylidae are another of the gastropod families with a world-wide distribution. In North America, they all have small cap-shaped (patelliform, ancyliform, limpet-shaped) shells in which the apices are on the right side, or tilted toward the right (Fig. 755b). Among freshwater limpets, such a shell has been derived from ancestors with sinistrally coiled shells, and in the Ancylidae the arrangement of the body morphology is always sinistral, i.e., the "gill" (pseudobranch), and the pulmonary, reproductive and excretory openings are all on the animal's left side. The two other North American freshwater snail families with members having patelliform shells, the Acroloxidae and the Lymnaeidae (Lancinae), are dextral in organization.

The Ancylidae seem to be closely related to the Planorbidae, but they differ from the latter in one conspicuous way: all ancylids have haemocyanin as their blood pigment rather than haemoglobin (which gives the planorbids their red body color). Within the Ancylidae, the North American genus *Rhodacmea* is most closely related to the Eurasian and North African genus *Ancylus*.

Among the ancylid subfamilies, the Ferrissinae have the widest distribution, both naturally and artificially. Pond species seem to be easily transported through human activities; riverine species are less tolerant.

Identification Key for the Ancylidae[99]

1 Shell elevated, apex in midline, tinged with pink or red inside and out, radially striate, with a notch-shaped depression evident in unworn specimens. Apertural lip broad and flat. Radular teeth in rows about 30 microns apart, with prominent inner cusps (Fig. 775)[100]. Penis simple, without a flagellum. In rivers in the southeastern states. Genus *Rhodacmea* 2

Shell elevated or depressed, apex in midline or to the right, the same color as the rest of the shell, finely radially striate or smooth. Apertural lip arched or flat, broad or narrow. Radular teeth in rows about 6-10 microns apart, without prominent inner cusps (Fig. 775)[100]. Penis with or without a flagellum. Widely distributed in running or standing water 4



FIG. 755. Shell terminology of freshwater limpets. a, A dextral shell (note that the apex is directed to the *left*); b, a sinistral shell (note that the apex is directed to the *right*). "Radial" sculpture on ancyliform shells corresponds to "spiral" sculpture on coiled shells. "Concentric" sculpture on ancyliform shells corresponds to "transverse" sculpture on coiled shells; on freshwater limpets it usually consists only of growth lines.



FIGS. 756-758. Shells of Ancylidae (Ancylinae). FIG. 756. *Rhodacmea cahawbensis* = *R. elatior*. FIG. 757. *R. filosa*. FIG. 758. *R. rhodacme* = *R. hinkleyi*. Measurement lines = 1 mm. Fig. 756 is from Walker (1917b).

2(1) Shell more or less ribbed with strong radiating lines extending from the apex to the apertural lip (Figs. 757, 759) . . . . . *Rhodacmea filosa* (Conrad)

Shell smooth, or nearly so 3

3(2) Shell moderately elevated, apex usually conspicuous in older specimens. Posterior slope straight or slightly concave; anterior slope straight or slightly convex (Figs. 758, 760) . . . . . *Rhodacmea hinkleyi* (Walker)

Shell very elevated, apex usually eroded in older specimens. Posterior slope straight or slightly convex, anterior slope clearly convex (Fig. 756) . . . . . *Rhodacmea elatior* (Anthony)

4(1) Shell usually elevated, but variable. Apex with fine radial striae, often eroded in older specimens. Aperture narrow to broadly ovate, entirely open or with a horizontal shelf-like septum closing the posterior part. Pseudobranch of one lobe, flat. Penis with a flagellum. Widely distributed in streams and standing water. Genus *Ferrissia* 5

Shell usually depressed. Apex smooth, with no trace of radial striae. Aperture ovate to subcircular, always open. Penis with or without a flagellum. Pseudobranch of two lobes, the lower of which is elaborately folded. In standing water, principally in eastern states and south 9

5(4) Shell thin, fragile, very much depressed, often a glossy red-brown color. Apex fairly prominent as a rounded bump in the right posterior quadrant. Length of shell to about 5 mm (Fig. 766). In streams in southern Alabama . . . . . *Ferrissia mcneili* Walker

Shell not as above, usually more elevated, color variable from straw-yellow to dark gray. Apex prominent to obtuse, in the midline or to the right. Length from 2 to 10 mm. Widely distributed in various habitats 6

6(5) Shell robust, to 7 mm long, elevated, aperture elliptical. Apex in midline or slightly to the right; anterior slope convex, posterior slope gently concave, lateral slopes approximately straight. Calcareous material often thick inside the shell (Figs. 761, 767). Many populations are smaller, especially those west of the Rocky Mountains. Widely distributed in North America in rivers and streams . . . . . *Ferrissia rivularis* (Say)

Shell not as above; habitat in standing water 7

7(6) Shell large, elevated, very narrow, length to 9 mm. Apex obtuse, in the midline; posterior slope flat or gently concave; lateral slopes straight or faintly concave. Apertural lip often arched. Canada and adjacent states, on vegetation in lakes . . . . . *Ferrissia parallelus* (Haldeman)

Shell in standing water, but not as above 8



759



760



761



762



763

FIGS. 759-763. Shells of Ancylidae (Ancylinae, Ferrissinae and Laevapecinae). FIG. 759. *Rhodacmea filosa*. FIG. 760. *R. rhodacme* = *R. hinkleyi*. FIG. 761. *Ferrissia rivularis*. FIG. 762. *Hebetancylus excentricus*. FIG. 763. *Laevapex fuscus*. Measurement lines = 1 mm.



FIGS. 764-771. Shells of Ancylidae (Ferrissinae and Laevapecinae). FIG. 764. *Ferrissia californica* = *F. fragilis*. FIG. 765. *F. shimeki* = *F. fragilis*. FIG. 766. *F. mcneili*. FIG. 767. *F. rivularis*. FIG. 768. *F. walkeri*. FIG. 769. *Hebetancylus excentricus*. FIG. 770. *Laevapex diaphanus*. FIG. 771. *L. fuscus*. Measurement lines = 1 mm. Fig. 766 is from Walker (1925b); Fig. 768 is from Pilsbry & Ferriss (1907).

8(7) Shell depressed or moderately elevated, less than 4 mm long, rarely exceeding 3.5 mm, with or without a shelf-like septum across the posterior part of the aperture. When non-septate, the aperture is distinctly oval, wider anteriorly. When septate, the shell is evenly elliptical. Secondary growth may be present (Figs. 764, 765). Widely distributed in eastern United States in ditches and other small bodies of standing water, often temporary, and usually stagnant . . . . . *Ferrissia fragilis* (Tryon)

Shell to 6 mm long, usually depressed; aperture clearly oval, wider anteriorly, septum never present. Apex subacute, often far in the right posterior quadrant. Anterior and left slopes convex, posterior and right slopes concave (Fig. 768). Widely distributed, reported from Arkansas, Michigan and southern California on vegetation and debris in ponds . . . . . *Ferrissia walkeri* (Pilsbry & Ferriss)

9(4) Apex subacute, distinctly eccentric, to the right of the midline (Figs. 762, 769). Penis with a long glandular flagellum terminating in a bulbous tip; preputium without pigment. Tentacles colorless. In southern Florida, and perhaps Texas, in canals, etc. . . . . *Hebetancylus excentricus* (Morelet)

Apex very obtuse, almost in the midline of the shell. Penis without a flagellum; preputium flecked with pigment spots. Tentacles with a central core of black pigment. Principally east of the Mississippi in ponds and river backwaters; occasionally in streams in south-central states. Genus *Laevapex* 10

10(9) Shell ovate, smooth or with fine raised riblets usually on the anterior slope. Apex behind the center of the shell (Figs. 763, 771). Widely distributed in eastern North America in still water on submerged vegetation or debris, typically in the backwater areas of rivers or in lakes . . . . . *Laevapex fuscus* (Adams)

Shell subcircular, smooth, often encrusted with dark material. Apex about in the middle of the shell (Fig. 770). In slowly flowing streams, south-central and eastern states . . . . . *Laevapex diaphanus* (Haldeman)

SUPPLEMENTAL NOTES*

[1]The name Neritidae has been credited consistently to Rafinesque (1815). However, the family name he used was Neritinia [=Neritinidae] ("Les Neritines"). In this family he listed two subfamilies and a number of generic names, which included *Neritina* and *Nerita* (both under "Famille. Neritinia").

[2]The following figures are by John L. Tottenham: Figs. 21-80, 125, 128-142, 188, 189, 191-200, 222-234, 249-259, 290, 295-308, 319-344, 355-367, 391-414, 426-457, 468-527, 550-572, 581-702, 705-714, 716-724, 728-750, 759-763. Figs. 1-15, 17, 20, 81, 82, 528, 573-577, 704, 725-727, 751, 755, 757, 758, 764, 765, 767, 769-773, 775 were prepared by J.B. Burch. Most of the figures by Tottenham and by Burch are copyrighted in Burch & Tottenham (1980). The remaining figures were taken from various published sources, and in each case credit is given in the legends beneath the figures. Figs. 83-106, 109-124, 201, 204-218, 220, 221, 247, 260, 262-267, 269, 275, 276, 278-281 and 284 are used with permission of the University of Florida Press.

[3]Shells of the genus *Tulotoma* are unique among North American Viviparidae by their usual nodular appearance, and by their oblique apertures with concave margins (Fig. 772).

Only one species of *Tulotoma* is recognized here, *T. magnifica* (Conrad), although a second species, *T. angulata* (Lea), is occasionally recognized, as well as a third, *T. coosaensis* (Lea). A fourth species has been named, *T. bimonilifera* (Lea), but it is clearly a synonym of *T. magnifica*. According to Goodrich (1944b), *T. coosaensis* is the smooth upstream form; *T. angulata* is transitional between it and the tuberculate *T. magnifica*. Although in museum collections *T. angulata* seems to intergrade completely with *T. magnifica*, the relationship between the two nominal species may not be so simple. Patterson (1965) found *T. angulata* to have one pair of chromosomes more than Pollister & Pollister (1940, 1943) reported for *T. magnifica*.



FIG. 772. Viviparid shells. *Tulotoma* is on the left. a = the columellar axis; b = the plane of the aperture.

[4]Clench (1962a: 277-280) listed 49 names for *Campeloma*, 34 of which he considered as synonyms of the 14 names he did not synonymize (one species listed (*Paludina humerosa* Anthony 1860) is not a viviparid, but a pleurocerid). Although not claiming for them the status of species (or subspecies), the names Clench did not synonymize were *brevispirum* F.C. Baker 1928, *crassula* Rafinesque 1819, *decampi* 'Currier' Binney 1865, *decisa* Say 1816, *exilis* Anthony 1860, *floridense* Call 1886, *genicula* Conrad 1834, *gibba* Currier 1867, *integra* Say 1821, *leptum* Mattox 1940, *lima* Anthony 1860, *milesi* Lea 1863, *regularis* Lea 1841 and *tannum* Mattox 1940. Clarke (1973: 220) considered "*Campeloma leptum* and *C. tannum* [to] differ from *C. decisum* and *C. integrum* by trivial characters only. They are certainly not distinct species but are simply slightly aberrant populations of *C. integrum* (and probably of *C. decisum*)." Further, Clarke (*loc. cit.*) suggested that *C. integrum*, as well as *C. milesi*, are the same as *C. decisum*. *Campeloma parthenum* Vail 1979 may be merely the southeastern form of *C. decisum*.

[5]The name *Paludina integra* Say 1821 has been applied commonly to a viviparid (as *Campeloma integrum* (Say

*The comments in this section refer to the superscript numbers located at various places in the text.

1821)) and to a hydrobioid (as *Cincinnatia integra* (Say 1821)). The shell length given by Say ("length ¼ inch") is undoubtedly a typographical error (? for 1¼ inch), since in his description Say compared *integra* to [*Campeloma*] *decisum* (cf. Say, 1821; Binney, 1865d; F.C. Baker, 1928c; Clarke, 1973). "*Amnicola integra* (Say)" of authors is *Cincinnatia cincinnatiensis* (Anthony).

[6]Vanatta (1935) distinguished *Lioplax pilsbryi choctawhatchensis* "from the typical form [*L. pilsbryi* Walker] by being smaller, but with similar sculpture. It is about the size of *L. subcarinata* Say, but is without the two spiral angles on the last whorl. It is smaller than *L. s. occidentalis* Pils. *L. cyclostomatiformis* Lea is narrower and smoother."

[7]Of *Lioplax subcarinata* and *L. sulculosa*, Clench & Turner (1955: 10) said, "This eastern species [*L. subcarinata*] of *Lioplax* is exceedingly close in its relationship to *L. sulculosa* Menke, the western form. Their characters differ mainly in degree. The shells of *L. subcarinata* Say are usually somewhat thinner and are proportionately a little more attenuate. It appears also that *Lioplax subcarinata* Say on the average is somewhat smaller, though selected examples of the largest specimens of both species are about equal in size. In addition, the umbilical opening of *L. sulculosa* is much larger."

[8]*Probythinella lacustris* (F.C. Baker) has gone under the name *emarginata* Küster 1852 (*Paludina*), but the latter apparently was based on the name *Lymnaeus* [*Stagnicola*] *emarginatus* Say 1821 (a lymnaeid), even though Küster described and figured a hydrobioid species. Küster's hydrobioid species did not receive a valid name until F.C. Baker (1928c) described the subspecies *lacustris*. Morrison (1947b) designated *Probythinella lacustris limafodens* Morrison 1947 as type species of *Probythinella* Thiele 1928.

[9]According to Morrison (1940a), *Somatogyrus tryoni* Pilsbry & F.C. Baker 1927 and *S. virginicus* (Walker 1904) should be transferred to *Clappia*.

[10]According to Pilsbry (1934b), there is no difference in the shell between the American genus *Fluminicola* Stimpson 1865 and the European *Lithoglyphus* Hartman 1821 [type species: *Paludina naticoides* C. Pfeiffer 1828], the distinction between the two genera "being in the form of the verge." Pilsbry saw "no advantage in recognizing *Lithoglyphus* in America, since its presence does not seem demonstrable" [at that time]. Taylor (1966a,b) combined the two genera, mentioning having examined the verge in most American species, but as yet none of the anatomical data have been published. Until it is shown conclusively that the European and American species are indeed congeneric, it would seem best to retain the well-known American name, *Fluminicola*.

[11]F.C. Baker (1928c), H.B. Baker (1964) and La Rocque (1968) placed Say's (1829) *Melania integra* in the genus *Somatogyrus*.

[12]*Somatogyrus virginicus* Walker is placed in the subgenus *Walkerilla* following Thompson (1969).

[13]*Pyrgulopsis letsoni* (Walker), *P. ozarkensis* Hinkley, *P. scalariformis* (Wolf) and *P. wabashensis* Hinkley may belong to the genus *Marstonia* (Thompson, 1977).

[14]*Marstonia lustrica* (Pilsbry 1890) is used here in the expectation that the International Commission on Zoological Nomenclature will rule favorably on H.B. Baker's (1960c) petition to suppress Say's (1821) *Paludina lustrica*. Otherwise, *Amnicola lustrica* Pilsbry 1890 is preoccupied by *Amnicola lustrica* (Say 1821), should the latter (actually a *nomen dubium*) be considered a member of the genus *Amnicola*.

[15]*Amnicola* Gould & Haldeman, as listed in Haldeman (1840, p. 3 and on inside back cover), has as its type species (by subsequent designation by Haldeman, 1840) *Paludina lustrica* Say 1821. Gould (1841a) gave the first detailed description of the genus and of the species *Amnicola porata* (Say 1821), and mentioned as included in the genus *Paludina limosa* Say 1817 and *Paludina lustrica* Say 1821, although the latter species was considered as only doubtfully belonging to *Amnicola* (H.B. Baker, 1960c). Later, Haldeman (1845) accepted *Amnicola* as described by Gould (1841a) (not as listed by Haldeman (1840)), placed the previous "*Amnicola lustrica*" Haldeman (not of Say) in the synonymy of *Amnicola limosa* (Say) and recognized *Amnicola lustrica* (Say) as a distinct species "closely allied to *A. lapidaria* [*Pomatiopsis lapidaria* (Say 1817)], of which it may possibly be the young." A year later, Herrmannsen (1846) designated Say's (1821) *Paludina porata* as type species for *Amnicola* Gould 1841, apparently being unaware of Haldeman's (1840) earlier introduction of *Amnicola* and designation of *Paludina lustrica* Say as its type species.

Since 1846, *Amnicola* has been used almost entirely as though *A. porata* (Say) were its type species, although, in fact, *A. lustrica*, which is a *nomen dubium* (and a *nomen oblitum* as well, according to H.B. Baker (1964)), is the validly designated type species. Unfortunately, the type specimen of Say's *Paludina lustrica* has been lost. To end the nomenclatoral controversy which surrounds *Amnicola*, H.B. Baker (1960c) requested that the International Commission on Zoological Nomenclature use its plenary powers to suppress the specific name *Paludina lustrica* Say 1821, and to place *Amnicola* Gould & Haldeman 1840, with *Paludina porata* Say 1821 as its type species, on the Official List of Generic Names in Zoology. (The Commission has not yet made a decision on this request.) Subsequently, Clarke (1973) selected a "neotype" for *Paludina lustrica* Say, which is also the same specimen H.B. Baker (1964) designated as the lectotype of *Amnicola walkeri* Pilsbry 1898. In my 1978 outline, I followed that system. Although such a procedure would provide a belated identity for "*Amnicola lustrica* (Say)" (i.e., it then would be the same as *A. walkeri*), apparently *A. walkeri* actually belongs to the subgenus *Lyogyrus* Gill 1863 (see Thompson, 1968), not to *Amnicola* s.s. as it has been perceived for some 130 years. Thus, *Lyogyrus* would become a junior subjective synonym of *Amnicola* s.s. and would contain the group of *A. walkeri/pupoidea*, and the group of *A. porata/limosa* would be left without a subgeneric name (unless the European *Marstoniopsis* should be shown to be congeneric). Therefore, it seems best to retain the customary concept of *Amnicola* (with *Paludina porata* Say 1821 as type species) in hope that the International Commission on Zoological Nomenclature will adopt H.B. Baker's (1960c) proposal. F.G. Thompson, who has done the most intensive recent work on North American Hydrobiidae (Thompson, 1968, 1969, 1977, 1979) has written (1974) in support of Baker's proposal.

[16]The reproductive anatomy has not been described, to my knowledge, of *aldrichi* Call & Beecher (and its subspecies), *bakeriana* Pilsbry, *clarkei* Pilsbry, *decisa* Haldeman, *missouriensis* Pilsbry and *proserpina* Hubricht, so their placement in the genus *Amnicola* is presumptive. Subsequent studies may alter the generic placement of these species.

[17]From drift debris of the Guadalupe River near New Braunfels, Texas, Pilsbry & Ferriss (1906) named "*Valvata*" *micra* and "*Valvata*" *micra nugax*, mentioning that they might prove to be "amnicoloid" snails comparable to *Horatia* Bourguignat or *Daudebardiella* Boettger in the Palaearctic fauna. Pilsbry (1916d) referred *micra* and *nugax* to the subgenus *Hauffenia* of the genus *Horatia*. Bole (1970) raised *Hauffenia* to the status of an independent genus, although still close to *Horatia*. Taylor (1975) placed *micra* in the genus *Hauffenia* and *nugax* in the genus *Horatia*. Hubricht (1940b) reported finding specimens of "*Horatia*" in an artesian well at the U.S. fish hatchery at San Marcos, Texas, and in a subterranean stream in Manitou Cave, near Fort Payne, Alabama.

[18]*Fontigens binneyana* (=*obtusa* Lea 1841 (*Paludina*), preoccupied by *Paludina obtusa* Troschel 1837) may prove to be a synonym of *Fontigens nickliniana* (Lea 1838).

[19]*Fontigens weberi* may be extinct. "*Fontigens weberi* was described as a recent species from a 'bone' specimen from West Lake, Everglades National Park. This species does not occur in the region at present, although shells of this species are common in Pliocene road fill near the lake" (Thompson, 1968: 12).

[20]Hubricht (1960) believes that *Pomatiopsis hinkleyi* Pilsbry is only a wet habitat form of *P. lapidaria* (Say).

[21]Following H.B. Baker (1963), I (1978, 1979) previously utilized the family name Paludomidae Gill 1871 instead of the recently commonly used Pleuroceridae Fischer 1885, the previously commonly used Strepomatidae Haldeman 1863 (based on an invalid manuscript name of Rafinesque), or Pachychilidae Troschel 1857 ("Of the 5 familial names prior to Pleuroceridae Fischer, 1885, all apparently are 'nomina oblita' except Paludominae Gill, 1871, which was used by Pilsbry as late as 1956"). In spite of its illegal or at least questionable nomenclatural status, Starobogatov (1970) used Pachychilidae Troschel, with Ceriphasiidae Gill 1863 and Pleuroceridae Fischer listed as synonyms, for all the North American pleurocerids (*Elimia, Gyrotoma, Io, Juga, Mudalia, Pleurocera,* etc.). The family name Pachychilidae is based on the Middle American *Pachychilus*. Starobogatov restricted the Paludomidae to Afro-Asian genera. On the other hand, Morrison (1954) placed the Asiatic *Paludomus* with the pleurocerids.

However, in spite of the above nomenclatural activity, there are as yet no really solid bases for adequately comparing *Pleurocera* and its allies with *Paludomus* and its related taxa or *Pachychilus* and its relatives. Until the necessary comparative studies have been completed and evaluated, perhaps it is best to retain the family name Pleuroceridae.

[22]A critical revision of the pleurocerids has not yet been made. The generic groups used here are based on classical shell characters, even though it is realized that these characters mostly seem to intergrade at one point or another.

Animal characteristics of value in pleurocerid systematics are currently so incompletely known that they cannot be used to precisely characterize biological generic groups or to assign the great majority of species to definite nomenclatural generic groups. *Pleurocera* is used as though *P. acuta* were its type species, in the expectation that the International Commission on Zoological Nomenclature will adopt the long-standing petition to preserve this usage. The identity of *Ellipstoma gibbosa* Rafinesque 1818 is too doubtful to give nomenclatural validity to *Ellipstoma* Rafinesque 1818.

[23]*Elimia* H. & A. Adams 1854 (type species *Melania acutocarinata* Lea 1841 = *Melania clavaeformis* Lea 1841) is used in place of its better known synonym *Goniobasis* Lea 1862 (type species *Goniobasis osculata* Lea 1862).

The classification in the genus *Elimia* presented here, and the distribution of the various recognized species and subspecies, is that of Goodrich (1930a, 1936, 1939d,e, 1940d, 1941a,b,c, 1942b, 1944d, 1945, 1950). No attempt has been made to assess the taxonomic validity of the species and subspecies.

[24]*Elimia perstriata decampi* (Lea) is "possibly only an aberrant form" (Goodrich, 1940d: 16).

[25]*Goniobasis* (=*Elimia*) *pilsbryi* Goodrich is a replacement name for *Melania* (=*Elimia*) *showalteri* Lea 1861, which is not *Lithasia* (=*Elimia*) *showalteri* Lea 1860.

[26]Goodrich (1941c: 20) said that *Elimia ampla* (Anthony) "may simply be an enlarged and conic phase of the [*E.*] *clara* of the transition zone."

[27]Goodrich (1944d: 44) thought that *Elimia ornata* (Lea) is probably a hybrid of *E. gerhardti* (Lea) and *E. caelatura* (Conrad).

[28]The genus *Gyrotoma* is now undoubtedly extinct, due to the biological destruction of the Coosa River. Goodrich (1924a) recognized 13 species in the genus, which he placed into five species groups. However, later (1944d: 46, 47) Goodrich was less certain about this arrangement. "In a study of this genus in 1924 with the unexampled H.H. Smith collections as a basis, the shape and depth of the sutural fissure were relied upon for differentiation among the species. The writer is not so sure, after twenty years, that the thirteen species then recognized by this standard are actually good species. For one thing, the range of the whole genus is only about one hundred and twenty miles of river. The habitats are shoals and reefs over which the currents are heavy. In all the forms, the operculum is large, thick and leatherly, the spiral lines nearly obsolete. The radulae, too, are alike. Considering how greatly a given species of *Goniobasis* may vary, and a member of *Pleurocera* more so, it is reasonable to suppose that variation in *Gyrotoma*, including its fissure, may be greater than was supposed in 1924. But in the absence of better information on the subject, the species are listed here as they were then recognized."

In general, I have disregarded the depth of the sutural fissure as a taxonomic character in *Gyrotoma*. Of the 13 species recognized by Goodrich, I have included six in the key: *G. excisum* (Lea), *G. lewisi* (Lea), *G. pagodum* (Lea), *G. pumilum* (Lea), *G. pyramidatum* Shuttleworth and *G. walkeri* Smith. *Gyrotoma hendersoni* Smith, which has a shallow fissure, is placed in the synonymy of *G. pumilum* (Lea), which has a deep fissure. *Gyrotoma alabamensis* (Lea), *G. amplum* (Anthony), *G. cariniferum* (Anthony), *G. incisum* (Lea), *G. laciniatum* (Lea) and *G. spillmani* (Lea) are placed in the synonymy of *G. excisum* (Lea). *Gyrotoma excisum* have deep sutural fissures, as do *G. alabamensis*, *G. cariniferum* and *G. laciniatum*. *Gyrotoma amplum*, *G. incisum* and *G. spillmani* have shallow fissures. These nominal species, here placed in synonymy, are illustrated in Figs. 435-440, 445.

Distributions (all in the Coosa River basin of Alabama) given by Goodrich (1944d) for *Gyrotoma* species are as follows:

G. alabamensis, Peckerwood Shoals, Talladega County, to Duncan's Riffle, Chilton County;
G. amplum, Talladega to Coosa County;
G. cariniferum, confined to a reef at Fort William Shoals, Talladega County, in swift water;
G. excisum, Three Island Shoals, Talladega County, to Wetumpka;
G. hendersoni, Fort William Shoals only;
G. incisum, Weduska Shoals to Wetumpka;
G. laciniatum, Fort William Shoals to Wetumpka;
G. lewisi, confined to two shoals of Talladega County;
G. pagodum, a lower river form; The Bar, Chilton County, to Wetumpka, Elmore County;
G. pumilum, Weduska Shoals, Shelby County, to Wetumpka;
G. pyramidatum, Ten Island Shoals, St. Clair County, to the mouth of Yellowleaf Creek, Shelby County ("the

first of the genus to appear in the river");
G. spillmani, known only from two shoals of Talladega County;
G. walkeri, Weduska Shoals to Butting Ram Shoals, Coosa County, a range of only a few miles.

[29]Displacing the well-described and well-known *Gyrotoma* Shuttleworth 1845 by the obscure and long forgotten "*Apella* 'Mighels' Anthony 1843" (e.g., see Turner, 1946; Clench, 1959a; Davis, 1977) would certainly be an injustice. *Apella* entered the literature in a sentence in a published (1843) letter from J.G. Anthony as follows. "I have, within two months past, received one species of this genus ["*Melatoma* Swainson"*] from Dr. Mighels, of Portland, Maine, under the name of '*Apella scissura*'." In 1860, after rejecting *Melatoma* as pertaining to a North American freshwater snail, Anthony stated, "In 1841 or 1842, Dr. J.W. Mighels sent me specimens of one species under the name of *Apella scissura*; but his generic name was never published, and his species, if not identical with any which Mr. Lea afterwards described seems to have been overlooked and forgotten." Anthony then adopted Shuttleworth's name *Gyrotoma*, which has been the recognized name (with the exceptions of the use of Lea's preoccupied *Schizostoma*, and of Turner's, Clench's and Davis' use of *Apella*) for the past 118 years. *Apella scissura* was and is still both a *nomen nudum* and a *nomen dubium*.

[30]*Io fluvialis* (Say) is the largest of the North American Pleuroceridae. It varies in shell form from the smooth *fluvialis* described by Say (1825), to spinose forms such as *spinosa* Lea and *turrita* Anthony. C.C. Adams (1915) treated admirably the monotypic genus *Io* and its geographic variation. He recognized 14 races or population forms of *I. fluvialis*: *angitremoides* C.C. Adams, *brevis* Anthony, *clinchensis* C.C. Adams, *fluvialis* Say, *loudonensis* C.C. Adams, *lyttonensis* C.C. Adams, *nolichuckyensis* C.C. Adams, *paulensis* C.C. Adams, *powellensis* C.C. Adams, *recta* Reeve, *spinosa* Lea, *turrita* Anthony[31], *unakensis* C.C. Adams and *verrucosa* Reeve. Several of these forms are illustrated on p. 136 (from Tryon, 1873b).

[31]*Io fluvialis* form *turrita* Anthony was reported (Clench, 1928) in the Little River, but this "purported finding has not been verified" (Goodrich, 1940d).

[32]*Leptoxis* s.s. of the Alabama river drainage is a variable group. Goodrich's (1922) monograph of them was one of his earliest publications on the Pleuroceridae. In it, clear-cut differences between most of the recognized taxa are not clearly expressed. Later (1941b, 1944d), Goodrich revised slightly his earlier concepts regarding a few of the species, but it would seem that he still recognized too many taxa. However, the Alabama *Leptoxis*, mostly confined to the Coosa river drainage, are undoubtedly now largely extinct, due to degradation of their habitats.

[33]*Leptoxis lirata* may be only a form of *L. showalteri* (Goodrich, 1944d).

[34]In shell characters, especially the nodulose shoulders, *Leptoxis crassa* seems closer to *Lithasia* s.s., and that is where I placed it in my 1979 list (Burch, 1979). However, in this manual *L. crassa* is placed with *Leptoxis* on radular characters (cf. Goodrich, 1931a, 1932d). *Leptoxis crassa* and its form *anthonyi* commonly have been assigned to the genus *Eurycaelon* on the belief that *anthonyi* was its type species. However, as pointed out by Morrison (1971), Neville (1885) designated *Goniobasis umbonata* Lea 1864 (=*Anculosa (Lithasia) geniculata* Haldeman 1840, *fide* Goodrich (1940d), Morrison (1971)) as the type species of *Eurycaelon*, which makes *Eurycaelon* a synonym of *Lithasia*. Morrison (1971) proposed *Athearnia* (type species *Anculosa anthonyi* Redfield 1854) as a replacement name, and this taxon is used here as one of the three subgenera of *Leptoxis*.

Although *Leptoxis crassa anthonyi* is given in the list of species (p. 44) as though it were a subspecies of *L. crassa*, it may not deserve such nomenclatural status. *Leptoxis crassa* s.s. is probably only a localized race or form (in much the same sense as those of *Io*, cf. C.C. Adams, 1915) of a much larger complex which customarily has gone under the nomenclaturally junior name *anthonyi*. In *L. crassa*, the lumpiness of the shoulders is strongly emphasized, becoming strong, well-developed tubercles. In *L. anthonyi*, the spire is generally not so depressed as in *crassa*, and the shoulder is often absent or not prominent and is commonly smooth or with only slight undulations. In both forms, the lower columella terminates in a flange.

[35]On shell characters, *Lithasia obovata* would seem to belong more naturally to the *Elimia/Pleurocera* group, and *L. geniculata pinguis* to *Leptoxis (Mudalia)*. However, these two species are placed with *Lithasia* because of their radular characters.

**Melatoma* Swainson 1840 is not the same as *Melatoma* Anthony 1843 (Gray, 1847; Anthony, 1860).

[36]The variability seen in *Lithasia salebrosa* (Conrad) would seem to include *L. geniculata* (Haldeman). Goodrich (1940d) separated the two, but (in 1941f) remarked that "the distinction between *geniculata* of the Cumberland River system and *salebrosa* of that of the Tennessee River is chiefly that the latter commonly has two or more rows of nodules." Specimens of *salebrosa* with but a single row of nodules do occur, but are not common. These have the conchological characters of *geniculata*. Several specimens of the single lot labelled "*Lithasia salebrosa*" from the "lower Cumberland River, Tennessee" in the Museum of Zoology collections (UMMZ 132477) have only a hint of a second row of nodules. The other specimens in this lot have only a single row at the shoulder of the whorls. Basically, they are *L. geniculata*.

Davis (1974) treated *Lithasia salebrosa* and *L. geniculata* as separate species, and listed the distribution of "*Io*" *salebrosa* as the Cumberland River and Caney Fork, and the Duck and Tennessee rivers. Goodrich (1940d) did not include the middle and upper Cumberland River, Caney Fork or the Duck River in the distribution of *L. salebrosa*; he reported *L. geniculata* in these streams. According to Davis (1974), "The one population found in the Duck River is not pure *salebrosa* as given in Fig. 45 by Tryon (1873). Two individuals were found in a population of over 200 snails where specimens reflected genetic mixtures of *geniculata, fuliginosa, geniculata* x *fuliginosa, fuliginosa* x *duttoniana*. Pure *salebrosa* is probably extinct."

According to Tryon (1873b), "Generally but one row of tubercles is developed on this species [*L. geniculata*], but occasionally a second and less prominent row is visible. The whorls are more shouldered, and the tubercles larger and less numerous than in *L. salebrosa*, Conrad. . . . Mr. Lea considers *geniculata* to be the same as *salebrosa*."

Curiously, some specimens of *Lithasia salebrosa* seem little different from *L. verrucosa* (Rafinesque). Further, *L. salebrosa subglobosa* (Lea) and some specimens of *L. geniculata* differ but little from *Leptoxis (Athearnia) crassa* (Haldeman), the latter also a species of the Tennessee river drainage. [Because of this close similarity, I (1979) previously included *Athearnia* in the synonymy of *Lithasia*.] The essential conchological difference separating *Leptoxis (Athearnia) crassa* from the *Lithasia salebrosa-geniculata* complex is the flange of the lower columellar lip of the aperture, perhaps a character of dubious generic value.

Davis (1974) treated *pinguis* Lea and *fuliginosa* Lea as headwaters and small rivers forms respectively of *geniculata*. Goodrich (1934a, 1941f) also discussed variation in this complex of races and forms. "*Lithasia geniculata* and *salebrosa* each has upstream or side-stream forms, distinguished by an elongation of the spire and an alteration of proportions of altitude to diameter, together with the curious characteristic of a development of nodulous sculpture, when that exists, at the periphery of the shell and not at the shoulder" (Goodrich, 1941f).

[37]The classification in the genus *Pleurocera* presented here, and the distribution of the various recognized species and subspecies, is that of Goodrich (1917, 1924b, 1927, 1928a,b, 1929b, 1930a, 1934c, 1935b, 1936, 1939d,e, 1940d, 1941b,c, 1942b, 1944d). No attempt has been made to assess the taxonomic validity of the species and subspecies.

[38]According to Goodrich (1940d), *Pleurocera currierianum* (Lea) is possibly only a depauperate form of *P. brumbyi* (Lea).

[39]Goodrich (1940d) thought that *Pleurocera viridulum* (Anthony) might be only a fast water modification of *P. pyrenellum*.

[40]The genus *Lymnaea* Lamarck 1799 has been used variously to include nearly all members of the Lymnaeidae (e.g., see Hubendick, 1951; Walter, 1969; Harman & Berg, 1971) or only *Lymnaea stagnalis*, its varieties, and several very closely related species (e.g., F.C. Baker, 1928c; Burch, 1979). In this latter system, the family contains a number of species groups (genera) equal in rank to *Lymnaea* s.s. A third system, more or less a compromise between the previous two, uses *Lymnaea* as a large inclusive genus, but recognizes various subgeneric groups within it. These subgenera correspond to the genera of the F.C. Baker scheme. As a convenience for species-group separation, the less conservative scheme is used here. Aside from convenience, there is some scientific justification for handling the lymnaeids in this fashion (Burch, Lindsay & LoVerde, 1971; Burch & Lindsay, 1973a).

[41]*Fossaria* Westerlund 1885 is used for the group of small lymnaeids rather than *Galba* auct. (which is only doubtfully the same as *Galba* Schrank 1803, type species *Galba pusilla* Schrank 1803 by monotypy; see Hesse, 1923; Pilsbry & Bequaert, 1927; F.C. Baker, 1928c; Clarke, 1973).

[42]The genus *Stagnicola* Leach (in Jeffreys) 1830 is based on the European *Buccinum palustre* Müller 1774. The work of Jackiewicz (1959) has shown that several distinct species have masqueraded under the name *palustris*. Just which anatomical type is represented by Müller's species is not known, and until that is settled, and it is determined

that such a species does indeed occur in North America, then it seems advisable not to use *S. palustris* here but the first name applied specifically to a North American *palustris*-like snail instead, i.e., Say's (1821) *Lymneus elodes.*

[43]The largest group of Lymnaeidae in North America are the stagnicoline lymnaeids, members of the genus *Stagnicola*. Their taxonomy, based largely on shell shape, has always been troublesome. Conditions of the water in which stagnicoline snails live can have some influence on the exact shape of their shells (ecophenotypic variation), whole populations exhibiting the abnormal characters when they occur. However, other cases of constant population differences seem to be due to small genetic differences between populations. The great problem in systematics of stagnicoline snails is in accurately assessing which characters are ecophenotypic and which are genetic, and of the genetic differences which are great enough to conclude that any particular population(s) is (are) distinct enough to deserve a binomial (or trinomial) name of its (their) own. Since there have been almost no experimental breeding studies to evaluate the taxonomic importance of any shell characters in *Stagnicola*, schemes for classifying the genus have all been quite subjective. Accordingly, systematic interpretations have varied widely, from the "splitters" to the "lumpers".

In reviewing North American *Stagnicola*, it seems to me that they fall into two general groups, the *Stagnicola elodes* group and the *Stagnicola catascopium/emarginata* group. Typically, species of the *Stagnicola elodes* group have an elongated, rather narrow, brown shell, and are inhabitants of quiet standing waters, such as ponds, pools, ditches, marshes, swamps, etc. The *Stagnicola catascopium/emarginata* group typically have compressed spires and subglobose body whorls, broader, light-colored shells, and are inhabitants of rivers and lakes.

Because of the fundamental uncertainties of their taxonomy, it is not easy to decide on a nomenclatural scheme for the stagnicolas. The one adopted here reflects a rather conservative approach.

[44]Hubendick (1951) recognized a separate subfamily, the Lancinae, for the limpet-shaped *Lanx*, in contrast to the subfamily Lymnaeinae, which included all other lymnaeids. However, whether or not a patelliform shell in the Lymnaeidae is, *per se*, enough to warrant the recognition of a subfamily, or whether sets of peculiar anatomical characteristics not related or only partially related to shell shape will eventually define subfamilies is not known at present. Walter (cf. 1969) mentioned certain close anatomical similarities of *Lanx* to "*Lymnaea catascopium* Say" (=*Stagnicola emarginata serrata* Haldeman). However, the use of anatomical characters for showing relationships in the Lymnaeidae needs to be reassessed (cf. Burch, Lindsay & LoVerde, 1971).

[45]It may not be worthwhile to distinguish between *Fisherola nuttalli nuttalli, F. nuttalli kootaniensis* and *F. nuttalli lancides*, but a more detailed study of *Fisherola* is needed to decide this. "*Fisherola lancides* is another subspecies of the Snake River, in which the apex is a little more anterior, but some of the original lot before me run close to *nuttalli*" (Pilsbry, 1925a). In describing *Fisherola lancides*, Hannibal (1912) gave the locality as "Snake River (H. Hemphill)." According to Henderson (1936c), "The Spokane River specimens obtained by Hemphill are doubtless the ones afterwards described from his specimens as *lancides*."

[46]Classification of the Physidae follows Te (1978). Subsequent to the preparation of this list, Te (1980) listed an "unnamed species" of *Physella (Physella)*, an "unnamed species" of *Physella (Costatella)*, an "unnamed subspecies" of *Physella (Physella) ancillaria* (Say 1825), an "unnamed morph" of *Physella (Costatella) osculans* (Haldeman 1841), and introduced as a subspecies of *Physella (Costatella) hendersoni* (Clench 1925) the *nomen nudum floridana* "Pilsbry ms."

[47]Species of the genus *Stenophysa* Martens 1898, native to Central America and Mexico, have been found in Texas (Te, 1978).

[48]The validity of *Gyraulus (Torquis) hornensis* is open to some doubt. It was named by F.C. Baker (1934e) for specimens that he had earlier (e.g., 1928c) called *Gyraulus arcticus* Beck (in Möller) 1842. Clarke (1973) placed *hornensis* in the synonymy of *G. deflectus.*

[49]If *Drepanotrema* and the Brazilian *Acrorbis* Odhner 1937 (type species: *Acrorbis petricola* Odhner 1937) are shown conclusively to belong to the same tribe, then apparently the earliest name for this taxon is Acrorbini Starobogatov 1958, predating Zilch's (1959) Drepanotremeae and Harry's (1962) Drepanotrematinae. (Starobogatov placed *Drepanotrema* in his Acrorbini (Starobogatov, 1970), and Harry (1962) placed *Acrorbis* in his Drepanotrematinae, but Zilch (1959) placed (questionably) *Acrorbis* in the tribe Segmentineae.)

[50]*Helisoma anceps* (Menke) exhibits considerable variation over its wide range, which has resulted in many varietal

names. An alphabetical list of names assigned to *Helisoma* s.s., with type localities in parentheses, follows: *anceps* Menke 1830 (Virginia), *anguistoma* Haldeman 1844 (no locality given, not figured, and specimen lost), *angulata* Rackett 1821 [preoccupied] (near Lake Huron), *angulatum* Wood 1828 [preoccupied] (from Haldeman (1844); I have not seen this reference), *antrosum* Conrad 1834 (Randon's Creek, near Clairborne, Alabama) (=*anceps* Menke 1830), *aroostookense* Pilsbry 1895 (East branch of Salmon Brook, Woodland, Aroostook County, Maine), *bartschi* F.C. Baker 1945 (Brook at Great Falls, Virginia), *bicarinatum* Say 1817 [preoccupied] (Delaware River), *cahni* F.C. Baker 1927 (Big Muskallonge Lake, Vilas County, Wisconsin), *corrugatum* "Currier" Walker 1909 (Perch Lake, Kent County, Michigan), *engonatum* Conrad 1835 (Albany, New York), *idahoense* F.C. Baker 1945 (Pend Oreille River, Sand Point, Idaho), *jordanense* Winslow 1823 (South Arm of Pine Lake, about two miles north of East Jordan, Charlevoix County, Michigan), *latchfordi* Pilsbry 1927 (Meach's Lake, Hull, Quebec), *major* Walker 1893 [preoccupied, renamed *percarinatum* Walker 1909] (Crystal Lake, Benzie County, Michigan), *minnesotense* F.C. Baker 1927 (Frontenac, Minnesota), *percarinatum* Walker 1909 [new name for *major* Walker 1909] (Crystal Lake, Benzie County, Michigan), *politum* F.C. Baker 1945 (Honeywell Creek, Carleton County, Ontario, Canada), *portagensis* F.C. Baker 1908 (Portage Lake, on Fish River, Aroostook County, Maine), *royalense* Walker 1909 (Siskowit Lake, Isle Royale, Lake Superior, Michigan), *rushi* F.C. Baker 1939 (Toad Island, Georgian Bay, Ontario, Canada), *sayi* F.C. Baker 1928 (Tomahawk Lake, Oneida County, Wisconsin), *shellense* F.C. Baker 1927 (Shell Lake, Washburn County, Wisconsin), *striatum* F.C. Baker 1902 (Pleistocene fossil from sewer excavation, eight feet below the surface of the ground, Cold Spring Park, Milwaukee, Wisconsin), *unicarinatum* Haldeman 1844 (Schuylkill River).

In Canada, Clarke (1973) recognized Walker's (1909e) variety *royalense* as a valid subspecies of *Helisoma anceps*. He considered *H. anceps rushi* F.C. Baker to be a synonym of *H. a. royalense*. Clarke considered ten other "subspecies" of *H. anceps* recorded from Canada: *anticostianum* F.C. Baker 1945 (a Pleistocene fossil), *aroostookense, cahni, latchfordi, percarinatum, politum, portagensis, sayi, striatum* and *unicarinatum*. He concluded (p. 443) that "it is probable that most of the 'subspecies' currently recognized [in the Canadian Interior Basin] are not geographically distinct and are taxonomically invalid but firm decisions on this must be deferred until analysis of more populations, including topotype populations can be made."

Helisoma eucosmium (Bartsch 1908) may be simply a form or juvenile of *H. anceps anceps* (Menke).

[51]The generic name *Carinifex* was first presented by Binney (1863), in combination with Lea's (1858a) *newberryi* (*Planorbis*), as a name without description in a pamphlet ("Smithsonian Miscellaneous Collection 000") containing a catalogue of North American Pulmonata. In 1865b, c, Binney described the genus and figured for the first time Lea's species *Carinifex newberryi*. In 1864c, Lea "provisionally" introduced the generic name *Megasystropha* for *newberryi*. The International Commission on Zoological Nomenclature in Opinion 432 [1956] suppressed the generic names *Carinifex* Binney 1863 and *Megasystropha* Lea 1864 in favor of *Carinifex* Binney 1865. *Carinifex* has been used for many years as a generic name for the *newberryi* group of North American planorbids. In subordinating it as a subgenus of *Helisoma*, I am following Henderson (1931b) and D.W. Taylor (1966a).

Whether there are more than one species of *Carinifex* is doubtful. "This [*Helisoma (Carinifex) newberryi*] has long been known as a very protean species, but conchologists have not been inclined to establish varietal names, as the variations are very numerous and intergrade thoroughly. If one begins naming them it is difficult to see where any lines may be satisfactorily drawn. It is doubtful whether the variations can be properly called even mutations. The variation is chiefly in the amount of elevation of the spire above the last whorl and a marked tendency toward scalariformity, with inevitable effect upon the general shape of the shell, and upon the width of the last whorl and of the umbilicus. The variation is so great and the gradation so minute that it is almost impossible to determine just what should be the normal form" (Henderson, 1931b). "I am disposed to look upon all of the described species and varieties of *Carinifex* as subspecies of a widely spread stock . . . " (Pilsbry, 1934a).

[52]Ten nominal species or subspecies are associated with *Menetus* s.s. in addition to its type species, *M. opercularis* Gould 1847. Two of the names are replacements for preoccupied names, *multilineatus* Vanatta 1899 for *oregonensis* Vanatta 1895 (*non oregonensis* Tryon 1865) and *cooperi* F.C. Baker 1940 for *planulatus* J.G. Cooper (in W. Cooper) 1859 (*non planulatus* Deshayes 1824). The other six names are *callioglyptus* Vanatta 1895, *centervillensis* Tryon 1871, *crassilabris* F.C. Baker 1945, *labiatus* F.C. Baker 1945, *planospirus* F.C. Baker 1945 and *portlandensis* F.C. Baker 1945. Whether any of these are more than forms or synonyms of *opercularis* is not presently known. The subgenus needs critical study. Dall (1905) was of the opinion that there was only one species, and, from my own limited observations, I agree. "The sculpture [of *M. opercularis*] is like that of [*Promenetus*] *exacuous*, the spiral sculpture being faint and sometimes absent in southern specimens, and tending to be emphasized in northern ones. As a rule the margin of the aperture is not thickened except in young specimens which have been overtaken by drought or winter before maturity. The keel is generally, but not always, present in southern shells, but those from Oregon and northward show a tendency to form a shell either without a noticable keel, or with the keel forming a margin to

a plane upper surface, rather than a median carina. When compared with Cooper's types in the National Museum Mr. Vanatta's *P.* ["*Planorbis*"] *callioglyptus* is seen to be identical. The variety *oregonensis* retains the typical form but has stronger spiral sculpture. I regard *P. centervillensis* of Tryon as a *P. planulatus* with the keel obsolete. What appear to be intergradational forms are numerous in the large series in the National Museum; though it would seem incredible to any one possessing only the extremes that they can belong to the same species" (Dall, 1905: 93).

[53]F.C. Baker (1945) said the following about his subgenus *Micromenetus*. "The group here separated as *Micromenetus* differs from typical *Menetus* in the size of the shell which is always much smaller, none exceeding 4 mm. in diameter. The form of the shell is lenticular and there is usually a peripheral carina more or less well developed. The penial gland has a duct which is almost three times as long as the gland and is attached to the inner wall of the preputium for the greater part of its length . . . In typical *Menetus*, this duct is short and enters the diaphragm directly without being attached to the wall of the preputium . . . The pseudobranch in *Micromenetus* is also very long and narrow while in typical *Menetus* it is short and wide . . .These are small differences, perhaps, but they appear constant. *Micromenetus* differs from both *Promenetus* and *Planorbula* in the shape of the penial gland. As far as examined the radulae of the two groups differ in formulae, that of *Menetus* being 20-1-20 while in *Micromenetus* it is 15-1-15."

[54]If it turns out that the eastern subgenus *Micromenetus* is represented by only one variable species, *M. dilatatus* (i.e., if the nominal species *M. brogniartianus* and *M. sampsoni* fall within the normal variation of *M. dilatatus*), as the western *Menetus* s.s. is represented by only the variable *M. opercularis*, then separating the two species each into a separate subgenus does not seem justified.

[55]How many species to recognize in the subgenus *Micromenetus* is difficult to decide without an intensive study of the group. Eight names for Recent planorbids are associated with the subgenus. *Menetus dilatatus* (Gould 1841) is the type species. Other names are *alabamensis* Pilsbry 1895, *brogniartianus* Lea 1842, *buchanensis* Lea 1841, *floridensis* F.C. Baker 1945, *lens* Lea 1838, *lenticularis* Lea 1844, *pennsylvanicus* Pilsbry 1916 and *sampsoni* 'Ancey' Sampson 1885. F.C. Baker (1945) listed *buchanensis, floridensis* and *pennsylvanicus* as subspecies of *M. dilatatus*. However, whether these are true subspecies or simply forms or synonyms is not known, but judging from Baker's 'splitting' in other groups they probably do not justify recognition by latinized names. Lea's *brogniartianus* and Pilsbry's *alabamensis* are both carinate forms at present not separable by their descriptions. They may prove to be only variations of *dilatatus*. From the specimens that I have observed, *M. sampsoni* differs from *dilatatus* by its rounder, less flared aperture and wider, shallower umbilicus. Whether or not these are constant characters is not known at present. Lea's *lens* (preoccupied) and *lenticularis* are synonyms of *brogniartianus*.

[56]Twelve Recent nominal specific or subspecific names are associated with *Planorbella* s.s.: *bella* Lea 1841, *campanulata* Say 1821, *canadensis* F.C. Baker & Cahn 1931, *collinsi* F.C. Baker 1939, *davisi* Winslow 1926, *michiganensis* F.C. Baker 1927, *minor* Dunker 1850, *multivolvis* Case 1847, *rideauensis* F.C. Baker 1945, *rudentis* Dall 1905, *smithi* F.C. Baker 1912 and *wisconsinensis* Winslow 1926. Say's *campanulata* is the type species of *Planorbella*. Binney (1865c) inspected Lea's type specimen of *bella*, an immature shell, and placed it in the synonymy of *P. campanulata*. F.C. Baker (1928c) synonymized *minor* with *P. campanulata* s.s. Clarke (1973) placed *canadensis, davisi, rudentis* and *wisconsinensis* in the synonymy of *P. campanulata* s.s.

[57]Dall (1905) proposed the section *Haldemanina* for Lea's (1858) *Planorbis wheatleyi*. F.C. Baker (1945) was "disposed to accept *Haldemanina* as a subgroup under *Planorbula*," and this arrangement was followed by Zilch (1959). However, Pilsbry & Ferriss (1906) considered *Haldemanina* to be an absolute synonym of *Planorbula*, and, on inspecting specimens of *Planorbula wheatleyi* in the Academy of Natural Sciences of Philadelphia, I am inclined to agree with them.

[58]Names associated with the genus *Promenetus* are *carus* Pilsbry & Ferriss 1906, *coloradoensis* F.C. Baker 1945, *exacuous* Say 1821, *harni* 'Pilsbry' Harn 1891, *hudsonicus* Pilsbry 1934, *hyalina* Lea 1838, *megas* Dall 1905, *rubellus* Sterki 1894, *umbilicatellus* Cockerell 1887 and *umbilicatus* J.W. Taylor 1885. Haldeman (1842-45 [1844]) and subsequent authors have considered *hyalina* to be a scalariform *P. exacuous*. Harn's (1891) *harni* is a *nomen nudum*, which Pilsbry (1899d) synonymized with *P. rubellus*. Cockerell's *umbilicatellus* is a replacement name for J.W. Taylor's *umbilicatus* (*non Planorbis umbilicatus* Müller 1774). F.C. Baker (1945) described (posthumously) *coloradoensis* as a member of the genus *Menetus*. H.B. Baker (1946) placed it with *Promenetus*. Hibbard & Taylor (1960) synonymized it, along with *hudsonicus, megas* and *rubellus*, with *exacuous*. In regard to *differentiae* as based on shell characters, I agree with the synonymies above. I have not seen the type specimens of *P. carus*, but I anticipate

that they will prove to be the same as *P. umbilicatellus.*

[59]D.W. Taylor (1960) erected a subgenus, *Phreatomenetus*, for *Promenetus umbilicatellus* (Cockerell) (type species), the Texan *P. carus* (Pilsbry & Ferriss) and the Central American and Caribbean *P. circumlineatus* (Shuttleworth). However, because of the small number of species known from *Promenetus* s.l. (only two of which have been studied anatomically), and the considerable variability which exists between species of Planorbidae, Clarke (1973) did not consider it prudent to recognize subgenera in the genus *Promenetus.*

[60]The genus *Vorticifex* is based on the fossil species *V. tryoni* Meek (in Dall) 1870. Living species are included in the subgenus *Parapholyx*. Separating the fossil species from the Recent ones by placing them in different subgenera may not be desirable. "The variability of the species [of *Vorticifex* s.l.], and the intergradations of form, are so great that no subordinate groupings within the genus seem practicable at this time" (Taylor, 1966a).

[61] *Vorticifex (Parapholyx) solida* (Dall) may not be specifically distinct from *V. (P.) effusa* (Lea).

[62]Walter (1970) was of the opinion that all four species of *Neoplanorbis* are only variants of *Amphigyra alabamensis* Pilsbry. I have not had time to investigate this.

[63]The species of *Fluminicola* are not dealt with in the identification key. A list of species with distributions can be found on pp. 22-23.

[64]In spite of the several publications which deal with the subgenus *Walkerilla*, it is still not well defined. For example, in proposing the subgenus, Thiele (1928) mentioned that the radula of its type species, *Somatogyrus (Walkerilla) coosaensis* Walker, has a central tooth with a finely serrated cutting edge (in his fig. 25 he shows a central tooth with a non-prominent central cusp flanked on each side by nine lateral cusps) and on each side a row of 8-10 basal denticles. The central tooth of *S. isogonus* (Say) he illustrated as having a prominent central cusp flanked by four lateral cusps, and a row of three basal cusps on each side. Yet Thompson (1969) illustrated *S. (W.) tenax* Thompson as having a relatively prominent central cusp flanked by six lateral cusps, and a row of three basal cusps on each side. Thompson (1969) figured the verge of *S. (W.) tenax* (it is a simple tapering structure with a single duct leading to its apex) and indicated that this type of verge is subgenerically distinct from that of *Somatogyrus* s.s. The sculpture of the apical whorls of *S. (W.) tenax* is also considered subgenerically distinct, and is described as "fine spiral striations which begin on first quarter of whorl as minute punctations, then become more intense and coalesce into distinct striations that terminate at the end of the apical whorl where the striations are slightly oblique."

[65]The species of *Somatogyrus* s.s. are not dealt with in the identification key. A list of species with distributions can be found on pp. 23-24.

[66]Much of the key on the southern, especially Floridanian, Hydrobiinae is based on the detailed studies of Thompson (1968, 1969).

[67]The monotypic genus *Hoyia* is distinguished by its radula (F.C. Baker, 1926a). Its anatomy has not been studied, so its subfamilial placement is presumptive. "The radula of [*Hoyia*] *sheldoni* is totally unlike that of any other American amnicoloid observed or published. The teeth are all very small, about a third the size of those of *Amnicola limosa*, and the denticulations are very fine, all teeth beyond the central being multicuspid, with the cusps of equal size" (F.C. Baker, 1928c).

[68]Taylor (1966b) characterizes *Tryonia* as follows: "Shell turriform, with more whorls, a narrower outline, smaller aperture, and a deeper suture than in most *Pyrgophorus*. The sculpture may consist only of growth line[s], or may be coarsely lirate, plicate, or reticulate. Spines of the shoulder of the shell (characteristic of *Pyrgophorus*) are unknown in *Tryonia*.

"Virtually all of the species are known by shell alone, so that no trenchant characterization of the genus is possible. *Tryonia cheatumi* is known to be ovoviviparous like *Pyrgophorus* (Pilsbry, 1935b[a])."

[69]The species of *Aphaostracon* are not dealt with in the identification key. A list of species with distributions can be found on pp. 19-20.

[70]The keys for the genera *Marstonia* and *Rhapinema* are from Thompson (1977).

[71]The species of *Cincinnatia, Fontelicella* s.s., *Natricola* and *Fontigens* are not dealt with in the identification key. Lists of species with distributions are given on pp. 25-26, 30-31.

[72]Distinguishing characters for *Amnicola* s.s. and *Lyogyrus* are from Thompson (1968). The species of neither of these two subgenera are dealt with in the identification key. Lists of species with distributions can be found on pp. 28-30.

[73]Pilsbry & Ferriss (1906) described small discoidal shells found in drift debris of the Guadalupe River in Texas as *Valvata micra* and *V. micra nugax*, but called attention to similarities of the shells to the Palaearctic hydrobiids *Horatia* Bourguignat and *Daudebardiella* Boettger. Pilsbry and Ferriss stated further that, until fresh specimens with soft parts or opercula were found, the taxonomic position of these tiny mollusks would remain uncertain. In 1916, Pilsbry placed them in the genus *Horatia* and the subgenus *Hauffenia* Pollonera. Bole (1970) separated *Hauffenia* as a genus distinct from *Horatia*, using characters of the seminal receptacle and operculum to distinguish the two taxa. Taylor (1975) placed *micra* in *Hauffenia* and *nugax* in *Horatia*. As yet, there are no published anatomical or opercular data on the American species, so it is not known to which, if either, genus they belong.

[74]There is considerable local variation in *Leptoxis* s.s., which has been responsible for the creation of many nominal species and a large synonymy. "It is clear to the eye [that] the *Anculosa* [=*Leptoxis* s.s.] of the main parts of the Cumberland and Tennessee rivers [are] higher in proportion to diameter than are shells of headwaters and tributaries. . . . In *Anculosa* [=*Leptoxis* s.s.], environmental polymorphism . . . is less simple than in the lithasias that have been studied. The main river anculosae follow the rule of having shorter spires than the upriver and tributary colonies. There is also another environmental modification. The body whorls of main river anculosae are higher in proportion to diameter than those of head and tributary waters. . . . The changes are irregularly progressive" (Goodrich, 1934a: 12, 15). "*A. subglobosa* Say is the headstream representative in the Tennessee River system. It is replaced downstream by *A.* [*Leptoxis*] *praerosa* Say in the main river, and those forms of *Anculosa* [*Leptoxis*] which penetrate the lower tributaries are, with only one or two exceptions, either this species or obvious offshoots of it. The group can be spoken of as the *subglobosa-praerosa* complex" (Goodrich, 1938: 4-6).

[75]Goodrich (1940d: 19) mentions that the radula of *Mudalia* ["*Nitocris*"] is distinctly different from that of *Leptoxis* s.s. ["the true *Anculosa*"]. As yet, I have not been able to confirm this. Any future study of the generic/subgeneric relationships of these two groups should include an inspection of their radulae with the scanning electron microscope.

[76]The shell of *Leptoxis taeniata* is quite variable in regard to spiral sculpturing, ranging from completely smooth to lirate. In the past, populations with lirate forms have been called *L. griffithiana* (Lea).

[77]Smooth shells may occur in various populations of *Leptoxis formosa*, but spiral striae are characteristic of the species.

[78]The species of *Elimia, Juga, Lithasia* s.s., *Angitrema* and *Pleurocera* are not dealt with in the identification key. Lists of species with distributions can be found on pp. 32-42, 44-48.

[79]The species of *Stagnicola* and *Lanx* s.s. are not dealt with in the identification key. A list of species with distributions can be found on pp. 51-53.

[80]Various lymnaeids are characterized by having radulae with either bicuspid or tricuspid lateral teeth. In the genus *Fossaria*, members of the subgenus *Fossaria* s.s. have tricuspid lateral teeth (Fig. 773a), whereas members of the subgenus *Bakerilymnaea* have bicuspid laterals (Fig. 773b). Because of possession of bicuspid lateral teeth (characteristic of North American *Stagnicola*), *Bakerilymnaea* was previously placed with the stagnicolas.

FIG. 773. Lymnaeid radular teeth. **a**, a central tooth and a *tricuspid* 1st lateral tooth; b, a central tooth and a *bicuspid* 1st lateral tooth.

[81]The relationships of the Alaskan representatives of the Holarctic *Fossaria truncatula* to Eurasian members of the species, as well as to the more eastern American fossarias, has not been critically studied.

[82]The shape of the shell of *Fossaria (Bakerilymnaea) hendersoni* from Colorado is quite similar to that of *F. (B.) sonomaensis*. Hibbard & Taylor (1960) considered the shell of *F. (B.) hendersoni* to fall within the range of variation of *F. (B.) cockerelli*. *F. (B.) sonomaensis* also may prove to be merely a morph of *cockerelli*, or of *bulimoides*, as suggested by Clarke (1973).

[83]The strong spiral striation of "*Galba*" *alberta* F.C. Baker suggests that this morph or species may belong to *Stagnicola* rather than to *Fossaria (Bakerilymnaea)*.

[84]The distinction between *Fossaria dalli* and *F. perplexa* seems a bit dubious. The latter has been reported from Washington (F.C. Baker & Henderson, 1929) and (as a morph of *bulimoides*) from California, Montana, Utah, Nevada and Arizona (Clarke, 1973).

[85]The Physidae are taken to genera in this key, except for *Aplexa* and *Stenophysa*, which are keyed to species. Lists of species with distributions can be found on pp. 53-57.

[86]North American snails of the genus *Aplexa* have generally been referred to the Eurasian species *A. hypnorum* (Linnaeus). Starobogatov & Streletzkaja (1967) and Te (1978, 1980) recognized the Western Hemisphere *Aplexa* as *A. elongata* (Say). Starobogatov & Streletzkaja reported *A. elongata* also in eastern Siberia.

[87]Couplets 5, 6 and 7 are from Walker (1908c).

[88]From Clarke (1973).

[89]F.C. Baker (1945) recognized only two species of *Planorbella* s.s., *P. campanulata* (Say) and *P. multivolvis* (Case), but for *campanulata* he recognized the nine subspecies [as *Helisoma (Planorbella) campanulatum*] listed below. [I have ommitted three subspecies known only as fossils.]

- *P. campanulata campanulata* (Say 1821). Vermont west to North Dakota, south to Ohio and Illinois, northward to Great Slave Lake (F.C. Baker, 1928c).
- *P. campanulata wisconsinensis* (Winslow 1926). Wisconsin, Michigan, and probably Quebec, Ontario and Manitoba (Winslow, 1926; F.C. Baker, 1928c).
- *P. campanulata davisi* (Winslow 1926). Michigan and Wisconsin (F.C. Baker, 1928c); New Hampshire (F.C. Baker, 1942c).
- *P. campanulata canadensis* (F.C. Baker & Cahn 1931). Lakes of northern Ontario (F.C. Baker & Cahn, 1931).
- *P. campanulata collinsi* (F.C. Baker 1939). Lake of the Woods District, western Ontario (F.C. Baker, 1939b).
- *P. campanulata michiganensis* (F.C. Baker 1927). Mud Lake, Roscommon County, Michigan (F.C. Baker, 1927e).
- *P. campanulata rudentis* (Dall 1905). Knee Lake, on Hayes River, Keewatin, northern Manitoba, Canada (Dall 1905; F.C. Baker & Cahn, 1931).
- *P. campanulata smithi* (F.C. Baker 1912). Douglas Lake, Michigan; ? also northern Wisconsin (F.C. Baker, 1928c).
- *P. campanulata rideauensis* (F.C. Baker 1945). Rideau River, Ottawa, Canada (F.C. Baker, 1945).

Clarke (1973) placed *rudentis* Dall, *wisconsinensis* Winslow, *davisi* Winslow and *canadensis* F.C. Baker & Cahn in the synonymy of *campanulata* s.s. He recognized *collinsi* F.C. Baker and also apparently *multivolvis* Case as subspecies of *campanulata*.

[90]Most of the nominal taxa within the subgenus *Pierosoma* are not critically enough defined, especially in regard to geographic, microgeographic and ecophenotypic variation, to present more than a very tentative taxonomy at this time. F.C. Baker (1945: 149) recognized [as "*Helisoma (Pierosoma)*"] 17 species plus an additional 10 subspecies for North America north of Mexico: *ammon* (Gould), *binneyi* (Tryon), *chautauquensis* F.C. Baker, *corpulenta corpulenta* (Say), *corpulenta vermilionensis* F.C. Baker, *horni* (Tryon), *magnifica* (Pilsbry), *multicostata multicostata* F.C. Baker, *multicostata whiteavesi* F.C. Baker, *occidentalis occidentalis* (Cooper), *occidentalis depressa* F.C. Baker, *oregonensis* (Tryon), *pilsbryi pilsbryi* (F.C. Baker), *pilsbryi infracarinata* (F.C. Baker), *plexata* (Ingersoll), *subcrenata*

subcrenata (Carpenter), *subcrenata disjecta* (Cooper), *tenuis californiensis* F.C. Baker, *tenuis sinuosa* (Bonnet), *traski* (Lea), *trivolvis trivolvis* (Say), *trivolvis fallax* (Haldeman), *trivolvis lenta* (Say), *trivolvis macrostoma* (Whiteaves), *trivolvis turgida* (Jeffreys), *truncata* (Miles) and *winslowi* (F.C. Baker). Baker (*op. cit.*) included *P. horni* and *P. plexata* as subspecies of *subcrenata* on plates 90, 92 and 93. He named additional taxa later in the same work: *randolphi* (a variety of *binneyi*), *columbiensis, kennicotti, preblei* (a variety of *pilsbryi*), *perdisjuncta* (a variety of *subcrenata*) and *marshalli* (a variety of *trivolvis*).

Clarke (1973) placed *fallax* Haldeman and *macrostoma* (Whiteaves) in the synonymy of *trivolvis* (Say), *horni* (Tryon) and *plexata* (Ingersoll) in the synonymy of *subcrenata* (Carpenter), *kennicotti* F.C. Baker and *preblei* F.C. Baker in the synonymy of *pilsbryi infracarinata* F.C. Baker, and *multicostata* F.C. Baker in the synonymy of *corpulenta* (Say). He (*op. cit.*) considered *subcrenata* to be a subspecies of *trivolvis.*

[91]The subgenus *Seminolina* was named by Pilsbry (1934a) to include "*Helisoma*" *scalare* (Jay 1839) (the type species), "*Helisoma*" *duryi* (Wetherby 1879) and its subspecies and forms, and the Pliocene "*Helisoma*" *conanti* (Dall 1890) and "*Helisoma*" *disstoni* (Dall 1890). He (p. 31) characterized them as, "Helisomas in which the external duct from penial gland to upper sac is short and adnate. Shell shaped like *Pierosoma* or with the spire produced on the left side and scalar, Physa-shaped. The smooth or malleate surface is not thread-striate, usually glossy." F.C. Baker (1945: 130, 134) further characterized the subgenus: "Shell . . . Large; sinistral, physa-shaped or planorboid, with every gradation between these forms, usually widely or deeply umbilicated; surface smooth, usually glossy, without the thread-like striae of *Pierosoma*. . . . *Seminolina* is a notable group of the subfamily Helisomatinae and one of the most variable genera as regards species. The physoid aspect of its type species, *Paludina scalaris* Jay, led the older conchologists to include it in *Physa* and the genus *Ameria* of the family [sic] Bulinidae. The largest species, [sic] *Helisoma duryi* (Wetherby), is perhaps more variable than any other species found in America, its extremes being from typical Physa-shaped to flatly discoidal shell. The elongation of the spire always produces a physoid aspect. The races of *duryi* blend into each other and often three forms will occur in the same lot, as *normale, intercalare*, and *duryi*."



FIG. 774. Diagram showing relations of Florida forms of *Planorbella* of the subgenus *Seminolina* (from Pilsbry, 1934a).

[92]The *Planorbella (Pierosoma) ammon* (Gould) group includes the nominal species *ammon* Gould 1855, *traski* Lea 1856, *binneyi* Tryon 1867, *occidentalis* Cooper 1870 and *columbiensis* F.C. Baker 1945. "*H. binneyi, H. ammon, H. occidentale,* and *H. traskii* are all closely related and may be found to belong to 1 species when the problem is investigated thoroughly" (Clarke, 1973: 465).

Henderson (1934a) discussed and figured the latter four nominal species. Of *P. ammon* he said, "An important character is the strong slope of the lateral outline, giving the shell somewhat the shape of a truncated cone. This is shared by most *Helisoma* species, but is more marked than usual in this species. Many much depressed specimens of similar diameter from California might easily be assigned to *ammon*, and there seem to be some intergrades, but I am inclined to believe there is no close relationship between them." Regarding *P. traski*, Henderson said, "The resemblance of this species to *binneyi* is notable, but it is more nearly barrel shaped, considerably higher proportionally, and the sculpture less pronounced, especially on the last whorl, where the striae are very fine, but just in front of the aperture they are coarser, and the apical whorls are deeply sunken. Young specimens of *ammon* from the same region much resemble *traskii*, but they soon begin to lose their barrel shape and take on the truncated cone shape of *ammon*, the carina is not so sharp and the apex not so deeply sunken." Henderson (*op. cit.*) selected a neotype for *P. occidentalis* from Klamath Lake, Oregon. "The neotype measures 27.5 mm. in diameter and 15 mm. in altitude just back of the slightly everted lip, approximating Cooper's maximum measurements. The last whorl is not carinate, but is shortly rounded above and more broadly below. . . . Though somewhat resembling *H. binneyi* (Tryon) in the strap-like whorls, *occidentalis* differs markedly in the less pronounced sculpture and the disappearance of the carina at an early stage of growth."

"*Helisoma columbiense* shows relationship to the *binneyi* group in its sculpture and the carination of its whorls. It differs from the members of that group in that it is of smaller size, has less relative axial height, its rib striae are less widely spaced and the whorls are usually more angulate. It differs from the *subcrenatum* group in having more regular and less widely spaced rib striae, in its angulated base and spire depression, and in the shape of the aperture" (F.C. Baker, 1945: 223).

[93]A second nominal species will also fit the diagnosis provided by the second halves of key couplets 40 and 46, *Planorbella (Pierosoma) tenuis* (Fig. 735). It is not clear to me just which shell features can be used to separate it from *P. (P.) trivolvis subcrenata*. My general impression is that *tenuis* is usually smaller and more finely sculptured than typical *subcrenata*. F.C. Baker (1934a) named a subspecies from Santa Clara County, California, *Helisoma tenue californiense*. "This race is widely distributed in California from Santa Clara County southward. *Helisoma tenue* is widely distributed in California and does not differ materially from the species as found in Mexico and Arizona." F.C. Baker (1945) figured "*Helisoma tenue sinuosum* (Bonnet)" from Arizona, Texas, New Mexico and Mexico.

[94]Key couplets 44 and 45 are from Clarke (1973).

[95]A second nominal species or subspecies will also fit the diagnosis provided by the second half of key couplet 44, *Planorbella (Pierosoma) winslowi* (F.C. Baker 1926). It is not clear to me just which shell features can be used to separate it from *P. (P.) corpulenta corpulenta*. F.C. Baker (1926b) named it originally as "a very distinct variety of *trivolvis*. It resembles *pilsbryi* in some respects, but is smaller, only about half the size of adult individuals of that variety, and the body whorl is sharply angulated and more flat-sided. It was first thought to represent a distinct species, but the presence of individuals varying toward *trivolvis* in the type lot, as well as in nearby waters, indicate a relationship to this large planorbid." Further, the shell characteristics of *P. (P.) winslowi* merge into *P. (P.) pilsbryi infracarinata*, which merge with *P. (P.) pilsbryi* s.s., which in turn seem to merge into *P. (P.) trivolvis*. (See note 96.)

[96]Clarke (1973: 459 ff.) recognized the subspecies *Helisoma* [=*Planorbella*] *(Pierosoma) pilsbryi infracarinatum* F.C. Baker 1932, but not without some hesitation. "Since reliable criteria are lacking for any new evaluation of the biological relationship between this taxon and the more southern *Helisoma* [=*Planorbella*] *pilsbryi* Baker, the most recent opinion (Baker, 1945: 138) is followed and the name *H. p. infracarinatum* is used. . . . Baker [1936b] . . . commented on the 'perplexing variation' in this subspecies. The variation is so great, in fact, that one is initially tempted to consider it analogous to the variation exhibited by *Gyraulus deflectus* and to regard *Helisoma* [=*Planorbella (Pierosoma)*] *pilsbryi infracarinatum* as a frequently occurring morphological variant of *H. trivolvis* (Say). . . . It is also possible that *Helisoma pilsbryi infracarinatum* is a morph which is intermediate between *H. corpulentum* (*s. str.*) and *H. trivolvis* (*s. str.*) derived from sporadic introgressive hybridization or representing a surviving parental stock from which *H. corpulentum* arose. The status of *H. p. infracarinatum* as a separate taxon requires additional research" (Clarke, 1973: 461-462).

Clarke (1973) placed *Helisoma kennicotti* F.C. Baker 1945 and *Helisoma pilsbryi preblei* F.C. Baker 1945 in the synonymy of *Planorbella (Pierosoma) pilsbryi infracarinata* (F.C. Baker).

From a comparison of authentic material of *pilsbryi* (paratypes, ANSP 140269) and *infracarinata* (paratypes, ANSP 158589), as well as *winslowi* F.C. Baker 1926 (paratypes, ANSP 158596), and considering variation seen in other museum lots and presented in the literature, I can see no compelling reason to separate the three forms taxonomically with latinized names. Further, I suspect that *Planorbella pilsbryi* is not taxonomically distinct from *P. trivolvis*.

The spire carinae in the form *infracarinata* have a tendency to be better developed than in *pilsbryi*; these carinae are rather prominent in the form *winslowi*, the shell of which also has well-developed basal carinae.

[97]F.C. Baker's opinions regarding *Planorbella (Pierosoma) trivolvis* (Say) changed over the years. In his final (1945) publication, he recognized the eight subspecies [as *Helisoma (Pierosoma) trivolvis*] listed below and gave various localities.

- *P. trivolvis trivolvis* (Say 1817). "This type of shell is abundantly distributed in the northeastern part of the United States from Maine westward to Nebraska. The southward extension of the typical form appears to be northern Illinois and Indiana, Ohio, Pennsylvania, and New Jersey" (F.C. Baker, 1936b). In 1945, Baker mentioned New York, Michigan and Wisconsin.
- *P. trivolvis fallax* (Haldeman 1844). Massachusetts, Maine.
- *P. trivolvis lenta* (Say 1834). Central Illinois, Tennessee, Louisiana, Texas.
- *P. trivolvis macrostoma* (Whiteaves 1863). Southern Canada, northern Wisconsin.
- *P. trivolvis turgida* (Jeffreys 1830) (=*intertexta* Sowerby 1878). South Carolina?, Florida, Alabama, Texas, Arkansas.
- *P. trivolvis marshalli* (F.C. Baker 1945). New Jersey, New York, Maryland, Virginia.
- *P. trivolvis chautauquensis* (F.C. Baker 1928). Chautauqua River, New York.
- *P. trivolvis holstonensis* (F.C. Baker 1945). Holston River, southwestern Virginia.

Clarke (1973) added *subcrenata* Carpenter 1857 as a subspecies of *Planorbella* ["*Helisoma*"] *trivolvis*, giving its distribution as California to British Columbia and Yukon Territory and east to Utah, Colorado, Minnesota and Manitoba. He placed *fallax* Haldeman and *macrostoma* Whiteaves in the synonymy of *trivolvis* s.s., and *horni* Tryon 1865 and *plexata* Ingersoll 1876 in the synonymy of *subcrenata*.

I doubt if it is advisable at this time to recognize varieties or subspecies of *Planorbella trivolvis*, at least until a careful study is undertaken and completed on this common and wide-spread complex of North American planorbid snails. However, if geographic subspecific names fit a need, then perhaps four can be tentatively adopted: *P. trivolvis* s.s. (northern North America east of the Rocky Mountains, south to Nebraska, northern Illinois, Pennsylvania and New Jersey), *P. trivolvis lenta* (central U.S.A. south of Nebraska and central Illinois to Texas (?) and Louisiana), *P. trivolvis turgida* (southeastern U.S.A., south of Pennsylvania and west to Alabama, Arkansas and possibly Texas), and *P. trivolvis subcrenata* (Rocky Mountain and Pacific states and provinces, possibly east in the north to Manitoba and Minnesota).

[98]Pilsbry (1934a) recognized six races of *Planorbella (Seminolina) duryi*: *duryi* s.s., *intercalaris* Pilsbry 1887, *preglabrata* Marshall 1926, *eudiscus* Pilsbry 1934, *normalis* Pilsbry 1934 and *seminolis* Pilsbry 1934. These, along with *P. (S.) scalaris*, are illustrated in Fig. 774.

Planorbella (Seminolina) duryi seminolis is the subspecies which is characterized by an everted spire of varying degrees. Higher spired individuals are very similar in appearance to *P. (S.) scalaris*, but the latter is narrower and generally less widely umbilicate. Also, in *P. (S.) duryi seminolis*, the "lower [i.e., anterior] margin of [the] aperture is generally advanced beyond [that of] the upper [i.e., posterior]" margin (except in exceptionally long individuals) (Pilsbry, 1934a: 35), whereas in *P. (S.) scalaris* the upper (posterior) margin of the aperture (when viewed from the spire end of the shell) projects further than the lower (anterior) margin.

[99]The identification key for the Ancylidae is adapted from Basch (1963).

[100]The lateral teeth of *Rhodacmea* are distinct from other North American ancylids by the possession of an "enormous mesocone, the blade-like cusp extending beyond the base, the ectocone is back of the mesocone, entirely separated from it and has several small cusps; there is no endocone" (Walker, 1918b) [Fig. 775].



FIG. 775. Central and 1st radular teeth of ancylid limpets. *Rhodacmea* is on the left.

GENERIC SYNONYMY

Acroluxus Keep 1887 = misspelling of *Acroloxus* Beck 1837. ("*Acroluxus Nuttalli*, Hald." in Keep (1887) = *Fisherola nuttalli* (Haldeman 1841).)

Alleghenya Clench & Boss 1967 = *Mudalia* Haldeman 1840.

Amarula Sowerby 1842 = *Thiara* Röding 1798.

Amblostoma Rafinesque (in Binney) 1865 = *Ambloxis* Rafinesque 1818, which is an unidentifiable name. Both names have the same type species, *A. eburnea* Rafinesque (in Binney) 1865.

Ambloxis Rafinesque 1818 = an unidentifiable name; occasionally mentioned as *possibly* being the same as *Campeloma* Rafinesque 1819.

Ambloxus Rafinesque 1831 = *Thiara* Röding 1798.

Ameria Dall 1870, preoccupied = *Seminolina* Pilsbry 1934.

Ampullaria Lamarck 1799 = *Pila* Röding 1798, a genus of Africa and Asia. In the earlier literature, species of *Pomacea* were erroneously assigned to the genus *Ampullaria*.

Ampullarius Montfort 1810 = *Pomacea* Perry 1810.

Anaplocamus Dall 1895 = *Mudalia* Haldeman 1840.

Anculosa Say 1821 = *Leptoxis* Rafinesque 1819.

Anculotus Say 1825 = emendation of *Anculosa* Say 1821 = *Leptoxis* Rafinesque 1819.

Ancylotus 'Say' Herrmannsen 1846 = emendation of *Anculosa* Say 1821 = *Leptoxis* Rafinesque 1819.

Ancylus Müller 1774 = a genus of the Palaearctic and Ethiopian regions. In the earlier literature, many or most ancylid species of the Western Hemisphere, as well as the patelliform Lymnaeidae, were erroneously assigned to the genus *Ancylus*.

Apella 'Mighels' Anthony 1843 = *Gyrotoma* Shuttleworth 1845. *Apella* is an invalid name based on an unknown species.

Aphella " 'Mighels' Anthony" Hannibal 1912 = misspelling of *Apella* 'Mighels' Anthony 1843 = *Gyrotoma* Shuttleworth 1845.

Armigerus Clessin 1884 = *Biomphalaria* Preston 1910. See Opinion 735 [1965] of the International Commission on Zoological Nomenclature.

Australorbis Pilsbry 1934 = *Biomphalaria* Preston 1910.

Bithinia Gray 1824 = *Bithynia* Leach (in Abel) 1818.

Bovillina Dall 1924 = *Orygoceras* Brusina 1882.

Bulimnaea 'Haldeman' Hubendick 1951 = misspelling of *Bulimnea* Haldeman 1841.

Bulimula Dall 1885 = *Bithynia* Leach (in Abel) 1818.

Bulimus Scopoli 1777, suppressed by the International Commission on Zoological Nomenclature, Opinion 475, 1957 = *Bithynia* Leach (in Abel) 1818.

Bulinus Müller 1781 = a planorbid genus of Africa, the Mediterranean region, the Middle East, and some of the Indian Ocean islands. In the earlier literature, it was occasionally used erroneously for members of the Physidae, including North American *Aplexa*.

Bythinella Moquin-Tandon 1856 = a European genus; it is not known to occur in North America.

Bythinia MacGillivray 1843 = *Bithynia* Leach (in Abel) 1818.

Callina Hannibal 1912 = *Viviparus* Montfort 1810.

Carnifex Keep 1893 = misspelling of *Carinifex* W.G. Binney 1865.

Ceratodes Guilding 1828 = *Marisa* Gray 1824.

Ceriphasia Swainson 1840 = *Pleurocera* Rafinesque 1818.

Chilocyclus Gill 1863 = *Pomatiopsis* Tryon 1862.

Cincinna Hübner 1810 = *Valvata* Müller 1774.
Cochliopa Stimpson 1865 = a genus of Panama; not found in North America (see Morrison, 1946).
Conchylium Cuvier 1816 = *Pomacea* Perry 1810.
Costella Meek 1876 = *Costatella* Dall 1870.
Cyclemis Rafinesque 1819, undeterminable = ? *Viviparus* Montfort 1810.
Cyclostoma Lamarck 1799 = *Epitonium* Röding 1798, a marine snail; *Cyclostoma* Draparnaud 1801 = *Pomatias* Studer 1789, a land snail. Some North American freshwater truncatelloid snails have previously been erroneously assigned this generic name.
Dentatus 'Beck' Gray 1847 = *Planorbula* Haldeman 1840.
Discus Haldeman 1840, preoccupied = *Planorbula* Haldeman 1840.
Ellipstoma Rafinesque 1818 = an unidentifiable name.
Euamnicola Crosse & Fischer 1891 = *Amnicola* Gould & Haldeman 1840.
Eurycaelon Lea 1864 = *Lithasia* Haldeman 1840.
Galba Schrank 1803 = a *nomen dubium*, based on an unidentifiable species (*Galba pusilla* Schrank 1803). In the past, *Galba* has been used unfortunately sometimes in place of *Fossaria* or *Stagnicola*.
Glottella Gray 1847 = *Angitrema* Haldeman 1841.
Goniobasis Lea 1862 = *Elimia* H. & A. Adams 1854. The type species of *Goniobasis* is *Goniobasis osculata* Lea 1862, selected by Hannibal (1912), which he said is the same as *Melania olivula* Conrad 1834. However, Goodrich (1936, 1941c) considered Lea's *osculata* to be a synonym of *Melania* ["*Goniobasis*"] *alabamensis* Lea 1861 and Conrad's *olivula* to be a distinct species. Both belong to the genus *Elimia*.
Gundlachia Pfeiffer 1849, type *G. ancyliformis* Pfeiffer 1849, by monotypy = a growth variant of *Ancylus havanensis* Pfeiffer 1839, which is a synonym of *Ancylus radiatus* Guilding 1829 (*fide* Harry & Hubendick, 1964). Not known to occur in the continental U.S.A. or Canada. Septate ancylids of North America (north of Mexico) are referable to the genus *Ferrissia*.
Haldemania Clessin 1880, preoccupied = *Ferrissia* Walker 1903.
Haldemania Tryon 1862 = *Lioplax* Troschel 1856.
Haldemanina Dall 1905 = *Planorbula* Haldeman 1840.
Helicosoma Agassiz 1846 = *Helisoma* Swainson 1840.
Hydrobia Hartmann 1821 = a genus of Europe; it does not occur in North American fresh waters. In the earlier literature, many species of freshwater truncatelloid snails of the Western Hemisphere were assigned erroneously to this genus.
Hydrognoma Gistel 1848 = *Thiara* Röding 1798.
Hypsogyra Lindholm 1927 = *Planorbella* Haldeman 1842.
Ibicornu Dall 1924 = *Orygoceras* Brusina 1882.
Incilicornu Dall 1924 = *Orygoceras* Brusina 1882.
Kincaidilla Hannibal 1912 = *Ferrissia* Walker 1903.
Laphrostoma Rafinesque 1815, *nomen nudum* = *Neritina* Lamarck 1816.
Lecythoconcha Annandale 1920 = *Cipangopaludina* Hannibal 1912.
Leptolimnea Swainson 1840, type species *Buccinum glabra* Müller 1774 = a European species.
Limnaea Blainville 1824 = *Lymnaea* Lamarck 1799.
Limnea Link 1807 = *Lymnaea* Lamarck 1799.
Limneus Draparnaud 1801 = *Lymnaea* Lamarck 1799.
Limnophysa Fitzinger 1833 = *Stagnicola* Leach (in Jeffreys) 1830.
Lithoglyphus Hartman 1821 = a European genus, possibly congeneric with the North American *Fluminicola* Stimpson 1865 (see note 10, p. 195).
Lithoparches Gistel 1848 = *Thiara* Röding 1798.
Lutella Haldeman 1840 = *Bithynia* Leach (in Abel) 1818.
Lymnaeus Cuvier 1817 = *Lymnaea* Lamarck 1799.
Lymneus Brard 1810 = *Lymnaea* Lamarck 1799.
Lymnula Rafinesque 1819 = *Lymnaea* Lamarck 1799.

Lymnulus Rafinesque (in Binney) 1865 = *Ambloxis* Rafinesque 1818, which is an unidentifiable name. Both names have the same type species, *A. eburnea* Rafinesque (in Binney) 1865.
Lymnus Montfort 1810 = *Lymnaea* Lamarck 1799.
Lythasia 'Lea' H. & A. Adams 1854 = spelling variation of *Lithasia* Haldeman 1840.
Macrolimen Lea 1862 = *Elimia* H. & A. Adams 1854.
Megara H. & A. Adams 1854 = *Angitrema* Haldeman 1841.
Megastropha Walker 1918 = misspelling of *Megasystropha* Lea 1864.
Megasystropha Lea 1864 = *Carinifex* Binney 1865. See Opinion 432 [1956] of the International Commission on Zoological Nomenclature.
Melacantha Swainson 1840 = *Thiara* Röding 1798.
Melafusus Swainson 1840 = *Io* Lea 1831.
Melania Lamarck 1799 = *Thiara* Röding 1798.
Melanidia Rafinesque 1815 = *Melania* Lamarck 1799 = *Thiara* Röding 1798.
Melantho Bowdich 1822 = *Campeloma* Rafinesque 1819.
Melas Montfort 1810 = *Thiara* Röding 1798.
Melasma H. & A. Adams 1854 = *Elimia* H. & A. Adams 1854.
Melatoma Anthony 1843 (not *Melatoma* Swainson 1840) = *Gyrotoma* Shuttleworth 1845.
Melatoma Swainson 1840 = a marine group.
Meseschiza Lea 1864 = *Angitrema* Haldeman 1841.
Meseshiza Lea 1876 = spelling error of *Meseschiza* = *Angitrema* Haldeman 1841.
Nasonia F.C. Baker 1928, preoccupied = *Bakerilymnaea* Weyrauch 1964.
Nauta Leach (in Turton) 1831 = *Aplexa* Fleming 1820.
Nautilus Linnaeus 1758 = a tetrabranch cephalopod. Used for *Gyraulus (Armiger) crista* (Linnaeus 1758) in the original species description.
Nerita Linnaeus 1758 = a marine genus, not found in North American fresh waters.
Nitocris H. & A. Adams 1854 = *Mudalia* Haldeman 1840.
Omphemis Rafinesque 1819, undeterminable = ? *Viviparus* Montfort 1810.
Omphiscola Rafinesque 1819 = an unidentifiable name.
Oxytrema Rafinesque 1819 = *nomen dubium.*
Paludestrina Orbigny 1839 = *Hydrobia* Hartmann 1821, a genus of Europe; it does not occur in North American fresh waters. In the earlier literature, many species of freshwater snails of the Western Hemisphere were listed under both of these generic names.
Paludina Lamarck (in Férussac) 1812 = *Viviparus* Montfort 1810.
Paradines Dall 1924 = *Vorticifex* Meek (in Dall) 1870.
Phreatomenetus Taylor 1960 = ? *Promenetus* F.C. Baker 1935.
Physina Rafinesque 1815 = *Physa* Draparnaud 1801.
Physodon Haldeman 1843 = *Physella* Haldeman 1843.
Planorbina Haldeman 1842 = *Biomphalaria* Preston 1910. See Opinion 735 [1965] of the International Commission on Zoological Nomenclature.
Planorbis Müller 1774 = a genus of the Palaearctic and Ethiopian regions. In the earlier literature, many or most planorbid species of the Western Hemisphere were assigned erroneously to the genus *Planorbis.*
Planorbulina Martens 1899, preoccupied = *Planorbula* Haldeman 1840.
Pleurovalvata Haas 1939 = *Valvata* Müller 1774.
Pompholycodea Lindholm 1927 = *Parapholyx* Hanna 1922.
Pompholyx Lea 1856, preoccupied = *Parapholyx* Hanna 1922.
Pomus H. & A. Adams 1856 = *Pomacea* Perry 1810.
Potamopyrgus Stimpson 1865 = a New Zealand genus; *P. jenkinsi* (Smith) has been introduced to and is widely distributed in Britain and Europe, but as yet no species of *Potamopyrgus* is known to occur in North America. North American species previously referred to *Potamopyrgus* are now assigned to other genera.
Pseudogalba F.C. Baker 1913 = *Fossaria* Westerlund 1885.

Pyrgula Cristofori & Jan 1832 = a genus of Europe; it does not occur in North American fresh waters. In earlier literature, some species of North American truncatelloid snails were assigned erroneously to this genus.
Rhodocephala Walker 1917 = *Rhodacmea* Walker 1917.
Scaphe 'Klein' Mörch 1852 = *Vitta* Mörch 1852, the North American subgenus of freshwater *Neritinia*.
Schizochilus Lea 1853 = *Gyrotoma* Shuttleworth 1845.
Schizostoma Lea 1843, preoccupied = *Gyrotoma* Shuttleworth 1845.
Segmentina Fleming 1817 = a genus of the Palaearctic region. In the earlier literature, species of *Planorbula* sometimes erroneously were assigned to the genus *Segmentina*.
Simpsonia F.C. Baker 1911, preoccupied = *Pseudogalba* F.C. Baker 1913 = *Fossaria* Westerlund 1885.
Spirodon 'Anthony' Tryon 1873 = *Mudalia* Haldeman 1840.
Stimpsonia Clessin 1878, preoccupied = *Fontigens* Pilsbry 1933.
Strepoma 'Rafinesque ms.' Haldeman 1863 = *Pleurocera* Rafinesque 1818.
Taphius H. & A. Adams 1855 = *Biomphalaria* Preston 1910. See Opinion 735 [1965] of the International Commission on Zoological Nomenclature.
Telescopella Gray 1847 = *Pleurocera* Rafinesque 1818.
Thomsonia Ancey 1886, preoccupied = *Seminolina* Pilsbry 1934.
Tiara Herrmannsen 1849, preoccupied = *Thiara* Röding 1798.
Tropidina H. & A. Adams 1854 = *Valvata* Müller 1774.
Trypanostoma Lea 1862 = *Pleurocera* Rafinesque 1818.
Tylotoma 'Haldeman' Fischer 1885 = emendation for *Tulotoma* Haldeman 1840.
Vancleavia F.C. Baker 1930 = *Probythinella* Thiele 1928.
Velletea Haldeman 1841 = spelling variation of *Velletia* Gray (in Turton) 1840 = *Acroloxus* Beck 1837. (*Ancylus (Velletea) nuttallii* Haldeman 1841 = *Fisherola nuttalli* (Haldeman 1841).)
Vivipara Sowerby 1813 = *Viviparus* Montfort 1810.
Viviparella Rafinesque 1815 = *Viviparus* Montfort 1810.

GLOSSARY

Abaxial. Directed away from the shell axis (i.e., the central line or central column of a coiled gastropod shell) outward.

Acroloxid. A common-name adjective referring to a member of the family Acroloxidae.

Acute. Sharp at the end.

Ampullariid. A common-name adjective referring to a member of the family Ampullariidae.

Ancylid. A common-name adjective referring to a member of the family Ancylidae.

Ancyliform. Limpet-shaped; patelliform; shaped like an obtuse cone (see Fig. 13).

Angular, angulate. Having an angle (or having the tendency to form an angle), rather than a round contour.

Angulation. Edge along which two surfaces in different planes meet at an angle.

ANSP. Abbreviation, usually associated with museum specimen catalogue numbers, for Academy of Natural Sciences of Philadelphia.

Aperture. The opening or "mouth" of a snail shell through which the head-foot protrudes when the snail is active.

Attenuate. Slender; elongated; long and narrow.

Auctorum (abbr. *auct.*). Of authors.

Auger-shaped. Shaped like an auger, i.e., with a flattened base terminating in a sharp, pointed twist.

Axial. Parallel to the axis or columella of a shell, i.e., transverse to the direction of the shell's spiral coil.

Base. The part of the shell opposite the apex. When a shell is held with the apex directed upward, the base is the "bottom" part of the shell. In regard to the natural position of the shell as carried by the snail, the "base" is the anterior end.

Bithyniid. A common-name adjective referring to a member of the family Bithyniidae.

Body whorl. The last complete whorl or volution of a spiral snail shell, measured from the outer lip back to a point immediately above the outer lip (Fig. 5). It is normally the largest whorl of the shell, and is called the body whorl because it encloses the greatest part of the snail's body.

Callus. A layer of calcareous material on a shell secreted by the snail's mantle.

Campanulate. Flared at the end; bell-shaped.

Canaliculate. Bearing a channel or groove.

Carina (pl. carinae). A sharp spiral edge, ridge or "keel" on the outer shell surface (see Fig. 6).

Carinate. Having one or more sharp spiral edges, ridges or keels on the outer shell surface.

Central tooth. The median or rachidian tooth of a transverse row of radular teeth. It is flanked by lateral teeth (see Fig. 772).

Channeled. Bearing a channel or groove.

Clavate. Club-shaped; growing gradually thicker toward one end.

Cleaver-like. Shaped like a butcher's cleaver, i.e., like a short, flat, broad cutting instrument.

Color bands. Revolving spiral stripes of a darker hue or different color from the ground or background color which occur on some species of gastropod shells.

Columella. The internal column around which the whorls revolve; the axis of a spiral shell.

Columellar lip. The apertural margin at the columellar region of a coiled gastropod shell (see Fig. 5).

Compressed. Refers to the spire of a gastropod shell which is relatively flattened, i.e., is not elongated.

Concentric. Having the same center, e.g., the nucleus, and expanding outward in parallel (i.e., equidistant) lines, as in the lines of growth of an operculum (Fig. 15c).

Continental Divide. The highland which divides the North American continent into two very large drainage regions, one in which the streams flow generally eastward into the Gulf of Mexico, Atlantic Ocean, Hudson Bay and the Arctic Ocean, and the other in which the streams flow generally westward into the Great Basin, the Gulf of California, the Pacific Ocean and the Bering Sea.

Corneous. Horn-like.

Costa (pl. costae). A transverse rib or rounded ridge of considerable size on the surface of a shell (see Fig. 6).

Costate. Refers to a shell in which the surface is sculptured with heavy, regular transverse ridges or ribs.

Crassate. Gross; thick; coarse; neither thin nor fine.

Crepidulaform. Shaped like *Crepidula*, i.e., limpet-like with a small, coiled apex.

Ctenidium. The characteristic respiratory appendage or gill of mollusks.

Cusp. The cutting blade or blades projecting from each tooth of the molluscan radula (see Fig. 72).

Cylindrical. Shaped like a cylinder; round in cross-section with nearly parallel sides (Fig. 8e).

Decollate. Cut off, i.e., as with the shell of some snails where the top several whorls of the spire break off or erode away.

Depauperate. Condition in which an individual, colony or race exhibits the outward manifestation of disease, accident or malnutrition, or a reaction to adverse environment. See depauperization.

Depauperization. The outward manifestation of disease, accident or malnutrition, or a reaction to inimical environment. It affects individual mollusks fairly frequently, but also it sometimes involves whole colonies and races. Symptoms of depauperization are dwarfing, lack of nacreous material (in certain bivalves), loose coiling and simplification of shell characters (Goodrich, 1939a).

Depressed. Flattened dorso-ventrally or postero-anteriorally, as the spire of a shell.

Elongate. Lengthened; extending length-wise; especially higher than wide.

Entire. Refers to the lip or peritreme of a shell that forms a continuous circle or oval, i.e., it is not broken by a space where it meets the parietal wall of the body whorl.

Fissure. A narrow slit.

Fusiform. Spindle-shaped, i.e., with a relatively thick middle and tapered to a point at each end (Fig. 8c).

Geniculate. Having a joint or bend.

Gibbous. Very convex or swollen; tumid.

Gradate. Arranged in steps, as a spire with shouldered whorls.

Growth lines. Minute lines on the outer shell surface indicating minor rest periods during growth (see Fig. 6). Not to be confused with the major "rest marks" or varices, caused by prolonged growth arrest (as during winter).

Heliciform. Shaped like *Helix*, i.e., with the characteristic shape of the majority of land snails, which have a somewhat depressed spire and whorls that increase regularly in diameter.

Hydrobiid. A common-name adjective referring to a member of the family Hydrobiidae.

Hyaline. Glassy; glossy and translucent or nearly transparent.

Imperforate. Refers to a spiral gastropod shell which has no opening or external cavity at its base. In such a case, the inner sides of the coiled whorls are appressed, leaving no cavity, or, if they are not appressed and a cavity is formed, then its opening is completely covered by a callus or the reflected columellar apertural lip.

Incised. Grooved; engraved (see Fig. 6).

Inflated. Refers to snail shells or individual whorls which are bulbous or swollen in appearance.

Labrum. The outer part of the apertural lip of a coiled gastropod shell, as opposed to the parietal or umbilical lip and the basal (anterior) lip.

Lateral teeth. The teeth on each side of the central or rachidian tooth in a transverse row of radular teeth (see Fig. 772).

Lira (pl. lirae). A ridge, specifically a spiral ridge on the outer surface of a snail shell (see Fig. 6).

Lirate. Refers to a shell with spiral ridges on its external surface.

Longitudinal. Refers to shell sculpturing that is at right angles to the spiral direction of the shell's coil; transverse (see Fig. 6).

Lymnaeid. A common-name adjective referring to a member of the family Lymnaeidae.

Malleated. Dented as if hit by a hammer (see Fig. 6).

Marginal teeth. The longitudinal rows of teeth at each edge of the molluscan radula.

MCZ. Abbreviation, usually associated with museum specimen catalogue numbers, for Museum of Comparative Zoology (Harvard University).

Median cusp. The middle cusp of a molluscan radular tooth, generally flanked by smaller lateral cusps.

Median tooth. The central or rachidian tooth of a transverse row of radular teeth. It is flanked by lateral teeth (see Fig. 772).

Micromelaniid. A common-name adjective referring to a member of the family Micromelaniidae.

Multispiral. Refers to an operculum in which there are numerous, very slowly enlarging spirals, coils or whorls (Fig. 15a).

Neritinid. A common-name adjective referring to a member of the family Neritinidae.

Neritiniform. Shaped like *Neritina*, i.e., subglobose or hemispherical, with few rapidly enlarging whorls, very reduced spire, and a heavily calloused and expanded parietal apertural margin (Fig. 14).

Nodule. A small knot, lump or irregularly shaped mass, such as the projections occurring on the shell surface of some freshwater snails (see Fig. 6).

Nomen dubium (pl. *nomina dubia*). A dubious name; one that cannot be applied with certainty to any known taxon.

Nomen nudum (pl. *nomina nuda*). A newly introduced species name without sufficient description to justify its acceptance in the zoological literature.

Nomen oblitum (pl. *nomina oblita*). A forgotten name. A name that has not been used as a senior synonym in the primary zoological literature for more than 50 years. Such a name has no validity in zoological nomenclature.

Nuchal lobe. One of the two right and left lobes at the anterior head-foot margin on either side of the mouth.

Nucleus. The first-formed (earliest) part of beginning of a shell or operculum (e.g., see Fig. 15d).

Oblique. Slanting; greater or less than a right angle; neither parallel with nor perpendicular to.

Obsolete. Obscure; indistinct; very rudimentary.

Obtuse. Blunt or rounded at the end, not acute or pointed.

Operculum (pl. opercula). A corneous or calcareous plate borne on the dorsal posterior foot of prosobranch snails which closes the aperture when the snail withdraws into its shell (Fig. 1).

Oval, ovate. In the shape of the longitudinal section of a hen's egg, i.e., oblong and curvilinear, with one end narrower than the other (Fig. 8b).

Ovoviviparus. Condition in which the young snails are formed within an egg, but hatch while still inside the mother snail, from which they emerge as young crawling snails.

Pagoda-like. Shaped like a pagoda, i.e., with a tapering, tower-like, storied, carinate shell spire (see Fig. 443).

Patelliform. Limpet-shaped; ancyliform; shaped like an obtuse cone (see Fig. 13).

Parietal. Pertains to the inside wall of the shell aperture (see Fig. 5).

Paucispiral. Refers to an operculum in which there are few rapidly enlarging spirals, coils or whorls (Fig. 15b).

Perforate. Refers to a spiral gastropod shell which has a very narrow perforation at its base, formed where the inner sides of the coiled whorls do not join.

Periostracum. The thin proteinaceous external layer covering most mollusk shells.

Periphery. The edges of a shell as seen in outline.

Peritreme. The peristome, apertural "lip" or apertural margin of a gastropod shell (does not include the parietal wall in shells without an entire (continuous) apertural margin).

Physid. A common-name adjective referring to a member of the family Physidae.

Physoid. Shaped like the shell of a member of the family Physidae, i.e., sinistral and with a raised spire.

Planispiral. Coiled in one plane (Fig. 12).

Planorbid. A common-name adjective referring to a member of the family Planorbidae.

Pleurocerid. A common-name adjective referring to a member of the family Pleuroceridae.

Plica (pl. plicae). A transverse or "vertical" ridge or "rib" on the outer shell surface.

Plicate. Bearing plicae, which are transverse or "vertical" ribs on a shell.

Plicate-striate. Refers to a shell having longitudinal (transverse) folds or ribs on its surface that are crossed by raised spiral lines.

Pomatiopsid. A common-name adjective referring to a member of the family Pomatiopsidae.

Ponderous. Very heavy; very thick.

Pseudobranch. A "false" or secondarily derived gill; a vascularized, fleshy outgrowth near the opening to the pulmonary cavity (pneumostone) of aquatic pulmonate snails which aids in respiration (see Fig. 3a). Not a true ctenidium.

Pyriform. Pear-shaped, i.e., large and round at one end and tapering at the other end.

Radula (pl. radulae). A rasp-like structure in the anterior end of the digestive tract of all mollusks except pelecypods which is used to scrape off food during feeding. The radula consists typically of a number of longitudinal and transverse rows of minute sharp "teeth", each with one or more cutting blades or "cusps" (see Fig. 18).

Revolving lines. A term sometimes used for spiral striae (see Fig. 6); occasionally also called "spirals".

Rimate. Refers to a coiled gastropod shell that has at its base a narrow "umbilical" opening that is partially closed by the expansion of the anterior columellar lip.

Rounded. Having a more or less evenly curved contour, in contrast to being angular.

Scalar. Pertaining to or like a flight of steps, i.e., a shell with elevated spire formed of right-angular whorls.

Scalariform. Shell form, usually pathologically produced, in which the whorls are disjoined or tend to become so.

Sculpture. The natural surface markings, other than those of color, usually found on snail shells, and often furnishing identifying marks for species recognition (see Fig. 6).

Sensu lato (abbr. *s.lat.* or *s.l.*). In the broad sense.

Sensu stricto (abbr. *s.str.* or *s.s.*). In the strict sense.

Shouldered. Refers to the appearance (in outline) of the posterior outer peripheral part of a whorl that is sharply rounded in contrast to the more even curvature of the rest of the shell (Fig. 11c).

Sic. Thus (to indicate exact transcription).

Sinuous. Wavy or S-shaped.

Spade-shaped. Shaped like a spade, i.e., like a broad, flat blade tapering rapidly at one end.

Spatulate. Shaped like a spatula, i.e., broad and oblong at one end, tapering rapidly near the center, and continuing as a narrower elongation at the other end.

Spindle-shaped. Fusiform; shaped like a spindle, i.e., with a relatively thick middle and tapered to a point at both ends (Fig. 8c).

Spiral. Winding, coiling or circling around a central axis; winding around a fixed point and continually receding from it; the form of the shell of most snails.

Spiral sculpture. Surface markings of a snail shell which pass continuously around the whorls more or less parallel to the suture (see Fig. 6).

Spire. The whorls of a snail shell, excepting the last or body whorl (Fig. 5). The spire is measured as the distance (parallel to the columella) from the suture where the apertural lip meets the body whorl to the shell apex.

Stria (pl. striae). A slight superficial spiral groove or furrow on the outer shell surface, or a fine spiral threadlike line or streak (see Fig. 6). Commonly used also, in a less precise sense, for raised spiral ridges on the shell surface.

Striate. Refers to a shell having spiral incised lines on its surface (see Fig. 6). Also used, less precisely, to describe shells with spiral raised lines, or for shells covered with fine transverse lines.

Subglobose. Nearly globular or spherical in shape (Fig. 8a).

Succiniform. *Succinea*-like, i.e., with a thin and fragile shell, which has a large oval aperture and body whorl and a small spire.

Suture. The line on the shell surface where two adjoining whorls meet (see Fig. 5).

Taxon (pl. taxa). Any taxonomic group, e.g., a race, subspecies, species, genus, family, order, etc.

Thiarid. A common-name adjective referring to a member of the family Thiaridae.

Transverse. At right angles to the spiral direction of the whorls; parallel to the columella or axis of the shell; in the same direction as (i.e., parallel to) the growth lines of a snail shell (see Fig. 6).

Truncatelloid. A common-name adjective referring to a member of the superfamily Truncatelloidea.

Tuberculate. Covered with tubercles or rounded knobs.

Tubercle. A nodule (see Fig. 6) or small eminence, such as a solid elevation occurring on the shell surface of some gastropods.

Tumid. Swollen or enlarged.

Turbinate, turbiniform. Shaped like a turban; refers to a shell in which the whorls decrease rapidly in diameter and taper broadly from a circular base to the apex (Fig. 8d).

Umbilicate. Refers to a spiral gastropod shell which has an opening or cavity at its base, and more specifically to one in which the opening is more than a very narrow perforation. This cavity is formed in those shells in which the inner sides of the coiled whorls do not join.

UMMZ. Abbreviation, usually associated with museum specimen catalogue numbers, for the University of Michigan Museum of Zoology (sometimes incorrectly cited as MZUM).

USNM. Abbreviation, usually associated with museum specimen catalogue numbers, for the United States National Museum (National Museum of Natural History).

Valvatid. A common-name adjective referring to a member of the family Valvatidae.

Viviparid. A common-name adjective referring to a member of the family Viviparidae.

Whorl (spelled "whirl" in early literature). One complete turn or coil of a spiral gastropod shell (see Fig. 5).

REFERENCES*

ABBOTT, R. Tucker. 1948. A potential snail host of oriental schistosomiasis in North America (*Pomatiopsis lapidaria*). *Proc. U.S. natl. Mus.*, 98(3222): 57-68, pl. 3.

ABBOTT, R. Tucker. 1952. A study of an intermediate snail host (*Thiara granifera*) of the oriental lung fluke (*Paragonimus*). *Proc. U.S. natl. Mus.*, 102(3292): 71-116, pls. 8, 9.

ABBOTT, R. Tucker. 1973. Spread of *Melanoides tuberculata* (Thiaridae). *Nautilus*, 87(1): 29.

ABDEL-MALEK, Emile T. 1952a. The preputial organ of snails in the genus *Helisoma*. *News Bull. annu. Rep. Am. malacol. Union 1951*, (18): 5-6.

ABDEL-MALEK, Emile T. 1952b. The preputial organ of snails in the genus *Helisoma* (Gastropoda: Pulmonata). *Am. midl. Nat.*, 48(1): 94-102, pls. 1, 2.

ABDEL-MALEK, Emile T. 1954. Morphological studies on the family Planorbidae (Mollusca: Pulmonata). I. Genital organs of *Helisoma trivolvis* (Say) (subfamily Helisomatinae F.C. Baker, 1945). *Trans. Am. microsc. Soc.*, 73(2): 103-124.

ADAMS, C.B. 1841a. Catalogue of the Mollusca of Middlebury, Vt., and vicinity, with observations. *Am. J. Sci.*, 40(2): 266-277.

ADAMS, C.B. 1841b. Descriptions of thirteen new species of New England shells. *Boston J. nat. Hist.*, 3(3): 318-332, pl. 3.

ADAMS, Chas. C. 1892. Mollusks as cat-fish food. *Nautilus*, 5(11): 127-128.

ADAMS, Chas. C. 1915. The variations and ecological distribution of the snails of the genus *Io*. *Mem. natl. Acad. Sci.*, 12(2), 2nd mem., 92 pp., pls. 1-60.

ADAMS, Henry & ADAMS, Arthur. 1863. Descriptions of new species of shells, chiefly from the Cumingian collection. *Proc. sci. Meetings zool. Soc. Lond., 1863*, pp. 428-435.

ADAMS, Lionel E. 1905. Conchological notes from the United States. *J. Conchol.*, 11(6): 165-168.

ADAMSTONE, F.B. 1924. The distribution and economic importance of the bottom fauna of Lake Nipigon with an appendix on the bottom fauna of Lake Ontario. *Univ. Toronto Stud., Publ. Ontario fish. res. Lab.*, (24): 33-100, pls. 1-4.

AGUAYO, C.G. 1949. Malacology and the official list of generic names. *Nautilus*, 63(1): 17-19.

AHLSTEDT, Steven. 1979. Recent mollusk transplants into the North Fork Holston River in southwestern Virginia. *Bull. Am. malacol. Union Inc. 1979*, 46: 21-23.

ALDRICH, T.H. 1905. A new species of *Somatogyrus* from south Alabama. *Nautilus*, 18(12): 140.

ALLEN, John A. 1911. *Lymnaea auricularia* Linné in Canada. *Nautilus*, 25(5): 60.

ALLISON, Leonard N. 1942. Trapping snails of the genus *Campeloma*. *Science*, 95(2457): 131-132.

AMERICAN MALACOLOGICAL UNION, INC. 1974. *How to study and collect shells*. A symposium. Am. malacol. Union, Box 394, Wrightsville Beach, N.C. 28480, U.S.A. pp. i-xi, 1-107.

ANCEY, C.F. 1887. Description of North American shells. *Conchol. Exc.*, 2(6): 79-80.

ANCEY, C.F. 1888. Étude monographique sur le genre *Pyrgulopsis*. *Bull. Soc. malacol. Fr.*, 5: 185-202. [Reprinted in *Sterkiana*, 1963, (10): 34-40.]

ANCEY, C.F. 1889. On Mr. Pilsbry's critics upon some American shells. *Nautilus*, 3(4): 39-42.

ANDERSON, Barbara Estel. 1966. Studies on the molluscan genus *Campeloma* Rafinesque, 1819. *Sterkiana*, (23): 9-18.

ANNANDALE, N. 1924. The evolution of the shell-sculpture in fresh-water snails of the family Viviparidae. *Proc. R. Soc.*, B, 96: 60-76.

ANTHONY, John G. 1840a. Descriptions of three new species of shells. *Boston J. nat. Hist.*, 3(1-2): 278-279, pl. 3, figs. 1-3.

ANTHONY, John G. 1840b. Descriptions of two new species of *Anculotus*. *Boston J. nat. Hist.*, 3(3): 394-395, pl. 3, figs. 4, 5.

ANTHONY, John G. 1841. [Published letter synonymizing *Anculotus kirtlandianus* Anthony with *Melania rogersii* Conrad.] *Proc. Boston Soc. nat. Hist.*, 1: 5.

ANTHONY, John G. 1843. [Published letter on *Schizostoma* and *Melatoma*.] *Proc. Acad. nat. Sci. Phila.*, 1: 251.

*See also Addenda on pp. 265-266.

ANTHONY, John G. 1854. Descriptions of new fluviatile shells of the genus *Melania* Lam., from the western states of North America. *Ann. Lyc. nat. Hist. N.Y.*, 6: 80-130, pls. 1-3.

ANTHONY, John G. 1855. Descriptions of new species of *Ancylus* and *Anculosa*, from the western states of North America. *Ann. Lyc. nat. Hist. N.Y.*, 6: 158-160, pl. 5, figs. 20-23.

ANTHONY, John G. 1860. Descriptions of new species of American fluviatile gasteropods. *Proc. Acad. nat. Sci. Phila.*, 12: 55-72.

ANTHONY, John G. 1862. Catalogue of shells from John G. Anthony, Cincinnati, Ohio. *Fifteenth rep. Cab. nat. Hist., Senate*, No. 116, pp. 15-21.

ANTHONY, John G. 1865. Descriptions of two new species of *Goniobasis*. *Am. J. Conchol.*, 1(1): 36, pl. 1, figs. 1-3.

ANTHONY, John G. 1866. Descriptions of new American fresh water shells. *Am. J. Conchol.*, 2(2): 144-147, pls. 6, 7.

ARCHER, Allan F. 1933. A new variety of *Viviparus contectoides* (W.G. Binn) from Jackson Co., Florida. *Nautilus*, 47(1): 18-20, pl. 3, figs. 1-3.

ARCHER, Allan F. 1939. The ecology of the Mollusca of the Edwin S. George Reserve, Livingston County, Michigan. *Occ. Pap. Mus. Zool. Univ. Mich.*, (398): 1-24.

ASHMUN, E.H. 1899. Collecting in Arizona and New Mexico. *Nautilus*, 13(2): 13-17.

ATHEARN, Herbert D. 1961a. Additions to the New Brunswick checklist. *Sterkiana*, (4): 33-34.

ATHEARN, Herbert D. 1961b. Some notes on the large freshwater snail *Tulotoma magnifica* Conrad. *Annu. Rep. Am. malacol. Union 1961*, (28): 5.

ATHEARN, Herbert D. 1968. Changes and reductions in our freshwater molluscan populations. *Annu. Rep. Am. malacol. Union 1967*, (34): 44-45.

ATHEARN, Herbert D. 1970. Discussion of Dr. Heard's paper [on rare and endangered eastern freshwater mollusks of the south Atlantic and Gulf drainages, in Am. malacol. Union Symp. Rare and Endangered Mollusks]. *Malacologia*, 10(1): 28-31.

ATHEARN, Herbert D. 1971. *Sphaerium simile* (Say) in Tennessee. *Nautilus*, 85(1): 37.

BAILEY, Albert. 1891. Shells of Erie Canal. *Nautilus*, 5(2): 23.

BAILY, Joshua L., Jr. 1929. Fresh water Mollusca in brackish water. *Nautilus*, 43(1): 34.

BAILY, Joshua L., Jr. 1931. Some data on growth, longevity, and fecundity in *Lymnaea columella* Say. *Biologia generalis*, 7(3): 407-428.

BAILY, Joshua L., Jr. 1934. The effect of alcohol on the growth of *Physa traskii*. *Rep. Am. malacol. Union 1934*, (3): 5.

BAILY, Joshua L., Jr. 1950. Some preliminary notes on the distribution of Mollusca in the lakes of the western states. *Nautilus*, 63(3): 73-78.

BAILY, Joshua L., Jr. 1956. Observations on the recently extinct mollusk fauna of Panamint Lake. *Nautilus*, 69(3): 100-103.

BAILY, Joshua L., Jr. & BAILY, Ruth Ingersoll. 1933. A new species of *Lymnaea* from Oregon. *Nautilus*, 47(1): 33-34, pl. 6, fig. 4.

BAILY, Joshua L., Jr. & BAILY, Ruth Ingersoll. 1951-52. Further observations on the Mollusca of the relict lakes in the Great Basin. *Nautilus*, 65(2): 46-53, pl. 4; 65(3): 85-93.

BAILY, Joshua L., Jr. & BAILY, Ruth Ingersoll. 1952. *Amnicola pilsbryana*, new name. *Nautilus*, 65(4): 144.

BAILY, Joshua L., Jr., PEARL, Raymond & WINSOR, Charles P. 1932. Variation in *Goniobasis virginica* and *Anculosa carinata* under natural conditions. *Biologia generalis*, 8(2): 607-630, pls. 14, 15.

BAILY, Joshua L., Jr., PEARL, Raymond & WINSOR, Charles P. 1933a. Variation in *Goniobasis virginica* and *Anculosa carinata* under natural conditions. II. The relation between size of the shells, and environmental factors. *Biologia generalis*, 9(1), lief. 2: 301-336, pls. 6-8.

BAILY, Joshua L., Jr., PEARL, Raymond & WINSOR, Charles P. 1933b. Variation in *Goniobasis virginica* and *Anculosa carinata* under natural conditions. III. Correlations, shape of shells, and conclusions. *Biologia generalis*, 9(2), lief. 3: 48-69.

BAILY, Joshua T., Jr. 1909. *Viviparus* in Philadelphia. *Nautilus*, 23(4): 60.

BAKER, Frank Collins. 1896. On the correct position of the aperture in *Planorbis*. *J. Cincinnati Soc. nat. Hist.*, 19(1): 45-48.

BAKER, Frank Collins. 1897a. On the pulsations of the molluscan heart. *J. Cincinnati Soc. nat. Hist.*, 19(2): 73-78.

BAKER, Frank Collins. 1897b. On a collection of mollusks from Grand Tower, Illinois. *Nautilus*, 11(3): 28-30.

BAKER, Frank Collins. 1897c. Notes on radulae. *J. Cincinnati Soc. nat. Hist.*, 19(3): 81-92, pls. 4, 5.

BAKER, Frank Collins. 1898a. The molluscan fauna of western New York. *Trans. Acad. Sci. St. Louis*, 8(5): 71-94.

BAKER, Frank Collins. 1898b. A day on the Chicago drainage canal. *Nautilus*, 12(6): 63-65.
BAKER, Frank Collins. 1899a. Notes on the mollusks of Lilycash Creek. *Nautilus*, 13(3): 30-31.
BAKER, Frank Collins. 1899b. Notes on the Mollusca of Owasco Lake, N.Y. *Nautilus*, 13(5): 57-59.
BAKER, Frank Collins. 1900a. On a collection of fresh-water shells from Rhode Island. *Nautilus*, 13(10): 112-113.
BAKER, Frank Collins. 1900b. The gross anatomy of *Limnaea emarginata* Say, variety *mighelsi* Binney. *Bull. Chicago Acad. Sci.*, 2(3): 191-211, pls. 1-6.
BAKER, Frank Collins. 1900c. A revision of the Physae of northeastern Illinois. *Nautilus*, 14(2): 16-24.
BAKER, Frank Collins. 1900d. Shell collecting near Rochester, N.Y. *Nautilus*, 14(6): 69-71.
BAKER, Frank Collins. 1901a. A revision of the limnaeas of northern Illinois. *Trans. Acad. Sci. St. Louis*, 11(1): 1-24, pl. 1.
BAKER, Frank Collins. 1901b. Snails of pond, river, and brook. *Birds and Nature*, 9: 128-133.
BAKER, Frank Collins. 1901c. New varieties of fresh water shells. *Nautilus*, 15(2): 17-18.
BAKER, Frank Collins. 1901d. The digitations of the mantle in *Physa*. *Bull. Chicago Acad. Sci.*, 2(4): 225-228, pls. 1, 2.
BAKER, Frank Collins. 1901e. Description of a new species of *Limnaea*. *Bull. Chicago Acad. Sci.*, 2(4): 229-230.
BAKER, Frank Collins. 1901f. The molluscan fauna of the Genesee River. *Am. Nat.*, 35(416): 659-664.
BAKER, Frank Collins. 1901g. *Limnaea auricularia* in America. *Nautilus*, 15(5): 59.
BAKER, Frank Collins. 1902a. *Planorbis bicarinatus striatus*, n. var. *Nautilus*, 15(10): 120.
BAKER, Frank Collins. 1902b. The Mollusca of the Chicago area. Part 2, The Gastropoda. *Bull. Chicago Acad. Sci., nat. Hist. Surv.*, 3(2): 131-410, index 8 pp., pls. 28-36.
BAKER, Frank Collins. 1903a. Shell collecting on the Mississippi. *Nautilus*, 16(9): 102-105.
BAKER, Frank Collins. 1903b. Pleistocene mollusks of White Pond, New Jersey. *Nautilus*, 17(4): 38-39.
BAKER, Frank Collins. 1904a. The mollusks of Cedar Lake, Indiana. *Nautilus*, 17(10): 112-113.
BAKER, Frank Collins. 1904b. New varieties of American limnaeas. *Nautilus*, 18(1): 10-12.
BAKER, Frank Collins. 1904c. The molluscan fauna of the Dells of Wisconsin. *Trans. Acad. Sci. St. Louis*, 14(2): 99-105.
BAKER, Frank Collins. 1904d. Notes on *Planorbis truncatus* Miles. *Trans. Acad. Sci. St. Louis*, 14(3): 107-110.
BAKER, Frank Collins. 1904e. New American lymnaeas. II. *Nautilus*, 18(6): 62-63.
BAKER, Frank Collins. 1905a. Critical notes on the smaller lymnaeas. *Nautilus*, 18(11): 125-127.
BAKER, Frank Collins. 1905b. New species of *Lymnaea*. *Nautilus*, 18(12): 141-142.
BAKER, Frank Collins. 1905c. The molluscan fauna of McGregor, Iowa. *Trans. Acad. Sci. St. Louis*, 15(3): 249-258.
BAKER, Frank Collins. 1905d. Notes on a collection of shells from Bass Lake, Indiana. *Nautilus*, 19(3): 27-28.
BAKER, Frank Collins. 1905e. A new species of *Lymnaea* from Ohio, with notes on *Lymnaea parva* Lea. *Nautilus*, 19(5): 51-53.
BAKER, Frank Collins. 1905f. Notes on the genitalia of *Lymnaea*. *Am. Nat.*, 39(465): 665-679.
BAKER, Frank Collins. 1906a. A new locality for *Planorbis nautileus* Linne. *Nautilus*, 19(10): 120.
BAKER, Frank Collins. 1906b. *Lymnaea hinkleyi* n. sp. *Nautilus*, 19(12): 142-143.
BAKER, Frank Collins. 1906c. Application of De Vries's mutation theory to the molluscs. *Am. Nat.*, 40(473): 327-334.
BAKER, Frank Collins. 1906d. Notes on a collection of mollusks from the vicinity of Alpena, Michigan. *Trans. Acad. Sci. St. Louis*, 16(2): 1-15, pl. 1.
BAKER, Frank Collins. 1906e. *Lymnaea danielsi* n. sp. *Nautilus*, 20(5): 55-56.
BAKER, Frank Collins. 1906f. A catalogue of the Mollusca of Illinois. *Bull. Ill. State Lab. nat. Hist.*, 7(6): 53-136, map, errata and addendum.
BAKER, Frank Collins. 1907a. Descriptions of new species of *Lymnaea*. *Nautilus*, 20(11): 125-127.
BAKER, Frank Collins. 1907b. Two new species of *Lymnaea*. *Nautilus*, 21(5): 52-55.
BAKER, Frank Collins. 1908a. Note on *Lymnaea desidiosa* Say. *Nautilus*, 22(2): 20-23.
BAKER, Frank Collins. 1908b. Suggestions for a natural classification of the family Lymnaeidae. *Science*, 27: 942-943.
BAKER, Frank Collins. 1908c. A new American *Planorbis*. *Nautilus*, 22(4-5): 45.
BAKER, Frank Collins. 1909a. A new species of *Lymnaea*. *Nautilus*, 22(12): 140-141.
BAKER, Frank Collins. 1909b. Note on *Planorbis binneyi* Tryon. *Nautilus*, 23(3): 41-42.
BAKER, Frank Collins. 1909c. Range of *Lymnaea umbilicata*. *Nautilus*, 23(6): 80.
BAKER, Frank Collins. 1909d. Mollusks from Kansas and Oklahoma. *Nautilus*, 23(7): 91-94.
BAKER, Frank Collins. 1910a. The ecology of the Skokie Marsh area, with special reference to the Mollusca. *Bull. Ill. State Lab. nat. Hist.*, 8(4): 437-499, pls. 6-24.

BAKER, Frank Collins. 1910b. A new variety of *Lymnaea stagnalis*. *Nautilus*, 23(9): 112-113; 23(10): 125-126.
BAKER, Frank Collins. 1910c. The ecology of the Skokie Marsh area with special reference to the Mollusca. *Trans. Ill. State Acad. Sci.*, 3: 106-108.
BAKER, Frank Collins. 1910d. Description of a new *Lymnaea*. *Nautilus*, 24(5): 58-60.
BAKER, Frank Collins. 1910e. Mollusks of Unionville, Connecticut. *Nautilus*, 24(6): 68-69.
BAKER, Frank Collins. 1911a. *The Lymnaeidae of North and Middle America, Recent and fossil.* Chicago Acad. Sci., Spec. Pub. 3, pp. i-xvi, 1-539, pls. 1-58.
BAKER, Frank Collins. 1911b. Mollusks of Wellesley Island and vicinity, St. Lawrence River. *Nautilus*, 25(6): 66-67.
BAKER, Frank Collins. 1911c. The molluscan fauna of Tomahawk Lake, Wisconsin. *Trans. Wis. Acad. Sci., Arts Lett.*, 17(1): 200-246, pls. 11-17.
BAKER, Frank Collins. 1912a. A new *Planorbis* from Michigan. *Nautilus*, 25(10): 118-120.
BAKER, Frank Collins. 1912b. Recent additions to the catalogue of Illinois Mollusca. *Trans. Ill. State Acad. Sci.*, 5: 143-145.
BAKER, Frank Collins. 1913a. A new *Lymnaea* from Montana. *Nautilus*, 26(10): 115-116.
BAKER, Frank Collins. 1913b. *Pseudogalba*, new name for *Simpsonia*. *Nautilus*, 26(10): 120.
BAKER, Frank Collins. 1913c. Notes on post-glacial Mollusca. I. Emmet County, Michigan. *Nautilus*, 27(1): 7-8.
BAKER, Frank Collins. 1913d. *Galba ferruginea* in Oregon. *Nautilus*, 27(2): 24.
BAKER, Frank Collins. 1913e. Mollusca of Wyoming County, N.Y. *Nautilus*, 27(5): 54.
BAKER, Frank Collins. 1913f. Notes on post-glacial Mollusca. II. Waukesha County, Wisconsin. *Nautilus*, 27(6): 68.
BAKER, Frank Collins. 1914a. Mollusks from Magician Lake, Cass and Van Buren counties, Michigan. *Nautilus*, 28(1): 8-10.
BAKER, Frank Collins. 1914b. Northern Idaho shells. *Nautilus*, 27(9): 104-106.
BAKER, Frank Collins. 1915a. On the classification of the Lymnaeidae. *Nautilus*, 29(2): 20-24.
BAKER, Frank Collins. 1915b. *Campeloma lewisii* Walker in Illinois. *Nautilus*, 29(3): 36.
BAKER, Frank Collins. 1915c. Mollusks from Berrien County, Michigan. *Nautilus*, 29(4): 47.
BAKER, Frank Collins. 1916a. The fresh-water Mollusca of Oneida Lake, New York. *Nautilus*, 30(1): 5-9.
BAKER, Frank Collins. 1916b. The relation of mollusks to fish in Oneida Lake. *N.Y. State Coll. For. Syracuse Univ., Tech. Pub.*, (4): 1-366, 1 map.
BAKER, Frank Collins. 1916c. Further notes on the post-glacial biota of Glacial Lake, Chicago. *Trans. Ill. State Acad. Sci.*, 7: 74-78.
BAKER, Frank Collins. 1917. Notes on *Acella haldemani* (Desh.) Binney. *Nautilus*, 30(12): 135-138.
BAKER, Frank Collins. 1918a. Further notes on the Mollusca of Oneida Lake, New York; the mollusks of lower South Bay. *Nautilus*, 31(3): 81-93.
BAKER, Frank Collins. 1918b. Notes on nidification in *Gillia* and *Amnicola*. *Nautilus*, 32(1): 19-23.
BAKER, Frank Collins. 1918c. The productivity of invertebrate fish food on the bottom of Oneida Lake, with special reference to mollusks. *N.Y. State Coll. For. Syracuse Univ., Tech. Pub.*, (9): 1-264, 2 unpaginated figs., 7 unpaginated tables.
BAKER, Frank Collins. 1918d. Post-glacial Mollusca from the marls of central Illinois. *J. Geol.*, 26(7): 659-671.
BAKER, Frank Collins. 1918e. The relation of shellfish to fish in Oneida Lake, New York. *N.Y. State Coll. For.*, Circ. 21, 17(27): 1-34, 12 pls.
BAKER, Frank Collins. 1919a. Description of a new species and variety of *Planorbis* from post-glacial deposits. *Nautilus*, 32(3): 1-3, pl. 7, figs. 4-6.
BAKER, Frank Collins. 1919b. Mollusks infested with parasitic worms. *Nautilus*, 32(3): 97-98.
BAKER, Frank Collins. 1919c. The ecology of North American Lymnaeidae. *Science*, 49(1274): 519-521.
BAKER, Frank Collins. 1919d. A new species of *Physa* from New York State. *Nautilus*, 33(1): 11-13.
BAKER, Frank Collins. 1919e. Fresh-water Mollusca from Colorado and Alberta. *Bull. Am. Mus. nat. Hist.*, 41(13): 527-539.
BAKER, Frank Collins. 1920a. Animal life and sewage in the Genesee River, New York. *Am. Nat.*, 54(631): 152-161.
BAKER, Frank Collins. 1920b. A new *Planorbis* from Illinois. *Nautilus*, 33(4): 123-125.
BAKER, Frank Collins. 1920c. *Physa smithiana*, new name for *Physa smithii*. *Nautilus*, 33(4): 142.
BAKER, Frank Collins. 1920d. Notes on a small collection of shells from Alaska. *Nautilus*, 34(2): 67-68.
BAKER, Frank Collins. 1920e. The effect of sewage and other pollution on animal life of rivers and streams. *Trans. Ill. State Acad. Sci.*, 13: 271-279.
BAKER, Frank Collins. 1922a. The importance of ecology in the interpretation of fossil faunas. *Ecology*, 2(4):

277-280.
BAKER, Frank Collins. 1922b. New species and varieties of Mollusca from Lake Winnebago, Wisconsin, with new records for this state. *Nautilus*, 35(4): 130-133; 36(1): 19-21.
BAKER, Frank Collins. 1922c. New lymnaeas from Wisconsin and Minnesota, with notes on shells from the latter state. *Nautilus*, 36(1): 22-25.
BAKER, Frank Collins. 1922d. The molluscan fauna of the Big Vermilion River, Illinois, with special reference to its modification as the result of pollution by sewage and manufacturing wastes. *Ill. biol. Monogr.*, 7(2): 1-126, pls. 1-15.
BAKER, Frank Collins. 1922e. Stream pollution. *Ecology*, 3(4): 340-341.
BAKER, Frank Collins. 1923a. Description of a new *Lymnaea* from Yellowstone Park. *Nautilus*, 36(4): 125-126.
BAKER, Frank Collins. 1923b. The use of molluscan shells by the Cahokia mound builders. *Trans. Ill. State Acad. Sci.*, 16: 328-334.
BAKER, Frank Collins. 1924a. A new *Physa* from California. *Nautilus*, 38(1): 15.
BAKER, Frank Collins. 1924b. The fauna of the Lake Winnebago region. A quantitative and qualitative survey with special reference to the Mollusca. *Trans. Wis. Acad. Sci., Arts Lett.*, 21: 109-146.
BAKER, Frank Collins. 1924c. Bottom and shore fauna in relation to fisheries. [Review.] *Ecology*, 5(4): 416-418.
BAKER, Frank Collins. 1926a. Nomenclatorial notes on American fresh-water Mollusca. *Trans. Wis. Acad. Sci., Arts Lett.*, 22: 192-205.
BAKER, Frank Collins. 1926b. New forms of *Planorbis* and *Lymnaea* with notes on other forms. *Nautilus*, 39(4): 116-121.
BAKER, Frank Collins. 1926c. Fresh-water Mollusca from White Lake, Oakland County, Michigan. *Nautilus*, 40(2): 49-52.
BAKER, Frank Collins. 1927a. New varieties of *Stagnicola* from Wisconsin and Wyoming. *Nautilus*, 40(3): 82-85.
BAKER, Frank Collins. 1927b. New varieties of *Helisoma antrosa* from Wisconsin and Minnesota. *Nautilus*, 40(3): 85-87.
BAKER, Frank Collins. 1927c. Description of a new species of *Lymnaea* from British Columbia. *Nautilus*, 41(1): 23-24.
BAKER, Frank Collins. 1927d. Molluscan associations of White Lake, Michigan: a study of a small inland lake from an ecological and systematic viewpoint. *Ecology*, 8(3): 353-370, pl. 4.
BAKER, Frank Collins. 1927e. A new variety of *Helisoma campanulata* from Michigan. *Nautilus*, 41(2): 49-51.
BAKER, Frank Collins. 1927f. The influence of a changed environment in the formation of new species and varieties. *Science*, 66(1714): 432.
BAKER, Frank Collins. 1928a. The American *Bithynia* not wholly an introduced species. *Trans. Ill. State Acad. Sci.*, 20: 56-62, 1 pl.
BAKER, Frank Collins. 1928b. Influence of a changed environment in the formation of new species and varieties. *Ecology*, 9(3): 271-283.
BAKER, Frank Collins. 1928c. The fresh water Mollusca of Wisconsin. *Wis. geol. nat. Hist. Surv.*, Bull. 70, Pt. 1, pp. i-xx, 1-507, pls. 1-28.
BAKER, Frank Collins. 1928d. The Mollusca of Chautauqua Lake, New York, with descriptions of a new variety of *Ptychobranchus* and of *Helisoma*. *Nautilus*, 42(2): 48-60.
BAKER, Frank Collins. 1929a. Certain anatomical features of the fresh-water mollusc, *Helisoma corpulenta* Say. *Trans. Am. microsc. Soc.*, 48(1): 44-47, pl. 8.
BAKER, Frank Collins. 1929b. Mollusca from Vermilion and Pelican lakes, Minnesota, with the description of a new variety of *Helisoma corpulenta*. *Nautilus*, 42(3): 95-97; 42(4): 131-136.
BAKER, Frank Collins. 1930a. The use of animal life by the mound-building Indians of Illinois. *Trans. Ill. State Acad. Sci.*, 22: 41-64, 5 pls.
BAKER, Frank Collins. 1930b. The molluscan fauna of the southern part of Lake Michigan and its relationship to old glacial Lake Chicago. *Trans. Ill. State Acad. Sci.*, 22: 186-194.
BAKER, Frank Collins. 1930c. On genus and species making. *Science*, 72(1854): 37-39.
BAKER, Frank Collins. 1930d. The variation of molluscan life during the Pleistocene and Recent time. *Nautilus*, 44(1): 21-24.
BAKER, Frank Collins. 1931a. Description of a new variety of *Valvata lewisii* Currier. *Nautilus*, 44(4): 119-121.
BAKER, Frank Collins. 1931b. The classification of the large planorbid snails of Europe and America. *Proc. zool. Soc. Lond.*, 1931(2): 575-592, pls. 1-6.

BAKER, Frank Collins. 1931c. Ecological relationship of the genus *Pomatiopsis* with special reference to *Pomatiopsis lapidaria. Ecology,* 12(3): 489-496.
BAKER, Frank Collins. 1932a. *Stagnicola apicina* and *Stagnicola walkeriana. Nautilus,* 45(4): 139-140.
BAKER, Frank Collins. 1932b. The ecology of Say's *Limnaeus elodes. Ecology,* 13(3): 286-289.
BAKER, Frank Collins. 1932c. New species and varieties of *Helisoma* and *Gyraulus* from Canada. *Nautilus,* 46(1): 6-9.
BAKER, Frank Collins. 1933a. The ecology of Say's *Limnaeus elodes. Nautilus,* 46(4): 139.
BAKER, Frank Collins. 1933b. Self fertilization and nidification in *Physa halei. Nautilus,* 47(1): 35.
BAKER, Frank Collins. 1933c. Studies on the bottom fauna of fresh-water lakes. *Science,* 78(2018): 190-191.
BAKER, Frank Collins. 1933d. New species of Lymnaeidae from British America. *J. Wash. Acad. Sci.,* 23(11): 520-523.
BAKER, Frank Collins. 1934a. Two new races and a new species of *Helisoma* from California. *Nautilus,* 47(4): 140-142.
BAKER, Frank Collins. 1934b. Two new Canadian lymnaeas. *Can. Field-Nat.,* 48(4): 69-70.
BAKER, Frank Collins. 1934c. The molluscan fauna of the great river valleys of Illinois. *Trans. Ill. State Acad. Sci.,* 26(3): 129.
BAKER, Frank Collins. 1934d. New Lymnaeidae from the United States and Canada: I. California, Oregon, and other western states. *Nautilus,* 48(1): 17-20.
BAKER, Frank Collins. 1934e. A new species of *Gyraulus* from Canada. *Can. Field-Nat.,* 48(8): 135.
BAKER, Frank Collins. 1935a. Land and freshwater Mollusca from North Star Lake and vicinity, Itasca County, Minnesota. *Am. midl. Nat.,* 16(3): 257-274.
BAKER, Frank Collins. 1935b. The generic position of *Planorbis umbilicatellus* with the description of a new group of Planorbidae. *Nautilus,* 49(2): 46-48.
BAKER, Frank Collins. 1935c. *Stagnicola elrodiana*, new name for *Limnaea montana* Elrod. *Nautilus,* 49(2): 64.
BAKER, Frank Collins. 1936a. The generic position of *Planorbis obstructus* Morelet. *Nautilus,* 59(3): 104.
BAKER, Frank Collins. 1936b. The freshwater mollusc *Helisoma corpulentum* and its relatives in Canada. *Natl. Mus. Can. Bull.,* (79): 1-37.
BAKER, Frank Collins. 1936c. New Lymnaeidae from the United States and Canada. II. Michigan, Minnesota, and Montana. *Nautilus,* 49(4): 127-140, pl. 7, figs. 7-9.
BAKER, Frank Collins. 1936d. The relationship of *Parapholyx. Nautilus,* 50(1): 30-31.
BAKER, Frank Collins. 1937. Mollusca from Prince Albert National Park, Saskatchewan. *Nautilus,* 50(4): 113-117.
BAKER, Frank Collins. 1939a. *Stagnicola bulimoides vancouverensis* nov. var. *Nautilus,* 52(4): 144.
BAKER, Frank Collins. 1939b. Land and freshwater Mollusca from western Ontario. *Can. J. Res.,* 17(D,4): 87-102.
BAKER, Frank Collins. 1940a. A new species of *Drepanotrema* and some preoccupied planorbid names. *Nautilus,* 54(3): 96-97, pl. 8.
BAKER, Frank Collins. 1940b. A new anatomical classification of the Planorbidae. *Am. malacol. Union 10th annu. Meet. 1940,* (11): 4-5.
BAKER, Frank Collins. 1941. A new race of *Parapholyx effusa. Nautilus,* 55(1): 16-17.
BAKER, Frank Collins. 1942a. *Lymnaea stagnalis* and *Lymnaea (Radix) auricularia. Nautilus,* 55(3): 105-106.
BAKER, Frank Collins. 1942b. Collecting and preserving fresh water snails. *Am. malacol. Union 11th annu. Meet. 1941,* (12): 5-9.
BAKER, Frank Collins. 1942c. Land and fresh water Mollusca of New Hampshire. *Am. midl. Nat.,* 27(1): 74-85.
BAKER, Frank Collins. 1945. *The molluscan family Planorbidae.* Univ. Illinois Press, Urbana. 530 pp.
BAKER, Frank Collins & BROOKS, Stanley T. 1935. New species and races of Lymnaeidae from Newfoundland. *Nautilus,* 49(1): 10-13.
BAKER, Frank Collins & CAHN, A.R. 1931. Freshwater Mollusca from central Ontario. *Natl. Mus. Can. Bull.,* (67): 41-64, pl. 2.
BAKER, Frank Collins & HENDERSON, Junius. 1929. *Fossaria perplexa* F.C. Baker and Junius Henderson. *Nautilus,* 42(3): 103-104.
BAKER, Frank Collins & HENDERSON, Junius. 1933. A new *Stagnicola* from Montana. *Nautilus,* 47(1): 30-32.
BAKER, H. Burrington. 1909. Key to the genera of Gastropoda of Michigan. *Eleventh Rep. Mich. Acad. Sci., 1909,* pp. 134-140.
BAKER, H. Burrington. 1910. Variation in *Lymnaea reflexa* Say, from Huron County. *Twelfth Rep. Mich. Acad. Sci., 1910,* pp. 60-63.

BAKER, H. Burrington. 1922. The Mollusca of Dickinson County, Michigan. *Occ. Pap. Mus. Zool. Univ. Mich.*, (111): 1-44, 1 map.
BAKER, H. Burrington. 1923. Notes on the radula of the Neritidae. *Proc. Acad. nat. Sci. Phila.*, 75: 117-178, pls. 9-16.
BAKER, H. Burrington. 1925. Anatomy of *Lanx*, a limpet-like lymnaeid mollusk. *Proc. Calif. Acad. Sci.*, 4th ser., 14(8): 143-169, pls. 11-14.
BAKER, H. Burrington. 1930. The Mollusca collected by the University of Michigan-Williamson Expedition in Venezuela. *Occ. Pap. Mus. Zool. Univ. Mich.*, (210): 1-94, pls. 1-33.
BAKER, H. Burrington. 1931. Notes on Physidae with descriptions of new species. By Wm. J. Clench (Occ. Papers Boston Soc. nat. Hist., 5: 301-315, 4 figs). [Review.] *Nautilus*, 44(3): 106.
BAKER, H. Burrington. 1942. *Goniobasis livescens* in Douglas Lake, Michigan. *Nautilus*, 56(1): 33-34.
BAKER, H. Burrington. 1946. Index to F.C. Baker's "The molluscan family Planorbidae." *Nautilus*, 59(4): 127-141.
BAKER, H. Burrington. 1947. *Amnicola* and *Euamnicola*. *Nautilus*, 60(3): 105-106.
BAKER, H. Burrington. 1956. Familial names for land operculates. *Nautilus*, 70(1): 28-31.
BAKER, H. Burrington. 1959. Measurements by C.B. Adams. *Nautilus*, 72(3): 106.
BAKER, H. Burrington. 1960a. Hydrobiidae or Truncatellidae? *Nautilus*, 74(1): 34-35.
BAKER, H. Burrington. 1960b. *Planorbina* (1843) vs. *Australorbis* (1934) vs. *Biomphalaria* (1910) vs. *Taphius* (1854). *Nautilus*, 74(1): 35-37.
BAKER, H. Burrington. 1960c. *Lustrica (Paludina)* Say, 1821 (Gastropoda): Proposed suppression under the plenary powers. Z.N.(S.) 730. *Bull. zool. Nomencl.*, 18(1/3): 146-148.
BAKER, H. Burrington. 1963. Paludomidae (Pleuroceridae). *Nautilus*, 77(1): 34-35.
BAKER, H. Burrington. 1964. Type land snails in the Academy of Natural Sciences of Philadelphia. Part III. Limnophile and thalassophile Pulmonata. Part IV. Land and fresh-water Prosobranchia. *Proc. Acad. nat. Sci. Phila.*, 116(4): 149-193.
BAKER, H. Burrington. 1967a. *Nitocris*. *Nautilus*, 80(3): 108.
BAKER, H. Burrington. 1967b. *Mudalia*. *Nautilus*, 81(1): 35.
BAKER, H. Burrington. 1967c. *Juga* and *Melasma*. *Nautilus*, 81(1): 36.
BARBOSA, Frederico S., BERRY, Elmer G., HARRY, Harold W., HUBENDICK, Bengt, MALEK, Emile A., PARAENSE, Wladimir Lobato, CHAMBERLAYNE, Earl C. & OLIVIER, Louis J. 1968. *A guide for the identification of the snail intermediate hosts of schistosomiasis in the Americas*. Pan Am. health Organ., Pan Am. sanit. Bur., Washington, D.C., Sci. Publ. No. 168, pp. i-ix, 1-122.
BARNES, D.W. 1823. On the genera *Unio* and *Alasmodonta*; with introductory remarks. *Am. J. Sci.*, 6(1): 107-127; 6(2): 258-280, pls. 1-8, 11-13. [Reprinted in *Sterkiana*, 1962, (6): 40-51; (7): 7-21, pls. 1-6.]
BARNHART, M. Christopher. 1978. Three introduced gastropods in Iowa. *Nautilus*, 92(3): 106-107.
BARTSCH, Paul. 1938. *Goniobasis* breeding. *Rep. Am. malacol. Union 1937*, (6): 3-5.
BARTSCH, Paul. 1949. Molluscan genetics: the role of hybridization, mutation, isolation, fixation and speciation in relation to taxonomy. *News Bull. annu. Rep. Am. malacol. Union 1948*, (15): 2-3.
BARTSCH, Paul. 1955. Hybridization among mollusks. *Annu. Rep. Am. malacol. Union 1954*, (21): 4.
BASCH, Paul F. 1959a. *Gundlachia* in Michigan. *Annu. Rep. Am. malacol. Union 1958*, (25): 13-14.
BASCH, Paul F. 1959b. Study on Ancylidae. *Annu. Rep. Am. malacol. Union 1958*, (25): 15.
BASCH, Paul F. 1959c. Status of the genus *Gundlachia* (Pulmonata, Ancylidae). *Occ. Pap. Mus. Zool. Univ. Mich.*, (602): 1-9.
BASCH, Paul F. 1959d. The anatomy of *Laevapex fuscus*, a freshwater limpet (Gastropoda: Pulmonata). *Misc. Pub. Mus. Zool. Univ. Mich.*, (108): 1-56.
BASCH, Paul F. 1959e. Studies on the development and reproduction of the fresh-water limpet, *Ferrissia shimekii* (Pilsbry). *Trans. Am. microsc. Soc.*, 78(3): 269-276.
BASCH, Paul F. 1960a. The Coosa revisited. *Annu. Rep. Am. malacol. Union 1959*, (26): 17.
BASCH, Paul F. 1960b. Anatomy of *Rhodacmea cahawbensis* Walker, 1917, a river limpet from Alabama. *Nautilus*, 73(3): 79-95.
BASCH, Paul F. 1962a. Radulae of North American ancylid snails. I. Subfamily Rhodacmeinae. *Nautilus*, 75(3): 97-101.
BASCH, Paul F. 1962b. Radulae of North American ancylid snails. II. Subfamily Neoplanorbinae. *Nautilus*, 75(4): 145-149.
BASCH, Paul F. 1962c. Radulae of North American freshwater limpet snails. III. *Ferrissia* and *Laevapex*. *Nautilus*, 76(1): 28-33.
BASCH, Paul F. 1963. A review of the Recent freshwater limpet snails of North America (Mollusca: Pulmonata).

Bull. Mus. comp. Zool. Harv. Coll., 129(8): 399-461.
BASCH, Paul F., BAINER, Philip & WILHM, Jerry. 1961. Some ecological characteristics of the molluscan fauna of a typical grassland situation in east central Kansas. *Am. midl. Nat.*, 66(1): 178-199.
BEASLEY, Clark W. & FULLINGTON, Richard W. 1978. Gastropods of Taylor County, Texas. *Sterkiana*, (69/70): 1-2.
BEAUCHAMP, Wm. M. 1887. Erosion of fresh-water shells. *Conchol. Exc.*, 1(9/10): 49.
BEAUCHAMP, Wm. M. 1891. Notes on familiar mollusks. *Nautilus*, 5(5): 52-53.
BEETLE, Dorothy E. 1954. Terrestrial and aquatic mollusks of Albany County, Wyoming. *Nautilus*, 67(4): 121-129.
BEETLE, Dorothy E. 1957. The Mollusca of Teton County, Wyoming. *Nautilus*, 71(1): 12-22.
BEETLE, Dorothy E. 1960a. Noteworthy records of Wyoming Mollusca. *Nautilus*, 73(4): 155-157.
BEETLE, Dorothy E. 1960b. Some mollusks from Manitoba, Canada. *Nautilus*, 74(2): 84.
BEETLE, Dorothy E. 1960c. Additional molluscan records for Albany County, Wyoming. *Nautilus*, 74(2): 84, iii.
BEETLE, Dorothy E. 1961a. Mollusca of the Big Horn Mountains. *Nautilus*, 74(3): 95-102.
BEETLE, Dorothy E. 1961b. A checklist of Wyoming Recent Mollusca. *Sterkiana*, (3): 1-9.
BEETLE, Dorothy E. 1962. Additions to Teton County, Wyoming, Mollusca. *Nautilus*, 76(2): 74.
BEETLE, Dorothy E. 1965. Molluscan fauna of some small ponds in Grand Teton National Park. *Nautilus*, 78(4): 125-130.
BEETLE, Dorothy E. 1968. *Laevapex fragilis* on the Outer Banks of North Carolina. *Nautilus*, 81(3): 107.
BEETLE, Dorothy E. 1970. A photographic record of snail activity. *Sterkiana*, (39): 1-8.
BEETLE, Dorothy E. 1973a. *Physa acuta* in Virginia. *Nautilus*, 87(1): 28.
BEETLE, Dorothy E. 1973b. Fresh-water mollusks from coastal Virginia. *Nautilus*, 87(1): 29.
BEETLE, Dorothy E. 1973c. A checklist of the land and freshwater mollusks of Virginia. *Sterkiana*, (49): 21-35.
BELL, Robert. 1859. On the natural history of the Gulf of St. Lawrence, and the distribution of the Mollusca of eastern Canada. *Can. Nat. Geol.*, 4: 197-220. [Pages on freshwater and land mollusks (pp. 212-216, 219-220) reproduced in *Sterkiana*, 1963, (10): 50-53.]
BELL, Robert. 1861. List of Recent land and fresh-water shells collected around lakes Superior and Huron in 1859-60. *Can. Nat. Geol.*, 6: 268-270. [Reprinted in *Sterkiana*, (8): 49-51.]
BEQUAERT, Joseph C. & MILLER, Walter B. 1973. *The mollusks of the arid Southwest, with an Arizona check list.* The Univ. Arizona Press, Tucson. xvi, 271 pp.
BERRY, Elmer G. 1931. Mollusca of Lamb's Canyon, Utah. *Nautilus*, 44(4): 113-114.
BERRY, Elmer G. 1938. The discovery of living *Pyrgulopsis letsoni* (Walker). *Rep. Am. malacol. Union 1937*, (6): 2.
BERRY, Elmer G. 1939. Conchology versus malacology in the family Amnicolidae. *Rep. Am. malacol. Union 1938*, (7): 8-9.
BERRY, Elmer G. 1943. The Amnicolidae of Michigan: distribution, ecology, and taxonomy. *Misc. Pub. Mus. Zool. Univ. Mich.*, (57): 1-68, pls. 1-9.
BERRY, Elmer G. 1947a. Members of the genus *Tropicorbis* found in Louisiana and Texas. *News Bull. annu. Rep. Am. malacol. Union 1946*, (13): 7.
BERRY, Elmer G. 1947b. [On collecting *Pomatiopsis chacei* in northern California.] *Min. conchol. Club. S. Calif.*, (73): 21.
BERRY, Elmer G. 1947c. Snails collected for the schistosomiasis investigations. *U.S. natl. Inst. Health Bull.*, (189): 55-69.
BERRY, Elmer G. 1948. A comparative study between members of the genera *Pomatiopsis* and *Oncomelania*. *News Bull. annu. Rep. Am. malacol. Union 1947*, (14): 14-15.
BERRY, Elmer G. 1949a. Morphological studies on *Tropicorbis havanensis* (Pfeiffer). *News Bull. annu. Rep. Am. malacol. Union 1948*, (15): 5-6.
BERRY, Elmer G. 1949b. Molluscacidal experiments in Brownsville, Texas. *News Bull. annu. Rep. Am. malacol. Union 1949*, (16): 3-4.
BERRY, Elmer G. 1949c. Recent investigations in medical malacology. *News Bull. annu. Rep. Am. malacol. Union 1949*, (16): 13-15.
BERRY, S. Stillman. 1909. The known Mollusca of San Bernardino County, California. *Nautilus*, 23(6): 73-79, pl. 6.
BERRY, S. Stillman. 1910. Mollusks of Unity, Maine. *Nautilus*, 24(6): 61-63.
BERRY, S. Stillman. 1913. A list of Mollusca from the Musselshell Valley, Montana. *Nautilus*, 26(11): 130-131.
BERRY, S. Stillman. 1916a. Notes on Mollusca of central Montana. *Nautilus*, 29(11): 124-128.
BERRY, S. Stillman. 1916b. Additional records of Mollusca from San Bernardino County, California. *Nautilus*, 30(4): 37-39.
BERRY, S. Stillman. 1916c. San Diego drift shells. *Nautilus*, 30(7): 83.

BERRY, S. Stillman. 1947. A new *Pyrgulopsis* from Oregon. *Nautilus*, 60(3): 76-78, pl. 7, figs. 6, 7 [correction to fig. nos. in *Nautilus*, 60(4): 148].

BICKEL, David. 1965. The role of aquatic plants and submerged structures in the ecology of a freshwater pulmonate snail, *Physa integra* Hald. *Sterkiana*, (18): 17-20.

BICKEL, David. 1966a. Stranded *Campeloma*. *Nautilus*, 79(3): 106-107.

BICKEL, David. 1966b. *Campeloma crassula* with reversed whorls. *Nautilus*, 79(3): 107-108.

BICKEL, David. 1967. Preliminary checklist of Recent and Pleistocene Mollusca of Kentucky. *Sterkiana*, (28): 7-20.

BICKEL, David. 1968a. *Goniobasis semicarinata* and *G. indianensis* in Blue River, Indiana. *Nautilus*, 81(4): 133-138.

BICKEL, David. 1968b. *Goniobasis curreyana lyoni*, a pleurocerid snail of west-central Kentucky. *Nautilus*, 82(1): 13-18.

BICKEL, David. 1968c. Checklist of the Mollusca of Tennessee. *Sterkiana*, (31): 15-39.

BILLUPS, A.C. 1902. *Angitrema verrucosa* at Lawrenceburg, Indiana. *Nautilus*, 16(7): 72.

BILLUPS, A.C. 1903. A proposed study of *Goniobasis*. *Nautilus*, 17(2): 22-24.

BINNEY, Wm.G. 1858. *The complete writings of Thomas Say, on the conchology of the United States*. H. Bailliere, New York. vi, 252 pp., pls. 1-75.

BINNEY, Wm. G. 1859. Catalogue of the terrestrial and fluviatile gasteropods inhabiting the continent of North America. First edition. *Smithson. misc. Collect.*, [unnumbered], pp. 1-15.

BINNEY, Wm. G. 1860a. [Geographical distribution of the terrestrial and fluviatile gastropods of North America.] *Proc. Acad. nat. Sci. Phila.*, 12: 49-50.

BINNEY, Wm. G. 1860b. Check lists of the shells of North America. No. 6. Fluviatile Gasteropoda. *Smithson. misc. Collect.*, 2(6): 7-13.

BINNEY, Wm. G. 1861. Catalogue of land and fresh water mollusks collected in British America by Messrs. Ross, Kennicott and Drexler, and deposited in the Smithsonian collection. *Proc. Acad. nat. Sci. Phila.*, 13: 330.

BINNEY, Wm. G. 1863. Synopsis of the species of air-breathing mollusks of North America. Smithsonian Institution ["Smithsonian Miscellaneous Collection 000"], Washington, D.C. pp. 1-12.

BINNEY, Wm. G. 1863-64. Bibliography of North American conchology previous to the year 1860. *Smithson. misc. Collect.*, 5: i-vii, 1-650 (Pt. 1, American authors); 9: i-iii, 1-298 (Pt. 2, 1864, Foreign authors).

BINNEY, Wm. G. 1865a. [Remarks on *Melania* and *Melantho*.] *Proc. Boston Soc. nat. Hist.*, 9: 327.

BINNEY, Wm. G. 1865b. Descriptions of new species of North American land and fresh water shells. *Am. J. Conchol.*, 1(1): 49-51, pl. 7.

BINNEY, Wm. G. 1865c. Land and fresh water shells of North America. Pt. 2. Pulmonata Limnophila and Thalassophila. *Smithson. misc. Collect.*, 7(2): i-ix, 1-161.

BINNEY, Wm. G. 1865d. Land and fresh-water shells of North America. Pt. 3. Ampullariidae, Valvatidae, Viviparidae, fresh-water Rissoidae, Cyclophoridae, Truncatellidae, fresh-water Neritidae, Helicinidae. *Smithson. misc. Collect.*, 7(3): i-viii, 1-120.

BINNEY, Wm.G. 1867. Notes sur quelques espèces de mollusques fluviatiles de l'Amérique du Nord. *J. Conchyliol.*, 15: 427-432.

BINNEY, Wm. G. 1870a. Note on *Vivipara lineata*, Valenciennes. *Ann. Lyc. nat. Hist. N.Y.*, 9: 295-297.

BINNEY, Wm. G. (Ed.). 1870b. *Report on the Invertebrata of Massachusetts*, published agreeably to an order of the Legislature. Second edition, comprising the Mollusca. By Augustus A. Gould. Wright and Potter, State Printers, Boston. v, 524 pp., pls. 16-27.

BINNEY, Wm. G. & BLAND, Thomas. 1870. Notes on lingual dentition of Mollusca. *Ann. Lyc. nat. Hist. N.Y.*, 9: 281-295.

BINNEY, Wm. G. & TRYON, George W., Jr. 1864. *The complete writings of Constantine Smaltz Rafinesque on Recent & fossil conchology*. Bailliere Bros., New York. 96 pp., index pp. 1-7, pls. 80-82.

BLATCHLEY, W.S. & DANIELS, L.E. 1903. On some Mollusca known to occur in Indiana. A supplementary paper to Call's catalogue. *27th annu. Rep. Dept. Geol. nat. Res. Indiana, 1902*, pp. 577-652, 678-680 (index), pls. 1-3.

BOER, H.H. & LEVER, J. 1959. On the anatomy of the circulatory system in *Ferrissia shimekii* (Ancylidae, Pulmonata); especially on the blood supply of the central nervous system. *K. Ned. Akad. Wet. (Amsterdam)*, ser. C, 62(1): 76-83.

BOETERS, Hans D. 1974. *Horatia* Bourguignat, *Plagigeyeria* Tomlin und *Litthabitella* Boeters. *Arch. Molluskenkd.*, 104(1/3): 85-92.

BOLE, J. 1970. Beitrag zur Kenntnis der Anatomie und Taxonomie der unterirdischen Hydrobiiden. *Diss. Acad. Sci. Art. Slovenica*, 13: 85-111.

BOVBJERG, R.V. 1965. Feeding and dispersal in the snail *Stagnicola reflexa* (Basommatophora: Lymnaeidae). *Malacologia,* 2(2): 199-207.
BRANSON, Branley A. 1960. *Gyraulus arizonensis* in Texas. *Nautilus,* 74(1): 37-38.
BRANSON, Branley A. 1961a. Recent Gastropoda of Oklahoma. Part II. Distribution, ecology and taxonomy of fresh-water species, with description of *Helisoma travertina* sp. nov. *Okla. State Univ. Pub.,* Arts & Sci. Stud., Biol. Stud. Ser., (6), 58(17): 1-72.
BRANSON, Branley A. 1961b. Notes on some gastropods from northern Louisiana. *Proc. La. Acad. Sci.,* 24: 24-30.
BRANSON, Branley A. 1962a. *Viviparus subpurpureus* in Oklahoma. *Nautilus,* 75(3): 124.
BRANSON, Branley A. 1962b. Some distribution records for Louisiana, Arkansas, and Missouri Gastropoda. *Sterkiana,* (7): 5-6.
BRANSON, Branley A. 1963. A note on molluscan zoogeography: distribution of *Gyraulus arizonensis* (Pilsbry and Ferriss). *Southwest. Nat.,* 8(1): 51-52.
BRANSON, Branley A. 1969a. Snail records from various southern, eastern and middle states. *Sterkiana,* (35): 1-4.
BRANSON, Branley A. 1969b. Distribution notes on western and southern snails. *Sterkiana,* (36): 21.
BRANSON, Branley A. 1970. *Juga, Oxytrema* and *Mudalia,* and a correction. *Sterkiana,* (39): 9-10.
BRANSON, Branley A. 1971a. Locality records for Oklahoma Mollusca. *Sterkiana,* (41): 35-40.
BRANSON, Branley A. 1971b. Variation in the shell of *Mudalia potosiensis* (Lea) (Pleuroceridae) from a single locality. *Nautilus,* 85(1): 26-30.
BRANSON, Branley A. 1977a. Freshwater and terrestrial Mollusca of the Olympic Peninsula, Washington. *Veliger,* 19(3): 310-330.
BRANSON, Branley A. 1977b. The Chinese Apple Snail, *Cipangopaludina chinensis,* on Orcas Island, Washington. *Nautilus,* 91(2): 76-77.
BRANSON, Branley A. & BATCH, Donald L. 1969. Notes on exotic mollusks in Kentucky. *Nautilus,* 82(3): 102-106.
BRANSON, Branley A. & BATCH, Donald L. 1971. Annotated distribution records for Kentucky Mollusca. *Sterkiana,* (43): 1-9.
BRANSON, Branley A. & HEARD, William R. 1959. Snails from upper peninsula of Alaska with feeding habits of Brooks Lake fishes. *Nautilus,* 73(1): 14-16.
BRANSON, Branley A. & PETERS, Lewis. 1964. *Limax maximus* and *Ferrissia meekiana* in Oklahoma. *Nautilus,* 77(3): 107-108.
BRETET, Roger & CARSWELL, Edwin J. 1952. A rich locality in the New York City area. *Nautilus,* 65(3): 100-102.
BRISCOE, M.S. 1941. The growth of the radula in *Goniobasis virginica* (Gmelin). *Rep. Am. malacol. Union 1940,* (9): 4.
BRISCOE, M.S. 1963. A survey of land and freshwater snails in Jefferson County, West Virginia. *Sterkiana,* (9): 41-48.
BROOKS, Stanley T. 1931. The gastropod family Pleuroceridae in Pennsylvania. *Nautilus,* 45(2): 58-64.
BROOKS, Stanley T. 1935. *Aplexa hypnorum pilsbryi* n. subsp. *Nautilus,* 48(3): 100.
BROOKS, Stanley T. 1936. Some molluscs from Utah. *Nautilus,* 50(1): 13-14.
BROOKS, Stanley T. & MacMILLAN, Gordon K. 1940. New Gastropoda from West Virginia. *Nautilus,* 53(3): 95-97, pl. 12, figs. 2, 3.
BROOKS, William Keith. 1907. The origin of the lung in *Ampullaria. Nautilus,* 21(2): 11-12.
BROOKS, William Keith & McGLONE, Bartjis. 1908. The origin of the lung in *Ampullaria. Carnegie Inst. Wash.,* (102): 95-104.
BROWN, Kenneth M. 1980. Life history evolution in temporary versus permanent ponds: *Lymnaea palustris* and *L. stagnalis. West. Soc. Malacol. annu. Rep.,* 12: 11.
BRYANT, Owen. 1908. A note on the type locality of *Paludestrina salsa,* at Cohasset, Mass. *Nautilus,* 22(8): 82.
BRYCE, George W., Jr. 1970. Rediscovery of the limpet, *Acroloxus coloradensis* (Basommatophora: Acroloxidae), in Colorado. *Nautilus,* 83(3): 105-108.
BUCCI, Dominique A. 1974. *Viviparus malleatus* in Montreal, Canada. *Nautilus,* 88(2): 55.
BURCH, John B. 1952. A preliminary list of the Mollusca of Hanover County, Virginia. *Nautilus,* 66(2): 60-63.
BURCH, John B. 1959. Chromosomes of aquatic pulmonate snails (Basommatophora). *Annu. Rep. Am. malacol. Union 1958,* (25): 9-10.
BURCH, John B. 1960a. Chromosomes of *Pomatiopsis* and *Oncomelania. Annu. Rep. Am. malacol. Union 1959,* (26): 15-16.
BURCH, John B. 1960b. *Some snails and slugs of quarantine significance to the United States.* Agric. res. Serv.,

U.S. Dept. Agric., 82(1): 1-73. [Reprinted in *Sterkiana*, 1960, (2): 13-53; Nav. med. Field res. Lab., Camp Lejeune, N.C., 1967, pp. 1-73.]

BURCH, John B. 1960c. The chromosomes of *Gyraulus circumstriatus*, a freshwater snail. *Nature*, 186(4723): 497-498.

BURCH, John B. 1960d. Chromosome morphology of aquatic pulmonate snails (Mollusca: Gastropoda). *Trans. Am. microsc. Soc.*, 79(4): 451-461.

BURCH, John B. 1960e. Chromosome studies of aquatic pulmonate snails. *Nucleus*, 3(2): 177-208, 4 pls.

BURCH, John B. 1962. Notes on the classification of freshwater limpets. *Annu. Rep. Am. malacol. Union 1962*, (29): 8-9.

BURCH, John B. 1965. Chromosome numbers and systematics in euthyneuran snails. *Proc. 1st Eur. malacol. Congr.*, 1962, pp. 215-241. Conchol. Soc. G. B. Irel. and Malacol. Soc. Lond.

BURCH, John B. 1967. *Some snails and slugs of quarantine significance to the United States.* Nav. med. Field res. Lab., Camp Lejeune, N.C. pp. 1-73.

BURCH, John B. 1968a. *Erinna newcombi* of Hawaii and *Limnaea onychia* of Japan. *Malacol. Rev.*, 1: 15-30.

BURCH, John B. 1968b. Morphology vs. immunology. *Annu. Rep. Am. malacol. Union 1968*, (35): 25-28.

BURCH, John B. 1973. A comparative study of some Polish and American Lymnaeidae: an assessment of phylogenetic characters. *Malacologia*, 14(1/2): 221-222.

BURCH, John B. 1974. Relationships of Eurasian-North American *Ancylus* to the North American *Rhodacmea* (Pulmonata: Ancylidae) as shown by apical shell sculpture. *Malacol. Rev.*, 7(1/2): 127-132.

BURCH, John B. 1975. Scanning electron microscope studies of the shell apices of freshwater limpets. *Malacol. Rev.*, 8(1/2): 117-118.

BURCH, John B. 1978. An outline of classification of the freshwater gastropods of North America (north of Mexico). *J. Conchyliol.*, 115(1/2): 1-9.

BURCH, John B. 1979. Genera and subgenera of Recent freshwater gastropods of North America (north of Mexico). *Malacol. Rev.*, 12(1/2): 97-100.

BURCH, John B. 1980. Further comment on the proposed designation of a type species for *Pleurocera* Rafinesque, 1818, Z.N.(S) 83. *Bull. zool. Nomencl.*, 36(4): 196-197.

BURCH, John B. & AYERS, P.A. 1973. Breeding experiments with *Stagnicola elodes* and *S. emarginata. Malacol. Rev.*, 6(1): 51-52.

BURCH, John B., BASCH, Paul F. & BUSH, Linda L. 1960. Chromosome numbers in ancylid snails. *Rev. Port. Zool. Biol. geral*, 2(3/4): 199-204.

BURCH, John B. & BUSH, Linda L. 1960. Chromosomes of *Physa gyrina* Say (Mollusca: Pulmonata). *J. Conchyliol.*, 100(2): 49-54.

BURCH, John B. & LINDSAY, G.K. 1968. An immunological approach to lymnaeid systematics. *Annu. Rep. Am. malacol. Union 1968*, (35): 22-23.

BURCH, John B. & LINDSAY, G.K. 1972. The systematic position of "*Lymnaea*" *bulimoides. Malacol. Rev.*, 5(1): 16-17.

BURCH, John B. & LINDSAY, G.K. 1973a. Taxonomic groupings in the Lymnaeidae. *Bull. Am. malacol. Union Inc. 1973*, (40): 15.

BURCH, John B. & LINDSAY, G.K. 1973b. Apparent hybrids from a laboratory culture of *Lymnaea stagnalis* and *Bulimnea megasoma. Malacol. Rev.*, 6(1): 61-62.

BURCH, John B., LINDSAY, G.K. & LoVERDE, P.T. 1971. A comparative study of some Polish and American Lymnaeidae: an assessment of phylogenetic characters. *Zool. zh.*, 50(8): 1158-1168.

BURCH, John B. & LoVERDE, P.T. 1974. Apical shell sculpture of some North American freshwater limpets (Mollusca: Basommatophora). *Occ. Pap. Moll.*, Harv. Univ., 5(50): 39-49.

BURCH, John B. & THOMPSON, Fred G. 1958. Taxonomic characters and infraspecific variation in mollusks. *Syst. Zool.*, 7(1): 48.

BURCH, John B. & TOTTENHAM, John L. 1980. North American freshwater snails. *Trans. POETS Soc.*, 1(3): 81-215.

BURCH, John B. & VAN DEVENDER, Amy Shrader. 1980. Identification of eastern North American land snails. The Prosobranchia, Opisthobranchia and Pulmonata (Actophila). *Trans. POETS Soc.*, 1(2): 33-80.

BURCH, Paul R. 1950. Mollusks. *In: The James River basin. Past, present and future.* Va. Acad. Sci., Richmond. 843 pp., 4 maps [pp. 129-137].

BURDSAL, L.A., DOHERTY, Wm., et al. 1876. *Publications of the O. G. B.* [Our Geological Boys]. *III. Mollusca of Cincinnati.* L.A. Burdsal, Printer, Cincinnati. 14 pp.

BURKY, Albert J. 1971. Biomass turnover, respiration, and interpopulation variation in the stream limpet *Ferrissia rivularis* (Say). *Ecol. Monogr.*, 41(3): 235-251.

CAFFREY, G.W. 1911. The molluscan fauna of Northampton County, Pennsylvania. *Nautilus*, 25(3): 26-29.
CALKINS, W.W. 1874. The land and fresh water shells, of La Salle County, Ills. *Proc. Ottawa Acad. nat. Sci.*, pp. 1-48.
CALKINS, W.W. 1878. Multiplication of species in the families Unionidae and Strepomatidae. *Valley Nat.*, 1: 3.
CALKINS, W.W. 1880. New freshwater Mollusca. *Valley Nat.*, 2(1): 6.
CALL, R. Ellsworth. 1880. Geology and natural history of Freemont County, Iowa. *In: History of Freemont County, Iowa.* Des Moines, Iowa. pp. 1-37 [Mollusca, pp. 34-36].
CALL, R. Ellsworth. 1882. Note on the geographical distribution of certain mollusks. *Am. Nat.*, 16: 400-402.
CALL, R. Ellsworth. 1883. Note on the genus *Campeloma* of Rafinesque. *Am. Nat.*, 17(6): 603-608.
CALL, R. Ellsworth. 1884. On the Quaternary and Recent Mollusca of the Great Basin, with descriptions of new forms. *Bull. U.S. geol. Surv.*, (11): 367-420, pls. 2-6.
CALL, R. Ellsworth. 1886a. On the genus *Campeloma* Rafinesque, with a revision of the species Recent and fossil. *Bull. Washburn Coll. Lab. nat. Hist.*, 1(5): 149-165, pls. 3-6.
CALL, R. Ellsworth. 1886b. Description of a new strepomatid mollusk of the genus *Goniobasis* (*G. ozarkensis*). *Bull. Washburn Coll. Lab. nat. Hist.*, 1(7): 189-190, pl. 7, figs. 1-10.
CALL, R. Ellsworth. 1886c. On certain Recent, Quaternary and new fresh water Mollusca. *Proc. Davenport Acad. nat. Sci.*, 5: 1-8, pl. 1.
CALL, R. Ellsworth. 1888a. On a new post-Pleiocene limnaeid. *Am. Geol.*, 1: 146-148, figs. 1-6.
CALL, R. Ellsworth. 1888b. On the gross anatomy of *Campeloma. Am. Nat.*, 22: 491-497, pl. 7, figs. 1-3.
CALL, R. Ellsworth. 1890. Notes on the gross anatomy of *Campeloma. Proc. Iowa Acad. Sci., 1887-89*, 1: 16-17.
CALL, R. Ellsworth. 1894a. A contribution to the knowledge of Indiana Mollusca. *Proc. Indiana Acad. Sci., 1893*, pp. 140-156.
CALL, R. Ellsworth. 1894b. On the geographic and hypsometric distribution of North American Viviparidae. *Am. J. Sci.*, 48: 132-141, 1 map.
CALL, R. Ellsworth. 1895. The Strepomatidae of the Falls of the Ohio. [Abstract.] *Proc. Indiana Acad. Sci., 1894*, pp. 140-143.
CALL, R. Ellsworth. 1896a. Second contribution to a knowledge of Indiana Mollusca. *Proc. Indiana Acad. Sci., 1895*, pp. 135-146.
CALL, R. Ellsworth. 1896b. On a small collection of mollusks from northern Indiana [also listed on reprint cover as "Indiana University Biological Station report on Mollusca"]. *Proc. Indiana Acad. Sci., 1895*, pp. 246-250.
CALL, R. Ellsworth. 1896c. Fishes and shells of the Falls of the Ohio. *In: Memorial history of Louisville, Kentucky.* Vol. 1, pp. 9-20.
CALL, R. Ellsworth. 1897. The hydrographic basins of Indiana and their molluscan fauna. *Proc. Indiana Acad. Sci., 1896*, pp. 247-258, 1 map.
CALL, R. Ellsworth. 1900. A descriptive illustrated catalogue of the Mollusca of Indiana. *24th annu. Rep. Dept. Geol. nat. Res. Indiana, 1899*, pp. 335-535, pls. 1-78.
CALL, R. Ellsworth & BEECHER, C.E. 1884. Notes on a Nevada shell (*Pyrgula nevadensis*). *Am. Nat.*, 18: 851-855.
CALL, R. Ellsworth & BEECHER, C.E. 1886. Description of a new rissoid mollusk (*Bythinella aldrichi*). *Bull. Washburn Coll. Lab. nat. Hist.*, 1(7): 190-192, pl. 7, figs. 11-14.
CALL, R. Ellsworth & PILSBRY, Harry [Henry] A. 1886. On *Pyrgulopsis*, a new genus of rissoid mollusk, with descriptions of two new forms. *Proc. Davenport Acad. nat. Sci.*, 5: 9-14, pl. 2.
CALOW, P. 1978. The evolution of life-cycle strategies in fresh-water gastropods. *Malacologia*, 17(2): 351-364.
CARPENTER, Horace F. 1887. The shell-bearing Mollusca of Rhode Island. *Conchol. Exc.*, 1(12): 76-78; 2(1): 2-3; 2(2): 18-20.
CARPENTER, Horace F. 1889a. *A catalogue of the shell-bearing Mollusca of Rhode Island.* Second edition. J.A. & R.A. Reid, Printers, Providence. 7 pp.
CARPENTER, Horace F. 1889b. The shell-bearing Mollusca of Rhode Island. *Nautilus*, 3(1): 11-12; 3(2): 21-23; 3(3): 32-33; 3(8): 92-95.
CARPENTER, Horace F. 1889c. Notes on *Valvata (Lyogyrus) brownii. Nautilus*, 3(6): 67.
CARPENTER, Horace F. 1902. The shell-bearing Mollusca of Rhode Island. *Nautilus*, 16(2): 17-18.
CARPENTER, Horace F. 1905. New locality for *Lyogyrus brownii* Carpenter. *Nautilus*, 19(4): 47-48.
CARPENTER, Philip P. 1872. The mollusks of western North America. *Smithson. misc. Collect.*, (252): i-xii, 1-325, index (108 pp.).
CASE, William. 1847. Description of two new species of shells. *Ann. Mag. nat. Hist.*, 19: 358.
CHACE, E.P. 1934. [*Lymnaea bulimoides bulimoides.*] *Rep. Am. malacol. Union 1934*, (3): 4.
CHAMBERLAIN, Norman A. 1958. Life history studies of *Campeloma decisum. Nautilus*, 72(1): 22-29, pl. 3, fig. 7.

CHAMBERLAIN [CHAMBERLIN], Ralph V. 1933. Observations on *Stagnicola kingi* (Meek), living and extinct. *Nautilus*, 46(3): 97-100.
CHAMBERLIN, Ralph V. & BERRY, Elmer G. 1929. Notes on the Mollusca of southeastern Utah. *Nautilus*, 42(4): 123-125.
CHAMBERLIN, Ralph V. & JONES, David. 1929. A descriptive catalogue of the Mollusca of Utah. *Bull. Univ. Utah*, 19(4): i-ix, 1-203, 1 map.
CHAMBERLIN, Ralph V. & ROSCOE, Ernest J. 1948. Check list of the Recent Utah Mollusca. *Bull. Univ. Utah*, 39(2): 1-16.
CHAMBERS, Steven M. 1978. An electrophoretically detected sibling species of "*Goniobasis floridensis*" (Mesogastropoda: Pleuroceridae). *Malacologia*, 17(1): 157-162.
CHEATUM, Elmer P. 1935. Gastropods of the Davis Mountains vicinity in West Texas. *Nautilus*, 48(4): 112-116, pl. 5.
CHEATUM, Elmer P. & FULLINGTON, Richard W. 1971. The aquatic and land Mollusca of Texas. Supplement. Keys to the families of the Recent land and fresh-water snails of Texas. *Dallas Mus. nat. Hist. Bull.*, (1): i-iii, 1-18.
CHEATUM, Elmer P., FULLINGTON, Richard W. & PRATT, Lloyd. 1972. Molluscan records from West Texas. *Sterkiana*, (46): 6-10.
CHENU, J.C. 1859. *Manuel de Conchyliologie et de Paléontologie Conchyliologique*. Librairie Victor Masson, Paris. Vol. 1. 508 pp. [pp. 286-296].
CLAMPITT, Philip T. 1970. Comparative ecology of the snails *Physa gyrina* and *Physa integra* (Basommatophora: Physidae). *Malacologia*, 10(1): 113-151.
CLAMPITT, Philip T. 1972. Seasonal migrations and other movements in Douglas Lake pulmonate snails. *Malacol. Rev.*, 5(1): 11-12.
CLAMPITT, Philip T. 1973. Substratum as a factor in the distribution of pulmonate snails in Douglas Lake, Michigan. *Malacologia*, 12(2): 379-399.
CLAMPITT, Philip T. 1974a. Seasonal migratory cycle and related movements of the fresh-water pulmonate snail, *Physa integra*. *Am. midl. Nat.*, 92(2): 275-300.
CLAMPITT, Philip T. 1974b. How fast is a snail's pace? Active and passive dispersal of *Physa integra* in Douglas Lake, Michigan. *Malacol. Rev.*, 8(1/2): 121.
CLAPP, Geo. H. 1913. *Gundlachia hjalmarsoni* Pfr. in the Rio Grande, Texas. *Nautilus*, 27(7): 77-78, pl. 4, figs. 6-8.
CLAPP, William F. 1913. *Radix auricularia* at Cambridge, Massachusetts. *Nautilus*, 26(10): 116-117.
CLAPP, William F. 1921. Eggs and young of the river limpet, *Ancylus fuscus* C.B. Adams. *Occ. Pap. Boston Soc. nat. Hist.*, 5: 5-10, pl. 2.
CLARK, Clarence F. 1962a. Records of gastropods collected in western Ohio. *Sterkiana*, (6): 15-22.
CLARK, Clarence F. 1962b. Records of gastropods collected in western Ohio -- a correction. *Sterkiana*, (7): 51.
CLARKE, Arthur H., Jr. 1961. Fresh-water mollusks of the James Bay watershed – a progress report. *Annu. Rep. Am. malacol. Union 1961*, (28): 11-12.
CLARKE, Arthur H., Jr. 1964. Freshwater mollusks of the Hudson Bay watershed, distribution patterns and determinant influences. *Annu. Rep. Am. malacol. Union 1964*, (31): 8-9.
CLARKE, Arthur H., Jr. 1967. Zoogeographic and evolutionary patterns in northern Lymnaeidae and Planorbidae. *Annu. Rep. Am. malacol. Union 1967*, (34): 21-22.
CLARKE, Arthur H., Jr. 1969. Some aspects of adaptive radiation in Recent freshwater mollusks. *Malacologia*, 9(1): 163.
CLARKE, Arthur H., Jr. 1970a. Discussion of Dr. Stansbery's paper [on rare and endangered eastern freshwater mollusks of the Mississippi and St. Lawrence River systems, in Am. malacol. Union Symp. Rare and Endangered Mollusks]. *Malacologia*, 10(1): 21-22.
CLARKE, Arthur H., Jr. 1970b. On *Acroloxus coloradensis* (Henderson) (Gastropoda, Basommatophora) in eastern Canada. *Publ. Zool., natl. Mus. Can.*, (2): i-vi, 1-13.
CLARKE, Arthur H., Jr. 1973. The freshwater mollusks of the Canadian Interior Basin. *Malacologia*, 13(1/2): 1-509.
CLARKE, Arthur H., Jr. 1976. Endangered freshwater mollusks of northwestern North America. *Bull. Am. malacol. Union Inc., 1976*, (43): 18-19.
CLARKE, Arthur H., Jr. 1978. The Asian Apple Snail, *Cipangopaludina chinensis* (Viviparidae) in Oneida Lake, New York. *Nautilus*, 92(3): 134.
CLARKE, Arthur H., Jr. 1979. Gastropods as indicators of trophic lake stages. *Nautilus*, 93(4): 138-142.
CLARKE, W.T., Jr. 1938. List of molluscs from drift debris of Paladora Creek, Texas. *Nautilus*, 52(1): 14-15.

CLENCH, William J. 1924. A new species of *Physa* from Texas. *Nautilus*, 38(1): 12-13.
CLENCH, William J. 1925a. Notes on the genus *Physa* with descriptions of three new species. *Occ. Pap. Mus. Zool. Univ. Mich.*, (161): 1-10, pl. 1.
CLENCH, William J. 1925b. A short list of land and freshwater mollusks from southeastern Texas. *Nautilus*, 39(1): 11-12.
CLENCH, William J. 1926a. Three new species of *Physa*. *Occ. Pap. Mus. Zool. Univ. Mich.*, (168): 1-6, pl. 1.
CLENCH, William J. 1926b. Some notes and a list of shells of Rio, Kentucky. *Nautilus*, 40(1): 7-12; 40(2): 65-67.
CLENCH, William J. 1928. *Io fluvialis turrita* Anthony. *Nautilus*, 42(1): 36.
CLENCH, William J. 1930a. Notes on Physidae with descriptions of new species. *Occ. Pap. Boston Soc. nat. Hist.*, 5: 301-315.
CLENCH, William J. 1930b. A note on the habitat of *Viviparus subpurpureus*. *Nautilus*, 44(2): 70.
CLENCH, William J. 1932a. Thomas Say – his first paper on American mollusks. *Nautilus*, 45(4): 112.
CLENCH, William J. 1932b. *Viviparus japonicus* v. Mart. *Nautilus*, 45(4): 136.
CLENCH, William J. 1934. *Physa acuta* Draparnaud in Massachusetts. *Nautilus*, 48(1): 33.
CLENCH, William J. 1935. *Physa ampullacea* 'Gould' Binney. *Nautilus*, 49(1): 30-32.
CLENCH, William J. 1937. *Physa canadensis* Whiteaves. *Nautilus*, 50(4): 143-144.
CLENCH, William J. 1940a. *Pyrgulopsis nevadensis* Stearns in Oregon. *Nautilus*, 53(4): 137.
CLENCH, William J. 1940b. Another record for *Viviparus malleatus* in Massachusetts. *Nautilus*, 54(2): 69-70.
CLENCH, William J. 1946. Republication of descriptions of North American mollusks by Verkrüzen, Kurtz and De Tar and Beecher. *Nautilus*, 60(2): 69-71.
CLENCH, William J. 1954. *Mesodon thyroidus* (Say) in Florida. *Nautilus*, 68(1): 23-24.
CLENCH, William J. 1955. *Melania cancellata* Say. *Nautilus*, 68(3): 107.
CLENCH, William J. 1959a. Mollusca. *In:* Edmondson, W.T. (Ed.), *Fresh-water biology*. Second edition. John Wiley and Sons, New York. 1248 pp. [pp. 1117-1160].
CLENCH, William J. 1959b. Method used by C.B. Adams in describing and measuring shells. *Nautilus*, 72(3): 105-106.
CLENCH, William J. 1962a. A catalogue of the Viviparidae of North America with notes on the distribution of *Viviparus georgianus* Lea. *Occ. Pap. Moll.*, 2(27): 261-287.
CLENCH, William J. 1962b. New records of the genus *Lioplax*. *Occ. Pap. Moll.*, 2(27): 288.
CLENCH, William J. 1965a. A new species of *Lithasia* from Mississippi. *Nautilus*, 79(1): 30-33.
CLENCH, William J. 1965b. A new species of *Clappia* from Alabama. *Nautilus*, 79(1): 33-34, fig. 2 (p. 32).
CLENCH, William J. 1965c. Supplement to the North American genus *Lioplax*. *Occ. Pap. Moll.*, 2(34): 426.
CLENCH, William J. 1969. *Melanoides tuberculata* (Muller) in Florida. *Nautilus*, 83(2): 72.
CLENCH, William J. & BOSS, Kenneth J. 1967. Freshwater Mollusca from James River, Va. and a new name for *Mudalia* of authors. *Nautilus*, 80(3): 99-102.
CLENCH, William J. & FULLER, Samuel L.H. 1965. The genus *Viviparus* (Viviparidae) in North America. *Occ. Pap. Moll.*, 2(32): 385-412.
CLENCH, William J. & RUSSELL, Henry D. 1940. Freshwater shells of New Hampshire. *Nautilus*, 54(2): 52-53.
CLENCH, William J. & TURNER, Ruth D. 1955. The North American genus *Lioplax* in the family Viviparidae. *Occ. Pap. Moll.*, 2(19): 1-20.
CLENCH, William J. & TURNER, Ruth D. 1956. Freshwater mollusks of Alabama, Georgia, and Florida from the Escambia to the Suwannee River. *Bull. Fla. State Mus. (biol. Sci.)*, 1(3): 97-239. [Index published privately by R.I. Johnson, 1972.]
CLENCH, William J. & TURNER, Ruth D. 1962. New names introduced by H.A. Pilsbry in the Mollusca and Crustacea. *Acad. nat. Sci. Phila. spec. Publ.*, (4): 1-218.
CLESSIN, S. 1878-86. Die Familie der Limnaeiden enthalten die Genera *Planorbis, Limnaeus, Physa* und *Amphipeplea*. *In: Systematisches Conchylien-Cabinet von Martini und Chemnitz*. Bauer und Raspe, Nürnberg. 1(17): 29-36a, 63-430, pls. 11-15, 17-55.
CLESSIN, S. 1880-82. Die Familie der Ancylinen. In Abbildungen nach der Natur mit Beschreibungen. *In: Systematisches Conchylien-Cabinet von Martini und Chemnitz*. Bauer und Raspe, Nürnberg. 1(6): 1-80, pls. 1-9.
CLESSIN, S. 1881a. Die Gruppe der *Limnaea truncatula* L. *Malakozool. Bl.*, n.s., 3: 77-85.
CLESSIN, S. 1881b. Diagnoses novarum specierum generis *Ancyli*. *Malakozool. Bl.*, n.s., 3: 159.
CLESSIN, S. 1882. Neue Arten. *Malakozool. Bl.*, n.s., 5: 187-193, pl. 4.
COCKERELL, T.D.A. 1902. Notes on *Limnaea*. *Nautilus*, 16(8): 95-96.
COCKERELL, T.D.A. 1939. *Physa* on the California islands. *Nautilus*, 52(4): 138.
COLTON, Harold Sellers. 1908. Some effects of environment on the growth of *Lymnaea columella* Say. *Proc. Acad. nat. Sci. Phila.*, 60: 410-448.

COLTON, Harold Sellers. 1915a. On classification in general and the genus *Lymnaea* in particular. *Nautilus*, 28(10): 116-119.

COLTON, Harold Sellers. 1915b. A provisional key to the subgenera and species of North American lymnaeids. *Nautilus*, 28(10): 119-120.

CONRAD, Timothy A. 1834a. *New fresh water shells of the United States, with coloured illustrations, and a monograph of the genus* Anculotus *of Say; also a synopsis of the American naiades.* Judah Dobson, Philadelphia. 76 pp., pls. 1-8.

CONRAD, Timothy A. 1834b. Description of some new species of fresh water shells from Alabama, Tennessee, & c. *Am. J. Sci. Arts*, 25(2): 338-343, pl. 1. [Reprinted in *Sterkiana*, 1963, (9): 49-52.]

CONRAD, Timothy A. 1835. Appendix [to *New fresh water shells of the United States, with coloured illustrations, and a monograph of the genus* Anculotus *of Say; also a synopsis of the American naiades.* 76 pp., 8 pls.]. *Additions to, and corrections of, the catalogue of species of American naiades, with descriptions of new species and varieties of fresh water shells.* Judah Dobson, Philadelphia. 8 pp., pl. 9.

CONRAD, Timothy A. 1849. Descriptions of new fresh water and marine shells. *Proc. Acad. nat. Sci. Phila.*, 4: 152-156.

CONRAD, Timothy A. 1855. Description of a new species of *Melania. Proc. Acad. nat. Sci. Phila.*, 7: 269.

CONRAD, Timothy A. 1866. Descriptions of American fresh-water shells. *Am. J. Conchol.*, 2(3): 278-279, pl. 15.

COOPER, J.G. 1867. *Geographical catalogue of the Mollusca found west of the Rocky Mountains, between latitudes 33° and 49° north.* Calif. geol. Surv., San Francisco. pp. 1-40.

COOPER, J.G. 1868. The shells of Montana. *Am. Nat.*, 2: 486-487.

COOPER, J.G. 1870. The west-coast fresh-water univalves, no. 1. *Proc. Calif. Acad. Sci.*, ser. 1, 4: 92-100.

COOPER, J.G. 1875. On shells of the west slope of North America. *Proc. Calif. Acad. Sci.*, ser. 1, 6: 14-27.

COOPER, J.G. 1892a. Catalogue of the land and fresh-water Mollusca of lower California. *Zoe*, 3: 12-25.

COOPER, J.G. 1892b. On land and fresh water Mollusca of lower California. No. 2. *Proc. Calif. Acad. Sci.*, ser. 2, 3: 207-217.

COOPER, J.G. 1893. On land and fresh water Mollusca of lower California. No. 3. *Proc. Calif. Acad. Sci.*, ser. 2, 3: 338-344, pls. 13, 14.

COOPER, William. 1859. Report upon the Mollusca collected on the survey. *U.S. War Dept. Pacific R.R. Surv.*, 12(book 2, pt. 3): 369-386.

COTTAM, Clarence. 1942. Supplementary notes on the food of the Limpkin. *Nautilus*, 55(4): 125-128.

COUPER, J. Hamilton. 1849. Catalogue of the Recent shells of Georgia. *In:* White, *Statistics of the State of Georgia, sketch of the geology of Georgia.* Appendix, pp. 37-41.

COUTHOUY, Joseph P. 1838. Descriptions of new species of Mollusca and shells, and remarks on several polypi found in Massachusetts Bay. *Boston J. nat. Hist.*, 2(1): 53-111, pls. 1-4.

CRABB, Edward Drane. 1927. Genetic experiments with pond snails *Lymnaea* and *Physa. Am. Nat.*, 61(672): 54-67.

CRABB, Edward Drane. 1929. Egg laying and birth of young in three species of Viviparidae. *Nautilus*, 42(4): 125-129.

CRAM, Eloise B. 1947. A study of native Planorbidae as possible intermediate hosts of *Schistosoma mansoni* in man. *News Bull. annu. Rep. Am. malacol. Union 1946*, (13): 4-5.

CRANDALL, O.A. 1892. *Physa heterostropha. Nautilus*, 6(2): 20-21.

CRANDALL, O.A. 1893. Post Pliocene shells. *Nautilus*, 6(9): 103.

CRANDALL, O.A. 1901. The America physae. *Nautilus*, 15(3): 25-30; 15(4): 42-45, pl. 2; 15(5): 54-58; 15(6): 69-71.

CRUTCHFIELD, Philip J. 1966. Positive rheotaxis in *Goniobasis proxima. Nautilus*, 79(3): 80-86.

CURRIER, A.O. 1867. Description of new species of fresh-water shells from Michigan. *Am. J. Conchol.*, 3(2): 112-113, pl. 6.

CURRIER, A.O. 1868. List of the shell-bearing Mollusca of Michigan, especially of Kent and adjoining counties. *Misc. Publ. Kent sci. Inst.*, (1): 1-12.

DALL, William Healey. 1870. On the genus *Pompholyx* and its allies, with a revision of the Lymnaeidae of authors. *Ann. Lyc. nat. Hist. N.Y.*, 9: 333-361, 1 pl.

DALL, William Healey. 1884. Report on the Mollusca of the Commander Islands, Bering Sea, collected by Leonhard Stejneger in 1882 and 1883. *Proc. U.S. natl. Mus.*, 7(22): 340-349.

DALL, William Healey. 1885. Notes on some Floridian land and fresh-water shells with a revision of the Auriculacea of the eastern United States. *Proc. U.S. natl. Mus.*, 8(16): 255-256; 8(17): 257-272; 8(18): 273-288; 8(19): 289; pls. 17, 18.

DALL, William Healey. 1889. *Paludina scalaris*, Jay. *Nautilus*, 3(1): 8.

DALL, William Healey. 1895. Diagnoses of new species of mollusks from the west coast of America. *Proc. U.S. natl. Mus.*, 18(1034): 7-20.
DALL, William Healey. 1896. Note on *Neritina showalteri* Lea. *Nautilus*, 10(2): 13-15.
DALL, William Healey. 1898. Description of a new *Ampullaria* from Florida. *Nautilus*, 12(7): 75-76.
DALL, William Healey. 1902. Illustrations and descriptions of new, unfigured, or imperfectly known shells, chiefly American, in the U.S. National Museum. *Proc. U.S. natl. Mus.*, 24(1264): 499-566, pls. 27-40.
DALL, William Healey. 1904a. *Gundlachia* and *Ancylus*. *Nautilus*, 17(9): 97-98.
DALL, William Healey. 1904b. Notes on the genus *Ampullaria*. *J. Conchol.*, 11(2): 50-55.
DALL, William Healey. 1905. Land and fresh water mollusks of Alaska and adjoining regions. *Smithson. Inst. Harriman Alaska Exped., 1899*, Alaska ser., 13: 1-171, pls. 1, 2.
DALL, William Healey. 1907. [*Planorbis magnificus* in aquaria.] *Nautilus*, 22(8): 90.
DALL, William Healey. 1910. A new Floridian *Amnicola*. *Nautilus*, 24(1): 2.
DALL, William Healey. 1911. Notes on *Gundlachia* and *Ancylus*. *Am. Nat.*, 45: 175-189.
DAVIS, Charles C. 1961. A study of the hatching process in aquatic invertebrates. I. The hatching process in *Amnicola limosa* (Gastropoda, Prosobranchia). II. Hatching in *Ranatra fusca* P. Beauvois (Hemiptera, Nepidae). *Trans. Am. microsc. Soc.*, 80(2): 227-234.
DAVIS, George Morgan. 1967. The systematic relationship of *Pomatiopsis lapidaria* and *Oncomelania hupensis formosana* (Prosobranchia: Hydrobiidae). *Malacologia*, 6(1/2): 1-143.
DAVIS, George Morgan. 1974. *Report on the rare and endangered status of a selected number of freshwater Gastropoda from southeastern U.S.A.* Rep. to U.S. Dept. Interior, Fish and Wildlife Serv., Washington, D.C. 51 pp., 25 maps.
DAVIS, George Morgan. 1977. Rare and endangered species: a dilemma. *Frontiers*, 41(4): 12-14.
DAVIS, George Morgan. 1979. The origin and evolution of the gastropod family Pomatiopsidae, with emphasis on the Mekong River Triculinae. *Monogr. Acad. nat. Sci. Phila.*, 20: vii, 120 pp.
DAWLEY, Charlotte. 1965. Checklist of freshwater mollusks of North Carolina. *Sterkiana*, (19): 35-39.
DAWSON, Jean. 1911. The biology of *Physa*. *Behav. Monogr.*, 1(4): 1-120, 5 pp. figs.
DAZO, Bonifacio Capili. 1961. Some studies on the genus *Io*. *Annu. Rep. Am. malacol. Union 1961*, (28): 6.
DAZO, Bonifacio Capili. 1965. The morphology and natural history of *Pleurocera acuta* and *Goniobasis livescens* (Gastropoda: Cerithiacea: Pleuroceridae). *Malacologia*, 3(1): 1-80; 3(3): 475-476 (1966).
DeCAMP, Wm. H. 1881. The shell-bearing Mollusca of Michigan. *Misc. Publ. Kent sci. Inst.*, (5): 1-15, pl. 1.
DE KAY, James E. 1843. *Zoology of New York. Pt. 5. Mollusca.* Carroll and Cook, Albany, New York. 271 pp., pls. 1-40.
DEXTER, Ralph W. 1955. Comparison of the gastropod fauna in the drainage systems of Champaign County, Illinois. *Annu. Rep. Am. malacol. Union 1954*, (21): 1-2.
DEXTER, Ralph W. 1959. Habitat change for *Ferrissia*. *Nautilus*, 72(4): 144-145.
DEXTER, Ralph W. 1961a. Changes in the gastropod populations in the Salt Fork of the Big Vermilion River in Illinois, 1918-1959. *Annu. Rep. Am. malacol. Union 1960*, (27): 24-25.
DEXTER, Ralph W. 1961b. Changes in the gastropod populations in the Salt Fork of the Big Vermilion River in Illinois, 1918-1959. *Sterkiana*, (3): 15-18.
DEXTER, Ralph W. 1961c. George J. Streator's cruise of the *Thomas Say* in 1885. *Sterkiana*, (5): 39-42.
DEXTER, Ralph W. 1963. Some interesting molluscan records of George J. Streator, 1881-1909. *Sterkiana*, (12): 9-14.
DEXTER, Ralph W. & SWART, Louis H. 1965. Changes in the gastropod populations of the West Branch of the Mahoning River in Ohio (1939-1963). *Sterkiana*, (17): 41-43.
DEXTER, Ralph W., SWART, Louis H. & DAVIS, A.F. 1952. Comparison of the mollusks in adjacent Ohio River and Lake Erie drainage systems. *News Bull. annu. Rep. Am. malacol. Union 1951*, (18): 2-3.
DILLON, Robert T. 1979. An electrophoretic and shell morphometric survey of *Goniobasis* from the upper New River and surrounding drainages. *Bull. Am. malacol. Union Inc. 1979*, 46: 67-68.
DOREMUS, Craig M. & HARMAN, Willard N. 1977. The effects of grazing by physid and planorbid freshwater snails on periphyton. *Nautilus*, 91(3): 92-96.
DUCH, Thomas M. 1976. Aspects of the feeding habitats of *Viviparus georgianus*. *Nautilus*, 90(1): 7-10.
DUNDEE, Dee Saunders. 1957. Aspects of the biology of *Pomatiopsis lapidaria* (Say) (Mollusca: Gastropoda: Prosobranchia). *Misc. Publ. Mus. Zool. Univ. Mich.*, (100): 1-37, pls. 1-14.
DUNDEE, Dee Saunders. 1974. Catalog of introduced molluscs of eastern North America (north of Mexico). *Sterkiana*, (55): 1-37.
DUNDEE, Dee Saunders & PAINE, Anna. 1977. Ecology of the snail, *Melanoides tuberculata* (Müller), intermediate host of the human liver fluke (*Opisthorchis sinensis*) in New Orleans, Louisiana. *Nautilus*, 91(1): 17-20.

EMERSON, William K. & JACOBSON, Morris K. 1976. *Guide to shells, land, freshwater, and marine, from Nova Scotia to Florida.* Alfred A. Knopf, New York. vii, 482, xviii (index) pp., 47 pls.
EMERY, D.L. 1943. Sinistral *Pomacea. Nautilus,* 57(2): 66.
EYERDAM, Walter J. 1925. Station of *Ferrissia oregonensis* (Clessin). *Nautilus,* 39(2): 71.
EYERDAM, Walter J. 1937. A concentrated population of *Lymnaea palustris nuttalliana. Nautilus,* 51(2): 72.
EYERDAM, Walter J. 1941. *Lymnaea auricularia* Linnaeus in western Washington and Kamchatka. *Nautilus,* 55(1): 18-19.
FARRER, W.J. 1892. *Planorbis multivolvis. Nautilus,* 6(3): 36.
FERRISS, James H. 1924. On the Rio Grande. *Nautilus,* 38(2): 37-43.
FLUCK, William Henry. 1932. *Valvata simplex* Gould. *Nautilus,* 46(1): 19-22.
FRAUENFELD, Georg Ritter von. 1862. Versuch einer Aufzählung der Arten der Gattung *Bithynia* Lch. und *Nematura* Bns. Nach der kaiserlichen und Cuming's Sammlung. *Verhandl. kais. konig. zool.-bot. Ges. Wien,* 1862, pp. 1145-1170.
FRAUENFELD, Georg Ritter von. 1863a. Die Arten der Gattung *Lithoglyphus* Mhlf., *Paludinella* Pf., *Assiminea* Gray, in der kaiserlichen und Cuming's Sammlung. *Verhandl. kais. konig. zool.-bot. Ges. Wien,* 13: 193-212.
FRAUENFELD, Georg Ritter von. 1863b. Vorläufige Aufzählung der Arten der Gattungen *Hydrobia* Htm. und *Amnicola* Gld. Hldm. in der kaiserlichen und in Cuming's Sammlung. *Verhandl. kais. konig. zool.-bot. Ges. Wien,* 13: 1017-1032.
FREAS, Dorothy L. 1950. *Viviparus chinensis. Nautilus,* 64(1): 35-36.
FRIED, Bernard & GOODCHILD, Chauncey G. 1963. Studies on the reproduction, growth, and survival of a planorbid snail, *Menetus dilatatus buchanensis* (Lea), an experimental host of *Spirorchis* sp. (Trematoda). *Trans. Am. microsc. Soc.,* 82(2): 143-149.
GAINEY, Louis F., Jr. 1976. Locomotion in the Gastropoda: functional morphology of the foot in *Neritina reclivata* and *Thais rustica. Malacologia,* 15(2): 411-431.
GETZ, Lowell L. 1971. On the occurrence of *Pomatiopsis cincinnatiensis* in Wisconsin. *Sterkiana,* (43): 20.
GILL, Theodore. 1863. Systematic arrangement of the mollusks of the family Viviparidae, and others, inhabiting the United States. *Proc. Acad. nat. Sci. Phila.,* 15: 33-40.
GILL, Theodore. 1871. Arrangement of the families of mollusks. *Smithson. misc. Collect.,* 10: i-xvi, 1-49 [ser. no. 227].
GOODRICH, Calvin. 1911. *Lymnaea auricularia* in Ohio. *Nautilus,* 25(1): 11-12.
GOODRICH, Calvin. 1913. Spring collecting in southwest Virginia. *Nautilus,* 27(7): 81-82; 27(8): 91-95.
GOODRICH, Calvin. 1914. Union of the Wabash and Maumee drainage systems. *Nautilus,* 27(11): 131-132.
GOODRICH, Calvin. 1916. A trip to islands in Lake Erie. *Ann. Carnegie Mus.,* 10: 527-531.
GOODRICH, Calvin. 1917. *Pleurocera subulare* Lea. *Nautilus,* 30(11): 122-124.
GOODRICH, Calvin. 1920. Goniobases of Ohio. *Nautilus,* 33(3): 73-74.
GOODRICH, Calvin. 1921a. Three new species of Pleuroceridae. *Occ. Pap. Mus. Zool. Univ. Mich.,* (91): 1-5, pl. 1.
GOODRICH, Calvin. 1921b. River barriers to aquatic animals. *Nautilus,* 35(1): 1-4.
GOODRICH, Calvin. 1921c. Anculosae north of the Alabama drainage. *Nautilus,* 35(1): 9-12.
GOODRICH, Calvin. 1921d. Something about *Angitrema. Nautilus,* 35(2): 58-59.
GOODRICH, Calvin. 1922. The Anculosae of the Alabama River drainage. *Misc. Publ. Mus. Zool. Univ. Mich.,* (7): 1-57, pls. 1-3.
GOODRICH, Calvin. 1923. Variations in *Goniobasis edgariana* Lea. *Nautilus,* 36(4): 115-119.
GOODRICH, Calvin. 1924a. The genus *Gyrotoma. Misc. Publ. Mus. Zool. Univ. Mich.,* (12): 1-29, pls. 1, 2.
GOODRICH, Calvin. 1924b. Some old pleurocerids and a new one. *Nautilus,* 38(2): 43-48.
GOODRICH, Calvin. 1927. Some misplaced pleurocerids. *Nautilus,* 41(2): 57-62.
GOODRICH, Calvin. 1928a. *Strephobasis*: a section of *Pleurocera. Occ. Pap. Mus. Zool. Univ. Mich.,* (192): 1-15, pls. 1, 2.
GOODRICH, Calvin. 1928b. The group of *Goniobasis catenaria. Nautilus,* 42(1): 28-32.
GOODRICH, Calvin. 1928c. *Goniobasis undulata. Nautilus,* 42(2): 67.
GOODRICH, Calvin. 1929a. The nomenclature of ecological varieties. *Nautilus,* 42(4): 114-118.
GOODRICH, Calvin. 1929b. The pleurocerid fauna of the Falls of the Ohio. *Nautilus,* 43(1): 1-17.
GOODRICH, Calvin. 1930a. Goniobases of the vicinity of Muscle Shoals. *Occ. Pap. Mus. Zool. Univ. Mich.,* (209): 1-25.
GOODRICH, Calvin. 1930b. *Pleurocera (Strephobasis) curtum* (Haldeman). *Nautilus,* 43(4): 141.
GOODRICH, Calvin. 1931a. The pleurocerid genus *Eurycaelon. Occ. Pap. Mus. Zool. Univ. Mich.,* (223): 1-9, pl. 1.
GOODRICH, Calvin. 1931b. Some conchological beginnings. *Nautilus,* 45(2): 41-47.
GOODRICH, Calvin. 1931c. Mollusks of Keweenaw County, Michigan. *Occ. Pap. Mus. Zool. Univ. Mich.,* (233): 1-

9, pl. 1.
GOODRICH, Calvin. 1932a. *The Mollusca of Michigan.* Mich. Handbook Ser., Mus. Zool. Univ. Mich., (5): 1-120, pls. i-vii.
GOODRICH, Calvin. 1932b. Collecting in Tennessee. *Nautilus,* 45(4): 136-137.
GOODRICH, Calvin. 1932c. Cress a possible introducer of species. *Nautilus,* 46(1): 35-36.
GOODRICH, Calvin. 1932d. The true position of Bruguiere's *Bulimus carinatus. Nautilus,* 46(2): 38-40.
GOODRICH, Calvin. 1933. Mollusks of Moose Factory. *Nautilus,* 47(1): 7-9.
GOODRICH, Calvin. 1934a. Studies of the gastropod family Pleuroceridae – I. *Occ. Pap. Mus. Zool. Univ. Mich.,* (286): 1-17, pl. 1.
GOODRICH, Calvin. 1934b. Studies of the gastropod family Pleuroceridae – II. *Occ. Pap. Mus. Zool. Univ. Mich.,* (295): 1-6, fig. 1.
GOODRICH, Calvin. 1934c. Studies of the gastropod family Pleuroceridae – III. *Occ. Pap. Mus. Zool. Univ. Mich.,* (300): 1-11.
GOODRICH, Calvin. 1935a. Studies of the gastropod family Pleuroceridae – IV. *Occ. Pap. Mus. Zool. Univ. Mich.,* (311): 1-11.
GOODRICH, Calvin. 1935b. Studies of the gastropod family Pleuroceridae – V. *Occ. Pap. Mus. Zool. Univ. Mich.,* (318): 1-12.
GOODRICH, Calvin. 1935c. A species of *Goniobasis* new to the Great Basin. *Nautilus,* 49(2): 66.
GOODRICH, Calvin. 1935d. A mollusk new to the Great Basin. *Rep. Am. malacol. Union 1935,* (4): 3.
GOODRICH, Calvin. 1936. *Goniobasis* of the Coosa River, Alabama. *Misc. Publ. Mus. Zool. Univ. Mich.,* (31): 1-60, pl. 1.
GOODRICH, Calvin. 1937a. *Goniobasis columbiensis* Whiteaves. *Nautilus,* 50(3): 82-84.
GOODRICH, Calvin. 1937b. Studies of the gastropod family Pleuroceridae – VI. *Occ. Pap. Mus. Zool. Univ. Mich.,* (347): 1-12.
GOODRICH, Calvin. 1938. Studies of the gastropod family Pleuroceridae – VII. *Occ. Pap. Mus. Zool. Univ. Mich.,* (376): 1-12.
GOODRICH, Calvin. 1939a. Aspects of depauperization. *Nautilus,* 52(4): 124-128.
GOODRICH, Calvin. 1939b. Certain mollusks of the Ogeechee River, Georgia. *Nautilus,* 52(4): 129-131.
GOODRICH, Calvin. 1939c. New opercula for old. *Nautilus,* 52(4): 140.
GOODRICH, Calvin. 1939d. Pleuroceridae of the St. Lawrence River basin. *Occ. Pap. Mus. Zool. Univ. Mich.,* (404): 1-4.
GOODRICH, Calvin. 1939e. Pleuroceridae of the Mississippi River basin exclusive of the Ohio River system. *Occ. Pap. Mus. Zool. Univ. Mich.,* (406): 1-4.
GOODRICH, Calvin. 1939f. Who or what? [*Publications of the O. G. B., III, Mollusca of Cincinnati,* 1876]. *Nautilus,* 53(2): 66-67.
GOODRICH, Calvin. 1939g. Part I. The gastropods and the Sphaeriidae. *In:* Goodrich, Calvin & van der Schalie, Henry, Aquatic mollusks of the Upper Peninsula of Michigan. *Misc. Publ. Mus. Zool. Univ. Mich.,* (43): 1-45, 2 maps [pp. 7-34].
GOODRICH, Calvin. 1940a. Sequence in the distribution of aquatic species. *Nautilus,* 53(3): 73-77.
GOODRICH, Calvin. 1940b. Another immigrant. *Nautilus,* 53(3): 105-106.
GOODRICH, Calvin. 1940c. Regarding O.G.B. [*Publications of the O. G. B., III, Mollusca of Cincinnati,* 1876]. *Nautilus,* 53(3): 106.
GOODRICH, Calvin. 1940d. The Pleuroceridae of the Ohio River drainage system. *Occ. Pap. Mus. Zool. Univ. Mich.,* (417): 1-21.
GOODRICH, Calvin. 1940e. Civilization and aquatic mollusks. *Nautilus,* 54(1): 6-10.
GOODRICH, Calvin. 1941a. Two new species of *Goniobasis. Occ. Pap. Mus. Zool. Univ. Mich.,* (426): 1-4, figs. 1, 2.
GOODRICH, Calvin. 1941b. Pleuroceridae of the small streams of the Alabama River system. *Occ. Pap. Mus. Zool. Univ. Mich.,* (427): 1-10.
GOODRICH, Calvin. 1941c. Distribution of the gastropods of the Cahaba River, Alabama. *Occ. Pap. Mus. Zool. Univ. Mich.,* (428): 1-30.
GOODRICH, Calvin. 1941d. *Lymnaea contracta* Currier. *Nautilus,* 54(4): 121-122.
GOODRICH, Calvin. 1941e. A sinistral albino. *Nautilus,* 54(4): 141.
GOODRICH, Calvin. 1941f. Studies of the gastropod family Pleuroceridae – VIII. *Occ. Pap. Mus. Zool. Univ. Mich.,* (447): 1-13.
GOODRICH, Calvin. 1942a. The American species of *Viviparus. Nautilus,* 55(3): 82-92.
GOODRICH, Calvin. 1942b. The Pleuroceridae of the Atlantic Coastal Plain. *Occ. Pap. Mus. Zool. Univ. Mich.,*

(456): 1-6.
GOODRICH, Calvin. 1942c. Observations upon a Florida form of *Viviparus*. *Nautilus*, 55(4): 115-118.
GOODRICH, Calvin. 1942d. Certain remarks about labels. *Nautilus*, 55(4): 119-120.
GOODRICH, Calvin. 1942e. The Pleuroceridae of the Pacific Coastal drainage, including the Western Interior basin. *Occ. Pap. Mus. Zool. Univ. Mich.*, (469): 1-4.
GOODRICH, Calvin. 1943. The Walker-Beecher paper of 1876 on the Mollusca of the Ann Arbor area. *Occ. Pap. Mus. Zool. Univ. Mich.*, (475): 1-26, 1 map.
GOODRICH, Calvin. 1944a. Pleuroceridae of the Great Basin. *Occ. Pap. Mus. Zool. Univ. Mich.*, (485): 1-11, figs. 1, 2.
GOODRICH, Calvin. 1944b. Certain operculates of the Coosa River. *Nautilus*, 58(1): 1-10.
GOODRICH, Calvin. 1944c. Pulmonates of the Coosa River. *Nautilus*, 58(1): 11-15.
GOODRICH, Calvin. 1944d. Pleuroceridae of the Coosa River basin. *Nautilus*, 58(2): 40-48.
GOODRICH, Calvin. 1945. *Goniobasis livescens* of Michigan. *Misc. Publ. Mus. Zool. Univ. Mich.*, (64): 1-36, pl. 1, fig. 1, 1 map.
GOODRICH, Calvin. 1950. *Goniobasis proxima* (Say). *Nautilus*, 63(3): 78-80.
GOODRICH, Calvin & VAN DER SCHALIE, Henry. 1944. A revision of the Mollusca of Indiana. *Am. midl. Nat.*, 32(2): 257-326.
GOULD, Augustus A. 1833. *Lamarck's genera of shells, with a catalogue of species: translated from the French.* Boston. xiii, 110 pp.
GOULD, Augustus A. 1837. List of shells found in Maine. *In:* Jackson, C.T., *First report on the geology of the state of Maine.* Augusta, Maine. p. 119.
GOULD, Augustus A. 1838. Dr. Gould's report [on molluscous and the other lower animals]. *In:* Reports of the commissioners on the zoological survey of the state. Massachusetts House Document No. 72. pp. 105-107.
GOULD, Augustus A. 1840. [Descriptions of thirteen new species of shells.] *Silliman's Am. J. Sci. Arts*, ser. 1, 38: 196-197.
GOULD, Augustus A. 1841a. *A report on the Invertebrata of Massachusetts, comprising the Mollusca, Annelida, and Radiata.* Folsom, Wells and Thurston, Cambridge, Massachusetts. pp. i-xiii, 1-373, 15 pls.
GOULD, Augustus A. 1841b. Results of an examination of the shells of Massachusetts, and their geographical distribution. *Boston J. nat. Hist.*, 3(4): 483-494.
GOULD [Augustus A.] 1843a. [Reversed *Paludina ponderosa*.] *Proc. Boston Soc. nat. Hist.*, 1: 32.
GOULD, Augustus A. 1843b. Notice of 'Monograph of the fresh-water univalve Mollusca of the U.S., by S.S. Haldeman. No. 6. Genus *Physa*'. [Review.] *Proc. Boston Soc. nat. Hist.*, 1: 132-133.
GOULD, Augustus A. 1847a. [Note to " 'Description of two new species of shells', by William Case."] *Silliman's Am. J. Sci. Arts*, ser. 2, 3: 101.
GOULD [Augustus A.] 1847b. [Descriptions of Limniadae, from the collection of the Exploring Expedition.] *Proc. Boston Soc. nat. Hist.*, 2: 210-212.
GOULD [Augustus A.] 1847c. [Descriptions of *Physa*, from the collection of the Exploring Expedition.] *Proc. Boston Soc. nat. Hist.*, 2: 214-215.
GOULD [Augustus A.] 1847d. [Descriptions of *Melania*, from the collection of the Exploring Expedition.] *Proc. Boston Soc. nat. Hist.*, 2: 222-225.
GOULD [Augustus A.] 1847e. [Specimens of *Planorbis multivolvis* Case, brought by Mr. C.J. Foster from the Lake Superior region.] *Proc. Boston Soc. nat. Hist.*, 2: 259.
GOULD [Augustus A.] 1847f. [Description of a new species of *Physa*, among the shells collected by Dr. C.T. Jackson on the shores of Lake Superior; together with two other new species of North American shells.] *Proc. Boston Soc. nat. Hist.*, 2: 262-263.
GOULD, Augustus A. 1848. [Descriptions of new shells.] *Proc. Boston Soc. nat. Hist.*, 3: 64-65.
GOULD, Augustus A. 1850a. [Remarks on specimens of *Lymnaea* from Lake Superior.] *Proc. Boston Soc. nat. Hist.*, 3: 181.
GOULD, Augustus A. 1850b. Catalogue of shells with descriptions of new species. *In:* Agassiz, Louis, *Lake Superior: physical character, vegetation, and animals, compared with those of other and similar regions.* Gould, Kendall and Lincoln, Boston. 428 pp. [pp. 243-245].
GOULD, Augustus A. 1850c. Descriptions of sixteen new species of *Melania*. *Proc. Boston Soc. nat. Hist.*, 3: 359-363.
GOULD, Augustus A. 1852-61. Mollusca and shells. Vol. 12. *In: United States Exploring Expedition, during the years 1838, 1839, 1840, 1841, 1842, under the command of Charles Wilkes, U.S.N.* [1852.] Gould and Lincoln, Boston. xv, 510 pp. [Addenda and corrigenda, 1856, pp. 499-509; Atlas, 1856 [1861], 16 pp., 52 pls; Philadelphia.]

GOULD, Augustus A. 1853. [Remarks on several attempts to bring living fresh water shells from Cuba and England to this country.] *Proc. Boston Soc. nat. Hist.*, 4: 287-288.

GOULD [Augustus A.] 1855a. New species of land and fresh-water shells from western (N.) America. *Proc. Boston Soc. nat. Hist.*, 5: 127-130.

GOULD, Augustus A. 1855b. Catalogue of shells collected in California by W.P. Blake, with descriptions of the new species. Appendix, art. 2, pp. 22-28, *of* Preliminary geological report, by W.P. Blake. *In:* Report of a reconnaissance and survey in California in connection with explorations for a practicable railway route from the Mississippi River to the Pacific Ocean in 1853 by Lt. R.S. Williamson. U.S. House of Representatives, Document 129, 33rd Congress, first session, Washington, D.C.

GOULD, Augustus A. 1857. Catalogue of the Recent shells, with descriptions of the new species. *U.S. War Dept. Pacific R.R. Surv.*, 5(2): 330-336, pl. 11.

GOULD, Augustus A. 1862. Otia conchologica: *Descriptions of shells and mollusks, from 1839 to 1862.* Gould and Lincoln, Boston. 256 pp.

GOULD, Augustus A. 1865a. [Remarks on erosion in mollusk shells.] *Proc. Boston Soc. nat. Hist.*, 9: 327.

GOULD, Augustus A. 1865b. Remarks on Mollusca and shells: 1. On the zoological regions; 2. Specific identity of shells; 3. Local aspect of species and characteristic forms of regions; 4. Analogous species in coordinate regions. *Edinb. new Philos. J.*, 56: 74-80.

GOULD, Augustus A. 1865c. [Remarks on the diversity among conchologists in the application of terms in the description of shells.] *Proc. Boston Soc. nat. Hist.*, 10: 77-78.

GOULD, Augustus A. 1866. On the correct application of certain terms used in the description of shells. *Am. J. Conchol.*, 2(2): 180-181.

GOULD, Augustus A. 1870. *Report on the Invertebrata of Massachusetts*, published agreeably to an order of the Legislature. Second edition, comprising the Mollusca. Edited by W.G. Binney. Wright and Potter, State Printers, Boston. v, 524 pp., pls. 16-27.

GRAY, J.E. 1847. A list of the genera of Recent Mollusca, their synonyma and types. *Proc. zool. Soc. Lond.*, 1847, pt. 15, pp. 129-219.

GREGG, Wendell O. 1923. Introduced species of *Lymnaea* in southern California. *Nautilus*, 37(1): 34.

GREGG, Wendell O. 1932. *Lymnaea (Fossaria) perplexa* in southern California. *Nautilus*, 45(4): 131-132.

GREGG, Wendell O. 1934. *Planorbis opercularis planulatus* in Los Angeles, California. *Nautilus*, 48(1): 33.

GREGG, Wendell O. 1941. *Fluminicola avernalis* and *Fluminicola avernalis carinifera*. *Nautilus*, 54(4): 117-118.

GREGG, Wendell O. & TAYLOR, Dwight W. 1965. *Fontelicella* (Prosobranchia: Hydrobiidae), a new genus of West American freshwater snails. *Malacologia*, 3(1): 103-110.

GUGLER, Carl W. 1969. Notes on *Valvata tricarinata* from central Nebraska. *Annu. Rep. Am. malacol. Union 1969*, (36): 61-62.

HAAS, Fritz. 1939. Malacological notes. *Zool. Ser. Field Mus. nat. Hist.*, 24(8): 93-103.

HALDEMAN, S. Stehman. 1841a. Observations on the melanians of Lamarck. *Am. J. Sci. Arts*, 41(1): 21-23.

HALDEMAN, S. Stehman. 1841b. Descriptions of two new freshwater shells. *Proc. Acad. nat. Sci. Phila.*, 1: 78-79.

HALDEMAN, S. Stehman. 1841c. Correction [to the article on melanians, vol. xli, p. 21]. *Am. J. Sci. Arts*, 42: 216.

HALDEMAN, S. Stehman. 1842. Description of five new species of freshwater shells. *J. Acad. nat. Sci. Phila.*, 8: 200-202.

HALDEMAN, S. Stehman. 1842-45 [1840-71]. *A monograph of the freshwater univalve Mollusca of the United States* [or *A monograph of the limniades or freshwater univalve shells of North America*]. J. Dobson, Philadelphia. 231 pp., 40 pls. [No. 1, 1840, *Paludina*, pp. 1-16, pls. 1-5; Suppl. to No. 1, 1840, pp. 1-3; No. 2, 1841, *Paludina*, pp. 17-32, pls. 6-10; No. 3, 1841, *Limnea*, pp. 1-16, pls. 1-5; No. 4, 1842, *Limnea*, pp. 17-32, pls. 6-10; No. 5, 1842, *Limnea*, pp. 33-55, pls. 11-15; No. 6, 1842 [1843], *Physa*, pp. 1-40, pls. 1-5; No. 7, 1844, *Planorbis*, pp. 1-32, pls. 1-4, *Ancylus*, pp. 1-14, pl. 1, index to Physadae, 2 pp.; No. 8, 1845, *Amnicola*, pp. 1-24, pl. 1, *Ampullaria*, pp. 1-11, pls. 1, 2, *Valvata*, pp. 1-11, pl. 1; No. 9, 1871, *Paludina*, pp. 33-36, pl. 11, index to Turbidae, corrections, contents, pp. 41-43.]

HALDEMAN, S. Stehman. 1844a. Enumeration of the Recent freshwater Mollusca which are common to North America and Europe; with observations on species and their distribution. *Boston J. nat. Hist.*, 4(4): 468-484.

HALDEMAN, S. Stehman. 1844b. Mollusca. *In: Rupp's History of Lancaster County.* Lancaster, Pennsylvania. pp. 479-482.

HALDEMAN, S. Stehman. 1847-48. G. *Leptoxis*. Leptoxe. Rafinesque. *In:* Chenu, J.C., *Illustrations conchyliologiques*. A. Franck, Paris. pp. 1-6, pls. 1-5.

HALDEMAN, S. Stehman. 1857. Mollusca. *In:* Baird, Spencer F. (Ed. and Transl.), J.G. Heck's *The iconographic encyclopaedia of science, literature, and art, systematically arranged.* Vol. 2. New York. pp. 83-90.

HALDEMAN, S. Stehman. 1863. On Strepomatidae as a name for a family of fluviatile Mollusca, usually confounded with *Melania*. *Proc. Acad. nat. Sci. Phila.*, 15: 273-274.
HALDEMAN, S. Stehman. 1865a. Description of two new species of *Goniobasis*. *Am. J. Conchol.*, 1(1): 37, pl. 1, figs. 4-7.
HALDEMAN, S. Stehman. 1865b. On the priority of *Leptoxis* as a generic name. *Am. J. Conchol.*, 1(4): 298.
HALDEMAN, S. Stehman. 1867. Description of a new species of *Limnea*. *Am. J. Conchol.*, 3(2): 194.
HAMILTON, Suzanne. 1979. *Bithynia tentaculata* – a replacement for North American pleurocerid snails? *Bull. Am. malacol. Union Inc. 1979*, 46: 68.
HAND, Edwin E. 1914. Sometimes locality adds interest to a shell. *Nautilus*, 27(12): 144.
HAND, Edwin E. 1928. Sinistral *Campeloma* (correspondence). *Nautilus*, 41(3): 106-107.
HANHAM, A.W. 1897. *Planorbis nautileus* L. *Nautilus*, 10(11): 130-131.
HANLEY, Sylvanus. 1855. Ipsa Linnaei conchylia. *The shells of Linnaeus, determined from his manuscript and collection. Also, an exact reprint of the* Vermes testacea *of the* 'Systema naturae' *and* 'Mantissa'. Williams & Norgate, London. 556 pp., 5 pls.
HANNA, G. Dallas. 1923. Note on *Lymnaea hemphilliana* (Baker). *Nautilus*, 37(1): 23-25.
HANNA, G. Dallas. 1925. *Zalophancylus*, a fish vertebra, not a mollusk. *Nautilus*, 39(1): 18-19.
HANNA, G. Dallas. 1930. *Pyrgulopsis nevadensis* (Stearns) in Oregon. *Nautilus*, 43(3): 103-104.
HANNA, G. Dallas & CLARK, H. Walton. 1925a. *Lymnaea auricularia* (Linn.) in California. *Nautilus*, 38(4): 125-127.
HANNA, G. Dallas & CLARK, H. Walton. 1925b. Additional records for *Lymnaea auricularia*. *Nautilus*, 39(2): 71.
HANNA, G. Dallas & HENDERSON, Junius. 1934. A review of the genus *Carinifex* Binney. *Rep. Am. malacol. Union 1934*, (3): 6.
HANNIBAL, Harold. 1908a. Formation of epiphragm by *Lymnaea palustris* (Müller). *Nautilus*, 22(3): 33.
HANNIBAL, Harold. 1908b. Exotic *Vivipara* in California. *Nautilus*, 22(3): 33-34.
HANNIBAL, Harold. 1910. Valvatidae of the western North America. *Nautilus*, 23(8): 104-107.
HANNIBAL, Harold. 1911a [1910]. Shells of lakes and streams. *In:* Keep, Josiah, *West Coast shells*. Whitaker and Ray-Wiggin, San Francisco. 346 pp. [pp. 299-318, pls. 1-3].
HANNIBAL, Harold. 1911b. Further notes on Asiatic viviparas in California. *Nautilus*, 25(3): 31-32.
HANNIBAL, Harold. 1912a. The aquatic molluscs of Southern California and adjacent regions, a transition fauna. *Bull. S. Calif. Acad. Sci.*, 10(2): 18-46, 1 pl.
HANNIBAL, Harold. 1912b. A synopsis of the Recent and Tertiary freshwater Mollusca of the Californian province, based upon an ontogenetic classification. *Proc. malacol. Soc. Lond.*, 10(2 & 3): 112-211, pls. 5-8.
HANNIBAL, Harold. 1914. Note on the classification of the Ancylidae. *Nautilus*, 28(2): 23-24.
HANNIBAL, Harold. 1918. Jura-Cretaceous stonewort and limneas, supposedly from Arkansas. *Science*, 48(1249): 578.
HANNING, Gary Wayne. 1977. Oviposition of *Pomacea paludosa* (Say) in Lake Okeechobee, Florida. *Bull. Am. malacol. Union Inc. 1977*, 44: 90-91.
HARMAN, Willard N. 1968a. Replacement of pleurocerids by *Bithynia* in polluted waters of central New York. *Nautilus*, 81(3): 77-83.
HARMAN, Willard N. 1968b. *Valvata piscinalis* in Cayuga Lake, N.Y. *Nautilus*, 81(4): 143-144.
HARMAN, Willard N. 1968c. Interspecific competition between *Bithynia* and pleurocerids. *Nautilus*, 82(2): 72-73.
HARMAN, Willard N. 1970. A local case of intra-population variation in *Lymnaea emarginata* (Say). *Annu. Rep. Am. malacol. Union Inc. 1970*, (37): 45-47.
HARMAN, Willard N. 1973. The Mollusca of Canadarago Lake and a new record for *Lasmigona compressa* (Lea). *Nautilus*, 87(4): 114.
HARMAN, Willard N. 1974. The effects of reservoir construction and canalization on the mollusks of the upper Delaware watershed. *Bull. Am. malacol. Union Inc. 1973*, (40): 12-14.
HARMAN, Willard N. & BERG, Clifford O. 1971. The freshwater snails of central New York, with illustrated keys to the genera and species. *Search, Agric., Entomol. (Ithaca) 2*, 1(4): 1-68.
HARPER, Francis. 1936. The distribution of the Limpkin and its staple food *Pomacea*. *Nautilus*, 50(2): 37-40.
HARRY, Harold W. 1962. A critical catalogue of the nominal genera and species of neotropical Planorbidae. *Malacologia*, 1(1): 33-53.
HARRY, Harold W. & HUBENDICK, Bengt. 1964. The freshwater pulmonate Mollusca of Puerto Rico. *Medd. Göteborgs Mus. Zool. Avd.*, (136): 1-77, 16 pls.
HEARD, William H. 1963. Reproductive features of *Valvata*. *Nautilus*, 77(2): 64-68.
HEARD, William H. 1968. Mollusca. *In:* Parrish, Fred K., *Keys to water quality indicative organisms (southeastern United States)*. Fed. water Pollution Control Admin., U.S. Dept. Interior. 208 pp. [pp. G1-G26].

HEARD, William H. 1970. Eastern freshwater mollusks (II). The south Atlantic and Gulf drainages. [Am. malacol. Union Symp. Rare and Endangered Mollusks.] *Malacologia*, 10(1): 23-31.
HEMPHILL, Henry. 1890. New forms of western limniades. *Nautilus*, 4(3): 25-27.
HENDERSON, Junius. 1910. [*Lymnaea bulimoides techella* Hald. in San Bernardino County.] *Nautilus*, 23(11): 144.
HENDERSON, Junius. 1912. *Lymnaea (Radix) auricularia* Linn. in Colorado. *Nautilus*, 26(7): 84.
HENDERSON, Junius. 1918. The nomenclature and systematic positions of some North American fossil and Recent mollusks. *Nautilus*, 32(2): 60-64.
HENDERSON, Junius. 1924. Mollusca of Colorado, Utah, Montana, Idaho and Wyoming. *Univ. Colo. Stud.*, 13(2): 65-223.
HENDERSON, Junius. 1925. *Physa cupraeonitens* and *Physa sayi warreniana* in hot springs. *Nautilus*, 38(3): 105-106.
HENDERSON, Junius. 1928a. Mollusca in their relation to the geological history of Soap Lake, Washington. *Nautilus*, 41(4): 118-120.
HENDERSON, Junius. 1928b. [Collecting in Idaho, Oregon and Washington.] *Nautilus*, 42(2): 64-65.
HENDERSON, Junius. 1929a. Some new forms of non-marine Mollusca from Oregon and Washington. *Nautilus*, 42(3): 80-82.
HENDERSON, Junius. 1929b. Some fossil fresh-water Mollusca from Washington and Oregon. *Nautilus*, 42(4): 119-123.
HENDERSON, Junius. 1929c. Non-marine Mollusca of Oregon and Washington. *Univ. Colo. Stud.*, 17(2): 47-190.
HENDERSON, Junius. 1930. *Ancylus coloradensis*, new name for *A. hendersoni* Walker, 1925, not 1908. *Nautilus*, 44(1): 31.
HENDERSON, Junius. 1931a. A new lymnaeid from Idaho. *Nautilus*, 44(3): 75-77, pl. 6, fig. 8.
HENDERSON, Junius. 1931b. Variation in *Carinifex newberryi* (Lea) and *Lymnaea utahensis* (Call). *Nautilus*, 44(3): 77-79, pl. 6.
HENDERSON, Junius. 1931c. *Lymnaea (Radix) auricularia* L. in Idaho. *Nautilus*, 44(4): 143.
HENDERSON, Junius. 1932a. *Carinifex jacksonensis*, new species, from Wyoming. *Nautilus*, 45(4): 133-134, pl. 11, fig. 6.
HENDERSON, Junius. 1932b. Prolonged aestivation of lymnaeids. *Nautilus*, 45(4): 140.
HENDERSON, Junius. 1932c. Comments on western Lymnaeidae. *Nautilus*, 45(4): 140-141.
HENDERSON, Junius. 1932d. The range of *Polygyra* and of *Goniobasis* in California. *Nautilus*, 46(1): 4-6.
HENDERSON, Junius. 1933. *Gyraulus vermicularis albolineatus*, new subspecies. *Nautilus*, 47(2): 78-79, pl. 6, fig. 3.
HENDERSON, Junius. 1934a. Some western fresh water mollusks. *Nautilus*, 47(3): 86-91, pl. 9.
HENDERSON, Junius. 1934b. Notes on western Lymnaeidae. *Nautilus*, 47(4): 122-124, pl. 14.
HENDERSON, Junius. 1935a. West American species of *Goniobasis*, with descriptions of new forms. *Nautilus*, 48(3): 94-99, pl. 4; 48(4): 130-134.
HENDERSON, Junius. 1935b. *Margaritifera* and *Fluminicola* in Wyoming. *Nautilus*, 48(3): 107.
HENDERSON, Junius. 1935c. Fossil non-marine Mollusca of North America. *Geol. Soc. Am., Spec. Pap.*, (3): i-vii, 1-313.
HENDERSON, Junius. 1935d. *Lymnaea auricularia* L. and *Ferrissia caurina* (Cooper) in Colorado. *Nautilus*, 49(2): 68-69.
HENDERSON, Junius. 1936a. Mollusca of Colorado, Utah, Montana, Idaho, and Wyoming – supplement. *Univ. Colo. Stud.*, 23(2): 81-145, pls. 1, 2.
HENDERSON, Junius. 1936b. *Physa ampullacea* Gould, *Physa nuttallii* Lea and *Physa columbiana* Hemphill. *Nautilus*, 49(3): 85-87.
HENDERSON, Junius. 1936c. The non-marine Mollusca of Oregon and Washington – supplement. *Univ. Colo. Stud.*, 23(4): 251-280, pls. 1, 2.
HENDERSON, Junius. 1936d. *Helisoma ammon* (Gould). *Nautilus*, 50(2): 41-42.
HERRINGTON, H.B. 1947. *Acella haldemani* in Ontario, Canada. *Nautilus*, 61(1): 20-25.
HERRMANN, Sheila A. & HARMAN, Willard N. 1975. Population studies on *Helisoma anceps* (Menke) (Gastropoda: Planorbidae). *Nautilus*, 89(1): 5-11.
HERRMANNSEN, A.N. 1846. *Indicis generum malacozoorum primordia.* Vol. 1. Theodori Fischeri, Cassellis. 637 pp.
HESSE, P. 1923. Kritische Fragmente. *Arch. Molluskenkd.*, 55: 193-198.
HIBBARD, Claude W. & TAYLOR, Dwight W. 1960. Two late Pleistocene faunas from southwestern Kansas. *Contrib. Mus. Paleontol. Univ. Mich.*, 16(1): 1-223, pls. 1-16.

HINDS, Richard Brinsley. 1844. Descriptions of new species of *Melania* collected during the voyage of H.M.S. Sulphur. *Ann. Mag. nat. Hist.*, 14(88): 8-11.
HINKLEY, Anson A. 1888. Notes on the Strepomatidae of Illinois. *Conchol. Exc.*, 2(7): 93-94.
HINKLEY, Anson A. 1894. Variations of *Pleurocera alveare* Conrad. *Nautilus*, 8(7): 82.
HINKLEY, Anson A. 1901. Notes on *Meseschiza grosvenorii* Lea. *Nautilus*, 15(1): 5-6.
HINKLEY, Anson A. 1903. A proposed study of *Goniobasis*. *Nautilus*, 17(3): 32-34.
HINKLEY, Anson A. 1906. Some shells of Mississippi and Alabama. *Nautilus*, 20(4): 40-44.
HINKLEY, Anson A. 1908. *Meseschiza grosvenorii*, Lea. *Nautilus*, 22(6): 56.
HINKLEY, Anson A. 1912a. Notes on *Physa gyrina*. *Nautilus*, 25(9): 108.
HINKLEY, Anson A. 1912b. *Anculosa minor*, n. sp. *Nautilus*, 26(4): 47-48, pl. 1, figs. 7, 8.
HINKLEY, Anson A. 1920. A new species of *Goniobasis*. *Occ. Pap. Mus. Zool. Univ. Mich.*, (78): 1-3.
HOFF, C. Clayton. 1940. Anatomy of the ancylid snail, *Ferrissia tarda* (Say). *Trans. Am. microsc. Soc.*, 59(2): 224-242.
HORST, Thomas J. & COSTA, Robert R. 1975. Seasonal migration and density patterns of the fresh water snail, *Amnicola limosa*. *Nautilus*, 89(2): 56-59.
HUBENDICK, Bengt. 1951. Recent Lymnaeidae. Their variation, morphology, taxonomy, nomenclature, and distribution. *Kungl. svenska Vet.-akad. Handl.*, ser. 4, 3(1): 1-223, pls. 1-5.
HUBENDICK, Bengt. 1955. Phylogeny in Planorbidae. *Trans. zool. Soc. Lond.*, 28(6): 453-542.
HUBENDICK, Bengt. 1962. Studies on *Acroloxus* (Moll. Basomm.). *Medd. Göteborgs Mus. Zool. Avd.*, (133): 1-68.
HUBENDICK, Bengt. 1964. Studies on Ancylidae. *Medd. Göteborgs Mus. Zool. Avd.*, (137): 1-72.
HUBENDICK, Bengt. 1969. A note on *Acroloxus coloradensis* (Henderson). *J. Conchyliol.*, 107(3): 109-118.
HUBRICHT, Leslie. 1940a. The Ozark amnicolas. *Nautilus*, 53(4): 118-122, pl. 14.
HUBRICHT, Leslie. 1940b. A subterranean snail from an artesian well. *Nautilus*, 54(1): 34-35.
HUBRICHT, Leslie. 1941. The cave Mollusca of the Ozark region. *Nautilus*, 54(4): 111-112.
HUBRICHT, Leslie. 1942. A new locality for *Amnicola proserpina* Hubricht. *Nautilus*, 55(3): 105.
HUBRICHT, Leslie. 1943. Notes on the sex ratios in *Campeloma*. *Nautilus*, 56(4): 138-139.
HUBRICHT, Leslie. 1954. The snails from two Indian shellmounds near Clarksville, Virginia. *Nautilus*, 67(3): 90-92.
HUBRICHT, Leslie. 1957. New species of *Fontigens* from the Shenandoah National Park. *Nautilus*, 71(1): 9-10.
HUBRICHT, Leslie. 1960. *Pomatiopsis lapidaria* on the southern Atlantic coastal plain, with remarks on the status of *P. praelonga* and *P. hinkleyi*. *Nautilus*, 74(1): 33-34.
HUBRICHT, Leslie. 1963a. *Lyogyrus granum* (Say) in Mississippi. *Nautilus*, 76(3): 112.
HUBRICHT, Leslie. 1963b. New species of Hydrobiidae. *Nautilus*, 76(4): 138-140, pl. 8, figs. A-F.
HUBRICHT, Leslie. 1963c. *Notogillia wetherbyi* (Dall) in Alabama. *Nautilus*, 76(4): 152.
HUBRICHT, Leslie. 1971. New Hydrobiidae from Ozark caves. *Nautilus*, 84(3): 93-96.
HUBRICHT, Leslie. 1976. The genus *Fontigens* from Appalachian caves (Hydrobiidae: Mesogastropoda). *Nautilus*, 90(2): 86-88.
HUBRICHT, Leslie. 1979. A new species of *Amnicola* from an Arkansas cave (Hydrobiidae). *Nautilus*, 93(4): 142.
HUNT, Burton P. 1958. Introduction of *Marisa* into Florida. *Nautilus*, 72(2): 53-55.
HUTCHISON, Lynn. 1947. Analysis of the activity of the fresh-water snail, *Viviparus malleatus* (Reeve). *Ecology*, 28(4): 335-345.
INABA, Akihiko. 1969. Cytotaxonomic studies of lymnaeid snails. *Malacologia*, 7(2/3): 143-168.
INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE. 1956. Opinion 432. Rejection, as an unpublished proof, of the paper by Binney (W.G.) dated "9 December 1863" and entitled *Synopsis of the species of air-breathing mollusks of North America* (confirmation of Ruling given in *Opinion* 87) and validation under the Plenary Powers of the generic name *Carinifex* Binney, 1865 (Class Gastropoda). *Opinions and declarations rendered by the International Commission on Zoological Nomenclature*, 14(16): 373-392.
INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE. 1957. Opinion 475. Validation under the Plenary Powers of the generic name *Bithynia* Leach, 1818 (Class Gastropoda) and matters associated therewith. *Opinions and declarations rendered by the International Commission on Zoological Nomenclature*, 16(17): 307-330.
INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE. 1965. Opinion 735. *Biomphalaria* Preston, 1910 (Gastropoda): grant under the Plenary Powers of precedence over *Planorbina* Haldeman, 1842, *Taphius* H. & A. Adams, 1855, and *Armigerus* Clessin 1884. *Bull. zool. Nomencl.*, 22(2): 94-99.
ISOM, Billy G., DENNIS, Sally D. & GOOCH, Charles. 1979. Rediscovery of some pleurocerids (Gastropoda) near Muscle Shoals. *Nautilus*, 93(2/3): 69-70.

JACOBSON, Morris K. 1969. On *Lymnaea pseudopinguis* (F.C. Baker). *Nautilus*, 82(3): 110-112.
JACOBSON, Morris K. & EMERSON, William K. 1961. *Shells from Cape Cod to Cape May.* Dover Publ., New York. xviii, 152 pp.
JAY, John C. 1839. *A catalogue of the shells, arranged according to the Lamarckian system; together with descriptions of rare species, contained in the collection of John C. Jay, M.D.* Third edition. Wiley & Putnam, New York. 126 pp., 10 pls.
JEWELL, Dorothea Dowd. 1931. Observations on reproduction in the snail *Goniobasis*. *Nautilus*, 44(4): 115-119.
JOHNSON, Bert M. 1952. Ciliary feeding in *Pomacea paludosa* (Say). *Nautilus*, 66(1): 3-5.
JOHNSON, Charles W. 1913. *Lymnaea (Radix) auricularia* in Charles River, Boston, Mass. *Nautilus*, 27(7): 83.
JOHNSON, Charles W. 1915a. *Viviparus malleatus* Reeve in Massachusetts. *Nautilus*, 29(3): 35-36.
JOHNSON, Charles W. 1915b. Fauna of New England. 13. List of the Mollusca. *Occ. Pap. Boston Soc. nat. Hist.*, (7): 1-231.
JOHNSON, Charles W. 1916a. *Viviparus malleatus* Reeve. *Nautilus*, 30(4): 48.
JOHNSON, Charles W. 1916b. [Publications of the O.G.B. III. Mollusca of Cincinnati. Cincinnati: L.A. Burdsal, Printer.] *Nautilus*, 30(5): 57.
JOHNSON, Charles W. 1916c. *Viviparus contectoides* Binn. in Boston, Mass. *Nautilus*, 30(6): 72.
JOHNSON, Charles W. 1918. *Viviparus malleatus* and *contectoides* in Massachusetts. *Nautilus*, 31(3): 107-108.
JOHNSON, Charles W. 1923. *Viviparus japonicus* v. Martens from another locality in Boston. *Nautilus*, 36(3): 105.
JOHNSON, Richard I. 1941. *Viviparus contectoides* Binney in Massachusetts. *Nautilus*, 54(4): 141.
JOHNSON, Richard I. 1973. Distribution of Hydrobiidae, a family of fresh and brackish water gastropods, in peninsular Florida. *Occ. Pap. Moll.*, 3(46): 281-303.
JOHNSON, Richard I. 1975. First paper on the conchology of the United States by an American author, Thomas Say, 1817. *J. Soc. Bibliogr. nat. Hist.*, 7(3): 265-267.
JOKINEN, Eileen H. 1977. The formation and structure of the shell varix in *Stagnicola elodes* (Say) (Gastropoda: Lymnaeidae). *Nautilus*, 91(1): 13-15.
JOKINEN, Eileen H. 1978. Habitats of two freshwater limpets (*Ferrissia*: Ancylidae) from New England. *Nautilus*, 92(4): 156-160.
JONES, W.C., Jr. & BRANSON, Branley A. 1964. The radula, genital system, and external morphology in *Mudalia potosiensis* (Lea) 1841 (Gastropoda: Prosobranchia: Pleuroceridae) with life history notes. *Trans. Am. microsc. Soc.*, 83(1): 41-62.
KATSIGIANIS, T.S. & HARMAN, Willard N. 1973. Variation in the radular teeth of *Helisoma anceps* (Menke). *Nautilus*, 87(1): 5-7.
KEEP, Josiah. 1887. *West Coast shells.* Bancroft Bros., San Francisco. 230 pp.
KEEP, Josiah. 1904. *West American shells.* Whitaker and Ray, San Francisco. 360 pp.
KEEP, Josiah. 1911 [1910]. *West Coast shells.* [Includes a chapter (chapt. 13, pp. 299-318, pls. 1-3) on "Shells of lakes and streams" by Harold Hannibal.] Whitaker and Ray-Wiggin, San Francisco. 346 pp.
KEEP, Josiah & BAILY, Joshua L., Jr. 1935. *West Coast shells.* Stanford Univ. Press. i-xi, 350 pp.
KEFERL, Eugene P. 1969. Some Mollusca of Cedar Bog, Champaign County, Ohio. *Annu. Rep. Am. malacol. Union 1969*, (36): 14-16.
KESLER, David H. 1980. Seasonal abundance of *Amnicola limosa* (Hydrobiidae) eggs and individuals in a Rhode Island pond. *Nautilus*, 94(1): 25-26.
KIRKLAND, R.J. 1899. Station of *Limnaea gracilis*. *Nautilus*, 12(10): 119-120 [see also correction in *Nautilus*, 12(11): 131].
KIRKLAND, R.J. 1900. Note on the habits of *Limnaea mighelsi* W.G. Binn. *Nautilus*, 14(1): 8-9.
KNIGHT, J. Brookes, COX, L.R., KEEN, A. Myra, BATTEN, R.L., YOCHELSON, E.L. & ROBERTSON, Robert. 1960. Systematic descriptions. *In:* Moore, Raymond C., (Ed.), *Treatise on Invertebrate Paleontology.* Pt. I, Mollusca 1, pp. i-xxiii, 1-351 [pp. 169-310].
KRIEGER, K.A. 1972. *Somatogyrus alcoviensis*, new gastropod species from Georgia (Hydrobiidae). *Nautilus*, 85(4): 120-125.
KRIEGER, K.A. & BURBANCK, W.D. 1976. Distribution and dispersal mechanisms of *Oxytrema* (=*Goniobasis*) *suturalis* Haldeman (Gastropoda: Pleuroceridae) in the Yellow River, Georgia, U.S.A. *Am. midl. Nat.*, 95(1): 49-63.
KUSHLAN, James A. 1975. Population changes of the Apple Snail, *Pomacea paludosa*, in the southern Everglades. *Nautilus*, 89(1): 21-23.
KUSHLAN, James A. 1978. Predation on Apple Snail eggs (*Pomacea*). *Nautilus*, 92(1): 57-58.
KÜSTER, H.C. 1852. Die Gattungen *Paludina, Hydrocaena* und *Valvata*. *In: Systematisches Conchylien-Cabinet von Martini und Chemnitz.* Bauer und Raspe, Nürnberg. 1(21): 1-96, pls. 1-14.

KÜSTER, H.C. 1862. Die Gattungen *Limnaeus, Amphipeplea, Chilina, Isidora* und *Physopsis. In: Systematisches Conchylien-Cabinet von Martini und Chemnitz.* Bauer und Raspe, Nürnberg. 1(17B): 1-77, pls. 1-12.

LAMARCK, M. le Chevalier de. 1822. *Histoire naturelle des animaux sans vertèbres, présentant les caractères généraux et particuliers de ces animaux, leur distribution, leur classes, leurs familles, leurs genres, et la citation des principales espèces qui s'y rapportent.* Vol. 6, Pt. 2, pp. 1-232. Chez l'auteur, au Jardin du Roi.

LANG, Bruce Z. 1968. Note on ecology of *Goniobasis proxima* in North Carolina. *Nautilus,* 82(1): 3-5.

LANG, Bruce Z. & DRONEN, Norman O. 1970. Eggs and attachment sites for egg capsules of *Valvata lewisi. Nautilus,* 84(1): 9-12.

LAPHAM, J.A. 1860. A list of shells of the state of Wisconsin. *Proc. Acad. nat. Sci. Phila.,* 12: 154-156.

LA ROCQUE, Aurèle. 1933a. A shell new to the Canadian list. *Can. Field-Nat.,* 47(5): 96.

LA ROCQUE, Aurèle. 1933b. Notes on *Helisoma latchfordi* Pils. and *Physa latchfordi* F.C. Baker. *Can. Field-Nat.,* 47(7): 134-135.

LA ROCQUE, Aurèle. 1934. Note on the range of *Valvata lewisi ontariensis* F.C. Baker. *Can. Field-Nat.,* 48(2): 39-40.

LA ROCQUE, Aurèle. 1948. Distributional notes on Canadian Mollusca. *Can. Field-Nat.,* 62(1): 36-37.

LA ROCQUE, Aurèle. 1953. Catalogue of the Recent Mollusca of Canada. *Natl. Mus. Can. Bull.,* (129): i-vii, 1-406.

LA ROCQUE, Aurèle. 1956. Variation of carinae in *Valvata tricarinata. Nautilus,* 70(1): 13-14.

LA ROCQUE, Aurèle. 1959a. Plans for a checklist of North American Mollusca. *Sterkiana,* (1): 19-22.

LA ROCQUE, Aurèle. 1959b. Checklist of Ohio Pleistocene and living Mollusca. *Sterkiana,* (1): 23-49.

LA ROCQUE, Aurèle. 1961a. Checklist of New Brunswick non-marine Mollusca. *Sterkiana,* (3): 40-42.

LA ROCQUE, Aurèle. 1961b. Checklist of Newfoundland non-marine Mollusca. *Sterkiana,* (3): 43-48.

LA ROCQUE, Aurèle. 1962. Checklist of the non-marine Mollusca of Quebec. *Sterkiana,* (7): 23-44.

LA ROCQUE, Aurèle. 1963. Late Cenozoic non-marine molluscan associations in eastern North America. *Sterkiana,* (11): 1-50; (12): 15-60; (13): 23-53; (14): 19-38.

LA ROCQUE, Aurèle. 1965. Changes in the molluscan fauna of White Lake, Lanark and Renfrew counties, Ontario, after thirty years. *Sterkiana,* (19): 1-2, 40.

LA ROCQUE, Aurèle. 1968. Pleistocene Mollusca of Ohio. *Div. geol. Surv. Dept. nat. Res. State of Ohio Bull.,* 62(3): xv-xxiv, 357-553, pls. 9-14.

LA ROCQUE, Aurèle. 1970-71. Un manuscrit inédit de l'Abbé Provancher sur les mollusques du Canada. *Sterkiana,* (37): 1-26; (41): 1-33.

LATCHFORD, F.R. 1914. *Valvata piscinalis* in Canada. *Nautilus,* 28(1): 10.

LATCHFORD, F.R. 1926. A new *Lymnaea. Can. Field-Nat.,* 40(3): 47-48.

LATCHFORD, F.R. 1927. *Lymnaea (Bulimnaea) megasoma* Say. *Nautilus,* 40(4): 136-137.

LEA, Isaac. 1831a. Observations on the naiades, and descriptions of new species of that and other families. *Trans. Am. philos. Soc.,* 4: 63-105, pls. 3-15.

LEA, Isaac. 1831b. Observations on the naiades, and descriptions of new species of that and other families. Supplement. *Trans. Am. philos. Soc.,* 4: 106-121, pls. 15-18.

LEA, Isaac. 1831c. Description of a new genus of the family Melaniana of Lamarck. *Trans. Am. philos. Soc.,* 4: 122-123.

LEA, Isaac. 1834. Observations on the naiades; and descriptions of new species of that and other families. *Trans. Am. philos. Soc.,* 5: 23-119, pls. 1-19.

LEA, Isaac. 1834-74. *Observations on the genus* Unio, ... etc. ... [titles vary], 1: 233 pp., 46 pls., 1829 [1834]; 2: 1-152, pls. 1-24, 1838; 3: 1-88, pls. 5-27, 1842; 4: 1-75, 13 pls., 1848; 5: 1-62, pls. 12-30, 1852; 9: 1-180, pls. 24-39, 1863; 11: 1-146, pls. 1-24, 1867; Index, vols. 1-11, pp. 1-63, 1867; 12: 1-105, pls. 29-54, Index, pp. 1-23, 1869; 13: 1-75, pls. 1-22, Index, vols. 1-13, pp. 1-29, 1874. Printed for the author, Philadelphia.

LEA, Isaac. 1838a. Description of new freshwater and land shells. *Trans. Am. philos. Soc.,* 6: 1-111, pls. 1-24.

LEA, Isaac. 1838b. *Melania cincinnatiensis. Proc. Am. philos. Soc.,* 1(5): 66.

LEA, Isaac. 1840. Descriptions of new fresh-water and land shells. *Proc. Am. philos. Soc.,* 1(13): 284-290.

LEA, Isaac. 1841a. Continuation of Mr. Lea's paper on new fresh water and land shells. *Proc. Am. philos. Soc.,* 2(16): 11-15.

LEA, Isaac. 1841b. On fresh water and land shells [cont.]. *Proc. Am. philos. Soc.,* 2(17): 30-34.

LEA, Isaac. 1841c. Continuation of Mr. Lea's paper on fresh water and land shells. *Proc. Am. philos. Soc.,* 2(19): 81-83.

LEA, Isaac. 1842a. [On the occurrence of *Melania altilis* on the shore of the Schuylkill near Philadelphia.] *Proc. Am. philos. Soc.,* 2(21): 150.

LEA, Isaac. 1842b. Description of new fresh water and land shells. *Trans. Am. philos. Soc.*, 8(2): 163-250, pls. 5-27.
LEA, Isaac. 1843. Continuation of paper on fresh water shells. *Proc. Am. philos. Soc.*, 2(24): 241-243.
LEA, Isaac. 1844. Continuation of Mr. Lea's paper on fresh water and land shells. *Trans. Am. philos. Soc.*, 9(1): 1-31.
LEA, Isaac. 1845. Descriptions of new fresh water and land shells. *Proc. Am. philos. Soc.*, 4(33): 162-168.
LEA, Isaac. 1848a. Description of new fresh water and land shells. *Trans. Am. philos. Soc.*, 10(1): 67-96, 100.
LEA, Isaac. 1848b. Catalogue of the genus *Melania*. *Trans. Am. philos. Soc.*, 10(1): 96-100.
LEA, Isaac. 1852. Description of a new genus (*Basistoma*) of the family Melaniana, together with some new species of American Melaniae. *Trans. Am. philos. Soc.*, 10: 295-302, pl. 30.
LEA, Isaac. 1853. Description of a new genus (*Basistoma*) of the family Melaniana, together with some new species of American Melaniae. *Proc. Am. philos. Soc.*, 5(48): 252.
LEA, Isaac. 1856. Description of new fresh-water shells from California. *Proc. Acad. nat. Sci. Phila.*, 8: 80-81.
LEA, Isaac. 1858a. Descriptions of a new *Helix* and two new planorbes. *Proc. Acad. nat. Sci. Phila.*, 10: 41.
LEA, Isaac. 1858b. Descriptions of twelve new species of uniones and other fresh-water shells of the United States. *Proc. Acad. nat. Sci. Phila.*, 10: 165-166.
LEA, Isaac. 1860a. Descriptions of four new species of Melanidae of the United States. *Proc. Acad. nat. Sci. Phila.*, 12: 93.
LEA, Isaac. 1860b. [Remarks on letter of Dr. Showalter, of Ala., on specimens of *Physa* (*gyrina*), Say, obtained in water supplied by an artesian well.] *Proc. Acad. nat. Sci. Phila.*, 12: 147.
LEA, Isaac. 1860c. Descriptions of fourteen new species of Schizostomae, Anculosae and Lithasiae. *Proc. Acad. nat. Sci. Phila.*, 12: 186-188.
LEA, Isaac. 1861a. Descriptions of new species of *Schizostoma*, *Anculosa*, and *Lithasia*. *Proc. Acad. nat. Sci. Phila.*, 13: 54-55.
LEA, Isaac. 1861b. Description of a new species of *Neritina*, from Coosa River, Alabama. *Proc. Acad. nat. Sci. Phila.*, 13: 55.
LEA, Isaac. 1861c. Description of a new genus (*Strephobasis*) of the family Melanidae, and three new species. *Proc. Acad. nat. Sci. Phila.*, 13: 96.
LEA, Isaac. 1861d. Descriptions of forty-nine new species of the genus *Melania*. *Proc. Acad. nat. Sci. Phila.*, 13: 117-123.
LEA, Isaac. 1861e. Description of seven new species of the genus *Io*. *Proc. Acad. nat. Sci. Phila.*, 13: 393-394.
LEA, Isaac. 1862a. Description of a new genus (*Trypanostoma*), of the family Melanidae, and of forty-five new species. *Proc. Acad. nat. Sci. Phila.*, 14: 169-175.
LEA, Isaac. 1862b. Description of a new genus (*Goniobasis*) of the family Melanidae and eighty-two new species. *Proc. Acad. nat. Sci. Phila.*, 14: 262-272.
LEA, Isaac. 1862c. Descriptions of eleven new species of Melanidae of the United States. *Proc. Acad. nat. Sci. Phila.*, 14: 272-273.
LEA, Isaac. 1863a. New Melanidae of the United States. *J. Acad. nat. Sci. Phila.*, 5: 217-356, pls. 34-39.
LEA, Isaac. 1863b. Descriptions of fourteen new species of Melanidae and one *Paludina*. *Proc. Acad. nat. Sci. Phila.*, 15: 154-156.
LEA, Isaac. 1864a. Description of a new genus of the family Melanidae. *Proc. Acad. nat. Sci. Phila.*, 16: 2-3.
LEA, Isaac. 1864b. Descriptions of eleven new indigenous species of Melanidae. *Proc. Acad. nat. Sci. Phila.*, 16: 3-5.
LEA, Isaac. 1864c. Description of and remarks on *Planorbis newberryi*. *Proc. Acad. nat. Sci. Phila.*, 16: 5.
LEA, Isaac. 1864d. Description of six new species of Succineidae of the United States. *Proc. Acad. nat. Sci. Phila.*, 16: 109-111.
LEA, Isaac. 1864e. Description of a new species of *Planorbis*. *Proc. Acad. nat. Sci. Phila.*, 16: 111.
LEA, Isaac. 1864f. Descriptions of thirteen new species of Melanidae of the United States. *Proc. Acad. nat. Sci. Phila.*, 16: 111-113.
LEA, Isaac. 1864g. Description of five new species of *Lymnaea* of North America. *Proc. Acad. nat. Sci. Phila.*, 16: 113.
LEA, Isaac. 1864h. Descriptions of twenty-four new species of *Physa* of the United States and Canada. *Proc. Acad. nat. Sci. Phila.*, 16: 114-116.
LEA, Isaac. 1866a. Description of two new species of the genus *Lithasia*. *Proc. Acad. nat. Sci. Phila.*, 18: 133.
LEA, Isaac. 1866b. New Unionidae, Melanidae, etc., chiefly of the United States. *J. Acad. nat. Sci. Phila.*, 6: 113-187, pls. 22-24.
LEA, Isaac. 1867. Descriptions of five new species of Unionidae and one *Paludina* of the United States. *Proc. Acad.*

nat. Sci. Phila., 19: 81.
LEA, Isaac. 1868a. Descriptions of twenty-six new species of Melanidae of the United States. *Proc. Acad. nat. Sci. Phila.*, 20: 151-153.
LEA, Isaac. 1868b. New Unionidae, Melanidae, etc., chiefly of the United States. *J. Acad. nat. Sci. Phila.*, 6: 303-343, pls. 46-54.
LEA, Isaac. 1869. Descriptions of six new species of fresh water shells. *Proc. Acad. nat. Sci. Phila.*, 21: 124-125.
LEA, Isaac. 1874. Supplement to Isaac Lea's paper on Unionidae. *J. Acad. nat. Sci. Phila.*, 8: 55-69, pls. 19-22.
LEONARD, A. Byron. 1959. Handbook of gastropods in Kansas. *Univ. Kans. Mus. nat. Hist. misc. Publ.*, (20): 1-224, 1 map.
LETSON, Elizabeth J. 1905. Check list of the Mollusca of New York. *N.Y. State ed. Dept. Bull.*, (341)[N.Y. State Mus. Bull. 88, zool. 11]: 1-112.
LEWIS, James. 1854a. [On *Lymnaea catascopium*.] *Proc. Boston Soc. nat. Hist.*, 5: 27-29.
LEWIS, James. 1854b. Notes on *Paludina decisa* Say. *Proc. Boston Soc. nat. Hist.*, 5: 52-53.
LEWIS, James. 1855. Remarks on *Cyclas* and *Lymnea*. *Proc. Boston Soc. nat. Hist.*, 5: 122-124.
LEWIS, James. 1863a. [Letter to Isaac Lea read before the Academy of Natural Sciences of Philadelphia.] *Proc. Acad. nat. Sci. Phila.*, 15: 26.
LEWIS, James. 1863b. On the identity of *Paludina subglobosa* Say with *Melania isogona* Say. *Proc. Boston Soc. nat. Hist.*, 9: 252.
LEWIS, James. 1869. Observations on *Melantho*. *Am. J. Conchol.*, 5: 33-36.
LEWIS, James. 1871. On the shells of the Holston River. *Am. J. Conchol.*, 7(2): 216-226.
LEWIS, James. 1876. Fauna of Alabama. 1. Fresh water and land snails. *Geol. Surv. Ala.*, pp. 1-40.
LINNAEUS, Carolus A. 1758. *Systema Naturae, per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis.* 10th edition. Laurentii Salvii, Holmiae. 824 pp.
LINNÉ [LINNAEUS], Caroli A. 1766. *Systema Naturae.* Editio Duodecima. Reprinted by Hanley, Sylvanus, 1855. Ipsa Linnaei conchylia. *The shells of Linnaeus, determined from his manuscript and collection. Also, an exact reprint of the* Vermes testacea *of the* 'Systema naturae' *and* 'Mantissa'. Williams & Norgate, London. 556 pp., 5 pls.
LONG, Bayard. 1912. *Lymnaea auricularia* near Philadelphia. *Nautilus*, 26(3): 27-29.
LoVERDE, P.T. & BURCH, J.B. 1973. The systematic position of *"Stagnicola (Bakerilymnaea) cubensis"*. *Malacol. Rev.*, 6(1): 44.
LUTZ, Louis & WEESE, A.O. 1951. Variation in *Io*, fifty years later. *Proc. Okla. Acad. Sci.*, 32: 1-3 [reprint pagination].
MACKIE, Gerry L., WHITE, David S. & ZDEBA, Thomas W. 1980. *A guide to freshwater mollusks of the Laurentian Great Lakes with special emphasis on the genus* Pisidium. Environ. Res. Lab., Off. Res. Dev. U.S. Environ. Protection Agency, Duluth, Minn. pp. i-vi, 1-144.
MacNAMARA, Mark C. & HARMAN, Willard N. 1975. Further studies of the Mollusca of the Otsego Lake area. *Nautilus*, 89(3): 87-90.
MAGRUDER, Samuel R. 1934. Notes on the life history of *Pleurocera canaliculatum* Say. *Nautilus*, 48(1): 26-28.
MAGRUDER, Samuel R. 1935. Record of a crystalline style in two fresh water gastropods. *Nautilus*, 48(3): 101-102.
MALEK, Emile A. 1969. Studies on "tropicorbid" snails (*Biomphalaria*: Planorbidae) from the Caribbean and Gulf of Mexico areas, including the southern United States. *Malacologia*, 7(2/3): 183-209.
MALEK, Emile A. & CHROSCIECHOWSKI, P. 1964. *Lymnaea (Pseudosuccinea) columella* from Venezuela, and notes on distribution of *Pseudosuccinea*. *Nautilus*, 78(2): 54-56.
MARSHALL, William B. 1931. *Monocondylaea costulata* Moricand. *Nautilus*, 45(2): 55-58.
MARSHALL, William B. 1933. *Bulimus tentaculatus* (L.) (*Bithynia tentaculata*) living in the Potomac River. *Nautilus*, 46(4): 141-142.
MATTOX, Norman Thomas. 1937. Oogenesis of *Campeloma rufum*, a parthenogenetic snail. *Z. Zellforsch. mikrosk. Anat.*, 27(4): 455-464.
MATTOX, Norman Thomas. 1938. Morphology of *Campeloma rufum*, a parthenogenetic snail. *J. Morphol.*, 62(2): 243-261.
MATTOX, Norman Thomas. 1940. Two new snails of the genus *Campeloma* from Ontario. *Nautilus*, 54(1): 12-17.
MAZYCK, William Gaillard. 1913. Catalog of the Mollusca of South Carolina. *Contrib. Charleston Mus.*, (2): i-xvi, 1-39.
McCLARY, Andrew. 1964. Surface inspiration and ciliary feeding in *Pomacea paludosa* (Prosobranchia: Mesogastropoda: Ampullariidae). *Malacologia*, 2(1): 87-104.
McCOY, Clarence J., Jr. 1964. On *Lymnaea auricularia* in Colorado. *Nautilus*, 78(2): 66.

McCRAW, Bruce M. 1952. Observations on the development of *Lymnaea palustris* Müller. *Can. J. Zool.*, 30: 378-386.
McCRAW, Bruce M. 1957. Studies on the anatomy of *Lymnaea humilis* Say. *Can. J. Zool.*, 35: 751-768.
McCRAW, Bruce M. 1959. The ecology of the snail, *Lymnaea humilis* Say. *Trans. Am. J. microsc. Soc.*, 78(1): 101-121.
McCRAW, Bruce M. 1970. Aspects of the growth of the snail *Lymnaea palustris* (Müller). *Malacologia*, 10(2): 399-413.
McDONALD, Sharon Lee Chalfant. 1969. The biology of *Lymnaea stagnalis* L. (Gastropoda: Pulmonata). *Sterkiana*, (36): 1-17.
McDONALD, Sharon Lee Chalfant. 1973. Activity patterns of *Lymnaea stagnalis* (L.) in relation to temperature conditions: a preliminary study. *Malacologia*, 14(1/2): 395-396.
McLEOD, M.J. & MOORE, J.D. 1978. Change in the gastropod *Io spinosa* (Pleuroceridae: Mollusca) in 70 years. *Am. midl. Nat.*, 99(1): 198-205.
McMAHON, Robert F. 1976a. Growth, reproduction and life cycle in six Texan populations of two species of freshwater limpets. *Am. midl. Nat.*, 95(1): 174-185.
McMAHON, Robert F. 1976b. Effluent-induced interpopulation variation in the thermal tolerance of *Physa virgata* Gould. *Comp. Biochem. Physiol.*, 55A: 23-28.
METCALF, Artie L. 1968. Mollusca of El Paso County, westernmost Texas. *Annu. Rep. Am. malacol. Union 1968*, (35): 32-33.
METCALF, Artie L. & SMARTT, Richard. 1972. Records of introduced mollusks: New Mexico and western Texas. *Nautilus*, 85(4): 144-145.
MEYER, Emmett. 1928. Finding a left-handed *Campeloma*. *Nautilus*, 41(3): 107.
MICHELSON, Edward H. 1961. On the generic limits in the family Pilidae. *Breviora*, (133): 1-10.
MICHELSON, Edward H. & DuBOIS, Lorin. 1974. *Lymnaea emarginata*, a possible agent for the control of the schistosome-snail host, *Biomphalaria glabrata*. *Nautilus*, 88(4): 101-108.
MIGHELS, J.W. & ADAMS, C.B. 1842. Descriptions of twentyfour species of the shells of New England. *Boston J. nat. Hist.*, 4(1): 37-54, pl. 4.
MILLER, Barry B. 1966. Five Illinoian molluscan faunas from the southern Great Plains. *Malacologia*, 4(1): 173-260.
MILLER, Barry B. & HIBBARD, Claude W. 1972. Recent Mollusca of Ellsworth County, Kansas. *Sterkiana*, (46): 11-14.
MORRIS, Percy A. 1939. *What shell is that?* A guide to the shell-bearing mollusks of eastern North America. D. Appleton-Century Co., New York and London. vii, 198 pp.
MORRISON, Joseph P.E. 1932. Studies on the life history of *Acella haldemani* ("Desh." Binney). *Trans. Wis. Acad. Sci. Arts Lett.*, 27: 397-414.
MORRISON, Joseph P.E. 1939. The type locality of *Cochliopa rowelli* (Tryon). *Nautilus*, 53(2): 67.
MORRISON, Joseph P.E. 1940a. A new species of *Fluminicola* with notes on "Colorado Desert" shells, and on the genus *Clappia*. *Nautilus*, 53(4): 124-127.
MORRISON, Joseph P.E. 1940b. Notes on the subgenera of *Valvata*. *Nautilus*, 53(4): 138-140.
MORRISON, Joseph P.E. 1941. Population studies on *Goniobasis*. *Rep. Am. malacol. Union 1940*, (9): 9.
MORRISON, Joseph P.E. 1942. Preliminary report on mollusks found in the shell mounds of the Pickwick Landing basin in the Tennessee River valley. *Bur. Am. Ethnol. Bull.*, (129): 339-392.
MORRISON, Joseph P.E. 1946. The nonmarine mollusks of San José Island, with notes on those of Pedro González Island, Pearl Islands, Panamá. *Smithson. misc. Collect.*, 106(6): 1-49, pls. 1-3.
MORRISON, Joseph P.E. 1947a. One hundred and six years of *Amnicola*. *Nautilus*, 60(3): 84-87.
MORRISON, Joseph P.E. 1947b. Notes on the genus *Probythinella* (Hydrobiinae). *Nautilus*, 61(1): 25-30.
MORRISON, Joseph P.E. 1949. The cave snails of eastern North America. *News Bull. annu. Rep. Am. malacol. Union 1948*, (15): 13-14.
MORRISON, Joseph P.E. 1952. World relations of the melanians. *News Bull. annu. Rep. Am. malacol. Union 1951*, 18: 6-9.
MORRISON, Joseph P.E. 1953. *Viviparus multilineatus* (Say) 1829, from Florida. *Nautilus*, 67(2): 56-58.
MORRISON, Joseph P.E. 1954. The relationships of Old and New World melanians. *Proc. U.S. natl. Mus.*, 103(3325): 357-394, pl. 11.
MORRISON, Joseph P.E. 1955a. Notes on the genera *Lanx* and *Fisherola* (Pulmonata). *Nautilus*, 68(3): 79-83.
MORRISON, Joseph P.E. 1955b. How many *Syncera* species are living in Death Valley? *Annu. Rep. Am. malacol. Union 1955*, (22): 29.
MORRISON, Joseph P.E. 1969. Zoogeography of hydrobiid cave snails. *Malacologia*, 9(1): 278.

MORRISON, Joseph P.E. 1970. Names for the subfamily Hydrobiinae. *Annu. Rep. Am. malacol. Union 1970*, (37): 7-8.

MORRISON, Joseph P.E. 1971. *Athearnia*, a new name for a genus of pleurocerid snails. *Nautilus*, 84(3): 110-111.

MORSE, Edward S. 1864. Observations on the terrestrial Pulmonifera of Maine, including a catalogue of all the species of terrestrial and fluviatile Mollusca known to inhabit the state. *J. Portland Soc. nat. Hist.*, 1(1): 1-63, pls. 1-10.

MORSE, Edward S. 1880. The gradual dispersion of certain mollusks in New England. *Bull. Essex Inst.*, 12: 3-8. [Reprinted in *Sterkiana*, (17): 7-9.]

MOZLEY, Alan. 1932. Notes on western Canadian Mollusca – *Planorbis campanulatus wisconsinensis* Winslow. *Nautilus*, 46(2): 59-62.

MOZLEY, Alan. 1933. A new variety of *Lymnaea* from the Rocky Mountain region of Canada. *Proc. malacol. Soc. Lond.*, 20(5): 241-243, pl. 22.

MÜLLER, Othone Friderico. 1774. *Vermium terrestrium et fluviatilium, seu animalium infusoriorum, helminthicorum et testaceorum, non marinorum, succincta historia.* Heineck et Faber, Havniae et Lipsiae. Vol. 2, xxxvi, 214 pp.

MURRAY, Harold D. 1964. *Tarebia granifera* and *Melanoides tuberculata* in Texas. *Annu. Rep. Am. malacol. Union 1964*, (31): 15-16.

MURRAY, Harold D. 1970. Discussion of Dr. Taylor's paper [on rare and endangered western freshwater mollusks, in Am. malacol. Union Symp. Rare and Endangered Mollusks]. *Malacologia*, 10(1): 33-34.

MURRAY, Harold D. 1976. *Melanoides tuberculata* (Muller), Las Moras Creek, Bracketville, Texas. *Bull. Am. malacol. Union Inc. 1975*, (42): 43.

MURRAY, Harold D. & ROY, Edward C., Jr. 1968. Checklist of freshwater and land mollusks of Texas. *Sterkiana*, (30): 25-42.

MURRAY, Harold D. & WOPSCHALL, Leon J. 1965. Ecology of *Melanoides tuberculata* (Müller) and *Tarebia granifera* (Lamarck) in south Texas. *Annu. Rep. Am. malacol. Union 1965*, (32): 25-26.

NEVILL, Geoffrey. 1885. *Hand list of the Mollusca in the Indian Museum, Calcutta. Pt. 2. Gastropoda. Prosobranchia-Neurobranchia (contd.).* Trustees of the Indian Museum, Calcutta. 306 pp.

NYLANDER, Olof O. 1899. [Notes on *Pisidium contortum, Pi. milium* and *Planorbis crista cristata*.] *Nautilus*, 13(5): 59-60.

NYLANDER, Olof O. 1901. Distribution of *Limnaea emarginata*, Say, and the var. *mighelsi*, Binney, in Fish River, Aroostook Co., Maine. Publ. by author. pp. 1-4, pls. 1-4.

NYLANDER, Olof O. 1902. The original locality of *Limnaea ampla* Mighels. *Nautilus*, 15(11): 127-129.

NYLANDER, Olof O. 1912. Lymnaeidae of Aroostook County, Maine. *Nautilus*, 25(9): 107-108.

NYLANDER, Olof O. 1914. *Lymnaea emarginata mighelsi* absent in its old localities. *Nautilus*, 28(8): 95-96.

NYLANDER, Olof O. 1921. The type localities of *Lymnaea emarginata* Say and *L. ampla* Mighels. *Nautilus*, 34(3): 77-80.

NYLANDER, Olof O. 1930. The *Valvata* of Aroostook Co., Maine. *Nautilus*, 44(1): 30-31.

NYLANDER, Olof O. 1942a. *Lymnaea stagnalis* in Chaleur Bay, Quebec. *Rep. Am. malacol. Union 1941*, (10): 1-2.

NYLANDER, Olof O. 1942b. *Lymnaea stagnalis* at Houlton, Maine. *Rep. Am. malacol. Union 1941*, (10): 2.

OUGHTON, John. 1938. *Valvata piscinalis* (Müller) in the Great Lakes. *Nautilus*, 52(1): 30-32; 52(2): 60-62.

OUGHTON, John. 1939. An observation on *Bulimnea megasoma*. *Nautilus*, 52(3): 106-107.

ORTMANN, Arnold E. 1913. The Alleghenian Divide, and its influence upon the freshwater fauna. *Proc. Am. philos. Soc.*, 52(210): 287-390, pls. 12-14.

OWEN, D.F. 1962. *Helisoma anceps* transported by a giant water bug. *Nautilus*, 75(3): 124-125.

PACE, Gary L. 1968. The distribution and habitats of *Carinifex* and *Parapholyx*. *Annu. Rep. Am. malacol. Union 1967*, (34): 32-33.

PACE, Gary L. 1971. The hold-fast function of the preputial organ in *Helisoma*. *Malacol. Rev.*, 4(1): 21-24.

PACE, Gary L. 1973a. Taxonomic studies on *Carinifex* and *Parapholyx* (Gastropoda: Planorbidae). *Echo*, (5): 39-40.

PACE, Gary L. 1973b. The freshwater snails of Taiwan (Formosa). *Malacol. Rev.*, suppl. 1, 118 pp.

PACE, Gary L., SZUCH, Ernest J. & DAPSON, Richard W. 1979. Depth distribution of three gastropods in New Mission Bay, Lake Michigan. *Nautilus*, 93(1): 31-36.

PARAENSE, W. Lobato. 1976. *Helisoma trivolvis* and some of its synonyms in the Neotropical region (Mollusca: Planorbidae). *Rev. Bras. Biol.*, 36(1): 187-204.

PATTERSON, C.M. 1963. Cytological studies of *Pomatiopsis* snails. *Annu. Rep. Am. malacol. Union 1963*, (30): 13-14.

PATTERSON, C.M. 1965. The chromosomes of *Tulotoma angulata* (Streptoneura: Viviparidae). *Malacologia*, 2(2): 259-265.
PATTERSON, C.M. 1967. Chromosome numbers and systematics in streptoneuran snails. *Malacologia*, 5(2): 111-125.
PENNAK, Robert W. 1953. *Fresh-water invertebrates of the United States.* First edition. Ronald Press, New York. 769 pp. [gastropods on pp. 667-693].
PENNAK, Robert W. 1978. *Freshwater invertebrates of the United States.* Second edition. John Wiley and Sons, New York. 803 pp. [mollusks on pp. 710-768].
PERRON, Frank & PROBERT, Thomas. 1973. *Viviparus malleatus* in New Hampshire. *Nautilus*, 87(3): 90.
PFEIFFER, Louis. 1849. Neue Molluskengattungen. *Z. Malakozool.*, 6(7): 97-105.
PILSBRY, Henry A. 1882. List of land and fresh-water Mollusca, found at Iowa City, Iowa. Printed by the author. 4 pp.
PILSBRY, Harry [Henry] A. 1886a. Notes on some eastern Iowa snails. *Am. Nat.*, 20(1): 75.
PILSBRY, Henry A. 1886b. Notes on some New Orleans fresh-water shells. *Conchol. Exc.*, 1(5): 20-21.
PILSBRY, Henry A. 1887a. Notes on the larger Florida planorbes. *Am. Nat.*, 21: 286-287.
PILSBRY, Harry [Henry] A. 1887b. Description of a new *Hydrobia*, with notes on other Rissoidae. *Proc. Davenport Acad. nat. Sci.*, 5: 33-34 [pl. 3, figs. 1-6, with separates].
PILSBRY, Henry A. 1888. *Lyogyrus*, Gill, and other American shells. *Conchol. Exc.*, 2(9): 113.
PILSBRY, Henry A. 1889a. New and little-known American molluscs. No. 1. *Proc. Acad. nat. Sci. Phila.*, 41: 81-89, pl. 3.
PILSBRY, Henry A. 1889b. Note upon Mr. Ancey's criticism. *Nautilus*, 3(4): 42-44.
PILSBRY, Henry A. 1889c. Recent additions to the United States snail fauna. *Nautilus*, 3(6): 61-64, pl. 1, figs. 1-4, 6-7, 9.
PILSBRY, Henry A. 1890a. *Limnaea columella* Say. *Nautilus*, 4(4): 47.
PILSBRY, Henry A. 1890b. A new species of fresh-water mollusk, by B. Shimek. [Review.] *Nautilus*, 4(4): 48.
PILSBRY, Henry A. 1890c. Remarks on certain goniobases. *Nautilus*, 4(5): 49-50.
PILSBRY, Henry A. 1890d. Preliminary notices of new Amnicolidae. *Nautilus*, 4(5): 52-53.
PILSBRY, Henry A. 1890e. Notices of new Amnicolidae. *Nautilus*, 4(6): 63-64.
PILSBRY, Henry A. 1890f. Descriptions of new West American land, fresh-water and marine shells, by Robt. E.C. Stearns. [Review.] *Nautilus*, 4(6): 72.
PILSBRY, Henry A. 1890g. New and little-known American mollusks, no. 3. *Proc. Acad. nat. Sci. Phila.*, 42: 296-302, pl. 5.
PILSBRY, Henry A. 1891a. Note on *Goniobasis catenaria* Say. *Nautilus*, 4(11): 124.
PILSBRY, Henry A. 1891b. Land and fresh-water mollusks collected in Yucatan and Mexico. *Proc. Acad. nat. Sci. Phila.*, 43: 310-334, pls. 14, 15.
PILSBRY, Henry A. 1891c. Mollusca from Nantucket, Mass. *Proc. Acad. nat. Sci. Phila.*, 43: 406-407.
PILSBRY, Henry A. 1892a. Preliminary notices of new forms of fresh water mollusks. *Nautilus*, 5(12): 142-143.
PILSBRY, Henry A. 1892b. *Goniobasis virginica* in Conn. *Nautilus*, 6(3): 36.
PILSBRY, Henry A. 1892c. On the amnicoloid genus *Lyogyrus*, with the description of a new species. *Nautilus*, 6(6): 61-62.
PILSBRY, Henry A. 1892d. Species identified. *Nautilus*, 6(7): 83.
PILSBRY, Henry A. 1892e. Note on *Lyogyrus*. *Nautilus*, 6(7): 83.
PILSBRY, Henry A. 1893. [Review of paper on shells collected by the Death Valley Expedition, by Dr. R.E.C. Stearns.] *Nautilus*, 7(3): 35-36.
PILSBRY, Henry A. 1894a. Critical list of mollusks collected in the Potomac Valley. *Proc. Acad. nat. Sci. Phila.*, 46: 11-31, pl. 1.
PILSBRY, Henry A. 1894b. Land and fresh-water shells of Allegheny County, Pa. *Nautilus*, 7(12): 135-136.
PILSBRY, Henry A. 1895a. New forms of American shells. *Nautilus*, 8(9): 102.
PILSBRY, Henry A. 1895b. New American fresh-water mollusks. *Nautilus*, 8(10): 114-116.
PILSBRY, Henry A. 1895c. *Bythinia tentaculata*. *Nautilus*, 8(12): 143.
PILSBRY, Henry A. 1895d. *Planorbis sampsoni* Ancey. *Nautilus*, 9(3): 36.
PILSBRY, Henry A. 1895e. *Planorbis centervillensis* Tryon. *Nautilus*, 9(3): 36.
PILSBRY, Henry A. 1895f. [Additional records of *Gundlachia* in North America.] *In:* Hedley, C., On the Australasian *Gundlachia*. *Nautilus*, 9(6): 61-68 [p. 61, figs. 1, 2].
PILSBRY, Henry A. 1895g. Review of our present knowledge of the molluscan fauna of Michigan. [Review.] *Nautilus*, 9(7): 82.
PILSBRY, Henry A. 1896a. New American Ancylidae. *Nautilus*, 9(12): 137-139.

PILSBRY, Henry A. 1896b. A new species of *Pomatiopsis*. *Nautilus*, 10(4): 37-38.
PILSBRY, Henry A. 1896c. *Limnaea bulimoides* Lea resisting drought. *Nautilus*, 10(8): 96.
PILSBRY, Henry A. 1897. *Campeloma decisum* Say, reversed. *Nautilus*, 10(10): 118.
PILSBRY, Henry A. 1898a. Notes on new and little-known Amnicolidae. *Nautilus*, 12(4): 42-44.
PILSBRY, Henry A. 1898b. Shells of Redding, Shasta Co., California. *Nautilus*, 12(5): 59-60.
PILSBRY, Henry A. 1899a. Catalogue of the Amnicolidae of the western United States. *Nautilus*, 12(11): 121-127.
PILSBRY, Henry A. 1899b. [*Planorbis dilatatus* in the Guadalupe River, Texas.] *Nautilus*, 12(12): 144.
PILSBRY, Henry A. 1899c. New Amnicolidae from Florida. *Nautilus*, 13(2): 20-22.
PILSBRY, Henry A. 1899d. *Planorbis rubellus* Sterki, and *P. harni* Pilsbry. *Nautilus*, 13(5): 51-52.
PILSBRY, Henry A. 1899e. A new *Ampullaria*. *Proc. Acad. nat. Sci. Phila.*, 51: 365.
PILSBRY, Henry A. 1899f. Mollusks collected by R.C. McGregor in northern California. *Nautilus*, 13(6): 64-67.
PILSBRY, Henry A. 1899g. *Physa cubensis* in Florida. *Nautilus*, 13(6): 70.
PILSBRY, Henry A. 1900a. Mollusca of the Great Smoky Mountains. *Proc. Acad. nat. Sci. Phila.*, 52: 110-150.
PILSBRY, Henry A. 1900b. Land snails from rejectamenta of the Rio Grande at Mesilla, New Mexico, and of the Gallinas R. at Las Vegas, N.M. *Nautilus*, 14(1): 9-10.
PILSBRY, Henry A. 1900c. Shells of Las Vegas, N.M. *Nautilus*, 14(4): 47.
PILSBRY, Henry A. 1900d. Notes on certain Mollusca of southwestern Arkansas. *Proc. Acad. nat. Sci. Phila.*, 52: 449-459.
PILSBRY, Henry A. 1903a. Mollusca of western Arkansas and adjacent states, with a revision of *Paravitrea*. *Proc. Acad. nat. Sci. Phila.*, 55: 193-214, pls. 9-11.
PILSBRY, Henry A. 1903b. The greatest American *Planorbis*. *Nautilus*, 17(7): 75-76.
PILSBRY, Henry A. 1903c. Shells of Douglas Co., central Washington. *Nautilus*, 17(7): 84.
PILSBRY, Henry A. 1904. A new Floridian *Amnicola*. *Nautilus*, 17(10): 113-114.
PILSBRY, Henry A. 1905a. *Planorbis alabamensis* and *dilatatus* in the Floridian Pliocene. *Nautilus*, 19(3): 34.
PILSBRY, Henry A. 1905b. Is *Cochliopa rowelli* a Californian shell? *Nautilus*, 19(8): 91-92.
PILSBRY, Henry A. 1905c. Land and fresh-water mollusks of Alaska and adjoining regions. [Review.] *Nautilus*, 19(8): 93-95.
PILSBRY, Henry A. 1906a. Shells of Grant, Valencia Co., New Mexico. *Nautilus*, 19(11): 130.
PILSBRY, Henry A. 1906b. Two new American genera of Basommatophora. *Nautilus*, 20(5): 49-51, pl. 3.
PILSBRY, Henry A. 1906c. Note on *Lepyrium*. *Nautilus*, 20(5): 51.
PILSBRY, Henry A. 1907. A new species of *Fluminicola*. *Nautilus*, 21(7): 75-76, pl. 11, fig. 4.
PILSBRY, Henry A. 1908a. Notes on Plate XI. *Nautilus*, 21(12): 133, pl. 11.
PILSBRY, Henry A. 1908b. *Valvata humeralis californica* n. subsp. *Nautilus*, 22(8): 82.
PILSBRY, Henry A. 1911. The Lymnaeidae of North and Middle America, Recent and fossil. [Review.] *Nautilus*, 25(1): 8-10.
PILSBRY, Henry A. 1912a. A new species of *Amnicola*. *Nautilus*, 26(1): 1, pl. 1, figs. 9, 10.
PILSBRY, Henry A. 1912b. [Review of "A synopsis of the Recent and Tertiary freshwater Mollusca of the Californian Province".] *Nautilus*, 26(6): 71.
PILSBRY, Henry A. 1913. Notes on *Gundlachia* Pfr. *Proc. Acad. nat. Sci. Phila.*, 65: 668-671, pl. 26, figs. 1-8. [Appendix to Baker, Fred, The land and fresh-water mollusks of the Stanford Expedition to Brazil. *Proc. Acad. nat. Sci. Phila.*, 65: 618-672, pls. 21-27.]
PILSBRY, Henry A. 1915. *Gundlachia* or *Navicella*? *Nautilus*, 28(11): 132.
PILSBRY, Henry A. 1916a. New species of *Amnicola* from New Mexico and Utah. *Nautilus*, 29(6): 111-112. [Figs. in *Nautilus*, 1917, 30(12): pl. 5, figs. 4, 8, 9.]
PILSBRY, Henry A. 1916b. *Goniobasis* in western Pennsylvania. *Nautilus*, 30(1): 4-5.
PILSBRY, Henry A. 1916c. New subspecies of *Viviparus* and *Campeloma*. *Nautilus*, 30(4): 41-43.
PILSBRY, Henry A. 1916d. Note on *Valvata micra* Pils. and Ferr. *Nautilus*, 30(7): 83-84.
PILSBRY, Henry A. 1916e. *Planorbis dilatatus* and *P. sampsoni*. *Nautilus*, 30(8): 95-96.
PILSBRY, Henry A. 1917a. Rafinesque's genera of fresh-water snails. *Nautilus*, 30(10): 109-114.
PILSBRY, Henry A. 1917b. A new *Hemphillia* and other snails from near Mt. Hood, Oregon. *Nautilus*, 30(10): 117-119, pl. 4, fig. 9.
PILSBRY, Henry A. 1917c. Amnicolidae from Oneida Lake, N.Y. *Nautilus*, 31(2): 44-46. [Reprinted in *Univ. Tech. Publ. 9, N.Y. State Coll. For.*, 1918, 18(2): 244-246.]
PILSBRY, Henry A. 1918. *Viviparus contectoides limi*. *Nautilus*, 32(2): 71.
PILSBRY, Henry A. 1923. Mollusca. *In:* Bishop, S.C. & Clarke, N.T., A scientific survey of Turner's Lake, Isle-au-Haut, Maine. Published privately. pp. 15-16.
PILSBRY, Henry A. 1925a. The family Lancidae distinguished from the Ancylidae. *Nautilus*, 38(3): 73-75.

PILSBRY, Henry A. 1925b [1926]. A fresh-water snail, *Physa zionis*, living under unusual conditions. *Proc. Acad. nat. Sci. Phila.*, 77: 325-328, pl. 11.
PILSBRY, Henry A. 1926. What constitutes publication? A question concerning nomenclature. *Science*, 64(1654): 248-249.
PILSBRY, Henry A. 1927a. A new subspecies of *Planorbis*. *Nautilus*, 40(3): 79-80.
PILSBRY, Henry A. 1927b. Genetic experiments with pond snails *Lymnaea* and *Physa*. [Review.] *Nautilus*, 40(4): 141.
PILSBRY, Henry A. 1927c. On *Pomacea* Perry (Ampullariidae). *Nautilus*, 41(2): 63-64.
PILSBRY, Henry A. 1931. A new race of *Neritina reclivata* Say. *Nautilus*, 45(2): 67-68, pl. 3, fig. 3.
PILSBRY, Henry A. 1932a. *Physa humerosa interioris* Ferriss, 1920. *Nautilus*, 45(4): 139, pl. 11, fig. 12.
PILSBRY, Henry A. 1932b. *Carinifex newberryi subrotunda*, n. subsp. *Nautilus*, 45(4): 139.
PILSBRY, Henry A. 1932c. *Bithynia tentaculata* (L.) in the Potomac. *Nautilus*, 46(2): 69.
PILSBRY, Henry A. 1933. Amnicolidae from Wyoming and Oregon. *Nautilus*, 47(1): 9-12, pl. 2, figs. 1-10.
PILSBRY, Henry A. 1934a. Review of the Planorbidae of Florida, with notes on other members of the family. *Proc. Acad. nat. Sci. Phila.*, 86: 29-66, pls. 7-11.
PILSBRY, Henry A. 1934b [1935]. Mollusks of the fresh-water Pliocene beds of the Kettleman Hills and neighboring oil fields, California. *Proc. Acad. nat. Sci. Phila.*, 86: 541-570, pls. 18-23.
PILSBRY, Henry A. 1935a. Western and southwestern Amnicolidae and a new *Humboldtiana*. *Nautilus*, 48(3): 91-94.
PILSBRY, Henry A. 1935b. *Lioplax subcarinata occidentalis*, new subspecies. *Nautilus*, 48(4): 143-144.
PILSBRY, Henry A. 1935c. Report on shells collected by E.B. Howard from lake bed southwest of Clovis, Roosevelt Co., New Mexico. *In:* Howard, Edgar Billings, Evidence of early man in North America. *Mus. J.* [Univ. Pa. Mus.], 24: 89-90.
PILSBRY, Henry A. 1937a. A Californian *Pomatiopsis*. *Nautilus*, 50(3): 84-85.
PILSBRY, Henry A. 1937b. The Japanese snail. *Aquarium*, Philadelphia, 7: 8.
PILSBRY, Henry A. 1943. The type of *Euamnicola* Crosse and Fischer. *Nautilus*, 57(2): 68-69.
PILSBRY, Henry A. 1944. Snails from Taylor County, Texas. *Nautilus*, 58(2): 69.
PILSBRY, Henry A. 1950a. New fountain snails from Florida. *Nautilus*, 64(2): 37-39, pl. 3, figs. 1-7.
PILSBRY, Henry A. 1950b. *Helisoma anceps* (Menke), 1830. *Nautilus*, 64(2): 68, pl. 4, fig. 44.
PILSBRY, Henry A. 1951. New species of *Amnicola* and *Streptostyla*. *Nautilus*, 64(4): 119-120, pl. 9, figs. 1-3.
PILSBRY, Henry A. 1952a. *Littoridina tenuipes* (Couper). *Nautilus*, 66(2): 50-54.
PILSBRY, Henry A. 1952b. Les types de la collection Lamarck au Muséum de Genève, III, by G. Mermod. [Review.] *Nautilus*, 66(2): 72.
PILSBRY, Henry A. 1953. The case of *Paludina multilineata* Say. *Nautilus*, 67(2): 58-61.
PILSBRY, Henry A. & BAKER, Frank Collins. 1927. Description of a new species of *Somatogyrus* from Wisconsin. *Nautilus*, 41(1): 24-27.
PILSBRY, Henry A. & BEQUAERT, Joseph C. 1927. The aquatic mollusks of the Belgian Congo, with a geographical and ecological account of Congo malacology. *Bull. Am. Mus. nat. Hist.*, 53(2): 69-602, maps 1-15, pls. 10-77.
PILSBRY, Henry A. & COCKERELL, T.D.A. 1900. Records of Mollusca from New Mexico. *Nautilus*, 14(8): 85-86.
PILSBRY, Henry A. & FERRISS, James H. 1906. Mollusca of the southwestern states. II. *Proc. Acad. nat. Sci. Phila.*, 58: 123-175, pls. 5-9.
PILSBRY, Henry A. & FERRISS, James H. 1907. Mollusca of the Ozarkian fauna. *Proc. Acad. nat. Sci. Phila.*, 1906, 58: 529-567, pls. 20-22.
PILSBRY, Henry A. & FERRISS, James H. 1909a. Mollusks from around Albuquerque, New Mexico. *Nautilus*, 22(10): 103-104.
PILSBRY, Henry A. & FERRISS, James H. 1909b [1910a]. Mollusca of the southwestern states. III. The Huachuca Mountains, Arizona. *Proc. Acad. nat. Sci. Phila.*, 61: 495-516, pls. 19-21.
PILSBRY, Henry A. & FERRISS, James H. 1910b. Mollusca of the southwestern states. IV. The Chiricahua Mountains, Arizona. *Proc. Acad. nat. Sci. Phila.*, 62: 44-147, pls. 1-14.
PILSBRY, Henry A. & FERRISS, James H. 1911. Mollusca of the southwestern states. V. The Grand Canyon and northern Arizona. *Proc. Acad. nat. Sci. Phila.*, 63: 174-199, pls. 12-14.
PILSBRY, Henry A. & FERRISS, James H. 1915. Mollusca of the southwestern states, VII: the Dragoon, Mule, Santa Rita, Baboquivari, and Tucson ranges, Arizona. *Proc. Acad. nat. Sci. Phila.*, 67: 363-418, pls. 8-15.
PILSBRY, Henry A. & FERRISS, James H. 1917. Mollusca of the southwestern states. VIII. The Black Range, New Mexico. *Proc. Acad. nat. Sci. Phila.*, 69: 83-107, pls. 7-10.

PILSBRY, Henry A. & FERRISS, James H. 1918 [1919]. Mollusca of the southwestern states. IX. The Santa Catalina, Rincon, Tortillita and Galiuro mountains. X. The mountains of the Gila headwaters. *Proc. Acad. nat. Sci. Phila.*, 70: 282-333, pls. 3-7.

PILSBRY, Henry A. & FERRISS, James H. 1924. List of species. *In:* Ferriss, James H., On the Rio Grande. *Nautilus*, 38(2): 37-43 [pp. 40-43].

PILSBRY, Henry A. & JOHNSON, Charles W. 1912. A new Floridian *Viviparus*. *Nautilus*, 26(2): pl. 3, figs. 6, 7; 26(4): 48.

PILSBRY, Henry A. & OLSSON, Axel A. 1951. The Lepyriidae, a new family of fresh-water snails (Gastropoda: Rissoacea). *Not. nat.*, Acad. nat. Sci. Phila., (233): 1-5.

PILSBRY, Henry A. & RHOADS, Samuel N. 1896. Contributions to the zoology of Tennessee. No. 4, mollusks. *Proc. Acad. nat. Sci. Phila.*, 48: 487-506.

PIP, Eva. 1975. Scalariformy in the pond snail, *Lymnaea stagnalis*. *Nautilus*, 89(2): 36-37.

PIP, Eva. 1977. Frequent scalariformy in a population of the pond snail *Lymnaea stagnalis*. *Nautilus*, 91(2): 52-54.

PIP, Eva. 1980. *Somatogyrus* (Gastropoda: Hydrobiidae) in Lake Winnipeg, Canada. *Nautilus*, 94(1): 4-5.

POLLISTER, A.W. & POLLISTER, P.F. 1940. Distribution of males in the genus *Campeloma*, with a note on the chromosome numbers in Viviparidae. *Anat. Rec.*, 78(suppl.): 128.

POLLISTER, A.W. & POLLISTER, P.F. 1943. The relation between centriole and centromere in atypical spermatogenesis of viviparid snails. *Ann. N.Y. Acad. Sci.*, 45: 1-48.

POTIEZ, V.L. & MICHAUD, A.L. 1838-44. *Galerie des mollusques, ou catalogue méthodique, descriptif et raisonné des mollusques et coquilles du Muséum de Douai.* J.-B. Baillière, Paris. Vol. 1, 1838, xxxvi, 562 pp.; Vol. 2, 1844, xxxvi-xliv, 511 pp.; atlas, pp. 1-56, pls. 1-70.

PRATT, William L. 1977. Hydrobiid snails of the Moapa Warm Spring complex, Nevada. *Annu. Rep. west. Soc. Malacol.*, 10: 7.

RACKETT, Thomas. 1821. Description of some shells found in Canada. *Trans. Linn. Soc. Lond.*, 13(1): 42-43, pl. 5, figs. 1, 2.

RADOMAN, Pavle. 1966. Die Gattungen *Pseudamnicola* und *Horatia*. *Arch. Molluskenkd.*, 95(5/6): 243-253.

RAFINESQUE, C.S. 1815. *Analyse de la nature, ou tableau de l'univers et des corps organises*. Palerme. 224 pp. [pp. 136-149].

RAFINESQUE, C.S. 1818a. Discoveries in natural history, made during a journey through the western region of the United States. *Am. mon. Mag. crit. Rev.*, 3: 354-355.

RAFINESQUE, C.S. 1818b. Farther account of discoveries in natural history, in the western states. *Am. mon. Mag. crit. Rev.*, 4: 39-42.

RAFINESQUE, C.S. 1818c. General account of the discoveries made in the zoology of the western states. *Am. mon. Mag. crit. Rev.*, 4: 106-107.

RAFINESQUE, C.S. 1819. De 70 nouveaux genres d'animaux découverts dans l'intérieur des Etats-Unis d'Amérique, durant l'année 1818. *J. Phys. Chim. Hist. nat.*, 88: 423-428.

RAFINESQUE, C.S. 1820. VIII. Clas. Apalosia.—The Mollusca. *Ann. Nature*, (1): 10-11.

RAFINESQUE, C.S. 1831. *Enumeration and account of some remarkable natural objects in the cabinet of Prof. Rafinesque, in Philadelphia.* C.S. Rafinesque, Philadelphia. pp. 1-4.

RAFINESQUE, C.S. 1864. *In:* Binney, Wm. G. & Tryon, George W., Jr., *The complete writings of Constantine Smaltz Rafinesque, on Recent and fossil conchology.* Bailliere Bros., New York. 96 pp., index pp. 1-7, pls. 80-82.

REDFIELD, John H. 1854. Descriptions of new species of shells. *Ann. Lyc. nat. Hist. N.Y.*, 6: 130-132, pl. 1.

REHDER, Harald A. 1942. A note on the genus *Anaplocamus* Dall. *Nautilus*, 56(2): 49-50.

RICHARDS, Charles S. 1962. Retarded development of the male reproductive system in a Florida *Gyraulus*. *Trans. Am. microsc. Soc.*, 81(4): 347-351.

RICHARDS, Horace G. 1929. Freshwater snails in brackish water. *Nautilus*, 42(4): 129-130.

RICHARDS, Horace G. & ADAMS, Joseph W. 1929. *Viviparus malleatus* in Philadelphia. *Nautilus*, 42(4): 142-143.

ROBERTSON, Imogene C.S. 1945. *Valvata piscinalis* (Müller) in Lake Erie. *Nautilus*, 59(1): 36.

ROBERTSON, Imogene C.S. & BLAKESLEE, Clifford L. 1948. The Mollusca of the Niagara Frontier region. *Bull. Buffalo Soc. nat. Sci.*, 19(3): i-xi, 1-191, pls. 1-14, 1 map.

ROBINS, Catherine H. 1970. The introduced freshwater snail, *Marisa*. *Annu. Rep. Am. malacol. Union 1970*, (37): 3.

ROSCOE, Ernest J. 1962. Additional Wisconsin records of *Viviparus contectoides*. *Nautilus*, 76(1): 36.

ROSCOE, Ernest J. 1963. Some goniobases in western United States. *Nautilus*, 77(2): 43-47, pl. 4, figs. 5-8.

ROSEWATER, Joseph. 1959. A pleurocerid index to past collecting in the Ohio and Tennessee river basins with some thoughts for the future. *Annu. Rep. Am. malacol. Union 1958*, (25): 23-24.

ROSEWATER, Joseph. 1960. Egg mass and gross embryology of *Pleurocera canaliculatum* (Say, 1821). *Annu. Rep. Am. malacol. Union 1959,* (26): 10-12.

ROSEWATER, Joseph. 1961. Preliminary observations on the nervous systems of some Pleuroceridae. *Annu. Rep. Am. malacol. Union 1961,* (28): 10-11.

ROSEWATER, Joseph. 1970. Another record of insect dispersal of an ancylid snail. *Nautilus,* 83(4): 144-145.

ROSSMÄSSLER, E.A. 1835. *Iconographie der Land- und Süsswasser-Mollusken, mit vorzüglicher Berücksichtigung der europäischen noch nicht abgebildeten Arten.* Arnoldische Buchhandung, Dresden und Leipzig. No. 1, pp. 1-132, 4 pls.

ROWAN, W.B. 1966. Autumn migration of *Helisoma trivolvis* in Montana. *Nautilus,* 79(3): 108, iii.

ROY, Edward C. 1963. Checklist of Pleistocene and living Mollusca of Wisconsin. *Sterkiana,* (10): 5-21.

RUDOLPH, Paul H. 1979. The strategy of copulation of *Stagnicola elodes* (Say) (Basommatophora: Lymnaeidae). *Malacologia,* 18: 381-389.

RUSSELL, Richard H. 1966. Lymnaeidae of western Montana. *Annu. Rep. Am. malacol. Union 1966,* (33): 26.

RUSSELL, Richard H. 1967. A new subspecies of *Lymnaea stagnalis* from Montana. *Nautilus,* 80(4): 125-126, pl. 9.

RUSSELL, Richard H. 1971a. The appearance of *Pseudosuccinea columella* (Say) in Arizona. *Nautilus,* 85(2): 71.

RUSSELL, Richard H. 1971b. Mollusca of Fish Springs, Juab County, Utah: rediscovery of *Stagnicola pilsbryi* (Hemphill, 1890). *Great Basin Nat.,* 31(4): 223-236.

RUSSELL, Richard H. 1972. The type locality of *Stagnicola montanensis* (Baker) 1913. *Nautilus,* 85(4): 145.

RUSSELL, Richard H. & BRUNSON, Royal Bruce. 1967a. A check-list of mollusks of Glacier National Park, Montana. *Sterkiana,* (26): 1-5.

RUSSELL, Richard H. & BRUNSON, Royal Bruce. 1967b. *Acroloxus coloradensis* from Montana. *Nautilus,* 81(1): 33.

SARGENT, H.E. 1894a. *Tulotoma* in the Tennessee drainage. *Nautilus,* 7(11): 130.

SARGENT, H.E. 1894b. *Amnicola olivacea* Pils. *Nautilus,* 8(7): 95-96.

SARGENT, H.E. 1895. *Ferussacia subcylindrica* and two new species in Jackson Co., Alabama. *Nautilus,* 8(9): 104-105.

SARGENT, H.E. 1896. Annotated list of the Mollusca found in the vicinity of Clearwater, Wright Co., Minnesota. *Nautilus,* 9(11): 125-128.

SAVAGE, Alice E. 1938. A comparison of the nervous system in normal and sinistral snails of the species *Campeloma rufum. Am. Nat.,* 72: 160-169.

SAY, Thomas. 1817a. Conchology. 15 pp., 4 pls. *In:* Nicholson, William, 1816-17, *American edition of the British encyclopedia, or dictionary of arts and sciences comprising an accurate and popular view of the present improved state of human knowledge.* First edition. Vol. 1, 1816; vols. 2-6, 1817 [vol. 2, B.....E]; no pagination. Samuel A. Mitchell and Horace Ames, Philadelphia.

SAY, Thomas. 1817b. Description of seven species of American fresh water and land snails, not noticed in the systems. *J. Acad. nat. Sci. Phila.,* 1(1): 13-16; 1(2): 17-18.

SAY, Thomas. 1817c. Descriptions of new species of land and fresh water shells of the United States. *J. Acad. nat. Sci. Phila.,* 1(1): 123-126.

SAY, Thomas. 1818a. Account of two new genera, and several new species, of fresh water and land shells. *J. Acad. nat. Sci. Phila.,* 1(2): 276-284.

SAY, Thomas. 1818b. Conchology. 15 pp., 4 pls. *In:* Nicholson, William, 1818, *American edition of the British encyclopedia, or dictionary of arts and sciences comprising an accurate and popular view of the present improved state of human knowledge.* Second edition. 12 vols. [vol. 4], no pagination. Mitchell, Ames and White, Philadelphia.

SAY, Thomas. 1819a. Conchology. 20 pp., 4 pls. *In:* Nicholson, William, 1819-21, *American edition of the British encyclopedia, or dictionary of arts and sciences comprising an accurate and popular view of the present improved state of human knowledge.* Third edition. Vols. 1-6, 1819 [vol. 4]; vols. 7-12, 1821; no pagination. Mitchell, Ames and White, Philadelphia.

SAY, Thomas. 1819b. Description of the land and fresh water shells of the United States. 14 pp. (unnumbered), 4 pls. [Reprint, with title page added and with introductory remarks deleted, of Say (1819a).]

SAY, Thomas. 1821. Descriptions of univalve shells of the United States. *J. Acad. nat. Sci. Phila.,* 2(1): 149-179.

SAY, Thomas. 1822a. An account of some of the marine shells of the United States. *J. Acad. nat. Sci. Phila.,* 2(2): 257-276.

SAY, Thomas. 1822b. Description of univalve terrestrial and fluviatile shells of the United States. *J. Acad. nat. Sci. Phila.,* 2(2): 370-381.

SAY, Thomas. 1824. *In: Narrative of an expedition to the source of the St. Peter's River, ... , under the command of Major Stephen H. Long.* Philadelphia. Vol. 2, appendix, pp. 256-265, pls. 14, 15.

SAY, Thomas. 1825. Descriptions of some new species of fresh water and land shells of the United States. *J. Acad. nat. Sci. Phila.*, 5(2): 119-131.

SAY, Thomas. 1829-31. Descriptions of some new terrestrial and fluviatile shells of North America. *New Harmony Dissem. useful Knowl.*, 2(15): 229-230 (1829); 2(16): 244-246 (1829); 2(17): 259-261 (1829); 2(19): 291-293 (1829); in newspaper form, Jan. 15, 1831, p. 26 (reprint) (1831).

SAY, Thomas. 1830-34. *American conchology, or descriptions of the shells of North America, illustrated by colored figures.* New Harmony, Indiana. 68 colored plates with legends. [Pt. 1, pls. 1-10, 1830; pt. 2, pls. 11-20, 1831; pt. 3, pls. 21-30, 1831; pt. 4, pls. 31-40, 1832; pt. 5, pls. 41-50, 1832; pt. 6, pls. 51-60, 1834; pt. 7, pls. 61-68, 1834.]

SAY, Thomas. 1840. *Descriptions of some new terrestrial and fluviatile shells of North America. 1829, 1830, 1831.* Lucy W. Say, New Harmony, Indiana. 26 pp.

SAY, Thomas. 1858. *In:* Binney, W.G., *The complete writings of Thomas Say, on the conchology of the United States.* H. Bailliere, New York. vi, 252 pp., pls. 1-75.

SCHMECK, Eugene H. 1942. *Viviparus malleatus* in Niagara River. *Nautilus*, 55(3): 102-103.

SHIMEK, B. 1890. A new species of fresh water mollusk. *Bull. Lab. nat. Hist. state Univ. Iowa*, 1(3/4): 214-215, pl. 3, figs. 5, 5a, 5b, 5c.

SHIMEK, B. 1935. *Ferrissia* in the lake region of Iowa. *Nautilus*, 49(2): 44-46.

SHOUP, Charles S. 1974. *A bibliography of the zoology of Tennessee and the Tennessee Valley region.* Tech. info. Center, Off. info. Serv., U.S. atomic Energy Comm., Oak Ridge, Tennessee. iv, 251 pp.

SHUTTLEWORTH, R.J. 1845. Über *Gyrotoma*, eine neue Gattung der Melaniana, Gasteropoda Pectinibranchiata. *Mitt. naturforsch. Ges. Bern*, (50): 85-88.

SIMPSON, Chas. T. 1888. *Gundlachia ancyliformis* Pfr. in Florida. *Conchol. Exc.*, 2(7): 96.

SIMPSON, Chas. T. 1895. *Pleurocera subulare* in water-mains. *Nautilus*, 9(4): 37-38.

SINCLAIR, Ralph M. 1969. The pleurocerid fauna of the Tennessee River Gastropoda: Prosobranchia. *Annu. Rep. Am. malacol. Union 1969*, (36): 45-47.

SINGLEY, J.A. 1889. Shell collecting in southern Texas. *Nautilus*, 3(5): 60.

SMALL, H.B. & SYMES, P.B. 1882. Report of the Conchological Branch for the season of 1882. *Trans. Ottawa field-nat. Club*, (3): 57-59. [Reprinted in *Sterkiana*, (8): 29-31.]

SMITH, Douglas G. 1975. The identity of *Planorbula jenksii* (H.F. Carpenter). *Nautilus*, 89(3): 96-98.

SMITH, Douglas G. 1976. Abnormal shells of *Gyraulus parvus* (Planorbidae). *Nautilus*, 90(4): 150.

SMITH, Douglas G. 1980. *Goniobasis virginica* (Gastropoda: Pleuroceridae) in the Connecticut River. *Nautilus*, 94(2): 50-54.

SMITH, Edgar. 1876a. Description of a new species of *Carinifex* from California. *Proc. sci. Meetings zool. Soc. Lond. 1875*, 43(34): 536-537, fig. on p. 539 [mislabelled *Diala leithii*; figs. on pp. 537 and 539 were mistakenly transposed].

SMITH, Edgar A. 1876b. Description of a new species of *Carinifex* from California. *Q. J. Conchol.*, 1: 150-151.

SMITH, Sanderson & PRIME, Temple. 1870. Report on the Mollusca of Long Island, N.Y., and of its dependencies. *Ann. Lyc. nat. Hist. N.Y.*, 9: 377-407.

SOGANDARES-BERNAL, Franklin & ABDEL-MALEK, Emile. 1961. *Pomatiopsis lapidaria* Say (Gastropoda: Amnicolidae) in Louisiana. *J. Parasitol.*, 47(5): 832.

SOLEM, Alan. 1961. Hydrobiid snails from Lake Pontchartrain, Louisiana. *Nautilus*, 74(4): 157-160.

SOWERBY, G.B. 1872. Monograph of the genus *Limnaea. In:* Reeve, Lovell, 1843-78, *Conchologia iconica.* Vol. 18, 32 pp., pls. 1-15. Reeve, Benham & Reeve, London.

SOWERBY, G.B. 1873-74. Monograph of the genus *Physa. In:* Reeve, Lovell, 1843-78, *Conchologica iconica.* Vol. 19, 27 pp., pls. 1-12. Reeve, Benham & Reeve, London.

SOWERBY, G.B. 1877a. Monograph of the genus *Ancylus. In:* Reeve, Lovell, 1843-78, *Conchologica iconica.* Vol. 20, 7 pp., pls. 1-3. Reeve, Benham & Reeve, London.

SOWERBY, G.B. 1877b. Monograph of the genus *Planorbis. In:* Reeve, Lovell, 1843-78, *Conchologica iconica.* Vol. 20, 31 pp., pls. 1-14. Reeve, Benham & Reeve, London.

SPOEL, S. van der. 1959. Age-determination by winter rings. *Basteria*, 23(3): 36-38.

STANSBERY, David H. 1970. Eastern freshwater mollusks (I). The Mississippi and St. Lawrence river systems. [Am. malacol. Union Symp. Rare and Endangered Mollusks.] *Malacologia*, 10(1): 9-22.

STANSBERY, David H. 1971. Rare and endangered freshwater mollusks in eastern United States. *In:* Jorgensen, S.E. & Sharp, R.W., *Proc. Symp. rare & endangered Moll. (naiads) of the U.S.* Bur. sport Fish. Wildl., Fish and Wildl. Serv., U.S. Dept. Int., Reg. 3, Twin Cities, Minnesota. 8, 79 pp. [pp. 5-18, 6 pls.].

STANSBERY, David H. 1972. The mollusk fauna of the North Fork Holston River at Saltville, Virginia. *Bull. Am. malacol. Union Inc. 1971*, (38): 45-46.

STANSBERY, David H. & CLENCH, William J. 1974. The Pleuroceridae and Unionidae of the North Fork Holston River above Saltville, Virginia. *Bull. Am. malacol. Union Inc. 1973*, (40): 33-36.
STANSBERY, David H. & CLENCH, William J. 1975. The Pleuroceridae and Unionidae of the Middle Fork Holston River in Virginia. *Bull. Am. malacol. Union Inc. 1974*, (41): 51-54.
STANSBERY, David H. & CLENCH, William J. 1978. The Pleuroceridae and Unionidae of the upper South Fork Holston River in Virginia. *Bull. Am. malacol. Union Inc. 1977*, (44): 75-78.
STANSBERY, David H. & STEIN, Carol B. 1976. Changes in the distribution of *Io fluvialis* (Say, 1825) in the upper Tennessee River system (Mollusca, Gastropoda, Pleuroceridae). *Bull. Am. malacol. Union Inc. 1976*, (43): 28-33.
STAROBOGATOV, Y.I. 1970. *Mollusk fauna and the zoogeographic partitioning of continental waterbodies of the globe.* [In Russian.] Nauka, Leningrad Br., Acad. Sci. U.S.S.R., Leningrad. 372 pp.
STEARNS, Robert E.C. 1881. Observations on *Planorbis. Proc. Acad. nat. Sci. Phila.*, 33: 92-110.
STEARNS, Robert E.C. 1883a. On the shells of the Colorado Desert and the region farther east. *Am. Nat.*, 17(2): 1014-1020.
STEARNS, Robert E.C. 1883b. Description of a new hydrobiinoid gasteropod from the mountain lakes of the Sierra Nevada, with remarks on allied species and the physiographical features of said region. *Proc. Acad. nat. Sci. Phila.*, 35: 171-176.
STEARNS, Robert E.C. 1889. Notes on *Physa triticea* of Lea; its relations and comments on the variation, etc., of physae. *Nautilus*, 3(5): 49-51.
STEARNS, Robert E.C. 1890. Descriptions of new West American land, fresh-water, and marine shells, with notes and comments. *In:* Scientific results of explorations by the U.S. Fish Commissions steamer Albatross. *Proc. U.S. natl. Mus.*, 13(813): 205-225, pls. 15-17.
STEARNS, Robert E.C. 1891. Notes on the sculpture of American limnaeas, etc. *Nautilus*, 4(11): 121-124.
STEARNS, Robert E.C. 1893. Report on the land and fresh-water shells collected in California and Nevada by the Death Valley Expedition, including a few additional species obtained by Dr. C. Hart Merriam and assistants in parts of the southwestern United States. *In: North American fauna.* No. 7, The Death Valley Expedition, a biological survey of parts of California, Nevada, Arizona and Utah. Div. of Ornithol. and Mammal., U.S. Dept. Agric. U.S. Gov. Print. Off., Washington, D.C. 402 pp., 13 pls., 5 maps [pp. 269-283].
STEARNS, Robert E.C. 1900. List of shells collected by Vernon Bailey in Heron and Eagle lakes, Minnesota, with notes. *Proc. U.S. natl. Mus.*, 22(1190): 135-138.
STEARNS, Robert E.C. 1901a. Japanese *Vivipara* in California. *Nautilus*, 15(8): 91.
STEARNS, Robert E.C. 1901b. The fossil fresh-water shells of the Colorado Desert, their distribution, environment, and variation. *Proc. U.S. natl. Mus.*, 24: 271-299, pls. 19-24.
STEPHENS, Kate. 1934. [Introduced "*Viviparus*", i.e., *Cipangopaludina*.] *Rep. Am. malacol. Union 1934*, (3): 5.
STERKI, V. 1891a. Food of Limnaeidae. *Nautilus*, 5(8): 94.
STERKI, V. 1891b. [Note on way *Planorbis trivolvis* Say and other related forms move on or in sand.] *Nautilus*, 5(8): 94-95.
STERKI, V. 1891c. [Observations on how snails move their odontophores.] *Nautilus*, 5(8): 95.
STERKI, V. 1907a. A few notes on Say's early writings and species. *Nautilus*, 21(3): 31-34.
STERKI, V. 1907b. A preliminary catalogue of the land and freshwater molluscs of Ohio. *Proc. Ohio state Acad. Sci.*, spec. pap. 12, 4(8): 367-402.
STIMPSON, William. 1851. *Shells of New England.* A revision of the synonymy of the testaceous mollusks of New England, with notes on their structure, and their geographical and bathymetrical distribution. Phillips, Sampson and Co., Boston. 58 pp., pls. 1, 2.
STIMPSON, William. 1863. Malacozoölogical notices, No. 1. *Proc. Boston Soc. nat. Hist.*, 9: 249-252.
STIMPSON, William. 1865a. Diagnoses of newly discovered genera of gasteropods, belonging to the sub-fam. Hydrobiinae, of the family Rissoidae. *Am. J. Conchol.*, 1(1): 52-54, pl. 8, fig. 1.
STIMPSON, William. 1865b. Researches upon the Hydrobiinae and allied forms; chiefly made upon materials in the museum of the Smithsonian Institution. *Smithson. misc. Collect.*, 7(4): i-iii, 1-59.
STIVEN, Alan E. & WALTON, Clarence R. 1967. Age and shell growth in the freshwater snail, *Goniobasis proxima* (Say). *Am. midl. Nat.*, 78(1): 207-214.
STREATOR, Geo. J. 1889. *Bythinia tentaculata*, Linn, in Ohio. *Nautilus*, 3(4): 46.
STRENG, L.H. 1896. A new variety of *Limnaea. Nautilus*, 9(11): 123.
SWAINSON, William. 1840. A treatise on malacology, or shells and shell-fish. *In:* Lardner, Dionysius, *The cabinet cyclopaedia.* Natural history. Longman, Orme, Brown, Green & Longmans, and John Taylor, London. 419 pp.
TAYLOR, Dwight W. 1950. Three new *Pyrgulopsis* from the Colorado Desert, California. *Leafl. Malacol.*, 1(7): 27-

33.
TAYLOR, Dwight W. 1952. Notes on the freshwater mollusks of Yellowstone Park, Wyoming. *Leafl. Malacol.*, 1(9): 43-49, pl. 1.
TAYLOR, Dwight W. 1954. A new Pleistocene fauna and new species of fossil snails from the High Plains. *Occ. Pap. Mus. Zool. Univ. Mich.*, (557): 1-16.
TAYLOR, Dwight W. 1960. Late Cenozoic molluscan faunas from the High Plains. *U.S. geol. Surv. profession. Pap. 337*, pp. i-iv, 1-94, pls. 1-4.
TAYLOR, Dwight W. 1964. Historical analysis of distribution of West American freshwater molluscs. [Abstract.] *Am. Zool.*, 4: 436-437.
TAYLOR, Dwight W. 1965. The study of Pleistocene nonmarine mollusks in North America. *In:* Wright, H.E., Jr. & Frey, David, *The Quaternary of the United States.* Princeton Univ. Press, Princeton, N.J. 922 pp. [pp. 597-611].
TAYLOR, Dwight W. 1966a. Summary of North American Blancan nonmarine mollusks. *Malacologia*, 4(1): 1-172.
TAYLOR, Dwight W. 1966b. A remarkable snail fauna from Coahuila, Mexico. *Veliger*, 9(2): 152-228.
TAYLOR, Dwight W. 1966c. A remarkable snail fauna from Coahuila, Mexico. [Abstract.] *Annu. Rep. Am. malacol. Union 1966*, (32): 70-72.
TAYLOR, Dwight W. 1967a. Freshwater mollusks collected by the United States and Mexican Boundary Surveys. *Veliger*, 10(2): 152-158.
TAYLOR, Dwight W. 1967b. Late Pleistocene nonmarine mollusks from the state of Puebla, Mexico. *Annu. Rep. Am. malacol. Union 1967*, (34): 76-78.
TAYLOR, Dwight W. 1970a. West American freshwater Mollusca, 1: bibliography of Pleistocene and Recent species. *San Diego Soc. nat. Hist. Mem.*, 4: 1-73, frontis.
TAYLOR, Dwight W. 1970b. Western freshwater mollusks [Editor's (Arthur H. Clarke) summary, in Am. malacol. Union symp. Rare and Endangered Mollusks]. *Malacologia*, 10(1): 33.
TAYLOR, Dwight W. 1974. The Tertiary gastropod *Orygoceras* found living. *Arch. Molluskenkd.*, 104(1-3): 93-96 [also published in *Soosiana*, 1974, 2: 37-43].
TAYLOR, Dwight W. 1975. Index and bibliography of Late Cenozoic freshwater Mollusca of western North America. Claude W. Hibbard Mem. Vol. 1. *Pap. Paleontol. Mus. Paleontol. Univ. Mich.*, (10): 1-384.
TAYLOR, Dwight W. & SOHL, N.F. 1962. An outline of gastropod classification. *Malacologia*, 1(1): 7-32.
TAYLOR, Dwight W., WALTER, Harold J. & BURCH, John B. 1963. Freshwater snails of the subgenus *Hinkleyia* (Lymnaeidae: *Stagnicola*) from the western United States. *Malacologia*, 1(2): 237-281.
TAYLOR, George W. 1892. Preliminary check-list of the land and fresh water Mollusca of Canada. *Ottawa Nat.*, 7: 33-37.
TAYLOR, George W. 1894. [Report of the appearance of *Paludina japonica* Mart. in the Chinese Market at Victoria, B.C.] *Nautilus*, 7(12): 144.
TAYLOR, George W. 1897. *Planorbis nautileus* L. in America. *Nautilus*, 10(12): 139-140.
TE, George A. 1974. Current taxonomic studies on the family Physidae. *Malacol. Rev.*, 8(1/2): 120-121.
TE, George A. 1975. Michigan Physidae, with systematic notes on *Physella* and *Physodon* (Basommatophora: Pulmonata). *Malacol. Rev.*, 8(1/2): 7-30.
TE, George A. 1978. *The systematics of the family Physidae (Basommatophora: Pulmonata).* Ph.D. Dissertation, Univ. of Michigan, Ann Arbor. 325 pp.
TE, George A. 1980. New classification system for the family Physidae. *Arch. Molluskenkd.*, 110(4/6): 179-184.
TE, George A. & MARDINLY, A. John. 1974. Scanning electron microscope study of radular teeth of *Physa gyrina hildrethiana* (Basommatophora: Physidae). *Malacol. Rev.*, 8(1/2): 119-120.
THIELE, Johannes. 1928. Revision des Systems der Hydrobiiden und Melaniiden. *Zool. Jahrb.*, 55(5/6): 351-402, pl. 8.
THIELE, Johannes. 1929. *Handbuch der systematischen Weichtierkunde.* Erster Teil, pp. 1-376. Gustav Fischer, Jena.
THOMPSON, Fred G. 1968. *The aquatic snails of the family Hydrobiidae of peninsular Florida.* Univ. Florida Press, Gainesville. ix, 268 pp.
THOMPSON, Fred G. 1969. Some hydrobiid snails from Georgia and Florida. *Q. J. Fla. Acad. Sci.*, 32(4): 241-265.
THOMPSON, Fred G. 1974. Comments on the proposed suppression of *Paludina lustrica* Say, 1821. Z.N.(S.) 730. *Bull. zool. Nomencl.*, 31(4): 170-171.
THOMPSON, Fred G. 1977. The hydrobiid snail genus *Marstonia. Bull. Fla. State Mus.* (biol. sci.), 21(3): 113-158.
THOMPSON, Fred G. 1979. The systematic relationships of the hydrobioid snail genus *Nymphophilus* Taylor 1966 and the status of the subfamily Nymphophilinae. *Malacol. Rev.*, 12(1/2): 41-49.
THOMPSON, Fred G. & McCALEB, John E. 1978. A new freshwater snail from a spring in eastern Alabama. *Am.*

midl. Nat., 100(2): 350-358.

TROSCHEL, F.H. 1856-93. *Das Gebiss der Schnecken zur Begründung einer natürlichen Classification.* Vol. 1, viii, 252 pp., pls. 1-20; Vol. 2, ix, 409 pp., pls. 1-32. Nicolaische Verlagsbuchhandl., Berlin. [Vol. 1, pt. 1, pp. 1-72, pls. 1-4, 1856; Vol. 1, pt. 2, pp. 73-112, pls. 5-8, 1857; Vol. 1, pt. 3, pp. 113-152, pls. 9-12, 1858; Vol. 1, pt. 4, pp. 153-196, pls. 13-16, 1861; Vol. 1, pt. 5, pp. 197-252, pls. 17-20, 1863; Vol. 2, pt. 1, pp. 1-48, pls. 1-4, 1865; Vol. 2, pt. 2, pp. 49-96, pls. 5-8, 1867; Vol. 2, pt. 3, pp. 97-132, pls. 9-12, 1869; Vol. 2, pt. 4, pp. 133-180, pls. 13-16, 1875; Vol. 2, pt. 5, pp. 181-216, pls. 17-20, 1878; Vol. 2, pt. 6, pp. 217-246, pls. 21-24, 1879; (Vol. 2, pt. 7, pp. 249-334, pls. 25-28, 1891, and Vol. 2, pt. 8, pp. 337-409, pls. 29-32, 1893, by Johannes Thiele).]

TRYON, George W., Jr. 1861a. Publications of Isaac Lea on Recent conchology. *In:* Tryon, George W., Jr., *A list of American writers on Recent conchology.* Bryson's, Philadelphia. [Reprint page nos. 1-13.]

TRYON, George W., Jr. 1861b. On the Mollusca of Harper's Ferry, Virginia. *Proc. Acad. nat. Sci. Phila.*, 13: 396-399.

TRYON, George W., Jr. 1862a. History of conchology in the United States. *Am. J. Sci. Arts*, 33: 13-32 [reprint pagination].

TRYON, George W., Jr. 1862b. Notes on American fresh water shells, with descriptions of two new species. *Proc. Acad. nat. Sci. Phila.*, 14: 451-452.

TRYON, George W., Jr. 1863a. Descriptions of new species of fresh water Mollusca, belonging to the families Amnicolidae, Valvatidae and Limnaeidae; inhabiting California. *Proc. Acad. nat. Sci. Phila.*, 15: 147-150, pl. 1, figs. 8-15.

TRYON, George W., Jr. 1863b. Description of a new species of *Pleurocera. Proc. Acad. nat. Sci. Phila.*, 15: 279-280, pl. 2, fig. 6.

TRYON, George W., Jr. 1863-65. Synonymy of the species of Strepomatidae, a family of fluviatile Mollusca, inhabiting North America. *Proc. Acad. nat. Sci. Phila.*, 15: 306-321 (pt. 1, 1863); 16: 24-48 (pt. 2, 1864); 16: 92-104 (pt. 3, 1864); 17: 19-36 (pt. 4, 1865).

TRYON, George W., Jr. 1865a. Description of new species of Strepomatidae. *Am. J. Conchol.*, 1(1): 38, pl. 1, figs. 8, 9.

TRYON, George W., Jr. 1865b. Observations on the genus *Io. Am. J. Conchol.*, 1(1): 41-44, pls. 3, 4.

TRYON, George W., Jr. 1865c. Catalogue of Mollusca, collected by Prof. D.S. Sheldon, at Davenport, Iowa. *Am. J. Conchol.*, 1(1): 68-70.

TRYON, George W., Jr. 1865d. On the structural characters of the so-called melanians of North America. [Review.] *Am. J. Conchol.*, 1(1): 78-79.

TRYON, George W., Jr. 1865e. Complete writings of Constantine Smaltz Rafinesque on Recent and fossil conchology. [Review.] *Am. J. Conchol.*, 1(1): 79-84, 1 table.

TRYON, George W., Jr. 1865f. Observations on the family Strepomatidae. *Am. J. Conchol.*, 1(2): 97-135, pl. 17.

TRYON, George W., Jr. 1865g. Catalogue of the species of *Physa*, inhabiting the United States. *Am. J. Conchol.*, 1(2): 165-173.

TRYON, George W., Jr. 1865h. Descriptions of new species of *Melania. Am. J. Conchol.*, 1(3): 216-218, pl. 22, figs. 1-4.

TRYON, George W., Jr. 1865i. Descriptions of new species of *Amnicola, Pomatiopsis, Somatogyrus, Gabbia, Hydrobia* and *Rissoa. Am. J. Conchol.*, 1(3): 219-222, pl. 22, figs. 5-13.

TRYON, George W., Jr. 1865j. Descriptions of new species of North American Limnaeidae. *Am. J. Conchol.*, 1(3): 223-231, pl. 22, figs. 14-17, pl. 23.

TRYON, George W., Jr. 1865k. Review of the goniobases of Oregon and California. *Am. J. Conchol.*, 1(3): 236-246, pl. 24.

TRYON, George W., Jr. 1865l. Catalogue of the species of *Limnaea* inhabiting the United States. *Am. J. Conchol.*, 1(3): 247-258.

TRYON, George W., Jr. 1865m. Synonymy of the species of Strepomatidae, a family of fluviatile Mollusca inhabiting North America. Part 4. [Review.] *Am. J. Conchol.*, 1(3): 271-272.

TRYON, George W., Jr. 1865n. *Synonymy of the species of Strepomatidae (melanians) of the United States; with critical observations on their affinities, and descriptions of land, fresh water and marine Mollusca.* Contrib. to conchol., vol. 3. Bailliere Bros., New York, etc. 100 pp., pls. 1, 2.

TRYON, George W., Jr. 1865-66. Monograph of the family Strepomatidae. *Am. J. Conchol.*, 1(4): 299-341 (1865); 2(1): 14-52 (1866); 2(2): 115-133 (1866).

TRYON, George W., Jr. 1866a. On an abnormal specimen of *Planorbis bicarinatus. Am. J. Conchol.*, 2(1): 3, pl. 1, fig. 4.

TRYON, George W., Jr. 1866b. Descriptions of new fresh-water shells of the United States. *Am. J. Conchol.*, 2(1):

4-7, pl. 2, figs. 1-6.
TRYON, George W., Jr. 1866c. Description of a new species of *Vivipara*. *Am. J. Conchol.*, 2(2): 108-110, pl. 10, fig. 2.
TRYON, George W., Jr. 1866d. Descriptions of new fluviatile Mollusca. *Am. J. Conchol.*, 2(2): 111-113, pl. 10, figs. 3-8.
TRYON, George W., Jr. 1866e. Observations on an abnormal specimen of *Physa gyrina*. *Am. J. Conchol.*, 2(2): 114.
TRYON, George W., Jr. 1866f. Note on the lingual dentition of the Strepomatidae. *Am. J. Conchol.*, 2(2): 134-135, pls. 11, 12.
TRYON, George W., Jr. 1866g. Researches upon the Hydrobiinae and allied forms. [Review.] *Am. J. Conchol.*, 2(2): 152-158.
TRYON, George W., Jr. 1867. Land and fresh water shells of North America, parts 2 and 3. [Review.] *Am. J. Conchol.*, 3(2): 195-198.
TRYON, George W., Jr. 1868. Notes on Mollusca collected by Dr. F.V. Hayden, in Nebraska. *Am. J. Conchol.*, 4(3): 150-151.
TRYON, George W., Jr. 1870-71. *A monograph of the fresh-water univalve Mollusca of the United States.* Conchol. Sect., Acad. nat. Sci. Phila., Philadelphia. 238 pp., 17 pls. [Pt. 1, Preface, Key to families, Viviparidae, Ampullariidae, Neritidae, Valvatidae, Amnicolidae, pp. 1-82, pls. 12-17, 1870; pt. 2, Physidae, Limnaeinae, *Limnaea*, pp. 83-124, pls. 16-18, 1871; pt. 3, *Physa, Bulinus*, Pompholinae, Planorbinae, pp. 125-186, pls. 6-9, 1871; pt. 4, Planorbinae, *Carinifex, Planorbis, Segmentina*, Ancylinae, pp. 187-238, pls. 2, 5-7, 1871.]
TRYON, George W., Jr. 1872a. Notes on Dr. James Lewis' paper, "On the shells of the Holston River." *Am. J. Conchol.*, 7(2): 86-88.
TRYON, George W., Jr. 1872b. Monograph of the fresh-water univalve Mollusca of the United States. [Review.] *Am. J. Conchol.*, 7(3): 202.
TRYON, George W., Jr. 1873a. [Land and fluviatile Mollusca from Utah.] *Proc. Acad. nat. Sci. Phila.*, 25: 285-286.
TRYON, George W., Jr. 1873b. Land and fresh-water shells of North America. Part IV. Strepomatidae (American melanians). *Smithson. misc. Collect.*, 16(1): i-lv, 1-435.
TRYON, George W., Jr. 1888. Monograph of the families Neritidae, Neritopsidae, Adeorbiidae, Cyclostrematidae, and Liotiidae. *Manual of conchology, structural and systematic*. Vol. 10: 1-160, pls. 1-36. Conchol. Sect., Acad. nat. Sci. Phila., Philadelphia.
TURNER, Hugh M. 1978. *Hebetancylus excentricus* (Morelet) (Pulmonata: Ancylidae) in Louisiana and a report of septum formation. *Nautilus*, 92(2): 83-85.
TURNER, H.W. 1883. Mollusca. *In: South Carolina: resources and population, institution and industries*. South Carolina State Board of Agriculture. Walker, Evans and Cogswell, Charleston. Pp. 298-304.
TURNER, Ruth D. 1946. John Gould Anthony, with a bibliography and catalogue of his species. *Occ. Pap. Moll.*, 1(8): 81-108.
TURNER, Ruth D. & CLENCH, William J. 1974. A new blind *Physa* from Wyoming with notes on its adaptation to the cave environment. *Nautilus*, 88(3): 80-85.
TUTHILL, Samuel J. 1962. A checklist of North Dakota Pleistocene and Recent Mollusca. *Sterkiana*, (8): 12-18.
TUTHILL, Samuel J. 1963. Corrections and additions to the checklist of North Dakota Pleistocene and Recent Mollusca. *Sterkiana*, (10): 29-30.
TUTHILL, Samuel J., CLAYTON, Lee & LAIRD, Wilson M. 1964. A comparison of a fossil Pleistocene molluscan fauna from North Dakota with a Recent molluscan fauna from Minnesota. *Am. midl. Nat.*, 71(2): 344-362.
VAIL, Virginia A. 1977. Comparative reproductive anatomy of 3 viviparid gastropods. *Malacologia*, 16(2): 519-540.
VAIL, Virginia A. 1978a. Observations on brood production in three viviparid gastropods. *Bull. Am. malacol. Union Inc. 1977*, (44): 90.
VAIL, Virginia A. 1978b. Seasonal reproductive patterns in 3 viviparid gastropods. *Malacologia*, 17(1): 73-97.
VAIL, Virginia A. 1979a. *Campeloma parthenum* (Gastropoda: Viviparidae), a new species from north Florida. *Malacol. Rev.*, 12(1/2): 85-86.
VAIL, Virginia A. 1979b. A preliminary revision of Florida *Lioplax* (Gastropoda: Viviparidae), with a description of *Lioplax talquinensis* sp. nov. *Malacol. Rev.*, 12(1/2): 87-88.
VAIL, Virginia A. 1979c. The species problem in *Campeloma* (Gastropoda: Viviparidae). *Bull. Am. malacol. Union Inc. 1979*, (46): 67.
VAILLANCOURT, Guy & COUTURE, Richard. 1975. Effets des variations de la température de l'eau sur quelques espèces de Physidae, Lymnaeidae, Planorbidae et Hydrobiidae (Gastropoda). *Sterkiana*, (57): 18-32.

VAILLANCOURT, Guy & COUTURE, Richard. 1977. Recolonization of *Bithynia tentaculata* (Linnaeus) (Mollusca, Gastropoda, Prosobranchia) in the zone affected by heated water overflows from the Gentilly nuclear plant, Quebec, Canada. *Sterkiana*, (65/66): 7-10.
VAILLANCOURT, Guy & DINGEMANN, M. 1972. Liste des gastéropodes du fleuve Saint-Laurent, région de Gentilly. *Sterkiana*, (48): 17-29.
VANATTA, E.G. 1895. Notes on the smaller American planorbes. *Nautilus*, 9(5): 52-55.
VANATTA, E.G. 1896. The geographic distribution of *Planorbis umbilicatellus*. *Nautilus*, 9(10): 117.
VANATTA, E.G. 1899. [*Planorbis opercularis* var. *oregonensis* Van. preoccupied, renamed *multilineatus*.] *Nautilus*, 13(4): 48.
VANATTA, E.G. 1902. *Planorbis parvus walkeri* n. var. *Nautilus*, 16(5): 58.
VANATTA, E.G. 1903. Notes on some shells from North Carolina. *Nautilus*, 16(9): 106.
VANATTA, E.G. 1911. *Planorbis bicarinatus* and *Pleurodonte angulata*. *Nautilus*, 24(12): 136-138.
VANATTA, E.G. 1912a. *Viviparus contectoides* in Philadelphia. *Nautilus*, 26(7): 84.
VANATTA, E.G. 1912b. *Cincinnatia* in the Delaware drainage. *Nautilus*, 26(8): 96.
VANATTA, E.G. 1915a. Two new varieties of *Valvata*. *Nautilus*, 28(9): 104-105.
VANATTA, E.G. 1915b. *Lymnaea (Pseudosuccinea) columella* Say in Oregon. *Nautilus*, 29(5): 60.
VANATTA, E.G. 1925. Newfoundland shells. *Nautilus*, 38(3): 92-93.
VANATTA, E.G. 1926. Texas land and freshwater shells. *Nautilus*, 40(1): 16-17.
VANATTA, E.G. 1927. Land and fresh-water shells from Newfoundland. *Nautilus*, 40(3): 112-114.
VANATTA, E.G. 1928. Shells from Live Oak Co., Texas, collected by Dr. Julia Gardner. *Nautilus*, 42(2): 66.
VANATTA, E.G. 1934. *Lyogyrus vanhyningi*, new species, pl. 13, fig. 9. *Nautilus*, 47(4): 149.
VANATTA, E.G. 1935. *Lioplax pilsbryi choctawhatchensis*, n. subsp. *Nautilus*, 49(2): 66-67.
VAN CLEAVE, Harley J. 1932. Studies on snails of the genus *Pleurocera*. I. The eggs and egg laying habits. *Nautilus*, 46(1): 29-34.
VAN CLEAVE, Harley J. 1933. Studies on snails of the genus *Pleurocera*. II. The young of *P. acuta*. *Nautilus*, 47(2): 48-49.
VAN CLEAVE, Harley J. 1936. Reversal of symmetry in *Campeloma rufum*, a fresh-water snail. *Am. Nat.*, 70: 567-573.
VAN CLEAVE, Harley J. & ALTRINGER, Dorothy A. 1937. Studies on the life cycle of *Campeloma rufum*, a fresh-water snail. *Am. Nat.*, 71: 167-184.
VAN CLEAVE, Harley J. & CHAMBERS, Ray. 1935. Studies on the life history of a snail of the genus *Lioplax*. *Am. midl. Nat.*, 16(6): 913-920.
VAN CLEAVE, Harley J. & LEDERER, Ludwig G. 1932. Studies on the life cycle of the snail *Viviparus contectoides*. *J. Morphol.*, 53: 499-522.
VAN CLEAVE, Harley J. & RICHEY, Emily McDavid. 1936. Studies on the radula in snails of the genus *Viviparus*. *Trans. Am. microsc. Soc.*, 60(2): 223-229.
VAN DER SCHALIE, Henry. 1940. Aestivation of *Lymnaea lanceata* (Gould). *Nautilus*, 53(4): 134-135.
VAN DER SCHALIE, Henry. 1953. The ecology and distribution of *Lymnaea (Bulimnea) megasoma* in Michigan. *Annu. Rep. Am. malacol. Union 1952*, (19): 14. [Reprinted in *Sterkiana*, 1963, (9): 31.]
VAN DER SCHALIE, Henry. 1964. Notes on the sex of *Campeloma*. *Annu. Rep. Am. malacol. Union 1964*, (31): 24-25.
VAN DER SCHALIE, Henry. 1965. Observations on the sex of *Campeloma* (Gastropoda: Viviparidae). *Occ. Pap. Mus. Zool. Univ. Mich.*, (641): 1-15.
VAN DER SCHALIE, Henry. 1973. The mollusks of the Duck River drainage in central Tennessee. *Sterkiana*, (52): 45-55.
VAN DER SCHALIE, Henry & BERRY, Elmer G. 1973. The role of temperature in the ecology and distribution of the snail, *Lymnaea stagnalis*. *Malacologia*, 14(1/2): 348.
VAN DER SCHALIE, Henry & DUNDEE, Dee Saunders. 1955. The distribution, ecology and life history of *Pomatiopsis cincinnatiensis* (Lea), an amphibious operculate snail. *Trans. Am. microsc. Soc.*, 74(2): 119-133.
VAN DER SCHALIE, Henry & DUNDEE, Dee Saunders. 1956a. The morphology of *Pomatiopsis cincinnatiensis* (Lea), an amphibious prosobranch snail. *Occ. Pap. Mus. Zool. Univ. Mich.*, (579): 1-17, pls. 1-7.
VAN DER SCHALIE, Henry & DUNDEE, Dee Saunders. 1956b. Studies on *Pomatiopsis* shells in Michigan. *Annu. Rep. Am. malacol. Union 1956*, (23): 22.
VAN DER SCHALIE, Henry & DUNDEE, Dee Saunders. 1959a. The egg-laying habits of *Pomatiopsis lapidaria* and *P. cincinnatiensis*; problems relating to the culture of schistosome intermediate hosts. *Annu. Rep. Am. malacol. Union 1958*, (25): 19-20.
VAN DER SCHALIE, Henry & DUNDEE, Dee Saunders. 1959b. Transect distribution of eggs of *Pomatiopsis*

lapidaria Say, an amphibious prosobranch snail. *Trans. Am. microsc. Soc.*, 78(4): 409-420.
VAN DER SCHALIE, Henry & GETZ, Lowell L. 1961. Comparisons of adult and young *Pomatiopsis cincinnatiensis* (Lea) in respect to moisture requirements. *Trans. Am. microsc. Soc.*, 80(2): 211-220.
VAN DER SCHALIE, Henry & GETZ, Lowell L. 1962a. Reproductive isolation in the snails, *Pomatiopsis lapidaria* and *P. cincinnatiensis*. *Am. midl. Nat.*, 68(1): 189-191.
VAN DER SCHALIE, Henry & GETZ, Lowell L. 1962b. Distribution and natural history of the snail *Pomatiopsis cincinnatiensis* (Lea). *Am. midl. Nat.*, 68(1): 203-231.
VAN DER SCHALIE, Henry & GETZ, Lowell L. 1962c. Morphology and development of the sex organs in the snail *Pomatiopsis cincinnatiensis* (Lea). *Trans. Am. microsc. Soc.*, 81(4): 332-340.
VAN DER SCHALIE, Henry, GETZ, Lowell L. & DAZO, Bonifacio C. 1962. Hybrids between American *Pomatiopsis* and Oriental *Oncomelania* snails. *Am. J. trop. Med. Hyg.*, 11(3): 418-420.
VAN DER SCHALIE, Henry & WALTER, Harold J. 1957. The egg-laying habits of *Pomatiopsis cincinnatiensis* (Lea). *Trans. Am. microsc. Soc.*, 76(4): 404-422.
VAN HYNING, T. 1904. A molluscan stampede. *Nautilus*, 18(3): 31-32.
WALKER, Bryant. 1879. Catalogue of the shell-bearing Mollusca of Michigan. *J. Conchol.*, 2: 325-337.
WALKER, Bryant. 1886. Conchological work in Michigan. *Collect. Sci. Mon.*, 1(1): 1-7.
WALKER, Bryant. 1888. On the re-discovery of *Planorbis multivalvis* Case, and *Planorbis truncatus* Miles. *J. Conchol.*, 5: 330-331.
WALKER, Bryant. 1892-93. The shell bearing Mollusca of Michigan. *Nautilus*, 6(2): 13-19 (1892); 6(3): 31-35, pl. 1 (1892); 6(4): 42-47 (1892); 6(6): 63-67 (1892); 6(12): 135-141 (1893).
WALKER, Bryant. 1894. Shells of the Saginaw Valley, Michigan. *Nautilus*, 7(11): 125-129.
WALKER, Bryant. 1895a. A review of our present knowledge of the molluscan fauna of Michigan. Printed for the author, Detroit. Pp. 1-27. [Reprinted in *Sterkiana*, 1965, (17): 10-25.]
WALKER, Bryant. 1895b. A month with the Michigan Fish Commission. *Nautilus*, 9(1): 2-6.
WALKER, Bryant. 1896. Report upon the Mollusca collected in the vicinity of Charlevoix, Michigan, in the summer of 1894. *Bull. Mich. Fish Comm.*, 6: 96-99.
WALKER, Bryant. 1897. *Planorbis nautileus* L. in America. *Nautilus*, 10(10): 117.
WALKER, Bryant. 1898. Mollusca contemporaneous with the mastodon. *Nautilus*, 11(11): 121-122.
WALKER, Bryant. 1900a. Changes with growth of *Lithasia obovata* (Say). *Nautilus*, 13(9): 97-98.
WALKER, Bryant. 1900b. The genuine *Planorbis corpulentus*, Say. *Nautilus*, 13(12): 133-138, pl. 3.
WALKER, Bryant. 1900c. The origin and distribution of the land and freshwater Mollusca of North America. *First Rep. Mich. Acad. Sci., 1894-1899*, pp. 43-61.
WALKER, Bryant. 1900d. Note on the habits of *Limnaea mighelsi* W.G. Binn. *Nautilus*, 14(1): 8-9.
WALKER, Bryant. 1900e. Supplemental note on *Planorbis corpulentus* Say. *Nautilus*, 14(3): 33-34.
WALKER, Bryant. 1901a. New varieties of *Physa ancillaria* Say. *Nautilus*, 14(9): 97-99.
WALKER, Bryant. 1901b. A new *Amnicola*. *Nautilus*, 14(10): 113-114.
WALKER, Bryant. 1901c. The synonymy of *Bythinella obtusa* Lea. *Nautilus*, 15(3): 30-32.
WALKER, Bryant. 1902a. A revision of the carinate valvatas of the United States. *Nautilus*, 15(11): 121-125.
WALKER, Bryant. 1902b. Notes on the marl fauna. *Rep. Mich. geol. Surv. 1901*, p. 137.
WALKER, Bryant. 1902c. The Mollusca of the Chicago area: Part II., The Gasteropoda. [Review.] *Nautilus*, 16(3): 33-36.
WALKER, Bryant. 1902d. Surface sculpture in *Ancylus*. *Nautilus*, 16(8): 85-88.
WALKER, Bryant. 1903a. On the specific validity of *Campeloma milesii* Lea. *Nautilus*, 16(11): 121-124, pl. 5.
WALKER, Bryant. 1903b. Notes on eastern American ancyli. *Nautilus*, 17(2): 13-19; 17(3): 25-31, pls. 1, 2.
WALKER, Bryant. 1903c. Appendix, on the shells of marls. *Geol. Surv. Mich.*, 8(3): 97-102.
WALKER, Bryant. 1904a. New species of *Somatogyrus*. *Nautilus*, 17(12): 133-142, pl. 5.
WALKER, Bryant. 1904b. Notes on eastern American ancyli. II. *Nautilus*, 18(2): 16-17; 18(3): 25-30, pls. 1, 2.
WALKER, Bryant. 1904c. Hints on collecting land and fresh water Mollusca. *J. appl. Microsc. lab. Methods*, 5: 1954-1961.
WALKER, Bryant. 1904d. Notes on eastern American ancyli. III. *Nautilus*, 18(7): 75-83, pls. 5, 6.
WALKER, Bryant. 1905a. List of shells from northwestern Florida. *Nautilus*, 18(12): 133-136, pl. 9, figs. 1-3, 6, 7.
WALKER, Bryant. 1905b. Recent progress in Michigan conchology with bibliography. *Bull. Mich. Acad. Sci.*, 2(2): 3-7.
WALKER, Bryant. 1906a. New and little known species of Amnicolidae. *Nautilus*, 19(9): 97-100, 114-117, pl. 5.
WALKER, Bryant. 1906b. Notes on *Valvata*. *Nautilus*, 20(3): 25-32, pl. 1.
WALKER, Bryant. 1907a. Mollusca of the southwestern states. II. By H.A. Pilsbry and J.H. Ferriss. Proc. Acad. nat. Science, Philadelphia, 1906, pp. 123-175. [Review.] *Nautilus*, 20(9): 106-108.

WALKER, Bryant. 1907b. A new species of *Segmentina*. *Nautilus*, 20(11): 122-124, pl. 7.
WALKER, Bryant. 1907c. Notes on *Gundlachia*. I. *Nautilus*, 21(2): 14-17, pl. 4.
WALKER, Bryant. 1907d. Notes on *Planorbis* – I. *Nautilus*, 21(6): 61-64, pl. 8.
WALKER, Bryant. 1907e. Marl shells from Cobalt. *Ottawa Nat.*, 21(9): 180.
WALKER, Bryant. 1908a. *Pomatiopsis robusta* n. sp. *Nautilus*, 21(9): 97.
WALKER, Bryant. 1908b. On certain immature anculosae. *Nautilus*, 21(10): 110-117, pl. 10.
WALKER, Bryant. 1908c. New species of Ancylidae. *Nautilus*, 21(11): 126-129, pl. 9, figs. 1-4, 17, 18.
WALKER, Bryant. 1908d. New species of Ancylidae. *Nautilus*, 21(12): 138-140, pl. 9, figs. 5-16.
WALKER, Bryant. 1908e. New Michigan lymnaeas. *Nautilus*, 22(1): 4-9, pls. 1, 2; 22(2): 16-19.
WALKER, Bryant. 1908f. *Lymnaea umbilicata* C.B. Adams in Canada. *Ottawa Nat.*, 22(5): 89-91.
WALKER, Bryant. 1909a. On the use of certain preoccupied names for European Mollusca. *J. Conchol.*, 12(9): 238-240.
WALKER, Bryant. 1909b. Annotated list of the Mollusca of Isle Royale, Michigan. *In:* Adams, Charles C. (Ed.), An ecological survey of Isle Royale, Lake Superior. xiv, 468 pp. [pp. 281-298, figs. 61-63]. *Rep. Board geol. Surv. 1908*.
WALKER, Bryant. 1909c. New Amnicolidae from Alabama. *Nautilus*, 22(9): 85-90, pl. 6.
WALKER, Bryant. 1909d. *Planorbis bicarinatus*. *Nautilus*, 22(11): 122.
WALKER, Bryant. 1909e. Notes on *Planorbis* II: *P. bicarinatus*. *Nautilus*, 23(1): 1-10, pls. 1, 2; 23(2): 21-32, pl. 3.
WALKER, Bryant. 1910. A new species for the United States fauna. *Nautilus*, 24(1): 11.
WALKER, Bryant. 1911a. The conchological survey of Michigan. *Thirteenth Rep. Mich. Acad. Sci., 1911*, pp. 116-120.
WALKER, Bryant. 1911b. A check-list of Michigan Mollusca. *Thirteenth Rep. Mich. Acad. Sci., 1911*, pp. 121-129.
WALKER, Bryant. 1915a. Results of the Mershon Expedition to the Charity Islands, Lake Huron. *Occ. Pap. Mus. Zool. Univ. Mich.*, (7): 1-7.
WALKER, Bryant. 1915b. On *Paludina coarctata* and *incrassata* Lea. *Nautilus*, 28(11): 121-127, pl. 5.
WALKER, Bryant. 1915c. Apical characters in *Somatogyrus* with descriptions of three new species. *Nautilus*, 29(4): 37-41; 29(5): 49-53, figs. 1-4.
WALKER, Bryant. 1915d. A list of shells collected in Arizona, New Mexico, Texas and Oklahoma by Dr. E.C. Case. *Occ. Pap. Mus. Zool. Univ. Mich.*, (15): 1-11.
WALKER, Bryant. 1916. The Mollusca collected in northeastern Nevada by the Walker-Newcomb expedition of the University of Michigan. *Occ. Pap. Mus. Zool. Univ. Mich.*, (29): 1-8, fig. 1.
WALKER, Bryant. 1917a. The type of *Pleurocera* Rafinesque. *Occ. Pap. Mus. Zool. Univ. Mich.*, (38): 1-10.
WALKER, Bryant. 1917b. A revision of the classification of the North American patelliform Ancylidae, with descriptions of new species. *Nautilus*, 31(1): 1-10, pls. 1-3.
WALKER, Bryant. 1917c. A sinistral *Ampullaria*. *Nautilus*, 31(1): 35-36.
WALKER, Bryant. 1917d. A list of shells from the east coast of Florida. *Nautilus*, 31(2): 53-57.
WALKER, Bryant. 1918a. The Mollusca. *In:* Ward, Henry Baldwin & Whipple, George Chandler, *Fresh-water biology*. John Wiley and Sons, New York. 1111 pp. [pp. 957-1020].
WALKER, Bryant. 1918b. A synopsis of the classification of the fresh-water Mollusca of North America, north of Mexico, and a catalogue of the more recently described species, with notes. *Misc. Pub. Mus. Zool. Univ. Mich.*, (6): 1-213.
WALKER, Bryant. 1920. A new fresh-water mollusk from Indiana. *Proc. U.S. natl. Mus.*, 57: 525.
WALKER, Bryant. 1920-21. *Ancylus obscurus* Haldeman and species referred to it. *Nautilus*, 33(3): 99-103 (1920); 34(3): 73-76, pl. 3 (1921).
WALKER, Bryant. 1921a. The life of the Pleistocene or glacial period by Frank Collins Baker. [Review.] *Ecology*, 2(1): 73-75.
WALKER, Bryant. 1921b. Hyperstrophy in the Ancylidae. *Twenty-second Rep. Mich. Acad. Sci., 1920*, pp. 123-124.
WALKER, Bryant. 1922. A list of shells from Goderich, Ontario. *Can. Field-Nat.*, 36: 132-133.
WALKER, Bryant. 1923a. Bourguignat's American species of *Ancylus*. *Nautilus*, 37(1): 7-16, pls. 1-3.
WALKER, Bryant. 1923b. The molluscan fauna of the Big Vermillion River, Illinois by Frank Collins Baker. [Review.] *Ecology*, 4(3): 317.
WALKER, Bryant. 1925a. New species of fresh-water operculates. *Nautilus*, 39(1): 5-8, pl. 1.
WALKER, Bryant. 1925b. New species of North American Ancylidae and Lancidae. *Occ. Pap. Mus. Zool. Univ. Mich.*, (165): 1-7, pls. 1-3.
WALKER, Bryant. 1929a. *Planorbis truncatus* Mighels in New York. *Nautilus*, 42(3): 104.

WALKER, Bryant. 1929b. *Gyraulus vermicularis hendersoni* n.v. *Nautilus*, 42(3): 104-105.
WALKER, Bryant & LANE, A.C. 1900. Recent shells (of the marl). *Rep. geol. Surv. Mich.*, 7(2): 247-252.
WALKER, Bryant & PILSBRY, Henry A. 1902. The Mollusca of the Mt. Mitchell region, North Carolina. *Proc. Acad. nat. Sci. Phila.*, 54: 413-442, pls. 24, 25.
WALKER, Bryant & RUTHVEN, A.G. 1906. Annotated list of the molluscs of the Porcupine Mountains and Isle Royale, Michigan. *In:* Adams, Charles C. (Ed.), An ecological survey in northern Michigan. *Rep. State Board geol. Surv. Mich. 1905.* 638 pp. [pp. 93-99].
WALL, Robert C. 1977. Seasonal movements of the pond snail, *Lymnaea catascopium*, in a northern lake. *Nautilus*, 91(2): 47-51.
WALTER, Harold J. 1959. The morphology of *Lymnaea emarginata serrata* Haldeman with remarks on the systematics of lymnaeids. *Annu. Rep. Am. malacol. Union 1958*, (25): 16-17.
WALTER, Harold J. 1968a. Evolution, taxonomic revolution, and zoogeography of the Lymnaeidae. *Annu. Rep. Am. malacol. Union 1968*, (35): 18-20.
WALTER, Harold J. 1968b. Comments on the paper by J. Burch and G. Lindsay. *Annu. Rep. Am. malacol. Union 1968*, (35): 23-25.
WALTER, Harold J. 1969. Illustrated biomorphology of the "*angulata*" lake form of the basommatophoran snail *Lymnaea catascopium* Say. *Malacol. Rev.*, 2(1): 1-102.
WALTER, Harold J. 1970. *Amphigyra*, miniplanorbs and microsculpture in Planorbidae systematics. *Annu. Rep. Am. malacol. Union 1970*, (37): 47-51.
WALTER, Harold J. 1972. Some occurrences and assemblages of aquatic Ohio Mollusca. *Bull. Am. malacol. Union Inc. 1971*, (38): 40-41.
WALTER, Harold J. & BURCH, John B. 1957. Key to the genera of freshwater gastropods (snails and limpets) occurring in Michigan. *Mus. Zool. Univ. Mich. Circ.*, (3): 1-8.
WASHBURN, Robert G. 1957. *Viviparus contectoides* in Wisconsin. *Nautilus*, 71(2): iii.
WEATHERBURN, Maryl. 1964. Radulae of Ottawa River snails. *Annu. Rep. Am. malacol. Union 1964*, (31): 13.
WENZ, W. 1938-44. Gastropoda, Prosobranchia. *In:* Schindewolf, Otto H., *Handbuch der Paläozoologie.* Bd. 6, Teil 1, vii, 1639 pp. Borntraeger, Berlin.
WESTERLUND, Carl Agardh. 1883. Malakologische Miscellen II. *Nachrichtenbl. dtsch. Malakozool. Ges.*, 15(11/12): 164-174.
WETHERBY, A.G. 1876. Remarks on the variation in form of the family Strepomatidae, with descriptions of new species. *Proc. Cincinnati Soc. nat. Hist. 1875*, (1): 1-12 [reprint pagination], 1 pl.
WETHERBY, A.G. 1877. Review of the genus *Tulotoma*, with remarks on the geographic distribution of the North American Viviparidae. *Q. J. Conchol.*, 1: 207-215.
WETHERBY, A.G. 1879. Notes on some new or little known North American Limnaeidae. *J. Cincinnati Soc. nat. Hist.*, 2: 93-100.
WEYRAUCH, Wolfgang K. 1964. Nomenklatorische Bemerkungen. *Arch. Molluskenkd.*, 93(3/4): 169.
WHEELER, H.E. 1937-38. Notes on "American conchology" by Thomas Say, with special reference to the seventh part edited by T.A. Conrad. *Nautilus*, 51(2): 43-50 (1937); 51(3): 91-97 (1938).
WHITEAVES, Joseph Frederick. 1862. On the land and fresh water Mollusca of lower Canada, with thoughts on the general geographical distribution of animals and plants throughout Canada. *Can. Nat. Geol.*, 6: 452-459. [Reprinted in *Sterkiana*, 1963, (10): 23-27.]
WHITEAVES, Joseph Frederick. 1863a. Trans-Atlantic sketches. - No. 1. On the Little Miami River, Waynesville, Warren County, Ohio. *Zoologist*, 1863 (Feb.): 8119-8124. [Reprinted in *Sterkiana*, 1964, (14): 45-48.]
WHITEAVES, Joseph Frederick. 1863b. On the land and fresh-water Mollusca of lower Canada. *Can. Nat. Geol.*, 8: 98-113. [Reprinted in *Sterkiana*, 1963, (10): 41-49.]
WHITEAVES, Joseph Frederick. 1905. Description of a new species of *Goniobasis*, from British Columbia. *Nautilus*, 19(6): 61-62, pl. 2, figs. 11, 12.
WHITTAKER, E.J. 1919. Notes on variation in *Planorbis campanulatus* Say, from Blue Sea Lake, Quebec. *Nautilus*, 32(4): 127-134.
WHITTEMORE, T.J. 1854. [Note accompanying James Lewis' communication on *Lymnaea catascopium*.] *Proc. Boston Soc. nat. Hist.*, 5: 29.
WINKLEY, Henry W. 1908. *Paludestrina salsa*, Pilsbry. *Nautilus*, 22(6): 53.
WINSLOW, Mina L. 1918. *Pleurobema clava* (Lam.) and *Planorbis dilatatus buchanensis* (Lea) in Michigan. *Occ. Pap. Mus. Zool. Univ. Mich.*, (51): 1-6, pl. 1.
WINSLOW, Mina L. 1921. Mollusca of North Dakota. *Occ. Pap. Mus. Zool. Univ. Mich.*, (98): 1-18.
WINSLOW, Mina L. 1923. Two new freshwater snails from Michigan. *Occ. Pap. Mus. Zool. Univ. Mich.*, (145): 1-4, pl. 1.

WINSLOW, Mina L. 1926a. The varieties of *Planorbis campanulatus* Say. *Occ. Pap. Mus. Zool. Univ. Mich.*, (180): 1-9, pls. 1, 2.
WINSLOW, Mina L. 1926b. A revised check list of Michigan Mollusca. *Occ. Pap. Mus. Zool. Univ. Mich.*, (181): 1-28.
WINSTON, Charles P. 1933. The eggs of *Goniobasis virginica* and *Anculosa carinata*. *J. Wash. Acad. Sci.*, 23: 34-36.
WOLF, John. 1869. Descriptions of three new species of shells. *Am. J. Conchol.*, 5(4): 198, pl. 17, figs. 1-3.
WOOD, Williard M. 1892. *Paludina japonica* Mart. for sale in the San Francisco Chinese markets. *Nautilus*, 5(10): 114-115.
WOODARD, T.M. 1934. Anatomy of the reproductive system of *Goniobasis laqueata* (Say). *J. Tenn. Acad. Sci.*, 9: 243-259.
WURTZ, Charles B. 1955. Fresh-water mollusks and stream pollution. *Annu. Rep. Am. malacol. Union 1955*, (22): 20-21.
WURTZ, Charles B. 1958. Dr. Pilsbry and fresh-water Mollusca. *Nautilus*, 71(3): 84-87.
ZILCH, A. 1959-60. Gastropoda, Euthyneura. *In:* Schindewolf, Otto H., *Handbuch der Paläozoologie*. Bd. 6, Teil 2, xii, 834 pp. Borntraeger, Berlin.

ADDENDA

ADAMS, Henry & ADAMS, Arthur. 1853-58. *The genera of Recent Mollusca; arranged according to their organization*. 3 vols. John Van Voorst, London. [1: 1-256 (1853); 1: 257-484 (1854a); 2: 1-92 (1854b); 2: 93-284 (1855); 2: 285-412 (1856); 2: 413-540 (1857); 2: 541-661 (1858a); 3: 138 pls. (1858b).]
BARBOSA, Frederico S., HUBENDICK, Bengt, MALEK, Emile T. A. & WRIGHT, C.A. 1961. The generic names *Australorbis, Biomphalaria, Platytaphius, Taphius* and *Tropicorbis* (Mollusca, Planorbidae). *Ann. Mag. nat. Hist.*, ser. 13, 4: 371-375.
BARTSCH, Paul. 1908. Notes on the fresh-water mollusk *Planorbis magnificus* and descriptions of two new forms of the same genus from the southern states. *Proc. U.S. natl. Mus.*, 33: 697-700, pl. 57.
BRANSON, Branley A. & BEASLEY, Clark W. 1967. Coiling abnormality in *Physa anatina*. *Nautilus*, 80(4): 144.
CALL, R. Ellsworth. 1885a. Contributions to a knowledge of the fresh-water Mollusca of Kansas. II. Fresh-water univalves. *Bull. Washburn Coll. Lab. nat. Hist.*, 1(2): 51-54.
CALL, R. Ellsworth. 1885b. Contributions to a knowledge of the fresh-water Mollusca of Kansas. IV. *Bull. Washburn Coll. Lab. nat. Hist.*, 1(4): 115-124.
CALL, R. Ellsworth. 1886d. Fifth contribution to a knowledge of the fresh-water Mollusca of Kansas. *Bull. Washburn Coll. Lab. nat. Hist.*, 1(4): 177-184.
CALL, R. Ellsworth. 1887. Sixth contribution to a knowledge of the fresh-water Mollusca of Kansas. *Bull. Washburn Coll. Lab. nat. Hist.*, 2(8): 11-24.
CHAMBERS, Steven M. 1980. Genetic divergence between populations of *Goniobasis* (Pleuroceridae) occupying different drainage systems. *Malacologia*, 20(1): 63-81.
CLENCH, William J. 1929. Freshwater shells of New England. *Bull. Boston Soc. nat. Hist.*, (52): 3-8.
CLENCH, William J. 1966. *Pomacea bridgesi* (Reeve) in Florida. *Nautilus*, 79(3): 105.
DILLON, Robert T., Jr. 1980. The *Goniobasis* of southern Virginia and northwestern North Carolina: genetic and shell morphometric relationships. *Malacologia*, 20(1): 83-98.
EYERDAM, Walter J. 1968. Fresh-water mollusks eaten by trout and other fish. *Nautilus*, 81(3): 103-104.
HARN, E.H. 1891. Western Pennsylvania shells. *Nautilus*, 4(12): 136-137.
IMLAY, Marc J. 1968. Resistance of fresh-water operculate snails to desiccation. *Nautilus*, 81(4): 138-140.
JACKIEWICZ, Maria. 1959. Badania nad zmiennaścia i stanowiskiem systematycznym. *Prace Komisji Biol., Pozn. Tow. Przyj. Nauk*, 19(3): 1-86, pls. 15-25.
JEFFREYS, John Gwyn. 1830. A synopsis of the testaceous pneumonobranchous Mollusca of Great Britain. *Trans. Linn. Soc. Lond.*, 16: 323-392.
JEFFREYS, John Gwyn. 1862. *British conchology. Or an account of the Mollusca which now inhabit the British Isles and the surrounding seas*. Vol. 1, i-cxiv, 1-341, pls. 1-8. John Van Voorst, London.
LA ROCQUE, Aurèle. 1966. Pleistocene Mollusca of Ohio. *Div. geol. Surv. Dept. nat. Res. State of Ohio Bull.*, 62(1): i-iii, 1-111.
METCALF, Artie L. 1967. *Late Quaternary mollusks of the Rio Grande Valley, Caballo Dam, New Mexico to El Paso, Texas*. Sci. Ser., no. 1, 62 pp. Texas Western Press, El Paso.
RHOADS, Samuel N. 1899. Annotated list of land and fresh-water shells recently collected in the vicinity of Miami, Florida. *Nautilus*, 13(4): 43-48.

STAROBOGATOV, Y.I. & STRELETSKAJA, E.A. 1967. Composition and zoogeographical characteristics of the freshwater malacofauna of eastern Siberia and the northern Far East. *In:* Mollusks and their role in biocenoses and composition of faunas. [In Russian.] *Akad. Nauk Soyuza Sov. sots. Resp., Trudy zool. Inst.*, 42: 221-268.

INDEX TO SCIENTIFIC NAMES

Acella, 5, 48, 146, 147, 148
  haldemani, 48, 146, 147, 148
Acmaeidae, 12
Acroloxidae, 4, 48, 66, 146, 147, 188
Acroloxoidea, 4, 66
Acroloxus, 4, 48, 146, 147, 210
  coloradensis, 48, 146, 147
  lacustris, 147
Acroluxus, 210
Acrorbini, 200
Acrorbis, 200
  petricola, 200
acuta acuta, Elimia, 32
acuta acuta, Pleurocera, 45, 142
acuta clavula, Elimia, 32
acuta, Costatella, 55, 165
acuta, Elimia, 32
acuta, Elimia acuta, 32
acuta hinkleyi, Pleurocera, 45
acuta lewisi, Pleurocera, 45, 142
acuta, Physella, 55, 165
acuta, Pleurocera, 45, 197
acuta, Pleurocera acuta, 45, 142
acutifilosa acutifilosa, Calibasis, 41
acutifilosa acutifilosa, Juga, 41
acutifilosa, Calibasis, 135
acutifilosa, Calibasis acutifilosa, 41
acutifilosa, Juga, 135
acutifilosa, Juga acutifilosa, 41
acutifilosa pittensis, Calibasis, 41
acutifilosa pittensis, Juga, 41
acutifilosa siskiyouensis, Calibasis, 41
acutifilosa siskiyouensis, Juga, 41
acutocarinata, Elimia, 130
acutocarinata, Melania, 197
aequicostatus, Hyalopyrgus, 20, 82, 85, 87, 98
aeruginosum, Antillorbis, 58, 171, 178
aeruginosum, Drepanotrema, 58, 171, 178
agarhecta, Marstonia, 27, 101, 102, 103
alabamense, Pleurocera canaliculatum, 46, 142
alabamensis, Amphigyra, 62, 171, 186, 203
alabamensis, Elimia, 36, 127, 211
alabamensis, Goniobasis, 211
alabamensis, Gyrotoma, 133, 197
alabamensis, Melania, 211
alabamensis, Menetus, 202
alabamensis, Micromenetus, 202
alba, Physella gyrina, 54, 162
albanyensis, Elimia boykiniana, 32
alberta, Bakerilymnaea, 49, 152, 205
alberta, Fossaria, 49, 152, 205
alberta, Galba, 205
albofilata, Physella gyrina aurea, 54, 162
alcoviensis, Somatogyrus, 23
aldrichi aldrichi, Amnicola, 28, 109
aldrichi, Amnicola, 196
aldrichi, Amnicola aldrichi, 28, 109
aldrichi antroecetes, Amnicola, 28
aldrichi insolita, Amnicola, 29
aldrichi, Somatogyrus, 93
Alleghenya, 210
alpenensis, Stagnicola, 155
alta, Lanx, 53, 153
altilis, Gillia, 23, 88, 99
alveare, Pleurocera, 45, 142, 143
Amarula, 210
Amblostoma, 210
  eburnea, 210
Ambloxis, 210
  eburnea, 210
Ambloxus, 210
Ameria, 206, 210
ammon, Helisoma, 205, 207
ammon, Pierosoma, 59, 176, 182, 205, 207
ammon, Planorbella, 59, 176, 182, 205, 207
Amnicola, 4, 28-30, 86, 108, 109, 111-115, 195, 196, 203, 204
  aldrichi, 196
  aldrichi aldrichi, 28, 109
  aldrichi antroecetes, 28
  aldrichi insolita, 29
  bakeriana, 29, 196
  browni, 29, 113
  clarkei, 29, 196
  cora, 29, 111
  dalli dalli, 29, 108, 113
  dalli johnsoni, 29, 86, 108, 109, 111
  decisa, 29, 111, 196
  grana, 29, 111, 113
  greggi, 29, 112, 113
  integra, 195
  lapidaria, 195
  limosa, 108, 111, 112, 195, 196, 203
  limosa limosa, 29, 113

limosa parva, 29
lustrica, 195, 196
missouriensis, 29, 196
pallida, 112
pilsbryi, 29, 112, 113
porata, 112, 195, 196
proserpina, 29, 196
pupoidea, 30, 111, 113, 196
retromargo, 30, 86, 111, 113
rhombostoma, 29, 108, 113
stygia, 29, 112
walkeri, 30, 111, 113, 114, 196
Amnicolinae, 4, 28, 85, 86, 108-114
amnicoloides, Somatogyrus, 23, 93
Amphigyra, 6, 62, 171, 186, 203
alabamensis, 62, 171, 186, 203
ampla, Elimia, 40, 132, 197
ampla, Leptoxis, 42, 122, 135
amplum, Gyrotoma, 133, 197
ampullacea, Physella gyrina, 54, 162
Ampullaria, 210
Ampullariidae, 3, 18, 66, 80-82
Ampullarioidea, 3, 65, 83
Ampullarius, 210
Anaplocamus, 210
anatina, Costatella virgata, 56, 166
anatina, Physella virgata, 56, 166
anceps anceps, Helisoma, 58, 174, 178, 201
anceps anticostianum, Helisoma, 201
anceps aroostookense, Helisoma, 201
anceps cahni, Helisoma, 201
anceps, Helisoma, 200-201
anceps, Helisoma anceps, 58, 174, 178, 201
anceps latchfordi, Helisoma, 201
anceps percarinatum, Helisoma, 201
anceps politum, Helisoma, 201
anceps portagensis, Helisoma, 201
anceps royalense, Helisoma, 58, 174, 201
anceps rushi, Helisoma, 201
anceps sayi, Helisoma, 201
anceps striatum, Helisoma, 201
anceps unicarinatum, Helisoma, 201
ancillaria, Physella, 53, 163, 200
Anculosa, 198, 204, 210
anthonyi, 198
geniculata, 198
praerosa, 204
subglobosa, 204
Anculotus, 210
Ancylidae, 6, 62, 67, 147, 160, 168, 188, 189, 191, 192, 208
ancyliformis, Gundlachia, 211
Ancylinae, 6, 62, 189, 191
Ancyloidea, 5, 66, 67
Ancylotus, 210
Ancylus, 188, 210, 211, 213
havanensis, 211
nuttallii, 213
radiatus, 211
Angitrema, 4, 44, 45, 123, 141, 142, 199, 204
armigera, 44, 141
curta, 44, 141
duttoniana, 45, 141
hubrichti, 45
jayana, 45, 142
lima, 45, 123, 142
verrucosa, 45, 142, 199
angitremoides, Io fluvialis, 132, 198
anguistoma, Helisoma, 201
angulata, Helisoma, 201
angulata, Tulotoma, 77, 194
angulatum, Helisoma, 201
annettae, Elimia, 35, 126
annuliferum, Pleurocera, 47, 144, 145
anthonyi, Anculosa, 198
anthonyi, Athearnia crassa, 44, 140, 198
anthonyi, Eurycaelon, 198
anthonyi, Leptoxis crassa, 44, 140, 198
anticostianum, Helisoma anceps, 201
Antillorbis, 5, 57, 58, 171, 178
aeruginosum, 58, 171, 178
Antrobia, 3, 21, 88, 92
culveri, 21, 88, 92
antroecetes, Amnicola aldrichi, 28
Antroselates, 3, 19, 83, 89
spiralis, 19, 83, 89
antrosum, Helisoma, 201
Apella, 198, 210
scissura, 198
Aphaostracon, 3, 19, 20, 82, 87, 89, 91, 98, 203
asthenes, 19, 87, 89
chalarogyrus, 19, 87, 89
hypohyalina, 19, 87, 89
monas, 19, 87, 89
pachynotus, 20, 87, 89
pycnus, 20, 87, 89
rhadinus, 20, 82, 87
theiocrenetus, 20, 87, 91
xynoelictus, 20, 87, 91
Aphella, 210
apicina, Stagnicola, 51
Aplexa, 5, 56, 167, 168, 205
elongata, 56, 167, 168, 205
elongata tryoni, 56, 167
hypnorum, 205
Aplexinae, 5, 56, 167
appressa, Lymnaea stagnalis, 50, 148, 154
arachnoidea arachnoidea, Elimia, 33, 116
arachnoidea, Elimia arachnoidea, 33, 116
arachnoidea spinella, Elimia, 33, 124
Archaeogastropoda, 12
archimedis, Pyrgulopsis, 27, 109, 110
arctica, Stagnicola, 51, 156
arcticus, Gyraulus, 200

arga, Marstonia, 27, 101-103
ariomus, Costatella hendersoni hendersoni, 55, 164
ariomus, Physella hendersoni hendersoni, 55, 164
arkansensis, Leptoxis, 43, 122, 139
arkansensis, Mudalia, 43, 122, 139
Armiger, 5, 57, 168, 177
  crista, 57, 168, 177
armigera, Angitrema, 44, 141
armigera armigera, Planorbula, 61, 173, 174, 185
armigera, Lithasia, 44, 141
armigera, Planorbula armigera, 61, 173, 174, 185
armigera wheatleyi, Planorbula, 61, 173, 185
Armigerus, 210
aroostookense, Helisoma, 201
aroostookense, Helisoma anceps, 201
Aspidobranchia, 12
asthenes, Aphaostracon, 19, 87, 89
aterina, Elimia, 38, 128
athearni, Elimia, 33
athearni, Physella gyrina, 54, 162
Athearnia, 4, 44, 120, 140, 198, 199
  crassa, 120, 198, 199
  crassa anthonyi, 44, 140, 198
  crassa crassa, 44, 140
atkaensis, Lymnaea, 50, 148, 154
augustina, Cincinnatia, 105
aurea albofilata, Physella gyrina, 54, 162
aurea, Physella gyrina, 54, 162
aureus, Somatogyrus, 23, 93, 99
auricularia, Radix, 51, 148, 155
Australorbis, 210
bakeri, Valvata tricarinata, 16, 68
bakeriana, Amnicola, 29, 196
Bakerilymnaea, 5, 49, 50, 149, 151, 152, 154, 160, 204, 205
  alberta, 49, 152, 205
  bulimoides, 49, 149, 151, 152, 154, 205
  bulimoides perplexa, 160, 205
  cockerelli, 49, 152, 154, 205
  cubensis, 50, 149, 152, 154, 160
  dalli, 50, 152, 154, 160, 205
  hendersoni, 50, 152, 205
  perplexa, 50, 152, 205
  perpolita, 50, 151
  sonomaensis, 50, 151, 152, 154, 205
  techella, 50, 152, 154
  vancouverensis, 50, 152
bartschi, Helisoma, 201
basalis, Valvata tricarinata, 16, 68
bayfieldensis, Physella gyrina, 54, 162
bella, Helisoma, 202
bella, Planorbella, 202
bellacrenata, Elimia, 32, 125
Bellamyinae, 3, 17, 77
bellula, Elimia, 37, 129
bentoniensis, Elimia, 34
berendti, Costatella virgata, 56, 166
berendti, Physella virgata, 56, 166
biangulatus, Somatogyrus, 23, 94
bicarinata bicarinata, Valvata, 15, 70
bicarinata morph *normalis, Valvata,* 15, 69, 70
bicarinata, Valvata, 69
bicarinata, Valvata bicarinata, 15, 70
bicarinatum, Helisoma, 201
binneyana, Fontigens, 30, 196
binneyi, Helisoma, 205, 207
binneyi, Pierosoma, 59, 205, 207
binneyi, Planorbella, 59, 205, 207
binneyi, Pomatiopsis, 31, 116, 117
binneyi randolphi, Helisoma, 206
binneyi randolphi, Pierosoma, 206
binneyi randolphi, Planorbella, 206
Biomphalaria, 5, 13, 58, 169, 173, 178
  glabrata, 58, 173, 178
  havanensis, 58, 173, 178
Biomphalariini, 5, 58, 178
Birgella, 4, 25, 97-100
  subglobosa, 25, 97-100
  subglobosa isogona, 25
Bithinia, 210
Bithynia, 3, 19, 81-83
  tentaculata, 81, 83
  tentaculata magnalacustris, 19, 82, 83
  tentaculata tentaculata, 19
Bithyniidae, 3, 19, 66, 81, 83
bonnevillensis, Stagnicola, 51, 156
Borysthenia, 67
  naticina, 67
bottimeri, Costatella, 55, 165
bottimeri, Paludestrina, 31, 114
bottimeri, Physella, 55, 165
boucardi, Physella, 53, 153, 159
Bovillina, 210
boykiniana albanyensis, Elimia, 32
boykiniana boykiniana, Elimia, 32, 116
boykiniana, Elimia, 32
boykiniana, Elimia boykiniana, 32, 116
boykiniana viennaensis, Elimia, 32, 116
brevis, Elimia, 33, 125
brevis, Io fluvialis, 136, 198
brevispira, Costatella integra, 56, 167
brevispira, Leptoxis, 138
brevispira, Physella integra, 56, 167
brevispirum, Campeloma, 17, 79, 194
brevissimus, Hyalopyrgus, 20, 91, 98
bridgesi, Pomacea, 19, 81
brogniartianus, Menetus, 59, 173, 181, 202
brogniartianus, Micromenetus, 59, 173, 181, 202
browni, Amnicola, 29, 113
browni, Lyogyrus, 29, 113
brumbyi, Pleurocera, 46, 145, 199
Buccinum, 199, 211
  glabra, 211
  palustre, 199

buchanensis, Menetus, 202
buchanensis, Menetus dilatatus, 202
buchanensis, Micromenetus, 202
bulbosa, Juga, 41, 135
bulbosa, Oreobasis, 41, 135
Bulimnaea, 210
Bulimnea, 5, 48, 146, 149
  megasoma, 48, 146, 149
bulimoides, Bakerilymnaea, 49, 149, 151, 152, 154, 205
bulimoides, Fossaria, 49, 149, 151, 152, 154, 205
bulimoides, Lymnaea, 149, 152
bulimoides perplexa, Bakerilymnaea, 160, 205
bulimoides perplexa, Fossaria, 160, 205
Bulimula, 210
Bulimus, 210
Bulinidae, 206
Bulinus, 210
bullula, Elimia, 38, 128
Bythinella, 31, 116, 210
  hemphilli, 31, 116
Bythinia, 210
caelatura caelatura, Elimia, 38, 130
caelatura, Elimia, 197
caelatura, Elimia caelatura, 38, 130
caelatura excellens, Elimia, 39, 130
caelatura georgiana, Elimia, 39, 130
caelatura infuscata, Elimia, 39, 131
caelatura lecontiana, Elimia, 39
caelatura luteocella, Elimia, 39, 130
caelatura stearnsiana, Elimia, 39, 130, 131
cahabensis, Clappia, 22, 88
cahawbensis cahawbensis, Elimia, 35, 127
cahawbensis, Elimia cahawbensis, 35, 127
cahawbensis fraterna, Elimia, 35
cahawbensis, Rhodacmea, 189
cahni, Helisoma, 201
cahni, Helisoma anceps, 201
Calibasis, 4, 41, 135
  acutifilosa, 135
  acutifilosa acutifilosa, 41
  acutifilosa pittensis, 41
  acutifilosa siskiyouensis, 41
  occata, 41, 135
californica, Ferrissia, 192
californica, Pomatiopsis, 31, 116, 117
californiense, Helisoma tenue, 206, 207
californiensis, Fontelicella, 26, 103
californiensis, Pierosoma tenuis, 206, 207
californiensis, Planorbella tenuis, 206, 207
Callina, 210
callioglyptus, Menetus, 201, 202
callioglyptus, Menetus opercularis, 180
callioglyptus, Planorbis, 202
campanulata campanulata, Planorbella, 59, 175, 182, 205
campanulata canadensis, Planorbella, 202, 205
campanulata collinsi, Planorbella, 59, 175, 202, 205
campanulata davisi, Planorbella, 202, 205
campanulata michiganensis, Planorbella, 202, 205
campanulata minor, Planorbella, 202
campanulata multivolvis, Planorbella, 205
campanulata, Planorbella, 202, 205
campanulata, Planorbella campanulata, 59, 175, 182, 205
campanulata rideauensis, Planorbella, 202, 205
campanulata rudentis, Planorbella, 202, 205
campanulata smithi, Planorbella, 202, 205
campanulata wisconsinensis, Planorbella, 202, 205
campanulatum campanulatum, Helisoma, 205
campanulatum, Helisoma, 202, 205
campanulatum, Helisoma campanulatum, 205
campanulatum canadense, Helisoma, 202, 205
campanulatum collinsi, Helisoma, 202, 205
campanulatum davisi, Helisoma, 202, 205
campanulatum michiganense, Helisoma, 202, 205
campanulatum minor, Helisoma, 202
campanulatum multivolvis, Helisoma, 205
campanulatum rideauense, Helisoma, 202, 205
campanulatum rudentis, Helisoma, 202, 205
campanulatum smithi, Helisoma, 202, 205
campanulatum wisconsinense, Helisoma, 202, 205
Campeloma, 3, 17, 18, 74-76, 78-81, 194, 195
  brevispirum, 17, 79, 194
  crassula, 17, 74, 78, 79, 194
  decampi, 17, 78, 79, 194
  decisum, 17, 74, 79, 81, 194, 195
  exilis, 17, 194
  floridense, 17, 78, 79, 194
  geniculum, 18, 78-80, 194
  gibbum, 17, 79, 194
  integrum, 17, 74, 194
  leptum, 17, 194
  lewisi, 17, 74
  limum, 18, 74, 78, 194
  milesi, 17, 74, 194
  obesum, 79
  parthenum, 18, 80, 81, 194
  regulare, 18, 74, 78, 80, 194
  subsolidum, 79
  tannum, 17, 79, 194
campestris, Planorbula, 61, 173, 185
canadense, Helisoma campanulatum, 202, 205
canadensis, Planorbella campanulata, 202, 205
canadensis, Stagnicola emarginata, 157
canaliculatum alabamense, Pleurocera, 46, 142
canaliculatum canaliculatum, Pleurocera, 45
canaliculatum excuratum, Pleurocera, 46, 143
canaliculatum filum, Pleurocera, 46, 142, 143
canaliculatum moriforme, Pleurocera, 46, 143
canaliculatum, Pleurocera, 45
canaliculatum, Pleurocera canaliculatum, 45
canaliculatum undulatum, Pleurocera, 46, 142, 143
Cancellariidae, 12

caperata, Hinkleyia, 52, 158
caperata, Stagnicola, 52, 158
capillaris, Elimia, 33, 124
carinata carinata, Leptoxis, 43, 122, 139
carinata carinata, Mudalia, 43, 122, 139
carinata, Leptoxis, 139
carinata, Leptoxis carinata, 43, 122, 139
carinata, Mudalia, 139
carinata, Mudalia carinata, 43, 122, 139
carinata nickliniata, Leptoxis, 43, 122, 139
carinata nickliniata, Mudalia, 43, 122, 139
carinatus, Neoplanorbis, 62, 171, 187
carinifera, Elimia, 32, 116
cariniferum, Gyrotoma, 133, 197
Carinifex, 5, 58, 59, 174, 175, 179, 180, 201
  newberryi, 201
  newberryi jacksonense, 59, 175, 179
  newberryi newberryi, 58, 175, 180
  newberryi newberryi ponsonbyi, 180
  newberryi occidentale, 59, 175, 179
carinocostata, Elimia, 34
Carnifex, 210
carus, Promenetus, 61, 202, 203
castanea, Elimia laqueata, 36, 128
castor, Marstonia, 27, 101-103
catascopium, Lymnaea, 200
catascopium, Stagnicola, 51, 156, 200
catenaria catenaria, Elimia, 33, 124
catenaria dislocata, Elimia, 33, 124, 125
catenaria, Elimia, 33
catenaria, Elimia catenaria, 33, 124
catenaria inclinans, Elimia, 33
catenaria postelli, Elimia, 33, 124
catenaria vanhyningiana, Elimia, 33, 124
centervillensis, Menetus, 201, 202
centervillensis, Planorbis, 202
Ceratodes, 210
Ceriphasia, 210
Ceriphasiidae, 196
Cerithioidea, 65
chacei, Pomatiopsis, 31, 117
chalarogyrus, Aphaostracon, 19, 87, 89
chautauquense, Helisoma, 205
chautauquense, Helisoma trivolvis, 208
chautauquensis, Pierosoma, 205
chautauquensis, Pierosoma trivolvis, 208
chautauquensis, Planorbella, 205
chautauquensis, Planorbella trivolvis, 208
cheatumi, Tryonia, 21, 90, 92, 203
Chilocyclus, 210
chiltonensis, Elimia, 37, 129
chinensis malleata, Cipangopaludina, 17, 75, 77
choctawhatchensis, Lioplax pilsbryi, 18, 76, 195
cimex, Drepanotrema, 58, 172, 179
cimex, Fossulorbis, 58, 172, 179
Cincinna, 211
Cincinnatia, 4, 25, 26, 97, 99-101, 103, 105, 110, 195, 204
  augustina, 105
  cincinnatiensis, 25, 97, 99, 100, 195
  comalensis, 25, 97
  floridana, 25, 100, 105
  fraterna, 25, 99, 100
  helicogyra, 25, 100, 103
  integra, 25, 195
  judayi, 97
  mica, 25, 100, 103
  monroensis, 25, 100
  parva, 25, 100, 103
  peracuta, 25, 103
  petrifons, 25, 100, 103
  ponderosa, 25, 101, 103
  vanhyningi, 26, 101, 105
  wekiwae, 26, 101, 103
cincinnatiensis, Cincinnatia, 25, 97, 99, 100, 195
cincinnatiensis, Pomatiopsis, 31, 115, 116
Cipangopaludina, 3, 17, 75, 77
  chinensis malleata, 17, 75, 77
  japonica, 17, 75, 77
circumlineatus, Promenetus, 203
circumstriatus, Gyraulus, 57, 172, 177
circumstriatus, Torquis, 57, 172, 177
clappi, Clappia, 92, 93
Clappia, 3, 22, 88, 92, 93, 195
  cahabensis, 22, 88
  clappi, 92, 93
  umbilicata, 22, 88, 92, 93
clara, Elimia, 40, 130, 197
clarkei, Amnicola, 29, 196
clathrata, Tryonia, 21, 90, 92
clausa, Elimia, 36, 126
clavaeformis, Elimia, 38, 130
clavaeformis, Melania, 197
clavula, Elimia acuta, 32
clenchi, Elimia, 32, 116
clinchensis, Io fluvialis, 198
clipeata, Leptoxis, 42, 121, 137
cochilaris, Elimia, 33
Cochliopa, 31, 114, 211
  texana, 31, 114
Cochliopina, 3, 22, 83, 84, 92
  riograndensis, 22, 83, 84, 92
cockerelli, Bakerilymnaea, 49, 152, 154, 205
cockerelli, Fossaria, 49, 152, 154, 205
cockerelli, Stagnicola, 152
collinsi, Helisoma campanulatum, 202, 205
collinsi, Planorbella campanulata, 59, 175, 202, 205
coloradensis, Acroloxus, 48, 146, 147
coloradoensis, Promenetus, 202
columbiana, Fluminicola, 22, 93
columbiana, Physella, 53, 159
columbiense, Helisoma, 206, 207
columbiensis, Pierosoma, 60, 206, 207
columbiensis, Planorbella, 60, 206, 207

columella, Pseudosuccinea, 50, 148, 155
comalensis, Cincinnatia, 25, 97
comalensis comalensis, Elimia, 33, 124
comalensis, Elimia comalensis, 33, 124
comalensis fontinalis, Elimia, 33, 125
comma, Elimia, 32
compacta, Leptoxis, 42, 122, 137
conanti, Helisoma, 206
conanti, Planorbella, 206
conanti, Seminolina, 206
Conchylium, 211
concolor, Costatella virgata, 56, 166
concolor, Physella virgata, 56, 166
conica, Spilochlamys, 28, 86, 104, 107, 108
Conidae, 12
conoidea, Costatella, 55, 153
conoidea, Physella, 55, 153
constrictus, Somatogyrus, 23, 94
contracta, Stagnicola, 51, 156
Conus, 12
cooperi, Menetus, 201
cooperi, Physella, 53, 159
cooperi planospirus, Menetus, 201
coosaensis, Leptoxis, 138
coosaensis, Somatogyrus, 24, 88, 93, 97, 99, 203
coosaensis, Tulotoma, 194
coosaensis, Walkerilla, 24, 88, 93, 97, 99, 203
cora, Amnicola, 29, 111
cornuarietis, Marisa, 18, 80, 81
corpulenta corpulenta, Pierosoma, 60, 176, 205, 207
corpulenta corpulenta, Planorbella, 60, 176, 205, 207
corpulenta, Leptoxis, 139
corpulenta, Mudalia, 139
corpulenta, Pierosoma, 206, 207
corpulenta, Pierosoma corpulenta, 60, 176, 205, 207
corpulenta, Planorbella, 206, 207
corpulenta, Planorbella corpulenta, 60, 176, 205, 207
corpulenta vermilionensis, Pierosoma, 60, 176, 205
corpulenta vermilionensis, Planorbella, 60, 176, 205
corpulenta whiteavesi, Pierosoma, 60, 176
corpulenta whiteavesi, Planorbella, 60, 176
corpulentum corpulentum, Helisoma, 205, 207
corpulentum, Helisoma, 206, 207
corpulentum, Helisoma corpulentum, 205, 207
corpulentum, Pleurocera, 47, 145
corpulentum, Strephobasis, 47, 145
corpulentum vermilionense, Helisoma, 205
corrugatum, Helisoma, 201
costata, Costatella, 55, 164
costata, Physella, 55, 164
Costatella, 5, 55, 56, 153, 164-167, 200
  acuta, 55, 165
  bottimeri, 55, 165
  conoidea, 55, 153
  costata, 55, 164
  cubensis cubensis, 55, 164
  cubensis peninsulae, 55, 164
  hendersoni, 55, 165
  hendersoni floridana, 200
  hendersoni hendersoni, 55, 164
  hendersoni hendersoni ariomus, 55, 164
  heterostropha halei, 55, 165
  heterostropha heterostropha, 55, 165
  heterostropha pomila, 55, 165
  humerosa, 55, 165
  integra brevispira, 56, 167
  integra integra, 56, 167
  integra integra walkeri, 56, 167
  johnsoni, 56, 165
  osculans, 56, 165, 200
  spelunca, 56, 166
  squalida, 56, 166
  virgata anatina, 56, 166
  virgata berendti, 56, 166
  virgata concolor, 56, 166
  virgata rhyssa, 56, 166
  virgata virgata, 56, 166
  virgata virgata parva, 56, 166
Costella, 211
costifera, Elimia, 36
costulata, Elimia laqueata, 36, 129
crandalli, Elimia potosiensis, 40
crassa anthonyi, Athearnia, 44, 140, 198
crassa anthonyi, Leptoxis, 44, 140, 198
crassa, Athearnia, 120, 199
crassa, Athearnia crassa, 44, 140
crassa crassa, Athearnia, 44, 140
crassa crassa, Leptoxis, 44, 140
crassa, Leptoxis, 120, 198, 199
crassa, Leptoxis crassa, 44, 140
crassilabris, Menetus, 201
crassilabris, Somatogyrus, 23, 94
crassula, Campeloma, 17, 74, 78, 79, 194
crassus, Somatogyrus, 23, 94
crenatella, Elimia, 33, 124
crispa, Elimia perstriata, 34
crista, Armiger, 57, 168, 177
crista, Gyraulus, 57, 168, 177
crista, Nautilus, 212
cryptica, Fontigens, 30, 114
cubensis, Bakerilymnaea, 50, 149, 152, 154, 160
cubensis, Costatella cubensis, 55, 164
cubensis cubensis, Costatella, 55, 164
cubensis cubensis, Physella, 55, 164
cubensis, Fossaria, 50, 149, 152, 154, 160
cubensis, Lymnaea, 149
cubensis peninsulae, Costatella, 55, 164
cubensis peninsulae, Physella, 55, 164
cubensis, Physella cubensis, 55, 164
culveri, Antrobia, 21, 88, 92
curreyana, Elimia, 36
currierianum, Pleurocera, 46, 145, 199
currierianus, Somatogyrus, 23, 94
curta, Angitrema, 44, 141

curta, Lithasia, 44, 141
curtum curtum, Pleurocera, 47, 145
curtum curtum, Strephobasis, 47, 145
curtum, Pleurocera curtum, 47, 145
curtum roanense, Pleurocera, 47, 146
curtum roanense, Strephobasis, 47, 146
curtum, Strephobasis curtum, 47, 145
curvicostata, Elimia, 35, 126
Cyclemis, 211
Cyclostoma, 211
cyclostoma, Fossaria, 48, 146, 150
cyclostomaformis, Lioplax, 18, 74, 76, 195
cylindracea, Elimia, 37, 129
cylindrica, Physella gyrina, 54, 162
dacryon, Rhapinema, 28, 86, 102, 107, 108
dallesensis, Juga hemphilli, 41, 135
dalli, Amnicola dalli, 29, 108, 113
dalli, Bakerilymnaea, 50, 152, 154, 160, 205
dalli dalli, Amnicola, 29, 108, 113
dalli, Fossaria, 50, 152, 154, 160, 205
dalli johnsoni, Amnicola, 29, 86, 108, 109, 111
danielsi, Valvata sincera, 16, 72, 73
Daudebardiella, 196, 204
davisi, Helisoma campanulatum, 202, 205
davisi, Planorbella campanulata, 202, 205
decampi, Campeloma, 17, 78, 79, 194
decampi, Elimia perstriata, 34, 124, 197
decampi, Fossaria, 48
decipiens, Somatogyrus, 23, 94
decisa, Amnicola, 29, 111, 196
decisum, Campeloma, 17, 74, 79, 81, 194, 195
deflectus, Gyraulus, 57, 172, 177, 200, 207
Dentatus, 211
depressa, Pierosoma occidentalis, 205
depressa, Planorbella occidentalis, 205
depressum, Helisoma occidentale, 205
depressus, Somatogyrus, 23, 99
depygis, Lithasia obovata, 141
deserta, Fontelicella, 26, 105
diaboli, Tryonia, 21, 90, 92
diaphanus, Laevapex, 63, 192, 193
dickinsoni, Elimia, 35, 126
dilatata, Leptoxis, 43, 122, 139
dilatata, Mudalia, 43, 122, 139
dilatatus buchanensis, Menetus, 202
dilatatus buchanensis, Micromenetus, 202
dilatatus floridensis, Menetus, 202
dilatatus floridensis, Micromenetus, 202
dilatatus, Menetus, 59, 173, 180, 181, 202
dilatatus, Micromenetus, 59, 173, 180, 181, 202
dilatatus pennsylvanicus, Menetus, 202
dilatatus pennsylvanicus, Micromenetus, 202
Diotocardia, 12
Discus, 211
disjecta, Pierosoma subcrenata, 206
disjecta, Planorbella subcrenata, 206
disjectum, Helisoma subcrenatum, 206
dislocata, Elimia catenaria, 33, 124, 125
disstoni, Helisoma, 206
disstoni, Planorbella, 206
disstoni, Seminolina, 206
Docoglossa, 12
downiei, Leptoxis, 137
Drepanotrema, 5, 57, 58, 168, 171, 172, 178, 200
  aeruginosum, 58, 171, 178
  cimex, 58, 172, 179
  kermatoides, 58, 172, 178
Drepanotrematinae, 200
Drepanotremeae, 200
Drepanotremini, 5, 57, 178, 179
duryi duryi, Helisoma, 206, 208
duryi duryi, Planorbella, 206, 208
duryi duryi, Seminolina, 206, 208
duryi eudiscus, Helisoma, 206, 208
duryi eudiscus, Planorbella, 206, 208
duryi eudiscus, Seminolina, 206, 208
duryi, Helisoma, 206, 208
duryi, Helisoma duryi, 206, 208
duryi intercalare, Helisoma, 206, 208
duryi intercalaris, Planorbella, 206, 208
duryi intercalaris, Seminolina, 206, 208
duryi normale, Helisoma, 206, 208
duryi normalis, Planorbella, 206, 208
duryi normalis, Seminolina, 206, 208
duryi, Planorbella, 61, 184, 188, 206, 208
duryi, Planorbella duryi, 206, 208
duryi preglabrata, Planorbella, 206, 208
duryi preglabrata, Seminolina, 206, 208
duryi preglabratum, Helisoma, 206, 208
duryi seminole, Helisoma, 206, 208
duryi, Seminolina, 61, 184, 188, 206, 208
duryi, Seminolina duryi, 206, 208
duryi seminolis, Planorbella, 184, 206, 208
duryi seminolis, Seminolina, 184, 206, 208
duttoniana, Angitrema, 45, 141
duttoniana, Lithasia, 45, 141
ebenum ebenum, Elimia, 35, 127
ebenum, Elimia, 35
ebenum, Elimia ebenum, 35, 127
ebenum emeryensis, Elimia, 35
eburnea, Amblostoma, 210
eburnea, Ambloxis, 210
eburnea, Lymnulus, 212
edgariana, Elimia, 33, 124
effusa, Parapholyx, 61, 174, 186, 203
effusa, Vorticifex, 61, 174, 186, 203
elatior, Rhodacmea, 62, 189, 190
Elimia, 4, 6, 32-40, 116, 123-132, 136, 196-198, 204, 211
  acuta, 32
  acuta acuta, 32
  acuta clavula, 32
  acutocarinata, 130
  alabamensis, 36, 127, 211

ampla, 40, 132, 197
annettae, 35, 126
arachnoidea arachnoidea, 33, 116
arachnoidea spinella, 33, 124
aterina, 38, 128
athearni, 33
bellacrenata, 32, 125
bellula, 37, 129
bentoniensis, 34
boykiniana, 32
boykiniana albanyensis, 32
boykiniana boykiniana, 32, 116
boykiniana viennaensis, 32, 116
brevis, 33, 125
bullula, 38, 128
caelatura, 197
caelatura caelatura, 38, 130
caelatura excellens, 39, 130
caelatura georgiana, 39, 130
caelatura infuscata, 39, 131
caelatura lecontiana, 39
caelatura luteocella, 39, 130
caelatura stearnsiana, 39, 130, 131
cahawbensis cahawbensis, 35, 127
cahawbensis fraterna, 35
capillaris, 33, 124
carinifera, 32, 116
carinocostata, 34
catenaria, 33
catenaria catenaria, 33, 124
catenaria dislocata, 33, 124, 125
catenaria inclinans, 33
catenaria postelli, 33, 124
catenaria vanhyningiana, 33, 124
chiltonensis, 37, 129
clara, 40, 130, 197
clausa, 36, 126
clavaeformis, 38, 130
clenchi, 32, 116
cochilaris, 33
comalensis comalensis, 33, 124
comalensis fontinalis, 33, 125
comma, 32
costifera, 36
crenatella, 33, 124
curreyana, 36
curvicostata, 35, 126
cylindracea, 37, 129
dickinsoni, 35, 126
ebenum, 35
ebenum ebenum, 35, 127
ebenum emeryensis, 35
edgariana, 33, 124
fascinans, 39, 131
flava, 35, 127
floridensis, 34
fusiformis, 34, 125
gerhardti, 35, 126, 127, 197
gibbera, 37, 129
hartmaniana, 35, 126
haysiana, 36, 127, 128
hydei, 36, 126
impressa, 34, 124
induta, 35
interrupta, 40, 132
interveniens, 36
jonesi, 39, 131
lachryma, 38, 129
laeta, 38, 129
laqueata, 36
laqueata castanea, 36, 128
laqueata costulata, 36, 129
laqueata laqueata, 36, 126, 129
laqueata tortum, 36
livescens, 37
livescens gracilior, 37
livescens haldemani, 37, 128
livescens livescens, 37, 129
macglameriana, 36, 127
murrayensis, 126
mutabilis, 37
mutabilis mutabilis, 37, 129
mutabilis timidus, 37
mutata, 129
nassula, 34, 125
olivula, 37, 38, 129, 211
ornata, 40, 197
osculata, 211
paupercula, 37, 128
perstriata crispa, 34
perstriata decampi, 34, 124, 197
perstriata perstriata, 34
pilsbryi, 38, 128, 197
plicatastriata, 34
porrecta, 34, 127
potosiensis, 40
potosiensis crandalli, 40
potosiensis ozarkensis, 40, 136
potosiensis plebius, 40, 136
potosiensis potosiensis, 40, 132
proxima, 39, 130
pupaeformis, 34, 125
pupoidea, 36, 128
pybasi, 37
pygmaea, 36, 127
rubella, 127
semicarinata, 38
showalteri, 38, 128, 197
simplex, 38, 39, 130
striatula, 34, 124
strigosa, 34, 126
symmetrica, 39, 131
taitiana, 37, 129
teres, 34, 126

troostiana, 34, 126
ucheensis, 32, 125
vanuxemiana, 38, 39, 131
varians, 35
variata, 38, 128
virginica, 39, 130
Ellipstoma, 197, 211
gibbosa, 197
elliptica, Physella gyrina gyrina, 54, 161
elodes, Lymnaeus, 200
elodes reflexa, Stagnicola, 156
elodes, Stagnicola, 51, 155, 156, 200
elongata, Aplexa, 56, 167, 168, 205
elongata tryoni, Aplexa, 56, 167
elrodi, Stagnicola, 51, 156
elrodiana, Stagnicola, 51, 155
emarginata canadensis, Stagnicola, 157
emarginata nashotahensis, Stagnicola, 157
emarginata, Paludina, 195
emarginata serrata, Stagnicola, 157, 200
emarginata, Stagnicola, 51, 157, 200
emarginatus, Lymnaeus, 195
emarginatus, Stagnicola, 195
emeryensis, Elimia ebenum, 35
Emmericiella, 83
engonatum, Helisoma, 201
Epitoniidae, 12
Epitonioidea, 12
Epitonium, 211
erythopoma, Fluminicola, 22
esterbrooki, Pleurocera, 144
Euamnicola, 211
eucosmium, Helisoma, 58, 174, 201
eucosmium vaughani, Helisoma, 58
eudiscus, Helisoma duryi, 206, 208
eudiscus, Planorbella duryi, 206, 208
eudiscus, Seminolina duryi, 206, 208
Eulimidae, 12
Eurycaelon, 198, 211
anthonyi, 198
exacuous, Promenetus, 61, 173, 186, 201, 202
excavatus, Somatogyrus, 23, 94, 95
excellens, Elimia caelatura, 39, 130
excentricus, Hebetancylus, 63, 191-193
excisum, Gyrotoma, 40, 119, 132, 133, 197
excuratum, Pleurocera canaliculatum, 46, 143
exigua, Fossaria, 49, 150, 151, 153
exilis, Campeloma, 17, 194
exilis, Stagnicola, 51, 155
fallax, Helisoma trivolvis, 206, 208
fallax, Pierosoma trivolvis, 206, 208
fallax, Planorbella trivolvis, 206, 208
fascinans, Elimia, 39, 131
Ferrissia, 6, 62, 63, 190-193
californica, 192
fragilis, 62, 192, 193
mcneili, 62, 190, 192
parallelus, 62, 190
rivularis, 63, 190-192
shimeki, 192
walkeri, 63, 192, 193
Ferrissinae, 6, 62, 188, 191, 192
filosa, Rhodacmea, 62, 189-191
filum, Pleurocera canaliculatum, 46, 142, 143
Fisherola, 5, 52, 147, 158, 200, 210, 213
lancides, 200
nuttalli, 147, 200, 210, 213
nuttalli kootaniensis, 52, 200
nuttalli lancides, 52, 158, 200
nuttalli nuttalli, 52, 200
flava, Elimia, 35, 127
florentiana, Lithasia salebrosa, 44, 141
floridana, Cincinnatia, 25, 100, 105
floridana, Costatella hendersoni, 200
floridana, Physella hendersoni, 200
floridense, Campeloma, 17, 78, 79, 194
floridensis, Elimia, 34
floridensis, Menetus, 202
floridensis, Menetus dilatatus, 202
floridensis, Micromenetus, 202
floridensis, Micromenetus dilatatus, 202
Fluminicola, 3, 22, 65, 84, 92, 93, 195, 203
columbiana, 22, 93
erythopoma, 22
fusca, 22, 92
hindsi, 22
merriami, 22, 93
minutissima, 22, 93
modoci, 22
nevadensis, 22, 93
nuttalliana, 22, 65, 92
seminalis, 22
turbiniformis, 22, 93
virens, 22
fluvialis angitremoides, Io, 132, 198
fluvialis brevis, Io, 136, 198
fluvialis clinchensis, Io, 198
fluvialis fluvialis, Io, 198
fluvialis, Io, 41, 118, 132, 198
fluvialis, Io fluvialis, 198
fluvialis loudonensis, Io, 198
fluvialis lyttonensis, Io, 198
fluvialis nolichuckyensis, Io, 198
fluvialis paulensis, Io, 198
fluvialis powellensis, Io, 198
fluvialis recta, Io, 136, 198
fluvialis spinosa, Io, 136, 198
fluvialis turrita, Io, 136, 198
fluvialis unakensis, Io, 198
fluvialis verrucosa, Io, 136, 198
Fontelicella, 4, 26, 103, 105, 106, 110, 204
californiensis, 26, 103
deserta, 26, 105
hendersoni, 26, 105

idahoensis, 26, 105, 106
intermedia, 26, 105
micrococcus, 26, 103, 106, 110
neomexicana, 26, 105
pilsbryana, 26
robusta, 26, 103, 106
stearnsiana, 26
Fontigens, 4, 30, 31, 110, 111, 114, 116, 196, 204
binneyana, 30, 196
cryptica, 30, 114
holsingeri, 30, 114
nickliniana, 30, 111, 116, 196
orolibas, 30, 114
tartarea, 30, 114
turritella, 30, 114
weberi, 31, 196
Fontigentinae, 4, 30, 85, 110, 111
fontinalis, Elimia comalensis, 33, 125
foremani, Leptoxis, 42, 121, 137
foremani, Pleurocera, 47, 144
formosa, Leptoxis, 42, 121, 204
Fossaria, 5, 48-50, 146, 149-154, 160, 199, 204, 205
alberta, 49, 152, 205
bulimoides, 49, 149, 151, 152, 154, 205
bulimoides perplexa, 160, 205
cockerelli, 49, 152, 154, 205
cubensis, 50, 149, 152, 154, 160
cyclostoma, 48, 146, 150
dalli, 50, 152, 154, 160, 205
decampi, 48
exigua, 49, 150, 151, 153
galbana, 48, 146, 150
hendersoni, 50, 152, 205
humilis, 48, 146, 149-151
modicella, 49, 150, 151, 153
modicella rustica, 151
obrussa, 49, 146, 150, 151, 153
parva, 49, 146, 149, 150, 152
peninsulae, 49, 150, 151, 153
perplexa, 50, 152, 205
perpolita, 50, 151
rustica, 49, 150, 151, 153
sonomaensis, 50, 151, 152, 154, 205
tazewelliana, 49, 146, 150
techella, 50, 152, 154
truncatula, 49, 149, 150, 154, 205
vancouverensis, 50, 152
Fossulorbis, 5, 58, 171, 172, 178, 179
cimex, 58, 172, 179
kermatoides, 58, 172, 178
fragilis, Ferrissia, 62, 192, 193
fraterna, Cincinnatia, 25, 99, 100
fraterna, Elimia cahawbensis, 35
fuliginosa, Lithasia geniculata, 44, 140, 199
fusca, Fluminicola, 22, 92
fuscus, Laevapex, 63, 191-193
fusiformis, Elimia, 34, 125
gabbi, Stagnicola, 52, 157
Galba, 149, 199, 205, 211
alberta, 205
pusilla, 149, 199, 211
galbana, Fossaria, 48, 146, 150
geniculata, Anculosa, 198
geniculata fuliginosa, Lithasia, 44, 140, 199
geniculata geniculata, Lithasia, 44, 140
geniculata, Io, 199
geniculata, Lithasia, 198, 199
geniculata, Lithasia geniculata, 44, 140
geniculata pinguis, Lithasia, 44, 123, 140, 198, 199
geniculum, Campeloma, 18, 78-80, 194
georgiana, Elimia caelatura, 39, 130
georgianus, Somatogyrus, 23, 95
georgianus, Viviparus, 16, 77, 78
gerhardti, Elimia, 35, 126, 127, 197
gibbera, Elimia, 37, 129
gibbosa, Ellipstoma, 197
gibbosa, Leptoxis, 138
gibbum, Campeloma, 17, 79, 194
Gillia, 3, 23, 88, 99
altilis, 23, 88, 99
glabra, Buccinum, 211
glabrata, Biomphalaria, 58, 173, 178
globosa, Physella, 53, 163
Glottella, 211
Goniobasis, 197, 198, 211
alabamensis, 211
olivula, 211
osculata, 197, 211
pilsbryi, 197
umbonata, 198
gouldi, Physella gyrina, 54, 162
gracilior, Elimia livescens, 37
gradatum, Pleurocera, 46, 144
grana, Amnicola, 29, 111, 113
grana, Lyogyrus, 29, 111, 113
granifera, Thiara, 32, 116, 118
gravis, Spilochlamys, 28, 104, 108, 109
greggi, Amnicola, 29, 112, 113
greggi, Lyogyrus, 29, 112, 113
griffithiana, Leptoxis, 204
Gundlachia, 211
ancyliformis, 211
Gymnoglossa, 12
Gyraulus, 5, 57, 168, 172, 177, 200, 207
arcticus, 200
circumstriatus, 57, 172, 177
crista, 57, 168, 177
deflectus, 57, 172, 177, 200, 207
hornensis, 57, 172, 177, 200
parvus, 57, 172, 177
gyrina alba, Physella, 54, 162
gyrina ampullacea, Physella, 54, 162
gyrina athearni, Physella, 54, 162
gyrina aurea albofilata, Physella, 54, 162

gyrina aurea, Physella, 54, 162
gyrina bayfieldensis, Physella, 54, 162
gyrina cylindrica, Physella, 54, 162
gyrina elliptica, Physella gyrina, 54, 161
gyrina gouldi, Physella, 54, 162
gyrina gyrina elliptica, Physella, 54, 161
gyrina gyrina hildrethiana, Physella, 54, 162
gyrina gyrina, Physella, 54, 161
gyrina hawni, Physella, 54, 162
gyrina hildrethiana, Physella gyrina, 54, 162
gyrina microstoma, Physella, 54, 163
gyrina, Physella gyrina, 54, 161
gyrina sayi, Physella, 54, 163
gyrina smithiana, Physella, 54, 163
Gyrotoma, 4, 40, 41, 118, 119, 132-135, 196-198
  alabamensis, 133, 197
  amplum, 133, 197
  cariniferum, 133, 197
  excisum, 40, 119, 132, 133, 197
  hendersoni, 134, 197
  incisum, 133, 197
  laciniatum, 133, 197
  lewisi, 40, 119, 134, 197
  pagodum, 40, 119, 134, 197
  pumilum, 41, 119, 134, 197
  pyramidatum, 41, 119, 134, 197
  spillmani, 133, 197, 198
  walkeri, 41, 119, 135, 197, 198
halcyon, Marstonia, 27, 101, 102, 107
haldemani, Acella, 48, 146-148
haldemani, Elimia livescens, 37, 128
Haldemania, 202, 211
Haldemanina, 211
halei, Costatella heterostropha, 55, 165
halei, Physella heterostropha, 55, 165
hanni, Lanx, 153
harni, Promenetus, 202
hartmaniana, Elimia, 35, 126
hastatum, Pleurocera unciale, 47, 145
Hauffenia, 4, 30, 113, 115, 196, 204
  micra, 30, 113, 115, 196, 204
  nugax, 196, 204
havanensis, Ancylus, 211
havanensis, Biomphalaria, 58, 173, 178
hawni, Physella gyrina, 54, 162
haysiana, Elimia, 36, 127, 128
Hebetancylus, 6, 63, 191-193
  excentricus, 63, 191-193
helicogyra, Cincinnatia, 25, 100, 103
Helicosoma, 211
Helisoma, 5, 58, 59, 174, 175, 178, 179, 200-202, 205-208
  ammon, 205, 207
  anceps, 200, 201
  anceps anceps, 58, 174, 178, 201
  anceps anticostianum, 201
  anceps aroostookense, 201
  anceps cahni, 201
  anceps latchfordi, 201
  anceps pericarinatum, 201
  anceps politum, 201
  anceps portagensis, 201
  anceps royalense, 58, 174, 201
  anceps rushi, 201
  anceps sayi, 201
  anceps striatum, 201
  anceps unicarinatum, 201
  anguistoma, 201
  angulata, 201
  angulatum, 201
  antrosum, 201
  aroostookense, 201
  bartschi, 201
  bella, 202
  bicarinatum, 201
  binneyi, 205, 207
  binneyi randolphi, 206
  cahni, 201
  campanulatum, 202, 205
  campanulatum campanulatum, 205
  campanulatum canadense, 202, 205
  campanulatum collinsi, 202, 205
  campanulatum davisi, 202, 205
  campanulatum michiganense, 202, 205
  campanulatum minor, 202
  campanulatum multivolvis, 205
  campanulatum rideauense, 202, 205
  campanulatum rudentis, 202, 205
  campanulatum smithi, 202, 205
  campanulatum wisconsinense, 202, 205
  chautauquense, 205
  columbiense, 206, 207
  conanti, 206
  corpulentum, 206, 207
  corpulentum corpulentum, 205, 207
  corpulentum vermilionense, 205
  corrugatum, 201
  disstoni, 206
  duryi, 206, 208
  duryi duryi, 206, 208
  duryi eudiscus, 206, 208
  duryi intercalare, 206, 208
  duryi normale, 206, 208
  duryi preglabratum, 206, 208
  duryi seminole, 206, 208
  engonatum, 201
  eucosmium, 58, 174, 201
  eucosmium vaughani, 58
  horni, 205, 206, 208
  idahoense, 201
  jordanense, 201
  kennicotti, 206, 208
  latchfordi, 201
  magnificum, 205

major, 201
minnesotense, 201
multicostatum multicostatum, 205
multicostatum whiteavesi, 205
multivolvis, 202, 205
newberryi, 201
newberryi jacksonense, 59, 175, 179
newberryi newberryi, 58, 175, 180
newberryi newberryi ponsonbyi, 180
newberryi occidentale, 59, 175, 179
occidentale, 207
occidentale depressum, 205
occidentale occidentale, 205
oregonense, 205
percarinatum, 201
pilsbryi, 207, 208
pilsbryi infracarinatum, 205-208
pilsbryi pilsbryi, 205
pilsbryi preblei, 206, 208
plexatum, 205, 206, 208
politum, 201
portagensis, 201
rushi, 201
sayi, 201
scalare, 206, 208
shellense, 201
striatum, 201
subcrenatum, 207, 208
subcrenatum disjectum, 206
subcrenatum horni, 206
subcrenatum perdisjunctum, 206
subcrenatum plexatum, 206
subcrenatum subcrenatum, 205
tenue, 207
tenue californiense, 206, 207
tenue sinuosum, 206, 207
traski, 206, 207
traskii, 207
trivolvis, 206-208
trivolvis chautauquense, 208
trivolvis fallax, 206, 208
trivolvis holstonense, 208
trivolvis lentum, 206, 208
trivolvis macrostomum, 206, 208
trivolvis marshalli, 206, 208
trivolvis subcrenatum, 206-208
trivolvis trivolvis, 206, 208
trivolvis turgidum, 206, 208
truncatum, 206
unicarinatum, 201
winslowi, 206-208
Helisomatinae, 206
Helisomini, 5, 58, 178-187
hemisphaericus, Laevapex, 63
hemphilli, Bythinella, 31, 116
hemphilli dallesensis, Juga, 41, 135
hemphilli hemphilli, Juga, 41, 135
hemphilli, Juga hemphilli, 41, 135
hendersoni ariomus, Costatella hendersoni, 55, 164
hendersoni ariomus, Physella hendersoni, 55, 164
hendersoni, Bakerilymnaea, 50, 152, 205
hendersoni, Costatella, 55, 165
hendersoni, Costatella hendersoni, 55, 164
hendersoni floridana, Costatella, 200
hendersoni floridana, Physella, 200
hendersoni, Fontelicella, 26, 105
hendersoni, Fossaria, 50, 152, 205
hendersoni, Gyrotoma, 134, 197
hendersoni hendersoni ariomus, Costatella, 55, 164
hendersoni hendersoni ariomus, Physella, 55, 164
hendersoni hendersoni, Costatella, 55, 164
hendersoni hendersoni, Physella, 55, 164
hendersoni, Lymnaea, 152
hendersoni, Natricola, 26, 105
hendersoni, Physella, 55, 165
hendersoni, Physella hendersoni, 55, 164
hendersoni, Somatogyrus, 23, 95
heterostropha, Costatella heterostropha, 55, 165
heterostropha halei, Costatella, 55, 165
heterostropha halei, Physella, 55, 165
heterostropha heterostropha, Costatella, 55, 165
heterostropha heterostropha, Physella, 55, 165
heterostropha, Physella heterostropha, 55, 165
heterostropha pomila, Costatella, 55, 165
heterostropha pomila, Physella, 55, 165
hildrethiana, Physella gyrina gyrina, 54, 162
hindsi, Fluminicola, 22
hinkleyi, Pleurocera acuta, 45
hinkleyi, Pomatiopsis, 31, 116, 117, 196
hinkleyi, Rhodacmea, 62, 189-191
hinkleyi, Somatogyrus, 23, 95
hinkleyi, Stagnicola, 52, 157
Hinkleyia, 5, 52, 158
caperata, 52, 158
montanensis, 52, 158
pilsbryi, 52, 158
holsingeri, Fontigens, 30, 114
holstonense, Helisoma trivolvis, 208
holstonensis, Pierosoma trivolvis, 208
holstonensis, Planorbella trivolvis, 208
horati, Valvata utahensis, 16, 70
Horatia, 4, 30, 114, 115, 196, 204
micra, 196, 204
nugax, 30, 114, 115, 196, 204
hordacea, Physella, 54, 161
hornensis, Gyraulus, 57, 172, 177, 200
hornensis, Torquis, 57, 172, 177, 200
horni, Helisoma, 205, 206, 208
horni, Helisoma subcrenatum, 206
horni, Pierosoma, 205, 206, 208
horni, Pierosoma subcrenata, 206
horni, Planorbella, 205, 206, 208
horni, Planorbella subcrenata, 206
Hoyia, 3, 20, 82, 90, 203

sheldoni, 20, 82, 90, 203
hubrichti, Angitrema, 45
hubrichti, Lithasia, 45
hudsonicus, Promenetus, 202
humeralis, Valvata, 15, 69, 72
humerosa, Costatella, 55, 165
humerosa, Paludina, 194
humerosa, Physella, 55, 165
humerosus, Somatogyrus, 23, 95
humilis, Fossaria, 48, 146, 149-151
humilis, Lymnaea, 149
hyalina, Promenetus, 202
Hyalopyrgus, 3, 20, 82, 85, 87, 91, 98
aequicostatus, 20, 82, 85, 87, 98
brevissimus, 20, 91, 98
hydei, Elimia, 36, 126
Hydrobia, 211
Hydrobiidae, 3, 7, 19, 65, 82-84, 89, 91-97, 99, 103, 105-107, 109, 111-116, 196
Hydrobiinae, 3, 19, 82, 85-88, 91, 92, 203
Hydrognoma, 211
hypnorum, Aplexa, 205
hypohyalina, Aphaostracon, 19, 87, 89
Hypsogyra, 211
Ibicornu, 211
idahoense, Helisoma, 201
idahoense, Stagnicola, 52, 157
idahoensis, Fontelicella, 26, 105, 106
idahoensis, Natricola, 26, 105, 106
imitator, Tryonia, 21, 90
impedita, Stagnicola, 156
impressa, Elimia, 34, 124
Incilicornu, 211
incisum, Gyrotoma, 133, 197
inclinans, Elimia catenaria, 33
induta, Elimia, 35
infracarinata, Pierosoma pilsbryi, 60, 182, 205-208
infracarinata, Planorbella pilsbryi, 60, 182, 205-208
infracarinata, Valvata tricarinata, 16, 68
infracarinatum, Helisoma pilsbryi, 205-208
infuscata, Elimia caelatura, 39, 131
insolita, Amnicola aldrichi, 29
integra, Amnicola, 195
integra brevispira, Costatella, 56, 167
integra brevispira, Physella, 56, 167
integra, Cincinnatia, 25, 195
integra, Costatella integra, 56, 167
integra integra, Costatella, 56, 167
integra integra, Physella, 56, 167
integra integra walkeri, Costatella, 56, 167
integra integra walkeri, Physella, 56, 167
integra, Melania, 195
integra, Paludina, 194, 195
integra, Physella integra, 56, 167
integra, Somatogyrus, 24
integra walkeri, Costatella integra, 56, 167
integra walkeri, Physella integra, 56, 167
integrum, Campeloma, 17, 74, 194
intercalare, Helisoma duryi, 206, 208
intercalaris, Planorbella duryi, 206, 208
intercalaris, Seminolina duryi, 206, 208
interioris, Juga, 42, 136
interioris, Oreobasis, 42, 136
intermedia, Fontelicella, 26, 105
interrupta, Elimia, 40, 132
intertextus, Viviparus, 16, 76, 77
interveniens, Elimia, 36
Io, 4, 41, 118, 132, 136, 196, 198, 199
fluvialis, 41, 118, 132, 198
fluvialis angitremoides, 132, 198
fluvialis brevis, 136, 198
fluvialis clinchensis, 198
fluvialis fluvialis, 198
fluvialis loudonensis, 198
fluvialis lyttonensis, 198
fluvialis nolichuckyensis, 198
fluvialis paulensis, 198
fluvialis powellensis, 198
fluvialis recta, 136, 198
fluvialis spinosa, 136, 198
fluvialis turrita, 136, 198
fluvialis unakensis, 198
fluvialis verrucosa, 136, 198
geniculata, 199
salebrosa, 199
isogona, Birgella subglobosa, 25
isogonus, Somatogyrus, 203
jacksonense, Carinifex newberryi, 59, 175, 179
jacksonense, Helisoma newberryi, 59, 175, 179
Janthinidae, 12
japonica, Cipangopaludina, 17, 75, 77
jayana, Angitrema, 45, 142
jayana, Lithasia, 45, 142
jenkinsi, Potamopyrgus, 212
jenksii, Planorbula, 185
jennessi, Physa, 53, 159
johnsoni, Amnicola dalli, 29, 86, 108, 109, 111
johnsoni, Costatella, 56, 165
johnsoni, Physella, 56, 165
jolietensis, Stagnicola, 155
jonesi, Elimia, 39, 131
jordanense, Helisoma, 201
judayi, Cincinnatia, 97
Juga, 4, 41, 42, 123, 135, 136, 196, 204
acutifilosa, 135
acutifilosa acutifilosa, 41
acutifilosa pittensis, 41
acutifilosa siskiyouensis, 41
bulbosa, 41, 135
hemphilli dallesensis, 41, 135
hemphilli hemphilli, 41, 135
interioris, 42, 136
laurae, 42, 136
nigrina, 42, 135

occata, 41, 135
plicifera, 41, 135
silicula, 41, 135
kennicotti, Helisoma, 206, 208
kennicotti, Pierosoma, 206, 208
kennicotti, Planorbella, 206, 208
kermatoides, Drepanotrema, 58, 172, 178
kermatoides, Fossulorbis, 58, 172, 178
Kincaidilla, 211
kingi, Stagnicola, 158
klamathensis, Lanx, 53, 147, 159
klamathensis, Walkerola, 53, 147, 159
kootaniensis, Fisherola nuttalli, 52, 200
labiatus, Menetus, 201
lachryma, Elimia, 38, 129
laciniatum, Gyrotoma, 133, 197
lacustris, Acroloxus, 147
lacustris limafodens, Probythinella, 195
lacustris, Probythinella, 21, 87, 88, 92, 195
laeta, Elimia, 38, 129
Laevapecinae, 6, 63, 191, 192
Laevapex, 6, 63, 191-193
diaphanus, 63, 192, 193
fuscus, 63, 191-193
hemisphaericus, 63
lancides, Fisherola, 200
lancides, Fisherola nuttalli, 52, 158, 200
Lancinae, 5, 52, 147, 153, 158, 159, 188, 200
Lanx, 5, 53, 147, 153, 159, 200, 204
alta, 53, 153
hanni, 153
klamathensis, 53, 147, 159
patelloides, 53, 153, 159
subrotundata, 53, 153
Laphrostoma, 211
lapidaria, Amnicola, 195
lapidaria, Pomatiopsis, 31, 115-117, 195, 196
laqueata castanea, Elimia, 36, 128
laqueata costulata, Elimia, 36, 129
laqueata, Elimia, 36
laqueata, Elimia laqueata, 36, 126, 129
laqueata laqueata, Elimia, 36, 126, 129
laqueata tortum, Elimia, 36
latchfordi, Helisoma, 201
latchfordi, Helisoma anceps, 201
latchfordi, Physella parkeri, 54, 164
laurae, Juga, 42, 136
laurae, Oreobasis, 42, 136
laurentiana, Stagnicola, 156
lecontiana, Elimia caelatura, 39
Lecythoconcha, 211
lens, Menetus, 202
lens, Micromenetus, 202
lenta, Pierosoma trivolvis, 60, 206, 208
lenta, Planorbella trivolvis, 60, 206, 208
lenticularis, Menetus, 202
lenticularis, Micromenetus, 202
lentum, Helisoma trivolvis, 206, 208
Lepetidae, 12
Leptolimnea, 211
Leptoxis, 4, 42-44, 119-123, 135, 137-140, 143, 198, 199, 204
ampla, 42, 122, 135
arkansensis, 43, 122, 139
brevispira, 138
carinata, 139
carinata carinata, 43, 122, 139
carinata nickliniata, 43, 122, 139
clipeata, 42, 121, 137
compacta, 42, 122, 137
coosaensis, 138
corpulenta, 139
crassa, 120, 198, 199
crassa anthonyi, 44, 140, 198
crassa crassa, 44, 140
dilatata, 43, 122, 139
downiei, 137
foremani, 42, 121, 137
formosa, 42, 121, 204
gibbosa, 138
griffithiana, 204
ligata, 42, 121, 137
lirata, 42, 121, 198
melanoides, 42, 122, 137
minor, 43, 122, 139
occultata, 42, 121, 137
picta, 42, 120, 137
plicata, 43, 122, 137
praerosa, 43, 120, 138, 204
showalteri, 43, 121, 138, 198
subglobosa, 138, 204
taeniata, 43, 120, 121, 138, 204
tintinabulum, 138
trilineata, 43, 122, 139
umbilicata, 43, 120, 143
virgata, 43, 123, 140
vittata, 43, 121, 138
leptum, Campeloma, 17, 194
Lepyrium, 3, 23, 84, 99
showalteri, 23, 84, 99
letsoni, Pyrgulopsis, 27, 108, 110, 195
lewisi, Campeloma, 17, 74
lewisi, Gyrotoma, 40, 119, 134, 197
lewisi lewisi, Valvata, 15, 72
lewisi morph *ontarioensis, Valvata*, 15, 67, 70, 71
lewisi, Pleurocera acuta, 45, 142
lewisi, Valvata, 71
lewisi, Valvata lewisi, 15, 72
ligata, Leptoxis, 42, 121, 137
lima, Angitrema, 45, 123, 142
lima, Lithasia, 45, 123, 142
limafodens, Probythinella lacustris, 195
Limnaea, 211
Limnea, 211

Limneus, 211
Limnophila, 4, 64
Limnophysa, 211
limosa, Amnicola, 108, 111, 112, 195, 196, 203
limosa, Amnicola limosa, 29, 113
limosa limosa, Amnicola, 29, 113
limosa, Paludina, 195
limosa parva, Amnicola, 29
limum, Campeloma, 18, 74, 78, 194
Lioplacinae, 3, 17, 74, 79, 80
Lioplax, 3, 18, 74;76, 80, 195
cyclostomaformis, 18, 74, 76, 195
pilsbryi, 195
pilsbryi choctawhatchensis, 18, 76, 195
pilsbryi pilsbryi, 18, 76, 80
subcarinata, 18, 76, 80, 195
subcarinata occidentalis, 195
sulculosa, 18, 76, 195
talquinensis, 18, 76, 80
lirata, Leptoxis, 42, 121, 198
Lithasia, 4, 44, 45, 123, 140-142, 197-199, 204
armigera, 44, 141
curta, 44, 141
duttoniana, 45, 141
geniculata, 198, 199
geniculata fuliginosa, 44, 140, 199
geniculata geniculata, 44, 140
geniculata pinguis, 44, 123, 140, 198, 199
hubrichti, 45
jayana, 45, 142
lima, 45, 123, 142
obovata, 44, 123, 141, 198
obovata depygis, 141
obovata pennsylvanica, 141
obovata sordida, 141
salebrosa, 199
salebrosa florentiana, 44, 141
salebrosa salebrosa, 44, 141
salebrosa subglobosa, 44, 141, 199
showalteri, 197
verrucosa, 45, 142, 199
Lithoglyphinae, 3, 21, 84-86, 92-97, 99, 100
Lithoglyphus, 195, 211
Lithoparches, 211
Littoridinops, 3, 20, 82, 85-87, 90, 91, 98
monroensis, 20, 82, 85, 87, 98
tenuipes, 20, 82, 86, 87, 91, 98
livescens, Elimia, 37
livescens, Elimia livescens, 37, 129
livescens gracilior, Elimia, 37
livescens haldemani, Elimia, 37, 128
livescens livescens, Elimia, 37, 129
lordi, Physella, 54, 161
loudonensis, Io fluvialis, 198
lustrica, Amnicola, 195, 196
lustrica, Marstonia, 27, 101, 104, 106, 107, 195
lustrica, Paludina, 195, 196
Lutella, 211
luteocella, Elimia caelatura, 39, 130
Lymnaea, 5, 50, 148, 149, 152, 154, 199, 200
atkaensis, 50, 148, 154
bulimoides, 149, 152
catascopium, 200
cubensis, 149
hendersoni, 152
humilis, 149
sonomaensis, 152
techella, 152
stagnalis, 199
stagnalis appressa, 50, 148, 154
stagnalis sanctaemariae, 50, 148, 154
Lymnaeidae, 5, 48, 66, 146, 147, 153-160, 188, 199, 200
Lymnaeinae, 5, 48, 146, 147, 153-158, 200
Lymnaeoidea, 5, 66
Lymnaeus, 195, 200, 211
elodes, 200
emarginatus, 195
Lymneus, 211
Lymnula, 211
Lymnulus, 212
eburnea, 212
Lymnus, 212
Lyogyrus, 4, 29, 30, 86, 111-115, 196, 204
browni, 29, 113
grana, 29, 111, 113
greggi, 29, 112, 113
pilsbryi, 29, 112, 113
pupoidea, 30, 111, 113
retromargo, 30, 86, 111, 113
walkeri, 30, 111, 113, 114
Lythasia, 212
lyttonensis, Io fluvialis, 198
macglameriana, Elimia, 36, 127
Macrolimen, 212
macrostoma, Pierosoma trivolvis, 206, 208
macrostoma, Planorbella trivolvis, 206, 208
macrostomum, Helisoma trivolvis, 206, 208
magbimonilifera, Tulotoma, 194
magnalacustris, Bithynia tentaculata, 19, 82, 83
magnalacustris, Physella, 54, 164
magnifica, Pierosoma, 60, 176, 183, 205
magnifica, Planorbella, 60, 176, 183, 205
magnifica, Tulotoma, 16, 75, 77, 194
magnificum, Helisoma, 205
major, Helisoma, 201
malleata, Cipangopaludina chinensis, 17, 75, 77
Marisa, 3, 18, 66, 80, 81
cornuarietis, 18, 80, 81
marmorata, Stenophysa, 57, 167, 168
marshalli, Helisoma trivolvis, 206, 208
marshalli, Pierosoma trivolvis, 206, 208
marshalli, Planorbella trivolvis, 206, 208
Marstonia, 4, 27, 101-104, 106, 107, 195, 204

agarhecta, 27, 101-103
arga, 27, 101-103
castor, 27, 101-103
halcyon, 27, 101, 102, 107
lustrica, 27, 101, 104, 106, 107, 195
ogmorphaphe, 27, 101, 104, 107
olivacea, 27, 104, 106
pachyta, 27, 101, 102, 107
winkleyi mozleyi, 106
Marstoniopsis, 196
maugeriae, Stenophysa, 57, 167, 168
mcneili, Ferrissia, 62, 190, 192
mediocarinata, Valvata tricarinata, 16, 68
Megara, 212
megas, Promenetus, 202
megasoma, Bulimnea, 48, 146, 149
Megastropha, 212
Megasystropha, 201, 212
newberryi, 201
Melacantha, 212
Melafusus, 212
Melania, 117, 195, 197, 211, 212
acutocarinata, 197
alabamensis, 211
clavaeformis, 197
integra, 195
olivula, 211
showalteri, 197
Melanidia, 212
Melaniidae, 117
Melanoides, 4, 31, 116, 117
tuberculata, 31, 116, 117
melanoides, Leptoxis, 42, 122, 137
Melantho, 212
Melas, 212
Melasma, 212
Melatoma, 198, 212
Menetus, 5, 59, 172, 173, 180, 181, 201, 202
alabamensis, 202
brogniartianus, 59, 173, 181, 202
buchanensis, 202
callioglyptus, 201, 202
centervillensis, 201, 202
cooperi, 201
cooperi planospirus, 201
crassilabris, 201
dilatatus, 59, 173, 180, 181, 202
dilatatus buchanensis, 202
dilatatus floridensis, 202
dilatatus pennsylvanicus, 202
floridensis, 202
labiatus, 201
lens, 202
lenticularis, 202
opercularis, 59, 173, 180, 201, 202
opercularis callioglyptus, 180
opercularis multilineatus, 201
opercularis oregonensis, 201, 202
pennsylvanicus, 202
planulatus, 201, 202
portlandensis, 201
sampsoni, 59, 172, 181, 202
mergella, Valvata, 15, 71, 72
merriami, Fluminicola, 22, 93
Meseschiza, 212
Meseshiza, 212
Mesogastropoda, 3, 12, 64
miamiensis, Pomacea, 81
mica, Cincinnatia, 25, 100, 103
michiganense, Helisoma campanulatum, 202, 205
michiganensis, Planorbella campanulata, 202, 205
micra, Hauffenia, 30, 113, 115, 196, 204
micra, Horatia, 196, 204
micra nugax, Valvata, 196, 204
micra, Valvata, 196, 204
Microamnicola, 4, 26, 103, 106, 110
micrococcus, 26, 103, 106, 110
micrococcus, Fontelicella, 26, 103, 106, 110
micrococcus, Microamnicola, 26, 103, 106, 110
Micromelaniidae, 3, 19, 65, 83, 89
Micromenetus, 5, 59, 172, 173, 180, 181, 202
alabamensis, 202
brogniartianus, 59, 173, 181, 202
buchanensis, 202
dilatatus, 59, 173, 180, 181, 202
dilatatus buchanensis, 202
dilatatus floridensis, 202
dilatatus pennsylvanicus, 202
floridensis, 202
lens, 202
lenticularis, 202
pennsylvanicus, 202
sampsoni, 59, 172, 181, 202
microstoma, Physella gyrina, 54, 163
microstriata, Physella, 54, 161
mighelsi, Stagnicola, 52, 149, 157
milesi, Campeloma, 17, 74, 194
minnesotense, Helisoma, 201
minor, Helisoma campanulatum, 202
minor, Leptoxis, 43, 122, 139
minor, Mudalia, 43, 122, 139
minor, Planorbella campanulata, 202
minutissima, Fluminicola, 22, 93
missouriensis, Amnicola, 29, 196
modestum, Pleurocera, 143
modicella, Fossaria, 49, 150, 151, 153
modicella rustica, Fossaria, 151
modoci, Fluminicola, 22
monas, Aphaostracon, 19, 87, 89
moniliferum, Pleurocera, 143
monroensis, Cincinnatia, 25, 100
monroensis, Littoridinops, 20, 82, 85, 87, 98
montanensis, Hinkleyia, 52, 158
montanensis, Stagnicola, 52, 158

moriforme, Pleurocera canaliculatum, 46, 143
mozleyi, Marstonia winkleyi, 106
Mudalia, 4, 43, 120, 122, 123, 139, 140, 196, 198, 204
arkansensis, 43, 122, 139
carinata, 139
carinata carinata, 43, 122, 139
carinata nickliniata, 43, 122, 139
corpulenta, 139
dilatata, 43, 122, 139
minor, 43, 122, 139
trilineata, 43, 122, 139
virgata, 43, 123, 140
multicostata multicostata, Pierosoma, 205
multicostata multicostata, Planorbella, 205
multicostata, Pierosoma multicostata, 205
multicostata, Planorbella multicostata, 205
multicostata whiteavesi, Pierosoma, 205
multicostata whiteavesi, Planorbella, 205
multicostatum, Helisoma multicostatum, 205
multicostatum multicostatum, Helisoma, 205
multicostatum whiteavesi, Helisoma, 205
multilineatus, Menetus opercularis, 201
multivolvis, Helisoma, 202, 205
multivolvis, Helisoma campanulatum, 205
multivolvis, Planorbella, 59, 175, 182, 202, 205
multivolvis, Planorbella campanulata, 205
murrayensis, Elimia, 126
mutabilis, Elimia, 37
mutabilis, Elimia mutabilis, 37, 129
mutabilis mutabilis, Elimia, 37, 129
mutabilis timidus, Elimia, 37
mutata, Elimia, 129
nana, Stiobia, 28, 98, 108, 113
nanus, Somatogyrus, 24, 95
nashotahensis, Stagnicola emarginata, 157
nasoni, Stagnicola, 158
Nasonia, 149, 212
nassula, Elimia, 34, 125
naticina, Borysthenia, 67
naticoides, Paludina, 195
Natricola, 4, 26, 103, 105, 106, 110, 204
hendersoni, 26, 105
idahoensis, 26, 105, 106
robusta, 26, 103, 106
Nauta, 212
Nautilus, 212
crista, 212
Neogastropoda, 12
neomexicana, Fontelicella, 26, 105
neopalustris, Stagnicola, 51, 155
Neoplanorbinae, 6, 62, 186, 187
Neoplanorbis, 6, 62, 171, 186, 187, 203
carinatus, 62, 171, 187
smithi, 62, 171, 187
tantillus, 62, 171, 186
umbilicatus, 62, 171, 187
Nerita, 194, 212
Neritidae, 194
Neritina, 3, 15, 67, 69, 194
reclivata, 67
reclivata palmae, 15, 67
reclivata reclivata, 15, 69
reclivata sphaera, 15, 67
Neritinacea, 3, 12, 64
Neritinia, 194, 213
Neritinidae, 3, 12, 15, 64, 67, 69, 194
Neritinoidea, 3, 64
nevadensis, Fluminicola, 22, 93
nevadensis nevadensis, Pyrgulopsis, 28, 109
nevadensis paiutica, Pyrgulopsis, 28
nevadensis, Pyrgulopsis, 107, 110
nevadensis, Pyrgulopsis nevadensis, 28, 109
newberryi, Carinifex, 201
newberryi, Carinifex newberryi, 58, 175, 180
newberryi, Helisoma, 201
newberryi, Helisoma newberryi, 58, 175, 180
newberryi jacksonense, Carinifex, 59, 175, 179
newberryi jacksonense, Helisoma, 59, 175, 179
newberryi, Megasystropha, 201
newberryi newberryi, Carinifex, 58, 175, 180
newberryi newberryi, Helisoma, 58, 175, 180
newberryi newberryi ponsonbyi, Carinifex, 180
newberryi newberryi ponsonbyi, Helisoma, 180
newberryi occidentale, Carinifex, 59, 175, 179
newberryi occidentale, Helisoma, 59, 175, 179
newberryi, Planorbis, 201
newberryi ponsonbyi, Carinifex newberryi, 180
newberryi ponsonbyi, Helisoma newberryi, 180
newfoundlandensis, Stagnicola, 156
nickliniana, Fontigens, 30, 111, 116, 196
nickliniata, Leptoxis carinata, 43, 122, 139
nickliniata, Mudalia carinata, 43, 122, 139
nigrina, Juga, 42, 135
nigrina, Oreobasis, 42, 135
Nitocris, 204, 212
nobile nobile, Pleurocera, 46, 145
nobile nodosa, Pleurocera, 46, 143
nobile, Pleurocera nobile, 46, 145
nodosa, Pleurocera nobile, 46, 143
nolichuckyensis, Io fluvialis, 198
normale, Helisoma duryi, 206, 208
normalis, Planorbella duryi, 206, 208
normalis, Seminolina duryi, 206, 208
normalis, Valvata bicarinata, 15, 69, 70
Notogillia, 4, 27, 86, 104, 107, 108
sathon, 27, 86, 104, 107
wetherbyi, 27, 104, 107, 108
nugax, Hauffenia, 196, 204
nugax, Horatia, 30, 114, 115, 196, 204
nugax, Valvata micra, 196, 204
nuttalli, Fisherola, 147, 200, 210, 213
nuttalli, Fisherola nuttalli, 52, 200
nuttalli kootaniensis, Fisherola, 52, 200
nuttalli lancides, Fisherola, 52, 158, 200

nuttalli nuttalli, Fisherola, 52, 200
nuttalli, Physella propinqua, 55, 163
nuttalli triticea, Physella propinqua, 55, 163
nuttalli venusta, Physella propinqua, 55, 163
nuttalliana, Fluminicola, 22, 65, 92
nuttallii, Ancylus, 213
nuttallii, Velletea, 213
nylanderi, Valvata sincera, 16, 72
Nymphophilinae, 4, 25, 85, 86, 88, 97, 99-101, 103, 105-109, 113, 114
obesum, Campeloma, 79
obovata depygis, Lithasia, 141
obovata, Lithasia, 44, 123, 141, 198
obovata pennsylvanica, Lithasia, 141
obovata sordida, Lithasia, 141
obrussa, Fossaria, 49, 146, 150, 151, 153
obtusa, Paludina, 196
obtusa, Valvata piscinalis, 16, 71, 72
obtusus, Somatogyrus, 24, 95
occata, Calibasis, 41, 135
occata, Juga, 41, 135
occidentale, Carinifex newberryi, 59, 175, 179
occidentale depressum, Helisoma, 205
occidentale, Helisoma, 207
occidentale, Helisoma newberryi, 59, 175, 179
occidentale, Helisoma occidentale, 205
occidentale occidentale, Helisoma, 205
occidentalis depressa, Pierosoma, 205
occidentalis depressa, Planorbella, 205
occidentalis, Lioplax subcarinata, 195
occidentalis occidentalis, Pierosoma, 205
occidentalis occidentalis, Planorbella, 205
occidentalis, Pierosoma, 60, 183, 207
occidentalis, Pierosoma occidentalis, 205
occidentalis, Planorbella, 60, 183, 207
occidentalis, Planorbella occidentalis, 205
occultata, Leptoxis, 42, 121, 137
ogmorphaphe, Marstonia, 27, 101, 104, 107
olivacea, Marstonia, 27, 104, 106
olivula, Elimia, 37, 38, 129, 211
olivula, Goniobasis, 211
olivula, Melania, 211
Omphemis, 212
Omphiscola, 212
Oncomelania, 115
ontarioensis, Valvata lewisi, 15, 67, 70, 71
opercularis callioglyptus, Menetus, 180
opercularis, Menetus, 59, 173, 180, 201, 202
opercularis multilineatus, Menetus, 201
opercularis oregonensis, Menetus, 201, 202
opercularis oregonensis, Planorbis, 202
optima, Parapholyx solida, 187
optima, Vorticifex solida, 187
oregonense, Helisoma, 205
oregonensis, Menetus opercularis, 201, 202
oregonensis, Pierosoma, 60, 175, 205
oregonensis, Planorbella, 60, 175, 205
oregonensis, Planorbis opercularis, 202
Oreobasis, 4, 41, 42, 135, 136
  bulbosa, 41, 135
  interioris, 42, 136
  laurae, 42, 136
  nigrina, 42, 135
ornata, Elimia, 40, 197
orolibas, Fontigens, 30, 114
oronoensis, Stagnicola, 52, 157
Orygoceras, 4, 27, 98, 106
osculans, Costatella, 56, 165, 200
osculans, Physella, 56, 165, 200
osculata, Elimia, 211
osculata, Goniobasis, 197, 211
Oxytrema, 212
ozarkensis, Elimia potosiensis, 40, 136
ozarkensis, Pyrgulopsis, 28, 110, 195
Pachychilidae, 196
Pachychilus, 196
pachynotus, Aphaostracon, 20, 87, 89
pachyta, Marstonia, 27, 101, 102, 107
pagodum, Gyrotoma, 40, 119, 134, 197
paiutica, Pyrgulopsis nevadensis, 28
pallida, Amnicola, 112
palmae, Neritina reclivata, 15, 67
Paludestrina, 31, 114, 212
  bottimeri, 31, 114
Paludina, 30, 194-196, 206, 212
  emarginata, 195
  humerosa, 194
  integra, 194, 195
  limosa, 195
  lustrica, 195, 196
  naticoides, 195
  obtusa, 30, 196
  porata, 195, 196
  scalaris, 206
Paludomidae, 196
Paludomus, 196
paludosa, Pomacea, 19, 81, 82
palustre, Buccinum, 199
palustris, Stagnicola, 199, 200
Paradines, 212
parallelus, Ferrissia, 62, 190
Parapholyx, 6, 61, 174, 186, 187, 203
  effusa, 61, 174, 186, 203
  solida, 61, 174, 203
  solida optima, 187
parkeri latchfordi, Physella, 54, 164
parkeri parkeri, Physella, 54, 164
parkeri, Physella parkeri, 54, 164
parthenum, Campeloma, 18, 80, 81, 194
parva, Amnicola limosa, 29
parva, Cincinnatia, 25, 100, 103
parva, Costatella virgata virgata, 56, 166
parva, Fossaria, 49, 146, 149, 150, 152
parva, Physella virgata virgata, 56, 166

parvulus, Somatogyrus, 24, 95
parvum, Pleurocera, 46, 143
parvus, Gyraulus, 57, 172, 177
parvus, Torquis, 57, 172, 177
Patellidae, 12
Patelloidea, 12
patelloides, Lanx, 53, 153, 159
paulensis, Io fluvialis, 198
paupercula, Elimia, 37, 128
peninsulae, Costatella cubensis, 55, 164
peninsulae, Fossaria, 49, 150, 151, 153
peninsulae, Physella cubensis, 55, 164
pennsylvanica, Lithasia obovata, 141
pennsylvanicus, Menetus, 202
pennsylvanicus, Menetus dilatatus, 202
pennsylvanicus, Micromenetus, 202
pennsylvanicus, Micromenetus dilatatus, 202
pennsylvanicus, Somatogyrus, 24, 88, 96
peracuta, Cincinnatia, 25, 103
percarinatum, Helisoma, 201
percarinatum, Helisoma anceps, 201
perconfusa, Valvata tricarinata, 16, 70
perdepressa perdepressa, Valvata, 16, 72
perdepressa, Valvata, 16, 71
perdepressa, Valvata perdepressa, 16, 72
perdepressa form *walkeri, Valvata,* 16, 70
perdisjuncta, Pierosoma subcrenata, 206
perdisjuncta, Planorbella subcrenata, 206
perdisjunctum, Helisoma subcrenatum, 206
perplexa, Bakerilymnaea, 50, 152, 205
perplexa, Bakerilymnaea bulimoides, 160, 205
perplexa, Fossaria, 50, 152, 205
perplexa, Fossaria bulimoides, 160, 205
perpolita, Bakerilymnaea, 50, 151
perpolita, Fossaria, 50, 151
perstriata crispa, Elimia, 34
perstriata decampi, Elimia, 34, 124, 197
perstriata, Elimia perstriata, 34
perstriata perstriata, Elimia, 34
petoskeyensis, Stagnicola, 52, 157
petricola, Acrorbis, 200
petrifons, Cincinnatia, 25, 100, 103
Petrophysa, 5, 56, 167, 168
  zionis, 56, 167, 168
Phreatomenetus, 203, 212
Physa, 5, 53, 159, 160, 206
  jennessi, 53, 159
  skinneri, 53, 159
Physella, 5, 53-56, 153, 159-168, 200
  acuta, 55, 165
  ancillaria, 53, 163, 200
  bottimeri, 55, 165
  boucardi, 53, 153, 159
  columbiana, 53, 159
  conoidea, 55, 153
  cooperi, 53, 159
  costata, 55, 164
  cubensis cubensis, 55, 164
  cubensis peninsulae, 55, 164
  globosa, 53, 163
  gyrina alba, 54, 162
  gyrina ampullacea, 54, 162
  gyrina athearni, 54, 162
  gyrina aurea, 54, 162
  gyrina aurea albofilata, 54, 162
  gyrina bayfieldensis, 54, 162
  gyrina cylindrica, 54, 162
  gyrina gouldi, 54, 162
  gyrina gyrina, 54, 161
  gyrina gyrina elliptica, 54, 161
  gyrina gyrina hildrethiana, 54, 162
  gyrina hawni, 54, 162
  gyrina microstoma, 54, 163
  gyrina sayi, 54, 163
  gyrina smithiana, 54, 163
  hendersoni, 55, 165
  hendersoni floridana, 200
  hendersoni hendersoni, 55, 164
  hendersoni hendersoni ariomus, 55, 164
  heterostropha halei, 55, 165
  heterostropha heterostropha, 55, 165
  heterostropha pomila, 55, 165
  hordacea, 54, 161
  humerosa, 55, 165
  integra brevispira, 56, 167
  integra integra, 56, 167
  integra integra walkeri, 56, 167
  johnsoni, 56, 165
  lordi, 54, 161
  magnalacustris, 54, 164
  microstriata, 54, 161
  osculans, 56, 165, 200
  parkeri latchfordi, 54, 164
  parkeri parkeri, 54, 164
  propinqua nuttalli, 55, 163
  propinqua nuttalli triticea, 55, 163
  propinqua nuttalli venusta, 55, 163
  propinqua propinqua, 54, 163
  spelunca, 56, 166
  squalida, 56, 166
  traski, 55, 161
  utahensis, 55, 161
  vinosa, 55, 164
  virgata anatina, 56, 166
  virgata berendti, 56, 166
  virgata concolor, 56, 166
  virgata rhyssa, 56, 166
  virgata virgata, 56, 166
  virgata virgata parva, 56, 166
  virginea, 55, 161
  zionis, 56, 167, 168
Physidae, 5, 53, 66, 147, 153, 159-167, 200, 205
Physina, 212
Physinae, 5, 53, 159, 161-167

Physodon, 212
picta, Leptoxis, 42, 120, 137
Pierosoma, 6, 59, 60, 175, 176, 182, 183, 188, 205-208
ammon, 59, 176, 182, 205, 207
binneyi, 59, 205, 207
binneyi randolphi, 206
chautauquensis, 205
columbiensis, 60, 206, 207
corpulenta, 206, 207
corpulenta corpulenta, 60, 176, 205, 207
corpulenta vermilionensis, 60, 176, 205
corpulenta whiteavesi, 60, 176
horni, 205, 206, 208
kennicotti, 206, 208
magnifica, 60, 176, 183, 205
multicostata multicostata, 205
multicostata whiteavesi, 205
occidentalis, 60, 183, 207
occidentalis depressa, 205
occidentalis occidentalis, 205
oregonensis, 60, 175, 205
pilsbryi, 60, 176, 207, 208
pilsbryi infracarinata, 60, 182, 205-208
pilsbryi pilsbryi, 60, 205
pilsbryi preblei, 206, 208
pilsbryi winslowi, 60
plexata, 205, 206, 208
subcrenata, 207, 208
subcrenata disjecta, 206
subcrenata horni, 206
subcrenata perdisjuncta, 206
subcrenata plexata, 206
subcrenata subcrenata, 205
tenuis, 60, 183, 207
tenuis californiensis, 206, 207
tenuis sinuosa, 206, 207
traski, 60, 206, 207
trivolvis, 188, 206-208
trivolvis chautauquensis, 208
trivolvis fallax, 206, 208
trivolvis holstonensis, 208
trivolvis lenta, 60, 206, 208
trivolvis macrostoma, 206, 208
trivolvis marshalli, 206, 208
trivolvis subcrenata, 60, 176, 183, 206-208
trivolvis trivolvis, 60, 184, 206, 208
trivolvis turgida, 60, 206, 208
truncata, 60, 176, 184, 206
winslowi, 206-208
Pila, 210
pilsbryana, Fontelicella, 26
pilsbryanus, Somatogyrus, 24, 96
pilsbryi, Amnicola, 29, 112, 113
pilsbryi choctawhatchensis, Lioplax, 18, 76, 195
pilsbryi, Elimia, 38, 128, 197
pilsbryi, Goniobasis, 197
pilsbryi, Helisoma, 207, 208
pilsbryi, Helisoma pilsbryi, 205
pilsbryi, Hinkleyia, 52, 158
pilsbryi infracarinata, Pierosoma, 60, 182, 205-208
pilsbryi infracarinata, Planorbella, 60, 182, 205-208
pilsbryi infracarinatum, Helisoma, 205-208
pilsbryi, Lioplax, 195
pilsbryi, Lioplax pilsbryi, 18, 76, 80
pilsbryi, Lyogyrus, 29, 112, 113
pilsbryi, Pierosoma, 60, 176, 207, 208
pilsbryi, Pierosoma pilsbryi, 60, 205
pilsbryi pilsbryi, Helisoma, 205
pilsbryi pilsbryi, Lioplax, 18, 76, 80
pilsbryi pilsbryi, Pierosoma, 60, 205
pilsbryi pilsbryi, Planorbella, 60, 205
pilsbryi, Planorbella, 60, 176, 207, 208
pilsbryi, Planorbella pilsbryi, 60, 205
pilsbryi preblei, Helisoma, 206, 208
pilsbryi preblei, Pierosoma, 206, 208
pilsbryi preblei, Planorbella, 206, 208
pilsbryi, Stagnicola, 52, 158
pilsbryi winslowi, Pierosoma, 60
pilsbryi winslowi, Planorbella, 60
pinguis, Lithasia geniculata, 44, 123, 140, 198, 199
piscinalis form *obtusa, Valvata*, 16, 71, 72
pittensis, Calibasis acutifilosa, 41
pittensis, Juga acutifilosa, 41
planogyrum, Pleurocera, 144
Planorbella, 5, 59-61, 174-176, 182-184, 188, 202, 205-208
ammon, 59, 176, 182, 205, 207
bella, 202
binneyi, 59, 205, 207
binneyi randolphi, 206
campanulata, 202, 205
campanulata campanulata, 59, 175, 182, 205
campanulata canadensis, 202, 205
campanulata collinsi, 59, 175, 202, 205
campanulata davisi, 202, 205
campanulata michiganensis, 202, 205
campanulata minor, 202
campanulata multivolvis, 205
campanulata rideauensis, 202, 205
campanulata rudentis, 202, 205
campanulata smithi, 202, 205
campanulata wisconsinensis, 202, 205
chautauquensis, 205
columbiensis, 60, 206, 207
conanti, 206
corpulenta, 206, 207
corpulenta corpulenta, 60, 176, 205, 207
corpulenta vermilionensis, 60, 176, 205
corpulenta whiteavesi, 60, 176
disstoni, 206
duryi, 61, 184, 188, 206, 208
duryi duryi, 206, 208
duryi eudiscus, 206, 208

duryi intercalaris, 206, 208
duryi normalis, 206, 208
duryi preglabrata, 206, 208
duryi seminolis, 184, 206, 208
horni, 205, 206, 208
kennicotti, 206, 208
magnifica, 60, 176, 183, 205
multicostata multicostata, 205
multicostata whiteavesi, 205
multivolvis, 59, 175, 182, 202, 205
occidentalis, 60, 183, 207
occidentalis depressa, 205
occidentalis occidentalis, 205
oregonensis, 60, 175, 205
pilsbryi, 60, 176, 207, 208
pilsbryi infracarinata, 60, 182, 205-208
pilsbryi pilsbryi, 60, 205
pilsbryi preblei, 206, 208
pilsbryi winslowi, 60
plexata, 205, 206, 208
scalaris, 61, 184, 188, 206, 208
subcrenata, 207, 208
subcrenata disjecta, 206
subcrenata horni, 206
subcrenata perdisjuncta, 206
subcrenata plexata, 206
subcrenata subcrenata, 205
tenuis, 60, 183, 207
tenuis californiensis, 206, 207
tenuis sinuosa, 206, 207
traski, 60, 206, 207
trivolvis, 188, 206-208
trivolvis chautauquensis, 208
trivolvis fallax, 206, 208
trivolvis holstonensis, 208
trivolvis lenta, 60, 206, 208
trivolvis macrostoma, 206, 208
trivolvis marshalli, 206, 208
trivolvis subcrenata, 60, 176, 183, 206-208
trivolvis trivolvis, 60, 184, 206, 208
trivolvis turgida, 60, 206, 208
truncata, 60, 176, 184, 206
winslowi, 206-208
Planorbidae, 5, 57, 66, 147, 160, 168, 177-188, 203
Planorbina, 212
Planorbinae, 5, 57, 177-187
Planorbini, 5, 57, 177
Planorbis, 201, 202, 212
callioglyptus, 202
centervillensis, 202
newberryi, 201
opercularis oregonensis, 202
planulatus, 202
umbilicatus, 202
wheatleyi, 202
Planorbula, 6, 61, 171, 173, 174, 185, 202
armigera armigera, 61, 173, 174, 185
armigera wheatleyi, 61, 173, 185
campestris, 61, 173, 185
jenksii, 185
wheatleyi, 202
Planorbulina, 212
planospirus, Menetus cooperi, 201
planulatus, Menetus, 201, 202
planulatus, Planorbis, 202
platyrachis, Pyrgophorus, 21, 86, 92, 98
plebius, Elimia potosiensis, 40, 136
Pleurocera, 4, 6, 45-47, 123, 142-146, 196-199, 204
acuta, 45, 197
acuta acuta, 45, 142
acuta hinkleyi, 45
acuta lewisi, 45, 142
alveare, 45, 142, 143
annuliferum, 47, 144, 145
brumbyi, 46, 145, 199
canaliculatum, 45
canaliculatum alabamense, 46, 142
canaliculatum canaliculatum, 45
canaliculatum excuratum, 46, 143
canaliculatum filum, 46, 142, 143
canaliculatum moriforme, 46, 143
canaliculatum undulatum, 46, 142, 143
corpulentum, 47, 145
currierianum, 46, 145, 199
curtum curtum, 47, 145
curtum roanense, 47, 146
esterbrooki, 144
foremani, 47, 144
gradatum, 46, 144
modestum, 143
moniliferum, 143
nobile nobile, 46, 145
nobile nodosa, 46, 143
parvum, 46, 143
planogyrum, 144
ponderosum, 143
postelli, 46, 144
prasinatum, 47, 144
pyrenellum, 46, 144, 145, 199
showalteri, 47, 144
striatum, 142
trochiformis, 46, 145
unciale, 47, 144
unciale hastatum, 47, 145
unciale unciale, 47, 144, 145
vestitum, 47, 144, 145
viridulum, 47, 145, 199
walkeri, 47, 146
Pleuroceridae, 4, 6, 32, 65, 116-118, 124-146, 196, 198
Pleurovalvata, 212
plexata, Pierosoma, 205, 206, 208
plexata, Pierosoma subcrenata, 206
plexata, Planorbella, 205, 206, 208
plexata, Planorbella subcrenata, 206

plexatum, Helisoma, 205, 206, 208
plexatum, Helisoma subcrenatum, 206
plicata, Leptoxis, 43, 122, 137
plicatastriata, Elimia, 34
plicifera, Juga, 41, 135
politum, Helisoma, 201
politum, Helisoma anceps, 201
Pomacea, 3, 19, 66, 81, 82
  bridgesi, 19, 81
  miamiensis, 81
  paludosa, 19, 81, 82
Pomatias, 211
Pomatiopsidae, 4, 31, 65, 115, 116
Pomatiopsis, 4, 31, 115-117, 195, 196
  binneyi, 31, 116, 117
  californica, 31, 116, 117
  chacei, 31, 117
  cincinnatiensis, 31, 115, 116
  hinkleyi, 31, 116, 117, 196
  lapidaria, 31, 115-117, 195, 196
pomila, Costatella heterostropha, 55, 165
pomila, Physella heterostropha, 55, 165
Pompholycodea, 212
Pompholyx, 212
Pomus, 212
ponderosa, Cincinnatia, 25, 101, 103
ponderosum, Pleurocera, 143
ponsonbyi, Carinifex newberryi newberryi, 180
ponsonbyi, Helisoma newberryi newberryi, 180
porata, Amnicola, 112, 195, 196
porata, Paludina, 195, 196
porrecta, Elimia, 34, 127
portagensis, Helisoma, 201
portagensis, Helisoma anceps, 201
portlandensis, Menetus, 201
postelli, Elimia catenaria, 33, 124
postelli, Pleurocera, 46, 144
Potamopyrgus, 212
  jenkinsi, 212
potosiensis crandalli, Elimia, 40
potosiensis, Elimia, 40
potosiensis, Elimia potosiensis, 40, 132
potosiensis ozarkensis, Elimia, 40, 136
potosiensis plebius, Elimia, 40, 136
potosiensis potosiensis, Elimia, 40, 132
powellensis, Io fluvialis, 198
praerosa, Anculosa, 204
praerosa, Leptoxis, 43, 120, 138, 204
prasinatum, Pleurocera, 47, 144
preblei, Helisoma pilsbryi, 206
preblei, Pierosoma pilsbryi, 206, 208
preblei, Planorbella pilsbryi, 206, 208
preglabrata, Planorbella duryi, 206, 208
preglabrata, Seminolina duryi, 206, 208
preglabratum, Helisoma duryi, 206, 208
Probythinella, 3, 21, 87, 88, 92, 195
  lacustris, 21, 87, 88, 92, 195
  lacustris limafodens, 195
producta, Pseudisidora, 147
Promenetus, 6, 61, 173, 186, 201-203
  carus, 61, 202, 203
  circumlineatus, 203
  coloradoensis, 202
  exacuous, 61, 173, 186, 201, 202
  harni, 202
  hudsonicus, 202
  hyalina, 202
  megas, 202
  rubellus, 202
  umbilicatellus, 61, 173, 186, 202, 203
  umbilicatus, 202
propinqua nuttalli, Physella, 55, 163
propinqua nuttalli triticea, Physella, 55, 163
propinqua nuttalli venusta, Physella, 55, 163
propinqua, Physella propinqua, 54, 163
propinqua propinqua, Physella, 54, 163
proserpina, Amnicola, 29, 196
Prosobranchia, 3, 12, 64, 160
protea, Tryonia, 21, 90, 92
proxima, Elimia, 39, 130
Pseudisidora, 147
  producta, 147
Pseudogalba, 149, 212
Pseudosuccinea, 5, 50, 148, 155
  columella, 50, 148, 155
Ptenoglossa, 12
Pulmonata, 4, 64
pumilis, Somatogyrus, 24, 96
pumilum, Gyrotoma, 41, 119, 134, 197
pupaeformis, Elimia, 34, 125
pupoidea, Amnicola, 30, 111, 113, 196
pupoidea, Elimia, 36, 128
pupoidea, Lyogyrus, 30, 111, 113
pusilla, Galba, 149, 199, 211
pybasi, Elimia, 37
pycnus, Aphaostracon, 20, 87, 89
pygmaea, Elimia, 36, 127
pygmaeus, Somatogyrus, 24, 96
pyramidatum, Gyrotoma, 41, 119, 134, 197
Pyramidellidae, 12
pyrenellum, Pleurocera, 46, 144, 145, 199
Pyrgophorus, 3, 21, 86, 90-92, 98, 203
  platyrachis, 21, 86, 92, 98
  spinosus, 21, 90, 91
Pyrgula, 213
Pyrgulopsis, 4, 27, 28, 107-110, 195
  archimedis, 27, 109, 110
  letsoni, 27, 108, 110, 195
  nevadensis, 107, 110
  nevadensis nevadensis, 28, 109
  nevadensis paiutica, 28
  ozarkensis, 28, 110, 195
  scalariformis, 28, 109, 110, 195
  wabashensis, 195

quadratus, Somatogyrus, 24, 96
Rachiglossa, 12
radiatus, Ancylus, 211
Radix, 5, 51, 148, 155
  auricularia, 50, 148, 155
randolphi, Helisoma binneyi, 206
randolphi, Pierosoma binneyi, 206
randolphi, Planorbella binneyi, 206
reclivata, Neritina, 67
reclivata, Neritina reclivata, 15, 69
reclivata palmae, Neritina, 15, 67
reclivata reclivata, Neritina, 15, 69
reclivata sphaera, Neritina, 15, 67
recta, Io fluvialis, 136, 198
reflexa, Stagnicola elodes, 156
regulare, Campeloma, 18, 74, 78, 80, 194
retromargo, Amnicola, 30, 86, 111, 113
retromargo, Lyogyrus, 30, 86, 111, 113
rhadinus, Aphaostracon, 20, 82, 87
Rhapinema, 4, 28, 86, 102, 107, 108, 204
  dacryon, 28, 86, 102, 107, 108
Rhipidoglossa, 12
rhodacme, Rhodacmea, 189, 191
Rhodacmea, 6, 62, 188-191, 209
  cahawbensis, 189
  elatior, 62, 189, 190
  filosa, 62, 189-191
  hinkleyi, 62, 189-191
  rhodacme, 189, 191
Rhodocephala, 213
rhombostoma, Amnicola, 29, 108, 113
rhyssa, Costatella virgata, 56, 166
rhyssa, Physella virgata, 56, 166
rideauense, Helisoma campanulatum, 202, 205
rideauensis, Planorbella campanulata, 202, 205
riograndensis, Cochliopina, 22, 83, 84, 92
Rissooidea, 65
rivularis, Ferrissia, 63, 190-192
roanense, Pleurocera curtum, 47, 146
roanense, Strephobasis curtum, 47, 146
robusta, Fontelicella, 26, 103, 106
robusta, Natricola, 26, 103, 106
royalense, Helisoma anceps, 58, 174, 201
rubella, Elimia, 127
rubellus, Promenetus, 202
rudentis, Helisoma campanulatum, 202, 205
rudentis, Planorbella campanulata, 202, 205
rushi, Helisoma, 201
rushi, Helisoma anceps, 201
rustica, Fossaria, 49, 150, 151, 153
rustica, Fossaria modicella, 151
salebrosa florentiana, Lithasia, 44, 141
salebrosa, Io, 199
salebrosa, Lithasia, 199
salebrosa, Lithasia salebrosa, 44, 141
salebrosa salebrosa, Lithasia, 44, 141
salebrosa subglobosa, Lithasia, 44, 141, 199
sampsoni, Menetus, 59, 172, 181, 202
sampsoni, Micromenetus, 59, 172, 181, 202
sanctaemariae, Lymnaea stagnalis, 50, 148, 154
sargenti, Somatogyrus, 24, 96
sathon, Notogillia, 27, 86, 104, 107
sayi, Helisoma, 201
sayi, Helisoma anceps, 201
sayi, Physella gyrina, 54, 163
scalare, Helisoma, 206, 208
scalariformis, Pyrgulopsis, 28, 109, 110, 195
scalaris, Paludina, 206
scalaris, Planorbella, 61, 184, 188, 206, 208
scalaris, Seminolina, 61, 184, 188, 206, 208
Scalidae, 12
Scaphe, 213
Schizochilus, 213
Schizostoma, 198, 213
scissura, Apella, 198
Segmentineae, 200
Segmentina, 213
semicarinata, Elimia, 38
seminalis, Fluminicola, 22
seminole, Helisoma duryi, 206, 208
seminolis, Planorbella duryi, 184, 206, 208
seminolis, Seminolina duryi, 184, 206, 208
shellense, Helisoma, 201
Seminolina, 6, 61, 175, 184, 188, 206, 208
  conanti, 206
  disstoni, 206
  duryi, 61, 184, 188, 206, 208
  duryi duryi, 206, 208
  duryi eudiscus, 206, 208
  duryi intercalaris, 206, 208
  duryi normalis, 206, 208
  duryi preglabrata, 206, 208
  duryi seminolis, 184, 206, 208
  scalaris, 61, 184, 188, 206, 208
serrata, Stagnicola emarginata, 157, 200
sheldoni, Hoyia, 20, 82, 90, 203
shimeki, Ferrissia, 192
showalteri, Elimia, 38, 128, 197
showalteri, Leptoxis, 43, 121, 138, 198
showalteri, Lepyrium, 23, 84, 99
showalteri, Lithasia, 197
showalteri, Melania, 197
showalteri, Pleurocera, 47, 144
silicula, Juga, 41, 135
simplex, Elimia, 38, 39, 130
simplex, Valvata tricarinata, 16, 70
Simpsonia, 149, 213
sincera form *danielsi, Valvata,* 16, 72, 73
sincera nylanderi, Valvata, 16, 72
sincera sincera, Valvata, 16, 68, 72, 73
sincera, Valvata sincera, 16, 68, 72, 73
sinuosa, Pierosoma tenuis, 206, 207
sinuosa, Planorbella tenuis, 206, 207
sinosum, Helisoma tenue, 206, 207

siskiyouensis, Calibasis acutifilosa, 41
siskiyouensis, Juga acutifilosa, 41
skinneri, Physa, 53, 159
smithi, Helisoma campanulatum, 202, 205
smithi, Neoplanorbis, 62, 171, 187
smithi, Planorbella campanulata, 202, 205
smithiana, Physella gyrina, 54, 163
solida optima, Parapholyx, 187
solida optima, Vorticifex, 187
solida, Parapholyx, 61, 174, 203
solida, Vorticifex, 61, 174, 203
Somatogyrus, 3, 23-25, 86, 88, 93-97, 99, 100, 195, 203
  alcoviensis, 23
  aldrichi, 93
  amnicoloides, 23, 93
  aureus, 23, 93, 99
  biangulatus, 23, 94
  constrictus, 23, 94
  coosaensis, 24, 88, 93, 97, 99, 203
  crassilabris, 23, 94
  crassus, 23, 94
  currierianus, 23, 94
  decipiens, 23, 94
  depressus, 23, 99
  excavatus, 23, 94, 95
  georgianus, 23, 95
  hendersoni, 23, 95
  hinkleyi, 23, 95
  humerosus, 23, 95
  integra, 24
  isogonus, 203
  nanus, 24, 95
  obtusus, 24, 95
  parvulus, 24, 95
  pennsylvanicus, 24, 88, 96
  pilsbryanus, 24, 96
  pumilus, 24, 96
  pygmaeus, 24, 96
  quadratus, 24, 96
  sargenti, 24, 96
  strengi, 24, 96
  substriatus, 24, 97
  tenax, 24, 86, 88, 99, 100, 203
  tennesseensis, 24, 97
  trothis, 24
  tryoni, 24, 195
  virginicus, 25, 88, 97, 195
  walkerianus, 24, 97
  wheeleri, 24, 97
sonomaensis, Bakerilymnaea, 50, 151, 152, 154, 205
sonomaensis, Fossaria, 50, 151, 152, 154, 205
sonomaensis, Lymnaea, 152
sordida, Lithasia obovata, 141
spelunca, Costatella, 56, 166
spelunca, Physella, 56, 166
sphaera, Neritina reclivata, 15, 67
spillmani, Gyrotoma, 133, 197, 198
Spilochlamys, 4, 28, 86, 104, 107-109
  conica, 28, 86, 104, 107, 108
  gravis, 28, 104, 108, 109
  turgida, 28, 104, 107
spinella, Elimia arachnoidea, 33, 124
spinosa, Io fluvialis, 136, 198
spinosus, Pyrgophorus, 21, 90, 91
spiralis, Antroselates, 19, 83, 89
Spirodon, 213
squalida, Costatella, 56, 166
squalida, Physella, 56, 166
stagnalis appressa, Lymnaea, 50, 148, 154
stagnalis, Lymnaea, 199
stagnalis sanctaemariae, Lymnaea, 50, 148, 154
Stagnicola, 5, 51, 52, 148, 149, 152, 155-158, 195, 199, 200, 204, 205
  alpenensis, 155
  apicina, 51
  arctica, 51, 156
  bonnevillensis, 51, 156
  caperata, 52, 158
  catascopium, 51, 156, 200
  cockerelli, 152
  contracta, 51, 156
  elodes, 51, 155, 156, 200
  elodes reflexa, 156
  elrodi, 51, 156
  elrodiana, 51, 155
  emarginata, 51, 157, 200
  emarginata canadensis, 157
  emarginata nashotahensis, 157
  emarginata serrata, 157, 200
  emarginatus, 195
  exilis, 51, 155
  gabbi, 52, 157
  hinkleyi, 52, 157
  idahoense, 52, 157
  impedita, 156
  jolietensis, 155
  kingi, 158
  laurentiana, 156
  mighelsi, 52, 149, 157
  montanensis, 52, 158
  nasoni, 158
  neopalustris, 51, 155
  newfoundlandensis, 156
  oronoensis, 52, 157
  palustris, 199, 200
  petoskeyensis, 52, 157
  pilsbryi, 52, 158
  traski, 51, 155
  utahensis, 52, 158
  walkeriana, 52, 158
  woodruffi, 52, 158
  wyomingensis, 156
stearnsiana, Elimia caelatura, 39, 130, 131

stearnsiana, Fontelicella, 26
Stenoglossa, 12
Stenophysa, 5, 57, 160, 167, 168, 200, 205
marmorata, 57, 167, 168
maugeriae, 57, 167, 168
Stimpsonia, 213
Stiobia, 4, 28, 98, 108, 113
nana, 28, 98, 108, 113
strengi, Somatogyrus, 24, 96
Strephobasis, 4, 47, 145, 146
corpulentum, 47, 145
curtum curtum, 47, 145
curtum roanense, 47, 146
walkeri, 47, 146
Strepoma, 213
Strepomatidae, 196
striatula, Elimia, 34, 124
striatum, Helisoma, 201
striatum, Helisoma anceps, 201
striatum, Pleurocera, 142
strigosa, Elimia, 34, 126
stygia, Amnicola, 29, 112
subcarinata, Lioplax, 18, 76, 80, 195
subcarinata occidentalis, Lioplax, 195
subcrenata disjecta, Pierosoma, 206
subcrenata disjecta, Planorbella, 206
subcrenata horni, Pierosoma, 206
subcrenata horni, Planorbella, 206
subcrenata perdisjuncta, Pierosoma, 206
subcrenata perdisjuncta, Planorbella, 206
subcrenata, Pierosoma, 207, 208
subcrenata, Pierosoma subcrenata, 205
subcrenata, Pierosoma trivolvis, 60, 176, 183, 206-208
subcrenata, Planorbella, 207, 208
subcrenata, Planorbella subcrenata, 205
subcrenata, Planorbella trivolvis, 60, 176, 183, 206-208
subcrenata plexata, Pierosoma, 206
subcrenata plexata, Planorbella, 206
subcrenata subcrenata, Pierosoma, 205
subcrenata subcrenata, Planorbella, 205
subcrenatum disjectum, Helisoma, 206
subcrenatum, Helisoma, 207, 208
subcrenatum, Helisoma subcrenatum, 205
subcrenatum, Helisoma trivolvis, 206-208
subcrenatum horni, Helisoma, 206
subcrenatum perdisjunctum, Helisoma, 206
subcrenatum plexatum, Helisoma, 206
subcrenatum subcrenatum, Helisoma, 205
subglobosa, Anculosa, 204
subglobosa, Birgella, 25, 97-100
subglobosa isogona, Birgella, 25
subglobosa, Leptoxis, 138, 204
subglobosa, Lithasia salebrosa, 44, 141, 199
subpurpureus, Viviparus, 17, 77, 78
subrotundata, Lanx, 53, 153
subsolidum, Campeloma, 79
substriatus, Somatogyrus, 24, 97
sulculosa, Lioplax, 18, 76, 195
symmetrica, Elimia, 39, 131
taeniata, Leptoxis, 43, 120, 121, 138, 204
Taenioglossa, 12
taitiana, Elimia, 37, 129
talquinensis, Lioplax, 18, 76, 80
tannum, Campeloma, 17, 79, 194
tantillus, Neoplanorbis, 62, 171, 186
Taphius, 213
tartarea, Fontigens, 30, 114
tazewelliana, Fossaria, 49, 146, 150
techella, Bakerilymnaea, 50, 152, 154
techella, Fossaria, 50, 152, 154
techella, Lymnaea, 152
Telescopella, 213
tenax, Somatogyrus, 24, 86, 88, 99, 100, 203
tenax, Walkerilla, 24, 86, 88, 99, 100, 203
tennesseensis, Somatogyrus, 24, 97
tentaculata, Bithynia, 81, 83
tentaculata, Bithynia tentaculata, 19
tentaculata magnalacustris, Bithynia, 19, 82, 83
tentaculata tentaculata, Bithynia, 19
tenue californiense, Helisoma, 206, 207
tenue, Helisoma, 207
tenue sinuosum, Helisoma, 206, 207
tenuipes, Littoridinops, 20, 82, 86, 87, 91, 98
tenuis californiensis, Pierosoma, 206, 207
tenuis californiensis, Planorbella, 206, 207
tenuis, Pierosoma, 60, 183, 207
tenuis, Planorbella, 60, 183, 207
tenuis sinuosa, Pierosoma, 206, 207
tenuis sinuosa, Planorbella, 206, 207
teres, Elimia, 34, 126
texana, Cochliopa, 31, 114
theiocrenetus, Aphaostracon, 20, 87, 91
Thiara, 4, 32, 116-118
granifera, 32, 116, 118
Thiaridae, 4, 31, 65, 116, 117
Thomsonia, 213
Tiara, 213
timidus, Elimia mutabilis, 37
tintinabulum, Leptoxis, 138
Torquis, 5, 57, 172, 177, 200
circumstriatus, 57, 172, 177
hornensis, 57, 172, 177, 200
parvus, 57, 172, 177
tortum, Elimia laqueata, 36
Toxoglossa, 12
traski, Helisoma, 206, 207
traski, Physella, 55, 161
traski, Pierosoma, 60, 206, 207
traski, Planorbella, 60, 206, 207
traski, Stagnicola, 51, 155
traskii, Helisoma, 207
tricarinata morph *bakeri, Valvata,* 16, 68
tricarinata morph *basalis, Valvata,* 16, 68

tricarinata morph *infracarinata, Valvata,* 16, 68
tricarinata morph *mediocarinata, Valvata,* 16, 68
tricarinata morph *perconfusa, Valvata,* 16, 70
tricarinata morph *simplex, Valvata,* 16, 70
tricarinata morph *tricarinata, Valvata,* 16, 68, 70
tricarinata morph *unicarinata, Valvata,* 16, 68
tricarinata, Valvata, 16, 67, 73
tricarinata, Valvata tricarinata, 16, 68, 70
trilineata, Leptoxis, 43, 122, 139
trilineata, Mudalia, 43, 122, 139
triticea, Physella propinqua nuttalli, 55, 163
trivolvis chautauquense, Helisoma, 208
trivolvis chautauquensis, Pierosoma, 208
trivolvis chautauquensis, Planorbella, 208
trivolvis fallax, Helisoma, 206, 208
trivolvis fallax, Pierosoma, 206, 208
trivolvis fallax, Planorbella, 206, 208
trivolvis, Helisoma, 206-208
trivolvis, Helisoma trivolvis, 206, 208
trivolvis holstonense, Helisoma, 208
trivolvis holstonensis, Pierosoma, 208
trivolvis holstonensis, Planorbella, 208
trivolvis lenta, Pierosoma, 60, 206, 208
trivolvis lenta, Planorbella, 60, 206, 208
trivolvis lentum, Helisoma, 206, 208
trivolvis macrostoma, Pierosoma, 206, 208
trivolvis macrostoma, Planorbella, 206, 208
trivolvis macrostomum, Helisoma, 206, 208
trivolvis marshalli, Helisoma, 206, 208
trivolvis marshalli, Pierosoma, 206, 208
trivolvis marshalli, Planorbella, 206, 208
trivolvis, Pierosoma, 188, 206-208
trivolvis, Pierosoma trivolvis, 60, 184, 206, 208
trivolvis, Planorbella, 188, 206-208
trivolvis, Planorbella trivolvis, 60, 184, 206, 208
trivolvis subcrenata, Pierosoma, 60, 176, 183, 206-208
trivolvis subcrenata, Planorbella, 60, 176, 183, 206-208
trivolvis subcrenatum, Helisoma, 206-208
trivolvis trivolvis, Helisoma, 206, 208
trivolvis trivolvis, Pierosoma, 60, 184, 206, 208
trivolvis trivolvis, Planorbella, 60, 184, 206, 208
trivolvis turgida, Pierosoma, 60, 206, 208
trivolvis turgida, Planorbella, 60, 206, 208
trivolvis turgidum, Helisoma, 206, 208
trochiformis, Pleurocera, 46, 145
troostiana, Elimia, 34, 126
Tropidina, 213
trothis, Somatogyrus, 24
truncata, Pierosoma, 60, 176, 184, 206
truncata, Planorbella, 60, 176, 184, 206
Truncatelloidea, 3, 65
truncatula, Fossaria, 49, 149, 150, 154, 205
truncatum, Helisoma, 206
tryoni, Aplexa elongata, 56, 167
tryoni, Somatogyrus, 24, 195
tryoni, Vorticifex, 203
Tryonia, 3, 21, 90-92, 203
cheatumi, 21, 90-92, 203
clathrata, 21, 90, 92
diaboli, 21, 90, 92
imitator, 21, 90
protea, 21, 90, 92
Trypanostoma, 213
tuberculata, Melanoides, 31, 116, 117
Tulotoma, 3, 16, 75, 77, 194
angulata, 77, 194
coosaensis, 194
magbimonilifera, 194
magnifica, 16, 75, 77, 194
turbiniformis, Fluminicola, 22, 93
turgida, Pierosoma trivolvis, 60, 206, 208
turgida, Planorbella trivolvis, 60, 206, 208
turgida, Spilochlamys, 28, 104, 107
turgidum, Helisoma trivolvis, 206, 208
Turridae, 12
turrita, Io fluvialis, 136, 198
turritella, Fontigens, 30, 114
Tylotoma, 213
ucheensis, Elimia, 32, 125
umbilicata, Clappia, 22, 88, 92, 93
umbilicata, Leptoxis, 43, 120, 143
umbilicatellus, Promenetus, 61, 173, 186, 202, 203
umbilicatus, Neoplanorbis, 62, 171, 187
umbilicatus, Planorbis, 202
umbilicatus, Promenetus, 202
umbonata, Goniobasis, 198
unakensis, Io fluvialis, 198
unciale hastatum, Pleurocera, 47, 145
unciale, Pleurocera, 47, 144
unciale, Pleurocera unciale, 47, 144, 145
unciale unciale, Pleurocera, 47, 144, 145
undulatum, Pleurocera canaliculatum, 46, 142, 143
unicarinata, Valvata tricarinata, 16, 68
unicarinatum, Helisoma, 201
unicarinatum, Helisoma anceps, 201
utahensis horati, Valvata, 16, 70
utahensis, Physella, 55, 161
utahensis, Stagnicola, 52, 158
utahensis utahensis, Valvata, 16, 68
utahensis, Valvata, 16, 73
utahensis, Valvata utahensis, 16, 68
Valvata, 3, 15, 16, 67-74, 196, 204
bicarinata, 69
bicarinata bicarinata, 15, 70
bicarinata morph *normalis,* 15, 69, 70
humeralis, 15, 69, 72
lewisi, 71
lewisi lewisi, 15, 72
lewisi morph *ontarioensis,* 15, 67, 70, 71
mergella, 15, 71, 72
micra, 196, 204
micra nugax, 196, 204
perdepressa, 16, 71

perdepressa perdepressa, 16, 72
perdepressa form *walkeri*, 16, 70
piscinalis form *obtusa*, 16, 71, 72
sincera form *danielsi*, 16, 72, 73
sincera nylanderi, 16, 72
sincera sincera, 16, 68, 72, 73
tricarinata, 16, 67, 73
tricarinata morph *bakeri*, 16, 68
tricarinata morph *basalis*, 16, 68
tricarinata morph *infracarinata*, 16, 68
tricarinata morph *mediocarinata*, 16, 68
tricarinata morph *perconfusa*, 16, 70
tricarinata morph *simplex*, 16, 70
tricarinata morph *tricarinata*, 16, 68, 70
tricarinata morph *unicarinata*, 16, 68
utahensis, 16, 73
utahensis morph *horati*, 16, 70
utahensis utahensis, 16, 68
virens, 16, 72, 73
winnebagoensis, 16, 70, 74
Valvatidae, 3, 15, 64, 67-69, 71, 73, 74
Valvatoidea, 3, 64
Vancleavia, 213
vancouverensis, Bakerilymnaea, 50, 152
vancouverensis, Fossaria, 50, 152
vanhyningi, Cincinnatia, 26, 101, 105
vanhyningiana, Elimia catenaria, 33, 124
vanuxemiana, Elimia, 38, 39, 131
varians, Elimia, 35
variata, Elimia, 38, 128
vaughani, Helisoma eucosmium, 58
Velletea, 213
nuttallii, 213
Velletia, 213
venusta, Physella propinqua nuttalli, 55, 163
Vermetoidea, 4, 65
vermilionense, Helisoma corpulentum, 205
vermilionensis, Pierosoma corpulenta, 60, 176, 205
vermilionensis, Planorbella corpulenta, 60, 176, 205
verrucosa, Angitrema, 45, 142, 199
verrucosa, Io fluvialis, 136, 198
verrucosa, Lithasia, 45, 142, 199
vestitum, Pleurocera, 47, 144, 145
viennaensis, Elimia boykiniana, 32, 116
vinosa, Physella, 55, 164
virens, Fluminicola, 22
virens, Valvata, 16, 72, 73
virgata anatina, Costatella, 56, 166
virgata anatina, Physella, 56, 166
virgata berendti, Costatella, 56, 166
virgata berendti, Physella, 56, 166
virgata concolor, Costatella, 56, 166
virgata concolor, Physella, 56, 166
virgata, Costatella virgata, 56, 166
virgata, Leptoxis, 43, 123, 140
virgata, Mudalia, 43, 123, 140
virgata parva, Costatella virgata, 56, 166
virgata parva, Physella virgata, 56, 166
virgata Physella virgata, 56, 166
virgata rhyssa, Costatella, 56, 166
virgata rhyssa, Physella, 56, 166
virgata virgata, Costatella, 56, 166
virgata virgata parva, Costatella, 56, 166
virgata virgata parva, Physella, 56, 166
virgata virgata, Physella, 56, 166
virginea, Physella, 55, 161
virginica, Elimia, 39, 130
virginicus, Somatogyrus, 25, 88, 97, 195
virginicus, Walkerilla, 25, 88, 97
viridulum, Pleurocera, 47, 145, 199
Vitta, 213
vittata, Leptoxis, 43, 121, 138
Vivipara, 213
Viviparella, 213
Viviparidae, 3, 16, 66, 74, 75, 77, 79, 80, 194
Viviparinae, 3, 16, 77
Viviparoidea, 65, 83
Viviparus, 3, 16, 17, 75-78
georgianus, 16, 77, 78
intertextus, 16, 76, 77
subpurpureus, 17, 77, 78
Vorticifex, 6, 61, 174, 186, 187, 203
effusa, 61, 174, 186, 203
solida, 61, 174, 203
solida optima, 187
tryoni, 203
wabashensis, Pyrgulopsis, 195
walkeri, Amnicola, 30, 111, 113, 114, 196
walkeri, Costatella integra integra, 56, 167
walkeri, Ferrissia, 63, 192, 193
walkeri, Gyrotoma, 41, 119, 135, 197, 198
walkeri, Lyogyrus, 30, 111, 113, 114
walkeri, Physella integra integra, 56, 167
walkeri, Pleurocera, 47, 146
walkeri, Strephobasis, 47, 146
walkeri, Valvata perdepressa, 16, 70
walkeriana, Stagnicola, 52, 158
walkerianus, Somatogyrus, 24, 97
Walkerilla, 3, 24, 25, 86, 88, 93, 97, 99, 100, 195, 203
coosaensis, 24, 88, 93, 97, 99, 203
tenax, 24, 86, 88, 99, 100, 203
virginicus, 25, 88, 97
Walkerola, 5, 53, 147, 159
klamathensis, 53, 147, 159
weberi, Fontigens, 31, 196
wekiwae, Cincinnatia, 26, 101, 103
wetherbyi, Notogillia, 27, 104, 107, 108
wheatleyi, Planorbis, 202
wheatleyi, Planorbula, 202
wheatleyi, Planorbula armigera, 61, 173, 185
wheeleri, Somatogyrus, 24, 97
whiteavesi, Helisoma multicostatum, 205
whiteavesi, Pierosoma corpulenta, 60, 176
whiteavesi, Pierosoma multicostata, 205

whiteavesi, Planorbella corpulenta, 60, 176
whiteavesi, Planorbella multicostata, 205
winkleyi mozleyi, Marstonia, 106
winnebagoensis, Valvata, 16, 70, 74
winslowi, Helisoma, 206-208
winslowi, Pierosoma, 206-208
winslowi, Pierosoma pilsbryi, 60
winslowi, Planorbella, 206-208
winslowi, Planorbella pilsbryi, 60
wisconsinense, Helisoma campanulatum, 202, 205
wisconsinensis, Planorbella campanulata, 202, 205
woodruffi, Stagnicola, 52, 158
wyomingensis, Stagnicola, 156
xynoelictus, Aphaostracon, 20, 87, 91
zionis, Petrophysa, 56, 167, 168
zionis, Physella, 56, 167, 168
Zygobranchia, 12

★ U.S. GOVERNMENT PRINTING OFFICE: 1982-559-092/3387