United States
Environmental Protection
Agency

Third External Review Draft of Air Quality Criteria for Particulate Matter (April, 2002):

Volume I

EPA/600/P-99/002aC
April 2002
Third External Review Draft

Air Quality Criteria for Particulate Matter

Volume I

National Center for Environmental Assessment-RTP Office
Office of Research and Development
U.S. Environmental Protection Agency
Research Triangle Park, NC

DISCLAIMER

This document is an external review draft for review purposes only and does not constitute U.S. Environmental Protection Agency policy. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

PREFACE

National Ambient Air Quality Standards (NAAQS) are promulgated by the United States Environmental Protection Agency (EPA) to meet requirements set forth in Sections 108 and 109 of the U.S. Clean Air Act (CAA). Sections 108 and 109 require the EPA Administrator (1) to list widespread air pollutants that reasonably may be expected to endanger public health or welfare; (2) to issue air quality criteria for them that assess the latest available scientific information on nature and effects of ambient exposure to them; (3) to set "primary" NAAQS to protect human health with adequate margin of safety and to set "secondary" NAAQS to protect against welfare effects (e.g., effects on vegetation, ecosystems, visibility, climate, manmade materials, etc.); and (5) to periodically (every 5 years) review and revise, as appropriate, the criteria and NAAQS for a given listed pollutant or class of pollutants.

The original NAAQS for particulate matter (PM), issued in 1971 as "total suspended particulate" (TSP) standards, were revised in 1987 to focus on protecting against human health effects associated with exposure to ambient PM less than 10 microns ($\leq 10\ \mu m$) that are capable of being deposited in thoracic (tracheobronchial and alveolar) portions of the lower respiratory tract. Later periodic reevaluation of newly available scientific information, as presented in the last previous version of this "Air Quality Criteria for Particulate Matter" document published in 1996, provided key scientific bases for PM NAAQS decisions published in July 1997. More specifically, the PM_{10} NAAQS set in 1987 (150 $\mu g/m^3$, 24-h; 50 $\mu g/m^3$, annual average) were retained in modified form and new standards (65 $\mu g/m^3$, 24-h; 15 $\mu g/m^3$, annual average) for particles $\leq 2.5\ \mu m$ ($PM_{2.5}$) were promulgated in July 1997.

This Third External Review Draft of revised Air Quality Criteria for Particulate Matter assesses new scientific information that has become available mainly between early 1996 through December 2001. The present draft is being released for public comment and review by the Clean Air Scientific Advisory Committee (CASAC) to obtain comments on the organization and structure of the document, the issues addressed, the approaches employed in assessing and interpreting the newly available information on PM exposures and effects, and the key findings and conclusions arrived at as a consequence of this assessment. Public comments and CASAC review recommendations will be taken into account in making any appropriate further revisions to this document for incorporation into a final draft. Evaluations contained in the present document will be drawn on to provide inputs to associated PM Staff Paper analyses prepared by EPA's Office of Air Quality Planning and Standards (OAQPS) to pose alternatives for consideration by the EPA Administrator with regard to proposal and, ultimately, promulgation of decisions on potential retention or revision of the current PM NAAQS.

Preparation of this document was coordinated by staff of EPA's National Center for Environmental Assessment in Research Triangle Park (NCEA-RTP). NCEA-RTP scientific staff, together with experts from other EPA/ORD laboratories and academia, contributed to writing of document chapters; and earlier drafts of this document were reviewed by experts from federal and state government agencies, academia, industry, and NGO's for use by EPA in support of decision making on potential public health and environmental risks of ambient PM. The document describes the nature, sources, distribution, measurement, and concentrations of PM in outdoor (ambient) and indoor environments. It also evaluates the latest data on human exposures to ambient PM and consequent health effects in exposed human populations (to support decision making regarding primary, health-related PM NAAQS). The document also evaluates ambient PM environmental effects on vegetation and ecosystems, visibility, and man-made materials, as well as atmospheric PM effects on climate change processes associated with alterations in atmospheric transmission of solar radiation or its reflectance from the Earth's surface or atmosphere (to support decision making on secondary PM NAAQS).

The NCEA of EPA acknowledges the contributions provided by authors, contributors, and reviewers and the diligence of its staff and contractors in the preparation of this document.

Air Quality Criteria for Particulate Matter

VOLUME I

EXECUTIVE SUMMARY E-1

1. INTRODUCTION 1-1

2. PHYSICS, CHEMISTRY, AND MEASUREMENT OF PARTICULATE MATTER 2-1

3. CONCENTRATIONS, SOURCES, AND EMISSIONS OF ATMOSPHERIC PARTICULATE MATTER 3-1
   - Appendix 3A: Spatial and Temporal Variability of the Nationwide AIRS $PM_{2.5}$ and $PM_{10-2.5}$ Data Sets 3A-1
   - Appendix 3B: Aerosol Composition Data from the Speciation Network 3B-1
   - Appendix 3C: Organic Composition of Particulate Matter 3C-1
   - Appendix 3D: Composition of Particulate Matter Source Emissions 3D-1

4. ENVIRONMENTAL EFFECTS OF PARTICULATE MATTER 4-1

5. HUMAN EXPOSURE TO PARTICULATE MATTER AND ITS CONSTITUENTS 5-1

Air Quality Criteria for Particulate Matter
(cont'd)

VOLUME II

6. DOSIMETRY OF PARTICULATE MATTER 6-1

7. TOXICOLOGY OF PARTICULATE MATTER IN HUMANS AND LABORATORY ANIMALS 7-1

8. EPIDEMIOLOGY OF HUMAN HEALTH EFFECTS FROM AMBIENT PARTICULATE MATTER8-1 8-1
   Appendix 8A: Short-Term PM Exposure-Mortality Studies: Summary Table 8A-1
   Appendix 8B: Particulate Matter-Morbidity Studies: Summary Tables 8B-1

9. INTEGRATIVE SYNTHESIS 9-1
   Appendix 9A: Key Quantitative Estimates of Relative Risk for Particulate Matter-Related Health Effects Based on Epidemiologic Studies of U.S. and Canadian Cities Assessed in the 1996 Particulate Matter Air Quality Criteria Document 9A-1

Table of Contents

Page

List of Tables . . . . . I-xii
List of Figures . . . . . I-xviii
Authors, Contributors, and Reviewers . . . . . I-xxx
U.S. Environmental Protection Agency Project Team for Development of
Air Quality Criteria for Particulate Matter . . . . . I-xli
Abbreviations and Acronyms . . . . . I-xliv

EXECUTIVE SUMMARY E-1

1. INTRODUCTION 1-1
1.1 LEGISLATIVE REQUIREMENTS 1-1
1.2 HISTORY OF PREVIOUS PM CRITERIA AND NAAQS REVIEWS 1-2
1.2.1 The 1997 PM NAAQS Revision 1-3
1.2.2 Coordinated Particulate Matter Research Program 1-5
1.3 CURRENT PM CRITERIA AND NAAQS REVIEW 1-10
1.3.1 Key Milestones 1-10
1.3.2 Methods and Procedures for Document Preparation 1-12
1.3.3 Approach 1-13
1.3.4 Key Human Health Issues of Concern 1-14
1.4 DOCUMENT ORGANIZATION AND CONTENT 1-17
REFERENCES 1-19

2. PHYSICS, CHEMISTRY, AND MEASUREMENT OF PARTICULATE MATTER . . . 2-1
2.1 PHYSICS AND CHEMISTRY OF PARTICULATE MATTER 2-2
2.1.1 Basic Concepts 2-2
2.1.2 Physical Properties and Processes 2-3
2.1.2.1 Definitions of Particle Diameter 2-3
2.1.2.2 Aerosol Size Distributions 2-4
2.1.2.3 Nuclei-Mode Particles 2-16
2.1.3 Chemistry of Atmospheric Particulate Matter 2-20
2.1.3.1 Chemical Composition and Its Dependence on Particle Size . . . 2-20
2.1.3.2 Primary and Secondary Particulate Matter 2-21
2.1.3.3 Particle-Vapor Partitioning 2-22
2.1.3.4 Atmospheric Lifetimes and Removal Processes 2-25
2.1.4 Summary 2-28
2.2 MEASUREMENT OF PARTICULATE MATTER 2-28
2.2.1 Particle Measurements of Interest 2-28
2.2.2 Issues in Measurement of Particulate Matter 2-31
2.2.2.1 Treatment of Semivolatile Components of Particulate Matter . . 2-32
2.2.2.2 Upper Cut Point 2-34
2.2.2.3 Cut Point for Separation of Fine-Mode and Coarse-Mode
Particulate Matter 2-35

Table of Contents
(cont'd)

Page

2.2.2.4 Treatment of Pressure, Temperature, and Relative Humidity . . . 2-38
2.2.2.5 Time Resolution 2-39
2.2.2.6 Accuracy and Precision 2-39
2.2.3 Measurement of Semivolatile Particulate Matter 2-41
2.2.3.1 Particulate Nitrates 2-43
2.2.3.2 Semivolatile Organic Compounds 2-46
2.2.3.3 Particle-Bound Water 2-59
2.2.4 U. S. Environmental Protection Agency Monitoring Methods 2-65
2.2.4.1 The Federal Reference Methods for Measurement of Equilibrated Mass for PM_{10}, $PM_{2.5}$, and $PM_{10-2.5}$ 2-65
2.2.5 Speciation Monitoring 2-78
2.2.6 Inorganic Elemental Analyses 2-80
2.2.6.1 Energy Dispersive X-ray Fluorescence (EDXRF). 2-81
2.2.6.2 Synchrotron Induced X-ray Fluorescence (S-XRF) 2-82
2.2.6.3 Proton Induced X-ray Emission (PIXE) 2-82
2.2.6.4 Proton Elastic Scattering Analysis (PESA) 2-83
2.2.6.5 Total Reflection X-ray Fluorescence (TRXRF) 2-83
2.2.6.6 Instrumental Neutron Activation Analysis (INAA) 2-84
2.2.6.7 Atomic Absorption Spectrophotometry (AAS) 2-85
2.2.6.8 Inductively Coupled Plasma with Atomic Emission Spectroscopy (ICP-AES) 2-86
2.2.6.9 Inductively Coupled Plasma with Mass Spectroscopy (ICP-MS) 2-86
2.2.6.10 Scanning Electron Microscopy (SEM) 2-87
2.2.7 Elemental and Organic Carbon in Particulate Matter 2-88
2.2.8 Ionic Species 2-98
2.2.9 Continuous Monitoring 2-98
2.2.9.1 Continuous Measurement of Mass 2-99
2.2.9.2 Continuous Measurement of Elemental and Organic Carbon 2-105
2.2.9.3 Continuous Measurements of Nitrate and Sulfate 2-107
2.2.9.4 Continuous Ion Chromatography of Water-Soluble Ions 2-109
2.2.9.5 Measurements of Individual Particles 2-109
2.2.9.6 Determination of Aerosol Surface Area in Real Time 2-111
2.2.9.7 Light Scattering 2-112
2.2.10 Low Flow Filter Samples for Multiday Collection of Particulate Matter 2-112
2.3 SUMMARY 2-114
2.3.1 Atmospheric Physics and Chemistry of Particles 2-114
2.3.2 Measurement of Atmospheric Particles 2-117
REFERENCES 2-121

Table of Contents
(cont'd)

Page

3. CONCENTRATIONS, SOURCES, AND EMISSIONS OF ATMOSPHERIC PARTICULATE MATTER 3-1
3.1 INTRODUCTION 3-1
3.2 PATTERNS AND TRENDS IN AMBIENT PM CONCENTRATIONS 3-2
3.2.1 Seasonal Variability in PM Concentrations 3-13
3.2.2 Diurnal (Circadian) Variability in PM Concentrations 3-27
3.2.3 Relations Among Particulate Matter in Different Size Fractions 3-29
3.2.4 Relations Between Mass and Chemical Component Concentrations 3-31
3.2.5 Spatial Variability in Particulate Matter and its Components 3-37
3.3 SOURCES OF PRIMARY AND SECONDARY PARTICULATE MATTER 3-56
3.3.1 Chemistry of Secondary PM Formation 3-59
3.3.2 The Long-Range Transport of Particulate Matter from Outside the United States 3-68
3.3.3 Source Contributions to Ambient PM Determined by Receptor Models 3-73
3.3.4 Emissions Estimates for Primary Particulate Matter, and Precursors to Secondary Particulate Matter (SO_2, NO_x, VOCs, and NH_3) in the United States 3-82
3.3.5 Uncertainties of Emissions Inventories 3-88
3.4 SUMMARY AND CONCLUSIONS 3-91
REFERENCES 3-95

APPENDIX 3A: Spatial and Temporal Variability of the Nationwide AIRS $PM_{2.5}$ and $PM_{10-2.5}$ Data Sets 3A-1
REFERENCES 3A-35

APPENDIX 3B: Aerosol Composition Data From The Speciation Network 3B-1
REFERENCES 3B-36

APPENDIX 3C: Organic Composition of Particulate Matter 3C-1
REFERENCES 3C-14

APPENDIX 3D: Composition of Particulate Matter Source Emissions 3D-1
REFERENCES 3D-26

4. ENVIRONMENTAL EFFECTS OF PARTICULATE MATTER 4-1
4.1 INTRODUCTION 4-1
4.2 IMPACTS ON VEGETATION AND ECOSYSTEMS 4-1
4.2.1 Particle Deposition 4-2
4.2.1.1 Wet Deposition 4-3

Table of Contents
(cont'd)

Page

4.2.1.2 Dry Deposition 4-5
4.2.1.3 Occult Deposition 4-30
4.2.1.4 Magnitude of Deposition 4-33
4.2.2 Effects on Vegetation and Ecosystems 4-44
4.2.2.1 Direct Effects of Particulate Matter on Individual Plant Species 4-46
4.2.2.2 Indirect Effects of Particulate Matter on Natural Ecosystems . . . 4-63
4.2.3 Urban Ecosystems 4-133
4.2.4 Ecosystem Goods and Services and Their Economic Valuation 4-137
4.3 EFFECTS ON VISIBILITY 4-141
4.3.1 Introduction 4-141
4.3.2 Factors Affecting Atmospheric Visibility 4-143
4.3.2.1 Optical Properties of the Atmosphere and Atmospheric Particles 4-143
4.3.2.2 Relative Humidity Effects on Particle Size and Light-Scattering Properties 4-149
4.3.3 Relationships Between Particles and Visibility 4-152
4.3.4 Photographic Modeling of Visibility Impairment 4-156
4.3.5 Visibility Monitoring Methods and Networks 4-158
4.3.6 Visibility Impairment: Trends and Current Conditions 4-161
4.4 EFFECTS ON MATERIALS 4-165
4.4.1 Corrosive Effects of Particles and Sulfur Dioxide on Man-Made Surfaces 4-166
4.4.1.1 Metals 4-166
4.4.1.2 Painted Finishes 4-168
4.4.1.3 Stone and Concrete 4-171
4.4.2 Soiling and Discoloration of Man-Made Surfaces 4-176
4.4.2.1 Stones and Concrete 4-177
4.4.2.2 Household and Industrial Paints 4-177
4.5 EFFECTS OF ATMOSPHERIC PARTICULATE MATTER ON GLOBAL CLIMATE CHANGE PROCESSES AND THEIR POTENTIAL HUMAN HEALTH AND ENVIRONMENTAL IMPACTS 4-178
4.5.1 Solar Ultraviolet Radiation Transmission Impacts on Human Health and the Environment: Atmospheric Particulate Matter Effects 4-180
4.5.1.1 Potential Effects of Increased Ultraviolet Radiation Transmission 4-180
4.5.1.2 Effects of Airborne Particles on Transmission of Solar Ultraviolet Radiation Through the Atmosphere 4-184
4.5.2 Global Warming Processes, Human Health and Environmental Impacts, and Roles of Atmospheric Particle 4-188

Table of Contents
(cont'd)

Page

4.5.2.1 Bases for Concern Regarding Global Warming and Climate Change 4-188
4.5.2.2 Airborne Particle Relationships to Global Warming and Climate Change 4-195
4.6 SUMMARY 4-200
4.6.1 Particulate Matter Effects on Vegetation and Ecosystems 4-200
4.6.2 Particulate Matter-Related Effects on Visibility 4-208
4.6.3 Particulate Matter-Related Effects on Materials 4-209
4.6.4 Effects of Atmospheric Particulate Matter on the Transmission of Solar Ultraviolet Radiation and Global Warming Processes 4-210
REFERENCES 4-212
APPENDIX 4A: Colloquial and Latin Names 4A-1

5. HUMAN EXPOSURE TO PARTICULATE MATTER AND ITS CONSTITUENTS 5-1
5.1 INTRODUCTION 5-1
5.1.1 Purpose 5-1
5.1.2 Particulate Matter Mass and Constituents 5-2
5.1.3 Relationship to Past Documents 5-3
5.2 STRUCTURE FOR THE CHAPTER 5-4
5.3 BASIC CONCEPTS OF EXPOSURE 5-5
5.3.1 Components of Exposure 5-5
5.3.2 Methods To Estimate Personal Exposure 5-7
5.3.2.1 Direct Measurement Methods 5-7
5 3.2.2 Indirect Methods (Modeling Methods) 5-8
5.3.2.3 Methods for Estimating Personal Exposure to Ambient Particulate Matter 5-15
5.4 SUMMARY OF PARTICULATE MATTER MASS DATA 5-20
5.4.1 Types of Particulate Matter Exposure Measurement Studies 5-20
5.4.2 Available Data 5-21
5.4.2.1 Personal Exposure Data 5-21
5.4.2.2 Microenvironmental Data 5-25
5.4.2.3 Reanalyses of Previously-Reported Particulate Matter Exposure Data 5-31
5.4.3 Factors Influencing and Key Findings on Particulate Matter Exposures 5-34
5.4.3.1 Relationship of Personal/Microenvironmental Particulate Matter with Ambient Particulate Matter 5-34
5.4.3.2 Factors That Affect Relationship between Personal Exposure and Ambient PM 5-53

Table of Contents
(cont'd)

Page

5.4.3.3 Impact of Ambient Sources on Exposures to Particulate Matter 5-70
5.4.3.4 Correlations of Particulate Matter with Other Pollutants 5-72
5.5 SUMMARY OF PARTICULATE MATTER CONSTITUENT DATA 5-76
5.5.1 Introduction 5-76
5.5.2 Monitoring Studies That Address Particulate Matter Constituents 5-77
5.5.3 Key Findings 5-77
5.5.3.1 Correlations of Personal and Indoor Concentrations with Ambient Concentrations of Particulate Matter Constituents 5-77
5.5.4 Factors Affecting Correlations Between Ambient Measurements and Personal or Microenvironmental Measurements of Particulate Matter Constituents 5-85
5.5.5 Limitations of Available Data 5-86
5.6 IMPLICATIONS OF USING AMBIENT PARTICULATE MATTER CONCENTRATIONS IN EPIDEMIOLOGIC STUDIES OF PARTICULATE MATTER HEALTH EFFECTS 5-87
5.6.1 Potential Sources of Error Resulting from Using Ambient Particulate Matter Concentrations in Epidemiologic Analyses 5-87
5.6.2 Associations Between Personal Exposures and Ambient Particulate Matter Concentrations 5-89
5.6.3 Role of Compositional Differences in Exposure Characterization for Epidemiology 5-93
5.6.4 Role of Spatial Variability in Exposure Characterization for Epidemiology 5-94
5.6.5 Analysis of Exposure Measurement Error Issues in Particulate Matter Epidemiology 5-95
5.6.5.1 Analysis of Exposure Measurement Errors in Time-Series Studies 5-96
5.6.5.2 Analysis of Exposure Measurement Errors in Long-Term Epidemiology Studies 5-99
5.6.5.3 Conclusions from Analysis of Exposure Measurement Errors on Particulate Matter Epidemiology 5-101
5.7 SUMMARY OF KEY FINDINGS AND LIMITATIONS 5-102
REFERENCES 5-110

List of Tables

Number Page

1-1 Key Scientific Uncertainties Related to the Source-To-Response Framework 1-8

1-2 Research Topics and Questions Recommended by National Research Council to be Addressed by Expanded Particulate Matter Research Program 1-9

1-3 Schedule for Development of the Current Revised Particulate Matter Air Quality Criteria Document 1-11

2-1 Comparison of Ambient Particles Fine Mode and Coarse Mode 2-29

2-2 Particulate Matter Components/Parameters of Interest for Health, Ecological, or Radiative Effects; for Source Category Apportionment Studies; or for Air Quality Modeling Evaluation Studies 2-30

2-3 Summary of Sensitivity Studies of WINS Impactor Performance 2-70

2-4 $PM_{2.5}$ Samplers Currently Designated as FRMs for $PM_{2.5}$ Mass Concentrations . . . 2-71

3-1 Distribution of Ratios of $PM_{2.5}$ to Pm_{10} and Correlations Between $PM_{2.5}$ and PM_{10}, $PM_{2.5}$ and $PM_{10-2.5}$, and $PM_{10-2.5}$ and PM_{10} Found at Collocated Monitoring Sites in Seven Aerosol Characteristic (EPA/HEI) Regions in 1999 3-30

3-2 Concentrations (in ng/m^3) of $PM_{2.5}$, $PM_{10-2.5}$, and Selected Elements (ng/m^3) in the $PM_{2.5}$ and $PM_{10-2.5}$ Size Ranges With Standard Deviations and Correlations Between Elements and $PM_{2.5}$ Mass in Philadelphia, PA 3-32

3-3 Concentrations (in ng/m^3) of $PM_{2.5}$, $PM_{10-2.5,}$ and Selected Elements in the $PM_{2.5}$ and $PM_{10-2.5}$ Size Range With Standard Deviations and Correlations (r) Between Elements and $PM_{2.5}$ and $PM_{10-2.5}$ Mass in Phoenix, AZ 3-33

3-4 Measures of the Spatial Variability of $PM_{2.5}$ Concentrations Within Selected Metropolitan Statistical Areas 3-38

3-5 Measures of the Spatial Variability of $PM_{10-2.5}$ Concentrations Within Selected Metropolitan Statistical Areas 3-46

3-6 Correlation Coefficients for Spatial Variation of $PM_{2.5}$ Mass and Different Sources for Pairs of Sampling Sites in the South Coast Air Basin (1986) 3-53

3-7 Correlation Coefficients for Spatial Variation of $PM_{2.5}$ Mass and Different Components for Pairs of Sampling Sites in Philadelphia (1994) 3-54

List of Tables
(cont'd)

Number Page

3-8 Constituents of Atmospheric Particles and Their Major Sources 3-57

3-9 Receptor Model Source Contributions to $PM_{2.5}$ 3-77

3-10 Receptor Model Source Contributions to PM_{10} 3-78

3-11 Emissions of Primary $PM_{2.5}$ by Various Sources in 1999 3-84

3-12 Emissions of Precursors to Secondary $PM_{2.5}$ Formation by Various Sources in 1999 3-85

3B-1 Summary Statistics for $PM_{2.5}$ Concentrations During February Through June 2000 Obtained by Collocated Federal Reference Method Samplers (in $\mu g/m^3$) 3B-2

3B-2 Summary Statistics for the Speciation Samplers at Bismarck, ND (in $\mu g/m^3$) 3B-3

3B-3 Summary Statistics for the Speciation Samplers at Boston, MA (in $\mu g/m^3$) 3B-6

3B-4 Summary Statistics for the Speciation Samplers at Bronx Botanical Garden, NY (in $\mu g/m^3$) 3B-8

3B-5 Summary Statistics for the Speciation Samplers at Chicago, IL (in $\mu g/m^3$) 3B-11

3B-6 Summary Statistics for the Speciation Samplers at Fresno, CA (in $\mu g/m^3$) 3B-14

3B-7 Summary Statistics for the Speciation Samplers at Houston, TX (in $\mu g/m^3$) 3B-16

3B-8 Summary Statistics for the Speciation Samplers at Lewis, FL (in $\mu g/m^3$) 3B-19

3B-9 Summary Statistics for the Speciation Samplers at Philadelphia, PA (in $\mu g/m^3$) . . 3B-21

3B-10 Summary Statistics for the Speciation Samplers at Phoenix, AZ (in $\mu g/m^3$) 3B-24

3B-11 Summary Statistics for the Speciation Samplers at Portland, OR (in $\mu g/m^3$) 3B-26

3B-12 Summary Statistics for the Speciation Samplers at Salt Lake City, UT (in $\mu g/m^3$) 3B-28

3B-13 Summary Statistics for the Speciation Samplers at St. Louis, MO (in $\mu g/m^3$) 3B-30

3B-14 Summary Statistics for the Speciation Samplers at Seattle, WA (in $\mu g/m^3$) 3B-33

List of Tables
(cont'd)

Number		Page
3C-1	Particulate Organic and Elemental Carbon Concentrations (in μg C/m^3) Based on Studies Published After 1995	3C-2
3C-2	Particulate Organic Compound Concentrations (in ng C/m^3) Based on Studies Published after 1990 at Selected Sites	3C-5
3D-1	Average Abundances of Major Elements in Soil and Crustal Rock	3D-2
3D-2	Summary of Particle-Phase Organic Compounds Present in Fine Particle Road Dust Sample	3D-6
3D-3	Composition of Fine Particles Released by Various Stationary Sources in the Philadelphia Area	3D-7
3D-4a	Organic and Elemental Carbon Fractions of Diesel and Gasoline Engine Particulate Matter Exhaust	3D-13
3D-4b	Contribution of Organic Carbon to Particulate Matter Carbon Emissions in Motor Vehicle Exhaust Collected from Vehicles Operated on Chassis Dynamometers	3D-14
3D-5	Emission Rates (mg/mi) for Constituents of Particulate Matter from Gasoline and Diesel Vehicles	3D-15
3D-6	Summary of Particle-Phase Organic Compounds Emitted from Motor Vehicles	3D-17
3D-7	Mass Emissions, Organic Carbon, and Elemental Carbon Emissions from Residential Combustion of Wood	3D-20
3D-8	Summary of Particle-Phase Organic Compounds Emitted from the Combustion of Wood in Fireplaces	3D-21
3D-9	Mean Aerosol Composition at Tropical Site (Sriwijaya University, Sumatra, Indonesia) Affected Heavily by Biomass Burning Emissions	3D-23
4-1	Relative Importance of Wet, Dry, Particulate, and Total Deposition to Three Forest Sites	4-4
4-2	Key Determinants of Dry Particulate Deposition to Vegetation	4-7

List of Tables
(cont'd)

Number Page

4-3 Reported Mean Deposition Velocities (V_d) for Sulfate, Chlorine, Nitrate, and Ammonium and Ion-Containing Particles 4-28

4-4 Representative Empirical Measurements of Deposition Velocity (V_d) for Particulate Deposition 4-29

4-5 Reported Mean Deposition Velocities for Potassium, Sodium, Calcium, and Magnesium Base Cation Containing Particles 4-29

4-6 Relative Magnitudes of Wet, Dry, and Occult Deposition of Nitrates (NO_3^-) and Sulfates (SO_4^{-2}) to Three Forest Sites Subject to Similar Gas- and Liquid-Phase Pollutant Concentrations During Spring and Summer 4-33

4-7 Mean Annual Nitrogen Deposition (equivalents/ha/yr) from Fine and Coarse Particles Compared to Total Nitrogen Deposition from all Sources to a Variety of Forest Ecosystems 4-36

4-8 Mean Annual Sulfate Deposition (equivalents/ha/yr) from Fine and Coarse Particles Compared to Total Sulfur Deposition from all Sources to a Variety of Forest Ecosystems 4-37

4-9 Mean Annual Base Cation Deposition (equivalents/ha/yr) from Fine and Coarse Particles Compared to Total Base Cation Deposition from all Sources to a Variety of Forest Ecosystems 4-38

4-10 Mean (±SE) Particle Size, Deposition Rates, and Derived Deposition Velocities (V_d) for Heavy Metal Deposition to the Upper Canopy (Inert Plates or Leaves) of an Upland Oak Forest 4-41

4-11 Total Heavy Metal Deposition to Temperate Latitude Forests 4-42

4-12 Annual Bulk Deposition Input of Seven Heavy Metals to the Hubbard Brook Experimental Forest (43° 56' N Latitude, 71° 45' W Longitude), White Mountain National Forest, NH, for the Period 1975 to 1991 (grams per hectare) 4-43

4-13 Ecosystem Services 4-65

4-14 Ecosystem Functions Impacted by Air Pollution Effects on Temperate Forest Ecosystems 4-71

4-15 Types of Plant Responses to Ultraviolet-B Radiation 4-84

List of Tables
(cont'd)

Number Page

4-16 Nitrogen-Saturated Forests in North America, Including Estimated N Inputs and Outputs 4-88

4-17 Primary Goods and Services Provided by Ecosystems 4-138

4-18 Corrosive Effects of Particulate Matter and Sulfur Dioxide on Metals 4-169

4-19 Corrosive Effects of Particulate Matter and Sulfur Dioxide on Stone 4-172

4-20 Examples of Impacts Resulting from Projected Changes in Extreme Climate Events 4-192

5-1 Classes of PM Exposure and Concentration Definitions 5-6

5-2 Activity Pattern Studies Included in the Consolidated Human Activity Database 5-11

5-3 Personal Exposure Models for PM 5-13

5-4 Summary of Recent Personal Exposure Studies 5-22

5-5 Summary of Recent Microenvironmental Measurement Studies 5-26

5-6 Papers Interpreting PM Exposure Studies 5-32

5-7 Personal Monitoring Studies for PM: Measured Concentrations and Correlation Coefficients 5-37

5-8 Mean Concentration for PM Mass Reported for the Baltimore (Williams et al., 2000a,b,c) and the Fresno (Evans et al., 2000) Studies 5-44

5-9 Daily-Average Correlation Coefficients Reported for the Baltimore (Williams et al., 2000a,b,c) and the Fresno (Evans et al., 2000) Studies 5-45

5-10 Regression Analysis Reported for Indoor/Outdoor Relationships for the Baltimore (Williams et al., 2000a,b,c) and the Fresno (Evans et al., 2000) Studies 5-45

5-11 Volume Mean Diameter and Maximum $PM_{2.5}$ Concentrations of Indoor Particle Sources 5-67

5-12 Correlations Between Personal $PM_{2.5}$ and Ambient Pollutant Concentrations 5-74

List of Tables
(cont'd)

Number Page

5-13 Correlations Between Hourly Personal $PM_{2.5}$ and Gaseous Pollutants 5-76

5-14 Studies That Have Measured Particulate Matter Constituents in Personal Exposure Samples 5-79

5-15 Studies That Have Measured Particulate Matter Constituents in Microenvironmental Samples 5-80

5-16 Summary Statistics for Personal, Indoor, and Outdoor Concentrations of Selected Aerosol Components in Two Pennsylvania Communities 5-82

5-17 Statistical Correlation of Outdoor (x) Versus Indoor (y) Concentration for Measured Species 5-84

List of Figures

Number Page

1-1 A general framework for integrating particulate-matter research 1-7

2-1 Number of particles as a function of particle diameter: (a) number concentrations are shown on a logarithmic scale to display the wide range by site and size; (b) number concentrations for the average urban distribution are shown on a linear scale for which the area under any part of the curve is proportional to particle number in that size range 2-6

2-2 Particle volume distribution as a function of particle diameter: (a) for the averaged rural and urban-influenced rural number distributions shown in Figure 2-1 and a distribution from south central New Mexico, and (b) for the averaged urban and freeway-influenced urban number distributions shown in Figure 2-1 2-7

2-3 Distribution of coarse (c), accumulation (a), and nuclei or ultrafine (n), mode particles by three characteristics, a) number (N), b) surface area (S) and c) volume (V) for the grand average continental size distribution 2-8

2-4 Volume size distribution, measured in traffic, showing fine-mode and coarse-mode particles and the nuclei and accumulation modes within the fine particle mode 2-9

2-5 An idealized distribution of ambient particulate matter showing fine-mode particles and coarse-mode particles and the fractions collected by size-selective samplers 2-11

2-6 Specified particle penetration (size-cut curves) through an ideal (no-particle-loss) inlet for five different size-selective sampling criteria 2-13

2-7 Comparison of penetration curves for two PM_{10} beta gauge samplers using cyclone inlets 2-15

2-8 Particle growth curves showing fully reversible hygroscopic growth of sulfuric acid (H_2SO_4) particles, deliquescent growth of ammonium sulfate [$(NH_4)_2 SO_4$] particles at about 80% relative humidity (RH), hygroscopic growth of ammonium sulfate solution droplets at RH greater than 80%, and hysteresis (the droplet remains supersaturated as the RH decreases below 80%) until the crystallization point is reached 2-25

2-9 Theoretical predictions and experimental measurements of growth of NH_4HSO_4 and particles at relative humidity between 95 and 100% 2-26

List of Figures
(cont'd)

Number		Page
2-10	Schematic showing major nonvolatile and semivolatile components of $PM_{2.5}$	2-33
2-11	Particulate matter concentrations in Spokane, WA, during the August 30, 1996 dust storm	2-37
2-12	Amount of ammonium nitrate volatilized from Teflon filters, expressed as a percentage of the measured $PM_{2.5}$ mass, for the SCAQS and CalTech studies, for spring and fall sampling periods	2-44
2-13	Aerosol water content expressed as a mass percentage, as a function of relative humidity	2-63
2-14	Schematic diagram of the sample collection portion of the $PM_{2.5}$ FRM sampler	2-67
2-15	Schematic view of the final design of the WINS	2-68
2-16	Evaluation of the final version of the WINS	2-69
2-17	Schematic diagram showing the principle of virtual impaction	2-77
2-18	This thermogram, for a sample containing rock dust (a carbonate source) and diesel exhaust, shows three traces that correspond to temperature, filter transmittance, and FID detector response	2-93
2-19	Comparison of mass measurements with collocated RAMS (real-time data), PC-BOSS (1-h data), FRM $PM_{2.5}$ sampler (average of 24-h data), and a conventional TEOM monitor (real-time data)	2-102
2-20	Size distribution of particles divided by chemical classification into organic, marine, and crustal	2-110
3-1a	1999-2000 county-wide average annual mean PM_{10} concentrations ($\mu g/m^3$)	3-6
3-1b	1999-2000 highest county-wide 98th percentile 24-h average PM_{10} concentrations ($\mu g/m^3$)	3-6
3-2	Nationwide trend in ambient PM_{10} concentration from 1989 through 1998	3-8
3-3	Trend in PM_{10} annual mean concentrations by EPA region, 1989-1998	3-8

List of Figures
(cont'd)

Number | Page

3-4a 1999-2000 county-wide average annual mean $PM_{2.5}$ concentrations ($\mu g/m^3$) 3-9

3-4b 1999-2000 highest county-wide 98th percentile 24-h average $PM_{2.5}$ concentrations ($\mu g/m^3$) 3-9

3-5 Collection of annual distribution of 24-h average $PM_{2.5}$ concentrations observed in U.S. and Canadian health studies conducted during the 1980's and early 1990's 3-10

3-6a 1999-2000 estimated county-wide average annual mean $PM_{10-2.5}$ concentrations ($\mu g/m^3$) 3-14

3-6b 1999-2000 estimated county-wide highest 98th percentile 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) 3-14

3-7a,b Quarterly distribution of 24-h average $PM_{2.5}$ concentrations for selected monitors in the (a) Columbia, SC; (b) Detroit, MI; (c) Chicago, IL; and (d) Los Angeles, CA MSAs. Values for the lowest, lower quartile, median, upper quartile and highest concentrations are shown in the figures 3-16

3-7c,d Quarterly distribution of 24-h average $PM_{2.5}$ concentrations for selected monitors in the (a) Columbia, SC; (b) Detroit, MI; (c) Chicago, IL; and (d) Los Angeles, CA MSAs. Values for the lowest, lower quartile, median, upper quartile and highest concentrations are shown in the figures 3-17

3-8 Concentrations of $PM_{2.5}$ and PM_{10} measured in the four MAACS cities 3-19

3-9a,b Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations for selected sites in the (a) Columbia, SC; (b) Detroit, MI; and (c) Los Angeles, CA metropolitan statistical areas 3-20

3-9c Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations for selected sites in the (a) Columbia, SC; (b) Detroit, MI; and (c) Los Angeles, CA metropolitan statistical areas 3-21

3-10 Frequency distribution of 24-h average $PM_{2.5}$ concentrations measured at the monitoring site at the Presbyterian home in southwestern Philadelphia from 1992 to 1995 3-22

3-11 Concentrations of 24-h average $PM_{2.5}$ measured at the EPA site in Phoenix, AZ from 1995 to 1997 3-24

List of Figures
(cont'd)

Number Page

3-12 Frequency distribution of 24-h average $PM_{2.5}$ concentrations measured at the EPA site in Phoenix, AZ from 1995 to 1997 3-24

3-13 Frequency distribution of 24-h average $PM_{2.5}$ measurements obtained from all California Air Resources Board dichotomous sampler sites from 1989 to 1998 . . . 3-25

3-14 Frequency distribution of 24-h average $PM_{10-2.5}$ concentrations obtained from all California Air Resource Board Dichotomous sampler sites from 1989 to 1998 . . . 3-25

3-15 Concentrations of 24-h average $PM_{2.5}$ measured at the Riverside-Rubidoux site from 1989 to 1998 3-26

3-16 Frequency distribution of 24-h average $PM_{2.5}$ concentrations measured at the Riverside-Rubidoux site from 1989 to 1994 3-26

3-17 Intraday variability of hourly average $PM_{2.5}$ concentrations across the United States 3-28

3-18 Occurrence of differences between pairs of sites in three metropolitan statistical areas 3-45

3-19 Intersite correlation coefficients for $PM_{2.5}$, PM_{10}, and $PM_{10-2.5}$ 3-49

3-20 $PM_{2.5}$ chemical components in downtown Los Angeles and Burbank (1986) have similar characteristics 3-51

3-21 Concentrations of $PM_{2.5}$ chemical components in Rubidoux and downtown Los Angeles (1986) 3-52

3-22 Monthly average Saharan dust components of the aerosol sampled in Miami, FL, from 1974 to 1996 3-70

3-23 $PM_{2.5}$ and PM_{10} concentrations measured at Chilliwack Airport, located in northwestern Washington State, just before and during the Asian desert dust episode of April and May 1998 3-71

3-24 Time series of 24-h average PM_{10} concentrations observed in the Rio Grande Valley during May 1998 3-72

3-25 PM_{10} concentrations observed in St. Louis, MO, during May 1998 3-72

List of Figures
(cont'd)

Number Page

3A-1 Philadelphia, PA-NJ MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-3

3A-2 Washington, DC MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-4

3A-3 Norfolk, VA MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-5

3A-4 Columbia, SC MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-6

3A-5 Atlanta, GA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-7

3A-6 Birmingham, AL MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-8

3A-7 Tampa, FL MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-9

3A-8 Cleveland, OH MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-10

List of Figures
(cont'd)

Number Page

3A-9 Pittsburgh, PA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-11

3A-10 Steubenville, OH-Weirton, WV MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-12

3A-11 Detroit MI MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-13

3A-12 Grand Rapids, MI MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-14

3A-13 Milwaukee, WI MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-15

3A-14 Chicago, IL MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-16

3A-15 Gary, IN MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-17

3A-16 Louisville, KY MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-18

List of Figures
(cont'd)

Number Page

3A-17 St. Louis, MO MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-19

3A-18 Baton Rouge, LA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-20

3A-19 Kansas City, KS-MO MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-21

3A-20 Dallas, TX MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-22

3A-21 Boise, ID MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-23

3A-22 Salt Lake City, UT MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-24

3A-23 Seattle, WA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-25

3A-24 Portland, OR MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-26

List of Figures
(cont'd)

Number | Page

3A-25 Los Angeles-Long Beach, CA MSA. (a) Locations of sampling sites by AIRS ID; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-27

3A-26 Riverside-San Bernadino, CA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-28

3A-27 San Diego, CA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (c) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given 3A-29

3A-28 Columbia, SC MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations; (c) Intersite correlation coefficients and number of measurements 3A-30

3A-29 Detroit, MI MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations; (c) Intersite correlation coefficients and number of measurements 3A-31

3A-30 Cleveland, OH MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations; (c) Intersite correlation coefficients and number of measurements 3A-32

3A-31 Steubenville, OH-Weirton, WV MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations; (c) Intersite correlation coefficients and number of measurements 3A-33

3A-32 St. Louis, MO-IL MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations; (c) Intersite correlation coefficients and number of measurements 3A-34

3D-1 Size distribution of particles generated in a laboratory resuspension chamber . . . 3D-3

3D-2 Size distribution of California source emissions, 1986 . . . . . 3D-4

3D-3 Chemical abundances for $PM_{2.5}$ emissions from paved road dust in Denver, CO 3D-5

List of Figures
(cont'd)

Number Page

3D-4 Chemical abundances for $PM_{2.5}$ emissions from wood burning in Denver, CO 3D-19

4-1 A simplified resistance catena representing the factors controlling deposition of particles to the surface 4-8

4-2 The relationship between deposition velocity of selected particulate materials and the distribution of the material between the coarse- and fine-aerosol fractions 4-9

4-3 The relationship between particle diameter and deposition velocity for particles 4-10

4-4 Vertical stratification of diverse, chemically speciated particles in a mixed oak forest 4-24

4-5 The relationship between particle size and concentration below a spruce canopy with wind velocity at a height of 16.8 m equaling 5 m s^{-1} 4-25

4-6 Mean (±SE) percent of total nitrogen, sulfur, or base cation deposition contributed by fine plus coarse particles 4-39

4-7 Effects of environmental stress on forest trees are presented on a hierarchial scale for the leaf, branch, tree, and stand levels of organization 4-67

4-8 Nitrogen cycle (dotted lines indicate processes altered by nitrogen saturation) 4-89

4-9 Diagrammatic overview of excess nitrogen (N) in North America 4-91

4-10 Schematic of sources and sinks of hydrogen ions in a forest 4-104

4-11 Calcium deposition in >2-μm particles, <2-μm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites 4-113

4-12 Magnesium deposition in >2-μm particles, <2-μm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites 4-114

4-13 Potassium deposition in >2-μm particles, <2-μm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites 4-115

4-14 Base cation deposition in >2-μm particles, <2-μm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites 4-116

List of Figures
(cont'd)

Number Page

4-15 Total cation leaching (total height of bar) balanced by sulfate and nitrate estimated from particulate deposition (assuming no ecosystem retention, particulate sulfur and nitrogen) and by other sources (both deposition and internal) of sulfate and nitrate (other sulfur and nitrogen sources) and by other anions in the Integrated Forest Study sites 4-117

4-16 Soil exchangeable Ca^{+2} pools and net annual export of Ca^{+2} (deposition minus leaching times 25 years) in the Integrated Forest Study sites 4-118

4-17 Soil exchangeable Mg^{+2} pools and net annual export of Mg^{+2} (deposition minus leaching times 25 years) in the Integrated Forest Study sites 4-118

4-18 Soil exchangeable K^{2+} pools and net annual export of K^{2+} (deposition minus leaching times 25 years) in the Integrated Forest Study sites 4-119

4-19a Simulated soil solution mineral acid anions and base cations in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition 4-122

4-19b Simulated soil solution mineral acid anions and base cations in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition 4-123

4-20a Simulated soil solution A1 and soil base saturation in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition 4-124

4-20b Simulated soil solution A1 and soil base saturation in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition 4-125

4-21a Simulated soil solution mineral acid anions and base cations in the Coweeta site with no change, 50% N and S deposition, and 50% base cation deposition 4-127

4-21b Simulated soil solution mineral acid anions and base cations in the Coweeta site with no change, 50% N and S deposition, and 50% base cation deposition 4-128

4-22 Relationship of plant nutrients and trace metals with vegetation 4-132

4-23 Linkages among various ecosystem goods and services (food, water, biodiversity, forest products) and other driving forces (climate change) 4-139

List of Figures
(cont'd)

Number Page

4-24 Light reflected from a target toward an observer 4-145

4-25 Light-scattering efficiency factor (per cross sectional area), Q, for a homogeneous sphere with an index of refraction of 1.50 as a function of the size parameter, $\alpha = \pi D/\lambda$ 4-148

4-26 Particle growth curve as a function of relative humidity showing deliquescent growth of ammonium sulfate [$(NH_4)_2 SO_4$] particles at the deliquescent point (A, about 80% relative humidity [RH]), reversible hygroscopic growth of ammonium sulfate solution droplets at RH greater than 80%, and hysteresis (the droplet remains supersaturated as the RH decreases below 80%) until the crystallization point (B, about 38% RH) is reached 4-150

4-27 Comparison of extinction (Mm^{-1}) and visual range (km) 4-154

4-28 Relative humidity adjustment factor, f(RH), for ammonium sulfate as a function of relative humidity 4-156

4-29a Aggregate visibility trends (in deciviews) for 10 eastern Class 1 areas 4-163

4-29b Aggregate visibility trends (in deciviews) for 26 western Class 1 areas 4-163

4-30a Eastern class I area aggregate trends in aerosol light extinction on the 20% haziest days, including trends by major aerosol component 4-164

4-30b Western class I area aggregate trends in aerosol light extinction on the 20% haziest days, including trends by major aerosol component 4-164

4-31 Processes involved in stratospheric ozone depletion because of man's production of CFCs, halons, and other trace gases are shown to the left 4-181

4-32 Estimated global mean radiative forcing exerted by gas and various particle phase species for the year 2000, relative to 1750 4-199

5-1 Comparison of correlation coefficients for longitudinal analyses of personal exposure versus ambient concentrations for individual subjects for $PM_{2.5}$ and sulfate 5-48

5-2 Personal exposure versus ambient concentrations for $PM_{2.5}$ and sulfate 5-50

List of Figures
(cont'd)

Number Page

5-3 Regression analyses of aspects of daytime personal exposure to PM_{10} estimated using data from the PTEAM study 5-52

5-4 Air-exchange rates measured in homes throughout the United States 5-59

5-5 Box plots of hourly air-exchange rates stratified by season in Boston, MA, during 1998 5-60

5-6 Geometric mean infiltration factor (indoor/outdoor ratio) for hourly nighttime, nonsource data for two seasons 5-60

5-7 Comparison of deposition rates from this study with literature values (adapted from Abt et al., 2000b) 5-62

5-8 Penetration efficiencies and deposition rates from models of nightly average data 5-63

5-9 Mean hourly indoor/outdoor particle concentration ratio from an unoccupied residence in Fresno, CA, during spring 1999 5-68

5-10 Personal versus outdoor $SO_4^=$ in State College, PA 5-83

5-11 Plots of nonambient exposure to PM_{10}, (a) daytime individual values from PTEAM data and (b) daily-average values from THEES data 5-92

Authors, Contributors, and Reviewers

EXECUTIVE SUMMARY

Principal Authors

Dr. William E. Wilson—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lester D. Grant—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Beverly Comfort—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. William Ewald—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. J.H.B. Garner—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Dennis J. Kotchmar—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Allan Marcus—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Joseph P. Pinto—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. James A. Raub—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

CHAPTER 1. INTRODUCTION

Principal Authors

Dr. Lester D. Grant—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Dennis J. Kotchmar—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Authors, Contributors, and Reviewers
(cont'd)

CHAPTER 2. PHYSICS, CHEMISTRY, AND MEASUREMENT OF PARTICULATE MATTER

Principal Authors

Dr. William Wilson—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Candis S. Claiborn—Washington State University, Laboratory for Atmospheric Research,
Department of Civil and Environmental Engineering, P.O. Box 642910, Pullman, WA 99164

Contributing Authors

Dr. Barbara J. Turpin, The State University of New Jersey, Rutgers, Department of
Environmental Sciences and Rutgers Cooperative Extension, New Brunswick, NJ 08901-8551

Dr. James J. Schauer, University of Wisconsin, College of Engineering, Department of Civil and
Environmental Engineering, Madison, WI 53706

Dr. Judith C. Chow, Desert Research Institute, 2215 Raggio Parkway, Reno, NV 89512

Dr. John G. Watson, Desert Research Institute, 2215 Raggio Parkway, Reno, NV 89512

Dr. Tom Cahill, University of California, Davis, One Shields Ave., Davis, CA 95616

Dr. Timothy Buckley—Johns Hopkins University, Department of Environmental Health
Sciences, 615 North Wolfe Street,, Baltimore, MD 21205

Ms. Lee Byrd—Office of Air Quality Planning and Standards (MD-14),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lyle Chinkin—Sonoma Technology, 1360 Redwood Way, Suite C,
Petaluma, CA 94549

Dr. Steven Colome—Integrated Environmental Services, 5319 University Drive, #430,
Irvine, CA 92612

Dr. Delbert Eatough—Brigham Young University, E 114 BNSN,
Department of Chemistry and Biochemistry, Provo, UT 84602

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers
(cont'd)

Dr. Edward O. Edney—National Exposure Research Laboratory (MD-84)
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. William Ewald—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Sheldon Friedlander—University of California at Los Angeles, Department of Chemical Engineering, 5531 Boelter Hall, Los Angeles, CA 90095

Dr. Judith Graham—National Exposure Research Laboratory (MD-75),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lynn Hildemann—Stanford University, Civil and Environmental Engineering Department, Stanford, CA 94305

Mr. Jim Homolya—Office of Air Quality Planning and Standards (MD-14),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Rudolf Husar—CAPITA, Washington University, Campus Box 1124,
One Brookings Drive, St. Louis, MO 63130

Dr. Charles W. Lewis—National Exposure Research Laboratory (MD-47),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Scott Mathias—Office of Air Quality Planning and Standards (MD-15)
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Tom McCurdy—National Exposure Research Laboratory (MD-56),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Frank McElroy—National Exposure Research Laboratory (MD-46),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Haluk Özkaynak—National Exposure Research Laboratory (MD-56),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Tom Pace—Office of Air Quality Planning and Standards (MD-14),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers
(cont'd)

Dr. Joseph Pinto—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Richard Poirot—VT Air Program, Building 3 South, 103 South Main Street,
Waterbury, VT 05671

Dr. Linda Sheldon—National Exposure Research Laboratory (MD-77),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Helen Suh—Harvard School of Public Health, 665 Huntington Avenue,
Boston, MA 02461

Mr. Robert Wayland—Office of Air Quality Planning and Standards (MD-15),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Russell Weiner—National Exposure Research Laboratory (MD-46)
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Dane Westerdahl—California Air Resources Board, 2020 L Street,
Sacramento, CA 95814

CHAPTER 3. CONCENTRATIONS, SOURCES, AND EMISSIONS OF ATMOSPHERIC PARTICULATE MATTER

Principal Author

Dr. Joseph P. Pinto—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Contributing Authors

Dr. Barbara J. Turpin, The State University of New Jersey, Rutgers, Department of
Environmental Sciences and Rutgers Cooperative Extension, New Brunswick, NJ 08901-8551

Dr. James J. Schauer, University of Wisconsin, College of Engineering, Department of Civil and
Environmental Engineering, Madison, WI 53706

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers

Dr. JoEllen Bandemeyer—Research Triangle Institute, PO Box 12194, Research Triangle Park, NC 27709

Dr. Timothy Buckley—Johns Hopkins University, Department of Environmental Health Sciences, 615 North Wolfe Street, Baltimore, MD 21205

Ms. Lee Byrd—Office of Air Quality Planning and Standards (MD-14), U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lyle Chinkin—Sonoma Technology, 1360 Redwood Way, Suite C, Petaluma, CA 94549

Dr. Candis S. Claiborn—Washington State University, Laboratory for Atmospheric Research, Department of Civil and Environmental Engineering, P.O. Box 642910, Pullman, WA 99164

Dr. Steven Colome—Integrated Environmental Services, 5319 University Drive, #430, Irvine, CA 92612

Mr. Tom Coulter—National Exposure Research Laboratory (MD-47), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Delbert Eatough—Brigham Young University, E 114 BNSN, Department of Chemistry and Biochemistry, Provo, UT 84602

Dr. Edward O. Edney—National Exposure Research Laboratory (MD-47), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. William Ewald—National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Terence Fitz-Simons—Office of Air Quality Planning and Standards (MD-14), U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Sheldon Friedlander—University of California at Los Angeles, Department of Chemical Engineering, 5531 Boelter Hall, Los Angeles, CA 90095

Dr. Judith Graham—National Exposure Research Laboratory (MD-75), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lynn Hildemann—Civil and Environmental Engineering Department, Stanford University, Stanford, CA 94305

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers
(cont'd)

Mr. Jim Homolya—Office of Air Quality Planning and Standards (MD-14),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Rudolf Husar—CAPITA, Washington University, Campus Box 1124,
One Brookings Drive, St. Louis, MO 63130

Dr. Allen S. Lefohn—A.S.L. & Associates, 111 North Last Chance Gulch, Helena, MT 59601

Dr. Charles W. Lewis—National Exposure Research Laboratory (MD-47),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Phil Lorang—Office of Air Quality Planning and Standards (MD-14),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Karen Magliano—California Air Resources Board, 2020 L Street,
Sacramento, CA 95814

Mr. Scott Mathias—Office of Air Quality Planning and Standards (MD-15)
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Tom McCurdy—National Exposure Research Laboratory (MD-56),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Frank McElroy—National Exposure Research Laboratory (MD-46),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Nehzat Mottallebi—California Air Resources Board, 2020 L Street,
Sacramento, CA 95814

Dr. Haluk Özkaynak—National Exposure Research Laboratory (MD-56),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Tom Pace—Office of Air Quality Planning and Standards (MD-14),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Richard Poirot—VT Air Program, Building 3 South, 103 South Main Street,
Waterbury, VT 05671

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers
(cont'd)

Mr. Win Setiawan—California Air Resources Board, 2020 L Street,
Sacramento, CA 95814

Dr. Linda Sheldon—National Exposure Research Laboratory (MD-77),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Helen Suh—Harvard School of Public Health, 665 Huntington Avenue,
Boston, MA 02461

Mr. Robert Wayland—Office of Air Quality Planning and Standards (MD-15),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Dane Westerdahl—California Air Resources Board, 2020 L Street, Sacramento, CA 95814

Dr. William Wilson—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

CHAPTER 4. ENVIRONMENTAL EFFECTS OF PARTICULATE MATTER

Principal Authors

Ms. Beverly Comfort—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. William Ewald—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. J.H.B. Garner—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lester D. Grant—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. David A. Grantz—University of California/Riverside, Kearney Agricultural Center,
9240 South Riverbend Avenue, Parlier, CA 93648

Authors, Contributors, and Reviewers
(cont'd)

Principal Authors
(cont'd)

Dr. Paul J. Hanson—Environmental Sciences Division, Oak Ridge National Laboratory, P.O. Box 2008, Bethel Valley Road, Building 1059, Oak Ridge, TN 37831-6422

Dr. Dale W. Johnson—Environmental and Resource Science, 1000 Valley Road, University of Nevada, Reno, NV 89512

Dr. Joseph P. Pinto—National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. William H. Smith—Yale University School of Forestry and Environmental Studies, 370 Prospect Street, New Haven, CT 06511

Contributors and Reviewers

Dr. Larry T. Cupitt—National Exposure Research Laboratory (MD-75), U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Debra Meyer Wefering—Duckterather Weg 61, Bergisch Gladbach, Germany 54169 (formerly with National Exposure Research Laboratory [MD-56], U.S. Environmental Protection Agency, Research Triangle Park, NC 27711)

Dr. Russell R. Dickerson—University of Maryland, Department of Meteorology, Stadium Drive, College Park, MD 20742

Dr. Anne Grambsch—National Center for Environmental Assessment (8601D), U. S. Environmental Protection Agency, Washington, DC 20036

Dr. Sagar V. Krupa—University of Minnesota, Department of Plant Pathology, 495 Borlaug Hall, 1991 Upper Buford Circle, St. Paul, MN 55108

Dr. Alan J. Krupnick—Quality of the Environment Division, Resources for the Future, 1616 P Street, NW, Washington, DC 20036

Mr. Paul T. Roberts—Sonoma Technology, Inc.,1360 Redwood Way - Suite C, Petaluma, CA 94954

Mr. John Spence—1206 Sturdivant Drive, Cary, NC 27511

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers
(cont'd)

Dr. Richard Zepp—National Exposure Research Laboratory (IOD),
U. S. Environmental Protection Agency, Athens, GA

CHAPTER 5. HUMAN EXPOSURE TO PARTICULATE MATTER AND ITS CONSTITUENTS

Principal Authors

Dr. David T. Mage—Institute for Survey Research, Temple University, Philadelphia, PA 19122-6099 (formerly with the National Exposure Research Laboratory (MD-56), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711)

Mr. Thomas McCurdy—National Exposure Research Laboratory (MD-56),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Linda S. Sheldon—National Exposure Research Laboratory (MD-56),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Halûk Özkaynak—National Exposure Research Laboratory (MD-56),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. William E. Wilson—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Contributing Authors

Dr. Janet Burke—National Exposure Research Laboratory (MD-56),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Roy Fortmann—National Exposure Research Laboratory (MD-56),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Gary Norris—National Exposure Research Laboratory (MD-47),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Authors, Contributors, and Reviewers
(cont'd)

Contributing Authors
(cont'd)

Dr. Anne Rea—National Exposure Research Laboratory (MD-56),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Alan Vette—National Exposure Research Laboratory (MD-56),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Contributors and Reviewers

Dr. Timothy Buckley—Johns Hopkins University, Department of Environmental Health Sciences, 615 North Wolfe Street,, Baltimore, MD 21205

Ms. Lee Byrd—National Exposure Research Laboratory (MD-75),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lyle Chinkin—Sonoma Technology, 1360 Redwood Way, Suite C, Petaluma, CA 94549

Dr. Steven Colome—Integrated Environmental Services, 5319 University Drive, #430,
Irvine, CA 92612

Dr. Delbert Eatough—Brigham Young University, E 114 BNSN,
Department of Chemistry and Biochemistry, Provo, UT 84602

Dr. Sheldon Friedlander—University of California at Los Angeles, Department of Chemical Engineering, 5531 Boelter Hall, Los Angeles, CA 90095

Dr. Judith Graham—National Exposure Research Laboratory (MD-75),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lester D. Grant—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Vic Hasselblad—29 Autumn Woods Drive, Durham, NC 27713

Dr. Lynn Hildemann—Civil and Environmental Engineering Department
Stanford University, Stanford, CA 94305

Mr. Jim Homolya—Office of Air Quality Planning and Standards (MD-14),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers
(cont'd)

Dr. Rudolf Husar—CAPITA, Washington University, Campus Box 1124,
One Brookings Drive, St. Louis, MO 63130

Dr. Dennis J. Kotchmar—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Charles W. Lewis—National Exposure Research Laboratory (MD-47),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Allan Marcus—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Frank McElroy—National Exposure Research Laboratory (MD-46),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Tom Pace—Office of Air Quality Planning and Standards (MD-14),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Joseph Pinto—National Center for Environmental Assessment (MD-52),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Richard Poirot—VT Air Program, Building 3 South, 103 South Main Street,
Waterbury, VT 05671

Mr. Harvey Richmond—Office of Air Quality Planning and Standards (MD-15),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Linda Sheldon—National Exposure Research Laboratory (MD-77),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Helen Suh—Harvard School of Public Health, 665 Huntington Avenue,
Boston, MA 02461

Mr. Robert Wayland—Office of Air Quality Planning and Standards (MD-15),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Dane Westerdahl—California Air Resources Board, 2020 L Street, Sacramento, CA 95814

Dr. Jim Xue—Harvard School of Public Health, 665 Huntington Avenue, Boston, MA 02115

U.S. ENVIRONMENTAL PROTECTION AGENCY
PROJECT TEAM FOR DEVELOPMENT OF AIR QUALITY CRITERIA FOR PARTICULATE MATTER

Executive Direction

Dr. Lester D. Grant—Director, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Scientific Staff

Dr. William E. Wilson—PM Team Leader and Air Quality Coordinator, Physical Scientist, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lawrence J. Folinsbee—Health Coordinator, Chief, Environmental Media Assessment Group, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711 (now deceased)

Dr. Dennis J. Kotchmar—Project Manager, Medical Officer, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Robert Chapman—Medical Officer, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Beverly Comfort—Health Scientist, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. William Ewald—Health Scientist, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. J.H.B. Garner—Ecological Scientist, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. David Mage—Physical Scientist, Institute for Survey Research, Temple University, Philadelphia, PA 19122-6099

Dr. Allan Marcus—Statistician, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. James McGrath—Visiting Senior Health Scientist, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

U.S. ENVIRONMENTAL PROTECTION AGENCY
PROJECT TEAM FOR DEVELOPMENT OF AIR QUALITY CRITERIA FOR PARTICULATE MATTER

(cont'd)

Scientific Staff
(cont'd)

Dr. Joseph P. Pinto—Physical Scientist, National Center for Environmental Assessment, U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. James A. Raub—Health Scientist, National Center for Environmental Assessment (MD-52), U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Technical Support Staff

Mr. Randy Brady—Deputy Directory, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Douglas B. Fennell—Technical Information Specialist, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Emily R. Lee—Management Analyst, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Diane H. Ray—Program Specialist, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Eleanor Speh—Office Manager, Environmental Media Assessment Branch, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Donna Wicker—Administrative Officer, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Richard Wilson—Clerk, National Center for Environmental Assessment (MD-52), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

U.S. ENVIRONMENTAL PROTECTION AGENCY
PROJECT TEAM FOR DEVELOPMENT OF AIR QUALITY CRITERIA FOR PARTICULATE MATTER

(cont'd)

Document Production Staff

Dr. Carol A. Seagle—Technical Editor, Computer Sciences Corporation,
2803 Slater Road, Suite 220, Morrisville, NC 27560

Ms. Diane G. Caudill—Graphic Artist, Computer Sciences Corporation,
2803 Slater Road, Suite 220, Morrisville, NC 27560

Ms. Carolyn T. Perry—Word Processor, Computer Sciences Corporation,
2803 Slater Road, Suite 220, Morrisville, NC 27560

Ms. Kelly Quiñones—Word Processor, InfoPro, Inc., 8405 Colesville Road, 2nd Floor,
Silver Spring, MD 20910

Technical Reference Staff

Mr. John A. Bennett—Technical Information Specialist, SANAD Support Technologies, Inc.,
11820 Parklawn Drive, Suite 400, Rockville, MD 20852

Ms. Sandra L. Hughey—Technical Information Specialist, SANAD Support Technologies, Inc.,
11820 Parklawn Drive, Suite 400, Rockville, MD 20852

Ms. Beth Olen—Records Management Technician, Reference Retrieval and Database Entry
Clerk, InfoPro, Inc., 8405 Colesville Road, 2nd Floor, Silver Spring, MD 20910

Abbreviations and Acronyms

σ_{abs}	light-absorption coefficient
σ_{ag}	light-absorption coefficient of gases
σ_{ap}	light-absorption coefficient of particles
σ_{ext}	light-extinction coefficient
σ_{g}	geometric standard deviation
σ_{scat}	light-scattering coefficient
σ_{sg}	light-scattering coefficient of gases
σ_{sp}	light-scattering coefficient of particles
4-POBN	α-(4-pyridyl-1-oxide)-N-tert-butylnitrone
A	alveolar
AAS	atomic absorption spectrophotometry
ACGIH	American Conference of Governmental Industrial Hygienists
AD	
ADS	annular denuder system
AES	atomic emission spectroscopy
AIRS	Aerometric Information Retrieval System
AM	alveolar macrophages
AQCD	Air Quality Criteria Document
AQI	Air Quality Index
AQRV	Air Quality Related Values
ARIES	Aerosol Research and Inhalation Epidemiology Study
ASOS	Automated Surface Observing System
ATDM	aerosol and toxic deposition model
ATOFMS	time-of-flight mass spectrometer
b	
B_a	absorption coefficient

BAD	brachial artery diameter
BAL	bronchoalveolar lavage
BALF	bronchoalveolar lavage fluid
BAUS	brachial artery ultrasonography
BC	black carbon (see also CB)
BW	bronchial wash
BYU	Bringham Young University
C	apparent contrast
Ca^{+2}	calcium
CAA	Clean Air Act
CAAM	continuous ambient mass monitor
CAMNET	
CAPs	concentrated ambient particles
CARB	California Air Resources Board
CASAC	Clean Air Scientific Advisory Committee
CASTNet	Clean Air Status and Trends Network
CAT	computer-aided tomography
CB	carbon black
C_B	base cation
CC	carbonate carbon
CCl_4	carbon tetrachloride
CCPM	continuous coarse particle monitor
CCSEM	computer-controlled scanning electron microscopy
CEN	European Standardization Committee
CF	Cystic Fibrosis
CFA	coal fly ash
CFCs	chlorofluorocarbons
CFD	computational fluid dynamics

CFR	Code of Federal Regulations
CH_2O	formaldehyde
CIF	charcoal-impregnated cellulose fiber
CL	chemiluminescence
CMAQ	Community Multi-Scale Air Quality
CMB	chemical mass balance
CMD	count mean diameter
CMP	copper smelter dust
CMSA	Consolidated Metropolitan Statistical Area
C_o	initial contrast
CO	carbon monoxide
CO CD	Air Quality Criteria Document for Carbon Monoxide
COPD	chronic obstructive pulmonary disease
CPC	condensation particle counter
CPZ	capsazepine
CR	concentration-response
CRP	Coordinated Research Program
CSIRO	
CSMCS	Carbonaceous Species Methods Comparison Study
CTM	chemistry-transport model
CV	coefficient of variation
D_{50}	
D_a	
DAQM	Denver Air Quality Model
DCFH	dichlorofluorescin
DE	deposition efficiencies
DE	diesel exhaust
DEF	Deferoxamine

DEP	diesel exhaust particles
DHR	dihydrorhodamine-123
DMS	dimethyl sulfide
DMTU	dimethylthiourea
DOFA	domestic oil fly ash
DPM	diesel particulate matter
DRG	dorsal root ganglia
dv	deciview index
EAD	electrical aerosol detector
EC	elemental carbon
ECAO	Environmental Criteria and Assessment Office
ECG	electrocardiogram
EDXRF	energy dispersive X-ray fluorescence
EGA	evolved gas analysis
EGF	epidermal growth factor
ELSIE	Elastic Light Scattering and Interactive Efficiency
ERK	extracellular receptor kinase
ESP	electrostatic precipitator
ESR	electron spin resonance
ET	extrathoacic
ETS	environmental tobacco smoke
EU	endotoxin units
EXPOLIS	
F	flux
FEF	forced expiratory flow
FEV_1	forced expiratory volume in 1 second
FID	flame ionization detection
FMD	flow-mediated dilation

FPD	flame photometric detector
FRM	Federal Reference Method
g SO_2	gaseous sulfur dioxide
GC	gas chromatography
GCMs	General Circulation Models
GCVTC	Grand Canyon Visibility Transport Commission
GG/MSD	gas chromatography/mass-selective detection
GHG	greenhouse gases
GMCSF	granulocyte macrophage colony stimulating factor
GMPD	geometric mean particle diameter
GSD	geometric standard deviation (see also σ_g)
GSH	glutathione
H_2SO_4	sulfuric acid
HAAQS	
HDM	house dust mite
HDS	honeycomb denuder/filter pack sampler
HEADS	Harvard-EPA Annular Denuder Sampler
HEI	Health Effects Institute
hivol	High blume sampler
HNO_3	nitric acid
HR	heart rate
HTGC-MS	high temperature gas chromotography-mass spectrometry
I	radiance
IκBα	inhibitory kappa B alpha
I_b	apparent radiance of the background
I_{bt}	transmitted radiance of the background
IC	ion chromatography
ICAM-1	intercellular adhesion molecule-1

ICP	inductively coupled plasma
ICRP	International Commission on Radiological Protection
Ie	equilibrium radiance or source function
IFS	Integrated Forest Study
IgE	immunoglobin E
IgG	immunoglobin G
IL	interleukin
IMPROVE	Interagency Monitoring of Protected Visual Environments
INAA	instrumental neutron activation analysis
IOVPS	integrated organic vapor/particle sampler
ip	intraperitoneal
I_p	path radiance
IPCC	Intergovernmental Panel on Climate Change
IPM	inhalable particulate matter
IPN	Inhalable Particulate Network
ISO	International Standards Organization
I_t	transmitted radiance
JNK	c-jun N-terminal kinase
J_{scp}	light scattering by coarse particles
J_{sfp}	light scattering by fine particles
J_{spd}	light scattering coefficient of particles under dry conditions
J_{spw}	light scattering coefficient of particles under humid conditions
K	Koschmieder constant
K^+	potassium ion
KOH	potassium hydroxide
LAI	leaf area indices
LFA-1	leukocyte function-associated antigen-1
LN	lymph nodes

LoS	low sulfur
lpm, Lpm, L/min	liters per minute
LPS	lipopolysaccharide
LWCA	liquid water content analyzer
MAA	mineral acid anion
MAACS	Metropolitan Acid Aerosol Characterization Study
MADPro	Mountain Acid Deposition Program
MAPK	mitogen-activated protein kinase
MAQSIP	page 3-83
MCM	mass concentrations monitor
MCT	monocrotaline
MEK	mitogen-activated protein kinase
MIP	macrophage inflammatory protein
Mm	megameters
MMAD	mean median aerodynamic diameter (see σ_g)
MMD	mass median diameter
MMPs	matrix metalloproteinases
MOUDI	micro-orifice uniform deposit impactor
MPL	multipath lung
MPO	myeloperoxidase
MS	mass spectroscopy
MSA	methane sulfonic acid
MSAs	metropolitan statistical areas
MSH	Mount St. Helens
MSP	
NAC	N-acetylcysteine (antioxidant)
NAL	nasal lavage fluid
NAMS	National Ambient Monitoring Stations

NaN_3	sodium azide
NAPAP	National Acid Precipitation Assessment Program
NAPRMN	
NARSTO	
NAST	National Assessment Synthesis Team
NCRPM	National Council on Radiation Protection and Measurements
ND	NIST diesel (also, not determined)
NDDN	National Dry Deposition Network
NDIR	nondispersive infrared spectrophotometry
NESCAUM	Northeast States for Coordinated Air Use Management
NF	nuclear factor
NF-κB	nuclear factor kappa B
NFRAQS	North Frontal Range Air Quality Study
NH_3	ammonia
NH_4^+	ammonium
$(NH_4)_2\ SO_4$	ammonium sulfate
$NH_4H_2SO_4$	ammonium acid sulfate
NHBE	normal human bronchial epithelial
NIOSH	
NIR	
NIST	National Institute of Standards and Technology
NMD	nitroglycerine-mediated dilation
NMD	number mean diameter
NMRI	Naval Medical Research Institute
NO	nitrogen oxide
NO_2	nitrogen dioxide
NO_3^-	nitrate
NOPL	naso-oro-pharyngo-laryngeal

NO_x	nitrogen oxides
NPP	net primary production
NRC	National Research Council
NuCM	nutrient cycling model
O_3	ozone
OAA	Ottowa ambient air
OAQPS	Office of Air Quality Planning and Standards
OAR	Office of Air and Radiation
OC	organic carbon
OFA	oil fly ask
OH^-	hydroxyl ion
ORD	Office of Research and Development
OVA	ovalbumin
p	partial pressure
p SO_4^{2-}	particulate sulfate
PAH	polynuclear aromatic hydrocarbon
PAHs	polycyclic aromatic hydrocarbons
PAN	peroxyacetyl nitrate
PAR	photosynthetically active radiation
PB	polymyxin-B
PBL	planetary boundary layer
PBY	
PC	pyrolitic carbon
PC	particle concentrator
PC-BOSS	Particulate Concentrator-Brigham Young University Organic Sampling System
PCA	principal component analysis
PCBs	polychloronated biphenyls
PCDD	polychlorinated dibenzo-*p*-dioxins

PCDF	polychlorinated dibenzofurans
PCM	particle composition monitor
pdf	probability density functions
PDGF	platelet-derived growth factor
PEM	Personal Environmental Monitor
PESA	proton elastic scattering analysis
PFA	
PIXE	proton induced X-ray emission
PM	particulate matter
PM AQCD	PM Air Quality Criteria Document
$PM_{(10-25)}$	coarse particulate matter
$PM_{2.5}$	fine particulate matter
PMF	positive matrix factorization
PMN	polymorphonuclear leukocytes
p^o	equilibrium vapor pressure
poly I:C	polyionosinic-polycytidilic acid
POP	persistent organic pollutant
PROBDET	Probability of Detection Algorithm
PTEAMS	
PTEP	PM_{10} Technical Enhancement Program
PTFE	polytetrafluoroethylene
PTFE	polytetrafluoroethylene
PUF	polyurethane foam
Q	respiratory flow rates
Qabs	efficiency of absorption
Qext	efficiency of extinction
Qscat	efficiency of scattering
r_a	aerodynamic resistance

RAAS	
RADM	Regional Acid Deposition Model
RAMS	Real-Time Air Monitoring System
RAMS	Regional Air Monitoring Study
RAPS	Regional Air Pollution Study
r_b	boundary layer resistance
r_c	canopy resistance
REMSAD	Regulatory Modeling System for Aerosols and Deposition
RFO	residual fuels oils
RH	relative humidity
ROFA	residual oil fly ash
ROFA	residual oil fly ash
ROME	Reactive and Optics Model Emissions
ROS	reactive oxygen species
RPM	respirable particulate matter
RPM	Regional Particulate Model
RTE	rat tracheal epithelial
RTP	Research Triangle Park
S	saturation ratio
SA	Sierra Anderson
SAD	small airway disease
SASS	
SCAQS	Southern California Air Quality Study
SCC	
SCOS	Southern California Ozone Study
sd	standard deviation
SEM	scanning electron microscopy
SES	sample equilibration system

SEV	Sensor Equivalent Visibility
SH	spontaneously hypertensive
SIP	State Implementation Plans
SIXE	synchrotron induced X-ray emission
SL	stochastic lung
SLAMS/NAMS	
SLAMS	State and Local Air Monitoring Stations
SLE	St. Louis encephalitis
SMPS	scanning mobility particle sizer
SO_2	sulfur dioxide
SO_4^{2-}	sulfate
SOA	
SOC	semivolatile organic compounds
SoCAB	South Coast Air Basin
SOD	superoxide dismutase
SOPM	secondary organic particulate matter
SP	Staff Paper
SPM	synthetic polymer monomers
SRI	
SRM	standard reference method
SSM	solid sampler module
Stk	Stokes number
SUVB	solar ultraviolet B radiation
SVOC	semivolatile organic compounds
SWMMC	Southwest Metropolitan Mexico City
T(CO)	core temperature
TB	tracheabronchial
TDF	total deposition fraction

TDMA	Tandem Differential Mobility Analyzer
TEOM	tapered element oscillating microbalance
TEOMs	
TIMP	tissue inhibitor of metaloproteinase
TLN	
TNF	tumor necrosis factor
TOFMS	aerosol time-of-flight mass spectroscopy
TOR	thermal/optical reflectance
TOT	thermal/optical transmission
TPM	thoracic particulate matter
TRXRF	total reflection X-ray fluorescence
TSI	
TSP	total suspended particulates
UAM-V	Urban Airshed Model Version V
UCM	unresolved complex mixture
ufCB	ultrafine carbon black
UFP	ultrafine fluorospheres
UNEP	United Nations Environment Programme
URG	University Research Glassware
USGCRP	U.S. Global Change Research Program
UVD	Utah Valley dust
VAPS	Versatile Air Pollution Samplers
VASM	Visibility Assessment Scoping Model
VBE	Japanese B encephalitis
VCAM-1	vascular cell adhesion molecule-1
V_d	deposition velocity
VDI	
VOC	volatile organic compounds

V_s	sedimentation velocity
V_t	turbulent diffusion velocity
V_t	tidal volume
WC	tungsten carbide
WEE	western equine encephalitis
WINS	Well Impactor Ninety-Six
WIS	Wistar
WKY	Wistar-Kyoto
WMO	World Meteorological Organization
Wo	single scattering albedo
WRAC	Wide Range Aerosol Classifier
X-XRF	synchrotron induced X-ray fluorescence
XAD	polystyrene-divinyl benzene
XRF	X-ray fluorescence
μ*	

EXECUTIVE SUMMARY

E.1 INTRODUCTION

E.1.1 Purpose of the Document

The purpose of this document, Air Quality Criteria for Particulate Matter, is to present air quality criteria for particulate matter (PM) in accordance with Clean Air Act (CAA) Sections 108 and 109, which govern establishment, review, and revision of U.S. National Ambient Air Quality Standards (NAAQS) as follows:

- Section 108 directs the U.S. Environmental Protection Agency (EPA) Administrator to list pollutants that may reasonably be anticipated to endanger public health or welfare and to issue air quality criteria for them. The air quality criteria are to reflect the latest scientific information useful in indicating the kind and extent of all identifiable effects on public health and welfare expected from the presence of the pollutant in ambient air.

- Section 109 directs the EPA Administrator to set and periodically revise, as appropriate, (a) primary NAAQS, which in the judgement of the Administrator, are requisite to protect public health, with an adequate margin of safety, and (b) secondary NAAQS which, in the judgement of the Administrator, are requisite to protect the public welfare from any known or anticipated adverse effects (e.g., impacts on vegetation, crops, ecosystems, visibility, climate, man-made materials, etc.).

- Section 109 of the CAA also requires periodic review and, if appropriate, revision of existing criteria and standards. Also, an independent committee of non-EPA experts, the Clean Air Scientific Advisory Committee (CASAC), is to provide the EPA Administrator advice and recommendations regarding the scientific soundness and appropriateness of criteria and NAAQS.

To meet these CAA mandates, this document assesses the latest scientific information useful in deriving criteria as scientific bases for decisions on possible revision of current PM NAAQS. A separate EPA PM Staff Paper will draw upon assessments in this document, together with technical analyses and other information, to identify alternatives for consideration by the EPA Administrator with regard to possible retention or revision of the PM NAAQS.

E.1.2 Organization of the Document

The present document is organized into nine chapters, as follows:

- This Executive Summary summarizes key points from the ensuing chapters.

- Chapter 1 provides a general introduction, including a brief summary of the history of the PM NAAQS and an overview of issues, methods and procedures used to prepare this document.

- Chapters 2, 3 and 5 provide background information on air quality and exposure aspects to help to place the succeeding discussions of PM effects into perspective.

- Chapter 4 deals with environmental effects of PM on vegetation and ecosystems, visibility, manmade materials, and climate.

- Human health issues related to PM are addressed in Chapter 6 (on dosimetry); Chapter 7 (on toxicology); and Chapter 8 (on community epidemiology).

- Chapter 9 provides an integrative synthesis of key points from the preceding chapters.

E.2 AIR QUALITY AND EXPOSURE ASPECTS

The document's discussion of air quality and exposure aspects considers chemistry and physics of atmospheric PM; analytical techniques for measuring PM mass, size, and chemical composition; sources of ambient PM in the United States; temporal/spatial variability and trends

in ambient U.S. PM levels; and ambient concentration-human exposure relationships. Key findings are summarized in the next several sections. Overall, the atmospheric science and air quality information provides further evidence substantiating the 1996 PM AQCD conclusion that distinctions between fine and coarse mode particles (in terms of emission sources, formation mechanisms, atmospheric transformation, transport distances, air quality patterns, and exposure relationships) warrant fine and coarse PM being viewed as separate subclasses of ambient PM.

E.2.1 Chemistry and Physics of Atmospheric Particles

- Airborne PM is not a single pollutant, but rather is a mixture of many subclasses of pollutants with each subclass containing many different chemical species. Atmospheric PM occurs naturally as fine-mode and coarse-mode particles that, in addition to falling into different size ranges, differ in formation mechanisms, chemical composition, sources, and exposure relationships.

- PM may be primary or secondary. PM is called primary if it exists in the same chemical form in which it was emitted or generated. PM is called secondary if it is formed through the atmospheric reaction of a precursor gas that forms a condensible product that in turn nucleates to form new particles or condenses on existing particles.

- Fine-mode PM is derived primarily from combustion material that has volatilized and then condensed to form primary PM or from precursor gases reacting in the atmosphere to form secondary PM. New fine-mode particles are formed by the nucleation of gas phase species; they grow by coagulation (existing particles combining) or condensation (gases condensing on existing particles). Fine particles are composed of freshly generated nuclei-mode particles, also called ultrafine or nanoparticles, and an accumulation mode (so-called because particles grow into and remain in that mode).

- Coarse-mode PM, in contrast, is formed by crushing, grinding, and abrasion of surfaces, which breaks large pieces of material into smaller pieces. These particles are then suspended by the wind or by anthropogenic activity. Energy considerations limit the break-up of large mineral particles and small particle aggregates generally to a minimum size of about 1 μm in diameter,

although biological material may exist or fragment into smaller sizes. Mining and agricultural activities are examples of anthropogenic sources of coarse-mode particles. Fungal spores, pollen, and plant and insect fragments are examples of natural bioaerosols also suspended as coarse-mode particles.

- Within atmospheric particle modes, the distribution of particle number, surface, volume, and mass by diameter is frequently approximated by lognormal distributions. Aerodynamic diameter, which depends on particle density and is defined as the diameter of a particle with the same settling velocity as a spherical particle with unit density (1 g/cm^3), is often used to describe particle size. Typical values of the mass median aerodynamic diameters (MMAD) are 0.05 to 0.07 μm for the nuclei mode, 0.3 to 0.7 μm for the accumulation mode, and 6 to 20 μm for the coarse mode. At high relative humidities or in air containing evaporating fog or cloud droplets, the accumulation mode may be split into a droplet mode (MMAD = 0.5 to 0.8 μm) and a condensation mode (MMAD = 0.2 to 0.3 μm).

- Research studies use impactors to determine mass and composition as a function of size over a wide range and particle counting devices to determine number of particles as a function of size. Such studies indicate an atmospheric bimodal distribution of fine and coarse particle mass with a minimum in the distribution between 1 and 3 μm aerodynamic diameter. Routine monitoring studies prior to 1999 generally measured thoracic PM, i.e., PM_{10} (upper size limited by a 50% cut at 10 μm aerodynamic diameter). Research studies and monitoring studies since 1999 measure fine PM, i.e., $PM_{2.5}$ (upper size limited by a 50% cut point at 2.5 μm aerodynamic diameter) and coarse thoracic PM, i.e, $PM_{10\text{-}2.5}$ the coarse fraction of PM_{10}, measured as the difference between PM_{10} and $PM_{2.5}$ mass measurements obtained at the same time and location and with similar inlets and other sampling and handling specifications. Cut points are not perfectly sharp for any of these PM indicators; some particles larger than the 50% cutpoint are collected and some particles smaller than the 50% cutpoint are not retained.

- The terms “fine” and “coarse” were originally intended to apply to the two major atmospheric particle distributions which overlap in the size range between 1 and 3 μm diameter. Now, fine has come to be often associated with the $PM_{2.5}$ fraction and coarse is often used to refer to

$PM_{10-2.5}$, coarse thoracic PM. However, $PM_{2.5}$ may also contain, in addition to the fine-particle mode, some of the lower-size tail of the coarse particle mode between about 1 and 2.5 μm aerodynamic diameter. Conversely, under high relative humidity conditions, the larger fine particles in the accumulation mode may also extend into the 1 to 3 μm aerodynamic diameter range.

- Four approaches are used to classify particles by size: (1) modes, based on formation mechanisms and the modal structure observed in the atmosphere, e.g., nuclei and accumulation modes (which comprise the fine-particle mode) and the coarse-particle mode; (2) cut point, based on the 50% cut point of the specific sampling device; (3) dosimetry, based on the ability of particles to enter certain regions of the respiratory tract; and (4) regulatory, based on instrument configuration or 50% cut-points, e.g., high volume sampler, PM_{10}, and $PM_{2.5}$.

E.2.2 Sources of Airborne Particles in the United States

- The chemical complexity of airborne particles requires that the composition and sources of a large number of primary and secondary components be considered. Major components of fine particles are: sulfate, strong acid, ammonium, nitrate, organic compounds, trace elements (including metals), elemental carbon, and water.

- Primary particles are emitted directly from sources. Secondary particles are formed from atmospheric reactions of sulfur dioxide (SO_2), nitrogen oxides (NO_x), and certain organic compounds. NO reacts with ozone (O_3) to form NO_2. SO_2 and NO_2 react with hydroxy radical (OH) during the daytime to form sulfuric and nitric acid. During the nighttime, NO_2 reacts with ozone and forms nitric acid through a sequence of reactions involving the nitrate radical (NO_3). These acids may react further with ammonia to form ammonium sulfates and nitrates. Some types of higher molecular weight organic compounds react with OH radicals, and olefinic compounds also react with ozone to form oxygenated organic compounds, which nucleate or can condense onto existing particles. SO_2 also dissolves in cloud and fog droplets, where it may react with dissolved O_3, H_2O_2, or, if catalyzed by certain metals, with O_2, yielding sulfuric acid or sulfates, that lead to PM when the droplet evaporates.

- Organic compounds constitute from 10 to 70% of dry $PM_{2.5}$ mass. Whereas the chemistry of particulate nitrate and sulfate formation has been relatively well studied, the chemistry of secondary organic particulate matter formation is still under active investigation. Although additional sources of secondary organic PM might still be identified, there appears to be a general consensus that biogenic compounds (monoterpenes, sesquiterpenes) and aromatic compounds (e.g., toluene and ethylbenzene) are the most significant precursors. Atmospheric transformations of the compounds, which are formed in the particle phase during the aging of particles, are still not adequately understood.

- Receptor modeling has proven to be a useful method for identifying contributions of different types of sources, especially for the primary components of ambient PM. Apportionment of secondary PM is more difficult because it requires consideration of atmospheric reaction processes and rates. Results from western U.S. sites indicate that fugitive dust, motor vehicles, and wood smoke are the major contributors to ambient PM samples there, whereas results from eastern U.S. sites indicate that stationary combustion, motor vehicles and fugitive dust are major contributors to ambient PM samples there. Sulfate and organic carbon are the major secondary components in the East, while nitrates and organic carbon are the major secondary components in the West.

E.2.3 Atmospheric Transport and Fate of Airborne Particles

- Primary and secondary fine particles have long lifetimes in the atmosphere (days to weeks) and travel long distances (hundreds to thousands of kilometers). They tend to be uniformly distributed over urban areas and larger regions, especially in the eastern United States. As a result, they are not easily traced back to their individual sources.

- Coarse particles normally have shorter lifetimes (minutes to hours) and generally only travel short distances (<10's of km). Therefore, coarse particles tend to be unevenly distributed across urban areas and tend to have more localized effects than fine particles. However, dust storms occasionally cause long range transport of small coarse-mode particles.

E.2.4 Airborne Particle Measurement Methods

- Measurements of ambient PM mass and chemical composition are needed to determine attainment of standards; to guide attainment of a standard (including determination of source categories and validation of air quality models); and to determine health, ecological, and radiative effects. A comprehensive approach requires a combination of analytical techniques to assess: (1) mass, (2) crustal and trace elements, (3) water-soluble ionic species including strong acidity, (4) elemental carbon, and (5) organic compounds.

- There are no calibration standards for suspended particle mass; therefore, the accuracy of particle mass measurements cannot be definitively determined. The precision of particle mass measurements can be determined by comparing results from collocated samplers. Mass concentration measurements with a precision of 10% or better have been obtained with collocated samplers of identical design. Field studies of EPA PM_{10} and $PM_{2.5}$ reference methods and reviews of field data from collocated PM_{10} and $PM_{2.5}$ samplers show high precision (better than ± 10%). The use of more careful techniques, including double weighing of filters, can provide higher precision and may be needed for precise determination of $PM_{10-2.5}$ by difference.

- Available technology allows accurate (± 10 to 15%) measurement of several of the major components of coarse and fine particles (crustal and trace elements, sulfates, nitrates, ammonium, and strong acidity). However, collection and measurement technologies for elemental carbon and organic carbon are not as well established. The split between elemental and organic carbon is operational, i.e., it is different for the two most frequently used measurement techniques. In addition, in order to estimate the mass of organic PM, the ratio of oxygen to carbon in organic PM must be estimated. It is higher for secondary organic than for primary organic PM, adding further to the uncertainty in organic and elemental carbon measurements.

- Semivolatile organic compounds and semivolatile ammonium compounds (such as NH_4NO_3) may be lost by volatilization during sampling. Such losses may be very important in woodsmoke impacted areas for organic compounds or in agricultural and other areas where low

sulfate and high ammonia lead to high NH_4NO_3 concentrations. New techniques are now in use for measurement of nitrates and new research techniques are being tested for measurement of mass of semivolatile organic compounds in PM and of the total (semivolatile plus non-volatile) PM mass. The Federal Reference Methods for PM_{10} and $PM_{2.5}$ give precise (± 10%) measurements of "equilibrated mass". However, the loss of semivolatile PM (ammonium nitrate and organic compounds) and the possible retention of some particle-bound water in current PM mass measurements contribute to uncertainly in the measurement of the mass of PM as it exists suspended in the atmosphere.

- Intercomparisons, using different techniques and samplers of different designs, coupled with mass balance studies (relating the sum of components to the measured mass), provide a method for gaining confidence in the reliability of PM measurements.

E.2.5 Ambient PM Concentrations in the U.S.: Regional Patterns and Trends

- Particle mass data have been collected at a number of rural, suburban, and urban sites across the United States by various local, state, and national programs. The data have been stored in the Aerometric Information Retrieval System (AIRS). Data have also been collected at remote sites as part of the IMPROVE and NESCAUM networks. An extensive analysis of this data was reported in the 1996 Air Quality Criteria Document for Particulate Matter (PM AQCD).

- The median $PM_{2.5}$ concentration across the United States during 1999 and 2000, the first two years of operation of the $PM_{2.5}$ FRM network, was 13 $\mu g/m^3$, with a 95th percentile value of 18 $\mu g/m^3$. The corresponding median $PM_{10-2.5}$ concentration was 10 $\mu g/m^3$, with a 95th percentile value of 21 $\mu g/m^3$.

- The spatial variability of $PM_{2.5}$ concentrations is characterized in this document, based on the availability of data at four or more sites within twenty-seven urban areas across the United States. Correlations of $PM_{2.5}$ concentrations between pairs of monitoring sites within the urban areas examined ranges from low to high. Highest correlations are found at site-pairs that are dominated by regional sources of secondary PM. Low correlations can be found if the sites are located in different air sheds or if at least one of the sites is affected more strongly by local,

primary sources. Although $PM_{2.5}$ concentrations may be highly correlated between sites, the concentrations themselves may not be uniform.

- Annual mean $PM_{2.5}$ concentrations within the urban areas examined are typically within about five $\mu g/m^3$ of each other. However, on a daily basis, absolute differences in $PM_{2.5}$ concentrations can be much larger. In approximately half of these urban areas, the 90th percentile difference in daily $PM_{2.5}$ concentrations is greater than 10 $\mu g/m^3$. Extreme values of concentration differences were greater than 100 $\mu g/m^3$ in a few cases. Caution should therefore be exercised in using these data to approximate community-wide exposures.

- The database for characterizing the spatial variability of $PM_{10-2.5}$ concentrations is not as extensive as it is for $PM_{2.5}$. Intersite correlations of $PM_{10-2.5}$ concentrations were lower than those for $PM_{2.5}$ in the few urban areas that had sufficient data for both $PM_{2.5}$ and $PM_{10-2.5}$. $PM_{10-2.5}$ concentrations also tended to be more variable, at least on a relative basis.

E.2.6 Human Exposure to PM

In community epidemiology studies of PM and other air pollutants, ambient concentrations are normally used as surrogates for personal exposure to pollutants of ambient origin. Since people spend most of their time indoors, and the indoor environment is protective for most ambient pollutants, it is important to understand the relationship between concentrations of ambient pollutants measured at community monitoring sites and the contributions of those concentrations to personal exposure. This is best done by considering separately (a) the relationship between concentrations at a community air-monitoring sites(s) and immediately outside an indoor environment, (b) the relationship between concentrations outside the indoor environment and the contributions of the outdoor pollutant to the indoor environment, and (c) the effect of activity patterns, i.e., time spent outdoors and in various indoor environments.

- Analyses of recent data from the $PM_{2.5}$ monitoring network show reasonable site-to-site correlation in most cities studied over distances of 20 to 50 km. This indicates that in such cities the concentration at a community air-monitoring site (or the average of several such sites) will provide an adequate representation of the concentration outside a home. Less information

is available regarding site-to-site correlations for $PM_{10-2.5}$, chemical components of PM (other than sulfate which has high site-to-site correlations), and contributions from specific source categories such as vehicular traffic-related PM. Even though site-to-site correlations may be high, annual or seasonal averages may show sizable differences.

- The relationship between outdoor air pollutant concentrations and indoor concentrations due to the concentration outdoors depends on the penetration factor (the fraction of the outdoor concentration which reaches the indoor environment), how rapidly the indoor air is diluted by outdoor air (measured as the air exchange rate), and the rate at which the ambient pollutant is deposited or removed in the indoor environment. The deposition rate for PM is highly dependent on particle size, being high for coarse and ultrafine particles but low for particles in the accumulation-mode size range (0.1 to 1.0 μm diameter). Thus, the infiltration factor (the ratio of the indoor concentration to the outdoor concentration) will be high for accumulation-mode particles and for $PM_{2.5}$ since most of the $PM_{2.5}$ mass will be in the accumulation mode.

- Exposure also depends on the amount of time people spend outdoors. The attenuation factor, in the case of PM, is defined as the ratio of the ambient PM exposure to the ambient PM concentration, and accounts for the difference in the time spent indoors and outdoors as well as the difference in exposure between indoors and outdoors.

- People are also exposed to particles and other pollutants generated indoors. It is not possible to measure ambient PM exposure directly; only the combination of ambient and nonambient PM exposure (total personal exposure to PM) can be measured. Ambient PM exposure must be inferred or estimated from measurements of ambient concentration and total personal exposure.

- Major indoor sources are smoking, other indoor combustion, cooking, cleaning, and general movement of people. Indoor particles are generated primarily in the ultrafine or coarse modes and therefore have shorter indoor lifetimes than ambient-infiltrated particles (particles that have penetrated indoors and remained suspended). The concentration of PM from indoor sources appears to be independent of ambient concentrations, since personal activities generally do not depend on ambient concentrations; however, this may change as more people are alerted to

high pollution days and stay indoors. If nonambient PM exposure is independent (not correlated with) ambient concentrations, a regression of measured personal exposures against ambient concentrations will provide the average attenuation coefficient (slope of the regression line) and the average nonambient concentration (the intercept). Such average values have been obtained in several studies.

- It is more difficult to estimate individual daily values of ambient PM exposures. This could be done for the PTEAM study because, in addition to ambient concentrations and personal exposures, data were available on air exchange rates and time outdoors and the penetration factor and deposition rate were estimated statistically. Ambient PM exposures can also be estimated by using the personal sulfate/ambient sulfate ratio as an estimate of the attenuation factor for $PM_{2.5}$. This technique assumes that there are both minimal indoor sources of sulfate and that the $PM_{2.5}$ and sulfate have similar particle size distributions.

- Most exposure studies measure one (or a few) subjects on one day and a different one (or a few) subjects on a different day. The highly variable nonambient exposure for different people results in a low correlation between ambient concentration and total personal exposure for this "pooled" data set. If a set of individuals each have their total personal exposure measured for enough days to provide a meaningful relationship, it is observed that some of them will have high correlations between ambient concentration and total personal exposure. The median correlations from such studies ("longitudinal") are higher than that for the "pooled" data set. If enough people are measured each day so that a meaningful daily average can be obtained, the correlation between ambient concentration and the daily average community PM exposure is high. Also, the correlation between ambient concentration and ambient PM exposure is high. Therefore, ambient PM concentration appears to provide an adequate indicator of ambient PM exposure for use in PM epidemiology studies, but such studies do not provide information on the health effects of nonambient pollution (i.e., indoor-generated pollution).

- As long as the nonambient PM exposure is not correlated with the ambient PM exposure, it will not bias the estimated health effect of PM. However, the effect per μg/ambient PM concentration will be biased low compared to the health effect per μg/ambient PM exposure by

the attenuation factor. This effect probably explains some of the heterogeneity in PM_{10} effects observed in multicity epidemiology studies, as indicated by a correlation of PM effects in different cities with air conditioning use in those cities (i.e., the higher the air conditioning use, the lower the health effect estimate per $\mu g/m^3$ of ambient PM).

- Exposure relationships also provide some insight into the issue of confounding. While the data base is small, concentrations of gaseous co-pollutants, NO_2, O_3, and SO_2 (and probably CO) are likely poorly correlated, and sometimes not significantly correlated, with personal exposure to the respective co-pollutant. However, they are frequently significantly correlated with both the ambient PM concentration and the ambient PM exposure. Thus, in a regression, where associations are found between gaseous co-pollutants and a health effect, it may be because they are a surrogate for PM rather than a confounder. That is, the health effect due to PM is transferred to the gaseous pollutant because of the positive correlation between the ambient concentration of the gas and the ambient PM exposure.

E.3 DOSIMETRY

Knowledge of the dose of particles delivered to a target site or sites in the respiratory tract is important for understanding possible health effects associated with human exposure to ambient PM and for extrapolating and interpreting toxicologic data obtained from studies of laboratory animals. Particles of different sizes are subject to large differences in regional respiratory tract deposition, translocation, clearance mechanisms and pathways, and consequent retention times. Key findings derived from the assessment of dosimetry information include:

- Respiratory tract deposition patterns are dependent on particle size, as indicated by the aerodynamic or thermodynamic diameter of the particles within the inspired air. Biologic effects may be a function not only of particle mass deposition but also of particle number; the total surface area of the particles; or the acidity, surface chemistry, or charge of the particles.

- Particles may be deposited in the extrathoracic (ET) region (i.e., mouth, nose, pharynx, and larynx); the conducting airways of the tracheobronchial (TB) region; and the alveolar (A)

region, where gas exchange occurs. There are differences in deposition mechanisms and dose distribution in each of these regions that are dependent on the physical characteristics of particles and on airway geometry.

- Particles deposit in the respiratory tract mainly by five mechanisms: (1) inertial impaction, (2) sedimentation, (3) diffusion, (4) electrostatic precipitation, and (5) interception. Impaction is an important deposition mechanism for particles >1μm in large extra- and intrathoracic airways at higher flows; sedimentation and diffusion are more important for particles >0.5 μm and <0.3 μm, respectively, at low flow rates in smaller airways. Particles between 0.3 and 0.5 μm in size are small enough to be little influenced by impaction or sedimentation and large enough to be minimally influenced by diffusion; and, so, they undergo the least respiratory tract deposition. Electrostatic precipitation is deposition related to particle charge; effects of charge on deposition are inversely proportional to particle size and airflow rate. The interception potential of any particle depends on its physical size rather than its aerodynamic size.

- Hygroscopicity, the propensity of a material for taking up and retaining moisture, is a property of some ambient particle species and affects respiratory tract deposition. Hygroscopicity generally increases deposition in the TB and A regions for particles with initial sizes larger than $\approx$0.5 μm or smaller than $\approx$0.01, but decreases deposition for intermediate sizes.

- The ET region acts as an efficient filter that reduces penetration of inhaled particles to the TB and A regions of the lower respiratory tract. Total respiratory tract deposition increases with particle size for particles >1.0 μm, is at a minimum for particles 0.3 to 0.5 μm, and increases as particle size decreases below that range.

- Enhanced particle retention occurs on carinal ridges in the trachea and through segmental bronchi; and deposition “hot spots” occur at airway bifurcations or branching points. Peak deposition sites shift from distal to proximal sites as a function of particle size, with greater surface dose in conducting airways than in the A region for all particle sizes. However, surface number dose (particles/cm^2/day) is much higher for fine particles than for coarse for typical bi-modal ambient aerosols.

- Extrathoracic deposition of ultrafine particles (<0.1 μm) is very high; as particle size decreases below 0.1 μm, particles tend to behave more like gases. Estimates of extrathoracic deposition range from 50% for oral breathing to >90% for nasal breathing. Within the thoracic region, the deposition distribution of ultrafine particles is highly skewed towards the proximal airway regions and resembles the deposition of coarse particles.

- Various host factors have been shown to influence particle deposition patterns, including airway dimensions (size and shape), breathing pattern (flow and volume), and the presence of obstructive or inflammatory airway disease. The ET deposition is higher with nose breathing than for mouth breathing, with increased ventilation rates associated with increasing levels of physical activity or exercise leading to more oronasal breathing and increased delivery of inhaled particles to TB and A regions in the lung. Gender and age differences in the homogeneity of deposition, as well as deposition rate, could affect susceptibility. Children, for example, would receive greater doses of particles per lung surface area than would adults. Also, obstructive airway diseases (such as asthma and chronic bronchitis) result in increased deposition of particles in the central airway region and distal lung regions receiving greater ventilation.

- Particles depositing on airway surfaces may be cleared from the respiratory tract completely or translocated to other sites within this system by regionally specific clearance mechanisms. Clearance is either absorptive (dissolution) or nonabsorptive (transport of intact particles). Deposited particles may be dissolved in body fluids, taken up by phagocytic cells, or transported by the mucociliary system. Retained particles tend to be small (<2.5 μm) and poorly soluble (e.g., silica, metals).

- Tracheobronchial clearance has both a fast and a slow component. In the fast phase particles deposited in the TB region clear out rapidly during the first several hours and continue to clear out for 24 hours. A small remaining portion may clear out over several days (slow phase). Translocation of poorly soluble PM to the lymph nodes takes a few days and is more rapid for smaller (< 2 μm) particles; elimination rates of these retained particles are on the order of years. People with COPD have increased particle retention partly because of increased initial

deposition and impaired mucociliary clearance and use cough to augment mucociliary clearance.

- Alveolar clearance takes months and years. Particles may be taken up by alveolar macrophages within 24 hours, but some phagocytosed macrophages translocate into the interstitium or lymphatics whereas some remain on the alveolar surface. Penetration of uningested particles into the interstitium increases with increasing particle load and results in increased translocation to lymph nodes.

- Acute effects of PM are probably best related to deposited dose, whereas chronic effects may be related to cumulative or retained dose. Retention of particles is a function of deposition site, clearance of particles by macrophages or the mucociliary system, and particle characteristics, especially solubility. Chronic effects may also arise from recurring cycles of pulmonary injury and repair.

- Mathematical and computational fluid dynamic models are available to predict deposition, clearance, and retention of particles in the respiratory tract. Although these models have become more sophisticated and versatile, validation of the models is still needed.

- A better understanding of species differences in deposition, translocation, and clearance of particles, especially ultrafine particles, is still needed. So are better models of extrapolation between animals used in inhalation studies and humans.

E.4 PARTICULATE MATTER HEALTH EFFECTS

E.4.1 Toxicology of Particulate Matter in Humans and Laboratory Animals

Toxicological research on ambient PM or combustion-related particles is used to address several related questions that are important toward an understanding of the cardiopulmonary effects that have been reported in PM-exposed human populations.

- Does exposure to PM at relevant ambient concentrations cause toxicological effects?
- What characteristics of PM contribute to the observed toxicity?
- What factors affect individual or subpopulation susceptibility to the effects of PM?
- What are the combined effects of PM and other pollutants in the ambient air?
- What mechanisms may be involved in the toxicological response to PM exposure?

Data on the toxicology of PM are derived from controlled inhalation exposure studies of humans and laboratory animals, intratracheal instillation studies in humans and animals, and ex vivo studies of human and animal cells grown in culture. The human or animal populations (cells) studied vary by age, health status, or other host factors. As seen in the previous section, deposition of PM in the respiratory tract depends on particle size and regional distribution. Potential biologic effects may be a function not only of particle mass deposition but also of particle number, the total surface area of the particles, particle acidity, and the surface chemistry, charge, and composition of the particle in addition to other exposure variables (e.g., duration, temperature, humidity, activity levels). Responses to PM in the respiratory tract also are dependent on the physiological status of the host, as well as the translocation of PM or PM constituents to other sites. Ex vivo studies provide important additional information regarding mechanisms of action of PM or PM constituents on cells or cellular components.

The data available in the previous 1996 PM AQCD and in other published documents were mainly from studies that investigated the respiratory effects of specific components of combustion-related particles from mobile or stationary sources (e.g., diesel particles, fly ash), ambient particles, or laboratory-derived surrogate particles (e.g., sulfuric acid droplets). In this document, more emphasis is placed on assessment of new data obtained from controlled studies of particles collected from emission sources or ambient samplers (e.g., impactors, diffusion denuders) and by the use of aerosol concentrators that provide a technique for exposing humans or laboratory animals by inhalation to concentrated ambient particles (CAPs). Key findings derived from the assessment of these effects include:

- Combustion-related particles (fly ash and urban air particles) from a large number of emission sources and ambient airsheds cause a spectrum of responses in the airways of laboratory animals and humans. These include inflammation, cellular injury, and increased permeability.

Soluble metal components (e.g., Cu, Fe, Ni, V, Zn) of combustion particles have been implicated in the responses, possibly related to oxidant production and release of intercellular signaling molecules (cytokines).

- Toxicological studies of aqueous extracts from ambient PM collected on filters in the Utah Valley around Provo, UT demonstrated increased pulmonary inflammatory effects after airway instillation exposure of humans and laboratory animals, and after direct exposure to cells in culture. Extract analysis of particle components acquired during operation of an open-hearth steel mill identified more sulfate, cationic salts (Ca, K, Mg), and metals (As, Cu, Fe, Mn, Ni, Pb, Sr, Zn). The inflammatory response was significantly reduced when the steel mill was closed, thus tending to corroborate epidemiology findings for the same time period, indicating that Utah Valley residents reported decreased hospital admissions for respiratory diseases.

- Cells primed by inflammatory mediators show increased cytokine responses to PM. Combustion-related particles may cause increased oxidant production, presumably related to metal components of particles, and damage to cells in vitro. Responses include impaired macrophage phagocytosis and altered permeability.

- Acute exposures to soluble transition metals can cause inflammatory responses in the respiratory tract of humans and laboratory animals. The effective exposure levels (mg/m^3) are typically much higher than typical ambient air metal concentrations ($<15\ \mu g/m^3$) in the U.S. atmosphere.

- Endotoxin, a lipopolysacharide associated with bacteria, and a common contaminant of ambient PM, also causes inflammation in humans and laboratory animals at concentrations ($>0.5\ \mu g$) that are much higher than typically found in the ambient air ($<0.5\ ng/m^3$).

- Human inhalation exposure to diesel exhaust particles causes increased acute sensory and respiratory symptoms, lung inflammation, and impairment of alveolar macrophage function. Effects in laboratory animals include pulmonary histopathology and chronic inflammation. These noncancer effects are thought to be due to the organic carbon constituents or to

metabolites of metal components of the particle. Except for diesel exhaust particles, no other research has been published on acute effects of organic carbon constituents which often comprise a substantial portion of ambient PM (10 to 60% of the total dry mass).

- Recent studies report systemic changes in rodents and dogs exposed to high concentrations of instilled and inhaled ambient PM and combustion-related particles, including alteration of heart rate (e.g., bradycardia, arrhythmia) and blood pressure, hypothermia, alterations in blood cells, and increased blood levels of endothelins (vasoactive substances) and fibrinogen (coagulation factors).

Mixtures

- Mixtures of ozone and PM (e.g., urban PM, diesel PM, sulfate aerosols, ultrafine carbon) may cause enhanced effects on lung cells, increased inflammation, and decrements in human lung function. In one controlled human study, a mixture of ozone and CAPs produced peripheral vasoconstriction, possibly caused by autonomic reflexes or as a result of increased circulating endothelins.

Mechanisms

- A number of studies indicate that increased production of inflammatory cytokines and reactive oxidant species (ROS) may play a role in PM-induced responses. The cytokine responses correlate with endotoxin, which is an important component of ambient coarse- and fine-mode PM. Catalysis of ROS is likely related to soluble metals in ambient PM and combustion-related particles.

- Somatosensory neurons of the autonomic nervous system (ANS) may also be affected by the inflammatory response to ambient PM, especially when there is epithelial airway damage. Pulmonary reflex responses through the ANS can have direct effects on the heart and may cause other systemic effects.

- Studies on ultrafine compared to fine-mode particles indicate a greater response to ultrafine particles in regards to airway inflammation, an effect that appears to be related to their greater

surface area. There also is some evidence to suggest that ultrafine PM may exit the lung and deposit in other organs, including the heart and liver.

- Combustion-related particles (e.g., oil fly ash) and urban PM can induce apoptosis (programmed cell death) of human alveolar macrophages.

- Other potential cellular and molecular mechanisms include changes in the expression of specific growth factors, adhesion molecules, stress proteins, matrix proteinases, transcription factors, and antioxidant enzymes.

Susceptibility

- Chemically or pharmacologically treated rat "models" of cardiopulmonary disease (e.g., hypertension, chronic bronchitis), as well as older rats, manifest increased cardiopulmonary responses, lung damage, and even death after exposure to ambient PM and combustion-related particles by intratracheal instillation or inhalation.

- Inhalation or instillation of combustion-related particles (e.g., oil fly ash) and diesel exhaust PM may augment the immune response to antigens in allergic animals or humans. These studies provide a plausible mechanism for an association between combustion-source PM exposure and exacerbation of asthma.

- Acid aerosols cause little or no changes in pulmonary function in healthy subjects; however, asthmatics may develop small, but potentially relevant increased airway responsiveness. New information relating acid aerosol exposure to cardiovascular effects in laboratory animals is interesting, but needs further investigation.

- Genetic susceptibility can plan a role in the response to inhaled or instilled particles.

E.4.2 Population Groups at Risk

Susceptibility can be affected by factors which influence dosimetry or the response of tissues to particle burdens. Host factors that may increase the susceptibility to PM include both

changes in physiologic factors affecting respiratory tract deposition and pathophysiologic factors affecting response.

- Susceptible groups most clearly at special risk for PM effects include the elderly and those with cardiopulmonary disease, based on available epidemiology findings.

- Epidemiology findings indicate that risk of mortality and morbidity due to lower respiratory disease (e.g. pneumonia) is increased by ambient PM exposure. This may be due to exacerbation by PM of already existing respiratory disease. PM may also increase susceptibility to infectious disease by decreasing clearance, impairing macrophage function, or through other specific and nonspecific effects on the immune system. The epidemiologic findings also indicate that individuals with preexisting infectious respiratory disease (e.g., pneumonia) are at increased risk for PM effects.

- Epidemiologic findings indicate that ambient PM exposures are also associated with increased risk for mortality and hospitalization due to cardiovascular causes. Cardiac arrhythmia has been hypothesized as being involved in mortality due to acute PM exposure. Thus, individuals with pre-existing cardiovascular disease(s) are likely a susceptible group at increased risk for ambient PM effects

- Studies of infants and children indicate that they are a potentially susceptible population. Panel studies on asthma and other respiratory conditions show exacerbation by PM exposure. Children are susceptible to respiratory effects associated with PM exposure from pre-natal and post-natal effects through exacerbation of asthma and respiratory symptoms in school age children.

E.4.3 Epidemiology Findings

Epidemiologic evidence concerning the mortality and morbidity effects of ambient PM has expanded greatly since the 1996 PM Air Quality Criteria Document (PM AQCD). The most important enhancements in information include:

- New multi-city studies of health endpoints using ambient PM_{10} mass concentrations and/or other ambient PM indicators (e.g. Black Smoke, Coefficient of Haze, etc).

- New studies on a variety of endpoints using ambient fine particle fraction ($PM_{2.5}$), a limited number using ambient coarse thoracic PM fraction ($PM_{10-2.5}$), and a few using ambient ultrafine particle ($PM_{0.1}$) mass concentrations;

- New studies in which the relationship of some health endpoints to ambient particle number concentrations were evaluated.

- Additional studies which evaluated the sensitivity of estimated PM effects to the inclusion of gaseous co-pollutants in the model.

- New studies evaluating the effects of specific source categories of air pollution based on empirical combinations (factor analysis).

- Further studies of cardiopulmonary endpoints associated with PM exposures. Cardiovascular, as well as respiratory, causes of death and hospitalization in older adults may be a significant component of PM-attributable mortality.

- New studies suggest that infants and children may represent an additional subgroup at special risk for ambient PM exposure effects. The new results most clearly indicate that children appear to be susceptible to respiratory effects associated with ambient PM exposures, including exacerbation of asthma and respiratory symptoms in school-age children.

- A few studies also report ambient PM to be associated with intrauterine growth reduction and low birth weight (known infant health risk factors) and excess infant mortality. However, no toxicologic evidence has yet been advanced to support biological plausibility of such effects due to ambient PM or to identify the pathophysiologic mechanisms involved.

- New analyses of American Cancer Society (ACS) data extended over more years and addressing criticisms of prior ACS analyses not only substantiate previous ACS study findings of PM associations with increased risk of cardiopulmonary-related mortality/morbidity but also find PM associations with increased risk of lung cancer.

- PM health effects have been reported to be associated with several different ambient PM size fractions (ultrafine, fine, coarse); individual chemical components (sulfate, nitrate, elemental carbon, organic carbon) and specific source categories (vehicular traffic, regional sulfate, vegetative burning, and fuel oil combustion).

- Various health effects may occur at different time scales for exposure to PM, from (a) short -term responses to hourly to daily exposures through (b) larger excess mortality associated with medium-term exposures (15 to 120 day averages) to (c) excess morbidity or mortality associated with long-term (multi-year) exposures.

- Because PM indicators and the gaseous co-pollutants, CO, NO_2, SO_2, and O_3, are frequently significantly correlated, the potential exists for the confounding of the adverse health effects attributable to $PM_{2.5}$ in short-term exposure studies by exposure to gaseous co-pollutants. This makes it difficult to apportion the risk among PM acting alone, PM acting in combination with gaseous co-pollutants, the gaseous co-pollutants *per se*, a specific source category, or the overall ambient pollutant mix. However, recent exposure studies suggest that the ambient concentrations of the gaseous co-pollutants, although frequently correlated with ambient PM concentrations, are not well correlated with the personal exposure to the respective gaseous co-pollutants. Therefore, the gaseous co-pollutants are not likely to be confounders, rather they are likely surrogates for PM (or specific source categories such as vehicular-traffic-related particles or regional sulfate). The low exposures to the reactive gaseous co-pollutants (NO_2, SO_2, O_3) as well as to CO, of people who spend most of their time indoors, relative to known toxic levels, also suggests that these gaseous pollutants are unlikely to be responsible for the health effects found to be associated with PM (although the gaseous pollutants may also independently exert effects on health, as well).

E.4.3.1 Ambient PM Mortality Effects

- New multi-city studies convincingly demonstrate the association of PM_{10} with health effects. The use of the same statistical model for many cities with different PM concentrations and compositions, and different correlations of PM and gaseous co-pollutants, strengthens the reliability of the associations and also demonstrates that the effects of PM are not confounded by the gaseous co-pollutants.

- Several studies have considered $PM_{2.5}$ and $PM_{10\text{-}2.5}$ as well as PM_{10}. In same cases $PM_{2.5}$, and in some cases $PM_{10\text{-}2.5}$, was more significant than PM_{10} and had a higher excess risk per $\mu g/m^3$ increase in PM. One study found both $PM_{2.5}$ and $PM_{10\text{-}2.5}$ to be statistically significant and to have excess risks that were higher than PM_{10} and that peaked on different lag days. Thus, epidemiologic studies also support the separation of PM_{10} into fine and thoracic coarse fraction components. However, there are some suggestions that the toxicity of $PM_{10\text{-}2.5}$ may not be due to the crustal or soil components per se, but rather more likely to toxic trace metals or organic compounds carried into the lungs on the coarse particles.

- Source apportionment techniques have been used to obtain groupings of PM components and gaseous co-pollutants (factors) with minimal correlation among factors. Use of these factors permits determination of associations with health effects with little potential for confounding among the factors. These factors can frequently be associated with specific source categories. However, the association of factors with source categories may be subjective and a factor may contain contributions from more than one source category.

- A vehicular traffic related factor has been identified in all four studies that examined that factor. Although not all studies measured all species, this factor appears to contain $PM_{2.5}$, CO, NO_2, EC, and OC, as well as specific elements Mn, Fe, Zn, and Pb, that might be emitted or resuspended by traffic. Epidemiology alone cannot apportion the health effect among these different components of the vehicular traffic related factor. However, the low potential for confounding of PM_{10} by the CO and NO_2 suggests that they may serve as surrogates for the PM component of the vehicular traffic-related factor.

- Independent associations, not confounded by gaseous co-pollutants have been found for nitrate (by single and multiple regression with gaseous co-pollutants) and for regional sulfate and a PM factor associated with vegetative burning (by use of source category factors). Thus, the major components of $PM_{2.5}$ by mass, sulfate, nitrate, and organic, have been found to have significant associations with mortality.

- A major concern has been that the effects attributed to PM may really be due to one or more of the gaseous co-pollutants (CO, NO_2, SO_2, O_3). Epidemiology cannot demonstrate causality, only association. However, multi-city studies in conjunction with exposure relationships have been able to demonstrate that the gaseous co-pollutants are more likely to be surrogates than confounders. This does not demonstrate that the gaseous co-pollutants do not have real effects, only that they are significantly correlated with the PM indicator but not with the personal exposure to the gaseous co-pollutants.

- The results of long-term exposure studies, indicative of increased mortality and/or morbidity risks being associated with exposures to ambient PM over many years, have been substantiated both by independent reanalyses and updated, extended analyses of more years of data and cases of mortality or morbidity. This includes much stronger evidence for ambient PM effects on cardiovascular and respiratory endpoints, as well as strong evidence for PM-related increases in lung cancer risks.

E.4.3.2 Ambient PM Morbidity Effects

Numerous epidemiologic studies in the United States and elsewhere have also demonstrated significant associations between ambient PM exposures indexed by a variety of indicators (PM_{10}, $PM_{2.5}$, $PM_{10-2.5}$, $SO_4^=$, H^+, BS, COH, TSP) and various acute and chronic morbidity outcomes. Such outcomes include, for example, hospital admissions, medical visits, increased respiratory symptoms, and decreased lung function.

- The ecologic time series studies add substantially to the body of available literature for effects of PM_{10} on acute CVD hospital admissions. Results for adult cardiovascular mortality are qualitatively consistent with those for hospital admissions. However, uncertainties regarding

the possible role of co-pollutants complicates interpretation with respect to independent PM_{10} effects. New studies evaluating longitudinal association between ambient PM and physiological measures of cardiovascular function or biochemical changes in the blood that may be associated with cardiac risk present a range of findings suggesting possible adverse effects of PM on cardiac rhythm and other biological functions.

- The results of new studies are generally consistent with regard to ambient PM associations of short-term exposures with respiratory-related hospital admissions/medical visits. The excess risk estimates fall most consistently in the range of 5 to 25% per 50 $\mu g/m^3$ PM_{10} increments, with those for asthma visits and hospital admissions tending to be somewhat higher than for COPD and pneumonia hospital admissions.

- The peak flow analyses results for asthmatics tend to show small decrements for both PM_{10} and $PM_{2.5}$. The effects of PM on respiratory symptoms in asthmatics tended to be positive, although they were much less consistent than the effects on lung function. Most PM_{10} studies showed increases in cough, phlegm, difficulty breathing, and bronchodilator use, although these increases were generally not statistically significant.

- Results of the PM_{10} peak flow analyses in non-asthmatic studies were inconsistent, with fewer studies reporting results in the same manner as for the asthmatic studies. The effects on respiratory symptoms in non-asthmatics were similar to those in asthmatics. Most studies showed that PM_{10} increases cough, phlegm, difficulty breathing, and bronchodilator use, although these increases were generally not statistically significant.

- Differences in peak flow and bronchitis symptoms and prevalence rates in children were found to be somewhat more closely associated with annual average H^+ concentrations than with other PM indicators. However, in studies demonstrating these effects, the acid levels were highly correlated with other fine-particle indicators.

- While numerous studies of PM related respiratory morbidity have been conducted using PM_{10} as an indicator, only a few studies have examined the effects of fine and coarse fraction particle

indicators separately. The $PM_{2.5}$ studies do show effects related to exposure to the fine fraction, but high correlations among $PM_{2.5}$, PM_{10}, and acid aerosols make it very difficult to attribute the effects to a single specific exposure indicator. A few studies also indicate that respiratory morbidity effects (increased respiratory symptoms) are associated with ambient coarse fraction ($PM_{10-2.5}$) concentrations.

E.5 ENVIRONMENTAL EFFECTS OF PM

E.5.1 Vegetation and Ecosystem Effects

- Deposition of particulate matter from the atmosphere has the potential to alter ecosystem structure and function. Human existence on this planet depends on the essential life-support services that ecosystem structure and functions provide. Concern has risen in recent years regarding the consequences of changing the biological diversity of ecosystems because human activities are creating disturbances that are altering the structure (complexity and stability) and functioning (producing changes in energy and water flow and nutrient cycling) of ecosystems.

- Human-induced changes in biotic diversity and alterations in the structure and functioning of ecosystems are the two most dramatic ecological trends in the past century. Biodiversity encompasses all levels of biological organization, including individuals, populations, species, and ecosystems. For this reason, there is a need to understand the effects of PM deposition on vegetation and ecosystems and biodiversity.

- Ecosystem functions maintain clean water, pure air, a green earth (biodiversity) and impart the following benefits: fixation of solar energy, absorption and breakdown of pollutants, cycling of nutrients, binding of soil, degradation of organic wastes, maintenance of a balance of atmospheric gases, regulation of radiation balance, and climate.

- The relationship between PM and effects on vegetation and ecosystems is dependent on the size, origin, and chemical constituents of the particles. Exposure to a given mass concentration

of PM may, depending on the particular mix of deposited particles, lead to widely differing phytotoxic responses.

- Atmospheric deposition of particles to ecosystems takes place via both wet and dry processes through the three major routes indicated below:
    (1) Precipitation scavenging in which particles are deposited in rain and snow
    (2) Fog, cloud-water, and mist interception
    (3) Dry deposition, a much slower, yet more continuous removal to surfaces.

- Deposition of heavy metal particles to ecosystems occurs by wet and dry processes. Dry deposition is considered more effective for coarse particles of natural origin and elements such as iron and manganese, whereas wet deposition generally is more effective for fine particles of atmospheric origin and elements such as cadmium, chromium, lead, nickel, and vanadium.

- The actual importance of wet versus dry deposition, however, is highly variable, depending on the type of ecosystem, location and elevation. The range of particle sizes, the variety of chemical constituents in airborne PM, and the diversity of canopy surfaces, have slowed progress in both prediction and measurement of dry particulate deposition. Wet deposition generally is confounded by fewer factors and has been easier to quantify.

- Particulate matter, when transferred from the atmosphere to plant surfaces, may cause direct effects when it (1) resides on the leaf, twig or bark surface for an extended period; (2) is taken up through the leaf surface; or produce indirect effects when (3) removed from the plant via suspension to the atmosphere, washing by rainfall, or by litter-fall with subsequent transfer to the soil.

- Deposition of PM on above-ground plant parts can have either a physical and or chemical impact, or both. The effects of “inert” PM are mainly physical, while the effects of toxic particles are both chemical and physical. The majority of the easily identified direct and indirect effects occur in severely polluted areas around heavily industrialized point sources (such as limestone quarries, cement kilns, iron, lead, and various smelting factories).

- Chemical effects include excessive alkalinity or acidity. Studies of the chemical additions of PM to foliage have found little or no effects on foliar processes unless exposure levels were significantly greater than would typically be expected in the ambient environment. The effects of dust deposited on plant surfaces or on soil are more likely to be associated with their chemistry than with the mass of deposited particles and are usually of more importance than any physical effects. The effects of limestone dust on plants can cause long-term changes in the structure, composition and function of the seedling-shrub and sapling strata of ecosystems in the vicinity of limestone quarries and processing plants.

- Secondary organics formed in the atmosphere and referred to as toxic substances, persistent organic pollutants (POPS), pesticides, hazardous air pollutants (HAPS), air toxics, and/or semivolatile organic compounds (SOCS), are chemical substances not controlled by National Ambient Air Quality Standards under Sect. 109 of the Clean Air Act (U.S. Code, 1994), but rather are controlled under Sect.112, Hazardous Air Pollutants. Mention of them is made in this document because many form or attach to particles in the atmosphere. As particles they become airborne and can be distributed over a wide area and impact remote ecosystems. Some are of concern to humans because they may reach toxic levels in food chains of animals as well as humans; others tend to decrease or maintain the same toxicity as they move through the food chain.

- The depletion of stratospheric ozone caused by the release of chloroflurocarbons (CFC's) and substances such as halides has resulted in heightened concern about potentially serious increases of solar UV-B (SUVB) reaching the earth's surface. Terrestrial vegetation is vulnerable to UV-B because of the need for sunlight during photosynthesis. Effects of UV-B on plant growth are likely to be incremental. However, plants grown in full sunlight (because they evolved under ambient UV-B radiation and have developed adaptive mechanisms) are not as sensitive as plants gown under weak visible light. Therefore, plant species vary enormously in their responses to UV-B exposures. In addition, large differences in response occur among genotypes within a species and dicotyledons are more sensitive than monocotyledons.

- Photosynthetic processes underlie the contributions of vegetation to nutrient cycling and energy flow. Light penetration into plant canopies limits photosynthetic productivity. Regional haze has been estimated to diminish surface visible solar radiation by approximately 8%. An assessment of the direct effects on crop production suggests that optimal crop yields are significantly affected by regional haze.

- Most PM deposited on vegetation eventually enters the soil environment, which is one of the most dynamic sites of biological interaction. The major impacts on vegetation and ecosystems are the indirect effects that occur through the soil and affect plant growth, vigor and reproduction. Changes in nutrient cycling and plant nutrient uptake determine plant and ecosystem responses.

- Bacteria and fungi in the soil have an important role in plant nutrition. Bacteria are essential components of the nitrogen and sulfur cycles that make these elements available for plant uptake. Fungi form mycorrhizae, a mutualistic symbiotic relationship with plant roots that is integral to the uptake of mineral nutrients. The impact of nitrates, sulfates and metals in PM is determined by their affect on the growth and functions of the bacteria and fungi involved in making nutrients available for plant uptake.

- Extensive evidence indicates that heavy metals deposited from the atmosphere to forests accumulate either in the richly organic forest floor, where the biological activity is the greatest, or in the soil layers immediately below. Accumulation of heavy metals in litter presents the greatest potential for altering nutrient cycling. Increased amounts of litter in metal-contaminated areas appear to result from reduced activity of microorganismal populations.

- Phytochelatins are intracellular metal-binding peptides that act as indicators of metal stress. Because they are produced by plants as a response to sublethal concentrations of heavy metals, they are indicators that heavy metals are involved in forest decline. Concentrations of phytochelatins were observed to increased with altitude as did forest decline and they also increased across the regions that showed increased levels of forest injury.

- The major impact of atmospheric PM on ecosystems is indirect and occurs in the soil environment through the deposition of nitrates and sulfates and the acidifying effects of the H+ ion associated with these compounds in wet and dry deposition. Most nitrate is not deposited or measured as PM but is a combination of wet and dry deposition.

- The deposition of nitrates, and the acidifying effects of the H^+ ion in association with nitrates and sulfates in precipitation, causes soil acidification, changes the Ca/Al ratio, affects the growth of soil bacteria and fungi, and alters biogeochemical cycling, all of which affects the growth of natural vegetation and forest trees. Leaching of nitrates and other minerals through runoff can also affect streams and coastal and aquatic wetlands and thus influence their ability to produce the products and services necessary for human society.

- Intensive research over nearly a decade indicates that, although the soils of most North American forests are nitrogen limited, severe symptoms of nitrogen saturation have been observed in: high-elevation spruce-fir forests of the Appalachian Mountains, hardwood watersheds near Parsons, WV; watersheds in the Los Angles Air Basin; high-elevation alpine watersheds in the Colorado Front Range; and a deciduous forest in Ontario, Canada.

- Nitrogen saturation results in a progressive syndrome of concurrent responses to long-term, chronic nitrogen deposition. As nitrogen reaches saturation in temperate-zone forests, there are decreases in nitrogen mineralization and increases in trends of foliar Mg:N and Ca:Al ratios. Preliminary evidence suggests some forests may decline in productivity and experience greater mortality as a result of chronic nitrogen deposition.

- Increases in soil nitrogen play a selective role. Plant succession patterns and biodiversity in some ecosystems are significantly affected by chronic nitrogen additions. Long-term nitrogen fertilization studies in both New England and Europe suggest that forests receiving chronic inputs of nitrogen may decline in productivity and experience greater mortality. Studies also suggest that declining coniferous forest stands with slow nitrogen cycling may be replaced by deciduous fast-growing forests which cycle nitrogen rapidly.

- Excess nitrogen inputs to unmanaged heathlands in the Netherlands has resulted in nitrophilous grass species replacing slower growing heath species. Over the past several decades the composition of plants in the forest herb layers had been shifting toward species commonly found in nitrogen-rich areas. It also was observed that the fruiting bodies of mycorrhizal fungi had decreased in number, indicating that formation of mycorrhizae were affected.

- The effects of excessive deposition of nitrogen, particularly NH_3 and NH_4^+, have led to changes in Dutch heathlands via: (1) acidification of the soil and the loss of cations of K^+, Ca^{2+} and Mg^{2+}; and (2) nitrogen enrichment, which results in increased plant growth rates and altered competitive relationships. Alteration of any of a number of parameters (e.g., increased nitrogen) can alter ecosystem structure and function.

- There is a major concern that soil acidification will result in nutrient deficiency. Growth of tree species can be affected when high aluminum to nutrient ratios limit uptake of calcium and magnesium. Calcium is essential in the formation of wood and the maintenance of cells, the primary plant tissues necessary for tree growth. Calcium must be dissolved in soil water to be taken up by plants. A major concern is that soil acidity will lead to nutrient deficiency.

- Acid deposition can increase the aluminum concentrations in soil water by lowering the pH in aluminum-rich soils through dissolution and ion-exchange processes. Aluminum in soil can be taken up by roots more readily than calcium because of its greater affinity for negatively charged surfaces. Tree species can be adversely affected if high Ca/Al ratios impair Ca and Mg uptake.

- Ecosystem processes and productivity, nitrogen mineralization rates, and nitrate leaching respond directly to human modification of ecosystems and to changes in atmospheric composition and climate.

E.5.2 Particulate Matter-Related Effects on Materials

Atmospheric PM and SO_2 exert effects on materials that are related both to aesthetic appeal and physical damage. Studies have demonstrated particles, primarily consisting of carbonaceous

compounds, cause soiling of commonly used building materials and culturally important items such as statues and works of art. Physical damage from the dry deposition of SO_2, particles, and the absorption or adsorption of corrosive agents on deposited particles can also result in the acceleration of the weathering of manmade building and naturally occurring cultural materials.

- The natural process of metal corrosion from exposure to environmental elements (wind, moisture, sun, temperature fluctuations, etc.) is enhanced by exposure to anthropogenic pollutants, in particular SO_2.

- Dry deposition of SO_2 enhances the effects of environmental elements on calcereous stones (limestone, marble, and carbonated cemented) by converting the calcium carbonate (calcite) in the stone to calcium sulphate dihydrate (gypsum). The rate of deterioration is determined by the SO_2 concentration, the stone's permeability and moisture content, and the deposition rate.

- Sulfur dioxide limits the life expectancy of paints by causing discoloration, loss of gloss, and loss of thickness of the paint film layer.

- A significant detrimental effect of particulate pollution is the soiling of painted surfaces and other building materials. Soiling is a degradation process requiring remediation by cleaning or washing, and depending on the soiled surface, repainting. Soiling decreases the reflectance of a material and reduces the transmission of light through transparent materials. Soiling may reduce the life usefulness of the material soiled.

E.5.3 Visibility

Chapter 4 of this document includes information supplementary to several other significant reviews of the science of visibility, including the 1991 report of the National Acid Precipitation assessment Program, the National Research Council's *Protecting Visibility in National Parks and Wilderness Areas* (1993), and EPA's 1995 *Interim Findings on the Status of Visibility Research*. The following points are made in Chapter 4 and/or in the above referenced documents.

- The relationships between air quality and visibility are well understood. Ambient fine particles are the major cause of visibility impairment. Significant scientific evidence exists showing that reducing fine particle concentrations will improve visibility.

- The National Research Council defines visibility qualitatively as "the degree to which the atmosphere is transparent to visible light." This definition may be expressed quantitatively in terms of contrast transmittance. The EPA has defined visibility impairment as a reduction in visual range (the farthest distance at which a large black object can be distinquished against the horizontal sky is the visual range) and/or atmospheric discoloration.

- Light, as it passes through the atmosphere from a scene to an observer, is both scattered and absorbed. The rate of loss of transmitted light intensity with distance is measured by the light-extinction coefficient, which may be expressed as the sum of the coefficients for: (a) light scattering due to gases; (b) light scattering due to particles; (c) light absorption by gases, and; (d) light absorption by particles. Light scattering by particles is the major component of light extinction. Light absorption by gases is almost entirely due to NO_2, and is typically significant only near NO_2 sources. Light absorption by particles is primarily caused by elemental carbon.

- Light scattering efficiency depends on particle size, falling off rapidly for particles below 0.3 or above 1.0 μm in diameter. Therefore, particles in the accumulation mode (of the fine particle mode) are most effective in scattering light and are more important in visibility degradation than either nuclei-mode or coarse-mode particles. Light absorption is not a strong function of particle size. Under exceptional circumstances, such as dust storms, coarse particles can dominate scattering.

- In addition to reducing the intensity of light carrying information about a scene (transmitted radiance), particles also scatter light into the observer's view. This extraneous light, called air light or path radiance, carries no information about the scene. The competition between these two sources of light, expressed as the ratio of transmitted radiance from the scene to path radiance, determines the contrast transmittance and the visual quality of the view.

- Visibility at any location is affected by air quality and non-air quality related effects. The visibility effects of atmospheric constituents are dependant upon not just the mass of pollutants, but on the size distribution and refractive index of particles, which are strongly influenced by relative humidity. Non-air quality effects include the angle between the sun and the observer's sight path, location of clouds, and reflectivity of the ground. These effects are independent of effects due to changes in atmospheric constituents. Lighting and scene effects can be accounted for by defining a range of these effects when estimating visibility changes due to air quality influences.

- The relationship between air pollution and the appearance of a scenic view is well understood. Models exist that, given an adequate description of the air quality and non-air quality variables, can produce a simulated photograph that accurately depicts a cloud-free scene as it would appear to a human observer.

- There are several potential quantitative indicators of visibility, including: (a) fine particle mass and composition (fine particle mass alone provides less of both types of information); (b) scattering by dried ambient particles; (c) scattering by particles under ambient conditions; (d) extinction (calculated from measurements of scattering plus absorption); (e) light extinction measured directly; and (f) contrast transmittance.

- A new index, the deciview *(dv)*, is now being used as a quantitative measure of haziness. It is related to the light extinction coefficient, b_{ext}, by $Haziness\ (dv) = 10\ ln(b_{ext}/10\ Mn)$. The deciview is more nearly linearly related to perceived changes in haze level than either visual range or light extinction. A change of 1 or 2 *dv* in uniform haze under many viewing conditions will be seen as a small but noticeable change in the appearance of a scene regardless of the initial haze condition.

- Visibility in the United States is best in the western, intermountain region. Visibility impairment or haziness is greatest in the eastern United States and southern California. Haziness in the eastern United States is caused primarily by atmospheric sulfate. Haziness in southern California is primarily caused by nitrate and organic PM. Nitrates contribute about

40% to the total light extinction in southern California. Nitrates account for 10 to 20% of the total extinction in other areas of the United States.

- Organics are the second largest contributors to light extinction in most areas in the United States. Organic carbon is the greatest cause of light extinction in the Pacific Northwest, Oregon, Idaho, and Montana, accounting for 40 to 45% of the total extinction. Organic carbon contributes between 15 to 20% to the total extinction in most of the western United States and 20 to 30% in the remaining areas of the United States.

- Light absorption by carbon is relatively insignificant but is highest in the Pacific Northwest (up to 15%) and in the eastern United States (3%).

- High dust concentrations transported from southern California and the subtropics have contributed to regional haze in the Grand Canyon and other Class I areas in the southwestern United States.

E.5.4 Global Change Processes and Their Potential Human Health and Environmental Impacts

Processes causing global change and their potential environmental and human health impacts have been accorded extensive attention during the past several decades, and they still continue to be of broad national and international concern. This is reflected by extensive research and assessment efforts undertaken since the mid-1970s by U.S. Federal Government Agencies (e.g., NOAA, EPA, CDC, etc.) or via U.S. Federal Interagency programs (e.g., the U.S. Global Change Research Program [USGCRP]). It is also reflected by analogous extensive research and assessment efforts undertaken by numerous other national governments or international collaborative activities, e.g., those coordinated by the Intergovernmental Panel on Climate Change (IPCC), established in the 1980s under the joint auspices of the World Meteorological Organization (WMO) and the United Nations Environment Programme (UNEP).

The present discussion of global climate change in Chapter 4 draws upon recent international assessments of (a) processes contributing to stratospheric ozone depletion and the status of progress towards ameliorating the problem (WMO, 1999) and (b) revised qualitative

and quantitative projections of likely consequent human health and environmental effects (UNEP, 1998, 2000) — with the findings and conclusions of these assessments being incorporated herein by reference. Also, Chapter 4 presents information on global warming and its potential impacts drawn mainly from extensive assessments contained in the Third Assessment Report of the IPPC (2001) and a series of reports by the USGCRP on projected impacts on the United States or subregions.

- Atmospheric particles play important roles in two key types of global change phenomena: (1) alterations in the amount of ultraviolet solar radiation (especially UV-B) penetrating through the Earth's atmosphere and reaching its surface, where it can exert a variety of effects on human health, plant and animal biota, and other environmental components; and (2) alterations in the amount of visible solar radiation transmitted through the Earth's atmosphere.

- Knowledge of factors controlling the transfer of solar radiation in the ultraviolet spectral range is needed for assessing potential biological and environmental impacts associated with exposure to UV-B radiation (290 to 315 nm). Knowledge of the effects of PM on the transfer of radiation in the visible and infrared spectral regions is needed for assessing the relationship between particles and global warming and its environmental and biological impacts.

PM Effects on Solar Ultraviolet Radiation Transmission Impacts

- The main types of deleterious effects hypothesized as likely to result from stratospheric ozone depletion and consequent increased SUVB penetration through the Earth's atmosphere are:
  (a) *Direct Human Health Effects*, such as skin damage (sunburn), leading to more rapid aging and increased incidence of skin cancer; ocular effects (retinal damage and increased cataract formation possibly leading to blindness); and suppression of some immune system components (possibly increasing susceptibility to certain infectious diseases or decreasing effectiveness of vaccinations).
  (b) *Agricultural/Ecological Effects*, mediated largely through altered biogeochemical cycling resulting in consequent damaging impacts on terrestrial plants (leading to possible reduced yields of rice, other food crops, and commercially important trees, as well as to biodiversity

shifts in natural terrestrial ecosystems); and deleterious effects on aquatic life (including reduced ocean zooplankton and phytoplankton, as important base components of marine food-chains supporting the existence of commercially important, edible fish and other seafood, as well as to other aquatic ecosystem shifts).

(c) *Indirect Human Health and Ecological Effects*, mediated through increased tropospheric ozone formation (and consequent exacerbation of surface-level, ozone-related health and ecological impacts) and alterations in the concentrations of other important trace species, most notably the hydroxyl radical and acidic aerosols.

(d) *Other Types of Effects*, such as faster rates of polymer weathering because of increased UV-B radiation and other effects on man-made commercial materials and cultural artifacts, secondary to climate change or exacerbation of air pollution problems.

- In contrast to negative impacts projected as likely to be associated with increased UV-B penetration to Earth's surface, some research results are suggestive of possible beneficial effects of increased UV-B radiation. For example, a number of U.S. and international studies report likely protective effects of UV-B radiation with regard to non-skin cancer incidence. These suggests potential relationships between sunlight, vitamin D, and reduced colon cancer and others provide evidence that UV-B radiation may be protective against several types of cancer and some other diseases.

- From among boundary layer pollutants (e.g., SO_2, O_3, PM), airborne particles are probably most effective in controlling the amount of SUV-B radiation reaching the Earth's surface. Fine particles are clearly more effective than coarse particles in this regard.

- Particles scatter and absorb solar radiation in the biologically important UV-B spectral region. The amount of SUV-B reaching the Earth's surface depends in a non-linear way on the content of scattering and absorbing material within airborne particles.

- Given the above considerations, quantification of projected effects of variations in atmospheric PM on human health or the environment because of the effects of particles on the transmission of solar UV-B radiation requires location-specific evaluations, taking into account composition,

concentration, and internal structure of the particles; temporal variations in atmospheric mixing height and depths of layers containing the particles the abundance of ozone and other absorbers within the planetary boundary layer and the free troposphere. The outcome of such modeling effects would likely vary from location to location in terms of increased or decreased surface level UV-B exposures because of location-specific changes in atmospheric PM concentrations or composition.

PM Effects on Global Warming Processes and Impacts

- During the 20th century global average surface temperatures increased by 0.6 ± 0.2°C. The decade of the 1990s was probably the warmest since 1861. The last few years have been among the warmest on record. The Intergovernmental Panel on Climate Change (IPCC) has attributed most of the warming observed over the past fifty years to human activities. The global average surface temperature is projected to rise by 1.4 to 5.8 °C from 1990 to 2100.

- There are health effects directly associated with climate change such as increased heat stress and cardiorespiratory failure due to rises in temperature. There are also health effects which are associated with changes in ecosystems and habitats of disease-carrying organisms that are the result of changes in climate variables such as temperature and humidity.

- Vectorborne diseases such as malaria and dengue fever may extend their ranges in the United States through the northward extension of habitats favorable to their development.

- Waterborne diseases may likely increase with increasing air and water temperatures, combined with heavy runoff events from agricultural and urban surfaces.

- The effects of climate change on air quality are also likely to be important, however, these effects are too uncertain to be predicted with any confidence at the present time. Likewise, little is known regarding changes in the effects of air quality on human health under a different climate.

- Particles, most notably those containing sulfate, exert a direct effect by scattering incoming solar radiation back to space. However, ‘black carbon’ in particles absorbs solar radiation and as these particles age, their capacity to absorb solar radiation may increase. Some recent estimates indicate that the effect of particles containing ‘black carbon’ may be twice as large as that given by the latest IPCC report and that the control of ‘black carbon’ emissions may be an effective means to slow ‘greenhouse warming’.

- Particles also exert an indirect effect on climate by causing an increase in the concentration of cloud droplets which is accompanied by a decrease in their size. The reduction in cloud droplet size tends to increase the liquid water content of the cloud, the lifetime of the cloud and the optical thickness of the cloud. As a result of these processes, clouds reflect more solar radiation back to space. Many of these effects have been observed; for example, cloud droplets in polluted areas tend to be smaller than those formed in clean areas. However, the magnitude of the overall effects on climate, although larger than the direct effects noted above, is more highly uncertain.

- On a globally averaged basis, radiative cooling due to anthropogenic particles may have substantially offset the radiative heating due to increases in atmospheric concentrations of greenhouse gases such as carbon dioxide, methane, and chlorofluorocarbons.

- Aerosol lifetimes are also much shorter than the time required for global mixing, therefore, aerosol radiative effects are most likely to exert their influence on a regional rather than on a global basis.

- The lifetimes of particles in the troposphere are short (days to weeks) compared to the above greenhouse gases (years to over 100 years). Therefore, aerosol concentrations will respond more rapidly to variations in emissions than will the greenhouse gases.

E.6 KEY CONCLUSIONS

- Epidemiologic studies show consistent positive associations of exposure to ambient PM with health effects, including mortality and morbidity. The observed associations of ambient PM exposure with health effects must be adjusted for the effects of other environmental or demographic factors, depending on whether the effects are acute or chronic, in order to quantitatively assess the role that may be attributed to PM exposure. Estimates of PM health effects have shown reasonable quantitative consistency in different studies, with only modest sensitivity to different methods of analysis. However, a clearer understanding of specific biologic mechanisms remains to be more fully established.

- Individuals with cardiovascular or pulmonary disease, especially if they are elderly, are more likely to suffer severe health effects (mortality or hospitalization) related to PM exposure than are healthy young adults. Children and asthmatics are also susceptible to certain PM effects, e.g., increased respiratory symptoms and decreased lung function. Smokers also constitute a population group at increased risk for ambient PM exposure effects.

- In human populations, daily personal exposures to ambient fine particles are reflected by daily ambient fine particle concentrations measured at a community air-monitoring site. This is consistent with the observed high correlations of personal sulfate exposures with ambient sulfate concentrations. Therefore, community air-monitoring site measurements of fine particle indicators can be useful in PM epidemiology studies. The relationship between personal exposure to thoracic coarse particles and the ambient concentration of thoracic coarse fraction particles is not as strong, making detection of effects due to coarse fraction particles harder to detect in epidemiological studies.

- Development of a comprehensive biologically-based exposure-dose-response model to aid health risk assessment requires further dosimetry data characterizing differences among species in percent deposition and regional deposition patterns including differences in inhalability, airway geometry, and clearance rates. More information is also required on mechanism(s) of clearance, pathological processes affecting deposition and clearance of particles, and factors which influence the response(s) of respiratory tract tissues to particle burden.

- The percent deposition and regional patterns of deposition depend strongly on particle size. Percent deposition is higher in smaller lungs (children; women), during exercise, and in the functioning parts of the lungs in people with compromised lungs.

- Estimation of public health impacts of ambient airborne particle exposures in the United States would most credibly require information from exposure-response relationships derived for particular U.S. urban areas, in combination with estimates of exposures to ambient particle concentrations for the general population and/or specific susceptible subgroups (e.g., the elderly) within those particular areas. At the present time, risk assessment is necessarily limited to use of available information from concentration-response relationships relating ambient concentrations to health effects in populations. In view of geographic differences in ambient PM mixtures and demographics, broad generalization and application of some single "best estimate" of relative risk for a given increment in concentration of a given particle indicator (e.g., PM_{10}, PM_{25}, etc.) would be subject to much uncertainty.

- Toxicology studies of PM using controlled inhalation exposure of humans and laboratory animals, intratracheal instillation in humans and animals, and exposure of human and animal cells grown in culture find numerous biological effects which may be related to adverse health effects. Newer studies are finding different biological effects for a variety of different particle components. Newer studies also are beginning to identify biological mechanisms whereby PM deposited in the lung can produce adverse effects on the cardiovascular and respiratory systems.

- Epidemiological studies indicate increased health risks associated with exposure to PM, alone or in combination with other air pollutants. PM-related increases in individual health risks are small, but likely significant from an overall public health perspective because of the large numbers of individuals in susceptible risk groups that are exposed to ambient PM.
- Numerous new studies, including multicity studies, continue to find a consistent association of PM_{10} exposure with mortality and various morbidity endpoints, thus substantiating the relationship of PM exposure with various health effects. However, new studies using $PM_{2.5}$ as an indicator find higher statistical significance and higher excess risk for $PM_{2.5}$ compared to

PM_{10}. Several studies have also observed statistically associations of $PM_{10-2.5}$ with health effects.

- Epidemiologic studies, in which factors identified with source categories or individual chemical components of PM have been used as indicators, also show significant associations with nitrate, sulfate, various indicators of elemental carbon, the organic component of PM, and some elements. Source category factors, attributed to PM associated with regional sulfate, vehicular traffic, vegetative combustion, and oil burning have also been found to be significantly associated with mortality.

- Data from multicity studies, comparisons of effects found in single and multiple regressions, exposure relationships, and toxicity suggest that the gaseous co-pollutants (CO, NO_3, SO_2,O_3) are not responsible for the relationships observed with PM indicators in community, time-series epidemiology. This does not indicate lack of an independent association of gaseous co-pollutants with health effects, but rather suggests that they may be surrogates for PM (due to significant correlations with PM) rather than confounders.

- Fine and thoracic coarse PM, indexed respectively by $PM_{2.5}$ and $PM_{10-2.5}$, should be considered as separate subclasses of PM. Considerations of emissions sources, atmospheric chemistry, physical behavior, exposure relationships, respiratory deposition, toxicologic findings, and epidemiologic observations argue for monitoring fine and thoracic coarse particles separately.

- Assessment of health risk in epidemiologic studies of ambient air pollutants, including PM, has relied largely on studies that focus on changes in health risks that occur in relation to normal changes in ambient air pollutant concentrations. Further evidence of the effects of air pollution on health may be deduced from intervention studies, i.e, studies of changes in health effects that occur when air pollution concentrations have been temporarily or permanently reduced through regulatory action, industrial shutdown, or other intervening factor(s). Only a few epidemiologic intervention studies are available, however, taken together, these studies lend confidence that further reduction of ambient air pollution exposures in the U.S. would benefit public health. It is likely that such reduction would bring about both respiratory and

cardiovascular heath benefits. Available studies also give reason to expect that further reductions in both particulate and gaseous air pollutants would benefit health. On balance, these studies suggest that selective reduction in ambient PM concentrations might well bring about greater benefit than would selective reduction in concentrations of other ambient criteria air pollutants. Furthermore, the experimental studies of Utah Valley filter extracts points to PM-associated metals as a likely cause or promoter of at least some of the health disorders associated with ambient PM.

- The weight of evidence, from exposure, dosimetry, toxicology, and epidemiology, leans toward the conclusion that PM, especially fine PM, is the primary contributor to a variety of adverse health effects associated with air pollution. However, there are difficult technical issues in separating the effects of fine and coarse particles and in separating particle effects from possible effects of gaseous co-pollutants.

1. INTRODUCTION

This document is an update of "Air Quality Criteria for Particulate Matter" published by the U.S. Environmental Protection Agency (EPA) in 1996, and it will serve as the basis for Congressionally-mandated periodic review of the National Ambient Air Quality Standards (NAAQS) for particulate matter (PM). The present document critically assesses the latest scientific information relative to determining the health and welfare effects associated with exposure to various concentrations of PM in ambient air. The document builds upon the assessment in the previous 1996 EPA Air Quality Criteria Document for Particulate Matter by focusing on assessment and integration of information most relevant to PM NAAQS criteria development, based on pertinent literature mainly available through December 2001, as well as some newly emerging studies published in early 2002. This introductory chapter presents a brief summary of legislative requirements and history of the PM NAAQS, provides an overview of issues addressed and procedures utilized in the preparation of the present document, and provides orientation to the general organizational structure of this document.

1.1 LEGISLATIVE REQUIREMENTS

As indicated in U.S. Code (1991), the U.S. Clean Air Act (CAA), Sections 108 and 109 (42 U.S.C. Sections 7408 and 7409) govern the establishment, review, and revision of National Ambient Air Quality Standards (NAAQS). Section 108a directs the EPA Administrator to list pollutants, which, in the Administrator's judgement, cause or contribute to air pollution which may reasonably be anticipated to endanger either public health or welfare and which are emitted by numerous (or diverse) stationary or mobile sources. The air quality criteria are to reflect the latest scientific information useful in indicating the kind and extent of exposure-related effects on public health and welfare that may be expected from the presence of the pollutant in ambient air.

Section 109(a) and (b) directs the Administrator of EPA to propose and promulgate "primary" and "secondary" NAAQS for pollutants identified under Section 108. Section 109(b)(1) defines a primary standard as a level of air quality, the attainment and maintenance of

which, in the judgement of the Administrator, based on the criteria and allowing for an adequate margin of safety, is requisite to protect the public health. A secondary standard, in contrast, is one which, again in the Administrator's judgement, is requisite to protect public welfare from any known or anticipated adverse effects associated with the presence of such pollutants. Welfare effects include effects on vegetation, crops, soils, water, animals, manufactured materials, visibility, weather, and climate, as well as damage to and deterioration of property, hazards to transportation, and effects on economic value and personal comfort and well-being (as per CAA Section 302h). Section 109(d) also requires periodic review and, as appropriate, revision of existing criteria and standards; and it requires an independent committee of non-EPA experts, the Clean Air Scientific Advisory Committee (CASAC), to provide advice and recommendations to the EPA Administrator regarding the scientific soundness and appropriateness of criteria and NAAQS for PM and other "criteria air pollutants" (e.g., ozone, nitrogen oxides, sulfur oxides, carbon monoxide, lead) regulated under CAA Sections 108-109.

1.2 HISTORY OF PREVIOUS PM CRITERIA AND NAAQS REVIEWS

"Particulate matter" is the generic term for a broad class of physically and chemically diverse substances that exist in ambient air as discrete particles (liquid droplets or solids) over a wide range of sizes. These airborne particles originate from a variety of stationary and mobile sources. Primary particles are emitted directly into ambient air; whereas secondary particles are formed in the atmosphere by transformation of gaseous emissions such as sulfur oxides (SO_x), nitrogen oxides (NO_x), and volatile organic compounds (VOCs). The physical and chemical properties of PM vary greatly with time, region, meteorology, and source category, thus complicating assessment of ambient PM health and welfare effects. Particles in ambient air are usually distributed bimodally in two somewhat overlapping size categories: (1) fine (diameter generally less than 2.5 μm) and (2) coarse (diameter generally greater than 2.5 μm). Particles in these two size fractions tend to differ in terms of sources of origin, composition, and behavior in the atmosphere and human respiratory tract.

EPA first promulgated primary and secondary NAAQS for PM on April 30, 1971 (Federal Register, 1971). These standards measured PM as "total suspended particulate" (TSP), which refers to ambient PM up to a nominal size of 25 to 45 micrometers (μm). The primary standards

for PM (measured as TSP) were 260 $\mu g/m^3$ (24-h average), not to be exceeded more than once per year, and 75 $\mu g/m^3$ (annual geometric mean). The secondary standard (measured as TSP) was 150 $\mu g/m^3$ (24-h average), not to be exceeded more than once per year.

EPA completed the next review of PM air quality criteria and standards in July 1987, revising the 1971 standards to protect against adverse health effects of inhalable airborne particles with an upper 50% cut-point of 10-μm aerodynamic diameter (PM_{10}), which can be deposited in the lower (thoracic) regions of the human respiratory tract (Federal Register, 1987). EPA established identical primary and secondary PM_{10} standards for two averaging times: 150 $\mu g/m^3$ (24-h average), with no more than one expected exceedance per year and 50 $\mu g/m^3$ (expected annual arithmetic mean), averaged over three years.

1.2.1 The 1997 PM NAAQS Revision

The last previous review of the air quality criteria and standards for PM was initiated in April 1994 by EPA announcing its intention to develop revised Air Quality Criteria for Particulate Matter (henceforth, the "PM Air Quality Criteria Document" or PM AQCD). Several workshops were held by EPA's Environmental Criteria and Assessment Office in Research Triangle Park, NC (ECAO-RTP) in November 1994 and January 1995 to discuss important new health effects information useful in preparing initial PM AQCD draft materials. Also, plans for review of the PM criteria and standards under a highly accelerated, court-ordered schedule were presented by EPA at a public meeting of the CASAC in December 1994. A court order entered in *American Lung Association v. Browner,* CIV-93-643-TUC-ACM (U.S. District Court of Arizona, 1995), as subsequently modified, required publication of EPA's final decision on the review of the PM NAAQS by July 19, 1997.

Several external review drafts of the revised PM AQCD were prepared by the RTP Division of EPA's newly created National Center for Environmental Assessment (i.e., by NCEA-RTP, the successor office to ECAO-RTP), and each were made available for public comment followed by CASAC review (at public meetings held in August 1995, December 1995, and February 1996). The CASAC came to closure on its review of the PM AQCD in early 1996, advising the EPA Administrator in a March 15, 1996 closure letter (Wolff, 1996) that "although our understanding of the health effects of PM is far from complete, a revised Criteria Document which incorporates the Panel's latest comments will provide an adequate review of the available

scientific data and relevant studies of PM." Revisions made in response to public and CASAC comments were then incorporated by NCEA-RTP, as appropriate, into the final 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). A PM Staff Paper (SP), prepared by the Office of Air Quality Planning and Standards (OAQPS) within EPA's Office of Air and Radiation (OAR), drew upon the 1996 PM AQCD and other exposure and risk assessments to pose options for PM NAAQS decisions. The PM SP also underwent public comment and CASAC review, with consequent revision to its July 1996 final form (U.S. Environmental Protection Agency, 1996b). The SP analyses served as key inputs to subsequently published proposals (Federal Register, 1996) for revision of the primary PM NAAQS.

Taking into account information and assessments presented in the 1996 PM AQCD and PM SP, advice and recommendations of CASAC, and public comments received on the proposed revisions, the EPA Administrator revised the PM NAAQS by adding new $PM_{2.5}$ standards and by revising the form of the 24-h PM_{10} standard (Federal Register, 1997a). Specifically, on July 18, 1997, the Administrator promulgated the following revisions to the PM NAAQS:

(1) The suite of PM standards was revised to include an annual primary $PM_{2.5}$ standard and a 24-h $PM_{2.5}$ standard.

(2) The 24-h $PM_{2.5}$ standard is met when the 3-year average of the 98th percentile of 24-h $PM_{2.5}$ concentrations at each population-oriented monitor within an area is less than or equal to 65 $\mu g/m^3$, with fractional parts of 0.5 or greater rounding up.

(3) The annual $PM_{2.5}$ standard is met when the 3-year average of the annual arithmetic mean $PM_{2.5}$ concentrations, from single or multiple community-oriented monitors, is less than or equal to 15 $\mu g/m^3$, with fractional parts of 0.05 or greater rounding up.

(4) The form of the 24-h PM_{10} (150 $\mu g/m^3$) standard was revised to be based on the 3-year average of the 99th percentile of 24-h PM_{10} concentrations at each monitor within an area.

(5) In addition, the Administrator retained the annual PM_{10} standard at the level of 50 $\mu g/m^3$, which is met when the 3-year average of the annual arithmetic mean PM_{10} concentrations at each monitor within an area is less than or equal to 50 $\mu g/m^3$, with fractional parts of 0.5 or greater rounding up.

The principal focus of the last review of the air quality criteria and standards for PM was on recent epidemiological evidence reporting associations between ambient concentrations of PM and a range of serious health effects. Special attention was given to several size-specific classes

of particles, including PM_{10} and the principal fractions of PM_{10}, referred to as the fine ($PM_{2.5}$) and coarse ($PM_{10-2.5}$) fractions. $PM_{2.5}$ refers to particles with an upper 50% cutpoint of 2.5 μm aerodynamic diameter. $PM_{10-2.5}$ refers to those particles with an upper 50% cutpoint of 10 μm and a lower 50% cut point of 2.5 μm aerodynamic diameter. In other words, the coarse fraction ($PM_{10-2.5}$) refers to the inhalable particles that remain if fine ($PM_{2.5}$) particles are removed from a sample of PM_{10} particles. As discussed in the 1996 PM AQCD, fine and coarse fraction particles can be differentiated by their sources and formation processes and by their chemical and physical properties, including behavior in the atmosphere. Detailed discussions of atmospheric formation, ambient concentrations, and health effects of ambient air PM, as well as quantitative estimates of human health risks associated with exposure to ambient air PM, can be found in the 1996 PM AQCD and the 1996 OAQPS SP (U.S. Environmental Protection Agency, 1996a,b).

1.2.2 Coordinated Particulate Matter Research Program

Shortly after promulgation of the 1997 PM NAAQS decisions, NCEA-RTP published a PM Health Risk Research Needs Document (U.S. Environmental Protection Agency, 1998a) that identified research needed to improve scientific information supporting future reviews of the PM NAAQS. The document aimed to provide a foundation for PM research coordination among Federal agencies and other research organizations and provided input to later National Research Council (NRC) PM research deliberations. The Office of Research and Development (ORD) of EPA also moved quickly to broaden its ongoing PM research activities by developing, in partnership with other Federal agencies, a coordinated interagency PM research program. This interagency program has and continues to focus mainly on expanding scientific knowledge of ambient PM exposure and health effects, as well as including development of improved monitoring methods and cost-effective mitigation strategies. The interagency effort also promotes substantially expanded coordination with other research organizations, including the Health Effects Institute (HEI) and other state-, university-, and industry-sponsored research groups. Beginning in the fall of 1997, public participation was and continues to be encouraged through workshops and review of program documentation.

In response to Congressional requirements in EPA's Fiscal Year 1998 Appropriation, the NRC established its Committee on Research Priorities for Airborne Particulate Matter in January 1998. This NRC PM Research Committee's charge is to identify the most important research

priorities relevant to setting particulate matter standards, to develop a conceptual plan for PM research, and to monitor research progress toward improved understanding of the relationship between PM and public health. The Committee issued its first report in early 1998 (National Research Council, 1998), a second one in 1999 (National Research Council, 1999), and a third one in 2001 (National Research Council, 2001).

In the above-noted series of reports, the NRC PM Research Committee recommended that expanded PM research efforts be planned and carried out in relation to a general conceptual framework as shown in Figure 1-1. That framework essentially calls for research aimed at: (a) identifying sources of airborne particles or gaseous precursor emissions and characterization of processes involved in atmospheric transformation, transport, and fate of ambient PM; (b) delineation of temporal and spatial patterns of air quality indicators (e.g., $PM_{2.5}$, $PM_{10-2.5}$, PM_{10} mass concentrations) of ambient PM and apportionment of observed variations in such ambient PM indicators to various emission sources; (c) characterization of human exposures to ambient PM as one important component of total personal exposure to particles, as modified by time-activity patterns and varying microenvironmental exposure to particles of indoor or ambient origin; (d) characterization of resulting respiratory tract deposition, clearance, retention, and disposition of inhaled particles, as determinants of dose to target tissues (e.g., locally in the lungs or via systemic translocation to the heart or other organs); (e) delineation of mechanisms of damage and repair plausibly leading to (f) human health responses, as extrapolated from or quantified by experimental animal or human exposure (toxicology) studies and/or observational (epidemiology) studies.

Research conducted under a PM Research Program structured in relation to the conceptual framework shown in Figure 1-1 would be expected (a) to reduce key scientific uncertainties regarding interrelationships between PM sources, ambient concentrations, exposures, dose to target tissues, and resulting health effects and (b) thereby improve the scientific underpinnings for both current and future periodic PM criteria/NAAQS reviews. Table 1-1 highlights some types of key uncertainties identified by the NRC PM Research Committee in relation to elements of the source-to-response conceptual framework illustrated in Figure 1-1. The NRC Committee went on to delineate a series of 10 research topics that they recommended be addressed in an expanded PM research program aimed at answering a set of broadly stated questions, as shown in Table 1-2.



Figure 1-1. A general framework for integrating particulate-matter research. Note that this figure is not intended to represent a framework for research management. Such a framework would include multiple pathways for the flow of information.

Source: National Research Council (2001), as modified from NRC (1983, 1994), Lioy (1990), and Sexton et al. (1992).

The EPA's PM Research Program is structured to address the topics shown in Table 1-2; and it includes, for example, studies to improve understanding of the formation and composition of fine PM, improved measurements and estimation of population exposures to ambient PM, the characteristics or components of PM that are responsible for its health effects, and the mechanisms by which these effects are produced. Specific EPA research efforts include controlled human exposure studies, in vivo and in vitro toxicology, epidemiology, atmospheric sciences including monitoring and modeling studies, development of data on emissions of fine particles from stationary and mobile sources, and identification and evaluation of risk management options. The results from these efforts, as well as related efforts by other Federal agencies and the general scientific community during the past several years, have substantially enhanced the scientific and technical bases for future decisions on the PM NAAQS, as well as for implementation of PM monitoring and control efforts that are beyond the scope of this document.

TABLE 1-1. KEY SCIENTIFIC UNCERTAINTIES RELATED TO THE SOURCE-TO-RESPONSE FRAMEWORK

Source → Concentration (or other indicator)
• Contribution of various emission sources to ambient and indoor particulate matter concentrations • Relative contribution of various sources to the most toxic components of particulate matter
Concentration (indicator) → Exposure
• Relationship between ambient (indoor) particulate matter and the composition of particles to which people are exposed • Contribution of ambient particulate matter to total personal exposure for: - Susceptible subpopulations - General population • Variation in relationship of ambient particulate matter concentrations to human exposure by place • Variation in contribution of ambient particulate matter concentrations to total human exposure over time • Covariance of particulate matter exposures with exposures to other pollutants • Relationships between outdoor ambient and personal exposures for particulate matter and copollutants
Exposure → Dose
• Relationship between inhaled concentration and dose of particulate matter and constituents at the tissue level in susceptible subjects - Asthma - Chronic obstructive pulmonary disease (COPD) - Heart disease - Age: infants and elderly - Others
Dose → Response
• Mechanisms linking morbidity and mortality to particulate matter dose to or via the lungs - Inflammation - Host defenses - Neural mechanisms

Source: National Research Council (2001).

TABLE 1-2. RESEARCH TOPICS AND QUESTIONS RECOMMENDED BY NATIONAL RESEARCH COUNCIL (NRC) TO BE ADDRESSED BY EXPANDED PM RESEARCH PROGRAM

RESEARCH TOPIC 1. OUTDOOR MEASURES VERSUS ACTUAL HUMAN EXPOSURES

- *What are the quantitative relationships between concentrations of particulate matter and gaseous copollutants measured at stationary outdoor air monitoring sites and the contributions of these concentrations to actual personal exposures, especially for subpopulations and individuals?*

RESEARCH TOPIC 2. EXPOSURES OF SUSCEPTIBLE SUBPOPULATIONS TO TOXIC PARTICULATE MATTER COMPONENTS

- *What are the exposures to biologically important constituents and specific characteristics of particulate matter that cause responses in potentially susceptible subpopulations and the general population?*

RESEARCH TOPIC 3. CHARACTERIZATION OF EMISSION SOURCES

- *What are the size distribution, chemical composition, and mass-emission rates of particulate matter emitted from the collection of primary-particle sources in the United States, and what are the emissions of reactive gases that lead to secondary particle formation through atmospheric chemical reactions?*

RESEARCH TOPIC 4. AIR-QUALITY MODEL DEVELOPMENT AND TESTING

- *What are the linkages between emission sources and ambient concentrations of the biologically important components of particulate matter?*

RESEARCH TOPIC 5. ASSESSMENT OF HAZARDOUS PARTICULATE MATTER COMPONENTS

- *What is the role of physicochemical characteristics of particulate matter in eliciting adverse health effects?*

RESEARCH TOPIC 6. DOSIMETRY: DEPOSITION AND FATE OF PARTICLES IN THE RESPIRATORY TRACT

- *What are the deposition patterns and fate of particles in the respiratory tract of individuals belonging to presumed susceptible subpopulations?*

RESEARCH TOPIC 7. COMBINED EFFECTS OF PARTICULATE MATTER AND GASEOUS POLLUTANTS

- *How can the effects of particulate matter be disentangled from the effects of other pollutants? How can the effects of long-term exposure to particulate matter and other pollutants be better understood?*

RESEARCH TOPIC 8. SUSCEPTIBLE SUBPOPULATIONS

- *What subpopulations are at increased risk of adverse health outcomes from particulate matter?*

RESEARCH TOPIC 9. MECHANISMS OF INJURY

- *What are the underlying mechanisms (local pulmonary and systemic) that can explain the epidemiological findings of mortality/morbidity associated with exposure to ambient particulate matter?*

RESEARCH TOPIC 10. ANALYSIS AND MEASUREMENT

- *To what extent does the choice of statistical methods in the analysis of data from epidemiological studies influence estimates of health risks from exposures to particulate matter? Can existing methods be improved? What is the effect of measurement error and misclassification on estimates of the association between air pollution and health?*

Source: National Research Council (2001).

1.3 CURRENT PM CRITERIA AND NAAQS REVIEW

1.3.1 Key Milestones

As with other NAAQS reviews, a rigorous assessment of relevant scientific information is to be presented in this updated, revised PM AQCD. As shown in Table 1-3, development of the document has involved substantial external peer review through (a) public workshops involving the general aerosol scientific community, (b) iterative reviews of successive drafts by CASAC, and (c) comments from the public. The final document will reflect input received through these reviews and will serve to evaluate and integrate the latest available scientific information to ensure that the review of the PM standards is based on rigorous evaluation of the available science. An earlier (October 1999) First External Review Draft of this updated document was released in the fall of 1999 for public comment and CASAC review. A Second External Review Draft (March 2001) took into account the earlier public comments and the December 1999 CASAC review. This Third External Review Draft similarly takes into account prior public comments and CASAC recommendations from its July 2001 review, and it includes consideration of relevant new peer-reviewed scientific studies published or accepted for publication mainly through December 2001, as well as some newly emerging key studies published in early 2002. Following a 60-day public comment period, it is to be reviewed by CASAC at a public meeting in July 2002.

After CASAC review of the First External Review Draft of this revised PM AQCD in December 1999, EPA's OAQPS started to prepare the associated PM Staff Paper. A preliminary draft SP was made available to the public and CASAC for review at their July 2001 meeting. The next draft PM SP will, to the extent possible, draw on the updated findings and conclusions from this Third Draft of the PM AQCD and will also undergo further public comment and CASAC review (scheduled for September 2002). Ultimately drawing on information in the final version of this newly revised PM AQCD, the PM SP will evaluate policy implications of the key studies and scientific findings contained in the AQCD, present related staff analyses of air quality and human health risk, and identify critical elements that EPA staff believes should be considered in reviewing the PM standards. The PM SP is intended to bridge the gap between the scientific review in the AQCD and the public health and welfare policy judgements required of

TABLE 1-3. SCHEDULE FOR DEVELOPMENT OF THE CURRENT REVISED PARTICULATE MATTER AIR QUALITY CRITERIA DOCUMENT (PM AQCD)

Major Milestones	Target Dates
PM NAAQS Review Plan to CASAC	October 1997
Prepare AQCD Development Plan	November 1997 to January 1998
Begin Literature Search	February 1998
Federal Register Call for Information/Sources Sought	April 1998
CASAC Meeting on AQCD Development Plan	May 1998
Prepare Workshop Drafts of Chapters	May to December 1998
Peer Review Workshop	April 1999
Prepare External Review Draft AQCD	March to September 1999
First External Review Draft	October 1999
Public Comment Period on Draft AQCD	October 1999 to January 2000
CASAC Meeting on First Draft AQCD	December 1999
Second External Review Draft	March 2001
Public Comment Period on Second Draft	April to July 2001
CASAC Meeting on Second Draft	July 2001
Third External Review Draft	April 2002
Public Comment Period on Third Draft	May to July 2002
CASAC Meeting on Third Draft	July 2002
Final PM AQCD	Oct/Nov 2002

the Administrator in reviewing the PM NAAQS. In doing so, the PM SP will include staff conclusions and recommendations of options for the Administrator's consideration.

Based on the final versions of the PM AQCD and the PM SP and on the advice of CASAC, the Administrator will propose to retain or revise the current PM NAAQS and provide opportunities for public comment and CASAC review of the proposed decisions. Taking into account public comments and CASAC advice, the Administrator will then make final decisions on the current PM NAAQS, which are now expected to be issued by December 31, 2003.

1.3.2 Methods and Procedures for Document Preparation

The procedures for developing this revised PM AQCD build on the experience and methods derived from the most recent previous PM, Ozone, and CO AQCD preparation efforts. Briefly, the respective responsibilities for production of the present PM AQCD are as follows. An NCEA-RTP PM team was formed and is responsible for developing and implementing the project plan for preparation of the PM AQCD, taking into account inputs from individuals in other EPA program and policy offices identified as part of the EPA PM Work Group. The resulting project plan (i.e., the PM Document Development Plan) was then discussed with CASAC (May 1998) and appropriately revised. An ongoing literature search has continued to be conducted to identify, to the extent possible, all pertinent PM literature published since early 1996. Additionally, EPA published (1) a request for information in the Federal Register asking for recently available research information on PM that may not yet be published and (2) a request for individuals with the appropriate type and level of expertise to contribute to the writing of PM AQCD materials to identify themselves (U.S. Environmental Protection Agency, 1998b). Specific authors of chapters or sections of the proposed document were selected on the basis of their expertise on the subject areas and their familiarity with the relevant literature; these include both EPA and non-EPA scientific experts. The project team defined critical issues and topics to be addressed by the authors and provided direction in order to emphasize evaluation of those studies most clearly identified as important for standard setting. It should be noted that materials contributed by non-EPA authors are incorporated and, at times, modified by EPA PM team staff to reflect internal and/or external review comments, e.g., by the public or CASAC, and that EPA is responsible for the ultimate content of the PM AQCD.

The main focus of this revised criteria document is the evaluation and interpretation of pertinent atmospheric science information, air quality data, human exposure information, and health and welfare effects information newly published since that assessed in the 1996 PM AQCD and likely to be useful in deriving criteria for PM NAAQS. Initial draft versions of AQCD chapters were evaluated via expert workshops and/or expert written peer reviews that focused on the selection of pertinent studies included in the chapters, the potential need for additional information to be added to the chapters, and the quality of the summarization and interpretation of the literature. The authors of the draft chapters then revised them on the basis of the workshop and/or written expert review recommendations. These and other integrative

summary materials were incorporated into the First External Review Draft of the PM AQCD (October 1999), which was released for public comment and reviewed at a December 1999 CASAC public meeting.

In order to foster timely presentation and publication of newly emerging PM research findings, EPA co-sponsored an Air and Waste Management Association International Speciality Conference, entitled "PM 2000: Particulate Matter and Health", which was held in January 2000 in Charleston, SC. The conference was co-sponsored in cooperation with several other government agencies and/or private organizations that also fund PM research. Topics covered included new research results concerning the latest advances in PM atmospheric sciences (e.g., PM formation, transport, transformation), PM exposure, PM dosimetry and extrapolation modeling, PM toxicology (e.g., mechanisms, laboratory animal models, human clinical responses), and PM epidemiology. The main purpose of the conference was to facilitate having the latest scientific information available in time for incorporation as quickly as possible into the Second External Review Draft of this revised PM AQCD. Hence, arrangements were made for scientists to submit written manuscripts on papers or posters presented at the PM 2000 Conference for expedited peer-review by several major journals, so that decisions on acceptance for publication could be made by mid-2000. The evaluations and findings set forth in the Second External Review Draft (March 2001) of the revised PM AQCD included consideration of such published PM 2000 papers and extensive additional information published elsewhere since the previous First External Review Draft; it also reflected public and CASAC comments on that First Draft. The Second External Review Draft (March 2001) was then reviewed by CASAC in July 2001. Further revisions incorporated into this Third External Review Draft (April 2002) reflect both public comment and CASAC review of the Second Draft, as well as assessment of additional pertinent information published since that addressed in the Second Draft. The final version of the newly revised PM AQCD will then incorporate changes made in response to public comments and CASAC review of this Third External Review Draft.

1.3.3 Approach

The approach to development of this revised PM AQCD is somewhat different from that used for previous criteria documents. Because the most recent prior document the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) provides an extensive discussion of most

topic areas, this new document focuses more specifically on critical issues that have been identified by the NRC as needing more research in order to improve the scientific bases (criteria) for PM NAAQS (see Section 1.2.2. above).

An initial step was to focus on selection of pertinent issues to include in the document as the basis for the development of PM NAAQS criteria. Preliminary issues were identified by the NCEA PM Team via input from other EPA program and policy offices, as derived from the 1996 PM AQCD and SP, their CASAC and public reviews, from the 1997 standard promulgation process, and from the 1998 PM Research Needs Document (alluded to in Section 1.2.2). Still, further identification and refinement of issues resulted from NRC review and reports on PM research priorities (also discussed in Section 1.2.2 above). The CASAC review of the PM AQCD Development Plan and public comments on earlier draft AQCD materials at various stages of their development have also played important roles in issue identification.

In developing draft materials for inclusion in the revised PM AQCD, detailed review of key new research was undertaken as a first step. However, instead of presenting a comprehensive review of all the literature, emphasis in this revised AQCD is placed on (1) first, the concise summary of key findings derived from previous PM criteria reviews and, then, (2) summarization and evaluation of the most pertinent new key information, with greater emphasis on more interpretive assessment—an approach reflecting CASAC recommendations. To aid in the development of a concise document, compilation of summary tables of relevant new literature published since completion of the previous 1996 PM AQCD and selective text discussion of that literature has been undertaken, with increased emphasis being placed in text discussions on interpretive evaluation and integration of key points derived from the newly summarized research results.

1.3.4 Key Human Health Issues of Concern

The present document reviews and assesses available data bearing on each of the broad topics or issues identified below:

(1) *Causality*. Evaluation of the evidence for or against a causal relationship between health outcomes and ambient PM and/or specific PM physical-chemical components.

- Specific components of interest include: (a) size classes, such as PM_{10}, $PM_{10-2.5}$, $PM_{2.5}$, and ultrafine particles, and (b) chemical components, such as transition metals, acidity, sulfates, nitrates, and organics.
- Expand review of foundations of causal inference for associated PM air pollution health effects.
- Assess new long-term PM exposure and health data to broaden interpretation of long-term exposure findings.
- Review data exploring potential mechanisms of response to PM physical-chemical characteristics, response pathways, and exposure-dose-response relationships (laboratory and clinical research).

(2) *Uncertainties.* In carrying out overall assessment, address the following types of uncertainty.

- Uncertainties between stationary PM monitoring instruments and personal exposure to PM of ambient origin, especially for susceptible groups and their related activity patterns. Specific topics include measurement error in outdoor monitors themselves, use of central monitors for estimates of community concentrations, and the use of community concentrations as a surrogate for personal exposure to particles of ambient origin.
- Uncertainties related to particulate matter size fraction, particle number, surface area, and content of semivolatile components.
- Uncertainties about the effects of long-term PM exposure, such as life shortening, and development and progression of disease.
- Uncertainties because of coexposure to other pollutants such as O_3, SO_2, CO, and NO_2, and because of meterological factors.
- Uncertainties because of potential confounding in epidemiologic studies (e.g., economic factors, demographic and lifestyle attributes, genetic susceptibility factors, occupational exposure, medical care).
- Uncertainty about shape of concentration-response (CR) relationships and associated community risks (linear and threshold models for CR).
- Uncertainty about methods for synthesis of health outcome studies and evaluation of sensitivity and confounding aspects, including but not limited to meta-analyses.

(3) *Biological Mechanisms of Action.* Evaluate data examining mechanisms underlying health outcomes of PM. Mechanistic information aids judgment about causality.

- New studies have examined mechanisms of action of PM constituents, including transition metals, airborne allergens, and the generation of reactive oxygen species. Different cell types have differing responses to PM components.
- Newly published studies also have identified potential mechanisms for the production of cardiac arrhythmias by PM constituents, especially in animal models of disease and suggest that particular attention should be accorded to PM metal constituents.
- Although many new animal toxicology studies involve instillation in the lung of previously collected particles and this technique is appropriate to study mechanisms of action, extrapolation to human equivalent exposure/doses is uncertain.
- Ongoing work on the effects of lung inflammation and PM phagocytosis on subsequent systemic effects, especially cardiac or vascular effects, is needed to provide further information on the relationship between inhaled pollutants and cardiac events.
- Interpretation of concentrated ambient particles (CAPs) studies. Newly available information is examined from toxicology studies using devices that concentrate (to variable extents) ambient PM to determine PM concentration-response relationships. Again, difficulties exist with regard to quantitative extrapolation to comparable human exposures to ambient PM.

(4) *Susceptible Populations.* Examine health outcome data to determine specific risk groups that are more susceptible than normal healthy adults to adverse effects from PM exposure.

- Preexisting respiratory or cardiovascular disease in conjunction with advanced age appear to be important factors in PM mortality susceptibility.
- For morbidity health endpoints, children and asthmatics potentially may display increased sensitivity to PM exposure. Data will be examined for coherence.
- Patterns of respiratory tract deposition, clearance, and retention in susceptible populations have been studied recently and provide evidence of differences in respiratory tract PM deposition for children and small-sized adults and for those with lung diseases.
- Animal models of lung disease exposed to PM constituents suggest a role for PM in cardiac death.

(5) *Environmental Effects.* Evaluate several types of PM welfare effects, as follows.

- Vegetation and ecosystem effects.
- Visibility effects.
- Materials damage.
- Role of PM in atmospheric radiative transfer and potential consequences for penetration of biologically harmful UVB to the earth's surface and for climate change.

(6) *Background Information Topics Useful in Evaluating Health Risks.*

- New monitoring methods, especially methods used in epidemiology studies.
- Indicator topics such as $PM_{2.5}$ versus $PM_{1.0}$; ultrafine; and $PM_{2.5}$ versus $PM_{10-2.5}$.
- New data patterns of daily and annual concentrations for $PM_{2.5}$, $PM_{10-2.5}$, and PM_{10}.

1.4 DOCUMENT ORGANIZATION AND CONTENT

The present draft document attempts to critically review and assess relevant scientific literature on PM mainly through December 2001, but does include some more recent studies published in early 2002. The material selected for review and comment in the text generally comes from the more recent literature published since early 1996, with emphasis on studies conducted at or near PM pollutant concentrations found in ambient air. Literature discussed in detail in the previous 1996 EPA PM AQCD (U.S. Environmental Protection Agency, 1996a) generally is not discussed in depth in this document. However, some limited treatment is included of the earlier studies judged to be potentially useful in deriving PM NAAQS. Key literature is presented mainly in tables; and the text mainly discusses overall interpretive points.

Primary emphasis is placed on consideration of published material that has undergone scientific peer review. However, in the interest of admitting new and important information expected to become available shortly, some material not yet fully published in the open literature but meeting other standards of scientific reporting (i.e., peer review, quality assurance) are now provisionally included. As noted earlier, emphasis has been placed on studies in the range of current ambient levels. However, studies examining effects of higher concentrations have been included if they contain unique data or documentation of a previously unreported effect or mechanism. In reviewing and summarizing the literature, an attempt has been made to present alternative points of view where scientific controversy exists.

The present document is basically organized to assess information related to topics along the same flow of issues presented in the NRC conceptual framework shown in Figure 1-1. It includes the Executive Summary and nine chapters presented in two volumes. Volume 1 contains the Executive Summary and this general introduction (Chapter 1). It also includes Chapters 2 and 3, which provide background information on physical and chemical properties of PM and related compounds; sources and emissions; atmospheric transport, transformation, and fate of PM; methods for the collection and measurement of PM; and U.S. ambient air PM concentrations. Next, Chapter 4 assesses PM environmental effects on vegetation and ecosystems, visibility, man-made materials, and climate-related effects (including effects on solar radiation), and includes limited information on economic impacts of some such welfare effects. Also included in Volume 1 is Chapter 5, which discusses factors affecting exposure of the general population to ambient PM.

The second volume contains Chapters 6 through 9. Chapters 6 evaluates information concerning dosimetry of inhaled particles in the respiratory tract. Chapter 7 assesses the toxicology of specific types of PM constituents and potential mechanisms of action, based on both laboratory animal studies and controlled human exposure studies. Chapter 8 discusses observational, i.e., epidemiological, studies. Lastly, Chapter 9 integrates key information on exposure, dosimetry, and critical health risk issues derived from studies reviewed in the prior chapters. That Integrative Synthesis chapter is basically organized in a manner to address the series of 10 issues (and, where appropriate, subissues) identified in the NRC PM Research Priorities Reports (National Research Council, 1998, 1999, 2001).

REFERENCES

Federal Register. (1971) National primary and secondary ambient air quality standards. F. R. (April 30) 36: 8186-8201.

Federal Register. (1987) Revisions to the national ambient air quality standards for particulate matter. F. R. (July 1) 52: 24,634-24,669.

Federal Register. (1996) National ambient air quality standards for particulate matter; proposed rule. F. R. (November 27) 61: 65638.

Federal Register. (1997a) National ambient air quality standards for particulate matter; final rule. F. R. (July 18) 62: 38,652-38,752.

Federal Register. (1997b) National ambient air quality standards for ozone; final rule. F. R. (July 18) 62: 38,856-38,896.

Federal Register. (1997c) Implementation of revised air quality standards for ozone and particulate matter. F. R. (July 18) 62: 38,421-38,422.

National Research Council. (1998) Research priorities for airborne particulate matter. I. Immediate priorities and a long-range research portfolio. Washington, DC: National Academy Press.

National Research Council. (1999) Research priorities for airborne particulate matter. II. Evaluating research progress and updating the portfolio. Washington, DC: National Academy Press.

National Research Council. (2001) Research priorities for airborne particulate matter. III. Early research progress. Washington, DC: National Academy Press.

U.S. Code. (1991) Clean Air Act, §108, air quality criteria and control techniques, §109, national ambient air quality standards. U. S. C. 42: §§7408-7409.

U.S. District Court of Arizona. (1995) American Lung Association v. Browner. West's Federal Supplement 884 F.Supp. 345 (No. CIV 93-643 TUC ACM).

U.S. Environmental Protection Agency. (1996a) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.

U.S. Environmental Protection Agency. (1996b) Review of the national ambient air quality standards for particulate matter: policy assessment of scientific and technical information. OAQPS staff paper. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA/452/R-96-013. Available from: NTIS, Springfield, VA; PB97-115406REB.

U.S. Environmental Protection Agency. (1998a) Particulate matter research needs for human health risk assessment to support future reviews of the national ambient air quality standards for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment; report no. EPA/600/R-97/132F.

U.S. Environmental Protection Agency. (1998b) Review of national ambient air quality standards for particulate matter. Commer. Bus. Daily: February 19. Available: http://cbdnet.access.gpo.gov/index.html [1999, November 24].

Wolff, G. T. (1996) Closure by the Clean Air Scientific Advisory Committee (CASAC) on the staff paper for particulate matter [letter to Carol M. Browner, Administrator, U.S. EPA]. Washington, DC: U.S. Environmental Protection Agency, Clean Air Scientific Advisory Committee; EPA-SAB-CASAC-LTR-96-008; June 13.

2. PHYSICS, CHEMISTRY, AND MEASUREMENT OF PARTICULATE MATTER

An extensive review of the physics and chemistry of particulate matter (PM) was included in Chapter 3 of the 1996 EPA document Air Quality Criteria for Particulate Matter (U.S. Environmental Protection Agency, 1996a). Chapter 2 of this new version of the PM Air Quality Criteria Document (PM AQCD) provides background information on the physics and chemistry of atmospheric particles that may be useful in reading subsequent sections and chapters. This PM AQCD follows the Risk Assessment Framework, as described in Chapter 1 (Section 1.2.2), beginning with sources and continuing to effects as shown in Figure 1-1. However, particular matter, unlike the other criteria pollutants (O_3, CO, NO_2, and Pb), is not a specific chemical entity but is a mixture of particles of different sizes, compositions, and properties. Therefore, it will be useful to present some background on the chemistry and physics of PM before entering the Risk Assessment Framework. This chapter includes new information that should be useful in understanding risk assessments of the effects of PM on human health and welfare. Emphasis is placed on differences between fine and coarse particles and differences between the nuclei mode and the accumulation mode within fine particles. PM information important for implementation of a standard, but not essential to the standard setting process, is not covered in this chapter. The reader is referred to the NARSTO Fine Particle Assessment (NARSTO, 2002) for information relevant to air quality management for PM.

PM is defined quantitatively by the measurement techniques. Therefore, before entering the Risk Assessment Framework, it will also be useful to discuss our understanding of the relationship between PM suspended in the atmosphere, PM inhaled by people, and PM measured by various sampling and analytical techniques. Chapter 4 of the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) contained a review of the state-of-the-art of PM measurement technology. Since that time, considerable progress has been made in understanding problems in the measurement of PM mass, chemical composition, and physical parameters. There also has been some progress in developing new and improved measurement techniques, especially for continuous measurements. Therefore, a more extensive survey on measurement

problems and on newly developed measurement techniques is included in Section 2.2. For more detail and older references, the reader is referred to Chapters 3 and 4 of the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a).

2.1 PHYSICS AND CHEMISTRY OF PARTICULATE MATTER

2.1.1 Basic Concepts

Atmospheric particles originate from a variety of sources and possess a range of morphological, chemical, physical, and thermodynamic properties. Examples include combustion-generated particles, such as diesel soot or fly ash; photochemically produced particles, such as those found in urban haze; salt particles formed from sea spray; and soil-like particles from resuspended dust. Some particles are liquid; some are solid. Others may contain a solid core surrounded by liquid. Atmospheric particles contain inorganic ions, metallic compounds, elemental carbon, organic compounds, and crustal compounds. Some atmospheric particles are hygroscopic and contain particle-bound water. The organic fraction is especially complex, containing hundreds of organic compounds. (Note: See Appendix 3C for information on the composition of the organic fraction and the concentration of specific organic compounds.) Primary particles are emitted directly from sources. Secondary particles are formed from gases through chemical reactions in the atmosphere involving atmospheric oxygen (O_2) and water vapor (H_2O); reactive species such as ozone (O_3); radicals such as the hydroxyl (•OH) and nitrate (•NO_3) radicals; and pollutants such as sulfur dioxide (SO_2), nitrogen oxides (NO_X), and organic gases from natural and anthropogenic sources. The particle formation process includes nucleation of particles from low-vapor pressure gases emitted from sources or formed in the atmosphere by chemical reactions, condensation of low-vapor pressure gases on existing particles, and coagulation of particles. Thus, any given particle may contain PM from many sources.

The composition and behavior of particles are fundamentally linked with those of the surrounding gas. Aerosol may be defined as a suspension of solid or liquid particles in air. The term aerosol includes both the particles and all vapor or gas phase components of air. However,

the term aerosol is sometimes used to refer to the suspended particles only. In this document, "particulate" is used only as an adjective, as in particulate matter.

A complete description of the atmospheric aerosol would include an accounting of the chemical composition, morphology, and size of each particle and the relative abundance of each particle type as a function of particle size (Friedlander, 1970). However, the physical and chemical characteristics of particles are usually measured separately. Size distributions by particle number, used to calculate surface area and volume distributions, often are determined by physical means, such as electrical mobility or light scattering of suspended particles. Chemical composition usually is determined by analysis of collected samples although some species can be measured *in situ*. The mass and average chemical composition of particles, segregated according to aerodynamic diameter by cyclones or impactors, can also be determined. However, recent developments in single particle analysis techniques, by electron microscopy with X-ray analysis of single particles (but not agglomerates) collected on a substrate or by mass spectroscopy of suspended particles passing through a sensing volume, provide elemental composition of individual particles by particle size and, thus, are bringing the description envisioned by Friedlander (1970) closer to reality.

2.1.2 Physical Properties and Processes

2.1.2.1 Definitions of Particle Diameter

The diameter of a spherical particle may be determined by optical or electron microscopy, by light scattering and Mie theory, by its electrical mobility, or by its aerodynamic behavior. However, atmospheric particles often are not spherical and vary in density. Therefore, their diameters are often described by an "equivalent" diameter (i.e., that of a unit density sphere that would have the same physical behavior). The aerodynamic diameter is important for particle transport, collection, and respiratory tract deposition. The aerodynamic diameter (D_a) depends on the density of the particle. It is defined as the diameter of a spherical particle with a density of 1 g/cm^3 but with a settling velocity equal to that of the particle in question. Consequently, particles with the same physical size and shape but different densities will have different aerodynamic diameters. Detailed definitions of the various sizes and their relationships are given in standard aerosol textbooks (e.g., Friedlander [1977], Reist [1984, 1993], Seinfeld and Pandis

[1998], Hinds [1999], Vincent [1989, 1995], Willeke and Baron [1993], Baron and Willeke [2001], and Fuchs [1964, 1989]).

2.1.2.2 Aerosol Size Distributions

Particle size, as indexed by one of the "equivalent" diameters, is an important parameter in determining the properties, effects, and fate of atmospheric particles. The atmospheric deposition rates of particles, and therefore, their residence times in the atmosphere, are a strong function of their aerodynamic diameters. The aerodynamic diameter also influences deposition patterns of particles within the lung. Because light scattering is strongly dependent on the optical particle size, the amount of light scattering per unit PM mass will be dependent on the size distribution of atmospheric particles. Therefore, the effects of atmospheric particles on visibility, radiative balance, and climate will be influenced by the size distribution of the particles. Studies using impactors or cyclones measure the particle-size distribution directly in aerodynamic diameter. The diameters of atmospheric particles range from 1 nm to 100 μm, spanning 5 orders of magnitude. A variety of different instruments, measuring a variety of equivalent diameters, are required to cover this range.

Older particle counting studies used optical particle counters to cover the range of 0.3 to 30 μm diameter. Diameters of particles below 0.5 μm were measured as mobility diameters. The particle diameters used in size distribution graphs from these studies usually are given as physical diameters rather than aerodynamic diameters. In recent years, aerodynamic particle sizers have been developed that give a direct measurement of the aerodynamic diameter in the range of approximately 0.7 to 10 μm diameter. These instruments have been used with electrical mobility analyzers that measure the mobility diameter of particles from 3-5 nm to approximately 0.5 μm (McMurry, 2000). Unfortunately, there is no agreed-upon technique for combining the various equivalent diameters. Some workers use various assumptions to combine the various measurements into one presentation; others report each instrument separately. Therefore, the user of size distribution data should be careful to determine exactly which equivalent diameter is reported. Aerodynamic diameter is the most widely used equivalent diameter. In this document D_p will be used for physical diameter and D_a for aerodynamic diameter.

Particle Size Distribution Functions

The distribution of particles with respect to size is an important physical parameter governing their behavior. Because atmospheric particles cover several orders of magnitude in particle size, size distributions often are expressed in terms of the logarithm of the particle diameter on the X-axis and the measured differential concentration on the Y-axis: $\Delta N/\Delta(\log D_p)$ = the number of particles per cm^3 of air having diameters in the size range from $\log D_p$ to $\log(D_p + \Delta D_p)$. Because logarithms do not have dimensions, it is necessary to think of the distribution as a function of $\log(D_p/D_{p0})$, where the reference diameter $D_{p0} = 1\ \mu m$ is not explicitly stated. If $\Delta N/\Delta(\log D_p)$ is plotted on a linear scale, the number of particles between D_p and $D_p + \Delta D_p$ is proportional to the area under the curve of $\Delta N/\Delta(\log D_p)$ versus $\log D_p$. Similar considerations apply to distributions of surface, volume, and mass. It has been found that atmospheric aerosol size distributions frequently may be approximated by a sum of log-normal distributions corresponding to the various modes or fractions. When approximated by a function, the distributions are usually given as $dN/d(\log D_p)$ rather than $\Delta N/\Delta(\log D_p)$.

Atmospheric Aerosol Size Distributions

Averaged atmospheric size distributions are shown in Figures 2-1 through 2-3 (Whitby, 1978; Whitby and Sverdrup, 1980). Figure 2-1 describes the number of particles as a function of particle diameter for rural, urban-influenced rural, urban, and freeway-influenced urban aerosols. For some of the same data, the particle volume distributions are shown in Figure 2-2. Figure 2-3 shows the number, surface, and volume distribution for the grand average continental size distribution. Volume, surface area, and sometimes number are shown on an arithmetic scale with the distributions plotted such that the volume, surface area, or number of particles in any specified size range is proportional to the corresponding area under the curve. These distributions show that most of the particles are quite small, below 0.1 μm; whereas most of the particle volume (and therefore most of the mass) is found in particles >0.1 μm.

An important feature of the mass or volume size distributions of atmospheric aerosols is their multimodal nature. Volume distributions, measured in ambient air in the United States, are almost always found to be bimodal with a minimum between 1 and 3 μm. The distribution of particles that are mostly larger than the minimum is termed “coarse.” The distribution of particles that are mostly smaller than the minimum is termed “fine.” Whitby and Sverdrup



Figure 2-1. Number of particles as a function of particle diameter: (a) number concentrations are shown on a logarithmic scale to display the wide range by site and size; (b) number concentrations for the average urban distribution are shown on a linear scale for which the area under any part of the curve is proportional to particle number in that size range.

Source: Whitby and Sverdrup (1980).

(1980), Whitby (1978), and Willeke and Whitby (1975) identified three modes: (1) nuclei, (2) accumulation, and (3) coarse. The three modes are most apparent in the freeway-influenced size distribution of Figure 2-2b, in the surface area distribution of Figure 2-3b, and in the in-traffic volume distribution of Figure 2-4. However, the nuclei mode, corresponding to particles below about 0.1 μm, may not be noticeable in volume or mass distributions. The middle mode, from 0.1 to 1 or 2 μm, is the accumulation mode. Fine particles include both the accumulation and the nuclei modes. The third mode, containing particles larger than 1 or 2 μm, is known as the coarse particle mode. The number concentrations of coarse particles are usually too small to be seen in arithmetic plots (Figures 2-1b and 2-3a) but can be seen in a logarithmic



Figure 2-2. Particle volume distribution as a function of particle diameter: (a) for the averaged rural and urban-influenced rural number distributions shown in Figure 2-1 and a distribution from south central New Mexico, and (b) for the averaged urban and freeway-influenced urban number distributions shown in Figure 2-1.

Source: Whitby and Sverdrup (1980) and Kim et al. (1993).

plot (Figure 2-1a). Whitby and Sverdrup (1980) observed that rural aerosols, not influenced by nearby sources, have a small accumulation mode and no observable nuclei mode. For urban aerosols, the accumulation and coarse particle modes are comparable in volume. The nuclei mode is small in volume, but it dominates the number distributions of urban aerosols. Whitby's conclusions were based on extensive studies of size distributions in a number of western and midwestern locations during the 1970s (Whitby, 1978; Whitby and Sverdrup, 1980).

No size-distribution studies of similar scope have been published since then. Newer results from particle counting and impactor techniques, including data from Europe (U.S. Environmental Protection Agency, 1996a) and Australia (Keywood et al., 1999, 2000), show similar results.



Figure 2-3. Distribution of coarse (c), accumulation (a), and nuclei- or ultrafine (n) -mode particles by three characteristics, a) number (N), b) surface area (S) and c) volume (V) for the grand average continental size distribution. DGV = geometric mean diameter by volume; DGS = geometric mean diameter by surface area; DGN = geometric mean diameter by number; D_p = particle diameter.

Source: Whitby (1978).



Figure 2-4. Volume size distribution, measured in traffic, showing fine-mode and coarse-mode particles and the nuclei and accumulation modes within the fine-particle mode. DGV (geometric mean diameter by volume, equivalent to volume median diameter) and σ_g (geometric standard deviation) are shown for each mode. Also shown are transformation and growth mechanisms (e.g., nucleation, condensation, and coagulation).

Source: Adapted from Wilson and Suh (1997).

Definitions of Particle Size Fractions

In the preceding discussion several subdivisions of the aerosol size distribution were identified. Aerosol scientists use four different approaches or conventions in the classification of particles by size: (1) modes, based on the observed size distributions and formation mechanisms; (2) cut point, usually based on the 50% cut point of the specific sampling device; (3) dosimetry or occupational health sizes, based on the entrance into various compartments of the respiratory system; and (4) legally specified, regulatory sizes for air quality standards.

Modal. The modal classification, first proposed by Whitby (1978), is shown in Figure 2-3. The nuclei mode can be seen clearly in the volume distribution only in traffic or near traffic or other sources of nuclei mode particles (Figure 2-4). The observed modal structure is frequently approximated by several log-normal distributions. Definitions of terms used to describe size distributions in modal terms are given below.

Coarse Mode: The distribution of particles with diameters mostly greater than the minimum in the particle mass or volume distributions, which generally occurs between 1 and 3 μm. These particles are usually mechanically generated (e.g., from wind erosion of crustal material).

Fine Mode: The distribution of particles with diameters mostly smaller than the minimum in the particle mass or volume distributions, which generally occurs between 1 and 3 μm. These particles are generated in combustion or formed from gases. The fine mode includes the accumulation mode and the nuclei mode.

Nuclei Mode: That portion of the fine particle mode with diameters below about 0.1 μm. The nuclei mode can be observed as a separate mode in mass or volume distributions only in clean or remote areas or near sources of new particle formation by nucleation. Toxicologists and epidemiologists use the term “ultrafine” to refer to particles in the nuclei-mode size range. Aerosol physicists and material scientists tend to use the term “nanoparticles” to refer to particles in this size range generated in the laboratory.

Accumulation Mode: That portion of the fine particle mode with diameters above about 0.1 μm. Accumulation-mode particles normally do not grow into the coarse mode. Nuclei-mode particles grow by coagulation (two particles combining to form one) or by condensation (low-equilibrium vapor pressure gas molecules condensing on a particle) and “accumulate” in this size range.

Over the years, the terms fine and coarse, as applied to particle sizes, have lost the precise meaning given in Whitby’s (1978) definition. In any given article, therefore, the meaning of fine

and coarse, unless defined, must be inferred from the author's usage. In particular, $PM_{2.5}$ and fine-mode particles are not equivalent. In this document, the term "mode" is used with fine and coarse when it is desired to specify the distribution of fine-mode particles or coarse-mode particles as shown in Figures 2-4 and 2-5.



Figure 2-5. An idealized distribution of ambient particulate matter showing fine-mode particles and coarse-mode particles and the fractions collected by size-selective samplers. (WRAC is the Wide Range Aerosol Classifier which collects the entire coarse mode [Lundgren and Burton, 1995].)

Source: Adapted from Wilson and Suh (1997).

Sampler Cut Point. Another set of definitions of particle size fractions arises from considerations of size-selective sampling. Size-selective sampling refers to the collection of particles below or within a specified aerodynamic size range, usually defined by the upper 50% cut point size, and has arisen in an effort to measure particle size fractions with some special significance (e.g., health, visibility, source apportionment, etc.). Dichotomous samplers split the particles into smaller and larger fractions that may be collected on separate filters. However, some fine particles ($\approx$10%) are collected with the coarse particle fraction. Cascade impactors use multiple size cuts to obtain a distribution of size cuts for mass or chemical composition measurements. One-filter samplers with a variety of upper size cuts also have been used.

Occupational Health or Dosimetric Size Cuts. The occupational health community has defined size fractions for use in the protection of human health. This convention classifies particles into inhalable, thoracic, and respirable particles according to their upper size cuts. However, these size fractions may also be characterized in terms of their entrance into various compartments of the respiratory system. Thus, inhalable particles enter the respiratory tract, including the head airways. Thoracic particles travel past the larynx and reach the lung airways and the gas-exchange regions of the lung. Respirable particles are a subset of thoracic particles that are more likely to reach the gas-exchange region of the lung. In the past exact definitions of these terms have varied among organizations. As of 1993, a unified set of definitions was adopted by the American Conference of Governmental Industrial Hygienists (ACGIH, 1994), the International Standards Organization (ISO), and the European Standardization Committee (CEN). The curves which define inhalable (IPM), thoracic (TPM), and respirable (RPM) particulate matter are shown in Figure 2-6.

Regulatory Size Cuts. In 1987, the NAAQS for PM were revised to use PM_{10}, rather than total suspended particulate matter (TSP), as the indicator for the NAAQS for PM (Federal Register, 1987). The use of PM_{10} as an indicator is an example of size-selective sampling based on a regulatory size cut (Federal Register, 1987). The selection of PM_{10} as an indicator was based on health considerations and was intended to focus regulatory concern on those particles small enough to enter the thoracic region of the human respiratory tract. The $PM_{2.5}$ standard set in 1997 is also an example of size-selective sampling based on a regulatory size cut (Federal



Figure 2-6. Specified particle penetration (size-cut curves) through an ideal (no-particle-loss) inlet for five different size-selective sampling criteria. Regulatory size cuts are defined in the Code of Federal Regulations; $PM_{2.5}$ (2001c), PM_{10} (2001a). $PM_{2.5}$ is also defined in the Federal Register (1997). Size-cut curves for inhalable particulate matter (IPM), thoracic particulate matter (TPM) and respirable particulate matter (RPM) size cuts are computed from definitions given by American Conference of Governmental and Industrial Hygienists (1994).

Register, 1997). The $PM_{2.5}$ standard was based primarily on epidemiological studies using concentrations measured with $PM_{2.5}$ samplers as an exposure index. However, the $PM_{2.5}$ sampler was not designed to collect respirable particles. It was designed to collect fine-mode particles because of their different sources (Whitby et al., 1974). Thus, the need to attain a $PM_{2.5}$ standard will tend to focus regulatory concern on control of sources of fine-mode particles.

Prior to 1987, the indicator for the NAAQS for PM was TSP. TSP is defined by the design of the High Volume Sampler (hivol) that collects all of the fine particles but only part of the

coarse particles (Figure 2-5). The upper cut-off size of the hivol depends on the wind speed and direction and may vary from 25 to 40 μm. The Wide Range Aerosol Classifier (WRAC) was designed specifically to collect the entire coarse mode (Lundgren and Burton, 1995).

An idealized distribution, showing the normally observed division of ambient aerosols into fine-mode particles and coarse-mode particles and the size fractions collected by the WRAC, TSP, PM_{10}, $PM_{2.5}$ and $PM_{10-2.5}$ samplers, is shown in Figure 2-5. PM_{10} samplers, as defined in Appendix J to Title 40 Code of Federal Regulations (40 CFR) Part 50 (Code of Federal Regulations, 2001a; Federal Register, 1987), collect all of the fine particles and part of the coarse particles. The upper cut point is defined as having a 50% collection efficiency at 10 ± 0.5 μm aerodynamic diameter. The slope of the collection efficiency curve is defined in amendments to 40 CFR, Part 53, (Code of Federal Regulations, 2001b). An example of a PM_{10} size-cut curve is shown in Figure 2-6.

An example of a $PM_{2.5}$ size-cut curve is also shown in Figure 2-6. The $PM_{2.5}$ size-cut curve, however, is defined by the design of the Federal Reference Method (FRM) Sampler. The basic design of the FRM is given in the Federal Register (1997, 1998) and as 40 CFR Part 50, Appendix L (Code of Federal Regulations, 2001c). Additional performance specifications are given in 40 CFR Parts 53 and 58 (Code of Federal Regulations, 2001b,d). Each actual $PM_{2.5}$ reference method, as represented by a specific sampler design and associated manual operational procedures, must be designated as a reference method under 40 CFR Part 53 in Section 1.2 of Appendix L (Code of Federal Regulations, 2001c). Thus there may be many somewhat different $PM_{2.5}$ FRMs (see Table 2-4).

Papers discussing PM_{10} or $PM_{2.5}$ frequently insert an explanation such as "PM_x (particles less than x μm diameter)" or "PM_x (nominally, particles with aerodynamic diameter $\leq$x μm)." While these explanations may seen easier than the more nearly correct than "PM_X, (particles collected with an upper 50% cut point of x μm aerodynamic diameter)," they are not entirely correct and may be misleading since they suggest an upper 100% cut point of x μm. Some countries use PM_{10} to refer not to samplers with a 50% cut at 10 μm D_a but samplers with 100% rejection of all particles greater than 10 μm D_a. Such samplers miss a fraction of coarse thoracic PM. A example is shown in Figure 2-7. PM_x, as defined by EPA, refers to a sampler with a penetration curve that collects 50% of x μm particles and excludes 50% of x μm particles. It also means that some particles > x are collected and not all particles < x are collected.



Figure 2-7. Comparison of penetration curves for two PM_{10} beta gauge samplers using cyclone inlets. The Wedding PM_{10} sampler uses the U.S. EPA definition of PM_x as x = 50% cut point. The Kimoto PM_{10} defines PM_x as x = the 100% cut point (or zero penetration).

Source: Tsai and Cheng (1996).

In an analysis reported in 1979, EPA scientists endorsed the need to measure fine and coarse particles separately (Miller et al., 1979). Based on the availability of a dichotomous sampler with a separation size of 2.5 μm D_a, they recommended 2.5 μm D_a as the cut point between fine and coarse particles. Because of the wide use of this cut point, the $PM_{2.5}$ fraction is frequently referred to as "fine" particles. However, although the $PM_{2.5}$ sample contains all of the fine particles, it may collect a small fraction of the coarse particles especially in dry areas or during dry conditions. A $PM_{10-2.5}$ size fraction may be obtained from a dichotomous sampler or by subtracting the mass collected by a $PM_{2.5}$ sampler from the mass collected by a PM_{10} sampler.

The resulting $PM_{10-2.5}$ mass, or $PM_{10-2.5}$, is sometimes called "coarse" particles. However, it would be more correct to call $PM_{2.5}$ an indicator of fine-mode particles (because it contains some coarse-mode particles) and $PM_{10-2.5}$ an indicator of the thoracic component of coarse-mode particles (because it excludes some coarse-mode particles below 2.5 μm D_a and above 10 μm D_a). It would be appropriate to call PM_{10} an indicator of thoracic particles. PM_{10} and thoracic PM, as shown in Figure 2-6, have the same 50% cut point. However, the thoracic cut is not as sharp as the PM_{10} cut; therefore, thoracic PM contains some particles between 10 and 30 μm diameter that are excluded from PM_{10}.

2.1.2.3 Nuclei-Mode Particles

As discussed in Chapter 7, Toxicology of Particulate Matter, and in Chapter 8, Epidemiology of Human Health Effects from Ambient Particulate Matter, some scientists argue that ultrafine (nuclei-mode) particles may pose potential health problems and that some health effects may be more closely associated with particle number or particle surface area than particle mass. Because nuclei-mode particles contribute the major portion of particle number and a significant portion of particle surface area, some additional attention will be given to nuclei-mode particles.

Formation and Growth of Fine Particles

Several processes influence the formation and growth of particles. New particles may be formed by nucleation from gas phase material. Particles may grow by condensation as gas phase material condenses on existing particles. Particles also may grow by coagulation as two particles combine to form one. Gas phase material condenses preferentially on smaller particles, and the rate constant for coagulation of two particles decreases as the particle size increases. Therefore, nuclei mode particles grow into the accumulation mode, but accumulation mode particles do not normally grow into the coarse mode (see Figure 2-4). More information and references on formation and growth of fine particles may be found in the 1996 AQCD PM (U.S. Environmental Protection Agency, 1996a).

Equilibrium Vapor Pressures

An important parameter in particle nucleation and in particle growth by condensation is the saturation ratio S, defined as the ratio of the partial pressure of a species, p, to its equilibrium vapor pressure above a flat surface, p_o: $S = p/p_o$. For either condensation or nucleation to occur, the species vapor pressure must exceed its equilibrium vapor pressure. For particles, the equilibrium vapor pressure is not the same as p_o. Two effects are important: (1) the Kelvin effect, which is an increase in the equilibrium vapor pressure above the surface due to its curvature (very small particles have higher vapor pressures and will not be stable to evaporation until they attain a critical size) and (2) the solute effect, which is a decrease in the equilibrium vapor pressure of the liquid due to the presence of other compounds in solution. Organic compounds may also be adsorbed on ultrafine carbonaceous particles.

For an aqueous solution of a nonvolatile salt, the presence of the salt decreases the equilibrium vapor pressure of the water over the droplet. This effect is in the opposite direction of the Kelvin effect, which increases the equilibrium vapor pressure above a droplet because of its curvature. The existence of an aqueous solution will also influence the vapor pressure of water-soluble species. The vapor pressure behavior of mixtures of several liquids or of liquids containing several solutes is complex.

New Particle Formation

When the vapor concentration of a species exceeds its equilibrium concentration (expressed as its equilibrium vapor pressure), it is considered condensable. Condensable species can either condense on the surface of existing particles or can form new particles. The relative importance of nucleation versus condensation depends on the rate of formation of the condensable species and on the surface or cross-sectional area of existing particles (McMurry and Friedlander, 1979). In ambient urban environments, the available particle surface area is sufficient to rapidly scavenge the newly formed condensable species. Formation of new particles (nuclei mode) is usually not important except near sources of condensable species. Wilson et al. (1977) report observations of the nuclei mode in traffic. New particle formation also can be observed in cleaner, remote regions. Bursts of new particle formation in the atmosphere under clean conditions usually occur when aerosol surface area concentrations are low (Covert et al., 1992). High concentrations of nuclei mode particles have been observed in regions with low particle

mass concentrations indicating that new particle formation is inversely related to the available aerosol surface area (Clarke, 1992).

Sources of Nuclei-Mode Particles

Nuclei mode particles are the result of nucleation of gas phase species to form condensed phase species with very low equilibrium vapor pressure. In the atmosphere there are four major classes of sources that yield particulate matter with equilibrium vapor pressures low enough to form nuclei mode particles:

(1) *Particles containing heavy metals*. Nuclei mode particles of metal oxides or other metal compounds are generated when metallic impurities in coal or oil are vaporized during combustion and the vapor undergoes nucleation. Metallic ultrafine particles also may be formed from metals in lubricating oil or fuel additives that are vaporized during combustion of gasoline or diesel fuels. Nuclei-mode metallic particles were discussed in Section 6.9 of the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a).

(2) *Elemental carbon or soot* (EC). EC particles are formed primarily by condensation of C_2 molecules generated during the combustion process. Because EC has a very low equilibrium vapor pressure, ultrafine EC particles can nucleate even at high temperatures (Kittelson, 1998; Morawska et al., 1998).

(3) *Sulfates and nitrates*. Sulfuric acid (H_2SO_4), or its neutralization products with ammonia (NH_3), ammonium sulfate ($(NH_4)_2SO_4$) or ammonium acid sulfate (NH_4HSO_4), are generated in the atmosphere by conversion of sulfur dioxide (SO_2) to H_2SO_4. As H_2SO_4 is formed, it can either nucleate to form new ultrafine particles, or it can condense on existing nuclei mode or accumulation mode particles (Clark and Whitby, 1975; Whitby, 1978). The possible formation of ultrafine NH_4NO_3 by reaction of NH_3 and nitric acid (HNO_3) vapor apparently has not been investigated.

(4) *Organic carbon*. Recent smog chamber studies and indoor experiments show that atmospheric oxidation of certain organic compounds found in the atmosphere can produce highly oxidized organic compounds with an equilibrium vapor pressure sufficiently low to result in nucleation (Kamens et al., 1999; Weschler and Shields, 1999).

Recent Measurements of Nuclei-Mode Particles

Instruments, developed during the past decade, permit measurement of size distributions down to 3 nm diameter particles. Use of these techniques have led to new information on the formation of new particles by nucleation. Such measurements have been carried out during intensive field measurement campaigns and also during continuous measurements in urban areas in several European cities and in the U.S. as a part of the Supersite program (McMurry et. al., 2000; Woo et al., 2001a). Nucleation has been observed in the outflows of convective clouds, downwind of coastal regions during low tide, over forests, downwind of certain biogenic emissions, and in urban areas. Nucleation events in outdoor air almost always occur during daylight, indicating that photochemistry plays a role in producing the gas phase precursors of new particles. There is strong evidence that sulfuric acid vapor often participates in nucleation. However, condensation of sulfuric acid and its associated water and ammonia typically can account for only 10% to 20% of the observed growth rates for freshly nucleated particles. Therefore, organic compounds may account for much of the formation and growth of freshly nucleated particles. Evidence of nucleation of organic particles comes largely from smog chamber studies (Kamens et al., 1999). Nucleation of organic particles may also occur indoors due to the reaction of infiltrated ozone with indoor terpenes from air fresheners or cleaning solutions (Weschler and Shields, 1999). The observation of bursts of nuclei-mode particles in Atlanta (Woo et al., 2001a), perhaps due to unusually high rates of production of condensible species, suggests that exposure to high concentrations of ultrafine or nuclei-mode particles may be a more frequent occurrence that previously expected.

Concentration of Nuclei-Mode Particles: A Balance Between Formation and Removal

Nuclei-mode particles may be removed by dry deposition or by growth into the accumulation mode. This growth takes place as other low vapor pressure material condenses on the particles or as nuclei-mode particles coagulate with themselves or with accumulation mode particles. Because the rate of coagulation would vary with the concentration of accumulation-mode particles, it might be expected that the concentration of nuclei-mode particles would increase with a decrease in accumulation-mode mass. On the other hand, the concentration of particles would be expected to decrease with a decrease in the rate of generation of particles by reduction in emissions of metal and carbon particles or a decrease in the rate of generation of

H_2SO_4 or condensable organic vapor. The rate of generation of H_2SO_4 depends on the concentration of SO_2 and hydroxyl radical (•OH), which is generated primarily by reactions involving ozone (O_3). Thus, reductions in SO_2 and O_3 would lead to a decrease in the rate of generation of H_2SO_4 and condensable organic vapor and to a decrease in the concentration of nuclei-mode particles. The balance between formation and removal is uncertain. However, these processes can be modeled using a general dynamic equation for particle size distribution (Friedlander, 1977) or by aerosol dynamics modules in newer air quality models (Binkowski and Shanker, 1995; Binkowski and Ching, 1995).

2.1.3 Chemistry of Atmospheric Particulate Matter

The major constituents of atmospheric PM are sulfate, nitrate, ammonium, and hydrogen ions; particle-bound water; elemental carbon; a great variety of organic compounds; and crustal material. Atmospheric PM also contains a large number of elements in various compounds and concentrations. More information and references on the composition of PM, measured in a large number of studies in the United States, may be found in 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). The composition and concentrations of PM are discussed in Chapter 3, Section 3.1, Patterns and Trends in Ambient $PM_{2.5}$ Concentrations. Ambient data for concentrations and composition of $PM_{2.5}$ are given in Appendices 3A, 3B, and 3C.

2.1.3.1 Chemical Composition and Its Dependence on Particle Size

Studies conducted in most parts of the United States indicate that sulfate, ammonium, and hydrogen ions; elemental carbon, secondary organic compounds and primary organic species from cooking and combustion; and certain transition metals are found predominantly in the fine particle mode. Crustal materials such as calcium, aluminum, silicon, magnesium, and iron are found predominately in the coarse particles. Some organic materials such as pollen, spores, and plant and animal debris are also found predominantly in the coarse mode. Some components such as potassium and nitrate may be found in both the fine and coarse particle modes but from different sources or mechanisms. Potassium in coarse particles comes from soil. Potassium also is found in fine particles in emissions from burning wood or cooking meat. Nitrate in fine particles comes primarily from the reaction of gas-phase nitric acid with gas-phase ammonia to

form particulate ammonium nitrate. Nitrate in coarse particles comes primarily from the reaction of gas-phase nitric acid with preexisting coarse particles.

2.1.3.2 Primary and Secondary Particulate Matter

Particulate material can be primary or secondary. PM is called "primary" if it is in the same chemical form in which it was emitted into the atmosphere. PM is called "secondary" if it is formed by chemical reactions in the atmosphere. Primary coarse particles are usually formed by mechanical processes. This includes material emitted in particulate form such as wind-blown dust, sea salt, road dust, and combustion-generated particles such as fly ash and soot. Primary fine particles are emitted from sources either directly as particles or as vapors that rapidly condense to form ultrafine or nuclei-mode particles. This includes soot from diesel engines, a great variety of organic compounds condensed from incomplete combustion or cooking, and compounds of As, Se, Zn, etc., that condense from vapor formed during combustion or smelting. The concentration of primary particles depends on their emission rate, transport and dispersion, and removal rate from the atmosphere.

Secondary PM is formed by chemical reactions of free, adsorbed, or dissolved gases. Most secondary fine PM is formed from condensable vapors generated by chemical reactions of gas-phase precursors. Secondary formation processes can result in either the formation of new particles or the addition of particulate material to pre-existing particles. Most of the sulfate and nitrate and a portion of the organic compounds in atmospheric particles are formed by chemical reactions in the atmosphere. Secondary aerosol formation depends on numerous factors including the concentrations of precursors; the concentrations of other gaseous reactive species such as ozone, hydroxyl radical, peroxy radicals, or hydrogen peroxide; atmospheric conditions including solar radiation and relative humidity (RH); and the interactions of precursors and pre-existing particles within cloud or fog droplets or in the liquid film on solid particles. As a result, it is considerably more difficult to relate ambient concentrations of secondary species to sources of precursor emissions than it is to identify the sources of primary particles.
A significant effort is currently being directed toward the identification and modeling of organic products of photochemical smog including the conversion of gases to particulate matter. More information of the transformation of precursor gases into secondary PM is given in Chapter 3, Section 3.3.1, Chemistry of Secondary PM Formation.

2.1.3.3 Particle-Vapor Partitioning

Several atmospheric aerosol species, such as ammonium nitrate and certain organic compounds, are semivolatile and are found in both gas and particle phases. A variety of thermodynamic models have been developed to predict the temperature and relative humidity dependence of the ammonium nitrate equilibria with gaseous nitric acid and ammonia. However, under some atmospheric conditions, such as cool, cold, or very clean air, the relative concentrations of the gas and solid phases are not accurately predicted by equilibrium considerations alone, and transport kinetics can be important. The gas-particle distribution of semivolatile organic compounds depends on the equilibrium vapor pressure of the compound, total particle surface area, particle composition, atmospheric temperature, and relative humidity. Although it generally is assumed that the gas-particle partitioning of semivolatile organics is in equilibrium in the atmosphere, neither the equilibria nor the kinetics of redistribution are well understood. Diurnal temperature fluctuations cause gas-particle partitioning to be dynamic on a time scale of a few hours and can cause semivolatile compounds to evaporate during the sampling process. The pressure drop across the filter can also contribute to the loss of semivolatile compounds. The dynamic changes in gas-particle partitioning caused by changes in temperature, pressure, and gas-phase concentration, both in the atmosphere and after collection, cause serious sampling problems that are discussed in Section 2.2.3, Measurement of Semivolatile Particulate Matter.

Equilibria with Water Vapor

As a result of the equilibrium of water vapor with liquid water in hygroscopic particles, many ambient particles contain liquid water (particle-bound water). Unless removed, this particle-bound water will be measured as a component of the particle mass. Particle-bound water is important in that it influences the size of the particles, and in turn, their light scattering properties and their aerodynamic properties, which are important for deposition to surfaces, to airways following inhalation, and in sampling instrumentation. The aqueous solution provides a medium for reactions of dissolved gases including reactions that do not take place in the gas phase. The aqueous solutions also may act as a carrier to convey soluble toxic species to the gas-exchange regions of the respiratory system, including species that would be removed by deposition in the upper airways if the particles had remained in the gas phase (Friedlander and

Yeh, 1998; Kao and Friedlander, 1995; Wilson, 1995). An extensive review of equilibrium with water vapor as it pertains to ambient aerosols was given in Chapter 3 of the 1996 PM AQCD (U.S. Environmental Protection, Agency, 1996a).

The interaction of particles with water vapor may be described briefly as follows. As relative humidity increases, particles of crystalline soluble salts, such as $(NH_4)_2SO_4$, NH_4HSO_4, or NH_4NO_3, undergo a phase transition to become aqueous solution particles. According to the phase rule, for particles consisting of a single component, this phase transition is abrupt, taking place at a relative humidity that corresponds to the vapor pressure of water above the saturated solution (the deliquescence point). With a further increase in relative humidity, the solution particle adds water (and the concentration of the solute decreases) so that the vapor pressure of the solution is maintained equal to that of the surrounding relative humidity; thus, the solution particle tends to follow the equilibrium growth curve. As relative humidity decreases, the solution particle follows the equilibrium curve to the deliquescence point. However, rather than crystallizing at the deliquescence relative humidity, the solute remains dissolved in a supersaturated solution to considerably lower relative humidities. Ultimately the solution particle abruptly loses its water vapor (efflorescence) and typically returns to the initial crystalline form.

For particles consisting of more than one component, the solid to liquid transition will take place over a range of relative humidities with an abrupt onset at the lowest deliquescence point of the several components and with subsequent growth as crystalline material in the particle dissolves according to the phase diagram for the particular multicomponent system. Under such circumstances, a single particle may undergo several more or less abrupt phase transitions until the soluble material is fully dissolved. At decreasing relative humidity, such particles tend to remain in solution to relative humidities well below the several deliquescence points. In the case of the sulfuric acid-ammonium sulfate-water system, the phase diagram is fairly well worked out. Mixed anion systems containing nitrate are more labile because of the equilibrium between particulate NH_4NO_3 and gaseous NH_3 and HNO_3. For particles of composition intermediate between NH_4HSO_4 and $(NH_4)_2SO_4$, this transition occurs in the range from 40% to below 10%, indicating that for certain compositions the solution cannot be dried in the atmosphere. At low relative humidities, particles of this composition would likely be present in the atmosphere as

supersaturated solution droplets (liquid particles) rather than as solid particles. Thus, they would exhibit hygroscopic rather than deliquescent behavior during relative humidity cycles.

Other pure compounds, such as sulfuric acid, are hygroscopic (i.e., they form aqueous solutions at any relative humidity and maintain a solution vapor pressure over the entire range of relative humidity). Soluble organic compounds may also contribute to the hygroscopicity of the atmospheric aerosol (Saxena et al., 1995; Saxena and Hildeman, 1996), but the equilibria involving organic compounds and water vapor, and, especially for mixtures of salts, organic compounds, and water, are not so well understood. These equilibrium processes may cause an ambient particle to significantly increase its diameter at relative humidities above about 40% (Figure 2-8). A particle can grow to five times its dry diameter as the RH approaches 100% (Figure 2-9). The Federal Reference Methods, for filter measurements of $PM_{2.5}$ and PM_{10} mass, require equilibration at a specified, low relative humidity (≈40% RH) after collection. This equilibration removes much of the particle-bound water and provides a stable PM mass (see Section 2.2 for details and references). Otherwise, particle mass would be a function of relative humidity, and the particle mass would be largely particle-bound water at higher relative humidities.

Continuous monitoring techniques generally attempt to remove particle-bound water before measurement, either by heating or dehumidification. Semivolatile material may be lost during sampling or equilibration; it is certainly lost when the collected sample is heated above ambient temperature. In addition to problems due to the loss of semivolatile species, recent studies have shown that significant amounts of particle-bound water are retained in particles collected on impaction surfaces even after equilibration and that the amount of retained particle-bound water increases with relative humidity during collection (Hitzenberger et al., 1997). Large increases in mass with increasing relative humidity were observed for the accumulation mode. The change in particle size with relative humidity also means that particle measurements such as surface area or volume, or composition as a function of size, should be made at the same RH in order for the results are to be comparable. These problems are addressed below in more detail, in Section 2.2 on Measurement of Particulate Matter.



Figure 2-8. Particle growth curves showing fully reversible hygroscopic growth of sulfuric acid (H_2SO_4) particles, deliquescent growth of ammonium sulfate [$(NH_4)_2$ SO_4] particles at the deliquescent point (A, about 80% relative humidity [RH]), reversible hygroscopic growth of ammonium sulfate solution droplets at RH greater than 80%, and hysteresis (the droplet remains supersaturated as the RH decreases below 80%) until the crystallization point (B, about 38% RH) is reached (adapted from National Research Council, 1993 and Tang, 1980).

Source: National Research Council (1993) adapted from Tang (1980).

2.1.3.4 Atmospheric Lifetimes and Removal Processes

The lifetimes of particles vary with size. Nuclei-mode particles rapidly grow into the accumulation mode. However, the accumulation mode does not grow into the coarse mode. Accumulation-mode fine particles are kept suspended by normal air motions and have very low deposition rates to surfaces. They can be transported thousands of km and remain in the atmosphere for a number of days. Coarse particles can settle rapidly from the atmosphere within



Figure 2-9. Theoretical predictions and experimental measurements of growth of NH_4HSO_4 particles at relative humidity between 95 and 100%.

Source: Li et al. (1992).

hours and normally travel only short distances. However, when mixed high into the atmosphere, as in dust storms, the smaller-sized coarse-mode particles may have longer lives and travel greater distances. Dry deposition rates are expressed in terms of a deposition velocity that varies with particle size, reaching a minimum between 0.1 and 1.0 μm aerodynamic diameter (e.g., Lin et al., 1994). Accumulation-mode particles are removed from the atmosphere primarily by cloud processes. Fine particles, especially particles with a hygroscopic component, grow as the relative humidity increases, serve as cloud condensation nuclei, and grow into cloud droplets. If the cloud droplets grow large enough to form rain, the particles are removed in the rain. Falling rain

drops impact coarse particles and remove them. Ultrafine or nuclei-mode particles are small enough to diffuse to the falling drop, be captured, and be removed in rain. Falling rain drops, however, are not nearly as effective in removing accumulation-mode particles as the cloud processes mentioned above. A more detailed discussion of particle deposition, including acid deposition, especially as it applies to deposition to vegetation, soil, and water surfaces, is given in Chapter 4, Environmental Effects of Particulate Matter. Acid deposition and PM are intimately related, first, because particles contribute to the acidification of rain and, secondly, because the gas phase species that lead to dry deposition of acidity are also precursors of particles. Therefore, reductions in SO_2 and NO_x emissions will decrease both acidic deposition and PM concentrations.

Sulfate, nitrate, and some partially oxidized organic compounds are hygroscopic and act as nuclei for the formation of cloud droplets. These droplets serve as chemical reactors in which (even slightly) soluble gases can dissolve and react. Thus, SO_2 can dissolve in cloud droplets and be oxidized to sulfuric acid by dissolved ozone or hydrogen peroxide. These reactions take place only in aqueous solution, not in the gas phase. Sulfur dioxide also may be oxidized by dissolved oxygen. This process will be faster if metal catalysts such as iron or manganese are present in solution. If the droplets evaporate, larger particles are left behind. If the droplets grow large enough, they will fall as rain; and the particles will be removed from the atmosphere with potential effects on the materials, plants, or soil on which the rain falls. (Similar considerations apply to dew.) Atmospheric particles that nucleate cloud droplets also may contain other soluble or nonsoluble materials such as metal salts and organic compounds that may add to the toxicity of the rain. Thus, the adverse effects of acid deposition on soils, plants, and trees as well as lakes, streams, and fish may be taken into account in setting secondary PM standards. Sulfuric acid, ammonium nitrate, ammonium sulfates, and organic particles also are deposited on surfaces by dry deposition. The utilization of ammonium by plants leads to the production of acidity. Therefore, dry deposition of particles can also contribute to the ecological damages caused by acid deposition. These effects are discussed in Chapter 4, Environmental Effects of Particulate Matter.

2.1.4 Summary

The physical and chemical properties of ultrafine mode, accumulation mode, and coarse mode particles are summarized in Table 2-1.

2.2 MEASUREMENT OF PARTICULATE MATTER

The 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) summarized sampling and analytical techniques for PM and acid deposition that had appeared in the literature since the earlier 1982 PM AQCD (U.S. Environmental Protection Agency, 1982). Excellent reviews have also been published by Chow (1995) and McMurry (2000). This section discusses problems in measuring PM; new techniques that attempt to alleviate these problems or measure problem species; Federal Reference Methods, speciation monitors, analytical methods for inorganic elements, organic and elemental carbon, and ionic species; and continuous and multiday monitors.

2.2.1 Particle Measurements of Interest

There are many PM components and parameters that are of interest across the various types of uses to which PM measurement data are applied. These uses include analyses of compliance with air quality standards and trends; source category apportionment studies, related to the develop of pollution reduction strategies and the validation of air quality models; studies related to health, ecological, and radiative effects; and characterization of current air quality for presentation to the public in the context of EPA's Air Quality Index. PM measurement components and parameters of specific interest for these various purposes are noted below and summarized in Table 2-2.

Particle measurements are needed to determine if a location is in compliance with air quality standards and to determine long-term trends in air quality patterns. For these purposes, precision of the measurements by a variety of measurement instruments in use is a critical consideration. Therefore, intercomparisons of various samplers, under a variety of atmospheric and air quality conditions, are important.

TABLE 2-1. COMPARISON OF AMBIENT PARTICLES, FINE MODE (Nuclei Mode Plus Accumulation Mode) AND COARSE MODE

	Fine		Coarse
	Nuclei	Accumulation	
Formed from:	Combustion, high-temperature processes, and atmospheric reactions		Break-up of large solids/droplets
Formed by:	Nucleation Condensation Coagulation	Condensation Coagulation Reactions of gases in or on particles Reactions of gases in or on particles Evaporation of fog and cloud droplets in which gases have dissolved and reacted	Mechanical disruption (crushing, grinding, abrasion of surfaces) Evaporation of sprays Suspension of dusts Reactions of gases in or on particles
Composed of:	Sulfates Elemental Carbon Metal compounds Organic compounds with very low saturation vapor pressure at ambient temperature	Sulfate, Nitrate, Ammonium, and Hydrogen ions Elemental carbon Large variety of organic compounds Metals: compounds of Pb, Cd, V, Ni, Cu, Zn, Mn, Fe, etc. Particle-bound water	Suspended soil or street dust Fly ash from uncontrolled combustion of coal, oil, and wood Nitrates/chlorides from HNO_3/HCl Oxides of crustal elements (Si, Al, It, Fe) $CaCO_3$, NaCl, sea salt Pollen, mold, fungal spores Plant and animal fragments Tire, brake pad, and road wear debris
Solubility:	Probably less soluble than accumulation mode	Largely soluble, hygroscopic, and deliquescent	Largely insoluble and nonhygroscopic
Sources:	Combustion Atmospheric transformation of SO_2 and some organic compounds High temperature processes	Combustion of coal, oil, gasoline, diesel fuel, wood Atmospheric transformation products of NO_x, SO_2, and organic compounds, including biogenic organic species (e.g., terpenes) High-temperature processes, smelters, steel mills, etc.	Resuspension of industrial dust and soil tracked onto roads and streets Suspension from disturbed soil (e.g., farming, mining, unpaved roads) Construction and demolition Uncontrolled coal and oil combustion Ocean spray Biological sources
Atmospheric half-life:	Minutes to hours	Days to weeks	Minutes to hours
Removal Processes:	Grows into accumulation mode	Forms cloud droplets and rains out Dry deposition	Dry deposition by fallout Scavenging by falling rain drops
Travel distance:	<1 to 10s of km	100s to 1000s of km	<1 to 10s of km (100s to 1000s in dust storms)

Source: Adapted from Wilson and Suh (1997).

TABLE 2-2. PARTICULATE MATTER COMPONENTS/PARAMETERS OF INTEREST FOR HEALTH, ECOLOGICAL, OR RADIATIVE EFFECTS; FOR SOURCE CATEGORY APPORTIONMENT STUDIES; OR FOR AIR QUALITY MODELING EVALUATION STUDIES

- Particle number
- Particle surface area
- Particle size distribution
- PM mass (fine PM mass [$PM_{2.5}$] and coarse thoracic PM mass [$PM_{10-2.5}$]) including both nonvolatile mass as measured by the current Federal Reference method and total mass (including semivolatile components such as ammonium nitrate and semivolatile organic compounds, but not particle-bound water)
- Ions (sulfate, nitrate, and ammonium)
- Strong acidity (H^+)
- Elemental carbon
- Organic carbon (total, nonvolatile, and semivolatile; functional groups and individual species)
- Transition metals (water soluble, bioavailable, oxidant generation)
- Specific toxic elements and organic compounds
- Crustal elements
- Bioaerosols
- Particle refractive index (real and imaginary)
- Particle density
- Particle size change with changes in relative humidity

In order to reduce pollution to attain a standard, local agencies and national research organizations need measurements to identify source categories and to develop and validate air quality models. For these purposes, PM parameters other than mass, such as chemical composition and size distribution, must also be measured. Moreover, measurements are needed with shorter time resolution in order to match changes in pollution with diurnal changes in the boundary layer.

A number of PM measurements are needed for use in epidemiological and exposure studies and to determine components of PM to guide planning and interpretation of toxicologic studies. For these purposes, size and chemical composition measurements are important, as is measurement across different time intervals. For epidemiologic studies of acute (i.e., short-term) PM exposures, 1-h or continuous measurements can provide important information beyond that provided by 24-h measurements. However, for epidemiologic studies of chronic PM exposures, measurements that integrate over longer intervals (e.g., a week to a month) are more relevant. For dosimetric studies and modeling, information will be needed on the particle size distribution and on the behavior of particles as the relative humidity and temperature are increased to those found in the respiratory system.

For studies of ecological effects and materials damage, measurements of particles and of the chemical components of particulate matter in rain, fog, and dew are needed (a) to understand the contributions of PM to soiling of surfaces and damage to materials and (b) to understand the wet and dry deposition of acidity and toxic substances to surface water, soil, and plants. Some differentiation into particles size is needed to determine dry deposition.

For studies of visibility impairment and radiative effects, information is needed that relates to how particles scatter and absorb light, including refractive index, ratio of scattering to absorption, size distribution, and change in particle size with change in relative humidity.

EPA's Air Quality Index is intended to provide the public with near real-time information of air quality in urban areas. For this purpose, PM measurements over short time intervals (e.g., 1-h) or continuous measurements are critical.

2.2.2 Issues in Measurement of Particulate Matter

The EPA decision to revise the PM standards by adding daily and yearly standards for $PM_{2.5}$ has led to a renewed interest in the measurement of atmospheric particles and also to a better understanding of the problems in developing precise and accurate measurements of particles. It is very difficult to measure and characterize particles suspended in the atmosphere; however, improvements in PM monitoring may be anticipated. EPA's PM standards are based, in part, on epidemiologic relationships between health effects and PM concentrations as measured with existing monitoring methods. As understanding of suspended PM has advanced and new monitoring information has become available, EPA has changed the indicator for the

PM NAAQS from TSP to PM_{10} to $PM_{2.5}$. During the current review consideration will be given to a standard for coarse PM.

The U.S. Federal Reference Methods (FRM) for $PM_{2.5}$ and PM_{10} provide relatively precise (±10 %) methods for determining the mass of material remaining on a Teflon filter after equilibration. However, numerous uncertainties remain as to the relationship between the mass and composition of material remaining on the filter, as measured by the FRMs, and the mass and composition of material that existed in the atmosphere as suspended PM. As a result, EPA defines accuracy for PM measurements in terms of agreement of a candidate sampler with a reference sampler. Therefore, intercomparisons of samplers become very important in determining how well various samplers agree and how various design choices influence what is actually measured.

There are six general areas where choices are made in the design and use of an aerosol sampler. These include (1) treatment of semivolatile components; (2) selection of an upper cut point; (3) separation of fine-mode and coarse-mode PM; (4) treatment of pressure, temperature, and relative humidity; (5) time resolution; and (6) assessment of the reliability of the measurement technique. In many cases, choices have been made without adequate knowledge or understanding of the consequences. As a result, measurement methods developed by different organizations may give different results when sampling the same atmosphere even though the techniques appear to be similar.

2.2.2.1 Treatment of Semivolatile Components of Particulate Matter

Current filtration-based mass measurements can experience significant evaporative losses, during and possibly after collection, of a variety of semivolatile components (i.e., species that exist in the atmosphere in dynamic equilibrium between the condensed phase and gas phase). Important examples include ammonium nitrate, semivolatile organic compounds, and particle-bound water. This problem is illustrated in Figure 2-10.

Possible approaches that have been used to address the problem of potentially lost semivolatile components include those that follow, which will be discussed in more detail in subsequent sections.

1. Collect/measure all components present in the atmosphere in the condensed phase except particle-bound water. (Examples: Brigham Young absorptive sampler and Harvard pressure



Figure 2-10. Schematic showing major nonvolatile and semivolatile components of $PM_{2.5}$. Semivolatile components are subject to partial to complete loss during equilibration or heating. The optimal technique would be to remove all particle-bound water but no ammonium nitrate or semivolatile organic PM.

drop monitor. Both require preconcentration of the accumulation mode and reduction of ambient humidity.)

2. Stabilize PM at a specified temperature high enough to remove all, or almost all, particle-bound water. This results in loss of much of the semivolatile PM. (Examples: tapered element oscillating microbalance (TEOM) operated at 50 ºC, beta gauge with heated inlet.)
3. Equilibrate collected material at fixed, near-room temperature and moderate relative humidity to reduce particle-bound water. Accept the loss of an unknown but possibly significant fraction of semivolatile PM. (Example: U.S. Federal Reference Method and most filter-weighing techniques.) Equilibration originally was designed to remove adsorbed water vapor

from glass fiber filters in order to maintain a stable filter weight. The designated RH (40%) was a compromise. If the RH is too low, electrostatic charging becomes a problem. The equilibration process does help provide a stable and reproducible mass. It also reduces the particle-bound water. However, it may not remove all particle-bound water.

The amount of semivolatile material lost is dependent on the concentration and composition of the semivolatile components and is, therefore, also dependent on season and location. The amount of semivolatile material lost has been found to be significant in air sheds with high nitrate, wood smoke, or secondary organic aerosols.

2.2.2.2 Upper Cut Point

The upper cut point of the high volume sampler varied with wind speed and direction. Newer PM samplers are usually designed to have an upper cut point and its standard deviation that are independent of wind speed and direction. Current PM samplers have upper cut points that are stable under normal operating conditions. However, problems may occur under unusual or adverse conditions. Ono et al. (2000) recently reported the results of a study in which several PM_{10} samplers were collocated and operated at various sites at Owens Lake, CA, a location with high concentrations of coarse PM. Samplers included the Partisol sampler, the TEOM, a dichotomous sampler, the Wedding high-volume sampler, and the Graseby high-volume sampler. They found that the TEOM and Partisol samplers agreed to within 6% on average. The dichotomous sampler and the Graseby and Wedding high-volume samplers, however, measured significantly lower PM_{10} concentrations than the TEOM (on average 10, 25, and 35% lower, respectively). These lower concentrations were attributed to a decrease in cut point at higher wind speeds and possibly when the inlet is dirty.

The choice of the cut point characteristics depends upon the application for the sampling device. A separation that simulates the removal of particles by the upper part of the human respiratory system might appear to be a good choice for both health risk and regulatory monitoring (i.e., measure what gets into the lungs). The ACGIH-ISO-CEN penetration curve for thoracic particles (particles able to pass the larynx and penetrate into the bronchial and alveolar regions of the lung) has a 50% cut point at 10 μm aerodynamic diameter (D_a). The U.S. PM_{10} separation curve is sharper than the thoracic penetration curve but has the advantage of reducing

the problem of maintaining the finite collection efficiency specified by the thoracic penetration curve for particles larger than 10 μm D_a. (See Section 2.1.2.2 and Figure 2-6).

2.2.2.3 Cut Point for Separation of Fine-Mode and Coarse-Mode Particulate Matter

As Table 2-1 showed, fine- and coarse-mode particles differ not only in size and morphology (e.g., smooth droplets versus rough solid particles), but also in formation mechanisms; sources; and chemical, physical, and biological properties. They also differ in concentration-exposure relationships, dosimetry (deposition in the respiratory system), toxicity, and health effects as observed by epidemiologic studies. Thus, it is desirable to measure fine-mode PM and coarse-mode PM separately in order to properly allocate health effects to either fine-mode or coarse-mode PM and to correctly determine sources by receptor modeling approaches. For example, sulfate in the fine-mode is associated with hydrogen or ammonium ions while sulfate in the coarse mode is associated with basic metal ions. Transition metals in the coarse mode are likely to be associated with soil and tend to be less soluble (and presumably less bioavailable) than transition metals in fresh combustion particles found in the fine mode.

The 2.5 μm D_a cut point was chosen in the early 1970s as the cut point for a new dichotomous sampler (Loo et al., 1976; Jaklevic et al., 1977) for use in the Regional Air Pollution Study in St. Louis, MO. At that time aerosol scientists were beginning to realize that there was a minimum between 1 and 3 μm in the distribution of particle size by volume (Whitby et al., 1974). The 2.5 μm cut point was subsequently used as an indicator of fine-mode PM in a number of studies, including the Harvard Six-City Studies of the relationships between mortality and PM concentrations (Dockery et al., 1993; Schwartz et al., 1996). A 2.5 μm cut point was also used in the Inhalable Particle Network (Suggs and Burton, 1983) which provided data for another major epidemiologic study of PM - mortality relationships using an American Cancer Society cohort (Pope et al., 1995). Therefore, at the time of the last review of the NAAQS for PM (U.S. Environmental Protection Agency, 1996a,b), there were a number of epidemiologic studies demonstrating a statistical relationship between $PM_{2.5}$ concentrations and mortality.

It is now understood that the intermodal region (1-2.5 μm) may contain either accumulation-mode or coarse-mode material and that the two modes may overlap in this region. The experimental information on the composition and source of the intermodal mass was discussed extensively in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a).

Depending on conditions, a significant amount of either accumulation- or coarse-mode material may be found in the intermodal region between 1 and 2.5 μm. The analysis demonstrated the important role of relative humidity in influencing the size of particles in both the accumulation and coarse modes.

As the relative humidity increases, hygroscopic accumulation-mode particles will increase in size due to accumulation of particle-bound water. At high relative humidities, some originally submicrometer accumulation-mode PM may be found with a D_a above 1 μm. At a relative humidity of 100%, such as found in fog and clouds, accumulation mode PM may exceed 2.5 μm D_a. What is not well understood is whether such particles will shrink to diameters below 1 μm as the RH decreases or whether reactions occurring in the wet particles will result in an increase in non-aqueous mass so that even at low RH the diameters would exceed 1 μm. On the other hand, at very low relative humidity, coarse-mode particles may be fragmented into smaller sizes, and small amounts of coarse-mode PM may be found with an D_a below 2.5 μm (Lundgren et al., 1984; Lundgren and Burton, 1995). Thus, a $PM_{2.5}$ sample will contain all of the fine-mode material except during periods of RH near 100 %. However, under conditions of low RH, it may also contain a small fraction of the coarse-mode PM. The selection of a cut point of 2.5 μm as a basis for EPA's 1997 NAAQS for fine particles (Federal Register, 1997) and its continued use in many health effects studies reflect the importance placed on more complete inclusion of fine-mode particles, while recognizing that intrusion of coarse-mode particles can occur under some conditions with this cut point.

In addition to the influence of relative humidity, in areas where winds cause high concentrations of windblown soil, there is evidence that a significant amount of coarse-mode PM may be found below 2.5 μm. An example, taken from data collected during the August 1996 dust storm in Spokane, WA, is shown in Figure 2-11. Note that the PM_{10} scale is 10 times that of the other size fractions. PM_1, although high in the morning, goes down as the wind increases and PM_{10}, $PM_{2.5}$, and $PM_{2.5-1}$ go up. During the peak of the dust storm, $PM_{2.5-1}$ was 88% of $PM_{2.5}$. For the 24-h period, $PM_{2.5-1}$ was 54% of $PM_{2.5}$. However, PM_1 was not affected by the intrusion of coarse-mode particles. Similar considerations probably apply to intrusions of dust transported from distance sources such as the Sahara and Gobi deserts (Husar et al., 2001).

A cut point of 1 μm could reduce the misclassification of coarse-mode material as fine, especially in a areas with high levels of wind blown soil, but under high RH conditions could



Figure 2-11. Particulate matter concentrations in Spokane, WA, during the August 30, 1996 dust storm.

Source: Claiborn et al. (2000).

result in some fine-mode material being misclassified as coarse. A reduction in RH, either intentionally or inadvertently, will reduce the size of the fine mode. A sufficient reduction in RH should yield a dry fine-particle mode with very little material above 1.0 μm. Studies of the changes in particle size with changes in relative humidity suggest that only a small fraction of accumulation mode particles will be above 1 μm in diameter at RH below 60%, but a substantial fraction will grow above 1 μm for RH above 80% (Hitzenberger et al., 1997; McMurry and Stolzenburg, 1989; U.S. Environmental Protection Agency, 1996a).

Under high relative humidity circumstances, a monitor using a 1.0 μm D_a cut point can achieve better modal separation if the air stream is dehumidified to some fixed humidity that would remove all or most particle-bound water without evaporating semivolatile components. New techniques which require reduction of RH prior to collection have been developed for measurement of fine particulate matter minus particle-bound water but including semivolatile nitrate and organic compounds. With such techniques, PM_1 measurements, in conjunction with concurrent $PM_{2.5}$ measurements, would be useful for exposure, epidemiologic, and source

apportionment studies, especially in areas where intrusion of coarse-mode particles into the intermodal range is likely.

2.2.2.4 Treatment of Pressure, Temperature, and Relative Humidity

There are a variety of techniques for defining (or ignoring) the pressure, temperature, and relative humidity during and after sampling. For example, the sample volume may be based on the mass or volumetric flow corrected to standard temperature and pressure (273 °K and 1 atm.) (current EPA technique for PM_{10}), or it may be on the volumetric flow at ambient conditions of temperature and pressure (current EPA technique for $PM_{2.5}$).

There are also a variety of options for the control of temperature during collection. The particles may be heated enough to remove much of the particle-bound water (i.e., TEOM at 50 °C); the particles may be heated several degrees, just enough to prevent condensation of water in sampling system; the particles and the sampler may be maintained near ambient temperature (±5 °C of ambient temperature is required for EPA FRMs); or the particles and sampler may be maintained at constant temperature inside a heated or air conditioned shelter. There are also options for control of temperature after collection: (a) no control (room temperature) or (b) ship and store at cool temperature (4 °C is the current EPA FRM requirement).

Consideration must also be given to relative humidity. Changes in relative humidity cause changes in particle size of hygroscopic or deliquescent particles. Changing relative humidity by adding or removing water vapor affects measurements of particle number, particle surface area, and particle size distribution and the amount of overlap of fine-mode and coarse-mode particles. Changing relative humidity by intentional or inadvertent changes in temperature also affects the amount of loss of ammonium nitrate and semivolatile organic compounds. Monitoring personnel should be aware of the various options for treatment of pressure, temperature, and relative humidity; make appropriate selections; and document which options are used.

Studies of relationships between personal/indoor/outdoor measurements present special problems. Indoor environments are typically dryer than outdoors and may be warmer or, if air-conditioned, cooler. These differences may change particle size and the amount of volatilization of semivolatile components. Such changes between indoors and outdoors will complicate the comparison of indoor to outdoor PM concentrations; the modeling of personal

exposure to all particles; and apportionment of personal exposure into particles of ambient origin, particles of indoor origin, and personal activity particles.

2.2.2.5 Time Resolution

The classic 24-hour filter collection technique is being supplemented by a variety of continuous monitors for various PM parameters. This process is being accelerated by the lower operational cost of continuous monitors and the availability of new continuous monitors for mass, number, and certain chemical components, as well as older methods based on beta attenuation or light scattering. Most epidemiologic studies have used 24-hour concentrations as exposure indicators. However, one epidemiologic study of chronic effects uses a filter sampler with a two-week collection period (Gauderman et al., 2000). Another recent study used 1-2 h concentrations (see Peters et al., 2000). Continuous methods are discussed in Section 2.2.5.

2.2.2.6 Accuracy and Precision

Precision is typically determined by comparison of collocated samplers or through replicate analyses; whereas accuracy is determined through the use of traceable calibration standards. Unfortunately, no standard reference calibration material or procedure has been developed for suspended, atmospheric PM. It is possible to determine the accuracy of certain components of the PM measurement system (e.g., flow control, inlet aspiration, $PM_{2.5}$ cut, weighing, etc.). The absolute accuracy for collecting a test aerosol can also be determined by isokinetic sampling in a wind tunnel. However, it is not currently feasible to provide a simulated atmospheric aerosol with naturally occurring semivolatile components. It is particularly challenging to develop an atmospheric aerosol calibration standard suitable for testing samplers in the field. Therefore, it is not possible at the present time to establish the absolute accuracy of a PM monitoring technique. Intercomparison studies may be used to establish the precision of identical monitors and the extent of agreement between different types of monitors. Such studies are important for establishing the reliability of PM measurements. Intercomparison studies have contributed greatly to our understanding of the problems in PM measurement. Such studies will be discussed as they apply to specific measurement problems, monitoring instruments, or analytical techniques.

Some measurement errors of concern in PM_{10} sampling, including those that arise due to uncertainty tolerances in cutpoint, particle bounce and reentrainment, impactor surface overloading, and losses to sampler internal surfaces, were discussed in detail in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). Other measurement errors of concern in $PM_{2.5}$ sampling arise not only because of our inability to assess accuracy in an absolute sense due to a lack of an atmospheric aerosol calibration standard, but also because of the inclusion in $PM_{2.5}$ of a small amount of coarse particles as discussed in Section 2.2.1.3 and because of problems associated with the definition of $PM_{2.5}$ as what remains after collection on a filter and equilibration rather than the mass of particles as they exist in the air. Still, it is possible to measure PM indicators with high precision. Detailed information on precision and quality assurance may be found on EPA's Technology Transfer Network website (U.S. Environmental Protection Agency, 2002). See discussion in Section 2.2.4.

Because of the difficulties associated with determining the accuracy of PM measurements, EPA has sought to make FRM measurements equivalent by specifying operating conditions and, in the case of $PM_{2.5}$ samplers, by specifying details of the sampler design. Thus, both the PM_{10} as well as the $PM_{2.5}$ standards are defined with consistency of measurement technique rather than with the accuracy of the true mass concentration measurement in mind (McMurry, 2000). It is acknowledged in the Federal Register (1997) that, "because the size and volatility of the particles making up ambient particulate matter vary over a wide range and the mass concentration of particles varies with particle size, it is difficult to define the accuracy of $PM_{2.5}$ measurements in an absolute sense…." Thus, accuracy is defined as the degree of agreement between a field $PM_{2.5}$ sampler and a collocated $PM_{2.5}$ reference method audit sampler (McMurry, 2000). The Federal Reference Method for $PM_{2.5}$ is discussed in Section 2.2.3.3. As mentioned earlier, volatilization of organic compounds and ammonium nitrate during sampling or post-sampling handling can lead to significant underestimation of the fine particulate mass concentration in some locations. Sources of error in the measurement of mass of $PM_{2.5}$ suspended in the atmosphere also arise because of adsorption or desorption of semivolatile vapors onto or from collected PM, filter media, or other sampler surfaces; neutralization of acid or basic vapors on either filter media or collected PM; and artifacts associated with particle-bound water.

During the past 25 years, there have been advancements in the generation and classification of monodisperse aerosols, as well as in the development of electron microscopy and imaging

analysis, that have contributed to the advancement in aerosol calibration (Chen, 1993). Still, one of the limitations in PM sampling and analysis remains the lack of primary calibration standards for evaluating analytical methods and for intercomparing laboratories. Klouda et al. (1996) examined the possibility of resuspending the National Institute of Science and Technology (NIST) Standard Reference Material 1649 (Urban Dust) in air for collection on up to 320 filters simultaneously using Standard Research International's dust generation and collection system. However, the fine component is not resuspended and the semivolatile component has evaporated. Consequently, this material is not a suitable standard for suspended PM. Little additional work in this area has been reported.

Methods validation was discussed in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a), and the usefulness of intercomparisons and "internal redundancy" was emphasized. For example, a number of internal consistency checks are applied to the IMPROVE network (Malm et al., 1994). These include mass balances, sulfur measurements by both proton induced X-ray emission (PIXE) and ion chromatography (IC), and comparison of organic matter by combustion and by proton elastic scattering analysis (PESA) analysis of hydrogen. Mass balances compare the gravimetrically determined mass with the mass calculated from the sum of the major chemical components (i.e., crustal elements plus associated oxygen, organic carbon, elemental carbon, sulfate, nitrate, ammonium, and hydrogen ions). Mass balances are useful validation techniques; however, they do not check for, or account for, artifacts associated with the absorption of gases during sampling or the loss of semivolatile material during sampling. The mass balance check may appear reasonable even if such artifacts are present because only the material collected on the filter is included in the balance.

2.2.3 Measurement of Semivolatile Particulate Matter

It is becoming increasingly apparent that the semivolatile component of PM may significantly affect the quality of the measurement and can lead to both positive and negative sampling artifacts. Loss of semivolatile species, like ammonium nitrate and many organic species, may occur during sampling because of changes in temperature, relative humidity, or composition of the aerosol or because of the pressure drop across the filter (McMurry, 2000). Gas phase organic species, both volatile and semivolatile, may adsorb onto or react with filter media or collected PM, leading to a positive sampling artifact. Quartz fiber filters have a large

specific surface area on which adsorption of gases can occur. A number of other types of filters (e.g., stretched Teflon membrane filters) have much smaller exposed surface areas (Turpin et al., 1994) and appear to be subject to less adsorption (Kirchstetter et al., 2001; Turpin et al., 1994). Tsai and Huang (1995) observed positive sulfate and nitrate artifacts on high-volume PM_{10} quartz filters and attributed the artifacts to interactions between acidic gases (SO_2, HONO, and HNO_3) and both the filter media (either glass fiber or quartz) and the coarse particles collected on the filter. Volatilization losses also have been reported to occur during sample transport and storage (Chow, 1995). Evaporative losses of particulate nitrates have been investigated in laboratory and field experiments (e.g., Wang and John, 1988) and in theoretical studies (Zhang and McMurry, 1992). The results of recent studies that focused on volatilization losses of particulate nitrates are discussed in more detail in Section 2.2.3.1.

The theory describing phase equilibria of semivolative organic compounds (SVOC) continues to be developed. Liang et al. (1997), Jang et al. (1997), and Strommen and Kamens (1997) have modeled the gas/particle partitioning of SVOC on inorganic, organic, and ambient smog aerosols.

The positive artifact associated with adsorption of organic vapors onto quartz filters has been examined in experiments in which two quartz fiber filters were deployed in series. The second quartz filter may indicate gaseous volatile organic compounds (VOC) adsorbed on both filters (positive artifact), SVOC evaporated from particles on the first filter and subsequently adsorbed on the second filter (negative artifact), or a combination of both effects. Unless the individual compounds are identified, the investigator does not know what to do with the loading value on the second filter (i.e., to add or subtract from the first filter loading value). Moreover, even if the individual compounds were identified on the back-up filter, the decision concerning adding or subtracting the back-up filter loading would not be straightforward.

The developing state of the art in which diffusion denuder technology is being applied to SVOC sampling (e.g., Eatough et al., 1993; Gundel et al., 1995), as well as for sampling of gas and particulate phase organic acids (Lawrence and Koutrakis, 1996a,b), holds promise for improving the understanding of SVOC sampling artifacts. In a denuder-based system, gas-phase organics are removed by diffusion to an adsorbent surface (e.g., activated carbon, special polymer resins, etc.). Particles then are collected on a filter downstream of the denuder and the remaining organic vapors (i.e., from denuder breakthrough and volatile losses from the collected

particles) are collected in an adsorbent downstream of the filter (e.g., charcoal or carbon-impregnated filters, polyurethane foam, or polystyrene-divinylbenzene resin [XAD]). The results of recent studies that have focused on treatment of both positive and negative sampling artifacts associated with SVOC are discussed in Section 2.2.3.2. Detailed information on the use of denuder systems to measure semivolatile species is provided in Section 2.2.3.3.

Finally, Eatough et al. (1999a) have reported on a batch sampler that attempts to correct simultaneously for volatilization losses of both nitrate and SVOC. These samplers are also discussed in more detail in Section 2.2.3.2.

2.2.3.1 Particulate Nitrates

It is well known that volatilization losses of particulate nitrates (e.g., Zhang and McMurry [1992]; see also Hering and Cass [1999] and references therein) occur during sampling on Teflon filters. The affect on the accuracy of atmospheric particulate measurements from these volatilization losses is more significant for $PM_{2.5}$ than for PM_{10}. The FRM for $PM_{2.5}$ will likely suffer loss of nitrates similar to that experienced with other simple filter collection systems. Sampling artifacts resulting from the loss of particulate nitrates represents a significant problem in areas such as southern California that experience high amounts of nitrates. Hering and Cass (1999) examined the errors in $PM_{2.5}$ mass measurements because of volatilization of particulate nitrate by looking at data from two field measurement campaigns conducted in southern California: (1) the Southern California Air Quality Study (SCAQS) (Lawson, 1990) and (2) the 1986 CalTech study (Solomon et al., 1992). In both these studies, side-by-side sampling of $PM_{2.5}$ was conducted. One sampler collected particles directly onto a Teflon filter. The second sampler consisted of a denuder to remove gaseous nitric acid followed by a nylon filter that absorbs the HNO_3 which evaporates from ammonium nitrate. In both studies, the denuder consisted of MgO-coated glass tubes (Appel et al., 1981). Fine particulate nitrate collected on the Teflon filter was compared to fine particulate nitrate collected on the denuded nylon filter. In both studies, the $PM_{2.5}$ mass lost because of volatilization of ammonium nitrate represented a significant fraction of the total $PM_{2.5}$ mass. The fraction of mass lost was higher during summer than during fall (17% versus 9% during the SCAQS study and 21% versus 13% during the CalTech study; Figure 2-12). In regard to percentage loss of nitrate, as opposed to percentage



Figure 2-12. Amount of ammonium nitrate volatilized from Teflon filters, expressed as a percentage of the measured $PM_{2.5}$ mass, for the SCAQS and CalTech studies, for spring and fall sampling periods.

Source: Herring and Cass (1999).

loss of mass discussed above, Hering and Cass (1999) found that nitrate remaining on the Teflon filter samples was on average 28% lower than that on the denuded nylon filters.

Hering and Cass (1999) also analyzed these data by extending the evaporative model developed by Zhang and McMurry (1987). The extended model utilized by Hering and Cass (1999) takes into account dissociation of collected particulate ammonium nitrate on Teflon filters into nitric acid and ammonia via three mechanisms: (1) scrubbing of nitric acid and ammonia in the sampler inlet (John et al., 1988 showed that clean PM_{10} inlet surfaces serve as an effective denuder for nitric acid), (2) heating of the filter substrate above ambient temperature by sampling, and (3) pressure drop across the Teflon filter. For the sampling systems modeled, the flow-induced pressure drop was measured to be less than 0.02 atm, and the corresponding change in vapor pressure was 2%, so losses driven by pressure drop were not considered to be significant in this work. Losses from Teflon filters were found to be higher during the summer compared to the winter, higher during the day compared to night, and reasonably consistent with modeled predictions.

Finally, during the SCAQS study, particulate samples also were collected using a Berner impactor and greased Tedlar substrates in size ranges from 0.05 to 10 μm in aerodynamic diameter. The Berner impactor $PM_{2.5}$ nitrate values were much closer to those from the denuded nylon filter than those from the Teflon filter with the impactor nitrate being approximately 2% lower than the nylon filter nitrate for the fall measurements and approximately 7% lower during the summer measurements. When the impactor collection was compared to the Teflon filter collection for a nonvolatile species (sulfate), the results were in agreement.

It should be noted that filters or collection surfaces were removed immediately after sampling and placed into vials containing a basic extraction solution during these intercomparison studies. Therefore, losses that might occur during handling, storage, and equilibration of filters or impaction surfaces were avoided. The loss of nitrate observed from Teflon filters and impaction surfaces in this study, therefore, is a lower limit compared to losses that might occur during the normal processes involved in equilibration and weighing of filters and impaction surfaces. Brook and Dann (1999) observed much higher nitrate losses during a study in which they measured particulate nitrate in Windsor and Hamilton, Ontario, Canada, by three techniques: (1) a single Teflon filter in a dichotomous sampler, (2) the Teflon filter in an annular denuder system (ADS), and (3) total nitrate including both the Teflon filter and the nylon back-up filter from the ADS. The Teflon filter from the dichotomous sampler averaged only 13% of the total nitrate. The Teflon filter from the ADS averaged 46% of the total nitrate. The authors concluded that considerable nitrate was lost from the dichotomous sampler filters during handling, which included weighing and x-ray fluorescence (XRF) measurement in a vacuum.

Kim et al. (1999) also examined nitrate sampling artifacts by comparing denuded and undenuded quartz and nylon filters, during the PM_{10} Technical Enhancement Program (PTEP) in the South Coast Air Basin of California. They observed negative nitrate artifacts (losses) for most measurements; however, for a significant number of measurements they observed positive nitrate artifacts. Kim et al. (1999) pointed out that random measurement errors make it difficult to measure true amounts of nitrate loss.

Several diffusion denuder samplers have been developed to account for the nitrate lost because of volatilization from filters, many of which were discussed in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). Eatough et al. (1999a) developed a high-volume diffusion denuder system in which diffusion denuder and particle concentrator

techniques were combined (see Section 2.2.3.2). The particle concentrator reduces the flow through the denuder so that the denuder can be operated for weeks without a loss of collection efficiency, making the sampler suitable for routine field sampling. The system was evaluated for the collection of fine particulate sulfate and nitrate in Riverside, CA (Eatough et al., 1999b). Concentrations of $PM_{2.5}$ nitrate obtained from the PC-BOSS agreed with those obtained using the Harvard-EPA Annular Denuder Sampler, HEADS (Koutrakis et al., 1988).

2.2.3.2 Semivolatile Organic Compounds

In addition to their contribution to suspended PM mass, SVOC are also of interest because of their possible health effects. SVOC include products of incomplete combustion such as polycyclic aromatic hydrocarbons (PAHs) and polycyclic organic matter, which has been identified as a hazardous air pollutant. PAHs also have been suggested as alternative particulate tracers for automobile emissions because the phase-out of organo-lead additives to gasoline means that lead is no longer a good tracer for automobiles (Venkataraman et al., 1994). PAHs also are emitted during biomass burning, including burning of cereal crop residues and wood fuels (Jenkins et al., 1996; Roberts and Corkill, 1998).

The positive quartz filter artifact was previously mentioned and has been discussed by others (Gundel et al., 1995; Turpin et al., 2000). It is also possible that some SVOC may desorb from the filter resulting in a negative artifact (Eatough et al., 1993; Tang et al., 1994; Eatough et al., 1995; Gundel et al., 1995; Cui et al., 1998; Pang et al., 2001; Finn et al., 2001). Semivolatile organic compounds can similarly be lost from Teflon filters because of volatilization, causing the $PM_{2.5}$ mass to be significantly underestimated (negative artifact). Like particulate nitrates, the FRM for $PM_{2.5}$ will suffer loss of SVOC, similar to the losses experienced with other simple filter collection systems. Most studies that have focused on the positive and negative sampling artifacts associated with SVOC compounds have utilized either diffusion denuder technology or placed an adsorbent media, such as a back-up quartz filter or a polyurethane foam adsorbent behind the main filter.

Using their multichannel diffusion denuder sampling system (BOSS), Eatough et al. (1995) reported that, for samples collected at the South Coast Air Quality Management District sampling site at Azusa, CA, changes in the phase distribution of SVOC could result in a loss on average of 35% of the particulate organic material. Cui et al. (1998) found that losses of SVOC

from particles in the Los Angeles Basin during the summer were greater during the night (average = 62%) than during the day (average = 42%).

The percent SVOC lost from the front filter in a filter-denuder system may be greater than that lost in a filter-only system such as the FRM. In a filter-denuder system, the gas-phase component of the SVOC is removed. The absence of the gas-phase causes the gas-particle equilibrium to shift so the SVOC collected on the filter may evaporate more rapidly in a filter-denuder system than in a filter-only collection system. To determine the fraction of SVOC lost from a Teflon filter in a filter-only system, it is necessary to compare the amount measured by a nondenuder system with that measured by a denuder system. At present, little information is available on the volatilization losses of SVOC. However, in one study (Pang et al., 2001), the total mass on denuded and undenuded filters were compared and found to be identical within error limits (R^2 = 0.816, slope = 0.961 ± 0.027 for total mass compared to R^2 = 0.940, slope = 0.986 ± 0.020 for sulfate). Pang et al. interpreted this result as suggesting that the major cause of loss of SVOC is the pressure drop across the filter.

Positive artifacts may occur during sample collection because of the adsorption of gases onto the filter materials (e.g., Gundel et al., 1995). Using a quartz filter behind a Teflon filter, Kim et al. (2001) estimated that on an annual average basis 30% of the $PM_{2.5}$ organic carbon concentration resulted from positive artifacts. There is a larger positive artifact because of greater adsorption of organic vapor onto quartz fiber filters than onto Teflon filters (Turpin et al., 1994; Chow et al., 1994, 1996; Eatough et al., 1996; Finn et al., 2001).

Kirchstetter et al. (2001) report that adsorptive properties of quartz fiber filters vary with lot number; therefore, front and back-up filters should be taken from the same lot. Recent literature suggests that a Teflon filter followed by a quartz back-up filter appears to provide a better estimate of the adsorption of gases on a quartz fiber front filter than does a quartz filter followed by a quartz backup and that the difference between these two adsorption estimates can be substantial for short durations (Novakov et al., 1997; Kirchstetter et al., 2001; Turpin et al., 2000). The typically lower organic carbon loadings on concurrently collected quartz followed by quartz filters relative to Teflon followed by quartz filters are believed to occur because adsorption on the quartz front filter acts to reduce the gas-phase concentration downstream until adsorption equilibrium has been achieved in the vicinity of the front quartz filter surface. Because Teflon filters have little affinity for organic vapors, this equilibrium occurs almost

instantaneously for Teflon filters, and the Teflon-quartz back-up filter is exposed to the ambient concentration of organic vapors from the beginning of the sampling period. It might be expected that the quantity of organic vapor adsorbed on quartz filters would depend on the organic composition and would vary by season and location.

Use of Denuder Systems To Measure Semivolatile Compounds

Phase distribution of semivolatile organic species has been the subject of several studies that have employed denuder technology (see Gundel et al., 1995; Gundel and Lane, 1999) to directly determine the phase distributions while avoiding some of the positive and negative sampling artifacts associated with using back-up quartz filters. In an ideal system with a denuder that is 100% efficient, the gas phase would be collected in the denuder and the particle phase would be the sum of the material collected on the filter and the adsorbent downstream. Denuder collection efficiency depends on the denuder surface area (+), the diffusivity (+) and vapor pressure (−) of the compound, the temperature (−) and flow rate (−) of the air stream, and the presence of competing species (−), including water vapor (Cui et al., 1998; Kamens and Coe, 1997; Lane et al., 1988). (The + and – symbols in parentheses indicate qualitatively the effect increasing each parameter would have on efficiency). In a system with a denuder collection efficiency less than 100%, the collection efficiency must be known to accurately attribute adsorbed organics from denuder breakthrough to the gas phase and adsorbed organics volatilized from collected particles to the particle phase. In calculating the overall phase distributions of SVOC PAH from a denuder system, the collection efficiency for each compound is needed.

The efficiency of silicone-grease-coated denuders for the collection of polynuclear aromatic hydrocarbons was examined by Coutant et al. (1992), who examined the effects of uncertainties in the diffusion coefficients and in the collisional reaction efficiencies on the overall phase distributions of SVOC PAH calculated using denuder technology. In their study, they used a single stage, silicone-grease-coated aluminum annular denuder with a filter holder mounted ahead of the denuder and an XAD trap deployed downstream of the denuder. In a series of laboratory experiments, they spiked the filter with a mixture of perdeuterated PAH, swept the system with ultra-high purity air for several hours, and then analyzed the filter and the XAD. They found that the effects of these uncertainties, introduced by using a single compound as a surrogate PAH (in their case, naphthalene) for validation of the denuder collection efficiency, are

less significant than normal variations because of sampling and analytical effects. Results on field studies using their sampling system have not been published.

For measuring particulate phase organic compounds, the denuder-based sampling system represents an improvement over the filter/adsorbent collection method (Turpin et al., 1993). Some researchers, however, have reported that denuder coatings themselves can introduce contamination (Mukerjee et al., 1997) and that the adsorbed species may be difficult to remove from the coating (Eatough et al., 1993).

In a study conducted in southern California (Eatough et al., 1995), the Brigham Young University Organic Sampling System (BOSS; Eatough et al., 1993) was used for determining POM composition, and a high-volume version (BIG BOSS; flow rate 200 L/min) was utilized for determining the particulate size distribution and the chemical composition of SVOC in fine particles. The BOSS, a multi-channel diffusion denuder sampling system, consists of two separate samplers (each operating at 35 L/min). The first sampler consists of a multi-parallel plate diffusion denuder with charcoal-impregnated filter papers as the collection surfaces followed by a two-stage quartz filter pack and a two-stage charcoal-impregnated filter pack. The second sampler operating in parallel with the first consists of a two-stage quartz filter pack, followed by the parallel plate denuder, followed by the two-stage charcoal-impregnated filter pack. The filter samples collected by the BOSS sampler were analyzed by temperature-programmed volatilization analysis. The second channel allows calculations of the efficiency of the denuder in removing gas-phase specifics that would be absorbed by the charcoal impregnated filter. Eatough et al. (1995) also operated a two-stage quartz filter pack alongside the BOSS sampler. The BIG BOSS system (Tang et al., 1994) consists of four systems (each with a flowrate of 200 L/min). Particle size cuts of 2.5, 0.8, and 0.4 μm are achieved by virtual impaction, and the sample subsequently flows through a denuder, then is split, with the major flow (150 L/min) flowing through a quartz filter followed by an XAD-II bed. The minor flow is sampled through a quartz filter backed by a charcoal-impregnated filter paper. The samples derived from the major flow (quartz filters and XAD-II traps) were extracted with organic solvents and analyzed by gas chromatography (GC) and GC-mass spectroscopy. The organic material lost from the particles was found to represent all classes of organic compounds.

Eatough et al. (1996) operated the BOSS sampler for a year at the IMPROVE site at Canyonlands National Park, UT, alongside the IMPROVE monitor and alongside a separate

sampler consisting of a two-stage quartz filter pack. They found that concentrations of particulate carbon determined from the quartz filter pack sampling system were low on average by 39%, and this was attributed to volatilization losses of SVOC from the quartz filters. In another study conducted with the BOSS in southern California, losses of 35% of the POM, on average, were found and attributed to losses of the SVOC during sampling (Eatough et al., 1995).

The denuder used in the various BOSS samplers consists of charcoal-impregnated cellulose fiber filter material. Denuder collection efficiencies of greater than 95% have been reported for organic gases that adsorb on quartz and charcoal-impregnated filters (Eatough et al., 1999a; Ding et al., 2002; Lewtas et al., 2001). However, because the mass concentration of gas phase species that adsorb on quartz and charcoal-impregnated filters is so much greater than the mass of semivolatile organic material in the particulate phase, it is necessary to measure and account for the inefficiency of the denuder in the BOSS samplers. To address this problem, Brigham Young University (BYU) developed a particle-concentrator (PC)-BOSS system (Ding et al., 2002; Eatough et al., 1999b; Lewtas et al., 2001; Modey et al., 2001; Pang et al., 2001, 2002a,b). The PC-BOSS includes a virtual impactor upstream of the denuder to improve the denuder collection efficiency by removing a majority of the gases from the aerosol flow. With this system, denuder collection efficiencies of greater than 99% have been reported for organic gases, SO_2(g), HNO_3(g) and other species that adsorb on quartz and charcoal-impregnated filters (Pang et al., 2001). Since the concentrations of semivolatile organic and other gases in the presence of the concentrated particles is not altered by this process, the gas-particle distribution should be little altered by the concentration process. The virtual impactor has a 50% cut point at 0.1 μm aerodynamic diameter. As a result, some particles in the 0.05 to 0.2 μm diameter size range will be removed in the major flow along with the majority of the gases. Therefore, the mass collection efficiency of the virtual impactor concentrator will be a function of the particle size distribution in the 0.05 to 0.1 μm size range. This collection efficiency is measured by comparing the concentration of nonvolatile components measured in the concentrated sample with that measured in an unconcentrated sample. The concentration efficiency varies from 50 to 75%. It is relatively constant over periods of weeks but varies by season and site, presumably as the particle size distribution changes. Previous studies at Harvard (Sioutas et al., 1995a,b) have shown that the composition of the sampled aerosol is little changed by the concentration process. The BYU studies listed above have shown that the concentration efficiencies for sulfate, organic

carbon (OC) and elemental carbon (EC) are comparable for a given sampling location. Furthermore, the concentrations of these species and of fine particulate nitrate determined using the PC-BOSS have been shown to be comparable to those determined using more conventional samplers for sulfate or EC or using simpler denuder systems for OC and nitrate.

Ding et al. (1998a) developed a method for the determination of total n-nitroso compounds in air samples and used the method to examine organic compounds formed from NO_x chemistry in Provo, UT (Ding et al., 1998b). In their method, n-nitroso compounds are selectively decomposed to yield nitric oxide, which is then detected using chemiluminescence. From the samples from Provo, they found that the majority of the n-nitroso and nitrite organic compounds that were present in fine particulate matter were semivolatile organic compounds that could be evaporated from the particles during sampling. They found particulate n-nitroso compound concentrations ranging between <1 and 3 nmoles/m^3 and gas-phase n-nitroso compound concentrations in the same range. Particulate organic nitrite concentrations were found in the range of <1 to ≈5 nmoles/m^3, and gas-phase concentrations as high as 10 nmoles/m^3 were found.

Turpin et al. (1993) developed a sampling system that corrects for the loss of semivolatile organic compounds during sampling by removal of most of the gas phase material from the particles in a diffusion separator sampling system. Unlike the previously mentioned systems, wherein the particulate phase is measured directly, in the system of Turpin et al. (1993) the gas-phase is measured directly. In the laminar flow system, ambient, particle-laden air enters the sampler as an annular flow. Clean, particle-free air is pushed through the core inlet of the separator. The clean air and ambient aerosol join downstream of the core inlet section, and flow parallel to each other through the diffusion zone. Because of the much higher diffusivities for gases compared to particles, the SVOC in the ambient air diffuses to the clean, core flow. The aerosol exits the separator in the annular flow, and the core flow exiting the separator now contains a known fraction of the ambient SVOC. Downstream of the diffusion separator, the core exit flow goes into a polyurethane foam (PUF) plug, where the SVOC is collected. The adsorbed gas phase on the PUF plug is extracted with supercritical fluid CO_2 and analyzed by gas chromatography/mass-selective detection (GC/MSD). The gas-phase SVOC is thus determined. Ultimately, to determine particulate phase SVOC concentrations, the total compound concentration will also be measured and the particulate phase obtained by difference. The system was tested for the collection of PAH. The diffusional transport of gas-phase PAHs and particle

concentrations agreed well with theory. Breakthrough was problematic for low molecular weight PAHs (MW < 160). Detection limits ranged from 20 to 50 pg of injected mass for all PAHs.

Gundel et al. (1995) recently developed a technique for the direct determination of phase distributions of semivolatile polycyclic aromatic hydrocarbons using annular denuder technology. The method, called the integrated organic vapor/particle sampler (IOVPS), uses a cyclone inlet with a 50% cutpoint of 2.5 μm at a sampling rate of 10 L/min. The airstream then goes through two or three sandblasted glass annular denuders that are coated with ground adsorbent resin material (XAD-4 was initially examined) that traps vapor-phase organics. The airstream subsequently passes through a filter, followed by a backup denuder. The denuder collection efficiency is high and compares well with predictions based on the diffusivity of the compounds. The denuder can also be extracted to obtain gas-phase concentrations directly (Gundel and Lane, 1999). Particle-phase PAHs are taken to be the sum of material on the filter and XAD adsorbent downstream after correction for denuder collection efficiency. The IOVPS was tested for sampling semivolatile PAH in laboratory indoor air and in environmental tobacco smoke (ETS). After exposure, the denuders, filters, and sorbent traps were extracted with cyclohexane (Gundel et al., 1995) and analyzed for PAHs from naphthalene to chrysene using dual-fluorescence detection (Mahanama et al., 1994). Recoveries from both denuders and filters were approximately 70% for 30 samples. Detection limits (defined as 3 times the standard deviation of the blanks) for gas-phase SVOC PAHs ranged from 0.06 ng for anthracene to 19 ng for 2-methylnaphthalene. The 95% confidence interval for reproduction of an internal standard concentration was 6.5% of the mean value. Relative precision, from a propagation of errors analysis or from the 95% confidence interval from replicate analyses of standard reference material SRM 1649 (urban dust/organics), was 12% on average (8% for naphthalene to 22% for fluorene). Sources of error included sampling flow rate, internal standard concentration, and co-eluting peaks. Gundel and Lane (1999) reported that roughly two-thirds of particulate PAH fluoranthene, pyrene, benz[*a*]anthracene, and chrysene were found on the postfilter denuders, so that it is likely that considerable desorption from the collected particles took place.

Solid adsorbent-based denuder systems have been investigated by other researchers as well. Bertoni et al. (1984) described the development of a charcoal-based denuder system for the collection of organic vapors. Risse et al. (1996) developed a diffusion denuder system to sample aromatic hydrocarbons. In their system, denuder tubes with charcoal coating and charcoal paper

precede a filter pack for particulate collection and an adsorption tube to capture particle blow-off from the filter sample. Breakthrough curves for benzene, toluene, ortho-xylene, and meta-xylene were developed for 60-, 90-, and 120-cm denuder tubes. The effects of relative humidity on the adsorption capacities of the denuder system were examined, and it was found that the capacity of the charcoal was not affected significantly by increases in relative humidity. The feasibility of outdoor air sampling with the system was demonstrated.

Krieger and Hites (1992) designed a diffusion denuder system that uses capillary gas chromatographic columns as the tubes for SVOC collection. The denuder was followed by a filter to collect particles, which in turn was followed by a PUF plug to collect organic material volatilizing off the filter. Denuder samples were analyzed by liquid solvent extraction (CH_2Cl_2) followed by GC-mass spectrometric analysis. The PUF plugs and filters were extracted with supercritical fluid extraction using supercritical N_2O. Using this system, an indoor air sample was found to contain primarily chlorinated biphenyls, ranging from trichlorobiphenyls (vapor pressures 10^{-3} – 10^{-4} Torr at 25 °C) to octachlorobiphenyls (10^{-6} – 10^{-7} Torr). This demonstrated that the sampler collects compounds with a wide range of volatility. They also found that on-line desorption is successful in maintaining good chromatographic peak shape and resolution. The entire method, from sample collection to the end of the chromatographic separation, took 2 h.

Organic acids in both the vapor and particulate phases may be important contributors to ambient acidity, as well as representing an important fraction of organic particulate matter. Lawrence and Koutrakis (1996a,b) used a modified Harvard/EPA annular denuder system (HEADS) to sample both gas and particulate phase organic acids in Philadelphia, PA, in the summer of 1992. The HEADS sampler inlet had a 2.1-μm cutpoint impactor (at 10 L/min), followed by two denuder tubes, and finally a filter pack with a Teflon filter. The first denuder tube was coated with potassium hydroxide (KOH) to trap gas phase organic acids. The second denuder tube was coated with citric acid to remove ammonia and thus to avoid neutralizing particle phase acids collected on the filter. The KOH-coated denuder tube was reported to collect gas phase formic and acetic acids at better than 98.5% efficiency and with precisions of 5% or better (Lawrence and Koutrakis, 1994). It was noted that for future field measurements of particulate organic acids, a Na_2CO_3-coated filter should be deployed downstream of the Teflon filter to trap organic acids that may evaporate from the Teflon filter during sampling.

Role of the Collection Media

The role of the collection media was recently examined in a study conducted in Seattle (Lewtas et al., 2001). In that study, the influence of denuder sampling methods and filter collection media on the measurement of SVOC associated with $PM_{2.5}$ was evaluated. Activated carbon and XAD collection media were used in diffusion denuders and impregnated back-up filters in two different samplers, the Versatile Air Pollution Sampler (VAPS) and the PC-BOSS. XAD-coated glass annular denuders and charcoal-impregnated cellulose fiber (CIF) filter denuders also were used. CIF filters also were compared to XAD-coated quartz filters as backup filter collection media. Lewtas et al. (2001) found that the two denuder types resulted in an equivalent measurement of particulate organic carbon and particle mass. The carbon-coated denuders in the BOSS sampler were more efficient than the XAD-coated denuders for the collection of the more highly volatile organic compounds (MHVOC). Lewtas et al. (2001) concluded that this MHVOC that is collected in the carbon-coated BOSS denuder does not contribute substantially to the particle mass or to the SVOC measured as OC on quartz filters. However, this MHVOC would be captured in the carbon impregnated filters placed behind the quartz filters, so that, in the XAD denuder configuration, the captured MHVOC would cause a higher OC concentration and an overestimation of the SVOC.

Some of the recent research in denuder technology also has focused on reduction in the size of the denuder, optimization of the residence time in the denuder, understanding the effect of diffusion denuders on the positive quartz filter artifact, identifying changes in chemical composition that occur during sampling, determining the effects of changes in temperature and relative humidity, and identifying possible losses by absorption in coatings.

Reducing the Size of Denuders

The typical denuder configuration is an annular diffusion denuder tube of significant length (e.g., 26.5 cm for 10 L/min [Koutrakis et al., 1988]). A more compact design based on a honeycomb configuration was shown to significantly increase the capacity (Koutrakis et al., 1993). However, in intercomparisons with an annular denuder/filter pack system (Koutrakis et al., 1988), significant losses of ammonia and nitric acid were observed for the honeycomb configuration and were attributed to the large inlet surface area and long sample residence time of the honeycomb design relative to the annular denuder system. Sioutas et al. (1996a)

subsequently designed a modified glass honeycomb denuder/filter pack sampler (HDS) with an inlet that minimizes vapor losses on the inlet surfaces. The modified HDS has reduced inlet surfaces and decreased residence time for sampled gases (NH_3 and HNO_3) compared to its predecessor (Sioutas et al., 1994a). Sioutas et al. (1996b) also tested various inlet materials (glass, PFA, and polytetrafluoroethylene [PTFE]) in laboratory tests and found that a PTFE Teflon coated inlet minimized loss of sampled gases (1 to 8% loss of HNO_3 observed, and –4 to 2% loss of NH_3 observed). The highest inlet losses were observed for HNO_3 lost to PFA surfaces (14 to 25%). The modified HDS was tested in laboratory and field tests and found to agree within 10% with the annular denuder system.

Residence Time in the Denuder

The efficiency of a diffusion denuder sampler for the removal of gas phase material can be improved by increasing the residence time of the sampled aerosol in the denuder. However, the residence time can only be increased within certain limits. Because the diffusion denuder reduces the concentration of gas-phase semivolatile organic material, semivolatile organic matter present in the particles passing through the denuder will be in a thermodynamically unstable environment and will tend to outgas SVOC during passage through the denuder. The residence time of the aerosol in the denuder, therefore, should be short enough to prevent significant loss of particulate phase SVOC to the denuder. Various studies have suggested that the residence time in the denuder should be less than about 2 s (Gundel and Lane, 1999; Kamens and Coe, 1997; Kamens et al., 1995). The residence times in the various denuder designs described by Gundel and Lane (1999) are from 1.5 to 0.2 s. The equilibria and evaporation rates are not as well understood for organic components as they are for NH_4NO_3 (Zhang and McMurry, 1987, 1992; Hering and Cass, 1999).

Effect of Diffusion Denuders on the Positive Quartz Filter Artifact

The adsorption of organic compounds by a second quartz filter has been shown to be reduced, but not eliminated, in samples collected in the Los Angeles Basin if a multi-channel diffusion denuder with quartz filter material as the denuder collection surface preceded the quartz filters (Fitz, 1990). This artifact can be further reduced by the use of activated charcoal as the denuder surface and the use of a particle concentrator to reduce the amount of gas phase organic

compounds relative to condensed phase organic compounds (Cui et al., 1997, 1998; Eatough, 1999). Recent experiments (Cotham and Bidleman, 1992; Cui et al., 1998; Eatough et al., 1995, 1996) have shown that the quartz filter artifact can result both from the collection of gas phase organic compounds and from the collection of semivolatile organic compounds lost from particles during sampling. Thus, results available to date suggest that both a "positive" and a "negative" artifact can be present in the determination of particulate phase organic compounds using two tandem quartz filters.

The importance of the adsorption of organic vapors on filters or PM relative to the volatilization of organic compounds from PM collected on a filter continues to be a topic of active debate. The relative importance of positive and negative artifacts will be different for denuded and undenuded filters; will depend on face velocity, sample loading, and the vapor pressures of the compounds of interest; and may vary with season and location because of variations in the composition of volatile and semivolatile organic material. Evidence exists for substantial positive and negative artifacts in the collection of organic PM.

Changes in Chemical Composition During Sampling

The use of sampling systems designed to correctly identify the atmospheric gas and particulate phase distributions of collected organic material has been outlined above. An additional sampling artifact that has received little consideration in the collection of atmospheric samples is the potential alteration of organic compounds as a result of the sampling process. These alterations appear to result from the movement of ambient air containing oxidants and other reactive compounds past the collected particles. The addition of NO_2 (<1ppm) or O_3 (<200 ppb) to the sampled air stream (0 to 5 °C) for a high-volume sampler reduced the concentrations of benzo[*a*]pyrene and benzo[*a*]anthracene from a few % to 38%, with the observed reduction increasing with increased concentration of the added gases (Brorström et al., 1983). Spiking a filter with an amine resulted in an increase in measured concentrations of nitrosamines in both the filter and a following XAD sorbent bed for a mid-volume sampler (Ding et al., 1998a,b). Similar results have been obtained for the exposure of a deuterated amine on a filter to NO_x (Pellizzari and Krost, 1984). When Tenax columns spiked with deuterated styrene and cyclohexene were exposed to ppm concentrations of ozone or halogens, oxygenated and halogenated compounds were formed (Pellizzari and Krost, 1984).

Similar oxidation of aldehydes and peroxyacetylnitrate (PAN) during sampling has been observed (Grosjean and Parmar, 1990). Collected PAH compounds can be oxygenated or nitrated on a filter (Davis et al., 1987; Lindskog and Brorstrom-Lunden, 1987), but 1-nitropyrene has been shown to be resistant to additional nitration (Grosjean, 1983). These various chemical transformations of collected organic compounds can be eliminated by removal of the gas phase oxidants, NO_x, HNO_3, etc., or by reaction or absorption prior to collection of the particles (Ding, 1998a,b; Grosjean and Parmar, 1990; Parmar and Grosjean, 1990; Pellizzari and Krost, 1984; Williams and Grosjean, 1990). The BOSS denuder should be effective in eliminating most of the chemical transformation artifacts because reactive gases are removed by the charcoal denuder that precedes the particle collection filter.

Temperature and Relative Humidity Effects

The problems of sampling artifacts associated with SVOC adsorption and evaporation are compounded by temperature and relative humidity effects (Pankow and Bidleman, 1991; Pankow et al., 1993; Falconer et al., 1995; Goss and Eisenreich, 1997). Effects of temperature on the partitioning of PAH were examined by Yamasaki et al. (1982), who found that the partition coefficient (PAH_{vapor}/PAH_{part}) was inversely related to temperature and could be described using the Langmuir adsorption concept. The dissociation of ammonium nitrate aerosol is also a function of temperature. Bunz et al. (1996) examined the dissociation and subsequent redistribution of NH_4NO_3 within a bimodal distribution using a nine-stage low-pressure Berner impactor followed by analysis by ion chromatography and found a strong temperature dependency on the redistribution. Bunz et al. (1996) found that at lower temperatures (below 10 °C) there was little change in the aerosol size distribution. At temperatures between 25 and 45 °C, however, the lifetime of NH_4NO_3 particles decreases by more than a factor of 10, and size redistribution, as measured by average ending particle diameter, increased more for higher temperatures than for lower temperatures.

The effects of relative humidity on the sorption of SVOC on particles are not well understood. In a series of laboratory experiments, Goss and Eisenreich (1997) examined the sorption of both nonpolar (hydrocarbons and chlorinated hydrocarbons) and polar (ethyl ether and acetone) volatile organic compounds onto combustion soot particles as a function of temperature and relative humidity. The soot particles used in their experiments were collected

from oil furnaces and contained 60% (w/w) iron sulfate (water-soluble fraction) and 9% (w/w) elemental and organic carbon. They found that, for all compounds, the sorption of VOC onto soot particles decreased with increasing relative humidity over the range of 10 to 95%. They also observed hysteresis in the relative humidity dependency, with sorption coefficients at a given relative humidity higher when the RH is being increased than when the RH is being decreased. The sorption coefficients were fit with an exponential function to the RH so that the slope of the regression line would provide a measure of the influence of relative humidity. Based on the magnitude of the slope, they concluded that the RH-dependency of sorption was stronger for water-soluble organic compounds.

In another study by Jang and Kamens (1998), humidity effects on gas-particle partitioning of SVOC were examined using outdoor environmental chambers and the experimentally determined partitioning coefficients were compared to theoretical values. They examined the partitioning of SVOC onto wood soot, diesel soot, and secondary aerosols and concluded that "the humidity effect on partitioning was most significant for hydrophobic compounds adsorbing onto polar aerosols." Although these two studies seem to be contradictory, on closer examination, it is difficult to compare the two studies for several reasons. The experiments conducted by Jang and Kamens (1998) were conducted in outdoor chambers at ambient temperatures and humidities. Their model was for absorptive partitioning of SVOC on *liquid-like* atmospheric particulate matter. In contrast, the results of Goss and Eisenreich (1997) were obtained from a gas chromatographic system operated at 70 °C higher than ambient conditions. The model of Goss and Eisenreich (1997) was for adsorptive partitioning of VOC on *solid-like* atmospheric particulate matter. In the study of Jang and Kamens (1998), calculated theoretical values for water activity coefficients for diesel soot were based on an inorganic salt content of 1 to 2%; whereas, the combustion particles studied by Goss and Eisenreich (1997) contained 60% water-soluble, inorganic salt content. Jang and Kamens (1998) obtained their diesel soot from their outdoor chamber, extracted it with organic solvent (mixtures of hexane and methylene chloride), and measured the organic fraction. The resulting salt content of 2% of the particulate matter studied in Jang and Kamens (1998) is enough to affect water uptake but presumably not to affect the sorption partitioning of organics.

Impactor Coatings

Impactors are used as a means to achieve a size cutpoint and as particle collection surfaces. Particles collected on impactors are exposed to smaller pressure drops than filter-collected particles, making them less susceptible to volatile losses (Zhang and McMurry, 1987). However, size resolution can be affected by bounce when samples are collected at low humidities (Stein et al., 1994). There are other sources of error inherent in some of the currently acceptable practices that could potentially affect particulate mass concentration measurements and that will surely become even more important as more emphasis is placed on chemical speciation. Allen et al. (1999a) reported that the practice of greasing impaction substrates may introduce an artifact from the absorption of semivolatile species from the gas phase by the grease because the grease could artificially increase the amount of PAHs and other organic compounds attributed to the aerosol. Allen et al. (1999a) offer several criteria to ensure that this absorption artifact is negligible, including selecting impaction oils in which analytes of interest are negligibly soluble and ensuring that species do not have time to equilibrate between the vapor and oil phases (criterion is met for nonvolatile species). They recommend using oiled impaction substrates only if the absorption artifact is negligible as determined from these criteria. Application of greases and impaction oils for preventing or reducing bounce when sampling with impactors is not suitable for carbon analysis because the greases contain carbon (Vasilou et al., 1999).

Kavouras and Koutrakis (2001) investigated the use of polyurethane foam (PUF) as a substrate for conventional inertial impactors. The PUF impactor substrate is not rigid like the traditional impactor substrate so particle bounce and reentrainment artifacts are reduced significantly. Kavouras and Koutrakis (2001) found that the PUF impaction substrate resulted in a much smaller 50% cut point at the same flow rate and Reynolds number. Moreover, the lower 50% cut point was obtained at a lower pressure drop than with the conventional substrate, which could lead to a reduction of artifact vaporization of semivolatile components.

2.2.3.3 Particle-Bound Water

It is generally desirable to collect and measure ammonium nitrate and semivolatile organic compounds. However, for many measurements of suspended particle mass, it is desirable to remove the particle-bound water before determining the mass. In other situations it may be important to know how much of the suspended particle's mass or volume results from particle-

bound water. The water content of PM is significant and highly variable. Moreover, there is significant hysteresis in the water adsorption-desorption pathways (Seinfeld and Pandis, 1998), further complicating the mass measurement. Figures 2-8 and 2-9 show the change in diameter of sulfate particles as a function of relative humidity. Figure 2-8 also shows the difference between deliquescence and crystallization points.

Pilinis et al. (1989) calculated the water content of atmospheric particulate matter above and below the deliquescent point. They predicted that aerosol water content is strongly dependent on composition and concluded from their calculations that liquid water could represent a significant mass fraction of aerosol concentration at relative humidities above 60%. Since then, a few researchers have attempted to measure the water content of atmospheric aerosol. Most techniques have focused on tracking the particle mass as the relative humidity is changed and are still in the development phase. There have been only a few demonstrations using actual ambient aerosol to date. Of interest, in particular, is the development of the Tandem Differential Mobility Analyzer (TDMA) and its applications in investigations of the effects of relative humidity on particle growth.

Lee et al. (1997) examined the influence of relative humidity on the size of atmospheric aerosol using a TDMA coupled with a scanning mobility particle sizer (SMPS). They reported that the use of the TDMA/SMPS system allowed for the abrupt size changes of aerosols at the deliquescence point to be observed precisely. They also reported that at relative humidities between 81 and 89% the water content of ammonium sulfate aerosols (by mass) ranged from 47 to 66%.

Andrews and Larson (1993) investigated the interactions of single aerosol particles coated with an organic film within a humid environment. Using an electrodynamic balance, they conducted laboratory experiments in which sodium chloride and carbon black particles were coated with individual organic surfactants (intended to simulate the surface-active, organic films that many atmospheric aerosol particles may exhibit) and their water sorption curves were examined. Their results showed that when ordinarily hydrophobic carbon black particles were coated with an organic surfactant, they sorbed significant amounts of water (20 to 40% of the dry mass of the particle).

Liang and Chan (1997) developed a fast technique using the electrodynamic balance to measure the water activity of atmospheric aerosols. In their technique, the mass of a levitated

particle is determined as the particle either evaporates or grows in response to a step change in the relative humidity. Their technique was demonstrated using laboratory experiments with NaCl, $(NH_4)_2SO_4$, $NaNO_3$, and $(NH_4)_2SO_4/NH_4NO_3$ solutions. They concluded that one of the advantages of their fast method is the ability to measure the water activity of aerosols containing volatile solutes such as ammonium chloride and some organics.

McInnes et al. (1996) measured aerosol mass concentration, ionic composition, and associated water mass of marine aerosol over the remote Pacific Ocean. The mass of particle-bound water was determined by taking the difference between the mass obtained at 48% RH and at 19% RH, assuming the aerosol particles were dry at 19% RH. Based on a comparison of the remote Pacific aerosol to aerosol collected at a site at the marine/continental interface of the Washington coast, the amount of water associated with the aerosol was observed to be a function of the ammonium to sulfate ratio. They found that the amount of water associated with the submicrometer aerosol comprised 29% of the total aerosol mass collected at 47% RH and 9% of the total mass at 35% RH.

Ohta et al. (1998) characterized the chemical composition of atmospheric fine particles (50% cut point of 2 μm) in Sapporo, Japan, and as part of their measurements, determined the water content using the Karl Fischer method (Meyer and Boyd, 1959). After exposing a Teflon filter, a portion of the filter was equilibrated at 30% RH for 24 h. Then the filter piece was placed in a water evaporator heated at 150 °C, vaporizing the particle-bound water. The vapor evolved was analyzed for water in an aqua-counter where it was titrated coulometrically in Karl Fischer reagent solution (containing iodine, sulfur, and methanol). The accuracy of the aqua-counter is ±1 mg. Using this technique, they determined that the water content of the particles ranged from 0.4 to 3.2% of the total particulate mass (at RH < 30%). This represents a smaller portion of water compared to their previous reported values (Ohta and Okita, 1990) that were determined by calculation at RH of 50%.

Speer et al. (1997) developed an aerosol liquid water content analyzer (LWCA) in which aerosol samples are collected on PTFE filters and then placed in a closed chamber in which the relative humidity is closely controlled. The aerosol mass is monitored using a beta-gauge, first as the relative humidity is increased from low RH to high RH, and then as the RH is decreased again. They demonstrated the LWCA on laboratory-generated aerosol and on an ambient $PM_{2.5}$ sample collected in Research Triangle Park, NC. The ambient aerosol sample was also analyzed

for chemical constituents. It is interesting to note that, although their laboratory-generated $(NH_4)_2SO_4$ aerosol demonstrated a sharp deliquescent point, their atmospheric aerosol, which was essentially $(NH_4)_2SO_4$, did not show a sharp deliquescent point.

Hygroscopic properties of aerosols have been studied from the viewpoint of their ability to act as condensation nuclei. The hygroscopic properties of fresh and aged carbon and diesel soot particles were examined by Weingartner et al. (1997) who found that fresh, submicron-size particles tended to shrink with increasing relative humidity because of a structural change. Lammel and Novakov (1995) found, in laboratory studies, that the hygroscopicity of soot particles could be increased by chemical modification and that the cloud condensation nucleation characteristics of diesel soot were similar to those of wood smoke aerosol.

The results of several of the above studies in which aerosol water content as a function of relative humidity was determined are summarized in Figure 2-13. In this figure, the results of Lee et al. (1997), McInnes et al. (1996), and Ohta et al. (1998) are included. Relative humidity ranged from 9%, at which the aerosol water content was assumed to be zero (McInnes et al., 1996), to 89%, at which the aerosol water content was determined to be 66% by mass (Lee et al., 1997). Koutrakis et al. (1989) and Koutrakis and Kelly (1993) also have reported field measurements of the equilibrium size of atmospheric sulfate particles as a function of relative humidity and acidity.

The effects of relative humidity on particle growth were also examined in several studies. Fang et al. (1991) investigated the effects of flow-induced relative humidity changes on particle cut sizes for aqueous sulfuric acid particles in a multi-nozzle micro-orifice uniform deposit impactor (MOUDI). Laboratory experiments were conducted in which polydisperse sulfuric acid aerosols were generated and the RH was adjusted. The aerosols were analyzed by a differential mobility analyzer. Fang et al. (1991) observed that for inlet RH less than 80%, the cut sizes for the sulfuric acid aerosols were within 5% of that for nonhygroscopic particles except at the stage for which the cut size was 0.047 μm where the cut size was 10.7% larger than the nonhygroscopic particle cut size. They concluded that flow-induced RH changes would have only a modest effect on MOUDI cut sizes at RH < 80%.

Hitzenberger et al. (1997) collected atmospheric aerosol in the size range of 0.06 to 15 μm in Vienna, Austria, using a nine-stage cascade impactor and measured the humidity-dependent water uptake when the individual impaction foils were exposed to high RH. They observed



Figure 2-13. Aerosol water content expressed as a mass percentage, as a function of relative humidity.

particle growth with varying growth patterns. Calculated extinction coefficients and single scattering albedo increased with humidity.

Hygroscopic properties, along with mixing characteristics, of submicrometer particles sampled in Los Angeles, CA, during the summer of 1987 SCAQS study and at the Grand Canyon, AZ, during the 1990 Navajo Generating Station Visibility Study were reported by Zhang et al. (1993). They used a tandem differential mobility analyzer (TDMA; McMurry and Stolzenburg, 1989) to measure the hygroscopic properties for particles in the 0.05- to 0.5-μm range. In their experimental technique, monodisperse particles of a known size are selected from the atmospheric aerosol with the first DMA. Then, the relative humidity of the monodisperse aerosol is adjusted, and the new particle size distribution is measured with the second DMA. At both sites, they observed that monodisperse particles could be classified according to “more” hygroscopic and “less” hygroscopic. Aerosol behavior observed at the two sites differed markedly. Within the experimental uncertainty (±2%) the “less” hygroscopic particles sampled in Los Angeles did not grow when the RH was increased to 90%; whereas at the Grand Canyon,

the growth of the "less" hygroscopic particles varied from day to day, but ranged from near 0 to 40% when the RH was increased to 90%. The growth of the "more" hygroscopic particles in Los Angeles was dependent on particle size (15% at 0.05 μm to 60% at 0.5 μm); whereas at the Grand Canyon, the "more" hygroscopic particles grew by about 50% with the growth not varying significantly with particle size. By comparison of the TDMA data to impactor data, Zhang et al. (1993) surmised that the more hygroscopic particles contained more sulfates and nitrates while the less hygroscopic particles contained more carbon and crustal components.

Although most of the work to date on the hygroscopic properties of atmospheric aerosols has focused on the inorganic fraction, the determination of the contribution of particle-bound water to atmospheric particulate mass is greatly complicated by the presence of organics. The effect of RH on adsorption of semivolatile organic compounds is discussed elsewhere in this chapter. Saxena et al. (1995) observed that particulate organic compounds also can affect the hygroscopic behavior of atmospheric particles. They idealized the organic component of aerosol as containing a hydrophobic fraction (high-molecular weight alkanes, alkanoic acids, alkenoic acids, aldehydes, and ketones) and a hydrophilic fraction (e.g., lower molecular weight carboxylic acids, dicarboxylic acids, alcohols, aldehydes, etc.) that would be likely to absorb water. They then analyzed data from a tandem differential mobility analyzer in conjunction with particle composition observations from an urban site (Claremont, CA) and from a nonurban site (Grand Canyon) to test the hypothesis that, by adding particulate organics to an inorganic aerosol, the amount of water absorbed would be affected, and the effect could be positive or negative, depending on the nature of the organics added. They further presumed that the particulate organic matter in nonurban areas would be predominantly secondary and thus hydrophilic, compared to the urban aerosol that was presumed to be derived from primary emissions and thus hydrophobic in nature. Their observations were consistent with their hypothesis, in that at the Grand Canyon, the presence of organics tended to increase the water uptake by aerosols; whereas at the Los Angeles site, the presence of organics tended to decrease water uptake.

Peng and Chan (2001) also recently studied the hygroscopic properties of nine water soluble organic salts of atmospheric interest using an electrodynamic balance operated at 25°C. Salts studied included sodium formate, sodium acetate, sodium succinate, sodium pyruvate, sodium methanesulfonate, sodium oxalate, ammonium oxalate, sodium malonate, and sodium maleate. They observed that hygroscopic organic salts have a growth factor of 1.76-2.18 from

RH=10-90%, comparable to that of typical hygroscopic inorganic salts such as NaCl and $(NH_4)_2SO_4$.

Nonequilibrium issues may be important for the TDMA, as well as for other methods of measuring water content. Although approach to equilibrium when the RH is increased is expected to be rapid for pure salts, it may be much slower for aerosols containing a complex mix of components (Saxena et al., 1995). For example, if an aerosol contains an organic film or coating, that film may impede the transport of water across the particle surface, thus increasing the time required for equilibrium (Saxena et al., 1995). Insufficient time to achieve equilibrium in the TDMA could result in underestimation of the water content.

2.2.4 U. S. Environmental Protection Agency Monitoring Methods

2.2.4.1 The Federal Reference Methods for Measurement of Equilibrated Mass for PM_{10}, $PM_{2.5}$, and $PM_{10-2.5}$

In 1997, EPA promulgated new standards for $PM_{2.5}$ to address fine-fraction thoracic particles and retained with minor revisions the 1987 PM_{10} standards to continue to address coarse-fraction thoracic particles (Federal Register, 1997). In partial response to numerous challenges to these standards, the U.S. Court of Appeals for the District of Columbia Circuit in *American Trucking Association v. EPA*, 175 F. 3d 1027 (U.S. Court of Appeals, D.C. Cir. 1999) found "ample support" for regulating coarse-fraction particles but revoked the revised PM_{10} standards (leaving in effect the 1987 PM_{10} standards) on the basis of PM_{10} being a "poorly matched indicator for coarse particulate pollution" because PM_{10} includes fine particles. Consistent with this specific aspect of the Court's ruling, which EPA did not appeal, EPA is now considering use of $PM_{10-2.5}$ as the indicator for coarse-fraction thoracic particles, in conjunction with $PM_{2.5}$ standards that address fine-fraction thoracic particles. Thus, EPA is now developing a Federal Reference Method for the measurement of $PM_{10-2.5}$.

2.2.4.1.1 PM_{10}

The FRM specified for measuring PM_{10} (Code of Federal Regulations, 2001a,b) has been discussed in previous PM AQCD's and will only be mentioned briefly. The PM_{10} FRM defines performance specifications for samplers in which particles are inertially separated with a penetration efficiency of 50% at an aerodynamic diameter (D_a) of 10 ± 0.5 μm. The collection

efficiency increases to ≈100% for smaller particles and drops to ≈ 0% for larger particles. Particles are collected on filters and mass concentrations are determined gravimetrically. Instrument manufacturers are required to demonstrate through field tests a measurement precision for 24-h samples of ± 5 $\mu g/m^3$ for PM_{10} concentrations below 80 $\mu g/m^3$ and 7% above this value. A number of samplers have been designated as PM_{10} reference samplers. The TEOM and several beta gauge samplers with 1-h time resolution have been designated as automated equivalent methods (U.S. Environmental Protection Agency, 2001).

2.2.4.1.2 $PM_{2.5}$

As opposed to the performance-based FRM standard for PM_{10}, the FRM for $PM_{2.5}$ (Code of Federal Regulations, 2001a) specifies certain details of the sampler design, as well as of sample handling and analysis, whereas other aspects have performance specifications (Noble et al., 2001). The $PM_{2.5}$ FRM sampler consists of a PM_{10} inlet/impactor, a $PM_{2.5}$ impactor with an oil-soaked impaction substrate to remove particles larger than 2.5 μm D_a, and a 47-mm PTFE filter with a particle collection efficiency greater than 99.7%. The sample duration is 24 h, during which time the sample temperature is not to exceed ambient temperatures by more than 5 °C. A schematic diagram of the $PM_{2.5}$ FRM sample collection system is shown in Figure 2-14. After collection, samples are equilibrated for 24 h at temperatures in the range of 20 to 23 °C (± 2 °C) and at relative humidities in the range of 30 to 40% (± 5%). The equilibration tends to reduce particle-bound water and stabilizes the filter plus sample weight. Filters are weighed before and after sampling under the same temperature and relative humidity conditions. For sampling conducted at ambient relative humidity less than 30%, mass measurements at relative humidities down to 20% are permissible (Code of Federal Regulations, 2001a).

The PM_{10} inlet specified for the $PM_{2.5}$ FRM is modified from a previous low flow-rate PM_{10} inlet that was acceptable in both EPA-designated reference and equivalent PM_{10} methods. The modification corrects a flaw that was reported for the previous sampler, in that under some meteorological conditions, the inlet may allow precipitation to penetrate the inlet. The modification includes a larger drain hole, a one-piece top plate, and louvers. Tolocka et al. (2001a) evaluated the performance of this modified inlet in a series of wind tunnel experiments. The modified inlet was found to provide a size out comparable to the original inlet, for both $PM_{2.5}$ and PM_{10} sampling. Since the modification did not change the characteristics of the size



Figure 2-14. Schematic diagram of the sample collection portion of the $PM_{2.5}$ FRM sampler.

Source: Noble et al. (2001).

cut, the modified inlet may be substituted for the original inlet as part of a reference or equivalent method for PM_{10} and $PM_{2.5}$ (Tolocka et al., 2001a).

WINS Impactor. Design and calibration of the EPA $PM_{2.5}$ Well Impactor Ninety-Six (WINS) is given by Peters et al. (2001a). The WINS impactor was designed to be deployed downstream of the Graseby-Andersen 246B PM_{10} inlet as part of a sampler operating at a flow rate of 16.7 L/m. The WINS is pictured in Figure 2-15. The $PM_{2.5}$ inlet consists of a single jet, round hole, with the jet exit directed toward an impaction surface that is comprised of a 37 mm diameter glass fiber filter immersed in 1 mL of low volatility diffusion pump oil (i.e., the well). Particles not having enough inertia to be removed by the impactor are captured downstream on the sample collection filter. This design was selected to minimize impactor overloading that



Figure 2-15. Schematic view of the final design of the WINS.

Source: Peters et al. (2001a).

would otherwise result in particle bounce. The oil wicks through the particulate deposit on the impactor to provide a continuously wetted surface for impaction. The penetration curve indicated a 50% cutpoint of 2.48 μm D_a with a geometric standard deviation of 1.18% (Figure 2-16).

The WINS separator was evaluated for its loading characteristics (Vanderpool et al., 2001) by monitoring the performance after repeated operation in an artificially generated, high concentration, coarse-mode aerosol composed of Arizona Test Dust, as well as in the field in Rubidoux, Phoenix, Philadelphia, Research Triangle Park, and Atlanta. In the wind tunnel experiments, the WINS performance was found to be a monotonic function of loading. A minus 5% bias in the $PM_{2.5}$ measurement resulted from a coarse particulate loading of approximately 16 mg. This negative bias was due to a slight reduction in the separator cutpoint. It was also



Figure 2-16. Evaluation of the final version of the WINS.

Source: Peters et al. (2001a).

found that the predictable results from the controlled laboratory experiments could not be extrapolated to field settings and that the WINS performance was more sensitive to the impactor loading in the field than it was in experiments with the single component aerosol. Significant particle bounce was not observed in either the laboratory or the field experiments. Vanderpool et al. (2001) conclude that their study supports the recommendation that the FRM WINS wells should be replaced after every 5 days of 24-h operation (U.S. Environmental Protection Agency, 1998).

A detailed sensitivity study of the WINS impactor was conducted (Vanderpool et al., 2001) in which the effects on the impactor performance of a number of parameters were examined. The results of this study are summarized in Table 2-3.

TABLE 2-3. SUMMARY OF SENSITIVITY STUDIES OF WINS IMPACTOR PERFORMANCE

Parameter	Amount of variance	Cutpoint variation	$PM_{2.5}$ mass concentration bias
Manufacturing tolerances on WINS components	Specified tolerances	0.05 micrometers	<1%
Flow control biases	4%	0.05 micrometers	Cutpoint shift partially offset volume bias
T and P measurement	Allowable ambient	± 0.02 micrometers	± 0.4%
Diffusion oil volume	0.75 ml to 3 ml	No effect	
Impactor loading	After 5 24 h events	−0.07 micrometers	<1.5%
Ambient P variations		Negligible	Negligible
Air Properties	0 C	2.40 micrometers	NA
Impactor oil crystallization		No effect	No effect
Impactor oil viscosity	−20 C	No effect	No effect
	−35 C	Need to change WINS more frequently than every 5 days	

Source: Vanderpool et al. (2001).

The regulations also allow for Class I, II, and III equivalent methods for $PM_{2.5}$ (Code of Federal Regulations, 2001c). Class I equivalent methods use samplers with relatively small deviations from the sampler described in the FRM. Class II equivalent methods include "all other $PM_{2.5}$ methods that are based upon 24-h integrated filter samplers that are subjected to subsequent moisture equilibration and gravimetric mass analysis." Class III equivalent methods include non-filter-based methods such as beta attenuation, harmonic oscillating elements, or nephelometry (McMurry, 2000). As of July 2001, 11 $PM_{2.5}$ samplers (listed in Table 2-4) had been tested and led to the conclusion that the PM_{10} sampling systems can be designed such that concentration measurements are precise to ±10%. For $PM_{2.5}$, cutpoint tolerances are not expected to affect the mass concentration as much as for PM_{10}, because the 2.5 μm D_a cutpoint generally occurs near a minimum in the mass distribution (e.g., Figure 2-5).

The $PM_{2.5}$ mass concentration will be affected, on the other hand, by other sampling issues mentioned but not discussed extensively in the previous 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). These issues have been discussed earlier in this chapter and include gas/particle, particle/particle, and particle/substrate interactions for sulfates and nitrates (e.g.,

TABLE 2-4. $PM_{2.5}$ SAMPLERS CURRENTLY DESIGNATED AS FRMs FOR $PM_{2.5}$ MASS CONCENTRATIONS

Sampler	Manufacturer	Description	Federal Register Reference
RAAS2.5-100	Andersen Instruments	FRM single	Vol. 63, p. 31991, 6/11/98
RAAS2.5-300	Andersen Instruments	FRM sequential	Vol. 63, p. 31991, 6/11/98
RAAS2.5-200	Andersen Instruments	FRM audit	Vol. 64, p. 12167, 3/11/99
Partisol 2000	Rupprecht & Patashnick	FRM single	Vol. 63, p. 18911, 4/16/98
Partisol-Plus 2025	Rupprecht & Patashnick	FRM sequential	Vol. 63, p. 18911, 4/16/98
Partisol 2000 audit	Rupprecht & Patashnick	FRM audit	Vol. 64, p. 19153, 4/19/99
PQ 200	BGI, Inc.	FRM single	Vol. 63, p. 18911, 4/16/98
PQ 200A	BGI, Inc.	FRM audit	Vol. 63, p. 18911, 4/16/98
605 CAPS	ThermoEnvironmental Instruments	FRM single	Vol. 63, p. 58036, 10/29/98
MASS 100	URC	FRM single	Vol. 65, p. 26603, 05/08/00
MASS 300	URC	FRM sequential	Vol. 65, p. 26603, 05/08/00

Source: Peters et al. (2001b); U.S. Environmental Protection Agency (2001).

Appel et al., 1984), volatilization losses of nitrates (Zhang and McMurry, 1992), semivolatile organic compound artifacts (e.g., Eatough et al., 1993), and relative humidity effects (e.g., Keeler et al., 1988).

Several studies now have been reported in which the FRM was collocated with other $PM_{2.5}$ samplers in intercomparison studies. During the Aerosol Research and Inhalation Epidemiology Study (ARIES), several $PM_{2.5}$ samplers were collocated at a mixed industrial-residential site near Atlanta, GA (Van Loy et al., 2000). These samplers included a standard $PM_{2.5}$ FRM, a TEOM with Nafion drier, a particulate composition monitor (PCM; Atmospheric Research and Analysis, Cary, NC), a medium-volume (113 L/min flow rate) fine particle ($PM_{2.5}$) and semivolatile organic compound (i.e., a filter followed by a solid adsorbent) sampler, operated by the Desert Research Institute, a HEADS sampler, and a dichotomous sampler for coarse PM. The PCM sampler has three channels, all of which have PM_{10} cyclone inlets. The first two channels both have two denuders preceding a 2.5-μm WINS impact and filter packs. The first denuder is coated with sodium carbonate to remove acid gases, and the second is coated with citric acid to

remove ammonia. The third channel has a carbon coated parallel-plate denuder preceding the WINS impactor. Measurements of 24-h mass from the FRM, PCM, and TEOM samplers, as well as reconstructed $PM_{2.5}$ mass (RPM), were compared for a 12-mo period. The slopes for the TEOM-FRM, PCM-FRM, and RPM-FRM correlations were 1.01, 0.94, and 0.91, respectively; whereas the y-intercepts for each were 0.68, 0.04, and 0.98. Particulate sulfate measurements on the FRM Teflon filter, the PCM Teflon filter, and PCM Nylon filter were nearly identical. Nitrate results from the three filters were much less consistent, with the FRM collecting substantially less nitrate than that collected on either the denuded nylon filter or a denuder followed by a Teflon-nylon filter sandwich. Particulate ammonia measurements were also compared and showed more scatter than the sulfate measurements but less than the nitrate measurements.

An intercomparison of both PM_{10} and $PM_{2.5}$ mass measurements was conducted during the 1998 Baltimore PM Study (Williams et al., 2000). PM monitors were collocated at a residential indoor, residential outdoor, and ambient monitoring site within Baltimore County, MD. PM samplers included TEOMs, $PM_{2.5}$ FRMs, cyclone-based inlets manufactured by University Research Glassware (URG), and Versatile Air Pollution Samplers (VAPS). The VAPS sampler is a dichotomous sampler operating at 33 L/min (one coarse particle channel at 3 L/min, and two fine particle channels at 15 L/min, each). In the configuration employed during this study, one fine particle channel was operated with a Teflon filter backed by a nylon filter and preceded by a sodium carbonate coated annular denuder; the second fine particle channel had a quartz filter preceded by a citric acid-coated annular denuder; and the coarse particle channel had a polycarbonate filter followed by a Zefluor filter for flow distribution. Differences in $PM_{2.5}$ mass concentrations between the samplers, although not large, were attributed to potential particle nitrate losses, denuder losses, and losses of SVOC for some samplers. Differences between coarse particulate mass concentrations, on the other hand, varied widely between the instruments.

In another intercomparison study, Tolocka et al. (2001b) examined the magnitude of potential sampling artifacts associated with the use of the FRM by collocating FRMs alongside other chemical speciation samplers at four U.S. cities. The locations included a high nitrate and carbon, low sulfate site (Rubidoux, CA); high crustal, moderate carbon and nitrate site (Phoenix); high sulfate, moderate carbon, and low nitrate (Philadelphia); and low $PM_{2.5}$ mass (Research Triangle Park, NC). The use of Teflon and heat-treated quartz filters was also

examined in this study. The Teflon filters collected less nitrate than the heat-treated quartz filters. Filters in samplers using denuders to remove organic gases collected less organic PM than filters in samplers without denuders.

Peters et al. (2001b) compiled the results of several field studies in which a number of FRM and other $PM_{2.5}$ samplers were intercompared. In addition to the FRM samplers listed in Table 2-3, other $PM_{2.5}$ samplers included the Sierra Instruments dichotomous sampler, the Harvard impactor, the IMPROVE sampler, and the Air Metrics saturation monitor. Results were compiled from $PM_{2.5}$ field studies conducted in Birmingham, Denver, Bakersfield, Phoenix, Research Triangle Park, Atlanta, and Rubidoux. Limited studies on precision for the non-FRM samplers suggest that the Harvard Impactor and dichotomous samplers had the lowest coefficient of variations (CV), with both under 10%. The CV for this study was calculated by dividing the sample standard deviation by the average concentration. The IMPROVE samplers had CVs between 10 and 12%, and the Air Metrics samplers had the highest observed CVs, over 15%. In intercomparisons with FRM samplers, the non-FRM samplers showed strong linear relationships in comparison to the FRM sampler; however, none of the comparisons passed the current EPA Subpart C equivalent method criteria, which EPA is in the process of revising.

Detailed information on precision of PM samplers used in monitoring networks may be found in EPA's Technology Transfer Network website (U.S. Environmental Protection Agency, 2002).

2.2.4.1.3 $PM_{10-2.5}$

Measurement techniques for $PM_{10-2.5}$ are somewhat more complex than those for $PM_{2.5}$ or PM_{10} because, for $PM_{10-2.5}$, it is necessary to isolate a size fraction between an upper 50% cut point of 10 μm D_a and a lower 50% cut point of 2.5 μm D_a.

The Difference Method. One approach to measurement of $PM_{10-2.5}$ is to make separate measurements of PM_{10} and $PM_{2.5}$ and take the difference of the resulting equilibrated masses. One problem is that, if either the $PM_{2.5}$ or the PM_{10} sampler fails, no $PM_{10-2.5}$ measurement can be obtained. In addition, errors in cut-point, flow rate, and filter weights (both before use and after collection and equilibration of particles) and uncertainties due to loss of semivolatile components of PM may occur for each size cut. Careful control of flow rate and equivalent treatment of PM_{10}

and $PM_{2.5}$ filters in terms of pressure drop across the filter and temperature of the filter during and after collection can improve precision and accuracy. Allen et al. (1999b) summarized several sampling issues to consider in measuring coarse particulate mass by difference, including the use of identical instrumentation (except cutpoints), filter media, filter face velocity, and ambient-filter temperature differences; common flow measurement devices; use of higher sampler flow rates (10 L/min minimum for 24-h sample is recommended); and avoiding excessive filter loading. The concern, expressed by Allen et al. (1999b), that the "pie-plate" inlet required by the final version of the $PM_{2.5}$ FRM might have a different cut point than the "flat-top" inlet of the PM_{10} FRM, has been addressed by a wind tunnel study which found both to have an appropriate PM_{10} cut point (Tolocka et al., 2001a).

Since the difference method requires weighing two filters, the key to obtaining high precision in the coarse mass measurement is precise measurements of filter weights. Allen et al. (1999b) discuss techniques for increasing the precision of the difference method by reducing errors in filter weights. These include proper temperature and humidity controls, use of a high quality microbalance, 100% replicate weighings, control of static charge, aging of new filters, weighing of a sufficient number of laboratory blank filters, and accounting for buoyancy errors caused by variability in barometric pressure. Allen et al. (1999b) emphasize the necessity of replicate weighing of filters and a third weighing if the difference between the first two weights exceeds a specified minimum. Lawless et al. (2001) investigated the magnitude of uncertainties attributed to fluctuations in some of these parameters (humidity, temperature, drafts, vibration, and electrostatic charges) and recommended methods for improving their control. Koistinen et al. (1999) and Hänninen et al. (2002) give a excellent discussion of the procedures developed to overcome problems associated with gravimetric measurements of $PM_{2.5}$ mass in the EXPOLIS (The Air Pollution Exposure Distributions Within Adult Urban Populations in Europe) Study. They discuss factors such as corrections for buoyancy, elimination of static charge, and increases in the mass of blank filters with time. The establishment of a temperature and humidity controlled room required for the equilibration and weighing of filters for the FRM is expensive. Allen et al. (2001) describe a more cost-effective technique that uses a chamber with relative humidity controlled at 34% relative humidity by a saturated aqueous solution of $MgCl_2$.

Allen et al. (1999b) recommend that, in reporting precision from collocated samplers both the (CV) and the square of the correlation coefficient (r^2) be reported. For a study in Boston with

27 pairs of mass data from collocated PM_{10} and $PM_{2.5}$ using standard weighing methods, they obtained a CV of 4.7% and an r^2 of 0.991 for $PM_{2.5}$, a CV of 4.4% and an r^2 of 0.994 for PM_{10}, and a CV of 15% and an r^2 of 0.88 for $PM_{10-2.5}$. By using duplicate weighings and other techniques suggested for improving precision, they obtained a CV of 1.3% and an r^2 of 0.998 for $PM_{2.5}$ in a study in Chicago with 38 collocated measurements. On the basis of the improvement in the CV for $PM_{2.5}$, they estimate that use of the recommended techniques for $PM_{10-2.5}$ by difference would have yielded a CV of 3.8% and an r^2 of 0.98 if they had been applied in the Chicago study.

This "difference" technique has been used to measure $PM_{10-2.5}$ in a number of studies. Currently, estimates of $PM_{10-2.5}$ are obtained by subtracting $PM_{2.5}$ from PM_{10} (both measured by FRM monitors). EPA is currently in the process of developing an FRM for $PM_{10-2.5}$.

Multistage Impaction. A second technique involves the use of impaction to isolate the size fraction between 2.5 and 10 μm D_a. In the impaction process, the air stream is first accelerated through a small hole (nozzle) or slit. The air stream is directed so that it "impacts" on a surface. Depending on the velocity and pressure of the air stream, particles smaller than a certain size will follow the air stream around the impactor surface. Larger particles will impact on the surface. In practice, impactors have 50% cut points similar to those for the rejection of larger particles in $PM_{2.5}$ and PM_{10} samples (Figure 2-6).

Multistage impactors are used to separate particles into several size fractions for the determination of mass and chemical composition as a function of size (Wang and John, 1988; Marple et al., 1991). The major problem with the use of impactors to separate the 10-2.5 μm D_a fraction of coarse particles (thoracic coarse PM) is bounce. Coarse particles tend to be dry, solid particles. When they hit a hard surface, they can bounce and be carried away with the air stream (e.g., Dzubay et al., 1976; Wesolowski et al., 1977; Rao and Whitby, 1978; Cheng and Yeh, 1979; Wang and John, 1987; John and Sethi, 1993). Various techniques have been used to reduce bounce. One technique is to use a porous substance such as a glass or quartz fiber filter (Chang et al., 1999) material or a polyurethane foam (Breum, 2000; Kavouras and Koutrakis, 2001). These techniques may result in less precise separation and yield a sample that must be extracted before chemical analyses can be performed. Another technique is to coat the impactor with a soft wax or grease (Rao and Whitby, 1977; Turner and Hering, 1987; Pak et al., 1992).

This can cause problems with weighing and chemical analyses. In addition, as the deposit of particles builds up, incoming particles may not hit the soft surface, but instead hit a previously collected hard particle and bounce off of it. The WINS impactor discussed earlier uses a filter in a well of low volatility oil to ensure a wetted surface at all times. However, such a technique, while appropriate for removing unwanted particles, would not yield a particle sample suitable for weighing or for chemical analyses.

Virtual Impaction. In an effort to alleviate the bounce problem, aerosol scientists have developed the "virtual" impactor (Loo et al., 1976; Jaklevic et al., 1977; Loo and Cork, 1988). A hole is placed in the impaction plate just below the accelerating jet. Two controlled flows allow a fraction, e.g., 10% (or another predetermined fraction, typically 5 - 20%), of the air to go through the hole and through a filter (minor flow). A 10% minor flow gives a coarse channel enrichment factor of 10. The remaining fraction (e.g., 90% of the airflow) follows a different path and goes through a second filter (major flow). The upper cutpoint is usually set by the inlet (e.g., 10 μm D_a). The flow rates, pressures, and distance from the nozzle to the virtual impactor surface can be varied to direct particles with an D_a greater than the lower cutpoint (i.e., > 2.5 μm) to go through the hole and be collected on the first filter and to direct smaller particles (i.e., < 2.5 μm) to flow around the impactor be collected on the second filter (Marple and Chien, 1980). This technique overcomes the problem of bounce. However, a fraction of the smaller particles, equal to the minor flow, will go through the virtual impaction opening with the air stream and be collected on the course particle filter. Thus, in order to determine the mass or composition of the coarse particles, it is necessary to determine the mass and composition of the fine particles and subtract the appropriate fraction from the mass or composition of the particles collected on the coarse particle filter. Virtual impactors that separate particles into two size fractions are known as dichotomous samples. Allen et al. (1999b) discuss potential errors in the dichotomous sampler caused by uncertainties in the coarse mass channel enrichment factor. An example of the separation into fine and coarse particles is shown in Figure 2-17.

The dichotomous sampler was developed for use in the Regional Air Monitoring Study (RAMS), part of the Regional Air Pollution Study (RAPS), conducted in St. Louis, Missouri in the mid-1970s (Loo et al., 1976). Dichotomous samplers were a new concept at that time, and there was concern that particle loses might be high at cut point sizes below 2.5 μm D_a.



Figure 2-17. Schematic diagram showing the principle of virtual impaction. The initial flow, Q_0, is split into a minor flow, Q_1, which carries the larger particles, which impact into the hole, to the coarse particle filter and a major flow, Q_2, which carries the smaller particles, which can follow the airflow, to the fine particle filter (Loo et al., 1976).

In subsequent years, the theory of virtual impaction has advanced. Now virtual impactors, with rectangular slits or round holes, are used to give cut point sizes as low as 0.15 μm D_a and are used to concentrate coarse, accumulation, and ultrafine mode particles for use in health studies (Solomon et al., 1983; Marple et al., 1990; Sioutas et al., 1994b,c,d). Dichotomous samplers were also used in a national network to measure $PM_{2.5}$ and $PM_{10-2.5}$ in the Harvard Six City Study (Dockery et al., 1993) and the Inhalable Particulate Network (Suggs and Burton, 1983).

2.2.5 Speciation Monitoring

Speciation Network and Monitoring

In addition to FRM sampling to determine compliance with PM standards, EPA requires states to conduct chemical speciation sampling primarily to determine source categories and trends (Code of Federal Regulations, 2001b). Source category apportionment calculations are discussed in Chapter 3. A $PM_{2.5}$ chemical speciation network has been deployed that consists of 54 core National Ambient Monitoring Stations (NAMS) and approximately 250 State and Local Air Monitoring Stations (SLAMS). In addition, over 100 IMPROVE (Interagency Monitoring of Protected Visual Environments) samplers located at regional background and transport sites can be used to fulfill SLAMS requirements. The overall goal of the speciation program is "to provide ambient data that support the Nation's air quality program objectives" (U.S. Environmental Protection Agency, 1999). Information and reports on EPA's speciation monitoring program may be found on EPA's Technology Transfer Network at http://www.epa.gov/ttn/amtic/pmspec.html. The NAMS speciation sites will provide routine chemical speciation data that will be used to develop annual and seasonal aerosol characterization, air quality trends analysis, and emission control strategies. The SLAMS speciation sites will further support the NAMS network and provide information for development of State Implementation Plans (SIPs).

At both NAMs and SLAMs sites, aerosol samples will be collected for analysis of trace elements, ions (sulfate, nitrate, ammonium, sodium, and potassium), and total carbon. The NAMS speciation sites will operate on a 1 in 3 day schedule, with 10 of these sites augmented with continuous speciation analyses for everyday operation. The SLAMS speciation sites will generally operate on a 1 in 6 day basis; however, many sites may be operated on a 1 in 3 day basis in locations where increased data collection is needed. The current samplers include three filters: (1) Teflon for equilibrated mass and elemental analysis by energy dispersive x-ray fluorescence (EDXRF), (2) a nitric acid denuded Nylon filter for ion analysis (ion chromatography), (3) a quartz fiber filter for elemental and organic carbon (but currently without any correction for positive or negative artifacts caused by adsorption of organic gases or the quartz filters or evaporation of semivolatile organic compounds from the collected particles); and (4) thermal optical analysis via NIOSH (National Institute for Occupational Safety and Health) method 5040 (Thermal Optical Transmission) [TOT]). There are several samplers that are

suitable for use in the NAMS/SLAMS network. These samples include an inlet cutpoint comparable to the WINS FRM; proven denuder technology for ions; and sampler face velocity and sample volume similar to that of the FRM with 46.2-mm diameter filters.

Since 1987, the IMPROVE network has provided measurements of ambient PM and associated light extinction in order to quantify PM chemical components that affect visibility at Federal Class 1 areas that include designated national parks, national monuments, and wilderness areas. Management of this network is a cooperative effort between U.S. EPA, federal land management agencies, and state governments. The IMPROVE program has established protocols for analysis of aerosol measurements that provide ambient concentrations for PM_{10}, $PM_{2.5}$, sulfates, nitrates, organic and elemental carbon, crustal material, and a number of other elements. Information on the IMPROVE program may be found at http://vista.cira.colostate.edu/improve.

IMPROVE aerosol monitoring consists of a combination of particle sampling and sample analysis. The IMPROVE sampler, which collects two 24-hour duration samples per week, simultaneously collects one sample of PM_{10} on a Teflon filter, and three samples of $PM_{2.5}$ on Teflon, nylon, and quartz filters. PM_{10} mass concentrations are determined gravimetrically from the PM_{10} filter sample, while $PM_{2.5}$ mass concentrations are determined gravimetrically from the $PM_{2.5}$ Teflon filter sample. The $PM_{2.5}$ Teflon filter sample is also used to determine concentrations of selected elements using particle-induced x-ray emission (PIXE), x-ray fluorescence (XRF), and Proton Elastic Scattering Analysis (PESA). The $PM_{2.5}$ nylon filter sample, which is preceded by a denuder to remove acidic gases, is analyzed to determine nitrate and sulfate aerosol concentrations using Ion Chromatography (IC). Finally, the $PM_{2.5}$ quartz filter sample is analyzed for organic and elemental carbon using the Thermal Optical Reflectance (TOR) method.

Several of the $PM_{2.5}$ size selectors developed for use in the EPA National $PM_{2.5}$ Chemical Speciation Trends network were recently evaluated by comparing their penetration curves under clean room experiments with that of the WINS impactor (Peters et al., 2001c). The corresponding speciation monitors were then compared to the FRM in four cities. The $PM_{2.5}$ inlets tested were the SCC 2.141 cyclone (6.7 L/min) that is in the Met One Instruments SASS sampler; the SCC 1.829 cyclone (5.0 L/min) that is proposed for use in the Rupprecht and Patashnik real-time sulfate/nitrate monitor; the AN 3.68 cyclone (24.0 L/min) that is in the

Andersen RAAS; and the spiral separator (7.0 Lpm) that was previously in the Met One SASS. The cutpoints of the SCC cyclones compared reasonably well with the WINS (2.52 and 2.44 micrometers for the SCC 2.141 and SCC 1.829, respectively, at their design flowrates), but both demonstrated a tail extending into the coarse particle mode. The AN inlet had the sharpest cutpoint curve, but the 50% cutpoint diameter was 2.7 μm D_a at its design flowrate. The spiral inlet had the shallowest cutpoint curve, and the 50% cut point was 2.69 and 2.67 μm D_a for an ungreased and greased inlet, respectively. The speciation samplers were also compared to the FRM sampler with WINS inlet under ambient conditions in four cities. The Andersen RAAS equipped with the AN 3.68 cyclone compared well to the FRM in all four cities, when compared on the basis of $PM_{2.5}$ mass, sulfate, and crustal concentrations. Greasing the spiral inlet in the Met One sampler improved the performance of that sampler, which tended to give much higher $PM_{2.5}$ concentrations than the FRM in cities with high crustal particulate matter.

2.2.6 Inorganic Elemental Analyses

In addition to the lighter elements, hydrogen, carbon, oxygen and nitrogen, the following 40 heavier elements are commonly found in ambient air samples: sodium, magnesium, aluminum, silicon, phosphorus, sulfur, chlorine, potassium, calcium, titanium, vanadium, chromium, manganese, iron, cobalt, nickel, copper, zinc, gallium, arsenic, selenium, bromine, rubidium, strontium, yttrium, zirconium, molybdenum, palladium, silver, cadmium, indium, tin, antimony, barium, lanthanum, gold, mercury, thallium, lead, and uranium. These often indicate air pollution sources and several of them are considered to be toxic (transition metals, water-soluble metals, and metals in certain valence states [e.g., Fe(II), Fe(III), Cr(III), Cr(VI), As(III), As(V)]). Measurement methods for the heavier elements include: (1) energy dispersive x-ray fluorescence (EDXRF); (2) synchrotron induced X-ray emission (S-XRF); (3) proton induced x-ray emission (PIXE); (4) proton elastic scattering analysis (PESA); (5) total reflection X-ray fluorescence (TRXRF); (6) instrumental neutron activation analysis (INAA); (7) atomic absorption spectrophotometry (AAS); (8) inductively coupled plasma with atomic emission spectroscopy (ICP-AES); (9) inductively coupled plasma with mass spectroscopy (ICP-MS); and (10) scanning electron microscopy (SEM). These methods differ with respect to detection limits, sample preparation, and cost (Chow, 1995). XRF and PIXE are the most commonly applied methods because they quantify more than 40 detectable elements, they are non-destructive, and

they are relatively inexpensive. Both were discussed in the previous 1996 PM AQCD. TRXRF and S-XRF are newer techniques with lower detection limits. AAS, ICP-AES, and ICP-MS are also appropriate for ionic measurements when the particles are extracted in deionized distilled water. PESA provides a means for measuring elements with lower atomic numbers from hydrogen to carbon.

2.2.6.1 Energy Dispersive X-ray Fluorescence (EDXRF)

EDXRF has usually been the method of choice for analysis of trace elements on filters. EDXRF is preferred for aerosol analysis over wavelength dispersive XRF because it allows fast and simultaneous analysis over the total spectrum, allowing for the analysis of numerous elements simultaneously. EDXRF can accommodate small sample sizes and requires little sample preparation or operator time after the samples are placed into the analyzer. It also leaves the sample intact after analysis; so, further analysis is possible. XRF irradiates a uniform particle deposit on the surface of a membrane filter with 1 to 50 kev x-rays that eject inner shell electrons from the atoms of each element in the sample (Dzubay and Stevens, 1975; Jaklevic et al., 1977; Billiet et al., 1980; Potts and Webb, 1992; Piorek, 1994; Bacon et al., 1995; deBoer et al., 1995; Holynska et al., 1997; Török et al., 1998; Watson et al., 1999). When a higher energy electron drops into the vacant lower energy orbital, a fluorescent x-ray photon is released. The energy of this photon is unique to each element, and the number of photons is proportional to the concentration of the element. Concentrations are quantified by comparing photon counts for a sample with those obtained from thin-film standards of known concentration (Dane et al., 1996). The previous 1996 PM AQCD included a detailed discussion of EDXRF.

Emitted x-rays with energies less than ~4 kev (affecting the elements sodium, magnesium, aluminum, silicon, phosphorus, sulfur, chlorine, and potassium) are absorbed in the filter, in a thick particle deposit, or even by large particles in which these elements are contained. Very thick filters also scatter much of the excitation radiation or protons, thereby lowering the signal-to-noise ratio for XRF and PIXE. For this reason, thin membrane filters with deposits in the range of 10 to 50 $\mu g/cm^2$ provide the best accuracy and precision for XRF and PIXE analysis (Davis et al., 1977; Haupt et al., 1995).

2.2.6.2 Synchrotron Induced X-ray Fluorescence (S-XRF)

S-XRF is a form of EDXRF in which the exciting x-rays are derived from a synchrotron. Bremmstrahlung x-rays are generated when energetic electrons (generally several GeV in energy) are forced by a magnetic field to make a bend in their path. The advantages of the technique are that an extremely high flux of x-rays may be obtained and that the x-rays are 100% polarized in the plane of the electron beam. The former allows for x-ray beams generally of 50 to 500 μm in diameter. However, the beams can be focused into x-ray microprobes, with spot sizes on the order of one μm diameter. The x-ray polarization allows for removal of most of the background normally found under the characteristic x-ray peaks, greatly improving sensitivity compared to other XRF techniques. The primary disadvantages are the limited number of synchrotrons, and that few synchrotrons have S-XRF capabilities. Thus, the technique has been relatively little used for PM, and then generally for special problems such as the smoke from the Kuwaiti oil fires (Cahill et al., 1992, Reid et al., 1994). However, with the increasing availability of S-XRF facilities dedicated to PM analysis, the first of which was the Advanced Light Source opened at Lawrence Berkeley National Laboratory last year, utilization of S-XRF for PM analysis is increasing.

2.2.6.3 Proton Induced X-ray Emission (PIXE)

PIXE differs from XRF analysis in the excitation source for producing fluorescence. The filter deposit is bombarded with high-energy protons to remove inner shell electrons and the resulting characteristic x-rays are analyzed as in XRF (Johansson, 1970, Cahill, 1981, 1985; Zeng et al., 1993). Small accelerators, generally Van de Graaffs, generate intense beams of low energy protons, generally of a few MeV in energy. These have the ability to remove electrons from inner shells of atoms of any element. Thus, PIXE can see a very wide range of elements in a single analysis. The cross section for producing x-rays using protons of a few MeV in energy tends to favor lighter elements, Na through Ca, but sensitivities for equivalent PIXE and multi-wavelength XRF analysis are roughly comparable. The technique has been widely used in the U.S. (Flocchini et al., 1976, Malm et al., 1994) and around the world, as many universities have the small accelerators needed for the method. Like S-XRF, the proton beams can be focused into μm size beams, but these have been relatively little used for PM. However, the mm

size beams used in both S-XRF and PIXE are well suited to analyzing the limited mass and small deposits that result from detailed particle size measurements by impactors. (Perry et al., 1999)

2.2.6.4 Proton Elastic Scattering Analysis (PESA)

With the routine availability of elemental analyses for all elements sodium and heavier, organic components remain the major unmeasured species for mass balance. For programs like IMPROVE (Malm et al., 1994), parallel filters are collected for separate organic and elemental carbon determinations. Aerosol programs that use PIXE can directly measure hydrogen simultaneously by scattering protons from Teflon filters that lack hydrogen (Cahill et al., 1989, 1992). Generally, results from organic matter by carbon combustion from quartz filters and organic matter by hydrogen from Teflon filters are in agreement, assuming certain assumptions about the chemical states of sulfates and nitrates are met (Malm et al., 1994, Cahill et al., 1996).

2.2.6.5 Total Reflection X-ray Fluorescence (TRXRF)

One of the limitations of the EDXRF method is the minimum detection limit, which may be high due to high background values (Streit et al., 2000). By implementation of x-ray optical geometries that use the total reflection of the primary radiation on flat surfaces, scattering on the substrate is reduced, so that detection limits can be reduced. This is the basis for the total reflection x-ray fluorescence (TRXRF) method (Aiginger and Streli, 1997). This modification to the EDXRF technique improves detection limits and avoids the need to correct for matrix effects. Despite its apparent advantages, TRXRF has not yet become widely in use for atmospheric aerosol analysis but has been used in the analysis of marine aerosol (Stahlschmidt et al., 1997) and at a high elevation site (Streit et al., 2000). Streit et al. sampled ambient air at the High Alpine Research Station (3580 m above sealevel) in the Bernese Alps, Switzerland, using a nine-stage, single-jet, low-pressure, cascade impactor equipped with quartz impactor plates coated with silicon oil diluted in 2-propanol. The typical sample volume for a weekly sample was 10 m^3. The quartz plates were analyzed directly by TRXRF. Streit et al. reported that the minimum detection limits, defined by the 3σ values of the blanks, ranged from 25 ng for S, decreased monotonically with increasing atomic number down to 5 pg for Rb, and decreased after that. The use of TRXRF is expected to increase as EDXRF users become aware of the

method. A relatively low-cost, add-on unit has been developed that would allow EDXRF users to test the TRXRF technique (Aiginger, 1991).

2.2.6.6 Instrumental Neutron Activation Analysis (INAA)

INAA irradiates a sample in the core of a nuclear reactor for few minutes to several hours, depending on the elements being quantified (Dams et al., 1970; Zoller and Gordon, 1970; Nadkarni, 1975; Landsberger, 1988; Olmez, 1989; Ondov and Divita, 1993). The neutron bombardment chemically transforms many elements into radioactive isotopes. The energies of the gamma rays emitted by these isotopes identify them and, therefore, their parent elements. The intensity of these gamma rays is proportional to the amount of the parent element present in the sample. Different irradiation times and cooling periods are used before counting with a germanium detector. In source apportionment studies, it is possible to use a combination of XRF and INAA to develop a relatively complete set of elemental measurements. Between these two analytical techniques, good sensitivity is possible for many elements, including most of the toxic metals of interest. In general, XRF provides better sensitivity for some metals (e.g., Ni, Pb, Cu, and Fe); whereas INAA provides better sensitivity for others (Sb, As, Cr, Co, Se, and Cd). Both methods provide similar detection limits for still other elements (V, Zn, and Mn). INAA does not quantify some of the abundant species in ambient particulate matter such as silicon, nickel, tin, and lead. While INAA is technically nondestructive, sample preparation involves folding the sample tightly and sealing it in plastic, and the irradiation process makes the filter membrane brittle and radioactive. These factors limit the use of the sample for subsequent analyses.

INAA has been used to examine the chemical composition of atmospheric aerosols in several studies either as the only method of analysis or in addition to XRF (e.g., Yatin et al., 1994; Gallorini, 1995). INAA has higher sensitivity for many trace species, and it is particularly useful in analyzing for many trace metals. Landsberger and Wu (1993) analyzed air samples collected near Lake Ontario for Sb, As, Cd, In, I, Mo, Si, and V using INAA. They demonstrated that using INAA in conjunction with epithermal neutrons and Compton suppression produces very precise values with relatively low detection limits.

Enriched rare-earth isotopes have been analyzed via INAA and used to trace sources of particulate matter from a coal-fired power plant (Ondov et al., 1992), from various sources in the San Joaquin Valley (Ondov, 1996), from intentially tagged (iridium) diesel emissions from

sanitation trucks (Suarez et al., 1996; Wu et al., 1998), and from iridium-tagged emissions from school buses (Wu et al., 1998).

An intercomparison was conducted in which 18 pairs of filters were sent to participants in the Coordinated Research Program (CRP) on Applied Research on Waste Using Nuclear Related Analytical Techniques (Landsberger et al., 1997). As part of that study, participants used PIXE, INAA, XRF, or AAS to analyze the samples. Many of the results for XRF and PIXE in the coarse fraction were observed to be biased low compared to INAA. The authors speculated that there is a systematic error because of self-attenuation of the x-rays resulting from the particle size effect.

2.2.6.7 Atomic Absorption Spectrophotometry (AAS)

AAS is applied to the residue of a filter extracted in a strong solvent to dissolve the solid material; the filter or a portion of it is also dissolved during this process (Ranweiler and Moyers, 1974; Fernandez, 1989; Jackson and Mahmood, 1994; Chow et al., 2000a). A few milliliters of this extract are injected into a flame where the elements are vaporized. Elements absorb light at certain wavelengths in the visible spectrum, and a light beam with wavelengths specific to the elements being measured is directed through the flame to be detected by a monochrometer. The light absorbed by the flame containing the extract is compared with the absorption from known standards to quantify the elemental concentrations. AAS requires an individual analysis for each element, and a large filter or several filters are needed to obtain concentrations for a large variety of elements. AAS is a useful complement to other methods, such as XRF and PIXE, for species such as beryllium, sodium, and magnesium which are not well-quantified by these methods. Airborne particles are chemically complex and do not dissolve easily into complete solution, regardless of the strength of the solvent. There is always a possibility that insoluble residues are left behind and that soluble species may co-precipitate on them or on container walls.

AAS was used to characterize the atmospheric deposition of trace elements Zn, Ni, Cr, Cd, Pb, and Hg to the Rouge River watershed by particulate deposition (Pirrone and Keeler, 1996). The modeled deposition rates were compared to annual emissions of trace elements that were estimated from the emissions inventory for coal and oil combustion utilities, iron and steel manufacturing, metal production, cement manufacturing, and solid waste and sewage sludge incinerators. They found generally good agreement between the trend observed in atmospheric

inputs to the river (dry + wet deposition) and annual emissions of trace elements, with R^2s varying from ≈0.84 to 0.98. Both atmospheric inputs and emissions were found to have followed downward trends for Pb. For the period of 1987 to 1992, steady increases were observed for Cd (major sources are municipal solid waste incineration, coal combustion, sludge incineration, and iron and steel manufacturing), Cr and Ni (major sources are iron and steel production and coal combustion), and Hg (major sources are coal, the contribution from which had decreased from 53 to 45%, and municipal, solid, and medical waste incineration, the contribution from which has increased).

2.2.6.8 Inductively Coupled Plasma with Atomic Emission Spectroscopy (ICP-AES)

ICP-AES introduces an extracted sample into an atmosphere of argon gas seeded with free electrons induced by high voltage from a surrounding Tesla coil (Fassel and Kniseley, 1974; McQuaker et al., 1979; Lynch et al., 1980; Harman, 1989; Tyler, 1992; Baldwin et al., 1994). The high temperatures in the induced plasma raise valence electrons above their normally stable states. When these electrons return to their stable states, a photon of light is emitted that is unique to the element that was excited. This light is detected at specified wavelengths to identify the elements in the sample. ICP-AES acquires a large number of elemental concentrations using small sample volumes with acceptable detection limits for atmospheric samples. As with AAS, this method requires complete extraction and destruction of the sample.

2.2.6.9 Inductively Coupled Plasma with Mass Spectroscopy (ICP-MS)

ICP-MS has been applied in the analysis of personal exposure samples (Tan and Horlick, 1986; Gray and Williams, 1987a,b; Nam et al., 1993; Munksgaard and Parry, 1998; Campbell and Humayun, 1999). Ion species generated from ICP and from the sample matrix can produce a significant background at certain masses resulting in formation of polyatomic ions that can limit the ability of ICP-MS to determine some elements of interest. Cool plasma techniques have demonstrated the potential to detect elements at the ultra-trace level (Nham et al., 1996) and to minimize common molecular ion interferences (Sakata and Kawabata, 1994; Turner, 1994; Plantz, 1996). Detection limits of ICP-MS using a one-second scan are typically in the range of 10^{-3} ng/m^3, which is an order of magnitude lower than other elemental analysis methods. The instrument can also be set up to analyze a wide dynamic range of aerosol concentrations. Isotope

analysis can also be performed with ICP-MS. Intercomparison studies are needed to establish the comparability of ICP-MS with other non-destructive filter analysis methods.

Keeler and Pirrone (1996) used ICP-MS to determine trace elements Cd, Mn, V, As, Se, and Pb in atmospheric fine particulate ($PM_{2.5}$) and total suspended particulate samples collected in two Detroit sites. The results were used in a deposition model with additional measurements using AAS to estimate the dry deposition flux of trace elements to Lake Erie.

2.2.6.10 Scanning Electron Microscopy (SEM)

Mamane et al. (2001) investigated the use of computer-controlled scanning electron microscopy (CCSEM) as a way of supplementing XRF analysis and providing automated analysis of particle size, chemistry, and particle classification. An ambient coarse particulate sample from Baltimore was collected on a polycarbonate filter for this analysis. CCSEM analyses were conducted for 2819 particles in 78 randomly selected fields of view during an unattended 8-h run. Mamane et al. confirmed the stability of the CCSEM instrument over several hours of operation. The physical properties of the sample such as particle diameter, mass loading per field, and particle number per field were well represented by analyzing approximately 360 particles with little additional information gained by analyzing more particles. Teflon filters are not well suited for SEM analyses. Analysis of fine PM is expected to pose analytical challenges not addressed in the present study (Mamane et al., 2000).

Nelson et al. (2000) applied Raman chemical imaging and SEM (Raman/SEM) to study the size, morphology, elemental and molecular composition, and molecular structure of fine particulate matter. In their study, filter compatibility was examined, and Raman/SEM chemical imaging was conducted for several standard materials as well as for ambient $PM_{2.5}$ samples. Polycarbonate was determined to be a suitable substrate for both SEM and Raman chemical imaging analysis.

Conner et al. (2001) used CCSEM with individual X-ray analysis to study the chemical and physical attributes of indoor and outdoor aerosols collected around a retirement home in Baltimore. The CCSEM technique was demonstrated to be capable of identifying spherical particles typical of combustion or other high temperature (presumably industrial) processes as well as pollens and spores. Indoor particles originating from cosmetics were also identified.

2.2.7 Elemental and Organic Carbon in Particulate Matter

Three classes of carbon are commonly measured in ambient aerosol samples collected on quartz-fiber filters: (1) organic, volatilized, or non-light absorbing carbon (organic carbon, OC); (2) elemental or light-absorbing carbon (elemental carbon, EC); and (3) carbonate carbon (CC). The sum of OC, EC, and CC in PM gives the total carbon (TC). Carbonate carbon (i.e., K_2CO_3, Na_2CO_3, $MgCO_3$, $CaCO_3$), which can be determined on a separate filter section by measurement of the carbon dioxide (CO_2) evolved upon acidification (Johnson et al., 1980), is usually on the order of 5% or less for particulate samples collected in urban areas (Appel, 1993). Black carbon (BC) refers to an estimate of EC measured by absorption of visible light. The 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) listed several filter-based, thermal methods for measuring OC and EC and described the thermal/optical reflectance (TOR) method that was noted, along with thermal manganese oxidation, to be one of the most commonly applied methods in the United States at the time. In thermal separation methods, thermally evolved OC and EC are oxidized to CO_2 and quantified either by nondispersive infrared detection or electrochemically or by reducing the CO_2 to CH_4 and quantifying CH_4 via flame ionization detection (FID). The various methods give similar results for TC, but not for EC or OC.

Chow and Watson (1998) summarize different carbon analysis methods along with their measurement principles. The definitions of organic and elemental carbon are operational (i.e., method dependent) and reflect the method and purpose of measurement. Elemental carbon is sometimes termed “soot”, “graphitic carbon”, or “black carbon.” For studying visibility reduction, light-absorbing carbon is a more useful concept than elemental carbon. For source apportionment by receptor models, several consistent but distinct fractions of carbon in both source and receptor samples are desired, regardless of their light-absorbing or chemical properties. Differences in ratios of the carbon concentrations in these fractions form part of the source profile that distinguishes the contribution of one source from the contributions of other sources (Watson et al., 1994a).

Light-absorbing carbon is not entirely graphitic carbon because there are many organic materials which absorb light (e.g., tar, motor oil, asphalt, coffee). Even the “graphitic” black carbon in the atmosphere has only a poorly developed graphitic structure with abundant surface chemical groups. “Elemental carbon” is a poor but common description of what is measured. For example, a substance of three-bond carbon molecules (e.g., pencil lead) is black and

completely absorbs light, but four-bond carbon in a diamond is completely transparent and absorbs very little light. Both are pure, elemental carbon.

Chow et al. (1993) document several variations of the thermal method for organic and elemental carbon. The thermal/optical reflectance (TOR), thermal/optical transmission (TOT), and thermal manganese oxidation (TMO) methods are most commonly used for the analysis of organic and elemental carbon in atmospheric PM. Filter transmission analysis is often performed to estimate particle light absorption which is proportional to the level of elemental carbon in the atmosphere. These methods are discussed in detail in the following subsections.

The thermal manganese oxidation (TMO) method (Mueller et al., 1982; Fung, 1990) uses manganese dioxide present and in contact with the sample throughout the analysis as the oxidizing agent. Temperature is relied upon to distinguish between organic and elemental carbon. Carbon evolving at 525 °C is classified as organic carbon, and carbon evolving at 850 °C is classified as elemental carbon. This method has been used in the SCENES (Subregional Cooperative Electric Utility, Department of Defense, National Park Services, and Environmental Protection Agency Study); (Sutherland and Bhardwaja, 1987; Mueller et al., 1986) visibility network, as well as in the SCAQS (Southern California Air Quality Study) (Chow et al., 1994a,b; Watson et al., 1993, 1994a,b).

The thermal/optical reflectance (TOR) method of carbon analysis developed by Huntzicker et al. (1982) has been adapted by several laboratories for the quantification of organic and elemental carbon in PM collected on quartz-fiber filters. Although the principle used by these laboratories is identical to that of Huntzicker et al. (1982), the details differ with respect to calibration standards, analysis time, temperature ramping, and volatilization/combustion temperatures.

In the most commonly applied version of the TOR method (Chow et al., 1993), a filter is submitted to volatilization at temperatures ranging from ambient to 550 °C in a pure helium atmosphere, then to combustion at temperatures between 550 °C to 800 °C in a 2% oxygen and 98% helium atmosphere with several temperature ramping steps. The carbon that evolves at each temperature is converted to methane and quantified with a flame ionization detector. The reflectance from the deposit side of the filter punch is monitored throughout the analysis. This reflectance usually decreases during volatilization in the helium atmosphere owing to the pyrolysis of organic material. When oxygen is added, the reflectance increases as the

light-absorbing carbon is combusted and removed. Organic carbon is defined as that which evolves prior to re-attainment of the original reflectance, and elemental carbon is defined as that which evolves after the original reflectance has been attained. By this definition, "organic carbon" is actually organic carbon that does not absorb light at the wavelength (632.8 nm) used, and "elemental carbon" is light-absorbing carbon (Chow et al., 1993).

The thermal/optical transmission (TOT) method applies to the same thermal/optical carbon analysis method except that transmission instead of reflectance of the filter punch is measured. The National Institute for Occupational Safety and Health (NIOSH) Method 5040 for monitoring elemental carbon as a marker for particulate diesel exhaust is based upon the TOT method (Birch and Cary, 1996). The TOT OC/EC method consists of a two-stage process with the first stage being conducted in a pure helium atmosphere at temperatures of 250, 500, 650, and 850 °C for a total of 4.5 minutes and the second stage conducted in a 2% oxygen/98% helium mix at temperatures of 650, 750, 850, and 940 °C for 4 minutes. A pyrolysis base correction is made based on the transmission measurement.

Chow et al. (1993) document several variations of the thermal (T), thermal/optical reflectance (TOR), thermal/optical transmission (TOT), and thermal manganese oxidation (TMO) methods for organic and elemental carbon. Comparisons among the results of the majority of these methods show that they yield comparable quantities of total carbon in aerosol samples, but the distinctions between organic and elemental carbon are quite different (Cadle and Groblicki, 1982; Cadle and Mulawa, 1990; Countess, 1990; Hering et al., 1990; Birch, 1998; Schmid et al., 2001). TOR was consistently higher than TMO for elemental carbon, especially in woodsmoke-dominated samples where the disparity was as great as sevenfold. For the sum of organic and elemental carbon, these methods reported agreement within 5% to 15% for ambient and source samples (Houck et al., 1989; Kusko et al., 1989; Countess, 1990; Shah and Rau, 1990) and within 3% on carefully prepared standards. Evaluation of these methods thus is a matter of assessing how they differentiate between organic and elemental carbon. The TMO method attributes more of the total carbon to organic carbon and less to elemental carbon than the TOR and TOT methods. None of the methods represents an ideal procedure for the separation of organic from elemental carbon.

In a methods comparison study (Countess, 1990), it was shown that it is necessary to minimize or correct for pyrolytically generated EC ("char") and that CC found in wood smoke

and automobile exhaust samples may interfere with some of the thermal methods. Recently, Lavanchy et al. (1999) reported on a study in which the operation of a catalytic oxidation system was modified in an attempt to minimize pyrolysis of OC and, at the same time, minimize the contribution of $CaCO_3$. The system uses two ovens, one at 340 °C and one at 650 °C. The filter sample is placed in a moveable sample boat. In order to minimize charring, the sample is first flash heated in the 650°C oven for 1 min. It is then inserted into the 340°C stage of the two-stage oven. In both steps, OC is oxidized to CO_2 in the presence of O_2. After 42 mins, the filter is moved into the second-stage oven. During this third step, EC is oxidized to CO_2 at 650°C for 32 min. This temperature is reported to be sufficient to completely oxidize EC, but with only about 1% of the $CaCO_3$ being vaporized (Lavanchy et al., 1999; Petzold et al., 1997). To test for charring, they challenged their system with atmospheric samples for which duplicates were analyzed via the German reference method (in which a solvent extraction is used to remove organics before combustion) for measuring OC and EC in atmospheric samples (Petzold and Niessner, 1996). Lavanchy et al. (1999) reported a high correlation ($R^2 = 0.97$) between their thermal oxidation method and the German reference method (VDI). The slope of the EC:EC VDI line was 0.92, and the intercept was –0.37 μg cm^{-2}. They also reported detection limits of 1.3 μg for EC and 1.8 μg for OC.

Pyrolytic char is corrected for in thermal-optical analysis. In thermal-optical methods (Birch and Cary, 1996; Chow et al., 1993), punches from a quartz sampling filter are inserted into the carbon analyzer and heated in a helium atmosphere to volatilize organic carbon. Then, the temperature is reduced, and oxygen is added to the carrier gas so that desorbed compounds are oxidized to CO_2, reduced to methane, and measured in a flame ionization detector. In order to account for the portion of the OC that is pyrolyzed, a He-Ne laser monitors the sample reflectance (or transmittance). As the pyrolysis occurs, the sample gets darker, and the reflectance decreases. As elemental carbon is removed, the filter lightens, and the reflectance increases until all carbon has been removed from the filter. The split between organic and elemental carbon is considered to be the point at which the reflectance regains its prepyrolysis value with material removed prior to this point being considered organic and that after, elemental.

The thermal/optical transmission (TOT) method is similar to the TOR with the primary difference being that light transmission rather than reflectance is monitored on the filter

throughout the analysis. The TOT method of Birch and Cary (1996) consists of a two-stage process with the first stage being conducted in a pure helium atmosphere and the second stage conducted in a 10% oxygen-helium mix. The temperature is raised to approximately 820 °C in the helium phase, during which organic and carbonate carbon are volatilized from the filter. In the second stage, the oven temperature is reduced then raised to about 860 °C. During the second stage, pyrolysis correction and EC measurement are made. Figure 2-18, an example of a TOT thermogram, shows temperature, transmittance, and FID response traces. Peaks are evident that correspond to OC, CC, EC, and pyrolitic carbon (PC). As can be seen in this figure, the high temperature in the first stage allows for decomposition of CC. The ability to quantify PC is particularly important in high OC/EC regions (like wood-smoke-impacted airsheds), allowing for the volatilization of any remaining complex organic compounds so they are not apportioned to the EC phase.

The NIOSH Method 5040 for monitoring elemental carbon as a marker for particulate diesel exhaust is based upon a TOT method analyzer (Birch and Cary, 1996); whereas the OC/EC method specified for the IMPROVE network is the TOR method (Chow et al., 2000b). Chow et al. (2000b) compared the OC, EC, and TC measurements from NIOSH and IMPROVE methods. The two methods use different temperature and atmospheric controls to separate OC and EC. In addition, the NIOSH (TOT) method uses light transmission through the filter and the IMPROVE (TOR) method uses light reflectance from the filter to measure pyrolyte carbon. The IMPROVE thermal protocol specifies organic carbon fractions at 120, 250, 450, and 550 ºC in a nonoxidizing atmosphere (He) and elemental organic fractions at 550, 700, and 800 ºC in an oxidizing atmosphere. The NIOSH method differs in its thermal protocol, which has organic carbon fractions at 250, 500, 650, and 850 ºC in a nonoxidizing atmosphere (also He) and elemental carbon fractions at 650, 750, and 850 ºC in an oxidizing atmosphere. The high temperature before addition of oxygen in the NIOSH method is necessary to quantify particulate carbonate, which evolves between 650 and 830 ºC (Birch and Cary, 1996). The two methods also differ in the specified residence times at each temperature setpoint. The residence times at each setpoint are typically longer for the IMPROVE analysis as compared to the NIOSH analysis.

Chow et al. (2000b) analyzed 60 quartz filter samples that represented a wide variety of aerosol compositions and concentrations. The TC measurements from each protocol were in good agreement with no statistically significant differences. A statistically significant difference



Figure 2-18. This thermogram, for a sample containing rock dust (a carbonate source) and diesel exhaust, shows three traces that correspond to temperature, filter transmittance, and FID detector response. Peaks correspond to organic (OC), carbonate (CC), pyrolytic (PC), and elemental (EC) carbon. The final peak is a methane calibration peak.

Source: Birch and Cary (1996).

was observed in the fraction of TC that is attributed to EC as determined by the IMPROVE and NIOSH thermal evolution protocols with the IMPROVE EC measurements typically higher than the NIOSH EC measurements. This difference was attributed to the 850 ºC temperature step in the oxidizing atmosphere in the NIOSH protocol. Chow et al. (2000b) compared the OC for each method and found that the two methods showed good agreement when the 850 ºC nonoxidizing temperature step in the NIOSH method was not included in determination of OC. There was also a difference between the reflectance and transmittance detection methods in the pyrolysis adjustment, although this difference was most noticeable for very black filters for which neither

reflectance nor transmittance was able to accurately detect further blackening by pyrolysis. Because OC and EC are operationally defined parameters, Chow et al. (2000b) pointed out that it is important to retain ancillary information when reporting EC and OC by these analytical methods, so that comparisons can be made among measurements taken at different sites using these two methods.

Further refinement of thermal techniques has resulted in the evolved gas analysis (EGA) method, described by Grosjean et al. (1994). This technique involves combustion of particulate matter samples in an oxidizing environment while the temperature is raised from 100 to 600 ^{o}C. The amount of evolved CO_2 contains information about the volatility of the organic aerosol compounds. Grosjean et al. (1994) present thermograms both for specific organic compounds (e.g., adipic acid) and for specific sources (e.g., vehicular traffic). They suggest that EGA may be useful for source apportionment applications. Kirchstetter et al. (2001) and Novakov et al. (1997) have also used EGA to provide insights regarding organic sampling artifacts.

A more recent international intercomparison on the analysis of carbonaceous aerosols on quartz fiber filters was organized by the Vienna University of Technology and involved seventeen laboratories and nine different thermal and optical methods (Schmid et al., 2001). All laboratories were sent punches from three 150-mm quartz fiber filters that had been exposed for 24 h near a high traffic street in Berlin. Five laboratories employed VDI 2465 methods that are official methods in Germany. Two of these laboratories used the VDI 2465/1 method that determines extractable organic carbon, non-extractable organic carbon, and elemental carbon. The solvent extraction step incorporates a 50:50 vol% mixture of toluene and 2-propanol for the removal of the extractable organic carbon. The filter is dried, and the non-extractable organic carbon is removed by thermal desorption under nitrogen at 500 °C. The remaining carbon on the filter, assumed to be elemental carbon, is combusted in an oxidizing atmosphere at 650 °C, and the CO_2 produced is detected by coulometry.

The other three laboratories using VDI 2465 methods incorporated the VDI 2465/2 method that separates the carbonaceous fractions of the aerosols due to their different thermal stabilities. The sample is first heated in an oxygen free inert gas (either helium or argon) at temperatures of 350 and 620 °C over a copper/cerium IV oxide catalyst to remove the organic carbon. The sample is heated at 700 °C in at least 20% oxygen to determine the elemental carbon, and the resulting CO_2 is detected by nondispersive infrared spectrophotometry (NDIR). A sixth

laboratory employed a thermal analyzer with a multistep program similar to the VDI 2465/2 method.

Four laboratories utilized different thermal procedures and techniques for detecting the emerging CO_2. Of these first nine laboratories, one adapted its technique to correct for pyrolytically formed char. The 10th and 11th laboratories used a thermal optical transmission method (the Sunset Laboratories Inc. instrument) with slightly different temperature programs and atmospheres. The 12th laboratory used a homemade version of the thermal optical transmission method. The 13th laboratory used thermal optical reflectance.

The 14th laboratory determined only total carbon using a Shimadzu TOC 5000 with a solid sampler module (SSM-500a). In this method, the sample is combusted at 900 °C over cobalt oxide and platinum catalysts and the evolved CO_2 is measured by NDIR. The 15th laboratory analyzed TC by catalytic combustion, using an elemental analyzer CE 440 (Leeman Labs, Inc.) with standard combustion analysis and thermal conductivity detection. Black carbon (BC) was determined by optical transmissiometry, using an aethalometer and an empirical constant of 19 $cm^2\ g^{-1}$. The 16th laboratory analyzed only BC using an integrating sphere. The 17th laboratory utilized a two-step thermal method, in which the organic material is removed under pure oxygen at 340 °C and the remaining carbon is determined by coulometric titration of the CO_2 evolved at 1100 °C in a carbon analyzer.

Good agreement of the TC results was obtained by all laboratories with only two outliers in the complete data set. The relative standard deviation between laboratories for the TC results were 6.7, 10.6, and 8.8% for the three samples. In contrast, the EC results were much more variable. The relative standard deviation between laboratories for the EC results were 36.6, 24.4, and 45.5% for the three samples. The VDI methods, especially the VDI 2465/2, were found to give generally higher amounts of EC than the thermal-optical methods. This trend was detected for all samples. The authors recognized that uncorrected thermal methods are prone to positive artifacts by charring during pyrolysis. They also noted that when using solvent extraction methods, the dissolution of polymeric aerosol constituents may not be successful. Both of these effects would lead to overestimation of the EC fraction. When the laboratories were grouped according to their methods, the relative standard deviations between laboratories was much smaller. This study demonstrates that the TC measurement can yield similar results from a variety of methods, but the EC measurement is highly dependent upon the method used. The

problems associated with the determination of EC are exacerbated by the lack of a standard reference material.

Elemental carbon can also be measured by optical absorption (OA), photoacoustic spectroscopy, and nonextractable mass (Chow et al., 1993). Optical absorption, assumed due entirely to elemental carbon, can be measured by determining light transmission through Teflon-membrane and quartz-fiber filters before and after sampling with a transmission densitometer. Informal intercomparisons among different filter transmission methods have shown high correlations of absorption, but differences of up to a factor of two in absolute values (Watson et al., 1988a,b). These differences are functions of the type of filter, filter loading, the chemical and physical nature of the deposit, the wavelengths of light used, calibration standards, and light diffusing methods. At the current time, there is no agreement on which combination most accurately represents light absorption in the atmosphere. This method is applied with the knowledge that absolute differences in absorption may be found between the measurements made on Teflon-membrane and quartz-fiber filters and with respect to absolute absorption measurements made on the same samples in other laboratories.

Black carbon (BC) is also used, in addition to the thermal and thermal/optical methods, for determining EC as a measure of soot (Penner and Novakov, 1996). Both EC and BC define a similar fraction of aerosol; but EC is determined based on thermal properties, whereas BC is based on light-absorption properties. Optical methods for determining BC tend to suffer from calibration problems (Hitzenberger et al., 1996). Lavanchy et al. (1999) compared their EC concentrations as determined from their catalytic thermal oxidation method to BC concentrations determined using an aethalometer operated at the same site; and they found that the instrumental calibration factor provided by the manufacturer was on the order of two times the calibration factor they determined (9.3 ± 0.4 m^2g^{-1}). It is possible to calculate a theoretical specific absorption coefficient (B_a) from Mie theory, given a known size distribution and refractive index. The B_a is defined as absorption per mass concentration and can be calculated given the sample filter area, the total deposited mass, and absorption signals for both the loaded and unloaded filters. Often, when no direct measurements are available, values of B_a on the order of 10 m^2g^{-1} have been used (Hitzenberger et al., 1996). Typically BC aerosols have values of B_a between 3 and 17 m^2g^{-1} (Hitzenberger et al., 1996).

Black carbon can be used as an indicator of particles from diesel engines. Therefore, it is important that accurate values for B_a are available. Hitzenberger et al. (1996) investigated the feasibility of using an integrating sphere photometer as an adequate measurement system for the BC content and the absorption coefficient. Based on samples collected during a 10-day period in May 1994, they determined that the usually assumed value of 10 m^2g^{-1} was also applicable to aerosol BC occurring in Vienna.

In another study (Hitzenberger et al., 1999), the integrating sphere method was compared to an aethalometer (Hansen et al., 1984), the thermal method of Cachier et al. (1989), and the thermal/optical method of Birch and Cary (1996). The absorption coefficients that were obtained from both the integrating sphere and the aethalometer were comparable. The BC mass concentrations obtained from the aethalometer were 23% of those obtained from the integrating sphere. Compared to the thermal method, the integrating sphere overestimated the BC mass concentrations by 21%. Compared to the thermal/optical method, the integrating sphere was within 5% of the 1:1 line. However, the data were not so well correlated.

In 1986, the Carbonaceous Species Methods Comparison Study (CSMCS) was conducted in Los Angeles. The CSMCS was mentioned in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). Hansen and McMurry (1990) compared two very dissimilar methods for aerosol elemental carbon. One involved collection of impactor samples backed by a quartz fiber afterfilter followed by EC analysis by oxidation in helium over a MnO_2 catalyst; the other real-time measurements using an aethalometer (an optical absorption technique). They found good agreement between these two very different methods. The CSMCS interlaboratory precision for total carbon was 4.2% (Turpin et al., 2000). However, because the split between OC and EC is operationally defined, there was substantial interlaboratory variability in OC and EC (e.g., 34% for EC [Turpin et al., 1990]). The implications for data analysis are twofold: (1) the analysis method used must be reported with particulate carbon data and (2) comparative analyses should not be conducted with data analyzed by more than one carbon analysis method unless the mutual compatibility of the methods has been demonstrated.

EC/OC Summary. With the limitations and precautions described above, laboratory analyses for the carbonaceous properties of collected particles have matured to the point where they can be performed with commercially-available instruments following established standard

operating procedures and with traceability to common standards. However, carbon analysis continues to be a subject of research, and carbon methods are currently being compared as part of other studies (e.g., the Atlanta Supersite). The state of the art for soot measurements continues to develop; and, although advances are being made, the definitions of EC and BC continue to be operational and determined by the method employed. Similarly, the distinction between OC and EC is defined operationally. Reports of EC/OC measurements should therefore include mention of the method with which the species were determined. Finally, if possible, all ancillary data should be retained, to allow later comparison to other methods.

2.2.8 Ionic Species

Ion chromatography (IC) is widely used for analyzing ionic species in the water-soluble portion of suspended PM. IC is the method of choice for the measurement of sulfate, nitrate, ammonium, sodium, and potassium ions for the NAMS program. Aerosol strong acidity, H^+, is determined by titration of a water solution of PM collected following a series of annular denuders to remove acid and basic gases with back-up filters to collect NH_3 and HNO_3 that volatilize from the PM during collection. The 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) discussed measurement of ions by IC (Section 4.3.3.1) and of strong acidity (Sections 3.3.1.1 and 4.3.3.1); so, no further details will be discussed here.

2.2.9 Continuous Monitoring

The EPA expects that many local environmental agencies will operate continuous PM monitors. All currently available continuous measurements of suspended particle mass share the problem of dealing with semivolatile PM components. So as not to include particle-bound water as part of the mass, the particle-bound water must be removed by heating or dehumidification. However, heating also causes loss of ammonium nitrate and semivolatile organic components. A variety of potential candidates for continuous measurement of mass or chemical components will be discussed in this section.

2.2.9.1 Continuous Measurement of Mass

Tapered Element Oscillating Microbalance (TEOM)

The advantages of continuous PM monitoring and the designation of the TEOM as an equivalent method for PM_{10}, have led to the deployment of the TEOM at a number of air monitoring sites. The TEOM also is being used to measure $PM_{2.5}$. The TEOM differs from the federal reference methods for particulate mass in that it does not require equilibration of the samples at a specified temperature and relative humidity. The TEOM samples at a constant temperature and is typically heated to some temperature higher than the ambient temperature (Meyer et al., 1995; Meyer and Rupprecht, 1996); whereas the FRM samples at the ambient temperature. Thus, the TEOM does not provide data equivalent to the FRM because of losses of volatile species. Volatilization losses in the TEOM sampler can be reduced by operating the instrument at 30 °C, rather than the typical 50 °C specified or by using a Nafion diffusion dryer instead of heating to dehumidify the particles.

This difference in operation and implications for fine particle measurements was examined by researchers at CSIRO Atmospheric Research in Australia (Ayers et al., 1999). That group compared 24-h mean $PM_{2.5}$ mass concentrations as determined by a TEOM and by two manual, gravimetric samplers (a low-volume filter sampler and a MOUDI sampler) in four Australian cities, on 15 days in the winter half-year. The TEOM was operated at 50 °C at one location and at 35 °C at the other three locations. A systematically low TEOM response in comparison to the integrated gravimetric methods was observed. In a comprehensive study, Allen et al. (1997) reported results in which TEOM data collected at 10 urban sites in the United States and Mexico were compared with 24 h integrated mass concentrations for both PM_{10} and $PM_{2.5}$. They collected a large data set that included both winter and summer seasons. Allen et al. (1997) concluded that, especially for urban areas, a significant fraction of PM_{10} could be semivolatile compounds that could be lost from the heated filter in the TEOM leading to a systematic difference between the TEOM and the EPA FRM for PM_{10}. They suggested that this difference is likely to be larger for $PM_{2.5}$ than it is for PM_{10} (Allen et al., 1997).

In a similar study conducted in Vancouver, British Columbia, the effect of equilibration temperature on PM_{10} concentrations from the TEOM was examined. Two collocated TEOM monitors operated at 30 and 50 °C, respectively, were operated in the Lower Fraser Valley in British Columbia for a period of approximately 17 months (Mignacca and Stubbs, 1999). A third

TEOM operating at 40 °C was operated for 2 months during this period. They found that, on average, the 1-h average PM_{10} from the TEOM operating at 30 °C was consistently greater than that from the TEOM operated at 50 °C. For the period during which the third TEOM was operated (at 40 °C), the PM_{10} from that instrument was between those values for the other two instruments. They also found that the differences in masses were proportional to the PM_{10} loading, and more strongly correlated to the PM_{10} from the TEOM operated at the lower temperature. They recommended that the TEOM monitors be operated at 40 ºC as opposed to operating at 50 °C in summer and 30 °C in winter, in order to avoid introducing a methodological seasonal bias.

A new sample equilibration system (SES) was developed to reduce losses of semivolatile species from the $PM_{2.5}$ TEOM by conditioning the sample stream to lower humidity and temperature (Meyer et al., 2000). The SES utilizes humidity sensors and a Nafion dryer designed for low particle loss. The dryer fits between the flow splitter that follows the size-selective inlet and the sensor unit. A dry purge gas flows over the exterior of the Nafion tubing and allows for self-regeneration. A TEOM with $PM_{2.5}$ inlet and equipped with an SES was operated at 30 °C alongside another TEOM operating at 50 °C without the SES in Albany, NY, over a 6-day period during a summertime high-temperature, high-relative-humidity episode. The SES maintained the sample air relative humidity under 30%, and the TEOM with the SES generally measured more mass than the other TEOM. The TEOM with SES also was operated alongside an FRM-type sampler for the period of June 6 through September 25, 1999. The correlation between the FRM and TEOM/SES showed a slope of 1.0293 and R^2 of 0.9352; whereas the correlation between the FRM and the TEOM without SES and operating at 50 °C showed a slope of 0.8612 and R^2 of 0.8209. The SES can be installed on existing TEOM monitors.

Patashnick et al. (2001) developed a differential TEOM system that is based on a pair of TEOM sensors, each of which is preceded by its own electrostatic precipitator (ESP) and downstream from a common size selective inlet. By alternately switching the ESPs on and off and out of phase with each other, the two sensors measure “effective mass” that includes both the nonvolatile component and the volatile component sampled by the TEOM, less the volatile component that vaporized during the sampling interval. On the sensor side with the ESP turned on, there is no particle collection on that filter so that only volatilization of previously collected particles continues. This allows a correction for the effective mass as measured by the first

sensor by subtracting out the volatilization artifact and leaving the nonvolatile and volatile components of the particulate matter. This system has yet to be well characterized for other biases or interferences such as reactions on the filters, particle collection efficiency of the ESPs, and particle and semivolatile material losses.

Real-Time Total Ambient Mass Sampler (RAMS)

The RAMS, a monitor based on diffusion denuder and TEOM monitor technology, has been developed, validated, and field tested for the real-time determination of total fine PM mass, including semivolatile PM (Eatough et al., 1999a; Obeidi and Eatough, 2002; Obedi et al., 2002; Pang et al., 2001). The RAMS measures the total mass of collected particles including semivolatile species with a TEOM monitor using a "sandwich filter." The sandwich contains a Teflon coated particle collection filter followed by a charcoal-impregnated filter (CIF) to collect any semivolatile species lost from the particles during sampling. Because the instrument measures total mass collected by the sandwich filter, all gas phase compounds that can be adsorbed by a CIF must be removed from the sampling stream prior to the TEOM monitor. Laboratory and field validation data indicate that the precision of fine PM mass determination is better than 10%. The RAMS uses a Nafion dryer to remove particle-bound water from the suspended particles and a particle concentrator to reduce the amount of gas phase organics that must be removed by the denuder. An example of data from the RAMS, the TEOM, and the PC-BOSS is shown in Figure 2-19. This figure also shows the $PM_{2.5}$ mass from the TEOM as being negative for the hours of 16 to 19. This likely results from the loss of volatile materials from the heated filter.

Continuous Ambient Mass Monitor (CAMM)

Koutrakis and colleagues (Koutrakis et al., 1996; Wang, 1997) have developed CAMM, a technique for the continuous measurement of ambient particulate matter mass concentration based on the measurement of pressure drop increase with particle loading across a membrane filter. Recently, Sioutas et al. (1999) examined the increase in pressure drop with increasing particle loading on Nuclepore filters. They tested filters with two pore diameters (2 and 5 μm) and filter face velocities ranging from 4 to 52 cm s^{-1} and examined the effects of relative humidity in the range of 10 to 50%. They found that, for hygroscopic ammonium sulfate



Figure 2-19. Comparison of mass measurements with collocated RAMS (real-time data), PC-BOSS (1-h data), FRM $PM_{2.5}$ sampler (average of 24-h data), and a conventional TEOM monitor (real-time data). The semivolatile fine particulate matter is sampled with the RAMS and PC-BOSS, but not with the TEOM monitor or the FRM $PM_{2.5}$ single filter sampler. The PC-BOSS provides information on both the nonvolatile component (NV) and the semivolatile organic component (SVOC).

Source: Eatough et al. (1999a).

particles, the change in pressure drop per unit time and concentration was a strong function of relative humidity, decreasing with increasing relative humidity. These results suggest that particulate concentration measurements, similar to the method of Koutrakis et al. (1996) that uses the pressure drop method, may be subject to additional uncertainties if used in an environment where the ambient relative humidity cannot be controlled accurately. The current version of the CAMM (Wang, 1997) uses a particle concentrator, a Nafion dryer, and frequently moves the

filter tape to avoid artifacts due to evaporation of semivolatile components from the active portion of the filter tape which would occur if the atmospheric concentration of the semivolatile components decreased.

The CAMMS was recently operated alongside a gravimetric PM method (the Harvard Impactor, or HI) in seven U.S. cities selected for their distinctly different ambient particulate compositions and densities. The correlation between the two methods was high, with an overall r^2 of 0.90 and average CAMM/HI ratio of 1.07 (Babich et al., 2000).

Beta-Gauge Techniques

The use of absorption of beta radiation as a indicator of particle mass has been used effectively to measure the mass of equilibrated particulate matter collected on Teflon filters (Jaklevic et al., 1981a; Courtney et al., 1982). The technique also has been used to provide near real-time measurements with time intervals on the order of an hour (Wedding and Weigand, 1993). However, real-time beta gauge monitors experience the same problems as other continuous or near real-time particulate matter mass monitoring techniques. Particle-bound water must be removed to reduce the sensitivity of the indicated mass to relative humidity. However, the simplest technique, mild heating, will remove a portion of the ammonium nitrate and the semivolatile organic compounds as well as the particle-bound water.

An intercomparison study of two beta gauges at three sites indicated that the Wedding beta gauge and the Sierra Anderson (SA) 1200 PM_{10} samplers were highly correlated, $r > 0.97$ (Tsai and Cheng, 1996). The Wedding beta gauge was not sensitive to relative humidity but yielded results approximately 7% lower. This suggests that the mild heating in the beta gauge causes losses comparable to those caused by equilibration, although the differences could result from slight differences in the upper cut points. The Kimoto beta gauge that was operated at ambient temperature was sensitive to relative humidity yielding significantly higher mass concentrations relative to the SA 1200 for RH > 80% than for RH < 80% even though the correlation with the SA 1200 was reasonable ($r = 0.94$ for RH > 80% and 0.83 for RH < 80%).

Piezoelectric Microbalance

Piezoelectric crystals have mechanical resonances that can be excited by applying an alternating electrical voltage to the crystal. As the resonance frequencies are well defined, such

crystals (quartz in particular) have found applications as secondary time and frequency standards in clocks and watches. As for all mechanical resonators, the resonance frequency is a function of mass. Therefore, by monitoring the resonance frequency in comparison with a second crystal, one can continuously measure the mass deposited on the crystal (Sem et al., 1977; Bowers and Chuan, 1989; Ward and Buttry, 1990; Noel and Topart, 1994). Comparison with a second crystal largely compensates for the effect of temperature changes on the resonance frequency.

The piezoelectric principle has been used to measure particle mass by depositing the particles on the crystal surface either by electrostatic precipitation or by impaction (Olin and Sem, 1971). The collection efficiency of either mechanism has to be determined as function of particle size to achieve quantitative measurements. In addition, the mechanical coupling of large particles to the crystal is uncertain. Both single and multi-stage impactors have been used (Olin and Sem, 1971; Fairchild and Wheat, 1984). Quartz crystals have sensitivities of several hundred hertz per microgram. This sensitivity results in the ability to measure the mass concentration of a typical 100 $\mu g/m^3$ aerosol to within a few percent in less than one minute (Olin and Sem, 1971).

Coarse Particle Mass

The RAMS and CAMM are only appropriate for fine particle measurements ($PM_{2.5}$ or PM_1). However, the TEOM, beta gauge, and piezoelectric microbalance may be used to measure either $PM_{2.5}$ or PM_{10} (or a sample with any specified upper 50% size cut). A pair of such samplers may be used to measure thoracic coarse PM mass concentration ($PM_{10\text{-}2.5}$) by difference between the PM_{10} and $PM_{2.5}$ concentrations. However, concerns have been raised concerning the quality of the data from such difference calculations and the resulting potential biases in exposure assessment and risk determinations (Wilson and Suh, 1997; White, 1998). Misra et al. (2001) describe the development and evaluation of a continuous coarse particle monitor (CCPM) that may provide direct measurements of coarse mode PM mass concentrations at short time intervals (on the order of 5-10 min). The basis of the CCPM is enrichment of the coarse particle concentrations through use of virtual impaction while maintaining fine particle concentrations at ambient levels. The resulting aerosol mixture is analyzed using a standard TEOM for which the response is now dominated by the enriched coarse PM mass. The coarse PM concentrations determined from the CCPM were compared to those obtained with a MOUDI, operating with only the 10- and 2.5-micron cutpoint stages, and a Partisol dichotomous sampler. The CCPM

coarse particulate concentrations were highly correlated with both the MOUDI ($R^2 = 0.88$) and the Partisol ($R^2 = 0.88$) coarse PM concentrations. By operating the CCPM at a coarse particle enrichment factor of 25, the coarse PM concentration can be determined *a priori* without determination of the fine particle concentration, so long as the fine-to-coarse particle concentration ratios are not unusually high (i.e., 4-6). Misra et al. (2001) also concluded from field experiments that the coarse particulate concentrations determined from the CCPM were independent of the ambient fine-to-coarse particulate concentration ratio due to the decrease in particle mass median diameter that should accompany fine-to-coarse particulate concentration ratios during stagnation conditions.

2.2.9.2 Continuous Measurement of Elemental and Organic Carbon

Testing and refinement of models that simulate aerosol concentrations from gas and particle emissions require air quality measurements of approximately 1-h time resolution to reflect the dynamics of atmospheric transport, dispersion, transformation, and removal. Below we describe instruments that have been used to collect and analyze atmospheric organic PM with better than 2-h resolution. These instruments were all present at the Atlanta Supersite experiment during the summer of 1999, and an intercomparison of results is underway.

Turpin et al. (1990) describe an *in situ*, time-resolved analyzer for particulate organic and elemental carbon that can operate on a time cycle as short as 90 min. This analyzer collects particulate matter on a quartz fiber filter mounted in a thermal-optical transmittance carbon analyzer (Turpin et al., 1990). A second quartz fiber filter behind a Teflon filter in a second sampling port may also be analyzed to provide an estimate of the positive sampling artifact (i.e., gas adsorption on the quartz sampling filter). The organic material in the collected PM is thermally desorbed from the filter at 650 °C and oxidized at 1000 °C over a MnO_2 catalyst bed. The evolved CO_2 is converted to methane over a nickel catalyst, and the methane is measured in a flame ionization detector. Then the elemental carbon is oxidized at 350 °C in a 98% He-2% O_2 atmosphere. Correction is made for pyrolytic conversion of some of the organic particulate matter. The instrument was operated with a 2-h resolution during SCAQS in 1987 (Turpin and Huntzicker, 1991;1995), as well as during CSMCS in 1986 (Turpin et al., 1990). By using elemental carbon as a tracer for primary, combustion-generated organic carbon, these authors estimated the contributions of primary sources (i.e., material emitted in particulate form) and

secondary sources (i.e., particulate material formed in the atmosphere) to the total atmospheric particulate organic carbon concentrations in these locations.

An automated carbon analyzer with 15-min to 1-h resolution is now commercially available (Rupprecht et al., 1995) and has been operated in several locations including the Atlanta Supersite. It collects samples on a 0.1-μm impactor downstream of an inlet with a 2.5-μm cutpoint. Use of an impactor eliminates gas adsorption that must be addressed when filter collection is used. However, this collection system may experience substantial particle bounce and loss of a sizable fraction of EC since some EC is in particles < 0.2 μm. It is possible that ongoing research, in which particle size is increased by humidification prior to impaction, may result in an improvement in collection efficiency. In the analysis step, carbonaceous compounds are removed by heating in filtered ambient air. Carbonaceous material removed below 340 °C is reported as organic carbon, and material removed between 340 and 750 °C is reported as elemental carbon. Turpin et al. (2000) comment that it would be more appropriate to report carbon values obtained by this method as "low-" and "high-temperature" carbon, because some organics are known to evolve at temperatures greater than 340 °C (e.g., organics from woodsmoke).

As discussed earlier, black carbon (BC), a carbon fraction very similar to EC, is most commonly measured using an aethalometer, a commercially available, automated, time-resolved instrument (i.e., 5- to 15-min sample duration) that measures the light attenuation of aerosol particles collected on a filter tape (Hansen et al., 1984). The concentration of elemental carbon is derived from the light absorption measured on a filter using an estimate of the specific absorption (m^2/g) of elemental carbon on the filter; the specific absorption value is derived from laboratory and atmospheric tests and is specified by the manufacturer. The specific absorption value could be expected to vary with location, season, and source mix. Comparisons in atmospheric experiments at some locations with EC values measured by thermal methods confirm that the aethalometer provides a statistically meaningful estimate of EC concentration (Allen et al., 1999c; Liousse et al., 1993). For instance, Allen et al. (1999c) found the following statistical relationship for Uniontown, PA, during summer 1990: black carbon (aethaometer) = 0.95*EC (thermal) – 0.2 (r^2 = 0.925, n not specified but appears to be >50, EC range from 0 to 9 $\mu g/m^3$). Another source of error in aethalometer measurements arises from the sampling procedure. Particles are trapped within a three-dimensional filter matrix. Therefore, scattering of

transmitted and reflected light may erroneously be attributed to absorption, thus causing errors in the BC calculation. Ballach et al. (2001) investigated immersing the filter in oil of a similar refractive index as a means to minimize the interferences due to light scattering effects from the filter, similar to a procedure common in microscopy. BC measurements determined using the oil immersion technique were compared to those from an integrating sphere, a polar photometer, and Mie calculations. Aerosols tests included several pure carbon blacks from different generating procedures that were used to calibrate the immersion technique, pure ammonium sulfate aerosol, and external and internal mixtures of ammonium sulfate with varying amounts of carbon blacks. The oil immersion technique was also tested on ambient air samples collected at two different sites in the cities of Frankfurt am Main and Freiburg, Germany. Optical measurements, both of blank and loaded filters, show that the oil immersion technique minimizes scattering losses. Ballach et al. (2001) found that site-related effects were reduced and that there was reasonably good agreement with the other optical techniques as well as with the Mie calculations.

Measurement of aerosol light absorption utilizing photoacoustic spectroscopy has been examined as a continuous method for measuring elemental carbon mass concentrations (Petzold and Niessner, 1996; Arnott et al., 1999; 2000). Like the aethalometer, this method measures light absorption; however, unlike most other light absorption methods, the photoacoustic technique does not require a filter. The photoacoustic spectrometer of Arnott and coworkers was demonstrated during the Northern Front Range Air Quality Study and compared to an aethalometer (Moosmuller et al., 1998). Neither the aethalometer nor the photoacoustic spectrometer measure elemental carbon mass directly. Because the photoacoustic spectrometer measures the absorption coefficient directly, the specific absorption efficiency must be known or assumed in order to determine elemental carbon mass. Assuming a light absorption efficiency of 10 m^2 g^{-1}, Arnott et al. (1999) reported a lower detection limit for light absorption of 0.4 M m^{-1} corresponding to a mass concentration of elemental carbon of approximately 40 ng^{-3}.

2.2.9.3 Continuous Measurements of Nitrate and Sulfate

Nitrate

An integrated collection and vaporization cell was developed by Stolzenburg and Hering (2000) that provides automated, 10-min resolution monitoring of fine particulate nitrate. In this system, particles are collected by a humidified impaction process and analyzed in place by flash

vaporization and chemiluminescent detection of the evolved nitrogen oxides. In field tests in which the system was collocated with two FRM samplers, the automated nitrate sampler results followed the results from the FRM, but were offset lower. The system also was collocated with a HEADS and a SASS speciation sampler (MetOne Instruments). In all these tests, the automated sampler was well correlated to other samplers with slopes near 1 (ranging from 0.95 for the FRM to 1.06 for the HEADS) and correlation coefficients ranging from 0.94 to 0.996.

During the Northern Front Range Air Quality Study in Colorado (Watson et al., 1998), the automated nitrate monitor captured the 12-minute variability in fine particle nitrate concentrations with a precision of approximately ±0.5 $\mu g/m^3$ (Chow et al., 1998). A comparison with denuded filter measurements followed by ion chromatographic analysis (Chow and Watson, 1999) showed agreement within ±0.6 $\mu g/m^3$ for most of the measurements, but exhibited a discrepancy of a factor of two for the elevated nitrate periods. More recent intercomparisons took place during the 1997 Southern California Ozone Study (SCOS97) in Riverside, CA. Comparisons with 14 days of 24-hour denuder-filter sampling gave a correlation coefficient of $R^2 = 0.87$ and showed no significant bias (i.e., the regression slope is not significantly different from 1). As currently configured, the system has a detection limit of 0.7 $\mu g/m^3$ and a precision of 0.2 $\mu g/m^3$.

Sulfate

Continuous methods for the quantification of aerosol sulfur compounds first remove gaseous sulfur (e.g., SO_2, H_2S) from the sample stream by a diffusion tube denuder followed by the analysis of particulate sulfur (Cobourn et al., 1978; Durham et al., 1978; Huntzicker et al., 1978; Mueller and Collins, 1980; Tanner et al., 1980). Another approach is to measure total sulfur and gaseous sulfur separately by alternately removing particles from the sample stream. Particulate sulfur is obtained as the difference between the total and gaseous sulfur (Kittelson et al., 1978). The total sulfur content is measured by a flame photometric detector (FPD) by introducing the sampling stream into a fuel-rich hydrogen-air flame (e.g., Stevens et al., 1969; Farwell and Rasmussen, 1976) that reduces sulfur compounds and measures the intensity of the chemiluminescence from electronically excited sulfur molecules (S_2*).

Because formation of S_2* requires two sulfur atoms, the intensity of the chemiluminescence is theoretically proportional to the square of the concentration of molecules that contain a single

sulfur atom. In practice, the exponent is between one and two and depends on the sulfur compound being analyzed (Dagnall et al., 1967; Stevens et al., 1971). Calibrations are performed using both particles and gases as standards. The FPD can also be replaced by a chemiluminescent reaction with ozone that minimizes the potential for interference and provides a faster response time (Benner and Stedman, 1989, 1990).

Capabilities added to the basic system include *in situ* thermal analysis and sulfuric acid speciation (Cobourn et al., 1978; Huntzicker et al., 1978; Tanner et al., 1980; Cobourn and Husar, 1982)). Sensitivities for particulate sulfur as low as 0.1 $\mu g/m^3$, with time resolution ranging from 1 to 30 min, have been reported. Continuous measurements of particulate sulfur content have also been obtained by on-line x-ray fluorescence analysis with resolution of 30 min or less (Jaklevic et al., 1981b). During a field-intercomparison study of five different sulfur instruments, Camp et al. (1982) reported four out of five FPD systems agreed to within ±5% during a one-week sampling period.

2.2.9.4 Continuous Ion Chromatography of Water-Soluble Ions

Dasgupta and Slanina have independently developed particle collection systems that grow particles by increasing the relative humidity and collect the particles in an aqueous solution suitable for injection into an ion chromatography (Simon and Dasgupta, 1995; Khlystov et al., 1995). Automation of these systems yield semi-continuous monitors for those ions that can be determined by ion chromatography. A similar system suing a particle size magnifier has been reported by Weber et al. (2001).

2.2.9.5 Measurements of Individual Particles

Recently, several researchers have developed instruments for real-time *in situ* analysis of single particles (e.g., Noble and Prather, 1996; Gard et al., 1997; Johnson and Wexler, 1995; Silva and Prather, 1997; Thomson and Murphy, 1994). Although the technique varies from one laboratory to another, the underlying principle is to fragment each particle into ions, using either a high-power laser or a heated surface and, then, a time-of-flight mass spectrometer (TOFMS) to measure the ion fragments in a vacuum. Each particle is analyzed in a suspended state in the air stream (i.e., without collection), avoiding sampling artifacts associated with impactors and filters. The technique is called aerosol time-of-flight mass spectrometry (ATOFMS). By measuring both

positive and negative ions from the same particle, information can be obtained about the chemical composition, not just the elemental composition, of individual particles of known aerodynamic diameter. This information is especially useful in determining sources of particles. Noble and Prather (1996) used ATOFMS to provide compositionally resolved particle-size distributions. Their instrument is capable of analyzing size and chemical composition of 50 to 100 particles/min at typical ambient concentrations and up to 600/min at high particle concentrations. An example of the type of information that can be determined is shown in Figure 2-20.



Figure 2-20. Size distribution of particles divided by chemical classification into organic, marine, and crustal.

Source: Noble and Prather (1998).

Because particles are analyzed individually, biases in particle sampling (the efficiency of particle transmission into the sensor chamber as a function of size; particle size measurement, and detection of particles prior to fragmentation) represent a major challenge for these

instruments. Moreover, the mass spectrometer has a relatively large variability in ion yields (i.e., identical samples would yield relatively large differences in mass spectrometer signals [Thomson and Murphy, 1994]); therefore, quantitation is inherently difficult (Murphy and Thomson, 1997). Quantitation will be even more challenging for complex organic mixtures because of the following two reasons: (1) a large number of fragments are generated from each molecule, and (2) ion peaks for organics can be influenced or obscured by inorganic ions (Middlebrook et al., 1998). Nonetheless, scientists have been successful in using these techniques to identify the presence of organics in atmospheric particles and laboratory-generated particles (i.e., as contaminants in laboratory-generated sulfuric acid droplets) as well as the identification of specific compound classes such as PAHs in combustion emissions (Castaldi and Senkan, 1998; Hinz et al., 1994; Middlebrook et al., 1998; Murphy and Thomson, 1997; Neubauer et al., 1998; Noble and Prather, 1998; Reilly et al., 1998; Silva and Prather, 1997). A new multivariate technique for calibration of ATOFMS using microorifice impactors shows promise for simplifying the calibration process (Fergenson et al., 2001). This calibration technique has been applied to gasoline and diesel particles to demonstrate the feasibility of using this technique for the source apportionment of gasoline and diesel particles in an atmospheric mixture (Song et al., 2001).

Until recently, ATOFMS systems have only been able to characterize particles that are larger than approximately 0.2 to 0.3 μm in diameter. Wexler and colleagues (Carson et al., 1997; Ge et al., 1998) have developed an ATOFMS instrument that is able to size, count, and provide chemical composition on individual particles ranging in size from 10 nm to 2 μm.

2.2.9.6 Determination of Aerosol Surface Area in Real Time

Aerosol surface area is an important aerosol property for health effects research. However, methods for on-line measurement of surface area are not widely available. Woo et al. (2001b) used three continuous aerosol sensors to determine aerosol surface area. They used a condensation particle counter (CPC, TSI, Inc., Model 3020), an aerosol mass concentration monitor (MCM, TSI, Inc., Model 8520), and an electrical aerosol detector (EAD, TSI, Inc., Model 3070) for measuring particle charge concentration. The three sensor signals were inverted to obtain the aerosol size distribution, using a log-normal size distribution model (by minimizing the difference between the measured signals and the theoretical values based upon a size

distribution model, the instrument calibration, and its theoretical responses). The log-normal function was then integrated to calculate the total surface area concentration. Woo et al. (2001b) demonstrated that this method can give near real-time measurements of aerosol surface area.

2.2.9.7 Light Scattering

A variety of types of nephelometers that integrate aerosol light scattering over various solid angles are available (McMurry, 2000). When used to measure visibility, e.g., to provide pilots with realtime data on visual range, it is desirable to include the light scattering due to particle-bound water. However, when used as an indicator of fine particle mass, it is desirable to exclude particle-bound water. This is frequently done by heating the ambient aerosol to a low reference relative humidity of 40%. However, this heating has the potential of also causing the loss of semivolatile components of the aerosol. The evaporation of ammonium nitrate aerosol in a heated nephelometer has been examined. Bergin et al. (1997) conducted laboratory experiments at low relative humidity (~10%) and as a function of temperature (27-47 °C), mean residence time in the nephelometer, and initial particle size distribution. The evaporation of ammonium nitrate aerosol was also modeled for comparison and was found to describe accurately the decrease in aerosol scattering coefficient as a function of aerosol physical properties and nephelometer operating conditions. Bergin et al. (1997) determined an upper limit estimate of the decrease in the aerosol light scattering coefficient at 450 nm due to evaporation for typical field conditions. The model estimates for their worst-case scenario suggest that the decrease in the aerosol scattering coefficient could be roughly 40%. Under most conditions, however, they estimate that the decrease in aerosol scattering coefficient is generally expected to be less than 20%.

2.2.10 Low Flow Filter Samples for Multiday Collection of Particulate Matter

For some purposes, such as demonstrating attainment of an annual standard or as an exposure indicator for epidemiologic studies of chronic health effects, 24-h measurements are not essential. Annual or seasonal averages may be adequate. Multiday sampling techniques can result in lower costs for weighing, chemical analysis, and travel time to change filters. The multiday sampler serves a second purpose. Most commercially available samplers are optimized

for collecting 24-h samples of the PM concentrations found in the U.S., Europe, or Japan. Many cities in other parts of the world have significantly higher PM concentrations. Under these conditions, the 16.7 L/min flow through 37 or 47 mm diameter filters may overload the filter and prevent the sampler from maintaining the prescribed flow rate for 24 h. A low flow sampler with a 0.4 L/min flow rate and a 47 mm diameter filter has been designed by Aerosol Dynamics, Inc. With this sampler, the sample collection time can be chosen to suit the ambient concentration level. This sampler, with a one-week collection period, has been used to characterize $PM_{2.5}$ in Beijing, PRC (He et al., 2001). With a two-week collection period, it is being used in a chronic epidemiologic study in southern California, USA (Gauderman, et al., 2000).

The sampler, as described by He et al. (2001), has three $PM_{2.5}$ channels. One channel collects PM on a Teflon filter for gravimetric mass measurement and elemental analysis by XRF. A second channel collects PM on a quartz filter for organic and elemental carbon analysis. A denuder to remove organic gases and a backup filter to collect semivolatile organic compounds may be added. The third channel uses a carbonate denuder to remove acid gases (HNO_3 and SO_2), a Teflon filter to collect PM for analysis of ions by ion chromatography, and a nylon filter to collect volatilized nitrate. The Teflon filter can also be weighed prior to extraction. Thus, the multiday sampler can provide the information needed for source apportionment by Chemical Element Balance techniques (Watson et al., 1990a,b; U.S. Environmental Protection Agency, 2002b).

Since PM is commonly sampled on less than daily schedules, the magnitude of sampling errors needs to be considered when quality issues are of concern. For monitoring sites with high day-to-day variability in PM concentrations, an integrated sample may provide a more accurate measurement of the annual average than can be obtained by less-than-everyday sampling schedules. Daily PM data from Spokane, WA were resampled to simulate common sampling schedules, and the sampling error was computed for regulatory and distribution statistics (Rumburg et al., 2001). Probability density functions (pdf’s) were fit to the annual daily data to determine the shape of the $PM_{2.5}$ concentration distributions. Pdf’s were also fit to the less than daily sampling schedules to determine if pdf's could be used to predict the daily high-concentration percentiles. There is an error when using a less than daily sampling schedule for all statistics. The error, expressed as a percentage difference from the everyday sampling, was as large as 1.7, 3.4, and 7.7% for the $PM_{2.5}$ mean for 1-in-2 day, 1-in-3 day, and 1-in-6 day

sampling, respectively. For the 98th percentile, the error was as great as 8.8, 18, and 67% for 1-in-2 day, 1-in-3 day, and 1-in-6 day sampling, respectively.

2.3 SUMMARY

2.3.1 Atmospheric Physics and Chemistry of Particles

Atmospheric particles originate from a variety of sources and possess a range of morphological, chemical, physical, and thermodynamic properties. The composition and behavior of aerosols are linked with those of the surrounding gas. Aerosol may be defined as a suspension of solid or liquid particles in air and includes both the particles and all vapor or gas phase components of air. However, the term aerosol often is used to refer to the suspended particles only.

A complete description of the atmospheric aerosol would include an accounting of the chemical composition, morphology, and size of each particle, and the relative abundance of each particle type as a function of particle size. Recent developments in single particle analysis techniques are bringing such a description closer to reality.

The diameter of a spherical particle may be determined geometrically, from optical or electron microscopy, by light scattering and Mie theory, or by a particle's behavior (e.g., electrical mobility or its aerodynamic behavior). However, the various types of diameters may be different, and atmospheric particles often are not spherical. Therefore, particle diameters are described by an "equivalent" diameter. Aerodynamic diameter (i.e., the diameter of a unit density sphere that would have the same terminal settling velocity as the real particle, symbol, D_a) is the most widely used equivalent diameter.

Atmospheric size distributions show that most atmospheric particles are quite small, below 0.1 μm; whereas most of the particle volume (and therefore most of the mass) is found in particles greater than 0.1 μm. An important feature of the mass or volume size distributions of atmospheric particles is their multimodal nature. Volume distributions, measured in ambient air in the United States, are almost always found to be bimodal with a minimum between 1.0 and 3.0 μm. The distribution of particles that are mostly larger than the minimum is termed the "coarse" mode. The distribution of particles that are mostly smaller than the minimum is termed

the “fine” mode. Fine-mode particles include both the accumulation mode and the nuclei mode. “Accumulation-mode” particles are that portion of the fine particle fraction with diameters above about 0.1 μm. The nuclei mode, that portion of the fine particle fraction with diameters below about 0.1 μm, can be observed as a separate mode in mass or volume distributions only in clean or remote areas or near sources of new particle formation by nucleation. Toxicologists and epidemiologists use “ultrafine” to refer to particles in the nuclei-mode size range. Aerosol physicists and material scientists tend to use “nanoparticles” to refer to particles in this size range.

The aerosol community uses four different approaches or conventions in the classification of particles by size: (1) modes, based on the observed size distributions and formation mechanisms; (2) cut point, usually based on the 50% cut point of the specific sampling device (i.e., the particle size at which 50% of the particles enter and 50% of the particles are rejected); (3) dosimetry or occupational sizes, based on the entrance into various compartments of the respiratory system; and (4) legally specified, regulatory sizes for air quality standards. Over the years, the terms fine and coarse as applied to particle sizes have lost the original precise meaning of fine mode and coarse mode. In any given article, therefore, the meaning of fine and coarse, unless defined, must be inferred from the author’s usage. In particular, $PM_{2.5}$ and fine-mode particles are not equivalent. In this document, the term “mode” is used with fine and coarse when it is desired to specify the distribution of fine-mode particles or coarse-mode particles as shown in Figures 2-4 and 2-5.

Several processes influence the formation and growth of particles. New particles may be formed by nucleation from gas phase material. Particles may grow by condensation as gas phase material condenses onto existing particles. Particles may also grow by coagulation as two particles combine to form one. Gas phase material condenses preferentially on smaller particles, and the rate constant for coagulation of two particles decreases as the particle size increases. Therefore, nuclei mode particles grow into the accumulation mode, but growth of accumulation mode particles into the coarse mode is rare.

The major constituents of atmospheric PM are sulfate, nitrate, ammonium, and hydrogen ions; particle-bound water; elemental carbon; a great variety of organic compounds; and crustal material. Atmospheric PM contains a large number of elements in various compounds and concentrations and hundreds of specific organic compounds. Particulate material can be primary

or secondary. PM is called primary if it is in the same chemical form in which it was emitted into the atmosphere. PM is called secondary if it is formed by chemical reactions in the atmosphere. Primary coarse particles are usually formed by mechanical processes; whereas primary fine particles are emitted from sources either directly as particles or as vapors that rapidly condense to form particles.

Most of the sulfate and nitrate and a portion of the organic compounds in atmospheric particles are secondary. Secondary aerosol formation depends on numerous factors including the concentrations of precursors; the concentrations of other gaseous reactive species such as ozone, hydroxyl radical, peroxy radicals, and hydrogen peroxide; atmospheric conditions, including solar radiation and relative humidity; and the interactions of precursors and preexisting particles within cloud or fog droplets or on or in the liquid film on solid particles. As a result, it is considerably more difficult to relate ambient concentrations of secondary species to sources of precursor emissions than it is to identify the sources of primary particles.

The lifetimes of particles vary with particle size. Coarse particles can settle rapidly from the atmosphere within minutes or hours and normally travel only short distances. However, when mixed high into the atmosphere, as in dust storms, the smaller-sized, coarse-mode particles may have longer lives and travel greater distances. Accumulation-mode fine particles are kept suspended by normal air motions and have a lower deposition velocity than coarse-mode particles. They can be transported thousands of kilometers and remain in the atmosphere for a number of days. Accumulation-mode particles are removed from the atmosphere primarily by cloud processes. Dry deposition rates are expressed in terms of a deposition velocity that varies with the particle size, reaching a minimum between 0.1 and 1.0 μm aerodynamic diameter.

PM is a factor in acid deposition. Particles serve as cloud condensation nuclei and contribute directly to the acidification of rain. In addition, the gas-phase species that lead to dry deposition of acidity are also precursors of particles. Therefore, reductions in SO_2 and NO_x emissions will decrease both acid deposition and PM concentrations. Sulfuric acid, ammonium nitrate, and organic particles also are deposited on surfaces by dry deposition and can contribute to ecological damage.

Particles also reduce visibility and affect radiative balance through scattering and absorption of light. The direct effects of particles in scattering and absorbing light and the indirect effects of particles on clouds impact climate change processes.

2.3.2 Measurement of Atmospheric Particles

The decision by the EPA to revise the PM standards by adding daily and yearly standards for $PM_{2.5}$ has led to a renewed interest in the measurement of atmospheric particles and to a better understanding of the problems in developing precise and accurate measurements of particles. Unfortunately, it is very difficult to measure and characterize particles suspended in the atmosphere.

PM monitoring is used to develop information to guide implementation of standards (i.e., to identify sources of particles; to determine whether or not a standard has been attained; and to determine health, ecological, and radiative effects). Federal Reference Methods (FRM) specify techniques for measuring PM_{10} and $PM_{2.5}$. Particles are collected on filters and mass concentrations are determined gravimetrically. The PM_{10} FRM sampler consists of a PM_{10} inlet/impactor and a 47-mm Teflon filter with a particle collection efficiency greater than 99.7%. The $PM_{2.5}$ FRM is similar except that it includes a $PM_{2.5}$ impactor with an oil-covered impaction substrate to remove particles larger than 2.5 μm. Both techniques provide relatively precise (±10%) methods for determining the mass of material remaining on a Teflon filter after equilibration. Despite considerable progress in measuring the atmospheric PM mass concentration, numerous uncertainties continue to exist as to the relationship between the mass and composition of material remaining on the filter as measured by the FRMs and the mass and composition of material that exists in the atmosphere as suspended PM. There is no reference standard for particles suspended in the atmosphere, nor is there an accepted way to remove particle-bound water without losing some of the semivolatile components of PM such as ammonium nitrate and semivolatile organic compounds. It also is difficult to cleanly separate fine-mode and coarse-mode PM. As a result, EPA defines accuracy for PM measurements in terms of agreement of a candidate sampler with a reference sampler. Therefore, intercomparisons of samplers become very important in determining how well various samplers agree and how various design choices influence what is actually measured.

Current filtration-based mass measurements lead to significant evaporative losses of a variety of semivolatile components (i.e., species that exist in the atmosphere in dynamic equilibrium between the condensed phase and gas phase) during and possibly after collection. Important examples include ammonium nitrate, semivolatile organic compounds, and particle-bound water. Loss of these components may significantly affect the quality of the measurement

and can lead to both positive and negative sampling artifacts. Negative artifacts resulting from loss of ammonium nitrate and semivolatile organic compounds may occur during sampling because of changes in temperature, relative humidity, or composition of the aerosol or because of the pressure drop across the filter. Negative artifacts also may occur during handling and storage because of evaporation. Positive artifacts occur when gas-phase compounds (H_2O, HNO_3, SO_2, and organic compounds) absorb onto or react with filter media or collected PM or when some particle-bound water is not removed.

Sampling systems for semivolatile PM components make use of denuders to remove the gas-phase fraction and absorptive filters to remove the condensed phase and retain any material that subsequently evaporates from the collected PM. The loss of particulate nitrate may be determined by comparing nitrate collected on a Teflon filter to that collected on a nylon filter (which absorbs nitric acid which evaporates from ammonium nitrate particles) preceded by a denuder to remove gas-phase nitric acid. In two studies in southern California, the $PM_{2.5}$ mass lost because of volatilization of ammonium nitrate was found to represent 10 to 20% of the total $PM_{2.5}$ mass and almost a third of the nitrate. Denuder/absorptive filter sampling systems also have been developed for measuring particulate phase organic compounds. This technique is an improvement over the filter/adsorbent collection method. However, the denuder systems currently discussed in the literature are not straightforward in their use, and the user must have a thorough understanding of the technology. The FRM for $PM_{2.5}$ will likely suffer loss of particulate nitrates and semivolatile organic compounds, similar to the losses experienced with other single filter collection systems.

It is generally desirable to collect and measure ammonium nitrate and semivolatile organic compounds as part of particulate matter mass. However, it is usually desirable to remove the particle-bound water before determining the mass. In some situations, it may be important to know how much of the suspended particle's mass or volume results from particle-bound water. Calculations and measurements indicate that aerosol water content is strongly dependent on relative humidity and composition. Particle-bound water can represent a significant mass fraction of the PM concentration at relative humidities above 60%. A substantial fraction of accumulation-mode PM is hygroscopic or deliquescent. The more hygroscopic particles tend to contain more sulfates, nitrates, and secondary organic compounds, while the less hygroscopic particles tend to contain more elemental carbon, primary organic compounds, and crustal

components. Fresh, submicron-size soot particles may tend to shrink with increasing relative humidity because of a structural change. The effects of relative humidity on the sorption of semivolatile organic compounds on particles are not well understood. The amount of water sorbed to an atmospheric aerosol may be affected by the presence of an organic film on the particle, which may impede the transport of water across the surface.

Fine-mode and coarse-mode particles differ not only in size, but also in formation mechanisms; sources; and chemical, physical, and biological properties. Fine and coarse particles overlap in the intermodal size range (1-2.5 μm D_a). As relative humidity increases, fine particles grow into this size range; as relative humidity decreases, more coarse particles may be suspended in this size range. It is desirable to measure fine-mode PM and coarse-mode PM separately in order to properly allocate health effects to either fine-mode PM or coarse-mode PM and to correctly determine sources by factor analysis or chemical mass balance. The selection of a cut point of 2.5 um as a basis for EPA's 1997 NAAQS for fine particles (Federal Register, 1997) and its continued use in many health effects studies reflects the importance placed on more complete inclusion of fine-mode particles, while recognizing that intrusion of coarse-mode particles can occur under some conditions with this cut point.

In addition to FRM sampling of equilibrated mass to determine compliance with PM standards, EPA requires states to conduct speciation sampling primarily to determine source categories and trends. The current speciation samplers collect $PM_{2.5}$ on three filters: (1) a Teflon filter for gravimetric determination of mass and for analysis of heavy elements by X-ray fluorescence; (2) a Nylon filter preceded by a nitric acid denuder for artifact-free determination of nitrate and measurement of other ionic species by ion chromatography; and (3) a quartz filter for measurement of elemental carbon (EC) and organic carbon (OC). In addition, IMPROVE (Interagency Monitoring of Protected Visual Environments) samplers provide information on regional PM background and transport. IMPROVE samplers, in addition to the three types of filters collected by the speciation samplers, also collect a PM_{10} sample. The IMPROVE and speciation networks use slightly different methods for determination of EC and OC. The two methods agree on total carbon but differ in the split of total carbon into EC and OC. Neither EC/OC method provides for any correction for positive or negative artifacts because of absorption of volatile organic compounds on the quartz filters or evaporation of semivolatile organic compounds from the collected particles.

The EPA expects that monitoring agencies will operate continuous PM monitors. EPA is in the process of providing guidance regarding appropriate continuous monitoring techniques. All currently available techniques for continuous measurements of suspended particle mass such as the integrating nephelometer, the beta-absorption monitor, and the Tapered Element Oscillating Microbalance (TEOM) share the problem of dealing with semivolatile PM components (in order not to include particle-bound water as part of the mass, the particle-bound water must be removed by heating or dehumidification). However, heating also causes ammonium nitrate and semivolatile organic compounds to evaporate. The TEOM monitor operates at a constant, but higher than ambient, temperature to remove particle-bound water. However, the FRM is required to operate at no more than 5 °C above the ambient temperature. Subsequently, much of the particle-bound water is removed during equilibration at 40% relative humidity. This difference in techniques for removal of particle-bound water causes differences in the measured mass concentration between TEOM and FRMs.

Several new techniques for continuous PM mass measurements are currently being field tested. The Real-Time Total Ambient Mass Sampler (RAMS) measures the total mass of collected particles including semivolatile species with a TEOM monitor using a "sandwich filter." The sandwich contains a Teflon-coated particle-collection filter followed by a charcoal-impregnated filter to collect any semivolatile species lost from the particles during sampling. The RAMS uses a Nafion dryer to remove particle-bound water from the suspended particles and a particle concentrator to reduce the quantity of gas phase organic compounds that must be removed by the denuder. The Continuous Ambient Mass Monitor (CAMM) estimates ambient particulate matter mass by measurement of the increase in the pressure drop across a membrane filter caused by particle loading. It also uses a Nafion dryer to remove particle-bound water. In addition to continuous mass measurement, a number of techniques for continuous measurement of sulfate, nitrate, or elements are being tested.

REFERENCES

Aiginger, H. (1991) Historical development and principles of total reflection X-ray fluorescence analysis (TXRF). Spectrochim. Acta 46B: 1313-1321.

Aiginger, H.; Streli, C. (1997) Total reflection X-ray fluorescence spectrometry (TXRF) for trace element analysis. Spectrosc. Eur. 9: 16-21.

Allen, G.; Sioutas, C.; Koutrakis, P.; Reiss, R.; Lurmann, F. W.; Roberts, P. T. (1997) Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas. J. Air Waste Manage. Assoc. 47: 682-689.

Allen, J. O.; Paschkewitz, J. S.; Plummer, E. F.; Lafleur, A. L.; Sarofim, A. F.; Smith, K. A. (1999a) Absorption of semi-volatile compounds in oiled impaction substrates: measurement of pyrene absorption. Aerosol Sci. Technol. 30: 16-29.

Allen, G. A.; Oh, J. A.; Koutrakis, P. (1999b) Techniques for high quality ambient coarse particulate mass measurements. J. Air Waste Manage. Assoc. 49: 133-141.

Allen, G. A.; Lawrence, J.; Koutrakis, P. (1999c) Field validation of a semi-continuous method for aerosol black carbon (aethalometer) and temporal patterns of summertime hourly black carbon measurements in southwestern PA. Atmos. Environ. 33: 817-823.

Allen, R.; Box, M.; Larson, T. V.; Liu, L. J. (2001) A cost-effective weighing chamber for particulate matter filters. J. Air Waste Manage. Assoc. 51: 1650-1653.

American Conference of Governmental Industrial Hygienists. (1994) 1994-1995 threshold limit values for chemical substances and physical agents and biological exposure indices. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.

Andrews, E.; Larson, S. M. (1993) Effect of surfactant layers on the size changes of aerosol particles as a function of relative humidity. Environ. Sci. Technol. 27: 857-865.

Appel, B. R. (1993) Atmospheric sample analysis and sampling artifacts. In: Willeke, K.; Baron, P. A., eds. Aerosol measurement: principles, techniques, and applications. New York, NY: Van Nostrand Reinhold; pp. 233-259.

Appel, B. R.; Tokiwa, Y.; Haik, M. (1981) Sampling of nitrates in ambient air. Atmos. Environ. 15: 283-289.

Appel, B. R.; Tokiwa, Y.; Haik, M.; Kothny, E. L. (1984) Artifact particulate sulfate and nitrate formation on filter media. Atmos. Environ. 18: 409-416.

Arnott, W. P.; Moosmüller, H.; Rogers, C. F.; Jin, T.; Bruch, R. (1999) Photoacoustic spectrometer for measuring light absorption by aerosol; instrument description. Atmos. Environ. 33: 2845-2852.

Arnott, W. P.; Moosmüller, H.; Walker, J. W. (2000) Nitrogen dioxide and kerosene-flame soot calibration of photoacoustic instruments for measurement of light absorption by aerosols. Rev. Sci. Instruments 71: 4545-4552.

Ayers, G. P.; Keywood, M. D.; Gras, J. L. (1999) TEOM vs. manual gravimetric methods for determination of $PM_{2.5}$ aerosol mass concentrations. Atmos. Environ. 33: 3717-3721.

Babich, P.; Davey, M.; Allen, G.; Koutrakis, P. (2000) Method comparisons for particulate nitrate, elemental carbon, and $PM_{2.5}$ mass in seven U.S. cities. J. Air Waste Manage. Assoc. 50: 1095-1105.

Bacon, J. R.; Chenery, S. R. N.; Ellis, A. T.; Fisher, A.; McMahon, A. W.; Potts, P. J.; Williams, J. G. (1995) Atomic spectrometry update–atomic mass spectrometry and x-ray fluorescence spectrometry. J. Anal. At. Spectrom. 10: 253R-310R.

Baldwin, D. P.; Zamzow, D. S.; D'Silva, A. P. (1994) Aerosol mass measurement and solution standard additions for quantitation in laser ablation-inductively coupled plasma atomic emission spectrometry. Anal. Chem. 66: 1911-1917.

Ballach, J.; Hitzenberger, R.; Schultz, E.; Jaeschke, W. (2001) Development of an improved optical transmission technique for black carbon (BC) analysis. Atmos. Environ. 35: 2089-2100.

Baron, P. A.; Willeke, K., eds. (2001) Aerosol measurement: principles, techniques, and applications. New York, NY: John Wiley & Sons.

Benner, R. L.; Stedman, D. H. (1989) Universal sulfur detection by chemiluminescence. Anal. Chem. 61: 1268-1271.

Benner, R. L.; Stedman, D. H. (1990) Field evaluation of the sulfur chemiluminescence detector. Environ. Sci. Technol. 24: 1592-1596.

Bergin, M. H.; Ogren, J. A.; Schwartz, S. E.; McInnes, L. M. (1997) Evaporation of ammonium nitrate aerosol in a heated nephelometer: implications for field measurements. Environ. Sci. Technol. 31: 2878-2883.

Bertoni, G.; Febo, A.; Perrino, C.; Possanzini, M. (1984) Annular active diffuse sampler: a new device for the collection of organic vapours. Ann. Chim. (Rome) 74: 97-104.

Billiet, J.; Dams, R.; Hoste, J. (1980) Multielement thin film standards for XRF analysis. X-Ray Spectr. 9: 206-211.

Binkowski, F. S.; Ching, J. K. S. (1995) Modeling fine particulate mass and visibility using the EPA regional particulate model. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development, National Exposure Research Laboratory; report no. EPA/600/A-95/143. Available from: NTIS, Springfield, VA; PB96-139100.

Binkowski, F. S.; Shankar, U. (1995) The regional particulate matter model. I. Model description and preliminary results. J. Geophys. Res. [Atmos.] 100: 26,191-26,209.

Birch, M. E. (1998) Analysis of carbonaceous aerosols: interlaboratory comparison. Analyst 123: 851-857.

Birch, M. E.; Cary, R. A. (1996) Elemental carbon-based method for occupational monitoring of particulate diesel exhaust: methodology and exposure issues. Analyst 121: 1183-1190.

Bowers, W. D.; Chuan, R. L. (1989) Surface acoustic-wave piezoelectric crystal aerosol mass monitor. Rev. Sci. Instrum. 60: 1297-1302.

Breum, N. O. (2000) The dust holding capacity of porous plastic foam used in particle size-selective sampling. J. Aerosol Sci. 31: 379-385.

Brook, J. R.; Dann, T. F. (1999) Contribution of nitrate and carbonaceous species to $PM_{2.5}$ observed in Canadian cities. J. Air Waste Manage. Assoc. 49: 193-199.

Brorström, E.; Grennfelt, P.; Lindskog, A. (1983) The effect of nitrogen dioxide and ozone on the decomposition of particle-associated polycyclic aromatic hydrocarbons during sampling from the atmosphere. Atmos. Environ. 17: 601-605.

Bunz, H.; Koyro, M.; Möhler, O.; Saathoff, H. (1996) Temperature dependence of the redistribution mechanisms for volatile components of atmospheric aerosol particles. J. Aerosol Sci. 27(suppl. 1): S653-S654.

Cachier, H.; Bremond, M.-P.; Buat-Ménard, P. (1989) Determination of atmospheric soot carbon with a simple thermal method. Tellus 41B: 379-390.

Cadle, S. H.; Groblicki, P. J. (1982) An evaluation of methods for the determination of organic and elemental carbon in particulate samples. In: Wolff, G. T.; Klimisch, R. L., eds. Particulate carbon: atmospheric life cycle. New York, NY: Plenum Press; pp. 89-109.

Cadle, S. H.; Mulawa, P.A. (1990) Atmospheric carbonaceous species measurement methods comparison study: General Motors results. Aerosol Sci. Technol. 12: 128-131.

Cadle, S. H.; Groblicki, P. J.; Mulawa, P. A. (1983) Problems in the sampling and analysis of carbon particles. Atmos. Environ. 17: 593-600.

Cahill, T. A. (1981) Innovative aerosol sampling devices based upon PIXE capabilities. Nucl. Instr. Meth. Phys. Res. 181: 473-480.

Cahill, T. A. (1985) Proton-induced x-ray emission (PIXE). In: Metals Handbook. Cleveland, OH: American Society for Metals.

Cahill, T. A.; Eldred, R. A.; Motallebi, N.; Malm, W. C. (1989) Indirect measurement of hydrocarbon aerosols across the United States by nonsulfate hydrogen-remaining gravimetric mass correlations. Aerosol Sci. Technol. 10: 421-429.

Cahill, T. A.; Wilkinson, K.; Schnell, R. (1992) Composition analyses of size-resolved aerosol samples taken from aircraft downwind of Kuwait, spring 1991. J. Geophys. Res. [Atmos.] 97: 14,513-14,520.

Cahill, T. A.; Wakabayashi, P.; James, T. A. (1996) Chemical states of sulfate at Shenandoah National Park during summer, 1991. Nucl. Instrum. Methods Phys. Res. B 109/110: 542-547.

Camp, D. C.; Stevens, R. K.; Cobourn, W. G.; Husar, R. B.; Collins, J. F.; Huntzicker, J. J.; Husar, J. D.; Jaklevic, J. M.; McKenzie, R. L.; Tanner, R. L.; Tesch, J. W. (1982) Intercomparison of concentration results from fine particle sulfur monitors. Atmos. Environ. 16: 911-916.

Campbell, A. J.; Humayun, M. (1999) Trace element microanalysis in iron meteorites by laser ablation ICPMS. Anal. Chem. 71: 939-946.

Carson, P. G.; Johnston, M. V.; Wexler, A. S. (1997) Laser desorption/ionization of ultra fine aerosol particles. Rapid Commun. Mass Spectrom. 11: 993-996.

Castaldi, M. J.; Senkan, S. M. (1998) Real-time, ultrasensitive monitoring of air toxics by laser photoionization time-of-flight mass spectroscopy. J. Air Waste Manage. Assoc. 48: 77-81.

Chang, M.; Kim, S.; Sioutas, C. (1999) Experimental studies on particle impaction and bounce: effects of substrate design and material. Atmos. Environ. 33: 2313-2322.

Chen, B. T. (1993) Instrument calibration. In: Willeke, K.; Baron, P. A., eds. Aerosol measurement: principles, techniques, and applications. New York, NY: Van Nostrand Reinhold; pp. 493-520.

Cheng, Y.-S.; Yeh, H.-C. (1979) Particle bounce in cascade impactors. Environ. Sci. Technol. 13: 1392-1396.

Chow, J. C. (1995) Measurement methods to determine compliance with ambient air quality standards for suspended particles. J. Air Waste Manage. Assoc. 45: 320-382.

Chow, J. C.; Watson, J. G. (1998) Guideline on speciated particulate monitoring (Draft 3). Research Triangle Park, NC: U.S. Environmental Protection Agency; Office of Air Quality Planning and Standards. Available: http://www.epa.gov/ttn/amtic/files/ambient/pm25/spec/drispec.pdf (20 March 2002).

Chow, J. C.; Watson, J. G. (1999) Ion chromatography. In: Landsberger, S.; Creatchman, M., eds. Elemental Analysis of Airborne Particles. Newark, NJ: Gordon and Breach; pp. 97-137.

Chow, J. C.; Watson, J. G.; Pritchett, L. C.; Pierson, W. R.; Frazier, C. A.; Purcell, R. G. (1993) The DRI thermal/optical reflectance carbon analysis system: description, evaluation and applications in U.S. air quality studies. In: Lodge, J. P., Jr. Fourth international conference on carbonaceous particles in the atmosphere; April 1991; Vienna, Austria. Atmos. Environ. Part A 27: 1185-1201.

Chow, J. C.; Watson, J. G.; Fujita, E. M.; Lu, Z.; Lawson, D. R.; Ashbaugh, L. L. (1994a) Temporal and spatial variations of PM2.5 and PM10 aerosol in the Southern California Air Quality Study. Atmos. Environ. 28: 2061-2080.

Chow, J. C.; Fujita, E. M.; Watson, J. G.; Lu, Z.; Lawson, D. R.; Ashbaugh, L. L. (1994b) Evaluation of filter-based aerosol measurements during the 1987 Southern California Air Quality Study. Environ. Monit. Assess. 30: 49-80.

Chow, J. C.; Zielinska, B.; Watson, J. G.; Fujita, E. M.; Richards, H. W.; Neff, W. D.; Dietrich, D.; Hering, S. V. (1998) Northern Front Range Air Quality Study. Volume A: Ambient measurements. Fort Collins, CO: Colorado State University, Cooperative Institute for Research in the Atmosphere. Available: http://www.nfraqs.colostate.edu/dri.html [4 February 2002].

Chow, J. C.; Watson, J. G.; Hackett, E. I.; Stone, R. H.; Hinsvark, B. A. (2000a) General atomic absorption procedure for trace metals in airborne material collected on filters. In: Lodge, J. P., ed. Methods of Air Sampling and Analysis. 4th ed. Pittsburgh, PA: Air and Waste Management Association.

Chow, J. C.; Watson, J. G.; Crow, D.; Lowenthal, D. H.; Merrifield, T. (2000b) Comparison of IMPROVE and NIOSH carbon measurements. Aerosol Sci. Technol. 34: 23-34.

Claiborn, C. S.; Finn, D.; Larson, T. V.; Koenig, J. Q. (2000) Windblown dust contributes to high $PM_{2.5}$ concentrations. J. Air Waste Manage. Assoc. 50: 1440-1445.

Clark, W. E.; Whitby, K. T. (1975) Measurements of aerosols produced by the photochemical oxidation of SO_2 in air. J. Colloid Interface Sci. 51: 477-490.

Clarke, A. D. (1992) Atmospheric nuclei in the remote free-troposphere. J. Atmos. Chem. 14: 479-488.

Cobourn, W. G.; Husar, R. B. (1982) Diurnal and seasonal patterns of particulate sulfur and sulfuric acid in St. Louis, July 1977-June 1978. Atmos. Environ. 16: 1441-1450.

Cobourn, W. G.; Husar, R. B.; Husar, J. D. (1978) Continuous *in situ* monitoring of ambient particulate sulfur using flame photometry and thermal analysis. In: Husar, R. B.; Lodge, J. P., Jr.; Moore, D. J., eds. Sulfur in the atmosphere: proceedings of the international symposium; September 1977; Dubrovnik, Yugoslavia. Atmos. Environ. 12: 89-98.

Code of Federal Regulations. (2001a) Appendix J to Part 50--reference method for the determination of particulate matter as PM_{10} in the atmosphere. C. F. R. 40: Part 50.

Code of Federal Regulations. (2001b) Ambient air monitoring reference and equivalent methods. C. F. R. 40: Part 53.

Code of Federal Regulations. (2001c) Appendix L to Part 50--reference method for the determination of fine particulate matter as $PM_{2.5}$ in the atmosphere. C. F. R. 40: Part 50.

Code of Federal Regulations. (2001d) Ambient air quality surveillance for particulate matter: final rule. C. F. R. 40: Part 58.

Conner, T.L.; Norris, G.A.; Landis, M.S.; Williams, R.W. (2001) Individual particle analysis of indoor, outdoor, and community samples from the 1998 Baltimore particulate matter study. Atmos. Environ. 35: 3935-3946.

Cotham, W. E.; Bidleman, T. F. (1992) Laboratory investigations of the partitioning of organochlorine compounds between the gas phase and atmospheric aerosols on glass fiber filters. Environ. Sci. Technol. 26: 469-478.

Countess, R. J. (1990) Interlaboratory analyses of carbonaceous aerosol samples. Aerosol Sci. Technol. 12: 114-121.

Courtney, W. J.; Shaw, R. W.; Dzubay, T. G. (1982) Precision and accuracy of a ß gauge for aerosol mass determinations. Environ. Sci. Technol. 16: 236-239.

Coutant, R. W.; Callahan, P. J.; Chuang, J. C.; Lewis, R. G. (1992) Efficiency of silicone-grease-coated denuders for collection of polynuclear aromatic hydrocarbons. Atmos. Environ. Part A 26: 2831-2834.

Covert, D. S.; Kapustin, V. N.; Quinn, P. K.; Bates, T. S. (1992) New particle formation in the marine boundary layer. J. Geophys. Res. [Atmos.] 97: 20,581-20,589.

Cui, W.; Machir, J.; Lewis , L.; Eatough, D. J.; Eatough, N. L. (1997) Fine particulate organic material at Meadview during the project MOHAVE summer intensive study. J. Air Waste Manage. Assoc. 47: 357-369.

Cui, W.; Eatough, D. J.; Eatough, N. L. (1998) Fine particulate organic material in the Los Angeles Basin - I: assessment of the high-volume Brigham Young University organic sampling system, BIG BOSS. J. Air Waste Manage. Assoc. 48: 1024-1037.

Dagnall, R. M.; Thompson, K. C.; West, T. S. (1967) Molecular emission in cool flames I. Analyst 92: 506-512.

Dams, R.; Robbins, J. A.; Rahn, K. A.; Winchester, J. W. (1970) Nondestructive neutron activation analysis of air pollution particulates. Anal. Chem. 42: 861-867.

Dane, A. D.; Timmermans, P. A. M.; Van Sprang, H. A.; Buydens, L. M. C. (1996) A genetic algorithm for model-free x-ray fluorescence analysis of thin film. Anal. Chem. 68: 2419-2425.

Davis, D. W.; Reynolds, R. L.; Tsou, G. C.; Zafonte, L. (1977) Filter attenuation corrections for x-ray fluorescence analysis of atmospheric aerosols. Anal. Chem. 49: 1990-1993.

Davis, C. S.; Fellin, P.; Otson, R. (1987) A review of sampling methods for polyaromatic hydrocarbons in air. J. Air Pollut. Control Assoc. 37: 1397-1408.

De Boer, D. K. G.; Leenaers, A. J. G.; Van Den Hoogenhof, W. W. (1995) Glancing-incidence x-ray analysis of thin-layered materials: a review. X-ray Spectrom. 24: 91-102.

Ding, Y.; Lee, M. L.; Eatough, D. J. (1998a) The determination of total nitrite and N-nitroso compounds in atmospheric samples. Int. J. Environ. Anal. Chem. 69: 243-255.

Ding, Y.; Cui, W.; Lee, M. L.; Eatough, D. J. (1998b) Fine particulate *N*-nitroso and nitrite organic compounds in the atmosphere. Appl. Occup. Environ. Hyg. 13: 432-438.

Ding, Y.; Pang, Y.; Eatough, D. J.; Eatough, N. L.; Tanner, R. L. (2002) High volume diffusion denuder sampler for the routine monitoring of fine particulate matter: II. field evaluation of the PC BOSS. Aerosol Sci. Technol.: in press.

Dockery, D. W.; Pope, C. A., III; Xu, X.; Spengler, J. D.; Ware, J. H.; Fay, M. E.; Ferris, B. G., Jr.; Speizer, F. E. (1993) An association between air pollution and mortality in six U.S. cities. N. Engl. J. Med. 329: 1753-1759.

Durham, J. L.; Wilson, W. E.; Bailey, E. B. (1978) Application of an SO_2-denuder for continuous measurement of sulfur in submicrometric aerosols. Atmos. Environ. 12: 883-886.

Dzubay, T. G.; Stevens, R. K. (1975) Ambient air analysis with dichotomous sampler and X-ray fluorescence spectrometer. Environ. Sci. Technol. 9: 663-668.

Dzubay, T. G.; Hines, L. E.; Stevens, R. K. (1976) Particle bounce errors in cascade impactors. Atmos. Environ. 10: 229-234.

Eatough, D. J. (1999) BOSS, the *B*righam Young University *O*rganic *S*ampling *S*ystem: determination of particulate carbonaceous material using diffision denuder sampling technology. In: Lane, D., ed. Gas and particle phase measurements of atmospheric organic compounds. New York, NY: Gordon and Breach Science Publishers. (Advances in environmental, industrial and process control technologies: v. 2).

Eatough, D. J.; Wadsworth, A.; Eatough, D. A.; Crawford, J. W.; Hansen, L. D.; Lewis, E. A. (1993) A multiple-system, multi-channel diffusion denuder sampler for the determination of fine-particulate organic material in the atmosphere. Atmos. Environ. Part A 27: 1213-1219.

Eatough, D. J.; Tang, H.; Cui, W.; Machir, J. (1995) Determination of the size distribution and chemical composition of fine particulate semi-volatile organic material in urban environments using diffusion denuder technology. In: Phalen, R. F.; Bates, D. V., eds. Proceedings of the colloquium on particulate air pollution and human mortality and morbidity, part II; January 1994; Irvine, CA. Inhalation Toxicol. 7: 691-710.

Eatough, D. J.; Eatough, D. A.; Lewis, L.; Lewis, E. A. (1996) Fine particulate chemical composition and light extinction at Canyonlands National Park using organic particulate material concentrations obtained with a multisystem, multichannel diffusion denuder sampler. J. Geophys. Res. [Atmos.] 101: 19,515-19,531.

Eatough, D. J.; Obeidi, F.; Pang, Y.; Ding, Y.; Eatough, N. L.; Wilson, W. E. (1999a) Integrated and real-time diffusion denuder sampler for $PM_{2.5}$. Atmos. Environ. 33: 2835-2844.

Eatough, D. J.; Pang, Y.; Eatough, N. L. (1999b) Determination of $PM_{2.5}$ sulfate and nitrate with a PC-BOSS designed for routine sampling for semi-volatile particulate matter. J. Air Waste Manage. Assoc. 49: 69-75.

Fairchild, C. I.; Wheat, L. D. (1984) Calibration and evaluation of a real-time cascade impactor. Am. Ind. Hyg. Assoc. J. 45: 205-211.

Falconer, R. L.; Bidleman, T. F.; Cotham, W. E. (1995) Preferential sorption of non- and mono-ortho-polychlorinated biphenyls to urban aerosols. Environ. Sci. Technol. 29: 1666-1673.

Fang, C. P.; McMurry, P. H.; Marple, V. A.; Rubow, K. L. (1991) Effect of flow-induced relative humidity changes on size cuts for sulfuric acid droplets in the microorifice uniform deposit impactor (MOUDI). Aerosol Sci. Technol. 14: 266-277.

Farwell, S. O.; Rasmussen, R. A. (1976) Limitations of the FPD and ECD in atmospheric analysis: a review. J. Chromatogr. Sci. 14: 224-234.

Fassel, V. A.; Kniseley, R. N. (1974) Inductively coupled plasma–optical emission spectroscopy. Anal. Chem. 46: 1110A-1120A.

Federal Register. (1987) Revisions to the national ambient air quality standards for particulate matter. F. R. (July 1) 52: 24,634-24,669.

Federal Register. (1997) National ambient air quality standards for particulate matter; final rule. F. R. (July 18) 62: 38,652-38,752.

Federal Register. (1998) National ambient air quality standards for particulate matter and revised requirements for designation of reference and equivalent methods for $PM_{2.5}$ and ambient air quality surveillance for particulate matter; correction. F. R. (February 17) 63: 7710-7718.

Fergenson, D. P.; Song, X.-H.; Ramadan, Z.; Allen, J. O.; Hughes, L. S.; Cass, G. R.; Hopke, P. K.; Prather, K. A. (2001) Quantification of ATOFMS data by multivariate methods. Anal. Chem. 73: 3535-3541.

Fernandez, F. J. (1989) Atomic absorption spectroscopy. In: Lodge, J. P., Jr., ed. Methods of Air Sampling and Analysis. Chelsea, MI: Lewis Publishers; pp. 83-89.

Finn, D.; Rumburg, B.; Claiborn, C.; Bamesberger, L.; Siems, W. F.; Koenig, J.; Larson, T.; Norris, G. (2001) Sampling artifacts from the use of denuder tubes with glycerol based coatings in the measurement of atmospheric particulate matter. Environ. Sci. Technol. 35: 40-44.

Fitz, D. R. (1990) Reduction of the positive organic artifact on quartz filters. Aerosol. Sci. Technol. 12: 142-148.

Flocchini, R. G.; Cahill, T. A.; Shadoan, D. J.; Lange, S. J.; Eldred, R. A.; Feeney, P. J.; Wolfe, G. W.; Simmeroth, D. C.; Suder, J. K. (1976) Monitoring California's aerosols by size and elemental composition. Environ. Sci. Technol. 10: 76-82.

Friedlander, S. K. (1970) The characterization of aerosols distributed with respect to size and chemical composition. J. Aerosol Sci. 1: 295-307.

Friedlander, S. K. (1977) Smoke, dust and haze: fundamentals of aerosol behavior. New York, NY: John Wiley & Sons.

Friedlander, S. K.; Yeh, E. K. (1998) The submicron atmospheric aerosol as a carrier of reactive chemical species: case of peroxides. Appl. Occup. Environ. Hyg. 13: 416-420.

Fuchs, N. A. (1964) The mechanics of aerosols. New York, NY: Pergamon Press.

Fuchs, N. A. (1989) The mechanics of aerosols. New York, NY: Dover Publications.

Gallorini, M. (1995) Trace element monitoring in atmospheric pollution processes by neutron activation analysis. Microchem. J. 51: 127-137.

Gard, E.; Mayer, J. E.; Morrical, B. D.; Dienes, T.; Fergenson, D. P.; Prather, K. A. (1997) Real-time analysis of individual atmospheric aerosol particles: design and performance of a portable ATOFMS. Anal. Chem. 69: 4083-4091.

Gauderman, W. J.; McConnell, R.; Gilliland, F.; London, S.; Thomas, D.; Avol, E.; Vora, H.; Berhane, K.; Rappaport, E. B.; Lurmann, F.; Margolis, H. G.; Peters, J. (2000) Association between air pollution and lung function growth in southern California children. Am. J. Respir. Crit. Care Med. 162: 1383-1390.

Ge, Z.; Wexler, A. S.; Johnston, M. V. (1998) Laser desorption/ionization of single ultrafine multicomponent aerosols. Environ. Sci. Technol. 32: 3218-3223.

Goss, K.-U.; Eisenreich, S. J. (1997) Sorption of volatile organic compounds to particles from a combustion source at different temperatures and relative humidities. Atmos. Environ. 31: 2827-2834.

Gray, A. L.; Williams, J. G. (1987a) Oxide and doubly charged ion response of a commercial inductively coupled plasma mass spectrometry instrument. J. Anal. At. Spectrom. 2: 81-82.

Gray, A. L.; Williams, J. G. (1987b) System optimisation and the effect on polyatomic, oxide and doubly charged ion response of a commercial inductively coupled plasma mass spectrometry instrument. J. Anal. At. Spectrom. 2: 599-606.

Grosjean, D. (1983) Polycyclic aromatic hydrocarbons in Los Angeles air from samples collected on Teflon, glass and quartz filters. Atmos. Environ. 17: 2565-2573.

Grosjean, D.; Parmar, S. S. (1990) Interferences from aldehydes and peroxyacetyl nitrate when sampling urban air organic acids on alkaline traps. Environ. Sci. Technol. 24: 1021-1026.

Grosjean, D.; Williams, E. L., II; Grosjean, E.; Novakov, T. (1994) Evolved gas analysis of secondary organic aerosols. Aerosol Sci. Technol. 21: 306-324.

Gundel, L. A.; Lane, D. A. (1999) Sorbent-coated diffusion denuders for direct measurement of gas/particle partitioning by semi-volatile organic compounds. In: Lane, D. A., ed. Gas and particle phase measurements of atmospheric organic compounds. Amsterdam, The Netherlands: Gordon and Breach Science Publishers. (Advances in environmental, industrial and process control technologies: v. 2).

Gundel, L. A.; Lee, V. C.; Mahanama, K. R. R.; Stevens, R. K.; Daisey, J. M. (1995) Direct determination of the phase distributions of semi-volatile polycyclic aromatic hydrocarbons using annular denuders. Atmos. Environ. 29: 1719-1733.

Hänninen, O. O.; Koistinen, K. J. (2002) Quantitative analysis of environmental factors in differential weighing of blank teflon filters. J. Air Waste Manage. Assoc. 52: 134-139.

Hansen, A. D. A.; McMurry, P. H. (1990) An intercomparison of measurements of aerosol elemental carbon during the 1986 carbonaceous species method comparison study. J. Air Waste Manage. Assoc. 40: 894-895.

Hansen, A. D. A.; Rosen, H.; Novakov, T. (1984) The aethalometer - an instrument for the real-time measurement of optical absorption by aerosol particles. Sci. Total Environ. 36: 191-196.

Harman, J. N., Chairman. (1989) Inductively coupled plasma emission spectroscopy. In: Lodge, J. P., Jr., ed. Methods of air sampling and analysis. 3rd ed. Chelsea, MI: Lewis Publishers, Inc.; pp. 89-92.

Haupt, O.; Klane, B.; Schaefer, C.; Dannecker, W. (1995) Preparation of quartz fibre filter standards for x-ray fluorescence analysis of aerosol samples. X-Ray Spectrom. 24: 267-275.

He, K.; Yang, F.; Ma, Y.; Zhang, Q.; Yao, X.; Chan, C. K.; Cadle, S.; Chan, T.; Mulawa, P. (2001) The characteristics of $PM_{2.5}$ in Beijing, China. Atmos. Environ. 35: 4959-4970.

Hering, S.; Cass, G. (1999) The magnitude of bias in the measurement of $PM_{2.5}$ arising from volatilization of particulate nitrate from Teflon filters. J. Air Waste Manage. Assoc. 49: 725-733.

Hering, S. V.; Appel, B. R.; Cheng, W.; Salaymeh, F.; Cadle, S. H.; Mulawa, P. A.; Cahill, T. A.; Eldred, R. A.; Surovik, M.; Fitz, D.; Howes, J. E.; Knapp, K. T.; Stockburger, L.; Turpin, B. J.; Huntzicker, J. J.; Zhang, X.-Q.; McMurry, P. H. (1990) Comparison of sampling methods for carbonaceous aerosols in ambient air. Aerosol Sci. Technol. 12: 200-213.

Hinds, W. C. (1999) Aerosol technology: properties, behavior, and measurement of airborne particles. 2nd ed. New York, NY: John Wiley & Sons.

Hinz, K.-P.; Kaufmann, R.; Spengler, B. (1994) Laser-induced mass analysis of single particles in airborne state. Anal. Chem. 66: 2071-2076.

Hitzenberger, R.; Dusek, U.; Berner, A. (1996) Black carbon measurements using an integrating sphere. J. Geophys. Res. [Atmos.] 101: 19,601-19,606.

Hitzenberger, R.; Berner, A.; Dusek, U.; Alabashi, R. (1997) Humidity-dependent growth of size-segregated aerosol samples. Aerosol Sci. Technol. 27: 116-130.

Hitzenberger, R.; Jennings, S. G.; Larson, S. M.; Dillner, A.; Cachier, H.; Galambos, Z.; Rouc, A.; Spain, T. G. (1999) Intercomparison of measurement methods for black carbon aerosols. Atmos. Environ. 33: 2823-2833.

Holyńska, B.; Ptasiński, J.; Maenhaut, W.; Annegarn, H. J. (1997) Energy-dispersive x-ray fluorescence spectrometer with capillary optics for the chemical analysis of atmospheric aerosols with high time resolution. J. Aerosol Sci. 28: 1455-1464.

Houck, J. E.; Chow, J. C.; Watson, J. G.; Simons, C. A.; Pritchett, L. C.; Goulet, J. M.; Frazier, C. A. (1989) Determination of particle size distribution and chemical composition of particulate matter from selected sources in California: volume I and executive summary (final report). Sacramento, CA: California Air Resources Board.

Huntzicker, J. J.; Hoffman, R. S.; Ling, C.-S. (1978) Continuous measurement and speciation of sulfur-containing aerosols by flame photometry. Atmos. Environ. 12: 83-88.

Husar, R. B.; Tratt, D. M.; Schichtel, B. A.; Falke, S. R.; Li, F.; Jaffe, D.; Gasso, S.; Gill, T.; Laulainen, N. S.; Lu, F.; Reheis, M. C.; Chun, Y.; Westphal, D.; Holben, B. N.; Gueymard, C.; McKendry, I.; Kuring, N.; Feldman, G. C.; McClain, C.; Frouin, R. J.; Merrill, J.; DuBois, D.; Vignola, F.; Murayama, T.; Nickovic, S.; Wilson, W. E.; Sassen, K.; Sugimoto, N.; Malm, W. C. (2001) The Asian dust events of April 1998. J. Geophys. Res. 106: 18,317-18,330.

Jackson, K. W.; Mahmood, T. M. (1994) Atomic absorption, atomic emission, and flame emission spectrometry. Anal. Chem. 66: 252R-279R.

Jaklevic, J. M.; Loo, B. W.; Goulding, F. S. (1977) Photon-induced X-ray fluorescence analysis using energy-dispersive detector and dichotomous sampler. In: Dzubay, T. G., ed. X-ray fluorescence analysis of environmental samples. Ann Arbor, MI: Ann Arbor Science Publishers, Inc.; pp. 3-18.

Jaklevic, J. M.; Gatti, R. C.; Goulding, F. S.; Loo, B. W. (1981a) A β-gauge method applied to aerosol samples. Environ. Sci. Technol. 15: 680-686.

Jaklevic, J. M.; Loo, B. W.; Fujita, T. Y. (1981b) Automatic particulate sulfur measurements with a dichotomous sampler and on-line x-ray fluorescence analysis. Environ. Sci. Technol. 15: 687-690.

Jang, M.; Kamens, R. M. (1998) A thermodynamic approach for modeling partitioning of semivolatile organic compounds on atmospheric particulate matter: humidity effects. Environ. Sci. Technol. 32: 1237-1243.

Jang, M.; Kamens, R. M.; Leach, K. B.; Strommen, M. R. (1997) A thermodynamic approach using group contribution methods to model the partitioning of semivolatile organic compounds on atmospheric particulate matter. Environ. Sci. Technol. 31: 2805-2811.

Jenkins, B. M.; Jones, A. D.; Turn, S. Q.; Williams, R. B. (1996) Particle concentrations, gas-particle partitioning, and species intercorrelations for polycyclic aromatic hydrocarbons (PAH) emitted during biomass burning. Atmos. Environ. 30: 3825-3835.

Johansson, T. B.; Askelsson, R.; Johansson, S. A. E. (1970) X-ray analysis: elemental trace analysis at the 10-(12)g level. Nucl. Instrum. Methods 84: 141-143.

John, W.; Sethi, V. (1993) Threshold for resuspension by particle impaction. Aerosol Sci. Technol. 19: 69-79.

John, W.; Wall, S. M.; Ondo, J. L. (1988) A new method for nitric acid and nitrate aerosol measurement using the dichotomous sampler. Atmos. Environ. 22: 1627-1635.

Johnson, M V.; Wexler, A. S. (1995) MS of individual aerosol particles. Anal. Chem. 67: 721A-726A.

Johnson, R. L.; Shah, J. J.; Huntzicker, J. J. (1980) Analysis of organic, elemental, and carbonate carbon in ambient aerosols. In: Verner, S. S. Sampling and analysis of toxic organics in the atmosphere: Proceedings from a symposium; August, 1979; Boulder, CO. Philadelphia, PA: American Society for Testing and Materials, pp. 111-119. [ASTM Special Technical Publication 721].

Kamens, R. M.; Coe, D. L. (1997) A large gas-phase stripping device to investigate rates of PAH evaporation from airborne diesel soot particles. Environ. Sci. Technol. 31: 1830-1833.

Kamens, R.; Odum, J.; Fan, Z.-H. (1995) Some observations on times to equilibrium for semivolatile polycyclic aromatic hydrocarbons. Environ. Sci. Technol. 29: 43-50.

Kamens, R.; Jang, M.; Chien, C.-J.; Leach, K. (1999) Aerosol formation from the reaction of α-pinene and ozone using a gas-phase kinetics-aerosol partitioning model. Environ. Sci. Technol. 33: 1430-1438.

Kao, A. S.; Friedlander, S. K. (1995) Frequency distributions of PM_{10} chemical components and their sources. Environ. Sci. Technol. 29: 19-28.

Kavouras, I. G.; Koutrakis, P. (2001) Use of polyurethane foam as the impaction substrate/collection medium in conventional inertial impactors. Aerosol Sci. Technol. 34: 46-56.

Keeler, G. J.; Pirrone, N. (1996) Atmospheric transport and deposition of trace elements to Lake Erie from urban areas. Water Sci. Technol. 33: 259-265.

Keeler, G. J.; Brachaczek, W. W.; Gorse, R. A., Jr.; Japar, S. M.; Pierson, W. R. (1988) Effect of ambient humidity on dichotomous sampler coarse/fine ratios. Atmos. Environ. 22: 1715-1720.

Keywood, M. D.; Ayers, G. P.; Gras, J. L.; Gillett, R. W.; Cohen, D. D. (1999) Relationships between size segregated mass concentration data and ultrafine particle number concentrations in urban areas. Atmos. Environ. 33: 2907-2913.

Keywood, M. D.; Ayers, G. P.; Gras, J. L.; Gillett, R. W.; Cohen, D. D. (2000) Size distribution and sources of aerosol in Launceston, Australia, during winter 1997. J. Air Waste Manage. Assoc. 50: 418-427.

Khlystov, A.; Wyers, G. P.; Slanina, J. (1995) The steam-jet aerosol collector. Atmos. Environ. 29: 2229-2234.

Kim, Y. J.; Boatman, J. F.; Gunter, R. L.; Wellman, D. L.; Wilkison, S. W. (1993) Vertical distribution of atmospheric aerosol size distribution over south-central New Mexico. Atmos. Environ. Part A 27: 1351-1362.

Kim, B. M.; Lester, J.; Tisopulos, L.; Zeldin, M. D. (1999) Nitrate artifacts during $PM_{2.5}$ sampling in the South Coast Air Basin of California. J. Air Waste Manage. Assoc. 49: 142-153.

Kim, B. M.; Cassmassi, J.; Hogo, H.; Zeldin, M. D. (2001) Positive organic carbon artifacts on filter medium during $PM_{2.5}$ sampling in the South Coast Air Basin. Aerosol. Sci. Technol. 34: 35-41.

Kirchstetter, T. W.; Corrigan, C. E.; Novakov, T. (2001) Laboratory and field investigation of the adsorption of gaseous organic compounds onto quartz filters. Atmos. Environ. 35: 1663-1671.

Kittelson, D. B. (1998) Engines and nanoparticles: a review. J. Aerosol Sci. 29: 575-588.

Kittelson, D. B.; McKenzie, R.; Vermeersch, M.; Dorman, F.; Pui, D.; Linne, M.; Liu, B.; Whitby, K. (1978) Total sulfur aerosol concentration with an electrostatically pulsed flame photometric detector system. Atmos. Environ. 12: 105-111.

Klouda, G. A.; Klinedinst, D. B.; Steel, E. B.; Benner, B. A., Jr.; Parish, H. J. (1996) Exploring a method to produce an urban dust particle filter standard. J. Aerosol Sci. 27(suppl. 1): S351-S352.

Koistinen, K. J.; Kousa, A.; Tenhola, V.; Hänninen, O.; Jantunen, M. J.; Oglesby, L.; Kuenzli, N.; Georgoulis, L. (1999) Fine particle ($PM_{2.5}$) measurement methodology, quality assurance procedures and pilot results of the *EXPOLIS* study. J. Air Waste Manage. Assoc. 49: 1212-1220.

Koutrakis, P.; Kelly, B. P. (1993) Equilibrium size of atmospheric aerosol sulfates as a function of particle acidity and ambient relative humidity. J. Geophys. Res. [Atmos.] 98: 7141-7147.

Koutrakis, P.; Wolfson, J. M.; Slater, J. L.; Brauer, M.; Spengler, J. D.; Stevens, R. K.; Stone, C. L. (1988) Evaluation of an annular denuder/filter pack system to collect acidic aerosols and gases. Environ. Sci. Technol. 22: 1463-1468.

Koutrakis, P.; Wolfson, J. M.; Spengler, J. D.; Stern, B.; Franklin, C. A. (1989) Equilibrium size of atmospheric aerosol sulfates as a function of the relative humidity. J. Geophys. Res. [Atmos.] 94: 6442-6448.

Koutrakis, P.; Sioutas, C.; Ferguson, S. T.; Wolfson, J. M.; Mulik, J. D.; Burton, R. M. (1993) Development and evaluation of a glass honeycomb denuder/filter pack system to collect atmospheric gases and particles. Environ. Sci. Technol. 27: 2497-2501.

Koutrakis, P.; Wang, P.-Y.; Wolfson, J. M.; Sioutas, C., inventors. (1996) Method and apparatus to measure particulate matter in gas. U.S. patent 5,571,945. November 5. Available: www.patents.ibm.com [1999, November 24].

Krieger, M. S.; Hites, R. A. (1992) Diffusion denuder for the collection of semivolatile organic compounds. Environ. Sci. Technol. 26: 1551-1555.

Kusko, B. H.; Cahill, T. A.; Eldred, R. A.; Matsuda, Y.; Miyake, H. (1989) Nondestructive analysis of total nonvolatile carbon by Forward Alpha Scattering Technique (FAST). Aerosol Sci. Technol. 10: 390-396.

Lammel, G.; Novakov, T. (1995) Water nucleation properties of carbon black and diesel soot particles. Atmos. Environ. 29: 813-823.

Landsberger, S. (1988) Improved methodology for the determination of the seven elemental tracer long-distance pollution signatures using thermal and epithermal neutron activation analysis. Anal. Chem. 60: 1842-1845.

Landsberger, S.; Wu, D. (1993) Improvement of analytical sensitivities for the determination of antimony, arsenic, cadmium, indium, iodine, molybdenum, silicon and uranium in airborne particulate matter by epithermal neutron activation analysis. J. Radioanal. Nucl. Chem. 167: 219-225.

Landsberger, S.; Wu, D.; Vermette, S. J.; Cizek, W. (1997) Intercomparison of IAEA airborne particulate matter reference material. J. Radioanal. Nucl. Chem. 215: 117-127.

Lane, D. A.; Johnson, N. D.; Barton, S. C.; Thomas, G. H. S.; Schroeder, W. H. (1988) Development and evaluation of a novel gas and particle sampler for semivolatile chlorinated organic compounds in ambient air. Environ. Sci. Technol. 22: 941-947.

Lavanchy, V. M. H.; Gäggeler, H. W.; Nyeki, S.; Baltensperger, U. (1999) Elemental carbon (EC) and black carbon (BC) measurements with a thermal method and an aethalometer at the high-alpine research station Jungfraujoch. Atmos. Environ. 33: 2759-2769.

Lawless, P. A.; Rodes, C. E.; Evans, G.; Sheldon, L.; Creason, J. (2001) Aerosol concentrations during the 1999 Fresno Exposure Studies as functions of size, season and meteorology. Aerosol Sci. Technol. 34: 66-74.

Lawrence, J. E.; Koutrakis, P. (1994) Measurement of atmospheric formic and acetic acids: methods evaluation and results from field studies. Environ. Sci. Technol. 28: 957-964.

Lawrence, J. E.; Koutrakis, P. (1996a) Measurement and speciation of gas and particulate phase organic acidity in an urban environment: 1. analytical. J. Geophys. Res. [Atmos.] 101: 9159-9169.

Lawrence, J.; Koutrakis, P. (1996b) Measurement and speciation of gas and particulate phase organic acidity in an urban environment: 2. speciation. J. Geophys. Res. [Atmos.] 101: 9171-9184.

Lawson, D. R. (1990) The Southern California Air Quality Study. J. Air Waste Manage. Assoc. 40: 156-165.
Lee, W.-M. G.; Shih, P.-M.; Wang, C.-S. (1997) The influence of relative humidity on the size of atmospheric aerosol. J. Environ. Sci. Health Part A: Environ. Sci. Eng. Toxic Hazard. Subst. Control 32: 1085-1097.
Lewtas, J.; Pang, Y.; Booth, D.; Reimer, S.; Eatough, D. J.; Gundel, L. A. (2001) Comparison of sampling methods for semi-volatile organic carbon associated with $PM_{2.5}$. Aerosol Sci. Technol. 34: 9-22.
Li, W.; Montassier, N.; Hopke, P. K. (1992) A system to measure the hygroscopicity of aerosol particles. Aerosol Sci. Technol. 17: 25-35.
Liang, Z.; Chan, C. K. (1997) A fast technique for measuring water activity of atmospheric aerosols. Aerosol Sci. Technol. 26: 255-268.
Liang, C.; Pankow, J. F.; Odum, J. R.; Seinfeld, J. H. (1997) Gas/particle partitioning of semivolatile organic compounds to model inorganic, organic, and ambient smog aerosols. Environ. Sci. Technol. 31: 3086-3092.
Lin, J. J.; Noll, K. E.; Holsen, T. M. (1994) Dry deposition velocities as a function of particle size in the ambient atmosphere. Aerosol Sci. Technol. 20: 239-252.
Lindskog, A.; Brorström-Lundén, E.; Alfheim, I; Hagen, I. (1987) Chemical transformation of PAH on airborne particles by exposure to NO_2 during sampling: a comparison between two filter media. Sci. Total Environ. 61: 51-57.
Liousse, C.; Cachier, H.; Jennings, S. G. (1993) Optical and thermal measurements of black carbon aerosol content in different environments: variation of the specific attenuation cross-section, sigma (σ). Atmos. Environ. 27A: 1203-1211.
Loo, B. W.; Cork, C. P. (1988) Development of high efficiency virtual impactors. Aerosol Sci. Technol. 9: 167-176.
Loo, B. W.; Jaklevic, J. M.; Goulding, F. S. (1976) Dichotomous virtual impactors for large scale monitoring of airborne particulate matter. In: Liu, B. Y. H., ed. Fine particles: aerosol generation, measurement, sampling, and analysis. New York, NY: Academic Press; pp. 311-350.
Lundgren, D. A.; Burton, R. M. (1995) Effect of particle size distribution on the cut point between fine and coarse ambient mass fractions. In: Phalen, R. F.; Bates, D. V., eds. Proceedings of the colloquium on particulate air pollution and human mortality and morbidity; January 1994; Irvine, CA. Inhalation Toxicol. 7: 131-148.
Lundgren, D. A.; Hausknecht, B. J.; Burton, R. M. (1984) Large particle size distribution in five U.S. cities and the effect on a new ambient particulate matter standard (PM_{10}). Aerosol Sci. Technol. 7: 467-473.
Lynch, A. J.; McQuaker, N. R.; Brown, D. F. (1980) ICP/AES analysis and the composition of airborne and soil materials in the vicinity of a lead/zinc smelter complex. J. Air Pollut. Control Assoc. 30: 257-260.
Mahanama, K. R. R.; Gundel, L. A.; Daisey, J. M. (1994) Selective fluorescence detection of polycyclic aromatic hydrocarbons in environmental tobacco smoke and other airborne particles. Int. J. Environ. Anal. Chem. 56: 289-309.
Malm, W. C.; Sisler, J. F.; Huffman, D.; Eldred, R. A.; Cahill, T. A. (1994) Spatial and seasonal trends in particle concentration and optical extinction in the United States. J. Geophys. Res. [Atmos.] 99: 1347-1370.
Mamane, Y.; Willis, R.; Conner, T. (2001) Evaluation of computer-controlled scanning electron microscopy applied to an ambient urban aerosol sample. Aerosol Sci. Technol. 34: 97-107.
Marple, V. A.; Chien, C. M. (1980) Virtual impactors: a theoretical study. Environ. Sci. Technol. 14: 976-985.
Marple, V. A.; Liu, B. Y. H.; Burton, R. M. (1990) High-volume impactor for sampling fine and coarse particles. J. Air Waste Manage. Assoc. 40: 762-767.
Marple, V. A.; Rubow, K. L.; Behm, S. M. (1991) A microorifice uniform deposit impactor (MOUDI): description, calibration, and use. Aerosol Sci. Technol. 14: 434-446.
McInnes, L. M.; Quinn, P. K.; Covert, D. S.; Anderson, T. L. (1996) Gravimetric analysis, ionic composition, and associated water mass of the marine aerosol. Atmos. Environ. 30: 869-884.
McMurry, P. H. (2000) A review of atmospheric aerosol measurements. Atmos. Environ. 34: 1959-1999.
McMurry, P. H.; Friedlander, S. K. (1979) New particle formation in the presence of an aerosol. Atmos. Environ. 13: 1635-1651.
McMurry, P. H.; Stolzenburg, M. R. (1989) On the sensitivity of particle size to relative humidity for Los Angeles aerosols. Atmos. Environ. 23: 497-507.
McMurry, P. H.; Woo, K. S.; Weber, R.; Chen, D.-R.; Pui, D. Y. H. (2000) Size distributions of 3-10 nm atmospheric particles: implications for nucleation mechanisms. Philos. Trans. R. Soc. London Ser. A 358: 2625-2642.
McQuaker, N. R.; Kluckner, P. D.; Chang, G. N. (1979) Calibration of an inductively coupled plasma–atomic emission spectrometer for the analysis of environmental materials. Anal. Chem. 51: 888-895.

Meyer, A. S., Jr.; Boyd, C. M. (1959) Determination of water by titration with coulometrically generated Karl Fischer reagent. Anal. Chem. 31: 215-219.

Meyer, M. B.; Rupprecht, E. G. (1996) Particulate matter sampling methods: the importance of standardization. J. Aerosol Sci. 27(suppl. 1): S349-S350.

Meyer, M. B.; Rupprecht, E.; Patashnick, H. (1995) Considerations for the sampling and measurement of ambient particulate matter. In: Particulate matter: health and regulatory issues: proceedings of an international specialty conference; April; Pittsburgh, PA. Pittsburgh, PA: Air & Waste Management Association; pp. 204-210. (A&WMA publication VIP-49).

Meyer, M. B.; Patashnick, H.; Ambs, J. L.; Rupprecht, E. (2000) Development of a sample equilibration system for the TEOM continuous PM monitor. J. Air Waste Manage. Assoc. 50: 1345-1349.

Middlebrook, A. M.; Murphy, D. M.; Thomson, D. S. (1998) Observations of organic material in individual marine particles at Cape Grim during the first aerosol characterization experiment (ACE 1). J. Geophys. Res. 103: 16475-16483.

Mignacca, D.; Stubbs, K. (1999) Effects of equilibration temperature on PM_{10} concentrations from the TEOM method in the Lower Fraser Valley. J. Air Waste Manage. Assoc. 49: 1250-1254.

Miller, F. J.; Gardner, D. E.; Graham, J. A.; Lee, R. E., Jr.; Wilson, W. E.; Bachmann, J. D. (1979) Size considerations for establishing a standard for inhalable particles. J. Air Pollut. Control Assoc. 29: 610-615.

Misra, C.; Geller, M. D.; Shah, P.; Sioutas, C.; Solomon, P. A. (2001) Development and evaluation of a continuous coarse (PM_{10}-$PM_{2.5}$) particle monitor. J. Air Waste Manage. Assoc. 51: 1309-1317.

Modey, W. K.; Pang, Y.; Eatough, N. L.; Eatough, D. J. (2001) Fine particulate ($PM_{2.5}$) composition in Atlanta, USA: assessment of the particle concentrator-Brigham Young University organic sampling system, PC-BOSS, during the EPA supersite study. Atmos. Environ. 35: 6493-6502.

Moosmüller, Hans; Arnott, W. P.; Rogers, C. F. (1998). Photoacoustic and filter measurements related to aerosol light absorption during the Northern Front Range Air Quality Study (Colorado 1996/1997). Journal of Geophysical Research 103, no. D21: 28,149-28,157.

Morawska, L.; Bofinger, N. D.; Kocis, L.; Nwankwoala, A. (1998) Submicrometer and supermicrometer particles from diesel vehicle emissions. Environ. Sci. Technol. 32: 2033-2042.

Mueller, P. K.; Collins, J. F. (1980) Development of a particulate sulfate analyzer. Palo Alto, CA: Electric Power Research Institute; report no. P-1382F.

Mueller, P. K.; Fung, K. K.; Heisler, S. L.; Grosjean, D.; Hidy, G. M. (1982) Atmospheric particulate carbon observations in urban and rural areas of the United States. In: Wolff, G. T.; Klimisch, R. L., eds. Particulate Carbon: Atmospheric Life Cycle. New York, NY: Plenum Publishing Corporation, 343-370.

Mukerjee, S.; Ellenson, W. D.; Lewis, R. G.; Stevens, R. K.; Somerville, M. C.; Shadwick, D. S.; Willis, R. D. (1997) An environmental scoping study in the lower Rio Grande Valley of Texas–III. residential microenvironmental monitoring for air, house dust, and soil. Environ. Int. 23: 657-673.

Munksgaard, N. C.; Parry, D. L. (1998) Lead isotope ratios determined by ICP-MS: monitoring of mining-derived metal particulates in atmospheric fallout, Northern Territory, Australia. Sci. Total Environ. 217: 113-126.

Murphy, D. M.; Thomson, D. S. (1997) Chemical composition of single aerosol particles at Idaho Hill: negative ion measurements. J. Geophys. Res. 102: 6353-6368.

Nadkarni, R. A. (1975) Multielement analysis of coal and coal fly ash standards by instrumental neutron activation analysis. Radiochem. Radioanal. Lett. 21: 161-176.

Nam, S.; Masamba, W. R. L.; Montaser, A. (1993) Investigation of helium inductively coupled plasma-mass spectrometry for the detection of metals and nonmetals in aqueous solutions. Anal. Chem. 65: 2784-2790.

NARSTO. (2002) Fine particle assessment (2nd review draft). Available: http://www.cgenv.com/Narsto/ (13 February 2002).

National Research Council. (1993) Protecting visibility in national parks and wilderness areas. Washington, DC: National Academy Press. 3v.

Nelson, M. P.; Zugates, C. T.; Treado, P. J.; Casucio, G. S.; Exline, D. L.; Schlaegle, S. F. (2000) Combining Raman chemical imaging scanning electron mocroscopy (SEM) to characterize ambient fine particulate. Aerosol Sci. Technol. 34: 108-117.

Neubauer, K. R.; Johnson, M. V.; Wexler, A. S. (1998) Humidity effects of the mass spectra of single aerosol particles. Atmos. Environ. 32: 2521-2529.

Nham, T. T. (1996) Cool plasma technique for the ultra-trace level determination of 56Fe, 40Ca, 39K, 7Li, 24Mg and 23Na for semi-conductor applications. Mulgrave, Victoria, Australia: Varian Australia Pty Ltd; ICP-MS instruments at work no. ICP-MS-7. Available: http://www.varianinc.com/cgi-bin/nav?varinc/docs/osi/icpms/atwork/icpms7&cid=680052 [18 January, 2002].

Noble, C. A.; Prather, K. A. (1996) Real-time measurement of correlated size and composition profiles of individual atmospheric aerosol particles. Environ. Sci. Technol. 30: 2667-2680.

Noble, C. A.; Prather, K. A. (1998) Single particle characterization of albuterol metered dose inhaler aerosol in near real-time. Aerosol Sci. Technol. 29: 294-306.

Noble, C. A.; Vanderpool, R. W.; Peters, T. M.; McElroy, F. F.; Gemmill, D. B.; Wiener, R. W. (2001) Federal reference and equivalent methods for measuring fine particulate matter. Aerosol Sci. Technol. 34: 457-464.

Noël, M. A.; Topart, P. A. (1994) High-frequency impedance analysis of quartz crystal microbalances. 1. General considerations. Anal. Chem. 66: 484-491.

Novakov, T.; Corrigan, C. E.; Penner, J. E.; Chuang, C. C.; Rosario, O.; Mayol Bracero, O. L. (1997) Organic aerosols in the Caribbean trade winds: a natural source? J. Geophys. Res. 102: 21307-21313.

Obeidi, F. D.; Eatough, D. J. (2002) Continuous measurement of semi-volatile fine particulate mass in Provo, UT. Aerosol Sci. Technol. 36: 191-203.

Obeidi, F.; Eatough, N. L.; Eatough, D. J. (2002) Use of the RAMS to measure semivolatile fine particulate matter at Riverside and Bakersfield, California. Aerosol Sci. Technol. 36: 204-216.

Ohta, S.; Okita, T. (1990) A chemical characterization of atmospheric aerosol in Sapporo. Atmos. Environ. Part A 24A: 815-822.

Ohta, S.; Hori, M.; Yamagata, S.; Murao, N. (1998) Chemical characterization of atmospheric fine particles in Sapporo with determination of water content. Atmos. Environ. 32: 1021-1025.

Olin, J. G.; Sem, G. J. (1971) Piezoelectric microbalance for monitoring the mass concentration of suspended particles. Atmos. Environ. 5: 653-668.

Olmez, I. (1989) Trace element signatures in groundwater pollution. In: Watson, J. G., ed. Receptor models in air resources management: an international specialty conference; February 1988; San Francisco, CA. Pittsburgh, PA: Air & Waste Management Association; pp. 3-11. (APCA transactions series: no. 14).

Ondov, J. M. (1996) Particulate tracers for source attribution: potential for application to California's San Joaquin Valley. J. Aerosol Sci. 27(suppl. 1): S687-S688.

Ondov, J. M.; Divita, F., Jr. (1993) Size spectra for trace elements in urban aerosol particles by instrumental neutron activation analysis. J. Radioanal. Nucl. Chem. 167: 247-258.

Ondov, J. M.; Kelly, W. R.; Holland, J. Z.; Lin, Z. C.; Wight, S. A. (1992) Tracing fly ash emitted from a coal-fired power plant with enriched rare-earth isotopes: an urban scale test. Atmos. Environ. Part B 26B: 453-462.

Ono, D. M.; Hardebeck, E.; Parker, J.; Cox, B. G. (2000) Systematic biases in measured PM_{10} values with U.S. Environmental Protection Agency-approved samplers at Owens Lake, California. J. Air & Waste Manage. Assoc. 50: 1144-1156.

Pak, S. S.; Liu, B. Y. H.; Rubow, K. L. (1992) Effect of coating thickness on particle bounce in inertial impactors. Aerosol Sci. Technol. 16: 141-150.

Pang, Y.; Ren, Y.; Obeidi, F.; Hastings, R.; Eatough, D. J.; Wilson, W. E. (2001) Semi-volatile species in $PM_{2.5}$: comparison of integrated and continuous samplers for $PM_{2.5}$ research or monitoring. J. Air Waste Manage. Assoc. 51: 25-36.

Pang, Y.; Eatough, N. L.; Wilson, J.; Eatough, D. J. (2002a) Effect of semivolatile material on PM2.5 measurement by the PM2.5 federal reference method sampler at Bakersfield, California. Aerosol Sci. Technol. 36: 289-299.

Pang, Y.; Eatough, N. L.; Eatough, D. J. (2002b) PM2.5 semi-volatile organic material at Riverside CA: implications for the PM2.5 federal reference method sampler. Aerosol Sci. Technol. 36: 277-288.

Pankow, J. F.; Bidleman, T. F. (1991) Effects of temperature, *TSP* and per cent non-exchangeable material in determining the gas-particle partitioning of organic compounds. Atmos. Environ. Part A 25: 2241-2249.

Pankow, J. F.; Storey, J. M. E.; Yamasaki, H. (1993) Effects of relative humidity on gas/particle partitioning of semivolatile organic compounds to urban particulate matter. Environ. Sci. Technol. 27: 2220-2226.

Parmar, S. S.; Grosjean, D. (1990) Laboratory tests of KI and alkaline annular denuders. Atmos. Environ. Part A 24: 2695-2698.

Patashnick, H.; Rupprecht, G.; Ambs, J. L.; Meyer, M. B. (2001) Development of a reference standard for particulate matter mass in ambient air. Aerosol Sci. Technol. 34: 42-45.

Pellizzari, E. D.; Krost, K. J. (1984) Chemical transformations during ambient air sampling for organic vapors. Anal. Chem. 56: 1813-1819.

Peng, C.; Chan, C.K. (2001) The water cycles of water-soluble organic salts of atmospheric importance. Atmos. Environ. 35: 1183-1192.

Penner, J. E.; Novakov, T. (1996) Carbonaceous particles in the atmosphere: a historical perspective to the fifth international conference on carbonaceous particles in the atmosphere. J. Geophys. Res. [Atmos.] 101: 19,373-19,378.

Perry, K. D.; Cahill, T. A.; Schnell, R. C.; Harris, J. M. (1999) Long-range transport of anthropogenic aerosols to the National Oceanic and Atmospheric Administration baseline station at Mauna Loa Observatory, Hawaii. J. Geophys. Res. Atmos. 104: 18,521-18,533.

Peters, A.; Liu, E.; Verrier, R. L.; Schwartz, J.; Gold, D. R.; Mittleman, M.; Baliff, J.; Oh, J. A.; Allen, G.; Monahan, K.; Dockery, D. W. (2000) Air pollution and incidence of cardiac arrhythmia. Epidemiology 11: 11-17.

Peters, T. M.; Vanderpool, R. W.; Wiener, R. W. (2001a) Design and calibration of the EPA $PM_{2.5}$ Well Impactor Ninety-Six (WINS), Aerosol Sci. Technol. 34: 389-397.

Peters, T. M.; Norris, G. A.; Vanderpool, R. W.; Gemmill, D. B.; Wiener, R. W.; Murdoch, R. W.; McElroy, F. F.; Pitchford, M. (2001b) Field performance of $PM_{2.5}$ federal reference method samplers, Aerosol Sci. Technol. 34: 433-443.

Peters, T. M.; Gussman, R. A.; Kenney, L. C.; Vanderpool, R. W. (2001c) Evaluation of $PM_{2.5}$ size selectors used in speciation samplers. Aerosol Sci. Technol. 34: 422-429.

Petzold, A., Niessner, R. (1996). Photoacoustic soot sensor for in-situ black carbon monitoring. Applied Physics B, no. 63: 191-197.

Petzold, A.; Kopp, C.; Niessner, R. (1997) The dependence of the specific attenuation cross-section on black carbon mass fraction and particle size. Atmos. Environ. 31: 661-672.

Pilinis, C.; Seinfeld, J. H.; Grosjean, D. (1989) Water content of atmospheric aerosols. Atmos. Environ. 23: 1601-1606.

Piorek, S. (1994) Modern, PC based, high resolution portable EDXRF analyzer offers laboratory performance for field, in-situ analysis of environmental contaminants. Nucl. Instr. Meth. Phys. Res. A 353: 528-533.

Pirrone, N.; Keeler, G. J. (1996) The Rouge River watershed pollution by trace elements: atmospheric depositions and emission sources. Water Sci. Technol. 33: 267-275.

Plantz, M. (1996) Common molecular ion interferences in ICP-MS. Mulgrave, Victoria, Australia: Varian Australia Pty Ltd; ICP-MS instruments at work no. ICP-MS-6. Available: http://www.varianinc.com/cgi-bin/nav?varinc/docs/osi/icpms/atwork/index&cid=680052 [18 January, 2002].

Pope, C. A., III; Thun, M. J.; Namboodiri, M. M.; Dockery, D. W.; Evans, J. S.; Speizer, F. E.; Heath, C. W., Jr. (1995) Particulate air pollution as a predictor of mortality in a prospective study of U.S. adults. Am. J. Respir. Crit. Care Med. 151: 669-674.

Potts, P. J.; Webb, P. C. (1992) X-ray fluorescence spectrometry. J. Geochem. Explor. 44: 251-296.

Ranweiler, L. E.; Moyers, J. L. (1974) Atomic absorption procedure for analysis of metals in atmospheric particulate matter. Environ. Sci. Technol. 8: 152-156.

Rao, A. K.; Whitby, K. T. (1977) Nonideal collection characteristics of single stage and cascade impactors. Am. Ind. Hyg. Assoc. J. 38: 174-177.

Rao, A. K.; Whitby, K. T. (1978) Non-ideal collection characteristics of inertial impactors--I. single-stage impactors and solid particles. J. Aerosol Sci. 9: 77-86.

Reid, J. S.; Cahill, T. A.; Wakabayashi, P. H.; Dunlap, M. R. (1994) Geometric/aerodynamic equivalent diameter ratios of ash aggregate aerosols collected in burning Kuwaiti well fields. Atmos. Environ. 28: 2227-2234.

Reilly, P. T. A.; Gieray, R. A.; Whitten, W. B.; Ramsey, J. M. (1998) Real-time characterization of the organic composition and size of individual diesel engine smoke particles. Environ. Sci. Technol. 32: 2672-2679.

Reist, P. C. (1984) Introduction to aerosol science. New York, NY: Macmillan Publishing Company.

Reist, P. C. (1993) Introduction to aerosol science. New York, NY: McGraw-Hill.

Risse, U.; Flammenkamp, E.; Kettrup, A. (1996) Determination of aromatic hydrocarbons in air using diffusion denuders. Fresenius J. Anal. Chem. 356: 390-395.

Roberts, R. A.; Corkill, J. (1998) Grass seed field smoke and its impact on respiratory health. J. Environ. Health 60 (June): 10-16.

Rumburg, B.; Alldredge, R.; Claiborn, C. (2001) Statistical distributions of particulate matter and the error associated with sampling frequency. Atmos. Environ. 35: 2907-2920.

Rupprecht, G.; Patashnick, H.; Beeson, D. E.; Green, R. N.; Meyer, M. B. (1995) A new automated monitor for the measurement of particulate carbon in the atmosphere. In: Particulate matter: health and regulatory issues: proceedings of an international specialty conference; April; Pittsburgh, PA. Pittsburgh, PA: Air & Waste Management Association; pp. 309-319. (A&WMA publication VIP-49).

Sakata, K.; Kawabata, K. (1994) Reduction of fundamental polyatomic ions in inductively coupled plasma mass spectrometry. Spectrochim. Acta 49B: 1027-1038.

Saxena, P.; Hildemann, L. M. (1996) Water-soluble organics in atmospheric particles: a critical review of the literature and applications of thermodynamics to identify candidate compounds. J. Atmos. Chem. 24: 57-109.

Saxena, P.; Hildemann, L. M.; McMurry, P. H.; Seinfeld, J. H. (1995) Organics alter hygroscopic behavior of atmospheric particles. J. Geophys. Res. [Atmos.] 100: 18,755-18,770.

Schmid, H.; Laskus, L.; Abraham, J. H.; Baltensperger, U.; Lavanchy, V.; Bizjak, M.; Burba, P.; Cachier, H.; Crow, D.; Chow, J.; Gnauk, T.; Even, A.; ten Brink, H. M.; Giesen, K.-P.; Hitzenberger, R.; Hueglin, C.; Maenhaut, W.; Pio, C.; Carvalho, A.; Putaud, J.-Philippe; Toom-Sauntry, D.; Puxbaum, H. (2001) Results of the "carbon conference" international aerosol carbon round robin test stage I. Atmos. Environ. 35: 2111-2121.

Schwartz, J.; Dockery, D. W.; Neas, L. M. (1996) Is daily mortality associated specifically with fine particles? J. Air Waste Manage. Assoc. 46: 927-939.

Seinfeld, J. H.; Pandis, S. N. (1998) Atmospheric chemistry and physics: from air pollution to climate change. New York, NY: John Wiley & Sons, Inc.

Sem, G. J.; Tsurubayashi, K.; Homma, K. (1977) Performance of the piezoelectric microbalance respirable aerosol sensor. Am. Ind. Hyg. Assoc. J. 38: 580-588.

Shah, J. J.; Rau, J. A. (1990) Carbonaceous methods comparison study: interlaboratory round robin interpretation results. Sacramento, CA: California Air Resources Board. Available: http://www.arb.ca.gov/research/apr/past/pm.htm (21 March 2002).

Silva, P. J.; Prather, K. A. (1997) On-line characterization of individual particles from automobile emissions. Environ. Sci. Technol. 31: 3074-3080.

Simon, P. K.; Dasgupta, P. K. (1995) Continuous automated measurement of the soluble fraction of atmospheric particulate matter. Anal. Chem. 67: 71-78.

Sioutas, C.; Koutrakis, P.; Wolfson, J. M. (1994a) Particle losses in glass honeycomb denuder samplers. Aerosol Sci. Technol. 21: 137-148.

Sioutas, C.; Koutrakis, P.; Burton, R. M. (1994b) Development of a low cutpoint slit virtual impactor for sampling ambient fine particles. J. Aerosol Sci. 25: 1321-1330.

Sioutas, C.; Koutrakis, P.; Burton, R. M. (1994c) A high-volume small cutpoint virtual impactor for separation of atmospheric particulate from gaseous pollutants. Particulate Sci. Technol. 12: 207-221.

Sioutas, C.; Koutrakis, P.; Olson, B. A. (1994d) Development and evaluation of a low cutpoint virtual impactor. Aerosol Sci. Technol. 21: 223-235.

Sioutas, C.; Koutrakis, P.; Burton, R. M. (1995a) A technique to expose animals to concentrated fine ambient aerosols. Environ. Health Perspect. 103: 172-177.

Sioutas, D.; Koutrakis, P.; Ferguson, S. T.; Burton, R. M. (1995b) Development and evaluation of a prototype ambient particle concentrator for inhalation exposure studies. Inhalation Toxicol. 7: 633-644.

Sioutas, C.; Koutrakis, P.; Wolfson, J. M.; Azaroff, L. S.; Mulik, J. D. (1996a) Field evaluation of a glass honeycomb denuder/filter pack system to collect atmospheric gases and particles. In: Gay, B. W.; Fuerst, R. G.; Jayanty, R. K. M., eds. Measurement of toxic and related air pollutants: proceedings of a conference; May, 1994; Durham, NC. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development; report no. EPA/600/R-94/136; pp. 426-435.

Sioutas, C.; Wang, P. Y.; Ferguson, S. T.; Koutrakis, P.; Mulik, J. D. (1996b) Laboratory and field evaluation of an improved glass honeycomb denuder/filter pack sampler. Atmos. Environ. 30: 885-895.

Sioutas, C.; Koutrakis, P.; Wang, P.-Y.; Babich, P.; Wolfson, J. M. (1999) Experimental investigation of pressure drop with particle loading in Nuclepore filters. Aerosol Sci. Technol. 30: 71-83.

Solomon, P. A.; Moyers, J. L.; Fletcher, R. A. (1983) High-volume dichotomous virtual impactor for the fractionation and collection of particles according to aerodynamic size. Aerosol Sci. Technol. 2: 455-464.

Solomon, P. A.; Salmon, L. G.; Fall, T.; Cass, G. R. (1992) Spatial and temporal distribution of atmospheric nitric acid and particulate nitrate concentrations in the Los Angeles area. Environ. Sci. Technol. 26: 1596-1601.

Song, X.-H.; Faber, N. M.; Hopke, P. K.; Suess, D. T.; Prather, K. A.; Schauer, J. J.; Cass, G. R. (2001) Source apportionment of gasoline and diesel by multivariate calibration based on single particle mass spectral data. Anal. Chim. Acta 446: 329-343.

Speer, R. E.; Barnes, H. M.; Brown, R. (1997) An instrument for measuring the liquid water content of aerosols. Aerosol Sci. Technol. 27: 50-61.

Stahlschmidt, T.; Schulz, M.; Dannecker, W. (1997) Application of total-reflection X-ray fluorescence for the determination of lead, calcium and zinc in size-fractionated marine aerosols. Spectrochim. Acta B 52: 995-1001.

Stein, S. W.; Turpin, B. J.; Cai, X.; Huang, P.-F.; McMurry, P. H. (1994) Measurements of relative humidity-dependent bounce and density for atmospheric particles using the DMA-impactor technique. Atmos. Environ. 28: 1739-1746.

Stevens, R. K.; O'Keeffe, A. E.; Ortman, G. C. (1969) Absolute calibration of a flame photometric detector to volatile sulfur compounds at sub-part-per-million levels. Environ. Sci. Technol. 3: 652-655.

Stevens, R. K.; Mulik, J. D.; O'Keefe, A. E.; Krost, K. J. (1971) Gas chromatography of reactive sulfur gases in air at the parts-per-billion level. Anal. Chem. 43: 827-831.

Stolzenburg, M. R.; Hering, S. V. (2000) Method for the automated measurement of fine particle nitrate in the atmosphere. Environ. Sci. Technol. 34: 907-914.

Streit, N.; Weingartner, E.; Zellweger, C.; Schwikowski, M.; Gäggeler, H. W.; Baltensperger, U. (2000) Characterization of size-fractionated aerosol from the Jungfraujoch (3580 m asl) using total reflection X-ray fluorescence (TXRF). Int. J. Environ. Anal. Chem. 76: 1-16.

Strommen, M. R.; Kamens, R. M. (1997) Development and application of a dual-impedance radial diffusion model to simulate the partitioning of semivolatile organic compounds in combustion aerosols. Environ. Sci. Technol. 31: 2983-2990.

Suarez, A. E.; Caffrey, P. F.; Ondov, J. M.; Thaung, K. C. (1996) Size distribution of Ir-tagged soot from diesel sanitation trucks. J. Aerosol Sci. 27(suppl. 1): S697-S698.

Suggs, J. C.; Burton, R. M. (1983) Spatial characteristics of inhalable particles in the Philadelphia metropolitan area. J. Air Pollut. Control Assoc. 33: 688-691.

Tan, S. H.; Horlick, G. (1986) Background spectral features in inductively coupled plasma mass spectrometry. Appl. Spectros. 40: 445-460.

Tang, I. N. (1980) Deliquescence properties and particle size change of hygroscopic aerosols. In: Willeke, K., ed. Generation of aerosols and facilities for exposure experiments. Ann Arbor, MI: Ann Arbor Science Publishers, Inc.; pp. 153-167.

Tang, H.; Lewis, E. A.; Eatough, D. J.; Burton, R. M.; Farber, R. J. (1994) Determination of the particle size distribution and chemical composition of semi-volatile organic compounds in atmospheric fine particles with a diffusion denuder sampling system. Atmos. Environ. 28: 939-947.

Tanner, R. L.; D'Ottavio, T.; Garber, R.; Newman, L. (1980) Determination of ambient aerosol sulfur using a continuous flame photometric detection system. I. Sampling system for aerosol sulfate and sulfuric acid. Atmos. Environ. 14: 121-127.

Thomson, D. S.; Murphy, D. M. (1994) Analyzing single aerosol particles in real time. Chemtech 24: 30-35.

Tolocka, M.P.; Peters, T. M.; Vanderpool, R. W.; Chen, F.-L.; Wiener, R. W. (2001a) On the modification of the low flow-rate PM_{10} dichotomous sampler inlet, Aerosol Sci. Technol. 34: 407-415.

Tolocka, M. P.; Solomon, P. A.; Mitchell, W.; Norris, G. A.; Gemmill, D. B.; Wiener, R. W.; Vanderpool, R. W.; Homolya, J. B.; Rice, J. (2001b) East vs. west in the US: chemical characteristics of $PM_{2.5}$ during the winter of 1999. Aerosol Sci. Technol. 34: 88-96.

Török, S. B.; Lábár, J.; Schmeling, M.; Van Grieken, R. E. (1998) X-ray spectrometry. Anal. Chem. 70: 495R-518R.

Tsai, C.-J.; Cheng, Y.-H. (1996) Comparison of two ambient beta gauge PM_{10} samplers. J. Air Waste Manage. Assoc. 46: 142-147.

Tsai, C.-J.; Huang, H.-Y. (1995) Atmospheric aerosol sampling by an annular denuder system and a high-volume PM_{10} sampler. Environ. Int. 21: 283-291.

Turner, D. B. (1994) Workbook of atmospheric dispersion estimates - an introduction to dispersion modeling. 2nd ed. Boca Raton, FL: Lewis Publishers.

Turner, J. R.; Hering, S. V. (1987) Greased and oiled substrates as bounce-free impaction surfaces. J. Aerosol Sci. 18: 215-224.

Turpin, B. J.; Huntzicker, J. J. (1991) Secondary formation of organic aerosol in the Los Angeles basin: a descriptive analysis of organic and elemental carbon concentrations. Atmos. Environ. Part A 25: 207-215.

Turpin, B. J.; Huntzicker, J. J. (1995) Identification of secondary organic aerosol episodes and quantitation of primary and secondary organic aerosol concentrations during SCAQS. Atmos. Environ. 29: 3527-3544.

Turpin, B. J.; Cary, R. A.; Huntzicker, J. J. (1990) An in situ, time-resolved analyzer for aerosol organic and elemental carbon. Aerosol Sci. Technol. 12: 161-171.

Turpin, B. J.; Liu, S.-P.; Podolske, K. S.; Gomes, M. S. P.; Eisenreich, S. J.; McMurry, P. H. (1993) Design and evaluation of a novel diffusion separator for measuring gas/particle distributions of semivolatile organic compounds. Environ. Sci. Technol. 27: 2441-2449.

Turpin, B. J.; Huntzicker, J. J.; Hering, S. V. (1994) Investigation of organic aerosol sampling artifacts in the Los Angeles basin. Atmos. Environ. 28: 3061-3071.

Turpin, B. J.; Saxena, P.; Andrews, E. (2000) Measuring and simulating particulate organics in the atmosphere: problems and prospects. Atmos. Environ. 34: 2983-3013.

Tyler, G. (1992) ICP vs. AA: how do you choose? Environ. Lab. 31-34.

U.S. Court of Appeals, District of Columbia Circuit. (1999) American Trucking Associations, Inc. vs. U.S. Environmental Protection Agency. 175 F3d 1027 (CA DC, 97-1440).

U.S. Environmental Protection Agency. (1982) Air quality criteria for particulate matter and sulfur oxides. Research Triangle Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; EPA report no. EPA-600/8-82-029aF-cF. 3v. Available from: NTIS, Springfield, VA; PB84-156777.

U.S. Environmental Protection Agency. (1996a) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.

U.S. Environmental Protection Agency. (1996b) Review of the national ambient air quality standards for particulate matter: policy assessment of scientific and technical information. OAQPS staff paper. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA/452/R-96-013. Available from: NTIS, Springfield, VA; PB97-115406REB.

U.S. Environmental Protection Agency (1998) Monitoring $PM_{2.5}$ in ambient air using designated reference of class I equivalent methods. In *Quality Assurance Handbook for Air Pollution Measurement Systems*, Volume II, Part II, Section 2.12.

U.S. Environmental Protection Agency. (1999) Particulate matter ($PM_{2.5}$) speciation guidance. Final draft. Edition 1. Research Triangle Park, NC: Office of Air Quality Planning and Standards. Available: www.epa.gov/ttn/amtic/files/ambient/pm25/spec/specfinl.pdf [2000, October 11].

U.S. Environmental Protection Agency. (2001) List of designated reference and equivalent methods. Available: http://www.epa.gov/ttn/amtic/files/ambient/criteria/repmlist.pdf [05 March 02].

U.S. Environmental Protection Agency. (2002a) Ambient monitoring technology information center (AMTIC). Washington, DC: Office of Air Quality Planning and Standards; Monitoring and Quality Assurance Group. Available: http://www.epa.gov/ttn/amtic/ (20 March 2002).

U.S. Environmental Protection Agency. (2002b) Support center for regulatory air models. Washington, DC: Office of Air Quality Planning and Standards. Available: http://www.epa.gov/scram001 (21 March 2002).

Van Loy, M.; Bahadori, T.; Wyzga, R.; Hartsell, B.; Edgerton, E. (2000) The Aerosol Research and Inhalation Epidemiology Study (ARIES): $PM_{2.5}$ mass and aerosol component concentrations and sampler intercomparisons. J. Air Waste Manage. Assoc. 50: 1446-1458.

Vanderpool, R. W.; Peters, T. M.; Natarajan, S.; Gemmill, D. B.; Wiener, R. W. (2001) Evaluation of the loading characteristics of the EPA WINS $PM_{2.5}$ separator. Aerosol Sci. Technol. 34: 444-456.

Vasilou, J. G.; Sorenson, D.; McMurry, P. H. (1999) Sampling at controlled relative humidity with a cascade impactor. Atmos. Environ. 30: 1049-1056.

Venkataraman, C.; Lyons, J. M.; Friedlander, S. K. (1994) Size distributions of polycyclic aromatic hydrocarbons and elemental carbon. 1. Sampling, measurement methods, and source characterization. Environ. Sci. Technol. 28: 555-562.

Vincent, J. H. (1989) Aerosol sampling: science and practice. New York, NY: Wiley.

Vincent, J. H. (1995) Aerosol science for industrial hygienists. Oxford, United Kingdom: Pergamon.

Wang, P. Y. (1997) Continuous aerosol mass measurement by flow obstruction. Boston, MA: Harvard University, School of Public Health.

Wang, H.-C.; John, W. (1987) Comparative bounce properties of particle materials. Aerosol Sci. Technol. 7: 285-299.

Wang, H.-C.; John, W. (1988) Characteristics of the Berner impactor for sampling inorganic ions. Aerosol Sci. Technol. 8: 157-172.

Ward, M. D.; Buttry, D. A. (1990) In situ interfacial mass detection with piezoelectric transducers. Science 249: 1000-1007.

Watson, J. G.; Chow, J. C.; Richards, L. W.; Neff, W. D.; Andersen, S. R.; Dietrich, D. L.; Houck, J. E.; Olmez, I. (1988a) The 1987-88 metro Denver brown cloud study volume II: measurements. Reno, NV: Desert Research Institute; final report no. 8810.1F2 (2 of 3).

Watson, J. G.; Chow, J. C.; Richards, L. W.; Neff, W. D.; Andersen, S. R.; Dietrich, D. L.; Houck, J. E.; Olmez, I. (1988b) The 1987-88 metro Denver brown cloud study volume III: data interpretation. Reno, NV: Desert Research Institute; final report no. 8810.1F3 (3 of 3).

Watson, J. G.; Robinson, N. F.; Chow, J. C.; Henry, R. C.; Kim, B. M.; Pace, T. G.; Meyer, E. L.; Nguyen, Q. (1990a) The USEPA/DRI chemical mass balance receptor model, CMB 7.0. Environ. Software 5: 38-49.

Watson, J. G.; Robinson, N. F.; Chow, J. C.; Henry, R. C.; Kim, B.; Nguyen, Q. T.; Meyer, E. L.; Pace, T. G. (1990b) Receptor model technical series, v. III (1989 revision): CMB7 user's manual. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; report no. EPA-450/4-90-004. Available: http://www.epa.gov/scram001/ [19 February 2002].

Watson, J. G.; Chow, J. C.; Lu, Z.; Fujita, E. M.; Lowenthal, D. H.; Lawson, D. R.; Ashbaugh, L. L. (1994) Chemical mass balance source apportionment of PM_{10} during the Southern California Air Quality Study. Aerosol Sci. Technol. 21: 1-36.

Watson, J. G.; Fujita, E. M.; Chow, J. C.; Zielinska, B.; Richards, L. W.; Neff, W.; Dietrich, D. (1998) Northern front range air quality study. Final report. Fort Collins, CO: Colorado State University, Cooperative Institute for Research in the Atmosphere. Available: http://www.nfraqs.colostate.edu/index2.html (16 Jan 2002).

Watson, J. G.; Chow, J. C.; Frazier, C. A. (1999) X-ray fluorescence analysis of ambient air samples. In: Landsberger, S.; Creatchman, M., eds. Elemental Analysis of Airborne Particles. Newark, NJ: Gordon and Breach; pp. 67-96.

Weber, R. J.; Orsini, D.; Daun, Y.; Lee, Y.-N.; kotz, P. J.; Brechtel, F. (2001) A particle-into-liquid collector for rapid measurement of aerosol bulk chemical composition. Aerosol Sci. Technol. 35: 718-727.

Wedding, J. B.; Weigand, M. A. (1993) An automatic particle sampler with beta gauging. J. Air Waste Manage. Assoc. 43: 475-479.

Weingartner, E.; Burtscher, H.; Baltensperger, U. (1997) Hygroscopic properties of carbon and diesel soot particles. Atmos. Environ. 31: 2311-2327.

Weschler, C. J.; Shields, H. C. (1999) Indoor ozone/terpene reactions as a source of indoor particles. Atmos. Environ. 33: 2301-2312.

Wesolowski, J. J.; John, W.; Devor, W.; Cahill, T. A.; Feeney, P. J.; Wolfe, G.; Flocchini, R. (1977) Collection surfaces of cascade impactors. In: Dzubay, T. G., ed. X-ray fluorescence analysis of environmental samples. Ann Arbor, MI: Ann Arbor Science Publishers, Inc.; pp. 121-131.

Whitby, K. T. (1978) The physical characteristics of sulfur aerosols. Atmos. Environ. 12: 135-159.

Whitby, K. T.; Sverdrup, G. M. (1980) California aerosols: their physical and chemical characteristics. In: Hidy, G. M.; Mueller, P. K.; Grosjean, D.; Appel, B. R.; Wesolowski, J. J., eds. The character and origins of smog aerosols: a digest of results from the California Aerosol Characterization Experiment (ACHEX). New York, NY: John Wiley & Sons, Inc.; pp. 477-517. (Advances in environmental science and technology: v. 9).

Whitby, K. T.; Charlson, R. E.; Wilson, W. E.; Stevens, R. K. (1974) The size of suspended particle matter in air. Science (Washington, DC) 183: 1098-1099.

White, W. H. (1998) Statistical considerations in the interpretation of size-resolved particulate mass data. J. Air & Waste Manage. Assoc. 48: 454-458.

Willeke, K.; Baron, P. A., eds. (1993) Aerosol measurement: principles, techniques and applications. New York, NY: Van Nostrand Reinhold Publishers.

Willeke, K.; Whitby, K. T. (1975) Atmosperic aerosols: size distribution interpretation. J. Air Pollut. Control Assoc. 25: 529-534.

Williams, E. L., II; Grosjean, D. (1990) Removal of atmospheric oxidants with annular denuders. Environ. Sci. Technol. 24: 811-814.

Williams, R.; Suggs, J.; Zweidinger, R.; Evans, G.; Creason, J.; Kwok, R.; Rodes, C.; Lawless, P.; Sheldon, L. (2000) The 1998 Baltimore Particulate Matter Epidemiology-Exposure Study: part 1. Comparison of ambient, residential outdoor, indoor and apartment particulate matter monitoring. J. Exposure Anal. Environ. Epidemiol. 10: 518-532.

Wilson, W. E. (1995) Aerosol exposure, physics, and chemistry. Inhalation Toxicol. 7: 769-772.

Wilson, W. E.; Suh, H. H. (1997) Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies. J. Air Waste Manage. Assoc. 47: 1238-1249.

Wilson, W. E.; Spiller, L. L.; Ellestad, T. G.; Lamothe, P. J.; Dzubay, T. G.; Stevens, R. K.; Macias, E. S.; Fletcher, R. A.; Husar, J. D.; Husar, R. B.; Whitby, K. T.; Kittelson, D. B.; Cantrell, B. K. (1977) General Motors sulfate dispersion experiment: summary of EPA measurements. J. Air Pollut. Control Assoc. 27: 46-51.

Woo, K. S.; Chen, D. R.; Pui, D. Y. H.; McMurry, P. H. (2001a) Measurement of Atlanta aerosol size distributions: observations of ultrafine particle events. Aerosol Sci. Technol. 34: 75-87.

Woo, K.-S.; Chen, D.-R.; Pui, D. Y. H.; Wilson, W. E. (2001b) Use of continuous measurements of integral aerosol parameters to estimate particle surface area. Aerosol Sci. Technol. 34: 57-65.

Wu, C. C.; Suarez, A. E.; Lin, Z.; Kidwell, C. B.; Borgoul, P. V.; Caffrey, P. F.; Ondov, J. M.; Sattler, B. (1998) Application of an Ir tracer to determine soot exposure to students commuting to school on Baltimore public buses. Atmos. Environ. 32: 1911-1919.

Yamasaki, H.; Kuwata, K.; Miyamoto, H. (1982) Effects of ambient temperature on aspects of airborne polycyclic aromatic hydrocarbons. Environ. Sci. Technol. 16: 189-194.

Yatin, M.; Tuncel, S. G.; Tuncel, G.; Aras, N. K. (1994) Trace element composition of atmospheric aerosols in Ankara, Turkey, determined by instrumental neutron activation analysis. J. Radioanal. Nucl. Chem. 181: 401-411.

Zeng, X.; Wu, X.; Yao, H.; Yang, F.; Cahill, T. A. (1993) PIXE-induced XRF with transmission geometry. Nucl. Instr. Meth. Phys. Res. B 75: 99-104.

Zhang, X. Q.; McMurry, P. H. (1987) Theoretical analysis of evaporative losses from impactor and filter deposits. Atmos. Environ. 21: 1779-1789.

Zhang, X.; McMurry, P. H. (1992) Evaporative losses of fine particulate nitrates during sampling. Atmos. Environ. Part A 26: 3305-3312.

Zhang, X. Q.; McMurry, P. H.; Hering, S. V.; Casuccio, G. S. (1993) Mixing characteristics and water content of submicron aerosols measured in Los Angeles and at the Grand Canyon. Atmos. Environ. Part A 27: 1593-1607.

Zoller, W. H.; Gordon, G. E. (1970) Instrumental neutron activation analysis of atmospheric pollutants utilizing Ge(Li) "gamma"-ray detectors. Anal. Chem. 42: 257-265.

3. CONCENTRATIONS, SOURCES, AND EMISSIONS OF ATMOSPHERIC PARTICULATE MATTER

3.1 INTRODUCTION

This chapter discusses topics covered in Chapter 5 (Sources and Emissions of Atmospheric Particles) and Chapter 6 (Environmental Concentrations) of the previous document, Air Quality Criteria for Particulate Matter or "1996 PM AQCD" (U.S. Environmental Protection Agency, 1996) and presents updates to these materials where available.

Information about concentrations, the composition, and the spatial and temporal variability of ambient particles across the United States is presented in Section 3.2. Ambient concentration data obtained during the first two years of operation of the recently deployed nationwide network of Federal Reference Method $PM_{2.5}$ monitors in twenty-seven metropolitan statistical areas (MSAs) are presented and analyzed in Appendix 3A. Initial data from the pilot method evaluation study for the national speciation network are presented in Appendix 3B. Results of field studies that have characterized the composition of organic compounds in ambient particles are summarized in Appendix 3C to complement the data for the inorganic composition of ambient particles presented in Appendix 6A of the 1996 PM AQCD and Appendix 3B of this document. Data for characterizing the daily and seasonal variability of $PM_{2.5}$ concentrations are discussed in Section 3.2.1, the intraday variability of $PM_{2.5}$ concentrations in Section 3.2.2, the relations among different size fractions in Section 3.2.3, the interrelations and correlations among PM components in Section 3.2.4, and the spatial variability of various PM components in Section 3.2.5.

Unlike gaseous criteria pollutants (SO_2, NO_2, CO, O_3), which are well-defined chemical entities, atmospheric particulate matter (PM) is composed of a variety of particles differing in size and chemical composition. Therefore, sources of each component of the atmospheric aerosol must be considered in turn. Differences in the composition of particles emitted by different sources also will lead to spatial and temporal heterogeneity in the composition of the atmospheric aerosol. The nature of the sources and the composition of the emissions from these sources are discussed in Section 3.3. The chemistry of formation of secondary PM from gaseous

precursors is discussed in Section 3.3.1. The long-range transport of PM from sources outside the United States is discussed in Section 3.3.2. Reviews of transport of PM and its precursors within the United States can be found in the NARSTO Fine Particle Assessment (NARSTO, 2002). More detailed information regarding sulfur and nitrogen species can be found in Hidy (1994). Estimates of contributions of various sources to ambient PM levels given by source apportionment studies also are presented in Section 3.3.3. More detailed information about the composition of emissions from various sources is given in Appendix 3D. Because PM is composed of both primary and secondary constituents, emissions of both the primary components and the gaseous precursors of secondary PM must be considered. Nationwide emissions estimates of primary PM and precursors to secondary PM are discussed in Section 3.3.4 and uncertainties in emissions estimates in Section 3.3.5.

The organization of topics in this chapter (ambient measurements, source characterization and apportionment, and emissions inventories) reflects, in a broad sense, the order in which these topics are addressed in scientific studies and, arguably, the increasing levels of uncertainty that are associated with these topics.

3.2 PATTERNS AND TRENDS IN AMBIENT PM CONCENTRATIONS

A significant amount of data for characterizing PM_{10} mass concentrations and trends exists, and that available up to about 1994 was presented in the 1996 PM AQCD. However, data sets for characterizing $PM_{2.5}$ and $PM_{10-2.5}$ mass or trends were not as extensive. Sources of data for $PM_{2.5}$ (fine) and $PM_{10-2.5}$ (coarse), which were discussed in the 1996 PM AQCD, include EPA's Aerometric Information Retrieval System (AIRS) (U.S. Environmental Protection Agency, 2000a), IMPROVE (Eldred and Cahill, 1994; Cahill, 1996), the California Air Resources Board (CARB) Data Base (California Air Resources Board, 1995), the Harvard Six-Cities Data Base (Spengler et al., 1986; Neas, 1996), and the Harvard Philadelphia Data Base (Koutrakis, 1995). The Inhalable Particulate Network (IPN) (Inhalable Particulate Network, 1985; Rodes and Evans, 1985) provided TSP, PM_{15}, and $PM_{2.5}$ data but only a small amount of PM_{10} data.

New sources of PM data include the recently deployed nationwide $PM_{2.5}$ compliance monitoring network, which provides mass measurements using a Federal Reference Method (FRM). This section summarizes data obtained during 1999 and 2000 by this network and

provides an approximate characterization of nationwide $PM_{10-2.5}$ concentrations by comparing PM_{10} to $PM_{2.5}$ measurements at sites where both types of compliance monitors are located. Various aspects of these data are presented in greater detail in Appendix 3A. In addition, a small number of recent studies in which daily mass and composition measurements are available for extended periods are discussed in this section. The results of quality assured aerosol composition data obtained by X-ray fluorescence (XRF) and by analyses of organic carbon (OC) and elemental carbon (EC) for thirteen urban areas from the methods evaluation study for the national $PM_{2.5}$ speciation network are presented in Appendix 3B.

Organic compounds contribute from 10 to 70% of the dry fine particle mass in the atmosphere (see Appendix 3C). However, organic PM concentrations, composition, and formation mechanisms are poorly understood. Particulate organic matter is an aggregate of hundreds of individual compounds spanning a wide range of chemical and thermodynamic properties (Saxena and Hildemann, 1996). Some of the organic compounds are "semivolatile" (i.e., they have atmospheric concentrations and saturation vapor pressures such that both gaseous and condensed phases exist in equilibrium in the atmosphere). The presence of semivolatile or multiphase organic compounds complicates the sampling process. Organic compounds originally in the gas phase may be absorbed on glass or quartz filter fibers and create a positive artifact. Conversely, semivolatile compounds originally present in the condensed phase may evaporate from particles collected on glass, quartz, or Teflon filters creating a negative artifact. In addition, no single analytical technique is currently capable of analyzing the entire range of organic compounds present in atmospheric PM. Rigorous analytical methods are able to identify only 10 to 20% of the organic PM mass on the molecular level (Rogge et al., 1993), and only about 50% of the condensed phase compounds could be identified in smog chamber studies of specific compounds (Forstner et al., 1997a,b). Measurement techniques are discussed in Section 2.2.3.2. Information on the identification and concentration of the many different organic compounds identified in atmospheric samples obtained during the 1990s is given in Appendix 3C.

Summary tables giving the results of 66 field studies that obtained data for the composition of particles in the $PM_{2.5}$, $PM_{10-2.5}$, or PM_{10} size ranges were presented in Appendix 6A of the 1996 PM AQCD. The summary tables include data for mass, organic carbon, elemental carbon, nitrate, sulfate, and trace elements. Data from the studies were presented for the eastern, western,

and central United States. It should be noted that these studies took place at various times and lasted for various durations over a 20-year period, and there may have been significant changes in the concentrations of many species between the times when these studies were conducted and now. These changes resulted from a number of factors (e.g., pollution controls, technological advances, land use changes, etc).

There were a number of discernible differences in the composition of particles across the United States evident in the data sets listed in Appendix 6A in the 1996 PM AQCD (cf. Figures 6-85a to 6-85c). However, these differences can only be discussed in the context of the uncertainties in the measurements of the main components (sulfate, organic carbon, elemental carbon, crustal material, ammonium, and nitrate). Sulfate, followed by crustal materials have the smallest uncertainties associated with their measurement among all the components listed. Sulfate constituted about 38% of $PM_{2.5}$ in the aerosol composition studies in the eastern United States and was the major identifiable component of $PM_{2.5}$, but it constituted only about 11% of $PM_{2.5}$ in the studies listed for the western United States. The contribution of crustal materials to $PM_{2.5}$ ranged from about 4% in the East to about 15% in the West. The contribution of unidentified material (possibly consisting mostly of water of hydration) ranged from 23% in the East to 0% in the West. The contribution of elemental carbon to $PM_{2.5}$ ranged from about 4% in the East to about 15% in the West. Organic compounds constituted about 21% of $PM_{2.5}$ in the eastern United States, ranging to about 39% for the studies listed in the western United States. However, uncertainties for organic carbon, elemental carbon, ammonium, and nitrate are larger than for sulfate and crustal material. A factor of 1.4 was used to account for the presence of oxygen and nitrogen in the organic compounds. This factor may vary among different areas and may represent the lowest reasonable estimate for an urban aerosol (Turpin and Lim, 2001). In addition, the samples collected in the studies were subject to a number of sampling artifacts involving the adsorption of gases and the evaporation of volatile components that either formed on the filters or were present in the ambient particles. The values reported for organic carbon and elemental carbon in filter samples depend strongly on the specific analysis method used (Chow et al., 2001).

Crustal materials constitute from 52% of $PM_{10-2.5}$ in the eastern United States to 70% of $PM_{10-2.5}$ in the studies in the western United States given in Appendix 6A of CD96. The fraction of unidentified material in $PM_{10-2.5}$ varied from 41% in the eastern United States to 27% in the

western United States. However, in the vast majority of these studies no attempt was made to characterize organic components or nitrate in the $PM_{10-2.5}$ size fraction. Even if analyses of total OC were available, they would not be able to distinguish between bioaerosols and simple organic compounds. Indeed, in many photomicrographs of $PM_{10-2.5}$ samples obtained by scanning electron microscopy, the fields of view were dominated by large numbers of pollens, plant and insect fragments, and microorganisms. Bioaerosols such as pollens, fungal spores, and most bacteria are expected to be found mainly in the coarse size fraction. However, allergens from pollens can also be found in fine particles (Monn, 2001). It should also be remembered that a small fraction (typically about 10%) of $PM_{2.5}$ is entrained into the flow of the channel of the dichotomous sampler that collects the $PM_{10-2.5}$ sample and that there may be errors invoked during the procedure used to account for this entrainment.

Data for the chemical composition of particles in a number of national parks and remote areas have been collected for a number of years by the IMPROVE network. Concentrations are reported for sulfate, nitrate, light absorbing carbon, organic carbon, and soil components. With the collection of compositional data by the speciation network, more synoptic (i.e., concurrent) coverage will be obtained for these constituents in continental background to urban environments across the United States.

PM_{10} Concentrations and Trends

Nationwide PM_{10} annual mean concentrations on a county-wide basis from the AIRS database for calendar years 1999 and 2000 are shown in Figure 3-1a. Concentrations in most areas of the country were below the level of the PM_{10} annual standard (50 $\mu g/m^3$) in 1999 and 2000. The median annual PM_{10} concentration was about 23 $\mu g/m^3$; and five percent of the countywide concentrations shown in Figure 3-1a were greater than 35 $\mu g/m^3$. The 98th percentile PM_{10} concentrations are shown in Figure 3-1b. Data from all monitors for the most recently available four consecutive quarters in 1999 and 2000 with at least eleven valid observations per quarter in a given county were averaged to produce Figure 3-1a; and data from the highest monitor in that county were used to produce Figure 3-1b. In these, and similar maps for $PM_{2.5}$ and $PM_{10-2.5}$, cut points were chosen at the median and 95th percentile concentrations. As shown by the blank areas on the maps, the picture is not complete because some monitoring locations did not record valid data for all four quarters or recorded fewer than 11 samples in one or more



Figure 3-1a. 1999-2000 county-wide average annual mean PM_{10} concentrations ($\mu g/m^3$).

Source: U.S. EPA Aerometric Information Retrieval System (12/19/2001).



Figure 3-1b. 1999-2000 highest county-wide 98th percentile 24-h average PM_{10} concentrations ($\mu g/m^3$).

Source: U.S. EPA Aerometric Information Retrieval System (12/19/2001).

quarters or counties simply did not have monitors. Similar considerations apply to the maps to be shown later for $PM_{2.5}$ and $PM_{10-2.5}$. It should also be noted that the area of counties can be much greater in the West than in the East. As a result, the density of monitors may appear to be greater in the West and air quality may appear to be worse over much larger areas in the West than in the East.

Nationwide trends in annual mean PM_{10} concentrations from 1990 through 1999 (based on data obtained at 153 rural sites, 375 suburban sites, and 408 urban sites reporting to AIRS) are shown in Figure 3-2 (U.S. Environmental Protection Agency, 2001). Though average concentration levels differ among sites, with higher levels at urban and suburban sites, the nationwide data set shows a decrease of 18% that occurred mainly during the first half of the record. PM_{10} concentrations then leveled off during the last few years of the record. Figure 3-3 shows the annual mean PM_{10} trend summarized by EPA region. Decreases in annual average PM_{10} concentrations from 1990 to 1999 were largest in the Northwest (10.3 $\mu g/m^3$) and smallest in the south central United States (3.2 $\mu g/m^3$). Analyses of available TSP measurements obtained since 1950 indicate that mean TSP concentrations could have declined two- to three-fold in urban areas between 1950 and 1980 (Lipfert, 1998).

$PM_{2.5}$ Concentrations and Trends

Nationwide annual mean $PM_{2.5}$ concentrations for 1999 and 2000 are shown in Figure 3-4a and 98th percentile concentrations are shown in Figure 3-4b. Quantities shown in Figure 3-4a and 3-4b were calculated for individual counties. Data from all monitors in a given county meeting the same minimum data completeness criteria for PM_{10} (given earlier) were averaged to produce Figure 3-4a, and results from the highest monitor were used to produce Figure 3-4b. The median $PM_{2.5}$ concentration nationwide was about 13 $\mu g/m^3$. Annual mean $PM_{2.5}$ concentrations were above 18 $\mu g/m^3$ at 5% of the sites, mainly in California and in the southeastern United States. The 98th percentile 24-h average concentrations were below 50 $\mu g/m^3$ at 95% of the sites sampled. Most of the sites with levels above 50 $\mu g/m^3$ are located in California.

Annual average $PM_{2.5}$ concentrations obtained as part of health studies conducted in various locations in the United States and Canada from the late 1980s to the early 1990s are shown in Figure 3-5 (Bahadori et al., 2000a). These studies include the Harvard six-cities study (Steubenville, OH; Watertown, MA; Portage, WI; Topeka, KS; St. Louis, MO; and Kingston-



Figure 3-2. Nationwide trend in ambient PM_{10} concentration from 1990 through 1999.

Source: U.S. Environmental Protection Agency (2001).



Figure 3-3. Trend in PM_{10} annual mean concentrations by EPA region, 1990 through 1999 ($\mu g/m^3$).

Source: U. S. Environmental Protection Agency (2001).



Figure 3-4a. 1999-2000 county-wide average annual mean $PM_{2.5}$ concentrations ($\mu g/m^3$).

Source: U.S. EPA Aerometric Information Retrieval System (12/19/2001).



Figure 3-4b. 1999-2000 highest county-wide 98th percentile 24-h average $PM_{2.5}$ concentrations ($\mu g/m^3$).

Source: U.S. EPA Aerometric Information Retrieval System (12/19/2001).



Figure 3-5. Collection of annual distribution of 24-h average $PM_{2.5}$ concentrations observed in U.S. and Canadian health studies conducted during the 1980's and early 1990's.

Source: Bahadori et al. (2000a).

DRAFT–DO NOT QUOTE OR CITE

Harriman, TN); PTEAMS (Riverside, CA); MAACS (Philadelphia, PA; Washington, DC; and Nashville, TN); South Boston Air Quality and Source Apportionment Study (Boston, MA); and NPMRMN (Phoenix, AZ). The remaining sites were part of the 24-cities study (Spengler et al., 1996).

Sufficient data are not yet available to permit the calculation of nationwide trends of $PM_{2.5}$ and $PM_{10-2.5}$; however, some general conclusions can be reached. Darlington et al. (1997) proposed that the consistent reductions in PM_{10} concentrations found in a wide variety of environments ranging from urban to rural may have resulted from common factors or controls that affected fine particles more strongly than coarse particles. This is because fine particles have longer atmospheric lifetimes than coarse particles and can be transported over longer distances and, hence, can affect larger areas. Apart from the IMPROVE network of monitoring sites located mainly in national parks, the longest time series of $PM_{2.5}$ concentration and composition data have been obtained by the California Air Resources Board (CARB). Their data show that annual average $PM_{2.5}$ concentrations decreased by about 50% in the South Coast Air Basin, 35% in the San Joaquin Valley, 30% in the San Francisco Bay Area, and 35% in the Sacramento Valley from 1990 to 1995 (Dolislager and Motallebi, 1999). $PM_{2.5}$ data were collected continuously from 1994 to 1998 as part of the children's health study in 12 communities in southern California (Taylor et al., 1998). Data obtained at all sites show decreases in $PM_{2.5}$ ranging from 2% at Santa Maria to 37% at San Dimas/Glendora from 1994 through 1998. These decreases were accompanied by decreases in major components such as nitrate, sulfate, ammonium, and acids. Based on the analysis of $PM_{2.5}$ data sets collected prior to 1990, Lipfert (1998) found that $PM_{2.5}$ concentrations could have decreased by about 5% per year from 1970 to 1990 in a number of urban areas. These declines were also found to be consistent with decreases in emissions from combustion sources over that time period.

Background $PM_{2.5}$ Concentrations

In common usage, the term "background concentrations" refers to concentrations observed in remote areas relatively unaffected by local pollution sources. However, as noted in Chapter 6 of the 1996 PM AQCD, several definitions of background concentrations are possible. In that document, the two definitions chosen as being most relevant for regulatory purposes are based on estimates of contributions from uncontrollable sources that can affect concentrations in the

United States. The first definition refers to the concentration resulting from anthropogenic and natural emissions outside North America and natural sources within North America. The second definition refers to the concentration resulting from natural sources only within and outside of North America. Because of long-range transport from anthropogenic source regions in North America, it is impossible to obtain background concentrations as defined above solely on the basis of direct measurement in remote areas in North America. However, these data can be used to place reasonable upper limits on what these concentrations could be. The range of values in the lowest 5th percentile annual mean $PM_{2.5}$ concentrations in the AIRS data base is from 2.8 $\mu g/m^3$ to 6.9 $\mu g/m^3$. This range of concentrations is consistent with the range of annual mean $PM_{2.5}$ concentrations at remote sites in the western United States obtained from 1996 through 1999 in the IMPROVE network. At most IMPROVE sites in the western United States, the mean concentration of $PM_{10-2.5}$ is higher than that of $PM_{2.5}$, and $PM_{2.5}$ concentrations are moderately correlated (r = 0.72) with $PM_{10-2.5}$ concentrations. In contrast, $PM_{2.5}$ concentrations are higher than those of $PM_{10-2.5}$ at IMPROVE sites in the eastern United States, and $PM_{2.5}$ concentrations are only weakly correlated (r = 0.26) with those of $PM_{10-2.5}$.

Annual average natural background concentrations of PM_{10} (according to definition 1) have been estimated to range from 4 to 8 $\mu g/m^3$ in the western United States and 5 to 11 $\mu g/m^3$ in the eastern United States. Corresponding $PM_{2.5}$ levels have been estimated to range from 1 to 4 $\mu g/m^3$ in the western United States and from 2 to 5 $\mu g/m^3$ in the eastern United States (U.S. Environmental Protection Agency, 1996). Although these values are broadly consistent with the data given above, the data discussed in the previous paragraph represent only upper limits to background concentrations because of possible contributions from long-range transport from anthropogenic sources within North America. Peak 24-h average natural background concentrations may be substantially higher than the annual or seasonal average natural background concentrations, especially within areas affected by wildfires and dust storms and long range transport from outside North America. Estimates of background concentrations according to definition 2 are not yet available. However, recent information about contributions to background concentrations that fall under definitions 1 and 2 because of long-range transport from sources outside the United States is given in Section 3.3.2.

$PM_{10-2.5}$ Concentrations

By using AIRS data for 1999 and 2000 obtained by the PM_{10} and $PM_{2.5}$ compliance networks, it is possible to construct a picture of the distribution of coarse PM across the country. This is accomplished by pairing data from 228 compliance monitoring sites where PM_{10} and $PM_{2.5}$ monitors are collocated and subtracting the mass concentrations of $PM_{2.5}$ from PM_{10}. Nationwide annual mean $PM_{10-2.5}$ concentrations calculated by this difference method are shown in Figure 3-6a. Annual mean $PM_{10-2.5}$ concentrations range from 1 to 48 $\mu g/m^3$, with a nationwide median concentration of about 10 $\mu g/m^3$; and 5% of the sites had mean concentrations greater than 20 $\mu g/m^3$. The higher values occur mainly in the western United States, particularly in California. The highest county-wide 98th percentile $PM_{10-2.5}$ concentrations based on this same data set are shown in Figure 3-6b. Highest values in the western United States are caused by dust raised locally either by natural means or by anthropogenic activity. Elevated dust levels are also found in southern Florida as the result of dust storms in North Africa (cf. Section 3.3.2). In many areas, combined errors in the $PM_{2.5}$ and PM_{10} measurements may be similar to or even greater than $PM_{10-2.5}$ concentrations. Because of this and other potential problems with this approach (cf. Section 3.2.1), these results should be viewed with caution.

3.2.1 Seasonal Variability in PM Concentrations

$PM_{2.5}$

Aspects of the spatial and temporal variability of $PM_{2.5}$ concentrations for 1999 and 2000 in a number of metropolitan areas across the United States are presented in this and following subsections. Data for multiple sites in 27 urban areas across the United States have been obtained from the AIRS data base and analyzed for their seasonal variations and for their spatial correlations and spatial uniformity in concentrations (Pinto, et al., 2002). Only 27 MSAs were included in the analyses based on the criteria that data be obtained on at least 15 days in each calendar quarter of either 1999 and 2000 or 2000 alone at four sites within that MSA. A number of aspects of the spatial and temporal variability of the 1999 $PM_{2.5}$ data set were presented in Rizzo and Pinto (2001), based in part on analyses given in Fitz-Simons et al. (2000).



Figure 3-6a. 1999-2000 estimated county-wide average annual mean $PM_{10-2.5}$ concentrations ($\mu g/m^3$).

Source: U.S. EPA Aerometric Information Retrieval System (12/19/2001).



Figure 3-6b. 1999-2000 estimated county-wide highest 98th percentile 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$).

Source: U.S. EPA Aerometric Information Retrieval System.

Information regarding the seasonal variability in $PM_{2.5}$ concentrations in four MSAs (Columbia, SC; Detroit, MI; Chicago, IL; Los Angeles-Long Beach, CA) in the United States is summarized in Figures 3-7a through 3-7d. These four urban areas were chosen to illustrate some general features of the spatial and temporal variability found in the United States. The figures show lowest, lower quartile, median, upper quartile, and highest concentrations for each calendar quarter of 1999 and 2000 for the Columbia, SC, and Los Angeles, CA MSAs and for 2000 for the Detroit, MI and Chicago, IL MSAs. For each monitoring site, the AIRS ID numbers, annual mean concentrations, the number of observations, and standard deviations are also shown. Data for multiple sites within these MSAs are shown to provide an indication of the degree of inter-site variability. Data for these MSAs and an additional twenty-three MSAs, criteria used for site selection, and additional descriptions of the data are given in Appendix 3A.

Annual mean $PM_{2.5}$ concentrations (based on two years data) at individual monitoring sites in the MSAs examined range from about 6 $\mu g/m^3$ to about 30 $\mu g/m^3$. The lowest values are found in rural portions of the MSAs examined, typically near the perimeter of the MSA. Annual mean concentrations tend to be higher in the Southeast than in the Northeast and higher in southern California compared to the Pacific Northwest (cf. Appendix 3A). However, average $PM_{2.5}$ concentrations tend to be lower in 1999 and 2000 in urban areas given in Appendix 3A compared to the concentrations observed during pollution-health outcome studies conducted in those five urban areas where these overlap (cf. Figure 3-5). It should be noted that there are no data demonstrating the comparability of the monitors used in the studies shown in Figure 3-5 and the FRM.

In four of the seven MSAs examined in the eastern United States (as in the Columbia, SC MSA, cf. Figure 3-7a), highest median concentrations occur at most sites during the third calendar quarter (i.e., summer months). There are exceptions to this pattern as shown for the Philadelphia, PA-NJ MSA (cf. Figure 3A-1). Highest median concentrations in the north-central United States tend to occur in the first or fourth quarters (i.e., winter months) as in the Detroit, MI and Chicago, IL MSA (cf. Figures 3-7b and 3-7c). Highest median concentrations occur during the fourth calendar quarter in MSAs in the western United States as in the Los Angeles, CA PMSA (cf. Figure 3-7d), although there are exceptions at individual sites in the Riverside, CA PMSA.



Figure 3-7a,b. Quarterly distribution of 24-h average $PM_{2.5}$ concentrations for selected monitors in the (a) Columbia, SC; (b) Detroit, MI; (c) Chicago, IL; and (d) Los Angeles, CA MSAs. Values for the lowest, lower quartile, median, upper quartile and highest concentrations are shown in the figures. The AIRS site ID number, annual concentration, number and standard deviation are shown above the figures for each site.





Figure 3-7c,d. Quarterly distribution of 24-h average $PM_{2.5}$ concentrations for selected monitors in the (a) Columbia, SC; (b) Detroit, MI; (c) Chicago, IL; and (d) Los Angeles, CA MSAs. Values for the lowest, lower quartile, median, upper quartile and highest concentrations are shown in the figures. The AIRS site ID number, annual concentration, number and standard deviation are shown above the figures for each site.

Source: Pinto et al. (2002).

Lowest median concentrations occur mainly during the first or fourth quarters at most sites in the eastern United States, with some occurring during the second quarter (cf. Appendix 3A). In moving westward, the seasonal pattern is not as distinct, with lowest median concentrations occurring in any quarter, but usually in the second or third quarter as in the Chicago, IL and Detroit, MI MSAs (cf. Figure 3-7b and 3-7c). In many of the MSAs examined, seasonal variations follow a similar pattern at all of the sites within the MSA, but in other MSAs there are noticeable differences in the seasonal pattern between sites. The large-scale differences in seasonal variability between MSAs tend to follow differences in the major categories of PM sources affecting the monitoring sites. Local heating by wood burning during the colder months is practiced more widely in the western United States than in the eastern United States. Hence, winter maxima and greater variability in $PM_{2.5}$ concentrations across sites are expected in the West due to the influence of the local sources. On the other hand, photochemical production of secondary PM, especially sulfate, occurs over wide areas in relatively homogeneous air masses during the summer months in the eastern United States. Because sulfates (along with associated cations and water) constitute the major fraction of summertime $PM_{2.5}$ in the East, there is greater uniformity in 3rd quarter PM concentrations within eastern MSAs (cf. Appendix 3A).

The highest values shown in the box plots in Figures 3-7a to 3-7d and in Figures 3A-1 to 3A-27 do not always follow the same seasonal pattern as do the median concentrations. These values likely reflect the existence of transient events such as forest fires (mainly in the West) or episodes of secondary PM production (mainly in the East). However, chemical analyses of filter samples or other evidence should be used to determine specific causes in particular locations.

There have been a few studies that have characterized $PM_{2.5}$ and PM_{10} concentrations in major urban areas. The Metropolitan Acid Aerosol Characterization Study (MAACS) (Bahadori et al., 2000b) characterized the levels and the spatial and temporal variability of $PM_{2.5}$, PM_{10}, and acidic sulfate concentrations in four cities in the eastern United States (Philadelphia, PA; Washington, D.C.; Nashville, TN; and Boston, MA). Seasonal variations in $PM_{2.5}$ and PM_{10} concentrations obtained during the course of this study are shown in Figure 3-8. The data for the four cities included in MAACS are presented as box plots showing the lowest, lowest tenth percentile, lowest quartile, median, highest quartile, highest tenth percentile, and highest $PM_{2.5}$ and PM_{10} values. Mean and highest $PM_{2.5}$ and PM_{10} concentrations are found during the summer in all four cities, although the seasonal pattern in Boston appears to be more nearly bimodal with



Figure 3-8. Concentrations of $PM_{2.5}$ and PM_{10} measured in the four MAACS cities. The data show the lowest, lowest tenth percentile, lowest quartile, median highest quartile, highest tenth percentile, and highest $PM_{2.5}$ 24-h average values. The dashed line shows the level of the annual $PM_{2.5}$ standard.

Source: Bahadori et al. (2000b).

an additional winter peak. This seasonal pattern, based on 2- to 3-year sampling periods for each city during 1992 through 1996, is in accord with that obtained from the FRM monitors in the NAMS and SLAMS network (cf. Appendix 3A).

$PM_{10-2.5}$

Data from the FRM $PM_{2.5}$ and PM_{10} compliance networks that could be used to characterize seasonal variations in $PM_{10-2.5}$ based on EPA minimum data completeness criteria (11 samples per calendar quarter) are available for 228 sites nationwide. Data for the seasonal variations in $PM_{10-2.5}$ concentrations for Columbia, SC and Detroit, MI are shown for 2000 and data for Los Angeles-Long Beach are shown for 1999 in Figures 3-9a,b,c. As can be seen by comparing the number of observations used in the calculation of values shown in Figures 3-7a,b,c,d and Figure 3-9a,b,c the number of days that could be used for calculating $PM_{10-2.5}$ concentrations is much less than that measured for $PM_{2.5}$. At least for the sites shown for Columbia, SC; and Detroit,



Figure 3-9a,b. Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations for selected sites in the (a) Columbia, SC; (b) Detroit, MI; and (c) Los Angeles, CA MSAs. Values for the lowest, lower quartile, median, upper quartile and highest concentrations are shown in the figures. The AIRS site ID number, annual concentration, number and standard deviation are shown above the figures for each site.



Figure 3-9c. Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations for selected sites in the (a) Columbia, SC; (b) Detroit, MI; and (c) Los Angeles, CA MSAs. Values for the lowest, lower quartile, median, upper quartile and highest concentrations are shown in the figures. The AIRS site ID number, annual concentration, number and standard deviation are shown above the figures for each site.

Source: Pinto et al. (2002).

MI the seasonal median maxima in concentrations occur during the second calendar quarter. It can readily be seen that a number of $PM_{10-2.5}$ concentrations are negative. (The negative estimates have been included in the calculation of mean concentrations.) There are a number of reasons for the negative concentration estimates, many of which arise because the ratios of $PM_{2.5}$ to PM_{10} are based on two independent measurements. Measurement imprecision plays a role when the ratios are large and concentrations are small. Differences in the behavior of semivolatile components in the two samplers could occur. The results may be due to errors in

sampler placement, field, laboratory, or data processing procedures. Therefore, caution should be exercised when attempting to interpret results for $PM_{10-2.5}$ based on the current network collocated $PM_{2.5}$ and PM_{10} monitors.

Frequency Distributions for $PM_{2.5}$ Data

Frequency distributions for $PM_{2.5}$ concentrations obtained in Philadelphia from 1992 through 1995 are shown in Figure 3-10 (data obtained by Bahadori et al., 2000b). Concentrations predicted from the log-normal distribution, using geometric mean values and standard deviations derived from the data, are also shown. In Philadelphia, the highest $PM_{2.5}$ values were observed when winds were from the southwest during sunny but hazy high pressure conditions. In contrast, the lowest values were found after significant rainstorms during all seasons of the year. Mean ± SD day-to-day concentration differences in the data set are 6.8 ± 6.5 $\mu g/m^3$ for $PM_{2.5}$ and 8.6 ± 7.5 $\mu g/m^3$ for PM_{10}. Maximum day-to-day concentration differences are 54.7 $\mu g/m^3$ for $PM_{2.5}$ and 50.4 $\mu g/m^3$ for PM_{10}.



Figure 3-10. Frequency distribution of 24-h average $PM_{2.5}$ concentrations measured at the Presbyterian home (PBY) monitoring site in southwestern Philadelphia from 1992 to 1995. Log-normal distribution fit to the data shown as solid line.

Different patterns are observed in data collected elsewhere in the United States. $PM_{2.5}$ concentrations obtained in Phoenix, AZ from 1995 through 1997 (Zweidinger et al., 1998) are summarized in Figure 3-11, and frequency distributions of $PM_{2.5}$ concentrations obtained in Phoenix are shown in Figure 3-12. Mean ±SD day-to-day concentration differences in this data set are 2.9 ± 3.0 $\mu g/m^3$ with a maximum day-to-day concentration difference of 23 $\mu g/m^3$. $PM_{2.5}$ and $PM_{10-2.5}$ data were obtained with dichotomous samplers at a number of sites in California on a sampling schedule of every 6 days from 1989 through 1998. Histograms showing the frequency distribution of the entire set of $PM_{2.5}$ and $PM_{10-2.5}$ concentrations obtained by the CARB network of dichotomous samplers from 1989 to 1998 are shown in Figures 3-13 and 3-14. Also shown are log-normal distributions generated using geometric means and standard deviations derived from the data as input. Although the data for both size fractions appear to be reasonably well simulated by the function, data obtained at individual locations may not be. Data showing the seasonal variability of $PM_{2.5}$ obtained at Riverside-Rubidoux are summarized in box plot form in Figure 3-15. The frequency distribution of $PM_{2.5}$ concentrations obtained at Riverside-Rubidoux from 1989 to 1994 is shown in Figure 3-16. It can be seen that the data are not as well fit by a log-normal distribution as are the data shown in Figure 3-10, partially as the result of a significant number of days when $PM_{2.5}$ concentrations are greater than 100 $\mu g/m^3$.

An examination of the data from the four MAACS cities, Phoenix, AZ, and Riverside, CA, indicates that substantial differences exist in aerosol properties between the eastern cities (MAACS) and the western cities (Phoenix, AZ; Riverside, CA). Fine-mode particles account for most of the PM_{10} mass observed in the MAACS cities and appear to drive the daily and seasonal variability in PM_{10} concentrations there. Coarse-mode particles represent a larger fraction of PM_{10} mass in Phoenix and Riverside and drive the seasonal variability in PM_{10} seen there. The average ratio of $PM_{2.5}$ to PM_{10} concentrations is much larger in the MAACS cities of Philadelphia (0.72); Washington, DC (0.74); and Nashville (0.63) than in either Phoenix (0.34) or Riverside (0.49). Differences between median and maximum concentrations in any size fraction are much larger at the Riverside site than at either the MAACS or Phoenix sites. Many of these differences could reflect the more sporadic nature of dust suspension at Riverside. In addition, the seasonal variability of $PM_{2.5}$ concentrations observed in Phoenix, AZ, and Riverside, CA, appears to be different from that observed in the MAACS cities. These



Figure 3-11. Concentrations of 24-h average $PM_{2.5}$ measured at the EPA site in Phoenix, AZ from 1995 to 1997. The data show the lowest, lowest tenth percentile, lowest quartile, median (black circles), highest quartile, highest tenth percentile, and highest $PM_{2.5}$ values.

Source: Zweidinger et al. (1998).



Figure 3-12. Frequency distribution of 24-h average $PM_{2.5}$ concentrations measured at the EPA site in Phoenix, AZ from 1995 to 1997.

Source: Zweidinger et al. (1998).



Figure 3-13. Frequency distribution of 24-h average $PM_{2.5}$ measurements obtained from all California Air Resources Board dichotomous sampler sites from 1989 to 1998.



Figure 3-14. Frequency distribution of 24-h average $PM_{10-2.5}$ concentrations obtained from all California Air Resource Board Dichotomous sampler sites from 1989 to 1998.



Figure 3-15. Concentrations of 24-h average $PM_{2.5}$ measured at the Riverside-Rubidoux site from 1989 to 1998. The data show the lowest, lowest tenth percentile, lowest quartile, median (black squares), highest quartile, highest tenth percentile, and highest $PM_{2.5}$ values.



Figure 3-16. Frequency distribution of 24-h average $PM_{2.5}$ concentrations measured at the Riverside-Rubidoux site from 1989 to 1994.

considerations demonstrate the hazards in extrapolating conclusions about the nature of variability in aerosol characteristics inferred at one location to another.

3.2.2 Diurnal (Circadian) Variability in PM Concentrations

The variability of PM concentrations on time scales shorter than a day can, in principle, be characterized by measurements made by continuous samplers (e.g, TEOMs and β-gauge monitors that are currently used to provide Air Quality Index [AQI] information to the public). A description of these methods was provided in Section 2.2.9. However, as shown in Chapter 2, continuous methods are subject to artifacts because, in large part, of heating of their inlets to remove water, which results in the loss of components such as ammonium nitrate and semivolatile organic compounds (cf. Sections 2.2.2.1 and 2.2.3 for further details concerning the chemistry of volatilizable components). Consequently, caution should be used in interpreting results obtained by these techniques.

The composite diurnal variation of $PM_{2.5}$ concentrations obtained throughout the continental United States by 31 TEOM and β-gauge monitors reporting to AIRS in 1999 is shown in Figure 3-17. As can be seen, there is a distinct pattern with maxima occurring during the morning and evening. Notable exceptions to this pattern occur in California, where broad nighttime maxima and daytime minima occur, which may be related to the use of β-gauges with unheated inlets there. It should be noted in examining the diurnal variations shown in Figure 3-17, that there is substantial day-to-day variability in the diurnal profile of $PM_{2.5}$ measured at the same location that is smoothed out after a suitably long averaging period is chosen. The large ratio of the interquartile range to the median values supports the view that there is substantial variability in the diurnal profiles.

The diurnal variability of PM components is determined by interactions between variations in emissions, the rates of photochemical transformations, and the vertical extent and intensity of turbulent mixing near the surface. Wilson and Stockburger (1990) characterized the diurnal variability of sulfate and lead in Philadelphia. At that time, Pb was emitted mainly by motor vehicles. Pollutants emitted mainly by motor vehicles, such as carbon monoxide, show two distinct peaks occurring during the morning and evening rush hours (see Chapter 3, U.S. Environmental Protection Agency, 2000b). Pollutants, such as sulfate, which are transported long distances in the free troposphere (i.e., above the planetary boundary layer), tend to be mixed



Figure 3-17. Intraday variability of hourly average $PM_{2.5}$ concentrations across the United States. Interquartile ranges, median and mean (+) values are shown. Values above the box plots refer to the number of observations during 1999.

Source: Fitz-Simons et al. (2000).

downward and have their highest concentrations during the afternoon when the intensity and vertical extent of turbulent mixing (and chemical oxidation) are greatest. Secondary aerosol components (such as secondary organic compounds) that are produced by photochemical reactions may have a daily maximum in the afternoon, similar to ozone. PM produced by residential heating (e.g., from wood burning), on the other hand, reach maximum levels during the night when inversions are near the surface.

Although the interquartile ranges for hour-to-hour changes in $PM_{2.5}$ concentrations shown in Figure 3-17 encompass several $\mu g/m^3$, extreme values for the hour-to-hour variations can be much larger (Fitz-Simons et al., 2000). The 98th percentile values for positive and negative excursions in concentration are all less than 20 $\mu g/m^3$. Maximum positive excursions were much larger, ranging from 27 $\mu g/m^3$ in the Northeast up to 220 $\mu g/m^3$ in the Southwest and with maximum excursions in other regions all less than 125 $\mu g/m^3$. It should be borne in mind that the hour-to-hour changes that are reported reflect the effects of a number of processes occurring

during passage through the sampler inlets and on the TEOM measurement elements. These factors add uncertainty to the interpretation of the hour-to-hour changes that are observed, as discussed in Chapter 2. However, because of the tendency of these monitoring instruments to lose material by evaporation, the concentrations reported during excursions probably represent lower limits to the true values that were present.

3.2.3 Relations Among Particulate Matter in Different Size Fractions

Relations Among $PM_{2.5}$, $PM_{10-2.5}$, and PM_{10}

Data obtained in 1999 by collocated $PM_{2.5}$ and PM_{10} FRM monitors have been used to calculate the ratio of $PM_{2.5}$ to PM_{10} concentrations and correlations among $PM_{2.5}$, $PM_{10-2.5}$, and PM_{10} concentrations. Results are shown in Table 3-1 for each of the seven aerosol characteristic regions identified in Chapter 6 of the 1996 PM AQCD. As can be seen from the table, the ratio of $PM_{2.5}$ to PM_{10} concentrations tends to be higher in the eastern United States than in the western United States. This general pattern and the values are consistent with that found for the studies included in Appendix 6A of 1996 PM AQCD. In that compilation based on the results of studies using dichotomous samplers, the mean ratio of $PM_{2.5}$ to PM_{10} was 0.75 in the East, 0.52 in the central United States, and 0.53 in the western United States. Although a large number of paired entries have been included in Table 3-1, seasonal variations and annual averages in a number of regions could not be determined from the data set because of data sparseness mainly during the early part of 1999. It also can be seen in Table 3-1 that the ratio of $PM_{2.5}$ to PM_{10} was greater than one for a few hundred measurements. There are a number of reasons for these results, as mentioned in Section 3.2.1 in the discussion on $PM_{10-2.5}$ concentrations.

Ultrafine Particle Concentrations

Data for characterizing the concentrations of ultrafine particles (<0.10 μm D_a) and the relations between ultrafine particles and larger particles are sparse. Although ultrafine particles dominate particle number concentrations, they make very minor contributions to $PM_{2.5}$ mass. For example, Cass et al. (2000) found that particles between 0.056 and 0.1 μm D_a contributed only 0.55 – 1.16 $\mu g/m^3$ at several sites in southern California. Perhaps the most extensive data set for ultrafine particle properties is that described by Woo et al. (2001) for a site located 10 km to the northwest of downtown Atlanta, GA. Size distributions from 3 to 2000 nm were measured

TABLE 3-1. DISTRIBUTION OF RATIOS OF $PM_{2.5}$ TO PM_{10} AND CORRELATIONS BETWEEN $PM_{2.5}$ AND PM_{10}, $PM_{2.5}$ AND $PM_{10-2.5}$, AND $PM_{10-2.5}$ AND PM_{10} FOUND AT COLLOCATED MONITORING SITES IN SEVEN AEROSOL CHARACTERISTIC (EPA/HEI) REGIONS IN 1999

				Percentiles							Correlations		
Region	**Mean**	**Sites**	**Values**	**95**	**90**	**75**	**50**	**25**	**10**	**5**	**$PM_{2.5}$:PM_{10}**	**$PM_{2.5}$:$PM_{10-2.5}$**	**$PM_{10-2.5}$:PM_{10}**
Northeast	0.70	45	1433	0.97	0.95	0.77	0.67	0.60	0.51	0.48	0.72[a]	0.02	0.71[a]
Southeast	0.70	76	2823	1.27	1.06	0.74	0.63	0.54	0.46	0.43	0.69[a]	−0.04[a]	0.69[a]
Industrial Midwest	0.70	92	4827	1.09	0.88	0.78	0.68	0.59	0.51	0.47	0.71[a]	0.17[a]	0.81[a]
Upper Midwest	0.53	39	1446	0.92	0.84	0.62	0.49	0.44	0.34	0.24	0.35[a]	−0.02	0.93[a]
Southwest	0.38	23	701	0.51	0.51	0.47	0.40	0.31	0.23	0.23	0.63[a]	0.49[a]	0.99[a]
Northwest	0.50	73	3300	0.67	0.65	0.56	0.49	0.44	0.39	0.36	0.69[a]	0.07[a]	0.77[a]
Southern California	0.47	36	1813	0.70	0.57	0.55	0.48	0.44	0.31	0.24	0.70[a]	0.19[a]	0.83[a]
		384	16,343										

[a]Results considered to be significantly different from zero at the $\alpha = 0.01$ level.

Source: U.S. EPA Aerometric Information Retrieval System.

DRAFT-DO NOT QUOTE OR CITE

every 12 minutes for 24 months beginning in August 1998. Approximately 89% of the total number of particles were found to be smaller than 100 nm; whereas 26% were found to be smaller than 10 nm. Concentrations tend to be lower during the summer than during the winter. No correlation was found between number concentration and either volume or surface area for particle sizes up to 2 μm. Because the total number of particles is concentrated in the smallest size ranges, these results also indicate that fine particle mass does not correlate with the number of ultrafine particles. The high time resolution of the measurements allows some inferences to be made about the possible sources of the ultrafine particles. The number of particles larger than 10 nm tends to peak during the morning rush hour (around 8 a.m.) and then to decrease through the day and to increase again after 6 p.m., consistent with a traffic-related source. Particles smaller than 10 nm tend to peak during the mid-afternoon, consistent with nucleation involving products of active photochemistry (McMurry et al., 2000). More direct relations between particle mass observed in different size ranges can be obtained using multi-stage impactors. Keywood et al. (1999) found a correlation between $PM_{2.5}$ and $PM_{0.15}$ of about 0.7; whereas they found correlations of about 0.96 between PM_1 and $PM_{2.5}$ and between $PM_{2.5}$ and PM_{10} based on samples collected by MOUDIs (Multiple Orifice Uniform Deposit Impactors) in six Australian cities.

3.2.4 Relations Between Mass and Chemical Component Concentrations

Time series of elemental composition data for $PM_{2.5}$ based on X-ray fluorescence (XRF) analyses have been obtained at a number of locations across the United States. Time series of components of the organic carbon fraction of the aerosol have not yet been obtained. The results of XRF analyses for the composition of the inorganic fraction of $PM_{2.5}$ and $PM_{10\text{-}2.5}$ are presented in Table 3-2 for Philadelphia, PA and in Table 3-3 for Phoenix, AZ. The frequency distribution for $PM_{2.5}$ concentration data collected at these sites were shown in Figures 3-10 and 3-11. All XRF analyses were performed at the same X-ray spectrometry facility operated by the U.S. Environmental Protection Agency in Research Triangle Park, NC. Data shown in the first column of Table 3-2 are based on analyses of filters collected over three years (April 1992 to April 1995, labeled a) at the PBY site in southwestern Philadelphia. These data and data for PM_{10} were collected using Harvard impactors. Data for $PM_{2.5}$ and $PM_{10\text{-}2.5}$ shown in the second and third columns were collected at the Castor Avenue Laboratory, operated by the City of Philadelphia from July 25 to August 14, 1994, using a modified dichomotous sampler (VAPS).

TABLE 3-2. CONCENTRATIONS (in ng/m^3) OF $PM_{2.5}$, $PM_{10-2.5}$, AND SELECTED ELEMENTS (ng/m^3) IN THE $PM_{2.5}$ AND $PM_{10-2.5}$ SIZE RANGES WITH STANDARD DEVIATIONS (SD) AND CORRELATIONS BETWEEN ELEMENTS AND $PM_{2.5}$ MASS IN PHILADELPHIA, PA*

n = 1105	Conc (ng/m^3) ± SD (unc)	r	n = 20	Conc (ng/m^3) ± SD (unc)	r	n = 20	Conc (ng/m^3) ± SD (unc)	r
$PM_{2.5}$ [1]	17 ± 0.9 (0.8) × 10^3	1	$PM_{2.5}$ [2]	29.8 ± 14.7 (1.1) × 10^3	1	$PM_{10-2.5}$ [2]	8.4 ± 2.9 (0.4) × 10^3	1
Al	4.0 ± 56 (31)	0.10	Al	109 ± 61 (21)	0.15	Al	325 ± 241 (99)	0.89
Si	116 ± 107 (21)	0.51	Si	191 ± 134 (26)	0.22	Si	933 ± 652 (231)	0.90
P	8.6 ± 14 (10)	0.31	P	15 ± 4.3 (2.7)	0.72	P	28 ± 9.4 (7.1)	0.78
S	2100 ± 1610 (143)	0.92	S	3190 ± 1920 (207)	0.91	S	38 ± 45 (71)	−0.15
Cl	5.1 ± 35 (3.4)	−0.01	Cl	23 ± 28 (5.5)	0.19	Cl	47 ± 48 (5.8)	−0.11
K	60.4 ± 45 (4.7)	0.50	K	68 ± 21 (6.4)	0.31	K	100 ± 66 (10)	0.81
Ca	47 ± 33 (4.2)	0.39	Ca	63 ± 33 (9.0)	−0.02	Ca	421 ± 192 (31)	0.81
Ti	4.9 ± 5.2 (4.1)	0.44	Ti	8.7 ± 4.7 (9.0)	0.47	It	30 ± 17 (5.6)	0.90
V	8.8 ± 8.7 (1.8)	0.37	V	9.7 ± 7.1 (2.9)	0.38	V	3.2 ± 2.2 (1.5)	0.66
Cr	0.7 ± 1.1 (0.7)	0.15	Cr	1.4 ± 1.2 (2.9)	0.09	Cr	1.0 ± 5.0 (0.9)	0.43
Mn	3.1 ± 2.2 (0.8)	0.39	Mn	3.2 ± 1.5 (1.6)	0.43	Mn	6.3 ± 4.1 (0.6)	0.90
Fe	109 ± 71 (10.5)	0.50	Fe	134 ± 49 (0.5)	0.48	Fe	352 ± 156 (24)	0.90
Co	0.1 ± 1.8 (1.4)	0.04	Co	0.8 ± 0.7 (8.5)	0.58	Co	−0.2 ± 0.5 (0.3)	−0.10
Ni	7.3 ± 8.4 (1.4)	0.22	Ni	8.5 ± 5.6 (0.3)	0.61	Ni	2.0 ± 1.4 (0.3)	0.08
Cu	4.8 ± 4.9 (1.1)	0.25	Cu	7.7 ± 3.8 (0.7)	0.22	Cu	14 ± 12 (1.1)	−0.05
Zn	36.9 ± 44 (3.7)	0.21	Zn	56 ± 37 (4.8)	0.22	Zn	52 ± 43 (4.7)	−0.03
As	0.6 ± 1.4 (1.2)	0.18	As	0.4 ± 1.0 (1.0)	−0.02	As	0 ± 0.5 (0.5)	0.07
Se	1.5 ± 1.3 (0.6)	0.63	Se	1.3 ± 0.8 (0.4)	0.65	Se	−0.1 ± 0.2 (0.2)	−0.24
Br	5.0 ± 11.7 (0.9)	0.11	Br	14 ± 12 (1.3)	0.21	Br	3.0 ± 2.5 (0.5)	−0.10
Pb	17.6 ± 22 (2.5)	0.19	Pb	28 ± 24 (2.4)	0.26	Pb	13 ± 11 (1.3)	0.10

[1]Data obtained at the Presbyterian home in Philadelphia from April 1992 to April 1995 with Harvard impactors.
[2]Data obtained at the Castor Avenue Laboratory, North Central Philadelphia from July 25 to August 14 with a modified dichotomous sampler.

*Note: Values in parentheses refer to analytical uncertainty (unc) in X-ray fluorescence determinations.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3-3. CONCENTRATIONS (in ng/m^3) OF $PM_{2.5}$, $PM_{10-2.5}$, AND SELECTED ELEMENTS IN THE $PM_{2.5}$ AND $PM_{10-2.5}$ SIZE RANGE WITH STANDARD DEVIATIONS (SD) AND CORRELATIONS (r) BETWEEN ELEMENTS AND $PM_{2.5}$ AND $PM_{10-2.5}$ MASS IN PHOENIX, AZ*

n = 164	Conc (ng/m^3) ± SD (unc)	r	n = 164	Conc (ng/m^3) ± SD (unc)	r
$PM_{2.5}$	11.2 ± 0.6 (0.6) × 10^3	1	$PM_{10-2.5}$	27.6 ± 14.8 × 10^3	1
Al	125 ± 77 (30)	0.23	Al	1879 ± 979 (547)	0.92
Si	330 ± 191 (48)	0.35	Si	535 ± 2825 (1347)	0.92
P	11 ± 7.8 (5.7)	0.52	P	37 ± 20 (17)	0.58
S	487 ± 254 (40)	0.16	S	131 ± 47 (26)	0.77
Cl	19 ± 44 (3.0)	0.13	Cl	208 ± 204 (24)	0.28
K	110 ± 63 (9.2)	0.67	K	561 ± 298 (62)	0.92
Ca	129 ± 72 (11)	0.51	Ca	1407 ± 755 (124)	0.90
It	11 ± 7.1 (2.7)	0.44	Ti	130 ± 71 (20)	0.90
V	0.7 ± 2.0 (2.2)	−0.28	V	2.0 ± 2.0 (1.5)	0.51
Cr	0.6 ± 0.9 (0.7)	0.41	Cr	2.6 ± 1.7 (0.7)	0.76
Mn	5.7 ± 4.3 (0.7)	0.64	Mn	29 ± 16 (3.0)	0.91
Fe	177 ± 113 (16)	0.80	Fe	1211 ± 674 (133)	0.90
Co	−0.4 ± 1.0 (1.0)	−0.01	Co	1.2 ± 2.2 (1.9)	0.38
Ni	0.6 ± 0.9 (0.5)	0.38	Ni	1.8 ± 1.4 (0.7)	0.70
Cu	5.2 ± 6.1 (1.5)	0.69	Cu	10.3 ± 9.0 (1.5)	0.58
Zn	17 ± 14.7 (1.8)	0.64	Zn	25 ± 16 (3.2)	0.64
As	1.9 ± 3.2 (0.6)	0.50	As	0.6 ± 0.8 (0.6)	0.41
Se	0.4 ± 0.8 (0.4)	0.40	Se	−0.02 ± 0.3 (0.3)	0.21
Br	3.8 ± 2.0 (0.6)	0.57	Br	0.8 ± 0.6 (0.4)	0.48
Pb	6.6 ± 6.6 (1.0)	0.69	Pb	4.6 ± 3.8 (1.1)	0.59

* Values in parenthesis refer to analytical uncertainty (unc) in X-ray fluorescence determinations.

Source: Zweidinger et al. (1998).

The samples at the Phoenix site were collected in 1996 and 1997 using the same type of dichotomous sampler used in the shorter term study in Philadelphia. These data are shown to give an idea of the range of concentrations found in studies conducted more recently than those

shown in Appendix 6A of the 1996 PM AQCD. The speciation network will at least provide more thorough coverage of the composition of particles in the $PM_{2.5}$ size range across the United States. Results from the pilot study for the speciation network are given in Appendix 3B.

As can be seen from inspection of Tables 3-2 and 3-3, the analytical uncertainty (given in parentheses next to concentrations) as a fraction of the absolute concentration is highly variable. It exceeds the concentration for a number of trace metals whose absolute concentrations are low; whereas it is very small for abundant elements such as sulfur.

Sulfur is the major element analyzed in the $PM_{2.5}$ size fraction in the two Philadelphia studies and is highly correlated with $PM_{2.5}$; however its abundance is roughly two orders of magnitude lower in the $PM_{10-2.5}$ size range and is negatively correlated with $PM_{10-2.5}$. Concentrations of the crustal elements: Al, Si, K, Ca, and Fe are much higher in the $PM_{10-2.5}$ size range than in the $PM_{2.5}$ size range and are well correlated with $PM_{10-2.5}$. A number of trace elements (e.g., Cr, Co, Ni, Cu, Zn, As, Se and Pb) are detectable in the two $PM_{2.5}$ data sets, and the concentrations of many of these elements are much greater than the uncertainty in their determination. Except for Co, As, and Se which are not detected in the $PM_{10-2.5}$ samples, the concentrations of many elements (Cr, Zn, and Pb) are comparable in the $PM_{2.5}$ and $PM_{10-2.5}$ size ranges. The concentration of Cu is significantly higher in the $PM_{10-2.5}$ size range, whereas the concentration of Ni is smaller in the $PM_{10-2.5}$ size range than in the $PM_{2.5}$ size range.

There are a number of distinct differences between the $PM_{2.5}$ sets for Philadelphia and Phoenix. For instance, sulfate and associated cations and water that would be expected to correspond to the measurement of S appear to constitute a major fraction of the composition of the PM in the Philadelphia data set; whereas they appear to constitute a much smaller fraction of the PM in the Phoenix data set. The highest $PM_{2.5}$ values were observed in Philadelphia during episodes driven by high sulfate abundances; whereas those in Phoenix were driven by raised soil dust. The concentration of S in Phoenix is much lower in the Phoenix $PM_{2.5}$ data set than in either Philadelphia $PM_{2.5}$ data set, and it is only weakly correlated with $PM_{2.5}$. As in Philadelphia, the concentration of S in Phoenix is higher in the $PM_{2.5}$ size range than in the $PM_{10-2.5}$ size range. Trace metals (e.g., Cr, Co, Ni, Cu, Zn, As, and Pb) are not well correlated ($0.04 < r < 0.25$) with $PM_{2.5}$ in the Philadelphia data set; whereas they are more variably correlated ($0.01 < r < 0.69$) with $PM_{2.5}$ in the Phoenix data set. The uncertainty in the concentration measurement most probably plays a role in determining a species' correlation with

$PM_{2.5}$, especially when the analytical uncertainty is high relative to concentration, as it is for a number of elements in the data shown in Tables 3-2 and 3-3. Concentrations of Al, Si, K, Ca, and Fe are again much higher in the $PM_{10-2.5}$ size range than in the $PM_{2.5}$ size range and are strongly correlated with $PM_{10-2.5}$ in both data sets.

There are also similarities in the $PM_{2.5}$ data sets for Philadelphia and Phoenix. Crustal elements are not as well correlated with $PM_{2.5}$ as they are with $PM_{10-2.5}$ in both data sets. The concentrations of trace metals (Cr, Ni, Cu, and Zn) in $PM_{2.5}$ are similar in Philadelphia and Phoenix. It can also be seen that their concentrations are of the same order of magnitude in both $PM_{2.5}$ and $PM_{10-2.5}$. Concentrations of Cu are noticeably higher in $PM_{10-2.5}$ than in $PM_{2.5}$ in both Philadelphia and Phoenix. These results are consistent with those of many monitoring studies shown in Appendix 6A of the 1996 PM AQCD, which also show that concentrations of these metals are of the same order of magnitude in both size fractions and that concentrations of Cu tend to be higher in $PM_{10-2.5}$ than in $PM_{2.5}$.

One study suggests that the partitioning of trace metals between the fine and coarse fractions varies with PM concentration. Salma et al. (2002) determined the size distribution of a number of trace elements at four sites characterizing environments ranging from the urban background to an urban traffic tunnel in Budapest, Hungary. S, K, V, Ni, Cu, Zn, As, and Pb were found mainly in the fine fraction at the urban background site; but their mass median aerodynamic diameters increased with increasing PM concentrations until they were all found mainly in the coarse fraction in the traffic tunnel. They also found that Na, Mg, Al, Si, P, Ca, It, Fe, Ga, Sr, Zr, Mo, and Ba were concentrated mainly in the coarse fraction at all four sites and that their mass median aerodynamic diameters increased with increasing PM concentrations.

The mean concentration of Pb observed in the methods evaluation study for the speciation network was only about 5 ng/m^3 in Philadelphia during the first half of 2000 (cf. Appendix 3B); whereas its concentration was about three times higher during the studies conducted during the early 1990s (Table 3-3). In a study conducted in the greater Philadelphia area during the summer of 1982, Dzubay et al. (1988) found concentrations of Pb of about 250 ng/m^3, or about fifty times higher than observed in 2000. The mean Pb concentration was about 3 ng/m^3 at the Phoenix site included as part of the same methods evaluation study for the speciation network; however, the mean Pb concentration was 39 ng/m^3 during an earlier study conducted during 1989 and 1990 in Phoenix (Chow et al., 1991). These changes in Pb concentrations are consistent with those in

many other urban areas for which monitoring studies have been conducted during the late 1970s and 1980s (cf. Appendix 6A of the 1996 PM AQCD) and for which there are data given in Appendix 3B. It should be remembered that the older studies were conducted while Pb was still used as a gasoline additive. The ratio of Pb in $PM_{2.5}$ to Pb in $PM_{10-2.5}$ was also much higher in the older studies than in the more recent ones, reflecting the importance of combustion as its source. Smaller decreases are apparent in the concentrations of other trace metals such as Cu, Ni, and Zn between studies conducted in the early 1980s and in the methods evaluation study for the speciation network conducted in 2000.

Some indication of the sources of metals such as Pb, Cu, Cd, and Zn in current, ambient $PM_{2.5}$ and $PM_{10-2.5}$ samples can be obtained by examining their sources in urban runoff. The sources of these elements in urban runoff were found to be the weathering of building surfaces, motor vehicle brake and tire wear, engine oil and lubricant leakage and combustion, and wet and dry atmospheric deposition (Davis et al., 2001). Once deposited on the ground, these elements can be resuspended with other material as $PM_{2.5}$ and $PM_{10-2.5}$, although research is needed into the mechanisms of how this is accomplished. Wind-abrasion on building siding and roofs (coatings such as Pb paint and building material such as brick, metal, and wood siding); brake wear (brake pads contain significant quantities of Cu and Zn); tire wear (Zn is used as a filler in tire production); and burning engine oil could all produce particles containing these metals, especially Zn.

Data for the chemical composition of ambient ultrafine particles are sparse. In a study conducted at several urban sites in Southern California, Cass et al. (2000) found that the composition of ultrafine particles ranged from 32 to 67% organic compounds, 3.5 to 17.5% elemental carbon, 1 to 18% sulfate, 0 to 19% nitrate, 0 to 9% ammonium, 1 to 26% metal oxides, 0 to 2% sodium, and 0 to 2% chloride. Thus carbon, in various forms, was found to be the major contributor to the mass of ultrafine particles. However, ammonium was found to contribute 33% of the mass of ultrafine particles at one site in Riverside. Iron was the most abundant metal found in the ultrafine particles. Chung et al. (2001) found that carbon was the major component of the mass of ultrafine particles in a study conducted during January of 1999 in Bakersfield, CA. However, in the study of Chung et al., the contribution of carbonaceous species (OC and EC) (typically 20 to 30%) was much lower than that found in the cities in Southern California. They found that calcium was the dominant cation, accounting for about 20% of the mass of ultrafine

particles in their samples. Sizable contributions from silicon (0 to 4%) and aluminum (6 to 14%) were also found. Further studies, including scanning electron microscopy, may be needed to quantify the role of coarse particle bounce from the upper stages of their MOUDI impactor.

Gone et al. (2000) measured the size distribution of trace elements from 0.056 μm to 1.8 μm D_a in Pasadena, CA and in the Great Smoky Mountains National Park, TN. They found that elements identified as being of anthropogenic origin had mass median diameters below 1 μm PM; whereas elements of crustal origin generally had a mass median diameter greater than 1 μm. Concentrations of trace metals were much higher in the accumulation mode than in the ultrafine mode in both study areas. In PM_1, 76% of Cr, 95% of Fe, 94% of Zn, 89% of As, and 79% of Cd at the Tennessee site were found in the accumulation mode; and 70% of Fe, 85% of Zn, 92% of As, and 84% of Cd were found in the accumulation mode in Pasadena. Fe was the most abundant metal found in the ultrafine particles. The abundance of crustal elements, such as Al, declined rapidly with decreasing particle size at both locations, and Al in PM_1 probably represented the lower tail of the coarse PM mode. However, on two days at Pasadena there were increases in the concentration of Al in ultrafine particles that were associated with increases in Sc and Sm. The latter two elements originate exclusively from crustal material (Gone et al., 2000).

3.2.5 Spatial Variability in Particulate Matter and its Components

$PM_{2.5}$

Aspects of the spatial variability of $PM_{2.5}$ concentrations on the urban scale are examined in this section. Intersite correlation coefficients for $PM_{2.5}$ can be calculated based on the results of FRM monitors placed at multiple sites within Metropolitan Statistical Areas (MSAs) across the United States. Pearson correlation coefficients (r) calculated for pairs of monitoring sites in the Columbia, SC; Detroit, MI; Chicago, IL; and Los Angeles, CA MSAs are shown in Table 3-4. The 90th percentile value, P_{90}, of the absolute differences (in $\mu g/m^3$) between the two sites is shown below r along with the coefficient of divergence (COD) in parentheses, and the number of observations used in the calculation of r, P_{90} and COD is given on the third line. The COD was used by Wongphatarakul et al. (1998) as a measure of the degree of similarity between two

TABLE 3-4. MEASURES OF THE SPATIAL VARIABILITY OF $PM_{2.5}$ CONCENTRATIONS WITHIN SELECTED METROPOLITAN STATISTICAL AREAS

(a) Columbia, SC

Site I.D. #	45-063-0005	45-063-0008	45-079-0007	45-079-0019
45-063-0005	1	0.882 (5.3, 0.121) 215	0.949 (3.9, 0.081) 204	0.93 (4.8, 0.099) 216
45-063-0008		1	0.933 (4.0, 0.082) 202	0.949 (3.3, 0.067) 216
45-079-0007				0.971 (2.7, 0.06) 203
45-079-0019				1
Mean	14.680	16.462	15.461	16.098
Obs	231	228	216	229
SD	6.760	7.121	6.900	7.148

Key
AIRS Site I.D.#
Pearson r
(90th %-ile difference in concentration, coefficient of divergence)
number of observations

(b) Detroit, MI

Site I.D. #	26-099-0009	26-125-0001	26-147-0005	26-163-0033	26-163-0036
26-099-0009	1	0.958 (4.9, 0.107) 83	0.952 (5.6, 0.127) 96	0.931 (12.7, 0.222) 98	0.926 (9.0, 0.177) 96
26-125-0001		1	0.939 (5.8, 0.121) 73	0.92 (12.3, 0.193) 77	0.917 (8.3, 0.151) 75
26-147-0005			1	0.876 (13.3, 0.222) 89	0.875 (8.9, 0.197) 88
26-163-0033				1	0.923 (7.1, 0.108) 89
26-163-0036					1
Mean	13.450	15.552	14.172	20.173	17.446
Obs	113	90	102	108	103
SD	7.922	9.223	8.771	10.475	9.626

Key
AIRS Site I.D.#
Pearson r
(90th %-ile difference in concentration, coefficient of divergence)
number of observations

TABLE 3-4 (cont'd). MEASURES OF THE SPATIAL VARIABILITY OF $PM_{2.5}$ CONCENTRATIONS WITHIN SELECTED METROPOLITAN STATISTICAL AREAS

(c) Chicago, IL

Site I.D. #	17-031-0014	17-031-0022	17-031-0050	17-031-0052	17-031-1016	17-031-2001	17-031-3301	17-031-4006	17-031-4201	17-043-4002	17-197-1002
17-031-0014	1	0.912 (4.4, 0.121) 96	0.946 (4.6, 0.077) 78	0.909 (6.6, 0.13) 100	0.921 (7.5, 0.143) 92	0.902 (5.6, 0.111) 98	0.927 (5.1, 0.104) 98	0.876 (5.8, 0.133) 88	0.936 (5.3, 0.139) 95	0.885 (5.7, 0.13) 95	0.774 (7.4, 0.158) 81
17-031-0022		1	0.92 (5.4, 0.113) 87	0.872 (6.5, 0.14) 108	0.866 (7.0, 0.141) 103	0.892 (5.7, 0.131) 104	0.879 (6.0, 0.132) 106	0.689 (7.9, 0.213) 92	0.86 (7.9, 0.197) 101	0.855 (7.2, 0.165) 100	0.79 (7.1, 0.17) 87
17-031-0050			1	0.941 (5.0, 0.094) 259	0.93 (7.8, 0.12) 83	0.955 (3.5, 0.082) 89	0.923 (5.3, 0.096) 91	0.75 (7.9, 0.176) 75	0.928 (6.2, 0.162) 247	0.922 (5.3, 0.117) 91	0.867 (7.6, 0.131) 87
17-031-0052				1	0.887 (7.9, 0.133) 105	0.885 (7.3, 0.125) 109	0.881 (7.0, 0.128) 110	0.797 (8.5, 0.177) 98	0.879 (9.6, 0.179) 310	0.836 (8.5, 0.154) 112	0.721 (10.2, 0.169) 108
17-031-1016					1	0.932 (7.3, 0.108) 99	0.898 (7.5, 0.124) 102	0.787 (10.0, 0.205) 92	0.915 (9.8, 0.2) 98	0.902 (9.5, 0.154) 95	0.84 (10.5, 0.173) 85
17-031-2001						1	0.931 (4.5, 0.084) 110	0.861 (5.9, 0.153) 93	0.943 (5.5, 0.14) 101	0.949 (4.3, 0.1) 99	0.893 (5.1, 0.118) 89
17-031-3301							1	0.823 (7.0, 0.158) 95	0.915 (6.4, 0.152) 103	0.953 (4.4, 0.092) 101	0.873 (5.8, 0.128) 91
17-031-4006								1	0.818 (7.3, 0.146) 92	0.865 (5.1, 0.124) 88	0.752 (7.6, 0.161) 78
17-031-4201									1	0.922 (4.8, 0.123) 106	0.809 (7.1, 0.157) 99
17-043-4002										1	0.921 (4.2, 0.099) 90
17-197-1002											1
Mean	15.823	17.933	16.996	18.295	20.277	16.790	16.889	15.268	14.283	15.215	15.994
Obs	104	113	274	346	108	113	115	101	327	116	112
SD	7.935	8.175	8.468	9.289	9.331	7.694	7.689	8.423	7.905	7.568	7.405

Key
AIRS Site I.D.#
Pearson r
(90th %-ile difference in concentration, coefficient of divergence)
number of observations

DRAFT-DO NOT QUOTE OR CITE

TABLE 3-4 (cont'd). MEASURES OF THE SPATIAL VARIABILITY OF $PM_{2.5}$ CONCENTRATIONS WITHIN SELECTED METROPOLITAN STATISTICAL AREAS

(d) Los Angeles, CA					
Site I.D. #	06-037-0002	06-037-1103	06-037-1601	06-037-4002	06-037-9002
06-037-0002	1	0.828	0.763	0.573	0.276
		(12.8, 0.192)	(17.3, 0.211)	(20.2, 0.263)	(28.0, 0.392)
		391	196	379	186
06-037-1103		1	0.88	0.752	0.328
			(11.8, 0.140)	(14.6, 0.191)	(26.4, 0.375)
			173	353	164
06-037-1601			1	0.859	0.363
				11.8, 0.174	31.0, 0.411
				171	181
06-037-4002				1	0.338
					(24.4, 0.356)
					157
06-037-9002					1
Mean	21.682	22.207	24.764	20.225	10.917
Obs	469	428	218	417	204
SD	13.923	13.840	14.056	12.994	5.043

Key
AIRS Site I.D.#
Pearson r
(90th %-ile difference in concentration, coefficient of divergence)
number of observations

Source: Pinto et al., (2002). Data from U.S. EPA Aerometric Information Retrieval System (AIRS).

aerosol data sets[1]. The annual mean concentrations, the number of observations used to calculate the annual average, and the standard deviation are shown directly beneath the correlation tables for each site. These analyses along with those for another 23 MSAs are given along with maps in

[1]The COD for this purpose is defined as follows:

$$COD_{jk} = \sqrt{\frac{1}{p}\sum_{i=1}^{p}\left(\frac{x_{ij} - x_{ik}}{x_{ij} + x_{ik}}\right)^2} \qquad (3\text{-}1)$$

where x_{ij} and x_{ik} represent the 24-h average $PM_{2.5}$ concentration for day i at site j and site k and p is the number of observations.

Appendix 3A. The four MSAs shown in Table 3-4 were chosen to illustrate different patterns of spatial variability across the United States. In addition, air pollution-health outcome studies have been performed in a few of these MSAs. It can be seen from inspection of Table 3-4 that correlation coefficients vary over a wide range in the MSAs shown. Correlations between sites in the Columbia, SC MSA and the Detroit, MI MSA are all high and span a relatively narrow range (0.88 to 0.97). Correlations between sites in the Chicago, IL MSA span a wider range (0.69 to 0.96). However, the correlations between sites in the Los Angeles-Long Beach MSA are much lower than in the three other MSAs and span an even wider range of values (0.28 to 0.88). The extension of these analyses to include the relevant CMSAs (consolidated MSA) would also produce a number of sites that are even less well correlated with each other in part because of the larger distances involved. Correlation coefficients between pairs of sites in the other 23 MSAs given in Appendix 3A fall within the range of values given in Table 3-4. Some indication of the performance of collocated monitors is given by inspection of the last two columns of Table 3A-10. These data were obtained by two collocated $PM_{2.5}$ monitors in the Steubenville, OH-Weirton, WV MSA. Values of r, P_{90}, and COD for these two monitors are 0.978, 2.5 $\mu g/m^3$, and 0.101.

There may be a regional pattern evident in the data given in Appendix 3A, data for which correlations tend to be higher between monitoring sites in MSAs in the eastern and central United States than between monitoring sites in the western United States. In a few MSAs (Milwaukee, WI; Norfolk, VA; Grand Rapids, MI; and Baton Rouge, LA), intersite correlations are all greater than 0.9. In several others (Philadelphia, PA; Columbia, SC; Steubenville, OH; Detroit, MI; Kansas City, KS-MO; and Dallas, TX), they are all greater than 0.8. Intersite correlations tend to be lower and to span a broader range in several cities such as Atlanta, GA; Seattle, WA; and Los Angeles, CA, in part due to the location of monitoring sites outside of the main urban area and in a different air shed. In many MSAs, there is a wide range in the intersite correlations that are found. For example, in the Seattle, WA MSA (Table 3A-23), values r of range from 0.41 to 0.95. Correlations between sites in the Atlanta, GA, Birmingham, AL, and Tampa, FL MSAs tend to be lower and span a broader range than do those for the other southern cities examined (Columbia, SC; Norfolk, VA; Baton Rouge, LA; and Dallas, TX). Likewise, correlations between a number of sites in western MSAs are higher than those in some eastern

MSAs. For example, correlations between monitors in the Pittsburgh, PA MSA tend to be lower than those in the Salt Lake City, UT MSA.

There are a number of factors that affect intersite correlations within MSAs. These include field measurement and laboratory analysis errors, placement of monitors close to active sources, placement of monitors in outlying areas, placement of monitors in locations that are isolated topographically from other monitors, placement of monitors in areas outside of local atmospheric circulation regimes (e.g., land-sea breezes), and transient local events (thunderstorms, sporadic emissions).

It should not be automatically assumed that distance between sites in urban areas is solely responsible for the spatial variability that is observed. In several areas such as Atlanta, GA; Seattle, WA; and Los Angeles-Long Beach, CA, there is at least one site that is remote from the others (by at least 100 km) and is physically separated from them by mountains and is really not part of the urban area. Correlations between concentrations at these sites and others tend to be lower than among the other sites, and concentration differences tend to be larger. However, in many MSAs, especially in the East, correlations are higher, and differences in concentrations are lower for sites that are located farthest apart. This situation arises because these sites are influenced more by the regional background of secondary PM rather than by local sources. Nor is there any set distance below which correlations and differences in concentrations tend towards some limiting values. In Gary, IN, for example, intersite correlations are lowest, and concentration differences are highest for the closest site pair.

Indications of land use (commercial, industrial, residential, agricultural, forest) and location of sites (urban/city center, suburban, rural) are given in the AIRS data base. Categories such as urban/city center can refer to very different conditions in Columbia, SC and Chicago, IL. Also, it should not be automatically assumed that concentrations measured at sites categorized as industrial are dominated by local emissions. The $PM_{2.5}$ monitoring sites are generally deployed to capture potential population exposures in a variety of environments as opposed to monitoring for compliance as it exists around local sources. It should be remembered that much of $PM_{2.5}$ is secondary in origin. The widespread formation of secondary PM coupled with the long lifetime of $PM_{2.5}$ ensures some measure of uniformity in the correlations of $PM_{2.5}$ across urban areas. Correlations between many site pairs classified as industrial can be high even though they are separated by large distances, as in the Seattle MSA.

Some indication of the variability of primary $PM_{2.5}$ produced by local sources can be obtained by examining the variability of carbon monoxide (CO), which is produced mainly by mobile sources (U.S. Environmental Protection Agency, 2000b) and by the variability in elemental carbon (EC) concentrations (Kinney et al., 2000). CO is relatively inert on the urban scale, and its distribution is governed by the spatial pattern of its emissions and the subsequent dispersion of these emissions and not by photochemistry. Carbon monoxide concentrations are at least a factor of three higher near urban centers than in surrounding rural areas within the four consolidated metropolitan statistical areas examined in the EPA document, Air Quality Criteria for Carbon Monoxide (CO AQCD) (U.S. Environmental Protection Agency, 2000b). Correlations of CO within the urban areas examined in that document were all low to moderate. Therefore, it might be expected that primary $PM_{2.5}$ produced by local traffic should be at least as heterogeneous as CO in a given urban area. EC is a significant component of diesel exhaust (cf. Appendix 3D). Kinney et al. (2000) measured EC and $PM_{2.5}$ concentrations at four sites located on sidewalks of streets characterized by varying exposures to diesel emissions in upper Manhattan (Harlem, NY). Whereas the mean $PM_{2.5}$ concentrations varied by about one-third from 37 to 47 $\mu g/m^3$ at the four sites, mean EC concentrations varied by a factor of four from 1.5 to 6.2 $\mu g/m^3$. The corresponding ratios of EC to $PM_{2.5}$ ranged from 0.039 to 0.14. Although EC constituted a relatively small fraction of $PM_{2.5}$ in this study, spatial variability in its sources (diesel and gasoline fueled vehicles, resuspended road dust, and cooking) contributed, on average, about one-third of the spatial variability observed in $PM_{2.5}$ concentrations. Further analyses are needed to determine whether the remaining variability could be attributed to other local and city-wide sources. Because the effects of emissions from local point sources on receptor sites depend strongly on wind direction, correlations involving contributions from sources can be much lower than from area sources (much as motor vehicle traffic) or from regionally dispersed sources (such as the photochemical production of secondary organic PM and sulfate).

The difference in mean $PM_{2.5}$ concentrations between the site with the lowest and the site with the highest mean concentration range in all MSAs included in Appendix 3A ranges from less than 1 $\mu g/m^3$ to about 7 $\mu g/m^3$, except for the Los Angeles MSA which shows larger differences. In the Los Angeles MSA, there is one monitoring site (Figure 3A-25a) that is separated from the remaining sites by the San Gabriel Mountains and has much lower mean

$PM_{2.5}$ concentrations, much smaller seasonal variability in concentrations, and much lower maximum concentrations than these other sites. However, the annual mean concentrations at all the other sites within the Los Angeles MSA are within 5 $\mu g/m^3$ of each other. Differences in annual mean concentrations are also larger between sites located in different MSAs but within the same CMSA. For example, in the consolidated MSA of Los Angeles-Riverside the range of annual mean $PM_{2.5}$ concentrations is extended from about 20 $\mu g/m^3$ in the urban area of Los Angeles county to about 29 $\mu g/m^3$ in Riverside County. Large differences in annual mean concentrations within a given area reflect differences in source or meteorological or unique topographic characteristics affecting sites; whereas very small differences found in some areas may only be the result of measurement imprecision.

Whereas high correlations of $PM_{2.5}$ provide an indication of the spatial uniformity in temporal variability (directions of changes) in $PM_{2.5}$ concentrations across urban areas, they do not imply uniformity in the $PM_{2.5}$ concentrations themselves. The 90th percentile difference in concentrations (P_{90}) and the coefficient of divergence are used here to give a more quantitative indication of the degree of spatial uniformity in $PM_{2.5}$ concentrations across urban areas. A COD of zero implies that both data sets are identical, and a COD of one indicates that two data sets are completely different. The calculation of the Pearson correlation coefficient, P_{90}, and COD allows for distinctions between pairs of sites to be made based on various combinations of these parameters. Figure 3-18 shows examples of the varying degree of heterogeneity in concentrations between pairs of sites that are highly correlated ($r > 0.9$ for all three site pairs). The increase in the spread of concentrations between the chosen site-pairs is reflected in increases in both P_{90} and COD. Pairs of sites showing low correlations, values of $P_{90} > 10$ $\mu g/m^3$, and CODs > 0.2, as in Los Angeles, CA (Table 3-5), indicate heterogeneity in both $PM_{2.5}$ concentrations and in their temporal variations. Note that the extended urban area or the CMSA includes Riverside County, as well as Los Angeles County. Even lower correlations and a greater degree of heterogeneity in $PM_{2.5}$ concentrations were found in the extended CMSA. Pairs of sites showing high correlations and CODs < 0.1 and P_{90}'s $\lesssim 5$ $\mu g/m^3$ (as in Columbia, SC) indicate homogeneity in both $PM_{2.5}$ concentrations and in their temporal variations. Presumably, sites such as these are more strongly affected by regional than to local sources. Pairs of sites showing high correlations ($r > 0.9$) and CODs > 0.2 and P_{90}'s $\gtrsim 10$ $\mu g/m^3$ (as in Detroit, MI) indicate heterogeneity in concentrations but homogeneity in their day to day changes.



Figure 3-18. Occurrence of differences between pairs of sites in three MSAs. The absolute differences in daily average $PM_{2.5}$ concentrations between sites are shown on the x-axis and the number of occurrences on the y-axis. The MSA, years of observations, AIRS site I.D. numbers for the site pairs, Pearson correlation coefficients (r), coefficients of divergence (COD), 90th percentile (P_{90}) difference in concentration between concurrent measurements are also shown.

Source: Pinto et al. (2002)

TABLE 3-5. MEASURES OF THE SPATIAL VARIABILITY OF $PM_{10-2.5}$ CONCENTRATIONS WITHIN SELECTED METROPOLITAN STATISTICAL AREAS

(a) Detroit, MI

Name	26-163-0001	26-163-0015		26-163-0025
26-163-0001	1	0.576		0.542
		53		50
26-163-0015		1		0.393
				51
26-163-0025				1
Mean	11.517	19.416		7.328
Obs	56	58		55
SD	10.262	15.611		7.638

(b) Chicago, IL

Name	17-031-1016	17-031-2001	17-031-3301	17-197-1002
17-031-1016	1	0.69	0.544	0.583
		49	51	43
17-031-2001		1	0.865	0.823
			54	44
17-031-3301			1	0.777
				46
17-197-1002				1
Mean	16.259	14.475	17.812	6.894
Obs	93	56	58	49
SD	18.972	12.137	13.641	10.217

(c) Los Angeles

Name	06-037-1002	06-037-1103	06-037-4002	06-037-9002
06-037-1002	1	0.79	0.83	0.59
		51	49	43
06-037-1103		1	0.79	.042
			53	46
06-037-4002			1	0.39
				47
06-037-9002				1
Mean	19.1	20.3	19.3	15.6
Obs	52	55	56	52
SD	10.58	8.4	9.2	12.9

Conversely, in the Tampa, FL MSA pairs of sites are only moderately correlated ($0.6 < r < 0.7$), but the distribution of concentrations is rather homogeneous (COD < 0.1) (cf. Appendix 3A). Thus, a number of different combinations of spatial uniformity in $PM_{2.5}$ concentrations and correlations of these concentrations are found.

Values of P_{90} for absolute differences in concentrations between sites span a wide range in the data set given in Appendix 3A. In many instances they can be quite low, only about a few $\mu g/m^3$; these cases are found mainly in the eastern United States. Values of P_{90} can be greater than 40 $\mu g/m^3$; these cases are found mainly in the western United States. Maximum differences in concentrations between sites can be much larger than shown in Figure 3-18 and have been larger than 100 $\mu g/m^3$ on several occasions in the Atlanta, GA and Los Angeles-Long Beach, CA MSAs. Rizzo and Pinto (2001) and Fitz-Simons et al. (2000) examined correlations between sites located even farther apart than those examined here based on the 1999 AIRS data set for $PM_{2.5}$. They found that in a number of MSAs, $PM_{2.5}$ concentrations are still well correlated ($r > 0.7$) to distances of 100 km or more. Leaderer et al. (1999) found $r = 0.49$ between sites outside of homes and a regional background monitor located from 1 to 175 km away in southwestern Virginia. $PM_{2.5}$ tends to be correlated over much larger areas in the East than in the West, mainly because the terrain tends to be flatter over wider areas in the East (Rizzo and Pinto, 2001).

There is also evidence for inter-annual variability in the spatial variability in $PM_{2.5}$ concentrations. The median year-to-year changes in inter-site r (0.03), P_{90} (-0.75 $\mu g/m^3$), and COD (-0.015) from 1999 to 2000 do not differ significantly from zero for all the site pairs considered in Appendix 3A. The year-to-year changes in the spatial variability of $PM_{2.5}$ concentrations in a number of MSAs such as the Columbia, SC; Grand Rapids, MI; Milwaukee, WI; Baton Rouge, LA; Kansas City, KS-MO; Boise, ID; and Portland, OR MSAs are similar and are smaller than those found in the Cleveland, OH; Salt Lake City, UT; and San Diego, CA MSAs. The ranges in these parameters are largest for a number of individual site-pairs, especially those involving sites that are remote from the others in their MSAs. In these MSAs (such as the Atlanta, GA; Los Angeles, CA; and Seattle, WA MSAs) there are sites that may be located in different air sheds from the remaining sites. Year-to-year changes in parameters describing spatial variability in $PM_{2.5}$ concentrations tend to be larger when sites in different counties within a given MSA are considered rather than when sites in the same county are

considered. There are a number of factors that can account for inter-annual variability in these parameters such as changes in patterns in the emissions of primary $PM_{2.5}$, in the transport and rates of transformation of secondary $PM_{2.5}$ precursors in field measurement and analysis procedures.

Some additional data for indicating the stability with respect to year to year changes in spatial variability are available from earlier studies. For example, a comparison of data obtained during the summers of 1992 and 1993 (Wilson and Suh, 1997) as shown in Figure 3-19 and data obtained during the summer of 1994 (Pinto et al., 1995) (cf. Table 3-8) in Philadelphia, PA suggests that inter-site correlations of $PM_{2.5}$ have remained high and that they changed very little between the two study periods.

$PM_{10-2.5}$

Intersite correlations of $PM_{10-2.5}$ concentrations obtained during the summers of 1992 and 1993 in Philadelphia, PA (Wilson and Suh, 1997) are shown in Figure 3-19. As can be seen, correlations of $PM_{10-2.5}$ are substantially lower than those for $PM_{2.5}$.

Intersite correlation coefficients can also be calculated for $PM_{10-2.5}$ based on the AIRS data set as shown in Table 3-5 for the Detroit, MI; Chicago, IL; and Los Angeles, CA MSAs. However, data for analyzing the spatial variability of $PM_{10-2.5}$ are more limited than for $PM_{2.5}$; therefore, fewer urban areas could be characterized in Appendix 3A (Figures 3A-28 to 3A-33). Whereas $PM_{2.5}$ concentrations were found to be highly correlated between sites in the Detroit, MI MSA (Table 3-4), estimated $PM_{10-2.5}$ concentrations are noticeably less well correlated. Likewise, correlations of $PM_{10-2.5}$ in the Chicago, IL MSA are also lower than those for $PM_{2.5}$. However correlations of $PM_{10-2.5}$ concentrations between several pairs of sites in the Los Angeles-Long Beach partial MSA are higher than those for $PM_{2.5}$.

The interpretation of these results is not straightforward, as concentrations of $PM_{10-2.5}$ are generated by taking the difference between collocated $PM_{2.5}$ and PM_{10} monitors. Consequently, caution must be exercised when viewing them. Errors in the measurement of $PM_{2.5}$ and PM_{10} may play a large role in reducing apparent correlations of $PM_{10-2.5}$ such that collocated $PM_{10-2.5}$ "measurements" may be expected to be poorly correlated (White, 1998). Indeed, several estimates are negative. The possible causes of these errors are essentially the same as those discussed in Section 3.2.1 with regard to the occurrence of $PM_{2.5}$ to PM_{10} ratios greater than one.



Figure 3-19. Intersite correlation coefficients for $PM_{2.5}$, PM_{10}, and $PM_{10-2.5}$.

Source: Wilson and Suh (1997).

There are also physical bases for expecting that $PM_{10-2.5}$ concentrations may be more variable than those for $PM_{2.5}$. $PM_{10-2.5}$ is mainly primary in origin, and its emissions are spatially and temporally heterogenous. Similar considerations apply to primary $PM_{2.5}$, but much of $PM_{2.5}$ is secondary, and sources of secondary PM are much less spatially and temporally variable. Dry deposition rates of particles depend strongly on particle size. Whereas all particles may be brought to the surface by turbulent motions in the atmosphere; gravitational settling becomes more important with increasing particle size. Gravitational settling can effectively limit the horizontal distance a particle can travel. For example, 10 μm D_a particles suspended in a

hypothetical 1 km deep planetary boundary layer can be removed within a few hours, but 1 μm D_a particles can remain suspended in the atmosphere for up to 100 to 1,000 times longer before being dry deposited. (Estimated atmospheric lifetimes were based on deposition velocities given in Lin et al. [1994] for typical wind speeds.) The findings of larger correlations of $PM_{10-2.5}$ between several site pairs in the Los Angeles basin and one other site pair in the St. Louis, MO-IL MSA (cf. Figure 3A-17 and 3A-30) are anomalous in light of the discussion above. However, these findings could have resulted from differences between the spatial and temporal behavior of sources of $PM_{2.5}$ and $PM_{10-2.5}$ in these locations. Because of negative values, CODs were not calculated.

PM Components

Three methods for comparing the chemical composition of aerosol databases obtained at different locations and times were discussed by Wongphatarakul et al. (1998). Log-log plots of chemical concentrations obtained at pairs of sampling sites accompanied by the coefficient of divergence (COD) were examined as a way to provide an easily visualized means of comparing two data sets[2]. Examples comparing downtown Los Angeles with Burbank and with Riverside-Rubidoux are shown in Figures 3-20 and 3-21, respectively. As the composition of two sampling sites become more similar, the COD approaches zero; as their compositions diverge, the COD approaches one. Correlation coefficients calculated between components can be used to show the degree of similarity between pairs of sampling sites.

In addition to calculating correlation coefficients for total mass or for individual components, correlation coefficients for characterizing the spatial variation of the contributions from given source types can also be calculated by averaging the correlation coefficients of the set of chemical components that represent the source type. Correlation coefficients showing the spatial relations among $PM_{2.5}$ (total) and contributions from different source categories obtained

[2]The COD for two sampling sites is defined as follows:

$$COD_{jk} = \sqrt{\frac{1}{p}\sum_{i=1}^{p}\left(\frac{x_{ij} - x_{ik}}{x_{ij} + x_{ik}}\right)^2}, \qquad (3\text{-}2)$$

where x_{ij} represents the average concentration for a chemical component i at site j, j and k represent two sampling sites, and p is the number of chemical components.



Figure 3-20. $PM_{2.5}$ chemical components in downtown Los Angeles and Burbank (1986) have similar characteristics. The spread in the data is shown by the bars.

Source: Wongphatarakul et al. (1998).



Figure 3-21. Concentrations of $PM_{2.5}$ chemical components in Rubidoux and downtown Los Angeles (1986). The diagram shows a significant spread in the concentrations for the two sites compared with downtown Los Angeles and Burbank (Figure 3-20).

Source: Wongphatarakul et al. (1998).

at various sites in the South Coast Air Basin (SoCAB) Study are shown in Table 3-6. In Wongphatarakul et al. (1998), crustal material (crustal), motor vehicle exhaust (mv), residual oil emissions (residual oil), and secondary PM (sec) were considered as source categories. Al, Si, Fe, and Ca were used as markers for crustal material (crustal). V and Ni were used as markers for fuel oil combustion (residual oil). Pb, Br, and Mn were used as markers for motor vehicle exhaust (mv), based on the lack of other, perhaps more suitable, tracers. NO_3^-, NH_4^+, and SO_4^{-2} represent secondary PM components (sec). The average of the correlation coefficients of marker elements within each source category are shown in Table 3-6. Values of r_{sec} and r_{mv} are much higher than those for $r_{crustal}$ and $r_{residual\ oil}$ throughout the SoCAB, suggesting a more uniform distribution of the contributions from secondary PM formation and automobiles than from crustal material and localized stationary sources.

TABLE 3-6. CORRELATION COEFFICIENTS FOR SPATIAL VARIATION OF $PM_{2.5}$ MASS AND DIFFERENT SOURCES FOR PAIRS OF SAMPLING SITES IN THE SOUTH COAST AIR BASIN (1986)

	r_{total}	$r_{crustal}$	r_{sec}	r_{mv}	$r_{residual\ oil}$
Hawthorne and Rubidoux	−0.027				
Long Beach and Rubidoux	0.051				
Anaheim and Rubidoux	0.066				
Downtown Los Angeles and Rubidoux	0.095				
Burbank and Rubidoux	0.120				
Hawthorne and Anaheim	0.760	0.034	0.768	0.492	0.170
Long Beach and Anaheim	0.852	0.075	0.888	0.504	0.150
Burbank and Anaheim	0.770	0.105	0.749	0.579	0.161
Downtown Los Angeles and Anaheim	0.827	0.143	0.804	0.556	0.233
Downtown Los Angeles and Hawthorne	0.808	0.568	0.854	0.669	0.533
Burbank and Hawthorne	0.704	0.599	0.790	0.688	0.491
Long Beach and Burbank	0.731	0.633	0.737	0.714	0.295
Long Beach and Hawthorne	0.880	0.649	0.909	0.861	0.482
Downtown Long Angeles and Long Beach	0.842	0.653	0.817	0.719	0.378
Downtown Los Angeles and Burbank	0.928	0.825	0.960	0.871	0.606

Source: Wongphatarakul et al. (1998).

Correlation coefficients in Philadelphia air for $PM_{2.5}$ (total), crustal components (Al, Si, Ca, and Fe), the major secondary component (sulfate), organic carbon (OC), and elemental carbon (EC) are shown in Table 3-7, based on data obtained at four sites. Because these data were obtained after Pb had been phased out of gasoline, a motor vehicle contribution could not be estimated from the data. Pb also is emitted by discrete point sources, such as the Franklin smelter. Concentrations of V and Ni were often beneath detection limits; so, the spatial variability in PM due to residual oil combustion were not estimated. Sulfate in aerosol samples collected in Philadelphia arises mainly from long-range transport from regionally dispersed sources (Dzubay et al., 1988). This conclusion is strengthened by the high correlations in sulfate between different monitoring sites and the uniformity in sulfate concentrations observed among the sites. Widespread area sources (e.g., motor vehicle traffic) also may emit pollutants that are correlated between sites provided that traffic patterns and emissions are similar throughout the area under consideration.

TABLE 3-7. CORRELATION COEFFICIENTS FOR SPATIAL VARIATION OF $PM_{2.5}$ MASS AND DIFFERENT COMPONENTS FOR PAIRS OF SAMPLING SITES IN PHILADELPHIA (1994)

	r_{tot}	$r_{crustal}$	r_{sec}	r_{OC}	r_{EC}	r_{Pb}
Castor Ave. and Roxboro	0.92	0.52	0.98	0.88	0.84	0.43
Castor Ave. and NE Airport	0.93	0.47	0.99	0.88	0.77	−0.07
Castor Ave. and Broad St.	0.93	0.57	0.99	0.85	0.89	0.11
Roxboro and NE Airport	0.98	0.67	0.98	0.83	0.82	0.20
Roxboro and Broad St.	0.95	0.90	0.98	0.86	0.79	0.47
NE Airport and Broad St.	0.95	0.69	0.99	0.84	0.63	0.11

Source: Pinto et al. (1995).

Landis et al. (2001) found relatively high correlations between $PM_{2.5}$ (r = 0.97), sulfate (r = 0.99), OC (r = 0.97), EC (r = 0.83), NaCl (r = 0.83), and nitrate (r = 0.83) measured at two sites located several km apart in the Baltimore, MD area. Concentrations of crustal material

(r = 0.63) and the sum of total metal oxides (r = 0.76) were not as well correlated. These results are consistent with those for another eastern city, Philadelphia, PA, given in Table 3-7. The results presented above for Philadelphia, PA; Baltimore, MD; and Los Angeles, CA, indicate that secondary PM components are more highly correlated than primary components and may be more highly correlated than total $PM_{2.5}$. These results suggest that the correlation of PM concentrations across an urban area may depend on the relative proportions of primary and secondary components of PM at individual sites. Sampling artifacts affecting the measurement of nitrate and organic carbon can obscure these relations and may depress correlations between sites.

Kao and Friedlander (1995) examined the statistical properties of a number of PM components in the South Coast Air Basin (Los Angeles area). They found that, regardless of source type and location within their study area, the concentrations of nonreactive, primary components of PM_{10} had approximately log-normal frequency distributions with constant values of the geometric standard deviations (GSDs). However, aerosol constituents of secondary origin (e.g., SO_4^{-2}, NH_4^+, and NO_3^-) were found to have much higher GSDs. Surprisingly, the GSDs of organic (1.87) and elemental (1.74) carbon were both found to be within 1 SD (0.14) of the mean GSD (1.85) for nonreactive primary species, compared to GSD's of 2.1 for sulfate, 3.5 for nitrate, and 2.6 for ammonium. These results suggest that most of the organic carbon seen in ambient samples in the South Coast Air Basin was of primary origin. Pinto et al. (1995) found similar results for data obtained during the summer of 1994 in Philadelphia. Further studies are needed to determine if these relations are valid at other locations and to what extent the results might be influenced by sampling artifacts such as the evaporation of volatile constituents during or after sampling.

Very few studies have compared aerosol composition in urban areas to that in nearby rural areas. One exception is Tanner and Parkhurst (2000), which indicates that sulfate constituted a larger fraction of fine particle mass at rural sites in the Tennessee Valley $PM_{2.5}$ monitoring network than did organic carbon. For urban sites, the situation was largely reversed, with organic carbon constituting a larger fraction of aerosol mass than sulfate. Systematic comparisons of urban-rural differences in aerosol properties will be facilitated in the future with the implementation of the national speciation network and the continued operation of the IMPROVE network.

3.3 SOURCES OF PRIMARY AND SECONDARY PARTICULATE MATTER

Information about the nature and relative importance of sources of ambient PM is presented in this section. Table 3-8 summarizes anthropogenic and natural sources for the major primary and secondary aerosol constituents of fine and coarse particles. Major sources of each constituent are shown in boldface type. Anthropogenic sources can be further divided into stationary and mobile sources. Stationary sources include fuel combustion for electrical utilities, residential space heating, and industrial processes; construction and demolition; metals, minerals, and petrochemicals; wood products processing; mills and elevators used in agriculture; erosion from tilled lands; waste disposal and recycling; and fugitive dust from paved and unpaved roads. Mobile or transportation-related sources include direct emissions of primary PM and secondary PM precursors from highway and off-highway vehicles and non-road sources. In addition to fossil fuel combustion, biomass in the form of wood is burned for fuel. Vegetation is burned to clear new land for agriculture and for building construction, to dispose of agricultural and domestic waste, to control the growth of animal or plant pests, and to manage forest resources (prescribed burning). Also shown are sources for precursor gases whose oxidation forms secondary particulate matter. A description of the atmospheric chemical processes producing secondary PM is given in Section 3.3.1.

In general, the sources of fine particulate matter are very different from those for coarse PM. Some of the mass in the fine size fraction has been formed during combustion from material that has volatilized in combustion chambers and then recondensed before emission into the atmosphere. By and large, however, most ambient $PM_{2.5}$ has been formed in the atmosphere from photochemical reactions involving precursor gases. PM formed by the first mechanism is referred to as primary, and PM formed by the second mechanism is referred to as secondary. $PM_{10\text{-}2.5}$ is mainly primary in origin as it is produced by the abrasion of surfaces or by the suspension of biological material. Because precursor gases undergo mixing during transport from their sources, it is difficult to identify individual sources of secondary constituents of PM. Transport and transformations of precursors can occur over distances of hundreds of kilometers. The coarse PM constituents have shorter lifetimes in the atmosphere, so their effects tend to be more localized. Only major sources for each constituent within each broad category shown at the top of Table 3-8 are listed. Not all sources are equal in magnitude. Chemical characterizations

TABLE 3-8. CONSTITUENTS OF ATMOSPHERIC PARTICLES AND THEIR MAJOR SOURCES[1]

	Sources					
	Primary (PM <2.5 μm)		Primary (PM >2.5 μm)		Secondary PM Precursors (PM <2.5 μm)	
Aerosol species	Natural	Anthropogenic	Natural	Anthropogenic	Natural	Anthropogenic
$SO_4^{=}$ Sulfate	Sea spray	Fossil fuel combustion	Sea spray	—	Oxidation of reduced sulfur gases emitted by the oceans and wetlands and SO_2 and H_2S emitted by volcanism and forest fires	Oxidation of SO_2 emitted from fossil fuel combustion
NO_3^- Nitrate	—	—	—	—	Oxidation of NO_x produced by soils, forest fires, and lighting	Oxidation of NO_x emitted from fossil fuel combustion and in motor vehicle exhaust
Minerals	Erosion and re-entrainment	Fugitive dust paved and unpaved roads, agriculture, and forestry	Erosion and re-entrainment	Fugitive dust, paved and unpaved road dust, agriculture, and forestry	—	—
NH_4^+ Ammonium	—	—	—	—	Emissions of NH_3 from wild animals, and undisturbed soil	Emissions of NH_3 from animal husbandry, sewage, and fertilized land
Organic carbon (OC)	Wild fires	Prescribed burning, wood burning, motor vehicle exhaust, and cooking	—	Tire and asphalt wear and paved road dust	Oxidation of hydrocarbons emitted by vegetation (terpenes, waxes) and wild fires	Oxidation of hydrocarbons emitted by motor vehicles, prescribed burning, and wood burning
Elemental carbon (EC)	Wild fires	Motor vehicle exhaust, wood burning, and cooking	—	Tire and asphalt wear and paved road dust	—	—
Metals	Volcanic activity	Fossil fuel combustion, smelting, and brake wear	Erosion, re-entrainment, and organic debris	—	—	—
Bioaerosols	Viruses and bacteria	—	Plant and insect fragments, pollen, fungal spores, and bacterial agglomerates	—	—	—

[1]Dash (–) indicates either very minor source or no known source of component.

DRAFT-DO NOT QUOTE OR CITE

of primary particulate emissions for a wide variety of natural and anthropogenic sources (as shown in Table 3-8) were given in Chapter 5 of the 1996 PM AQCD. Summary tables of the composition of source emissions presented in the 1996 PM AQCD and updates to that information are provided in Appendix 3D. The profiles of source composition were based in large measure on the results of various studies that collected signatures for use in source apportionment studies.

Natural sources of primary PM include windblown dust from undisturbed land, sea spray, and plant and insect debris. The oxidation of a fraction of terpenes emitted by vegetation and reduced sulfur species from anaerobic environments leads to secondary PM formation. Ammonium (NH_4^+) ions, which play a major role in regulating the pH of particles, are derived from emissions of ammonia (NH_3) gas. Source categories for NH_3 have been divided into emissions from undisturbed soils (natural) and emissions that are related to human activities (e.g., fertilized lands, domestic and farm animal waste). There is ongoing debate about characterizing emissions from wild fires (i.e., unwanted fire) as either natural or anthropogenic. Wildfires have been listed in Table 3-8 as natural in origin, but land management practices and other human actions affect the occurrence and scope of wildfires. For example, fire suppression practices allow the buildup of fire fuels and increase the susceptibility of forests to more severe and infrequent fires from whatever cause, including lightning strikes. Similarly, prescribed burning is listed as anthropogenic, but can viewed as a substitute for wildfires that would otherwise occur eventually on the same land.

The transformations that gaseous precursors to secondary PM formation undergo after being emitted from the sources shown in Table 3-8 are described in Section 3.3.1. Aspects of the transport of primary PM and secondary PM, including the transport of material from outside the United States, are described in Section 3.3.2. A brief introduction to the deposition of particles is also given in Section 3.3.2, and a more detailed discussion of deposition processes is presented in Chapter 4. Methods to infer contributions from different source categories to ambient PM using receptor models and the results of these modeling efforts are given in Section 3.3.3. Estimates of emissions of primary PM and precursors to secondary PM from major sources are presented in Section 3.3.4. A discussion of the uncertainties associated with these emissions is given in Section 3.3.5.

3.3.1 Chemistry of Secondary PM Formation

The precursors to secondary PM have natural and anthropogenic sources, just as primary PM has natural and anthropogenic sources. Whereas the major atmospheric chemical transformations leading to the formation of particulate nitrate and sulfate have been relatively well understood; those involving the formation of secondary aerosol organic carbon are still under investigation. A large number of organic precursors are involved; many of the kinetic details still need to be determined; and many of the actual products of the oxidation of hydrocarbons have yet to be identified.

Formation of Sulfates and Nitrates

A substantial fraction of the fine particle mass, especially during the warmer months of the year, is secondary sulfate and nitrate formed as the result of atmospheric reactions. Such reactions involve the gas phase conversion of SO_2 to H_2SO_4 initiated by reaction with OH radicals and aqueous-phase reactions of SO_2 with H_2O_2, O_3, or O_2 (catalyzed by Fe and Mn). These heterogeneous reactions may occur in cloud and fog droplets or in films on atmospheric particles. NO_2 can be converted to HNO_3 by reaction with OH radicals during the day. At night, NO_2 also is oxidized to nitric acid by a sequence of reactions initiated by O_3 that produce nitrate radicals (NO_3) and dinitrogenpentoxide (N_2O_5) as intermediates. Both H_2SO_4 and HNO_3 react with atmospheric ammonia (NH_3). Gaseous NH_3 reacts with gaseous HNO_3 to form particulate NH_4NO_3. Gaseous NH_3 reacts with H_2SO_4 to form acidic HSO_4^- (in NH_4 HSO_4) as well as SO_4^{-2} in $(NH_4)_2SO_4$. In addition, acid gases such as SO_2 and HNO_3 may react with coarse particles to form coarse secondary PM containing sulfate and nitrate. Examples include reactions with basic compounds resulting in neutralization (e.g., $CaCO_3 + 2\ HNO_3 \rightarrow Ca\ (NO_3)_2 + H_2CO_3\uparrow$) or with salts of volatile acids resulting in release of the volatile acid (e.g., $SO_2 + 2NaCl + H_2O \rightarrow Na_2SO_3 + 2HCl\uparrow$).

If particulate NH_4NO_3 coagulates with an acidic sulfate particle (H_2SO_4 or HSO_4^-), gaseous HNO_3 will be released, and the NH_3 will increase the neutralization of the acidic sulfate. Thus, in the eastern United States, where PM tends to be acidic, sulfate is usually a larger fraction of PM mass than nitrate. However, in the western United States, where higher NH_3 and lower SO_2 emissions permit complete neutralization of H_2SO_4, the concentration of nitrate may be higher than that of sulfate. As SO_2 concentrations in the atmosphere in the eastern United States are

reduced, the NH_3 left in the atmosphere after neutralization of H_2SO_4 will be able to react with HNO_3 to form NH_4NO_3. Therefore, a reduction in SO_2 emissions, especially without a reduction in NO_x emissions, could lead to an increase in NH_4NO_3 concentrations (West et al., 1999; Ansari and Pandis, 1998). Thus, possible environmental effects of NH_4NO_3 are of interest for both the western and eastern United States.

Chemical reactions of SO_2 and NO_x within plumes are an important source of H^+, SO_4^{-2}, and NO_3^-. These conversions can occur by gas-phase and aqueous-phase mechanisms. In power-plant or smelter plumes containing SO_2 and NO_x, the gas-phase chemistry depends on plume dilution, sunlight, and volatile organic compounds either in the plume or in the ambient air mixing into and diluting the plume. For the conversion of SO_2 to H_2SO_4 in the gas-phase in such plumes during summer midday conditions in the eastern United States, the rate typically varies between 1 and 3% h^{-1} but in the cleaner western United States rarely exceeds 1% h^{-1}. For the conversion of NO_x to HNO_3, the gas-phase rates appear to be approximately three times faster than the SO_2 conversion rates. Winter rates for SO_2 conversion are approximately an order of magnitude lower than summer rates.

The contribution of aqueous-phase chemistry to particle formation in point-source plumes is highly variable, depending on the availability of the aqueous phase (wetted aerosols, clouds, fog, and light rain) and the photochemically generated gas-phase oxidizing agents, especially H_2O_2 for SO_2 chemistry. The in-cloud conversion rates of SO_2 to SO_4^{-2} can be several times larger than the gas-phase rates given above. Overall, it appears that SO_2 oxidation rates to SO_4^{-2} by gas-phase and aqueous-phase mechanisms may be comparable in summer, but aqueous-phase chemistry may dominate in winter. Further details concerning the chemistry of SO_2 and NO_x in power plant plumes can be found in Hewitt (2001).

In the western United States, markedly higher SO_2 conversion rates have been reported in smelter plumes than in power plant plumes. The conversion occurs predominantly by a gas-phase mechanism. This result is attributed to the lower NO_x in smelter plumes. In power plant plumes, NO_2 depletes OH radicals and competes with SO_2 for OH radicals.

In urban plumes, the upper limit for the gas-phase SO_2 conversion rate appears to be about 5% h^{-1} under the more polluted conditions. For NO_2, the rates appear to be approximately three times faster than the SO_2 conversion rates. Conversion rates of SO_2 and NO_x in background air are comparable to the peak rates in diluted plumes. Neutralization of H_2SO_4 formed by SO_2

conversion increases with plume age and background NH_3 concentration. If the NH_3 concentrations are more than sufficient to neutralize H_2SO_4 to $(NH_4)_2SO_4$, the HNO_3 formed from NO_x conversions may be converted to NH_4NO_3.

Formation of Secondary Organic Particulate Matter (SOPM)

Atmospheric reactions involving volatile organic compounds such as alkanes, alkenes, aromatics, cyclic olefins, and terpenes (or any reactive organic gas that contains at least seven carbon atoms) yield organic compounds with low saturation vapor pressures at ambient temperature. Such reactions may occur in the gas phase, in fog or cloud droplets (Graedel and Goldberg, 1983; Faust, 1994), or possibly in aqueous aerosols (Aumont et al., 2000). Reaction products from the oxidation of reactive organic gases also may nucleate to form new particles or condense on existing particles to form secondary organic PM (SOPM). Organic compounds with two double bounds may react to form dicarboxylic acids, which, with four or more carbon atoms, also may condense. Both biogenic and anthropogenic sources contribute to primary and secondary organic particulate matter (Grosjean, 1992; Hildemann et al., 1996; Mazurek et al., 1997; Schauer et al., 1996). Oxalic acid was the most abundant organic acid found in $PM_{2.5}$ in California (Poore, 2000).

Although the mechanisms and pathways for forming inorganic secondary particulate matter are fairly well known, those for forming SOPM are not as well understood. Ozone and the OH radical are thought to be the major initiating reactants. However, HO_2 and NO_3 radicals also may initiate reactions and organic radicals may be nitrated by HNO_2, HNO_3, or NO_2. Pun et al. (2000) discuss formation mechanisms for highly oxidized, multifunctional organic compounds. The production of such species has been included in a photochemical model by Aumont et al. (2000), for example. Understanding the mechanisms of formation of secondary organic PM is important because SOPM can contribute in a significant way to ambient PM levels, especially during photochemical smog episodes. Experimental studies of the production of secondary organic PM in ambient air have focused on the Los Angeles Basin. Turpin and Huntzicker (1991, 1995) and Turpin et al. (1991) provided strong evidence that secondary PM formation occurs during periods of photochemical ozone formation in Los Angeles and that as much as 70% of the organic carbon in ambient PM was secondary in origin during a smog episode in 1987. Schauer et al. (1996)

estimated that 20 to 30% of the total organic carbon PM in the <2.1 μm size range in the Los Angeles airshed is secondary in origin on an annually averaged basis.

Pandis et al. (1992) identified three mechanisms for formation of SOPM: (1) condensation of oxidized end-products of photochemical reactions (e.g., ketones, aldehydes, organic acids, and hydroperoxides), (2) adsorption of semivolatile organic compounds (SVOC) onto existing solid particles (e.g., polycyclic aromatic hydrocarbons), and (3) dissolution of soluble gases that can undergo reactions in particles (e.g., aldehydes). The first and third mechanisms are expected to be of major importance during the summertime when photochemistry is at its peak. The second pathway can be driven by diurnal and seasonal temperature and humidity variations at any time of the year. With regard to the first mechanism, Odum et al. (1996) suggested that the products of the photochemical oxidation of reactive organic gases are semivolatile and can partition themselves onto existing organic carbon at concentrations below their saturation concentrations. Thus, the yield of SOPM depends not only on the identity of the precursor organic gas but also on the ambient levels of organic carbon capable of absorbing the oxidation products.

Haagen-Smit (1952) first demonstrated that hydrocarbons irradiated in the presence of NO_x produce light scattering aerosols. The aerosol forming potentials of a wide variety of individual anthropogenic and biogenic hydrocarbons were compiled by Pandis et al. (1992), based mainly on estimates made by Grosjean and Seinfeld (1989) and data from Pandis et al. (1991) for β-pinene and from Izumi and Fukuyama (1990) for aromatic hydrocarbons. Zhang et al. (1992) examined the oxidation of α-pinene. Pandis et al. (1991) found no aerosol products formed in the photochemical oxidation of isoprene, although they and Zhang et al. (1992) found that the addition of isoprene to reaction mixtures increased the reactivity of the systems studied. Further details about the oxidation mechanisms and secondary organic PM yields from various reactive organic gases are given in the above studies. Estimates of the production rate of secondary organic PM in the Los Angeles airshed are provided in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996).

More recently, Odum et al. (1997a,b) have found that the aerosol formation potential of whole gasoline vapor can be accounted for solely by summing the contributions of the individual aromatic compounds in the fuel. In general, data for yields for secondary organic PM formation can be broken into two distinct categories. The oxidation of toluene and aromatic compounds containing ethyl or propyl groups (i.e., ethylbenzene, ethyltoluene, n-propylbenzene) produced

higher yields of secondary organic PM than did the oxidation of aromatic compounds containing two or more methyl groups (i.e., xylenes, di-, tri-, tetra-methylbenzenes). Yields in the first group ranged from about 7 to 10%; and in the second group, they ranged from 3 to 4% for organic carbon concentrations between 13 and 100 $\mu g/m^3$. Reasons for the differences in secondary organic PM yields found between the two classes of compounds are not clear.

There have been a few recent studies that have examined the composition of secondary organic PM. Edney et al., (2001) carried out a smog chamber study to investigate the formation of multi-functional oxygenates from photooxidation of toluene. The experiments were carried out by irradiating toluene/propylene/NO_x/air mixtures in a smog chamber operated in the dynamic mode and analyzing the collected aerosol by positive chemical ionization GC-MS after derivatization of the carbonyl oxidation products. The results of the GC-MS analyses were consistent with the formation of semivolatile multi-functional oxygenates, including hydroxy diones as well as triones, tetraones, and pentaones. The authors also suggested that some of these compounds could be present in SOPM in the form of polymers.

Jang and Kamens (2001a) employed a number of analytical approaches, including GC-MS detection of volatile derivatives of carbonyl, hydroxy, and acid compounds in SOPM formed in the irradiation of toluene/NO_x mixtures. A wide range of substituted aromatics, nonaromatic ring-retaining and ring-opening products were detected. Newly identified ring opening oxycarboxylic acids detected included: glyoxylic acid; methylglyoxylic acid; 4-oxo-2-butenoic acid; oxo-C_5-alkenoic acids; dioxopentenoic acids; oxo-C_7-alkadienoic acids; dioxo-C_6-alkenoic acids; hydroxydioxo-C_7-alkenoic acids; and hydroxytrioxo-C_6-alkanoic acids. Other newly identified compounds included methylcyclohexenetriones; hydroxymethylcyclohexenetriones; 2-hydroxy-3-penten-1,5-dial, hydroxyoxo-C_6-alkenals; hydroxy-C_5-triones, hydroxydioxo-C_7-alkenals; and hydroxy-C_6-tetranones. Included among these compounds were a number of the hydroxy polyketones detected by Edney et al., (2001). Recent laboratory and field studies support the concept that nonvolatile and semivolatile oxidation products from the photooxidation of biogenic hydrocarbons contribute significantly to ambient PM concentrations in both urban and rural environments. The oxidation of a variety of biogenic hydrocarbons emitted by trees and plants, such as terpenes (α-pinene, β-pinene, Δ^3-carene, sabinene, α-terpinene, γ-terpinene, terpinolene, myrcene, and ocimene) and sesequiterpenes (β-caryophyllene and α-humulene) could form SOPM. Vegetation also emits oxygenated organic compounds such as alcohols,

acetates, aldehyde, ketones, ethers, and esters (Winer et al., 1992). However, their contribution to SOPM, remains uncertain. Hoffmann et al. (1997) found SOPM yields of ≈5% for open-chain biogenic hydrocarbons such as ocimene and linalool; 5 to 25% for monounsaturated cyclic monoterpenes such as α-pinene; Δ-3 carene and terpinene-4-ol; and ≈40% for a cyclic monoterpene with an endocyclic and an exocyclic double bond such as d-limonene. Secondary organic PM yields of close to 100% were observed during the photochemical oxidation of one sesquiterpene, trans-caryophyllene. These results were all obtained for initial hydrocarbon mixing ratios of 100 ppb, which are much higher than found in the atmosphere.

Kamens et al. (1999) observed SOPM yields of 20 to 40% for α-pinene. Using information on the composition of secondary PM formed from α-pinene (Jang and Kamens, 1999), they were able to calculate formation rates with a kinetic model including formation mechanisms for O_3 + α-pinene reaction products.

Griffin et al. (1999) introduced the concept of incremental aerosol reactivity, the change in the secondary organic aerosol mass produced (in $\mu g/m^3$) per unit change of parent organic reacted (in ppb), as a measure of the aerosol-forming capability of a given parent organic compound in a prescribed mixture of other organic compounds. They measured the incremental aerosol reactivity for a number of aromatic and biogenic compounds for four initial mixtures. Incremental aerosol reactivity ranged from 0.133 to 10.352 μgm^{-3} ppb^{-1} and varied by almost a factor of two depending on the initial mixture.

A number of multifunctional oxidation products produced by the oxidation of biogenic hydrocarbons have been identified in laboratory studies (Yu et al., 1998; Glasius et al., 2000; Christoffersen et al., 1998; Koch et al., 2000; and Leach et al., 1999). Many of these compounds have subsequently been identified in field investigations (Yu et al., 1999; Kavouras et al., 1998, 1999a,b; Pio et al., 2001; and Castro et al., 1999). Most studies of the formation of secondary organic aerosol formation from terpenes have focused on their reactions with ozone. There have been many fewer studies dealing with the oxidation of terpenes initiated by OH radicals. Larson et al. (2001) found that the major aerosol products produced ultimately from the reaction of OH radicals with mono-terpenes with endocyclic double bonds (α-pinene, 3-carene) were C_{10} kato-carboxylic acids (such as pinonic and caronic acids); whereas the major products from the oxidation of mono-terpenes with exocyclic double bonds (β-pinene) were C_9-dicarboxylic acids (such as pinic acid), and the major product from the oxidation of limonene (which has both

endo- and exocyclic double bonds) was 3-acetyl-6-oxo-heptanal (Kato-limonaldehyde). A large number of related aldehydes, ketones and acids were also found in their experiments. However, the total yields of condensable products are much lower than for the corresponding reactions with ozone. For example, yields of C_9-dicarboxylic acids, C_{10}-hydroxy-keto-carboxylic acids, and C_{10}-hydroxy-Kato-aldehydes from the reaction of ozone with mono-terpenes with endocyclic double bonds ranged form 3% to 9%; whereas they ranged only from 0.4 to 0.6% in the reaction with OH radicals. Likewise, the reaction of mono-terpenes with exocyclic double bonds with ozone produced much higher yields (1% to 4%) of C_8- and C_9-dicarboxylic acids than did their reaction with OH radicals (0.2% to 0.3%). Apart from the complex products noted above, it should be remembered that much simpler products, such as formaldehyde and formic acid, are also formed in much larger yields form the same reactants (e.g., Winterhalter et al., 2000). Compounds such as these also contribute to the formation of secondary organic aerosol according to the mechanisms given in Pandis et al. (1992) and mentioned earlier in this section.

It is worth noting that the dicarboxylic acids and hydroxy-Kato-carboxylic acids have very low vapor pressures and may act as nucleating species in OH- and O_3- terpene reactions (Larson et al., 2001). The rate coefficient for reaction of α-pinene with OH radicals is approximately a factor of 10^6 greater than for its reaction with O_3, based on data given in Atkinson (1994). The daytime average concentration of O_3 is typically a factor of 10^6 greater than that for OH radicals in polluted boundary layers; whereas the above mentioned yields of aerosol products are roughly a factor of ten greater in the O_3-initiated reaction than in the corresponding OH radical reaction. The foregoing suggests that the O_3-initiated reaction may be more important than the OH initiated reaction for the formation of aerosol products. Because ambient ozone is present at night and it penetrates indoors, new particles may also be generated under these conditions. For example, Wainman et al. (2000) found that ozone can react with limonene released by air fresheners in indoor environments to produce substantial quantities of submicron particles. The corresponding reaction involving OH radicals at night and in indoor environments is expected to be negligible by comparison because of the very low OH concentrations present in these environments. Although much progress has been made in determining the importance of anthropogenic and biogenic hydrocarbons for the formation of secondary organic PM, further investigations are needed to accurately assess their overall contributions to $PM_{2.5}$ concentrations.

Reactions of organic compounds either in particles or on the surface of particles have only recently come under study. Tobias and Ziemann (2000) reported evidence for the formation of relatively stable low volatility peroxy hemiacetals from reactions of hydroperoxides with aldehydes and ketones on the surface of secondary organic particles. Shortly after the publication of these results, Jang and Kamens (2001a) suggested, based on results of their laboratory investigations of SOPM formation from irradiation of toluene/propylene/NO_x /air mixtures, that carbonyls and hydroxy compounds (either within or on the surface of aromatic SOPM) could react together to form larger and less volatile hemiacetals and acetals. They also proposed that dissolved carbonyls could undergo further reactions leading to the formation of a polymer, a mechanism that has also been suggested by Edney et al. (2001). Each of these mechanisms that also could be catalyzed by the presence of acids involves converting, through heterogenous reactions, volatile compounds into much less volatile compounds, a mechanism that could contribute to SOPM yields in aromatic and possibly biogenic systems.

As a first step in addressing these issues, Jang and Kamens (2001a) carried out a series of laboratory screening experiments to assess whether volatile carbonyl compounds absorbed into particles could undergo further chemical reactions forming low vapor pressure compounds. Experiments were carried out whereby carbonyls were introduced in Teflon bags in the dark in the presence of a seed aerosol containing either ammonium sulfate or a mixture of ammonium sulfate and sulfuric acid. The increase in the aerosol volume was then measured using a scanning mobility particle sizer. The carbonyls employed for the study included glyoxal, hexanal, and octanal. Increased organic aerosol yields were found in the presence of the ammonium sulfate seed aerosol for each of the carbonyls, with the highest yield being found for octanol followed in decreasing order by glyoxal and then octanal. The presence of the acidified sulfate salt significantly increased the yields even further. In a number of other experiments, 1-decanol was added to the carbonyl-aerosol system to investigate the possible formation of hemiacetals and/or acetals. Again, the volume of aerosol increased in both the presence of ammonium sulfate aerosol and the acidified salt with a significantly larger yield found in the presence of acidity.

To explain their findings for acid-catalyzed carbonyl reactions, Jang and Kamens (2001a,b) proposed a chemical mechanism in which the dissolved carbonyl first undergoes a protonization reaction forming an adduct that can react with water to form its hydrate (1,1-dihydroxy gem-diol). The adducts can then react with OH groups of the gem-diol forming higher molecular

weight and less volatile dimers that are subject to further reactions. In principal, this process, which the authors refer to as a "zipping reaction" can lead to the formation of polymers. However, because the individual reactions are reversible, the process can also be reversed by an unzipping reaction. The zipping process could serve as an important mechanism for SOPM formation by converting volatile oxidation products including glyoxal and methyl glyoxal into low volatility compounds. On the other hand, the unzipping process that could take place during the workup of the aerosol samples could be responsible for the detection of high volatile oxidation products in SOPM, including glyoxal and methyl glyoxal that has been reported by Edney et al. (2001), Cocker et al. (2001), and Jang and Kamens (2001a). While these processes may take place in the absence of significant acidity, the experimental results suggest the processes are likely enhanced by acid-catalyzed reactions.

Sampling and characterizing PM in the ambient atmosphere and in important microenvironments is required to address important issues in exposure, toxicology, and compliance. Currently, it is not possible to fully quantify the concentration, composition, or sources of the organic components. Many of the secondary organic aerosol components are highly oxidized, difficult to measure, multifunctional compounds. Additional laboratory studies are needed to identify such compounds, strategies need to be developed to sample and measure such compounds in the atmosphere, and models of secondary organic aerosol formation need to be improved and added to air quality models in order to address issues related to human exposure.

A high degree of uncertainty is associated with all aspects of the calculation of secondary organic PM concentrations. This is compounded by the volatilization of organic carbon from filter substrates during and after sampling as well as potential positive artifact formation from the absorption of gaseous hydrocarbon on quartz filters. Significant uncertainties always arise in the interpretation of smog chamber data because of wall reactions. Limitations also exist in extrapolating the results of smog chamber studies to ambient conditions found in urban airsheds and forest canopies. Concentrations of terpenes and NO_x are much lower in forest canopies (Altshuller, 1983) than the levels commonly used in smog chamber studies. The identification of aerosol products of terpene oxidation has seldom been a specific aim of field studies, making it difficult to judge the results of model calculations of secondary organic PM formation. Uncertainties also arise because of the methods used to measure biogenic hydrocarbon emissions.

Khalil and Rasmussen (1992) found much lower ratios of terpenes to other hydrocarbons (e.g., isoprene) in forest air than were expected based on their relative emissions strengths and rate coefficients for reaction with OH radicals and O_3. They offered two explanations: (1) either the terpenes were being removed rapidly by some heterogeneous process or (2) emissions were enhanced artificially by feedbacks caused by the bag enclosures they used. If the former consideration is correct, then the production of aerosol carbon from terpene emissions could be substantial; if the latter is correct, then terpene emissions could have been overestimated by the techniques used.

3.3.2 The Long-Range Transport of Particulate Matter from Outside the United States

Apart from sources within the continental United States, particulate matter can be brought in by long-range transport from sources outside the United States. For example, the transport of PM from uncontrolled biomass burning in Central America and southern Mexico resulted in anomalously high PM levels observed in southern Texas and generally elevated PM concentrations throughout the entire central and southeastern United States during the spring and early summer of 1998. Windblown dust from individual dust storms in the Sahara desert has been observed in satellite images as plumes crossing the Atlantic Ocean and reaching the southeast coast of the United States (e.g., Ott et al., 1991). Dust transport from the deserts of Asia across the Pacific Ocean also occurs (Prospero, 1996). Most dust storms in the deserts of China occur in the spring following the passage of strong cold fronts after the snow has melted and before a surface vegetation cover has been established. Strong winds and unstable conditions result in the rapid transport of dust to altitudes of several kilometers, where it is transported by strong westerly winds out over the Pacific Ocean (Duce, 1995). Satellite images were used to track the progress of a dust cloud from the Gobi desert to the northwestern United States during the spring of 1998 (Husar et al., 2000).

Satellite images obtained at visible wavelengths cannot track mineral dust across the continents because of a lack of contrast between the plume and the underlying surface. Other means must be used to track the spread of North African dust through the eastern United States. Perry et al. (1997) used two criteria ($PM_{2.5}$ soil concentration > 3 $\mu g/m^3$ and Al/Ca > 3.8) to distinguish between soil of local origin from soil originating in North Africa in characterizing the

sources of PM in aerosol samples collected in the IMPROVE (Interagency Monitoring of Protected Visual Environments) network. North African dust has been tracked as far north as Illinois (Gatz and Prospero, 1996) and Maine (Perry et al., 1997). The analysis of Perry et al. (1997) indicates that incursions of Saharan dust into the continental United States have occurred, on average, about three times per year from 1992 to 1995. These events persist for about 10 days on average, principally during the summer. Large scale dust events typically cover from 15 to 30% of the area of the continental United States and result in increases of $PM_{2.5}$ levels of 8.7 ± 2.3 $\mu g/m^3$ throughout the affected areas, with mean maximum dust contributions of 19.7 ± 8.4 $\mu g/m^3$ during these events and a peak contribution of 32 $\mu g/m^3$ to 24-h average $PM_{2.5}$ levels.

As can be expected, the frequency of dust events is highest in the southeastern United States. About half of these events are observed only within the state of Florida, and these events are associated with dense hazes in Miami (Figure 3-22) during the summer (Prospero et al., 1987). North African dust is the dominant aerosol constituent in southern Florida during the summer; whereas soil dust constitutes only a minor fraction of PM during the remainder of the year (Prospero, 1999). Approximately one-third to one-half of the mass of the particles reaching southern Florida have aerodynamic diameters less than 2.5 micrometers (Prospero et al., 2001). During episodes when daily total dust concentrations ranged up to 100 $\mu g/m^3$, it can be seen that daily $PM_{2.5}$ concentrations of up to 50 $\mu g/m^3$ could have resulted in Miami, FL.

Husar et al. (2000) documented the transport of dust from the Gobi and Taklimakan deserts to North America during April 1998. The PM_{10} concentration averaged over 150 stations in Washington, Oregon, California, Nevada, and Idaho reporting data to AIRS was 65 $\mu g/m^3$ between April 26 and May 1, compared to about 20 $\mu g/m^3$ during the rest of April and May. Data from several networks indicated that PM_{10} concentrations were over 100 $\mu g/m^3$ in central British Columbia, Washington State, and Oregon. The highest PM concentrations observed were 120 $\mu g/m^3$ for PM_{10} and 50 $\mu g/m^3$ for $PM_{2.5}$ at Chilliwack Airport in northwestern Washington State (Figure 3-23). Aircraft measurements made over the northwestern United States were consistent with a mass median diameter of the dust being between 2 and 3 μm.

Desert dust deposited over oceans provides nutrients to marine ecosystems (Savoie and Prospero, 1980). Desert dust deposited on nutrient depleted soils also provides nutrients, as in Hawaiian rain forests (Chadwick et al., 1999). Microorganisms, including various species and



Figure 3-22. Monthly average Saharan dust components of the aerosol sampled in Miami, FL, from 1974 to 1996. Approximately one-third to one-half of Saharan dust is in the $PM_{2.5}$ size range.

Source: Prospero (1999).

genera of fungi and bacteria, have been found attached to African dust particles in the U.S. Virgin Islands (Griffin et al., 2001). The fungus, *Aspergillus sydowii*, which has been connected to the death of coral reefs, has been identified in air samples collected in the Caribbean during African dust transport events (Smith et al., 1996; Shinn et al., 2000). Measurements of the composition of Saharan dust in Miami indicate enhancements of nitrate, non-sea-salt sulfate, ammonium, and trace metals over concentrations expected for clean marine air, suggesting pollution emitted in Europe and North Africa as sources (Prospero, 1999). It is likely that many other constituents will be found associated with dust from outside North America as more measurements are made. It should be noted that, as North African dust and associated material



Figure 3-23. $PM_{2.5}$ and PM_{10} concentrations measured at Chilliwack Airport, located in northwestern Washington State, just before and during the Asian desert dust episode of April and May 1998.

Source: U.S. EPA Aerometric Information Retrieval System (AIRS).

are transported northward through the United States during the summer, they are added to the mixture of primary and secondary PM generated domestically.

Biomass burning for agricultural purposes occurs normally during the spring of each year in Central America and southern Mexico. During the spring of 1998, fires burned uncontrollably because of abnormally hot and dry conditions associated with the intense El Niño of 1997 to 1998. PM_{10} concentrations observed in the southern Rio Grande Valley were elevated substantially during the passage northward of the biomass burning plume produced by these fires as shown in Figure 3-24. Elevated PM_{10} concentrations also were found as far north as St. Louis, MO (Figure 3-25). As can be seen from Figure 3-24 and Figure 3-25, the elevations in PM



Figure 3-24. Time series of 24-h average PM_{10} concentrations observed in the Rio Grande Valley during May 1998.

Source: U.S. EPA Aerometric Information Retrieval System (AIRS).



Figure 3-25. PM_{10} concentrations observed in St. Louis, MO, during May 1998.

Source: U.S. EPA Aerometric Information Retrieval System (AIRS).

concentrations were limited in duration. Uncontrolled wildfires occur in the United States every year, but their effects on air quality throughout the United States still need to be evaluated systematically. These fires can be widespread, and the frequency of their occurrence can vary markedly from year to year. For example, approximately 26,000 km^2 were consumed during 2000, but only a small fraction of this area was burnt during 2001 in the western United States. Wildfires also occur in the boreal forests of northwestern Canada. Wotawa and Trainer (2000) suggested that the plume from fires occurring in the Northwest Territories of Canada in early July 1995 may have extended throughout most of the eastern United States, resulting in elevated levels of CO and ozone. Simple scaling of their calculated excess CO concentrations because of the fires, by the ratio of emission factors of $PM_{2.5}$ to CO, indicates that the excess $PM_{2.5}$ concentrations in the plume may have ranged from about 5 $\mu g/m^3$ in the Southeast and increasing to close to 100 $\mu g/m^3$ in the northern Plains States.

3.3.3 Source Contributions to Ambient PM Determined by Receptor Models

Receptor models are perhaps the primary means used to estimate the contributions of different source categories to PM concentrations at individual monitoring sites. Dispersion models (i.e., three-dimensional chemistry and transport models) are formulated in a prognostic manner (i.e., they attempt to predict species concentrations using a tendency equation that includes terms based on emissions inventories, atmospheric transport, chemical transformations, and deposition). Receptor models are diagnostic in their approach (i.e., they attempt to derive source contributions based either on ambient data alone or in combination with data from the chemical composition of sources). These methods have the advantage that they do not invoke all of the uncertainties inherent in emissions inventories or in parameterizing atmospheric transport processes in grid point models.

There are two main approaches to receptor modeling. Receptor models such as the chemical mass balance (CMB) model (Watson et al., 1990a) relate source category contributions to ambient concentrations based on analyses of the composition of ambient particulate matter and source emissions samples. This technique has been developed for apportioning source categories of primary particulate matter and was not formulated to include the processes of secondary particulate matter formation. In the second approach, various forms of factor analysis are used, which rely on the analysis of time series of compositional data from ambient samples to derive

both the composition of sources and the source contributions. Standard approaches such as factor analysis or Principal Component Analysis (PCA) can apportion only the variance and not the mass in an aerosol composition data set. The other techniques described below, PMF and UNMIX do apportion mass, however. Positive matrix factorization (PMF) is a recently developed multivariate technique (Paatero and Tapper, 1993 and 1994) that overcomes many of the limitations of standard techniques, such as principal components analysis (PCA), by allowing for the treatment of missing data and data near or below detection limits. This is accomplished by weighting elements inversely according to their uncertainties. Standard methods such as PCA weight elements equally regardless of their uncertainty. Solutions also are constrained to yield non-negative factors. Both the CMB and the PMF approaches find a solution based on least squares fitting and minimize an object function. Both methods provide error estimates for the solutions based on estimates of the errors in the input parameters. It should be remembered that the error estimates often contain subjective judgments. For a complete apportionment of mass, all of the major sources affecting a monitoring site must be sampled for analysis by CMB; whereas there is no such restriction in the use of PMF.

Among other approaches, the UNMIX model takes a geometric approach that exploits the covariance of the ambient data to determine the number of sources, the composition and contributions of the sources, and the uncertainties (Henry, 1997). A simple example may help illustrate the approach taken by UNMIX. For example, in a two-element scatter plot of ambient Al and Si, a straight line and a high correlation for Al versus Si can indicate a single source for both species (soil), while the slope of the line gives information on the composition of the soil source. In the same data set, iron may not plot on a straight line against Si, indicating other sources of Fe in addition to soil. More importantly, the Fe-Si scatter plot may reveal a lower edge. The points defining this edge represent ambient samples collected on days when the only significant source of Fe was soil. Success of the UNMIX model hinges on the ability to find these “edges” in the ambient data from which the number of sources and the source compositions are extracted. UNMIX uses principal component analysis to find edges in m-dimensional space, where m is the number of ambient species. The problem of finding edges is more properly described as finding hyperplanes that define a simplex. The vertices at which the hyperplanes intersect represent pure sources from which source compositions can be determined. However, there are measurement errors in the ambient data that “fuzz” the edges making them difficult to

find. UNMIX employs an "edge-finding" algorithm to find the best edges in the presence of error. UNMIX does not make explicit use of errors or uncertainties in the ambient concentrations, unlike the methods outlined above. This is not to imply that the UNMIX approach regards data uncertainty as unimportant, but rather that the UNMIX model results implicitly incorporate error in the ambient data. The underlying philosophy is that the uncertainties are often unquantifiable, and hence it is best to make no *a priori* assumptions about what they are.

In addition to chemical speciation data, Norris et al. (1999) showed that meteorological indices could prove useful in identifying sources of particulate matter that are responsible for observed health effects (specifically asthma) associated with exposure to particulate matter. They examined meteorology associated with elevated pollution events in Spokane and Seattle, WA, and identified a "stagnation index" that was associated with low wind speeds and increases in concentrations of combustion-related pollutants. Their factor analysis also identified a meteorological index (low relative humidity and high temperatures) that was associated with increases in soil-derived particulate matter, as well as a third factor (low temperatures and high relative humidity) that was associated with increases in concentrations of particulate sulfate and nitrate species (Norris, 1998).

Ondov (1996) examined the feasibility of using sensitive isotopic and elemental tracer materials to determine the contributions of petroleum-fueled sources of PM_{10} in the San Joaquin Valley, CA. Costs of these experiments are affected not only by the tracer materials cost, but also by the sensitivities of the analytical methods for each, as well as the background levels of the tracers. Suarez et al. (1996) used iridium as a tracer to tag emissions from diesel-burning sanitation trucks in Baltimore and determined the size distribution of soot from the trucks.

A number of specialty conference proceedings, review articles, and books have been published that provide greater detail about source category apportionment receptor models then described in the 1996 PM AQCD. A review of the various methods used to apportion PM in ambient samples among its source categories was given in Section 5.5.2 of the 1996 PM AQCD. The collection of the source category characterization profiles shown in Appendix 3D has been motivated in many cases by the need to use them in receptor modeling applications.

The results of several source apportionment studies are discussed in this section to provide an indication of the relative importance of different sources of particulate matter across the

United States. First, results obtained mainly by using the chemical mass balance (CMB) approach for estimating contributions to $PM_{2.5}$ from different source categories at monitoring sites in the United States are discussed and presented in Table 3-9. More recent results using the PMF approach are included for Phoenix, AZ. Results obtained at a number of monitoring sites in the central and western United States by using the CMB model for PM_{10} are shown in Table 3-10. The sampling sites represent a variety of different source characteristics within different regions of Arizona, California, Colorado, Idaho, Illinois, Nevada, and Ohio. Definitions of source categories also vary from study to study. The results of the PM_{10} source apportionment studies were given in the 1996 PM AQCD and are presented here to allow easy comparison with results of $PM_{2.5}$ source apportionment studies. Chow and Watson (2002) present a detailed comparison of numerous studies using the CMB model performed mainly after 1995.

There are several differences between the broadly defined source categories shown at the tops of Tables 3-9 and 3-10. These differences reflect the nature of sources that are important for producing fine and coarse particulate matter shown in Table 3-8. They also are related to improvements in the ability to distinguish between sources of similar nature (e.g., diesel and gasoline vehicles, meat cooking, and vegetation burning). The use of organic tracers allows motor vehicle emissions to be broken down into contributions from diesel and gasoline vehicles. In studies where this distinction cannot be made, the source type is listed as 'total motor vehicles' in the tables. The studies that were reported to be able to distinguish gasoline from diesel fueled vehicles found that gasoline vehicles make significant, and sometimes the dominant, contributions to ambient $PM_{2.5}$ concentrations. Meat cooking is also distinguished from vegetation burning in more recent studies, although both are considered to be part of biomass burning. Vegetation burning consists of contributions from residential fuel wood burning, wildfires, prescribed burning, and burning of agricultural and other biomass waste. Miscellaneous sources of fine particles include contributions from combustion sources; whereas miscellaneous sources of coarse particles consist of contributions from soil and sea spray and industrial processing of geological material (e.g., cement manufacturing). Although a large number of elements and chemical components are used to differentiate among source categories and although there can be a large number of source types affecting a given site, only a few broadly defined source types are needed to account for most of the mass of $PM_{2.5}$ and PM_{10}.

TABLE 3-9. RECEPTOR MODEL SOURCE CONTRIBUTIONS TO $PM_{2.5}$

Sampling Site	Measured $PM_{2.5}$ Concentration	% Contribution[3] Total Motor Vehicles	Diesel	Gasoline Vehicles	Road Dust, Soil	Vegetation Burning	Secondary Sulfate	Secondary Nitrate	Misc. Source 1	Misc. Source 2	Misc. Source 3	Misc. Source 4	Total % Allocated
Pasadena, CA 1982[1]	28.2	—	18.8	5.7	12.4	9.6	20.9	7.4	5.3[a]	9.2[b]	8.5[c]	1.1[d]	98.9
Downtown LA, CA 1982[1]	32.5	—	35.7	6.5	11.1	5.8	20.3	9.2	3.7[a]	9.2[b]	5.2[c]	0.6[d]	107.3
West LA, CA 1982[1]	24.5	—	18.0	5.7	12.2	11.0	24.1	7.8	4.1[a]	9.4[b]	8.2[c]	1.6[d]	102.1
Rubidoux, CA 1982[1]	42.1	—	12.8	0.7	13.1	1.2	13.8	24.7	4.5[a]	12.1[b]	4.5[c]	0.5[d]	87.9
Sacramento, CA Winter 1991-96[2]	39.5	24.5[e]	—	—	1.2	18.1	4.5	36.6	—	—	—	—	84.9
Bakersfield, CA Winter 1996[3]	52	16[e]	—	—	<3	20	7	34	—	—	—	—	<80
Fresno, CA Winter 1996[3]	63	13[e]	—	—	<3	19	5	32	—	—	—	—	<85
Philadelphia, PA Summer 1982[4]	27.0	8.5[e]	—	—	4.4	—	81.9[f]	—	2.2[g]	1.9[h]	0.4[i]	—	99.3
Camden, NJ Summer 1982[4]	28.3	9.2[e]	—	—	3.2	—	81.3[f]	0.4	2.5[g]	2.5[h]	0.7[i]	—	99.8
Clarksboro, NJ Summer 1982[4]	26.0	5.8[e]	—	—	2.7	—	84.6[f]	—	0.8[g]	1.5[h]	0.4[i]	—	95.8
Grover City, IL ENE[j] 1986-87[5]		—	—	—	2.3	—	83.2[f]	—	9.7[k]	3.0[l]	1.2[g]	—	99.4
Grover City, IL SSW[j] 1986-87[5]		—	—	—	—	—	59.0[f]	—	11.6[k]	11.9[l]	4.1[g]	4.6[m]	91.2
Grover City, IL WNW[j] 1986-87[5]		2.4[e]		—	5.1	—	88.5[f]	—	2.8[k]	—	—	—	98.8
Grover City, IL NNW[j] 1986-87[5]		—	—	—	3.1	—	86.6[f]	—	3.4[l]	3.0[n]	—	—	96
Reno, NV Sumer 1998[6]	7.8	68[e]	—	—	14.5	4	11	2	0.6[q]	—	—	—	100.1
Phoenix, AZ Summer 1995-98[7]	8.3	—	10.9	36.2	1.8	15.0	—	—	20.8[n]	4.9[r]	6.7[s]	3.6[q]	99.9
Phoenix, AZ Winter 1995-98[7]	13.8	—	14.5	38.9	1.1	8.9	—	—	9.5[n]	4.5[r]	18.7[s]	4.1[q]	100.2

[1]Schauer et al., 1996
[2]Motallebi, 1999
[3]Magliano et al., 1998
[4]Dzubay et al, 1988
[5]Glover et al., 1991
[6]Gillies et al., 2000
[7]Ramadan et al., 2000

[a]Secondary and other organic compounds
[b]Secondary ammonium
[c]Meat cooking
[d]Vegetative detritus
[e]Value represents sum of diesel and gasoline vehicle exhaust
[f]Including associated cations and water

[g]Incinerators
[h]Oil fly ash
[i]Fluidized catalyst cracker
[j]Wind direction
[k]Lead smelter
[l]Iron works
[m]Copper smelter

[n]Coal power plant
[o]As ammonium sulfate
[p]As ammonium nitrate
[q]Sea salt
[r]Wood burning
[s]Nonferrous smelting

DRAFT-DO NOT QUOTE OR CITE

TABLE 3-10. RECEPTOR MODEL SOURCE CONTRIBUTIONS TO PM_{10}

Sampling Site	Measured PM_{10} Concentration	% Contribution										Total % Allocated
		Primary Geological	Primary Construction	Primary Motor Vehicle Exhaust	Primary Vegetative Burning	Secondary Ammonium Sulfate	Secondary Ammonium Nitrate	Misc. Source 1	Misc. Source 2	Misc. Source 3	Misc. Source 4	
Craycroft, AZ Winter 1989-1990[y]	23.4	55.6	0.0	35.5	0.0	3.0	2.6	5.1[a]	0.0	0.0	0.0	101.8
Hayden 1, AZ 1986[z]	105.0	4.8	1.9[b]	0.0	0.0	3.8	0.0	70.5[c]	4.8[d]	1.0[e]	0.0	86.6
Hayden 2, AZ 1986[1]	59.0	35.6	6.8[b]	0.0	0.0	6.8	0.0	47.5[c]	0.0	1.7[e]	0.0	98.4
Rillito, AZ 1988[2]	79.5	53.7	17.4[b]	1.5[f]	0.0	0.0	0.0	14.6[g]	0.0	0.0	0.0	87.2
Bakerfield, CA 1988-1989[3]	79.6	53.9	2.0	9.7	8.2	6.9	16.0	1.3[m]	1.9[n]	0.8[k]	0.0	100.7
Crows Landing, CA 1988-1989[3]	52.5	61.3	0.0	4.2	6.5	5.3	12.4	1.0[m]	1.9[n]	2.3[k]	0.0	94.9
Fellows, CA 1988-1989[3]	54.6	53.1	2.6	3.8	6.2	9.3	13.7	12.8[m]	2.6[n]	2.6[k]	0.0	106.7
Fresno, CA 1988-1989[3]	71.5	44.5	0.0	9.5	7.1	5.0	14.5	0.4[m]	1.9[n]	0.1[k]	0.0	83
Indio, CA[4]	58.0	56.9	5.2	7.6	12.2	6.2	7.1	0.3[j]	1.7[h]	0.0	0.0	97.2
Kern Wildlife Refuge, CA 1988-1989[3]	47.8	31.6	4.2	4.6	8.4	6.9	3.1	1.0[m]	3.1[n]	1.5[k]	0.0	64.4
Long Beach, CA 1986[5]	51.9	39.9	0.0	9.8[i]	0.0	15.4	17.7	0.2[j]	3.9[h]	12.3[k]	0.0	63.2
Long Beach, CA Summer 1987[6]	46.1	24.1	0.0	13.7	0.0	23.6	1.7	0.2[j]	4.8[h]	0.0	0.0	68.1
Long Beach, CA Fall 1987[6]	96.1	11.8	0.0	44.5	0.0	4.0	24.1	0.0[j]	2.8[h]	0.0	0.0	87.2
Riverside, CA 1988[7]	64.0	50.9	0.0	10.9	0.0	7.5	33.4	0.5[j]	2.0[h]	1.7[o]	0.0	106.9
Rubidoux, CA 1986[5]	87.4	49.3	4.6	6.4[i]	0.0	7.3	24.4	0.3[j]	1.1[h]	6.8[k]	0.0	100.2
Rubidoux, CA Summer 1987[6]	114.8	30.4	3.9	15.1	0.0	8.3	23.9	0.0[j]	4.4[h]	0.0	0.0	86
Rubidoux, CA Fall 1987[6]	112.0	17.1	14.4	27.1	0.0	1.9	28.2	0.0[j]	1.0[h]	0.0	0.0	89.7
Rubidoux, CA 1988[7]	87.0	55.2	0.0	11.7	0.0	6.1	24.9	0.6[j]	1.7[h]	6.6[o]	0.0	106.8
San Nicolas Island, CA Summer 1987[6]	17.4	9.2	0.0	5.2	0.0	21.3	2.9	0.0[j]	24.7[h]	0.0	0.0	63.3
Stockton, CA 1989[3]	62.4	55.1	0.8	8.3	7.7	5.0	11.2	1.1[m]	2.9[n]	0.0[k]	0.0	92.1
Pocatello, ID 1990[8]	100.0	8.3	7.5[q]	0.1	0.0	0.0	0.0	0.0	0.0	84.1[r]	0.0	100
S. Chicago, IL 1986[9]	80.1	34.0	3.0	3.5	0.0	19.2[s]	—	18.9[t]	2.7[u]	0.0	0.0	81.3
S.E. Chicago, IL 1988[10]	41.0	35.9[v]	0.0	2.2[f]	0.0	18.8	—	2.0[t]	0.7[h]	2.7[w]	18.8[g]	81.1
Reno, NV 1986-87[11]	30.0	49.7	0.0	33.3	6.3	4.3	2.0	0.0	0.0	0.0	0.0	95.6

DRAFT-DO NOT QUOTE OR CITE

TABLE 3-10 (cont'd). RECEPTOR MODEL SOURCE CONTRIBUTIONS TO PM_{10}

Sampling Site	Measured PM_{10} Concentration	% Contribution										Total % Allocated
		Primary Geological	Primary Construction	Primary Motor Vehicle Exhaust	Primary Vegetative Burning	Secondary Ammonium Sulfate	Secondary Ammonium Nitrate	Misc. Source 1	Misc. Source 2	Misc. Source 3	Misc. Source 4	
Sparks, NV 1986-87[11]	41.0	36.8	0.0	28.3	32.7	6.6	2.2	0.0	0.0	0.5[k]	0.0	107.1
Follansbee, WV 1991[12]	66.0	15.2	0.0	53.0	0.0	24.2	—	14.1[t]	0.0	0.0	0.0	106.5
Mingo, OH 1991[12]	60.0	20.0	0.0	23.3	6.8	25.0	—	5.7[t]	18.3[x]	0.0	0.0	99.1
Steubenville, OH 1991[12]	46.0	18.0	0.0	30.4	1.7	30.4	—	8.3[t]	10.9[x]	0.0	0.0	99.7

[1]Chow et al., 1992a
[2]Garfield; Ryan et al., 1988
[3]Jail; Ryan et al., 1988
[4]Thanukos et al., 1992
[5]Chow et al., 1992b
[6]Kim et al., 1992
[7]Gray et al., 1988
[8]Watson et al., 1994
[9]Chow et al., 1992c
[10]Houck et al., 1992
[11]Hopke et al., 1988
[12]Vermette et al., 1992
[13]Chow et al., 1988
[14]Skidmore et al., 1992
[a]Smelter background aerosol
[b]Cement plant sources, including kiln stacks, gypsum pile, and kiln area
[c]Copper ore
[d]Copper tailings
[e]Copper smelter building
[f]Heavy-duty diesel exhaust emission
[g]Background aerosol
[h]Marine aerosol, road salt, and sea salt plus sodium nitrate
[i]Motor vehicle exhaust from diesel and leaded gasoline
[j]Residual oil combustion
[k]Secondary organic carbon
[l]Biomass burning
[m]Primary crude oil
[n]$NaCl + NaNO_3$
[o]Lime
[p]Road sanding material
[q]Asphalt industry
[r]Phosphorus/phosphate industry
[s]Regional sulfate
[t]Steel mills
[u]Refuse incinerator
[v]Local road dust, coal yard road dust, and steel haul road dust
[w]Incineration
[x]Unexplained mass

DRAFT-DO NOT QUOTE OR CITE

At any given site, #5 source types account for >65% of the mass of $PM_{2.5}$ (Table 3-9); and #5 source types account for >65% of the mass of PM_{10} (Table 3-10).

Secondary sulfate is the dominant component of $PM_{2.5}$ samples collected in the studies of Dzubay et al. (1988) and Glover et al. (1991). Both studies found that sulfate at their monitoring site arose from regionally dispersed sources. Sulfate also represents the major component of $PM_{2.5}$ found in monitoring studies in the eastern United States shown in Appendix 6A of the 1996 PM AQCD. Primary and secondary organic components also make major contributions to $PM_{2.5}$. Contributions from road dust and soils are relatively minor, typically constituting less than 10% of $PM_{2.5}$ in the studies shown in Table 3-9. Studies in the western United States shown in Table 3-9 have found larger contributions from motor vehicles, fugitive dust, and ammonium nitrate. The most notable difference in the relative importance of major source categories of $PM_{2.5}$ shown in Table 3-9 and PM_{10} shown in Table 3-10 involves crustal material, (e.g., soil, road dust), which represents about 40% on average of the total mass of PM_{10} in the studies shown in Table 3-10. The fraction is higher at sites located away from specific sources such as sea spray or smelters. Emissions of crustal material are concentrated mainly in the $PM_{10-2.5}$ size range.

In Table 3-10, primary motor vehicle exhaust contributions account for up to 40% of average PM_{10} at many of the sampling sites. Vehicle exhaust contributions are also variable at different sites within the same study area. The mean value and the variability of motor vehicle exhaust contributions reflects the proximity of sampling sites to roadways and traffic conditions during the time of sampling. Many studies were conducted during the late 1980s, when a portion of the vehicle fleet still used leaded gasoline. Pb and Br in motor vehicle emissions facilitated the distinction of motor vehicle contributions from other sources. Vehicles using leaded fuels have higher emission rates than vehicles using unleaded fuels. Pb also poisons automobile exhaust catalysts and produces adverse human health effects. As a result, Pb has been eliminated from vehicle fuels. However, organic species such as n-pentacosane through n-nonacosene, cholestanes, ergostanes, sitostanes, and hopanes have replaced Pb as a source marker for motor vehicle emissions (e.g., Schauer and Cass, 2000). In their comprehensive review of CMB modeling studies undertaken since 1995, Chow and Watson (2002) note that in twenty-two studies fossil fuel combustion was found to be a large contributor to $PM_{2.5}$ and PM_{10}

concentrations, with most of the contributions to primary PM originating from the exhaust of diesel and gasoline vehicles.

Marine aerosol is found, as expected, at coastal sites such as Long Beach (average 3.8% of total mass) and San Nicolas Island (25%). These contributions to PM_{10} are relatively variable and are larger at the more remote sites. Individual values reflect proximity to local sources. Of great importance are the contributions from secondary ammonium sulfate in the eastern United States and ammonium nitrate in the western United States. Secondary ammonium sulfate is especially noticeable at sites in California's San Joaquin Valley (Bakersfield, Crows Landing, Fellows, Fresno, and Stockton) and in the Los Angeles area.

Because many source apportionment studies address problems in compliance with the National Ambient Air Quality Standards and other air quality standards, samples selected for chemical analysis are often biased toward the highest PM_{10} mass concentrations in the studies shown in Table 3-10. Thus, the average source contribution estimates shown in Table 3-10 are probably not representative of annual averages. For example, the study by Motallebi (1999) considered only days when the PM_{10} concentration was greater than 40 $\mu g/m^3$. Quoted uncertainties in the estimated contributions of the individual sources shown in Tables 3-9 and 3-10 range from 10 to 50%. Errors can be much higher when the chemical source profiles for different sources are highly uncertain or are too similar to distinguish one source from another.

Very few source apportionment studies using the CMB modeling technique have examined the spatial variability of source contributions at different sites within an urban area. As can be seen from Table 3-9, Dzubay et al. (1988) found a uniform distribution of sulfate among the NE Airport in Philadelphia, PA; downtown Camden, NJ; and Clarksboro, NJ, during the summer of 1982. The farthest distance between two monitoring sites (NE Airport and Clarksboro) was approximately 40 km. Magliano et al. (1998) examined the spatial variability of PM_{10} source contributions at a number of sites in Fresno and Bakersfield, CA, during the winter of 1995-1996 and reported values for 1 day, December 27, 1995. During that day, mobile sources contributed from 13.0 to15.8 $\mu g/m^3$, vegetation burning contributed from 5.1 to 11.1 $\mu g/m^3$, ammonium sulfate contributed 2.4 to 3.4 $\mu g/m^3$, and ammonium nitrate contributed 19.3 to 24.6 $\mu g/m^3$ to PM_{10} at the sites in Bakersfield. Mobile sources contributed 13.9 to 22.5 $\mu g/m^3$, vegetation burning contributed 8.2 to 15.7 $\mu g/m^3$, ammonium sulfate contributed 1.8 to 2.3 $\mu g/m^3$, and ammonium nitrate contributed 14.5 to 18.9 $\mu g/m^3$ at the sites in Fresno. All of these components

are expected to be found mainly in the $PM_{2.5}$ size fraction. As can be seen, source contributions at different sites varied by factors of 1.2 to 2.2 in Bakersfield and by factors of 1.3 to 1.9 in Fresno on that day.

The receptor modeling methods outlined above do not explicitly include consideration of the distances between PM sources and the receptor site. Information about the relative importance of sources as a function of distance may be available from examination of data obtained by continuous monitoring methods. For example, concentration spikes are expected to be the result of transport from nearby sources, because turbulent mixing in the atmosphere would not allow them to persist for very long. Short duration spikes in the time series of concentrations are assumed to result from emissions from local sources (0.1 to 1 km away) in this method. Contributions from sources located further away are determined by comparisons between baselines measured at different sites. Details such as these are also lost in integrated 24-h samples. Watson and Chow (2001) used time series of black carbon (BC) obtained by aetholometers over five minute intervals to estimate the contributions from sources located < 1 km away, 1 to 5 km away, and > 5 km away from a monitoring site in downtown Mexico City. They found that most of the BC was produced by sources scattered throughout the city and that sources located less than 1 km away from the site contributed only about 10% to BC concentrations even in the presence of local sources such as buses and trucks.

3.3.4 Emissions Estimates for Primary Particulate Matter, and Precursors to Secondary Particulate Matter (SO_2, NO_x, VOCs, and NH_3) in the United States

In principle, source contributions to ambient PM also could be estimated on the basis of predictions made by chemistry-transport models (CTM) or even on the basis of emissions inventories alone. Uncertainties in emissions inventories have arguably been regarded as representing the largest source of uncertainty in CTMs (Calvert et al., 1993). Apart from uncertainties in emission inventories, a number of other factors limit the ability of an emissions-inventory driven CTM to determine the effects of various sources on particle samples obtained at a particular location. CTM predictions represent averages over the area of a grid cell, which in the case of CMAQ (Community Model for Air Quality) and MAQSIP (Multiscale Air Quality Simulation Platform), ranges from 16 km^2 (4 km × 4 km) to 1296 km^2 (36 km × 36 km). CMAQ

and MAQSIP constitute the CTMs within the overall Models3 framework, which also includes emissions processors, the meteorological model, and modules for decision support. The contributions of sources to pollutant concentrations at a monitoring site are controlled strongly by local conditions that cannot be resolved by an Eulerian grid-cell model. Examples would be the downward mixing of tall stack emissions and deviations from the mean flow caused by buildings. The effect of local sources at a particular point in the model domain may not be predicted accurately because their emissions would be smeared over the area of a grid cell or if the local wind fields at the sampling point deviated significantly from the mean wind fields calculated by the model. CTMs also have problems in predicting pollutant concentrations because of uncertainties in vertical mixing and in predicting concentrations of pollutants from stationary combustion sources resulting from uncertainties in estimates of plume rise. CTMs are an integral part of air quality management programs and are reviewed in the NARSTO Fine Particle Assessment (NARSTO, 2002).

Estimated emissions of primary $PM_{2.5}$ from different sources in the United States are summarized in Table 3-11, and estimated emissions of precursors to the formation of secondary $PM_{2.5}$ (SO_2, NO_x, VOCs, and NH_3) are summarized in Table 3-12. These estimates are given to provide a rough overview of the relative importance of major PM sources in the United States. The emissions estimates are based on information presented in the EPA National Air Pollutant Emission Trends Report, 1900-1999 (U.S. Environmental Protection Agency, 2001), to which the reader is referred for detailed tables showing trends in $PM_{2.5}$ emissions from a number of source categories from 1990 to 1999. Detailed descriptions of the methodology for constructing emissions inventories for criteria pollutants, quality assurance procedures, and examples of calculations of emissions can be found in U. S. Environmental Protection Agency (1999). Although uncertainties associated with the estimates in the National Air Pollutant Emission Trends Report are not given therein, a discussion of uncertainties in emissions estimates is given in Section 3.3.5.

For the sake of completeness, an attempt was made to supplement the information given in the emissions tables in the Trends Report, which concentrates mainly on anthropogenic emissions, with information about emissions from natural sources. Details regarding the composition of the emissions of primary $PM_{2.5}$ from the source categories shown in Table 3-11 are summarized in Appendix 3D, where available. Fugitive dust emissions are estimated to

TABLE 3-11. EMISSIONS OF PRIMARY $PM_{2.5}$ BY VARIOUS SOURCES IN 1999

Source	Emissions (10^9 kg/y)	Major PM Components	Notes
On-road vehicle exhaust	0.21	Organic compounds, elemental carbon	Exhaust emissions from diesel (72%) and gasoline vehicles (28%).
Non-road vehicle exhaust	0.37	Organic compounds, elemental carbon	Exhaust emissions from off-road diesel (57%) and gasoline vehicles (20%); ships and boats (10%); aircraft (7%); railroads (6%).
Fossil fuel combustion	0.36	Crustal elements, trace metals	Fuel burning in stationary sources such as power plants (33%); industries (39%); businesses and institutions (25%); residences (3%).
Industrial processes	0.35	Metals, crustal material, organic compounds	Metals processing (29%); mineral products (27%); chemical mfg. (11%); other industries (33%).
Biomass burning	1.2	Organic compounds, elemental carbon	Managed burning (47%); residential wood burning (28%); agricultural burning (7%); wildfires (18%).
Waste disposal	0.48	Organic compounds, trace metals	Open burning (91%); incineration (9%).
Fugitive dust	3.3	Crustal elements	Dust raised by vehicles on paved (19%) and unpaved roads (40%); construction (15%), dust from raising crops (24%) and livestock (2%).
Windblown dust	NA[1]	Crustal elements	Dust raised by wind on bare land.
Other	0.02	Organic compounds, elemental carbon	Structural fires
Total	6.2		

[1]NA = not available.

Source: Adapted from U. S. Environmental Protection Agency (2001).

constitute over 50 percent of nationwide primary $PM_{2.5}$ emissions, according to Table 3-11. However, there are a number of issues concerning the methods for obtaining relevant emissions factor data for fugitive dust in field studies, as discussed in Section 3.3.5. An estimate of the production of $PM_{2.5}$ from wind erosion on natural surfaces was not included in Table 3-11 because this source is highly sporadic, occurs during periods of high winds and, thus, the resulting emissions are too highly uncertain to be included. As can be seen from a comparison of entries in Tables 3-11 and 3-12, estimates of emissions of potential precursors to secondary PM formation are considerably larger than those for estimates of primary $PM_{2.5}$ emissions in the

TABLE 3-12. EMISSIONS OF PRECURSORS TO SECONDARY $PM_{2.5}$ FORMATION BY VARIOUS SOURCES IN 1999

Precursor	Emissions (10^9 kg/y)	Secondary PM Component	Notes
SO_2	17	Sulfate	Exhaust from on-road (2%) and non-road (5%) engines and vehicles; fossil fuel combustion by electrical utilities, industries, other sources (85%); various industrial processes (7%); and other minor sources (1%).
NO_x[1,2]	26	Nitrate	Exhaust from on-road (34%) and non-road (22%) engines and vehicles; fossil fuel combustion by electrical utilities, industries, other sources (39%); lightning (4%); soils (4%); and other minor sources (5%).
Anthropogenic VOCs	16	Various mainly unidentified compounds of 'OC'	Evaporative and exhaust emissions from on-road (29%) and non-road (18%) vehicles; evaporation of solvents and surface coatings (27%); biomass burning (9%); storage and transport of petroleum and volatile compounds (7%); chemical and petroleum industrial processes (5%); other sources (5%).
Biogenic VOCs[1]	44	Various mainly unidentified compounds of 'OC'	Approximately 98% emitted by vegetation. Isoprene (35%); monoterpenes (25%); all other reactive and non-reactive compounds (40%).
NH_3	45	Ammonium	Exhaust from on-road and non-road engines and vehicles (5%); chemical manufacturing (3%); waste disposal, recycling, and other minor sources (5%); livestock (82%); and fertilizer application (18%).

[1]Includes estimates of natural sources from Guenther et al. (2000).
[2]Emissions expressed in terms of NO_2.

Source: Adapted from U. S. Environmental Protection Agency (2001).

United States. The emissions of SO_2, NO_x, and NH_3 should be multiplied by factors of 1.5, 1.35, and 1.07, respectively, to account for their chemical form in the aerosol phase. Estimating a factor for VOCs is somewhat less straight forward. Turpin and Lim (2001) recommends a factor of 2 to account for the conversion of VOC precursors to oxygen and nitrogen containing compounds in the aerosol phase. These factors are all greater than 1 and further underscore the potential importance of secondary PM precursor emissions relative to primary PM emissions. However, the emissions of precursors cannot be translated directly into rates of PM formation. Dry deposition and precipitation scavenging of some of these gaseous precursors and their intermediate oxidation products occur before they are converted to PM in the atmosphere.

In addition, some fraction of these gases are transported outside of the domain of the continental United States before being oxidized. Likewise, emissions of these gases from areas outside the United States can result in the transport of their oxidation products into the United States.

As discussed in Section 3.3.1, the photochemical oxidation of sulfur dioxide leads to the production of sulfate; whereas that of nitrogen oxide leads ultimately to particulate-phase nitrite and nitrate. Due to uncertainties it is difficult to calculate the rates of formation of secondary organic particulate matter (SOPM) from the emissions of VOC precursors. Smog chamber and laboratory studies discussed in Section 3.3.1 indicate that anthropogenic aromatic compounds and biogenic terpenoid compounds have the highest potential for forming secondary organic particulate matter; and as can be seen from Table 3C-1, the dominant compounds tend to be those derived from these categories. Each of the source categories capable of emitting VOCs shown in Table 3-12 has components capable of forming SOPM, although in small yields (ranging typically up to several per cent, cf. Section 3.3.1). The oxidation of lighter organic compounds leads ultimately to the formation of CO and CO_2. As discussed by Pandis et al. (1991) and in Section 3.3.1, soluble gas phase compounds, such as formaldehyde (CH_2O), other aldehydes, organic acids, etc. formed during the oxidation of a wide variety of hydrocarbons, can be incorporated into suspended particles. Although isoprene is a major component of biogenic emissions, its oxidation has not been found to result in the formation of new particles; whereas the oxidation of monoterpenes has. However, it should be remembered that soluble gas phase species such as CH_2O are formed during the oxidation of isoprene.

The emissions estimates shown in this section are based on annual totals. However, annual averages do not reflect the variability of a number of emissions categories on shorter time scales. Residential wood burning in fireplaces and stoves, for example, is a seasonal practice that reaches its peak during cold weather. Cold weather also affects motor vehicle exhaust particulate matter emissions, both in terms of chemical composition and emission rates (e.g., Watson et al., 1990b; Huang et al., 1994). Agricultural activities such as planting, fertilizing, and harvesting are also seasonal. Forest fires occur mainly during the local dry season and during periods of drought. Maximum dust production by wind erosion in the United States occurs during the spring; whereas the minimum occurs during the summer (Gillette and Hanson, 1989). Efforts are being made to account for the seasonal variations of emissions in the nationwide emissions

inventories. Techniques for calculating emissions of criteria pollutants on a seasonal basis are given in U. S. Environmental Protection Agency (1999).

Trends in nationwide, annual average concentrations of PM_{10}, and precursor gases (SO_2, NO_2, and VOC) over the 10 years from 1989 to 1998 are shown in Table 3-13. As can be seen from Table 3-13, there have been substantial decreases in the ambient concentrations of PM_{10}, SO_2, and NO_2. Not enough data are available to define trends in concentrations of VOCs. There also have been substantial decreases in the emissions of all the species shown in Table 3-13, except for NO_2, although its average ambient concentration has decreased by 14%. These entries suggest that decreases in the average ambient concentration of PM_{10} could have been produced by both decreases in emissions of primary PM_{10} and the formation of secondary PM_{10}. The large reductions in ambient SO_2 concentrations have resulted in reductions in sulfate formation that would have been manifest in $PM_{2.5}$ concentrations on the regional scale in the East and Midwest, where sulfate has constituted a larger fraction of $PM_{2.5}$ than in the West. Likewise, reductions in NO_2 concentrations would have had a more noticeable effect on $PM_{2.5}$ concentrations in the West than in the East, because nitrate is a larger component of the aerosol in the West.

TABLE 3-13. NATIONWIDE CHANGES IN AMBIENT CONCENTRATIONS AND EMISSIONS OF PM_{10} AND GASEOUS PRECURSORS TO SECONDARY PARTICULATE MATTER FROM 1990 TO 1999

	% Change 1990-1999	
	Ambient Concentration	Emissions
PM_{10}	−18%	−15%
$PM_{2.5}$ (1992 to 1999)	Urban east −2% Rural east −5 Rural west −15%	−17%
$SO_4^{=}/SO_2$	−36% (sulfate)	−20% (SO_2)
NO_3^{-}/NO_x	−10% (nitrate)	+5% (NO_x)
VOC	—	−14%

Source: U. S. Environmental Protection Agency (2000d).

Trends in aerosol components (i.e., nitrate, sulfate, carbon, etc.) are needed for a more quantitative assessment of the effects of changes in emissions of precursors. Aerosol nitrate and sulfate concentrations obtained at North Long Beach and Riverside, CA, tracked downward trends in NO_x concentrations. SO_2 and sulfate concentrations have both decreased; however, the rate of decline of sulfate has been smaller than that of SO_2, indicating the long range transport of sulfate from outside the air shed may be an important source in addition to the oxidation of locally generated SO_2. There are a number of reasons why pollutant concentrations do not track estimated reductions in emissions. Some of these reasons are related to atmospheric effects such as meteorological variability and secular changes in the rates of photochemical transformations and deposition (U.S. Environmental Protection Agency, 2000c). Other reasons are related to uncertainties in ambient measurements and in emissions inventories.

3.3.5 Uncertainties of Emissions Inventories

As described in the 1996 PM AQCD, it is difficult to assign uncertainties quantitatively to entries in emissions inventories. Methods that can be used to verify or place constraints on emissions inventories are sparse. In general, the overall uncertainty in the emissions of a given pollutant includes contributions from all of the terms used to calculate emissions (i.e., activity rates, emissions factors, and control device efficiencies). Additional uncertainties arise during the compilation of an emissions inventory because of missing sources and computational errors. The variability of emissions can cause errors when annual average emissions are applied to applications involving shorter time scales.

Activity rates for well-defined point sources (e.g., power plants) should have the smallest uncertainty associated with their use because emissions are monitored continuously in many cases accurate production records need to be kept. On the other hand, activity rates for a number of very dispersed fugitive sources are difficult to quantify. Emissions factors for easily measured fuel components that are released quantitatively during combustion (e.g., CO_2, SO_2) should be the most reliable. Emissions of components formed during combustion are more difficult to characterize, as the emissions rates are dependent on factors specific to individual combustion units and on combustion stage (i.e., smoldering or active). Although the AP-42 emissions factors (U.S. Environmental Protection Agency, 1995) contain extensive information for a large number of source types, these data are very limited in the number of sources sampled. The efficiency of

control devices is determined by their design, their age, their maintenance history, and operating conditions. It is virtually impossible to assign uncertainties in control device performance because of these factors. It should be noted that the largest uncertainties occur for those devices that have the highest efficiencies (>90%). This occurs because the efficiencies are subtracted from one, and small errors in assigning efficiencies can lead to large errors in emissions.

Ideally, an emissions inventory should include all major sources of a given pollutant. This may be an easy task for major point sources. However, area sources of both primary PM and precursors to secondary PM formation are more difficult to characterize than point sources; and, thus, they require special emphasis when preparing emission inventories. Further research is needed to better characterize the sources of pollutants to reduce this source of uncertainty. Errors also can arise from the misreporting of data, and arithmetic errors can occur in the course of compiling entries from thousands of individual sources. A quality assurance program is required to check for outliers and arithmetic errors.

Because of the variability in emissions rates, there can be errors in the application of inventories developed on an annually averaged basis (as are the inventories shown in Tables 3-11 and 3-12) to episodes occurring on much shorter time scales. As an example, most modeling studies of air pollution episodes are carried out for periods of a few days.

Uncertainties in annual emissions were estimated to range from 4 to 9% for SO_2 and from 6 to 11% for NO_x in the 1985 NAPAP inventories for the United States (Placet et al., 1991). Uncertainties in these estimates increase as the emissions are disaggregated both spatially and temporally. The uncertainties quoted above are minimum estimates and refer only to random variability about the mean assuming that the variability in emissions factors was adequately characterized and that extrapolation of emissions factors to sources other than those for which they were measured is valid. The estimates do not consider the effects of weather or variations in operating and maintenance procedures.

Fugitive dust sources, as mentioned above, are extremely difficult to quantify; and stated emission rates may represent only order-of-magnitude estimates. Although crustal dust emissions constitute about 50% of the total primary $PM_{2.5}$ inventory, they constitute less than about 15% of the source strengths inferred from the receptor modeling studies shown in Table 3-9. However, it should be remembered that secondary components (sulfate, nitrate, and some fractions of organic carbon) often account for most of the mass of ambient $PM_{2.5}$ samples.

Although mineral dust sources represent the major category in Table 3-11, their contributions are distributed much more widely than are those from combustion sources. Watson and Chow (1999) reexamined the methodology used to determine emissions of fugitive dust. The standard methods use data obtained by particle monitors stacked at several elevations from 1 to 2 m up to 7 to 10 m above the surface. However, small-scale turbulent motions and variable winds characterize atmospheric flow patterns immediately adjacent to the surface (Garratt, 1994). The depth of this turbulent layer is determined by surface roughness elements; and, if particle monitors are sampling within this layer, there is a high probability of particles being entrained in turbulent eddies and redepositing on the ground within a very short distance. In addition to the source sampling problem referred to above, it should be remembered that dust often is raised in remote areas far removed from population centers. Precipitation or scavenging by cloud droplets and dry deposition removes particles during transport from the source area. In addition, gravitational settling can be an important loss mechanism for particles larger than a few micrometers in aerodynamic diameter.

As rough estimates, uncertainties in emissions estimates could be as low as 10% for the best characterized source categories; whereas emissions figures for windblown dust should be regarded as order-of-magnitude estimates. The application of emissions inventories to the estimation of source contributions at monitoring sites is also limited by the effects of local topography and meteorology. For example, Pinto et al. (1998) found that the contribution of power plants and residential space heating to $PM_{2.5}$ concentrations in northwestern Bohemia are comparable on the basis of CMB receptor modeling. However, according to the emissions inventories, the contribution from power plants should have been roughly an order of magnitude larger than that from residential space heating. The difference between the two methods can be explained by noting that mixing of the emissions from the power plants downward to the surface is inhibited by strong surface inversions that develop during the winter season in this area.

There have been few field studies designed to test emissions inventories observationally. The most direct approach would be to use aircraft to obtain cross-sections of pollutants upwind and downwind of major urban areas. The computed mass flux through a cross section of the urban plume can then be equated to emissions from the city chosen. This approach has been attempted on a few occasions, but results have been ambiguous because of contributions from fugitive sources, variable wind flows, and logistic difficulties.

3.4 SUMMARY AND CONCLUSIONS

The recently deployed $PM_{2.5}$ FRM network has returned data for a large number of sites across the United States. Annual mean $PM_{2.5}$ concentrations range from about 5 $\mu g/m^3$ to about 30 $\mu g/m^3$. In the eastern United States, the data from 1999 and 2000 indicate that highest quarterly mean concentrations and maximum concentrations occur during the summer. In the western United States, highest quarterly mean values and maximum values occur mainly during the winter at a number of sites, although there were exceptions to these general patterns. These findings are generally consistent with those based on longer term data sets such as MAAQS in the eastern United States and the CARB network of dichotomous samplers in California. $PM_{2.5}$ and PM_{10} concentrations in a number of urban areas have generally declined over the past few decades. However, they have leveled off in the past few years.

Differences in annual mean $PM_{2.5}$ concentrations between monitoring sites in urban areas examined are typically less than 6 or 7 $\mu g/m^3$. However, on individual days, differences in 24-h average $PM_{2.5}$ concentrations can be much larger. Some sites in metropolitan areas are highly correlated with each other, but other sites are not due to the presence of local sources, topographic barriers, etc. Although $PM_{2.5}$ concentrations at sites within an MSA can be highly correlated, there still can be significant differences in their concentrations on any given day. Consequently, additional measures should be used to characterize the spatial variability of $PM_{2.5}$ concentrations. The degree of spatial uniformity in $PM_{2.5}$ concentrations in urban areas varies across the country. These factors should be considered in using data obtained by the $PM_{2.5}$ FRM network to approximate community-scale human exposure, and caution should be exercised in extrapolating conclusions obtained in one urban area to another. $PM_{2.5}$ to PM_{10} ratios were generally higher in the East than in the West, and values for this ratio are consistent with those found in numerous earlier studies presented in the 1996 PM AQCD.

Data for $PM_{10-2.5}$ are not as abundant as they are for $PM_{2.5}$, and their interpretation is complicated by the difference method used to determine their concentrations. The more sporadic nature of sources of $PM_{10-2.5}$ and its shorter atmospheric lifetime tend to result in lower correlations for $PM_{10-2.5}$ than for $PM_{2.5}$ concentrations. Errors in measurement of $PM_{2.5}$ and PM_{10} also result in lower spatial correlations of $PM_{10-2.5}$. Calculated concentrations of $PM_{10-2.5}$ are occasionally negative as reflected by $PM_{2.5}$ to PM_{10} ratios greater than one. Because analytical errors are generally larger for individual species than for total mass, similar problems arise in

their determination in $PM_{10-2.5}$ samples by the difference approach. Some, but not all of these problems could be resolved by the use of dichotomous samplers that also provide a direct sample of $PM_{10-2.5}$ for compositional analyses.

Estimates of concentrations of individual species in $PM_{10-2.5}$ samples were limited to those obtained by dichotomous samplers. Generally, concentrations of most elements differ for $PM_{2.5}$ and $PM_{10-2.5}$. However, the available data suggest that concentrations of many metals are of the same order of magnitude in both size fractions. This is in marked contrast to the situation twenty years ago, when uncontrolled combustion sources were prevalent. At that time, concentrations of many metals, especially lead, were much higher than today in fine-mode particles, and their concentrations were much higher in the fine-mode than in the coarse-mode. No substantive conclusions about contemporary concentrations and composition of ultrafine particles ($0.1\ \mu m < D_a$) can be drawn for the nation as a whole, because of a lack of data.

Ambient PM contains both primary and secondary components. The results of ambient monitoring studies and receptor modeling studies indicate that $PM_{2.5}$ is dominated by secondary components in the eastern United States. Depending on the origin of OC in ambient samples, $PM_{2.5}$, on average, may also be dominated by secondary components throughout the rest of the United States. Primary constituents represent smaller but still important components of $PM_{2.5}$. Crustal materials, which are primary constituents, constitute the largest measured fraction of $PM_{10-2.5}$ throughout the United States. Data for the concentration of bioaerosols in both the $PM_{2.5}$ and $PM_{10-2.5}$ size ranges are sparse. Data collected in several airsheds, including the Los Angeles Basin, Bakersfield and Fresno, CA; and Philadelphia, PA, suggest that secondary PM components are more uniformly distributed than are primary PM components. Compositional data obtained at multiple sites in other urban areas are sparse.

Because of the complexity of the composition of ambient $PM_{2.5}$ and $PM_{10-2.5}$, sources are best discussed in terms of individual constituents of both primary and secondary $PM_{2.5}$ and $PM_{10-2.5}$. Each of these constituents can have anthropogenic and natural sources, as shown in Table 3-8. The distinction between natural and anthropogenic sources is not always obvious. Although windblown dust might seem to be the result of natural processes, highest emission rates are associated with agricultural activities in areas that are susceptible to periodic drought. Examples include the dust bowl region of the midwestern United States and the Sahel of Africa. There is also ongoing debate about characterizing wild fires as either natural or anthropogenic.

Land management practices and other human actions affect the occurrence and scope of wild fires. Similarly, prescribed burning can be viewed as anthropogenic, or as a substitute for wild fires that would otherwise occur eventually on the same land.

Over the past decade, a significant amount of research has been carried out to improve the understanding of the atmospheric chemistry of secondary organic PM formation. Although additional sources of SOPM might still be identified, there appears to be a general consensus that biogenic compounds (monoterpenes, sesquiterpenes) and aromatic compounds (toluene, ethylbenzene) are the most significant SOPM precursors. A large number of compounds have been detected in biogenic and aromatic SOPM, although the chemical composition of these two categories has not been fully established, especially for aromatic SOPM. Transformations that occur during the aging of particles are still inadequately understood. There are still large gaps in the current understanding of a number of key processes related to the partitioning of semivolatile compounds between the gas phase and ambient particles containing organic compounds, liquid water, and inorganic salts and acids. In addition, there is a general lack of reliable analytical methods for measuring multifunctional oxygenates in the gas and aerosol phases.

The results of receptor modeling studies throughout the United States indicate that the combustion of fossil and biomass fuels is the major source of measured ambient $PM_{2.5}$. Fugitive dust, found mainly in the $PM_{10\text{-}2.5}$ range size, represents the largest source of measured ambient PM_{10} in many locations in the western United States. Quoted uncertainties in the source apportionment of constituents in ambient aerosol samples typically range from 10 to 50%. It is apparent that a relatively small number of source categories, compared to the total number of chemical species that typically are measured in ambient monitoring-source receptor model studies, are needed to account for the majority of the observed mass of PM in these studies.

As seen in Table 3-8, emissions of mineral dust, organic debris, and sea spray are concentrated mainly in the coarse fraction of PM_{10} (>2.5 μm aerodynamic diameter). A small fraction of this material is in the $PM_{2.5}$ size range (< 2.5 μm aerodynamic diameter). Still, $PM_{2.5}$ concentrations of crustal material can be appreciable, especially during dust events. It also should be remembered that from one-third to one-half of the Saharan dust reaching the United States is in the $PM_{2.5}$ size range. Emissions from combustion sources (mobile and stationary sources and biomass burning) are also predominantly in the $PM_{2.5}$ size range.

Although most emphasis in this chapter has been on sources within the United States, it should be remembered that sources outside the United States contribute to ambient PM levels that can, at times, exceed the ambient NAAQS level for PM. Dust is frequently transported from northern Africa to the eastern United States. This dust often produces dense haze during the summer in southern Florida. Bioaerosols and pollutants are also transported with the dust. Large-scale dust storms in the deserts of central Asia recently have been found to contribute to PM levels in the northwestern United States on an episodic basis. Uncontrolled biomass burning in central America and Mexico may have contributed to elevated PM levels that exceeded the daily NAAQS level for PM in Texas. Wildfires throughout the United States, Canada, Mexico, and Central America all contribute to background concentrations of PM in the United States.

REFERENCES

Altshuller, A. P. (1983) Review: natural volatile organic substances and their effect on air quality in the United States. Atmos. Environ. 17: 2131-2165.

Ansari, A. S.; Pandis, S. N. (1998) Response of inorganic PM to precursor concentrations. Environ. Sci. Technol. 32: 2706-2714.

Atkinson, R. (1994) Gas-phase tropospheric chemistry of organic compounds. Washington, DC: American Chemical Society. (J. Phys. Chem. Ref. Data Monograph no. 2).

Aumont, B.; Madronich, S.; Bey, I.; Tyndall, G. S. (2000) Contribution of secondary VOC to the composition of aqueous atmospheric particles: a modeling approach. J. Atmos. Chem. 35: 59-75.

Bahadori, T.; Suh, H.; Koutrakis, P. (2000a) A cross-study analysis of ambient concentrations of fine mass. J. Air Waste Manage. Assoc.: in press.

Bahadori, T.; Suh, H.; Koutrakis, P. (2000b) $PM_{2.5}$ and PM_{10} concentrations in four metropolitan cities in eastern United States. J. Air Waste Manage. Assoc.: in press.

Cahill, T. A. (1996) [Data from IMPROVE data base.] Fort Collins, CO: Air Resource Specialists, Inc.

California Air Resources Board. (1995) [Data base provided by John R. Holmes, Chief, Research Division.] Sacramento, CA: California Air Resources Board.

Calvert, J. G.; Heywood, J. B.; Sawyer, R. F.; Seinfeld, J. H. (1993) Achieving acceptable air quality: some reflections on controlling vehicle emissions. Science (Washington, DC) 261: 37-45.

Cass, G. R. ; Hughes, L. A.; Bhave, P.; Kleeman, M. J.; Allen, J. O.; Salmon, L. G. (2000) The chemical composition of atmospheric ultrafine particles. Philos. Trans. R. Soc. London Ser. A 358: 2581-2592.

Castro, L. M.; Pio, C. A.; Harrison, R. M.; Smith, D. J. T. (1999) Carbonaceous aerosol in urban and rural European atmospheres: estimation of secondary organic carbon concentrations. Atmos. Environ. 33: 2771-2781.

Chadwick, O. A.; Derry, L. A.; Vitousek, P. M.; Huebert, B. J.; Hedin, L. O. (1999) Changing sources of nutrients during four million years of ecosystem development. Nature (London, UK) 397: 491-497.

Chow, J. C.; Watson, J. G. (2002) Review of $PM_{2.5}$ and PM_{10} apportionment for fossil fuel combustion and other sources by the chemical mass balance receptor model. Energy Fuels: in press.

Chow, J. C.; Watson, J. G.; Frazier, C. A.; Egami, R. T.; Goodrich, A.; Ralph, C. (1988) Spatial and temporal source contributions to PM_{10} and $PM_{2.5}$ in Reno, NV. In: Mathai, C. V.; Stonefield, D. H., eds. PM_{-10}: implementation of standards, an APCA/EPA international specialty conference; February; San Francisco, CA. Pittsburgh, PA: Air Pollution Control Association; pp. 438-457. (APCA transactions series TR-13).

Chow, J. C.; Watson, J. G.; Richards, L. W.; Haase, D. L.; McDade, C.; Dietrich, D. L.; Moon, D.; Sloane, C. (1991) The 1989-90 Phoenix PM_{10} study. Volume II: source apportionment. Final report. Phoenix, AZ: Arizona Department of Environmental Quality; Desert Research Institute document no. 8931.6F1.

Chow, J. C.; Watson, J. G.; Lowenthal, D. H.; Frazier, C. A.; Hinsvark, B. A.; Prichett, L. C.; Neuroth, G. R. (1992a) Wintertime PM_{10} and $PM_{2.5}$ chemical compositions and source contributions in Tucson, Arizona. In: Chow, J. C.; Ono, D. M., eds. PM_{10} standards and nontraditional particulate source controls, an A&WMA/EPA international specialty conference, v. I; January; Scottsdale, AZ. Pittsburgh, PA: Air & Waste Management Association; pp. 231-243. (A&WMA transactions series no. 22).

Chow, J. C.; Watson, J. G.; Lowenthal, D. H.; Solomon, P. A.; Magliano, K. L.; Ziman, S. D.; Richards, L. W. (1992b) PM_{10} source apportionment in California's San Joaquin Valley. Atmos. Environ. Part A 26: 3335-3354.

Chow, J. C.; Liu, C. S.; Cassmassi, J.; Watson, J. G.; Lu, Z.; Pritchett, L. C. (1992c) A neighborhood-scale study of PM_{10} source contributions in Rubidoux, California. Atmos. Environ. Part A 26: 693-706.

Chow, J. C.; Watson, J. G.; Edgerton, S. A.; Vega, E.; Ortiz, E. (2001) Spatial differences in outdoor PM_{10} mass and aerosol composition in Mexico City. J. Air Waste Manage. Assoc.: submitted.

Christoffersen, T. S.; Hjorth, J.; Horie, O.; Jensen, N. R.; Kotzias, D.; Molander, L. L.; Neeb, P.; Ruppert, L.; Winterhalter, R.; Virkkula, A.; Wirtz, K.; Larsen, B. R. (1998) cis-Pinic acid, a possible precursor for organic aerosol formation from ozonolysis of α-pinene. Atmos. Environ. 32: 1657-1661.

Chung, A.; Herner, J. D.; Kleeman, M. J. (2001) Detection of alkaline ultrafine atmospheric particles at Bakersfield, California. Environ. Sci. Technol. 35: 2184-2190.

Cocker, D. R., III; Clegg, S. L.; Flagan, R. C.; Seinfeld, J. H. (2001) The effect of water on gas-particle partitioning of secondary organic aerosol. Part I: α-pinene/ozone system. Atmos. Environ. 35: 6049-6072.

Darlington, T. L.; Kahlbaum, D. F.; Heuss, J. M.; Wolff, G. T. (1997) Analysis of PM_{10} trends in the United States from 1988 through 1995. J. Air Waste Manage. Assoc. 47: 1070-1078.

Davis, A. P.; Shokouhian, M.; Ni, S. (2001) Loading estimates of lead, copper, cadmium, and zinc in urban runoff from specific sources. Chemosphere 44: 997-1009.

Dolislager, L. J.; Motallebi, N. (1999) Characterization of particulate matter in California. J. Air Waste Manage. Assoc. 49: PM-45-56.

Duce, R. A. (1995) Sources, distributions, and fluxes of mineral aerosols and their relationship to climate. In: Charlson, R. J.; Heintzenberg, J., eds. Aerosol forcing of climate: report of the Dahlem workshop on aerosol forcing of climate; April 1994; Berlin, Federal Republic of Germany. Chichester, United Kingdom: John Wiley & Sons, Ltd.; pp. 43-72.

Dzubay, T. G.; Stevens, R. K.; Gordon, G. E.; Olmez, I.; Sheffield, A. E.; Courtney, W. J. (1988) A composite receptor method applied to Philadelphia aerosol. Environ. Sci. Technol. 22: 46-52.

Edney, E. O.; Driscoll, D. J.; Weathers, W. S.; Kleindienst, T. E.; Conver, T. S.; McIver, C. D.; Li, W. (2001) Formation of polyketones in irradiated toluene/propylene/NO_x/air mixtures. Aerosol Sci. Technol. 35: 998-1008.

Eldred, R. A.; Cahill, T. A. (1994) Trends in elemental concentrations of fine particles at remote sites in the United States of America. Atmos. Environ. 28: 1009-1019.

Faust, B. C. (1994) Photochemistry of clouds, fogs, and aerosols. Environ. Sci. Technol. 28: 217A-222A.

Fitz-Simons, T. S.; Mathias, S.; Rizzo, M. (2000) Analyses of 1999 PM data for the PM NAAQS review. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; November 17. Available: http://www.epa.gov/oar/oaqps/pm25/analyses.html [2 April, 2002].

Forstner, H. J. L.; Flagan, R. C.; Seinfeld, J. H. (1997a) Molecular speciation of secondary organic aerosol from photooxidation of the higher alkenes: 1-octene and 1-decene. Atmos. Environ. 31: 1953-1964.

Forstner, H. J. L.; Flagan, R. C.; Seinfeld, J. H. (1997b) Secondary organic aerosol from the photooxidation of aromatic hydrocarbons: molecular composition. Environ. Sci. Technol. 31: 1345-1358.

Garratt, J. R. (1994) The atmospheric boundary layer. Cambridge, United Kingdom: Cambridge University Press. (Cambridge atmospheric and space science series).

Gatz, D. F.; Prospero, J. M. (1996) A large silicon-aluminum aerosol plume in central Illinois: North African Desert dust? Atmos. Environ. 30: 3789-3799.

Gillette, D. A.; Hanson, K. J. (1989) Spatial and temporal variability of dust production caused by wind erosion in the United States. J. Geophys. Res. [Atmos.] 94: 2197-2206.

Gillies, J. A.; Gertler, A. W.; Lowenthal, D.; Abu-Allaban, M.; Jennison, B.; Goodrich, A. (2000) Enhanced $PM_{2.5}$ source apportionment using chemical mass balance receptor modeling and scanning electron microscopy. Aerosol Sci. Technol.: submitted.

Glasius, M.; Lahaniati, M.; Calogirou, A.; Di Bella, D.; Jensen, N. R.; Hjorth, J.; Kotzias, D.; Larsen, B. R. (2000) Carboxylic acids in secondary aerosols from oxidation of cyclic monoterpenes by ozone. Environ. Sci. Technol. 34: 1001-1010.

Glover, D. M.; Hopke, P. K.; Vermette, S. J.; Landsberger, S.; D'Auben, D. R. (1991) Source apportionment with site specific source profiles. J. Air Waste Manage. Assoc. 41: 294-305.

Gone, J.-K.; Olmez, I.; Ames, M. R. (2000) Size distribution and probable sources of trace elements in submicron atmospheric particulate material. J. Radioanal. Nucl. Chem. 244: 133-139.

Graedel, T. E.; Goldberg, K. I. (1983) Kinetic studies of raindrop chemistry: 1. Inorganic and organic processes. J. Geophys. Res. C: Oceans Atmos. 88: 10,865-10,882.

Gray, H. A.; Landry, B.; Liu, C. S.; Henry, R. C.; Cooper, J. A.; Sherman, J. R. (1988) Receptor modeling for PM_{10} source apportionment in the South Coast Air Basin of California. In: Mathai, C. V.; Stonefield, D. H., eds. PM_{-10}: implementation of standards, an APCA/EPA international specialty conference; February; San Francisco, CA. Pittsburgh, PA: Air Pollution Control Association; pp. 399-418. (APCA transactions series TR-13).

Griffin, R. J.; Cocker, D. R., III; Seinfeld, J. H. (1999) Incremental aerosol reactivity: application to aromatic and biogenic hydrocarbons. Environ. Sci. Technol. 33: 2403-2408.

Griffin, D. W.; Garrison, V. H.; Herman, J. R.; Shinn, E. A. (2001) African desert dust in the Caribbean atmosphere: microbiology and public health. Aerobiologia 17: 203-213.

Grosjean, D. (1992) In situ organic aerosol formation during a smog episode: estimated production and chemical functionality. Atmos. Environ. Part A 26: 953-963.

Grosjean, D.; Seinfeld, J. H. (1989) Parameterization of the formation potential of secondary organic aerosols. Atmos. Environ. 23: 1733-1747.

Guenther, A.; Geron, C.; Pierce, T.; Lamb, B.; Harley, P.; Fall, R. (2000) Natural emissions of non-methane volatile organic compounds, carbon monoxide, and oxides of nitrogen from North America. Atmos. Environ. 34: 2205-2230.

Haagen-Smit, A. J. (1952) Chemistry and physiology of Los Angeles smog. Ind. Eng. Chem. 44: 1342-1346.

Henry, R. C. (1997) History and fundamentals of multivariate air quality receptor models. Chemom. Intell. Lab. Syst. 37: 37-42.

Hewitt, C. N. (2001) The atmospheric chemistry of sulphur and nitrogen in power station plumes. Atmos. Environ. 35: 1155-1170.

Hidy, G. M. (1994) Atmospheric sulfur and nitrogen oxides: eastern North American source-receptor relationships. San Diego, CA: Academic Press, Inc.

Hildemann, L. M.; Rogge, W. F.; Cass, G. R.; Mazurek, M. A.; Simoneit, B. R. T. (1996) Contribution of primary aerosol emissions from vegetation-derived sources to fine particle concentrations in Los Angeles. J. Geophys. Res. [Atmos.] 101: 19,541-19,549.

Hoffmann, T.; Odum, J. R.; Bowman, F.; Collins, D.; Klockow, D.; Flagan, R. C.; Seinfeld, J. H. (1997) Formation of organic aerosols from the oxidation of biogenic hydrocarbons. J. Atmos. Chem. 26: 189-222.

Hopke, P. K.; Wlaschin, W.; Landsberger, S.; Sweet, C.; Vermette, S. J. (1988) The source apportionment of PM_{10} in south Chicago. In: Mathai, C. V.; Stonefield, D. H., eds. PM_{-10}: implementation of standards, an APCA/EPA international specialty conference; February; San Francisco, CA. Pittsburgh, PA: Air Pollution Control Association; pp. 484-494. (APCA transactions series TR-13).

Houck, J. E.; Rau, J. A.; Body, S.; Chow, J. C. (1992) Source apportionment: Pocatello, Idaho PM_{10} nonattainment area. In: Chow, J. C.; Ono, D. M., eds. PM_{10} standards and nontraditional particulate source controls, an A&WMA/EPA international specialty conference, v. I; January; Scottsdale, AZ. Pittsburgh, PA: Air & Waste Management Association; pp. 219-230. (A&WMA transactions series no. 22).

Huang, X.; Olmez, I.; Aras, N. K.; Gordon, G. E. (1994) Emissions of trace elements from motor vehicles: potential marker elements and source composition profile. Atmos. Environ. 28: 1385-1391.

Husar, R. B.; Schichtel, B. A.; Falke, S. R.; Li, F.; Wilson, W. E.; Pinto, J.; Malm, W. C.; Fox, D. G.; Feldman, G. C.; McClain, C.; Kuring, N.; Holben, B. N.; Vermote, E. F.; Herman, J. R.; Elvidge, C. D. (2000) The impact of the 1998 Central American smoke on the atmospheric environment of eastern North America. J. Geophys. Res.: submitted.

Inhalable Particulate Network. (1985) [Inhalable Particulate Network Database.] Research Triangle Park, NC: U.S. Environmental Protection Agency, National Exposure Research Laboratory; Jose M. Sune, contact.

Izumi, K.; Fukuyama, T. (1990) Photochemical aerosol formation from aromatic hydrocarbons in the presence of NO_x. Atmos. Environ. Part A 24: 1433-1441.

Jang, M.; Kamens, R. M. (1999) Newly characterized products and composition of secondary aerosols from the reaction of α-pinene with ozone. Atmos. Environ. 33: 459-474.

Jang, M.; Kamens, R. M. (2001a) Atmospheric secondary aerosol formation by heterogeneous reactions of aldehydes in the presence of a sulfuric acid aerosol catalyst. Environ. Sci. Technol. 35: 4758-4766.

Jang, M.; Kamens, R. M. (2001b) Characterization of secondary aerosol from the photooxidation of toluene in the presence of NO_x and 1-propene. Environ. Sci. Technol. 35: 3626-3639.

Kamens, R.; Jang, M.; Chien, C.-J.; Leach, K. (1999) Aerosol formation from the reaction of α-pinene and ozone using a gas-phase kinetics-aerosol partitioning model. Environ. Sci. Technol. 33: 1430-1438.

Kao, A. S.; Friedlander, S. K. (1995) Frequency distributions of PM_{10} chemical components and their sources. Environ. Sci. Technol. 29: 19-28.

Kavouras, I. G.; Mihalopoulos, N.; Stephanou, E. G. (1998) Formation of atmospheric particles from organic acids produced by forests. Nature (London) 395: 683-686.

Kavouras, I. G.; Mihalopoulus, N.; Stephanou, E. G. (1999a) Formation and gas/particle partitioning of monoterpene photooxidation products over forests. Geophys. Res. Lett. 26: 55-58.

Kavouras, I. G.; Mihalopoulus, N.; Stephanou, E. G. (1999b) Secondary organic aerosol formation vs primary organic aerosol emission: in situ evidence for the chemical coupling between monoterpene acidic photooxidation products and new particle formation over forests. Environ. Sci. Technol. 33: 1028-1037.

Keywood, M. D.; Ayers, G. P.; Gras, J. L.; Gillett, R. W.; Cohen, D. D. (1999) Relationships between size segregated mass concentration data and ultrafine particle number concentrations in urban areas. Atmos. Environ. 33: 2907-2913.

Khalil, M. A. K.; Rasmussen, R. A. (1992) Forest hydrocarbon emissions: relationships between fluxes and ambient concentrations. J. Air Waste Manage. Assoc. 42: 810-813.

Kim, B. M.; Zeldin, M. D.; Liu, C. S. (1992) Source apportionment study for state implementation plan development in the Coachella Valley. In: Chow, J. C.; Ono, D. M., eds. PM_{10} standards and nontraditional particulate source controls: an A&WMA/EPA international specialty conference, v. II; January; Scottsdale, AZ. Pittsburgh, PA: Air & Waste Management Association; pp. 979-991. (A&WMA transactions series no. 22).

Kinney, P. L.; Aggarwal, M.; Northridge, M. E.; Janssen, N. A. H.; Shepard, P. (2000) Airborne concentrations of PM2.5 and diesel exhaust particles on Harlem sidewalks: a community-based pilot study. Environ. Health Perspect. 108: 213-218.

Koch, S.; Winterhalter, R.; Uherek, E.; Kolloff, A.; Neeb, P.; Moortgat, G. K. (2000) Formation of new particles in the gas-phase ozonolysis of monoterpenes. Atmos. Environ. 34: 4031-4042.

Koutrakis, P. (1995) [$PM_{2.5}$ and PM_{10} data for Philadelphia, 1992-1995]. Boston, MA: Harvard School of Public Health.

Landis, M. S.; Norris, G. A.; Williams, R. W.; Weinstein, J. P. (2001) Personal exposures to $PM_{2.5}$ mass and trace elements in Baltimore, MD, USA. Atmos. Environ. 35: 6511-6524.

Larsen, B. R.; Di Bella, D.; Glasius, M.; Winterhalter, R.; Jensen, N. R.; Hjorth, J. (2001) Gas-phase OH oxidation of monoterpenes: gaseous and particulate products. J. Atmos. Chem. 38: 231-276.

Leach, K. B.; Kamens, R. M.; Strommen, M. R.; Jang, M. (1999) Partitioning of semivolatile organic compounds in the presence of a secondary organic aerosol in a controlled atmosphere. J. Atmos. Chem. 33: 241-264.

Leaderer, B. P.; Naeher, L.; Jankun, T.; Balenger, K.; Holford, T. R.; Toth, C.; Sullivan, J.; Wolfson, J. M.; Koutrakis, P. (1999) Indoor, outdoor, and regional summer and winter concentrations of PM_{10}, $PM_{2.5}$, SO_4^{2-}, H^+, NH_4^+, NO_3^-, NH_3, and nitrous acid in homes with and without kerosene space heaters. Environ. Health Perspect. 107: 223-231.

Lin, J. J.; Noll, K. E.; Holsen, T. M. (1994) Dry deposition velocities as a function of particle size in the ambient atmosphere. Aerosol Sci. Technol. 20: 239-252.

Lipfert, F. W. (1998) Trends in airborne particulate matter in the United States. Appl. Occup. Environ. Hyg. 13: 370-384.

Magliano, K. L.; Ranzieri, A.; Solomon, P.; Watson, J. (1998) Chemical mass balance modeling of data from the 1995 integrated monitoring study. Sacramento, CA: California Air Resources Board; California Regional PM10/PM2.5 Air Quality Study.

Mazurek, M.; Masonjones, M. C.; Masonjones, H. D.; Salmon, L. G.; Cass, G. R.; Hallock, K. A.; Leach, M. (1997) Visibility-reducing organic aerosols in the vicinity of Grand Canyon National Park: properties observed by high resolution gas chromatography. J. Geophys. Res. [Atmos.] 102: 3779-3793.

McMurry, P. H.; Woo, K. S.; Weber, R.; Chen, D.-R.; Pui, D. Y. H. (2000) Size distributions of 3 to 10 nm atmospheric particles: implications for nucleation mechanisms. Philos. Trans. R. Soc. London Ser. A 358: 2625-2642.

Monn, C. (2001) Exposure assessment of air pollutants: a review on spatial heterogeneity and indoor/outdoor/personal exposure to suspended particulate matter, nitrogen dioxide and ozone. Atmos. Environ. 35: 1-32.

Motallebi, N. (1999) Wintertime $PM_{2.5}$ and PM_{10} source apportionment at Sacramento, California. J. Air Waste Manage. Assoc. 49: PM-25-34.

NARSTO. (2002) Fine particle assessment (2nd review draft). Available: http://www.cgenv.com/Narsto/ (13 February 2002).

Neas, L. M. (1996) [Data provided for 1986 from Harvard six cities data base.] Boston, MA: Harvard School of Public Health.

Norris, G. (1998) Air pollution and the exacerbation of asthma in an arid, western, US city [dissertation]. Seattle, WA: University of Washington. Available from: University Microfilms International, Ann Arbor, MI; publication no. 9836228.

Norris, G.; Young-Pong, S. N.; Koenig, J. Q.; Larson, T. V.; Sheppard, L.; Stout, J. W. (1999) An association between fine particles and asthma emergency department visits for children in Seattle. Environ. Health Perspect. 107: 489-493.

Odum, J. R.; Hoffmann, T.; Bowman, F.; Collins, D.; Flagan, R. C.; Seinfeld, J. H. (1996) Gas/particle partitioning and secondary organic aerosol yields. Environ. Sci. Technol. 30: 2580-2585.

Odum, J. R.; Jungkamp, T. P. W.; Griffin, R. J.; Flagan, R. C.; Seinfeld, J. H. (1997a) The atmospheric aerosol-forming potential of whole gasoline vapor. Science (Washington, DC) 276: 96-99.

Odum, J. R.; Jungkamp, T. P. W.; Griffin, R. J.; Forstner, H. J. L.; Flagan, R. C.; Seinfeld, J. H. (1997b) Aromatics, reformulated gasoline, and atmospheric organic aerosol formation. Environ. Sci. Technol. 31: 1890-1897.

Ondov, J. M. (1996) Particulate tracers for source attribution: potential for application to California's San Joaquin Valley. J. Aerosol Sci. 27(suppl. 1): S687-S688.

Ott, S.-T.; Ott, A.; Martin, D. W.; Young, J. A. (1991) Analysis of a trans-Atlantic Saharan dust outbreak based on satellite and GATE data. Mon. Weather Rev. 119: 1832-1850.

Paatero, P.; Tapper, U. (1993) Analysis of different modes of factor analysis as least squares fit problems. Chemom. Intelligent Lab. Syst. 18: 183-194.

Paatero, P.; Tapper, U. (1994) Positive matrix factorization: a non-negative factor model with optimal utilization of error estimates of data values. Environmetrics 5: 111-126.

Pandis, S. N.; Paulson, S. E.; Seinfeld, J. H.; Flagan, R. C. (1991) Aerosol formation in the photooxidation of isoprene and β-pinene. Atmos. Environ. Part A 25: 997-1008.

Pandis, S. N.; Harley, R. A.; Cass, G. R.; Seinfeld, J. H. (1992) Secondary organic aerosol formation and transport. Atmos. Environ. Part A 26: 2269-2282.

Perry, K. D.; Cahill, T. A.; Eldred, R. A.; Dutcher, D. D.; Gill, T. E. (1997) Long-range transport of North African dust to the eastern United States. J. Geophys. Res. [Atmos] 102: 11,225-11,238.

Pinto, J. P.; Pleasant, M.; Kellogg, R.; Torok, Sz.; Stiles, D.; Willis, R. D.; Koutrakis, P.; Allen, G. A. (1995) Chemical characterization of the ambient aerosol in Philadelphia. Presented at: Particulate matter: health and regulatory issues: an A&WMA international specialty conference; April; Pittsburgh, PA. Pittsburgh, PA: Air & Waste Management Association; submitted abstract.

Pinto, J. P.; Stevens, R. K.; Willis, R. D.; Kellogg, R.; Mamane, Y.; Novak, J.; Šantroch, J.; Beneš, I.; Leniček, J.; Bureš, V. (1998) Czech air quality monitoring and receptor modeling study. Environ. Sci. Technol. 32: 843-854.

Pinto, J. P.; Lefohn, A. S.; Shadwick, D. S. (2002) Aspects of the spatial variability of $PM_{2.5}$ concentrations within urban areas of the United States. Environ. Sci. Technol.: submitted.

Pio, C.; Alves, C.; Duarte, A. (2001) Organic components of aerosols in a forested area of central Greece. Atmos. Environ. 35: 389-401.

Placet, M.; Battye, R. E.; Fehsenfeld, F. C.; Bassett, G. W. (1991) Emissions involved in acidic deposition processes. In: Irving, P. M., ed. Acidic deposition: state of science and technology, volume I: emissions, atmospheric processes, and deposition. Washington, DC: The U.S. National Acid Precipitation Assessment Program. (State of science and technology report no. 1).

Poore, M. W. (2000) Oxalic acid in $PM_{2.5}$ particulate matter in California. J. Air Waste Manage. Assoc. 50: 1874-1875.

Prospero, J. M. (1996) The atmospheric transport of particles to the ocean. In: Ittekkot, V.; Honjo, S.; Depetris, P. J., eds. Particle flux in the ocean. New York, NY: John Wiley & Sons, Ltd.; pp. 19-52. (SCOPE report no. 57).

Prospero, J. M. (1999) Long-range transport of mineral dust in the global atmosphere: impact of African dust on the environment of the southeastern United States. Proc. Natl. Acad. Sci. U.S.A. 96: 3396-3403.

Prospero, J. M.; Nees, R. T.; Uematsu, M. (1987) Deposition rate of particulate and dissolved aluminum derived from Saharan dust in precipitation at Miami, Florida. J. Geophys. Res. [Atmos.] 92: 14,723-14,731.

Prospero, J. M.; Olmez, I.; Ames, M. (2001) Al and Fe in PM2.5 and PM10 suspended particles in south-central Florida: the impact of the long range transport of African mineral dust. J. Water Air Soil Pollut. 125: 291-317.

Pun, B. K.; Seigneur, C.; Grosjean, D.; Saxena, P. (2000) Gas-phase formation of water-soluble organic compounds in the atmosphere: a retrosynthetic analysis. J. Atmos. Chem. 35: 199-223.

Ramadan, Z.; Song, X.-H.; Hopke, P. K. (2000) Identification of sources of Phoenix aerosol by positive matrix factorization. J. Air Waste Manage. Assoc. 50: 1308-1320.

Rizzo, M.; Pinto, J. P. (2001) Initial characterization of fine particulate matter (PM2.5) collected by the National Federal Reference Monitoring Network. Presented at: 94th annual conference & exhibition of the Air &Waste Management Association; June, Orlando, FL. Pittsburgh, PA: Air & Waste Management Association.

Rodes, C. E.; Evans, E. G. (1985) Preliminary assessment of 10 μm particulate sampling at eight locations in the United States. Atmos. Environ. 19: 293-303.

Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993) Sources of fine organic aerosol. 2. Noncatalyst and catalyst-equipped automobiles and heavy-duty diesel trucks. Environ. Sci. Technol. 27: 636-651.

Ryan, W. M.; Badgett-West, C. R.; Holtz, D. R.; Peters, T. A.; Cooper, J. A.; Ono, D. (1988) Reconciliation of receptor and dispersion modeling impacts of PM_{10} in Hayden, Arizona. In: Mathai, C. V.; Stonefield, D. H., eds. PM_{-10}: implementation of standards, an APCA/EPA international specialty conference; February; San Francisco, CA. Pittsburgh, PA: Air Pollution Control Association; pp. 419-429. (APCA transactions series TR-13).

Salma, I.; Maenhaut, W.; Záray, G. (2002) Comparative study of elemental mass size distributions in urban atmospheric aerosol. J. Aerosol Sci. 33: 339-356.

Savoie, D. L.; Prospero, J. M. (1980) Water-soluble potassium, calcium, and magnesium in the aerosols over the tropical North Atlantic. J. Geophys. Res. 85: 385-392.

Saxena, P.; Hildemann, L. M. (1996) Water-soluble organics in atmospheric particles: a critical review of the literature and applications of thermodynamics to identify candidate compounds. J. Atmos. Chem. 24: 57-109.

Schauer, J. J.; Cass, G. R. (2000) Source apportionment of wintertime gas-phase and particle-phase air pollutants using organic compounds as tracers. Environ. Sci. Technol. 34: 1821-1832.

Schauer, J. J.; Rogge, W. F.; Hildemann, L. M.; Mazurik, M. A.; Cass, G. R. (1996) Source apportionment of airborne particulate matter using organic compounds as tracers. Atmos. Environ. 30: 3837-3855.

Shinn, E. A.; Smith, G. W.; Prospero, J. M.; Betzer, P.; Hayes, M. L.; Garrison, V.; Barber, R. T. (2000) African dust and the demise of Caribbean coral reefs. Geophys. Res. Lett. 27: 3029-3032.

Skidmore, L. W.; Chow, J. C.; Tucker, T. T. (1992) PM_{10} air quality assessment for the Jefferson County, Ohio air quality control region. In: Chow, J. C.; Ono, D. M., eds. PM_{10} standards and nontraditional particulate source controls, an A&WMA/EPA international specialty conference, v. II; January; Scottsdale, AZ. Pittsburgh, PA: Air and Waste Management Association; pp. 1016-1031. (A&WMA transactions series no. 22).

Smith, G. T.; Ives, L. D.; Nagelkerken, I. A.; Ritchie, K. B. (1996) Caribbean sea fan mortalities. Nature (London) 383: 487.

Spengler, J. D.; Briggs, S. L. K.; Özkaynak, H. (1986) Relationship between TSP measurements and size-fractionated particle mass measurements in six cities participating in the Harvard Air Pollution Health Study. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; EPA cooperative agreement no. CR-812667-01-0; December 5.

Spengler, J. D.; Koutrakis, P.; Dockery, D. W.; Raizenne, M.; Speizer, F. E. (1996) Health effects of acid aerosols on North American children: air pollution exposures. Environ. Health Perspect. 104: 492-499.

Suarez, A. E.; Caffrey, P. F.; Ondov, J. M.; Thaung, K. C. (1996) Size distribution of Ir-tagged soot from diesel sanitation trucks. J. Aerosol Sci. 27(suppl. 1): S697-S698.

Tanner, R. L.; Parkhurst, W. J. (2000) Chemical composition of fine particles in the Tennessee Valley region. J. Air Waste Manage. Assoc. 50: 1299-1307.

Taylor, C. A., Jr.; Stover, C. A.; Westerdahl, F. D. (1998) Speciated fine particle (<2.5 μm aerodynamic diameter) and vapor-phase acid concentrations in southern California. Presented at: 91st annual meeting & exhibition of the Air & Waste Management Association; June; San Diego, CA. Pittsburgh, PA: Air & Waste Management Association; paper no. WA7401.

Thanukos, L. C.; Miller, T.; Mathai, C. V.; Reinholt, D.; Bennett, J. (1992) Intercomparison of PM_{10} samplers and source apportionment of ambient PM_{10} concentrations at Rillito, Arizona. In: Chow, J. C.; Ono, D. M., eds. PM_{10} standards and nontraditional particulate source controls, an A&WMA/EPA international specialty conference, v. I; January; Scottsdale, AZ. Pittsburgh, PA: Air & Waste Management Association; pp. 244-261. (A&WMA transactions series no. 22).

Tobias, H. J.; Ziemann, P. J. (2000) Thermal desorption mass spectrometric analysis of rganic aerosol formed from reactions of 1-tetradecene and O_3 in the presence of alcohols and carboxylic acids. Environ. Sci. Technol. 34: 2105-2115.

Turpin, B. J.; Huntzicker, J. J. (1991) Secondary formation of organic aerosol in the Los Angeles basin: a descriptive analysis of organic and elemental carbon concentrations. Atmos. Environ. Part A 25: 207-215.

Turpin, B. J.; Huntzicker, J. J. (1995) Identification of secondary organic aerosol episodes and quantitation of primary and secondary organic aerosol concentrations during SCAQS. Atmos. Environ. 29: 3527-3544.

Turpin, B. J.; Lim, H.-J. (2001) Species contributions to PM2.5 mass concentrations: revisiting common assumptions for estimating organic mass. Aerosol Sci. Technol. 35: 602-610.

Turpin, B. J.; Huntzicker, J. J.; Larson, S. M.; Cass, G. R. (1991) Los Angeles summer midday particulate carbon: primary and secondary aerosol. Environ. Sci. Technol. 25: 1788-1793.

U.S. Environmental Protection Agency. (1995) Compilation of air pollutant emission factors. Volume I: stationary point and area sources. 5th ed. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. AP-42-ED-5. Available from: NTIS, Springfield, VA; PB95-196028.

U.S. Environmental Protection Agency. (1996) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.

U.S. Environmental Protection Agency. (1999) Getting started: emissions inventory methods for PM-2.5. Research Triangle Park, NC: Office of Air Quality Planning and Standards, Emission Factor and Inventory Group; EIIP document series - volume IX. Available: http://www.epa.gov/ttn/chief/eiip/techreport/volume09/index.html [4 April, 2002].

U.S. Environmental Protection Agency. (2000a) Aerometric Information Retrieval System (AIRS). Research Triangle Park, NC: Office of Air Quality Planning and Standards. Available: www.epa.gov/ttn/airs/ [2000, October 6].

U.S. Environmental Protection Agency. (2000b) Air quality criteria for carbon monoxide. Research Triangle Park, NC: National Center for Environmental Assessment; report no. EPA/600/P-99/001F. Available: www.epa.gov/ncea/co/ [2000, October 6].

U.S. Environmental Protection Agency. (2000c) National air pollutant emission trends, 1900-1998. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA-454/R-00-002. Available: http://www.epa.gov/ttn/chief/trends/trends98/ [2001, January 30].

U.S. Environmental Protection Agency. (2000d) National air quality and emissions trends report, 1998. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA/454/R-00-003. Available from: NTIS, Springfield, VA; PB2000-104960.

U.S. Environmental Protection Agency. (2001) National air quality and emissions trends report, 1999. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA/454/R-01-004. Available: http://www.epa.gov/oar/aqtrnd99/ [11 March, 2002].

Vermette, S. J.; Williams, A. L.; Landsberger, S. (1992) PM_{10} source apportionment using local surface dust profiles: examples from Chicago. In: Chow, J. C.; Ono, D. M., eds. PM_{10} standards and nontraditional particulate source controls, an A&WMA/EPA international specialty conference, v. I; January; Scottsdale, AZ. Pittsburgh, PA: Air & Waste Management Association; pp. 262-278. (A&WMA transactions series no. 22).

Wainman, T.; Zhang, J.; Weschler, C. J.; Lioy, P. J. (2000) Ozone and limonene in indoor air: a source of submicron particle exposure. Environ. Health Perspect. 108: 1139-1145.

Watson, J. G.; Chow, J. C. (1999) Reconciling urban fugitive dust emissions inventory and ambient source contribution estimates: summary of current knowledge and needed research [draft]. Reno, NV: Desert Research Institute; DRI document no. 6110.4D2.

Watson, J. G.; Chow, J. C. (2001) Estimating middle-, neighborhood-, and urban-scale contributions to elemental carbon in Mexico City with a rapid response aethalometer. J. Air Waste Manage. Assoc. 51: 1522-1528.

Watson, J. G.; Robinson, N. F.; Chow, J. C.; Henry, R. C.; Kim, B. M.; Pace, T. G.; Meyer, E. L.; Nguyen, Q. (1990a) The USEPA/DRI chemical mass balance receptor model, CMB 7.0. Environ. Software 5: 38-49.

Watson, J. G.; Chow, J. C.; Pritchett, L. C.; Houck, J. A.; Ragazzi, R. A.; Burns, S. (1990b) Chemical source profiles for particulate motor vehicle exhaust under cold and high altitude operating conditions. Sci. Total Environ. 93: 183-190.

Watson, J. G.; Chow, J. C.; Lu, Z.; Fujita, E. M.; Lowenthal, D. H.; Lawson, D. R.; Ashbaugh, L. L. (1994) Chemical mass balance source apportionment of PM_{10} during the Southern California Air Quality Study. Aerosol Sci. Technol. 21: 1-36.

West, J. J.; Ansari, A. S.; Pandis, S. N. (1999) Marginal $PM_{2.5}$: nonlinear aerosol mass response to sulfate reductions in the eastern United States. J. Air Waste Manage. Assoc. 49: 1415-1424.

White, W. H. (1998) Statistical considerations in the interpretation of size-resolved particulate mass data. J. Air Waste Manage. Assoc. 48: 454-458.

Wilson, W. E.; Stockburger, L. (1990) Diurnal variations in aerosol composition and concentration. In: Masuda, S.; Takahashi, K., eds. Aerosols: science, industry, health and environment: proceedings of the third international aerosol conference, v. 2; September; Kyoto, Japan. Oxford, United Kingdom: Pergamon Press plc; pp. 962-965.

Wilson, W. E.; Suh, H. H. (1997) Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies. J. Air Waste Manage. Assoc. 47: 1238-1249.

Winer, A. M.; Arey, J.; Atkinson, R.; Aschmann, S. M.; Long, W. D.; Morrison, C. L.; Olszyk, D. M. (1992) Emission rates of organics from vegetation in California's Central Valley. Atmos. Environ. Part A 26: 2647-2659.

Winterhalter, R.; Neeb, P.; Grossmann, D.; Kolloff, A.; Horie, O.; Moortgat, G. (2000) Products and mechanisms of the gas phase reaction of ozone with β-pinene. J. Atmos. Chem. 35: 165-197.

Wongphatarakul, V.; Friedlander, S. K.; Pinto, J. P. (1998) A comparative study of $PM_{2.5}$ ambient aerosol chemical databases. Environ. Sci. Technol. 32: 3926-3934.

Woo, K.-S.; Chen, D.-R.; Pui, D. Y. H.; McMurry, P. H. (2001) Measurement of Atlanta aerosol size distributions: observations of ultrafine particle events. Aerosol Sci. Technol. 34: 75-87.

Wotawa, G.; Trainer, M. (2000) The influence of Canadian forest fires on pollutant concentrations in the United States. Science (Washington, DC) 288: 324-328.

Yu, J.; Cocker, D. R., III; Griffin, R. J.; Flagan, R. C.; Seinfeld, J. H. (1998) Gas-phase ozone oxidation of monoterpenes: gaseous and particulate products. J. Atmos. Chem. 34: 207-258.

Yu, J.; Griffin, R. J.; Cocker, D. R., III; Flagan, R. C.; Seinfeld, J. H. (1999) Observation of gaseous and particulate products of monoterpene oxidation in forest atmospheres. Geophys. Res. Lett. 26: 1145-1148.

Zhang, S.-H.; Shaw, M.; Seinfeld, J. H.; Flagan, R. C. (1992) Photochemical aerosol formation from α-pinene and β-pinene. J. Geophys. Res. [Atmos.] 97: 20,717-20,729.

Zweidinger, R.; Purdue, L.; Fitzgerald, K.; Carmichael, L.; Kellog, R. (1998) NERL PM research monitoring platforms: Baltimore, Fresno and Phoenix, data report. Research Triangle Park, NC: U.S. Environmental Protection Agency, National Exposure Research Laboratory; report no. EPA/600/R-98/138. Available from: NTIS, Springfield, VA; PB99-105660.

APPENDIX 3A

Spatial and Temporal Variability of the Nationwide AIRS $PM_{2.5}$ and $PM_{10-2.5}$ Data Sets

Aspects of the spatial and temporal variability of 24-h average $PM_{2.5}$ concentrations for 1999 and 2000 in a number of metropolitan statistical areas (MSAs) across the United States are presented in this Appendix. Data for multiple sites in 27 urban areas have been obtained from the AIRS data base and analyzed for their seasonal variations, for their spatial correlations, and for their spatial uniformity (Pinto et al., 2002). A number of aspects of the spatial and temporal variability of the $PM_{2.5}$ data set from 1999 were presented in Rizzo and Pinto (2001), based in part on analyses given in Fitz-Simons et al. (2000).

Quality assured measurements for at least fifteen days during each calendar quarter for 1999 and 2000, or for 2000 alone, at a minimum of four monitoring sites in a given MSA were required for their inclusion in the analyses given in this appendix. The Baton Rouge, LA MSA, which had only three sites meeting this criterion, was an exception. Data from Baton Rouge were included for the sake of geographic coverage. Typically, at least 200 measurements were available for each monitoring site chosen. Monitoring sites were chosen without consideration of the land use type used to characterize their locations.

Because of changes in monitoring strategies, funding levels etc., there were year to year changes in monitoring sites meeting the above criteria in a number of MSAs. Data for the Philadelphia, PA, Norfolk, VA, Pittsburgh, PA, Detroit, MI, Chicago, IL, Louisville, KY, St. Louis, MO, and the Dallas, TX MSAs have been analyzed only for the year 2000 because of a lack of consistent coverage in 1999.

Information about seasonal and spatial variability in $PM_{2.5}$ concentrations within 27 MSAs across the United States are provided in the accompanying figures (Figures 3A-1 to 3A-27). Underneath the value for r, the 90th percentile values of the absolute difference in $PM_{2.5}$ concentrations (in $\mu g/m^3$) and the coefficient of divergence (COD) are given in parentheses.

Underneath these two measures of spatial variability, the numbers of observations used in the calculations of the statistics in part *c* of each figure are given.

The COD was defined mathematically and used earlier in Chapter 3 as a measure of the degree of similarity between two data sets. A COD of zero implies that values in both data sets are identical, and a COD of one indicates that two data sets are completely different. Values of P_{90} provide a measure in absolute terms of differences in concentrations between sites, and CODs provide a relative measure of these differences. The maximum number of days of coincident data from paired sites were used to calculate correlation coefficients, values for P_{90}, and CODs. The correlation coefficients were also calculated by using only concurrent measurements obtained at all of the monitoring sites within urban areas meeting the above selection criteria. The correlation coefficients that were calculated differed only in the third significant figure between the two methods.

Information about the spatial and temporal variability of 24-h average $PM_{10-2.5}$ concentrations is summarized in Figures 3A-28 to 3A-32. Data are shown for 2000 for all MSAs, except the Los Angeles-Long Beach MSA, for which data are shown for 1999. A schematic map showing locations of sampling sites within each MSA is given in part *a*, at the top of each figure. Also included in the map are major highways and a distance scale. A key giving the AIRS site ID #'s is shown alongside each map. Box plots showing lowest, lower quartile, median, upper quartile and highest $PM_{2.5}$ concentrations for each calendar quarter are shown in part *b* of each figure. AIRS site ID #'s, annual mean concentrations, the number of observations, and the standard deviation of the data are shown above the box plots. Finally, in part *c* of each figure, statistics characterizing the spatial variability in $PM_{2.5}$ concentrations are given. For each site-pair, the Pearson correlation coefficient (r) is provided. Underneath each value for r, the number of observations is given.

Philadelphia, PA MSA

a.



	AIRS Site ID
Site A	34-007-0003
Site B	34-007-1007
Site C	42-045-0002
Site D	42-101-0004
Site E	42-101-0136

b.

AIRS ID#	3400700031	3400710071	4204500021	4210100041	4210101361
Mean	14.941	15.427	15.992	14.823	14.718
Obs	108	103	112	284	278
SD	8.524	8.749	8.265	8.537	8.295



c.

Site	A	B	C	D	E
A	1	0.964 (3.3, 0.082) 95	0.868 (6.3, 0.155) 98	0.88 (4.8, 0.129) 81	0.868 (5.4, 0.147) 80
B		1	0.849 (6.9, 0.158) 94	0.894 (3.7, 0.135) 77	0.857 (6.4, 0.148) 79
C			1	0.868 (5.0, 0.149) 85	0.818 (6.6, 0.154) 83
D				1	0.918 (4.9, 0.13) 246
E					1

Figure 3A-1. Philadelphia, PA-NJ MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.



Figure 3A-2. Washington, DC MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.



Figure 3A-3. Norfolk, VA MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Columbia, SC MSA

a.



	AIRS Site ID
Site A	45-063-0005
Site B	45-063-0008
Site C	45-079-0007
Site D	45-079-0019

b.



c.

Site	A	B	C	D
A	1	0.882 (5.3, 0.121) 215	0.949 (3.9, 0.081) 204	0.93 (4.8, 0.099) 216
B		1	0.933 (4.0, 0.082) 202	0.949 (3.3, 0.067) 216
C			1	0.971 (2.7, 0.06) 203
D				1

Figure 3A-4. Columbia, SC MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

a.



	AIRS Site ID
Site A	13-063-0091
Site B	13-089-0002
Site C	13-089-2001
Site D	13-121-0032
Site E	13-121-0039
Site F	13-121-1001
Site G	13-223-0003

b.



c.

Site	A	B	C	D	E	F	G
A	1	0.808 (8.3, 0.161) 148	0.828 (8.1, 0.115) 167	0.795 (7.1, 0.150) 158	0.82 (8.2, 0.131) 159	0.886 (7.7, 0.140) 162	0.543 (13.9, 0.219) 158
B		1	0.77 (8.2, 0.158) 466	0.669 (8.3, 0.178) 440	0.855 (9.9, 0.174) 150	0.841 (9.3, 0.181) 148	0.639 (12.5, 0.224) 155
C			1	0.727 (6.5, 0.143) 474	0.861 (9.0, 0.129) 167	0.802 (10.6, 0.192) 165	0.574 (12.0, 0.207) 166
D				1	0.815 (9.4, 0.157) 160	0.819 (9.5, 0.182) 162	0.527 (10.9, 0.218) 161
E					1	0.785 (10.5, 0.188) 158	0.574 (14.0, 0.242) 160
F						1	0.547 (9.9, 0.213) 155
G							1

Figure 3A-5. Atlanta, GA MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.



Figure 3A-6. Birmingham, AL MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.



Figure 3A-7. Tampa, FL MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

a.



	AIRS Site ID
Site A	39-035-0013
Site B	39-035-0038
Site C	39-035-0060
Site D	39-035-0065
Site E	39-035-0066
Site F	39-035-1002
Site G	39-085-1001

b.



c.

Site	A	B	C	D	E	F	G
A	1	0.93 (6.9, 0.110) 194	0.946 (3.8, 0.135) 206	0.939 (5.3, 0.096) 206	0.92 (6.8, 0.150) 192	0.847 (9.7, 0.192) 198	0.872 (10.7, 0.211) 197
B		1	0.904 (7.1, 0.164) 192	0.869 (10.3, 0.156) 189	0.793 (13.8, 0.227) 184	0.791 (14.1, 0.220) 195	0.793 (14.9, 0.240) 184
C			1	0.912 (5.2, 0.156) 198	0.868 (8.3, 0.204) 189	0.823 (9.6, 0.222) 197	0.849 (11.8, 0.240) 192
D				1	0.94 (5.5, 0.161) 185	0.868 (6.3, 0.179) 197	0.874 (8.2, 0.216) 192
E					1	0.812 (5.7, 0.193) 192	0.883 (6.6, 0.166) 185
F						1	0.85 (6.9, 0.196) 187
G							1

Figure 3A-8. Cleveland, OH MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Pittsburgh, PA MSA

a.



	AIRS Site ID
Site A	42-003-0008
Site B	42-003-0064
Site C	42-125-0200
Site D	42-125-5001

b.



c.

Site	A	B	C	D
A	1	0.767 (14.2, 0.162) 217	0.937 (4.1, 0.089) 70	0.876 (7.4, 0.154) 213
B		1	0.58 (21.3, 0.238) 95	0.575 (20.5, 0.272) 297
C			1	0.905 (5.9, 0.138) 98
D				1

Figure 3A-9. Pittsburgh, PA MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Steubenville, OH-Weirton, WV MSA

a.



	AIRS Site ID
Site A	39-081-0016
Site B	39-081-1001
Site C	54-009-0005
Site D	54-029-0011
Site E	54-029-0011

b.



c.

Site	A	B	C	D	E
A	1	0.831 (9.9, 0.186) 166	0.85 (8.7, 0.172) 196	0.843 (8.2, 0.187) 201	0.805 (9.9, 0.199) 194
B		1	0.846 (6.6, 0.171) 166	0.86 (9.2, 0.183) 174	0.816 (9.2, 0.207) 161
C			1	0.883 (7.6, 0.164) 220	0.873 (7.9, 0.179) 204
D				1	0.978 (2.5, 0.101) 210
E					1

Figure 3A-10. Steubenville, OH-Weirton, WV MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Detroit, MI MSA

a.



	AIRS Site ID
Site A	26-099-0009
Site B	26-125-0001
Site C	26-147-0005
Site D	26-163-0033
Site E	26-163-0036

b.



c.

Site	A	B	C	D	E
A	1	0.958 (4.9, 0.107) 83	0.952 (5.6, 0.127) 96	0.931 (12.7, 0.222) 98	0.926 (9.0, 0.177) 96
B		1	0.939 (5.8, 0.121) 73	0.92 (12.3, 0.193) 77	0.917 (8.3, 0.151) 75
C			1	0.876 (13.3, 0.222) 89	0.875 (8.9, 0.197) 88
D				1	0.923 (7.1, 0.108) 89
E					1

Figure 3A-11. Detroit MI MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.



Figure 3A-12. Grand Rapids, MI MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Milwaukee, WI MSA

a.



	AIRS Site ID
Site A	55-079-0010
Site B	55-079-0026
Site C	55-079-0043
Site D	55-079-0051
Site E	55-079-0059
Site F	55-079-0099
Site G	55-133-0027
Site H	55-133-0034

b.



c.

Site	A	B	C	D	E	F	G	H
A	1	0.934 (3.3, 0.111) 654	0.966 (3.7, 0.089) 213	0.974 (3.0, 0.086) 209	0.96 (3.0, 0.094) 217	0.977 (3.9, 0.085) 228	0.96 (3.5, 0.089) 451	0.961 (3.3, 0.108) 227
B		1	0.947 (3.7, 0.112) 210	0.947 (3.8, 0.116) 206	0.944 (3.8, 0.112) 215	0.963 (3.8, 0.110) 225	0.95 (3.7, 0.111) 441	0.932 (3.6, 0.130) 224
C			1	0.945 (4.4, 0.121) 188	0.95 (4.7, 0.100) 187	0.964 (3.6, 0.094) 198	0.948 (3.9, 0.106) 212	0.929 (5.3, 0.141) 200
D				1	0.954 (3.9, 0.111) 194	0.956 (4.5, 0.114) 212	0.967 (3.6, 0.093) 207	0.964 (3.5, 0.111) 209
E					1	0.963 (3.5, 0.098) 203	0.959 (3.5, 0.095) 211	0.96 (3.7, 0.123) 207
F						1	0.963 (4.4, 0.099) 219	0.95 (5.5, 0.133) 221
G							1	0.973 (2.9, 0.101) 220
H								1

Figure 3A-13. Milwaukee, WI MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Chicago, IL MSA

a.



	AIRS Site ID
Site A	17-031-0014
Site B	17-031-0022
Site C	17-031-0050
Site D	17-031-0052
Site E	17-031-1016
Site F	17-031-2001
Site G	17-031-3301
Site H	17-031-4006
Site I	17-031-4201
Site J	17-043-4002
Site K	17-197-1002

b.

AIRS ID#	1703100141	1703100221	1703100501	1703100521	1703110161	1703120011	1703133011	1703140061	1703142011	1704340021	1719710021
Mean	15.823	17.933	16.996	18.295	20.277	16.790	16.889	15.268	14.283	15.215	15.994
Obs	104	113	274	346	108	113	115	101	327	116	112
SD	7.935	8.175	8.468	9.289	9.331	7.694	7.689	8.423	7.905	7.568	7.405



c.

Site	A	B	C	D	E	F	G	H	I	J	K
A	1	0.912 (4.4, 0.121) 96	0.946 (4.6, 0.077) 78	0.909 (6.6, 0.13) 100	0.921 (7.5, 0.143) 92	0.902 (5.6, 0.111) 98	0.927 (5.1, 0.104) 98	0.876 (5.8, 0.133) 88	0.936 (5.3, 0.139) 95	0.885 (5.7, 0.13) 95	0.774 (7.4, 0.158) 81
B		1	0.92 (5.4, 0.113) 87	0.872 (6.5, 0.14) 108	0.866 (7.0, 0.141) 103	0.892 (5.7, 0.131) 104	0.879 (6.0, 0.132) 106	0.689 (7.9, 0.213) 92	0.86 (7.9, 0.197) 101	0.855 (7.2, 0.165) 100	0.79 (7.1, 0.17) 87
C			1	0.941 (5.0, 0.094) 259	0.93 (7.8, 0.012) 83	0.955 (3.5, 0.082) 89	0.923 (5.3, 0.096) 91	0.75 (7.9, 0.176) 75	0.928 (6.2, 0.162) 247	0.922 (5.3, 0.117) 91	0.867 (7.6, 0.131) 87
D				1	0.887 (7.9, 0.133) 105	0.885 (7.3, 0.125) 109	0.881 (7.0, 0.128) 110	0797 (8.5, 0.177) 98	0.879 (9.6, 0.179) 310	0.836 (8.5, 0.154) 112	0.721 (10.2, 0.169) 108
E					1	0.932 (7.3, 0.108) 99	0.898 (7.5, 0.124) 102	0.787 (10.0, 0.205) 92	0.915 (9.8, 0.2) 98	0.902 (9.5, 0.154) 95	0.84 (10.5, 0.173) 85
F						1	0.931 (4.5, 0.084) 110	0.861 (5.9, 0.153) 93	0.943 (5.5, 0.14) 101	0.949 (4.3, 0.1) 99	0.893 (5.1, 0.118) 89
G							1	0.823 (7.0, 0.158) 95	0.915 (6.4, 0.152) 103	0.953 (4.4, 0.092) 101	0.873 (5.8, 0.128) 91
H								1	0.818 (7.3, 0.146) 92	0.865 (5.1, 0.124) 88	0.752 (7.6, 0.161) 78
I									1	0.922 (4.8, 0.123) 106	0.809 (7.1, 0.157) 99
J										1	0.921 (4.2, 0.099) 90
K											1

Figure 3A-14. Chicago, IL MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Gary, IN MSA

a.



	AIRS Site ID
Site A	18-089-0006
Site B	18-089-0022
Site C	18-089-1016
Site D	18-127-0024

0 20 40 km

b.



c.

Site	A	B	C	D
A	1	0.666 (8.6, 0.205) 502	0.849 (5.9, 0.164) 512	0.864 (7.2, 0.214) 167
B		1	0.568 (11.5, 0.220) 532	0.525 (14.9, 0.265) 169
C			1	0.838 (7.9, 0.218) 177
D				1

Figure 3A-15. Gary, IN MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Louisville, KY MSA

a.



	AIRS Site ID
Site A	18-019-0005
Site B	18-043-1004
Site C	21-111-0043
Site D	21-111-0044

b.



c.

Site	A	B	C	D
A	1	0.768 (7.8, 0.131) 78	0.661 (8.4, 0.139) 66	0.414 (11.2, 0.23) 89
B		1	0.818 (6.3, 0.148) 59	0.592 (9.0, 0.222) 76
C			1	0.775 (9.3, 0.143) 232
D				1

Figure 3A-16. Louisville, KY MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

St. Louis, MO MSA

a.



	AIRS Site ID
Site A	17-119-0023
Site B	17-119-1007
Site C	17-119-3007
Site D	17-163-0010

b.



c.

Site	A	B	C	D
A	1	0.79 (11.1, 0.145) 111	0.76703 (12.2, 0.176) 107	0.67 (12.5, 0.192) 100
B		1	0.842 (6.0, 0.121) 107	0.786 (7.9, 0.149) 101
C			1	0.812 (7.8, 0.151) 105
D				1

Figure 3A-17. St. Louis, MO MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Baton Rouge, LA MSA

a.



	AIRS Site ID
Site A	22-033-0002
Site B	22-033-0009
Site C	22-121-0001

b.



c.

Site	A	B	C
A	1	0.938 (2.9, 0.082) 215	0.948 (3.2, 0.078) 207
B		1	0.96 (2.5, 0.071) 663
C			1

Figure 3A-18. Baton Rouge, LA MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Kansas City, KS-MO, MSA

a.



	AIRS Site ID
Site A	20-091-0008
Site B	20-091-0009
Site C	29-047-0005
Site D	29-047-0026
Site E	29-047-0041
Site F	29-095-2002

b.

AIRS ID#	2009100081	2009100091	2904700051	2904700261	2904700411	2909520021
Mean	11.727	11.262	11.122	12.738	12.034	13.682
Obs	212	214	214	669	623	235
SD	5.986	5.860	5.910	6.300	5.958	6.609



c.

Site	A	B	C	D	E	F
A	1	0.94 (1.9, 0.105) 200	0.918 (4.1, 0.124) 184	0.895 (4.0, 0.130) 201	0.902 (4.1, 0.116) 182	0.893 (5.8, 0.142) 204
B		1	0.9 (3.9, 0.143) 189	0.869 (4.23, 0.150) 201	0.862 (3.6, 0.143) 187	0.865 (6.1, 0.170) 208
C			1	0.943 (3.9, 0.101) 202	0.929 (3.7, 0.114) 181	0.884 (7.4, 0.169) 206
D				1	0.915 (3.8, 0.115) 578	0.933 (5.0, 0.118) 220
E					1	0.908 (5.7, 0.135) 197
F						1

Figure 3A-19. Kansas City, KS-MO MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

a.



	AIRS Site ID
Site A	48-085-0005
Site B	48-113-0020
Site C	48-113-0035
Site D	48-113-0050
Site E	48-113-0057
Site F	48-113-0069
Site G	48-113-0087

b.



c.

Site	A	B	C	D	E	F	G
A	1	0.876 (3.5, 0.119) 97	0.906 (4.3, 0.108) 90	0.935 (4.1, 0.109) 89	0.872 (5.6, 0.13) 83	0.939 (3.3, 0.1) 106	0.949 (2.7, 0.084) 76
B		1	0.934 (3.4, 0.086) 91	0.929 (3.5, 0.088) 246	0.877 (5.3, 0.108) 83	0.917 (3.0, 0.089) 298	0.932 (3.6, 0.089) 83
C			1	0.967 (2.0, 0.052) 84	0.926 (2.8, 0.08) 79	0.969 (2.2, 0.059) 98	0.956 (3.2, 0.086) 72
D				1	0.918 (2.6, 0.078) 81	0.974 (2.2, 0.056) 273	0.948 (3.6, 0.101) 74
E					1	0.939 (2.1, 0.072) 95	0.915 (4.9, 0.114) 71
F						1	0.97 (3.0, 0.075) 89
G							1

Figure 3A-20. Dallas, TX MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

a.



	AIRS Site ID
Site A	16-010-0011
Site B	16-001-0017
Site C	16-027-0001
Site D	16-027-0005

b.



c.

Site	A	B	C	D
A	1	0.889 (4.4, 0.155) 194	0.944 (3.8, 0.118) 229	0.882 (6.0, 0.158) 198
B		1	0.837 (6.6, 0.184) 199	0.75 (8.9, 0.231) 203
C			1	0.948 (4.1, 0.109) 203
D				1

Figure 3A-21. Boise, ID MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Salt Lake City, UT MSA

a.



	AIRS Site ID
Site A	49-011-0001
Site B	49-035-0003
Site C	49-035-0012
Site D	49-035-3006
Site E	49-035-3007
Site F	49-057-0007

b.

AIRS ID#	4901100011	4903500031	4903500121	4903530061	4903530071	4905700071
Mean	8.504	11.413	13.447	10.604	11.101	7.765
Obs	237	231	228	644	221	215
SD	7.963	10.812	11.374	10.285	10.411	4.921



c.

Site	A	B	C	D	E	F
A	1	0.821 (8.8, 0.236) 221	0.881 (12.6, 0.281) 219	0.927 (6.7, 0.204) 211	0.89 (8.5, 0.225) 208	0.883 (3.9, 0.160) 203
B		1	0.834 (7.6, 0.207) 214	0.866 (5.4, 0.189) 205	0.828 (6.1, 0.198) 204	0.769 (8.8, 0.213) 203
C			1	0.92 (7.1, 0.201) 203	0.961 (5.0, 0.173) 200	0.825 (11.8, 0.257) 199
D				1	0.936 (5.1, 0.167) 194	0.883 (7.9, 0.180) 190
E					1	0.868 (8.7, 0.211) 188
F						1

Figure 3A-22. Salt Lake City, UT MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Seattle, WA MSA

a.



	AIRS Site ID
Site A	53-033-0017
Site B	53-033-0021
Site C	53-033-0057
Site D	53-033-0080
Site E	53-061-1007

b.

AIRS ID#	5303300171	5303300211	5303300571	5303300801	5306110071
Mean	5.963	11.053	12.070	9.165	11.437
Obs	200	708	699	459	223
SD	3.664	7.542	7.293	5.482	8.743



c.

Site	A	B	C	D	E
A	1	0.413 (15.0, 0.378) 194	0.412 (14.9, 0.419) 191	0.501 (10.4, 0.309) 178	0.487 (15.3, 0.372) 184
B		1	0.952 (3.8, 0.148) 679	0.932 (6.9, 0.150) 441	0.818 (6.5, 0.157) 217
C			1	0.918 (7.7, 0.204) 443	0.778 (8.4, 0.199) 214
D				1	0.773 (7.9, 0.195) 199
E					1

Figure 3A-23. Seattle, WA MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.





c.

Site	A	B	C	D
A	1	0.799 (6.5, 0.238) 319	0.921 (4.5, 0.1935) 309	0.813 (4.4, 0.170) 313
B		1	0.747 (4.0, 0.146) 631	0.894 (4.3, 0.162) 362
C			1	0.826 (4.6, 0.18) 349
D				1

Figure 3A-24. Portland, OR MSA. *(a)* Locations of sampling sites by AIRS ID#; (*b*) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; (*c*) Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Los Angeles-Long Beach, CA MSA

a.



	AIRS Site ID
Site A	06-037-0002
Site B	06-037-1103
Site C	06-037-1601
Site D	06-037-4002
Site E	06-037-9002

b.



c.

Site	A	B	C	D	E
A	1	0.828 (12.8, 0.192) 391	0.763 (17.3, 0.211) 196	0.573 (20.2, 0.263) 379	0.276 (28.0, 0.392) 186
B		1	0.88 (11.8, 0.140) 173	0.752 (14.6, 0.191) 353	0.328 (26.4, 0.375) 164
C			1	0.859 (11.8, 0.174) 171	0.363 (31.0, 0.411) 181
D				1	0.338 (24.4, 0.356) 157
E					1

Figure 3A-25. Los Angeles-Long Beach, CA MSA. *(a)* Locations of sampling sites by AIRS ID; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

Riverside-San Bernadino, CA MSA

a.



	AIRS Site ID
Site A	06-065-1003
Site B	06-065-8001
Site C	06-071-0025
Site D	06-071-2002
Site E	06-071-9004

b.



c.

Site	A	B	C	D	E
A	1	0.925 (7.0, 0.118) 174	0.799 (15.9, 0.228) 173	0.93 (10.7, 0.140) 184	0.885 (10.0, 0.138) 164
B		1	0.721 (20.2, 0.252) 174	0.888 (13.3, 0.152) 186	0.847 (11.9, 0.147) 158
C			1	0.824 (15.1, 0.229) 180	0.73 (18.1, 0.247) 162
D				1	0.91 (9.2, 0.122) 173
E					1

Figure 3A-26. Riverside-San Bernadino, CA MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.

San Diego, CA MSA

a.



	AIRS Site ID
Site A	06-073-0001
Site B	06-073-0003
Site C	06-073-1002
Site D	06-073-1007

b.

AIRS ID#	0607300011	0607300031	0607310021	0607310071
Mean	14.153	15.788	16.041	16.530
Obs	185	579	518	541
SD	7.320	8.059	8.563	14.153



c.

Site	A	B	C	D
A	1	0.722 (10.1, 0.197) 144	0.728 (10.1, 0.200) 127	0.787 (11.1, 0.177) 139
B		1	0.818 (7.4, 0.156) 436	0.699 (11.1, 0.212) 441
C			1	0.693 (11.9, 0.226) 385
D				1

Figure 3A-27. San Diego, CA MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{2.5}$ concentrations; *(c)* Intersite correlation statistics, for each data pair, the correlation coefficient, (P_{90}, coefficient of divergence) and number of measurements are given.



Figure 3A-28. Columbia, SC MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations.



Figure 3A-29. Detroit, MI MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations; *(c)* Intersite correlation coefficients and number of measurements.

Cleveland. OH MSA

a



	AIRS Site ID
Site A	39-035-0013
Site B	39-035-0038
Site C	39-035-0045
Site D	39-035-0060
Site E	39-035-0065
Site F	39-085-1001

b



c

Site	A	B	C	D	E	F
A	1	0.602 90	0.715 44	0.691 43	0.612 47	0.442 43
B		1	0.67 42	0.735 42	0.705 43	0.426 42
C			1	0.636 44	0.724 47	0.529 43
D				1	0.719 44	0.216 44
E					1	0.215 45
F						1

Figure 3A-30. Cleveland, OH MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations; *(c)* Intersite correlation coefficients and number of measurements.



Figure 3A-31. Steubenville, OH-Weirton, WV MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations; *(c)* Intersite correlation coefficients and number of measurements.

St. Louis, MO-IL MSA

a



	AIRS Site ID
Site A	17-119-0023
Site B	17-119-3007
Site C	17-163-0010
Site D	29-510-0086

b



c

Site	A	B	C	D
A	1	0.698 51	0.731 47	0.679 48
B		1	0.82 50	0.83 51
C			1	0.837 48
D				1

Figure 3A-32. St. Louis, MO-IL MSA. *(a)* Locations of sampling sites by AIRS ID#; *(b)* Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations; *(c)* Intersite correlation coefficients and number of measurements.

REFERENCES

Fitz-Simons, T. S.; Mathias, S.; Rizzo, M. (2000) Analyses of 1999 PM data for the PM NAAQS review. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; November 17. Available: http://www.epa.gov/oar/oaqps/pm25/analyses.html [2 April, 2002].

Pinto, J. P.; Lefohn, A. S.; Shadwick, D. S. (2002) Aspects of the spatial variability of $PM_{2.5}$ concentrations within urban areas of the United States. Environ. Sci. Technol.: submitted.

Rizzo, M.; Pinto, J. P. (2001) Initial characterization of fine particulate matter (PM2.5) collected by the National Federal Reference Monitoring Network. Presented at: 94th annual conference & exhibition of the Air & Waste Management Association; June, Orlando, FL. Pittsburgh, PA: Air & Waste Management Association.

APPENDIX 3B

Aerosol Composition Data from the Speciation Network

Data from thirteen sites designed to evaluate the suitability of various aerosol sampling devices for obtaining $PM_{2.5}$ composition data are summarized in this appendix. Three types of aerosol sampling devices were used in this study, which lasted from February 2000 through July 2000. A network consisting of 54 core sites across the United States has been implemented to provide a consistent data set for the characterization and evaluation of trends in PM components. This network has been used as a model for the deployment of a more comprehensive network, consisting of approximately 250 additional sites. Data obtained from the three sampling devices are shown for each site. A complete description of the data, techniques used to analyze the filters, and the results of the evaluation of the performance of the sampling devices (including a number of caveats regarding the data) can be found in Coutant and Stetzer (2001) and the analyses of data in Coutant et al. (2001).

Summary statistics for concentrations of $PM_{2.5}$ are given in Table 3B-1. Data are presented for all of the sites used in the pilot study for the speciation network in Tables 3B-2 through 14. Entries in the tables give the AIRS ID for each site; the number of samples (N); the mean, minimum and maximum 24-h $PM_{2.5}$ and component concentrations; and the minimum detection limit for each constituent in the data sets for each site. Numbers given in parenthesis next to the sampler indicate the POC code for identifying samplers in AIRS. Mass was determined gravimetrically; anions and cations ammonium (through sulfate) were determined by ion chromatography; carbonaceous species were determined by the thermal optical reflectance method; and trace elements (aluminum through zirconium) were determined by X-ray fluorescence spectrometry. There is a residual unknown portion ranging from <1 $\mu m/m^3$ to 4 $\mu g/m^3$, depending on the site. This residual is based on a comparison of the mass measured gravimetrically with that determined by summing the contributions from measured components.

TABLE 3B-1. SUMMARY STATISTICS FOR $PM_{2.5}$ CONCENTRATIONS DURING FEBRUARY THROUGH JUNE 2000 OBTAINED BY COLLOCATED FRM SAMPLERS (in $\mu g/m^3$)

Site	N	Mean	Max	Min
Bismarck, ND (380150003)	60	5.97	14.30	2.50
Boston, MA (250250042)	34	12.53	28.70	5.10
Bronx Botanical Garden, NY (360050083)	62	13.87	39.00	4.70
Chicago, IL (170310050)	67	16.39	35.80	3.10
Fresno, CA (060190008)	86	11.12	50.00	4.00
Houston, TX (482011039)	34	12.24	21.90	5.90
Lewis, FL (120571075)	59	12.50	26.70	2.87
Philadelphia, PA (421010004)	51	13.93	42.50	3.70
Salt Lake City, UT (490353006)	35	6.52	23.70	2.50
Seattle, WA (530330080)	61	7.37	25.00	1.90
St. Louis, MO (295100085)	68	15.15	36.80	3.10

TABLE 3B-2. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT BISMARCK, ND (in $\mu g/m^3$)

	Bismarck, ND (380150003)									
	Met One (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	23	6.35392	10.9262	3.40788	0.10400	22	4.80160	8.88982	3.04179	0.04000
Ammonium (88301)	21	0.45558	1.18190	0.14508	0.017	21	0.55753	1.17094	0.29308	0.00700
Sodium Ion (88302)	20	0.08233	0.17132	0.02898	0.03	20	0.03012	0.09578	0.00408	0.01200
Potassium Ion (88303)	0	—	—	—	0.014	10	0.05161	0.07882	0.03138	0.00600
Nitrate (88306)	21	0.52373	2.03068	0.09675	0.00800	21	0.42761	1.75630	0.06708	0.00300
Sulfate (88403)	21	1.39787	3.32569	0.75503	0.012	21	1.31732	2.01230	0.71669	0.00500
Organic Carbon (88305)	25	2.43559	4.19042	1.56074	0.146	24	1.46333	3.29766	0.55629	0.05900
Elemental Carbon (88307)	25	0.23671	0.68209	0.02249	0.146	24	0.22107	0.72424	0.06229	0.05900
Total Carbon	25	2.67230	4.85158	1.67502	NA	24	1.68439	4.02190	0.79034	NA
Aluminum (88104)	17	0.03782	0.34114	0.00082	0.01088	14	0.02811	0.27570	0.00132	0.00436
Antimony (88102)	14	0.00645	0.01436	0.00103	0.01476	15	0.00275	0.00608	0.00104	0.00592
Arsenic (88103)	15	0.00179	0.00394	0.00023	0.00247	14	0.00060	0.00132	0.00019	0.00099
Barium (88107)	23	0.05272	0.09574	0.00336	0.05876	21	0.02441	0.04242	0.00217	0.02360
Bromine (88109)	17	0.00153	0.00322	0.00035	0.00199	21	0.00137	0.00278	0.00005	0.00080
Cadmium (88110)	7	0.00403	0.01319	0.00056	0.0105	10	0.00189	0.00330	0.00033	0.00421
Calcium (88111)	23	0.05576	0.23228	0.01600	0.00347	22	0.03126	0.16318	0.00899	0.00139
Carbonate Carbon (88308)	0	—	—	—	0.146	0	—	—	—	0.059
Cerium (88117)	13	0.02152	0.07436	0.00460	0.08603	10	0.01459	0.03447	0.00523	0.03450
Cesium (88118)	12	0.01574	0.04227	0.00035	0.03689	7	0.00620	0.01554	0.00028	0.01480
Chlorine (88115)	12	0.00231	0.00499	0.00069	0.00578	4	0.00122	0.00207	0.00033	0.00232
Chromium (88112)	9	0.00046	0.00138	0.00011	0.00159	4	0.00015	0.00023	0.00005	0.00063
Cobalt (88113)	0	—	—	—	0.00141	2	0.00016	0.00019	0.00014	0.00056
Copper (88114)	16	0.00081	0.00203	0.00011	0.00135	18	0.00042	0.00099	0.00005	0.00054

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-2 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT BISMARCK, ND (in $\mu g/m^3$)

	Bismarck, ND (380150003)									
	Met One (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Europium (88121)	1	0.00300	0.00300	0.00300	0.01124	4	0.00178	0.00292	0.00080	0.00451
Gallium (88124)	23	0.00275	0.00414	0.00023	0.00331	20	0.00115	0.00184	0.00028	0.00133
Gold (88143)	14	0.00216	0.00601	0.00023	0.00501	16	0.00072	0.00207	0.00005	0.00201
Hafnium (88127)	12	0.01164	0.02674	0.00024	0.02605	12	0.00372	0.00857	0.00014	0.01050
Indium (88131)	10	0.00443	0.00912	0.00093	0.01128	11	0.00197	0.00508	0.00013	0.00452
Iridium (88133)	18	0.00413	0.00780	0.00024	0.00594	18	0.00120	0.00240	0.00010	0.00238
Iron (88126)	23	0.05132	0.26884	0.01337	0.00196	22	0.03335	0.19338	0.00932	0.00079
Lanthanum (88146)	12	0.03345	0.05805	0.00572	0.06947	6	0.01280	0.02524	0.00532	0.02790
Lead (88128)	21	0.00382	0.01036	0.00012	0.00549	21	0.00228	0.00471	0.00071	0.00220
Magnesium (88140)	10	0.01462	0.05475	0.00106	0.01841	11	0.00722	0.01714	0.00217	0.00738
Manganese (88132)	20	0.00232	0.00990	0.00035	0.00231	21	0.00165	0.00631	0.00019	0.00092
Mercury (88142)	14	0.00227	0.00448	0.00035	0.00437	13	0.00099	0.00155	0.00014	0.00175
Molybdenum (88134)	9	0.00117	0.00453	0.00025	0.00477	11	0.00111	0.00212	0.00005	0.00191
Nickel (88136)	16	0.00295	0.02075	0.00011	0.00125	14	0.00038	0.00085	0.00005	0.00050
Niobium (88147)	11	0.00102	0.00287	0.00012	0.00420	11	0.00063	0.00141	0.00019	0.00168
Potassium (88180)	23	0.03065	0.15682	0.00138	0.00341	22	0.02871	0.13414	0.00207	0.00137
Rubidium (88176)	8	0.00118	0.00254	0.00011	0.00217	11	0.00037	0.00094	0.00005	0.00087
Samarium (88162)	1	0.00024	0.00024	0.00024	0.00617	4	0.00111	0.00245	0.00047	0.00247
Scandium (88163)	1	0.00012	0.00012	0.00012	0.00243	5	0.00021	0.00047	0.00005	0.00097
Selenium (88154)	12	0.00156	0.00281	0.00011	0.00212	17	0.00057	0.00122	0.00010	0.00085
Silicon (88165)	23	0.13816	0.84236	0.02970	0.00753	22	0.08587	0.60907	0.01615	0.00302
Silver (88166)	12	0.00522	0.01336	0.00108	0.01048	18	0.00204	0.00448	0.00010	0.00420
Sodium (88184)	18	0.06076	0.14392	0.00081	0.05107	16	0.04287	0.11010	0.01328	0.02050

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-2 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT BISMARCK, ND (in $\mu g/m^3$)

	Bismarck, ND (380150003)									
	Met One (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Tin (88160)	23	0.01887	0.02888	0.01044	0.01787	22	0.00843	0.01309	0.00579	0.00717
Titanium (88161)	23	0.00358	0.01910	0.00046	0.00208	22	0.00228	0.01281	0.00037	0.00083
Vanadium (88164)	0	—	—	—	0.00150	1	0.00005	0.00005	0.00005	0.00060
Wofram (88186)	14	0.00837	0.02148	0.00162	0.01380	15	0.00297	0.00537	0.00061	0.00554
Yttrium (88183)	6	0.00123	0.00264	0.00012	0.00304	2	0.00073	0.00123	0.00023	0.00122
Zinc (88167)	13	0.00321	0.01717	0.00083	0.00145	18	0.00206	0.00556	0.00014	0.00058
Zirconium (88185)	9	0.00083	0.00241	0.00011	0.00359	7	0.00051	0.00104	0.00014	0.001

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-3. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT BOSTON, MA (in $\mu g/m^3$)

	Boston, MA (250250042)														
	Andersen (5)					Andersen (6)					URG (7)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	25	10.6683	24.8748	4.45285	0.04	28	10.995	25.9611	4.38247	0.04000	27	10.3092	25.3771	4.29292	0.04000
Ammonium (88301)	21	0.9294	2.36412	0.07001	0.01500	25	0.90881	3.499	0.06	0.015	25	1.24094	3.4547	0.21059	0.00700
Sodium Ion (88302)	22	0.15548	0.59535	0.0189	0.02800	26	0.20178	1.02994	–.00291	0.02800	25	0.12490	0.3809	0.0219	0.01200
Potassium Ion (88303)	3	0.08220	0.0985	0.0632	0.013	1	0.16434	0.16434	0.16434	0.01300	16	0.05144	0.07342	0.023	0.00600
Nitrate (88306)	22	0.94089	4.15629	0.12428	0.008	26	0.72730	3.31728	0.09602	0.00800	25	0.90214	4.49158	0.10802	0.003
Sulfate (88403)	22	2.61927	6.60445	0.45984	0.011	26	2.6825	7.94791	0.25376	0.01100	25	3.06590	9.01206	0.45664	0.00500
Organic Carbon (88305)	5	3.82282	6.18258	2.27753	0.13400	8	4.6666	6.00087	2.53827	0.13400	9	3.46889	5.24618	2.15679	0.05900
Elemental Carbon (88307)	5	0.94296	1.85341	0.12801	0.134	8	0.9856	1.66607	0.48448	0.13400	9	0.90155	1.53401	0.33213	0.05900
Total Carbon	5	4.76578	8.03600	2.40554	NA	8	5.65214	7.21195	3.34141	NA	9	4.37044	6.78019	3.12194	NA
Aluminum (88104)	14	0.02451	0.14572	0.00151	0.00436	20	0.02292	0.15922	0.00087	0.00436	16	0.00595	0.01846	0.00094	0.00436
Antimony (88102)	17	0.00238	0.00639	0.00049	0.00592	15	0.00343	0.00579	0.00022	0.00592	21	0.00312	0.00824	0.00047	0.00592
Arsenic (88103)	19	0.00094	0.00329	0.00004	0.00099	19	0.00108	0.00235	0.00031	0.00099	13	0.00090	0.00273	0.00014	0.00099
Barium (88107)	25	0.02831	0.04840	0.00602	0.02360	28	0.03	0.05487	0.01400	0.02360	26	0.02501	0.05663	0.00424	0.02360
Bromine (88109)	25	0.00249	0.00895	0.00028	0.00080	26	0	0.01	0.00087	0.00080	27	0.00279	0.00692	0.00108	0.00080
Cadmium (88110)	9	0.00169	0.00464	0.00014	0.00421	15	0.00187	0	0.00009	0.00421	10	0.00154	0.00278	0.00047	0.00421
Calcium (88111)	25	0.05016	0.16804	0.01590	0.00139	28	0.04912	0.1648	0.01466	0.00139	27	0.03094	0.11255	0.01083	0.00139
Carbonate Carbon (88308)	0	—	—	—	0.13400	0	—	—	—	0.13400	0	—	—	—	0.05900
Cerium (88117)	16	0.01037	0.03822	0.00046	0.03450	12	0.01043	0.02062	0	0.03450	14	0.01102	0.02806	0.00104	0.03450
Cesium (88118)	14	0.00376	0.00922	0.00082	0.01480	15	0.00560	0.01054	0	0.01480	16	0.00548	0.01874	0.00066	0.01480
Chlorine (88115)	24	0.06815	1.18279	0.00004	0.00232	22	0.01822	0.07545	0.00111	0.00232	18	0.02582	0.32562	0.00066	0.00232
Chromium (88112)	16	0.00050	0.00348	0.00004	0.00063	20	0.00070	0.00424	0.00010	0.00063	15	0.00023	0.00085	0.00005	0.00063
Cobalt (88113)	1	0.00004	0.00004	0.00004	0.00056	1	0.00004	0.00004	0.00004	0.00056	0	—	—	—	0.00056
Copper (88114)	25	0.00222	0.00499	0.00068	0.00054	28	0.00244	0.00605	0.00094	0.00054	27	0.00171	0.00433	0.00047	0.00054
Europium (88121)	0	—	—	—	0.00451	1	0.00028	0.00028	0.00028	0.00451	1	0.00043	0.00043	0.00043	0.00451
Gallium (88124)	21	0.00088	0.00226	0.00018	0.00133	23	0.00075	0.00186	0.00018	0.00133	22	0.00086	0.00170	0.00014	0.00133
Iron (88126)	25	0.07623	0.25559	0.03201	0.00079	28	0.08281	0.25009	0.03017	0.00079	27	0.04838	0.09559	0.02228	0.00079
Lanthanum (88146)	10	0.01342	0.03916	0.00014	0.02790	16	0.01184	0.02467	0	0.02790	12	0.00787	0.02496	0.00259	0.02790
Lead (88128)	25	0.00343	0.00636	0.00155	0.00220	28	0.00378	0.01053	0.00062	0.00220	27	0.00337	0.00721	0.00137	0.00220

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-3 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT BOSTON, MA (in $\mu g/m^3$)

	Boston, MA (250250042)														
	Andersen (5)					Andersen (6)					URG (7)				
Parameter	N	Mean	Max	Min	MDL	N	Mean	Max	Min	MDL	N	Mean	Max	Min	MDL
Magnesium (88140)	12	0.01609	0.05037	0.00018	0.00738	20	0.01027	0.02878	0.00071	0.00738	13	0.00906	0.02392	0.00047	0.00738
Manganese (88132)	25	0.00185	0.00767	0.00004	0.00092	26	0.00175	0.00386	0.00048	0.00092	24	0.00102	0.00254	0.00005	0.00092
Mercury (88142)	15	0.00107	0.00226	0.00004	0.00175	17	0.00085	0.00196	0.00004	0.00175	9	0.00075	0.00226	0.00010	0.00175
Molybdenum (88134)	11	0.00073	0.00139	0.00018	0.00191	13	0.00072	0.00176	0.00019	0.00191	12	0.00082	0.00184	0.00010	0.00191
Nickel (88136)	25	0.00284	0.00810	0.00091	0.00050	28	0.00279	0.00823	0.00086	0.00050	27	0.00408	0.03146	0.00057	0.00050
Niobium (88147)	9	0.00040	0.00117	0.00009	0.00168	10	0.00069	0.00191	0.00010	0.00168	13	0.00043	0.00113	0.00005	0.00168
Phosphorous (88152)	0	—	—	—	0.00251	1	0	0.00067	0.00067	0.003	0	—	—	—	0.00251
Potassium (88180)	25	0.03758	0.08191	0.01023	0.00137	28	0.04	0.07416	0.01093	0.00137	27	0.03177	0.06259	0.00315	0.00137
Rubidium (88176)	11	0.00029	0.00076	0.00004	0.00087	10	0.00021	0.00051	0.00005	0.00087	6	0.00022	0.00047	0.00005	0.00087
Samarium (88162)	0	—	—	—	0.00247	1	0.00094	0.00094	0.00094	0.00247	1	0.00019	0.00019	0.00019	0.00247
Scandium (88163)	2	0.00032	0.00045	0.00019	0.00097	3	0.00036	0.00062	0.00023	0.00097	3	0.00038	0.00052	0.00023	0.00097
Selenium (88154)	19	0.00093	0.00321	0.00004	0.00085	20	0.00104	0.00255	0.00009	0.00085	20	0.00101	0.00315	0.00014	0.00085
Silicon (88165)	25	0.09181	0.51655	0.01870	0.00302	28	0.09564	0.41927	0.02776	0.00302	27	0.05312	0.13214	0.01926	0.00302
Silver (88166)	14	0.00236	0.00470	0.00023	0.00420	20	0.00173	0.00416	0.00057	0.00420	16	0.00184	0.00386	0.00014	0.00420
Sodium (88184)	22	0.17809	1.08304	0.00229	0.02050	24	0.15575	0.42302	0.00183	0.02050	25	0.10689	0.32783	0.00777	0.02050
Strontium (88168)	13	0.00066	0.00152	0.00004	0.00101	16	0.00045	0.00125	0	0.00101	16	0.00045	0.00094	0.00005	0.00101
Sulfur (88169)	25	0.93332	2.66932	0.17688	0.00265	28	1.02389	2.93344	0.22349	0.00265	27	1.00354	2.77815	0.16127	0.00265
Tantalum (88170)	24	0.00737	0.01585	0.00113	0.00784	26	0.00707	0.01392	0.00073	0.00784	26	0.007	0.01601	0.00033	0.008
Terbium (88172)	3	0.00166	0.00299	0.00014	0.00302	6	0.00083	0.00134	0.00042	0.00302	3	0.00069	0.00146	0	0.00302
Tin (88160)	25	0.00739	0.01171	0.00131	0.00717	28	0.00785	0.01296	0.00373	0.00717	27	0.00765	0.01267	0.002	0.00717
Titanium (88161)	25	0.00437	0.01595	0.00121	0.00083	27	0.00475	0.01558	0.00109	0.00083	26	0.00306	0.00588	0.001	0
Vanadium (88164)	25	0.00297	0.00913	0.00062	0.00060	28	0.00323	0.01841	0.00043	0.00060	27	0.00376	0.01955	0	0.00060
Wofram (88186)	9	0.00189	0.00416	0.00037	0.00554	11	0.00297	0.00881	0.00004	0.00554	8	0.00145	0.00235	0.00033	0.006
Yttrium (88183)	7	0.00041	0.00073	0.00009	0.00122	8	0.00033	0.00060	0.00004	0.00122	7	0.00050	0.00094	0.00010	0.00122
Zinc (88167)	25	0.00974	0.01784	0.00276	0.00058	28	0.00955	0.01855	0	0	27	0.00898	0.01709	0.00165	0
Zirconium (88185)	11	0.00147	0.01059	0.00009	0.00144	10	0	0.00165	0.00004	0.00144	14	0.00072	0.00165	0.00005	0.001

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-4. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT BRONX BOTANICAL GARDEN, NY (in $\mu g/m^3$)

	Bronx Botanical Garden, NY (360050083)														
	Andersen (5)					Met One (6)					Met One (7)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	25	12.5018	35.3799	4.13505	0.04000	36	14.3136	40.0400	4.67955	0.10400	37	15.6399	43.3368	5.34188	0.10400
Ammonium (88301)	31	1.5786	5.55365	0.11312	0.01500	35	1.54307	5.57605	0.19379	0.01700	36	1.51177	5.55544	0.15741	0.01700
Sodium Ion (88302)	31	0.16773	0.63440	0.02340	0.02800	35	0.14766	0.50700	0.00990	0.03000	36	0.16675	0.53988	0.02833	0.03000
Potassium Ion (88303)	4	0.12953	0.16503	0.09966	0.01300	5	0.13031	0.18103	0.09126	0.01400	4	0.11317	0.12365	0.10041	0.01400
Nitrate (88306)	31	1.05845	4.24427	0.12851	0.00800	35	1.12762	4.30423	0.12779	0.00800	36	1.15671	4.29506	0.12589	0.00800
Sulfate (88403)	31	4.19607	13.9566	0.60446	0.01100	35	3.90576	13.7889	0.65552	0.01200	36	3.80892	13.0896	0.58524	0.01200
Organic Carbon (88305)	6	4.20397	9.39840	2.11015	0.13400	26	4.23325	8.87590	1.67583	0.14600	26	4.20562	8.81401	1.53944	0.14600
Elemental Carbon (88307)	6	1.30671	1.87039	0.69807	0.134	26	1.31710	3.14339	0.28686	0.14600	26	1.32068	2.70728	0.27614	0.14600
Total Carbon	6	5.51068	11.2688	2.80822	NA	26	5.55035	11.0917	2.52934	NA	26	5.52630	10.6248	2.70700	NA
Aluminum (88104)	20	0.00919	0.03014	0.00125	0	20	0.01804	0.07911	0.00136	0.01088	26	0.01255	0.06945	0.00120	0.01088
Antimony (88102)	19	0.00362	0.00667	0.00031	0.01	20	0.00729	0.01789	0.00012	0.01476	30	0.00724	0.02246	0.00024	0.01476
Arsenic (88103)	18	0.00113	0.00233	0.00013	0.00099	23	0.00183	0.00412	0.00012	0.00247	28	0.00211	0.00471	0.00024	0.00247
Barium (88107)	25	0.02598	0.04590	0.00535	0.02360	34	0.06317	0.09658	0.01405	0.05876	36	0.06711	0.16887	0.00636	0.05876
Bromine (88109)	25	0.00253	0.01058	0.00017	0.00080	33	0.00275	0.00813	0.00035	0.00199	34	0.00286	0.01289	0.00024	0.00199
Cadmium (88110)	10	0.00172	0.00312	0.00022	0.00421	22	0	0.00965	0.00023	0.01050	12	0.00369	0.00751	0.00024	0.01050
Calcium (88111)	25	0.03839	0.09992	0.01449	0.00139	36	0.048	0.10880	0.01593	0.00347	37	0.04810	0.12002	0.01662	0.00347
Carbonate Carbon (88308)	0	—	—	—	0.13400	0	—	—	—	0.14600	0	—	—	—	0.14600
Cerium (88117)	12	0.00897	0.02442	0.00184	0.03450	22	0.02991	0.0857	0.00318	0.08603	19	0.03198	0.08526	0.00045	0.08603
Cesium (88118)	9	0.00655	0.01627	0.00130	0.01480	15	0.00803	0.0307	0.00034	0.03689	21	0.02048	0.04350	0.00036	0.03689
Chlorine (88115)	16	0.00975	0.08329	0.00017	0.00232	25	0.00744	0.0296	0.00023	0.00578	20	0.00729	0.04759	0.00012	0.00578
Chromium (88112)	18	0.00034	0.00109	0.00009	0.00063	13	0.00081	0.002	0	0.00159	18	0.00056	0.00131	0.00012	0.00159
Cobalt (88113)	1	0.00043	0.00043	0.00043	0.00056	3	0.00175	0.00457	0	0.00141	5	0.00077	0.00301	0.00012	0.00141
Copper (88114)	25	0.00283	0.00753	0.00086	0.00054	33	0.00309	0.00697	0	0.00135	36	0.00311	0.01203	0.00024	0.00135
Europium (88121)	0	—	—	—	0.00451	1	0.00170	0.00170	0.002	0.01124	1	0.00012	0.00012	0.00012	0.01124
Gallium (88124)	18	0.00096	0.00176	0.00052	0.00133	33	0.00218	0.00539	0	0.00331	31	0.00276	0.00529	0.00011	0.00331

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-4 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT BRONX BOTANICAL GARDEN, NY (in $\mu g/m^3$)

	Bronx Botanical Garden, NY (360050083)														
	Andersen (5)					Met One (6)					Met One (7)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Iron (88126)	25	0.09133	0.21643	0.04547	0.00079	36	0.10337	0.23509	0.03960	0	37	0.10434	0.26049	0.04346	0.00196
Lanthanum (88146)	15	0.01467	0.03096	0.00103	0.02790	25	0.02741	0.08614	0.00068	0.069	14	0.03950	0.07094	0.00381	0.06947
Lead (88128)	25	0.00419	0.00853	0.00097	0.00220	36	0.00582	0.01097	0.00103	0.01	36	0.00583	0.01209	0.00096	0.00549
Magnesium (88140)	15	0.00477	0.01027	0.00109	0.00738	14	0.02012	0.05336	0.00069	0.018	18	0.01857	0.05926	0.00290	0.01841
Manganese (88132)	24	0.00132	0.00360	0.00004	0.00092	32	0.00211	0.00596	0.00011	0	27	0.00221	0.00507	0.00012	0.00231
Mercury (88142)	14	0.00080	0.00186	0.00009	0.00175	21	0.00203	0.00773	0.00011	0.00437	18	0.00196	0.00457	0.00012	0.00437
Molybdenum (88134)	15	0.00089	0.00215	0.00021	0.00191	15	0.00182	0.00577	0.00023	0.00477	18	0.00267	0.00531	0.00060	0.00477
Nickel (88136)	25	0.01197	0.04563	0.00392	0.00050	36	0.01722	0.05531	0.00417	0.00125	37	0.02253	0.18701	0.00424	0.00125
Niobium (88147)	7	0.00076	0.00187	0.00013	0.00168	20	0.00162	0.00451	0.00023	0.00420	16	0.00174	0.00480	0.00048	0.00420
Phosphorous (88152)	1	0.00561	0.00561	0.00561	0.00251	2	0.00383	0.00471	0.00295	0.00627	0	—	—	—	0.00627
Potassium (88180)	25	0.03778	0.13856	0.00946	0.00137	36	0.03594	0.13477	0.00091	0.00341	37	0.03734	0.13799	0.00313	0.00341
Rubidium (88176)	12	0.00023	0.00065	0.00004	0.00087	10	0.00047	0.00147	0.00011	0.00217	8	0.00051	0.00108	0.00012	0.00217
Samarium (88162)	0	—	—	—	0	0	—	—	—	0.01	0	—	—	—	0.006
Scandium (88163)	5	0.00016	0.00052	0.00004	0.00097	7	0.00108	0.00233	0.00011	0.00243	8	0.00110	0.00241	0.00036	0.00243
Selenium (88154)	19	0.00076	0.00243	0.00013	0.00085	23	0.00134	0.00373	0.00011	0.00212	22	0.00131	0.00344	0.00024	0.00212
Silicon (88165)	25	0.07495	0.27206	0.02708	0.00302	36	0.09172	0.34880	0.00403	0.00753	37	0.09781	0.34934	0.02746	0.00753
Silver (88166)	10	0.00213	0.00364	0.00073	0.00420	20	0.00358	0.01155	0.00011	0.01048	24	0.00483	0.01050	0.00024	0.01048
Sodium (88184)	18	0.07192	0.28986	0.00021	0.02050	30	0.12093	0.37960	0.00632	0.05107	28	0.13348	0.49236	0.00215	0.05107
Strontium (88168)	20	0.00062	0.00184	0.00009	0.00101	12	0.00208	0.01566	0.00023	0.00251	13	0.00209	0.01520	0.00012	0.00251
Sulfur (88169)	25	1.20653	4.55104	0.20827	0.00265	36	1.34117	5.03203	0.09461	0.00662	37	1.44442	5.16369	0.26699	0.00662
Tantalum (88170)	23	0.00708	0.01545	0.00013	0.00784	35	0.02138	0.05584	0.00410	0.01954	35	0.02006	0.04054	0.00024	0.01954
Terbium (88172)	6	0.00068	0.00103	0.00034	0.00302	4	0.00165	0.00232	0.00068	0.00752	4	0.00156	0.00216	0.00109	0.00752
Tin (88160)	25	0.00758	0.01144	0.00154	0.00717	36	0.01890	0.03306	0.00856	0.01787	37	0.02168	0.03268	0.00687	0.01787
Titanium (88161)	25	0.00407	0.01006	0.00009	0.00083	36	0.00552	0.02499	0.00080	0.00208	37	0.00523	0.01265	0.00132	0.00208
Vanadium (88164)	25	0.00315	0.00948	0.00084	0.00060	31	0.00383	0.01016	0.00011	0.00150	36	0.00361	0.00894	0.00024	0.00150

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-4 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT BRONX BOTANICAL GARDEN, NY (in $\mu g/m^3$)

	Bronx Botanical Garden, NY (360050083)														
	Andersen (5)					Met One (6)					Met One (7)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Wofram (88186)	4	0.00112	0.00280	0.00030	0.00554	17	0.00617	0.01523	0.00059	0.01380	15	0.00390	0.01099	0.00036	0.01380
Yttrium (88183)	9	0.00048	0.00103	0.00004	0.00122	11	0.00126	0.00271	0.00023	0.00304	12	0.00085	0.00182	0.00012	0.00304
Zinc (88167)	25	0.02100	0.06214	0.00540	0.00058	36	0.02445	0.11719	0.00380	0.00145	37	0.02493	0.10768	0.00421	0.00145
Zirconium (88185)	8	0.00059	0.00104	0.00013	0.00144	16	0.00752	0.07212	0.00023	0.00359	12	0.0012	0.00275	0.00012	0.004

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-5. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT CHICAGO, IL (in $\mu g/m^3$)

	Chicago, IL (170310050)									
	Andersen (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	33	15.6892	33.8161	4.16636	0.04000	35	15.4128	33.1750	3.45877	0.04000
Ammonium (88301)	31	1.86837	5.17987	0.19935	0.01500	32	1.94182	5.49383	0.23548	0.00700
Sodium Ion (88302)	31	0.11463	0.32829	0.01819	0.02800	32	0.07970	0.37386	0.02158	0.01200
Potassium Ion (88303)	12	0.15903	0.55075	0.04076	0.01300	22	0.10708	0.53020	0.03710	0.00600
Nitrate (88306)	31	2.1707	8.51846	0.17247	0.00800	32	1.90254	8.80715	0.07974	0.00300
Sulfate (88403)	31	3.94298	9.26518	0.70557	0.011	32	4.06138	9.26054	0.67398	0.00500
Organic Carbon (88305)	33	4.15846	6.74101	1.40580	0.134	35	3.02862	6.39614	0.87813	0.05900
Elemental Carbon (88307)	33	1.17651	2.72362	0.35479	0.134	35	0.97663	2.02174	0.33258	0.05900
Total Carbon	33	5.33496	9.13711	1.86054	NA	35	4.00525	7.39630	1.21071	NA
Aluminum (88104)	27	0.02286	0.17028	0.0016	0	22	0.023	0.13164	0.00090	0.00436
Antimony (88102)	21	0.00279	0.00614	0.001	0.01	23	0	0.00768	0.00047	0.00592
Arsenic (88103)	25	0.00123	0.00363	0	0.00099	26	0	0.00292	0.00005	0.00099
Barium (88107)	33	0.03239	0.08344	0.00270	0.024	35	0.029	0.06951	0.00631	0.02360
Bromine (88109)	33	0.00279	0.00800	0.00046	0	34	0	0.00782	0.00037	0.00080
Cadmium (88110)	20	0.00160	0.00410	0.00014	0	21	0.00142	0.004	0.00010	0.00421
Calcium (88111)	33	0.11438	0.33888	0.02419	0	35	0.09139	0.28217	0.01956	0.00139
Carbonate Carbon (88308)	0	—	—	—	0.13400	0	—	—	—	0.05900
Cerium (88117)	15	0.00891	0.02859	0.00046	0.03450	19	0.019	0.04235	0	0.03450
Cesium (88118)	17	0.00915	0.01388	0.00004	0.01480	12	0.01	0.01621	0	0.01480
Chlorine (88115)	28	0.02258	0.32760	0.00018	0.00232	20	0.033	0.27325	0	0.00232
Chromium (88112)	30	0.00094	0.00727	0.00004	0.00063	27	0.00099	0.007	0	0.00063
Cobalt (88113)	1	0.00028	0.00028	0.00028	0.00056	0	—	—	—	0.00056
Copper (88114)	33	0.00312	0.01038	0.00075	0.00054	35	0.00286	0.00979	0	0.00054

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-5 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT CHICAGO, IL (in $\mu g/m^3$)

	Chicago, IL (170310050)									
	Andersen (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Indium (88131)	19	0.00155	0.00499	0.00009	0.00452	18	0.00185	0.00410	0	0.00452
Iridium (88133)	18	0.00129	0.00298	0.00014	0.00238	18	0.00112	0.00414	0	0.00238
Iron (88126)	33	0.16423	0.47993	0.03366	0.00079	35	0.14520	0.45138	0.03	0.0008
Lanthanum (88146)	18	0.01258	0.02731	0.00037	0.02790	23	0.01227	0.03647	0	0.0279
Lead (88128)	33	0.00864	0.02078	0.00065	0.00220	35	0.00818	0.01871	0	0.00220
Magnesium (88140)	22	0.01423	0.08985	0.00068	0.00738	25	0.00866	0.03057	0	0.00738
Manganese (88132)	33	0.00769	0.03205	0.00056	0.00092	35	0.00656	0.02807	0.00057	0.0009
Mercury (88142)	17	0.00068	0.00230	0.00004	0.00175	14	0.00096	0.00212	0.00019	0.00175
Molybdenum (88134)	16	0.00103	0.00235	0.00014	0.00191	18	0.00121	0.00565	0.00014	0.00191
Nickel (88136)	30	0.00136	0.01287	0.00018	0.00050	32	0.00175	0.01361	0.00014	0.00050
Niobium (88147)	13	0.00102	0.00232	0.00042	0.00168	11	0.00053	0.00094	0.00010	0.00168
Phosphorous (88152)	0	—	—	—	0	0	—	—	—	0.00251
Potassium (88180)	33	0.09270	0.58173	0.01370	0.00137	35	0.08544	0.56626	0.01272	0.00137
Rubidium (88176)	17	0.00040	0.00111	0.00004	0.00087	13	0.00046	0.00127	0.00005	0.00087
Samarium (88162)	2	0.00027	0.00032	0.00023	0.00247	1	0.00061	0.00061	0.00061	0.00247
Scandium (88163)	2	0.00055	0.00065	0.00046	0.00097	3	0.00031	0	0	0.001
Selenium (88154)	32	0.00146	0.00414	0.00014	0.00085	32	0.00165	0.005	0.00014	0.0009
Silicon (88165)	33	0.13614	0.66607	0.02395	0.00302	35	0.11101	0.48048	0.02072	0.00302
Silver (88166)	23	0.00197	0.00464	0.00051	0.00420	27	0.00186	0.004	0.00010	0.0042
Sodium (88184)	13	0.05172	0.14020	0.00698	0.02050	18	0.05275	0.16799	0.00434	0.02050
Strontium (88168)	27	0.00106	0.00747	0.00004	0.00101	23	0.00104	0.00608	0.00005	0.00101
Sulfur (88169)	33	1.40800	2.85499	0.30327	0.00265	35	1.47025	4.02403	0.29744	0.00265
Tantalum (88170)	30	0.00659	0.01553	0.00005	0.00784	31	0.00586	0.01234	0.00005	0.00784

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-5 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT CHICAGO, IL (in $\mu g/m^3$)

	Chicago, IL (170310050)									
	Andersen (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Terbium (88172)	12	0.00217	0.00538	0.00005	0.00302	10	0.00139	0.00396	0	0.00302
Tin (88160)	33	0.00963	0.01947	0.00341	0.00717	35	0.00985	0.01611	0	0.00717
Titanium (88161)	32	0.00456	0.01272	0.00041	0.00083	34	0.00423	0.00979	0.00108	0.0008
Vanadium (88164)	15	0.00090	0.00302	0.00004	0.00060	18	0.00080	0.00348	0.00005	0.0006
Wofram (88186)	3	0.00256	0.00322	0.00218	0.00554	4	0.00099	0.00283	0.00014	0.00554
Yttrium (88183)	11	0.00060	0.00156	0.00004	0.00122	12	0.00029	0.00057	0.00005	0.00122
Zinc (88167)	33	0.04496	0.14491	0.00238	0.00058	35	0.04233	0.13774	0.00188	0.00058
Zirconium (88185)	18	0.00061	0.00139	0.00004	0.00144	19	0.00075	0.00179	0	0.00144

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-6. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT FRESNO, CA (in $\mu g/m^3$)

	Fresno, CA (060190008)									
	Met One (5)					Andersen (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	25	10.4804	16.7861	6.08247	0.10400	25	9.11039	18.7161	5.11270	0.04000
Ammonium (88301)	20	0.73724	1.64233	0.21479	0.01700	20	0.59866	1.23704	0.16523	0.01500
Sodium Ion (88302)	20	0.25999	0.62721	0.10849	0.03000	20	0.21312	0.40863	0.07708	0.02800
Potassium Ion (88303)	7	0.20847	0.48186	0.05730	0.01400	4	0.14457	0.37056	0.06370	0.01300
Nitrate (88306)	20	1.41616	3.58681	0.36860	0.00800	20	1.27893	3.04898	0.51775	0.00800
Sulfate (88403)	20	1.77563	2.78369	0.60374	0.01200	20	1.55442	2.46932	0.53107	0.01100
Organic Carbon (88305)	24	4.71732	8.20203	2.96153	0.14600	24	4.45785	7.47617	2.78310	0.13400
Elemental Carbon (88307)	24	0.51751	0.95420	0.12369	0.14600	24	0.52888	0.99233	0.10721	0.13400
Total Carbon	24	5.23483	8.77901	3.10442	NA	24	4.98674	8.11572	2.95225	NA
Aluminum (88104)	25	0.03524	0.09154	0.00596	0.01088	25	0.03758	0.08153	0.00173	0.00436
Antimony (88102)	16	0.00731	0.01520	0.00011	0.01476	18	0.00209	0.00750	0.00026	0.00592
Arsenic (88103)	12	0.00147	0.00357	0.00035	0.0025	16	0.00066	0.00105	0.00010	0.00099
Barium (88107)	24	0.06186	0.09844	0.01265	0.05876	25	0.02736	0.06863	0.00179	0.02360
Bromine (88109)	23	0.00238	0.00460	0.00024	0.002	25	0.002	0.00579	0.00085	0.00080
Cadmium (88110)	8	0.00604	0.01011	0.00150	0.01050	11	0.001	0.00287	0.00023	0.00421
Calcium (88111)	25	0.05638	0.09913	0.02704	0.00347	25	0.0546	0.08277	0.02980	0.00139
Carbonate Carbon (88308)	0	--	--	--	0.14600	0	--	--	--	0.13400
Cerium (88117)	18	0.03076	0.06598	0.00150	0.08603	15	0.0107	0.0417	0.00029	0.03450
Cesium (88118)	14	0.01431	0.06182	0.00011	0.03689	12	0.005	0.0141	0.00005	0.01480
Chlorine (88115)	18	0.01254	0.08483	0.00139	0.00578	22	0.009	0.0674	0.00014	0.00232
Chromium (88112)	13	0.00064	0.00151	0.00011	0.00159	20	0	0.004	0	0.00063
Cobalt (88113)	0	--	--	--	0.00141	2	0.00029	0	0	0.00056
Copper (88114)	24	0.00309	0.01162	0.00080	0.00135	25	0.00351	0.013	0	0.00054
Europium (88121)	2	0.00431	0.00461	0.00401	0.01124	2	0.00163	0.002	0.0012	0.00451
Gallium (88124)	23	0.00180	0.00466	0.00023	0.00331	20	0.00091	0.003	0.00010	0.00133

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-6 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT FRESNO, CA (in $\mu g/m^3$)

	Fresno, CA (060190008)									
	Met One (5)					Andersen (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Iron (88126)	25	0.08889	0.14245	0.04555	0.00196	25	0.08660	0.14179	0.03598	0.00079
Lanthanum (88146)	14	0.03589	0.08080	0.01508	0.06947	14	0.01508	0.03212	0.00256	0.02790
Lead (88128)	25	0.00395	0.00795	0.00035	0.00549	25	0.00285	0.00991	0.00033	0.00220
Magnesium (88140)	13	0.01738	0.05313	0.00184	0.01841	17	0.01300	0.05139	0.00014	0.00738
Manganese (88132)	23	0.00205	0.00441	0.00023	0.00231	25	0.00212	0.00430	0.00020	0.00092
Mercury (88142)	10	0.00132	0.00322	0.00023	0.00437	14	0.00106	0.00192	0.00014	0.00175
Molybdenum (88134)	11	0.00194	0.00391	0.00011	0.00477	10	0.00098	0.00223	0.00010	0.00191
Nickel (88136)	24	0.00933	0.02900	0.00011	0.00125	19	0.00061	0.00140	0.00010	0.00050
Niobium (88147)	7	0.00085	0.00150	0.00011	0.00420	11	0.00078	0.00172	0.00015	0.00168
Phosphorous (88152)	0	—	—	—	0.0063	0	—	—	—	0.00627
Potassium (88180)	25	0.07233	0.41635	0.02418	0.00341	25	0.07447	0.41798	0.02572	0.00137
Rubidium (88176)	8	0.00127	0.00220	0.00011	0.00217	6	0.00022	0.00067	0.00005	0.00087
Samarium (88162)	0	—	—	—	0.00617	1	0.00020	0.00020	0.00020	0.00247
Scandium (88163)	5	0.00067	0.00115	0.00023	0.00243	4	0.00026	0.00047	0.00014	0.00097
Selenium (88154)	19	0.00191	0.00587	0.00011	0.00212	20	0.00169	0.00407	0.00030	0.00085
Silicon (88165)	25	0.18161	0.32125	0.09166	0.00753	25	0.17293	0.27442	0.05783	0.00302
Silver (88166)	13	0.00419	0.00808	0.00011	0.01048	14	0.00212	0.00577	0.00005	0.00420
Sodium (88184)	20	0.14560	0.43413	0.02406	0.05107	20	0.10596	0.45392	0.00715	0.02050
Strontium (88168)	8	0.00240	0.00518	0.00057	0.00251	15	0.00093	0.00619	0.00010	0.00101
Sulfur (88169)	25	0.58123	1.22825	0.23819	0.00662	25	0.54945	1.22229	0.21920	0.00265
Tantalum (88170)	23	0.01896	0.03739	0.00413	0.01954	24	0.00822	0.01775	0.00029	0.00784
Terbium (88172)	0	—	—	—	0.00752	7	0.00134	0.00263	0.00033	0.00302
Tin (88160)	25	0.02086	0.03423	0.01152	0.01787	25	0.00890	0.01476	0.00371	0.00717
Zinc (88167)	23	0.00615	0.06497	0.00011	0.00145	25	0.02414	0.08440	0.00733	0.00058
Zirconium (88185)	11	0.00127	0.00290	0.00011	0.00359	11	0.00100	0.00371	0	0.00144

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-7. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT HOUSTON, TX (in $\mu g/m^3$)

	Houston, TX (482011039)									
	Andersen (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	25	14.1882	23.8011	7.31156	0.04000	24	11.9393	17.6740	5.16860	0.04000
Ammonium (88301)	23	0.80390	3.32458	0.06796	0.01500	22	1.26974	2.67911	0.33023	0.00700
Sodium Ion (88302)	23	0.56265	1.78638	0.07091	0.02800	22	0.34266	1.09133	0.05162	0.01200
Potassium Ion (88303)	14	0.15733	0.33064	0.10210	0.01300	20	0.12180	0.38460	0.04691	0.00600
Nitrate (88306)	23	0.84851	2.50829	0.29509	0.00800	22	0.74211	2.95220	0.24085	0.00300
Sulfate (88403)	23	3.84944	10.6928	1.06609	0.01100	22	3.89597	7.13703	1.15058	0.00500
Organic Carbon (88305)	20	2.45022	3.65958	1.56175	0.13400	19	1.84457	3.74028	0.72794	0.05900
Elemental Carbon (88307)	20	0.39000	0.7091	0.04994	0.13400	19	0.31709	0.55817	0.07787	0.05900
Total Carbon	20	2.84022	4.05657	1.6586	NA	19	2.16165	4.24789	0.85714	NA
Aluminum (88104)	18	0.21171	1.22376	0.001	0.004	20	0.13097	0.92532	0.00023	0.00436
Antimony (88102)	15	0.00334	0.00619	0.001	0.006	16	0.00292	0.00852	0.00038	0.00592
Arsenic (88103)	20	0.00103	0.00266	0	0	20	0.00095	0.00221	0.00010	0.00099
Barium (88107)	25	0.02703	0.04721	0.00449	0.0236	23	0.02388	0.03965	0.00400	0.02360
Bromine (88109)	25	0.00428	0.01236	0.00045	0	24	0.00364	0.01304	0.00010	0.00080
Cadmium (88110)	9	0.00224	0.00665	0.00022	0.004	11	0.001	0.00475	0.00019	0.00421
Calcium (88111)	25	0.10541	0.41857	0.03970	0.001	24	0.0667	0.26045	0.02006	0.00139
Carbonate Carbon (88308)	0	—	—	—	0.13400	0	—	—	—	0.05900
Cerium (88117)	12	0.00861	0.01814	0.00221	0.03450	11	0.01	0.0282	0.00170	0.03450
Cesium (88118)	12	0.00522	0.01497	0.00009	0.01480	11	0.005	0.009	0.00023	0.01480
Chlorine (88115)	23	0.15774	1.16485	0.00077	0.00232	20	0.0988	0.60174	0.00014	0.00232
Chromium (88112)	21	0.00067	0.00160	0.00014	0.00063	19	0	0	0	0.00063
Cobalt (88113)	0	—	—	—	0.00056	1	0	0.00028	0.0003	0.00056
Copper (88114)	25	0.00228	0.01601	0.00045	0.00054	24	0.002	0.02350	0.0002	0.00054

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-7 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT HOUSTON, TX (in $\mu g/m^3$)

	Houston, TX (482011039)									
	Andersen (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Europium (88121)	0	—	—	—	0.005	0	—	—	—	0.00451
Gallium (88124)	22	0.00113	0.00193	0.00031	0.00133	19	0.001	0.00198	0.0001	0.00133
Gold (88143)	9	0.00074	0.00187	0.00004	0.00201	9	0.001	0.00268	0.0001	0.00201
Hafnium (88127)	9	0.00536	0.01202	0.00033	0.01050	14	0.003	0.00810	0.0001	0.0105
Indium (88131)	10	0.00168	0.00379	0.00039	0.00452	9	0.002	0.004	0.00066	0.00452
Iridium (88133)	19	0.00124	0.00330	0.00005	0.00238	15	0.00145	0.003	0.00023	0.00238
Iron (88126)	25	0.12309	0.71419	0.02203	0.00079	24	0.08674	0.47667	0.01385	0.00079
Lanthanum (88146)	17	0.01132	0.02234	0.00060	0.02790	10	0.01324	0.02119	0.00250	0.0279
Lead (88128)	25	0.00283	0.00703	0.00037	0.00220	24	0.00277	0.00494	0.00071	0.00220
Magnesium (88140)	11	0.02578	0.06027	0.00975	0.00738	10	0.02440	0.05015	0.00010	0.00738
Manganese (88132)	25	0.00330	0.01214	0.00041	0.00092	20	0.00233	0.00945	0.0001	0.00092
Mercury (88142)	16	0.00086	0.00201	0.00022	0.00175	10	0.00071	0.00198	0.0001	0.00175
Molybdenum (88134)	10	0.00102	0.00275	0.00009	0.00191	15	0.00122	0.00410	0	0.00191
Nickel (88136)	25	0.00155	0.00306	0.00030	0.00050	24	0.00239	0.00866	0.0002	0.00050
Niobium (88147)	14	0.00061	0.00219	0.00009	0.00168	12	0.00089	0.00198	0.00010	0.00168
Phosphorous (88152)	0	—	—	—	0.003	0	—	—	—	0.00251
Potassium (88180)	25	0.12207	0.29581	0.03480	0.00137	24	0.10719	0.34553	0.03123	0.00137
Rubidium (88176)	12	0.00049	0.00114	0.00010	0.00087	13	0.00054	0.002	0	0.00087
Samarium (88162)	0	—	—	—	0.002	0	—	—	—	0.00247
Scandium (88163)	2	0.00012	0.00013	0.00010	0.00097	1	0.00014	0	0.0001	0.00097
Selenium (88154)	20	0.00078	0.00161	0.00009	0.00085	18	0.00060	0.002	0	0.00085
Silicon (88165)	25	0.36800	2.30769	0.03828	0.00302	24	0.27270	1.72228	0.02576	0.00302
Silver (88166)	13	0.00225	0.00504	0.00041	0.00420	16	0.00188	0.004	0	0.00420

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-7 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT HOUSTON, TX (in $\mu g/m^3$)

	Houston, TX (482011039)									
	Andersen (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Terbium (88172)	3	0.00107	0.00135	0.00055	0.00302	4	0.00124	0.00307	0.0001	0.00302
Tin (88160)	25	0.00817	0.01260	0.00372	0.00717	24	0.00737	0.01304	0.00217	0.00717
Titanium (88161)	25	0.01120	0.06742	0.00197	0.00083	24	0.00785	0.04432	0.00141	0.0008
Vanadium (88164)	24	0.00311	0.00814	0.00074	0.00060	23	0.00313	0.00815	0.00014	0.0006
Wofram (88186)	10	0.00282	0.00591	0.00020	0.00554	13	0.00281	0.00669	0.00066	0.00554
Yttrium (88183)	5	0.00043	0.00086	0.00009	0.00122	7	0.00040	0.00085	0.00005	0.00122
Zinc (88167)	25	0.00657	0.02032	0.00060	0.00058	24	0.00560	0.01926	0.00033	0.0006
Zirconium (88185)	11	0.00072	0.00196	0.00014	0.00144	10	0	0.00113	0.00010	0.00144

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-8. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT LEWIS, FL (in $\mu g/m^3$)

	Lewis, FL (120571075)									
	MetOne (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	34	14.6122	35.7215	6.96379	0.10400	32	12.7384	32.0973	6.50163	0.04000
Ammonium (88301)	35	1.10695	3.45870	0.03176	0.01700	33	1.68153	3.80897	0.53032	0.00700
Sodium Ion (88302)	34	0.31849	1.14027	0.11746	0.03000	33	0.15983	0.45793	0.02519	0.01200
Potassium Ion (88303)	11	0.18475	0.9625	0.04954	0.01400	27	0.10138	0.91296	0.03434	0.00600
Nitrate (88306)	35	0.56896	1.48237	0.03221	0.00800	33	0.50700	1.46923	0.15998	0.00300
Sulfate (88403)	35	4.43481	12.1344	1.71479	0.01200	33	4.47554	9.88074	1.86214	0.00500
Organic Carbon (88305)	37	3.46126	8.02624	1.42506	0.146	35	2.50716	6.87372	1.27824	0.05900
Elemental Carbon (88307)	37	0.55103	1.27681	0.14487	0.146	35	0.41304	0.72688	0.16214	0.05900
Total Carbon	37	4.01229	8.56808	1.94724	NA	35	2.92020	7.04766	1.45098	NA
Aluminum (88104)	30	0.03213	0.23294	0.00168	0.01088	23	0.02566	0.22656	0.00259	0.00436
Antimony (88102)	26	0.00484	0.01885	0.0003	0.01476	16	0.00348	0.00896	0.00099	0.00592
Arsenic (88103)	26	0.00247	0.00569	0.00023	0.00247	24	0.00162	0.00527	0.00028	0.00099
Barium (88107)	34	0.05813	0.12715	0.00716	0.05876	32	0.02719	0.07907	0.00099	0.02360
Bromine (88109)	33	0.00317	0.00817	0.00032	0.00199	32	0.00342	0.00947	0.00071	0.00080
Cadmium (88110)	20	0.00322	0.01160	0.00023	0.0105	16	0.00221	0.00457	0.00047	0.00421
Calcium (88111)	34	0.07670	0.13849	0.03215	0.00347	32	0.05308	0.08769	0.019	0.00139
Carbonate Carbon (88308)	0	—	—	—	0.14600	0	—	—	—	0.059
Cerium (88117)	21	0.02962	0.07865	0.00089	0.08603	15	0.00855	0.01756	0.00104	0.0345
Cesium (88118)	15	0.01518	0.03517	0.00078	0.03689	13	0.00447	0.01410	0.00010	0.0148
Chlorine (88115)	21	0.03747	0.29395	0.00134	0.00578	18	0.01529	0.13924	0.00047	0.00232
Chromium (88112)	14	0.00076	0.00156	0.00011	0.00159	20	0.0003	0.00137	0.00005	0.00063
Cobalt (88113)	3	0.00045	0.00101	0.00011	0.00141	3	0.0002	0.00033	0.00005	0.00056
Copper (88114)	30	0.00221	0.01852	0.00011	0.00135	32	0.00172	0.01761	0.00028	0.00054
Europium (88121)	1	0.00023	0.00023	0.00023	0.01124	1	0.0007	0.0007	0.00066	0.00451
Iron (88126)	34	0.05842	0.17794	0.02057	0.00196	32	0.04877	0.16403	0.01763	0.00079
Lanthanum (88146)	16	0.03758	0.07532	0.00011	0.06947	23	0.01039	0.02383	0.00080	0.02790
Gallium (88124)	31	0.00261	0.00605	0.00045	0.00331	29	0.00105	0.00311	0.00023	0.00133

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-8 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT LEWIS, FL (in $\mu g/m^3$)

	Lewis, FL (120571075)									
	MetOne (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Lead (88128)	34	0.00557	0.02700	0.00122	0.00549	32	0.00438	0.02519	0.00080	0.00220
Magnesium (88140)	14	0.02254	0.04844	0.00805	0.01841	16	0.00912	0.02081	0.00160	0.00738
Manganese (88132)	28	0.00194	0.00470	0.00011	0.00231	31	0.00139	0.00325	0.00028	0.00092
Mercury (88142)	23	0.00206	0.00513	0.00011	0.00437	14	0.00053	0.00151	0.00005	0.00175
Molybdenum (88134)	15	0.00210	0.00537	0.00023	0.00477	18	0.00088	0.00193	0.00005	0.00191
Nickel (88136)	31	0.00554	0.05896	0.00011	0.00125	30	0.00203	0.01554	0.00028	0.00050
Niobium (88147)	16	0.00136	0.00274	0.00011	0.00420	14	0.00062	0.00151	0.00005	0.00168
Phosphorous (88152)	0	—	—	—	0.00617	0	—	—	—	0.00251
Potassium (88180)	34	0.07957	0.89347	0.01909	0.00341	32	0.08082	0.89697	0.02027	0.00137
Rubidium (88176)	13	0.00079	0.00246	0.00011	0.00217	14	0.00038	0.00108	0.00005	0.00087
Samarium (88162)	3	0.00052	0.00112	0.00011	0.00617	0	—	—	—	0.00247
Scandium (88163)	7	0.00067	0.00145	0.00011	0.00243	2	0.00014	0.00014	0.00014	0.00097
Selenium (88154)	22	0.00110	0.00268	0.00011	0.00212	24	0.00058	0.00162	0.00005	0.00085
Silicon (88165)	34	0.14964	0.58150	0.05025	0.00753	32	0.12164	0.49329	0.03804	0.00302
Silver (88166)	22	0.00449	0.00902	0.00045	0.01048	24	0.00184	0.00444	0.00014	0.00420
Sodium (88184)	31	0.17202	0.46907	0.00903	0.05107	29	0.11949	0.41652	0.01269	0.02050
Strontium (88168)	13	0.00189	0.01060	0.00022	0.00251	16	0.00140	0.01224	0.00019	0.00101
Sulfur (88169)	34	1.57397	4.30113	0.64698	0.00662	32	1.45578	3.33434	0.67286	0.00265
Tantalum (88170)	34	0.02010	0.04911	0.00136	0.01954	31	0.00867	0.01990	0.00146	0.00784
Terbium (88172)	3	0.00052	0.00101	0.00011	0.00752	4	0.00065	0.00132	0.00005	0.00302
Tin (88160)	34	0.01868	0.03130	0.00291	0.01787	32	0.00838	0.01551	0.00198	0.00717
Zinc (88167)	33	0.00568	0.01264	0.00033	0.00145	32	0.00625	0.01263	0.00165	0.00058

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-9. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT PHILADELPHIA, PA (in $\mu g/m^3$)

	Philadelphia, PA (421010004)									
	Andersen (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	37	14.7207	43.1932	3.75162	0.04000	37	13.3653	39.0898	3.66728	0.04000
Ammonium (88301)	38	1.60087	4.62702	0.20477	0.01500	38	1.89732	5.62864	0.36506	0.00700
Sodium Ion (88302)	37	0.18607	0.73099	0.03102	0.02800	38	0.07003	0.19235	0.01947	0.01200
Potassium Ion (88303)	7	0.18764	0.37397	0.06598	0.01300	21	0.07957	0.27421	0.03753	0.00600
Nitrate (88306)	38	1.38838	3.69438	0.21257	0.00800	38	1.23693	4.35089	0.14346	0.00300
Sulfate (88403)	38	4.34017	13.8852	0.9297	0.011	38	4.43424	14.2670	0.86455	0.00500
Organic Carbon (88305)	37	4.17646	10.5209	1.41139	0.134	37	3.20367	8.35341	0.97135	0.05900
Elemental Carbon (88307)	37	0.83466	1.89372	0.18759	0.13400	37	0.66304	1.53473	0.26893	0.05900
Total Carbon	37	5.01112	11.2164	1.59898	NA	37	3.86671	8.75990	1.24027	NA
Aluminum (88104)	30	0.01787	0.07417	0.00163	0.0044	21	0.0101	0.03047	0.00010	0.00436
Antimony (88102)	26	0.00354	0.00813	0.00049	0.0059	21	0.003	0.00725	0.00014	0.00592
Arsenic (88103)	30	0.00101	0.00241	0.0001	0.001	29	0.00111	0.003	0.00005	0.00099
Barium (88107)	37	0.02638	0.04363	0.00443	0.0236	37	0.02628	0.047	0.00212	0.02360
Bromine (88109)	37	0.00340	0.00828	0.00070	0.00080	37	0.00334	0.00810	0.0003	0.00080
Cadmium (88110)	18	0.00180	0.00483	0	0.0042	20	0.002	0.00414	0.00005	0.00421
Calcium (88111)	37	0.05694	0.15631	0.01194	0.0014	37	0.0342	0.10905	0.00946	0.00139
Carbonate Carbon (88308)	0	—	—	—	0.134	0	—	—	—	0.05900
Cerium (88117)	18	0.01002	0.01922	0.00071	0.03450	16	0.01305	0.02825	0.00179	0.03450
Cesium (88118)	17	0.00563	0.01665	0.00010	0.0148	17	0.00648	0.01738	0.00080	0.01480
Chlorine (88115)	27	0.00771	0.06483	0.00010	0.0023	16	0.00495	0.01695	0.00033	0.00232
Chromium (88112)	28	0.00107	0.00386	0.00004	0.001	27	0.00055	0.00160	0.00005	0.00063
Cobalt (88113)	0	—	—	—	0.001	2	0.00016	0.00028	0.00005	0.00056
Copper (88114)	37	0.00453	0.01252	0.00034	0.001	37	0.003	0.00579	0.00080	0.00054
Europium (88121)	2	0.00044	0.00052	0.00035	0.0045	0	—	—	—	0.00451
Gallium (88124)	29	0.00091	0.00228	0.00010	0.0013	31	0	0.00212	0.00005	0.00133

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-9 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT PHILADELPHIA, PA (in $\mu g/m^3$)

	Philadelphia, PA (421010004)									
	Andersen (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Iron (88126)	37	0.10270	0.22846	0.02972	0.00079	37	0.0662	0.15637	0.02608	0.00079
Lanthanum (88146)	18	0.00968	0.02769	0.00023	0.02790	17	0.00693	0.0167	0.00037	0.02790
Lead (88128)	37	0.00556	0.01189	0.00161	0.00220	37	0.00513	0.0106	0.00165	0.00220
Magnesium (88140)	18	0.00773	0.01847	0.00053	0.00738	17	0.00738	0.0208	0.00010	0.00738
Manganese (88132)	37	0.00268	0.00700	0.00029	0.00092	35	0.00171	0.00372	0.00033	0.00092
Mercury (88142)	19	0.00092	0.00232	0.00005	0.00175	20	0.00084	0.00155	0.00005	0.00175
Molybdenum (88134)	22	0.00120	0.00289	0.00004	0.00191	16	0.00098	0.00325	0.00005	0.00191
Nickel (88136)	35	0.00441	0.02189	0.00005	0.00050	37	0.00413	0.01714	0.00005	0.00050
Niobium (88147)	18	0.00078	0.00167	0.00010	0.00168	24	0	0.002	0.00005	0.00168
Phosphorous (88152)	0	—	—	—	0.0025	0	—	—	—	0.00251
Potassium (88180)	37	0.05987	0.29381	0.00959	0.00137	37	0.0506	0.30154	0.00588	0.00137
Rubidium (88176)	15	0.00037	0.00085	0.00005	0.00087	12	0.00050	0.001	0.0001	0.00087
Samarium (88162)	0	—	—	—	0.00247	1	0.00010	0	0.0001	0.00247
Scandium (88163)	6	0.00033	0.00057	0.00010	0.00097	12	0.00035	0.00071	0	0.001
Selenium (88154)	31	0.00113	0.00330	0.00010	0.00085	32	0.00108	0.003	0.00005	0.0009
Silicon (88165)	37	0.11766	0.41821	0.02285	0.00302	37	0.07632	0.32847	0.01992	0.00302
Silver (88166)	23	0.00186	0.00503	0.00015	0.00420	25	0.00201	0.00424	0.00028	0.00420
Sodium (88184)	23	0.06349	0.19720	0.00069	0.02050	27	0.04800	0.17957	0.00410	0.02050
Strontium (88168)	26	0.00091	0.00515	0.00010	0.00101	22	0.00097	0.00447	0.00010	0.00101
Sulfur (88169)	37	1.50616	5.23630	0.32794	0.00265	37	1.49876	5.21337	0.30435	0.00265
Tantalum (88170)	35	0.00715	0.01651	0.00019	0.00784	35	0.00765	0.01624	0.00014	0.00784
Terbium (88172)	4	0.00087	0.00266	0.00005	0.00302	4	0.00072	0.00108	0.00019	0.00302
Tin (88160)	37	0.00913	0.01527	0.00189	0.00717	37	0.00864	0.01450	0.00146	0.00717

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-9 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT PHILADELPHIA, PA (in $\mu g/m^3$)

	Philadelphia, PA (421010004)									
	Andersen (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Titanium (88161)	37	0.00590	0.01489	0.00124	0.00083	36	0.00373	0.01191	0.00099	0.00083
Vanadium (88164)	30	0.00358	0.01202	0.00005	0.00060	29	0.00354	0.01140	0.00019	0.00060
Wofram (88186)	10	0.00210	0.00355	0.00043	0.00554	6	0.00254	0.00452	0.00058	0.00554
Yttrium (88183)	11	0.00039	0.00123	0.00005	0.00122	12	0.00036	0.00122	0.00010	0.00122
Zinc (88167)	37	0.01588	0.04560	0.00190	0.00058	37	0.01372	0.04045	0.00113	0.0006

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-10. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT PHOENIX, AZ (in $\mu g/m^3$)

	Phoenix, AZ (040139997)														
	URG (5)					URG (6)					MetOne (7)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	31	7.90504	14.0493	3.75000	0.04000	31	7.92975	13.054	3.54388	0.04000	28	9.77348	15.1378	5.53264	0.10400
Ammonium (88301)	33	0.40373	0.73590	0.14333	0.00700	31	0.3968	0.7288	0.14380	0.00700	30	0.39832	0.76107	0.12791	0.01700
Sodium Ion (88302)	33	0.14516	0.37517	0.05387	0.01200	31	0.1347	0.35604	0.03440	0.01200	30	0.17580	0.39862	0.05086	0.03000
Potassium Ion (88303)	28	0.08571	0.22077	0.0381	0.00600	25	0.088	0.2091	0.03403	0.00600	8	0.14045	0.19476	0.07246	0.01400
Nitrate (88306)	33	0.44349	1.53715	0.18007	0.003	31	0.50471	2.1011	0.16563	0.00300	30	0.60367	2.02319	0.20640	0.00800
Sulfate (88403)	33	1.22486	2.16162	0.46270	0.005	31	1.19059	2.0969	0.4645	0.00500	30	1.26266	2.27230	0.57097	0.01200
Organic Carbon (88305)	33	3.32080	7.27046	1.44303	0.059	30	3.49197	6.43129	1.74887	0.05900	25	4.51393	8.86971	2.67486	0.14600
Elemental Carbon (88307)	33	0.61949	1.37649	0.14996	0.05900	30	0.6383	1.3136	0.12856	0.05900	25	0.74349	1.66706	0.23419	0.14600
Total Carbon	33	3.94028	8.64695	1.59299	NA	30	4.1302	7.7449	1.87742	NA	25	5.25742	10.2845	3.12239	NA
Aluminum (88104)	31	0.12266	0.50254	0.00833	0.004	31	0.11959	0.5118	0.01	0.00436	28	0.17061	0.59217	0.00815	0.01088
Antimony (88102)	17	0.00275	0.00501	0.00019	0.00592	18	0.00317	0.01	0	0.00592	14	0.00492	0.01123	0.00011	0.01476
Arsenic (88103)	20	0.00092	0.00268	0.00010	0.00099	20	0.00079	0	0.00005	0	20	0.00158	0.00559	0.00023	0.00247
Barium (88107)	31	0.02565	0.06113	0.00085	0.02360	30	0.02804	0.049	0.00938	0.02360	27	0.06177	0.12311	0.00034	0.05876
Bromine (88109)	31	0.00392	0.00825	0.00108	0.00080	31	0.00407	0.00811	0	0.00080	28	0.00322	0.00638	0.00034	0.00199
Cadmium (88110)	13	0.00179	0.00433	0.00010	0.00421	15	0.00124	0.00311	0	0.00421	14	0.00527	0.01050	0.00034	0.01050
Calcium (88111)	31	0.13114	0.33690	0.02783	0.00139	31	0.13075	0.36095	0.031	0.00139	28	0.18478	0.42720	0.03091	0.00347
Carbonate Carbon (88308)	0	—	—	—	0.05900	0	—	—	—	0.05900	0	—	—	—	0.14600
Cerium (88117)	17	0.00735	0.02552	0.00028	0.03450	18	0.01519	0.03043	0	0.03450	14	0.02084	0.04110	0.00023	0.08603
Cesium (88118)	14	0.00607	0.01577	0.00066	0.01480	12	0.01011	0.01709	0	0.01480	14	0.01244	0.02864	0.00160	0.03689
Chlorine (88115)	31	0.06039	0.22815	0.00160	0.00232	31	0.05382	0.26950	0	0.00232	26	0.04426	0.16390	0.00219	0.00578
Chromium (88112)	20	0.00031	0.00080	0.00005	0.00063	22	0.00030	0.00061	0	0	14	0.00068	0.00160	0.00011	0.00159
Cobalt (88113)	1	0.00019	0.00019	0.00019	0.00056	1	0.00005	0.00005	0.00005	0	0	—	—	—	0.00141
Copper (88114)	31	0.00256	0.00725	0.00043	0.00054	31	0.00286	0.00782	0.00094	0.00054	28	0.00334	0.00813	0.00088	0.00135
Europium (88121)	3	0.00024	0.00033	0.00014	0.00451	1	0.00189	0.00189	0.00189	0.00451	3	0.00202	0.00379	0.00069	0.01124
Gallium (88124)	29	0.00094	0.00222	0.00005	0.00133	27	0.00074	0.00221	0.00010	0.00133	24	0.00239	0.00551	0.00045	0.00331
Gold (88143)	13	0.00084	0.00226	0.00010	0.00201	12	0.00082	0.00198	0.00010	0.00201	16	0.00155	0.00637	0.00011	0.00501
Lanthanum (88146)	16	0.01360	0.02868	0.00071	0.02790	13	0.00990	0.02482	0.00033	0.02790	10	0.03140	0.06490	0.00239	0.06947
Lead (88128)	31	0.00312	0.00636	0.00160	0.00220	31	0.00346	0.00806	0.00005	0.00220	25	0.00493	0.01395	0.00145	0.00549

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-10 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT PHOENIX, AZ (in $\mu g/m^3$)

	Phoenix, AZ (040139997)														
	URG (5)					URG (6)					MetOne (7)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Magnesium (88140)	20	0.01391	0.03612	0.00221	0.00738	12	0.01396	0.03523	0.00410	0.00738	16	0.02083	0.05143	0.00139	0.01841
Manganese (88132)	31	0.00310	0.00777	0.00085	0.00092	31	0.00333	0.00730	0.00099	0.00092	26	0.00435	0.00887	0.00045	0.00231
Mercury (88142)	18	0.00085	0.00184	0.00010	0.00175	17	0.00077	0.00146	0.00010	0.00175	19	0.00209	0.00525	0.00023	0.004
Molybdenum (88134)	18	0.00073	0.00212	0.00005	0.00191	10	0.00108	0.00311	0.00019	0.00191	9	0.00200	0.00454	0.00034	0.00477
Nickel (88136)	29	0.00045	0.00132	0.00005	0.00050	26	0.00050	0.00118	0.00005	0.00050	25	0.00370	0.04763	0.00023	0.00125
Niobium (88147)	9	0.00069	0.00113	0.00014	0.00168	14	0.00066	0.00184	0.00010	0.00168	15	0.00146	0.00307	0.00055	0.00420
Phosphorous (88152)	0	—	—	—	0.00251	1	0.00099	0.00099	0.00099	0.00251	3	0.00291	0.00477	0.00089	0.00627
Potassium (88180)	31	0.10542	0.23577	0.02623	0.00137	31	0.10696	0.28344	0.02926	0.00137	28	0.11410	0.27725	0.02838	0.00341
Rubidium (88176)	17	0.00051	0.00118	0.00010	0.00087	16	0.00041	0.00108	0.00005	0.00087	13	0.00094	0.00323	0.00011	0.00217
Samarium (88162)	1	0.00023	0.00023	0.00023	0.00247	2	0.00069	0.00128	0.00010	0.00247	1	0	0.00114	0.00114	0.00617
Scandium (88163)	5	0.00016	0.00028	0.00010	0.00097	1	0.00023	0.00023	0.00023	0.00097	6	0.00057	0.00103	0.00023	0.00243
Selenium (88154)	18	0.00042	0.00090	0.00005	0.00085	19	0.00041	0.00104	0.00005	0.00085	7	0.00078	0.00125	0.00011	0.00212
Silicon (88165)	31	0.34686	1.16593	0.07738	0.00302	31	0.34468	1.23567	0.08035	0.00302	28	0.47685	1.41560	0.07708	0.00753
Silver (88166)	16	0.00197	0.00499	0.00014	0.00420	19	0.00146	0.00513	0.00023	0.00420	20	0.00502	0.00958	0.00023	0.01048
Sodium (88184)	14	0.07840	0.17328	0.00608	0.02050	14	0.07773	0.17248	0.00023	0.02050	13	0.09012	0.25108	0.01047	0.05107
Strontium (88168)	29	0.00202	0.01026	0.00010	0.00101	29	0.00233	0.00923	0.00005	0.00101	22	0.00194	0.01174	0.00011	0.00251
Sulfur (88169)	31	0.39759	0.68547	0.14079	0.00265	31	0.40492	0.71819	0.14642	0.00265	28	0.39844	0.72924	0.20112	0.00662
Tantalum (88170)	29	0.00822	0.01644	0.00071	0.00784	31	0.00699	0.01423	0.00090	0.00784	27	0.01807	0.04400	0.00295	0.01954
Terbium (88172)	9	0.00116	0.00334	0.00023	0.00302	12	0.00119	0.00416	0.00005	0.00302	1	0.00205	0.00205	0.00205	0.00752
Tin (88160)	31	0.00683	0.01319	0.00221	0.00717	31	0.00776	0.01413	0	0.00717	28	0.01739	0.02781	0.00544	0.01787
Titanium (88161)	31	0.00879	0.02152	0.00212	0.00083	31	0.00860	0.02388	0	0.00083	28	0.01185	0.02736	0.00045	0.00208
Vanadium (88164)	17	0.00063	0.00184	0.00010	0.00060	14	0.00076	0.00146	0	0.00060	6	0.00065	0.00228	0.00011	0.00150
Wofram (88186)	12	0.00316	0.00800	0.00090	0.00554	18	0.00215	0.00504	0	0.01	19	0.00576	0.01920	0.00011	0.01380
Yttrium (88183)	9	0.00041	0.00123	0.00010	0.00122	12	0.00064	0.00179	0.00005	0	8	0.00061	0.00102	0.00023	0.00304
Zinc (88167)	31	0.00544	0.01262	0.00047	0.00058	31	0.00658	0.01861	0.00033	0	28	0.00552	0.01601	0.00023	0.001
Zirconium (88185)	16	0.00060	0.00141	0.00010	0.00144	17	0.00060	0.00127	0.00010	0	14	0	0.00324	0.00011	0.004

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-11. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT PORTLAND, OR (in $\mu g/m^3$)

	Portland, OR (410510080)									
	Andersen (5)					MetOne (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	37	10.3055	61.0285	3.14779	0.04000	33	11.1644	39.9458	4.27350	0.104
Ammonium (88301)	33	0.34810	1.09408	0.04244	0.01500	26	0.43002	1.17956	0.05355	0.017
Sodium Ion (88302)	35	0.24866	0.71557	0.09839	0.02800	31	0.26273	0.56101	0.11850	0.03
Potassium Ion (88303)	12	1.46477	16.2389	0.06684	0.01300	9	1.21993	9.6779	0.10714	0.014
Nitrate (88306)	35	0.78165	3.13843	0.19594	0.00800	31	0.86400	3.39907	0.26691	0.008
Sulfate (88403)	35	1.81885	19.2367	0.33448	0.01100	31	1.80200	12.7212	0.40446	0.012
Organic Carbon (88305)	29	4.62628	11.8141	1.94468	0.13400	26	4.81523	10.6439	2.12891	0.14600
Elemental Carbon (88307)	29	0.69398	1.62679	0.19218	0.13400	26	0.71215	2.12536	0.13641	0.146
Total Carbon	29	5.32026	12.4246	2.18600	NA	26	5.52738	11.4524	2.26533	NA
Aluminum (88104)	34	0.04028	0.75854	0.00277	0.0044	29	0.04276	0.66708	0.00083	0.01088
Antimony (88102)	25	0.0046	0.054	0.00032	0.00592	19	0.00775	0.02758	0.00011	0.01476
Arsenic (88103)	33	0.00161	0.00967	0	0.00099	26	0.00256	0.00881	0.00035	0.00247
Barium (88107)	36	0.05910	1.29088	0.00587	0.02360	32	0.0773	0.70015	0.01070	0.05876
Bromine (88109)	34	0.00170	0.00585	0.0002	0.00080	25	0.00221	0.00611	0.00024	0.00199
Cadmium (88110)	22	0.00252	0.00466	0.00036	0.0042	18	0.004	0.01283	0.00024	0.01050
Calcium (88111)	36	0.03677	0.10647	0.00857	0.0014	33	0.04712	0.33552	0.01378	0.00347
Carbonate Carbon (88308)	0	—	—	—	0.13400	0	—	—	—	0.146
Cerium (88117)	17	0.00733	0.02011	0.00064	0.03450	18	0.0199	0.0547	0.00270	0.08603
Cesium (88118)	21	0.00601	0.01333	0.00071	0.01480	13	0.0161	0.03794	0.00107	0.03689
Chlorine (88115)	35	0.13788	3.28566	0.00087	0.00232	29	0.10324	1.90968	0.0009	0.00578
Chromium (88112)	27	0.00126	0.00445	0.00009	0.00063	24	0.00131	0.007	0.0001	0.00159
Cobalt (88113)	1	0.00018	0.00018	0.00018	0.00056	0	—	—	—	0.00141
Copper (88114)	37	0.01073	0.26328	0.00050	0.00054	29	0.00999	0.16719	0.0006	0.00135
Europium (88121)	1	0.00094	0.00094	0.00094	0.00451	2	0.00053	0	0.0004	0.01124
Gallium (88124)	25	0.00120	0.00303	0.00014	0.00133	25	0.00240	0.006	0.00011	0.00331
Iron (88126)	37	0.06334	0.24453	0.00726	0.00079	33	0.06408	0.22954	0.00624	0.00196
Lanthanum (88146)	21	0.00934	0.02025	0.00045	0.02790	23	0.01952	0.04803	0.00011	0.06947
Lead (88128)	36	0.01578	0.36292	0.00063	0.00220	32	0.01447	0.22048	0.00035	0.00549

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-11 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT PORTLAND, OR (in $\mu g/m^3$)

	Portland, OR (410510080)									
	Andersen (5)					MetOne (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Magnesium (88140)	24	0.04275	0.86179	0.00073	0.00738	22	0.04338	0.66761	0.00035	0.01841
Manganese (88132)	36	0.00389	0.01948	0.00041	0.00092	30	0.00432	0.01992	0.00024	0.00231
Mercury (88142)	21	0.00070	0.00157	0.00004	0.00175	15	0.00264	0.00475	0.00071	0.00437
Molybdenum (88134)	24	0.00080	0.00261	0.00004	0.00191	16	0.00216	0.00615	0.00011	0.00477
Nickel (88136)	36	0.00184	0.00617	0.00022	0.00050	32	0.00931	0.04438	0.00024	0.00125
Niobium (88147)	19	0.00063	0.00224	0.00004	0.00168	14	0.00205	0.00482	0.00011	0.00420
Phosphorous (88152)	0	—	—	—	0.0025	0	—	—	—	0.00627
Potassium (88180)	37	0.42873	13.7123	0.01234	0.00137	33	0.32038	8.76512	0.00603	0.00341
Rubidium (88176)	16	0.00034	0.00071	0.00004	0.00087	10	0.00118	0.00225	0.00012	0.00217
Samarium (88162)	1	0.00018	0.00018	0.00018	0.00247	1	0.00095	0.00095	0.00095	0.00617
Scandium (88163)	8	0.00024	0.00073	0.00004	0.00097	6	0.00095	0.00201	0.00035	0.00243
Selenium (88154)	23	0.00065	0.00234	0.00009	0.00085	19	0.00134	0.00294	0.00012	0.00212
Silicon (88165)	36	0.07451	0.27605	0.01087	0.00302	32	0.07510	0.25940	0.00978	0.00753
Silver (88166)	24	0.00204	0.00462	0.00013	0.00420	20	0.00408	0.01006	0.00024	0.01048
Sodium (88184)	36	0.15270	0.46990	0.01399	0.02050	32	0.19653	0.51494	0.02382	0.05107
Strontium (88168)	24	0.01156	0.25846	0.00004	0.00101	13	0.01220	0.14383	0.00011	0.00251
Sulfur (88169)	37	0.61466	6.08678	0.13558	0.00265	33	0.61745	4.09887	0.13664	0.00662
Tantalum (88170)	33	0.00763	0.01849	0.00023	0.00784	31	0.02026	0.04933	0.00188	0.01954
Terbium (88172)	7	0.00046	0.00141	0.00004	0.00302	2	0.00035	0.00058	0.00011	0.00752
Tin (88160)	37	0.00892	0.01603	0.00280	0.00717	33	0.01832	0.02711	0.00613	0.01787
Titanium (88161)	36	0.00930	0.18710	0.00042	0.00083	33	0.00832	0.11765	0.00118	0.00208
Vanadium (88164)	27	0.00121	0.00462	0.00009	0.00060	16	0.00143	0.00417	0.00012	0.00150
Wofram (88186)	9	0.00228	0.00478	0.00018	0.00554	15	0.00621	0.01540	0.00105	0.01380
Yttrium (88183)	14	0.00039	0.00081	0.00004	0.00122	10	0.00126	0.00247	0.00011	0.00304
Zinc (88167)	37	0.01258	0.10924	0.00123	0.00058	32	0.00985	0.05763	0.0001	0.00145

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-12. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT SALT LAKE CITY, UT (in $\mu g/m^3$)

	Salt Lake City, UT (490353006)									
	Met One (5)					Andersen (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	33	8.60543	23.3333	3.83420	0.10400	35	8.23244	24.7030	3.84018	0.04
Ammonium (88301)	30	0.41737	2.75328	0.07100	0.01700	31	0.38550	2.03933	0.04100	0.01500
Sodium Ion (88302)	30	0.10337	0.3047	0.02387	0.03000	32	0.09332	0.20916	0.01739	0.02800
Potassium Ion (88303)	5	0.26540	0.42790	0.1226	0.01400	6	0.30784	0.59569	0.12049	0.01300
Nitrate (88306)	30	1.07903	8.80071	0.18073	0.00800	32	0.91318	7.00222	0.13652	0.00800
Sulfate (88403)	30	0.93003	1.78049	0.45655	0.012	32	0.87752	1.58005	0.36993	0.01100
Organic Carbon (88305)	22	4.30412	6.37857	2.35082	0.146	24	4.09455	9.02102	2.13018	0.13400
Elemental Carbon (88307)	22	0.69880	1.39183	0.24718	0.146	24	0.65719	1.71189	0.16748	0.13400
Total Carbon	22	5.00292	7.53968	2.70476	NA	24	4.75174	9.94744	2.29766	NA
Aluminum (88104)	33	0.07483	0.39225	0.00071	0.01088	34	0.07018	0.30535	0.00228	0.00436
Antimony (88102)	20	0.00626	0.01536	0.00011	0.01476	22	0.0033	0.00780	0.00022	0.0059
Arsenic (88103)	24	0.00210	0.00714	0.00011	0.00247	29	0.0012	0.00615	0.00013	0.00099
Barium (88107)	32	0.06479	0.16508	0.00687	0.05876	35	0.03573	0.12361	0.00095	0.0236
Bromine (88109)	31	0.00310	0.01029	0.00024	0.00199	34	0.0038	0.01245	0.00080	0.00080
Cadmium (88110)	17	0.00546	0.01265	0.00011	0.01050	17	0.00164	0.0039	0.00004	0.00421
Calcium (88111)	33	0.16492	0.58893	0.03144	0.00347	35	0.15305	0.40597	0.02674	0.00139
Carbonate Carbon (88308)	0	—	—	—	0.14600	0	—	—	—	0.13400
Cerium (88117)	17	0.02896	0.07303	0.00105	0.08603	17	0.01145	0.02567	0.001	0.03450
Cesium (88118)	12	0.01934	0.04359	0.00827	0.03689	18	0.00719	0.01370	0.0016	0.0148
Chlorine (88115)	32	0.01382	0.10220	0.00116	0.00578	35	0.02276	0.1633	0.00193	0.0023
Chromium (88112)	13	0.00105	0.00633	0.00011	0.00159	24	0.00029	0.00065	0	0.001
Cobalt (88113)	2	0.00092	0.00103	0.00081	0.00141	0	—	—	—	0.001
Copper (88114)	32	0.00397	0.01427	0.00092	0.00135	35	0.00456	0.01507	0.0016	0.001
Europium (88121)	2	0.00093	0.00163	0.00024	0.01124	3	0.00072	0.00157	0	0.0045
Gallium (88124)	29	0.00252	0.00572	0.00011	0.00331	22	0.00080	0.00182	0.00004	0.0013
Gold (88143)	20	0.00197	0.00550	0.00011	0.00501	8	0.00090	0.00207	0	0.002
Lanthanum (88146)	19	0.03008	0.06026	0.00129	0.06947	21	0.01262	0.02497	0.00040	0.02790
Lead (88128)	33	0.00755	0.04106	0.00162	0.00549	35	0.00654	0.06542	0.00005	0.00220
Magnesium (88140)	22	0.01825	0.03693	0.00058	0.01841	27	0.01395	0.05318	0.00088	0.00738

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-12 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT SALT LAKE CITY, UT (in $\mu g/m^3$)

	Salt Lake City, UT (490353006)									
	Met One (5)					Andersen (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Manganese (88132)	32	0.00347	0.00819	0.00046	0.00231	35	0.00328	0.00889	0.00058	0.00092
Mercury (88142)	20	0.00186	0.00508	0.00011	0.00437	18	0.00070	0.00179	0.00004	0.00175
Molybdenum (88134)	12	0.00191	0.00627	0.00011	0.00477	16	0.00087	0.00182	0.00019	0.00191
Nickel (88136)	28	0.01309	0.17676	0.00011	0.00125	23	0.00038	0.00193	0.00004	0.00050
Niobium (88147)	13	0.00224	0.00523	0.00023	0.00420	20	0.00068	0.00175	0.00009	0.00168
Phosphorous (88152)	5	0.00340	0.00687	0.00011	0.00627	3	0.00309	0.00387	0.00203	0.00251
Potassium (88180)	33	0.08496	0.44816	0.00721	0.00341	35	0.09978	0.60087	0.00969	0.00137
Rubidium (88176)	16	0.00091	0.00209	0.00011	0.00217	13	0.00052	0.00160	0.00004	0.00087
Samarium (88162)	1	0.00222	0.00222	0.00222	0.00617	1	0.00058	0.00058	0.00058	0.00247
Scandium (88163)	7	0.00113	0.00302	0.00011	0.00243	2	0.00020	0.00036	0.00004	0.00097
Selenium (88154)	17	0.00092	0.00187	0.00011	0.00212	23	0.00042	0.00101	0.00004	0.00085
Silicon (88165)	33	0.23857	0.95256	0.02825	0.00753	35	0.22324	0.77713	0.03143	0.00302
Silver (88166)	22	0.00719	0.02222	0.00046	0.01048	20	0.00197	0.00494	0.00004	0.00420
Sodium (88184)	15	0.05035	0.18042	0.00035	0.05107	5	0.02972	0.05233	0.01463	0.02050
Strontium (88168)	19	0.00173	0.00696	0.00024	0.00251	30	0.00151	0.01017	0.00004	0.00101
Sulfur (88169)	33	0.29801	0.64844	0.10711	0.00662	35	0.29797	0.67114	0.08485	0.00265
Tantalum (88170)	33	0.01857	0.03691	0.00024	0.01954	31	0.00568	0.00986	0.00040	0.00784
Terbium (88172)	7	0.00091	0.00208	0.00024	0.00752	6	0.00066	0.00274	0.00009	0.00302
Tin (88160)	33	0.01952	0.03399	0.00210	0.01787	35	0.00818	0.01321	0.00381	0.00717
Titanium (88161)	33	0.00898	0.02477	0.00223	0.00208	35	0.00742	0.02181	0.00200	0.00083
Vanadium (88164)	2	0.00076	0.00105	0.00046	0.00150	3	0.00028	0.00058	0.00009	0.00060
Wofram (88186)	19	0.00772	0.01560	0.00011	0.01380	14	0.00217	0.00438	0.00004	0.00554
Yttrium (88183)	12	0.00133	0.00248	0.00024	0.00304	12	0.00055	0.00127	0.00005	0.00122
Zinc (88167)	33	0.00705	0.02960	0.00070	0.00145	35	0.00827	0.03078	0.0015	0.00058

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-13. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT ST. LOUIS, MO (in $\mu g/m^3$)

	St. Louis, MO (295100085)									
	Met One (5)					Andersen (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	35	15.9460	36.8015	5.61856	0.10400	37	15.5672	33.9177	6.51278	0.04000
Ammonium (88301)	38	1.75246	6.37668	0.32688	0.01700	40	1.64614	5.48037	0.16619	0.01500
Sodium Ion (88302)	38	0.13497	0.36566	0.01987	0.03000	40	0.13138	0.42148	0.02548	0.02800
Potassium Ion (88303)	10	0.15767	0.28898	0.07387	0.014	15	0.15195	0.36176	0.07339	0.01300
Nitrate (88306)	38	1.73075	6.63161	0.25581	0.008	40	1.69734	6.56552	0.24213	0.00800
Sulfate (88403)	38	4.23020	17.5918	1.33765	0.012	40	4.15895	16.8408	1.16105	0.01100
Organic Carbon (88305)	24	4.55208	6.92448	2.60132	0.14600	25	4.34840	6.71026	2.54373	0.13400
Elemental Carbon (88307)	24	0.80882	1.57296	0.23018	0.14600	25	0.85087	1.74821	0.45032	0.13400
Total Carbon	24	5.36090	7.87283	3.16637	NA	25	5.19927	7.86998	3.20344	NA
Aluminum (88104)	30	0.04142	0.51741	0.00119	0.01088	26	0.09593	1.81318	0.00285	0.00436
Antimony (88102)	17	0.00737	0.02042	0.00178	0.01476	27	0.00319	0.0074	0.00049	0.00592
Arsenic (88103)	28	0.00227	0.00507	0.00035	0.00247	34	0.00150	0.0035	0.00005	0.00099
Barium (88107)	35	0.06549	0.11098	0.01269	0.05876	36	0.02959	0.05213	0.00066	0.02360
Bromine (88109)	33	0.00431	0.05302	0.00046	0.002	37	0.00426	0.05280	0.001	0.00080
Cadmium (88110)	17	0.00447	0.01002	0.00023	0.0105	20	0.00180	0.00792	0	0.00421
Calcium (88111)	35	0.11230	0.21063	0.01564	0.00347	37	0.13481	0.58229	0.04329	0.00139
Carbonate Carbon (88308)	0	—	—	—	0.146	0	—	—	—	0.13400
Cerium (88117)	23	0.02320	0.05889	0.00143	0.08603	25	0.01048	0.02772	0	0.03450
Cesium (88118)	14	0.01361	0.02719	0.00153	0.03689	20	0.00601	0.01447	0	0.01480
Chlorine (88115)	16	0.01529	0.10653	0.00046	0.0058	26	0.02162	0.30948	0	0.00232
Chromium (88112)	23	0.00143	0.00979	0.00012	0.0016	31	0.00136	0.01104	0	0.001
Cobalt (88113)	2	0.00023	0.00035	0.00011	0.0014	2	0.00014	0.00023	0.00004	0.001
Copper (88114)	34	0.01640	0.19018	0.00036	0.0014	37	0.03913	0.71688	0.001	0.001
Europium (88121)	1	0.00166	0.00166	0.00166	0.01124	2	0.00187	0.00304	0.001	0.00451
Gallium (88124)	28	0.00208	0.00425	0.00011	0.0033	15	0.00071	0.00167	0	0.00133

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-13 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT ST. LOUIS, MO (in $\mu g/m^3$)

	St. Louis, MO (295100085)									
	Met One (5)					Andersen (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Gold (88143)	8	0.00091	0.00199	0.00011	0.00501	1	0.00029	0.00029	0	0.00201
Hafnium (88127)	13	0.01301	0.03328	0.00024	0.02605	19	0.00433	0.01215	0	0.0105
Indium (88131)	14	0.00435	0.01265	0.00059	0.01128	13	0.00178	0.00291	0.00057	0.0045
Iridium (88133)	17	0.00356	0.00684	0.00058	0.00594	10	0.00087	0.00232	0.00014	0.0024
Iron (88126)	35	0.16336	0.55651	0.02843	0.00196	37	0.18826	1.06552	0.05701	0.00079
Lanthanum (88146)	17	0.03044	0.07101	0.00472	0.06947	21	0.01341	0.02849	0.00108	0.02790
Lead (88128)	35	0.01644	0.05871	0.00235	0.00549	37	0.01609	0.08643	0.00227	0.00220
Magnesium (88140)	16	0.01989	0.03903	0.00070	0.01841	17	0.01052	0.02089	0.00038	0.00738
Manganese (88132)	34	0.00958	0.05922	0.00118	0.00231	37	0.009	0.05653	0.00097	0.00092
Mercury (88142)	18	0.00198	0.00389	0.00035	0.00437	15	0.001	0.00184	0.00005	0.00175
Molybdenum (88134)	15	0.00160	0.00417	0.00024	0.00477	19	0.001	0.0021	0.00010	0.00191
Nickel (88136)	34	0.01684	0.10784	0.00189	0.00125	34	0.00135	0.01017	0.00004	0.001
Niobium (88147)	17	0.00180	0.00341	0.00047	0.00420	12	0.00047	0.00100	0.00009	0.00168
Phosphorous (88152)	1	0.07769	0.07769	0.07769	0.00627	1	0.05253	0.05253	0.05253	0.00251
Potassium (88180)	35	0.07724	0.30465	0.00952	0.00341	37	0.09665	0.70515	0.02424	0.00137
Rubidium (88176)	12	0.00106	0.00178	0.00011	0.00217	10	0.00044	0.00092	0.00004	0.00087
Samarium (88162)	0	—	—	—	0.00617	2	0.00111	0.00116	0.00105	0.00247
Scandium (88163)	4	0.00088	0.00165	0.00011	0.00243	3	0.00012	0.00023	0.00004	0.00097
Selenium (88154)	27	0.00161	0.00389	0.00024	0.00212	34	0.00129	0.00458	0.00010	0.00085
Silicon (88165)	35	0.15945	1.04966	0.03410	0.00753	37	0.23356	3.30811	0.03851	0.00302
Silver (88166)	24	0.00508	0.01003	0.00011	0.01048	24	0.00205	0.00527	0.00018	0.00420
Sodium (88184)	28	0.08976	0.21981	0.00187	0.05107	12	0.06837	0.25709	0.00548	0.02050
Strontium (88168)	17	0.00115	0.00248	0.00011	0.00251	24	0.00101	0.00979	0.00013	0.00101

DRAFT-DO NOT QUOTE OR CITE

TABLE 3 B-13 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT ST. LOUIS, MO (in $\mu g/m^3$)

	St. Louis, MO (295100085)									
	Met One (5)					Andersen (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Titanium (88161)	34	0.00639	0.02906	0.00071	0.00208	35	0.00826	0.09360	0.00111	0.00083
Vanadium (88164)	8	0.00097	0.00295	0.00011	0.00150	13	0.00100	0.00403	0.00005	0.00060
Wofram (88186)	12	0.00674	0.01858	0.00047	0.01380	3	0.00094	0.00115	0.00057	0.00554
Yttrium (88183)	11	0.00104	0.00201	0.00058	0.00304	14	0.00057	0.00121	0.00020	0.00122
Zinc (88167)	35	0.02435	0.07338	0.00107	0.00145	37	0.04568	0.47977	0.00864	0.00058
Zirconium (88185)	15	0.00183	0.00579	0.00035	0.00359	20	0.00088	0.0037	0.00004	0.0014

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-14. SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT SEATTLE, WA (in $\mu g/m^3$)

	Seattle, WA (530330080)									
	Met One (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
$PM_{2.5}$ Mass (88101)	40	9.4676	25.1163	3.29897	0.10400	41	7.58131	23.7103	3.04205	0.04000
Ammonium (88301)	32	0.45565	1.86358	0.07057	0.01700	41	0.51602	1.97357	0.09762	0.00700
Sodium Ion (88302)	39	0.25671	0.57632	0.06332	0.03000	41	0.16680	0.48493	0.04035	0.01200
Potassium Ion (88303)	5	0.55895	2.31058	0.04885	0.01400	27	0.14516	2.33559	0.01906	0.00600
Nitrate (88306)	39	0.83427	3.02535	0.19552	0.008	41	0.68270	2.60029	0.12613	0.00300
Sulfate (88403)	39	1.52949	4.19391	0.54916	0.01200	41	1.47833	4.30718	0.44612	0.005
Organic Carbon (88305)	39	3.82378	9.31148	1.70716	0.146	41	2.50795	7.09910	0.93538	0.05900
Elemental Carbon (88307)	39	0.86143	2.78426	0.14378	0.146	41	0.61197	1.63835	0.14849	0.05900
Total Carbon	39	4.68521	12.0957	2.13806	NA	41	3.11992	8.73745	1.31881	NA
Aluminum (88104)	33	0.02074	0.17279	0.00024	0.0109	29	0.0176	0.20840	0.00023	0.00436
Antimony (88102)	28	0.00646	0.01577	0.00058	0.0148	21	0.00311	0.007	0.00028	0.00592
Arsenic (88103)	30	0.00244	0.00892	0.00035	0.002	29	0.00144	0.004	0.00010	0.00099
Barium (88107)	39	0.05726	0.21693	0.00094	0.0588	39	0.03005	0.15435	0.00655	0.02360
Bromine (88109)	36	0.00196	0.00560	0.00011	0.00199	41	0.00199	0.00428	0.00033	0.00080
Cadmium (88110)	19	0.00495	0.01156	0.00105	0.01050	19	0.002	0.00551	0.0006	0.00421
Calcium (88111)	40	0.04455	0.11359	0.01447	0.00347	40	0.0337	0.09934	0.00725	0.00139
Carbonate Carbon (88308)	0	—	—	—	0.14600	0	—	—	—	0.059
Cerium (88117)	23	0.02206	0.05807	0.00071	0.08603	21	0.01	0.02387	0.00038	0.0345
Cesium (88118)	16	0.01369	0.04476	0.00011	0.03689	19	0.00779	0.0257	0.00090	0.0148
Chlorine (88115)	34	0.05698	0.44694	0.00058	0.00578	39	0.03522	0.30344	0.00005	0.00232
Chromium (88112)	35	0.00212	0.00981	0.00011	0.00159	39	0.00174	0.009	0.00010	0.0006
Cobalt (88113)	4	0.00053	0.00106	0.00011	0.00141	3	0.00020	0	0.00010	0.0006
Copper (88114)	38	0.00350	0.04612	0.00011	0.00135	41	0.00317	0.0449	0.00010	0.0005
Europium (88121)	3	0.00160	0.00364	0.00024	0.01124	3	0.00077	0.001	0.00034	0.00451
Gallium (88124)	33	0.00218	0.00525	0.00024	0.00331	32	0.00085	0.002	0.00005	0.00133

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-14 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT SEATTLE, WA (in $\mu g/m^3$)

	Seattle, WA (530330080)									
	Met One (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Gold (88143)	20	0.00237	0.00647	0.00035	0.00501	18	0.00097	0.003	0.00023	0.00201
Hafnium (88127)	14	0.00826	0.02125	0.00093	0.02605	17	0.00431	0.0125	0.00023	0.01050
Indium (88131)	17	0.00475	0.01196	0.00047	0.01128	20	0.00194	0.005	0.00005	0.00452
Iridium (88133)	21	0.00326	0.00941	0.00024	0.00594	20	0.00117	0.002	0.00023	0.00238
Iron (88126)	40	0.07195	0.32351	0.01493	0.00196	41	0.05825	0.28782	0.01153	0.00079
Lanthanum (88146)	20	0.02539	0.05843	0.00520	0.06947	24	0.01351	0.02547	0.00523	0.0279
Lead (88128)	39	0.00756	0.06970	0.00093	0.00549	41	0.00662	0.07830	0.00184	0.00220
Magnesium (88140)	19	0.02235	0.11371	0.00093	0.01841	28	0.01466	0.18727	0.00085	0.00738
Manganese (88132)	40	0.00471	0.02406	0.00035	0.00231	39	0.00430	0.0226	0.00014	0.00092
Mercury (88142)	22	0.00157	0.00459	0.00011	0.00437	18	0.00108	0.00353	0.00019	0.00175
Molybdenum (88134)	26	0.00193	0.00515	0.00024	0.00477	28	0.00156	0.00513	0.0001	0.00191
Nickel (88136)	38	0.00667	0.08607	0.00071	0.00125	41	0.00286	0.01191	0.00038	0.00050
Niobium (88147)	16	0.00156	0.00470	0.00024	0.00420	19	0.00068	0.00141	0.00005	0.00168
Phosphorous (88152)	3	0.00121	0.00200	0.00071	0.00627	0	—	—	—	0.00251
Potassium (88180)	40	0.09503	2.16303	0.00745	0.00341	41	0.09413	2.26987	0.01046	0.00137
Rubidium (88176)	13	0.00071	0.00200	0.00011	0.00217	15	0.00037	0.00094	0.00014	0.00087
Samarium (88162)	1	0.00093	0.00093	0.00093	0.00617	1	0.00005	0.00005	0.00005	0.00247
Scandium (88163)	10	0.00090	0.00247	0.00011	0.00243	3	0.00025	0.00047	0.00010	0.00097
Selenium (88154)	22	0.00097	0.00199	0.00024	0.00212	25	0.00044	0.00090	0.00005	0.00085
Silicon (88165)	40	0.06449	0.25778	0.01234	0.00753	40	0.05035	0.22823	0.00852	0.00302
Silver (88166)	26	0.00550	0.01152	0.00035	0.01048	30	0.00166	0.00457	0.00005	0.00420
Sodium (88184)	37	0.20378	0.51094	0.03422	0.05107	38	0.15366	0.39380	0.02369	0.02050
Strontium (88168)	16	0.00375	0.04213	0.00011	0.00251	25	0.00252	0.04359	0.00005	0.00101

DRAFT-DO NOT QUOTE OR CITE

TABLE 3B-14 (cont'd). SUMMARY STATISTICS FOR THE SPECIATION SAMPLERS AT SEATTLE, WA (in $\mu g/m^3$)

	Seattle, WA (530330080)									
	Met One (5)					URG (6)				
Parameter	**N**	**Mean**	**Max**	**Min**	**MDL**	**N**	**Mean**	**Max**	**Min**	**MDL**
Titanium (88161)	40	0.00403	0.02453	0.00047	0.00208	38	0.00321	0.02604	0.0007	0.00083
Vanadium (88164)	25	0.00465	0.01665	0.00024	0.00150	30	0.00412	0.01516	0.00010	0.00060
Wofram (88186)	28	0.00603	0.01537	0.00035	0.01380	19	0.00252	0.00706	0.00014	0.00554
Yttrium (88183)	10	0.00087	0.00152	0.00011	0.00304	18	0.00072	0.00250	0.00005	0.00122
Zinc (88167)	40	0.00741	0.03048	0.00035	0.00145	41	0.00823	0.03061	0.00179	0.0006
Zirconium (88185)	18	0.00146	0.00423	0.00011	0.00359	23	0.00073	0.00240	0	0.00144

*The blank spaces mean there are no non-zero, valid measurements.

DRAFT-DO NOT QUOTE OR CITE

REFERENCES

Coutant, B.; Stetzer, S. (2001) Evaluation of $PM_{2.5}$ speciation sampler performance and related sample collection and stability issues: final report. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; report no. EPA-454/R-01-008. Available: http://www.epa.gov/ttn/amtic/pmspec.html [5 April, 2002].

Coutant, B.; Zhang, X.; Pivetz, T. (2001) Summary statistics and data displays for the speciation minitrends study: final report. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; contract no. 68-D-98-030.

APPENDIX 3C

Organic Composition of Particulate Matter

Although organic compounds typically constitute approximately 10 to 70% of the total dry fine particle mass in the atmosphere, organic PM concentrations, composition, and formation mechanisms are poorly understood. This is because particulate organic matter is an aggregate of hundreds of individual compounds spanning a wide range of chemical and thermodynamic properties (Saxena and Hildemann, 1996). The presence of multiphase or "semivolatile" compounds complicates collection of organic particulate matter. Furthermore, no single analytical technique currently is capable of analyzing the entire range of compounds present. Rigorous analytical methods frequently identify only 10 to 20% of the organic mass on the molecular level (Rogge et al., 1993). The data shown in Appendix 3C are meant to complement the data given for the inorganic components of particles in Appendix 6A of the 1996 PM AQCD (U. S. Environmental Protection Agency, 1996). Table 3C-1 lists a number of recent urban and some rural measurements of particulate organic and elemental carbon in μg of carbon/m^3 (μg C/m^3). Emphasis is placed on measurements published after 1995. The analysis method and artifact correction procedure, if any, are indicated. Table 3C-2 presents information on recent (post-1990) studies concerning concentrations (in ng C/m^3) of particulate organic compounds found at selected U.S. sites.

TABLE 3C-1. PARTICULATE ORGANIC AND ELEMENTAL CARBON CONCENTRATIONS (in μg C/m^3) BASED ON STUDIES PUBLISHED AFTER 1995

Reference	Location	Dates	OC Mean (Max)	EC Mean (Max)	TC Mean (Max)	Avg. Time	Notes
URBAN $PM_{2.5}$							
Offenberg and Baker (2000)	Chicago, IL	July 1994; Jan 1995			2.2 (3.8)	12 h	PM_{12}; Imp; TOT
					1.7		$PM_{1.4}$; Imp; TOT
Allen et al. (1999)	Uniontown, PA	July-Aug 1990	(0.8-8.4)[a]	(0.4-3.5)[a]		3 h	$PM_{2.5}$; DQQ; TOR[b]
				1.3 (3.1)		10 min	Aeth
Pedersen et al. (1999)	Boston, MA	Jan-Dec 1995	5.8	1.7		24 h	$PM_{2.0}$; Q; TOT
	Reading, MA (suburban)		4.0	0.7			
	Quabbin, MA (rural)		2.8	0.5			
	Rochester, NY (urban)		3.3	0.7			
	Brockport, NY (rural)		2.7	0.5			
IMPROVE (2000)	Washington, DC	1994-1998	3.4	1.1		24 h	$PM_{2.5}$; QQ; TOR
	Seattle, WA		1.8	0.3			
Lewtas et al. (2001)	Seattle, WA	Apr-May 1999	8.0	1.4		23 h	$PM_{2.5}$; DQA; EGA[c]
Khwaja (1995)	Schenectady, NY	Oct 24-26, 1991			23.2 (49.9)	6 h	$PM_{1.0}$; Q; Th
Christoforou et al. (2000)	Azusa, CA	Jan-Dec 1993	9.4	1.3		24 h	$PM_{2.1}$; Q; TOR
	Long Beach, CA		8.9	1.8			
	Central, LA		12.3	2.7			
	Rubidoux, LA		9.7	1.5			
	San Nicolas, LA		1.6	1.5			
Turpin and Huntzicker (1995)	Claremont, CA	Jun-Sept	na (29.4)	na (9.0)		2 h	$PM_{2.5}$; Q+TQ; TOT[d]
	Long Beach, CA	Nov-Dec 1987	na (62.6)	na (24.6)		2-6 h	

DRAFT-DO NOT QUOTE OR CITE

TABLE 3C-1 (cont'd). PARTICULATE ORGANIC AND ELEMENTAL CARBON CONCENTRATIONS (in μg C/m^3) BASED ON STUDIES PUBLISHED AFTER 1995

Reference	Location	Dates	OC Mean (Max)	EC Mean (Max)	TC Mean (Max)	Avg. Time	Notes
RURAL $PM_{2.5}$							
Klinedinst and Currie (1999)	Welby, CO Brighton, CO	Dec 1996-Jan 1997	5.6 (13.4) 3.6 (6.4)	3.3 (8.1) 1.9 (3.6)		6 h	$PM_{2.5}$; Q; TOR
Andrews et al. (2000)	Look Rock, Smoky Mountains, TN	July-Aug 1995	2.2 2.7 1.2	0.4 0.1 0.2		12 h (day)	$PM_{2.1}$; QQ; TOR[e] $PM_{2.1}$; Q+TQ; TOR[d] $PM_{1.8}$; Imp; TMO
Malm and Gebhart (1996)	Tahoma Wood, WA	June-Aug 1990	2.6 (7.4)	0.7 (2.2)		12 h	$PM_{2.5}$; QQ; TOR[f]
IMPROVE (2000)	Three Sisters Wilderness, OR Rocky Mountains, CO Brigantine, NJ Acadia, MA Jefferson: James River Face Wilderness, VA Glacier, MT	1994-1998	0.9 1.0 2.0 1.2 3.8 2.4	0.2 0.2 0.5 0.2 0.7 0.4		24 h	$PM_{2.5}$; QQ; TOR
Hegg et al. (1997)	150 km East of Mid-Atlantic Coast (0.02-4 km altitude)	July 1996			2.9 (5.4)		$PM_{1.0}$; QQ ; EGA[e]
Cui et al. (1997)	Meadview, AZ	Aug 6-15, 1992			3.0	12 h	$PM_{2.5}$; VDQA; EGA[c]
Chow et al. (1996)	Point Reyes, CA Altamont Pass, CA Pacheco Pass, CA Crows Landing, CA Academy, CA Button-Willow, CA Edison, CA Caliente, CA Sequoia, CA Yosemite, CA	July-Aug 1990	1.5 (2.7) 4.8 (7.2) 3.2 (6.1) 7.4 (12.7) 5.9 (8.7) 6.4 (10.6) 10.0 (12.8) 7.4 (10.7) 5.3 (7.0) 12.1 (25.8)	0.4 (0.6) 2.6 (3.9) 1.0 (1.3) 1.8 (2.5) 1.4 (2.4) 1.9 (2.7) 2.9 (4.1) 3.3 (4.4) 1.6 (3.0) 1.9 (3.5)	5-7 $hPM_{2.5}$; Q+TQ; TOR[g]		

DRAFT-DO NOT QUOTE OR CITE

TABLE 3C-1 (cont'd). PARTICULATE ORGANIC AND ELEMENTAL CARBON CONCENTRATIONS (in μg C/m^3) BASED ON STUDIES PUBLISHED AFTER 1995

Reference	Location	Dates	OC Mean (Max)	EC Mean (Max)	TC Mean (Max)	Avg. Time	Notes
RURAL $PM_{2.5}$							
Malm and Day (2000)	Grand Canyon, AZ	July-Aug 1998	1.1 (1.6)	0.10 (0.3)		24 h	$PM_{2.5}$; QQ; TOR[f]
PM_{10}							
Omar et al. (1999)	Bondville, IL	Jan-Dec 1994	2.6	0.2		24-48 h	PM_{10}; Q; TOR
Gertler et al. (1995)	Bullhead City, AZ	Sept 1988-Oct 1989	6.0 (16.0)	1.9 (4.0)		24 h	PM_{10}; Q; TOR
Chow et al. (1996)	Santa Barbara, CA (urban)	Jan-Dec 1989			8.8	24 h	PM_{10}; Q; TOR
	Santa Maria, CA (urban)				4.6		
	Santa Ynez, CA (airport)				3.5		
	Gaviota, CA (rural SB)				3.4		
	Watt Road, CA (rural SB)				2.1		
	Anacapa Island, CA				3.1		
Lioy and Daisey (1987)		1982:					PM_{15}; Q
	Newark, NJ	Summer	4.1	3.0			
		Winter	5.9	3.3			
	Elizabeth, NJ	Summer	2.1	1.7			
		Winter	7.1	2.3			
	Camden, NJ	Summer	2.2	1.3			
		Winter	5.2	2.0			

A limited amount of rural data is presented. In some cases, total carbon (TC = OC + EC) is reported. OC concentrations must be multiplied by the average molecular weight per carbon weight to convert to mass of particulate organic compounds. The location and dates over which sampling occurred are provided. Averaging time refers to the sampling duration. Sampling method: Q – quartz fiber filter; QQ – two quartz fiber filters in series; Q+TQ – a quartz fiber filter in one port and a Teflon followed by a quartz filter in a parallel port; Imp – cascade impactor; DQQ – denuder followed by two quartz fiber filters; DQA – denuder followed by quartz fiber filter and adsorbent; VDQA – virtual impactor inlet followed by denuder, quartz filter, and adsorbent. Analysis method is reported as follows: TOR – thermal optical reflectance; TOT – thermal optical transmittance; TMO – thermal MnO_2 oxidation; EGA – evolved gas analysis; Th – Thermal analysis; Aeth – Aethalometer. na – data not available.

[a]Range is provided. It should be noted that samples were collected only during elevated pollution episodes and are not representative of average concentrations.
[b]Particulate OC was considered to be the sum of front and back quartz fiber filters.
[c]Sum of adsorbent and filter after correction for inlet losses and denuder efficiency.
[d]Corrected for adsorption by subtracting the Teflon-quartz back-up filter.
[e]Reported concentrations are corrected for adsorption by subtracting the quartz (TQ or QQ) back-up filter.
[f]Sampler contained two quartz fiber filters in series, but publication did not indicate whether the quartz back-up filter was subtracted to correct for adsorption.
[g]Corrected for adsorption using Micro-Orifice Uniform Deposit Impactor (MOUDI) data from a collocated sampler.

DRAFT-DO NOT QUOTE OR CITE

TABLE 3C-2. PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m^3) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993)[a] Jan-Dec 1982 (annual average) $PM_{2.1}$		**Schauer and Cass (2000)** Dec 26-28, 1995 (pollution episode) $PM_{2.5}$		**Veltkamp et al. (1996)** July 24-Aug 4, 1989 no precut	**Khwaja (1995)** October 1991 (semiurban) no precut	**Allen et al. (1997)** Summer 1994 (urban) $PM_{1.9}$	**Fraser et al. (1998)** Sept 8-9, 1993 (urban)
	Los Angeles, CA	**Pasadena, CA**	**Fresno, CA**	**Bakersfield, CA**	**Niwot Ridge, CO**	**Schenectady, NY**	**Kenmore Square, Boston, MA**	**Los Angeles Basin, CA**
n-Alkanes								
n-tricosane	6.7	5.4			19.23 (57.7)			
n-tetracosane	6.4	4.7	42.3	12.7	6.04 (21.1)			
n-pentacosane	11.2	9.5	41.2	14.2	7.77 (21.3)			
n-hexacosane	8.2	4.3	29.9	10.7	2.08 (12.7)			
n-heptacosane	6.7	5.6	25.0	10.8	5.62 (15.1)			
n-octacosane	3.1	2.5	12.3	5.24	1.26 (9.0)			
n-nonacosane	7.1	4.7	33.8	23.6	7.70 (20.6)			
n-triacontane	2.7	2.5	7.39	4.27	0.76 (4.6)			
n-hentriacontane	12.6	9.6	16.1	9.66	5.24 (17.9)			
n-dotriacontane	1.5	1.5	2.61	3.50	0.41 (2.1)			
n-tritriacontane	2.1	2.3	5.02	3.31	1.49 (5.5)			
n-tetratriacontane	0.58	0.68						
Total n-alkanes	**68.9**	**53.3**	**215.6**	**98.0**	**57.9**			
n-Alkanoic Acids								
n-nonanoic acid	6.6	5.3						
n-decanoic acid	2.0	2.4	0.711	0.164				
n-undecanoic acid	2.8	6.0						
n-dodecanoic acid	5.3	7.0	0.905	0.803				
n-tridecanoic acid	4.3	4.9	6.17	1.78				
n-tetradecanoic acid	19.7	22.2	9.42	4.01				
n-pentadecanoic acid	5.3	6.1	33.7	5.63				
n-hexadecanoic acid (palmitic acid)	140.5	127.4	166	54.4				
n-heptadecanoic acid	4.7	5.2	13.6	3.77				
n-octadecanoic acid (stearic acid)	59.2	50.0	60.0	24.1				
n-nonadecanoic acid	1.1	1.1	10.7	2.58				
n-eicosanoic acid	5.1	6.1	41.2	10.4				
n-heneicosanoic acid	2.1	2.3	20.8	6.46				
n-docosanoic acid	8.7	9.9	160	43.1				
n-tricosanoic acid	2.0	2.5	32.1	9.71				
n-tetracosanoic acid	11.8	16.5	205	78.0				

DRAFT-DO NOT QUOTE OR CITE

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m^3) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993)[a] Jan-Dec 1982 (annual average) $PM_{2.1}$		**Schauer and Cass (2000)** Dec 26-28, 1995 (pollution episode) $PM_{2.5}$		**Veltkamp et al. (1996)** July 24-Aug 4, 1989 no precut	**Khwaja (1995)** October 1991 (semiurban) no precut	**Allen et al. (1997)** Summer 1994 (urban) $PM_{1.9}$	**Fraser et al. (1998)** Sept 8-9, 1993 (urban)
	Los Angeles, CA	**Pasadena, CA**	**Fresno, CA**	**Bakersfield, CA**	**Niwot Ridge, CO**	**Schenectady, NY**	**Kenmore Square, Boston, MA**	**Los Angeles Basin, CA**
n-Alkanoic Acids (cont'd)								
n-pentacosanoic acid	1.3	1.6	15.4	6.59				
n-hexacosanoic acid	5.6	9.3	174	81.3				
n-heptacosanoic acid	0.49	0.81	2.56	2.38				
n-octacosanoic acid	2.7	4.9	21.3	9.65				
n-nonacosanoic acid	0.33	0.57	1.46	2.11				
n-triacontanoic acid	1.0	2.2	4.32	5.79				
Total n-alcanoic acids	**292.6**	**294.3**	**979.3**	**352.7**				
n-Alkenoic Acids								
n-9-hexadecenoic acid			18.8	3.96				
n-9-octadecenoic acid	24.8	26.0	27.1	3.96				
n-9,12-octadecane-dienoic acid			13.6	1.83				
Total n-alkenoic acids	**24.8**	**26.0**	**59.5**	**9.75**				
n-Alkanals								
1-octanal					3.26 (14.4)			
n-nonanal	5.7	9.5	19.4	3.01	29.01 (62.8)			
n-decanal					23.58 (71.2)			
n-dodecanal					6.01 (16.4)			
n-tridecanal					6.50 (25.8)			
n-tetradecanal					9.62 (30.7)			
n-pentadecanal					12.47 (113.6)			
n-hexadecanal					17.45 (49.3)			
n-heptadecanal					24.09 (88.9)			
n-octadecanal					1.84 (11.7)			
Total n-alkanals	**5.7**	**9.5**	**19.4**	**3.01**	**133.8**			

DRAFT-DO NOT QUOTE OR CITE

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m^3) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993)[a] Jan-Dec 1982 (annual average) $PM_{2.1}$		**Schauer and Cass (2000)** Dec 26-28, 1995 (pollution episode) $PM_{2.5}$		**Veltkamp et al. (1996)** July 24-Aug 4, 1989 no precut	**Khwaja (1995)** October 1991 (semiurban) no precut	**Allen et al. (1997)** Summer 1994 (urban) $PM_{1.9}$	**Fraser et al. (1998)** Sept 8-9, 1993 (urban)
	Los Angeles, CA	**Pasadena, CA**	**Fresno, CA**	**Bakersfield, CA**	**Niwot Ridge, CO**	**Schenectady, NY**	**Kenmore Square, Boston, MA**	**Los Angeles Basin, CA**
n-Alkanols								
1-decanol					8.66 (64.1)			
1-dodecanol					21.29 (61.7)			
1-tetradecanol					13.59 (41.4)			
1-pentadecanol					4.50 (30.1)			
1-hexadecanol					27.42 (141.1)			
Total n-alkanols					**75.5**			
Aliphatic Dicarboxylic Acids								
oxalic acid (C_2)						198 (360)		
malonic acid (propanedioic)	32.7	44.4				84 (107)		
methylmalonic acid (methylpropanedioic)			2.13	nd				
malonic acid (2-butenedioic)	0.66	1.3						
succinic acid (butanedioic)	66.5	51.2				102 (167)		
methylsuccinic acid (methylbutanedioic)	18.0	15.0	24.0	8.80				
glutaric acid (pentanedioic)	32.3	28.3	21.3	10.5				
methylglutaric acid (methylpentanedioic)	19.3	16.6						
hydroxybutanedioic acid	14.3	16.0						
adipic acid (hexanedioic)	14.1	14.1	3.39	3.07				
pimelic acid (heptanedioic)			2.22	1.03				
suberic acid (octanedioic)	3.4	4.1	4.41	13.4				
axelaic acid (nonanedioic)	29.0	22.8	19.9	8.22				
Total aliphaitc dicarboxylic acids	**230.3**	**213.8**	**77.4**	**45.0**		**384**		

DRAFT-DO NOT QUOTE OR CITE

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m^3) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993)[a] Jan-Dec 1982 (annual average) $PM_{2.1}$		**Schauer and Cass (2000)** Dec 26-28, 1995 (pollution episode) $PM_{2.5}$		**Veltkamp et al. (1996)** July 24-Aug 4, 1989 no precut	**Khwaja (1995)** October 1991 (semiurban) no precut	**Allen et al. (1997)** Summer 1994 (urban) $PM_{1.9}$	**Fraser et al. (1998)** Sept 8-9, 1993 (urban)
	Los Angeles, CA	**Pasadena, CA**	**Fresno, CA**	**Bakersfield, CA**	**Niwot Ridge, CO**	**Schenectady, NY**	**Kenmore Square, Boston, MA**	**Los Angeles Basin, CA**
Ketocarboxylic Acids								
pyruvic acid (C_3)						59 (103)		
glyoxylic acid (C_2)						44 (68)		
Total ketocarboxylic acids						**103**		
Diterpenoid/Resin Acids								
dehydroabietic acid	23.6	22.6	98.5	8.01				
abietic acid			30.4	0.784				
13-isopropyl-5α-podocarpa-6,8,11,13-tetraen-16-oic acid	0.63	1.2						
8,15-pimaradien-18-oic acid	0.44	0.57	0.48	0.03				
pimaric acid	2.3	4.8	9.97	0.735				
isopimaric acid	1.3	2.3	127	7.95				
7-oxodehydroabietic acid	3.4	4.1	6.68	1.43				
abieta-6,8,11,13,15-pentaen-18-oic acid			11.8	2.43				
abieta-8,11,13,15-tetraen-18-oic acid			2.62	0.251				
sandaracopimaric acid	1.6	2.2	8.91	0.525				
Total diterpenoid acids	**33.3**	**37.6**	**296.4**	**22.15**				
Aromatic Polycarboxylic Acids								
1,2-benzene-dicarboxylic acid (phthalic acid)	60.0	55.7	9.16	6.78				
1,3-benzene-dicarboxylic acid	3.4	2.9	3.41	1.98				

DRAFT-DO NOT QUOTE OR CITE

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m^3) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993)[a] Jan-Dec 1982 (annual average) $PM_{2.1}$		**Schauer and Cass (2000)** Dec 26-28, 1995 (pollution episode) $PM_{2.5}$		**Veltkamp et al. (1996)** July 24-Aug 4, 1989 no precut	**Khwaja (1995)** October 1991 (semiurban) no precut	**Allen et al. (1997)** Summer 1994 (urban) $PM_{1.9}$	**Fraser et al. (1998)** Sept 8-9, 1993 (urban)
	Los Angeles, CA	**Pasadena, CA**	**Fresno, CA**	**Bakersfield, CA**	**Niwot Ridge, CO**	**Schenectady, NY**	**Kenmore Square, Boston, MA**	**Los Angeles Basin, CA**
Aromatic Polycarboxylic Acids (cont'd)								
1,4-benzene-dicarboxylic acid	2.8	1.5	5.16	4.48				
benzene tricarboxylic acids			14.4	8.77				
4-methyl-1,2-benzenedicarboxylic acid	27.8	28.8						
1,2,4-benzene-tricarboxylic acid (trimellitic acid)	0.52	0.84						
1,3,5-benzene-tricarboxylic acid (trimesic acid)	20.6	17.2						
1,2,4,5-benzene-tetracarboxylic acid (pyromellitic acid)	0.74	0.80						
Total aromatic polycarboxylic acids	**115.9**	**107.7**	**32.1**	**22.0**				
Polycyclic Aromatic Hydrocarbons								
retene	0.07	0.06	6.02	0.563				
fluoranthene	0.15	0.13	2.52	0.553				0.07 (0.26)
acephenanthrylene			0.834	0.302				0.02 (0.05)
pyrene	0.26	0.17	3.28	0.564				0.07 (0.26)
C_1-202 MW PAH			11.7	3.80				0.07 (0.36)
C_2-202 MW PAH								0.03 (0.32)
benz[*a*]anthracene	0.29	0.25	13.8	2.49				0.15 (1.09)
cyclopenta[*cd*]pyrene	0.23	0.41	1.90	0.496				0.14 (1.02)
benzo[*ghi*]-fluoranthene	0.39	0.30	6.05	1.25				0.20 (0.97)
C_1-226 MW PAH			10.1	1.48				0.14 (0.97)
chrysene/triphenyline	0.61	0.43	7.70	1.50				0.34 (1.62)

DRAFT-DO NOT QUOTE OR CITE

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m^3) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993)[a] Jan-Dec 1982 (annual average) $PM_{2.1}$		**Schauer and Cass (2000)** Dec 26-28, 1995 (pollution episode) $PM_{2.5}$		**Veltkamp et al. (1996)** July 24-Aug 4, 1989 no precut	**Khwaja (1995)** October 1991 (semiurban) no precut	**Allen et al. (1997)** Summer 1994 (urban) $PM_{1.9}$	**Fraser et al. (1998)** Sept 8-9, 1993 (urban)
	Los Angeles, CA	**Pasadena, CA**	**Fresno, CA**	**Bakersfield, CA**	**Niwot Ridge, CO**	**Schenectady, NY**	**Kenmore Square, Boston, MA**	**Los Angeles Basin, CA**
Polycyclic Aromatic Hydrocarbons (cont'd)								
C_1-228 MW PAH			17.6	5.35				0.34 (2.16)
C_2-228 MW PAH								0.09 (0.46)
benz[*e*]acephen-anthrylene								0.20 (1.00)
benzo[*k*]fluoranthene	1.15	1.20	8.69	2.13				0.22 (1.07)
benzo[*b*]fluoranthene	1.23	0.85	10.7	2.48				
benzo[*j*]fluoranthene			3.62	0.499				0.02 (0.10)
benzo[*e*]pyrene	0.97	0.93	7.20	1.98				0.22 (1.00)
benzo[*a*]pyrene	0.42	0.44	8.23	1.77				0.14 (0.80)
perylene			1.50	0.246				0.05 (0.51)
methyl-substituted 252 MW PAH								0.10 (0.88)
indeno[*1,2,3-cd*]-pyrene	0.37	0.42	6.84	2.56				0.29 (1.38)
indeno[*1,2,3-cd*]-fluoranthene	1.05	1.09	1.36	0.764				0.10 (0.46)
benzo[*ghi*]perylene	4.47	4.43	9.75	3.49				0.77 (4.23)
anthanthrene			0.180	0.131				
coronene								
Total polycyclic aromatic hydrocarbons	**11.66**	**11.10**	**139.57**	**34.40**				**3.77**
Oxygenated PAHs/ Polycyclic Aromatic Ketones/Quinones								
1,4-naphthoquinone								
1-acenaphthenone							0.26	
9-fluorenone							2.07	0.29 (1.04)
1,8-naphthalic anhydride							1.77	0.41 (1.65)
phenanthrenequinone							0.43	
phenalen-9-one								0.53 (2.23)
anthracene-9,10-dione								0.36 (1.14)
methylanthracene-9,10-dione								0.09 (0.24)
11H-benzo[*a*]fluoren-11-one							1.03	

DRAFT-DO NOT QUOTE OR CITE

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m^3) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993)[a] Jan-Dec 1982 (annual average) $PM_{2.1}$		**Schauer and Cass (2000)** Dec 26-28, 1995 (pollution episode) $PM_{2.5}$		**Veltkamp et al. (1996)** July 24-Aug 4, 1989 no precut	**Khwaja (1995)** October 1991 (semiurban) no precut	**Allen et al. (1997)** Summer 1994 (urban) $PM_{1.9}$	**Fraser et al. (1998)** Sept 8-9, 1993 (urban)
	Los Angeles, CA	**Pasadena, CA**	**Fresno, CA**	**Bakersfield, CA**	**Niwot Ridge, CO**	**Schenectady, NY**	**Kenmore Square, Boston, MA**	**Los Angeles Basin, CA**
Oxygenated PAHs/ Polycyclic Aromatic Ketones/Quinones (cont'd)								
7H-benzo[*c*]fluoren-7-one							0.37	
11H-benzo[*b*]fluoren-11-one							0.85	
1H-phenalen-1-one			7.96	0.588				
benzanthrone							1.18	
5,12-naphthacene-quinone							0.32	
7H-benz[*de*]-anthracen-7-one	0.81	0.84	7.80	1.48				
benz[*de*]anthracene-7-dione								0.20 (1.00)
benz[*a*]anthracene-7,12-dione	0.21	0.25						0.09 (0.31)
cyclopenta[*def*]phen-anthrone								0.05 (0.14)
benzo[*cd*]pyren-6-one	0.80	1.24						0.54 (2.47)
6H-benzo[*cd*]pyrene-6-one							1.34	
benzo[*a*]pyrene-6,12-dione							0.096	
Total polycyclic aromatic ketones/quinones	**1.82**	**2.33**	**15.76**	**2.07**			**9.72**	**2.56**
Steroids								
cholesterol	**nd**	**1.9**						
Substituted Phenols								
p-benzenediol			3.46	nd				
m-benzenediol			7.59	nd				
hydroxybenzaldehydes			2.64	0.604				
Total substituted phenols			**13.69**	**0.604**				

DRAFT-DO NOT QUOTE OR CITE

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m^3) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993)[a] Jan-Dec 1982 (annual average) $PM_{2.1}$		**Schauer and Cass (2000)** Dec 26-28, 1995 (pollution episode) $PM_{2.5}$		**Veltkamp et al. (1996)** July 24-Aug 4, 1989 no precut	**Khwaja (1995)** October 1991 (semiurban) no precut	**Allen et al. (1997)** Summer 1994 (urban) $PM_{1.9}$	**Fraser et al. (1998)** Sept 8-9, 1993 (urban)
	Los Angeles, CA	**Pasadena, CA**	**Fresno, CA**	**Bakersfield, CA**	**Niwot Ridge, CO**	**Schenectady, NY**	**Kenmore Square, Boston, MA**	**Los Angeles Basin, CA**
Guaiacol and Substituted Guaiacols								
guaiacol			0.889	0.832				
4-methylguaiacol			0.606	0.387				
trans-isoeugenol			1.45	1.04				
vanillin			26.8	6.05				
acetovanillone			3.23	0.705				
guaiacyl acetone			10.8	4.29				
coniferyl aldehyde			47.0	nd				
Total guaiacol and substituted guaiacols			**90.78**	**13.30**				
Syringol and Substituted Syringols								
syringol			1.16	0.845				
4-methylsyringol			1.72	1.77				
4-ethylsyringol			2.28	2.39				
4-propylsyringol			0.871	nd				
4-propenylsyringol			4.38	1.40				
syringaldehyde			135	44.5				
acetosyringone			171	55.7				
acetonylsyringol			406	68.1				
propionylsyringol			32.1	16.2				
butyrylsyringol			15.3	6.18				
sinapyl aldehyde			15.9					
Total syringol and substituted syringols			**785.7**	**197.1**				
Sugars								
levoglucosan			7590	1100				
other sugars			1070	171				
Total sugars			**8660**	**1271**				

DRAFT-DO NOT QUOTE OR CITE

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m^3) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993)[a] Jan-Dec 1982 (annual average) $PM_{2.1}$		**Schauer and Cass (2000)** Dec 26-28, 1995 (pollution episode) $PM_{2.5}$		**Veltkamp et al. (1996)** July 24-Aug 4, 1989 no precut	**Khwaja (1995)** October 1991 (semiurban) no precut	**Allen et al. (1997)** Summer 1994 (urban) $PM_{1.9}$	**Fraser et al. (1998)** Sept 8-9, 1993 (urban)
	Los Angeles, CA	**Pasadena, CA**	**Fresno, CA**	**Bakersfield, CA**	**Niwot Ridge, CO**	**Schenectady, NY**	**Kenmore Square, Boston, MA**	**Los Angeles Basin, CA**
Other Compounds								
divanillyl			19.4	3.18				
divanillyl methane			2.39	nd				
vanillylmethylguaiacol			3.24	0.568				
Total other			**25.0**	**3.75**				
N-Containing Compounds								
3-methoxypyridine	0.86	1.4						
isoquinoline	1.1	1.1						
1-methoxypyridine	0.27	0.24						
1,2-dimethoxy-4-nitro-benzene	1.8	3.9						
dihydroxynitrobenzene								1.62 (10.52)
Total N-containing compounds	**4.03**	**6.64**						**1.62**
Total Quantified Organic Compound Mass	**789**	**764**	**11410**	**2075**	**267**	**487**	**10**	**8**
Total Organic Compound Mass			**55700**	**18700**				
Percent of Organic Mass Quantified	**8-15% (a)**	**8-15% (a)**	**20%**	**11%**		**<3%**		
Percent of Organic Mass Extractable and Elutable	**45-60% (a)**	**45-60% (a)**	**30%**	**21%**				

Mean values are provided with maximum concentrations in parentheses.

[a]Rogge et al. (1993) summarized these percentages for all four Los Angeles Basin sampling sites (West LA, Downtown LA, Pasadena, and Rubidoux). Only Downtown LA and Pasadena data are shown here.

DRAFT-DO NOT QUOTE OR CITE

REFERENCES

Allen, J. O.; Dookeran, N. M.; Taghizadeh, K.; Lafleur, A. L.; Smith, K. A.; Sarofim, A. F. (1997) Measurement of oxygenated polycyclic aromatic hydrocarbons associated with a size-segregated urban aerosol. Environ. Sci. Technol. 31: 2064-2070.

Allen, G. A.; Lawrence, J.; Koutrakis, P. (1999) Field validation of a semi-continuous method for aerosol black carbon (aethalometer) and temporal patterns of summertime hourly black carbon measurements in southwestern PA. Atmos. Environ. 33: 817-823.

Andrews, E.; Saxena, P.; Musarra, S.; Hildemann, L .M.; Koutrakis, P.; McMurry, P. H.; Olmez, I.; White, W. H. (2000) Concentration and composition of atmospheric aerosols from the 1995 SEAVS experiment and a review of the closure between chemical and gravimetric measurements. J. Air Waste Manage. Assoc. 50: 648-664.

Chow, J. C.; Watson, J. G.; Lu, Z.; Lowenthal, D. H.; Frazier, C. A.; Solomon, P. A.; Thuillier, R. H.; Magliano, K. (1996) Descriptive analysis of $PM_{2.5}$ and PM_{10} at regionally representative locations during SJVAQS/AUSPEX. In: Parrish, D.; Trainer, M.; Rao, S. T.; Solomon, P. A., eds. A&WMA international specialty conference on regional photochemical measurements and modeling, part 2; November 1993; San Diego, CA. Atmos. Environ. 30: 2079-2112.

Christoforou, C. S.; Salmon, L. G.; Hannigan, M. P.; Solomon, P. A.; Cass, G. R. (2000) Trends in fine particle concentration and chemical composition in southern California. J. Air Waste Manage. Assoc. 50: 43-53.

Cui, W.; Machir, J.; Lewis , L.; Eatough, D. J.; Eatough, N. L. (1997) Fine particulate organic material at Meadview during the project MOHAVE summer intensive study. J. Air Waste Manage. Assoc. 47: 357-369.

Fraser, M. P.; Cass, G. R.; Simoneit, B. R. T.; Rasmussen, R. A. (1998) Air quality model evaluation data for organics. 5. C_6-C_{22} nonpolar and semipolar aromatic compounds. Environ. Sci. Technol. 32: 1760-1770.

Gertler, A. W.; Lowenthal, D. A.; Coulombe, W. G. (1995) PM_{10} source apportionment study in Bullhead City, Arizona. J. Air Waste Manage. Assoc. 45: 75-82.

Hegg, D. A.; Livingston, J.; Hobbs, P. V.; Novakov, T.; Russell, P. (1997) Chemical apportionment of aerosol column optical depth off the mid-Atlantic coast of the United States. J. Geophys. Res. 102: 25,293-25,303.

IMPROVE: interagency monitoring of protected visual environments [database]. (2000) [Data on particulate organic and elemental carbon concentrations after 1995]. Fort Collins, CO: National Park Service (NPS); Cooperative Institute for Research in the Atmosphere (CIRA). Available at: http://vista.cira.colostate.edu/improve/ [2001, January 26].

Khwaja, H. (1995) Atmospheric concentrations of carboxylic acids and related compounds at a semiurban site. Atmos. Environ. 29: 127-139.

Klinedinst, D. B.; Currie, L. A. (1999) Direct quantification of $PM_{2.5}$ fossil and biomass carbon within the northern front range air quality study's domain. Environ. Sci. Technol. 33: 4146-4154.

Lewtas, J.; Pang, Y.; Booth, D.; Reimer, S.; Eatough, D. J.; Gundel, L. A. (2001) Comparison of sampling methods for semi-volatile organic carbon associated with $PM_{2.5}$. Aerosol. Sci. Technol. 34: 9-22.

Lioy, P. J.; Daisey, J. M. (1987) Toxic air pollution: a comprehensive study of non-criteria air pollutants. Chelsea, MI: Lewis Publishers.

Malm, W. C.; Day, D. E. (2000) Optical properties of aerosols at Grand Canyon National Park. Atmos. Environ. 34: 3373-3391.

Malm, W. C.; Gebhart, K. A. (1996) Source apportionment of organic and light-absorbing carbon using receptor modeling techniques. Atmos. Environ. 30: 843-855.

Offenberg, J. H.; Baker, J. E. (2000) Aerosol size distributions of elemental and organic carbon in urban and over-water atmospheres. Atmos. Environ. 34: 1509-1517.

Omar, A. H.; Biegalski, S.; Larson, S. M.; Landsberger, S. (1999) Particulate contributions to light extinction and local forcing at a rural Illinois site. Atmos. Environ. 33: 2637-2646.

Pedersen, D. U.; Durant, J. L.; Penman, B. W.; Crespi, C. L.; Hemond, H. F.; Lafleur, A. L.; Cass, G. R. (1999) Seasonal and spatial variations in human cell mutagenicity of respirable airborne particles in the northeastern United States. Environ. Sci. Technol. 33: 4407-4415.

Rogge, W. F.; Mazurek, M. A.; Hildemann, L. M.; Cass, G. R.; Simoneit, B. R. T. (1993) Quantification of urban organic aerosols at a molecular level: identification, abundance and seasonal variation. Atmos. Environ. Part A 27: 1309-1330.

Saxena, P.; Hildemann, L. M. (1996) Water-soluble organics in atmospheric particles: a critical review of the literature and applications of thermodynamics to identify candidate compounds. J. Atmos. Chem. 24: 57-109.

Schauer, J. J.; Cass, G. R. (2000) Source apportionment of wintertime gas-phase and particle-phase air pollutants using organic compounds as tracers. Environ. Sci. Technol. 34: 1821-1832.

Turpin, B. J.; Huntzicker, J. J. (1995) Identification of secondary organic aerosol episodes and quantitation of primary and secondary organic aerosol concentrations during SCAQS. Atmos. Environ. 29: 3527-3544.

U.S. Environmental Protection Agency. (1996) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.

Veltkamp, P. R.; Hansen, K. J.; Barkley, R. M.; Sievers, R. E. (1996) Principal component analysis of summertime organic aerosols at Niwot Ridge, Colorado. J. Geophys. Res. [Atmos.] 101: 19,495-19,504.

APPENDIX 3D

Composition of Particulate Matter Source Emissions

This appendix includes discussions of the elemental composition of emissions from various source categories discussed in Table 3-8. Discussions in this appendix incorporate material dealing with the inorganic components of source emissions from Chapter 5 of the 1996 PM AQCD (U. S. Environmental Protection Agency, 1996), updates to that material, and material describing the composition of organic components in source emissions. Primary emphasis is placed in the discussions on the composition of $PM_{2.5}$ sources.

Soil and Fugitive Dust

The compositions of soils and average crustal material are shown in Table 3D-1 (adapted from Warneck, 1988). Two entries are shown as representations of average crustal material. Differences from the mean soil composition shown can result from local geology and climate. Major elements in both soil and crustal profiles are Si, Al, and Fe, which are found in the form of various minerals. In addition, organic matter constitutes a few percent, on average, of soils. In general, the soil profile is similar to the crustal profiles, except for the depletion of soluble elements such as Ca, Mg, Na, and K. It should be noted that the composition of soils from specific locations can vary considerably from these global averages, especially for elements like Ca, Mg, Na, and K.

Fugitive dust emissions arise from paved and unpaved roads, building construction and demolition, parking lots, mining operations, storage piles, feed lots, grain handling, and agricultural tilling, in addition to wind erosion. Figure 3D-1 shows examples of size distributions in dust from paved and unpaved roads, agricultural soil, sand and gravel, and alkaline lake bed sediments, which were measured in a laboratory resuspension chamber as part of a study in California (Chow et al., 1994). This figure shows substantial variation in particle size among some of these fugitive dust sources. The $PM_{1.0}$ abundance (6.9%) in the total suspended PM (TSP) from alkaline lake bed dust is twice its abundance in paved and unpaved road dust. Approximately 10% of the TSP is in the $PM_{2.5}$ fraction and approximately 50% of TSP is in the PM_{10} fraction. The sand/gravel dust sample shows that 65% of the mass is in particles

TABLE 3D-1. AVERAGE ABUNDANCES OF MAJOR ELEMENTS IN SOIL AND CRUSTAL ROCK

	Elemental Abundances (ppmw)		
		Crustal Rock	
Element	Soil (1)	(2)	(3)
Si	330,000	277,200	311,000
Al	71,300	81,300	77,400
Fe	38,000	50,000	34,300
Ca	13,700	36,300	25,700
Mg	6,300	20,900	33,000
Na	6,300	28,300	31,900
K	13,600	25,900	29,500
Ti	4,600	4,400	4,400
Mn	850	950	670
Cr	200	100	48
V	100	135	98
Co	8	25	12

Source: (1) Vinogradov (1959); (2) Mason (1966); (3) Turekian (1971), Model A; as quoted in Warneck (1988).

larger than the PM_{10} fraction. The $PM_{2.5}$ fraction of TSP is approximately 30 to 40% higher in alkaline lake beds and sand/gravel than in the other soil types. The tests were performed after sieving and with a short (<1 min) waiting period prior to sampling. It is expected that the fraction of $PM_{1.0}$ and $PM_{2.5}$ would increase with distance from a fugitive dust emitter as the larger particles deposit to the surface faster than do the smaller particles.

The size distribution of samples of paved road dust obtained from a source characterization study in California is shown in Figure 3D-2. As might be expected, most of the emissions are in the coarse size mode. The chemical composition of paved road dust obtained in Denver, CO, during the winter of 1987-1988 is shown in Figure 3D-3. The chemical composition of paved road dust consists of a complex mixture of particulate matter from a wide variety of sources. Hopke et al. (1980) found that the inorganic composition of urban roadway dust in samples from



Figure 3D-1. Size distribution of particles generated in a laboratory resuspension chamber.

Source: Chow et al. (1994).

Urbana, IL, could be described in terms of contributions from natural soil, automobile exhaust, rust, tire wear, and salt. Automobile contributions arose from exhaust emissions enriched in Pb; from rust as Fe; tire wear particles enriched in Zn; brake linings enriched in Cr, Ba, and Mn; and cement particles derived from roadways by abrasion. In addition to organic compounds from combustion and secondary sources, road dust also contains biological material such as pollen and fungal spores.

Very limited data exist for characterizing the composition in organic compounds found in resuspended paved road dust and soil dust. The only reported measurements are from Rogge et al. (1993a) and Schauer and Cass (2000), which consist of data for the fine particle fraction. The resuspended road dust sample analyzed by Rogge et al. (1993a) was collected in Pasadena, CA, during May of 1988. The sample analyzed by Schauer and Cass (2000) is a composite



Figure 3D-2. Size distribution of California source emissions, 1986.

Source: Houck et al. (1989, 1990).



Figure 3D-3. Chemical abundances for $PM_{2.5}$ emissions from paved road dust in Denver, CO. Solid bars represent fractional abundances, and the error bars represent variability in species abundances. Error bars represent detection limits when there are no solid bars.

Source: Watson and Chow (1994).

sample collected at several sites in the Central Valley of California in 1995. In both cases, road dust samples were resuspended in the laboratory. Samples were drawn through a $PM_{2.0}$ cyclone upstream of the collection substrate to remove particles with aerodynamic diameters greater than 2.0 μm. It is unclear if these samples are representative of road dust in other locations of the United States. Table 3D-2 summarizes the organic compounds measured in these road dust samples.

TABLE 3D-2. SUMMARY OF PARTICLE-PHASE ORGANIC COMPOUNDS PRESENT IN FINE PARTICLE ROAD DUST SAMPLE

Source	Compound Class	Contribution to Particulate Mass (%)	Dominant Contributors to Emissions of Compound Class
Pasadena Road Dust (Rogge et al., 1993a)	n-Alkanes	0.13	C_{17}, C_{19}, C_{21}
	n-Alkanoic acids	0.37	Palmitic acid and stearic acid
	n-Alkenoic acids	0.028	Oleic acid and linoleic Acid
	Petroleum biomarkers	0.017	Hopanes and steranes
	PAH	0.0059	No dominant compounds
	n-Alkanals	0.046	Octacosanol and triacontanal
	n-Alkanols	0.021	Hexacosanol and octacosanol
San Joaquin Valley Road Dust (Schauer and Cass, 2000)	n-Alkanes	0.023	No dominant compounds
	n-Alkanoic acids	0.23	Palmitic acid and stearic acid
	n-Alkenoic acids	0.095	Oleic acid, linoleic acid, and hexadecenoic acid

Stationary Sources

The elemental composition of primary particulate matter emitted in the fine fraction from a variety of power plants and industries in the Philadelphia area is shown in Table 3D-3 as a representative example of emissions from stationary fossil combustion sources (Olmez et al., 1988). Entries for the coal fired power plant show that Si and Al, followed by sulfate, are the major primary constituents produced by coal combustion; whereas fractional abundances of elemental carbon were much lower and organic carbon species were not detected. Sulfate is the major particulate constituent released by the oil fired power plants examined in this study; and, again, elemental and organic carbon are not among the major species emitted. Olmez et al. (1988) also compared their results to a number of similar studies and concluded that their data could have much wider applicability to receptor model studies in other areas with some of the same source types. The high temperature of combustion in power plants results in the almost complete oxidation of the carbon in the fuel to CO_2 and very small amounts of CO. Combustion conditions in smaller boilers and furnaces allow the emission of unburned carbon and sulfur in

TABLE 3D-3. COMPOSITION OF FINE PARTICLES RELEASED BY VARIOUS STATIONARY SOURCES IN THE PHILADELPHIA AREA

Species (Units)	Eddystone Coal-Fired Power Plant	N	Oil-Fired Power Plants				Secondary Al Plant	N	Fluid Cat. Cracker	N	Municipal Incinerator	N
			Eddystone	N	Schuylkill	N						
C-v (%)	ND		2.7 ± 1.2	3	0.75 ± 0.63	4	1.6 ± 1.5	2	ND		0.57 ± 0.26	4
C-e (%)	0.89 ± 0.12	3	7.7 ± 1.5	3	0.22 ± 0.17	4	0.18 ± 0.10	2	0.16 ± 0.05	3	3.5 ± 0.2	4
NH_4 (%)	1.89 ± 0.19	3	3.5 ± 1.6	3	3.7 ± 1.7	4	2.2 ± 0.9	2	0.43 ± 0.22	3	0.36 ± 0.07	4
Na (%)	0.31 ± 0.03	3	3.0 ± 0.8	3	3.3 ± 0.8	3	16.3 ± 0.8	1	0.38 ± 0.05	3	6.6 ± 3.5	3
Al (%)	14 ± 2	3	0.45 ± 0.09	3	0.94 ± 0.08	3	1.74 ± 0.09	1	6.8 ± 1.2	3	0.25 ± 0.10	3
Si (%)	21.8 ± 1.6	9	1.9 ± 0.6	9	2.6 ± 0.4	11	3.1 ± 2.2	2	9.8 ± 20.0	9	1.7 ± 0.3	10
P (%)	0.62 ± 0.10	9	1.5 ± 0.4	9	1.0 ± 0.2	11	0.45 ± 0.27	2	ND		0.63 ± 0.12	10
S (%)	3.4 ± 0.6	9	11 ± 2	9	13 ± 1	11	3 ± 4	2	4.2 ± 12.6	9	2.9 ± 0.8	10
SO_4 (%)	11.9 ± 1.2	3	40 ± 4	3	45 ± 7	4	5.9 ± 2	2	38 ± 4	3	6.8 ± 2.3	4
Cl (%)	0.022 ± 0.11	3	0.019 ± 0.009	2	ND		21 ± 4	1	ND		29 ± 5	3
K (%)	1.20 ± 0.09	9	0.16 ± 0.05	9	0.21 ± 0.03	11	10.9 ± 1.5	2	0.031 ± 0.005	9	7.6 ± 2.3	10
Ca (%)	1.4 ± 0.5	3	3.6 ± 1.0	3	2.3 ± 1.0	3	0.12 ± 0.09	2	0.030 ± 0.004	9	0.23 ± 0.10	10
Sc (ppm)	42 ± 2	3	0.17 ± 0.02	3	0.47 ± 0.02	3	0.092 ± 0.039	1	2.7 ± 0.4	3	0.11 ± 0.02	1
Ti (%)	1.1 ± 0.2	3	0.040 ± 0.044	9	0.12 ± 0.02	11	0.024 ± 0.003	2	0.38 ± 0.1	3	0.030 ± 0.015	10
V (ppm)	550 ± 170	3	11500 ± 3000	3	20,000 ± 3000	3	36 ± 7	1	250 ± 70	3	8.6 ± 5.3	2
Cr (ppm)	390 ± 120	3	235 ± 10	3	230 ± 70	3	410 ± 20	1	59 ± 8	3	99 ± 31	3
Mn (ppm)	290 ± 15	3	380 ± 40	3	210 ± 50	3	120 ± 15	1	14 ± 3	3	165 ± 40	3
Fe (%)	7.6 ± 0.4	3	1.6 ± 0.2	3	1.7 ± 0.4	3	0.31 ± 0.02	1	0.20 ± 0.03	9	0.22 ± 0.05	3
Co (ppm)	93 ± 10	3	790 ± 150	3	1100 ± 200	3	13 ± 2	1	15 ± 2	3	3.7 ± 0.8	3
Ni (ppm)	380 ± 50	9	15000 ± 5000	9	19000 ± 2000	11	300 ± 100	2	220 ± 30	9	290 ± 40	10

DRAFT–DO NOT QUOTE OR CITE

TABLE 3D-3 (cont'd). COMPOSITION OF FINE PARTICLES RELEASED BY VARIOUS STATIONARY SOURCES IN THE PHILADELPHIA AREA

Species (units)	Eddystone Coal-Fired Power Plant	N	Oil-Fired Power Plants				Secondary Al Plant	N	Fluid Cat. Cracker	N	Municipal Incinerator	N
			Eddystone	N	Schuylkill	N						
Cu (ppm)	290 ± 20	9	980 ± 320	9	1100 ± 500	11	450 ± 200	2	14 ± 8	9	1300 ± 500	3
Zn (%)	0.041 ± 0.005	3	1.3 ± 0.3	3	0.78 ± 0.30	3	0.079 ± 0.006	1	0.0026 ± 0.0007	3	10.4 ± 0.5	3
As (ppm)	640 ± 80	3	33 ± 6	1	50 ± 16	3	15 ± 6	1	ND		64 ± 34	3
Se (ppm)	250 ± 20	3	26 ± 9	3	23 ± 7	3	66 ± 3	1	15 ± 1	3	42 ± 16	3
Br (ppm)	35 ± 8	3	90 ± 60	9	45 ± 17	11	630 ± 70	2	5.6 ± 1.8	9	2300 ± 800	10
Rb (ppm)	190 ± 80	1	ND		ND		97 ± 38	1	ND		230 ± 50	2
Sr (ppm)	1290 ± 60	9	160 ± 50	9	280 ± 70	11	ND		36 ± 6	9	87 ± 14	10
Zr (ppm)	490 ± 190	9	140 ± 180	9	100 ± 120	11	ND		130 ± 50	2	ND	
Mo (ppm)	170 ± 60	2	930 ± 210	3	1500 ± 300	3	ND		ND		240 ± 130	10
Ag (ppm)	ND		ND		ND		ND		ND		71 ± 15	3
Cd (ppm)	ND		ND		ND		ND		ND		1200 ± 700	3
In (ppm)	0.71 ± 0.04	2	ND		ND		ND		ND		4.9 ± 1.4	3
Sn (ppm)	ND		320 ± 230	9	200 ± 80	11	550 ± 540	2	ND		6700 ± 1900	10
Sb (ppm)	[a]		370 ± 410	3	1020 ± 90	3	6100 ± 300	1	7.7 ± 1.5	3	1300 ± 1000	3
Cs (ppm)	9.2 ± 0.9	2	ND		ND		ND		ND		5.9 ± 3.0	3
Ba (ppm)	ND		1960 ± 100	3	2000 ± 500	3	ND		290 ± 90	2	ND	
La (ppm)	120 ± 10	3	130 ± 30	3	450 ± 30	3	19 ± 2	1	3300 ± 500	3	1.1 ± 0.5	1
Ce (ppm)	180 ± 10	2	89 ± 23	3	360 ± 20	3	ND		2700 ± 400	3	ND	
Nd (ppm)	80 ± 26	3	28 ± 5	2	230 ± 20	3	ND		1800 ± 250	3	ND	
Sm (ppm)	23 ± 2	3	3.7 ± 0.7	3	20.5 ± 1.5	3	ND		170 ± 20	3	ND	

DRAFT–DO NOT QUOTE OR CITE

TABLE 3D-3 (cont'd). COMPOSITION OF FINE PARTICLES RELEASED BY VARIOUS STATIONARY SOURCES IN THE PHILADELPHIA AREA

Species (units)	Eddystone Coal-Fired Power Plant	N	Oil-Fired Power Plants				Secondary Al Plant	N	Fluid Cat. Cracker	N	Municipal Incinerator	N
			Eddystone	N	Schuylkill	N						
Eu (ppm)	5.1 ± 0.5	3	ND		0.65 ± 0.23	3	ND		4.9 ± 0.7	3	ND	
Gd (ppm)	ND		ND		ND		ND		71 ± 10	3	ND	
Tb (ppm)	3.3 ± 0.3	3	ND		0.90 ± 0.29	3	ND		8.9 ± 1.3	3	ND	
Yb (ppm)	10.3 ± 0.5	1	ND		ND		ND		3.7 ± 0.4	3	ND	
Lu (ppm)	ND		ND		ND		ND		0.59 ± 0.17	3	ND	
Hf (ppm)	5.8 ± 0.8	3	0.39 ± 0.07	1	ND		ND		0.99 ± 0.08	3	ND	
Ta (ppm)	ND		ND		ND		ND		0.56 ± 0.10	3	ND	
W (ppm)	20 ± 8	1	60 ± 5	2	ND		ND		ND		ND	
Au (ppm)	ND		0.054 ± 0.017	2	ND		ND		ND		0.56 ± 0.27	3
Pb (%)	0.041 ± 0.004	9	1.8 ± 0.6	9	1.0 ± 0.2	11	0.081 ± 0.014	2	0.0091 ± 0.0021	9	5.8 ± 1.2	10
Th (ppm)	24 ± 2	3	1.9 ± 0.5	2	ND		ND		6.2 ± 0.7	3	ND	
% mass	24 ± 2	6	93.5 ± 2.5	6	96 ± 2	6	81 ± 10	2	97 ± 2	7	89 ± 2	7

[a]Omitted because of sample contamination.

N = Number of samples.
ND = Not detected.
The "% mass" entries give the average percentage of the total emitted mass found in the fine fraction.

Source: Adapted from Olmez et al. (1988).

DRAFT–DO NOT QUOTE OR CITE

more reduced forms such as thiophenes and inorganic sulfides. A number of trace elements are greatly enriched over crustal abundances in different fuels, such as Se in coal and V, Zn, and Ni in oil. In fact, the higher V content of the fuel oil than in coal could help account for the higher sulfate seen in the profiles from the oil-fired power plant compared to the coal-fired power plant because V at combustion temperatures found in power plants is known to catalyze the oxidation of reduced sulfur species. During combustion at lower temperatures, the emission of reduced sulfur species also occurs. For example, Huffman et al. (2000) identified sulfur species emitted by the combustion of several residual fuels oil (RFO) in a fire tube package boiler that is meant to simulate conditions in small institutional and industrial boilers. They found that sulfur was emitted not only as sulfate (26 to 84%), but as thiophenes (13 to 39%) with smaller amounts of sulfides and elemental S. They also found that Ni, V, Fe, Cu, Zn, and Pb are present mainly as sulfates in emissions. Linak et al. (2000) found, when burning RFO, that the fire tube package boiler produced particles with a bimodal size distribution in which about 0.2% of the mass was associated with particles smaller than 0.1 μm AD, with the rest of the mass lying between 0.5 and 100 μm. Miller et al. (1998) found that larger particles consisted mainly of cenospheric carbon; whereas trace metals and sulfates were found concentrated in the smaller particles in a fire tube package boiler. In contrast, when RFO was burning in a refractory-lined combustor that is meant to simulate combustion conditions in a large utility residual oil fired boiler, Linak et al. (2000) found that particles were distributed essentially unimodally, with a mean diameter of about 0.1 μm.

Apart from emissions in the combustion of fossil fuels, trace elements are emitted as the result of various industrial processes such as steel and iron manufacturing and nonferrous metal production (e.g., for Pb, Cu, Ni, Zn, and Cd). As may be expected, emissions factors for the various trace elements are highly source-specific (Nriagu and Pacyna, 1988). Inspection of Table 3D-3 reveals that the emissions from the catalytic cracker and the oil-fired power plant are greatly enriched in rare-earth elements such as La compared to other sources.

Emissions from municipal waste incinerators are heavily enriched in Cl, arising mainly from the combustion of plastics and metals that form volatile chlorides. The metals can originate from cans or other metallic objects, and some metals such as Zn and Cd are also additives in plastics or rubber. Many elements such as S, Cl, Zn, Br, Ag, Cd, Sn, In, and Sb are enormously enriched compared to their crustal abundances. A comparison of the trace elemental composition

of incinerator emissions in Philadelphia, PA (shown in Table 3D-3), with the composition of incinerator emissions in Washington DC, and Chicago, IL (Olmez et al., 1988), shows agreement for most constituents to better than a factor of two.

Very limited data exist for characterizing the chemical composition of organic compounds present in particulate emissions from industrial-scale stationary fuel combustion. Oros and Simoneit (2000) have presented the abundance and distribution of organic constituents in coal smokes that have been burned under laboratory conditions. This work provides the basis for further investigation addressing the emissions of coal fired boilers.

Rogge et al. (1997a) measured the composition of the organic constituents in the particulate matter emissions from a 50 billion kj/h boiler that was operating at 60% capacity and was burning number 2 distillate fuel oil. The fine carbon particulate matter emissions from this boiler over five tests were composed of an average of 14% organic carbon and 86% elemental carbon (Hildemann et al., 1991). Significant variability in the distribution of organic compounds present in the emissions from two separate tests was observed. Most of the identified organic mass consisted of n-alkanonic acids, aromatic acids, n-alkanes, PAH, oxygeanted PAH, and chlorinated compounds. It is unclear if these emissions are representative of typical fuel oil combustion units in the United States. Rogge et al. (1997b) measured the composition of hot asphalt roofing tar pots, and Rogge et al. (1993b) measured the composition of emissions from home appliances that use natural gas.

Motor Vehicles

Exhaust emissions of particulate matter from gasoline powered motor vehicles and diesel powered vehicles have changed significantly over the past 25 years (Sawyer and Johnson, 1995; Cadle et al., 1999). These changes have resulted from reformulation of fuels, the wide application of exhaust-gas treatment in gasoline-powered motor vehicles, and changes in engine design and operation. Because of these evolving tailpipe emissions, along with the wide variability of emissions between vehicles of the same class (Hildemann et al., 1991; Cadle et al., 1997; Sagebiel et al., 1997; Yanowitz et al., 2000), well-defined average emissions profiles for the major classes of motor vehicles have not been established. Two sampling strategies have been employed to obtain motor vehicle emissions profiles: (1) the measurement of exhaust emissions from vehicles operating on dynamometers and (2) the measurement of integrated

emissions of motor vehicles driving through roadway tunnels. Dynamometer testing can be used to measure vehicle emissions operating over an integrated driving cycle and allows the measurement of emissions from individual vehicles. However, dynamometer testing requires considerable resources and usually precludes testing a very large number of vehicles. In contrast, a large number of vehicles can be readily sampled in tunnels; however, vehicles driving through tunnels operate over limited driving conditions, and the measurements represent contributions from a large number of vehicle types. As a result, except in a few cases, tunnel tests have not been effective at developing chemically speciated particulate matter emissions profiles for individual motor vehicle classes. Rather, several studies have measured the contribution of both organic and elemental carbon to the particulate matter emissions from different classes of motor vehicles operating on chassis dynamometers.

The principal components emitted by diesel and gasoline fueled vehicles are organic carbon (OC) and elemental carbon (EC) as shown in Tables 3D-4a and 4b. As can be seen, the variability among entries for an individual fuel type is large and overlaps that found between different fuel types. On average, the abundance of elemental carbon is larger than that of organic carbon in the exhaust of diesel vehicles; whereas organic carbon is the dominant species in the exhaust of gasoline fueled vehicles. Per vehicle mile, total carbon emissions from light and heavy duty diesel vehicles can range from 1 to 2 orders of magnitude higher than those from gasoline vehicles.

As might be expected, most of the PM emitted by motor vehicles is in the $PM_{2.5}$ size range. Particles in diesel exhaust are typically trimodal (consisting of a nuclei mode, an accumulation mode, and a coarse mode) and are log-normal in form (Kittelson, 1998). More than 90% of the total number of particles are in the nuclei mode, which contains only about 1 to 20% of the particle mass with a mass median diameter of about 0.02 μm; whereas the accumulation mode (with a mass median diameter of about 0.25 μm) contains most of the mass with a smaller fraction (5 to 20%) contained in the coarse mode. Kerminin et al. (1997), Bagley et al. (1998), and Kleeman et al. (2000) also have shown that gasoline and diesel fueled vehicles produce particles that are mostly less than 2.0 μm in diameter. Cadle et al. (1999) found that 91% of PM emitted by in-use gasoline vehicles in the Denver area was in the $PM_{2.5}$ size range, which increased to 97% for “smokers” (i.e., light-duty vehicles with visible smoke emitted from their tailpipes) and 98% for light-duty diesels. Durbin et al. (1999) found that about 92% of the PM

TABLE 3D-4a. ORGANIC AND ELEMENTAL CARBON FRACTIONS OF DIESEL AND GASOLINE ENGINE PARTICULATE MATTER EXHAUST

	Organic Carbon	Elemental Carbon
Heavy-duty diesel engines[a]	19 ± 8%	75 ± 10%
Heavy-duty diesel engines (SPECIATE)[b]	21 - 36%	52 - 54%
Light-duty diesel engines[c]	30 ± 9%	61 ± 16%
Light-duty diesel engines (SPECIATE)[b]	22 - 43%	51 - 64%
Gasoline engines (hot stabilized)[a]	56 ± 11%	25 ± 15%
Gasoline engines ("smoker" and "high emitter")[a,c]	76 ± 10%	7 ± 6%
Gasoline engines (cold start)[a]	46 ± 14%	42 ± 14%

[a]Fujita et al. (1998) and Watson et al. (1998).
[b]U.S. EPA SPECIATE database.
[c]Norbeck et al. (1998).

Source: U.S. Environmental Protection Agency (2002).

was smaller than 2.5 μm for smokers and diesels. The mass median diameter of the PM emitted by the gasoline vehicles sampled by Cadle et al. (1999) was about 0.12 μm and increased to 0.18 μm for smokers and diesels. Corresponding average emissions rates of $PM_{2.5}$ found by Cadle et al. (1999) were 552 mg/mile for diesels; 222 mg/mile for gasoline smokers; and 38 mg/mile for other gasoline vehicles. The values for gasoline smokers and for diesels appear to be somewhat lower than those given in Table 3D-5; whereas the value for other gasoline vehicles falls in the range given for low and medium gasoline vehicle emissions.

Examples of data for the trace elemental composition of the emissions from a number of vehicle classes obtained December 1997 in Colorado, as part of the North Frontal Range Air Quality Study (NFRAQS), are shown in Table 3D-5. As can be seen from Table 3D-5, emissions of total carbon (TC), which is equal to the sum of organic carbon (OC) and elemental carbon (EC), from gasoline vehicles are highly variable. Gillies and Gertler (2000) point out that there is greater variability in the concentrations of trace elements and ionic species than for OC and EC among different source profiles (e.g., SPECIATE, Lawson and Smith [1998], Norbeck et al. [1998]). They suggest that this may arise because emissions of trace elements are not related only to the combustion process, but also to their abundances in different fuels and lubricants and

TABLE 3D-4b. CONTRIBUTION OF ORGANIC CARBON TO PARTICULATE MATTER CARBON EMISSIONS IN MOTOR VEHICLE EXHAUST COLLECTED FROM VEHICLES OPERATED ON CHASSIS DYNAMOMETERS

	Year of Tests	Test Cycle	Number of Vehicles	OC % of Total Carbon	Notes
GASOLINE POWERED VEHICLES					
Light-duty vehicles	1996-97	FTP	195[a]	70	A
High-CO/VOC-emitting smokers	1994	IM-240	7	91	B
High-CO/VOC-emitting nonsmokers	1994	IM-240	15	76	B
Catalyst-equipped vehicles	Mid-1980s	FTP	7	69	C
Noncatalyst vehicles	Mid-1980s	FTP	6	89	C
DIESEL VEHICLES					
Light-duty diesel vehicles	1996-1997	FTP	195[a]	40	A
Medium-duty diesel vehicles	1996	FTP	2	50[b]	D
Heavy-duty diesel vehicles	1992	[c]	6	42	E
Heavy-duty diesel vehicles	Mid-1980s	[c]	2	45	C

Notes:
A. From Cadle et al. (1999). Average of summer and winter cold start emissions.
B. From Sagebiel et al. (1997). Hot start testing of vehicles identified as either high emitters of carbon monoxide or volatile organic compounds (VOCs).
C. From Hildemann et al. (1991). Cold start tests.
D. From Schauer et al. (1999). Hot start tests of medium duty vehicles operating on an FTP cycle.
E. From Lowenthal et al. (1994). Only includes measurement of vehicles powered by diesel fuel operated without an exhaust particulate trap.

[a]A total of 195 light duty vehicles were tested that include both gasoline powered vehicles and diesel powered vehicles.
[b]Fraction of particulate matter consisting of organic carbon was measured with and without an organics denuder upstream of particulate filter. Results reported here represent measurement without an organics denuder for consistency with other measurements. Using an organics denuder, the organic carbon comprised 39% of the particulate matter carbon.
[c]Driving cycle comprised of multiple idle, steady acceleration, constant speed, deceleration steps (see reference for more details).

to wear and tear during vehicle operation. Emissions from gasoline smokers are comparable to those from light-duty diesel vehicles. Thus, older, poorly maintained gasoline vehicles could be significant sources of $PM_{2.5}$ (Sagebiel et al., 1997; Lawson and Smith, 1998), in addition to being

TABLE 3D-5. EMISSION RATES (mg/mi) FOR CONSTITUENTS OF PARTICULATE MATTER FROM GASOLINE AND DIESEL VEHICLES

	Gasoline Vehicles				Diesel Vehicles	
	Low	Medium	High	Smoker	Light Duty	Heavy Duty
TC	9.07 ± 0.75	41.30 ± 1.68	207.44 ± 7.29	456.38 ± 16.80	373.43 ± 13.75	1570.69 ± 58.24
OC	6.35 ± 0.54	26.02 ± 1.31	95.25 ± 4.28	350.24 ± 15.27	132.01 ± 5.82	253.94 ± 16.12
EC	2.72 ± 0.52	15.28 ± 0.99	112.19 ± 5.82	106.14 ± 5.42	241.42 ± 12.11	1316.75 ± 55.33
NO_3^-	0.039 ± 0.027	0.057 ± 0.028	0.141 ± 0.031	0.964 ± 0.051	1.474 ± 0.071	1.833 ± 1.285
$SO_4^=$	0.158 ± 0.036	0.518 ± 0.043	0.651 ± 0.052	2.160 ± 0.137	2.902 ± 0.165	3.830 ± 1.286
Na	0.060 ± 0.063	0.023 ± 0.111	0.052 ± 0.092	0.000 ± 0.000	0.000 ± 0.000	1.288 ± 2.160
Mg	0.036 ± 0.022	0.068 ± 0.027	0.041 ± 0.033	0.000 ± 0.000	0.000 ± 0.000	1.061 ± 0.729
Al	0.083 ± 0.016	0.078 ± 0.016	0.057 ± 0.014	0.000 ± 0.000	0.000 ± 0.000	0.321 ± 0.543
Si	0.066 ± 0.008	0.279 ± 0.011	0.714 ± 0.012	0.000 ± 0.000	0.000 ± 0.000	8.018 ± 0.221
P	0.035 ± 0.004	0.152 ± 0.007	0.113 ± 0.007	0.000 ± 0.000	0.000 ± 0.000	0.407 ± 0.136
S	0.085 ± 0.006	0.442 ± 0.009	0.822 ± 0.022	2.515 ± 0.116	2.458 ± 0.124	3.717 ± 0.111
Cl	0.024 ± 0.012	0.038 ± 0.012	0.081 ± 0.020	0.140 ± 0.117	0.228 ± 0.114	0.881 ± 0.221
K	0.010 ± 0.009	0.019 ± 0.009	0.031 ± 0.035	0.033 ± 0.386	0.000 ± 0.426	0.064 ± 0.248
Ca	0.060 ± 0.010	0.212 ± 0.011	0.210 ± 0.030	0.362 ± 0.250	0.150 ± 0.304	0.716 ± 0.107
Fe	0.143 ± 0.004	0.756 ± 0.005	1.047 ± 0.010	2.438 ± 0.054	0.515 ± 0.057	0.376 ± 0.055
Ni	0.001 ± 0.004	0.005 ± 0.004	0.011 ± 0.005	0.008 ± 0.017	0.014 ± 0.018	0.002 ± 0.057
Cu	0.002 ± 0.004	0.016 ± 0.003	0.021 ± 0.005	0.071 ± 0.018	0.024 ± 0.021	0.001 ± 0.062
Zn	0.048 ± 0.003	0.251 ± 0.004	0.265 ± 0.023	0.188 ± 0.272	0.000 ± 0.299	0.707 ± 0.032
Br	0.001 ± 0.002	0.016 ± 0.002	0.079 ± 0.003	0.047 ± 0.012	0.003 ± 0.014	0.012 ± 0.050
Ba	0.013 ± 0.136	0.009 ± 0.138	0.011 ± 0.299	0.380 ± 2.175	0.428 ± 2.390	0.493 ± 3.108
Pb	0.007 ± 0.006	0.085 ± 0.005	0.255 ± 0.008	0.345 ± 0.032	0.153 ± 0.033	0.008 ± 0.154

Source: Lawson and Smith (1998).

significant sources of gaseous pollutants (e.g., Calvert et al., 1993). Durbin et al. (1999) point out that although smokers constitute only 1.1 to 1.7% of the light-duty fleet in the South Coast Air Quality Management District in California, they contribute roughly 20% of the total PM

emissions from the light-duty fleet. In general, motor vehicles that are high emitters of hydrocarbons and carbon monoxide also will tend to be high emitters of PM (Sagebiel et al.,1997; Cadle et al., 1997). Particle emission rates, even in newer vehicles, also are correlated with vehicle acceleration; and emissions occur predominantly during periods of heavy acceleration (Maricq et al., 1999).

Although the data shown in Table 3D-5 indicate that S (mainly in the form of sulfate) is a minor component of $PM_{2.5}$ emissions, S may be the major component of the ultrafine particles that are emitted by either diesel or internal combustion engines (Gertler et al., 2000). It is not clear what the source of the small amount of Pb seen in the auto exhaust profile is. It is extremely difficult to find suitable tracers for automotive exhaust because Pb has been removed from gasoline. However, it also should be remembered that restrictions in the use of leaded gasoline have resulted in a dramatic lowering of ambient Pb levels.

Several tunnel studies have measured the distribution of organic and elemental carbon in the integrated exhaust of motor vehicle fleets comprising several classes of motor vehicles (Pierson and Brachaczek, 1983; Weingartner et al., 1997a; Fraser et al., 1998a). The study by Fraser et al. (1998a) found that organic carbon constituted 46% of the carbonaceous PM emissions from the vehicles operating in the Van Nuys tunnel in Southern California in the Summer of 1993. Although diesel vehicles constituted only 2.8% of the vehicles measured by Fraser et al. (1998a), the contribution of the organic carbon to the total particulate carbon emissions obtained in the Van Nuys tunnels is in reasonable agreement with the dynamometer measurements shown in Table 3D-4b.

Very few studies have reported comprehensive analyses of the organic composition of motor vehicle exhaust. The measurements by Rogge et al. (1993c) are the most comprehensive but are not expected to be the best representation of current motor vehicle emissions because these measurements were made in the mid-1980s. Measurements reported by Fraser et al. (1999) were made in a tunnel study conducted in 1993 and represent integrated diesel and gasoline powered vehicle emissions. In addition, exhaust emissions from two medium-duty diesel vehicles operating over an FTP cycle were analyzed by Schauer et al. (1999). A unique feature of both the measurements by Faser et al. (1999) and Schauer et al. (1999) is that they include the quantification of unresolved complex mixture (UCM), which comprises aliphatic and cyclic hydrocarbons that cannot be resolved by gas chromatography (Schauer et al., 1999). Schauer

et al. (1999) have shown that all of the organic compound mass in their diesel exhaust samples could be extracted and eluted by CG/MS techniques even though not all of the organic compound mass can identified on a single compound basis. Table 3D-6 summarizes the composition of motor vehicle exhaust measured by Fraser et al. (1999) and Schauer et al. (1999).

TABLE 3D-6. SUMMARY OF PARTICLE-PHASE ORGANIC COMPOUNDS EMITTED FROM MOTOR VEHICLES

Source	Compound Class	Contribution to Particulate Mass (%)	Dominant Contributors to Emissions of Compound Class
Gasoline and diesel-powered vehicles driving through the Van Nuys Tunnel (Fraser et al., 1999)[a]	n-Alkanes	0.009	C_{21} through C_{29}
	Petroleum biomarkers	0.078	Hopanes and steranes
	PAH	0.38	No dominant compound
	Aromatic acids	0.29	Benzenedicarboxylic acids
	Aliphatic acids	0.21	Palmitic and stearic acids
	Substituted aromatic	0.042	No dominant compound
	UCM[b]	23.0	
Medium-duty diesel vehicles operated over an FTP Cycle (Schauer et al., 1999)	n-Alkanes	0.22	C_{20} through C_{28}
	Petroleum biomarkers	0.027	Hopanes and steranes
	PAH	0.54	No dominant compound
	Aliphatic acids	0.24	n-Octadecanoic acid
	Aromatic acids	0.014	Methylbenzoic acid
	Saturated cycloalkanes	0.037	C_{21} through C_{25}
	UCM[b]	22.2	

[a]Includes emissions of brake wear, tire wear, and resuspension of road dust associated with motor vehicle traffic.
[b]Unresolved complex mixture.

Several studies have measured the distribution of polycyclic aromatic hydrocarbons (PAHs) in motor vehicles exhaust from on-road vehicles (Westerholm et al., 1991; Lowenthal et al., 1994; Venkataraman et al., 1994; Westerholm and Egeback, 1994; Reilly et al., 1998; Cadle

et al., 1999, Weingartner et al., 1997b; Marr et al., 1999). Cadle et al. (1999) found high molecular weight PAHs (PAHs with molecular weights greater than or equal to 202 g/mole) constitute 0.1 to 7.0% of the particulate matter emissions from gasoline powered and diesel powered light duty vehicles. It is important to note, however, that PAHs with molecular weights of 202 (fluoranthene, acephenanthrylene, and pyrene), 226 (benzo[ghi]fluoranthene and cyclopenta[cd]pyrene), and 228 (benz[*a*]anthracene, chrysene, and triphenylene) exist in both the gas-phase and particle-phase at atmospheric conditions (Fraser et al., 1998b) although those with molecular weight of 228 are predominantly associated with particles, with only traces in the gas-phase (Arey et al., 1987). Excluding these semivolatile PAHs, the contribution of nonvolatile PAHs to the particulate matter emitted from the light-duty vehicles sampled by Cadle et al. (1999) ranges from 0.013 to 0.18%. These measurements are in good agreement with the tunnel study conducted by Fraser et al. (1999) and the heavy-duty diesel truck and bus exhaust measurements by Lowenthal et al. (1994), except that the nonvolatile PAH emissions from the heavy duty diesel vehicles tested by Lowenthal et al. (1994) were moderately higher, making up approximately 0.30% of the particulate matter mass emissions.

Biomass Burning

In contrast to the mobile and stationary sources discussed earlier, emissions from biomass burning in wood stoves and forest fires are strongly seasonal and can be highly episodic within their peak emissions seasons. The burning of fuelwood is confined mainly to the winter months and is acknowledged to be a major source of ambient air particulate matter in the northwestern United States during the heating season. Forest fires occur primarily during the driest seasons of the year in different areas of the country and are especially prevalent during prolonged droughts. PM produced by biomass burning outside the United States (e.g., in Central America during the spring of 1988) also can affect ambient air quality in the United States.

An example of the composition of fine particles ($PM_{2.5}$) produced by wood stoves is shown in Figure 3D-4. These data were obtained in Denver during the winter of 1987-1988 (Watson and Chow, 1994). As was the case for motor vehicle emissions, organic and elemental carbon are the major components of particulate emissions from wood burning. It should be remembered that the relative amounts shown for organic carbon and elemental carbon vary with the type of stove, the stage of combustion, and the type and condition of the fuelwood. Fine particles are



Figure 3D-4. Chemical abundances for $PM_{2.5}$ emissions from wood burning in Denver, CO. Solid bars represent fractional abundances, and the error bars represent variability in species abundances. Error bars represent detection limits when there are no solid bars.

Source: Watson and Chow (1994).

dominant in smoke studies of wood burning emissions. For instance, the mass median diameter of wood particles was found to be about 0.17 μm in a study of the emissions from burning hardwood, softwood, and synthetic logs (Dasch, 1982).

Kleeman et al. (1999) showed that the particles emitted by the combustion of wood in fireplaces are predominately less than 1.0 μm in diameter, such that the composition of fine PM ($PM_{2.5}$) emitted from fireplace combustion of wood is representative of the total particulate matter emissions from this source. Hildemann et al. (1991) and McDonald et al. (2000) reported that smoke from fireplace and wood stove combustion consists of 48% to 71% OC and 2.9% to 15% EC. Average elemental and organic carbon contents for these measurements are shown in Table 3D-7. It should be noted that the two methods used for the measurements shown in Table 3D-7 have been reported to produce different relative amounts of OC and EC for wood

TABLE 3D-7. MASS EMISSIONS, ORGANIC CARBON, AND ELEMENTAL CARBON EMISSIONS FROM RESIDENTIAL COMBUSTION OF WOOD

Wood Type	Combustion Type	Average Mass Emission Rate (g kg^{-1} of wood burned)	Number of Tests	Percent Organic Carbon[a]	Percent Elemental Carbon[a]	References
Softwood	Fireplace	13.0	2	48.4	5.2	Hildemann et al. (1991)
Softwood	Fireplace	5.14	5	58.5	15.0	McDonald et al. (2000)
Hardwood	Fireplace	5.28	3	48.4	2.9	Hildemann et al. (1991)
Hardwood	Fireplace	5.66	5	63.2	7.0	McDonald et al. (2000)
Hardwood	Wood Stove	3.96	8	71.2	9.0	McDonald et al. (2000)

[a]Hildemann et al. (1991) used the method described by Birch and Cary (1996) to measure EC and McDonald et al. (2000) used the method reported by Chow et al. (1993) to measure OC.

smoke samples but show good agreement for total carbon (OC + EC) measurements (Chow et al., 1993).

Hawthorne et al. (1988) and Hawthorne et al. (1989) measured gas-phase and particle-phase derivatives of guaiacol (2-methoxyphenol), syringol (2,6-dimethoxyphenol), phenol, and catechol (1,2-benzenediol) in the downwind plume of 28 residential wood stoves and fireplaces. Rogge et al. (1998) reported a broad range of particle-phase organic compounds in the wood smoke samples collected by Hildemann et al. (1991), which include n-alkanes, n-alkanoic acids, n-alkenoic acids, dicarboxylic acids, resin acids, phytosterols, polycyclic aromatic hydrocarbons (PAH), and the compounds reported by Hawthorne et al. (1989). Supplementing these measurements, McDonald et al. (2000) reported the combined gas-phase and particle-phase emissions of PAH and the compounds quantified by Hawthorne et al. (1989). The measurements by Rogge et al. (1998), which represent a comprehensive data set of the organic compounds present in wood smoke aerosol, are summarized in Table 3D-8. It should be noted, however, that these nearly 200 compounds account for only approximately 15 to 25% of the organic carbon particle mass emitted from the residential combustion of wood. Simoneit et al. (1999) have shown that levoglucosan constitutes a noticeable portion of the organic compound mass not identified by Rogge et al. (1998). In addition, Elias et al. (1999) used high-temperature gas chromatography/mass spectrometry (HTGC-MS) to measure high-molecular-weight organic

TABLE 3D-8. SUMMARY OF PARTICLE-PHASE ORGANIC COMPOUNDS EMITTED FROM THE COMBUSTION OF WOOD IN FIREPLACES*

Biomass Type	Compound Class	Contribution to Particulate Mass (%)	Dominant Contributors to Emissions of Compound Class
Fireplace combustion of softwood	n-Alkanes	0.039	C_{21} through C_{31}
	n-Alkanoic acids	0.45	C_{16}, C_{18}, C_{20}, C_{21}, C_{22}, C_{24}
	n-Alkenoic acids	0.12	Oleic and linoleic acid
	Dicarboxylic acids	0.36	Malonic acid
	Resin acids	1.28	Abietic, dehydroabietic, isopimaric, pimaric, and sandaracopimaric acids
	Substituted phenols	3.30	Benzenediols and guaiacols
	Phytosterols	0.37	-Sitosterol
	PAH	0.092	Fluoranthene and pyrene
	Oxygenated PAH	0.019	1H-phenalen-1-one
Fireplace combustion of hardwood	n-Alkanes	0.044	C_{21} through C_{29}
	n-Alkanoic acids	1.33	C_{16}, C_{22}, C_{24}, C_{26}
	n-Alkenoic acids	0.049	Oleic and linoleic acid
	Dicarboxylic acids	0.42	Succinic acid
	Resin acids	0.11	Dehydroabietic acid
	Substituted phenols	8.23	Benzediols, guaiacols, and syringols
	Phytosterols	0.21	-sitosterol
	PAH	0.13	No dominant compounds
	Oxygenated PAH	0.020	1H-phenalen-1-one

*Note: Measurements were made using a dilution sampler and no semivolatile organic compound sorbent.

Source: Rogge et al. (1998).

compounds in smoke from South American leaf and stem litter biomass burning. These compounds cannot be measured by the analytical techniques employed by Rogge et al. (1998) and, therefore, are strong candidates to make up some of the unidentified organic mass in the wood smoke samples analyzed by Rogge et al. (1998). These compounds, which include

triterpenyl fatty acid esters, wax esters, triglycerides, and high-molecular-weight n-alkan-2-ones, are expected to be present in North American biomass smoke originating from agricultural burning, forest fires, grassland fires, and wood stove/fireplace smoke.

Measurements of aerosol composition, size distributions, and aerosol emissions factors have been made in biomass burning plumes, either on towers (Susott et al., 1991) or aloft on fixed-wing aircraft (e.g., Radke et al., 1991) or on helicopters (e.g., Cofer et al., 1988). As was found for wood stove emissions, the composition of biomass burning emissions is strongly dependent on the stage of combustion (i.e., flaming, smoldering, or mixed), and the type of vegetation (e.g., forest, grassland, scrub). Over 90% of the dry mass in particulate biomass burning emissions is composed of organic carbon (Mazurek et al., 1991). Ratios of organic carbon to elemental carbon are highly variable, ranging from 10:1 to 95:1, with the highest ratio found for smoldering conditions and the lowest for flaming conditions. Emissions factors for total particulate emissions increase by factors of two to four in going from flaming to smoldering stages in the individual fires studied by Susott et al. (1991).

Particles in biomass burning plumes from a number of different fires were found to have three distinguishable size modes: (1) a nucleation mode, (2) an accumulation mode, and (3) a coarse mode (Radke et al., 1991). Based on an average of 81 samples, approximately 70% of the mass was found in particles <3.5 μm in aerodynamic diameter. The fine particle composition was found to be dominated by tarlike, condensed hydrocarbons; and the particles were usually spherical in shape. Additional information for the size distribution of particles produced by vegetation burning is shown in Figure 3D-2.

An example of ambient data for the composition of $PM_{2.5}$ collected at a tropical site that was heavily affected by biomass burning is shown in Table 3D-9. The samples were collected during November of 1997 on the campus of Sriwijaya University, which is located in a rural setting on the island of Sumatra in Indonesia (Pinto et al., 1998). The site was subjected routinely to levels of $PM_{2.5}$ well in excess of the U.S. NAAQS as a result of the Indonesian biomass fires from the summer of 1997 through the spring of 1998. As can be seen from a comparison of the data shown in Table 3D-9 with those shown in Figure 3D-4, there are a number of similarities and differences (especially with regard to the heavy metal content) in the abundances of many species. The abundances of some crustal elements (e.g., Si, Fe) are higher

TABLE 3D-9. MEAN AEROSOL COMPOSITION AT TROPICAL SITE (SRIWIJAYA UNIVERSITY, SUMATRA, INDONESIA) AFFECTED HEAVILY BY BIOMASS BURNING EMISSIONS[a]

Component	Abundance (%)	Component	Abundance (%)
OC	76	Cr	BD[b]
EC	1.2	Mn	BD[b]
SO_4^{-2}	11	Fe	3.9×10^{-2}
Al	BD[b]	Ni	$<3.8 \times 10^{-5}$
Si	9.3×10^{-2}	Cu	4.8×10^{-4}
Cl	4.4	Zn	3.1×10^{-3}
K	0.7	As	6.4×10^{-4}
Ca	4.5×10^{-2}	Se	2.8×10^{-4}
Ti	4.2×10^{-3}	Br	3.6×10^{-2}
V	BD[b]	Pb	3.1×10^{-3}

[a]The mean $PM_{2.5}$ concentration during the sampling period (November 5 through 11, 1997) was 264 $\mu g/m^3$.
[b]Beneath detection limit.

Source: Pinto et al. (1998).

in Table 3D-9 than in Figure 3D-4, perhaps reflecting additional contributions of entrained soil dust.

Limited emissions data that include organic compound speciation information have been reported for agricultural burning (Jenkins et al., 1996), forest fires (Simoneit, 1985), and grassland burning (Standley and Simoneit, 1987). Jenkins et al. (1996) present PAH emissions factors for the combustion of cereals (barley, corn, rice, and wheat), along with PAH emissions factors for wood burning. Profiles of organic compounds in emissions from meat cooking (Rogge et al., 1991) and cigarette smoke (Rogge et al., 1994) also have been obtained.

Natural Sources

Although sea-salt aerosol production is confined to salt water bodies, it is included here because many marine aerosols can exert a strong influence on the composition of the ambient aerosol in coastal areas. In some respects, the production of sea-salt aerosols is like that of windblown dust, in that both are produced by wind agitation of the surface. The difference between the two categories arises because sea-salt particles are produced from the bursting of air bubbles rising to the sea surface. Air bubbles are formed by the entrainment of air into the water by breaking waves. The surface energy of a collapsing bubble is converted to kinetic energy in the form of a jet of water that can eject drops above the sea surface. The mean diameter of the jet drops is about 15% of the bubble diameter (Wu, 1979). Bubbles in breaking waves range in size from a few μm to several mm in diameter. Field measurements by Johnson and Cooke (1979) of bubble size spectra show maxima in diameters at around 100 μm, with the bubble size distribution varying as $(d/d_0)^{-5}$ with $d_0 = 100$ μm.

Because sea-salt particles receive water from the surface layer, which is enriched in organic compounds, aerosol drops are composed of this organic material in addition to sea salt (about 3.5% by weight in seawater). Na^+ (30.7%), Cl^- (55.0%), SO_4^{-2} (7.7%), Mg^{2+} (3.6%), Ca^{2+} (1.2%), K^+ (1.1%), HCO_3^- (0.4%), and Br^- (0.2%) are the major ionic species by mass in seawater (Wilson, 1975). The composition of the marine aerosol also reflects the occurrence of displacement reactions that enrich sea-salt particles in SO_4^{-2} and NO_3^- while depleting them of Cl^- and Br^-.

Sea salt is concentrated in the coarse size mode with a mass median diameter of about 7 μm for samples collected in Florida, the Canary Islands, and Barbados (Savoie and Prospero, 1982). The size distribution of sulfate is distinctly bimodal. Sulfate in the coarse mode is derived from sea water, but sulfate in the submicron aerosol arises from the oxidation of dimethyl sulfide (CH_3SCH_3) or DMS. DMS is produced during the decomposition of marine micro-organisms. DMS is oxidized to methane sulfonic acid (MSA), a large fraction of which is oxidized to sulfate (e.g., Hertel et al., 1994).

Apart from sea spray, other natural sources of particles include the suspension of organic debris and volcanism. Profiles of organic compounds in vegetative detritus have been obtained by Rogge et al. (1993d). Particles are released from plants in the form of seeds, pollen, spores, leaf waxes, and resins, ranging in size from 1 to 250 μm (Warneck, 1988). Fungal spores and

animal debris, such as insect fragments, also are to be found in ambient aerosol samples in this size range. Although material from all the foregoing categories may exist as individual particles, bacteria usually are found attached to other dust particles (Warneck, 1988). Smaller bioaerosol particles include viruses, individual bacteria, protozoa, and algae (Matthias-Maser and Jaenicke, 1994). In addition to natural sources, other sources of bioaerosol include industry (e.g., textile mills), agriculture, and municipal waste disposal (Spendlove, 1974). The size distribution of bioaerosols has not been characterized as well as it has for other categories of airborne particles.

Trace metals are emitted to the atmosphere from a variety of sources such as sea spray, wind-blown dust, volcanoes, wildfires and biotic sources (Nriagu, 1989). Biologically mediated volatilization processes (e.g., biomethylation) are estimated to account for 30 to 50% of the worldwide total Hg, As, and Se emitted annually; whereas other metals are derived principally from pollens, spores, waxes, plant fragments, fungi, and algae. It is not clear, however, how much of the biomethylated species are remobilized from anthropogenic inputs. Median ratios of the natural contribution to globally averaged total sources for trace metals are estimated to be 0.39 (As), 0.15 (Cd), 0.59 (Cr), 0.44 (Cu), 0.41 (Hg), 0.35 (Ni), 0.04 (Pb), 0.41 (Sb), 0.58 (Se), 0.25 (V), and 0.34 (Zn), suggesting a significant natural source for many trace elements. It should be noted, however, that these estimates are based on emissions estimates that have uncertainty ranges of an order of magnitude.

REFERENCES

Arey, J.; Zielinska, B.; Atkinson, R.; Winer, A. M. (1987) Polycyclic aromatic hydrocarbon and nitroarene concentrations in ambient air during a wintertime high-NO_x episode in the Los Angeles basin. Atmos. Environ. 21: 1437-1444.

Bagley, S. T.; Gratz, L. D.; Johnson, J. H.; McDonald J. F. (1998) Effects of an oxidation catalytic converter and a biodiesel fuel on the chemical, mutagenic, and particle size characteristics of emissions from a diesel engine. Environ. Sci. Technol. 32: 1183-1191.

Birch, M. E.; Cary, R. A. (1996) Elemental carbon-based method for occupational monitoring of particulate diesel exhaust: methodology and exposure issues. Analyst 121: 1183-1190.

Cadle, S. H.; Mulawa, P. H.; Ball, J.; Donase, C.; Weibel, A.; Sagebiel, J. C.; Knapp, K. T.; Snow, R. (1997) Particulate emission rates from in-use high-emitting vehicles recruited in Orange county, California. Environ. Sci. Technol. 31: 3405-3412.

Cadle, S. H.; Mulawa, P.; Hunsanger, E. C.; Nelson, K.; Ragazzi, R. A.; Barrett, R.; Gallagher, G. L.; Lawson, D. R.; Knapp, K. T.; Snow, R. (1999) Light-duty motor vehicle exhaust particulate matter measurement in the Denver, Colorado, area. J. Air Waste Manage. Assoc. 49: PM-164-174.

Calvert, J. G.; Heywood, J. B.; Sawyer, R. F.; Seinfeld, J. H. (1993) Achieving acceptable air quality: some reflections on controlling vehicle emissions. Science (Washington, DC) 261: 37-45.

Chow, J. C.; Watson, J. G.; Pritchett, L. C.; Pierson, W. R.; Frazier, C. A.; Purcell, R. G. (1993) The DRI thermal/optical reflectance carbon analysis system: description, evaluation and applications in U.S. air quality studies. In: Lodge, J. P., Jr. Fourth international conference on carbonaceous particles in the atmosphere; April 1991; Vienna, Austria. Atmos. Environ. Part A 27: 1185-1201.

Chow, J. C.; Watson, J. G.; Houck, J. E.; Pritchett, L. C.; Rogers, C. F.; Frazier, C. A.; Egami, R. T.; Ball, B. M. (1994) A laboratory resuspension chamber to measure fugitive dust size distributions and chemical compositions. Atmos. Environ. 28: 3463-3481.

Cofer, W. R., III; Levine, J. S.; Sebacher, D. I.; Winstead, E. L.; Riggin, P. J.; Brass, J. A.; Ambrosia, V. G. (1988) Particulate emissions from a mid-latitude prescribed chaparral fire. J. Geophys. Res. [Atmos.] 93: 5207-5212.

Dasch, J. M. (1982) Particulate and gaseous emissions from wood-burning fireplaces. Environ. Sci. Technol. 16: 639-645.

Durbin, T. D.; Smith, M. R.; Norbeck, J. M.; Truex, T. J. (1999) Population density, particulate emission characterization, and impact on the particulate inventory of smoking vehicles in the South Coast Air Quality Management District. J. Air Waste Manage. Asssoc. 49: 28-38.

Elias, V. O.; Simoneit, B. R. T.; Pereira, A. S.; Cabral, J. A.; Cardoso, J. N. (1999) Detection of high molecular weight organic tracers in vegetation smoke samples by high-temperature gas chromatography-mass spectrometry. Environ. Sci. Technol. 33: 2369-2376.

Fraser, M. P.; Cass, G. R.; Simoneit, B. R. T. (1998a) Gas-phase and particle-phase organic compounds emitted from motor vehicle traffic in a Los Angeles roadway tunnel. Environ. Sci. Technol. 32: 2051-2060.

Fraser, M. P.; Cass, G. R.; Simoneit, B. R. T.; Rasmussen, R. A. (1998b) Air quality model evaluation data for organics. 5. C6-C22 nonpolar and semipolar aromatic compounds. Environ. Sci. Technol. 32: 1760-1770.

Fraser, M. P.; Cass, G. R.; Simoneit, B. R. T. (1999) Particulate organic compounds emitted from motor vehicle exhaust and in the urban atmosphere. Atmos. Environ. 33: 2715-2724.

Fujita, E.; Watson, J. G.; Chow, J. C.; Robinson, N. F.; Richards, L. W.; Kumar, N. (1998) Northern front range air quality study, volume C: source apportionment and simulation methods and evaluation. Reno, NV: Desert Research Institute; prepared for Colorado State University, Cooperative Institute for Research in the Atmosphere.

Gertler, A. W.; Abu-Allaban, M.; Coulombe, W.; Gilles, J. A.; Pierson, W. R.; Rogers, C. F.; Sagebiel, J. C.; Tarnay, L. (2000) Preliminary results of a tunnel study to characterize mobile source particulate emissions. Presented at: PM2000: particulate matter and health—the scientific basis for regulatory decision-making, specialty conference & exhibition; January; Charleston, SC. Pittsburgh, PA: Air & Waste Management Association.

Gillies, J. A.; Gertler, A. W. (2000) Comparison and evaluation of chemically speciated mobile source $PM_{2.5}$ particulate matter profiles. J. Air Waste Manage. Assoc. 50: 1459-1480.

Hawthorne, S. B.; Miller, D. J.; Barkley, R. M.; Krieger, M. S. (1988) Identification of methoxylated phenols as candidate tracers for atmospheric wood smoke pollution. Environ. Sci. Technol. 22: 1191-1196.

Hawthorne, S. B.; Krieger, M. S.; Miller, D. J.; Mathiason, M. B. (1989) Collection and quantitation of methoxylated phenol tracers for atmospheric pollution from residential wood stoves. Environ. Sci. Technol. 23: 470-475.

Hertel, O.; Christensen, J.; Hov, Ø. (1994) Modelling of the end products of the chemical decomposition of DMS in the marine boundary layer. Atmos. Environ. 28: 2431-2449.

Hildemann, L. M.; Markowski, G. R.; Jones, M. C.; Cass, G. R. (1991) Submicrometer aerosol mass distributions of emissions from boilers, fireplaces, automobiles, diesel trucks, and meat-cooking operations. Aerosol Sci. Technol. 14: 138-152.

Hopke, P. K.; Lamb, R. E.; Natusch, D. F. S. (1980) Multielemental characterization of urban roadway dust. Environ. Sci. Technol. 14: 164-172.

Houck, J. E.; Chow, J. C.; Watson, J. G.; Simons, C. A.; Pritchett, L. C.; Goulet, J. M.; Frazier, C. A. (1989) Determination of particle size distribution and chemical composition of particulate matter from selected sources in California: volume I and executive summary (final report). Sacramento, CA: California Air Resources Board.

Houck, J. E.; Goulet, J. M.; Chow, J. C.; Watson, J. G.; Pritchett, L. C. (1990) Chemical characterization of emission sources contributing to light extinction. In: Mathai, C. V., ed. Visibility and fine particles: an A&WMA/EPA international specialty conference; October 1989; Estes Park, CO. Pittsburgh, PA: Air & Waste Management Association; pp. 437-446. (A&WMA transactions series no. TR-17).

Huffman, G. P.; Huggins, F. E.; Shah, N.; Huggins, R.; Linak, W. P.; Miller, C. A.; Pugmire, R. J.; Meuzelaar, H. L. C.; Seehra, M. S.; Manivannan, A. (2000) Characterization of fine particulate matter produced by combustion of residual fuel oil. J. Air Waste Manage. Assoc. 50: 1106-1114.

Jenkins, B. M.; Jones, A. D.; Turn, S. Q.; Williams, R. B. (1996) Particle concentrations, gas-particle partitioning, and species intercorrelations for polycyclic aromatic hydrocarbons (PAH) emitted during biomass burning. Atmos. Environ. 30: 3825-3835.

Johnson, B. D.; Cooke, R. C. (1979) Bubble populations and spectra in coastal waters: a photographic approach. J. Geophys. Res. C: Oceans Atmos. 84: 3761-3766.

Kerminen, V. M.; Makela, T. E.; Ojanen, C. H.; Hillamo, R. E.; Vilhunen, J. K.; Rantanen, L.; Havers, N.; Von Bohlen, A.; Klockow, D. (1997) Characterization of the particulate phase in the exhaust from a diesel car. Environ. Sci. Technol. 31: 1883-1889.

Kittelson, D. B. (1998) Engines and nanoparticles: a review. J. Aerosol Sci. 29: 575-588.

Kleeman, M. J.; Schauer, J. J.; Cass, G. R. (1999) Size and composition distribution of fine particulate matter emitted from wood burning, meat charbroiling, and cigarettes. Environ. Sci. Technol. 33: 3516-3523.

Kleeman, M. J.; Schauer, J. J.; Cass, G. R. (2000) Size and composition distribution of fine particulate matter emitted from motor vehicles. Environ. Sci. Technol. 34: 1132-1142.

Lawson, D. R.; Smith, R. E. (1998) The northern front range air quality study: a report to the Governor and General Assembly. Fort Collins, CO: Colorado State University; December.

Linak, W. P.; Miller, C. A.; Wendt, J. O. L. (2000) Comparison of particle size distributions and elemental partitioning from the combustion of pulverized coal and residual fuel oil. J. Air Waste Manage. Assoc. 50: 1532-1544.

Lowenthal, D. H.; Zielinska, B.; Chow, J. C.; Watson, J. G.; Gautam, M.; Ferguson, D. H.; Neuroth, G. R.; Stevens, K. D. (1994) Characterization of heavy-duty diesel vehicle emissions. Atmos. Environ. 28: 731-743.

Maricq, M. M.; Podsiadlik, D. H.; Chase, R. E. (1999) Gasoline vehicle particle size distributions: comparison of steady state, FTP, and US06 measurements. Environ. Sci. Technol. 33: 2007-2015.

Marr, L. C.; Kirchstetter, T. W.; Harley, R. A.; Miguel, A. H.; Hering, S. V.; Hammond, S. K. (1999) Characterization of polycyclic aromatic hydrocarbons in motor vehicle fuels and exhaust emissions. Environ. Sci. Technol. 33: 3091-3099.

Mason, B. (1966) Principles of geochemistry. 3rd ed. New York, NY: John Wiley & Sons, Inc.

Matthias-Maser, S.; Jaenicke, R. (1994) Examination of atmospheric bioaerosol particles with radii > 0.2 μm. J. Aerosol Sci. 25: 1605-1613.

Mazurek, M. A.; Cofer, W. R., III; Levine, J. S. (1991) Carbonaceous aerosols from prescribed burning of a boreal forest ecosystem. In: Levine, J. S., ed. Global biomass burning: atmospheric, climatic, and biospheric implications. Cambridge, MA: MIT Press; pp. 258-263.

McDonald, J. D.; Zielinska, B.; Fujita, E. M.; Sagebiel, J. C.; Chow, J. C.; Watson, J. G. (2000) Fine particle and gaseous emission rates from residential wood combustion. Environ. Sci. Technol. 34: 2080-2091.

Miller, C. A.; Linak, W. P.; King, C.; Wendt, J. O. L. (1998) Fine particle emissions from heavy fuel oil combustion in a firetube package boiler. Combust. Sci. Technol. 134: 477-502.

Norbeck, J. M.; Durbin, T. D.; Truex, T. J. (1998) Measurement of primary particulate matter emissions from light-duty motor vehicles. Riverside, CA: University of California, College of Engineering, Center for Environmental Research and Technology; prepared for Coordinating Research Council, Inc. and South Coast Air Quality Management District.

Nriagu, J. O. (1989) A global assessment of natural sources of atmospheric trace metals. Nature (London) 338: 47-49.

Nriagu, J. O.; Pacyna, J. M. (1988) Quantitative assessment of worldwide contamination of air, water and soils by trace metals. Nature (London) 333: 134-139.

Olmez, I.; Sheffield, A. E.; Gordon, G. E.; Houck, J. E.; Pritchett, L. C.; Cooper, J. A.; Dzubay, T. G.; Bennett, R. L. (1988) Compositions of particles from selected sources in Philadelphia for receptor modeling applications. JAPCA 38: 1392-1402.

Oros, D. R.; Simoneit, B. R. T. (2000) Identification and emission rates of molecular tracers in coal smoke particulate matter. Fuel 79: 515-536.

Pierson, W. R.; Brachaczek, W. W. (1983) Particulate matter associated with vehicles on the road. II. Aerosol Sci. Technol. 2: 1-40.

Pinto, J. P.; Grant, L. D.; Hartlage, T. A. (1998) Report on U.S. EPA air monitoring of haze from S.E. Asia biomass fires. Research Triangle Park, NC: U.S. Environmental Protection Agency, National Center for Environmental Assessment; report no. EPA/600/R-98/071.

Radke, L. F.; Hegg, D. A.; Hobbs, P. V.; Nance, J. D.; Lyons, J. H.; Laursen, K. K.; Weiss, R. E.; Riggan, P. J.; Ward, D. E. (1991) Particulate and trace gas emissions from large biomass fires in North America. In: Levine, J. S., ed. Global biomass burning: atmospheric, climatic, and biospheric implications. Cambridge, MA: MIT Press; pp. 209-224.

Reilly, P. T. A.; Gieray, R. A.; Whitten, W. B.; Ramsey, J. M. (1998) Real-time characterization of the organic composition and size of individual diesel engine smoke particles. Environ. Sci. Technol. 32: 2672-2679.

Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1991) Sources of fine organic aerosol. 1. Charbroilers and meat cooking operations. Environ. Sci. Technol. 25: 1112-1125.

Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993a) Sources of fine organic aerosol. 3. Road dust, tire debris, and organometallic brake lining dust: roads as sources and sinks. Environ. Sci. Technol. 27: 1892-1904.

Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993b) Sources of fine organic aerosol. 5. Natural gas home appliances. Environ. Sci. Technol. 27: 2736-2744.

Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993c) Sources of fine organic aerosol. 2. Noncatalyst and catalyst-equipped automobiles and heavy-duty diesel trucks. Environ. Sci. Technol. 27: 636-651.

Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993d) Sources of fine organic aerosol. 4. Particulate abrasion products from leaf surfaces of urban plants. Environ. Sci. Technol. 27: 2700-2711.

Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1994) Sources of fine organic aerosol. 6. Cigarette smoke in the urban atmosphere. Environ. Sci. Technol. 28: 1375-1388.

Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1997a) Sources of fine organic aerosol. 8. Boilers burning no. 2 distillate fuel oil. Environ. Sci. Technol. 31: 2731-2737.

Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1997b) Sources of fine organic aerosol. 7. Hot asphalt roofing tar pot fumes. Environ. Sci. Technol. 31: 2726-2730.

Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1998) Sources of fine organic aerosol. 9. Pine, oak, and synthetic log combustion in residential fireplaces. Environ. Sci. Technol. 32: 13-22.

Sagebiel, J. C.; Zielinska, B.; Walsh, P. A.; Chow, J. C.; Cadle, S. H.; Mulawa, P. A.; Knapp, K. T.; Zweidinger, R. B.; Snow, R. (1997) PM-10 exhaust samples collected during IM-240 dynamometer tests of in-service vehicles in Nevada. Environ. Sci. Technol. 31: 75-83.

Savoie, D. L.; Prospero, J. M. (1982) Particle size distribution of nitrate and sulfate in the marine atmosphere. Geophys. Res. Lett. 9: 1207-1210.

Sawyer, R. F.; Johnson, J. H. (1995) Diesel emissions and control technology. Diesel exhaust: a critical analysis of emissions, exposure, and health effects. Cambridge, MA: Health Effects Institute.

Schauer, J. J.; Cass, G. R. (2000) Source apportionment of wintertime gas-phase and particle-phase air pollutants using organic compounds as tracers. Environ. Sci. Technol. 34: 1821-1832.

Schauer, J. J.; Kleeman, M. J.; Cass, G. R.; Simoneit, B. R. T. (1999) Measurement of emissions from air pollution sources. 2. C_1 through C_{30} organic compounds from medium duty diesel trucks. Environ. Sci. Technol. 33: 1578-1587.

Simoneit, B. R. T. (1985) Application of molecular marker analysis to vehicular exhaust for source reconciliations. Int. J. Environ. Anal. Chem. 22: 203-233.

Simoneit, B. R. T.; Schauer, J. J.; Nolte, C. G.; Oros, D. R.; Elias, V. O.; Fraser, M. P.; Rogge, W. F.; Cass, G. R. (1999) Levoglucosan, a tracer for cellulose in biomass burning and atmospheric particles. Atmos. Environ. 33: 173-182.

Spendlove, J. C. (1974) Industrial, agricultural, and municipal microbial aerosol problems. In: Current problems in aerobiology: proceedings of the thirtieth general meeting of the Society for Industrial Microbiology; August 1973; Evanston, IL. Dev. Ind. Microbiol. 15: 20-27.

Standley, L. J.; Simoneit, B. R. T. (1987) Characterization of extractable plant wax, resin, and thermally matured components in smoke particles from prescribed burns. Environ. Sci. Technol. 21: 163-169.

Susott, R. A.; Ward, D. E.; Babbitt, R. E.; Latham, D. J. (1991) The measurement of trace emissions and combustion characteristics for a mass fire. In: Levine, J. S., ed. Global biomass burning: atmospheric, climatic, and biospheric implications. Cambridge, MA: MIT Press; pp, 245-257.

Turekian, K. K. (1971) Elements, geochemical distribution of. In: Encyclopedia of science and technology, v. 4. New York, NY: McGraw-Hill Book Company; pp. 627-630.

U.S. Environmental Protection Agency. (1996) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.

U.S. Environmental Protection Agency. (2002) Health assessment document for diesel engine exhaust. Washington, DC: Office of Research and Development; report no. EPA/600/8-90/057.

Venkataraman, C.; Lyons, J. M.; Friedlander, S. K. (1994) Size distributions of polycyclic aromatic hydrocarbons and elemental carbon. 1. Sampling, measurement methods, and source characterization. Environ. Sci. Technol. 28: 555-562.

Vinogradov, A. P. (1959) The geochemistry of rare and dispersed chemical elements in soils. 2nd ed. New York, NY: Consultants Bureau, Inc.

Warneck, P. (1988) Chemistry of the natural atmosphere. New York, NY: Academic Press, Inc.

Watson, J. G.; Chow, J. C. (1994) Clear sky visibility as a challenge for society. Annu. Rev. Energy Environ. 19: 241-266.

Watson, J. G.; Fujita, E. M.; Chow, J. C.; Zielinska, B.; Richards, L. W.; Neff, W.; Dietrich, D. (1998) Northern fron range air quality study final report. Reno, NV: Desert Research Institute; prepared for Colorado State University, Cooperative Institute for Research in the Atmosphere.

Weingartner, E.; Burtscher, H.; Baltensperger, U. (1997a) Hygroscopic properties of carbon and diesel soot particles. Atmos. Environ. 31: 2311-2327.

Weingartner, E.; Keller, C.; Stahel, W. A.; Burtscher, H.; Baltensperger, U. (1997b) Aerosol emission in a road tunnel. Atmos. Environ. 31: 451-462.

Westerholm, R.; Egebäck, K.-E. (1994) Exhaust emissions from light- and heavy-duty vehicles: chemical composition, impact of exhaust after treatment, and fuel parameters. In: Symposium of risk assessment of urban air: emissions, risk identification, and risk quantitation; May-June 1992; Stockholm, Sweden. Environ. Health Perspect. 102(suppl. 4) 13-23.

Westerholm, R. N.; Almén, J.; Li, H.; Rannug, J. U.; Egebäck, K.-E.; Grägg, K. (1991) Chemical and biological characterization of particulate-, semivolatile-, and gas-phase-associated compounds in diluted heavy-duty diesel exhausts: a comparison of three different semivolatile-phase samplers. Environ. Sci. Technol. 25: 332-338.

Wilson, T. R. S. (1975) Salinity and the major elements of sea water. In: Riley, J. P.; Skirrow, G., eds. Chemical oceanography: v. 1. 2nd ed. London, United Kingdom: Academic Press Inc.; pp. 365-413.

Wu, J. (1979) Spray in the atmospheric surface layer: review and analysis of laboratory and oceanic results. J. Geophys. Res. C: Oceans Atmos. 84: 1693-1704.

Yanowitz, J.; McCormick, R. L.; Grabowski, M. S. (2000) In-use emissions from heavy-duty diesel vehicles. Environ. Sci. Technol. 34: 729-740.

4. ENVIRONMENTAL EFFECTS OF PARTICULATE MATTER

4.1 INTRODUCTION

Several later chapters (Chapters 5 through 8) of this document assess the latest available information on determinants of human exposures to particulate matter; the dosimetry of particle deposition, clearance, and retention in human respiratory tract; toxicologic evaluations of pathophysiologic effects of PM and underlying mechanisms of action; and epidemiologic analyses of health effects associated with human exposures to ambient PM. The human exposure and health-related findings assessed in those chapters provide key elements of the scientific bases to support decision making regarding review of the primary PM National Ambient Air Quality Standards (PM NAAQS). This chapter, in contrast, assesses information pertinent to decision making regarding secondary standards aimed at protecting against welfare effects of PM. More specifically, this chapter assesses environmental effects of atmospheric PM, including PM effects on vegetation and ecosystems, effects on visibility, and on man-made materials, as well as relationships of ambient PM to global climate change processes.

4.2 IMPACTS ON VEGETATION AND ECOSYSTEMS

The PM NAAQS first set in 1971 were specified in terms of total suspended particulates (TSP), which included both fine and coarse mode particles (the latter ranging up to 25 to 40 μm in size). The 1987 revision of the PM NAAQS to PM_{10} standards focused attention on those particles ($\leq$10 μm mean aerometic diameter) capable of being deposited in lower (thoracic) portions of the human respiratory tract. The subsequent 1997 PM NAAQS revisions retained the PM_{10} standards and added fine particle ($PM_{2.5}$) standards (both specified in terms of mass concentrations of particles undifferentiated in terms of their specific chemical composition). The effects of PM on vegetation and ecosystems as a basis for a secondary standard were not considered as part of the 1997 PM NAAQS revisions. Vegetation and ecosystem effects of ambient PM evaluated in this chapter are dependent not simply on PM size-related mass

concentration, but rather on exposure of plants to PM components differentiated by chemical composition as well.

This section deals with PM deposition and effects on individual plants in natural habitats and terrestrial ecosystems. Except for the deposition of nitrogen and sulfur-containing compounds and their effects exerted via acidic precipitation, information concerning the effects of deposition of other specific substances as PM on crops is not readily available. An extensive overall discussion of the effects of acidic deposition is presented in the U.S. National Acid Precipitation Assessment Program (NAPAP) Biennial Report to Congress: An Integrated Assessment (National Science and Technology Council, 1998). The effects of gaseous sulfur oxides and nitrogen oxides on crops are discussed in detail in EPA criteria documents for those substances (U.S. Environmental Protection Agency, 1982, 1993). Detailed discussion of ecological effects of acidic precipitation and nitrate deposition on aquatic ecosystems can also be found in the EPA Nitrogen Oxides Air Quality Criteria Document (U.S. Environmental Protection Agency, 1993). Neither nitrate nor sulfate deposition on crops is discussed in this chapter, as they are frequently added in fertilizers. Lead effects on crops, vegetation, and ecosystems are discussed in the EPA document, Air Quality Criteria for Lead (U.S. Environmental Protection Agency, 1986). Also, the effects of "certain pesticides, metal compounds, chlorinated organic compounds, and nitrogen compounds" are discussed in Deposition of Air Pollutants to the Great Waters, Third Report to Congress (U.S. Environmental Protection Agency, 2000a).

4.2.1 Particle Deposition

This subsection reviews interactions between vegetation and the fine (<2.5 μm) and coarse (>2.5 μm) components of airborne particulate matter (PM) that lead to deposition. Particulate matter has not been defined by chemical nature, structure, or source; it has been defined mainly by size fraction. While size is related to mode and magnitude of deposition to vegetated landscapes and may be a useful surrogate for chemical constitution (Whitby, 1978; U.S. Environmental Protection Agency, 1996a), the size classes have no specific relevance to vegetation. Both fine- and coarse-mode particles may affect plants. An evaluation of particulate deposition to plants and vegetated surfaces is presented because the determinants of deposition

ultimately determine the magnitude of both contact effects and soil mediated effects of PM on vegetation.

Particulate matter deposition to vegetation is not well understood. A recent review emphasizes semivolatile organics and early work on radio nuclide deposition (Smith and Jones, 2000). Atmospheric deposition of particles to ecosystems takes place via both wet and dry processes through three major routes: (1) precipitation scavenging in which particles are deposited in rain and snow; (2) fog, cloud-water, and mist interception (i.e., "occult" deposition); and (3) much slower dry deposition. Unlike gaseous dry deposition, neither the solubility of the particle, nor the physiological activity of the surface are likely to be of first order importance in determining particulate dry deposition velocity (V_d). Factors that contribute to surface wetness or stickiness may be critical determinants of deposition efficiency. Available tabulations of deposition velocity are highly variable and suspect. Recent evidence indicates that all three modes of deposition (wet, occult, and dry) must be considered in determining inputs to water sheds or ecosystems, because each may dominate over specific intervals of time or space.

4.2.1.1 Wet Deposition

Wet deposition results from the incorporation of atmospheric particles and gases into cloud droplets and their subsequent precipitation as rain or snow, or from the scavenging of particles and gases by raindrops or snowflakes as they fall (Lovett, 1994). Precipitation scavenging, in which particles are incorporated in hydrometeors and deposited in the resulting rain and snow, includes rainout (within-cloud incorporation by nucleation) and washout (below-cloud scavenging by impaction). Wet deposition generally is confounded by fewer factors than dry or occult deposition and has been easier to quantify. Total inputs from wet deposition to vegetative canopies can be significant (Table 4-1) although not all wet deposition involves particle scavenging because gaseous pollutants also dissolve in raindrops during precipitation events (Lovett, 1994). This contribution is obscured during measurements because wet deposition is measured simply by chemical analysis of total precipitation collected in clean, non-reactive buckets. Exclusion of dry deposited material (as opposed to dissolved gaseous species) requires closure or covering of the vessels except during periods of precipitation.

Wet deposition is not affected by surface properties as much as is dry or occult deposition although leaves retain liquid and solubilized PM according to their surface properties of

TABLE 4-1. RELATIVE IMPORTANCE OF WET, DRY, PARTICULATE, AND TOTAL DEPOSITION TO THREE FOREST SITES[a]

	Deposition							
	Total Nitrogen[b]				Total Sulfur[c]			
Site	Wet (%)	Dry (%)	Particle (%)	Total (kg ha^{-1})	Wet (%)	Dry (%)	Particle (%)	Total (kg ha^{-1})
Duke Forest	75	25	0.11	9.87	64	33	2.7	17.20
Cary Forest	71	20	0.94	5.80	76	20	4.2	7.60
Austin Forest	71	29	0.58	6.57	83	13	4.3	7.79

[a]Data from Allen et al. (1994). Sampling was by triple filter pack so that fine-mode particles could be sampled preferentially. An average particle deposition velocity of 0.9 cm s^{-1} was derived as in Hicks et al. (1987).
[b]Wet nitrogen consists of NO_3^- and NH_4^+; dry nitrogen consists of vapor phase HNO_3 and NO_2; and particulate nitrogen consists of NO_3^-.
[c]Wet sulfur consists of SO_4^{-2}, dry sulfur consists of vapor phase SO_2, and particulate sulfur consists of $pSO_4^{-2.}$

wettability, exposure, and roughness. Wet deposition is largely a function of precipitation amount and ambient pollutant concentrations. Any material deposited in precipitation to the upper stratum of foliage will likely be intercepted by several foliar surfaces before reaching the soil, because extensive vegetative canopies typically develop leaf area indices (LAI; ratio of projected leaf area to ground area) much in excess of 1.

Landscape characteristics may also be important. Forested hillsides receive four- to six-fold greater inputs of wet deposition than short vegetation in nearby valleys. This is due to a variety of orographic effects (Unsworth and Wilshaw, 1989) and closer aerodynamic coupling to the atmosphere of tall forest canopies than of the shorter canopies in the valleys. This leads to more rapid foliar drying, which reduces the residence time but concentrates more quickly the solubilized particulate materials available for foliar uptake on the cuticular surface; and concentration increases the thermodynamic driving force for foliar uptake (Fowler et al., 1991; Unsworth, 1984; Schönherr and Huber, 1977). Humidity and temperature conditions following wet deposition strongly influence the extent of biological effects, reflecting the competing effects of drying versus concentrating the solutions and influencing the rate of metabolic uptake of surface solutes (Swietlik and Faust, 1984). The net consequence of these factors on direct physical effects of wet deposited PM on leaves is not known.

Rainfall introduces wet deposition and redistributes throughout the canopy previously dry-deposited particulate material, particularly coarse particles which are preferentially deposited in the upper foliage (Peters and Eiden, 1992). Both effects scale the likelihood of foliar contact and potential direct PM effects on vegetation nearly linearly with canopy leaf area. The concentrations of suspended and dissolved materials are typically highest at the onset and decline with duration of individual precipitation events (Lindberg and McLaughlin, 1986; Hansen et al., 1994). Sustained rainfall removes much of the accumulation of dry-deposited PM from foliar surfaces, reducing direct foliar effects and combining the associated chemical burden with the wet-deposited material (Lovett and Lindberg, 1984; Lovett, 1994) for transfer to the soil. Intense rainfall may contribute substantial total particulate inputs to vegetated land surfaces, mostly via the soil, but is less effective as a source of directly bioavailable or injurious pollutants to foliar surfaces. This washing effect, combined with differential foliar uptake and foliar leaching of different chemical constituents of PM, alter the composition of the rainwater that reaches the soil. Low intensity precipitation events, in contrast, may be of greater significance for direct effects of foliar-deposited particulate pollutants to foliar surfaces. Because of the short duration and limited atmospheric cleansing, the concentration of PM in the final precipitation that remains in contact with foliar surfaces may be high. Additionally, such events may hydrate some previously dry-deposited particles without removing them and thereby facilitate their foliar uptake.

This combination of dry deposition to foliage and subsequent wet removal enhances the soil pathway for PM effects, first by enhancing dry deposition relative to adjacent unvegetated surfaces and then by accelerating passage along with wet deposited material of the deposited PM by throughfall and stemflow to the soil where important soil-mediated, ecosystem-level biogeochemical cycles of major, minor, and trace elements may be affected.

4.2.1.2 Dry Deposition

Dry deposition of atmospheric particles to plant and soil is a much slower process than wet or occult deposition, but it acts nearly continuously and affects all exposed surfaces (Hicks, 1986). In dry deposition, particles at the large end of the spectrum (i.e., > 5 μm diameter) are deposited mainly by gravitational sedimentation and inertial impaction. Smaller particles, especially those with diameters between ≈ 0.2 and 2 μm, are not readily dry-deposited and tend to travel long distances in the atmosphere until their eventual deposition, most likely by

incorporation into precipitation. This long-distance transport of fine aerosols is largely responsible for the regional nature of acid deposition (Lovett, 1994). A major conclusion from atmospheric deposition research is the realization that dry deposition is usually a significant and, in some cases, a dominant portion of total atmospheric deposition to an ecosystem (Lovett, 1994). Plant parts of all types, including those not currently physiologically active, along with exposed soil and water surfaces, receive steady deposits of dry dusts, elemental carbon encrustations, grease films, tarry acidic coatings, and heterogeneous secondary particles formed from gaseous precursors (U.S. Environmental Protection Agency, 1982). The range of particle sizes, the diversity of canopy surfaces, and the variety of chemical constituents in airborne PM have slowed progress in both prediction and measurement of dry particulate deposition. Particulate deposition is a complex, poorly characterized process controlled primarily by atmospheric stability, macro- and micro-surface roughness, particle diameter, and surface characteristics (Table 4-2; Hosker and Lindberg, 1982). Deposition of particles suspended regionally and throughout the full depth of the planetary boundary layer (PBL) is controlled by different mechanisms within the three distinct atmospheric transport zones above the surface. In the lower atmosphere, fine particles are transported by turbulent eddies of mechanical and convective origin. In the relatively unstirred, laminar boundary layer surrounding individual surface elements, Brownian diffusion dominates. Near the surface, actual deposition and contact with the surface is mediated by impaction (El-Shobokshy, 1985).

Deposition fluxes may be calculated from measurements, estimates, or modeled values of mass concentration (C) at a specified measurement height and the total conductance or deposition velocity (V_d) from this height to the surface (Eq. 4-1; Hicks et al., 1987). These modeling techniques are closely allied with the micrometeorological techniques used to measure such fluxes. The flux (F) may be inferred as:

$$F = V_d * (C_z - C_o), \qquad (4\text{-}1)$$

where F is flux to the surface, C_z is the particle concentration at measurement height z, C_o is the particle concentration at receptor sites in the canopy (usually assumed equal to 0), and V_d is the overall deposition velocity. The flux is controlled by V_d and C_z.

TABLE 4-2. KEY DETERMINANTS OF DRY PARTICULATE DEPOSITION TO VEGETATION

Ambient concentration	Proximity/strength of source Timing/intensity of precipitation
Atmospheric conditions	Wind speed/turbulence Stability/mixing height Temperature/humidity
Aerosol properties	Chemical reactivity/solubility Aerodynamic diameter/diffusivity/sedimentation Biological availability
Vegetation characteristics	Roughness/plant-branch spacing/flexibility Roughness/leaf shape/pubescence Salt/organic exudates/dew

Adapted from Lindberg and McLaughlin (1986).

Vertical transport of particles through the lower atmosphere to the vicinity of the vegetation elements is by turbulence and sedimentation, such that:

$$V_d = V_t + V_s, \tag{4-2}$$

in which V_t (inner, left hand pathway of Figure 4-1) is a turbulent diffusion term, and V_s is a sedimentation term that dominates deposition of very coarse particles (Figure 4-2) and increases with particle size (Figure 4-3; dotted line). Sedimentation may be considered a pathway parallel to turbulent transport (Figure 4-2), but this is an over simplification. V_s affects the concentration of particles near the surface where eddy transport may occur and also governs the redeposition of some fraction of the particles lost to resuspension or rebound following deposition by impaction. For this reason, V_s is included (Figure 4-1) in the composite surface resistance term ($R_a R_{cp} V_s$) as well as in the parallel sedimentation term.

For submicron particles for which sedimentation is negligible (Hicks et al., 1987; Monteith and Unsworth, 1990; Wesely, 1989), the Ohm's Law Analogy (resistance catena) analogous to



Figure 4-1. A simplified resistance catena representing the factors controlling deposition of particles to the surface. Vegetation-specific parameters are not explicitly considered. Modified after Hicks et al. (1987).

that used to describe transport of heat, momentum, or gases may be adequate, as:

$$V_d = V_t = [r_a + r_b + r_c]^{-1}, \qquad (4\text{-}3)$$

where V_t is the deposition velocity due to turbulent transport of particles or other entities through the atmosphere; r_a is aerodynamic resistance (inverse of conductance or velocity) associated with the efficiency of turbulent transport above the canopy; r_b is the boundary layer resistance associated with diffusional transport through the still air layer immediately adjacent to canopy elements; and r_c is canopy resistance associated with physiological control of leaf porosity largely stromata in the leaf surface. Significant departures from the analogy arise near the surface (Chamberlain, 1975; Sehmel, 1980), as particles that were transported efficiently by turbulent eddies are slowed substantially in the laminar boundary layer that reduces the efficiency of



Figure 4-2. The relationship between deposition velocity of selected particulate materials and the distribution of the material between the coarse- and fine-aerosol fractions. Data from Foltescu et al. (1994). Ranges for Mn and Fe are from Davidson and Wu (1989).

impaction. The preservation of momentum in this zone declines with decreasing diameter; however, this is offset by an increase in Brownian diffusivity with decreasing diameter (Figure 4-3). Aerodynamic streamlines are parallel to the surface of each roughness element, so that deposition ultimately depends on diffusion to the surface. The transition from impaction to diffusion is likely blurred in the presence of leaf pubescence extending beyond the boundary layer. These conflicting trends lead to a broad range over which empirical measurements of V_d and particle size are relatively independent (Figure 4-3), further demonstrating the importance of the quasilaminar boundary layer (Lamaud et al., 1994; Shinn, 1978).

The aerodynamic term (r_a) decreases with increasing wind speed, turbulence, and friction velocity and increases with measurement height and atmospheric stability. It describes the



Figure 4-3. The relationship between particle diameter and deposition velocity for particles. Values measured in wind tunnels by Little and Wiffen (1977) over short grass with wind speed of 2.5 m s^{-1} closely approximate the theoretical distribution determined by Peters and Eiden (1992) for a tall spruce forest. These distributions reflect the interaction of Brownian diffusivity (descending dashed line), which decreases with particle size and sedimentation velocity (ascending dotted line from Stokes Law), which increases with particle size. Intermediate-sized particles (≈0.1 to 1.0 μm) are influenced strongly by both, and deposition is independent of size.

capacity of turbulent eddies to transport material, momentum, and heat between the measurement height and the roughness height of the surface. Coarse particles may not be carried efficiently by the high frequency eddies near the surface and may fall more rapidly than they diffuse by either Brownian or turbulent process. Thus the relevance of r_a breaks down as V_s increases. Indeed because V_s (Eq. 4-2) is independent of a concentration gradient, the electrical analogy is a theoretically flawed approximate approach (Venkatram and Pleim, 1999).

Deposition Velocity

Because the final stage of deposition for particles involves either impaction following deceleration through a quasi-laminar boundary layer or diffusion through this boundary layer, its effective depth is a critical determinant of V_d (Wiman et al., 1985; Peters and Eiden, 1992). The term corresponding to the boundary layer resistance for gases (r_b; equation 4-3) incorporates the absence of form drag for gases. This parameter decreases with increasing turbulence and particle diffusivity but is poorly characterized for gases, depending critically on canopy morphology, vertical wind profiles, and gust penetration, and is of extremely limited usefulness for particles.

Once delivered by turbulent transport or sedimentation to the vicinity of vegetative surface elements, a variety of particle size-dependent mechanisms come into play, some differing substantially from those governing gaseous deposition. The concepts of r_b (the still air or boundary layer resistance) and r_c (the canopy or surface resistance) are not generally applicable to deposition of polydisperse particles. Because of the roles of momentum and bounce-off and complication by reentrainment back into the airstream following deposition of a particle to the surface, the factors determining the effective r_b and r_c for particle deposition are not as independent as for gases. They are replaced in some resistance formulations (e.g., Hicks et al., 1987) by the term, r_{cp}, that combines near-surface and surface effects and by a mathematically derived composite term, $R_aR_{cp}V_s$, that combines atmospheric, surface, and sedimentation effects (Figure 4-1). This latter term was insignificant for the submicron sulfate component considered originally in its derivation (Hicks et al., 1987) but scales with the square of particle diameter so that its general applicability to polydisperse particles is unclear. In general, transport between the turbulent air column and the leaf surface through the laminar boundary layer remains difficult to describe (Lindberg and McLaughlin, 1986).

Current estimates of regional particulate dry deposition (e.g., Edgerton et al., 1992; Brook et al., 1999) infer fluxes from the product of (variable and uncertain) measured or modeled particulate concentrations and (even more variable and uncertain) measured or modeled estimates of dry deposition velocity parameterized for a variety of specific surfaces (e.g., Brook et al., 1999). However, even for specific sites and well defined particles, uncertainties in F are largest in the values of V_d, which are typically characterized by the large ranges and variances described in Section 4.2.2.2 and other sources (e.g., Bytnerowicz et al., 1987a,b, Hanson and Lindberg, 1991, for nitrogen-containing particles; McMahon and Denison, 1979, Hicks et al., 1987, for

general treatment). The nature of the vegetative cover to which particulate deposition occurs has a moderate to substantial effect on the components of V_d. The surface resistance (Hicks et al., 1987) is a significant and highly site-specific component of total resistance that is difficult to predict along with site, seasonal, and diurnal effects on the atmospheric components of total resistance.

Early models of dry particulate deposition to vegetation (e.g., U.S. Environmental Protection Agency, 1982; Chamberlain, 1975; Davidson and Friedlander, 1978; Garland, 1978; Little and Wiffen, 1977; McMahon and Denison, 1979; Sehmel, 1980; Sehmel and Hodgson, 1976; and Slinn, 1977, 1978) used this paradigm (e.g., Eq. 4-3) to deal with transport to the near-surface regime explicitly including conventional micrometeorological and particle size considerations. Alternative modeling treatments have attempted to parameterize the geometry of vegetative receptor surfaces and within-canopy micrometeorology (Wiman and Ågren, 1985; Peters and Eiden, 1992). Chemical reactivity, particle shape and density, rates of physiological sequestration, and reentrainment by gusts of wind remain to be addressed. Modeling the deposition of particles to vegetation is at a relatively early stage of development, and it is not currently possible to identify a best or most generally applicable modeling approach. These approaches have been further elaborated with canopy-specific choices among the available models and with specific incorporation of capture efficiencies by Brook et al. (1999).

Methods of Measuring Dry Deposition

Methods of measuring dry deposition of particles are more restricted than for gaseous species and fall into two major categories (Davidson and Wu, 1990). Surface extraction or washing methods characterize the accumulation of particles on natural receptor surfaces of interest or on experimental surrogate surfaces. These techniques rely on methods designed specifically to remove only surface-deposited material (Lindberg and Lovett, 1985). Total surface rinsate may be equated to accumulated deposition or to the difference in concentrations in rinsate between exposed and control (sheltered) surfaces and may be used to refine estimates of deposition (John et al., 1985; Dasch, 1987). In either case, foliar extraction techniques may underestimate deposition to leaves because of uptake and translocation processes that remove pollutants from the leaf surface (Taylor et al., 1988; Garten and Hanson, 1990). Foliar extraction methods also cannot distinguish sources of chemicals (e.g., N) deposited as gases from those

deposited as particles (e.g., nitric acid [HNO_3] or nitrate [NO_3^-] from nitrogen dioxide [NO_2], or ammonia [NH_3] from ammonium [NH_4^+]; Bytnerowicz et al., 1987a,b; Dasch, 1987; Lindberg and Lovett, 1985; Van Aalst, 1982). Despite these limitations, these methods are often used in the development of in-canopy deposition models (McCartney and Aylor, 1987).

Deposition of pollutants by wet deposition is relatively straightforward to determine through analysis of precipitation samples. Dry deposition of pollutants, on the other hand, is more difficult to measure. The National Dry Deposition Network (NDDN) was established in 1986 to document the magnitude, spatial variability, and trends in dry deposition across the United States. Currently, the network operates as a component of the Clean Air Status and Trends Network (CASTNet) (Clarke et al., 1997).

Dry deposition is not measured directly, but is determined by a inferential approach (i.e., fluxes are calculated as the product of measured ambient concentration and a modeled deposition velocity). This method is appealing and widely used because atmospheric concentrations are relatively easy to measure when compared to dry deposition fluxes, and models have been developed to calculate deposition velocities (Lovett, 1994). Ambient pollutant concentrations, meteorological conditions, and land use data required for the inferential model are routinely collected at CASTnet dry deposition sites. Chemical species include ozone, sulfate, nitrate, ammonium, sulfur dioxide, and nitric acid. The temporal resolution for the ambient concentration measurements and dry deposition flux calculations is hourly for ozone and weekly for the other chemical substances (Clarke et al., 1997). Isotopic labeling of dry deposited PM (e.g., sulfate with ^{35}S) prior to experimental surface exposures and extractions (Garten et al., 1988) can provide more precise differentiation between the deposition rates of related chemical species (e.g., sulfate [SO_4^{-2}] from sulfur dioxide [SO_2]).

At the whole-canopy level, natural surface washing by rainfall may be used to estimate dry deposition of PM and gases during the preceeding dry period (Cape et al., 1992; Davidson and Wu, 1990; Draaijers and Erisman, 1993; Erisman, 1993; Fahey et al., 1988; Lindberg and Lovett, 1992; Lovett and Lindberg, 1993; Reiners and Olson, 1984; Sievering, 1987). Collection and analysis of stem flow and throughfall provides useful estimates of particulate deposition when compared to directly sampled precipitation. The method is most precise for strictly PM deposition when gaseous deposition is a small component of the total dry deposition and when leaching or uptake of compounds of interest out of or into the foliage (i.e., N, S, base cations) is

not a significant fraction of the total depositional flux (Davidson and Wu, 1990; Draaijers and Erisman, 1993; Lindberg and Lovett, 1992; Lovett and Lindberg, 1993). Throughfall sampling of sulfate deposition (Garten et al., 1988; Lindberg and Garten, 1988; Lindberg et al. 1990) often suggests substantial foliar exchange. Other throughfall studies (e.g., Erisman, 1993; Fahey et al., 1988) may lack sufficient specificity for dry particle deposition.

Deposition to surrogate surfaces deployed in extensive plant canopies provides a measure of particle deposition to the surrounding foliage or soil surfaces. For example, a uniform population of submicron particles (gold colloid, 0.8 μm) were deposited similarly to leaves of *Phaseolus vulgaris* and to upward facing inert surfaces (Klepper and Craig, 1975). However, comparison of dry deposition of particles to foliage and to inert surrogate surfaces (polycarbonate Petri dishes; Lindberg and Lovett, 1985) in a deciduous forest demonstrated greater accumulation on the inert surfaces; with both surfaces having accumulated particles of a similar range of sizes. These persistent differences in deposition/accumulation remain to be fully characterized and hinder efforts to use these surrogate techniques to provide quantitative estimates of deposition.

Micrometeorological methods employ an eddy covariance, eddy accumulation, or flux gradient protocol in contrast to washing or extracting of receptor surfaces to quantify dry deposition. These techniques require measurements of PM concentrations and of atmospheric transport processes. They are currently well developed for ideal conditions of flat, homogeneous, and extensive landscapes and for chemical species for which accurate and rapid sensors are available. Recent progress has expanded the range of such species and extended these techniques to more complex terrain (McMillen, 1988; Hicks et al., 1984; Wesely and Hicks, 1977).

The eddy covariance technique measures vertical fluxes of gases and fine particles directly from calculations of the mean covariance between the vertical component of wind velocity and pollutant concentration (Wesely et al., 1982). It is particularly limited by a requirement for sensors capable of acquiring concentration data at 5-20 Hz. For the flux-gradient or profile techniques, vertical fluxes are calculated from a concentration difference and an eddy exchange coefficient determined at discrete heights (Erisman et al., 1988; Huebert et al., 1988). Businger (1986), Baldocchi (1988), and Wesely and Hicks (1977) evaluate the benefits and pitfalls of these micrometeorological flux measurements for gases. Most measurements of eddy transport of PM have used chemical sensors (rather than mass or particle counting) to focus on specific PM components. These techniques have not been well developed for generalized particles and may

be less suitable for coarse PM_{10} transported efficiently in high frequency eddies (Gallagher et al., 1988) for the same reasons that limit mathematical description of particle deposition above.

Factors Affecting Dry Deposition

Ambient Concentration. The ambient concentration of particles (C_z; Eq. 4-1), the parameter for which there is the most data (for example, see Chapter 3, this document), is at best an indicator of exposure. However, it is the amount of PM actually entering the immediate plant environment that determines the biological effect. The linkage between ambient concentration and delivery to vegetation is the deposition velocity (V_d), as noted above (Eq. 4-1). C_z is determined by regional and local emission sources, regional circulation, and weather. It may be locally sensitive to removal from the atmosphere by deposition, but the effect is generally small. Current ambient $PM_{2.5}$ concentrations common to non-urban ecosystems are generally well below 50 μg/m. Mean annual NO_3^- concentrations across the eastern United States ranged from 0.2 to 3.9 μg/m. Summer-time mean sulfur concentrations in western states did not exceed 1 μg/m (3 μg/m SO_4^{-2}), but mean levels in the eastern states were commonly greater than 2 μg/m, especially in the southern Appalachians (Edgerton et al., 1992; Eldred and Cahill, 1994).

Deposition increases linearly with concentration of many materials over a broad range. This allows atmospheric cleansing to take place and accounts for the greater surface impact of pollutants during pollution episodes. A serious limitation of the V_d formulation used to infer deposition of specific chemical species that exist in a range of particle sizes is an appropriate specification of their concentration. Most sulfur emissions are readily oxidized to sulfite, bisulfite, and sulfate. In the presence of atmospheric ammonia, particulate ammonium sulfate is formed. However, this material is hygroscopic will increase in mass and diameter in the presence of high humidity and alter its deposition behavior. Similarly, coalescence of small particles into larger aggregates and adsorption of gaseous pollutants onto existing coarse particles complicate the association of particle size with concentration of individual chemical species.

Distance and the resulting residence time in the atmosphere control the relative concentrations of surface reactive materials (NO, SO_2) of secondary particles that take some time to form in the atmosphere and of coarse particles that exhibit high rates of deposition by sedimentation near the source. These interacting processes affect the time required for formation of secondary particles by gas-to-particle conversion reactions and result in a greater ratio of dry

to wet deposition near emission sources where gaseous sulfur dioxide (gSO_2) deposition predominates than at greater distances where rainout of particulate SO_4^{-2} (pSO_4^{-2}) may dominate (Barrie et al., 1984) and dry deposition of pSO_4^{-2} may be greater than of gSO_2 . The effect of gas-to-particle conversion on dry deposition of a specific chemical species can be substantial because V_d for SO_2 is approximately 0.33 ± 0.17 cm s^{-1}; whereas it is approximately 0.16 ± 0.08 cm s^{-1} for SO_4^{-2}. These phase conversions impact both C_z and the effective V_d which together control dry depositional fluxes (Eq. 4-1). The neutralization of acidic gaseous and particulate species by alkaline coarse particles has been described in arid regions, but it may be more prevalent in urban New York, where coarse particles are observed to be neutral with alkaline cations approximately balancing gaseous acidic species (Lovett et al., 2000). The deposition of the acidic materials in the urban environment is likely enhanced by incorporation into these previously formed coarse particles.

Similarly, the ratio of coarse to fine particle concentrations determines the effective V_d for chemically speciated particles (Figure 4-2). This reflects the size-dependent deposition processes that govern delivery of PM to receptor surfaces (Fig. 4-3). For example, SO_4^{-2} was found predominantly on fine submicron particles; whereas potassium ion (K^+), calcium (Ca^{+2}), and nitrate (NO_3^-) were associated most often with coarse particles larger than 2 μm (Lindberg and Lovett, 1985). However, concentrations of particulate S and K^+ within a coniferous canopy were strongly correlated (Wiman and Lannefors, 1985), suggesting a primary source of coarse-mode sulfur particles. Furthermore, NO_3^- and SO_4^{-2}-containing fine particles readily coalesce with coarse particles derived from sea spray or primary geologic material (Wu and Okada, 1994; Milford and Davidson, 1987). Gaseous N and S species may undergo gas to particle conversion directly onto such preexisting coarse particles. As a result, marine and continental particle size spectra for both N and S differ substantially, with a peak in the coarse mode generally apparent near marine sources (Milford and Davidson, 1987). The issue for NO_3^- is further confounded by uncertain discrimination between gaseous and particulate species in current sampling methods. The substantial effect of particle size on V_d (Figure 4-3) implies a need for size resolution as well as chemically speciated ambient particulate concentrations even within the PM_{10} fraction.

Particle Effects on V_d. Particle size is a key determinant of V_d as noted above; but, unfortunately, the size spectra may be quite complex. The particles in the study of Lindberg and

Lovett (1985) at Walker Branch Watershed had median diameters ranging from 3 to 5 μm; but approximately 25% of the particles had diameters less than 1 μm (0.2 to 0.3 μm), and 5 to 20% of the particles were much larger aggregates. The aggregated particles are significant in that chemically they reflect their fine particle origins, but physically they behave like large particles and deposit by sedimentation. Direct observations with SEM demonstrate that particle morphology can be highly variable. Many submicron particles can be observed on trichomes (leaf hairs), although most particles are in the 5 to 50 μm diameter range. Large aggregated particles in excess of 100 μm also are seen, with carbonaceous aggregate particles being especially common (Smith, 1990a). Trichomes are especially efficient particle receptors; however, trichomes are reduced in size by "weathering" and occasionally are completely broken off during the growing season.

In the size range around 0.1 to 1.0 μm, where V_d is relatively independent of particle diameter (Fig. 4-3), deposition is controlled by macroscopic roughness properties of the surface and by the stability and turbulence of the atmospheric surface layer. The resistance catena (Figure 4-1) is less useful in this size range and, in some treatments, has been abandoned entirely (e.g., Erisman et al., 1994; Eq. 4-4). Impaction and interception dominate over diffusion, and the V_d is considerably (up to two orders of magnitude; Figure 4-3) lower than for particles either smaller or larger (Shinn, 1978). The deposition velocity may be parameterized in this size range as a function of friction velocity,

$$V_d = (a/b)u^*, \qquad (4\text{-}4)$$

where a depends on atmospheric stability and b depends on surface roughness (Wesely et al., 1985; Erisman et al., 1994). Similar formulations have been presented in terms of turbulence (standard deviation of wind direction) and wind speed (e.g., Wesely et al., 1983), both determinants of u*.

Deposition of particles between 1 and 10 μm diameter, including the coarse mode of PM_{10}, is strongly dependent on particle size (Shinn, 1978). Larger particles within this size range are collected more efficiently at typical wind speeds than are smaller particles (Clough, 1975), suggesting the importance of impaction. Impaction is related to wind speed, the square of particle diameter, and the inverse of receptor diameter as a depositing particle fails to follow the

streamlines of the air in which it is suspended around the receptor. When particle trajectory favors a collision, increasing wind speed or ratio of particle size to receptor cross section makes collision nearly certain; and, as these parameters become very small, the probability of collision becomes negligible. However, the shape parameter for the more common range of situations between these extremes remains poorly characterized (Peters and Eiden, 1992; Wiman and Ågren, 1985).

As particle size increases above 1 μm, deposition is governed increasingly by sedimentation (Figure 4-3) with a correspondingly declining influence of turbulence and impaction. Particles between approximately 10 and 24 μm (Gallagher et al., 1988) are both small enough to be transported efficiently by turbulent eddies to the surface and large enough to impact with sufficient momentum to overcome boundary layer effects. These particles deposit highly efficiently and relatively independently of particle size.

Deposition of the largest suspended particles (e.g., >50 μm) is governed, through sedimentation and the corresponding terminal settling velocity (V_s), almost entirely by size. These particles are not transported efficiently by small-scale eddies near the surface.

Theoretically based models for predicting particle deposition velocities have been published by Bache (1979a,b), Davidson et al. (1982), Noll and Fang (1989), Slinn (1982), and Wiman (1985). These models deal primarily with low canopies or individual elements of canopy surfaces. Wiman and Ågren (1985) have developed an aerosol deposition model that specifically treats the problem of particle deposition to forests where turbulence plays a particularly important role, especially at roughness transitions such as forest edges. They found that deposition of supermicron particles is controlled by complex interactions among particle size and concentration, forest structure, and aerodynamics; whereas deposition of fine particles (submicron) is controlled by particle concentration and forest structure.

At the present time, empirical measurements of V_d for fine particles under wind tunnel and field conditions are often several-fold greater than predicted by available theory (Unsworth and Wilshaw, 1989). A large number of transport phenomena, including streamlining of foliar obstacles, turbulence structure near surfaces, and various phoretic transport mechanisms remain poorly parameterized in current models. The discrepancy between measured and predicted values of V_d may reflect such model limitations or experimental limitations in specification of the effective size and number of receptor obstacles, as suggested by Slinn (1982).

Available reviews (Davidson and Wu, 1990; McMahon and Denison, 1979; Nicholson, 1988; Sehmel, 1980; Slinn, 1982; U.S. Environmental Protection Agency, 1982, 1996a) suggest the following generalizations: (1) particles larger than 10 μm exhibit a variable V_d between $\approx$0.5 and 1.1 cm s^{-1} depending on friction velocities; whereas a minimum particle V_d of 0.03 cm s^{-1} exists for particles in the size range 0.1 to 1.0 μm; (2) the V_d of particles is approximately a linear function of friction velocity; and (3) deposition of particles from the atmosphere to a forest canopy is from 2 to 16 times greater than deposition in adjacent open terrain (i.e., grasslands or other vegetation of low stature).

Leaf Surface Effects on V_d. The term r_c (Equation 4-3) reflects the chemical, physical, or physiological characteristics of the surface that govern its ability to capture, denature, or otherwise remove particulate material from the atmospheric surface layer. For gases, relevant surface properties involve the physiological state of the vegetation, including stomatal opening and mesophyll antioxidant activity, and the chemical reactivity of the exposed surface with the specific gas. For particles, relevant surface properties involve stickiness, microscale roughness, and cross-sectional area which determine the probability of impaction and bounce (e.g., Shinn, 1978). The chemical composition of the PM is not usually a primary determinant of deposition velocity. At the microscopic scale where Van der Waals forces may determine particle bounce and reentrainment, the chemical properties of both surface and particle may be significant but remain poorly understood.

Stickiness may itself depend on previous deposition of deliquescent particles that prolong leaf wetness, on the wettability of foliar surfaces, and on the presence of sticky residues such as honeydew deposited by aphids. These factors increase deposition by decreasing bounce-off, wind reentrainment, and, to some extent, wash-off by precipitation.

The distribution of particles on and the efficiency of deposition to vegetation also varies based on leaf shape and plant part. Particles are more prevalent on the adaxial (upper [facing the twig]) surface than on the abaxial (lower [away from the twig]) surface. Peripheral leaf areas tend to be the cleanest with most particles accumulating in the midvein, central portion of leaves.

The rough area surrounding the stomatal pores was not found to be a preferential site for particle deposition or retention (Smith and Staskawicz, 1977). Most particles were located near veins with smaller particles localized on the trichomes. The greatest particulate loading on

dicotyledonous leaves is frequently on the adaxial surface at the base of the blade, just above the petiole junction. It is probable that precipitation washing plays an important role in this distribution pattern. Lead particles accumulated to a larger extent on older than younger needles and twigs of white pine, indicating that wind and rain were insufficient to fully wash the foliage. Fungal mycelia (derived from windborne spores) were frequently observed in intimate contact with other particles on other leaves (Smith and Staskawicz, 1977), which may reflect shelter by the particles minimizing reentrainment of the spores, mycelia development near sources of soluble nutrients provided by the particles, or codeposition. This pattern is significant and could yield further insight into deposition mechanisms.

Leaves with complex shapes collect more particles than those with regular shapes. Conifer needles are more effective than broad leaves in accumulating particles. The edge to area ratio (Woodcock, 1953) is also a key determinant of salt deposition to individual artificial leaves. A strong negative correlation was observed under wind tunnel conditions between the area of individual leaves and deposition of coarse particles (Little, 1977). Small twigs and branches were more effective particle collectors than were large branches and trunks of trees (Smith, 1984). Lead particles accumulated 20-fold more on woody stems than on leaves of white pine (*Pinus strobus*), even though leaves displayed a 10-fold greater total area (Heichel and Hankin, 1976). Deposition is heaviest at tips of individual leaves.

Rough, pubescent broadleaf discs collected coarse (5.0-μm) particles up to sevenfold more efficiently than glabrous leaf discs (Little, 1977). Laminae, petioles, and stems, all differed in collection efficiency. Pubescent leaves of sunflower (*Helianthus annuus*) collected coarse particles nearly an order of magnitude more efficiently than the glabrous leaves of tulip poplar (*Liriodendron tulipifera*) under wind tunnel conditions (Wedding et al., 1975). Rough pubescent leaves of nettle (*Urtica dioica*) were more effective in capturing coarse PM_{10} than were the densely tomentose leaves of poplar (*Populus alba*) or smooth leaves of beech (*Fagus sylvatica*).

Conifer needles are more efficient than broad leaves in collecting particles by impaction. This reflects the small cross section of the needles relative to larger leaf laminae of broadleaves and the greater penetration of wind into conifer than broadleaf canopies (below). Conifers were more effective in removing coarse ($\approx$20 μm) particles of ragweed pollen from the atmosphere than were broadleaf trees (Steubing and Klee, 1970) and in intercepting the even coarser particles of rain (Smith, 1984). Conifers are also more effective in retaining and accumulating particles

against reentrainment by wind and removal by rain, particularly on foliar surfaces where they are likely to be most biologically active. Seedlings of white pine (*Pinus strobus*) and oak (*Quercus rubra*) each initially retained between a quarter (pine) and a third (oak) of very coarse (88 to 175 μm) ^{134}Cs-tagged quartz particles applied under field conditions (Witherspoon and Taylor, 1971). After 1 h, the pine retained over 20% of the ^{134}Cs-tagged particles; whereas the oak retained only approximately 3%. Long-term retention of the particles was concentrated at the base of the fascicles in pine and near the surface roughness caused by the vascular system on leaves of oak. The sheltered locations available in the conifer foliage contribute substantially to greater retention of particles. For similar reasons, grasses also are efficient particle collectors (Smith and Staskawicz, 1977) with long-term retention mostly in the ligule and leaf sheath.

Wind tunnel studies also demonstrated equivalent deposition properties of 3.36-μm particles of dense lead chloride and 6.77-μm particles of less dense uranine dye. These particles were shown to be aerodynamically equivalent, substantiating the use of aerodynamic diameter as a classification parameter for particle deposition.

Canopy Surface Effects on V_d. In general, surface roughness contributes to greater particulate deposition. As a result, V_d is typically greater for a forest than for a field or nonwoody wetland and greater for a field than for a water surface. The contrasting transport properties and deposition velocities of different size particles lead to predictable patterns of deposition. For coarse particles, the upwind leading edges of forests, hedge rows, and individual plants, as well as of individual leaves, are primary sites of deposition. Impaction at high wind speed and the sedimentation that follows the reduction in wind speed and carrying capacity of the air in these areas lead to preferential deposition of larger particles.

Air movement is slowed in proximity to vegetated surfaces. Resulting log profiles of wind and pollutant concentrations in the near-surface turbulent boundary layer above canopies reflect surface characteristics of roughness length, friction velocity, and displacement height. Plasticity, streamlining, and oscillations of foliar elements also alter the aerodynamic roughness and the level of within-canopy turbulence. Canopies of uneven age or with a diversity of species are typically aerodynamically rougher and receive larger inputs of pollutants than do smooth, low, or monoculture vegetation (Garner et al., 1989; Wiman and Ågren, 1985). Canopies on slopes facing the prevailing winds and individual plants on the windward edges of discontinuities in

vegetative cover over which roughness increases receive larger inputs of pollutants than more sheltered, interior canopy regions. For example, some 80% of coarse particulate sea salt was deposited on the upwind edge of a hedgerow (Edwards and Claxton, 1964), and the concentration of ragweed (*Ambrosia spp.*) pollen was reduced by 80% within 100 m of the leading edge of a forest (Neuberger et al., 1967).

Beier et al. (1992) and Beier (1991) discussed two methods for estimating the dry deposition of base cations to forest edges: (1) a difference method between measured precipitation and throughfall concentrations of base cations, and (2) a calculation method based on known ratios of Na^+ deposition in wet and dry forms (Ulrich, 1983). A combination of these two approaches produced the best estimates of SO_4^{-2}, Ca^{+2}, Mg^{+2}, and K^+ particle deposition. Using these methods, particulate SO_4^{-2} (Beier, 1991) and particulate Ca^{+2}, Mg^{+2}, and K^+ (Beier et al., 1992) were found to decrease by an order of magnitude from the forest edge to the forest interior. A number of authors also have shown that particle deposition is elevated at forest edges when compared to a uniform forest canopy (Draaijers et al., 1988; Grennfelt, 1987; Lindberg and Owens, 1993), and Draaijers et al. (1992) reported that differences are likely to exist between forest types because of variable canopy structure. Draaijers et al. (1988) further emphasized that enhanced particle deposition at or near forest edges is strongly dependent on the velocity and wind direction during observations.

The factors leading to horizontal gradients are confounded by time- and distance-related sedimentation, geologic dust (mostly around 7 μm aerodynamic diameter) being collected on stems of wild oats (*Avena spp.*; Davidson and Friedlander, 1978) and on eastern white pine (*Pinus strobus*; Heichel and Hankin, 1972; Smith 1973) downwind of roadways. Rapid sedimentation of coarse crustal particles suggests that potential direct effects may be restricted to roadway margins, forest edges, and, because of the density of unpaved roads in agricultural areas, crop plants.

Simulated deposition to an ecologically complex, mixed canopy was considerably higher than to a pure spruce stand in which most of the leaf area was concentrated in regions of low wind speed. Limitations to the application of these models to predict deposition over large regions include a limited understanding both of the nature of microscopic particle-surface interactions and of the effects of complex terrain and species composition on macroscopic transport processes.

Macroscopic turbulent transport processes, related to r_a, at successive layers through the canopy can be separated from microscopic processes, related to r_b and r_c (or r_{cp}), at each deposition surface (e.g., Peters and Eiden, 1992; Wiman and Lannefors, 1985). The macroscopic approach deals with deposition as the product of a turbulent diffusion coefficient and a concentration (C_z) at each canopy layer, both of which vary with particle size and with height (Z) in the canopy. The microscale parameters involve those factors that determine absorption of a particle at each surface as captured imperfectly by r_c. Shelter effects caused by the crowding of foliar elements within the canopy can be ignored if the wind speed within each canopy layer is specified. This approach requires knowledge of the vertical distribution of particle concentration and foliage density in the canopy airspace along with profiles of wind speed or turbulence.

Once introduced into a forest canopy, elements associated with course particles tend to decrease markedly with canopy depth; whereas elements associated with fine particles do not (Lovett and Lindberg, 1992).

Trace elements and alkaline earth elements are enriched below the canopies of both southern (Lindberg et al., 1986) and northern (Eaton et al., 1973) hardwood forests. Vertical gradients in concentration of coarse particles and of elements associated with coarse particles were observed in a mixed conifer/birch forest canopy (Wiman and Lannefors, 1985; Wiman et al., 1985) and in a mixed oak forest (e.g., Ca^{+2}, Figure 4-4A; Lovett and Lindberg, 1992). The highly reactive gas HNO_3 also exhibited a vertical gradient, but with a steep decline at the top of the canopy (Figure 4-4B). Lovett and Lindberg (1992) studied concentration profiles of various gases and particles within an closed canopy forest and concluded that coarse particle concentrations associated with elements like Ca^{+2} would decrease markedly with depth in the canopy, but they found only minor reductions with depth in the concentrations of fine aerosols containing SO_4^{-2}, NH_4^+, and H^+. These data suggest that all foliar surfaces within a forest canopy are not exposed equally to particle deposition: upper canopy foliage would receive maximum exposure to coarse and fine particles, but foliage within the canopy would receive primarily fine aerosol exposures. Fine-mode particles (e.g., sulfate [SO_4^{-2}], Figure 4-4C) and unreactive gases typically do not exhibit such vertical profiles, suggesting that uptake is smaller in magnitude and more evenly distributed throughout the canopy. In multilayer canopies, simultaneous reentrainment and deposition may effectively uncouple deposition from local concentration.



Figure 4-4. Vertical stratification of diverse, chemically speciated particles in a mixed oak forest. Nitric acid vapor, as a highly reactive, nonparticulate species, is shown for comparison. The horizontal dotted line indicates canopy height, and Δ indicates the percent depletion from above to below the canopy. Data from Table 2-4 of Lovett and Lindberg (1992).

Polydisperse size distributions of many chemical species effectively prevent the use of a single estimate of V_d for any element if highly accurate results are required.

Although gradients (Figures 4-4 and 4-5) may be related to local V_d within the canopy (Bennett and Hill, 1975), the absence of a gradient may reflect either low rates of deposition or very high rates relative to turbulent replenishment from above the canopy (Tanner, 1981). Below- or within-canopy emissions may confound interpretation of vertical gradients. Linear gradients of the gaseous pollutants hydrogen fluoride (HF) and ozone (O_3) reflected large uptake rates; whereas small gradients in NO suggested little uptake by foliage (Bennett and Hill, 1973,



Figure 4-5. The relationship between particle size and concentration below a spruce canopy with wind velocity at a height of 16.8 m equaling 5 m s^{-1}. Modified from Peters and Eiden (1992).

1975). However, soil efflux of NO could have complicated the latter interpretation. The lack of a vertical gradient and a peak near the top of the active canopy in particulate K^+ (Figure 4-4D) was interpreted as evidence for a biogenic source within the deciduous forest canopy with moderate rates of deposition (Lindberg et al., 1986; Lovett and Lindberg, 1992).

The size dependence of this vertical stratification of particulate concentration (see Figure 4-5) is reflected in current simulation models (Wiman et al., 1985; Peters and Eiden, 1992). The model of Wiman and Ågren (1985) predicts a uniform vertical distribution of fine-mode particles and a pronounced vertical gradient of coarse-mode particles which is in agreement with observations (Lovett and Lindberg, 1992).

Simulation of the horizontal deposition patterns at the windward edge of a spruce forest downwind of an open field with the canopy between 1 and 25 m above the ground indicated that deposition was maximal at the forest edge where wind speed and impaction were greatest. Simulation of the vertical deposition pattern was more complex. Deposition was not greatest at

the top of the canopy where wind speed was highest, but at z = 0.75 h where the balance between leaf area (obstacles for impaction) and wind speed (momentum for impaction) was optimal, although neither parameter alone was maximal. Simulated deposition in this spruce forest increased considerably with increasing LAI at the forest edge, where wind speed was insensitive to LAI but the number of obstacles increased. Inside the forest, where both wind speed and impaction increasingly were attenuated by increasing LAI, deposition increased only marginally in spite of the increase in obstacle frequency.

To scale surface-specific measurements of particle deposition to forest or crop canopies, conversions of the following type have been suggested:

$$V_{d.canopy} = V_{d.surface} * \text{scaling factor}, \qquad (4\text{-}5)$$

with empirical scaling factors proposed by Lindberg et al. (1988).

To appropriately scale surface-specific measurements of particle deposition to landscapes, one must consider the complexity of grassland, crop, and forest canopies in order to avoid serious over- or under-estimates of particle deposition. Individual species exposed to similar ambient concentrations may receive a range of particulate loading that is more closely related to foliar damage than the ambient concentration (Vora and Bhatnagar, 1987).

Both uptake and release of specific constituents of PM may co-occur within a single canopy (e.g., K^+; Lovett and Lindberg, 1992). The leaf cuticular surface is a region of dynamic exchange processes through leaching and uptake. Exchange occurs with epiphytic microorganisms and bark and through solubilization and erosion of previously deposited PM. Vegetation emits a variety of particulate and particulate precursor materials. Terpenes and isoprenoids predominate and, on oxidation, become condensation nuclei for heterogeneous particle formation. Salts and exudates on leaves and other plant parts continually are abraded and suspended as particles, as are plant constituents from living and dead foliage (Rogge et al., 1993a). Soil minerals, including radioactive strontium, nutrient cations and anions, and trace metals are transferred to the active upper foliage and then to the atmosphere in this way. Although not representing a net addition to an ecosystem, particle release from vegetation is a mechanism for redistributing chemical pollutants derived from the soil or prior deposition within a canopy, potentially enhancing direct effects and confounding estimates of V_d.

Range of Deposition Velocity

As noted in the previous criteria document (U.S. Environmental Protection Agency, 1982) and in McMahon and Denison (1979), estimates of V_d for PM_{10} particles to vegetation are variable and suggest a minimum between 0.1 and 1.0 μm as predicted from first principles (Monteith and Unsworth, 1990; Sehmel, 1980). Determinations in wind tunnels with passive collectors and micrometeorological methods tend to converge in this range. The range of V_d for sulfate from passive collectors was found to be from 0.147 to 0.356 cm s^{-1}; and, from eddy covariance techniques, a mean V_d of 0.27 cm s^{-1} was observed (Dolske and Gatz, 1984). Micrometeorological techniques over grass (Wesely et al., 1985); indirect, inert collector techniques within an oak forest (Lindberg and Lovett, 1985); and many other empirical determinations (e.g. McMahon and Denison, 1979; Table 4-3) generally support this range. Over aerodynamically smooth snow (Duan et al., 1988; Table 4-4), measurements of V_d were an order of magnitude smaller. Very coarse particles, often non-size-specified primary geologic material, frequently exhibit V_d greater than 1.0 cm s^{-1} (e.g., Clough, 1975). The increase in V_d with decreasing size below 0.1 μm is probably hidden in most empirical determinations of V_d, because the total mass in this fraction is very small despite the large number of individual particles. Table 4-5 shows published estimates of V_d with variability estimates for fine particles of specified aerodynamic diameters dominated by a range of chemical species.

Ibrahim et al. (1983) evaluated the deposition of ammonium sulfate particles to a range of surfaces and found that particles having a mean diameter of 0.7 μm had deposition velocities ranging from 0.039 to 0.096 cm s^{-1}. Larger particles (having mean diameters of 7 μm) had greater deposition velocities (between 0.096 and 0.16 cm s^{-1}). The authors further concluded that the hygroscopic nature of the sulphate particle could increase its size and enhance deposition near sources of water, e.g., snow. Using eddy correlation approaches, Hicks et al. (1989) found a mean daily V_d for sulfur-containing PM to be 0.6 cm s^{-1}. However, they suggested that the V_d value could be as high as 1 cm s^{-1} during the day and near zero at night.

Lindberg et al. 1990 found a wide discrepancy between deposition velocities for NO_3^- between study sites in Oak Ridge, TN ($\approx$2 cm s^{-1}) and Göttingen, Germany ($\approx$0.4 cm s^{-1}). They suggest that the increased V_d at Oak Ridge could be explained by the primary occurrence of NO_3^- in coarse particles that exhibit greater V_d than fine particles (Davidson et al., 1982). Large values

TABLE 4-3. REPORTED MEAN DEPOSITION VELOCITIES (V_d) FOR SULFATE, CHLORINE, NITRATE, AND AMMONIUM AND ION-CONTAINING PARTICLES

Chemical Species/ Surface	V_d (cm s^{-1})[a]	Method	Reference
SO_4^{-2}			
Inert plates	0.13	Extraction	Lindberg and Lovett (1985)
Inert plates	≈0.14	Extraction	Lindberg et al. (1990)
Inert plates	0.14	Extraction	Davidson and Wu (1990)[a]
Inert bucket	0.51	Extraction	Davidson and Wu (1990)[a]
Foliage	0.29	Extraction	Davidson and Wu (1990)[a]
Chaparral	0.15	Extraction	Bytnerowicz et al. (1987a)
Grass canopy	0.10	Gradient	Allen et al. (1991)
Grass canopy	0.07	Gradient	Nicholson and Davies (1987)
Pine foliage	0.07	Extraction	Wiman (1981)
Plant canopies	0.50	Gradient	Davidson and Wu (1990)[a]
Grass canopy	0.22	Eddy covariance	Weseley et al. (1985)
Cl^-			
Inert bucket	3.1	Extraction	Dasch and Cadle (1985)
Inert bucket	5	Extraction	Dasch and Cadle (1986)
Beech canopy	1	Throughfall	Höfken et al. (1983)
Spruce canopy	1.9	Throughfall	Höfken et al. (1983)
NO_3^-			
Inert plates	0.4-2	Extraction	Lindberg et al. (1990)
Ceanothus	0.4	Extraction	Bytnerowicz et al. (1987a)
Glycine max	0.24	Extraction	Dolske (1988)
Ligustrum	0.1 - 0.5	Extraction	John et al. (1985)
Quercus	0.7 - 1.1	Extraction	Dasch (1987)
Quercus			
summer	0.55	Throughfall	Lovett and Lindberg (1984)
winter	0.7	Throughfall	Lovett and Lindberg (1984)
Quercus			
summer	0.3	—	Lovett and Lindberg (1986)
winter	0.1	—	Lovett and Lindberg (1986)
Pinus	0.5 - 1.3	Extraction	Dasch (1987)
Pasture	0.7 - 0.8	Gradient	Huebert et al. (1988)
Ulmus	1.1	Extraction	Dasch (1987)
NH_4^+			
Calluna/Molina	0.18	Gradient	Duyzer et al. (1987)
Ceanothus	0.4	Extraction	Bytnerowicz et al. (1987a)
Kalmia	0.03- 0.14	Extraction	Tjepkema et al. (1981)
Pinus	0.01 - 0.06	Extraction	Dasch (1987)

[a]These data represent the mean of data by measurement technique as reported in the cited reference. The reader is referred to the referenced articles for information on the specific cations contributing to the means.

TABLE 4-4. REPRESENTATIVE EMPIRICAL MEASUREMENTS OF DEPOSITION VELOCITY (V_d) FOR PARTICULATE DEPOSITION

V_d x ± SE (cm s^{-1})	Particle Size (μm)	Method	Reference
0.034 ± 0.014 0.021 ± 0.005	0.15 - 0.30 0.5 - 1.0	Eddy covariance with optical counter, flat snow surface	Duan et al. (1988)
0.1 ± 0.03	0 - 2	Profile, fine SO_4^{2-}, short grass	Allen et al. (1991)
0.22 ± 0.06	0.1 - 2.0	Eddy covariance with flame photometer plus denuder, 40-cm grass, fine SO_4^{2-}	Wesely et al. (1985)
0.13 ± 0.02 0.75 ± 0.24 1.1 ± 0.1	(SO_4^{2-}) (K^+) (Ca^{+2})	Inert surface collectors (petri dish) in oak forest	Lindberg and Lovett (1985)
0.9 2.5 9.4	2.75 5.0 8.5	Wind tunnel to pine shoots; polystyrene beads; within- "canopy" wind speed, 2.5 m s^{-1}	Chamberlain and Little (1981)

TABLE 4-5. REPORTED MEAN DEPOSITION VELOCITIES FOR POTASSIUM, SODIUM, CALCIUM, AND MAGNESIUM BASE CATION CONTAINING PARTICLES

Chemical Species/Surface	V_d (cm s^{-1})	Method	Reference
K^+			
Inert plates	0.75	Extraction	Lindberg and Lovett (1985)
Inert bucket	0.51 - 2.4	Extraction	Dasch and Cadle (1985)
Na^+			
Inert bucket	1.7 - 2.9	Extraction	Dasch and Cadle (1985)
Inert plate	0.8 - 8.2	Extraction	
Ca^{+2}			
Inert plates	1.1	Extraction	Lindberg and Lovett (1985)
Inert plates	≈2	Extraction	Lindberg et al. (1990)
Inert bucket	1.7 - 3.2	Extraction	McDonald et al. (1982)
Mg^{+2}			
Inert bucket	1.1 - 2.7	Extraction	Dasch and Cadle (1985)

of V_d for base-cation-containing particles (>1 cm s^{-1}) suggest their occurrence in coarse particles (Lindberg and Lovett, 1985).

The several attempts to estimate V_d for SO_4^{-2}, NO_3^-, and NH_4^+ with the throughfall mass balance approach (Davidson and Wu, 1990; Gravenhorst et al., 1983; Höfken and Gravenhorst, 1982) have produced higher V_d values that are considered suspect. They have not been included in Tables 4-3 and 4-5. Overestimates of V_d for SO_4^{-2} and NO_3^- particles derived from throughfall mass balance approaches may be the result of gaseous SO_2 and HNO_3 gaseous deposition to foliar surfaces (Lindberg and Lovett, 1985). A similar contribution of NH_3 deposition may lead to erroneously high V_d values for NH_4^+ when the throughfall method is attempted in areas of high NH_3 concentrations. Dolske's (1988) reported V_d values for NO_3^- deposition to soybean ranged from 0.4 to 31 with a mean of 0.24 cm s^{-1}. However, because Dolske's leaf extraction measurements included a component of HNO_3 vapor, the V_d values may represent more than deposition caused by aerosol nitrate alone.

4.2.1.3 Occult Deposition

Gaseous pollutant species may dissolve in the suspended water droplets of fog and clouds. The stability of the atmosphere and persistence of the droplets often allow a condition of gas/liquid phase equilibrium to develop. This allows knowledge of air mass history or ambient concentrations of specific pollutants to be used to estimate fog or cloud water concentrations. Further estimates of the deposition velocity of the polluted droplets allows calculation of depositional fluxes. Unfortunately, interception of fog or cloud droplets by plant parts or other receptor surfaces remains difficult both to predict and to measure. Fog formation influences the total atmospheric burden and deposition of particulate matter (Pandis and Seinfeld, 1989) by accreting and removing particles from the air, by facilitating particle growth through aqueous oxidation reactions, and by enhancing deposition as noted. Aqueous condensation may occur onto preexisting fine particles, and such particles may coalesce or dissolve in fog or cloud droplets. Material transported in fog and cloud water and intercepted by vegetation escapes detection by measurement techniques designed to quantify either dry or wet deposition; hence it is hidden (i.e., “occult”) from the traditional measurements.

Low elevation radiation fog has different formation and deposition characteristics from high elevation cloud or coastal fog water droplets. A one dimensional deposition model has

recently been described for a radiation fog episode (Von Glasow and Bott, 1999). A substantially greater concentration of key polluting species (eg. NO_3^-, SO_4^{-2}, organics) may be observed in smaller than in larger droplets in fog (Collett et al., 1999). Acidity differences exceeding 1 pH unit were also observed in the San Joaquin Valley winter radiation fog with smaller particles being more acidic. This has implications for aqueous phase oxidation of sulfur and nitrogen compounds, in particular, while sulfur oxidation by ozone (the dominant reaction in this environment even during winter) is well known in typically acidic fog droplets. However, the alkaline larger droplets in the San Joaquin Valley could lead to greater nitrate production through aqueous ozonation reactions (Collett et al., 1999). The size class distinctions have substantial implications for deposition of particulate pollutant species in the fog droplets due to the larger V_d for impaction and occult deposition of the larger fog particles.

Acidic cloud water deposition has been associated with forest decline in industrialized areas of the world (Anderson et al., 1999). Clouds can contain high concentrations of acids and other ions. The four most prevalent ions found in cloud water samples, in decreasing order of concentration, were usually sulfate (SO_4^{2-}), hydrogen (H^+), ammonium (NH_4^+), and nitrate (NO_3^-). The concentrations of these major ions tend to co-vary within within cloud events and typically there was an inverse relationship between LWC of the cloud and ionic concentration of the cloud water. The acidity of cloud water typically is 5 to 20 times more acid than rain water. This can increase by more than 50% pollutant deposition and exposure of vegetation and soils at high-elevation sites when compared with rainfall and dry deposition.

The widespread injury to mountain forests in document since the 1970s in West Germany and other parts of Europe and more recently in the Appalachian Mountains has been attributed to exposure to the cloud water reducing cold tolerance of red spruce. Forest injury also has been attributed to increased leaching of cations and amino acids, increased deposition of nitrogen and aluminum toxicity resulting from acidic deposition and the combined effect of acidic precipitation, acid fog, oxidants, and heavy metals (Anderson et al., 1999).

The Mountain Acid Deposition Program (MADPro) was initiated in 1993 as part of the Clean Air Status and Trends Network (CASTnet). MADPro monitoring efforts focused on the design and implementation of an automated cloud water collection system in combination with continuous measurement of cloud liquid water content (LWC) and meteorological parameters relevant to the cloud deposition process.

Results from the MADPro automated cloud water collectors at three selected mountain sites (Whiteface Mt., NY; Whitetop Mt. Va; and Clingman's Dome, TN) taken hourly from nonprecipitating clouds during non-freezing seasons of the year from 1994 to 1997 were promptly analyzed for pH, conductivity, and concentration of dissolved ions give an indication of exposures at the three sites. Cloud liquid water content were measured at each site. The mean cloud water frequencies and LWC were higher at Whiteface Mountain, NY, that in the southern Appalachians. The four most prevalent ions found in cloud water samples, in order of decreasing concentrations, usually were sulfate (SO_4^{2-}), hydrogen (H^+), ammonium (NH_4^+), and nitrate (NO_3^-). The concentrations of these ions tended to co-vary within cloud events and typically there was an inverse relationship between LWC of the cloud and ionic concentration of the cloud water. Highest ionic concentrations were seen in mid-summer during the sampling season. Ionic concentrations of samples from southern sites were significantly higher than samples from Whiteface Mountain, however, further analysis indicated that this observation was due at least in part to North to South differences in the LWC of clouds (Anderson et al., 1999).

Several factors make occult deposition particularly effective for delivery of dissolved and suspended materials to vegetation. Concentrations of particulate-derived materials are often many-fold higher in cloud or fog water than in precipitation or ambient air in the same area due to orographic effects and from gas-liquid partitioning coefficients of specific chemical species. Fog and cloud water deliver PM in a hydrated and, therefore, bioavailable form to foliar surfaces. Previously dry-deposited PM may also become hydrated through delinquence or by dissolving in the film of liquid water from fog deposition. The presence of fog itself maintains conditions of high relative humidity and low radiation, thus reducing evaporation and contributing to the persistence of these hydrated particles on leaf surfaces. Deposition of fog water is very efficient (Fowler et al., 1991) with a V_d (fog 10 - 24 μm; Gallagher et al., 1988), essentially equal to the aerodynamic conductance for momentum transfer $(r_a)^{-1}$. This greatly enhances deposition by sedimentation and impaction of submicron aerosol particles that exhibit very low V_d prior to fog droplet formation (Fowler et al., 1989). The near equivalence of V_d and $(r_a)^{-1}$ simplifies calculation of fog water deposition and reflects the absence of vegetative physiological control over surface resistance. Fog particles outside this size range may exhibit V_d below $(r_a)^{-1}$. For smaller particles, this decline reflects the increasing influence of still air and boundary layer effects on impaction as particle size and momentum decline (Figure 2-1). For larger particles

momentum is sufficient to overcome these near surface limitations, but V_d may decline as turbulent eddy transport to the surface becomes inefficient with increasing inertia (Gallagher et al., 1988). The deposition to vegetation for PM in fog droplets is directly proportional to wind speed, droplet size, concentration, and fog density (liquid water content per volume air), although the latter two may be inversely related.

In some areas, typically along foggy coastlines or at high elevations, occult deposition represents a substantial fraction of total deposition to foliar surfaces (Fowler et al., 1991, Table 4-6).

TABLE 4-6. RELATIVE MAGNITUDES OF WET, DRY, AND OCCULT DEPOSITION OF NITRATES (NO_3^-) AND SULFATES (SO_4^{-2}) TO THREE FOREST SITES SUBJECT TO SIMILAR GAS- AND LIQUID-PHASE POLLUTANT CONCENTRATIONS DURING SPRING AND SUMMER[a]

	Deposition (kg ha^{-1})[b]					
	Wet		Dry		Occult	
Site[c]	NO_3^-	SO_4^{-2}	NO_3^-	SO_4^{-2}	NO_3^-	SO_4^{-2}
Keilder Forest, UK 300 m Fog 11%	3	13	4	<1	1	7
Whitetop Mt., VA, USA 1,682 m Fog 35%	5	14	3	<1	40	120
Mt. Mitchell, NC, USA 2,006 m	5	—	3[d]	—	18	—

[a]Adapted from Unsworth and Wilshaw (1989), summarizing data of Fowler et al. (1989), Mueller (1988), and Aneja and Murthy (1994).
[b]Averaging periods and methods differ between sites.
[c]Elevation above sea level.
[d]Only 0.7% of dry deposition was particulate.

4.2.1.4 Magnitude of Deposition

Dry deposition of PM is most effective for coarse particles including primary geologic material and for elements such as iron and manganese. Wet deposition is most effective for fine particles of atmospheric (secondary) origin (e.g., nitrogen and sulfur, Table 4-6) and elements

such as cadmium, chromium, lead, nickel, and vanadium (Reisinger, 1990; Smith, 1990a,b,c; Wiman and Lannefors, 1985). The occurrence of occult deposition is more restricted. The relative magnitudes of the different deposition modes varies with ecosystem type, location, elevation, and chemical burden of the atmosphere. For the Walker Branch Watershed, a deciduous forest in rural eastern Tennessee, dry deposition constituted a major fraction of the total annual atmospheric input of cadmium and zinc (≈20%), lead (≈55%), and manganese (≈90%). Whereas wet deposition fluxes during precipitation events exceeded dry deposition fluxes by one to four orders of magnitude (Lindberg and Harriss, 1981), dry deposition was nearly continuous. Immersion of high-elevation forests in cloud water may occur for 10% or more of the year, significantly enhancing transfer of PM and dissolved gases to the canopy. Occult deposition in the Hawaiian Islands dominated total inputs of inorganic N (Heath and Huebert, 1999). Much of this N was volcanically derived during the generation of volcanic fog in part through reactions with seawater. In this humid climate, the dominance of occult rather than wet deposition is notable.

High-elevation forests receive larger particulate deposition loadings than equivalent low elevation sites. Higher wind speeds enhance the rate of aerosol impaction. Orographic effects enhance rainfall intensity and composition and increase the duration of occult deposition. Coniferous species in these areas with needle-shaped leaves also enhance impaction and retention of PM delivered by all three deposition modes (Lovett, 1984).

In more arid regions, such as the western United States, the importance of dry deposition may be larger. In the San Gabriel Mountains of southern California, for example, while annual deposition of SO_4^{-2} (partly of marine origin) was dominated by wet deposition (Fenn and Kiefer, 1999), deposition of NO_3^- was dominated by dry deposition, as was that of NH_4^+ at two of three sites. Similarly, at a series of low elevation sites in southern California (Padgett et al., 1999), dry deposition of NO_3^- was dominated by dry deposition. In both cases, however, the contribution of gaseous HNO_3 was probably substantial.

Nitrates, Sulfates and Cations

Much particulate sulfate and nitrate is found on particles in the 0.1- to 1.0-μm size range (U.S. Environmental Protection Agency, 1982), but most of these and of base cation and heavy metal inputs to forested ecosystems results from the deposition of larger particles (Lindberg and

Lovett, 1985; Lindberg et al., 1982). The influence of aerodynamic diameter is particularly critical for nitrogen species, because they exist as a wide range of particle sizes in the atmosphere (Milford and Davidson, 1987). For example, at many sites in North America, NO_3^- is characterized by a bimodal size distribution with modes above and below 1 μm. The supermicron particles are often the result of reactions between HNO_3 and coarse alkaline aerosols (Wolff, 1984) as, for example, in the San Joaquin Valley of California (Lindberg et al., 1990). Although the annual deposition of NH_4^+ is distributed similarly among the fine and coarse particles, particulate NO_3^- is found predominantly in the coarse-particle fraction (Table 4-7). Similar to the pattern for NH_4^+, the estimated annual deposition of SO_4^{-2} particles occurs in both the fine- and coarse-particulate fractions (Table 4-8), while base cation deposition is virtually restricted to contributions from coarse particles (Table 4-9).

Although the annual chemical inputs to ecosystems from particle deposition is significant by itself, it is important to compare it with the total chemical inputs from all sources of atmospheric deposition (i.e., precipitation, particles, and gaseous dry deposition). Figure 4-6 shows the mean percentage contribution of NO_3^- and NH_4^+, SO_4^{-2}, and base cation-containing particles to the total nitrogen, sulfur, and base cation deposition load to forest ecosystems (derived from Tables 4-7 through 4-9). Although the mean contribution of particulate deposition to cumulative nitrogen and sulfur deposition is typically less than 20% of annual inputs from all atmospheric sources, particulate inputs of base cations average half the total base cations entering forest ecosystems from the atmosphere.

An extensive comparison of particle to total chemical deposition is provided by the Integrated Forest Study (IFS) (Johnson and Lindberg, 1992a; Lovett, 1994; Lovett and Lindberg, 1993; Lindberg and Lovett, 1992; Ragsdale et al., 1992). Other similar data sets are available (Kelly and Meagher, 1986; Miller et al., 1993; Lindberg et al., 1986, 1990). These data in (Tables 4-7 through 4-9) clearly indicate that the contribution of coarse and fine aerosols to deposition to forest ecosystems is strongly dependent on the chemical species.

Dry deposition is an important flux of sulfur and nitrogen compounds at all of the IFS sites and ranges from 9 to 59% of total (wet + dry + cloud) deposition for sulfur, 25% to 70% for NO_3^-, and, 2% to 33% for NH_4^+. For only NH_4^+ is wet deposition consistently greater than dry deposition (Lovett, 1994).

TABLE 4-7. MEAN ANNUAL NITROGEN DEPOSITION (equivalents/ha/yr) FROM FINE AND COARSE PARTICLES COMPARED TO TOTAL NITROGEN DEPOSITION FROM ALL SOURCES TO A VARIETY OF FOREST ECOSYSTEMS

		Dry Particle Deposition						
		Fine		Coarse				
Region/Forest Type	Location	NO_3^-	NH_4^+	NO_3^-	NH_4^+	Total	Annual Total Nitrogen[a]	References[b]
North America								
Douglas fir	Washington	2	9	58	27	96	345	1,2
Loblolly pine	Georgia	2	27	36	37	102	647	1,2
Loblolly pine	North Carolina	3	35	89	66	193	997	1,2
Loblolly pine	Tennessee	0.8	18	27	7	53	699	1,2
Loblolly pine	Tennessee	0.2	14	8	1	23	410[c]	3
Slash pine	Florida	6	16	105	4	131	431	1,2
White pine	North Carolina	1	22	23	23	69	510	1,2
Red spruce	Maine	2	14	64	102	182	545	1,2
Red spruce	North Carolina	3	74	133	43	253	1,939	1,2
Red spruce	New York	1	9	5	2	17	1,136	1,2
Alder	Washington	1	5	58	27	91	339	1,2
Maple/beech	New York	0.3	5	37	13	55	567	1,2
Oak	Tennessee					307	857	4
Oak	Tennessee	1	36	83	8	128	720	5
Europe								
Norway spruce	Norway	8	21	52	29	110	775	1,2
Norway spruce	Germany	21	62	56	4	143	1,250[c]	3

[a]Includes deposition from precipitation, gases, and particles.
[b]1 = Johnson and Lindberg (1992a), 2 = Lovett (1992) and Lovett and Lindberg (1993), 3 = Lindberg et al. (1990), 4 = Kelly and Meagher (1986), 5 = Lindberg et al. (1986).
[c]Includes only the growing season from April to October 1987.

TABLE 4-8. MEAN ANNUAL SULFATE DEPOSITION (equivalents/ha/yr) FROM FINE AND COARSE PARTICLES COMPARED TO TOTAL SULFUR DEPOSITION FROM ALL SOURCES TO A VARIETY OF FOREST ECOSYSTEMS

		Dry Particle Deposition				
Region/Forest Type	Location	Fine	Coarse	Total	Annual Sulfur Deposition[a]	References[b]
North America						
Douglas fir	Washington	12	62	74	320	1,2
Loblolly pine	Georgia	47	74	121	776	1,2
Loblolly pine	North Carolina	57	59	116	1,050	1,2
Loblolly pine	Tennessee	27	69	96	941	1,2
Loblolly pine	Tennessee	22	44	66	570[c]	3
Slash pine	Florida	25	129	154	514	1,2
White pine	North Carolina	33	55	88	552	1,2
Red spruce	Maine	34	115	149	585	1,2
Red spruce	North Carolina	135	161	296	2,214	1,2
Red spruce	New York	10	4	14	1,096	1,2
Alder	Washington	6	62	68	325	1,2
Maple/beech	New York	7	32	39	488	1,2
Oak forest	Tennessee	70	190	260	1,600	4
Europe						
Norway spruce	Germany	58	97	155	1,100[c]	3
Norway spruce	Norway	19	79	98	663	1,2

[a]Includes deposition from precipitation, gases, and particles.
[b]1 = Johnson and Lindberg (1992a), 2 = Lindberg (1992) and Lindberg and Lovett (1992), 3 = Lindberg et al. (1990); 4 = Lindberg et al. (1986).
[c]Includes only the growing season from April to October 1987.

DRAFT-DO NOT QUOTE OR CITE

TABLE 4-9. MEAN ANNUAL BASE CATION DEPOSITION (equivalents/ha/yr) FROM FINE AND COARSE PARTICLES COMPARED TO TOTAL BASE CATION DEPOSITION FROM ALL SOURCES TO A VARIETY OF FOREST ECOSYSTEMS

		Dry Base Cation Particles Deposition[a]				
Region/Forest Type	Location	Fine	Coarse	Total	Total Annual Deposition[b]	References[c]
North America						
Douglas fir	Washington	9	180	189	670	1,2
Loblolly pine	Georgia	17	130	147	300	1,2
Loblolly pine	North Carolina	30	340	370	780	1,2
Loblolly pine	Tennessee	6.1	220	226	410	1,2
Mixed Hardwood	Tennessee	9.5[d]	121[d]	131[d]	899[d]	3
Slash pine	Florida	17	600	617	1,200	1,2
White pine	North Carolina	4	150	154	280	1,2
Red spruce	Maine	5	150	155	240	1,2
Red spruce	North Carolina	25	440	465	1,500	1,2
Red spruce	New York	2	36	38	230	1,2
Alder	Washington	4	180	184	660	1,2
Maple/beech	New York	1	84	85	200	1,2
Oak forest	Tennessee	11[e]	312[e]	323[e]	452[e]	3
Europe						
Spruce	Germany[d]					3
Spruce	Norway	6	180	186	390	1,2

[a]Includes K^+, Na^+, Ca^{+2}, and Mg^{+2}.
[b]Includes deposition from precipitation, gases, and particles.
[c]1 = Ragsdale et al. (1992), 2 = Johnson and Lindberg (1992a), 3 = Lindberg et al. (1986).
[d]Includes H^+, K^+, Na^+, Ca^{+2}, Mg^{+2} for the growing season from April to October 1987.
[e]Includes only Ca^{+2} and K^+.



Figure 4-6. Mean (±SE) percent of total nitrogen, sulfur, or base cation deposition contributed by fine plus coarse particles. Data are means from Tables 4-7 through 4-9.

After emission from their sources, air pollutants are transformed and transported by atmospheric processes (i.e, atmospheric meteorology) until deposited from the atmosphere to an aquatic or terrestrial ecosystem. As a result, ground-level concentrations of an air pollutant depend on the proximity to the sources, prevailing meteorology, and nature and extent of atmospherical reactions between the source and the receptor (Holland et al., 1999). A more direct relationship exists between source strength and downwind ambient concentrations for primary air pollutants (e.g., SO_2) than for secondary pollutants (e.g., sulfate, SO_4^{2-}). Interaction of the chemical and physical atmospheric processes and source locations for all of the pollutants have a tendency to produce data patterns that show large spatial and temporal variability.

Holland et al. (1999) analyzed CASTnet monitoring data and using generative additive models (GAM) estimated the form and magnitude of trends of airborne concentrations of SO_2, SO_4^{2-}, and nitrogen from 1989 to 1995 at 34 rural long-term CASTnet monitoring sites in the eastern United States. These models provide a highly flexible method for describing potential

nonlinear relationships between concentrations, meteorology, seasonality, and time (e.g., how weekly SO_2 varies as a function of temperature). For most of the 34 sites in the eastern United State, estimates of change in SO_2 concentrations showed a decreasing functional form in 1989-1990, followed by a relatively stable period during 1991-1993), then a sharper decline beginning in 1994 (Holland et al., 1999).

Regional trends of seasonal and annual wet deposition and precipitation-weighted concentrations (PWCs) of sulfate in the United States over the period 1980-1995 were developed by Shannon (1999) from monitoring date and scaled to a mean of unity. In order to reduce some effects of year to year climatological variability, the unitless regional deposition and PWC trends were averaged (hereafter referred to a CONCDEP). During the 16 year period examined in the study, estimated aggregate emissions of SO_2 in the United States and Canada fell approximately 12% from about 1980 to 1982, it remained roughly level for a decade and then fell approximately another 15% from 1992 to 1995 — for an overall decrease of about 18%. Eastern regional trends of sulfate concentrations and deposition and their average CONCDEP, also exhibited patterns of initial decrease, near steady state, and final decrease with year to-to-year variability. The overall relative changed in CONCDEPs are greater than the changes in SO_2 emissions.

Concentrations and calculated deposition (concentration times amount of water) of SO_4^{-2} at the Hubbard Brook Experiment Forest (HBEF) in the White Mountains of central New Hampshire have been measured since June of 1964 (Likens et al., 2001). These measurements represent the longest continuous record of precipitation chemistry in North America. The long-term measurements generally concur with those of Shannon (1999) discussed above. Major declines in emissions of SO_2 have been observed during recent decades in the eastern United States and have been correlated with significant decreases in SO_4^{-2} concentrations in precipitation (Shannon, 1999).

Deposition of sulfates and nitrates are very clearly linked to emissions. Reduction in emissions must occur before concentrations can be reduced below current levels (Likens, et al., 2001). Deposition is the key variable as sensitive ecosystems in the eastern North America have not yet shown improvement in response to decreased emissions of SO_2 (Driscoll et al., 1989; Likens et al., 1996). Clearly, additions of other chemicals, such as nitric acid and base cations, must be considered in addition to sulfur when attempting to resolve the acid rain problem

(Likens et al., 1996, 1998). The effects of sulfur and nitrogen deposition on ecosystems are discussed in Section 4.2.2.2.

The long-term record indicates that a reduction in the deposition of basic cations (Ca^{+2}, $Mg^{2+,}$ K^{+},Na^{+}) in bulk precipitation was associated with significant declines in sulfate deposition cited above for the HBEF region (Driscoll et al., 1989). Decreases in streamwater concentrations of basic cations have decreased simultaneously, suggesting that streamwater concentrations of basic cations are relatively responsive to changes in atmospheric inputs. Regardless of the cause, the decline in atmospheric influx of basic cations could have important effects on nutrient availability as well as on the acid/base status of soil and drainage water (Driscoll et al., 1989).

Trace Elements

Deposition velocities for fine particles to forest surfaces have been reported in the range of 1 to 15 cm s^{-1} (Smith, 1990a). For example, total, annual heavy metal deposition amounts are highly variable depending on specific forest location and upwind source strength (Table 4-10). Lindberg et al. (1982) quantified the dry deposition of heavy metals to inert surfaces and to leaves of an upland oak forest. As noted for other chemical species, V_d was highly dependent on particle size and chemical species (Table 4-11) with the larger particles depositing more efficiently.

TABLE 4-10. MEAN (±SE) PARTICLE SIZE, DEPOSITION RATES, AND DERIVED DEPOSITION VELOCITIES (V_d) FOR HEAVY METAL DEPOSITION TO THE UPPER CANOPY (INERT PLATES OR LEAVES) OF AN UPLAND OAK FOREST

Metal	Particle Size (μm)	Deposition Rate (pg cm^{-2} h^{-1})	V_d (cm s^{-1})
Manganese	3.4 ± 0.7	91 ± 23	6.4 ± 3.6
Cadmium	1.5 ± 0.7	0.3 ± 0.1	0.37 ± 0.18
Zinc	0.9 ± 0.2	6 ± 1	0.38 ± 0.1
Lead	0.5	23 ± 8	0.06 ± 0.01

Source: Lindberg et al. (1982).

TABLE 4-11. TOTAL HEAVY METAL DEPOSITION TO TEMPERATE LATITUDE FORESTS

Heavy Metal	Forest Deposition kg ha^{-1} $year^{-1}$ (Range)
Cadmium	0.002 - 0.02
Copper	0.016 - 0.24
Lead	0.099 - 1000
Nickel	0.014 - 0.15
Zinc	0.012 - 0.178

Source: Smith (1990c).

Preferential association of heavy metals with fine particles allows these particles to escape emission controls. Metal removal efficiencies for baghouse filters are typically 95 to 99% for all but mercury, but fine particle capture is much less efficient. Wet scrubber efficiency varies with design and pressure drop, typically 50 to 90% (McGowan et al., 1993). Fine particles also have the longest atmospheric residence times and, therefore, can be carried long distances. Depending on climate conditions and topography, fine particles may remain airborne for days to months and may be transported 1,000 to 10,000 km or more from their source. This long-distance transport and subsequent deposition qualify heavy metals as regional- and global-scale air pollutants. Ecosystems immediately downwind of major emissions sources (such as power generating, industrial, or urban complexes) may receive locally heavy inputs. Mass balance budgets (inputs and outputs) of seven heavy metals (cadmium, copper, iron, lead, manganese, nickel, and zinc) have been determined at the Hubbard Brook Experimental Forest (White Mountain National Forest) in New Hampshire. This forest is about 120 km northwest of Boston and relatively distant from major sources of heavy metal emissions. However, continental air masses that have passed over centers of industrial and urban activity also frequently follow storm tracks over northern New England. The resulting annual input for the seven heavy metals at Hubbard Brook for 1975 to 1991 is presented in Table 4-12. Inputs of most of the heavy metal species remained relatively constant over the 16-year period. The 44-fold decrease in lead deposition is correlated with removal of lead from motor vehicle fuels.

TABLE 4-12. ANNUAL[a] BULK DEPOSITION[b] INPUT OF SEVEN HEAVY METALS TO THE HUBBARD BROOK EXPERIMENTAL FOREST (43° 56' N LATITUDE, 71° 45' W LONGITUDE), WHITE MOUNTAIN NATIONAL FOREST, NH, FOR THE PERIOD 1975 TO 1991[c] (grams per hectare)

	Cadmium	Copper	Iron	Lead	Manganese	Nickel	Zinc
1975	2.5	18.2	832.0	352.0	100.0	29.4	175.0
1976	3.0	11.6	1,214.0	359.0	199.0	18.0	182.0
1977	40.0	10.0	372.0	195.0	39.0	8.0	116.0
1978	11.0	26.0	234.0	141.0	74.0	13.0	95.0
1979	16.0	16.0	207.0	155.0	172.0	12.0	278.0
1982	8.0	14.0	178.0	70.0	49.0	13.0	54.0
1983	8.0	17.0	206.0	57.0	52.0	7.5	76.0
1984	7.5	18.6	217.0	56.6	85.0	7.7	73.8
1985	6.3	9.0	174.0	41.0	53.8	7.0	54.8
1986	4.5	6.8	128.0	25.9	58.4	9.4	54.7
1987	2.3	4.5	16.0	17.2	55.8	10.7	45.2
1988	2.2	4.7	145.0	12.5	65.6	8.2	47.0
1989	1.6	3.3	160.0	11.9	74.2	7.4	57.2
1990	2.5	10.5	124.0	11.8	42.4	8.0	47.1
1991	2.1	15.0	134.0	8.3	72.0	9.6	55.5

[a]Mean of monthly totals.
[b]Total input, including both wet and dry deposition.
[c]Data not available for 1980 and 1981.

Trace element investigations conducted in roadside, industrial, and urban environments have demonstrated that impressive burdens of particulate heavy metals accumulate on vegetative surfaces. Lead deposition to roadside vegetation (prior to its removal from fuel) was 5 to 20, 50 to 200, and 100 to 200 times lead deposition to agricultural crops, grasses, and trees, respectively, in non-roadside environments. In an urban setting, it has been estimated that the leaves and twigs of a 30-cm (12-in.) diameter sugar maple remove 60, 140, 5800, and 820 mg of cadmium, chromium, lead, and nickel, respectively, during the course of a single growing season (Smith, 1973).

Effective deposition of PM is required before biological effects on plants or ecosystems can occur. It is clear that substantially improved techniques for monitoring and predicting deposition will be required to characterize these effects with certainty.

Semivolatile Organics

Organic compounds partition between gas and particle phases, and particulate deposition depends largely on the particle sizes available for adsorption (Pankow, 1987; Smith and Jones, 2000). Dry deposition of organic materials (eg. dioxins, dibenzofurans, polycyclic aromatics) is often dominated by the coarse fraction, even though mass loading in this size fraction may be small (Lin et al., 1993) relative to the fine PM fraction. For example, measurements in Bavaria in both summer and winter revealed that >80% of organics were in the fine (<1.35 μm) fraction (Kaupp and McLachlan, 1999). Nevertheless, in most cases, calculated values of dry deposition were dominated by the material adsorbed to coarse particles. Wet deposition, in contrast, was dominated by the much larger amount of material associated with fine particles. In this environment (where monthly precipitation is about 50 mm in winter and summer), wet deposition dominated, with dry deposition accounting for only 14 to 25% of total deposition (Kaupp and McLachlan, 1999). Lower relative contents of more volatile species in summer than winter (Kaupp and McLachlan, 1999) indicate the critical importance of gas-particle phase interconversions in determining deposition.

4.2.2 Effects on Vegetation and Ecosystems

Exposure to a given mass concentration of airborne PM may lead to widely differing phytotoxic responses, depending on the particular mix of deposited particles. The most common and useful subdivision of PM, derived from the typical bimodal distribution of atmospheric particles, is into fine and coarse particles (Wilson and Suh, 1997). The smallest particle at or near 1.0 to 2.5 μm generally is taken as the division between fine and coarse, although this is not an absolute and is subject to some shift (e.g., with changing ambient humidity). However, the typical the rule of thumb, as used in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a), is that fine PM nominally falls in the range of 0 to 2.5 μm and coarse-mode PM in the range of 2.5 to 10.0 μm.

Effects of particulate deposition on individual plants or ecosystems are difficult to characterize because of the complex interactions among biological, physicochemical, and climatic factors. Most direct effects other than regional effects associated with global changes occur in the severely polluted areas surrounding industrial point sources, such as limestone quarries, cement kilns, and metal smelting facilities. Fine particles have greater distribution. Experimental applications of PM constituents to foliage typically elicit little response at the more common ambient concentrations. The diverse chemistry and size characteristics of ambient PM and the lack of clear distinction between effects attributed to phytotoxic particles and to other air pollutants further confound understanding of the direct effects on foliar surfaces. The majority of documented toxic effect of particles on vegetation reflect their chemical content (acid/base, trace metal, nutrient), surface properties, or salinity. Studies of direct effects of particles on vegetation have not yet advanced to the stage of reproducible exposure experiments. Difficulties in experimental application of ambient particles to vegetation have been discussed by Olszyk et al. (1989). It is now clear that many phytotoxic gases are deposited more readily, assimilated more rapidly, and lead to greater direct injury to vegetation than do most common particulate materials (Guderian, 1986). The dose-specific responses (dose-response curves) obtained in early experiments following the exposure of plants to phytotoxic gases generally have not been observed following the application of particles.

Unlike gaseous dry deposition, neither the solubility of the particles nor the physiological activity of the surface is likely to be of first order of importance in determining deposition velocity (V_d). Factors that contribute to surface wetness and stickiness may be critical determinants of sticking efficiency. Available tabulation of deposition velocities are highly variable and suspect. High-elevation forests receive larger particle deposition loadings than equivalent lower elevations sites because of higher wind speeds and enhanced rates of aerosol impaction; orographic effects on rainfall intensity and composition; increased duration of occult deposition; and, in many areas, the dominance of coniferous species with needle-shaped leaves (Lovett, 1984). Recent evidence indicates that all three modes of deposition (wet, occult, and dry) must be considered in determining inputs to ecosystems or watersheds, because each may dominate over specific intervals of space.

Coarse-mode particles are primary in nature, having been produced and emitted from a point or area source as a fully formed particle. They generally range in size from ca. 2.5 to

100 μm. This material is created by abrasion and may be subsequently suspended by wind or mechanical means. Suspended geologic material contains the chemical and, potentially, the biological signature of the soil from which it derives (e.g., it may be dominated by iron, silica, aluminum, and/or calcium). Additional anthropogenically derived coarse-mode PM derives from fly ash, automobile tires and brake linings, and industrial effluent associated with crushing and grinding operations. Coarse-mode particles also include biogenically derived organic materials (e.g., fragments of plants and insects, pollen, fungal spores, bacteria, and viruses in marine aerosols).

In general, fine-mode PM is secondary in nature, having condensed from the vapor phase or been formed by chemical reaction from gaseous precursors in the atmosphere. These particles exist in a nucleation mode (having a mass median aerodynamic diameter or MMAD of about 0.06 μm) and may grow by coagulation of existing particles or by condensation of additional gases onto existing particles into an accumulation mode (about 0.5 μm). Sulfur and nitrogen oxides (SO_x and NO_x), as well as volatile organic gases, are common precursors for fine PM and are often neutralized with ammonium cations as particulate salts. Condensation of volatilized metals and products of incomplete combustion also are common precursors. Reactions of many of these materials with an oxidizing atmosphere lead to high secondary PM concentrations during the summer months in many parts of the United States.

Atmospheric PM may affect vegetation directly following deposition on foliar surfaces or indirectly by changing the soil chemistry or through changes in the amount of radiation reaching the Earth's surface through PM-induced climate change processes. Indirect effects, however, are usually the most significant because they can alter nutrient cycling and inhibit plant nutrient uptake. The possible direct responses to PM deposition are considered in this section, and the indirect responses are discussed in the later sections on ecosystems.

4.2.2.1 Direct Effects of Particulate Matter on Individual Plant Species

Particles transferred from the atmosphere to foliar surfaces may reside on the leaf, twig, or bark surface for extended periods; be taken up through the leaf surface; or be removed from the plant via resuspension to the atmosphere, washing by rainfall, or litter-fall with subsequent transfer to the soil. Any PM deposited on above-ground plant parts may exert physical or chemical effects. The effects of "inert" PM are mainly physical; whereas those of toxic particles

are both chemical and physical. The effects of dust deposited on plant surfaces or soil are more likely to be associated with their chemistry than simply with the mass of deposited particles and may be more important than any physical effects (Farmer, 1993). Nevertheless, vegetative surfaces represent filtration and reaction/exchange sites (Tong, 1991; Youngs et al., 1993).

Effects of Coarse Particles

Coarse particles, ranging in size from 2.5 to 100 μm, are chemically diverse, dominated by local sources, and typically deposited near their source because of their sedimentation velocities. Airborne coarse particles are derived from road, cement kiln, and foundry dust; fly ash; tire particles and brake linings; soot and cooking oil droplets; biogenic materials (e.g, plant pollen, fungal spores, bacteria and viruses); abraded plant fragments; sea salt; and hydrated deliquescent particles of otherwise fine aerosol. In many rural areas and some urban areas, the majority of mass in the coarse particle mode is in the elements silicon, aluminum, calcium, and iron, suggesting a crustal origin as fugitive dust from disturbed land, roadways, agriculture tillage, or construction activities. Rapid sedimentation of coarse particles tends to restrict their direct effects on vegetation largely to roadsides and forest edges.

Physical Effects—Radiation. Dust can cause physical and chemical effects. Deposition of inert PM on above-ground plant organs may result in an increase in radiation received, a rise in leaf temperature and the blockage of stomata. Increased leaf temperature, heat stress, reduced net photosynthesis, and leaf chlorosis, necrosis, and abscission were reported by Guderian (1986). Road dust decreased the leaf temperature on *Rhododendron catawbiense* by ca. 4 °C (Eller, 1977); whereas foundry dust caused an 8.7 °C increase in leaf temperature of black poplar (*Populus nigra*) under the conditions of the experiment (Guderian, 1986). Deciduous (broad) leaves exhibited larger temperature increases because of particle loading than did conifer (needle) leaves, a function of poorer coupling to the atmosphere. Inert road dust caused a three- to four-fold increase in the absorption coefficient of leaves of English ivy (Eller, 1977; Guderian, 1986) for near infrared radiation (NIR; 750 to 1350 nm). Little change in absorption occurred for photosynthetically active radiation (PAR; 400 to 700 nm). The increase in NIR absorption was equally at the expense of reflectance and transmission in these wavelengths. The net energy budget increased by ca. 30% in the dust-affected leaves. Deposition of coarse particles increased

leaf temperature and contributed to heat stress; reduced net photosynthesis; and caused leaf chlorosis, necrosis, and abscission (Dässler et al., 1972; Parish, 1910; Guderian, 1986; Spinka, 1971).

Starch storage in dust-affected leaves increased with dust loading under high (possibly excessive) radiation, but decreased following dust deposition when radiation was limiting. These modifications of the radiation environment had a large impact on single-leaf utilization of light. The boundary layer properties, determined by leaf morphology and environmental conditions, strongly influenced the direct effects of particle deposition on radiation heating (Eller, 1977; Guderian, 1986) and on gas exchange as well. Brandt and Rhoades (1973) attributed the reduction in growth of trees to crust formation from limestone dust on the leaves. Crust formation reduced photosynthesis and the formation of carbohydrates needed for normal growth, induced premature leaf-fall, damaged leaf tissues, inhibited growth of new tissue, and reduced starch storage. Dust may decrease photosynthesis, respiration, and transpiration; and it may allow penetration of phytotoxic gaseous pollutants, thereby causing visible injury symptoms and decreased productivity. Permeability of leaves to ammonia increased with increasing dust concentrations and decreasing particle size (Farmer, 1993).

Dust also has been reported to physically block stomata (Krajíčková and Mejstřík, 1984). Stomatal clogging by particulate matter from automobiles, stone quarries, and cement plants was also studied by Abdullah and Iqbal (1991). The percentage of clogging was low in young leaves when compared with old and mature leaves and the amount of clogging varied with species and locality. The maximum clogging of stomata observed was about 25%. The authors cited no evidence that stomatal clogging inhibited plant functioning. The heaviest deposit of dust is usually on the upper surface of broad-leaved plants; whereas the majority of the stomata are on the lower surface where stomatal clogging would be less likely.

Chemical Effects. The chemical composition of PM is usually the key phytotoxic factor leading to plant injury. Cement-kiln dust on hydration liberates calcium hydroxide that can penetrate the epidermis and enter the mesophyll; and, in some cases, this has caused the leaf surface alkalinity to reach a pH of 12. Lipid hydrolysis, coagulation of the protein compounds, and ultimately plasmolysis of the leaf tissue result in reduction in the growth and quality of plants (Guderian, 1986). In experimental studies, application of cement kiln dust of known

composition for 2 to 3 days yielded dose-response curves between net photosynthetic inhibition or foliar injury and dust application rate (Darley, 1966). Lerman and Darley (1975) determined that leaves must be misted regularly to produce large effects. Alkalinity was probably the essential phytotoxic property of the applied dusts.

Salinity. Particulate matter enters the atmosphere from oceans following the mixing of air into the water column and the subsequent bursting of bubbles at the surface. The effervescence of bubbles on the surface of the ocean forcefully ejects droplets of sea water into the air. These droplets, concentrated by evaporation, are carried inland by wind and deposited on the seaward side of coastal plants (Boyce, 1954). This occurs largely in the surf line (i.e., near land and potentially sensitive terrestrial receptors). This process can be a significant source of sulfate, sodium, chloride, and trace elements (as well as living material) in the atmospheric aerosol impacting coastal vegetation. Sea-spray particles (Taback et al., 1979) are approximately 24% greater in size than 10 μm, and 54% are between 3 and 10 μm. Thus, approximately only 20% are fine (0 to 2.5 μm) particles; and deposition by sedimentation and impaction is concentrated near the coast. Airborne concentrations of this marine PM decrease quickly with distance inland from the surfline both by deposition and dilution within atmospheric mixed layer (McKay et al., 1994; Nelis et al., 1994). Near-shore sediments with associated pollutants present in coastal runoff may be suspended in the surf and reentrained into the air. This can be a substantial source of microorganisms and of radionuclides to coastal vegetation (Nelis et al, 1994; McKay et al., 1994).

Sea-salt particles can serve as nuclei for the absorption and subsequent reaction of other gaseous and particulate air pollutants. Both nitrate and sulfate from the atmosphere have been found to associate with coarse and fine sea-salt particles (Wu and Okada, 1994). Direct effects on vegetation reflect these inputs, as well as classical salt injury caused by the sodium and chloride that constitute the bulk of these particles. Foliar accumulation of airborne salt particles may lead to foliar injury, thusly affecting the species composition in coastal environments (Smith, 1984).

The effects of winds and sea spray on coastal vegetation has been reported in the literature since the early 1800s (Boyce, 1954). However, there has been a difference of opinion as to whether the injury to coastal vegetation resulted from windblown aerial salts or from mechanical

injury (i.e., sand blasting) due to wind alone. Though the significance of sea water dashed on fore dunes and rocky coasts had been recognized by several authors, Wells and Shunk (1937, 1938) and Wells (1939) were the first to recognize the importance of salt spray in coastal ecology. Wells and Shunk (1937) reported that salt spray carried over dunes was the most important factor influencing growth form, zonation, and succession in coastal dunes. Salt spay injury was recorded 1.25 miles inland on the North Carolina coast. On the basis of observations in the Cape Fear area, they determined that the shape of coastal "wind form" shrubs were the result of sea spray carried by high winds. They found injury on shrubs only near the coast while those at a greater distances inland showed no injury whatsoever after a strong southeast wind that persisted for a period of nineteen hours during cloudy weather and abundant soil moisture.

To determine the cause of injury, injured and uninjured shoots were titrated for chlorides. A marked difference was observed between the injured and uninjured shoots (Wells and Shunk, 1937, 1938). Experimental spraying of shoots of woody plants with seawater resulted in a pattern of injury similar to the injury observed on seaside shrubs. The absence of the more inland species, such as persimmon (*Diosporos virginiana* L.), turkey oak (*Quercus laevis* Walt.), longleaf pine (*Pinus palustris* Mill., *P. australis* Michx.), and wire grass (*Aristida stricta* Michx.), was explained on the basis of intolerance of these species to salt spray. The dominance of live oak (*Quercus virginiana* Mill.), as a practically pure stand on Smith Island (also known as Bald Head Island), NC and along the eastern and southern NC coast, was determined by Wells (1939) to be due to the tree's tolerance to salt spray. He termed the long term stabilization of the live oak stand as a new type of climax, the "salt spray climax." The later work of Oosting and Billings (1942) near Beaufort, NC corroborated the findings of Wells and Shunk, 1937, 1938).

The report by Boyce (1954) is probably the most extensive on salt-spray communities. Dune sands in many coastal areas have been shown to have extremely low concentrations of dissolved salts. Studies have indicated that the salt content of the coastal dunes of Virginia, Massachusetts, and California did not exceed the maximum occurring in ordinary cultivated soils. Oosting and Billings (1942) found no correlation between soil salinity and plant distribution on the North Carolina coast. Surface crusts of sand dunes have been shown to have high concentrations of chlorides which could be attributed to sea spray, while concentrations of chlorides in underlying layers was low. The surface layer, however, varied with exposure of the dunes to oceanic winds (Boyce, 1954).

Boyce (1954), Wells (1939), and Wells and Shunk (1938) concluded on the basis of their studies that necrosis and death of plant tissues results from the high deposition of salt spray and high accumulation of the chloride ion in the plant tissues. Very little salt is taken up by plant roots; most enters through the aerial organs. Leaves of plants exposed to salt spray show a distinct pattern of injury (Wells and Shunk, 1938). Necrotic areas first appear at the leaf tips and upper margins and then progress slowly in an inverted "V" toward the petiole. This leaf injury pattern was verified experimentally. Mechanical injury resulting from leaves and twigs beating against each another in the wind causes the formation of small lesions through which salt can enter. After entry into the plant, the chloride ion is rapidly translocated to the apices of the leaves and twigs where it accumulates to injurious concentrations and results in the death of only a portion of the plant. The differential deposition and translocation of the chloride ion results in the death of the seaward leaves and twigs. The result is the continued growth of the uninjured branches in an inland direction. As a result, the canopy angle varies with the intensity of the spray (Boyce, 1954).

Little or no mineral ions are available in the silicate sands of the of coastal dunes. As a consequence, plants obtain the mineral ions needed for growth from the salt spray. Seawater contains all of the mineral ions required for growth, except nitrogen and phosphorus. The amount of nitrogen and phosphorus in seawater varies over a wide range (Boyce, 1954). Experiments indicated that available nitrogen in sea spray was a conditioning factor. Low nitrogen availability increased the tolerance of dune species to salt spray. Increasing the availability of nitrogen resulted in a different pattern of plant zonation and distribution. Dicotyledonous species were restricted to areas of lower spray intensity. The severity of chloride injury was associated more with the amount of available potassium than with the concentration of chlorides within the limits of 280-360 mg Cl/liter (Boyce, 1954).

Other sources of phytotoxic saline PM include aerosols from cooling towers and roadway deicing salt. Cooling towers used to dissipate waste heat from steam-electric power generating facilities may emit salt if brackish water or saltwater is used as a coolant (McCune et al., 1977; Talbot, 1979). Foliar injury is related to salt droplets deposited by sedimentation or impaction from cooling tower drift. The distance of the salt drift determines the amount of deposition and location of injury. Environmental conditions most conducive to injury were absence of precipitation, which can wash salt off leaves, and high relative humidity (RH; Talbot, 1979).

Increased injury is associated with wind speed and salt concentrations. Typical toxicity symptoms from acute exposures include marginal foliar necrosis and lesions; shoot-tip dieback; leaf curl; and interveinal necrosis (McCune et al., 1977). Based on experimental data, Grattan et al. (1981) observed that, to cause injury, salt deposited on leaf surfaces must dissolve and be absorbed into leaf tissue. Their work also indicated the importance of RH in foliar uptake. If RH remained below 70%, even heavy deposition of salt did not induce injury in peppers, soybeans, and tomatoes.

Injury to vegetation from the application of deicing salt was related to salt spray blown or drifting from the highways by Hofstra and Hall (1971) and Viskari and Kärenlampi (2000). The most severe injury was observed nearest to the highways. The results presented in these studies agrees with that of Wyttenbach et al. (1989), who observed that conifers planted near roadway margins in the eastern United States often exhibit foliar injury due to toxic levels of saline aerosols deposited from deicing solutions. Piatt and Kranse (1974) demonstrated that road and site factors influence the spread of deicing salt into forested areas. The slope away from the road influenced the distance from the road where injury was observed. The percent slope was correlated with the distance.

Effects of Fine Particles

Fine PM is generally secondary in nature, having condensed from the vapor phase or been formed by chemical reaction from gaseous precursors in the atmosphere, and is generally smaller than 1 to 2.5 μm. Nitrogen and sulfur oxides, volatile organic gases, condensates of volatilized metals, and products of incomplete combustion are common precursors for fine PM. Reactions of many of these materials with an oxidizing atmosphere contribute to high secondary PM concentrations during summer months in many U.S. areas. The conclusion reached in the 1982 PM AQCD (U.S. Environmental Protection Agency, 1982), that sufficient data were not available for adequate quantification of dose-response functions for direct effects of fine aerosols on vegetation, continues to be true today. Only a few studies on the direct effects of acid aerosols have been completed (U. S. Environmental Protection Agency, 1982). The major effects are indirect and occur through the soil (Section 4.3).

Nitrogen. Nitrate is observed in both fine and coarse particles. Nitrates from atmospheric deposition represent a substantial fraction of total nitrogen inputs to southeastern forests (Lovett and Lindberg, 1986). However, much of this is contributed by gaseous nitric acid vapor, and a considerable amount of the particulate nitrate is taken up indirectly through the soil. Garner et al. (1989) estimated deposition of nitrogen to forested landscapes in eastern North America at 10 to 55 kg/ha/year for nitrate and 2 to 10 kg/ha/year for ammonium. About half of these values were ascribed to dry deposition.

Atmospheric additions of particulate nitrogen in excess of vegetation needs are lost from the system, mostly as leachate from the soil as nitrate. Managed agricultural ecosystems may be able to utilize deposited particulate nitrogen more efficiently than native ecosystems, although many cultivated systems also lose considerable nitrogen as nitrate in runoff, deep drainage, or soil water. It has proven difficult to quantify direct foliar fertilization by uptake of nitrogen from ambient particles.

There is no doubt that foliar uptake of nitrate can occur, as clearly shown by the efficacy of foliar fertilization in horticultural systems. Potassium nitrate was taken up by leaves of deciduous fruit trees (Weinbaum and Neumann, 1977) and resulted in increased foliar nitrogen concentrations. Not all forms of nitrogen are absorbed equally, nor are all equally benign. Following foliar application of 2600 ppm of nitrogen as $Ca(NO_3)_2$, $(NH_4)_2SO_4$, or $(NH_2)_2CO$ to apple canopies (Rodney, 1952; Norton and Childers, 1954), leaf nitrogen levels were observed to increase to similar levels; but calcium nitrate and ammonium sulfate caused visible foliar injury, whereas urea did not. Urea is generally the recommended horticultural foliar fertilizer.

The mechanism of uptake of foliarly deposited nitrate is not well established. Nitrate reductase is generally a root-localized enzyme. It is generally not present in leaves, but is inducible there. Induction typically occurs when the soil is heavily enriched in NO_3^-. As the root complement of nitrate reductase becomes overloaded, unreduced nitrate reaches the leaves through the transpiration stream. Nitrate metabolism has been demonstrated in leaf tissue (Weinbaum and Neumann, 1977) following foliar fertilization. Residual nitrate reductase activity in leaves may be adequate to assimilate typical rates of particulate nitrate deposition. Uptake of nitrate may be facilitated by codeposited sulfur (Karmoker et al., 1991; Turner and Lambert, 1980).

Nitrate reductase is feedback-inhibited by its reaction product NH_4^+. The common atmospheric aerosol, NH_4NO_3, therefore may be metabolized in two distinct biochemical steps: first the ammonium (probably leaving nitric acid) and then the nitrate. Losses of nitric acid by volatization during this process, if they occur, have not been quantified.

Direct foliar effects of particulate nitrogen have not been documented. Application of a variety of fine nitrogenous aerosol particles (0.25 μm) ranging from 109 to 244 $\mu g/m^3$ nitrogen with or without 637 $\mu g/m^3$ sulfur caused no consistent short-term (2- to 5-h) effect on gas exchange in oak, maize, or soybean leaves (Martin et al., 1992).

Although no evidence exists for the direct transfer of nutrient particulate aerosols into foliage, a few studies give insights into the potential for ammonium and nitrate transfer into leaves. Fluxes of both NO_3^- and NH_4^+, measured in wet deposition and in throughfall plus stemflow in forests, commonly indicate higher fluxes of nitrogen above the canopy (Parker, 1983; Lindberg et al., 1987; Sievering et al., 1996), indicating net foliar uptake. Lovett and Lindberg (1993) reported a linear relationship between inorganic nitrogen fluxes in deposition and throughfall, suggesting that uptake may be considered passive to some extent.

Garten and Hanson (1990) studied the movement of ^{15}N-labeled nitrate and ammonium across the cuticles of red maple (*Acer rubrum*) and white oak (*Quercus alba*) leaves when applied as an artificial rain mixture. Brumme et al. (1992), Bowden et al. (1989), and Vose and Swank (1990) have published similar data for conifers. These studies show the potential for nitrate and ammonium to move into leaves, where it may contribute to normal physiological processes (e.g., amino acid production; Wellburn, 1990). Garten (1988) showed that internally translocated ^{35}S was not leached readily from tree leaves of yellow poplar (*Liriodendron tulipifera*) and red maple (*Acer rubrum*), suggesting that SO_4^{-2} would not be as mobile as the nitrogen-containing ions discussed by Garten and Hanson (1990). Further, when the foliar extraction method is used, it is not possible to distinguish sources of chemicals deposited as gases or particles (e.g., nitric acid [HNO_3], nitrogen dioxides [NO_2], nitrate [NO_3^-]), or sources of ammonium (deposited as ammonia [NH_3] or ammonium ion [NH_4^+]) (Garten and Hanson, 1990).

Particle deposition contributes only a portion of the total atmospheric nitrogen deposition reaching vegetation; but, when combined with gaseous and precipitation-derived sources, total nitrogen deposition to ecosystems has been identified as a possible causal factor leading to changes in natural ecosystems (See Section 4.2.3).

Sulfur. Anthropogenic sulfur emissions are >90% SO_2. Most of the remaining emission of sulfur is directly as sulfate (U.S. Environmental Protection Agency, 1996a). Sulfur dioxide is hydrophilic and is rapidly hydrated and oxidized to sulfite and bisulfite and then to sulfate, which is approximately 30-fold less phytotoxic. The ratio of SO_4^{-2}/SO_2 increases with aging of the air mass and, therefore, with distance from the source. Sulfate is sufficiently hygroscopic in humid air that it may exist significantly in the coarse particulate fraction. Because dilution of both SO_2 and particulate SO_4^{-2} occurs with distance from the source, it is unusual for damaging levels of particulate sulfate to be deposited. Gas to particle conversion in this case is of benefit to vegetation.

Sulfur is an essential plant nutrient. Low dosages of sulfur serve as a fertilizer, particularly for plants growing in sulfur-deficient soil (Hogan et al., 1998). However, current levels of sulfate deposition reportedly exceed the capacity of most vegetative canopies to immobilize the sulfur (Johnson, 1984). Nitrogen uptake in forests may be regulated loosely by sulfur availability, but sulfate additions in excess of needs do not typically lead to injury (Turner and Lambert, 1980).

There are few field demonstrations of foliar sulfate uptake (Krupa and Legge, 1986, 1998). Sulfate in throughfall is often enriched above levels in precipitation. The relative importance of foliar leachate and prior dry-deposited sulfate particles remains difficult to quantify (Cape et al., 1992). Leaching rates are not constant and may respond to levels of other pollutants, including acids. Uptake and foliar retention of gaseous and particulate sulfur are confounded by variable rates of translocation and accessibility of deposited materials to removal and quantification by leaf washing. Following soil enrichment with $^{35}SO_4^{-2}$ in a Scots pine forest, the apparent contribution of leachate to throughfall was only a few percent following an initial burst of over 90% because of extreme disequilibrium in labeling of tissue sulfate pools (Cape et al., 1992).

Olszyk et al. (1989) provide information on the effects of multiple pollutant exposures including particles (NO_3^-, 142 $\mu g/m^3$; NH_4^+, 101 $\mu g/m^3$; SO_4^{-2}, 107 $\mu g/m^3$). They found that only gaseous pollutants produced direct (harmful) effects on vegetation for the concentrations documented, but the authors hypothesized that long-term accumulation of the nitrogen and sulfur compounds contributed from particle deposition might have effects on plant nutrition over long periods of time. Martin et al. (1992) exposed oak (*Quercus macrocarpa*), soybean (*Glycine max*), and maize (*Zea mays*) plants to acute exposures (2 to 5 h) of aerosols (0.25 μm) containing

only nitrate (109 $\mu g/m^3$), ammonium and nitrate (244 and 199 $\mu g/m^3$), or ammonium and sulfate (179 and 637 $\mu g/m^3$). They found that these exposures, which exceeded the range of naturally occurring aerosol concentrations, had little effect on foliar photosynthesis and conductance. Martin et al. (1992) concluded that future investigations should focus on the effects of particles on physiological characteristics of plants following chronic exposures.

Acidic Deposition. The effects of acidic deposition have been accorded wide attention in the media and elsewhere (Altshuller and Linthurst, 1984; Hogan et al., 1998). Probably the most extensive assessment of acidic deposition processes and effects is the NAPAP Biennial Report to Congress: An Integrated Assessment (National Science and Technology Council, 1998). Concern regarding the effects of acidic deposition on crops and forest trees has resulted in extensive monitoring and research. Exposures to acidic rain or clouds can be divided into "acute" exposures to higher ionic concentrations (several μmol/L) and "chronic" long-term repeated exposures to lower concentrations (Cape, 1993). Pollutant concentrations in rainfall have been shown to have little capacity for producing direct effects on vegetation (Altshuller and Linthurst, 1984; Hogan et al., 1998); however, fog and clouds, which may contain solute concentrations up to 10 times those found in rain, have the potential to cause direct effects. More than 80% of the ionic composition of most cloud water is made up of four major pollutant ions: H^+, NH_4^+, NO_3^-, and SO_4^{-2}. Ratios of hydrogen to ammonium and sulfate to nitrate vary from site to site with all four ions usually present in approximately equal concentrations. Available data from plant effect studies suggest that hydrogen and sulfate ions are more likely to cause injury than ions containing nitrogen (Cape, 1993).

The possible direct effects of acidic precipitation on forest trees have been evaluated by experiments on seedlings and young trees. The size of mature trees makes experimental exposure difficult, therefore necessitating extrapolations from experiments on seedlings and saplings; however, such extrapolations must be used with caution (Cape, 1993). Both conifers and deciduous species have shown significant effects on leaf surface structures after exposure to simulated acid rain or acid mist at pH 3.5. Some species have shown subtle effects at pH 4 and above. Visible lesions have been observed on many species at pH 3 and on sensitive species at pH 3.5 (Cape, 1993). The relative sensitivities of forest vegetation to acidic precipitation based

on macroscopic injury have been ranked as follows: herbaceous dicots > woody dicots > monocots > conifers (Percy, 1991).

Huttunen (1994) described the direct effects of acid rain or acidic mist on epicuticular waxes whose ultrastructure is affected by plant genotype and phenotype. The effects of air pollutants on epicuticular waxes of conifers have received greater study than the waxes of other species. Leaf age and the shorter life span of broad-leaved trees make them less indicative of the effects of acid precipitation. Many experimental studies indicate that epicuticular waxes that function to prevent water loss from plant leaves can be destroyed by acid rain in a few weeks (Huttunen, 1994). This function is crucial in conifers because of their longevity and evergreen foliage. Microscopic observations of epicuticular wax structures have, for a long time, suggested links between acidic deposition and aging. In Norway spruce (*Picea abies*), acid rain causes not only the aging of needles (which in northern conditions normally last from 11 to 14 years) to be shortened, but also accelerates the erosion rate of the waxes as the needles age.

The effects of acidic precipitation and fog on red spruce (*Picea rubens*) have been studied extensively (Schier and Jensen, 1992). Visible foliar injury of the needles in the form of a reddish-brown discoloration has been observed on red spruce seedlings experimentally exposed to acidic mist, but this visible symptom has not been observed in the field. Ultrastructural changes in the epicuticular wax were observed both experimentally and on spruce growing at high elevations. Laboratory studies indicate that visible injury usually does not occur unless the pH is 3 or less (Schier and Jensen, 1992). Cape (1993) reported that, when compared with other species, red spruce seedlings appeared to be more sensitive to acid mist. From studies of conifers and a review of the literature, Huttunen (1994) concluded that acidic precipitation causes direct injury to tree foliage and indirect effects through the soil. The indirect effects of acidic precipitation are discussed in Section 4.3.

Based on a review of the many studies in the literature involving field and controlled laboratory experiments on crops, Cape (1993) drew a number of conclusions concerning the direct effects of acidic precipitation on crops:

- foliar injury and growth reduction occurs below pH 3;
- allocation of photosynthate is altered, with increased shoot to root ratios;
- expanded and recently expanded leaves are most susceptible, and injury occurs first to epidermal cells;

- leaf surface characteristics such as wettability, buffering capacity, and transport of material across the leaf surface contribute to susceptibility and differ among species;
- data obtained from experiments in greenhouses or controlled environmental chambers cannot be used to predict effects on plants grown in the field;
- quantitative data from experimental exposures cannot be extrapolated to field exposures because of differences and fluctuations in concentrations, durations, and frequency of exposure;
- there are large differences in response within species;
- timing of exposure in relation to phenology is of utmost importance;
- plants may be able to recover from or adapt to injurious exposures; and
- sequential exposure to acidic precipitation and gaseous pollutants is unlikely to be more injurious than exposure to individual pollutants.

Studies by Chevone et al. (1986), Krupa and Legge (1986), and Blaschke (1990) differ with the last conclusion of Cape listed above. Their studies indicate that interactions between acidic deposition and gaseous pollutants do occur. Acidity affects plant responses to both O_3 and SO_2. Chevone et al. (1986) observed increased visible injury on soybean and pinto bean when acid aerosol exposure preceded O_3 exposure; whereas linear decreases in dry root weight of yellow poplar occurred as acidity increased with simultaneous exposures to O_3 and simulated acid rain. Krupa and Legge (1986) also noted increased visible injury to pinto bean plants when aerosol exposure preceded O_3 exposure. In none of the studies cited above did acid rain per se produce significant growth changes. In contrast, Blaschke (1990) observed a decrease in ectomycorrhizal frequency and short root distribution caused by acid rain exposure in combination with either SO_2 or O_3.

Trace Elements. All but 10 of the 90 elements that comprise the inorganic fraction of the soil occur at concentrations of less than 0.1% (1000 μg/g) and are termed "trace" elements. Trace elements with a density greater than 6 g cm^{-3}, referred to as "heavy metals," are of particular interest because of their potential toxicity for plant and animals. Although some trace metals are essential for vegetative growth or animal health, they are all toxic in large quantities. Combustion processes produce metal chlorides that tend to be volatile and metal oxides that tend to be nonvolatile in the vapor phase (McGowan et al., 1993). Most trace elements exist in the

atmosphere in particulate form as metal oxides (Ormrod, 1984). Aerosols containing trace elements derive predominantly from industrial activities (Ormrod, 1984). Generally, only cadmium, chromium, nickel, and mercury are released from stacks in the vapor phase (McGowan et al., 1993). Concentrations of heavy metals in incinerator fly ash increase with decreasing particle size.

Vegetational surfaces, especially the foliage, present a major reaction and filtration surface to the atmosphere and act to accumulate particles deposited via wet and dry processes described in Section 4.2.1 (Tong, 1991; Youngs et al., 1993). The chemical constituents of particles deposited on foliar surfaces may be taken up through the leaf surface. The greatest particle loading is usually on the adaxial (upper) leaf surface where particles accumulate in the mid-vein, center portion of the leaves. Additionally, the mycelium of fungi becomes particularly abundant on leaf surfaces as the growing season progresses and is in intimate association with deposited particles (Smith, 1990c).

Investigations of trace elements present along roadsides and in industrial and urban environments indicate that impressive burdens of particulate heavy metals can accumulate on vegetative surfaces. Foliar uptake of available metals could result in metabolic effects in above-ground tissues. Only a few metals, however, have been documented to cause direct phytotoxicity in field conditions. Copper, zinc, and nickel toxicities have been observed most frequently. Low solubility, however, limits foliar uptake and direct heavy metal toxicity because trace metals must be brought into solution before they can enter into leaves or bark of vascular plants. In those instances when trace metals are absorbed, they are frequently bound in leaf tissue and are lost when the leaf drops off (Hughes, 1981). Trace metals in mixtures may interact to cause a different plant response when compared with a single element; however, there has been little research on this aspect (Ormrod, 1984). In experiments using chambers, Marchwinska and Kucharski (1987) studied the effects of SO_2 alone and in combination with PM components (Pb, Cd, Zn, Fe, Cu, and Mn) obtained from a zinc smelter bag filter. The combined effects of SO_2 and PM further increased the reduction in yield of beans caused by SO_2; whereas the combination, though severely injuring the foliage, produced little effect on carrots and parsley roots except after long-term exposures (when there was a decrease in root weight).

Trace metal toxicity of lichens has been demonstrated in relatively few cases. Nash (1975) documented zinc toxicity in the vicinity of a zinc smelter near Palmerton, PA. Lichen species

richness and abundance were reduced by approximately 90% in lichen communities at Lehigh Water Gap near the zinc smelter when compared with those at Delaware Water Gap. Zinc, cadmium, and sulfur dioxide were present in concentrations toxic to some species near the smelter; however, toxic zinc concentrations were detected farther away than the detectable limits of sulfur dioxide (Nash, 1975). Experimental data suggest that lichen tolerance to Zn and Cd falls between 200 and 600 ppm (Nash, 1975).

Though there has been no direct evidence of a physiological association between tree injury and exposure to metals, heavy metals have been implicated because their deposition pattern is correlated with forest decline. The role of heavy metals has been indicated by phytochelatin measurements. Phytochelatins are intracellular metal-binding peptides that act as specific indicators of metal stress. Because they are produced by plants as a response to sublethal concentrations of heavy metals, they can indicate that heavy metals play a role in forest decline (Gawel et al., 1996). Concentrations of heavy metals increased with altitude, as did forest decline, and increased concentrations across the study region that show increased levels of forest injury, as well.

Phytochelatin concentrations were measured in red spruce and balsam fir (*Abies balsamea*) needles throughout the 1993 growing season at 1000 m on Whiteface Mountain in New York. Mean foliar concentrations in red spruce were consistently higher than in balsam fir from June until August, with the greatest and most significant difference occurring at the peak of the growing season in mid-July. In July, the phytochelatin concentrations were significantly higher than at any other time measured. Balsam fir did exhibit this peak, but maintained a consistently low level throughout the season. Both the number of dead red spruce trees and phytochelatin concentrations increased sharply with elevation (Gawel et al., 1996). The relationship between heavy metals and the decline of forests in northeastern United States was further tested by sampling red spruce stands showing varying degrees of decline at 1000 m on nine mountains spanning New Hampshire, Vermont, and New York. The collected samples indicated a systematic and significant increase in phytochelatin concentrations associated with the extent of tree injury. The highest phytochelatin concentrations were measured during 1994 from sites most severely affected by forest decline in the Green Mountains, VT, and the Adirondack Mountains, NY. These data strongly imply that metal stress causes tree injury and contributes to forest decline in the northeastern United States (Gawel et al., 1996).

One potential direct effect of heavy metals is on the activity of microorganisms and arthropods resident on and in the leaf surface ecosystem. The fungi and bacteria living on and in the surfaces of leaves play an important role in the microbial succession that prepares leaves for decay and litter decomposition after their fall (U.S. Environmental Protection Agency, 1996b).

Numerous fungi were consistently isolated from foliar surfaces at various crown positions from London plane trees growing in roadside environments in New Haven, CT. Those existing primarily as saprophytes included *Aureobasidium pullulans, Chaetomium sp., Cladosporium sp., Epicoccum sp.*, and *Philaphora verrucosa*. Those existing primarily as parasites included *Gnomonia platani*, *Pestalotioposis sp.*, and *Pleurophomella sp.* The following cations were tested in vitro for their ability to influence the growth of these fungi: cadmium, copper, manganese, aluminum, chromium, nickel, iron, lead, sodium, and zinc. Results indicated variable fungal response with no correlation between saprophytic or parasitic activity and sensitivity to heavy metals. Both linear extension and dry weight data indicated that the saprophytic *Chaetomum sp.* was very sensitive to numerous metals. *Aureobasidium pullulans*, *Epicoccum sp.*, and especially *P.verrucosa*, on the other hand, appeared to be much more tolerant. Of the parasites, *G. platani* appeared to be more tolerant than *Pestalotiopsis sp.* and *Pleurophomella sp.* Metals exhibiting the broadest spectrum growth suppression were iron, aluminum, nickel, zinc, manganese, and lead (Smith and Staskawicz, 1977; Smith, 1990c). These in vitro studies employed soluble compounds containing heavy metals. Trace metals probably occur naturally on leaf surfaces as low-solubility oxides, halides, sulfates, sulfides, or phosphates (Clevenger et al., 1991; Koslow et al., 1977). In the event of sufficient solubility and dose, however, changes in microbial community structure on leaf surfaces because of heavy metal accumulation are possible.

Organic Compounds. Fine particles in the atmosphere reacting with volatilized chemical compounds are partitioned between the gas and particle phases, depending on the liquid phase vapor pressure at the ambient atmospheric temperature, the surface area of the particles per unit volume of air, the nature of the particles and of the chemical being adsorbed; and they can be removed by wet and dry deposition (McLachlan, 1996a). Materials as diverse as DDT, polychlorinated biphenyls (PCBs), and polynuclear aromatic hydrocarbons (PAHs) are being deposited from the atmosphere on rural as well as urban landscapes (Kylin et al., 1994). Motor

vehicles emit particles to the atmosphere from several sources in addition to the tailpipe. Rogge et al. (1993b) inventoried the organic contaminants associated with fine particles (diameter $\leq 2.0\ \mu m$) in road dust, brake-lining-wear particles, and tire tread debris. More than 100 organic compounds were identified in these samples, including n-alkanols, benzoic acids, benzaldehydes, polyalkylene glycol ethers, PAHs, oxy-PAH, steranes, hopanes, natural resins, and other compound classes. A large number of PAHs, ranging from naphthalene ($C_{10}H_8$) to 5- and 6-ring and higher PAHs, their alkyl-substituted analogues, and their oxygen- and nitrogen-containing derivatives are emitted from motor vehicle sources (Seinfeld, 1989).

Plants may be used as environmental monitors to compare the deposition of PAH, persistent organic pollutants (POPs), or semivolatile organic components (SOCs) between sites (e.g., urban versus rural; Wagrowski and Hites, 1997; Ockenden et al., 1998; McLachlan, 1999). Vegetation can be used qualitatively to indicate organic pollutant levels as long as the mechanism of accumulation is considered. The substance may enter the plant via the roots or, as mentioned above, be deposited as a particle onto the waxy cuticle of leaves or be taken up through the stomata. The pathways are a function of the chemical and physical properties of the pollutant such as its lipophilicity, water solubility, vapor pressure (which controls the vapro-particle partitioning) and Henry's law constant; environmental conditions such as ambient temperature and the organic content of the soil; and the plant species, which controls the surface area and lipids available for accumulation (Simonich and Hites, 1995). Ockenden et al. (1998) have observed that, for lipophilic POPs, atmospheric transfer to plant has been the main avenue of accumulation. Plants can differentially accumulate POPs. Results have shown differences between species with higher concentrations in the lichen (*Hypogymnia physiodes*) than in Scots pine needles (*Pinus sylvestris*). Even plants of the same species, because they have different growth rates and different lipid contents (depending on the habitat in which they are growing), have different rates of sequestering pollutants. These facts confound data interpretations and must be taken into account when considering their use as passive samplers.

Vegetation itself is an important source of hydrocarbon aerosols. Terpenes, particularly α-pinene, β-pinene, and limonene, released from tree foliage may react in the atmosphere to form submicron particles. These naturally generated organic particles contribute significantly to the blue haze aerosols formed naturally over forested areas (Geron et al., 2000).

The low water solubility with high lipoaffinity of many of these organic xenobiotics strongly control their interaction with the vegetative components of natural ecosystems. The cuticles of foliar surfaces are covered with a wax layer that helps protect plants from moisture and short-wave radiation stress. This epicuticular wax, consisting mainly of long-chain esters, polyesters, and paraffins, has been demonstrated to accumulate lipophilic compounds. Organic air contaminants in the particulate or vapor phase are absorbed to and accumulate in the epicuticular wax of vegetative surfaces (Gaggi et al., 1985; Kylin et al., 1994). Direct uptake of organic contaminants through the cuticle or the vapor-phase uptake through the stomates are characterized poorly for most trace organics. The phytotoxicity and toxicity of organic contaminants to soil microorganisms is not well studied (Foster, 1991).

Summary

Particulate matter transferred from the atmosphere may be deposited on above-ground plant parts and may exert physical or chemical effects or both. The effects of dust deposited on plant surfaces are more likely to be associated with their chemistry than simply with the mass of deposited particles. Studies of the effects of chemicals in PM deposited on foliage have found little or no effects on foliar processes unless exposure levels were significantly higher than typically would be experienced in the ambient environment. The majority of easily identified direct effects, other than climate, occur in severely polluted areas around heavily industrialized point sources such as limestone quarries, cement kilns, and smelting facilities for iron, lead, or various other metals. The direct effects of PM on foliar surfaces are confounded by the chemical nature and size characteristics of ambient airborne particles and the absence of a clear distinction between effects of PM on foliar surfaces and effects attributed to forms of air pollutants. Most documented toxic effects of particles on vegetation are associated with their acidity, trace metal content, nutrient content, surfactant properties, or salinity.

4.2.2.2 Indirect Effects of Particulate Matter on Natural Ecosystems

All life on this planet is dependent on the chemical energy in the form of carbon compounds to sustain their life processes. Terrestrial vegetation, via the process of photosynthesis, provides approximately half of the carbon that annually cycles between the Earth and the atmosphere (Chapin and Ruess, 2001). Plants do not live alone. They are members of

ecosystems, structurally complex communities comprised of populations of plant, animals (including humans), insects, and microorganisms that interact with one another and with their non-living (abiotic) chemical and physical environment. Ecosystems are dynamic, self-adjusting, self-maintaining, complex and adaptive systems, in which patterns at the higher levels of organization emerge from interactions and selection processes at localized levels (Levin, 1998).

Ecosystem components must have an adequate supply of energy, mineral nutrients, and water to maintain themselves and function properly. Sunlight is the energy source for most ecosystems. The energy obtained by plants (the producers) from sunlight during photosynthesis and chemical nutrients (e.g., nitrogen, phosphorus, sulfur) taken up from the soil are transferred to other species (the consumers) within the ecosystem through food webs. The movement of chemical nutrients through an ecosystem is cyclic, as the nutrients are used or stored and eventually returned to the soil by decomposer organisms. Energy, on the other hand, is transferred from organism to organism through an ecosystem in food webs and, finally, is dissipated into the atmosphere as heat (Odum, 1993). The flows of energy and cycling of nutrients provide the interconnectedness between ecosystem parts and transforms the community from a random collection of species into an integrated whole, an ecosystem, in which the biotic and abiotic parts are interrelated (Levin, 1998).

Macroscopic ecosystem properties (such as structure, diversity-productivity relationships, and patterns of nutrient and energy movement) that emerge from the interactions among the various components may feed back to influence subsequent development of those interactions. The relationship between structure and function is a fundamental one in ecosystem science. Ecosystem structure refers to the component species, their biodiversity, abundance, mass, and arrangement within an ecosystem. Ecosystem functions (energy flow, nutrient flux, water and material flow) are characterized by the way in which the components (e.g., plants, animals, and microorganisms) interact and the effect their activities have on the physical and chemical environment. Elucidating these interactions across scales is fundamental to understanding the relationships between biodiversity and ecosystem functioning (Levin, 1998).

Both ecosystem structure and functions play an essential role in providing ecosystem services. Human existence on this planet depends on the life-support services provided by ecosystem structure and functions (Daily, 1997). Ecosystem functions are characterized by the way components interact. These are the functions that maintain clean water, pure air, a vegetated

earth, and a balance of organisms, the functions that enable humans to survive. They are the dynamics of ecosystems. The benefits they impart include absorption and breakdown of pollutants, cycling of nutrients, binding of soil, degradation of organic waste, maintenance of a balance of gases in the air, regulation of radiation balance, climate, and the fixation of solar energy (Table 4-13; Westman, 1977; Daily, 1997). Economic benefits and values associated with ecosystem functions and services, and the need to preserve them because of their value to human life, are discussed by Costanza et al. (1997) and (Pimentel et al., 1997). Services usually are not considered to be items with market value.

TABLE 4-13. ECOSYSTEM SERVICES

- Purification of air and water – functions
- Mitigation of floods and droughts – structure and functions
- Detoxification and decomposition of wastes – functions
- Generation and renewal of soil and soil fertility – functions
- Pollination of crops and natural vegetation – functions
- Control of the vast majority of potential agricultural pests – functions
- Dispersal of seeds and translocation of nutrients – functions
- Maintenance of biodiversity, from which humanity has derived key elements of its agricultural, medicinal, and industrial enterprises
- Partial stabilization of climate
- Moderation of temperature extremes and the force of winds and waves
- Support of diverse human cultures
- Providing of aesthetic beauty and intellectual stimulation that lift the human spirit

Source: Daily (1997).

Concern has risen in recent years regarding the consequences of changing the biological diversity of ecosystems (Tilman, 2000; Ayensu et al., 1999; Wall, 1999; Hooper and Vitousek, 1997; Chapin et al., 1998). The concerns arise because human activities are creating disturbances that are causing the loss of biodiversity, altering the complexity and stability of ecosystems, and producing changes in nutrient cycling and the structure and function of ecosystems (Pimm, 1984; Levin, 1998; Chapin et al., 1998; Peterson et al., 1998; Tilman, 1996;

Tilman and Downing, 1994; Wall, 1999; Daily and Ehrlich, 1999). Changes in ecosystem structure and function affect the ecosystem services vital to human life.

There are few ecosystems on earth today that are not influenced by humans (Freudenburg and Alario, 1999; Vitousek et al., 1997; Matson et al. 1997; Noble and Dirzo, 1997). The scientific literature is filled with publications discussing the importance of ecosystem structure and function. Eco-risk, complexity, stability, biodiversity, resilience, sustainability, management, risk assessment, and ecosystem health, are frequently discussed topics. The deposition of particulate matter from the atmosphere has the potential to alter ecosystem structure and function by altering nutrient cycling and changing biodiversity. There is a need, therefore, to understand how ecosystems respond to both natural and anthropogenic stresses and, especially, the ways that anthropogenic stresses are impacting ecosystem services and products. Specifically, understanding the ecological effects of PM deposition is as important as quantifying the human health effects of PM deposition.

Ecosystem Response to Stresses

Ecosystem response to stress begins at the population level. Population change, however, begins with the response of individual plants or animals. Plant responses, both structural and functional, must be scaled in both time and space and propagated from the individual to the more complex levels of community interaction to produce observable changes in an ecosystem (see Figure 4-7). In an ecosystem, at least three levels of biological interaction are involved: (1) the individual plant and its environment; (2) the population and its environment; and (3) the biological community composed of many species and its environment (Billings, 1978). Individual organisms within a population vary in their ability to withstand the stress of environmental change. The response of individual organisms within a population is based on their genetic constitution (genotype), stage of growth at time of exposure to stress, and the microhabitat in which they are growing (Levin, 1998). The range within which these organisms can exist and function determines the ability of the population to survive. Those able to cope with the stresses survive and reproduce. Competition among the different species results in succession (community change over time) and, ultimately, produces ecosystems composed of populations of plant species that have the capability to tolerate the stresses (Rapport and Whitford, 1999; Guderian, 1985).



For a given level, the dot associated with a line begins with a process (e.g.photosynthesis for #1 under leaf) and ends with the associated structure (e.g., the needle).

Evaluating Impacts Within a Level of Organization

Leaf Level	Carbon exchange-1 Carbon pools-2 Needle number and size-3 Needle retention/abscission-4	Tree Level	Height and diameter growth-10 Crown shape and size-11 Tree vigor-12 Mortality-13
Branch Level	Carbon allocation-5 Branch growth-6 Branch morphology-7 Branch vigor-8 Branch retention-9	Stand Level	Productivity-14 Mortality-15 Species composition-16

Evaluating Interactions Between Different Levels of Organization



The diagonal arrow indicates the interaction between any two levels of organization.
The types of interaction are due to the properties of variability and compensation.

A - Refers to the interaction between the leaf and branch levels, where, for example, variability at the branch level determines leaf quantity, and compensation at the leaf level in photosynthesis may compensate for the reduction in foliage amount.

B - Refers to the interaction between the branch and the tree, where variability in branches determines initial interception, branch vigor, and branch location in the crown; compensation may be related to increased radiation reaching lower branches.

C - Refers to the interaction between the tree and the stand. Both genetic and environmental variability, inter- and intraspecific compensations, and tree historical and competitive synergisms are involved.

Figure 4-7. Effects of environmental stress on forest trees are presented on a hierarchial scale for the leaf, branch, tree, and stand levels of organization. The evaluation of effects within a level of organization are indicated by horizontal arrows. The evaluation of interactions between different levels of organization are indicated by diagonal arrows.

Source: Hinckley et al. (1992).

The number of species in a community usually increases during succession in unpolluted atmospheres. Productivity, biomass, community height, and structural complexity increase. Severe stresses, on the other hand, divert energy from growth and reproduction to maintenance and return succession to an earlier stage (Waring and Schlesinger, 1985). Ecosystems are subject to natural periodic stresses, such as drought, flooding, fire, and attacks by biotic pathogens (e.g., fungi, insects). Extremely severe natural perturbations return succession to an earlier stage; reduce ecosystem structure and functions (i.e., produce a scarcity of life forms and extinguish symbiotic interactions); disrupt the plant processes of photosynthesis and nutrient uptake, carbon allocation, and transformation that are directly related to energy flow and nutrient cycling; shorten food chains; and reduce the total nutrient inventory (Odum, 1993). This transformation, however, sets the stage for recovery that permits the perturbed ecosystem to adapt to changing environments (Holling, 1986). Therefore, these perturbations are seldom more than a temporary setback, and recovery can be rapid (Odum, 1969).

In contrast, anthropogenic stresses usually are severe, debilitating stresses. Severely stressed ecosystems do not recover readily, but may be further degraded (Odum, 1969; Rapport and Whitford, 1999). Anthropogenic stresses can be classified into four main groups: (1) physical restructuring (e.g., changes resulting from land use); (2) introduction of exotic species; (3) over harvesting; and (4) discharge of toxic substances into the atmosphere, onto land, and into water. Ecosystems lack the capacity to adapt to the above stresses and maintain their normal structure and functions unless the stress is removed (Rapport and Whitford, 1999). These stresses result in a process of degradation marked by a decrease in biodiversity, reduced primary and secondary production, and a lower capacity to recover and return to its original state. In addition, there is an increased prevalence of disease, reduced nutrient cycling, increased dominance of exotic species, and increased dominance by smaller, short-lived opportunistic species (Odum, 1985; Rapport and Whitford, 1999). Discharge of toxic substances into the atmosphere, onto land, and into water can cause acute and chronic stresses; and, once the stress is removed, a process of succession begins that can ultimately return the ecosystem to a semblance of its former structure. Air pollution stresses, if acute, are usually short term and the effects soon visible. Chronic stresses, on the other hand, are long-term stresses whose effects occur at different levels of ecosystem organization and appear only after long-term exposures, as in the

case of acidic deposition in the northeast or ozone in California (Shortle and Bondietti, 1992; U.S. Environmental Protection Agency, 1996b).

The possible effects of air pollutants on ecosystems have been categorized by Guderian (1977) as follows:

(1) accumulation of pollutants in the plant and other ecosystem components (such as soil and surface- and groundwater),
(2) damage to consumers as a result of pollutant accumulation,
(3) changes in species diversity because of shifts in competition,
(4) disruption of biogeochemical cycles,
(5) disruption of stability and reduction in the ability of self-regulation,
(6) breakdown of stands and associations, and
(7) expansion of denuded zones.

How changes in these functions can result from PM deposition and influence ecosystems is discussed in the following text. It should be remembered that, although the effects of PM are being emphasized, the vegetational components of ecosystems also are responding to multiple stresses from other sources.

Ecosystem Response to Direct Plant Effects

The presence of PM in the atmosphere may affect vegetation directly, following physical contact with the foliar surface (Section 4.2), but in most cases, the more significant effects are indirect. These effects may be mediated by suspended PM (i.e., through effects on radiation and climate) and by particles that pass through the vegetative canopies to the soil. Particulate matter, as considered in this chapter is a heterogeneous mixture of particles differing in size, origin, and chemical constituents, and their effects vary depending on the chemical nature of PM being deposited on vegetation or soil. Particulate inputs and ecosystem cycling of key elements are considered below.

The majority of studies dealing with direct effects of particulate dust and trace metals on vegetation have focused on responses of individual plant species and were conducted in the laboratory or in controlled environments (Saunders and Godzik, 1986). A few have considered

the effects of particles on populations, communities, and ecosystems. Most of these focused on ecosystems in industrialized areas heavily polluted by deposits of both chemically inert and active dusts. Effects can result from direct deposition or indirectly by deposition onto the soil. Reductions in growth, yield, flowering, and reproduction of plants from particulate deposition have been reported (Saunders and Godzik, 1986). Sensitivities of individual species have been associated with changes in composition and structure of natural ecosystems.

Evidence from studies of effects of PM deposition, specifically chemically inert and active dusts indicates that, within a population, plants exhibit a wide range of sensitivity, which is the basis for the natural selection of tolerant individuals. Rapid evolution of certain populations of tolerant species at sites with heavy trace element and nitrate deposition has been observed (Saunders and Godzik, 1986). Tolerant individuals present in low frequencies in populations when growing in unpolluted areas have been selected for tolerance at both the seedling and adult stages when exposed to trace metal or nitrate deposition (Ormrod, 1984; U.S. Environmental Protection Agency, 1993). Chronic pollutant injury to a forest community may result in the loss of sensitive species, loss of tree canopy, and maintenance of a residual cover of pollutant-tolerant herbs or shrubs that are recognized as successional species (Table 4-14; Smith, 1974).

Responses of ecosystems to stresses (unless severe or catastrophic) are difficult to determine because the changes are subtle (Garner, 1991). This is particularly true of responses to particles. Changes in the soil may not be observed until accumulation of the pollutant has occurred for 10 or more years, except in the severely polluted areas around heavily industrialized point sources (Saunders and Godzik, 1986). In addition, the presence of other co-occurring pollutants makes it difficult to attribute the effects to PM alone. In other words, the potential for alteration of ecosystem function and structure exists but is difficult to quantify, especially when there are other pollutants present in the ambient air that may produce additive or synergistic responses even though PM concentrations may not be elevated.

Physical Effects

The direct effects of limestone dust on plants and ecosystems has been known for many years. Long-term changes in the structure and composition of the seedling-shrub and sapling strata of an experimental site near limestone quarries and processing plants in Giles County in southwestern Virginia were reported by Brandt and Rhoades (1972, 1973). Dominant trees in the

TABLE 4-14. ECOSYSTEM FUNCTIONS IMPACTED BY AIR POLLUTION EFFECTS ON TEMPERATE FOREST ECOSYSTEMS

Forest Soil and Vegetation: Activity and Response	Ecosystem Consequence and Impact
1. Forest tree reproduction, alteration, or inhibition	1. Altered species composition
2. Forest nutrient cycling, alteration a. Reduced litter decomposition b. Increased plant and soil leaching and soil weathering c. Disturbance of microbial symbioses	2. Reduced growth, less biomass
3. Forest metabolism a. Decreased photosynthesis b. Increased respiration c. Altered carbon allocation	3. Reduced growth, less biomass
4. Forest stress, alteration a. Phytophagous insects, increased or decreased activity b. Microbial pathogens, increased or decreased activity c. Foliar damage increased by direct air pollution influence	4. Altered ecosystem stress: increased or decreased insect infestations; increased or decreased disease epidemics; and reduced growth, less biomass, and altered species composition

Source: Smith (1974).

control area, a part of the oak-chestnut association of the eastern deciduous forests of eastern North America, were chestnut oak *(Quercus prinus)*, red oak *(Q. rubra)*, and red maple *(Acer rubrum)*. An abundance of uniformly distributed saplings and seedlings were visible under the tree canopy, and herbs appeared in localized areas in canopy openings. Chestnut oak dominated the area, and the larger trees were 60 to 80 years old. The dusty site was dominated by white oak *(Q. alba)*; whereas red oak and tulip poplar *(Liriodendron tulipifera)* were subcodominants. The largest trees were 100 years old and had necrotic leaves, peeling bark, and appeared to be in generally poor condition except for tulip poplar (which thrived in localized areas). The site contained a tangled growth of seedlings and shrubs, a few saplings, and a prevalence of green briar *(Smilax spp.)* and grape *(Vitis spp)*. The sapling strata in the area was represented by red maple, hickory *(Carya spp.)*, dogwood *(Cornus florida)*, and hop-hornbeam *(Ostrya virginiana)*. Saplings of none of the leading dominant trees were of importance in this stratum. The most obvious form of vegetation in the seedling-shrub stratum, because of their

tangled appearance, were dogwood, hop-hornbeam, redbud *(Cercis canadensis)*, and sugar maple *(Acer saccarum)*.

Crust formation reduced photosynthesis, induced premature leaf fall and destruction of leaf tissues, inhibited growth of new tissue, and reduced the formation of carbohydrate needed for normal growth and storage (Brandt and Rhoades, 1973). The authors (Brandt and Rhoades, 1972), citing Odum (1969), also stated that one result of the accumulation of toxic pollutants in the biosphere (as the result of human activities) is the simplification of both plant and animal communities. In plant communities, structure is determined by sampling various strata within the community. Each stratum comprises a particular life form (e.g., herbs, seedlings, saplings, trees). Dust accumulation favored growth of some species and limited others. For example, sugar maple was more abundant in all strata of the dusty site when compared with the control site where it was present only as a seedling. The growth of tulip poplar, dogwood, hop-hornbeam, black haw *(Viburnum prunifolium)*, and redbud (*C. canadensis*) appeared to be favored by the dust. Growth of conifers and acidophiles such as rhododendron (*Rhododendron maximum*), however, was limited. Although dust accumulation began in 1945, the heaviest accumulation occurred between 1967 and 1972 during the time of the study.

Changes in community composition were associated closely with changes in the growth of the dominant trees. Decrease in density of seedlings and saplings and in mean basal area, as well as lateral growth of red maple, chestnut oak, and red oak, occurred in all strata. On the other hand, all of these characteristics increased in tulip poplar, which was a subordinate species before dust accumulation began but had assumed dominance at the time of the study. Reduction in growth of the dominant trees had apparently given tulip poplar competitive advantage because of its ability to tolerate dust. Changes in soil alkalinity occurred because of the heavy deposition of limestone dust; however, the facilities necessary for critical analysis of the soils were not available. From the foregoing, it is obvious that PM physical effects in the vicinity of limestone quarries and processing plants can affect ecosystems.

Changes in ecosystem structure resulting from exposures to sea salt were cited previously (Section 4.3.1.1). The dominance of live oak (*Quercus virginiana*) as a practically pure stand on Smith Island (Bald Head), NC and along the eastern and southern coast of North Carolina has been explained as due to its tolerance to salt spray. The absence of more inland species is

attributed to their intolerance to salt spray. Wells (1939) termed the long-term stabilization of live oak as "salt spray climax," a new type of climax.

Acidic Deposition

The effects of acidic deposition have been discussed in several previous reports. The 1982 EPA document, *Air Quality Criteria for Particulate Matter and Sulfur Oxides*, devoted a chapter to the effects of acidic deposition (U.S. Environmental Protection Agency, 1982). In 1984, EPA published *The Acidic Deposition Phenomenon and Its Effects* (Altshuller and Linthurst, 1984), and, in 1991, NAPAP published the result of its extensive study, *Acidic Deposition: State of Science and Technology* (Irving, 1991). The major effects of acidic deposition occur through the soil and are discussed under indirect effects. However, included among the direct responses of forest trees to acidic deposition are increased leaching of nutrients from foliage; accelerated weathering of leaf cuticular surfaces; increased permeability of leaf surfaces to toxic materials, water, and disease agents; and altered reproductive processes (Altshuller and Linthurst, 1984).

Trace Elements

Possible direct responses of trace elements on vegetation result from their deposition and residence on the phyllosphere (i.e., leaf surfaces). Fungi and other microorganisms living on the leaves of trees and other vegetation play an important role in leaf decomposition after litterfall (Miller and McBride, 1999; Jensen, 1974; Millar, 1974). Possible impacts of heavy metals on nutrient cycling and their effects on leaf microflora appear not to have been studied.

A trace metal must be brought into solution before it can enter into the leaves or bark of vascular plants. Low solubility limits entry. In those instances when trace metals are absorbed, they frequently are bound in the leaf tissue and then are lost from the plant when the leaf drops off (Hughes, 1981) are transferred to the litter layer where they can affect litter decomposition, an important source of soil nutrients. Changes in litter decomposition processes influence nutrient cycling in the soil and limit the supply of essential nutrients. Both Cotrufo et al. (1995) and Nikliňska et al. (1998) point out that heavy metals affect forest litter decomposition. Cotrufo et al. (1995) observed that decomposition of oak leaves containing Fe, Zn, Cu, Cr, Ni, and Pb was influenced strongly during the early stages by metal contamination. Fungal mycelium was significantly less abundant in litter and soil in contaminated sites when compared with control

sites. Nikliňska et al. (1998) stated that toxic effects of heavy metals on soil respiration rate have been reported by many scientists, and that, in polluted environments, this results in accumulation of undecomposed organic matter. However, they state that results of experiments should identify the most important "natural" factors affecting soil/litter sensitivity because the effects of heavy metals on respiration rates depend on the dose of heavy metals, the type of litter, types of metals deposited, and the storage time before respiration tests are made.

Trace metals, particularly heavy metals (e.g., cadmium, copper, lead, chromium, mercury, nickel, zinc) have the greatest potential for influencing forest growth (Smith, 1991). Experimental data indicate that the broadest spectrum of growth suppression of foliar microflora resulted from iron, aluminum, and zinc. These three metals also inhibited spore formation, as did cadmium, chromium, manganese, and nickel (see Smith, 1990c). In the field, the greatest injury occurs from pollution near mining, smelting, and other industrial sources (Ormrod, 1984). Direct metal phytotoxicity can occur only if the metal can move from the surface into the leaf or directly from the soil into the root.

Organic Compounds

Secondary organic compounds formed in the atmosphere, the effects of some of which are discussed below, have been referred to under the following terms: toxic substances, pesticides, hazardous air pollutants (HAPS), air toxics, semivolatile organic compounds (SOCs), and persistent organic pollutants (POPS). Again, it should be noted that the chemical substances denoted by such headings are not criteria air pollutants controlled by the NAAQS under Section 109 of the Clean Air Act (U.S. Code, 1991), but rather are controlled under Sect.112, Hazardous Air Pollutants. Their possible effects on humans and ecosystems are discussed in a number of government documents and in many other publications. They are mentioned here because many of the chemical compounds are partitioned between gas and particle phases in the atmosphere. As particles, they can become airborne, be distributed over a wide area, and affect remote ecosystems. Some of the chemical compounds are of concern because they may reach toxic levels in food chains of both animals and humans; whereas others tend to decrease or maintain the same toxicity as they move through the food chain. Some examples of movement through food chains are provided below.

Many chemical compounds from a variety of anthropogenic sources are released into the ambient air (see Section 4.2.1). In the atmosphere, the emitted compounds initially go through a mixing process, and the airborne particles then are distributed over a wide area and ultimately deposited on ecosystem components. Atmospheric deposition of polychlorinated dibenzo-*p*-dioxins and dibenzofurans (PCDD/Fs), as an example, can be divided into three different forms: (1) dry gaseous, (2) dry particle-bound, and (3) wet deposition. Dry particle-bound deposition occurs when the PM containing the pollutant is deposited on the plant surface; whereas wet deposition ranges from hail through rain to fog and dew fall (McLachlan, 1996b).

Human exposure to PCDD/Fs has been demonstrated to be caused almost exclusively by the ingestion of animal fat from fish, meat, and dairy products. Almost half of human exposure to PCDD/Fs is caused by consumption of beef and dairy products (McLachlan,1996b). Cattle obtain most of their PCCD/Fs though grass. Therefore, the grass–cattle–milk/beef pathway is critical for human exposure. It has been shown that root uptake/translocation is an insignificant pathway of PCDD/Fs to aerial plant parts. Wet and dry particle deposition are the most important for the accumulation of the higher chlorinated cogeners in vegetation. The persistence of PCDD/Fs in plants has not been investigated extensively; however, biodegradation probably does not occur in that these compounds are found primarily in the lipophilic cuticle and are very resistant to microbial degradation (McLachlan, 1996b). Feed contaminated with soil containing the pollutant can be another source of exposure of beef and dairy cattle, as well as chickens. The PCDD/Fs are near a steady state in milk cows and laying hens; however, animals raised for meat production (such as beef cattle and pigs) may accumulate them. The beef cattle and pigs cannot excrete the contaminants in a lipid-rich matrix such as milk or eggs. Thus, all of the PCDD/Fs, ingested are stored in the body. In agricultural food chains, there is a biodilution of PCDD/Fs, with the fugacity decreasing by up to three orders of magnitude between the air and cows milk (McLachlan, 1996b). Fürst et al. (1993), based on surveys to determine the factors that influence the presence of PCDD/PCDF in cows milk, earlier concluded that regardless of which pathway, soil → grass → cow or air → grass → cow, it was the congener of the chemical that was most important.

Chlorinated persistent organic pollutants (POPS), such as PCBs, PCDFs, and PCDDs, can be transported as particles through the atmosphere from industrial and agricultural sources; be brought down via wet and dry deposition in remote regions, such as the Arctic; and have been

detected in all levels of the Arctic food chain (Oehme et al., 1995). High concentrations of PCB (1 to 10 ppm) were found in seals, but the concentrations increased to 10 to 100 ppm in polar bears. The polar bear is the top predator in the Arctic and feeds preferentially on ringed seals and, to a lesser extent, on other seal species. Bioconcentration factors of organochlorines in the Arctic food web, reaching 10^7 for fish and seals, are biomagnified in polar bears (Oehme et al., 1995). Polychlorinated dibenzo-*p*-dioxins and polychlorinated dibenzofurans have also been found in seals (Oehme et al., 1995). Milk taken from anaesthetized polar bears was also found to contain PCDD/PCDF. Very little is known regarding the intake of milk by polar bear cubs. However, estimates of the intake of milk containing detectable levels of PCDD/PCDF and PCB and the additional consumption of seal blubber confirm that these pollutants are passed on to the next generation (Oehme et al., 1995).

Section 112 of the CAA, provides the legislative basis for U.S. hazardous air pollutant (HAP) programs. In response to mounting evidence that air pollution contributes to water pollution, Congress included Section 112m (*Atmospheric Deposition to Great Lakes and Coastal Waters*) in the 1990 CAA Amendments that direct the EPA to establish a research program on atmospheric deposition of HAPS to the "Great Waters".

Actions taken by EPA and others to evaluate and control sources of Great Waters pollutants of concern appear to have positively affected trends in pollutant concentrations measured in air, sediment, and biota. Details concerning these effects may be found in "Deposition of Air Pollutants to the Great Waters", Third Report to Congress (U. S. Environmental Protection Agency, 2000a). The Third Report (EPA-453/R-00-005, June 2000), like the First and Second Reports to Congress, focuses on 15 pollutants of concern, including pesticides, metal compounds, chlorinated organic compounds, and nitrogen compounds. The new scientific information in the Third Report supports and builds on three broad conclusions presented in the previous two EPA Reports to Congress and discussed below.

(1) Atmospheric deposition from human activities can be a significant contributor of toxic chemicals and nitrogen compounds to the Great Waters. The relative importance of atmospheric loading for a particular chemical in a water body depends on many factors (e.g., characteristics of the water body, properties of the chemical, and the kind and amount of atmospheric deposition versus or water discharges).

(2) A plausible link exists between emissions of toxic pollutants of concern into the air above the Great Waters; the deposition of these pollutants (and their transformation products); and the concentrations of these pollutants found in the water, sediments, and biota, especially fish and shellfish. For mercury, fate and transport modeling and exposure assessments predict that the anthropogenic contribution to the total amount of methylmercury in fish is, in part, the result of anthropogenic mercury releases from industrial and combustion sources increasing mercury body burdens (i.e., concentrations) in fish. Also, the consumption of fish is the dominant pathway of exposure to methylmercury for fish-consuming humans and wildlife. However, what is known about each stage of this process varies with each pollutant (for instance, the chemical species of the emissions and its transformation in the atmosphere).

(3) Airborne emissions from local as well as distant sources, from both within and outside the United States, contribute pollutant loadings to waters through atmospheric deposition. Determining the relative roles of particular sources—local, regional, national, and possibly global, as well as anthropogenic, natural, and reemission of pollutants—contributing to specific water bodies is complex, requiring careful monitoring, atmospheric modeling, and other analytical techniques.

Ecosystem Response to Indirect Effects of Particulate Matter

The presence of PM in the atmosphere directly affects vegetation following physical contact with foliar surfaces (as discussed above), but in many cases the more significant effects are indirect. These effects may be mediated by suspended PM (i.e., through effects on radiation and climate) and by particles that pass through vegetative canopies to reach the soil. Effects mediated in the atmosphere are considered briefly below and in greater detail later, under Section 4.5. Indirect plant responses are chiefly soil-mediated and depend primarily on the chemical composition of the individual elements deposited in PM. The individual elements must be bioavailable to have an effect. The soil environment, composed of mineral and organic matter, water, air, and a vast array of bacteria, fungi, algae, actinomycetes, protozoa, nematodes, and arthropods, is one of the most dynamic sites of biological interactions in nature (Wall and Moore, 1999; Alexander, 1977). The quantity of organisms in soils varies by locality. Bacteria and fungi are usually most abundant in the rhizosphere, the soil around plant roots that all

mineral nutrients must pass through. Bacteria and fungi benefit from the nutrients in the root exudates (chiefly sugars) in the soil and, in turn, they play an essential role by making mineral nutrients available for plant uptake (Wall and Moore, 1999; Rovira and Davey, 1974). Their activities create chemical and biological changes in the rhizosphere by decomposing organic matter and making inorganic minerals available for plant uptake. Bacteria are essential in the nitrogen and sulfur cycles and make these elements available for plant uptake and growth (see Section 4.3.3). Fungi are directly essential to plant growth. Attracted to the roots by the exudates, they develop mycorrhizae, a mutualistic, symbiotic relationship, that is integral in the uptake of the mineral nutrients (Allen, 1991). The impact in ecosystems of PM, particularly nitrates, sulfates, and metals, is determined by their effects on the growth of the bacteria involved in nutrient cycling and the fungi involved in plant nutrient uptake.

Atmospheric Turbidity: Effects on Vegetative Processes. Photosynthetic processes underlie the contribution of vegetative surfaces to nutrient and energy cycling. Photosynthesis and the heat-driven process of water cycling depend on net receipts and characteristics of the radiation environment. These characteristics may be altered substantially when the atmosphere becomes turbid because of particulate loading. Which wavelengths are of interest depends on the vegetation process under consideration. Canopy temperature and water relations are particularly sensitive to long-wave, infrared radiation; whereas primary photosynthetic charge separations depend on short-wave radiation in the visible and photosynthetically active range (0.4 to 0.7 μm).

Effects of anthropogenic aerosols on the radiation environment at the Earth's surface are difficult to assess. The residence time of suspended particles varies with size and environmental conditions (seconds to months or years), and concentrations are spatially and temporally variable. In particularly polluted urban and near-urban areas, unambiguous particulate effects on radiation and local climate may be observed. Visibility was degraded by 50% in a large plume originating in the St. Louis urban area during the midweek, midday period (Pueschel, 1993). In contrast, visibility was reduced by only 20% on weekends when traffic and industrial emissions were reduced. The area affected by the plume from the St. Louis urban area includes highly productive agricultural land.

Empirical relationships between the mass of specific components of the aerosol and radiation scattering have been developed (e.g., Pueschel, 1993), from which regional visibility

(or radiation attenuation) isopleths can be constructed if appropriate mass data are available. These estimates support trends observed by direct measurement of turbidity (e.g., Flowers et al., 1969; U.S. Environmental Protection Agency, 1982).

Sulfates, nitrates, and elemental carbon dominate effects on visibility, in part, because they frequently dominate the mass profiles and, in part, because they exhibit particularly large absorption coefficients (see Section 4.3). Absorption by particles containing carbon may range from 5 to 10% in rural areas to up to 50% in urban areas (U.S. Environmental Protection Agency, 1982). In west-coast cities with contrasting particulate sources and loadings, the common component that related PM to visibility degradation was sulfate between 0.65 and 3.6 μm (Barone et al., 1978). For example, in Los Angeles, sulfate and nitrate had similar effects on visibility (White, 1976), despite the dominance of nitrate from transportation sources in the aerosol, although this is changing with controls on point sources of sulfate (Farber et al., 1994).

A long-term global trend of increasing atmospheric optical depth has not been documented (Bolle et al., 1986; Pueschel, 1993) although seasonal and regional effects are substantial. The classic study by Flowers et al. (1969) demonstrated large regional distinctions in turbidity across the United States. Typically, the western deserts, plains, and Rocky Mountains exhibited low mean annual turbidity; whereas the more humid and densely vegetated eastern half of the country exhibited much greater turbidities. In the mid-1970s, visible range in the mountainous southwest exceeded 110 km and radiation attenuation was ca. 2.6%; whereas, in the East, visible range was below 24 km and radiation attenuation was ca. 10%. Visibility in the eastern United States has decreased generally since the 1940s (Flowers et al., 1969; Trijonis and Shapland, 1979; U.S. Environmental Protection Agency, 1982). Correlative trends in visibility degradation and emissions of sulfur oxides suggest that particulate sulfate may account for much of the turbidity.

These trends are typical of urban industrial areas around the world. Turbidity has increased above Mexico City (Binenko and Harshvardhan, 1993) since the 1911 to 1928 period. During this early period, a single annual peak of turbidity coincided with the end of the dry period, and natural sources dominated. By 1957 to 1962, the number of annual peaks had increased as anthropogenic sources came to dominate. During this period, atmospheric transmission of direct-beam solar radiation decreased by about 10% (Binenko and Harshvardhan, 1993). Visibility in the Los Angeles basin has improved very slightly in the past decades (Farber et al., 1994) as sulfate emissions have been controlled by regulation. The composition of the aerosol has

changed, particularly in inland areas, as the former dominance of sulfate shifts to a preponderance of secondary organics.

Particles interact with solar radiation through scattering and absorption. Absorption of short-wavelength solar radiation reduces the amount of radiation reaching the Earth's surface and leads to atmospheric heating. If the absorbing particles re-radiate in the infrared range, then some of this energy is lost as long-wave re-radiation to space. This loss mechanism is minimized because most of the anthropogenic aerosol in the troposphere resides in the planetary boundary layer (Bolle et al., 1986), even within the lower 500 m (Binenko and Harshvardhan, 1993) where the temperature is similar to that of the surface. Some of this energy is captured at the surface as down-welling infrared radiation.

These wavelengths directly affect canopy temperatures and influence transpirational water use by vegetation. The presence of absorbing aerosols reduces the ratio of photosynthetically active radiation to total radiation received at the surface, potentially reducing photosynthetic water use efficiency. The net effect of aerosol absorption on the surface depends on the relative magnitudes of the particulate absorption coefficients in the visible and infrared area and on the albedo of the Earth's surface. In general, absorption is not a dominant particulate effect.

Scattering of radiation dominates the effects of particulate loading on visibility and turbidity. Non-absorbing, scattering aerosols raise the overall albedo of the atmosphere and reduce the amount of radiation reaching the surface by the amount reflected or backscattered to space. As atmospheric turbidity increases, so does the scattering of light, including forward scattering of photosynthetically active radiation that intercepts the Earth's surface (Hoyt, 1978).

The largest effect is described by Mie-scattering theory. Forward scattering reduces the intensity of direct radiation by disrupting the solar beam, thereby increasing the path length and probability of absorption and by increasing the intensity of diffuse (sky) radiation. In a clear atmosphere, diffuse radiation may be on the order of 10% of total solar radiation (Choudhury, 1987). However, in highly turbid, humid conditions, this fraction may increase, even up to 100% of solar radiation in extreme cases. The direct-to-diffuse-radiation ratio is highest at solar noon and lowest near dawn or dusk when the path length through the atmosphere is longest.

Particle scattering is wavelength dependent, causing objects to appear blue- or red- tinged depending on viewing and illumination angles and on the light quality, the alteration of which is a minor contributor to photosynthetic light-use efficiency. The wavelength dependence of

scattering decreases rapidly from extreme sensitivity for very fine particles to little dependence at 10 μm. Equations relating scattering at a reference wavelength to scattering at wavelengths of interest are rigorously applicable only to spherical particles but may be extended to nonspherical particles of equal volume (Janzen, 1980).

World Meteorological Organization data summarized in U.S. Environmental Protection Agency (1982) indicated that turbidity in the eastern United States commonly resulted in radiation losses of ca. 3.5% because of backscattered radiation and ca. 3.5% because of absorption, with a resulting total reduction of incident radiation to ca. 93% of total solar radiation. However, 28% of the radiation reaching the surface was converted from direct radiation to diffuse, or sky, radiation. Under more polluted conditions, losses were ca. 9% backscattered and 9% absorbed, reducing total radiation to 82% of total solar radiation and converting 72% from direct beam to diffuse radiation. Photosynthetically active radiation (0.4 to 0.7 μm) typically is enriched in diffuse radiation relative to total or direct beam radiation.

Altered Radiative Flux: Effects on Vegetative Processes. Canopy photosynthesis is typically a nearly linear function of incident radiation, overcoming saturation exhibited by individual leaves by distributing the light throughout the multilayer canopy. Light penetration into canopies limits photosynthetic productivity (Rosenberg et al., 1983). The uppermost leaves of many canopies are at or above light saturation for photosynthetic processes. The simplest radiative transfer functions describing plant canopies relate total down-welling radiation (direct plus diffuse radiation measured above the canopy) to radiation interception at each leaf level through a Beer's Law analogy. The expected exponential decline in radiation through the canopy depends only on total radiation and a bulk canopy extinction coefficient that depends on leaf size, orientation, and distribution, as well as on reflectance and absorption in wavelengths of interest. These simplified models predict radiation distribution adequately for homogeneous canopies. Turbidity affects canopy processes only by attenuating the total radiation impinging on the canopy surface.

In more complex, and more realistic, canopy-response models (e.g., Choudhury, 1987), radiation is considered in its direct and diffuse components. Foliar interception by canopy elements is considered for both up- and down-welling radiation (a two-stream approximation). In this case, the effect of atmospheric PM on turbidity affects canopy processes both by radiation

attenuation and by influencing the efficiency of radiation interception throughout the canopy through conversion of direct to diffuse radiation (Hoyt, 1978). Diffuse radiation is more uniformly distributed throughout the canopy and increases canopy photosynthetic productivity by distributing radiation to lower leaves. The treatment of down-welling direct-beam radiation in the two-stream approach remains an elaboration of the simplified Beer's Law analogy with solar angle, leaf area distribution, and orientation individually parameterized (Choudhury, 1987). Diffuse down-welling radiation is a function of diffuse and direct radiation at the top of the canopy and penetration within the canopy according to cumulative leaf area density and foliage orientation. Up-welling (diffuse) radiation results from scattering and reflectance of both direct and diffuse down-welling radiation within the canopy and by the soil.

The altered distribution between diffuse and direct radiation affects photosynthesis in upper, exposed leaves as a function of leaf angle and in total canopy photosynthesis as a function of penetration of radiation within the canopy. This depends on canopy structure, leaf optical properties, and leaf area density, as well as on solar angle and atmospheric turbidity. Absorption of radiation by particles heats the upper atmosphere and results in reduced vertical temperature gradients. This could reduce the intensity of atmospheric turbulent mixing. The magnitude of such potential effects on turbulent transport within canopies remains unknown although damping of eddy transport could inhibit canopy gas exchange. Suppressed tropospheric mixing also could intensify local temperature inversions and increase the severity of pollution episodes (Pueschel, 1993) with direct inhibitory effects on photosynthetic processes.

The most significant effect of aerosols on vegetation is probably through their role as cloud condensation nuclei because clouds have a substantial effect on radiation receipts at the surface. An important characteristic of fine particles is their ability to affect the flux of solar radiation passing through the atmosphere directly, by scattering and absorbing solar radiation, and indirectly, by acting as cloud condensation nuclei which in turn influence the optical properties of clouds (Chameides et al., 1999). Regional haze has been estimated to diminish surface solar visible radiation by approximately 8%. Crop yields have been reported as being sensitive to the amount of sunlight received. The potentially significant effect of regional haze on the yield of crops because of reduction in solar radiation has been examined by Chameides et al. (1999). Using a case study approach, Chameides et al. (1999), studied the effects of regional haze on crop production in China where regional haze is especially severe. A rudimentary assessment of

the direct effect of atmospheric aerosols on agriculture suggests that optimal crop yields of approximately 70% of the crops are being depressed by at least 3 to 5% by regional scale air pollution and its associated haze (Chameides et al., 1999).

Effects of Solar Ultraviolet Radiation. The transmission of solar UV-B radiation through the earth's atmosphere is controlled by ozone, clouds and particles. The depletion of stratospheric ozone, caused by the release of chlorofluorcarbons (CFCs) and other substances such as halides, has resulted in heightened concern about potentially deleterious increases in the amount of solar UV-B (SUVB) radiation reaching the Earth's surface. One salient consideration is that, although CFC levels in the stratosphere have reached peak levels and now are beginning to fall as a result of the signing of the Montreal Protocol, the problem will likely continue well into the future because of the length of time it takes ozone-depleting molecules to reach the stratosphere (Greenberg, 1997).

The vulnerability of terrestrial plants to UV-B results from their requirement for sunlight for photosynthesis. Each 1% decline in stratospheric ozone has been predicted to decrease crop yield by 1% (Greenberg et al., 1997). In addition to inhibiting photosynthesis, UV-B radiation triggers numerous responses in plants, e.g.: membrane, protein, and DNA damage; delayed maturation; diminished growth; activation of chemical stress; flavonoid synthesis; and leaf thickening (Table 4-15). It is not known which of the injury and damage effects are most detrimental to plant growth (Table 4-15). Effects of increased UV-B on plant growth are likely to be incremental. Because plants evolved under the selective pressure of ambient UV-B radiation in sunlight, they have developed adaptive mechanisms (Greenberg et al., 1997). Although inhibition of photosynthesis is a detrimental growth effect, flavonoid synthesis represents acclimation. Plants growing under full light have been shown to be protected against UV-B effects, but not when growing under weak visible light (Björn, 1996). A common adaptation is alteration in leaf transmission properties, which results in attenuation of UV-B in the epidermis before it can reach the leaf interior.

Plant species vary enormously in their response to UV-B exposures, and large differences in response occur among different genotypes within a species. In general, dicotyledonous plants are more sensitive than monocotyledons from similar environments. In addition, plant responses may differ depending on stage of development. Therefore, extrapolation of experimental

TABLE 4-15. TYPES OF PLANT RESPONSES TO ULTRAVIOLET-B RADIATION[a]

Acclimation and Morphological Responses	Damage and Injury Responses
Altered biomass distribution	Altered gene expression
Altered leaf cell division	Degradation of auxin
Cotyledon curling	Degradation of chlorophyll and carotenoids
Increased DNA repair	Degradation of proteins
Increased flavonoid biosynthesis	Diminished biomass
Increased leaf thickness	Epidermal collapse
Increased leaf number	Inhibition of growth
Increased number of tillars	Inhibition of photosynthesis
Leaf wrinkling	Increased stomatal conductance
Reduced leaf area	Lower seed yield
Reduced hypocotyl growth	Oxidation of DNA
Reduced shoot height	Peroxidation of lipids
Reduced stomatal density	Prymidine dimer formation

[a]Entries in alphabetical order.

responses from seedlings to mature plants must be made with caution (Björn, 1996). The above facts are especially important when considering the effects of UV-B on agricultural plants. For example, among soybeans and rice, there are varieties for which growth and crop yield are severely decreased by increased UV-B radiation and other varieties that are not affected or may even be stimulated. On the other hand, the growth of the same sensitive soybeans when grown under water stress was not inhibited. Many crop plants grown in temperate regions originated in more tropical areas, hence, a gene pool for more resistant varieties is likely to exist (Björn, 1996). Crop plants, unlike forest trees and vegetation in natural ecosystems, are only exposed for one generation; and, thus, it may be possible to readily change the genotype if a variety proves to be sensitive.

Trees, forests, and perennial evergreen plants are long-lived when compared to agricultural systems, making it possible for UV-B exposure impacts to accumulate with time. Saplings and young and small trees react differently when compared to mature trees; also, on evergreen trees, needles of different ages respond differently (Björn, 1996). Breeding and testing trees is a slow process; and, for this reason, much care needs to be taken when planting large areas with trees of a single species and one provenance (e.g., Stika Spruce [*Picea sitchensis*] in Britain). The response of only a few broad-leaved trees have been studied. The most investigated genus has been loblolly pine (*Pinus taeda*; Björn, 1996).

A few studies indicate that the photomorphogenesis (changes in leaf thickness under UV-B that results in a transition from shade to sun leaves, Table 4-15) and the variable responses of native plants in ecosystems to UV-B exposures results in changes in interactions between various plants species, changes between plants and other organisms, and between plants and their abiotic environment. These preliminary studies suggest that in natural ecosystems, composed of many different plant species, with complex interactions between plants and between plants and other organisms, effects of UV-B may develop that cannot be determined from experiments on single plant species. The effects of UV-B on natural plant systems, therefore, should be of greater concern than on agricultural crops (Björn, 1996).

Effects of Nitrogen Deposition. Nitrogen has long been recognized as the nutrient most important for plant growth. Nitrogen is of overriding importance in plant metabolism and, to a large extent, governs the utilization of phosphorus, potassium, and other nutrients. Most of the nitrogen in soils is associated with organic matter. Typically, the availability of nitrogen via the nitrogen cycle controls net primary productivity, and possibly, the decomposition rate of plant litter. Photosynthesis is influenced by nitrogen uptake in that ca. 75% of the nitrogen in a plant leaf is used during the process of photosynthesis. The nitrogen-photosynthesis relationship is, therefore, critical to the growth of trees and other plants (Chapin et al., 1987). Plants usually obtain nitrogen directly from the soil through their roots by absorbing NH_4^+ or NO_3^-, or it is formed by symbiotic organisms in the roots. Plants, however, vary in their ability to absorb ammonium and nitrate (Chapin et al., 1987).

Because nitrogen is not readily available and is usually in shortest supply, it is the chief element in agricultural fertilizers. Atmospherically deposited nitrogen also can act as a fertilizer

in soil low in nitrogen. Not all plants, however, are capable of utilizing extra nitrogen. Inputs of nitrogen to natural ecosystems that alleviate deficiencies and increase growth of some plants can alter competitive relationships and alter species composition and diversity (Ellenberg, 1987; Kenk and Fischer, 1988; U.S. Environmental Protection Agency, 1993).

The effect of increasing nitrogen inputs (e.g., NO_x, nitrates, nitric acid) on the nitrogen cycle in forests, wetlands, and aquatic ecosystems is discussed in detail elsewhere (U.S. Environmental Protection Agency, 1993, 1997a; Garner, 1994; World Health Organization, 1997). The sources and forms of organic nitrogen in the atmosphere are poorly studied, and the concentrations are rarely measured, except in precipitation. Possible sources include particulate-entrained material from soils and vegetation (e.g., pollen, soil dust and spores) and reaction products of nitrogen oxides with organic compounds (e.g., peroxyacetyl nitrate, PAN; Lovett, 1992). The most important effects of nitrogen deposition are accumulation of nitrogen compounds resulting in the enhanced availability of nitrate or ammonium, soil-mediated effects of acidification, and increased susceptibility to stress factors (Bobbink et al., 1998). A major concern is "nitrogen saturation," the result of the deposition of large amounts of particulate nitrates. Nitrogen saturation results when additions to soil background nitrogen (nitrogen loading) exceeds the capacity of plants and soil microorganisms to utilize and retain nitrogen (Aber et al., 1989, 1998; Garner, 1994; U.S. Environmental Protection Agency, 1993). Under these circumstances, disruptions of ecosystem functioning may result (Hornung and Langan, 1999).

The growth of most forests in North America is limited by the nitrogen supply. Severe symptoms of nitrogen saturation, however, have been observed in high-elevation, nonaggrading spruce-fir ecosystems in the Appalachian Mountains, as well as in the eastern hardwood watersheds at Fernow Experimental Forest near Parsons, WV. Mixed conifer forests and chaparral watersheds with high smog exposure in the Los Angeles Air Basin also are nitrogen saturated and exhibit the highest stream water NO_3^- concentrations for wildlands in North America (Bytnerowicz and Fenn, 1996; Fenn et al., 1998). Forests in southern California, the southwestern Sierra Nevada in Central California, and the Front Range in northern Colorado have all been exposed to highly elevated nitrogen deposition, and nitrogen saturated watersheds have been reported in the above mentioned areas. Annual nitrogen additions through deposition (6-11 kg ha^{-1} y^{-1}as through fall) in the southwestern Sierra Nevada are similar to nitrogen storage

(4 kg ha^{-1} y^{-1}) in vegetation growth increment of western forests suggesting that current nitrogen deposition rates may be near the assimilation capacity of the overstory vegetation. Ongoing urban expansion will increase the potential for nitrogen saturation of forests from urban sources (e.g., Salt Lake City, Seattle, Tucson, Denver, central and southern California) unless there are improved emission controls (Fenn et al., 1998).

Not all forest ecosystems react in the same manner to nitrogen deposition. High-elevation alpine watersheds in the Colorado Front Range (Bowman, 2000) and a deciduous forest in Ontario, Canada, also are naturally saturated even though nitrogen deposition has been moderate ($\approx$8 kg ha^{-1} y^{-1}). The nitrogen saturated forests in North America, including estimated inputs and outputs, are shown in Table 4-16 (Fenn et al., 1998). The Harvard Forest hardwood stand in Massachusetts, however, has absorbed >900 kg N/ha without significant NO_3^- leaching during a nitrogen amendment study of 8 years (Table 4-16; Fenn et al.,1998). Johnson et al. (1991a) reported that measurements showing the leaching of nitrates and aluminum (Al^{+3}) from high elevation forests in the Great Smoky Mountains indicate that these forests have reached saturation.

Possible ecosystem responses to nitrate saturation, as postulated by Aber and coworkers (Aber et al., 1989), include (1) a permanent increase in foliar nitrogen and reduced foliar phosphorus and lignin caused by the lower availability of carbon, phosphorus, and water; (2) reduced productivity in conifer stands because of disruptions of physiological function; (3) decreased root biomass and increased nitrification and nitrate leaching; and (4) reduced soil fertility, resulting from increased cation leaching, increased nitrate and aluminum concentrations in streams, and decreased water quality. Saturation implies that some resource other than nitrogen is limiting biotic function.

Water and phosphorus for plants and carbon for microorganisms are the resources most likely to be the secondary limiting factors. The appearance of nitrogen in soil solution is an early symptom of excess nitrogen. In the final stage, disruption of forest structure becomes visible (Garner, 1994).

Changes in nitrogen supply can have a considerable effect on an ecosystem's nutrient balance (Waring, 1987). Large chronic additions of nitrogen influence normal nutrient cycling and alter many plant and soil processes involved in nitrogen cycling (Aber et al., 1989). Among the processes affected are (1) plant uptake and allocation, (2) litter production,

TABLE 4-16. NITROGEN-SATURATED FORESTS IN NORTH AMERICA, INCLUDING ESTIMATED N INPUTS AND OUTPUTS

Location	Forest Type	Elevation (m)	N Input (kg ha^{-1} $year^{-1}$)	N Output (kg ha^{-1} $year^{-1}$)	Reference
Adirondack Mts. northeastern New York	Northern hardwoods or hardwood/ conifer mix	396-661	9.3[a]	Stage 1 N loss[b]	Driscoll and Van Dreason (1993)
Catskill Mts., southeastern New York	Mainly hardwood; some eastern hemlock	335-675	10.2[a]	Stage 1 and 2 N loss[b]	Stoddard (1994)
Turkey Lakes Watershed, Ontario, Canada	Sugar maple and yellow birch	350-400	7.0-7.7 (as throughfall)	17.9-23.6	Foster et al. (1989); Johnson and Lindberg (1992a)
Whitetop Mt., southwestern Virginia	Red spruce	1650	32[c]	47[c]	Joslin and Wolfe (1992); Joslin et al. (1992)
Fernow, West Virginia	Mixed hardwood	735-870	15-20	6.1	Gilliam et al. (1996); Peterjohn et al. (1996)
Great Smoky Mts. National Park, Tennessee	American beech	1600	3.1[d]	2.9	Johnson and Lindberg (1992b)
Great Smoky Mts. National Park, Becking Site, North Carolina	Red spruce	1800	10.3[d]	19.2	Johnson et al. (1991a)
Great Smokey Mts. National Park, Tower Site, North Carolina	Red spruce	1740	26.6	20.3	Johnson et al. (1991a)
Front Range, Colorado	Alpine tundra, subalpine conifer	3000-4000	7.5-8.0	7.5	Williams et al. (1996)
San Dimas, San Gabriel Mts. southern California	Chapparral and grasslands	580-1080	23.3[e]	0.04-19.4	Riggan et al. (1985)
Camp Paivika, San Bernadino Mts., southern California	Mixed conifer	1600	30	7-26[f]	Fenn et al. (1996)
Klamath Mts, northern California	Western coniferous	NA	Mainly geologic[g]	NA[g]	Dahlgren (1994)
Thompson Forest, Cascade Mts., Washington	Red alder	220	4.7 plus > 100 as N_2 fixation	38.9	Johnson and Lindberg (1992b)

[a]Estimated total N deposition from wet deposition data is from Driscoll et al. (1991) for the Adirondacks, and from Stoddard and Murdoch (1991) for the Catskills. Total deposition was estimated based on the wet deposition/total N deposition ratio (0.56) at Huntington Forest in the Adirondacks (Johnson and Lindberg, 1992b). Nitrogen deposition can be higher in some areas, especially at high-elevation sites such as Whiteface Mountain (15.9 kg ha^{-1} $year^{-1}$; Johnson and Lindberg, 1992b).
[b]Stage 1 and 2 of N loss according to the watershed conceptual model of Stoddard (1994). Nitrogen discharge (kg ha^{-1} $year^{-1}$) data are not available; only stream water NO_3^- concentration trend data were collected.
[c]Values appear high compared to other sites, especially N leaching losses. Joslin and Wolfe (1992) concede that "there is considerable uncertainty associated with the estimates of atmospheric deposition and leaching fluxes." However, elevated NO_3^- concentrations in soil solution, and lack of a growth response to N fertilization (Joslin and Wolfe, 1994) support the hypothesis that the forest at Whitetop Mountain is N saturated.
[d]Estimated total N deposition from throughfall data. Total deposition was estimated based on the throughfall/total N deposition ration (0.56) from the nearby Smokies Tower site (Johnson and Lindberg, 1992b).
[e]Annual throughfall deposition to the chaparral ecosystem.
[f]Nitrogen output is from unpublished streamwater data (M.E. Fenn and M.A. Poth, 1999). The low value represents a year of average precipitation, and the high value is for 1995, when precipitation was nearly double the long-term average. Nitrogen output includes N export in streamwater and to groundwater.
[g]Annual input and output data are not known, although N deposition in this forest is probably typical for much of the rural western United States (2-3 kg N ha^{-1} $year^{-1}$ (Young et al., 1988). Excess N is from weathering of ammonium in mica schist bedrock. The ammonium was rapidly nitrified, leading to high NO_3^- concentrations in soil solution (Dahlgren, 1994).

DRAFT-DO NOT QUOTE OR CITE

(3) immobilization (includes ammonification [the release of ammonia] and nitrificatrion [conversion of ammonia to nitrate during decay of litter and soil organic matter]), and (4) nitrate leaching and trace gas emissions (Figure 4-8; Aber et al., 1989).



Figure 4-8. Nitrogen cycle (dotted lines indicate processes altered by nitrogen saturation).

Source: Garner (1994).

Subsequent studies have shown that, although there was an increase in nitrogen mineralization initially (i.e., the conversion of soil organic matter to nitrogen in available form [see item 3 above]), nitrogen mineralization rates were reduced under nitrogen-enriched

conditions. Also, studies suggest that soil microbial communities change from predominantly fungal (mycorrhizal) communities to those dominated by bacteria during saturation (Aber et al., 1998).

Because the competitive equilibrium of plants in any community is finely balanced, the alteration of one of a number of environmental parameters, (e.g., continued nitrogen additions), can change the vegetation structure of an ecosystem (Bobbink, 1998; Skeffington and Wilson, 1988). Increases in soil nitrogen play a selective role. When nitrogen becomes more readily available, plants adapted to living in an environment of low nitrogen availability will be replaced by plants capable of using increased nitrogen because they have a competitive advantage.

Plant succession patterns and biodiversity are affected significantly by chronic nitrogen additions in some North American ecosystems (Figure 4-9). The location of nitrogen saturated ecosystems in North America, and the steps leading to nitrogen saturation, are indicated on the map in Figure 4-9. Conceptual models of regional nitrogen saturation are located in New England, the Colorado alpine ecosystems and in California forests. Fenn et al. (1998) report that long-term nitrogen fertilization studies in both New England and Europe, as well, suggest that some forests receiving chronic inputs of nitrogen may decline in productivity and experience greater mortality. Long-term fertilization experiments at Mount Ascutney, Vermont, suggest that declining coniferous forest stands with slow nitrogen cycling may be replaced by deciduous fast-growing forests that cycle nitrogen rapidly (Fenn et al., 1998).

Competition among species can result in changes in community composition; therefore, it is one of the most notable responses to environmental change (Bowman, 2000). Nitrogen saturation, the result of increased deposition in the alpine tundra of Niwot Ridge in the Front Range of the Southern Rockies in Colorado has changed nitrogen cycling and provided the potential for replacement in plant species by more competitive, faster growing species (Bowman and Steltzer, 1998; Bowman, 2000; Baron et al., 2000). Plants growing in an alpine tundra, as is true of other plants growing in low resource environments (e.g., infertile soil, shaded understory, deserts), have been observed to have certain similar characteristics: a slow grow rate, low photosynthetic rate, a low capacity for nutrient uptake and low soil microbial activity (Bowman and Steltzer, 1998; Bowman, 2000). An important feature of such plants is that they continue to grow slowly and tend to respond even less when provided with an optimal supply and balance of resources (Pearcy et al., 1987; Chapin, 1991). Plants adapted to cold, moist environments grow



Figure 4-9. Diagrammatic overview of excess nitrogen (N) in North America.

Source: Fenn et al. (1998).

more leaves than roots as the relative availability of nitrogen increases; however, other nutrients may soon become limiting. These patterns of vegetative development affect their capacity to respond to variation in available resources and to environmental stresses such as frost, high winds, and drought. Preformation of buds 3-4 years in advance of emergence, reduced cell numbers, and high biomass allocation to belowground organs also limits the ability of many alpine plants to respond to variations in their environment (Bowman, 2000). However, significant interspecific genetic variation influences the capacity of the alpine species to respond to changes in resource availability. The capacity of subalpine and boreal species in particular, and gymnosperms in general, to reduce nitrates in either roots or leaves appears to be limited. In addition, the ability of trees to use nitrogen varies with the age of the tree and the density of the stand (Waring, 1987).

In experimental studies of nitrogen deposition conducted by Wedin and Tilman (1996) over a 12-year period on Minnesota grasslands, plots dominated by native warm-season grasses shifted to low-diversity mixtures dominated by cool-season grasses at all but the lowest rates of nitrogen addition. Grasslands with high nitrogen retention and carbon storage rates were the most vulnerable to loss of species and major shifts in nitrogen cycling. The shift to low-diversity mixtures was associated with the decrease in biomass carbon to nitrogen (C:N) ratios, increased nitrogen mineralization, increased soil nitrate, high nitrogen losses, and low carbon storage (Wedin and Tilman, 1996). Naeem et al. (1994) experimentally demonstrated (under controlled environmental conditions) that the loss of biodiversity, genetic resources, productivity, ecosystem buffering against ecological perturbation, and loss of aesthetic and commercially valuable resources also may alter or impair ecosystems services.

The long-term effects of increased nitrogen deposition have been studied in several western and central European plant communities: lowland heaths, species-rich grasslands, mesotrophic fens, ombrotrophic bogs, upland moors, forest-floor vegetation, and freshwater lakes (Bobbink, 1998). Large changes in species composition have been observed in regions with high nitrogen loadings or infield experiments after years of nitrogen addition (Bobbink et al., 1998). The increased input of nitrogen gradually increased availability of nitrogen in the soil, and its retention because of low rates of leaching and denitrification resulted in faster litter decomposition and rate of mineralization. Faster growth and greater height of nitrophilic species enables these plants to shade out the slower growing species, particularly those in oligotrophic or

mesotrophic conditions (Bobbink, 1998; Bobbink et al., 1998). Excess nitrogen inputs to unmanaged heathlands in the Netherlands has resulted in nitrophilous grass species replacing slower growing heath species (Roelofs et al., 1987; Garner, 1994). Van Breemen and Van Dijk (1988) noted that over the past several decades the composition of plants in the forest herb layers has been shifting toward species commonly found on nitrogen-rich areas. It also was observed that the fruiting bodies of mycorrhizal fungi had decreased in number.

Other studies in Europe point out the effects of excessive nitrogen deposition on mixed-oak forest vegetation along a deposition gradient largely controlled by soil acidity, nitrogen supply, canopy composition, and location of sample plots (Brunet et al., 1998; Falkengren-Grerup, 1998). Results of the study, using multivariate methods, suggest that nitrogen deposition has affected the field-layer vegetation directly by increased nitrogen availability and, indirectly, by accelerating soil acidity. Time series studies indicate that 20 of the 30 field-layer species (nonwoody plants) that were associated most closely with high nitrogen deposition increased in frequency in areas with high nitrogen deposition during the past decades. Included in the field-layer species were many generally considered nitrophilous; however, there were several acid tolerant species (Brunet et al, 1998). Falkengren-Grerup (1998), in an experimental study involving 15 herbs and 13 grasses, observed that species with a high nitrogen demand and a lesser demand for other nutrients were particularly competitive in areas with acidic soils and high nitrogen deposition. The grasses grew better than herbs with the addition of nitrogen. It was concluded that, at the highest nitrogen deposition, growth was limited for most species by the supply of other nutrients; and, at the intermediate nitrogen concentration, the grasses were more efficient than the herbs in utilizing nitrogen. Nihlgård (1985) suggested that excessive nitrogen deposition may contribute to forest decline in other specific regions of Europe. Also, Schulze (1989), Heinsdorf (1993), and Lamersdorf and Meyer (1993) attribute magnesium deficiencies in German forests, in part, to excessive nitrogen deposition.

The carbon to nitrogen (C:N) ratio of the forest floor can also be changed by nitrogen deposition over time. This change appears to occur when the ecosystem becomes nitrogen saturated (Gundersen et al., 1998a). Long-term changes in C:N status have been documented in Central Europe and indicate that nitrogen deposition has changed the forest floor. In Europe, low C:N ratios coincide with high deposition regions (Gundersen et al., 1998a). A strong decrease in forest floor root biomass has been observed with increased nitrogen availability. Roots and the

associated mycorrhizae appear to be an important factor in the accumulation of organic matter in the forest floor at nitrogen limited sites. If root growth and mycorrhizal formation are impaired by nitrogen deposition, the stability of the forest floor may be affected by stimulating turnover and decreasing the root litter input to the forest floor and thus decrease the nitrogen that can be stored in the forest floor pool (Gundersen et al., 1998b). Nitrogen-limited forests have a high capacity for deposited nitrogen to be retained by the plants and microorganisms competing for available nitrogen (Gundersen et al., 1998b). Nitrate leaching has been correlated significantly with nitrate status but not with nitrate depositions. Forest floor C:N ratio has been used as a rough indicator of ecosystem nitrogen status in mature coniferous forests and the risk of nitrate leaching; analyses of European databases indicated an empirical relationship between forest floor C:N ratio and nitrate leaching (Gundersen et al., 1998a). Nitrate leaching was observed when the deposition received was more than 10 kg N/ha. All of the data sets supported a threshold at which nitrate leaching seems to increase at a C:N ratio of 25. Therefore, to predict the rate of changes in nitrate leaching, it is necessary to be able to predict the rate of changes in the forest floor C:N ratio. Decreased foliar and soil nitrogen and soil C:N ratios, as well as changes in nitrogen mineralization rates, have been observed when comparing responses to nitrogen deposition in forest stands east and west of the Continental Divide in the Colorado Front Range (Baron et al., 2000; Rueth and Baron, 2002). Understanding the variability in forest ecosystem response to nitrogen input is essential in assessing pollution risks (Gundersen et al., 1998a).

The plant root is an important region of nutrient dynamics. The rhizosphere includes the soil that surrounds and is influenced by plant roots (Wall and Moore, 1999). The mutualistic relationship between plant roots, fungi, and microbes is critical for the growth of the organisms involved. The plant provides shelter and carbon; whereas the symbiont provides access to a limiting nutrients such as nitrogen and phosphorus. As indicated above, changes in soil nitrogen influence the mycorrhizal-plant relationship. Mycorrhizal fungal diversity is associated with above-ground plant biodiversity, ecosystem variability, and productivity (Wall and Moore, 1999). Aber et al. (1998) showed a close relationship between mycorrhizal fungi and the conversion of dissolved inorganic nitrogen to soil nitrogen. During nitrogen saturation, soil microbial communities change from being fungal, and probably being dominated by mycorrhizae, to being dominated by bacteria. The loss of mycorrhizal function has been hypothesized as the key

process leading to increased nitrification and nitrate mobility. Increased nitrate mobility leads to increased cation leaching and soil acidification (Aber et al., 1998).

The interrelationship of above- and below-ground flora is illustrated by the natural invasion of heathlands by oaks (*Quercus robur*). Soils are dynamic entities, the features of which can change like the rest of the ecosystem with age and management. The soil-forming factors under the heath have been vegetation typed during the last 2000 years; whereas the invasion by oaks has been taking place for only a few decades. Clearly changes in the ground floor and soil morphology takes place when trees colonize heath (Nielsen et al., 1999). The distribution of roots also changed under the three different vegetation types. Under both heather and the Sitka spruce plantation, the majority of roots are confined to the uppermost horizons; whereas under oak, the roots are distributed more homogeneously. There was also a change in the C:N ratio when heather was replaced by oaks. Also, the spontaneous succession of the heath by oaks changed the biological nutrient cycle into a deeper vertical cycle when compared to the heath where the cycle is confined to the upper soil horizons. Soils similar to those described in this study (Jutland, Denmark) with mainly an organic buffer system seem to respond quickly to changes in vegetation (Nielsen et al., 1999).

The affects of changes in root to shoot relationships in plants were observed in studies of the coastal sage scrub community in southern California which is composed of the drought-deciduous shrubs *Artemisia californica*, *Encelia farinosa*, and *Eriogonum fasciculatum*. The coastal sage scrub in California has been declining in land area and in shrub density over the past 60 years and is being replaced in many areas by Mediterranean annual grasses (Allen et al., 1998; Padgett et al., 1999; Padgett and Allen, 1999). Nitrogen deposition was considered as a possible cause. Up to 45 kg/ha/yr are deposited in the Los Angeles Air Basin (Bytnerowicz and Fenn, 1996). Tracts of land set aside as reserves, which in many cases in southern California are surrounded by urbanization, receive large amounts of nitrogenous compounds from polluted air. The coastal sage scrub is of particular interest because some 200 sensitive plant species and several federally listed animal species are found in the area (Allen et al., 1998). Because changes in plant community structure often can be related to increases in the availability of a limiting soil nutrient or other resource, experiments were conducted to determine whether increased nitrogen availability was associated with the significant loss in native shrub cover. Studies indicated that the three native perennial shrubs (*Artemisia californica*, *Eriogonum fasciculatum*, and *Encelia*

farinosa tended to be more nitrophilous than the two exotic annual grasses (*Bromus rubens*, *Avena fatua*) and the weedy pod mustard (*Brassica geniculata*). These results contrast with most models dealing with the adaptation of perennial species to stressful environments (Padgett and Allen, 1999). If nitrogen were the only variable between the invasive annuals and native shrubs, neither shrubs nor grasses have a particular advantage. However, additional studies indicated that the decline in the coastal sage scrub was not associated with its inability to compete with the grasses, but rather with changes in the arbuscular mycorrhizal community in the soil (Edgerton-Warburton and Allen, 2000). Nitrogen enrichment of the soils induced a shift in the arbuscular mycorrhizal community composition. Larger-spored fungal species (*Scutellospora* and *Gigaspora*), due to a failure to sporulate, decreased in number with a concomitant proliferation of small-spored species of *Glomus aggregatum*, G. *leptotichum*, and G. *geosporum*, indicating a strong selective pressure for the smaller spores species of fungi (Edgerton-Warburton and Allen, 2000). These results demonstrate that nitrogen enrichment of the soil significantly alters the arbuscular mycorrhizal species composition and richness and markedly decreases the overall diversity of the arbuscular mycorrhizal community. The decline in coastal sage scrub species can, therefore, directly be linked to the decline of the arbuscular mycorrhizal community (Edgerton-Warburton and Allen, 2000).

In addition to excess nitrogen deposition effects on terrestrial ecosystems of the types noted above (e.g., dominant species shifts and other biodiversity impacts), direct atmospheric nitrogen deposition and increased nitrogen inputs via runoff into streams, rivers, lakes, and oceans can have notable impacts on aquatic ecosystems as well. One illustrative example is recently reported research (Paerl et al., 2001) characterizing impacts of nitrogen deposition on the Pamlico Sound, NC, estuarine complex, which serves as a key fisheries nursery supporting an estimated 80% of commercial and recreational finfish and shellfish catches in the southeastern U.S. Atlantic coastal region. Such direct atmospheric nitrogen deposition onto waterways feeding into the Pamlico Sound or onto the sound itself and indirect nitrogen inputs via runoff from upstream watersheds contribute to conditions of severe water oxygen depletion; formation of algae blooms in portions of the Pamlico Sound estuarine complex; altered fish distributions, catches, and physiological states; and the incidence of disease. Under extreme conditions of especially high rainfall rate events (e.g., hurricanes) affecting watershed areas feeding into the sound, the effects of nitrogen runoff (in combination with excess loadings of metals or other

nutrients) can be massive—e.g., creation of the widespread “dead-zone” affecting large areas of the Pamlico Sound for many months after hurricane Fran in 1996 and hurricanes Dennis, Floyd, and Irene in 1999 impacted eastern North Carolina.

Nitrogen saturation of a high elevation watershed in the southern Appalachian Mountains was observed to affect streamwater chemistry. The Great Smoky Mountains in the southeastern United States receive high total atmospheric deposition of sulfur and nitrogen (2,200 Eq/ha/yr of total sulfur and approximately 1,990 Eq/ha/yr of total nitrogen). A major portion of the atmospheric loading is from dry and cloud deposition. Extensive surveys conducted in October 1993 and March 1994 indicated that stream pH values were near or below pH 5.5 and that the Acid Neutralizing Capacity (ANC) was below 50 μeq/L at high elevations. Analysis of streamwater indicated that nitrate was the dominant anion (Flum and Nodvin, 1995; Nodvin et al., 1995). The study was expanded to the watershed scale with monitoring of precipitation, thoughfall, stream hydrology, and stream chemistry. Nitrogen saturation of the watershed resulted in extremely high exports of nitrate and promoted both chronic and episodic stream acidification in which the nitrate was the dominant ion. Significant exports of base cation was also observed. Nitrification of the watershed soils resulted in elevations of soil solution aluminum concentrations to levels known to inhibit calcium uptake in red spruce (Nodvin et al., 1995).

In the Northeast, nitrogen is the element most responsible for eutrophication in coastal waters of the region (Jaworski et al., 1997). There has been a 3 to 8-fold increase in nitrogen flux from 10 watersheds in the Northeastern United States since the early 1900s. These increases are associated with nitrogen oxide emissions from combustion which have increased 5-fold. Riverine nitrogen fluxes have been correlated with atmospheric deposition onto their landscapes and also with nitrogen oxides emissions into their airsheds. Data from 10 benchmark watersheds with good historical records, indicate that ca. 36-80% of the riverine total nitrogen export, with an average of 64%, was derived directly or indirectly from nitrogen oxide emissions (Jaworski et al., 1997).

Excessive nitrogen loss is a symptom of terrestrial ecosystem dysfunction and results in the degradation of water quality and potentially deleterious effects on terrestrial and aquatic ecosystems (Fenn and Poth, 1999). Data from a number of hydrologic, edaphic, and plant indicators indicate that the mixed conifer forests and chaparral systems directly exposed to air

pollution from greater Los Angeles are nitrogen saturated. Preliminary data suggests that symptoms of nitrogen saturation are evident in mixed conifer or chaparral sites receiving atmospheric deposition of 20 to 25 kg/N/ha/y (Fenn et al, 1996). Available data clearly indicate that ecosystems with a Mediterranean climate have a limited capacity to retain nitrogen within the terrestrial system (Fenn and Poth, 1999). A 3-year study of streamwater NO_3^- concentrations along nitrogen deposition gradients in the San Bernardino Mountains in southern California evaluated streamwater quality and whether the streamwater concentrations covaried with nitrogen deposition across pollution gradients in the San Bernardino Mountains. Streamwater NO_3^- concentrations at Devil Canyon in the San Gabriel Mountains northeast of Los Angeles are the highest reported in North America for forested watersheds (Fenn and Poth, 1999). Five of the six streams monitored maintained elevated NO_3^- throughout the year. Peak nitrate concentrations ranged from 40 to 350 μmol/L. In the San Gorgonio Wilderness, an area of low to moderate deposition where 12 streams were sampled, only the five that had the greatest air pollution exposure had high NO_3^- concentrations. The results of the study suggested a strong association between levels of NO_3^- export in streamwater and the severity of chronic nitrogen deposition to the terrestrial watersheds. However, nitrogen processing within terrestrial and aquatic systems, even in areas with high nitrogen deposition, determine streamwater NO_3^- concentrations (Fenn and Poth, 1999). The Fernow Experimental Forest in West Virginia, the Great Smoky Mountains National park in Tennessee, and watersheds in southwestern Pennsylvania are the only undisturbed forested sites in North America known to have streamwater NO_3^- concentrations within the range of values found at Devil Canyon (Fenn and Poth, 1999).

Effects of Sulfur Deposition. Sulfur is an essential plant nutrient and, as such, is a major component of plant proteins. The most important source of sulfur is sulfate taken up from the soil by plant roots even though plants can utilize atmospheric SO_2 (Marschner, 1995). The availability of organically bound sulfur in soils depends largely on microbial decomposition, a relatively slow process. The major factor controlling the movement of sulfur from the soil into vegetation is the rate of release from the organic to the inorganic compartment (May et al., 1972; U.S. Environmental Protection Agency, 1982; Marschner, 1995). Sulfur plays a critical role in agriculture as an essential component of the balanced fertilizers needed to grow and increase

worldwide food production (Ceccotti and Messick, 1997). Atmospheric deposition is an important component of the sulfur cycle. This is true not only in polluted areas where atmospheric deposition is very high, but also in areas of low sulfur input. Additions of sulfur into the soil in the form of SO_4^{-2} could alter the important organic-sulfur/organic-nitrogen relationship involved in protein formation in plants. The biochemical relationship between sulfur and nitrogen in plant proteins and the regulatory coupling of sulfur and nitrogen metabolism indicate that neither element can be assessed adequately without reference to the other. Sulfur deficiency reduces nitrate reductase and, to a similar extent, also glutamine synthetase activity. Nitrogen uptake in forests, therefore, may be loosely regulated by sulfur availability, but sulfate additions in excess of needs do not necessarily lead to injury (Turner and Lambert, 1980; Hogan et al., 1998).

Only two decades ago, there was little information comparing sulfur cycling in forests with other nutrients, especially nitrogen. With the discovery of deficiencies in some unpolluted regions (Kelly and Lambert, 1972; Humphreys et al., 1975; Turner et al., 1977; Schnug, 1997) and excesses associated with acidic deposition in other regions of the world (Meiwes and Khanna, 1981; Shriner and Henderson, 1978; Johnson et al., 1982a,b), interest in sulfur nutrition and cycling in forests has heightened. General reviews of sulfur cycling in forests have been written by Turner and Lambert (1980), Johnson (1984), Mitchell et al. (1992a,b), and Hogan et al. (1998). The salient elements of the sulfur cycle as it may be affected by changing atmospheric deposition are summarized by Johnson and Mitchell (1998). Sulfur has become the most important limiting factor in European agriculture because of the desulfurization of industrial emissions (Schnug, 1997).

Most studies dealing with the impacts of sulfur deposition on plant communities have been conducted in the vicinity of point sources and have investigated above-ground effects of SO_2 or acidifying effects of sulfate on soils (Krupa and Legge, 1998; Dreisinger and McGovern, 1970; Legge, 1980; Winner and Bewley, 1978a,b; Laurenroth and Michunas, 1985; U.S. Environmental Protection Agency, 1982). Krupa and Legge (1986), however, observed a pronounced increase in foliar sulfur concentrations in all age classes of needles of the hybrid pine lodgepole x jack pine (*Pinus contorta* x *P. banksiana*). This vegetation had been exposed to chronic low concentrations of sulfur dioxide (SO_2) and hydrogen sulfide (H_2S) for more than 20 years and, then, to fugitive sulfur aerosol. Observations under the microscope showed no sulfur deposits on

the needle surfaces and led to the conclusion that the sulfur in the needles was derived from the soil. The oxidation of elemental sulfur and the generation of protons is well known for the soils of Alberta, Canada. This process is mediated by bacteria of the *Thiobacillus* sp. As elemental sulfur gradually is converted to protonated SO_4, it can be leached downward and readily taken up by plant roots. The activity of *Thiobacillus* sp. is stimulated by elemental sulfur additions (Krupa and Legge, 1986).

Effects of Acidic Deposition on Forest Soils. Acidic deposition over the past quarter of a century has emerged as a critical environmental stress that affects forested landscapes and aquatic ecosystems in North America, Europe, and Asia (Driscoll et al., 2001). Acidic deposition can originate from transboundary air pollution and affect large geographic areas. It is composed of ions, gases, particles derived from gaseous emissions of sulfur dioxide (SO_2), nitrogen oxides (NO_x), ammonia (NH_3), and particulate emissions of acidifying and neutralizing compounds and is highly variable across space and time. It links air pollution to diverse terrestrial and aquatic ecosystems and alters the interactions of the hydrogen ion (H^+) and many elements (e.g., sulfur, nitrogen, calcium, magnesium, and aluminum). Acidic deposition contributes directly and indirectly to biological stress and the degradation of ecosystems and has played a major role in recent acidification of soil in some areas of Europe and, to a more limited extent, eastern North America (Driscoll et al., 2001).

Substantial and previously unsuspected changes in soils have been observed in polluted areas of eastern North America, the United Kingdom, Sweden, and Central Europe and in less polluted regions of Australia and western North American (reviewed by Johnson et al., 1999 and by Huntington, 2000). In some cases, trends were toward more acidic soils (e.g., Markewitz et al., 1998), and, in others, there were no consistent trends, with some soils showing increases and some showing decreases at different sampling times, and some showing no change (e.g., Johnson and Todd, 1998; Trettin et al., 1999; Yanai et al., 1999).

Significant changes in soil chemistry have occurred at many sites in the eastern United States during recent decades. Patterns of change in tree ring chemistry, principally at high elevations sites in the eastern United States, reflect the changing inputs of regional pollutants to forests. A temporal sequence of changes in uptake patterns, and possibly in tree growth, would be expected if significant base cation mobilization and depletion of base cations from eastern

forest soils has occurred. Temporal changes in the chemistry of tree rings of red spruce were examined as indicators of historical changes in the chemical environment of red spruce.

Analysis of changes in wood chemistry from samples across several sites indicated that there have been substantial departures from the expected linear decreases in calcium accumulation patterns in wood. A region-wide calcium increase above expected levels followed by decreasing changes in wood calcium suggest that calcium mobilization began possibly 30 to 40 years ago and has been followed by reduced accumulation rates in wood, presumably associated with decreasing calcium availability in soil (Bondietti and McLaughlin, 1992). The period of calcium mobilization coincides with a region-wide increase in growth rate of red spruce; whereas the period of decreasing levels of calcium in wood corresponds temporally with patterns of decreasing radial growth at high elevation sites throughout the region during the past 20 to 30 years. The decline in wood calcium suggests that calcium loss may have increased to the point at which base saturation of soils has been reduced. Increases in aluminum and iron typically occur as base cations are removed from the soils by tree uptake (Bondietti and McLaughlin, 1992). The changes are spatially and temporally consistent with changes in the emissions of SO_2 and NO_2 across the region and suggest that increased acidification of soils has occurred.

Studies by Shortle and Bondietti (1992) support the view that changes in soil chemistry in eastern North America forest sites occurred many decades ago, “before anybody was looking.” Sulfur and nitrogen emissions began increasing in eastern North America in the 1920s and continued to increase into the 1980s when sulfur began to decrease but nitrogen emissions did not (Garner et al., 1989). Shortle and Bondietti (1992) present evidence that, from the late 1940s into the 1960s, the mor humus (organic) layer of acid-sensitive forest sites in eastern North America underwent a significant change that resulted in the loss of exchangeable essential base cations and interrupted the critical base nutrient cycles between mature trees and the root-humus complex. The timing of the effect appears to have coincided with the period when the SO_x and NO_x emissions in eastern North America subject to long-range transport were increasing the most rapidly (see above; Shortle and Bondietti, 1992). Although forest ecosystems other than the high-elevation spruce-fir forests are not currently manifesting symptoms of injury directly attributable to acid deposition, less sensitive forests throughout the United States are experiencing gradual losses of base cation nutrients, which in many cases will reduce the quality

of forest nutrition over the long term (National Science and Technology Council, 1998). In some cases it may not even take decades because these forests already have been receiving sulfur and nitrogen deposition for many years. The current status of forest ecosystems in different U.S. geographic regions varies, as does their sensitivity to nitrogen and sulfur deposition. Variation in potential future forest responses or sensitivity are caused, in part, by differences in deposition of sulfur and nitrogen, ecosystem sensitivities to sulfur and nitrogen additions, and responses of soils to sulfur and nitrogen inputs (National Science and Technology Council, 1998).

Acidic deposition has played a major role in recent soil acidification in some areas of Europe and, to a more limited extent, eastern North America. Examples include the study by Hauhs (1989) at Lange Bramke, Germany, which indicated that leaching was of major importance in causing substantial reduction in soil-exchangeable base cations over a 10-year period (1974-1984). Soil acidification and its effects result from the deposition of nitrate (NO_3^-) and sulfate (SO_4^{-2}) and the associated hydrogen (H^+) ion. The effects of excessive nitrogen deposition on soil acidification and nutrient imbalances have been well established in Dutch forests (Van Breemen et al., 1982; Roelofs et al., 1985; Van Dijk and Roelofs, 1988). For example, Roelofs et al. (1987) proposed that NH_3 /NH_4^+ deposition leads to heathland changes via two modes: acidification of the soil and the loss of cations K^+, Ca^{+2}, and Mg^{+2}; and nitrogen enrichment that results in “abnormal” plant growth rates and altered competitive relationships. Nihlgård (1985) suggested that excessive nitrogen deposition may contribute to forest decline in other specific regions of Europe. Falkengren-Grerup (1987) noted that, during about 50 years, unexpectedly large increases in growth of beech (*Fagus sylvatica L.*) were associated with decreases in pH and exchangeable cations in some sites in southernmost Sweden.

Likens et al. (1996, 1998) suggested that soils are changing at the Hubbard Brook Watershed, NH, because of a combination of acidic deposition and reduced base cation deposition. They surmised, based on long-term trends in streamwater data, that large amounts of calcium and magnesium have been lost from the soil-exchange complex over a 30-year period from approximately 1960 to 1990. The authors speculate that the declines in base cations in soils may be the cause of recent slowdowns in forest growth at Hubbard Brook. In a follow-up study, however, Yanai et al. (1999) found no significant decline in calcium and magnesium concentrations in forest floors at Hubbard Brook over the period 1976 to 1997. They also found both gains and losses in forest floor calcium and magnesium between 1980 and 1990 in a

regional survey. Thus, they concluded that "forest floors in the region are not currently experiencing rapid losses of base cations, although losses may have preceded the onset of these three studies." The biogeochemistry of calcium at Hubbard Brook is discussed in detail by Likens et al. (1998).

Hydrogen ions entering a forest ecosystem first encounter the forest canopy, where they are often exchanged for base cations that then appear in throughfall (Figure 4-10 depicts a model of H^+ sources and sinks). Base cations leached from the foliage must be replaced through uptake from the soil, or foliage cations will be reduced by the amounts leached. In the former case, the acidification effect is transferred to the soil where H^+ is exchanged for a base cation at the root-soil interface. Uptake of base cations or NH_4^+ by vegetation or soil microorganisms causes the release of H^+ in order to maintain charge balance; uptake of nutrients in anionic form (NO_3^-, SO_4^{-2}, PO_4^{-3}) causes the release of OH^- in order to maintain charge balance. Thus, the net acidifying effect of uptake is the difference between cation and anion uptake. The form of ions taken up is known for all nutrients but nitrogen because either NH_4^+ or NO_3^- can be utilized. In that nitrogen is a nutrient taken up in greatest quantities, the uncertainty in the ionic form of nitrogen utilized creates great uncertainty in the overall H^+ budget for soils (Johnson 1992).

The cycles of base cations differ from those of N, P, and S in several respects. The fact that calcium, potassium, and magnesium exist primarily as cations in solution, whereas N, P, and S exist primarily as anions, has major implications for the cycling of the nutrients and the effects of acid deposition on these cycles. The most commonly accepted model of base cation cycling in soils is one in which base cations are released by weathering of primary minerals to cation exchange sites where they are available for either plant uptake or leaching (Figure 4-10). The introduction of H^+ by atmospheric deposition or by internal processes will affect the fluxes of Ca, K, and Mg via cation exchange or weathering processes. Therefore, soil leaching is often of major importance in cation cycles, and many forest ecosystems show a net loss of base cations (Johnson, 1992).

Two basic types of soil change are involved: (1) a short-term intensity type change resulting from the concentrations of chemicals in soil water and (2) a long-term capacity change based on the total content of bases, aluminum, and iron stored in the soil (Reuss and Johnson, 1986; Van Breemen et al., 1983). Changes in intensity factors can have a rapid affect on the chemistry of soil solutions. Increases in the amounts of sulfur and nitrogen in acidic deposition



Figure 4-10. Schematic of sources and sinks of hydrogen ions in a forest (from Taylor et al., 1994).

can cause immediate increases in acidity and mobilization of aluminum in soil solutions. Increased aluminum concentrations and an increase in the Ca/Al ratio in soil solution have been linked to a significant reduction in the availability of essential base cations to plants, an increase in plant respiration, and increased biochemical stress (National Science and Technology Council, 1998).

Rapid changes in intensity resulting from the addition of increased amounts of nitrogen or sulfur in acidic deposition can have a rapid effect on the chemistry of soil solutions by increasing the acidity and mobilizing aluminum. Increased concentrations of aluminum and an increase in the ratio of calcium to aluminum in soil solution have been linked to a significantly reduced availability of essential cations to plants.

Capacity changes are the result of many factors acting over long time periods. The content of base cations (calcium, magnesium, sodium, and potassium) in soils results from additions from the atmospheric deposition, decomposition of vegetation, and geologic weathering. Loss of base cations may occur through plant uptake and leaching. Increased leaching of base cations may result in nutrient deficiencies in soils as has been happening in some sensitive forest ecosystems (National Science and Technology Council, 1998).

Aluminum toxicity is a possibility in acidified soils. Atmospheric deposition (or any other source of mineral anions) can increase the concentration of Al, especially Al^{3+}, in soil solution without causing significant soil acidification (Johnson and Taylor, 1989). Aluminum can be brought into soil solution in two ways: (1) by acidification of the soil and (2) by an increase in the total anion and cation concentration of the soil solution. The introduction of mobile, mineral acid anions to an acid soil will cause increases in the concentration of aluminum in the soil solution, but extremely acid soils in the absence of mineral acid anions will not produce a solution high in aluminum. An excellent review of the relationships among the most widely used cation-exchange equations and their implications for the mobilization of aluminum into soil solution is provided by Reuss (1983).

A major concern has been that soil acidity would lead to nutrient deficiency. Calcium is essential for root development and the formation of wood, and it plays a major role in cell membrane integrity and cell wall structure. Aluminum concentrations in the soil can influence forest tree growth in regions where acidic deposition and natural acidifying processes increase soil acidity. Acidic deposition mobilizes calcium and magnesium, which are essential for root development and stem growth. Mobilized aluminum can also bind to fine root tips of red spruce, further limiting calcium and magnesium uptake (Shortle and Smith, 1988; Shortle et al., 1997).

There is abundant evidence that aluminum is toxic to plants. Upon entering tree roots, it accumulates in root tissues (Thornton et al., 1987; Vogt et al., 1987a, b). Reductions in calcium uptake have been associated with increases in aluminum uptake (Clarkson and Sanderson, 1971). A number of studies suggest that the toxic effect of aluminum on forest trees could be caused by Ca^{+2} deficiency (Shortle and Smith, 1988; Smith, 1990a). Mature trees have a high calcium requirement relative to agriculture crops (Rennie, 1955). Shortle and Smith (1988) attributed the decline of red spruce in eight stands across northern New England from Vermont to Maine to an imbalance of Al^{3+} and Ca^{+2} in fine root development.

To be taken up from the soil by roots, calcium must be dissolved in soil water (Lawrence and Huntington, 1999). Aluminum in soil solution reduces calcium uptake by competing for binding sites in the cortex of fine roots. Tree species may be adversely affected if high aluminum to nutrient ratios create a nutrient deficiency by limiting uptake of calcium and magnesium (Shortle and Smith, 1988; Garner, 1994). Acid deposition, by lowering the pH of aluminum-rich soil, can increase aluminum concentrations in soil water through dissolution and ion exchange processes. Aluminum is more readily taken up than is calcium because of its greater affinity for negatively charged surfaces. When present in the forest floor, aluminum tends to displace adsorbed calcium and causes it to be more readily leached. The continued buildup of aluminum in the forest floor layer, where nutrient uptake is greatest, can lower efficiency of calcium uptake when the ratio of calcium to aluminum in soil water is less than one (Lawrence and Huntington, 1999). Reduction in calcium uptake suppresses cambial growth and reduces the rate of wood (annual ring) formation, decreases the amount of functional sapwood and live crown, and predisposes trees to disease and injury from stress agents when the functional sapwood becomes less that 25% of cross-sectional stem area (Smith, 1990a). A 1968 Swedish report to the United Nations postulated a decrease in forest growth of ca. 1.5% per year when the ratio of calcium to aluminum in soil water is less than one (Lawrence and Huntington, 1999). The concern that acidification and nutrient deficiency may result in forest decline remains today.

Acidic deposition has been firmly implicated as a causal factor in northeastern high-elevation decline of red spruce (DeHayes et al., 1999). The frequency of freezing injury of red spruce has increased over the past 40 years, a period that coincides with increase emissions of sulfur and nitrogen oxides and acidic deposition (DeHayes et al., 1999). Studies indicate that there is a significant positive association between cold tolerance and foliar calcium in trees exhibiting deficiency in foliar calcium. Most of the calcium in conifer needles is insoluble calcium oxalate and pectate crystals, which are of little physiological importance. It is the labile calcium ions in equilibrium within the plasma membrane that are of major physiological importance (DeHayes et al., 1999). The membrane-associated pool of calcium (mCa), although a relatively small fraction of total foliar ion pools, strongly influences the response of cells to changing environmental conditions. The plant plasma membrane plays a critical role in mediating cold acclimation and low-temperature injury. Leaching of calcium associated with acidic deposition is considered to be the result of cation exchange due to exposure to the H^+ ion.

The studies of DeHayes et al. (1999) demonstrate that the direct deposition of acidic deposition on needles represents a unique environmental stress, in that it preferentially removes mCa which is not readily replaced in autumn. They propose that direct deposition on red spruce foliage preferentially displaces calcium ions specifically associated with plasma membranes of mesophyll cells resulting in the reduction of mCa and the destabilizing of plasma membranes and depletion of messenger calcium. Further, DeHayes et al.(1999) state that their studies raise the strong possibility that acid rain alteration of the mCa and membrane integrity is not unique to red spruce but has been demonstrated in many other northern temperate forest tree species including yellow birch (*Betula alleghaniensis*), white spruce (*Picea glaucus*), red maple (*Acer rubrum*) eastern white pine (Pinus strobus), and sugar maple (*Acer saccharum*). Assessments of mCa, membrane integrity, and the effects of other secondary stresses have not yet been made for these species.

Seasonal and episodic acidification of surface waters have been observed in the eastern United States, Canada and Europe (Hyer et al., 1995). In the Northeast, the Shenandoah National Park in Virginia, and the Great Smoky Mountains, episodic acidification has been associated with the nitrate ion (Driscoll et al., 2001; Hyer et al., 1995 ; Eshleman et al., 1995). The short-term acid episodes occur during spring snowmelts and large precipitation events (Driscoll et al., 2001). Episodic acidification of surface waters has usually been considered to be a transient loss of acid neutralizing capacity associated with snowmelt/rainfall runoff and, as such, represents short-term (hours to weeks) effects considered to be distinguishable from chronic long-term (years to centuries) changes in acidity. Studies of both episodic and chronic acidification of surface waters indicate that acidification can have long-term adverse effects on fish populations, declines of species richness, abundance of zooplankton, and macroinvertebrates (Driscoll et al., 2001; Eshleman et al., 1995). Nitrogen saturation of soils and the slow release of nitrates has inhibited the recovery of acid sensitive systems (Driscoll et al., 2001). The acidification of aquatic ecosystems and the effects on aquatic biota has been discussed in greater detail in the Air Quality Criteria for Nitrogen Oxides (U.S. Environmental Protection Agency, 1993).

Air pollution is not the sole cause of soil change. High rates of acidification are occurring in less polluted regions of the western United States and Australia because of internal soil processes, such as tree uptake of nitrate and nitrification associated with excessive nitrogen fixation (Johnson et al., 1991b). Many studies have shown that acidic deposition is not a

necessary condition for the presence of extremely acid soils, as evidenced by their presence in unpolluted, even pristine, forests of the northwestern United States and Alaska (Johnson et al., 1991b). Soil can become acidic when H^+ ions attached to NH_4^+ or HNO_3 remain in the soil after nitrogen is taken up by plants. For example, Johnson et al. (1982b) found significant reductions in exchangeable K^+ over a period of only 14 years in a relatively unpolluted Douglas fir Integrated Forest Study (IFS) site in the Washington Cascades. The effects of acid deposition at this site were negligible relative to the effects of natural leaching (primarily carbonic acid) and nitrogen tree uptake (Cole and Johnson, 1977). Even in polluted regions, numerous studies have shown the importance of tree uptake of NH_4^+ and NO_3^- in soil acidification. Binkley et al. (1989) attributed the marked acidification (pH decline of 0.3 to 0.8 units and base saturation declines of 30 to 80%) of abandoned agricultural soil in South Carolina over a 20-year period to NH_4^+ and NO_3^- uptake by a loblolly pine plantation.

An interesting example of uptake effects on soil acidification is that of Al uptake and cycling (Johnson et al., 1991b). Aluminum accumulation in the leaves of coachwood (*Ceratopetalum apetalum*) in Australia has been found to have a major effect on the distribution and cycling of base cations (Turner and Kelly, 1981). The presence of C. *apetalum* as a secondary tree layer beneath brush box (*Lophostemon confertus*) was found to lead to increased soil exchangeable Al^{3+} and decreased soil exchangeable Ca^{2+} (Turner and Kelly, 1981). The constant addition of aluminum-rich litter fall obviously has had a substantial effect on soil acidification, even if base cation uptake is not involved directly.

Given the potential importance of particulate deposition for base cation status of forest ecosystems, the findings of Driscoll et al. (1989, 2001) and Hedin et al. (1994) are especially relevant. Driscoll et al. (1989, 2001) noted a decline in both SO_4^{-2} and base cations in both atmospheric deposition and stream water over the past two decades at Hubbard Brook Watershed, NH. The decline in SO_4^{-2} deposition was attributed to a decline in emissions, and the decline in stream water SO_4^{-2} was attributed to the decline in sulfur deposition.

Hedin et al. (1994) reported a steep decline in atmospheric base cation concentrations in both Europe and North America over the past 10 to 20 years. The reductions in SO_2 emissions in Europe and North America in recent years have not been accompanied by equivalent declines in net acidity related to sulfate in precipitation. These current declines in sulfur deposition have, in varying degrees, been offset by declines in base cations and may be contributing "to the increased

sensitivity of poorly buffered systems." Analysis of the data from the IFS supports the authors' contention that atmospheric base cation inputs may seriously affect ecosystem processes. Johnson et al. (1994b) analyzed base cation cycles at the Whiteface Mountain IFS site in detail and concluded that Ca losses from the forest floor were much greater than historical losses, based on historical changes in forest floor Ca observed in an earlier study. Further, the authors suggest that the difference between historical and current net loss rates of forest floor Ca may be caused by sharply reduced atmospheric inputs of calcium after about 1970 and may be exacerbated by sulfate leaching (Johnson et al., 1994b).

The calcium/aluminum molar ratio has been suggested as a valuable ecological indicator of an approximate threshold beyond which the risk of forest injury from Al stress and nutrient imbalances increases (Cronan and Grigal, 1995). The Ca/Al ratio also can be used as an indicator to assess forest ecosystem changes over time in response to acidic deposition, forest harvesting, or other process that contribute to acid soil infertility. This ratio, however, may not be a reliable indicator of stress in areas with both high atmospheric deposition of ammonium and magnesium deficiency via antagonism involving ammonium rather than aluminum and in areas with soil solutions with calcium concentrations greater than 500 micromoles per liter (National Science and Technology Council, 1998). Cronan and Grigal (1995), based on a review of the literature, have made the following estimates for determining the adverse impact of acidic deposition on tree growth or nutrition:

- forests have a 50% risk of adverse impacts if the Ca/Al ration is 1.0,
- the risk is 75% if the ratio is 0.5, and
- the risk approaches 100% if the ratio is 0.2.

The Ca/Al ratio of soil solution provides only an index of the potential for Al stress. Cronan and Grigal (1995) state that the overall uncertainty of the Ca/Al ratio associated with a given probability ratio is considered to be approximately ±50%. Determination of thresholds for potential forest impacts requires the use of the four successive measurement endpoints in the soil, soil solution, and plant tissue listed below.

(1) Soil base saturation less than 15% of effective cation exchange capacity,
(2) Soil solution Ca/Al molar ratio less than 1.0 for 50% risk,
(3) Fine roots tissue Ca/Al molar ratio less than 0.2 for 50% risk, and
(4) Foliar tissue Ca/Al molar ratio less than 12.5 for 50% risk.

The application of the Ca/Al ratio indicator for assessment and monitoring of forest health risks has been recommended for sites or in geographic regions where the soil base saturation <15%.

Critical Loads. In Europe, the critical load concept generally has been accepted as the basis for abatement strategies to reduce or prevent injury to the functioning and vitality of forest ecosystems caused by long-range transboundary acidic deposition (Lokke, et al., 1996). The critical load has been defined as a "quantitative estimate of an exposure to one or more pollutants below which significant harmful effects on specified sensitive elements of the environment do not occur according to present knowledge" (Lokke et al., 1996). A biological indicator, a chemical criterion, and a critical value are the elements used in the critical load concept. The biological indicator is the organism used to indicate the status of the receptor ecosystem; the chemical criterion is the parameter that results in harm to the biological indicator; and the critical value is the value of the chemical criterion below which no significant harmful response occurs to the biological indicator (Lokke et al., 1996). Trees, and sometimes other plants, are used as the biological indicators in the case of critical loads for forests. The critical load calculation using the current methodology, is essentially an acidity/alkalinity mass balance calculation. The chemical criterion must be expressible in terms of alkalinity. Initially, the Ca/Al ratio was used; but, recently, the (Ca+Mg+K)/Al ratio has been used (Lokke et al., 1996).

Ideally, changes in acidic deposition should result in changes in the status of the biological indicator used in the critical load calculation. However, the biological indicator is the integrated response to a number of different stresses. Furthermore, there are other organisms more sensitive to acid deposition than trees. At high concentrations, Al^{3+} is known to be toxic to plants, inhibiting root growth and, ultimately, plant growth and performance (Lokke et al., 1996; National Science and Technology Council, 1998). Sensitivity to Al varies considerably between species and within species because of changes in nutritional demands and physiological status that are related to age and climate. Experiments have shown that there are large variations in Al sensitivity, even among ecotypes.

Mycorrhizal fungi as possible biological indicators have been suggested by Lokke et al. (1996) because they are intimately associated with tree roots, depend on plant assimilates, and play an essential role in plant nutrient uptake influencing the ability of their host plants to tolerate different anthropogenically generated stresses. Mycorrhizas and fine roots are an extremely

dynamic component of below-ground ecosystems and can respond rapidly to stress. They have a relatively short life span, and their turnover appears to be strongly controlled by environmental factors. Changes in mycorrhizal species composition or the loss of dominant mycorrhizal species in areas where diversity is already low may lead to increased susceptibility of plant to stress (Lokke et al., 1996). Stress affects the total amount of carbon fixed by plants and modifies carbon allocation to biomass, symbionts, and secondary metabolites. The physiology of carbon allocation has also been suggested as an indicator of anthropogenic stress (Andersen and Rygiewicz, 1991). Because mycorrhizal fungi are dependent for their growth on the supply of assimilates from the host plants, stresses that shift the allocation of carbon reserves to the production of new leaves at the expense of supporting tissues will be reflected rapidly in decreased fine root and mycorrhizzal biomass (Winner and Atkinson, 1986). Decreased carbon allocation to roots also affects soil carbon and rhizosphere organisms. Soil dwelling animals are important for decomposition, soil aeration, and nutrient redistribution in the soil. They contribute to decomposition and nutrient availability mainly by increasing the accessibility of dead plant material to microorganisms. Earthworms decrease in abundance and in species number in acidified soils (Lokke et al., 1996).

Effects of Wet and Dry Deposition on Biogeochemical Cycling—The Integrated Forest Study. The Integrated Forest Study (IFS) (Johnson and Lindberg, 1992a) has provided the most extensive data set available on wet and dry deposition and deposition effects on the cycling of elements in forest ecosystems. The overall patterns of deposition and cycling have been summarized by Johnson and Lindberg (1992a), and the reader is referred to that reference for details. The following is a summary of particulate deposition, total deposition, and leaching in the IFS sites.

Particulate deposition in the IFS was separated at the 2-μm level; a decision was made to include total particulate deposition in this analysis and may include the deposition of particles larger than 10 μm.

Particulate deposition contributes considerably to the total impact of base cations to most of the IFS sites. On average, particulate deposition contributes 47% to total calcium deposition (range: 4 to 88%), 49% of total potassium deposition (range: 7 to 77%), 41% to total magnesium deposition (range: 20 to 88%), 36% to total sodium deposition (range: 11 to 63%), and 43% to

total base cation deposition (range: 16 to 62%). Of the total particulate deposition, the vast majority (>90%) is >2 μm.

Figures 4-11 through 4-14 summarize the deposition and leaching of calcium, magnesium, potassium, and total base cations for the IFS sites. As noted in the original synthesis (Johnson and Lindberg, 1992a), some sites show net annual gains of base cations (i.e., total deposition > leaching), some show losses (total deposition < leaching), and some are approximately in balance. Not all cations follow the same pattern at each site. For example, calcium shows net accumulation at the Coweeta, TN, Durham (Duke), NC, and Florida sites (Figure 4-11). Potassium shows accumulation at the Duke, Florida, Douglas-fir; red alder, Thompson, WA, Huntington Forest, NY, and Whiteface Mountain, NY, sites (Figure 4-13). Magnesium accumulated only at the Florida sites (Figure 4-12); only at the Florida site, is there a clear net accumulation of total base cations (Figure 4-14).

As noted previously, the factors affecting net calcium accumulation or loss include the soil-exchangeable cation composition; base cation deposition rate; the total leaching pressure because of atmospheric sulfur and nitrogen inputs, as well as natural (carbonic and organic) acids; and biological demand (especially for potassium). At the Florida site, which has a very cation-poor, sandy soil (derived from marine sand), the combination of all these factors leads to net base cation accumulation from atmospheric deposition (Johnson and Lindberg, 1992a). The site showing the greatest net base cation losses, the red alder stand in Washington state, is one that is under extreme leaching pressure by nitrate produced because of excessive fixation by that species (Van Miegroet and Cole, 1984). In the red spruce site in the Smokies, the combined effects of SO_4^{-2} and NO_3^- leaching are even greater than in the red alder site (Figure 4-15), but a considerable proportion of the cations leached from this extremely acid soil consist of H^+ and Al^{+3} rather than of base cations (Johnson and Lindberg, 1992a). Thus, the red spruce site in the Smokies is approximately in balance with respect to calcium and total base cations, despite the very high leaching pressure at this site (Figures 4-11 and 4-14).

The relative importance of particulate base cation deposition varies widely with site and cation and is not always related to the total deposition rate. The proportion of calcium deposition in particulate form ranges from a low of 4% at the Whiteface Mountain site to a high of 88% at the Maine site (Figure 4-11). The proportion of potassium deposition as particles ranges from 7% at the Smokies site to 77% at the Coweeta site (Figure 4-13), and the proportion of total base



Figure 4-11. Calcium deposition in >2-μm particles, <2-μm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites. CP = *Pinus strobus*, Coweeta, TN; DL = *Pinus taeda*, Durham (Duke), NC; GS = *Pinus taeda*, B. F. Grant Forest, GA; LP = *Pinus taeda*, Oak Ridge, TN; FS = *Pinus eliottii*, Bradford Forest, FL; DF = *Psuedotsuga menziesii*, Thompson, WA; RA = *Alnus rubra*; Thompson WA; NS = *Picea abies*, Nordmoen, Norway; HF = northern hardwood, Huntington Forest, NY; MS = *Picea rubens*, Howland, ME; WF = *Picea rubens*, Whiteface Mountain, NY; and ST = *Picea rubens*, Clingman's Dome, NC.

cation deposition ranges from 16% at the Whiteface site to 62% at the Maine site (Figure 4-14). Overall, particulate deposition at the site in Maine accounted for the greatest proportion of calcium, magnesium, potassium, and base cation deposition (88, 88, 57, and 62%, respectively), even though total deposition was relatively low. At some sites, the relative importance of particulate deposition varies considerably by cation. At the Whiteface Mountain site, particulate deposition accounts for 4, 20, and 40% of calcium, magnesium, and potassium deposition, respectively. At the red spruce site in the Smokies, particulate deposition accounts for 46, 26%, 7% of calcium, magnesium, and potassium deposition, respectively.



Figure 4-12. Magnesium deposition in >2-μm particles, <2-μm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites. See Figure 4-11 for site abbreviations.

As indicated in the IFS synthesis, SO_4^{-2} and NO_3^- leaching often are dominated by atmospheric sulfur and nitrogen (Johnson and Lindberg, 1992a). The exceptions to this are in cases where natural nitrogen inputs are high (i.e.,the nitrogen-fixing red alder stand), as are NO_3^- leaching rates even though nitrogen deposition is low, and where soils adsorb much of the atmospherically deposited SO_4^{-2} thus reducing SO_4^{-2} leaching compared to atmospheric sulfur input.

Sulfate and NO_3^- leaching have a major effect on cation leaching in many of the IFS sites (Johnson and Lindberg, 1992a). Figure 4-15 shows the total cation leaching rates of the IFS sites and the degree to which cation leaching is balanced by SO_4^{-2} + NO_3^- deposition. The SO_4^{-2} and NO_3^- fluxes are subdivided further into that proportion potentially derived from particulate sulfur and nitrogen deposition (assuming no ecosystem retention, a maximum effect) and other sulfur and nitrogen sources (wet and gaseous deposition, internal production).



Figure 4-13. Potassium deposition in >2-μm particles, <2-μm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites. See Figure 4-11 for site abbreviations.

As noted in the IFS synthesis, total SO_4^{-2} and NO_3^- inputs account for a large proportion (28 to 88%) of total cation leaching in most sites. The exception is the Georgia loblolly pine site, where there were high rates of HCO_3^- and Cl^- leaching (Johnson and Lindberg, 1992a). The role of particulate sulfur and nitrogen deposition in this leaching is generally very small (<10%), however, even if it is assumed that there is no ecosystem sulfur or nitrogen retention.

It was noted previously in this chapter that the contribution of particles to total deposition of nitrogen and sulfur at the IFS sites is lower than that for base cations. On average, particulate deposition contributes 18% to total nitrogen deposition (range: 1 to 33%) and 17% to total sulfur deposition (range: 1 to 30%). Particulate deposition contributes only a small amount to total H^+ deposition (average = 1%; range: 0 to 2%). (It should be noted, however, that particulate H^+ deposition in the > 2 μm fraction was neglected.)



Figure 4-14. Base cation deposition in >2-μm particles, <2-μm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites. See Figure 4-11 for site abbreviation.

Based on the IFS data, it appears that the particulate deposition has a greater effect on base cation inputs to soils than on base cation losses associated with inputs of sulfur, nitrogen, and H^+. It cannot be determined what fraction of the mass of these particles are <10 μm, but only a very small fraction is <2 μm. These inputs of base cations have considerable significance, not only to the base cation status of these ecosystems, but also to the potential of incoming precipitation to acidify or alkalize the soils in these ecosystems. As noted above, the potential of precipitation to acidify or alkalize soils depends on the ratio of base cations to H^+ in deposition, rather than simply on the inputs of H^+ alone. In the case of calcium, the term "lime potential" has been applied to describe this ratio; the principle is the same with respect to magnesium and potassium. Sodium is a rather special case, in that it is a poorly absorbing cation and leaching tends to balance input over a relatively short term.

Net balances of base cations tell only part of the story as to potential effects on soils; these net losses or gains must be placed in the perspective of the soil pool size. One way to express



Figure 4-15. Total cation leaching (total height of bar) balanced by sulfate and nitrate estimated from particulate deposition (assuming no ecosystem retention, particulate sulfur and nitrogen) and by other sources (both deposition and internal) of sulfate and nitrate (other sulfur and nitrogen sources) and by other anions in the Integrated Forest Study sites. See Figure 4-11 for legend.

this perspective is to simply compare soil pool sizes with the net balances. This comparison is made for exchangeable pools and net balances for a 25-year period in Figures 4-16 to 4-18. It readily is seen that net leaching losses of cations pose no threat in terms of depleting soil-exchangeable Ca^{+2}, K^{+}, or Mg^{+2} within 25 years at the Coweeta, Duke, Georgia, Oak Ridge, or Douglas-fir sites. However, there is a potential for significant depletion at the red alder, Whiteface Mountain (magnesium), and Smokies red spruce sites.

The range of values for soil-exchangeable turnover is very large, reflecting variations in both the size of the exchangeable pool and the net balance of the system. Soils with the highest turnover rates are those most likely to experience changes in the shortest time interval, other things being equal. Thus, the Whiteface Mountains, Smokies, and Maine red spruce sites; the Thompson red alder site; and the Huntington Forest northern hardwood site appear to be most sensitive to change. The actual rates, directions, and magnitudes of changes that may occur in



Figure 4-16. Soil exchangeable Ca^{+2} pools and net annual export of Ca^{+2} (deposition minus leaching times 25 years) in the Integrated Forest Study sites. See Figure 4-11 for site abbreviations.



Figure 4-17. Soil exchangeable Mg^{+2} pools and net annual export of Mg^{+2} (deposition minus leaching times 25 years) in the Integrated Forest Study sites. See Figure 4-11 for site abbreviations.



Figure 4-18. Soil exchangeable K^{2+} pools and net annual export of K^{2+} (deposition minus leaching times 25 years) in the Integrated Forest Study sites. See Figure 4-11 for site abbreviations.

these soils (if any) will depend on weathering inputs and vegetation outputs, in addition to deposition and leaching. It is noteworthy that each of the sites listed above as sensitive has a large store of weatherable minerals, whereas many of the other soils, with larger exchangeable cation reserves, have a small store of weatherable minerals (e.g., Coweeta white pine, Duke loblolly pine, Georgia loblolly pine, and Oak Ridge loblolly pine) (Johnson and Lindberg, 1992a; April and Newton, 1992).

Base cation inputs are especially important to the Smokies red spruce site because of potential aluminum toxicity and calcium and magnesium deficiencies. Johnson et al. (1991a) found that soil solution aluminum concentrations occasionally reached levels found to inhibit calcium uptake and cause changes in root morphology in solution culture studies of red spruce (Raynal et al., 1990). In a follow-up study, Van Miegroet et al. (1993) found a slight but significant growth response to calcium and magnesium fertilizer in red spruce saplings near the Smokies red spruce site. Joslin et al. (1992) reviewed soil and solution characteristics of red spruce in the southern Appalachians, and it appears that the IFS site is rather typical.

Wesselink et al. (1995) reported on the complicated interactions among changing deposition and soils at this site (including repeated sampling of soil exchangeable base cation pools) from 1969 to 1991 and compared these results with those of a simulation model. They identified three basic stages of change in this ecosystem. During Stage 1, there was increased deposition of sulfur and constant deposition of base cations, causing increased base cation leaching and reduced base saturation in the soils. During Stage II, sulfur deposition is reduced, and soil solution sulfate and base cation leaching decline accordingly, but base saturation continues to decrease. During Stage III, two alternative scenarios are introduced: (a) sulfur deposition continues to decline, whereas base cation deposition says constant; or (b) both sulfur and base cation deposition decline. Under Stage III-a, sulfate and base cation leaching continue to decline, and base saturation begins to increase as base cations displace exchangeable aluminum and cause it to transfer to the gibbsite pool. Under Stage III-b, this recovery in base saturation is over-ridden by the reduction in base cation deposition.

The IFS project, for the first time, accurately quantifies atmospheric deposition inputs to nutrient cycles using state-of-the-art techniques to measure wet and dry deposition. The principal aim of the project was to determine the effects of atmospheric deposition on nutrient status of a variety of forest ecosystems and to determine if these effects are in any way related to current or potential forest decline. Acidic deposition is having a significant effect on nutrient cycling in most of the forest ecosystems studied in the IFS project. The exceptions were the relatively unpolluted Douglas fir, red alder, and Findley Lakes in Washington state. The nature of the effects, however, varies from one location to another (Johnson,1992). In all but the relatively unpolluted Washington sites, atmospheric deposition was having a significant, often overwhelming effect on cation leaching from the soils. In general, nutrient budget data from IFS and the literature suggest that the susceptibility of southeastern sites to base cation depletion from soils and the development of cation deficiencies by that mechanism appears to be greater than in northern sites (Johnson, 1992).

Atmospheric deposition may have affected significantly the nutrient status of some IFS sites through the mobilization of Al. Soil solution Al levels in the Smokies sites approach and sometimes exceed levels noted to impede cation uptake in solution culture studies. It is therefore possible that the rates of base cation uptake and cycling in these sites have been reduced because of soil solution Al. To the extent that atmospheric deposition has contributed to these elevated

soil solution Al levels, it has likely caused a reduction in base cation uptake and cycling rates at these sites. Nitrate and sulfate are the dominant anions in the Smokies sites, and nitrate pulses are the major cause of Al pulses in soil solution (Johnson, 1992). The connection between Al mobilization and forest decline is not clear. The decline in red spruce certainly has been more severe in the Northeast than in the Southeast, yet all evidence indicates that Al mobilization is most pronounced in the southern Appalachians. However, at the Whiteface Mountain site selected for study because it was in a state of decline, soil solution levels were lower than in the Smokies, which are not in a visibly obvious state of decline (there was no dieback other than the fir killed by the balsam wooly adelgid, no needle yellowing). Thus, Al mobilization constitutes a situation worthy of further study (Johnson, 1992).

The simple calculations shown above give some idea of the importance of particulate deposition in these forest ecosystems, but they cannot account for the numerous potential feedbacks between vegetation and soils nor for the dynamics through time that can influence the ultimate response. One way to examine some of these interactions and dynamics is to use simulation modeling. The nutrient cycling model (NuCM) has been developed specifically for this purpose and has been used to explore the effects of atmospheric deposition, fertilization, and harvesting on some of the IFS sites (Johnson et al., 1993). The NuCM model is a stand-level model that incorporates all major nutrient cycling processes (uptake, translocation, leaching, weathering, organic matter decay, and accumulation).

Johnson et al. (1999) used the NuCM model to simulate the effects of reduced S, N, and base cation (C_B) deposition on nutrient pools, fluxes, soil, and soil solution chemistry in two contrasting southern Appalachian forest ecosystems: the red spruce and Coweeta hardwood sites from the IFS project. The scenarios chosen for these simulations included “no change;” 50% N and S deposition; 50% C_B deposition; and 50% N, S, and C_B deposition (50% N, S, C_B). The NuCM simulations suggested that, for the extremely acid red spruce site, S and N deposition is the major factor affecting soil solution Al concentrations and C_B deposition is the major factor affecting soil solution C_B concentrations. The effects of S and N deposition were largely through changes in soil solution SO_4^{-2} and NO_3^- and, consequently, mineral acid anion (MAA) concentrations rather than through changes in soils. This is illustrated in Figures 4-19a,b and 4-20a,b, which show simulated soil solution mineral acid anions, base cations, Al, and soil base saturation in B horizon from in the red spruce site. The 50% S and N scenario caused reductions



Figure 4-19a. Simulated soil solution mineral acid anions and base cations in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition. Redrawn from Johnson et al. (1999).

DRAFT-DO NOT QUOTE OR CITE



Figure 4-19b. Simulated soil solution mineral acid anions and base cations in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition. Redrawn from Johnson et al. (1999).

DRAFT-DO NOT QUOTE OR CITE



Figure 4-20a. Simulated soil solution A1 and soil base saturation in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition. Redrawn from Johnson et al. (1999).

DRAFT-DO NOT QUOTE OR CITE



Figure 4-20b. Simulated soil solution A1 and soil base saturation in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition. Redrawn from Johnson et al. (1999).

DRAFT-DO NOT QUOTE OR CITE

in soil solution SO_4^{-2}, NO_3^- and, therefore, MAA concentrations, as expected. This, in turn, caused short-term reductions in base cation concentrations. However, by the end of the 24-year simulation, base cations in the 50% S, N scenario were nearly as high as in the no change scenario because base saturation had increased and the proportion of cations as Al decreased. The 50% C_B scenario had virtually no effect on soil solution SO_4^{-2}, NO_3^- and, therefore, MAA concentrations, as expected, but did cause a long-term reduction in base cation concentrations. This was caused by a long-term reduction in base saturation (Figure 4-20a,b). Thus, the effects of C_B deposition were solely through changes in soils rather than through changes in soil solution MAA, as postulated by Driscoll et al. (1989). In the less acidic Coweeta soil, base saturation was high and little affected by the scenario (cited above); Al was unimportant; and S and N deposition had a much greater effect than C_B deposition in all respects (Figure 4-21a,b).

In summary, Johnson et al. (1999) found that the results of the red spruce simulations support the hypothesis of Driscoll et al. (1989) in part: C_B deposition can have a major effect on C_B leaching through time in an extremely acid system. This effect occurred through changes in the soil exchanger and not through changes in soil solution MAA concentration. On the other hand, S and N deposition had a major effect on Al leaching at the Noland Divide site. This occurred primarily because of changes in soil solution MAA concentration. At the less acidic Coweeta site, C_B deposition had a minor effect on soils and soil solutions; whereas S and N deposition had delayed but major effects on C_B leaching because of changes in SO_4^{-2} and MAA concentrations.

Effects of Trace Elements

Trace metals are natural elements that are ubiquitous in small (trace) amounts in soils, ground water, and vegetation. Many are essential elements required for growth by plants and animals as micronutrients. Naturally occurring surface mineralizations can produce metal concentrations in soils and vegetation that are as high, or higher, than those in the air and deposited near man-made sources (Freedman and Hutchinson, 1981). The occurrence and concentration of trace metals in any ecosystem component depend on the sources of the metal (i.e., via the soil or as a particulate). Even when air pollution is the primary source, continued deposition can result in the accumulation of trace metals in the soil (Martin and Coughtrey,



Figure 4-21a. Simulated soil solution mineral acid anions and base cations in the Coweeta site with no change, 50% N and S deposition, and 50% base cation deposition. Redrawn from Johnson et al. (1999).

DRAFT-DO NOT QUOTE OR CITE



Figure 4-21b. Simulated soil solution mineral acid anions and base cations in the Coweeta site with no change, 50% N and S deposition, and 50% base cation deposition. Redrawn from Johnson et al. (1999).

1981). Many metals are deposited into soils by chemical processes and are not available to plants (Saunders and Godzik, 1986).

When aerial deposition is the primary source of metal particles, both the chemical form and particle size deposited determine the heavy metal concentration in the various ecosystem components (Martin and Coughtrey, 1981). Human activities introduce heavy metals into the atmosphere and have resulted in the deposition of antimony, cadmium, chromium, copper, lead, molybdenum, nickel, silver, tin, vanadium, and zinc (Smith, 1990c). Extensive evidence indicates that heavy metals deposited from the atmosphere to forests accumulate either in the richly organic forest floor or in the soil layers immediately below, areas where the activity in roots and soil is greatest. The greater the depth of soil, the lower the metal concentration. The accumulation of metal in the soil layers where the biological activity is greatest, therefore, has the potential to be toxic to roots and soil organisms and to interfere with nutrient cycling (Smith, 1990e). The shallow rooted plant species are those most likely to take up metals from the soil (Martin and Coughtrey, 1981). Though all metals can be directly toxic at high levels, only toxicity from copper, nickel, and zinc have been documented frequently. Toxicity of cadmium, cobalt, and lead has been seen only under unusual conditions (Smith, 1990c). Exposures at lower concentrations have the potential, over the long-term, for interfering with the nutrient-cycling processes when they affect mycorrhizal function.

Biological accumulation of metals through the plant-herbivore and litter-detrivore chains can occur. A study of the accumulation of cadmium, lead, and zinc concentrations in earthworms suggested that cadmium and zinc were concentrated, but not lead. Studies indicate that heavy metal deposition onto the soil, via food chain accumulation, can cause excessive levels and toxic effects in certain animals. Cadmium appears to be relatively mobile within terrestrial food chains; however, the subsequent mobility of any metal after it is ingested by a herbivorous animal depends on the site of accumulation within body tissues. Although food chain accumulation may not in itself cause death, it can reduce the breeding potential in a population (Martin and Coughtrey, 1981).

In actual case studies, it was observed that the deposition of copper and zinc particles around a brassworks resulted in an accumulation of incompletely decomposed litter. In one study, litter accumulation was reported up to 7.4 km from the stack of a primary smelter in southeastern Missouri. Similar results were reported around a metal smelter at Avonmouth,

England. In the latter case, litter accumulation was associated closely with concentrations specifically of cadmium, as well as with those of lead, copper, and zinc (Martin and Coughtrey, 1981). Experimental data (using mesh bags containing litter) supports the hypothesis that reduced decomposition occurs close to heavy metal sources.

Accumulations of metals emitted in particulate matter also were reported in soil litter close to a metal smelter at Palmerton, PA, in 1975 and 1978. The continued presence of cadmium, lead, zinc, and copper in the upper soil horizons (layers) were observed 6 years after the smelter terminated operation in 1980. Metal levels were highest near the smelter. The relationship of decreasing amounts of metal in body tissues also held true for amphibians and mammals. Levels of cadmium in kidneys and liver of white-tailed deer (*Odocoileus virginaus*) were five times higher at Palmerton than in those collected 180 km southwest downwind. The abnormal amounts of metal in the tissues of terrestrial vertebrates and the absence or low abundance of wildlife at Palmerton indicated that ecological processes within 5 km of the zinc smelter continued to be markedly influenced even 6 years after its closing (Storm et al., 1994).

Accumulation of heavy metals in litter presents the greatest potential for interference with nutrient cycling. Accumulation of metals in the litter occurs chiefly around brass works and lead and zinc smelters. There is some evidence that invertebrates inhabiting soil litter do accumulate metals. Earthworms from roadsides were shown to contain elevated concentrations of cadmium, nickel, lead, and zinc; however, interference with earthworm activity was not cited (Martin and Coughtrey, 1981). It has been shown, however, that when soils are acidic, earthworm abundance decreases, and bioaccumulation of metals from soil may increase exponentially with decreasing pH (Lokke et al, 1996). Organisms that feed on earthworms living in soils with elevated levels of Cd, Ni, Pb, and Z for extended periods could accumulate lead and zinc to toxic levels (Martin and Coughtrey, 1981). Increased concentrations of heavy metals have been found in a variety of small mammals living in areas with elevated heavy metal concentrations in the soils.

Studies by Babich and Stotsky (1978) support the concept that increased accumulation of litter in metal-contaminated areas is due to the effects on the microorganismal populations. Cadmium toxicity to microbial populations was observed to decrease and prolong logarithmic rates of microbial increase, to reduce microbial respiration and fungal spore formation and germination, to inhibit bacterial transformation, and to induce abnormal morphologies. Also, the effects of cadmium, copper, nickel, and zinc on the symbiotic activity of fungi, bacteria, and

actinomycetes were reported by Smith (1991). The formation of mycorrhizae by *Glomus mosseae* with onions was reduced when zinc, copper, nickel, or cadmium was added to the soil. The relationship of the fungus with white clover, however, was not changed. It was suggested that the effect of heavy metals on vesicular-arbuscular mycorrhizal fungi will vary from host to host (Gildon and Tinker, 1983). Studies with ericoid plants indicated that, in addition to *Calluna vulgaris*, mycorrhizae also protect *Vaccinium macrocarpa* and *Rhodendron ponticum* from heavy metals (Bradley et al., 1981). Heavy metals tend to accumulate in the roots, and shoot toxicity is prevented.

The effects of sulfur deposition on litter decomposition in the vicinity of smelters also must be considered. Metal smelters emit SO_2 as well as heavy metals. Altered litter decomposition rates have been well documented near SO_2 sources (Prescott and Parkinson, 1985). The presence of sulfur in litter has been associated with reduced microbial activity (Bewley and Parkinson, 1984). Additionally, the effects on symbiotic activity of fungi, bacteria, and actinomycetes were reported by Smith (1990d).

The potential pathways of accumulation of trace metals in terrestrial ecosystems, as well as the possible consequences of trace metal deposition on ecosystem functions, is summarized in Figure 4-22. The generalized trophic levels found in an ecosystem and the various physiological and biological processes that could be affected by trace metals are shown in the figure. Reduction in physiological processes can affect productivity, fecundity, and mortality (Martin and Coughtrey, 1981). Therefore, any effects on structure and function of an ecosystem are likely to occur through the soil and litter (Tyler, 1972).

Certain species of plants are tolerant of metal contaminated soils (e.g., soils from mining activities) (Antonovics et al., 1971). Certain species of plants also have been used as bioindicators of metals (e.g., *Astragalus* is an accumulator of selenium). The sources of both macroelements and trace metals in the soil of the Botanical Garden of the town of Wroclow, Poland, were determined by measuring the concentrations of the metals in Rhododendron *catawbiense, Ilex aquifolium,* and *Mahonia aquifolium* growing in the garden and comparing the results with the same plant species growing in two other botanical gardens in nonpolluted areas. Air pollution deposition was determined as the source of metals in plants rather than the soil (Samecka-Cymerman and Kempers, 1999).



Figure 4-22. Relationship of plant nutrients and trace metals with vegetation. Compartments (roman numerals) represent potential storage sites; whereas arrows (arabic numerals) represent potential transfer routes.

The effects of lead in ecosystems are discussed in the *Air Quality Criteria for Lead* (U.S. Environmental Protection Agency, 1986). Studies have shown that there is cause for concern in three areas in which ecosystems may be extremely sensitive to lead: (1) delay of decomposition because the activity of some decomposer microorganisms and invertebrates is inhibited by lead, (2) subtle shifts toward plant populations tolerant of lead, and (3) lead in the soil and on the surfaces of vegetation which may circumvent the processes of biopurification. The problems cited above arise because lead is deposited on the surface of vegetation, accumulates in the soil, and is not removed by the surface and ground water of the ecosystem (U.S. Environmental Protection Agency, 1986).

4.2.3 Urban Ecosystems

Humans dominate Earth's ecosystems. Their influence on the environment has been pervasive for thousands of years. Evidence has been accumulating from anthropological and archeological research that human influence has been pervasive for thousands of years (Grimm et al., 2000). Major human impacts on the environment probably began as early as 12,000 to 15,000 years ago. The Earth abounds with both subtle and pronounced evidence of the influence of humans on natural ecosystems. Clearly, human actions have continued to dramatically alter the functioning of ecosystems of which they are a part, and, it is equally clear that humans now influence virtually all ecosystems. Nowhere has human action been more intense than in cities, suburbs, exurbs and in the supporting hinterlands (Grimm et al., 2000). This fact has lead to much recent interest in the study of ecological systems.

Vitousek et al. (1997) point out that understanding a human-dominated planet requires that the human dimensions of global change—the social, cultural, and other drivers of human actions—need to be included within ecological analyses. Therefore, humans must be integrated into models for a complete understanding of extant ecological systems. Development of more realistic models for ecological systems will lead to greater success in finding solutions to environmental problems.

In the past, ecological plant or animal studies conducted in urban settings used traditional ecological approaches and considered humans as agents of disturbance. Although the term urban ecosystem has been used to describe human-dominated ecosystems, it does not adequately take into account the developmental history, sphere of influence, and potential impacts required in order to understand the true nature of an urban ecosystem (McIntyre, et al., 2000). Urban ecology, because urbanization is both an ecological and a social phenomenon, implicitly recognizes the role humans play in developing unique systems. Therefore, if urban ecology is to be a truly interdisciplinary field, it is crucial that it integrate both social and natural sciences into the study of urban ecosystems (McIntyre, et al., 2000).

Although the study of ecological phenomena in urban environments is not a new area of science, the concept of the city as an ecosystem is relatively new for the field of ecology (Grimm et al., 2000). There is a wealth of information on the terrestrial components of urban ecological systems. However, much of it is organized from the perspective of ecology *in* cities while the more comprehensive perspective identified as ecology *of* cities is needed (Pickett et al., 2001).

The basic questions addressed by the literature of ecology *in* cities are how do ecological patterns and processes differ in cities as compared with other environments? What is the effect of the city (i.e., a concentration of human population and activities) on the ecology of organisms inside and outside of its boundary and influence? The concept of ecology *of* cities has to do with how aggregated parts make up the whole, i.e., how cities process energy or matter relative to their surroundings (Grimm et al., 2000). The latter concept includes primary production, species richness, biogeophysical budgets, ecosystem patterns and processes, and an open definition of urban ecosystems that incorporates the exchanges of materials and influence between cities and surrounding landscapes (Pickett et al., 2001). If ecosystems are to be understood, there is a need for a new integrative ecology that explicitly incorporates human decisions, culture, institutions, and economic systems (Grimm et al., 2000). This fact makes an ecological approach to land use planning not only necessary but essential to maintain long-term sustainability of ecosystem benefits, services, and resources (Zipperer et al., 2000). The ecological and social effects of "edge city" need to be studied as they may be greater than the previous patterns of suburbanization. The classical ecosystem approach and a patch dynamic approach are needed to understand and manage the dynamics of urban and urbanizing ecosystems (Zipperer et al., 2000).

There has been little work on the rates of atmospheric deposition to urban ecosystems despite the large body of knowledge on concentrations and chemical reactions of air pollutants in cities. A search of the abundant literature produced no references that dealt with the effects of PM deposition. Lovett et al. (2000), however, reported that urban ecosystems are likely to be subjected to large rates of deposition of anthropogenic pollutants. Decades of research on urban air quality indicate that cities are often sources of nitrogen oxides, sulfur oxides, and dust, among many other pollutants. Some of these air pollutants are major plant nutrients (e.g., nitrogen) and may be affecting nutrient cycles in plant-dominated areas in and around cities. Studying the deposition rates of atmospheric pollutants in urban areas can provide a quantitative estimate of the amounts of gaseous and particulate air pollutants that are removed by urban vegetation.

To determine the patterns of atmospheric deposition and throughfall in the vicinity of a large city, Lovett et al. (2000) measure bulk deposition, oak forest throughfall, and particulate dust at sites along a transect within and to the north of New York City. The gases and particles in urban air can increase atmospheric deposition within and downwind of the city. They observed that concentrations and fluxes of NO_3^-, NH_4^+, Ca^{+2}, Mg^{+2}, SO_4^{-2}, and Cl^- in throughfall

all declined significantly with distance from the city, while hydrogen ion concentration and flux increased significantly with distance from the city. Most of the change in concentrations and fluxes occurred within 45 km of the city. Also, it was observed that throughfall nitrogen was twice as high in the urban areas when compared with the suburban and rural sites. Most of the dry deposition of nitrate was from gaseous nitrogen oxides. As mentioned above, the effects of the atmospheric deposition of the particulate pollutants was not mentioned.

McDonnell et al. (1997) in a 10-year study of ecosystem processes along an urban-rural gradient included plant litter dynamics and nitrogen cycling of two key components of a forest ecosystem: litter decomposition and heavy metal levels in soil and foliar litter. Foliar litter decomposition integrates many features of the abiotic and biotic environment. It is an important site of heavy metal incorporation into ecosystems and provides a both a habitat and a resource for fungi, bacteria, and invertebrates. Litter decomposition integrates the effects of resource quality, environmental factors, and activities of decomposer organisms on nutrient cycling and serves as an easily measured indicator of the effect of urbanization on an important ecosystem function. McDonnell et al. (1997) noted that levels of heavy metals in the foliar litter in urban forest soils were higher than in rural. The levels in urban forest stands approached or exceeded the levels reported to affect soil invertebrates, macrofungi, and soil microbial processes. The urban forests exhibited reduced fungal biomass and microarthropod densities when compared to rural stands. These results supported the concept that urban forests have depauperate communities because of anthropogenic stress resulting from poor air quality due to high levels of SO_2, sulfate, ozone and nitrogen; elevated levels of soil and forest floor heavy metals; and low water availability such as those caused by hydrophobic soils (McDonnell et al.,1997). Thus, forests at the urban end of the gradient exhibited reduced fungal and microarthropod populations and poorer leaf quality than the more rural forests. The potential effect of these conditions on the ecosystem processes of decomposition and nitrogen cycling in urban forests appeared to be ameliorated by two other anthropogenic factors: increased average temperatures caused by the heat island effect and the introduction and successful colonization of earthworms in the urban forests (McDonnell et al.,1997).

McDonnell et al. (1997) observed that the changes in forest nitrogen dynamics were related to increased anthropogenic nitrogen deposition in an urban environment. The studies of Aber et al. (1989) in the northeastern United States on forest nitrogen dynamics demonstrated that

elevated nitrogen deposition over many years results in increased nitrification and the mineralization of more nitrogen than can be taken up by plants and microorganisms. Nitrification can precipitate decreases in fine root biomass and increases in nitrate leaching below the root zone. These effects of nitrogen deposition were not related to inputs from a specific source such as PM.

There have also been studies of heavy metal deposition in or near cities. However, the studies do not cite the effects of metals in the soil. Pouyat and McDonnell (1991) discuss heavy metal accumulations in forest soils along an urban-rural gradient in southeastern New York. Variations in the amounts of Zn, Cu, Ni, and Cd appeared indicative of a pattern of atmospheric deposition near point sources (Section 4.3.2.6). The concentrations of heavy metals in forest floor and soils corresponded closely with the urban-rural land use gradient. Again, as in the study by Lovett et al. (2000), the pollutants were highest near the urban end of the gradient and declined toward rural sites with Pb, Ni and Cu highest near the urban end.

The air quality of the region around East St. Louis has been a source of concern due the industries in the area (Kaminski and Landsberger, 2000a). Industries include ferrous and nonferrous metal smelters (Pb, Zn, Cu, and Al), coal-fired power plants, producers of organic and inorganic chemicals, municipal waste incinerators, and petroleum refineries. The city also is located in the path of diverse plumes from refineries to the north, coal-fired power plants to the west, and nonferrous smelters to the south. Due to years of exposure to the industrial emissions, concerns have arisen with the community about the environmental impact. Concentrations of heavy metals and metalloids (As, Cd, Cu, Hg, Pb, Sb, Zn) in the soil provided a basis for analysis (Kaminski and Landsberger, 2000b). The dual aims of these studies was (1) to make an accurate technical assessment of the extent of the pollutants on the soil surface as well as the extent of the depth to determine possibilities of remediation and (2) to determine the leaching dynamics of heavy metals to determine possible effects on biota uptake or groundwater contamination. The effects on biota are not mentioned; however, the soils in the area acted as a sink and there was little groundwater mobility (Kaminski and Landsberger, 2000b). The possible effects of heavy metals in soils is discussed in the previous section (4.3.2.6) on trace metals.

The above assessment of new information leads to the clear conclusion that atmospheric PM at levels currently found in the United States have the potential to alter ecosystem structure

and function in ways that may reduce their ability to meet societal needs. The possible direct effects of airborne PM on individual plants were discussed in Section 4.2.2.1.

4.2.4 Ecosystem Goods and Services and Their Economic Valuation

Human existence on this planet depends on ecosystems and the services and products they provide. The essential services and products provided by the planet's collective biodiversity (the earth's flora, fauna, and microorganisms) are clean air, clean water, clean soil, and clean energy (Table 4-17). Today, governments around the world tend to pursue a "bottom line" that is driven by an economy that is disconnected from the natural world and is fundamentally destructive of local ecosystems (Suzuki, 1997). For this reason, human society needs to be reconnected to the biologically diverse ecosystems so that they realize that they are a part of the natural world (Suzuki, 1997). There is a need to understand that biodiversity encompasses all levels of biological organization, including individuals, populations, species, and ecosystems (Wilson, 1997). Human-induced changes in biotic diversity and alterations in the structure and functioning of ecosystems are the two most dramatic ecological trends in the past century (Vitousek et al., 1997). Ecosystem processes such productivity, nitrogen mineralization rate, and nitrate leaching respond directly to human modification of ecosystems and to changes in atmospheric composition and climate (Chapin et al., 1997). Habitat conversion, changes in land use, and the introduction of exotic species result in changes in biota, reduced genetic/species diversity, and leads to a homogenization of the global biota. These biotic changes will influence ecosystems processes sufficiently to alter the future state of the world's ecosystems and the services they provide (Chapin et al., 1997).

Though *Homo sapiens* is only one of perhaps 5-30 million animal species on earth, it controls a disproportionate share of the planet's resources. Humans are co-opting approximately 40% of the present net primary production (NPP) of organic material each year. NPP is the amount of energy remaining after subtracting the respiration of primary producers (mostly plants) from the total amount of energy (mostly solar) that is fixed biologically and provides the basis for the maintenance, growth and reproduction of all consumer and decomposers. It is the total food resource of earth (Vitousek et al., 1986).

The number, biodiversity, structure, and functions of ecosystem populations, provide ecosystem products (goods) and services (Figure 4-23). For any given population, the number of

TABLE 4-17. PRIMARY GOODS AND SERVICES PROVIDED BY ECOSYSTEMS

Ecosystem	Goods	Services
Agroecosystems	• Food crops • Fiber corps • Crop genetic resources	• Maintain limited watershed functions (infiltration, flow control, and partial soil protection) • Provide habitat for birds, pollinators, and soil organisms important to agriculture • Sequester atmospheric carbon • Provide employment
Coastal ecosystems	• Fish and shellfish • Fishmeal (animal feed) • Seaweeds (for food and industrial use) • Salt • Genetic resources	• Moderate storm impacts (mangroves, barrier islands) • Provide wildlife (marine and terrestrial) habitat and breeding areas/hatcheries/nurseries • Maintain biodiversity • Dilute and treat wastes • Provide harbors and transportation routes • Provide human and wildlife habitat • Provide employment • Contribute aesthetic beauty and provide recreation
Forest ecosystems	• Timber • Fuelwood • Drinking and irrigation water • Fodder • Nontimber products (vines, bamboos, leaves, etc.) • Food (honey, mushrooms fruit, and other edible plants; game) • Genetic resources	• Remove air pollutants, emit oxygen • Cycle nutrients • Maintain array of watershed functions (infiltration, purification, flow control, soil stabilization) • Maintain biodiversity • Sequester atmospheric carbon • Moderate weather extremes and impacts • Generate soil • Provide employment • Provide human and wildlife habitat • Contribute aesthetic beauty and provide recreation
Freshwater	• Drinking and irrigation water • Fish • Hydroelectricity • Genetic resources	• Buffer water flow (control timing and volume) • Dilute and carry away wastes • Cycle nutrients • Maintain biodiversity • Provide aquatic habitat • Provide transportation corridor • Provide employment • Contribute aethetic beauty and provide recreation
Grassland ecosystems	• Livestock (food, game, hides, and fiber) • Drinking and irrigation water • Genetic resources	• Maintain array of watershed functions (infiltration, purification, flow control, and soil stabilization) • Cycle nutrients • Remove air pollutants and emit oxygen • Maintain biodiversity • Generate soil • Sequester atmospheric carbon • Provide human and wildlife habitat • Provide employment • Contribute aesthetic beauty and provide recreation

Source: World Resources (2000-2001).



Figure 4-23. Linkages among various ecosystem goods and services (food, water, biodiversity, forest products) and other driving forces (climate change) (modified from Ayensu et al., 1999).

individuals, the genetic variation between individuals, and the area occupied affects ecosystem functioning and the delivery of ecosystem services and other benefits provided by that population (Hughes, et al., 1997). Loss or altering of population diversity means loss of the benefits described in Table 4-17 and, in particular, with time, the loss of the life-support systems on which humanity relies (Hughes et al., 1997).

Attempts have been made to calculate the value of biodiversity and the world's ecosystem services and natural capital (Pimentel et al., 1997; Costanza et al., 1997). Pimentel et al. (1997) estimated economic and environmental benefits for services contributed from all biota in the United States, including their genes, at $319 billion per year. Costanza et al. (1997) have estimated the total value of ecosystem services by biome for the entire bioshere and concluded that ecosystems provide at least $33 trillion worth of services annually. Approximately 63% of

the estimated value is contributed by marine ecosystems ($20.9 trillion per year), most of which comes from coastal ecosystems ($10.6 trillion per year). About 38% of the estimated value comes from terrestrial ecosystems, mainly from forests ($4.7 trillion per year) and wetlands ($4.9 trillion per year). Costanza et al. (1997) state that it may never be possible to make a precise estimate of the services provided by ecosystems; however, their estimates indicate the relative importance of ecosystem services, not their true value considering that the loss of ecosystem services can affect human existence.

Heal (2000), however, feels that attempts to value ecosystems and their services are misplaced: "Economics cannot estimate the importance of natural environments to society: only biology can do that" (Heal, 2000). The role of economics is to help design institutions that will provide incentives to the public and policy makers for the conservation of important natural systems and for mediating human impacts on the biologically diverse ecosystems and the biosphere so that they are sustainable. The approach of Harwell et al. (1999) also deals with the need to understand human effects on ecosystems so that ecosystem management can define what ecological conditions are desired. Further, they state that the establishment of ecological goals involves a close linkage between scientists and decision makers in which science informs decision makers and the public by characterizing the ecological conditions that are achievable under particular management regimes. Decision makers then can make choices that reflect societal values including issues of economics, politics, and culture. For management to achieve their goals, the general public, scientific community, resource managers, and decision makers need to be routinely apprised of the condition or integrity of ecosystems so that ecological goals may be established (Harwell et al., 1999).

Though usually considered as toxic pollutants locally (Section 4.3.2.3), secondary organics as PM can become airborne and distributed over a wide area and affect ecosystems remote from the source. Some of the chemical compounds may reach toxic levels in the food chains of human and other animals. However, other compounds tend either to decrease or maintain the same level of toxicity. The effects of toxicity on the animal population can alter the functioning of the ecosystem. The major impacts of airborne PM on ecosystems, however, are through the indirect effects on plant populations that occur through the soil and affect the cycling of nutrients necessary for plant growth, vigor, and maintenance of biodiversity as discussed in Section 4.2.2.2. By altering the cycling of nitrogen, nitrogen deposition changes the biodiversity of

ecosystems and their functioning and, by altering the vigor of forest tree stands, alters forest succession. Additionally, nitrogen deposition in combination with the deposition of sulfur in the form of acid rain alters the biogeochemical cycling of soil mineral nutrients and changes the biodiversity and functioning of forest ecosystems. The changes in the ability of forest vegetation and soil microorganisms to utilize nutrients results in the leaching of nitrates and other minerals from the soils. The nitrate and mineral runoff affects streams and coastal and aquatic ecosystems and, thus, influences the services important to human life provided by these ecosystems as well (Table 4-17).

4.3 EFFECTS ON VISIBILITY

4.3.1 Introduction

Visibility may be thought of as the degree to which the atmosphere is transparent to visible light (National Research Council, 1993). The beauty of scenic vistas in many parts of the U.S. is often diminished by haze that reduces contrast, washes out colors, and renders distant landscape features indistinct or invisible. This degradation of visibility is due primarily to the scattering and absorption of light by fine particles suspended in the atmosphere. One quantitative measure of visibility, used traditionally by meteorologists, is the visual range, defined as the farthest distance at which a large black object can be distinguished against the horizon sky (U.S. Environmental Protection Agency, 1979).

In August 1977, Congress amended the Clean Air Act (CAA) to establish as a national goal "the prevention of any future and remedying of any existing impairment of visibility in mandatory Class I Federal areas (many national parks and wilderness areas), which impairment results from manmade air pollution" (Title I Part C Section 169A, U.S. Code [1990]). The 1977 Amendments also included provisions requiring applicants for new major source permits to assess the potential for their projects to cause adverse impacts on the air quality-related values, including visibility, in nearby Class I areas. In 1980, the EPA established regulatory requirements under Section 169A to address Class I protection from "reasonably attributable" visibility impairment, i.e., visibility impairment attributable to a single source or small group of sources.

The CAA, as amended in 1990 (section 169B), required the U.S. Environmental Protection Agency to conduct research on regional visibility impairment and to establish the Grand Canyon Visibility Transport Commission (GCVTC). The GCVTC was charged with assessing and providing recommendations to help preserve clear days and improve visibility in the 16 national parks and wilderness areas located on the Colorado Plateau. The GCVTC was mandated to provide recommendations to the U.S. Environmental Protection Agency for the reduction of visibility impairment due to regional haze, described as any perceivable change in visibility (light extinction, visual range, contrast, or coloration) from that which would have existed under natural conditions that is caused predominantly by a combination of many anthropogenic sources over a wide geographical area (U.S. Environmental Protection Agency, 1999a). In July 1999, the U.S. Environmental Protection Agency published the Regional Haze Rule (Federal Register, 1999). The regulation established a program for the improvement and protection of visibility in the 156 protected Class I parks and wilderness areas, including the establishment of baseline and current visibility conditions and the tracking of changes in visibility conditions over time. Implementation of the regional haze regulations is supported by the U.S. Environmental Protection Agency's $PM_{2.5}$ monitoring network and an expanded Interagency Monitoring of Protected Visual Environments (IMPROVE) network. A discussion on the $PM_{2.5}$ monitoring network and the IMPROVE network appears elsewhere in this section (National Park Service, 1998; Evans and Pitchford, 1991; U.S. Environmental Protection Agency, 2000b; U.S. Environmental Protection Agency, 2001).

The objective of the visibility discussion in this section is to provide a brief description of the fundamentals of atmospheric visibility and to summarize the linkage between particulate matter and visibility. Visibility is an effect of air quality and, unlike the particulate matter concentration, is not a property of an element of volume in the atmosphere. Visibility can be quantified only for a sight path and depends on the illumination of the atmosphere and the direction of view. However, the concentration of particles in the atmosphere plays a key role in determining visibility. Therefore, visibility impairment may be controlled by control of particle concentrations. The relationships between particles, other factors, and visibility impairment are described in this section. For a more detailed discussion on visibility, the reader is referred to the 1996 Air Quality Criteria for Particulate Matter (PM AQCD) (U.S. Environmental Protection Agency, 1996a), the Recommendations of the Grand Canyon Visibility Transport Commission

(Grand Canyon Visibility Transport Commission, 1996), the National Research Council (National Research Council, 1993), the National Acid Precipitation Assessment Program (Trijonis et al., 1991), Interim Findings on the Status of Visibility Research (U.S. Environmental Protection Agency, 1995a), and reports summarizing visibility science and data from the IMPROVE visibility monitoring network (Malm et al., 2000; Sisler, 1996; Sisler et al., 1993).

4.3.2 Factors Affecting Atmospheric Visibility

The visual perceptions of a distant object is influenced by a large number of factors including human vision, various characteristics of the atmosphere (e.g., atmospheric illumination, path and transmitted radiance, contrast, and optical properties), and atmospheric pollution. Detailed discussion of this full range of topics can be found in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) and other general references (e.g., Malm, 1999). This section focuses only on those topics that have generally been addressed by more recent research, including atmospheric illumination, the optical properties of gases and particles in the atmosphere, and the effects of relative humidity on the optical properties of particles.

4.3.2.1 Optical Properties of the Atmosphere and Atmospheric Particles

Atmospheric particles and gases attenuate image-forming light as it travels from a viewed object to an observer. The fractional attenuation of light per unit distance is known as the light extinction coefficient. The light extinction coefficient, b_{ext}, is expressed in units of one over length, for example inverse kilometers (km^{-1}) or inverse megameters (Mm^{-1}). The light extinction coefficient can be expressed as the sum of the light scattering and light absorption coefficients of particles and gases.

$$b_{ext} = b_{ap} + b_{ag} + b_{sg} + b_{sp} \qquad (4\text{-}6)$$

where the subscripts *p* and *g* signify particles and gases, and *s* and *a* signify scattering and absorption.

Transmitted Radiance and Path Radiance

The appearance of a distant object is determined by light from two sources: the light reflected from the object itself and the light reflected by the intervening atmosphere. Light reflected by the object is attenuated by scattering and absorption as it travels through the atmosphere toward the observer. The portion that reaches the observer is the transmitted radiance. During the daytime, the sight path is illuminated by the direct rays of the sun, diffuse skylight, light that has been reflected from the surface of the Earth, etc. Some of this illumination is scattered toward the observer by the air molecules and particulate matter in the sight path. The accumulation of the light scattered into the sight path is the path radiance or air light. The path radiance is significantly influenced by the illumination of the sight path. However, not all of the light scattered into the sight path reaches the observer.

The transmitted radiance carries the information about the object. The path radiance only carries information about the intervening atmosphere and is often quite featureless. When the transmitted radiance is dominant, visibility is good. Conversely, when the path radiance is dominant, visibility is poor. In a dense fog, the transmitted radiance from nearby objects can be seen, but the transmitted radiance from more distant objects is completely overwhelmed by the path radiance (i.e., the light scattered by the fog). Distant objects are lost in the white (or gray) of the fog (Gazzi et al., 2001).

Figure 4-24 illustrates the radiance seen by an observer looking at a hillside or through the aperture of a measurement instrument. The radiance that enters the eye of the observer (or the aperture of a measurement instrument) is known as the apparent radiance (i.e., the sum of the transmitted and path radiance). It is the competition between the transmitted radiance and the path radiance that determines visibility.

Light Absorption and Scattering by Gases

In the ambient atmosphere the only visible light absorbing gas of any consequence is nitrogen dioxide (NO_2), which absorbs primarily blue light, thus causing a yellow or brown color if in sufficient concentration across a sight path. Usually the absorption by NO_2 is much smaller than the scattering by particles that are typically present in polluted environments, such as urban areas. The most common exception to this situation of relatively small NO_2 absorption is in effluent plumes from combustion facilities where the particles are effectively removed but the



Figure 4-24. Light reflected from a target toward an observer. The intervening atmosphere scatters a portion of this light out of the sight path and scatters light from the sun into the sight path. Some particles and gases also absorb a portion of the light from the target. The light scattered into the sight path increases with distance from the target, whereas the light transmitted from the target decreases with distance from the target. The visual range is the closest distance between the target and the observer at which the transmitted light no longer can be distinguished from the light scattered into the sight path.

Source: Watson and Chow (1994).

nitrogen oxide (NO), which can convert rapidly to NO_2, is not removed. Except for such particle- depleted NO plumes, the light absorption coefficient for gases is usually ignored in determinations of the light extinction coefficient.

Scattering by gases in the atmosphere is described by the Rayleigh scattering theory (vandeHulst, 1981) and is referred to as Rayleigh scattering. The magnitude of the Rayleigh scattering depends on the gas density of the atmosphere and varies from about 9 Mm^{-1} to 11 Mm^{-1} for most locations of interest, depending primarily on site elevation. To simplify

comparisons of light extinction coefficient values among sites at a variety of elevations, a standard value of 10 Mm^{-1} is often used for the Rayleigh scattering component (Malm, 2000).

Light Absorption by Particles

Absorption by particles is principally caused by elemental carbon (also referred to as soot or light-absorbing carbon), which usually results from incomplete combustion of fossil fuels. Some minerals in crustal particles also absorb light and can be a significant factor during fugitive dust episodes.

Most particle absorption data are determined by measuring light transmission of particles captured on filter media. Absorption estimates made in this way are sensitive to the filter substrate used, the optical configuration of the transmission measurement, particle loading on the filter, and particle scattering albedo, with the result that there are significant uncertainties for measurements of filtered particles (Horvath, 1993). Another approach to estimating aerosol light absorption is by subtracting concurrent light scattering measurements, using a nephelometer, from light extinction measurements, using a transmissometer. Substantial uncertainties in this difference approach result from the assumption that the point measurement of scattering is representative of the scattering over a long path (1 to 10 km), typically required for transmissometers measurements. A recently field-tested prototype photoacoustic spectrometer designed to determine absorption of suspended aerosol and an enclosed-folded path transmissometer offers hope of resolving the problems of the filter-based and difference approaches to the measurement of light absorption by particles (Arnott et al., 1999).

The relationship between elemental carbon concentration and particle absorption can be calculated using Mie equations for particles with known size distribution, particle density, index of refraction, shapes, and for various internal mixtures with non-absorbing aerosol materials (Fuller et al., 1999). While such application of this theory can provide a range of absorption efficiencies for various model aerosol distributions, it is rare that sufficiently detailed particle characterization data for ambient aerosols are available. Also, although elemental carbon is the strongest and most common of the absorbing particles, light absorption by elemental carbon particles can be reduced when the particle is covered by other chemical species (Dobbins et al., 1994) or may be enhanced when coated with a non-absorbing refractive material such as ammonium sulfate (Fuller et al., 1999).

More commonly, estimates of elemental carbon absorption efficiency are empirically determined from the ratios of or the slopes of regression analysis fits to absorption coefficient and corresponding elemental carbon concentration measurements. Use of the regression approach permits the inclusion of crustal component concentrations as a second dependent parameter, so that crustal absorption can also be estimated. Uncertainties in the absorption efficiency determined empirically are a combination of the measurement uncertainties for the absorption coefficients, elemental carbon concentrations, and where used, the crustal concentrations. In reviews of estimates of elemental carbon light absorption mass efficiency (i.e., the absorption coefficient per carbon mass concentration), Horvath (1993) and Liousse et al. (1993) found values ranging from 2 to 17 m^2/g. Moosmüller et al. (1998) showed that by limiting the absorption coefficient estimates to those using photoacoustic methods, the absorption efficiency shows a wavelength dependence, with highest values (17 m^2/g) at the shortest wavelength used ($\lambda = 0.42\ \mu m$) and lowest values (3 m^2/g) at the longest wavelengths used ($\lambda = 0.8\ \mu m$). The center of the visible light wavelength ($\lambda = 0.53\ \mu m$) yielded elemental carbon absorption efficiencies values of about 10 m^2/g, which is a commonly used value for elemental carbon absorption efficiency. Fuller et al. (1999) suggested that isolated spheres of light absorbing carbon have a specific absorption of less than 10 m^2/g. Light absorption by carbon particles only will be greater than 10 m^2/g if the particles are internally mixed and the occluding particles are sufficiently large. Absorption values for graphitic and amorphous carbon for primary sizes typical of diesel soot are around 5 m^2/g.

Light Scattering by Particles

Particle scattering tends to dominate light extinction, except under pristine atmospheric conditions when Rayleigh scattering by gas molecules is the largest contributor. The Mie equation can be used to calculate particle scattering for aerosols of known size distribution, particle density, index of refraction, shape, and for various known internal component mixtures (Fuller et al., 1999). Unlike particle absorption, which is principally associated with elemental carbon, all particles scatter light.

Light-scattering by particles has been reported to account for 68 to 86% of the total extinction coefficient in several cities in California (Eldering et al., 1994). When light-scattering increases, visibility is impaired because of a decrease in the transmitted radiance and an increase

in the path radiance. The single most important factor that determines the amount of light scattered by a particle is its size, as shown in Figure 4-25. The maximum single particle scattering efficiency (i.e., scattering per cross-sectional area of the particle) is associated with particles with diameters of about the wavelength of visible light (centered at 0.53 μm). For particles that are small compared to the wavelength of light, the single particle scattering efficiency is low. For particles larger than the wavelength, the single particle scattering efficiency initially decreases with diameter and then fluctuates around a value of two as size increases. However, a larger particle always scatters more light than a smaller particle because particle cross-sectional area increases faster with diameter than the decrease in single particle scattering efficiency at any point on the scattering efficiency curve. The mass scattering efficiency (i.e., the scattering per mass concentration) peaks for particles that are about 0.5 μm to 0.8 μm in diameter. Smaller particles are much less efficient at scattering light, while larger particles have mass that increases with particle size faster than the increase in the amount of light they scatter.



Figure 4-25. Light-scattering efficiency factor (per cross sectional area), Q, for a homogeneous sphere with an index of refraction of 1.50 as a function of the size parameter, $\alpha = \pi D/\lambda$.

Source: Penndorf (1958).

Use of the Mie equation to calculate light scattering or the light scattering efficiency of particles in the atmosphere is severely limited by the general lack of sufficiently detailed particle characterization data. At a minimum, size-resolved particle composition data (e.g., aerosol collected on an 8-stage impactor) are needed to permit meaningful Mie scattering calculations. The chemical composition provides clues to the appropriate particle density and index of refraction, while the size distribution is inferred by fitting a distribution function to the concentration for each stage. Assumptions are still necessary to address the particle component mixture characteristics of the aerosol. Resulting scattering calculations can be compared to directly measured particle extinction to assess the reasonableness of the Mie calculations.

Reported calculated scattering efficiencies for sulfates range from 1.2 to 5.6 m^2/g. Sulfate scattering efficiencies have been reported to increase by a factor of two when the size distribution went from 0.15 to 0.5 μm (McMurry et al., 1996). Calculated scattering efficiencies for carbon particles ranged from 0.9 to 8.1 m^2/g. A scattering efficiency of 1.0 and 0.6 m^2/g was reported for soil and coarse mass, respectively (U.S. Environmental Protection Agency, 1996a; Sisler and Malm, 2000).

Integrating nephelometers directly measure ambient particle scattering. A variety of nephelometer configurations that can include the use of unrestricted or size selective inlets and the control of sample air temperature and relative humidity permit the composite scattering properties of ambient aerosol to be directly observed (Day et al., 1997). When sample-controlled nephelometer data are combined with collocated particle speciation data, composite particle scattering efficiency values for ambient aerosol can be empirically derived (Malm et al., 2000).

4.3.2.2 Relative Humidity Effects on Particle Size and Light-Scattering Properties

The ability of some commonly occurring chemical components of atmospheric aerosol to absorb water from the vapor phase has a significant impact on particle light scattering. Hygroscopic particulate materials, which typically include sulfuric acid, the various ammonium sulfate salts, ammonium nitrate, and sodium chloride, change size by the accumulation and loss of water as they maintain equilibrium with the vapor phase as a function of changes in relative humidity. For some materials (e.g., sulfuric acid), the growth is continuous and reversible over the entire range of relative humidity. For other materials, water absorption begins abruptly for a dry particle at a specific relative humidity known as the deliquescent point (e.g., ~80% for

ammonium sulfate) and continues as relative humidity increases. There is a hysteresis effect with these materials in that, once wet, the relative humidity can be reduced below the deliquescent point until crystallization occurs at a substantially lower relative humidity (e.g., ~30% for ammonium sulfate). Figure 4-26 shows the water vapor growth curve for ammonium sulfate.



Figure 4-26. Particle growth curve as a function of relative humidity showing deliquescent growth of ammonium sulfate [$(NH_4)_2 SO_4$] particles at the deliquescent point (A, about 80% relative humidity [RH]), reversible hygroscopic growth of ammonium sulfate solution droplets at RH greater than 80%, and hysteresis (the droplet remains supersaturated as the RH decreases below 80%) until the crystallization point (B, about 38% RH) is reached.

Source: Adapted from National Research Council (1993) and Tang (1980).

Water growth behavior for hygroscopic materials commonly found in atmospheric aerosol in pure form or in some mixtures is generally well known as a result of laboratory measurements (Tang and Munkelwitz, 1994; Tang, 1997). Models that calculate water growth of mixtures from

known solubility properties of many common water-soluble chemicals have long been available (Zdanovskii, 1948) and have been successfully applied to determine growth for particles with known composition (Saxena and Peterson, 1981; Pilinis et al., 1995; Saxena et al., 1993).

The water growth of individual ambient particles can be directly measured using a humidity-controlled tandem differential mobility analyzer or TDMA (McMurry and Stolzenburg, 1989; Zhang et al., 1993). Inferences can be made about the mixtures of soluble and insoluble particle components by comparing TDMA measured growth and size-resolved aerosol composition data with water growth model predictions (Pitchford and McMurry, 1994; Zhang et al., 1993; Saxena et al., 1995). A practical limitation of TDMA measurements in investigating aerosol optical properties is that particles with diameter greater than 0.5 μm are not well measured by this approach.

Accounting for water growth of atmospheric aerosols is important for visibility because particles containing hygroscopic or deliquescent materials change size and index of refraction, and hence scattering efficiency, with changing relative humidity. The nonlinear nature of particle growth curves for hygroscopic aerosols means that substantial light scattering changes result from modest relative humidity changes under humid conditions (relative humidity > 90%). The magnitude of the water growth effect on light scattering for ambient aerosols can be directly measured with humidity-controlled nephelometer measurements (Day et al., 1997). Measurements of water growth effects on scattering are compared to results of water growth and Mie scattering models applied to size-resolved composition data using various mixture assumptions to infer average mixture and other aerosol characteristics (Malm et al., 2000).

While the importance of inorganic hygroscopic particles is well understood, the role of organic compounds in particle water growth has been the subject of recent investigations. In their interpretation of TDMA and particle composition data from two locations, Saxena et al. (1995) made the case that organic components of the aerosol enhanced water absorption by particles at a remote desert location and retarded water absorption at an urban location. They speculated that the latter might be due to hydrophobic organic material coatings on inorganic hygroscopic particles.

While some of the thousands of organic compounds that are in atmospheric aerosols are known to be hygroscopic and while a significant fraction of the organic aerosol material is known to be water soluble, there is a general lack of water absorption data for most organic

compounds. The incomplete water solubility data, combined with incomplete data on the abundance of the numerous organic compounds in ambient aerosols, means that organic water growth model calculations are not a reasonable approach to assessing the importance of water growth by organic aerosol components in the atmosphere. To overcome this constraint, Saxena et al. (1995) compared organic concentration to the difference between total aerosol water measured by TDMA and model-estimated water for the inorganic hygroscopic aerosol components. One the other hand, Pitchford and McMurry (1994) using the same remote location data set showed that on six of the eight sampling days water uptake by the sulfates and nitrates could account for all of the measured water absorption.

Swietlicki et al. (1999) made TDMA measurements in northern England and found that growth takes place in two modes, one mode being less hygroscopic that the other. They concluded that growth could be attributed to the inorganic content of the aerosol. Cocker et al. (2001) measured hygroscopic properties of Pasadena, California aerosol and concluded that growth factors increased when forest fires were present. McDow et al. (1995) measured water uptake by diesel soot, automobile exhaust, and wood smoke particles. They found all three emission types absorbed water, with the wood smoke sample weight increasing by about 10% as sample relative humidity was increased; whereas diesel soot sample weight increased by only 2% to 3%. Chughtai et al. (1999) examined hydration characteristics of a number of anthropogenic and natural organic materials. They found surface water adsorption to increase with age and surface oxidation. Analysis of humidity controlled and size-resolved chemistry data from Great Smoky Mountains and Grand Canyon National Parks (Malm et al., 1997; Malm and Kreidenweis, 1996; Malm et al., 2000) show that, to within the measurement uncertainty and modeling assumptions, ambient organic aerosol are at most weakly hygroscopic.

A more detailed discussion of the effects of relative humidity on the size distribution of ambient particles appears in Chapter 2 of this document.

4.3.3 Relationships Between Particles and Visibility

Visibility, referring to the appearance of scenic elements in an observer's line of sight, depends on more than the optical characteristics of the atmosphere. Numerous scene and lighting characteristics are important to this broad definition of visibility. However, under a variety of

viewing conditions, visibility reduction or haziness is directly related to the extinction coefficient, which, as shown above, is related to the concentrations of ambient particles.

Often visibility conditions are communicated in terms of the visual range, which is commonly taken to be the greatest distance that a large dark object (e.g., a mountain in shadow) can be seen against the background sky (Middleton, 1952). Visual range was developed and continues to function well as an aid in military operations and transportation safety. An inverse relationship between visual range and the light extinction coefficient, know as the Koschmeider constant, was developed using a number of restrictive assumptions about lighting, scene, and atmospheric conditions.

$$Visual\ Range = 3.912/\ b_{ext} \qquad (4\text{-}7)$$

where visual range is in kilometers, b_{ext} is in km^{-1}, and a threshold contrast of 2% is assumed. If b_{ext} is in Mm^{-1}, the Koschmeider constant becomes 3,912.

A new index of haziness, expressed in deciview (dv) units, is also very simply related to the light extinction coefficient (Pitchford and Malm,1994).

$$Haziness\ (dv) = 10\ ln(\ b_{ext}/10\ Mm^{-1}) \qquad (4\text{-}8)$$

An important characteristic of this visibility index is that it is more nearly linearly related to perceived changes in haze level than either visual range or light extinction. A change of 1 or 2 dv in uniform haze under many viewing conditions will be seen as a small but noticeable change in the appearance of a scene regardless of the initial haze condition.

Figure 4-27 illustrates the relationship of light extinction in Mm^{-1}, deciview index, and visual range in kilometers. Although the deciview is related to extinction, it is scaled in such a way that is perceptually correct (Fox et al., 1999).

Comparisons of paired light extinction coefficient (or scattering coefficient) and particle mass concentration data reveal a definite but noisy linear relationship. In general such a relationship can be improved by either restricting the data to periods of low relative humidity or by empirically adjusting for the nonlinear effects of water growth using relative humidity data (White and Roberts, 1977; Malm, 1989). Where particle speciation data for the major aerosol



Figure 4-27. Comparison of extinction (Mm^{-1}) and visual range (km).

Source: Fox et al. (1999).

components are available, the relationship between particles and light extinction can be further improved by treating the individual major components separately.

Most routine aerosol monitoring programs and many special study visibility characterization programs were designed to measure the major aerosol components (Malm et al., 1994; Tombach and Thurston, 1994; Watson et al., 1990); they were not designed to determine the microphysical and chemical characteristics of these species. However, the inherent limitations of estimating aerosol optical properties from bulk aerosol measurements have been addressed, at least in part, by a number of authors. For instance, Ouimette and Flagan (1982) have shown from basic theoretical considerations that if an aerosol is mixed externally (i.e., separate particles contain the major aerosol components), or if in an internally mixed aerosol the index of refraction is not a function of composition or size and the aerosol density is independent of volume, then

$$b_{sp} = \sum \alpha_i m_i \qquad (4\text{-}9)$$

where α_i is the specific mass scattering efficiency and m_i is the mass of the individual aerosol species.

Sloane (1983, 1984, 1986), Sloane and Wolff (1985), and more recently Lowenthal et al. (1995) and Malm and Kreidenweiss (1997) have shown that differences in estimated specific scattering between external and internal model assumptions are usually less than about 10%. In the absence of detailed microphysical and chemical information of ambient particles, the

above studies demonstrate that a reasonable estimate of aerosol extinction can be achieved by assuming each species is externally mixed.

The latest IMPROVE Program report (Malm, 2000) includes calculated aerosol light extinction for each of the five major fine fraction particle ($PM_{2.5}$) components plus coarse fraction mass ($PM_{10-2.5}$) and sums them for an estimate of total light extinction in Mm^{-1} using the following algorithm:

$$
\begin{aligned}
b_{ext} = {} & (3)\, f\,(RH)\, [SULFATE] + \\
& (3)\, f\,(RH)\, [NITRATE] \\
& +(4)\, [ORGANIC\ CARBON] \\
& +(10)\, [LIGHT\ ABSORBING\ CARBON] \\
& +(1)\, [SOIL] \\
& +(0.6)\, [COARSE\ PM] \\
& +10\ (for\ Rayleigh\ scattering\ by\ gases)
\end{aligned}
\tag{4-10}
$$

where each PM term is the product of a constant dry extinction efficiency for that species, the mass concentration of the species, and, for sulfate and nitrate, an adjustment factor that is a function of relative humidity to account for their hygroscopic behavior. The relative humidity adjustment term for sulfate and nitrate, shown in Figure 4-28, is based upon the ammonium sulfate growth curve, shown in Figure 4-26, smoothed between the upper and lower curves of the hysteresis loop for the relative humidity range of 30-80%.

The extinction efficiencies for soil and coarse mass used in this algorithm are taken from a literature review by Trijonis et al. (1987). The extinction efficiency for light absorbing (elemental) carbon of 10 m^2/g is consistent with the value reported by Moosmüller et al. (1998) corresponding to $\lambda = 0.53$ in the middle of the visible light spectrum. The dry extinction efficiencies of 3 m^2/g for sulfate and nitrate species and 4 m^2/g for organic species are based on literature reviews by Trijonis et al. (1991) and by White (1991). Trijonis' best estimate for sulfates is 2.5 m^2/g with an uncertainty of a factor of 2, while White's average low and high estimates for the rural West are 3.0 and 3.7 m^2/g, respectively. For organics Trijonis estimates a dry extinction efficiency of 3.75 m^2/g with an uncertainty of a factor of 2, and White's range for



Figure 4-28. Relative humidity adjustment factor, f(RH), for ammonium sulfate as a function of relative humidity.

Source: Malm et al. (2000).

the rural West is 1.8 to 4.1 m^2/g. Malm et al. (1996) and Malm (2000) used this algorithm to successfully reconstruct scattering at a total of eleven IMPROVE monitoring sites.

Malm (2000) used additional sophisticated aerosol size, composition, and microphysical data from a special study at the Great Smoky Mountain National Park to compare the performance of a number of models for calculating light extinction. He found that the simplist approach adequately predicted for periods of low light scattering but under-predicted by about 30% during periods of high sulfate concentration. The greatest improvement over the simple model was obtained by including the degree of sulfate ammoniation in the model, which produced better estimates of extinction coefficient over the entire range.

4.3.4 Photographic Modeling of Visibility Impairment

None of the visibility indices communicate visibility associated with various aerosol conditions as well as directly seeing their effects on a scene. However, photographic modeling

for the representation of haze can be useful in portraying changes in visibility specifically due to changes in air pollutant concentrations. Photographic modeling holds constant the effects of sun angle, cloud cover, and relative humidity and is a cost-effective method of evaluating various air quality scenarios. This is difficult to do with actual photographs because of the range of possible conditions in the same scene over multiple days; and, over time, photographs can be expensive to produce. One of the limitations in using photographic models for representation of haze is that haze is assumed to be uniformly distributed throughout the scene and selected conditions are idealized, so the full range of conditions that occur in a scene are not represented.

Eldering et al. (1996) proposed the use of a model that uses simulated photographs from satellite and topographic images to evaluate the effect of atmospheric aerosols and gases on visibility. Use of this model requires ground-based photography and size distribution and chemical composition of atmospheric aerosols, NO_2 concentration, temperature, and relative humidity for a clear day, for comparison purposes. Light extinction and sky color are then calculated based on differences in aerosol size distribution, NO_2 concentration, temperature, and relative humidity. The images created represent natural landscape elements.

Molenar et al. (1994) provides a discussion of existing visual air quality simulation methods based on techniques under development for the past 20 years. A photograph taken on a very clean, cloud-free day serves as the base image. The photograph is taken during the season and at the same time of day as the scene to be modeled. The light extinction represented by the scene is derived from aerosol and optical data associated with the day the image was taken, or it is estimated from contrast measurements of features in the image. The image is then digitized to assign an optical density to each picture element (pixel) for the wavelength bands of interest. A detailed topographic map and an interactive image-processing display system is used to determine the specific distance, elevation angle, and azimuth angle for each element in the picture with respect to the observer's position.

Various models are employed to allow the presentation of different air quality scenarios. The output from atmospheric aerosol models (e.g., extinction, scattering coefficients, single scattering albedo, and scattering phase matrix) is incorporated into radiative transfer models to calculate the changes in radiant energy (path radiance, image radiance, sky radiance, terrain radiance) caused by scattering and absorption by gases and particles as it passes through the

atmosphere. Atmospheric aerosol models are also use to model the effect of relative humidity on the visual air quality (Molenar et al., 1994).

Molenar et al. (1994) has developed a system call WinHaze that permits the viewing of computer-generated uniform hazes superimposed on digitized scenic photographs of both remote and urban scenes. The program simulates changes in visual air quality imagery from user-specified changes in optical parameters (e.g., σ_{ext}, visual range, or deciview values) or aerosol species concentrations. WinHaze includes imaging for various Class I national parks and wilderness areas and Boston, MA; Dallas, TX; Denver, CO; Fort Collins, CO; Phoenix, AZ; and Tucson, AZ. The computer software is available through the IMPROVE website (http://vista.cira.colostate.edu/ improve/).

4.3.5 Visibility Monitoring Methods and Networks

Visibility monitoring studies measure the properties of the atmosphere either at the sampler inlets (point measurements), as is the case with air quality measurements, or by determining the optical properties of a sight path through the atmosphere (path measurements). Instrumental methods for measuring visibility are generally of three types: (1) direct measurement of light extinction of a sight path using a transmissometer, (2) measurement of light scattering at one location using an integrating nephelometer, and (3) measurement of ambient aerosol mass concentration and composition (Mathai, 1995).

The largest instrumental visibility monitoring network in the United States is the Automated Surface Observing System (ASOS). The Automated Surface Observing System has been commissioned by the National Weather Service, Federal Aviation Administration, and Department of Defense at more than 900 airports. The system is designed to objectively measure the clarity of the air versus the more subjective evaluations of human observations. The system provides real-time data for airport visibility.

The visibility sensor, instead of measuring how far one can see, measures the clarity of the air using a forward scatter visibility meter. The clarity is then converted to what would be perceived by the human eye using a value called Sensor Equivalent Visibility (SEV). Values derived from the sensor are not affected by terrain, location, buildings, trees, lights, or cloud layers near the surface. The amount of moisture, dust, snow, rain, and particles in the light beam will affect the amount of light scattered. The sensor transmits 1-min values based on rolling

10-min periods. The value provides a generally accurate and representative visibility measurement within a 2 to 3 mile radius of the site. The forward scatter meter was found to correlate fairly well with extinction coefficient measurements from the Optec Transmissometer (National Weather Service, 1998).

Visibility data from the ASOS network is reported in terms of visual range in increments of 1/4 to 1 statute mile. Visual range conditions exceeding 10 miles are truncated to 10 miles for real-time reporting purposes. Data is not extensively archived at ASOS locations. However, researchers have been able to download the raw data directly from certain sites. In addition, since 1998, the raw visibility data (including light extinction measurements corresponding to visual ranges exceeding 10 miles) have been archived for a number of sites. These data are available from the National Climatic Data Center.

The ASOS data may be useful for aiding in the characterization of visibility conditions in urban and suburban areas across the country. It also may be useful in future analyses to better understand the effects of fine PM on visibility in Class I areas. The Agency is in the process of analyzing ASOS data for a limited number of sites to determine how well it correlates to particulate matter monitoring results. In addition, the analysis will evaluate annual averages and seasonal, monthly, and daily visibility conditions (U.S. Environmental Protection Agency, 2001). The Agency expects that the results of these analyses will be available for inclusion in the final PM AQCD.

The largest monitoring network that includes both visibility and aerosol conditions is the Interagency Monitoring of Protected Visual Environments (IMPROVE) network. This network was formed in 1987 as a collaborative effort between Federal, regional, and state organizations responsible for protection of visibility in the 156 mandatory Class I Federal areas (national parks and wilderness areas) and other areas of interest to land management agencies, states, tribes, and other organizations (National Park Service, 1998; U.S. Environmental Protection Agency, 1996a; U.S. Environmental Protection Agency, 1995b; Eldred et al., 1997; Perry et al., 1997; Sisler and Malm, 2000; U.S. Environmental Protection Agency, 1999b). It is predominantly a rural-based network, with more than 140 sites across the country. The primary monitoring objectives of the IMPROVE program are to document current visibility conditions in the mandatory Class I areas, identify anthropogenic chemical species and emission sources of visibility impairment through the collection of speciated $PM_{2.5}$ data, and document long-term trends for assessing progress

towards elimination of anthropogenic visibility impairment. The IMPROVE network has also been involved in visibility related research, including the advancement of visibility monitoring instrumentation and analysis techniques and visibility monitoring and source attribution field studies (National Park Service, 1998; Evans and Pitchford, 1991).

Visibility monitoring under the IMPROVE network can be divided into three categories: aerosol, optical, and scene. Twenty-four hour $PM_{2.5}$ and PM_{10} aerosol samples are collected at least every third day utilizing filter-based aerosol technology. The $PM_{2.5}$ samples are analyzed to determined the mass concentration of the major particulate constituents (sulfates, nitrates, organic carbon compounds, elemental carbon, chlorides, and crustal elements) and for elements that indicate sources of visibility-impairing particles (trace elements and ions). Optical monitoring provides a direct measurement of light scattering and absorption. Color photographic imaging documents the appearance of the scene under a variety of air quality and illumination conditions (U.S. Environmental Protection Agency, 1999b). It is anticipated that all data generated by the IMPROVE network will be added to the AIRS database.

The U.S. Environmental Protection Agency has deployed a new national monitoring network (Federal Reference Method Monitoring network) designed to assess $PM_{2.5}$ concentrations and composition. As of early 2001, 1,108 monitoring sites were in operation (including more than 250 urban sites) and 1,044 sites had reported data to the Aerometric Informational Retrieval System (AIRS). Analyses of these data are expected to provide a more complete understanding of visibility conditions, in particular urban visibility, across the country. The $PM_{2.5}$ monitoring effort has been coordinated with visibility monitoring efforts currently in place to maximize the benefits of all of the monitoring programs (U.S. Environmental Protection Agency, 1997b; U.S. Environmental Protection Agency, 2000b; U. S. Environmental Protection Agency, 2001).

The Northeast States for Coordinated Air Use Management (NESCAUM) has established a real-time visibility monitoring network (CAMNET) using digital photographic imaging. Currently, there are digital photographic imaging for five urban locations (Boston, MA; Burlington, VT; Hartford, CT; Newark, NJ; and New York City, NY), and two rural locations (Acadia National Park, ME and Mt. Washington, NH). The visibility images are updated every 15 minutes. Near real-time air pollution and meteorological data are updated every hour. Archived images will be available to understand the visual effects of particulate matter air

pollution in the Northeast. CAMNET may be accessed at www.hazecam.net (Northeast States for Coordinated Air Use Management, 2002; Leslie, 2001).

4.3.6 Visibility Impairment: Trends and Current Conditions

In the United States, visibility impairment is caused by particles primarily composed of sulfates, nitrates, organic compounds, carbon soot, and crustal dust. Generally, sulfate is the major component responsible for visibility impairment in the eastern United States. However, nitrates, organic compounds, carbon soot, and crustal material are significant contributors to visibility impairment in some locations (Sisler and Cahill, 1993).

Trends in Visibility Impairment

Trends in visibility impairment or haziness often are used as indicators of trends in fine particles. Observations of visual range, obtained by the National Weather Service and available through the National Climatic Data Center of the National Oceanic and Atmospheric Administration, provide one of the few truly long-term, daily records of impairment related to air pollution. After some manipulation, including correction for relative humidity effects, the visual range data can be used as an indicator of fine mode particle pollution. The data reduction process and analyses of resulting trends have been reported by Schichtel et al. (2001), Husar et al. (1994), Husar and Wilson (1993), and Husar et al. (1981).

There are many statistical approaches to estimating trends. These approaches include simple correlation and regression analyses, time series analyses, and methods based on non-parametric statistics. A discussion and comparison of the methods for the detection of linear trends is provided in Hess et al. (2001). Schimek (1981) provides a discussion of nonlinear trends. In its annual air quality trends report, the U. S. Environmental Protection Agency characterized trends using a non-parametric regression analysis approach commonly referred to as the Theil test (U.S. Environmental Protection Agency, 1998; Hollander and Wolfe, 1973).

Generally, visibility impairment is greatest in the eastern United States and southern California. Visibility impairment or haziness in the southeastern United States, caused largely by sulfate formed from SO_2, is greatest in the humid summer months because of the ability of sulfate to absorb atmospheric water vapor, followed by the spring and fall, and winter. Summer haziness increased in the southeastern United States from the 1950s to 1980 along with

increasing SO_2 emissions. A decrease in haziness between 1980 and 1995 corresponded with a similar decrease in sulfur emissions (Schichtel et al., 2001). A statistically significant increase in summer sulfate concentrations was noted in two Class I areas in the eastern United States (Shenandoah and the Great Smoky Mountains) from 1982 to 1992 (Eldred et al., 1993; Cahill et al., 1996). During that time period, the majority of the Southwest showed decreasing sulfur (Eldred et al., 1993; Eldred and Cahill, 1994). The increasing sulfate concentrations were later shown to correlate with an increased trend in hazy days at those two locations (Iyer et al., 2000).

The U.S. Environmental Protection Agency's National Air Quality and Emission Trends Report summarized the regional trends and current conditions in 35 Class I areas and one urban area (Washington, DC) using data from the IMPROVE network (U.S. Environmental Protection Agency, 2001). The visibility trends analysis is an aggregate of 10 eastern Class I areas and 26 western Class I areas. Trends were presented for annual average values for the clearest ("best") 20% , middle ("typical") 20%, and haziest ("worst") 20% of the days monitored each year. The visibility trends, given in changes in deciview values, for the eastern and western sites are illustrated in Figures 4-29a and 4-29b. From the figures it can be seen that the haziest days in the West are equivalent to the best days in the East. In the East there was a 16% (1.5 deciview) improvement in haziness on the clearest days since 1992. Improvements in visibility were noted in the East for the haziest days; however, based on monitoring data for 1999, visibility remains significantly impaired with a visual range of 23 km for the haziest days compared to a mean visual range of 84 km for the clearest days. A 25 and14% improvement in visibility impairment was seen for the clearest and middle days in the West, respectively. Conditions for the haziest days degraded by 18.5% (1.7 deciviews) between 1997 and 1999, but were relatively unchanged compared to 1990 conditions (U.S. Environmental Protection Agency, 2001).

Figures 4-30a and 4-30b illustrate aggregate trends in aerosol light extinction, including trends by major aerosol component for the haziest 20% of days monitored for the 10 eastern Class I areas from 1992 to 1999 and the haziest 20% of days monitored for the 26 western Class I areas from 1990 to 1999. The report also includes a number of maps characterizing aerosol light extinction and key components at 36 IMPROVE sites (all rural except Washington, DC) for 1997 through 1999 (U.S. Environmental Protection Agency, 2001).



Figure 4-29a. Aggregate visibility trends (in deciviews) for 10 eastern Class 1 areas.



Figure 4-29b. Aggregate visibility trends (in deciviews) for 26 western Class 1 areas.

Source: U.S. Environmental Protection Agency (2001).



Figure 4-30a. Eastern class I area aggregate trends in aerosol light extinction on the 20% haziest days, including trends by major aerosol component.



Figure 4-30b. Western class I area aggregate trends in aerosol light extinction on the 20% haziest days, including trends by major aerosol component.

Source: U.S. Environmental Protection Agency (2001).

Current Conditions

Current visibility conditions have been well-characterized for Class I areas using updated data from the IMPROVE network (U.S. Environmental Protection Agency, 2001; Malm et al., 2000; IMPROVE, 1998). During recent decades, daytime visibility conditions at all major airports throughout the United States were recorded hourly by human observation. These data were used to determine current visibility conditions and visibility trends in the United States, as well as the spatial distribution of visibility conditions (Trijonis et al., 1991). The use of airport human observation is being replaced by an automated observing system, Automated Surface Observing System (ASOS). More than 900 airports are currently commissioned. Additionally, the U.S. Environmental Protection Agency has deployed a new national monitoring network to assess $PM_{2.5}$ concentrations and composition. Visibility conditions for urban and suburban areas will become more widely available as data from the national $PM_{2.5}$ speciation monitoring network and the ASOS airport visibility network are further analyzed.

4.4 EFFECTS ON MATERIALS

Effects of air pollution on materials are related to both aesthetic appeal and physical damage. Studies have demonstrated that particles, primarily consisting of carbonaceous compounds, cause soiling of commonly used building materials and culturally important items, such as statutes and works of art. Physical damage from the dry deposition of air pollutants, such as PM (especially sulfates and nitrates) and SO_2, and the absorption or adsorption of corrosive agents on deposited particles also can result in the acceleration of naturally occurring weathering processes of man-made building and cultural materials.

In the atmosphere, PM may be “primary,” existing in the same form in which it was emitted, or “secondary,” formed by the chemical reactions of free, absorbed, or dissolved gases. The major constituents of atmospheric PM are sulfate, nitrate, ammonium, and hydrogen ions; particle-bound water; elemental carbon; a great variety of organic compounds; and crustal material. A substantial fraction of the fine particle mass, particularly during the warmer months, is secondary sulfate and nitrate. Sulfates may be formed by the gas-phase conversion of SO_2 to H_2SO_4 by OH radicals and aqueous-phase reactions of SO_2 with H_2O_2, O_3, or O_2. During the day, NO_2 may be converted to nitric acid (HNO_3) by reacting with OH radicals. Nitrogen dioxide also

can be oxidized to HNO_3 by a sequence of reactions initiated by O_3. A more detailed discussion of the atmospheric chemistry of PM appears in Chapter 2 of this document.

4.4.1 Corrosive Effects of Particles and Sulfur Dioxide on Man-Made Surfaces

Limited new studies have been published that better define the role of air pollution in materials damage. This section briefly summarizes information on exposure particle related effects on materials and sulfur-containing pollutants (formed by the chemical reactions of SO_2 with other atmospheric pollutants) addressed in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) and presents relevant information published since completion of that document. The effects of nitrates on man-made building materials and naturally occurring cultural materials was discussed in the earlier EPA Nitrogen Oxides Criteria Document (U.S. Environmental Protection Agency, 1993).

4.4.1.1 Metals

Metals undergo natural weathering processes in the absence of environmental pollutants. The additive effect of pollutants on the natural weathering processes depend on the nature of the pollutant, and the deposition rate (the uptake of a pollutant by the material's surface), and the presence of moisture. The influence of the metal-protective corrosion film, the presence of other surface electrolytes, the orientation of the metal surface, the presence of surface moisture, and the variability in the electrochemical reactions will also contribute to the effect of pollutant exposure on metal surfaces.

Several studies demonstrate the importance of the duration of surface wetness (caused by dew and fog condensation and rain) on metals. Surface moisture facilitates the deposition of pollutants, especially SO_2, and promotes corrosive electrochemical reactions on metals (Haynie and Upham, 1974; Sydberger and Ericsson, 1977). Of critical importance is the formation of hygroscopic salts on the metal that increases the duration of surface wetness and, thereby, enhances the corrosion process.

Pitchford and McMurry (1994) and Zhang et al. (1993) demonstrated particle size-related effects of relative humidity. The effect of temperature on the rate of corrosion is complex. Under normal temperature conditions, temperature would not have an affect on the rate of

corrosion; but when the temperature decreases, the relative humidity increases and the diffusivity decreases. The corrosion rate decreases as the temperature approaches freezing because ice prohibits the diffusion of SO_2 to the metal surface and minimizes electrochemical processes (Haynie, 1980; Biefer, 1981; Sereda, 1974).

The metal protective corrosion film (i.e., the rust layer on metal surfaces) provides some protection against further corrosion. The effectiveness of the corrosion film in slowing down the corrosion process is affected by the solubility of the corrosion layer and the concentration and deposition rate of pollutants. If the metal-protective corrosion film is insoluble, it may add some protection against acidic pollutants. An atmospheric corrosion model that considers the formation and dissolution of the corrosion film on galvanized steel was proposed by Spence et al. (1992). The model considers the effects of SO_2, rain acidity, and the time of wetness on the rate of corrosion. Although the model does not specifically characterize particle effects, the contribution of particulate sulfate was considered in model development.

Whether suspended particles actually impact on the corrosion of metals is not clear. Several studies suggest that suspended particles will promote the corrosion of metals (Goodwin et al., 1969; Barton, 1958; Sanyal and Singhania, 1956; Baedecker et al., 1991); however, other studies have not demonstrated a correlation between particle exposure and metal corrosion (Mansfeld, 1980; Edney et al., 1989). Walton et al. (1982) suggested that catalytic species within several species in fly ash promote the oxidation of SO_x to a corrosive state. Still other researchers indicate that the catalytic effect of particles is not significant and that the corrosion rate is dependent on the conductance of the thin-film surface electrolytes during periods of wetness. Soluble particles likely increase the solution conductance (Skerry et al., 1988; Askey et al., 1993).

The corrosion of most ferrous metals (iron, steel, and steel alloys) is increased by increasing SO_2 exposure. Steels are susceptible to corrosion when exposed to SO_2 in the absence of protective organic or metallic coatings. Studies on the corrosive effects of SO_2 on steel indicate that the rate of corrosion increases with increasing SO_2 and is dependent on the deposition rate of the SO_2 (Baedecker et al., 1991; Butlin et al., 1992a). The corrosive effects of SO_2 on aluminum is exposure-dependent, but appears to be insignificant (Haynie, 1976; Fink et al., 1971; Butlin et al., 1992a). The rate of formation of the patina (protective covering) on copper can take as long as 5 years and is dependent on the SO_2 concentration, deposition rate,

temperature, and relative humidity (Simpson and Horrobin, 1970). Further corrosion is controlled by the availability of copper to react with deposited pollutants (Graedel et al., 1987). Butlin et al. (1992a), Baedecker et al. (1991), and Cramer et al. (1989) reported an average corrosion rate of 1 μm/year for copper; however, less than a third of the corrosion was attributed to SO_2 exposure, suggesting that the rate of patina formation was more dependent on factors other than SO_2. A recent report by Strandberg and Johansson (1997) showed relative humidity to be the primary factor in copper corrosion and patina formation. The results of the studies on particles and SO_2 corrosion of metals are summarized in Table 4-18.

4.4.1.2 Painted Finishes

Exposure to air pollutants affect the durability of paint finishes by promoting discoloration, chalking, loss of gloss, erosion, blistering, and peeling. Evidence exists that indicates particles can damage painted finishes by serving as carriers for corrosive pollutants (Cowling and Roberts, 1954) or by staining and pitting of the painted surfaces (Fochtman and Langer, 1957; Wolff et al., 1990).

The erosion rate of oil-based house paint has been reported to be enhanced by exposure to SO_2 and high humidity. In a study by Spence et al. (1975), an erosion rate of 36.71 ± 8.03 μm/year was noted for oil-based house paint samples exposed to SO_2 (78.6 $\mu g/m^3$), O_3 (156.8 $\mu g/m^3$), and NO_2 (94 $\mu g/m^3$) and low humidity (50%). The erosion rate increased with increased SO_2 and humidity. The authors concluded that SO_2 and humidity accounted for 61% of the erosion. Acrylic coil coating and vinyl coil coating shows less pollutant-related erosion. Erosion rates range from 0.7 to 1.3 μm/year and 1.4 to 5.3 μm/year, respectively. Similar findings on SO_2-related erosion of oil-based house paints and coil coatings have been reported by other researchers (Davis et al., 1990; Yocom and Grappone, 1976; Yocom and Upham, 1977; Campbell et al., 1974). Several studies suggest that the effect of SO_2 is caused by its reaction with extender pigments such as calcium carbonate and zinc oxide (Campbell et al., 1974; Xu and Balik, 1989; Edney, 1989; Edney et al., 1988, 1989). However, Miller et al. (1992) suggested that calcium carbonate acts to protect paint substrates. Another study indicated that exposure to SO_2 can increase the drying time of some paints by reacting with certain drying oils and will compete with the auto-oxidative curing mechanism responsible for crosslinking the binder (Holbrow, 1962).

TABLE 4-18. CORROSIVE EFFECTS OF PARTICULATE MATTER AND SULFUR DIOXIDE ON METALS

Metal	Exposure Conditions	Comments	Source
Mild Steel Galvanized Steel	Specimens exposed to SO_2 and O_3 under natural and artificial conditions, and to NO_2 under natural conditions. SO_2 concentrations ranged from 2.1 to 60 $\mu g/m^3$. Annual average concentrations were about 20 $\mu g/m^3$. Meteorological conditions were unaltered. Specimens exposed at 29 sites for 2 years for mild steel and 1 y for galvanized steel.	Steel corrosion was dependent on long-term SO_2 exposure. The corrosion rate was about 50 μm/year for mild steel specimens for most industrial sites, but ranged from 21 to 71 μm/year. The corrosion rate ranged from 1.45 to 4.25 μm/year for galvanized steel. The authors concluded that rainfall also may have a significant effect on galvanized steel based on a corrosion rate of 3.4 μm/year seen at a very wet site.	Butlin et al. (1992a)
Zinc	Rolled zinc specimens exposed at various sites around the country (rural, industrialized, marine) for up to 20 years. Actual pollutant exposures not reported.	The highest corrosion rates were associated with industrialized environments and marine environments in direct contact with salt spray.	Showak and Dunbar (1982)
Zinc	Specimens exposed at 5 sites for 1 to 5 years. Average SO_2 concentrations ranged from 2 ± 4 to 15 ± 17 ppb (5.2 ± 10.4 to 39.3 ± 44.5 $\mu g/m^3$). PM concentrations ranged from 14 to 60 $\mu g/m^3$. Highest pollutant concentrations recorded at 1 year exposure site.	Average corrosion rate ranged from 0.63 to 1.33 μm/y. The highest corrosion was noted in the most industrialized area. However, the corrosion rates did not differ significant regardless of the SO_2 concentration, suggesting that SO_2 exposure may not be the dominant factor in zinc corrosion.	Baedecker et al. (1991) Cramer et al. (1989)
Carbon Steel Weathering Steel	See Baedecker et al. (1991) above for exposure conditions.	Average corrosion rate for samples exposed for 5 years ranged from 6.6 to 12.8 μm/year for carbon steel and 3.7 to 5.0 μm/year for weathering steel. Highest corrosion rate noted for samples exposed for 1 year.	Baedecker et al. (1991) Cramer et al. (1989)
Aluminum	See Baedecker et al. (1991) above for exposure conditions.	Corrosion rate was very low at all sites and ranged from 0.036 to 0.106 μm/year.	Baedecker et al. (1991)
Aluminum	See Butlin et al. (1992a) above for exposure conditions.	Corrosion greater on the under side of specimens, possibly because of lack of washoff and increased PM in area. Maximum corrosion rate was 0.85 μm/year. Pit depths of up to 72 μm were noted after 2 years of exposure.	Butlin et al. (1992a)

DRAFT-DO NOT QUOTE OR CITE

TABLE 4-18 (cont'd). CORROSIVE EFFECTS OF PARTICULATE MATTER AND SULFUR DIOXIDE ON METALS

Metal	Exposure Conditions	Comments	Source
Copper	See Baedecker et al. (1991) above for exposure conditions.	Average corrosion rate for 3- and 5-year exposures was about 1 μm/year but the soluble portion was less than a third of that which could be contributed to SO_2 exposure. Dry deposition of SO_2 was not as important in patina formation as wet deposition of H^+.	Baedecker et al. (1991)
Copper	See Butlin et al. (1992a) above for exposure conditions.	Majority of test sites showed a corrosion rate of 1 ± 0.2 μm/year. The corrosion rate was 1.48 μm/year at the site receiving the most rainfall. The lowest corrosion rate, 0.66 μm/year, was associated with low rainfall, low SO_2.	Butlin et al. (1992a)
Copper	Specimens exposed to 4 to 69 ppb (10.4 to 180.7 $\mu g/m^3$) and 1.0 ppm (2,618.7 $\mu g/m^3$) SO_2 for 20h at various relative humidities.	SO_2 had no effect on copper when relative humidity was $<$75%. Increasing relative humidity increases patina formation in presence of trace SO_2. No SO_2-related effects were noted on copper specimens exposed to high SO_2 regardless of the percent relative humidity.	Strandberg and Johansson (1997)
Copper	Specimens exposed artificially to 0.49 ± 0.01 ppm (187 ± 3.8 $\mu g/m^3$) SO_2 for 4 weeks at 70 and 90% relative humidity.	Corrosive effect of SO_2 on copper increased with increasing relative humidity.	Eriksson et al. (1993)

DRAFT-DO NOT QUOTE OR CITE

4.4.1.3 Stone and Concrete

Numerous studies suggest that air pollutants can enhance the natural weathering processes on building stone. The development of crusts on stone monuments have been attributed to the interaction of the stone's surface with sulfur-containing pollutants, wet or dry deposition of atmospheric particles, and dry deposition of gypsum particles from the atmosphere. Because of a greater porosity and specific surface, mortars have a greater potential for reacting with environmental pollutants (Zappia et al., 1998). Details on these studies are discussed in Table 4-19. The stones most susceptible to the deteriorating effects of sulfur-containing pollutants are the calcareous stones (limestone, marble, and carbonated cement). Exposure-related damage to building stones result from the formation of salts in the stone that are subsequently washed away during rain events leaving the stone surface more susceptible to the effects of pollutants. Dry deposition of sulfur-containing pollutants promotes the formation of gypsum on the stone's surface. Gypsum is a gray to black crusty material comprised mainly of calcium sulfate dihydrate from the reaction of calcium carbonate (calcite) in the stone with atmospheric SO_2 and moisture (relative humidities exceeding 65%). Approximately 99% of the sulfur in gypsum is sulfate because of the sulfation process caused by the deposition of SO_2 aerosol. Sulfites also are present in the gypsum layer as an intermediate product (Sabbioni et al., 1996; Ghedini et al., 2000; Gobbi et al., 1998; Zappia et al., 1998). Gypsum is more soluble than calcite and is known to form on limestone, sandstones, and marble when exposed to SO_2. Gypsum also has been reported to form on granite stone by replacing silicate minerals with calcite (Schiavon et al., 1995). Gypsum occupies a larger volume than the original stone, causing the stone's surface to become cracked and pitted. The rough surface serves as a site for deposition of airborne particles.

The dark colored gypsum is caused by surface deposition of carbonaceous particles (noncarbonate carbon) from combustion processes occurring in the area (Sabbioni, 1995; Saiz-Jimenez, 1993; Ausset et al., 1998), trace metals contained in the stone, dust, and numerous other anthropogenic pollutants. After analyzing damaged layers of several stone monuments, Zappia et al. (1993) found that the dark-colored damaged surfaces contained 70% gypsum and 20% noncarbonate carbon. The lighter colored damaged layers were exposed to rain and contained 1% gypsum and 4% noncarbonate carbon. It is assumed that rain removes reaction products, permitting further pollutant attack of the stone monument, and likely redeposits some

TABLE 4-19. CORROSIVE EFFECTS OF PARTICULATE MATTER AND SULFUR DIOXIDE ON STONE

Stone	Exposure Conditions	Comments	Source
Vermont marble	Runoff water was analyzed from seven summer storms. SO_2 concentration stated to be low.	Between 10 to 50% of calcium in runoff water estimated from gypsum formation from dry deposition of SO_2.	Schuster et al. (1994)
Marble sandstone	Analysis of runoff water for five slabs test exposed to ambient conditions at a angle of 30° to horizontal.	Pollutant exposure related erosion was primarily caused by dry deposition of SO_2 and nitric acid between rain events and wet deposition of hydrogen ion. Recession estimates ranged from 15 to 30 μm/year for marble and 25 to 45 μm/year for limestone. A large portion of the erosion results from the reaction of CO_2 with the calcium in the stone.	Baedecker et al. (1992)
Limestone	Ambient air conditions. Exposure ranged from 70 to 1065 days. Averaged pollutant exposure ranged from 1.4 to 20.4 ppb (3.7 to 53.4 $\mu g/m^3$) SO_2; 4.1 to 41.1 ppb NO_x; 2.4 to 17.4 ppb (4.5 to 32.7 $\mu g/m^3$) NO_2; 10.1 to 25.6 ppb (19.8 to 50.2 $\mu g/m^3$) O_3.	Increased stone weight loss with increased SO_2. Rainfall did not significantly affect stone degradation. Stone loss associated with SO_2 exposure estimated to be 24 μm/year. Slight trend in decreasing stone loss with increasing length of exposure.	Webb et al. (1992)
Portland limestone White Mansfield dolomitic sandstone Monk's Park limestone	Experimental tablets exposed under sheltered and unsheltered ambient air conditions. Exposure for l and 2 years.	Significant correlations existed between the mean annual SO_2 concentration, rainfall volume, and hydrogen ion loading and the weight changes.	Butlin et al. (1992b)
Sandstones (calcite and noncalcite stones)	Ambient air; low concentrations of sulfates, SO_2, and nitrates; RH sufficient to produce condensation on stones rarely occurred.	Insignificant differences in erosion rate found between calcite and noncalcite sandstone. Moisture affected the rate of pollutant deposition and enhanced susceptibility to pollutant related erosion. Rain events given as primary factor affecting stone erosion. Pollutant related erosion judged to be insignificant.	Petuskey et al. (1995)
Limestones Sandstones Marble Granite Basalt	Ambient air; urban and rural locations in Mediterranean.	Crusts on stones were found to contain two layers; top layer, usually black in color, composed of gypsum between 40 and 400 μm thick. Innermost layer, ranging from brown to orange in color, primarily consisted of calcite, between 10 and 600 μm thick. Gypsum-rich layer thought to be the result of sulphation of the calcitic layer by atmospheric pollutants or dry or wet deposition of atmospheric dust.	Garcia-Vallès et al. (1998)
Portland limestone Massangis Jaune Roche limestone White Mansfield dolomitic	Samples exposed to SO_2, NO_2, and NO at 10 ppmv, both with and without O_3 and under dry (coming to equilibrium with the 84% RH) or wetted with CO_2-equilibrated deionized water conditions. Exposure was for 30 days.	In the absence of moisture, little reaction is seen. SO_2 is oxidized to sulfates in the presence of moisture. The effect is enhanced in the presence of O_3. Massangis Jaune Roche limestone was the least affected by the pollutant exposure. Crust lined pores of specimens exposed to SO_2.	Haneef et al. (1993)

DRAFT-DO NOT QUOTE OR CITE

TABLE 4-19 (cont'd). CORROSIVE EFFECTS OF PARTICULATE MATTER AND SULFUR DIOXIDE ON STONE

Stone	Exposure Conditions	Comments	Source
Monk's Park Portland limestone	Samples exposed for 2 mo under both sheltered and unsheltered conditions. Mean daily atmospheric SO_2 concentration was 68.7 $\mu g/m^3$ and several heavy rainfalls.	Significant amounts of gypsum were noted on the Portland stone. Sheltered stones also showed soiling by carbonaceous particles and other combustion products. Etch holes and deep etching was noted in some of the exposed unsheltered samples.	Viles (1990)
Carrara marble Travertine Tranistone	Sample exposed in laboratory to 3 ppm SO_2 and 95% RH at 25 °C for 150 days. Samples were coated with three carbonaceous particle samples from combustion sources, and with activated carbon and graphite.	Exposure to particles from combustion processes enhanced sulfation of calcareous materials by SO_2 because of metal content of particles.	Sabbioni et al. (1996)
Carrara marble Georgia marble	Samples exposed in sheltered ambient environment for 6, 12, or 20 mo.	Carrara marble found to be more reactive with SO_2 than Georgia marble possibly because of the compactness of the Georgia marble. Greater effects noted when samples were also exposed to NO_2.	Yerrapragada et al. (1994)
Carrara marble	Samples exposed for 6 mo (cold and hot conditions) in ambient environment. PM concentrations ranged from 57.3 to 116.7 $\mu g/m^3$ (site 1) and 88 to 189.8 $\mu g/m^3$ (site 2). Some exposures also were associated with high SO_2, NO, and NO_2.	Pollutant exposed samples showed increased weight gain over that expected from natural weathering processes. There was a blackening of stone samples exposed to carbonaceous rich particulate matter.	Realini et al. (1995)
Monk's Park limestone Portland limestone	Samples artificially exposed to fly-ash containing 1309.3 $\mu g/m^3$ SO_2 (0.5 ppm), at 95% RH and 25 °C for 81 or 140 days. Fly-ash samples from five different sources were used in study.	Exposure to fly-ash did not enhance oxidation of SO_2 to sulfates. Mineral oxides in fly ash contributed to sulphation of $CaCO_3$.	Hutchinson et al. (1992)
Lime mortar Pozzolan mortar Cement mortar	Samples exposed to 7,856 $\mu g/m^3$ (3 ppm) SO_2 at 100% RH and 25 °C for 30, 60, or 90 days; samples sprayed with bidistilled water every 7 days to simulate rainfall.	Exposure to SO_2 produced significant quantities of calcium sulfite and calcium sulfate on specimens; however, the amount produced was dependent of the porosity, specific surface, and alkalinity of the sample.	Zappia et al. (1994)
Limestone Travertine marble	Samples exposed under actual ambient air conditions at two locations in Rome. Monitoring data obtained for SO_2, NO, NO_2, and total suspended particulates (TSP) but not reported. Exposure was for four seasons.	TSP exposure increased the cleaning frequency for stone monuments. Monuments are soiled proportionately overtime, based on brightness values. Horizontal surfaces showed higher graying values because of particle sediment.	Lorusso et al. (1997)

TABLE 4-19 (cont'd). CORROSIVE EFFECTS OF PARTICULATE MATTER AND SULFUR DIOXIDE ON STONE

Stone	Exposure Conditions	Comments	Source
Limestone Quartz-cemented sandstone Calcite-cemented sandstone Granite Brick	Samples from structures exposed for varying periods of time under ambient air conditions. Samples selected because of black layer on surface.	Black layers were found to be primarily comprised of iron compounds, quartz, silicate, soot, and dirt.	Nord and Ericsson (1993)
Limestone Sandstone	Samples of ancient grey crust formed between 1180 and 1636 on the Church of Saint Trophime in Arks and formed between 1530 and 1187 on the Palazz d'Accursio in Bolonga.	Crust samples contained calcite, soil dust, carbonaceous particles, and gypsum crystals.	Ausset et al. (1998)
Carrara marble Travertine marble Trani limestone Portland limestone Lime mortar Pozzolan mortar Cement mortar	Samples of the stones and mortars were representative of those used in the past and currently for new construction and restorations. Samples were exposed for 6, 12, and 24 mo under ambient conditions in Milan.	Mortars were more reactive than the stones. Of the mortars, cement and pozzolan mortar were more reactive than the lime mortar. Carrara marble was the least reactive of the stones. The maximum amount of degradation was found in areas sheltered from rain.	Zappia et al. (1998)
Lime mortar	Sample of black crust taken from Zamboni Tower Gate.	Exposure to environmental pollutants caused the formation of two separate layers on the mortar: an outer thin surface black crust composed of gypsum and carbonaceous particles and the inner composed of products from the dissolution and sulphation of the carbonate matrix in the mortar.	Sabbioni et al. (1998)
Carrara marble	Samples of crust removed from Milan General Hospital, built around 1937.	Gypsum main component of crust followed by carbonaceous particles and iron oxides. Estimated rate of crust formation was 2-5 μm/year. Total amount of gypsum formed over the lifetime of exposure was 5 to 13 mg/cm^2, an estimated 0.2 mg/cm^2/year.	Bugini et al. (2000)

DRAFT-DO NOT QUOTE OR CITE

of the reaction products at rain runoff sites on the stone. Following sulfur compounds, carbon was reported to be the next highest element in dark crust on historical monuments in Rome. Elemental carbon and organic carbon accounted for 8 and 39% of the total carbon in the black crust samples. The highest percentage of carbon, carbonate carbon, was caused by the carbonate matrix in the stones. The high ratio of organic carbon to elemental carbon indicates the presence of a carbon source other than combustion processes (Ghedini et al., 2000). Cooke and Gibbs (1994) suggested that stones damaged during times of higher ambient pollution exposure likely would continue to exhibit a higher rate of decay, termed the "memory effect," than newer stones exposed under lower pollution conditions. Increased stone damage also has been associated with the presence of sulfur oxidizing bacteria and fungi on stone surfaces (Garcia-Vallès et al., 1998; Young, 1996; Saiz-Jimenez, 1993; Diakumaku et al., 1995).

Dissolution of gypsum on the stone's surface initiates structural changes in the crust layer. Garica-Vallès et al. (1998) proposed a double mechanism: the dissolution of the gypsum, in the presence of sufficient moisture, followed by recrystallization inside fissures or pores. In the event of limited moisture, the gypsum is dissolved and recrystallizes at its original location. According to the authors, this would explain the gypsum-rich crustal materials on stone surfaces sheltered from precipitation.

Moisture was found to be the dominant factor in stone deterioration for several sandstones (Petuskey et al., 1995). Dolske (1995) reported that the deteriorative effects of sulfur-containing rain events, sulfates, and SO_2 on marble were largely dependent on the shape of the monument or structure rather than the type of marble. The author attributed the increased fluid turbulence over a nonflat vertical surface versus a flat surface to the increased erosion. Sulfur-containing particles also have been reported to enhance the reactivity of Carrara marble and Travertine and Trani stone to SO_2 (Sabbioni et al., 1992). Particles with the highest carbon content had the lowest reactivity.

The rate of stone deterioration is determined by the pollutant and the pollutant concentration, the stone's permeability and moisture content, and the pollutant deposition velocity. Dry deposition of SO_2 between rain events has been reported to be a major causative factor in pollutant-related erosion of calcareous stones (Baedecker et al., 1991; Dolske, 1995; Cooke and Gibbs, 1994; Schuster et al., 1994; Hamilton et al., 1995; Webb et al., 1992). Sulfur dioxide deposition increases with increasing relative humidity (Spiker et al., 1992), but the

pollutant deposition velocity is dependent on the stone type (Wittenburg and Dannecker, 1992), the porosity of the stone, and the presence of hygroscopic contaminants.

Although it is clear from the available information that gaseous pollutants, in particular dry deposition of SO_2, will promote the decay of some types of stones under the specific conditions, carboneous particles (noncarbonate carbon) may help to promote the decay process by aiding in the transformation of SO_2 to a more acidic species (Del Monte and Vittori, 1985). Several authors have reported enhanced sulfation of calcareous material by SO_2 in the presence of particles containing metal oxides (Sabbioni et al., 1996; Hutchinson et al., 1992).

4.4.2 Soiling and Discoloration of Man-Made Surfaces

Ambient particles can cause soiling of man-made surfaces. Soiling has been defined as the deposition of particles of less than 10 μm on surfaces by impingement. Soiling generally is considered an optical effect, that is, soiling changes the reflectance from opaque materials and reduces the transmissions of light through transparent materials. Soiling can represent a significant detrimental effect requiring increased frequency of cleaning of glass windows and concrete structures, washing and repainting of structures, and, in some cases, reduction in the useful life of the object. Particles, in particular carbon, also may help catalyze chemical reactions that result in the deterioration of materials during exposure.

It is difficult to determine the accumulated particle levels that cause an increase in soiling; however, soiling is dependent on the particle concentration in the ambient environment, particle size distribution, and the deposition rate and the horizontal or vertical orientation and texture of the surface being exposed (Haynie, 1986). The chemical composition and morphology of the particles and the optical properties of the surface being soiled will determine the time at which soiling is perceived (Nazaroff and Cass, 1991). Carey (1959) reported that the average observer could observe a 0.2% surface coverage of black particles on a white background. A recent study suggests that it would take a 12% surface coverage by black particles before there is 100% accuracy in identifying soiling (Bellan et al., 2000). The rate at which an object is soiled increases linearly with time; however, as the soiling level increases, the rate of soiling decreases. The buildup of particles on a horizontal surface is counterbalanced by an equal and opposite depletion process. The depletion process is based on the scouring and washing effect of wind and rain (Schwar, 1998).

4.4.2.1 Stones and Concrete

Most of the research evaluating the effects of air pollutants on stone structures have concentrated on gaseous pollutants. The deposition of the sulfur-containing pollutants are associated with the formation of gypsum on the stone (see Section 4.4.1.3). The dark color of gypsum is attributed to soiling by carbonaceous particles from nearby combustion processes. A lighter gray colored crust is attributed to soil dust and metal deposits (Ausset et al., 1998; Camuffo, 1995; Moropoulou et al., 1998). Realini et al. (1995) found the formation of a dark gypsum layer and a loss of luminous reflection in Carrara marble structures exposed for 1 year under ambient air conditions. Dark areas of gypsum were found by McGee and Mossitti (1992) on limestone and marble specimens exposed under ambient air conditions for several years. The black layers of gypsum were located in areas shielded from rainfall. Particles of dirt were concentrated around the edges of the gypsum formations. Lorusso et al. (1997) attributed the need for frequent cleaning and restoration of historic monuments in Rome to exposure to total suspended particulates. They also concluded that, based on a decrease in brightness (graying), surfaces are soiled proportionately over time; however, graying is higher on horizontal surfaces because of sedimented particles. Davidson et al. (2000) evaluated the effects of air pollution exposure on a limestone structure on the University of Pittsburgh campus using estimated average TSP levels in the 1930s and 1940s and actual values for the years 1957 to 1997. Monitored levels of SO_2 were available for the years 1980 to 1998. Based on the available data on pollutant levels and photographs, it was thought that soiling began while the structure was under construction. With decreasing levels of pollution, the soiled areas have been slowly washed away, the process taking several decades, leaving a white, eroded surface. Studies describing the effects of particles on stone surfaces are discussed in Table 4-9.

4.4.2.2 Household and Industrial Paints

Few studies are available that evaluate the soiling effects of particles on painted surfaces. Particles composed of elemental carbon, tarry acids, and various other constituents are responsible for soiling of structural painted surfaces. Coarse-mode particles (>2.5 μm) initially contribute more soiling of horizontal and vertical painted surfaces than do fine-mode particles (<2.5 μm), but are more easily removed by rain (Haynie and Lemmons, 1990). The accumulation of fine particles likely promotes remedial action (i.e., cleaning of the painted

surfaces). Coarse-mode particles are primarily responsible for soiling of horizontal surfaces. Rain interacts with coarse particles, dissolving the particle and leaving stains on the painted surface (Creighton et al., 1990; Haynie and Lemmons, 1990). Haynie and Lemmons (1990) proposed empirical predictive equations for changes in surface reflectance of gloss-painted surfaces that were exposed protected and unprotected from rain and oriented horizontally and vertically.

Early studies by Parker (1955) and Spence and Haynie (1972) demonstrated an association between particle exposure and increased frequency of cleaning of painted surfaces. Particle exposures also caused physical damage to the painted surface (Parker, 1955). Unsheltered painted surfaces are initially more soiled by particles than sheltered surfaces but the effect is reduced by rain washing. Reflectivity is decreased more rapidly on glossy paint than on flat paint (Haynie and Lemmons, 1990). However, surface chalking of the flat paint was reported during the exposure. The chalking interfered with the reflectance measurements for particle soiling. Particle composition measurements that were taken during exposure of the painted surfaces indicated sulfates to be a large fraction of the fine mode and only a small fraction of the coarse mode. Although no direct measurements were taken, fine mode particles likely also contained large amounts of carbon and possibly nitrogen or hydrogen (Haynie and Lemmons, 1990).

4.5 EFFECTS OF ATMOSPHERIC PARTICULATE MATTER ON GLOBAL CLIMATE CHANGE PROCESSES AND THEIR POTENTIAL HUMAN HEALTH AND ENVIRONMENTAL IMPACTS

Processes causing global climate change and their potential environmental and human health impacts have been accorded extensive attention during the past several decades, and they still continue to be of broad national and international concern. This is reflected by extensive research and assessment efforts undertaken since the mid-1970s by U.S. Federal Government Agencies (e.g., NOAA, EPA, CDC, etc.) or via U.S. Federal Interagency programs (e.g., the U.S. Global Climate Change Research Program [USGCRP]). It is also reflected by analogous extensive research and assessment efforts undertaken by numerous other national governments or international collaborative activities, e.g., those coordinated by the Intergovernmental Panel on

Climate Change (IPCC), established in the 1980s under the joint auspices of the World Meteorological Organization (WMO) and the United Nations Environment Programme (UNEP).

Atmospheric particles play important roles in two key types of global climate change processes: (1) alterations in the amount of ultraviolet solar radiation (especially UV-B) penetrating through the Earth's atmosphere and reaching its surface, where it can exert a variety of effects on human health, plant and animal biota, and other environmental components; and (2) alterations in the amount of visible solar radiation transmitted through the Earth's atmosphere. Particles both absorb and reflect solar radiation back into space. The absorption of solar radiation by particles, together with trapping of infrared radiation emitted by the Earth's surface by certain gases, enhances heating of the Earth's surface and lower atmosphere (i.e., the widely-known "greenhouse effect") and leads to consequent "global warming" impacts on human health and the environment. Atmospheric particles also play a lesser role by absorbing infrared radiation emitted by the Earth's surface.

The effects of atmospheric PM on the transmission of electromagnetic radiation emitted by the sun at ultraviolet and visible wavelengths and by the earth at infrared wavelengths depend on radiative properties (extinction efficiency, single scattering albedo, and asymmetry parameter) of the particles, which depend, in turn, on the size and shape of the particles, the composition of the particles, and the distribution of components within individual particles. In general, the radiative properties of particles are size- and wavelength-dependent. In addition, the extinction cross-section tends to be at a maximum when the particle radius is similar to the wavelength of the incident radiation. Thus, fine particles present mainly in the accumulation mode would be expected to exert a greater influence on the transmission of electromagnetic radiation than would coarse particles. The composition of particles can be crudely summarized in terms of the broad classes identified in Chapter 2 of this document. These include fine particles consisting mainly of (a) nitrate, sulfate, mineral dust, elemental carbon, organic carbon compounds (e.g., PAHs), and (b) metals derived from high temperature combustion or smelting processes. The major sources of these components are shown in Table 3-9 of Chapter 3 in this document.

Knowledge of factors controlling the transfer of solar radiation in the ultraviolet spectral range is needed for assessing potential biological and environmental impacts associated with exposure to UV-B radiation (290 to 315 nm). Knowledge of the effects of PM on the transfer of radiation in the visible and infrared spectral regions is needed for assessing the relationship

between particles and global warming and its environmental and biological impacts. Important conceptual aspects and factors related to solar ultraviolet radiation processes and effects are first summarized below and atmospheric PM roles discussed. This is followed by a summary of global warming processes, their potential human health and environmental impacts, and their potential relationships to atmospheric PM.

4.5.1 Solar Ultraviolet Radiation Transmission Impacts on Human Health and the Environment: Atmospheric Particulate Matter Effects

4.5.1.1 Potential Effects of Increased Ultraviolet Radiation Transmission

The transmission of solar UV-B radiation through the earth's atmosphere is controlled by ozone, clouds, and particles. The depletion of stratospheric ozone caused by the release of anthropogenically produced chlorine (Cl)-and bromine (Br)-containing compounds has resulted in heightened concern about potentially serious increases in the amount of solar UV-B radiation (SUVB) reaching the Earth's surface. SUVB is also responsible for initiating the production of OH radicals that oxidize a wide variety of volatile organic compounds, some of which can deplete stratospheric ozone (e.g., CH_3Cl, CH_3Br), absorb terrestrial infrared radiation (e.g., CH_4), and contribute to photochemical smog formation (e.g., C_2H_4 , C_5H_8).

Increased penetration of SUVB to the Earth's surface as the result of stratospheric ozone depletion continues to be of much concern because of projections of consequent increased surface-level SUVB exposure and associated potential negative impacts on human health, plant and animal biota, and man-made materials. Several summary overviews (Kripke, 1989; Grant, 1989; Kodama and Lee, 1994; Van der Leun et al., 1995, 1998) of salient points related to stratospheric ozone depletion and bases for concern provide a concise introduction to the subject, as does Figure 4-31. As shown to the left in Figure 4-31, stratophospheric ozone depletion results from: (a) anthropogenic emissions of certain trace gases having long atmospheric residence times, e.g., chlorofluorocarbons (CFCs), carbon tetrachloride (CCl_4), and Halon 1211 (CF_2Cl Br) and 1301 (CF_3Br)—which have atmospheric residence times of 75 to 100 years, 50 years, 25 years, and 110 years, respectively; (b) their tropospheric accumulation and gradual transport, over decades, up to the stratosphere, where (c) they photolyze to release Cl and Br that catalyze ozone destruction; leading to (d) stratospheric ozone depletion. Such ozone depletion is most marked over Antarctica during spring in the Southern Hemisphere, to a less marked but still



Figure 4-31. Processes involved in stratospheric ozone depletion because of man's production of CFCs, halons, and other trace gases are shown to the left. The types of effects caused by stratospheric ozone depletion and consequent increased UV-B penetration to the Earth's surface are hypothesized to include both direct effects on human health (e.g., increased cancer rates, immune suppression, etc.) and other terrestrial and aquatic ecological effects resulting from increased UV-B alterations of biogeochemical cycles.

significant extent over the Arctic Polar Region during late winter and spring in the Northern Hemisphere, and to a lesser extent, over mid-latitude regions during any season.

Given the long time involved in transport of such gases to the stratosphere and their long residence times there, any effects already seen on stratospheric ozone are likely caused by the atmospheric loadings of trace gases from anthropogenic emissions over the past few decades.

Those gases already in the atmosphere may continue to exert stratospheric ozone depletion effects well into the 21st century. Shorter-lived gases, such as CH_3Br, also exert significant ozone depletion effects.

The main types of deleterious effects hypothesized as likely to result from stratospheric ozone depletion and consequent increased SUVB penetration through the Earth's atmosphere include the following.

(1) *Direct Human Health Effects*, such as skin damage (sunburn), leading to more rapid aging and increased incidence of skin cancer; ocular effects (retinal damage and increased cataract formation possibly leading to blindness); and suppression of some immune system components (contributing to skin cancer induction and spread to nonirradiated skin areas, as well as possibly increasing susceptibility to certain infectious diseases or decreasing effectiveness of vaccinations).

(2) *Agricultural/Ecological Effects*, mediated largely through altered biogeochemical cycling resulting in consequent damaging impacts on terrestrial plants (leading to possible reduced yields of rice, other food crops, and commercially important trees, as well as to biodiversity shifts in natural terrestrial ecosystems); and deleterious effects on aquatic life (including reduced ocean zooplankton and phytoplankton, as important base components of marine food-chains supporting the existence of commercially important, edible fish and other seafood, as well as to other aquatic ecosystem shifts).

(3) *Indirect Human Health and Ecological Effects*, mediated through increased tropospheric ozone formation (and consequent exacerbation of surface-level, ozone-related health and ecological impacts) and alterations in the concentrations of other important trace species, most notably the hydroxyl radical and acidic aerosols.

(4) *Other Types of Effects*, such as faster rates of polymer weathering because of increased UV-B radiation and other effects on man-made commercial materials and cultural artifacts, secondary to climate change or exacerbation of air pollution problems.

Extensive qualitative and quantitative characterizations of stratospheric ozone depletion processes and projections of their likely potential impacts on human health and the environment have been the subjects of periodic (1988, 1989, 1991, 1994, 1998) international assessments carried out under WMO and UNEP auspices since the 1987 signing of the Montreal Protocol on Substances that Deplete the Ozone Layer. For detailed up-to-date information, the reader is

referred to recent international assessments of (a) processes contributing to stratospheric ozone depletion and the status of progress towards ameliorating the problem (WMO, 1999) and (b) revised qualitative and quantitative projections of likely consequent human health and environmental effects (UNEP, 1998, 2000) — with the findings and conclusions of these assessments being incorporated herein by reference.

Of considerable importance is the growing recognition, as reflected in these newer assessments, of impacts of enhanced solar radiation on biogeochemical cycles (see, for example, Zepp et al., 1998, and earlier discussions in this chapter in Section 4.2). As noted in the Zepp et al. paper, the effects of UV-B radiation (both in magnitude and direction) on trace gas (e.g., CO) emissions and mineral nutrient cycling are species specific and can affect a variety of processes. These include, for example, changes in the chemical composition of living plant tissue, photodegradation of dead plant matter (e.g., ground litter), release of CO from vegetation previously charred by fire, changes in microbial decomposer communities, and effects on nitrogen-fixing microorganisms and plants. Also, studies of natural acquatic ecosystems indicate that organic matter is the primary determinant of UV-B penetration through water. Changes in the amount and composition of organic matter, caused by enhanced UV-B penetration, affect the transmission of solar ultraviolet and visible radiation through the water column. These changes in light quality broadly impact the effects of UV-B on aquatic biogeochemical cycles. Enhanced UV-B levels have both positive and negative impacts on microbial activities in aquatic ecosystems that can affect nutrient cycling and the uptake or release of greenhouse gases. Thus, there are emerging complex issues regarding interactions and feedbacks between climate change and changes in terrestrial and marine biogeochemical cycles because of increased UV-B penetration to the Earth's surface.

In contrast to the above types of negative impacts projected as likely to be associated with increased UV-B penetration to Earth's surface, some research results are suggestive of possible beneficial effects of increased UV-B radiation. For example, a number of U.S. and international studies have focused on the protective effects of UV-B radiation with regard to non-skin cancer incidence. One of the first of these studies investigated potential relationships between sunlight, vitamin D and colon cancer (Garland and Garland, 1980). More recent studies continue to provide evidence that UV-B radiation may be protective against several types of cancer and some other diseases. For example, Grant (2002) has conducted a number of ecologic-type

epidemiologic studies, which suggest that UV-B radiation, acting through the production of vitamin D, is a risk-reduction factor for mortality due to several types of cancer, including cancer of the breast, colon, ovary, and prostate, as well as non-Hodgkin lymphoma. Other related studies that provide evidence for protective effects of UV-B radiation include: Gorham et al. (1989); Gorham et al. (1990); Garland et al. (1990); Hanchette and Schwartz (1992); Ainsleigh (1993); Lefkowitz and Garland (1994); Hartge et al. (1996); and Freedman et al. (1997).

As noted in the above detailed international assessments, since the signing of the Montreal Protocol, much progress has been made in reducing emissions of ozone depleting gases, leading to estimates that the maximum extent of stratospheric ozone depletion has likely leveled off during recent years, and this is expected to be followed by gradual lessening of the problem and its impacts during the next half-century. However, the assessments also note that the modeled projections are subject to considerable uncertainty (see, for example, UNEP, 2000). Varying potential roles of atmospheric particles, discussed below, are among numerous salient factors complicating predictive modeling efforts.

4.5.1.2 Effects of Airborne Particles on Transmission of Solar Ultraviolet Radiation Through the Atmosphere

A given amount of ozone in the lower troposphere has been shown to absorb more solar radiation than an equal amount of ozone in the stratosphere because of the increase in its effective optical path produced by Rayleigh scattering in the lower atmosphere (Brühl and Crutzen, 1988). The effects of particles are more complex. The impact of particles on the SUVB flux throughout the boundary layer are highly sensitive to the altitude of the particles and to their single scattering albedo. Even the sign of the effect can reverse as the composition of the particle mix changes from scattering to absorbing types (e.g., from sulfate to elemental carbon or PAHs) (Dickerson et al., 1997). In addition, scattering by particles also may increase the effective optical path of absorbing molecules, such as ozone, in the lower atmosphere.

The effects of particles present in the lower troposphere on the transmission of SUVB have been examined both by field measurements and by radiative transfer model calculations. The presence of particles in urban areas modifies the spectral distribution of solar irradiance at the surface. Shorter wavelength radiation (i.e., in the ultraviolet) is attenuated more than visible radiation (e.g., Peterson et al., 1978; Jacobson, 1999). Wenny et al. (1998) also found greater

attenuation of SUVB than SUVA (315 to 400 nm). However, this effect depends on the nature of the specific particles involved and, therefore, is expected to depend strongly on location. Lorente et al. (1994) observed an attenuation of SUVB ranging from 14 to 37%, for solar zenith angles ranging from about 30º to about 60º, in the total (direct and diffuse) SUVB reaching the surface in Barcelona during cloudless conditions on very polluted days (aerosol scattering optical depth at 500 nm, $0.46 \lesssim \tau_{500\,\mathrm{nm}} \lesssim 1.15$) compared to days on which the turbidity of urban air was similar to that for rural air ($\tau_{500\,\mathrm{nm}} \lesssim 0.23$). Particle concentrations that can account for these observations can be estimated roughly by combining Koschmeider's relation for expressing visual range in terms of extinction coefficient with one for expressing the mass of $PM_{2.5}$ particles in terms of visual range (Stevens et al., 1984). By assuming a scale height (i.e., the height at which the concentration of a substance falls off to 1/e of its value at the surface) of 1 km for $PM_{2.5}$, an upper limit of 30 μg/ m^3 can be derived for the clear case and between 60 and 150 μg/m^3 for the polluted case. Estupiñán et al. (1996) found that summertime haze under clear sky conditions attenuates SUVB between 5 and 23% for a solar zenith angle of 34°, compared to a clear sky day in autumn. Mims (1996) measured a decrease in SUVB by about 80% downwind of major biomass burning areas in Amazonia in 1995. This decrease in transmission corresponded to optical depths at 340 nm ranging from three to four. Justus and Murphey (1994) found that SUVB reaching the surface decreased by about 10% because of changes in aerosol loading in Atlanta, GA, from 1980 to 1984. Also, higher particle levels in Germany (48 °N) may be responsible for greater attenuation of SUVB than in New Zealand (Seckmeyer and McKenzie, 1992).

In a study of the effects of nonurban haze on SUVB transmission, Wenny et al. (1998) derived a very simple regression relation between the measured aerosol optical depth at 312 nm,

$$\ln(\text{ SUVB transmission at solar noon}) = -0.1422\,\tau_{312\,\mathrm{nm}} - 0.138,\ R^2 = 0.90, \qquad (4\text{-}11)$$

and the transmission of SUVB to the surface. In principle, values of $\tau_{312\,\mathrm{nm}}$ could be found from knowledge of the aerosol optical properties and visual range values. Wenny et al. (1998) also found that absorption by particles accounted for 7 to 25% of the total (scattering + absorption) extinction. Relations such as the above one are strongly dependent on local conditions and should not be used in other areas without knowledge of the differences in aerosol properties.

Although all of the above studies reinforce the idea that particles play a major role in modulating the attenuation of SUVB, none included measurements of ambient PM concentrations, so direct relations between PM levels and SUVB transmission could not be determined.

Vuilleumier et al. (2001) concluded that variations in aerosol scattering and absorption were responsible for 97% of the variability in the optical depth measured at seven wavelengths from 300 to 360 nm at Riverside, CA from 1 July to 1 November, 1997. Similar measurements made at Mt. Wilson, located above the main surface haze layer, showed that 80% of the variations in optical depth were still driven by variations in aerosol scattering and absorption. The remainder of the variability in optical depth was attributed mainly to variability in ozone under clear-sky conditions. However, these results cannot be extrapolated to other locations because these effects are coupled and non-linear and are not straightforward. They depend on the concentrations of these species and on the physical and chemical characteristics of the particles. Hence, any quantitative statements regarding the relative importance of particles and ozone will be location-specific.

Liu et al. (1991) estimated, roughly, the overall effects on atmospheric transmission of SUVB of increases of anthropogenic airborne particles that have occurred since the beginning of the industrial revolution. Based on (a) estimates of the reduction in visibility from about 95 km to about 20 km over nonurban areas in the eastern United States and in Europe, (b) calculations of optical properties of airborne particles found in rural areas to extrapolate the increase in extinction at 550 to 310 nm, and (c) radiative transfer model calculations, Liu et al. concluded that the amount of SUVB reaching Earth's the surface likely has decreased from 5 to 18% since the beginning of the industrial revolution. This was attributed mainly to scattering of SUVB back to space by sulfate containing particles. Radiative transfer model calculations have not been done for urban particles.

Although aerosols are expected to decrease the flux of SUVB reaching the surface, scattering by particles is expected to result in an increase in the actinic flux within and above the aerosol layer. However, when the particles significantly absorb SUVB, a decrease in the actinic flux is expected. Actinic flux is the radiant energy integrated over all directions at a given wavelength incident on a point in the atmosphere, and is the quantity needed to calculate rates of photolytic reactions in the atmosphere. Blackburn et al. (1992) measured attenuation of the photolysis rate of ozone and found that aerosol optical depths near unity at 500 nm reduced

ozone photolysis rate by as much as a factor of two. Dickerson et al. (1997) showed that the photolysis rate for NO_2 , a key parameter for calculating the overall intensity of photochemical activity, could be increased within and above a scattering aerosol layer extending from the surface, although it would be decreased at the surface. This effect is qualitatively similar to what is seen in clouds, where photolysis rates are increased in the upper layers of a cloud and above the cloud (Madronich, 1987). For a simulation of an ozone episode that occurred during July 1995 in the Mid-Atlantic region, Dickerson et al. (1997) calculated ozone increases of up to 20 ppb compared to cases that did not include the radiative effects of particles in urban airshed model (UAM-IV) simulations. In contrast, Jacobson (1998) found that particles may have caused a 5 to 8% decrease in O_3 levels during the Southern California Air Quality Study in 1987. Absorption by organic compounds and nitrated inorganic compounds was hypothesized to account for the reductions in UV radiation intensity.

The photolysis of ozone in the Hartley bands also leads to production of electronically excited oxygen atoms, $O(^1D)$ that then react with water vapor to form OH radicals. Thus, enhanced photochemical production of ozone is accompanied by the scavenging of species involved in greenhouse warming and stratospheric depletion. However, these effects may be neutralized or even reversed by the presence of absorbing material in the particles. Any evaluation of the effects of particles on photochemical activity therefore will depend on the composition of the particles and also will be location-specific.

Also complicating any straightforward evaluation of UV-B penetration to specific areas of the Earth's surface are the influences of clouds, as discussed by Erlick et al. (1998), Frederick et al. (1998), and Soulen and Fredrick (1999). The transmission of solar UV and visible radiation is highly sensitive to cloud type and cloud amount and the extent of their external or internal mixing with cloud droplets. Even in situations of very low atmospheric PM (e.g., over Antarctica), interannual variations in cloudiness over specific areas can be as important as ozone levels in determining UV surface irradiation, with net impacts varying from a month or season to another (Soulen and Fredrick, 1999). Evaluations of the effects of changes in the transmission of solar UV-B radiation to the surface have been performed usually for cloud-free or constant cloudiness conditions.

Given the above considerations, quantification of projected effects of variations in atmospheric PM on human health or the environment because of the effects of particles on the

transmission of solar UV-B radiation requires location-specific evaluations, taking into account composition, concentration, and internal structure of the particles; temporal variations in atmospheric mixing height and depths of layers containing the particles the abundance of ozone and other absorbers within the planetary boundary layer and the free troposphere. The outcome of such modeling effects would likely vary from location to location in terms of increased or decreased surface level UV-B exposures because of location-specific changes in atmospheric PM concentrations or composition. For example, to the extent that any location-specific scattering by airborne PM were to affect the directional characteristics of UV radiation at ground level, and thereby enhance radiation incident from low angles (Dickerson, 1997), the biological effectiveness (whether deleterious or beneficial) of resulting ground-level UV-B exposures could be enhanced. Airborne PM also can reduce the ground-level ratio of photorepairing radiation (UV-A and short-wavelength visible) to damaging UV-B radiation. Lastly, PM deposition is a major source of PAHs in certain freshwater lakes and coastal areas, and the adverse effects of solar UV are enhanced by the uptake of PAHs by aquatic organisms. Thus, although airborne PM may, in general, tend to reduce ground-level UV-B, its net effect in some locations may be to increase UV damage to certain aquatic and terrestrial organisms, as discussed by Cullen and Neale (1997).

4.5.2 Global Warming Processes, Human Health and Environmental Impacts, and Roles of Atmospheric Particle

4.5.2.1 Bases for Concern Regarding Global Warming and Climate Change

Various trace gases emitted by man's activities, including several noted above as contributing to stratospheric ozone depletion, can act as "greenhouse gases" (GHG). That is, as their tropospheric concentrations increase, they retard the escape of infrared radiation from the earth's surface and thereby contribute to the trapping of heat near the surface (the "greenhouse effect") and, ultimately, to consequent global warming and climate change. Much concern has evolved with regard to increases in the naturally very low concentrations in the atmosphere of some of these gases, especially carbon dioxide (CO_2), nitrous oxide (N_2O), methane (CH_4), chloroflurocarbons (CFCs), and tropospheric ozone (O_3).

Atmospheric processes involved in mediating global warming and its likely consequent effects have been reviewed extensively (United Nations Environment Programme, 1986; World

Meteorological Organization, 1988; U.S. Environmental Protection Agency, 1987; IPCC, 1996, 1998, 2001a,b; NAST, 2000) and more concisely summarized by others (e.g., Patz et al., 2000a,b). The reader is referred to such reviews for more detailed information than that concisely summarized below. The main focus here is first (a) to provide a brief summary of key points regarding processes involved and types of effects projected as likely to be associated with global warming and climate change and, then, (b) to discuss salient considerations regarding potential impacts of atmospheric PM on such processes and effects.

The Third Assessment Report of the IPCC (2001a) discusses observed past changes in the climate system of the Earth. Of particular note is the calculation stated in that IPCC report indicating that the global average temperature (i.e., the average of near surface air temperatures over land and sea surface temperatures) has increased by 0.6 ± 0.2 °C over the course of the 20th Century. Globally, the decade of the 1990's was likely the warmest of the Century and 1998 the warmest year in the instrumental record (IPCC, 2001a). New analyses of proxy data for the Northern Hemisphere also indicate that the rise in temperature over the 20th Century is likely the largest of any century during the past 1,000 years (IPCC, 2001a). However, the projecting of future trends in global average temperature and regional climate impacts is difficult and fraught with many uncertainties.

All of the above noted assessments and summaries emphasize that estimating likely future global warming trends and associated climate change caused by greenhouse gases is extremely complex, with modeling results being highly dependent on key assumptions about the rates of future increases in various gases and numerous other factors (including particle effects). Modeling of the magnitude of the warming directly associated with radiative forcing by greenhouse gases (without feedback enhancement) projects temperature increases ranging from 1.4 °C to 5.8 °C over the period 1990 to 2100 (IPCC, 2001a). This range does not include uncertainties in the modeling of radiative forcing (e.g., aerosol radiative forcing). Feedbacks that likely would increase temperatures further are expected to occur. Increased water vapor (trapping heat) and snow and ice melting (reducing reflection of radiation back into space) are two examples of such feedback factors expected to increase temperatures. However, major uncertainties exist with regard to feedbacks between global warming and clouds, which could either amplify or, perhaps, reduce a temperature rise. Taking assumptions about rates of increase (or decrease) in GHG concentrations, consequent initial warming effects, feedback effects, and

accompanying uncertainties into account, numerous modeling efforts have attempted to project likely future trends in global warming. Despite the complexity and uncertainties inherent in such modeling efforts, all typically agree that some global warming has occurred and will continue to occur during the coming decades, but the ranges of quantitative estimates vary considerably depending on specific assumptions incorporated into the models. Thus, for example, "low" scenarios assuming stabilization or reductions in GHG emissions (resulting from implementation of the 1987 Montréal Protocol) project lower temperature changes than other scenarios assuming higher rates of increase in GHG emissions or differing feedback-effect patterns.

Given the wide range of estimates of global warming trends and patterns of associated climate change emerging from modeling efforts, the estimation of likely human health and ecological effects associated with global warming on any quantitative basis is extremely difficult. The onset of any notable global warming effect is also important, with various analyses indicating that global temperatures for the past century have been rising (and now appear to be beyond average levels within the range of variation seen with cycles of global warming or cooling over the past several centuries before marked anthropogenic emissions of greenhouse gases occurred). Also posing difficulties for the quantitative estimation of human health and other effects are expected wide regional variations in temperature and climate characteristics (e.g., rain and snowfall amounts) that may be projected reasonably to result from various global warming trend scenarios. Lastly, it should be noted that, despite general warming trends in long-term average temperatures, wide extremes in both high *and* low temperatures also are expected to occur more frequently in some areas.

Special reports of the IPCC Working Group II on impacts of climate change (IPCC, 1998, 2001b) assess global warming processes and identify several types of vulnerabilities likely to occur because of climate change resulting from global warming. Such general types of vulnerabilities include impacts on terrestrial and aquatic ecosystems, hydrology and water resources, food and fiber production, coastal systems, and human health. The executive summaries of these IPCC (1998, 2001b) reports provide helpful overviews of key points regarding projected global warming processes, likely climate change patterns, and their consequent impacts in terms of the types of vulnerabilities noted above.

The IPCC (1998, 2001b) reports indicate that human activities resulting in emissions of long-lived GHCs are projected by general circulation models (GCMs) as likely to lead to marked

global and regional changes in temperature, precipitation and other climate variables. The average rate of warming is projected to be more rapid than any seen in the past 10,000 years, although regional changes could differ substantially from mean global rates. This is expected to result in increases in global mean sea level; prospects for more extreme weather events, floods, and droughts in some areas; and consequent changes in soil moisture. The most recent IPCC Reports (2001a,b) highlight GCM modeling results, based on various scenarios of current and plausible future emissions of GHGs and aerosols and the range of sensitivities of climate change to atmospheric levels (and residence time) of GHGs, which project mean annual global surface temperature increases leading by 2100 to global mean sea level rise of 0.09 to 0.88 m above 1990 levels and significant changes in spatial and temporal patterns of precipitation.

Human health, ecosystems, and socioeconomic sectors (e.g., hydrology and water resources, food and fiber production, etc.) are also projected by IPCC (1998, 2001b) to be vulnerable to the magnitude and rate of climate change and extremes (see Table 4-20). Wide variations in the courses and net impacts of climate change in different geographic areas can be expected. In general, projected climate change impacts can be expected to represent additional stresses on those natural ecosystems and human societal systems already impacted by increasing resource demands, unsustainable resource management practices, and pollution—with wide variation likely across regions and nations in their ability to cope with consequent alterations in ecological balances, in availability of adequate food, water, and clean air, and in human health and safety. However, although many regions are likely to experience severe adverse impacts (some possibly irreversible) of climate change, some climate change impacts may be locally beneficial in some regions.

The 1998 IPCC special report regarding the assessment of different types of vulnerabilities to climate change included projections of likely impacts for each of 10 different geographic regions of the Earth, including those projected for two regions (North America and Polar) of most relevance to the continental United States and Alaska. Probably of most note are findings indicating that (a) the characteristics of subregions and sectors of North America suggest that neither impacts of climate change nor response options will be uniform, and (b) many systems of North America are moderately to highly sensitive to climate change, with the range of estimated effects encompassing potential substantial damage or, conversely, some potential for beneficial outcomes. The most vulnerable continental United States sectors and regions include long-lived

TABLE 4-20. EXAMPLES OF IMPACTS RESULTING FROM PROJECTED CHANGES IN EXTREME CLIMATE EVENTS

Projected changes during the 21st Century in Extreme Climate Phenomena and their Likelihood[a]	Representative Examples of Projected Impacts[b] *(all high confidence of occurrence in some areas[c])*
Simple Extremes	
Higher maximum temperatures; more hot days and heat waves[d] over nearly all land areas (*very likely*[a])	• Increased incidence of death and serious illness in older age groups and urban poor • Increased heat stress in livestock and wildlife • Shift in tourist destinations • Increased risk of damage to a number of crops • Increased electric cooling demand and reduced energy supply reliability
Higher (increasing) minimum temperatures; fewer cold days, frost days, and cold waves[d] over nearly all land areas (*very likely*[a])	• Decreased cold-related human morbidity and mortality • Decreased risk of damage to a number of crops, and increased risk to others • Extended range and activity of some pest and disease vectors • Reduced heating energy demand
More intense precipitation events (*very likely*[a] over many years)	• Increased flood, landslide, avalance, and mudslide damage • Increased soil erosion • Increased flood runoff could increase recharge of some floodplain aquifers • Increased pressure on government and private flood insurance systems and disaster relief
Complex Extremes	
Increased summer drying over most mid-latitude continental interiors and associated risk of drought (*likely*[a])	• Decreased crop yields • Increased damage to building foundations caused by ground shrinkage • Decreased water resource quantity and quality • Increased risk of forest fire
Increase in tropical cyclone peak wind intensities, mean and peak precipitation intensities (*likely*[a] over some areas)[e]	• Increased risk to human life, risk of infections, disease epidemics, and many other risks • Increased coastal erosion and damage to coastal buildings and infrastructure • Increased damage to coastal ecosystems such as coral reefs and mangroves
Intensified droughts and floods associated with El Niño events in many different regions (*likely*[a]) (see also under droughts and intense precipitation events)	• Decreased agricultural and rangeland productivity in drought- and flood-prone regions • Decreased hydro-power potential in drought-prone regions
Increased Asian summer monsoon precipitation variability (*likely*[a])	• Increased flood and drought magnitude and damages in temperate and tropical Asia
Increased intensity of mid-latitude storms (little agreement between current models)[d]	• Increased risks to human life and health • Increased property and infrastructure losses • Increased damage to coastal ecosystems

[a]Likelihood refers to judgmental estimates of confidence used by TAR WGI: *very likely* (90-99% chance); *likely* (66-90% chance). Unless otherwise stated, information on climate phenomena is taken from the Summary for Policymakers, TAR WGI. TAR WGI = Third Assessment Report of Working Group 1 (IPCC, 2001a).
[b]These impacts can be lessened by appropriate response measures.
[c]High confidence refers to probabilities between 67 and 95%.
[d]Information from TAR WGI, Technical Summary.
[e]Changes in regional distribution of tropical cyclones are possible but have not been established.

Source: IPCC (2001b).

natural forest ecosystems in the East and interior West, water resources in the southern plains, agriculture in the Southeast and southern plains, northern ecosystems and habitats, estuaries and beaches in developed areas, and low-latitude cool and cold water fisheries. Other sectors or subregions may benefit from warmer temperatures or increased CO_2 fertilization (e.g., west coast coniferous forests; some western rangelands; reduced energy costs for heating in northern latitudes; reduced road salting and snow-clearance costs; longer open-water seasons in norther channels and ports; and agriculture in the northern latitudes, the interior West, and the west coast). For Alaska, substantial shifts in ecosystems (with possible major declines or loss of some sensitive species like bear and caribou or of other ice-dependent animals) may occur in parallel to beneficial effects such as opening of ice-bound water transportation routes or possible expanded agricultural viability secondary to longer growing seasons. On the other hand, for North America, the potential for mainly deleterious direct or indirect effects on human health is likely to increase (e.g., increased mortality directly linked to temperature extremes, increases in incidence and spread of vector-borne infectious diseases, impacts secondary to sea-level rise, and impacts secondary to increased tropospheric air pollution.

More detailed evaluations of possible global climate change impacts on various U.S. geographic areas are being conducted by the United States Global Change Research Program (USGCRP). An overview report on the assessment results and key findings from a series of workshops convened by the USGCRP National Assessment Synthesis team (NAST) has been prepared (NAST, 2000). Overall key findings from the USGCRP (NAST, 2000) report are noted below.

(1) *Increased Warming*. Assuming continued growth in world GHG emissions, the primary climate models used in the USGCRP assessment project that temperatures in the United States will rise by 5 to 9 °F (3 to 5 °C) on average during the next 100 years. A wide range of outcomes is possible.

(2) *Differing Regional Impacts*. Climate change will vary widely across the United States. Temperature increases will vary somewhat from region to region. Heavy and extreme precipitation events are likely to become more frequent, yet some regions will get drier. The potential impacts of climate change will vary widely across the nation.

(3) *Vulnerable Ecosystems*. Many ecosystems are highly vulnerable to the projected rate and magnitude of climate change. A few, such as alpine meadows in the Rocky Mountains and

some barrier islands, are likely to disappear entirely in some areas; and others, such as some forests of the Southeast, are likely to experience major species shifts or break up into a mosaic of grasslands, woodlands, and forests. Goods and services lost through disappearance or fragmentation of certain ecosystems are likely to be costly or impossible to replace.

(4) *Widespread Water Concerns*. Water is an issue in every region, but the nature of the vulnerabilities varies, with different nuances in each. Drought is an important concern in every region. Floods and water quality are concerns in many regions. Snowpack changes are especially important in the West, the Pacific Northwest, and Alaska.

(5) *Secure Food Supply*. At the national level, the U.S. agriculture sector is likely to be able to adapt to climate change. Overall, U.S. crop productivity is very likely to increase over the next few decades, but the gains will not be uniform across the nation. Falling prices and competitive pressures are very likely to stress some farmers, while benefiting consumers.

(6) *Near-Term Increases in Forest Growth*. Forest productivity is likely to increase over the next several decades in some areas as trees respond to higher CO_2 levels. Over the longer term, changes in larger scale processes such as fire, insects, droughts, and disease will possibly decrease forest productivity. Also, climate change is likely to cause long-term shifts in forest species (e.g., distribution of sugar maple stands more northward, out of the United States).

(7) *Increased Damage in Coastal and Permafrost Areas*. Climate change and the resulting rise in sea level are likely to exacerbate threats to building, roads, powerlines, and other infrastructure in climatically sensitive places. For example, infrastructure damage is related to permafrost melting in Alaska and to sea level rise and storm surges in low-lying coastal areas.

(8) *Adaptation Determines Health Outcomes.* A range of negative health impacts is possible from climate change, but adaptation is likely to help protect much of the U.S. population. Maintaining our nation's public health and community infrastructure, from water treatment systems to emergency shelters, will be important for minimizing the impacts of waterborne diseases, heat stress, air pollution, extreme weather events, and diseases transmitted by insects, ticks, and rodents.

(9) *Other Stresses Magnified by Climate Change*. Climate change will very likely magnify the cumulative impacts of other stresses, such as air and water pollution and habitat destruction caused by human development patterns. For some systems, such as coral reefs, the combined effects of climate change and other stresses are very likely to exceed a critical threshold, bringing large, possibly irreversible impacts.

(10) *Uncertainties Remain and Surprises Are Expected*. Significant uncertainties remain in the science underlying regional climate changes and their impacts. Further research is needed to improve understanding and predictive ability about societal and ecosystem impacts and to provide the public with additional useful information about adaptation strategies. However, it is likely that some aspects and impacts of climate change will be totally unanticipated as complex systems respond to ongoing climate change in unforeseeable ways.

For more specific information on the types of effects projected as likely to occur in the United States, the reader is referred to the USGSRC Report (NAST, 2000), several subsidiary regional reports (MARAT, 2000; Yarnal et al., 2000; NERAG, 2001; GLRAG, 2000), and the health assessment report (Bernard, et al., 2001).

Findings from the USGCRP (NAST, 2000) report and subsidiary regional reports illustrate well the considerable uncertainties and difficulties in projecting likely climate change impacts on regional or local scales. The findings also reflect well the mixed nature of projected potential climate change impacts (combinations of mostly deleterious, but other possible beneficial effects) for U.S. regions and their variation across the different regions. Difficulties in projecting region-specific climate change impacts are complicated by the need to evaluate potential effects of local- or regional-scale changes in key air pollutants not only on global scale temperature trends but also in terms of potentially more local- or regional- scale impacts on temperature and precipitation patterns. Of much importance for this are varying roles played by atmospheric particles.

4.5.2.2 Airborne Particle Relationships to Global Warming and Climate Change

Atmospheric particles both scatter and absorb incoming solar radiation at visible light wavelengths. The scattering of solar radiation back to space leads to a decrease in transmission of visible radiation to the Earth's surface and, hence, to a decrease in the heating rate of the

surface and the atmosphere. The absorption of either incoming solar radiation or outgoing terrestrial infrared radiation by atmospheric particles results in heating of the lower atmosphere. Interactions of atmospheric particles with electromagnetic radiation from the visible through the infrared spectral regions are responsible for their direct effects on climate, which are the result of the same physical processes responsible for visibility degradation. Visibility reduction is caused by particle scattering in all directions, whereas climate effects result mainly from scattering in the upward direction. The net effect of the above processes can be expressed as a radiative forcing, which is the change in the average net radiation at the top of the troposphere because of a change in solar (shortwave, or visible) or terrestrial (longwave, or infrared) radiation (Houghton et al., 1990). The radiative forcing drives the climate to respond, but because of uncertainties in a number of feedback mechanisms involving climate response, radiative forcing is used as a first-order estimate of the potential importance of various substances. Sulfate particles scatter solar radiation effectively and do not absorb at visible wavelengths, whereas they absorb weakly at infrared wavelengths (IPCC, 2001). Nitrate particles exhibit grossly similar properties. The effects of mineral dust particles are complex; they weakly absorb solar radiation but their overall effect on solar radiation depends on particle size and the reflectivity of the underlying surface. They absorb infrared radiation and thus contribute to greenhouse warming (Tegen et al., 1996). Organic carbon particles mainly reflect solar radiation, whereas elemental carbon and other black carbon particles (e.g., PAHs with H:C ratios of ≤ 0.3) are strong absorbers of solar radiation (IPCC, 2001). However, the optical properties of carbonaceous particles are modified if they become coated with water or sulfuric acid. Particles containing black carbon also can exert a direct effect after deposition onto surfaces that are more reflective (e.g., snow and ice). In this case, additional solar radiation is absorbed by the surface; conversely, more reflective particles deposited on a dark surface result in additional solar radiation being reflected back to space.

Anthropogenic (Twomey, 1974; Twomey, 1977) and biogenic (Charlson et al., 1987) sulfate particles also exert indirect effects on climate by serving as cloud condensation nuclei, which results in changes in the size distribution of cloud droplets by producing more particles with smaller sizes. The same mass of liquid water in smaller particles leads to an increase in amount of solar radiation that clouds reflect back to space because the total surface area of the cloud droplets is increased. This has been supported by satellite observations indicating that the effective radius of cloud droplets is smaller in the Northern Hemisphere than in the Southern

Hemisphere (Han et al., 1994). Smaller cloud droplets also have a lower probability of precipitating and, thus, have a longer lifetime than larger ones. Although the effects of sulfate have been considered most widely, interactions with other aerosol components also may be important. Novakov and Penner (1993) have provided evidence that carbonaceous particles can modify the nucleation properties of sulfate particles.

The amount of solar radiation incident on the earth-atmosphere system, or the solar constant, is 1370 W m^{-2}, or 342.5 W m^{-2} on a globally averaged basis (calculated by dividing the solar constant by 4). The addition of sulfate and organic carbon as airborne PM results in enhanced scattering and net cooling, whereas the addition of particles containing elemental carbon results in absorption of solar and terrestrial radiation and net heating. The estimated radiative forcing because of the scattering of solar radiation back to space caused mainly by sulfate particles is −0.4 W m^{-2}; −0.2 W m^{-2} for biomass-burning aerosols; −0.1 W m^{-2} for fossil fuel organic carbon; and +0.2 W m^{-2} for fossil fuel black carbon. Uncertainties in these quantities are about a factor of two (IPCC, 2001a). The uncertainty range reflects uncertainties in the emissions of SO_2, the amount of SO_2 that is oxidized to sulfate, the atmospheric lifetime of sulfate, and the optical properties of the sulfate particles. These values may be compared to the radiative forcing exerted by greenhouse gases of about + 2.4 W m^{-2}, with an uncertainty factor of 1.15 from the preindustrial era (ca. 1750) to 2000. Since the beginning of the 20th century, the mean surface temperature of the earth has increased by about 0.6 °C (IPCC, 2001a). Estimates of the indirect effects of particles range from 0 to −2.0 W m^{-2} (IPCC, 2001a). Because of a lack of quantitative knowledge, no central value could be given. Therefore, on a globally averaged basis, the direct and indirect effects of anthropogenic sulfate particles likely have offset partially the warming effects caused by increases in levels of greenhouse gases (Charlson et al., 1992).

Much of the work investigating the effects of particles on climate has focused on sulfate particles. However, particles containing elemental carbon (EC) from fossil fuel combustion and biomass burning or mineral dust may exert radiative forcing, with spatial distributions very different than for sulfate. Tegen et al. (1996) and Tegen and Lacis (1996) used a global scale three-dimensional model to evaluate the radiative forcing caused by mineral dust particles. Tegen and Lacis (1996) found that the sign and the magnitude of the radiative forcing depends on the height distribution of the dust and the effective radius of the particles. In particular, for a dust layer extending from 0 km to 3 km, positive radiative forcing at visible wavelengths is found for

particle radii greater than 1.8 μm, whereas negative forcing is found for smaller particles. They calculated a global mean radiative forcing caused by mineral dust from all sources of 0.14 W m^{-2} and from mineral dust from lands disturbed by human activity of 0.09 W m^{-2}. This value represents a near cancellation between a much larger solar forcing of −0.25 W m^{-2} and a thermal forcing of 0.34 W m^{-2}. Uncertainty factors could not be estimated for these calculations because they were judged to be largely unknown. Haywood and Shine (1995) estimated a global mean radiative forcing of 0.1 W m^{-2}, with an uncertainty factor >3, caused by the absorption of solar radiation by EC released by fossil fuel combustion. The IPCC (1995) estimated a global mean radiative forcing of −0.1 W m^{-2} caused by particles produced by biomass burning, with an uncertainty factor of three. The global mean radiative forcing exerted by particles would then be −0.5 W m^{-2}, with an uncertainty of about a factor of 2.4. Figure 4-32 summarizes estimates of global mean radiative forcing exerted by greenhouse gases and various types of particles.

Deviations from the global mean values can be very large on the regional scale. For instance, Tegen et al. (1996) found that local radiative forcing exerted by dust raised from disturbed lands ranges from −2.1 W m^{-2} to 5.5 W m^{-2} over desert areas and their adjacent seas. The largest regional values of radiative forcing caused by anthropogenic sulfate are about −3 W m^{-2} in the eastern United States, south central Europe, and eastern China (Kiehl and Briegleb, 1993). These regional maxima in aerosol forcing are at least a factor of 10 greater than their global mean values shown in Figure 4-32. By comparison, regional maxima in forcing by the well-mixed greenhouse gases are only about 50% greater than their global mean value (Kiehl and Briegleb, 1993). Thus, the estimates of local radiative forcing by particles also are large enough to completely cancel the effects of greenhouse gases in many regions and to cause a number of changes in the dynamic structure of the atmosphere that still need to be evaluated. A number of anthropogenic pollutants whose distributions are highly variable are also effective greenhouse absorbers. These gases include O_3 and, possibly, HNO_3 , C_2H_4 , NH_3, and SO_2, all of which are not commonly considered in radiative forcing calculations (Wang et al. 1976). High ozone values are found downwind of urban areas and areas where there is biomass burning. However, Van Dorland et al. (1997) found that there may not be much cancellation between the radiative effects for ozone and for sulfate, because both species have different seasonal cycles and show significant differences in their spatial distribution.



Figure 4-32. Estimated global mean radiative forcing exerted by gas and various particle phase species for the year 2000, relative to 1750.

Source: IPCC (2001a).

Observational evidence for the climatic effects of particles is sparse. Haywood et al. (1999) found that the inclusion of anthropogenic aerosols results in a significant improvement between calculations of reflected sunlight at the top of the atmosphere and satellite observations in oceanic regions close to sources of anthropogenic PM.

Uncertainties in calculating the direct effect of airborne particles arise from a lack of knowledge of their vertical and horizontal variability, their size distribution, chemical composition and the distribution of components within individual particles. For instance, gas-phase sulfur species may be oxidized to form a layer of sulfate around existing particles in continental environments, or they may be incorporated in sea-salt particles (e.g., Li-Jones and Prospero, 1998). In either case, the radiative effects of a given mass of the sulfate will be much lower than if pure sulfate particles were formed. It also must be stressed that the overall radiative

effect of particles at a given location is not simply determined by the sum of effects caused by individual classes of particles because of interactions between particles with different radiative characteristics and with gases.

Calculations of the indirect effects of particles on climate are subject to much larger uncertainties than are calculations of their direct effects, reflecting uncertainties in a large number of chemical and microphysical processes in describing the effects of sulfate on the size distribution and number of droplets within a cloud. A complete assessment of the radiative effects of PM will require supercomputer calculations that incorporate the spatial and temporal behavior of particles of varying composition that have been emitted or formed from precursors emitted from different sources. Refining values of model input parameters (such as improving emissions estimates) may be as important as improving the models per se in calculations of direct radiative forcing (Pan et al., 1997) and indirect radiative forcing (Pan et al., 1998) caused by sulfate. However, uncertainties associated with the calculation of radiative effects of particles likely will remain much larger than those associated with well-mixed greenhouse gases.

This means that, although on a global scale atmospheric particles likely exert an overall net effect of slowing global warming, much uncertainty would apply to any modeling efforts aimed at projecting net effects on global warming processes, resulting climate change, and any consequent human health or environmental effects because of location-specific increases or decreases in anthropogenic emissions of atmospheric particles or their precursors. For example, any net impacts of regional sulfates in reducing global-climate-change-induced increases in local temperatures may well be offset partially by local surface level heating because of carbonaceous particles from diesel emissions or coal combustion energy generation being deposited on snow or ice covered surfaces or contributing to more rapid evaporation or rainout of water from overhead clouds.

4.6 SUMMARY

4.6.1 Particulate Matter Effects on Vegetation and Ecosystems

Particulate matter (PM) deposition on vegetation and ecosystems has been defined mainly by size fraction, not by chemical composition, structure, or source. Though size is related to

mode and magnitude of deposition to vegetated landscapes and may be a useful surrogate for chemical composition, the size classes have little specific relevance to vegetation.

Deposition of PM on vegetation and ecosystems is not well understood. Atmospheric deposition of particles takes place via both wet and dry processes via three major routes: (1) precipitation and scavenging in which particles are deposited in rain and snow; (2) occult (fog, cloud water, and mist interception); and (3) the much slower dry deposition. All three modes of deposition must be considered when determining inputs to ecosystems or water sheds because each may dominate over specific intervals of time of space.

Wet deposition is generally confounded by fewer factors than the other two methods and has been easier to quantify. Total inputs by wet deposition can be significant; however, not all wet deposition involves particle scavenging because gaseous pollutants also dissolve in rain drops during precipitation events. This contribution is obscured because wet deposition is measured simply by chemical analysis of total precipitation collected in clean non-reactive buckets. Wet deposition is largely function of precipitation amount and ambient pollutant concentrations. Surface properties are relatively unimportant except for landform features that alter local distribution of precipitation (orographic effects). Wet deposition is most effective for fine particles of atmospheric (secondary) origin and elements such as cadmium, chromium, lead, nickel, and vanadium.

Dry deposition depends more strongly on surface properties, such as micrometeorological roughness, which determine impaction and reentrainment of individual particles, and on particle size distribution in the atmosphere. Vegetation discontinuities, such as forest edges and margins of cultivated fields, may be subject to increased deposition of PM. Dry deposition of atmospheric particles to plant and soil is a much slower processes than either wet or occult deposition and is most effective for coarse particles including primary geologic material, and for elements such as iron and manganese. It is nearly continuous and affects all types of plant parts, including those not currently physiologically active, along with exposed soil and water surfaces, received steady deposits of dry dusts, elemental carbon encrustations, grease films, tarry acidic coatings and heterogeneous secondary particles for from gaseous precursors.

Occult deposition is of more restricted occurrence than either of the above. Occult deposition of cloud and fog water droplets containing PM may be determined by both atmospheric and surface features. Formation of fog may accelerate deposition by transforming

fine PM with low deposition velocities, into larger hydrometeors with correspondingly larger deposition velocities.

The sources and forms of nitrogen in the atmosphere are poorly studied, and the concentrations are rarely measured, except in precipitation. The influence of aerodynamic diameter is particularly critical for the deposition of nitrogen species because they exist as a wide range of particle sizes in the atmosphere. For example, at many North American sites NO_3^- is characterized by bimodal distribution, with modes above and below 1 μm. Although the annual deposition for NH_4^+ is distributed similarly among the fine and coarse particles, particulate NO_3^- is found predominately in coarse particle fractions. Similar to the pattern for NH_4^+, the estimated annual deposition of SO_4^{-2} particles occurs in both the fine and coarse particulate fractions. Base cation deposition is virtually restricted to contributions from coarse particles.

The ambient concentration of particles, the parameter for which there is most data (Chapter 3), is at best a surrogate indicator of exposure. The amount entering the immediate plant environment, deposition to the plant surfaces or soil in the vicinity of the roots, determines the biological effect.

Annual amounts of total heavy metal deposition are highly variable depending on specific forest location and upwind source strength. Depending on climate conditions and topography, fine particles may remain airborne for days to months and may be transported 1,000 to 10, 000 km or more from their source. This long-distant transport and subsequent deposition qualify heavy metals as regional- and global-scale air pollutants. Ecosystems immediately downwind of major emissions sources such as power generating, industrial, or urban complexes may receive locally heavy inputs. Mass balance budgets of seven heavy metals (cadmium, copper, iron, manganese, nickel, and zinc) have been determined at the Hubbard Brook Experimental Forest approximately 120 km north of Boston and relatively distant from major source of heavy metals. Investigations of trace metals conducted in roadside, industrial, and urban environments have demonstrated that impressive burdens of particulate heavy metal accumulate on vegetative surfaces. Theory and measurement techniques for wet and occult deposition processes are well advanced. In contrast, dry deposition of particles has remained difficult to measure and to model. Further advances in quantification of PM deposition will require development of improved analytical treatments of dry deposition, and increased chemical speciation of size classed PM.

Human existence on this planet depends on ecosystems and the services and products they provide. Both ecosystem structure and function play an essential role in providing societal benefits. Society derives two types of benefits from the structural aspects of an ecosystem: (1) products with market value such as fish, minerals, forage, forest products, biomass fuels, natural fiber, and many pharmaceuticals, and the genetic resources of valuable species (e.g., plants for crops and timber and animals for domestication); and (2) the use and appreciation of ecosystem for recreation, aesthetic enjoyment, and study.

Ecosystem functions that maintain clean water, pure air, a green earth, and a balance of creatures, are functions that enable humans to survive. They are the dynamics of ecosystems. The benefits they impart include absorption and breakdown of pollutants, cycling of nutrients, binding of soil, degradation of organic waste, maintenance of a balance of gases in the air, regulation of radiation balance, climate, and the fixation of solar energy. Concern has risen in recent years concerning the integrity of ecosystems because there are few ecosystems on Earth today that are not influenced by humans. For this reason, the deposition of PM and its impact on vegetation and ecosystems is of great importance.

The PM whose effects on vegetation and ecosystems are considered in this chapter is not a single pollutant but represents a heterogeneous mixture of particles differing in origin, size, and chemical constituents. The effects of exposure to a given mass concentration of PM of particular size (measured as PM_{10}; $PM_{2.5}$, etc.) may, depending on the particular mix of deposited particles, lead to widely differing phytotoxic responses. This has not been characterized adequately.

Atmospheric deposition of particles to ecosystems takes place via both wet and dry processes through the three major routes indicated below.

(1) Precipitation scavenging, in which particles are deposited in rain and snow

(2) Fog, cloud water, and mist interception

(3) Dry deposition, a much slower, yet more continuous removal to surfaces

Deposition of heavy metal particles to ecosystems occurs by wet and dry processes. Dry deposition is considered more effective for coarse particles of natural origin and elements such as iron and manganese, whereas wet deposition generally is more effective for fine particles of atmospheric origin and elements such as cadmium, chromium, lead, nickel, and vanadium. The actual importance of wet versus dry deposition, however, is highly variable, depending on the type of ecosystem, location, and elevation.

Deposition of PM on above-ground plant parts can have either a physical and or chemical impact, or both. Particles transferred from the atmosphere to plant surfaces may cause direct effects if they (1) reside on the leaf, twig, or bark surface for an extended period; (2) be taken up through the leaf surface; or (3) are removed from the plant via resuspension to the atmosphere, washing by rainfall, or litter-fall with subsequent transfer to the soil.

Chemical effects include excessive alkalinity or acidity. The effects of "inert" PM are mainly physical, whereas the effects of toxic particles are both chemical and physical. The effects of dust deposited on plant surfaces or on soil are more likely to be associated with their chemistry than with the mass of deposited particles and are usually of more importance than any physical effects. The majority of the easily identifiable direct and indirect effects, other than climate-change impacts, occur in severely polluted areas around heavily industrialized point sources such as limestone quarries; cement kilns; and iron; lead, and various smelting factories. Studies of the direct effects of chemical additions to foliage in particulate deposition have found little or no effects of PM on foliar processes; however, both conifers and deciduous species have shown significant effects on leaf surface structures after exposure to simulated acid rain or mist at pH 3.5. Many experimental studies indicate that epicuticular waxes (which function to prevent water loss from plant leaves) can be destroyed by acid rain in a few weeks. This function is particularly crucial in conifers because of the longevity of evergreen foliage.

Though there has been no direct evidence of a physiological association between tree injury and exposure to metals, heavy metals have been implicated because their deposition pattern is correlated with forest decline. The role of heavy metals has been indicated by phytochelatin measurements. Phytochelatins are intracellular metal-binding peptides that act as indicator of metal stress. Because they are produced by plants as a response to sublethal concentrations of heavy metals, they can be used to indicate that heavy metals are involved in forest decline. Concentrations of the phytochelatins increased with altitude, as did forest decline, and they also increased across regions showing increased levels of forest injury.

Secondary organics formed in the atmosphere have been referred to under the following terms: toxic substances, pesticides, hazardous air pollutants (HAPS), air toxics, semivolatile organic compounds (SOCs), and persistent organic pollutants (POPS). The chemical substances listed under the above headings are not criteria pollutants controlled by NAAQS as cited under CAA Sections 108 and 109 (U.S. Code, 1991), but rather are controlled under CAA Sect.112,

Hazardous Air Pollutants. Their possible effects in the environment on humans and ecosystems are discussed in many other government documents and publications. They are mentioned in this chapter because, in the atmosphere many of the chemical compounds are partitioned between gas and particle phases and are deposited as particulate matter. As particles, they become airborne and can be distributed over a wide area and impact remote ecosystems. Some of the chemical compounds are of concern to humans because they may reach toxic levels in food chains of both animals and humans, whereas others tend to decrease or maintain the same toxicity as they move through the food chain.

An important characteristic of fine particles is their ability to affect the flux of solar radiation passing through the atmosphere directly, by scattering and absorbing solar radiation, and indirectly, by acting as cloud condensation nuclei that, in turn, influence the optical properties of clouds. Regional haze has been estimated to diminish surface solar visible radiation by approximately 8%. Crop yields have been reported as being sensitive to the amount of sunlight received, and crop losses have been attributed to increased airborne particle levels in some areas of the world.

The transmission of solar UV-B radiation through the Earth's atmosphere is controlled by ozone, clouds, and particles. The depletion of stratospheric ozone caused by the release of chlorofluorcarbons and other ozone-depleting substances has resulted in heightened concern regarding potentially serious increases in the amount of solar UV-B (SUVB) radiation reaching the Earth's surface. Plant species vary enormously in their response to UV-B exposures, and large differences in response also occur among different genotypes within a species. In general, dicotyledonous plants are more sensitive than monocotyledons from similar environments. In addition, plant responses may differ depending on stage of development. Because plants evolved under the selective pressure of ambient UV-B radiation in sunlight, they have developed adaptive mechanisms. Although inhibition of photosynthesis is a detrimental growth effect, flavonoid synthesis represents acclimation. Plants growing under full light have been shown to be protected against UV-B effects but not when growing under weak visible light. A common adaptation is alteration in leaf transmission properties, which results in attenuation of UV-B in the epidermis before it can reach the leaf interior.

Indirect effects of PM are considered of greatest significance because their deposition on the soil has altered nutrient cycling and inhibited nutrient uptake and changed the functioning,

structure and biodiversity of ecosystems. Indirect effects occur through the soil and result from the deposition of heavy metals, nitrates, sulfates, or acidic precipitation and their impact on the soil microbial community. The soil environment is one of the most dynamic sites of biological interaction in nature. Bacteria in the soil are essential components of the nitrogen and sulfur cycles that make these elements available for plant uptake. Fungi form mycorrhizae, a mutualistic symbiotic relationship, that is integral in mediating plant uptake of mineral nutrients. Changes in the soil environment that influence the role of the bacteria and fungi in nutrient cycling and availability determine plant and ecosystem response.

Major impacts of PM on soil environments occur through deposition of nitrates and sulfates and the acidifying effect of the H^+ ion associated with these compounds in wet and dry deposition. Although the soils of most of North American forest ecosystems are nitrogen limited, there are some forests that exhibit severe symptoms of nitrogen saturation. They include the high-elevation, spruce-fir ecosystems in the Appalachian Mountains; the eastern hardwood watersheds at the Fernow Experimental Forest near Parsons, WV; the mixed conifer forest and chaparral watershed with high smog exposure in the Los Angeles Air Basin; the high-elevation alpine watersheds in the Colorado Front Range; and a deciduous forest in Ontario, Canada.

Nitrogen saturation results when additions to soil background nitrogen (nitrogen loading) exceed the capacity of plants and soil microorganisms to utilize and retain nitrogen. An ecosystem no longer functions as a sink under these circumstances. Possible ecosystem responses to nitrate saturation, as postulated by Aber and his coworkers, include (1) a permanent increase in foliar nitrogen and reduced foliar phosphorus and lignin because of the lower availability of carbon, phosphorus, and water; (2) reduced productivity in conifer stands caused by disruptions of physiological function; (3) decreased root biomass and increased nitrification and nitrate leaching; (4) reduced soil fertility, the results of increased cation leaching, increased nitrate and aluminum concentrations in streams, and decreased water quality. Saturation implies that some resource other than nitrogen is limiting biotic function. Water and phosphorus for plants and carbon for microorganisms are the resources most likely to be the secondary limiting factors. The appearance of nitrogen in soil solution is an early symptom of excess nitrogen. In the final stage, disruption of forest structure becomes visible.

Changes in nitrogen supply can have a considerable impact on an ecosystem's nutrient balance. Increases in soil nitrogen play a selective role. Plant succession patterns and

biodiversity are affected significantly by chronic nitrogen additions in some ecosystems. Long-term nitrogen fertilization studies in both New England and Europe suggest that some forests receiving chronic inputs of nitrogen may decline in productivity and experience greater mortality. For example, long-term fertilization experiments at Mount Ascutney, VT, suggest that declining coniferous forest stands with slow nitrogen cycling may be replaced by deciduous fast-growing forests that cycle nitrogen rapidly. Excess nitrogen inputs to unmanaged heathlands in the Netherlands also have been found to result in nitrophilous grass species replacing slower growing heath species. Over the past several decades, the composition of plants in the forest herb layers had been shifting toward species commonly found on nitrogen-rich areas. It also was observed that the fruiting bodies of mycorrhizal fungi had decreased in number.

Notable impacts of excess nitrogen deposition also have been observed with regard to aquatic systems. For example, atmospheric nitrogen deposition into soils in watershed areas feeding into estuarine sound complexes (e.g., the Pamlico Sound of North Carolina) appear to contribute to excess nitrogen flows in runoff (especially during and after heavy rainfall events such as hurricanes). Together with excess nitrogen runoff from agricultural practices or other uses (e.g., fertilization of lawns or gardens), massive influxes of such nitrogen into watersheds and sounds can lead to dramatic decreases in water oxygen and increases in algae blooms that can cause extensive fish kills and damage to commercial fish and sea food harvesting.

Acidic deposition has played a major role in soil acidification in some areas of Sweden, elsewhere in Europe, and in eastern North America. Soil acidification and its effects result from deposition of nitrates, sulfates, and associated H^+ ion. A major concern is that soil acidity will lead to nutrient deficiency. Growth of tree species can be affected when high aluminum-to-nutrient ratios limit uptake of calcium and magnesium and create a nutrient deficiency. Calcium is essential in the formation of wood and the maintenance of cells (the primary plant tissues necessary for tree growth), and it must be dissolved in soil water to be taken up by plants. Acidic deposition can increase aluminum concentrations in soil water by lowering the pH in aluminum-rich soils through dissolution and ion-exchange processes. Aluminum in soil can then be taken up by roots more readily than calcium because of its greater affinity for negatively charged surfaces. Tree species can be adversely affected if altered Ca/Al ratios impair Ca or Mg uptake.

Overall, then, PM produced by human activities has the potential to cause the loss of ecosystem biodiversity in ways that reduces the ability of ecosystems to provide the services that

society requires to sustain life. The major impacts of PM on ecosystems are the indirect effects that occur through the soil and affect plant growth, vigor, and reproduction. Mineral nutrient cycling can be altered by the deposition of heavy metals. The deposition of nitrogen and sulfur and the acidifying effects of the two in association with the H^+ ion in precipitation also alter biogeochemical cycling, cause soil acidification, alter the Ca/Al ratio, and impact the growth of vegetation and forest trees, in particular. Leaching of nitrates and other minerals through runoff can impact coastal and aquatic wetlands and, thus, influence their ability to produce the products and services necessary for existence of human society.

4.6.2 Particulate Matter-Related Effects on Visibility

Visibility is defined as the degree to which the atmosphere is transparent to visible light and the clarity and color fidelity of the atmosphere. Visual range is the farthest distance a black object can be distinquished against the horizontal sky. Visibility impairment is any humanly perceptible change in visibility. For regulatory purposes, visibility impairment, characterized by light extinction, visual range, contrast, and coloration, is classified into two principal forms: (1) “reasonably attributable” impairment, attributable to a single source or small group of sources, and (2) regional haze, any perceivable change in visibility caused by a combination of many sources over a wide geographical area.

Visibility is measured by human observation, light scattering by particles, the light extinction-coefficient and parameters related to the light-extinction coefficient (visual range and deciview scale), the light scattering coefficient, and fine PM concentrations. The air quality within a sight path will affect the illumination of the sight path by scattering or absorbing solar radiation before it reaches the Earth’s surface. The rate of energy loss with distance from a beam of light is the light extinction coefficient. The light extinction coefficient is the sum of the coefficients for light absorption by gases (σ_{ag}), light scattering by gases (σ_{sg}), light absorption by particles (σ_{ap}), and light scattering by particles (σ_{sp}). Atmospheric particles are frequently divided into fine and coarse particles. Corresponding coefficients for light scattering and absorption by fine and coarse particles are σ_{sfp} and σ_{afp} and σ_{scp} and σ_{acp}, respectively. Visibility within a sight path longer than approximately 100 km (60 mi) is affected by change in the optical properties of the atmosphere over the length of the sight path.

Visibility impairment is associated with airborne particle properties, including size distributions (i.e., fine particles in the 0.1- to 1.0-μm size range) and aerosol chemical composition, and with relative humidity. With increasing relative humidity, the amount of moisture available for absorption by particles increases, thus causing the particles to increase in both size and volume. As the particles increase in size and volume, the light scattering potential of the particles also generally increases. Visibility impairment is greatest in the eastern United States and Southern California. In the eastern United States, visibility impairment is caused primarily by light scattering by sulfate aerosols and, to a lesser extent, by nitrate particles and organic aerosols, carbon soot, and crustal dust. Haziness in the southeastern United States, caused by increased atmospheric sulfate, has increased by ca. 80% since the 1950s and is greatest in the summer months, followed by the spring and fall, and winter. Light scattering by nitrate aerosols is the major cause of visibility impairment in southern California. Nitrates contribute about 40% to the total light extinction in southern California and accounts for 10 to 20% of the total extinction in other U.S. areas.

Organic particles are the second largest contributors to light extinction in most U.S. areas. Organic carbon is the greatest cause of light extinction in the Pacific Northwest, Oregon, Idaho, and Montana, accounting for 40 to 45% of the total extinction. Also, organic carbon contributes between 15 to 20% to the total extinction in most of the western United States and 20 to 30% in the remaining U.S. areas.

Coarse mass and soil, primarily considered “natural extinction”, is responsible for some of the visibility impairment in northern California and Nevada, Oregon, southern Idaho, and western Wyoming. Dust transported from southern California and the subtropics has been associated with regional haze in the Grand Canyon and other southwestern U.S. class I areas.

4.6.3 Particulate Matter-Related Effects on Materials

Building materials (metals, stones, cements, and paints) undergo natural weathering processes from exposure to environmental elements (wind, moisture, temperature fluctuations, sun light, etc.). Metals form a protective film that protects against environmentally induced corrosion. The natural process of metal corrosion from exposure to natural environmental elements is enhanced by exposure to anthropogenic pollutants, in particular SO_2, rendering the protective film less effective.

Dry deposition of SO_2 enhances the effects of environmental elements on calcereous stones (limestone, marble, and cement) by converting calcium carbonate (calcite) to calcium sulfate dihydrate (gypsum). The rate of deterioration is determined by the SO_2 concentration, the stone's permeability and moisture content, and the deposition rate; however, the extent of the damage to stones produced by the pollutant species apart from the natural weathering processes is uncertain. Sulfur dioxide also has been found to limit the life expectancy of paints by causing discoloration and loss of gloss and thickness of the paint film layer.

A significant detrimental effect of particle pollution is the soiling of painted surfaces and other building materials. Soiling changes the reflectance of a material from opaque and reduces the transmission of light through transparent materials. Soiling is a degradation process that requires remediation by cleaning or washing, and, depending on the soiled surface, repainting. Available data on pollution exposure indicates that particles can result in increased cleaning frequency of the exposed surface and may reduce the life usefulness of the material soiled. Attempts have been made to quantify the pollutants exposure levels at which materials damage and soiling have been perceived. However, to date, insufficient data are available to advance our knowledge regarding perception thresholds with respect to pollutant concentration, particle size, and chemical composition.

4.6.4 Effects of Atmospheric Particulate Matter on the Transmission of Solar Ultraviolet Radiation and Global Warming Processes

Extensive potential future impacts on human health and the environment are projected to occur because of increased transmission of solar ultraviolet radiation (UV-B) through the Earth's atmosphere, secondary to stratospheric ozone depletion resulting from anthropogenic emissions of chlorofluorcarbons (CFCs), halons, and certain other gases. However, the estimation of the likely future extent of detrimental effects caused by increased penetration of solar UV-B to the Earth's surface is complicated by atmospheric particle effects, which vary depending on size and composition of particles that can differ substantially over different geographic areas and from season to season over the same area. Also, atmospheric particles greatly complicate projections of future trends in global warming processes because of emissions of greenhouse gases; consequent increases in global mean temperature, and resulting changes in regional and local

weather patterns; and mainly deleterious (but some beneficial) location-specific human health and environmental impacts.

The physical processes (i.e., scattering and absorption) responsible for airborne particle effects on transmission of solar ultraviolet and visible radiation are the same as those responsible for visibility degradation. Scattering of solar radiation back to space and absorption of solar radiation determine the effects of an aerosol layer on solar radiation. The transmission of solar UV-B radiation is affected strongly by atmospheric particles. Measured attenuations of UV-B under hazy conditions range up to 37% of the incoming solar radiation. Measurements relating variations in PM mass directly to UV-B transmission are lacking. Particles also can affect the rates of photochemical reactions occurring in the atmosphere. Depending on the amount of absorbing substances in the particles, photolysis rates either can be increased or decreased.

In addition to direct climate effects through the scattering and absorption of solar radiation, particles also exert indirect effects on climate by serving as cloud condensation nuclei, thus affecting the abundance and vertical distribution of clouds. The direct and indirect effects of particles appear to have significantly offset the global warming effects caused by the buildup of greenhouse gases because the onset of the Industrial Revolution, on a globally averaged basis. However, because the lifetime of particles is much shorter than that required for complete mixing within the Northern Hemisphere, the climate effects of particles generally are felt much less homogeneously than are the effects of long-lived greenhouse gases.

Any effort to model the impacts of local alterations in particle concentrations on projected global climate change or consequent local and regional weather patterns would be subject to considerable uncertainty. This also would be the case for any projections of impacts of location-specific airborne PM alterations on potential human health or environmental effects associated with either increased atmospheric transmission of solar UV radiation or global warming secondary to accumulation of stratospheric ozone-depleting substances or “greenhouse gases.”

REFERENCES

Abdullah, U. M.; Iqbal, M. Z. (1991) Response of automobile, stone and cement particulate matters on stomatal clogging of plants. Geobios (Jodhpur, India) 18: 196-202.

Aber, J. D.; Nadelhoffer, K. J.; Steudler, P.; Melillo, J. M. (1989) Nitrogen saturation in northern forest ecosystems: excess nitrogen from fossil fuel combustion may stress the biosphere. Bioscience 39: 378-386.

Aber, J.; McDowell, W.; Nadelhoffer, K.; Magill, A.; Berntson, G.; Kamakea, M.; McNulty, S.; Currie, W.; Rustad, L.; Fernandez, I. (1998) Nitrogen saturation in temperate forest ecosystems. BioScience 48: 921-934.

Ainsleigh, H. G. (1993) Beneficial effects of sun exposure on cancer mortality. Prev. Med. 22: 132-140.

Alexander, M. (1977) Introduction to soil microbiology. 2nd ed. New York, NY: John Wiley & Sons; pp. 225-230; 239-246; 272-286; 355-356.

Allen, M. F. (1991) The ecology of mycorrhizae. Cambridge, MA: Cambridge University Press.

Allen, A. G.; Harrison, R. M.; Nicholson, K. W. (1991) Dry deposition of fine aerosol to a short grass surface. Atmos. Environ. Part A 25: 2671-2676.

Allen, E. R.; Cabrera, N.; Kim, J.-C. (1994) Atmospheric deposition studies in rural forested areas of the southeastern United States. Presented at: 87th annual meeting & exhibition of the Air & Waste Management Association; June; Cincinnati, OH. Pittsburgh, PA: Air & Waste Management Association; paper no. 94-WP88.02.

Allen, E. B.; Padgett, P. E.; Bytnerowicz, A.; Minich, R. (1998) Nitrogen deposition effects on coastal sage vegetation of southern California. USDA Forest Service Gen. Tech. Rep. PSW-GTR-166, pp. 131-139.

Altshuller, A. P.; Linthurst, R. A., eds. (1984) Acidic deposition phenomenon and its effects: critical assessment review papers; volume I, atmospheric sciences; volume II, effects sciences. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development; EPA report nos. EPA/600/8-83/016aF-bF. Available from: NTIS, Springfield, VA; PB85-100030 and PB85-100048.

Andersen, C. P.; Rygiewicz, P. T. (1991) Stress interactions and mycorrhizal plant response: understanding carbon allocation priorities. Environ. Pollut. 73: 217-244.

Anderson, J. B.; Baumgardner, R. E.; Mohnen, V. A.; Bowser, J. J. (1999) Cloud chemistry in the eastern United States, as sampled from three high-elevation sites along the Appalachian Mountains. Atmos. Environ. 33: 5105-5114.

Aneja, V. P.; Murthy, A. B. (1994) Monitoring deposition of nitrogen-containing compounds in a high-elevation forest canopy. Air Waste 44: 1109-1115.

Antonovics, J.; Bradshaw, A. D.; Turner, R. G. (1971) Heavy metal tolerance in plants. In: Cragg, J. B., ed. Advances in ecological research. Volume 7. London, United Kingdom: Academic Press; pp. 1-85.

April, R.; Newton, R. (1992) Mineralogy and mineral weathering. In: Johnson, D. W.; Lindberg, S. E., eds. Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study. New York, NY: Springer-Verlag; pp. 378-425. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies: analysis and synthesis: v. 91).

Arnott, W. P.; Moosmüller, H.; Rogers, C. F.; Jin, T.; Bruch, R. (1999) Photoacoustic spectrometer for measuring light absorption by aerosol; instrument description. Atmos. Environ. 33: 2845-2852.

Askey, A.; Lyon, S. B.; Thompson, G. E.; Johnson, J. B.; Wood, G. C.; Sage, P. W.; Cooke, M. J. (1993) The effect of fly-ash particulates on the atmospheric corrosion of zinc and mild steel. Corros. Sci. 34: 1055-1081.

Ausset, P.; Bannery, F.; Del Monte, M.; Lefevre, R. A. (1998) Recording of pre-industrial atmospheric environment by ancient crusts on stone monuments. Atmos. Environ. 32: 2859-2863.

Ayensu, E.; Van R. Claasen, D.; Collins, M.; Dearing, A.; Fresco, L.; Gadgil, M.; Gitay, H.; Glaser, G.; Juma, C.; Krebs, J.; lenton, R.; Lubchenco, J.; McNeely, J. A.; Mooney, H. A.; Pinstrup-Andersen, P.; Ramos, M.; Raven, P.; Reid, W. V.; Samper, C.; Sarukhan, J.; Schei, P.; Tundisi, J. G.; Watson, R. T.; Guanhua, X.; Zakri, A. H. (1999) International ecosystem assessment. Science 286: 685-686.

Babich, H.; Stotzky, G. (1978) Effects of cadmium on the biota: influence of environmental factors. In: Perlman, D., ed. Advances in applied microbiology, volume 23. New York, NY: Academic Press; pp. 55-117.

Bache, D. H. (1979a) Particle transport within plant canopies - I. a framework for analysis. Atmos. Environ. 13: 1257-1262.

Bache, D. H. (1979b) Particulate transport within plant canopies - II. prediction of deposition velocities. Atmos. Environ. 13: 1681-1687.

Baedecker, P. A.; Edney, E. O.; Moran, P. J.; Simpson, T. C.; Williams, R. S. (1991) Effects of acidic deposition on materials. In: Irving, P. M., ed. Acidic deposition: state of science and technology, volume III: terrestrial, materials, health and visibility effects. Washington, DC: The U.S. National Acid Precipitation Assessment Program. (State of science and technology report no. 19).

Baedecker, P. A.; Reddy, M. M.; Reimann, K. J.; Sciammarella, C. A. (1992) Effects of acidic deposition on the erosion of carbonate stone--experimental results from the U.S. National Acid Precipitation Assessment Program (NAPAP). Atmos. Environ. Part B 26: 147-158.

Baldocchi, D. (1988) A multi-layer model for estimating sulfur dioxide deposition to a deciduous oak forest canopy. Atmos. Environ. 22: 869-884.

Baron, J. S.; Rueth, H. M.; Wolfe, A. M.; Nydick, K. R.; Allstott, E. J.; Minear, J. T.; Moraska, B. (2000) Ecosystem responses to nitrogen deposition in the Colorado front range. Ecosystems 3: 352-368.

Barone, J. B.; Cahill, T. A.; Eldred, R. A.; Flocchini, R. G.; Shadoan, D. J.; Dietz, T. M. (1978) A multivariate statistical analysis of visibility degradation at four California cities. Atmos. Environ. 12: 2213-2221.

Barrie, L. A.; Anlauf, K.; Wiebe, H. A.; Fellin, P. (1984) Acidic pollutants in air and precipitation at selected rural locations in Canada. In: Hicks, B. B., ed. Deposition both wet and dry: [papers presented at the American Chemical Society acid rain symposium]; March 1982; Las Vegas, NV. Boston, MA: Butterworth Publishers; pp. 15-35. (Teasley, J. I., ed. Acid precipitation series: v. 4).

Barton, K. (1958) The influence of dust on atmospheric corrosion of metals. Werkst. Korros. 8/9: 547-549.

Beier, C. (1991) Separation of gaseous and particulate dry deposition of sulfur at a forest edge in Denmark. J. Environ. Qual. 20: 460-466.

Beier, C.; Gundersen, P.; Rasmussen, L. (1992) A new method for estimation of dry deposition of particles based on throughfall measurements in a forest edge. Atmos. Environ. Part A 26: 1553-1559.

Bellan, L. M.; Salmon, L. G.; Cass, G. R. (2000) A study on the human ability to detect soot deposition onto works of art. Environ. Sci. Technol. 34: 1946-1952.

Bennett, J. H.; Hill, A. C. (1973) Absorption of gaseous air pollutants by a standardized plant canopy. J. Air Pollut. Control Assoc. 23: 203-206.

Bennett, J. H.; Hill, A. C. (1975) Interactions of air pollutants with canopies of vegetation. In: Mudd, J. B.; Kozlowski, T. T., eds. Responses of plants to air pollution. New York, NY: Academic Press; pp. 273-306.

Bernard, S. M.; McGeehin, M. G.; Patz, J. A., eds. (2001) Health: the potential consequences of climate variability and change. Environ. Health Perspect. 109(suppl. 2): 175-233.

Bewley, R. J. F.; Parkinson, D. (1984) Effects of sulphur dioxide pollution on forest soil microorganisms. Can. J. Microbiol. 30: 179-185.

Biefer, G. J. (1981) Atmospheric corrosion of steel in the Canadian Arctic. Mater. Perform. 20: 16-19.

Billings, W. D. (1978) Plants and the ecosystem. 3rd ed. Belmont, CA: Wadsworth Publishing Company, Inc.; pp. 1-62, 83-108.

Binenko, V. I.; Harshvardhan, H. (1993) Aerosol effects in radiation transfer. In: Jennings, S. G., ed. Aerosol effects on climate. Tucson, AZ: University of Arizona Press; pp. 190-232.

Binkley, D.; Valentine, D.; Wells, C.; Valentine, U. (1989) An empirical analysis of the factors contributing to 20-year decrease in soil pH in an old-field plantation of loblolly pine. Biogeochemistry 8: 39-54.

Bjorn, L. O. (1996) Effects of ozone depletion and increased UV-B on terrestrial ecosystems. Int. J. Environ. Stud. 51: 217-243.

Blackburn, T. E.; Bairai, S. T.; Stedman, D. H. (1992) Solar photolysis of ozone to singlet D oxygen atoms. J. Geophys. Res. 97: 10,109-10,117.

Blaschke, H. (1990) Mycorrhizal populations and fine root development on Norway spruce exposed to controlled doses of gaseous pollutants and simulated acidic rain treatments. Environ. Pollut. 68: 409-418.

Bobbink, R. (1998) Impacts of tropospheric ozone and airborne nitrogenous pollutants on natural and semi-natural ecosystems: a commentary. New Phytol. 139: 161-168.

Bobbink, R.; Hornung, M.; Roelofs, J. G. M. (1998) The effects of air-borne nitrogen pollutants on species diversity in natural and semi-natural European vegetation. J. Ecol. 86: 717-738.

Bolle, H.-J.; Seiler, W.; Bolin, B. (1986) Other greenhouse gases and aerosols: assessing their role for atmospheric radiative transfer. In: Bolin, B.; Doos, B. R.; Jager, J.; Warrick, R. A., eds. The greenhouse effect, climatic change, and ecosystems. Chichester, United Kingdom: John Wiley and Sons; pp. 157-203. (SCOPE 29).

Bondietti, E. A.; McLaughlin, S. B. (1992) Evidence of historical influences of acidic deposition on wood and soil chemistry. In: Johnson, D. W.; Lindberg, S. E., eds. Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study. New York, NY: Springer-Verlag: pp. 358-377. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies analysis and synthesis: v. 91).

Bowden, R. D.; Geballe, G. T.; Bowden, W. B. (1989) Foliar uptake of ^{15}N from simulated cloud water by red spruce (*Picea rubens*) seedlings. Can. J. For. Res. 19: 382-386.

Bowman, W. D. (2000) Biotic controls over ecosystem response to environmental change in alpine tundra of the Rocky Mountains. Ambio 29: 396-400.

Bowman, W. D.; Steltzer, H. (1998) Positive feedbacks to anthropogenic nitrogen deposition in Rocky Mountain alpine tundra. Ambio 27: 514-517.

Boyce, S. G. (1954) The salt spray community. Ecol. Monogr. 24: 29-67.

Bradley, R.; Burt, A. J.; Read, D. J. (1981) Mycorrhizal infection and resistance to heavy metal toxicity in Calluna vulgaris. Nature (London) 292: 335-337.

Brandt, C. J.; Rhoades, R. W. (1972) Effects of limestone dust accumulation on composition of a forest community. Environ. Pollut. 3: 217-225.

Brandt, C. J.; Rhoades, R. W. (1973) Effects of limestone dust accumulation on lateral growth of forest trees. Environ. Pollut. 4: 207-213.

Brook, J. R.; Zhang, L.; Di-Giovanni, F.; Padro, J. (1999) Description and evaluation of a model of deposition velocities for routine estimates of air pollutant dry deposition over North America. Part I: model development. Atmos. Environ. 33: 5037-5051.

Brühl, C.; Crutzen, P. J. (1988) On the disproportionate role of tropospheric ozone as a filter against solar UV-B radiation. Geophys. Res. Lett. 16: 703-706.

Brumme, R.; Leimcke, U.; Matzner, E. (1992) Interception and uptake of NH_4 and NO_3 from wet deposition by above-ground parts of young beech (*Fagus silvatica L.*) trees. Plant Soil 142: 273-279.

Brunet, J.; Diekmann, M.; Falkengren-Grerup, U. (1998) Effects of nitrogen deposition on field layer vegetation in south Swedish oak forests. Environ. Pollut. (Oxford, U.K.) 102(suppl. 1): 35-40.

Bugini, R.; Tabasso, M. L.; Realini, M. (2000) Rate of formation of black crusts on marble. A case study. J. Cult. Heritage 1: 111-116.

Businger, J. A. (1986) Evaluation of the accuracy with which dry deposition can be measured with current micrometeorological techniques. J. Clim. Appl. Meteorol. 25: 1100-1124.

Butlin, R. N.; Coote, A. T.; Devenish, M.; Hughes, I. S. C.; Hutchens, C. M.; Irwin, J. G.; Lloyd, G. O.; Massey, S. W.; Webb, A. H.; Yates, T. J. S. (1992a) Preliminary results from the analysis of metal samples from the National Materials Exposure Programme (NMEP). Atmos. Environ. Part B 26: 199-206.

Butlin, R. N.; Coote, A. T.; Devenish, M.; Hughes, I. S. C.; Hutchens, C. M.; Irwin, J. G.; Lloyd, G. O.; Massey, S. W.; Webb, A. H.; Yates, T. J. S. (1992b) Preliminary results from the analysis of stone tablets from the National Materials Exposure Programme (NMEP). Atmos. Environ. Part B 26: 189-198.

Bytnerowicz, A.; Fenn, M. E. (1996) Nitrogen deposition in California forests: a review. Environ. Pollut. 92: 127-146.

Bytnerowicz, A.; Miller, P. R.; Olszyk, D. M. (1987a) Dry deposition of nitrate, ammonium and sulfate to a *Ceanothus crassifolius* canopy and surrogate surfaces. Atmos. Environ. 21: 1749-1757.

Bytnerowicz, A.; Miller, P. R.; Olszyk, D. M.; Dawson, P. J.; Fox, C. A. (1987b) Gaseous and particulate air pollution in the San Gabriel Mountains of southern California. Atmos. Environ. 21: 1805-1814.

Cahill, T. A.; Eldred, R. A.; Wakabayashi, P. H. (1996) Trends in fine particle concentrations at Great Smoky Mountains National Park. For presentation at: Annual meeting of the Air & Waste Management Association; June; Nashville, TN. Pittsburgh, PA: Air & Waste Management Association; submitted paper no. 96-MP1A.05.

Campbell, G. G.; Schurr, G. G.; Slawikowski, D. E.; Spence, J. W. (1974) Assessing air pollution damage to coatings. J. Paint Technol. 46: 59-71.

Camuffo, D. (1995) Physical weathering of stones. In: Saiz-Jimenez, C., ed. The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation; January 1994; Sevilla, Spain. Sci. Total Environ. 167: 1-14.

Cape, J. N. (1993) Direct damage to vegetation caused by acid rain and polluted cloud: definition of critical levels for forest trees. Environ. Pollut. 82: 167-180.

Cape, J. N.; Sheppard, L. J.; Fowler, D.; Harrison, A. F.; Parkinson, J. A.; Dao, P.; Paterson, I. S. (1992) Contribution of canopy leaching to sulphate deposition in a Scots pine forest. Environ. Pollut. 75: 229-236.

Carey, W. F. (1959) Atmospheric deposits in Britain: a study of dinginess. Int. J. Air Pollut. 2: 1-26.
Ceccotti, S. P.; Messick, D. L. (1997) A global review of crop requirements, supply, and environmental impact on nutrient sulphur balance. In: Cram, W. J.; De Kok, L. J.; Stulen, I.; Brunold, C.; Rennenberg, H., eds. Sulphur metabolism in higher plants. Leiden, The Netherlands: Backhuys Publishers; pp. 155-163.
Chamberlain, A. C. (1975) The movement of particles in plant communities. In: Monteith, J. L., ed. Vegetation and the atmosphere: v. 1, principles. London, United Kingdom: Academic Press; pp. 155-203.
Chamberlain, A. C.; Little, P. (1981) Transport and capture of particles by vegetation. In: Grace, J.; Ford, E. D.; Jarvis, P. G., eds. Plants and their atmospheric environment: the 21st symposium of the British Ecological Society; 1979; Edinburgh, United Kingdom. Oxford, United Kingdom: Blackwell Scientific Publications; pp. 147-173.
Chameides, W. L.; Yu, H.; Liu, S. C.; Bergin, M.; Zhou, X.; Mearns, L.; Wang, G.; Kiang, C. S.; Saylor, R. D.; Luo, C.; Huang, Y.; Steiner, A.; Giorgi, F. (1999) Case study of the effects of atmospheric aerosols and regional haze on agriculture: an opportunity to enhance crop yields in China through emission controls? Proc. Natl. Acad. Sci. U. S. A. 96: 13626-13633.
Chapin, F. S., III. (1991) Integrated responses of plants to stress: a centralized system of physiological responses. Bioscience 41: 29-36.
Chapin, F. S., III; Ruess, R. W. (2001) The roots of the matter. Nature (London) 411: 749-752.
Chapin, F. S., III; Bloom, A. J.; Field, C. B.; Waring, R. H. (1987) Plant responses to multiple environmental factors. BioScience 37: 49-57.
Chapin, F. S., III; Walker, B. H.; Hobbs, R. J.; Hooper, D. U.; Lawton, J. H.; Sala, O. E.; Tilman, D. (1997) Biotic control over functioning ecosystems. Science 277: 500-504.
Chapin, F. S., III; Sala, O. E.; Burke, I. C.; Grime, J. P.; Hooper, D. U.; Lauenroth, W. K.; Lombard, A.; Mooney, H. A.; Mosier, A. R.; Naeem, S.; Pacala, S. W.; Roy, J.; Steffen, W. L.; Tilman, D. (1998) Ecosystem consequences of changing biodiversity. BioScience 48: 45-52.
Charlson, R. J.; Lovelock, J. E.; Andreae, M. O.; Warren, S. G. (1987) Oceanic phytoplankton, atmospheric sulfphur, cloud albedo and climate. Nature (London) 326: 655-661.
Charlson, R. J.; Schwartz, S. E.; Hales, J. M.; Cess, R. D.; Coakley, J. A., Jr.; Hansen, J. E.; Hofmann, D. J. (1992) Climate forcing by anthropogenic aerosols. Science (Washington, DC) 255: 423-430.
Chevone, B. I.; Herzfeld, D. E.; Krupa, S. V.; Chappelka, A. H. (1986) Direct effects of atmospheric sulfate deposition on vegetation. J. Air Pollut. Control Assoc. 36: 813-815.
Choudhury, B. J. (1987) Relationships between vegetation indices, radiation absorption, and net photosynthesis evaluated by a sensitivity analysis. Remote Sens. Environ. 22: 209-233.
Chughtai, A. R.; Williams, G. R.; Atteya, M. M. O.; Miller, N. J.; Smith, D. M. (1999) Carbonaceous particle hydration. Atmos. Environ. 33: 2679-2687.
Clarke, J. F.; Edgerton, E. S.; Martin, B. E. (1997) Dry deposition calculations for the clean air status and trends network. Atmos. Environ. 31: 3667-3678.
Clarkson, D. T.; Sanderson, J. (1971) Inhibition of the uptake and long-distance transport of calcium by aluminum and other polyvalent cations. J. Exp. Bot. 22: 837-851.
Clevenger, T. E.; Saiwan, C.; Koirtyohann, S. R. (1991) Lead speciation of particles on air filters collected in the vicinity of a lead smelter. Environ. Sci. Technol. 25: 1128-1133.
Clough, W. S. (1975) The deposition of particles on moss and grass surfaces. Atmos. Environ. 9: 1113-1119.
Cocker, D. R., III; Whitlock, N. E.; Flagen, R. C.; Seinfield, J. H. (2001) Hygroscopic properties of Pasadena, California aerosol. Aerosol Sci. Technol. 35: 637-647.
Cole, D. W.; Johnson, D. W. (1977) Atmospheric sulfate additions and cation leaching in a Douglas fir ecosystem. Water Resour. Res. 13: 313-317.
Collett, J. L., Jr.; Hoag, K. J.; Sherman, D. E.; Bator, A.; Richards, L. W. (1999) Spatial and temporal variations in San Joaquin Valley fog chemistry. Atmos. Environ. 33: 129-140.
Cooke, R. U.; Gibbs, G. B. (1994) Crumbling heritage? Studies of stone weathering in polluted atmospheres. Atmos. Environ. 28: 1355-1356.
Costanza, R.; d'Arge, R.; De Groot, R.; Farber, S.; Grasso, M.; Hannon, B.; Limburg, K.; Naeem, S.; O'Neill, R. V.; Paruelo, J.; Raskin, R. G.; Sutton, P.; Van Den Belt, M. (1997) The value of the world's ecosystem services and natural capital. Nature (London) 387: 253-259.
Cotrufo, M. F.; De Santo, A. V.; Alfani, A.; Bartoli, G.; De Cristofaro, A. (1995) Effects of urban heavy metal pollution on organic matter decomposition in *Quercus ilex L.* woods. Environ. Pollut. 89: 81-87.

Cowling, J. E.; Roberts, M. E. (1954) Paints, varnishes, enamels, and lacquers. In: Greathouse, G. A.; Wessel, C. J., eds. Deterioration of materials: causes and preventive techniques. New York, NY: Reinhold Publishing Corporation; pp. 596-645.

Cramer, S. D.; McDonald, L. G.; Bhagia, G.; Flinn, D. R.; Linstrom, P. J.; Carter, J. P. (1989) Effects of acid deposition on the atmospheric corrosion of structural metals. Research Triangle Park, NC: U.S. Environmental Protection Agency; internal report. Incorporated in: Irving, P. M., ed. Acidic Deposition: State of Science and Technology. Volume III: Terrestrial, Materials, Health and Visibility Effects; report 19, section 2. Washington, DC: National Acid Precipitation Assessment Program.

Creighton, P. J.; Lioy, P. J.; Haynie, F. H.; Lemmons, T. J.; Miller, J. L.; Gerhart, J. (1990) Soiling by atmospheric aerosols in an urban industrial area. J. Air Waste Manage. Assoc. 40: 1285-1289.

Cronan, C. S.; Grigal, D. F. (1995) Use of calcium/aluminum ratios as indicators of stress in forest ecosystems. J. Environ. Qual. 24: 209-226.

Cullen, J. J.; Neale, P. J. (1997) Biological weighting functions for describing the effects of ultraviolet radiation on aquatic ecosystems. In: Häder, D.-P., ed. The effects of ozone depletion on aquatic ecosystems. Austin, TX: Academic Press, pp. 97-118.

Dahlgren, R. A. (1994) Soil acidification and nitrogen saturation from weathering of ammonium-bearing rock. Nature 368: 838-841.

Daily, G. C. (1997) Introduction: what are ecosystem services? In: Daily, G. C., ed. Nature's services: societal dependence on natural ecosystems. Washington, DC: Island Press; pp. 1-10.

Daily, G. C.; Ehrlich, P. R. (1999) Managing Earth's ecosystems: an interdisciplinary challenge. Ecosystems 2: 277-280.

Darley, E. F. (1966) Studies on the effect of cement-kiln dust on vegetation. J. Air Pollut. Control Assoc. 16: 145-150.

Dasch, J. M. (1987) Measurement of dry deposition to surfaces in deciduous and pine canopies. Environ. Pollut. 44: 261-277.

Dasch, J. M.; Cadle, S. H. (1985) Wet and dry deposition monitoring in southeastern Michigan. Atmos. Environ. 19: 789-796.

Dasch, J. M.; Cadle, S. H. (1986) Dry deposition to snow in an urban area. Water Air Soil Pollut. 29: 297-308.

Dässler, H.-G.; Ranft, H.; Rehn, K.-H. (1972) Zur Widerstandsfähigkeit von Gehölzen gegenüber Fluorverbindungen und Schwefeldioxid [The susceptibility of woody plants exposed to fluorine compounds and SO_2]. Flora (Jena) 161: 289-302.

Davidson, C. I.; Friedlander, S. K. (1978) A filtration model for aerosol dry deposition: application to trace metal deposition from the atmosphere. J. Geophys. Res. C: Oceans Atmos. 83: 2343-2352.

Davidson, C. I.; Wu, Y.-L. (1989) Dry deposition of trace elements. In: Pacyna, J.M.; Ottar, B., eds. Control and fate of atmospheric trace metals: proceedings of the NATO advanced research workshop on fate and control of toxic metals in the atmosphere; September 1988; Oslo, Norway. Boston, MA: Kluwer Academic Publishers; pp. 147-202. (NATO ASI series C, mathematical and physical sciences: v. 268).

Davidson, C. I.; Wu, Y.-L. (1990) Dry deposition of particles and vapors. In: Lindberg, S. E.; Page, A. L.; Norton, S. A., eds. Acidic precipitation: v. 3, sources, deposition, and canopy interactions. New York, NY: Springer-Verlag; pp. 103-216.

Davidson, C. I.; Miller, J. M.; Pleskow, M. A. (1982) The influence of surface structure on predicted particle dry deposition to natural grass canopies. Water Air Soil Pollut. 18: 25-43.

Davidson, C. I.; Tang, W.; Finger, S.; Etyemezian, V.; Striegel, M. F.; Sherwood, S. I. (2000) Soiling patterns on a tall limestone building: changes over 60 years. Environ. Sci. Technol. 34: 560-565.

Davis, G. D.; Shaw, B. A.; Arah, C. O.; Fritz, T. L.; Moshier, W. C.; Simpson, T. C.; Moran, P. J.; Zankel, K. L. (1990) Effects of SO_2 deposition on painted steel surfaces. Surf. Interface Anal. 15: 107-112.

Day, D. E.; Malm, W. C.; Kreidenweis, S. M.; Tree, R. M. (1997) Aerosol light scattering measurements: a comparison of differently configured optec nephelometers. In: Visual air quality: aerosols and global radiation balance: proceedings of a specialty conference sponsored by the Air & Waste Management Association and the American Geophysical Union; September; Bartlett, NH. Pittsburgh, PA: Air & Waste Management Association; pp. 952-962. (A&WMA publication VIP-76).

DeHayes, D. H.; Schaberg, P. G.; Hawley, G. J.; Strimbeck, G. R. (1999) Acid rain impacts on calcium nutrition and forest health. Bioscience 49: 789-800.

Del Monte, M.; Vittori, O. (1985) Air pollution and stone decay: the case of Venice. Endeavour 9: 117-122.

Diakumaku, E.; Gorbushina, A. A.; Krumbein, W. E.; Panina, L.; Soukharjevski, S. (1995) Black fungi in marble and limestones--an aesthetical, chemical and physical problem for the conservation of monuments. In: Saiz-Jimenez, C., ed. The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation; January 1994; Sevilla, Spain. Sci. Total Environ. 167: 295-304.

Dickerson, R. R.; Kondragunta, S.; Stenchikov, G.; Civerolo, K. L.; Doddridge, B. G.; Holben, B. N. (1997) The impact of aerosols on solar ultraviolet radiation and photochemical smog. Science (Washington, DC) 278: 827-830.

Dobbins, R. A.; Mulholland, G. W.; Bryner, N. P. (1994) Comparison of a fractal smoke optics model with light extinction measurements. Atmos. Environ. 28: 889-897.

Dolske, D. A. (1988) Dry deposition of airborne sulfate and nitrate to soybeans. Environ. Pollut. 53: 1-12.

Dolske, D. A. (1995) Deposition of atmospheric pollutants to monuments, statues, and buildings. Sci. Total Environ. 167: 15-31.

Dolske, D. A.; Gatz, D. F. (1984) Field intercomparison of sulfate dry deposition monitoring and measurement methods: preliminary results. In: Hicks, B. B., ed. Deposition both wet and dry: [papers presented at the American Chemical Society acid rain symposium]; March 1982; Las Vegas, NV. Boston, MA: Butterworth Publishers; pp. 121-131. (Teasley, J. I., ed. Acid precipitation series: v. 4).

Draaijers, G. P. J.; Erisman, J. W. (1993) Atmospheric sulphur deposition to forest stands: throughfall estimates compared to estimates from inference. Atmos. Environ. Part A 27: 43-55.

Draaijers, G. P. J.; Ivens, W. P. M. F.; Bleuten, W. (1988) Atmospheric deposition in forest edges measured by monitoring canopy throughfall. Water Air Soil Pollut. 42: 129-136.

Draaijers, G. P. J.; van Ek, R.; Meijers, R. (1992) Research on the impact of forest stand structure on atmospheric deposition. Environ. Pollut. 75: 243-249.

Dreisinger, B. R.; McGovern, P. C. (1970) Monitoring atmospheric sulphur dioxide and correlating its effects on crops and forests in the Sudbury area. In: Linzon, S. N., ed. Impact of air pollution on vegetation conference; April; Toronto, ON, Canada. Pittsburgh, PA: Air Pollution Control Association.

Driscoll, C. T.; Van Dreason, R. (1993) Seasonal and long-term temporal patterns in the chemistry of Adirondack lakes. Water Air Soil Pollut. 67: 319-344.

Driscoll, C. T.; Wyskowski, B. J.; DeStaffan, P.; Newton, R. M. (1989) Chemistry and transfer of aluminum in a forested watershed in the Adirondack region of New York, USA. In: Lewis, T. E., ed. Environmental chemistry and toxicology of aluminum. Chelsea, MI: Lewis Publishers, Inc.; pp. 83-105.

Driscoll, C. T.; Newton, R. M.; Gubala, C. P.; Baker, J. P.; Christensen, S. W. (1991) Adirondack mountains. In: Charles, D. F., ed. Acidic deposition and aquatic ecosystems: regional case studies. New York, NY: Springer-Verlag; pp. 133-202.

Driscoll, C. T.; Lawrence, G. B.; Bulger, A. J.; Butler, T. J.; Cronan, C. S.; Eagar, C.; Lambert, K. F.; Likens, G. E.; Stoddard, J. L.; Weathers, K. C. (2001) Acidic deposition in the northeastern United States: sources and inputs, ecosystem effects, and management strategies. BioScience 51: 180-198.

Duan, B.; Fairall, C. W.; Thomson, D. W. (1988) Eddy correlation measurements of the dry deposition of particles in wintertime. J. Appl. Meteorol. 27: 642-652.

Duyzer, J. H.; Bouman, A. M. M.; Van Aalst, R. M.; Diederen, H. S. M. A. (1987) Assessment of dry deposition fluxes of NH_3 and NH_4^+ over natural terrains. In: Asman, W. A. H.; Diederen, H. S. M. A., eds. Ammonia and acidification: proceedings of a symposium of the European Association for the Science of Air Pollution (EURASAP); April; Bilthoven, The Netherlands. European Association for the Science of Air Pollution; pp. 97-106.

Eaton, J. S.; Likens, G. E.; Bormann, F. H. (1973) Throughfall and stemflow chemistry in a northern hardwood forest. J. Ecol. 61: 495-508.

Edgerton, E. S.; Lavery, T. F.; Boksleitner, R. P. (1992) Preliminary data from the USEPA dry deposition network: 1989. Environ. Pollut. 75: 145-156.

Edgerton-Warburton, L. M.; Allen, E. B. (2000) Shifts in arbuscular mycorrhizal communities along an anthropogenic gradient nitrogen deposition gradient. Ecol. Appl. 10: 484-496.

Edney, E. O. (1989) Paint coatings: controlled field and chamber experiments. Research Triangle Park, NC: U.S. Environmental Protection Agency, Atmospheric Research and Exposure Assessment Laboratory; report no. EPA/600/3-89/032. Available from: NTIS, Springfield, VA; PB89-189849.

Edney, E. O.; Stiles, D. C.; Corse, E. W.; Wheeler, M. L.; Spence, J. W.; Haynie, F. H.; Wilson, W. E., Jr. (1988) Field study to determine the impact of air pollutants on the corrosion of galvanized steel. Mater. Perform. 27: 47-50.

Edney, E. O.; Cheek, S. F.; Corse, E. W.; Spence, J. W.; Haynie, F. H. (1989) Atmospheric weathering caused by dry deposition of acidic compounds. J. Environ. Sci. Health Part A 24: 439-457.

Edwards, R. S.; Claxton, S. M. (1964) The distribution of air-borne salt of marine origin in the Aberystwyth area. J. Appl. Ecol. 1: 253-263.

El-Shobokshy, M. S. (1985) The dependence of airborne particulate deposition on atmospheric stability and surface conditions. Atmos. Environ. 19: 1191-1197.

Eldering, A.; Cass, G. R.; Moon, K. C. (1994) An air monitoring network using continuous particle size distribution monitors: connecting pollutant properties to visibility via Mie scattering calculations. Atmos. Environ. 28: 2733-2749.

Eldering, A.; Hall, J. R.; Hussey, K. J.; Cass, G. R. (1996) Visibility model based on satellite-generated landscape data. Environ. Sci. Tech. 30: 361-370.

Eldred, R. A.; Cahill, T. A. (1994) Trends in elemental concentrations of fine particles at remote sites in the United States of America. Atmos. Environ. 28: 1009-1019.

Eldred, R. A.; Cahill, T. A.; Feeney, P. J. (1993) Comparison of independent measurements of sulfur and sulfate in the IMPROVE network. Presented at: 86th annual meeting and exhibition of the Air & Waste Management Association; June; Denver, CO. Pittsburgh, PA: Air & Waste Management Association; paper no. 93-RA-110.02.

Eldred, R. A.; Cahill, T. A.; Flocchini, R. G. (1997) Composition of $PM_{2.5}$ and PM_{10} aerosols in the IMPROVE network. In: Special issue on aerosols and atmospheric optics–part I, [published papers from the international specialty conference; September 1994; Snowbird, UT]. J. Air Waste Manage. Assoc. 47: 194-203.

Ellenberg, H. (1987) Floristic changes due to eutrophication. In: Asman, W. A. H.; Diederen, S. M. A., eds. Ammonia and acidification: proceedings of a symposium of the European Association for the Science of Air Pollution (EURASAP); April; Bilthoven, The Netherlands. European Association for the Science of Air Pollution; pp. 301-308.

Eller, B. M. (1977) Road dust induced increase of leaf temperature. Environ. Pollut. 13: 99-107.

Eriksson, P.; Johansson, L.-G.; Strandberg, H. (1993) Initial stages of copper corrosion in humid air containing SO_2 and NO_2. J. Electrochem. Soc. 140: 53-59.

Erisman, J. W. (1993) Acid deposition onto nature areas in the Netherlands; part II. Throughfall measurements compared to deposition estimates. Water Air Soil Pollut. 71: 81-99.

Erisman, J.-W.; Vermetten, A. W. M.; Asman, W. A. H.; Waijers-IJpelaan, A.; Slanina, J. (1988) Vertical distribution of gases and aerosols: the behaviour of ammonia and related components in the lower atmosphere. Atmos. Environ. 22: 1153-1160.

Erisman, J. W.; Van Pul, A.; Wyers, P. (1994) Parametrization of surface resistance for the quantification of atmospheric deposition of acidifying pollutants and ozone. Atmos. Environ. 28: 2595-2607.

Erlick, C.; Frederick, J. E.; Saxena, V. K.; Wenny, B. N. (1998) Atmospheric transmission in the ultraviolet and visible: aerosols in cloudy atmospheres. J. Geophys. Res. 103: 31,541-31,556.

Eshleman, K. N.; Miller-Marshall, L. M.; Webb, J. R. (1995) Long-term changes in episodic acidification of streams in Shenandoah National Park, Virginia (U.S.A.). Water Air Soil Pollut. 85: 517-522.

Estupiñán, J. B.; Raman, S.; Crescenti, G. H.; Streicher, J. J.; Barnard, W. F. (1996) Effects of clouds and haze on UV-B radiation. J. Geophys. Res. 101: 16,807-16,816.

Evans, E. G.; Pitchford, M. (1991) Federal visibility monitoring in the eastern United States. Presented at: 84th annual meeting and exhibition of the Air & Waste Management Association; June; Vancouver, BC, Canada. Pittsburgh, PA: Air & Waste Management Association; paper no. 91-49.6.

Fahey, T. J.; Yavitt, J. B.; Joyce, G. (1988) Precipitation and throughfall chemistry in *Pinus contorta* ssp. *latifolia* ecosystems, southeastern Wyoming. Can. J. For. Res. 18: 337-345.

Falkengren-Grerup, U. (1987) Long-term changes in pH of forest soils in southern Sweden. Environ. Pollut. 43: 79-90.

Falkengren-Grerup, U. (1998) Nitrogen response of herbs and graminoids in experiments with simulated acid soil solution. Environ. Pollut. 102(suppl. 1): 93-99.

Farber, R. J.; Welsing, P. R.; Rozzi, C. (1994) PM_{10} and ozone control strategy to improve visibility in the Los Angeles Basin. Atmos. Environ. 28: 3277-3283.

Farmer, A. M. (1993) The effects of dust on vegetation--a review. Environ. Pollut. 79: 63-75.

Federal Register. (1999) Regional haze regulations; final rule. F. R. (July 1) 64: 35,713-35,774.

Fenn, M. E.; Keifer, J. W. (1999) Throughfall deposition of nitrogen and sulfur in a Jeffrey pine forest in the San Gabriel Mountains, Southern California. Environ. Pollut. 104: 179-187.

Fenn, M. E.; Poth, M. A. (1999) Temporal and spatial trends in streamwater nitrate concentrations in the San Bernardino Mountains, southern California. J. Environ. Qual. 28: 822-836.

Fenn, M. E.; Poth, M. A.; Johnson, D. W. (1996) Evidence for nitrogen saturation in the San Bernardino Mountains in southern California. For. Ecol. Manage. 82: 211-230

Fenn, M. E.; Poth, M. A.; Aber, J. D.; Baron, J. S.; Bormann, B. T.; Johnson, D. W.; Lemly, A. D.; McNulty, S. G.; Ryan, D. F.; Stottlemyer, R. (1998) Nitrogen excess in North American ecosystems: predisposing factors, ecosystem responses, and management strategies. Ecol. Appl. 8: 706-733.

Fink, F. W.; Buttner, F. H.; Boyd, W. K. (1971) Final report on technical-economic evaluation of air-pollution corrosion costs on metals in the U.S. Columbus, OH: Battelle Memorial Institute, Columbus Laboratories; EPA contract no. CPA 70-86; EPA report no. APTD-0654. Available from: NTIS, Springfield, VA; PB-198-453.

Flowers, E. C.; McCormick, R. A.; Kurfis, K. R. (1969) Atmospheric turbidity over the United States, 1961-1966. J. Appl. Meteorol. 8: 955-962.

Flum, T.; Nodvin, S. C. (1995) Factors affecting streamwater chemistry in the Great Smoky Mountains, USA. Water Air Soil Pollut. 85: 1707-1712.

Fochtman, E. G.; Langer, G. (1957) Automobile paint damaged by airborne iron particles. J. Air Pollut. Control Assoc. 6: 243-247.

Foltescu, V. L.; Isakson, J.; Selin, E.; Stikans, M. (1994) Measured fluxes of sulphur, chlorine and some anthropogenic metals to the Swedish west coast. Atmos. Environ. 28: 2639-2649.

Foster, J. R. (1991) Effects of organic chemicals in the atmosphere on terrestrial plants. In: Moser, T. J.; Barker, J. R.; Tingey, D. T., eds. Ecological exposure and effects of airborne toxic chemicals: an overview. Corvallis, OR: U.S. Environmental Protection Agency, Environmental Research Laboratory; pp. 60-89; EPA report no. EPA-600/3-91-001. Available from: NTIS, Springfield, VA; PB91-148460.

Foster, N. W.; Nicolson, J. A.; Hazlett, P. W. (1989) Temporal variation in nitrate and nutrient cations in drainage waters from a deciduous forest. J. Environ. Qual. 18: 238-244.

Fowler, D.; Cape, J. N.; Unsworth, M. H. (1989) Deposition of atmospheric pollutants on forests. Philos. Trans. R. Soc. London B 324: 247-265.

Fowler, D.; Duyzer, J. H.; Baldocchi, D. D. (1991) Inputs of trace gases, particles and cloud droplets to terrestrial surfaces. Proc. R. Soc. Edinburgh Sect. B: Biol. Sci. 97: 35-59.

Fox, D. G.; Malm, W. C.; Mitchell, B.; Fisher, R. W. (1999) Where there's fire, there's smoke: fine particulate and regional haze. EM November: 15-24.

Frederick, J. E.; Qu, Z.; Booth, C. R. (1998) Ultraviolet radiation at sites on the Antarctic coast. J. Photochem. Photobiol. B 68: 183-190.

Freedman, B.; Hutchinson, T. C. (1981) Sources of metal and elemental contamination of terrestrial environments. In: Lepp, N. W., ed. Effect of heavy metal pollution on plants. Volume 2: metals in the environment. London, United Kingdom: Applied Science Publishers; pp. 35-94.

Freedman, D. M.; Zahm, S. H.; Dosemeci, M. (1997) Residential and occupational exposure to sunlight and mortality from non-Hodgkin's lymphoma: composite (threefold) case-control study. Br. Med. J. BMJ 314: 1451-1455.

Freudenburg, W. R.; Alario, M. (1999) What ecologists can learn from nuclear scientists. Ecosystems 2: 286-291.

Fuller, K. A.; Malm, W. C.; Kreidenweis, S. M. (1999) Effects of mixing on extinction by carbonaceous particles. J. Geophys. Res. 104: 15,941-15,954.

Fürst, P.; Krause, G. H. M.; Hein, D.; Delschen, T.; Wilmers, K. (1993) PCDD/PCDF in cow's milk in relation to their levels in grass and soil. Chemosphere 27: 1349-1357.

Gaggi, C.; Bacci, E.; Calamari, D.; Fanelli, R. (1985) Chlorinated hydrocarbons in plant foliage: an indication of the tropospheric contamination level. Chemosphere 14: 1673-1686.

Gallagher, M. W.; Choularton, T. W.; Morse, A. P.; Fowler, D. (1988) Measurements of the size dependence of cloud droplet deposition at a hill site. Q. J. R. Meteorol. Soc. 114: 1291-1303.

Garcia-Vallès, M.; Vendrell-Saz, M.; Molera, J.; Blazquez, F. (1998) Interaction of rock and atmosphere: patinas on Mediterranean monuments. Environ. Geol. 36(1-2): 137-149.

Garland, J. A. (1978) Dry and wet removal of sulphur from the atmosphere. Atmos. Environ. 12: 349-362.

Garland, C.; Garland, F. (1980) Do sunlight and vitamin D reduce the likelihood of colon cancer? Int. J. Epidemiol. 9: 227-231.

Garland, F. C.; Garland, C. F.; Gorham, E. D.; Young, J. F. (1990) Geographic variation in breast cancer mortality in the United States: a hypothesis involving exposure to solar radiation. Prev. Med. 19: 614-622.

Garner, J. H. B. (1991) Ozone exposure and nitrogen loading in the forests of eastern North America. In: Berglund, R. L.; Lawson, D. R.; McKee, D. J., eds. Tropospheric ozone and the environment: papers from an international conference; March 1990; Los Angeles, CA. Pittsburgh, PA: Air & Waste Management Association; pp. 289-310. (A&WMA transactions series no. TR-19).

Garner, J. H. B. (1994) Nitrogen oxides, plant metabolism, and forest ecosystem response. In: Alscher, R. G.; Wellburn, A. R., eds. Plant responses to the gaseous environment: molecular, metabolic and physiological aspects, [3rd international symposium on air pollutants and plant metabolism]; June 1992; Blacksburg, VA. London, United Kingdom: Chapman & Hall; pp. 301-314.

Garner, J. H. B.; Pagano, T.; Cowling, E. B. (1989) An evaluation of the role of ozone, acid deposition, and other airborne pollutants in the forests of eastern North America. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station; general technical report SE-59.

Garten, C. T., Jr. (1988) Fate and distribution of sulfur-35 in yellow poplar and red maple trees. Oecologia 76: 43-50.

Garten, C. T., Jr.; Hanson, P. J. (1990) Foliar retention of ^{15}N-nitrate and ^{15}N-ammonium by red maple (*Acer rubrum*) and white oak (*Quercus alba*) leaves from simulated rain. Environ. Exp. Bot. 30: 333-342.

Garten, C. T., Jr.; Bondietti, E. A.; Lomax, R. D. (1988) Contribution of foliar leaching and dry deposition to sulfate in net throughfall below deciduous trees. Atmos. Environ. 22: 1425-1432.

Gawel, J. E.; Ahner, B. A.; Friedland, A. J.; Morel, F. M. M. (1996) Role for heavy metals in forest decline indicated by phytochelatin measurements. Nature 381: 64-65.

Gazzi, M.; Georgiadis, T.; Vicentini, V. (2001) Distant contrast measurements through fog and thick haze. Atmos. Environ. 35: 5143-5149.

Geron, C.; Rasmussen, R.; Arnts, R. R.; Guenther, A. (2000) A review and synthesis of monoterpene speciation from forests in the United States. Atmos. Environ. 34: 1761-1781.

Ghedini, N.; Gobbi, G.; Sabbioni, C.; Zappia, G. (2000) Determination of elemental and organic carbon on damaged stone monuments. Atmos. Environ. 34: 4383-4391.

Gildon, A.; Tinker, P. B. (1983) Interactions of vesicular-arbuscular mycorrhizal infection and heavy metals in plants: I. the effects of heavy metals on the development of vesicular-arbuscular mycorrhizas. New Phytol. 95: 247-261.

Gilliam, F. S.; Adams, M. B.; Yurish, B. M. (1996) Ecosystem nutrient responses to chronic nitrogen inputs at Fernow Experimental Forest, West Virginia. Can. J. For. Res. 26: 196-205.

Gobbi, G.; Zappia, G.; Sabbioni, C. (1998) Sulphite quantification on damaged stones and mortars. Atmos. Environ. 32: 783-789.

Goodwin, J. E.; Sage, W.; Tilly, G. P. (1969) Study of erosion by solid particles. Proc. Inst. Mech. Eng. 184: 279-292.

Gorham, E. D.; Garland, C. F.; Garland, F. C. (1989) Acid haze air pollution and breast and colon cancer mortality in 20 Canadian cities. Can. J. Public Health 80: 96-100.

Gorham, E. D.; Garland, F. C.; Garland, C. F. (1990) Sunlight and breast cancer incidence in the USSR. Int. J. Epidemiol. 19: 820-824.

Graedel, T. E.; Nassau, K.; Franey, J. P. (1987) Copper patinas formed in the atmosphere—I. introduction. Corros. Sci. 27: 639-657.

Grand Canyon Visibility Transport Commission. (1996) Report of the Grand Canyon Visibility Transport Commission to the United States Environmental Protection Agency: recommendations for improving western vistas. Denver, CO: Grand Canyon Visibility Transport Commission.

Grant, L. D. (1989) Health effects issues associated with regional and global air pollution problems. In: Proceedings of the changing atmosphere: implications for global security; June, 1988; Toronto, Canada. Geneva, Switzerland: World Meteorological Organization; report no. WMO-No. 710.

Grant, W. B. (2002) An estimate of premature cancer mortality in the U.S. due to inadequate doses of solar ultraviolet-B radiation. Cancer 6: 1867-1875.

Grattan, S. R.; Maas, E. V.; Ogata, G. (1981) Foliar uptake and injury from saline aerosol. J. Environ. Qual. 10: 406-409.

Gravenhorst, G.; Hoefken, K. D.; Georgii, H. W. (1983) Acidic input to a beech and spruce forest. In: Beilke, S.; Elshout, A. J., eds. Acid deposition: proceedings of the CEC workshop organized as part of the concerted action "Physico-chemical behaviour of atmospheric pollutants;" September 1982; Berlin, Federal Republic of Germany. Dordrecht, The Netherlands: D. Reidel Publishing Company; pp. 155-171.

Great Lakes Regional Assessment Group (GLRAG). (2000) Preparing for a changing climate: the potential consequences of climate variability and change. Great Lakes overview. Washington, DC: U.S. Environmental Protection Agency; Office of Research and Development; U.S. Global Change Research Program (USGCRP) and Ann Arbor, MI: University of Michigan; Atmospheric, Oceanic and Space Sciences Department.

Greenberg, B. M.; Wilson, M. I.; Huang, X.-D.; Duxbury, C. L.; Gerhardt, K. E.; Gensemer, R. W. (1997) The effects of ultraviolet-B radiation on higher plants. In: Wang, W.; Gorsuch, J. W.; Hughes, J. S., eds. Plants for Environmental Studies. New York, NY: Lewis Publishers; pp. 1-35.

Grennfelt, P. (1987) Deposition processes for acidifying compounds. Environ. Technol. Lett. 8: 515-527.

Grimm, N. B.; Grove, J. M.; Pickett, S. T. A.; Redman, C. L. (2000) Integrated approaches to long-term studies of urban ecological systems. Bioscience 50: 571-584.

Guderian, R. (1977) Accumulation of pollutants in plant organs. In: Air pollution: phytotoxicity of acidic gases and its significance in air pollution control. Berlin, Germany: Springer-Verlag; pp. 66-74. (Ecological studies: v. 22).

Guderian, R., ed. (1985) Air pollution by photochemical oxidants: formation, transport, control, and effects on plants. New York, NY: Springer-Verlag. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies: analysis and synthesis: v. 52).

Guderian, R. (1986) Terrestrial ecosystems: particulate deposition. In: Legge, A. H.; Krupa, S. V., eds. Air pollutants and their effects on the terrestrial ecosystem. New York, NY: John Wiley & Sons; pp. 339-363. (Advances in environmental science and technology: v. 18).

Gundersen, P.; Callesen, I.; De Vries, W. (1998a) Nitrate leaching in forest ecosystems is related to forest floor C/N ratios. Environ. Pollut. 102(suppl. 1): 403-407.

Gundersen, P.; Emmett, B. A.; Kjonaas, O. J.; Koopmans, C. J.; Tietema, A. (1998b) Impact of nitrogen deposition on nitrogen cycling in forests: a synthesis of NITREX data. For. Ecol. Manage. 101: 37-55.

Hamilton, R. S.; Revitt, D. M.; Vincent, K. J.; Butlin, R. N. (1995) Sulphur and nitrogen particulate pollutant deposition on to building surfaces. In: Saiz-Jimenez, C., ed. The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation; January 1994; Sevilla, Spain. Sci. Total Environ. 167: 57-66.

Han, Q.; Rossow, W. B.; Lacis, A. A. (1994) Near-global survey of effective droplet radii in liquid water clouds using ISCCP data. J. Clim. 7: 465-497.

Hanchette, C. L.; Schwartz, G. G. (1992) Geographic patterns of prostate cancer mortality. Cancer 70: 2861-2869.

Haneef, S. J.; Johnson, J. B.; Thompson, G. E.; Wood, G. C. (1993) Simulation of the degradation of limestones and dolomitic sandstone under dry deposition conditions. In: Corrosion control for low-cost reliability: preceedings [of the] 12th international corrosion congress. Volume 2: process industries plant operation. Houston, TX: National Association of Corrosion Engineers; pp. 700-710.

Hansen, K.; Draaijers, G. P. J.; Ivens, W. P. M. F.; Gundersen, P.; van Leeuwen, N. F. M. (1994) Concentration variations in rain and canopy throughfall collected sequentially during individual rain events. Atmos. Environ. 28: 3195-3205.

Hanson, P. J.; Lindberg, S. E. (1991) Dry deposition of reactive nitrogen compounds: a review of leaf, canopy and non-foliar measurements. Atmos. Environ. Part A 25: 1615-1634.

Hartge, P.; Devessa, S. S.; Graumen, D.; Fears, T. R.; Fraumeni, J. F. (1996) Non-Hodgkin's lymphoma and sunlight. J. Natl. Cancer Inst. 88: 298-300.

Harwell, M. A.; Myers, V.; Young, T.; Bartuska, A.; Gassman, N.; Gentile, J. H.; Harwell, C. C.; Appelbaum, S.; Barko, J.; Causey, B.; Johnson, C.; McLean, A.; Smola, R.; Templet, P.; Tosini, S. (1999) A framework for an ecosystem integrity report card. BioScience 49: 543-556.

Hauhs, M. (1989) Lange Bramke: an ecosystem study of a forested catchment. In: Adriano, D. C.; Havas, M., eds. Acidic precipitation: volume 1, case studies. New York, NY: Springer-Verlag; pp. 275-305. (Adriano, D. C.; Salomons, W., eds. Advances in environmental science).

Haynie, F. H. (1976) Air pollution effects on stress induced intergranular corrosion of 7005-T53 aluminum alloy. In: Stress corrosion–new approaches. Philadelphia, PA: American Society for Testing and Materials; pp. 32-43. (ASTM special technical publication 610).

Haynie, F. H. (1980) Theoretical air pollution and climate effects on materials confirmed by zinc corrosion data. In: Sereda, P. J.; Litvan, G. G., eds. Durability of building materials and components: proceedings of the 1st international conference; August 1978; Ottawa, ON, Canada. Philadelphia, PA: American Society for Testing and Materials; pp. 157-175. (ASTM special technical publication 691).

Haynie, F. H. (1986) Environmental factors affecting corrosion of weathering steel. In: Baboian, R., ed. Materials degradation caused by acid rain: developed from the 20th state-of-the-art symposium of the American Chemical Society; June 1985; Arlington, VA. Washington, DC: American Chemical Society; pp. 163-171. (ACS symposium series 318).

Haynie, F. H.; Lemmons, T. J. (1990) Particulate matter soiling of exterior paints at a rural site. Aerosol Sci. Technol. 13: 356-367.

Haynie, F. H.; Upham, J. B. (1974) Correlation between corrosion behavior of steel and atmospheric pollution data. In: Coburn, S. K., ed. Corrosion in natural environments: presented at the 76th annual meeting American Society for Testing and Materials; June 1973; Philadelphia, PA. Philadelphia, PA: American Society for Testing and Materials; pp. 33-51. (ASTM special technical publication 558).

Haywood, J. M.; Shine, K. P. (1995) The effect of anthropogenic sulfate and soot aerosol on the clear sky planetary radiation budget. Geophys. Res. Lett. 22: 603-606.

Haywood, J. M.; Ramaswamy, V.; Soden, B. J. (1999) Tropospheric aerosol climate forcing in clear-sky satellite observations over the oceans. Science (Washington, DC) 283: 1299-1303.

Heal, G. (2000) Valuing ecosystem services. Ecosystems 3: 24-30.

Heath, J. A.; Huebert, B. J. (1999) Cloudwater deposition as a source of fixed nitrogen in a Hawaiian montane forest. Biogeochemistry 44: 119-134.

Hedin, L. O.; Granat, L.; Likens, G. E.; Buishand, T. A.; Galloway, J. N.; Butler, T. J.; Rodhe, H. (1994) Steep declines in atmospheric base cations in regions of Europe and North America. Nature (London) 367: 351-354.

Heichel, G. H.; Hankin, L. (1972) Particles containing lead, chlorine, and bromine detected on trees with an electron microprobe. Environ. Sci. Technol. 6: 1121-1122.

Heichel, G. H.; Hankin, L. (1976) Roadside coniferous windbreaks as sinks for vehicular lead emissions. J. Air Pollut. Control Assoc. 26: 767-770.

Heinsdorf, D. (1993) The role of nitrogen in declining Scots pine forests (*Pinus sylvestris*) in the lowland of East Germany. Water Air Soil Pollut. 69: 21-35.

Hess, A.; Iyer, H.; Malm, W. (2001) Linear trend analysis: a comparison of methods. Atmos. Environ. 35: 5211-5222.

Hicks, B. B. (1986) Differences in wet and dry particle deposition parameters between North America and Europe. In: Lee, S. D.; Schneider, T.; Grant, L. D.; Verkerk, P. J., eds. Aerosols: research, risk assessment and control strategies; proceedings of the second U.S.-Dutch international symposium; May 1985; Williamsburg, VA. Chelsea, MI: Lewis Publishers; pp. 973-982.

Hicks, B. B.; Matt, D. R.; McMillen, R. T.; Womack, J. D.; Shetter, R. E. (1984) Eddy fluxes of nitrogen oxides to a deciduous forest in complex terrain. In: Samson, P. J., ed. Transactions: the meteorology of acid deposition, an APCA specialty conference; October 1983; Hartford, CT. Pittsburgh, PA: Air Pollution Control Association; pp. 189-201.

Hicks, B. B.; Baldocchi, D. D.; Meyers, T. P.; Hosker, R. P., Jr.; Matt, D. R. (1987) A preliminary multiple resistance routine for deriving dry deposition velocities from measured quantities. Water Air Soil Pollut. 36: 311-330.

Hicks, B. B.; Matt, D. R.; McMillen, R. T.; Womack, J. D.; Wesely, M. L.; Hart, R. L.; Cook, D. R.; Lindberg, S. E.; de Pena, R. G.; Thomson, D. W. (1989) A field investigation of sulfate fluxes to a deciduous forest. J. Geophys. Res. [Atmos.] 94: 13,003-13,011.

Hinckley, T.; Ford, D.; Segura, G.; Sprugel, D. (1992) Key processes from tree to stand level. In: Wall, G., ed. Implications of climate change for Pacific Northwest forest management. Waterloo, ON, Canada: University of Waterloo, Department of Geography; pp. 33-43. (Department of Geography publication series: occasional paper no. 15).

Höfken, K. D.; Gravenhorst, G. (1982) Deposition of atmospheric aerosol particles to beech- and spruce forest. In: Georgii, H.-W.; Pankrath, J., eds. Deposition of atmospheric pollutants: proceedings of a colloquium; November 1981; Oberursel/Taunus, Federal Republic of Germany. Dordrecht, The Netherlands: D. Reidel Publishing Company; pp. 191-194.

Höfken, K. D.; Meixner, F. X.; Ehhalt, D. H. (1983) Deposition of atmospheric trace constituents onto different natural surfaces. In: Pruppacher, H. R.; Semonin, R. G.; Slinn, W. G. N., eds. Precipitation scavenging, dry deposition, and resuspension: v. 2, dry deposition and resuspension: proceedings of the fourth international conference; November-December 1982; Santa Monica, CA. New York, NY: Elsevier; pp. 825-835.

Hofstra, G.; Hall, R. (1971) Injury on roadside trees: leaf injury on pine and white cedar in relation to foliar levels of sodium and chloride. Can. J. Bot. 49: 613-622.
Hogan, G. D.; Rennenberg, H.; Fink, S. (1998) Role and effect of sulfur in tree biology. In: Maynard, D. G., ed. Sulfur in the environment. New York, NY: Marcel Dekker, Inc.; pp. 173-217. (Books in soils, plants, and the environment: v. 67).
Holbrow, G. L. (1962) Atmospheric pollution: its measurement and some effects on paint. J. Oil Colour Chem. Assoc. 45: 701-718.
Holland, D. M.; Principe, P. P.; Sickles, J. E., II. (1999) Trends in atmospheric sulfur and nitrogen species in the eastern United States for 1989-1995. Atmos. Environ. 33: 37-49.
Hollander, M.; Wolfe, D. A. (1973) Nonparametric statistical methods. New York: John Wiley & Sons.
Holling, C. S. (1986) The resilience of terrestrial ecosystems: local surprise and global change. In: Clark, W. C.; Munn, R. E., eds. Sustainable development of the biosphere. Cambridge, UK: Cambridge University; pp. 292-317.
Hooper, D. U.; Vitousek, P. M. (1997) The effects of plant composition and diversity on ecosystem processes. Science 277: 1302-1305.
Hornung, M.; Langan, S. J. (1999) Nitrogen deposition, sources, impacts and responses in natural and semi-natural ecosystems. In: Bangan, S. J., ed. Impact of nitrogen deposition on natural ecosystems and semi-natural ecosystems. Dordrect, Netherlands: Kluwer Academic Publishers; pp. 1-14. [Environmental Pollution, no.3].
Horvath, H. (1993) Atmospheric light absorption–a review. Atmos. Environ. Part A 27: 293-317.
Hosker, R. P., Jr.; Lindberg, S. E. (1982) Review: atmospheric deposition and plant assimilation of gases and particles. Atmos. Environ. 16: 889-910.
Houghton, J. T.; Jenkins, G. J.; Ephraums, J. J., eds. (1990) Climate change: the IPCC scientific assessment. Cambridge, MA: Cambridge University Press; p. 55.
Hoyt, D. V. (1978) A model for the calculation of solar global insolation. Sol. Energy 21: 27-35.
Huebert, B. J.; Luke, W. T.; Delany, A. C.; Brost, R. A. (1988) Measurements of concentrations and dry surface fluxes of atmospheric nitrates in the presence of ammonia. J. Geophys. Res. [Atmos.] 93: 7127-7136.
Hughes, M. K. (1981) Cycling of trace metals in ecosystems. In: Lepp, N. W., ed. Effect of heavy metal pollution on plants. Volume 2: metals in the environment. London, United Kingdom: Applied Science Publishers; pp. 95-118.
Hughes, J. B.; Daily, G. C.; Ehrlich, P. R. (1997) The loss of population diversity and why it matters. In: Raven, P. H., ed. Nature and Human Society: the quest for a sustainable world. Washington, DC: National Academy Press; pp. 71-83.
Humphreys, F. R.; Lambert, M. J.; Kelly, J. (1975) The occurrence of sulphur deficiency in forests. In: McLachlan, K. D., ed. Sulphur in Australasian agriculture. Sydney, Australia: Sydney University Press; pp. 154-162.
Huntington, T.G. (2000) The potential for calcium depletion in forest ecosystems of southeastern United States: review and analysis. Global Biogeochem. Cycles 14: 623-638.
Husar, R. B.; Wilson, W. E. (1993) Haze and sulfur emission trends in the eastern United States. Environ. Sci. Technol. 27: 12-16.
Husar, R. B.; Holloway, J. M.; Patterson, D. E.; Wilson, W. E. (1981) Spatial and temporal pattern of eastern U.S. haziness: a summary. Atmos. Environ. 15: 1919-1928.
Husar, R. B.; Elkins, J. B.; Wilson, W. E. (1994) U.S. visibility trends, 1960-1992, regional and national. Presented at: 87th annual meeting of the Air & Waste Management Association; June; Cincinnati, OH. Pittsburgh, PA: Air & Waste Management Association; paper no. 94-MP3.05.
Hutchinson, A. J.; Johnson, J. B.; Thompson, G. E.; Wood, G. C.; Sage, P. W.; Cooke, M. J. (1992) The role of fly-ash particulate material and oxide catalysts in stone degradation. Atmos. Environ. 26A: 2795-2803.
Huttunen, S. (1994) Effects of air pollutants on epicuticular wax structure. In: Percy, K. E.; Cape, J. N.; Jagels, R.; Simpson, C. J., eds. Air pollutants and the leaf cuticle. New York, NY: Springer-Verlag; pp. 81-96. (NATO ASI series).
Hyer, K. E.; Webb, J. R.; Eshleman, K. N. (1995) Episodic acidification of three streams in Shenandoah National Park, Virginia, USA. Water Air Soil Pollut. 85: 523-528.
Ibrahim, M.; Barrie, L. A.; Fanaki, F. (1983) An experimental and theoretical investigation of the dry deposition of particles to snow, pine trees and artificial collectors. Atmos. Environ. 17: 781-788.

IMPROVE: interagency monitoring of protected visual environments [database]. (1998) [Data on gross chemical composition of PM2.5 particles in rural areas of the eastern and western United States as of 1998]. Fort Collins, CO: National Park Service (NPS); Cooperative Institute for Research in the Atmosphere (CIRA). Available at: http://vista.cira.colostate.edu/improve/ [2001, January 26].

Intergovernmental Panel on Climate Change (IPCC). (1995) Climate change 1994: radiative forcing of climate change and an evaluation of the IPCC IS92 emission scenarios. Cambridge, United Kingdom: University Press.

Intergovernmental Panel on Climate Change (IPCC). (1996) Climate change 1996: contribution of working group I to the second assessment of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom: Cambridge University Press; p. 572.

Intergovernmental Panel on Climate Change (IPCC). (1998) The regional impacts of climate change: an assessment of vulnerability. Cambridge, United Kingdom: Cambridge University Press.

Intergovernmental Panel on Climate Change (IPCC). (2001a) Climate change 2001: the scientific basis. Contribution of working group I to the third assessment report of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom: Cambridge University Press.

Intergovernmental Panel on Climate Change (IPCC). (2001b) Climate change 2001: impacts, adaptation, and vulnerability. Contribution of working group II to the third assessment report of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom: Cambridge University Press.

Irving, P. M., ed. (1991) Acidic deposition: state of science and technology: volumes I-IV. Washington, DC: The U.S. National Acid Precipitation Assessment Program.

Iyer, H.; Patterson, P.; Malm, W. C. (2000) Trends in the extremes of sulfur concentration distributions. J. Air Waste Manage. Assoc. 50: 802-808.

Jacobson, M. Z. (1998) Studying the effects of aerosols on vertical photolysis rate coefficient and temperature profiles over an urban airshed. J. Geophys. Res. [Atmos.] 103: 10,593-10,604.

Jacobson, M. Z. (1999) Isolating nitrated and aromatic aerosols and nitrated aromatic gases as sources of ultraviolet light absorption. J. Geophys. Res. 104 (D3): 3527-3542.

Janzen, J. (1980) Extinction of light by highly nonspherical strongly absorbing colloidal particles: spectrophotometric determination of volume distributions for carbon blacks. Appl. Opt. 19: 2977-2985.

Jaworski, N. A.; Howarth, R. W.; Hetling, L. J. (1997) Atmospheric deposition of nitrogen oxides onto the landscape contributes to coastal eutrophication in the northeast United States. Environ. Sci. Technol. 31: 1995-2004.

Jensen, V. (1974) Decomposition of angiosperm tree leaf litter. In: Dickinson, C. H.; Pugh, G. J. F., eds. Biology of plant litter decomposition: v. I. London, United Kingdom: Academic Press; pp. 69-104.

John, W.; Wall, S. M.; Ondo, J. L. (1985) Dry acid deposition on materials and vegetation: concentrations in ambient air. Berkeley, CA: California Department of Health Services, Air and Industrial Hygiene Laboratory; report no. CA/DOH/AIHL/SP-34. Available from: NTIS, Springfield, VA; PB85-241206.

Johnson, D. W. (1984) Sulfur cycling in forests. Biogeochemistry 1: 29-43.

Johnson, D. W. (1992) Base cation distribution and cycling. In: Johnson, D. W.; Lindberg, S. E., eds. Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study. New York, NY: Springer-Verlag, Inc.; pp. 275-333. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies, analysis and synthesis: v. 91).

Johnson, D. W.; Lindberg, S. E., eds. (1992a) Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study. New York, NY: Springer-Verlag, Inc. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies: analysis and synthesis: v. 91).

Johnson, D. W.; Lindberg, S. E., eds. (1992b) Nitrogen chemistry, deposition, and cycling in forests. In: Johnson, D. W.; Lindberg, S. E., eds. Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study. New York, NY: Springer-Verlag, Inc.; pp. 150-213. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies: analysis and synthesis: v. 91).

Johnson, D. W.; Mitchell, M. J. (1998) Responses of forest ecosystems to changing sulfur inputs. In: Maynard, D. G., ed. Sulfur in the environment. New York, NY: Marcel Dekker, Inc.; pp. 219-262. [Books in Soils, Plants, and the Environment, v. 67].

Johnson, D. W.; Taylor, G. E. (1989) Role of air pollution in forest decline in eastern North America. Water Air Soil Pollut. 48: 21-43.

Johnson, D. W.; Todd, D. E., Jr. (1998) Harvesting effects on long-term changes in nutrient pools of mixed oak forest. Soil Sci. Soc. Amer. J. 62: 1725-1735.

Johnson, D. W.; Henderson, G. S.; Huff, D. D.; Lindberg, S. E.; Richter, D. D.; Shriner, D. S.; Todd, D. E.; Turner, J. (1982a) Cycling of organic and inorganic sulphur in a chestnut oak forest. Oecologia 54: 141-148.

Johnson, D. W.; Cole, D. W.; Bledsoe, C. S.; Cromack, K.; Edmonds, R. L.; Gessel, S. P.; Grier, C. C.; Richards, B. N.; Vogt, K. A. (1982b) Nutrient cycling in forests of the Pacific Northwest. In: Edmonds, R. L., ed. Analysis of coniferous forest ecosystems in the Western United States. Stroudsburg, PA: Hutchinson Ross Publishing Company; pp. 186-232. (US/IBP synthesis series: v. 14).

Johnson, D. W.; Van Miegroet, H.; Lindberg, S. E.; Todd, D. E.; Harrison, R. B. (1991a) Nutrient cycling in red spruce forests of the Great Smoky Mountains. Can. J. For. Res. 21: 769-787.

Johnson, D. W.; Cresser, M. S.; Nilsson, S. I.; Turner, J.; Ulrich, B.; Binkley, D.; Cole, D. W. (1991b) Soil changes in forest ecosystems: evidence for and probable causes. Proc. R. Soc. Edinburgh Sect. B: Biol. Sci. 97B: 81-116.

Johnson, D. W.; Swank, W. T.; Vose, J. M. (1993) Simulated effects of atmospheric sulfur deposition on nutrient cycling in a mixed deciduous forest. Biogeochemistry 23: 169-196.

Johnson, A. H.; Friedland, A. J.; Miller, E. K.; Siccama, T. G. (1994) Acid rain and soils of the Adirondacks. III. Rates of soil acidification in a montane spruce-fir forest at Whiteface Mountain, New York. Can. J. For. Res. 24: 663-669.

Johnson, D.W.; Susfalk, R.B.; Brewer, P.,F.; Swank, W.T. (1999) Simulated effects of reduced sulfur, nitrogen, and base cation deposition on soils and solutions in southern Appalachian forests. J. Environ. Qual 28: 1336-1346.

Joslin, J. D.; Wolfe, M. H. (1992) Red spruce soil chemistry and root distribution across a cloud water deposition gradient. Can. J. For. Res. 22: 893-904.

Joslin, J. D.; Wolfe, M. H. (1994) Foliar deficiencies of mature Southern Appalachian red spruce determined from fertilizer trials. Soil Sci. Soc. Am. J. 58: 1572-1579.

Joslin, J. D.; Kelly, J. M.; Van Miegroet, H. (1992) Soil chemistry and nutrition of North American spruce-fir stands: evidence for recent change. J. Environ. Qual. 21: 12-30.

Justus, C. G.; Murphey, B. B. (1994) Temporal trends in surface irradiance at ultraviolet wavelengths. J. Geophys. Res. 99 (D1): 1389-1394.

Kaminski, M. D.; Landsberger, S. (2000a) Heavy metals in urban soils of East St. Louis, IL, part I: total concentration of heavy metals in soils. J. Air Waste Manage. Assoc. 50: 1667-1679.

Kaminski, M. D.; Landsberger, S. (2000b) Heavy metals in urban soils of East St. Louis, IL, part II: leaching characteristics and modeling. J. Air Waste Manage. Assoc. 50: 1680-1687.

Karmoker, J. L.; Clarkson, D. T.; Saker, L. R.; Rooney, J. M.; Purves, J. V. (1991) Sulphate deprivation depresses the transport of nitrogen to the xylem and the hydraulic conductivity of barley (*Hordeum vulgare L.*) roots. Planta 185: 269-278.

Kaupp, H.; McLachlan, M. S. (1999) Atmospheric particle size distributions of polychlorinated debenzo-*p*-dioxins and dibenzofurans (PCDD/Fs) and polycyclic aromatic hydrocarbons (PAHs) and their implications for wet and dry deposition. Atmos. Environ. 33: 85-95.

Kelly, J.; Lambert, M. J. (1972) The relationship between sulphur and nitrogen in the foliage of *Pinus radiata*. Plant Soil 37: 395-407.

Kelly, J. M.; Meagher, J. F. (1986) Nitrogen input/output relationships for three forested watersheds in eastern Tennessee. In: Correll, D. L., ed. Watershed research perspectives. Washington, DC: Smithsonian Institution Press; pp. 360-391.

Kenk, G.; Fischer, H. (1988) Evidence from nitrogen fertilisation in the forests of Germany. Environ. Pollut. 54: 199-218.

Kiehl, J. T.; Briegleb, B. P. (1993) The relative roles of sulfate aerosols and greenhouse gases in climate forcing. Science (Washington, DC) 260: 311-314.

Klepper, B.; Craig, D. K. (1975) Deposition of airborne particulates onto plant leaves. J. Environ. Qual. 4: 495-499.

Kodama, Y.; Lee, S. D., eds. (1994) The 13th UOEH international symposium and the 2nd pan Pacific cooperative symposium on impact of increased UV-B exposure on human health and ecosystem; October 1993; Kitakyushu, Japan. Kitakyushu, Japan: University of Occupational and Environmental Health.

Koslow, E. E.; Smith, W. H.; Staskawicz, B. J. (1977) Lead-containing particles on urban leaf surfaces. Environ. Sci. Technol. 11: 1019-1021.

Krajíčková, A.; Mejstřík, V. (1984) The effect of fly-ash particles on the plugging of stomata. Environ. Pollut. 36: 83-93.

Kripke, M. L. (1989) Health effects of stratospheric ozone depletion: an overview. In: Schneider, T.; Lee, S. D.; Wolters, G. J. R.; Grant, L. D. eds. Atmospheric ozone research and its policy implications: proceedings of the Third US-Dutch International Symposium; May, 1988; Nijmegen, The Netherlands. Amsterdam, The Netherlands: Elsevier, pp. 795-802. [Studies in Environmental Science 35].

Krupa, S. V.; Legge, A. H. (1986) Single or joint effects of coarse and fine particulate sulfur aerosols and ozone on vegetation. In: Lee, S. D.; Schneider, T.; Grant, L. D.; Verkerk, P. J., eds. Aerosols: research, risk assessment and control strategies, proceedings of the second U.S.-Dutch international symposium; May 1985; Williamsburg, VA. Chelsea, MI: Lewis Publishers, Inc.; pp. 879-887.

Krupa, S. V.; Legge, A. H. (1998) Sulphur dioxide, particulate sulphur and its impacts on a boreal forest ecosystem. In: Ambasht, R. S., ed. Modern trends in ecology and environment. Leiden, The Netherlands: Backhuys Publishers; pp. 285-306.

Kylin, H.; Grimvall, E.; Östman, C. (1994) Environmental monitoring of polychlorinated biphenyls using pine needles as passive samplers. Environ. Sci. Technol. 28: 1320-1324.

Lamaud, E.; Chapuis, A.; Fontan, J.; Serie, E. (1994) Measurements and parameterization of aerosol dry deposition in a semi-arid area. Atmos. Environ. 28: 2461-2471.

Lamersdorf, N. P.; Meyer, M. (1993) Nutrient cycling and acidification of a northwest German forest site with high atmospheric nitrogen deposition. For. Ecol. Manage. 62: 323-354.

Laurenroth, W. K.; Milchunas, D. G. (1985) SO_2 effects on plant community function. In: Winner, W. E.; Mooney, H. A.; Goldstein, R. A., eds. Sulfur dioxide and vegetation: physiology, ecology, and policy issues. Stanford, CA: Stanford University Press; pp. 454-477.

Lawrence, G. B.; Huntington, T. G. (1999) Soil-Calcium depletion linked to acid rain and forest growth in the eastern United States. Reston, VA: U.S. Geological Survey. Available: http://btdqs.usgs.gov/acidrain [1999, November 23].

Lefkowitz, E. S.; Garland, C. F. (1994) Sunlight, vitamin D, and ovarian cancer mortality rates in U.S. women. Int. J. Epidemiol. 23: 1133-1136.

Legge, A. H. (1980) Primary productivity, sulfur dioxide, and the forest ecosystem: an overview of a case study. In: Miller, P. R., ed. Proceedings of symposium on effects of air pollutants on Mediterranean and temperate forest ecosystems; June; Riverside, CA. Berkeley, CA: U.S. Department of Agriculture, Pacific Southwest Forest and Range Experiment Station; pp. 51-62; Forest Service general technical report no. PSW-43. Available from: NTIS, Springfield, VA; PB81-133720.

Lerman, S. L.; Darley, E. F. (1975) Particulates. In: Mudd, J. B.; Kozlowski, T. T., eds. Responses of plants to air pollution. New York, NY: Academic Press; pp. 141-158.

Leslie, M. (2001) Having a bad air day. Science 294: 2255.

Levin, S. A. (1998) Ecosystems and the biosphere as complex adaptive systems. Ecosystems 1: 431-436.

Li-Jones, X.; Prospero, J. M. (1998) Variations in the size distribution of non-sea-salt sulfate aerosol in the marine boundary layer at Barbados: impact of African dust. J. Geophys. Res. 103: 16,073-16,084.

Likens, G. E.; Driscoll, C. T.; Buso, D. C. (1996) Long-term effects of acid rain: response and recovery of a forest ecosystem. Science (Washington, DC) 272: 244-246.

Likens, G. E.; Driscoll, C. T.; Buso, D. C.; Siccama, T. G.; Johnson, C. E.; Lovett, G. M.; Fahey, T. J.; Reiners, W. A.; Ryan, D. F.; Martin, C. W.; Bailey, S. W. (1998) The biogeochemistry of calcium at Hubbard Brook. Biogeochemistry 41: 89-173.

Likens, G. E.; Butler, T. J.; Buso, D. C. (2001) Long- and short-term changes in sulfate deposition: effects of the 1990 Clean Air Act Amendments. Biogeochemistry 52: 1-11.

Lin, J.-M.; Fang, G.-C.; Holsen, T. M.; Noll, K. E. (1993) A comparison of dry deposition modeled from size distribution data and measured with a smooth surface for total particle mass, lead and calcium in Chicago. Atmos. Environ. Part A 27: 1131-1138.

Lindberg, S. E. (1992) Atmospheric deposition and canopy interactions of sulfur. In: Johnson, D. W.; Lindberg, S. E., eds. Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study. New York, NY: Springer-Verlag, Inc.; pp. 74-90. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies: analysis and synthesis: v. 91).

Lindberg, S. E.; Garten, C. T., Jr. (1988) Sources of sulphur in forest canopy throughfall. Nature (London) 336: 148-151.

Lindberg, S. E.; Harriss, R. C. (1981) The role of atmospheric deposition in an eastern U.S. deciduous forest. Water Air Soil Pollut. 16: 13-31.

Lindberg, S. E.; Lovett, G. M. (1985) Field measurements of particle dry deposition rates to foliage and inert surfaces in a forest canopy. Environ. Sci. Technol. 19: 238-244.
Lindberg, S. E.; Lovett, G. M. (1992) Deposition and forest canopy interactions of airborne sulfur: results from the integrated forest study. Atmos. Environ. Part A 26: 1477-1492.
Lindberg, S. E.; McLaughlin, S. B. (1986) Air pollutant interactions with vegetation: research needs in data acquisition and interpretation. In: Legge, A. H.; Krupa, S. V., eds. Air pollutants and their effects on the terrestrial ecosystem. New York, NY: John Wiley & Sons, Inc.; pp. 449-503. (Nriagu, J. O., ed. Advances in environmental science and technology: v. 18).
Lindberg, S. E.; Owens, J. G. (1993) Throughfall studies of deposition to forest edges and gaps in montane ecosystems. Biogeochemistry 19: 173-194.
Lindberg, S. E.; Harriss, R. C.; Turner, R. R. (1982) Atmospheric deposition of metals to forest vegetation. Science (Washington, DC) 215: 1609-1611.
Lindberg, S. E.; Lovett, G. M.; Richter, D. D.; Johnson, D. W. (1986) Atmospheric deposition and canopy interactions of major ions in a forest. Science (Washington, DC) 231: 141-145.
Lindberg, S. E.; Lovett, G. M.; Meiwes, K.-J. (1987) Deposition and forest canopy interactions of airborne nitrate. In: Hutchinson, T. C.; Meema, K. M., eds. Effects of atmospheric pollutants on forests, wetlands and agricultural ecosystems. Berlin, Federal Republic of Germany: Springer-Verlag; pp. 117-130. (NATO as 1 series: v. G16).
Lindberg, S. E.; Lovett, G. M.; Schaefer, D. A.; Bredemeier, M. (1988) Dry deposition velocities and surface-to-canopy scaling factors for aerosol calcium from forest canopy throughfall. J. Aerosol Sci. 19: 1187-1190.
Lindberg, S. E.; Bredemeier, M.; Schaefer, D. A.; Qi, L. (1990) Atmospheric concentrations and deposition of nitrogen and major ions in conifer forests in the United States and Federal Republic of Germany. Atmos. Environ. Part A 24: 2207-2220.
Liousse, C.; Cachier, H.; Jennings, S. G. (1993) Optical and thermal measurements of black carbon aerosol content in different environments: variation of the specific attenuation cross-section, sigma (σ). Atmos. Environ. Part A 27: 1203-1211.
Little, P. (1977) Deposition of 2.75, 5.0 and 8.5 μm particles to plant and soil surfaces. Environ. Pollut. 12: 293-305.
Little, P.; Wiffen, R. D. (1977) Emission and deposition of petrol engine exhaust Pb--I. deposition of exhaust Pb to plant and soil surfaces. Atmos. Environ. 11: 437-447.
Liu, S. C.; McKeen, S. A.; Madronich, S. (1991) Effect of anthropogenic aerosols on biologically active ultraviolet radiation. Geophys. Res. Lett. 18: 2265-2268.
Lokke, H.; Bak, J.; Falkengren-Grerup, U.; Finlay, R. D.; Ilvesniemi, H.; Nygaard, P. H.; Starr, M. (1996) Critical loads of acidic deposition for forest soils: is the current approach adequate. Ambio 25: 510-516.
Lorente, J.; Redaño, A.; de Cabo, X. (1994) Influence of urban aerosol on spectral solar irradiance. J. Appl. Meteorol. 33: 406-415.
Lorusso, S.; Marabelli, M.; Troili, M. (1997) Air pollution and the deterioration of historic monuments. J. Environ. Pathol. Toxicol. Oncol. 16: 171-173.
Lovett, G. M. (1984) Atmospheric deposition to forests. In: Breece, L.; Hasbrouck, S., eds. Forest responses to acidic deposition: proceedings of the U. S.-Canadian conference; August 1983; Orono, ME. Orono, ME: University of Maine, Orono, Land and Water Resources Center; pp. 7-18.
Lovett, G. M. (1992) Atmospheric deposition and canopy interactions of nitrogen. In: Johnson, D. W.; Lindberg, S. E., eds. Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study. New York, NY: Springer-Verlag, Inc.; pp. 152-166. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies: analysis and synthesis: v. 91).
Lovett, G. M. (1994) Atmospheric deposition of nutrients and pollutants in North America: an ecological perspective. Ecol. Appl. 4: 629-650.
Lovett, G. M.; Lindberg, S. E. (1984) Dry deposition and canopy exchange in a mixed oak forest as determined by analysis of throughfall. J. Appl. Ecol. 21: 1013-1027.
Lovett, G. M.; Lindberg, S. E. (1986) Dry deposition of nitrate to a deciduous forest. Biogeochemistry 2: 137-148.
Lovett, G. M.; Lindberg, S. E. (1992) Concentration and deposition of particles and vapors in a vertical profile through a forest canopy. Atmos. Environ. Part A 26: 1469-1476.
Lovett, G. M.; Lindberg, S. E. (1993) Atmospheric deposition and canopy interactions of nitrogen in forests. Can. J. For. Res. 23: 1603-1616.

Lovett, G. M.; Traynor, M. M.; Pouyat, R. V.; Carreiro, M. M.; Zhu, W.-X.; Baxter, J. W. (2000) Atmospheric deposition to oak forest along an urban-rural gradient. Environ. Sci. Technol. 34: 4294-4300.

Lowenthal, D. H.; Rogers, C. F.; Saxena, P.; Watson, J. G.; Chow, J. C. (1995) Sensitivity of estimated light extinction coefficients to model assumptions and measurement errors. Atmos. Environ. 29: 751-766.

Madronich, S. (1987) Photodissociation in the Atmosphere. 1. Actinic flux and the effects of ground reflections and clouds. J. Geophys. Res. [Atmos.] 92: 9740-9752.

Malm, W. C. (1989) Atmospheric haze: its sources and effects on visibility in rural areas of the continental United States. Environ. Monit. Assess. 12: 203-225.

Malm, W. C. (1999) Introduction to visibility. Fort Collins, CO: Colorado State University; Cooperative Institute for Research in the Atmosphere (CIRA); NPS Visibility Program. Available: http://www.epa.gov/air/visibility/introvis.pdf [2 April 2002].

Malm, W. C. (2000) Spatial and seasonal patterns and temporal variability of haze and its constituents in the United States. Report III. Fort Collins, CO: Cooperative Institute for Research in the Atmosphere, Colorado State University. Available: http://vista.cira.colostate.edu/improve/Publications/Reports/2000/2000.htm [22 March, 2002].

Malm, W. C.; Kreidenweis, S. M. (1997) The effects of models of aerosol hygroscopicity on the apportionment of extinction. Atmos. Environ. 31: 1965-1976.

Malm, W. C.; Sisler, J. F.; Huffman, D.; Eldred, R. A.; Cahill, T. A. (1994) Spatial and seasonal trends in particle concentration and optical extinction in the United States. J. Geophys. Res. [Atmos.] 99: 1347-1370.

Malm, W. C.; Molenar, J. V.; Eldred, R. A.; Sisler, J. F. (1996) Examining the relationship among atmospheric aerosols and light scattering and extinction in the Grand Canyon area. J. Geophys. Res. [Atmos.] 101: 19,251-19,265.

Malm, W. C.; Day, D.; Kreidenweis, S. M. (1997) Comparison of measured and reconstructed scattering during an intensive field study at Great Smoky Mountains National Park. Presented at: 90th annual meeting and exhibition of the Air & Waste Management Association; June; Toronto, ON, CA. Pittsburgh, PA: Air & Waste Management Association; 97-WA70.02.

Malm, W. C.; Day, D. E.; Kreidenweis, S. M. (2000) Light scattering characteristics of aerosols as a function of relative humidity: part I--a comparison of measured scattering and aerosol concentrations using the theoretical models. J. Air Waste Manage. Assoc. 50: 686-700.

Malm, W. C.; Day, D. E.; Kreidenweis, S. M. (2000) Light scattering characteristics of aerosols as a function of relative humidity: Part II–a comparison of measured scattering and aerosol concentrations using the statistical models. J. Air Waste Manage. Assoc. 50: 701-709.

Mansfeld, F. (1980) Regional air pollution study: effects of airborne sulfur pollutants on materials. Research Triangle Park, NC: U.S. Environmental Protection Agency, Environmental Sciences Research Laboratory; EPA report no. EPA-600/4-80-007. Available from: NTIS, Springfield, VA; PB81-126351.

Marchwińska, E.; Kucharski, R. (1987) The combined influence of SO_2 and heavy metal-containing particulates on beans, carrots and parsley. Environ. Monit. Assess. 8: 11-25.

Markewitz, D.; Richter, D.D.; Allen, L. H.; Urrego, J. B. (1998) Three decades of observed soil acidification in the Calhoun Experimental Forest: Has acid rain made a difference? Soil Sci. Soc. Amer. J. 62: 1428-1439.

Marschner, H. (1995) Sulfur. In: Mineral nutrition of higher plants. 2nd ed. London, United Kingdom: Academic Press; pp. 255-265.

Martin, M. H.; Coughtrey, P. J. (1981) Impact of metals on ecosystem function and productivity. In: Lepp, N. W., ed. Effect of heavy metal pollution on plants: volume 2, metals in the environment. Barking, United Kingdom: Applied Science Publishers; pp. 119-158. (Mellanby, K., ed. Pollution monitoring series).

Martin, C. E.; Gravatt, D. A.; Loeschen, V. S. (1992) Photosynthetic responses of three species to acute exposures of nitrate- and sulphate-containing aerosols. Atmos. Environ. Part A 26: 381-391.

Mathai, C. V. (1995) The Grand Canyon visibility. EM (December): 20-31.

Matson, P. A.; Parton, W. J.; Power, A. G.; Swift, M. J. (1997) Agricultural intensification and ecosystem properties. Science (Washington, DC) 277: 504-509.

May, P. F.; Till, A. R.; Cumming, M. J. (1972) Systems analysis of ^{35}sulphur kinetics in pastures grazed by sheep. J. Appl. Ecol. 9: 25-49.

McCartney, H. A.; Aylor, D. E. (1987) Relative contributions of sedimentation and impaction to deposition of particles in a crop canopy. Agric. For. Meteorol. 40: 343-358.

McCune, D. C.; Silberman, D. H.; Mandl, R. H.; Weinstein, L. H.; Freudenthal, P. C.; Giardina, P. A. (1977) Studies on the effects of saline aerosols of cooling tower origin on plants. J. Air Pollut. Control Assoc. 27: 319-324.

McDonald, R. L.; Unni, C. K.; Duce, R. A. (1982) Estimation of atmospheric sea salt dry deposition: wind speed and particle size dependence. J. Geophys. Res. C: Oceans Atmos. 87: 1246-1250.

McDonnell, M. J.; Pickett, S. T. A.; Groffman, P.; Bohlen, P.; Pouyat, R. V.; Zipperer, W. C.; Parmelee, R. W.; Carreiro, M. M.; Medley, K. (1997) Ecosystem processes along an urban-to-rural gradient. Urban Ecosyst. 1: 21-36.

McDow, S. R.; Vartiainen, M.; Sun, Q.; Hong, Y.; Yao, Y.; Kamens, R. M. (1995) Combustion aerosol water content and its effect on polycyclic aromatic hydrocarbon reactivity. Atmos. Environ. 29: 791-797.

McGee, E. S.; Mossotti, V. G. (1992) Gypsum accumulation on carbonate stone. Atmos. Environ. Part B 26: 249-253.

McGowan, T. F.; Lipinski, G. E.; Santoleri, J. J. (1993) New rules affect the handling of waste fuels. Chem. Eng. (March): 122-128.

McIntyre, N. E.; Knowles-Yánez, K.; Hope, D. (2000) Urban ecology as an indisciplinary field: differences in the use of "urban" between the social and natural sciences. Urban Ecosyst. 4: 5-24.

McKay, W. A.; Garland, J. A.; Livesley, D.; Halliwell, C. M.; Walker, M. I. (1994) The characteristics of the shore-line sea spray aerosol and the landward transfer of radionuclides discharged to coastal sea water. Atmos. Environ. 28: 3299-3309.

McLachlan, M. S. (1996a) Bioaccumulation of hydrophobic chemicals in agricultural food chains. Environ. Sci. Tech. 30: 252-259.

McLachlan, M. S. (1996b) Biological uptake and transfer of polychlorinated dibenzo-p-dioxins and dibenzofurans. In: Hester, R. E.; Harrison, R. M., eds. Chlorinated organic micropollutants. Cambridge, United Kingdom: Royal Society of Chemistry; pp. 31-52. (Issues in environmental science and technology).

McLachlan, M. S. (1999) Framework for the interpretation of measurements of SOCs in plants. Environ. Sci. Technol. 33: 1799-1804.

McMahon, T. A.; Denison, P. J. (1979) Empirical atmospheric deposition parameters--a survey. Atmos. Environ. 13: 571-585.

McMillen, R. T. (1988) An eddy correlation technique with extended applicability to non-simple terrain. Boundary Layer Meteorol. 43: 231-245.

McMurry, P. H.; Stolzenburg, M. R. (1989) On the sensitivity of particle size to relative humidity for Los Angeles aerosols. Atmos. Environ. 23: 497-507.

McMurry, P. H.; Zhang, X.; Lee, C.-T. (1996) Issues in aerosol measurement for optics assessments. J. Geophys. Res. [Atmos.] 101: 19,189-19,197.

Meiwes, K. J.; Khanna, P. K. (1981) Distribution and cycling of sulphur in the vegetation of two forest ecosystems in an acid rain environment. Plant Soil 60: 369-375.

Mid-Atlantic Regional Assessment Team (MARAT). (2000) Preparing for a changing climate: the potential consequences of climate variability and change. Mid-Atlantic overview. Washington, DC: U.S. Environmental Protection Agency; Office of Research and Development; U.S. Global Change Research Program (USGCRP) and University Park, PA: Pennsylvania State University.

Middleton, W. E. K. (1952) Vision through the atmosphere. Toronto, ON, Canada: University of Toronto Press.

Milford, J. B.; Davidson, C. I. (1987) The sizes of particulate sulfate and nitrate in the atmosphere--a review. JAPCA 37: 125-134.

Millar, C. S. (1974) Decomposition of coniferous leaf litter. In: Dickinson, C. H.; Pugh, G. J. F., eds. Biology of plant litter decomposition: volume I. New York, NY: Academic Press; pp. 105-128.

Miller, P. R.; McBride, J. R., eds. (1999) Oxidant air pollution impacts in the Montane forests of southern California: a case study of the San Bernardino Mountains. New York, NY: Springer-Verlag. (Ecological Studies: v. 134.)

Miller, W. C.; Fornes, R. E.; Gilbert, R. D.; Speer, A.; Spence, J. (1992) Removal of $CaCO_3$ extender in residential coatings by atmospheric acidic deposition. In: Measurement of toxic and related air pollutants: proceedings of the 1992 U.S. EPA/A&WMA international symposium. Pittsburgh, PA: Air & Waste Management Association; pp. 129-134. (A&WMA publication VIP-25).

Miller, E. K.; Panek, J. A.; Friedland, A. J.; Kadlecek, J.; Mohnen, V. A. (1993) Atmospheric deposition to a high-elevation forest at Whiteface Mountain, New York, USA. Tellus Ser. B 45: 209-227.

Mims, F. M., III. (1996) Significant reduction of UVB caused by smoke from biomass burning in Brazil. Photochem. Photobiol. 64: 814-816.

Mitchell, M. J.; Harrison, R. B.; Fitzgerald, J. W.; Johnson, D. W.; Lindberg, S. E.; Zhang, Y.; Autry, A. (1992a) Sulfur distribution and cycling in forest ecosystems. In: Johnson, D. W.; Lindberg, S. E., eds. Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study. New York, NY: Springer-Verlag; pp. 90-129. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies: analysis and synthesis: v. 91).

Mitchell, M. J.; Foster, N. W.; Shepard, J. P.; Morrison, I. K. (1992b) Nutrient cycling in Huntington Forest and Turkey Lakes deciduous stands: nitrogen and sulfur. Can. J. For. Res. 22: 457-464.

Molenar, J. V.; Malm, W. C.; Johnson, C. E. (1994) Visual air quality simulation techniques. Atmos. Environ. 28: 1055-1063.

Monteith, J. L.; Unsworth, M. H. (1990) Principles of environmental physics. 2nd ed. London, United Kingdom: Edward Arnold; pp. 165-175.

Moosmüller, H.; Arnott, W. P.; Rogers, C. F. (1998) Photoacoustic and filter measurements related to aerosol light absorption during the Northern Front Range Air Quality Study (Colorado 1996.1997). J. Geophys. Res. 103: 28,149-28,157.

Moropoulou, A.; Bisbikou, K.; Torfs, K.; Van Grieken, R.; Zezza, F.; Macri, F. (1998) Origin and growth of weathering crusts on ancient marbles in industrial atmosphere. Atmos. Environ. 32: 967-982.

Mueller, S. F. (1988) Chemical deposition to high-elevation spruce-fir forests in the eastern United States. Muscle Shoals, AL: Tennessee Valley Authority, Air Quality Branch.

Naeem, S.; Thompson, L. J.; Lawler, S. P.; Lawton, J. H.; Woodfin, R. M. (1994) Declining biodiversity can alter the performance of ecosystems. Nature 368: 734-737.

Nash, T. H., III. (1975) Influence of effluents from a zinc factory on lichens. Ecol. Monogr. 45: 183-198.

National Assessment Synthesis Team (NAST). (2000) Climate change impacts on the United States: the potential consequences of climate variability and change. Overview. Washington, DC: U.S. Global Change Research Program.

National Park Service. (1998) IMPROVE program goals and objectives. Washington, DC: National Park Service. Available: www2.nature.nps.gov/ard/impr/index.htm [24 November 1999].

National Research Council. (1993) Protecting visibility in national parks and wilderness areas. Washington, DC: National Academy Press. 3v.

National Science and Technology Council. (1998) National acid precipitation assessment program biennial report to Congress: an integrated assessment; executive summary. Silver Spring, MD: U.S. Department of Commerce, National Oceanic and Atmospheric Administration. Available: www.nnic.noaa.gov/CENR/NAPAP/NAPAP_96.htm [24 November 1999].

National Weather Service. (1998) Visibility. Washington, DC: National Weather Service, National Oceanic and Atmospheric Administration. Available: www.nws.noaa.gov/asos/vsby.htm [24 November 1999].

Nazaroff, W. W.; Cass, G. R. (1991) Protecting museum collections from soiling due to the deposition of airborne particles. Atmos. Environ. Part A 25: 841-852.

Nelis, P. M.; Branford, D.; Unsworth, M. H. (1994) A model of the transfer of radioactivity from sea to land in sea spray. Atmos. Environ. 28: 3213-3223.

Neuberger, H.; Hosler, C. L.; Kocmond, W. C. (1967) Vegetation as aerosol filter In: Tromp, S. W.; Weihe, W. H., eds. Biometeorology, v. 2. pp. 693-702.

New England Regional Assessment Group (NERAG). (2001) Preparing for a changing climate: the potential consequences of climate variability and change. New England regional overview. Washington, DC: U.S. Environmental Protection Agency; Office of Research and Development; U.S. Global Change Research Program (USGCRP) and Durham, NH: University of New Hampshire; Institute for the Study of Earth, Oceans, and Space.

Nicholson, K. W. (1988) The dry deposition of small particles: a review of experimental measurements. Atmos. Environ. 22: 2653-2666.

Nicholson, K. W.; Davies, T. D. (1987) Field measurements of the dry deposition of particulate sulphate. Atmos. Environ. 21: 1561-1571.

Nielsen, K. E.; Ladekarl, U. L.; Nornberg, P. (1999) Dynamic soil processes on heathland due to changes in vegetation to oak and Sitka spruce. For. Ecol. Manage. 114: 107-116.

Nihlgård, B. (1985) The ammonium hypothesis--an additional explanation to the forest dieback in Europe. Ambio 14: 2-8.

Niklińska, M.; Laskowski, R.; Maryański, M. (1998) Effect of heavy metals and storage time on two types of forest litter: basal respiration rate and exchangeable metals. Ecotoxicol. Environ. Safety 41: 8-18.
Noble, I. R.; Dirzo, R. (1997) Forests as human-dominated ecosystems. Science (Washington, DC) 277: 522-525.
Nodvin, S. C.; Van Miegroet, H.; Lindberg, S. E.; Nicholas, N. S.; Johnson, D. W. (1995) Acidic deposition, ecosystem processes, and nitrogen saturation in a high elevation southern Appalachian watershed. Water Air Soil Pollut. 85: 1647-1652.
Nord, A. G.; Ericsson, T. (1993) Chemical analysis of thin black layers on building stone. Stud. Conserv. 38: 25-35.
Northeast States for Coordinated Air Use Management (NESCAUM). (2002) CAMNET: realtime air pollution and visibility monitoring. Boston, MA: NESCAUM. Available at: http://www.hazecam.net. (23 January 2002).
Norton, R. A.; Childers, N. F. (1954) Experiments with urea sprays on the peach. Proc. Am. Soc. Hortic. Sci. 63: 23-31.
Novakov, T.; Penner, J. E. (1993) Large contribution of organic aerosols to cloud-condensation-nuclei concentrations. Nature (London) 365: 823-826.
Ockenden, W. A.; Stinnes, E.; Parker, C.; Jones, K. C. (1998) Observations on persistent organic pollutants in plants: implications for their use as passive air samplers and for POP cycling. Environ. Sci. Technol. 32: 2721-2726.
Odum, E. P. (1969) The strategy of ecosystem development: an understanding of ecological succession provides a basis for resolving man's conflict with nature. Science (Washington, DC) 164: 262-270.
Odum, E. P. (1985) Trends expected in stressed ecosystems. BioScience 35: 419-422.
Odum, E. P. (1993) The ecosystem. In: Ecology and our endangered life-support systems. 2nd ed. Sunderland, MA: Sinauer Associates, Inc.; pp. 38-67.
Oehme, M.; Biseth, A.; Schlabach, M.; Wiig, Ø. (1995) Concentrations of polychlorinated dibenzo-*p*-dioxins, dibenzofurans and non-*ortho* substituted biphenyls in polar bear milk from Svalbard (Norway). Environ. Pollut. 90: 401-407.
Olszyk, D. M.; Bytnerowicz, A.; Takemoto, B. K. (1989) Photochemical oxidant pollution and vegetation: effects of gases, fog, and particles. Environ. Pollut. 61: 11-29.
Oosting, H. J.; Billings, W. D. (1942) Factors effecting vegetational zonation on coastal dunes. Ecology 23: 131-142.
Ormrod, D. P. (1984) Impact of trace element pollution on plants. In: Treshow, M., ed. Air pollution and plant life. Chichester, United Kingdom: John Wiley & Sons; pp. 291-319.
Ouimette, J. R.; Flagan, R. C. (1982) The extinction coefficient of multicomponent aerosols. Atmos. Environ. 16: 2405-2420.
Padgett, P.; Allen, E. B. (1999) Differential responses to nitrogen fertilization in native shrubs and exotic annuals common to Mediterranean coastal sage scrub of California. Plant Ecol. 144: 93-101.
Padgett, P. E.; Allen, E. B.; Bytnerowicz, A.; Minich, R. A. (1999) Changes in soil inorganic nitrogen as related to atmospheric nitrogenous pollutants in southern California. Atmos. Environ. 33: 769-781.
Paerl, H. W.; Bales, J. D.; Ausley, L. W.; Buzzelli, C. P.; Crowder, L. B.; Eby, L. A.; Go, M.; Peierls, B. L.; Richardson, T. L.; Ramus, J. S. (2001) Ecosystem impacts of three sequential hurricanes (Dennis, Floyd, and Irene) on the United States' largest lagoonal estuary, Pamlico Sound, NC. Proc. Nat. Acad. Sci. U. S. A. 98: 5655-5611.
Pan, W.; Tatang, M. A.; McRae, G. J.; Prinn, R. G. (1997) Uncertainty analysis of direct radiative forcing by anthropogenic sulfate aerosols. J. Geophys. Res. 102 (D18): 21,915-21,924.
Pan, W.; Tatang, M. A.; McRae, G. J.; Prinn, R. G. (1998) Uncertainty analysis of indirect radiative forcing by anthropogenic sulfate aerosols. J. Geophys. Res. 103 (D4): 3815-3823.
Pandis, S. N.; Seinfeld, J. H. (1989) Mathematical modeling of acid deposition due to radiation fog. J. Geophys. Res. [Atmos.] 94: 12,911-12,923.
Pankow, J. F. (1987) Review and comparative analysis of the theories on partitioning between the gas and aerosol particulate phases in the atmosphere. Atmos. Environ. 21: 2275-2283.
Parish, S. B. (1910) The effect of cement dust on citrus trees. Plant World 13: 288-291.
Parker, A. (1955) The destructive effects of air pollution on materials. Colliery Guardian 191: 447-450.
Parker, G. G. (1983) Throughfall and stemflow in the forest nutrient cycle. In: Macfadyen, A.; Ford, E. D., eds. Advances in ecological research, v. 13; pp. 57-120.
Patz, J. A.; Engelberg, D.; Last, J. (2000a) The effects of changing weather on public health. Ann. Rev. Public Health 21: 271-307.

Patz, J. A.; McGeehin, M. A.; Bernard, S. M.; Ebi, K. L.; Epstein, P. R.; Grambsch, A.; Gubler, D. J.; Reiter, P.; Romieu, I.; Rose, J. B.; Samet, J. M.; Trtanj, J. (2000b) The potential health impacts of climate variability and change for the United States: executive summary of the report of the health sector of the U.S. national assessment. Environ. Health Perspect. 108: 367-376.

Pearcy, R. W.; Björkman, O.; Caldwell, M. M.; Keeley, J. E.; Monson, R. K.; Strain, B. R. (1987) Carbon gain by plants in natural environments. BioScience 37: 21-29.

Penndorf, R. B. (1958) An approximation method to the Mie theory for colloidal spheres. J. Phys. Chem. 62: 1537-1542.

Percy, K. E. (1991) Effects of acid rain on forest vegetation: morphological and non-mensurational growth effects. In: Rennie, P. J.; Robitaille, G., eds. Effects of Acid Rain on Forest Resources: Proceedings of the Conference ; June, 1983; Sainte-Foy, Quebec. Ottawa, Canada: Canadian Forestry Service; pp. 97-111.

Perry, K. D.; Cahill, T. A.; Eldred, R. A.; Dutcher, D. D.; Gill, T. E. (1997) Long-range transport of North African dust to the eastern United States. J. Geophys. Res. [Atmos] 102: 11,225-11,238.

Peterjohn, W. T.; Adams, M. B.; Gilliam, F. S. (1996) Symptoms of nitrogen saturation in two central Appalachian hardwood forest ecosystems. Biogeochemistry 35: 507-522.

Peters, K.; Eiden, R. (1992) Modelling the dry deposition velocity of aerosol particles to a spruce forest. Atmos. Environ. Part A 26: 2555-2564.

Peterson, J. T.; Flowers, E. C.; Rudisill, J. H. (1978) Urban-rural solar radiation and atmospheric turbidity measurements in the Los Angeles basin. J. Appl. Meteorol. 17: 1595-1609.

Peterson, G.; Allen, C. R.; Holling, C. S. (1998) Ecological resilience, biodiversity, and scale. Ecosystems 1: 6-18.

Petuskey, W. T.; Richardson, D. A.; Dolske, D. A. (1995) Aspects of the deterioration of sandstone masonry in Anasazi dwelling ruins at Mesa Verde National Park, Colorado, USA. Sci. Total Environ. 167: 145-159.

Piatt, J. R.; Krause, P. D. (1974) Road and site characteristics that influence road salt distribution and damage to roadside aspen trees. Water Air Soil Pollut. 3: 301-304.

Pickett, S. T. A.; Cadenasso, M. L.; Grove, J. M.; Nilon, C. Y.; Pouyat, R. V.; Zipperer, W. C.; Costanza, R. (2001) Urban ecological systems: linking terrestrial ecological, physical, and socioeconomic components of metropolitan areas. Annu. Rev. Ecol. Syst. 32: 127-157.

Pilinis, C.; Pandis, S. N.; Seinfield, J. H. (1995) Sensitivity of direct climate forcing by atmospheric aerosols to aerosol size and composition. J. Geophys. Res. [Atmos.] 100: 18739-18754.

Pimentel, D.; Wilson, C.; McCullum, C.; Huang, R.; Dwen, P.; Flack, J.; Tran, Q.; Saltman, T.; Cliff, B. (1997) Economic and environmental benefits of biodiversity. BioScience 47: 747-757.

Pimm, S. L. (1984) The complexity and stability of ecosystems. Nature (London) 307: 321-326.

Pitchford, M. L.; Malm, W. C. (1994) Development and applications of a standard visual index. Atmos. Environ. 28: 1049-1054.

Pitchford, M. L.; McMurry, P. H. (1994) Relationship between measured water vapor growth and chemistry of atmospheric aerosol for Grand Canyon, Arizona, in winter 1990. Atmos. Environ. 28: 827-839.

Pouyat, R. V.; McDonnell, M. J. (1991) Heavy metal accumulations in forest soils along an urban-rural gradient in southeastern New York, USA. Water Air Soil Pollut. 57/58: 797-808.

Prescott, C. E.; Parkinson, D. (1985) Effects of sulphur pollution on rates of litter decomposition in a pine forest. Can. J. Bot. 63: 1436-1443.

Pueschel, R. F. (1993) Potential climatic effects of anthropogenic aerosols. In: Jennings, S. G., ed. Aerosol effects on climate. Tucson, AZ: University of Arizona Press; pp. 110-132.

Ragsdale, H. L.; Lindberg, S. E.; Lovett, G. M.; Schaefer, D. A. (1992) Atmospheric deposition and throughfall fluxes of base cations. In: Johnson, D. W.; Lindberg, S. E., eds. Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study. New York, NY: Springer Verlag; pp. 235-253. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies: analysis and synthesis: v. 91).

Rapport, D. J.; Whitford, W. G. (1999) How ecosystems respond to stress: common properties of arid and aquatic systems. BioScience 49: 193-203.

Raynal, D. J.; Joslin, J. D.; Thornton, F. C.; Schaedle, M.; Henderson, G. S. (1990) Sensitivity of tree seedlings to aluminum: III. Red spruce and loblolly pine. J. Environ. Qual. 19: 180-187.

Realini, M.; Negrotti, R.; Appollonia, L.; Vaudan, D. (1995) Deposition of particulate matter on stone surfaces: an experimental verification of its effects on Carrara marble. In: Saiz-Jimenez, C., ed. The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation; January 1994; Sevilla, Spain. Sci. Total Environ. 167: 67-72.

Reiners, W. A.; Olson, R. K. (1984) Effects of canopy components on throughfall chemistry: an experimental analysis. Oecologia 63: 320-330.
Reisinger, L. M. (1990) Analysis of airborne particles sampled in the southern Appalachian Mountains. Water Air Soil Pollut. 50: 149-162.
Rennie, P. J. (1955) The uptake of nutrients by mature forest growth. Plant Soil 7: 49-95.
Reuss, J. O. (1983) Implications of the calcium-aluminum exchange system for the effect of acid precipitation on soils. J. Environ. Qual. 12: 591-595.
Reuss, J. O.; Johnson, D. W. (1986) Acid deposition and the acidification of soils and waters. New York, NY: Springer-Verlag. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies: analysis and synthesis: v. 59).
Riggan, P. J.; Lockwood, R. N.; Lopez, E. N. (1985) Deposition and processing of airborne nitrogen pollutants in Mediterranean-type ecosystems of southern California. Environ. Sci. Technol. 19: 781-789.
Rodney, D. R. (1952) The entrance of nitrogen compounds through the epidermis of apple leaves. Proc. Am. Soc. Hortic. Sci. 59: 99-102.
Roelofs, J. G. M.; Kempers, A. J.; Houdijk, A. L. F. M.; Jansen, J. (1985) The effect of air-borne ammonium sulphate on *Pinus nigra* var. *maritima* in the Netherlands. Plant Soil 84: 45-56.
Roelofs, J. G. M.; Boxman, A. W.; Van Dijk, H. F. G. (1987) Effects of airborne ammonium on natural vegetation and forests. In: Asman, W. A. H.; Diederen, H. S. M. A., eds. Ammonia and acidification: proceedings [of a] symposium of the European Association for the Science of Air Pollution (EURASAP); April; Bilthoven, The Netherlands. European Association for the Science of Air Pollution; pp. 266-276.
Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993a) Sources of fine organic aerosol. 4. Particulate abrasion products from leaf surfaces of urban plants. Environ. Sci. Technol. 27: 2700-2711.
Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993b) Sources of fine organic aerosol. 3. Road dust, tire debris, and organometallic brake lining dust: roads as sources and sinks. Environ. Sci. Technol. 27: 1892-1904.
Rosenberg, N. J.; Blad, B. L.; Verma, S. B. (1983) Microclimate: the biological environment. 2nd ed. New York, NY: John Wiley & Sons.
Rovira, A. D.; Davey, C. B. (1974) Biology of the rhizosphere. In: Carson, E. W., ed. The plant root and its environment: proceedings of an institute; July, 1971; Blacksburg, VA. Charlottesville, VA: University Press of Virginia; pp. 153-204.
Rueth, H.; Baron, J. S. (2002) Differences in Englemann spruce forest biogeochemistry east and west of the Continental Divide in Colorado, USA. Ecosystems 5: 45-57.
Sabbioni, C. (1995) Contribution of atmospheric deposition to the formation of damage layers. In: Jimenez-Saiz, C., ed. The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation; January 1994; Sevilla, Spain. Sci. Total Environ. 167: 49-56.
Sabbioni, C.; Zappia, G.; Gobbi, G. (1992) Carbonaceous particles on carbonate building stones in a simulated system. In: Proceedings of the 1992 European aerosol conference; September; Oxford, United Kingdom. J. Aerosol Sci. 23(suppl. 1): S921-S924.
Sabbioni, C.; Zappia, G.; Gobbi, G. (1996) Carbonaceous particles and stone damage in a laboratory exposure system. J. Geophys. Res. [Atmos.] 101: 19,621-19,627.
Sabbioni, C.; Zappia, G.; Ghedini, N.; Gobbi, G.; Favoni, O. (1998) Black crusts on ancient mortars. Atmos. Environ. 32: 215-223.
Saiz-Jimenez, C. (1993) Deposition of airborne organic pollutants on historic buildings. Atmos. Environ. Part B 27: 77-85.
Samecka-Cymerman, A.; Kempers, A. J. (1999) Bioindication of heavy metals in the town Wro⁄caw (Poland) with evergreen plants. Atmos. Environ. 33: 419-430.
Sanyal, B.; Singhania, G. K. (1956) Atmospheric corrosion of metals: part I. J. Sci. Ind. Res. Sect. B 15: 448-455.
Saunders, P. J. W.; Godzik, S. (1986) Terrestrial vegetation--air pollutant interactions: nongaseous air pollutants. In: Legge, A. H.; Krupa, S. V., eds. Air pollutants and their effects on the terrestrial ecosystem. New York, NY: John Wiley & Sons; pp. 389-394. (Advances in environmental science and technology: v. 18).
Saxena, P.; Peterson, T. W. (1981) Thermodynamics of multicomponent electrolytic aerosols. J. Colloid Interface Sci. 79: 496-510.
Saxena, P.; Mueller, P. K.; Kim, Y. P.; Seinfeld, J. H.; Koutrakis, P. (1993) Coupling thermodynamic theory with measurements to characterize acidity of atmospheric particles. Aerosol Sci. Technol. 19: 279-293.

Saxena, P.; Hildemann, L. M.; McMurry, P. H.; Seinfeld, J. H. (1995) Organics alter hygroscopic behavior of atmospheric particles. J. Geophys. Res. [Atmos.] 100: 18,755-18,770.

Schiavon, N.; Chiavari, G.; Schiavon, G.; Fabbri, D. (1995) Nature and decay effects of urban soiling on granitic building stones. In: Jimenez-Saiz, C., ed. The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation; January 1994; Sevilla, Spain. Sci. Total Environ. 167: 87-101.

Schichtel, B. A.; Husar, R. B.; Falke, S. R.; Wilson, W. E. (2001) Haze trends over the United States, 1980-1995. Atmos. Environ. 35: 5205-5210.

Schier, G. A.; Jensen, K. (1992) Atmospheric deposition effects on foliar injury and foliar leaching in red spruce. In: Eager, C.; Adams, M. B., eds. Ecology and decline of red spruce in the eastern United States. New York, NY: Springer-Verlag; pp. 271-294. [Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. D.; Remmert, H., eds. Ecological Studies, Analysis and Synthesis Ecological Studies, v. 96].

Schimek, M. G. (1981) Smoothing and regression. New York, NY: Wiley.

Schnug, E. (1997) Significance of sulphur for the quality of domesticated plants. In: Cram, W. J.; De Kok, L. J.; Stulen, I.; Brunold, C.; Rennenberg, H., eds. Sulphur metabolism in higher plants: molecular, ecophysiological and nutritional aspects. Leiden, The Netherlands: Backhuys Publishers; pp. 109-130.

Schönherr, J.; Huber, R. (1977) Plant cuticles are polyelectrolytes with isoelectric points around three. Plant Physiol. 59: 145-150.

Schulze, E.-D. (1989) Air pollution and forest decline in a spruce (*Picea abies*) forest. Science (Washington, DC) 244: 776-783.

Schuster, P. F.; Reddy, M. M.; Sherwood, S. I. (1994) Effects of acid rain and sulfur dioxide on marble dissolution. Mater. Perform. 33: 76-80.

Schwar, M. J. R. (1998) Nuisance dust deposition and soiling rate measurements. Environ. Technol. 19: 223-229.

Seckmeyer, G.; McKenzie, R. L. (1992) Increased ultraviolet radiation in New Zealand (45 degrees S) relative to Germany (48 degrees N). Nature 359: 135-137.

Sehmel, G. A. (1980) Particle and gas dry deposition: a review. Atmos. Environ. 14: 983-1011.

Sehmel, G. A.; Hodgson, W. H. (1976) Predicted dry deposition velocities. In: Atmosphere-surface exchange of particulate and gaseous pollutants (1974): proceedings of a symposium; September 1974; Richland, WA. Oak Ridge, TN: Energy Research and Development Administration; pp. 399-422. (ERDA symposium series: 38). Available from: NTIS, Springfield, VA; CONF-740921.

Seinfeld, J. H. (1989) Urban air pollution: state of the science. Science 243: 745-752.

Sereda, P. J. (1974) Weather factors affecting corrosion of metals. In: Corrosion in natural environments: three symposia presented at the seventy-sixth annual meeting [of the] American Society for Testing and Materials; June 1973; Philadelphia, PA. Philadelphia, PA: American Society for Testing and Materials; pp. 7-22. (ASTM special technical publication 558).

Shannon, J. D. (1999) Regional trends in wet deposition of sulfate in the United States and SO2 emissions from 1980 through 1995. Atmos. Environ. 33: 807-816.

Shinn, J. H. (1978) A critical survey of measurements of foliar deposition of airborne sulfates and nitrates. Presented at: 71st annual meeting of the Air Pollution Control Association; June; Houston, TX. Pittsburgh, PA: Air Pollution Control Association; paper no. 78-7.2.

Shortle, W. C.; Bondietti, E. A. (1992) Timing, magnitude, and impact of acidic deposition on sensitive forest sites. Water Air Soil Pollut. 61: 253-267.

Shortle, W. C.; Smith, K. T. (1988) Aluminum-induced calcium deficiency syndrome in declining red spruce. Science (Washington, DC) 240: 1017-1018.

Shortle, W. C.; Smith, K. T.; Minocha, R.; Lawrence, G. B.; David, M. B. (1997) Acidic deposition, cation mobilization, and biochemical indicators of stress in healthy red spruce. J. Environ. Qual. 26: 871-876.

Showak, W.; Dunbar, S. R. (1982) Effect of 1 percent copper addition on atmospheric corrosion of rolled zinc after 20 years' exposure. In: Dean, S. W.; Rhea, E. C., eds. Atmospheric corrosion of metals: a symposium; May 1980; Denver, CO. Philadelphia, PA: American Society for Testing and Materials; pp. 135-162. (ASTM special technical publication 767).

Shriner, D. S.; Henderson, G. S. (1978) Sulfur distribution and cycling in a deciduous forest watershed. J. Environ. Qual. 7: 392-397.

Sievering, H. (1987) Dynamics of sulfur exchange at the air/forest canopy interface: a review of throughfall inferred deposition rates. Global Biogeochem. Cycles 1: 233-249.

Sievering, H.; Rusch, D.; Marquez, L. (1996) Nitric acid, particulate nitrate and ammonium in the continental free troposphere: nitrogen deposition to an alpine tundra ecosystem. Atmos. Environ. 30: 2527-2537.

Simonich; S. L.; Hites, R. A. (1995) Organic pollutant accumulation in vegetation. Environ. Sci. Technol. 29: 2905-2914.

Simpson, J. W.; Horrobin, P. J., eds. (1970) The weathering and performance of building materials. New York, NY: Wiley-Interscience.

Sisler, J. F. (1996) Spatial and seasonal patterns and long term variability of the composition of the haze in the United States: an analysis of data from the IMPROVE network. Fort Collins, CO: Colorado State University, Cooperative Institute for Research in the Atmosphere; EPA air docket A096-56, document no. VI-B-09-(ee).

Sisler, J. F.; Cahill, T. A. (1993) Spacial and temporal patterns and the chemical composition of the haze and its impact on visibility in Alaska. In: Proceedings [of the] 86th annual meeting & exhibition of the Air & Waste Management Association. Volume 8: general environmental topics; June; Denver, CO. Pittsburgh, PA: Air & Waste Management Association; 93-MP-4.03.

Sisler, J. F.; Malm, W. C. (2000) Interpretation of trends of $PM_{2.5}$ and reconstructed visibility from the IMPROVE network. J. Air Waste Manage. Assoc. 50: 775-789.

Sisler, J. F.; Huffman, D.; Lattimer, D. A.; Malm, W. C.; Pitchford, M. L. (1993) Spatial and temporal patterns and the chemical composition of the haze in the United States: an anlysis of data from the IMPROVE network, 1988-1991. Fort Collins, CO: Colorado State University, Cooperative Institute for Research in the Atmosphere (CIRA).

Skeffington, R. A.; Wilson, E. J. (1988) Excess nitrogen deposition: issues for consideration. Environ. Pollut. 54: 159-184.

Skerry, B. S.; Alavi, A.; Lindgren, K. I. (1988) Environmental and electrochemical test methods for the evaluation of protective organic coatings. J. Coat. Technol. 60: 97-106.

Slinn, W. G. N. (1977) Some approximations for the wet and dry removal of particles and gases from the atmosphere. Water Air Soil Pollut. 7: 513-543.

Slinn, W. G. N. (1978) Parameterizations for resuspension and for wet and dry deposition of particles and gases for use in radiation dose calculations. Nucl. Saf. 19: 205-219.

Slinn, W. G. N. (1982) Predictions for particle deposition to vegetative canopies. Atmos. Environ. 16: 1785-1794.

Sloane, C. S. (1983) Optical properties of aerosols–comparison of measurements with model calculations. Atmos. Environ. 17: 409-416.

Sloane, C. S. (1984) Optical properties of aerosols of mixed composition. Atmos. Environ. 18: 871-878.

Sloane, C. S. (1986) Effect of composition on aerosol light scattering efficiencies. Atmos. Environ. 20: 1025-1037.

Sloane, C. S.; Wolff, G. T. (1985) Prediction of ambient light scattering using a physical model responsive to relative humidity: validation with measurements from Detroit. Atmos. Environ. 19: 669-680.

Smith, W. H. (1973) Metal contamination of urban woody plants. Environ. Sci. Technol. 7: 631-636.

Smith, W. H. (1974) Air pollution - effects on the structure and function of the temperate forest ecosystem. Environ. Pollut. 6: 111-129.

Smith, W. H. (1984) Pollutant uptake by plants. In: Treshow, M., ed. Air pollution and plant life. New York, NY: John Wiley & Sons, Ltd.; pp. 417-450.

Smith, W. H. (1990a) Forests as sinks for air contaminants: soil compartment. In: Air pollution and forests: interactions between air contaminants and forest ecosystems. 2nd ed. New York, NY: Springer-Verlag; pp. 113-146. (Springer series on environmental management).

Smith, W. H. (1990b) Forests as sinks for air contaminants: vegetative compartment. In: Air pollution and forests: interactions between air contaminants and forest ecosystems. 2nd ed. New York, NY: Springer-Verlag; pp. 147-180. (Springer series on environmental management).

Smith, W. H. (1990c) Forest nutrient cycling: toxic ions. In: Air pollution and forests: interactions between air contaminants and forest ecosystems. 2nd ed. New York, NY: Springer-Verlag; pp. 225-268. (Springer series on environmental management).

Smith, W. H. (1990d) Forest biotic agent stress: air pollutants and disease caused by microbial pathogens. In: Air pollution and forests: interactions between air contaminants and forest ecosystems. 2nd ed. New York, NY: Springer-Verlag; pp. 366-397. (Springer series on environmental management).

Smith, W. H. (1991) Air pollution and forest damage. Chem. Eng. News 69(45): 30-43.

Smith, K. E. C.; Jones, K. C. (2000) Particles and vegetation: implications for the transfer of particle-bound organic contaminants to vegetation. Sci. Total Environ. 246: 207-236.

Smith, W. H.; Staskawicz, B. J. (1977) Removal of atmospheric particles by leaves and twigs of urban trees: some preliminary observations and assessment of research needs. Environ. Manage. (N. Y.) 1: 317-330.

Soulen, P. E.; Frederick, J. E. (1999) Estimating biologically active UV irradiance from satellite radiance measurements: a sensitivity study. J. Geophys. Res. 104: 4117-4126.

Spence, J. W.; Haynie, F. H. (1972) Paint technology and air pollution: a survey and economic assessment. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Programs; publication no. AP-103. Available from: NTIS, Springfield, VA; PB-210736.

Spence, J. W.; Haynie, F.; Upham, J. B. (1975) Effects of gaseous pollutants on paints: a chamber study. J. Paint Technol. 47: 57-63.

Spence, J. W.; Haynie, F. H.; Lipfert, F. W.; Cramer, S. D.; McDonald, L. G. (1992) Atmospheric corrosion model for galvanized steel structures. Corrosion 48: 1009-1019.

Spiker, E. C.; Comer, V. J.; Hosker, R. P.; Sherwood, S. I. (1992) Dry deposition of SO_2 on limestone and marble: the role of humidity. In: Rodrigues, J. D.; Henriques, F.; Jeremias, F. T., eds. Proceedings of the 7th international congress on deterioration and conservation of stone; June; Lisbon, Portugal. Lisbon, Portugal: Laboratório Nacional de Engenharia Civil; pp. 397-406.

Spinka, J. (1971) Vliv znečištěného ovzduší na ovocné stromy a zeleninu [Effects of polluted air on fruit trees and legumes]. Ziva 19: 13-15.

Steubing, L.; Klee, R. (1970) Vergleichende Untersuchungen zur Staubfilterwirkung von Laub- und Nadelgehölzen [Comparative investigations on the dust-filter effect of broad-leaved and coniferous trees]. Angew. Bot. 44: 73-85.

Stevens, R. K.; Dzubay, T. G.; Lewis, C. W.; Shaw, R. W., Jr. (1984) Source apportionment methods applied to the determination of the origin of ambient aerosols that affect visibility in forested areas. Atmos. Environ. 18: 261-272.

Stoddard, J. L. (1994) Long-term changes in watershed retention of nitrogen: its causes and aquatic consequences. In: Baker, L. A., ed. Environmental chemistry of lakes and reservoirs. Washington, DC: American Chemical Society; pp. 223-284. (Advances in chemistry series no. 237).

Stoddard, J. L.; Murdoch, P. S. (1991) Catskill Mountains. In: Charles, D. F., ed. Acidic deposition and aquatic ecosystems: regional case studies. New York, NY: Springer-Verlag; pp. 237-271.

Storm, G. L.; Fosmire, G. J.; Bellis, E. D. (1994) Persistence of metals in soil and selected vertebrates in the vicinity of the Palmerton zinc smelters. J. Environ. Qual. 23: 508-514.

Strandberg, H.; Johansson, L.-G. (1997) The formation of black patina on copper in humid air containing traces of SO_2. J. Electrochem. Soc. 144: 81-89.

Suzuki, D. T. (1997) Barriers to perception: from a world of interconnection to fragmentation. In: Raven, P. H., ed. Nature and Human Society: the quest for a sustainable world. Washington, DC: National Academy Press; pp. 11-21.

Swietlicki, E.; Zhou, J. C.; Berg, O. H.; Martinsson, B. G.; Frank, G.; Cederfelt, S. I.; Desek, U.; Berner, A.; Birmilli, W.; Wiedensohler, A.; Yuskiewicz, B.; Bower, K. N. (1999) A closure study of sub-micrometer aerosol particle hygroscopic behaviour. Atmos. Res. 50: 205-240.

Swietlik, D.; Faust, M. (1984) Foliar nutrition of fruit crops. Hortic. Rev. 6: 287-355.

Sydberger, T.; Ericsson, R. (1977) Laboratory testing of the atmospheric corrosion of steel. Werkst. Korros. 28: 154-158.

Taback, H. J.; Brienza, A. R.; Macko, J.; Brunetz, N. (1979) Fine particle emissions from stationary and miscellaneous sources in the south coast air basin: appendix. Sacramento, CA: California Air Resources Board; contract no. A6-191-30.

Talbot, J. J. (1979) A review of the potential biological impacts of cooling tower salt drift. Atmos. Environ. 13: 395-405.

Tang, I. N. (1980) Deliquescence properties and particle size change of hygroscopic aerosols. In: Willeke, K., ed. Generation of aerosols and facilities for exposure experiments. Ann Arbor, MI: Ann Arbor Science Publishers, Inc.; pp. 153-167.

Tang, I. N. (1997) Thermodynamic and optical properties of mixed-salt aerosols of atmospheric importance. J. Geophys. Res. [Atmos.] 102: 1883-1893.

Tang, I. N.; Munkelwitz, H. R. (1994) Water activities, densities, and refractive indices of aqueous sulfates and sodium nitrate droplets of atmospheric importance. J. Geophys. Res. [Atmos.] 99: 18,801-18,808.

Tanner, R. L. (1981) An ambient experimental study of phase equilibrium in the atmospheric system: aerosol H^+, NH_4^+, SO_4^{2-}, NO_3^- --NH_3(g), HNO_3(g). In: Fogler, H. S., ed. Chemical reactors: based on a symposium sponsored by the Division of Industrial and Engineering Chemistry at the second chemical congress of the North American continent (180th ACS national meeting); August 1980; Las Vegas, NV. Washington, DC: American Chemical Society; pp. 527-532. (ACS symposium series 168).

Taylor, G. E., Jr.; Hanson, P. J.; Baldocchi, D. D. (1988) Pollutant deposition to individual leaves and plant canopies: sites of regulation and relationship to injury. In: Heck, W. W.; Taylor, O. C.; Tingey, D. T., eds. Assessment of crop loss from air pollutants. New York, NY: Elsevier Applied Science; pp. 227-257.

Tegen, I.; Lacis, A. A. (1996) Modelling of particle size distribution and its influence on the radiative properties of mineral dust aerosol. J. Geophys. Res. Atmos. 101: 19,237-19,244.

Tegen, I.; Lacis, A. A.; Fung, I. (1996) The influence on climate forcing of mineral aerosols from disturbed soils. Nature (London) 380: 419-422.

Thornton, F. C.; Schaedle, M.; Raynal, D. J. (1987) Effects of aluminum on red spruce seedlings in solution culture. Environ. Exp. Bot. 27: 489-498.

Tilman, D. (1996) Biodiversity: population versus ecosystem stability. Ecology 77: 350-363.

Tilman, D. (2000) Causes, consequences and ethics of biodiversity. Nature 405: 208-211.

Tilman, D.; Downing, J. A. (1994) Biodiversity and stability in grasslands. Nature (London) 367: 363-365.

Tjepkema, J. D.; Cartica, R. J.; Hemond, H. F. (1981) Atmospheric concentration of ammonia in Massachusetts and deposition on vegetation. Nature (London) 294: 445-446.

Tombach, I.; Thurston, S. A. (1994) The quality of the SCENES measurements: the roles of data quality goals and evolving technology. In: Aerosols and atmospheric optics: radiative balance and visual air quality: proceedings of the international specialty conference, volume A; September; Snowbird, UT. Pittsburgh, PA: Air & Waste Management Association; pp. 21-32. (A&WMA publication VIP-41).

Tong, S. T. Y. (1991) The retention of copper and lead particulate matter in plant foliage and forest soil. Environ. Int. 17: 31-37.

Trettin, C.C.; Johnson, D.W.; Todd, D.E., Jr. (1999) Forest nutrient and carbon pools at Walker Branch watershed: changes during a 21-year-period. Soil Sci. Soc. Amer. J. 63: 1436-1448.

Trijonis, J.; Shapland, D. (1979) Existing visibility levels in the United States: isopleth maps of visibility in suburban/nonurban areas during 1974-76. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; report no. EPA-450/5-79-010. Available from: NTIS, Springfield, VA; PB80-156177.

Trijonis, J. C.; Pitchford, M.; McGown, M. (1987) Preliminary extinction budget results from the RESOLVE program. In: Bhardwaja, P. S., ed. Visibility protection: research and policy aspects, an APCA international specialty conference; September 1986; Grand Teton National Park, WY. Pittsburgh, PA: Air Pollution Control Association; pp. 872-883. (APCA transactions series no. TR-10).

Trijonis, J. C.; Malm, W. C.; Pitchford, M.; White, W. H.; Charlson, R.; Husar, R. (1991) Visibility: existing and historical conditions--causes and effects. In: Irving, P. M., ed. Acidic deposition: state of science and technology, volume III: terrestrial, materials, health and visibility effects. Washington, DC: The U.S. National Acid Precipitation Assessment Program. (State of science and technology report no. 24).

Turner, J.; Kelly, J. (1981) Relationships between soil nutrients and vegetation in a north coast forest, New South Wales. Aust. For. Res. 11: 201-208.

Turner, J.; Lambert, M. J. (1980) Sulfur nutrition of forests. In: Shriner, D. S.; Richmond, C. R.; Lingberg, S. E., eds. Atmospheric sulfur deposition: environmental impact and health effects, proceedings of the second life sciences symposium; October 1979; Gatlinburg, TN. Ann Arbor, MI: Ann Arbor Science Publishers, Inc.; pp. 321-333.

Turner, J.; Lambert, M. J.; Gessel, S. P. (1977) Use of foliage sulphate concentrations to predict response to urea application by Douglas-fir. Can. J. For. Res. 7: 476-480.

Twomey, S. (1974) Pollution and the planetary albedo. Atmos. Environ. 8: 1251-1256.

Twomey, S. (1977) Atmospheric aerosols. Amsterdam, The Netherlands: Elsevier Scientific Publishing Company; p. 237.

Tyler, G. (1972) Heavy metals pollute nature, may reduce productivity. Ambio 1: 52-59.

U.S. Code. (1977) Clean Air Act, "section"169A, visibility protection for federal class I areas. U. S. C. 42: "section"7491.

U.S. Code. (1990) Clean Air Act, "section"169B, visibility. U. S. C. 42: "section"7492.

U.S. Code. (1991) Clean Air Act, section 108, air quality criteria and control techniques, section 109, national ambient air quality standards. U. S. C. 42: sections 7408-7409.

U.S. Environmental Protection Agency. (1979) Protecting visibility: an EPA report to Congress. Research Triangle Park, NC: Office of Air Quality Planning and Standards; EPA report no. EPA-450/5-79-008. Available from: NTIS, Springfield, VA; PB80-220320.

U.S. Environmental Protection Agency. (1982) Air quality criteria for particulate matter and sulfur oxides. Research Triangle Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; EPA report no. EPA-600/8-82-029aF-cF. 3v. Available from: NTIS, Springfield, VA; PB84-156777.

U.S. Environmental Protection Agency. (1986) Air quality criteria for lead. Research Triangle Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; EPA report no. EPA-600/8-83/028aF-dF. 4v. Available from: NTIS, Springfield, VA; PB87-142378.

U.S. Environmental Protection Agency. (1987) Assessing the risks of trace gases that can modify the stratosphere. Washington, DC: U.S. Environmental Protection Agency, Office of Air Radiation; report no. EPA 400/1-87/001A-H.

U.S. Environmental Protection Agency. (1993) Air quality criteria for oxides of nitrogen. Research Triangle Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; report nos. EPA/600/8-91/049aF-cF. 3v. Available from: NTIS, Springfield, VA; PB95-124533, PB95-124525, and PB95-124517.

U.S. Environmental Protection Agency. (1995a) Interim findings on the status of visibility research. Research Triangle Park, NC: Office of Research and Development; report no. EPA/600/R-95/021.

U.S. Environmental Protection Agency. (1995b) Testing of meteorological and dispersion models for use in regional air quality modeling. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA-454/R-95-005. Available from: NTIS, Springfield, VA; PB95-215638.

U.S. Environmental Protection Agency. (1996a) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.

U.S. Environmental Protection Agency. (1996b) Air quality criteria for ozone and related photochemical oxidants, v. I-III. Research Triangle Park, NC: Office of Research and Development; report no. EPA/600/P-93/004aF-cF. Available from: NTIS, Springfield, VA; PB96-185582. PB96-185590, and PB96-185608. Available: www.epa.gov/ncea/ozone.htm [25 February 2002].

U.S. Environmental Protection Agency. (1997a) Nitrogen oxides: impacts on public health and the environment. Washington, DC: Office of Air and Radiation; August. Available: www.epa.gov/ttncaaa1/t1/reports/noxrept.pdf [1999, November 24].

U.S. Environmental Protection Agency. (1997b) Summary of the particulate monitoring program. Washington, DC: Office of Air and Radiation.

U.S. Environmental Protection Agency. (1998) Monitoring PM2.5 in ambient air using designated reference of class I equivalent methods. In: Quality Assurance Handbook for Air Pollution Measurement Systems, V. II, Part II, Section 2.12. Research Triangle Park, NC: Research and Development, Environmental Monitoring Systems Laboratory.

U.S. Environmental Protection Agency. (1999a) Protection of visibility. C. F. R. 40: "Section" 51.300-51.309.

U.S. Environmental Protection Agency. (1999b) Visibility monitoring guidance. Research Triangle Park, NC: Emissions, Monitoring, and Analysis Division.

U.S. Environmental Protection Agency. (2000a) Deposition of air pollutants to the great waters. Third report to Congress. [Executive Summary]. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; report no. EPA-453/R-00-005.

U.S. Environmental Protection Agency. (2000b) Latest findings on national air quality: 1999 status and trends. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; report no. EPA-454/F-00-002.

U.S. Environmental Protection Agency. (2001) National air quality and emissions trends report, 1999. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA/454/R-01-004. Available: http://www.epa.gov/oar/aqtrnd99/ [11 March, 2002].

Ulrich, B. (1983) Interaction of forest canopies with atmospheric constituents: SO_2, alkali and earth alkali cations and chloride. In: Ulrich, B.; Pankrath, J., eds. Effects of accumulation of air pollutants in forest ecosystems: proceedings of a workshop; May 1982; Göttingen, Federal Republic of Germany. Dordrecht, The Netherlands: D. Reidel Publishing Company; pp. 33-45.

United Nations Environment Programme (UNEP). (1986) Report of the international conference on the assessment of the role of carbon dioxide and of other greenhouse gases in climate variations and associated impacts; October 1985; Villach, Austria. Geneva, Switzerland: World Meteorological Organization; WMO no. 661.

United Nations Environment Programme (UNEP). (1998) Environmental effects of ozone depletion: 1998 assessment. J. Photochem. Photobiol. B 46: 1-4.

United Nations Environment Programme (UNEP). (2000) Environmental effects of ozone depletion: interim summary. Available at: http://www.gcrio.org/ozone/unep2000summary.html [9 April 2002].

Unsworth, M. H. (1984) Evaporation from forests in cloud enhances the effects of acid deposition. Nature (London) 312: 262-264.

Unsworth, M. H.; Wilshaw, J. C. (1989) Wet, occult and dry deposition of pollutants on forests. Agric. For. Meteorol. 47: 221-238.

Van Aalst, R. M. (1982) Dry deposition of NOx. In: Schneider, T.; Grant, L., eds. Air pollution by nitrogen oxides. Amsterdam, The Netherlands: Elsevier Scientific Publishing Company; pp. 263-270.

Van Breemen, N.; Van Dijk, H. F. G. (1988) Ecosystem effects of atmospheric deposition of nitrogen in The Netherlands. In: Dempster, J. P.; Manning, W. J., eds. Excess nitrogen deposition. Environ. Pollut. 54: 249-274.

Van Breemen, N.; Burrough, P. A.; Velthorst, E. J.; Van Dobben, H. F.; De Wit, T.; Ridder, T. B.; Reijnders, H. F. R. (1982) Soil acidification from atmospheric ammonium sulphate in forest canopy throughfall. Nature (London) 299: 548-550.

Van Breemen, N.; Mulder, J.; Driscoll, C. T. (1983) Acidification and alkalinization of soils. Plant Soil 75: 283-308.

Van de Hulst, H. C. (1957, reprint 1981) Light scattering by small particles. New York, NY: Wiley. (Structure of matter series).

Van der Leun, J. C.; Tang, X.; Tevini, M. (1995) Environmental effects of ozone depletion: 1994 assessment. Ambio 24: 138.

Van der Leun, J. C.; Tang, X.; Tevini, M., eds. (1998) Environmental effects of ozone depletion: 1998 assessment. J. Photochem. Photobiol. B: Biology 46(1-3): 1-108.

Van Dijk, H. F. G.; Roelofs, J. G. M. (1988) Effects of excessive ammonium deposition on the nutritional status and condition of pine needles. Physiol. Plant. 73: 494-501.

Van Dorland, R.; Dentener, F. J.; Lelieveld, J. (1997) Radiative forcing due to tropospheric ozone and sulfate aerosols. J. Geophys. Res. [Atmos.] 102: 28,079-28,100.

Van Miegroet, H.; Cole, D. W. (1984) The impact of nitrification on soil acidification and cation leaching in red alder ecosystem. J. Environ. Qual. 13: 586-590.

Van Miegroet, H.; Johnson, D. W.; Todd, D. E. (1993) Foliar response of red spruce saplings to fertilization with Ca and Mg in the Great Smoky Mountains National Park. Can. J. For. Res. 23: 89-95.

Venkatram, A.; Pleim, J. (1999) The electrical analogy does not apply to modeling dry deposition of particles. Atmos. Environ. 33: 3075-3076.

Viles, H. A. (1990) The early stages of building stone decay in an urban environment. Atmos. Environ. 24A: 229-232.

Viskari, E.-L.; Kärenlampi, L. (2000) Roadside Scots pine as an indicator of deicing salt use - a comparative study from two consecutive winters. Water Air Soil Pollut. 122: 405-419.

Vitousek, P. M.; Ehrlich, P. R.; Chrlich, A. H.; Matson, P. A. (1986) Human appropriation of the products of photosynthesis. Bioscience 36: 368-373.

Vitousek, P. M.; Mooney, H. A.; Lubchenco, J.; Melillo, J. M. (1997) Human domination of Earth's ecosystems. Science (Washington, DC) 277: 494-499.

Vogt, K. A.; Dahlgren, R.; Ugolini, F.; Zabowski, D.; Moore, E. E.; Zasoski, R. (1987a) Aluminum, Fe, Ca, Mg, K, Mn, Cu, Zn and P in above- and belowground biomass. I. *Abies amabilis* and *Tsuga mertensiana*. Biogeochemistry 4: 277-294.

Vogt, K. A.; Dahlgren, R.; Ugolini, F.; Zabowski, D.; Moore, E. E.; Zasoski, R. (1987b) Aluminum, Fe, Ca, Mg, K, Mn, Cu, Zn and P in above- and belowground biomass. II. Pools, and circulation in a subalpine *Abies amabilis* stand. Biogeochemistry 4: 295-311.

Von Glasow, R.; Bott, A. (1999) Interaction of radiation fog with tall vegetation. Atmos. Environ. 33: 1333-1346.

Vora, A. B.; Bhatnagar, A. R. (1987) Comparative study of dust fall on the leaves in high pollution and low pollution area of Ahmedabad: V. Caused foliar injury. J. Environ. Biol. 8: 339-346.

Vose, J. M.; Swank, W. T. (1990) Preliminary estimates of foliar absorption of ^{15}N-labeled nitric acid vapor (HNO_3) by mature eastern white pine (*Pinus strobus*). Can. J. For. Res. 20: 857-860.

Vuilleumier, L.; Harley, R. A.; Brown, N. J.; Slusser, J. R.; Kolinski, D.; Bigelow, D. S. (2001) Variability in ultraviolet total optical depth during the Southern California Ozone Study (SCOS97). Atmos. Environ. 35: 1111-1122.

Wagrowski, D. M.; Hites, R. A. (1997) Polycyclic aromatic hydrocarbon accumulation in urban, suburban, and rural vegetation. Environ. Sci. Technol. 31: 279-282.

Wall, D. H. (1999) Biodiversity and ecosystem functioning. BioScience 49: 107-108.

Wall, D. H.; Moore, J. C. (1999) Interactions underground: soil biodiversity, mutualism, and ecosystem processes. Bioscience 49: 109-117.

Walton, J. R.; Johnson, J. B.; Wood, G. C. (1982) Atmospheric corrosion initiation by sulphur dioxide and particulate matter: II. characterisation and corrosivity of individual particulate atmospheric pollutants. Br. Corros. J. 17: 65-70.

Wang, W. C.; Yung, Y. L.; Lacis, A. A.; Mo, T.; Hansen, J. E. (1976) Greenhouse effects due to man-made perturbations of trace gases. Science (Washington, DC) 194: 685-690.

Waring, R. H. (1987) Nitrate pollution: a particular danger to boreal and subalpine coniferous forests. In: Fujimori, T.; Kimura, M., eds. Human impacts and management of mountain forests: [proceedings of a symposium]. Ibaraki, Japan: Forestry and Forest Products Research Institute; pp. 93-105.

Waring, R. H.; Schlesinger, W. H. (1985) The carbon balance of trees. In: Forest ecosystems: concepts and management. Orlando, FL: Academic Press, Inc.; pp. 7-37.

Watson, J. G.; Chow, J. C. (1994) Clear sky visibility as a challenge for society. Annu. Rev. Energy Environ. 19: 241-266.

Watson, J. G.; Robinson, N. F.; Chow, J. C.; Henry, R. C.; Kim, B. M.; Pace, T. G.; Meyer, E. L.; Nguyen, Q. (1990) The USEPA/DRI chemical mass balance receptor model, CMB 7.0. Environ. Software 5: 38-49.

Webb, A. H.; Bawden, R. J.; Busby, A. K.; Hopkins, J. N. (1992) Studies on the effects of air pollution on limestone degradation in Great Britain. Atmos. Environ. Part B 26: 165-181.

Wedding, J. B.; Carlson, R. W.; Stukel, J. J.; Bazzaz, F. A. (1975) Aerosol deposition on plant leaves. Environ. Sci. Technol. 9: 151-153.

Wedin, D. A.; Tilman, D. (1996) Influence of nitrogen loading and species composition on the carbon balance of grasslands. Science 274: 1720-1723.

Weinbaum, S. A.; Neumann, P. M. (1977) Uptake and metabolism of ^{15}N-labeled potassium nitrate by French prune *(Prunus domestica L.)* leaves and the effects of two surfactants. J. Am. Soc. Hortic. Sci. 102: 601-604.

Wellburn, A. R. (1990) Tansley review no. 24: Why are atmospheric oxides of nitrogen usually phytotoxic and not alternative fertilizers? New Phytol. 115: 395-429.

Wells, B. W. (1939) A new forest climax: the salt spray climax of Smith Island, North Carolina. Torrey Bot. Club Bull. 66: 629-634.

Wells, B. W.; Shunk, I. V. (1937) Seaside shrubs: wind forms vs. spray forms. Science 85: 499.

Wells, B. W.; Shunk, I. V. (1938) Salt spray: an important factor in coastal ecology. Torrey Bot. Club Bull. 65: 485-492.

Wenny, B. N.; Schafer, J. S.; DeLuisi, J. J.; Saxena, V. K.; Barnard, W. F.; Petropavlovskikh, I. V.; Vergamini, A. J. (1998) A study of regional aerosol radiative properties and effects on ultraviolet-B radiation. J. Geophys. Res. [Atmos.] 103: 17,083-17,097.

Wesely, M. L. (1989) Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models. Atmos. Environ. 23: 1293-1304.

Wesely, M. L.; Hicks, B. B. (1977) Some factors that affect the deposition rates of sulfur dioxide and similar gases on vegetation. J. Air Pollut. Control Assoc. 27: 1110-1116.

Wesely, M. L.; Eastman, J. A.; Stedman, D. H.; Yalvac, E. D. (1982) An eddy-correlation measurement of NO_2 flux to vegetation and comparison to O_3 flux. Atmos. Environ. 16: 815-820.

Wesely, M. L.; Cook, D. R.; Hart, R. L. (1983) Fluxes of gases and particles above a deciduous forest in wintertime. Boundary Layer Meteorol. 27: 237-255.

Wesely, M. L.; Cook, D. R.; Hart, R. L.; Speer, R. E. (1985) Measurements and parameterization of particulate sulfur dry deposition over grass. J. Geophys. Res. [Atmos.] 90: 2131-2143.

Wesselink, L. G.; Meiwes, K.-J.; Matzner, E.; Stein, A. (1995) Long-term changes in water and soil chemistry in spruce and beech forests, Solling, Germany. Environ. Sci. Technol. 29: 51-58.

Westman, W. E. (1977) How much are nature's services worth? Measuring the social benefits of ecosystem functioning is both controversial and illuminating. Science (Washington, DC) 197: 960-964.
Whitby, K. T. (1978) The physical characteristics of sulfur aerosols. Atmos. Environ. 12: 135-159.
White, W. H. (1976) Reduction of visibility by sulphates in photochemical smog. Nature (London) 264: 735-736.
White, W. H. (1990) The components of atmospheric light extinction: a survey of ground-level budgets. Atmos. Environ. Part A 24: 2673-2679.
White, W. H. (1991) Contributions to light extinction. In: Irving, P. M., ed. Acidic deposition: state of science and technology, v. III: terrestrial, materials, health and visibility effects; report 24. Washington, DC: National Acid Precipitation Assessment Program; pp. 85-102.
White, W. H.; Roberts, P. T. (1977) On the nature and origins of visibility-reducing aerosols in the Los Angeles air basin. Atmos. Environ. 11: 803-812.
Williams, M. W.; Baron, J. S.; Caine, N.; Sommerfeld, R.; Sanford, R. (1996) Nitrogen saturation in the Rocky Mountains. Environ. Sci. Technol. 30: 640-646.
Wilson, E. O. (1997) The creation of biodiversity. In: Raven, P. H., ed. Nature and human society: the quest for a sustainable world: proceedings of the 1997 forum on biodiversity; October; sponsored by the National Research Council. Washington, DC: National Academy Press.
Wilson, W. E.; Suh, H. H. (1997) Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies. J. Air Waste Manage. Assoc. 47: 1238-1249.
Wiman, B. (1981) Aerosol collection by Scots pine seedlings: design and application of a wind tunnel method. Oikos 36: 83-92.
Wiman, B. L. B. (1985) A non-stationary model for sulphur aerosol depletion and deposition in a young spruce stand. Tellus Ser. B 37: 230-242.
Wiman, B. L. B.; Agren, G. I. (1985) Aerosol depletion and deposition in forests--a model analysis. Atmos. Environ. 19: 335-347.
Wiman, B. L. B.; Lannefors, H. O. (1985) Aerosol characteristics in a mature coniferous forest--methodology, composition, sources and spatial concentration variations. Atmos. Environ. 19: 349-362.
Wiman, B. L. B.; Agren, G. I.; Lannefors, H. O. (1985) Aerosol concentration profiles within a mature coniferous forest--model versus field results. Atmos. Environ. 19: 363-367.
Winner, W. E.; Atkinson, C. J. (1986) Absorption of air pollution by plants, and consequences for growth. Trends Ecol. Evol. 1: 15-18.
Winner, W. E.; Bewley, J. D. (1978a) Terrestrial mosses as bioindicators of SO_2 pollution stress: synecological analysis and the index of atmospheric purity. Oecologia 35: 221-230.
Winner, W. E.; Bewley, J. D. (1978b) Contrasts between bryophyte and vascular plant synecological responses in an SO_2-stressed white spruce association in central Alberta. Oecologia 33: 311-325.
Witherspoon, J. P.; Taylor, F. G., Jr. (1971) Retention of 1--44 μ simulated fallout particles by soybean and sorghum plants. Health Phys. 21: 673-677.
Wittenburg, C.; Dannecker, W. (1992) Dry deposition and deposition velocity of airborne acidic species upon different sandstones. In: Proceedings of the 1992 European aerosol conference; September; Oxford, United Kingdom. J. Aerosol Sci. 23(suppl. 1): S869-S872.
Wolff, G. T. (1984) On the nature of nitrate in coarse continental aerosols. Atmos. Environ. 18: 977-981.
Wolff, G. T.; Collins, D. C.; Rodgers, W. R.; Verma, M. H.; Wong, C. A. (1990) Spotting of automotive finishes from the interactions between dry deposition of crustal material and wet deposition of sulfate. J. Air Waste Manage. Assoc. 40: 1638-1648.
Woodcock, A. H. (1953) Salt nuclei in marine air as a function of altitude and wind force. J. Meteorol. 10: 362-371.
World Health Organization. (1997) Nitrogen oxides. 2nd ed. Geneva, Switzerland: World Health Organization. (Environmental health criteria 188).
World Meteorological Organization. (1988) Developing policies for responding to climatic change: a summary of the discussions and recommendations of workshops; September-October 1987; Villach, Austria; and November 1987; Bellagio, Austria. Geneva, Switzerland: World Meteorological Organization; report no. WMO/TD; no. 225. [World Climate Impact Programme series report no. WCIP-1].
World Resources Institute. (2000) World resources 2000-2001: people and ecosystems: the fraying web of life. Washington, DC: World Resources Institute.
Wu, P.-M.; Okada, K. (1994) Nature of coarse nitrate particles in the atmosphere--a single particle approach. Atmos. Environ. 28: 2053-2060.

Wyttenbach, A.; Tobler, L.; Bajo, S. (1989) Na, Cl and Br in needles of Norway spruce (P. abies) and in the aerosol adhering to the needles. Toxicol. Environ. Chem. 19: 25-33.

Xu, J. R.; Balik, C. M. (1989) Infrared attenuated total reflectance study of latex paint films exposed to aqueous sulfur dioxide. J. Appl. Polym. Sci. 38: 173-183.

Yanai, R. D.; Siccama, T. G.; Arthur, M.. A.; Federer, C. A.; Friedland, A. J. (1999) Accumulation and depletion of base cations in forest floors in the northeastern United States. Ecology 80: 2774-2787.

Yarnel, B.; Kalkstein, L. S.; Scheraga, J. D. (2000) Mid-Atlantic regional assessment of climate change impacts. Clim. Res. (CR Special 7) 14: 153-269.

Yerrapragada, S. S.; Jaynes, J. H.; Chirra, S. R.; Gauri, K. L. (1994) Rate of weathering of marble due to dry deposition of ambient sulfur and nitrogen dioxides. Anal. Chem. 66: 655-659.

Yocom, J. E.; Grappone, N. (1976) Effects of power plant emissions on materials. Palo Alto, CA: Electric Power Research Institute; report no. EPRI/EC-139. Available from: NTIS, Springfield, VA; PB-257 539.

Yocom, J. E.; Upham, J. B. (1977) Effects of economic materials and structures. In: Stern, A. C., ed. Air pollution. 3rd ed. Volume II. New York, NY: Academic Press; pp. 93-94.

Young, P. (1996) Pollution-fueled "biodeterioration" threatens historic stone: researchers are probing the role microorganisms play in the decay of buildings and artwork. Environ. Sci. Technol. 30: 206A-208A.

Young, J. R.; Ellis, E. C.; Hidy, G. M. (1988) Deposition of air-borne acidifiers in the western environment. J. Environ. Qual. 17: 1-26.

Youngs, W. D.; Rutzke, M.; Gutenmann, W. H.; Lisk, D. J. (1993) Nickel and vanadium in foliage in the vicinity of an oil-fired power plant. Chemosphere 27: 1269-1272.

Zappia, G.; Sabbioni, C.; Gobbi, G. (1993) Non-carbonate carbon content on black and white areas of damaged stone monuments. Atmos. Environ. 27A: 1117-1121.

Zappia, G.; Sabbioni, C.; Pauri, M. G.; Gobbi, G. (1994) Mortar damage due to airborne sulfur compounds in a simulation chamber. Mater. Struct. 27: 469-473.

Zappia, G.; Sabbioni, C.; Riontino, C.; Gobbi, G.; Favoni, O. (1998) Exposure tests of building materials in urban atmosphere. Sci. Total Environ. 224: 235-244.

Zdanovskii, A. B. (1948) New methods of calculating solubilities of electrolytes in multicomponent systems. Zh. Fiz. Khim. 22: 1475-1485.

Zepp, R. G.; Callaghan, T. V.; Erickson, D. J. (1998) Effects of enhanced solar ultraviolet radiation on biogeochemical cycles. J. Photochem. Photobiol. B 46: 69-82.

Zhang, X. Q.; McMurry, P. H.; Hering, S. V.; Casuccio, G. S. (1993) Mixing characteristics and water content of submicron aerosols measured in Los Angeles and at the Grand Canyon. Atmos. Environ. Part A 27: 1593-1607.

Zipperer, W. C.; Wu, J.; Pouyat, R. V.; Pickett, S. T. A. (2000) The application of ecological principles to urban and urbanizing landscapes. Ecol. Appl. 10: 685-688.

Appendix 4A

Colloquial and Latin Names

Alder, hazel	*Alnus serrulata* (Aiton) Willdenow
Alder, red	*Alnus rubra* Bong.
Bean, common	*Phaseolus vulgaris* L.
Beech	*Fagus sylvatica* L.
Birch, yellow	*Betula alleghaniensis* Britt.
Blackhaw	*Viburnum prunifolium* L.
Brush box	*Lophostemon confertus* (R. BR.) P.G. Wilson & Waterhouse
Ceanothus, hoaryleaf	*Ceanothus crassifolius* Torry
Chaparral	*Ceanothus crassifolius*
Coachwood	*Ceratopetalum apetalum*, D.Don
Corn	*Zea mays* L.
Dogwood, flowering	*Cornus florida* L.
Elm	*Ulmus* spp.
Fir, balsam	*Abies balsamea* (L) Mill.
Fir, Douglas	*Pseudotsuga menziesii* (Mirb.) Franco.
Fir, fraser	*Abies fraseri* (Pursh.) Poir
Grape	*Vitis* spp.
Grass, red brome	*Bromus rubens* L.
Grass, purple moor	*Molina caerulea* (L.) Moench.
Greenbriar	*Smilax* spp.
Gum, sweet	*Liquidambar styraciflua* L.
Haw, black	*Viburnum prunifolium* l.
Heather, Scottish	*Calluna vulgaris* Salisb.

Hickory	*Carya* spp.
Hophornbeam	*Ostrya virginiana* (Mill.) Koch
Ivy, English	*Hedera helix* L.
Laurel, mountain	*Kalmia latifolia* L.
Lichen, monks hood	*Hypogymnia physiodes*
Maize	*Zea mays* L.
Maple, red	*Acer rubrum* L.
Maple	*Acer* spp.
Maple, sugar	*Acer saccharum* Marsh.
Mustard, small podded	*Brassica geniculata* L.
Nettle, stinging	*Urtica dioica* L.
Oak, bur	*Quercus macrocarpa* Michx.
Oak, English	*Quercus rober* L.
Oak, chestnut	*Quercus prinus* = Q. *montana* Willd.
Oak, live	*Quercus virginiana* Mill.
Oak, northern red	*Quercus rubra* L.
Oak, turkey	*Quercus laevis* Walt.
Oak, white	*Quercus alba* L.
Oak	*Quercus* spp.
Oats, domestic	*Avena sativa* L.
Oats, wild	*Avena fatua* L
Persimmon, common	*Diosporos virginiana* L.
Pine, eastern white	*Pinus strobus* L.
Pine, jack pine	Pinus *banksiana* Lamb.
Pine, loblolly	*Pinus taeda* L.
Pine, lodgepole	*Pinus contorta* Loud.
Pine, Scots (Scotch)	*Pinus sylvestris* L.
Pine, slash	*Pinus elliotti* Englem.

Pine, lodgepole x jack pine	*Pinus contorta* (Douglas ex Loud) x P. *banksiana* Lamb.
Poplar, black	*Populus nigra L.*
Poplar, white	*Populus alba* L.
Poplar, yellow or tulip	*Liriodendron tulipifera* L.
Privet	*Ligustrum* spp.
Purple Moor Grass	*Molina caerulea* (L.) Moench.
Ragweed	*Ambrosia* spp.
Rhododendron, Catawba	*Rhododendron catawbiense* Michx.
Rhododendron, rosebay	*Rhododendron maximum* L.
Sage, coastal	*Artemisia californica* Less.
Scottish Heather	*Calluna vulgaris* Salisb.
Soybean	*Glycine max* (L.) Merr.
Spruce, Norway	*Picea abies* (L.) Karst.
Spruce, red	*Picea rubens* Sarg.
Spruce, sitka	*Picea sitchensis* (Bong.) Carr.
Spruce, white	*Picea glauca* (Moench.) Voss.
Sunflower	*Helianthus annuus* L.
Sweetgum	*Liquidambar styraciflua* L.
Tulip poplar	*Liriodendron tulipifera* L.

EUMYCOTA-FUNGI

Zygomycota

Vesicular Arbuscular Mycorrhizae

Scutellospora

Gigaspora

Glomus agrigatum

Glomus leptototicum

Ascomycotina

Chaetomium sp.

Fungi Imperfecti

Aureobasidium pullulans

Cladosporium sp.

Epicoccum sp.

Pestalotiopsis

Phialophora verrucosa

Pleurophomella =Sirodothis

5. HUMAN EXPOSURE TO PARTICULATE MATTER AND ITS CONSTITUENTS

5.1 INTRODUCTION

5.1.1 Purpose

Exposure is defined as the contact by an individual with a pollutant for a specific duration of time at a visible external boundary (modified from Duan 1982, 1991). For airborne particulate matter (PM), the breathing zone is considered the point of contact; and the lung and heart are the target organs of concern. An individual's exposure is measured as the PM air concentration in his/her breathing zone over time. Understanding exposure is important, because it is the individual who experiences adverse health effects associated with elevated PM concentrations. Human exposure data and models provide the link between ambient monitoring data or atmospheric models and lung deposition models to enable estimates of the source- air- exposure-dose relationship for input into dose-response assessments for PM from ambient sources.

The goal of this chapter is to provide current information on the development of human exposure data and models. This includes information on the relationships between PM measured at ambient sites and personal exposures to PM from both ambient and nonambient sources and the factors that effect these relationships. Human exposure data and models presented in this chapter provide the critical link between ambient monitoring data, PM dosimetry, and toxicological studies and epidemiological studies presented in other chapters. Specific objectives of this chapter are fourfold:

(1) To provide an overall conceptual framework of exposure science as applied to PM, including the identification and evaluation of factors that determine personal PM exposure

(2) To provide a concise summary and review of recent data (since 1996) and findings from pertinent PM exposure studies

(3) To characterize quantitative relationships between ambient air quality measurements (mass, chemical components, number, etc.), as determined by a community monitoring site, and total personal PM exposure, as well as its ambient and nonambient components

(4) To evaluate the implications of using ambient PM concentrations as a surrogate for personal exposure in epidemiological studies of PM health effects

The U.S. Environmental Protection Agency's (EPA's) regulatory authority for PM applies primarily to ambient air and those sources that contribute to ambient PM air concentrations. Thus, a major emphasis must be to develop an understanding of exposure to PM from ambient sources. However, personal exposure to total PM may result from exposure to PM from both ambient and nonambient sources, and it is likely that both ambient and nonambient components will have adverse health effects. Ultimately, it will be necessary to account for both in order to fully understand the relationship between PM and health effects. In addition, an individual's personal exposure to ambient, nonambient, and total PM would provide useful information for studies where health outcomes are tracked individually.

5.1.2 Particulate Matter Mass and Constituents

Current EPA PM regulations are based on mass as a function of aerodynamic size. However, EPA also measures the chemical composition of PM in both monitoring and research studies. The composition of PM is variable and (as discussed in Chapters 7 and 8) adverse health effects may be related to PM characteristics other than mass. Since PM from ambient and nonambient sources also may have different physical and chemical characteristics, they may also have different health effects. Ultimately, to understand and control health impacts caused by PM exposures from all sources, it is important to quantify and understand exposure to those chemical constituents responsible from various sources for the adverse health effects.

The National Research Council (NRC) recognized the distinction between measuring exposure to PM mass and to chemical constituents when setting *Research Priorities for Airborne Particulate Matter I: Immediate Priorities and a Long-range Research Portfolio* (NRC, 1998). Specifically, NRC Research Topic 1 recommends evaluating the relationship between outdoor measures versus actual human exposure for PM mass. The NRC Research Topic 2 recommends evaluating exposures to biologically important constituents and specific characteristics of PM that cause responses in potentially susceptible subpopulations and the general population. It also was recognized by the NRC that "a more targeted set of studies under this research topic (#2) should await a better understanding of the physical, chemical, and biological properties of airborne particles associated with the reported mortality and morbidity outcomes" (NRC, 1999).

The NRC also stated that the later studies "should be designed to determine the extent to which members of the population contact these biologically important constituents and size fraction of concern in outdoor air, outdoor air that has penetrated indoors, and air pollutants generated indoors" (NRC, 1999). Thus, exposure studies should include contributions from all sources. The emphasis in this chapter on PM mass reflects the current state of the science. Where available, data also have been provided on chemical constituents, although in most cases, the data are limited. As recognized by the NRC, a better understanding of exposures to PM chemical constituents from multiple sources will be required to more fully identify, understand, and control those sources of PM with adverse health effects and to accurately define the relationship between PM exposure and health outcomes due to either short-term or chronic exposures.

5.1.3 Relationship to Past Documents

Early versions of PM criteria documents did not emphasize total human exposure but rather focused almost exclusively on outdoor air concentrations. For instance, the 1969 Air Quality Criteria for Particulate Matter (National Air Pollution Control Administration, 1969) did not discuss either *exposure* or *indoor concentrations*. The 1982 EPA PM Air Quality Criteria Document (PM AQCD), however, provided some discussion of indoor PM concentrations, reflecting an increase in microenvironmental and personal exposure studies (U.S. Environmental Protection Agency, 1982). The new data indicated that personal activities, along with PM generated by personal and indoor sources (e.g., cigarette smoking), could lead to high indoor levels and high personal exposures to total PM. Some studies reported indoor concentrations that exceeded PM concentrations found in the air outside the monitored microenvironments or at nearby monitoring sites. Between 1982 and 1996, many more studies of personal and indoor PM exposure demonstrated that, in most inhabited domestic environments, indoor PM concentrations and personal PM exposures of the residents were greater than ambient PM concentrations measured simultaneously (e.g., Sexton et al., 1984; Spengler et al., 1985; Clayton et al., 1993). As a result, the NRC (1991) recognized the potential importance of indoor sources of contaminants (including PM) in causing adverse health outcomes.

The 1996 AQCD (U.S. Environmental Protection Agency, 1996) reviewed the human PM exposure literature through early 1996, mainly to evaluate the use of ambient monitors as surrogates for PM exposure in epidemiology studies. Many of the studies cited showed poor

correlations between personal exposure or indoor measurements of PM and outdoor or ambient site measurements. Conversely, Janssen et al. (1995) and Tamura et al. (1996a) showed that in the absence of major nonambient sources, total PM exposures to individuals tracked through time were highly correlated with ambient PM concentrations. Analyses of these latter two studies led to consideration of ambient and nonambient exposures as separate components of total personal exposure. As a result, the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996), for the first time, distinguished between ambient and nonambient PM personal exposure. This chapter builds on the work of the 1996 PM AQCD by further evaluating the ambient and nonambient components of PM, as well as reporting research that evaluates the relationship between ambient concentrations and total, ambient, and nonambient personal exposure.

5.2 STRUCTURE FOR THE CHAPTER

The chapter is organized to provide information on the principles of exposure, review the existing literature, and summarize key findings and limitations in the information; the specific sections are described below.

- Section 5.3 discusses the basic concepts of exposure, including definitions, methods for estimating exposure, and methods for estimating ambient components of exposure.
- Section 5.4 presents PM mass data, including a description of the key available studies, the relationship of PM exposures with ambient concentrations, and factors that affect the relationship.
- Section 5.5 presents data on PM constituents, including a description of the key available studies, the relationship with ambient concentrations, and factors that affect the relationship.
- Section 5.6 discusses the implications of using ambient PM concentrations in epidemiological studies of PM health effects.
- Section 5.7 summarizes key findings and limitations of the information.

5.3 BASIC CONCEPTS OF EXPOSURE

5.3.1 Components of Exposure

The total exposure of an individual over a discrete period of time includes exposures to many different particles from various sources while in different microenvironments (μ**e**'s). Duan (1982) defined a microenvironment as "a [portion] of air space with homogeneous pollutant concentration." It also has been defined as a volume in space, for a specific time interval, during which the variance of concentration within the volume is significantly less than the variance between that microenvironment and surrounding ones (Mage, 1985). In general, people pass through a series of microenvironments, including outdoor, in-vehicle, and indoor microenvironments, as they go through time and space. Thus, total *daily* exposure for a single individual to PM can be expressed as the sum of various exposures for the microenvironments that the person occupies in the day (modified from National Research Council, 1991).

In a given microenvironment, particles may originate from a wide variety of sources. For example, in an indoor microenvironment, PM may be generated by (1) indoor activities, (2) outdoor PM entering indoors, (3) the chemical interaction of outdoor air pollutants and indoor air or indoor sources, (4) transport from another indoor microenvironment, or (5) personal activities. All of these disparate sources have to be accounted for when estimating total human exposure to PM.

An analysis of personal exposure to PM mass (or constituent compounds) requires definition and discussion of several classes of particles and exposure. In this chapter, PM metrics may be described in terms of exposure or as an air concentration. PM also may be described according to both its source (i.e., ambient, nonambient) and the microenvironment where exposure occurs. Table 5-1 provides a summary of the terms used in this chapter, the notation used for these terms, and their definition. These terms are used throughout this chapter and provide the terminology for evaluating personal exposure to total PM and to PM from ambient and nonambient sources.

The 1997 NAAQS were developed largely on the basis of evidence from epidemiological studies that found relatively consistent associations between outdoor particulate matter concentrations and observed health effects. Thus an emphasis in this chapter is on determinations of personal exposure to PM of ambient origin and the relationship between the

TABLE 5-1. CLASSES OF PARTICULATE MATTER EXPOSURE AND CONCENTRATION DEFINITIONS

Term	Notation	Definition
		General Definitions
Concentration	C	Air concentration of PM in a given microenvironment, expressed in $\mu g/m^3$
Personal Exposure	E	Contact at visible external boundaries of an individual with a pollutant for a specific duration of time; quantified by the amount of PM available in concentration units ($\mu g/m^3$) at the oral/nasal contact boundary for a specified time period (**Δt**). General term for any exposure variable.
Microenvironment	μ**e**	Volume in space, for a specific time interval, during which the variance of concentration within the volume is significantly less than the variance between that μ**e** and surrounding μ**e**s
		Concentration Variables
Ambient PM	C_a	PM in the atmosphere measured at a community ambient monitoring site either emitted into the atmosphere directly (primary PM) or formed in it (secondary PM). Major sources of PM species are industry, motor vehicles, commerce, domestic emissions such as wood smoke, and natural wind-blown dust or soil.
Ambient-Outdoor PM	C_{ao}	Ambient PM in an outdoor microenvironment
Indoor PM	C_i	All PM found indoors
Ambient-Indoor PM	C_{ai}	Ambient PM that has infiltrated indoors (i.e., has penetrated indoors and remains suspended)
Primary Indoor-generated PM	C_{pig}	Primary PM generated indoors
Secondary Indoor-generated PM	C_{sig}	Secondary PM generated by outdoor vapors reacting with indoor vapors
		Exposure Variables
Personal Exposure to Indoor-Generated PM	E_{pig}	Sum of personal exposure resulting from primary indoor-generated PM
Personal Exposure to Indoor-Formed PM	E_{sig}	Sum of personal exposure resulting from secondary indoor-generated PM
Personal Exposure to Personal-Activity PM	E_{pact}	Small-scale PM-generating activities that primarily influence exposure of the person performing the activity itself
Personal Exposure to Nonambient PM	E_{nonag}	Sum of personal exposure to indoor-generated and personal activity PM $E_{nonag} = E_{sig} + E_{sig} + E_{pact}$
Personal Exposure to Ambient-Generated PM	E_{ag}	Sum of personal exposure caused by ambient-outdoor and ambient indoor PM (does not include resuspended ambient PM previously deposited indoors)
Personal Exposure to Total PM	E_t	Sum of all personal exposures to ambient and nonambient PM $E_t = E_{pig} + E_{sig} + E_{pact} + E_{ag} = E_{nonag} + E_{ag}$

PM concentrations measured at ambient sites and personal exposure to PM. Although this is an emphasis, it should be kept in mind that every particle that deposits in the lung becomes part of a dose delivered to the individual. It is likely that the nonambient component of total exposure also has health effects which would not be detected using community time-series epidemiology studies. Since both ambient and nonambient components of PM exposure may have partial influence on the ultimate dose and the health outcome, both components should be understood and accounted for when assessing risk from PM and its constituents.

5.3.2 Methods To Estimate Personal Exposure

Personal exposure may be estimated using either direct or indirect approaches. *Direct approaches* measure the contact of the person with the chemical concentration in the exposure media over an identified period of time. Direct measurement methods include personal exposure monitors (PEMs) for PM that are worn continuously by individuals as they encounter various microenvironments and perform their daily activities. *Indirect approaches* use models and available information on concentrations of chemicals in microenvironments, the time individuals spend in those microenvironments, and personal PM generating activities to estimate personal exposure. This section describes the methods to directly measure personal exposures and microenvironmental concentrations, as well as the models used to estimate exposure. Several approaches to estimate personal exposure to ambient PM also are described.

5.3.2.1 Direct Measurement Methods

5.3.2.1.1 Personal Exposure Monitoring Methods

In theory, personal exposure to total PM is measured by sampling the concentration of PM in inhaled air entering the nose or mouth. Practically, it is defined as that PM collected by a PEM worn by a person and sampling from a point near the breathing zone (but not impacted by exhaled breath). PEMs for PM use measurement techniques similar to those used for ambient PM. The PEM is a filter-based mass measurement of a particle size fraction (PM_{10} or $PM_{2.5}$), usually integrated over either a 24- or 12-h period at flow rates of 2 to 4 L/min using battery-operated pumps. PEMs must be worn by study participants and, therefore, they must be quiet, compact, and battery-operated. These requirements limit the type of pumps and the total sample volume that can be collected. Generally, small sample volumes limit personal exposure

measurements to PM mass and a few elements detected by XRF. In most studies, $PM_{2.5}$ and PM_{10} have not been collected concurrently; thus, for personal exposure, there are very few data available by which to estimate coarse thoracic PM (i.e., $PM_{10-2.5}$).

Other methods used for ambient PM also have been adapted for use as a personal exposure monitor. For example, a personal nephelometer that measures particle number within a specific particle size range using light scattering has been used in personal exposure studies to obtain real-time measurements of PM.

5.3.2.1.2 Microenvironmental Monitoring Methods

Direct measurements of microenvironmental PM concentrations, which are used with models to estimate personal exposure to PM, also use methods similar to those for ambient PM. These methods differ from PEMs in that they are stationary with respect to the microenvironment (such as a stationary PEM). Microenvironmental monitoring methods include filter-based mass measurements of particle size fractions (PM_{10}, $PM_{2.5}$), usually integrated over either a 24- or 12-h period. Flow rates vary between various devices from 4 to 20 L/min. Larger sample volumes allow more extensive chemical characterization to be conducted on microenvironmental samples. Because more than one pumping system can be used in a microenvironment, $PM_{2.5}$ and PM_{10} can be collected simultaneously. Other continuous ambient PM measurement methods that have been utilized for microenvironmental monitoring are the Tapered Element Oscillating Microbalance (TEOM) and nephelometers. Various continuous techniques for counting particles by size (Climet, LASX, SMPS, APS) also have also been used. Measurement techniques are discussed in Chapter 2.

5 3.2.2 Indirect Methods (Modeling Methods)

5.3.2.2.1 Personal Exposure Models

Exposure modeling for PM mass ($PM_{2.5}$ and $PM_{10-2.5}$) and chemical constituents is a relatively new field facing significant methodological challenges and input data limitations. Exposure models typically use one of two general approaches: (1) a time-series approach that estimates microenvironmental exposures sequentially as individuals go through time or (2) a time-averaged approach that estimates microenvironmental exposures using average microenvironmental concentrations and the total time spent in each microenvironment. Although

the time-series approach to modeling personal exposures provides the appropriate structure for accurately estimating personal exposures (Esmen and Hall, 2000; Mihlan et al., 2000), a time-averaged approach typically is used when the input data needed to support a time-series model are not available. In addition, the time-varying dose profile of an exposed individual can be modeled only by using the time-series approach (McCurdy, 1997, 2000). We define the *personal exposure* of an individual to a chemical in air to be (Lioy, 1990; NRC, 1991)

$$E = \int_{t=t_1}^{t=t_2} C(t)\, dt, \tag{5-1}$$

where

E is the personal exposure during the time period from t_1 to t_2, and

C(t) is the concentration near the nose and mouth not impacted by exhaled air, at time t.

Even though the processes that lead to exposure are nonlinear in nature, personal exposure models are often used to combine microenvironmental concentration data with human activity pattern data in order to estimate personal exposures. Activity pattern data and information on size, age, gender, and health status can be used to estimate inhalation rate. Time-averaged models also can be used to estimate personal exposure for an individual or for a defined population. Total personal exposure models estimate exposures for all of the different microenvironments in which a person spends time, and total average personal exposure is calculated from the sum of these microenvironmental exposures:

$$E = \sum_j E_j = \frac{1}{T}\left(\sum_j \overline{C}_j t_j\right), \tag{5-2}$$

where E_j is the personal exposure in each microenvironment, *j* (Duan, 1982). Example microenvironments include outdoors, indoors at home, indoors at work, and in transit. Each microenvironmental exposure, E_j, is calculated from the average concentration in microenvironment *j*, $\overline{C}_j$, weighted by the time spent in microenvironment *j*, t_j. T is the sum of t_j

over all *j*. It is important to note that, although measurement data may be an average concentration over some time period (i.e., 24 h), significant variations in PM concentrations can occur during that time period. Thus, an error may be introduced if real-time concentrations are highly variable, and an average concentration for a microenvironment is used to estimate exposure when the individual is in that microenvironment for only a fraction of the total time. This model has been applied to concentration data in a number of studies (Ott, 1984; Ott et al., 1988, 1992; Miller et al., 1998; Klepeis et al., 1994; Lachenmyer and Hidy, 2000).

Microenvironmental concentrations used in the exposure models can be measured directly or estimated from one or more microenvironmental models. Microenvironmental models vary in complexity, from a simple indoor/outdoor ratio to a multi-compartmental mass-balance model. A discussion of microenvironmental models is presented below in Section 5.3.2.2.2.

On the individual level, the time spent in the various microenvironments is obtained from time/activity diaries that are completed by the individual. For population-based estimates, the time spent in various microenvironments is obtained from human activity databases. Many of the largest human activity databases have been consolidated by EPA's National Exposure Research Laboratory (NERL) into one comprehensive database called the Consolidated Human Activity Database (CHAD). CHAD contains over 22,000 person-days of 24-h activity data from 11 different human activity pattern studies (McCurdy et al., 2000). Population cohorts with diverse characteristics can be constructed from the activity data in CHAD and used for exposure analysis and modeling (McCurdy, 2000). These databases can also be used to estimate inhalation rates based on activity levels, age, gender, and weight. Table 5-2 is a summary listing of the human activity studies in CHAD.

Methodologically, personal exposure models can be divided into three general types: (1) statistical models based on empirical data obtained from one or more personal monitoring study, (2) simulation models based upon known or assumed physical relationships, and (3) physical-stochastic models that include Monte Carlo or other techniques to explicitly address variability and uncertainty in model structure and input data (Ryan, 1991; MacIntosh et al., 1995). The attributes, strengths, and weaknesses of these model types are discussed by Ryan (1991), National Research Council (1991), Frey and Rhodes (1996), and Ramachandran and Vincent (1999). A recent summary review of the logic of exposure modeling is found in Klepeis (1999).

TABLE 5-2. ACTIVITY PATTERN STUDIES INCLUDED IN THE CONSOLIDATED HUMAN ACTIVITY DATABASE (CHAD)

Study Name	Calendar Time Period of the Study	Age[1]	Days[2]	Diary		Rate[5]	Documentation or Reference	Notes
				Type[3]	Time[4]			
Baltimore	Jan-Feb 1997 Jul-Aug 1998	65+	391	Diary; 15-min blocks	24h Standard	No	Williams et al. (2000a,b)	Multiple days, varying from 5-15; part of a $PM_{2.5}$ PEM study
CARB: Adolescents and Adults	Oct 1987-Sept 1988	12 - 94	1,762	Retrospective	24h Standard	No	Robinson et al. (1989) Wiley et al. (1991a)	
CARB: Children	Apr 1989-Feb 1990	0 - 11	1,200	Retrospective	24h Standard	No	Wiley et al. (1991b)	
Cincinnati (EPRI)	Mar-Apr and Aug 1985	0 - 86	2,614	Diary	24h; nominal 7 p.m.-7 a.m.	Yes	Johnson (1989)	3 consecutive days; 186 P-D removed[7]
Denver (EPA)	Nov 1982-Feb 1983	18 - 70	805	Diary	24h; nominal 7 p.m.-7 a.m.	No	Akland et al. (1985) Johnson (1984)	Part of CO PEM[6] study; 2 consec. days; 55 P-D removed[7]
Los Angeles: Elem. School Children	Oct 1989	10 - 12	51	Diary	24h Standard	Yes	Spier et al. (1992)	7 P-D removed[7]
Los Angeles: High School Adoles.	Sept-Oct 1990	13 - 17	43	Diary	24h Standard	Yes	Spier et al. (1992)	23 P-D removed[7]
National: NHAPS-A[8]	Sept 1992-Oct 1994	0 - 93	4,723	Retrospective	24h Standard	No[9]	Klepeis et al. (1995) Tsang and Klepeis (1996)	A national random-probability survey
National: NHAPS-B[8]	As above	0 - 93	4,663	Retrospective	24h Standard	No[9]	As above	As above
University of Michigan: Children	Feb-Dec1997	0 - 13	5,616	Retrospective	24h Standard	No	Institute for Social Research (1997)	2 days of data: one is a weekend day
Valdez, AK	Nov 1990-Oct 1991	11 - 71	401	Retrospective	Varying 24-h period	No	Goldstein et al. (1992)	4 P-D removed[7]
Washington, DC (EPA)	Nov 1982-Feb 1983	18 - 98	699	Diary	24h; nominal 7 p.m.-7 a.m.	No	Akland et al. (1985) Hartwell et al. (1984)	Part of a CO PEM[6] study; 6 P-D removed[7]

Notes: [1]All studies included both genders. The age range depicted is for the subjects actually included; in most cases, there was not an upper limit for the adult studies. Ages are inclusive. Age 0 = babies < 1 year old.
[2]The actual number of person-days of data in CHAD after the "flagging" and removal of questionable data. See the text for a discussion of these procedures.
[3]Retrospective: a "what did you do yesterday" type of survey; also known as an ex post survey. Diary: a "real-time" paper diary that a subject carried as he or she went through the day.
[4]Standard = midnight-to-midnight.
[5]Was activity-specific breathing rate data collected?
[6]PEM = a personal monitoring study. In addition to the diary, a subject carried a small CO or $PM_{2.5}$ monitor throughout the sampling period.
[7]P-D removed = The number of person-days of activity pattern data removed from consolidated CHAD because of missing activity *and* location information; completeness criteria are listed in the text.
[8]National Human Activity Pattern Study; A = the air version; B = the water version. The activity data obtained on the two versions are identical.
[9]A question was asked regarding which activities (within each 6-h time block in the day) involved "heavy breathing", lifting heavy objects, and running hard.

DRAFT-DO NOT QUOTE OR CITE

Personal exposure models that have been developed for PM are summarized in Table 5-3. The regression-based models (Johnson et al., 2000; Janssen et al., 1997; Janssen et al., 1998a) were developed for a specific purpose (i.e., to account for the observed difference between personal exposure and microenvironmental measurements) and are based on data from a single study, which limits their utility for broader purposes. Other types of models in Table 5-3 were limited by a lack of data for the various model inputs. For example, ambient PM monitoring data is not generally of adequate spatial and temporal resolution for these models. Lurmann and Korc (1994) used site-specific coefficient of haze (COH) information to stochastically develop a time series of 1-h PM_{10} data from every sixth day 24-h PM_{10} measurements. A mass-balance model typically was used for indoor microenvironments when sufficient data were available, such as for a residence. For most other microenvironments, indoor/outdoor ratios were used because of the lack of data for a mass-balance model. In addition, only the deterministic model PMEX included estimation of inhaled dose from activity-specific breathing rate information. Data from recent PM personal exposure and microenvironmental measurement studies will help facilitate the development of improved personal exposure models for PM.

An integrated human exposure source-to-dose modeling system that will include exposure models to predict population exposures to environmental pollutants, such as PM, currently is being developed by EPA/NERL. A first-generation population exposure model for PM, called the Stochastic Human Exposure and Dose Simulation (SHEDS-PM) model, recently has been developed. The SHEDS-PM model uses a 2-stage Monte Carlo sampling technique previously applied by MacIntosh et al. (1995) for benzene exposures. This technique allows for separate characterization of variability and uncertainty in the model predictions (to predict the distribution of total exposure to PM for the population of an urban/metropolitan area and to estimate the contribution of ambient PM to total PM exposure). Results from a case study using data from Philadelphia have been reported (Burke et al., 2001). Recently, the SHEDS model has been extended by EOHSI scientists to provide estimates of integrated PM doses for different regions of the lung for the Philadelphia case study population (Vyas et al, 2002). The inhalation model uses dosimetry equations that account for anatomic, metabolic, and physical variability information specified in the ICRP and HUMTRM models. These efforts are still preliminary but critical for generating population based exposure and dose estimates by utilizing the available dosimetric information and models described in Section 6. Ultimately, comprehensive

TABLE 5-3. PERSONAL EXPOSURE MODELS FOR PARTICULATE MATTER

Study Citation	Model Name	Model Type	Microenvironments or Predictors	Output	Notes
Time-Series Models:					
Hayes and Marshall (1999)	PMEX	Deterministic	Indoors: residential, work, school Outdoors: near roadway, other Motor vehicle	Inhaled dose of PM_{10} Hourly for 24 h By age/gender groups Source contributions	Used IAQM Used human activity data with activity-specific breathing rate info.
Johnson et al. (2000)		Regression-based	Auto travel, roadside, ETS, food prep. grilling, high ambient PM	$PM_{2.5}$ exposure 24-h average	Developed from scripted activity study (Chang et al., 2000)
Klepeis et al. (1994)		Stochastic	ETS, cooking, cleaning, attached garage, wood burning	Respirable particle ($PM_{3.5}$) exposure	
Lurmann and Korc (1994)	REHEX-II	Stochastic	12 residential with different sources, restaurant/bar, nonresidential indoors, in transit, outdoors	Distribution of PM_{10} exposure for population Three averaging times (1 h, 24 h, season)	Fixed I/O ratio of 0.7 for indoors w/o sources and 1.2 for in transit Reduced form mass balance model for indoors with PM sources
Koontz and Niang (1998)	CPIEM	Stochastic	Indoors: residence, office, industrial plant, school, public building, restaurant/lounge, other Outdoors, in vehicle	Distribution of PM_{10} exposure for population	Used California activity pattern and breathing rate data. Used either a mass balance model or I/O ratio distribution for indoor microenvironments. Indoor sources included.
Time-Averaged Models:					
Clayton et al. (1999a)	SIM	Stochastic		Distribution of annual $PM_{2.5}$ exposures	Based on 3-day ambient measurements
Janssen et al. (1997)		Regression-based	Smoking parent, ETS exposure, outdoor physical activity	Accounts for difference between personal and microenvironmental PM_{10}	Children only
Janssen et al. (1998a)		Regression-based	Number of cigarettes smoked, hours of ETS exposure, residence on busy road, time in vehicle	Accounts for difference between personal and microenvironmental PM_{10}	Adults only
Ott et al. (2000)	RCS	Statistical	Not separated	Distribution of PM_{10} exposure for population	A random-component superposition (RCS) model that uses distribution of ambient PM_{10} and estimated nonambient PM_{10} concentrations. Results for Ontario, Canada not corrected for 72-h compared to 24-h averaging time in Riverside, CA and Phillipsburg, NJ.
Burke et al. (2001)	SHEDS-PM	Stochastic	Outdoors, indoors: residence, office, stores, school, in vehicle, restaurant/lounge,	$PM_{2.5}$ exposure distributions for population, by age, gender, smoking and employment status; $PM_{2.5}$ exposure uncertainty predictions. Percent contribution from PM of ambient origin to total personal exposures	A 2-stage Monte-Carlo simulation model for predicting population distribution of daily- average personal exposures to PM. Model has been applied to Philadelphia using spatially and temporally interpolated $PM_{2.5}$ ambient measurements from 1992-1993 and 1990 census data. Does not consider $PM_{2.5}$ exposure from active smoking or exposure in subways.
Chao and Tung (2001)	None	Mass Balance with Empirical corrections	Indoors in unoccupied residences in Hong Kong	Predictions of ambient PM in indoor microenvironments	Model makes corrections for nonideal mixing (residence with multiple compartments with limited intermixing).

DRAFT-DO NOT QUOTE OR CITE

evaluation of PM pollution and health data will utilize both exposure and dose metrics generated for subgroups of concern.

5.3.2.2.2 Microenvironmental Models

The mass balance model has been used extensively in exposure analysis to estimate PM concentrations in indoor microenvironments (Calder, 1957; Sexton and Ryan, 1988; Duan, 1982, 1991; McCurdy, 1995; Johnson, 1995; Klepeis et al., 1995; Dockery and Spengler, 1981; Ott, 1984; Ott et al., 1988, 1992, 2000; Miller et al., 1998; Mage et al., 1999; Wilson et al., 2000). The mass balance model describes the infiltration of particles from outdoors into the indoor microenvironment, the removal of particles in indoor microenvironments, and the generation of particles from indoor sources:

$$V \; dC_i \; / dt = v \, P \, C_a - v \, C_i - kVC_i + Q_i \, , \qquad (5\text{-}3)$$

where V = volume of the well-mixed indoor air (cubic meters),
C_i = concentration of indoor PM;
ν = volumetric air exchange rate between indoors and outdoors (cubic meters per hour);
P = penetration ratio, the fraction of ambient (outdoor) PM that is not removed from ambient air during its entry into the indoor volume;
C_a = concentration of PM in the ambient air (micrograms per cubic meter);
k = removal rate (per hour); and
Q_i = indoor sources of particles (micrograms per hour).

Q_i contains a variety of indoor, particle-generating sources, including: combustion or mechanical processes; condensation of vapors formed by combustion or chemical reaction; suspension from bulk material; and resuspension of previously deposited PM. The removal rate, *k*, includes dry deposition to interior surfaces by diffusion, impaction, electrostatic forces, and gravitational fallout. It may include other removal processes, such as filtration by forced air heating, ventilation, or air-conditioning (HVAC), or by independent air cleaners. All parameters

except V are functions of time. P and *k* also are functions of particle aerodynamic diameter ν, and house characteristics. All variables in Equation 5-3 will have distributions within the population and, in some cases, may change by a factor of 5 to 10. It is important to determine the distribution of these variables. Sensitivity and uncertainty analyses are necessary when attempting to explain the results.

In addition to the mass balance model, a number of single-source or single-microenvironment models exist. However, most are used to estimate personal exposures to environmental tobacco smoke (ETS). These models include both empirically based statistical models and physical models based on first principles; some are time-averaged, whereas others are time-series. These models evaluate the contribution of ETS to total PM exposure in an enclosed microenvironment and can be applied as activity-specific components of total personal exposure models. Examples of ETS-oriented personal exposure models are Klepeis (1999), Klepeis et al. (1996, 2000), Mage and Ott (1996), Ott (1999), Ott et al. (1992, 1995), and Robinson et al. (1994).

5.3.2.3 Methods for Estimating Personal Exposure to Ambient Particulate Matter

In keeping with the various components of PM exposure described above in Section 5.3.1, personal exposure to PM can be expressed as the sum of exposure to particles from different sources summed over all microenvironments in which exposure occurs. Total personal exposure may be expressed as:

$$E_t = E_{ag} + E_{pig} + E_{pact} + E_{sig}$$
$$E_t = E_{ag} + E_{nonag}, \qquad (5\text{-}4)$$

where E_t is the total personal exposure to ambient and nonambient PM, E_{ag} is personal exposure to ambient PM (the sum of ambient PM while outdoors and ambient PM that has infiltrated indoors, while indoors), E_{pig} is personal exposure to indoor-generated PM, E_{pact} is personal exposure to PM from personal activity, E_{sig} is exposure to indoor-formed PM, and E_{nonag} is personal exposure to nonambient PM. Again, this is a linear simplification of personal exposures and ignores possible synergisms or interaction among indoor and outdoor pollutants. Although

personal exposure to ambient and nonambient PM cannot be measured directly, they can be calculated or estimated from other measurement data. Approaches for estimating these components of PM exposure are described in the following section.

5.3.2.3.1 Mass Balance Approach

Ambient-Indoor Concentrations of Particulate Matter

The mass balance model described above (Equation 5-3) has been used to estimate PM concentrations in indoor microenvironments. This model also may be used to estimate ambient-indoor (C_{ai}) and indoor-generated (C_{pig}) PM concentrations. The mass balance model can be solved for C_{ai} and C_{pig} assuming equilibrium conditions, and assuming that all variables remain constant (Ott et al., 2000; Dockery and Spengler, 1981; Koutrakis et al., 1992). By substituting $dC_{ai} + dC_{pig}$ for dC_i in Equation 5-3 and assuming dC_{ai} and $dC_{pig} = 0$, ambient-indoor PM (C_{ai}) and indoor-generated PM (C_{pig}), at equilibrium, are given by

$$C_{ai} = (C_{ao} / P\,a) \Big/ (a + k) \tag{5-5}$$

$$C_{pig} = \left(Q_i / V\right) \Big/ (a + k)\,, \tag{5-6}$$

where $a = \nu/V$, the number of air exchanges per hour. Equations 5-5 and 5-6 assume equilibrium conditions and, therefore, are valid only when the parameters k, a, C_{ao}, and Q_i are not changing rapidly and when the Cs are averaged over several hours. It should be understood that equilibrium is a simplification of indoor microenvironments that are occupied by residents. This assumption of equilibrium may only represent a virtual set of individuals or populations at risk. Under certain conditions (e.g., air-conditioned homes, homes with HVAC or air cleaners that cycle on and off, ambient pollutants with rapidly varying concentrations), nonequilibrium versions of the mass balance model (Ott et al., 2000; Freijer and Bloeman, 2000; Isukapalli and Georgopoulos, 2000) are likely to provide a more accurate estimate of C_{ai} and C_{pig}. However, the equilibrium model provides a useful, if simplified, example of the basic relationships (Ott et al., 2000).

Equation 5-5 may be rearranged further to give C_{ai}/C_{ao}, the equilibrium fraction of ambient PM that is found indoors, defined as the infiltration factor (F_{INF}) (Dockery and Spengler, 1981).

$$F_{INF} = \frac{C_{ai}}{C_{ao}} = \frac{P\,a}{a + k} \tag{5-7}$$

The penetration ratio (P) and the decay rate (*k*) can be estimated using a variety techniques. A discussion of these variables and estimation techniques is given in Section 5.4.3.2.2. Because both P and *k* are a function of particle aerodynamic diameter, air exchange rate, and housing characteristics, F_{INF} also will be a function of these parameters. As a result F_{INF} may present substantial variability within a population. Distributions of this parameter should be estimated to understand the uncertainty and variability associated with estimating exposure to PM of ambient origin.

Personal Exposure to Ambient-Generated Particulate Matter

Personal exposure to ambient-generated PM (E_{ag}) may be estimated using ambient-indoor PM concentration (C_{ai}) from the mass balance model, ambient outdoor PM concentrations (C_{ao}) and information on the time an individual spent in the various microenvironments. Mathematically, this may be expressed as

$$\begin{aligned} E_{ag} &= yC_{ao} + (1-y)C_{ai} \\ &= y\,C_{ao} + (1-y)\,C_{ao}\left[\frac{Pa}{(a+k)}\right], \end{aligned} \tag{5-8}$$

where *y* is the fraction of time that an individual spent outdoors, and $(1 - y)$ is the fraction of time spent indoors.

It is convenient to express personal exposure to ambient generated PM (E_{ag}) as the product of the ambient PM concentration (C_{ao} or C_a) and a personal exposure or attenuation factor. Following the usage in several recent papers (Zeger et al., 2000; Dominici et al., 2000; Ott et al.,

2000), the symbol α will be used for this attenuation factor. Equation 5-8 can be rearranged to obtain an expression for α:

$$\alpha = \frac{E_{ag}}{C_{ao}} = y + (1 - y)\left[\frac{P\,a}{a + k}\right]. \qquad (5\text{-}9)$$

Substituting equation 5-7 in equation 5-9 gives a relationship for α in terms of the infiltration factor F_{INF} and the fraction of time spent in the various microenvironments:

$$\alpha = y + (1 - y)\, F_{INF}. \qquad (5\text{-}10)$$

Thus, personal exposures to ambient PM (E_{ag}) may be calculated from measurable quantities:

$$E_{ag} = \alpha\, C_{ao}. \qquad (5\text{-}11)$$

The factor α can be measured directly or calculated from measured or estimated values of the parameters *a*, *k*, and P and the time spent in various microenvironments from activity pattern diaries (Wilson et al., 2000). Since α depends on housing factors and lifestyle factors, air exchange rate, and PM deposition rate, it could vary to a certain extent from region to region and from season to season. Consequently, predicted exposures based on these physical modeling concepts will provide exposure distributions derived conceptually as resulting from housing, lifestyles, and meteorological considerations. For any given population the coefficient α may represent substantial intra- and inter-personal variability, based on personal activities, housing characteristics, particle size, and composition. Distributions of α should be determined using population studies in order to evaluate the uncertainty and variability associated with model exposures.

The use of a mass balance model to separate personal exposure into two components because of exposure to ambient and nonambient concentrations is not novel. This approach, based on Equation 5-4 as given in Duan (1982) and called superposition of component

concentrations, has been applied using multiple microenvironments estimate exposures to carbon monoxide (Ott, 1984; Ott et al., 1988, 1992), volatile organic compounds (Miller et al., 1998), and particles (Koutrakis et al., 1992; Klepeis et al., 1994). However, in these studies, and in most of the exposure literature, the ambient and nonambient components are added to yield a personal exposure from all sources of the pollutant. The use of the mass balance model, ambient concentrations, and exposure parameters to estimate exposure to ambient-generated PM and exposure to indoor-generated PM separately as different classes of exposure has been discussed in Wilson and Suh (1997) and in Wilson et al. (2000).

5.3.2.3.2 Tracer Species as Surrogates of Ambient-Generated Particulate Matter

The ratio of personal exposure to ambient concentration for a PM component that has no indoor sources has often been used to calculate α (Wilson et al., 2000). Sulfate, in particular, is often used as a marker of outdoor air in indoor microenvironments (Jones et al., 2000). It is found primarily in the $PM_{2.5}$ fraction of the aerosol (Cohen et al., 2000) and is formed in the ambient air via photochemical oxidation of gaseous sulfur dioxide arising from the primary emissions from the combustion of fossil fuels containing sulfur. It also arises from the direct emissions of sulfur-containing particles from nonanthropogenic sources (e.g., volcanic activity, wind-blown soil). In the indoor environment, the only common sources of sulfate may be resuspension by human activity of deposited PM containing ammonium sulfates or soil sulfates that were tracked into the home. In some homes an unvented kerosene heater using a high-sulfur fuel may be a major contributor during winter (Leaderer et al., 1999). Use of matches to light cigarettes or gas stoves can also be a source of sulfates. Studies that have used sulfate as a surrogate for ambient PM are discussed in Section 5.4.3.1 (i.e., Oglesby et al., 2000; Sarnat et al., 2000; Ebelt et al., 2000). When there are no indoor sources of fine-mode sulfates, one may deduce that the ambient-to-personal relationship found for sulfates probably would be the same as that for particulate matter of the same aerodynamic size range and physical/chemical properties. This assumption has not been validated, however; and ambient PM with different physical or chemical characteristics may not behave similarly to sulfate.

5.3.2.3.3 Source-Apportionment Techniques

Source apportionment techniques provide a method for determining personal exposure to PM from specific sources. If a sufficient number of samples are analyzed with sufficient compositional detail, it is possible to use statistical techniques to derive source category signatures, identify indoor and outdoor source categories, and estimate their contribution to indoor and personal PM. Daily contributions from sources that have no indoor component can be used as tracers to generate exposure to ambient PM of similar aerodynamic size or directly as exposure surrogates in epidemiologic analyses. Studies that have used source-apportionment are discussed in Section 5.4.3.3 (i.e., Özkaynak and Thurston, 1987; Yakovleva et al., 1999; Mar et al. 2000; Laden et al., 2000).

5.4 SUMMARY OF PARTICULATE MATTER MASS DATA

5.4.1 Types of Particulate Matter Exposure Measurement Studies

A variety of field measurement studies have been conducted to quantify personal exposure to PM mass, to measure microenvironmental concentrations of PM, to evaluate relationships between personal exposure to PM and PM air concentrations measured at ambient sites, and to evaluate factors that affect exposure. In general, exposure measurement studies are of two types, depending on how the participants are selected for the study. In a *probability* study, participants are selected using a probability sampling design where every member of the defined population has a known, positive probability of being included into the sample. Probability study results can be used to make statistical inferences about the target population. In a *purposeful* or *nonprobability* design, any convenient method may be used to enlist participants and the probability of any individual in the population being included in the sample is unknown. Participants in purposeful samples (also referred to as a “convenience” samples) may not have same the characteristics that would lead to exposure as the rest of the unsampled population. Thus, results of purposeful studies apply only to the subjects sampled on the days that they were sampled and not to other periods of time. Although such studies may report significant differences, confidence intervals, and *p* values, they do not have inferential validity (Lessler and Kalsbeek,1992). Purposeful studies may have generalizability (external validity). The extent of

generalizability is a matter of judgement based on study participant characteristics. Purposeful studies of PM personal exposure can provide data to develop relationships on important exposure factors and useful information for developing and evaluating either statistical or physical/chemical human exposure models.

Regardless of the sampling design (probability or purposeful) there are two general categories of study design that can be used to measure personal exposure to PM and evaluate the relationship between personal PM exposure levels and ambient PM concentrations measured simultaneously: longitudinal and cross-sectional. These are discussed in Section 5.4.3.1.1.

5.4.2 Available Data

5.4.2.1 Personal Exposure Data

Table 5-4 gives an overview of the personal exposure studies that have been conducted and are reviewed in this section. This includes studies that have been reported since the 1996 PM AQCD. Major studies that were reported before that time also have been included to provide a comprehensive evaluation of data in this area. Table 5-4 gives information on the sampling and study designs, the study population, the season, number of participants, PM exposure metric, and the PM size fraction measured.

Although there are a number of studies listed in the table, the data available to evaluate longitudinal relationships and the factors that influence these are limited. Few are based on probability sampling designs that allow study results to be inferred to the general population and to develop distributional data or exposures and the factors that affect exposure. Unfortunately, none of these probability studies uses a longitudinal study design. This limits our ability to provide population estimates and distributional data on the relationship between personal PM exposures and ambient site measurements. In addition, most of the probability studies of PM exposure were conducted during a single season; thus, variations in ambient concentrations, air exchange rates, and personal activities are not accounted for across seasons. In these cases, study results are only applicable to a specific time period. Longitudinal studies, on the other hand, generally have small sample sizes and use a purposeful sampling design. Some studies did not include ambient site measurements to allow comparisons with the exposure data, and approximately half of these studies monitored $PM_{2.5}$. Only one or two studies measured both PM_{10} and $PM_{2.5}$ to provide information on $PM_{10-2.5}$.

TABLE 5-4. SUMMARY OF RECENT PM PERSONAL EXPOSURE STUDIES

Study Design	Study Location and Population	No. of Subjects	Study Period	Age	Days per Subject	PM Exposure[a] Metrics	PM Size Measured[b]	Co-Pollutant Metrics	Reference
Probability Studies									
Pooled	Riverside, CA, PTEAM	178	Fall 1990	10-70	1 (12 h)	P, I, O, A	PM_{10}		Clayton et al. (1993) Özkaynak et al. (1996a,b)
Pooled	Basel, Switzerland, EXPOLIS	50	1997		1 (48 h)	P, I, P	$PM_{2.5}$	VOC, CO, NO_2, S, K, Pb, Br, Ca	Oglesby et al. (2000) Jantunen et al. (1998)
Pooled	Toronto, Canada	732	9/ 1995 - 8/1996	16+	3	P, I, O, A	$PM_{2.5}$ (12 mo) PM_{10} (3 mo)		Clayton et al. (1999a), Pellizzari et al. (1999)
Pooled	Mexico City	66	1992	< 65	1	P, I, O	PM_{10}		Santos-Burgoa et al. (1998)
Purposeful Studies									
Longitudinal	Wageningen, Netherlands, school children	13	1995	10-12	6	P, A, School	$PM_{2.5}$, PM_{10}		Janssen et al. (1999a)
Longitudinal	Amsterdam (Am), Helsinki (Hls), elderly angina or coronary heart disease	41 (Am) 49 (Hls)	Winter 1998 Spring 1999	50-84	22 (Am) 27 (Hls)	P, I, O	$PM_{2.5}$		Janssen et al. (2000)
Longitudinal	Baltimore, elderly healthy and COPD	21	7-8/1998	72-93	5-22	P, I, O , A	$PM_{2.5}$, PM_{10}	CO, O_3, NO_2, SO_2	Williams et al. (2000a,b)
Longitudinal	Fresno I Fresno II (elderly)	5 16	Feb, 1999 Apr-May 1999	60+	24 24	P, I, O, A P, I, O, A	$PM_{2.5}$, PM_{10} $PM_{2.5}$, PM_{10}	CO, O_3	Evans et al. (2000)
Longitudinal	Los Angeles, elderly COPD subjects	30	Summer/ Fall 1996	56 - 83	4	P, I, O	$PM_{2.5}$		Linn et al. (1999)
Longitudinal	Boston, COPD subjects	18	Winter 1996-7 Summer 1996		12	P, I, O, A	$PM_{2.5}$, PM_{10}		Rojas-Bracho et al. (2000)
Longitudinal	Nashville, TN, COPD subjects	10	Summer 1995	36-88	6	P, I, O	$PM_{2.5}$, PM_{10}		Bahadori et al. (2001)
Longitudinal	Vancouver, British Columbia, COPD	16	April-Sept, 1998	54-86	7	P, A	$PM_{2.5}$, PM_{10}		Ebelt et al. (2000)
Longitudinal	Amsterdam and Wageningen, Neth., school children	45	1994, 1995	10-12	4-8	P, A, School	PM_{10}		Janssen et al. (1997)

DRAFT-DO NOT QUOTE OR CITE

TABLE 5-4 (cont'd). SUMMARY OF RECENT PM PERSONAL EXPOSURE STUDIES (SINCE 1996)

Study Design	Study Location and Population	No. of Subjects	Study Period	Age	Days per Subject	PM Exposure[a] Metrics	PM Size Measured[b]	Co-Pollutant Metrics	Reference
Purposeful Studies (cont'd)									
Longitudinal	Amsterdam, adults	37	1994	51-70	5-8	P, I, A	PM_{10}		Janssen et al. (1998a)
Longitudinal	Baltimore, elderly subjects	15	Summer 1998, Spring 1999	75 ± 6.8	12	P	$PM_{2.5}$, PM_{10}	O_3, NO_2, SO_2 VOCs	Sarnat et al. (2000)
Longitudinal	Baltimore, elderly, COPD, children	56	Summer 1998, Winter 1999	Adults: 75±6.8 Children: 9-13 COPD: 65±6.6	12	P, I, O, A	$PM_{2.5}$	O_3, NO_2, SO_2, CO, EC,/OC. VOC	Sarnat et al. (2000)
Longitudinal	Tokyo, Japan, elderly housewives	18	1992		3	P, I, O, A	SPM	NO_2	Tamura et al. (1996a)
Longitudinal	Osaka, Japan	26	Fall 1990-1995		Multiple days	P, I, O	PM_2,PM_{2-10},$PM_{>10}$		Tamura et al. (1996b)
Pooled	Milan, Italy, office workers	100	Spring/summer and winter		1	P, Home, Office, Commuting	PM_{10}	NO_2, CO, VOCs	Carrer et al. (1998)
Probability Sample, Pooled	Indianapolis, IN	240	1996	16 - ?	One 72-h sample/subject	P, I, A, O	$PM_{2.5}$, PM_{10}	Mn, Al, Ca	Pellizzari et al. (2001)
Pooled	Banská Bystrica, Slovakia	49	1997-1998	15 - 59	1	P, I, O, A	PM_{10}, $PM_{2.5}$	SO4=, nicotine	Brauer et al. (2000)
Longitudinal	Wageningen, NL	13	??????	12 - 14	5 - 8	P, A, I at school	$PM_{2.5}$, PM_{10}	None	Janssen et al. (1999a)
Longitudinal µE diary	Mpala, Kenya	252	1996-1998	5 - 75	2 years	I	Undefined Optical MIE	CO	Ezzati and Kammen (2001)
Individual	Brunei	5	1998	Adult	1	P , PI, PO	Undefined Optical MIE	None	Muraleedharan TR, Radojevic M (2000).
Longitudinal	London, UK	10	1997	9-11	5day/season 3 seasons	P, I, O	$PM_{2.5}$, PM_{10}	None	Wheeler et al. 2000
Pooled	Zurich, CH	10	1998	Adults	12h/day for 3days	P, I, O	Pollen	None	Riediker et al. 2000

[a]All based on gravimetric measurements.
[b]P = personal, I = indoors, O = outdoors, A = ambient.

DRAFT-DO NOT QUOTE OR CITE

Four large-scale probability studies that quantify personal exposure to PM under normal ambient source conditions have been reported in the literature. These include the EPA's Particle Total Exposure Assessment Methodology (PTEAM) study (Clayton et al., 1993; Özkaynak et al., 1996a,b); the Toronto, Ontario, study (Clayton et al., 1999a and Pellizzari et al., 1999); the Air Pollution Exposure Distribution within Adult Urban Populations in Europe (EXPOLIS) exposure study (Jantunen et al., 1998, 2000; Oglesby, et al., 2000); and a study of a small, highly polluted, area in Mexico City (Santos-Burgoa et al., 1998). Only preliminary results have been reported for the EXPOLIS study. A fifth study conducted in Kuwait during the last days of the oil-well fires (Al-Raheem et al., 2000) is not reported here because the ambient PM levels were not representative of normal ambient source conditions.

Recent longitudinal exposure studies have focused on potentially susceptible subpopulations such as the elderly with preexisting respiratory and heart diseases (hypertension, chronic obstructive pulmonary disease, and congestive heart disease). This is in keeping with epidemiologic studies that indicate mortality associated with high levels of ambient $PM_{2.5}$ is greatest for elderly people with cardiopulmonary disease (U.S. Environmental Protection Agency, 1996). Longitudinal studies were conducted in the Netherlands by Janssen (1998) and Janssen et al. (1997, 1998a,b, 1999b,c) on purposefully selected samples of adults (50 to 70 years old) and children (10 to 12 years old). Several additional studies have focused on nonsmoking elderly populations in Amsterdam and Helsinki (Janssen et al., 2000), Tokyo (Tamura et al., 1996a), Baltimore (Liao et al., 1999; Williams et al., 2000a,b,c), and Fresno, CA (Evans et al. 2000). These cohorts were selected because of the low incidence of indoor sources of PM (such as combustion or cooking). This should allow an examination of the relationship between personal and ambient PM concentrations without the large influences caused by smoking, cooking, and other indoor particle-generating activities. The EPA has a research program focused on understanding PM exposure characteristics and relationships. Within the program, longitudinal studies are being conducted on elderly participants with underlying heart and lung disease (COPD, patients with cardiac defibrillator, and myocardial infarction), an elderly environmental justice cohort, and asthmatics. These studies are being conducted in several cities throughout the United States and over several seasons. Only preliminary data are currently available, and results are not reported in this document.

A series of studies by Phillips et al. (1994, 1996, 1997a,b, 1998a,b, 1999) examined personal ETS exposure in several European cities. Participants varied by age and occupation. Respirable Particulate Matter (RSP) concentrations were reported. These studies are not included in Table 5-4, because of their focus on ETS exposure (which is not the focus of this chapter). A small personal exposure study in Zurich, Switzerland, was reported by Monn et al., (1997) for PM_{10}. This study also is not listed in Table 5-4, because indoor and outdoor measurements were not taken simultaneously with the personal measurements and other details of the study were not published.

5.4.2.2 Microenvironmental Data

Usually, personal PM monitoring is conducted using integrated measurements over a 12- or 24-h period. As such, total PM exposure estimates based on PEM measurements do not capture data from individual microenvironments. Recent studies have examined PM concentrations in various microenvironments using a number of different types of instruments ranging from filter-based to continuous particle monitors. Details on the instruments used, measurements collected, and findings of these studies according to microenvironment (residential indoor, nonresidential indoor, and traffic-related) are summarized in Table 5-5. Those studies which collected microenvironmental data as part of a personal exposure monitoring study are summarized in Table 5-4. In general, the studies listed in Table 5-5 are relatively small, purposeful studies designed to provide specific data on the factors that effect microenvironmental concentration of PM from both ambient and nonambient sources.

Recently published studies have used various types of continuous monitors to examine particle concentrations in specific microenvironments and resulting from specific activities. Continuous particle monitors such as the SMPS, APS, and Climet have been used to measure particle size distributions in residential microenvironments (Abt et al., 2000a; Long et al., 2000a; Wallace et al., 1997; Wallace, 2000a; McBride et al., 1999; Vette et al., 2001). These studies have been able to assess penetration efficiency for ambient particles to indoor microenvironments, as well as penetration factors and deposition rates. Continuous instruments are also a valuable tool for assessing the impact of particle resuspension caused by human activity. A semi-quantitative estimate of PM exposure can be obtained using personal nephelometers that measure PM using light-scattering techniques. Recent PM exposure studies

TABLE 5-5. SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction	Summary of Measurements	Notes/Findings
Residential Indoor: Nonsmoking Homes					
Abt et al. (2000a) Boston, MA	2 homes, 2 seasons, 6 days	SMPS APS		Detailed indoor/outdoor traces of PM in various size classes for different air exchange rates (< 1h $^{-1}$ to > 2 h^{-1})	Major indoor sources of PM: cooking, cleaning, human activity
Long et al. (2000a) Boston, MA	9 homes, 2 seasons	SMPS TEOM	0.02-10	Continuous PM distributions and size distributions obtained for indoor and outdoor air	Sources of fine particles: cooking and outdoor particles; Sources of coarse particles: cooking, cleaning, indoor activities. 50% of particles by volume generated by indoor events were ultrafine particles.
Anuszewski et al. (1998) Seattle, WA	9 homes, 18 days	Nephelometer (radiance)	2.5	Simultaneous indoor and outdoor PM measured continuously; 1-h avg time, I/O = 0.98; air exchange rate: 0.7-1.7 h^{-1}	Homes contained asthmatic children, heavy wood burning. Dominant source of fine particles was outdoor air.
Leaderer et al. (1999) Southwest, VA	58 homes, summer		10	24 h mean: Regional air 26.0 ± 11.5 μg/m^3 (n = 47); Outdoor homes 28.0 ± 17.7 μg/m^3 (n = 43); Indoor w/ AC 28.9 ± 18.7 μg/m^3 (n = 49); Indoor w/o AC 33.3 ± 14.2 μg/m^3 (n = 8)	Epidemiological study of maternal and infant health effects associated with indoor air pollution
			2.5	24 h mean: Regional air 20.2 ± 9.9 μg/m^3 (n = 50); Outdoor homes 21.8 ± 14.8 μg/m^3 (n = 43); Indoor w/ AC 18.7 ± 13.2 μg/m^3 (n = 49); Indoor w/o AC 21.1 ±7.5 μg/m^3 (n = 9)	Indoor PM concentrations were lower for homes with air conditioning (AC) than non-air-conditioned homes.
Wallace et al. (1997), Wallace (2000b) Reston, VA	1 home, 4 years	SMPS Climet PAHs Black carbon	6 size bins; 100 size channels 0.01- 0.4 μm	Time activity data, whole-house air exchange rates Continuous carbon monoxide: descriptive data for monitored pollutants; size profiles for six indoor particle sources	0.3- to 0.5-μm particles linked to outdoor concentrations, frying, broiling; 0.5- to 2.5-μm particles related to cooking events; >2.5-μm particles influenced by physical movement.
Howard-Reed et al. (2000) Fresno, CA Baltimore, MD	15 participants	Nephelometer (personal MIE) PEM	0.1-10 2.5	Continuous (15-min avg) PM and time activity data; 24-h PM mass; participants from Baltimore and Fresno PM panel studies. Descriptive statistics from each study for five microenvironments	Time-series plots of personal nephelometer data showed that each participant's PM exposure consisted of a series of short-term peaks, imposed on a background caused by ambient PM concentrations.
Rea et al. (2001) Baltimore, MD Fresno, CA	15 participants	Nephelometer (personal MIE) PEM	0.1-10 2.5 and 10	Continuous (15-min avg) PM and time activity data; 24-h PM mass; Modeled PM mass and time activity data to apportion time spent in a location. Good comparison with nephelometer mass (6-20%)	54 ± 31% of average daily $PM_{2.5}$ exposure occurred indoor residences, where participants spent 83 ± 10% of their time. A significant portion of $PM_{2.5}$ exposure occurred where participants spent 4-13% of their time.

DRAFT-DO NOT QUOTE OR CITE

TABLE 5-5 (cont'd). SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction	Summary of Measurements	Notes/Findings
Residential Indoor: Nonsmoking Homes (cont'd)					
Quintana et al. (2000) San Diego, CA	Asthmatic children indoor and outdoor 9 homes	Nephelometer (personal MIE) Harvard impactors TEOM	0.1-10 2.5 and 10	Indoor and outdoor measurements collected using passive, active, and active heated nephelometers for comparison to PM mass measurements.	Nephelometer correlates best with $PM_{2.5}$ vs. Indoor $PM_{2.5}$ r = 0.66 vs. indoor PM_{10} r = 0.13 vs. outdoor $PM_{2.5}$ r = 0.42 vs. outdoor PM_{10} r = 0.20
Chang et al. (2000) Baltimore, MD	1 person performing predetermined activities	"Roll around" monitor (RAS) ($PM_{2.5}$, CO, VOC, O_3, NO_2, SO_2)	2.5	1-h personal exposures measured simultaneously. Personal and ambient concentrations were compared.	1-h personal O_3 exposures were significantly lower in indoor than outdoor microenvironments. 1-h personal CO exposures were highest in vehicles. Personal and ambient $PM_{2.5}$ correlations were strongest for outdoor microenvironments and those with high air exchange rates (i.e., vehicles).
Lioy et al. (1999) NA	10 vacuum cleaners		0.3-0.5	Vacuum cleaners ranged in collection efficiency from 29-99%.	Substantial fine particle emissions from motors with emission rates from 0.028 - 128.8 μg/min.
Ezzati and Kammen (2001) Mpala , Kenya	55 Native Huts 2- years	MiniRam (MIE)	Not specified. Optical Device detects particles 1-10 um, but it is not PM_{10}	Measured PM surrounding wood fires in unvented huts. PM measures were up to 8000 µg/m3, but uncalibrated against wood smoke	Exposures were related to ARI
Chao and Tung (2001) Hong Kong	5 unoccupied homes measured indoors and outdoors, along with air exchange rates	Dust-Trak (TSI)	$PM_{2.5}$ real time, calibrated against an Andersen Mark II	In the limit as air exchange goes to zero, there appears to be a r esidual source, perhaps from drafts or thermal effects. Above air exchange rates of 4.5/hr penetration goes to 1, but indoor turbulence resuspends previously settled $PM_{2.5}$	Developed an excellent model for ambient PM infiltration in the absence of anthropogenic indoor sources.
Fischer et al. (2000) Amsterdam, NL	Measured traffic related differences of PM and VOCs, indoor/outdoor in 18 paired homes at varying distances from traffic.	Harvard Impactors	$PM_{2.5}$ and PM_{10}. EC was measured by reflectance of the $PM_{2.5}$ filters. PAH also measured as indicator of Diesel traffic	Outdoor PM_{10} and $PM^{2.5}$ were approximately 15-20% higher at higher traffic streets than at the quiet streets on the same days. However, much larger differences were found for PAH and EC which are traffic specific	"This [study] supports the use [of] traffic related pollution mapping as an exposure proxy in large-scale epidemiologic studies into health effects of motorized traffic emissions."

DRAFT-DO NOT QUOTE OR CITE

TABLE 5-5 (cont'd). SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction	Summary of Measurements	Notes/Findings
Residential Indoor: Nonsmoking Homes (cont'd)					
Kingham et al. (2000) Huddersfield, UK	Measured PM at ten homes of non-smokers, < 50 m and >300m from traffic.	Harvard Impactors	$PM_{2.5}$ and PM_{10} and PAH. EC measured by filter reflectance .	Median Indoor/outdoor ratio =1 (no indoor combustion sources)	Found an absence of a spatial gradient, perhaps due to wind direction effects (e.g. sometimes upwind and sometimes downwind of traffic)
Morawska et al. (2001) Brisbane, Australia	Measured PM indoors and outdoors at 16 homes while residents were absent. Air exchange rate estimated, not measured.	Scanning mobility particle sizer, aerodynamic particle sizer, and a TSI dust-trak	Submicron PM, Supramicron PM, $PM_{2.5}$	For supra and sub micron particles, indoor = outdoor for normal ventilation conditions of > 2 air changes/hour.	Average outdoor PM concentrations are good estimates of average indoor concentrations of PM of ambient origin for air exchange rates of > 0.5/h
Residential Indoor: Other Home Types					
Brauer et al. (1996) Mexico	22 rural Mexican homes (smoking and nonsmoking)	Inertial impactor Nephelometer (Radiance)	10 2.5	Indoor $PM_{2.5}$: 132-555 $\mu g/m^3$ PM_{10}: 282-768 $\mu g/m^3$. Outdoor $PM_{2.5}$: 37 $\mu g/m^3$ PM_{10}: 68 $\mu g/m^3$; I/O $PM_{2.5}$: 1.8-12.4; PM_{10}: 4.7-10.0	Variety of cooking fuels used Nephelometer data were highly correlated with $PM_{2.5}$ and PM_{10} indoors (r = ≈0.87-0.95)
Jenkins et al. (1996 a,b) 16 U.S. Cities	Smoking and nonsmoking homes	Fluoropore membrane filters	Particle phase ETS markers	Mean $PM_{3.5}$ concentrations were 17-20 $\mu g/m^3$ in smoking homes over nonsmoking homes	
McBride et al. (1999) NA	Combustion source (incense) and walking (1 room, carpeted)	Met-One laser particle counter		Ratios of particle counts a 1.0 and 5.7 m from the combustion source/activity were obtained	Proximity to source may help explain the existence of a personal cloud.
Vette et al. (2001) Fresno, CA	Detached semioccupied residence	SMPS LASX	0.01-2.5	Temporal relationships between indoor and outdoor aerosol concentrations evaluated; penetration factors and deposition rates estimated. Fresno panel study empty residence	Diurnally variable indoor/outdoor aerosol concentration ratios because of resuspension from daytime activities. Penetration factors ranged from 0.5 to 0.9.

DRAFT-DO NOT QUOTE OR CITE

TABLE 5-5 (cont'd). SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction	Summary of Measurements	Notes/Findings
Nonresidential Microenvironments					
Bohadana et al. (2000)	Manufacturing plant, woodworkers		Not given	443 personal time-weighted average occupations samples of airborne dust	
Donham et al. (2000) San Francisco, CA	34 poultry workers	NIOSH Method 0600 monitors probed respirators	5	Total dust sampled indoor respiratory masks. Personal monitoring: 630 ± 980 $\mu g/m^3$ (n = 210) ranging from 10-7,730 $\mu g/m^3$	Respirable dust constituted about 10% of total dust measured.
Klepeis et al. (1996) San Francisco, CA	Airport lounge, ETS	TSI 8510 piezobalance	3.5	Estimated cigarette emission rate of 1.43 mg/min/cigarette.	Personal exposures to ETS can be modeled in these types of microenvironments.
Nieuwenhuijsen et al. (1999)	Agricultural activities		4	Average respirable fraction: 4.5 mg/m^3	
Teschke et al. (1999)	Wood production wood finishing wood construction workers		≈50	1,632 observations from 1979-1997. Arithmetic mean exposure: 7.93 mg/m^3 Geometric mean exposure: 1.86 mg/m^3	
Baek et al. (1997) Korea	Indoor and outdoor smoking restaurants		3.5	Indoor concentrations: 33-475 $\mu g/m^3$ Outdoor concentrations: 12-172 $\mu g/m^3$ I/O: 2.4	No significant correlation between indoor and outdoor measurements.
Ott et al. (1996) California	Bar before and after smoking prohibited	Piezobalance	3.5	Smoking permitted: indoor 26.3-182 $\mu g/m^3$; outdoor <5-67 $\mu g/m^3$ Smoking prohibited: indoor 4-82 $\mu g/m^3$; outdoor 2-67 $\mu g/m^3$	I/O nonsmoking: 2.2. I/O smoking: 3.4
Houseman et al. (2001) Boston, MA	Indoor and outdoor restaurants, stores	TSI DUSTRAK	10	Indoor restaurants: 14-278 $\mu g/m^3$ Outdoor restaurants: 7-281 $\mu g/m^3$ Indoor stores: 12-206 $\mu g/m^3$ Outdoor stores: 7-281 $\mu g/m^3$	Avg I/O for restaurants: 2.3 Not known if the restaurants allowed smoking In stores, indoor and outdoor measurements were correlated, avg I/O: 0.83
Brauer and Mannetje (1998) Vancouver, BC	indoor restaurants, various smoking policies		2.5 10	Nonsmoking: $PM_{2.5}$ 7-65 $\mu g/m^3$; PM_{10} <10-74 $\mu g/m^3$ Restricted smoking (>40% nonsmoking) $PM_{2.5}$ 11-163 $\mu g/m^3$; PM_{10} 24-89 $\mu g/m^3$ Unrestricted smoking: $PM_{2.5}$ 47-253 $\mu g/m^3$; PM_{10} 51-268 $\mu g/m^3$	
Lee and Chang (1999) Hong Kong	indoor and outdoor 5 classrooms		10	Indoor PM_{10}: 30-470 $\mu g/m^3$ Outdoor PM_{10}: 20-617 $\mu g/m^3$	

DRAFT-DO NOT QUOTE OR CITE

TABLE 5-5 (cont'd). SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction	Summary of Measurements	Notes/Findings
Traffic-Related Microenvironments					
Praml and Schierl (2000) Munich, Germany	Trams and buses, rural and urban	Continuous millipore polycarbonate filter	10	n = 201 4-h trips, mean concentration 155 $\mu g/m^3$ range: 13-686 $\mu g/m^3$ I/O: 2.8	Tram > circular bus route > radial bus route Day > night
Alm et al. (1999) Kuopio, Finland	9-km commuter route, rush hours 1/mo	Climet	6 channels	Windows closed, vents open air exchange rate 36-47 h^{-1}	Morning commutes were generally higher than afternoon commutes; relationships determined between PM and wind speed and vehicle speed
Monn et al. (1997) Switzerland	Spatial scale from a city street	Harvard μe monitor	10	48- or 72-h avg times; horizontal distance from street: 0, 15, 50, and 80 m; vertical distance from street: 20 m	Mean PM_{10} 27.3 ± 3.0 $\mu g/m^3$ No vertical gradient (0-20 m) and horizontal gradient (0-80 m) in distance from road, each about 13%. No significant differences between wet and dry periods.
Rodes et al. (1998) Saevanenso, Los Angeles, CA	In-vehicle, various road types, 2-h trips		2.5 10	Air exchange rates measured at various ventilation settings and speeds. Monitoring vehicle followed a diesel bus or truck.	Vehicles in front of the monitored vehicle accounted for most of the in-vehicle commuting exposure; average I/O: 0.6-0.8 h^{-1} for $PM_{2.5}$; carpool lane concentrations were 30-60% lower than noncarpool lane concentrations
Roorda-Knape et al. (1998) van Vliet et al. (1997) Netherlands	Gradient in distance from roadway	Harvard impactor	2.5 10 Black smoke	PM monitoring at 50, 100, 150, and 300 m from roadway; 1-week avg time	No concentration gradient with increasing distance from the roadways for $PM_{2.5}$ and PM_{10}; concentration gradient did exist for black smoke, also found an effect with wind direction
Houseman et al. (2001) Boston, MA	Indoor and outdoor vehicles buses, subways	TSI DUSTRAK	PM_{10}	Vehicle concentrations ranged from 33-170 $\mu g/m^3$. Outdoor vehicle concentrations ranged from 40-144 $\mu g/m^3$. Bus concentrations: 17-268 $\mu g/m^3$; outdoor 10-203 $\mu g/m^3$ Subway: 28-174 $\mu g/m^3$; outdoor 8-203 $\mu g/m^3$	The average in-vehicle to outdoor ratio was 0.99. Average I/O: 3; subway values were correlated with outdoor concentrations.
Brauer et al. (1999) Vancover, BC	Commuting environments	APC-1000		PM < 5: greatest concentrations by combustion powered vehicles PM > 5: greatest concentrations by bicycling and buses	
Janssen et al. (1997) Netherlands	Background and roadway		2.5 10	$PM_{2.5}$ background: 21-35 $\mu g/m^3$; roadway 23-43 $\mu g/m^3$ PM_{10} background:13-32 and 29-62 $\mu g/m^3$; roadway 16-56 and 30-75 $\mu g/m^3$	Average roadway/background ratio: 3 for $PM_{2.5}$ and PM_{10}. Average increase in concentration at the roadway 7.2-12.7 $\mu g/m^3$.

DRAFT-DO NOT QUOTE OR CITE

have used personal nephelometers (1 min avg time) to measure PM continuously (Howard-Reed et al., 2000; Quintana et al., 2000) in various microenvironments. These data have been used to identify the most important ambient and nonambient sources of PM, to provide an estimate of source strength, and to compare modeled time activity data and PEM 24-h mass data to nephelometer measurements (Rea et al., 2001). Several studies also have examined PM exposure in vehicles using both continuous and filter-based techniques.

5.4.2.3 Reanalyses of Previously-Reported Particulate Matter Exposure Data

Several papers that have been published recently that reanalyzed and interpreted the data collected in previous PM exposure studies are summarized in Table 5-6. These reanalyses are directed towards understanding the personal cloud, the variability in total PM exposure, and the personal exposure-to-ambient concentration relationships for PM. Results of these reanalyses are summarized in Table 5-6 and given in more detail in Section 5.4.3. Brown and Paxton (1998) determined that the high variability in personal exposure to PM makes the personal-to-ambient PM relationship difficult to predict. Wallace (2000b) used data from a number of studies to test two hypotheses: elderly COPD patients have (1) smaller personal clouds and (2) higher correlations between personal exposure and ambient concentrations, compared to healthy elderly, children, and the general population. The analysis by Wallace (2000a) and three subsequent longitudinal studies (Williams 2000a,b,c; Ebelt et al., 2000; Sarnat et al., 2000) support hypothesis 1 but not hypothesis 2. Özkaynak and Spengler (1996) show that at least 50% of personal PM_{10} exposure for the general population is because of ambient particles that significantly contribute to inhaled particles. Wilson and Suh (1997) conclude that fine and coarse particles should be treated as separate classes of pollutants because of differences in characteristics and potential health effects. Wilson et al. (2000) give a review of what they call the “exposure paradox” and determine that personal PM needs to be divided into different classes according to source type, and that correlations between personal and ambient PM will be higher when nonambient sources of PM are removed from the personal PM concentration. Mage (1998) conducted analysis using the PTEAM data and showed that the average person in PTEAM (Riverside, CA in the fall) was exposed to >75% of ambient $PM_{2.5}$ and >64% of ambient PM_{10}. Mage et al. (1999) use an algorithm to fill in missing data and outliers to analyzed data sets and

TABLE 5-6. PAPERS REPORTING REANALYSES OF PARTICULATE MATTER EXPOSURE STUDIES

Reference	Study Cited	Objectives/Hypotheses	Findings
Wallace (2000a)	PTEAM (Özkaynak et al., 1990; Spengler et al., 1989; Wiener 1988, 1989; Wiener et al., 1990) THEES (Lioy et al., 1990) Nashville COPD (Bahadori et al., 2001) Amsterdam COPD (Janssen et al., 1997, 1998a) Boston COPD (Rojas-Bracho et al., 2000)	Examines the differences between pooled and longitudinal correlations in personal and ambient (or outdoor) data for $PM_{2.5}$ and PM_{10}. Discusses the personal cloud for $PM_{2.5}$ and PM_{10}. Hypothesizes that COPD patients have (1) smaller personal clouds (supported) and (2) higher correlations of personal exposure with outdoor concentrations because of reduced mobility (not supported).	Median longitudinal correlation coefficient is much higher than the pooled correlation coefficient for the same data sets. Personal cloud for PM_{10}: 3-67 $\mu g/m^3$; $PM_{2.5}$ 6-27 $\mu g/m^3$. Personal cloud for elderly COPD was much smaller (PM_{10}: 6-11 $\mu g/m^3$; $PM_{2.5} \approx 6$ g/m^3) than for other healthy populations (PM_{10}: 27-56 $\mu g/m^3$; $PM_{2.5}$: 11-27 $\mu g/m^3$) of elderly, children, and the general population. However, higher correlations of personal exposure with ambient concentrations were not observed.
Özkaynak and Spengler (1996)	Dockery and Spengler, 1981 PTEAM (Özkaynak et al., 1996a,b) Netherlands (Janssen et al., 1995)	Uses statistical modeling techniques to examine the relationship between ambient PM concentrations and personal exposures. Data analysis involves use of air exchange rates, penetration factors, and indoor/outdoor ratios, as well as examining exposure in various microenvironments (traveling, working, outdoors, indoors) activities (exposure to smoke, cooking), and source strengths.	The important components of personal exposures are received during contact with indoor sources, mainly in homes and work places. Ambient aerosols contribute about 50% or more to the personal PM_{10} exposures of the general population. The contribution of ambient aerosols to the total toxicity of inhaled particles is significant.
Brown and Paxton (1998)	THEES (Lioy et al., 1990) PTEAM pilot (Wallace, 1996) Boston and Nashville COPD (Rojas-Bracho et al., 2000); Bahadori et al., 2001)	Cross-sectional and longitudinal regression analysis on data sets.	Individual personal PM exposure is subject to high variability, which makes the personal-to-ambient PM relationship difficult to predict.
Wilson and Suh (1997)	Philadelphia (Burton et al., 1996; Suggs and Burton, 1983) EPA AIRS database	Determines the utility of fine and coarse PM concentrations as indicators of time-series epidemiology with regard to day-to-day variability, area uniformity, and indoor/outdoor PM ratios. Necessary to treat personal exposure to ambient PM and personal exposure to nonambient PM as separate components of total personal PM exposure.	Fine and coarse particles should be considered separate classes of pollutants. Fixed-site ambient fine-particle measurements likely give a reasonable indication of the variability in the concentration of ambient fine particles across the community. Coarse-particle measurements most likely will not.
Wilson et al. (2000)	New Jersey (Lioy et al., 1990) Japan (Tamura et al., 1996a) PTEAM (Clayton et al., 1993; Özkaynak et al., 1996a,b) Netherlands (Janssen, 1998a; Suh et al., 1992)	Synoptic review of the "exposure paradox": low correlations between personal exposure and ambient PM concentrations in spite of the existence of statistical association between ambient PM and epidemiological health effects. Uses personal exposure equation, mass balance, regression analysis, and deductive logic.	Personal PM exposure needs to be divided into different classes according to source type: exposure to ambient PM (outdoor and indoors) and exposure to nonambient PM (indoor source and personal activity). Correlations are higher between personal exposure and ambient PM concentrations when PM exposures from nonambient sources are removed.
Mage et al. (1999)	Japan (Tamura et al., 1996a) State College (Suh et al., 1995) Netherlands (Janssen et al., 1997, 1998a, 1999a) New Jersey (Lioy et al., 1990) PTEAM (Clayton et al., 1993; Özkaynak et al., 1996a,b)	Examines the influence of nonambient PM on total PM concentrations and how it may confound the outdoor/personal PM relationship. Missing data and outlier values created using an algorithm. Linear regression analysis of subsequent data sets.	Variation in daily personal exposure for subjects with similar lifestyles and no ETS exposure are driven by variations in ambient PM concentrations. Exposure to ambient PM is highly correlated in time with ambient PM concentrations measured at a community site. Indoor PM does not confound the relationship between daily mortality and ambient PM.
Mage (1998)	PTEAM (Clayton et al., 1993; Özkaynak et al., 1993, 1996a,b)	Uses a reduced-form mass-balance model to predict the average fraction of ambient PM the average person is exposed to.	On average, a person is exposed to >75% of ambient $PM_{2.5}$ and >64% of ambient PM_{10} measured by the community monitor.

DRAFT-DO NOT QUOTE OR CITE

TABLE 5-6 (cont'd). PAPERS REPORTING REANALYSES OF PARTICULATE MATTER EXPOSURE STUDIES

Reference	Study Cited	Objectives/Hypotheses	Findings
Monn (2001)	Multiple Literature Review	To make an objective review of literature published since 1996 as an implicit update to the 1996 USEPA PM AQCD. Emphasis on European studies.	"It is important to note that a personal measurement does not *a priori* provide more valid data than a stationary ambient measurement, i.e. a personal sample in a study investigating effects from outdoor combustion particles is often influenced by sources other than outdoor sources and may thus confound the exposure-effect outcome." "Despite some lack of correlation between personal (PM10) and outdoor values, outdoor fine particle concentrations were strongly associated with mortality and morbidity indicating that outdoor sources (e.g. vehicular emissions) emit the toxic entity (Dockerey et al., 1993; Schwartz et al., 1996)."
Rotko et al. (2000a)	Jantunen et al. (1998) Carrer et al. (1997) Koistinen et al. (1999)	To make a comparison of exposure relationships between the six EXPOLIS European cities (Athens, Basel, Grenoble, Helsinki, Milan, Prague).	Demographic Bias exists because women and more-educated individuals are more likely to respond to survey. Socioeconomic bias exists in low SES subjects less likely to participate in diary keeping and exposure monitoring. Weighting is required for inter-city comparisons. Selection bias is not a problem for characterizing physical factors influencing personal exposure
Rotko et al. (2000b)	Rotko et al. (2000a), Jantunen et al. (1998)	To determine sociodemographic influences of exposure in Helsinki	Distinct male vs female differences: Males had higher exposures to PM2.5, related to ETS, and a larger variance between sociodemographic groupings. No sociodemographic differences existed in outdoor PM2.5 concentrations. Lower occupational status contributed to greater P{M2.5 exposures than higher (professional) occupational status.

DRAFT-DO NOT QUOTE OR CITE

show that variation in daily personal exposures for subjects with similar activity patterns and no ETS exposure are driven by variation in ambient PM concentrations.

5.4.3 Factors Influencing and Key Findings on Particulate Matter Exposures

5.4.3.1 Relationship of Personal/Microenvironmental Particulate Matter with Ambient Particulate Matter

Understanding the relationship between ambient site measurements and personal exposure to PM is important for several reasons. First, it allows us to determine when and for which PM constituents it is appropriate to use ambient measurements as surrogates for exposure in epidemiological studies. Second, it provides information that may improve surrogate exposure measurements and, hence, increase the power of epidemiological studies. Finally, since compliance with the NAAQS is based on ambient monitoring, it can be used to understand the impact of regulation on exposures to PM and its constituents and, hence, can help link the impact of regulations to health outcomes. Many of the studies, summarized above in Table 5-4, have analyzed this relationship using measurements of personal PM exposures and ambient PM concentrations. Of primary interest are the PM concentrations measured in ambient, indoor, outdoor, and personal exposure samples, the statistical correlations between measurements, and the attenuation and/or infiltration factors developed for personal exposure and indoor microenvironments. Attenuation and infiltration factors are discussed in Section 5.3.2.3.1. Information on correlation analysis are provided below.

5.4.3.1.1 Types of Correlations

The three types of correlation data that will be discussed in this section are longitudinal, “pooled”, and daily-average correlations. *Longitudinal correlations* are calculated when data from a study includes measurements over multiple days for each subject (longitudinal study design). Longitudinal correlations describe the temporal relationship between daily personal PM exposure or microenvironment concentration and daily ambient PM concentration for each individual subject. The longitudinal correlation coefficient, r, may differ for each subject, and an analysis of the variability in r across subjects can be performed with this type of data. Typically, the median r is reported along with the range across subjects in the study. There are two types of cross-sectional correlations: pooled and daily average. *Pooled correlations* are calculated when

a study involves one or only a few measurements per subject and different subjects are studied on subsequent days. Pooled correlations combine individual subject/individual day data for the correlation calculation. Pooled correlations describe the relationship between daily personal PM exposure and daily ambient PM concentration across all subjects in the study. For some studies, the multiple days of measurements for each subject were assumed to be independent (after autocorrelation and sensitivity analysis) and combined together in the correlation calculation (Ebelt et al., 2000). *Daily-average correlations* are calculated by averaging exposure across subjects for each day. Daily-average correlations then describe the relationship between the daily average exposure and daily ambient PM concentration.

Pooled correlations have been simulated from longitudinal data by using a random-sampling procedure to select a random day from each subject's measurements for use in the correlation. This procedure was repeated many times, and statistics such as the mean and standard deviation of the pooled correlation coefficient were reported (Janssen et al., 1997, 1998a, 1999c).

The type of correlation analysis can have a substantial effect on the resulting correlation coefficient. Mage et al. (1999) mathematically demonstrated that very low correlations between personal exposure and ambient concentrations could be obtained when people with very different nonambient exposures are pooled, even though their individual longitudinal correlations are high. The longitudinal studies conducted by Tamura et al. (1996a) and Janssen et al. (1997, 1998a, 1999c) determined that the longitudinal correlations between personal exposure and ambient PM concentrations were higher than the correlations obtained from a pooled data set. Wallace (2000a) reviewed a number of longitudinal studies and found that the median longitudinal correlation coefficient was higher than the pooled correlation coefficient for the same data (see Tables 1 and 2, Wallace, 2000a).

Mage et al. (1999) examined three longitudinal exposure data sets where several subjects were measured each day. They showed that by averaging daily exposures across subjects, daily-average correlations could be obtained. These were all higher than the median longitudinal correlations. Williams et al. (2000a,b) and Evans et al. (2000) have also reported higher correlation coefficients for daily-average correlations compared to longitudinal correlations. The higher correlations found between daily-average personal exposures and ambient PM concentrations, as opposed to lower correlations found between individual exposures and

ambient PM levels, recently have been attributed to the statistical process of averaging (Ott et al., 2000). Personal exposures include contributions from nonambient as well as ambient PM concentrations. When several subjects are measured on the same day the mean variability due to variations in nonambient exposures are reduced due to averaging. Therefore, the correlation between personal exposure and ambient concentrations increases as the number of subjects measured daily increases. Ott et al. (2000), using the theory on which their Random Component Superposition (RCS) model is based, predict expected correlations above 0.9 for the PTEAM study and above 0.70 for the New Jersey study (Lioy et al., 1990) if 25 subjects had been measured daily in each study.

5.4.3.1.2 Correlation Data from Personal Exposure Studies

Measurement data and correlation coefficients for the personal exposure studies described in Section 5.4.2.1 are summarized in Table 5-7. All data are based on mass measurements. The studies are grouped by the type of study design, longitudinal or pooled. For each study in Table 5-7, summary statistics for the total personal PM exposure measurements are presented, as well as statistics for residential indoor, residential outdoor, and ambient PM concentrations, where available. The correlation coefficient (r) between total personal PM exposures and ambient PM concentrations also are presented and classified as longitudinal or pooled correlations. When reported, *p*-values for the correlation coefficients are included. Correlation coefficients between personal, indoor, outdoor, and ambient also are reported, when available.

5.4.3.1.3 Correlations Between Personal Exposures, Indoor, Outdoor, and Ambient Measurements

Longitudinal and pooled correlations between personal exposure and ambient or outdoor PM concentrations varied considerably between study and study subjects. Most studies report longitudinal correlation coefficients that range from <0 to ≈1, indicating that an individual's activities and residence type may have a significant effect on total personal exposure to PM. General population studies tend to show lower correlations because of the higher variation in the levels of PM generating activities. In contrast, the absence of indoor sources for the populations in several of the longitudinal studies resulted in high correlations between personal exposure and ambient PM within subjects over time for these populations. But even for these studies,

TABLE 5-7. PERSONAL MONITORING STUDIES FOR PARTICULATE MATTER: MEASURED CONCENTRATIONS AND CORRELATION COEFFICIENTS

		Measured Concentration Levels ($\mu g/m^3$)						Personal-Ambient [2] Correlation Coefficients (r)			Other Correlation Coefficients (r)	
Size Fraction	Avg. Time	Statistic	Sample Size[1]	Personal	Residential Indoor	Residential Outdoor	Ambient	Type[3]	Value (Range)	Notes	Type[3]	Value (Range)
Longitudinal Studies												
Ebelt et al. (2000) – Vancouver, BC												
$PM_{2.5}$	24 h	$\bar{x}$ ± SD Range	106	18.2 ± 14.6 2- 91			11.4 ± 4.1 4 - 29	Median L P	0.48 (-0.68-0.83) 0.15	n = 16 COPD subjects		
Evans et al. (2000) – Fresno, CA												
$PM_{2.5}$	24 h	$\bar{x}$ Range	24	13.3 1 - 24	9.7 4 - 17	20.5 4 - 52	21.7 6 - 37	P	0.41[4]	Fresno-1 study	P_{p-i} P_{p-o}	0.81[4] 0.80[4]
$PM_{2.5}$	24 h	$\bar{x}$ Range	12	11.1 7 - 16	8.0 4 - 12	10.1 5 - 20	8.6 4 - 16	P	0.84[4]	Fresno-2 study	P_{p-i} P_{p-o}	0.95[4] 0.80[4]
Janssen et al. (1997) – Netherlands												
PM_{10}	24 h	$\bar{x}$ ± SD Range	301	105.2 ± 28.7 57 - 195			38.5 ± 5.6 25 - 56	Median L Median L Median L Mean P Mean P Mean P	0.63 (0.1-0.9) 0.63 0.59 0.28 (0.12)[5] 0.45 (0.16)[5] 0.20 (0.19)[5]	n = 45 school children With nonsmoking parents With smoking parents All With nonsmoking parents With smoking parents		
Janssen et al. (1998a) – Netherlands												
PM_{10}	24 h	$\bar{x}$ ± SD Range	262	61.7 ± 18.3 38 - 113	35.0 ± 9.4 19 - 65		41.5 ± 4.3 32 - 50	Median L Median P Median P	0.50 (-0.41-0.92) 0.50 (0.07-0.83)[5] 0.34 (-0.09-0.67)[5]	n = 37 adults No ETS exposure All	Med. L_{p-i} Med. L_{i-a}	0.72 (-0.10-0.98) 0.73 (-0.88-0.95)
Janssen et al. (1999c) – Netherlands												
$PM_{2.5}$	24 h	$\bar{x}$ ± SD Range	77	28.3 ± 11.3 19 - 60			17.1 ± 2.8 14 - 22	Median L Median P	0.86 (-0.11-0.99) 0.41 (-0.28-0.93)[5]	n = 13 school children		
$PM_{2.5}$	24 h	$\bar{x}$ ± SD Range	55	24.4 ± 4.9 19 - 33			17.1 ± 2.6 15 - 22	Median L Median P	0.92 0.82[5]	With nonsmoking parents		
$PM_{2.5}$	24 h	$\bar{x}$ ± SD Range	22	37.0 ± 17.4 21 - 60			17.1 ± 3.7 14 - 21			With smoking parents		

DRAFT-DO NOT QUOTE OR CITE

TABLE 5-7 (cont.). PERSONAL MONITORING STUDIES FOR PM: MEASURED CONCENTRATIONS AND CORRELATION COEFFICIENTS

		Measured Concentration Levels ($\mu g/m^3$)						Personal-Ambient [2] Correlation Coefficients (r)			Other Correlation Coefficients (r)	
Size Fraction	Avg. Time	Statistic	Sample Size[1]	Personal	Residential Indoor	Residential Outdoor	Ambient	Type[3]	Value (Range)	Notes	Type[3]	Value (Range)
Janssen et al. (2000) – Netherlands												
$PM_{2.5}$	24 h	$\bar{x}$ ± SD Range	338	24.3 ± 25.7 9 - 134	28.6 ± 41.8 9 - 239		20.6 ± 4.0 13 - 31	Median L Median L	0.79 (-0.41-0.98) 0.85	n = 36 elderly w/CV disease No ETS exposures	Med. L_{p-i} Med. L_{i-a}	0.91 (-0.28-1.0) 0.84(-0.00-0.98)
Janssen et al. (2000) – Finland												
$PM_{2.5}$	24 h	$\bar{x}$ ± SD Range	336	10.8 ± 4.4 4 - 33	11.0 ± 4.0 3 - 27		12.6 ± 2.0 10 - 18	Median L	0.76 (-0.12-0.97)	n = 46 elderly w/CV disease	Med. L_{p-i} Med. L_{i-a}	0.89 (0.14-1.0) 0.70 (-0.15-0.94)
Linn et al. (1999) – Los Angeles												
$PM_{2.5}$	24 h	$\bar{x}$ ± SD Range	60	23.8 ± 15.1 4 - 65	23.5 ± 15.3 4 - 92	24.8 ± 14.5 4 - 63		P	0.26[6]		P_{i-a} P_{o-a}	0.26[6] 0.47[6]
PM_{10}	24 h	$\bar{x}$ ± SD Range	59	34.8 ± 14.8 5 - 85	32.6 ± 15.6 9 - 105	39.8 ± 18.3 7 - 97	33 ± 15 9 - ??	P	0.22[6]		P_{i-a} P_{o-a}	0.32[6] 0.66[6]
Rojas-Bracho et al. (2000) – Boston												
$PM_{2.5}$	12 h	$\bar{x}$ ± SD Range	224	21.6 ± 13.6 1 - 128	17.5±14.1 2 - 73	14.2 ± 11.2 1 - 57		Median L	0.61 (0.10-0.93)[6]	n = 17 adults	Med. L_{p-i} Med. L_{i-o}	0.87[6] 0.74[6]
PM_{10}	12 h	$\bar{x}$ ± SD Range	225	37.2 ± 22.8 9 - 211	31.9 ± 25.2 2 - 329	22.2 ± 18.7 3 - 76		Median L	0.35 (0.0-0.72)[6]		Med. L_{p-i} Med. L_{i-o}	0.71[6] 0.50[6]
$PM_{10-2.5}$	12 h	$\bar{x}$ ± SD Range	222	15.6 ± 14.6 -11 - 103	14.5 ± 9.2 -3 - 255	8.1 ± 6.8 -2 - 64		Median L	0.30 (0.0-0.97)[6]		Med. L_{p-i} Med. L_{i-o}	0.42[6] 0.20[6]
Sarnat et al. (2000) – Baltimore												
$PM_{2.5}$	24 h	$\bar{x}$ ± SD $\bar{x}$ ± SD	37 36	26.7 ± 13.7 18.5 ± 11.2			25.2 ± 11.5 5.6 ± 49.0	Median L Median L P P P P	0.76 (-0.21-0.95)[7] 0.25 (-0.38-0.81)[7] 0.89[8] 0.75[8] 0.50[8] 0.44[8]	n = 15 adults; summer n = 15 adults; winter High ventilation; summer Med. ventilation; summer Low ventilation; summer WINTER		
PM_{10}	24 h	$\bar{x}$ ± SD $\bar{x}$ ± SD	37 36	33.9 ± 11.7 28.0 ± 16.5			34.0 ± 12.8 7.5 ± 73.2	Median L Median L	0.64 (0.08-0.86)[7] 0.53 (-0.79-0.89)[7]	SUMMER WINTER		

DRAFT-DO NOT QUOTE OR CITE

TABLE 5-7 (cont.). PERSONAL MONITORING STUDIES FOR PM: MEASURED CONCENTRATIONS AND CORRELATION COEFFICIENTS

		Measured Concentration Levels ($\mu g/m^3$)						Personal-Ambient [2] Correlation Coefficients (r)			Other Correlation Coefficients (r)	
Size Fraction	Avg. Time	Statistic	Sample Size[1]	Personal	Residential Indoor	Residential Outdoor	Ambient	Type[3]	Value (Range)	Notes	Type[3]	Value (Range)
Sarnat et al. (2000) – Baltimore (cont'd)												
$PM_{10-2.5}$	24 h	$\bar{x} \pm SD$ $\bar{x} \pm SD$	37 36	7.2 ± 4.0 9.6 ± 7.9			8.4 ± 2.3 -1.3 ± 24.2	Median L Median L	0.11 (-0.60-0.64)[7] 0.32 (-0.48-0.68)[7]	SUMMER WINTER		
Tamura et al. (1996a) – Tokyo												
PM_{10}	48 h							P	0.83	n = 7 elderly adults		
Williams et al. (2000a,b) – Baltimore												
$PM_{2.5}$	24 h	$\bar{x}$ Range	23	13.0 7 - 25	9.4 4 - 19	22.0 7 - 52	22.0 8 - 59	Median L P	0.80 (0.38-0.98)[6] 0.89[4]	n = 21 elderly adults	P_{p-i} P_{p-o} P_{i-o} P_{i-a} P_{o-a}	0.90[4] 0.95[4] 0.94[4] 0.87[4] 0.96[4]
PM_{10}	24 h	$\bar{x}$ Range	28		11.0 4 - 23	30.0 13 - 66	29.9 13 - 74				P_{i-o} P_{i-a} P_{o-a}	0.82[4] 0.81[4] 0.94[4]
$PM_{10-2.5}$	24 h	$\bar{x}$ Range	26		1.0 -3 - 5	8.0 -2 - 16	8.0 1 - 15				P_{i-o} P_{i-a} P_{o-a}	0.18[4] 0.08[4] 0.45[4]
Pooled Studies												
Bahadori (1998) – Nashville												
$PM_{2.5}$	12 h	$\bar{x} \pm SD$ Range	30	21.7 ± 10.5 10 - 67	15.5 ± 6.6 5 - 40	23.4 ± 6.8 3 - 61		P	0.09	n = 10 COPD subjects; daytime	P_{p-i} P_{i-o}	0.72 0.31
PM_{10}	12 h	$\bar{x} \pm SD$ Range	30	33.0 ± 16.9 5 - 88	21.6 ± 10.7 9 - 77	32.5 ± 8.1 7 - 76		P	-0.08	n = 10 COPD subjects; daytime	P_{p-i} P_{i-o}	0.43 0.06
Pellizzari et al. (1999) – Toronto												
$PM_{2.5}$	3 d	$\bar{x}$	922	28.4	21.1	15.1		P	0.23	n = 178; n for indoor, outdoor lower than personal	P_{p-i} P_{i-o}	0.79 0.33
PM_{10}	3 d	$\bar{x}$	141	67.9	29.8	24.3				No correlations reported		

DRAFT-DO NOT QUOTE OR CITE

TABLE 5-7 (cont.). PERSONAL MONITORING STUDIES FOR PM: MEASURED CONCENTRATIONS AND CORRELATION COEFFICIENTS

				Measured Concentration Levels ($\mu g/m^3$)				Personal-Ambient [2] Correlation Coefficients (r)			Other Correlation Coefficients (r)	
Size Fraction	Avg. Time	Statistic	Sample Size[1]	Personal	Residential Indoor	Residential Outdoor	Ambient	Type[3]	Value (Range)	Notes	Type[3]	Value (Range)
Oglesby et al. (2000) – EXPOLIS Basel												
$PM_{2.5}$	48 h	$\bar{x}$ ± SD	44	23.7 ± 17.1		19.0 ± 11.7		P	0.07	All		
			20	17.5 ± 13.0		17.7 ± 7.1		P	0.21	No ETS exposure		
Santos-Burgoa et al. (1998) – Mexico City												
PM_{10}	24 h	$\bar{x}$ ± SD	66	97 ± 44	99 ± 50			P	0.26		P_{p-i} P_{i-a}	0.47 0.23
Tamura et al. (1996b) – Osaka												
PM_2	48 h							P	0.74			
PM_{10}	48 h							P	0.67			
Pellizzari et al., 2001 – Indianapolis												
$PM_{2.5}$	72-h	Median	250	23	18	18	18	P	0.102	Between the Logarithms of concentrations	P vs outdoor P vs Indoor	0.138 0.923
Brauer et al.,2000 – Banska BystricaSO4												
$PM_{2.5}$ PM_{10}	24-h	Mean	PM_{10} Summer	122	79		35	P PM_{10}	r^2<0.17	Multivar. With nicotine	P indoor PM2.5	r^2=0.15 r^2=0.23
			PM_{10} Winter	120	66		45				P	
			$PM_{2.5}$ Summer	88	55		22				Personal SO4 vs Amb. SO4	
			$PM_{2.5}$ Winter	69	53		32					
			SO_4 Winter	6.5	4.6		5.7					

Abbreviations used:
Avg. = Averaging (time)
Conc. = Concentration
CV = Cardiovascular
d = Day
ETS = Environmental tobacco smoke
h = Hour
i-a = Indoor-ambient correlation
i-o = Indoor-outdoor correlation
L = Longitudinal correlation
Med. = Median
o-a = Outdoor-ambient correlation
P = Pooled correlation
p-i = Personal-indoor correlation
p-o = Personal-outdoor correlation
SD = Standard deviation
Stat. = Statistic
$\bar{x}$ = Mean

Notes:
1. Sample size is for personal concentrations; indoor, outdoor and ambient sample sizes may differ.
2. Correlation coefficient is for personal-residential outdoor if no ambient concentration data reported.
3. See text for description of types of correlations.
4. Daily-averaged correlation (values for individual subjects averaged for each day).
5. Pooled correlations estimated using a Monte Carlo sampling procedure, n = 1000. If mean P is shown, then SD given; if median P is shown, then range is given.
6. Obtained from a regression equation; $r = \sqrt{(R^2)}$.
6. Spearman rank correlations.
7. Calculated, $r = \sqrt{(R^2)}$, from R^2 from a mixed model regression.

DRAFT-DO NOT QUOTE OR CITE

correlations varied by individual, depending on their activities and the microenvironments that they occupied.

Probability Studies

In the Toronto study (Pellizzari et al., 1999), pooled correlations were derived for personal, indoor, outdoor, and fixed site ambient measurements. This study was conducted in Toronto on a probability sample of 732 participants who represented the general population, 16 years and older. The study included between 185 and 203 monitoring periods with usable PM data for personal, residential indoor, and outdoor measurements. For PM_{10}, measurements, the mean concentrations were 67.9 $\mu g/m^3$ for personal, 29.8 $\mu g/m^3$ for indoor air, and 24.3 $\mu g/m^3$ for outdoor air samples. For $PM_{2.5}$, the mean concentrations were 28.4 $\mu g/m^3$ for personal, 21.1 $\mu g/m^3$ for indoor air, and 15.1 $\mu g/m^3$ for outdoor air samples. A low but significant correlation ($r = 0.23$, $p < 0.01$) was reported between personal exposure and ambient measurements. The correlations between indoor concentrations and the various outdoor measurements of $PM_{2.5}$ ranged from 0.21 to 0.33. The highest correlations were for outdoor measurements at the residences with the ambient measurements made at the roof site (0.88) and the other fixed site (0.82). Pellizzari et al. (1999) state that much of the difference among the data for personal/indoor/outdoor PM

> . . . can be attributed to tobacco smoking, since all variables reflecting smoking . . . were found to be highly correlated with the personal (and indoor) particulate matter levels, relative to other variables that were measured . . . none of the outdoor concentration data types (residential or otherwise) can adequately predict personal exposures to particulate matter. (p. 729)

Using a Random Component Superposition Statistical Model, Ott et al. (2000) calculated an attenuation factor of 0.6144 for personal exposure for PM_{10}. The mean nonambient exposure component for PM_{10} was estimated as 52.62 $\mu g/m^3$ with a standard deviation of 84.82 $\mu g/m^3$. Although the data were available for $PM_{2.5}$, similar calculations were not made.

PM_{10} data from the PTEAM study were analyzed using the same approach (Ott et al., 2000). For PTEAM, an attenuation factor of 0.5546 was calculated for personal exposure. Infiltration factors were calculated for each residence with an average of 0.5594 and a standard

deviation of 0.1476. Values ranged from a minimum of 0.19 to a maximum of 0.87 showing the substantial variability that can be seen between homes depending upon the housing characteristics and operation of the HVAC system. The mean nonambient exposure component for PM_{10} was estimated as 59.23 $\mu g/m^3$ with a standard deviation of 45.85 $\mu g/m^3$.

Santos-Burgoa et al. (1998) describe a 1992 study of personal exposures and indoor concentrations to a randomly sampled population near Mexico City. The sample of 66 monitored subjects included children, students, office and industrial workers, and housewives. None of the people monitored were more than 65 years old. The mean 24-h personal exposure and indoor concentrations were 97 ± 44 (SD) and 99 ± 50 $\mu g/m^3$, respectively, with an $\mathbf{r}_{Personal/Ambient} = 0.26$ ($p = 0.099$). Other correlations of interest were $\mathbf{r}_{Personal/Indoor} = 0.47$ ($p = 0.002$) and $\mathbf{r}_{Indoor/Ambient} = 0.23$ ($p = 0.158$). A strong statistical association was found between personal exposure and socioeconomic class ($p = 0.047$) and a composite index of indoor sources at the home ($p = 0.039$).

Correlation analysis for personal exposure has not yet been reported for EXPOLIS. Some preliminary results (Jantunen et al., 2000) show that, in Basel and Helsinki, a single ambient monitoring station was sufficient to characterize the ambient $PM_{2.5}$ concentration in each city. Using microenvironmental concentration data collected while the subjects were at home, at work, and outdoors, they calculated the sum of the time-weighted-averages of these data and found the results closely match the personal $PM_{2.5}$ exposure data collected by the monitors carried by most of the subjects, with a few subjects (mostly smokers) being noticeable exceptions.

Longitudinal Studies

A number of longitudinal studies using a purposeful sampling design have been conducted and reported in the literature since 1996. A number of these studies (Janssen et al., 1998a, 1999b, 2000; Williams et al., 2000b; Evans et al., 2000) support the previous work by Janssen et al. (1995) and Tamura et al. (1996a) and demonstrate that, for individuals with little exposure to nonambient sources of PM, correlations between total PM exposure and ambient PM measurements are high. Other studies (Ebelt et al., 2000; Sarnat et al., 2000) show strong correlations for the SO_4^{-2} component of $PM_{2.5}$ but poorer correlations for $PM_{2.5}$ mass. Still other studies show only weak correlations (Rojas-Bracho et al., 2000; Linn et al., 1999; Bahadori et al., 2001). Even when strong longitudinal correlations are demonstrated for individuals in a study,

the variety of living conditions may lead to variations in attenuating factors or the fraction of ambient PM contributing to personal exposure. Groups with similar living conditions, especially if measurements are conducted during one season, may have similar α and, therefore, very high correlations between personal exposure and ambient concentrations, even for pooled correlations. However, when studies contain subjects with homes of very different ventilation characteristics or cover more than one season, variations in α can be high across subjects, thus, showing poor pooled correlations even in the absence of indoor sources.

Elderly Subjects. Janssen et al. (2000) continued their longitudinal studies with measurements of personal, indoor, and outdoor concentrations of $PM_{2.5}$ for elderly subjects with doctor-diagnosed angina pectoris or coronary heart disease. Studies were conducted in Amsterdam and Helsinki, Finland, in the winter and spring of 1998 and 1999. In the Amsterdam study, with 338 to 417 observations, the mean $PM_{2.5}$ concentrations were 24.3, 28.6, and 20.6 $\mu g/m^3$ for personal, indoor, and outdoor samples, respectively. If the measurements with ETS in the home were excluded, the mean indoor concentration dropped to 16 $\mu g/m^3$, which was lower than outdoors. In the Helsinki study, the mean $PM_{2.5}$ concentrations were 10.8 $\mu g/m^3$ for personal, 11.0 $\mu g/m^3$ for indoor air, and 12.6 $\mu g/m^3$ outdoor air samples. The authors note that for this group of subjects, personal exposure, indoor concentrations, and ambient concentrations of $PM_{2.5}$ were highly correlated within subjects over time. Median Pearson's correlation coefficients between personal exposure and outdoor concentrations were 0.79 in Amsterdam and 0.76 in Helsinki. The median Pearson's r for the indoor/outdoor relationship was 0.85 for the Amsterdam study, excluding homes with ETS. The correlation for indoors versus outdoors was 0.70 for all homes.

Results from the correlation analysis can be used to estimate infiltration factors and penetration factors for these two groups of subjects. In Amsterdam, the attenuation factor was 0.43 and the infiltration factor was 0.47. Very similar results were seen in Helsinki for the attenuation factor (0.45) and the infiltration factor (0.51).

A series of PM personal monitoring studies involving elderly subjects was conducted in Baltimore County, MD, and Fresno, CA. The first study was a 17-day pilot (January-February 1997) to investigate daily personal and indoor $PM_{1.5}$ concentrations, and outdoor $PM_{2.5}$ and $PM_{2.5\text{-}10}$ concentrations experienced by nonsmoking elderly residents of a retirement community located near Baltimore (Liao et al., 1999; Williams et al., 2000c). The 26 residents were aged

65 to 89 (mean = 81), and 69% of them reported a medical condition, such as hypertension or coronary heart disease. In addition, they were quite sedentary; less than 5 h day^{-1}, on average, was spent on ambulatory activities. Because most of the residents ate meals in a communal dining area, the average daily cooking time in the individual apartments was only 0.5 h (range 0 to 4.5 h). About 96% of the residents' time was spent indoors (Williams et al., 2000c). Personal monitoring, conducted for five subjects, yielded longitudinal correlation coefficients between ambient concentrations and personal exposure ranging from 0.00 to 0.90.

The Baltimore main study and the Fresno study were conducted using similar monitoring techniques and study design. Concentrations measured in these studies are summarized in Table 5-8. For $PM_{2.5}$, personal exposure and indoor air concentrations are similar for all three studies even though outdoor air concentrations for Fresno in the winter are only half of those measured for Fresno in the Winter and Baltimore in the summer. This result is presumably due to high penetration efficiencies in the spring in Fresno when the weather was warm and participants kept the windows and doors of their homes open. These data also show that even when correlations are high, the use of an ambient monitor as a surrogate for exposure in epidemiological studies can bias the strength of the health effect found, due to differing exposure levels.

TABLE 5-8. MEAN CONCENTRATION FOR PM MASS REPORTED FOR THE BALTIMORE (Williams et al., 2000a,b,c) AND THE FRESNO (Evans et al., 2000) STUDIES

	$PM_{2.5}$ Concentration ($\mu g/m^3$)			PM_{10} Concentration ($\mu g/m^3$)		
Study	**Personal**	**Indoors**	**Outdoors**	**Personal**	**Indoors**	**Outdoors**
Baltimore	13.0 ± 4.2	10.5 ± 4.9	22.0 ± 12.0	—	13.5 ± 6.3	30.0 ± 13.7
Fresno-Winter	13.3 ± 5.9	9.7 ± 5.0	20.5 ± 13.4	—	15.1 ± 4.1	28.2 ± 15.9
Fresno-Spring	11.1 ± 2.8	8.0 ± 1.8	10.1 ± 3.2	37.3	16.7 ± 3.1	28.7 ± 6.6

Calculated Correlation Coefficients are summarized in Tables 5-9 and 5-10. In Table 5-9, results for Baltimore show excellent daily average correlations for both $PM_{2.5}$ and PM_{10}. These

TABLE 5-9. DAILY- AVERAGE CORRELATION COEFFICIENTS REPORTED FOR THE BALTIMORE (WILLIAMS ET AL., 2000a,b,c) AND THE FRESNO (EVANS ET AL., 2000) STUDIES

	$PM_{2.5}$ r^2			PM_{10} r^2
Study	**Ambient/Outdoor**	**Personal/Ambient**	**Personal/Indoors**	**Ambient/Outdoor**
Baltimore	0.92	0.80 (0.14-0.80)[a]	0.98 (0.20-0.99)[a]	0.89
Fresno-Winter	0.48	—	—	0.48
Fresno-Spring	0.53	0.70	0.77	0.61

[a]Range for individual participants.

TABLE 5-10. REGRESSION ANALYSIS REPORTED FOR INDOOR/OUTDOOR RELATIONSHIPS FOR THE BALTIMORE (WILLIAMS ET AL., 2000a,b,c) AND THE FRESNO (EVANS ET AL., 2000) STUDIES

	Daily Average			Individual		
Study	**r^2**	**slope**	**Intercept ($\mu g/m^3$)**	**r^2**	**slope**	**Intercept ($\mu g/m^3$)**
Baltimore	0.92	0.39	1.5	0.73 ± 0.16	0.43 ± 0.15	0.9 ± 2.6
Fresno-Winter	0.86	nr	nr	0.55 ± 0.25	0.25 ± 0.17	4.4 ± 3.2
Fresno-Spring	0.56	nr	nr	0.39 ± 0.21	0.49 ± 0.38	3.0 ± 3.7

results represent primarily the behavior of fine particle regional sulfate for a group of participants who have few indoor or personal sources. However even for this group, there was a wide range of individual correlation coefficients. The Fresno data, on the other hand, shows much poorer daily average correlations. Of special note are the poorer correlation for the ambient to outdoor residential monitor. This could be due to the higher concentrations of nitrate in the samples. In addition, the residential site may have be influenced by highway traffic.

The correlation analysis in Table 5-10 shows correlation coefficients as well as the slope (infiltration factor) and the intercept (indoor concentration due to nonambient sources) for the Baltimore and Fresno studies. These data show strongest correlations for Baltimore, with very

low indoor concentrations from nonambient sources. Correlations for Fresno are not as strong, with higher concentrations from nonambient sources. The infiltration factors for Baltimore and Fresno-Spring time are very similar at approximately 0.5. The infiltration factors for Fresno-Winter are considerably lower.

Subjects with COPD. Linn et al. (1999) describe a 4-day longitudinal assessment of personal $PM_{2.5}$ and PM_{10} exposures (on alternate days) in 30 COPD subjects aged 56 to 83; concurrent indoor and outdoor monitoring were conducted at their residences. This study occurred in the summer and autumn of 1996 in the Los Angeles area. PM_{10} data from the nearest fixed-site monitoring station to each residence also was obtained. Pooled correlations for personal exposure to outdoor measurements were 0.26 and 0.22 for $PM_{2.5.}$ and PM_{10}, respectively. Correlations of day-to-day changes in $PM_{2.5}$ and PM_{10} measured outside the homes and correlated with concurrent PM_{10} measurements at the nearest ambient monitoring location gave R^2 values of 0.22 and 0.44, respectively. Correlations of day to day changes in PM mass measured indoors correlated with outdoor measurements at the homes gave R^2 values of 0.27 and 0.19 for PM_{10} and $PM_{2.5}$, respectively.

Personal, indoor, and outdoor $PM_{2.5,}$ $PM_{10,}$ and $PM_{2.5\text{-}10}$ correlations were reported by Rojas-Bracho et al. (2000) for a study conducted in Boston, MA, on 18 individuals with COPD. Both the mean and median personal exposure concentrations were higher than the indoor concentrations, which were higher than outdoor concentrations for all three PM measurement parameters. Geometric mean indoor/outdoor ratios were 1.4 ± 1.9 for PM_{10}, 1.3 ± 1.8 for $PM_{2.5}$, and 1.5 ± 2.7 for $PM_{2.5\text{-}10}$. Median longitudinal R^2s between personal exposure and ambient PM measurements were 0.12 for PM_{10}, 0.37 for $PM_{2.5}$ and 0.07 for $PM_{2.5\text{-}10}$. The relationship between the indoor and outdoor concentrations was strongest for $PM_{2.5}$, with a median R^2 of 0.55 and 11 homes having significant R^2 values. For $PM_{10,}$ the median R^2 value was 0.25, with significant values for eight homes. Only five homes had significant indoor/outdoor associations for $PM_{2.5\text{-}10}$, with an insignificant median R^2 value of 0.04. The poor correlations for $PM_{10\text{-}2.5}$ are a result of poorer penetration efficiencies, higher decay rates, and spatial inhomogeneities.

Bahadori et al. (2001) report a pilot study of the PM exposure of 10 nonrandomly chosen chronic obstructive pulmonary disease (COPD) patients in Nashville, TN, during the summer of 1995. Each subject alternately carried a personal $PM_{2.5}$ or PM_{10} monitor for a 12-h daytime period (8 a.m. to 8 p.m.) for 6 consecutive days. These same pollutants were monitored

simultaneously indoors and outdoors at their homes. All of the homes were air-conditioned and had low air exchange rates (mean = 0.57 h^{-1}), which may have contributed to the finding that mean indoor $PM_{2.5}$ was 66% of the mean ambient $PM_{2.5}$. This can be contrasted with the PTEAM study in Riverside, CA, where no air conditioners were in use and the mean indoor $PM_{2.5}$ was 98% of the mean ambient $PM_{2.5}$ (Clayton et al., 1993). Data sets were pooled for correlation analysis. Resulting pooled correlations between personal and outdoor concentrations were r = 0.09 for $PM_{2.5}$ and r = −0.08 for PM_{10}.

5.4.3.1.4 Sulfate as a Surrogate for Personal Exposure to Ambient Particulate Matter

A study, conducted in Vancouver, involving sixteen COPD patients aged 54 to 86, reported low median longitudinal (r = 0.48) and pooled (r = 0.15) correlation coefficients between personal exposures and ambient concentrations of $PM_{2.5}$ (Ebelt et al., 2000). However, the mean correlation between personal exposure to sulfate and ambient concentrations of sulfate was much higher (r = 0.96). Since typically there are minimal indoor sources of sulfate, the relationship between ambient concentrations and personal exposures to sulfate would not be weakened by variability in an indoor-generated sulfate component, as for example in the case for $PM_{2.5}$ for which there are many primary indoor sources as well as some secondary indoor sources. Correlations of ambient concentrations vs. personal exposures for $PM_{2.5}$ and sulfate are compared in Figure 5-1.

Another study, conducted in Baltimore, MD, involved 15 nonsmoking adult subjects (>64 years old) who were monitored for 12 days during summer 1998 and winter 1999 (Sarnat et al., 2000). All subjects (nonrandom selection) were retired, physically healthy, and lived in nonsmoking private residences. Each residence, except one, was equipped with central air-conditioning; however, not all residences used air-conditioning throughout the summer. The average age of the subjects was 75 years (±6.8 years). Sarnat et al. (2000) reported higher longitudinal and pooled correlations for $PM_{2.5}$ during summer than winter. Similar to Ebelt et al. (2000), Sarnat et al. (2000) reported stronger associations between personal exposure to SO_4^{2-} and ambient concentrations of SO_4^{2-} than for total personal $PM_{2.5}$ exposure and ambient $PM_{2.5}$ concentrations. The ranges of correlations are shown in Figure 5-1 along with similar data from Ebelt et al. (2000).



Figure 5-1. Comparison of correlation coefficients for longitudinal analyses of personal exposure versus ambient concentrations for individual subjects for $PM_{2.5}$ and sulfate.

The higher correlation coefficients and the narrower range of the correlation coefficient for sulfate suggest that removing indoor-generated and personal activity PM from total personal PM would result in a higher correlation with ambient concentrations. If there are indeed no indoor sources, a personal exposure measurement for sulfate gives the ambient exposure of sulfate; the ratio of personal sulfate to ambient sulfate gives the attenuation coefficient on an individual, daily basis; and the attenuation coefficient times the ambient $PM_{2.5}$ concentration gives the individual, daily values of ambient $PM_{2.5}$ exposures (Wilson et al., 2000).

This technique applies only to the non-volatile components of fine PM, as measured by $PM_{2.5}$. It requires that the sulfate concentration be large enough so that it can be measured with reasonable accuracy. It does not require that sulfate be correlated with $PM_{2.5}$ or the non-sulfate components of $PM_{2.5}$ since the sulfate data is used to estimate the attenuation coefficient, not

$PM_{2.5}$. The technique does require that there be minimal indoor sources of sulfate, as indicated by a near-zero intercept for the regression, and that the size distribution of $PM_{2.5}$ and sulfate be similar.

Sarnat et al. (2001) subsequently extended the Baltimore study to include 20 older adults, 21 children, and 15 individuals with COPD for a total of 56 subjects. In both studies (Sarnat et al., 2000, 2001), they used their personal and ambient sulfate data to estimate the ambient $PM_{2.5}$ exposure. They used this information in mixed model analysis (mixed models account for differences among individual subjects) but did not report correlations between ambient $PM_{2.5}$ exposure and ambient $PM_{2.5}$ concentrations.

However, Sarnat et al. (2001) did report slopes from the mixed model analyses. The t-statistic for the slope of ambient exposure versus ambient concentration as compared to total personal exposure versus ambient concentration increased from 9.96 to 11.12 (total exposure vs. ambient concentration) for the summer period and 4.36 to 19.88 (ambient exposure vs. ambient concentration) for the winter period.

The study conducted by Sarnat et al. (2000) also illustrates the importance of ventilation on personal exposure to PM. During the summer, subjects recorded the ventilation status of every visited indoor location (e.g., windows open, air-conditioning use). As a surrogate for the air-exchange rate, personal exposures were classified by the fraction of time the windows were open while a subject was in an indoor environment (F_v). Sarnat et al. (2000) report regression analyses for personal exposure on ambient concentration for total $PM_{2.5}$ and for sulfate for each of the three ventilation conditions. Personal exposure to sulfate may be taken as a surrogate for personal exposure to ambient accumulation-mode PM in the absence of indoor sulfate sources. Figure 5-2 shows a comparison of the regressions and indicates how the use of a sulfate tracer as a surrogate for PM of ambient origin improves the correlation coefficient. The improvement is especially pronounced for the lowest ventilation conditions. For the lowest ventilation condition, R^2 improves from 0.25 to 0.72.

5.4.3.1.5 Personal Exposure to Ambient and Nonambient Particulate Matter

The utility of treating personal exposure to ambient PM, E_{ag}, and personal exposure to nonambient PM, E_{nonag}, as separate and distinct components of total personal exposure to PM, E_t, was pointed out by Wilson and Suh (1997). The PTEAM study measured, in addition to indoor,



Figure 5-2. Personal exposure versus ambient concentrations for $PM_{2.5}$ and sulfate. (Slope estimated from mixed models).

Source: Sarnat et al. (2000).

outdoor, and personal PM, the air exchange rate for each home and collected information on the time spent in various indoor and outdoor μ**e**. This information is available for 147, 12-h daytime periods. With this information, it is possible to estimate the daytime E_{ag} and E_{nonag} as described in Section 5.3.2.3.1. Various examples of this information have been reported (Mage et al., 1999; Wilson et al., 2000). Graphs showing the relationships between ambient concentration and the various components of personal exposure (E_t, E_{ag}, and E_{nonag}) are shown in Figure 5-3. The correlation coefficient for the pooled data set improves from r = 0.377 for E_t versus C_a (Figure 5-3a) to r = 0.856 for E_{ag} versus C_a (Figure 5-3b) because of the removal of the E_{nonag} , which, as shown in Figure 5-3c, is highly variable and independent of C_a. The correlation between E_{ag} and C_a is less than 1 because of the day-to-day variation in α_{it}. The regression analysis with E_t total PM gives $\overline{\alpha}$ = 0.711 and $\overline{N}$ = 81.6 μg/m^3. The regression analysis with E_{ag} gives $\overline{\alpha}$ = 0.625. The regression with E_{nonag} gives $\overline{N}$ = 79.2 μg/m^3. The finite intercept in the regression with E_{ag} must be attributed to bias or error in some of the measurements. No studies, other than PTEAM, have provided the quantity of data on E_t , C_a , C_i, and *a* required to conduct an analysis comparable to that shown in Figure 5-3. It should be noted that the PTEAM study was conducted in southern California in the fall, when house were open and air exchange rates were high and relatively uniform. These are best case conditions for showing high correlations between ambient site measurements and personal correlations.

The RCS model introduced by Ott et al.(2000) presents a modeling framework to determine the contribution of ambient PM_{10} and indoor-generated PM_{10} on personal exposures in large urban metropolitan areas. The model has been tested using personal, indoor and outdoor PM_{10} data from three urban areas (Riverside, CA; Toronto; and Phillipsburg, NJ). Results suggest that it is possible to separate the ambient and nonambient PM contributions to personal exposures on a community-wide basis. However, as discussed in the paper, the authors make some assumptions that require individual consideration in each-city specific application of the model for exposure or health effects investigations. Primarily, housing factors, air-conditioning, seasonal differences, and complexities in time-activity profiles specific to the cohort being studied have to be taken into account prior to adopting the model to a given situation. Finally, this and other available exposure-based analyses presented here does not yet predict the relative contribution of indoor and outdoor PM to particle mass burden to the lung as a function of



Figure 5-3. Regression analyses of aspects of daytime personal exposure to PM_{10} estimated using data from the PTEAM study. (a) Total personal exposure to PM, E_t, regressed on ambient concentration, C_a. (b) Personal exposure to ambient PM, E_{ag} regressed on C_a. (c) Personal exposure to nonambient PM, E_{nonag} regressed on C_a.

Source: Data taken from Clayton et al. (1993).

human activities and different microenvironmental sources and concentrations of PM and its copollutants.

5.4.3.2 Factors That Affect Relationship between Personal Exposure and Ambient PM

A number of factors will affect the relationship between personal exposure and PM measured at ambient-site community monitors. Spatial variability in outdoor microenvironments and penetration into indoor microenvironments will influence the relationship for ambient-generated PM, air-exchange rates, and decay rates in indoor microenvironments will influence the relationship for both ambient-generated and total PM, whereas personal activities will influence the relationship for total PM but not ambient-generated PM. Information on these effects is presented in detail in the following section.

5.4.3.2.1 Spatial Variability and Correlations Over Time

Chapter 3 (Section 3.2.3) presents information on the spatial variability of PM mass and chemical components at fixed-site ambient monitors; for purposes of this chapter, this spatial variability is called an "ambient gradient". The data presented in Section 3.2.3 indicate that ambient gradients of PM and its constituents exist in urban areas to a greater or lesser degree. This gradient, and any that may exist between a fixed-site monitor and the outdoor μ**e** near where people live, work, and play, obviously affects the exposure. The purpose of this section is to review the available data on ambient monitor-to-outdoor microenvironmental concentration gradients, or relationships, that have been measured by researchers since 1996. These analyses below are, in general, consistent with the previous studies covered in the 1996 PM AQCD. A few outdoor-to-outdoor monitoring studies also are included to highlight relationships among important μ**e** categories. To assess spatial variability or gradients, the spatial correlations in the data are usually analyzed. However, it should be noted that high temporal correlation between two monitoring locations does not imply low spatial variability or low ambient gradients. High temporal correlation between two sites indicates that changes in concentrations at one site can be estimated from data at another site. Results presented below are consistent with previous studies assessed in the 1996 PM AQCD.

Oglesby et al. (2000), in a paper on the EXPOLIS-EAS study, conclude that very little spatial variability exists in Basel, Switzerland, between PM levels measured at fixed site

monitors and the participant's outdoor μ**e**. The authors report a high correlation between home outdoor $PM_{2.5}$ levels (48-h measurements beginning and ending at 8:00 a.m.) and the corresponding 24-h average PM_4 (time-weighted values calculated from midnight to midnight) measured at a fixed monitoring station ($n = 38$, $r_{sp} = 0.96$, $p < 0.001$). They considered each home outdoor monitor as a temporary fixed monitor and concluded that "the $PM_{2.5}$ level measured at home outdoors . . . represents the fine particle level prevailing in the city of Basel during the 48-h measuring period . . . ."

In a study conducted in Helsinki, Finland, Buzorius et al. (1999) conclude that a single monitor may be used to adequately describe the temporal variations in concentration across the metropolitan area. Particle size distributions were measured using a differential mobility particle sizer (DMPS; Wintlmayer) coupled with a condensation particle counter (CPC TSI 3010, 3022) at four locations including the official air monitoring station, which represented a "background" site. The monitoring period varied between 2 weeks and 6 mo for the sites and data were reported for 10-min and 1-, 8-, and 24-h averages. As expected, temporal variation decreased as the averaging time increased. The authors report that particle number concentration varied in magnitude with local traffic intensity. Linear correlation coefficients computed for all possible site-pairs and averaging times showed that the correlation coefficient improved with increasing averaging time. Using wind speed and direction vectors, lagged correlations were calculated and were generally higher than the "raw" data correlations. Weekday correlations were higher than weekend correlations as "traffic provides relatively uniform spatial distribution of particulate matter" (p. 565). The authors conclude that, even for time periods of 10 min and 1 h, sampling at one station can describe temporal variations across relatively large areas of the city with a correlation coefficient >0.7.

Dubowsky et al. (1999) point out that, although the variation of $PM_{2.5}$ mass concentration across a community may be small, there may be significant spatial variations of specific components of the total mass on a local scale. An example is given of a study of concentrations of polycyclic aromatic hydrocarbons (PAH) at three indoor locations in a community: (1) an urban and (2) a semi-urban site separated by 1.6 km, and (3) a suburban site located further away. The authors found the geometric mean PAH concentrations at these three locations varied respectively as 31:19:8 ng/m^3, and suggest that the local variations in traffic density were responsible for this gradient. Note that these concentrations are 1,000 times lower than the total

PM mass concentration, so that such a small gradient would not be detectable for total $PM_{2.5}$ mass measurements on the order of 25 μg m^{-3}.

The THEES study reported by Waldman et al. (1991) measured indoor, outdoor, and personal BaP levels and found that the outdoor BaP was the same at all outdoor sites across the three sampling periods. This study showed the seasonal differences vs. BaP levels and exposures due to indoor and outdoor sources and individual activities.

Leaderer et al. (1999) monitored 24-h PM_{10}, $PM_{2.5}$, and sulfates during the summers of 1995 and 1996 at a regional site in Vinton, VA (6 km from Roanoke, VA). One similar 24-h measurement was made outdoors at residences in the surrounding area, at distances ranging from 1 km to >175 km from the Vinton site, at an average separation distance of 96 km. The authors reported significant correlations for $PM_{2.5}$ and sulfates between the residential outdoor values and those measured at Vinton on the same day. In addition, the mean values of the regional site and residential site $PM_{2.5}$ and sulfates showed no significant differences in spite of the large distance separations and mountainous terrain intervening in most directions. However, for the concentrations of $PM_{2.5-10}$, estimated as PM_{10}-$PM_{2.5}$, no significant correlation among these sites was found ($n = 30$, $r = -0.20$).

Lillquist et al. (1998) found no significant gradient in PM_{10} concentrations in Salt Lake City, UT, when levels were low, but a gradient existed when levels were high. PM_{10} concentrations were measured outdoor at three hospitals using a Minivol 4.01 sampler (Airmetrics, Inc.) operating at 5 L min^{-1} and at the Utah Department of Air Quality (DAQ) ambient monitoring station located between 3 and 13 km from the hospitals for a period of about 5 mo.

Pope et al. (1999) monitored ambient PM_{10} concentrations in Provo, UT (Utah Valley), during the same time frame the following year and reported nearly identical concentrations at three sites separated by 4 to 12 km. Pearson correlation coefficients for the data were between 0.92 and 0.96. The greater degree of variability in the Salt Lake City PM_{10} data relative to the Provo data may be related to the higher incidence of wind-blown crustal material in Salt Lake City. Pope et al. (1999) reported that increased health effects in the Utah Valley were associated with stagnation and thermal inversions trapping anthropogenically derived PM_{10}, whereas, no increases in health effects were observed when PM_{10} levels were increased during events of wind blown crustal material.

Väkevä et al. (1999) found significant vertical gradients in submicron particles existed in an urban street canyon of Lahti, Finland. Particle number concentrations were measured using a TSI screen diffusion battery and a condensation particle counter at 1.5 and 25 m above the street at rooftop level. The authors found a fivefold decrease in concentration between the two sampling heights and attributed the vertical gradient to dilution and dispersion of pollutants emitted at street level.

White (1998**)** suggests that the higher random measurement error for the coarse PM fraction compared to the error for the fine PM fraction may be responsible for a major portion of the apparent greater spatial variability of coarse ambient PM concentration compared to fine ambient PM concentration in a community (e.g., Burton et al., 1996; Leaderer et al., 1999). When $PM_{2.5}$ and PM_{10} are collected independently, and the coarse fraction is obtained by difference ($PM_{2.5-10} = PM_{10}-PM_{2.5}$), then the expected variance in the coarse fraction is influenced by the variances of the PM_{10} and $PM_{2.5}$ measurements. When a dichotomous sampler collects $PM_{2.5}$ and $PM_{2.5-10}$ on two separate filters, the coarse fraction also is expected to have a larger error than the fine fraction. There is a possible error caused by loss of mass below the cut-point size and a gain of mass above the cut-point size that is created by the asymmetry of the product of the penetration times PM concentration about the cut-point size. Because a dichotomous PM sampler collects coarse mass using an upper and lower cut-point, it is expected to have a larger variance than for fine mass collected using only one cut-point.

Wilson and Suh (1997) conclude that $PM_{2.5}$ and PM_{10} concentrations are correlated more highly across Philadelphia than are $PM_{2.5-10}$ concentrations. Ambient monitoring data from 1992 to 1993 was reviewed for $PM_{2.5}$, $PM_{2.5-10}$, and PM_{10}, as well as for $PM_{2.5}$ and $PM_{2.5-10}$ dichotomous data for 212 site-years of information contained in the AIRS database (U.S. Environmental Protection Agency, 2000). The authors also observed that PM_{10} frequently was correlated more highly with $PM_{2.5}$ than with $PM_{2.5-10}$. The authors note that $PM_{2.5}$ constitutes a large fraction of PM_{10}, and that this is the likely reason for the strong agreement between $PM_{2.5}$ and PM_{10}. Similar observations were made by Keywood et al. (1999) in six Australian cities. The authors reported that PM_{10} was more highly correlated with $PM_{2.5}$ than with coarse PM ($PM_{2.5-10}$), suggesting that "variability in PM_{10} is dominated by variability in $PM_{2.5}$."

Lippmann et al. (2000) examined the site-to-site temporal correlations in Philadelphia (1981 to 1994) and found the ranking of median site-to-site correlation was O_3 (0.83), PM_{10}

(0.78), TSP (0.71), NO_2 (0.70), CO (0.50), and SO_2 (0.49). The authors explain that O_3 and a fraction of TSP and PM_{10} (e.g., sulfate) are secondary pollutants that would tend to be distributed spatially more uniformly within the city than primary pollutants such as CO and SO_2, which are more likely to be influenced by local emission sources. Lippman et al. (2000) conclude: "Thus, spatial uniformity of pollutants may be due to area-wide sources, or to transport (e.g., advection) of fairly stable pollutants into the urban area from upwind sources. Relative spatial uniformity of pollutants would therefore vary from city to city or region to region."

5.4.3.2.2 Physical Factors Affecting Indoor Microenvironmental Particulate Matter Concentrations

Several physical factors affect ambient particle concentrations in the indoor microenvironment, including air exchange, penetration, and particle deposition. Combined, these factors are critical variables that describe ambient particle dynamics in the indoor μ**e** and, to a large degree, significantly affect an individual's personal exposure to ambient-generated particles while indoors. The relationship between ambient outdoor particles and ambient particles that have infiltrated indoors is given by:

$$C_{ai} / C_{ao} = Pa / (a + k), \qquad (5\text{-}12)$$

where C_{ai} and C_{ao} are the concentration of ambient indoor and outdoor particles, respectively; *P* is the penetration factor; *a* is the air exchange rate; and *k* is the particle deposition rate (as discussed in Section 5.3.2.3.1, use of this model assumes equilibrium conditions and assumes that all variables remain constant). Particle penetration is a dimensionless quantity that describes the fraction of ambient particles that effectively penetrates the building shell. "Air exchange" is a term used to describe the rate at which the indoor air in a building or residence is replaced by outdoor air. The dominant processes governing particle penetration are air exchange and deposition of particles as they traverse through cracks and crevices and other routes of entry into the building. Although air-exchange rates have been measured in numerous studies, very few field data existed prior to 1996 to determine size-dependent penetration factors and particle deposition rates. All three parameters (P, *a*, and *k*) may vary substantially depending on building type, region of the country, and season. In the past several years, researchers have made

significant advancements in understanding the relationship between particle size and penetration factors and particle deposition rates. This section will highlight the studies that have been conducted to better understand physical factors affecting indoor particle dynamics.

Air-Exchange Rates

The air-exchange rate, *a*, in a residence varies depending on a variety of factors, including geographical location, age of the building, the extent to which window and doors are open, and season. Murray and Burmaster (1995) used measured values of *a* from households throughout the United States to describe empirical distributions and to estimate univariate parametric probability distributions of air-exchange rates. Figure 5-4 shows the results classified by season and region. In general, *a* is highest in the warmest region and increases from the coldest to the warmest region during all seasons. Air-exchange rates also are quite variable within and between seasons, as well as between regions (Figure 5-4). Data from the warmest region in summer should be viewed cautiously as many of the measurements were made in Southern California in July, when windows were more likely to be open than in other areas of the country where air-conditioning is used. Use of air-conditioning generally results in lowering air-exchange rates. In a separate analyses of these data, Koontz and Rector (1995) suggested that a conservative estimate for air exchange in residential settings would be 0.18 h^{-1} (10th percentile) and a typical air exchange would be 0.45 h^{-1} (50th percentile).

These data provide reasonable experimental evidence that *a* varies by season in locations with distinct seasons. As a result, infiltration of ambient particles may be more efficient during warmer seasons when windows are likely to be opened more frequently and air-exchange rates are higher. This suggests that the fraction of ambient particles present in the indoor μ**e** would be greater during warmer seasons than colder seasons. For example, in a study conducted in Boston, MA, participants living in non-air-conditioned homes kept the windows closed except during the summer (Long et al., 2000a). This resulted in higher and more variable air-exchange rates in summer than during any other season (Figure 5-5). During nighttime periods, when indoor sources are negligible, the indoor/outdoor concentration ratio or infiltration factor may be used to determine the relative contribution of ambient particles in the indoor microenvironment. Particle data collected during this study (Figure 5-6) shows the indoor/outdoor concentration ratios by particle size. Data show that, for these nine homes in Boston, the fraction of ambient



Figure 5-4. Air-exchange rates measured in homes throughout the United States. Climatic regions are based on heating-degree days: Coldest region ≥ 7000, Colder region = 5500 to 6999, Warmer region = 2500 to 4999, and Warmest region ≤ 2500 heating-degree days.

Based on data from Murray and Burmaster (1995).

particles penetrating indoors is higher during summer when air exchange rates were higher than fall (Long et al., 2000b).

Particle Deposition Rates and Penetration Factors

Physical factors affecting indoor particle concentrations, including particle deposition rates, *k*, and penetration factors, *P*, are possibly the most uncertain and variable quantities. Although *k* can be modeled with some success, direct measurements are difficult and results often vary from study to study. Particle deposition rates vary considerably depending on particle size because of the viscous drag of air on the particles hindering their movement to varying degrees. The nature



Figure 5-5. Box plots of hourly air-exchange rates stratified by season in Boston, MA, during 1998.

Source: Long et al. (2000a).



Figure 5-6. Geometric mean infiltration factor (indoor/outdoor ratio) for hourly nighttime, nonsource data for two seasons. Box plots of air exchange rates are shown as inserts for each plot. (Boston, 1998)

Source: Long et al. (2000b).

and composition of particles also affect deposition rates. Surface properties of particles, such as their electrostatic properties, can have a significant influence on deposition rates. In addition, thermophoresis can also affect *k*, but probably to a lesser degree in the indoor μ**e** because temperatures generally vary over a small range. Combined, these effects can produce order of magnitude variations in *k* between particles of different size and, in the case of electrophoresis and thermophoresis, particles of the same size.

Particle penetration efficiency into indoor microenvironments depends on particle size and air exchange rates. Penetration varies with particle size because of the size-dependent deposition of particles caused by impaction, interception, and diffusion of particles onto surfaces as they traverse through cracks and crevices. Penetration also is affected by air exchange rates. When air exchange rates are high, P approaches unity because the majority of ambient particles have less interaction with the building shell. In contrast, when air exchange rates are low, P is governed by particle deposition as particles travel through cracks and crevices.

Significant advancements have been made in the past few years to better characterize particle deposition rates and penetration factors. Several new studies, including two in which semi-continuous measurements of size distributions were measured indoors and outdoors, have produced new information on these quantities, which are key to understanding the contributions of ambient PM to indoor PM concentrations (Equation 5-10).

Studies involving semi-continuous measurements of indoor and outdoor particle size distributions have been used to estimate *k* and P as a function of particle size (Vette et al., 2001; Long et al., 2000b; Abt et al., 2000b). These studies each demonstrated that the indoor/outdoor concentration ratios (C_{ai}/C_{ao} in Equation 5-12) were highest for accumulation mode particles and lowest for ultrafine and coarse-mode particles. Various approaches were used to estimate size-specific values for *k* and P. Vette et al. (2001) and Abt et al. (2000b) estimated *k* by measuring the decay of particles at times when indoor levels were significantly elevated. Vette et al. (2001) estimated P using measured values of *k* and indoor/outdoor particle measurements during nonsource nighttime periods. Long et al. (2000b) used a physical-statistical model, based on Equation 5-10, to estimate *k* and P during nonsource nighttime periods. The results for *k* reported by Long et al. (2000b) and Abt et al. (2000b) are compared with other studies in Figure 5-7. Although not shown in Figure 5-7, the results for *k* obtained by Vette et al. (2001) were similar to the values of *k* reported by Abt et al. (2000b) for particle sizes up to 1 μm.



[a]Decay rates represent Summary Estimates from the four houses examined.
[b]Decay rates are based on sulfate and are presented as <2.5 µm.
Estimates were computed using a surface-to-volume ratio of 2 m^{-1} (Koutrakis *et al.*, 1992).
[c]Data represent $PM_{2.5}$
[d]Particle sizes are the midpoint of the ranges examined.
[e]Decay rates presented are estimates of k for nightly average data from all nine study homes.
[f]Decay rates are theorectically modeled deposition values for smooth indoor surfaces and homogeneous and isotropically turbulent air flow.
Presented curves assume typical room dimensions (3 m x 4 m x 5 m) and a friction velocity of 1.0 cm/s.

Figure 5-7. Comparison of deposition rates from this study with literature values (adapted from Abt et al., 2000b). Error bars represent standard deviations for same-study estimates.

Source: Adapted from Long et al. (2000b).

Results for P by Long et al. (2000b) show that penetration was highest for accumulation-mode particles and decreased substantially for coarse-mode particles (Figure 5-8). The results for P reported by Vette et al. (2001) show similar trends, but are lower than those reported by Long et al. (2000b). This likely is because of lower air-exchange rates in the Fresno, CA, residence (a ≈ 0.5 h^{-1}; Vette et al., 2001) than the Boston, MA, residences (a > 1 h^{-1}; Long et al., 2000b). These data for P and k illustrate the role that the building shell may provide in increasing the



Figure 5-8. Penetration efficiencies and deposition rates from models of nightly average data. Error bars represent standard errors. (Boston, 1998, winter and summer)

Source: Long et al. (2000b).

concentration of particles because of indoor sources and reducing the concentration of indoor particles from ambient sources, especially for homes with low air-exchange rates.

Compositional Differences Between Indoor-Generated and Ambient-Generated Particulate Matter

Wilson et al. (2000) discuss the differences in composition between particles from indoor and outdoor sources. They note that, because of the difficulty in separating indoor PM into ambient and nonambient PM, there is little direct experimental information on the composition differences between the two. Although experimental data are limited, Wilson et al. (2000) suggest the following.

Photochemistry is significantly reduced indoors; therefore, most secondary sulfate [H_2SO_4, NH_4HSO_4, and $(NH_4)_2SO_4$] and nitrate (NH_4NO_3) found indoors come from ambient sources. Primary organic emissions from incomplete combustion may be similar, regardless of the source. However, atmospheric reactions of polyaromatic hydrocarbons and other organic compounds produce highly oxygenated and nitrated products, so these species are also of ambient origin. Gasoline, diesel fuel, and vehicle lubricating oil all contain naturally present metals or metal additives. Coal and heavy fuel oil also contain more metals and nonmetals, such as selenium and arsenic, than do materials such as wood or kerosene burned inside homes. Environmental tobacco smoke (ETS), however, with its many toxic components, is primarily an indoor-generated pollutant.

Particles generated indoors may have different chemical and physical properties than those generated by anthropogenic ambient sources. Siegmann et al. (1999) have demonstrated that elemental carbon in soot particles generated indoors have different properties than in those generated outdoors by automotive or diesel engines. In the United States, combustion-product PM in the ambient/outdoor air generally is produced by burning fossil fuels (e.g., coal, gasoline, fuel oil) and wood, whereas combustion-product PM from indoor sources is produced by biomass burning (e.g., tobacco, wood, foods, etc.). However, some indoor sources of PM (such as cigarette smoking, meat cooking, and coal burning) occur both indoors and outdoors and may constitute an identifiable portion of measured ambient PM (Cha et al., 1996; Kleeman and Cass, 1998).

Indoor Air Chemistry

Gas- and aerosol-phase chemical reactions in the indoor microenvironment are responsible for secondary particle formation and modification of existing particles. This process could be complex and may influence the interpretation of exposures to indoor generated particles in instances when particles are generated by outdoor gases reacting with gases indoors to produce fresh particles. For example, homogeneous gas phase reactions involving ozone and terpenes (specifically d-limonene, α-terpinene, and α-pinene) have been identified as an important source of submicron particles (Weschler and Shields, 1999). Terpenes are present in several commonly available household cleaning products and d-limonene has been identified in more than 50% of the buildings monitored in the BASE study (Hadwen et al., 1997). Long et al. (2000a) found that when PineSol (primary ingredient is α-pinene) was used indoors, indoor $PM_{2.5}$ mass

concentrations increased by 3 to 32 μg m^{-3} (indoor ozone concentrations unknown, but ambient ozone concentrations were 44 to 48 ppb). Similarly, a 10-fold increase in number counts of 0.1 to 0.2 μm particles was observed in an experimental office containing supplemented d-limonene and normally encountered indoor ozone concentrations (< 5 to 45 ppb), resulting in an average increase in particle mass concentration of 2.5 to 5.5 μg m^{-3} (Weschler and Shields, 1999). Ozone appears to be the limiting reagent as particle number concentration varied proportionally to ozone concentrations (Weschler and Shields, 1999). Other studies showed similar findings (e.g., Jang and Kamens, 1999; Wainman et al., 2000).

Indoor Sources of Particles

The major sources of indoor PM in nonsmoking residences and buildings include suspension of PM from bulk material, cooking, cleaning, and the use of combustion devices, such as stoves and kerosene heaters. Human and pet activities also lead to PM detritus production (from tracked-in soil, fabrics, skin and hair, home furnishings, etc.), which is found ubiquitously in house dust deposited on floors and other interior surfaces. House dust and lint particles may be resuspended indoors by agitation (cleaning) and turbulence (HVAC systems, human activities, etc.). Ambient particles that have infiltrated into the indoor **μe** also may be resuspended after deposition to indoor surfaces. Typically, resuspension of particles from any source involves coarse-mode particles (>1 μm); particles of smaller diameter are not resuspended efficiently. On the other hand, cooking produces both fine- and coarse-mode particles, whereas combustion sources typically produce fine-mode particles.

Environmental tobacco smoke (ETS) is also a major indoor source of PM. It is, however, beyond the scope of this chapter to review the extensive literature on ETS. A number of articles provide source strength information for cigarette or cigar smoking (e.g., Daisey et al., 1998 and Nelson et al., 1998).

A study conducted on two homes in the Boston metropolitan area (Abt et al., 2000a) showed that indoor PM sources predominate when air exchange rates were <1 h^{-1}, and outdoor sources predominate when air exchange rates were $\geq$2 h^{-1}. The authors attributed this to the fact that when air-exchange rates were low (<1 h^{-1}), particles released from indoor sources tend to accumulate because particle deposition is the mechanism governing particle decay and not air exchange. Particle deposition rates are generally <1 h^{-1}, especially for accumulation-mode

particles. When air-exchange rates were higher ($\geq 2\ h^{-1}$), infiltration of ambient aerosols and exfiltration of indoor-generated aerosols occur more rapidly, reducing the impact of indoor sources on indoor particle levels. The study also confirmed previous findings that the major indoor sources of PM are cooking, cleaning, and human activity. They discuss the size characteristics of these ubiquitous sources and report the following.

> The size of the particles generated by these activities reflected their formation processes. Combustion processes (oven cooking, toasting, and barbecuing) produced fine particles and mechanical processes (sauteing, frying, cleaning, and movement of people) generated coarse particles. These activities increased particle concentrations by many orders of magnitude higher than outdoor levels and altered indoor size distributions. (Abt et al., 2000a; p. 43)

They also note that variability in indoor PM for all size fractions was greater than for outdoor PM, especially for short averaging times (2 to 33 times higher).

In a separate study conducted in nine nonsmoking homes in the Boston area, Long et al. (2000a) concluded that the predominant source of indoor fine particles was infiltration of outdoor particles, and that cooking activities were the only other significant source of fine particles. Coarse particles, however, had several indoor sources, such as cooking, cleaning, and various indoor activities. This study also concluded that more than 50% of the particles (by volume) generated during indoor events were ultrafine particles. Events that elevated indoor particle levels were found to be brief, intermittent, and highly variable, thus requiring the use of continuous instrumentation for their characterization. Table 5-11 provides information on the mean volume mean diameter (VMD) for various types of indoor particle sources. The differences in mean VMD confirm the clear separation of source types and suggest that there is very little resuspension of accumulation-mode PM. In addition, measurements of organic and elemental carbon indicated that organic carbon had significant indoor sources, whereas elemental carbon was primarily of ambient origin.

Vette et al. (2001) found that resuspension was a significant indoor source of particles >1 μm, whereas fine- and accumulation-mode particles were not affected by resuspension. Figure 5-9 shows the diurnal variability in the indoor/outdoor aerosol concentration ratio from an unoccupied residence in Fresno. The study was conducted in the absence of common indoor

TABLE 5-11. VOLUME MEAN DIAMETER (VMD) AND MAXIMUM $PM_{2.5}$ CONCENTRATIONS OF INDOOR PARTICLE SOURCES [a,b]

		Size Statistics		$PM_{2.5}$	
Particle Source	N	Indoor Activity Mean VMD (μm)	Background[a,e] Mean VMD (μm)	Maximum Mean	Concentration[c,d] SD
Cooking					
Baking (Electric)	8	0.189[f]	0.221[f]	14.8	7.4
Baking (Gas)	24	0.107[f]	0.224[f]	101.2	184.9
Toasting	23	0.138[f]	0.222[f]	54.9	119.7
Broiling	4	0.114[f]	0.236[f]	29.3	43.4
Sautéing	13	0.184[f], 3.48[g]	0.223[f], 2.93[g]	65.6	95.4
Stir-Frying	3	0.135[f]	0.277[f]	37.2	31.4
Frying	20	0.173[f]	0.223[f]	40.5	43.2
Barbecuing	2	0.159[f]	0.205[f]	14.8	5.2
Cleaning					
Dusting	11	5.38[g]	3.53[g]	22.6	22.6
Vacuuming	10	3.86[g]	2.79[g]	6.5	3.9
Cleaning with Pine Sol	5	0.097[f]	0.238[f]	11.0	10.2
General Activities					
Walking Vigorously (w/Carpet)	15	3.96[g]	3.18[g]	12.0	9.1
Sampling w/Carpet	52	4.25[g]	2.63[g]	8.0	6.6
Sampling w/o Carpet	26	4.28[g]	2.93[g]	4.8	3.0
Burning Candles	7	0.311[f]	0.224[f]	28.0	18.0

Notes:
[a]All concentration data corrected for background particle levels.
[b]Includes only individual particle events that were unique for a given time period and could be detected above background particle levels.
[c]PM concentrations in $\mu g/m^3$.
[d]Maximum concentrations computed from 5-min data for each activity.
[e]Background data are for time periods immediately prior to the indoor event.
[f]Size statistics calculated for $PV_{0.02-0.5}$ using SMPS data.
[g]Size statistics calculated for $PV_{0.7-10}$ using APS data.

Source: Long et al. (2000a).

particle sources such as cooking and cleaning. The data in Figure 5-9 show the mean indoor/outdoor concentration ratio for particles >1 μm increased dramatically during daytime hours. This pattern was consistent with indoor human activity levels. In contrast, the mean



Figure 5-9. Mean hourly indoor/outdoor particle concentration ratio from an unoccupied residence in Fresno, CA, during spring 1999.

Source: Vette et al. (2001).

indoor/outdoor concentration ratio for particles <1 μm (fine- and accumulation-mode particles) remain fairly constant during both day and night.

5.4.3.2.3 Time/Activity Patterns

Total exposure to PM is the sum of various microenvironmental exposures that an individual encounters during the day and will depend on the microenvironments occupied. As discussed previously, PM exposure in each microenvironment is the sum of exposures from ambient sources (E_{ag}), indoor sources (E_{pig}), and personal activities (E_{pact}). E_{ag} and E_{pig} are determined by the microenvironments in which an individual spends time; whereas E_{pact} is determined by the personal activities that he/she conducts while in those microenvironments. As mentioned before, PM exposures and its components are variable across the population; and,

thus, each are distributions rather than point estimates. A thorough analyses of these distributions would require a comprehensive sensitivity and uncertainty analysis.

Determining microenvironments and activities that contribute significantly to human exposure begins with establishing human activity pattern information for the general population, as well as subpopulations. Personal exposure and time activity pattern studies have shown that different populations have varying time activity patterns and, accordingly, different personal PM exposures. Both characteristics will vary greatly as a function of age, health status, ethnic group, socioeconomic status, season, and region of the country. Collecting detailed time activity data can be very burdensome on participants but is clearly valuable in assessing human exposure and microenvironments. For modeling purposes, human activity data frequently come from general databases that are discussed below.

The gathering of human activity information, often called “time-budget” data, started in the 1920s; however, their use for exposure assessment purposes only began to be emphasized in the 1980s. Many of the largest U.S. human activity databases have been consolidated by EPA’s National Exposure Research Laboratory’s (NERL) into one comprehensive database containing over 22,000 person-days of 24-h activity known as the Consolidated Human Activity Database, or CHAD (Glen et al., 1997; McCurdy et al., 2000). The information in CHAD is accessible for constructing population cohorts of people with diverse characteristics that are useful for analysis and modeling (McCurdy, 2000). See Table 5-2 for a summary listing of human activity studies in CHAD. Most of the databases in CHAD are available elsewhere, including the National Human Activity Pattern Survey (NHAPS), California’s Air Resources Board (CARB), and the University of Michigan’s Institute for Survey Research data sets.

Although CHAD provides a very valuable resource for time and location data, there is less information on PM-generating personal activities. In addition, very few of the time-activity studies have collected longitudinal data within a season or over multiple seasons. Such longitudinal data are important in understanding potential variability in activities and how they impact correlations between PM exposure and ambient site measurements for both total PM and PM of ambient origin.

5.4.3.3 Impact of Ambient Sources on Exposures to Particulate Matter

Different sources may generate ambient PM with different aerodynamic and chemical characteristics, which may, in turn, result in different health responses. Thus, to fully understand the relationship between PM exposure and health outcome, exposure from difference sources should be identified and quantified. Source apportionment techniques provide a method for determining personal exposure to PM from specific sources. Daily contributions from sources that have no indoor component can be used as tracers to generate exposure estimates for ambient PM of similar aerodynamic size or directly as exposure surrogates in epidemiologic analyses. The most recent EPA PM Research Needs Document (U. S. Environmental Protection Agency, 1998) recommended use of source apportionment techniques to determine daily time-series of source categories for use in community, time-series epidemiology.

A number of epidemiological studies (discussed more fully in Chapter 8) have evaluated relationships between health outcomes and sources of particulate matter determined from measurements at a community monitor. These studies suggest the importance of examining sources and constituents of indoor, outdoor, and personal PM. For example, Özkaynak and Thurston (1987) evaluated the relationship between PM sources and mortality in 36 Standard Metropolitan Statistical Areas (SMSAs). Particulate matter samples from EPA's Inhalable Particle (IP) Network were analyzed for SO_4^{2-} and NO_3^- by automated colorimetry, and elemental composition was determined with X-ray fluorescence (XRF). Mass concentrations from five PM source categories were determined from multiple regression of absolute factor scores on the mass concentration: (1) resuspended soil, (2) auto exhaust, (3) oil combustion, (4) metals, and (5) coal combustion.

In another study, Mar et al. (2000) applied factor analysis to evaluate the relationship between PM composition (and gaseous pollutants) in Phoenix. In addition to daily averages of $PM_{2.5}$ elements from XRF analysis, they included in their analyses organic and elemental carbon in $PM_{2.5}$ and gaseous species emitted by combustion sources (CO, NO_2, and SO_2). They identified five factors classified as (1) motor vehicles, (2) resuspended soil, (3) vegetative burning, (4) local SO_2, and (5) regional sulfate.

Also, Laden et al. (2000) applied specific rotation factor analysis to particulate matter composition (XRF) data from six eastern cities (Ferris et al., 1979). Fine PM was regressed on

the recentered scores to determine the daily source contributions. Three main sources were identified: (1) resuspended soil (Si), (2) motor vehicle (Pb), and (3) coal combustion (Se).

Source apportionment or receptor modeling has been applied to the personal exposure data to understand the relationship between personal and ambient sources of particulate matter. Application of source apportionment to ambient, indoor, and personal PM composition data is especially useful in sorting out the effects of particle size and composition. If a sufficient number of samples are analyzed with sufficient compositional detail, it is possible to use statistical techniques to derive source category signatures, identify indoor and outdoor source categories and estimate their contribution to indoor and personal PM.

Positive Matrix Factorization (PMF) has been applied to the PTEAM database by Yakovleva et al. (1999). The authors utilize mass and XRF elemental composition data from indoor and outdoor $PM_{2.5}$ and personal, indoor, and outdoor PM_{10} samples. PMF is an advance over ordinary factor analysis because it allows measurements below the quantifiable limit to be used by weighting them by their uncertainty. This effectively increases the number of species that can be used in the model. The factors used by the authors correspond to general source categories of PM, such as outdoor soil, resuspended indoor soil, indoor soil, personal activities, sea-salt, motor vehicles, nonferrous metal smelters, and secondary sulfates. PMF, by identifying not only the various source factors but also apportioning them among the different monitor locations (personal, indoor, and outdoor), was able to quantify an estimate of the contribution of resuspended indoor dust to the personal cloud (15% from indoor soil and 30% from resuspended indoor soil). Factor scores for these items then were used in a regression analysis to estimate personal exposures (Yakovleva et al., 1999).

The most important contributors to PM_{10} personal exposure were indoor soil, resuspended indoor soil, and personal activities; these accounted for approximately 60% of the mass (Yakovleva et al., 1999). Collectively, they include personal cloud PM, smoking, cooking, and vacuuming. For both $PM_{2.5}$ and PM_{10}, secondary sulfate and nonferrous metal operations accounted for another 25% of PM mass. Motor vehicle exhausts, especially starting a vehicle inside of an attached garage, accounted for another 10% of PM mass. The authors caution that these results may not apply to other geographic areas, seasons of the year, or weather conditions.

Simultaneous measurement of personal (PM_{10}) and outdoor measurements ($PM_{2.5}$ and PM_{10}) were evaluated as a three-way problem with PMF, which allowed for differentiation of

source categories based on their variation in time and type of sample, as well as their variation in composition. By use of this technique, it was possible to identify three sources of coarse-mode, soil-type PM. One was associated with ambient soil, one with indoor soil dispersed throughout the house, and one with soil resulting from the personal activity of the subject.

Two other source apportionment models have been applied to ambient measurement data and can be used for the personal exposure studies. The effective variance weighted Chemical Mass Balance (CMB) receptor model (Watson et al., 1984, 1990, 1991) solves a set of linear equations that incorporate the uncertainty in the sample and source composition. CMB requires the composition of each potential source of PM and the uncertainty for the sources and ambient measurements. Source apportionment with CMB can be conducted on individual samples; however, composition of each of the sources of PM must be known. An additional source apportionment model, UNMIX (Henry et al., 1994) is a multivariate source apportionment model. UNMIX is similar to PMF, but does not use explicitly the measurement uncertainties. Because measurement uncertainties are not used, only species above the detection limit are evaluated in the model. UNMIX provides the number of sources and source contributions and requires a similar number of observations as PMF.

The Yakovleva et al. (1999) study demonstrates that source apportionment techniques also could be very useful in determining parameters needed for exposure models and for determining exposure to ambient-generated PM. Exposure information, similar to that obtained in the PTEAM study, but including other PM components useful for definition of other source categories (e.g., elemental [EC] and organic carbon [OC]; organic tracers for elemental carbon from diesel vehicle exhaust, gasoline vehicle exhaust, and wood combustion; nitrate; Na; Mg and other heavy metal tracers; and, also, gas-phase pollutants) would be useful as demonstrated in the use of EC/OC and gas-phase pollutants by Mar et al. (2000).

5.4.3.4 Correlations of Particulate Matter with Other Pollutants

Several epidemiological studies have included the gaseous pollutants CO, NO_2, SO_2, and O_3 along with PM_{10} or $PM_{2.5}$ in the analysis of the statistical association of health responses with pollutants. In a recent study, the personal exposure to O_3 and NO_2 were determined, as well as that to $PM_{2.5}$ and $PM_{2.5\text{-}10}$ for a cohort 15 elderly subjects in Baltimore, MD, although measured personal exposures to O_3, NO_2, and SO_2 were below their respective LOD for 70% of the

samples. Spearman correlations for 14 subjects in summer and 14 subjects in winter are given in Table 5-12 for relationships between personal $PM_{2.5}$ and ambient concentrations of $PM_{2.5}$, $PM_{2.5-10}$, O_3, and NO_2. In contrast to ambient concentrations, neither personal exposure to total $PM_{2.5}$ nor $PM_{2.5}$ ambient origin was correlated significantly with personal exposures to the co-pollutants, $PM_{2.5-10}$, nonambient $PM_{2.5}$, O_3, NO_2, and SO_2. Personal-ambient associations for $PM_{2.5-10}$, O_3, NO_2, and SO_2 were similarly weak and insignificant. Based on these results, Sarnat et al. (2000) conclude that the potential for confounding of $PM_{2.5}$ by O_3, NO_2, or $PM_{10-2.5}$ appears to be limited, because, despite significant correlations observed among ambient pollutant concentrations, the correlations among personal exposures were low.

Sarnat et al. (2001) further evaluated the role of gaseous pollutants in particulate matter epidemiology by extending the measurements taken on the earlier adult cohort of 20 individuals in Baltimore by including additional PM and gaseous pollutant measurements that were collected during the same 1998-1999 period from 15 individuals with chronic obstructive pulmonary disease (COPD) and from 21 children. 24-h average personal exposures for $PM_{2.5}$, O_3, SO_2 and NO_2, and corresponding ambient concentrations for $PM_{2.5}$, O_3, SO_2 , NO_2 and CO for all 56 subjects were collected over 12 consecutive days. Results from correlation and regression analysis of the personal and ambient data showed that personal $PM_{2.5}$ and personal gaseous pollutant exposures were generally not correlated. The analysis also showed that ambient $PM_{2.5}$ concentrations had significant associations with personal $PM_{2.5}$ exposures in both seasons. On the other hand, ambient gaseous pollutant concentrations were not correlated with their corresponding personal exposure concentrations. However, ambient gaseous concentrations were found to be strongly associated with personal $PM_{2.5}$ exposures, suggesting that ambient gaseous concentrations for O_3, NO_2, SO_2 are acting as surrogates, as opposed to confounders of $PM_{2.5}$, in the estimation of PM health effects based on multi pollutant models. This study did not measure personal CO and also did not find a significant association between summertime ambient CO and personal $PM_{2.5}$ (a significant winter time association, however, was found). Personal EC and SO_4 were also measured during the winter for the cohort of COPD patients only. The analysis of this subset of the data showed that personal SO_4 was significantly and negatively associated with ambient O_3 and SO_2, and personal EC was significantly associated with ambient O_3, NO_2 and CO. The authors interpret these findings as suggesting that O_3 is primarily a surrogate for secondary particle exposures, whereas ambient CO and NO_2 are primarily surrogates for particles

TABLE 5-12. CORRELATIONS BETWEEN PERSONAL $PM_{2.5}$ AND AMBIENT POLLUTANT CONCENTRATIONS[1]

		Personal $PM_{2.5}$ vs. Ambient:				Personal $PM_{2.5}$ of Ambient Origin vs. Ambient:		
SUMMER	Subject	$PM_{2.5}$	O_3	NO_2	$PM_{2.5-10}$	O_3	NO_2	$PM_{2.5-10}$
	SA1	0.55	0.15	0.38	-0.12	0.27	*0.71*	0.15
	SA2	*0.85*	0.31	*0.66*	0.57	0.21	0.64	0.68
	SA5	*0.89*	0.18	*0.82*	0.64	0.33	*0.81*	0.79
	SB1	*0.65*	0.40	-0.15	0.38	*0.89*	*-0.74*	-0.03
	SB2	-0.21	-0.62	*0.81*	0.15	0.26	0.08	0.33
	SB3	*0.82*	0.55	-0.14	-0.04	0.52	-0.20	0.00
	SB4	*0.73*	*0.62*	-0.34	-0.12	0.45	-0.29	-0.14
	SB5	*0.73*	0.45	-0.42	0.23	0.36	-0.48	0.33
	SB6	0.53	0.15	-0.38	0.12	-0.03	-0.57	0.32
	SC1	*0.95*	*0.78*	*0.66*	*0.65*	*0.83*	*0.63*	0.57
	SC2	*0.78*	*0.68*	0.36	0.51	*0.66*	*0.65*	*0.76*
	SC3	*0.85*	*0.78*	*0.73*	*0.68*	*0.69*	*0.71*	*0.80*
	SC4	*0.78*	*0.66*	0.59	*0.70*	0.50	0.50	0.51
	SC5	0.55	0.51	0.32	0.43	0.34	0.33	0.27
WINTER	WA1	0.22	-0.18	-0.26	-0.05	*-0.78*	-0.04	-0.24
	WA2	-0.38	-0.07	-0.36	-0.70	-0.15	-0.15	0.02
	WA4	-0.18	*0.67*	-0.22	-0.29	-0.33	0.20	0.00
	WA5	0.22	-0.43	*0.61*	0.50	*-0.72*	-0.09	0.40
	WB1	*0.80*	-0.84	*0.77*	0.41	*-0.87*	0.53	*0.66*
	WB2	*0.62*	-0.32	*0.59*	0.09	*-0.76*	*0.59*	*0.59*
	WB3	0.55	-0.45	*0.62*	0.04	*-0.77*	0.56	0.60
	WB4	-0.12	-0.01	0.34	-0.10	*-0.80*	*0.68*	0.48
	WC1	*0.74*	-0.62	-0.15	0.44	*-0.64*	0.02	0.69
	WC2	*0.79*	-0.88	0.17	*0.77*	*-0.87*	0.25	*0.71*
	WC3	0.28	-0.42	0.03	0.57	*-0.77*	0.30	-0.45
	WC4	0.19	-0.84	0.50	0.45	*-0.72*	0.22	0.67
	WC5	*0.81*	-0.62	0.08	*0.81*	*-0.76*	0.05	0.42
	WC6	0.01	-0.03	*0.65*	0.37	*-0.75*	0.19	-0.45
Median	Summer	0.76	0.48	0.37	0.41	0.41	0.42	0.33
Median	Winter	0.25	-0.43	0.26	0.39	-0.76	0.21	0.45

[1]Correlations represent Spearman's r values; italicized values indicate significance at the $\alpha = 0.05$ level.

Source: Sarnat et al. (2000).

from traffic. Sarnat et.(2001) caution that these findings were found in only one location and various physical and personal factors, such as ventilation, time spent outdoors, and household characteristics could affect the strength of the reported associations for certain individuals and cohorts, even though the qualitative results found are unlikely to change.

A newly developed Roll-Around System (RAS) was used to evaluate the hourly relationship between gaseous pollutants (CO, O_3, NO_2, SO_2, and VOCs) and PM (Chang et al., 2000). Exposures were characterized over a 15-day period for the summer and winter in Baltimore, based on scripted activities to simulate activities performed by older adults (65+ years of age). Spearman rank correlations were reported for $PM_{2.5}$, O_3, CO, and toluene for both the summer and winter and the correlations are given for each microenvironment in Table 5-13: indoor residence, indoor other, outdoor near roadway, outdoor away from road, and in vehicle. No significant relationships ($p < 0.05$) were found between hourly $PM_{2.5}$ and O_3. Significant relationships were found between hourly $PM_{2.5}$ and CO: indoor residence, winter; indoor other, summer and winter; and outdoor away from roadway, summer. Significant relationships also were found between hourly $PM_{2.5}$ and toluene: indoor residence, winter; indoor other, winter; and in vehicle, winter. The significant relationships between CO and $PM_{2.5}$ in the winter may be caused by reduced air-exchange rates that could allow them to accumulate (Chang et al., 2000). Although no significant correlation was found between in vehicle $PM_{2.5}$ and CO, toluene, which is a significant component of vehicle exhaust (Conner et al., 1995), was correlated significantly to $PM_{2.5}$ in the winter.

Carrer et al. (1998) present data on the correlations among personal and microenvironmental PM_{10} exposures and concentrations and selected environmental chemicals that were monitored simultaneously (using a method that was not described). These chemicals were nitrogen oxides (NO_x), carbon monoxide (CO), and total volatile organic compounds (TVOC), benzene, toluene, xylene, and formaldehyde. The Kendall τ correlation coefficient was used; only results significant at $p < 0.05$ are mentioned here. Significant associations were found only between the following pairs of substances (τ shown in parentheses): personal PM_{10} (24 h) and NO_x (0.34), CO (0.34), TVOC (0.18), toluene (0.19), and xylene (0.26); office PM_{10} and NO_x (0.31); home PM_{10} and NO_x (0.24), CO (0.24), toulene (0.17), and xylene (0.25). Surprisingly, because most of the chemical substances are associated with motor vehicular emissions, there

TABLE 5-13. CORRELATIONS BETWEEN HOURLY PERSONAL $PM_{2.5}$ AND GASEOUS POLLUTANTS

	Indoor Residence		Indoor Other		Outdoor Near Roadway		Outdoor Away from road		In Vehicle	
	N	r_s	N	r_s	N	r_s	N	r_s	N	r_s
$PM_{2.5}$ vs. O_3										
Summer	35	0.29	16	-0.14	10	0.05	12	0.45	37	0.21
Winter	56	0.05	37	-0.06	11	-0.28	7	0.04	34	-0.10
$PM_{2.5}$ vs. CO										
Summer	41	0.25	19	0.59[a]	13	0.14	12	0.62	46	0.23
Winter	59	0.43[a]	39	0.62[a]	13	0.37	8	0.41	37	0.10
$PM_{2.5}$ vs. Toluene										
Summer	46	0.23	21	-0.14	14	0.26	14	0.02	48	0.12
Winter	66	0.38[a]	47	0.44[a]	17	0.40	8	0.48	42	0.43[a]

[a]Correlations represent Spearman's r values; italicized values indicate significance at the $\alpha = 0.05$ level.

Source: Chang et al. (2000).

was no significant correlation between "commuting PM_{10}" and any of the substances (Carrer et al., 1998).

5.5 SUMMARY OF PARTICULATE MATTER CONSTITUENT DATA

5.5.1 Introduction

Atmospheric PM contains a number of chemical constituents that may be of significance with respect to the human exposure and health effects. These constituents may be either components of the ambient particles or bound to the surface of particles. They may be elements, inorganic species, or organic compounds. A limited number of studies have collected data on concentrations of elements, acidic aerosols, and polycyclic aromatic hydrocarbons (PAHs) in ambient, personal, and microenvironmental PM samples. But, there have not been extensive analyses of the constituents of PM in personal or microenvironmental samples. Data from

relevant studies are summarized in this section. The summary does not address bacteria, bioaerosols, viruses, or fungi (e.g., Owen et al., 1992; Ren et al., 1999).

5.5.2 Monitoring Studies That Address Particulate Matter Constituents

A limited number of studies have measured the constituents of PM in personal or microenvironmental samples. Relevant studies published in recent years are summarized in Tables 5-11 and 5-12 for personal exposure measurements of PM and microenvironmental samples, respectively. Studies that measured both personal and microenvironmental samples are included in Table 5-11.

The largest database on personal, microenvironmental, and outdoor measurements of PM elemental concentrations is the PTEAM study (Özkaynak et al., 1996b). The results are highlighted in the table and discussed below. The table shows that a number of studies have measured aerosol acidity, sulfate, ammonia, and nitrate concentrations. Also, a number of studies have measured PAHs, both indoors and outdoors. Other than the PAHs, there is little data on organic constituents of PM.

5.5.3 Key Findings

5.5.3.1 Correlations of Personal and Indoor Concentrations with Ambient Concentrations of Particulate Matter Constituents

The elemental composition of PM in personal samples was measured in the PTEAM study, the first probability-based study of personal exposure to particles. A number of important observations, made from the PTEAM data collected in Riverside, CA, are summarized by Özkaynak et al. (1996b). Population-weighted daytime personal exposures averaged $150 \pm 9\ \mu g/m^3$, compared to concurrent indoor and outdoor concentrations of $95 \pm 6\ \mu g/m^3$. The personal exposure measurements suggested that there was a “personal cloud” of particles associated with personal activities. Daytime personal exposures to 14 of the 15 elements measured in the samples were considerably greater than concurrent indoor or outdoor concentrations, with sulfur being the only exception.

The PTEAM data also showed good agreement between the concentrations of the elements measured outdoors at the backyard of the residences with the concentrations measured at the central site in the community. The agreement was excellent for sulfur. Although the particle and

element mass concentrations were higher in personal samples than for indoor or outdoor samples, a nonlinear mass-balance method showed that the penetration factor was nearly 1 for all particles and elements.

Similarly to the PTEAM results, recent measurements of element concentrations in NHEXAS showed elevated concentrations of As and Pb in personal samples relative to indoor and outdoor samples (Clayton et al., 1999b). The elevated concentrations of As and Pb were consistent with elevated levels of PM in personal samples (median particle exposure of 101 $\mu g/m^3$), compared to indoor concentrations (34.4 $\mu g/m^3$). There was a strong association between personal and indoor concentrations and indoor and outdoor concentrations for both As and Pb. However, there were no central site ambient measurements for comparison to the outdoor or indoor measurements at the residences.

Manganese (Mn) concentrations were measured in $PM_{2.5}$ samples collected in Toronto (Crump, 2000). The mean $PM_{2.5}$ Mn concentrations were higher outdoors than indoors. But the outdoor concentrations measured at the participant's homes were lower than those measured at two fixed locations. Crump (2000) suggested that the difference in the concentrations may have been because the fixed locations were likely closer to high-traffic areas than were the participant's homes.

Studies of acidic aerosols and gases typically measure strong acidity (H^+), SO_4^{2-}, NH^{4+}, and NO^{3-}. The relationship between the concentrations of these ions and the relationship between indoor and outdoor concentrations have been addressed in a number of studies during which personal samples, microenvironmental, and outdoor samples have been collected, as shown in Tables 5-14 and 5-15. Key findings from these studies include those shown below.

- Acid aerosol concentrations measured at the residences in the Uniontown, PA, study were significantly different from those measured at a fixed ambient site located 16 km from the community. But, Leaderer et al. (1999) reported that the regional ambient air monitoring site in Vinton, VA, provided a reasonable estimate of indoor and outdoor sulfate measurements during the summer at homes without tobacco combustion.
- Approximately 75% of the fine aerosol indoors during the summer was associated with outdoor sources based on I/O sulfate ratios measured in the Leaderer et al. (1999) study.
- Personal exposures to strong acidity (H^+) were lower than corresponding outdoor levels measured in studies by Brauer et al. (1989, 1990) and Suh et al. (1992). But the personal

TABLE 5-14. STUDIES THAT HAVE MEASURED PARTICULATE MATTER CONSTITUENTS IN PERSONAL EXPOSURE SAMPLES

PM Constituent	Study Name/Reference	Study Location	Population Size/No. of Samples	Summary of Results
Elements	PTEAM/Özkaynak et al. (1996b)	Riverside, CA	178 adults	Outdoor air was the major source for most elements indoors, providing 70 to 100% of the observed indoor concentrations for 12 of the 15 elements. Correlation coefficients for central monitoring site versus outdoor at the residences were 0.98 for sulfur and 0.5 to 0.9 for other elements (except copper).
As and Pb	NHEXAS/Clayton et al. (1999b)	EPA Region 5	167 samples	Personal As and Pb levels higher than indoor or outdoor levels. No community ambient site for comparison.
Mn	Pellizzari et al. (1998, 1999) Clayton et al. (1999a), Crump (2000)	Toronto	925 personal samples	Mean $PM_{2.5}$ Mn higher outdoors than indoors. But $PM_{2.5}$ Mn concentrations higher at two fixed locations than at participants' homes.
Acid Aerosol Constituents	Sarnat et al. (2000)	Baltimore, MD	20 adults	High correlations between personal and ambient sulfate measurements in summer and winter.
	Brauer et al. (1989)	Boston, MA	—	Personal exposures to aerosol strong acidity slightly lower than concentrations measured at stationary site.
	Suh et al. (1992)	Uniontown, PA	24 children for 2 days	Personal exposures to H^+ and SO_4^{-2} lower than outdoor levels, but higher than indoor microenvironmental levels; personal NH_4^+ and NO_3^- higher than indoor or outdoor levels.
	Suh et al. (1993a,b)	State College, PA	47 children	Results similar to Uniontown, PA, study.
	Suh et al. (1994)			Results indicate strong neutralization of acidity indoors.
	Waldman and Liang (1993), Waldman et al. (1990)	Georgia and New Jersey	Hospital, daycares	Indoor sulfate levels were 70 to 100% of outdoor levels. Indoor ammonia levels 5- to 50-times higher than outdoors. Indoors, acid aerosols were largely neutralized.
PAHs	Zmirou et al. (2000)	Grenoble, France	38 adults	Ambient air concentrations close to traffic emissions were 1.1- to 3.5-times higher than personal exposure concentrations.

DRAFT-DO NOT QUOTE OR CITE

TABLE 5-15. STUDIES THAT HAVE MEASURED PARTICULATE MATTER CONSTITUENTS IN MICROENVIRONMENTAL SAMPLES

PM Constituent	Study Name/Reference	Study Location	Population Size/No. of Samples	Summary of Results
Acid Aerosol Constituents	Jones et al. (2000)	Birmingham, England	12 residences	Sulfate I/O ratios ranged from 0.7 to 0.9 for three PM size fractions.
	Patterson and Eatough (2000)	Lindon, UT	One school	Ambient sulfate, SO_2, nitrate, soot, and total particle number showed strong correlations with indoor exposure, although ambient $PM_{2.5}$ mass was not a good indicator of total $PM_{2.5}$ exposure.
	Leaderer et al. (1999)	Virginia and Connecticut	232 homes	The regional ambient air monitoring site provided a reasonable estimate of indoor and outdoor sulfate at nonsmokers homes. I/O sulfate ratio of 0. 74 during summer. Ammonia concentrations were an order of magnitude higher indoors than outdoors. Nitrous acid levels higher indoors than outdoors.
	Brauer et al. (1990)	Boston, MA	11 homes	Outdoor levels of H^+, SO_2, HNO_3, and SO_4^{-2} exceeded indoor levels in winter and summer. I/O ratios of H^+ lower than I/O ratios of SO_4^{-2} indicated neutralization of the acidity by ammonia.
PAHs	Chuang et al. (1999)	Durham, NC	24 homes	Measurements with continuous monitor; PAH levels generally higher indoors than outdoors.
	Dubowsky et al. (1999)	Boston, MA	3 buildings	PAHs indoors attributable to traffic, cooking, and candle-burning.
	Sheldon et al. (1993a,b)	Placerville and Roseville, CA	280 homes	Mass balance model used to estimate source strengths for PAH sources such as smoking, wood-burning and cooking.
PAHs and phthalates	PTEAM/Özkaynak et al. (1996b), Sheldon et al. (1993c)	Riverside, CA	120 homes	12-h I/O ratios for particulate-phase PAHs ranged from 1.1 to 1.4 during the day and 0.64 to 0.85 during night. The concentrations of phthalates and the number of samples with detectable phthalates were higher indoors than outdoors.

DRAFT-DO NOT QUOTE OR CITE

exposure levels measured by Suh et al. (1992) were higher than the indoor microenvironmental levels.

- Personal exposures to NH^{4+}, and NO^{3-} were reported by Suh et al. (1992) to be lower than either indoor or outdoor levels.
- Personal exposures to SO_4^{2-} were also lower than corresponding outdoor levels, but higher than the indoor microenvironmental levels (Suh et al., 1992; 1993a,b), as shown in Table 5-16.

The fact that the personal and indoor H^+ concentrations were substantially lower than outdoor concentrations suggests that a large fraction of aerosol strong acidity is neutralized by ammonia. Ammonia is emitted in relatively high concentrations in exhaled breath and sweat. The difference between indoor and outdoor H^+ concentrations in the Suh et al. (1992, 1993a,b) studies was also much higher than the difference for indoor and outdoor SO_4^{2-}, indicative of neutralization of the H^+. Results of the Suh et al. (1992, 1993a,b) studies also showed substantial interpersonal variability of H^+ concentrations that could not be explained by variation in outdoor concentrations.

Similar results for ammonia were reported by Waldman and Liang (1993). They reported that levels of ammonia in institutional settings that they monitored were 10– to 50– times higher than outdoors, and that acid aerosols were largely neutralized. Leaderer et al. (1999) reported that ammonia concentrations during both winter and summer in residences were an order of magnitude higher indoors than outdoors, consistent with results of other studies and the presence of sources of ammonia indoors.

Sulfate aerosols appear to penetrate indoors effectively. Waldman et al. (1990) reported I/O ratios of 0.7 to 0.9 in two nursing care facilities and a day-care center. Sulfate I/O ratios were measured for three particle size fractions in 12 residences in Birmingham, England, by Jones et al. (2000). The sulfate I/O ratios were 0.7 to 0.9 for PM < 1.1 μm, 0.6 to 0.8 for PM 1.1 to 2.1 μm, and 0.7 to 0.8 for PM 2.1 to 10 μm. Suh et al. (1993b) reported that personal and outdoor sulfate concentrations were highly correlated, as depicted in Figure 5-10.

Indoor/outdoor relationships were measured for a number of $PM_{2.5}$ components and related species in Lindon, UT, during January and February of 1997 by Patterson and Eatough (2000). Outdoor samples were collected at the Utah State Air Quality monitoring site. Indoor samples

TABLE 5-16. SUMMARY STATISTICS FOR PERSONAL, INDOOR, AND OUTDOOR CONCENTRATIONS OF SELECTED AEROSOL COMPONENTS IN TWO PENNSYLVANIA COMMUNITIES

			Concentration (nmol m^{-3})		
Aerosol	Home Type	Sample Site (In/Out)[a]	Indoor (12 h) GM ± GSD[b]	Outdoor (24 h) GM ± GSD[b]	Personal (12 h) GM ± GSD[b]
State College					
NO^{3-}	A/C Homes[c]	53/71	2.1 ± 2.7	1.4 ± 2.1	—
	Non-A/C	254/71	3.2 ± 2.3	1.4 ± 2.1	—
SO_4^{2-}	A/C Homes	56/75	61.8 ± 2.5	109.4 ± 2.4	—
	Non-A/C	259/75	96.7 ± 2.5	109.4 ± 2.4	—
	All Homes[d]	214/76	69.1 ± 2.6	91.0 ± 2.5	71.5 ± 2.4
NH_4^+	All Homes	314/155	154.7 ± 2.8	104.4 ± 2.3	—
H^+	A/C Homes	28/74	4.2 ± 4.3	82.5 ± 2.6	
	Non-A/C	230/74	11.2 ± 3.1	82.5 ± 2.6	
	All Homes[e]	163/75	9.1 ± 3.5	72.4 ± 2.9	18.4 ± 3.0
Uniontown					
SO_4^{2-}	All Homes[e]	91/46	87.8 ± 2.1	124.9 ± 1.9	110.3 ± 1.8
NH_4^+	All Homes[e]	91/44	157.2 ± 2.8	139.4 ± 2.1	167.0 ± 2.0
H^+	All Homes[e]	91/46	13.7 ± 2.5	76.6 ± 2.7	42.8 ± 2.2

[a]In/Out = Indoor sample site/outdoor sample site.
[b]GM ± GSD = Geometric mean ± geometric standard deviation.
[c]A/C Homes = Homes that had air-conditioning (A/C); this does not imply that it was on during the entire sampling period.
Non-A/C = Homes without air conditioning.
[d]The sample size (n) for the personal monitoring = 209.
[e]n = 174 for personal monitoring.

Source: Suh et al. (1992, 1993a,b).

were collected in the adjacent Lindon Elementary School. The infiltration factors, C_{ai}/C_{ao}, given by the slope of the regression lines (Table 5-17), were low (0.27 for sulfate and 0.12 for $PM_{2.5}$), possibly because of removal of particles in the air heating and ventilation system. The authors concluded that the data indicate that indoor $PM_{2.5}$ mass may not always be a good indicator of exposure to ambient combustion material caused by the influence of indoor sources of particles. However, ambient sulfate, SO_2, nitrate, soot, and total particulate number displayed strong correlations with indoor exposure. Ambient $PM_{2.5}$ mass was not a good indicator of indoor $PM_{2.5}$ mass exposure.



Figure 5-10. Personal versus outdoor $SO_4^=$ in State College, PA. Open circles represent children living in air conditioned homes; the solid line is the 1:1 line.

Source: Suh et al. (1993b).

Oglesby et al. (2000) conducted a study to evaluate the validity of fixed-site fine particle concentration measurements as exposure surrogates for air pollution epidemiology. Using 48-h EXPOLIS data from Basel, Switzerland, they investigated the personal exposure/outdoor concentration relationships for four indicator groups: (1) $PM_{2.5}$ mass, (2) sulfur and potassium for regional air pollution, (3) lead and bromine for traffic-related particles, and (4) calcium for crustal particles. The authors reported that personal exposures to $PM_{2.5}$ mass were not correlated

TABLE 5-17. STATISTICAL CORRELATION OF OUTDOOR (x) VERSUS INDOOR (y) CONCENTRATION FOR MEASURED SPECIES (Units are nmol m^{-3}, except for soot and metals, which are $\mu g/m^3$ and absorption units m^{-3}, respectively.)[a]

Species	Slope	Intercept	r^2	Average Outdoors
SO_2 All Samples	0.0272 ± 0.0023	0.34 ± 0.13	0.73	38
SO_2 Day Samples	0.0233 ± 0.0037	0.75 ± 0.26	0.62	56
SO_2 Night Samples	0.0297 ± 0.0029	0.099 ± 0.075	0.82	20
Sulfate All Samples	0.267 ± 0.024	−0.14 ± 0.48	0.70	16
Sulfate Day Samples	0.261 ± 0.034	0.40 ± 0.66	0.71	16
Sulfate Night Samples	0.282 ± 0.035	−0.84 ± 0.68	0.70	16
Nitrate All Samples	0.0639 ± 0.0096	0.9 ± 1.5	0.54	134
Nitrate Day Samples	0.097 ± 0.0096	−0.4 ± 1.4	0.88	126
Nitrate Night Samples	0.047 ± 0.011	1.5 ± 1.8	0.44	139
Soot Day Samples	0.43 ± 0.25	3.5 ± 1.7	0.43	6
Soot Night Samples	0.33 ± 0.13	0.00 ± 0.55	0.69	4
Total Acidity All Samples	0.04 ± 0.73	0.42 ± 0.23	0.00	0.2
Metals All Samples	0.10 ± 0.30	0.0014 ± 0.0042	0.01	0.0042

[a]Lindon Elementary School, Lindon, UT, January and February 1997.

Source: Patterson and Eatough (2000).

to corresponding home outdoor levels (n = 44, r = 0.07). In the study group reporting neither relevant indoor sources nor relevant activities, personal exposures and home outdoor levels of sulfur were highly correlated (n = 40, r = 0.85). These results are consistent with spatially homogeneous regional pollution and higher spatial variability of traffic and crustal materials.

PAHs have been measured in studies by EPA and the California Air Resources Board. PAH results from a probability sample of 125 homes in Riverside are discussed in reports by Sheldon et al. (1992a,b) and Özkaynak et al. (1996b). Data for two sequential 12-h samples were reported for PAHs by ring size (3 to 7) and for individual phthalates. The results are summarized below.

- The particulate-phase 5- to 7-ring species had lower relative concentrations than the more volatile 3- to 4-ring species.

- The 12-h indoor/outdoor ratios for the 5- to 7-ring species ranged from 1.1 to 1.4 during the day and from 0.64 to 0.85 during the night (Sheldon et al., 1993a).
- An indoor air model used to calculate indoor "source strengths" for the PAHs showed that smoking had the strongest effect on indoor concentrations.

Results from a larger PAH probability study in 280 homes in Placerville and Roseville (Sheldon et al., 1993a,b) were similar to the 125-home study. The higher-ring, particle-bound PAH's had lower indoor and outdoor concentrations than the lower-ring species. For most PAHs, the I/O ratio was greater than 1 for smoking and smoking/fireplace homes and less than 1 for fireplace-only, wood stove, wood stove/gas heat, gas heat, and "no source" homes.

A study of PAHs in indoor and outdoor air was conducted in 14 inner-city and 10 rural low-income homes near Durham, NC, in two seasons (winter and summer) in 1995 (Chuang et al., 1999). Fine-particle-bound PAH concentrations measured with a real-time monitor were usually higher indoors than outdoors (2.47 ± 1.90 versus 0.53 ± 0.58 $\mu g/m^3$). Higher indoor levels were seen in smoker's homes compared with nonsmoker's homes, and higher outdoor and indoor PAH levels were seen in urban areas compared with rural areas.

In a study reported by Dubowsky et al. (1999), the weekday indoor PAH concentrations attributable to traffic (indoor source contributions were removed) were 39 ± 25 ng/m^3 in a dormitory that had a high air exchange rate because of open windows and doors, 26 ± 25 ng/m^3 in an apartment, and 9 ± 6 ng/m^3 in a suburban home. The study showed that both outdoor—especially motor vehicular traffic—and indoor sources contributed to indoor PAH concentrations. BaP concentrations were measured in the THEES study (Waldman et al., 1991). A comprehensive analysis of the data showed considerable seasonal variability of indoor and outdoor sources and resultant changes in personal exposures to BaP.

5.5.4 Factors Affecting Correlations Between Ambient Measurements and Personal or Microenvironmental Measurements of Particulate Matter Constituents

The primary factors affecting correlations between personal exposure and ambient air PM measurements have been discussed in Section 4.3.2. These include air-exchange rates, particle penetration factors, decay rates and removal mechanisms, indoor air chemistry, indoor sources, and freshly-generated particles indoors. The importance of these factors varies for different PM

constituents. For acid aerosols, indoor air chemistry is particularly important as indicated by the discussion of the neutralization of the acidity by ammonia, which is present at higher concentrations indoors because of the presence of indoor sources. For SVOCs, including PAHs and phthalates, the presence of indoor sources will impact substantially the correlation between indoor and ambient concentrations (Özkaynak et al., 1996b; Sheldon et al., 1993b). Penetration factors for PM will impact correlations between indoors and outdoors for most elements, except Pb, which may have significant indoor sources in older homes. Indoor air chemistry, decay rates, and removal mechanisms may affect soot and organic carbon. Furthermore, reactions between indoor and outdoor gases and particles may also produce freshly generated aerosols indoors. These factors must be fully evaluated when attempting to correlate ambient, personal, and indoor PM concentrations.

5.5.5 Limitations of Available Data

The previous discussion demonstrates that there is very limited data available that can be used to compare personal, microenvironmental, and ambient air concentrations of PM constituents. Because of resource limitations, PM constituents have not been measured in many studies of PM exposure. There are little data on freshly generated aerosols indoors. Although there is some data on acid aerosols, the comparisons between the personal and indoor data generally have been with outdoor measurements at the participant's residences, not with community ambient air measurement sites. The relationship between personal exposure and indoor levels of acid aerosols is not clear because of the limited database. The exception is sulfate, for which there appears to be a strong correlation between indoor and ambient concentrations.

With the exception of PAHs, there are practically no data available to relate personal or indoor concentrations with outdoor or ambient site concentrations of SVOCs, which may be generated from a variety of combustion and industrial sources. The relationship between exposure and ambient concentrations of particles from specific sources, such as diesel engines, has not been determined.

Although there is an increasing amount of research being performed to measure PM constituents in different PM size fractions, the current data are inadequate to adequately assess the relationship between indoor and ambient concentrations of most PM constituents. Another

area where additional information has to be developed is the PM exposures that are derived from outdoor vapors (ov) reacting (rxn) with indoor vapors (iv). This is a source that could also vary with outdoor PM, for example, when the (ov) is ozone.

5.6 IMPLICATIONS OF USING AMBIENT PARTICULATE MATTER CONCENTRATIONS IN EPIDEMIOLOGIC STUDIES OF PARTICULATE MATTER HEALTH EFFECTS

In this section, the exposure issues that relate to the interpretation of the findings from epidemiologic studies of PM health effects are examined. This section examines the errors that may be associated with using ambient PM concentrations in epidemiologic analyses of PM health effects. First, implications of associations found between personal exposure and ambient PM concentrations are reviewed. This is discussed separately in the context of either community time-series studies or long-term, cross-sectional studies of chronic effects. Next, the role of compositional and spatial differences in PM concentrations are discussed and how these may influence the interpretation of findings from PM epidemiology. Finally, using statistical methods, an evaluation of the influence of exposure measurement errors on PM epidemiology studies is presented.

5.6.1 Potential Sources of Error Resulting from Using Ambient Particulate Matter Concentrations in Epidemiologic Analyses

Measurement studies of personal exposures to PM are still few and limited in spatial, temporal, and demographic coverage. Consequently, with the exception of a few longitudinal panel studies, most epidemiologic studies of PM health effects rely on ambient community monitoring data giving 24-h average PM concentration measurements. Moreover, because of limited sampling for $PM_{2.5}$, many of these epidemiologic studies had to use available PM_{10} or in some instances had to rely on historic data on other PM measures or indicators, such as TSP, $SO_4^{=}$, IP_{15}, RSP, COH, KM, etc. A critical question often raised in the interpretation of results from acute or chronic epidemiologic community-based studies of PM is whether the use of ambient stationary site PM concentration data influences or biases the findings from these studies. Because the health outcomes are measured on individuals, the epidemiologists might

prefer to use personal exposure measurements (total, ambient, or nonambient) instead of surrogates, such as ambient PM concentration measurements collected at one or more ambient monitoring sites in the community. Use of ambient concentrations could lead to misclassification of individual exposures and to errors in the epidemiologic analysis of pollution and health data depending on the pollutant and on the mobility and lifestyles of the population studied. Ambient monitoring stations can be some distance away from the individuals and can represent only a fraction of all likely outdoor microenvironments that individuals come in contact with during the course of their daily lives. Furthermore, most individuals are quite mobile and move through multiple microenvironments (e.g., home, school, office, commuting, shopping, etc.) and engage in diverse personal activities at home (e.g, cooking, gardening, cleaning, smoking). Some of these microenvironments and activities may have different sources of PM and result in distinctly different concentrations of PM than that monitored by the fixed-site ambient monitors. Consequently, exposures of some individuals will be classified incorrectly if only ambient monitoring data are used to estimate individual level exposures to PM. Thus, bias or loss of precision in the epidemiologic analysis may result from improper assessment of exposures using data routinely collected by the neighborhood monitoring stations.

Because individuals are exposed to particles in a multitude of indoor and outdoor microenvironments during the course of a day, concern over error introduced in the estimation of PM risk coefficients using ambient, as opposed to personal, PM measurements has received considerable attention recently from exposure analysts, epidemiologists, and biostatisticians. Some exposure analysts contend that, for community time-series epidemiology to yield information on the statistical association of a pollutant with a health response, there must be an association between personal exposure to a pollutant and the ambient concentration of that pollutant because people tend to spend around 90% time indoors and are exposed to both indoor and outdoor-generated PM (cf. Wallace, 2000b; Brown and Paxton, 1998; Ebelt et al., 2000). Consequently, numerous findings reported in the epidemiologic literature on significant associations between ambient PM concentrations and various morbidity and mortality health indices, in spite of the low correlations between ambient PM and concentrations and measures of personal exposure, has been described by some exposure analysts as an exposure paradox (Lachenmyer and Hidy, 2000, Wilson et al., 2000).

To resolve the so-called exposure paradox, several types of analyses need to be considered. The first type of analysis has to examine the correlations between ambient PM concentrations and personal exposures that are relevant to most of the existing PM epidemiology studies using either pooled, daily-average, or longitudinal exposure data. The second approach has to study the degree of correlations between the two key components of personal PM exposures (i.e., exposures caused by ambient-generated PM and exposures caused by nonambient PM) with ambient or outdoor PM concentrations, for each of the three types of exposure study design. Yet, even with these two approaches, it may still be difficult to examine complex synergisms which, in some situations, may preclude simple decoupling of in indoor and outdoor particles either in terms of exposure or total dose delivered to the lung. In addition, several factors influencing either the exposure or health response characterization of the subjects have to be addressed. These include such factors as:

- spatial variability of PM components,
- health or sensitivity status of subjects,
- variations of PM with other co-pollutants,
- co-generation of fine and ultrafine particles from outdoor air and indoor gaseous pollutants,
- formal evaluation of exposure errors in the analysis of health data, and
- how the results may depend on the variations in the design of the epidemiologic study.

To facilitate the discussion of these topics, a brief review of concepts pertinent to exposure analysis issues in epidemiology is presented.

5.6.2 Associations Between Personal Exposures and Ambient Particulate Matter Concentrations

As defined earlier in Sections 5.3 and 5.4, personal exposures to PM result from an individual's exposures to PM in many different types of microenvironments (e.g., outdoors near home, outdoors away from home, indoors at home, indoors at office or school, commuting, restaurants, malls, other public places, etc.). Total personal exposures (E_t) that occur in these indoor and outdoor microenvironments can be classified as those resulting from PM of outdoor origin (E_{ag}) and those primarily generated by indoor sources and personal activities (E_{nonag} = E_{pig}+E_{pact}). The associations between personal exposures and ambient PM concentrations that

have been reported from various personal exposure monitoring studies under three broad categories of study design: (1) longitudinal, (2) daily-average, or (3) pooled exposure studies are summarized below.

In the previous Sections 5.4.3.1.2 and 5.4.3.1.3, some recent studies mainly conducted in the United States, and involving children, the elderly, and subjects with COPD were reviewed, and they indicated that both intra- and interindividual variability in the relationships between personal exposures and ambient PM concentrations were observed. A variety of different physical, chemical, and personal or behavioral factors were identified by the original investigators that seem to influence the magnitude and the strength of the associations reported.

Clearly, for cohort studies in which individual daily health response are obtained, individual longitudinal PM personal exposure data (including ambient-generated and nonambient components) provide the appropriate indicators. In this case, health responses of each individual can be associated with the total personal exposure, the ambient-generated exposure, or the nonambient exposure of each individual. Also, the relationships of personal exposure indicators with ambient concentration can be investigated. In the case of community time-series epidemiology, however, it is not feasible to obtain experimental measurements of personal exposure for the millions of people over time periods of years that are needed to investigate the relationship between air pollution and infrequent health responses such as deaths or even hospital admissions. The epidemiologist must work with the aggregate number of health responses occurring each day and a measure of the ambient concentration that is presumed to be representative of the entire community. The relationship of PM exposures of the potentially susceptible groups to monitored ambient PM concentrations depends on their activity pattern and level, residential building and HVAC factors (which influence the infiltration factor), status of exposure to ETS, amount of cooking or cleaning indoors, and seasonal factors, among others. Average personal exposures of these special subgroups to ambient-generated PM are correlated well with ambient PM concentrations regardless of individual variation in the absence of major microenvironmental sources.

Even though both E_{ag} and E_{nonag} contributes to daily baseline PM dose received by the lung, there seem to be clear differences in the relationships of ambient (E_{ag}) and nonambient (E_{nonag}) exposure with ambient concentration (C_a). Various researchers have shown that E_{nonag} is independent of C_a, but that E_{ag} is a function of C_a. Wilson et al. (2000) explains the difference

based on different temporal patterns that affect PM concentrations. "Concentrations of ambient PM are driven by meteorology and by changes in the emission rates and locations of emission sources, while concentrations of nonambient PM are driven by the daily activities of people." Still, although E_{nonag} may not correlate with ambient C_a or E_{ag}, it will nevertheless add to the daily baseline dose received by the lung.

Ott et al. (2000) also discuss the reasons for assuming that E_{nonag} is independent of E_{ag} and C_a. They show that the nonambient component of total personal exposure is uncorrelated with the outdoor concentration data. Ott et al. (2000) show the $\overline{E}_{nonag}$ is similar for three population-based exposure studies, including two large probability-based studies, the PTEAM study conducted in Riverside (Clayton et al., 1993; Thomas et al., 1993; Özkaynak et al., 1996a,b) and a study in Toronto (Pelizzarri et al., 1999; Clayton et al., 1999a), as well as a nonprobability-based study, conducted in Phillipsburg (Lioy et al., 1990). Based on these three studies, they conclude that $\overline{E}_{nonag}$ and the distribution of$(E_{nonag})_{it}$ can be treated as constant from city to city, where i refers to a specific individual and t to a specific day..

Dominici et al. (2000) examined a larger database consisting of five different PM exposure studies and concluded that $\overline{E}_{nonag}$ can be treated as relatively constant from city to city. If $(E_{nonag})_t$ were constant, this would imply that it would have a zero correlation with $(C_a)_t$. However, this hypothesis of constant $(E_{nonag})_{it}$ has not been established fully because only a few studies have obtained the data needed to estimate $(E_{nonag})_{it}$. Although $\overline{E}_{nonag}$ is independent of C_a , it may not be independent of α. Sarnat et al. (2000) show that $\overline{E}_{nonag}$ goes up as the ventilation rate (and α) goes down. Lachenmeyer and Hidy (2000) also show, by comparing winter and summer regression equations, that as the slope (α) goes down, the intercept $(\overline{E}_{nonag})$ goes up.

Mage et al. (1999) assume that the PM_{10} concentration component from indoor sources (such as smoking, cooking, cleaning, burning candles, and so on) is not correlated with the outdoor concentration. They indicate that this lack of correlation is expected, because people are unaware of ambient concentrations and do not necessarily change their smoking or cooking activities as outdoor PM_{10} concentrations vary, an assumption supported by other empirical analyses of personal exposure data. For the PTEAM data set, Mage et al. (1999) have shown that E_{pig} and C_a have r near zero ($R^2 = 0.005$). Wilson et al. (2000) have shown the C_{ai} and C_{pig} also

have r near zero ($R^2 = 0.03$). Figure 5-11 shows the relationship of estimated $(E_{nonag})_{it}$ and E_{nonag} with C_a (calculated by EPA from PTEAM and THEES data).



Figure 5-11. Plots of nonambient exposure to PM_{10}, (a) daytime individual values from PTEAM data and (b) daily-average values from THEES data.

Source: Data taken from (a) Clayton et al. (1993) and (b) Lioy et al. (1990).

Based on these results it is reasonable to assume that ordinarily E_{nonag} has no relationship with C_a in the absence of sufficient study-specific data on the complex interactions between indoor and outdoor gases and particles producing fresh particles indoors. Therefore, in linear nonthreshold models of PM health effects, E_{nonag} is not expected to contribute to the relative risk determined in a regression of health responses on C_a. Furthermore, in time-series analysis of pooled or daily heath data, it is expected that E_{ag} rather than E_t will have the stronger association with C_a.

5.6.3 Role of Compositional Differences in Exposure Characterization for Epidemiology

The majority of the available data on PM exposures and relationships with ambient PM have come from a few large-scale studies, such as PTEAM, or longitudinal studies on selected populations, mostly the elderly. Consequently, for most analyses, exposure scientists and statisticians had to rely on PM_{10} or $PM_{2.5}$ mass data, instead of elemental or chemical compositional information on individual or microenvironmental samples. In a few cases, researchers have examined the factors influencing indoor outdoor ratios or penetration and deposition coefficients using elemental mass data on personal, indoor, and outdoor PM data (e.g., Özkaynak et al. 1996a,b; Yakovleva et al. 1999). These results have been informative in terms of understanding relative infiltration of different classes of particle sizes and sources into residences (e.g., fossil fuel combustion, mobile source emissions, soil-derived, etc.). Clearly, in the accumulation-mode, particles associated with stationary or mobile combustion sources have greater potential for penetration into homes and other microenvironments than do crustal material. The chemical composition of even these broad categories of source classes may have distinct composition and relative toxicity. Moreover, when particles and reactive gases are present indoors in the presence of other pollutants or household chemicals, they may react to form additional or different compounds and particles with yet unknown physical, chemical, and toxic composition (Wainman et al. 2000). Thus, if indoor-generated and outdoor-generated PM were responsible for different types of health effects, or had significantly different toxicities on a per unit mass basis, it would be then be important that E_{ag} and E_{nonag} should be separated and treated as different species, much like the current separation of PM_{10} into $PM_{2.5}$ and $PM_{10\text{-}2.5}$. These complexities in personal exposure profiles may introduce nonlinearities and other

statistical challenges in the selection and fitting of concentration-response models. Unfortunately, PM health effects models have not yet been able to meaningfully consider such complexities. The relationships of toxicity to the chemical and physical properties of PM are discussed in Chapter 7.

It is important also to note that individuals spend time in places other than their homes and outdoors. Many of the interpretations reported in the published literature on factors influencing personal PM_{10} exposures, as well as in this chapter, come from the PTEAM study. The PTEAM study was conducted 10 years ago in one geographic location in California, during one season, and most residences had very high and relatively uniform air-exchange rates. Nonhome indoor microenvironments were not monitored directly during the PTEAM study. Commuting exposures from traffic or exposures in a variety of different public places or office buildings could not be assessed directly. Nonresidential buildings may have lower or higher ambient infiltration rates depending on the use and type of the mechanical ventilation systems employed. Because the source and chemical composition of particulate matter effecting personal exposures in different microenvironments vary by season, day-of-the-week, and time of day, it is likely that some degree of misclassification of exposures to PM toxic agents of concern will be introduced when health effects models use only daily-average mass measures such as PM_{10} or $PM_{2.5}$. Because of the paucity of currently available data on many of these factors, it is impossible to ascertain at this point the magnitude and severity of these more complex exposure missclassification problems in the interpretation of results from PM epidemiology.

5.6.4 Role of Spatial Variability in Exposure Characterization for Epidemiology

Chapter 3 (Section 3.2.3) and Chapter 5 (Section 5.3) present information on the spatial variability of PM mass and chemical components at fixed-site ambient monitors; for purposes of this chapter, this spatial variability is called an “ambient gradient.” Any gradient that may exist between a fixed-site monitor and the outdoor microenvironments near where people live, work, and play, obviously affects the concentration profile actually experienced by people as they go about their daily lives.

However, the evidence so far indicates that PM concentrations, especially fine PM (mass and sulfate), generally are distributed uniformly in most metropolitan areas. This reduces the

potential for exposure misclassification because of outdoor spatial gradients when a limited number of ambient PM monitors are used to represent population average ambient exposures in time-series or cross-sectional epidemiologic studies of PM. This topic is further discussed below in Section 5.6.5. However, as discussed earlier, the same assumption is not necessarily true for different components of PM, because source-specific and other spatially nonuniform pollutant emissions could alter the spatial profile of individual PM components in a community. For example, particulate and gaseous pollutants emitted from motor vehicles tend to be higher near roadways and inside cars. Likewise, acidic and organic PM species may be location- and time-dependent. Furthermore, human activities are complex, and if outdoor PM constituent concentration profiles are either spatially or temporally variable, it is likely that exposure misclassification errors could be introduced in the analysis of PM air pollution and health data.

5.6.5 Analysis of Exposure Measurement Error Issues in Particulate Matter Epidemiology

The effects of exposure misclassification on relative risk estimates of disease using classical 2×2 contingency design (i.e., exposed/nonexposed versus diseased/nondiseased) have been studied extensively in the epidemiologic literature. It has been shown that the magnitude of the exposure-disease association (e.g., relative risk) because of either misclassification of exposure or disease alone (i.e., nondifferential misclassification) biases the effect results toward the null, and differential misclassification (i.e, different magnitudes of disease misclassification in exposed and nonexposed populations) can bias the effect measure toward or away from the null value relative to the true measure of association (Shy et al., 1978; Gladen and Rogan, 1979; Copeland et al., 1977; Özkaynak et al., 1986). However, the extension of these results from contingency analysis design to multivariate (e.g., log-linear regression, Poissson, logit) models typically used in recent PM epidemiology has been more complicated. Recently, researchers have developed a framework for analyzing measurement errors typically encountered in the analysis of time-series mortality and morbidity effects from exposures to ambient PM (cf. Zeger et al., 2000; Dominici et al., 2000; Samet et al., 2000). Some analysis in the context of cross-sectional epidemiology have also been conducted (e.g. Navidi et al., 1999).

The appropriateness of using ambient PM concentration as an exposure metric in the context of epidemiologic analysis of health effects associated with exposure to PM recently has

been examined by a number of investigators (cf. Zeger et al., 2000; Dominici et al., 2000; Navidi et al., 1999; Özkaynak and Spengler, 1996). In the following section, the error analysis model framework developed in Zeger et al. (2000) will be discussed in the context of time-series epidemiology. After which, issues and implications of exposure errors to findings from long-term/chronic or cross-sectional epidemiology will be discussed briefly.

5.6.5.1 Analysis of Exposure Measurement Errors in Time-Series Studies

The discussion presented in this section is further examined in Chapter 8 under the context of implications of exposure errors to results and interpretation of findings from PM epidemiology. The discussion presented in this section also focuses more on the potential for exposure misclassification biases on the estimated regression slopes rather than on the more subtle issues, such as those dealing with "effect modification" discussed further in Chapter 8.

Zeger et al. (2000) provide a useful framework for analyzing exposure error in community time-series epidemiology. This framework, coupled with results from recent exposure studies, makes it possible to clarify some important questions regarding relationships among the three aspects of personal exposure (1) total personal, (2) personal caused by ambient PM, and (3) personal resulting from nonambient PM and ambient concentration. Consider the regression of a health response (i.e., mortality rate on day t, Y_t, against the ambient concentration of PM on day t, C_t). In analyzing pollution-level data on mortality and air pollution, log-linear regressions of the form:

$$Y_t = \exp\left[s(t) + C_t\beta_c + u_t\beta_u\right] \tag{5-12}$$

are fit, where Y_t is the expected mortality rate; *s(t)* is an arbitrary but smooth function of time, introduced to control for the confounding of longer trends and seasonality; C_t, is the average of multiple monitor measurements of ambient pollution measurement for day *t*; and u_t are other possible confounders such as temperature and dew point on the same or previous day. Each coefficient, β, in Equation 5-12 gives the expected change in the health response, Y, because of a unit change in its corresponding variable.

However, instead of Equation 5-12, Zeger et al. (2000) suggest that the analyst would like to know the corresponding relationship for personal exposure rather than ambient concentration,

$$Y_t = \exp[s(t) + E_t\beta_E + u_t\beta_u]. \tag{5-13}$$

Zeger et al. (2000) do not differentiate among the three aspects of personal or community exposure. To understand the error in β caused by using ambient concentrations instead of personal exposure in the regression analysis, it is necessary to examine the relationship between β_c, based on a unit change in the ambient concentration, C, and β_E, based on a unit change in one of the three aspects of personal exposure, E. In considering the consequences for β_c, as an estimate of β_E, of having a measure of ambient pollution C_t, rather than actual personal exposure E_{it}, it is convenient to express the desired pollution measurement, E_{it}, as C_t plus three error terms:

$$E_{it} = C_t + (E_{it} - \overline{E}_t) + (\overline{E}_t - C_t^*) + (C_t^* - C_t). \tag{5-14}$$

Here $\overline{E}_t$ represents the daily, community-average personal exposure. The first term, $(E_{it} - \overline{E}_t)$, is the error resulting from having only aggregated or community-averaged exposure rather than individual-level exposure data. The second term, $(E_t - C_t^*)$, is the difference between the average personal exposure and the true ambient pollutant level, and the third term, $(C^*_t - C_t)$, represents the difference between the true and the measured ambient concentration.

In the evaluation of these error terms, two types of measurement error often are considered in the context of epidemiology. The classical error model assumes that measurement error, $(C_t\text{-}E_t)$, depends on ambient measurements [simply referred to as C_t here instead of $(C_a)_t$]. The Berkson error model assumes that the measurement error is dependent on the true value or the personal exposure (E_t). The regression coefficient (β_C), estimated from the health effects model in the Berkson error case, gives an unbiased estimate of β_E. In the classical error case, β_C is a biased estimate of β_E, and the degree of bias depends on the correlation between the measurement error and C_t. The measurement error analysis of Zeger et al. (2000) includes three components: (1) an individual's deviation from the risk-weighted average personal exposure;

(2) the difference between the average personal exposure and the true ambient level; and (3) the difference between the measured and the true ambient levels, which include the spatial variation of outdoor PM and instrument sampling error. Zeger et al. (2000) conclude that the first and third components are of the Berkson type and, therefore, are likely to have smaller effects on the relative risk estimates for PM. However, the second component can be a source of substantial bias if, for example, there are short-term associations of the contributions of indoor sources with ambient concentrations. Recent analysis of PTEAM data (Mage et al., 1999; Wilson et al., 2000) and theoretical considerations (Ott et al., 2000) indicate that it is unlikely that nonambient exposures will be correlated with the ambient concentration (even though total lung dose will be influenced both by ambient and nonambient PM sources and concentrations). Therefore, this type of bias is unlikely. However, if the community average exposure to ambient PM is less than the ambient concentration, the risk regression coefficient, β_C, will be biased low. According to Carroll et al. (1995), $\beta_C = \alpha\ \beta_E$, where β_C is the percentage increase in risk because of a unit increase in ambient concentration, and β_E is the estimated percentage increase in risk because of a unit increase in the community-average personal exposure to ambient PM. Both Zeger et al. (2000) and Dominici et al. (2000) examine the nature of error with this second component. Both of these analyses conclude that the error introduced because of measured differences between the average personal exposure and ambient levels can bias the regression coefficients. In both cases they find the β_C is close to $\alpha\ \beta_E$.

This framework analysis demonstrates the importance of the daily community-average exposure, $\overline{E}_t$, in community time-series epidemiology. It is $\overline{E}_t$, not the random, pooled values of $E_{i,t}$, that need to have a statistically significant correlation with C_t for proper interpretation of community time-series epidemiology studies based on ambient monitoring data, as discussed further in Wilson et al. (2000) and Mage et al. (1999).

A critical assumption in the above analysis is that the risk varies linearly with C or E (i.e., β_c and β_E are constant). This assumption does not permit a threshold (a concentration below which there is no effect). It also includes the assumption that the appropriate metric for determination of a health response is the 24-h average PM mass concentration. Zeger et al. (2000) show that the likely consequence of using ambient concentrations instead of the risk-weighted average personal exposure measures is to underestimate the pollution effects. According to Zeger et al. (2000) the largest biases in inferences about the mortality-personal

exposure relative risk will occur because of more complex errors between ambient concentration and daily-average personal exposure measures. It is important to note that both the Zeger et al. (2000) and the Dominici et al. (2000) error analyses used personal PM_{10} data from the PTEAM study data. However, effects of measurement error estimates may differ by particle size and composition. It is possible that $PM_{2.5}$, ultrafine particle measures, or another component of PM, may better reflect personal exposures to PM of outdoor origin. Finally, the seasonal or temporal variations in the measurement errors and correlations between different PM concentration measures and gaseous co-pollutants (e.g. SO_2, CO, NO_2, O_3) could influence the error analysis results reported by the investigators cited above.

5.6.5.2 Analysis of Exposure Measurement Errors in Long-Term Epidemiology Studies

The Six Cities (Dockery et al., 1993) and ACS (Pope et al., 1995) studies have played an important role in assessing the health effects from long-term exposures to particulate pollution. Even though these studies often have been considered as chronic epidemiologic studies, it is not easy to differentiate the role of historic exposures from those of recent exposures on chronic disease mortality. In the Six Cities study, fine particles and sulfates were measured at the community level, and the final analysis of the database used six city-wide average ambient concentration measurements. This limitation also applies to the ACS study but has less impact because of the larger number of cities considered in that study. In a HEI-sponsored reanalysis of the Six Cities and the ACS data sets, Krewski et al. (2000) attempted to examine some of the exposure misclassification issues either analytically or through sensitivity analysis of the aerometric and health data. The HEI reanalysis project also addressed exposure measurement error issues related to the Six Cities study. For example, the inability to account for exposures prior to the enrollment of the cohort hampered accurate interpretation of the relative risk estimates in terms of acute versus chronic causes. Although the results seem to suggest past exposures are more strongly associated with mortality than recent exposures, the measurement error for long-term averages could be higher, thus influencing these interpretations. For example, Krewski et al. (2000), using the individual mobility data available for the Six Cities cohort, analyzed the mover and nonmover groups separately. The relative risk of fine particle effects on all-cause mortality was shown to be higher for the nonmover group than for the mover group, suggesting the possibility of higher exposure misclassification biases for the movers. The issue

of using selected ambient monitors in the epidemiologic analyses also was investigated by the ACS and Six Cities studies reanalysis team. Krewski et al.(2000) presented the sensitivity of results to choices made in selecting stationary or mobile-source-oriented monitors. For the ACS study, reanalysis of the sulfate data using only those monitors designated as residential or urban, and excluding sites designated as industrial, agricultural, or mobile did not change the risk estimates appreciably. On the other hand, application of spatial analytic methods designed to control confounding at larger geographic scales (i.e., between cities) caused changes in the particle and sulfate risk coefficients. Spatial adjustment may account for differences in pollution mix or PM composition, but many other cohort-dependent risk factors will vary across regions or cities in the United States. Therefore, it is difficult to interpret these findings solely in terms of spatial differences in pollution composition or relative PM toxicity until further research is concluded.

Another study that has examined the influence of measurement errors in air pollution exposure and health effects assessments is the one reported by Navidi et al. (1999). This study developed techniques to incorporate exposure measurement errors encountered in long-term air pollution health effects studies and tested them on the data from the University of Southern California Children's Health Study conducted in 12 communities in California. These investigators developed separate error analysis models for direct (i.e., personal sampling) and indirect (i.e., microenvironmental) personal exposure assessment methods. These models were generic to most air pollutants, but a specific application was performed using a simulated data set for studying ozone health effects on lung function decline in children. Because the assumptions made in their microenvironmental simulation modeling framework were similar to those made in estimating personal PM exposures, it is useful to consider the conclusions from Navidi et. al. (1999). According to Navidi et al. (1999), neither the microenvironmental nor the personal sampler method produces reliable estimates of the exposure-response slope (for O_3) when measurement error is uncorrected. Because of nondifferential measurement error, the bias was toward zero under the assumptions made in Navidi et al. (1999) but could be away from zero if the measurement error was correlated with the health response. A simulation analysis indicated that the standard error of the estimate of a health effect increases as the errors in exposure assessment increase (Navidi et al., 1999). According to Navidi et al. (1999), when a fraction of the ambient level in a microenvironment is estimated with a standard error of 30%, the standard

error of the estimate is 50% higher than it would be if the true exposures were known. It appears that errors in estimating ambient PM indoor/ambient PM outdoor ratios have much more influence on the accuracy of the microenvironmental approach than do errors in estimating time spent in these microenvironments.

5.6.5.3 Conclusions from Analysis of Exposure Measurement Errors on Particulate Matter Epidemiology

Personal exposures to PM are influenced by a number of factors and sources of PM located in both indoor and outdoor microenvironments. However, PM resulting from ambient sources does penetrate into indoor environments, such as residences, offices, public buildings, etc., in which individuals spend a large portion of their daily lives. The correlations between total personal exposures and ambient or outdoor PM concentrations can vary depending on the relative contributions of indoor PM sources to total personal exposures. Panel studies of both adult and young subjects have shown that, in fact, individual correlations of personal exposures with ambient PM concentrations could vary person to person, and even day to day, depending on the specific activities of each person. Separation of PM exposures into two components, ambient-generated PM and nonambient PM, would reduce uncertainties in the analysis and interpretation of PM health effects data. Nevertheless, because ambient-generated PM is an integral component of total personal exposures to PM, statistical analysis of cohort-average exposures are strongly correlated with ambient PM concentrations when the size of the underlying population studied is large. Using the PTEAM study data, analysis of exposure measurement errors, in the context of time-series epidemiology, also has shown that errors or uncertainties introduced by using surrogate exposure variables, such as ambient PM concentrations, could lead to biases in the estimation of health risk coefficients. These then would need to be corrected by suitable calibration of the PM health risk coefficients. Correlations between the PM exposure variables and other covariates (e.g., gaseous co-pollutants, weather variables, etc.) also could influence the degree of bias in the estimated PM regression coefficients. However, most time-series regression models employ seasonal or temporal detrending of the variables, thus reducing the magnitude of this cross-correlation problem (Özkaynak and Spengler 1996).

Ordinarily, exposure measurement errors are not expected to influence the interpretation of findings from either the cross-sectional or time-series epidemiologic studies that have used ambient concentration data if they include sufficient adjustments for seasonality and key confounders. Clearly, there is no question that better estimates of exposures to components of PM of health concern are beneficial. Composition of PM may vary in different geographic locations and different exposure microenvironments. Compositional and spatial variations could lead to further errors in using ambient PM measures as surrogates for exposures to PM. Even though the spatial variability of PM ($PM_{2.5}$ in particular) mass concentrations in urban environments seems to be small, the same conclusions drawn above regarding the influence of measurement errors may not necessarily hold for all of the PM toxic components. Again, the expectation based on statistical modeling considerations is that these exposure measurement errors or uncertainties will most likely reduce the statistical power of the PM health effects analysis, making it difficult to detect a true underlying association between the correct exposure metric and the health outcome studied. However, until more data on exposures to toxic agents of PM become available, existing studies on PM exposure measurement errors must be relied on; thus, at this time, the working hypothesis is that the use of ambient PM concentrations as a surrogate for exposures is not expected to change the principal conclusions from PM epidemiologic studies, utilizing community average health and pollution data.

5.7 SUMMARY OF KEY FINDINGS AND LIMITATIONS

Exposure Definitions and Components

- Personal exposure (E) to PM mass or its constituents results when individuals come in contact with particulate pollutant concentrations (C) in locations or microenvironments **(μe)** that they frequent during a specific period of time. Various PM exposure metrics can be defined according to its source (i.e., ambient, nonambient) and the microenvironment where exposure occurs.
- Personal exposure to PM results from an individual's exposure to PM in many different types of microenvironments (e.g., outdoors near home, outdoors away from home, indoors at home, indoors at office or school, commuting, restaurants, malls, other public places, etc.). Thus, total

daily exposure to PM for a single individual (E_t) can be expressed as the sum of various microenvironmental exposures that the person encounters during the course of a day.

- In a given μ**e**, particles may originate from a wide variety of sources. In an indoor microenvironment, PM may be generated from within as a result of PM generating activities (e.g., cooking, cleaning, smoking, resuspending PM from PM resulting from both indoor and outdoor sources that had settled out), from outside (outdoor PM entering through cracks and openings in the structure), and from the chemical interaction of pollutants from outdoor air with indoor-generated pollutants.
- The total daily exposure to PM for a single individual (E_t) also can be expressed as the sum of contributions of ambient-generated (E_{ag}) and nonambient-generated (E_{nonag}) PM (i.e., $E = E_{ag} + E_{nonag}$). E_{nonag}, in turn, is composed of PM generated by indoor sources (E_{pig}) and PM generated by personal activities (E_{pact}) (i.e., $E_{nonag} = E_{pig} + E_{pact}$). E_{ag} is composed of exposures to ambient PM concentrations while outdoors, $\sum_t C_a \Delta t_a$, and ambient PM that has infiltrated indoors, $\sum_t C_{a\,i} \Delta t_i$ while indoors (i.e., $E_{ag} = \sum_t C_a \Delta t_a + \sum_t C_{a\,i} \Delta t_i$). However, within a large population group, there will be distributions of E_t and its components (E_{ag}, E_{nonag}) due to variations in human activities and microenvironmental concentrations and sources each individual encounters.
- Exposure models are useful tools for examining the importance of sources, microenvironments, and physical and behavioral factors that influence personal exposures to PM. However, development and evaluation of population exposure models for PM and its components has been limited. Improved modeling methodologies and new model input data are needed.

Factors Affecting Concentrations and Exposures to Particulate Matter

- Concentrations of PM indoors are affected by several factors and mechanisms: ambient concentrations outdoors; air exchange rates; particle penetration factors; particle production from indoor sources and indoor air chemistry; and indoor particle decay rates and removal mechanisms caused by physical processes or resulting from mechanical filtration, ventilation or air-conditioning devices.
- Average personal exposures to PM mass and its constituents are influenced by microenvironmental PM concentrations and by how much time is spent by each individual in

these various indoor and outdoor microenvironments. Nationwide, individuals, on average, spend nearly 90% of their time indoors (at home and in other indoor locations) and about 6% of their time outdoors.

- Personal exposures are associated with both indoor as well as outdoor sources; the personal exposure/outdoor concentration ratios present substantial intra- and inter-personal variability; although this variability was originally thought to be mainly due to the presence of personal and microenvironmental sources, the results from recent exposure studies suggest that it is the varying impact of the outdoor particles on indoor environments that is mainly responsible for the observed intra- and inter-variability in exposure/outdoor concentration ratios
- Home characteristics may be the most important factor that effects the relationship between the average population exposures and ambient concentrations. Air exchange rate seems to be an important home characteristic surrogate that can explain a large fraction of the observe inter- and intra-personal variability. These findings explain why longitudinal studies (many repeated measurements per person) provide stronger correlations between personal exposure and outdoor concentrations than cross-sectional studies (few repeated measurements per individual).
- Since home characteristics is the most important factor affecting personal exposures, one would expect that correlations between average population exposures and outdoor concentrations will vary by season and geography.
- The relative size of personal exposure to ambient-generated PM relative to nonambient-generated PM depends on the ambient concentration, the infiltration rate of outdoor PM into indoor microenvironments, the amount of PM generated indoors (e.g., ETS, cooking and cleaning emissions), and the amount of PM generated by personal activity sources. Infiltration rates primarily depend on air-exchange rate, size-dependent particle penetration across the building membrane, and size-dependent removal rates. All of these factors vary over time and across subjects and building types.
- The relationship between PM exposure, dose, and health outcome could depend on the concentration, composition, and toxicity of the PM originating from different sources. Application of source apportionment techniques to indoor and outdoor $PM_{2.5}$ and personal, indoor, and outdoor PM_{10} composition data have identified the following general source

categories: outside soil, resuspended indoor soil, indoor soil, personal activities, sea-salt, motor vehicles, nonferrous metal smelters, and secondary sulfates.

- There have been only a limited number of studies that have measured the physical and chemical constituents of PM in personal or microenvironmental samples. Available data on PM constituents indicate that
  - S personal and indoor sulfate measurements often are correlated highly with outdoor and ambient sulfate concentration measurements;
  - S for acid aerosols, indoor air chemistry is particularly important because of the neutralization of the acidity by ammonia, which is present at higher concentrations indoors because of the presence of indoor sources of ammonia;
  - S for SVOCs, including PAHs and phthalates, the presence of indoor sources will substantially impact the relation between indoor and ambient concentrations;
  - S penetration and decay rates are a functions of size and will cause variations in the attenuation factors as a function of particle size; infiltration rates will be higher for PM_1 and $PM_{2.5}$ than for PM_{10}, $PM_{10\text{-}2.5}$ or ultrafine particles; and
  - S Indoor air chemistry may increase indoor concentrations of organic PM.
- Even though there is an increasing amount of research being performed to measure PM constituents in different PM size fractions, with few exceptions (i.e., sulfur or sulfates), the current data are inadequate to adequately assess the relationship between personal, indoor, and ambient concentrations of most PM constituents.

Correlations Between Personal Exposures, Indoor, Outdoor, and Ambient Measurements

- Most of the available personal data on PM measurements and information on the relationships between personal and ambient PM come from a few large-scale studies, such as the PTEAM study, or the longitudinal panel studies, which have been conducted on selected populations, such as the elderly.
- Panel and cohort studies that have measured PM exposures and concentrations typically have reported their results in terms of three types of correlations: (1) longitudinal, (2) pooled, and (3) daily-average correlations between personal and ambient or outdoor PM.
- The type of correlation analysis performed can have a substantial effect on the resulting correlation coefficient. Low correlations with ambient concentrations could result when people

with very different nonambient exposures are pooled, even though temporally, their individual personal exposures may be correlated highly with ambient concentrations.

- Recent studies conducted by EPA of the elderly subjects living in a retirement facility in Baltimore and a group of elderly living in Fresno produced higher correlation coefficients between personal and ambient PM for daily-average correlations compared to longitudinal correlations. This supports earlier analyses showing the daily-average correlations are higher than pooled correlations.
- Longitudinal and pooled correlations between personal exposure and ambient or outdoor PM concentrations reported by various investigators varied considerably among the different studies and in each study between the study subjects. Most studies report longitudinal correlation coefficients that range from close to zero to near one, indicating that individual's activities and residence type may have a significant effect on total personal exposures to PM.
- Longitudinal studies that measured sulfate found high correlations between personal and ambient sulfate.
- In general, probability-based population studies tend to show low pooled correlations because of the high differences in levels of nonambient PM generating activities from one subject to another. In contrast, the absence of indoor sources for the populations in several of the longitudinal panel studies resulted in high correlations between personal exposure and ambient PM within subjects over time for these populations. But even for these studies, correlations varied by individual depending on their activities and microenvironments that they occupied.

Potential Sources of Error Resulting from Using Ambient Particulate Matter Concentrations in Epidemiologic Analyses

- There is, as yet, no clear consensus among exposure analysts as to how well ambiently measured PM concentrations represent a surrogate for personal exposure to total PM or to ambient-generated PM.
- Measurement studies of personal exposures to PM are still few and limited in spatial, temporal, and demographic coverage. Consequently, with the exception of a few longitudinal panel studies, most epidemiologic studies on PM health effects have relied on daily-average PM concentration measurements obtained from ambient community monitoring data as a surrogate for the exposure variable.

- Because individuals are exposed to particles in a multitude of indoor and outdoor microenvironments during the course of a day, concerns about error introduced in the estimation of PM risk coefficients using ambient, as opposed to personal PM measurements, have been raised.
- Total personal exposures to PM could vary from person to person, and even day to day, depending on the specific activities of each person. Separation of PM exposures into two components, ambient-generated PM and nonambient-generated PM, would reduce potential uncertainties in the analysis and interpretation of PM health effects data.
- Available data indicate that PM mass concentrations, especially fine PM, typically are distributed uniformly in most metropolitan areas, thus reducing the potential for exposure misclassification because of spatial variability when a limited number of ambient PM monitors are used to represent population average ambient exposures in community time-series or long-term, cross-sectional epidemiologic studies of PM.
- Even though the spatial variability of PM (in particular, $PM_{2.5}$) mass concentrations in urban environments seems to be small, the same conclusions drawn above regarding the influence of measurement errors may not necessarily hold for all of the PM components.
- There are important differences in the relationship of ambient PM concentrations (C_a) with exposures to ambient PM (E_{ag}), and with exposures to nonambient PM (E_{nonag}). Various researchers have shown that E_{ag} is a function of C_a, and that concentrations of ambient PM are driven by meteorology, by changes in source emission rates, and in locations of emission sources relative to the measurement site. However, E_{nonag} is independent of C_a, because concentrations of nonambient PM are driven by the daily activities of people.
- Because personal exposures also include a contribution from ambient concentrations, the correlation between daily-average personal exposure and the daily-average ambient concentration increases as the number of subjects measured daily increases. An application of a Random Component Superposition (RCS) model has shown that the contributions of ambient PM_{10} and indoor-generated PM_{10} to community mean exposure can be decoupled in modeling urban population exposure distributions.
- If linear nonthreshold models are assumed in time-series analysis of daily-average ambient PM concentrations and community health data, E_{nonag} is not expected to contribute to the relative risk estimates determined by regression of health responses on C_a.

- Using the PTEAM study data, analysis of exposure measurement errors in the context of time-series epidemiology has shown that errors or uncertainties introduced by using surrogate exposure variables, such as ambient PM concentrations, could lead to biases in the estimation of health risk coefficients.
- Because sources and chemical composition of particulate matter affecting personal exposures in different microenvironments vary, by season, day-of-the-week, and time of day, it is likely that some degree of misclassification of exposures to PM toxic agents of concern will be introduced when health effects models use only daily-average mass measures such as PM_{10} or $PM_{2.5}$. Because of the paucity of currently available data on many of these factors, it is impossible to ascertain at this point the significance of these more complex exposure misclassification problems in the interpretation of results from PM epidemiology.
- Exposure measurement errors may depend on particle size and composition. $PM_{2.5}$ better reflects personal exposure to PM of outdoor origin than PM_{10}. It is possible that various ultrafine particle measures, or other components of PM may be better exposure indicators for epidemiologic studies.
- Seasonal or temporal variations in the measurement errors and their correlations between different PM concentration measures and co-pollutants (e.g., SO_2, CO, NO_2, O_3) could influence the error analysis results but not likely the interpretation of current findings.
- Multi-pollutant personal exposure studies have suggested that ambient concentrations of gaseous copollutants serve as surrogates of personal exposures to particles rather than as confounders.
- Ordinarily, PM exposure measurement errors are not expected to influence the interpretation of findings from either the community time-series or long-term epidemiologic studies that have used ambient concentration data if they include sufficient adjustments for seasonality and key personal and geographic confounders.
- In the context of long-term epidemiologic studies, it appears that the errors introduced in estimating ambient PM indoor/ambient PM outdoor ratios have much more influence on the accuracy of the microenvironmental exposure estimation approach than do errors in estimating time spent in these microenvironments.

- To reduce exposure misclassification errors in PM epidemiology, conducting new cohort studies of sensitive populations with better real-time techniques for exposure monitoring and further speciation of indoor-generated, ambient, and personal PM mass are essential.
- Based on statistical modeling considerations, it is expected that existing PM exposure measurement errors or uncertainties most likely will reduce the statistical power of the PM health effects analysis, thus making it difficult to detect a true underlying association between the correct exposure metric and the health outcome studied.
- Although exposure measurement errors for fine particles are not expected to influence the interpretation of findings from either the community time-series or the long-term, cross-sectional epidemiologic studies that have used ambient concentration data, they may underestimate the strength of the impact. Sufficient data are not available to evaluate the impact of exposure measurement error for other PM species or size fractions.

REFERENCES

Abt, E.; Suh, H. H.; Allen, G.; Koutrakis, P. (2000a) Characterization of indoor particle sources: a study conducted in the metropolitan Boston area. Environ. Health Perspect. 108: 35-44.

Abt, E.; Suh, H. H.; Catalano, P.; Koutrakis, P. (2000b) The relative contribution of outdoor and indoor particle sources to indoor concentrations. Environ. Sci. Technol. 34: 3579-3587.

Akland, G. G.; Hartwell, T. D.; Johnson, T. R.; Whitmore, R. W. (1985) Measuring human exposure to carbon monoxide in Washington, D.C., and Denver, Colorado, during the winter of 1982-1983. Environ. Sci. Technol. 19: 911-918.

Al-Raheem, M. Y. A.; El-Desouky, M.; Mage, D. T.; Kollander, M.; Wallace, L. A.; Spengler, J. D. (2000) Personal exposures to PM and gaseous air pollutants at the end of the 1991 Kuwait oil fires. Presented at: PM2000: particulate matter and health—the scientific basis for regulatory decision-making, specialty conference & exhibition; January; Charleston, SC. Pittsburgh, PA: Air & Waste Management Association.

Alm, S.; Jantunen, M. J.; Vartiainen, M. (1999) Urban commuter exposure to particle matter and carbon monoxide inside an automobile. J. Exposure Anal. Environ. Epidemiol. 9: 237-244.

Anuszewski, J.; Larson, T. V.; Koenig, J. Q. (1998) Simultaneous indoor and outdoor particle light-scattering measurements at nine homes using a portable nephelometer. J. Exposure Anal. Environ. Epidemiol. 8: 483-493.

Baek, S.-O.; Kim, Y.-S.; Perry, R. (1997) Indoor air quality in homes, offices and restaurants in Korean urban areas--indoor/outdoor relationships. Atmos. Environ. 31: 529-544.

Bahadori, T.; Suh, H. H.; Rojas, L.; Koutrakis, P. (1999) Characterization of personal PM10 and PM2.5 exposures for individuals with COPD: results from a pilot study in Nashville, TN. J. Exposure Anal. Environ. Epidemiol.: in press.

Bahadori, T.; Suh, H. H.; Rojas-Bracho, L.; Koutrakis, P. (2001) Personal exposure to particulate matter of individuals with chronic obstructive pulmonary disease (COPD). J. Exposure Anal. Environ. Epidemiol.: accepted.

Bohadana, A. B.; Massin, N.; Wild, P.; Toamain, J.-P.; Engel, S.; Goutet, P. (2000) Symptoms, airway responsiveness, and exposure to dust in beech and oak wood workers. Occup. Environ. Med. 57: 268-273.

Brauer, M.; Mannetje, A. (1998) Restaurant smoking restrictions and environmental tobacco smoke exposure. Am. J. Public Health 88: 1834-1836.

Brauer, M.; Koutrakis, P.; Spengler, J. D. (1989) Personal exposures to acidic aerosols and gases. Environ. Sci. Technol. 23: 1408-1412.

Brauer, M.; Koutrakis, P.; Keeler, G. J.; Spengler, J. D. (1990) Indoor and outdoor concentrations of acidic aerosols and gases. In: Indoor air '90: precedings of the 5th international conference on indoor air quality and climate, volume 2, characteristics of indoor air; July-August; Toronto, ON, Canada. Ottawa, ON, Canada: International Conference on Indoor Air Quality and Climate, Inc.; pp. 447-452.

Brauer, M.; Bartlett, K.; Regalado-Pineda, J.; Perez-Padilla, R. (1996) Assessment of particulate concentrations from domestic biomass combustion in rural Mexico. Environ. Sci. Technol. 30: 104-109.

Brauer, M.; Hirtle, R. D.; Hall, A. C.; Yip, T. R. (1999) Monitoring personal fine particle exposure with a particle counter. J. Exposure Anal. Environ. Epidemiol. 9: 228-236.

Brauer, M.; Hrubá, F.; Mihalíková, E.; Fabiánová, E.; Miskovic, P.; Plziková, A.; Lendacká, M.; Vanderberg, J.; Cullen, A. (2000) Personal exposure to particles in Banská Bystrica, Slovakia. J. Exposure Anal. Environ. Epidemiol. 10: 478-487.

Brown, K. G.; Paxton, M. B. (1998) Predictability of personal exposure to airborne particulate matter from ambient concentrations in four communities. In: Vostal, J. J., ed. Health effects of particulate matter in ambient air. Proceedings of an international conference; 1997; Prague, Czech Republic. Pittsburgh, PA: Air & Waste Management Association; pp. 333-350. (A&WMA publication VIP-80).

Burke, J.; Zufall, M.; Özkaynak, H. (2001) A population exposure model for particulate matter: case study results for $PM_{2.5}$ in Philadelphia, PA. J. Exposure Anal. Environ. Epidemiol.11: 470-489.

Burton, R. M.; Suh, H. H.; Koutrakis, P. (1996) Spatial variation in particulate concentrations within metropolitan Philadelphia. Environ. Sci. Technol. 30: 400-407.

Buzorius, G.; Hämeri, K.; Pekkanen, J.; Kulmala, M. (1999) Spatial variation of aerosol number concentration in Helsinki city. Atmos. Environ. 33: 553-565.

Calder, K. L. (1957) A numerical analysis of the protection afforded by buildings against BW aerosol attack. Fort Detrick, MD: Office of the Deputy Commander for Scientific Activities; BWL tech. study no. 2.

Carrer, P.; Alcini, D.; Cavallo, D.; Bollini, D.; Ghalandar, R.; Lovato, L.; Ploia, E.; Vercelli, F.; Visigalli, F.; Maroni, M. (1998) Personal exposure to PM_{10} of subjects living in Milan. In: Vostal, J. J., ed. Health effects of particulate matter in ambient air. Proceedings of an international conference; 1997; Prague, Czech Republic. Pittsburgh, PA: Air & Waste Management Association; pp. 359-365. (A&WMA publication VIP-80).

Carroll, R. J.; Ruppert, D.; Stefanski, L. A. (1995) Measurement error in nonlinear models. London, United Kingdom: Chapman & Hall. (Cox, D. R.; Hinkley, D. V.; Keiding, N.; Reid, N.; Rubin, D. B.; Silverman, B. W., eds. Monographs on statistics and applied probability: v. 63).

Cha, Q.; Chen, Y.-Z.; Du, Y.-X. (1996) Influence of indoor air pollution on quality of outdoor air in city of Guangzhou, China. In: Indoor air '96: the 7th international conference on indoor air quality, volume I; July; Nagoya, Japan. Tokyo, Japan: Institute of Public Health; pp. 1131-1135.

Chang, L.-T.; Koutrakis, P.; Catalano, P. J.; Suh, H. H. (2000) Hourly personal exposures to fine particles and gaseous pollutants - results from Baltimore, Maryland. J. Air Waste Manage. Assoc. 50: 1223-1235.

Chao, C. Y. H.; Tung, T. C. (2001) An empirical model for outdoor contaminant transmission into residential buildings and experimental verification. Atmos. Environ. 35: 1585-1596.

Chuang, J. C.; Callahan, P. J.; Lyu, C. W.; Wilson, N. K. (1999) Polycyclic aromatic hydrocarbon exposures of children in low-income families. J. Exposure Anal. Environ. Epidemiol. 2: 85-98.

Clayton, C. A.; Perritt, R. L.; Pellizzari, E. D.; Thomas, K. W.; Whitmore, R. W.; Wallace, L. A.; Özkaynak, H.; Spengler, J. D. (1993) Particle total exposure assessment methodology (PTEAM) study: distributions of aerosol and elemental concentrations in personal, indoor, and outdoor air samples in a southern California community. J. Exposure Anal. Environ. Epidemiol. 3: 227-250.

Clayton, C. A.; Pellizzari, E. D.; Rodes, C. E.; Mason, R. E.; Piper, L. L. (1999a) Estimating distributions of long-term particulate matter and manganese exposures for residents of Toronto, Canada. Atmos. Environ. 33: 2515-2526.

Clayton, C. A.; Pellizzari, E. D.; Whitmore, R. W.; Perritt, R. L., Quackenboss, J. J. (1999b) National human exposure assessment survey (NHEXAS): distributions and associations of lead, arsenic, and volatile organic compounds in EPA Region 5. J. Exposure Anal. Environ. Epidemiol. 9: 381-392.

Cohen, B. S.; Li, W.; Xiong, J. Q.; Lippmann, M. (2000) Detecting H^+ in ultrafine ambient aerosol using iron nano-film detectors and scanning probe microscopy. Appl. Occup. Environ. Hyg. 15: 80-89.

Conner, T. L.; Lonneman, W. A.; Seila, R. L. (1995) Transportation-related volatile hydrocarbon source profiles measured in Atlanta. J. Air Waste Manage. Assoc. 45: 383-394.

Copeland, K. T.; Checkoway, H.; McMichael, A. J.; Holbrook, R. H. (1977) Bias due to misclassification in the estimation of relative risk. Am. J. Epidemiol. 105: 488-495.

Crump, K. S. (2000) Manganese exposures in Toronto during use of the gasoline additive, methylcyclopentadienyl manganese tricarbonyl. J. Exposure Anal. Environ. Epidemiol. 10: 227-239.

Daisey, J. M.; Mahanama, K. R. R.; Hodgson, A. T. (1998) Toxic volatile organic compounds in simulated enviromental tobacco smoke: emission factors for exposure assessment. J. Exposure Anal. Environ. Epidemiol. 8: 313-334.

Dockery, D. W.; Spengler, J. D. (1981) Indoor-outdoor relationships of respirable sulfates and particles. Atmos. Environ. 15: 335-343.

Dockery, D. W.; Pope, C. A., III; Xu, X.; Spengler, J. D.; Ware, J. H.; Fay, M. E.; Ferris, B. G., Jr.; Speizer, F. E. (1993) An association between air pollution and mortality in six U.S. cities. N. Engl. J. Med. 329: 1753-1759.

Dominici, F.; Zeger, S. L.; Samet, J. (2000) A measurement error model for time-series studies of air pollution and mortality. Biostatistics 1: 157-175.

Donham, K. J.; Cumro, D.; Reynolds, S. J.; Merchant, J. A. (2000) Dose-response relationships between occupational aerosol exposures and cross-shift declines of lung function in poultry workers: recommendations for exposure limits. J. Occup. Environ. Med. 42: 260-269.

Duan, N. (1982) Models for human exposure to air pollution. Environ. Int. 8: 305-309.

Duan, N. (1991) Stochastic microenvironment models for air pollution exposure. J. Exposure Anal. Environ. Epidemiol. 1: 235-257.

Dubowsky, S. D.; Wallace, L. A.; Buckley, T. J. (1999) The contribution of traffic to indoor concentrations of polycyclic aromatic hydrocarbons. J. Exposure Anal. Environ. Epidemiol. 9: 312-321.

Ebelt, S. T.; Petkau, A. J.; Vedal, S.; Fisher, T. V.; Brauer, M. (2000) Exposure of chronic obstructive pulmonary disease patients to particulate matter: relationships between personal and ambient air concentrations. J. Air Waste Manage. Assoc. 50: 1081-1094.

Esmen, N. A.; Hall, T. A. (2000) Theoretical investigation of the interrelationship between stationary and personal sampling in exposure estimation. Appl. Occup. Environ. Hyg. 15: 114-119.

Evans, G. F.; Highsmith, R. V.; Sheldon, L. S.; Suggs, J. C.; Williams, R. W.; Zweidinger, R. B.; Creason, J. P.; Walsh, D.; Rodes, C. E.; Lawless, P. A. (2000) The 1999 Fresno particulate matter exposure studies: comparison of community, outdoor, and residential PM mass measurements. J. Air Waste Manage. Assoc. 50: 1887-1896.

Ezzati, M.; Kammen, D. M. (2001) Quantifying the effects of exposure to indoor air pollution from biomass combustion on acute respiratory infections in developing countries. Environ. Health Perspect. 109: 481-488.

Ferris, B. G., Jr.; Speizer, F. E.; Spengler, J. D.; Dockery, D.; Bishop, Y. M. M.; Wolfson, M.; Humble, C. (1979) Effects of sulfur oxides and respirable particles on human health: methodology and demography of populations in study. Am. Rev. Respir. Dis. 120: 767-779.

Fischer, P. H.; Hoek, G.; Van Reeuwijk, H.; Briggs, D. J.; Lebret, E.; Van Wijnen, J. H.; Kingham, S.; Elliott, P. E. (2000) Traffic-related differences in outdoor and indoor concentrations of particles and volatile organic compounds in Amsterdam. Atmos. Environ. 34: 3713-3722.

Freijer, J. I.; Bloemen, H. J. T. (2000) Modeling relationships between indoor and outdoor air quality. J. Air Waste Manage. Assoc. 50: 292-300.

Frey, H. C.; Rhodes, D. S. (1996) Characterizing, simulating and analyzing variability and uncertainty: an illustration of methods using an air toxic emissions example. Hum. Ecol. Risk Assess. 2: 762-797.

Gladen, B.; Rogan, W. J. (1979) Misclassification and the design of environmental studies. Am. J. Epidemiol. 109: 607-616.

Glen, G.; Lakkadi, Y.; Tippett, J. A.; del Valle-Torres, M. (1997) Development of NERL/CHAD: the National Exposure Research Laboratory consolidated human activity database. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development; contract no. 68-D5-0049.

Goldstein, B. R.; Tardiff, G.; Hoffnagle, G.; Kester, R. (1992) Valdez health study: summary report. Ankorage, AK: Alyeska Pipeline Service Company.

Hadwen, G. E.; McCarthy, J. F.; Womble, S. E.; Girman, J. R.; Brightman, H. S. (1997) Volatile organic compound concentrations in 41 office buildings in the continental United States. In: Woods, J. E.; Grimsrud, D. T.; Boschi, N., eds. Proceedings of healthy buildings/indoor air quality international conference; September; Bethesda, MD. Washington, DC: Healthy Buildings/IAQ. [Healthy Buildings - International Conference; v. 2, pp. 465-470.]

Hartwell, T. D.; Clayton, C. A.; Michie, R. M., Jr.; Whitmore, R. W.; Zelon, H. S.; Whitehurst, D. A.; Akland, G. G. (1984) Study of carbon monoxide exposures of residents of Washington, D.C. Research Triangle Park, NC: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory; report no. EPA-600/D-84-177. Available from: NTIS, Springfield, VA; PB84-230051.

Hayes, S. R.; Marshall, J. R. (1999) Designing optimal strategies to attain the new U.S. particulate matter standards: some initial concepts. J. Air Waste Manage. Assoc. 49: PM192-PM198.

Henry, R. C.; Lewis, C. W.; Collins, J. F. (1994) Vehicle-related source compositions from ambient data: the GRACE/SAFER method. Environ. Sci. Technol. 28: 823-832.

Houseman, E. A.; Ryan, L.; Levy, J.; Richardson, D.; Spenger, J. D. (2001) Autocorrelation in real-time continuous monitoring of microenvironments. J. Appl. Stat.: in press.

Howard-Reed, C.; Rea, A. W.; Zufall, M. J.; Burke, J. M.; Williams, R. W.; Suggs, J. C.; Sheldon, L. S.; Walsh, D.; Kwock, R. (2000) Use of a continuous nephelometer to measure personal exposure to particles during the U.S. Environmental Protection Agency Baltimore and Fresno panel studies. J. Air Waste Manage. Assoc. 50: 1125-1132.

Institute for Social Research. (1997) Child development supplement to the panel study of income dynamics. Ann Arbor, MI: University of Michigan. Available: www.isr.umich.edu/src/child-development/home.html [2001, January 31].

Isukapalli, S. S.; Georgopoulos, P. G. (2000) Modeling of outdoor/indoor relationships of gas phase pollutants and particulate matter. Presented at: PM2000: particulate matter and health—the scientific basis for regulatory decision-making, specialty conference & exhibition; January; Charleston, SC. Pittsburgh, PA: Air & Waste Management Association.

Jang, M.; Kamens, R. M. (1999) Newly characterized products and composition of secondary aerosols from the reaction of α-pinene with ozone. Atmos. Environ. 33: 459-474.

Janssen, N. (1998) Personal exposure to airborne particles: validity of outdoor concentrations as a measure of exposure in time series studies [dissertation]. Wageningen, Netherlands: Landbouwuniversiteit Wageningen.

Janssen, N. A. H.; Hoek, G.; Harssema, H.; Brunekreef, B. (1995) A relation between personal and ambient . Epidemiology 6(suppl.): S45.

Janssen, N. A. H.; Hoek, G.; Harssema, H.; Brunekreef, B. (1997) Childhood exposure to PM_{10}: relation between personal, classroom, and outdoor concentrations. Occup. Environ. Med. 54: 888-894.

Janssen, N. A. H.; Hoek, G.; Brunekreef, B.; Harssema, H.; Mensink, I.; Zuidhof, A. (1998a) Personal sampling of particles in adults: relation among personal, indoor, and outdoor air concentrations. Am. J. Epidemiol. 147: 537-547.

Janssen, N. A. H.; Hoek, G.; Harssema, H.; Brunekreef, B. (1998b) Personal sampling of airborne particles: method performance and data quality. J. Exposure Anal. Environ. Epidemiol. 8: 37-49.

Janssen, N. A. H.; Hoek, G.; Harssema, H.; Brunekreef, B. (1999a) Personal exposure to fine particles in children correlates closely with ambient fine particles. Arch. Environ. Health 54: 95-101.

Janssen, N. A. H.; Hoek, G.; Brunekreef, B.; Harssema, H. (1999b) Mass concentration and elemental composition of PM_{10} in classrooms. Occup. Environ. Med. 56: 482-487.

Janssen, L. H. J. M.; Buringh, E.; van der Meulen, A.; van den Hout, K. D. (1999c) A method to estimate the distribution of various fractions of PM_{10} in ambient air in the Netherlands. Atmos. Environ. 33: 3325-3334.

Janssen, N. A. H.; De Hartog, J. J.; Hoek, G.; Brunekreef, B.; Lanki, T.; Timonen, K. L.; Pekkanen, J. (2000) Personal exposure to fine particulate matter in elderly subjects: relation between personal, indoor, and outdoor concentrations. J. Air Waste Manage. Assoc. 50: 1133-1143.

Jantunen, M. J.; Hänninen, O.; Katsouyanni, K.; Knöppel, H.; Kuenzli, N.; Lebret, E.; Maroni, M.; Saarela, K.; Srám, R.; Zmirou, D. (1998) Air pollution exposure in European cities: the *EXPOLIS* study. J. Exposure Anal. Environ. Epidemiol. 8: 495-518.

Jantunen, M. J.; Kousa, A.; Hanninen, O.; Koistinen, K.; Sram, R.; Lanki, T.; Lioy, P. J. (2000) Long term population PM exposures correlate well with outdoor air levels - short term individual exposures do not. Presented at: PM2000: particulate matter and health—the scientific basis for regulatory decision-making, specialty conference & exhibition; January; Charleston, SC. Pittsburgh, PA: Air & Waste Management Association.

Jenkins, R. A.; Palausky, M. A.; Counts, R. W.; Guerin, M. R.; Dindal, A. B.; Bayne, C. K. (1996a) Determination of personal exposure of non-smokers to environmental tobacco smoke in the United States. Lung Cancer 14(suppl. 1): S195-S213.

Jenkins, R. A.; Palausky, A.; Counts, R. W.; Bayne, C. K.; Dindal, A. B.; Guerin, M. R. (1996b) Exposure to environmental tobacco smoke in sixteen cities in the United States as determined by personal breathing zone air sampling. J. Exposure Anal. Environ. Epidemiol. 6: 473-502.

Johnson, T. (1984) A study of personal exposure to carbon monoxide in Denver, Colorado. Research Triangle Park, NC: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory; report no. EPA-600/4-84-014.

Johnson, T. (1989) Human activity patterns in Cincinnati, Ohio [final report]. Palo Alto, CA: Electric Power Research Institute; project no. RP940-06; report no. EN-6204.

Johnson, T. R. (1995) Recent advances in the estimation of population exposure to mobile source pollutants. J. Exposure Anal. Environ. Epidemiol. 5: 551-571.

Johnson, T.; Long, T.; Ollison, W. (2000) Prediction of hourly microenvironmental concentrations of fine particles based on measurements obtained from the Baltimore scripted activity study. J. Exposure Anal. Environ. Epidemiol. 10: 403-411.

Jones, N. C.; Thornton, C. A.; Mark, D.; Harrison, R. M. (2000) Indoor/outdoor relationships of particulate matter in domestic homes with roadside, urban and rural locations. Atmos. Environ. 34: 2603-2612.

Keywood, M. D.; Ayers, G. P.; Gras, J. L.; Gillett, R. W.; Cohen, D. D. (1999) Relationships between size segregated mass concentration data and ultrafine particle number concentrations in urban areas. Atmos. Environ. 33: 2907-2913.

Kingham, S.; Briggs, D.; Elliott, P.; Fischer, P.; Lebret, E. (2000) Spatial variations in the concentrations of traffic-related pollutants in indoor and outdoor air in Huddersfield, England. Atmos. Environ. 34: 905-916.

Kleeman, M. J.; Cass, G. R. (1998) Source contributions to the size and composition distribution of urban particulate air pollution. Atmos. Environ. 32: 2803-2816.

Klepeis, N. E. (1999) An introduction to the indirect exposure assessment approach: modeling human exposure using microenvironmental measurements and the recent national human activity pattern survey. Environ. Health Perspect. Suppl. 107(2): 365-374.

Klepeis, N. E.; Ott, W. R.; Switzer, P. (1994) A Total Human Exposure Model (THEM) for Respirable Suspended Particles (RSP). Presented at: 87th annual meeting & exhibition of the Air & Waste Management Association; June; Cincinnati, OH. Pittsburgh, PA: Air & Waste Management Association; paper no. 94-WA75A.03. Available from: NTIS, Springfield, VA; PB94-197415.

Klepeis, N. E.; Ott, W. R.; Switzer, P. (1995) Modeling the time series of respirable suspended particles and carbon monoxide from multiple smokers: validation in two public smoking lounges. Presented at: 88th annual meeting and exhibition of the Air & Waste Management Association; June; San Antonio, TX. Pittsburgh, PA: Air & Waste Management Association; paper no. 95-MPH.03.

Klepeis, N. E.; Ott, W. R.; Switzer, P. (1996) A multiple-smoker model for predicting indoor air quality in public lounges. Environ. Sci. Technol. 30: 2813-2820.

Klepeis, N. E.; Switzer, P.; Ott, W. R. (2000) Estimating exposure to environmental tobacco smoke particles using 24-hour time-activity recall diaries. Presented at: PM2000: particulate matter and health—the scientific basis for regulatory decision-making, specialty conference & exhibition; January; Charleston, SC. Pittsburgh, PA: Air & Waste Management Association.

Koontz, M. D.; Rector, H. E. (1995) Estimation of distributions for residential air exchange rates: final report. Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics; contract nos. 68-D9-0166 and 68-D3-0013.

Koontz, M. D.; Niang, L. L. (1998) California population indoor exposure model (CPIEM): version 1.4F. Final report A933-157: user's guide. Sacramento, CA: California Air Resources Board; GEOMET report no. IE-2632.

Koutrakis, P.; Briggs, S. L. K.; Leaderer, B. P. (1992) Source apportionment of indoor aerosols in Suffolk and Onondaga Counties, New York. Environ. Sci. Technol. 26: 521-527.

Krewski, D.; Burnett, R. T.; Goldberg, M. S.; Hoover, K.; Siemiatycki, J.; Jerrett, M.; Abrahamowicz, M.; White, W. H. (2000) Reanalysis of the Harvard Six Cities study and the American Cancer Society study of particulate air pollution and mortality: a special report of the Institute's Particle Epidemiology Reanalysis Project. Cambridge, MA: Health Effects Institute.

Lachenmyer, C.; Hidy, G. M. (2000) Urban measurements of outdoor-indoor $PM_{2.5}$ concentrations and personal exposure in the deep south. Part I. Pilot study of mass concentrations for nonsmoking subjects. Aerosol Sci. Technol. 32: 34-51.

Laden, F.; Neas, L. M.; Dockery, D. W.; Schwartz, J. (2000) Association of fine particulate matter from different sources with daily mortality in six U.S. cities. Environ. Health Perspect. 108: 941-947.

Leaderer, B. P.; Naeher, L.; Jankun, T.; Balenger, K.; Holford, T. R.; Toth, C.; Sullivan, J.; Wolfson, J. M.; Koutrakis, P. (1999) Indoor, outdoor, and regional summer and winter concentrations of PM_{10}, $PM_{2.5}$, SO_4^{2-}, H^+, NH_4^+, NO_3^-, NH_3, and nitrous acid in homes with and without kerosene space heaters. Environ. Health Perspect. 107: 223-231.

Lee, S. C.; Chang, M. (1999) Indoor air quality investigations at five classrooms. Indoor Air 9: 134-138.

Lessler, J. T.; Kalsbeek, W. D. (1992) Nonsampling error in surveys. New York, NY: John Wiley & Sons, Inc.; p. 7.

Liao, D.; Creason, J.; Shy, C.; Williams, R.; Watts, R.; Zweidinger, R. (1999) Daily variation of particulate air pollution and poor cardiac autonomic control in the elderly. Environ. Health Perspect. 107: 521-525.

Lillquist, D. R.; Lee, J. S.; Ramsay, J. R.; Boucher, K. M.; Walton, Z. L.; Lyon, J. L. (1998) A comparison of indoor/outdoor PM_{10} concentrations measured at three hospitals and a centrally located monitor in Utah. Appl. Occup. Environ. Hyg. 13: 409-415.

Linn, W. S.; Gong, H., Jr.; Clark, K. W.; Anderson, K. R. (1999) Day-to-day particulate exposures and health changes in Los Angeles area residents with severe lung disease. J. Air Waste Manage. Assoc. 49: PM108-PM115.

Lioy, P. J. (1990) Assessing total human exposure to contaminants: a multidisciplinary approach. Environ. Sci. Technol. 24: 938-945.

Lioy, P. J.; Waldman, J. M.; Buckley, T.; Butler, J.; Pietarinen, C. (1990) The personal, indoor and outdoor concentrations of PM-10 measured in an industrial community during the winter. Atmos. Environ. Part B 24: 57-66.

Lioy, P. J.; Wainman, T.; Zhang, J.; Goldsmith, S. (1999) Typical household vacuum cleaners: the collection efficiency and emissions characteristics for fine particles. J. Air Waste Manage. Assoc. 49: 200-206.

Lippmann, M.; Ito, K.; Nádas, A.; Burnett, R. T. (2000) Association of particulate matter components with daily mortality and morbidity in urban populations. Cambridge, MA: Health Effects Institute; research report no. 95.

Long, C. M.; Suh, H. H.; Koutrakis, P. (2000a) Characterization of indoor particle sources using continuous mass and size monitors. J. Air Waste Manage. Assoc. 50: 1236-1250.

Long, C. M.; Suh, H. H.; Koutrakis, P. (2000b) Using time- and size-resolved particulate data to explore infiltration and deposition behavior. Environ. Sci. Technol.: accepted.

Lurmann, F. W.; Korc, M. E. (1994) User's guide to the regional human exposure (REHEX) model. San Francisco, CA: Bay Area Air Quality Management District; draft report no. STI-93150-1414-DR.

MacIntosh, D. L.; Xue, J.; Özkaynak, H.; Spengler, J. D.; Ryan, P. B. (1995) A population-based exposure model for benzene. J. Exposure Anal. Environ. Epidemiol. 5: 375-403.

Mage, D. T. (1985) Concepts of human exposure assessment for airborne particulate matter. Environ. Int. 11: 407-412.

Mage, D. T. (1998) How much PM of ambient origin are people exposed to? In: Measurement of toxic and related air pollutants. Volume II: proceedings of a specialty conference; September; Cary, NC. Pittsburgh, PA: Air & Waste Management Association; pp. 666-681. (A&WMA publication VIP-85).

Mage, D. T.; Ott, W. R. (1996) Accounting for nonuniform mixing and human exposure in indoor environments. In: Tichenor, B. A., ed. Characterizing sources of indoor air pollution and related sink effects. Philadelphia, PA: American Society for Testing and Materials; pp. 263-278.

Mage, D.; Wilson, W.; Hasselblad, V.; Grant, L. (1999) Assessment of human exposure to ambient particulate matter. J. Air Waste Manage. Assoc. 49: 174-185.

Mar, T. F.; Norris, G. A.; Koenig, J. Q.; Larson, T. V. (2000) Associations between air pollution and mortality in Phoenix, 1995-1997. Environ. Health Perspect. 108: 347-353.

McBride, S. J.; Ferro, A. R.; Ott, W. R; Switzer, P.; Hildemann, L. M. (1999) Investigations of the proximity effect for pollutants in the indoor environment. J. Exposure Anal. Environ. Epidemiol. 9: 602-621.

McCurdy, T. (1995) Estimating human exposure to selected motor vehicle pollutants using the NEM series of models: lessons to be learned. J. Exposure Anal. Environ. Epidemiol. 5: 533-550.

McCurdy, T. (1997) Modeling the dose profile in human exposure assessments: ozone as an example. Rev. Toxicol. 1: 3-23.

McCurdy, T. (2000) Conceptual basis for multi-route intake dose modeling using an energy expenditure approach. J. Exposure Anal. Environ. Epidemiol. 10: 86-97.

McCurdy, T.; Glen, G.; Smith, L.; Lakkadi, Y. (2000) The National Exposure Research Laboratory's Consolidated Human Activity Database. J. Exposure Anal. Environ. Epidemiol. 10: 566-578.

Mihlan, G. J.; Todd, L. A.; Truong, K. N. (2000) Assessment of occupational exposure patterns by frequency-domain analysis of time series data. Appl. Occup. Environ. Hyg. 15: 120-130.

Miller, S. L.; Branoff, S.; Nazaroff, W. W. (1998) Exposure to toxic air contaminants in environmental tobacco smoke: an assessment for California based on personal monitoring data. J. Exposure Anal. Environ. Epidemiol. 8: 287-311.

Monn, C. (2001) Exposure assessment of air pollutants: a review on spatial heterogeneity and indoor/outdoor/personal exposure to suspended particulate matter, nitrogen dioxide and ozone. Atmos. Environ. 35: 1-32.

Monn, Ch.; Carabias, V.; Junker, M.; Waeber, R.; Karrer, M.; Wanner, H. U. (1997) Small-scale spatial variability of particulate matter <10 µm (PM_{10}) and nitrogen dioxide. Atmos. Environ. 31: 2243-2247.

Morawska, L.; He, C.; Hitchins, J.; Gilbert, D.; Parappukkaran, S. (2001) The relationship between indoor and outdoor airborne particles in the residential environment. Atmos. Environ. 35: 3463-3473.

Muraleedharan, T. R.; Radojevic, M. (2000) Personal particle exposure monitoring using nephelometry during haze in Brunei. Atmos. Environ. 34: 2733-2738.

Murray, D. M.; Burmaster, D. E. (1995) Residential air exchange rates in the United States: empirical and estimated parametric distributions by season and climatic region. Risk Anal. 15: 459-465.

National Air Pollution Control Administration. (1969) Air quality criteria for particulate matter. Washington, DC: U.S. Department of Health, Education, and Welfare, Public Health Service; NAPCA publication no. AP-49. Available from: NTIS, Springfield, VA; PB-190251/BA.

National Research Council. (1991) Human exposure assessment for airborne pollutants: advances and opportunities. Washington, DC: National Academy of Sciences.

National Research Council. (1998) Research priorities for airborne particulate matter. I. Immediate priorities and a long-range research portfolio. Washington, DC: National Academy Press.

National Research Council. (1999) Research priorities for airborne particulate matter. II. Evaluating research progress and updating the portfolio. Washington, DC: National Academy Press.

Navidi, W.; Thomas, D.; Langholz, B.; Stram, D. (1999) Statistical methods for epidemiologic studies of the health effects of air pollution. Cambridge, MA: Health Effects Institute; research report no. 86.

Nelson, P. R.; Conrad, F. W.; Kelly, S. P.; Maiolo, K. C.; Richardson, J. D.; Ogden, M. W. (1998) Composition of environmental tobacco smoke (ETS) from international cigarettes part II: nine country follow-up. Environ. Int. 24: 251-257.

Nieuwenhuijsen, M. J.; Noderer, K. S.; Schenker, M. B.; Vallyathan, V.; Olenchock, S. (1999) Personal exposure to dust, endotoxin and crystalline silica in California agriculture. Ann. Occup. Hyg. 43: 35-42.

Oglesby, L.; Künzli, N.; Röösli, M.; Braun-Fahrländer, C.; Mathys, P.; Stern, W.; Jantunen, M.; Kousa, A. (2000) Validity of ambient levels of fine particles as surrogate for personal exposure to outdoor air pollution—results of the European EXPOLIS-EAS study (Swiss Center Basel). J. Air Waste Manage. Assoc. 50: 1251-1261.

Ott, W. R. (1984) Exposure estimates based on computer generated activity patterns. J. Toxicol. Clin. Toxicol. 21: 97-128.

Ott, W. R. (1999) Mathematical models for predicting indoor air quality from smoking activity. Environ. Health Perspect. 107(suppl. 2): 375-381.

Ott, W.; Thomas, J.; Mage, D.; Wallace, L. (1988) Validation of the simulation of human activity and pollutant exposure (SHAPE) model using paired days from the Denver, CO, carbon monoxide field study. Atmos. Environ. 22: 2101-2113.

Ott, W. R.; Mage, D. T.; Thomas, J. (1992) Comparison of microenvironmental CO concentrations in two cities for human exposure modeling. J. Exposure Anal. Environ. Epidemiol. 2: 249-267.

Ott, W. R.; Klepeis, N. E.; Switzer, P. (1995) Modeling environmental tobacco smoke in the home using transfer functions. Presented at: 88th annual meeting and exhibition of the Air & Waste Management Association; June; San Antonio, TX; paper no. 95-WP84B.03. Research Triangle Park, NC: U.S. Environmental Protection Agency, Atmospheric Research And Exposure Assessment Laboratory; report no. EPA/600/A-95/079. Available from: NTIS, Springfield, VA; PB95-225488.

Ott, W.; Switzer, P.; Robinson, J. (1996) Particle concentrations inside a tavern before and after prohibition of smoking: evaluating the performance of an indoor air quality model. J. Air Waste Manage. Assoc. 46: 1120-1134.

Ott, W.; Wallace, L.; Mage, D. (2000) Predicting particulate (PM_{10}) personal exposure distributions using a random component superposition (RCS) statistical model. J. Air Waste Manage. Assoc. 50: 1390-1406.

Owen, M. K.; Ensor, D. S.; Sparks, L. E. (1992) Airborne particle sizes and sources found in indoor air. Atmos. Environ. Part A 26: 2149-2162.

Özkaynak, H.; Spengler, J. (1996) The role of outdoor particulate matter in assessing total human exposure. In: Wilson, R.; Spengler, J., eds. Particles in our air: concentrations and health effects. Cambridge, MA: Harvard University Press.

Özkaynak, H.; Thurston, G. D. (1987) Associations between 1980 U.S. mortality rates and alternative measures of airborne particle concentration. Risk Anal. 7: 449-461.

Özkaynak, H.; Ryan, P. B.; Spengler, J. D.; Laird, N. M. (1986) Bias due to misclassification of personal exposures in epidemiologic studies of indoor and outdoor air pollution. In: Berglund, B.; Berglund, U.; Lindvall, T.; Spengler, J.; Sundell, J., eds. Indoor air quality: papers from the third international conference on indoor air quality and climate; August 1984; Stockholm, Sweden. Environ. Int. 12: 389-393.

Özkaynak, H.; Spengler, J. D.; Ludwig, J. F.; Butler, D. A.; Clayton, C. A.; Pellizzari, E.; Wiener, R. W. (1990) Personal exposure to particulate matter: findings from the Particle Total Exposure Assessment Methodology (PTEAM) prepilot study. In: Indoor air '90: precedings of the 5th international conference on indoor air quality and climate, volume 2, characteristics of indoor air; July-August; Toronto, ON, Canada. Ottawa, ON, Canada: International Conference on Indoor Air Quality and Climate, Inc.; pp. 571-576.

Özkaynak, H.; Spengler, J. D.; Xue, J.; Koutrakis, P.; Pellizzari, E. D.; Wallace, L. (1993) Sources and factors influencing personal and indoor exposures to particles, elements and nicotine: findings from the particle TEAM pilot study. In: Jantunen, M.; Kalliokoski, P.; Kukkonen, E.; Saarela, K.; Seppänen, O.; Vuorelma,

H., eds. Indoor air '93: proceedings of the 6th international conference on indoor air quality and climate, v. 3, combustion products, risk assessment, policies; July; Helsinki, Finland. Helsinki, Finland: Indoor Air '93; pp. 457-462.

Özkaynak, H.; Xue, J.; Spengler, J.; Wallace, L.; Pellizzari, E.; Jenkins, P. (1996a) Personal exposures to airborne particles and metals: results from the particle TEAM study in Riverside, California. J. Exposure Anal. Environ. Epidemiol. 6: 57-78.

Özkaynak, H.; Xue, J.; Weker, R.; Butler, D.; Koutrakis, P.; Spengler, J. (1996b) The particle TEAM (PTEAM) study: analysis of the data: final report, volume III . Research Triangle Park, NC: U.S. Environmental Protection Agency, Atmospheric Research and Exposure Assessment Laboratory; report no. EPA/600/R-95/098. Available from: NTIS, Springfield, VA; PB97-102495.

Patterson, E.; Eatough, D. J. (2000) Indoor/outdoor relationships for ambient $PM_{2.5}$ and associated pollutants: epidemiological implications in Lindon, Utah. J. Air Waste Manage. Assoc. 50: 103-110.

Pellizzari, E. D., Mason, R. E.; Clayton, C. A.; Thomas, K. W.; Cooper, S.; Piper. L.; Rodes, C.; Goldberg, M.; Roberds, J.; Michael, L. (1998) Manganese exposure study (Toronto). Final report. Research Triangle Park, NC: Research Triangle Institute, Analytical and Chemical Sciences; report no. RTI/6312/02-01 F.

Pellizzari, E. D.; Clayton, C. A.; Rodes, C. E.; Mason, R. E.; Piper, L. L.; Fort, B.; Pfeifer, G.; Lynam, D. (1999) Particulate matter and manganese exposures in Toronto, Canada. Atmos. Environ. 33: 721-734.

Pellizzari, E. D.; Clayton, C. A.; Rodes, C. E.; Mason, R. E.; Piper, L. L.; Fort, B.; Pfeifer, G.; Lynam, D. (2001) Particulate matter and manganese exposures in Indianapolis, Indiana. J. Exposure Anal. Environ. Epidemiol. 11: 423-440.

Phillips, K.; Howard, D. A.; Browne, D.; Lewsley, J. M. (1994) Assessment of personal exposures to environmental tobacco smoke in British nonsmokers. Environ. Int. 20: 693-712.

Phillips, K.; Bentley, M. C.; Howard, D. A.; Alván, G. (1996) Assessment of air quality in Stockholm by personal monitoring of nonsmokers for respirable suspended particles and environmental tobacco smoke. Scand. J. Work Environ. Health 22(suppl. 1): 1-24.

Phillips, K.; Bentley, M. C.; Howard, D. A.; Alván, G.; Huici, A. (1997a) Assessment of air quality in Barcelona by personal monitoring of nonsmokers for respirable suspended particles and environmental tobacco smoke. Environ. Int. 23: 173-196.

Phillips, K.; Howard, D. A.; Bentley, M. C.; Alván, G. (1997b) Assessment of air quality in Turin by personal monitoring of nonsmokers for respirable suspended particles and environmental tobacco smoke. Environ. Int. 23: 851-871.

Phillips, K.; Bentley, M. C.; Howard, D. A.; Alván, G. (1998a) Assessment of air quality in Paris by personal monitoring of nonsmokers for respirable suspended particles and environmental tobacco smoke. Environ. Int. 24: 405-425.

Phillips, K.; Howard, D. A.; Bentley, M. C.; Alván, G. (1998b) Assessment of environmental tobacco smoke and respirable suspended particle exposures for nonsmokers in Lisbon by personal monitoring. Environ. Int. 24: 301-324.

Phillips, K.; Howard, D. A.; Bentley, M. C.; Alván, G. (1999) Assessment of environmental tobacco smoke and respirable suspended particle exposures for nonsmokers in Basel by personal monitoring. Atmos. Environ. 33: 1889-1904.

Pope, C. A., III; Thun, M. J.; Namboodiri, M. M.; Dockery, D. W.; Evans, J. S.; Speizer, F. E.; Heath, C. W., Jr. (1995) Particulate air pollution as a predictor of mortality in a prospective study of U.S. adults. Am. J. Respir. Crit. Care Med. 151: 669-674.

Pope, C. A., III; Hill, R. W.; Villegas, G. M. (1999) Particulate air pollution and daily mortality on Utah's Wasatch Front. Environ. Health Perspect.: 107: 567-573.

Praml, G.; Schierl, R. (2000) Dust exposure in Munich public transportation: a comprehensive 4-year survey in buses and trams. Int. Arch. Occup. Environ. Health 73: 209-214.

Quintana, P. J. E.; Samimi, B. S.; Kleinman, M. T.; Liu, L.-J.; Soto, K.; Warner, G. Y.; Bufalino, C.; Valencia, J.; Francis, D.; Hovell, M. H.; Delfino, R. J. (2000) Evaluation of a real-time passive personal particle monitor in fixed site residential indoor and ambient measurements. J. Exposure Anal. Environ. Epidemiol. 10: 437-445.

Ramachandran, G.; Vincent, J. H. (1999) A Bayesian approach to retrospective exposure assessment. Appl. Occup. Environ. Hyg. 14: 547-557.

Rea, A. W.; Zufall, M. J.; Williams, R. W.; Sheldon, L. S.; Howard-Reed, C. (2001) The influence of human activity patterns on personal PM exposure: a comparative analysis of filter-based and continuous particle measurements. J. Air Waste Manage. Assoc. 51: 1271-1279.

Ren, P.; Jankun, T. M.; Leaderer, B. P. (1999) Comparisons of seasonal fungal prevalence in indoor and outdoor air and in house dusts of dwellings in one northeast American county. J. Exposure Anal. Environ. Epidemiol. 9: 560-568.

Riediker, NM.; Keller, S.; Wuthrich, B.; Koller, T.; Monn, C. (2000) Personal pollen exposure compared to stationary measurements. J. Invest. Allergol. Clin. Immunol. 10: 200-203.

Robinson, J. P.; Wiley, J. A.; Piazza, T.; Garrett, K.; Cirksena, K. (1989) Activity patterns of California residents and their implications for potential exposure to pollution. Sacramento, CA: California Air Resources Board report no. CARB-A6-177-33.

Robinson, J. P.; Switzer, P.; Ott, W. (1994) Microenvironmental factors related to Californians' potential exposure to environmental tobacco smoke (ETS). Research Triangle Park, NC: U.S. Environmental Protection Agency, Atmospheric Research and Exposure Assessment Laboratory; report no. EPA/600/R-94/116. Available from: NTIS, Springfield, VA; PB94-214814.

Rodes, C.; Sheldon, L.; Whitaker, D.; Clayton, A.; Fitzgerald, K.; Flanagan, J.; DiGenova, F.; Hering, S.; Frazier, C. (1998) Measuring concentrations of selected air pollutants inside California vehicles [final report]. Sacramento, CA: California Environmental Protection Agency, Air Resources Board; contract no. 95-339.

Rojas-Bracho, L.; Suh, H. H.; Koutrakis, P. (2000) Relationships among personal, indoor, and outdoor fine and coarse particle concentrations for individuals with COPD. J. Exposure Anal. Environ. Epidemiol. 10: 294-306.

Roorda-Knape, M. C.; Janssen, N. A. H.; De Hartog, J. J.; Van Vliet, P. H. N.; Harssema, H.; Brunekreef, B. (1998) Air pollution from traffic in city districts near major motorways. Atmos. Environ. 32: 1921-1930.

Rotko, T.; Oglesby, L.; Künzli, N.; Jantunen, M. J. (2000a) Population sampling in European air pollution exposure study, *EXPOLIS*: comparisons between the cities and representativeness of the samples. J. Exposure Anal. Environ. Epidemiol. 10: 355-364.

Rotko, T.; Koistinen, K.; Hänninen, O.; Jantunen, M. (2000b) Sociodemographic descriptors of personal exposure to fine particles ($PM_{2.5}$) in *EXPOLIS* Helsinki. J. Exposure Anal. Environ. Epidemiol. 10: 385-393.

Ryan, P. B. (1991) An overview of human exposure modeling. J. Exposure Anal. Environ. Epidemiol. 1: 453-474.

Samet, J. M.; Dominici, F.; Zeger, S. L.; Schwartz, J.; Dockery, D. W. (2000) National morbidity, mortality, and air pollution study. Part I: methods and methodologic issues. Cambridge, MA: Health Effects Institute; research report no. 94.

Santos-Burgoa, C.; Rojas-Bracho, L.; Rosas-Pérez, I.; Ramírez-Sánchez, A.; Sánchez-Rico, G.; Mejía-Hernández, S. (1998) Modelaje de exposición a partículas en población general y riesgo de enfermedad respiratoria [Particle exposure modeling in the general population and risk of respiratory disease]. Gac. Med. Mex. 134: 407-417.

Sarnat, J. A.; Koutrakis, P.; Suh, H. H. (2000) Assessing the relationship between personal particulate and gaseous exposures of senior citizens living in Baltimore, MD. J. Air Waste Manage. Assoc. 50: 1184-1198.

Sarnat, J. A.; Schwartz, J.; Catalano, P. J.; Suh, H. H. (2001) Gaseous pollutants in particulate matter epidemiology: confounders or surrogates? Environ. Health Perspect. 109: 1053-1061.

Schwartz, J.; Dockery, D. W.; Neas, L. M. (1996) Is daily mortality associated specifically with fine particles? J. Air Waste Manage. Assoc. 46: 927-939.

Sexton, K.; Ryan, P. B. (1988) Assessment of human exposure to air pollution: methods, measurements, and models. In: Watson, A. Y.; Bates, R. R.; Kennedy, D., eds. Air pollution, the automobile, and public health. Washington, DC: National Academy Press; pp. 207-238.

Sexton, K.; Spengler, J. D.; Treitman, R. D. (1984) Personal exposure to respirable particles: a case study in Waterbury, Vermont. Atmos. Environ. 18: 1385-1398.

Sheldon, L.; Clayton, A.; Jones, B.; Keever, J.; Perritt, R.; Smith, D.; Whitaker, D.; Whitmore, R. (1992a) Indoor pollutant concentrations and exposures. Sacramento, CA: California Air Resources Board; report no. ARB/R-92/486.

Sheldon, L.; Clayton, A.; Keever, J. Perritt, R.; Whitaker, D. (1992b) PTEAM: monitoring of phthalates and PAHs in indoor and outdoor air samples in Riverside, California, volume II [final report]. Sacramento, CA: California Environmental Protection Agency, State Air Resources Board, Research Division; contract no. A933-144. Available from: NTIS, Springfield, VA; PB93-205649/XAB.

Sheldon, L.; Whitaker, D.; Keever, J.; Clayton, A.; Perritt, R. (1993a) Phthalates and PAHs in indoor and outdoor air in a southern California community. In: Jantunen, M.; Kalliokoski, P.; Kukkonen, E.; Saarela, K.; Seppänen, O.; Vuorelma, H., eds. Indoor air '93: proceedings of the 6th international conference on indoor air quality and climate. Volume 3: combustion products, risk assessment, policies; July; Helsinki, Finland. Helsinki, Finland: Indoor Air '93; pp. 109-114.

Sheldon, L.; Clayton, A.; Keever, J.; Perritt, R.; Whitaker, D. (1993b) Indoor concentrations of polycyclic aromatic hydrocarbons in California residences. Sacramento, CA: Air Resources Board; ARB contract no. A033-132.

Sheldon, L.; Clayton, A.; Perritt, R.; Whitaker, D. A.; Keever, J. (1993c) Indoor concentrations of polycyclic aromatic hydrocarbons in California residences and their relationship to combustion source use. In: Jantunen, M.; Kalliokoski, P.; Kukkonen, E.; Saarela, K.; Seppänen, O.; Vuorelma, H., eds. Indoor air '93: proceedings of the 6th international conference on indoor air quality and climate. Volume 3: combustion products, risk assessment, policies; July; Helsinki, Finland. Helsinki, Finland: Indoor Air '93; pp. 29-34.

Shy, C. M.; Kleinbaum, D. G.; Morgenstern, H. (1978) The effect of misclassification of exposure status in epidemiological studies of air pollution health effects. Bull. N. Y. Acad. Med. 54: 1155-1165.

Siegmann, K.; Scherrer, L.; Siegmann, H. C. (1999) Physical and chemical properties of airborne nanoscale particles and how to measure the impact on human health. J. Mol. Struct. (Theochem) 458: 191-201.

Spengler, J. D.; Treitman, R. D.; Tosteson, T. D.; Mage, D. T.; Soczek, M. L. (1985) Personal exposures to respirable particulates and implications for air pollution epidemiology. Environ. Sci. Technol. 19: 700-707.

Spengler, J. D.; Özkaynak, H.; Ludwig, J.; Allen, G.; Pellizzari, E. D.; Wiener, R. (1989) Personal exposures to particulate matter: instruments and methodologies P-TEAM. In: Measurement of toxic and related air pollutants: proceedings of the 1989 U.S. EPA/A&WMA international symposium; May; Raleigh, NC. Pittsburgh, PA: Air & Waste Management Association; pp. 449-463. (A&WMA publication VIP-13).

Spier, C. E.; Little, D. E.; Trim, S. C.; Johnson, T. R.; Linn, W. S.; Hackney, J. D. (1992) Activity patterns in elementary and high school students exposed to oxidant pollution. J. Exposure Anal. Environ. Epidemiol. 2: 277-293.

Suggs, J. C.; Burton, R. M. (1983) Spatial characteristics of inhalable particles in the Philadelphia metropolitan area. J. Air Pollut. Control Assoc. 33: 688-691.

Suh, H. H.; Spengler, J. D.; Koutrakis, P. (1992) Personal exposures to acid aerosols and ammonia. Environ. Sci. Technol. 26: 2507-2517.

Suh, H. H.; Koutrakis, P.; Spengler, J. D. (1993a) Indoor and outdoor acid aerosol and gas concentrations. In: Jantunen, M.; Kalliokoski, P.; Kukkonen, E.; Saarela, K.; Seppänen, O.; Vuorelma, H., eds. Indoor air '93: proceedings of the 6th international conference on indoor air quality and climate. Volume 3: combustion products, risk assessment, policies; July; Helsinki, Finland. Helsinki, Finland: Indoor Air '93; pp. 257-262.

Suh, H. H.; Koutrakis, P.; Spengler, J. D. (1993b) Validation of personal exposure models for sulfate and aerosol strong acidity. J. Air Waste Manage. Assoc. 43: 845-850.

Suh, H. H.; Koutrakis, P.; Spengler, J. D. (1994) The relationship between airborne acidity and ammonia in indoor environments. J. Exposure Anal. Environ. Epidemiol. 4: 1-23.

Suh, H. H.; Allen, G. A.; Koutrakis, P.; Burton, R. M. (1995) Spatial variation in acidic sulfate and ammonia concentrations within metropolitan Philadelphia. J. Air Waste Manage. Assoc. 45: 442-452.

Tamura, K.; Ando, M.; Sagai, M.; Matsumoto, Y. (1996a) Estimation of levels of personal exposure to suspended particulate matter and nitrogen dioxide in Tokyo. Environ. Sci. (Tokyo) 4: 37-51.

Tamura, K.; Ando, M.; Miyazaki, T.; Funasaka, K.; Kuroda, K. (1996b) Indoor concentrations and personal exposure to particulate matter at roadside houses. In: Indoor air '96: the 7th international conference on indoor air quality, volume I; July; Nagoya, Japan. Tokyo, Japan: Institute of Public Health; pp. 599-602.

Teschke, K.; Marion, S. A.; Vaughan, T. L.; Morgan, M. S.; Camp, J. (1999) Exposures to wood dust in U.S. industries and occupations, 1979 to 1997. Am. J. Ind. Med. 35: 581-589.

Thomas, K. W.; Pellizzari, E. D.; Clayton, C. A.; Whitaker, D. A.; Shores, R. C.; Spengler, J.; Özkaynak, H.; Froehlich, S. E.; Wallace, L. A. (1993) Particle total exposure assessment methodology (PTEAM) 1990 study: method performance and data quality for personal, indoor, and outdoor monitoring. J. Exposure Anal. Environ. Epidemiol. 3: 203-226.

Tsang, A. M.; Klepeis, N. E. (1996) Descriptive statistics tables from a detailed analysis of the national human activity pattern survey (NHAPS) data. Las Vegas, NV: U.S. Environmental Protection Agency, Office of Research and Development; report no. EPA/600/R-96/148.

U.S. Environmental Protection Agency. (1982) Air quality criteria for particulate matter and sulfur oxides. Research Triangle Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; EPA report no. EPA-600/8-82-029aF-cF. 3v. Available from: NTIS, Springfield, VA; PB84-156777.

U.S. Environmental Protection Agency. (1996) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.

U.S. Environmental Protection Agency. (1998) Particulate matter research needs for human health risk assessment to support future reviews of the national ambient air quality standards for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment; report no. EPA/600/R-97/132F.

U.S. Environmental Protection Agency. (2000) Aerometric Information Retrieval System (AIRS). Research Triangle Park, NC: Office of Air Quality Planning and Standards. Available: www.epa.gov/ttn/airs/ [2000, October 6].

Väkevä, M.; Hämeri, K.; Kulmala, M.; Lahdes, R.; Ruuskanen, J.; Laitinen, T. (1999) Street level versus rooftop concentrations of submicron aerosol particles and gaseous pollutants in an urban street canyon. Atmos. Environ. 33: 1385-1397.

Van Vliet, P.; Knape, M.; De Hartog, J.; Janssen, N.; Harssema, H.; Brunekreef, B. (1997) Motor vehicle exhaust and chronic respiratory symptoms in children living near freeways. Environ. Res. 74: 122-132.

Vette, A. F.; Rea, A. W.; Lawless, P. A.; Rodes, C. E.; Evans, G.; Highsmith, V. R.; Sheldon, L. (2001) Characterization of indoor-outdoor aerosol concentration relationships during the Fresno PM exposure studies. Aerosol Sci. Technol.: accepted.

Vyas, V. M.; Sun, Q.; Chandrasekar, A.; Wang, S.; Georgopoulos, P. G.; Burke, J.; Xue, J.; Özkaynak, H. (2002) Source-to-dose analysis of population exposures to PM. Presented at: Eleventh annual meeting of the International Society of Exposure Analysis; November 2001; Charleston, SC.

Wainman, T.; Zhang, J.; Weschler, C. J.; Lioy, P. J. (2000) Ozone and limonene in indoor air: a source of submicron particle exposure. Environ. Health Perspect. 108: 1139-1145.

Waldman, J. M.; Liang, C. S. K. (1993) Penetration and neutralization of acid aerosols at institutional settings. In: Jantunen, M.; Kalliokoski, P.; Kukkonen, E.; Saarela, K.; Seppänen, O.; Vuorelma, H., eds. Indoor air '93: proceedings of the 6th international conference on indoor air quality and climate. Volume 3: combustion products, risk assessment, policies; July; Helsinki, Finland. Helsinki, Finland: Indoor Air '93; pp. 263-268.

Waldman, J. M.; Liang, C. S. K.; Menon, P. (1990) Indoor exposures to acidic aerosols at child and elderly care facilities. In: Indoor air '90: precedings of the 5th international conference on indoor air quality and climate, volume 2, characteristics of indoor air; July-August; Toronto, ON, Canada. Ottawa, ON, Canada: International Conference on Indoor Air Quality and Climate, Inc.; pp. 607-612.

Waldman, J. M.; Lioy, P. J.; Greenberg, A.; Butler, J. P. (1991) Analysis of human exposure to benzo(a)pyrene via inhalation and food ingestion in the Total Human Environmental Exposure Study (THEES). J. Exposure Anal. Environ. Epidemiol. 1: 193-225.

Wallace, L. (1996) Indoor particles: a review. J. Air Waste Manage. Assoc. 46: 98-126.

Wallace, L. (2000a) Real-time monitoring of particles, PAH, and CO in an occupied townhouse. Appl. Occup. Environ. Hyg. 15: 39-47.

Wallace, L. (2000b) Correlations of personal exposure to particles with outdoor air measurements: a review of recent studies. Aerosol Sci. Technol. 32: 15-25.

Wallace, L.; Quackenboss, J.; Rodes, C. (1997) Continuous measurements of particles, PAH, and CO in an occupied townhouse in Reston, VA. In: Measurement of toxic and related air pollutants. Volume II: proceedings of a specialty conference; April-May; Research Triangle Park, NC. Pittsburgh, PA: Air & Waste Management Association, pp. 860-871. (A&WMA publication VIP-74).

Watson, J. G.; Cooper, J. A.; Huntzicker, J. J. (1984) The effective variance weighting for least squares calculations applied to the mass balance receptor model. Atmos. Environ. 18: 1347-1355.

Watson, J. G.; Robinson, N. F.; Chow, J. C.; Henry, R. C.; Kim, B. M.; Pace, T. G.; Meyer, E. L.; Nguyen, Q. (1990) The USEPA/DRI chemical mass balance receptor model, CMB 7.0. Environ. Software 5: 38-49.

Watson, J. G.; Chow, J. C.; Pace, T. G. (1991) Chemical mass balance. In: Hopke, P. K., ed. Receptor modeling for air quality management. New York, NY: Elsevier; pp. 83-116. (Data handling in science and technology—volume 7).

Weschler, C. J.; Shields, H. C. (1999) Indoor ozone/terpene reactions as a source of indoor particles. Atmos. Environ. 33: 2301-2312.

Wheeler, A. J.; Williams, I.; Beaumont, R. A.; Hamilton, R. S. (2000) Characterisation of particulate matter sampled during a study of children's personal exposure to airborne particulate matter in a UK urban environment. Environ. Monit. Assess. 65: 69-77.

White, W. H. (1998) Statistical considerations in the interpretation of size-resolved particulate mass data. J. Air Waste Manage. Assoc. 48: 454-458.

Wiener, R. W. (1988) Measurement and evaluation of personal exposure to aerosols. In: Measurement of toxic and related air pollutants: proceedings of the 1988 EPA/APCA international symposium; May; Research Triangle Park, NC. Pittsburgh, PA: Air Pollution Control Association; pp. 84-88. (APCA publication VIP-10).

Wiener, R. W. (1989) Particle total exposure assessment methodology--an overview of planning and accomplishments. In: Measurement of toxic and related air pollutants: proceedings of the 1989 U.S. EPA/A&WMA international symposium; May; Raleigh, NC. Pittsburgh, PA: Air & Waste Management Association; pp. 442-448. (A&WMA publication VIP-13).

Wiener, R. W.; Wallace, L.; Pahl, D.; Pellizzari, E.; Whittaker, D.; Spengler, J.; Özkaynak, H. (1990) Review of the particle TEAM 9 home field study. In: Measurement of toxic and related air pollutants: proceedings of the 1990 EPA/A&WMA international symposium, v. 2; May; Raleigh, NC. Pittsburgh, PA: Air & Waste Management Association; pp. 452-460. (A&WMA publication VIP-17).

Wiley, J. A.; Robinson, J. P.; Piazza, T.; Garrett, K.; Cirksena, K.; Cheng, Y.-T.; Martin, G. (1991a) Activity patterns of California residents. Final report. Sacramento, CA: California Air Resources Board; report no. ARB/R93/487. Available from: NTIS, Springfield, VA.; PB94-108719.

Wiley, J. A.; Robinson, J. P.; Cheng, Y.-T.; Piazza, T.; Stork, L.; Pladsen, K. (1991b) Study of children's activity patterns: final report. Sacramento, CA: California Air Resources Board; report no. ARB-R-93/489.

Williams, R.; Suggs, J.; Zweidinger, R.; Evans, G.; Creason, J.; Kwok, R.; Rodes, C.; Lawless, P.; Sheldon, L. (2000a) The 1998 Baltimore Particulate Matter Epidemiology-Exposure Study: part 1. Comparison of ambient, residential outdoor, indoor and apartment particulate matter monitoring. J. Exposure Anal. Environ. Epidemiol. 10: 518-532.

Williams, R.; Suggs, J.; Creason, J.; Rodes, C.; Lawless, P.; Kwok, R.; Zweidinger, R.; Sheldon, L. (2000b) The 1998 Baltimore particulate matter epidemiology-exposure study: part 2. Personal exposure assessment associated with an elderly study population. J. Exposure Anal. Environ. Epidemiol.: 10: 533-543.

Williams, R.; Creason, J.; Zweidinger, R.; Watts, R.; Sheldon, L.; Shy, C. (2000c) Indoor, outdoor, and personal exposure monitoring of particulate air pollution: the Baltimore elderly epidemiology-exposure pilot study. Atmos. Environ. 34: 4193-4204.

Wilson, W. E.; Suh, H. H. (1997) Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies. J. Air Waste Manage. Assoc. 47: 1238-1249.

Wilson, W. E.; Mage, D. T.; Grant, L. D. (2000) Estimating separately personal exposure to ambient and nonambient particulate matter for epidemiology and risk assessment: why and how. J. Air Waste Manage. Assoc. 50: 1167-1183.

Yakovleva, E.; Hopke, P. K.; Wallace, L. (1999) Receptor modeling assessment of particle total exposure assessment methodology data. Environ. Sci. Technol. 33: 3645-3652.

Zeger, S. L.; Thomas, D.; Dominici, F.; Samet, J. M.; Schwartz, J.; Dockery, D.; Cohen, A. (2000) Exposure measurement error in time-series studies of air pollution: concepts and consequences. Environ. Health Perspect. 108: 419-426.

Zmirou, D.; Masclet, P.; Boudet, C.; Dor, F.; Déchenaux, J. (2000) Personal exposure to atmospheric polycyclic aromatic hydrocarbons in a general adult population and lung cancer risk assessment. J. Occup. Environ. Med. 42: 121-126.