




Distribution of Phytoplankton in Iowa Lakes

Working
Paper 695



DISTRIBUTION OF PHYTOPLANKTON IN IOWA LAKES

by

L. R. Williams, S. C. Hern, V. W. Lambou
F. A. Morris*, M. K. Morris*, and W. D. Taylor

Water and Land Quality Branch
Monitoring Operations Division
Environmental Monitoring and Support Laboratory
Las Vegas, Nevada 89114

*Department of Biological Sciences
University of Nevada, Las Vegas
Las Vegas, Nevada 89154

WORKING PAPER NO. 695

NATIONAL EUTROPHICATION SURVEY
OFFICE OF RESEARCH AND DEVELOPMENT
U.S. ENVIRONMENTAL PROTECTION AGENCY

November 1978

DISCLAIMER

This report has been reviewed by the Environmental Monitoring and Support Laboratory-Las Vegas, U.S. Environmental Protection Agency, and approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

FOREWORD

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nationwide threat of accelerated eutrophication to freshwater lakes and reservoirs. The Survey was designed to develop, in conjunction with State environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and State management practices relating to point source discharge reduction and nonpoint source pollution abatement in lake watershed.

The Survey collected physical, chemical, and biological data from 815 lakes and reservoirs throughout the contiguous United States. To date, the Survey has yielded more than two million data points. In-depth analyses are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's freshwater lakes.

CONTENTS

| | <u>Page</u> |
|---|-------------|
| Foreword | iii |
| Introduction | 1 |
| Materials and Methods | 3 |
| Lake and Site Selection | 3 |
| Sample Preparation | 3 |
| Examination | 4 |
| Quality Control | 5 |
| Results | 6 |
| Nygaard's Trophic State Indices | 6 |
| Palmer's Organic Pollution Indices | 6 |
| Species Diversity and Abundance Indices | 8 |
| Species Occurrence and Abundance | 10 |
| Literature Cited | 11 |
| Appendix A. Phytoplankton Species list for the State of Iowa | 12 |
| Appendix B. Summary of Phytoplankton Data | 16 |

INTRODUCTION

The collection and analysis of phytoplankton data were included in the National Eutrophication Survey in an effort to determine relationships between algal characteristics and trophic status of individual lakes.

During spring, summer, and fall of 1974, the Survey sampled 179 lakes in 10 States. Over 700 algal species and varieties were identified and enumerated from the 573 water samples examined.

This report presents the species and abundance of phytoplankton in the 15 lakes sampled in the State of Iowa (Table 1). The Nygaard's Trophic State (Nygaard 1949), Palmer's Organic Pollution (Palmer 1969), and species diversity and abundance indices are also included.

TABLE 1. LAKES SAMPLED IN THE STATE OF IOWA

| STORET No. | Lake Name | County |
|------------|---------------------|-------------------------|
| 1901 | Lake Ahquabi | Warren |
| 1902 | Big Creek Reservoir | Polk |
| 1903 | Black Hawk Lake | Sac |
| 1904 | Clear Lake | Cerro Gordo |
| 1905 | Lake Darling | Washington |
| 1906 | Lost Island Lake | Palo Alto |
| 1907 | Lake McBride | Johnson |
| 1908 | Prairie Rose Lake | Shelby |
| 1909 | Rathbun Reservoir | Appanoose, Wayne, Lucas |
| 1910 | Red Rock Reservoir | Marion |
| 1911 | Rock Creek Lake | Jasper |
| 1912 | Silver Lake | Worth |

(Continued)

TABLE 1. LAKES SAMPLED IN THE STATE OF IOWA (Continued)

| STORET No. | Lake Name | County |
|------------|-------------------|------------|
| 1913 | Spirit Lake | Dickinson |
| 1914 | Viking Lake | Montgomery |
| 1915 | West Lake Okoboji | Dickinson |

MATERIALS AND METHODS

LAKE AND SITE SELECTION

Lakes and reservoirs included in the Survey were selected through discussions with State water pollution agency personnel and U.S. Environmental Protection Agency Regional Offices (U.S. Environmental Protection Agency 1975). Screening and selection strongly emphasized lakes with actual or potential accelerated eutrophication problems. As a result, the selection was limited to lakes:

- (1) impacted by one or more municipal sewage treatment plant outfalls either directly into the lake or by discharge to an inlet tributary within approximately 40 kilometers of the lake;
- (2) 40 hectares or larger in size; and
- (3) with a mean hydraulic retention time of at least 30 days.

Specific selection criteria were waived for some lakes of particular State interest.

Sampling sites for a lake were selected based on available information on lake morphometry, potential major sources of nutrient input, and on-site judgment of the field limnologist (U.S. Environmental Protection Agency 1975). Primary sampling sites were chosen to reflect the deepest portion of each major basin in a test lake. Where many basins were present, selection was guided by nutrient source information on hand. At each sampling site, a depth-integrated phytoplankton sample was taken. Depth-integrated samples were uniform mixtures of water from the surface to a depth of 15 feet (4.6 meters) or from the surface to the lower limit of the photic zone representing 1 percent of the incident light, whichever was greater. If the depth at the sampling site was less than 15 feet (4.6 meters), the sample was taken from just off the bottom to the surface. Normally, a lake was sampled three times in 1 year, providing information on spring, summer, and fall conditions.

SAMPLE PREPARATION

To preserve the sample 4 milliliters (ml) of Acid-Lugol's solution (Prescott 1970) were added to each 130-ml sample from each site at the time of collection. The samples were shipped to the Environmental Monitoring and Support Laboratory, Las Vegas, Nevada, where equal volumes from each site

were mixed to form two 130-ml composite samples for a given lake. One composite sample was put into storage and the other was used for the examination.

Prior to examination, the composite samples were concentrated by the settling method. Solids were allowed to settle for at least 24 hours prior to siphoning off the supernate. The volume of the removed supernate and the volume of the remaining concentrate were measured and concentrations determined. A small (8-ml) library subsample of the concentrate was then taken. The remaining concentrate was gently agitated to resuspend the plankton and poured into a capped, graduated test tube. If a preliminary examination of a sample indicated the need for a more concentrated sample, the contents of the test tube were further concentrated by repeating the settling method. Final concentrations varied from 15 to 40 times the original.

Permanent slides were prepared from concentrated samples after analysis was complete. A ring of clear Karo® corn syrup with phenol (a few crystals of phenol were added to each 100 ml of syrup) was placed on a glass slide. A drop of superconcentrate from the bottom of the test tube was placed in the ring. This solution was thoroughly mixed and topped with a coverglass. After the syrup at the edges of the coverglass had hardened, the excess was scraped away and the mount was sealed with clear fingernail polish. Permanent diatom slides were prepared by drying sample material on a coverglass, heating in a muffle furnace at 400° C for 45 minutes, and mounting in Hyrax®. Finally, the mounts were sealed with clear fingernail polish.

Backup samples, library samples, permanent sample slides, and Hyrax®-mounted diatom slides are being stored and maintained at the Environmental Monitoring and Support Laboratory-Las Vegas.

EXAMINATION

The phytoplankton samples were examined with the aid of binocular compound microscopes. A preliminary examination was performed to precisely identify and list all forms encountered. The length of this examination varied depending on the complexity of the sample. An attempt was made to find and identify all of the forms present in each sample. Often forms were observed which could not be identified to species or to genus. Abbreviated descriptions were used to keep a record of these forms (e.g., lunate cell, blue-green filament, Navicula #1). Diatom slides were examined using a standard light microscope. If greater resolution was essential to accurately identify the diatoms, a phase-contrast microscope was used.

After the species list was compiled, phytoplankton were enumerated using a Neubauer Counting Chamber with a 40X objective lens and a 10X ocular lens. All forms within each field were counted. The count was continued until a minimum of 100 fields had been viewed, or until the dominant form had been observed a minimum of 100 times.

® Registered trademark

QUALITY CONTROL

Project phycologists performed internal quality control intercomparisons regularly on 7 percent of the species identification and counts. Although an individual had primary responsibility for analyzing a sample, taxonomic problems were discussed among the phycologists.

Additional quality control checks were performed on the Survey samples by Dr. G. W. Prescott of the University of Montana at the rate of 5 percent. Quality control checks were made on 75 percent of these samples to verify species identifications while checks were made on the remaining 25 percent of the samples to verify genus counts. Presently, the agreement between quality control checks for species identification and genus enumerations is satisfactory.

RESULTS

A phytoplankton species list for the State is presented in Appendix A. Appendix B summarizes all of the phytoplankton data collected from the State by the Survey. The latter is organized by lake, and includes an alphabetical phytoplankton species list with concentrations for individual species given by sampling date. Results from the application of several indices are presented (Nygaard's Trophic State, Palmer's Organic Pollution, and species diversity and abundance). Each lake has been assigned a four-digit STORET number. (STORET (STOrage and RETrieval) is the U.S. Environmental Protection Agency's computer system which processes and maintains water quality data.) The first two digits of the STORET number identify the State; the last two digits identify the lake.

NYGAARD'S TROPHIC STATE INDICES

Five indices devised by Nygaard (1949) were proposed under the assumption that certain algal groups are indicative of levels of nutrient enrichment. These indices were calculated in order to aid in determining the surveyed lakes' trophic status. As a general rule, Cyanophyta, Euglenophyta, centric diatoms, and members of the Chlorococcales are found in waters that are eutrophic (rich in nutrients), while desmids and many pennate diatoms generally cannot tolerate high nutrient levels and so are found in oligotrophic waters (poor in nutrients).

In applying the indices to the Survey data, the number of taxa in each major group was determined from the species list for each sample. The ratios of these groups give numerical values which can be used as a biological index of water richness. The five indices and the ranges of values established for Danish Lakes by Nygaard for each trophic state are presented in Table 2. The appropriate symbol, (E) eutrophic and (O) oligotrophic, follows each calculated value in the tables in Appendix B. A question mark (?) following a calculated value in these tables was entered when that value was within the range of both classifications.

PALMER'S ORGANIC POLLUTION INDICES

Palmer (1969) analyzed reports from 165 authors and developed algal pollution indices for use in rating water samples with high organic pollution. Two lists of organic-pollution-tolerant forms were prepared, one containing 20 genera, the other, 20 species (Tables 3 and 4). Each form was assigned a pollution index number ranging from 1 for moderately tolerant forms to 6 for

TABLE 2. NYGAARD'S TROPHIC STATE INDICES ADAPTED FROM HUTCHINSON (1967)

| Index | Calculation | Oligotrophic | Eutrophic |
|---------------|---|--------------|-----------|
| Myxophycean | <u>Myxophyceae</u> Desmideae | 0.0-0.4 | 0.1-3.0 |
| Chlorophycean | <u>Chlorococcales</u> Desmideae | 0.0-0.7 | 0.2-9.0 |
| Diatom | <u>Centric Diatoms</u> Pennate Diatoms | 0.0-0.3 | 0.0-1.75 |
| Euglenophyte | <u>Euglenophyta</u> <u>Myxophyceae + Chlorococcales</u> | 0.0-0.2 | 0.0-1.0 |
| Compound | <u>Myxophyceae + Chlorococcales +</u> <u>Centric Diatoms + Euglenophyta</u> Desmideae | 0.0-1.0 | 1.2-25 |

TABLE 3. ALGAL GENUS POLLUTION INDEX
(Palmer 1969)

| Genus | Pollution Index |
|-----------------------|-----------------|
| <u>Anacystis</u> | 1 |
| <u>Ankistrodesmus</u> | 2 |
| <u>Chlamydomonas</u> | 4 |
| <u>Chlorella</u> | 3 |
| <u>Closterium</u> | 1 |
| <u>Cyclotella</u> | 1 |
| <u>Euglena</u> | 5 |
| <u>Gomphonema</u> | 1 |
| <u>Lepocinclis</u> | 1 |
| <u>Melosira</u> | 1 |
| <u>Micractinium</u> | 1 |
| <u>Navicula</u> | 3 |
| <u>Nitzschia</u> | 3 |
| <u>Oscillatoria</u> | 5 |
| <u>Pandorina</u> | 1 |
| <u>Phacus</u> | 2 |
| <u>Phormidium</u> | 1 |
| <u>Scenedesmus</u> | 4 |
| <u>Stigeoclonium</u> | 2 |
| <u>Synedra</u> | 2 |

TABLE 4. ALGAL SPECIES POLLUTION INDEX (Palmer 1969)

| Species | Pollution Index |
|--------------------------------|-----------------|
| <u>Ankistrodesmus falcatus</u> | 3 |
| <u>Arthrospira jenneri</u> | 2 |
| <u>Chlorella vulgaris</u> | 2 |
| <u>Cyclotella meneghiniana</u> | 2 |
| <u>Euglena gracilis</u> | 1 |
| <u>Euglena viridis</u> | 6 |
| <u>Gomphonema parvulum</u> | 1 |
| <u>Melosira varians</u> | 2 |
| <u>Navicula cryptocephala</u> | 1 |
| <u>Nitzschia acicularis</u> | 1 |
| <u>Nitzschia palea</u> | 5 |
| <u>Oscillatoria chlorina</u> | 2 |
| <u>Oscillatoria limosa</u> | 4 |
| <u>Oscillatoria princeps</u> | 1 |
| <u>Oscillatoria putrida</u> | 1 |
| <u>Oscillatoria tenuis</u> | 4 |
| <u>Pandorina morum</u> | 3 |
| <u>Scenedesmus quadricauda</u> | 4 |
| <u>Stigeoclonium tenue</u> | 3 |
| <u>Synedra ulna</u> | 3 |

extremely tolerant forms. Palmer based the index numbers on occurrence records and/or where emphasized by the authors as being especially tolerant of organic pollution.

In analyzing a water sample, any of the 20 genera or species of algae present in concentrations of 50 per milliliter or more are recorded. The pollution index numbers of the algae present are totaled, providing a genus score and a species score. Palmer determined that a score of 20 or more for either index can be taken as evidence of high organic pollution, while a score of 15 to 19 is taken as probable evidence of high organic pollution. Lower figures suggest that the organic pollution of the sample is not high, that the sample is not representative, or that some substance or factor interfering with algal persistence is present and active.

SPECIES DIVERSITY AND ABUNDANCE INDICES

"Information content" of biological samples is being used commonly by biologists as a measure of diversity. Diversity in this connection means the degree of uncertainty attached to the specific identity of any randomly selected individual. The greater the number of taxa and the more equal their proportions, the greater the uncertainty, and hence, the diversity (Pielou 1966). There are several methods of measuring diversity, e.g., the formulas given by Brillouin (1962) and Shannon and Weaver (1963). The method which is appropriate depends on the type of biological sample on hand.

Pielou (1966) classifies the types of biological samples and gives the measure of diversity appropriate for each type. The Survey phytoplankton samples are what she classifies as larger samples (collections in Pielou's terminology) from which random subsamples can be drawn. According to Pielou, the average diversity per individual (H) for these types of samples can be estimated from the Shannon-Wiener formula (Shannon and Weaver 1963):

$$H = -\sum_{i=1}^S p_i \log_x p_i$$

where P is the proportion of the i th taxon in the sample, which is calculated from n_i/N ; n_i is the number of individuals per milliliter of the i th taxon; N is the total number of individuals per ml; and S is the total number of taxa. However, Basharin (1959) and Pielou (1966) have pointed out that H calculated from the subsample is a biased estimator of the sample H , and if this bias is to be accounted for, we must know the total number of taxa present in the sample since the magnitude of this bias depends on it.

Pielou (1966) suggests that if the number of taxa in the subsample falls only slightly short of the number in the larger sample, no appreciable error will result in considering S , estimated from the subsample, as being equal to the sample value. Even though considerable effort was made to find and identify all taxa, the Survey samples undoubtedly contain a fair number of rare phytoplankton taxa which were not encountered.

In the Shannon-Wiener formula, an increase in the number of taxa and/or an increase in the evenness of the distribution of individuals among taxa will increase the average diversity per individual from its minimal value of zero. Sager and Hasler (1969) found that the richness of taxa was of minor importance in determination of average diversity per individual for phytoplankton and they concluded that phytoplankton taxa in excess of the 10 to 15 most abundant ones have little effect on H. This was verified by our own calculations. Our counts are in number per milliliter and since logarithms to the base 2 were used in our calculations, H is expressed in units of bits per individual. When individuals of a taxon were so rare that they were not counted, a value of 1/130 per milliliter or 0.008 per milliliter was used in the calculations since at least one individual of the taxon must have been present in the collection.

A Survey sample for a given lake represents a composite of all phytoplankton collected at different sampling sites on the lake during a given sampling period. Since the number of samples (M) making up a composite is a function of both the complexity of the lake sampled and its size, it should affect the richness-of-taxa component of the diversity of our phytoplankton collections. The maximum diversity (MaxH) (i.e., when the individuals are distributed among the taxa as evenly as possible) was estimated from $\log_2 S$ (Pielou 1966), while the minimum diversity (MinH), was estimated from the formula:

$$\text{MinH} = -\frac{S-1}{N} \log_2 \frac{1}{N} - \left[\frac{N - (S-1)}{N} \right] \log_2 \left[\frac{N - (S-1)}{N} \right]$$

given by Zand (1976). The total diversity (D) was calculated from H_N (Pielou 1966). Also given in Appendix B are L (the mean number of individuals per taxa per milliliter) and K (the number of individuals per milliliter of the most abundant taxon in the sample).

The evenness component of diversity (J) was estimated from H/MaxH (Pielou 1966). Relative evenness (RJ) was calculated from the formula:

$$RJ = \frac{H-\text{MinH}}{\text{MaxH}-\text{MinH}}$$

given by Zand (1976). Zand suggests that RJ be used as a substitute for both J and the redundancy expression given by Wilhm and Dorris (1968). As pointed out by Zand, the redundancy expression given by Wilhm and Dorris does not properly express what it is intended to show, i.e., the position of H in the range between MaxH and MinH. RJ may range from 0 to 1; being 1 for the most even samples and 0 for the least even samples.

Zand (1976) suggests that diversity indices be expressed in units of "sits", i.e., in logarithms to base S (where S is the total number of taxa in the sample) instead of in "bits", i.e., in logarithms to base 2. Zand points out that the diversity index in sits per individual is a normalized number ranging from 1 for the most evenly distributed samples to 0 for the least evenly distributed samples. Also, it can be used to compare different samples, independent of the number of taxa in each. The diversity in bits per

individual should not be used in direct comparisons involving various samples which have different numbers of taxa. Since MaxH equals $\log S$, the expression in sites is equal to $\log S$, or 1. Therefore diversity in sites per individual is numerically equivalent to J, the evenness component for the Shannon-Wiener formula.

SPECIES OCCURRENCE AND ABUNDANCE

The alphabetic phytoplankton species list for each lake, presented in Appendix B, gives the concentrations of individual species by sampling date. Concentrations are in cells, colonies, or filaments (CEL, COL, FIL) per milliliter. An "X" after a species name indicates that the species identified in the preliminary examination was in such a low concentration that it did not appear in the count. A blank space indicates that the organism was not found in the sample collected on that date. Column S is used to designate the examiner's subjective opinion of the five dominant taxa in a sample, based upon relative size and concentration of the organism. The percent column (%C) presents, by abundance, the percentage composition of each taxon.

LITERATURE CITED

- Basharin, G. P. 1959. On a statistical estimate for the entropy of a sequence of independent random variables, pp. 333-336. In: Theory of Probability and Its Applications (translation of "Teoriya Veroyatnosei i ee Premeneniya"). N. Artin (ed). 4. Society for Industrial and Applied Mathematics, Philadelphia.
- Brillouin, L. 1962. Science and Information Theory (2nd ed.). Academic Press, New York. 351 pp.
- Hutchinson, G. E. 1967. A Treatise on Limnology. II. Introduction to Lake Biology and the Limnoplankton. John Wiley and Sons, Inc., New York. 1,115 pp.
- Nygaard, G. 1949. Hydrobiological studies of some Danish ponds and lakes. II. (K danske Vidensk. Selsk.) Biol. Sci. 7:293.
- Palmer, C. M. 1969. A composite rating of algae tolerating organic pollution. J. Phycol. 5:78-82.
- Pielou, E. C. 1966. The measurement of diversity in different types of biological collections. J. Theor. Biol. 13:131-144.
- Prescott, G. W. 1970. How to Know the Freshwater Algae. William C. Brown Company, Dubuque. 348 pp.
- Sager, P. E., and A. D. Hasler. 1969. Species diversity in lacustrine phytoplankton. I. The components of the index of diversity from Shannon's formula. Amer. Natur. 103(929):51-59.
- Shannon, C. E., and W. Weaver. 1963. The Mathematical Theory of Communication. University of Illinois Press, Urbana. 117 pp.
- U.S. Environmental Protection Agency. 1975. National Eutrophication Survey Methods 1973-1976. Working Paper No. 175. Environmental Monitoring and Support Laboratory, Las Vegas, Nevada, and Corvallis Environmental Research Laboratory, Corvallis, Oregon. 91 pp.
- Withm, V. L., and T. C. Dorris. 1968. Biological parameters for water quality criteria. Bio-Science. 18:477.
- Zand, S. M. 1976. Indexes associated with information theory in water quality. J. Water Pollut. Contr. Fed. 48(8):2026-2031.

APPENDIX A
PHYTOPLANKTON SPECIES FOR THE STATE OF IOWA

| | |
|---|---|
| <i>Achmanthes</i> sp. | <i>Cosmarium clepsydra</i> |
| <i>Actinastrum hantzschii</i> | v. <i>nanum</i> |
| <i>Amphora</i> sp. | <i>Crucigenia apiculata</i> |
| <i>Anabaena plantonica</i> | <i>Crucigenia quadrata</i> |
| <i>Anabaena spiroides</i> | <i>Crucigenia rectangularis</i> |
| <i>Anabaenopsis circularis</i> | <i>Crucigenia tetrapedia</i> |
| <i>Anabaenopsis raciborskii</i> | <i>Cryptomonas erosa</i> |
| <i>Ankistrodesmus falcatus</i> | <i>Cryptomonas marssonii</i> |
| <i>Aphanizomenon flos-aquae</i> | <i>Cryptomonas ovata</i> |
| <i>Aphanizomenon flos-aquae</i> v. <i>gracile</i> | <i>Cryptomonas reflexa</i> |
| <i>Aphanocapsa delicatissima</i> | <i>Cyclotella meneghiniana</i> |
| <i>Aphanocapsa elachista</i> | <i>Cylindrospermum stagnale</i> ? |
| <i>Aphanocapsa elachista</i> v. <i>planctonica</i> | <i>Cymatopleura elliptica</i> |
| <i>Aphanothecce castagnei</i> | <i>Cymatopleura solea</i> |
| <i>Aphanothecce clathrata</i> | <i>Cymbella affinis</i> |
| <i>Arthrodesmus</i> sp. | <i>Cymbella triangulum</i> |
| <i>Asterionella formosa</i> | <i>Dactylococcopsis irregularis</i> |
| <i>Botryococcus braunii</i> | <i>Diatoma vulgare</i> |
| <i>Caloneis amphisbaena</i> | <i>Dictyosphaerium pulchellum</i> |
| <i>Caloneis lewisii</i> v. <i>inflata</i> | <i>Dinobryon cylindricum</i> |
| <i>Carteria</i> sp. | <i>Dinobryon divergens</i> |
| <i>Ceratium hirundinella</i> | <i>Dinobryon sertularia</i> |
| <i>Ceratium hirundinella</i> v. <i>furcoides</i> | <i>Dinobryon sociale</i> |
| <i>Ceratium hirundinella</i> v. <i>robustum</i> | <i>Elakothrix gelatinosa</i> |
| <i>Chilomonas</i> ? <i>paramaecium</i> | <i>Epithemia turgida</i> |
| <i>Chlamydomonas pseudopertyi</i> | <i>Errerella</i> ? <i>bornhemiensis</i> |
| <i>Chlorogonium</i> sp. | <i>Euastrum denticulatum</i> |
| <i>Chroococcus dispersus</i> | <i>Eudorina elegans</i> |
| <i>Chroococcus dispersus</i> v. <i>minor</i> | <i>Euglena acus</i> |
| <i>Chroococcus limneticus</i> | <i>Euglena deses</i> |
| <i>Chroomonas nordstedtii</i> | <i>Euglena gracilis</i> |
| <i>Closteriopsis longissima</i> | <i>Euglena oxyuris</i> |
| <i>Closterium gracile</i> ? | <i>Euglena sciotensis</i> ? |
| <i>Cocconeis placentula</i> | <i>Euglena spiroides</i> |
| <i>Cocconeis placentula</i> v. <i>euglypta</i> | <i>Euglena tripteris</i> |
| <i>Coelastrum cambricum</i> | <i>Eunotia gracilis</i> |
| <i>Coelastrum cambricum</i> v. <i>intermedium</i> | <i>Fragilaria construens</i> |
| <i>Coelastrum reticulatum</i> | <i>Fragilaria cotonensis</i> |
| <i>Coelastrum sphaericum</i> | <i>Fragilaria intermedia</i> |
| <i>Coelosphaerium minutissimum</i> | <i>Franceia</i> sp. |
| <i>Coelosphaerium naegelianum</i> | <i>Glenodinium aciculiferum</i> |
| | <i>Glenodinium gymnodinium</i> |
| | v. <i>biscutelliforme</i> |
| | <i>Glenodinium oculatum</i> |
| | <i>Gloeocystis vesiculosa</i> |
| | <i>Golenkinia</i> sp. |
| | <i>Gomphonema acuminatum</i> |
| | v. <i>coronata</i> |

| | |
|---|--|
| <i>Gomphonema constrictum</i> | <i>Pediastrum duplex</i> |
| <i>Gomphonema parvulum</i> | <i>Pediastrum duplex</i> |
| <i>Gomphosphaeria</i> | v. <i>clathratum</i> |
| <i>Gymnodinium album</i> | <i>Pediastrum duplex</i> |
| <i>Gymnodinium ordinatum</i> | v. <i>reticulatum</i> |
| <i>Gyrosigma</i> sp. | <i>Pediastrum kawraiskyi</i> |
| <i>Kirchneriella elongata</i> | <i>Pediastrum simplex</i> |
| <i>Kirchneriella lunaris</i> | <i>Pediastrum tetras</i> |
| v. <i>irregularis</i> | <i>Pediastrum tetras</i> |
| <i>Lagerheimia citriformis</i> | v. <i>tetraodon</i> |
| <i>Lepocinclus acuta</i> ? | <i>Peridinium cinctum</i> |
| <i>Lepocinclus butschlii</i> | <i>Peridinium inconspicuum</i> |
| <i>Lyngbya circumcreta</i> | <i>Peridinium quadrident</i> |
| <i>Lyngbya limnetica</i> | <i>Peridinium umbonatum</i> |
| <i>Mallomonas acaroides</i> | <i>Phacus acuminatus</i> |
| <i>Mastogloia</i> sp. | <i>Phacus caudatus</i> |
| <i>Melosira distans</i> | <i>Phacus longicauda</i> |
| <i>Melosira granulata</i> | <i>Phacus megalopsis</i> |
| <i>Melosira granulata</i> | <i>Phacus nordstedtii</i> |
| v. <i>angustissima</i> | <i>Phacus orbicularis</i> ? |
| <i>Melosira granulata</i> | <i>Phacus pleuronectes</i> |
| v. <i>angustissima</i> f. <i>spiralis</i> | <i>Phacus pseudonordstedtii</i> |
| <i>Melosira islandica</i> | <i>Phacus pyrum</i> |
| <i>Melosira varians</i> | <i>Phacus tortus</i> |
| <i>Merismopedia minima</i> | <i>Phormidium mucicola</i> |
| <i>Merismopedia punctata</i> | <i>Pinnularia</i> sp. |
| <i>Merismopedia tenuissima</i> | <i>Polyedriopsis</i> sp. |
| <i>Mesostigma viridis</i> | <i>Pteromonas angulosa</i> |
| <i>Microcystis aeruginosa</i> | <i>Quadrigula</i> sp. |
| <i>Microcystis incerta</i> | <i>Raphidiopsis curvata</i> |
| <i>Microcystis stagnalis</i> | <i>Rhoicosphenia</i> sp. |
| <i>Navicula cryptocephala</i> | <i>Rhopalodia gibba</i> |
| <i>Navicula cuspidata</i> | <i>Scenedesmus abundans</i> |
| <i>Navicula cuspidata</i> | <i>Scenedesmus acuminatus</i> |
| v. <i>ambigua</i> | <i>Scenedesmus arcuatus</i> |
| <i>Navicula exigua</i> | <i>Scenedesmus bicaudatus</i> |
| <i>Navicula zanoni</i> | <i>Scenedesmus bijuga</i> |
| <i>Nitzschia acicularis</i> | v. <i>flexuosus</i> |
| <i>Nitzschia filiformis</i> | <i>Scenedesmus denticulatus</i> |
| <i>Nitzschia holsatica</i> | <i>Scenedesmus dimorphus</i> |
| <i>Nitzschia palea</i> | <i>Scenedesmus granulatus</i> |
| <i>Nostoc</i> sp. | v. <i>longispina</i> f. <i>granulata</i> |
| <i>Oocystis submarina</i> | <i>Scenedesmus intermedius</i> |
| <i>Oscillatoria agardhii</i> | <i>Scenedesmus intermedius</i> |
| <i>Oscillatoria limnetica</i> | v. <i>bicaudatus</i> |
| <i>Oscillatoria subbrevis</i> | <i>Scenedesmus protuberans</i> |
| <i>Pandorina morum</i> | <i>Scenedesmus quadricauda</i> |
| <i>Pediastrum bicaudata</i> | <i>Scenedesmus quadricauda</i> |
| v. <i>longecornutum</i> | v. <i>biornatus</i> f. <i>giganticus</i> |
| <i>Pediastrum boryanum</i> | |

| | |
|---|-------------------------------------|
| <i>Scenedesmus quadricauda</i> | <i>Tetraedron muticum</i> |
| v. <i>quadrispina</i> | <i>Tetraedron regulare</i> |
| <i>Scenedesmus reniforme</i> | <i>Tetraedron regulare</i> |
| <i>Schizochlamys ? compacta</i> | v. <i>incus</i> |
| <i>Schroederia setigera</i> | <i>Tetrastrum elegans</i> |
| <i>Spermatozoopsis</i> sp. | <i>Tetrastrum heteracanthum</i> |
| <i>Sphaerocystis schroeteri</i> | <i>Tetrastrum staurogeniaeforme</i> |
| <i>Spirogyra</i> sp. | <i>Trachelomonas atomaria</i> |
| <i>Staurastrum chaetocerus</i> | <i>Trachelomonas ensifera</i> |
| <i>Staurastrum tetracerum</i> | v. <i>javanica</i> |
| <i>Stauroneis</i> | <i>Trachelomonas fluviatilis</i> |
| <i>Stephanodiscus astraea</i> | <i>Trachelomonas gibberosa</i> |
| <i>Stephanodiscus hantzschii</i> ? | <i>Trachelomonas intermedia</i> |
| <i>Stephanodiscus niagarae</i> | <i>Trachelomonas oblonga</i> |
| <i>Surirella angustata</i> | <i>Trachelomonas planctonica</i> |
| <i>Synedra acus</i> | <i>Trachelomonas pulchella</i> |
| <i>Synedra ulna</i> | <i>Trachelomonas scabra</i> |
| <i>Tetraedron caudatum</i> | <i>Trachelomonas urceolata</i> |
| <i>Tetraedron caudatum</i> v. <i>longecornutum</i> | <i>Trachelomonas volvocina</i> |
| <i>Tetraedron constrictum</i> | <i>Treubaria</i> sp. |
| <i>Tetraedron hastatum</i> | <i>Westella botryoides</i> ? |
| <i>Tetraedron minimum</i> | |
| v. <i>scrobiculatum</i> | |

APPENDIX B. SUMMARY OF PHYTOPLANKTON DATA

This appendix was generated by computer. Because it was only possible to use upper case letters in the printout, all scientific names are printed in upper case and are not italicized.

The alphabetic phytoplankton lists include taxa without species names (e.g., EUNOTIA, EUNOTIA #1, FLAGELLATE, FLAGELLATES, MICROCYSTIS INCERTA ?, CHLOROPHYTAN COCCOID CELLED COLONY). When species determinations were not possible, symbols or descriptive phrases were used to separate taxa for enumeration purposes. Each name on a list, however, represents a unique species different from any other name on the same list, unless otherwise noted, for counting purposes.

Numbers were used to separate unidentified species of the same genus. A generic name listed alone is also a unique species. A question mark (?) is placed immediately after the portion of a name which was assigned with uncertainty. Numbered, questioned, or otherwise designated taxa were established on a lake-by-lake basis; therefore NAVICULA #2 from lake A cannot be compared to NAVICULA #2 from lake B. Pluralized categories (e.g., FLAGELLATES, CENTRIC DIATOMS, SPP.) were used for counting purposes when taxa could not be properly differentiated on the counting chamber.

LAKE NAME: ACQUABI
STORET NUMBER: 1901

NYGAARD TROPHIC STATE INDICES

| DATE | 04 17 74 | 07 02 74 | 09 25 74 |
|---------------|----------|----------|----------|
| MYXOPHYCEAN | 2.00 E | 6.00 E | 3.50 E |
| CHLOROPHYCEAN | 0.01 0 | 1.00 E | 3.00 F |
| EUGLENOPHYTE | 0.50 E | 0.43 E | 0.15 ? |
| DIATOM | 0.50 E | 0.75 E | 2.00 F |
| COMPOUND | 5.01 E | 13.0 E | 9.50 F |

PALMER'S ORGANIC POLLUTION INDICES

| DATE | 04 17 74 | 07 02 74 | 09 25 74 |
|---------|----------|----------|----------|
| GENUS | 05 | 09 | 06 |
| SPECIES | 33 | 22 | 22 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

| DATE | 04 17 74 | 07 02 74 | 09 25 74 |
|----------------------------------|----------|----------|----------|
| AVERAGE DIVERSITY | H | 2.47 | 2.62 |
| NUMBER OF TAXA | S | 14.00 | 21.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 1.00 | 1.00 |
| MAXIMUM DIVERSITY | MAXH | 3.81 | 4.39 |
| MINIMUM DIVERSITY | MINH | 0.11 | 0.13 |
| TOTAL DIVERSITY | D | 3324.62 | 5067.08 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 1346.00 | 1934.00 |
| EVENNESS COMPONENT | J | 0.65 | 0.60 |
| RELATIVE EVENNESS | RJ | 0.64 | 0.59 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 96.14 | 92.10 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 475.00 | 575.00 |
| | | | 769.00 |

| TAXA | FORM | 06 17 74 | | | 07 02 74 | | | 09 25 74 | | |
|-----------------------------|------|----------|------|--------------------|----------|------|--------------------|----------|------|--------------------|
| | | IS | %C | ALGAL UNITS PER ML | IS | %C | ALGAL UNITS PER ML | IS | %C | ALGAL UNITS PER ML |
| ANABAENA | FIL | | | | 11 | 29.7 | 575 | | | |
| ANABAENA #1 | FIL | | | | | | X | 4 | 4.7 | 77 |
| ANABAENA PLANCTONICA | FIL | | | | | | | | | X |
| APHANIZOMENON FLOS-AQUAE | FIL | 4 | 11.7 | 158 | 2.7 | | 52 | 3 | 9.3 | 154 |
| ASTERIONELLA FORMOSA | CEL | 3 | 14.7 | 198 | | | | | | |
| CALONEIS LEWISIT | | | | | | | | | | |
| V. INFLATA | CEL | | | | | | X | | | |
| CENTRIC DIATOM | CEL | 2 | 20.6 | 277 | 4 | 16.2 | 316 | | | |
| CERATIUM HIRUNDINELLA | CEL | | | | | | X | | | |
| CLOSTERIUM #1 | CEL | | | X | | | X | | | X |
| COCCOID CHRYSOPHYTAN | CFL | | | X | | | | | | |
| COELASTRUM CAMBRICUM | COL | | | | | | | | | X |
| COELOSPHAERIUM MINUTISSIMUM | COL | | | | | | | | | X |
| COELOSPHAERIUM NAEGLIANUM | COL | | | | | | | 5 | 2.3 | 38 |
| CRUCIGENIA APICULATA | COL | | | | | | | 5 | 2.3 | 38 |
| CRYPTOMONAS EROSA | CEL | | 5.9 | 79 | | | X | | | |
| CRYPTOMONAS REFLEXA | CEL | | | X | | | | | | |
| EUDORINA ELEGANS | COL | | | | | 2.7 | 52 | | | |
| FLAGELLATE #1 | CEL | | 3.0 | 40 | | | | | | |
| FRAGILARIA | CEL | | | X | | | X | | | |
| LEPOCINCIS | CFL | | | | | | X | | | |
| LYNGBYA | FIL | | | | | | | | | X |
| MALLOMONAS ACAROIDES | CEL | | | X | | | X | | 2.3 | 38 |
| MELOSIRA GRANULATA | CEL | | | | | | | | | X |
| MELOSIRA GRANULATA | | | | | | | | | | |
| V. ANGUSTISSIMA | CEL | | | | | | | 2 | 23.3 | 384 |
| MELOSIRA VARIANS | CEL | | | | | | X | | | |
| MICROCYSTIS AERUGINOSA | COL | | | | 3 | 5.4 | 105 | | | |
| NITZSCHIA #1 | CEL | 15 | 8.8 | 119 | | | | | | |
| NITZSCHIA #2 | CEL | | | | | | | | | X |
| NITZSCHIA #3 | CEL | | | | | | X | | | |
| NITZSCHIA #4 | CEL | | | | | 2.7 | 52 | | | |
| NOSTOC | CEL | | | | | | X | | | |
| ONCYSTIS | CEL | | | | | | | | | X |
| OSCILLATORIA AGARDHII | FIL | | | | | | | 1 | 46.5 | 768 |
| OSCILLATORIA LINNETICA | FIL | | | X | | 2.7 | 52 | | | |
| PEDIASTRUM DUPLEX | | | | | | | | | | |
| V. RETICULATUM | COL | | | | | | | | 2.3 | 38 |
| PTEROMONAS ANGULOSA | CEL | | | | | | | | | X |
| SCENEDESmus ARCUATUS | COL | | | | | | | | | |
| SCHROFDERIA SETIGERA | CEL | | | | 5 | 5.4 | 105 | | | |
| SPHAEROCYSTIS SCHROETERI | COL | | | | | | | | | X |
| STAURASTRUM CHAETOCERUS | CEL | | | | | | | | | X |
| SYNEDRA ULNA | CEL | 11 | 35.3 | 475 | 2.7 | | 52 | | | |
| TRACHELOMONAS ATOMARIA | CEL | | | X | | | | | 4.7 | 77 |
| TRACHELOMONAS INTERMEDIA | CEL | | | | | | | | 2.3 | 38 |
| TRACHELOMONAS OBLONGA | CEL | | | | | | | | | |
| TRACHELOMONAS PLANTONICA | CEL | | | | 2 | 29.7 | 575 | | | |
| TOTAL | | | | | 1346 | | | 1934 | | 1650 |

LAKE NAME: BIG CREEK
STORE NUMBER: 1907

NYGAARD TROPHIC STATE INDICES

DATE 04 18 74 07 09 74 09 25 74

| | | | |
|---------------|--------|--------|--------|
| MYXOPHYCEAN | 0/0 O | 01/1 E | 3.00 E |
| CHLOROPHYCEAN | 02/0 F | 0/0 O | 2.00 E |
| EUGLENOPHYTE | 1.00 E | 1.00 E | 0/05 ? |
| DIATOM | 0.43 E | 0.50 E | 1.01 E |
| COMPOUND | 06/0 E | 03/0 E | 7.00 F |

PALMER'S ORGANIC POLLUTION INDICES

DATE 04 18 74 07 09 74 09 25 74

| | | | |
|---------|----|----|----|
| GENUS | 36 | 33 | 71 |
| SPECIES | 03 | 00 | 00 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE 04 18 74 07 09 74 09 25 74

| | | | | |
|----------------------------------|------|----------|---------|---------|
| AVERAGE DIVERSITY | H | 3.80 | 1.20 | 1.58 |
| NUMBER OF TAXA | S | 19.00 | 7.00 | 20.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 2.00 | 2.00 | 2.00 |
| MAXIMUM DIVERSITY | MAXH | 4.25 | 2.81 | 4.32 |
| MINIMUM DIVERSITY | MINH | 0.02 | 0.06 | 0.14 |
| TOTAL DIVERSITY | N | 13274.40 | 1317.60 | 2629.12 |
| TOTAL NUMBER OF INDIVIDUALS/ML | V | 16593.00 | 1098.00 | 1664.00 |
| EVENNESS COMPONENT | J | 0.19 | 0.43 | 0.37 |
| RELATIVE EVENNESS | RJ | 0.19 | 0.42 | 0.35 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 873.32 | 156.86 | 93.20 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 14887.00 | 595.00 | 1167.00 |

LAKE NAME: BIG CREEK
STORET NUMBER: 1902

CONTINUED

| TAXA | FORM | 04 18 74 | | | 07 09 74 | | | 09 25 74 | | |
|-----------------------------|------|----------|------|--------------------|----------|-------|--------------------|----------|-------|--------------------|
| | | IS | VC | ALGAL UNITS PER ML | IS | VC | ALGAL UNITS PER ML | IS | VC | ALGAL UNITS PER ML |
| ANKistrodeSMUS FALCATUS | CEL | 15 | 1.51 | 250 | | | X | | 2.41 | 40 |
| APHANIZOMENON FLOS-AQUAE | FIL | | | | | | | | | |
| APHAНОTHECE CASTAGNEI | COL | | | | 11 | 54.21 | 595 | | 2.41 | 40 |
| ASTERIONELLA FORMOSA | CEL | | | | | | | | | |
| CERATIUM HIRUNDINELLA | | | | | | | | | | |
| F. ROBUSTUM | CEL | | | | | | | | | |
| CHILOMONAS ? PARAMAECIUM | CEL | | | | | | | | | |
| CHLAMYDOMONAS | CEL | | 1.3 | 208 | | | | | | |
| CHLOROPHYTAN FLAGELLATE | CEL | | | | | | | | | |
| COCconeis PLACENTULA | | | | | | | | | | |
| V. EUGLYPTA | CEL | | | | | | | | | |
| COELASTRUM CAMBRICUM | COL | | | | | | | | | |
| COSMARIA #1 | CEL | | | | | | | | | |
| CRYPTOMONAS #1 | CEL | | | | | | | | | |
| CRYPTOMONAS #2 | CEL | | 0.5 | 83 | | | | | 2.41 | 40 |
| CRYPTOMONAS EROSA | CEL | | | | | | | | | |
| CRYPTOMONAS REFLEXA | CEL | | | | | | | | | |
| CYNATOPLEURA SOLEA | CEL | | | | | | | | | |
| DINORRYON SERTULARIA | CEL | | 0.81 | 125 | | | | | | |
| EUDORINA ELEGANS | COL | | | | | | | | 1.21 | 20 |
| EUGLENA | CEL | | | | | | | | | |
| FLAGELLATE #1 | CEL | 14 | 2.3 | 374 | 31 | 4.21 | 46 | | 2.41 | 40 |
| FLAGELLATE #7 | CEL | 13 | 1.8 | 291 | | | | | | |
| FRAGILARIA CROTONENSIS | CEL | | | | | | | 21 | 15.41 | 257 |
| GLENODINIUM ACICULIFERUM | CFL | | 0.51 | 83 | | | | | | |
| GLENODINIUM OCULATUM | CEL | | 0.3 | 42 | | | | | | |
| GLOEOCYSTIS VESICULOSA | COL | | | | | | | | | |
| COMPHONEMA | CFL | | 0.31 | 42 | | | | | | |
| GYMNODINIUM ORDINATUM | CFL | 12 | 1.0 | 166 | | | | | | |
| MELOSIRA | CEL | | 0.31 | 42 | | | | | | |
| MELOSIRA GRANULATA | CFL | | | | | | | 11 | 70.11 | 1167 |
| NAVICULA | CFL | | | | | | | | | |
| NITZSCHIA #1 | CEL | | | | | | | | | |
| OSCILLATORIA LIMNETICA | FIL | | | | | | | | 1.21 | 20 |
| PANDORINA MORUM | COL | | | | | | | | | |
| PEDIASTRUM DUPLEX | COL | | | | | | | | | |
| PHACUS CRIBULARIS ? | CEL | | | | | | | | | |
| RHOICOSPHEMIA | CEL | | | | | | | | | |
| SCENEDESmus QUADRICAUDA | COL | | | | | | | | | |
| V. QUADRISPINA | | | | | | | | | | |
| STEPHANODISCUS ASTRAEA | CEL | | | | 21 | 41.61 | 457 | 31 | 2.41 | 40 |
| STEPHANODISCUS HANTZSCHII ? | CFL | 11 | 89.7 | 14887 | x | 11 | | 11 | | |
| TRACHELOMONAS | CFL | | | | | | | | | |
| TOTAL | | | | 16593 | | | 1998 | | 1664 | |

LAKE NAME: BLACK HAWK
STORFT NUMBER: 1902

NYGAARD TROPHIC STATE INDICES

| | DATE | 04 19 74 | 07 03 74 | 09 25 74 |
|---------------|------|----------|----------|----------|
| MYXOPHYCEAN | | 04/0 E | 2.00 E | 7.00 E |
| CHLOROPHYCEAN | | 16/0 E | 4.67 E | 6.00 E |
| EUGLENOPHYTE | | 0.10 ? | 0.10 ? | 0.23 E |
| DIATOM | | 0.80 E | 0.22 ? | 1.67 E |
| COMPOUND | | 26/0 E | 9.00 F | 18.5 F |

PALMER'S ORGANIC POLLUTION INDICES

| | DATE | 04 19 74 | 07 03 74 | 09 25 74 |
|---------|------|----------|----------|----------|
| GENUS | | 16 | 06 | 27 |
| SPECIES | | 11 | 04 | 06 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

| | DATE | 04 19 74 | 07 03 74 | 09 25 74 |
|----------------------------------|------|-----------|----------|-----------|
| AVERAGE DIVERSITY | H | 2.77 | 2.38 | 3.05 |
| NUMBER OF TAXA | S | 39.00 | 39.00 | 47.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 2.00 | 2.00 | 2.00 |
| MAXIMUM DIVERSITY MAXH | | 5.29 | 5.29 | 5.55 |
| MINIMUM DIVERSITY MINH | | 0.01 | 0.02 | 0.01 |
| TOTAL DIVERSITY | D | 163690.38 | 60052.16 | 165663.80 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 59094.00 | 25232.00 | 54316.00 |
| EVENNESS COMPONENT | J | 0.52 | 0.45 | 0.55 |
| RELATIVE EVENNESS | RJ | 0.53 | 0.45 | 0.55 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 1515.23 | 646.97 | 1155.66 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 20231.00 | 9323.00 | 28092.00 |

| TAXA | FORM | 04 19 74 | | | 07 03 74 | | | 09 25 74 | | |
|----------------------------|------|----------|------|--------------------|----------|------|--------------------|----------|-------|--------------------|
| | | S | ZC | ALGAL UNITS PER ML | S | ZC | ALGAL UNITS PER ML | S | ZC | ALGAL UNITS PER ML |
| ACTINASTRUM | CEL | 1 | 0.2 | 106 | | | | 3.1 | 1697 | |
| ANABAENA | FIL | | | X | 1.7 | | 417 | 1.31 | 679 | |
| ANABAENOPSIS CIRCULARIS | FIL | | | | | | | 1.41 | 1019 | |
| ANABAENOPSIS RACIBORSKII | FIL | | | | | | | | | |
| ANKISTRODESMUS FALCATUS | CEL | 15 | 4.9 | 2875 | 11 | 36.9 | 9323 | 0.61 | 239 | |
| APHANIZOMENON FLOS-AQUAE | CEL | | | | | | | | | |
| APHANIZOMENON FLOS-AQUAE | FIL | | | | | | | 31 | 4.71 | 2566 |
| V. GRACILE | COL | | | | | | | 0.61 | 339 | |
| APHANOCPASA ELACHISTA | COL | | | | | | | 0.51 | 756 | |
| APHANOTHECE | COL | | | | | | | | | |
| ASTERIONELLA FORMOSA | CEL | | | | | | X | | | |
| CERATIUM HIRUNDINELLA | CEL | | | | | | | | | |
| F. FURCOIDES | CEL | | | | | | X | | | |
| CHLAMYDOMONAS | CFL | | | X | | | | 0.61 | 339 | |
| CHROOCOCCUS DISPERSUS | CFL | | | | | | X | | | |
| CHROOCOCCUS DISPERSUS | COL | | | | | | | | | |
| V. MINOR | COL | | | | | | | | | X |
| CLOSTERIOPSIS | CEL | | 0.4 | 213 | | | | | | |
| CLOSTERIOPSIS ? LONGISSIMA | CEL | | | | | | X | | | |
| CLOSTRIUM | CEL | | | | | | X | | | |
| CLOSTERIUM #1 | CEL | | | | 0.4 | | 93 | | | |
| CLOSTRIUM #2 | CEL | | | | | | | | | X |
| CLOSTRIUM SPP. | CEL | | | | | | | 0.31 | 170 | |
| COCCONEIS | CEL | | | X | | | | | | X |
| COELASTRUM CAMBRICUM | COL | | | X | | 0.21 | 46 | | | |
| COSMARIUM | CFL | | | | | | X | | | |
| CRUCIGENIA TETRAPEDIA | COL | | | | | 4.6 | 1160 | | | |
| CRYPTOMONAS | CEL | | | | | 1.31 | 325 | 0.61 | 130 | |
| CRYPTOMONAS EROSA | CEL | | | X | | | | | | |
| CRYPTOMONAS REFLEXA | CEL | | | X | | | X | | | |
| CRYPTOMONAS SPP. | CEL | 31 | 5.4 | 3194 | | | | | | |
| CYCLOTELLA MENEGHINIANA | CEL | | 1.4 | 852 | | | | | | |
| CYMATOPLEURA SOLEA | CEL | | 0.4 | 213 | | | X | | | |
| CYMBELLA TRIANGULUM | CEL | | | | 15 | 1.5 | 371 | | | |
| DACTYLOCOPCOPSIS | CEL | | 1.6 | 958 | | | | | | |
| DICTYOSPHAERIUM | COL | | | X | | | | 0.51 | 255 | |
| ELAKATOThRIX | CEL | | | | | | | 7.6 | 310 | |
| ELAKATOThRIX GELATINOSA | COL | | | | 0.61 | | 139 | | | |
| EPITHEMIA TURGIDA | CEL | | | | | | X | | | |
| EPREPPELLA ? BORNHEMENSIS | CEL | | | | | | X | | | |
| EUGLENA | CEL | | | X | | | | 0.21 | 85 | |
| FLAGELLATE #1 | CEL | 14 | 13.5 | 7986 | | | | 4.8 | 3140 | |
| FLAGELLATES | CEL | | 3.1 | 1810 | | | | | | |
| KTRCHNERIELLA ? | CEL | | | | | | | 1.31 | 679 | |
| LEPOCINCLIS | CEL | | | | | | | 7.31 | 170 | |
| MELOSIRA #4 | CEL | | 1.1 | 639 | | 0.4 | 93 | | | |
| MELOSIRA GRANULATA | CEL | | | X | | | | | | X |
| MELOSIRA GRANULATA | CEL | | | | | | | | | |
| V. ANGSTISSIMA | CEL | | | | | | | 0.81 | 425 | |
| MEPISOPEDIA MINIMA | COL | | 1.4 | 852 | | | | 151.71 | 24197 | |
| MEPISOPEDIA TENUISSIMA | COL | | | | | | | | | |
| MESOSTIGMA VIRIDIS | CEL | | 0.21 | 106 | | 0.61 | 139 | | | |
| MICROCYSTIS AERUGINOSA | COL | | | | | | | 0.31 | 170 | |
| MICROCYSTIS INCERTA | COL | | | | | | X | 7.81 | 424 | |
| NAVICULA | CEL | | 0.5 | 319 | | | | 14 | 3.61 | 1957 |
| NAVICULA #1 | CEL | | | | | | | | | |
| NITZSCHIA ACICULARIS | CEL | 11 | 34.2 | 20231 | | | X | 0.51 | 255 | |
| NITZSCHIA PALEA | CEL | | 0.7 | 426 | | | | 0.61 | 339 | |
| NOSTOC | CEL | | | | | | X | | | |
| OXYCYSTIS | CEL | | | | | | | | | |
| OSCILLATORIA | CFL | | | X | 3 | 22.2 | 5612 | | | |
| OSCILLATORIA #1 | FIL | | | | | | | 12 | 7.21 | 3924 |
| OSCILLATORIA #2 | FIL | | | | | | | | | X |
| PEDIASTRUM BORYANUM | COL | | | | | | X | | | |
| PEDIASTRUM DUPLEX | COL | | 0.2 | 106 | | | | | | |
| PEDIASTRUM DUPLEX | COL | | | | | | | | | |
| V. RETICULATUM | COL | | | | | | X | | | X |
| PEDIASTRUM KAWRAITSKYI | COL | | | | | | X | | | |
| PEDIASTRUM TETRAS | COL | | | | | | X | | | |
| PHACUS | CEL | | 0.4 | 213 | | | | | | |
| PHACUS CAUDATUS | CEL | | | | | | | 0.21 | 85 | |
| PHACUS MEGALOPLIS | CEL | | | | | | | 0.51 | 255 | |
| PHACUS PYRUM ? | CEL | | | | | | | 0.31 | 170 | |
| PINNULARIA | CEL | | | | | | X | | | |
| QUADRIGULA | CEL | | | X | | | | | | |
| SCENEDESMUS | COL | | | | | | X | | | |
| SCENEDESMUS ABUNDANS | COL | | | | | | | | | X |
| SCENEDESMUS ACUMINATUS | COL | | 0.2 | 106 | | | | | | |
| SCENEDESMUS DIMORPHUS | COL | | 0.4 | 213 | | | X | | | |
| SCENEDESMUS INTERMEDIUS | COL | | 0.7 | 426 | | | | 0.61 | 339 | |
| SCENEDESMUS PROTUBERANS | COL | | | X | | | | | | |

LAKE NAME: BLACK HAWK
STORE NUMBER: 1903 -

CONTINUED

| TAXA | FORM | 04 19 74 | | | 07 03 74 | | | 09 25 74 | | |
|------------------------------|------|----------|------|--------------------|----------|------|--------------------|----------|-------|--------------------|
| | | IS | %C | ALGAL UNITS PER ML | IS | %C | ALGAL UNITS PER ML | IS | %C | ALGAL UNITS PER ML |
| SCENEDESMUS QUADRICAUDA | COL | | | X | | 0.6 | 139 | | | |
| SCENEDESMUS SPP. | COL | | | | | | | 0.5 | 755 | |
| SCHROEDERIA SETIGERA | CEL | | | X | | 0.4 | 93 | | 3.3 | 1762 |
| SPHAEROCYSTIS SCHROETERI | COL | | | | 4 | 25.6 | 6447 | | | |
| STEPHANODISCUS #1 | CEL | | | | | | | 0.3 | 170 | |
| STEPHANODISCUS #2 | CEL | | | | | | | 15 | 4.5 | 2461 |
| STEPHANODISCUS ASTRAEA | CEL | 2 | 28.1 | 16611 | | 2 | 3.3 | 835 | | |
| STEPHANODISCUS NIAGARAE | CFL | | | | | | | X | | |
| SURIRELLA #1 | CEL | | | | | | | X | | |
| SURIRELLA #2 | CFL | | | | | | | | | |
| TETRASTRUM ELEGANS | COL | | | X | | | | | 0.2 | 84 |
| TETRASTRUM HETFRACANTHUM | COL | | | X | | | | | | |
| TETRASTRUM SPP. | COL | 1 | 1.1 | 639 | | | | | | |
| TETRASTRUM STAUROGENTAEFORME | COL | | | X | | | | | | |
| TRACHELOMONAS #1 | CEL | | | | | | | | | |
| TRACHELOMONAS #2 | CEL | | | | | | | | 0.2 | 84 |
| TRACHELOMONAS FLUVIATILIS | CEL | 1 | 1 | | | | X | | | |
| TOTAL | | | | 59094 | | | 25232 | | 54316 | |

LAKE NAME: CLEAR
STORE NUMBER: 1904

NYGAARD TROPHIC STATE INDICES

| | DATE | 04 18 74 | 07 03 74 | 09 23 74 |
|---------------|------|----------|----------|----------|
| MYXOPHYCEAN | | 3.50 E | 3.00 E | 12.0 F |
| CHLOROPHYCEAN | | 5.50 F | 5.50 E | 12.0 F |
| EUGLENOPHYTE | | 0.04 ? | 0/17 ? | 0/24 ? |
| DIATOM | | 0.50 E | 0.43 E | 1.00 E |
| COMPOUND | | 10.5 E | 10.0 F | 26.0 F |

PALMER'S ORGANIC POLLUTION INDICES

| | DATE | 04 18 74 | 07 03 74 | 09 23 74 |
|---------|------|----------|----------|----------|
| GENUS | | 14 | 06 | 02 |
| SPECIES | | 76 | 04 | 00 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

| | DATE | 04 18 74 | 07 03 74 | 09 23 74 |
|----------------------------------|------|----------|----------|----------|
| AVERAGE DIVERSITY | H | 2.94 | 2.78 | 2.22 |
| NUMBER OF TAXA | S | 33.00 | 39.00 | 32.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 3.00 | 3.00 | 3.00 |
| MAXIMUM DIVERSITY | MAXH | 5.04 | 5.29 | 5.32 |
| MINIMUM DIVERSITY | MINH | 0.03 | 0.04 | 0.03 |
| TOTAL DIVERSITY | O | 44667.42 | 39815.16 | 35453.40 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 15193.00 | 14322.00 | 15970.00 |
| EVENNESS COMPONENT | J | 0.58 | 0.53 | 0.44 |
| RELATIVE EVENNESS | RJ | 0.59 | 0.53 | 0.45 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 460.39 | 367.23 | 499.06 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 5340.00 | 5919.00 | 8132.00 |

| TAXA | FORM | 04 18 74 | | 07 03 74 | | 09 23 74 | | | | |
|----------------------------------|------|----------|------|--------------------|-------|----------|--------------------|--------|-------|--------------------|
| | | IS | SC | ALGAL UNITS PER ML | IS | SC | ALGAL UNITS PER ML | IS | SC | ALGAL UNITS PER ML |
| ACTINASTRUM | CEL | | | X | | | | | | |
| ANARAENA PLANTONICA | FIL | | | | 4 | 3.9 | 477 | | | |
| ANARAENA SPIROIDES | FIL | | | | | | | | | X |
| ANISTRODESMUS FALCATUS | CEL | 0.9 | 140 | | | | | | | |
| APHANOcapsa DELICATISSIMA | COL | | | | | | 1222.9 | 3653 | 705 | |
| APHANOcapsa ELACHISTA | COL | | | | | | 1.4 | | | |
| APHANOThCE ? | COL | | | X | | | | | | |
| ASTERIONELLA FORMOSA | CEL | 4 | 5.8 | 884 | | 0.8 | 119 | 0.4 | 59 | |
| CERATIUM HIRUNDINELLA | CEL | | | | | | X | | | |
| CHLAMYDOMONAS | CEL | | | | | | | | | X |
| CHROMOCOCCUS LIMNETICUS | COL | | | | 0.8 | 119 | | | | |
| CHROOMONAS NORSTENTII | CEL | | | | | | | | | |
| COELOSPHAERIUM | COL | | | | | 1311.2 | 1609 | | | |
| COELOSPHAERIUM NAEGELIANUM | COL | | | X | | | | | | X |
| COSMARIAUM | CEL | | | | | | X | | | |
| COSMARIAUM #2 | CEL | | | | | | | | | X |
| CRUCIGENIA QUADRATA | COL | 0.3 | 47 | | | | | 0.7 | 118 | |
| CRUCIGENIA REC TANGULARIS | COL | | | | | | | 0.7 | 118 | |
| CRYPTOMONAS | CEL | 3.4 | 511 | | | | | 151 | 236 | |
| CYANOPHYTA FILAMENT | FIL | | | | | | | 1.5 | 236 | |
| CYMBELLA #1 | CEL | | | | | | | | | |
| CYMBELLA #2 | CEL | | | | | | | | | |
| CYMBELLA spp. | CEL | | | | | | | | | |
| DICTYOSPHAERIUM PULCHELLUM | COL | | | X | 1.2 | 179 | | | | |
| DINOBRYON CYLINDRICUM | CEL | | | | 0.4 | 60 | | | | |
| DINOBRYON DIVERGENS | CEL | 1135.1 | 5340 | | | | | | | X |
| DINOBRYON spp. | CEL | | | | | 1.2 | 179 | | | |
| ELAKATOOTHRIX | CFL | 2.8 | 419 | | | | | X | | |
| FLAGELLATE #1 | CEL | 51 | 5.2 | 793 | | 0.8 | 119 | 2.9 | 471 | |
| FRAGILARIA #1 | CEL | | | | | | X | | | |
| FRAGILARIA CONSTRIENS | CEL | | | | | | X | | | |
| FRAGILARIA CROTONENSIS | CEL | | | | | | X | | | |
| GLENDONIUM GYMNOCLINUM | CEL | 1313.8 | 2091 | | 3.7 | 536 | | | | X |
| V. BISCUTELLIFORME | CEL | | | | | | X | | | |
| GOMPHONEMA CONSTRICUTUM | CEL | | | | | | X | | | |
| GOMPHOSPHAERIA | COL | 0.3 | 47 | | | | | | | |
| GYNNOCLINUM | CEL | 0.6 | 93 | | | | | | | |
| KIRCHNERIELLA | CEL | 1.2 | 187 | | | | | | | |
| LYNGBYA | FIL | | | | 1.7 | 238 | | | | |
| LYNGBYA #1 | FIL | | | 2.4 | 370 | | | | | X |
| LYNGBYA LIMNETICA | FIL | | | 0.9 | 140 | | | | | |
| MELOSIRA | CEL | 222.6 | 3432 | | | | | | | |
| MELOSIRA GRANULATA | CEL | | | | | | X | | | |
| MELOSIRA ISLANDICA | CEL | | | | | | X | 4111.0 | 1886 | |
| MELOSIRA spp. | CEL | | | | | | | | | |
| MERISMOPEDIA MINIMA | COL | 0.3 | 47 | | 1.1 | 238 | | | | X |
| MICROCYSTIS ? STAGNALIS | COL | | | | 222.9 | 3277 | | | | |
| MICROCYSTIS #1 | COL | | | | | | | | | X |
| MICROCYSTIS AERUGINOSA | COL | | | | | | | 312.6 | 412 | |
| MICROCYSTIS INCERTA | COL | | | | | | | 0.7 | 118 | |
| MICROCYSTIS STAGNALIS | COL | | | | | | | 150.9 | 8132 | |
| OMICYSTIS | CEL | | | | 51 | 3.7 | 536 | 0.4 | 59 | |
| OSCILLATORIA | FIL | 0.9 | 140 | | | | | | | |
| PEDIASTRUM #1 | COL | 0.3 | 46 | | | 0.4 | 60 | | | |
| PEDIASTRUM BORYANUM | COL | | | | | | | | | |
| PEDIASTRUM DUPLEX | COL | | | | | | | | | |
| PEDIASTRUM DUPLEX | COL | | | | | | | | | |
| V. CLATHRATUM | COL | | | | | | | | | |
| PEDIASTRUM KAWPAISKYI | COL | | | X | | | X | 0.4 | 59 | |
| PERIDINUM CINCTUM | CEL | | | X | | | X | | | |
| RAPHIDIOPSIS CURVATA | FIL | | | | 1.2 | 179 | | | | |
| SCENEDESMUS #2 | COL | | | | 0.4 | 60 | | | | |
| SCENEDESMUS BIJUGA | COL | | | | | | | | | |
| V. FLEXUOSUS | COL | | | | | | X | | | |
| SCENEDESMUS DIMORPHUS | COL | | | X | | | | | | |
| SCENEDESMUS QUADRICAUDA | COL | | | | 0.4 | 60 | | | | |
| SCENEDESMUS QUADRICAUDA | COL | | | | | | | | | |
| V. RIORNATUS F. GIGANTICUS | COL | | | | | | | | | X |
| SCENEDESMUS spp. | COL | 0.9 | 140 | | | | | | | |
| SCHROEDERIA SETIGERA | CEL | 0.6 | 93 | | 1.2 | 179 | | 0.7 | 118 | |
| SPHAEROCYSTIS | COL | | | | | | | | | |
| STATOSPORE (DINOBRYON CIVERGENS) | CEL | | | X | | | X | | | |
| STAURASTRUM | CEL | | | X | | | | | | |
| STAURASTRUM #1 | CEL | | | X | | | | | | |
| STAURASTRUM #3 | CEL | | | X | | | | | | |
| STAURONETIS | CEL | | | X | | | | | | |
| STEPHANODISCS ASTRAEA | CEL | 0.6 | 93 | | 1.2 | 179 | | | | |
| SYNEDRA ULNA | CFL | 0.6 | 93 | | | | | | | |
| TETRAEDRON CAUDATUM | CFL | | | | | | | | | |
| TRACHELOMONAS VOLVOCINA | CEL | 1 | 0.31 | 47 | 1 | 1 | | | | |
| TOTAL | | | | | 15193 | | 14322 | | 15970 | |

LAKE NAME: DARLING
STORET NUMBER: 1905

NYGAARD TROPHIC STATE INDICES

DATE 04 18 74 09 24 74

| | | |
|---------------|--------|--------|
| MYXOPHYCEAN | 02/0 E | 1.00 E |
| CHLOROPHYCEAN | 02/0 E | 1.50 E |
| EUGLENOPHYTE | 0.75 E | 0.40 E |
| DIATOM | 0.50 E | 01/0 E |
| COMPOUND | 08/0 E | 4.00 E |

PALMER'S ORGANIC POLLUTION INDICES

DATE 04 18 74 09 24 74

| | | |
|---------|----|----|
| GENUS | 09 | 02 |
| SPECIES | 03 | 00 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE 04 18 74 09 24 74

| | | | |
|----------------------------------|------|---------|----------|
| AVERAGE DIVERSITY | H | 2.45 | 1.45 |
| NUMBER OF TAXA | S | 19.00 | 15.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 2.00 | 2.00 |
| MAXIMUM DIVERSITY | MAXH | 4.25 | 3.91 |
| MINIMUM DIVERSITY | MINH | 0.09 | 0.03 |
| TOTAL DIVERSITY | D | 6220.55 | 10682.15 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 2539.00 | 7367.00 |
| EVENNESS COMPONENT | J | 0.58 | 0.37 |
| RELATIVE EVENNESS | RJ | 0.57 | 0.37 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 133.63 | 491.13 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 1367.00 | 5665.00 |

04 18 74 09 24 74

| TAXA | FORM | ALGAL UNITS PER ML | | | ALGAL UNITS PER ML | | |
|---|------|--------------------|------|------|--------------------|------|-----|
| | | IS | ZC | | IS | ZC | |
| ANABAENA PLANKTONICA | FIL | 1 | 1 | 1 | | | X |
| ANASTRODESMUS FALCATUS | CEL | 121 | 7.71 | 195 | | | |
| APHAENIZUM NON FLOS-AQUAE | FIL | 1 | 1 | | 1176.91 | 5665 | |
| CLOSTERIUM | CEL | 1 | 1 | | 146.0.51 | 40 | |
| COCCOID CELL | CEL | 14 | 1.4 | 36 | | | |
| COELASTRUM CARBILUM | COL | 1 | 1 | | 0.81 | 59 | |
| COSMAGIUM | CEL | 1 | 1 | | 0.31 | 20 | |
| CRYPTOCRUMAS EROSA | CEL | | | X | 121.7.81 | 574 | |
| CRYPTOCRUMAS SPP. | CEL | 1153.81 | 1367 | | | | |
| CYANOPHYTAN FILAMENT | FIL | 1 | 0.71 | 18 | | | |
| CYCLOTELLA | CEL | 1 | 3.91 | 89 | | | |
| DINGBYTOM DIVERGENS | CEL | 1 | 1 | X | | | |
| EUGLENA #1 | CEL | 1 | 1 | X | | | |
| EUGLENA #2 | CEL | 1 | 1 | 2 | | | |
| EUGLENA SPP. | CEL | 138 | 2.11 | 53 | | | |
| FLAGELLATE #1 | CEL | 1 | 3.91 | 89 | | 4.31 | 317 |
| FLAGELLATE #2 | CEL | 111.21 | 284 | | | | |
| GLENODINIUM GYANOQUINUM V. BISCUTELLIFORME | CEL | 1 | 1 | | | | X |
| KIRCHHEIJELLA | CEL | 1 | 2.11 | 53 | | | |
| LEPOCINCLIS #1 | CEL | 1 | 2.11 | 53 | | 0.81 | 59 |
| LEPOCINCLIS #2 | CEL | 1 | 2.11 | 53 | | | |
| RELUSIBA GRANULATA | CEL | 1 | 1 | | 0.61 | 59 | |
| NAVICULA #2 | CEL | 1 | 1 | X | | | |
| NAVICULA SPP. | CEL | 1 | 1.41 | 36 | | | |
| NAVICULA ZANONI | CEL | 1 | 1 | X | | | |
| PALMELLOID CHLOROPHYTAN COLONY | COL | 1 | 1 | | 0.61 | 59 | |
| PERIDINIUM UMBONATUM | CEL | 151 | 2.11 | 53 | | | |
| QUADRIGULA | COL | 1 | 1 | | 0.31 | 20 | |
| RAPHIDIOPSIS CURVATA | FIL | 1 | 0.41 | 213 | | | |
| SCHROEDERIA SETIGERA | CEL | 1 | 1 | | 3.21 | 238 | |
| STEPHANUDISCUS | CEL | 1 | 1 | | 158.2.11 | 158 | |
| TRACHELUMUNAS | CEL | 1 | 1 | | 131.1.31 | 99 | |
| TOTAL | | | | 2939 | | 7367 | |

LAKE NAME: LOST ISLAND
STORET NUMBER: 1906

NYGAARD TROPHIC STATE INDICES

| | DATE | 09 23 74 | 04 24 76 |
|---------------|------|----------|----------|
| MYXOPHYCEAN | | 15/0 E | 2.67 E |
| CHLOROPHYCEAN | | 13/0 E | 3.00 E |
| EUGLENOPHYTE | | 0/28 ? | 0.29 E |
| DIATOM | | 2.00 E | 0.19 ? |
| COMPOUND | | 34/0 E | 8.67 E |

PALMER'S ORGANIC POLLUTION INDICES

| | DATE | 09 23 74 | 04 24 76 |
|---------|------|----------|----------|
| 'GENUS | | 12 | 26 |
| SPECIES | | 01 | 07 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

| | DATE | 09 23 74 | 04 24 76 |
|----------------------------------|------|-----------|----------|
| AVERAGE DIVERSITY | H | 3.50 | 4.47 |
| NUMBER OF TAXA | S | 41.00 | 61.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 2.00 | 1.00 |
| MAXIMUM DIVERSITY MAXH | | 5.36 | 5.93 |
| MINIMUM DIVERSITY MINH | | 0.02 | 0.10 |
| TOTAL DIVERSITY | D | 136552.50 | 39680.19 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 39015.00 | 8877.00 |
| EVENESS COMPONENT | J | 0.65 | 0.75 |
| RELATIVE EVENESS | RJ | 0.66 | 0.75 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 951.59 | 145.52 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 7769.00 | 1614.00 |

| TAXA | FORM | 09 23 74 | | | 09 24 76 | | |
|---------------------------|------|----------|-------|--------|----------|------|--------|
| | | IS | ZC | PER ML | IS | ZC | PER ML |
| ACTINIASTRUM | CEL | | | | | 1.31 | 115 |
| ANABAENA | FIL | | | | | X | |
| ANABAENA #1 | FIL | | 0.71 | 256 | | | |
| ANABAENA #2 | FIL | | 2.41 | 939 | | | |
| ANOMISTRUDES MUS FALCATUS | CEL | | | | 151 | 7.81 | 692 |
| APHANIZONE NON FLOS-AQUAE | FIL | 151 | 5.51 | 2134 | | | |
| APHANOCAPS A 7 | CUL | 141 | 19.91 | 7769 | | | |
| APHANOTHECE CLATHRATA | COL | | 2.01 | 768 | | | |
| ASTERIONELLA FORMOSA | CEL | | | | | 0.71 | 58 |
| CARTERIA | CEL | | | | | 1.31 | 115 |
| CENTRIC DIATOMS | CEL | | 1.31 | 912 | | | |
| CHRODODUCUS DISPERSUS | CGL | | | X | | | |
| COCCONECIS PLACENTULA | CFL | | | | | 3.61 | 317 |
| CORLASTRUM CAMBICUM | CGL | | | | | X | |
| COELOSPHAERIUM | COL | | 3.41 | 171 | | | |
| COSMARIA | CEL | | | | | 0.31 | 29 |
| CRYPTODIOMAS | CEL | | 0.41 | 171 | | | |
| CRYPTODIOMAS ERUSA | CEL | | | | | X | |
| CRYPTODIOMAS REFLEXA | CEL | | | | | X | |
| CRYPTODIOMAS SPP. | CEL | | | | 11 | 9.71 | 865 |
| CYANOPHYTA FILAMENT | FIL | | | | | 1.31 | 114 |
| CYCLUTELLA | CEL | | | | | 1.01 | 144 |
| CYCLUTELLA RENEGMHNJANA | CEL | | | X | | | |
| CYMBELLA AFFIMIS | CEL | | | | | X | |
| CYMBELLA SPP. | CEL | | | | | 3.21 | 288 |
| DINOBRYON DIVERGENS | CEL | | | | | 4.91 | 432 |
| ELAKATOTOM IX | CEL | | | X | | | |
| EPITHIMERIA | CEL | | | | | 1.31 | 86 |
| EUGLENA SPP. | CEL | | | | | 1.91 | 173 |
| EUNGRIA #1 | CEL | | | | | X | |
| EUNOTIA GRACILIS | CEL | | | | | X | |
| EUNOTIA SPP. | CEL | | | | | 0.31 | 29 |
| FLAGELLATE #1 | CEL | | 1.31 | 598 | 18.21 | 1014 | |
| FLAGELLATE #3 | CEL | | | | | 2.61 | 231 |
| FRAGILARIA #2 | CEL | | | | | 2.61 | 231 |
| FRAGILARIA CROTONENSIS | CEL | | | | 141 | 6.51 | 576 |
| FRAGILARIA INTERMEDIA | CEL | | | | | X | |
| GOMPHOMEREA ACURINATUM | CEL | | | | | | |
| V. CURUNATA | CEL | | | | | | |
| GOMPHOMEREA CLINSTRICTUM | CEL | | | | | | |
| GOMPHOMEREA PARVULUM | CEL | | | | | | |
| GOMPHOMEREA SPP. | CEL | | | | | | |
| GYRUSIGMA | CEL | | | | | 0.31 | 29 |
| KIRCHNERIELLA | CEL | | 0.41 | 341 | | 4.91 | 403 |
| KIRCHNERIELLA #1 | CEL | | | X | | | |
| KIRCHNERIELLA ELONGATA | CEL | | | X | | | |
| LAGERNA JINA CITRIFORMIS | CEL | | | X | | | |
| LIMNOVA CIRCURCRETA | FIL | | 3.91 | 1537 | | | |
| LIMNOVA LINNETICA | FIL | 111 | 15.81 | 6147 | | | |
| MALLOMMAS | CEL | | | X | | | |
| RASMOGLIA | CEL | | | | | 1.31 | 86 |
| RELOSIRA | CEL | | | X | | | |
| RELOSIRA GRANULATA | CEL | | | X | | | |
| RELOSIRA GRANULATA | CEL | | | X | | | |
| V. ANGUSTISSIMA | CEL | | | X | | | |
| RELOSIRA ISLANDICA | CEL | | | | | | |
| RELOSIRA SPP. | CEL | 131 | 14.91 | 9805 | | | |
| PERISOPEDIA | COL | | | | | | |
| PERISOPEDIA MINIMA | COL | | 10.11 | 3927 | | 1.01 | 86 |
| PERISOPEDIA TENUISSIMA | COL | | | X | | | |
| MICROCYSTIS | COL | | | | | | |
| MICROCYSTIS AERUGINOSA | COL | | 3.31 | 1281 | | 1.31 | 115 |
| MICROCYSTIS INCERTA | COL | | | X | | | |
| NAVICULA CUSPIDATA | COL | | | | | | |
| NAVICULA CUSPIDATA | CEL | | | | | | |
| V. ARDUGA | CEL | | | | | | |
| NAVICULA SPP. | CEL | | | | | | |
| MITZSCHEA ACICULARIS | CEL | | 2.01 | 768 | | | |
| MITZSCHEA MULSATICA | CEL | 121 | 21.21 | 4354 | | | |
| MITZSCHEA PALEA | CEL | | | | | | |
| OOCYSTIS | CEL | | 0.41 | 171 | | | |
| OSCILLATORIA | FIL | | 2.41 | 939 | | 2.31 | 202 |
| PEDIASTRUM BORYANUM | COL | | | X | | | |
| PEDIASTRUM DUPLEX | COL | | | X | | | |
| PEDIASTRUM KRAVATSKYI | COL | | | X | | | |
| PEDIASTRUM TETRAS | COL | | | X | | | |
| PHACUS ACURINATUS | CEL | | | | | | |
| PHACUS MEGALOPLYSIS | CEL | | | | | 0.31 | 29 |
| PHACUS SPP. | CEL | | | | | 0.31 | 29 |
| PHORMIDIUM #2 | FIL | | | | | 0.31 | 29 |
| PHORMIDIUM MUCICOLA | COL | | | X | | | |
| PINNULARIA | CEL | | | | | 0.71 | 58 |
| RAPIDIDIOPSIS | FIL | | | | | 4.91 | 432 |
| RHOPALOGIA GIBBA | CEL | | | | | X | |
| SCHEDESMUS ABUNDANS | COL | | | | | 0.71 | 56 |

LAKE NAME: LUST ISLAND
STORET NUMBER: 1906

CONTINUED

| TAXA | FORM | 09 23 74 | | | 04 24 76 | | |
|-------------------------|------|----------|------|--------------------|----------|------|--------------------|
| | | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML |
| SCENEDESMEUS ACUMINATUS | COL | | | X | | | |
| SCENEDESMEUS BIJUGA | COL | | | | | 2.61 | 231 |
| SCENEDESMEUS QUADRICUDA | COL | | | | | 1.34 | 115 |
| SPERMATOZOOPSIS | CEL | | | | | 0.71 | 58 |
| SPIRULIGRA | CEL | | | | | 131 | 29 |
| STAUROSTRUM | CEL | | | | | | X |
| STAUROSTRUM #2 | CEL | | | | | | X |
| STEPHANODISCUS | CEL | | | X | | | |
| STEPHANODISCUS ASTREA | CEL | | | | | 21 | 202 |
| SYNEURA | CEL | | | | | | |
| SYNEOPA ACUS | CFL | | 0.71 | 256 | | | 0.71 |
| TETRAEDRUM CAUDATUM | CEL | | | X | | | |
| TETRAEDRUM MASTATUM | CEL | | | X | | | |
| TETRASTRUM ELEGANS | COL | | 3.41 | 171 | | | |
| TOTAL | | | | | 39015 | | 8677 |

LAKE NAME: MCBRIDE
STUDY NUMBER: 1907

NYGAARD TROPHIC STATE INDICES

| DATE | 04 18 74 | 07 03 74 | 09 24 74 |
|---------------|----------|----------|----------|
| MYXOPHYCEAN | C1/0 E | 3.00 E | 4.00 E |
| CHLOROPHYCEAN | C2/0 E | 4.00 E | 13.0 E |
| EUGLENOPHYTE | 0.33 E | 0.43 E | 0.24 E |
| DIATOM | 3.00 E | 3.00 E | 0.87 E |
| COMPOUND | G7/0 E | 13.0 E | 26.0 E |

PALMER'S ORGANIC POLLUTION INDICES

| DATE | 04 18 74 | 07 03 74 | 09 24 74 |
|---------|----------|----------|----------|
| GENUS | 00 | 02 | 10 |
| SPECIES | 00 | 00 | 00 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

| DATE | 04 18 74 | 07 03 74 | 09 24 74 |
|----------------------------------|----------|----------|----------|
| AVERAGE DIVERSITY | H | 1.52 | 1.77 |
| NUMBER OF TAXA | S | 17.00 | 20.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 3.00 | 3.00 |
| MAXIMUM DIVERSITY | MAXH | 4.09 | 4.32 |
| MINIMUM DIVERSITY | MINH | 0.02 | 0.04 |
| TOTAL DIVERSITY | D | 20988.18 | 10651.86 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 13808.00 | 6018.00 |
| EVENNESS COMPONENT | J | 0.37 | 0.41 |
| RELATIVE EVENNESS | RJ | 0.37 | 0.41 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 812.24 | 300.90 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 8889.00 | 4077.00 |

| TAXA | FORM | 04 18 74 | | | 07 03 74 | | | 09 24 74 | | |
|-----------------------------|------|----------|------|--------------------|----------|------|--------------------|----------|------|--------------------|
| | | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML |
| ACHMANTHES | CEL | 1 | 1 | | | | | | | x |
| AMAGAEMA | FIL | 1 | 1 | | | | x | | | |
| ANKISTRODESMUS FALCATUS | CEL | 1 | 1 | x | | | | | | x |
| APHANIZOMON FLUSS-AQUAE | FIL | 1 | 1 | 0.91 | 119 | 1 | 0.71 | 43 | | |
| ASTERIONELLA FORMOSA | CEL | 191 | 4.31 | 593 | | | x | | | |
| CENTRIC DIATOM | CEL | 11164.41 | 8889 | | | | | | | |
| CHLOROCUCCALEAN CELL | CEL | 1 | 1 | | | | | 0.41 | 37 | |
| COCCONEIS | CEL | 1 | 1 | | | | | | | x |
| COELUSPHAEARIUM NAEGELIANUM | COL | 1 | 1 | | | | x | | | |
| CRUCIGENIA APICULATA | COL | 1 | 1 | | | | | | | x |
| CRUCIGENIA TETRAPEDIA | COL | 1 | 1 | | | | | | | x |
| CRYPTOMMAS | CEL | 1 | 1 | | | | | 131 | 6.51 | 635 |
| CRYPTOMMAS ERUSA | CEL | 1 | 1 | x | | | | | | |
| CRYPTOMMAS KARSSONII | CEL | 1 | 1 | x | | | x | | | |
| CRYPTOMMAS REFLEXA | CEL | 1 | 1 | x | 121 | 3.31 | 198 | 151 | 0.41 | 37 |
| CRYPTOMMAS SPP. | CEL | 141 | 4.71 | 652 | | | | | | |
| CYCLOTELLA MENEGHINIANA | CEL | 1 | 1 | | | | | | | x |
| DACTYLOCOPCOPSIS | CEL | 1 | 1 | | | | | 1.91 | 187 | |
| DICTYOSPMAERIUM | COL | 1 | 1 | | | | | | | x |
| DINOBRYON SOCIALE | CEL | 1 | 1 | x | | 1.31 | 79 | | | |
| EUASTRUM DENTICULATUM | CEL | 1 | 1 | | | | | | | x |
| EUDORINA ELEGANS | COL | 1 | 1 | | | 1.31 | 79 | | | |
| EUGLENA | CEL | 1 | 1 | | | | | | | x |
| EUGLENA SCIOTENSIS ? | CEL | 1 | 1 | | | | | | | x |
| FLAGELLATE #1 | CEL | 1 | 1 | x | | | x | | | |
| FLAGELLATES | CEL | 12122.31 | 3081 | 5.31 | 317 | | 112.61 | 1232 | | |
| FRANCEJA | CEL | 1 | 1 | | | | | 0.41 | 37 | |
| GLENODINIUM | CEL | 1 | 1 | | | | | 0.41 | 37 | |
| GYMNODINIUM ALBULUM | CEL | 1 | 1 | x | | | | | | |
| LEPUCIMCLIS | CEL | 1 | 1 | | 151 | 2.01 | 119 | 1 | | |
| LUMATE CELL | CEL | 1 | 1 | | | | | | | x |
| MELOSIRA #1 | CEL | 1 | 1 | | | | | 141 | 6.11 | 597 |
| MELOSIRA DISTANS | CEL | 1 | 1 | | | | | 11380.71 | 3770 | |
| MELOSIRA GRANULATA | CEL | 1 | 1 | x | 1167.71 | 4077 | 121 | 2.31 | 224 | |
| MELOSIPA GRANULATA | CEL | 1 | 1 | | | | | | | |
| V. ANGUSTISSIMA | CEL | 1 | 1 | | | | | | | |
| MELOSIRA GRANULATA | CEL | 1 | 1 | | | | | | | |
| V. ANGUSTISSIMA F. SPHALIS | CEL | 1 | 1 | | | | | 1.11 | 112 | |
| MERISIOPEDIA TENUISSIMA | COL | 1 | 1 | | | | | 0.41 | 37 | |
| MESOSTIGMA | CEL | 1 | 1 | | | | | | | x |
| NAVICULA | CEL | 1 | 1 | | | | | 0.41 | 37 | |
| MITZSCHIA | CEL | 1 | 1 | | | | | 0.81 | 75 | |
| OCYSTIS | CEL | 1 | 1 | | | | | | | |
| OSCILLATORIA ? | CEL | 1 | 1 | | 3113.21 | 292 | | | | x |
| PAMOURINA RUGUM | FIL | 1 | 1 | | | | | | | |
| PEDIASTRUM BICAUDATA | COL | 1 | 1 | x | | | | | | |
| V. LONGECORNUTUM | COL | 1 | 1 | | | | | | | x |
| PEDIASTRUM DUPLEX | COL | 1 | 1 | | | | | | | |
| V. CLATHRATUM | COL | 1 | 1 | | | | x | | | |
| PEDIASTRUM SIMPLEX | COL | 1 | 1 | | | | | 0.81 | 659 | |
| PENNATE DIATOMS | CEL | 1 | 1 | | | | | | | |
| PHACUS | CEL | 1 | 1 | | | | | | | |
| PHACUS LUNGICAUDA | CEL | 1 | 1 | x | | | x | | | |
| PHACUS PYRUM | CEL | 1 | 1 | | | | | | | x |
| PTEROPORAS ANGULOSA | CEL | 1 | 1 | | | | | 0.81 | 75 | |
| SCENEDESmus DENTICULATUS | COL | 1 | 1 | | | | | | | x |
| SCENEDESmus QUADRICAUDA | COL | 1 | 1 | x | | | | | | |
| V. ? | COL | 1 | 1 | | | | | | | x |
| SCENEDESmus SPP. | CUL | 1 | 1 | | | | | 1.91 | 187 | |
| SCHROEDERIA SETIGERA | CEL | 1 | 1 | | | | | | | |
| STAUROSTRUM | CEL | 1 | 1 | | | | | | | |
| STEPHANO DISCUS | CEL | 1 | 1 | | 0.71 | 43 | | | | |
| STEPHANO DISCUS NIAGARAE | CEL | 131 | 3.41 | 674 | | | | 3.01 | 373 | |
| SUBIRELLA | CEL | 1 | 1 | | | | | | | x |
| SYNODRA | CEL | 1 | 1 | | | | | 0.51 | 635 | |
| SYNODRA ACUS ? | CEL | 1 | 1 | | | | | | | x |
| TETRAECRUM MINIMUM | CEL | 1 | 1 | | | | | 0.81 | 75 | |
| V. SCROBICULATUM | CEL | 1 | 1 | | | | | | | x |
| TETRAEDRUM MITICUM | CEL | 1 | 1 | | | | | | | x |
| TETRASTRUM HETERACANTHUM | COL | 1 | 1 | | | | | | | |
| TETRASTRUM STURUGENIAEFORME | COL | 1 | 1 | | | | | 1.11 | 112 | |
| TRACHELOMMA | CEL | 1 | 1 | | | | | | | |
| TOTAL | | | | 13808 | | | 6018 | | 9743 | |

LAKE NAME: PRAIRIE ROSE
STORET NUMBER: 1908

NYGAARD TROPHIC STATE INDICES

DATE 04 17 74 07 02 74 09 25 74

| | | | |
|---------------|--------|--------|--------|
| MYXOPHYCEAN | 2.60 E | 1.67 E | 1.67 E |
| CHLOROPHYCEAN | 3.00 E | 2.00 E | 1.67 E |
| EUGLENOPHYTE | 0.40 E | 0.18 ? | 0.30 E |
| DIATOM | 0.71 E | 2.00 E | 2.00 E |
| COMPOUND | 4.50 E | 5.00 E | 5.00 E |

PALMER'S ORGANIC POLLUTION INDICES

DATE 04 17 74 07 02 74 09 25 74

| | | | |
|---------|----|----|----|
| GENUS | 06 | 11 | 05 |
| SPECIES | 00 | 00 | 00 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE 04 17 74 07 02 74 09 25 74

| | | | | |
|----------------------------------|------|----------|---------|---------|
| AVERAGE DIVERSITY | H | 2.33 | 3.10 | 2.41 |
| NUMBER OF TAXA | S | 36.00 | 22.00 | 22.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 2.00 | 2.00 | 2.00 |
| MAXIMUM DIVERSITY | MAXH | 5.17 | 4.46 | 4.46 |
| MINIMUM DIVERSITY | MINH | 0.04 | 0.11 | 0.12 |
| TOTAL DIVERSITY | D | 30956.38 | 7483.40 | 5333.33 |
| TOTAL NUMBER OF INDIVIDUALS/ML. | N | 13286.00 | 2414.00 | 2213.00 |
| EVENNESS COMPONENT | J | 0.45 | 0.70 | 0.54 |
| RELATIVE EVENNESS | RJ | 0.45 | 0.69 | 0.53 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 369.06 | 109.73 | 100.59 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 7006.00 | 522.00 | 681.00 |

LAKE NAME: PRAIRIE ROSE
STREETER NUMBER: 1908

CONTINUED

| TAXA | FORM | 04 17 74 | | | 07 02 74 | | | 09 25 74 | | |
|----------------------------|------|----------|------|--------------------|----------|------|--------------------|----------|-----|--------------------|
| | | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML |
| ANABAENA | FIL | | | x | | | | | | |
| ANABAENA SPIROIDES | FIL | | | | 1820.5 | | 494 | | | x |
| ANGISTRODESmus? | CEL | 3.3 | 435 | | 215.9 | | 385 | 1.4 | 43 | |
| APHANIZOMENON FLOS-AQUAE | FIL | | | | | | x | | | |
| APHANOCAPS4 | COL | | | | | | x | | | |
| ASTERIONELLA FORRosa | CEL | 13152.7 | 7006 | | | | | | | |
| BOTRYOCOCCUS BRAUNII | COL | | | | | | x | | | |
| CENTRIC DIATOM | CEL | 16.2 | 2198 | | | | x | | | |
| CERATIUM MIRUNDINELLA | CEL | | | | 1.1 | | 27 | | | |
| CHLANTDOMUNAS? | CEL | | | x | | | | | | |
| CLOSTERIOPSIS LONGISSIMA | CEL | | | | | | | 1130.8 | 681 | |
| CLOSTERIUM #1 | CEL | | | | 1.1 | | 27 | | | |
| CLOSTERIUM #2 | CEL | | | | | | x | | | |
| CLOSTERIUM #3 | CEL | | | x | | | | | | |
| COCCOID CHRYSOPHYTAN | CEL | | | x | | | | | | |
| COELASTRUM CAMBRICUM | COL | 0.2 | 23 | | | | x | | | x |
| COELOSPHAERIUM MAEGELIANUM | COL | | | x | 1.1 | | 27 | | | x |
| CGSMAPTIUM | CEL | | | | | | x | | | |
| CRYPTOMUNAS | CEL | | | | 151.9.1 | | 220 | | | |
| CRYPTOMUNAS EROSA | CEL | 21.2.5 | 337 | | | | | | | |
| CRYPTOMUNAS EROSA? | CEL | | | | | | | 13126.9 | 596 | |
| CRYPTOMUNAS HARSSUNII | CEL | 1.5 | 193 | | | | | | | |
| CRYPTOMUNAS HARSSUNII? | CEL | | | | | | | 151.7.7 | 170 | |
| CRYPTOMUNAS REFLEXA | CEL | 0.4 | 48 | | | | | | | |
| CYNATUPLEURA ELLIPTICA | CEL | | | x | | | | | | |
| DICTYOSPHAERIUM PULCHELLUM | COL | 0.2 | 23 | | | | x | | | |
| DINOBRYON DIVERGENS | CEL | 6.3 | 835 | | | | | | | x |
| EUGLENA | CEL | | | x | | | 3.4 | | | |
| EUGLENA FI | CEL | | | x | | | 82 | | | |
| EUGLENA UXTURIS | CEL | | | | | | | | | x |
| FLAGELLATE #1 | CEL | 19111.2 | 1487 | | 1.1 | | 27 | | | |
| FRAGILARIA CRUTONENSIS | CEL | 0.5 | 68 | | 111.4 | | 275 | 12117.3 | 383 | |
| GYROSIGMA? | CEL | | | x | | | | | | |
| HALLOPONAS | CEL | | | x | | | | | | |
| MELLOSIRA DISTANS | CEL | | | x | | | | | | |
| MELLOSIRA GRANULATA | CEL | 0.4 | 48 | | 13121.6 | | 922 | | | x |
| MICROCYSTIS VARIANS | CEL | 0.3 | 46 | | | | | | | x |
| MICROCYSTIS AERUGINOSA | COL | | | | | | | | | |
| MITZSCMIA | CEL | 0.2 | 23 | | | | | | | x |
| MITZSCMIA #1 | CEL | 0.5 | 71 | | | | | | | |
| OCUCYSTIS | CEL | 1.0 | 138 | | | | | | | x |
| OSCILLATORIA | FIL | | | x | | | | | | |
| OSCILLATORIA AGARDHII | FIL | | | x | | | | | | |
| OSCILLATORIA SUBREVIS | FIL | | | | 148.8.0 | | 192 | 14111.5 | 299 | |
| PEDIASTRUM DUPLEX | COL | | | | | | x | | | |
| PEDIASTRUM DUPLEX | | | | | | | | | | |
| V. CLATHRATUM | | | | | | | | | | |
| PEDIASTRUM DUPLEX | COL | | | x | | | | | | x |
| V. RETICULATUM | COL | | | | | | x | | | |
| PEDIASTRUM SIMPLEX | COL | | | | | | | | | x |
| PENNATE DIATOM | CEL | 0.2 | 23 | | | | | | | x |
| PHACUS TORTUS | CEL | | | x | | | | | | |
| PHACUS TORTUS? | CEL | | | x | | | | | | |
| SCENEDEDESmus QUADRICAUDA | COL | | | x | | | | | | x |
| SCHIZOCHELMATIS COMPACTA | COL | | | | | | | | | |
| STAURASTRUM | CEL | | | | 1.1 | | 27 | | | x |
| STAURASTRUM CHAETOCERUS | CEL | | | x | | | | | | |
| STEPHANODISCUS ASTRaea | CEL | 141.2.4 | 324 | | 3.4 | | 82 | 3.6 | 85 | |
| TETRAEDRON CONSTRICTUM | CEL | | | | | | x | | | |
| TRACHELOMUNAS | CEL | | | | 1.1 | | 27 | | | |
| TRACHELOMUNAS? | CEL | | | x | | | | | | |
| TRACHELOMUNAS VOLVOCINA | CEL | | | x | | | | | | x |
| TOTAL | | | | 13286 | | 2414 | | 2213 | | |

LAKE NAME: RATHBUN
STORET NUMBER: 1909

NYGAARD TROPHIC STATE INDICES

| DATE | 04 19 74 | 07 03 74 | 09 24 74 |
|---------------|----------|----------|----------|
| MYXOPHYCEAN | 1.00 E | 01/0 E | 6.00 E |
| CHLOROPHYCEAN | 6.00 E | 04/0 E | 9.00 E |
| EUGLENOPHYTE | 0.86 E | 0.60 E | 0.53 E |
| DIATOM | 0.60 E | 1.67 E | 1.00 E |
| COMPOUND | 16.0 E | 13/0 E | 28.0 E |

PALMER'S ORGANIC POLLUTION INDICES

| DATE | 04 19 74 | 07 03 74 | 09 24 74 |
|---------|----------|----------|----------|
| GENUS | 03 | 01 | 09 |
| SPECIES | 03 | 00 | 00 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

| DATE | 04 19 74 | 07 03 74 | 09 24 74 |
|----------------------------------|----------|----------|----------|
| AVERAGE DIVERSITY | H | 0.63 | 3.47 |
| NUMBER OF TAXA | S | 27.00 | 21.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 6.00 | 6.00 |
| MAXIMUM DIVERSITY | MAXH | 4.75 | 4.39 |
| MINIMUM DIVERSITY | MINH | 0.01 | 0.22 |
| TOTAL DIVERSITY | D | 21306.60 | 3612.27 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 33820.00 | 1041.00 |
| EVENNESS COMPONENT | J | 0.13 | 0.79 |
| RELATIVE EVENNESS | RJ | 0.14 | 0.78 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 1252.59 | 49.57 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 30898.00 | 190.00 |
| | | | 818.00 |

| TAXA | FORM | 04 19 74 | | | 07 03 74 | | | 09 24 74 | | |
|------------------------------|------|----------|-------|--------------------|----------|-----|--------------------|----------|------|--------------------|
| | | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML |
| ACHMANTHES ? | CEL | | | | | | X | | | |
| ACTINASTRUM | CEL | | | X | | | | | | |
| ANABAENA | FIL | | | | | | | | | |
| AMERISTRODESMUS FALCATUS | CEL | 14 | 2.31 | 766 | | | X | | | X |
| ASTERIOMELLA FORMOSA | CEL | | | X | | | | | | X |
| CARTERIA | CEL | | | | | | | | | X |
| CHRODORONAS ? NORDSTEDTII | CEL | | | | | | | | | X |
| CLUSTERIUM | CEL | | | | | | | | | X |
| COELASTRUM CAMBRICUM | CEL | | | X | | | | | | |
| V. INTERMEDIUM | COL | | | | | | | 1.91 | 50 | |
| CRYPTODRONAS | CEL | 0.61 | 192 | 12113.61 | 142 | | 121 | 9.31 | 248 | |
| CRYPTODRONAS SPP. | CEL | | | | | | | | | |
| CYANOPHYTA FILAMENT | FIL | | | | 4.51 | 47 | | 0.91 | 25 | |
| CYANOPHYTA FILAMENT #1 | FIL | | | | | | | 0.91 | 25 | |
| CYANOPHYTA FILAMENT #2 | FIL | | | | | | | | | |
| CYCLOTELLA MENEGHINIANA | CEL | | | | 4.51 | 47 | | | | |
| DICTYOSPHAERIUM PULCHELLUM | COL | | | X | | | | | | |
| DIMOFLAGELLATE | CEL | | | X | | | | | | |
| EUGLENA | CEL | | | X | | | | 2.81 | 74 | |
| EUGLENA ACUS | CEL | | | X | | | | | | X |
| EUGLENA DESES | CEL | | | | | | X | | | |
| EUGLENA SPP. | CEL | | | | | | | | | |
| EUGLENA TRIPTERIS | CEL | | | | 4.51 | 47 | | | | |
| FLAGELLATE #1 | CEL | 121 | 2.81 | 958 | 6.81 | 71 | | 5.61 | 149 | |
| FLAGELLATE #2 | CEL | | | | 12113.61 | 142 | | | | X |
| FLAGELLATE #3 | CEL | | | | | | | 12115.71 | 421 | |
| FRAGILARIA ? | CEL | | | | | | | | | X |
| GYMNOBINIUM ? ORIGINATUM | CEL | | | X | | | | | | |
| KIRCHNERIELLA | CEL | | | | 9.11 | 95 | | 1.91 | 50 | |
| LEPOCINCLIS | CEL | | | | | | | 0.91 | 25 | |
| MALLOMONAS | CEL | | | | | | | 0.91 | 25 | |
| MELOSIRA | CEL | | | | | | | | | |
| MELOSIRA DISTANS | CEL | 131 | 1.41 | 479 | 1516.81 | 71 | | | | X |
| MELOSIRA GRANULATA | CEL | | | | 15118.31 | 190 | | | | X |
| MELOSIRA ISLANDICA | CEL | | | | | | | | | X |
| MELOSIRA SPP. | CEL | | | | | | | 12130.51 | 818 | |
| MELOSIRA VARIANS | CEL | | | | 4.51 | 47 | | 1.91 | 50 | |
| MERISMOPEDIA TENUISSIMA | COL | | | | | | | 1.91 | 50 | |
| NAVICULA | CEL | | | | | | | 1.91 | 50 | |
| NAVICULA ZANONI | CEL | | | | | | X | | | |
| NITZSCHIA | CEL | | | | 4.51 | 47 | | | | X |
| OSCILLATORIA | FIL | | | X | | | | | | |
| PALMELLIOID COLONY | COL | | | | | | X | | | |
| PANDORINA MOGUM | COL | | | X | | | | | | |
| PEDIASTRUM TETRAS | COL | | | | | | | | | X |
| V. TETRAODON | COL | | | X | | | | | | |
| PENNATE DIATOM | CEL | | | | | | | | | |
| PENNATE DIATOM #1 | CEL | | | X | | | | | | |
| PENNATE DIATOM #2 | CEL | | | X | | | | | | |
| PENNATE DIATOM #3 | CEL | | | X | | | | | | |
| PERIDINIUM | CEL | | | | | | | 0.91 | 25 | |
| PERIDINIUM QUADRIDENTS | CEL | | | | | | | | | X |
| PHACUS | CEL | | | | | | | | | X |
| PHACUS CAUDATUS | CEL | | | | | | | | | X |
| PHACUS MEGALOPSIS | CEL | | | X | | | | | | |
| PHACUS NORDSTEDTII | CEL | | | X | | | | | | |
| RAPHIDIOPSIS | FIL | | | | | | | 4.61 | 124 | |
| RAPHIDIOPSIS CURVATA | FIL | | | | | | | | | |
| SCENEDESmus | COL | 0.11 | 48 | | | | | | | |
| SCENEDESmus BICAUDATUS | COL | | | | | | X | | | |
| SCENEDESmus BIJUGA | COL | | | | | | | | | |
| SCENEDESmus QUADRICAUDA | COL | | | | | | | | | X |
| SCHREUERIA | CEL | | | | | | | | | |
| SCHREUERIA SETIGERA | CEL | | | | | | X | | | |
| STEPHANODISCUS | CEL | 11191.41 | 30898 | 1316.81 | 71 | | 1617.61 | 198 | | |
| STEPHANODISCUS #2 | CEL | | | | | | | | | X |
| STEPHANODISCUS ASTREA | CEL | | | | | | | | | X |
| SYNEDRA | CEL | | | | | | | | | |
| SYNEDRA ULNA | CEL | | | X | | | | | | X |
| TETRAEDRON CONSTRUCTUM | CFL | | | | | | | | | |
| TETRAEDRON MUTICUM | CEL | | | | | | | | | X |
| TETRAEDRUM HETERACANTHUM | COL | 0.61 | 192 | | | | | 1.91 | 50 | |
| TETRASTRUM STAURUGENIAEFORME | COL | | | | | | | 1212.61 | 74 | |
| TRACHELOMONAS | CEL | 151 | 0.81 | 287 | | | | | | |
| TRACHELOMONAS ? | CEL | | | | 2.31 | 24 | | | | |
| TRACHELOMONAS PLANCTONICA | CEL | | | | | | | | | X |
| TRACHELOMONAS PULCELLA | CEL | | | | | | | | | X |
| TRACHELOMONAS SCABRA | CEL | | | | | | | | | X |
| TRACHELOMONAS SPP. | CEL | | | | | | | 1317.41 | 198 | |
| TOTAL | | | | 33820 | | | 1041 | | 2679 | |

LAKE NAME: RED RUCK
STORE NUMBER: 1910

NYGAARD TRUPHIC STATE INDICES

| DATE | 04 18 74 | 07 08 74 | 09 24 74 |
|---------------|----------|----------|----------|
| MYXOPHYCEAN | 2.00 E | 07/0 E | 3.00 E |
| CHLOROPHYCEAN | 5.00 E | 14/0 E | 7.00 E |
| EUGLENOPHYTE | 0.57 E | 0.24 E | 0.10 ? |
| DIATOM | 0.50 E | 0.83 E | 1.50 E |
| COMPOUND | 14.0 E | 31/0 E | 17.0 E |

PALMER'S ORGANIC POLLUTION INDICES

| DATE | 04 18 74 | 07 08 74 | 09 24 74 |
|---------|----------|----------|----------|
| GENUS | 19 | 19 | 21 |
| SPECIES | 00 | 07 | 04 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

| DATE | 04 18 74 | 07 08 74 | 09 24 74 |
|------------------------------------|----------|----------|----------|
| AVERAGE DIVERSITY H | 1.55 | 4.10 | 2.59 |
| NUMBER OF TAXA S | 25.00 | 42.00 | 25.00 |
| NUMBER OF SAMPLES COMPOSITED M | 2.00 | 5.00 | 3.00 |
| MAXIMUM DIVERSITY MAXH | 4.64 | 5.39 | 4.64 |
| MINIMUM DIVERSITY MINH | 0.03 | 0.14 | 0.05 |
| TOTAL DIVERSITY D | 20884.70 | 16014.60 | 16143.47 |
| TOTAL NUMBER OF INDIVIDUALS/ML N | 13474.00 | 3906.00 | 6233.00 |
| EVENNESS COMPONENT J | 0.33 | 0.76 | 0.56 |
| RELATIVE EVENNESS RJ | 0.33 | 0.76 | 0.56 |
| MEAN NUMBER OF INDIVIDUALS/TAXA L | 538.96 | 93.00 | 249.32 |
| NUMBER/ML OF MOST ABUNDANT TAXON K | 9795.00 | 454.00 | 3213.00 |

| TAXA | FORM | 64 18 74 | | | 67 06 74 | | | 09 24 74 | | |
|------------------------------------|------|----------|------|--------------------|----------|------|--------------------|----------|------|--------------------|
| | | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML |
| ACTINASTRUM | CEL | | | | | | | | | |
| ACTINASTRUM HANTZSCHII | CEL | | | X | | | | | | X |
| AMABAENA | FIL | | | | 134 | 4.71 | 182 | | | |
| ANISTRUDESMUS ? | CEL | 2.4 | | 325 | | | | | | |
| APHANIZUMENON FLOS-AQUAE | FIL | | | | 16 | 5.61 | 227 | | | |
| APHANGCAPSA | COL | | | | | | X | | | |
| ARTHRODESMSUS | CEL | | | X | | | | | | |
| ASTERIONELLA FORMOSA | CEL | | | X | | | X | | | |
| CALONEIS AMPHIBAENA | CEL | | | X | | | | | | |
| CENTRIC DIATOM | CEL | 2172.7 | | 9795 | | | | | | |
| CHLOROPHYTAN COCCOID CELLED COLONY | COL | 1.6 | | 216 | | | | | | |
| CLOSTERIUM | CEL | | | | | | | | | X |
| COELASTAUM CAMBRICUM | COL | | | | | | | | | |
| V. INTERMEDIUM | COL | | | | | | X | | | |
| COELUSPNAERIUM | COL | | | 151 | 7.01 | 273 | | | | |
| CRUCIGENIA TETRAPEDIA | COL | | | | 1.21 | | 45 | | | |
| CRYPTOMMAS | CEL | | | | 3.51 | | 130 | | | |
| CRYPTOMMAS EROSA | CEL | 1.6 | 3.61 | 487 | | | | 1.01 | | 64 |
| CRYPTOMMAS EROSA ? | CEL | | | X | | | X | | | |
| CRYPTOMMAS REFLEXA | CEL | | | | | | | | | |
| CYCLOTELLA | CEL | | | | | | | 1151.51 | | 3213 |
| CYCLOTELLA MEGHENINIANA | CEL | | | | 3.51 | | 136 | | | |
| DACTYLOCOCCOPSIS | CEL | | | | 4.71 | | 182 | | | |
| DACTYLOCOCCOPSIS IRREGULARIS | CEL | | | | | | | 3.21 | | 193 |
| DICTYOSPHAERIUM PULCHELLUM | COL | | | | | | | 1.01 | | 64 |
| DINGBYRON SOCIALE | CEL | 0.41 | | 54 | | | | | | |
| EUGLENA | CEL | | | | | | | | | |
| EUGLENA #1 | CEL | | | X | | | | | | |
| EUGLENA #2 | CEL | | 0.8 | 108 | | | | | | |
| FLAGELLATE #1 | CEL | | | | | | | 151 | 5.21 | 323 |
| FLAGELLATE #2 | CEL | | | | 5.81 | | 227 | | | |
| GLENOIDIUM OCULATUM | CEL | | | | | | X | | | |
| KIRCHMENIELLA | CEL | | | | 2.31 | | 91 | | | |
| LUNATE CELL | CEL | | | | | | | | | X |
| MELOSIRA | CEL | | | | | | | | | X |
| MELOSIRA #4 | CEL | | | | | | | 131 | 7.21 | 450 |
| MELOSIRA DISTANS | CEL | | | | 2.31 | | 91 | | | 257 |
| MELOSIRA GRANULATA | CEL | | | X | | | X | | | |
| PELUSIRA GRANULATA | CEL | | | | | | | | | |
| V. ANGUSTISSIMA | CFL | | | | | | X | | | X |
| HERISOPEDIA TENUISSIMA | COL | | | | 2.31 | | 91 | | | 257 |
| MAVICULA | CEL | 0.41 | | 54 | | | | | | |
| MAVICULA CRYPTOCEPHALA | CFL | | | | | | X | | | |
| MAVICULA EXIGUA | CEL | | | | | | X | | | |
| NETZSCHIA #1 | CEL | | | | | | | 3.11 | | 193 |
| NETZSCHIA #2 | CEL | | | | | | | | | X |
| NETZSCHIA ACICULARIS | CEL | | | | 11.61 | | 454 | | | X |
| NETZSCHIA FILIFORMIS | CEL | | | | | | X | | | |
| NETZSCHIA HOLSATICA | CEL | | | | | | X | | | |
| OSCILLATORIA | FIL | 151 | 1.2 | 162 | 1211.61 | | 454 | | | |
| OSCILLATORIA LIRNETICA | FIL | | | X | | | | 12112.41 | | 771 |
| PALMELLUID COLONY | COL | | | | 1.21 | | 45 | | | |
| PEDIASTRUM | COL | | | | | | X | | | |
| PEDIASTRUM DUPLEX | COL | | | | | | | | | X |
| PEDIASTRUM DUPLEX | COL | | | | | | | | | |
| V. CLATHRATUM | COL | | | | | | X | | | |
| PEDIASTRUM DUPLEX | COL | | | X | | | | | | |
| V. RETICULATUM | COL | | | | | | | | | |
| PEDIASTRUM TETRAS | COL | | | | | | | | | |
| V. TETRAUDON | COL | | | | | | | | | X |
| PHACUS CAUDATUS | CEL | | | X | | | | | | |
| PHACUS MEGALOPSIS | CEL | | | | | | X | | | |
| PHACUS PSEUDOMORDSTEDTII | CFL | 0.41 | | 54 | | | | | | |
| SCENEDESMSUS ABUNDANS | COL | | | | | | | | | X |
| SCENEDESMSUS ACURINATUS | COL | | | | 1.21 | | 45 | | | |
| SCENEDESMSUS BIJUGA | COL | | | | 2.31 | | 91 | | | |
| SCENEDESMSUS BIJUGA | COL | | | | | | | | | |
| V. FLEALVUS | COL | | | | | | X | | | |
| SCENEDESMSUS GIGRAPHUS | COL | | | X | | | | 1.01 | | 64 |
| SCENEDESMSUS GRANULATUS | COL | | | | | | X | | | |
| V. LUNGISPINA F. GRANULATA | COL | | | | | | X | | | |
| SCENEDESMSUS QUADRICAUDA | COL | | | | 2.31 | | 91 | | | 129 |
| SCENEDESMSUS REINIFORME | COL | | | | | | X | | | |
| SCHOGGERIA SETIGERA | CEL | | | | | | X | | | |
| STEPHANODISCUS ASTREA | CEL | 11 | 0.2 | 1245 | 41 | 6.11 | 318 | | | X |
| SURIRELLA ANGUSTATA | CEL | | | X | | | | | | |
| SYNEDRA | CEL | 131 | 7.21 | 974 | | | | 3.11 | | 193 |
| SYNEDRA ULNA | CEL | | | X | | | | | | |
| TETRASTRUM STAUROGENIAEFORME | COL | | | | 4.71 | | 182 | | | |
| TRACHELOMMAS ENSIFERA | CEL | | | | | | X | | | |
| V. JAVANICA | CEL | | | | | | X | | | |
| TRACHELOMMAS URCEOLATA | CEL | | | | | | X | | | |

TOTAL

13474

3906

6233

LAKE NAME: RUCK CREEK
STORET NUMBER: 1911

NYGAARD TRUPHIC STATE INDICES

| | DATE | 04 19 74 | 07 03 74 | 09 24 74 |
|---------------|------|----------|----------|----------|
| MYXOPHYCEAN | | 0.50 E | 0/03 U | 2.50 E |
| CHLOROPHYCEAN | | 1.00 E | 0/03 U | 2.50 E |
| EUGLENOPHYTE | | 0.67 E | 12/0 E | 0.30 E |
| DIATOM | | 0.43 E | 0.75 E | 0.67 E |
| COMPOUND | | 4.00 E | 5.00 E | 7.50 E |

PALMER'S ORGANIC POLLUTION INDICES

| | DATE | 04 19 74 | 07 03 74 | 09 24 74 |
|---------|------|----------|----------|----------|
| GENUS | | 07 | 10 | 36 |
| SPECIES | | 03 | 00 | 00 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

| | DATE | 04 19 74 | 07 03 74 | 09 24 74 |
|----------------------------------|------|----------|----------|----------|
| AVERAGE DIVERSITY | H | 0.69 | 2.31 | 1.7e |
| NUMBER OF TAXA | S | 24.00 | 30.00 | 26.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 2.00 | 2.00 | 2.00 |
| MAXIMUM DIVERSITY | MAXH | 4.58 | 4.91 | 4.70 |
| MINIMUM DIVERSITY | MINH | 0.02 | 0.23 | 0.08 |
| TOTAL DIVERSITY | D | 12482.10 | 3411.87 | 7835.52 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 18040.00 | 1477.00 | 4452.00 |
| EVENNESS COMPONENT | J | 0.15 | 0.47 | 0.37 |
| RELATIVE EVENNESS | RJ | 0.15 | 0.45 | 0.37 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 753.75 | 49.23 | 171.23 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 16387.00 | 502.00 | 2759.00 |

| TAXA | FORM | 04 19 74 | | | 07 03 74 | | | 09 24 74 | | |
|----------------------------|------|----------|-------|--------------------|----------|------|--------------------|----------|------|--------------------|
| | | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML |
| ANABAENA #1 | FIL | | | | | | | | | X |
| ANABAENA #2 | FIL | | | | | | | | | X |
| APHAENIZOMENON FLOS-AQUAE | FIL | 0.31 | 52 | | | | | 1162.00 | | 2759 |
| ASTERIUMELLA FURNOSA | CEL | 0.21 | 35 | | | | | | | |
| CENTRIC DIATOM | CEL | | | 13128.00 | 413 | | | | | |
| CHLAMYDOMONAS | CEL | 151 | 0.71 | 122 | | | | 147.00 | | 313 |
| CHLAMYDOMONAS PSEUDOPENTTI | CEL | | | | | | | | | |
| CLOSTERIUM #1 | CEL | | | | | | | | | X |
| CLOSTERIUM #2 | CEL | | | | | | | | | X |
| CLOSTERIUM #3 | CEL | | | | | | | | | |
| CLOSTERIUM #4 | CEL | | | | | | | | | |
| COELASTRUM CAMBRICUM | COL | | | | | | | | | X |
| COELOSPHEARIUM MAGELLANUM | COL | | | | | | | | | X |
| CRUCIGENIA QUADRATA | COL | | | | | | | | | |
| CRYPTOMONAS EROSA | CEL | 131 | 2.21 | 400 | 11134.00 | 502 | | 2.81 | | 125 |
| CRYPTOMONAS OVATA | CEL | | | | | | | | | X |
| CRYPTOMONAS REFLEXA | CEL | 121 | 1.01 | 174 | | | | | | X |
| CRYPTOMONAS SPP. | CEL | | | | | | | 12115.51 | | 690 |
| CYCLOTELLA | CEL | 11190.61 | 16387 | | | | | | | |
| CYMATOPLEURA SOLLA | CEL | | | X | | | | | | |
| CYMBELLA | CEL | | | | | | | | | X |
| DIMOBRYTON SOCIALE | CEL | | | | | | | | | |
| EUDORINA ELEGANS | CEL | | | | | | | | | |
| EUGLENA | CEL | | | | | | | | | |
| EUGLENA #1 | CEL | | | | | | | | | |
| EUGLENA ACUS | CEL | | | | | | | | | |
| EUGLENA GRACILIS | CEL | | | | | | | | | |
| EUGLENA OXYURIS | CEL | | | | | | | | | |
| EUGLENA SPIROIDES | CEL | | | | | | | | | |
| EUGLENA TRIPTERIS | CEL | | | | | | | | | |
| FLAGELLATE #1 | CEL | 3.91 | 712 | | 4.01 | 59 | 15 | 9.91 | | 439 |
| GLENODINIUM GYMNOGINIUM | | | | | | | | | | |
| V. BISCUTELLIFORME | | | | | | | | | | |
| GLOLOCYSTIS ? | CEL | | | | | | | | | X |
| GYMNOGINIUM | CEL | 0.31 | 52 | | | | | | | |
| GYROSIGMA | CEL | | | | | | | | | |
| GYROSIGRA ? | CEL | | | | | | | | | |
| LEPOCINCLIS #1 | CEL | | | X | | | | | | |
| LEPOCINCLIS ACUTA ? | CEL | | | | | | | | | X |
| LEPOCINCLIS BUTSCHLII | CEL | | | | | | | | | |
| LEPOCINCLIS SPP. | CEL | | | | | | | | | |
| MALLOMONAS ACAROIDES | CEL | | | | | | | | | |
| NELOSIRA GRANULATA | CEL | | | X | 41 | 4.01 | 59 | | | |
| NELOSIRA GRANULATA | | | | | | | | | | |
| V. ANGUSTISSIMA | CEL | | | | | | | | | |
| NICHOCYSTIS AEGUGINOSA | CEL | | | | | | | | | |
| NAVICULA ZAMONI ? | CEL | | | X | | | | | | |
| NITZSCHIA #1 | CEL | | | | | | | | | |
| NITZSCHIA #2 | CEL | | | | | | | | | |
| NITZSCHIA SPP. | CEL | | | | | | | | | |
| OCCYSTIS | CEL | | | | | | | | | |
| PANDINIA MURUM | COL | | | | | | | | | |
| PEDIASTRUM DUPLEX | | | | | | | | | | |
| V. RETICULATUM | COL | | | | | | | | | |
| PHACUS ? | COL | | | | | | | | | |
| PHACUS LONGICAUDA ? | CEL | | | X | | | | | | |
| PHACUS NEGLAPSIS | CEL | | | | | | | | | |
| PHACUS PLEURUMECTES | CEL | | | | | | | | | |
| SCENEDESMIUS INTERMEDIUS | | | | | | | | | | |
| V. BICAUDATUS | COL | 0.11 | 17 | | 1 | | | | | |
| SCHROEDERIA SLITIGERA | CEL | | | X | | | | | | |
| STAURASTRUM CHAETOCERUS | CFL | | | X | | | | | | |
| STEPHANODISCUS ASTREA | CFL | 0.21 | 35 | | | | | | | |
| SUPIRELLA | CEL | 0.11 | 17 | | | | | | | |
| SYNEURA | CEL | | | | | | | | | |
| SYNEURA ACUS | CEL | | | | | | | | | |
| SYNEURA ULNA | CEL | 148 | 0.41 | 70 | | | | 131 | 1.41 | 63 |
| TETRAEDROM CUNSTRUCTUM | CEL | | | | | | | | | |
| TRACHELUMONAS GIBBEROSA | CEL | | | | | | | | | |
| TRACHELUMONAS VOLVUCINA | CEL | | | | | | | | | |
| TOTAL | | | | 18690 | | 1477 | | 4452 | | |

LAKE NAME: SILVER
STORE NUMBER: 1912

NYGAARD TROPHIC STATE INDICES

| | DATE | 07 03 74 | 09 23 74 |
|---------------|------|----------|----------|
| MYXOPHYCEAN | | 4.00 E | 3.00 E |
| CHLOROPHYCEAN | | 7.67 E | 3.50 E |
| EUCLENOPHYTE | | 0.06 ? | 0/26 ? |
| DIATOM | | 0.40 E | 0.75 E |
| COMPOUND | | 13.0 E | 7.25 E |

PALMER'S ORGANIC POLLUTION INDICES

| | DATE | 07 03 74 | 09 23 74 |
|---------|------|----------|----------|
| GENUS | | 13 | 08 |
| SPECIES | | 07 | 04 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

| | DATE | 07 03 74 | 09 23 74 |
|----------------------------------|------|-----------|-----------|
| AVERAGE DIVERSITY | H | 2.61 | 1.08 |
| NUMBER OF TAXA | S | 52.00 | 40.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 2.00 | 1.00 |
| MAXIMUM DIVERSITY | MAXH | 5.70 | 5.32 |
| MINIMUM DIVERSITY | MINH | 0.01 | 0.00 |
| TOTAL DIVERSITY | D | 215792.19 | 229099.32 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 82679.00 | 212129.00 |
| EVENNESS COMPONENT | J | 0.46 | 0.20 |
| RELATIVE EVENNESS | RJ | 0.46 | 0.21 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 1589.98 | 5303.23 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 45876.00 | 180490.00 |

LAKE NAME: SILVER
STORE NUMBER: 1912

CONTINUED

| TAXA | FORM | 67 J 3 74 | | | 59 23 74 | | |
|------------------------------------|------|-----------|-------|--------------------|----------|--------|--------------------|
| | | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML |
| ACTINASTRUM | COL | 1 | 0.51 | 386 | 1 | 0.51 | 1062 |
| ANABALNOPSIS CIRCULARIS | FIL | 1 | 1 | | 1 | 0.51 | 1062 |
| ANKISTRODESMUS FALCATUS | CEL | 1 | 0.11 | 97 | 1 | 1 | X |
| APHANIZUMENCH FLOS-AQUAE | FIL | 1113.21 | 10914 | 121 4.81 | 13192 | | |
| APHANOCAPSA DELICATISSIMA | COL | 1 | 1 | | 1 | 0.61 | 1274 |
| APHANOCAPS A ELACHISTA | COL | 1 | 0.11 | 97 | 1 | 0.31 | 637 |
| APHANTHECE CLATHRATA | COL | 1 | 1 | | 1 | 0.11 | 212 |
| CENTRIC ULATOR | CEL | 1 | 0.11 | 97 | 1 | 1 | |
| CERATIUM HIRUNDINELLA | CEL | 1 | 0.11 | 97 | 1 | 1 | |
| CHLOROGUNIUM | CEL | 1 | 0.11 | 97 | 1 | 1 | X |
| CHROOCUCUS | COL | 1 | 0.11 | 97 | 1 | 1 | |
| CHROOCUCUS #1 | COL | 1 | 1 | X | 1 | 1 | |
| CHROOCOCCUS #2 | COL | 1 | 1 | | 1 | 1 | X |
| COELASTRUM SPHAERICUM | CUL | 1 | 1 | 1 | 1 | 1 | |
| COSMARIA #1 | CEL | 1 | 0.11 | 97 | 1 | 1 | X |
| CGSMARIA CLEPSYSGRA | CEL | 1 | 0.11 | 97 | 1 | 1 | |
| V. NANUM | CEL | 1 | 0.11 | 97 | 1 | 1 | X |
| CRYPTOMUNAS | CEL | 1 | 0.91 | 773 | 1 | 0.21 | 425 |
| CYMATOPLEURA SOLEA | CEL | 1 | 1 | X | 1 | 1 | X |
| DICTYOSPHERIUM PULCELLUM | COL | 1 | 1 | X | 1 | 1 | |
| EPITHEMIA | CEL | 1 | 1 | X | 1 | 1 | |
| EUGLENA | CEL | 1 | 1 | X | 1 | 1 | |
| FLAGELLATES | CEL | 1 | 1.61 | 1352 | 1 | 0.11 | 212 |
| FRAGILARIA | CEL | 1 | 1.41 | 1159 | 1 | 1 | X |
| GLENDODINIUM GYMNODINIUM | CEL | 1 | 1 | X | 1 | 1 | |
| GOLENKINIA | CEL | 1 | 1.81 | 1449 | 1 | 1 | X |
| KIRCHNERIELLA LUNARIS | CEL | 1 | 0.51 | 386 | 1 | 1 | |
| V. IRREGULARIS | CEL | 1 | 0.11 | 97 | 1 | 1 | |
| LAGERHEIMIA | CEL | 1 | 1 | X | 1 | 1 | |
| LYNGBYA | FIL | 1 | 1 | | 1 | 1 | |
| LYNGBYA & OSCILLATORIA | FIL | 12155.51 | 45876 | 1 | 1 | 1 | |
| LYNGBYA CIRCUNCRETA | FIL | 1 | 2.01 | 1642 | 1 | 2.11 | 4459 |
| LYNGBYA LINNETICA | FIL | 1 | 1 | | 1 | 1 | X |
| LYNGBYA LINNETICA & OSCILLATUNA LI | FIL | 1 | 1 | | 1185.11 | 183490 | |
| MALLDONAS ACARDIUS | CEL | 1 | 1 | X | 1 | 1 | |
| MELOSIRA | CEL | 140 | 6.11 | 5022 | 151 | 0.31 | 637 |
| MERISMOPEDIA PUNCTATA | COL | 1 | 1 | X | 1 | 1 | |
| MERISMOPEDIA TENUISSIMA | COL | 1 | 1 | X | 1 | 0.11 | 212 |
| MICROCYSTIS AERUGINOSA | COL | 151 | 2.11 | 1738 | 131 | 2.01 | 5946 |
| MICRUCYSTIS INCERTA | COL | 1 | 0.91 | 773 | 1 | 1.11 | 2335 |
| MITZSCHIA HULSATICA | CEL | 1 | 0.91 | 773 | 1 | 1 | X |
| PEDIASTRUM BORTANUM | COL | 1 | 0.21 | 193 | 1 | 1 | X |
| PEDIASTRUM DUPLEX | COL | 1 | 1 | X | 1 | 1 | |
| V. CLATHRATUM | COL | 1 | 1 | X | 1 | 1 | |
| PEDIASTRUM DUPLEX | COL | 1 | 1 | X | 1 | 1 | X |
| V. RETICULATUM | COL | 1 | 1 | X | 1 | 1 | |
| PHACUS MEGALOPSIS | CEL | 1 | 1 | X | 1 | 1 | |
| PHORMIDIUM NUCICOLA | COL | 1 | 1 | X | 1 | 1 | X |
| POLYEDRIOPSIS ? | CEL | 1 | 0.11 | 97 | 1 | 1 | |
| SCENEDESmus ABUNDANS | COL | 2.61 | 2125 | 1 | 0.61 | 1274 | |
| SCENEDESmus ACUMINATUS | COL | 1 | 1 | X | 1 | 1 | X |
| SCENEDESmus BIJUGA | COL | 0.81 | 676 | 1 | 1 | | |
| SCENEDESmus BIJUGA V. FLEXUOSUS | COL | 1 | 1 | | 0.21 | 425 | |
| SCENEDESmus DIMORPHUS | COL | 1 | 0.11 | 97 | 1 | 0.21 | 425 |
| SCENEDESmus QUADRICAUDA | COL | 0.71 | 579 | 1 | 0.21 | 425 | |
| SCHROEDERIA SETIGERA | CEL | 1 | 1 | X | 1 | 1 | |
| SPHAEROCYSTIS SCHNEITERI | COL | 1 | 1 | X | 1 | 1 | |
| STAURASTRUM #1 | CEL | 0.11 | 97 | 1 | 1 | 1 | X |
| STAURASTRUM TETRACERUM | CEL | 1 | 1 | | 1 | 1 | |
| STEPHANO-DISCUS ASTREA | CEL | 1 | 1.21 | 906 | 1 | 1 | X |
| SYNEURA | CEL | 131 | 5.41 | 4443 | 1 | 0.51 | 1062 |
| TETRAEDRON CAUDATUM | 1 | 1 | 1 | 1 | 1 | 1 | |
| V. LONGECORNUTUM | CEL | 1 | 0.41 | 290 | 1 | 1 | |
| TETRAEDRON MASTATUM | CEL | 1 | 0.11 | 97 | 1 | 1 | X |
| TETRAEDRON MINIMUM | 1 | 1 | 1 | 1 | 1 | 1 | |
| V. SCROBICULATUM | CEL | 1 | 0.11 | 97 | 1 | 1 | X |
| TETRAEDRON REGULARE | CEL | 1 | 1 | | 1 | 1 | |
| TETRAEDRON REGULARE V. INCUS | CEL | 1 | 1 | | 0.21 | 425 | |
| TREUBARIA | CEL | 1 | 1 | X | 1 | 1 | X |
| WESTELLA BOTYOIDES ? | CEL | 1 | 1 | X | 1 | 1 | |
| TOTAL | | | | 82679 | | 212129 | |

LAKE NAME: SPIRIT
STORET NUMBER: 1913

NYGAARD TROPHIC STATE INDICES

| | DATE | 04 23 74 | 07 09 74 | 09 23 74 |
|---------------|------|----------|----------|----------|
| MYXOPHYCEAN | | 01/0 E | 06/0 E | 4.00 E |
| CHLOROPHYCEAN | | 01/0 E | 04/0 E | 3.00 E |
| EUGLENOPHYTE | | 0/02 ? | 0/10 ? | 0.08 ? |
| DIATOM | | 0.12 ? | 0.33 E | 0.50 E |
| COMPOUND | | 03/0 E | 12/0 E | 14.0 E |

PALMER'S ORGANIC POLLUTION INDICES

| | DATE | 04 23 74 | 07 09 74 | 09 23 74 |
|---------|------|----------|----------|----------|
| GENUS | | 01 | 01 | 07 |
| SPECIES | | 00 | 00 | 00 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

| | DATE | 04 23 74 | 07 09 74 | 09 23 74 |
|----------------------------------|------|----------|----------|----------|
| AVERAGE DIVERSITY | H | 2.40 | 2.36 | 2.98 |
| NUMBER OF TAXA | S | 17.00 | 20.00 | 19.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 3.00 | 3.00 | 3.00 |
| MAXIMUM DIVERSITY | MAXH | 4.09 | 4.32 | 4.25 |
| MINIMUM DIVERSITY | MINH | 0.13 | 0.08 | 0.36 |
| TOTAL DIVERSITY | D | 3686.40 | 6855.80 | 11055.80 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 1536.00 | 2905.00 | 3710.00 |
| EVENNESS COMPONENT | J | 0.59 | 0.55 | 0.70 |
| RELATIVE EVENNESS | RJ | 0.58 | 0.54 | 0.70 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 90.35 | 145.25 | 195.26 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 672.00 | 1569.00 | 1182.00 |

| TAXA | FORM | 04 23 74 | | | 07 09 74 | | | 09 23 74 | | |
|----------------------------|------|----------|----|--------------------|----------|----|--------------------|----------|-----|--------------------|
| | | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML |
| ANABAENA PLACONICA | FIL | 1 | 1 | 111 8.01 | 232 | 1 | 111 19.81 | 734 | 1 | |
| APHANIZONEUM ? FLOS-AQUAE | FIL | 1 | 1 | 121 9.01 | 262 | 1 | 121 8.81 | 326 | 1 | |
| APHANIZONEUM FLOS-AQUAE | FIL | 1 | 1 | 1 | 1 | 1 | 121 8.81 | 326 | 1 | |
| APHANOCAPSA ? | COL | 1 | 1 | 131 54.01 | 1569 | 1 | 1 | 1 | 1 | |
| APHANOTHECE ? | COL | 1 | 1 | 6.21 | 96 | 1 | 1 | 1 | 1 | |
| APHANOTHECE ? CLATHRATA | COL | 1 | 1 | 1 | 1 | 1 | 131 31.91 | 1182 | 1 | |
| ASTEKIGMELLA FORMUSA | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | X | 1 |
| CLUSTERIA | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | X | 1 |
| COCCONEIS PLACENTULA | CEL | 1 | 1 | 1 | 2.01 | 58 | 1 | 1 | X | 1 |
| COELOSPHAERIUM NAEGELIANUM | COL | 1 | 1 | 1 | 1 | 1 | 161 7.71 | 285 | 1 | |
| CRYPTOMNAS EROSA | CEL | 1 | 1 | 9.41 | 144 | 1 | 145 5.01 | 144 | 1 | |
| CYANOPHYTA FILAMENT | FIL | 1 | 1 | 1 | 2.01 | 58 | 1 | 1 | 1 | |
| CYLINDRUSPEPNM STAGNALE ? | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | X | 1 |
| CYBELLA | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| DINOBRYUM SOCIALE | CEL | 1 | 1 | 151 19.61 | 240 | 1 | 1 | 1 | 1 | |
| DINOFLAGELLATE CYST | CEL | 1 | 1 | 141 3.21 | 48 | 1 | 1 | 1 | 1 | |
| EPITHERIA | CEL | 1 | 1 | 1 | X | 1 | 1 | 1 | 1 | |
| EUGLENA ? | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | X | 1 |
| EUNOTIA ? | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| FLAGELLATE #1 | CEL | 1 | 1 | 121 43.71 | 672 | 1 | 151 4.01 | 116 | 1 | 5.51 |
| FLAGELLATE #2 | CEL | 1 | 1 | 131 15.61 | 240 | 1 | 1 | 1 | 1 | 204 |
| FRAGILARIA #1 | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| FRAGILARIA #2 | CEL | 1 | 1 | 1 | X | 1 | 1 | 1 | 1 | |
| FRAGILARIA CROTONENSIS | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| FRAGILARIA INTERMEDIA | CEL | 1 | 1 | 1 | X | 1 | 1 | 1 | 1 | |
| MALLUMUNAS | CEL | 1 | 1 | 1 | X | 1 | 1 | 1 | 1 | |
| MELOSIRA GRANULATA | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| MELOSIRA ISLANUICA | CEL | 1 | 1 | 1 | X | 1 | 1 | 1 | 1 | |
| MERISMOPEDIA MINIMA | COL | 1 | 1 | 1 | 1 | 1 | 4.01 | 116 | 1 | 4.41 |
| MICROCYSTIS AERUGINOSA | COL | 1 | 1 | 1 | 1 | 1 | 1 | 151 5.51 | 163 | 204 |
| NAVICULA | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| NITZSCHIA FILIFORMIS | CEL | 1 | 1 | 1 | X | 1 | 1 | 1 | 1 | |
| OOCYSTIS | CEL | 1 | 1 | 1 | 1 | 1 | 3.01 | 87 | 1 | 1 |
| OSCILLATORIA | FIL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | X |
| PEDIASTRUM BORYANUM | COL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| PEDIASTRUM SIMPLEX | COL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| PHORMIDIUM MUCICOLA | COL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| SCHROEDERIA SETIGERA | CEL | 1 | 1 | 3.11 | 46 | 1 | 9.01 | 262 | 1 | 3.31 |
| STEPHANODISCUS ASTRAEA | CEL | 1 | 1 | 1 | 1 | 1 | 1 | X | 1 | 122 |
| SYNEDRA ULMA | CEL | 1 | 1 | 1 | X | 1 | 1 | 1 | 1 | 82 |
| TOTAL | | | | 1536 | | | 2905 | | | 3710 |

LAKE NAME: VIKING
STORET NUMBER: 1914

NYGAARD TROPHIC STATE INDICES

DATE 04 17 74 07 02 74 09 25 74

| | | | |
|---------------|--------|--------|--------|
| MYXOPHYCEAN | 01/0 E | 5.00 E | 1.25 E |
| CHLOROPHYCEAN | 01/0 E | 3.00 E | 0.50 ? |
| EUGLENOPHYTE | 0/02 ? | 0.12 ? | 0.14 ? |
| DIATOM | 0/01 ? | 1.00 E | 1.00 E |
| COMPOUND | 02/0 E | 10.0 E | 2.25 E |

PALMER'S ORGANIC POLLUTION INDICES

DATE 04 17 74 07 02 74 09 25 74

| | | | |
|---------|----|----|----|
| GENUS | 00 | 05 | 02 |
| SPECIES | 00 | 00 | 00 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE 04 17 74 07 02 74 09 25 74

| | | | | |
|----------------------------------|----|----------|---------|---------|
| AVERAGE DIVERSITY | H | 0.56 | 2.47 | 2.22 |
| NUMBER OF TAXA | S | 7.00 | 16.00 | 17.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 1.00 | 1.00 | 1.00 |
| MAXIMUM DIVERSITY MAXH | | 2.81 | 4.00 | 4.39 |
| MINIMUM DIVERSITY MINH | | 0.00 | 0.05 | 0.05 |
| TOTAL DIVERSITY | D | 30812.68 | 9862.71 | 9446.10 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 55023.00 | 3993.00 | 4255.00 |
| EVENNESS COMPONENT | J | 0.20 | 0.62 | 0.54 |
| RELATIVE EVENNESS | RJ | 0.20 | 0.62 | 0.54 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 7860.43 | 249.56 | 250.29 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 49981.00 | 1719.00 | 1684.00 |

| TAXA | FORM | 4 17 74 | | | 5 02 74 | | | 09 25 74 | | | |
|----------------------------|------|----------|-------|--------------------|---------|------|--------------------|----------|------|--------------------|----|
| | | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML | |
| ANABACINA PLANTONICA | FIL | 1 | 1 | 1 | 1 | 1 | 11117.51 | 730 | 1 | 1.41 | 61 |
| ANABACINA SPICIDES | FIL | 1 | 1 | 1 | 1 | 1 | 1 | 151 | 2.91 | 122 | |
| APHAENIZUMENON FLUSS-AQUAE | FIL | 121 | 6.61 | 3617 | 121 | 6.1 | 641 | 121 | 9.41 | 398 | |
| ASTERIUMELLA FUKUOSA | CEL | 11190.81 | 49981 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| CERATIUM HYPUNDINELLA | CEL | 1 | 1 | 1 | 1 | 0.71 | 29 | 1 | 1 | x | |
| CLOSTELIUM GRACILE # | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| V. ELONGATUM | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 31 | |
| COELOSPHEARIUM NAEGETIANUM | COL | 1 | 1 | 1 | 148 | 3.71 | 148 | 11139.61 | 1685 | | |
| COSMIDIUM | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| CRYPTOCRUMAS | CEL | 141 | 3.41 | 219 | 1 | 1 | 1 | 1 | 1 | 2.21 | |
| CRYPTOCRUMAS EROSA | CEL | 151 | 6.61 | 151 | 6.61 | 262 | 1 | 1 | 1 | 92 | |
| DICRYOSPHAERIUM PULCHELLUM | COL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| DINOBATON CYLINDRICUM | CEL | 151 | 6.61 | 329 | 1 | 1 | 1 | 1 | 1 | 1 | |
| FLAGELLATE #1 | CEL | 131 | 1.41 | 767 | 1 | 1 | 2.21 | 87 | 1 | 1.41 | |
| FRAGILARIA CROTONENSIS | CEL | 1 | 1 | 13143.11 | 1719 | 1 | 1 | 1 | 1 | 61 | |
| GYNGGUINUM | CEL | 1 | 1 | 0.21 | 110 | 1 | 1 | 1 | 1 | 1 | |
| LEPOCINCLIS | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| MELOSIRA | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| MELUSIKA GRANULATA | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 2.91 | |
| MICROCYSTIS AERUGINOSA | COL | 1 | 1 | 1 | 1 | 1 | 1 | 131 | 4.31 | 184 | |
| OCYCTYSIS | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| OSCILLATORIA | FIL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| PEDIASTRUM DUPLEX | COL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| V. CLATHRATUM | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| PERIODINUM INCONSPICUUM | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| SCHRUDEKIA SETIGERA | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| SPHAEROCYSTIS SCHRETERI | COL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| STAURASTRUM #1 | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| STAURASTRUM #2 | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| TRACHELOMUNAS VULVOCINA | CEL | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | x | |
| TOTAL | | | | 59023 | | | 3993 | | 4255 | | |

LAKE NAME: WEST LAKE OKOBJEE
STORET NUMBER: 1915

NYGAARD TROPHIC STATE INDICES

| | DATE | 04 22 74 | 07 09 74 | 09 23 74 |
|---------------|------|----------|----------|----------|
| MYXOPHYCEAN | | 01/0 E | 06/0 E | 10.0 E |
| CHLOROPHYCEAN | | 0/0 D | 06/0 E | 7.00 E |
| EUGLENOPHYTE | | 1.00 E | 0/12 ? | 0/17 ? |
| DIATOM | | 0.37 E | 1.00 E | 0.57 E |
| COMPOUND | | 05/0 E | 15/0 E | 21.0 E |

PALMER'S ORGANIC POLLUTION INDICES

| | DATE | 04 22 74 | 07 09 74 | 09 23 74 |
|---------|------|----------|----------|----------|
| GENUS | | 02 | 01 | 02 |
| SPECIES | | 02 | 00 | 00 |

SPECIES DIVERSITY AND ABUNDANCE INDICES

| | DATE | 04 22 74 | 07 09 74 | 09 23 74 |
|----------------------------------|------|----------|----------|----------|
| AVERAGE DIVERSITY | H | 1.95 | 2.02 | 3.34 |
| NUMBER OF TAXA | S | 21.00 | 20.00 | 34.00 |
| NUMBER OF SAMPLES COMPOSITED | M | 3.00 | 3.00 | 3.00 |
| MAXIMUM DIVERSITY | MAXH | 4.39 | 4.32 | 5.09 |
| MINIMUM DIVERSITY | MINH | 0.09 | 0.12 | 0.43 |
| TOTAL DIVERSITY | D | 5924.10 | 4970.14 | 2889.10 |
| TOTAL NUMBER OF INDIVIDUALS/ML | N | 3038.00 | 1897.00 | 865.00 |
| EVENNESS, COMPONENT | J | 0.44 | 0.61 | 0.66 |
| RELATIVE EVENNESS | RJ | 0.44 | 0.60 | 0.63 |
| MEAN NUMBER OF INDIVIDUALS/TAXA | L | 144.67 | 94.85 | 25.44 |
| NUMBER/ML OF MOST ABUNDANT TAXON | K | 1395.00 | 642.00 | 164.00 |

| TAXA | FORM | 04 22 74 | | | 07 09 74 | | | 09 23 74 | | |
|-----------------------------|------|----------|------|--------------------|----------|----|--------------------|----------|----|--------------------|
| | | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML | IS | ZC | ALGAL UNITS PER ML |
| AMPHORA | CEL | | | | | | | | | X |
| ANABAENA | FIL | | | | | | | | | X |
| APHAENIZUM NON FLGS-AQUAE | FIL | | | 2815.31 | | | 57 | | | |
| APHAENICAPS A DELICATISSIMA | FIL | | | | | | 285 | | | |
| APHAENICAPS ELACHISTA | COL | | | | | | 151 | 7.21 | | 62 |
| V. PLANCTONICA | COL | | | | | | 471 | | | |
| APHAENICAPS Z CLATHRATA | COL | | | | | | | | | |
| ASTERIONELLA FORMUSA | CEL | | 1.C | 31 | | | | | | |
| BOTRYOCUCUS BRAUNII | COL | | | | | | | | | |
| CLOSTERIUM | CEL | | | | | | | | | |
| COCCOID CHRYSOPHYTAN | CEL | | | | | | | | | |
| COCCINEIS | CEL | | | | | | | | | |
| COELASTRUM RETICULATUM | COL | | | | | | | | | |
| COELOSPHAERIUM MAEGELLANUM | COL | | | | | | | | | |
| CRYPTOMMAS | CEL | | | | | | | | | |
| CRYPTOMMAS EROSA | CEL | | 1.01 | 31 | | | | | | |
| CRYPTOMMAS REFLEXA | CEL | | | | | | | | | |
| CYCLUTELLA | CEL | | | | | | | | | |
| CYNAGLEPLURA SOLEA | CEL | | | | | | | | | |
| CYNBELLA | CEL | | | | | | | | | |
| DIATOMA VULGARE | CEL | | | | | | | | | |
| DINOBRUTUM CYLINDRICUM | CEL | | | | | | | | | |
| EUDORINA | CEL | | | | | | | | | |
| FLAGELLATE #1 | CEL | | | | | | | | | |
| FLAGELLATE #3 | CEL | | | | | | | | | |
| FRAGILARIA | CEL | | | | | | | | | |
| GYMNODINIUM ALBULUM | CEL | | 1.01 | 62 | | | | | | |
| GYROSTIGMA | CEL | | | | | | | | | |
| LUNATE CELL | CEL | | | | | | | | | |
| HELOSIRA GRANULATA | CEL | | | | | | | | | |
| HELOSIRA GRANULATA ? | CEL | | | | | | | | | |
| HELOSIRA GRANULATA | CEL | | | | | | | | | |
| V. ANGUSTISSIMA | CEL | | | | | | | | | |
| HELOSIRA VARIANS | CEL | | | | | | | | | |
| MICROCYSTIS AERUGINOSA | CEL | | | | | | | | | |
| MICROCYSTIS INCERTA | COL | | | | | | | | | |
| NAVICULA | CEL | | | | | | | | | |
| NITZSCHIA ? #3 | CEL | | | | | | | | | |
| NITZSCHIA #1 | CEL | | | | | | | | | |
| NITZSCHIA #2 | CEL | | | | | | | | | |
| NITZSCHIA #3 | CEL | | | | | | | | | |
| OOCYSTIS | CEL | | | | | | | | | |
| OOCYSTIS SUBRARINA | CEL | | | | | | | | | |
| OSCILLATORIA | FIL | | | | | | | | | |
| OSCILLATORIA AGARUMIL | FIL | | | | | | | | | |
| PEDIASIUM | COL | | | | | | | | | |
| PEDIASIUM BURKANUM | COL | | | | | | | | | |
| PENNATE DIATOM | CEL | | | | | | | | | |
| PERIDINUM | CEL | 1.01 | 31 | | | | | | | |
| PHORRIDIUM MUCICOLA | COL | | | | | | | | | |
| SCHROEDERIA SETIGERA | CEL | | | | | | | | | |
| SPHAEROCYSTIS SCHNEIDERI | CEL | | | | | | | | | |
| STEPHANODISCUS | CEL | | | | | | | | | |
| STEPHANODISCUS ASTREA | CEL | | | | | | | | | |
| SYNEDRA | CEL | | | | | | | | | |
| TETRAEDRUM REGULARE | CEL | | | | | | | | | |
| TRACHELONAS VOLVOCINA | CEL | | | X | | | | | | |
| TOTAL | | | | 3036 | | | 1897 | | | 869 |