


**U.S. ENVIRONMENTAL PROTECTION AGENCY
NATIONAL EUTROPHICATION SURVEY
WORKING PAPER SERIES**



A COMPENDIUM OF LAKE AND RESERVOIR
DATA COLLECTED BY THE NATIONAL
EUTROPHICATION SURVEY IN THE
WESTERN UNITED STATES
WORKING PAPER NO. 477

CORVALLIS ENVIRONMENTAL RESEARCH LABORATORY - CORVALLIS, OREGON
and
ENVIRONMENTAL MONITORING & SUPPORT LABORATORY - LAS VEGAS, NEVADA

A COMPENDIUM OF LAKE AND RESERVOIR
DATA COLLECTED BY THE NATIONAL
EUTROPHICATION SURVEY IN THE
WESTERN UNITED STATES
WORKING PAPER NO. 477

September, 1978

INTRODUCTION

The National Eutrophication Survey (NES) was initiated in 1972 by the U.S. Environmental Protection Agency (EPA) to investigate the nationwide threat of accelerated eutrophication to freshwater lakes and reservoirs. In conjunction with State environmental agencies, the Survey developed information on nutrient sources, inputs, and impacts on selected freshwater lakes and reservoirs throughout the contiguous United States. In total, over 800 lakes and reservoirs, 4,200 tributaries and lake outlets, and 1,000 sewage treatment plants were included in the sampling programs which involved a joint field effort by EPA personnel, the National Guard of each State, operators of municipal and industrial waste treatment plants, and personnel of the respective State agency responsible for water pollution control activities. For details of the procedures and methods used in the geographical area encompassed in this report, refer to NES Working Paper No. 175, "National Eutrophication Survey Methods, 1973-1976".

One of the primary outputs of the NES program is the individual lake or reservoir report in which are summarized the trophic condition; the nutrient sources, loads, and controllability; and the limiting nutrient. Each report also includes all of the NES data pertaining to the water body, the drainage area, and the nutrient point sources. To make the NES data accessible to many users, data in each lake report have been summarized and compiled in this report which includes information on the water bodies sampled during the fourth year of the Survey.

(1975). Geographically, this compendium includes data on lakes and reservoirs in Arizona, California, Colorado, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

Compendia of data on water bodies in other areas of the U.S. have been prepared. Working Paper No. 474 includes data on water bodies sampled in 1972 (northeast and part of north-central U.S.); Working Paper No. 475 provides data on water bodies sampled in 1973 (eastern, north-central, and southeastern States); and Working Paper No. 476 includes the data obtained in 1974 (central and plains States).

COMPENDIUM COMPONENTS

In the compendium heading for each water body, the identifiers are given (NAME, STATE, principal COUNTY, STORET NO., WORKING PAPER NO., and NTIS ACCESSION NO.). Following the NAME, the trophic condition of the water body is shown in parentheses.

The trophic condition is based on an assessment of the data collected during the sampling year supplemented by results of past studies, if any, and communications with State personnel. Each water body was categorized as "OLIGOTROPHIC" (low nutrient levels and productivity), "MESOTROPHIC" (moderate nutrient levels and productivity), "EUTROPHIC" (high nutrient levels and productivity), or "HYPEREUTROPHIC" (very high nutrient levels and productivity). For large water bodies, two or more trophic categories maybe indicated; e.g., the major tributary embayments (nearest the nutrient sources) may be eutrophic while the main portion of the water body is mesotrophic.

Following the heading, the data are arranged in five categories:

- I. MORPHOMETRY
- II. PHYSICAL AND CHEMICAL CHARACTERISTICS
- III. BIOLOGICAL CHARACTERISTICS
- IV. NUTRIENT LOADING CHARACTERISTICS
- V. NON-POINT SOURCE NUTRIENT EXPORT

Each of these categories contains related information as discussed below. If data were not obtained, a series of asterisks is shown.

I. MORPHOMETRY

The morphometric data were compiled from the literature and/or from information provided by State and Federal personnel.

LAKE TYPE - either of NATURAL origin or resulting from stream IMPOUNDMENT.

DRAINAGE AREA (SQ KM) - the total drainage area (measured to the outlet) in square kilometers.

SURFACE AREA (SQ KM) - the area of the water surface in square kilometers.

MEAN DEPTH (METERS) - the volume of the water body, in cubic meters, divided by the surface area in square meters.

TOTAL INFLOW (CMS) - the mean of the inflows of all tributaries and the immediate drainage in cubic meters per second.

RETENTION TIME (YEARS or DAYS) - a mean value determined by dividing the lake volume, in cubic meters, by the mean annual outflow in cubic meters per unit of time. Note that the outflow maybe less than the total inflow because of evaporation; withdrawals for irrigation, public water supply, or other uses are included in the outflow.

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

These data are based on the sampling of each water body in the spring, summer, and fall of 1975. Depending on size, from one to many

sites were sampled, and multiple depths usually were sampled at each site. For every parameter in this category, except Secchi disc depth, the median value is reported. The median represents the middle value of all sampling stations, times, and depths. The mean of the SECCHI DISC depths at all stations and all sampling times is given.

MEDIAN ALKALINITY (MG/L) - total alkalinity, as CaCO_3 , in milligrams per liter.

MEDIAN CONDUCTIVITY (UMHOS) - specific conductance at 25°C in micromhos per centimeter.

MEAN SECCHI DISC (METERS) - the mean limit of visibility of a standard Secchi disc in meters.

MEDIAN TOTAL PHOSPHORUS (MG/L) - as P in milligrams per liter.

MEDIAN DISSOLVED PHOSPHORUS (MG/L) - as P in milligrams per liter.

MEDIAN INORGANIC NITROGEN (MG/L) - nitrate + nitrite + ammonia, as N, in milligrams per liter.

MEDIAN TOTAL NITROGEN (MG/L) - Kjeldahl nitrogen + nitrate + nitrite, as N.

III. BIOLOGICAL CHARACTERISTICS

MEAN CHLOROPHYLL A (UG/L) - the mean concentration of all samples, in micrograms per liter.

ALGAL ASSAY CONTROL YIELDS (MG/L-DRY WT) - for most of the water bodies are based on two or more values, in milligrams per liter dry weight, obtained from samples collected during the first (spring) and last (fall) samplings. If more than one sample was taken, the values shown represent the range of yields, and the number following in parentheses indicates the number of samples assayed. The test organism was Selenastrum capricornutum Printz.

LIMITING NUTRIENT (no units) - may be determined by two procedures which are (1) the growth response of Selenastrum capricornutum to the addition of various amounts of phosphorus or nitrogen or (2) the ratio of inorganic nitrogen to dissolved phosphorus determined from the sampling data. When the inorganic nitrogen

to dissolved phosphorus ratio is 14/1 or greater, the water body is considered phosphorus limited, whereas ratios of less than 14/1 are considered indicative of nitrogen limitation.

The LIMITING NUTRIENT at each sampling time is given. Except for the first and last sampling dates, the limiting nutrient is based on N/P ratios. The limiting nutrient for the first and last sampling date generally is based on algal assay results. However, if no value is shown for the ALGAL ASSAY CONTROL YIELD, the limiting nutrient for those dates is determined by the N/P ratio. Where "ND" is shown, nutrient data were not obtained, and the limiting nutrient cannot be determined.

SUMMARY OF PHYTOPLANKTON DATA - the COUNT of individuals, filaments, or colonies per milliliter of sample for each of the five most numerous genera on the date shown. The sum of the units of other genera present in the sample, but not specified, is also included.

IV. NUTRIENT LOADING CHARACTERISTICS

Nutrient loads of significant tributaries and the water body outlet(s) were calculated using the results of analyses of from 12 to 14 samples collected from each stream by the State National Guard monthly for a one-year period and stream flow estimates as provided by the U.S. Geological Survey through an interagency agreement. The nutrient loads of the unsampled portion of the drainage areas were estimated from the measured nutrient loads in the sampled streams in the area. Nutrient loads of all streams and the unsampled drainage area were estimated on the basis of a year of average or "normal" stream flow to minimize the influence of extreme hydrological events that may have occurred during the sampling year.

Sewage treatment plant nutrient loads were determined from results of analyses of from 5 to 14 monthly effluent samples and corresponding flow data provided by plant operators or by State agency personnel. For sewage treatment plants which were not sampled and those from which fewer than five samples were received, nutrient discharges were estimated on the basis of the population served by the facility.

For details of sampling procedures and methods of calculation, refer to NES Working Paper No. 175.

A. INPUT - an estimate of all external inputs of nitrogen and phosphorus to the water body.

POINT SOURCE MUNICIPAL (KG/YR) - an estimate of annual nitrogen and phosphorus inputs from municipal sewage treatment plants in kilograms per year.

POINT SOURCE INDUSTRIAL (KG/YR) - an estimate of annual nitrogen and phosphorus inputs from industrial waste treatment plants in kilograms per year.

POINT SOURCE SEPTIC TANKS (KG/YR) - an estimate of annual nitrogen and phosphorus inputs from septic tanks within approximately 90 meters of the shoreline in kilograms per year. If a value is shown for nitrogen but not phosphorus, the estimated phosphorus input was less than 5 kg.

NON-POINT SOURCE (KG/YR) - an estimate of the annual nitrogen and phosphorus inputs from tributaries, immediate drainage, and precipitation in kilograms per year.

TOTAL LOADING (KG/YR) - the sum of all external nitrogen and phosphorus inputs.

LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) - the total loading for the sampling year divided by the lake surface area ($\frac{\text{kg/yr}}{\text{km}^2} \times 10^{-3}$) in grams per square meter of surface area per year.

B. OUTPUT - an estimate of the annual nitrogen and phosphorus discharged through the lake OUTLET(S) (KG/YR) in kilograms per year. Asterisks indicate little or no outlet flow (and output) during the sampling year.

PERCENT RETENTION - the percentage of incoming nitrogen or phosphorus retained in the lake annually:

$$\left(\frac{\text{Input load}-\text{output load}}{\text{input load}} \times 100\% \right).$$

V. NON-POINT SOURCE NUTRIENT EXPORT

STREAM NAME -

MEAN FLOW (CMS) - the mean stream flow in a year of average hydrology in cubic meters per second.

DRAINAGE AREA (SQ KM) - the drainage area of the stream in square kilometers.

MEAN TOTAL P (MG/L) - the mean concentration of total phosphorus in the stream at the sampling site during the year of sampling.

MEAN TOTAL N (MG/L) - the mean concentration of total nitrogen in the stream at the sampling site during the year of sampling.

TOTAL P EXPORT (KG/SQ KM/YR) - the total phosphorus load of the stream (after subtracting known point-source loads) divided by the drainage area, in kilograms per square kilometer per year. Asterisks indicate the phosphorus was less than 0.1 kg if a value is shown for nitrogen.

TOTAL N EXPORT (KG/SQ KM/YR) - the total nitrogen load of the stream (after subtracting known point-source loads) divided by the drainage area, in kilograms per square kilometer per year.

AVAILABILITY OF WORKING PAPERS

Compendium users desiring more detailed information may obtain a copy of the report on the water body of interest. Requests to the National Eutrophication Survey should include the NAME of the water body and the WORKING PAPER NO. as shown in the compendium heading.

Requests may be addressed to:

National Eutrophication Survey, EPA
Corvallis Environmental Research Laboratory
200 S.W. 35th Street
Corvallis, OR 97330

Only limited numbers of the Working Papers can be provided by NES.

When these are exhausted, Working Papers can be obtained from:

National Technical Information Service
Department of Commerce
Springfield, VA 22161

The NTIS accession number is shown in the compendium heading. A blank or a series of zeros indicates numbers that were not available at the time of this printing; these accession numbers can be obtained from NES at the address shown on page 7.

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN ARIZONA

NAME - BIG LAKE
 COUNTY - APACHE
 STORET NO. - 0401

(EUTROPHIC)
 WORKING PAPER NO. 726, NTIS ACCESSION NO. PB-279 258/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 1.94 | 4.4 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 77. | 101. | 2.9 | 0.032 | 0.007 | 0.090 | 0.820 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|-------|------------------------------------|
| 2.9 | 0.2 - 9.5 (2) | | (6/19/75) N (10/ 6/75) N |

SUMMARY OF PHYTOPLANKTON DATA
 6/19/75 10/ 6/75

| GENERA | COUNT | GENERA | COUNT |
|---------------|------------|----------------|-------------|
| CRYPTOMONAS | 92 | APHAENOTHECE | 480 |
| SCHROEDERIA | 23 | MELOSIRA | 137 |
| CYSTS | 23 | CHROOMONAS | 137 |
| TRACHELOMONAS | 23 | ASTERIONELLA | 69 |
| EUGLENA | 23 | COELOSPHAERIUM | 69 |
| OTHER | 0 | OTHER | 113 |
| TOTAL | 184 | TOTAL | 1005 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | **** | ***** |
| NITROGEN | ***** | **** | ***** |

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN ARIZONA

NAME - FOOLS HOLLOW LAKE (EUTROPHIC)

COUNTY - NAVAJO

STORET NO. - 0402

WORKING PAPER NO. 727, NTIS ACCESSION NO. PB-279 256/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 282.30 | 0.57 | 7.0 | 0.080 | 31.4 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 81. | 152. | 0.8 | 0.059 | 0.014 | 0.090 | 0.640 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (3 / 4 / 75) N | LIMITING NUTRIENT AT SAMPLING TIME (6 / 19 / 75) N | (10 / 6 / 75) N |
|------------------------------|---------------------------------------|---------------------------|---|-------------------|
| 10.7 | 0.3 - 13.0 (2) | | | |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/ 4 / 75 | 6 / 19 / 75 | | 10 / 6 / 75 | |
|--------------|-------------|--------------|-------------|----------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 2587 | MELOSIRA | 1681 | MELOSIRA | 7673 |
| CRYPTOMONAS | 1053 | COELASTRUM | 193 | APHANIZOMENON | 310 |
| CYCLOTELLA | 511 | OOCYSTIS | 193 | TRACHELOMONAS | 194 |
| MELOSIRA | 421 | CRYPTOMONAS | 154 | CRYPTOMONAS | 116 |
| FLAGELLATES | 391 | SCENEDESMUS | 135 | ANKISTRODESMUS | 116 |
| OTHER | 1142 | OTHER | 696 | OTHER | 466 |
| TOTAL | 6105 | TOTAL | 3052 | TOTAL | 8875 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 65. | ***** | 5. | 145. | 215. |
| NITROGEN | 140. | ***** | 140. | 3565. | 3845. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | 0.38 |
| NITROGEN | ***** | ***** | 6.7 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SHOW LOW CREEK (A2) | 0.060 | 239.6 | 0.052 | 1.417 | -0.4 | 9. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------------|------------------------|------------------------|
| SHOW LOW CREEK (A3) | 0.064 | 1.219 |
| UNNAMED CREEK (B1) | 0.230 | 1.265 |
| UNNAMED CREEK (C1) | 0.067 | 1.257 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN ARIZONA

NAME - LAKE HAVASU (MESOTROPHIC)
 COUNTY - MOHAVE, AZ; SAN BERNADINO, CA
 STORET NO. - 0403 WORKING PAPER NO. 728, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 463091.80 | 77.30 | 9.2 | 306.020 | 32.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 136. | 995. | 2.0 | 0.015 | 0.005 | 0.170 | 0.540 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 3.9 | 0.1 - 0.3 (4) | (2/27/75) P (6/13/75) P (11/19/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 2/27/75 | 6/13/75 | 11/19/75 |
|--|---------|---------|----------|
|--|---------|---------|----------|

| | | | | | |
|--------------|------------|---------------|-------------|--------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| OSCILLATORIA | 495 | FRAGILARIA | 514 | RAPHIDIOPSIS | 624 |
| CHROOMONAS | 180 | CYCLOTELLA | 206 | CRYPTOMONAS | 381 |
| CRYPTOMONAS | 90 | TRACHELOMONAS | 128 | CYCLOTELLA | 173 |
| SYNEDRA | 90 | CHROOMONAS | 103 | CHROOMONAS | 139 |
| CYCLOTELLA | 90 | OOCYSTIS | 77 | LYNGBYA | 139 |
| OTHER | 45 | OTHER | 128 | OTHER | 276 |
| TOTAL | 990 | TOTAL | 1156 | TOTAL | 1732 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 7490. | ***** | 25. | 274205. | 281720. |
| NITROGEN | 19060. | ***** | 1015. | 8054025. | 8074100. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 190275. | 32. | 3.64 |
| NITROGEN | 7826805. | 3. | 104.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| COLORADO RIVER | 305.290 | 446256.8 | 0.036 | 0.831 | 0.6 | 18. |
| GENE WASH | 0.010 | 17.6 | 0.021 | 0.603 | 0.3 | 12. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------------------|------------------------|------------------------|
| COLORADO RIVER AQUEDUCT | 0.036 | 0.862 |
| BILL WILLIAMS RIVER | 0.026 | 1.275 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN ARIZONA

NAME - LUNA LAKE
 COUNTY - APACHE
 STORET NO. - 0404

WORKING PAPER NO. 729, NTIS ACCESSION NO. PB-281 462/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 93.50 | 0.30 | 2.5 | 0.110 | 107.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 97. | 220. | 2.6 | 0.182 | 0.131 | 0.050 | 0.720 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (6.2 - 13.3 (2) | LIMITING NUTRIENT AT SAMPLING TIME (4/30/75) N | LIMITING NUTRIENT AT SAMPLING TIME (6/18/75) N | LIMITING NUTRIENT AT SAMPLING TIME (10/ 6/75) N |
|------------------------------|---------------------------------------|-----------------------------|---|---|--|
| 3.4 | | | | | |

SUMMARY OF PHYTOPLANKTON DATA
4/30/75 6/18/75 10/ 6/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------------|---------------|-------------|--------------|------------|
| CRYPTOMONAS | 2990 | OSCILLATORIA | 420 | CRYPTOMONAS | 34 |
| CHROOMONAS | 1627 | ANABAENA | 262 | OSCILLATORIA | 34 |
| CYCLOTELLA | 308 | FRAGILARIA | 210 | NITZSCHIA | 34 |
| SYNEDRA | 132 | APHANIZOMENON | 105 | | |
| ANKISTRODESmus | 88 | CRYPTOMONAS | 105 | | |
| OTHER | 44 | OTHER | 0 | OTHER | 0 |
| TOTAL | 5189 | TOTAL | 1102 | TOTAL | 102 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|--------------------------------------|---------------------------------------|---|-----------------------------|--------------------------|
| PHOSPHORUS | 105. | ***** | ***** | 465. | 570. |
| NITROGEN | 285. | ***** | ***** | 4240. | 4525. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 540. | 5. | 1.90 |
| NITROGEN | 2320. | 49. | 15.1 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SAN FRANCISCO RIVER | 0.100 | 75.4 | 0.165 | 0.992 | 5. | 42. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN ARIZONA

NAME - LYMAN LAKE
 COUNTY - APACHE
 STORET NO. - 0405
 WORKING PAPER NO. 730, NTIS ACCESSION NO. PB-281 570/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 2046.10 | 5.67 | 6.7 | 0.800 | 2.4 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 169. | 307. | 0.4 | 0.099 | 0.056 | 0.060 | 0.430 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (5.8 - 8.6 (2) | LIMITING NUTRIENT AT SAMPLING TIME (4/30/75) N | LIMITING NUTRIENT AT SAMPLING TIME (6/19/75) N | LIMITING NUTRIENT AT SAMPLING TIME (10/ 6/75) N |
|------------------------------|---------------------------------------|----------------------------|---|---|--|
| 2.6 | | | | | |

SUMMARY OF PHYTOPLANKTON DATA

| | 4/30/75 | 6/19/75 | | 10/ 6/75 | |
|-------------|---------|----------------|-------|----------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CRYPTOMONAS | 62 | CHROOMONAS | 234 | CHROOCOCCUS | 276 |
| COCCONEIS | 31 | STEPHANODISCUS | 39 | CHROOMONAS | 184 |
| NITZSCHIA | 31 | | | STEPHANODISCUS | 46 |
| PHACUS | 31 | | | CRYPTOMONAS | 46 |
| OTHER | 0 | OTHER | 0 | ANABAENA | 46 |
| TOTAL | 155 | TOTAL | 273 | OTHER | 45 |
| | | | | TOTAL | 643 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 1425. | ***** | ***** | 8300. | 9725. |
| NITROGEN | 3535. | ***** | 35. | 44355. | 47925. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 3445. | 65. | 1.72 |
| NITROGEN | 43090. | 10. | 8.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-----------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| LITTLE COLORADO RIVER | 0.730 | 1934.7 | 0.328 | 1.516 | 4. | 19. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN ARIZONA

NAME - LAKE MOHAVE (MESOTROPHIC)
 COUNTY - MOHAVE, AZ; CLARK, NV
 STORET NO. - 0406 WORKING PAPER NO. 731, NTIS ACCESSION NO. PB-282 827/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 438486.80 | 105.63 | 18.9 | 314.950 | 73.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 137. | 884. | 3.3 | 0.017 | 0.010 | 0.240 | 0.600 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (4.4) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|------------------|------------------------------------|
| | | 0.2 - 0.5 (5) | (2/26/75) P |
| | | | (6/12/75) P |
| | | | (12/ 2/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 2/26/75 | 6/12/75 | 12/ 3/75 |
|--|---------|---------|----------|
|--|---------|---------|----------|

| | | | | | |
|--------------|-------------|--------------|-------------|--------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CYCLOTELLA | 802 | CHROOMONAS | 401 | CHROOMONAS | 378 |
| CHROOMONAS | 698 | CYSTS | 218 | RAPHIDIOPSIS | 310 |
| OSCILLATORIA | 419 | OOCYSTIS | 182 | CYCLOTELLA | 206 |
| CRYPTOMONAS | 349 | FRAGILARIA | 146 | SCENEDESmus | 103 |
| SYNEDRA | 314 | CERATIUM | 36 | SYNEDRA | 69 |
| OTHER | 0 | OTHER | 37 | OTHER | 69 |
| TOTAL | 2582 | TOTAL | 1020 | TOTAL | 1135 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 5. | 252955. | 252960. |
| NITROGEN | ***** | ***** | 175. | 12216250. | 12216425. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 203395. | 20. | 2.39 |
| NITROGEN | 8797855. | 28. | 115.7 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| COLORADO RIVER | 314.740 | 434601.8 | 0.033 | 2.016 | 0.6 | 28. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN ARIZONA

NAME - LAKE PLEASANT (EUTROPHIC)
 COUNTY - MARICOPA, YAVAPAI
 STORET NO. - 0407 WORKING PAPER NO. 732, NTIS ACCESSION NO. PB-281 402/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | ***** | 14.16 | 13.7 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 173. | 492. | 1.3 | 0.027 | 0.004 | 0.040 | 0.520 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (3 / 5 / 75) P | LIMITING NUTRIENT AT SAMPLING TIME (6 / 17 / 75) N | (11 / 17 / 75) P |
|------------------------------|---------------------------------------|---------------------------|---|--------------------|
| 9.8 | 0.4 - 1.5 (2) | | | |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/ 5/75 | 6/17/75 | 11/15/75 |
|--|---------|---------|----------|
|--|---------|---------|----------|

| | | | | | |
|----------------|-------------|------------------|-------------|------------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| PERIDINIUM | 540 | CENTRIC DIATOM | 804 | DACTYLOCoccOPSIS | 1518 |
| ANKISTRODESMUS | 295 | TETRAEDRON | 739 | TETRASTRUM | 875 |
| CHROOMONAS | 147 | MELOSIRA | 565 | MELOSIRA | 875 |
| CRYPTOMONAS | 98 | DACTYLOCoccOPSIS | 544 | NITZSCHIA | 669 |
| MELOSIRA | 98 | CRUCIGENIA | 261 | CRUCIGENIA | 437 |
| OTHER | 198 | OTHER | 1239 | OTHER | 3188 |
| TOTAL | 1376 | TOTAL | 4152 | TOTAL | 7562 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

1
5

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN ARIZONA

NAME - LAKE POWELL (OLIGOTROPHIC--MESOTROPHIC)
COUNTY - COCONINO, AZ; GARFIELD, KANE, SAN JUAN, UT
STORED NO. - 0408 WORKING PAPER NO. 733, NTIS ACCESSION NO. PB-281 514/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 279459.00 | 653.16 | 51.0 | 393,800 | 2.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 130. | 580. | 4.5 | 0.010 | 0.009 | 0.410 | 0.600 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L-DRY WT) (4/15/75) P (8/14/75) P (12/2/75) P

SUMMARY OF PHYTOPLANKTON DATA

4/16/75 8/14/75 12/ 1/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|-------------|-------|----------------|-------|-------------|-------|
| CHROOMONAS | 513 | FRAGILARIA | 816 | CHROOMONAS | 486 |
| FRAGILARIA | 428 | CHROOMONAS | 371 | FRAGILARIA | 333 |
| CRYPTOMONAS | 257 | CENTRIC DIATOM | 185 | CRYPTOMONAS | 26 |
| GLENODINIUM | 29 | NAVICULA | 148 | TETRAEDRON | 26 |
| | | SKELETONEMA | 148 | SCENEDESMUS | 26 |
| OTHER | 0 | OTHER | 279 | OTHER | 0 |
| TOTAL | 1227 | TOTAL | 1947 | TOTAL | 897 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | ***** | **** | ***** |
| NITROGEN | ***** | **** | ***** |

***** LAKE SAMPLING ONLY *****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN ARIZONA

NAME - RAINBOW LAKE
 COUNTY - NAVAJO
 STORET NO. - 0409

WORKING PAPER NO. 734, NTIS ACCESSION NO. PB-281 518/A8

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 26.20 | 0.32 | 4.6 | 0.290 | 171.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 118. | 179. | 1.5 | 0.046 | 0.009 | 0.045 | 0.620 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 16.4 | 4.0 - 4.8 (2) | (3/ 4/75) N (6/19/75) N (10/ 1/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/ 4/75 | 6/19/75 | | 10/ 6/75 | |
|----------------|--------------|--------------|--------------|--------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CYCLOTELLA | 52327 | FRAGILARIA | 17976 | FRAGILARIA | 3821 |
| CHROOMONAS | 706 | EPITHEMIA | 618 | CHROOMONAS | 303 |
| CRYPTOMONAS | 680 | CRYPTOMONAS | 353 | CRYPTOMONAS | 243 |
| OOCYSTIS | 419 | CHROOMONAS | 265 | EPITHEMIA | 121 |
| ANKISTRODESMUS | 419 | COCCONEIS | 132 | COCCONEIS | 30 |
| OTHER | 1125 | OTHER | 398 | OTHER | 62 |
| TOTAL | 55676 | TOTAL | 19742 | TOTAL | 4580 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 10. | ***** | 10. | 675. | 695. |
| NITROGEN | 340. | ***** | 435. | 10595. | 11370. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 340. | 51. | 2.17 |
| NITROGEN | 6000. | 47. | 35.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| WALNUT CREEK | 0.250 | 21.8 | 0.074 | 1.012 | 26. | 396. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------|------------------------|------------------------|
| TROUT CREEK | 0.060 | 1.147 |
| GOOSEBERRY CREEK | 0.061 | 0.767 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN ARIZONA

NAME - THEODORE ROOSEVELT LAKE (MESOTROPHIC)
 COUNTY - GILA
 STORET NO. - 0410 WORKING PAPER NO. 735, NTIS ACCESSION NO. PB-281 516/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 15084.20 | 60.70 | 28.0 | 31.960 | 1.9 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 124. | 795. | 1.8 | 0.020 | 0.008 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--------------------------------------|
| 4.1 | 0.2 - 1.7 (4) | (3/ 4/75) N (6/17/75) N (11/18/75) N |

SUMMARY OF PHYTOPLANKTON DATA
3/ 5/75 6/17/75 11/18/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|------------------|-------------|-----------------|-------------|------------------|-------------|
| CHROOMONAS | 1343 | PENNATE DIATOMS | 488 | CHROOMONAS | 639 |
| DACTYLOCOCCOPSIS | 564 | TETRAEDRON | 234 | KIRCHNERIELLA | 486 |
| CRYPTOMONAS | 322 | ANKISTRODESMUS | 195 | DACTYLOCOCCOPSIS | 256 |
| GLENODINIUM | 161 | CHROOMONAS | 156 | SCENEDESMUS | 102 |
| ANKISTRODESMUS | 54 | ANABAENOPSIS | 117 | FRAGILARIA | 102 |
| OTHER | 189 | OTHER | 313 | OTHER | 563 |
| TOTAL | 2633 | TOTAL | 1503 | TOTAL | 2148 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 15050. | ***** | ***** | 111440. | 126490. |
| NITROGEN | 47245. | ***** | ***** | 630815. | 678060. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 22380. | 82. | 2.08 |
| NITROGEN | 401180. | 41. | 11.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SALT RIVER | 25.010 | 11152.5 | 0.142 | 0.574 | 9. | 42. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------|------------------------|------------------------|
| PINAL CREEK | 0.110 | 0.879 |
| PINTO CREEK | 0.828 | 1.518 |
| SALOME CREEK | 0.026 | 0.421 |
| TONTO CREEK | 0.061 | 0.529 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN ARIZONA

NAME - SAN CARLOS RESERVOIR (EUTROPHIC)
COUNTY - GILA, GRAHAM, PINAL
STORET NO. - 0411 WORKING PAPER NO. 736, NTIS ACCESSION NO. PB-281 517/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 33374.70 | 52.61 | 15.3 | 12-220 | 2.9 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 16. | 1258. | 0.6 | 0.056 | 0.009 | 0.060 | 0.520 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L-DRY WT) (3/4/75) N (6/18/75) N (11/18/75) N

SUMMARY OF PHYTOPLANKTON DATA

| | 3/ 4/75 | | 6/18/75 | | 11/18/75 |
|------------------|-------------|------------------|-------------|------------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| ANKISTRODESMUS | 2045 | LYNGBYA | 2463 | CHROOMONAS | 6312 |
| CRYPTOMONAS | 1402 | CARTERIA | 1351 | DACTYLOCOCCOPSIS | 207 |
| DACTYLOCOCCOPSIS | 584 | DACTYLOCOCCOPSIS | 530 | EUGLENA | 89 |
| KIRCHNERIELLA | 467 | CENTRIC DIATOM | 424 | MESOSTIGMA | 59 |
| CHROOMONAS | 380 | CERATIUM | 265 | CHLAMYDOMONAS | 59 |
| OTHER | 1256 | OTHER | 1191 | OTHER | 327 |
| TOTAL | 6134 | TOTAL | 6224 | TOTAL | 7053 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 1565. | ***** | ***** | 490415. | 491980. |
| NITROGEN | 3680. | ***** | ***** | 1073140. | 1076820. |

B. OUTPUT

| | OUTLET(S) | PERCENT | LAKE SURFACE AREA | LOADING RATE |
|------------|-----------|-----------|-------------------|--------------|
| | (KG/YR) | RETENTION | (G/SQ M/YR) | |
| PHOSPHORUS | 10165. | 98. | | 9.35 |
| NITROGEN | 239960. | 78. | | 20.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| GILA RIVER | 10.090 | 29707.3 | 0.222 | 2.040 | 15. | 31. |
| SAN CARLOS RIVER | 1.620 | 2659.3 | 0.283 | 1.132 | 8. | 22. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LAKE AMADOR (EUTROPHIC)
COUNTY - AMADOR
STORET NO. - 0601 WORKING PAPER NO. 739, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 151.30 | 1.56 | 17.4 | 1,236 | 261.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN (METERS) | SECCHI DISC | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------|-------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 73. | 175. | 2.3 | 0.040 | 0.020 | 0.390 | 0.680 | |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L-DRY WT) (3/14/75) P (6/26/75) P

SUMMARY OF PHYTOPLANKTON DATA

3/14/75 6/26/75 11/12/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------|-------------|-------|-------------|-------|
| CENTRIC DIATOM | 68733 | FRAGILARIA | 5304 | MELOSIRA | 1064 |
| CHROOMONAS | 561 | SCENEDESMUS | 3700 | CHROOMONAS | 172 |
| ASTERIONELLA | 421 | CHROOMONAS | 1375 | GLENODINUM | 144 |
| STEPHANODISCUS | 224 | CRYPTOMONAS | 360 | CRYPTOMONAS | 144 |
| MELOSIRA | 112 | PERIDINIUM | 262 | CERATIUM | 29 |
| OTHER | 112 | OTHER | 361 | OTHER | 28 |
| TOTAL | 70163 | TOTAL | 11362 | TOTAL | 1581 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 2610. | ***** | ***** | 1205. | 3815. |
| NITROGEN | 8065. | ***** | ***** | 71915. | 79980. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 1800. | 53. | 2.45 |
| NITROGEN | 68335. | 15. | 51.3 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| JACKSON CREEK | 0.899 | 109.8 | 0.301 | 1.904 | 8. | 469. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - BOCA RESERVOIR (OLIGOTROPHIC)
 COUNTY - NEVADA
 STORET NO. - 0602 WORKING PAPER NO. 740, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 445.50 | 3.97 | 12.8 | 6.024 | 112.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 37. | 50. | 3.2 | 0.012 | 0.003 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--------------------------------------|
| 1.7 | 0.2 - 0.8 (2) | (6/ 9/75) N (6/30/75) P (11/ 4/75) P |

SUMMARY OF PHYTOPLANKTON DATA
6/ 9/75 6/30/75 11/ 4/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|------------|----------------|------------|--------------|-------------|
| TABELLARIA | 120 | CHROOMONAS | 114 | TABELLARIA | 646 |
| OSCILLATORIA | 90 | STEPHANODISCUS | 57 | FRAGILARIA | 258 |
| CHROOMONAS | 90 | ASTERIONELLA | 57 | CHROOMONAS | 194 |
| CENTRIC DIATOM | 60 | TABELLARIA | 28 | CRYPTOMONAS | 161 |
| MELOSIRA | 60 | OTHER | 0 | ASTERIONELLA | 129 |
| OTHER | 62 | | | OTHER | 0 |
| TOTAL | 482 | TOTAL | 256 | TOTAL | 1388 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 3380. | 3380. |
| NITROGEN | ***** | ***** | ***** | 78165. | 78165. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 2675. | 21. | 0.85 |
| NITROGEN | 72570. | 7. | 19.7 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| LITTLE TRUCKEE RIVER | 5.010 | 378.1 | 0.017 | 0.353 | 6. | 133. |
| DRY CREEK | 0.367 | 16.1 | 0.019 | 0.439 | 11. | 318. |
| EAST BOCA CANYON CREEK | 0.139 | 6.0 | 0.061 | 0.887 | 78. | 1546. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LAKE BRITTON (EUTROPHIC)
COUNTY - SHASTA
STORET NO. - 0603 WORKING PAPER NO. 741, NTIS ACCESSION

1/AE

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 11929.50 | 5.12 | 9.8 | 97,520 | 6.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN (METERS) | SECCI DISC | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------|------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 74. | 135. | 1.3 | | 0.067 | 0.047 | 0.115 | 0.315 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME

SUMMARY OF PHYTOPLANKTON DATA
3/26/75

| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
|----------------|-------|------------|-------|-------------|-------|
| CENTRIC DIATOM | 189 | MELOSIRA | 1000 | FRAGILARIA | 138 |
| CHROOMONAS | 54 | ANABAENA | 437 | OOCYSTIS | 79 |
| | | SYNEDRA | 219 | CHAETOCEROS | 39 |
| | | FRAGILARIA | 125 | CHROOMONAS | 39 |
| | | CHROOMONAS | 125 | CRYPTOMONAS | 20 |
| OTHER | 0 | OTHER | 187 | OTHER | 0 |
| TOTAL | 243 | TOTAL | 2093 | TOTAL | 315 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | 2960. | 5. | 220350. | 22315. |
| NTIROGEN | ***** | 8510. | 175. | 2399770. | 2408455. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE |
|------------|----------------------|----------------------|----------------------------------|--------------|
| PHOSPHORUS | 226420. | LOSS | 43.62 | |
| NITROGEN | 2454865. | LOSS | 470.4 | |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| PIT RIVER | 66.540 | 10328.9 | 0.076 | 0.900 | 15. | 182. |
| CLARK CREEK | 1.080 | 50.2 | 0.033 | 0.340 | 22. | 231. |
| BURNNEY CREEK | 5.860 | 471.4 | 0.037 | 1.000 | 15. | 392. |
| HAT CREEK | 23.220 | 942.8 | 0.075 | 0.422 | 55. | 323. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------|------------------------|------------------------|
| CLAYTON CREEK | 0.022 | 1.150 |

* FISH HATCHERIES.

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LAKE CASITAS
 COUNTY - VENTURA
 STORET NO. - 0604
 WORKING PAPER NO. 742, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 89.40 | 10.97 | 28.6 | 0.522 | 19.1 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 143. | 520. | 2.5 | 0.029 | 0.014 | 0.050 | 0.280 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 3.2 | ***** | (3/ 7/75) N (6/24/75) N (11/13/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/ 7/75 | 6/24/75 | 11/13/75 | |
|----------------|---------|------------|----------|------|
| GENERA | COUNT | GENERA | COUNT | |
| CHROOMONAS | 268 | DINOBYRON | 2973 | |
| ANKISTRODESMUS | 239 | CHROOMONAS | 125 | |
| SCENEDESMUS | 89 | | | |
| OTHER | 0 | OTHER | 0 | |
| TOTAL | 596 | TOTAL | 3098 | |
| | | | TOTAL | 2677 |

20
80

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 25. | 395. | 420. |
| NITROGEN | ***** | ***** | 865. | 23435. | 24300. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 445. | LOSS | 0.04 |
| NITROGEN | 14865. | 39. | 2.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| COYOTE CREEK | 0.191 | 34.2 | 0.012 | 0.495 | 2. | 87. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|----------------------|------------------------|------------------------|
| SANTA ANA CREEK | 0.014 | 1.222 |
| ROBLES-CASITAS CANAL | 0.020 | 0.530 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LAKE CROWLEY
 COUNTY - MONO
 STORET NO. - 0605

WORKING PAPER NO. 743, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | ***** | 21.38 | 10.6 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 108. | 239. | 3.2 | 0.046 | 0.034 | 0.045 | 0.275 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 5.8 | 2.7 | (6/10/75) N (6/30/75) N (11/ 5/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 6/10/75 | 6/30/75 | 11/ 5/75 |
|--|---------|---------|----------|
|--|---------|---------|----------|

| | | | | | |
|--------------|-------------|--------------|-------------|----------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 1049 | CHROOMONAS | 892 | APHANIZOMENON | 421 |
| ASTERIONELLA | 504 | ANABAENA | 803 | FRAGILARIA | 225 |
| DINOBYRON | 378 | ASTERIONELLA | 758 | CHROOMONAS | 168 |
| FRAGILARIA | 252 | SCHROEDERIA | 580 | OSCILLATORIA | 140 |
| CRYPTOMONAS | 84 | CRYPTOMONAS | 312 | STEPHANODISCUS | 112 |
| OTHER | 0 | OTHER | 668 | OTHER | 29 |
| TOTAL | 2267 | TOTAL | 4013 | TOTAL | 1095 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - DON PEDRO RESERVOIR (MESOTROPHIC)
COUNTY - TUOLUMNE
STORED NO. - 0606 WORKING PAPER NO. 744. NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 3957.90 | 52.45 | 47.7 | 46,868 | 1.7 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|---------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 19. | 44. | 3.0 | 0.013 | 0.004 | 0.060 | 0.250 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

SUMMARY OF PHYTOPLANKTON DAT

| | 3/11/75 | COUNT | 6/25/75 | COUNT | 11/12/75 | COUNT |
|-------------|---------|-------|----------------|-------|-------------|-------|
| GENERAL | | | GENERAL | | GENERAL | |
| CHROOMONAS | 1543 | | MELOSIRA | 2260 | CHROOMONAS | 171 |
| MELOSIRA | 450 | | CHROOMONAS | 231 | MELOSIRA | 114 |
| CRYPTOMONAS | 107 | | CENTRIC DIATOM | 138 | CRYPTOMONAS | 38 |
| | | | CERATIUM | 92 | FRAGILARIA | 19 |
| | | | ASTERIONELLA | 92 | PERIDINIUM | 19 |
| OTHER | 0 | | OTHER | 231 | OTHER | 0 |
| TOTAL | 2100 | | TOTAL | 3044 | TOTAL | 361 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 10625. | ***** | 5. | 44305. | 54935. |
| NITROGEN | 23565. | ***** | 195. | 1312590. | 1336350. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 26600. | 52. | 1.05 |
| NITROGEN | 1328520. | 1. | 25.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT-SOURCE NUTRIENT STREAM NAME | EXPORT MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------------------------------|------------------------|-----------------------|---------------------|---------------------|------------------------------|------------------------------|
| TUOLUMNE RIVER | 27.750 | 2393.2 | 0.022 | 0.775 | 8. | 283. |
| ROGER CREEK | 0.028 | 10.7 | 0.056 | 1.822 | 5. | 150. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------------|------------------------|------------------------|
| N. FK., TUOLUMNE RIVER | 0.027 | 0.910 |
| HATCH CREEK | 0.026 | 1.601 |
| MOCCASIN CREEK | 0.040 | 0.957 |
| SULLIVAN CREEK | 0.061 | 1.189 |
| WOODS CREEK * | 0.704 | 2.797 |
| TURNBACK CREEK * | 0.441 | 1.798 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LAKE ELSINORE (HYPEREUTROPHIC)
 COUNTY - RIVERSIDE
 STORET NO. - 0607 WORKING PAPER NO. 745, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 10.52 | ***** | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 940. | 5545. | 0.3 | 0.469 | 0.092 | 0.120 | 3.035 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD | LIMITING NUTRIENT AT SAMPLING TIME | |
|------------------------------|---------------------------------------|--------------|------------------------------------|--------------|
| 70.6 | 6.3 - 16.5 (3) | (3/10/75) N | (6/23/75) N | (11/13/75) N |

SUMMARY OF PHYTOPLANKTON DATA
 3/10/75 6/23/75 11/13/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|---------------|----------------|--------------|---------------------|---------------|
| CHLAMYDOMONAS | 99798 | CHLAMYDOMONAS | 10045 | CHLOROPHYTAN CELLS | 79650 |
| ANKISTRODESMUS | 34004 | ANABAENOPSIS | 5281 | CHLOROCOCCALAN CELL | 25519 |
| CYCLOTELLA | 2213 | ANKISTRODESMUS | 2485 | ANKISTRODESMUS | 12663 |
| CRYPTOMONAS | 2213 | CYCLOTELLA | 414 | NITZSCHIA | 11406 |
| CHLOROGONIUM | 805 | OSCILLATORIA | 362 | PENNATE DIATOMS | 7733 |
| OTHER | 402 | OTHER | 674 | OTHER | 4544 |
| TOTAL | 139435 | TOTAL | 19261 | TOTAL | 141515 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - FALLEN LEAF LAKE (OLIGOTROPHIC)
 COUNTY - EL DORADO
 STORET NO. - 0608 WORKING PAPER NO. 746, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 43.30 | 5.71 | 1.5 | 1.510 | 64.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 12. | 14. | 12.1 | 0.007 | 0.005 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|---|
| 0.8 | ***** | (3/18/75) N (6/27/75) N (11/4/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/18/75 | 6/27/75 | | 11/4/75 | |
|---------------|---------|------------|-------|-----------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CRYPTOMONAS | 205 | SYNEDRA | 20 | CHROOMONAS | 43 |
| NITZSCHIA | 160 | CYCLOTELLA | 10 | DINOFLAGELLATES | 21 |
| DINOBYRON | 160 | DINOBYRON | 10 | | |
| CYCLOTELLA | 114 | | | | |
| CHLAMYDOMONAS | 91 | | | | |
| OTHER | 114 | OTHER | 0 | OTHER | 0 |
| TOTAL | 844 | TOTAL | 40 | TOTAL | 64 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 50. | 585. | 635. |
| NITROGEN | ***** | ***** | 1810. | 36445. | 38255. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 510. | 20. | 0.11 |
| NITROGEN | 28475. | 26. | 6.7 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| GLEN ALPINE CREEK | 1.037 | 28.0 | 0.011 | 0.699 | 13. | 805. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LAKE HENNESSEY
 COUNTY - NAPA
 STORET NO. - 0609

WORKING PAPER NO. 747, NTIS ACCESSION NO. PB-

/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 134.70 | 3.52 | 10.5 | 1.353 | 1.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| ALKALINITY(MG/L) | MEDIAN 131. | MEDIAN 247. | MEAN SECCHI DISC (METERS) | MEDIAN 0.027 | MEDIAN 0.012 | MEDIAN 0.060 | MEDIAN 0.350 |
|---------------------|----------------|----------------|------------------------------|-----------------|-----------------|-----------------|-----------------|
| CONDUCTIVITY(UMHOS) | | | 2.1 | | | | |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | 4.5 | ALGAL ASSAY CONTROL (MG/L--DRY WT) | 5.4 | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|-----|---------------------------------------|-----|--|
| | | | | (3/13/75) N (6/26/75) N (11/11/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/13/75 | 6/26/75 | | 11/11/75 | |
|----------------|---------|---------------|-------|----------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 2039 | GLOEOCYSTIS | 780 | CLOSTERIUM | 262 |
| ASTERIONELLA | 105 | APHANIZOMENON | 654 | APHANIZOMENON | 75 |
| STEPHANODISCUS | 35 | CHROOMONAS | 503 | CRYPTOMONAS | 37 |
| SYNEDRA | 18 | MELOSIRA | 151 | CHROOMONAS | 37 |
| CRYPTOMONAS | 18 | COELASTRUM | 101 | ANKISTRODESmus | 37 |
| OTHER | 18 | OTHER | 328 | OTHER | 0 |
| TOTAL | 2233 | TOTAL | 2517 | TOTAL | 448 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 5155. | 5155. |
| NITROGEN | ***** | ***** | ***** | 61695. | 61695. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION LOSS | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|------------------------------|---|
| PHOSPHORUS | 7760. | | 1.46 |
| NITROGEN | 39955. | 35. | 17.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| CONN CREEK | 0.483 | 33.4 | 0.232 | 1.325 | 106. | 604. |
| CHILES CREEK | 0.266 | 47.1 | 0.045 | 1.239 | 8. | 221. |
| CLEAR CREEK | 0.354 | 35.5 | 0.048 | 1.174 | 15. | 369. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------|------------------------|------------------------|
| UNNAMED CREEK H1 | 0.054 | 1.587 |
| SAGE CREEK | 0.082 | 1.803 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LAKE MENSHEW (EUTROPHIC)
COUNTY - SAN DIEGO
STORET NO. - 0610 WORKING PAPER NO. 748. NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 533.50 | 24.36 | 6.1 | 0.964 | 5.7 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 170. | 540. | 1.0 | 0.138 | 0.073 | 0.070 | 0.620 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L-DRY WT) (3/7/75) N (6/23/75) N (11/13/75) N

SUMMARY OF PHYTOPLANKTON DATA

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------|------------------------|-------|----------------|-------|
| MALLOMONAS | 7701 | ANACYSTIS(MICROCYSTIS) | 987 | OSCILLATORIA | 681 |
| CHROOMONAS | 553 | ANKISTRODESMUS | 816 | CHROOMONAS | 367 |
| ANKISTRODESMUS | 277 | GLENODINIUM | 258 | ANKISTRODESMUS | 314 |
| CRYPTOMONAS | 184 | CHROOMONAS | 258 | PEDIASTRUM | 209 |
| MELOSIRA | 184 | KIRCHNERIELLA | 215 | MELOSIRA | 209 |
| OTHER | 323 | OTHER | 728 | OTHER | 577 |
| TOTAL | 9222 | TOTAL | 3262 | TOTAL | 2357 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT A. INPUT

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 2115. | 2115. |
| NITROGEN | ***** | ***** | ***** | 70365. | 70365. |

B. OUTPUT

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 5510. | LOSS | | 0.09 |
| NITROGEN | 39460. | 44. | | 2.9 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SAN LUIS REY RIVER | 0.156 | 87.0 | ***** | ***** | 4.* | 85.* |
| W. FK. SAN LUIS REY R. | 0.125 | 66.0 | 0.029 | 1.403 | 2. | 84. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------------|------------------------|------------------------|
| CARISTA CREEK | 0.057 | 1.572 |
| UNNAMED STREAM E-1 | 0.020 | 0.860 |
| UNNAMED STREAM F-1 | 0.047 | 1.559 |
| SPARLING CREEK | 0.065 | 1.232 |
| UNNAMED STREAM H-1 | 0.071 | 1.507 |

* ESTIMATED

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - IRON GATE RESERVOIR (EUTROPHIC)
 COUNTY - SISKIYOU
 STORET NO. - 0611 WORKING PAPER NO. 749, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 11838.90 | 3.39 | 21.4 | 74,483 | 11.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 71. | 150. | 1.5 | 0.184 | 0.124 | 0.690 | 1.520 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (2) | LIMITING NUTRIENT AT SAMPLING TIME | |
|------------------------------|---------------------------------------|----------------|------------------------------------|----------------|
| 6.2 | 26.7 - 33.2 | (3/27/75) N | (7/16/75) N | (10/31/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/27/75 | 7/16/75 | 10/31/75 |
|--|---------|---------|----------|
|--|---------|---------|----------|

| | | | | | |
|-----------------|-------------|---------------|------------|----------------|------------|
| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
| CRYPTOMONAS | 1976 | SCHROEDERIA | 435 | OSCILLATORIA | 133 |
| STEPHANODISCUS | 1297 | SYNEDRA | 73 | CRYPTOMONAS | 80 |
| CHROOMONAS | 247 | CRYPTOMONAS | 73 | APHANIZOMENON | 53 |
| PENNATE DIATOMS | 62 | MELOSIRA | 73 | STEPHANODISCUS | 53 |
| EUGLENA | 62 | APHANIZOMENON | 36 | CHROOMONAS | 27 |
| OTHER | 60 | OTHER | 0 | OTHER | 0 |
| TOTAL | 3704 | TOTAL | 690 | TOTAL | 346 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 372800. | 372800. |
| NITROGEN | ***** | ***** | ***** | 4085210. | 4085210. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 345290. | 7. | 109.97 |
| NITROGEN | 4944425. | LOSS | 1205.1 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| KLAMATH RIVER | 70,400 | 11181.0 | 0.163 | 1.766 | 33. | 356. |
| FALL CREEK | 0.230 | 37.8 | 0.033 | 0.693 | 6. | 133. |
| CAMP CREEK | 0.283 | 43.3 | 0.059 | 0.970 | 12. | 200. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------|------------------------|------------------------|
| JENNY CREEK | 0.036 | 0.686 |
| SCOTHCH CREEK | 0.036 | 1.117 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LOPEZ RESERVOIR
 COUNTY - SAN LUIS OBISPO
 STORET NO. - 0614

WORKING PAPER NO. 750, NTIS ACCESSION NO. PB-

/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 175.30 | 3.80 | 16.6 | 0.336 | 7.3 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 337. | 493. | 3.3 | 0.371 | 0.343 | 0.090 | 0.540 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 8.7 | 4.5 - 12.9 (2) | (3/10/75) N (6/25/75) N (11/12/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/10/75 | 6/25/75 | 11/12/75 |
|--|---------|---------|----------|
|--|---------|---------|----------|

| | | | | | |
|----------------|------------|----------------|------------|---------------|------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CERATIUM | 534 | CHROOMONAS | 298 | CHROOMONAS | 191 |
| FRAGILARIA | 140 | APHANIZOMENON | 216 | APHANIZOMENON | 127 |
| STEPHANODISCUS | 56 | SCHROEDERIA | 162 | CRYPTOMONAS | 127 |
| CRYPTOMONAS | 56 | FRAGILARIA | 81 | CERATIUM | 64 |
| EUGLENA | 28 | STEPHANODISCUS | 54 | OOCYSTIS | 42 |
| OTHER | 0 | OTHER | 55 | OTHER | 21 |
| TOTAL | 814 | TOTAL | 866 | TOTAL | 572 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 5. | 4270. | 4275. |
| NITROGEN | ***** | ***** | 280. | 11705. | 11985. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 2835. | 34. | 1.12 |
| NITROGEN | 7865. | 34. | 3.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| ARROYO GRANDE CREEK | 0.044 | 35.2 | 0.789 | 1.419 | 31. | 56. |
| LOPEZ CRFFK | 0.146 | 55.9 | 0.197 | 0.464 | 16. | 38. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LAKE MARY
 COUNTY - MONO
 STORET NO. - 0615
 WORKING PAPER NO. 751, NTIS ACCESSION NO. PB-282 633/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 0.42 | 8.2 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 21. | 33. | 5.2 | 0.010 | 0.002 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|------------------------------------|
| 2.6 | ***** | (6/30/75) P (11/5/75) N |

SUMMARY OF PHYTOPLANKTON DATA
6/30/75 11/5/75

| GENERA | COUNT | GENERA | COUNT |
|--------------|-------------|--------------|------------|
| SYNEDRA | 568 | ASTERIONELLA | 215 |
| ASTERIONELLA | 487 | DINOBYRON | 132 |
| CRYPTOMONAS | 81 | CRYPTOMONAS | 83 |
| DINOBYRON | 41 | CHROOMONAS | 66 |
| CHROOMONAS | 41 | SYNEDRA | 17 |
| OTHER | 41 | OTHER | 0 |
| TOTAL | 1259 | TOTAL | 513 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | **** | ***** |
| NITROGEN | ***** | **** | ***** |

**** LAKE SAMPLING ONLY ****

32

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LAKE MENDOCINO (MESO-EUTROPHIC)
COUNTY - MENDOCINO
STORET NO. - 0616 WORKING PAPER NO. 752, NTIS ACCESSION NO. PB-282 598/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 271.90 | 7.92 | 19.1 | 10.952 | 168.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|-----------------------------|--------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 76. | 104. | 1.6 | 0.020 | 0.008 | 0.050 | 0.320 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (3/12/75) N (6/26/75) N (11/11/75) N

| | | | | |
|-----|-------|--|--|--|
| 3.1 | ***** | | | |
|-----|-------|--|--|--|

SUMMARY OF PHYTOPLANKTON DATA
3/12/75 6/26/75

| | | | | | |
|------------|-------|----------------|-------|----------------|-------|
| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
| MELOSIRA | 313 | CHROOMONAS | 1188 | CHROOMONAS | 1103 |
| CHROOMONAS | 57 | ANKISTRODESmus | 64 | OSCILLATORIA | 227 |
| | | CRYPTOMONAS | 64 | CRYPTOMONAS | 194 |
| OTHER | 0 | OTHER | 0 | MELOSIRA | 8 |
| TOTAL | 370 | TOTAL | 1316 | STEPHANODISCUS | 8 |
| | | | | OTHER | 115 |
| | | | | TOTAL | 1799 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| ***** | ***** | ***** | 31360. | 31360. |
| ***** | ***** | ***** | 393180. | 393180. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE |
|------------|----------------------|----------------------|----------------------------------|--------------|
| PHOSPHORUS | 19285. | 39. | 3.96 | |
| NITROGEN | 275870. | 30. | 49.6 | |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| EAST FORK RUSSIAN RIVER | 10.480 | 238.8 | 0.079 | 0.984 | 118. | 1457. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------|------------------------|------------------------|
| COLD CREEK | 0.082 | 0.931 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - NICASIO RESERVOIR (EUTROPHIC)
 COUNTY - MARIN
 STORET NO. - 0617 WORKING PAPER NO. 753, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 93.20 | 3.42 | 8.1 | 1.353 | 256.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 67. | 140. | 0.4 | 0.055 | 0.013 | 0.345 | 0.690 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 6.6 | ***** | (3/13/75) N (6/26/75) P (11/11/75) N |

SUMMARY OF PHYTOPLANKTON DATA

3/13/75 6/26/75 11/11/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------|----------------|-------|---------------|-------|
| CRYPTOMONAS | 362 | STEPHANODISCUS | 3156 | CRYPTOMONAS | 297 |
| CHROOMONAS | 226 | CHROOMONAS | 298 | PEDIASTRUM | 231 |
| STEPHANODISCUS | 90 | APHANIZOMENON | 149 | ASTERIONELLA | 99 |
| | | CRUCIGENIA | 124 | MELOSIRA | 66 |
| | | APHANOCAPSA | 99 | TRACHELOMONAS | 33 |
| OTHER | 0 | OTHER | 250 | OTHER | 66 |
| TOTAL | 678 | TOTAL | 4076 | TOTAL | 792 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 2055. | 2055. |
| NITROGEN | ***** | ***** | ***** | 70440. | 70440. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 2575. | LOSS | 0.60 |
| NITROGEN | 68470. | 3. | 20.6 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| NICASIO CREEK | 0.456 | 31.3 | 0.047 | 1.474 | 22. | 677. |
| HALLECK CREEK | 0.328 | 22.5 | 0.046 | 1.684 | 21. | 774. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LOWER OTAY RESERVOIR (EUTROPHIC)
COUNTY - SAN DIEGO
STORET NO. - 0618 WORKING PAPER NO. 754, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 256.40 | 4.58 | 15.2 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 151. | 875. | 1.3 | 0.058 | 0.013 | 0.180 | 0.920 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A (UG/L) ALGAL ASSAY CONTROL YIELD (MG/L-DRY WT) LIMITING NUTRIENT AT SAMPLING TIME
 15.9 ***** (3/7/75) P (6/20/75) N (11/14/75) P AND N

SUMMARY OF PHYTOPLANKTON DATA
3/7/75 6/

| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
|--------------|-------|------------------|-------|------------------|-------|
| ASTERIONELLA | 2170 | APHANIZOMENON | 370 | OSCILLATORIA | 6873 |
| COELASTRUM | 457 | DACTYLOCOCCOPSIS | 148 | CHLAMYDOMONAS | 1430 |
| CRYPTOMONAS | 381 | DINORRYON | 74 | DACTYLOCOCCOPSIS | 440 |
| SCENEDESmus | 304 | OSCILLATORIA | 74 | SCENEDESmus | 440 |
| OOCYSTIS | 266 | ANABAENA | 74 | CRYPTOMONAS | 385 |
| OTHER | 381 | OTHER | 75 | OTHER | 329 |
| TOTAL | 3959 | TOTAL | 815 | TOTAL | 9897 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|---|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |
| R. OUTPUT | | | | | |
| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) | | |
| PHOSPHORUS | ***** | **** | ***** | | |
| NITROGEN | ***** | **** | ***** | | |

***** LAKE SAMPLING ONLY *****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LAKE PILLSHURY (EUTROPHIC)
 COUNTY - LAKE
 STORET NO. - 0619 WORKING PAPER NO. 755, NTIS ACCESSION NO. PB-282 646/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 748.50 | 8.10 | 14.3 | 16.067 | 87.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 60. | 86. | 0.8 | 0.022 | 0.008 | 0.060 | 0.230 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 6.4 | 0.6 | (3/12/75) N (6/26/75) P AND N (11/11/75) N |

SUMMARY OF PHYTOPLANKTON DATA
3/12/75 6/26/75 11/11/75

| GENERA SYNEDRA EUGLENA | COUNT 10 10 | GENERA ANARAENA CRYPTOMONAS TRACHELOMONAS | COUNT 1861 64 64 | GENERA MELOSIRA CRYPTOMONAS DACTYLOCOCCOPSIS CHROOMONAS | COUNT 331 37 37 36 |
|------------------------------|-------------------|--|---------------------------|---|--------------------------------|
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 20 | TOTAL | 1989 | TOTAL | 441 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 10. | 26005. | 26015. |
| NITROGEN | ***** | ***** | 425. | 455255. | 455680. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 30910. | LOSS | 3.21 |
| NITROGEN | 599395. | LOSS | 56.3 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| EEL RIVER | 7.530 | 362.6 | 0.051 | 0.720 | 36. | 640. |
| RICE FORK | 4.720 | 224.6 | 0.026 | 0.534 | 20. | 349. |
| SQUAW VALLEY CREEK | 0.215 | 9.1 | 0.052 | 0.845 | 119. | 1769. |
| SMOKEHOUSE CREEK | 0.775 | 32.9 | 0.061 | 0.653 | 53. | 528. |
| SALMON CREEK | 0.771 | 32.6 | 0.036 | 1.045 | 31. | 965. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - SANTA MARGARITA LAKE (EUTROPHIC)
 COUNTY - SAN LUIS OBISPO
 STORET NO. - 0620 WORKING PAPER NO. 756. NTIS ACCESSION NO. PB-282 779/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 290.10 | 3.18 | 10.1 | 0.747 | 1.6 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 125. | 291. | 2.5 | 0.037 | 0.014 | 0.070 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 9.1 | 8.0 | (3/10/75) N (6/25/75) N (11/12/75) N |

SUMMARY OF PHYTOPLANKTON DATA
3/10/75 6/25/75 11/12/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|--------------|-------|--------------|-------|------------------|-------|
| ASTERIONELLA | 4920 | MOUGEOTIA | 2181 | DACTYLOCOCCOPSIS | 304 |
| CRYPTOMONAS | 109 | SYNEDRA | 488 | CERATIUM | 152 |
| | | DINOBYRON | 163 | STEPHANODISCUS | 122 |
| | | ANABAENOPSIS | 130 | DINOBYRON | 91 |
| | | OSCILLATORIA | 130 | GLENODINIUM | 61 |
| OTHER | 0 | OTHER | 98 | OTHER | 121 |
| TOTAL | 5029 | TOTAL | 3190 | TOTAL | 851 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 5. | 1535. | 1540. |
| NITROGEN | ***** | ***** | 210. | 12880. | 13090. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 935. | 39. | 0.48 |
| NITROGEN | 14135. | LOSS | 4.1 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SALINAS RIVER | 0.501 | 182.1 | 0.039 | 0.259 | 3. | 22. |
| TORO CREEK | 0.017 | 24.8 | 0.102 | 0.688 | 5. | 16. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------|------------------------|------------------------|
| ALAMO CREEK | 0.102 | 0.688 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - SHASTA LAKE (MESOTROPHIC)
 COUNTY - SHASTA
 STORET NO. - 0621 WORKING PAPER NO. 757, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 16630.40 | 119.40 | 46.5 | 295.380 | 217.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 57. | 98. | 3.0 | 0.021 | 0.015 | 0.060 | 0.240 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 4.1 | ***** | (3/26/75) N (7/14/75) N (11/ 3/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/26/75 | 7/14/75 | | 11/ 3/75 |
|--|---------|---------|--|----------|
|--|---------|---------|--|----------|

| | | | | | |
|----------------|-------------|-----------------------|------------|--------------|------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| STEPHANODISCUS | 1475 | CRYPTOMONAS | 85 | MELOSIRA | 637 |
| ASTERIONELLA | 413 | SYNEDRA | 42 | SYNEDRA | 80 |
| PERIDINIUM | 59 | STEPHANODISCUS | 42 | CHROOMONAS | 80 |
| CRYPTOMONAS | 30 | CHLOROPHYTAN COLONIES | 42 | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 1977 | TOTAL | 211 | TOTAL | 797 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 4760. | ***** | ***** | 359855. | 364615. |
| NITROGEN | 12800. | ***** | ***** | 11399335. | 11412135. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 316715. | 13. | 3.05 |
| NITROGEN | 13283340. | LOSS | 95.6 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SACRAMENTO RIVER | 40.950 | 1100.7 | 0.033 | 1.584 | 34. | 1847. |
| SALT CREEK C1 | 0.854 | 35.0 | 0.020 | 0.982 | 15. | 756. |
| MCLOUD RIVER | 26.750 | 1585.1 | 0.021 | 0.815 | 11. | 434. |
| SQUAW CREEK B1 | 6.310 | 158.0 | 0.014 | 0.587 | 18. | 739. |
| PIT RIVER | 172.120 | 12823.1 | 0.050 | 1.165 | 21. | 493. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|----------------|------------------------|------------------------|
| NOSONI CREEK | 0.018 | 1.365 |
| SQUAW CREEK F1 | 0.019 | 0.876 |
| SALT CREEK H1 | 0.022 | 1.153 |
| POTEM CREEK | 0.018 | 0.765 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - SHAVER LAKE (MESOTROPHIC)
COUNTY - FRESNO
STORET NO. - 0622 WORKING PAPER NO. 758. NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 75.40 | 8.39 | 19.9 | 7,649 | 252.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| PHYSICAL AND CHEMICAL CHARACTERISTICS | | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|---------------------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | 3.9 | 0.014 | 0.004 | 0.060 | 0.250 |
| 13. | 18. | | | | | |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A (UG/L) ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) LIMITING NUTRIENT AT SAMPLING TIME
 1.7 1.0 (6/4/75) P (6/24/75) N (11/13/75) N

SUMMARY OF PHYTOPLANKTON DATA

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------|-------------|-------|----------------|-------|
| SYNEDRA | 27 | CHROOMONAS | 193 | CRYPTOMONAS | 153 |
| PERIDINIUM | 20 | MALLOMONAS | 97 | ANKISTRODESMUS | 44 |
| ASTERIONELLA | 20 | CRYPTOMONAS | 32 | ACHNANTHES | 22 |
| ANKISTRODESMUS | 7 | DINOBRYON | 32 | | |
| | | SYNEDRA | 32 | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 74 | TOTAL | 386 | TOTAL | 219 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|------------------|
| PHOSPHORUS | ***** | ***** | 10. | 870. | 880. |
| NITROGEN | ***** | ***** | 400. | 37295. | 37695. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 3620. | LOSS | | 0.10 |
| NITROGEN | 66825. | LOSS | | 4.5 |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------------|------------------------|------------------------|
| N. FK. STEVENSON CREEK | 0.025 | 1.565 |
| MARKWOOD CREEK | 0.019 | 0.740 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - SILVER LAKE (OLIGOTROPHIC)
 COUNTY - MONO
 STORET NO. - 0623 WORKING PAPER NO. 759. NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 0.45 | 10.5 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 23. | 33. | 3.7 | 0.012 | 0.003 | 0.055 | 0.225 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 1.8 | ***** | (6/10/75) N (6/30/75) P (11/ 5/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 6/10/75 | 6/30/75 | 11/ 5/75 | | |
|--------------|---------|--------------|----------|-------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 343 | CHROOMONAS | 228 | CHROOMONAS | 57 |
| DINOBYRON | 115 | DINOBYRON | 114 | CRYPTOMONAS | 45 |
| ASTERIONELLA | 38 | SYNEDRA | 114 | DINOBYRON | 11 |
| CRYPTOMONAS | 38 | ASTERIONELLA | 76 | | |
| SYNEDRA | 38 | | | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 572 | TOTAL | 532 | TOTAL | 113 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - TULLOCK RESERVOIR (EUTROPHIC)
COUNTY - CALAVERAS, TUOLUMNE
STORED NO. - 0624 WORKING PAPER NO. 760, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 2533.30 | 5.10 | 16.5 | 51.590 | 19.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN (METERS) | SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------|-------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 33. | 65. | 1.7 | 0.025 | 0.009 | 0.060 | 0.280 | |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L-DRY WT) (3/14/75) N (6/25/75) N (11/12/75) P

SUMMARY OF PHYTOPLANKTON DATA
3/14/75 6/

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------|----------------|-------|----------------|-------|
| ACHNANTHES | 3864 | ASTERIONELLA | 432 | OSCILLATORIA | 225 |
| ASTERIONELLA | 2006 | STEPHANODISCUS | 118 | GLENODINIUM | 112 |
| SYNEDRA | 1783 | FRAGILARIA | 79 | ASTERIONELLA | 37 |
| STEPHANODISCUS | 743 | OSCILLATORIA | 79 | CRYPTOMONAS | 37 |
| CRYPTOMONAS | 297 | DINOBRYON | 79 | STEPHANODISCUS | 37 |
| OTHER | 744 | OTHER | 195 | OTHER | 113 |
| TOTAL | 9437 | TOTAL | 982 | TOTAL | 561 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NOVEMBER

| POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS ***** | ***** | 10. | 37410. | 37420. |
| NITROGEN ***** | ***** | 425. | 1528520. | 1528945. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 32540. | 13. | 7.34 |
| NITROGEN | 1506550. | 1. | 299.8 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| STANISLAUS RIVER | 49.840 | 2343.9 | 0.022 | 0.916 | 15. | 614. |
| BLACK CREEK | 0.242 | 49.5 | 0.022 | 1.473 | 3. | 227. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------|------------------------|------------------------|
| GREEN SPRING RUN | 0.054 | 1.515 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - LOWER TWIN LAKE (MESOTROPHIC)
 COUNTY - MONO
 STORET NO. - 0626 WORKING PAPER NO. 761, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 101.30 | 1.52 | 15.2 | 1.699 | 157.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 26. | 25. | 6.4 | 0.014 | 0.003 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 2.9 | ***** | (3/19/75) N (7/ 1/75) P (11/ 6/75) P |

SUMMARY OF PHYTOPLANKTON DATA

3/19/75 7/ 1/75 11/ 6/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|------------|----------------|------------|--------------|------------|
| CHROOMONAS | 275 | MELOSIRA | 81 | FRAGILARIA | 72 |
| MELOSIRA | 78 | CRYPTOMONAS | 33 | CRYPTOMONAS | 72 |
| SYNEDRA | 78 | STEPHANODISCUS | 16 | CHROOMONAS | 72 |
| STEPHANODISCUS | 39 | SYNEDRA | 16 | TABELLARIA | 36 |
| ASTERIONELLA | 39 | FRAGILARIA | 16 | | |
| OTHER | 40 | OTHER | 0 | OTHER | 0 |
| TOTAL | 549 | TOTAL | 162 | TOTAL | 252 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 10. | 825. | 835. |
| NITROGEN | ***** | ***** | 360. | 48840. | 49200. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 695. | 17. | 0.55 |
| NITROGEN | 51060. | LOSS | 32.4 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| ROBINSON CREEK | 1.303 | 76.4 | 0.015 | 0.730 | 8. | 474. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------|------------------------|------------------------|
| UNNAMED CREEK B1 | 0.018 | 1.142 |

42

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN CALIFORNIA

NAME - UPPER TWIN LAKE (MESOTROPHIC)
 COUNTY - MONO
 STORET NO. - 0625 WORKING PAPER NO. 762, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 76.40 | 1.07 | 14.3 | 1.303 | 136.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 25. | 32. | 5.1 | 0.015 | 0.004 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 3.3 | ***** | (6/10/75) N (7/ 1/75) P (11/ 6/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 6/10/75 | 7/ 1/75 | 11/ 6/75 |
|--|---------|---------|----------|
|--|---------|---------|----------|

| | | | | | |
|----------------|-------|----------------|-------|-------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| MELOSIRA | 122 | SYNEDRA | 265 | MELOSIRA | 231 |
| STEPHANODISCUS | 61 | ASTERIONELLA | 166 | CRYPTOMONAS | 115 |
| TABELLARIA | 61 | STEPHANODISCUS | 33 | DINOBYRON | 38 |
| ANABAENA | 31 | MELOSIRA | 33 | | |
| | | CHROOMONAS | 33 | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 275 | TOTAL | 530 | TOTAL | 384 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 5. | ***** | 5. | 885. | 895. |
| NITROGEN | 235. | ***** | 130. | 32675. | 33040. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 615. | 31. | 0.84 |
| NITROGEN | 36200. | LOSS | 30.9 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| ROBINSON CREEK | 0.889 | 54.4 | 0.021 | 0.767 | 11. | 395. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - BARKER RESERVOIR (MESOTROPHIC)
 COUNTY - BOULDER
 STORET NO. - 0801 WORKING PAPER NO. 765, NTIS ACCESSION NO. PB-271 470/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 99.20 | 1.70 | 8.3 | 1.608 | 104.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 18. | 29. | 2.1 | 0.023 | 0.006 | 0.045 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L.) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|-------------------------------|---------------------------------------|---|
| 5.3 | ***** | (5/ 7/75) P AND N (8/26/75) N. (10/10/75) N |

SUMMARY OF PHYTOPLANKTON DATA
 5/ 7/75 8/26/75 10/10/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|--------------|--------------|--------------|-------------|---------------|-------------|
| MELOSIRA | 9734 | FRAGILARIA | 1381 | CHLAMYDOMONAS | 6478 |
| CHROOMONAS | 721 | SPONDYLOSUM | 563 | MELOSIRA | 748 |
| FRAGILARIA | 613 | CHROOMONAS | 256 | CHROOMONAS | 208 |
| CRYPTOMONAS | 216 | MELOSIRA | 104 | OSCILLATORIA | 125 |
| MALLOMONAS | 72 | ANABAENA | 51 | CRYPTOMONAS | 42 |
| OTHER | 108 | OTHER | 203 | OTHER | 81 |
| TOTAL | 11464 | TOTAL | 2558 | TOTAL | 7682 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 455. | ***** | ***** | 550. | 1005. |
| NITROGEN | 1640. | ***** | ***** | 29880. | 31520. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 860. | 14. | 0.59 |
| NITROGEN | 26340. | 16. | 18.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| MIDDLE BOULDER CREEK | 1.550 | 93.8 | 0.016 | 0.623 | 5. | 288. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - BARR LAKE
COUNTY - ADAMS
STORET NO. - 0802

(FUTROPHIC)

WORKING PAPER NO. 766. NTIS ACCESSION NO. PB-278 507/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | ***** | 7.57 | 5.2 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 162 | 581 | 1.2 | 0.930 | 0.730 | 1.090 | 2.960 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

• BIOLOGICAL CHARACTERISTICS (LAKE)
 MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (5/ 5/75) N (8/26/75) N (10/10/75) N
 28.8 42.8 - 186.3 (2)

SUMMARY OF PHYTOPLANKTON DATA
5/ 5/75 8/

| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
|-------------|-------|-------------------------|-------|-------------------------|-------|
| CHROOMONAS | 1148 | APHANIZOMENON | 3265 | APHANIZOMENON | 1012 |
| CRYPTOMONAS | 836 | GOMPHOSPHAERIA | 96 | MELOSIRA | 156 |
| SCHROEDERIA | 313 | ANACYSTIS (MICROCYSTIS) | 96 | ANACYSTIS (MICROCYSTIS) | 78 |
| CYSTS | 52 | PHORMIDIUM | 48 | NITZSCHIA | 78 |
| OTHER | 0 | OTHER | 0 | CHROOMONAS | 78 |
| TOTAL | 2349 | TOTAL | 3505 | OTHER | 0 |
| | | | | | 1402 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE |
|------------|----------------------|----------------------|----------------------------------|--------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P | MEAN TOTAL N |
|---------------------|--------------|--------------|
| | (MG/L) | (MG/L) |
| SPEER CANAL | 1.664 | 6.565 |
| O'BRIAN CANAL | 1.960 | 7.701 |
| DENVER-HUDSON CANAL | 2.441 | 6.851 |
| UNNAMED DITCH (C1) | 0.270 | 1.855 |
| UNNAMED STREAM (D1) | 1.611 | 5.861 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - BLUE MESA RESERVOIR (MESOTROPHIC)
COUNTY - GUNNISON
STORET NO. - 0803 WORKING PAPER NO. 767, NTIS ACCESSION NO. PB-271 658/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 8873.30 | 36.62 | 31.1 | 47,647 | 325.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN (METERS) | SECCHI DISC | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------|----------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 91. | 149. | 2.6 | | 0.019 | 0.005 | 0.040 | 0.320 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

SUMMARY OF PHYTOPLANKTON DATA

| GENERA | COUNT | GENERA | COUNT |
|-------------------------|-------------|---------------|------------|
| APHANIZOMENON | 1324 | APHANIZOMENON | 439 |
| CHROOMONAS | 461 | CHROOMONAS | 18 |
| CRYPTOMONAS | 230 | FRAGILARIA | 9 |
| FRAGILARIA | 115 | CRYPTOMONAS | 9 |
| ANACYSTIS (MICROCYSTIS) | 115 | SCHROEDERIA | 4 |
| OTHER | 231 | OTHER | 0 |
| TOTAL | 2476 | TOTAL | 481 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 3160. | ***** | 10. | 94390. | 97560. |
| NITROGEN | 11655. | ***** | 355. | 1187820. | 1199830. |

b. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 30730. | 69. | 2.66 | |
| NITROGEN | 817720. | 32. | 32.8 | |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT SOURCE NUTRIENT EXPORT | STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------------------------|--------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| | GUNNISON RIVER | 24.780 | 5524.5 | 0.071 | 1.115 | 9. | 143. |
| | LAKE FORK GUNNISON RIVER | 7.430 | 955.7 | 0.068 | 0.723 | 15. | 172. |
| | CEBOLLA CREEK | 3.130 | 955.7 | 0.075 | 0.488 | 8. | 49. |
| | WILLOW CREEK | 0.007 | 186.5 | 0.223 | 1.493 | 43. | 380. |
| | SOAP CREEK | 1.950 | 181.3 | 0.064 | 0.379 | 24. | 144. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------------|------------------------|------------------------|
| CEBOLLA CREEK (C2) | 0.115 | 0.907 |
| STEUBEN CREEK | 0.067 | 0.893 |
| EAST ELK CREEK | 0.187 | 0.736 |
| RED CREEK | 0.130 | 1.005 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - CHERRY CREEK LAKE (EUTROPHIC)
 COUNTY - ARAPAHOE
 STORET NO. - 0804 WORKING PAPER NO. 768, NTIS ACCESSION NO. PB-271 473/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 997.10 | 3.24 | 5.2 | 0.208 | 3.6 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 177. | 422. | 0.8 | 0.054 | 0.007 | 0.040 | 0.770 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 23.3 | 3.2 | (5/ 7/75) N (8/22/75) N (10/ 9/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/ 7/75 | 8/22/75 | 10/ 9/75 | |
|----------------|---------|----------------|----------|-------|
| GENERA | COUNT | GENERA | COUNT | |
| CHROOMONAS | 841 | FRAGILARIA | 920 | |
| CRYPTOMONAS | 342 | DIATOMA | 920 | |
| STEPHANODISCUS | 156 | NAVICULA | 307 | |
| NAVICULA | 31 | STEPHANODISCUS | 184 | |
| ANKISTRODESMUS | 31 | CRYPTOMONAS | 184 | |
| OTHER | 280 | OTHER | 367 | |
| TOTAL | 1681 | TOTAL | 2882 | |
| | | | | COUNT |
| | | | | 373 |
| | | | | 248 |
| | | | | 124 |
| | | | | 41 |
| | | | | 41 |
| | | | | 84 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 5. | 4940. | 4945. |
| NITROGEN | ***** | ***** | 245. | 26095. | 26340. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 745. | 85. | 1.53 |
| NITROGEN | 6870. | 74. | 8.1 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| CHERRY CREEK | 0.196 | 932.4 | 0.788 | 3.615 | 5. | 24. |
| COTTONWOOD CREEK | 0.008 | 20.1 | 0.044 | 0.656 | 0.5 | 8. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - DILLON RESERVOIR (OLIGOTROPHIC)
 COUNTY - SUMMIT
 STORET NO. - 0806 WORKING PAPER NO. 769, NTIS ACCESSION NO. PB-278 484/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 867.60 | 12.76 | 24.6 | 9.610 | 1.3 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 42. | 86. | 8.1 | 0.009 | 0.002 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|------------------------------------|
| 3.2 | ***** | (8/25/75) P (10/ 9/75) P |

SUMMARY OF PHYTOPLANKTON DATA
8/25/75 10/ 9/75

| GENERA | COUNT | GENERA | COUNT |
|--------------|-------------|--------------|-------------|
| FLAGELLATES | 1537 | FRAGILARIA | 3297 |
| CRYPTOMONAS | 713 | ASTERIONELLA | 1486 |
| ASTERIONELLA | 604 | FLAGELLATES | 650 |
| FRAGILARIA | 384 | CRYPTOMONAS | 279 |
| DINOBRYON | 384 | CYSTS | 232 |
| OTHER | 0 | OTHER | 93 |
| TOTAL | 3622 | TOTAL | 6037 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 2365. | ***** | 10. | 5395. | 7770. |
| NITROGEN | 15635. | ***** | 315. | 191310. | 207260. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 2225. | 71. | 0.61 |
| NITROGEN | 174050. | 16. | 16.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BLUE RIVER | 2.440 | 313.4 | 0.012 | 0.444 | 1. | 85. |
| TENMILE CREEK | 2.630 | 240.4 | 0.022 | 0.850 | 10. | 302. |
| MINERS CREEK | 0.270 | 17.3 | 0.011 | 0.527 | 5. | 266. |
| SNAKE RIVER | 2.080 | 201.0 | 0.011 | 0.385 | 3. | 121. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------|------------------------|------------------------|
| MEADOW CREEK | 0.021 | 0.714 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - GRAND LAKE (MESOTROPHIC)
 COUNTY - GRAND *** INCLUDES TRIBS AND LOADS FOR SHADOW MTN LAKE (0813)
 STORET NO. - 0807 WORKING PAPER NO. 770. NTIS ACCESSION NO. PB-272 233/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 479.10 | 2.05 | 41.3 | 13,465 | 92.0 * |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 19. | 19. | 3.4 | 0.013 | 0.003 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (2) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--------------|------------------------------------|
| 4.9 | 0.2 - | 0.4 (2) | (8/26/75) P |
| | | | (10/10/75) P |

SUMMARY OF PHYTOPLANKTON DATA
8/26/75 10/10/75

| GENERA | COUNT | GENERA | COUNT |
|--------------|-------------|---------------|------------|
| ASTERIONELLA | 523 | SPONDYLOSIMUM | 339 |
| ANABAENA | 262 | CRYPTOMONAS | 136 |
| CRYPTOMONAS | 131 | MELOSIRA | 136 |
| MELOSIRA | 87 | ASTERIONELLA | 102 |
| OTHER | 0 | ANABAENA | 34 |
| | | OTHER | 33 |
| TOTAL | 1003 | TOTAL | 780 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 920. | ***** | 45. | 9455. | 10420. |
| NITROGEN | 3695. | ***** | 1715. | 397275. | 402685. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 5980. | 43. | 1.38* |
| NITROGEN | 199480. | 50. | 53.5* |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| NORTH INLET | 1.990 | 118.9 | 0.020 | 0.673 | 11. | 355. |
| ECHO CREEK | 1.441 | 70.4 | 0.011 | 0.461 | 7. | 298. |
| COLORADO RIVER | 2.500 | 264.2 | 0.030 | 1.056 | 9. | 315. |
| GRANBY PUMP CANAL | 7.310 | ***** | 0.022 | 1.039 | ***** | ***** |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------------|------------------------|------------------------|
| LITTLE COLUMBINE CREEK | 0.032 | 0.370 |
| ONAHU CREEK | 0.022 | 0.657 |
| UNNAMED CREEK | 0.019 | 0.451 |

* COMBINED GRAND-SHADOW MOUNTAIN LAKES.

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - SHADOW MOUNTAIN LAKE (MESOTROPHIC)
 COUNTY - GRAND *** SEE GRAND LAKE (0807) FOR LOADING VALUES
 STORET NO. - 0813 WORKING PAPER NO. 770, NTIS ACCESSION NO. PB-272 233/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 471.60 | 5.48 | 3.8 | 13.465 | 92.0 * |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 27. | 33. | 1.8 | 0.020 | 0.003 | 0.040 | 0.300 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--------------------------------------|
| 5.7 | ***** | (8/26/75) P AND N (10/10/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA
8/26/75 10/10/75

| GENERA | COUNT | GENERA | COUNT |
|------------------------|-------------|-----------------|-------------|
| APHANIZOMENON | 630 | MELOSIRA | 706 |
| PHORMIDIUM | 533 | ASTERIONELLA | 246 |
| ANACYSTIS(MICROCYSTIS) | 388 | CRYPTOMONAS | 246 |
| CHROOMONAS | 291 | CHROOMONAS | 184 |
| NITZSCHIA | 242 | PENNATE DIATOMS | 184 |
| OTHER | 486 | OTHER | 214 |
| TOTAL | 2570 | TOTAL | 1780 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

50

* COMBINED GRAND-SHADOW MOUNTAIN LAKES.

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - GREEN MOUNTAIN RESERVOIR (OLIGOTROPHIC)
 COUNTY - SUMMIT
 STORET NO. - 0808 WORKING PAPER NO. 771, NTIS ACCESSION NO. PB-272 235/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 1551.40 | 8.60 | 22.2 | 13.460 | 164.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 54. | 107. | 2.8 | 0.010 | 0.002 | 0.040 | 0.230 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|------------------------------------|
| 5.8 | 0.1 | (8/25/75) P (10/ 9/75) P |

SUMMARY OF PHYTOPLANKTON DATA
8/25/75 10/ 9/75

| GENERA | COUNT | GENERA | COUNT |
|--------------|-------------|--------------|------------|
| FRAGILARIA | 3470 | CYCLOTELLA | 635 |
| CYCLOTELLA | 132 | FRAGILARIA | 212 |
| ASTERIONELLA | 132 | CRYPTOMONAS | 60 |
| CRYPTOMONAS | 88 | | |
| OTHER | 0 | OTHER | 0 |
| TOTAL | 3822 | TOTAL | 907 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 895. | ***** | 10. | 5680. | 6585. |
| NITROGEN | 22350. | ***** | 355. | 152755. | 175460. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 6390. | 3. | 0.77 |
| NITROGEN | 168530. | 4. | 20.4 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BLUE RIVER | 9.640 | 1323.5 | 0.016 | 0.435 | 3. | 78. |
| BLACK CREEK | 0.862 | 47.9 | 0.014 | 0.364 | 8. | 223. |
| OTTER CREEK | 0.131 | 21.8 | 0.075 | 0.481 | 16. | 94. |
| CATARACT CREEK | 0.537 | 36.3 | 0.011 | 0.317 | 6. | 154. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - HOLBROOK RESERVOIR (EUTROPHIC)

COUNTY - OTERO

STORET NO. - 0809

WORKING PAPER NO. 772, NTIS ACCESSION NO. PB-271 554/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | ***** | 2.72 | 2.5 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN (METERS) | SECCHI DISC | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------|-------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 85. | 1812. | 0.3 | | 0.329 | 0.028 | 0.070 | 3.020 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A (UG/L) ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) LIMITING NUTRIENT AT SAMPLING TIME (5/6/75) N (8/22/75) P (10/7/75) N

SUMMARY OF PHYTOPLANKTON DATA
5/6/75

| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
|----------------|--------------|--------------|--------------|----------------|--------------|
| OSCILLATORIA | 36005 | ANABAENOPSIS | 34071 | ANABAENOPSIS | 31245 |
| SYNEDRA | 16440 | OSCILLATORIA | 11005 | OSCILLATORIA | 13077 |
| FLAGELLATES | 6083 | ANABAENA | 2111 | NITZSCHIA | 7487 |
| STEPHANODISCUS | 2137 | CRYPTOMONAS | 2111 | CRYPTOMONAS | 2595 |
| CRYPTOMONAS | 1973 | SCENEDESMUS | 1508 | STEPHANODISCUS | 799 |
| OTHER | 4185 | OTHER | 1808 | OTHER | 1099 |
| TOTAL | 66823 | TOTAL | 52614 | TOTAL | 56302 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

NITROGEN & OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | **** | ***** |
| NITROGEN | ***** | **** | ***** |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------------|------------------------|------------------------|
| OUTLET DITCH (A1) | 0.210 | 1.955 |
| DITCH (A2) | 0.440 | 3.630 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - LAKE MEREDITH (EUTROPHIC)
COUNTY - CROWLEY
STORET NO. - 0810 WORKING PAPER NO. 773, NTIS ACCESSION NO. PB-271 478/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | ***** | 13.03 | 2.5 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN (METERS) | SECCHI DISC TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------|-------------------------------|--------------------------|--------------------------|--------------------------|
| 100. | 7055. | 0.3 | 0.397 | 0.098 | 0.110 | 4.630 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

• BIOLOGICAL CHARACTERISTICS (CONT.)
 MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L-DRY WT) (5/6/75) N (8/22/75) N (10/7/75) N

SUMMARY OF PHYTOPLANKTON DATA

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------|-------------------------|--------|----------------|--------|
| OSCILLATORIA | 14778 | MERISMOPEDIA | 85437 | OSCILLATORIA | 932405 |
| CYLINDROCYSTIS | 6584 | OSCILLATORIA | 63972 | ANABAENOPSIS | 13091 |
| FLAGELLATES | 3438 | ANABAENA | 11477 | COELOSPHAERIUM | 2036 |
| APHANOCAPSAS | 1756 | ANABAENOPSIS | 3400 | CYLINDROCYSTIS | 1455 |
| EUGLENA | 1171 | ANACYSTIS (MICROCYSTIS) | 1913 | APHANOTHECE | 1164 |
| OTHER | 4609 | OTHER | 7013 | OTHER | 3782 |
| TOTAL | 32336 | TOTAL | 173212 | TOTAL | 953933 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NOTRIENT A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

NITROGEN B. OXIDE

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | ***** | **** | ***** | ***** |
| NITROGEN | ***** | **** | ***** | ***** |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN NUTRIENT CONCENTRATIONS IN UNAGED STREAMS | |
|----------------------|--|------------------------|
| | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
| LAKE MEREDITH OUTLET | 0.120 | 5.405 |
| LAKE MEREDITH INLET | 0.071 | 4.174 |
| UNNAMED DITCH (B1) | 0.220 | 3.267 |
| UNNAMED DITCH (C1) | 0.153 | 3.717 |
| UNNAMED DITCH (D1) | 0.070 | 7.827 |
| BOB CREEK | 0.053 | 4.912 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - MILTON RESERVOIR

(EUTROPHIC)

COUNTY - WELD

STORET NO. - 0811

WORKING PAPER NO. 774, NTIS ACCESSION NO. PB-271 469/AB

I. MORPHOMETRY

| LAKE TYPE IMPOUNDMENT | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|--------------------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| ***** | 8.41 | 7.3 | ***** | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| ALKALINITY(MG/L) 324. | MEDIAN CONDUCTIVITY(UMHOS) 1078. | MEAN SECCHI DISC (METERS) 1.8 | MEDIAN TOTAL P(MG/L) 0.846 | MEDIAN ORTHO P(MG/L) 0.808 | MEDIAN INORG N(MG/L) 2.280 | MEDIAN TOTAL N(MG/L) 3.530 |
|--------------------------|--|-------------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
|--------------------------|--|-------------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) 5.9 | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) 7.2 | LIMITING NUTRIENT AT SAMPLING TIME (5/ 6/75) N (8/26/75) N (10/10/75) N |
|-------------------------------------|--|--|
|-------------------------------------|--|--|

SUMMARY OF PHYTOPLANKTON DATA

| 5/ 6/75 | 8/26/75 | 10/10/75 | | | |
|-------------|---------|--------------|-------|-----------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| PEDIASTRUM | 102 | MELOSIRA | 590 | CHROOMONAS | 272 |
| CRYPTOMONAS | 102 | FRAGILARIA | 553 | NITZSCHIA | 136 |
| CHROOMONAS | 102 | CHROOMONAS | 406 | CYCLOTELLA | 91 |
| SCHROEDERIA | 51 | ASTERIONELLA | 369 | PENNATE DIATOMS | 91 |
| SCENEDESMUS | 51 | ANABAENA | 332 | APHANIZOMENON | 45 |
| OTHER | 0 | OTHER | 257 | OTHER | 89 |
| TOTAL | 408 | TOTAL | 2507 | TOTAL | 724 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS ***** | ***** | ***** | ***** | ***** |
| NITROGEN ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|----------------------|----------------------|---|
| PHOSPHORUS ***** | ***** | ***** |
| NITROGEN ***** | ***** | ***** |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------------|------------------------|------------------------|
| GILMORE DITCH | 0.980 | 3.286 |
| HEEBE CANAL | 0.501 | 3.730 |
| PLATTE VALLEY CANAL | 1.357 | 6.067 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN COLORADO

NAME - NAVAJO RESERVOIR (MESO-EUTROPHIC)
 COUNTY - ARCHULETA, CO; SAN JUAN, RIO ARRIBA, NM
 STORET NO. - 0812 WORKING PAPER NO. 775, NTIS ACCESSION NO. PB-278 545/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 8443.40 | 63.25 | 33.3 | 49,460 | 1.4 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 71. | 166. | 0.5 | 0.036 | 0.013 | 0.050 | 0.365 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 2.2 | ***** | (4/30/75) P AND N (8/18/75) P AND N (9/30/75) N |

SUMMARY OF PHYTOPLANKTON DATA
8/18/75 9/30/75

| GENERA | COUNT | GENERA | COUNT |
|---------------|------------|--------------|------------|
| CRYPTOMONAS | 253 | CHROOMONAS | 605 |
| APHANIZOMENON | 63 | CRYPTOMONAS | 287 |
| OTHER | 0 | OTHER | 0 |
| TOTAL | 316 | TOTAL | 892 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 2510. | ***** | 10. | 155570. | 158090. |
| NITROGEN | 7205. | ***** | 360. | 1769425. | 1776990. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 27970. | 82. | 2.50 |
| NITROGEN | 1403140. | 21. | 28.1 |

| V. NON-POINT-SOURCE NUTRIENT EXPORT | STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------------------------------|-------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SAN JUAN RIVER | | 17.440 | 3185.7 | 0.102 | 1.137 | 17. | 203. |
| PIEDRA RIVER | | 14.720 | 1629.1 | 0.076 | 0.972 | 35. | 294. |
| SAMBrito CREEK | | 0.064 | 46.6 | 0.121 | 1.488 | 6. | 65. |
| UNNAMED STREAM (D1) | | 0.010 | 20.7 | 0.159 | 1.579 | 18. | 273. |
| LOS PINOS RIVER | | 5.620 | 1320.9 | 0.070 | 0.973 | 7. | 127. |
| UNNAMED STREAM (F1) | | 0.052 | 18.6 | 0.140 | 1.333 | 31. | 115. |
| UNNAMED STREAM (H1) | | 0.040 | 17.4 | 0.015 | 0.831 | 1. | 70. |
| SPRING CREEK | | 0.336 | 150.2 | 0.324 | 1.880 | 14. | 116. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - AMERICAN FALLS RESERVOIR (EUTROPHIC)
 COUNTY - BANNOCK, BINGHAM, POWER
 STORET NO. - 1601 WORKING PAPER NO. 776, NTIS ACCESSION NO. PB-272 237/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 35224.00 | 277.03 | 9.2 | 182.140 | 109.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 150. | 324. | 0.9 | 0.105 | 0.035 | 0.080 | 0.575 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 15.4 | 0.2 - 16.0 (2) | (5/15/75) P AND N (8/5/75) N (9/18/75) N |

SUMMARY OF PHYTOPLANKTON DATA
 5/15/75 8/ 5/75 9/18/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|--------------|--------------|---------------|-------------|---------------|-------------|
| CYCLOTELLA | 4856 | APHANIZOMENON | 3684 | APHANIZOMENON | 745 |
| ASTERIONELLA | 2147 | FRAGILARIA | 1505 | CHROOMONAS | 348 |
| CHROOMONAS | 1227 | MELOSIRA | 415 | CRYPTOMONAS | 248 |
| DIATOMA | 562 | NAVICULA | 208 | FRAGILARIA | 149 |
| NITZSCHIA | 460 | CYMBELLA | 156 | CYCLOTELLA | 99 |
| OTHER | 1279 | OTHER | 207 | OTHER | 100 |
| TOTAL | 10531 | TOTAL | 6175 | TOTAL | 1689 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS 58095. | 108130. | ***** | 443790. | 610015. |
| NITROGEN 271125. | 442190. | 55. | 7465330. | 8179700. |

B. OUTPUT

| OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|----------------------|----------------------|--|
| PHOSPHORUS 577365. | 5. | 2.20 |
| NITROGEN 12267495. | LOSS | 29.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SNAKE RIVER | 144.900 | 29292.9 | 0.075 | 1.184 | 11. | 185. |
| UNNAMED STREAM | 1.210 | 271.9 | 0.364 | 2.312 | 19. | 134. |
| DANIELSON CREEK | 1.700 | 253.3 | 0.042 | 1.660 | 9. | 356. |
| PORTNEUF RIVER | 15.800 | 3367.0 | 0.503 | 3.115 | 26. | 252. |
| BANNOCK CREEK | 1.090 | 1020.5 | 0.381 | 1.816 | 13. | 61. |
| SPRING CREEK | 13.580 | 69.9 | 0.042 | 1.333 | 130. | 8172. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------------|------------------------|------------------------|
| UNNAMED STREAM C1 | 0.209 | 1.715 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - CASCADE RESERVOIR (EUTROPHIC)
 COUNTY - VALLEY
 STORET NO. - 1602 WORKING PAPER NO. 777, NTIS ACCESSION NO. PB-272 231/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 1621.30 | 107.24 | 8.1 | 27.260 | 339.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|-----------------------------|--------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 18. | 32. | 2.2 | 0.032 | 0.009 | 0.060 | 0.320 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (0.6 - 1.2 (3)) | LIMITING NUTRIENT AT SAMPLING TIME (6/ 4/75) N | (8/ 1/75) N | (9/16/75) P AND N |
|------------------------------|---------------------------------------|--------------------------|---|-------------|-------------------|
| 8.1 | 0.6 - 1.2 (3) | | | | |

SUMMARY OF PHYTOPLANKTON DATA
 6/ 4/75 8/ 1/75 9/16/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------------|---------------|-------------|---------------|-------------|
| ASTERIONELLA | 5069 | FRAGILARIA | 1206 | APHANIZOMENON | 979 |
| DINOBYRON | 870 | APHANIZOMENON | 827 | CRYPTOMONAS | 367 |
| SYNEDRA | 605 | MELOSIRA | 276 | FRAGILARIA | 245 |
| CENTRIC DIATOM | 151 | CHROOMONAS | 138 | CHROOMONAS | 82 |
| MELOSIRA | 151 | APHANOTHECE | 103 | MELOSIRA | 82 |
| OTHER | 417 | OTHER | 277 | OTHER | 40 |
| TOTAL | 7263 | TOTAL | 2827 | TOTAL | 1795 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 1520. | ***** | ***** | 39025. | 40545. |
| NITROGEN | 4600. | ***** | 90. | 516985. | 521675. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 25985. | 36. | 0.38 |
| NITROGEN | 829205. | LOSS | 4.9 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| NORTH FORK PAYETTE RIVER | 10.190 | 440.3 | 0.033 | 0.527 | 20. | 389. |
| GOLD FORK RIVER | 5.910 | 388.5 | 0.057 | 0.437 | 26. | 216. |
| BOULDER CREEK | 0.950 | 99.7 | 0.076 | 0.597 | 22. | 205. |
| LAKE FORK CREEK | 3.350 | 198.4 | 0.022 | 0.363 | 31. | 210. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------|------------------------|------------------------|
| ROCK CREEK | 0.034 | 0.486 |
| MUD CREEK | 0.052 | 0.939 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - COEUR D'ALENE LAKE (MESOTROPHIC)
 COUNTY - BENEWAH, KOOTENAI
 STORET NO. - 1603 WORKING PAPER NO. 778, NTIS ACCESSION NO. PB-278 518/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 9945.60 | 129.50 | 2.3 | 179.380 | 19.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 23. | 43. | 3.0 | 0.017 | 0.005 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 10.4 | ***** | (4 / 4/75) P AND N (7/22/75) N (9 / 9/75) P |

SUMMARY OF PHYTOPLANKTON DATA

4/ 4/75 7/22/75 9/ 9/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|--------------|--------------|---------------|-------------|---------------|-------------|
| MELOSIRA | 9594 | MELOSIRA | 389 | APHANIZOMENON | 471 |
| ASTERIONELLA | 592 | APHANIZOMENON | 272 | TABELLARIA | 471 |
| SYNEDRA | 296 | FRAGILARIA | 272 | CRYPTOMONAS | 118 |
| FRAGILARIA | 85 | ANABAENA | 233 | MELOSIRA | 47 |
| CRYPTOMONAS | 42 | CHROOMONAS | 117 | SYNEDRA | 47 |
| OTHER | 42 | OTHER | 233 | OTHER | 95 |
| TOTAL | 10651 | TOTAL | 1516 | TOTAL | 1249 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 13580. | ***** | 250. | 165985. | 179815. |
| NITROGEN | 44875. | ***** | 9325. | 3342530. | 3396730. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 109525. | 39. | 1.39 |
| NITROGEN | 4354550. | LOSS | 26.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| WOLF LODGE CREEK | 1.480 | 153.1 | 0.018 | 0.580 | 5. | 162. |
| COEUR D'ALENE RIVER | 72.370 | 3833.2 | 0.049 | 0.638 | 26. | 389. |
| ST. JOE RIVER | 96.110 | 4480.7 | 0.020 | 0.447 | 13. | 331. |
| BENEWAH CREEK | 0.710 | 137.0 | 0.048 | 0.569 | 7. | 106. |
| PLUMMER CREEK | 0.310 | 112.4 | 0.054 | 1.180 | 4. | 109. |
| LAKE CREEK | 0.660 | 68.6 | 0.052 | 1.842 | 14. | 534. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------------|------------------------|------------------------|
| FERNAN LAKE OUTLET | 0.042 | 0.743 |
| ROCKFORD CREEK | 0.053 | 1.180 |
| MICA CREEK | 0.034 | 0.471 |

518

| | | |
|-----------------------|-------|-------|
| MURPHY CREEK | 0.170 | 7.690 |
| ROSE CREEK M1 | 0.192 | 3.307 |
| ROSE CREEK N1 | 0.241 | 3.151 |
| SOUTH FORK ROCK CREEK | 0.173 | 3.384 |
| COUGAR CREEK | 0.070 | 0.941 |
| COTTONWOOD CREEK | 0.492 | 4.316 |
| SQUAW CREEK | 0.060 | 1.036 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - DWORSHAK RESERVOIR (OLIGOTROPHIC)
 COUNTY - CLEARWATER
 STORET NO. - 1604 WORKING PAPER NO. 779. NTIS ACCESSION NO. PB-272 257/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 6319.60 | 76.89 | 55.5 | 196.780 | 305.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 18. | 18. | 2.5 | 0.010 | 0.009 | 0.080 | 0.260 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (4.2 - 0.9 (4)) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--------------------------|--|
| 2.4 | 0.2 - | 0.9 (4) | (4/7/75) P AND N (7/23/75) N (9/11/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA
 4/7/75 7/23/75 9/11/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|-------------|-------|----------------|-------|----------------|-------|
| MELOSIRA | 1036 | MALLOMONAS | 286 | COELOSPHAERIUM | 579 |
| CHROOMONAS | 192 | OOCYSTIS | 239 | MELOSIRA | 241 |
| CRYPTOMONAS | 154 | ANKISTRODESMUS | 95 | APHANIZOMENON | 96 |
| | | MELOSIRA | 48 | ANKISTRODESMUS | 48 |
| OTHER | 0 | OTHER | 0 | ASTERIONELLA | 48 |
| TOTAL | 1382 | TOTAL | 668 | OTHER | 97 |
| | | | | | |
| | | | | | |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 570. | ***** | ***** | 114745. | 115315. |
| NITROGEN | 1700. | ***** | ***** | 3246250. | 3247950. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 84025. | 27. | 1.50 |
| NITROGEN | 1892860. | 42. | 42.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| N FORK CLEARWATER RIVER | 106.920 | 3522.4 | 0.019 | 0.590 | 17. | 534. |
| ELK CREEK | 7.300 | 240.4 | 0.031 | 0.234 | 27. | 231. |
| REEDS CREEK | 4.900 | 161.6 | 0.036 | 0.303 | 33. | 262. |
| BREAKFAST CREEK | 11.000 | 334.1 | 0.017 | 0.417 | 17. | 569. |
| LITTLE N FK CLEARWATER R | 23.000 | 678.6 | 0.018 | 0.421 | 18. | 462. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-----------------|------------------------|------------------------|
| CRANBERRY CREEK | 0.031 | 0.643 |
| SWAMP CREEK | 0.029 | 0.532 |
| WEITAS CREEK | 0.033 | 0.393 |
| ISABELLA CREEK | 0.013 | 0.516 |

60

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - HAUSER LAKE
 COUNTY - KOOTENAI
 STORET NO. - 1605

EUTROPHIC

WORKING PAPER NO. 780, NTIS ACCESSION NO. PB-272 256/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 2.43 | 1.2 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 24. | 42. | 3.4 | 0.028 | 0.013 | 0.075 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALgal ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 11.1 | 0.6 - 11.6 (3) | (6/ 3/75) N (7/23/75) N (9/10/75) N (10/23/75) N |

| SUMMARY OF PHYTOPLANKTON DATA | | | | | | | |
|-------------------------------|---------|--------------|---------|---------------|--------|-------------|-------|
| | 6/ 3/75 | 7/23/75 | 9/10/75 | | | | |
| GENERA | COUNT | GENERA | GENERA | COUNT | GENERA | GENERA | COUNT |
| CRYPTOMONAS | 459 | CHROOMONAS | 260 | APHANIZOMENON | 2225 | CRYPTOMONAS | 217 |
| ANKISTRODESUS | 153 | TABELLARIA | 186 | MELOSIRA | 805 | CHROOMONAS | 186 |
| UNKNOWN CELLS | 92 | ASTERIONELLA | 74 | EUGLENA | 284 | MELOSIRA | 186 |
| NITZSCHIA | 61 | CRYPTOMONAS | 74 | CHLAMYDOMONAS | 189 | SCHROEDERIA | 186 |
| | | DINOBYRON | 74 | TABELLARIA | 189 | MALLOMONAS | 62 |
| OTHER | 0 | OTHER | 112 | OTHER | 568 | OTHER | 29 |
| TOTAL | 765 | TOTAL | 780 | TOTAL | 4260 | TOTAL | 866 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - HAYDEN LAKE (OLIGO-MESOTROPHIC)
 COUNTY - KOOTENAI
 STORET NO. - 1606 WORKING PAPER NO. 781, NTIS ACCESSION NO. PB-272 232/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 161.40 | 17.00 | 7.3 | 2.000 | 2.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 30. | 32. | 6.5 | 0.010 | 0.003 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD 2.8 | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|---------------|--|
| | | 0.2 - 1.5 (2) | (4/4/75) P (7/23/75) N (9/10/75) P |

SUMMARY OF PHYTOPLANKTON DATA
4/4/75 7/23/75 9/10/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------------|----------------|------------|----------------|------------|
| DINOBRYON | 510 | DINOBRYON | 333 | STEPHANODISCUS | 80 |
| MELOSIRA | 198 | CYCLOTELLA | 212 | CHROOMONAS | 32 |
| ASTERIONELLA | 170 | ANABAENA | 30 | APHANOTHECE | 16 |
| ANKISTRODESMUS | 113 | ANKISTRODESMUS | 30 | | |
| CYCLOTELLA | 113 | | | | |
| OTHER | 58 | OTHER | 0 | OTHER | 0 |
| TOTAL | 1162 | TOTAL | 605 | TOTAL | 128 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 180. | 1315. | 1495. |
| NITROGEN | ***** | ***** | 6785. | 55215. | 62000. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | ***** | **** | 0.09 |
| NITROGEN | ***** | **** | 3.6 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| HAYDEN CREEK | 0.910 | 73.8 | 0.016 | 0.479 | 6. | 190. |
| MOKINS CREEK | 0.250 | 20.2 | 0.030 | 0.811 | 11. | 436. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------------|------------------------|------------------------|
| YELLOWBANKS CREEK | 0.018 | 2.100 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - ISLAND PARK RESERVOIR (EUTROPHIC)
 COUNTY - FREMONT
 STORET NO. - 1607 WORKING PAPER NO. 782, NTIS ACCESSION NO. PB-278 522/A8

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 1245.80 | 31.54 | 5.0 | 13,560 | 109.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 71. | 120. | 2.8 | 0.034 | 0.012 | 0.050 | 0.320 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (3) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--------------|---|
| 9.3 | 3.6 - | 6.6 (3) | (7/30/75) N (9/3/75) N (10/20/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| 7/30/75 | 9/3/75 | 10/20/75 |
|---------|--------|----------|
|---------|--------|----------|

| | | | | | |
|------------------------|------------|---------------|-------------|--------------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| APHANIZOMENON | 454 | APHANIZOMENON | 3254 | FRAGILARIA | 596 |
| CHROOMONAS | 165 | CHROOMONAS | 38 | APHANIZOMENON | 553 |
| CRYPTOMONAS | 83 | MELOSIRA | 38 | MELOSIRA | 213 |
| CYANOPHYTON FILAMENTS | 83 | | | CHRYSOPHYTAN CELLS | 170 |
| ANACYSTIS(MICROCYSTIS) | 83 | | | CRYPTOMONAS | 85 |
| OTHER | 82 | OTHER | 0 | OTHER | 0 |
| TOTAL | 950 | TOTAL | 3330 | TOTAL | 1617 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 30. | 9975. | 10005. |
| NITROGEN | ***** | ***** | 1065. | 142730. | 143795. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 22550. | LOSS | 0.32 |
| NITROGEN | 459510. | LOSS | 4.6 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| HENRYS FORK | 9.140 | 676.0 | 0.022 | 0.585 | 9. | 78. |
| HOTEL CREEK | 0.520 | 38.3 | 0.041 | 0.589 | 18. | 233. |
| SHEEP CREEK | 0.080 | 32.4 | 0.056 | 0.781 | 4. | 61. |
| ICEHOUSE CREEK | 0.560 | 19.7 | 0.037 | 0.651 | 22. | 423. |
| SHERIDAN CREEK | 0.850 | 282.3 | 0.061 | 0.668 | 3. | 60. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - LAKE LOWELL (EUTROPHIC)
 COUNTY - CANYON
 STORET NO. - 1608 WORKING PAPER NO. 783, NTIS ACCESSION NO. PB-278 528/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT ***** | 39.80 | 5.9 | 10.160 | 262.0 | |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 90. | 199. | 0.6 | 0.070 | 0.015 | 0.070 | 0.690 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (4 / 8 / 75) N | LIMITING NUTRIENT AT SAMPLING TIME (8 / 1 / 75) N | (9 / 16 / 75) P AND N |
|------------------------------|---------------------------------------|---------------------------|--|-------------------------|
| 25.4 | 0.4 - 1.0 (2) | | | |

SUMMARY OF PHYTOPLANKTON DATA
4/ 8/75 8/ 1/75 9/16/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|--------------|----------------|-------------|----------------|-------------|
| FRAGILARIA | 9120 | APHANIZOMENON | 5622 | MELOSIRA | 5126 |
| ASTERIONELLA | 5267 | MELOSIRA | 861 | APHANIZOMENON | 267 |
| MELOSIRA | 636 | FRAGILARIA | 208 | ANKISTRODESMUS | 160 |
| STEPHANODISCUS | 600 | STEPHANODISCUS | 124 | COSMARIUM | 107 |
| SYNEDRA | 141 | CHROOMONAS | 84 | OOCYSTIS | 107 |
| OTHER | 390 | OTHER | 306 | OTHER | 107 |
| TOTAL | 16154 | TOTAL | 7205 | TOTAL | 5874 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 10. | 32090. | 32100. |
| NITROGEN | ***** | ***** | 340. | 282160. | 282500. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 18195. | 43. | 0.81 |
| NITROGEN | 304500. | LOSS | 7.1 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| UNNAMED CANAL C1 | 0.420 | ***** | 6.100 | 1.812 | ***** | ***** |
| NEW YORK CANAL | 9.740 | ***** | 0.072 | 0.761 | ***** | ***** |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------|------------------------|------------------------|
| UNNAMED CANAL E1 | 0.083 | 1.806 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - MAGIC RESERVOIR
 COUNTY - CAMAS, BLAINE
 STOCK NO. - 1609

WORKING PAPER NO. 785. NTIS ACCESSION NO. PB-278 553/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 4144.00 | 15.80 | 15.0 | 15.420 | 175.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 94. | 166. | 2.5 | 0.062 | 0.020 | 0.130 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (1.3 - 8.3 (2)) | LIMITING NUTRIENT AT SAMPLING TIME (5/15/75) P AND N (8/5/75) N (9/17/75) N |
|------------------------------|---------------------------------------|--------------------------|---|
| 7.3 | | | |

SUMMARY OF PHYTOPLANKTON DATA
 5/15/75 8/5/75 9/17/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------------|--------------|------------|---------------|-------------|
| STEPHANODISCUS | 4099 | CHROOMONAS | 132 | APHANIZOMENON | 3297 |
| CHROOMONAS | 592 | CRYPTOMONAS | 75 | CERATIUM | 230 |
| CRYPTOMONAS | 42 | FRAGILARIA | 75 | CHROOMONAS | 153 |
| | | ASTERIONELLA | 56 | MELOSIRA | 153 |
| | | ANABAENA | 38 | CRYPTOMONAS | 38 |
| OTHER | 0 | OTHER | 56 | OTHER | 0 |
| TOTAL | 4733 | TOTAL | 432 | TOTAL | 3871 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 8770. | ***** | 5. | 31975. | 40750. |
| NITROGEN | 32255. | ***** | 110. | 376630. | 408995. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 16845. | 59. | 2.58 |
| NITROGEN | 359260. | 12. | 25.9 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BIG WOOD RIVER | 9.290 | 2147.1 | 0.046 | 0.684 | 4. | 83. |
| CAMAS CREEK | 5.290 | 1678.3 | 0.077 | 1.384 | 11. | 83. |
| POISON CREEK | 0.040 | 18.7 | 0.092 | 0.830 | 49. | 597. |
| ROCK CREEK | 0.320 | 105.7 | 0.097 | 1.160 | 11. | 118. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------|------------------------|------------------------|
| LAVA CREEK | 0.053 | 0.577 |
| CAMP CREEK | 0.071 | 0.688 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - PALISADES RESERVOIR (MESOTROPHIC)
 COUNTY - BONNEVILLE, ID; LINCOLN, WY
 STORET NO. - 1610 WORKING PAPER NO. 786, NTIS ACCESSION NO. PB-272 236/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 13571.60 | 61.31 | 28.2 | 181.050 | 108.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 112. | 225. | 3.9 | 0.024 | 0.007 | 0.080 | 0.270 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|---------------|--|
| 2.1 | 0.3 - 1.8 (3) | (8/ 5/75) N | (9/18/75) P AND N (10/20/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA

| 8/ 5/75 | 9/18/75 | 10/20/75 |
|---------|---------|----------|
|---------|---------|----------|

| | | | | | |
|---------------|-------|---------------|-------|---------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 339 | APHANIZOMENON | 237 | APHANIZOMENON | 638 |
| ASTERIONELLA | 242 | CHROOMONAS | 142 | CHROOMONAS | 478 |
| CRYPTOMONAS | 97 | ASTERIONELLA | 47 | CRYPTOMONAS | 478 |
| CERATIUM | 48 | | | MELOSIRA | 319 |
| CHLAMYDOMONAS | 48 | | | ASTERIONELLA | 53 |
| OTHER | 98 | OTHER | 0 | OTHER | 53 |
| TOTAL | 872 | TOTAL | 426 | TOTAL | 2019 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 165. | ***** | 10. | 383070. | 383245. |
| NITROGEN | 4895. | ***** | 460. | 3585375. | 3590730. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 126270. | 67. | 6.25 |
| NITROGEN | 2920000. | 19. | 58.6 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SNAKE RIVER | 128.990 | 8984.7 | 0.041 | 0.594 | 29. | 261. |
| BEAR CREEK | 2.210 | 199.7 | 0.036 | 0.356 | 19. | 122. |
| BIG ELK CREEK | 1.960 | 153.3 | 0.027 | 0.426 | 11. | 172. |
| INDIAN CREEK | 0.390 | 100.5 | 0.037 | 0.494 | 10. | 131. |
| MCCUY CREEK | 2.310 | 279.7 | 0.017 | 0.289 | 4. | 77. |
| SALT RIVER | 21.420 | 2198.9 | 0.091 | 1.423 | 23. | 375. |
| GREYS RIVER | 18.470 | 1160.3 | 0.116 | 0.343 | 49. | 179. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------|------------------------|------------------------|
| LITTLE ELK CREEK | 0.020 | 0.259 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - PAYETTE LAKE (MESOTROPHIC)
 COUNTY - VALLEY
 STORET NO. - 1611 WORKING PAPER NO. 784, NTIS ACCESSION NO. PB-277 931/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 373.00 | 21.60 | 5.7 | 10.580 | 140.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 12. | 18. | 6.8 | 0.013 | 0.007 | 0.060 | 0.240 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD 0.2 - 0.2 (2) | LIMITING NUTRIENT AT SAMPLING TIME (6/ 4/75) P AND N (8/ 1/75) N (9/16/75) P |
|------------------------------|---------------------------------------|------------------------|---|
| 4.6 | | | |

SUMMARY OF PHYTOPLANKTON DATA
6/ 4/75 8/ 1/75 9/16/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|--------------|-------------|--------------|-------------|--------------|-------------|
| ASTERIONELLA | 1170 | ASTERIONELLA | 1505 | TABELLARIA | 629 |
| MELOSIRA | 1080 | MELOSIRA | 1254 | MELOSIRA | 449 |
| SYNEDRA | 225 | TABELLARIA | 502 | ASTERIONELLA | 180 |
| CHROOMONAS | 90 | CHROOMONAS | 351 | CRUCIGENIA | 45 |
| CRYPTOMONAS | 45 | SYNEDRA | 151 | | |
| OTHER | 0 | OTHER | 150 | OTHER | 0 |
| TOTAL | 2610 | TOTAL | 3913 | TOTAL | 1303 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 15. | 4080. | 4095. |
| NITROGEN | ***** | ***** | 535. | 197510. | 198045. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 4030. | 2. | 0.19 |
| NITROGEN | 118875. | 40. | 9.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| NORTH FORK PAYETTE RIVER | 7.580 | 255.9 | 0.013 | 0.413 | 11. | 508. |
| HORSE CREEK | 0.290 | 12.8 | 0.012 | 0.578 | 8. | 430. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------|------------------------|------------------------|
| ELIP CREEK | 0.012 | 0.282 |
| FALL CREEK | 0.015 | 0.504 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - LOWER TWIN LAKES (EUTROPHIC)
 COUNTY - KOOTENAI
 STORET NO. - 1612 WORKING PAPER NO. 787, NTIS ACCESSION NO. PB-278 554/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|-----------------------|----------------------|---------------------|--------------------|------------------------|
| IMPOUNDMENT | 133.40 | 16.35 | 2.3 | 2,140 | 1.8 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|--------------------------|-----------------------------|---------------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| 15. | 22. | 3.3 | 0.016 | 0.009 | 0.050 | 0.320 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|---------------------------|--|--|
| 2.3 | 0.4 - 0.9 (3) | (6/ 3/75) N (7/23/75) N (9/10/75) P AND N (10/24/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA

| | 6/ 3/75 | 7/23/75 | 9/10/75 | 10/24/75 | |
|--------------|-------------|--------------------|-------------|--------------|-------------|
| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
| DINOBYRON | 1214 | MELOSIRA | 875 | MERISMOPEDIA | 3081 |
| ASTERIONELLA | 187 | FRAGILARIA | 175 | CHROOMONAS | 171 |
| FLAGELLATES | 140 | ANKISTRODESMUS | 131 | MELOSIRA | 171 |
| ANABAENA | 47 | ASTERIONELLA | 87 | CRYPTOMONAS | 128 |
| NITZSCHIA | 47 | CHLOROPHYTAN CELLS | 87 | OOCYSTIS | 86 |
| OTHER | 45 | OTHER | 176 | OTHER | 129 |
| TOTAL | 1680 | TOTAL | 1531 | TOTAL | 3766 |
| | | | | | |
| | | | | | |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|--------------------------------|---------------------------------|-----------------------------------|--------------------------|-----------------------|
| PHOSPHORUS | ***** | ***** | 120. | 1790. | 1910. |
| NITROGEN | ***** | ***** | 4465. | 47180. | 51645. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|-------------------|-------------------|--|
| PHOSPHORUS | 235. | 88. | 0.12 |
| NITROGEN | 7610. | 85. | 3.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------|-----------------|-----------------------|---------------------|---------------------|------------------------------|------------------------------|
| SHOVE CREEK | 0.110 | 6.6 | 0.023 | 0.642 | 8. | 298. |
| FISH CREEK | 0.610 | 37.8 | 0.030 | 0.558 | 15. | 227. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------------|---------------------|---------------------|
| UNNAMED STREAM B1 | 0.037 | 0.570 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN IDAHO

NAME - UPPER TWIN LAKES
 COUNTY - KOOTENAI
 STORET NO. - 1613

WORKING PAPER NO. 787, NTIS ACCESSION NO. PB-278 554/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 133.40 | 16.35 | 2.3 | 2.140 | 1.8 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 14. | 20. | 3.3 | 0.017 | 0.004 | 0.045 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L-DRY WT) | YIELD (5.0 0.3 - 2.1 (3)) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|--------------------------------------|---------------------------------------|---|
| | | | (6/ 3/75) N (7/23/75) N (9/10/75) P (10/24/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA

| | 6/ 3/75 | 7/23/75 | 9/10/75 | 10/24/75 | |
|----------------|---------|----------------|---------|----------------|------|
| GENERA | COUNT | GENERA | COUNT | GENERA | |
| DINOBYRON | 788 | CRYPTOMONAS | 291 | MELOSIRA | 932 |
| CHROOMONAS | 338 | CHROOMONAS | 97 | FRAGILARIA | 883 |
| ANKISTRODESMUS | 113 | STICHOCOCCUS | 97 | EPITHEMIA | 343 |
| ASTERIONELLA | 113 | CHROOCOCCUS | 48 | LYNGBYA | 294 |
| CRYPTOMONAS | 113 | COELOSPHAERIUM | 48 | ANKISTRODESMUS | 196 |
| OTHER | 167 | OTHER | 0 | OTHER | 1769 |
| TOTAL | 1632 | TOTAL | 581 | TOTAL | 4417 |
| | | | | | 2735 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 120. | 1790. | 1910. |
| NITROGEN | ***** | ***** | 4465. | 47180. | 51645. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 235. | 88. | 0.12 |
| NITROGEN | 7610. | 85. | 3.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SHOVE CREEK | 0.110 | 6.6 | 0.023 | 0.642 | 8. | 298. |
| FISH CREEK | 0.610 | 37.8 | 0.030 | 0.558 | 15. | 227. |

69

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - CANYON FERRY RESERVOIR (EUTROPHIC)
 COUNTY - BROADWATER, LEWIS & CLARK
 STORET NO. - 3001 WORKING PAPER NO. 790. NTIS ACCESSION NO. PB-270 605/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 41191.30 | 142.45 | 17.4 | 155.493 | 186.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 127. | 235. | 1.4 | 0.047 | 0.029 | 0.170 | 0.440 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (5.2 - 7.4 (3) | LIMITING NUTRIENT AT SAMPLING TIME (5/28/75) N | (7/31/75) N | (9/ 4/75) P AND N (10/22/75) N |
|------------------------------|---------------------------------------|---------------------------|---|---------------|------------------------------------|
| 5.8 | | | | | |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/28/75 | 7/31/75 | 9/ 4/75 | 10/22/75 |
|------------|---------|---------------|---------|----------------|
| GENERA | COUNT | GENERA | COUNT | GENERA |
| CHROOMONAS | 411 | FRAGILARIA | 2667 | FRAGILARIA |
| MELOSIRA | 376 | APHANIZOMENON | 1731 | MELOSIRA |
| FRAGILARIA | 274 | NITZSCHIA | 187 | CHROOMONAS |
| DIATOMA | 205 | CHROOMONAS | 187 | ASTERIONELLA |
| CYMBELLA | 137 | SYNEDRA | 140 | STEPHANODISCUS |
| OTHER | 753 | OTHER | 94 | OTHER |
| TOTAL | 2156 | TOTAL | 5006 | TOTAL |
| | | | | |
| | | | | |
| | | | | |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 1305. | ***** | 105. | 327900. | 329310. |
| NITROGEN | 4485. | ***** | 3900. | 4178515. | 4186900. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 183030. | 44. | 2.31 |
| NITROGEN | 3857500. | 8. | 29.4 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| MISSOURI RIVER | 150.450 | 39738.4 | 0.078 | 0.870 | 8. | 100. |
| MAGPIE CREEK | 0.105 | 63.4 | 0.026 | 0.427 | 1. | 22. |
| BEAVER CREEK | 0.222 | 148.7 | 0.077 | 0.540 | 4. | 28. |
| CONFEDERATE CREEK | 0.440 | 149.2 | 0.030 | 0.710 | 2. | 66. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-----------------|------------------------|------------------------|
| AVALANCHE CREEK | 0.086 | 0.650 |
| WHITE CREEK | 0.050 | 0.330 |
| MANLEY DITCH | 0.240 | 0.715 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - CLARK CANYON RESERVOIR (EUTROPHIC)
 COUNTY - BEAVERHEAD
 STORET NO. - 3002 WORKING PAPER NO. 791, NTIS ACCESSION NO. PB-269 846/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 6011.40 | 19.97 | 15.8 | 10.180 | 1.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 264. | 385. | 2.6 | 0.049 | 0.027 | 0.160 | 0.550 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 2.4 | ***** | (5/28/75) N (7/31/75) N (9/ 3/75) N (10/20/75) N |

SUMMARY OF PHYTOPLANKTON DATA
 5/28/75 7/31/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|--------------|------------|--------------|------------|---------------|-------------|---------------|------------|
| CHROOMONAS | 175 | SCHROEDERIA | 371 | APHANIZOMENON | 1634 | APHANIZOMENON | 327 |
| CRYPTOMONAS | 145 | ANARAENA | 53 | CHROOMONAS | 218 | CHROOMONAS | 210 |
| FUNOTIA | 87 | CHROOMONAS | 53 | CRYPTOMONAS | 145 | CRYPTOMONAS | 117 |
| SCHROEDERIA | 58 | | | SCHROEDERIA | 36 | SCHROEDERIA | 23 |
| HANTZSCHIA | 58 | | | | | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 523 | TOTAL | 477 | TOTAL | 2033 | TOTAL | 677 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 5. | 19935. | 19940. |
| NITROGEN | ***** | ***** | 140. | 404850. | 404990. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 11775. | 41. | 1.00 |
| NITROGEN | 311590. | 23. | 20.3 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| RED ROCK RIVER | 6.650 | 4006.7 | 0.041 | 1.065 | 2. | 56. |
| HORSE PRAIRIE CREEK | 3.080 | 1320.9 | 0.075 | 1.150 | 6. | 85. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - FLATHEAD LAKE
COUNTY - FLATHEAD, LAKE
STORET NO. - 3003

WORKING PAPER NO. 792. NTIS ACCESSION NO. PB-271 466/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 18854.20 | 475.60 | 13.5 | 331.730 | 225.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|---------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 89. | 121. | 5.9 | 0.008 | 0.004 | 0.050 | 0.230 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L-DRY WT) (5/29/75) N (7/25/75) P AND N (9/8/75) P

SUMMARY OF PHYTOPLANKTON DATA

SUMMARY OF PATIENTS FOR LARRYON DATA 5/29/75 7/25/75 9/ 8/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|--------------|-------|----------------|-------|----------------|-------|
| ASTERIONELLA | 67 | DINOBRYON | 300 | CENTRIC DIATOM | 37 |
| TABELLARIA | 38 | CENTRIC DIATOM | 128 | CHROOMONAS | 37 |
| DINOBRYON | 29 | TABELLARIA | 43 | NITZSCHIA | 37 |
| CRYPTOMONAS | 10 | ASTERIONELLA | 43 | | |
| CHROOMONAS | 10 | OOCYSTIS | 43 | | |
| OTHER | 2 | OTHER | 0 | OTHER | 0 |

NUTRIENT LOADING CHARACTERISTICS/LAKE

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 26290. | ***** | 465. | 146700. | 173455. |
| NITROGEN | 73820. | ***** | 17540. | 6342010. | 6433370. |

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 114370. | 34. | 0.36 | |
| NITROGEN | 5510465. | 14. | 13.5 | |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT-SOURCE NUTRIENT EXPORT | STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------------------------|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| | FLATHEAD RIVER | 298.370 | 16058.0 | 0.015 | 0.582 | 7. | 337. |
| | DAYTON CREEK | 0.112 | 179.0 | 0.031 | 0.788 | 0.6 | 16. |
| | PROCTOR CREEK | 0.007 | 12.1 | 0.021 | 0.876 | 0.4 | 16. |
| | STONER CREEK | 0.338 | 58.5 | 0.022 | 0.638 | 4. | 116. |
| | SWAN RIVER | 32.880 | 1888.1 | 0.017 | 0.374 | 9. | 205. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------------|------------------------|------------------------|
| BOHANNON CREEK | 0.013 | 0.399 |
| FORREY CREEK | 0.018 | 0.689 |
| TEEPEE CREEK | 0.019 | 0.394 |
| HELLROARING CREEK | 0.014 | 0.377 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - GEORGETOWN LAKE (EUTROPHIC)
 COUNTY - DEER LODGE, GRANITE
 STORET NO. - 3004 WORKING PAPER NO. 793, NTIS ACCESSION NO. PB-270 600/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 137.30 | 11.20 | 3.4 | 1.297 | 342.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 101. | 162. | 3.4 | 0.022 | 0.011 | 0.040 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|------------------------------------|
| 7.0 | 3.6 | (7/29/75) N (9/4/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA
7/29/75 9/4/75

| GENERA | COUNT | GENERA | COUNT |
|--------------|-------------|------------------------|-------------|
| FRAGILARIA | 732 | CHROOMONAS | 641 |
| OOCYSTIS | 628 | FRAGILARIA | 529 |
| CRYPTOMONAS | 384 | OSCILLATORIA | 84 |
| FLAGELLATES | 384 | ANABAENA | 56 |
| TETRAEDRON | 314 | ANACYSTIS(MICROCYSTIS) | 56 |
| OTHER | 174 | OTHER | 110 |
| TOTAL | 2616 | TOTAL | 1476 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 55. | 725. | 780. |
| NITROGEN | ***** | ***** | 2125. | 33970. | 36095. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 1230. | LOSS | 0.07 |
| NITROGEN | 37545. | LOSS | 3.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| NORTH FORK FLINT CREEK | 0.144 | 37.3 | 0.021 | 0.863 | 3. | 105. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|----------------|------------------------|------------------------|
| BLODGETT GULCH | 0.097 | 0.823 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - HERGEN LAKE
 COUNTY - GALLATIN
 STORET NO. - 3005

WORKING PAPER NO. 794, NTIS ACCESSION NO. PB-270 604/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 2408.70 | 51.27 | 8.2 | 28.853 | 172.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 76. | 179. | 3.4 | 0.022 | 0.020 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|---|
| 4.1 | 1.7 | (7/30/75) N (9/3/75) N (10/20/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| 7/30/75 | 9/3/75 | 10/20/75 |
|---------|--------|----------|
|---------|--------|----------|

| | | | | | |
|---------------|------------|----------------|------------|------------------------|------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 171 | FRAGILARIA | 316 | MELOSIRA | 428 |
| FRAGILARIA | 128 | CHROOMONAS | 281 | ANACYSTIS(MICROCYSTIS) | 86 |
| APHANIZOMENON | 86 | ASTERIONELLA | 105 | SCHROEDERIA | 86 |
| ANABAENA | 86 | CRYPTOMONAS | 70 | CHROOMONAS | 86 |
| APHANOTHECE | 86 | STEPHANODISCUS | 35 | GOMPHOSPHAERIA | 43 |
| OTHER | 42 | OTHER | 105 | OTHER | 169 |
| TOTAL | 599 | TOTAL | 912 | TOTAL | 898 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 55. | 24120. | 24175. |
| NITROGEN | ***** | ***** | 1985. | 661015. | 663000. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 20550. | 15. | 0.47 |
| NITROGEN | 691505. | LOSS | 12.9 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| MADISON RIVER | 13.940 | 1157.7 | 0.023 | 0.774 | 13. | 211. |
| WATKINS CREEK | 0.098 | 29.5 | 0.009 | 0.338 | 1. | 35. |
| SOUTH FORK MADISON RIVER | 3.920 | 211.6 | 0.020 | 0.761 | 12. | 445. |
| DUCK CREEK | 1.350 | 305.4 | 0.033 | 0.948 | 5. | 132. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------|------------------------|------------------------|
| DENNY CREEK | 0.016 | 0.750 |
| COUGAR CREEK | 0.029 | 0.556 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - KOOCANUSA RESERVOIR (MESOTROPHIC)
COUNTY - LINCOLN, MT; BRITISH COLUMBIA, CAN.
STORED NO. - 3006 WORKING PAPER NO. 795, NTIS ACCESSION NO. PB-270 633/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 23271.10 | 188.18 | 38.4 | 308.610 | 271.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN (METERS) | SECCHI DISC TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------|-------------------------------|--------------------------|--------------------------|--------------------------|
| 96. | 173. | 4.1 | 0.045 | 0.044 | 0.100 | 0.300 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

• BIOLOGICAL CHARACTERISTICS (LAKE)
 MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L-DRY WT) (6/2/75) N (7/24/75) N (9/5/75) N

SUMMARY OF PHYTOPLANKTON DATA

| | 6/ 2/75 | 7/24/75 | 9/ 5/75 | | |
|----------------|---------|--------------|---------|---------------|-------|
| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
| CHROOMONAS | 339 | ASTERIONELLA | 355 | CHROOMONAS | 124 |
| PANDORINA | 135 | CHROOMONAS | 148 | ANABAENA | 62 |
| CRYPTOMONAS | 135 | CRYPTOMONAS | 89 | APHANIZOMENON | 31 |
| CENTRIC DIATOM | 135 | | | ASTERIONELLA | 31 |
| NITZSCHIA | 68 | | | | |
| OTHER | 35 | OTHER | 0 | OTHER | 0 |
| TOTAL | 847 | TOTAL | 592 | TOTAL | 248 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| 1835. | ***** | ***** | 24755. | 26590. |
| 9120. | ***** | ***** | 406850. | 415970. |

NITROGEN

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 228840. | LOSS | 1.22* |
| NITROGEN | 3725205. | LOSS | 19.86 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT-SOURCE NUTRIENT EXPORT STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BRISTOW CREEK | 0.440 | 66.8 | 0.017 | 0.439 | 3. | 76. |
| BIG CREEK | 4.310 | 361.8 | 0.015 | 0.279 | 7. | 81. |
| TOBACCO RIVER | 6.520 | 1139.6 | 0.040 | 0.743 | 11. | 146. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------------|------------------------|------------------------|
| PINKHAM CREEK | 0.024 | 0.659 |
| CRIPPLE HORSE CREEK | 0.017 | 0.567 |
| CANYON CREEK | 0.031 | 0.734 |
| JACKSON CREEK | 0.018 | 0.336 |
| BARRON CREEK | 0.015 | 0.378 |
| WARLAND CREEK | 0.037 | 0.657 |

MCGUIRE CREEK

0.014

0.312

* ESTIMATE BASED ON OUTLET LOADS.

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - MARY RONAN LAKE
 COUNTY - LAKE
 STORET NO. - 3007

WORKING PAPER NO. 796, NTIS ACCESSION NO. PB-270 635/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 76.80 | 6.15 | 8.5 | 0.110 | 17.4 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 68. | 103. | 3.3 | 0.020 | 0.006 | 0.040 | 0.520 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 4.7 | ***** | (6/ 2/75) N (7/28/75) N (9/ 5/75) N (10/22/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 6/ 2/75 | 7/28/75 | 9/ 5/75 | 10/22/75 |
|--|---------|---------|---------|----------|
|--|---------|---------|---------|----------|

| | | | | | | | |
|------------------------|-------------|---------------|------------|---------------|-------------|------------------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| DINOBRYON | 2787 | APHANIZOMENON | 245 | CHROOMONAS | 373 | APHANIZOMENON | 851 |
| CRYPTOMONAS | 190 | CRUCIGENIA | 163 | APHAOTHECE | 217 | CRYPTOMONAS | 675 |
| ANACYSTIS(MICROCYSTIS) | 127 | Cryptomonas | 163 | ANABAENA | 155 | ANACYSTIS(MICROCYSTIS) | 294 |
| APHANIZOMENON | 95 | CHROOMONAS | 163 | Cryptomonas | 155 | CHROOMONAS | 264 |
| FRAGILARIA | 95 | ANABAENA | 82 | APHANIZOMENON | 93 | FRAGILARIA | 175 |
| OTHER | 126 | OTHER | 124 | OTHER | 32 | OTHER | 441 |
| TOTAL | 3420 | TOTAL | 940 | TOTAL | 1025 | TOTAL | 2700 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 15. | 1105. | 1120. |
| NITROGEN | ***** | ***** | 535. | 16840. | 17375. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 490. | 56. | 0.18 |
| NITROGEN | 8520. | 51. | 2.8 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| FREELAND CREEK | 0.060 | 31.8 | 0.042 | 0.377 | 11. | 84. |
| DONALDSON CREEK | 0.046 | 5.7 | 0.025 | 0.502 | 54. | 835. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - LAKE MCDONALD (OLIGOTROPHIC)
COUNTY - FLATHEAD
STORE NO. - 3008 WORKING PAPER NO. 797. NTIS ACCESSION NO. PB-270 634/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 443.90 | 32.63 | 45.7 | 14,957 | 3.2 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 54. | 6. | 7.9 | 0.006 | 0.002 | 0.180 | 0.370 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (6/1/75) P (7/28/75) P

SUMMARY OF PHYTOPLANKTON DATA
6/ 1/75 7/28

| | COUNT | | COUNT |
|--------------|-----------|--------------|-----------|
| GENERAL | | GENERAL | |
| DINOBYRON | 56 | CYCLOTELLA | 57 |
| GYMNOCLADIA | 14 | | |
| OTHER | 0 | OTHER | 0 |
| TOTAL | 70 | TOTAL | 57 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NOTES

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 135. | ***** | 70. | 9295. | 9500. |
| NITROGEN | 400. | ***** | 2635. | 437040. | 440075. |

NITROGEN

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 5035. | 47. | | 0.29 |
| NITROGEN | 364395. | 17. | | 13.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT SOURCE NUTRIENT EXPORT STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| MCDONALD CREEK | 12.650 | 283.9 | 0.014 | 0.769 | 26. | 1223. |
| FISH CREEK | 0.700 | 39.6 | 0.015 | 0.806 | 11. | 473. |
| SNYDER CREEK | 0.472 | 13.8 | 0.024 | 0.736 | 21. | 814. |
| SPRAGUE CREEK | 0.165 | 16.5 | 0.013 | 0.691 | 4. | 241. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

MEAN NUTRIENT CONCENTRATIONS IN SELECTED STREAMS
 STREAM NAME MEAN TOTAL P MEAN TOTAL N
 (MG/L) (MG/L)

| | | |
|---------------|-------|-------|
| JACKSON CREEK | 0.020 | 0.722 |
| FERN CREEK | 0.014 | 0.867 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - NELSON RESERVOIR (EUTROPHIC)
 COUNTY - PHILLIPS
 STORET NO. - 3009 WORKING PAPER NO. 798, NTIS ACCESSION NO. PB-270 603/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT ***** | 18.45 | 5.7 | 2.290 | 1.8 | |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 175. | 562. | 1.1 | 0.029 | 0.007 | 0.075 | 0.555 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 7.2 | ***** | (5/30/75) N (7/24/75) P AND N (9/8/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/30/75 | 7/24/75 | 9/8/75 |
|--|---------|---------|--------|
|--|---------|---------|--------|

| | | | | | |
|----------------------|--------------|----------------------|--------------|-------------------------|---------------|
| GENERA CHROOMONAS | COUNT 347 | GENERA CHROOMONAS | COUNT 523 | GENERA APHANIZOMENON | COUNT 3800 |
| CENTRIC DIATOM | 261 | APHANIZOMENON | 486 | FLAGELLATES | 192 |
| CRYPTOMONAS | 87 | ANKISTRODESMUS | 75 | CHROOMONAS | 171 |
| ANKISTRODESMUS | 43 | | | OSCILLATORIA | 128 |
| NITZSCHIA | 43 | | | SCHROEDERIA | 107 |
| OTHER | 0 | OTHER | 0 | OTHER | 43 |
| TOTAL | 781 | TOTAL | 1084 | TOTAL | 4441 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 15. | 7620. | 7635. |
| NITROGEN | ***** | ***** | 535. | 154895. | 155430. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 1340. | 82. | 0.41 |
| NITROGEN | 81395. | 48. | 8.4 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| NELSON CANAL | 2.290 | ***** | 0.101 | 1.869 | ***** | ***** |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - SEELEY LAKE
 COUNTY - MISSOULA
 STORET NO. - 3010

WORKING PAPER NO. 799, NTIS ACCESSION NO. PB-270 636/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 378.40 | 3.49 | 18.3 | 5.881 | 127.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 47. | 60. | 3.5 | 0.015 | 0.010 | 0.040 | 0.225 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (*-----*) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 2.2 | (MG/L--DRY WT) | (5/28/75) N (7/29/75) N (9/4/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/28/75 | 7/29/75 | | 9/4/75 | |
|--------------|-------------|--------------|-------------|--------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| DINOBRYON | 826 | CHROOMONAS | 341 | DINOBRYON | 425 |
| MELOSIRA | 425 | FRAGILARIA | 289 | FRAGILARIA | 297 |
| CRYPTOMONAS | 142 | MELOSIRA | 210 | CRYPTOMONAS | 170 |
| FLAGELLATES | 142 | DINOBRYON | 105 | FLAGELLATES | 170 |
| FRAGILARIA | 71 | ASTERIONELLA | 26 | ANABAENA | 85 |
| OTHER | 93 | OTHER | 53 | OTHER | 382 |
| TOTAL | 1699 | TOTAL | 1024 | TOTAL | 1529 |

08

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 40. | 3340. | 3380. |
| NITROGEN | ***** | ***** | 1490. | 74575. | 76065. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 2105. | 38. | 0.97 |
| NITROGEN | 115975. | LOSS | 21.8 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| CLEARWATER RIVER | 4.000 | 273.2 | 0.014 | 0.356 | 8. | 178. |
| DEER CREEK | 0.763 | 51.3 | 0.018 | 0.504 | 14. | 255. |
| SEELEY CREEK | 0.111 | 13.2 | 0.015 | 0.427 | 4. | 138. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------|------------------------|------------------------|
| RICE CREEK | 0.013 | 0.528 |
| AUGGIE CREEK | 0.027 | 0.623 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - SWAN LAKE
 COUNTY - LAKE
 STORET NO. - 3011

(MESOTROPHIC)
 WORKING PAPER NO. 800, NTIS ACCESSION NO. PB-270 602/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 1737.90 | 10.85 | 21.9 | 34.338 | 85.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| ALKALINITY(MG/L) | MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|------------------|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 85. | 124. | | 5.5 | 0.010 | 0.004 | 0.050 | 0.230 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 3.3 | ***** | (6 / 2 / 75) N (7 / 28 / 75) P (9 / 5 / 75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 6/ 2 / 75 | 7/ 28 / 75 | 9/ 5 / 75 | | |
|--------------|------------|--------------|------------|--------------|------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| DINOBYRON | 545 | ASTERIONELLA | 290 | CHROOMONAS | 238 |
| GYMNODINIUM | 95 | CHROOMONAS | 290 | CRYPTOMONAS | 119 |
| CRYPTOMONAS | 47 | | | ASTERIONELLA | 119 |
| ASTERIONELLA | 47 | | | CYCLOTELLA | 40 |
| MELOSIRA | 47 | | | | |
| OTHER | 72 | OTHER | 0 | OTHER | 0 |
| TOTAL | 853 | TOTAL | 580 | TOTAL | 516 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 55. | 16910. | 16965. |
| NITROGEN | ***** | ***** | 1990. | 768200. | 770190. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 9885. | 42. | 1.56 |
| NITROGEN | 894800. | LOSS | 71.0 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SWAN RIVER | 32.310 | 1416.7 | 0.013 | 0.606 | 10. | 450. |
| JOHNSON CREEK | 0.183 | 31.3 | 0.012 | 0.572 | 2. | 104. |
| BOND CREEK | 0.477 | 19.6 | 0.012 | 0.582 | 9. | 491. |
| HALL CREEK | 0.358 | 12.1 | 0.012 | 0.592 | 9. | 555. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------|------------------------|------------------------|
| SIXMILE CREEK | 0.015 | 0.780 |
| GROOM CREEK | 0.017 | 0.795 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - TALLY LAKE
COUNTY - FLATHEAD
STORET NO. - 3012

WORKING PAPER NO. 801. NTIS ACCESSION NO. PB-269 843/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 527.10 | 5.37 | 68.9 | 4,250 | 2.8 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 94. | 109. | 4.1 | 0.011 | 0.004 | 0.050 | 0.240 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

• BIOLOGICAL CHARACTERISTICS (CONT.)
 MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (.MG/L--DRY WT)
 2.1 ***** (6 / 2/75) N (7/28/75) P (9 / 5/75) P

SUMMARY OF PHYTOPLANKTON DATA

| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
|----------------|-------|--------------|-------|----------------|-------|
| DINOBRYON | 322 | CHROOMONAS | 106 | FRAGILARIA | 1107 |
| SYNEDRA | 290 | OSCILLATORIA | 35 | DINOBRYON | 420 |
| FLAGELLATES | 258 | | | CHROOMONAS | 191 |
| ASTERIONELLA | 226 | | | CENTRIC DIATOM | 153 |
| STEPHANODISCUS | 193 | | | CRYPTOMONAS | 38 |
| OTHER | 129 | OTHER | 0 | OTHER | 38 |
| TOTAL | 1418 | TOTAL | 141 | TOTAL | 1947 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 2730. | 2730. |
| NITROGEN | ***** | ***** | 80. | 184530. | 184610. |

NITROGEN

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 2035. | 25. | 0.51 | |
| NITROGEN | 170390. | A. | 34.4 | |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| LOGAN CREEK | 3.620 | 437.2 | 0.018 | 1.215 | 5. | 343. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - TIBER RESERVOIR (MESO-EUTROPHIC)
 COUNTY - LIBERTY, TOOLE
 STORET NO. - 3013 WORKING PAPER NO. 802, NTIS ACCESSION NO. PB-270 601/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 12760.90 | 50.59 | 15.3 | 28.170 | 318.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 133. | 402. | 1.3 | 0.018 | 0.004 | 0.180 | 0.430 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 2.8 | 0.3 - 0.3 (2) | (5/30/75) P AND N (7/25/75) P (9/9/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA

| 5/30/75 | 7/25/75 | 9/9/75 | | | |
|----------------|------------|--------------|-------------|--------------|------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 417 | NITZSCHIA | 1390 | SYNEDRA | 214 |
| ANKISTRODESmus | 179 | FRAGILARIA | 348 | CHROOMONAS | 143 |
| SYNEDRA | 89 | CHROOMONAS | 139 | OSCILLATORIA | 107 |
| CRYPTOMONAS | 30 | SCHROEDERIA | 139 | | |
| TRACHELOMONAS | 30 | CRYPTOMONAS | 70 | | |
| OTHER | 0 | OTHER | 69 | OTHER | 0 |
| TOTAL | 745 | TOTAL | 2155 | TOTAL | 464 |

88
CC

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 4390. | ***** | ***** | 27735. | 32125. |
| NITROGEN | 18755. | ***** | 70. | 1730285. | 1749110. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 13855. | 57. | 0.64 |
| NITROGEN | 740560. | 58. | 34.6 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| MARIAS RIVER | 27.180 | 9665.9 | 0.017 | 1.427 | 2. | 167. |
| WILLOW CREEK | 0.383 | 2177.2 | 0.140 | 1.874 | 2. | 21. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - TONGUE RIVER RESERVOIR (EUTROPHIC)

COUNTY - BIG HORN

STORET NO. - 3014

WORKING PAPER NO. 803, NTIS ACCESSION NO. PB-270 646/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 4584.30 | 14.15 | 6.1 | 14.150 | 78.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 199. | 541. | 0.7 | 0.051 | 0.008 | 0.050 | 0.620 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 16.9 | ***** | (5/23/75) N (8/29/75) N (10/15/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA

5/23/75 8/29/75

10/15/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------------|---------------|-------------|----------------|-------------|
| CHROOMONAS | 2447 | EUGLENA | 946 | STEPHANODISCUS | 670 |
| EUGLENA | 1106 | GLENODINIUM | 908 | CRYPTOMONAS | 591 |
| CRYPTOMONAS | 804 | CHROOMONAS | 870 | FRAGILARIA | 552 |
| STEPHANODISCUS | 737 | STIPITOCOCCUS | 416 | CHROOMONAS | 197 |
| ASTERIONELLA | 168 | ASTERIONELLA | 378 | NAVICULA | 118 |
| OTHER | 269 | OTHER | 794 | OTHER | 354 |
| TOTAL | 5531 | TOTAL | 4312 | TOTAL | 2482 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 14775. | ***** | ***** | 118755. | 133530. |
| NITROGEN | 50200. | ***** | ***** | 610205. | 660405. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 16490. | 88. | 9.44 |
| NITROGEN | 618385. | 6. | 46.7 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| TONGUE RIVER | 14.150 | 3825.4 | 0.192 | 1.184 | 26. | 132. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|----------------|------------------------|------------------------|
| DEER CREEK | 0.063 | 1.399 |
| SQUIRREL CREEK | 0.127 | 1.719 |
| YOUNGS CREEK | 0.099 | 1.430 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN MONTANA

NAME - WHITEFISH LAKE (OLIGOTROPHIC)
COUNTY - FLATHEAD
STORET NO. - 3016 WORKING PAPER NO. 804, NTIS ACCESSION NO. PB-270 637/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 325-80 | 13.56 | 32.9 | 5,570 | 2.5 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS
 MEDIAN MEDIAN MEAN SECCHI DISC MEDIAN MEDIAN MEDIAN MEDIAN MEDIAN
 ALKALINITY(MG/L) CONDUCTIVITY(UMHOS) (METERS) TOTAL P(MG/L) ORTHO P(MG/L) INORG N(MG/L) TOTAL N(MG/L)
 87. 106. 5.3 0.008 0.003 0.040 0.220

III. BIOLOGICAL CHARACTERISTICS (LAKE)

• BIOLOGICAL CHARACTERISTICS (YEAR)
 MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L-DRY WT) (6/27/75) N (7/28/75) P (8/5/75) P

SUMMARY OF PHYTOPLANKTON DATA
6/ 2/75 7

| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
|----------------|-------|--------------|-------|---------------|-------|
| FRAGILARIA | 259 | FRAGILARIA | 524 | DINOBRYON | 314 |
| SYNEDRA | 216 | DINOBRYON | 489 | SPHAEROCYSTIS | 146 |
| STEPHANODISCUS | 86 | ASTERIONELLA | 105 | SCHROEDERIA | 35 |
| ASTERIONELLA | 86 | CRYPTOMONAS | 70 | | |
| CHROOMONAS | 86 | SCENEDESMUS | 70 | | |
| OTHER | 2 | OTHER | 174 | OTHER | 0 |
| TOTAL | 735 | TOTAL | 1432 | TOTAL | 486 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 55. | 5800. | 5855. |
| NITROGEN | ***** | ***** | 2145. | 157215. | 159360. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE |
|------------|----------------------|----------------------|----------------------------------|--------------|
| PHOSPHORUS | 2580. | 56. | | 0.43 |
| NITROGEN | 87710. | 45. | | 11.8 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT SOURCE NUTRIENT | EXPORT | MEAN FLOW | DRAINAGE AREA | MEAN TOTAL P | MEAN TOTAL N | TOTAL P EXPORT | TOTAL N EXPORT |
|---------------------------|--------|-----------|---------------|--------------|--------------|----------------|----------------|
| STREAM NAME | | (CMS) | (SQ KM) | (MG/L) | (MG/L) | (KG/SQ KM/YR) | (KG/SQ KM/YR) |
| LAZY CREEK | | 0.790 | 43.5 | 0.017 | 0.716 | 9. | 379. |
| SWIFT CREEK | | 3.480 | 198.1 | 0.027 | 0.816 | 21. | 483. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------------|------------------------|------------------------|
| BEAVER CREEK | 0.012 | 0.643 |
| HELL ROARING CREEK | 0.014 | 0.558 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEVADA

NAME - LAHONTAN RESERVOIR (EUTROPHIC)
COUNTY - CHURCHILL, LYON
STORED NO. - 3202 WORKING PAPER NO. 807. NTIS ACCESSION NO. PB-281 900/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 9375.80 | 19.69 | 6.7 | 16,960 | 91.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 86. | 141. | 0.7 | 0.198 | 0.148 | 0.350 | ***** |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (3/17/75) N (7/9/75) N (11/7/75) N

SUMMARY OF PHYTOPLANKTON DATA

| SUMMARY OF PHYTOPLANKTON DATA | | | | | |
|-------------------------------|-------------|--------------|-------------|----------------|-------------|
| | 3/17/75 | | 7/ 9/75 | | 11/ 7/75 |
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CENTRIC DIATOM | 684 | FRAGILARIA | 3646 | NITZSCHIA | 316 |
| NITZSCHIA | 650 | CHROOMONAS | 413 | CENTRIC DIATOM | 158 |
| CHROOMONAS | 308 | MELOSIRA | 413 | CHROOMONAS | 158 |
| CHLAMYDOMONAS | 171 | ASTERIONELLA | 69 | NAVICULA | 158 |
| DACTYLOCOCCOPSIS | 68 | SCHROEDERIA | 69 | MELOSIRA | 105 |
| OTHER | 103 | OTHER | 0 | OTHER | 264 |
| TOTAL | 1984 | TOTAL | 4610 | TOTAL | 1159 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENTS

| • INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 5. | 187620. | 187625. |
| NITROGEN | ***** | ***** | 210. | 854200. | 854410. |

NITROGEN & DILUTION

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 97485. | 48. | 9.53 |
| NITROGEN | 749841. | 12. | 43.4 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| CARSON RIVER | 12.410 | 4229.5 | 0.284 | 1.260 | 31. | 130. |
| TRUCKEE CANAL | 4.350 | ***** | 0.233 | 1.316 | ***** | ***** |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

MEAN NUTRIENT CONCENTRATIONS IN UNGAUL STREAMS
 STREAM NAME MEAN TOTAL P MEAN TOTAL N
 (MG/L) (MG/L)
 UNNAMED CREEK C1 0.438 2.500

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEVADA

NAME - LAKE MEAD (OLIGO-MESOTROPHIC)
 COUNTY - CLARK, NV; MOHAVE, AZ
 STORE NO. - 3201 WORKING PAPER NO. 808, NTIS ACCESSION NO. PB-

/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 434601.80 | 592.88 | 59.1 | 377.340 | 3.5 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 136. | 815. | 5.9 | 0.016 | 0.005 | 0.340 | 0.550 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|-------|--|
| 3.1 | 0.1 - 0.7 (9) | | (2/24/75) P (6/11/75) P (11/20/75) P (12/11/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 2/24/75 | 6/11/75 | | 11/20/75 | | 12/ 1/75 | |
|-------------|---------|----------------|-------|----------------|-------|--------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 511 | CHROOMONAS | 110 | RAPHIDIOPSIS | 1056 | CHROOMONAS | 170 |
| CRYPTOMONAS | 46 | RHOICOSPHEENIA | 44 | CHROOMONAS | 624 | RAPHIDIOPSIS | 43 |
| TETRAEDRON | 46 | | | ANABAENOPSIS | 240 | | |
| | | | | CENTRIC DIATOM | 240 | | |
| | | | | CRYPTOMONAS | 96 | | |
| OTHER | 0 | OTHER | 0 | OTHER | 96 | OTHER | 0 |
| TOTAL | 603 | TOTAL | 154 | TOTAL | 2352 | TOTAL | 213 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 322055. | ***** | 10. | 3370770. | 3692835. |
| NITROGEN | 961520. | ***** | 375. | 25918510. | 26880405. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 247325. | 93. | 6.23 |
| NITROGEN | 11996385. | 55. | 45.3 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| COLORADO RIVER | 370.210 | 378139.9 | 0.233 | 1.909 | 8. | 64. |
| MUDY RIVER | 0.400 | 21238.0 | 0.375 | 1.800 | 0.2 | 1. |
| VIRGIN RIVER | 4.820 | 15255.1 | 1.047 | 1.795 | 13. | 20. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEVADA

NAME - RYE PATCH RESERVOIR (EUTROPHIC)
 COUNTY - PERSHING
 STORET NO. - 3204 WORKING PAPER NO. 809, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 41699.00 | 46.14 | 4.6 | 7.170 | 1.2 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 349. | 736. | 0.8 | 0.095 | 0.039 | 0.050 | 0.620 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALgal ASSAY CONTROL (MG/L--DRY WT) | YIELD (6 / 2/75) N | LIMITING NUTRIENT AT SAMPLING TIME (7/11/75) N | (11/ 7/75) N |
|------------------------------|---------------------------------------|-------------------------|---|--------------|
| 4.9 | 0.5 - 6.2 (4) | | | |

SUMMARY OF PHYTOPLANKTON DATA
6/ 1/75 7/11/75 11/ 7/75

| GENERA CENTRIC DIATOM | COUNT | GENERA CRYPTOMONAS | COUNT | GENERA CHROOMONAS | COUNT |
|--------------------------|-------|-----------------------|-------|----------------------|-------|
| | 30 | NITZSCHIA | 34 | SCHROEDERIA | 889 |
| | | | 34 | CHLAMYDOMONAS | 727 |
| | | | | OOCYSTIS | 566 |
| | | | | PANDORINA | 242 |
| OTHER | 0 | OTHER | 0 | OTHER | 162 |
| TOTAL | 30 | TOTAL | 68 | TOTAL | 2748 |

88

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 54495. | 54495. |
| NITROGEN | ***** | ***** | ***** | 291585. | 291585. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE |
|------------|----------------------|----------------------|----------------------------------|--------------|
| PHOSPHORUS | 27770. | 49. | | 1.18 |
| NITROGEN | 201970. | 31. | | 6.3 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| HUMBOLDT RIVER | 6.610 | 40663.0 | 0.397 | 1.372 | 1. | 6. |
| PITT-TAYLOR DIV. CANAL | 0.450 | ***** | 0.485 | 1.378 | ***** | ***** |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------------------|------------------------|------------------------|
| PITT TAYLOR RES. OUTLET | 0.203 | 1.360 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEVADA

NAME - LAKE TAHOE (OLIGOTROPHIC)
 COUNTY - CARSON CITY, DOUGLAS, WASHOE, NV; EL DORADO, PLACER, CA
 STORET NO. - 3205 WORKING PAPER NO. 810, NTIS ACCESSION NO. PB-282 053/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 1307.90 | 501.82 | 307.0 | 17.850 | 477.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 46. | 76. | 12.8 | 0.005 | 0.003 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (0.1 - 0.2 (6) | LIMITING NUTRIENT AT SAMPLING TIME (3/18/75) N | (6/27/75) N | (11/ 4/75) P |
|------------------------------|---------------------------------------|---------------------------|---|---------------|----------------|
| 0.6 | 0.1 - 0.2 (6) | | | | |

SUMMARY OF PHYTOPLANKTON DATA

| 3/18/75 | 7/ 2/75 | 11/ 4/75 | | | |
|----------------|---------|-----------|-------|----------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| NITZSCHIA | 35 | NITZSCHIA | 302 | CENTRIC DIATOM | 6 |
| STEPHANODISCUS | 35 | | | CYMBELLA | 6 |
| OTHER | 0 | OTHER | 0 | GYMNODINIUM | 3 |
| TOTAL | 70 | TOTAL | 302 | OTHER | 0 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 20675. | 20675. |
| NITROGEN | ***** | ***** | ***** | 831030. | 831030. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 3440. | 83. | 0.04 |
| NITROGEN | 76385. | 91. | 1.7 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| UPPER TRUCKEE RIVER | 3.350 | 141.9 | 0.020 | 0.499 | 14. | 398. |
| WARD CREEK | 0.880 | 25.1 | 0.016 | 0.354 | 16. | 412. |
| THIRD CREEK | 0.230 | 15.5 | 0.029 | 0.434 | 16. | 200. |
| TROUT CREEK | 1.340 | 104.6 | 0.035 | 0.478 | 14. | 120. |
| TAYLOR CREEK | 1.510 | 46.9 | 0.015 | 0.547 | 13. | 474. |
| GENERAL CREEK | 0.750 | 19.9 | 0.017 | 0.453 | 19. | 530. |
| MCKINNEY CREEK | 0.500 | 13.2 | 0.012 | 0.398 | 12. | 500. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------------------|------------------------|------------------------|
| UNNAMED CREEK C2 | 0.010 | 0.507 |
| MILL CREEK F1 | 0.055 | 0.688 |
| SECRET HARBOR CREEK | 0.028 | 1.006 |
| SLAUGHTERHOUSE CANYON CR | 0.052 | 1.082 |

| | | |
|-----------------------|-------|-------|
| GRIFF CREEK | 0.024 | 0.534 |
| EDGEWOOD CREEK | 0.039 | 0.374 |
| HEAVENLY VALLEY CREEK | 0.024 | 0.980 |
| LONELY GULCH CREEK | 0.020 | 0.870 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEVADA

NAME - TOPAZ LAKE (EUTROPHIC)
 COUNTY - DOUGLAS, NV; MONO, CA
 STORET NO. - 3206 WORKING PAPER NO. 811, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 1080.00 | 9.71 | 15.8 | 6.810 | 264.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| ALKALINITY(MG/L) | MEDIAN 74. | MEDIAN 99. | MEAN SECCHI DISC (METERS) | MEDIAN 3.1 | MEDIAN TOTAL P(MG/L) 0.058 | MEDIAN ORTHO P(MG/L) 0.042 | MEDIAN INORG N(MG/L) 0.160 | MEDIAN TOTAL N(MG/L) 0.440 |
|------------------|---------------|---------------|------------------------------|---------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
|------------------|---------------|---------------|------------------------------|---------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) 7.5 | YIELD 6.4 - 8.3 (2) | LIMITING NUTRIENT AT SAMPLING TIME (3/19/75) N | (7/ 1/75) N | (11/ 5/75) N |
|------------------------------|--|------------------------|--|--------------|--------------|
|------------------------------|--|------------------------|--|--------------|--------------|

SUMMARY OF PHYTOPLANKTON DATA

| | 3/19/75 | 6/30/75 | 7/ 1/75 | 11/ 5/75 | |
|-----------------|---------|---------------|---------|---------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CENTRIC DIATOM | 198 | FRAGILARIA | 5059 | FRAGILARIA | 4766 |
| FRAGILARIA | 159 | ASTERIONELLA | 670 | APHANIZOMENON | 801 |
| DICTYOSPHAERIUM | 119 | CHROOMONAS | 335 | ASTERIONELLA | 521 |
| MELOSIRA | 119 | APHANIZOMENON | 274 | CHROOMONAS | 320 |
| CHROOMONAS | 79 | CRYPTOMONAS | 30 | MELOSIRA | 160 |
| OTHER | 39 | OTHER | 32 | OTHER | 80 |
| TOTAL | 713 | TOTAL | 6400 | TOTAL | 6648 |
| | | | | | 560 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 13180. | 13180. |
| NITROGEN | ***** | ***** | 10. | 125860. | 125870. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 6195. | 53. | 1.36 |
| NITROGEN | 195470. | LOSS | 13.0 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| UNNAMED CANAL R1 | 6.750 | 1048.9 | 0.060 | 0.506 | 12. | 109. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------|------------------------|------------------------|
| CALIFORNIA CREEK | 0.115 | 0.900 |
| SLINKARD CREEK | 0.123 | 0.715 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEVADA

NAME - UPPER PAHRANAGAT (EUTROPHIC)
COUNTY - LINCOLN
STORED NO. - 3207 WORKING PAPER NO. 812, NTIS ACCESSION NO. PB-282 471/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 6845.40 | 0.97 | 0.6 | 0.260 | 57.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 605. | 775. | 0.8 | 0.173 | 0.026 | 0.125 | 0.800 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

• BIOLOGICAL CHARACTERISTICS (CONT.)
 MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (5/6/75) N (8/21/75) N (11/21/75) N

SUMMARY OF PHYTOPLANKTON DATA

| | 5/ 6/75 | | 8/21/75 | | 11/21/75 |
|----------------|---------|-----------------|---------|-----------------|----------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 425 | MERISMOPEDIA | 37834 | PENNATE DIATOMS | 899 |
| SCHROEDERIA | 196 | EUGLENA | 946 | NAVICULA | 623 |
| CRYPTOMONAS | 163 | CYCLOTELLA | 883 | CHROOMONAS | 254 |
| ANKISTRODESmus | 65 | PENNATE DIATOMS | 568 | CYLINDROTHECA | 231 |
| QUADRIGULA | 65 | PHACUS | 252 | NITZSCMIA | 231 |
| OTHER | 34 | OTHER | 1324 | OTHER | 436 |
| TOTAL | 948 | TOTAL | 41807 | TOTAL | 2674 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT A. INPUT

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 1265. | 1265. |
| NITROGEN | ***** | ***** | ***** | 11295. | 11295. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 360. | 72. | | 1.30 |
| NITROGEN | 5525. | 51. | | 11.6 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| WHITE RIVER R1 | 0.220 | 6765.1 | 0.150 | 1.402 | 0.1 | 1. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEVADA

NAME - WALKER LAKE
 COUNTY - MINERAL
 STORET NO. - 3211

WORKING PAPER NO. 813, NTIS ACCESSION NO. PB-282 339/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 153.78 | 24.1 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 3760. | 7094. | 2.4 | 0.601 | 0.574 | 0.080 | 2.020 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 3.4 | ***** | (3/17/75) N (7/11/75) N (11/6/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/17/75 | 7/9/75 | | 11/6/75 | |
|-----------------|---------|-------------|-------|----------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| PENNATE DIATOMS | 27 | CHROOCOCCUS | 19 | NAVICULA | 6 |
| CHROOMONAS | 16 | | | | |
| TETRAEDRON | 5 | | | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 48 | TOTAL | 19 | TOTAL | 6 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

***** LAKE SAMPLING ONLY *****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEVADA

NAME - WASHOE LAKE (EUTROPHIC)
COUNTY - WASHOE
STORE NO. - 3208 WORKING PAPER NO. 814, NTIS ACCESSION NO. PB-282 432/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 215.00 | 16.59 | 1.5 | 0.870 | 5.7 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 359. | 445. | 0.0 | 0.403 | 0.268 | 0.130 | 1.340 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

• BIOLOGICAL CHARACTERISTICS (LAKE)
 MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (3/17/75) N (6/27/75) N (11/10/75) N

| | | | | |
|------|-----|--------------|--------------|--------------|
| 11.6 | 4.4 | (3/17/75) N | (6/27/75) N | (11/10/75) N |
|------|-----|--------------|--------------|--------------|

SUMMARY OF PHYTOPLANKTON DATA

| | 3/17/75 | COUNT | 6/27/75 | COUNT | 11/11/75 | COUNT |
|-------------|---------|-------|-------------|-------|------------|-------|
| GENERAL | | | GENERAL | | GENERAL | |
| MELOSIRA | 671 | | MELOSIRA | 859 | MELOSIRA | 1111 |
| CHROOMONAS | 79 | | CRYPTOMONAS | 136 | EPITHEMIA | 115 |
| CRYPTOMONAS | 79 | | EUGLENA | 136 | FRAGILARIA | 115 |
| RHOPALODIA | 79 | | RHOPALODIA | 136 | STAURONEIS | 77 |
| | | | FRAGILARIA | 90 | SURIRELLA | 38 |
| OTHER | 0 | | OTHER | 315 | OTHER | 229 |
| TOTAL | 908 | | TOTAL | 1672 | TOTAL | 1685 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 1735. | 1735. |
| NITROGEN | ***** | ***** | ***** | 34770. | 34770. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE |
|------------|----------------------|----------------------|----------------------------------|--------------|
| PHOSPHORUS | 1770. | LOSS | | 0.10 |
| NITROGEN | 8875. | 74. | | 2.1 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| OPHIR CREEK | 0.130 | 16.8 | 0.035 | 0.402 | 8. | 115. |
| FRANKTOWN CREEK | 0.240 | 39.6 | 0.046 | 0.617 | 8. | 66. |
| WINTERS CREEK | 0.020 | 4.7 | 0.034 | 0.555 | 4. | 79. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------|------------------------|------------------------|
| GALENA CREEK | 0.029 | 0.376 |
| JONES CREEK | 0.024 | 0.497 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEVADA

NAME - WILD HORSE RESERVOIR (EUTROPHIC)
 COUNTY - ELKO
 STORET NO. - 3209 WORKING PAPER NO. 815, NTIS ACCESSION NO. PB-282 427/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 541.30 | 12.14 | 7.3 | 1.210 | 2.4 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 76. | 100. | 0.3 | 0.114 | 0.065 | 0.320 | 0.920 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (7.7 - 15.4 (3) | LIMITING NUTRIENT AT SAMPLING TIME (5/30/75) N | (8 / 1/75) N | (11 / 6/75) N |
|------------------------------|---------------------------------------|-----------------------------|---|----------------|-----------------|
| 75.5 | | | | | |

SUMMARY OF PHYTOPLANKTON DATA
5/30/75 8/ 1/75 11/ 6/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------------|---------------|-------------|---------------|------------|
| CENTRIC DIATOM | 8958 | GLOEOTRICHIA | 541 | CHROOMONAS | 196 |
| FRAGILARIA | 444 | FRAGILARIA | 496 | APHANIZOMENON | 33 |
| CRYPTOMONAS | 121 | APHANIZOMENON | 270 | TRACHELOMONAS | 33 |
| PHORMIDIUM | 121 | APHANOTHECE | 90 | SCHROEDERIA | 33 |
| LYNGBYA | 81 | COCCONEIS | 90 | | |
| OTHER | 80 | OTHER | 225 | OTHER | 0 |
| TOTAL | 9805 | TOTAL | 1712 | TOTAL | 295 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 7985. | 7985. |
| NITROGEN | ***** | ***** | ***** | 133465. | 133465. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 10415. | LOSS | 0.66 |
| NITROGEN | 61640. | 54. | 11.0 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| OWYHEE RIVER | 0.540 | 145.0 | 0.157 | 1.317 | 17. | 256. |
| DEEP CREEK | 0.200 | 42.5 | 0.086 | 1.271 | 12. | 183. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------|------------------------|------------------------|
| PENROD CREEK | 0.087 | 1.000 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEVADA

NAME - WILSON RESERVOIR (EUTROPHIC)
 COUNTY - ELKO
 STORET NO. - 3210 WORKING PAPER NO. 816, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 240.90 | 3.35 | 3.9 | 0.270 | 5.1 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 72. | 96. | 7.7 | 0.049 | 0.016 | 0.120 | 0.520 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 10.0 | 1.6 - 10.0 (2) | (6/ 1/75) N (8/ 1/75) N (11/ 6/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| 5/31/75 | 8/ 1/75 | 11/ 6/75 | | | |
|-------------|---------|-------------|-------|-----------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CRYPTOMONAS | 1609 | FLAGELLATES | 210 | PENNATE DIATOMS | 446 |
| CHROOMONAS | 623 | GYMNODINIUM | 84 | FRAGILARIA | 377 |
| NITZSCHIA | 260 | TETRAEDRON | 42 | CHROOMONAS | 343 |
| SYNEDRA | 52 | SCENEDESMUS | 42 | APHANIZOMENON | 137 |
| OTHER | 0 | OTHER | 0 | EPITHHEMIA | 69 |
| TOTAL | 2544 | TOTAL | 378 | OTHER | 207 |
| | | | | | 1579 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 700. | 700. |
| NITROGEN | ***** | ***** | 10. | 10510. | 10520. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 700. | 0. | 0.21 |
| NITROGEN | 15490. | LOSS | 3.1 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| WILSON CREEK | 0.260 | 208.8 | 0.227 | 1.680 | 3. | 29. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|----------------|------------------------|------------------------|
| COLUMBIA CREEK | 0.063 | 0.600 |
| BULL CREEK | 0.096 | 0.924 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEW MEXICO

NAME - ALAMOGORDO RESERVOIR (EUTROPHIC)
 COUNTY - DE BACA, GUADALUPE
 STORET NO. - 3501 WORKING PAPER NO. 817, NTIS ACCESSION NO. PB-278 517/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 11370.10 | 18.49 | 8.1 | 2.770 | 1.7 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 108. | 1959. | 0.8 | 0.025 | 0.003 | 0.050 | 0.320 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 5.9 | 0.4 | (5/ 1/75) N (8/20/75) P (10/ 2/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/ 1/75 | 8/20/75 | | 10/ 2/75 | |
|----------------|---------|--------------|-------|------------------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 759 | RAPHIDIOPSIS | 395 | CHROOMONAS | 979 |
| ANKISTRODESMUS | 715 | NITZSCHIA | 395 | FLAGELLATES | 793 |
| RAPHIDIOPSIS | 223 | OSCILLATORIA | 346 | DACTYLOCOCOPSIS | 746 |
| CRYPTOMONAS | 40 | OOCYSTIS | 197 | ANACYSTIS(MICRUCYSTIS) | 746 |
| | | CRYPTOMONAS | 148 | MERISMOPEDIA | 670 |
| OTHER | 0 | OTHER | 840 | OTHER | 2828 |
| TOTAL | 1737 | TOTAL | 2321 | TOTAL | 6762 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 2625. | ***** | 20. | 25415. | 28060. |
| NITROGEN | 9920. | ***** | 800. | 106885. | 117605. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 3580. | 87. | 1.52 |
| NITROGEN | 68495. | 42. | 6.4 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| PECOS RIVER | 2.540 | 10282.3 | 0.292 | 1.067 | 2. | 8. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEW MEXICO

NAME - BLUEWATER LAKE (MESO-EUTROPHIC)
COUNTY - MCKINLEY, VALENCIA
STORE NO. - 3502 WORKING PAPER NO. 81B. NTIS ACCESSION NO. PB-278 497/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 520.60 | 7.07 | 6.7 | 0.260 | 5.8 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| PHYSICAL AND CHEMICAL CHARACTERISTICS | | | | | | | |
|---------------------------------------|-------------------------------|------------------|-------------|-------------------------|-------------------------|-------------------------|-------------------------|
| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN (METERS) | SECCHI DISC | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
| 126. | 349. | 0.5 | | 0.036 | 0.012 | 0.140 | 0.700 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

• BIOLOGICAL CHARACTERISTICS (LEAK)
 MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT)
 3.9 0.2 - 1.2 (?) (5/5/75) N (8/19/75) N (10/1/75) P

SUMMARY OF PHYTOPLANKTON DATA

| | 5/ 5/75 | 8/19/75 | | 10/ 1/75 | |
|---------------|---------|---------------|-------|---------------|-----|
| GENERAL | COUNT | GENERAL | COUNT | GENERAL | |
| FLAGELLATES | 570 | APHANIZOMENON | 1213 | CHROOMONAS | 432 |
| CYCLOTELLA | 342 | CHLAMYDOMONAS | 152 | APHANIZOMENON | 351 |
| CHROOMONAS | 171 | CHROOMONAS | 152 | CRYPTOMONAS | 54 |
| SCHROEDERIA | 171 | CRYPTOMONAS | 114 | | |
| CHLAMYDOMONAS | 114 | CYCLOTELLA | 76 | | |
| OTHER | 229 | OTHER | 74 | OTHER | 0 |
| TOTAL | 1597 | TOTAL | 1781 | TOTAL | 837 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NOVEMBER

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 590. | 590. |
| NITROGEN | ***** | ***** | 45. | 14620. | 14665. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 475. | 19. | | 0.08 |
| NITROGEN | 9350. | 36. | | 2.1 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT SOURCE NUTRIENT EXPORT | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BLUEWATER CREEK | 0.090 | 195.3 | 0.040 | 0.748 | 0.6 | 11. |
| COTTONWOOD CREEK | 0.090 | 167.8 | 0.092 | 0.995 | 1. | 15. |
| UNNAMED STREAM (C1) | 0.010 | 15.8 | 0.040 | 0.996 | 0.6 | 17. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEW MEXICO

NAME - CONCHAS RESERVOIR (MESOTROPHIC)
 COUNTY - SAN MIGUEL
 STORET NO. - 3503 WORKING PAPER NO. 819. NTIS ACCESSION NO. PB-278 540/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 19189.30 | 38.83 | 11.8 | 4.760 | 3.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 152. | 751. | 1.2 | 0.020 | 0.004 | 0.040 | 0.350 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (0.1 - 0.4 (3) | LIMITING NUTRIENT AT SAMPLING TIME (5 / 1 / 75) N | (8 / 21 / 75) N | (10 / 2 / 75) P |
|------------------------------|---------------------------------------|---------------------------|--|-------------------|-------------------|
| 3.3 | | | | | |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/ 1/75 | 8/21/75 | 10/ 2/75 | | |
|----------------|---------|----------------|----------|----------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 915 | CHROOMONAS | 402 | CHROOMONAS | 118 |
| ANKISTRODESMUS | 335 | OOCYSTIS | 183 | CRYPTOMONAS | 59 |
| OOCYSTIS | 134 | ANKISTRODESMUS | 110 | TRACHELOMONAS | 29 |
| ASTERIONELLA | 45 | OSCILLATORIA | 73 | SCENEDESMUS | 29 |
| | | CRYPTOMONAS | 37 | ANKISTRODESMUS | 29 |
| OTHER | 0 | OTHER | 72 | OTHER | 30 |
| TOTAL | 1429 | TOTAL | 877 | TOTAL | 294 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 40. | 11410. | 11450. |
| NITROGEN | ***** | ***** | 1520. | 179890. | 181410. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 690. | 94. | 0.29 |
| NITROGEN | 624795. | LOSS | 4.7 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| CANADIAN RIVER | 3.840 | 15578.8 | 0.084 | 0.847 | 0.6 | 7. |
| CONCHAS RIVER | 0.400 | 1354.6 | 0.010 | 0.807 | 0.1 | 12. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------|------------------------|------------------------|
| CONCHAS CANAL | 0.019 | 0.683 |
| TREMENTINA CREEK | 0.420 | 2.020 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEW MEXICO

NAME - EAGLE NEST LAKE (EUTROPHIC)
 COUNTY - COLFAX
 STORET NO. - 3504 WORKING PAPER NO. 820, NTIS ACCESSION NO. PB-278 548/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 432.50 | 9.82 | 9.9 | 0.580 | 5.3 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 149. | 275. | 1.1 | 0.181 | 0.132 | 0.070 | 0.520 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (2.0 - 14.0 (2)) | LIMITING NUTRIENT AT SAMPLING TIME (5/ 6/75) N | LIMITING NUTRIENT AT SAMPLING TIME (8/21/75) N | LIMITING NUTRIENT AT SAMPLING TIME (10/ 7/75) N |
|------------------------------|---------------------------------------|---------------------------|---|---|--|
| 13.4 | | | | | |

SUMMARY OF PHYTOPLANKTON DATA
5/ 6/75 8/21/75 10/ 7/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|-------------|-------|----------------|-------|---------------|-------|
| CRYPTOMONAS | 3042 | APHANIZOMENON | 389 | APHANIZOMENON | 3639 |
| CHROOMONAS | 1703 | CHROOMONAS | 130 | CHROOMONAS | 1178 |
| | | STEPHANODISCUS | 32 | OSCILLATORIA | 659 |
| | | CRYPTOMONAS | 32 | CRYPTOMONAS | 173 |
| | | SCHROEDERIA | 32 | SCHROEDERIA | 35 |
| OTHER | 0 | OTHER | 66 | OTHER | 0 |
| TOTAL | 4745 | TOTAL | 681 | TOTAL | 5684 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 2630. | 2630. |
| NITROGEN | ***** | ***** | 65. | 42210. | 42275. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 3120. | LOSS | 0.27 |
| NITROGEN | 32270. | 24. | 4.3 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| MORENO CREEK | 0.220 | 191.1 | 0.152 | 1.444 | 7. | 68. |
| SIXMILE CREEK | 0.070 | 27.2 | 0.077 | 1.219 | 7. | 110. |
| CIENEGUILA CREEK | 0.200 | 145.0 | 0.152 | 1.691 | 5. | 73. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEW MEXICO

NAME - ELEPHANT BUTTE RESERVOIR (EUTROPHIC)
COUNTY - SIERRA, SOCORRO
STORET NO. - 3505 WORKING PAPER NO. 821, NTIS ACCESSION NO. PB-278 498/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 76275.40 | 148.05 | 18.3 | 38.310 | 2.2 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 133. | 600. | 0.6 | 0.083 | 0.052 | 0.110 | 0.440 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (5/2/75) N (8/19/75) N (10/3/75) N

| | | | | |
|-----|----------------|------------|-------------|-------------|
| 6.8 | 2.4 - 10.6 (2) | (5/2/75) N | (8/19/75) N | (10/3/75) N |
|-----|----------------|------------|-------------|-------------|

SUMMARY OF PHYTOPLANKTON DATA

| | 5/ 2/75 | | 8/19/75 | | 10/ 3/75 | |
|----------------|---------|------------------------|---------|-----------------|----------|--|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT | |
| STEPHANODISCUS | 722 | STEPHANODISCUS | 437 | PENNATE DIATOMS | 41 | |
| CHROOMONAS | 611 | CARTERIA | 328 | CHROOMONAS | 21 | |
| COELASTRUM | 56 | CHROOMONAS | 273 | | | |
| ANKISTRODESMUS | 56 | ANACYSTIS(MICROCYSTIS) | 109 | | | |
| | | SCHROEDERIA | 109 | | | |
| OTHER | 0 | OTHER | 300 | OTHER | 0 | |
| TOTAL | 1445 | TOTAL | 1556 | TOTAL | 62 | |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 540360. | ***** | 5. | 2028905. | 2569270. |
| NITROGEN | 1440480. | ***** | 150. | 5804095. | 7244725. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 83595. | 97. | | 17.35 |
| NITROGEN | 1141900. | 84. | | 48.9 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT SOURCE NUTRIENT EXPORT STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--|-----------------|-----------------------|---------------------|---------------------|------------------------------|------------------------------|
| RIO GRANDE | 37.070 | 71742.9 | 1.902 | 5.690 | 28. | 78. |
| UNNAMED TRIBUTARY (C1) | 0.010 | 5.7 | 0.028 | 0.541 | 18. | 757. |
| UNNAMED STREAM (E1) | 0.010 | 41.3 | 0.013 | 0.744 | 0.1 | 6. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-----------------------|------------------------|------------------------|
| RIO GRANDE CONVEYANCE | 0.883 | 2.587 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEW MEXICO

NAME - EL VADO RESERVOIR (MESO-EUTROPHIC)
 COUNTY - RIO ARriba
 STORET NO. - 3506 WORKING PAPER NO. 822, NTIS ACCESSION NO. PB-278 500/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 2012.40 | 13.07 | 18.3 | 8.530 | 326.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 80. | 170. | 0.8 | 0.034 | 0.014 | 0.140 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (2.4 / 2) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--------------------|---|
| 2.2 | 0.3 - | 2.4 (2) | (5 / 5 / 75) P AND N (8 / 19 / 75) N (10 / 1 / 75) P AND N |

| SUMMARY OF PHYTOPLANKTON DATA | | | | | |
|-------------------------------|---------|---------------|-------|---------------|-------|
| | 5/ 5/75 | 8/19/75 | | 10/ 1/75 | |
| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
| CRYPTOMONAS | 154 | CHROOMONAS | 490 | APHANIZOMENON | 234 |
| CHROOMONAS | 92 | APHANIZOMENON | 31 | CHROOMONAS | 176 |
| CYCLOTELLA | 31 | | | CRYPTOMONAS | 79 |
| NITZSCHIA | 31 | | | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 308 | TOTAL | 521 | TOTAL | 489 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 1375. | ***** | 5. | 33540. | 34920. |
| NITROGEN | 3000. | ***** | 115. | 251875. | 254990. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 16285. | 53. | 2.67 |
| NITROGEN | 269615. | LOSS | 19.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| RIO CHAMA | 6.180 | 1243.2 | 0.108 | 0.968 | 19. | 153. |
| WILLOW CREEK | 1.940 | 499.9 | 0.057 | 0.929 | 7. | 60. |
| BOULDER CREEK | 0.130 | 188.8 | 2.430 | 4.077 | 25. | 72. |

102

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEW MEXICO

NAME - LAKE McMILLAN (EUTROPHIC)
 COUNTY - EDDY
 STORET NO. - 3507 WORKING PAPER NO. 823, NTIS ACCESSION NO. PB-278 495/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 44004.10 | 23.05 | 2.1 | 6.950 | 81.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 104. | 3998. | 0.3 | 0.097 | 0.009 | 0.045 | 0.470 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME (5/ 1/75) P AND N (8/20/75) N (10/ 2/75) N |
|------------------------------|---|--|
| 14.1 | 0.2 - 3.2 (2) | (5/ 1/75) P AND N (8/20/75) N (10/ 2/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| 5/ 1/75 | | | 8/20/75 | | | 10/ 2/75 | | |
|--------------|-------------|--------------|-------------|--------------|-------------|----------|-------|--|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT | GENERA | COUNT | |
| ACHNANTHES | 2733 | NITZSCHIA | 770 | NITZSCHIA | 2718 | | | |
| CYCLOTELLA | 1997 | OSCILLATORIA | 147 | OSCILLATORIA | 614 | | | |
| LUNATE CELLS | 1682 | EUGLENA | 147 | ANABAENA | 380 | | | |
| CHROOMONAS | 420 | CRYPTOMONAS | 37 | EUGLENA | 146 | | | |
| OOCYSTIS | 315 | CYCLOTELLA | 37 | BINUCLEARIA | 146 | | | |
| OTHER | 210 | OTHER | 36 | OTHER | 380 | | | |
| TOTAL | 7357 | TOTAL | 1174 | TOTAL | 4384 | | | |

103

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 13610. | ***** | ***** | 10625. | 24235. |
| NITROGEN | 40810. | ***** | ***** | 430120. | 470930. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 16860. | 30. | 1.05 |
| NITROGEN | 641260. | LOSS | 20.4 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| PECOS RIVER | 6.690 | 42597.2 | 0.123 | 1.681 | 0.2 | 8. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------------------|------------------------|------------------------|
| OVERFLOW OUTLET | 0.043 | 1.605 |
| PECOS RIVER (W CHANNEL) | 0.230 | 1.910 |
| UNNAMED CREEK (D1) | 0.022 | 5.920 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN NEW MEXICO

NAME - UTE RESERVOIR
 COUNTY - QUAY
 STOCK NO. - 3509

WORKING PAPER NO. 824, NTIS ACCESSION NO. PB-278 494/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 28852.60 | 16.79 | 8.0 | 3.020 | 1.4 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 274. | 843. | 1.3 | 0.021 | 0.004 | 0.040 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (0.2 - 0.2 (2)) | LIMITING NUTRIENT AT SAMPLING TIME (5/ 2/75) N | LIMITING NUTRIENT AT SAMPLING TIME (8/20/75) N | LIMITING NUTRIENT AT SAMPLING TIME (10/ 3/75) P AND N |
|------------------------------|---------------------------------------|--------------------------|---|---|--|
| 3.2 | | | | | |

SUMMARY OF PHYTOPLANKTON DATA
5/ 2/75 8/20/75 10/ 3/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|-------------|-------|------------------|-------|--------------|-------|
| CHROOMONAS | 490 | CHROOMONAS | 280 | CRYPTOMONAS | 142 |
| CRYPTOMONAS | 286 | SYNEDRA | 224 | SYNEDRA | 85 |
| FRAGILARIA | 82 | MERISMOPEDIA | 224 | OOCYSTIS | 57 |
| CYCLOTELLA | 41 | CRYPTOMONAS | 205 | MERISMOPEDIA | 57 |
| OTHER | 0 | DACTYLOCOCCOPSIS | 149 | CLOSTERIUM | 57 |
| TOTAL | 899 | OTHER | 298 | OTHER | 113 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS ***** | ***** | ***** | 11225. | 11225. |
| NITROGEN ***** | ***** | ***** | 61230. | 61230. |

B. OUTPUT

| OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|----------------------|----------------------|--|
| PHOSPHORUS 1095. | 90. | 0.67 |
| NITROGEN 184455. | LOSS | 3.6 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| ROMERO DRAW | 0.030 | 45.3 | 0.082 | 1.550 | 2. | 32. |
| PAJARITO CREEK | 0.250 | 1263.9 | 1.259 | 3.415 | 8. | 21. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN OREGON

NAME - BROWNLEE RESERVOIR (EUTROPHIC)
 COUNTY - BAKER, OR; WASHINGTON, ID
 STORET NO. - 4101 WORKING PAPER NO. 827. NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 188033.90 | 56.66 | 31.6 | 535.556 | 39.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 137. | 309. | 1.8 | 0.079 | 0.043 | 0.560 | 0.910 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|---|
| 16.2 | ***** | (4/ 9/75) P AND N (8/ 4/75) P AND N (9/15/75) N |

SUMMARY OF PHYTOPLANKTON DATA
4/ 9/75 8/ 4/75 9/15/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|--------------|--------------|-------------|---------------|-------------|
| STEPHANODISCUS | 9749 | FRAGILARIA | 1312 | MELOSIRA | 2705 |
| ASTERIONELLA | 603 | MELOSIRA | 150 | APHANIZOMENON | 312 |
| SURIRELLA | 235 | CHROOMONAS | 150 | SCHROEDERIA | 35 |
| NAVICULA | 168 | CRYPTOMONAS | 37 | NAVICULA | 35 |
| MELOSIRA | 101 | | | | |
| OTHER | 267 | OTHER | 0 | OTHER | 0 |
| TOTAL | 11123 | TOTAL | 1649 | TOTAL | 3087 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 4550. | 50570. | 990. | 1890725. | 1946835. |
| NITROGEN | 61170. | 160530500. | 440500401. | 30973275. | 632065346. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 1023210. | 47. | 34.36 |
| NITROGEN | 18218785. | 97. | 11155.4 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SNAKE RIVER | 507.920 | 180004.9 | 0.129 | 1.852 | 10. | 168. |
| EAGLE CREEK | 7.670 | 507.6 | 0.045 | 0.667 | 29. | 341. |
| POWDER RIVER | 6.980 | 3548.3 | 0.149 | 1.197 | 11. | 93. |
| ROCK CREEK | 0.075 | 42.2 | 0.079 | 0.918 | 5. | 54. |
| BURNT RIVER | 3.900 | 2830.9 | 0.157 | 1.131 | 9. | 54. |
| LITTLE ROCK CREEK | 0.041 | 31.9 | 0.096 | 1.063 | 4. | 46. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|----------------|------------------------|------------------------|
| BROWNLEE CREEK | 0.065 | 0.883 |
| DENNELL CREEK | 0.031 | 1.274 |
| TRAIL CREEK | 0.097 | 0.928 |

HENLEY CREEK

0.110

0.548

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN OREGON

NAME - DIAMOND LAKE (MESO-EUTROPHIC)
COUNTY - DOUGLAS
STORET NO. - 4102 WORKING PAPER NO. 828, NTIS ACCESSION NO. PB-281 809/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 13.00 | 6.9 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 22. | 26. | 5.2 | 0.028 | 0.011 | 0.040 | 0.370 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

SUMMARY OF PHYTOPLANKTON DATA
7/16/75 10/31/

| GENERA | COUNT | GENERA | COUNT |
|--------------|-------|----------------|-------|
| ASTERIONELLA | 2113 | STEPHANODISCUS | 402 |
| ANABAENA | 31 | CYCLOTELLA | 331 |
| CHROOMONAS | 31 | CHROOMONAS | 47 |
| | | CYMBELLA | 24 |
| | | EPITHEMIA | 24 |
| OTHER | 0 | OTHER | 22 |
| TOTAL | 2175 | TOTAL | 850 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

NITROGEN B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | **** | ***** | ***** |

***** LAKE SAMPLING ONLY *****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN OREGON

NAME - HELLS CANYON RESERVOIR (EUTROPHIC)
 COUNTY - BAKER, WALLA OR; ADAMS, IDAHO, ID
 STORET NO. - 4103 WORKING PAPER NO. 829. NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 189846.90 | 10.12 | 20.4 | 553.950 | 4.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 138. | 263. | 1.8 | 0.068 | 0.045 | 0.640 | 0.830 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 18.7 | ***** | (4/ 8/75) P (8/ 4/75) N (9/15/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 4/ 8/75 | 8/ 4/75 | 9/15/75 |
|----------------|---------|--------------|----------------|
| GENERA | COUNT | GENERA | GENERA |
| STEPHANODISCUS | 7332 | FRAGILARIA | CHROOMONAS |
| ASTERIONELLA | 296 | MELOSIRA | MELOSIRA |
| MELOSIRA | 148 | ASTERIONELLA | STEPHANODISCUS |
| FRAGILARIA | 111 | SKELETONEMA | CRYPTOMONAS |
| SURIRELLA | 111 | CRYPTOMONAS | SCHROEDERIA |
| OTHER | 334 | OTHER | OTHER |
| TOTAL | 8332 | TOTAL | TOTAL |
| | | 11588 | 1517 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 942285. | 942285. |
| NITROGEN | ***** | ***** | ***** | 25888790. | 25888790. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 1017465. | LOSS | 93.11 |
| NITROGEN | 21630255. | 16. | 2558.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SNAKE RIVER | 532.660 | 188810.9 | ***** | ***** | 5. | 134. |
| INDIAN CREEK | 1.940 | 102.8 | ***** | ***** | 39. | 670. |
| PINE CREEK | 10.860 | 595.7 | ***** | ***** | 51. | 616. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------|------------------------|------------------------|
| KINNEY CREEK | 0.042 | 0.654 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN OREGON

NAME - HILLS CREEK RESERVOIR (EUTROPHIC)
 COUNTY - LANE
 STORET NO. - 4104 WORKING PAPER NO. 830, NTIS ACCESSION NO. PB-281 857/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 1007.50 | 11.07 | 39.2 | 33.040 | 152.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 25. | 35. | 1.6 | 0.038 | 0.027 | 0.060 | 0.230 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 2.3 | ***** | (3/28/75) N (7/16/75) N (10/30/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/28/75 | 7/16/75 | | 10/30/75 | |
|-------------|---------|--------------|-------|----------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 219 | FRAGILARIA | 1569 | MELOSIRA | 4118 |
| CRYPTOMONAS | 17 | CRYPTOMONAS | 297 | ASTERIONELLA | 223 |
| | | ASTERIONELLA | 212 | FRAGILARIA | 148 |
| | | CHROOMONAS | 42 | CHLAMYDOMONAS | 111 |
| OTHER | 0 | OTHER | 0 | CENTRIC DIATOM | 111 |
| TOTAL | 236 | TOTAL | 2120 | TOTAL | 4786 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 40540. | 40540. |
| NITROGEN | ***** | ***** | 30. | 201335. | 201365. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 36130. | 11. | 3.66 |
| NITROGEN | 256375. | LOSS | 18.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| MIDDLE FK. WILLAMETTE R. | 22.370 | 668.2 | 0.040 | 0.172 | 43. | 179. |
| HILLS CREEK | 4.310 | 136.5 | 0.033 | 0.222 | 32. | 226. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-----------------|------------------------|------------------------|
| BUCK CREEK | 0.041 | 0.230 |
| COFFEEPOT CREEK | 0.050 | 0.274 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN OREGON

NAME - LAKE Owyhee
 COUNTY - MALHEUR
 STORET NO. - 4105

WORKING PAPER NO. 831, NTIS ACCESSION NO. PB-281 799/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 28904.40 | 56.25 | 24.6 | 27.180 | 1.7 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 83. | 150. | 0.5 | 0.095 | 0.064 | 0.425 | 0.780 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 3.4 | 8.2 | (4/ 8/75) N (8/ 1/75) N (9/16/75) N |

SUMMARY OF PHYTOPLANKTON DATA
 4/ 8/75 8/ 1/75 9/16/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|---------------|------------|----------------|------------|---------------|------------|
| CYCLOTELLA | 207 | COELOSPHAERIUM | 304 | APHANIZOMENON | 409 |
| CHROOMONAS | 207 | CRYPTOMONAS | 154 | CHROOMONAS | 204 |
| CHLAMYDOMONAS | 69 | APHANIZOMENON | 101 | | |
| PHORMIDIUM | 69 | PHORMIDIUM | 101 | | |
| CRYPTOMONAS | 34 | SCENEDESMUS | 51 | | |
| OTHER | 68 | OTHER | 0 | OTHER | 0 |
| TOTAL | 654 | TOTAL | 711 | TOTAL | 613 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 30. | 99555. | 99585. |
| NITROGEN | ***** | ***** | 1065. | 1098735. | 1099800. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 101020. | LOSS | 1.77 |
| NITROGEN | 1017855. | 7. | 19.6 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| OWYHEE RIVER | 25.520 | 20720.0 | 0.115 | 1.211 | 4. | 47. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------|------------------------|------------------------|
| DRY CREEK | 0.086 | 0.870 |
| BIRCH CREEK | 0.102 | 1.241 |
| CROOKED CREEK | 0.047 | 1.372 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN OREGON

NAME - OXBOW RESERVOIR (EUTROPHIC)
 COUNTY - BAKER, OR; ADAMS, ID
 STORET NO. - 4106 WORKING PAPER NO. 832, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 188810.90 | 5.66 | 11.4 | 532,660 | 1.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 131. | 265. | 1.9 | 0.071 | 0.040 | 0.690 | 0.910 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 10.3 | ***** | (4/ 9/75) P (8/ 4/75) N (9/15/75) N |

| SUMMARY OF PHYTOPLANKTON DATA | | | | | |
|-------------------------------|---------|----------------|-------|-------------|-------|
| 4/ 9/75 | 8/ 4/75 | 9/15/75 | | | |
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| STEPHANODISCUS | 5621 | FRAGILARIA | 433 | MELOSIRA | 807 |
| ASTERIONELLA | 300 | STEPHANODISCUS | 31 | CYCLOTELLA | 70 |
| MELOSIRA | 257 | | | CRYPTOMONAS | 35 |
| SURIRELLA | 43 | | | | |
| CHROOMONAS | 43 | | | | |
| OTHER | 44 | OTHER | 0 | OTHER | 0 |
| TOTAL | 6308 | TOTAL | 464 | TOTAL | 912 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 1038645. | 1038645. |
| NITROGEN | ***** | ***** | ***** | 18480225. | 18480225. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 895060. | 14. | 183.51 |
| NITROGEN | 25222530. | LOSS | 3265.1 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SNAKE RIVER | 526,270 | 188033.9 | 0.062 | 1.512 | 5. | 97. |
| WILDHORSE RIVER | 4,920 | 458.4 | 0.038 | 0.727 | 20. | 331. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN OREGON

NAME - SUTTLE LAKE (MESO-EUTROPHIC)
COUNTY - JEFFERSON
STORET NO. - 4107 WORKING PAPER NO. 833, NTIS ACCESSION NO. PB-281 870/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 57.50 | 1.09 | 9.9 | 1,447 | 87.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

P H Y S I C A L A N D C H E M I C A L C H A R A C T E R I S T I C S

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 27. | 47. | 10.3 | 0.031 | 0.020 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A (UG/L) ALGAL ASSAY CONTROL YIELD (MG/L-DRY WT) LIMITING NUTRIENT AT SAMPLING TIME (3/28/75) N (7/16/75) N (10/31/75) N

SUMMARY OF PHYTOPLANKTON DATA
3/28/75 7/

| | | | | | |
|------------------|-------|--------------|-------|----------------|-------|
| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
| MELOSIRA | 1374 | ANABAENA | 1263 | STEPHANODISCUS | 637 |
| DACTYLOCOCCOPSIS | 153 | ASTERIONELLA | 90 | FRAGILARIA | 495 |
| SYNEDRA | 38 | | | CHROOMONAS | 283 |
| | | | | CRYPTOMONAS | 106 |
| | | | | ASTERIONELLA | 35 |
| OTHER | 0 | OTHER | 0 | OTHER | 142 |
| TOTAL | 1565 | TOTAL | 1353 | TOTAL | 1698 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT A. INPUT

| • INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 10. | 1845. | 1855. |
| NITROGEN | ***** | ***** | 345. | 18820. | 19165. |

NITROGEN B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 1355. | 27. | | 1.70 |
| NITROGEN | 29455. | LOSS | | 17.6 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BLUE LAKE OUTLET | 1.350 | 46.9 | 0.034 | 0.364 | 32. | 313. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN OREGON

NAME - WALDO LAKE
 COUNTY - LANE
 STORET NO. - 4108

(OLIGOTROPHIC)

WORKING PAPER NO. 834, NTIS ACCESSION NO. PB-281 871/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 26.70 | 35.6 | ***** | 21.2 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|---------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 10. | 3. | 15.2 | 0.005 | 0.006 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|------------------------------------|
| 0.4 | 0.1 | (7/16/75) N (10/30/75) P |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|---|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |
| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) | | |
| PHOSPHORUS | ***** | **** | ***** | | |
| NITROGEN | ***** | **** | ***** | | |

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - BEAR LAKE (OLIGOTROPHIC)
COUNTY - RICH, UT; BEAR LAKE, ID
STORED NO. - 4901 WORKING PAPER NO. 836, NTIS ACCESSION NO. PB-279 105/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 1030.80 | 171.59 | 10.2 | 2,546 | 21.9 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| PHYSICAL AND CHEMICAL CHARACTERISTICS | | | | | | | |
|---------------------------------------|-------------------------------|------------------|-------------|-------------------------|-------------------------|-------------------------|-------------------------|
| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN (METERS) | SECCHI DISC | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
| 374. | 543. | 6.3 | | 0.011 | 0.003 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

• BIOLOGICAL CHARACTERISTICS (CONT'D)
 MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (5/14/75) N (8/6/75) P AND N (9/19/75) P AND N

SUMMARY OF PHYTOPLANKTON DATA

| | 5/14/ 0 | 8/ 6/ 0 | 9/19/ 0 | COUNT |
|----------------|---------|---------------|---------|-------|
| GENERAL | | | | |
| ANKISTRODESMUS | 1003 | LAGERHEIMIA | 95 | |
| CHROOMONAS | 56 | KIRCHNERIELLA | 95 | |
| | | CRYPTOMONAS | 48 | |
| | | OSCILLATORIA | 48 | |
| OTHER | 0 | OTHER | 0 | |
| TOTAL | 1059 | TOTAL | 286 | TOTAL |
| | | | | 632 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

Review

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 95. | 14645. | 14740. |
| NITROGEN | ***** | ***** | 3515. | 467505. | 471020. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 4925. | 67. | 0.09 |
| NITROGEN | 124190. | 74. | 2.7 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BIG CREEK | 0.530 | 246.0 | 0.084 | 1.213 | 6. | 82. |
| INDIAN CREEK | 0.006 | 11.7 | 0.321 | 3.530 | 5. | 57. |
| N. EDEN CANYON | 0.317 | 137.3 | 0.092 | 0.573 | 7. | 42. |
| UNNAMED CREEK E1 | 0.137 | 59.6 | 0.181 | 0.980 | 13. | 71. |
| SWAN CREEK | 0.510 | 10.1 | 0.027 | 0.394 | 43. | 627. |
| FISH HAVEN CREEK | 0.096 | 31.1 | 0.041 | 0.777 | 4. | 76. |
| LITTLE CREEK | 0.200 | 93.2 | 0.030 | 0.918 | 2. | 62. |
| PUMPED RETURN FLOW | 6.720 | ***** | 0.044 | 1.460 | ***** | ***** |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - DEER CREEK RESERVOIR (EUTROPHIC)
 COUNTY - WASATCH
 STORET NO. - 4903 WORKING PAPER NO. 837. NTIS ACCESSION NO. PB-281 899/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 1450.40 | 9.66 | 19.9 | 10.676 | 211.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 142. | 303. | 1.8 | 0.038 | 0.006 | 0.215 | 0.585 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 9.1 | 1.0 - 3.3 (2) | (5/12/75) P (8/11/75) P AND N (9/19/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/12/75 | 8/11/75 | 9/19/75 | | |
|----------------|---------|---------------|---------|-------|------|
| GENERA | COUNT | GENERA | COUNT | | |
| STEPHANODISCUS | 5047 | FRAGILARIA | 7571 | | |
| CHROOMONAS | 3426 | SPHAEROCYSTIS | 55 | | |
| FLAGELLATES | 2100 | | | | |
| CRYPTOMONAS | 700 | | | | |
| ASTERIONELLA | 74 | | | | |
| OTHER | 0 | OTHER | 0 | | |
| TOTAL | 11347 | TOTAL | 7626 | | |
| | | | | TOTAL | 3222 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 7360. | 4195. | ***** | 12295. | 23850. |
| NITROGEN | 29710. | 47000. | ***** | 234645. | 311355. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 15605. | 35. | 2.47 |
| NITROGEN | 242720. | 22. | 32.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| PROVO RIVER | 8.230 | 976.4 | 0.065 | 0.885 | 9. | 204. |
| DECKER CREEK | 0.011 | 4.7 | 0.059 | 0.324 | 6.* | 51.* |
| SNAKE CREEK | 1.300 | 82.9 | 0.090 | 1.282 | 7.* | 52.* |
| DANIELS CREEK | 0.100 | 126.9 | 0.255 | 1.242 | 6. | 35. |
| MAIN CREEK | 0.375 | 186.5 | 0.119 | 1.049 | 7. | 68. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------|------------------------|------------------------|
| SPRING CREEK | 0.300 | 1.492 |

* ESTIMATED

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - ECHO RESERVOIR (EUTROPHIC)
 COUNTY - SUMMIT
 STORET NO. - 4904 WORKING PAPER NO. 838, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 1895.90 | 5.95 | 15.3 | 7.687 | 134.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 172. | 333. | 1.3 | 0.047 | 0.012 | 0.170 | 0.460 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|-------|--|
| 7.0 | ***** | | (5/12/75) N (8/7/75) P AND N (9/18/75) N |

SUMMARY OF PHYTOPLANKTON DATA
5/12/75 8/7/75 9/18/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|--------------|----------------|------------|---------------|-------------|
| CHROOMONAS | 9336 | CHROOMONAS | 305 | FRAGILARIA | 918 |
| CENTRIC DIATOM | 1221 | CRYPTOMONAS | 153 | CHROOMONAS | 598 |
| FRAGILARIA | 1103 | FRAGILARIA | 61 | CRYPTOMONAS | 160 |
| CRYPTOMONAS | 906 | ASTERIUNELLA | 31 | APHANIZOMENON | 40 |
| ASTERIONELLA | 118 | ANKISTRODESmus | 31 | ASTERIONELLA | 40 |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 12684 | TOTAL | 581 | TOTAL | 1756 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 3785. | ***** | ***** | 11115. | 14900. |
| NITROGEN | 10495. | ***** | ***** | 306120. | 316615. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 10165. | 32. | 2.50 |
| NITROGEN | 178965. | 43. | 53.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| WEBER RIVER | 5.640 | 1134.4 | 0.045 | 1.211 | 7. | 190. |
| CHALK CREEK | 1.740 | 655.3 | 0.032 | 1.177 | 3. | 99. |
| UNNAMED CREEK D1 | 0.080 | 15.5 | 0.129 | 2.037 | 21. | 332. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - FISH LAKE (MESOTROPHIC--EUTROPHIC)
 COUNTY - SEVIER
 STORET NO. - 4906 WORKING PAPER NO. 839, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 57.00 | 10.12 | 25.9 | 0.143 | 58.5 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 63. | 87. | 8.8 | 0.023 | 0.004 | 0.040 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|--|------------------------------------|
| 12.5 | ***** | (8/12/75) N (9/25/75) N |

SUMMARY OF PHYTOPLANKTON DATA

8/12/75 9/25/75

| GENERA | COUNT | GENERA | COUNT |
|------------------------|-------------|---------------|-------------|
| FRAGILARIA | 2281 | CHLAMYDOMONAS | 1103 |
| ANACYSTIS(MICROCYSTIS) | 2082 | OOCYSTIS | 531 |
| CHROOCOCCUS | 760 | CHROMONAS | 286 |
| NAVICULA | 198 | KIRCHNERIELLA | 123 |
| DINOBYRON | 132 | LAGERHEIMIA | 123 |
| OTHER | 925 | OTHER | 203 |
| TOTAL | 6378 | TOTAL | 2369 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 40. | 455. | 495. |
| NITROGEN | ***** | ***** | 1455. | 13490. | 14945. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 210. | 58. | 0.05 |
| NITROGEN | 3925. | 74. | 1.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| JORGENSEN CREEK | 0.056 | 8.3 | 0.067 | 0.464 | 14. | 99. |
| BOWERY CREEK | 0.018 | 2.6 | 0.060 | 0.662 | 13. | 144. |
| TWIN CREEK | 0.009 | 1.3 | 0.057 | 0.631 | 12. | 138. |
| DOCTOR CREEK | 0.004 | 7.0 | 0.061 | 0.606 | 1. | 11. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|----------------|------------------------|------------------------|
| ANDERSON CREEK | 0.035 | 0.122 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - HUNTINGTON LAKE
 COUNTY - EMERY
 STORET NO. - 4907

WORKING PAPER NO. 840, NTIS ACCESSION NO. PB-281 873/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 0.93 | 5.2 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 161. | 394. | 2.7 | 0.013 | 0.005 | 0.040 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 1.9 | ***** | (5/13/75) N (8/12/75) N (9/24/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/13/75 | 8/12/75 | 9/24/75 | | |
|----------------|---------|-------------|---------|-------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| ANKISTRODESMUS | 182 | CYCLOTELLA | 47 | CRYPTOMONAS | 333 |
| CHROOMONAS | 182 | APHANOCAPSA | 39 | CHROOMONAS | 133 |
| CRYPTOMONAS | 91 | ACHNANTHES | 31 | MALLOMONAS | 33 |
| SYNEDRA | 45 | CRUCIGENIA | 31 | | |
| | | CRYPTOMONAS | 16 | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 500 | TOTAL | 164 | TOTAL | 499 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

118

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - JOES VALLEY RESERVOIR (MESOTROPHIC)
 COUNTY - EMERY
 STORET NO. - 4908 WORKING PAPER NO. 841, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 378.10 | 4.74 | 16.3 | 2,838 | 316.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 195. | 273. | 2.5 | 0.012 | 0.003 | 0.045 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 2.5 | 0.2 | (5/13/75) N (8/12/75) P (9/24/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/13/75 | 8/12/75 | 9/24/75 | |
|------------------------|---------|------------------------|------------------------|------|
| GENERA | COUNT | GENERA | GENERA | |
| FLAGELLATES | 1354 | CYCLOTELLA | ANACYSTIS(MICROCYSTIS) | |
| ANACYSTIS(MICROCYSTIS) | 1113 | DINOBYRON | FRAGILARIA | |
| CHROOMONAS | 242 | CHROOMONAS | DINOBYRON | |
| CARTERIA | 48 | ANACYSTIS(MICROCYSTIS) | CRYPTOMONAS | |
| CRYPTOMONAS | 48 | CRYPTOMONAS | | |
| OTHER | 0 | OTHER | OTHER | |
| TOTAL | 2805 | TOTAL | 2965 | |
| | | | TOTAL | 1179 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 3335. | 3335. |
| NITROGEN | ***** | ***** | ***** | 82995. | 82995. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|--|
| PHOSPHORUS | 2310. | 31. | 0.70 |
| NITROGEN | 87240. | LOSS | 17.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| LOWRY CREEK | 1.470 | 186.5 | 0.033 | 0.649 | 8. | 161. |
| LITTLES CREEK | 0.113 | 16.6 | 0.124 | 0.877 | 27. | 188. |
| SEELY CREEK | 1.200 | 93.2 | 0.026 | 1.132 | 11. | 460. |
| SWANSEY CREEK | 0.023 | 16.3 | 0.230 | 1.223 | 10. | 54. |
| NORTH DRAGON CREEK | 0.002 | 38.8 | 0.272 | 1.861 | 0.4 | 3. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - LOWER BOWNS RESERVOIR (MESOTROPHIC)
 COUNTY - GARFIELD
 STORET NO. - 4902 WORKING PAPER NO. 842, NTIS ACCESSION NO. PB-281 872/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | ***** | 0.36* | 0.1 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 70. | 120. | 4.2 | 0.031 | 0.006 | 0.040 | 0.530 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 5.6 | ***** | (5/9/75) P (8/13/75) N (9/25/75) N |

SUMMARY OF PHYTOPLANKTON DATA
5/9/75 8/13/75 9/25/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|-------------|-------|------------------------|-------|--------------|-------|
| CRYPTOMONAS | 286 | GLOEOTRICHIA | 231 | CHROOMONAS | 970 |
| CHROOMONAS | 24 | CHROOMONAS | 103 | APHANOCAPSA | 562 |
| | | ANABAENA | 51 | CRYPTOMONAS | 357 |
| | | ANACYSTIS(MICROCYSTIS) | 26 | STAURASTRUM | 153 |
| OTHER | 0 | OTHER | 0 | GLOEOTRICHIA | 102 |
| TOTAL | 310 | TOTAL | 411 | TOTAL | 2144 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | **** | ***** |
| NITROGEN | ***** | **** | ***** |

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - LYNN RESERVOIR (EUTROPHIC)
 COUNTY - BOX ELDER
 STORET NO. - 4905 WORKING PAPER NO. 843, NTIS ACCESSION NO. PB-282 343/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | ***** | ***** | ***** | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| ALKALINITY(MG/L) | MEDIAN 144. | MEDIAN 318. | MEAN SECCHI DISC (METERS) | MEDIAN 2.1 | MEDIAN TOTAL P(MG/L) 0.121 | MEDIAN ORTHO P(MG/L) 0.052 | MEDIAN INORG N(MG/L) 0.200 | MEDIAN TOTAL N(MG/L) 0.440 |
|------------------|----------------|----------------|------------------------------|---------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
|------------------|----------------|----------------|------------------------------|---------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) 39.6 | YIELD 12.4 | LIMITING NUTRIENT AT SAMPLING TIME (5/15/75) N | (8/ 5/75) N | (9/17/75) N |
|------------------------------|---|---------------|--|--------------|--------------|
|------------------------------|---|---------------|--|--------------|--------------|

SUMMARY OF PHYTOPLANKTON DATA
5/15/75 8/ 5/75 9/17/75

| | | | | | |
|---------------|-------|-------------|-------|---------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| NAVICULA | 239 | MELOSIRA | 386 | APHANIZOMENON | 7130 |
| PHORMIDIUM | 159 | SCHROEDERIA | 15 | SYNEDRA | 4045 |
| PANDORINA | 106 | | | DIATOMA | 548 |
| CRYPTOMONAS | 80 | | | ANABAENA | 548 |
| TRACHELOMONAS | 27 | | | ASTERIONELLA | 480 |
| OTHER | 132 | OTHER | 0 | OTHER | 2058 |
| TOTAL | 743 | TOTAL | 401 | TOTAL | 14809 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - MINERSVILLE RESERVOIR (EUTROPHIC)
 COUNTY - BEAVER
 STORET NO. - 4909 WORKING PAPER NO. 846. NTIS ACCESSION NO. PB-282 022/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 1320.90 | 4.01 | 5.6 | 1.037 | 266.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 284. | 550. | 1.4 | 0.192 | 0.107 | 0.060 | 1.070 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALgal ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 33.6 | 2.9 | (5/8/75) N (8/12/75) N (9/25/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/8/75 | 8/12/75 | 9/25/75 |
|--|--------|---------|---------|
|--|--------|---------|---------|

| | | | | | |
|------------------|---------------|----------------------|---------------|-------------------------|----------------|
| GENERA SYNURA | COUNT 3728 | GENERA PHORMIDIUM | COUNT 3118 | GENERA APHANIZOMENON | COUNT 11379 |
| STEPHANODISCUS | 270 | APHANIZOMENON | 432 | STEPHANODISCUS | 646 |
| ASTERIONELLA | 90 | GLOEOTRICHIA | 432 | TRACHELOMONAS | 50 |
| EUNOTIA | 45 | CHROOMONAS | 192 | | |
| CRYPTOMONAS | 45 | CRYPTOMONAS | 192 | | |
| OTHER | 45 | OTHER | 238 | OTHER | 0 |
| TOTAL. | 4223 | TOTAL | 4604 | TOTAL | 12075 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 1020. | 2435. | ***** | 2360. | 5815. |
| NITROGEN | 2400. | 23585. | 35. | 15970. | 41990. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 4450. | 23. | 1.45 |
| NITROGEN | 36375. | 13. | 10.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BEAVER RIVER | 0.910 | 704.5 | 0.188 | 1.253 | 3. | 14. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - MOON LAKE (OLIGOTROPHIC)
COUNTY - DUCHESNE
STORET NO. - 4910 WORKING PAPER NO. 847, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 290.10 | 3.12 | 14.1 | 3,826 | 140.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

MEAN SECCHI DISC
ALKALINITY(MG/L) CONDUCTIVITY(UMHOS) MEAN (METERS) TOTAL P(MG/L) ORTHO P(MG/L) INORG N(MG/L) TOTAL N(MG/L)

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L-DRY WT) (8/11/75) P (9/23/75) P

SUMMARY OF PHYTOPLANKTON DATA

| GENERA | COUNT | GENERA | COUNT |
|-----------------|-------|-----------------|-------|
| CHROOMONAS | 69 | CYCLOTELLA | 251 |
| ASTERIONELLA | 34 | PENNATE DIATOMS | 126 |
| PENNATE DIATOMS | 34 | ASTERIONELLA | 42 |
| OTHER | 0 | DINOBRYON | 42 |
| TOTAL | 137 | OTHER | 0 |
| | | TOTAL | 461 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 1630. | 1630. |
| NITROGEN | ***** | ***** | ***** | 145445. | 145445. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 1610. | 1. | 0.52 | |
| NITROGEN | 65495. | 55. | 46.6 | |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT SOURCE NUTRIENT EXPORT | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| LAKE FORK RIVER | 3.280 | 202.0 | 0.012 | 1.247 | 6. | 639. |
| FISH CREEK | 0.177 | 25.9 | 0.014 | 0.760 | 3. | 164. |
| BROWN DUCK CREEK | 0.264 | 36.3 | 0.025 | 0.761 | 6. | 175. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - NAVAJO LAKE
 COUNTY - KANE
 STORET NO. - 4911

(OLIGOTROPHIC)
 WORKING PAPER NO. 848. NTIS ACCESSION NO. PB-277 940/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 2.50 | 3.7 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| ALKALINITY(MG/L) | MEDIAN MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|------------------|---|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 67. | 99. | 3.3 | 0.016 | 0.003 | 0.040 | 0.470 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|------------------------------------|
| 2.0 | ***** | (8/13/75) N (9/25/75) P |

SUMMARY OF PHYTOPLANKTON DATA
8/13/75 9/25/75

| GENERA | COUNT | GENERA | COUNT |
|---------------|-------------|---------------|-------------|
| FLAGELLATES | 1204 | KIRCHNERIELLA | 437 |
| KIRCHNERIELLA | 262 | TETRAEDRON | 312 |
| CHROOMONAS | 105 | COSMARIUM | 156 |
| CRYPTOMONAS | 79 | OOCYSTIS | 156 |
| PERIDINIUM | 52 | ELAKATOTHRIX | 62 |
| OTHER | 104 | OTHER | 33 |
| TOTAL | 1806 | TOTAL | 1156 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

124

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - NEWCASTLE RESERVOIR (EUTROPHIC)
 COUNTY - IRON
 STORET NO. - 4912 WORKING PAPER NO. 849, NTIS ACCESSION NO. PB-281 874/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | ***** | 0.66 | 7.2 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 195. | 573. | 1.8 | 0.051 | 0.009 | 0.040 | 0.470 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 12.5 | ***** | (5/8/75) N (8/13/75) N (9/26/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/8/75 | 8/13/75 | 9/26/75 | |
|----------------|--------|--------------|---------------|------|
| GENERA | COUNT | GENERA | GENERA | |
| STEPHANODISCUS | 6117 | ELAKATOTHRIX | ASTERIONELLA | |
| NITZSCHIA | 1001 | CYCLOTELLA | OOCYSTIS | |
| CHROOMONAS | 278 | OOCYSTIS | FRAGILARIA | |
| GLENODINIUM | 56 | EUDORINA | SYNEDRA | |
| SCHROEDERIA | 56 | | APHANIZOMENON | |
| OTHER | 0 | OTHER | OTHER | |
| TOTAL | 7508 | TOTAL | 2134 | |
| | | | TOTAL | 8632 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

125

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - OTTER CREEK RESERVOIR (EUTROPHIC)

COUNTY - PIUTE

STORET NO. - 4913

WORKING PAPER NO. 850. NTIS ACCESSION NO. PB-277 939/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 984.20 | 10.20 | 6.3 | 1.613 | 1.4 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 187. | 365. | 1.2 | 0.067 | 0.033 | 0.040 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 11.8 | 2.5 - 4.2 (2) | (5/ 9/75) N (8/13/75) N (9/25/75) N |

SUMMARY OF PHYTOPLANKTON DATA

5/ 9/75 8/13/75

9/25/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|---------------|-------------|--------------|-------------|--------------|------------|
| CYCLOTELLA | 5323 | FRAGILARIA | 4919 | CRYPTOMONAS | 446 |
| FLAGELLATES | 1331 | ANABAENA | 642 | ANABAENA | 74 |
| CHROOMONAS | 1065 | CERATIUM | 214 | CERATIUM | 74 |
| CHLAMYDOMONAS | 887 | MELOSIRA | 214 | | |
| CRYPTOMONAS | 444 | CRYPTOMONAS | 53 | | |
| OTHER | 486 | OTHER | 106 | OTHER | 0 |
| TOTAL | 9536 | TOTAL | 6148 | TOTAL | 594 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 6415. | 6415. |
| NITROGEN | ***** | ***** | ***** | 91475. | 91475. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 4305. | 33. | 0.63 |
| NITROGEN | 54970. | 40. | 9.0 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| OTTER CREEK | 0.352 | 854.7 | 0.121 | 2.265 | 2. | 29. |
| EAST FORK CANAL | 1.230 | ***** | 0.123 | 1.369 | ***** | ***** |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - PANGUITCH LAKE
COUNTY - GARFIELD
STORET NO. - 4914

WORKING PAPER NO. 851, NTIS ACCESSION NO. PB-281 875/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 121.70 | 4.99 | 6.4 | 0.748 | 2.2 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

MEAN AND CHLORINE CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN (METERS) | SECCHI DISC | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------|-------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 98. | 172. | 1.9 | | 0.071 | 0.010 | 0.040 | 1.020 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (8/13/75) N (9/25/75) N

SUMMARY OF PHYTOPLANKTON DATA
8/13/75

| GENERA | COUNT | GENERA | COUNT |
|-------------------------|-------|---------------|-------|
| APHANIZOMENON | 3127 | APHANIZOMENON | 8979 |
| FRAGILARIA | 324 | | |
| ANACYSTIS (MICROCYSTIS) | 118 | | |
| SPHAEROCYSTIS | 88 | | |
| CRYPTOMONAS | 88 | | |
| OTHER | 90 | OTHER | 0 |
| TOTAL | 3835 | TOTAL | 8979 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NOVEMBER

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|-----------------------------|
| PHOSPHORUS | ***** | ***** | 20. | 1795. | 1815. |
| NITROGEN | ***** | ***** | 695. | 25160. | 25855. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 1350. | 26. | | 0.36 |
| NITROGEN | 22895. | 11. | | 5.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT SOURCE NUTRIENT EXPORT | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BLUE SPRING CREEK | 0.400 | 41.4 | 0.067 | 0.983 | 20. | 300. |
| IPSON CREEK | 0.208 | 28.5 | 0.098 | 0.933 | 23. | 215. |
| CLEAR CREEK | 0.092 | 31.1 | 0.050 | 0.285 | 5. | 27. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - PELICAN LAKE
 COUNTY - UNTAH
 STORET NO. - 4915

WORKING PAPER NO. 852. NTIS ACCESSION NO. PB-277 937/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 25.90 | 6.88 | 3.0 | 0.126 | 36.8 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 282. | 897. | 1.6 | 0.044 | 0.004 | 0.050 | 0.920 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 6.3 | 1.9 | (5/13/75) N (8/ 7/75) P (9/23/75) P |

| SUMMARY OF PHYTOPLANKTON DATA | | | | | |
|-------------------------------|-------------|--------------|-------------|------------------------|--------------|
| 5/13/75 | COUNT | GENERA | 8/ 7/75 | COUNT | GENERA |
| CHLAMYDOMONAS | 801 | TETRAEDRON | 376 | CHLOROPHYTAN CELLS | 70495 |
| FRAGILARIA | 616 | FRAGILARIA | 282 | ANACYSTIS(MICROCYSTIS) | 2287 |
| CHROOMONAS | 493 | CHROOMONAS | 235 | DACTYLOCOCOPSIS | 1334 |
| CRYPTOMONAS | 246 | CRYPTOMONAS | 188 | MERISMOPEDIA | 953 |
| SCENEDESMUS | 62 | OOCYSTIS | 141 | OSCILLATORIA | 667 |
| OTHER | 0 | OTHER | 377 | OTHER | 284 |
| TOTAL | 2218 | TOTAL | 1599 | TOTAL | 76020 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 915. | 915. |
| NITROGEN | ***** | ***** | ***** | 13820. | 13820. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 45. | 95. | 0.13 |
| NITROGEN | 700. | 95. | 2.0 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| UNNAMED CANAL C1 | 0.123 | ***** | 0.200 | 1.609 | ***** | ***** |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - PINEVIEW RESERVOIR (EUTROPHIC)
COUNTY - WEHRS
STORET NO. - 4916 WORKING PAPER NO. 853. NTIS ACCESSION NO. PB-277 936/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 802.90 | 10.12 | 13.4 | 6,343 | 248.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| PHYSICAL AND CHEMICAL CHARACTERISTICS | | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|---------------------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | 1.6 | 0.028 | 0.006 | 0.300 | 0.740 |
| 110. | 197. | | | | | |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (5/14/75) P (8/7/75) P (9/23/75) P

SUMMARY OF PHYTOPLANKTON DATA

| | 5/14/75 | COUNT | 8/ 7/75 | COUNT | 9/23/75 | COUNT |
|---------------|---------|-------|-------------|-------|----------------|-------|
| GENERA | | | GENERA | | GENERA | |
| CHROOMONAS | 2552 | | FLAGELLATES | 681 | FRAGILARIA | 332 |
| CRYPTOMONAS | 1411 | | CRYPTOMONAS | 306 | APHANIZOMENON | 166 |
| CHLAMYDOMONAS | 489 | | CHROOMONAS | 211 | CHROOMONAS | 166 |
| SCHROEDERIA | 326 | | CYSTS | 118 | CRYPTOMONAS | 165 |
| FLAGELLATES | 217 | | SCHROEDERIA | 118 | STEPHANODISCUS | 41 |
| OTHER | 217 | | OTHER | 69 | OTHER | 0 |
| TOTAL | 5212 | | TOTAL | 1503 | TOTAL | 870 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 25. | 7215. | 7240. |
| NITROGEN | ***** | ***** | 855. | 214310. | 215165. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 7775. | LOSS | | 0.72 |
| NITROGEN | 189145. | 12. | | 21.3 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| N&S BR., S. FK. OGDEN R. | 2.200 | 440.3 | 0.036 | 0.977 | 6. | 154. |
| MIDDLE FK. OGDEN RIVER | 0.589 | 85.5 | 0.020 | 0.650 | 4. | 141. |
| GERSTEN CREEK | 0.130 | 17.1 | 0.076 | 0.897 | 18. | 215. |
| N. FK. OGDEN RIVER | 1.000 | 160.6 | 0.024 | 1.243 | 5. | 244. |
| SPRING CREEK | 0.270 | 19.4 | 0.056 | 1.467 | 24. | 644. |
| UNNAMED CANAL H1 | 0.066 | ***** | 0.032 | 1.048 | ***** | ***** |
| GROUND WATER | 1.558 | ***** | ***** | ***** | ***** | ***** |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - PIUTE RESERVOIR (EUTROPHIC)
 COUNTY - PIUTE
 STORET NO. - 4917 WORKING PAPER NO. 854, NTIS ACCESSION NO. PB-281 880/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 6319.60 | 10.15 | 10.1 | 4.917 | 239.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 269. | 394. | 0.4 | 0.047 | 0.007 | 0.150 | 0.680 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|---|
| 25.3 | 0.2 | (5/ 9/75) P (8/13/75) P (9/24/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/ 9/75 | 8/13/75 | 9/24/75 | | |
|--------------|--------------|--------------|-------------|--------------|-------------|
| GENERA | COUNT | GENERA | COUNT | | |
| DIATOMA | 10007 | MALLOMONAS | 489 | CYCLOTELLA | 1635 |
| DINOBYRON | 1943 | CRYPTOMONAS | 383 | MALLOMONAS | 1372 |
| CYCLOTELLA | 426 | CHROOMONAS | 349 | CRYPTOMONAS | 739 |
| CRYPTOMONAS | 80 | OOCYSTIS | 70 | EUGLENA | 369 |
| GYMNODINIUM | 80 | STAURASTRUM | 70 | SCHROEDERIA | 369 |
| OTHER | 79 | OTHER | 105 | OTHER | 897 |
| TOTAL | 12615 | TOTAL | 1466 | TOTAL | 5381 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 20655. | 20655. |
| NITROGEN | ***** | ***** | ***** | 273815. | 273815. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 10855. | 47. | 2.03 |
| NITROGEN | 204370. | 25. | 27.0 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SEVIER RIVER | 2.860 | 3185.7 | 0.174 | 1.893 | 5. | 54. |
| E FORK SEVIER RIVER | 1.870 | 2952.6 | 0.070 | 1.405 | 1. | 28. |
| WEST CANAL | 0.036 | ***** | 0.061 | 1.245 | ***** | ***** |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - PORCUPINE RESERVOIR (EUTROPHIC)
 COUNTY - CACHE
 STORET NO. - 4918 WORKING PAPER NO. 855. NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | ***** | 0.77 | 20.1 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 163. | 238. | 1.5 | 0.025 | 0.011 | 0.110 | 0.410 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 7.9 | 2.0 | (5/14/75) P (8/ 6/75) N (9/23/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/14/75 | 8/ 6/75 | 9/23/75 |
|--|---------|---------|---------|
|--|---------|---------|---------|

| | | | | | |
|-------------|-------|-------------|-------|--------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 3196 | DINOBYRON | 1147 | CYCLOTELLA | 887 |
| DINOBYRON | 761 | CYCLOTELLA | 688 | MELOSIRA | 242 |
| CRYPTOMONAS | 203 | CRYPTOMONAS | 191 | DINOBYRON | 202 |
| | | OOCYSTIS | 76 | ASTERIONELLA | 161 |
| | | PERIDINIUM | 38 | CRYPTOMONAS | 121 |
| OTHER | 0 | OTHER | 40 | OTHER | 121 |
| TOTAL | 4160 | TOTAL | 2180 | TOTAL | 1734 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

131

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - PRUESS LAKE
 COUNTY - MILLARD
 STORET NO. - 4919

WORKING PAPER NO. 856. NTIS ACCESSION NO. PB-281 895/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 1.46 | 5.8 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 455. | 949. | 0.2 | 0.057 | 0.008 | 0.140 | 0.920 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 4.5 | ***** | (5/8/75) N (8/13/75) N (9/26/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/8/75 | 8/13/75 | 9/26/75 | | |
|------------------|--------|----------------|---------|---------------|-------|
| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
| DACTYLOCOCCOPSIS | 2732 | SCHROEDERIA | 154 | SCHROEDERIA | 72 |
| CHROOMONAS | 102 | SPHAEROCYSTIS | 61 | SPHAEROCYSTIS | 48 |
| | | STEPHANODISCUS | 31 | CRYPTOMONAS | 24 |
| | | | | APHANIZOMENON | 24 |
| | | | | EUGLENA | 24 |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 2834 | TOTAL | 246 | TOTAL | 192 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

***** LAKE SAMPLING ONLY *****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - SEVIER BRIDGE RESERVOIR (EUTROPHIC)
 COUNTY - JUAB, SANPETE
 STORET NO. - 4920 WORKING PAPER NO. 857. NTIS ACCESSION NO. PB-279 104/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 13260.80 | 44.52 | 6.5 | 5.923 | 1.6 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 369. | 1996. | 1.3 | 0.026 | 0.008 | 0.355 | 1.010 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 18.2 | 0.4 - 0.4 (2) | (5/12/75) P (8/12/75) P (9/24/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/12/ 0 | 8/12/ 0 | 9/24/ 0 | | |
|--------------|-------------|---------------|-------------|--------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| ELAKATOTHRIX | 1191 | CRUCIGENIA | 2309 | CYCLOTELLA | 937 |
| SYNEDRA | 496 | OOCYSTIS | 553 | NITZSCHIA | 386 |
| CHROOMONAS | 347 | CRYPTOMONAS | 553 | TETRASTRUM | 331 |
| CRYPTOMONAS | 248 | APHANIZOMENON | 520 | CRYPTOMONAS | 331 |
| DIATOMA | 149 | FRAGILARIA | 390 | SKELETONEMA | 331 |
| OTHER | 100 | OTHER | 1400 | OTHER | 1268 |
| TOTAL | 2531 | TOTAL | 5725 | TOTAL | 3584 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 2885. | ***** | ***** | 30535. | 33420. |
| NITROGEN | 8275. | ***** | ***** | 733870. | 742145. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 4035. | 88. | 0.75 |
| NITROGEN | 258715. | 65. | 16.7 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| SEVIER RIVER | 5.720 | 12742.8 | 0.167 | 3.690 | 2. | 52. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - STARVATION RESERVOIR (MESOTROPHIC)
 COUNTY - DUCHESNE
 STORET NO. - 4921 WORKING PAPER NO. 858. NTIS ACCESSION NO. PB-281 888/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 2745.40 | 11.17 | 19.9 | 5.834 | 1.8 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 290. | 523. | 2.7 | 0.016 | 0.004 | 0.040 | 0.320 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|---|
| 5.7 | ***** | (5/13/75) N (8/11/75) N (9/24/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/13/75 | 8/11/75 | 9/24/75 | |
|----------------|---------|-------------|---------|-----|
| GENERA | COUNT | GENERA | COUNT | |
| STEPHANODISCUS | 1179 | OOCYSTIS | 165 | |
| FRAGILARIA | 937 | CHROOMONAS | 110 | |
| ASTERIONELLA | 514 | DINOHRYON | 82 | |
| DIATOMA | 30 | CRYPTOMONAS | 55 | |
| OTHER | 0 | OTHFR | 0 | |
| TOTAL | 2660 | TOTAL | 412 | |
| | | | TOTAL | 376 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 30175. | 30175. |
| NITROGEN | ***** | ***** | ***** | 380310. | 380310. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 2770. | 91. | 2.70 |
| NITROGEN | 87865. | 77. | 34.0 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| STRAWBERRY RIVER | 4.110 | 2382.8 | 0.184 | 1.659 | 10. | 89. |
| DUCHESNE RIVER DIVERSION | 1.714 | ***** | 0.045 | 1.000 | ***** | ***** |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - STEINAKER RESERVOIR (MESOTROPHIC)
 COUNTY - UNTAH
 STORET NO. - 4922 WORKING PAPER NO. 859. NTIS ACCESSION NO. PB-281 876/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 54.40 | 3.32 | 14.0 | 0.804 | 2.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 100. | 238. | 4.7 | 0.011 | 0.005 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 1.8 | ***** | (5/13/75) N (8/7/75) P (9/22/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/13/75 | 8/7/75 | 9/22/75 |
|--|---------|--------|---------|
|--|---------|--------|---------|

| | | | | | |
|----------------|-------------|--------------|------------|--------------|------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 533 | APHANOCAPSA | 253 | APHANOCAPSA | 201 |
| CRYPTOMONAS | 267 | CHROOMONAS | 95 | CHROOMONAS | 134 |
| ANKISTRODESMUS | 200 | SCHROEDERIA | 32 | ASTERIONELLA | 34 |
| CYCLOTELLA | 33 | | | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 1033 | TOTAL | 380 | TOTAL | 369 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 365. | 365. |
| NITROGEN | ***** | ***** | ***** | 18235. | 18235. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION LOSS | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|------------------------------|---|
| PHOSPHORUS | 460. | 0. | 0.11 |
| NITROGEN | 18165. | 0. | 5.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| STEINAKER FEEDER CANAL | 0.754 | ***** | 0.012 | 0.585 | ***** | ***** |
| STEINAKER DITCH | 0.010 | ***** | 0.020 | 0.343 | ***** | ***** |
| STEINAKER DRAW | 0.013 | 17.1 | ***** | ***** | 0.3* | 12.* |
| SPRING CREEK | 0.013 | 16.6 | ***** | ***** | 0.3* | 12.* |

* ESTIMATED

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - TROPIC RESERVOIR (MESOTROPHIC)
 COUNTY - GARFIELD
 STORET NO. - 4923 WORKING PAPER NO. 860, NTIS ACCESSION NO. PB-281 012/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 217.60 | 0.73 | 3.0 | 0.463 | 56.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 170. | 218. | 1.9 | 0.021 | 0.006 | 0.050 | 0.420 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 7.2 | 0.2 | (5/ 8/75) P (9/25/75) N (8/14/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/ 8/75 | 8/14/75 | | 9/25/75 | |
|-------------------------|-------------|--|---------------|--|-------------------------|
| GENERAL OSCILLATORIA | COUNT 14 | GENERAL GLOEOTRICHIA CHROOMONAS ANARAENA STAURASTRUM | COUNT 4660 | GENERAL CHROOMONAS CRYPTOMONAS PEDIASTRUM | COUNT 80 40 20 |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 14 | TOTAL | 4856 | TOTAL | 140 |

136

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 295. | 295. |
| NITROGEN | ***** | ***** | ***** | 8940. | 8940. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 260. | 12. | 0.40 |
| NITROGEN | 10810. | LOSS | 12.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| EAST FORK SEVIER RIVER | 0.440 | 191.7 | 0.023 | 0.392 | 1. | 40. |
| BADGER CREEK | 0.007 | 14.0 | 0.016 | 0.566 | 0.4 | 11. |
| KING CREEK | 0.005 | 5.4 | 0.014 | 0.923 | 0.9* | 26.* |

* ESTIMATED

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - UTAH LAKE (HYPEREUTROPHIC)
COUNTY - UTAH
STORED NO. - 4924 WORKING PAPER NO. 861, NTIS ACCESSION NO. PB-277 932/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 6876.40 | 396.60 | 2.1 | 17.180 | 2.5 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 166. | 1118. | 0.2 | 0.132 | 0.012 | 0.320 | 1.160 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (5/12/75) P AND N (8/8/75) N (9/19/75) P AND N
 72.0 6.7 - 13.8 (2)

SUMMARY OF PHYTOPLANKTON DATA

| | 5/12/75 | COUNT | GENERA | 8/ 8/75 | COUNT | GENERA | 9/19/75 | COUNT |
|-------------|---------|-------|--------------|---------|-------|--------------|---------|-------|
| CHROOMONAS | 1618 | | ANABAENA | 9639 | | ANABAENA | | 6854 |
| CRYPTOMONAS | 917 | | MELOSIRA | 455 | | OSCILLATORIA | | 219 |
| SCHROEDERIA | 890 | | OSCILLATORIA | 342 | | CRYPTOMONAS | | 165 |
| EUGLENA | 405 | | CHROOMONAS | 304 | | GLENODINIUM | | 137 |
| PTEROMONAS | 216 | | CRYPTOMONAS | 152 | | SCENEDESMUS | | 82 |
| OTHER | 998 | | OTHER | 151 | | OTHER | | 275 |
| TOTAL | 5044 | | TOTAL | 11043 | | TOTAL | | 7732 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 125700. | 7645. | ***** | 68550. | 201895. |
| NITROGEN | 523420. | 460570. | ***** | 1333575. | 2317565. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 139995. | 31. | | 0.51 |
| NITROGEN | 817185. | 65. | | 5.8 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT SOURCE NUTRIENT EXPORT STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BENJAMIN SLOUGH | 1.320 | 326.3 | 0.339 | 2.888 | 22. | 297. |
| UNNAMED CANAL E1 | 0.132 | 12.4 | 0.191 | 2.218 | 64. | 745. |
| SPANISH FORK | 2.075 | 1813.0 | 0.171 | 1.362 | 6. | 49. |
| DRY CREEK | 0.420 | 98.4 | 0.745 | 5.690 | 15.* | 350.* |
| HOBBLE CREEK | 0.617 | 297.8 | 0.054 | 1.529 | 4. | 100. |
| SPRING CREEK K1 | 0.250 | 24.1 | 1.607 | 6.125 | 38.* | 851.* |
| MILL RACE CREEK | 1.240 | 44.0 | 1.354 | 5.773 | 303. | 779. |
| PROVO RIVER | 5.296 | 1787.1 | 0.053 | 0.936 | 3. | 74. |
| UNNAMED STREAM N1 | 0.330 | 3.4 | 1.727 | 4.796 | 351.* | 7962.* |
| UNNAMED STREAM P1 | 0.970 | 24.6 | 0.064 | 1.467 | 143.* | 3234.* |
| AMERICAN FORK CREEK | 0.038 | 163.2 | 0.024 | 1.462 | 0.2 | 11. |
| PLEASANT GROVE WST DITCH | 0.750 | 98.4 | 0.329 | 3.362 | 23. | 540. |
| UNNAMED STREAM Y1 | 0.111 | 3.6 | 0.113 | 3.475 | 110. | 3379. |

| | | | | | | |
|-------------------|-------|-------|-------|-------|-------|--------|
| SPRING CREEK 41 | 0.520 | 18.1 | 0.045 | 1.958 | 104.* | 2357.* |
| UNNAMED STREAM 51 | 0.023 | 1.0 | 0.115 | 2.601 | 85. | 1885. |
| UNNAMED STREAM 61 | 0.010 | 1.8 | 0.204 | 2.980 | 36. | 522. |
| UNNAMED STREAM 71 | 0.210 | 1.0 | 0.031 | 2.329 | 205. | 15425. |
| CURRENT CREEK 81 | 0.298 | 999.7 | 0.088 | 1.514 | 0.8 | 14. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------------|------------------------|------------------------|
| MILL RACE CREEK L2 | 0.073 | 1.234 |
| SOUTH FORK | 0.026 | 0.476 |
| CURRENT CREEK (82) | 0.042 | 1.331 |

* ESTIMATED

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN UTAH

NAME - WILLARD RESERVOIR (EUTROPHIC)
 COUNTY - BOX ELDER
 STORET NO. - 4925 WORKING PAPER NO. 862, NTIS ACCESSION NO. PB-279 254/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 62.20 | 40.47 | 5.9 | 2,570 | 3.3 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 171. | 989. | 1.1 | 0.044 | 0.009 | 0.060 | 0.620 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 7.6 | ***** | (5/14/75) N (8/ 6/75) N (9/23/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/14/ 0 | 8/ 6/ 0 | 9/23/ 0 |
|--|---------|---------|---------|
|--|---------|---------|---------|

| | | | | | |
|----------------|-------|----------------|-------|---------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| ANKISTRODESMUS | 2412 | MELOSIRA | 2987 | APHANIZOMENON | 1104 |
| CRYPTOMONAS | 36 | FRAGILARIA | 1873 | CRYPTOMONAS | 28 |
| | | CRYPTOMONAS | 253 | | |
| | | OOCYSTIS | 202 | | |
| | | ANKISTRODESMUS | 202 | | |
| OTHER | 0 | OTHER | 51 | OTHER | 0 |
| TOTAL | 2448 | TOTAL | 5568 | TOTAL | 1132 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 110. | ***** | ***** | 8250. | 8360. |
| NITROGEN | 1485. | ***** | ***** | 210285. | 211770. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 7380. | 12. | 0.21 |
| NITROGEN | 124710. | 41. | 5.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| WILLARD CANAL | 2.250 | ***** | 0.101 | 2.132 | ***** | ***** |
| UNNAMED STREAM B1 | 0.184 | 12.6 | 0.056 | 1.294 | 26. | 596. |
| UNNAMED STREAM C1 | 0.020 | 2.3 | 0.047 | 2.061 | 13. | 565. |
| UNNAMED STREAM D1 | 0.011 | 0.3 | 0.024 | 1.786 | 33. | 2067. |
| UNNAMED STREAM E1 | 0.006 | 1.6 | 0.020 | 2.191 | 3. | 259. |
| UNNAMED STREAM F1 | 0.042 | 4.9 | 0.020 | 2.881 | 5. | 779. |
| UNNAMED STREAM G1 | 0.057 | ***** | 0.050 | 1.748 | ***** | ***** |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - AMERICAN LAKE
 COUNTY - PIERCE
 STORET NO. - 5301

WORKING PAPER NO. 864. NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 65.80 | 4.70 | 15.4 | 0.160 | 38.2 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 41. | 69. | 4.0 | 0.027 | 0.007 | 0.105 | 0.450 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (4.7 - 5.3 (2)) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--------------------------|--|
| 4.8 | 0.7 | 5.3 (2) | (4/ 1/75) P (7/17/75) N (10/29/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 4/ 1/75 | 7/17/75 | 10/29/75 | |
|----------------|-------------|-----------------|--------------|-------------|
| GENERA | COUNT | GENERA | COUNT | |
| FRAGILARIA | 435 | CYCLOTELLA | 529 | |
| CHROOMONAS | 373 | CHROOMONAS | 511 | |
| MELOSIRA | 373 | MELOSIRA | 91 | |
| CRYPTOMONAS | 155 | OOCYSTIS | 73 | |
| STEPHANODISCUS | 93 | ANKISTRODESMIUS | 55 | |
| OTHER | 124 | OTHFR | 220 | |
| TOTAL | 1553 | TOTAL | 1479 | |
| | | | TOTAL | 1782 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 35. | 445. | 480. |
| NITROGEN | ***** | ***** | 1330. | 12835. | 14165. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | **** | 0.10 |
| NITROGEN | ***** | **** | 3.0 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL-N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| MURRAY CREEK | 0.160 | 28.2 | 0.026 | 0.558 | 6. | 127. |

140

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - BANKS LAKE
 COUNTY - GRANT, DOUGLAS
 STORET NO. - 5302

(MESOTROPHIC)
 WORKING PAPER NO. 865, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | ***** | 100.77 | 12.4 | ***** | 175.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 60. | 120. | 3.4 | 0.021 | 0.007 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD | LIMITING NUTRIENT AT SAMPLING TIME | |
|------------------------------|---------------------------------------|-------------|------------------------------------|-------------------|
| 7.4 | 0.2 - 0.7 (4) | (4/ 3/75) N | (7/22/75) N | (9/10/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA

| | 4/ 3/75 | 7/22/75 | 9/10/75 | | |
|--------------|-------------|----------------|-------------|---------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| ASTERIONELLA | 3034 | MELOSIRA | 690 | ASTERIONELLA | 624 |
| TABELLARIA | 358 | ASTERIONELLA | 248 | TABELLARIA | 451 |
| LYNGBYA | 167 | ANABAENA | 221 | FRAGILARIA | 416 |
| DINOBRYON | 119 | ANKISTRODESmus | 55 | CHROOMONAS | 243 |
| CHROOMONAS | 96 | CHROOMONAS | 55 | APHANIZOMENON | 173 |
| OTHER | 168 | OTHER | 111 | OTHER | 348 |
| TOTAL | 3942 | TOTAL | 1380 | TOTAL | 2255 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|----------------|------------------------|------------------------|
| PAYNES GULCH | 0.259 | 5.734 |
| NORTHRUP CREEK | 0.196 | 2.219 |
| FEEDER CANAL | 0.039 | 0.406 |
| RUSHO CREEK | 1.120 | 8.019 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - LAKE CHELAN (OLIGOTROPHIC)
COUNTY - CHELAN
STOKET NO. - 5303 WORKING PAPER NO. 866, NTIS ACCESSION N

1 AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 2393.00 | 133.80 | 8.8888 | 61,940 | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN (METERS) | SECCHI DISC | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------|-------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 21. | 37. | 9.9 | | 0.005 | 0.003 | 0.070 | 0.250 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

• BIOLOGICAL CHARACTERISTICS (CONT'D)
 MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (4/21/75) P (7/21/75) N (9/11/75) P

SUMMARY OF PHYTOPLANKTON DAT

SUMMARY OF PHYTOPLANKTON DATA 7/21/75 9/11/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|--------------|------------|-----------------------|------------|--------------|-----------|
| ASTERIONELLA | 374 | TABELLARIA | 262 | TABELLARIA | 12 |
| TABELLARIA | 166 | ASTERIONELLA | 105 | | |
| CYCLOTELLA | 146 | CHLOROPHYTAN COLONIES | 79 | | |
| CYMBELLA | 21 | APHANOThCE | 26 | | |
| OOCYSTIS | 21 | | | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 728 | TOTAL | 472 | TOTAL | 12 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| 455. | ***** | 145. | 34095. | 34695. |
| 1360. | ***** | 5390. | 1320505. | 1327255. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 24225. | 30. | | 0.26 |
| NITROGEN | 303120. | 77. | | 9.9 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NONPOINT-SOURCE NUTRIENT STREAM NAME | EXPORT MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------------------------|------------------------|-----------------------|---------------------|---------------------|------------------------------|------------------------------|
| STEHEKIN RIVER | 40.470 | 891.0 | 0.014 | 0.635 | 20. | 910. |
| RAILROAD CREEK | 5.780 | 172.0 | 0.011 | 0.553 | 12. | 586. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------------|------------------------|------------------------|
| STINK CREEK | 0.045 | 1.129 |
| POISON CREEK | 0.020 | 0.248 |
| TWENTY FIVE MILE CREEK | 0.014 | 0.276 |
| GRADE CREEK | 0.020 | 0.130 |
| PRINCE CREEK | 0.011 | 0.360 |
| MITCHELL CREEK | 0.026 | 0.720 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - DIAMOND LAKE
 COUNTY - PEND OREILLE
 STORET NO. - 5304

WORKING PAPER NO. 867. NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 3.07 | 8.1 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 38. | 53. | 5.0 | 0.014 | 0.010 | 0.060 | 0.530 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|--|
| 14.5 | 0.4 | (6/ 4/75) N (7/23/75) N (9/16/75) P (10/24/75) P |

| SUMMARY OF PHYTOPLANKTON DATA | | | | | |
|-------------------------------|------------|--------------|-------------|--------------|-------------|
| | 6/ 4/75 | 7/23/75 | | 9/16/75 | |
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| DINOBYRON | 306 | PHORMIDIUM | 969 | CHROOCOCCUS | 366 |
| CYCLOTELLA | 167 | GLOEOCAPSA | 588 | | |
| OOCYSTIS | 56 | ANABAENA | 554 | CYCLOTELLA | 284 |
| CHROOMONAS | 56 | EUNOTIA | 554 | EUNOTIA | 244 |
| CRYPTOMONAS | 28 | COSMARUM | 242 | NAVICULA | 244 |
| OTHER | 83 | OTHER | 2147 | OTHER | 2440 |
| TOTAL | 696 | TOTAL | 5054 | TOTAL | 3578 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------------|------------------------|------------------------|
| MOON CREEK (OUTLET) | 0.030 | 0.855 |

14
cc

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - GREEN LAKE
 COUNTY - KING
 STORET NO. - 5305

WORKING PAPER NO. 868, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 1.04 | 3.8 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEDIAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|-----------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 22. | 33. | 2.2 | 0.027 | 0.009 | 0.050 | 0.360 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 3.0 | 0.8 - 2.9 (2) | (4/1/75) N (7/18/75) N (10/29/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 4/1/75 | 7/18/75 | 10/29/75 | | |
|----------------|--------|------------------------|----------|--------------------|------|
| GENERA | COUNT | GENERA | COUNT | | |
| ASTERIONELLA | 3960 | ANACYSTIS(MICROCYSTIS) | 341 | ASTERIONELLA | 676 |
| APHANIZOMENON | 2487 | OSCILLATORIA | 341 | CHRYSOPHYTAN CELLS | 204 |
| FRAGILARIA | 1301 | CYCLOTELLA | 189 | ANABAENA | 157 |
| FLAGELLATES | 287 | CHROOMONAS | 114 | FRAGILARIA | 157 |
| CENTRIC DIATOM | 57 | CLOSTERIUM | 38 | APHANIZOMENON | 94 |
| OTHER | 192 | OTHER | 37 | OTHER | 158 |
| TOTAL | 8244 | TOTAL | 1060 | TOTAL | 1446 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

**** LAKE SAMPLING ONLY ****

VII

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - KEECHELUS LAKE
 COUNTY - KITTITAS
 STORET NO. - 5306 WORKING PAPER NO. 869, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 141.70 | 10.36 | 18.7 | 10.200 | 220.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEDIAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|-----------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 15. | 24. | 5.6 | 0.007 | 0.002 | 0.040 | 0.220 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 1.4 | 0.2 - 0.3 (3) | (7/21/75) P AND N (9/12/75) P (10/28/75) P |

SUMMARY OF PHYTOPLANKTON DATA
 7/21/75 9/12/75 10/28/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|--------------|------------|--------------------|------------|--------------|-----------|
| CYCLOTELLA | 466 | CHLOROPHYTAN CELLS | 195 | CHROOMONAS | 24 |
| CHROOMONAS | 89 | DINOBRYON | 195 | DINOBRYON | 24 |
| MELOSIRA | 89 | MELOSIRA | 130 | | |
| CRYPTOMONAS | 22 | CRYPTOMONAS | 22 | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 666 | TOTAL | 542 | TOTAL | 48 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 4525. | 4525. |
| NITROGEN | ***** | ***** | ***** | 95980. | 95980. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 1685. | 63. | 0.44 |
| NITROGEN | 40835. | 57. | 9.3 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| MEADOW CREEK | 1.470 | 20.4 | 0.008 | 0.159 | 38. | 404. |
| GOLD CREEK | 2.610 | 36.3 | 0.007 | 0.196 | 28. | 835. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------|------------------------|------------------------|
| ROARING CREEK | 0.007 | 0.248 |
| COLD CREEK | 0.007 | 0.265 |

145

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - MAYFIELD LAKE
 COUNTY - LEWIS
 STORET NO. - 5307

(MESOTROPHIC)
 WORKING PAPER NO. 870. NTIS ACCESSION NO. PB-

/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 3626.00 | 8.90 | 17.6 | 188.350 | 9.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 24. | 30. | 2.5 | 0.014 | 0.007 | 0.100 | 0.280 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD 0.1 - 0.2 (2) | LIMITING NUTRIENT AT SAMPLING TIME (3/28/75) P AND N (7/17/75) N (10/30/75) P |
|------------------------------|---------------------------------------|------------------------|---|
| 4.2 | | | |

| SUMMARY OF PHYTOPLANKTON DATA | | | | | |
|-------------------------------|---------|--------------|-------|--------------|-------|
| | 3/28/75 | 7/17/75 | | 10/30/75 | |
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| MELOSIRA | 297 | ASTERIONELLA | 1223 | ASTERIONELLA | 145 |
| CRYPTOMONAS | 81 | TABELLARIA | 100 | CHROOMONAS | 29 |
| CHROOMONAS | 27 | CRYPTOMONAS | 25 | | |
| PENNATE DIATOMS | 27 | | | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 432 | TOTAL | 1348 | TOTAL | 174 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 935. | ***** | 10. | 100135. | 101080. |
| NITROGEN | 4105. | ***** | 320. | 1590505. | 1594930. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION LOSS | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|------------------------------|--|
| PHOSPHORUS | 102690. | | 11.36 |
| NITROGEN | 1548220. | 3. | 174.2 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| COWLITZ RIVER | 155.860 | 2988.9 | 0.015 | 0.224 | 26. | 383. |
| KLICKITAT CREEK | 0.660 | 26.9 | 0.053 | 1.448 | 24. | 1249. |
| TILTON RIVER | 23.980 | 365.2 | 0.025 | 0.406 | 46. | 856. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------|------------------------|------------------------|
| SILVER CREEK | 0.032 | 0.892 |
| WINSTON CREEK | 0.038 | 0.750 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - MEDICAL LAKE
 COUNTY - SPOKANE
 STORE NO. - 5308

WORKING PAPER NO. 871, NTIS ACCESSION NO. PB-

/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 0.64 | 9.7 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 1380. | 1479. | 2.5 | 0.275 | 0.166 | 0.225 | 2.900 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 16.4 | 9.0 - 30.5 (3) | (6/ 3/75) N (7/23/75) N (9/11/75) N (10/24/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 6/ 3/75 | 7/23/75 | 9/11/75 | 10/24/75 | |
|-------------|---------|------------------------|---------|------------------------|-------|
| GENERA | COUNT | GENERA | GENERA | GENERA | COUNT |
| CHROOMONAS | 769 | NITZSCHIA | 2267 | ANACYSTIS(MICROCYSTIS) | 1560 |
| SCHROEDERIA | 678 | ANACYSTIS(MICROCYSTIS) | 599 | PHORMIDIUM | 1147 |
| PHORMIDIUM | 271 | CRYPTOMONAS | 385 | FRAGILARIA | 551 |
| CRYPTOMONAS | 226 | CHROOMONAS | 342 | CRYPTOMONAS | 229 |
| OTHER | 0 | APHANOCAPSA | 257 | PENNATE DIATOMS | 229 |
| TOTAL | 1944 | OTHER | 384 | OTHER | 277 |
| | TOTAL | | TOTAL | TOTAL | |
| | | | 4234 | 3993 | 2467 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - MOSFS LAKE
 COUNTY - GRANT
 STORET NO. - 5309
 WORKING PAPER NO. 872, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 7990.10 | 27.58 | 5.9 | 4.560 | 289.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| ALKALINITY (MG/L) | MEDIAN 183. | MEDIAN 377. | MEAN SECCHI DISC 0.9 | MEDIAN 0.115 | MEDIAN 0.038 | MEDIAN 0.150 | MEDIAN 1.040 |
|----------------------|----------------|----------------|-------------------------|-----------------|-----------------|-----------------|-----------------|
| CONDUCTIVITY (UMHOS) | | | | TOTAL P (MG/L) | ORTHO P (MG/L) | INORG N (MG/L) | TOTAL N (MG/L) |
| | | | | | | | |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (4.7 - 14.8 (4)) | LIMITING NUTRIENT AT SAMPLING TIME (4/2/75) P AND N (7/21/75) N (9/10/75) N |
|------------------------------|---------------------------------------|---------------------------|---|
| 29.1 | | | |

SUMMARY OF PHYTOPLANKTON DATA
4/2/75 7/21/75 9/10/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|--------------|---------------|--------------|---------------|--------------|
| STEPHANODISCUS | 37485 | APHANIZOMENON | 5406 | APHANIZOMENON | 7171 |
| SCENEDESMUS | 813 | OOCYSTIS | 2806 | OOCYSTIS | 850 |
| CRYPTOMONAS | 642 | SCENEDESMUS | 1471 | CRYPTOMONAS | 702 |
| CHROOMONAS | 428 | MELOSIRA | 753 | MELOSIRA | 481 |
| MELOSIRA | 342 | CHROOMONAS | 479 | OSCILLATORIA | 444 |
| OTHER | 1241 | OTHFR | 1472 | OTHER | 923 |
| TOTAL | 40951 | TOTAL | 12387 | TOTAL | 10571 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 9905. | ***** | 20. | 21220. | 31145. |
| NITROGEN | 21210. | ***** | 780. | 335925. | 357915. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 17230. | 45. | 1.13 |
| NITROGEN | 206055. | 42. | 13.0 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-----------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| CRAH CREEK | 2.010 | 5770.5 | 0.099 | 1.734 | 0.8 | 16. |
| ROCKY COULEE WASTEWAY | ***** | 764.0 | 0.122 | 1.904 | 5. | 79. |
| RUCKY FORD CREEK | 2.550 | 1217.3 | 0.139 | 1.752 | 9. | 112. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - OZETTE LAKE
 COUNTY - CLALLAM
 STORET NO. - 5310

(OLIGO-MESOTROPHIC)
 WORKING PAPER NO. 873, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 29.54 | 39.6 | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 10. | 29. | 2.5 | 0.010 | 0.009 | 0.110 | 0.290 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 1.2 | 0.1 | (3/31/75) N (7/18/75) N (10/29/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/31/75 | 7/18/75 | | 10/29/75 | |
|-------------|---------|-----------------|-------|----------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 120 | MELOSIRA | 719 | STEPHANODISCUS | 66 |
| CYCLOTELLA | 60 | FLAGELLATES | 332 | CHROOMONAS | 66 |
| MELOSIRA | 60 | PENNATE DIATOMS | 166 | SYNEDRA | 22 |
| CRYPTOMONAS | 30 | CRYPTOMONAS | 111 | CLOSTERIUM | 22 |
| | | ANKISTRODESmus | 111 | OSCILLATORIA | 22 |
| OTHER | 0 | OTHER | 165 | OTHER | 24 |
| TOTAL | 270 | TOTAL | 1604 | TOTAL | 222 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|----------------|------------------------|------------------------|
| OZETTE RIVER | 0.015 | 0.344 |
| UMBRELLA CREEK | 0.015 | 0.236 |
| BIG RIVER | 0.022 | 0.433 |
| CROOKED CREEK | 0.015 | 0.282 |
| TROUT CREEK | 0.041 | 0.454 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - SAMMAMISH LAKE (MESO-EUTROPHIC)

COUNTY - KING
STORET NO. - 5311

WORKING PAPER NO. 874. NTIS ACCESSION NO. PB- /AB

1 AE

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 253.00 | 19.82 | 17.7 | 6.410 | 1.8 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY (MG/L) | MEDIAN CONDUCTIVITY (UMHOS) | MEAN (METERS) | MEDIAN TOTAL P (MG/L) | MEDIAN ORTHO P (MG/L) | MEDIAN INORG N (MG/L) | MEDIAN TOTAL N (MG/L) |
|-----------------------------|--------------------------------|------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| 39. | 81. | 3.2 | 0.015 | 0.006 | 0.210 | 0.540 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (3/31/75) P (7/17/75) P (10/28/75) P

SUMMARY OF PHYTOPLANKTON DATA

3/31/75 7/17/75 10/28/75

| GENERAL | COUNT | GENERAL | COUNT | GENERAL | COUNT |
|----------------|-------|-----------------------|-------|---------------|-------|
| MELOSIRA | 5024 | APHANOCAPSA | 1081 | APHANOCAPSA | 683 |
| FRAGILARIA | 1136 | APHANOTHECE | 765 | FRAGILARIA | 683 |
| STEPHANODISCUS | 723 | CHROOMONAS | 90 | CHROOMONAS | 528 |
| ASTERIONELLA | 551 | BOTRYOCOCCUS | 45 | CRYPTOMONAS | 186 |
| SYNEDRA | 344 | CHLOROPHYTAN COLONIES | 45 | APHANIZOMENON | 31 |
| OTHER | 344 | OTHER | 135 | OTHER | 63 |
| TOTAL | 8122 | TOTAL | 2161 | TOTAL | 2174 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT

| • INPUT | POINT SOURCE MUNICIPAL (KG/YP) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YP) | NON-POINT SOURCE (KG/YP) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 110. | 6020. | 6130. |
| NITROGEN | ***** | ***** | 4165. | 336010. | 340175. |

8. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE |
|------------|----------------------|----------------------|----------------------------------|--------------|
| PHOSPHORUS | 4450. | 27. | 0.31 | |
| NITROGEN | 129275. | 52. | 17.2 | |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| NON-POINT SOURCE NUTRIENT | MEAN FLOW | DRAINAGE AREA | MEAN TOTAL P | MEAN TOTAL N | TOTAL P EXPORT | TOTAL N EXPORT |
|---------------------------|-----------|---------------|--------------|--------------|----------------|----------------|
| | (CMS) | (SQ KM) | (MG/L) | (MG/L) | (KG/SQ KM/YR) | (KG/SQ KM/YR) |
| ISSAQAH CREEK | 4.140 | 141.7 | 0.036 | 1.309 | 33. | 1259. |
| TIBBETTS CREEK | 0.250 | 13.0 | 0.023 | 1.907 | 11. | 1441. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------|------------------------|------------------------|
| UNNAMED CREEK B1 | 0.025 | 1.686 |
| UNNAMED CANAL C1 | 0.065 | 0.735 |

13

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - LAKE WHATCOM
 COUNTY - WHATCOM
 STORET NO. - 5312

(MESOTROPHIC)
 WORKING PAPER NO. 875, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 145.30 | 20.25 | 1.8 | 4.770 | 95.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 17. | 39. | 5.4 | 0.009 | 0.009 | 0.320 | 0.510 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 3.4 | 0.3 - 0.4 (2) | (3/31/75) P (7/18/75) P (10/29/75) P |

SUMMARY OF PHYTOPLANKTON DATA

| | 3/31/75 | 7/18/75 | | 10/29/75 | |
|----------------|---------|------------|-------|--------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| MELOSIRA | 735 | TABELLARIA | 733 | ASTERIONELLA | 184 |
| ASTERIONELLA | 331 | SYNEDRA | 193 | FRAGILARIA | 184 |
| CHROOMONAS | 257 | CYCLOTELLA | 38 | SPONDYLOSIUM | 184 |
| STEPHANODISCUS | 184 | MALLOMONAS | 38 | CHROOMONAS | 147 |
| TABELLARIA | 110 | OTHER | 0 | APHANOCAPSA | 73 |
| OTHER | 0 | | | OTHER | 109 |
| TOTAL | 1617 | TOTAL | 1002 | TOTAL | 881 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 75. | 2515. | 2590. |
| NITROGEN | ***** | ***** | 2785. | 164600. | 167385. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 1480. | 43. | 0.13 |
| NITROGEN | 89620. | 46. | 8.3 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| ULSEN CREEK | 0.300 | 9.8 | 0.019 | 0.787 | 17. | 1112. |
| BEAVER CREEK | 0.580 | 20.0 | 0.022 | 0.838 | 18. | 1270. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WASHINGTON

NAME - LOWER GRANITE RESERVOIR (EUTROPHIC)

COUNTY - GARFIELD, WHITMAN

STORET NO. - 5313

WORKING PAPER NO. 876, NTIS ACCESSION NO. PB-

/AB

I. MORPHOMETRY

| LAKE TYPE IMPOUNDMENT | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|--------------------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| ***** | 36.42 | 13.7 | ***** | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOH) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 59. | 140. | 1.6 | 0.033 | 0.022 | 0.150 | 0.330 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (1.1 - 3.9 (4) | LIMITING NUTRIENT AT SAMPLING TIME (7/23/75) N | LIMITING NUTRIENT AT SAMPLING TIME (9/12/75) N |
|------------------------------|---------------------------------------|----------------------------|--|--|
| 4.9 | | | | |

SUMMARY OF PHYTOPLANKTON DATA
7/23/75 9/12/75

| GENERA | COUNT | GENERA | COUNT |
|----------------|-------------|--------------|-------------|
| SKELETONEMA | 630 | MELOSIHA | 810 |
| CENTRIC DIATOM | 540 | CHROOMONAS | 778 |
| FRAGILARIA | 210 | SKELETONEMA | 195 |
| CHROOMONAS | 60 | CYCLOTELLA | 162 |
| NITZSCHIA | 60 | NITZSCHIA | 130 |
| OTHER | 150 | OTHER | 812 |
| TOTAL | 1650 | TOTAL | 2887 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| ***** | ***** | ***** | ***** | ***** |
| PHOSPHORUS | | | | |
| NITROGEN | | | | |

B. OUTPUT

| OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|----------------------|----------------------|---|
| ***** | ***** | ***** |
| PHOSPHORUS | | |
| NITROGEN | | |

**** LAKE SAMPLING ONLY ****

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - BIG SANDY RESERVOIR (EUTROPHIC)
 COUNTY - SUBLETTE, SWEETWATER
 STORET NO. - 5601 WORKING PAPER NO. 881, NTIS ACCESSION NO. PB-276 825/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 1139.60 | 8.34 | 5.8 | 2,462 | 273.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 41. | 98. | 0.3 | 0.087 | 0.020 | 0.060 | 0.430 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|---|
| 4.4 | ***** | (5/19/75) N (9/2/75) N (10/16/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/19/75 | 9/2/75 | 10/16/75 | | |
|----------------|---------|---------------|----------|-------------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CENTRIC DIATOM | 568 | APHANIZOMENON | 577 | CHROOMONAS | 1050 |
| CHROOMONAS | 512 | CHROOMONAS | 241 | CRYPTOMONAS | 131 |
| CRYPTOMONAS | 57 | CRYPTOMONAS | 120 | SCHROEDERIA | 98 |
| SURIKELLA | 28 | SCHROEDERIA | 48 | | |
| NITZSCHIA | 28 | | | | |
| OTHER | 0 | OTHER | 0 | OTHER | 0 |
| TOTAL | 1193 | TOTAL | 986 | TOTAL | 1279 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | 9980. | 9980. |
| NITROGEN | ***** | ***** | ***** | 133855. | 133855. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 8485. | 15. | 1.20 |
| NITROGEN | 99505. | 26. | 16.0 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| HIG SANDY RIVER | 2,450 | 834.0 | 0.114 | 1.131 | 9. | 110. |

150

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - BOULDER LAKE
 COUNTY - SWEETWATER
 STORET NO. - 5602

WORKING PAPER NO. 882. NTIS ACCESSION NO. PB-277 049/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| NATURAL | 336.70 | 7.03 | 12.2 | 5.724 | 181.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 16. | 3. | 3.5 | 0.008 | 0.002 | 0.040 | 0.235 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|-------|------------------------------------|
| 2.5 | ***** | | (8/28/75) N (10/17/75) P |

SUMMARY OF PHYTOPLANKTON DATA
8/28/75 10/17/75

| GENERA | COUNT | GENERA | COUNT |
|---------------|-------|---------------|-------|
| CHLAMYDOMONAS | 3806 | SPHAEROCYSTIS | 104 |
| CHROOMONAS | 327 | APHANOTHECE | 78 |
| CRYPTOMONAS | 59 | ELAKATOTHRIX | 52 |
| SPHAEROCYSTIS | 30 | CRYPTOMONAS | 26 |
| OTHER | 0 | CHROOMONAS | 26 |
| TOTAL | 4222 | OTHER | 0 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS ***** | ***** | 5. | 4800. | 4805. |
| NITROGEN ***** | ***** | 90. | 137030. | 137120. |

B. OUTPUT

| OUTLET(S) (KG/YR) | PERCFTN RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|----------------------|----------------------|---|
| PHOSPHORUS 3105. | 35. | 0.68 |
| NITROGEN 176470. | LOSS | 19.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|---------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BOULDER CREEK | 5.180 | 297.8 | 0.022 | 0.676 | 12. | 371. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------|------------------------|------------------------|
| UNNAMED CREEK HI | 0.063 | 1.108 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - BOYSEN RESERVOIR (EUTROPHIC)
 COUNTY - FREMONT
 STORET NO. - 5603 WORKING PAPER NO. 883, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 19945.60 | 89.84 | 10.4 | 45.717 | 270.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 129. | 454. | 0.9 | 0.037 | 0.014 | 0.140 | 0.340 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|---|
| 6.3 | ***** | (5/19/75) P AND N (9/2/75) P AND N (10/17/75) N |

SUMMARY OF PHYTOPLANKTON DATA
5/19/75 9/2/75

| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
|----------------|-------------|---------------|-------------|---------------|------------|
| FRAGILARIA | 2777 | APHANIZOMENON | 1129 | CHROOMONAS | 350 |
| ASTERIONELLA | 2529 | CHROOMONAS | 635 | NAVICULA | 225 |
| CHROOMONAS | 397 | CRYPTOMONAS | 212 | APHANIZOMENON | 125 |
| CENTRIC DIATOM | 248 | ANABAENA | 71 | CRYPTOMONAS | 75 |
| CRYPTOMONAS | 124 | DINOBYRON | 47 | SYNEDRA | 50 |
| OTHER | 100 | OTHER | 93 | OTHER | 100 |
| TOTAL | 6175 | TOTAL | 2187 | TOTAL | 925 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS 18750. | ***** | ***** | 301270. | 320020. |
| NITROGEN 64780. | ***** | ***** | 2573715. | 2638495. |

B. OUTPUT

| OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|----------------------|----------------------|--|
| PHOSPHORUS 30360. | 91. | 3.56 |
| NITROGEN 1277070. | 52. | 29.4 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| WIND RIVER | 39.900 | 11370.1 | 0.180 | 1.188 | 22. | 137. |
| BADWATER CREEK | 0.650 | 2092.7 | 0.630 | 1.749 | 3. | 14. |
| DITCH (D1) | 0.040 | ***** | 0.060 | 0.597 | ***** | ***** |
| DITCH (E1) | 0.059 | ***** | 0.054 | 1.351 | ***** | ***** |
| FIVEMILE CREEK | 4.200 | 1082.6 | 0.151 | 3.590 | 23. | 450. |
| MUDY CREEK | 0.531 | 859.9 | 0.212 | 1.440 | 2. | 27. |
| WYOMING CANAL | 0.119 | ***** | 0.087 | 0.899 | ***** | ***** |
| COTTONWOOD CREEK | 0.055 | 427.3 | 0.118 | 1.351 | 4. | 57. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|--------------|------------------------|------------------------|
| POISON CREEK | 0.617 | 2.251 |

155

BIRDSEYE CREEK

0.037

1.611

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - LAKE DE SMET
 COUNTY - JOHNSON
 STORET NO. - 5604

(MESO-EUTROPHIC)
 WORKING PAPER NO. 884. NTIS ACCESSION NO. PB-277 050/AB

I. MORPHOMETRY

| LAKE TYPE IMPOUNDMENT | DRAINAGE AREA (SQ KM) 25.90 | SURFACE AREA (SQ KM) 10.74 | MEAN DEPTH (METERS) 12.8 | TOTAL INFLOW (CMS) 0.167 | RETENTION TIME (YEARS) ***** |
|--------------------------|-----------------------------------|----------------------------------|--------------------------------|--------------------------------|------------------------------------|
|--------------------------|-----------------------------------|----------------------------------|--------------------------------|--------------------------------|------------------------------------|

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) 167. | MEDIAN CONDUCTIVITY(UMHOS) 629. | MEAN SECCHI DISC (METERS) 2.3 | MEDIAN TOTAL P(MG/L) 0.033 | MEDIAN ORTHO P(MG/L) 0.006 | MEDIAN INORG N(MG/L) 0.040 | MEDIAN TOTAL N(MG/L) 0.620 |
|------------------------------------|---------------------------------------|-------------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
|------------------------------------|---------------------------------------|-------------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) 11.2 | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) ***** | LIMITING NUTRIENT AT SAMPLING TIME (5/22/75) N | (8/29/75) N | (10/15/75) N |
|--------------------------------------|--|--|--------------|--------------|
|--------------------------------------|--|--|--------------|--------------|

SUMMARY OF PHYTOPLANKTON DATA

| | 5/22/75 | 8/29/75 | 10/15/75 | |
|----------------|---------|------------------------|----------|------|
| GENERA | COUNT | GENERA | COUNT | |
| CENTRIC DIATOM | 5052 | MELOSIRA | 1749 | |
| FRAGILARIA | 1160 | FLAGELLATES | 875 | |
| DIATOMA | 414 | APHANIZOMENON | 807 | |
| STEPHANODISCUS | 373 | ANABAENA | 673 | |
| FLAGELLATES | 124 | ANACYSTIS(MICROCYSTIS) | 673 | |
| OTHER | 165 | OTHER | 807 | |
| TOTAL | 7288 | TOTAL | 5584 | |
| | | | TOTAL | 3773 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) ***** | POINT SOURCE INDUSTRIAL (KG/YR) ***** | POINT SOURCE SEPTIC TANKS (KG/YR) 5. | NON-POINT SOURCE (KG/YR) 770. | TOTAL LOADING (KG/YR) 775. |
|------------|--|---|--|-------------------------------------|----------------------------------|
| PHOSPHORUS | | | 215. | 18075. | 18290. |
| NITROGEN | | | | | |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 1030. | LOSS | 0.07 |
| NITROGEN | 11835. | 35. | 1.7 |

V. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------------------|------------------------|------------------------|
| LITTLE PINEY CREEK DIV. | 0.025 | 0.680 |
| SHELL CREEK | 0.110 | 1.230 |

157

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - FLAMING GORGE RESERVOIR (MESOTROPHIC--EUTROPHIC)
 COUNTY - SWEETWATER, WY; DAGGETT, UT
 STORET NO. - 5605 WORKING PAPER NO. 885, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| IMPOUNDMENT | 39109.00 | 137.67 | 33.9 | 66.182 | 2.3 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 160. | 514. | 3.4 | 0.014 | 0.003 | 0.605 | 0.900 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|---------|---|
| 5.6 | 0.2 | 0.4 (5) | (5/15/75) P AND N (8/7/75) P (9/22/75) P AND N |

SUMMARY OF PHYTOPLANKTON DATA

| 5/15/75 | 8/7/75 | 9/22/75 | | | |
|----------------|-------------|---------------|-------------|---------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| STEPHANODISCUS | 1981 | CHROOMONAS | 1331 | APHANIZOMENON | 1086 |
| FRAGILARIA | 775 | APHANIZOMENON | 951 | CRYPTOMONAS | 827 |
| DIATOMA | 258 | FRAGILARIA | 428 | CHROOMONAS | 465 |
| CHROOMONAS | 258 | CRYPTOMONAS | 333 | OSCILLATORIA | 103 |
| NAVICULA | 43 | MELOSIRA | 190 | | |
| OTHER | 0 | OTHER | 428 | OTHER | 0 |
| TOTAL | 3315 | TOTAL | 3661 | TOTAL | 2481 |

158

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 40405. | ***** | ***** | 145835. | 186240. |
| NITROGEN | 95255. | ***** | ***** | 2731180. | 2826435. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 24365. | 87. | 1.35 |
| NITROGEN | 2883495. | LOSS | 20.5 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| GREEN RIVER | 48.400 | 25900.0 | 0.060 | 1.027 | 2. | 58. |
| CART CREEK | 0.402 | 72.5 | 0.023 | 0.602 | 4. | 100. |
| W FURK EAGLE CREEK | 0.088 | 8.0 | 0.018 | 0.792 | 6. | 286. |
| UPPER MARSH CREEK | 0.076 | 116.5 | 0.092 | 1.590 | 2. | 34. |
| BIRCH SPRING DRAW | 0.128 | 62.2 | 0.086 | 20.570 | 4.* | 131.* |
| HENRYS FORK | 2.360 | 1346.8 | 0.062 | 0.918 | 4. | 52. |
| BLACKS FORK | 10.500 | 8029.0 | 0.124 | 1.308 | 9. | 68. |
| SAGE CREEK | 0.280 | 303.0 | 0.166 | 2.659 | 5. | 104. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------|------------------------|------------------------|
| CURRENT CREEK | 0.047 | 1.568 |

| | | |
|--------------------|-------|-------|
| MIDDLE MARSH CREEK | 0.081 | 1.528 |
| SPRING CREEK | 0.958 | 2.462 |
| UNNAMED CREEK (M1) | 0.018 | 0.843 |

* ESTIMATED

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - FREMONT LAKE (OLIGOTROPHIC)
COUNTY - SUHLETTÉ
STORET NO. - 5606 WORKING PAPER NO. 886, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | 295.30 | 20.23 | 24.4 | 6,300 | 2.5 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| PHYSICAL AND CHEMICAL CHARACTERISTICS | | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|---------------------------------------|---------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| ALKALINITY(MG/L) | CONDUCTIVITY(UMHOS) | 13.3 | 0.006 | 0.002 | 0.040 | 0.220 |
| 14. | 1. | | | | | |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L-DRY WT) (8/28/75) P (10/17/75) P

SUMMARY OF PHYTOPLANKTON DATA

| GENERA | COUNT | GENERA | COUNT |
|---------------|-------------|--------------------------|-------------|
| CHLAMYDOMONAS | 3336 | CHLAMYDOMONAS | 3181 |
| ASTERIONELLA | 128 | ASTERIONELLA | 468 |
| TARELLARIA | 86 | CHRYSOPHYTAN FLAGELLATES | 281 |
| RHIZOSOLENIA | 43 | TARELLARIA | 140 |
| DINOHYRON | 43 | CHROOMONAS | 94 |
| OTHER | 0 | OTHER | 0 |
| TOTAL | 3636 | TOTAL | 4164 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 15. | 2345. | 2360. |
| NITROGEN | ***** | ***** | 570. | 224065. | 224635. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE |
|------------|----------------------|----------------------|----------------------------------|--------------|
| PHOSPHORUS | 1870. | 21. | 0.12 | |
| NITROGEN | 28185. | 87. | 11.1 | |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| PINE CREEK | 5.090 | 196.8 | 0.009 | 0.841 | 7. | 735. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|-------------|------------------------|------------------------|
| FAIR CREEK | 0.011 | 0.588 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - GLENDO RESERVOIR (EUTROPHIC)
 COUNTY - CONVERSE, PLATTE
 STORET NO. - 5607

WORKING PAPER NO. 887, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 50523.10 | 53.97 | 18.3 | 47.113 | 264.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 148. | 525. | 1.0 | 0.045 | 0.014 | 0.315 | 0.695 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (0.6 - 5.3 (3) | LIMITING NUTRIENT AT SAMPLING TIME (5/22/75) P | (8/27/75) P | (10/14/75) P |
|------------------------------|---------------------------------------|---------------------------|--|--------------|--------------|
| 8.5 | | | | | |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/22/75 | 8/27/75 | 10/14/75 | | |
|----------------|-------------|---------------|-------------|----------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| DIATOMA | 645 | APHANIZOMENON | 2407 | NAVICULA | 440 |
| STEPHANODISCUS | 469 | NAVICULA | 193 | MELOSIRA | 439 |
| CRYPTOMONAS | 293 | DIATOMA | 48 | DIATOMA | 275 |
| ASTERIONELLA | 176 | RHOICOSPHENIA | 48 | CHROOMONAS | 275 |
| NAVICULA | 59 | SCENEDESMUS | 48 | STEPHANODISCUS | 165 |
| OTHER | 116 | OTHER | 0 | OTHER | 163 |
| TOTAL | 1758 | TOTAL | 2744 | TOTAL | 1757 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

| A. INPUT | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 2680. | ***** | 5. | 172270. | 174955. |
| NITROGEN | 8920. | ***** | 105. | 2243640. | 2252665. |

| B. OUTPUT | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 49205. | 72. | 3.24 |
| NITROGEN | 1742630. | 23. | 41.7 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| NORTH PLATTE RIVER | 47.060 | 48813.7 | 0.117 | 1.476 | 4. | 45. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|------------------|------------------------|------------------------|
| WHISKEY GULCH | 0.837 | 4.717 |
| ELKHORN CREEK | 0.010 | 0.298 |
| COTTONWOOD CREEK | 0.150 | 1.231 |
| MUDDY CREEK | 0.068 | 1.772 |
| WILLOW CREEK | 0.084 | 1.903 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - OCEAN LAKE
 COUNTY - FREMONT
 STORET NO. - 5609 WORKING PAPER NO. 889, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 24.68 | 4.2 | 1.205 | 4.4 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) 200. | MEDIAN CONDUCTIVITY(UMHOS) 1400. | MEAN SECCHI DISC (METERS) 0.5 | MEDIAN TOTAL P(MG/L) 0.043 | MEDIAN ORTHO P(MG/L) 0.004 | MEDIAN INORG N(MG/L) 0.040 | MEDIAN TOTAL N(MG/L) 0.820 |
|------------------------------------|--|-------------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
|------------------------------------|--|-------------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) 7.5 | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) ***** | LIMITING NUTRIENT AT SAMPLING TIME (5/19/75) N (9/ 2/75) P (10/16/75) N |
|-------------------------------------|--|--|
|-------------------------------------|--|--|

SUMMARY OF PHYTOPLANKTON DATA

| | 5/19/75 | 9/ 2/75 | | 10/16/75 |
|--------------|---------|---------------|-------|------------------------|
| GENERA | COUNT | GENERA | COUNT | GENERA |
| CRYPTOMONAS | 981 | OOCYSTIS | 609 | CHLOROPHYTAN FILAMENTS |
| TABELLARIA | 235 | CRYPTOMONAS | 578 | ASTERIONELLA |
| NITZSCHIA | 157 | APHANIZOMENON | 183 | CRYPTOMONAS |
| SCHROEDERIA | 118 | SCENEDESMUS | 91 | CHROOMONAS |
| CHLOROGONIUM | 78 | ANABAENA | 30 | ANKISTRODESmus |
| OTHER | 196 | OTHER | 61 | OTHER |
| TOTAL | 1765 | TOTAL | 1552 | TOTAL |
| | | | | 6778 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|--------------------------------------|---------------------------------------|---|-----------------------------|--------------------------|
| PHOSPHORUS | 70. | ***** | ***** | 6465. | 6535. |
| NITROGEN | 225. | ***** | ***** | 92050. | 92275. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 710. | 89. | 0.26 |
| NITROGEN | 34435. | 63. | 3.7 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| PILOT DRAIN | 0.550 | ***** | 0.210 | 1.999 | ***** | ***** |
| SAND BUTTE AQUEDUCT (E1) | 0.059 | ***** | 0.186 | 1.283 | ***** | ***** |
| SAND BUTTE AQUEDUCT (E2) | 0.033 | ***** | 0.047 | 0.413 | ***** | ***** |
| UNNAMED CREEK (F1) | 0.480 | ***** | 0.119 | 1.618 | ***** | ***** |
| UNNAMED CREEK (H1) | 0.050 | ***** | 0.089 | 1.729 | ***** | ***** |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - SEMINOE RESERVOIR (MESOTROPHIC-EUTROPHIC)

COUNTY - CARBON

STORET NO. - 5610

WORKING PAPER NO. 890. NTIS ACCESSION NO. PB-277 036/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 28979.50 | 48.77 | 25.6 | 43.309 | 342.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEDIAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|-----------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 107. | 334. | 1.3 | 0.030 | 0.007 | 0.130 | 0.480 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD (4.2 - 4.4 (4)) | LIMITING NUTRIENT AT SAMPLING TIME (5/19/75) P AND N (8/27/75) P AND N (10/16/75) P AND N |
|------------------------------|---------------------------------------|--------------------------|--|
| 2.5 | | | |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/19/75 | 8/27/75 | | 10/16/75 | |
|-----------------|------------|---------------|------------|---------------|-------------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT |
| CHROOMONAS | 306 | APHANIZOMENON | 675 | CHROOMONAS | 558 |
| GUMPHONEMA | 109 | CRYPTOMONAS | 75 | APHANIZOMENON | 512 |
| SYNEURA | 66 | CHROOMONAS | 75 | | |
| CENTRIC DIATOM | 66 | ASTERIONELLA | 25 | | |
| PENNATE DIATOMS | 66 | EPITHEMIA | 25 | | |
| OTHER | 87 | OTHFR | 0 | OTHER | 0 |
| TOTAL | 700 | TOTAL | 875 | TOTAL | 1070 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS 4205. | ***** | ***** | 137495. | 141700. |
| NITROGEN 16940. | ***** | ***** | 1671510. | 1688450. |

B. OUTPUT

| OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|----------------------|----------------------|---|
| PHOSPHORUS 39285. | 72. | 2.91 |
| NITROGEN 1634305. | 3. | 34.6 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|--------------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| NORTH PLATTE RIVER | 31.120 | 21067.1 | 0.070 | 1.066 | 4. | 50. |
| HURT CREEK | 0.042 | 103.6 | 0.191 | 13.929 | 3. | 112. |
| SAYLOR CREEK | 0.160 | 54.4 | 0.094 | 1.236 | 11. | 122. |
| AUSTIN CREEK | 0.359 | 69.9 | 0.063 | 0.939 | 17. | 203. |
| MEDICINE ROW RIVER | 4.410 | 6055.4 | 0.144 | 1.835 | 5. | 50. |
| TRROBLESON CREEK | 0.242 | 134.7 | 0.183 | 1.849 | 11. | 108. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|---------------|------------------------|------------------------|
| O'BRIEN CREEK | 0.072 | 0.971 |

16A

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - SODA LAKE
 COUNTY - SURETTE
 STORET NO. - 5611

WORKING PAPER NO. 891, NTIS ACCESSION NO. PB- /AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (YEARS) |
|-----------|--------------------------|-------------------------|------------------------|-----------------------|---------------------------|
| NATURAL | ***** | 1.11 | ***** | ***** | ***** |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 2090. | 2472. | 2.9 | 0.063 | 0.014 | 0.040 | 2.520 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL YIELD (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---|------------------------------------|
| 5.6 | 6.8 | (8/28/75) N (10/20/75) N |

SUMMARY OF PHYTOPLANKTON DATA
8/28/75

| GENERAL | COUNT |
|--------------------------|------------|
| CYANOPHYTAN COCCOID COL. | 221 |
| ELAKATOTHRIX | 85 |
| APHANIZOMENON | 17 |
| OTHER | 0 |
| TOTAL | 323 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | ***** | ***** | ***** |
| NITROGEN | ***** | ***** | ***** | ***** | ***** |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | ***** | **** | ***** |
| NITROGEN | ***** | ***** | ***** |

**** LAKE SAMPLING ONLY ****

16

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - VIVA NAUGHTON RESERVOIR (EUTROPHIC)
 COUNTY - LINCOLN
 STORET NO. - 5612

WORKING PAPER NO. 892. NTIS ACCESSION NO. PB-277 037/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 624.20 | 5.90 | 8.9 | 3.923 | 170.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 145. | 241. | 1.8 | 0.065 | 0.024 | 0.120 | 0.440 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|--|
| 25.1 | ***** | (8/ 6/75) N (9/18/75) N (10/16/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 8/ 6/75 | 9/18/75 | 10/16/75 |
|--|---------|---------|----------|
|--|---------|---------|----------|

| | | | | | |
|-------------------------|---------------|-------------------------|---------------|-------------------------|---------------|
| GENERA APHANIZOMENON | COUNT 4430 | GENERA APHANIZOMENON | COUNT 4073 | GENERA APHANIZOMENON | COUNT 2135 |
| CHROOMONAS | 292 | SCHROEDERIA | 41 | CRYPTOMONAS | 137 |
| ANACYSTIS(MICROCYSTIS) | 243 | | | CHROOMONAS | 172 |
| OTHER | 390 | OTHER | 0 | STEPHANODISCUS | 34 |
| TOTAL | 5355 | TOTAL | 4114 | OTHER | 0 |
| | | | | TOTAL | 2478 |

IV. NUTRIENT LOADING CHARACTERISTICS(LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | ***** | ***** | 10. | 3940. | 3950. |
| NITROGEN | ***** | ***** | 320. | 125965. | 126285. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 5290. | LOSS | 0.67 |
| NITROGEN | 19115. | 85. | 21.4 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| HAMS FORK | 3.740 | 530.9 | 0.029 | 0.936 | 7. | 210. |
| TRAIL CREEK | 0.055 | 7.5 | 0.085 | 1.699 | 19. | 393. |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - WOODRUFF NARROWS RES. (EUTROPHIC)
 COUNTY - UNTIA
 STORET NO. - 5613 WORKING PAPER NO. 893, NTIS ACCESSION NO. PB-277 047/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 2030.60 | 7.08 | 4.9 | 7.069 | 58.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 142. | 229. | 0.8 | 0.069 | 0.019 | 0.105 | 0.620 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

| MEAN CHLOROPHYLL A (UG/L) | ALGAL ASSAY CONTROL (MG/L--DRY WT) | YIELD | LIMITING NUTRIENT AT SAMPLING TIME |
|------------------------------|---------------------------------------|-------|--|
| 13.0 | ***** | | (5/16/75) N (8/7/75) N (9/18/75) N (10/16/75) N |

SUMMARY OF PHYTOPLANKTON DATA

| | 5/16/75 | 8/7/75 | 9/18/75 | 10/16/75 |
|----------------|---------|--------------|---------|----------------|
| GENERA | COUNT | GENERA | COUNT | GENERA |
| FLAGELLATES | 7723 | MERISMOPEDIA | 676 | ANABAENA |
| CHROOMONAS | 1296 | SCHROEDERIA | 386 | STEPHANODISCUS |
| CENTRIC DIATOM | 810 | CHROOMONAS | 145 | APHANIZOMENON |
| CRYPTOMONAS | 216 | ANABAENA | 97 | MELOSIRA |
| SURIRELLA | 54 | CRYPTOMONAS | 97 | |
| OTHER | 55 | OTHER | 48 | OTHER |
| TOTAL | 10154 | TOTAL | 1449 | TOTAL |
| | | | | 2247 |
| | | | | TOTAL |
| | | | | 533 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 6685. | ***** | ***** | 35970. | 42655. |
| NITROGEN | 17260. | ***** | ***** | 505895. | 523155. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|---|
| PHOSPHORUS | 17420. | 59. | 6.02 |
| NITROGEN | 215255. | 59. | 73.9 |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|-------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BEAR RIVER | 7.060 | 1947.7 | 0.122 | 1.730 | 18. | 246. |

VI. MEAN NUTRIENT CONCENTRATIONS IN UNGAGED STREAMS

| STREAM NAME | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) |
|----------------------|------------------------|------------------------|
| WHITNEY CANYON CREEK | 0.157 | 1.810 |
| RED CANYON CREEK | 0.230 | 1.441 |

COMPENDIUM OF NATIONAL EUTROPHICATION SURVEY LAKES IN WYOMING

NAME - YELLOWTAIL RESERVOIR (MESOTROPHIC--EUTROPHIC)
COUNTY - BIGHORN, WY; BIGHORN, CARBON, MT
STORED NO. - 5614 WORKING PAPER NO. 894, NTIS ACCESSION NO. PB-279 255/AB

I. MORPHOMETRY

| LAKE TYPE | DRAINAGE AREA (SQ KM) | SURFACE AREA (SQ KM) | MEAN DEPTH (METERS) | TOTAL INFLOW (CMS) | RETENTION TIME (DAYS) |
|-------------|--------------------------|-------------------------|------------------------|-----------------------|--------------------------|
| IMPOUNDMENT | 50937.50 | 51.34 | 26.8 | 106.180 | 158.0 |

II. PHYSICAL AND CHEMICAL CHARACTERISTICS

PHYSICAL AND CHEMICAL CHARACTERISTICS

| MEDIAN ALKALINITY(MG/L) | MEDIAN CONDUCTIVITY(UMHOS) | MEAN SECCHI DISC (METERS) | MEDIAN TOTAL P(MG/L) | MEDIAN ORTHO P(MG/L) | MEDIAN INORG N(MG/L) | MEDIAN TOTAL N(MG/L) |
|----------------------------|-------------------------------|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 139. | 54. | 3.4 | 0.026 | 0.017 | 0.310 | 0.500 |

III. BIOLOGICAL CHARACTERISTICS (LAKE)

MEAN CHLOROPHYLL A ALGAL ASSAY CONTROL YIELD LIMITING NUTRIENT AT SAMPLING TIME
 (UG/L) (MG/L--DRY WT) (5/21/75) P (8/29/75) P AND N (10/17/75) P

SUMMARY OF PHYTOPLANKTON DATA

| | 5/21/75 | | 8/29/ 0 | | 10/17/ 0 | |
|--------------|---------|------------------------|---------|--------------|----------|-------|
| GENERA | COUNT | GENERA | COUNT | GENERA | COUNT | COUNT |
| FRAGILARIA | 357 | APHANIZOMENON | 1728 | CHROOMONAS | 363 | 363 |
| MELOSIRA | 268 | SKELETONEMA | 241 | NAVICULA | 161 | 161 |
| ASTERIONELLA | 223 | CARTERIA | 201 | SYNEDRA | 81 | 81 |
| LYNGBYA | 134 | ANACYSTIS(MICROCYSTIS) | 201 | NITZSCHIA | 81 | 81 |
| NAVICULA | 134 | CHROOMONAS | 201 | OSCILLATORIA | 40 | 40 |
| OTHER | 359 | OTHER | 201 | OTHER | 100 | 100 |
| TOTAL | 1475 | TOTAL | 2773 | TOTAL | 826 | 826 |

IV. NUTRIENT LOADING CHARACTERISTICS (LAKE)

NUTRIENT A. INPUT

| | POINT SOURCE MUNICIPAL (KG/YR) | POINT SOURCE INDUSTRIAL (KG/YR) | POINT SOURCE SEPTIC TANKS (KG/YR) | NON-POINT SOURCE (KG/YR) | TOTAL LOADING (KG/YR) |
|------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------|--------------------------|
| PHOSPHORUS | 6055. | 2125. | ***** | 1040045. | 1048225. |
| NITROGEN | 15835. | 48840. | ***** | 5282760. | 5347435. |

B. OUTPUT

| | OUTLET(S) (KG/YR) | PERCENT RETENTION | LAKE SURFACE AREA (G/SQ M/YR) | LOADING RATE (G/SQ M/YR) |
|------------|----------------------|----------------------|----------------------------------|-----------------------------|
| PHOSPHORUS | 66965. | 94. | 20.42 | |
| NITROGEN | 5064620. | 5. | 104.2 | |

V. NON-POINT-SOURCE NUTRIENT EXPORT

| STREAM NAME | MEAN FLOW (CMS) | DRAINAGE AREA (SQ KM) | MEAN TOTAL P (MG/L) | MEAN TOTAL N (MG/L) | TOTAL P EXPORT (KG/SQ KM/YR) | TOTAL N EXPORT (KG/SQ KM/YR) |
|----------------|--------------------|--------------------------|------------------------|------------------------|---------------------------------|---------------------------------|
| BIGHORN RIVER | 64,660 | 40831.3 | 0.322 | 1.513 | 16. | 76. |
| SHOSHONE RIVER | 29,480 | 6086.5 | 0.413 | 2.081 | 57. | 309. |
| DRY HEAD CREEK | 0.300 | 197.6 | 0.035 | 1.296 | 6. | 69. |
| CROOKED CREEK | 0.300 | 300.4 | 0.100 | 1.108 | 2. | 29. |