United States
Environmental Protection
Agency

Office of
Pesticides and Toxic Substances
Washington DC 20460

August 1981

Pesticides

EPA

Temephos

O, O'-(thiodi-4,1-phenylene) bis(O,O-dimethyl phosphorothioate)

Pesticide Registration Standard

TEMEPHOS

Acknowledgements

I. Special Pesticide Review Division (SPRD)

David Brooks	Project Manager
Benjamin Lemlich	Project Manager

II. Technical Support Team

A. Hazard Evaluation Division (HED)

Raymond Kent	Residue Chemist
John Brantner	Toxicologist
Robert Holst	Plant Physiologist
Richard Stevens	Wildlife Ecologist
Hudson Boyd	Environmental Chemist

B. Benefits and Field Studies Division (BFSD)

William Gross	Entomologist
Russell Scarato	Economist
E. David Thomas	Supervisory Entomologist, Section Head

III. Special Acknowledgement

A. Special Pesticide Review Division (SPRD)

Joan Warshawsky	Section Head
Barbara Moore	Secretary
Tonda Hicks	Clerk Typist

TABLE OF CONTENTS

Pages

I. How to Register Under a Registration Standard..... 1

A. Organization of the Standard..... 1
B. Purpose of the Standard..... 1
C. Requirements to Reregister Under the Standard.... 3
D. "Product-Specific" Data and "Generic" Data..... 3
E. Data Compensation Requirements Under FIFRA 3(c)(1)(D)..... 5
F. Obtaining Data to Fill Data Gaps; FIFRA 3(c)(2)(B)..... 6
G. Amendments to the Standard..... 7

II. Regulatory Position..... 9

A. Introduction..... 9
B. Description of Chemical..... 9
C. Regulatory Position..... 9
D. Regulatory Rationale..... 9
E. Criteria for Registration Under the Standard..... 11
F. Tolerance Reassessment..... 22

III. Summary of Data Requirements and Data Gaps..... 24

A. Table III.A Generic Data Requirements... 25
B. Table III.B Product-Specific Data Requirements for Manufacturing-Use Products..... 25
C. Table III.C Product-Specific Data Requirements for End-Use Products..... 26

IV. Product Chemistry..... 55

A. Introduction..... 55
B. Chemical Identity..... 55
C. Manufacturing Process..... 55
D. Product Analytical Methods and Data..... 57
E. Physical/Chemical Properties..... 57
F. Summary of Data Gaps..... 59

V. Environmental Fate..... 60

A. Use Summary..... 60
B. Environmental Fate Profile..... 60
C. Exposure Profile..... 63
D. Summary of Data Gaps..... 65

VI. Toxicology..... 66

A. Toxicology Profile: Technical Temephos..... 66
B. Toxicology Profile: Manufacturing-Use Temephos... 67
C. Toxicology Profile: End-Use Temephos..... 68
D. Human Hazard Assessment..... 70
E. Summary of Data Gaps..... 71

VII. Residue Chemistry..... 73

A. Introduction..... 73
B. Metabolism of Temephos in Plants..... 73
C. Metabolism of Temephos in Animals..... 75
D. Analytical Methods..... 75
E. Residue in Plants..... 79
F. Storage/Stability..... 80
G. Residue of Temephos in Animals..... 80
H. Summary of Data Gaps..... 81

VIII. Ecological Effects..... 83

A. Ecological Effects Profile..... 83
B. Hazard Assessment..... 84
C. Summary of Data Gaps..... 85

IX. Efficacy..... 86

A. Efficacy Profile..... 86
B. Factors Influencing Efficacy..... 86
C. Use Sites..... 86
D. Summary of Data Gaps..... 88

X. Confidential Annex..... 89

XI. Bibliography..... 90

A. Guide to Use of Bibliography..... 90

B. Section 1: Citations Considered to be Part of the Data Base Supporting Registrations Under the Standard

C. Section 2: Citations Judged to be Inappropriate for Use in Developing this Standard

D. Section 3: Standard Reference Material

I. How to Register Under a Registration Standard

A. Organization of the Standard

The first part of this document explains the purpose of a Registration Standard and summarizes the legal principles involved in registering or reregistering under a Standard. The second part presents the Agency's regulatory position and rationale. The third part sets forth the requirements, in tabular form, that must be met to obtain or retain registration for products covered by this particular Registration Standard. In the remaining parts, the Agency reviews the available data by scientific discipline, discusses the Agency's concerns with the identified potential hazards, and logically develops the conditions and requirements that would reduce those hazards to acceptable levels.

B. Purpose of the Standard

Section 3 of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) provides that "no person in any State may distribute, sell, offer for sale, hold for sale, ship, deliver for shipment, or receiveand (having so received) deliver or offer to deliver, to any person any pesticide which is not registered with the Administrator [of EPA]." To approve the registration of a pesticide, the Administrator must find, pursuant to Section 3(c)(5) that.

> (A) its composition is such as to warrant the proposed claims for it;
>
> (B) its labeling and other material required to be submitted comply with the requirements of this Act;
>
> (C) it will perform its intended function without unreasonable adverse effects on the environment; and
>
> (D) when used in accordance with widespread and commonly recognized practice it will not generally cause unreasonable adverse effects on the environment.

In making these findings, the Agency reviews a wide range of data which registrants are required to submit, and assesses the risks associated with the use of the proposed pesticide. However, the established approach to making these findings has been found to be defective on two counts.

First, the Environmental Protection Agency (EPA) and its predecessor agency, the United States Department of Agriculture (USDA), routinely reviewed registration applications on a "product-by-product" basis, evaluating each product-specific application somewhat independently. In the review of products containing similar components, there was little opportunity for a retrospective review of the full range of pertinent data available in Agency files and in the public literature. Thus the "product-by-product" approach was often inefficient and

sometimes resulted in inconsistent or incomplete regulatory judgments.

Second, over the years, as a result of inevitable and continuing advances in scientific knowledge, methodology, and policy, the data base for many pesticides came to be considered inadequate by current scientific and regulatory standards. Given the long history of pesticide regulation in several agencies, it is even likely that materials may have been lost from the data files. When EPA issued new requirements for registration in 1975 (40 CFR 162) and proposed new guidelines for hazard testing in 1978 (43 FR 29686, July 10, 1978 and 43 FR 37336, August 22, 1978) and in 1980 (45 FR 72948, November 3, 1980), many products that had already been registered for years were being sold and used without the same assurances of human and environmental safety as was being required for new products. Because of this inconsistency, Congress directed EPA to reregister all previously registered products, so as to bring their registrations and their data bases into compliance with current requirements [see FIFRA Section 3(g)].

Facing the enormous job of re-reviewing and calling-in new data for the approximately 35,000 current registrations, and realizing the inefficiencies of the "product-by-product" approach, the Agency decided that a new, more effective method of review was needed.

A new review procedure has been developed. Under it, EPA publishes documents called Registration Standards, each of which discusses a particular pesticide active ingredient. Each Registration Standard summarizes all the data available to the Agency on a particular active ingredient and its current uses, and sets forth the Agency's comprehensive position on the conditions and requirements for registration of all existing and future products which contain that active ingredient. These conditions and requirements, all of which must be met to obtain or retain full registration or reregistration under Section 3(c)(5) of FIFRA, include the submission of needed scientific data which the Agency does not now have; compliance with standards of toxicity, composition, labeling, and packaging; and satisfaction of the compensation provisions of FIFRA Section 3(c)(1)(D).

The Standard will also serve as a tool for product classification. As part of the registration of a pesticide product, EPA may classify each product for "general use" or "restricted use" [FIFRA Section 3(d)]. A pesticide is classified for "restricted use" when some special regulatory restriction is needed to ensure against unreasonable adverse effects to man or the environment. Many such risks of unreasonable adverse effects can be lessened if expresslydesigned label precautions are strictly followed. Thus, the special regulatory restriction for a "restricted use" pesticide is usually a requirement that it be applied only by, or under the supervision of, an applicator who has been certified by the State or Federal government as being competent to use pesticides safely, responsibly, and in

accordance with label directions. A restricted use pesticide can have other regulatory restrictions [40 CFR 162.11(c)(5)] instead of, or in addition to, the certified applicator requirement. These other regulatory restrictions may include such actions as seasonal or regional limitations on use, or a requirement for the monitoring of residue levels after use. A pesticide classified for "general use", or not classified at all, is available for use by any individual who is in compliance with State or local regulations. The Registration Standard review compares information about potential adverse effects of specific uses of the pesticide with risk criteria listed in 40 CFR 162.11(c), and thereby determines whether a product needs to be classified for "restricted use". If the Standard does classify a pesticide for "restricted use", this determination is stated in Part II of this Standard.

C. Requirement to Reregister Under the Standard

FIFRA Section 3(g), as amended in 1978, directs EPA to reregister all currently registered products as expeditiously as possible. Congress also agreed that reregistration should be accomplished by the use of Registration Standards.

Each registrant of a currently registered product to which this Standard applies, and who wishes to continue to sell or distribute his product in commerce, must apply for reregistration. His application must contain proposed labeling that complies with this Standard.

EPA will issue a notice of intent to cancel the registration of any currently registered product to which this Standard applies if the registrant fails to comply with the procedures for reregistration set forth in the Guidance Package which accompanies this Standard.

D. "Product-Specific" Data and "Generic" Data

In the course of developing this Standard, EPA has determined the types of data needed for evaluation of the properties and effects of products to which the Standard applies, in the disciplinary areas of Product Chemistry, Environmental Fate, Toxicology, Residue Chemistry, Ecological Effects, and Efficacy. These determinations are based primarily on the data Guidelines proposed in 1978 (43 FR 29686, July 10, 1978, and 43 FR 37336, August 22, 1978), and in 1980 (45 FR 72948, November 3, 1980), as applied to the use patterns of the products to which this Standard applies. Where it appeared that data from a normally applicable Guidelines requirement were actually unnecessary to evaluate these products, the Standard indicates that the requirement has been waived. On the other hand, in some cases, studies not required by the Guidelines may be needed because of the particular composition or use pattern of products the Standard covers; if so, the Standard explains the Agency's reasoning. Data guidelines have not yet been proposed for the Residue Chemistry discipline, but the requirements for such data

have been in effect for some time and are, the Agency believes, relatively familiar to registrants. Data which the Agency has found needed to evaluate the registrability of some products covered by the Standard may not be needed for the evaluation of other products, depending upon the composition, formulation type, and intended uses of the product in question. The Standard states which data requirements apply to which product use categories (see Part III).

The various kinds of data normally required for registration of a pesticide product can be divided into two basic groups:

1) data that are "product-specific" i.e., data that relate only to the properties or effects of a product with a particular composition (or a group of products with closely similar composition); and

2) "generic" data that pertain to the properties or effects of a particular ingredient, and thus are relevant to an evaluation of the risks of all products containing that ingredient (or all such products having a certain use pattern), regardless of any such product's unique composition or use.

The Agency requires certain "product-specific" data for each product to characterize the product's particular composition and physical/chemical properties (Product Chemistry), and to characterize the product's acute toxicity (which is a function of its total composition). The applicant for registration or re-registration of any product, whether it is a manufacturing-use or end-use product, and without regard to its intended use pattern, must submit or cite enou ! of this kind of data to allow EPA to evaluate the product. For such purposes, "product-specific" data on any product other than the applicant's are irrelevant, unless the other product is similar in composition to the applicant's. (Where it has been found practicable to group similar products for purposes of evaluating, the Standard indicates which guideline requirement that support similar products.) "Product-specific" data on the efficacy of particular end-use products are also required where the formulation may affect public health or where failure of efficacy could cause public health problems (see 44 FR 27932, May 11, 1979).

All other data needed to evaluate pesticide products concern the properties or effects of a particular ingredient of products (normally a pesticidally active ingredient, but in some cases a pesticidally inactive, or "inert" ingredient). Some data in this "generic" category are required to evaluate the properties and effects of all products containing that ingredient e.g., the acute LD_{50} of the active ingredient in its technical or purest grade (see proposed 40 CFR 163.81-1(a), 43 FR 37355).

Other "generic" data are required to evaluate all products which both contain a particular ingredient and are intended for certain uses (see proposed 40 CFR 163.82-1, 43 FR 37363, which requires subchronic oral testing of the active ingredient with respect to certain use patterns only). Where a particular data requirement is use pattern dependent, it will apply to each end-use product which is to be labeled for that use pattern (except where such end-use product is formulated from a registered manufacturing-use product permitting such formulations) and to each manufacturing-use product with labeling that allows it to be used to make end-use products with that use pattern. Thus, for example, a subchronic oral dosing study is needed to evaluate the safety of any manufacturing-use product that legally could be used to make an end-use, food crop pesticide. However, if an end-use product's label specified that it was for use only in ways that did not involve a food/feed exposure and did not involve repeated human exposure via the oral route, then the subchronic oral study would not be relevant to the evaluation of the manufacturing-use product.

If a registrant of a currently registered manufacturing-use or end-use product wishes to avoid the costs of data compensation under FIFRA Section 3(c)(1)(D) or data generation under Section 3(c)(2)(B) for "generic" data that are required only with respect to some use patterns, he may elect to delete those use patterns from his labeling at the time he reregisters his products. An applicant for registration of a new product under this Standard may similarly request approval for only certain use patterns.

E. Data Compensation Requirements Under FIFRA 3(c)(1)(D)

Under FIFRA Section 3(c)(1)(D), an applicant for registration, reregistration or amended registration must offer to pay compensation for certain existing data the Agency has used in developing the Registration Standard. Compensation must be offered for all data which are described by all the following criteria:

1) the data were first submitted to EPA or to its predecessor agency, USDA, or FDA on or after January 1, 1970;

2) the data were submitted to EPA (or USDA or FDA) by some other applicant or registrant in support of an application for an experimental use permit or a tolerance, an amendment adding a new use to a registration, or for reregistration, or to support or maintain in effect an existing registration;

3) the data are relevant to the Agency's decision to register or reregister the applicant's product under the Registration Standard, taking into account the applicant's product composition and intended use pattern(s);

4) the data are determined by EPA to be valid and usable in reaching regulatory conclusions; and

5) the data are not those for which the applicant has been exempted by FIFRA Section 3(c)(2)(D) from the study to offer to pay compensation. (This exemption applies to the "generic" product, not to "product-specific" data. The exemption is available only to an applicant whose end-use product is formulated from another registered manufacturing-use product containing that active ingredient.)

An applicant for reregistration of an already registered product under this Standard, or for registration of a new product under this Standard, accordingly must determine which of the data used by EPA in developing the Standard must be the subject of an offer to pay compensation, and must submit with his application the appropriate statements evidencing his compliance with FIFRA Section 3(c)(1)(D).

An applicant would never be required to offer to pay for "product-specific" data submitted by another firm. In many, if not in most cases, data which are specific to another firm's product will not suffice to allow EPA to evaluate the applicant's product, that is, will not be useful to the Agency in determining whether the applicant's product is registrable. There may be cases, however, where, because of close similarities between the composition of two or more products, another firm's data may suffice to allow EPA to evaluate some or all of the "product-specific" aspects of the applicant's product. In such a case, the applicant may choose to cite that data instead of submitting data from tests on his own product, and if he chooses that option, he would have to comply with the offer-to-pay requirements of Section 3(c)(1)(D) for that data.

Each applicant for registration or reregistration of a manufacturing-use product, and each applicant for registration or reregistration of an end-use product, who is not exempted by FIFRA Section 3(c)(2)(D), must comply with the Section 3(c)(1)(D) requirements. This is with respect to each item of "generic" data that relates to his product's intended uses.

A detailed description of the procedures an applicant must follow in applying for reregistration (or new registration) under this Standard is found in the Guidance Package for this Standard.

F. Obtaining Data to Fill Data Gaps; FIFRA 3(c)(2)(B)

Some of the kinds of data EPA needs for its evaluation of the properties and effects of products to which this Standard applies have not been submitted to the Agency (or, if submitted, have been found to have deficiencies rendering them inadequate for making registration oriented decisions) and have not been located in the published literature search that EPA conducted in preparation of this Standard. Such instances of missing but required data are referred to in this Standard as "data gaps".

FIFRA Section 3(c)(2)(B), added to FIFRA by the Congress in 1978, authorizes EPA to require registrants to whom a data

requirement applies, to generate (or otherwise produce) data to fill such "gaps" and submit those data to EPA. EPA must allow a reasonably sufficient period for this to be accomplished. If a registrant fails to take appropriate and timely steps to fill the data gaps identified by a Section 3(c)(2)(B) order, his product registration may be suspended until the data are submitted. A mechanism is provided whereby two or more registrants may agree to share in the costs of producing data for which they are both responsible.

The Standard lists, in Part III, the "generic" data gaps and "product-specific" data gaps for manufacturing- and end-use products. It also notes the classes of products to which these data gaps pertain. The Standard also points out that to be registered under the Standard, a product must be supported by certain required "product-specific" data on one currently registered product, but may lack such data on another. Only those Standards which apply to a very small number of currently registered products will attempt to state definitively the "product-specific" data gaps on a "product-by-product" basis. (Although the Standard will, in some cases, note which data EPA does possess that would suffice to satisfy certain "product-specific" data requirements for a category of products with similar composition characteristics.)

As part of the process of reregistering currently registered products, EPA will issue FIFRA Section 3(c)(2)(B) directives requiring the registrants to take appropriate steps to fill all identified data gaps whether those data in question are "product-specific" or "generic" data gaps. The Guidance Package for this Standard details the steps that must be taken by registrants to comply with Section 3(c)(2)(B).

In addition, FIFRA Section 6(a)(2) requires the registrant to submit factual information raising concerns of possible unreasonable adverse effects of a pesticide. The registrant should notify the Agency of interim results of studies in progress if those results show possible adverse effects.

G. Amendments to the Standard

Applications for registration which propose uses or formulations that are not presently covered by the Standard, or which present product compositions, hazard data, toxicity levels, or labeling that do not meet the requirements of the Standard, will automatically be considered by the Agency to be requests for amendments to the Standard. In response to such applications, the Agency may request additional data to support the proposed amendment to the Standard, or may deny the application for registration on the grounds that the proposed product would cause unreasonable adverse effects to the environment. In the former case, when additional data have been satisfactorily supplied, and providing that the data do not indicate the potential for unreasonable adverse effects, the Agency will then amend the Standard to cover the new registration.

Each Registration Standard is based upon all data and information available to the Agency's reviewers on a particular date prior to the publication date. This "cut-off" date is stated at the beginning of the Part II.C. Any subsequent data submissions and any approved amendments will be incorporated into the Registration Standard by means of addenda, which are available for inspection at EPA in Washington, D.C., or copies may be requested from the Agency. When all of the present "data gaps" have been filled and the submitted data have been reviewed, the Agency will revise the Registration Standard. Thereafter, when the Agency determines that the internally maintained addenda have significantly altered the conditions for registration under the Standard, the document will be updated and reissued for publication.

While the Registration Standard discusses only the uses and hazards of products containing the designated active ingredient(s), the Agency is also concerned with the potential hazards of some inert ingredients and impurities. Independent of the development of any one Standard, the Agency has initiated the evaluation of some inert pesticide ingredients. Where the Agency has identifiedinert ingredients of concern in a specific product to which the Standard applies, these ingredients will be indicated in the Guidance Package.

II. Regulatory Position

A. Introduction

This part presents the Agency's regulatory position and rationale based on an evaluation of all registered products containing temephos as the sole active ingredient with the same use patterns described in this Standard. After briefly describing the chemical, this part presents the regulatory position and rationale, the criteria by which applicants for registration of temephos products will be approved, labeling considerations and requirements related to the tolerance reassessment, if applicable. A summary of the data requirements is contained in Part III. Discussion of the data upon which this regulatory position is based is presented in each of the disciplinary chapters, Parts IV through IX.

B. Description of Chemical

Temephos (O,O'-(thiodi-4,1-phenylene)bis(O,O-dimethyl phosphorothioate) is used primarily as a mosquito larvicide. An alternative chemical name for the compound is O,O,O',O'-tetramethyl O,O'-thiodi-p-phenylene phosphorothioate. Temephos is also known by the trade names "Abate®", "Abathion®", "Abat®", and "Biothion®." The Chemical Abstracts Registry (CAS) number for temephos is 3383-96-8 and the EPA Shaughnessy number is 059001.

C. Regulatory Position

Temephos, as described in this Standard, may be registered for sale, distribution, reformulation and use in the United States. The Agency has considered the limited amount of scientific data obtained from the open literature as of August, 1981, and the data submitted to the Agency by the registrants through the time of publication of this Standard. Based on review of these limited data, the Agency finds that none of the risk criteria found in Section 162.11(a) of Title 40 of the U.S. Code of Federal Regulations (CFR) was met or exceeded for temephos and that temephos does not appear to cause unreasonable adverse effects when applied in accordance with proper label directions and precautions. Temephos products currently registered may be reregistered subject to the conditions imposed for data requirements. New products may be registered under this Standard and are subject to the same requirements. Revised label requirements will be addressed in the Guidance Package which accompanies this Standard and in Part II.E of this Standard (see 40 CFR 162.10 for a detailed description of standard labeling requirements).

D. Regulatory Rationale

A review of the available data regarding the manufacturing-use and end-use products of temephos shows that much information that could be used to support registration and reregistration is not available. The Agency has the prerogative to not only request information to satisfy the data Guidelines proposed in

1978 and 1980, but may also require additional testing. These additional tests may be requested by the Agency if it believes that particular concerns regarding the health or environmental effects of a pesticide must be evaluated. In some cases, this additional testing may be deferred pending the Agency's development of specific test protocols or the completion of other related tests (e.g., tiered testing).

It is the policy of the Agency not to automatically request guideline information just because a particular guideline exists. Each element of the guideline is considered individually with regard to the pesticide's uses, exposure and risk. If the Agency concludes that specific elements of the data guidelines are required to prepare a comprehensive Registration Standard, the data will be requested.

The Agency has concluded that it should continue registration for this product for the following reasons:

1) No significant adverse effects have been uncovered in the review of studies pertinent to temephos. These studies include some which meet guideline data requirements and some which, while failing to meet guideline requirements, provide a level of qualitative information. As a result of examining these studies, the Agency has concluded that there is no immediate cause for regulatory concern.

2) There is evidence that technical grade temephos causes only a low level of acute oral and dermal health effects. The potential for eye irritation is minimal and no adverse effects have been identified from analysis of the chronic feeding study. Furthermore, the results of one study evaluating the oncogenicity of temephos are negative.

3) The Pesticide Incident Monitoring System's Report #125 of February 1979, reports only three accidents with temephos. In one instance, 20 cattle died in a field sprayed with temephos at twice the maximum registered rate. Investigation did not produce sufficient evidence to link the registered use of the pesticide to the incident. In the second event, damaged containers of temephos contaminated the interior of a truck/trailer carrying the pesticide. In the third incident, a drum of temephos exploded and burned while being heated with a heating strap during the manufacturing of end-use products. In all these incidents there is no record of human injury or deleterious effects to the human system that could be attributed to the health and environmental effects of temephos, or effects to animals other than by misuse.

   Of these three events, none involve the pesticidal nature of temephos and the level of accidents reported involving the registered temephos use patterns are generally very low. However, due to the use of temephos in isolated aquatic areas, the Agency realizes that the incidence of accidents may not be indicative of potential hazard.

4) Based on 1976 data, it is estimated that the total number of ground applicators potentially exposed to temephos approximates less than 500; the number of applicators, mixers, loaders and flaggers potentially exposed as a result of aerial application also is less than 500*. The majority of temephos used in mosquito abatement programs is applied aerially as emulsifiable concentrate or as granular formulations. During aerial application, the pilot is usually not exposed to the spray or granules, the aircraft cockpits are usually sealed and are ventilated with a positive air flow system, and the pesticide delivery system is closed. Consequently, the number of persons potentially exposed to temephos during the application process is minimal, especially during aerial application where the potential for human exposure appears slight. Exposure to the general population is also considered minimal because temephos is used primarily in marshlands, swamps and tidal areas where human population density is generally low.

5) In accordance with FIFRA, the Agency's policy is not to routinely cancel the registration of products or to withhold registration merely for the lack of data (see Sections 3(c)(2)(B) and 3(c)(7) of FIFRA). Rather, the publication of this Standard provides a mechanism for identifying data needs, and registration of temephos under this Standard allows for the upgrading of labels during the period in which the required data are being generated. When these data are received, they will be reviewed by the Agency and the registerability of the chemical will be reassessed.

E. Criteria for Registration Under the Standard

To be subject to this Standard, temephos products must meet the following conditions:

- contain temephos as the sole active ingredient** and have the same use patterns described in this Standard;

- bear required labeling; and

---

* Geomet Technologies, Rockville, Md., Contract No. 68015155 September 4, 1980.

** Note: Since petroleum distillates have little or no insecticidal activity in temephos products (e.g., emulsifiable concentrates), the Agency has determined that petroleum distillates should be considered as part of the inert ingredients. If the registrant wishes to declare petroleum distillates as active ingredients, supporting data must accompany the application for reregistration.

- conform to the acute toxicity limits, product composition, and use pattern requirements stated below.

The applicant for registration or reregistration of temephos products subject to this Standard must comply with all terms and conditions described in this Standard, including a commitment to fill data gaps on a time schedule specified by the Agency as indicated in Part III and, when applicable, offer to pay compensation to the extent required by Sections 3(c)(1)(D) and 3(c)(2)(D) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, 7 U.S.C. 136(c)(1)(D) and 136 (c)(2)(D). As discussed in Part I and in the Guidance Package, applicants for registration of temephos products under this Standard must contact the Agency for specific instructions, including updated information on data requirements, and companies whose data may be cited and to whom compensation must be offered.

1. Manufacturing-Use Temephos

a. Acceptable Ranges and Limits

i. Product Composition Standard

To be covered under this Standard, manufacturing-use temephos products must conform to the requirements stated below:

Active Ingredient: Any percentage acceptable.

ii. Acute Toxicity Limits

The Agency will consider registration of manufacturing-use temephos products in the toxicity categories indicated by a "yes":

	I	II	III	IV
Acute oral toxicity	yes	yes	yes	yes
Acute dermal toxicity	yes	yes	yes	yes
Acute inhalation toxicity	yes	yes	yes	yes
Primary eye irritation	yes	yes	yes	yes
Primary dermal irritation	yes	yes	yes	yes

iii. Use Patterns

To be covered under this Standard, manufacturing-use temephos products must be labeled to allow for formulation into end-use insecticides which are intended only for:

1) outdoor, nondomestic, nonfood aquatic areas;
2) domestic use on cats and dogs, and in their living quarters;
3) domestic flea collars for cats and dogs; and
4) outdoor, nondomestic, food use on orange and tangerine trees (California and Arizona only).

iv. Required labeling

All manufacturing-use temephos products must bear appropriate labeling as specified in 40 CFR 162.10. The Agency may, after review of data to be submitted under this Standard, impose additional label requirements.

Efficacy Statements

All manufacturing-use temephos products must list on the label the intended end-uses of formulated products produced from the manufacturing-use products. In accordance with data to be submitted or cited, all temephos product labels must bear one or a combination of the following statements:

1) "For Formulation into End-Use Insecticide Products Intended Only as an Insecticide for Nondomestic, Nonfood, Outdoor Aquatic Uses";

2) "For Formulation into End-Use Insecticide Products Intended Only as an Insecticide for Nondomestic, Outdoor, Food Use";

3) "For Formulation into End-Use Insecticide Products Intended for Cat and Dog Flea Collars"; and/or

4) "For Formulation into End-Use Insecticide Products Intended for Direct Application on Cats and Dogs, and in their Living Quarters."

Ecological Effects Statements

The following statements must appear under the heading "Environmental Hazards" [see Section 162.10(h)(2(ii) of the guidelines]:

> This pesticide is toxic to birds and fish. Do not discharge directly into lakes, streams, ponds, or public waters unless in accordance with an NPDES permit. For guidance, contact your Regional Office of the Environmental Protection Agency.

Additional fish and wildlife labeling may be required upon review of the requested studies (see Tables III.A-5 in Part III).

Product Chemistry Statements

The ingredient statement for manufacturing-use products must list the active ingredient as:

Temephos [O,O'-(thiodi-4,1-phenylene)bis(O,O-dimethyl phosphorothioate)].....N% (min.)

There is currently no special hazard precautionary labeling for the manufacturing-use products. However, as required physical/chemical property data gaps are filled, the Agency may impose additional labeling requirements.

2. End-Use Temephos: Emulsifiable Concentrates

a. Acceptable Ranges and Limits

i. Product Composition Standard

To be covered under this Standard, emulsifiable concentrate products must conform to the requirements stated below:

Active ingredient: Any percentage acceptable.

Inert ingredient: Only those inert ingredients which are exempt from the requirements of a tolerance (see 40 CFR Section 180.1001) are acceptable for use on food crops. There are no restrictions on inert ingredients when used for nonfood purposes.

ii. Acute Toxicity Limits

The Agency will consider registration of emulsifiable concentrate temephos products in the toxicity categories indicated by "yes". The Agency will consider registration of products in toxicity category I when all of the acute toxicity data and other required data appropriate for an amended registration are satisfied. After receipt and review of these data, the Agency may require other appropriate data (e.g., special or subchronic

studies), impose additional labeling restrictions, or deny registration.

	I	II	III	IV
Acute oral toxicity	amendment required	yes	yes	yes
Acute dermal toxicity	amendment required	yes	yes	yes
Acute inhalation toxicity	amendment required	yes	yes	yes
Primary eye irritation	amendment required	yes	yes	yes
Primary dermal irritation	amendment required	yes	yes	yes

iii. Use Patterns

To be registered under this Standard, emulsifiable concentrate temephos products must be labeled as an insecticide for non-domestic use for one or more of the following uses: 1) food uses, orange and tangerine trees (California and Arizona only); and 2) nonfood crop aquatic uses, intermittently flooded areas, temporary rain pools, catch basins, standing water, woodland pools, shallow lakes and lake margins, log ponds, sloughs, stagnant water, marshes, tidal water, and swamps.

iv. Required labeling

All end-use emulsifiable concentrate temephos products must bear appropriate labeling as specified in 40 CFR 162.10. The Agency may, after review of data to be submitted under this Standard, impose additional label requirements.

Product Chemistry Statements

Th ingredient statement for end-use emulsifiable concentrate products must list the active ingredient as:

Temephos [O,O'-(thiodi-4,1-phenylene)bis(O,O-dimethyl phosphorothioate)].....N% (min.)

There is currently no special hazard precautionary labeling for the end-use emulsifiable concentrate products. However, as required physical/chemical property data gaps are filled, the Agency may impose additional labeling requirements.

Efficacy Statements

Appropriate product dilution rates for application [e.g., milliliters product (liquid) per square feet] must be provided for spot treatment of areas such as catch basins, containers of

water (e.g., old tires), and other water-filled sunken areas where mosquitoes may breed.

The following statement must appear on all temephos emulsifiable concentrates under the heading "Directions for Use":

> The higher rate should be used in areas known to have organic phosphate resistant mosquitoes.

Ecological Effects Statements

For those emulsifiable concentrates registered for use on tangerine and orange trees (California and Arizona only), the following labeling must appear under the heading "Environmental Hazards" [see Section 162.10(h)(2)(ii) of the guidelines]:

> This pesticide is toxic to birds and fish. Use with care when applying in areas frequented by wildlife or adjacent to any body of water. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area. Do not contaminate water by cleaning of equipment or disposing of wastes.

Under "Directions for Use" for the emulsifiable concentrate for aquatic uses, the following labeling must be included:

> When using in lakes, ponds, tidal marshes, and estuaries, apply only in shallow water at the edge of these areas.

For those emulsifiable concentrates registered for aquatic uses, the following statements must be included on the labeling and appear under the headings "Environmental Hazards":

> This pesticide is toxic to birds and fish. Fish and other aquatic organisms in waters treated with this product may be killed. You must consult your State Fish and Game Agency before applying this product to waters or wetlands. Do not contaminate water by cleaning of equipment or disposing of wastes.

Additional fish and wildlife statements may be required upon review of the requested studies (see Table III.A-5 in Part III).

Residue Chemistry Statements

The labeling for formulations used on citrus bears two restrictions: 1) "on bearing trees the product is to be applied at prebloom or at petal fall only after harvest of mature fruit. This product should not be applied to any varieties of citrus when immature fruit are present during prebloom and petal fall. Do not apply during bloom period." and 2) "use restricted to Arizona and California". These restrictions must continue to appear on the labeling for the food crop use.

In lieu of submitting required data on residues in cover crops in temephos-treated citrus orchards, the following statement must appear on the labeling:

> Do not feed or allow livestock to graze on cover crops grown in treated orchards.

Products for use on aquatic sites (e.g., catch basins, shallow lakes and lake margins, sloughs, marshes, tidal water, swamps) must continue to bear the restriction against use of the product on crops for food, forage, or pasture. Because of the large number of aquatic sites on which temephos may be used, residues may appear in potable water, fish, irrigated crops and livestock. Upon receipt and evaluation of data on the degradation of temephos in water and fish, the following labeling statements may be required:

> Do not apply to the following aquatic sites: 1) potable water, 2) water used for irrigating crops, 3) water used for watering livestock, and 4) water containing edible fish. Do not drain treated water into a flowing stream that is a potential source of water for one or more of the above uses.

In lieu of labeling restrictions, appropriate tolerances for potable water, fish, shellfish, irrigated crops, meat, milk, poultry and eggs may be necessary. Other labeling restrictions may be imposed as the required residue data gaps are filled.

3. End-Use Temephos: Granular

a. Acceptable Ranges and Limits

i. Product Composition Standard

To be covered under this Standard, granular temephos products must conform to the requirements stated below:

Active ingredient: Any percentage acceptable.

Inert ingredient: There are no restrictions on inert ingredients when used for nonfood purposes.

ii. Acute Toxicity Limits

The Agency will consider registration of granular temephos products in the toxicity categories indicated by "yes". The Agency will consider registrations of products in toxicity category I when all of the acute toxicity data and other required data appropriate for an amended registration are satisfied. After receipt and review of these data, the Agency may require other appropriate data (e.g., special or subchronic

studies), impose additional labeling restrictions, or deny registration.

	I	II	III	IV
Acute oral toxicity	amendment required	yes	yes	yes
Acute dermal toxicity	amendment required	yes	yes	yes
Acute inhalation toxicity	amendment required	yes	yes	yes
Primary eye irritation	amendment required	yes	yes	yes
Primary dermal irritation	amendment required	yes	yes	yes

iii. Use Patterns

To be registered under this Standard, granular temephos products must be labeled as an insecticide for one or more of the following nonfood crop aquatic uses: sloughs, stagnant water, log ponds, temporary rain pools, intermittently flooded areas, catch basins, lakes, standing water, shallow ponds, woodland pools, tidal water, marshes, swamps, water high in organic matter, highly polluted water, marshlands and intertidal zones of sandy beaches.

iv. Required Labeling

All granular temephos products must bear appropriate labeling as specified in 40 CFR 161.10. The Agency may, after review of data to be submitted under this Standard, impose additional label requirements.

Product Chemistry Statements

The ingredient statement for end-use granular products must list the active ingredient as:

Temephos [O,O'-(thiodi-4,1-phenylene)bis(O,O-dimethyl phosphorothioate)].....N% (min.)

There is currently no special hazard precautionary labeling for the end-use granular products. However, as required physical/chemical property data gaps are filled, the Agency may impose additional labeling requirements.

Efficacy Statements

Appropriate product dilution rates for application [e.g., grams product (weight) per square feet] must be provided for spot treatments of areas such as catch basins, containers of water

(e.g., old tires), and other water-filled sunken areas where mosquitoes may breed.

The following statement must appear on all temephos granular products under the heading "Directions for Use":

> The higher rate should be used in areas known to have organic phosphate resistant mosquitoes.

Ecological Effects Statements

The following statements must appear on granular temephos products for aquatic uses under the heading "Environmental Hazards":

> This pesticide is toxic to birds and fish. Fish and other aquatic organisms in water treated with this product may be killed. You must consult your State Fish and Game Agency before applying this product to waters or wetlands. Do not contaminate water by cleaning of equipment or disposing of wastes.

Additional fish and wildlife labeling may be required when the Agency reviews the requested studies (see Table III.A-5 in Part III).

Residue Chemistry Statements

Products for use on aquatic sites (e.g., sloughs, catch basins, lakes, shallow ponds, tidal water, marshes, swamps, water high in organic content, highly polluted water, marshlands, intertidal zones of sandy beaches) must continue to bear the restriction against use of the product on crops for food, forage, or pasture. Because of the large number of aquatic sites on which temephos may be used, residues may appear in potable water, fish, irrigated crops and livestock. Upon receipt and evaluation of data on the degradation of temephos in water and fish, the following labeling statements may be required:

> Do not apply to the following aquatic sites: 1) potable water, 2) water used for irrigating crops, 3) water used for watering livestock, and 4) water containing edible fish. Do not drain treated water into a flowing stream that is a potential source of water for one or more of the above uses.

In lieu of labeling restrictions, appropriate tolerances for potable water, fish, shellfish, irrigated crops, meat, milk, poultry and eggs may be necessary. Other labeling restrictions may be imposed as the required residue data gaps are filled.

4. End-Use Temephos: Dusts

a. Acceptable Ranges and Limits

i. Product Composition Standard

To be covered under this Standard temephos dust products must conform to the requirements stated below:

Active ingredient: Any percentage acceptable.

Inert ingredients: There are no restrictions on inert ingredients when used for nonfood purposes.

ii. Acute Toxicity Limits

The Agency will consider registration of temephos dust products in the toxicity categories indicated by "yes". The Agency will consider registration of products in toxicity categories I and II when all of the acute toxicity data and other required data appropriate for an amended registration are satisfied. After receipt and review of these data, the Agency may require other appropriate data (e.g., special or subchronic studies), impose additional labeling restrictions, or deny registration.

	I	II	III	IV
Acute oral toxicity	amendment required	amendment required	yes	yes
Acute dermal toxicity	amendment required	amendment required	yes	yes
Acute inhalation toxicity	amendment required	amendment required	yes	yes
Primary eye irritation	amendment required	amendment required	yes	yes
Primary dermal irritation	amendment required	amendment required	yes	yes

iii. Use Patterns

To be registered under this Standard, temephos dust products must be labeled as an insecticide for the following domestic uses: direct application on dogs and cats, and in their living quarters.

iv. Required Labeling

All temephos dust products must bear appropriate labeling as specified in 40 CFR 162.10. The Agency may, after review of data to be submitted under this Standard, impose additional label requirements.

Product Chemistry Statements

The ingredient statement for end-use dust products must list the active ingredient as:

Temephos [O,O'-(thiodi-4,1-phenylene)bis(O,O-dimethyl phosphorothioate)].....N% (min.)

There is currently no special hazard precautionary labeling for the end-use dust products. However, as required physical/chemical property data gaps are filled, the Agency may impose additional labeling requirements.

5. End-Use Temephos: Impregnated Material

a. Acceptable Ranges and Limits

i. Product Composition Standard

To be covered under this Standard, impregnated temephos products must conform to the requirements stated below:

Active ingredient: Any percentage acceptable.

Inert ingredients: There are no restrictions on inert ingredients used for nonfood purposes.

ii. Acute Toxicity Limits

The Agency will consider registration of impregnated temephos products in the toxicity categories indicated by "yes". The Agency will consider registration of products in toxicity categories I and II when all of the acute toxicity data and other required data appropriate for an amended registration are satisfied. After receipt and review of these data, the Agency may require other appropriate data (e.g., special or subchronic

studies), impose additional labeling restrictions, or deny registration.

	I	II	III	IV
Acute oral toxicity	amendment required	amendment required	yes	yes
Acute dermal toxicity	amendment required	amendment required	yes	yes
Acute inhalation toxicity	amendment required	amendment required	yes	yes
Primary eye irritation	amendment required	amendment required	yes	yes
Primary dermal irritation	amendment required	amendment required	yes	yes

iii. Use Patterns

To be registered under this Standard, a product impregnated with temephos must be labeled as an insecticide for the following domestic uses: cat and dog flea collars.

iv. Required Labeling

All temephos-impregnated cat and dog collar products must bear appropriate labeling as specified in 40 CFR 162.10. The Agency may, after review of data to be submitted under this Standard, impose additional label requirements.

Product Chemistry Statements

The ingredient statement for end-use impregnated material products must list the active ingredient as:

Temephos [O,O'-(thiodi-4,1-phenylene)bis(O,O-dimethyl phosphorothioate)].....N% (min.)

There is currently no special hazard precautionary labeling for the end-use impregnated material products. However, as required physical/chemical property data gaps are filled, the Agency may impose additional labeling requirements.

F. Tolerance Reassessment

To provide for the use of temephos on orange and tangerine trees in Arizona and California, a tolerance of 0.1 ppm has been established for combined negligible residues of temephos and its sulfoxide in or on the raw agricultural commodity, citrus fruits (40 CFR 180.170). Based on the established tolerance, the theoretical maximum residue contribution (TMRC) of temephos to the human diet is calculated to be 0.0057 mg/1.5 kg diet/day. However, because there is some uncertainty about the nature of

the residue on temephos-treated citrus, the tolerance from which the TMRC is calculated may not be adequate (the current method for detecting temephos residues does not detect residues other than temephos and its sulfoxide). A tolerance reassessment is consequently not possible at this time (August, 1981). A reassessment will be performed after the Agency receives and reviews the data listed in Table III.A-4, Residue Chemistry (Part VII) and Table III.A-3, Toxicology (Part VI)*.

Codex Maximum Residue Limits (MRL's) have not been established for temephos. Currently, there are no Canadian or Mexican tolerances for temephos.

---

* The Agency requires the submittal of these data (i.e., Residue Chemistry and Toxicology) to maintain the tolerance which provides for temephos use on imported citrus as well as on citrus growing in the United States. If these data (see Part III, Table III.A, e.g., residue, toxicity) are not provided to the Agency to evaluate the possible hazards to man and other nontarget species from the potential effects of its use, the Agency will propose revocation of the tolerance under the Federal Insecticide, Fungicide and Rodenticide Act, 40 CFR Section 180.001 and the Food and Drug Cosmetic Act, 21 CFR Section 408(e).

III. Summary of Data Requirements

Applicants for registration of manufacturing-use and end-use temephos products must cite or submit the information identified as required in the tables in this part. The tables applicable to end-use products indicate whether the product to be tested is the technical grade or the formulation. Data generated on one formulation may be used to satisfy the data requirement for a substantially similar formulation based on the chemical composition as indicated by the Confidential Statement of Formula (CSF). Information on which product-specific data requirements are already satisfied is available in the Guidance Package and in Tables III.A through C in this Standard.

Preceeding each requirement, with the exception of Residue Chemistry and Efficacy*, is listed the Proposed Guidelines which describe the type of data and when the data are required to be submitted. Justification for the test requirement is provided in the Guidelines (see 43 FR 29696 of July 10, 1978; 43 FR 37336 of August 22, 1978; and 45 FR 72948 of November 3, 1980). A discussion of why data additional to that already submitted are necessary, or why data usually required are not necessary for this chemical, are explained in footnotes to the tables. The specified data requirements are the minimum that will be required. Areas where additional data may be required as the result of tiered testing are indicated (see tables for Toxicology, Residue Chemistry, and Ecological Effects in this part).

Bibliographic citations in the following tables indicate that the listed study accomplishes one of the following:

1) provides all the information required by the guideline requirement;

2) provides partial information required by the guideline requirement; and

3) provides specific information that is elaborated on by an accompanying footnote.

Bibliographic citations noted in Parts IV through IX which are not listed in this part are provided as reference and background material to the reader; they do not relate directly to or satisfy the guideline requirements.

The word "yes" under the heading "Does EPA have data to partially or totally satisfy this requirement?" means that the requirement has been satisfied, unless otherwise noted.

---

* There are currently no Proposed Guidelines for Efficacy. Refer to Part I.D for discussion of registrants' familiarity of data requirements for Residue Chemistry.

In some cases the data requirements are imposed by the use patterns and the use patterns appear in the tables under the heading "Are data required by this Standard?" (see Tables III.A-2, 3, 4, and 5). Where only the word "yes" appears under this heading, the data requirements are required for all currently registered products, regardless of use pattern.

A. Table III.A Generic Data Requirements

This table includes those data that pertain to the properties or effects of temephos as an active ingredient. Thus, these data are relevant to an evaluation of the risks of all products containing temephos. Providing data to fill indicated gaps in the data base is the primary responsibility of the registrant(s) of manufacturing-use temephos. Registrants of end-use products which are not exempted by FIFRA Section 3(c)(2)(D) are also responsible for the submission of these data.

However, applicants for the registration of end-use products containing temephos are advised that if the Agency does not receive commitments within the time specified in the tables from manufacturing-use temephos registrant(s) to fill data gaps identified for the manufacturing-use product (Table III.A), manufacturing-use product registrations will be suspended. Formulators must then bear the burden of supplying these data in both Table III.A and B if continued registration of their product is desired.

Applicants for the registration or reregistration of manufacturing-use temephos products must acknowledge reliance on existing data which fill indicated data requirements under FIFRA Section 3(c)(1)(D). These data are listed under the column "Bibliographic Citation" in Table III.A.

B. Table III.B Product-Specific Data Requirements for Manufacturing-Use Products

This table includes those data that relate only to the properties or effects of a product with a specific composition (or substantially similar composition). Thus, these data are required of each product (or substantially similar product) to characterize the product's particular composition and physical/chemical properties, and to characterize the product's acute toxicity.

Product compostion data, physical/chemical properties and acute toxicity are required for each manufacturing-use product. Providing data to fulfill these requirements is the responsibility of each applicant for the registration or reregistration of a manufacturing-use temephos product. If the Agency has data which fulfills this requirement for a particular product(s), this is indicated in the table and in the Guidance Package accompanying this Standard.

Product-specific data need not be acknowledged under FIFRA Section 3(c)(1)(D) unless the Agency has established that one

product is substantially similar to another product for which the Agency has received acceptable data.

C. Table III.C Product-Specific Data Requirements for End-Use Products

This table includes those data that relate only to the properties or effects of a product with a specific composition (or substantially similar composition). Thus, these data are required of each product (or substantially similar product) to characterize the product's particular composition and physical/chemical properties, and to characterize the product's acute toxicity.

Product compostion data, physical/chemical properties and acute toxicity are required for each end-use product. Providing data to fultill these requirements is the responsibility of each applicant for the registration or reregistration of an end-use temephos product. If the Agency has data which fulfills this requirement for a particular product(s), then this is indicated in the table and in the Guidance Package accompanying this Standard.

Product-specific data may be acknowledged under FIFRA Section 3(c)(1)(D) only if the Agency has established that one product is substantially similar to another product for which the Agency has received acceptable data.

Table III.A-1 Generic Data Requirements for Temephos, Product Chemistry, (see Part IV)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.61-3	Product Identity and Disclosure of Ingredients	yes	Technical grade of active ingredient	partial[1/]	000003268, 000001216	yes/8 months
163.61-8(1)	Color	yes	Technical grade of active ingredient	yes	000001379	no
163.61-8(2)	Odor	yes	Technical grade of active ingredient	no	-	yes/8 months
163.61-8(3)	Melting Point	yes	Technical grade of active ingredient	yes	000001379	no
163.61-8(4)	Solubility	yes	Technical grade of active ingredient	partial[2/]	000001379	yes/8 months
163.61-8(5)	Stability	yes	Technical grade of active ingredient	yes	000001379	no
163.61-8(6)	Octanol/Water partition coefficient	yes	Techncial grade of active ingredient	no	-	yes/8 months

August, 1981

Table III.A-1 Generic Data Requirements for Temephos, Product Chemistry (see Part IV) (cont'd)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.61-8(7)	Physical State	yes	Technical grade of active ingredient	yes	000003264	no
163.61-8(8)	Density or Specific Gravity	yes	Technical grade of active ingredient	yes	000001379	no
163.61-8(9)	Boiling Point	yes	Technical grade of active ingredient	no	-	yes/8 months
163.61-8(10)	Vapor Pressure	yes	Technical grade of active ingredient	no	-	yes/8 months
163.61-8(11)	pH	yes	Technical grade of active ingredient	no	-	yes/8 months

1/ The impurities exceeding 0.1% of the technical temephos product weight must be determined.

2/ A numerical value for solubility of temephos in organic solvents must be determined.

August, 1981

Table III.A-2 Generic Data Requirements for Temephos, Environmental Fate (see Part V)

Guidelines Citation	Name of Test	Are Data Required for this Standard?		Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.62-7(b)	Hydrolysis		yes	Radiolabeled analytical grade or technical grade of active ingredient	no	-	yes/8 months
163.62-7(c)	Photodegradation	Food crops Aquatic	yes yes	Radiolabeled analytical grade or technical grade of active ingredient	no	-	yes/12 months
163.62-8(b)	Aerobic soil metabolism	Food crops	yes	Radiolabeled analytical grade or technical grade of active ingredient	no	-	yes/36 months
163.62-8(d)	Anaerobic aquatic metabolism	Aquatic	yes	Radiolabeled analytical grade or technical grade of active ingredient	no	-	yes/18 months
163.62-8(e)	Aerobic aquatic metabolism	Aquatic	yes	Radiolabeled analytical grade or technical grade of active ingredient	no	-	yes/18 months
163.62-8(f)	Microbial metabolism (2) effects of microbes on pesticides	Food crops Aquatic	yes yes	1/	no	-	1/

August, 1981

Table III.A-2 Generic Data Requirements for Temephos, Environmental Fate (see Part V) (cont'd)

Guidelines Citation	Name of Test	Are Data Required for this Standard?		Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.62-8(f)	Microbial metabolism (cont'd)					-	
	(3) effects of pesticides on microbes	Food crops Aquatic	yes yes	1/	no	-	1/
163.62-8(g)	Activated sludge metabolism		yes	1/	no	-	1/
163.62-9(b)	Leaching 2/	Food crops	yes	Radiolabeled analytical grade or technical grade of active ingredient	no	-	yes/12 months
163.62-9(d)	Adsorption/ 3/ desorption	Food crops Aquatic	yes yes	Radiolabeled analytical grade or technical grade of active ingredient	no	-	yes/12 months
163.82-9(e)	Water dispersal	Aquatic	yes	Representative formulated product 4/	no	-	yes/14 months
163.62-10(b)	Terrestrial field dissipation						
	(2) Tree fruit and nut crop uses	Food crops	yes	Representative formulated emulsifiable concentrate (43-60%) made from each MUP (preferably the highest concentrated (ai) product	no	-	yes/28 months

Table III.A-2 Generic Data Requirements for Temephos, Environmental Fate (see Part V) (cont'd)

Guidelines Citation	Name of Test	Are Data Required for this Standard?		Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.62-10(c)	Aquatic field dissipation (2) Aquatic noncrop aquatic	Aquatic	yes	Representative formulated product 4/	no	-	yes/14 months
163.62-10(d)	Terrestrial/ aquatic (forest) field dissipation	Aquatic	yes	Representative formulated product 4/	no	-	yes/14 months
163.62-10(e)	Aquatic impact uses						
	(1) Direct discharge	Aquatic	5/				
	(2) Indirect discharge	Aquatic	6/				
163.62-11(c)	Accumulation in irrigated crops	Aquatic	yes	Representative formulated product 4/	no	-	yes/14 months
163.62-11(d)	Fish accumulation	Food crops Aquatic	yes yes	Radiolabeled analytical grade or technical grade of active ingredient	partial 7/	005000978	yes/12 months
163.62-11(e)	Special studies; accumulation in aquatic noncrop uses	Aquatic	yes	Technical grade of active ingredient	no	-	yes/18 months

August, 1981

Table III.A-2 Generic Data Requirementsfor Temephos, Environmental Fate (see Part V) (cont'd)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.62-13	Disposal and storage	yes	Technical grade of active ingredient and all formulations	no	-	1/

1/ The requirement for submission of these data is currently being reserved pending the review and modification of testing protocols.

2/ For terrestrial noncrop uses, orchard crop uses, field vegetable crop uses, and forestry uses, the mobility of the test substance and its degradates in soil shall be assessed either by soil thin-layer chromatography, soil column, or batch equilibrium (adsorption/desorption) procedures.

3/ For domestic outdoor uses, greenhouse uses, aquatic uses, and aquatic impact-direct discharge uses, the mobility of the test substance and its degradates in soil shall be assessed only by the batch equilibrium (adsorption/desorption) procedure.

4/ Any currently registered formulated emulsifiable concentrate which contains 43-60% temephos and any formulated granular containing 1%, 2% or 5% temephos made from each manufacturing-use product (MUP).

5/ Data necessary to meet this requirement will be provided by data from 163.62-7(b) and (c); 163.62-8(d), (e), (f), and (g); 163.62-9(d); 163.62-10(c); and 163.62-11(d).

6/ Data necessary to meet this requirement will be provided by data from 163.62-7(b) and 163.62-8(g).

7/ Additional studies are required to measure the effects of temephos on fish in a flow through environment.

August, 1981

Table III.A-3 Generic Data Requirements for Temephos, Toxicology (see Part VI)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.81-1	Acute Oral Toxicity	yes	Technical grade of active ingredient	yes	000001365, 000001354, 000001368, 005000974	no
163.81-2	Acute Dermal Toxicity	yes	Technical grade of active ingredient	yes	000001354, 005000974, 000001252	no
163.81-3	Acute Inhalation Toxicity	yes	Technical grade of active ingredient	partial	000001252	yes/8 months[1/]
163.81-4	Primary Eye Irritation	yes	Technical grade of active ingredient	yes	000001352	no
163.81-5	Primary Dermal Irritation	yes	Technical grade of active ingredient	no	-	no[2/]
163.81-6	Dermal Sensitization	yes	Technical grade of active ingredient	no	-	yes/8 months

August, 1981

Table III.A-3 Generic Data Requirements for Temephos, Toxicology (see Part VI) (cont'd)

Guidelines Citation	Name of Test	Are Data Required for this Standard?		Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.81-7	Acute Delayed Neurotoxicity		yes	Technical grade of active ingredient	no	-	yes/24 months[3/]
163.82-1	Subchronic Oral Toxicity	Food crops	yes	Technical grade of active ingredient	yes	000001239, 005001356, 000001240, 000001252	no
163.82-5	Subchronic Neurotoxicity		yes	Technical grade of active ingredient	no	-	yes/18 months[4/]
163.83-1	Chronic Feeding Study	Food crops	yes	Technical grade of active ingredient	yes	000001386	no
163.83-2	Oncogenicity	Food crops	yes	Technical grade of active ingredient	partial[5/]	000001386	yes/36 months
163.83-3	Teratogenicity	Food crops Pets and their living quarters Flea collars	yes yes yes	Technical grade of active ingredient	no	-	yes/36 months

August, 1981

Table III.A-3 Generic Data Requirements for Temephos, Toxicology (see Part VI) (cont'd)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.82-4	Reproduction	Food crops yes	Technical grade of active ingredient	yes	000001388	no
163.84-1 163.84-2 163.84-3 163.84-4	Mutagenicity	Food crops yes Pets and their living quarters yes Flea collars yes	Technical grade of active ingredient	no	-	yes/24 months
163.85-1	Metabolism	Food crops yes	Radiolabeled analytical grade	yes	000001376, 005000858 005000806, 000001343	no

1/ This study will be considered valid if data are submitted to support the statement in the study that the test concentration was at a saturation level. If this information is not provided or considered invalid, further testing will be required.

2/ Since the technical formulation is not irritating to the eyes of test animals, skin irritation is assumed to be unlikely. The data for 163.81-5 are not required.

3/ This test is required because temephos is expected to cause cholinesterase depression.

4/ The requirement for this test is contingent upon the results of the acute delayed neurotoxicity test (see 163.81-7).

5/ The above cited oncogenicity study, while acceptable, must be supplemented by a confirmatory second study (18 months) using the mouse as the test species.

August, 1981

Table III.A-4 Generic-Data Requirements for Temephos, Residue Chemistry (see Part VII)

Guidelines Citation	Name of Test	Are Data Required for this Standard?		Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
-	Metabolism in Plants	Food crops	yes	Technical grade of active ingredient 1/	partial 2/	000001377	yes/24 months
-	Metabolism in Animals	Food crops	yes	Technical grade of active ingredient 1/	yes	000001403, 000001343, 000001344	no
-	Storage Stability	Food crops	yes	Technical grade of active ingredient	no	-	yes/24 months
-	Analytical Methods	Food crops	yes	Technical grade of active ingredient	yes	005000638	no 3/
-	Residue Data:						
	1. Mature Citrus	Food crops	yes	Technical grade of active ingredient	partial	000001402	yes/24 months 4/
	2. Citrus by-products	Food crops	yes	Technical grade of active ingredient	no	-	yes/24 months 4/

August, 1981

Table III.A-4 Generic-Data Requirements for Temephos, Residue Chemistry (see Part VII) (cont'd)

Guidelines Citation	Name of Test	Are Data Required for this Standard?		Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
-	Residue Data: (cont'd)						
	3. Cover Crops	Food crops	yes	Technical grade of active ingredient	no	-	yes/24 months 5/

1/ The ^{14}C-active ingredient should be used.

2/ Data are available on the metabolism of temephos applied to bean leaves. Data are required for 1) the fate of temephos in citrus (or a demonstration that the fate of temephos in a variety of food crops is similar); 2) the uptake of temephos from soil; and 3) the fate of temephos under field conditions.

3/ If other toxicologically significant metabolites are found during the course of plant metabolism studies, analytical methods for their determination must be submitted.

4/ These test data will not be required if metabolism data show that residues other than temephos and temephos sulfoxide are insignificant in plants.

5/ In lieu of acquiring residue data on cover crops in temephos-treated citrus orchards, a restriction against the grazing or feeding of temephos-treated cover crops to livestock may be imposed.

August, 1981

Table III.A-5 Generic Data Requirements for Temephos, Ecological Effects (see Part VIII)

Guidelines Citation	Name of Test	Are Data Required for this Standard?		Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.71-1	Avian Single Dose Oral LD_{50}	Food crops	yes	Technical grade of active ingredient	partial[1/]	005000975, GS0006004	yes/16 months
		Aquatic	yes				
163.71-2	Subacute Dietary toxicity (avian dietary LC_{50})	Food crops	yes	Technical grade of active ingredient	yes	000001384, 000022923	no
		Aquatic	yes				
163.72-1	Fish Acute LC_{50}	Food crops	yes	Technical grade of active ingredient	yes	005000819, GS0006006, GS0006005	no
		Aquatic	yes	Emulsifiable Concentrate	yes[2/]	000001337, GS0006005	no
				Granular	no	-	yes/8 months
163.72-2	Acute Toxicity to Aquatic Inverte-brates	Food crops	yes	Technical grade of active ingredient	yes	GS0006005	no
		Aquatic	yes	Emulsifiable Concentrate	no	-	yes/8 months
				Granular	yes[3/]	GS0006007	no
163.72-3	Acute Toxicity to Estuarine and Marine Organism	Food crops	yes	Technical grade of active ingredient	no	-	yes/16 months
		Aquatic	yes	Emulsifiable Concentrate	no	-	yes/8 months
				Granular	no	-	yes/8 months

August, 1981

Table III.A-5 Generic Data Requirements for Temephos, Ecological Effects (see Part VIII) (cont'd)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.72-4	Embryolarvae and Life-cycle Studies of Fish and Aquatic Invertebrates	Food crops yes Aquatic yes	4/	no	-	4/
163.72-5	Aquatic Organism Toxicity and Residue Studies	Food crops yes Aquatic yes	4/	no	-	4/
163.72-6	Simulated or Actual Field Testing for Aquatic Organisms	Food crops yes Aquatic yes	5/	no	-	5/
163.122-1	Seed Germination/ Seedling Emergence and Vegetative Vigor (Tier I)	Food crops yes Aquatic yes	Technical grade of active ingredient	no	-	yes 6/
163.122-2	Growth and Reproduction of Aquatic Plants (Tier I)	Food crops yes Aquatic yes	Technical grade of active ingredient	no	-	yes 6/

1/ Additional data are needed on dose response, age of test species and specific LD_{50} values.

2/ For the 43% emulsifiable concentrate.

3/ For 5% or less granular formulations.

4/ Data are being reserved pending results of tests pertaining to Guidelines 163.72-1, 2 and 3.

5/ Data are being reserved pending results of tests pertaining to Guidelines 163.72-4 and 5.

6/ A 48 month time period will be allowed from the time of acceptance of the Non-Target Plant Hazard Evaluation Guidelines.

Table III.B-1 Product-Specific Requirements for Manufacturing-Use Products, Product Chemistry[1/] (see Part IV)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.61-3	Product Identity and Disclosure of Ingredients	yes	each manufacturing-use product	yes[2/]	-	no
163.61-4	Description of manufacturing process	yes	each manufacturing-use product	no	-	yes/8 months
163.61-5	Discussion on formation of unintentional ingredients	yes	each manufacturing-use product	no	-	yes/8 months
163.61-6	Declaration and certification of ingredients limits	yes	each manufacturing-use product	no	-	yes/8 months
163.61-7	Product analytical methods and data	yes	each manufacturing-use product	partial[3/]	005000638	yes/8 months
163.61-8(7)	Physical State	yes	each manufacturing-use product	yes[2/]	-	no
163.61-8(8)	Density or Specific Gravity	yes	each manufacturing-use product	partial[4/]	-	yes/8 months

August, 1981

Table III.B-1 Product-Specific Requirements for Manufacturing-Use Products, Product Chemistry[1/] (see Part IV) (cont'd)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.61-8(11)	pH	yes	each manufacturing-use product	no	-	yes/8 months
163.61-8(12)	Storage Stability	yes	each manufacturing-use product	no	-	yes/8 months
163.61-8(13)	Flammability	yes	each liquid manufacturing-use product	partial[5/]	-	yes/8 months
163.61-8(14)	Oxidizing or Reducing Action	yes	each manufacturing-use product	no	-	yes/8 months
163.61-8(15)	Explosiveness	yes	each manufacturing-use product	no	-	yes/8 months
163.61-8(16)	Miscibility	yes	each liquid manufacturing-use product	partial[6/]	-	yes/8 months
163.61-8(17)	Viscosity Coefficient	yes	each liquid manufacturing-use product	no	-	yes/8 months
163.61-8(18)	Corrosion Characteristics	yes	each manufacturing-use product	no	-	yes/8 months

August, 1981

Table III.B-1 Product-Specific Requirements for Manufacturing-Use Products, Product Chemistry[1/] (see Part IV) (cont'd)

1/ Even though some of the requirements have been partially or completely fulfilled for some products, the references cannot be cited to support other products, except for 163.61-7. These requirements must be resubmitted by each applicant at the time of registration or reregistration.

2/ The Agency has information to support all currently registered manufacturing-use products [information was provided in the Confidential Statements of Formula (CSF)].

3/ The analytical method for the active ingredient is of sufficient detail to satisfy the Agency's requirement, but validation data and results of analysis must be submitted. Analytical methods and validation data must be submitted for the impurities for each manufacturing-use product.

4/ The Agency has information to support EPA Reg. No. 241-242 (CSF). This data must be submitted for other manufacturing-use products.

5/ The Agency has information to support EPA Reg. No. 241-242 (CSF). This data must be submitted for other manufacturing-use products.

6/ The Agency has information to support EPA Reg. No. 241-242 (MRID #000001379). This data must be submitted for other manufacturing-use products.

August, 1981

Table III.B-2 Data Requirements for Manufacturing-Use Products, Toxicology (see Part VI)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.81-1	Acute Oral Toxicity	yes	25% concentrate dust 1/	yes	000001307	no
			70% concentrate 1/	yes	000001352	no
163.81-2	Acute Dermal Toxicity	yes	25% concentrate dust 1/	yes	000001307	no
			70% concentrate 1/	yes	000001352	no
163.81-3	Acute Inhalation Toxicity	yes	-	no	-	no 2/
163.81-4	Primary Eye Irritation	yes	25% concentrate dust 1/	yes	000001307	no
			70% concentrate 1/	yes	000001352	no
163.81-5	Primary Dermal Irritation	yes	25% concentrate dust 1/	yes 3/	000001352	no
			70% concentrate 1/	yes	000001352	no
163.81-6	Dermal Sensitization	yes	25% concentrate dust 1/	no	-	no 2/
			70% concentrate 1/			

1/ Each manufacturing-use product or substantially similar product based on Confidential Statement of Formula.

2/ Based on Confidential Statements of Formula; testing of the technical grade of the active ingredient will be sufficient to satisfy this requirement.

3/ The study conducted on the 70% concentrate (MRID #000001352) is also adequate to support the testing requirements on the 25% concentrated dust product.

August, 1981

Table III.C-1 Product-Specific Data Requirements for End-Use Temephos Products, Product Chemistry[1/] (see Part IV)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.61-3	Product Identity and Disclosure of ingredients	yes	each product	yes[2/]	-	no
163.61-4	Description of manufacturing process	yes	each product	partial[3/]	000001267	yes/8 months
163.61-5	Discussion of formation of unintentional ingredients	yes	each product	no	-	yes/8 months
163.61-6	Declaration and certification of ingredient limits	yes	each product	no	-	yes/8 months
163.61-7	Product analytical methods and data	yes	each product	partial[4/]	005000638	yes/8 months
163.61-8(1)	Color	yes	each product	partial[5/]	-	yes/8 months
163.61-8(2)	Odor	yes	each product	partial[5/]	-	yes/8 months

August, 1981

Table III.C-1 Product-Specific Data Requirements for End-Use Temephos Products, Product Chemistry[1/] (see Part IV) (cont'd)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.61-8(7)	Physical State	yes	each product	yes[2/]	-	no
163.61-8(8)	Density or Specific Gravity	yes	each product	partial [6/]	-	yes/8 months
163.61-8(11)	pH	yes	each product	no	-	yes/8 months
163.61-8(12)	Storage Stability	yes	each product	no	-	yes/8 months
163.61-8(13)	Flammability	yes	each liquid product	partial[7/]	-	yes/8 months
163.61-8(14)	Oxidizing or Reducing Action	yes	each product	no	-	yes/8 months[8/]
163.61-8(15)	Explosiveness	yes	each product	no	-	yes/8 months[8/]
163.61-8(16)	Miscibility	yes	each liquid	no	-	yes/8 months

August, 1981

Table III.C-1 Product-Specific Data Requirements for End-Use Temephos Products, Product Chemistry[1/] (see Part IV) (cont'd)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.61-8(17)	Viscosity Coefficient	yes	each liquid product	partial[9/]	-	yes/8 months
163.61-8(18)	Corrosion characteristics	yes	each product	no	-	yes/8 months[8/]

1/ Even though some of the requirements have been partially or completely fulfilled for some products, references cannot be cited to support other products, except for 163.61-7. These requirements must be resubmitted by each applicant at the time of registration or reregistration.

2/ The Agency has information to support all currently registered end-use products [information was provided in the Confidential Statement of Formula (CSF)].

3/ The Agency has partial information to support flea collar products, but lacks information on the amount of starting materials (MRID #000001267). This data must be submitted for other end-use products.

4/ The analytical method for the active ingredient is of sufficient detail to satisfy the Agency's requirement, but validation data and results of analysis must be submitted.

5/ The Agency has information to support EPA Reg. No. 241-132 (MRID #000001254). This data must be submitted for other end-use products.

6/ The Agency has information to support EPA Reg. No. 241-223 (CSF) and EPA Reg. No. 802-495 (MRID #000001379). This data must be submitted for other end-use products.

7/ The Agency has information to support EPA Reg. No. 241-132 (CSF) and EPA Reg. No. 241-223 (CSF). This data must be submitted for other end-use products.

8/ Based on the Confidential Statement of Formula, these data will not be required on the currently registered impregnated materials.

9/ The Agency has information to support EPA Reg. No. 241-132 (MRID #000001379). This data must be submitted for other end-use products.

August, 1981

Table III.C-2.a. Product-Specific Data Requirements for End-Use Temephos Products, Toxicology (see Chapter VI)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition Impregnated Material (40.7%)	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
-	Impregnated Material Test 1/	yes	each product	no	-	yes 2/

1/ Study must be conducted on animal at risk (e.g., dog, cat) at the following dose levels: vehicle control, the proposed use level, and five times the proposed use level. Cholinesterase activities must be measured and animals observed for signs of dermal irritation. See Guideline sections 163.86-1 for general guidance and 163.82-1 for specific guidance for conducting toxicology studies.

2/ Additional test protocols may be developed, especially in the area of child safety requirements.

August, 1981

Table III.C-2.b. Product-Specific Data Requirements for End-Use Temephos Products, Toxicology (see Part VI)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition Emulsifiable Concentrate (43-60%)	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.81-1	Acute Oral Toxicity	yes	1/	yes	000001354, 000001353	no
163.81-2	Acute Dermal Toxicity	yes	1/	yes	000001354, 000001353	no
163.81-3	Acute Inhalation Toxicity	yes	1/	yes	000001354	no
163.81-4	Primary Eye Irritation	yes	1/	no	-	yes/8 months
163.81-5	Primary Dermal Irritation	yes	1/	yes	000001354	no
163.81-6	Dermal Sensitization	yes	1/	no	-	no 2/

1/ Each product or substantially similar product, based on Confidential Statement of Formula.

2/ Based on Confidential Statement of Formula, testing of the technical grade of the active ingredient will be sufficient to satisfy this requirement.

August, 1981

Table III.C-2.c. Product-Specific Data Requirements for End-Use Temephos Products, Toxicology (see Part VI)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition Granular (1, 2, & 5%)	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.81-1	Acute Oral Toxicity	yes	1/	yes	000001308, 000001233	no
163.81-2	Acute Dermal Toxicity	yes	1/	yes	000001308, 000001233	no
163.81-3	Acute Inhalation Toxicity	yes	1/	no	-	no 2/
163.81-4	Primary Eye Irritation	yes	1/	yes	000001308, 000001233	no
163.81-5	Primary Dermal Irritation	yes	1/	yes	000001233	no
163.81-6	Dermal Sensitization	yes	1/	no	-	no 2/

1/ Each end-use product or substantially similar product, based on Confidential Statement of Formula.

2/ Based on Confidential Statement of Formula, testing of the technical material will be sufficient to satisfy this requirement.

August, 1981

Table III.C-2.d. Product-Specific Data Requirements for End-Use Temephos Products, Toxicology (see Part VI)

Guidelines Citation	Name of Test	Are Data Required for this Standard?	Composition Dust (2%)	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
163.81-1	Acute Oral Toxicity	yes	1/	yes	000001308, 000001233	no
163.81-2	Acute Dermal Toxicity	yes	1/	yes	000001308, 000001233	no
163.81-3	Acute Inhalation Toxicity	yes	1/	no	-	no 2/
163.81-4	Primary Eye Irritation	yes	1/	yes	000001308, 000001233	no
163.81-5	Primary Dermal Irritation	yes	1/	yes	000001233	no
163.81-6	Dermal Sensitization	yes	1/	no	-	no 2,3/
163.82-2	21-Day Dermal Sensitization	yes	1/	no	-	yes/8 months 4/

1/ Each end-use product or substantially similar product, based on Confidential Statement of Formula.

2/ Based on Confidential Statements of Formula, testing of the technical material will be sufficient to satisfy this requirement.

3/ In addition to the guideline data (163.81-6), additional test protocols may be developed, especially in the area of child safety requirements.

4/ For dog and cat use, testing is required and must include an adequate cholinesterase analysis of the animals at risk.

August, 1981

Table III.C-3 Product-Specific Products Data Requirements for End-Use Temephos, Efficacy (see Part IX)

Aquatic Sites	Name of Pest	Are Data Required for this Standard?	Composition and Active Ingredient/ Acre (AI/A)	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
Sloughs, stagnant water, log ponds, temporary rain pools, intermittently flooded areas, catch basins, lakes, standing water, shallow ponds, woodland ponds	Mosquito larvae	yes	1,2 and 5% Granular (0.05 to 0.1 lb AI/A)	yes	000001338, 000003267, 000001400, 005000925	no
Tidal water, marshes, swamps, water high in organic matter	Mosquito larvae	yes	1,2 and 5% Granular (0.1 to 0.2 lb AI/A)	yes	000001245, 000001398, 000001397	no
Highly polluted water	Mosquito larvae	yes	2 and 5% Granular (0.5 lb AI/A)	yes	005000637, 005001004, 005000730	no
Marshlands and intertidal zones of sandy beaches	Biting midge larvae (punky and sand fly larvae)	yes	2 and 5% Granular (0.2 lb AI/A)	yes	000001301, 005000928, 000001302	no

August, 1981

Table III.C-3 Product-Specific Products Data Requirements for End-Use Temephos, Efficacy (see Part IX) (cont°d)

Aquatic Sites	Name of Pest	Are Data Required for this Standard?	Composition and Active Ingredient/ Acre (AI/A)	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
Intermittently flooded areas, temporary rain pools, catch basins, standing water, wood-land pools, shallow lakes and lake margins, log ponds, sloughs, stagnant water, marshes, tidal waters, swamps	Mosquito larvae	yes	4 lbs emulsifiable concentrate (0.016 to 0.047 lb AI/A)	yes	000001236, 000001245, 000001247, 000001249, 000001317, 000001326, 000001331, 000001347, 000001349, 000001350, 000001351, 000001399, 000004967, 000004968, 005000710	no

August, 1981

Table III.C-3 Product-Specific Products Data Requirements End-Use Temephos, Efficacy (see Part IX) (cont'd)

Pet and Pet Quarters Sites	Name of Pest	Are Data Required for this Standard?	Composition and Active Ingredient/ Acre (AI/A)	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
Cats: Direct body application and living quarters	fleas	yes	2% Dust	yes	000001217, 000001219, 000001224, 000001225, 000001226, 000001227, 000001228, 000001234, 005000784, 005000792, 005001005	no
Cats: pets collars	fleas	yes	40.7% Impregnated Material	yes	000001260, 000001261, 000001262, 000001270, 000001271, 000001272, 000001273, 000001274, 000001276, 005000793	no
Dogs: direct body application and living quarters	fleas	yes	2% Dust	yes	000001217, 000001218, 000001224, 000001225, 000001226, 000001227, 000001228, 000001234, 005000784, 005000792, 005001005	no

August, 1981

Table III.C-3 Product-Specific End-Use Products Data Requirements for End-Use Temephos, Efficacy (see Part IX) (cont'd)

Pet and Pet Quarters Sites	Name of Pest	Are Data Required for this Standard?	Composition and Active Ingredient/ Acre (AI/A)	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation (MRID #)	Must Additional Data be Submitted under FIFRA 3(c)(2)(B)? If so, months allowed for submission from published date of standard.
Dogs: pet collars	fleas	yes	40.7% Impregnated Material	yes	000001260, 000001261, 000001262, 000001268, 000001270, 000001271, 000001272, 000001273, 000001274, 000001275, 000001290, 005000793	no

August, 1981

IV. Product Chemistry

A. Introduction

The Agency published "Proposed Guidelines for Registering Pesticides" which explains the minimum product chemistry requirements for registering pesticide products in the United States (see 43 FR 29696, July 10, 1978). The Agency requires the following information on chemical identity, manufacturing process, active ingredient limits in pesticide products, product analytical methods and data, and physical/chemical properties in order to assess a pesticide's product chemistry.

B. Chemical Identity

The Agency requires identifying information including chemical names, product names, and numerical codes of all substances known or assumed to be present in pesticide products. As discussed in Part II.B, the common name accepted for temephos by the American National Standards Institute (ANSI) is O,O'-(thiodi-4,1-phenylene)bis(O,O-dimethyl phosphorothioate). The empirical formula for temephos is $C_{16}H_{20}O_6P_2S_3$; it has a molecular weight of 466.4; its chemical structure is shown in Figure 1. An alternative chemical name for the compound is O,O,O',O'-tetramethyl O,O'-thiodi-p-phenylene phosphorothioate and is also known by such trade names as Abate®, Abathion®, Abat® and Biothion®. The Chemical Abstracts Registry (CAS) number for temephos is 3383-96-8, the American Cyanamid internal code number is 52160 (prefixed by AC-, CL-, or EL-), and the EPA Shaughnessy number is 059001. Thus, the nomenclature of temephos has been adequately defined.

C. Manufacturing Process

A detailed description of the manufacturing process is required because the route by which a pesticide is synthesized determines the nature and amount of potentially toxic impurities. Technical temephos is made by reacting dimethylchlorothiophosphate with an alkali salt of 4,4'-dihydroxyphenyl sulfide [4,4'-thiodiphenol] (Pasarela and Orloski, 1973, MRID #005000638). Detailed information (American Cyanamid Co., 1975?, MRID #000001278 and American Cyanamid Co., 1978, MRID #GS0006001) on technical temephos manufacture is Confidential Business Information and is appropriately safeguarded in Agency files. However, the manufacturing process of technical temephos is not detailed enough to satisfy the Agency's requirements. Data on the composition of the technical product indicate that there are several classes of impurities present and identification of individual compounds within a class is difficult (American Cyanamid Co., 19??, MRID #000003268, and American Cyanamid Co., 1974, MRID #000001216). The impurities exceeding 0.1% of the technical temephos product weight must be determined.

The Agency also needs data on the manufacturing process of the other manufacturing-use products and most end-use products. A process description has been submitted on the temephos-



Empirical Formula:

$C_{16}H_{20}O_6P_2S_3$

Molecular Weight:

466.4

FIG. 1

containing flea collars (American Cyanamid Co., 1976, MRID #000001267), but the amounts of starting materials are lacking.

D. Product Analytical Methods and Data

Spectrophotometric and gas chromatographic methods for analysis of temephos in the technical product and in formulations have been published (Pasarela and Orloski, 1973, MRID #005000638). For analysis by ultraviolet spectrophotometry, solid samples are extracted with methylene chloride, the solvent evaporated, and the residue dissolved in a mixture of ethyl acetate and hexane. Liquid samples are dissolved directly in the ethyl acetate-hexane solvent. Samples are cleaned up on silica gel columns, the column eluate evaporated to dryness, and the residue dissolved in methanol. The absorbance of the methanol solution at 274 nanometer (nm) wavelenght is determined and the temephos content determined by comparison to a standard carried through the clean-up procedure.

Precision of the method is claimed to be ± 1.0% for technical temephos, ± 0.7% for 6E emulsifiable concentrates, and ± 0.2% for 1% through 3% granular formulations (Pasarela and Orloski, 1973, MRID #005000638).

For gas-chromatographic analysis of products, liquid samples or methylene chloride extracts of solid samples are diluted with methylene chloride, the solutions cleaned up on columns of acidic alumina, and the column eluate analyzed for temephos on a gas chromatographic column of OV-22® silicone on 60-80 mesh Gas-Chrom Q® equipped with a flame ionization detector. An internal standardization technique is used to determine the content of temephos. Claimed precision for the method when peak areas are determined by electronic integration is ± 1.0% for the technical product, ± 0.7% for 6E emulsifiable concentrates, and ± 0.02% for 1% through 3% granular formulations (Pasarela and Orloski, 1973, MRID #005000638).

For determination of temephos in flea collars, one gram of collar is dissolved in tetrahydrofuran and polymeric material precipitated by addition of hexane. The solution is cleaned up on a silica gel thin-layer chromatography plate, temephos eluted from the plate with methanol, and the concentration of temephos in the methanol solution determined by spectrophotometry (Fiala, 1975, MRID #000001269)

Methods for the determination of active ingredient in temephos containing products have been published, but validation data and results of analysis obtained by these methods have not been submitted.

E. Physical/Chemical Properties

A small amount of data is available on the physical/chemical properties of temephos. Data which are not available but which are required to be submitted are listed in the Tables III.A through C in Part III. All of the following physical/chemical

property data were developed by the American Cyanamid Company. The year of publication and Master Record Identification (MRID) numbers are as follows: 19??, #000003264; 1972?, #000001379; 1974?, #000001223; 1974?, #000001222: and 1969, #000001254. Each property listed below has only the MRID number referenced or references are made to the Confidential Statement of Formula (CSF). The following available data for temephos are either for the purified grade, technical grade, or formulated product as noted.

Color: The purified grade is a white crystalline solid (MRID #000001379); the technical grade is a brown viscous liquid (MRID #000001379); and the end-use product, EPA Reg. No. 241-132, is an amber liquid (MRID #000001254).

Odor: End-use product, EPA Reg. No. 241-132, is described as having a "typical aromatic petroleum odor" (MRID #000001254).

Melting Point: Purified grade melts at 30.0°-30.5°C; technical may crystallize below 15°C (MRID #000001379).

Density or Specific Gravity: The specific gravity of technical temephos is 1.3 at 25°C (MRID #000001379). The density of a manufacturing-use product, Abate® 70% concentrate (EPA Reg. No. 241-242), is 1.21 gms/cm^3 (CSF). The density of two end-use products is 1.2 gms/cm^3 (CSF) for a 6E formulation (EPA Reg. No. 241-223) and 1.13 gms/cm^3 for a 4E formulation (EPA Reg. No. 802-495) [MRID #000001379].

Physical State: Purified grade is a crystalline solid and technical grade is a viscous liquid (MRID #000003264). Formulations currently registered are liquids, granules, dusts or impregnated materials (CSF).

Stability: Technical grade appears to be indefinitely stable at room temperature and moderately stable to hydrolysis in contact with aqueous alkali. No hydrolysis was observed at pH 8 at room temperature for several weeks or at pH 11 at 40°C for several hours; hydrolysis at high pH for prolonged periods of time may be expected (MRID #000001379).

Solubility: Technical grade is soluble in water at 25 ppb at 25°C (MRID #000001379); also soluble in the following solvents*: acetonitrile, carbon tetrachloride, chloroform, ether, ethylene dichloride, lower (molecular weight) alkyl ketones, and aromatic hydrocarbons. It is insoluble in hexane and methyl cyclohexane (MRID #000001379).

Storage Stability: The stability of end-use products stored under typical conditions is the basis for over formulation or

---

* The Agency requires a numerical value for solubility of temephos in organic solvents.

for assigning expiration dates, and for assessing changes over time of such physical/chemical properties as pH, viscosity, color, odor, and density. The Agency has information on the 25% dust concentrate, EPA Reg. No. 241-235 (MRID #000001223) and the 2% powder, EPA Reg. No. 241-234 (MRID #000001222). However, the storage conditions of these two formulations are not sufficiently detailed.

Flammability: Data on flammability determines the type of flammability warnings to be placed on labels of pesticide products. The manufacturing-use product, Abate® 70% concentrate insecticide (EPA Reg. No. 241-242), has a flash point of 84°C T.O.C. (CSF); and the end-use products have a flash point of 43°C T.C.C. and >85°C T.O.C. (CSF), and >80°C T.O.C. (CSF) for Biothion® 6E Insecticide (EPA Reg. No. 241-223) and Abate® 4E Insecticide (EPA Reg. No. 241-132), respectively.

Specific Viscosity: Technical grade (liquid), EPA Reg. No. 241-220, has a viscosity of 3,000 centipoises (cp) at 5°C, 300 cp at 25°C, 180 cp at 40°C, and 30 cp at 80°C (MRID #00001379). The viscosity of the liquid end-use product (EPA Reg. No. 241-132) is approximately 36 cp at 25°C (MRID #000001379).

F. Summary of Data Gaps

Though some information is available to satisfy the Guidelines' requirements on description of manufacturing process, discussion on formation of unintentional ingredients, declaration and certification of ingredient limits, and product analytical methods, additional data are required in all instances for the manufacturing-use and end-use products (see Tables III.B-1 and C-1 for specific-data requirements).

There is little or no information for technical temephos regarding odor, octanol/water partition, pH, solubility in organic solvents, boiling point*, vapor pressure, and product identity and disclosure of ingredients.

There is little or no information for temephos manufacturing-use products regarding density or specific gravity, pH, storage stability, flammability (liquid), oxidizing or reducing action, explosiveness, miscibility (liquid), viscosity coefficient (liquids), and corrosion characteristics. There is also little or no information for temephos end-use products regarding color, odor, density or specific gravity, pH, storage stability, flammibility (liquid), oxidizing or reduction action**, explosiveness**, miscibility (liquids), viscosity coefficient (liquids), and corrosion characteristics**.

---

* Data may be submitted indicating that the boiling point cannot be determined.

** Based on the Confidetial Statement of Formula, the data are not required for the currently registered impregnated materials.

V. Environmental Fate

A. Use Summary

Temephos is primarily a larvicide for use against the larval stage of mosquitoes, gnats, and biting midges (punkies and sand flies). The only registered food use (Federal Registration) is for the control of thrips on oranges and tangerines in Arizona and California. It is also registered for control of fleas on cats and dogs, and in their living quarters.

The major areas of application for temephos are considered aquatic such as intermittently flood areas, marshes, lake margins, ponds, pools, sloughs, standing water and sandy beaches.

The percent of temephos in the various federally registered end-use formulations are as follows: emulsifiable concentrates: 43, 45.2, 48.5, 50.3 and 60%; granular formulations: 1, 2 and 5% temephos; dust formulation: 2%; and impregnation in pet collar material: 40.7%.

The granular and emulsifiable concentrate formulations are used predominately for the control of mosquitoes, gnats and biting midges. The rates of application range from 0.05 to 0.5 lb of active ingredient (ai)/acre for the granular form and from 0.016 to 0.047 lb ai/acre for the emulsifiable concentrate. The granular formulations contain, by weight, 1-5% of the active ingredient, whereas the emulsifiable concentrate contains 4 lbs of the active ingredient per gallon.

Higher application rates are registered for polluted aquatic sites and those sites that contain increased levels of organic content. The dust formulations and the impregnated pet collar have an unspecified rate of application. The emulsifiable concentrate registrated for use on orange and tangerine trees has a 1.0 lb ai/acre aerial or ground application rate.

The Agency estimates that the total usage of temephos as an active ingredient is 100,000-200,000 lbs, annually. The quantity used for non-aquatic site treatment (e.g., food use), which currently is exported, is quite minor, probably less than 2%.

B. Environmental Fate Profile

The available data are insufficient to fully assess the environmental fate of temephos [O,O'-(thiodi-4,1-phenylene)bis(O,O-dimethyl phosphorothioate)].

1. Hydrolysis

Available preliminary data from a microbial degradation study (Daorai and Menzer, 1977, MRID #005000796) on the hydrolysis of radiolabeled temephos indicate a degradation rate of about 10%

after 10 days in distilled water (pH near neutral), resulting in the following five degradation products:

1) O,O,O',O'-tetramethyl O,O'-sulfinyldi-p-phenylene phosphorothioate (S-SO-S);

2) O-(p-(p-hydroxyphenylthio)phenyl) O,O-dimethyl ester with O',O'-dimethyl phosphate (S-S-O);

3) O-(p-(p-hydroxyphenylsulfonyl)phenyl) O,O-dimethyl ester with O',O'-dimethyl phoshate ($S-SO_2-O$);

4) O-(p-(p-hydroxyphenylthio)phenyl) O,O-dimethyl phosphorothioate (S-S-OH); and

5) O,O,O',O'-tetramethyl O,O'-thio-p-phenylene phosphate (O-S-O).

The first degradation product listed was predominant. These data indicate a relatively slow rate of hydrolysis for temephos. However, the data that are available on hydrolysis is considered insufficient because the Daoria and Menzer findings (1977, MRID #005000796) are not conclusive.

2. Microbiological - Effects of Microbes on Pesticides

Evidence regarding the bacterial degradation of temephos (1% Abate®) indicated that previous soil enrichment with temephos may increase the degradation rate of the compound to seven different products. Daorai and Menzer (1977, MRID #005000796) found that soil containing 1% temephos, plus an unidentified bacterium (possibly a Pseudomonas) previously cultured in the presence of temephos, resulted in 47% degradation of the applied compound (1% Abate®) to unidentified water-soluble metabolites of temephos in 10 days, whereas the presence of a second bacterium not previously cultured in the presence of temephos, produced only 28% degradation of the compound to the water-soluble metabolites over the same period of time. Five of the seven degradation products from both bacterial cultures were the same as those formed by hydrolysis. Two additional degradation products formed by both bacterial cultures were unidentified. This study provided some preliminary information on the microbial degradation of temephos by two unidentified bacterial species; however, further studies with a series of microbes indigenous to temephos treatment sites are necessary before an adequate assessment of microbial metabolism of temephos can be made.

3. Anaerobic and Aerobic Aquatic Metabolism

Temephos was degraded relatively rapidly in samples of outdoor waters (Drevenkor et al., 1977, MRID #005000718). According to preliminary data, only 40% and 57% of the temephos added at about 0.7 ppm as a granular formulation remained after 26 days in polluted brook and puddle water, respectively. Soluble temephos was not detected in the puddle water samples at the end

of the test period, and only about 40% of the temephos detected in the brook water at the end of testing (26 days) was in the soluble state. This study provides only preliminary information on aquatic metabolism since metabolites were not analyzed and some minor problems may have been caused by interfering substances.

4. Microbiological - Effects of Pesticides on Microbes

Preliminary data indicate that the effect of temephos on microbial populations is generally slight. Bacterial populations in waste water exposed to an unspecified formulation of temephos at 1 ppm exhibited minimal effects over a 48-hour test period. At higher concentrations (1,000-50,000 ppm), temephos decreased bacterial populations 49-87% after 24 hours (Steelman et al., 1967, MRID #005001102). Temephos concentrations of 20 ppb and 500 ppb increased the rate of nitrogen fixation in mixed plankton cultures (94% of the culture consisted of the blue-green alga Aphanizomenon flos-aquae) to about 160% and 390%, respectively above controls (Wurtsbaugh and Appersen, 1978, MRID #005011182). Other data demonstrated that temephos at 2.5 ppm in Danube River water within 24 hours of exposure had a slight inhibitory effect (10-30% growth inhibition) on major phytoplankton populations, including green algae, diatoms, and euglenoids. Temephos at higher concentrations resulted in greater inhibitions. Conversely, temephos (1% in sand granules) at 2.5 ppm caused a 100% increase in blue-green algal populations (Tevan and Lang, 1975, MRID #005003734). These studies provided partial information concerning the effect of temephos on nitrogen fixation by blue-green algae and the effect of temephos on populations of aquatic algae. Further studies are needed for temephos that use other soil microorganisms and address other parameters, including oxygen consumption, carbon dioxide evolution, nitrogen cycle reactions and enzyme activities. The Agency also recommends that additional data be acquired on the effects of temephos at higher concentrations (above 2.5 ppm) on nontarget algae populations (eucaryotic algae) since the available data and the use patterns indicate that temephos may have an adverse impact on these organisms, which constitute the majority of primary producers in aquatic environments.

5. Field Dissipation - Aquatic

Preliminary data from one study (Henry et al., 1971, MRID #005000794) indicated that temephos (technical grade) concentrations decreased quickly from an initial level of 0.1 ppm to steady-state concentrations of 0.02-0.04 ppm within 48 hours in a micro-marsh at temperatures of 10-35°C. In another study (Vrochinskii, 1976, MRID #005003887), the half-life of an unspecified formulation of temephos applied at about 3.5 ppm to a model reservoir (18-20°C, pH 7.4 to pH 8, dissolved O_2 at about 6-8 ppm) was 9 days. By 113 days, 95% of the applied temephos had dissipated. In both studies, the degradation products were not identified but indicated a moderately low degree of persistence for parent temephos in the aquatic environment.

6. Fish Accumulation

Preliminary data indicated that temephos (50% emulsifiable concentrate) accumulated in Indopacific milkfish (Chanos chanos) exposed to the compound at about 0.35 ppm had a relatively low bioaccumulation factor of approximately 2 over an unspecified time period (Tsai, 1978, MRID #005000978). Rapid depuration occurred as the temephos residue level declined over a 14-day period, from about 0.7 ppm to about 0.02 ppm, in both whole and edible fish tissues.

In summary, temephos is hydrolized relatively slowly; however, limited data suggest in aquatic environments, temephos degrades rapidly to at least seven different products, five known and two unidentified. The environmental fate of these products is unknown. This pesticide generally has limited effects on bacterial populations, but may have an adverse impact on eucaryotic algae. Growth and nitrogen fixation is stimulated by temephos in some blue-green algae exposed at low concentrations. In a static flow study, temephos exhibited a low accumulation potential and rapid depuration in fish. Additional studies are required to measure the effects of temephos on fish in a flow through environment.

C. Exposure Profile

1. Introduction

Granular (G) and emulsifiable concentrate (EC) formulations of temephos are registered for control of aquatic larvae (mosquitoes, sand flies, midges, gnats, etc.) at a variety of sites (sandy beaches, catch basins, lakes, marshes, ponds, etc.). The G and EC formulations are applied at rates ranging from 0.05 to 0.5 lb ai/acre and from 0.016 to 0.047 lb ai/acre, respectively. These formulations are primarily applied by aircraft, although hand sprayers, mechanical ground sprayers, broadcast and ultra low volume (ULV) ground applications are also used.

A dust formulation and temephos in impregnated material (flea collars) are registered for control of fleas on dogs and cats (the dust formulation is also used in pet living quarters and bedding).

2. All Formulations (except impregnated materials)

A comprehensive exposure profile cannot be generated at the present time (August, 1981) because of the scarcity of data. However, from the data which are available, the following profile can be developed:

> The principal region of exposure is expected to be the northeastern quarter of the United States and Florida where the greatest use of temephos occurs. As with most pesticides, the greatest human exposure will probably be to workers engaged in mixing, loading, and working in treatment operations. However, quantitative

data are not available to estimate the degree of such occupational exposure nor to accidental exposures, i.e., the public which may be within or surrounding the area of application, or human exposures resulting from the domestic use. The potential exposure by inhalation of volatilized temephos cannot be assessed due to the lack of vapor pressure and volatility data on the manufacturing-use product or formulations. Similarly, the lack of soil mobility data prevents an assessment of the potential for human dietary exposure through ground water contamination. Degradation of temephos by soil bacteria has been demonstrated, but the unknown environmental fate of the temephos degradation products prevents an adequate determination of how important this degradation is in eliminating use-associated hazards. Field dissipation data do suggest a slight persistence of temephos in the aquatic environment.

a. Granular Formulations

Granular formulations of temephos are applied primarily via aircraft. The potential for human exposure (especially respiratory and ocular) during loading operations is recognized, although available information indicates that such exposure is minimized by the use of protective clothing and a respirator or surgical mask during this operation*. The potential for pilot exposure during application is minimized by positive air flow ventilated aircraft cabins and by closed insecticide delivery systems. Drift from aerial application of these formulations is not considered an important mode of exposure for persons, livestock, and wildlife outside the application site due to the weight of the granules.

Applicator exposure during ground applications may be higher than during aerial applications where, as discussed above, the pilot is protected from the insecticide. However, the reported use of rubber boots, gloves, coveralls, and respirators by ground applicators can reduce exposure during loading and application*.

The potential exposure of nontarget organisms, especially wildlife, through contact with treated surface waters must also be considered, although preliminary environmetnal fate data indicate that such exposure might be limited by the dissipation of temephos in aquatic systems. The aquatic field dissipation of temephos has been demonstrated to vary between a half-life of 9 days in a model reservoir to less than 48 hours in micro-marsh ponds.

The labeling for these products recommends "avoid contact with eyes, skin and clothing" and "avoid breathing spray mist or dust."

b. Emulsifiable Concentrate Formulations

Exposures resulting from the use of emulsifiable concentrates of temephos in aquatic areas are expected to be similar to those discussed previously for granular formulations. Although a potential for dermal exposure via splashing of the EC formulations during dilution and tank mix operations is recognized which could be further reduced by use of a closed transfer system, such exposure would be minimized by the use of protective equipment described previously. The potential for spray drift from the use of emulsifiable concentrate formulations is especially high because these formulations are commonly applied from aircraft as ULV sprays.

c. Impregnated Materials

Little exposure to humans is expected from such formulations. Exposure that does occur is likely to be dermal in nature, resulting from handling flea collars. No exposure to wildlife or livestock is expected from these formulations.

d. Dust Formulations

Little exposure to humans, wildlife, or livestock is expected to result from using these formulations for controlling fleas on cats and dogs, and their living quarters.

D. Summary of Data Gaps

The data gaps that are to be filled are hydrolysis, photodegradation, aerobic aquatic metabolism, fish accumulation, leaching, adsorption/desorption, water dispersal, terrestrial field dissipation (tree fruit and nut crop use), aquatic field dissipation (aquatic noncrop use), terrestrial/aquatic (forest field dissipation), accumulation in irrigated crops, special studies (accumulation in aquatic noncrop uses), aerobic soil metabolism, and anaerobic aquatic metabolism.

There are data gaps relating to disposal and storage, microbial metabolism and activated sludge. However, pending review and modification of the testing protocol, it is currently not necessary to satisfy these data gaps. Further testing details are in Part III.

VI. Toxicology

A. Toxicology Profile: Technical Temephos

Temephos is an organophosphorus pesticide. Low doses of temephos would be expected to inhibit cholinesterase activity without necessarily causing acute signs. In vivo, temephos has been shown to inhibit cholinesterase activity. In rats, subchronic oral doses as low as 6 ppm resulted in depressed erythrocyte cholinesterase activity (Levinskas, 1965, MRID #000001239). In dogs, subchronic oral doses of 700/500 ppm caused depressed erythrocyte, brain, and plasma cholinesterase activities (Hutchinson and Levinskas, 1965, MRID #000001356). Subchronic oral dosing with temephos caused no effect on survival, food consumption, or tissue and organ histopathology in rats given up to 350 ppm in the diet, and dogs given up to 700/500 ppm in the diet for 90 days (Levinskas, 1965, MRID #000001239 and Hutchinson and Levinskas, 1965, MRID #000001240). Weight gain was reduced in rats given 350 ppm for 90 days. Cholinesterase activity was depressed after subchronic oral dosing. In another subchronic oral study (American Cyanamid Co., 1964, MRID #000001252), male rats were fed diets containing 0, 250, 500, and 1000 ppm (0, 0.025, 0.05, and 0.1%) technical temephos for 30 days. The No Observable Effect Level (NOEL) for systemic effects was 250 ppm and the Lowest Effect Level (LEL) was 500 ppm; the effect being weight gain depression. Cholinesterase depression (i.e., brain, plasma, and erythrocyte) was noted in all dose groups.

The relatively high acute oral LD_{50} (0.77-13.0 g/kg) of technical temephos in rats suggests a low acute hazard in humans (American Cyanamid Co., 1966?, MRID #000001354; Gaines et al., 1967, MRID #000001368; Gaines, 1969, MRID #005000974 and McNerney and Levinskas, 1967, MRID #000001365). The large range in LD_{50} values might be attributed to differences in sensitivity of the strains of rats tested. Signs of toxicity in rats were characteristic of organophosphate intoxication (i.e., tremors, salivation, diarrhea, tearing, depression, and weakness).

A moderate dermal hazard in humans is suggested by the relatively low acute dermal LD_{50} in rabbits (0.97 g/kg to greater than 4 g/kg) [American Cyanamid Co., 1966?, MRID #000001354 and Gaines, 1969, MRID #005000974]. In another study which provided supplemental information using male rabbits, the LD_{50} was estimated to be 1.02 g/kg (American Cyanamid Co., 1964, MRID #000001252). No tests of primary dermal irritation or an adequate subchronic dermal studies were available on the technical temephos.

In an acute inhalation study, albino CF Nelson rats (six males) were exposed to air saturated with technical temephos (concentration unspecified) for 8-hours (American Cyanamid Co., 1964, MRID #000001252). Deaths did not occur and toxic signs were not noticed; suggesting that technical temephos has a low acute inhalation toxicity potential. However, until data aresubmitted

to support the statement that the test concentration was at the saturation level, additional testing will be required.

Temephos has a very low potential for eye irritation in humans based on a test in rabbits. Instillation of this material into the conjuctival sac of rabbits resulted in no eye irritation (American Cyanamid Co., 1974, MRID #000001352).

In Sprague-Dawley rats, dietary administration of technical temephos at doses of 10, 100, and 300 ppm for 2 years had no adverse effects; however cholinesterase activities were not measured (Underwood, 1977, MRID #000001386). This study can be used to assess the chronic and oncogenic potential of temephos. An oncogenic study using mice will be required on the technical product (see Table III.A-3 in Part III for Toxicology).

There are no adequate studies available on the teratogenic and mutagenic effects of temephos. In a reproduction study, no treatment-related effects occurred in the reproductive capacity of rats given dosages of 25 ppm or 125 ppm temephos. Furthermore, no effects were noted in the offspring of these rats (McNerney et al., 1968, MRID #000001388).

Metabolism of temephos is relatively rapid after oral dosing. In rats and guinea pigs, about 95% of a single oral dose of temephos was eliminated by fecal and urinary routes within 96 hours. The average half-life of temephos was reported as 10 hours in rat blood (Blinn, 1966, MRID #000001376 and Blinn, 1969, MRID #005000858) and the residues of temephos in tissue were relatively low; adipose tissue contained the highest levels (1-2 ppm). Metabolites identified in urine and feces were 4,4'-thiodiphenol, and to a lesser extent, 4,4'-sulfinyldiphenol, either free or as glucuronide or sulfate conjugates (Blinn, 1969, MRID #005000858).

When temephos was applied as a single dermal dose to lactating goats, approximately 33% of the absorbed dose was excreted in urine, 50% was excreted in the feces, and 12-15% was present as tissue residues at 14 days (Chiu and Eisner, 1975, MRID #000001343). The metabolites identified in the urine and feces of goats (either free or as conjugates) were 4,4'-sulfinyldiphenol and temephos sulfoxide which were also indentified in the rat. In addition, temephos sulfone and monooxon were found in feces. Residues of about 0.3 ppm were found in fat. Shafik (1970, MRID #005000806) demonstrated that a temephos metabolite 4,4'-thiodiphenol reached its peak level in rats after a 3 day daily dosing of 9.6 mg/Abate®. The metabolite peaked at 300 ug/day and then dropped to 10 ug/day after 14 days.

B. Toxicology Profile: Manufacturing-Use Temephos

Using a 25% temephos dust concentrate, very low acute oral hazard and a low acute dermal hazard are expected. This is because the oral LD_{50} in male rats is 6.20 g/kg body weight and the dermal LD_{50} in male rabbits is greater than 5 g/kg (Fischer, 1974, MRID #000001307).

Applying a 70% concentrate of Abate®, the acute oral LD_{50} in male rats was 2.03 g/kg body weight and the dermal LD_{50} in male rabbits was 1.89 g/kg body weight (American Cyanamid Co., 1974, MRID #000001352).

The 25% temephos dust concentrate is expected to have a very low eye irritation potential in humans because instillation of this material in rabbit eyes resulted in only mild conjunctival irritation at 24 hours (Fischer, 1974, MRID #000001307). The primary eye irritation test using a 70% concentrate on rabbits showed negative results (American Cyanamid Co., 1974, MRID #00001352).

In a primary dermal irritation study, mild to moderate erythema was observed 72 hours after a 70% concentrate formulated intermediate was applied to abraded rabbit skin (American Cyanamid Co., 1974, MRID #000001352). This study is also sufficient to support the toxicity of the 25% dust concentrate.

C. Toxicology Profile: End-Use Temephos

1. Emulsifiable Concentrate

Low acute oral toxicity and low acute dermal toxicity are expected in numans for Abate® mosquito larvicide (one gallon contains 4 lbs of 51.5% temephos) based on the relatively high oral LD_{50} in rats (3.08-3.53 ml/kg) and dermal LD_{50} in rabbits (1.77-2.03 ml/kg) [American Cyanamid Co., 1966?, MRID #000001354].

Acute oral LD_{50} of Biothion® 6E (containing 69.2% temephos by weight or 60% active ingredient) in male rats (1.40 g/kg) suggests a low acute oral hazard potential in humans. Similarly, the acute dermal LD_{50} in male rabbits (5.60 g/kg) indicates low dermal hazard (American Cyanamid Co., 1971; MRID #000001353). Placing Biothion® 6E in the conjunctival sac of rabbits caused mild to moderate conjunctival irritation and mild corneal irritation (American Cyanamid Co., 1971, MRID #000001353). However, this study is not sufficient to classify the toxicity of the EC formulations because the observation time for development of corneal opacity in the test animals did not exceed 72 hours.

In an acute inhalation study using the 4 lbs/gallon emulsifiable concentrate, Wistar rats were exposed to 2.68 mg/liter (nominal concentration) for 1 hour and were observed for 7 hours following exposure (American Cyanamid Co., 1966?, MRID #000001354). The animals did not die but they did exhibit annoyance and discomfort. This study demonstrated that the 4 lbs/gallons EC has a low acute inhalation toxicity potential.

In a primary dermal irritation study, application of 0.5 ml of the 4 lbs/gallons emulsifiable concentrate on intact and abraded rabbit skin resulted in moderate to severe erythema and edema which persisted at the 72 hour reading (American Cyanamid Co., 1966?, MRID #00001354). This study indicates that the

4 lbs/gallon EC formulations cause moderate to severe dermal irritation.

2. Dust Formulation

Using a 2% Abate® dusting powder, the oral LD_{50} in male rats was greater than 5 g/kg body weight and using a 2.44% (by weight) temephos powder, the oral LD_{50} in male rats was greater than 10 g/kg body weight (American Cyanamid Co., 1971, MRID #000001308 and American Cyanamid Co., 1973, MRID #000001233). Both dusts tested on male rabbits showed a dermal LD_{50} greater than 5 g/kg body weight.

A test in rabbits using the 2% dusting powder revealed no eye irritation, and tests using the 2.44% (by weight) formulation revealed only mild conjunctival irritation after 4 hours. These dusts were also tested in an acute dermal study on intact and abraded skin of rabbits and showed only mild dermal irritation (American Cyanamid Co., 1971, MRID #000001308 and American Cyanamid Co., 1973, MRID #000001233). In the primary dermal irritation study, application of 0.5 grams of an aqueous paste of a 2% dust resulted in mild erythema at 24 through 72 hours; indicating a very low potential for dermal irritation (American Cyanamid Co., 1973, MRID #000001233).

Regarding the subchronic dermal toxicity test, dogs, using 2% temephos powder with a dosage of 80 mg/kg body weight once a week for 3 weeks, or 20 mg/kg body weight for five days a week for three weeks, resulted in no adverse effects including no effects on cholinesterase activity and growth. Cats dosed with the same powder with a dosage of 80 mg powder per kilo of body weight one or five times a week for three weeks showed some inhibition of plasma and erythrocyte cholinesterase activity. In these tests, however, sample size were quite limited and changes in various parameters (e.g., other toxic effects) might not have been detected (Wang et al., 1974, MRID #000001232).

In humans, single-dose dusting with a 2% formulation of temephos in pyrax powder did not result in depresson of whole blood cholinesterase values (Steinberg et al., 1970, MRID #005001274). Patch tests with this formulation indicated no evidence of skin irritation or sensitization in humans (Steinberg et al., 1970, MRID #005001984). Sleeve-tests with 1% or 2% temephos in pyrophyllite powder resulted in no ill effects (Cole et al., 1971, MRID #000001312). Ingestion of increasing doses of temephos (2-256 mg/day) over four weeks or daily doses of 64 mg/day for four weeks did not result in alterations in red blood cell or plasma cholinesterase activity or in clinical symptoms (Laws, 1975, MRID #000001375).

3. Granular Formulations

No toxicity data were available on granular formulations containing temephos; however, based on the similarities in the composition of dust and granular formulations, as indicated by the Confidential Statements of Formulae, the studies cited for

the dust formulation can be used to support the registrations of the granular formulations.

4. Impregnated Materials

No toxicity data were available on dog and cat flea collars impregnated with temephos. Since the temephos is impregnated into the material and inhalation and dermal sensitization data will be required on the technical grade of active ingredient, the Agency will only require a special impregnated material study (see Table III.C-2.a for study requirements). This study must include an adequate cholinesterase analysis of the animals at risk (i.e., cat and dog). Also, based on the results of the dermal sensitization study required for the technical grade of active ingredient, additional test protocols may be developed, especially in the area of child safety requirements.

D. Human Hazard Assessment

Exposure to temephos can be experienced by the dermal, inhalation or dietary routes. It is difficult to assess the degree of exposure to temephos by inhalation or by ground water contamination, due to insufficient environmental chemistry data.

1. Technical Temephos

Temephos is an organophosphorus pesticide which at low doses inhibits cholinesterase activity in animals without necessarily causing symptoms of acute toxicity. Technical grade temephos exhibits low acute oral and dermal toxicity to animals and its potential for eye irritation is very low based on animal data. Weight gain and cholinesterase activity were depressed after subchronic dosing of rats (350 ppm in the diet, 90 days) and dogs (700/500 ppm). No effects on reproduction were seen when rats were given up to 125 ppm temephos in their diet. No chronic or oncogenic effects were seen when rats were fed up to 300 ppm temephos. Metabolism studies of temephos indicated that its half-life was about 10 hours in rat blood. Tissue residues were relatively low; adipose tissue had the highest level.

2. Manufacturing-use Products

Based on animal studies, a very low acute oral and a low acute dermal hazard is expected for humans. Based on eye irritation tests in rabbits, the Agency concludes that when results are extrapolated to humans, the eye irritation is expected to be low.

3. End-Use Products

a. Emulsifiable Concentrate (EC) Formulations

The acute oral and dermal toxicity of Abate® mosquito larvicide (51.5% temephos) is expected to be low, based on animal data. This formulation is expected to be moderately irritating to human eyes, and to be a moderate to severe dermal irritant.

Biothion® 6E (69.2% temephos by weight) is similar to Abate® mosquito larvicide in its acute oral toxicity and dermal toxicity effects. Its ability to cause eye irritation is similar to Abate® mosquito larvicide (51.5% temephos). Dermal exposure to humans is possible via splashing during dilution and the tank mixing of these formulations. The acute inhalation test provided information that the emulsifiable concentrate (EC) formulation has a low inhalation toxicity potential. The primary dermal irritation test revealed a moderate to severe dermal irritation potential.

b. Dust Formulations

Based on animal toxicity studies using 2% temephos powder, it is expected that for humans, temephos has the potential to cause very low levels of acute oral toxicity, acute dermal toxicity, primary dermal irritation and eye irritation.

Subchronic dermal dosing using dogs showed no effect, whereas cats showed some inhibition of cholinesterase activity. When temephos was used on humans for single dose dusting, patch tests, sleeve-tests and oral ingestion, no effects were evident.

In a study pertaining to the application of a 2% Abate®/98% pyrax dust formulation for lice control, it was suggested that the operator of the dusting equipment use some form of respiratory protection to minimize exposure. However, this is an empirical evaluation and not based on any quantitative risk/benefit analysis.

d. Granular Formulations

Granular formulations are applied by using aircraft, thus exposure to people outside the application site due to spray drift may be a concern. In general, applicator exposure during ground applications could be reduced by use of protective clothing and respirators. However, this is an empirical evaluation and not based on any quantitative risk/benefit analysis.

e. Impregnated Material

Because of the data gap for the impregnated materials, the Agency cannot adequately assess the hazard potential to domestic animals and humans from this formulation. However, information available on the technical and emulsifiable concentrate formulations of temephos indicate that a low hazard to domestic animals and humans is expected.

E. Summary of Data Gaps

The generic data gaps for technical grade temephos are acute inhalation toxicity, dermal sensitization, acute delayed neurotoxicity, teratogenicity, mutagenicity and oncogenicity (mouse).

Oncogenicity (in the mouse), teratogenicity, and mutagenicity tests are required because temephos is registered for use on food crops. Some of these studies (see Table III.A-3) are also required for the domestic uses of temephos (i.e., for direct application on cats, dogs, and in their living quarters; dogs' and cats' flea collars). The data required for the food crop use is also required to maintain the established tolerance. If these studies are not provided to maintain the tolerance, the Agency will implement procedures under FIFRA Section 180.001 and Food and Drug Cosmetic Act (FDCA) Section 408(e) to revoke the temephos-established tolerance.

The data gap for end-use temephos products are the 21-day dermal sensitization test for the dust formulation and an impregnated material study for the flea collar products on the animals at risks (i.e., cat and dog). In addition to the dermal sensizition test being conducted on the technical grade of active ingredient, test protocols may also be developed, especially in the area of child safety requirements. The data gap for the emulsifiable concentrate is the primary eye irritation test.

VII. Residue Chemistry

A. Introduction

Temephos is an insecticide with registered uses on 1) dogs, cats, and in their living quarters for control of fleas; 2) orange and tangerine trees to combat thrips, and 3) aquatic sites to control larvae of mosquitoes, "gnats", and certain biting midges.

Foliar treatment with a 60% emulsifiable concentrate of temephos is used for control of thrips on orange and tangerine trees. Following harvest of mature fruit, the formulation is applied at one pound per acre at prebloom or at petal fall, but not during bloom. The formulation may also be applied to nonbearing trees. The formulation is diluted in 15 to 20 gallons of water per acre for aerial application and 200 gallons per acre for ground application. There is no specified frequency of application on the label and the use of temephos on citrus is limited to Arizona and California. With the exception of the use on citrus, the above uses of temephos are not food uses; however, the likelihood exists that application of temephos to aquatic sites could result in contamination of drinking water.

B. Metabolism of Temephos in Plants

Whenever a pesticide is proposed for use on agricultural crops, the Agency requires data on the fate of the pesticide in plants.

Data from a single study indicated that a variety of degradation products of temephos are formed in bean plants (Blinn, 1966, MRID #000001377). In this study, tritiated temephos, with the label ortho to the sulfide linkage, was applied to the primary leaves of three-week old lima bean plants. The plants were grown in a greenhouse and sampled 10, 21, 28 days after treatment. Treated leaves and non-treated plant parts were analyzed separately for metabolites or degradation products.

Total radioactivity on treated leaves remained constant throughout the test period; 94.9, 84.4, and 96.5% of the applied dose were recovered 10, 21, and 28 days respectively after treatment. Most of the radioactivity on treated leaves was methanol extractable; less than 4% was not extracted from the 28 day sample. Only traces of radioactivity were found in treated plant tissue indicating little or no migration from treated areas.

The distribution of temephos degradation products with time is shown in Table VII.A (Blinn, 1966, MRID #000001377). Fraction A contained temephos sulfone and the symmetrical dioxygen analog of temephos. Fraction B contained water soluble conjugates. Hydrolysis products of mixture B contained 4,4'-thiodiphenol and the sulfinyl and sulfonyl derivatives of 4,4'-thiodiphenol. The mono-oxygen analog of temephos sulfone was found in very small amounts.

TABLE VII.A. The Distribution of Temephos Degradation Products in Bean Leaves[a].

	% OF APPLIED DOSE/DAYS AFTER APPLICATION		
Residue	10	21	28
Temephos	85.5	67.3	71.5
Temephos sulfoxide	8.3	4.9	4.3
Fraction A	----	1.1	0.6
Fraction B	----	8.2	16.7

a. Blinn, 1966 (MRID #000001377).

The studies concerning temephos metabolism in plants cannot be considered adequate for the fate of temephos in plants; hence, further studies on degradation of the compound in other plants, uptake from soil, and fate of temephos under actual field conditions must be undertaken.

C. Metabolism of Temephos in Animals

When use of a pesticide on agricultural crops results in residues in products for animal feed, or when a pesticide is applied to food animals, the Agency requires data on the fate of the pesticide in food animals.

Data from a feeding study indicate that temephos and its sulfoxide are interconvertible in animals. The sulfoxide was found in milk of cows fed 20 ppm temephos, and temephos was present in milk of cows fed 5 ppm of the sulfoxide (American Cyanamid Co., 1971?, MRID #000001403).

The sulfoxide was detected in fat and kidney of cattle as a result of dermal application of 0.1% temephos emulsion. Temephos and the sulfoxide were both highest in fat (Manuel, 1975, MRID #000001344).

Chiu and Eisner (1975, MRID #000001343) sprayed three lactating goats with a ^{14}C-temephos formulation, and the goats were sacrificed 2, 7, and 14 days after treatment. The distribution of residues in the treated animals is shown in Table VII.B. The distribution of residues in solid tissues, expressed in ppm, is shown in Table VII.C. Residues in leg muscle, brain, and blood were at or below the sensitivity of the analytical method (0.01 ppm). The tissues of the goat sacrificed on day 14 were analyzed for metabolites of temephos. Residues are tabulated in Table VII.D. Other metabolites found in small amounts include: 4,4'-sulfinyldiphenol; 4,4'-sulfonyldiphenol; glycosides of the sulfinyl- and sulfonyldiphenols, and of 4,4'-thiodiphenol; and mono- and dioxygen analogs of temephos, temephos sulfoxide, and temephos sulfone.

The data on metabolism of temephos in animals are considered adequate to support the current uses of temephos. Current analytical methods for residues of temephos in animals are capable of measuring temephos, temephos sulfoxide, and temephos sulfone.

D. Analytical Methods

A method for the determination of temephos in water, oysters, rice grain, rice foliage (Blinn and Pasarela, 1966, MRID #005000857), and in cottonseed (American Cyanamid Co., 1966, MRID #000001363) has been submitted to the Agency. Samples are extracted with organic solvents, cleaned up on Florisil columns, and hydrolyzed to liberate 4,4'-thiodiphenol, which is then determined colorimetrically. The method is limited because it cannot determine temephos metabolites.

TABLE VII.B. The Distribution of Temephos Residues in Treated Lactating Goats[a].

	% INITIAL DOSE RECOVERED (mg)		
	DAY OF SACRIFICE		
Sample	2	7	14
Hair, extracted	3.10	8.60	18.70
Hide, extracted	46.80	51.70	45.20
Urine	0.80	2.70	3.70
Feces	0.02	3.00	5.60
Milk	0.05	0.05	0.20
Blood	0.05	0.03	0.04
Intestinal Wash	0.30	0.05	0.06
Rumen Content	<0.01	<0.01	<0.01
Solid Tissues	1.70	1.40	2.10
Total Recovery	52.70	67.40	75.50
Initial Dose	141 mg	123 mg	123 mg

a. Chiu and Eisner, 1975 (MRID #000001343).

TABLE VII.C. The Distribution of Temephos Residues in Solid Tissues from Treated Lactating Goats[a].

	PPM (EXPRESSED AS TEMEPHOS)		
	DAYS AFTER TREATMENT		
TISSUE	2	7	14
Back fat	0.35	0.28	0.29
Omental fat	0.06	0.38	0.36
Liver	0.05	0.04	0.06
Kidney	0.06	0.06	0.03
Heart	0.04	0.07	0.07
Tenderloin muscle	0.05	0.02	0.05

a. Chiu and Eisner, 1975 (MRID #000001343).

TABLE VII.D. The Distribution of Temephos Metabolites in Tissues of a Treated Goat Sacrificed on Day 14[a].

	% OF EXTRACTABLE RADIOACTIVITY		
TISSUE	TEMEPHOS	SULFOXIDE	SULFONE
Hair	68	6.6	0.6
Hide	67	1.6	1.5
Back fat	74	18	---
Omental fat	3	76	10
Milk	5	75	15
Liver	-	39	45
Feces	5	20	10

a. Chiu and Eisner, 1975 (MRID #000001343).

A gas-liquid chromatographic (GLC) method for determination of residues in a variety of samples has been published by Pasarela and Orloski (1973, MRID #005000638). This method has been validated for a wide variety of samples including pasture grass, alfalfa, citrus, broccoli, lettuce, snapbeans, carrots, peanuts, cottonseed, milk, cattle tissues, shrimp, oysters, crabs, fish, and water. Results from 162 samples simultaneously fortified with temephos and sulfoxide at 0.0001 ppm to 5 ppm, indicated that 1) temephos recovery averaged 87% with individual values varying from 55% to 129% and 2) recovery of temephos sulfoxide averaged 88% with values varying from 36% to 129%.

Validation data for the GLC method for various crops are also reported in other studies; alfalfa (Devine, 1969, MRID #000001382), cottonseed (Higham, 1969, MRID #000001297), citrus (Manuel and Devine, 1970, MRID #000001402), cattle tissues (Manuel, 1975, MRID #000001344; Orloski, 1969, MRID #000001392), and milk (American Cyanamid Company, 1971, MRID #000001403).

The claimed sensitivity of the GLC method for most crops is 0.05 ppm, each, for temephos and it sulfoxide; however, control values as high as 0.25 ppm for temephos and 0.44 ppm sulfoxide have been reported (Devine, 1969, MRID #000001382). The method has been tested on cottonseed in two Food and Drug Administration laboratories: successfully in one laboratory (Zee, 1971, MRID #GS0006002), and unsuccessfully in the other (Gibson, 1970, MRID #GS0006003).

Of the temephos methods available, the GLC method of Pasarela and Orloski (1973, MRID #005000638) is satisfactory for obtaining residue data. However, the method determines only temephos and temephos sulfoxide, whereas metabolism data show that temephos sulfone and other degradation products are often minor components of the residue. If these minor residues are found to be toxicologically significant, new methods or modifications to existing methods will be required for their analysis. No analytical method for temephos has been published in the Pesticide Analytical Manual (PAM).

E. Residues in Plants

Whenever a pesticide is proposed for use on an agricultural crop, the Agency requires data on the amount of residue on the crop resulting from the proposed use.

Residue data reflecting application of temephos to citrus, the only currently registered food use for temephos, have been submitted to the Agency (Manuel and Devine, 1970, MRID #000001402). Temephos formulated as Biothion® 6E or as Biothion® 50-WP was applied to oranges, lemons, and grapefruit at sites in Arizona and California. The formulations were applied by airplane or ground equipment following petal fall at 1 or 2 pounds of temephos per acre. Whole fruit was analyzed during maturation, and peel and pulp were analyzed separately at harvest for residues of temephos and temephos sulfoxide. The GLC method of

Pasarela and Orloski (1973, MRID #005000638) was used for residue analysis.

Of 22 samples of immature fruit, seven samples contained detectable (0.05 ppm) temephos up to 0.22 ppm. Three of the samples contained detectable (0.05 ppm) temephos sulfoxide up to 0.08 ppm.

Levels of temephos and temephos sulfoxide in 44 samples of peel and pulp obtained at harvest did not exceed the method sensitivity (0.05 ppm) with one exception: a sample of peel from lemons treated at 2 pounds per acre contained 0.14 ppm temephos and 0.1 ppm sulfoxide (Manuel and Devine, 1970, MRID #000001402). Data on residues in dried citrus pulp or molasses have not been submitted. Also, there are no data on residues in cover crops growing in temephos-treated orchards.

F. Storage/Stability

None of the residue studies has been accompanied by information on sample storage between harvest and analysis. A published study indicates that temephos in fish is partially oxidized to temephos sulfoxide after several weeks of storage in formalin, but is not altered after three weeks of storage at -20°C (Miles et al., 1976, MRID #005000795). The Agency requests that registrants indicate their method of sample storage, and establish that the temephos residue profile is not altered during sample storage.

G. Residues of Temephos in Animals

When use of a pesticide on agricultural crops results in residues in items of animal feed, or when a pesticide is directly applied to animals used for food, the Agency requires data on whether residues are transferred to meat, milk, poultry and eggs.

In one feeding study (Orloski, 1969, MRID #000001392), calves in groups of three were fed, via capsule, equal amounts of temephos and temephos sulfoxide at 0, 0.05, 0.15, and 0.5 ppm each in the diet. After 21 consecutive days of treatment, the animals were slaughtered and residues in muscle, liver, kidney, and fat analyzed by the GLC method of Pasarela and Orloski (1973, MRID #005000638). Temephos or the sulfoxide were not detected in any sample.

In another study (American Cyanamid Co., 1971, MRID #000001403), cows were administered temephos in the diet at 0.3, 0.5, 1.0, 2, 5, 10, 15, and 20 ppm per day. Another cow was fed temephos sulfoxide at 5 ppm per day. Treatment lasted 12 days and milk was sampled during the treatment period and up to 34 days after treatment ceased. Samples were analyzed by the Pasarela and Orloski method (1973, MRID #005000638). Temephos sulfoxide levels did not exceed the method sensitivity (0.01 ppm claimed for milk) except in the cow fed 5 ppm sulfoxide (maximum residue was 0.04 ppm) and a cow fed 20 ppm temephos (0.16 ppm residue was 0.04 ppm) and a cow fed 20 ppm temephos (0.16 ppm

maximum). Residues of temephos did not exceed 0.02 ppm in cows fed up to 0.5 ppm temephos; maximum residue in cows fed 5 and 20 ppm were 0.22 and 2.23 ppm, respectively. The cow fed the sulfoxide contained more temephos (0.18 ppm maximum) than sulfoxide (0.04 maximum). Temephos residues 34 days after cessation of treatment exceeded 0.01 ppm only in the cow fed 20 ppm of the pesticide (0.09 ppm).

Manuel (1975, MRID #000001344) sprayed Holstein steers to the point of runoff with 0.1% temephos emulsifiable concentrate. At 7, 14, 28, 42 and 56 days after treatment, animals in groups of three were slaughtered, and residues in muscle, liver, kidney, and fat were determined by the method of Pasarela and Orloski (1973, MRID #005000638). Temephos was not detected (<0.05 ppm) in any sample of muscle or liver, but was detected (0.13 and 0.08 ppm) in kidney of two of the three animals slaughtered seven days after treatment. Temephos averaged 1.19 ppm in the fat of the three animals killed at 7 days, and declined thereafter with a half-life of 6 days. The sulfoxide was not detected in muscle, liver, or kidney, but averaged 0.39 and 0.09 ppm in the fat of steers killed 7 and 14 days after treatment, respectively.

Chiu and Eisner (1975, MRID #000001343) sprayed three lactating goats with a ^{14}C-temephos formulation at a rate of 10 mg/kg and a fourth goat was sprayed with a non-radioactive formulation. Milk was collected daily during the study, and animals were sacrificed 2, 7, 14 days after treatment. Milk and tissues were analyzed for total radioactivity and results of the study are tabulated in Table VII.D.

H. Summary of Data Gaps

The data gaps for temephos regarding residue chemistry are:

1) data on the metabolism of temephos in citrus*. As a result of the required metabolism studies, if metabolites other than temephos sulfoxide are found to be present in toxicologically significant amounts then the following second tier tests are needed:

a) an analytical method, or methods, for the determination of the newly identified metabolites, and

b) data on residues of the metabolities in mature citrus and in citrus by-products (if finite residues in mature citrus are found).

---

* As an alternative to data from citrus, registrants may submit data demonstrating that metabolism of temephos in a variety of food crops is similar.

2) data on conditions of sample storage between harvest and analysis, and

3) data on residues in cover crops in temephos-treated* citrus orchards.

Registrants of new food use formulations or new uses for currently registered food use formulations must fill applicable data gaps and must provide residue data, including control values and recovery data, reflecting proposed uses.

---

* In lieu of acquiring this residue data, a restriction against grazing or feeding of livestock on temephos-treated cover crops may be imposed.

VIII. Ecological Effects

A. Ecological Effects Profile

1. Technical Temephos

Most algal growth was unaffected with 100 ppb temephos in solution even though the photosynthetic process was affected at very low levels (2-58 ppb; Derby and Ruber, 1971, MRID #000001335). Nitrogen fixation in blue-green algae was unaffected at 500 ppb (Wurtsbaugh and Apperson, 1978, MRID #005000807). Two species of Spartina were unaffected with 1 oz of temephos per acre (3 applications over 3 months) [Busch, 1975, MRID #005001001].

Laboratory studies for temephos show high acute toxicity to birds. LD_{50} values for the most representative avian species are in the neighborhood of 30 to 80 mg/kg (Tucker and Haegele, 1971, MRID #005000975 and Tucker and Crabtree, 1970, MRID #GS0006004). These studies, however, are not considered sufficient to meet guideline requirements and additional data are needed on dose response, age of test species and specific LD_{50} values. Comparative subacute dietary toxicity studies show that avian dietary LC_{50} values range from <30 to 76 ppm for passerines (bluejay, sparrow, cardinal) [Hill, 1971, MRID #000001384]; 92 to 260 ppm for upland game birds (bobwhite, quail, ringneck) [Hill et al., 1975, MRID #000002923]; and up to 894 ppm for waterfowl (mallard) [Hill et al., 1975, MRID #000022923].

The 96-hour LC_{50} values for stripped bass and rainbow trout were shown to be 1.0 and 3.49 ppm, respectively (Korn and Earnest, 1974, MRID #005000819 and Fish Pesticide Research Lab., 1971, MRID #GS0006006, respectively). Bluegill sunfish had a 96-hour LC_{50} of 21.8 ppm (Fish Pesticide Reseach Lab., 1964, 1965, 1970, MRID #GS0006005).

There is sufficient information to characterize the toxicity of temephos as moderately toxic to certain estuarine and marine organisms. The 24-hour LC_{50} and EC_{50} (impairment of predator escape response) values for the fiddler crab are 9.12 ppm and 4.31 ppm, respectively (Ward and Busch, 1976, MRID #005000957). The studies provide supplemental information* and the uses of temephos indicate the need for estuarine or marine testing in order to complete a hazard evaluation.

With acute 96-hour LC_{50} values of 0.01 to 0.082 ppm, temephos is highly toxic to freshwater aquatic invertebrates (Fish Pesticide Research Lab., 1964, 1965, 1970, MRID GS0006005).

---

* Studies were not conducted on appropriate test species, see proposed guidelines Section 163.72-3.

Field and artifical stream studies (% ai unknown) indicated that temephos is likely to cause, albeit only temporarily, toxic effects to freshwater aquatic invertebrates in the field. The impact was less severe with sequential or partial application of temephos (Ali and Mulla, 1978, MRID #005000841; Thayer and Ruber, 1976, MRID #005000947; and Ruber and Kocor, 1976, MRID #005000946).

2. Formulations

a. Granular

A 5% granular formulation of temephos was demonstrated to be highly toxic to freshwater aquatic invertebrates (McCann and Pitcher, 1975, MRID #GS0006007). McCann and Pitcher (1975, MRID #GS0006007) determined that the 48-hour LC_{50} to daphnia was 0.54 ppm in one test and 2.5 ppm in another. Field tests using 2% granular temephos at 0.1 lb ai/acre showed that temephos is likely to cause toxic effects to some estuarine and marine organisms (Ward et al., 1976, MRID #005001768). The uses of temephos indicate the need for estuarine or marine testing in order to complete a hazard evaluation.

b. Emulsifiable Concentrate

A 43% emulsifiable concentrate was tested on rainbow trout (Kennedy, 1970, MRID #000001337) and bluegill sunfish (Fish Pesticide Research Lab., 1964, 1965, 1970, MRID #GS0006005) and was shown to be of high acute toxicity to fish. The 96-hour LC_{50} for the rainbow trout (Kennedy, 1970, MRID #000001337) was 0.158 ppm and 1.14 ppm for the bluegill sunfish (Fish Pesticide Research Lab., 1964, 1965, 1970, MRID #GS0006005). It has been noted that the 4E formulation is approximately twenty times more toxic to fish that the technical.

Emulsifiable concentrate temephos (4 lbs/gal) is of low to moderate toxicity to bees (Johansen and Eves, 1965, MRID #000001361; Johansen, 1966, MRID #000001358; and Johansen, 1972, MRID #005000837). However, laboratory and field studies using various formulations showed that temephos is toxic to a number of nontarget freshwater aquatic insects (Roberts and Miller, 1970, MRID #005000774; Muirhead-Thomson, 1978, MRID #005000797; Helson and West, 1978, MRID #005000828; Muirhead-Thomson, 1978, MRID #005000798; and Wallace et al., 1973, MRID #005000821).

There is also sufficient information to characterize the 5% EC temephos as moderately toxic to certain reptiles and amphibians; however this formulation is not currently registered. The 24-hour LC_{50} (5% EC) for a frog and salamander was 4.18 ppm and 3.97 ppm, respectively (Hatorri, 1974, MRID #005003886).

B. Hazard Assessment

Due to the absence of appropriate environmental fate data and essential nontarget toxicity data, an Ecological Effects Hazard

Assessment cannot be made for manufacturing-use or end-use products of temephos. Preliminary data indicate that the end-use emulsifiable concentrate and the granular products when applied according to directions for aquatic uses as a mosquito larvicide may exceed the Restricted Use criteria [see FIFRA Section 162.11 (c)(2)(iii)(C)]*. However, the Agency cannot propose finalized classification of these products for all intended use sites until the Agency receives and reviews the requested data in Table III.A-5 in Part III.

C. Summary of Data Gaps

The data gaps for technical grade of active ingredient in temephos products are the Avian Single Dose Oral LD_{50} on one waterfowl (preferably the mallard) or on an upland game bird (preferably the bobwhite quail or the ring-necked pheasant); the 96-hour LC_{50} for shrimp, crab and an estuarine or marine fish, and the 48-hour LC_{50} for oyster embryo-larvae or the 96-hour EC_{50} shell deposition data for a representative mollusc; seed germination/seedling emergence and vegetative vigor (Tier I); and growth and reproduction of aquatic plants (Tier I).

The data gaps for the end-use temephos products (emulsifiable concentrate and granular formulations) are: the 96-hour LC_{50} for shrimp, crab and an estuarine or marine fish, and the 48-hour LC_{50} for oyster embryo-larvae or the 96-hour EC_{50} shell deposition data for a representative mollusc (both formulations); acute toxicity to aquatic invertebrates (emulsifiable concentrate only); and fish acute LC_{50} (granular only).

---

* The maximum calculated concentration (0.036 to 0.073 ppm) immediately following direct application to a 6-inch layer of water is greater than one tenth (0.0158 ppm) the acute LC_{50} (0.158 ppm) for aquatic organisms as represented by rainbow trout.

IX. Efficacy

A. Efficacy Profile

The review of temephos data and the identification of data gaps are limited to efficacy data only as it related to public health applications. In all other instances the Agency has provided for the waiver of efficacy data submission as a part of the registration process (44 FR 27932, May 11, 1979).

B. Factors Influencing Efficacy

Several factors appear to impact the efficacy of temephos. The relative susceptibility of the target species is one of the more important factors. Other factors are achieving and maintaining a toxic concentration of the active ingredient at the treatment site for a sufficient period of time to reach the required level of control.

Factors which influence achieving and maintaining a toxic concentration of the active ingredient at the treatment site are: 1) the rate of release of active ingredient (impacting on how fast the toxic concentration is achieved) and how long the toxic level is maintained (as indicated by Bender, 1964, MRID #000001245; Wall and Marganian, 1971, MRID #005000928; Mulla and Chaudhury, 1968, MRID #005000750; American Cyanamid Co., 197?, MRID #000001260; and American Cyanamid Co., 1976, MRID #000001272) and 2) the organic content and/or pollution level of the water at the treatment site (Bender, 1965, MRID #000001398 and Hagmann, 1966, MRID #005000637).

C. Use Sites

1. Non-Agricultural Aquatic Sites

The test data reviewed by the Agency clearly indicates that temephos granular products containingl, 2 or 5% active ingredient are efficacious for use as a mosquito and biting midge larvicide. The data supports the mosquito larviciding at differential rates, these range from a low rate for temporary rainpools to the high rate for highly polluted waters. Also, effective control of biting midge larvae has been demonstrated in intertidal zones of sandy beaches and marshlands.

The data also support the use of emulsifiable concentrate products containing 4 lbs active ingredient per gallon for control of mosquito larvae. The low rates are for aquatic areas of clean open water and the high rates are for aquatic areas high in organic matter. No claims are made for utilizing emulsifiable concentrate products in highly polluted waters.

Both the emulsifiable concentrate and granular products were applied by air and ground equipment. The efficacy of these products are independent of the method of application.

2. Pets and Pet Living Quarters

The data clearly indicate that direct application of products containing 2% active ingredient and pet collars containing 40.7% active ingredient are efficacious for control of fleas on cats and dogs.

No data have been submitted for evaluating temephos dust for control of fleas in pet-living quarters. However, the studies on pet flea control shows that temephos dust is effective against fleas. Consequently, dust data will be considered adequate to demonstrate the efficacy of dust against fleas on cats, dogs, and in their living quarters.

3. Target Pests

The names currently used on registered temephos labels are general terms which include many different insect species. For example, the term "gnat" includes many species and groups of innocuous flies which, in the adult stage, create problems because of their large numbers, their attraction to people or animals, and their annoyance to man and animals. The midges (Family: Chironomidae) are included in this group. However, the term "gnats" specifically excludes biting gnats (Family: Simuliidae, the buffalo gnats/black flies and turkey gnats) and biting midges (Family: Ceratopogonidae).

The pest species found on registered temephos labels that fall within the Agency's area of public health concern are as follows: mosquito larvae, punky and sandy fly larvae (biting midges, Family: Ceratopogonidae), and fleas.

a. Mosquito Larve*

The term mosquito includes several hundred insect species comprising the Family Culicidae within the Order Diptera. Laboratory studies using representative North American species have demonstrated that there is essentially no difference to the susceptibility of domestic species of mosquito larvae (Craven, and Steelman, 1968, MRID #005000634; Yap et al., 1968, MRID #005000648; Gahan et al., 1966, MRID #005000711; Jones et al.,1976, MRID #005000739; Pelsue et al., 1972, MRID #005000758; Boike and Rathburn, 1969, MRID #005000920; Mount et al., 1971, MRID #005000930; Boike and Rathburn 1972, MRID #005001115; and McFarland and Pelsue, 1971, MRID #005001767).

---

* Appropriate dilution rates [e.g., grams of product (weight) or mls of product (liquid) per square feet] must be provided on labels for spot treatments of areas such as, catch basins, containers of water (e.g., old tires) and other water-filled sunken areas where mosquitoes may breed.

Resistance to temephos and cross resistance to other organophosphate larvicides have been reported. The resistance and cross resistance varies with the locality and the mosquito species (Womeldorf et al., 1970, MRID #005000666; Lewallen and Peters, 1966, MRID #005000710; Sutherland and Evans, 1976, MRID #005000740; Gillies, et al., 1968, MRID #005000749; Womeldorf et al., 1972, MRID #005000757; Georghiou et al., 1975, MRID #005000761; Georghiou et al., 1975, MRID #005000888; and Gillies et al., 1975, MRID #005000994). In areas where phosphate resistant mosquitoes are known to exist, it has been shown that up to 0.05 lb active ingredient per acre of the emulsifiable concentrate may be required to achieve adequate control (Bowman, 1965, MRID #000001236). Therefore, the following statement should appear on all temephos emulsifiable concentrate and granular products: "The higher rate should be used in areas known to have organicphosphate resistant mosquitoes."

b. Biting Midge Larvae (Punky and Sand Fly Larvae)

Few studies are available on the efficaciousness of temephos for the control of biting midges. In field studies effective control of Culicoides melleus and C. hollensis control was obtained when 2% temephos granules were applied at the rate of 0.2 lb active ingredient per acre. These studies were conducted in the intertidal zones of sandy river banks, sandy beaches and on soft mud of tidal marshlands.

c. Fleas

Temephos, in end-use products as a 2% dust and as a 40.7% impregnated collar, has been demonstrated to be effective in controlling the flea (Ctenocephalides felis) on dogs and cats. The available data adequately demonstrate the efficaciousness of temephos against fleas infesting both dogs and cats. The dust data will also be considered adequate to demonstrate the control of fleas in both dog and cat living quarters.

Label claims concerning the duration of activity of impregnated collars are formulation dependent and will be required for each formulation registered.

Effective control has been demonstrated for these pests (refer to Table III.C-3 in Part III for references).

D. Summary of Data Gaps

There are no major data gaps pertaining to efficacy regarding temephos.

X. Confidential Annex

Persons wishing to read the contents of this annex, which contains Confidential Business Information, are requested to contact the United States Environmental Protection Agency for further details.

XI. Bibliography

A. Guide to Use of This Bibliography

1. Content of Bibliography

This bibliography contains citations of all the studies reviewed by EPA in arriving at the positions and conclusions stated elsewhere in this standard. The bibliography is divided into 3 sections: 1) citations that contributed information useful to the review of the chemical and considered to be part of the data base supporting registrations under the standard; 2) citations examined and judged to be inappropriate for use in developing the standard; and 3) standard reference material. Primary sources for studies in this bibliography have been the body of data submitted to EPA and its predecessor agencies in support of past regulatory decisions, and the published technical literature.

2. Units of Entry

The unit of entry in this bibliography is called a "study". In the case of published materials, this corresponds closely to an article. In the case of unpublished materials submitted to the agency, the Agency has sought to identify documents at a level parallel to a published article from within the typically larger volumes in which they were submitted. The resulting "studies" generally have a distinct title (or at least a single subject), can stand alone for purposes of review, and can be described with a conventional bibliographic citation. The Agency has attempted also to unite basic documents and commentaries upon them, treating them as a single study.

3. Identification of Entries

The entries in this bibliography are sorted by author, date of the document, and title. Each entry bears, to the left of the citation proper, an nine-digit identifier. This number is unique to the citations, and should be used at any time specific reference is required. This number is called the "Master Record Identifier", or "MRID". It is not related to the six-digit "Accession Number" which has been used to identify volumes of submitted data; see paragraph 4(d)(4) below for a further explanation.

4. Form of the Entry

In addition to the Master Record Identifier (MRID), each entry consists of a bibliographic citation containing standard elements followed, in the case of materials submitted to EPA, by a description of the earliest known submission. The bibliographic conventions used reflect the standards for the American National Standards Institute (ANSI), expanded to provide for certain special needs. Some explanatory notes of specific elements follow:

a. Author

Whenever the Agency could confidently identify one, the Agency has chosen to show a personal author. When no individual was identified, the Agency has shown an identificable laboratory or testing facility as author. As a last resort, the Agency has shown the first known submitter as author.

b. Document Date

When the date appears as four digits with no question marks, the Agency took it directly from the document. When a four-digit date is followed by a question mark, the bibliographer deduced the date from evidence in the document. When the date appears as (19??), the Agency was unable to determine or estimate the date of the document.

c. Title

This is the third element in the citation. In some cases it has been necessary for the Agency bibliographers to create or enhance a document title. Any such editorial insertions are contained between square brackets.

d. Trailing Parenthesis

For studies submitted to us in the past, the trailing parenthesis include (in addition to any self-explanatory text) the following elements describing the earliest known submission.

1) Submission Date. Immediately following the word 'received' appears the date of the earliest known submission.

2) Administrative Number. The next element, immediately following the word 'under', is the registration number, experimental permit number, petition number, or other administrative number associated with the earliest known submission.

3) Submitter. The third element is the submitter, following the phrase 'submitted by'. When authorship is defaulted to the submitter, this element is omitted.

4) Volume Identification. The final element in the trailing parenthesis identifies the EPA accession number of the volume in which the original submission of the study appears. The six-digit accession number follows the symbol 'CDL', standing for "Company Data Library". This accession number is in turn followed by an alphabetic suffix which shows the relative position of the study within the volume. For example,

within accession number 123456, the first study would be 123456-A; the second, 123456-B; the 26th, 123456-Z; and the 27th 123456-AA.

TEMOPHOS BIBLIOGRAPHY

Section 1

Citations Considered To Be Part Of The Data Base Supporting Registration Under The Standard

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Product Chemistry (Part IV)

MRID	Citation
000003264	American Cyanamid Company (19??) Chemical and Physical Properties of Abate. (Unpublished study received Feb 7, 1967 under 7G0566; CDL:092854-G)
000003268	American Cyanamid Company (19??) Chemical and Physical Properties of Biothion Insecticide. (Unpublished study received Apr 7, 1971 under 1F1156; CDL:093476-E)
000001254	American Cyanamid Company (1969) Modern Mosquito Control with Abate® Mosquito Larvicide and Insecticide and Cythion® Insecticide, the Premium Grade Malathion. (Unpublished study received 1969 under unknown admin. no.; CDL:128976-A)
000001379	American Cyanamid Company (1972?) [Chemical and Physical Properties of Abate Manufacturing Concentrate Insecticide and Formulations]. (Unpublished study received Jan 12, 1972 under 2270-554; CDL:025308-A)
000001222	American Cyanamid Company (1974?) Stability of Temephos Powder. (Unpublished study received Mar 6, 1974 under 241-234; CDL: 026901-M)
000001223	American Cyanamid Company (1974?) Stability of 25% Temephos Powder Concentrate. (Unpublished study received Mar 6, 1974 under 241-234; CDL:026901-N)
000001216	American Cyanamid Company (1974) Description of Processing Procedure: [Temephos]. (Unpublished study received Mar 6, 1974 under 241-234; CDL:026901-D)
000001278	American Cyanamid Company (1975?) The Specification for Raw Materials used in the Manufacture of Abate®. (Unpublished study received Nov 19, 1975 under 241-246; CDL:224604-C)
000001267	American Cyanamid Company (1976) Manufacturing Procedure for 40.7% Temephos Flea Collar. (Unpublished study received Feb 10, 1976 under 241-251; CDL:229333-H)
GS0006001	American Cyanamid Company (1978) Abate Technical Insecticide. (Letter from J. Robt Gray to Marilyn Mautz, Registration Division, EPA under 241-220; submitted by American Cyanamid Co., Princeton, N.J.)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Product Chemistry (Part IV) (Cont'd)

MRID	Citation
000001269	Fiala, R. (1975) Analytical Method for Temephos Collars. Method M-656 dated Dec 17, 1975. (Unpublished study received Feb 10, 1976 under 241-251; submitted by American Cyanamid Co., Princeton, N.J.; CDL:229333-J)
005000638	Pasarela, N.R.; Orloski, E.J. (1973) Abate insecticide. Pages 119-146, In Analytical Methods for Pesticides and Plant Growth Regulators. Edited by G. Zweig. Vol. VII. Thin-layer and Liquid Chromatography. Pesticides of International Importance. Edited by J. Sherma and G. Zweig. New York: Academic Press.

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Environmental Fate (Part V)

MRID	Citation
005000796	Daorai, A.; Menzer, R.E. (1977) Behavior of Abate in microorganisms isolated from polluted water. Archives of Environmental Contamination and Toxicology 5(2):229-240.
005000718	Drevenkar, V.; Fink, K.; Kovac, V.; Radic, B.; Vasilic, Z. (1977) The fate of pesticides in aquatic environment. III. Thin-layer chromatographic method for the determination of Abate. Arhiv za Higijenu Rada Toksikologiju 28(2):163-169.
005000794	Henry, R.A.; Schmit, J.A.; Dieckman, J.F.; Murphey, F.J. (1971) Combined high speed liquid chromatography and bioassy for the evaluation and analysis of an organophosphorus larvicide. Analytical Chemistry 43(8):1053-1057.
005001102	Steelman, C.D.; Colmer, A.R.; Cabes, L.; Barr, H.T.; Tower, B.A. (1967) Relative toxicity of selected insecticides to bacterial populations in waste disposal lagoons. Journal of Economic Entomology 60(2):467-468.
005003734	Tevan, L.; Lang, F. (1975) Az Abate szunyoglarvairtoszer hatasa a vizi oekoszisztemara. [Effect of Abate, a mosquito larva insecticide, on the ecosystem of water.] Egeszsegtudomany.
005000978	Tsai, S. (1978) Control of chironomids in milkfish (Chanos) Transactions of American Fisheries Society 107(3):493-499.
005003887	Vrochinskii, K.K. (1976) Model'naya vodnaya ekosistema kak test opredeleniya opasnosti pestitsida dlya vodoema. [Model of an aquatic ecosystem as a test for determining the extent to which a pesticide is harmful to a water body.] Izvestiya Gosudarstvennogo Nauchno-Issledovatel'skogo Instituta Ozernogo i Rechnogo Rybnogo Khozyaistva. [Bulletin of the State Scientific Research Institute of Lake and River Fisheries.] 109:88-93.
005011182	Wurtsbaugh, W.A.; Apperson, C.S. (1978) Effects of mosquito control insecticides on nitrogen fixation and growth of blue-green algae in natural plankton associations. Bulletin of Environmental Contamination and Toxicology 19:641-647.

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Toxicology (Part VI)

MRID	Citation
000001252	American Cyanamid Company (1964) Toxicity Data [on] Abate 52,160: Report No. 64-113. (Unpublished study received 1964 under unknown admin. no.; CDL:222026-A)
000001354	American Cyanamid Company (1966?) Toxicity Data: Phosphorothioic Acid, O,O,O',O' Tetramethyl O,O' Thiodi-p-Phenylene Ester (Crystalline) [and Abate Mosquito Larvicide]. (Unpublished study received Feb 3, 1967 under 7G0566; CDL:090712-A)
000001308	American Cyanamid Company (1971) Toxicity Data: Abate 2% Dusting Powder: Report No. A-71-152. (Unpublished study received Sep 1, 1972, under 241-230; CDL:002063-E)
000001353	American Cyanamid Company (1971) Toxicity Data: Biothion 6-E Insecticide: Report A-71-137. (Unpublished study received Sep 9, 1975 under 6G1666; CDL:095075-R)
000001233	American Cyanamid Company (1973) Temephos Toxicity Data: Report No. A-73-141. (Unpublished study received Mar 6, 1974 under 241-234; CDL:026901-X)
000001352	American Cyanamid Company (1974) Toxicity Data: Abate 70% Concentrate (AC2233-107): Report A-74-6. (Unpublished study received Sep 9, 1975 under 6G1666; CDL:095075-Q)
000001376	Blinn, R.C. (1966) Abate® Insecticide: The Metabolic Fate of Phenylene Phosphorothioate in the Rat: Project no. 35-54-3-11-83. (Unpublished study received Feb 7, 1967 under 7G0566; submitted by American Cyanamid Co., Agricultural Div., Princeton, N.J.; CDL:092854-H)
005000858	Blinn, R.C. (1969) Metabolic fate of Abate insecticide in the rat. Journal of Agricultural and Food Chemistry 17(1):118-122.
000001343	Chiu, T.Y.; Eisner, S.K. (1975) Temephos: Metabolism of ^{14}C-Temephos Dermally Sprayed on Lactating Goats: Project no. 2-755. (Unpublished study received Sep 9, 1975 under 6G1666; submitted by American Cyanamid Co., Princeton, N.J.; CDL: 095075-N)
000001312	Cole M.M.; Hirst, J.M.; Gilbert, I.H.; Adams, C.T. (1971) Sleeve tests of insecticides for control of body lice in 1969-70. Journal of Economic Entomology 64(3):761-762. (Also In unpublished submission received Sep 1, 1972 under 241-230; submitted by American Cyanamid Co.; Princeton, N.J.; CDL:002063-K)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Toxicology (Part VI) (Cont'd)

MRID	Citation
000001307	Fischer, J. (1974) Toxicity Data: Temephos, 25% Dust Concentrate: Report No. A-74-120. (Unpublished study received Oct 21, 1974 under 241-235; CDL:100873-A)
005000974	Gaines, T.B. (1969) Acute toxicity of pesticides. Toxicology and Applied Pharmacology 14(3):515-534.
000001368	Gaines, T.B.; Kimbrough, R.; Laws, E.R., Jr. (1967) Toxicology of Abate in laboratory animals. Archives of Environmental Health 14(?/Feb):283-288. (Also In unpublished submission received Jul 17, 1967 under 7G0566; submitted by American Cyanamid Co., Princeton, N.J.; CDL:090713-I)
000001240	Hutchison, E.B.; Levinskas, G.J. (1965) CL 52,160: Ninety-Day Repeated Feeding to Dogs: Report No. 65-19. Method dated Mar 9, 1965 entitled: Modified Michel Method for Assay of Cholinesterase Activity. (Unpublished study received May 27, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-S)
000001356	Hutchinson, E.B.; McNerney, J.M.; Levinskas, G.J. (1966) Abate (CL 52,160): Report no. 66-162. Undated method entitled: Modified Michel Method for Assay of Cholinesterase Activity. (Unpublished study received Feb 3, 1967 under 7G0566; submitted by American Cyanamid Co., Agricultural Div., Princeton, N.J.; CDL:090712-D)
000001375	Laws, E.R., Jr.; Morales, F.R.; Hayes, W.J., Jr.; Romney Joseph, C. (1967) Toxicology of Abate in volunteers. Archives of Environmental Health 14(?/Feb):289-291. (Also In unpublished submission received Feb 7, 1967 under 7G0566; submitted by American Cyanamid Co., Agricultural Div., Princeton, N.J.; CDL:092854-E)
000001239	Levinskas, G.J. (1965) Report on CL 52,160: Ninety-Day Feeding to Albino Rats: Report No. 65-18. Method dated Mar 9, 1965 entitled: Modified Michel Method for Assay of Cholinesterase Activity. (Unpublished study received May 27, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-R)
000001365	McNerney, J.M.; Levinskas, G.J. (1967) Abate Mosquito Larvicide: Single Oral Dose Toxicity to Rats and Joint Toxic Action with Other Pesticides: Report nos. 67-45 and 67-169. (Unpublished study received Jul 17, 1967 under 7G0566; submitted by American Cyanamid Co., Princeton, N.J.; CDL: 090713-B)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Toxicology (Part VI) (Cont'd)

MRID	Citation
000001388	McNerney, J.M.; Ribelin, W.E.; Levinskas, G.J. (1968) Abate® Mosquito Larvicide: Successive Generation Studies in Rats: Report no. 68-9. (Unpublished study received May 24, 1970 under 0F0913; submitted by American Cyanamid Co., Agricultural Div., Princeton, N.J.; CDL:091568-K)
005000806	Shafik, M.T. (1970) The determination of 4,4'-thiodiphenol in human and rat urine as an indication of exposure to low levels of Abate. Bulletin of Environmental Contamination and Toxicology 5(4):311-316.
005001984	Steinberg, M.; Rowe, S.E.; Boldt, R.E. (1970) Assessment of Relative Toxicity of Candidate Louse Toxicant ENT-27165 O,O,O',O'-Tetramethyl O,O'-Thiodi-P-Phenylene Phosphorothioate (Abate): Evaluation of Animal and Human Exposure Data and Recommendations Concerning a Field Test Program. Edgewood Arsenal, Md.: U.S. Army Environmental Hygiene Agency. (USAEHA study no. 33-001-70; available from: NTIS, Springfield, VA; AD 869 265)
005001274	Steinberg, M.; Godke, R.A.; Miller, T.A. (1970) Human exposure to 2 percent ENT-27165 (O,O,O',O'-tetramethyl O,O'-thiodi-p-phenylene phosphorothioate) Abate in pyrax powder and 2 percent ENT-27041 (4-benzothienyl-N-methylcarbamate) Mobam in pyrax powder evaluations of findings from volunteer use of louse toxicant with recommendation for standardization, January-March, 1970. Edgewood Arsenal, Md.: U.S. Army Environmental Hygiene Agency. (USAEHA Toxicologic and Entomologic special study no. 99-006-70/71: available from: NTIS, Springfield, VA; AD-877 508)
000001386	Underwood, P.C. (1977) Two-Year Chronic Toxicity and Carcinogenesis Study of Temephos in the Rat: Report No. 7354. (Unpublished study received Apr 4, 1978 under 241-220; prepared by Pharmacopathics Research Laboratories, Inc.; submitted by American Cyanamid Co., Princeton, N.J.; CDL:233402-B; 235935)
000001232	Wang, G.T.; Colavita, J.H.; Phillips, A.A. (1974) Temephos: Safety of 2% Temephos Powder When Applied Liberally on Cats Five Days a Week for Three Weeks: Project No. 30-0960-3-755. (Unpublished study received Mar 6, 1974 under 241-234; submitted by American Cyanamid Co., Princeton, N.J.; CDL:026901-W)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Residue Chemistry (Part VII)

MRID	Citation
000001363	American Cyanamid Company (1966) The Determination of Abate Residues in Cottonseed. Method dated Apr 12, 1966. (Unpublished study received Feb 3, 1967 under 7G0566; CDL:090712-U)
000001403	American Cyanamid Company (1971?) Cattle Feeding Studies: [Abate: Residues in Cattle Tissues and Milk]. Undated method entitled: Determination of Abate in Bovine Muscle, Kidney, and Liver; Undated method entitled: Determination of Abate in Bovine Fat; Method M-126 dated Oct 12, 1970. (Unpublished study received Apr 23, 1971 under 1F1156; CDL:090947-AD)
000001377	Blinn, R.C. (1966) Abate® Insecticide: The Metabolic Fate of O,O,O',O'-Tretramethyl O,O'-Thiodi-p-Phenylene Phosphorothioate on Bean Leaves: Project no. 35--54-3-11-83. (Unpublished study received Feb 7, 1967 under 7G0566; submitted by American Cyanmid Co., Agricultural Div., Princeton, N.J.; CDL:092854-I)
005000857	Blinn, R.C.; Paserala, N.R. (1966) Colorimetric determination of Abate residues from several environmental conditions. Journal of Agricultural and Food Chemistry 14(2):152-156.
000001343	Chiu, T.Y.; Eisner, S.K. (1975) Temephos: Metabolism of ^{14}C-Temephos Dermally Sprayed on Lactating Goats: Project no. 2-755. (Unpublished study received Sep 9, 1975 under 6G1666; submitted by American Cyanamid Co., Princeton, N.J.; CDL: 095075-N)
000001382	Devine, J.M. (1969) Abate and Abate Sulfoxide Residues in Alfafa. (Unpublished study received Jun 27, 1969 under unknown admin. no.; prepared by Syracuse Univ. Research Corp.; submitted by Allied Chemical Corp., Morristown, N.J.; CDL:120715-A)
GS0006003	Gibson, M. (1970) Pesticide Petition No. 6F0913 Method Tryout, Abate on Cottonseed. (Letter from Maxwell Gibson, DHEW, Laboratory Director, to George Beusch, Pesticide Evaluation Branch, Office of Pesticide Products Safety Bureau of Foods, BF-216, on Apr 15, 1970)
000001297	Higham, J.M. (1969) Abate® (CL 52,160) and Abate Sulfoxide (CL 105,103) Residues in Cottonseed: Report No. C-198. Undated method. (Unpublished study received Nov 24, 1969 under 241-223; submitted by American Cyanamid Co., Princeton, N.J.; CDL:026905-B)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Residue Chemistry (Part VII) (Cont'd)

MRID	Citation
000001344	Manuel, A.J. (1975) Temephos (CL 52,160) and Temephos Sulfoxide (CL 105,103) Residues in Cattle Tissues: Report no. C-650. Method M-508 dated Apr 19, 1974. (Unpublished study received Sep 9, 1975 under 6G1666; sumbitted by American Cyanamid Co., Princeton, N.J.; CDL:095075-0)
000001402	Manuel, A.J.; Devine, J. (1970) Biothion 6E: Abate® and Abate Sulfoxide Residues in Citrus (Oranges, Lemons, Grapefruit) from Studies Conducted in California and Arizona: Report No. C-246. Method M-145 dated Dec 15, 1970. (Unpublished study received Apr 23, 1971 under 1F1156; CDL:090947-AS)
005000795	Miles, J.W.; Dale, W.E.; Churchill, F.C. (1976) Storage and analysis of samples of water, fish, and mud from environments contaminated with Abate. Archives of Environmental Contamination and Toxicology 5(1):29-41.
000001392	Orloski, E.J. (1969) Abate® and Abate Sulfoxide Residues in Cattle Tissues: Report no. C-206. Method dated Aug 28, 1969 entitled: Determination of Abate in Bovine Muscle, Kidney and Liver; method dated Aug 28, 1969 entitled: Determination of Abate in Bovine Fat. (Unpublished study including letter dated Aug 28, 1969 from A. Post to Edward J. Orloski and MRI Project no. 5-582-C final report, received May 24, 1970 under 0F0913; prepared in cooperation with Midwest Research Institute, submitted by American Cyanamid Co., Agricultural Div., Princeton, N.J.; CDL:091568-AB)
005000638	Pasarela, N.R.; Orloski, E.J. (1973) Abate insecticide. Pages 119-146, In Analytical Methods for Pesticides and Plant Growth Regulators. Edited by G. Zweig. Vol. VII. Thin-layer and Liquid Chromatography. Pesticides of International Importance. Edited by J. Sherma and G. Zweig. New York: Academic Press.
GS0006002	Zee, K.T. (1971) Method Trail for Abate and Abate Sulfoxide on Cotton Seed (Unpublished study received 1971 under 0F0913)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Ecological Effects (Part VIII)

MRID	Citation
005000841	Ali, A.; Mulla, M.S. (1978) Effects of chironomid larvicides and diflubenzuron on nontarget invertebrates in residential recreational lakes. Environmental Entomology 7(1):21-27.
005001001	Busch, D.A. (1975) The failure of temephos and chloropyrifos to affect the short-term productivity of Spartina alterniflora and S. patens on a New Jersey salt marsh. Pages 50-55, In Proceedings of the Sixty-second Annual Meeting of the New Jersey Mosquito Contronl Association; Mar 11-14, 1975, Atlantic City, New Jersey. new Brunswick, N.J.: New Jersey Mosquito Extermination Association.
000001335	Derby, S.B.S.; Ruber, E. (1971) Primary production: Depression of oxygen evolution in algal cultures by organophosphorus insecticides. Bulletin of Environmental Contamination & Toxicology 5(6):533-558. (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-AE)
GS0006005	Fish-Pesticide Reserach Lab. Unpublished Reports dated 1964, 1965 and 1970. Acute Toxicity, Summary Sheets for Abate 43% and 86%. Submitted by Dr. M.T. Finley, USDI, Columbia, MO., 11-14-79.
GS0006006	Fish-Pesticide Research Lab. 1971. Acute toxicity Summary Sheet for Abate 90%. Submitted by Dr. M.T. Finley, U.S.D.I., Columbia, Mo., 11-6-79.
005003886	Hattori, K. (1974) Satchuzai difenpos (Abeito) no ryoseiryui ni taisuru dokusei. [Toxicity of difenphos (Abate) in the amphibia.] Hokkaidoritsu Eisei Kenkyusho Ho. [Report of the Hokkaido Institute of Public Health.] 24:152-153.
005000828	Helson, B.V.; West, A.S. (1978) Particulate formulations of Abate and methoxychlor as black fly larvicides: their selective effects on stream fauna. The Canadian Entomologist 110(6):591-602.
000001384	Hill, E.F. (1971) Toxicity of selected mosquito larvicides to some common avian species. Journal of Wildlife Management 35(4): 757-762. (Also In unpublished submission received Feb 11, 1976 under 201-225; submitted by Shell Chemical Co., Washington, D.C.; CDL:222966-A)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Ecological Effects (Part VIII) (Cont'd)

MRID	Citation
000022923	Hill, E.F.; Heath, R.G.; Spann, J.W.; et al. (1975) Lethal Dietary Toxicities of Environmental Pollutants to Birds: Special Scientific Report--Wildlife No. 191. (U.S. Dept. of the Interior, Fish and Wildlife Service, Patuxent Wildlife Research Center; unpublished report)
000001358	Johansen, C. (1966) Abate--Toxicity to Bees--Lygus Control. (Unpublished study received Feb 3, 1967 under 7G0566 prepared by Washington State Univ., submitted by American Cyanamid Co., Princeton, N.J.; CDL:090712-P)
000001361	Johansen, C.; Eves, J. (1965) Bee Poisoning Investigations, 1965. (Unpublished study including table 12, received Feb 3, 1967 under 7G0566; prepared by Washington State Univ., submitted by American Cyanamid Co., Princeton, N.J.; CDL:090712-S)
005000837	Johansen, C.A. (1972) Toxicity of field-weathered insecticide residues to four kinds of bees. Environmental Entomology 1(3):393-394.
000001337	Kennedy, H.D. (1970) Acute Toxicity of Pesticides to Fish: [Temephos]: Annual Progress Report: 1970. (unpublished study received Nov 26, 1971 under 241-132; prepared by U.S. Fish and Wildlife Service, Fish-Pesticide Research Laboratory, submitted by U.S. Cyanamid Co., Princeton, N.J.; CDL:129448-AG)
005000819	Korn, S.; Earnest, R. (1974) Acute toxicity of twenty insecticides to striped bass, Morone saxatilis. California Fish and Game 60(3):128-131.
GS0006007	McCann F.A and F.G Pitcher, 1975. Biological Laboratories Report. Abate. USEPA, Chemical and Biological Investigation Branch, ARC-East, Beltsville, Md. (Test No 840,836)
005000798	Muirhead-Thomson, R.C. (1978) Lethal and behavioral impact of chlorpyrifos methyl and temephos on select stream macroinvertebrates: experimental studies on downstream drift. Archives of Environmental Contamination and Toxicology 7(2):139-147.
005000797	Muirhead-Thomson, R.C. (1978) Relative susceptibility of stream macroinvertebrates to temephos and chlorpyrifos, determined in laboratory continuous-flow systems. Archives of Environmental Contamination and Toxicology 7(2):129-137.

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Ecological Effects (Part VIII) (Cont'd)

MRID	Citation
005000774	Roberts, D.R.; Miller, T.A. (1970) Effects of Polymer Formulations of Dursban and Abate on Non-target Organism Populations, April-October, 1970. Edgewood Arsenal, Md.: U.S. Army Environmental Hygiene Agency. (Entomological special study no. 31-004-71; available from: NTIS, Springfield, Va.: AD-729 342)
005000946	Ruber, E.; Kor, R. (1976) The measurement of upstream migration in a laboratory stream as an index of potential side-effects of temephos and chlorpyriphos on Gammarus fesciatus (Amphipoda, Crustacea). Mosquito News 36(4):424-429.
005000947	Thayer, A.; Ruber, E. (1976) Previous feeding history as a factor in the effects of temephos and chlorpyrifos on migration of Gammarus fasciatus (Amphipoda, Crustacea). Mosquito News 36(4):429-432.
GS0006004	Tucker, R.K.; Crabtree, D.G. (1970) Handbook of Toxicity of Pesticides to Wildlife. Pages 12-13, Bureau of Sport Fisheries and Wildlife, Denver Wildlife Research Center, Publication No. 84.
005000975	Tucker, R.K.; Haegele, M.A. (1971) Comparative acute oral toxicity of pesticides to six species of birds. Toxicology and Applied Pharmacology 20(1):57-65.
005000821	Wallace, R.R.; West, A.S.; Downe, A.E.R.; Hynes, H.B.N. (1973) The effects of experimental blackfly (Diptera, Simuliidae) larviciding with Abate, Dursban, and Methoxychlor on stream invertebrates. The Canadian Entomologist 105(6):817-831.
005000957	Ward, D.V.; Busch, D.A. (1976) Effects of temephos, an organophosphorous insecticide, on survival and escape behaviour of the marsh fiddler crab Uca pugnax. Oikos 27(2):331-335.
005001768	Ward, D.V.; Howes, B.L.; Ludwig, D.F. (1976) Interactive effects of predation pressure and insecticide (temephos) toxicity on populations of the marsh fiddler crab Uca pugnax. Marine Biology 35(2):119-126.
005000807	Wurtsbaugh, W.A.; Apperson, C.S. (1978) Effects of mosquito control insecticides on nitrogen fixation and growth of blue-green algae in natural plankton associations. Bulletin of Environmental Contamination and Toxicology 19(6):641-647.

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Efficacy (Part IX)

MRID	Citation
000001226	Ah, H.S.; Clark, J.D. (1974?) Evaluation of 2% Temephos Powder Formulation as Ectoparasiticide Against the Adult Cat Flea (Unpublished study received Mar 6, 1974 under 241-234; prepared by Univ. of Georgia, Dept. of Parasitology and Laboratory Animal Medicine Unit; submitted by American Cyanamid Co., Princeton, N.J.; CDL:02691-Q)
000001273	Ah, H.S.; Hanson, W.L. (1975) Effectiveness of CL52,160-Impregnated study received Feb 10, 1976 under 241-251; prepared by the Univ. of Georgia, Dept. of Parasitology, submitted by American Cyanamid Co., Princeton, N.J.; CDL:229333-L)
000001270	Ah, H.S.; Hanson, W.L. (1976) Effectiveness of CL52,160-Impregnated Collars in Controlling Fleas on Dogs and Cats: Part II Final Report. (Unpublished study received Aug 24, 1964 under 241-251; prepared by the Univ. of Georgia, Dept. of Parasitology, submitted by American Cyanamid Co.; Princeton, N.J.; CDL:229331-D)
000001397	American Cyanamid Company (1965) Summary of Data on Mosquito and Midge Larvae Control with Abate Granular: Table I. (Unpublished study received Dec 1, 1965 under 241-132; CDL:001793-A)
000001351	American Cyanamid Company (1969) First Periodic Report: Status of the Experimental Program Regarding Temporary Permit No. 241-EXP-47G Which Involves ULV Applications of Abate 4-E Insecticide for Control of Mosquito Larvae on Nonagricultural Lands. (Unpublished study received Jul 3, 1969 under 241-EX-47; CDL:123154-D)
000001217	American Cyanamid Company (1974?) [Efficacy Data of Temephos on Cat Flea (Ctenocephalides felis) on Both Dogs and Cats]. Summary of studies 026901-E, 026901-O through 026901-R and 026901-Y. (Unpublished study received Mar 6, 1974 under 251-234; CDL:026901-E)
000001228	American Cyanamid Company (1974?) Therapeutic and Residual Activity of 2% Temephos Powder Following a Single Topical Application on Dogs and Cats. (Unpublished study received Mar 6, 1974 under 241-234; CDL:026901-S)
000001290	American Cyanamid Company (1975?) Efficacy Studies: [Temephos]. (Unpublished study received Mar 6, 1975 under 241-EX-69; CDL:223485-C)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Efficacy (Part IX) (Cont'd)

MRID	Citation
000001262	American Cyanamid Company (1976?) [A Review of Performance Data of 40.7% Temephos Flea Collar]. Summary of Studies 229333-K thrugh 229333-Q. (Unpublished study received Feb 10, 1976 under 241-251; CDL:229333-B)
000001261	American Cyanamid Company (1976?) [Performance Data: Temephos]. Summary of Studies 229331-C through 229331-G and 229331-R. (Unpublished study received Aug 24, 1976 under 241-251; CDL:229331-B)
000001268	American Cyanamid Company (1976?) Stability: [Temephos]. (Unpublished study received Feb 10, 1976 under 241-251; CDL:229333-I)
000001260	American Cyanamid Company (1976?) Temephos in Vitro Diffusion Test. (Unpublished study received Aug 24, 1976 under 241-251; CDL:229331-A)
000001272	American Cyanamid Company (1976) Summary of Laboratory Performance of Temephos Flea collars on Dogs and Cats in Different Geographical Locations. (Unpublished study received Aug 24, 1976 under 241-251; CDL:229331-H)
000001247	Armstrong, R. (1964) Larvicide Tests on Spring Brood of Aedes Mosquitoes, 1964: [Temephos]. (Unpublished study including 2 tests and letter, received May 27, 1965 under 241-132; prepared by East Middlesex Mosquito Control Project, submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-AC)
005001004	Axtell, R.C.; Rutz, D.A.; Edwards, T.D. (1975) Chemical control of Culex pipiens quinquefasciatus in swine waste lagoons. Pages 234-235, In Proceedings of the Sixty-second Annual Meeting of the New Jersey Mosquito Control Association; Mar 11-14, 1975, Atlantic City, New Jersey. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005000730	Axtell, R.C.; Rutz, D.A.; Overcash, M.R.; Humenik, F.J. (1975) Mosquito production and control in animal waste lagoons. Pages 15-18,21, In Proceedings of the Third International Symposium on Managing Livestock Wastes; Apr 21-24, University of Illinois, Urbana-Champaign, Illinois. St. Joseph, Mich.: American Society of Agricultural Engineers. (ASAE publication PROC-275)
005000793	Baker, N.F.; Miller, J.E. (1977) Temephos collars for control of fleas on dogs and cats. American Journal of Veterinary Research 38(8):1187-1190.

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Efficacy (Part IX) (Cont'd)

MRID	Citation
000001245	Bender, E.K. (1964) Monthly Report--October, 1964: 52,160--Mosquito Larvicide--Maryland. (Unpublished study received May 27, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-AA)
000001400	Bender, E.K. (1965) Monthly Report--May, 1965: Abate Granular--Excellent on Woodland Pools for Mosquito Larvae--Delaware. (Unpublished study received Dec 1, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001793-E)
000001398	Bender, E.K. (1965) Monthly Report--September, 1965: Abate 2% G--Four Weeks Control of Culex pipiens Larvae--New Jersey. (Unpublished study received Dec 1, 1965 under 241-132; submitted by American Cyanamid Co., by American Cyanamid Co., Princeton, N.J.; CDL:001793-C)
000001276	Berger, H.; Colavita, J.H.; Miller, R.F. (1975) Temephos: Effectiveness and Safety of 40.71% Temephos Flea Collars on Cats Maintained in Runs: Project No. 340-0960-3-755. (Unpublished study received Feb 10, 1976 under 241-251; submitted by American Cyanamid Co., Princeton, N.J.; CDL:229333-P)
000001275	Berger, H.; Colavita, J.H.; Miller, R.F. (1975) Temephos: Effectiveness and Safety of 40.71% Temephos Flea Collars on Dogs Maintained in Runs: Project No. 30-0960-3-755. (Unpublished study received Feb 10, 1976 under 241-251; submitted by American Cyanamid Co., Princeton, N.J.; CDL:229333-O)
005000920	Boike, A.H., Jr.; Rathburn, C.B., Jr. (1969) Laboratory tests of the susceptibility of mosquito larvae to insecticides in Florida, 1968. Mosquito News 29(3):392-395.
005001115	Boike, A.H., Jr.; Rathburn, C.B., Jr. (1972) The susceptibility of mosquito larvae to insecticides in Florida, 1969-1971. Mosquito News 32(3):328-331.
000001236	Bowman, J.S. (1965?) Summary of Efficacy Data with Abate (EI 52160). (Unpublished study received May 27, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL: 001791-B)
005000634	Craven, B.R.; Steelman, C.D. (1968) Relative susceptibility of Psorophora confinnis (Lynch-Arribalzaga) larvae in the rice producing area of southern Louisiana to selected insecticides. Mosquito News 28(4):596-597.

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Efficacy (Part IX) (Cont'd)

MRID	Citation
000001349	Delaware Highway Department (1969?) Abate ULV Usage in Delaware. (Unpublished study received Apr 8, 1970 under 241-EX-47; prepared by Delaware Highway Dept., Mosquito Control Div., submitted by American Cyanamid Co., Princeton, N.J.; CDL: 123154-B)
000001347	Delaware, University of (1968) LV Abate Field [Tests]. (Unpublishe study including letter dated Oct 28, 1966 from R.W. Lake to John Barker, received Feb 28, 1969 under 241-EX-47; prepared by Univ. of Delaware, College of Agricultural Sciences, Dept. of Entomology and Applied Ecology, submitted by American Cyanamid Co., Princeton, N.J.; CDL:123153-D)
000001350	Delaware, University of (1970) Field Evaluation of Mosquito Insecticides: Section II-A: [Temephos]. (Unpublished study received Apr 8, 1970 under 241-EX-47; prepared by Univ. of Delaware, College of Agricultural Sciences, Dept. of Entomology and Applied Ecology, submitted by American Cyanamid Co., Princeton, N.J.; CDL:123154-C)
000001399	Dobson, J.B. (1965) Monthly Report--August, 1965: Insecticide Development: Abate. (Unpublished study received Dec 1, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001793-D)
000001225	[Fitzgerald, P.R.] (1973?) The Effect of Compound CL52,160 (Temephos) on the Flea Ctenocephalides felis on Dogs and Cats. (Unpublished study received Mar 6, 1974 under 241-234; prepared by Univ. of Illinois, submitted by American Cyanamid Co., Princeton, N.J.; CDL:026901-P)
005000711	Gahan, J.B.; Smith, C.N.; Glancey, B.M. (1966) Resistance in Florida and countermeasures involving chemicals. Mosquito News 26(3):330-337.
005000761	Georghiou, G.P.; Ariaratnam, V.; Pasternak, M.E.; Lin. C.S. (1975) Evidence of organophosphorus multiresistance in Culex pipiens quinquefasciatus and Culex pipiens pipiens in California. Pages 41-44, In Proceedings and Papers of the Annual Conference Visalia, Calif.: CMCA.
005000888	Georghiou, G.P.; Ariaratnam, V.; Pasternak, M.E.; Lin, C.S. (1975) Organophosphorus multiresistance in Culex pipiens quinquefasciatus in California. Journal of Entomology 68(4):461-467.

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Efficacy (Part IX) (Cont'd)

MRID	Citation
005000749	Gillies, P.A.; Womeldorf, D.J.; White, K.E. (1968) Cross-tolerance of California Aedes nigromaculis (Ludlow) larvae to EPN, Abate, and Dursban. Pages 85, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 36. Visalia, Calif.: CMCA.
005000994	Gillies, P.A.; Zboray, E.P.; Womeldorf, D.J.; White, K.E. (1975) Insecticide susceptibility of mosquitoes in California: occurrence of organophosphorus resistance in Culex the Forty-third Annual Conference of the California Mosquito Control Associationl Jan 26-29, 1975, Redding, California. Edited by T.D. Mulhern. Visalia, Calif.: CMCA Press.
000001234	Goulding, R.L. (1974) A study on the Effectiveness of Protex (CL 52,160) 2% Dust Against the Cat Flea Ctenocephalides felis (Bouche') on Dogs and Cats. (Unpublished study received Mar 6, 1974 under 241-234; prepared by Oregon State Univ., Dept. of Entomology, submitted by American Cyanamid Co., Princeton, N.J.; CDL:026901-Y)
000001227	Greve, J.H. (1973?) Evaluation of CL 52,160 (Temephos) Against Ctenocephalides felis in Dogs and Cats. (Unpublished study received Mar 6, 1974 under 241-234; prepared by Iowa State Univ., Dept. of Veterinary Pathology, submitted by American Cyanamid Co., Princeton, N.J.; CDL:026901-R)
005000637	Hagmann, L.E. (1966) Limited field tests on the use of Abate for the control of Culex pipiens L. Pages 173-175, In Proceedings of the Fifty-third Annual Meeting of the New Jersey Mosquito Extermination Association; March, 1966. New Brunswick, N. J.: New Jersey Mosquito Extermination Association.
000001219	Hamilton, D.R.; Schrider, M.; Cosgrove, W. (1973) Initial and Residual Activity of CL 52,160 (Temephos) in 2% Dust Formulation on Cats Experimentally Infested with Ctenocephalides felis: Project No. 3-755. (Unpublished study received Mar 6, 1974 under 241-234; submitted by American Cyanamid Co., Princeton, N.J.; CDL:026901-G)
005000739	Jones, G.E.; Carroll, D.F.; Wills, W. (1976) Susceptibility of Pennsylvania mosquito larvae to Abate, Dursban and Baytex. Pages 161-164, In Proceedings of the Sixty-third Annual Meeting of the New Jersey Mosquito Extermination Association. Vol. 63. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Efficacy (Part IX) (Cont'd)

MRID	Citation
000004967	Kauffman, G. (1964) Field Test of Liquid Insecticide Applied by Aircraft for Mosquito Control: Rept 8-B. (Unpublished study including letter dated Nov 13, 1964 from A.O. Jensen to Jim Bowman, received May 27, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-AG)
000001338	Lake, R.W.; Murphey, F.J.; Stachecki, C.J. (1967) Field trials with granular Abate for control of early spring Aedes mosquitoes. Pages 179-180, In Proceedings of Fifty-fourth Annual Meeting; 1967, Atlantic City, N.J. Atlantic City, N.J.: New Jersey Mosquito Extermination Association. (Delaware Agricultural Experiment Station miscellaneous paper No. 568; Univ. of Delaware, Dept. of Entomology and Applied Ecology. Publication No. 375; also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-AH)
005000710	Lewallen, L.L.; Peters, R.F. (1966) Control of California mosquitoes resistant to organophosphorus insecticides. Mosquito News 26(3):328-330.
000003267	Lewis, L.F.; Darrow, D.I. (1965) Results of Laboratory and Field Tests with American Cyanamid Compound 52160 (ENT 27165) against Mosquitoes. (Unpublished study received Dec 1, 1965 under 241-132; prepared by U.S. Agricultural Research Service, Entomology Research Div., Insects Affecting Man and Animals Research Branch, submitted by American Cyanamid Co., Princeton, N.J.; CDL:001793-F)
000001271	Lyles, D.I.; Johnson, W.E., Jr. (1976) Final Report: Effectiveness of the CL 52,160 Impregnated Collars in Controlling Ctenocephalides Felis on Dogs and Cats. (Unpublished study received Aug 24, 1976 under 241-251; prepared by Tuskegee Institute, Dept. of Pathology and Parasitology, submitted by American Cyanamid Co., Princeton, N.J.; CDL:229331-E)
000001317	Mallack, J.; Langford, G.S.; Sanderson, A.E., Jr. (1971?) Mosquito Control in a Landfill with Abate 4-E ULV and Observations on Fish, Shrimp and Blue Crabs in the Adjacent Estuary. (Maryland State Board of Agriculture. Contribution no. 28; unpublished study received Nov 26, 1971 under 241-132; prepared by Univ. of Maryland, State Board of Agriculture in cooperation with Maryland Dept. of Water Resources, submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-E)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Efficacy (Part IX) (Cont'd)

MRID	Citation
005001767	McFarland, G.C.; Pelsue, F.W. (1971) "Public health protection chemicals" resistance in Culex species in the Southeast Mosquito Abatement District. Pages 20-22, In Proceedings--Annual Meeting of the Utah Mosquito Abatement Association. Vol. 24.
000001224	Miller, J.E.; Baker, N.F. (1973?) Insecticidal Activity of Temephos Against Ctenocephalides felis on Dogs and Cats. (Unpublished study received Mar 6, 1974, under 241-234; prepared by Univ. of California--Davis, Dept. of Veterinary Microbiology, submitted by American Cyanamid Co., Princeton, N.J.; CDL:026901-O)
005000792	Miller, J.E.; Baker, N.F. (1975) Insecticidal activity of temephos against Ctenocephalides felis on dogs and cats. American Journal of Veterinary Research 36(9):1281-1283.
000001331	Moore, J.B.; Breeland, S.G. (1967) Field evaluation of two mosquito larvicides, Abate and Dursban, against Anopheles News 27(1):105-111. (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-Z)
005000930	Mount, G.A.; Dame, D.A.; Lofgren, C.S. (1971) Susceptibility of a Florida strain of Aedes taeniorhynchus (Weidemann) to insecticides. Mosquito News 31(3):438-440.
000004968	Mulhern, T.; Wyman, E.; Frolli,R.; Lewis, E. (1965) Field Test of Liquid Insecticide Applied by Aircraft for Mosquito Control. (Unpublished study including letter dated Nov 10, 1964 from R.F. Peters and T.D. Mulhern to Arthur O. Jensen, received May 27, 1965 under 241-132; prepared by California Dept. of Public Health, Bureau of Vector Control, Mosquito Control, submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-AH)
005000750	Mulla, M.S.; Chaudhury, M.F.B. (1968) Release of new mosquito larvicides into water from granular formulations. Pages 86-89, In Proceedings and Papers of the Conference of the California Mosquito Control Association. Vol. 36. Visalia, Calif.: CMCA.
005000758	Pelsue, F.W.; McFarland, G.C.; Gillies, P.A. (1972) Public health protection chemical resistance in larval Culex pipiens Mosquito Abatement District. Pages 25-29, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 40. Visalia, Calif.:CMCA.

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Efficacy (Part IX) (Cont'd)

MRID	Citation
000001326	Porter, C.H.; Gojmerac, W.L. (1969) Field observations with Abate and Bromophos: Their effect on mosquitoes and aquatic arthropods in a Wisconsin park. Mosquito News 29(4):617-620. (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-U)
000001218	Schrider, M.; Cosgrove, W. (1974) Activity of Temephos 2% Powder for the Control of Ctenocephalides felis Infestations on Dogs: Project No. 3-755. (Unpublished study received Mar 6, 1974 under 241-234; submitted by American Cyanamid Co., Princeton, N.J., CDL:026901-F)
005001005	Schrider, M.; Poeschel, G.; Doscher, M.; Hamilton, D. (1974) The use of temephos in controlling Ctenocephalides felis. Pages 1006-1007, In Proceedings of the Third International Congress of Parasitology; Aug 25-31, Munich, Germany. Vol.2. Munich[?]: Facta Publication.
005000784	Schrider, M.S.; Poeschel, G.P.; Annand, A.M., inventors; American Cyanamid Co., assignee (1976) Method of controlling fleas and ticks in cats and dogs. U.S. patent 3,968,207. Jul 6. 10 p. Int. Cl.2* A61K31/66.
000001249	Smith, C.N.; Roos, J. (1965) Results of Tests with American Cyanamid 52,160 from American Cyanamid Company. (Unpublished study received May 27, 1965 under 241-132; prepared by U.S. Agricultural Research Service, Entomology Research Div., Insects Affecting Man and Animals Research Branch, submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-AJ)
005000740	Sutherland, D.J.; Evans, E.S., Jr. (1976) Insecticide resistance in New Jersey mosquitoes: a five-year summary. Pages 176-181, In Proceedings of the Sixty-third Annual Meeting of the New Jersey Mosquito Extermination Association. Vol. 63. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005000925	Tawfik, M.S.; Gooding, R.H. (1970) Dursban and Abate clay granules for larval mosquito control in Alberta. Mosquito News 30(3):461-464.
000001274	Tuskegee Institute (1975) Insecticidal Activity of "Temephos Flea Collar" Against Ctenocephalides felis on Cats and Dogs: Partial Report. (Unpublished study received Feb 10, 1976 under 241-251; submitted by American Cyanamid Co.; CDL:229333-M)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

Efficacy (Part IX) (Cont'd)

MRID	Citation
005000928	Wall, W. J., Jr.; Marganian, V.M. (1971) Control of Culicoides mellus (COQ.) (Diptera: Ceratopogonidae) with granular organophosphorus pesticides and the direct effect on other fauna. Mosquito News 31(2):209-214.
000001302	Wall, W.J., Jr. (1966) Experimental Tests with Abate in Small Plots to Control C. melleus Larvae: Preliminary Report. (Unpublished study received Apr 13, 1967 under 241-151; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001812-B)
000001301	Wall, W.J., Jr. (1967) Experimental Control with Abate and Sarolex in Small Plots to Control Culicoides hollensis (Canithorax) Larvae: Preliminary Report. (Unpublished study received Apr 13, 1967 under 241-151; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001812-A)
005000757	Womeldorf, D.J.; Gillies, P.A.; White, K.E. (1972) Insecticide susceptibility of mosquitoes in California illustrated distribution of organophosphorus resistance in larval Aedes. Pages 17-21, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 40. Visalia, Calif.:CMCA.
005000666	Womeldorf, D.J.; Washino, R.K.; White, K.E.; Gieke, P.A. (1970) Insecticide susceptibility of mosquitoes in California: response of Anopheles freeborni Aitken larvae to organophosphorus compounds. Mosquito News 30(3):375-382.
005000648	Yap, H.H.; Cutkomp, L.K.; Buzicky, A.W. (1968) Insecticidal tests against Mansonia perturbans (Walker). Mosquito News 28(4):504-506.

TEMOPHOS BIBLIOGRAPHY

Section 2

Citations Judged To Be Inappropriate For Use In Developing This Standard

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005021394	Abban, E.K.; Samman, J. (1980) Preliminary observations on the effect of the insect larvicide Abate on fish catches in the River Oti, Ghana. Environmental Pollution 21(4):307-311.
005009711	Abelentseva, G.M.; Sedykh, A.S.; Popov, P.V. (1975) Toksichnost' insektitsidov dlya lichinok komara Culex pipiens molestus F. Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 44(6):687-690.
005000652	Abul-Nasr, S.; El-Sherif, S.I.; Isa, A.L.; Aly, A.S. (1975) Efficiency of certain insecticides for the control of the maizeleaf-miner Pseudonpomyza spicata Mall (Diptera, Agromyzidae). Agricultural Research Review 53(1):81-85.
005001103	Adair, H.M.; Kincade, R.J.; Laster, M.L.; Brazzel, J.R. (1967) Low-volume aerial spraying of several insecticides for cotton-insect control. Journal of Economic Entomology 60(4):1121-1127.
005001018	Ajami, A.M. (1975) Inhibitors of ester hydrolysis as synergists for biological activity of Cecropia juvenile hormone. Journal of Insect Physiology 21(5):1017-1025.
005015343	Akhundov, V.Y.; Akhundov, K.F.; Mamedova, A.P. (1974) Vliyanie abata (difosa) na obshchii sanitarnyi rezhim vodoemov. [Effect of abate (diphos) on the general sanitary conditions of water. Part 2.] Izvestiya Akademii Nauk Azerbaidzhanskoi SSR, Seriya Biologicheskikh Nauk. [Proceedings of the Academy of Sciences of the Azerbaidzhan SSR, Series on Biological Sciences.] (2):112-117.
005002674	Akhundov, V.Y.; Akhundov, K.F.; Mamedova, A.P. (1976) K obosnovaniyu predel'no dopustimoi kontsentratsii abata v vode vodoemov. [Substantiation of the maximum permissible concentration of Abate in reservoir water.] Gigiena i Sanitariya. [Hygiene and Sanitation.] (10):101-103.
005009173	Alford, A.L.; Keith, L.H. (1971) Catalog of Pesticide NMR Spectra. Athens, Ga.: U.S. Environmental Protection Agency, Southeast Water Laboratory. (Available from: NTIS, Springfield, VA; PB-206 144)
005014578	Alford, A.L.; Keith, L.H. (1971) Catalog of Pesticide NMR Spectra. Washington, D.C.: U.S. Environmental Protection Agency, Office of Research and Monitoring. (Water pollution control research series; available from: NTIS, Springfield, VA; PB-206 144)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000890	Ali, A.; Mulla, M.S. (1976) Insecticidal control of chironomid midges in the Santa Ana River water spreading system, Orange County, California. Journal of Economic Entomology 69(4):509-513.
005000892	Ali, A.; Mulla, M.S. (1977) Chemical control of nuisance midges in the Santa Ana River Basin, southern California. Journal of Economic Entomology 70(2):191-195.
005000893	Ali, A.; Mulla, M.S. (1977) The IGR diflubenzuron and organophosphorus insecticides against nuisance midges in man-made residential-recreational lakes. Journal of Economic Entomology 70(5):571-577.
005015566	Ali, A.; Mulla, M.S. (1978) Chemical control of nuisance midges in residential-recreational lakes. Folia Entomologica Mexicana 18:188.
005012191	Ali, A.; Mulla, M.S. (1978) Declining field efficacy of chlorpyrifos against chironomid midges and laboratory evaluation of substitute larvicides. Journal of Economic Entomology 71(5):778-782.
005007457	Ali, A.; Mulla, M.S.; Pfuntner, A.R.; Luna, L.L. (1978) Pestiferous midges and their control in a shallow residential-recreational lake in southern California. Mosquito News 38(4):528-535.
005000924	Altman, R.M.; Keenan, C.M.; Pearson, W.G. (1970) Control of Culicoides sand flies, Fort Kobbe, Canal Zone, in 1968. Mosquito News 30(2):235-240.
000003265	American Cyanamid Company (19??) Chemical and Physical Properties of Biothion Insecticide. (Unpublished study received May 13, 1971 under 0F0913; CDL:093212-E)
000004971	American Cyanamid Company (19??) Chemical and Physical Properties of Biothion®. (Unpublished study received Jan 14, 1969 241-EX-39; CDL:123143-E)
000002212	American Cyanamid Company (19??) Chemical and Physical Properties of Biothion®: 8.A. Data Sheet. (Unpublished study received Feb 7, 1967 under 7G0566; CDL:092854-C)
000003263	American Cyanamid Company (1964) Toxicity Data: Abate 52,160. Report No. 64-114. (Unpublished study received Oct 14, 1964 under unknown admin. no.; CDL:222025-A)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000002208	American Cyanamid Company (1964) Toxicity Data: Phosphorothioic Acid, O,O,O',O'-Tetramethyl O,O'-Thiodi-p-Phenylene Ester: Report No. 64-112. (Unpublished study received May 27, 1965 under 241-132; CDL:001791-N)
000001396	American Cyanamid Company (1965?) Chemical and Physical Properties of EI 52160. (Unpublished study received Jun 6, 1965 under 241-132; CDL:001792)
000001237	American Cyanamid Company (1965?) Feeding Studies with CL 52,160 and CL 43,913 in Chicks: [Temephos]. (Unpublished study received May 27, 1965 under 241-132; CDL:001791-I)
000001235	American Cyanamid Company (1965?) Physical and Chemical Properties: (Unpublished study received ?,?,1965 under 241-132; CDL:001791-A)
000001241	American Cyanamid Company (1965) Determination of Residues of EL 52160 in Stream and Pond Water and Mud: Report No. C-74. Method dated Mar 10, 1965 entitled: Colorimetric Procedure for the Determination of CL 52,160 Residues in Water and Mud. (Unpublished study received May 27, 1965 under 241-132; CDL:001791-T)
000001378	American Cyanamid Company (1965) Effects of Aerial Application of Granular Formulations of Three Systemic Carbamate Insecticides and Several Standard Sprays on Beneficial Predators in an Alfalfa Seed Field, Walla Walla, Washington, 1965: Table 2: (Unpublished study received 1965 under 7G0566; CDL:090712-W)
000002210	American Cyanamid Company (1966) Toxicity Data: Abate: Report No. 66-16. (Unpublished study received Feb 10, 1966 under unknown admin. no.; CDL:222022-A)
000003266	American Cyanamid Company (1966) Toxicity Data: Abate Mosquito Larvicide: Report 66-14. (Unpublished study received May 24, 1970 under 0F0913; CDL:091568-AJ)
000001393	American Cyanamid Company (1966) Toxicity Data: Solvent System for Abate Mosquito Larvicide: Report 66-15. (Unpublished study received May 24, 1970 under 0F0913; CDL:091568-AI)
000001253	American Cyanamid Company (1968) Abate®: Mosquito Larvicide and Insecticide: Technical Information. (Unpublished study received Mar 1, 1968 under unknown admin. no.; CDL:128980-A)
000001300	American Cyanamid Company (1969?) Reasonable Grounds in Support of the Petition [for Temephos]. (Unpublished study received Nov 24, 1969 under 241-223; CDL:026905-O)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000001387	American Cyanamid Company (1969?) Review of Experimental Work on the Toxicity of Biothion Insecticide (Also Known as Abate Insecticide). (Unpublished study received May 24, 1970 under 0F0913; CDL:091568-A)
000001346	American Cyanamid Company (1969?) Summary on the Toxicology of Abate. (Unpublished study received Feb 28, 1969 under 241-EX-47; CDL:123153-B)
000001296	American Cyanamid Company (1969) Summary of Lygus Control with Biothion on Cotton. (Unpublished study received Nov 24, 1969 under 241-223; CDL:026905-A)
000001348	American Cyanamid Company (1970?) Proposed Experimental Program When Applied as ULV or Granular Formulation]. (Unpublished study including experimental permit request, received Apr 8, 1970 under 241-EX-47; CDL:123154-A)
000001215	American Cyanamid Company (1970) Abate® Insecticide: Determination by Gas Liquid Chromatography in Formulations and in the Technical Product. Method M-180 dated Apr 5, 1970. (Unpublished study received Mar 6, 1974 under 241-234; CDL:026901-C)
000001304	American Cyanamid Company (1970) Determination of Abate® Insecticide in the Technical Product and Its Formulations by Ultraviolet Spectrophotometry: Recommended Method of Analysis. Method M-131 dated Nov 6, 1970. (Unpublished study received Sep 1, 1972 under 241-230; CDL:002063-B)
000001295	American Cyanamid Company (1970) Summary of Experiments Using Biothion on Non-Bearing Citrus and at Petal-Fall on Bearing Citrus to Control Citrus Thrips. (Unpublished study received 1970 under 241-223; CDL:002007-A)
000001405	American Cyanamid Company (1971?) Reasonable Grounds in Support of the Petition: [Biothion]. (Unpublished study received Apr 7, 1971 under 1F1156; CDL:093476-H)
000004970	American Cyanamid Company (1971?) Review of the Literature on the Effects of Abate on Non-Target Organisms. (Unpublished study received Mar 12, 1975 under 241-242; CDL:165068-A)
000001333	American Cyanamid Company (1971?) Review of the Literature on the Effects of Abate on Non-Target Organisms. (Unpublished study received Nov 26, 1971 under 241-132; CDL:129448-AB)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000001401	American Cyanamid Company (1971?) Summary of the Toxicity Data on Biothion: (Other Designations: Abate®; CL 52,160). (Unpublished study received Apr 23, 1971 under 1F1156; CDL: 090947-C)
000001306	American Cyanamid Company (1972?) American Cyanamid's Central Medical Department Toxicity Reports: Summary of the Toxicity Data on Biothion (Other Designations: Abate®; CL 52,160). (Unpublished study received Sep 1, 1972 under 241-230; CDL: 082063-D)
000001303	American Cyanamid Company (1972?) Chemical and Physical Properties of Abate® Insecticide, [Formulation and Stability]. (Unpublished study received Sep 2, 1972 under 241-230; CDL:002063-A).
000001305	American Cyanamid Company (1972?) Review of Performance Data: lished study received Sep 1, 1972 under 241-230; CDL:002063-C)
000001292	American Cyanamid Company (1974?) [Analytical Methods: Ultraviolet Spectrophotometry and Gas-Liquid Chromatography of Abate]. (Unpublished study received Aug 26, 1974 under 241-242; CDL:121115-B)
000001291	American Cyanamid Company (1974?) [Chemical and Physical Properties, Manufacturing and Formulations of Abate]. (Unpublished study received Aug 26, 1974 under 241-242; CDL:121115-A)
000001214	American Cyanamid Company (1974?) Chemical and Physical Properties of Temephos. (Unpublished study received Mar 6, 1974 under 241-234; CDL:026901-B)
000001293	American Cyanamid Company (1974?) Comparative Acute Toxicities of Some Insecticides. (Unpublished study including toxicology chart, received Aug 26, 1974 under 241-242; CDL:121115-C)
000001221	American Cyanamid Company (1974?) Temephos 25% Powder Concentrate. (Unpublished study received Mar 6, 1974 under 241-234; CDL:026901-L)
000001220	American Cyanamid Company (1974?) 2% Temephos Powder. (Unpublished study received Mar 6, 1974 under 241-234; CDL:026901-K)
000001294	American Cyanamid Company (1974) [Toxicity Data of Abate 70% Concentrate, Abate Technical, 90% and Panasol AN-2: Report A-74-6]. (Unpublished study received Aug 26, 1974 under 241-242; CDL:140124-A)
000001340	American Cyanamid Company (1975?) Chemical and Physical Properties of Temephos. (Unpublished study received Sep 9, 1975 under 6G1666; CDL:095075-A)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000001345	American Cyanamid Company (1975?) Preliminary Study to Show Effectiveness of Temephos for Control of Lice on Cattle. (Unpublished study received Sep 9, 1975 under 6G1666; CDL:095075-P)
000001279	American Cyanamid Company (1975?) Review of the Literature on the Effects of Abate on Non-Target Organisms. (Unpublished study received Nov 19, 1975 under 241-246; CDL:224604-E)
000001289	American Cyanamid Company (1975?) Toxicity and Safety Data [of Temephos]. (Unpublished study received Mar 6, 1975 under 241-EX-69; CDL:223485-B)
000001341	American Cyanamid Company (1975?) Toxicity Data on Formulated Product: [Temephos]. (Unpublished study received Sep 9, 1975 under 6G1666; CDL:095075-B)
000001342	American Cyanamid Company (1975) Metabolism and Tissue Residue Data: [Temephos]. Summary of studies 095075-N through 095095-O. (Unpublished study received Sep 9, 1975 under 6G1666; CDL: 095075-M)
000001265	American Cyanamid Company (1976?) Description of the Manufacturing of Temephos Technical. (Unpublished study received Feb 10, 1976 under 241-251; CDL:229333-F)
000001266	American Cyanamid Company (1976?) Ingredients in Temephos Technical. (Unpublished study received Feb 10, 1976 under 241-251; CDL:229333-G)
000001268	American Cyanamid Company (1976?) Stability: [Temephos]. (Unpublished study received Feb 10, 1976 under 241-251; CDL:229333-I)
000001264	American Cyanamid Company (1976?) Summary of the Toxicity Data on Biothion (Other Designations: Abate®; CL 52,160). (Unpublished study received on Feb 10, 1976 under 241-251; CDL:229333-E)
000001263	American Cyanamid Company (1976?) Temephos Safety and Toxicity Data. (Unpublished study received Feb 10, 1976 under 241-251; CDL:229333-C)
005003579	American Cyanamid Company. Agricultural Division (1972) Modern Mosquito Control with Cythion Insecticide, the Premium Grade Malathion. 3rd ed. Princeton, N.J.: American Cyanamid.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000001383	Amvac Chemical Corporation (1971) Release of Toxicants from Various Granular Formulations as Tested Against 4th Stage Larvae of Culex Quinquefasciatus in the Laboratory. (Unpublished study received Jul 30, 1971 under 5481-110; CDL:220268-A)
000001359	Anderson, L.D.; Atkins, E.L., Jr. (1966) 1965 Research on the Effect of Pesticides on Honey Bees. (Unpublished study including table 2, received Feb 3, 1967 under 7G0566; prepared by Univ. of California--Riverside, Entomology Dept., submitted by American Cyanamid Co., Princeton, N.J.; CDL:090712-Q)
005000875	Anderson, L.D.; Nakakihara, H. (1968) Toxicity of pesticides to corn earworm on sweet corn in Southern California, 1962-1967. Journal of Economic Entomology 61(6):1477-1482.
005003375	Andersson, S. (1975) Forosk med kemisk bekampning av betbladlusen i akerbona 1969-1970. [Field experiments with chemical control of black bean aphid in field beans 1969-70.] Meddelanden, Statens Vaxtskyddsanstalt. [Contributions, National Swedish Institute for Plant Protection.] 16(165):101-113.
005000962	Angerhofer, R.A.; Weeks, M.H.; Pope, C.R. (1978) Prenatal response of rabbits to various Abate formulations. Pharmacologist 20(3):201.
000028589	Anon. (1965) Effects of Pesticides on Fish and Wildlife: 1964 Research Findings of the Fish & Wildlife Service. N.P. (Fish and Wildlife Circular 226; p. 73 only; also In unpublished submission received Mar 18, 1980 under 6F1851; submitted Shell Chemical Co., Washington, D.C.; CDL:099341-AE)
000001380	Anon. (1968) Ground Equipment and Insecticides for Mosquito Control. Revised. Fresno, Calif: American Mosquito Control Association. (AMCA Bulletin No. 2; pages 20-21 only; also In unpublished submission received Jun 21, 1972 under 4816-133I; submitted by Fairield American Corp., Medina, N.Y.; CDL:026117-A)
000004409	Anon. (1968) Washington State Pesticide Application Handbook--1968. N.P. (p. 29 only: also In unpublished submission received Dec 29, 1970 under 707-59; submitted by Rohm & Haas Co., Philadelphia, Pa.; CDL:024155-A)
005001036	Anon. (1969) An organophosphorus round-up: part II. Food and Cosmetics Toxicology 7(1):79-83.
005003088	Anon. (1971) Public health pesticides. Pest Control 39(3):13-16,19-22,24,26,28-30,32,34,36,38,40,42,44-47,50.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005002613	Anon. (1973) Grand strategy against onchocerciasis. WHO Chronicle 27(12):540.
005002612	Anon. (1976) Dengue fever surveillance in some countries of Asia and the South-West Pacific. Weekly Epidemiological Record 51(15):113-116.
005018947	Anon. (1978) Shuang-liu-lin de can-xico ce-ding ju gi sha wen-you-chong xico-li de you-jiu. [Studies on the residue determination of Abate and its usage in mosquito larvae control.] K'un Ch'ung Hsueh Pao. [Chinese Journal of Entomology.] 21(1):49-54.
005000658	Anon. (1978) Solid progress in the Onchocerciasis Control Programme in the Volta River Basin area (OCP). WHO Chronicle 32(3):127-128.
005008225	Anon. (1979) Pesticides for use in public health: a WHO report. Pesticides 13(1):15-19.
000015963	Ansolabehere, M.J.; Puente, J.M.; Lauck, J.E.; et al. (1976) [Effectiveness of Orthene for Insect Control on Cabbage]. (Unpublished study including letters dated Oct 18, 1971 from K.A. Sorensen to Emil A. Moherek and Jul 7, 1972 from M. Shepard to Emil A. Moherek and from T.E. Skelton to Emil A. Moherek, received Nov 10, 1976 under 239-2418; submitted by Chevron Chemical Co., Richmond, Calif.; CDL:095574-B)
005000760	Apperson, C.S. (1974) Culex tarsalis resistance to organophosphorus insecticides in retrospect. Pages 106, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 42. Visalia, Calif.: CMCA.
005000763	Apperson, C.S. (1975) Biological activity of insecticides against Culicoides variipennis (Coquillett) (Diptera: Ceratopogonidae). Pages 118-119, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 43. Visalia, Calif.: CMCA.
005000901	Apperson, C.S.; Georghiou, G.P. (1974) Comparative resistance to insecticides in populations of three sympatric species of mosquitoes in the Coachella Valley of California. Journal of Medical Entomology 11(5):573-576.
005000791	Apperson, C.S.; Georghiou, G.P. (1975) Changes in cross-resistance spectrum resulting from methyl parathion selection of Culex Hygiene 24(4):698-703.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005005416	Arkatova, N.N.; Perlova, T.G.; Shvetsova-Shilovskaya, K.D. (1975) Izuchenie kinetiki gidroliza difosa. [Study of diphos hydrolysis kinetics.] Khimicheskie Sredstva Zashchity Rastenii. [Chemical Agents for Plant Protection.] 5:8-12.
000012498	Armbrust, E.J.; Gyrisco, G.G. (1965) Alfalfa Weevil Control Studies in New York. (Unpublished study received Dec 28, 1967 under 8F0648; prepared by Cornell Univ., Dept. of Entomology, submitted by Geigy Chemical Corp., Ardsley, N.Y.; CDL:091129-E)
005001106	Armstrong, R.K. (1970) Comparison of three insecticides for mosquito larval control on Okinawa. Mosquito News 30(1):1-5.
005000996	Armstrong, R.L. (1971) The use of insecticides by the East Middlesex Mosquito Control Project, Waltham, Massachusetts. Pages 47-54, In Proceedings of the Fifty-eighth Annual Meeting of the New Jersey Mosquito Extermination Association; Mar 17-19, 1971, Atlantic City, New Jersey. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005000131	Ashworth, R.B. (1973) Pesticides in public health. Chemistry and Industry (London) 14:691-693.
005000836	Asibey, E.O.A. (1977) The blackfly dilemma. Environmental Conservation 4(4):291-295.
000011695	Atkins, E.L., Jr. (1969) Biology and Control of Orangeworms Attacking Citrus, with Reference to Orange Tortrix, Pyroderces. (Unpublished study received Jan 16, 1974 under unknown admin. no.; prepared by Univ. of California--Riverside, Agricultural Experiment Station, Dept. of Entomology, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:122290-I)
000014038	Atkins, E.L., Jr. (1969) Product Performance Report. (Unpublished study received Mar 5, 1970 under 0F0956; prepared by Univ. of California--Riverside, Dept. of Entomology; submitted by Chevron Chemical Co., Richmond, Calif.; CDL:093266-AJ)
000011697	Atkins, E.L., Jr. (1970) Biology and Control of Orangeworms Attacking Citrus, with Reference to Orange Tortrix, Pyroderces. (Unpublished study received Jan 16, 1974 under unknown admin. no.; prepared by Univ. of California--Riverside, Agricultural Experiment Station, Div. of Economic Entomology, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:122290-K)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000009181	Atkins, E.L., Jr.; Anderson, L.D.; Greywood, E.A. (1969) Effect of Pesticides on Apiculture: Project No. 1499. (Unpublished study received Jul 29, 1976 under 352-342; prepared by Univ. of California--Riverside, Dept. of Entomology, submitted by E.I. du Pont de Nemours & Co., Wilmington, Del.; CDL:224800-C)
000018842	Atkins, E.L., Jr.; Anderson, L.D.; Greywood, E.A. (1969) Effect of Pesticides on Apiculture: Project No. 1499; Research Report CF-7501. (Unpublished study received May 8, 1971 under 1F1174; prepared by Univ. of California--Riverside, Dept. of Entomology, submitted by Ciba Agrochemical Co., Summit, N.J.; CDL:090973-B)
005009359	Atkins, E.L., Jr.; Anderson, L.D.; Greywood, E.A. (1970) Research on the effect of pesticides on honey bees 1968-69: part II. American Bee Journal 110(11):426-429.
005001109	Aziz, S.A. (1973) Toxicity of certain insecticide standards against the southern armyworm. Journal of Economic Entomology 66(1):68-70.
005010317	Babad, H.; Herbert, W.; Goldberg, M.C. (1968) Nuclear magnetic resonance studies of phosphorus(V) pesticides: Part I. Chemical shifts of protons as a means of identification of pesticides. Analytica Chimica Acta 41(2):259-268.
000012539	Bacon, O.G. (1965) Ecology, Biology and Control of Insects and Mites Affecting the Seed Production of Alflafa [sic], Clovers and Other Small Seeded Legumes in California: Project No. H-1735. (Unpublished study received May 8, 1970 under 0F0892; prepared by Univ. of California--Davis, Agricultural Experiment Station, Dept. of Entomology, submitted by Geigy Chemical Corp., Ardsley, N.Y.; CDL:091543-J)
000008985	Bacon, O.G. (1969) Lygus Bug Control Experiment #1. (Unpublished study received Apr 27, 1971 under 1F1159; prepared by Univ. of California--Davis, Agricultural Experiment Station, Dept. of Entomology, submitted by E.I. du Pont de Nemours & Co., Inc., Wilmington, Del.; CDL:090952-J)
000001285	Bacon, O.G.; Riley, W.D. (1965) Ecology, Biology, and Control of Insects and Mites Affecting the Seed Production of Alfalfa, Clovers and other Small Seeded Legumes in California: Project No. H-1735. (Unpublished study including table 5, received Feb 10, 1967 under unknown admin. no.; prepared by Univ. of California, Agricultural Experimental Station, submitted by American Cyanamid Co., Princeton, N.J.; CDL: 122856-E)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000664	Bailey, D.L.; LaBrecque, G.C.; Meifert, D.W.; Bishop, P.M. (1968) Insecticides in dry sugar baits against two strains of house flies. Journal of Economic Entomology 61(3):743-747.
005000926	Bailey, D.L.; LaBrecque, G.C.; Whitfield, T.L. (1970) Slow-release and emulsifiable formulations of Dursban and Abate for controlling larvae of Culex pipiens quinquefasciatus Say. Mosquito News 30(3):465-467.
005000826	Baldwin, W.F.; Gross, H.P.; Wilson, M.L.; Keill, D.J.; Stuart, R.J.; Sebastien, R.J.; Knight, A.G.; Chant, G.D.; Knight, P.A.; West, A.S. (1977) Supression of black fly populations in Deep River, Ontario. The Canadian Entomologist 109(2):249-254.
005009989	Ball, H.J. (1969) Topical toxicity of insecticides to Diabrotica 62(5):1172-1174.
005000817	Bang, Y.H.; Gratz, N.; Pant, C.P. (1972) Suppression of a field population of Aedes aegypti by malathion thermal fogs and Abate larvicide. Bulletin of the World Health Organization 46(4):554-558.
005000816	Bang, Y.H.; Pant, C.P. (1972) A field trial of Abate larvicide for the control of Aedes aegypti in Bangkok, Thailand. Bulletin of the World Health Organization 46(3):416-425.
005000808	Bang, Y.H.; Tonn, R.J. (1969) Effectiveness of different insecticides and formulations against Aedes aegypti larvae in ant traps in Bangkok, Thailand. Bulletin of the World Health Organization 41(2):320-324.
005000969	Bang, Y.H.; Tonn, R.J.; Jatanasen, S. (1972) Pilot studies of Abate as a larvicide for control of Aedes aegypti in Bangkok, Thailand. Southeast Asian Journal of Tropical Medicine and Public Health 3(1):106-115.
005000701	Barker, J.S. (1966) Abate...a new mosquito larvicide. Pages 160-162, In Proceedings of the 52nd Mid-Year Meeting, Chemical Specialties Manufacturers Association. New York: Chemical Specialties Manufacturers Assoc.
005000918	Barnes, W.W.; Webb, A.B. (1968) A field evaluation of Abate briquettes in woodland pools. Mosquito News 28(3):458-561.
005000631	Barnes, W.W.; Webb, A.B.; Savage, L.B. (1967) Laboratory evaluation of Abate and malathion insecticidal briquettes. Mosquito News 27(4):448-490.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000008236	Bass, M.H.; Guthrie, C.D.; Cobb, P.P.; et al. (1973?) Annual Report for Period Jan 1 to Dec 31, 1972: Biology and Control of Insect Pests of Soybeans and Peanuts: Ala-274, RRF Ala-295, S-74. (Unpublished study received Jun 29, 1976 under 201-347; prepared by Auburn Univ., Zoology-Entomology Dept., submitted by Shell Chemical Co., Washington, D.C.; CDL:232405-C)
000015974	Bass, M.H.; Johnson, S. (1972) Granulate Cutworm Control on Peanuts. (Unpublished study received Jun 29, 1976 under 201-347; prepared by Auburn Univ., submitted by Shell Chemical Co., Washington, D.C.; CDL:232408-O)
005001662	Beckman, H.; Garber, D. (1969) Recovery of 65 organophosphorus pesticides from Florisil with a new solvent elution system. Journal of the Association of Official Analytical Chemists 52(2):286-293.
005001012	Beckman, H.; Gauer, W.O. (1966) The characteristics and operation parameters of a thermionic emission detector, selective and sensitive to phosphorus. Bulletin of Environmental Contamination and Toxicology 1(4):149-158.
005000643	Bedford, E.C.G. (1971) The effect of Abate and Delnav on populations of red scale, Aonidiella aurantii (Mask.) in citrus orchards under integrated control. Journal of the Entomological Society of Southern Africa 34(1):159-178.
005001020	Bedford, E.C.G. (1975) The present status of the integrated control of citrus pests in South Africa. Pages 203-215, In Proceedings of the First Congress of the Entomological Society of Southern Africa; 1974, Stellenbosch, South Africa. Edited by H.J.R. Durr, J.H. Giliomee and S. Neser. Pretoria, South Africa: Entomological Society of Southern Africa.
005000831	Begum, A.; Begum, J. (1973) Relative toxicity of some insecticides to the fourth instar larvae of Aedes aegypti (L.) (Diptera, Culicidae). Dacca University Studies XXI(B):1-7.
005001014	Bellett, E.M.; Westlake, W.E.; Gunther, F.A. (1967) A method for obtaining the emission spectra of organic compounds utilizing the microwave emission detector for the gas chromatograph. Bulletin of Environmental Contamination and Toxicology 2(5):255-263.
000001246	Bender, E.K. (1964) Monthly Report--October, 1964: AC 52,160--Failure to Give Adequate Control of Mosquito Larvae--Delaware. (Unpublished study received May 27, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-AB)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000012500	Bennett, S.E. (1964) 1963-64 Progress Report on Alfalfa Weevil Research. (Unpublished study received Dec 28, 1967 under 8F0648; prepared by Univ. of Tennessee, submitted by Geigy Chemical Corp., Ardsley, N.Y.; CDL:091129-G)
005001663	Benson, W.R. (1969) The chemistry of pesticides. Annals of the New York Academy of Sciences 160(1):7-29.
005016099	Beroza, M.; Bowman, M.C. (1971) Instrumentation in determination of organophosphorus terminal residues. Pages 79-92, In Pesticide Terminal Residues: Invited Papers from the International Symposium on Pesticide Terminal Residues; Feb 17-19, 1971, Tel Aviv, Israel. Edited by A.S. Tahori. London: Butterworths.
005001019	Berry, J.F.; Cevallos, W.H. (1966) Lipid class and fatty acid composition of peripheral nerve from normal and organophosphorus-poisoned chickens. Journal of Neurochemistry 13(2):117-124.
000007046	Berry, R.E. (1970) Control of Insect Pests in Peppermint. (Unpublished study received Jun 13, 1972 under 3E1303; prepared by Oregon State Univ., Dept. of Entomology, submitted by Interregional Research Project No. 4, New Brunswick, N.J.; CDL: 092205-F)
000016032	Berry, R.E. (1972) Control of Insect Pests on Peppermint. (Unpublished study received Jun 29, 1976 under 201-347; prepared by Oregon State Univ., Dept. of Entomology, submitted by Shell Chemical Co., Washington, D.C.; CDL:232410-AP)
005004742	Bertrand, J.Y. (1976) Action de l'abate, larvicide anti-simulies, sur quelques elements de la faune non cible du Bandama (Lamto, Cote d'Ivoire). [Action of abate, an anti-sandfly larvicide, on some elements of the non-target fauna of the Bandama (Lamto, Ivory Coast.] Bulletin de l'Ecologie 7(4):445-459.
005000874	Bickley, W.E.; Ludlam, K.W. (1968) Insecticidal control of Glyptotendipes barbipes in two Maryland sewage lagoons. Journal of Economic Entomology 61(5):1454.
005000829	Birmingham, B.C.; Colman, B. (1977) The effect of two organophosphate insecticides on the growth of freshwater algae. Canadian Journal of Botany 55(11):1453-1456.
005001023	Biros, F.J.; Ryan, J.F., III (1974) Determination of metastable transitions in the mass spectra of pesticides by direct analysis of daughter ions. Pages 113-134, In Mass Spectrometry and NMR Spectroscopy in Pesticide Chemistry. New York, N.Y.: Plenum Press.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005017000	Biryukova, N.P. (1976) Dinamika difosa v vode, rosteniyakh i rybe pri obra-botke vodoyemov protiv lichinok komarov. [Dynamics of diphos in water, plants and fish in the process of treatment of reservoirs for control of mosquito larvae.] Trudy Vsesoyuznyi Nauchno-Issledovatel'skii Institut Veterinarnoi Sanitarii. Veterinary Sanitation.] 56:118-123.
005003735	Biryukova, N.P. (1977) Effektivnost' difosa protiv vshei krupnogo rogatogo skota. [The effectiveness of diphos against cattle lice.] Trudy, Vsesoyuznyi Nauchno-Issledovatel'skii Institut Veterinarnoi i Sanitarii. [Transactions, All-Union Scientific Research Institute of Veterinary Sanitation.] 58:70-73.
005021800	Biryukova, N.P.; Moryakov, Y.F.; Nepoklonov, A.A. (1976) Opredelenie ostatochnykh kolichestv difosa. [Determination of dyfos residues.] Veterinariya (Moscow). [Veterinary Science.] (11):103-104.
005007007	Biryukova, N.P.; Volik, G.N. (1976) Ispytanie difosa protiv lichinok komarov (predvaritel'noe soobshchenie). [Testing of Diphos against mosquito larvae (preliminary report).] Trudy, Vsesoyuznyi Nauchno-Issledovatel'skii Institut Veterinarnoi Sanitarii. [Transactions, All-Union Scientific Research Institute of Veterinary Sanitation.] 52:182-185.
000012501	Bishop, J.L. (1964) Alfalfa Weevil Control: Spring Treatments. (Unpublished study received Dec 28, 1967 under 8F0648; prepared by Virginia Polytechnic Institute and State Univ., Dept. of Entomology, submitted by Geigy Chemical Corp., Ardsley, N.Y.; CDL:091129-H)
005018040	Blanchet, P.F. (1979) Use of hexane for the quantitative recovery of organophosphorus pesticides from aqueous solution: a field method and a preservation technique. Journal of Chromatography 179(1):123-129.
000001367	Blinn, R.C. (1967?) Abate Insecticide: The Fate of O,O,O',O'-Tetramethyl O,O'-Thiodi-p-Phenylene Phosphorothioate on Bean Leaves. (Unpublished study received Jul 17, 1967 under 7G0566; submitted by American Cyanamid Co., Princeton, N.J.; CDL:090713-F)
005000704	Blinn, R.C. (1968) Abate insecticide - The fate of O,O,O',O'-tetramethyl-O,O'-thiodi-p-phenylene phosphorodithioate on bean leaves. Journal of Agricultural and Food Chemistry 16(3):441-445.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005018164	Blinn, R.C. (1971) Infrared microtechniques useful for the identification of pesticides at the microgram level. Pages 81-94, In Pesticide Identification at the Residue Level. Washington, D.C.: American Chemical Society Publications. (Advances in chemistry series no. 104)
005000627	Blinn, R.C.; Pasarela, N.R. (1966) Colorimetric determination of Abate residues from several environmental conditions. Agricultural and Food Chemistry 14(2):152-156.
005000777	Booth, D.C. (1975) Pesticide Monitoring of State Forest Insect Control Operations in New York State (Public Service Legislative Studies Program). Albany, New York: New York State Assembly Public ServiceLegislative Studies Program. (Available from: NTIS, Springfield, VA.; PB-268 733)
000007973	Booth, G.M. (1978) Relative Safety Evaluation of the Impact of Flit-MLO, Baygon, Fenthion, Abate, Dursban, Malathion, Parathion, Altosid, and Dimilin on the Aquatic Ecosystem. (Unpublished study received Mar 15, 1978 under 148-1259; prepared by Brigham Young Univ., Dept. of Zoology, submitted by Thompson-Hayward Chemical Co., Kansas City, Kans.; CDL:096962-C)
000001404	Bowman, J.S.; Orloski, E.J. (1966) Abate® insecticide residues in streams and ponds treated for control of mosquito larvae. Mosquito News 26(4):557-561. (Also In unpublished submission received Apr 23, 1971 under 1F1156; submitted by American Cyanamid Co., Agricultural Div., Princeton, N.J.; CDL:090947-AG)
005001037	Bowman, M.C.; Beroza, M. (1970) GLC retention times of pesticides and metabolites containing phosphorus and sulfur on four thermally stable columns. Journal of the Association of Official Analytical Chemists 53(3):499-508.
005000876	Bowman, M.C.; Ford, H.R.; Lofgren, C.S.; Weidhaas, D.E. (1968) Residues of Abate: analysis in mosquito larvae and larvicide suspensions by flame photometric gas chromatography. Journal of Economic Entomology 61(6):1586-1589.
005000904	Brandt, R.L.; Arlian, L.G. (1976) Mortality of house dust mites, Dermatophagoides farinae and D. pteronyssinus, exposed to dehydrating conditions of selected pesticides. Journal of Medical Entomology 13(3):327-331.
005000993	British Crop Protection Council (1974) Pesticide Manual: Basic Information on the Chemicals Used as Active Components of Pesticides. Edited by H. Martin and C.R. Worthing. 4th ed. London, England: British Crop Protection Council.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005021406	British Standards Institution (1974) Recommended Common Names for Pesticides. London, England: British Standards Institution. (British Standard 1831 (1969), supplement no. 3 (1974))
005000914	Brooks, C.D.; Smith, E.A.; Miles, J.W. (1967) Accumulative effects of repeated Abate granular treatments in water storage drums. Mosquito News 27(2):164-171.
005000714	Brooks, G.D.; Schoof, H.F.; Smith, E.A. (1966) Evaluations of five formulations of Abate against Aedes aegypti, Savannah, Georgia, 1965. Mosquito News 26(4):580-582.
005000630	Brooks, G.D.; Smith, E.A.; Miles, J.W. (1967) Accumulative effects of repeated Abate granular treatments in water storage drums. Mosquito News 27(2):164-171.
005002139	Brown, A.W.A. (1972) Methods at present available for Stegomyia control. Cahiers ORSTOM, Serie Entomologie Medicale et Parasitologie X(2):163-164.
005000736	Brown, A.W.A. (1974) Chemical control. Pages 10-15, In Proceedings of the Annual Meeting of the Utah Mosquito Abatement Association. Vol. 27.
005009664	Brown, T.M.; DeVries, D.H.; Brown, A.W.A. (1978) Induction of resistance to insect growth regulators. Journal of Economic Entomology 71(2):223-229.
005002914	Buei, K.; Ito, S. (1974) [On the susceptibility of insecticides against the vector mosquito of Japanese encephalitis, Culex tritaeniorhynchus summorosus Dyar.] Eisei Dobutsu.
005002917	Buitendag, C.H. (1976) Sitruswasdopluis: verdere aspekte in verband met insekmiddelbeheer. [Citrus waxy scale: further aspects in relation to insecticide control.] Citrus and Subtropical Fruit Journal (510):5-10.
000006853	Burts, W. (1965) Report No. Port 65-66. (Unpublished study received Jul 5, 1968 under 8F0668; submitted by Chipman Chemical Co., Inc., Burlingame, Calif.; CDL:092027-BM)
005000998	Busch, D.A. (1974) Methods of evaluating effects of mosquito larvicide (Abate) on non-target salt-marsh organisms. Pages 125-129, In Proceedings of the Sixty-first Annual Meeting of the New Jersey Mosquito Extermination Association; Mar 13-15, 1974, Atlantic City, New Jersey. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000979	Butcher, J.; Boyer, M.; Fowle, C.D. (1975) Impact of Dursban and Abate on microbial numbers and some chemical properties of standing ponds. Water Pollution Research in Canada 10:33-41.
005014420	Butler, G.L. (1977) Algae and pesticides. Pages 19-62, In Residues of Pesticides and Other Contaminants in the Total Environment. Edited by F.A. Gunther. New York: Springer. (Residue reviews, vol. 66)
005000884	Cameron, E.A.; Campbell, R.L.; Adams, L.E. (1973) Insecticidal control of Adelges abietis in Pennsylvania. Journal of Economic Entomology 66(3):811-812.
005000840	Campbell, R.C.; Denno, R.F. (1976) The effect of temephos and chlorpyrifos on the aquatic insect community of a New Jersey salt marsh. Environmental Entomology 5(3):477-483.
005000780	Cardarelli, N.F.; Hess, J.C., inventors; B.F. Goodrich Co., assignee (1970) Larvizide Kautschukmischungen und ihre Verwendung. [Larvicidal rubber compositions and their application.] German (Fed. Rep.) offenlegungsschrift 1,954,807. Jul 2. 25 p. Int. Cl. 2* A01n.17/08; Int. Cl. 2* C07f.9/18.
005000929	Carestia, R.R.; Roan, C.C.; Loy, V.A. (1971) Field evaluation of mosquito control in sewage oxidation ponds using plaster of paris impregnated briquettes. Mosquito News 31(4):492-498.
005000741	Carey, W.E.; Iadevaia, R.; Van de Burgt, L.; Helrich, K. (1976) Abate residues in salt marsh substrates. Pages 186-189, In Proceedings of the Sixty-third Annual Meeting of New Jersey Mosquito Extermination Association. Vol. 63. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005003870	Carlson, E.C. (1967) Control of sunflower moth larvae and their damage to sunflower seeds. Journal of Economic Entomology 60(4):1068-1071.
005004410	Carlson, E.C. (1968) Control of sunflower moth larvae and damage to sunflower seeds. California Agriculture 22(7):6-8.
005001035	Casida, J.E.; Lykken, L. (1969) Metabolism of organic pesticide chemicals in higher plants. Annual Review of Plant Physiology 20:607-636.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000771	Catangui, F.P.; Gutierrez, R.C.; Abrenica, D.D.; Collins, R.T. (1973) The effects of Abate as a larvicide against Anopheles 38-45, In Vector Control in Southeast Asia. Proceedings of the First Southeast Asian Ministers of Education Organization Workshop; Aug 17-18, 1972, Singapore. Edited by Chan Yow-Cheong, Chan Kai-Lok and Ho Beng-Chuan. Bangkok, Thailand: Southeast Asian Ministers of Education Organization.
005002853	Cavalcaselle, B. (1974) Prove di lotta contro le larve di Neodiprion sertifer (Geoff.) con alcuni insetticidi chimici e con una sospensione virale. [Tests in the control of larvae of Neodiprion sertifer (Geoff.) with some chemical insecticides and a viral suspension.] Cellulosa e Carta. [Cellulose and Paper.] 25(12):27-32.
005000856	Cecil, H.C.; Bitman, J.; Harris, S.J. (1973) Effects of different classes of pesticides on pentobarbital anesthesia and toxicity in Japanese quail. Journal of Agricultural and Food Chemistry 21(1):114-117.
005006880	Center for Disease Control (1972) Public health pesticides. Pest Control 40(4):35-38,45-46,48,50,52,54-65.
005010816	Chalfant, R.B. (1969) Control of the cabbage looper and diamondback moth caterpillar on cabbage and collards in southern Georgia. Journal of the Georgia Entomological Society 4(2):65-74.
005000706	Champ, B.R.; Campbell-Brown, M.J. (1970) Insecticide resistance in
005010816	Chalfant, R.B. (1969) Control of the cabbage looper and diamondback moth caterpillar on cabbage and collards in southern Georgia. Journal of the Georgia Entomological Society 4(2):65-74.
005000706	Champ, B.R.; Campbell-Brown, M.J. (1970) Insecticide resistance in Australian Tribolium castaneum (Herbst) (Coleoptera, Tenebrionidae)-II: malathion resistance in eastern Australia. Journal of Stored Products Research 6(2):111-131.
005000967	Chance, M.M. (1970) A review of chemical control methods for blackfly larvae (Diptera: Simuliidae). Quaestiones Entomologicae 6(3):287-292.
000022420	Chapman, R.K. (1967) [Leafhopper and Flea Beetle Control on Potatoes and Beans]. (Unpublished study received May 2, 1970 under 0F0886; submitted by E.I. du Pont de Nemours & Co., Inc., Wilmington, Del.; CDL:091530-W)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000012459	Chapman, R.K. (1967) Cooperator Research Report: Potato Leafhopper and Flea Beetle Control on Potatoes. (Unpublished study received Nov 10, 1972 under 3F1323; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:093552-AS)
005005002	Chauvet, G.; Ravaonjanahary, C.; Brunhes, J. (1971) Sensibilite et resistance a divers insecticides organophosphores chez Culex insecticides of Culex pipiens fatigans Wied. in urban areas of Madagascar.] Comptes Rendus des Seances de la Societe de Biologie de Madagascar 165(2):444-448.
005016999	Chen, Y.P.; Sudderuddin, K.I. (1978) Toxicological studies of insecticides on Culex quinquefasciatus Say and Aedes aegypti (L.). Southeast Asian Journal of Topical Medicine and Public Health 9(3):378-383.
005000822	Cheng, H.H. (1973) Further field evaluation of insecticides for control of the dark-sided cutworm (Lepidoptera: Noctuidae) on tobacco in Ontario. The Canadian Entomologist 105(10):1351-1357.
005000769	Cherry, E.T. (1974) Effect of plant bugs on cotton in west Tennessee. Pages 21-23, In Tennessee Farm and Home Science, Progress Report. Vol. 90. Knoxville, Tenn.: University of Tennessee, Agricultural Experiment Station.
005000697	Cheymol, J.; Chbier, P.; Thanh, T.N.; Thizy, A.; Demozay, D.; Pillon, D. (1967) Toxicite comparee (Mammiferes, Insectes, Acariens) de nouveaux derives S-substitues des acides S,S-dimethylphosphorique et thiophosphorique. [Comparative toxicity (Mammifera, Insecta, Acarina), of new S substituted derivatives of S,S-demethylphosphoric and thiophosphoric acids.] Comptes Rendus Hebdomadaires des Seances de l'Acadamie des Sciences, Serie D: Sciences Naturelles 265(3):291-294.
000006872	Chipman Chemical Company, Incorporated (1965) Report No. BB 65-131. (Unpublished study received Jul 5, 1968 under 8F0668; CDL: 092027-CM)
005000642	Chiu, T.Y.; Blinn, R.C. (1975) Metabolism of O,O,O',O'-tetramethyl-O,O'-thiodi-p-phenylene phosphorothioate (temephos) dermally sprayed onto lactating goats. Pages 2, In Abstracts of Papers, 170th National Meeting, American Chemical Society; Aug 24-26, 1975, Chicago, Illinois. Baltimore, Md.: Port City Press.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005021679	Chung, J.H. (1978) Yugi inje nongyaguro inhan pihae. [Hazards due to organic phosphorus pesticides.] Taehan Uihak Hyophoe Chapchi. [Journal of the Korean Medical Association.] 21(5):359-364.
000014256	Ciarletta, J.A.; Smith, R.L.; Sleesman, J.; et al. (1975) Monitor®: Efficacy, Phytotoxicity and Yield Effects on Peppers. (Unpublished study including report nos. 31525, 31659, 31726, 31735, 32153, 36254, 36258, 36259, 36337, 36342, 36345, 36346, 39118, 39140, 39799, 40064, 40729, 40946, 41098, 42715, 42880, 42916, 43225, 43379 and 44930, received Aug 24, 1978 under 3125-280; submitted by Mobay Chemical Corp., Agricultural Div., Kansas City, Mo.; CDL:097315-A)
005000885	Cline, R.E.; Hall, M.H. (1973) Larvicidal activities of N,N'-dialkylalkylenediamines and other fatty diamines in oil films and water dispersions. Journal of Economic Entomology 66(3):697-702.
005000886	Cline, R.E.; Miles, J.W.; Churchill, F.C., II (1974) Mosquito larvicidal activities of oil solutions and emulsifiable formulations containing lipophilic N-alkyl and N,N'-dialkylalkylenediamines. Journal of Economic Entomology 67(5):625-628.
000012522	Clower, D.F. (1965) [Insect Control Experiments]. (Unpublished study received Dec 28, 1967 under 8F0648; prepared by Northeast Louisiana Experiment Station, submitted by Geigy Chemical Corp., Ardsley, N.Y.; CDL:091129-AO)
005001011	Coahran, D.R. (1966) A sodium flame detector of increased stability for phosphorus-containing pesticides. Bulletin of Environmental Contamination and Toxicology 1(4):141-148.
005000977	Cohen, S.D.; Murphy, S.D. (1970) Dose and time relationships for carboxylesterase inhibition and malathion potentiation by EPN, Abate, and parathion. Toxicology and Applied Pharmacology 17(1):303-304.
005000906	Cohen, S.D.; Murphy, S.D. (1971) Carboxylesterase inhibition as an indicator of malathion potentiation in mice. Journal of Pharmacology and Experimental Therapeutics 176(3):733-742.
005000966	Cohen, S.D.; Murphy, S.D. (1972) Inactivation of malaoxon by mouse liver. Proceedings of the Society for Experimental Biology and Medicine 139(4):1385-1389.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000733	Cole, M.M.; Clark, P.H.; Grothaus, R.H. (1973) Some candidate louse powders. Pages 179-180, In Proceedings of the International Symposium on the Control of Lice and Louse-born Diseases; Dec 4-6, 1972, Washington, D.C. Washington, D.C.: Pan American Health Organization. (Scientific publication no. 263)
005000646	Cole, M.M.; Hirst, J.M.; Gilbert, I.H.; Adams, C.T. (1971) Sleeve tests of insecticides for control of body lice in 1969-70. Journal of Economic Entomology 64(3):761-762.
000001310	Cole, M.M.; Hirst, J.M.; McWilliams, J.G.; Gilbert, I.H. (1969) Sleeve tests of insecticidal powders for control of body lice. Journal of Economic Entomology 62(1):198-200. (Also In unpublished submission received Sep 1, 1972 under 241-230; submitted by American Cyanamid Co., Princeton, N.J.; CDL: 002063-I)
005000933	Coombes, L.E.; Lee, J.T.; Meisch, M.V. (1973) Effectiveness of ground ULV aerosols against larvae of Psoropohora
005000940	Cooney, J.C.; Pickard, E. (1974) Field tests with Abate and Dursban insecticides for control of floodwater mosquitoes in the Tennessee Valley region. Mosquito News 34(1):12-22.
005016727	Coosemans, M.; Mouchet, J.; Dejardin, J.; Barathe, J.; Sannier, C. (1978) Doses diagnostiques de la resistance d' Aedes to determine the resistance of Aedes aegypti to organophosphorus insecticides.] Annales de la Societe Belge de Medecine Tropicale 58(3):219-230.
000001334	Cope, O.B. (1965) Sport fishery investigations. Pages 51-55, In Effects of Pesticides on Fish and Wildlife: 1964 Research Findings of the Fish and Wildlife Service. By U.S. Dept. of Interior, Fish and Wildlife Service. Washington, D.C.: U.S. D.I., Fish and Wildlife Service. (Circular 226; also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-AD)
005002495	Cousserans, M. (1977) Le probleme de la demoustication. [Mosquito control problems.] Travaux de la Societe de Pharmacie de Montpellier 37(1):45-53.
005000866	Craven, B.R.; Steelman, C.D. (1968) Studies on a biological and a chemical method of controlling the dark rice field mosquito in Louisiana. Journal of Economic Entomology 61(5):1333-1336.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005019280	Cress, D.; Ruppel, R.; Wallner, W.; Jones, A.; Bird, G.; Meggitt, W.; Putnam, A. (1976) Pesticides Manual: Classification, Toxicities, Formulation, Handling, Application. East Lansing, Mich.: Michigan State University, Cooperative Extension Service. (Michigan Cooperative Extension Service bulletin no. E-751)
005019989	Crisp, C.E. (1972) The molecular design of systemic insecticides and organic functional groups in translocation. Pages 211-264, In Pesticide Chemistry, Proceedings of the International IUPAC Congress of Pesticide Chemistry, 2nd; Feb 22-26, 1971, Tel Aviv, Israel. Vol. 1. Edited by A.S. Tahori. New York: Gordon & Breach.
005000724	Cyanamid International, Research and Development Department (1974) Abate: Insecticide in Public Health Programs. Wayne, N.J.: Cyanamid International.
005000853	Dale, W.E.; Miles, J.W. (1969) Gas chromatographic determination of Abate using flame photometric and electron-capture detectors. Journal of Agricultural and Food Chemistry 17(1):60-62.
005000861	Dale, W.E.; Miles, J.W. (1976) Partition chromatographic separation of pesticide residues from fats. Journal of the Association of Official Analytical Chemists 59(1):165-168.
005000728	Dale, W.E.; Miles, J.W.; Guerrant, G.O. (1975) Monitoring of residues of Abate in streams treated for Simulium control. Pages 780-783, In Environmental Quality and Safety, Supplement. Vol. III: Pesticides. International Union of Pure and Applied Chemistry, Third International Congress; Jul 3-9,1974, Helsinki, Finland. Edited by Frederick Coulston and Friedhelm Korte. Stuttgart, West Germany: George Thieme Publishers.
005000990	Dang-Que, N.; Van-Rung, D.; Chow, C.Y. (1974) Aedes mosquito surveillance in the Republic of Vietnam. Southeast Asian Journal of Tropical Medicine and Public Health 5(4):569-573.
005000991	Danish Pest Infestation Laboratory (1971) Statens Skadedyrlaboratorium Arsberetning-Danish Pest Infestation Laboratory Annual Report 1970. Lyngby, Denmark: Statens Skadedyrlaboratorium.
005000992	Danish Pest Infestation Laboratory (1973) Statens Skadedyrlaboratorium Arsberetning-Danish Pest Infestation Laboratory Annual Report 1972. Lyngby, Denmark: Statens Skadedyrlaboratorium.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000941	Darwazeh, H.A.; Mulla, M.S. (1974) Toxicity of herbicides and mosquito larvicides to the mosquito fish Gambusia affinis . Mosquito News 34(2):214-219.
005005417	Davidson, L.M.; Peairs, R.H. (1966) Applied insect control: chemical. Pages 72-100, In Insect Pests of Farm, Garden and Orchard. 6th ed. New York: John Wiley.
005007499	Davies, J.B.; Le Berre, R.; Walsh, J.F.; Cliff, B. (1979) Onchocerciasis and Simulium control in the Volta River Basin. Mosquito News 38(4):466-472.
005005418	Dejoux, C.; Elouard, J.M. (1977) Action de l'Abate sur les invertebres aquatiques: cinetique de decrochement a court et moyen terme. [Action of Abate on aquatic invertebrates: short and moderate term detachment kinetics.] Cahiers ORSTOM, Serie Hydrobiologie XI(3):217-230.
005000719	De Lavaur, E.; Arnold, A. (1977) Pesticides and wildlife: results of toxicological analysis of game from 1974 to 1976. A translation of: Phytiatrie-Phytopharmacie 26(3):159-168.
000001315	Delaware, University of (1969) Effects of Abate (ULV) on Salt Marsh Organisms. (Unpublished study received Nov 26, 1971 under 241-132,; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-C)
000006746	DeLay, D.L.; Tuttle, D. (1967) Report No. PA 67-59. (Unpublished study received Feb 16, 1970 under 0F0948; prepared in cooperation with Univ. of Arizona, submitted by Rhodia, Inc., New Brunswick, N.J.; CDL:092026-BG)
000006234	DeLay, D.L.; Tuttle, D.M. (1969) Zolone EC--Lemons: Final Report No. PA 69-40. (Unpublished study received Aug 19, 1971 under 2F1193; prepared in cooperation with Univ. of Arizona, Yuma Experiment Station, submitted by Rhodia, Inc., New Brunswick, N.J.; CDL:093518-M)
000006098	DeLay, D.L.; Tuttle, D.M. (1970) Zolone EC--Lemon: Final Report No. PA 70-48. (Unpublished study received Aug 19, 1971 under 2F1193; prepared in cooperation with Univ. of Arizona, Yuma Experiment Station, submitted by Rhodia, Inc., New Brunswick, N.J.; CDL:093518-T)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000006099	DeLay, D.L.; Tuttle, D.M. (1970) Zolone EC--Lemons: Final Report No. PA 70-49. (Unpublished study received Aug 19, 1971 under 2F1193; prepared in cooperation with Univ. of Arizona, submitted by Rhodia, Inc., New Brunswick, N.J.; CDL:093518-U)
005001000	Denno, R.F. (1974) Initial studies of Abate in a salt marsh ecosystem: sampling the non-target insect component. Pages 138-144, In Proceedings of the Sixty-first Annual Meeting of the New Jersey Mosquito Extermination Association; Mar 13-15, 1974, Atlantic City, New Jersey. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005001021	Depalma, A.E.; Plummer, N. (1968) Industrial exposure groups and controls. Pages 1-15, In American Cyanamid Contract Study: Work Unit 1. Wayne, New Jersey: American Cyanamid Company.
005000982	DePew, L.J. (1975) Field evaluation of insecticides to control pale western cutworm in winter wheat. Journal of Economic Entomology 68(1):85-86.
000001394	Devine, J.W. (1970) Abate & Abate Sulfoxide Residues in Alfafa; Period Covered: 2/22/70--2/27/70. (Unpublished study received Mar 9, 1970 under unknown admin. no.; prepared by Syracuse Univ. Research Corp., submitted by American Cyanamid Co., Princeton, N.J.; CDL:121114-A)
005000942	Didia, V.; LaSalle, R.; Liem, K. (1975) The effects of Abate 2G mosquito larvicide on selected non-target organisms collected from forested temporary pools. Mosquito News 35(2):227-228.
005011888	Dimitri, M.S.; Falkehag, S.I., inventors; Westvaco Corp., assignee (1975) Composites of lignin and biologically active materials. U.S. patent 3,929,453. Dec 30. 4 p. Int. Cl. 2* A 01N 9/12; U.S. Cl. 71/101.
005000877	Dixon, R.D.; Brust, R.A. (1971) Field testing of insecticides used in mosquito control, and a description of the bioassay technique used in temporary pools. Journal of Economic Entomology 64(1):11-14.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005001009	Dolan, J.W.; Seiber, J.N. (1977) Chlorine-selective detection for liquid chromatography with a Coulson electrolytic conductivity detector. Analytical Chemistry 49(2):326-331.
000011863	Dominick, C.B. (1965) Insect Control in Tobacco. (Unpublished study received Dec 28, 1971 under 100-501; prepared by Virginia Polytechnic Institute and State Univ., Bright Tobacco Research Station, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL: 000539-B)
005008368	Done, A.K. (1979) The toxic emergency: the great equalizers II. Anticholinesterases. Emergency Medicine 11(6):167-168,173-175.
005000997	Dorer, R.E. (1973) The use of plaster of paris insecticidal briquettes in Virginia. Pages 142-143, In Proceedings of the Sixtieth Annual Meeting of the New Jersey Mosquito Extermination Association; Mar 14-16, 1973, Atlantic City, New Jersey. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
000012505	Dorsey, C.K.; Quinn, D.O. (1964) Alfalfa Weevil Control. (Unpublished study received Dec 28, 1967 under 8F0648; prepared by West Virginia Univ., Agricultural Experiment Station, Dept. of Entomology, submitted by Geigy Chemical Corp., Ardsley, N.Y.; CDL:091129-L)
005000959	Drake, J.L.; Crowder, L.A.; Ware, G.W. (1971) Effects of X-irradiation on the toxicity of organophosphate insecticides to the house fly. Pesticide Biochemistry and Physiology 1(3/4):373-380.
005000732	Dukhanina, N.N.; Quadeer, A.; Polevoi, N.I.; Yakoobi, G.H.; Artemiev, M.M.; Nushin, M.K.; Ermishev, Y.V.; Pivnenko, V.G.; Flerova, O.A.; Jalilov, H.D.; Kulkova, T.A.; Pereyaslavzav, G.P.; Parvin Nushin, A.; Mohammad, K.; Basir, A.; Jamil, T.U. (1973) Epidemiological justification for supplementary anti malaria measures in the Malaria Eradication Programme of northeastern Afghanistan. Pages 209-210, In Ninth International Congress on Tropical Medicine and Malaria; Oct 14-21, 1973, Athens, Greece. Vol. II. Abstracts of Communications. Edited by George J. Papaevangelou. Athens, Greece: Congress Publications Committee.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005004740	Duport, M.; Sandescu, I.; Paun-Durbaca, S.; Gheorghiu, G.; Visan, M. (1971) Etude comparative de l'efficacite de divers insecticides sur des larves de Culicidae . Experiences de laboratoire et de terrain. [Comparative study of the effectiveness of different insecticides on Culicidae larval forms. Laboratory and field experiments.] Archives Roumaines de Pathologie Experimentale et de Microbiologie 30(3/4):427-434.
005000922	Eliason, D.A.; Kilpatrick, J.W.; Babbitt, M.F. (1970) Evaluation of the effectiveness of the ultra low volume aerial application of insecticides against Aedes aegypti (L.) in Florida. Mosquito News 30(3):430-436.
005001278	Ellis, R.A. (1976) Emergency measures and mosquito control operations during the 1975 western encephalomyelitis outbreak in Manitoba. Canadian Journal of Public Health, Supplement one 67:59-60.
005019326	Ennin, M.A.; Franklin, C.S. (1979) Some sub-cellular effects of an organophosphorus insecticide, Abate. British Journal of Pharmacology 66(1):72-73.
000001238	Esposito, R.G.; Levinskas, G.J. (1964) CL 52,160: Three-Week Dermal Applications to Rats: Report No. 64-123. (Unpublished study received May 27, 1965 under 241-132; submitted by American Cyanamid Co., Princeton N.J.; CDL:001791-Q)
000001280	Esposito, R.G.; Morici, I.J.; Szepsy, T.L.; Levinskas, G.J. (1965) CL 43,913 and CL 52,160: Seven-Day Repeated Feeding to Young Coturnix Quail: Report No. 65-44. (Unpublished study received Nov 19, 1975 under 241-246; submitted by American Cyanamid Co., Princeton, N.J.; CDL:224604-F)
005016098	Esser, H.O. (1971) Terminal residues of organophosphorus insecticides in animals. Pages 33-56, In Pesticide Terminal Residues: Invited Papers from the International Symposium on Pesticide Terminal Residues; Feb 17-19, 1971, Tel Aviv, Israel. Edited by A.S. Tahori. London: Butterworths.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000013990	Ewart, W. (1966) Summary of 1966 Citrus Thrips Control. (Unpublished study received Feb 3, 1977 under 201-274; prepared by Univ. of California--Riverside, submitted by Shell Chemical Co., Washington, D.C.; CDL:095827-K)
000013989	Ewart, W. (1967) Summary of 1967 Citrus Thrips Control. (Unpublished study received Feb 3, 1977 under 201-274; prepared by Univ. of California--Riverside, submitted by Shell Chemical Co., Washington, D.C.; CDL:095827-J)
000013988	Ewart, W. (1968) Summary of 1968 Citrus Thrips Control. (Unpublished study received Feb 3, 1977 under 201-274; prepared by Univ. of California--Riverside, submitted by Shell Chemical Co., Washington, D.C.; CDL:095827-I)
000017226	Ewart, W. (1974) Effectiveness of Nudrin 1.8EC Sprays Applied 5-?-74 for the Control of the Scirtothrips citri on Mature Navel Orange Trees in Tulare County, California. (Unpublished study received Jun 29, 1976 under 201-347; submitted by Shell Chemical Co., Washington, D.C.; CDL:232411-U)
005007048	Fairchild, E.J., ed. (1977) Agricultural Chemicals and Pesticides: A Subfile of the NIOSH Registry of Toxic Effects of Chemical Substances. Cincinnati, Ohio: National Institute for Occupational Safety and Health. (Pagination includes 46 pages numbered i-xvi; available from: NTIS, Springfield, VA: PB-274 748)
000001336	Fales, J.H.; Spangler, P.J.; Bodenstein, O.F.; Mills, G.D.; Durbin, C.G., Jr. (1968) Laboratory and field evaluations of Abate® against a backswimmer, Notonecta Undulata Say (Hemiptera: Notonectidae). Mosquito News 28(1):77-81. (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-AF)
005002611	Febles, C.R.; Romney, H. (1976) Follow-up on dengue--Puerto Rico. Morbidity and Mortality 25(9):65-66.
000001277	Fischer, J.E. (1975) Toxicity Data: Temephos 2% Aerosol: Report No. A75-58. (Unpublished study received Nov 19, 1975 under 241-246; submitted by American Cyanamid Co., Princeton, N.J.; CDL:224604-B)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005002610	Fishman, M.J.; Robinson, B.P. (1969) Water analysis. Analytical Chemistry 41(5):323A-360R.
005000834	Fitzpatrick, G. (1975) Impact of temephos and other mosquito larvicides on the salt marsh snail Melampus bidentatus Say (Basommatophora: Ellobiidae). Dissertation Abstracts International B 36(5):2045-2046.
005001002	Fitzpatrick, G.; Sutherland, D.J. (1975) Impact of Abate and Dursban larvicidal formulations on the salt marsh pulmonate snail Melampus bidentatus Say (Basommatophora: Ellobiidae). Pages 139, In Proceedings of the Sixty-second Annual Meeting of the New Jersey Mosquito Control Association; Mar 11-14, 1975, Atlantic City, New Jersey. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005000905	Fitzpatrick, G.; Sutherland, D.J. (1975) Temephos residues in the salt marsh snail Melampus bidentatus Say (Bassommatophora: Ellobiidae). Journal of the New York Entomological Society 83(4):267-268.
005000912	Fitzpatrick, G.; Sutherland, D.J. (1978) Effects of the organophosphorous insecticides temephos (Abate) and chlorpyrifos (Dursban) on populations of the salt-marsh snail Melampus bidentatus. Marine Biology 46(1):23-28.
005003972	Fitzpatrick, G.; Sutherland, D.J. (1978) Effects of the organophosphorus insecticides temephos (Abate) and chlorpyrifos (Dursban) on populations of the salt-marsh snail Melampus
005000936	Fontaine, R.E.; Rosen, P. (1973) Evaluation of Abate insecticide formulations as larvicides against Anopheles gambiae in northern Nigeria. Mosquito News 33(3):428-440.
005001340	Forgash, A.J. (1974) Mini-panel: initial studies of Abate in a salt-marsh ecosystem--introductory remarks. Pages 122-125, In Proceedings of Sixty-first Annual Meeting--New Jersey Mosquito Extermination Association; Mar 13-15, Atlantic City, New Jersey. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005000964	Forgash, A.J. (1976) A summary of studies of the impact of temephos and chlorpyrifos on the saltmarsh environment. Pages 94-98, In Proceedings of the Sixty-third Annual Meeting of the New Jersey Mosquito Extermination Association; Aug 10-12, Cherry Hill, New Jersey. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000001244	Foster, J.R. (1964) Larvicide Tests. (Unpublished study received May 27, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-Z)
005007632	Fox, I.; Rivera, G.A.; Umpierre, C.C. (1966) Toxicity of various insecticides to Xenopsylla cheopis. American Journal of Tropical Medicine and Hygiene 15(4):611-613.
005001110	Frank, A.M.; Sjogren, R.D. (1978) Effect of temephos and chlorpyrifos on Crustacea. Mosquito News 38(1):138-139.
005011171	Fuchs, T.W.; Harding, J.A.; Dupnik, T. (1972) Insecticidal Control of the Variegated Cutworm on Cantaloupes. College Station, Tex.: Texas Agricultural Experiment Station. (Texas Agricultural Experiment Station progress report no. 3135)
005010208	Fuchs, T.W.; Harding, J.A.; Dupnik, T. (1973) Control of Loopers on Cabbage in the Lower Rio Grande Valley of Texas. College Station, Tex.: Texas Agricultural Experiment Station. (Texas Agricultural Experiment Station progress report PR-3244)
005000636	Gahan, J.B.; Glancey, B.M.; Lofgren, C.S. (1965) New larvicides showing promise against Anopheles mosquitoes. Pages 156-159, In Proceedings of the Fifty-second Annual Meeting of the New Jersey Mosquito Extermination Association; 1965, Atlantic City, New Jersey. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005000641	Gaines, T.B.; Kimbrough, R.; Laws, E.R., Jr. (1967) Toxicology of Abate in laboratory animals. Archives of Environmental Health 14:283-288.
005000903	Garris, G.I.; Noblet, R. (1975) Notes on parasitism of black flies (Diptera: Simulidae) in streams treated with Abate. Journal of Medical Entomology 12(4):481-482.
005000981	Geevarghese, G.; Dhanda, V.; Rao, P.N.R.; Deobhankar, R.B. (1977) Field trials for the control of Aedes aegypti with Abate in Poona city and suburbs. Indian Journal of Medical Research 65(4):466-473.
005000654	Georgala, M.B. (1974) Thrips control: slow progress with new treatment. The Citrus and Sub-Tropical Fruit Journal 484:6-8,13.
005000790	Georghiou, G.P. (1972) Studies on resistance to carbamate and organophosphorus insecticides in Anopheles albimanus. American Journal of Tropical Medicine and Hygiene 21(5):797-806.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000747	Gillies, P.A.; Womeldorf, D.J.; White, K.E. (1968) Interpretation of resistance in California Aedes nigromaculis larvae. Pages 79-81, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 36. Visalia, Calif.: CMCA.
005000756	Gillies, P.A.; Womeldorf, D.J.; White, K.E. (1971) Insecticide susceptibility of Aedes melanimon and Aedes vexans larvae in California. Pages 108-111, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 39. Visalia, Calif.: CMCA.
005000839	Gillott, M.A.; Floyd, G.L.; Ward, D.V. (1975) The role of sediment as a modifying factor in pesticide-algae interactions. Environmental Entomology 4(4):621-624.
005000916	Glancey, B.M.; Moussa, M.A.; Scanlon, J.E.; Lofgren, C.S. (1968) Abate and Dursban against Aedes aegypti (L.) breeding in concrete water jars in Bangkok, Thailand. Mosquito News 28(2):205-206.
005004741	Goncharenko, N.G. (1975) O toksichnosti abata pri ingalyatsionnom postuplenii v organism. [On the toxicity of abate during inhalation entrance into the organism.] Vrachebnoe Delo.
005000900	Gould, D.J.; Mount, G.A.; Scanlon, J.E.; Ford, H.R.; Sullivan, M.F. (1970) Ecology and control of dengue vectors on an island in the Gulf of Thailand. Journal of Medical Entomology 7(4):499-508.
005000789	Gould, D.J.; Mount, G.A.; Scanlon, J.E.; Sullivan, M.F.; Winter, P.E. (1971) Dengue control of an island in the Gulf of Thailand. I. Results of an Aedes aegypti control program. American Journal of Tropical Medicine and Hygiene 20(5):705-714.
000001362	Granular Formulation of Three Systemic Carbamate Insecticides Residues in Cottonseed: Report No. C-124. (Unpublished study that includes updated report no. C-140 dated Jan 30, 1967 by Marion Moyer and E.J. OrLoski, and colorimetric procedure, received Feb 3, 1967 under 241-223; submitted by American Cyanamid Co.; CDL:090712-S)
005014221	Gras, G.; Rioux, J.A. (1968) Toxicite de quelques insecticides pour Aedes caspius (Pallus), Aedes detritus (Haliday), Culex de la Societe de Pharmacie de Montpellier 28(4):301-322.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000635	Gras, G.; Rioux, J.A. (1969) Laboratory evaluation of some organophosphorus compounds against the larvae of Aedes (o.) detrius (Haliday), Ades (o.) Caspius (Pallas) and Culex pipiens pipiens L. Mosquito News 29(2):202-209.
005002916	Gulyas, M.; Lang, F. (1972) Kiserletes vizsgalatok az Abate nevu szelektiv szunyoglarva-irtoszerrel. [Experimental investigations with Abate as a selective mosquito larvicide.] Parasitologia Hungarica 5:305-316.
005001030	Gunther, F.A. (1971) Residue Reviews. New York, N.Y.: Springer-Verlag.
005001276	Haji Noordin, H.J.B.; Zachariah, V.J. (1976) Aedes aegypti conrtol in Brunei. Brunei Museum Journal 3(4):188-192
000013982	Hall, D.H. (1973) Bidrin Thrip Control Trial on Oranges. (Unpublished study received Feb 3, 1977 under 201-274; prepared in cooperation with Progressive Farming, submitted by Shell Chemical Co., Washington, D.C.; CDL:095827-C)
005000754	Hansen, C.P. (1971) Midge control in California. Pages 51, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 39. Visalia, Calif.: CMCA.
000008892	Harding, J.A. (1971) Field Tests on Spring, 1971, Tomatoes Evaluating Spray Applications of Insecticides for Insect Control. (Unpublished study received Dec 20, 1974 under 352-342; prepared by Texas A & M Univ., Lower Rio Grande Valley Research and Extension Center, submitted by E.I. du Pont de Nemours & Co., Wilmington, Del.; CDL:022993-L)
000008885	Harding, J.A. (1971) Field Tests on 1971 Fall Tomatoes Evaluating Various Insecticides for Heliothis spp. and Tomato Pinworm Control. (Unpublished study received Dec 20, 1974 under 352-342; prepared by Texas A & M Univ., Lower Rio Grande Valley Research and Extension Center, submitted by E.I. du Pont de Nemours & Co., Wilmington, Del.; CDL:022993-B)
005000887	Harlan, D.P. (1974) Whitefringed beetles: control of larvae in nursery containers and phytotoxicity of Abate. Journal of Economic Entomology 67(6):804-805.
005000647	Harris, C.R.; Svec, H.J.; Sans, W.W. (1973) Toxicological studies on cutworms. IX. Laboratory and microplot field studies on effectiveness and persistence of some experimental insecticides used for control of the darksided cutworm. Journal of Economic Entomology 66(1):199-203.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000824	Harris, C.R.; Turnbull, S.A. (1975) Laboratory studies on the toxicity of insecticides to the bertha armyworm (Amestra Entomologist 107(8):865-872.
005000898	Hatfield, C.T.; Johansen, P.H. (1972) Effects of four insecticides on the ability of Atlantic salmon parr (Salmo salar) to learn and retain a simple conditioned response. Journal of the Fisheries Research Board of Canada 29(3):315-321.
005001251	Hayes, W.J., Jr. (1969) Pesticides and human toxicity. Annals of the New York Academy of Sciences 160(1):40-54.
005000770	Heath, R.G.; Spann, J.W.; Hill, E.F.; Kreitzer, J.F. (1972) Comparative Dietary Toxicities of Pesticides to Birds. Washington, D.C.: U.S. Fish and Wildlife Service. (Special scientific report: Wildlife, no. 152)
005003231	Herricks, E.E.; Buikema, A.S. (1977) Effects of pollution on freshwater invertebrates. Journal of the Water Pollution Control Federation 49(6):1493-1506.
000006596	Hiddleson, L.; Hofmaster, R.N.; Waterfield, R.L. (1968) Zolone EC--Potatoes: Final Report No. BB 65-51. (Unpublished study including final report no. BB 65-51A, received Dec 7, 1972 under 3F1337; prepared in cooperation with Virginia Truck and Ornamentals Research Station, Eastern Shore Branch, submitted by Rhodia, Inc., New Brunswick, N.J.; CDL:094849-Q)
000007601	Higgins, D.; Bass, M.H. (1970) Control of the Beet Armyworm, Spodoplera exigua (hub.) on Peanuts. (Unpublished report received Apr 26, 1971 under 1F1158; submitted by E.I. du Pont de Nemours & Co., Inc., Wilmington, Del.; CDL:090951-A)
000015976	Higgins, D.; Bass, M.H. (1970) Control of the Beet Armyworm, Spodoplera exqua (hub.) on Peanuts. (Unpublished report received Jun 29, 1976 under 201-347; prepared by Auburn Univ., submitted by Shell Chemical Co., Washington, D.C.; CDL:232408-Q)
005000911	Hill, E.F. (1972) Avoidance of lethal dietary concentrations of insecticide by house sparrows. Journal of Wildlife Management 36(2):635-639.
000001311	Hirst, J.M.; Cole, M.M.; Gilbert, I.H.; Adams, C.T. (1970) Further sleeve tests of new powders for control of body lice. Journal of Economic Entomology 63(3):861-862. (Also In unpublished submission received Sep 1, 1972 under 241-230; submitted by American Cyanamid Co., Princeton, N.J.; CDL:002063-J)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000705	Hitchcock, S.W.; Anderson, J.F. (1968) Field-plot tests with insecticides for control of Chironomous atrella. Journal of Economic Entomology 61(1):16-19.
000001370	Hodges, J.L. (1965) Insecticide Treatments and Schedules. (Unpublished study received Jul 17, 1967 under 7G0566; prepared by Univ. of Arizona Cotton Research Center?, submitted by American Cyanamid Co., Princeton, N.J.; CDL:090713-K)
000006989	Hofmaster, R.N. (1965) Report No. BB-65-51. (Unpublished study received on unknown date under 7G0575; prepared by Virginia Truck and Ornamentals Research Station, submitted by Chipman Chemical Co., Inc., Burlingame, Calif.; CDL:090736-Y)
000011928	Hofmaster, R.N.; Waterfield, R.L. (1965) Foliage Treatments to Control Colorado Potato Beetles. (Unpublished study including letter dated Oct 15, 1965 from R.N. Hofmaster to E.R. Cowett, received Nov 10, 1972 under 3F1323; prepared by Virginia Truck Experiment Station, Eastern Shore Branch, Tidewater Virginia Vegetable Research Station, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:093552-AN)
005001013	Holland, H.T.; Coppage, D.L.; Butler, P.A. (1967) Use of fish brain acetylcholinesterase to monitor pollution by organophosphorus pesticides. Bulletin of Environmental Contamination and Toxicology 2(3):156-162.
005000854	Howe, L.H., III; Petty, C.F. (1969) Thin-layer chromatographic quantitation of Abate (O,O,O',O'-tetramethyl-O,O'-thiodi-p-phenylene phosphorothioate) residues in water. Journal of Agricultural and Food Chemistry 17(2):401-402.
000001355	Hutchison, E.B.; Levinskas, G.J. (1966) Abate® 4E Mosquito Larvicide: Five-Day Repeated Applications to Intact Rabbit Skin: Report no. 66-167. Undated method entitled: Modified Michel Method for Assay of Cholinesterase Activity. (Unpublished study received Feb 3, 1967 under 7G0566; submitted by American Cyanamid Co., Agricultural Div., Princeton, N.J.; CDL:090712-C)
005012519	Inoue, Y.; Mihara, J. (1974) Jikken niyoru usukari yochu sachuzai no koryoku hyoka. [Evaluation of the effect of insecticides on Cricotopus bicinctus.] Eisei Dobutsu. [Sanitary Zoology.] 24(4):283.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005006246	Inoue, Y.; Mihara, M. (1975) Waga kuni ni okeru fukai konchu to shite yusurika ni kansuru kenkyu: 1. Moderu shiken ni yoru sushu yuki-rin-satchuzai no sesujiyusurika gorei jochu ni taisuru koryoku hyoka. [Studies on the harmful Japanese Chironomid midges: I. Larvicidal effects of some organophosphorus insecticides against the last instar larvae of Chironomus yoshimatsui Martin and Sublette.] Eisei Dobutsu.
005004387	Isa, A.L.; Awadallah, W.H.; El Tantawy, A.M. (1972) The chemical control of Sesamia cretica Led. on corn. Agricultural Research Review 50(1):51-59.
005007006	Ivanova, G.B.; Roslavtseva, S.A.; Evtyushina, T.M.; Polyakova, V.K. (1972) Kharakteristika shtamma komnatnoi mukhi (Musca a coumaphos-resistant housefly strain (Musca domestica L.).] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 41(2):175-180.
005000707	Jakob, W.L. (1965) Laboratory test of larvicides for mosquito control in potable waters. Mosquito News 25(3):316-319.
005000812	Jamnback, H. (1969) Field tests with larvicides other than DDT for control of black fly (Diptera simuliidae) in New York. Bulletin of the World Health Organization 40(4):635-638.
005000737	Jamnback, H.A.; Means, R.G. (1968) Formulation as a factor influencing the effectiveness of Abate in control of black flies (Diptera: simuliidae). Pages 89-94, In Proceedings of the Fifty-fifth Annual Meeting of the New Jersey Mosquito Extermination Association. Vol. 55. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005001016	Jamnbeck, H.; Frempong-Boadu, J. (1966) Testing blackfly larvicides in the laboratory and in streams. Bulletin of the World Health Organization 34(3):405-421.
000007824	Jefferson, R.N.; Morishita, F.S. (1967?) Summary of Results, 1966 & 1967: Project 1338-A. (Unpublished study received Jul 23, 1968 under 352-342; prepared by Univ. of California--Riverside, submitted by E.I. du Pont de Nemours & Co., Wilmington, Del.; CDL:003000-E)
005000803	Jeppson, L.R. (1974) Pest management in citrus orchards. Bulletin of the Entomological Society of America 20(3):221-222.
005000895	Jeppson, L.R.; McMurtry, J.A.; Mead, D.W.; Jesser, M.J.; Johnson, H.G. (1975) Toxicity of citrus pesticides to some predaceous phytoseiid mites. Journal of Economic Entomology 68(5):707-710.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005008989	Johansen, C.; Eves, J. (1967) Toxicity of Insecticides to the Alkali Bee and the Alfalfa Leafcutting Bee. Pullman, Wash.: Washington State University Agricultural Research Station. (Washington Agricultural Research Station circular no. 475)
005000764	Johnson, C.R. (1976) Further investigations into the culture and winter maintenance of Gambusia affinis affinis (Pisces: Poeciliidae). Pages 52, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 44. Visalia, Calif.: CMCA.
005000766	Johnson, C.R. (1977) The effects of field applied rates of five organophosphorus insecticides on thermal tolerance, orientation, and survival in Gambusia affinis affinis (Pisces: Poeciliidae). Pages 56-58, In Proceedings and Papers of the Annual Conference of the California Mosquito and Vector Control Association. Vol. 45. Visalia, Calif.: CMCA.
005018646	Johnson, C.R. (1978) The effect of five organophosphorus insecticides on survival and temperature tolerance in the copepod, Macrocyclops albidus (Copepoda: Cyclopidae). Zoological Journal of the Linnean Society 64(1):59-62.
005018647	Johnson, C.R. (1978) The effects of sublethal concentrations of five organophosphorus insecticides on temperature tolerance, reflexes, and orientation in Gambusia affinis affinis (Pisces: Poeciliidae). Zoological Journal of the Linnean Society 64(1):63-70.
005008878	Johnson, C.R. (1979) The effects of sublethal concentrations of five organophosphorus insecticides on temperature tolerance, reflexes, and orientation in Gambusia affinis affinis (Pisces: Poeciliidae). Sport Fishery Abstracts 24(2):112.
005014957	Johnson, C.R.; Prine, J.E. (1976) The effects of sublethal concentrations of organophosphorus insecticides and an insect growth regulator on temperature tolerance in hydrated and dehydrated juvenile western toads, Bufo boreas. Comparative Biochemistry and Physiology 53A(2):147-149.
005000830	Johnson, C.R.; Prine, J.E. (1976) The effects of sublethal concentrations of organophosphorus insecticides and an insect growth regulator on temperature tolerance in hydrated and dehydrated juvenile western toads, Bufo boreas. Comparative Biochemistry and Physiology 53(A):147-149.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005001008	Johnson, W.P.; Alford, B.T.; Simkins, K.L.; Wang, G.T. (1975) Over- and multidose safety tests with temephos 2% ectoparasiticide on dogs and cats. Pages 244-245, In 20th World Veterinary Congress. Summaries, vol. 1. Thessaloniki, Greece.
005004417	Jones, K.H.; Sanderson, D.M.; Noakes, D.N. (1968) Acute toxicity data for pesticides (1968). World Review of Pest Control 7(3):135-143.
005000949	Joseph, S.R. (1977) Temporary mosquito control in Maryland. Mosquito News 37(1):27-30.
005001025	Kaemmerer, K.; Buntenkoetter, S. (1973) The problem of residues in meat of edible domestic animals after application or intake of organophosphate esters. Residue Reviews 46:1-240.
005016161	Kaemmerer, K.; Buntenkoetter, S. (1973) The problem of residues in meat of edible domestic animals after application or intake of organophosphate esters. Pages 1-240, In The Problem of Residues in Meat. Edited by F.A. Gunther. New York: Springer. (Residue reviews, vol. 46)
005000958	Kamel, O.M.; Nassif, M.; Nagy, M. (1976) Laboratory and field studies on the use of temephos against body lice in Egypt. Pans 22(3):350-353.
005000745	Kanakkanatt, S.V.; Cardarelli, N.F. (1975) A new concept in blackfly control. Pages 222-229, In Proceedings of the 1975 International Controlled Release Pesticide Symposium; Sep 8-10, Dayton, Ohio. Dayton, Ohio: Wright State University, College of Science and Engineering.
005002510	Katz, M.; Pederson, G.L.; Yoshinaka, M.; Sjolseth, D. (1969) Effects of pollution on fish life. Journal of the Water Pollution Control Federation 41(6):994-1016.
005000938	Kay, B.H.; Ferguson, K.J.; Morgan, R.N.C. (1973) Control of salt-marsh mosquitoes with Abate insecticide at Coombabah Lakes, Queensland, Australia. Mosquito News 33(4):529-535.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005004489	Keiser, I.; Kobayashi, R.M.; Schneider, E.L.; Tomikawa, I. (1973) Laboratory assessment of 73 insecticides against the oriental fruit fly, melon fly, and Mediterranean fruit fly. Journal of Economic Entomology 66(4):837-839.
005000850	Kerdpibule, V.; Hirakoso, S. (1971) Susceptibilities to several insecticides in adult house flies collected from various districts in Thailand. Japanese Journal of Experimental Medicine 41(6):541-545.
005000985	Khalil, H.M.; Rifaat, M.A.; Gad, A.M.; Sadek, S. (1974) Filarial infectivity rate of Culex pipiens molestus subjected to sublethal concentrations of insecticides Abate and Sevin and distribution of infective filaria larvae in mosquito body regions. Journal of the Egyptian Public Health Association XLIX(4/5):221-224.
005000768	Khan, M.A.Q.; Khan, H.M.; Sutherland, D.J. (1974) Ecological and health effects of the photolysis of insecticides. Pages 333-355, In Survival in Toxic Environments. Edited by M.A.Q. Khan and John P. Bederka, Jr. New York: Academic Press.
005003885	Khristova, T.; Tsanev, I.; Dorovski, D.; Todorova, M.; Bunkov, M.; Smilova, D. (1974) Sredstva az borba s larvi na komari. Mikrobiologiya i Infektsiozni Bolesti. [Epidemiology, Microbiology and Infectious Diseases.] 11(4):381-388.
005000921	Kilpatrick, J.W.; Eliason, D.A.; Babbitt, M.F. (1970) Studies of the potential effectiveness of ultra low volume aerial applications of insecticides against Aedes aegypti (L.) larvae. Mosquito News 30(2):250-258.
005000902	Kissam, J.B.; Noblet, R.; Garris, G.I. (1975) Large-scale aerial treatment of an endemic area with Abate granular larvicide to control black flies (Diptera: Simuliidae) and suppress Entomology 12(3):359-362.
005000894	Kissam, J.B.; Noblet, R.; Moore, H.S., IV (1973) Simulium: field evaluation of Abate larvacide for control in an area endemic for Leucocytozoon smithi of turkeys. Journal of Economic Entomology 66(2):426-428.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000001287	Knapp, F.W. (1976) Temephos Field Studies for Cattle Lice Control. (Unpublished study received Jun 2, 1976 under 241-EX-77; prepared by Univ. of Kentucky, submitted by American Cyanamid Co., Princeton, N.J.; CDL:226521-C)
005002851	Kovacs, A.; Zambelli, N.; Baldini, G.P.; Komblas, K.N. (1968) Dursban: su evaluacion como larvicida de mosquitos. Biokemia 15:10-13.
005007008	Kovban, V.Z. (1975) Poisk effektivoi bor'by s moshchkami. [The search for effective control of black flies.] Pages 225-227, In Problemy Parazitologii, Materialy Nauchnoi Konferentsii Parazitologov Ukrainskoi SSR, 8th; Donetsk. Scientific Conference of Parasitologists of the Ukrainian SSR. Part 1.] Edited by T.D. Peck. Kiev, USSR: Naukova Dumka.
005012182	Krijgsman, W.; Kamp, C.G. van de (1976) Analysis of organophosphorus pesticides by capillary gas chromatography with flame photometric detection. Journal of Chromatography 117(1):201-205.
005000864	Krijgsman, W.; Van de Kamp, C.G. (1976) Analysis of organophosphorus pesticides by capillary gas chromatography with flame photometric detection. Journal of Chromatography 117(1):201-205.
005000617	Kurinnyi, A.I. (1975) [Comparative study of the cytogenetic effect of certain organophosphorus pesticides.] Genetika.
005000709	Kutz, F.W.; Burbutis, P.P. (1966) Evaluation of some new insecticides as mosquito larvicides. Mosquito News 26(1):73-77.
005000951	Kutz, F.W.; Strassman, S.C. (1977) Human uninary metabolites of organophosphate insecticides following mosquito adulticiding. Mosquito News 37(2):211-218.
005000896	Lacey, L.A.; Mulla, M.S. (1977) A new bioassay unit for evaluating larvicides against blackflies. Journal of Economic Entomology 70(4):453-456.
005004480	Lang, J.H.; Bry, R.E.; Boatright, R.E. (1976) Organophosphorus insecticides as potential wool protectants. Soap, Cosmetics, Chemical Specialties 52(9):40,42,44.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005002918	Lavagnino, A.; Smiraglia, C.B. (1970) Trattamento di focolai di Aedes mariae con larvicidi (Abate, Dursban) in supporto di polietilene. [Treatment of Aedes mariae with larvicides (Abate, Dursban) in support of polyethylene.] Igiene Moderna.
005007515	Lavaur, E. de; Arnold, A. (1977) Pesticides et faune sauvage: resultats des analyses toxicologiques effectuees sur le gibier de 1974 a 1976. [Pesticides and wildlife: results of toxicological analyses performed on game from 1974 to 1976.] Phytiatrie-Phytopharmacie 26(3):159-167.
005000700	Laws, E.R., Jr.; Morales, F.R.; Hayes, W.J., Jr.; Joseph, C.R. (1967) Toxicology of Abate in volunteers. Archives of Environmental Health 14(2):289-291.
005000810	Laws, E.R., Jr.; Sedlak, V.A.; Miles, J.W.; Joseph, C.R.; Lacomba, J.R.; Rivera, A.D. (1968) Field study of the safety of Abate for treating potable water and observations on the effectiveness of a control programme involving both Abate and malathion. Bulletin of the World Health Organization 38(3):439-445.
000001374	Laws, E.R., Jr.; Sedlak, V.A.; Miles, J.W.; Romney Joseph, C.; Lacomba, J.R.; Diaz Rivera, A. (1968) Field study of the safety of Abate for treating potable water and observations on the effectiveness of a control programme involving both Abate and Malathion. Bulletin de l'Organisation mondiale de la Sante; Bulletin of the World Health Organization 38(??):439-445. (Also submitted by American Cyanamid Co., Princeton, N.J.; CDL:090713-O)
005009712	Lee, C.M. (1963) [Determination of the structure of organophosphorus compounds by refractometry.] Huaxue Tongbao.
005001116	Lee, C.W. (1973) Aerial spraying trials in West Africa for blackfly control. PANS 19(2):190-192.
005000988	Lee, C.W.; Parker, J.D.; Philippon, B.; Baldry, D.A.T. (1975) A prototype rapid release system for the aerial application of larvicide to control Simulium damnosum Theo. PANS 21(1):92-102.
005000987	Lee, D.; Moore, C.G. (1973) Mosquito studies during an interepidemic outbreak of dengue in Puerto Rico. Mosquito News 33(4):506-509.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000007085	Lee, L.; Bass, M.H. (1967) Peanut Insects: Annual Report for Period January 1 to December 31, 1967: State Research 13-008. (Unpublished study received Apr 26, 1971 under 1F1158; submitted by E.I. du Pont de Nemours & Co., Inc., Wilmington, Del.; CDL: 090951-K)
005014595	Leesch, J.G. (1970) The Chemical and Biochemical Basis for the Selective Toxicity of Abate to Mosquito Larvae [Doctoral Dissertation]. Riverside, Calif.: University of California. (Available from: University Microfilms, Ann Arbor, MI; order no. 71-20,246)
005000779	Leesch, J.G. (1971-72) The chemical and biochemical basis for the selective toxicity of Abate to mosquito larvae. Dissertation Abstracts International B 32(2):995.
005000960	Leesch, J.G.; Fukuto, T.R. (1972) The metabolism of Abate in mosquito larvae and houseflies. Pesticide Biochemistry and Physiology 2(2):223-235.
000007897	Leigh, T.F. (1966) LD_{50} Values and ?Values for 34 Insecticides When Applied Topically to Adult Male and Female Lygus hesperus Knight: Shafter, California: 1960-1966: Table 1. (Unpublished study including letter dated Sep 19, 1967 from T.F. Leigh to Donald L. Burgoyne, received Apr 16, 1971 under 1G1144; prepared by Univ. of California--Davis, Agricultural Experiment Station, Dept. of Entomology, submitted by E.I. du Pont de Nemours & Co., Inc., Wilmington, Del.; CDL:090931-B)
000007898	Leigh, T.F.; Jackson, C.E. (1969) Evaluation of Insecticides for Control of Lygus hesperus Knight on Cotton. (Unpublished study received Apr 16, 1971 under 1G1144; prepared by Univ. of California--Davis, submitted by E.I. du Pont de Nemours & Co., Inc., Wilmington, Del.; CDL:090931-C)
005000891	Leigh, T.F.; Jackson, C.E.; Wynholds, P.F.; Cota, J.A. (1977) Toxicity of selected insecticides applied topically to Lygus hesperus. Journal of Economic Entomology 70(1):42-44.
005000725	Leveque, C.; Odei, M.; Thomas, M.P. (1977) The Onchocerciasis Control Programme and the monitoring of its effect on the riverine biology of the Volta River Basin. Pages 133-143, In Ecological Effects of Pesticides. Edited by F.H. Perring and K. Mellandy. New York: Academic Press. (Linnean Society symposium series no. 5)
000001339	Levinskas, G.J. (1965?) Seven-Day Repeated Feeding to Young Coturnix Quail: CL 43,913 and CL 52,160. (Unpublished study received Jun 18, 1968 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:105218-F)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000973	Levinskas, G.J.; Shaffer, C.B. (1970) Toxicity of Abate, a mosquito larvicide, and its sulfoxide. Toxicology and Applied Pharmacology 17(1):301-302.
005000645	Levy, R.; Miller, T.W., Jr. (1977) Susceptibility of the mosquito nematode Romanomermis culicivorax (Mermithidae) to pesticides and growth regulators. Environmental Entomology 6(3):447-448.
005000644	Levy, R.; Miller, T.W., Jr. (1978) Tolerance of the planarian Dugesia dorotocephala to high concentrations of pesticides and growth regulators. Entomophaga 23(1):31-34.
005001033	Lewis, A.E.; Brody, T.M. (1969) Clinical and laboratory aspects of the diagnosis and management of organophosphate toxicity. University of Michigan Medical Center Journal XXXV(1):18-22.
005000961	Lewis, L.F.; Christenson, D.M. (1968) Abate, Dursban studied in northwest project. Pest Control 36(7):38-40.
005000943	Lewis, L.F.; Christenson, D.M. (1975) Residual activity of temephos, chlorpyrifos, DDT, fenthion, and malathion against Aedes sierrensis (Ludlow) in fabricated treeholes. Mosquito News 35(3):381-384.
005000762	Lewis, L.F.; Christenson, D.M. (1975) Residual tests with Abate, chlorpyrifos, DDT, fenthion, and malathion against Aedes sierrensis (Ludlow) in fabricated treeholes. Page84, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 43. Visalia, Calif.: CMCA.
005000708	Lewis, L.F.; Christenson, D.M.; Eddy, G.W. (1965) Control of mosquito larvae in Williamette Valley, Oregon log ponds. Mosquito News 25(4):400-402.
005000945	Liem, K.K.; LaSalle, R.N. (1976) Effects of Abate 2G and Abate coexisting with mosquito larvae in woodland depressions. Mosquito News 36(2):202-203.
000006100	Lillie, D.T.; Tuttle, D.M.; Arvizo, G.L. (1971) Zolone EC--Oranges: Final Report No. FD 71-15. (Unpublished study received Aug 19, 1971 under 2F1193; prepared in cooperation with Univ. of Arizona, submitted by Rhodia, Inc., New Brunswick, N.J.; CDL: 093518-X)
000006860	Lincoln, C. (1966) Report No. PAS 66-79. (Unpublished study received Jul 5, 1968 under 8F0668; submitted by Chipman Chemical Co., Inc., Burlingame, Calif.; CDL:092027-BY)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000007905	Lincoln, C. (1966) Small Plot Tests, Southeast Branch Station, 1966. (Unpublished study received Apr 16, 1971 under 1G1144; prepared by Univ. of Arkansas, submitted by E.I. du Pont de Nemours & Co., Inc., Wilmington, Del.; CDL:090931-S)
000007972	Livingston, R.J. (1978) Comparative Review of the Environmental Impact of Various Insecticides Used for Mosquito Control in the U.S. (Unpublished study received Mar 15, 1978 under 148-1259; submitted by Thompson-Hayward Chemical Co., Kansas City, Kans.; CDL:096962-B)
000001286	Lloyd, J.E. (1976) Temephos for Lice Control on Range Cattle. (Unpublished study received Jun 2, 1976 under 241-EX-77; prepared by Univ. of Wyoming, submitted by American Cyanamid Co., Princeton, N.J.; CDL:226521-B)
005000628	Lofgren, C.S.; Pennington, N.; Young, W. (1966) Evaluation of insecticides against two species of Culex mosquitoes on Okinawa. Mosquito News 26(1):52-59.
005000715	Lofgren, C.S.; Scanlon, J.E.; Israngura, V. (1967) Evaluation of insecticides against Aedes aegypti (L.) and Culex pipiens Thailand. Mosquito News 27(1):16-21.
000027094	Lowe, J.I. (1964) Gulf Shellfish: Effects of Pesticides. (Unpublished study including letter dated Jul 28, 1964 from J.I. Lowe to H.E. Fairchild, received Jul 28, 1964 under unknown admin. no.; prepared by U.S. Fish and Wildlife Service, Bureau of Commercial Fisheries, submitted by Ciba-Geigy, Greensboro, N.C.; CDL:131211-A)
005000818	Lyons, G.R.L. (1973) The control of guineaworm with Abate: a trial in a village of North-West Ghana. Bulletin of the World Health Organization 49(2):215-216.
005000731	MacDonald, W.W.; Pant, C.P. (1973) Control of Aedes aegypti, the principal vector of dengue haemorrhagic fever. Pages 9-10, In Ninth International Congress on Tropical Medicine and Malaria; Oct 14-21, Athens, Greece. Vol I. Abstracts of Invited Papers. Edited by George J. Papaevangelou. Athens, Greece: Congress Publications Committee.
005000773	Macko, J.A., Jr.; Harvey, J.G. (1977) Inhalation Hazard Evaluation of Pyrax/Insecticide Dust During a Simulated Delousing Exposure, June 1977-December 1977. Abderdeen Proving Ground, Md.: U.S. Army Environmental HygieneAgency. (USAEHA study no. 51-1302-78; available from: NTIS, Springfield, VA; AD-A049 245)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000917	Madhukar, B.V.R.; Pillai, M.K.K. (1968) Insecticide susceptibility studies in Indian strains of Aedes aegypti L. Mosquito News 28(2):222-225.
005000815	Madhukar, B.V.R.; Pillai, M.K.K. (1970) Development of organophosphorus resistance in Indian strains of Aedes aegypti (L.). Bulletin of the World Health Organization 43(5):735-742.
005000983	Mahdi, A.; Taha, A.M.; Ezz El-Arab, M.A. (1967) Preliminary small scale field trial for evaluation of aerial spraying of Malathion L. V. and Abate against A. pharoensis in the Nile delta 1966. Journal of the Egyptian Public Health Association 42(1):8-13.
005000723	Makarova, S.V.; Khmel'nitskaya, V.N.; Eliseeva, M.A. (1974) Photometric determination of oranophosphorus pesticides after separation by thin-layer chromatography. A translation of: Zavodskaya Laboratoriya. [Industrial Laboratory.] 40(7):792-795.
005018094	Malinin, O.A. (1979) Opredelenie fosfororganicheskikh pestitsidov. (Moscow). [Veterinary Science.] (1):72-74.
005000742	Mallack, J.; Langford, G.S.; Sanderson, A.E., Jr. (1971) Mosquito control in a landfill with Abate and observations on fish, shrimp and blue crabs in the adjacent estuary. Pages 110-117, In Proceedings of the Fifty-eighth Annual Meeting of the New Jersey Mosquito Extermination Association. Vol. 58. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005000752	Mallars, J.; Brumbaugh, L.R. (1969) Field observations of organophosphate drip systems applied to waste water stabilization lagoons in San Joaquin County. Pages 138-140, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 37. Visalia, Calif.: CMCA.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005002137	Manouchehri, A.V.; Khademi, M.; Eshghi, N. (1975) Comparative tests of five organo-phosphorus insecticides against lab-bred and wild caught larvae of Anopheles stephensi mysorensis in southern Iran. Iranian Journal of Public Health 3(4):179-182.
005000778	Marei, A.H.S. (1970) Granular Formulations of Abate and Its Release in Water; Doctoral Dissertation. New Brunswick, N.J..: Rutgers University. (Available from: University Microfilms, Ann Arbor, Michigan; order no. 71-3074)
005000823	Martel, P.; Harris, C.R.; Svec, H.J. (1975) Toxicological studies on the carrot weevil, Listronotus oregonensis (Coleoptera: Curculionidae). The Canadian Entomologist 107(5):471-475.
005014445	Mashourian, R.S. (1977) Abati yev defosi hamematakar dununakudyuna tavari artaken mshagman depkum. [Comparative toxicological effect of Abate and Diphos on cattle when applied externally.] Izvestiya Sel'skokhozyaistvennykh Nauk. [Bulletin of Agricultural Sciences.] 20(11):102-106.
005000657	Mason, J.; Hobbs, J.H. (1973) A study of the epidemiology of malaria in a high-incidence coastal area of El Salvador, C. A. Revista del Instituto de Investigaciones Medicas 2(1):51-54.
005021700	Matsumura, F.; Madhukar, B.V. (1980) Exposure to insecticides. Pharmacology and Therapeutics 9:27-49.
005015192	Matthiessen, P.; Johnson, J.S. (1978) Accumulation of the organophosphate blackfly larvicide Abate (temephos) in Sarotherodon mossambicus, with reference to the larvicidal control of Simulium damnosum. Journal of Fish Biology 13(5):575-586.
005000785	Mayberry, R.M.; Savage, J. (1978) Mutagenic activity of several pesticides using the Salmonella test and Saccharomyces Annual Meeting of the American Society for Microbiology; May 14-19, Las Vegas, Nevada.
000001330	Mayer, F.L., Jr.; Walsh, D.F. (1970) Annual Progress Report: 1970: Pesticide-Induced Changes in Pond Ecology: [Abate]. (Unpublished study received Nov 26, 1971 under 241-132; prepared by U.S. Fish and Wildlife Service, Fish-Pesticide Research Laboratory, submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-Y)
000001314	McCain, J.C. (1971) Final Report: Acute Fish Toxicity Study--Rainbow Trout: Project No. 362-123. (Unpublished study received Nov 26, 1971 under 241-132; prepared by TRW/Hazleton Laboratories submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-B)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000698	McCarty, R.T.; Haufler, M.; McBeth, C.A., Jr. (1968) Toxicologic effects of a phosphorothioic acid insecticide in sheep. Journal of the American Veterinary Medical Association 152(3):279-281.
005000862	McCully, K.A. (1978) Report on phosphated pesticides. Journal of the Association of Official Analytical Chemists 61(2):364-368.
005005244	McCully, K.A. (1979) Report on organophosphorus pesticides. Journal of the Association of Official Analytical Chemists 62(2):383-387.
005000931	McDonald, J.L. (1972) Evaluations of Abate for mosquito control in polluted water. Mosquito News 32(4):627-631.
000001364	McNerney, J.M.; Levinskas, G.J. (1967) Abate® Mosquito Larvicide: Demyelination Studies in White Leghorn Hens: Report no. 67-25. (Unpublished study received Jul 17, 1967 under 7G0566; submitted by American Cyanamid Co., Princeton, N.J.; CDL:090713-A)
000001366	McNerney, J.M.; Ribelin, W.E.; Levinskas, G.J. (1968) Abate Report no. 68-9. (Unpublished study received on unknown date under 7G0566; submitted by American Cyanamid Co., Princeton, N.J.; CDL:090713-D)
005011352	Mel'nikov, N.N. (1971) O perspektivnom assortimente insektitsidov. Khozyaistve. [Chemistry in Agriculture.] 9(3):194-198.
005002678	Mestres, R.; Chevallier, C. (1971) Penetration des insecticides dans les sols submerges. Deuxieme partie: Etude au laboratoire de la penetration de l'abate. [Penetration of insecticides in submerged soil. Part 2: Laboratory study of the penetration of abate.] Travaux de la Societe de Pharmacie de Montpellier 31(2):125-132.
005000721	Mestres, R.; Chevallier, C.; Rioux, J.A.; Cousserans, J.; Sinegre, G. (1971) Penetration of two insecticides into halomorphous soils temporarily submerged. A translation of: Travaux de la Societe de Pharmacie de Montpellier 31(2):159-166.
005000953	Micks, D.W.; Rougeau, D. (1977) Organophosphorus tolerance in Culex quinquefasciatus in Texas. Mosquito News 37(2):233-239.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000726	Miles, J.W.; Dale, W.E. (1975) Field extraction and laboratory analysis of residues of Abate in streams treated for simulium control. Pages 100-104, In Environmental Quality and Safety Supplement. Vol. III: Pesticides. International Union of Pure and Applied Chemistry, Third International Congress; Jul 3-9,1974, Helsinki, Finland. Edited by Frederick Coulston and Friedhelm Korte. Stuttgart, West Germany: George Thieme Publishers.
005000859	Miles, J.W.; Dale, W.E. (1978) Gas chromatographic determination of organophosphorus pesticides by in-block methylation. Journal of Agricultural and Food Chemistry 26(2):480-482.
000001381	Miles, J.W.; Woehst, J.E. (1967) Formulations for controlled release of Abate in water. (Unpublished paper presented at the 153rd National Meeting American Chemical Society; Apr 9-14, 1967; Miami Beach, Fla.; received Apr 1967 under 4822-80; prepared by U.S. Public Health Service, National Communicable Disease Center, Technical Development Laboratories, submitted by S.C. Johnson & Sons, Inc., Racine, Wis.; CDL:028367-A)
005004574	Miles, J.W.; Woehst, J.E. (1969) Formulations for controlled release of Abate in water. Pages 183-191, In Pesticidal Formulations Research. Washington, D.C.: American Chemical Society. (Advances in chemistry series)
005000783	Milionis, J.P.; Spicer, L.D., inventors; American Cyanamid Co., assignee (1976) Compositions resineuses renfermant de O'O,O'-thio di-p-phenylenephosphorothioate O,O,O',O'-tetramethyle. [Resin compositions containing O,O,O',O'-tetramethyl-O,O-thio di-p-phenylenephosphorothioate.] Belgian brevet d'invention 832,841. Mar 1. 11 p.
005000863	Miller, C.W.; Funes, F.M. (1971) Alkali-flame gas chromatography of Abate. Journal of Chromatography 59(1):161-162.
005000639	Miller, T.A. (1971) Field Evaluation of Encapsulated Formulations of Malathion (3MCAP-M) and Abate (3MCAP-A) as Mosquito Larvicides, July-October, 1971. Edgewood Arsenal, Md.: U.S. Army Environmental Hygiene Agency. (Entomological special study no. 44-003-72; available from: NTIS, Springfield, VA: AD-741 347)
005009412	Miller, T.A. (1972) Field Evaluation of Encapsulated Formulations of Malathion (3MCAP-M) and Abate (3MCAP-A) as Mosquito Larvicides, July-October 1971. Edgewood Arsenal, Md.: U.S. Army Environmental Hygiene Agency. (U.S. Army Environmental Hygiene Agency entomological special study no. 44-003-72; available from: NTIS, Springfield, VA; AD-741 347)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000934	Miller, T.A.; Lawson, M.A.; Nelson, L.L.; Young, W.W. (1973) Polymer formulations of mosquito larvicides. VII. Laboratory and field evaluation of encapsulated formulations of malathion and Abate. Mosquito News 33(3):413-417.
005001113	Mills, C. (1975) Arkansas mosquito control. Journal of the Arkansas Medical Society 71(9):292-293.
005002496	Milne, D.L.; De Villiers, E.A. (1975) Danger periods for applying certain pesticides to citrus. The Citrus and Subtropical Fruit Journal 496:17-18.
005001985	Milne, D.L.; de Villiers, E.A. (1976) Soil systematics against citrus thrips: results of the 1974-75 field trials. Australian Citrus News 50:10-11.
005015715	Ministro della Sanita (1979) Quantita massime di residui di antiparassitari consentite negli alimenti. [Maximum amount of permissable antiparasitic residues permitted in foods.] Industria Conserve. [Canning Industry.] 54(1):77-84.
005000972	Mitchell, C.J.; Chen, P.S. (1974) Susceptibility and resistance of four Culex species in Taiwan to certain insecticides. Journal of the Formosan Medical Association 73(4):185-195.
005001827	Mitchell, C.J.; Chen, P.S.; Chapman, H.C. (1974) Exploratory trials utilizing a mermithid nematode as a control agent for Culex mosquitoes in Taiwan. Journal of the Formosan Medical Association 73(5):241-254.
005000655	Mitchell, C.J.; Chen, P.S.; Okuno, T. (1974) Preliminary studies on the control of Japanese encephalitis vectors in Taiwan by chemical means. Journal of the Formosan Medical Association 73(6):295-312.
005002855	Mitrofanov, A.M.; Krivtsova, E.N.; Timofeeva, L.S.; Shogam, S.M. (1973) Izuchenie granulirovannykh fosfororganicheskikh larvitsidov v bor'be s komarami p. Aedes. [Study of granulated organophosphorus larvicides for control of Aedes mosquitos.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 42(3):315-320.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005002676	Mitrofanov, A.M.; Petruchuk, O.E.; Tulupova, A.M. (1977) Razrabotka metodov bor'by s gnusom v usloviyakh severnoi taigi na stroitel'stve zapadnogo uchastka Baikalo-Amurskoi magistrali v 1975 g. (na primere pos. magistral'nogo. [Development of methods for conrolling Aedes mosquitoes in the northern Taiga during construction of the western section of the Baikal-Amur Railway in 1975 (as illustrated by the Magistral'nyi Settlement).] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 46(6):721-727.
005012248	Miyamoto, J. (1971) Yukirin sacchuzai to kankyo hozen. Kagaku. [Scientific Pest Control.] 36(III):135-158.
000015275	Moherek, E.A.; Smith, J.W.; Berberet, R.; et al. (1976) [The Efficacy of Orthene 75 S as an Insect Control for Peanuts]. (Unpublished study received Jun 29, 1976 under 239-EX-81; submitted by Chevron Chemical Co., Richmond, Calif.; CDL:095985-D)
000001318	Moody, D.S.; Laster, M.L. (1968) Field Studies of the Effect of Abate 4-E ULV on Quail. (Unpublished study received Nov 26, 1971 under 241-132; prepared by Mississippi State Univ., Dept. of Entomology, submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-F)
005000656	Moore, C.G. (1977) Insecticide avoidance by ovipositing Aedes
005000965	Moore, C.G.; Jimenez-Garcia, J. (1974) Insecticide avoidance by ovipositing Aedes aegypti. Pages 52, In Proceedings of the Papers of the Annual Conference of the California Mosquito Control Association. Vol. 42. Visalia, Calif.: CMCA.
005000661	Moore, C.G.; Jiminez-Garcia, J. (1974) Insecticide avoidance by ovipositing Aedes aegypti. Pages 52, In Proceedings and Papers of the Forty-second Annual Conference of the California Mosquito Control Association; Feb 24-27, 1974, Anaheim, California. Edited by T.D. Mulhern. Visalia, Calif.: CMCA Press.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000007084	Morgan, L.W.; Canerday, T.D.; French, J.C.; et al. (1969) Control of Foliage Feeding Insects on Peanuts: State Project 2-110: (USDA 10,353). (Unpublished study received Apr 26, 1971 under 1F1158; prepared by Univ. of Georgia, Coastal Plain Experiment Station, submitted by E.I. du Pont de Nemours & Co., Inc.; CDL: 090951-J)
005000882	Morgan, L.W.; French, J.C. (1971) Granulate cutworm control in peanuts in Georgia. Journal of Economic Entomology 64(4):937-939.
005010806	Morgan, L.W.; Todd, J.W. (1975) Insecticidal baits for control of corn earworm on peanuts and soybeans and green cloverworm on soybeans. Journal of the Georgia Entomological Society 10(1):18-25.
000001357	Morici, I.J.; Levinskas, G.J. (1965) CL 43,913 and CL 52,160: Seven-Day Repeated Feeding to Young Coturnix Quail: Report No. 65-44. (Unpublished study received Feb 3, 1967 under C7G0566; submitted by American Cyanamid Co., Princeton, N.J.; CDL:090712-O)
005001277	Morishita, F.S.; Jefferson, R.N.; Humphrey, W.A.; Besemer, S.T. (1969) Control of thrips on chrysanthemums grown for cut flowers. California Agriculture 23(7):16-17.
005000852	Moriya, K.; Yabe, T.; Harada, F. (1969) Resistance levels in larvae of Culex tritaeniorhynchus summorosus in Kanagawa Prefecture. Eisei Dobutsu. [Japanese Journal of Sanitary Zoology.] 20(3):161-167.
005000827	Morris, O.N. (1977) Compatibility of 27 chemical insecticides with Bacillus thuringiensis Var. kurstaki. The Canadian Entomologist 109(6):855-864.
005002673	Mouchet, J.; DeJardin, J.; Barathe, J.; Sannier, C.; Sales, S. (1972) Doses discriminatives pour la resistance d'Aedes quelques elements susceptibles de modifier les resultats des tests. [Dose for detecting the resistance of Aedes aegypti to organophosphorus insecticides and study of some elements capable of modifying test results.] Cahiers ORSTOM, Serie Entomologie Medicale et Parasitologie X(1):77-83.
005011370	Mouchet, J.; Dejardin, J.; Subra, R. (1968) Sensibilite aux insecticides de Culex pipiens fatigans en Afrique de l'Ouest. Western Africa.] Medecine Tropicale (Marseille) 28(3):374-387.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005004497	Mouchet, J.; Pichon, G.; Gayral, P.; Hamon, J. (1971) Sensibilite et resistance aux insecticides d'Aedes aegypti en Afrique de l'Ouest et methodes de controle de ce vecteur. [Sensitivity and resistance of Aedes aegypti in western Africa to insecticides and methods for vector control.] Bulletin of the World Health Organization 45(3):394-404.
005002852	Mouchet, J.; Quelennec, G.; Berl, D.; Sechan, Y.; Grebaut, S. (1977) Methodologie pour tester la sensibilite aux insecticides des larves de Simulium damnosum s.l. [Methodology for testing insecticide susceptibility of Simulium damnosum s.l. larvae.] Cahiers ORSTOM, Serie Entomologie Medicale et Parasitologie XV(1):55-66.
005007799	Mouchet, J.; Quiroga, M. (1976) La resistance aux insecticides chez les Culicines. [The resistance to insecticides of the Culicines.] Cahiers ORSTOM, Serie Entomologie Medicale et Parasitologie XIV(2):111-123.
005000865	Mount, G.A.; Hirst, J.M.; McWilliams, J.G.; Lofgren, C.S.; White, S.A. (1968) Insecticides for control of the lone star tick tested in the laboratory and as high- and ultra-low-volume sprays in wooded areas. Journal of Economic Entomology 61(4):1005-1007.
005001248	Moye, H.A. (1975) High speed liquid chromatography of pesticides. Journal of Chromatographic Science 13(6):268-279.
000001242	Moyer, M.; Orloski, E.J. (1965) Residues of CL 52,160 in Rice Grain and Plants: Report No. C-73. (Unpublished study received May 27, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-U)
005002857	Mozes, I.A. (1977) Morfologicheskie izmeneniya v organakh dykhaniya pri smertel'nykh otravleniyakh fosfororganicheskimi soedineniyami i ikh sudebnomeditsinskoe znachenie. [poisonings with organophosphorus compounds and their medicolegal significance.] Sudebno-Meditsinskaya Ekspertiza.
005000956	Muirhead-Thomson, J. (1974) Blackfly and the environment. New Scientist 62(898):384-385.
005000950	Muirhead-Thomson, R.C. (1977) Comparative tolerance levels of black fly (Simulium) larvae to Permethrin (NRDC 143) and temephos. Mosquito News 37(2):172-179.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005019464	Muirhead-Thomson, R.C. (1979) Experimental studies on macroinvertebrate predator-prey impact of pesticides. The reactions of Rhyacophila and Hydrosyche (Trichoptera) larvae to Simulium larvicides. Canadian Journal of Zoology 57(11):2264-2270.
005000955	Muirhead-Thomson, R.C.; Merryweather, J. (1969) Effect of larvicides on Simulium eggs. Nature (London) 221(5183):858-859.
005000667	Mulla, M.S. (1964) Development of new mosquito larvicides. Pest Control 32(9):26,28,32,34.
000001332	Mulla, M.S. (1966) Toxicity of new organic insecticides to mosquito fish and some other aquatic organisms. Mosquito News 26(1): 87-91. (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-AA)
005000944	Mulla, M.S.; Barnard, D.R.; Norland, R.L. (1975) Chironomid midges and their control in Spring Valley Lake, California. Mosquito News 35(3):389-395.
005001273	Mulla, M.S.; Darwazeh, H.A. (1978) Effectiveness and residual activity of polymer formulations of mosquito larvicides. Pages 123-125, In Proceedings and Papers--Annual Conference--California Mosquito Vector Control Association. Vol. 46. Visalia, Calif.: CMCA.
005000767	Mulla, M.S.; Darwazeh, H.A.; Dhillon, M.S. (1977) Cemetery mosquitoes and their control with organophosphorus larvicides and the insect growth regulator methoprene. Pages 162-165, In Proceedings and Papers of the Annual Conference of the California Mosquito and Vector Control Association. Vol. 45. Visalia, Calif.: CMCA.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000011048	Mulla, M.S.; Darwazeh, H.A.; Dhillon, M.S. (1977) Cemetery mosquitoes and their control with organophosphorus larvicides and the insect growth regulator Methoprene. Pages 162-165, In Proceedings and Papers of the Forty-Fifth Annual Conference of the California Mosquito and Vector Control Association, Inc.; Feb 13-16, 1977, Visalia, Calif. Visalia, Calif.: C.M.V.C.A., Inc. (Also In unpublished submission received Dec 6, 1977 under 20954-13; submitted by Zoecon Corp., Palo Alto, Calif.; CDL: 232439-M)
005000870	Mulla, M.S.; Darwazeh, H.A.; Geib, A.F.; Westlake, W.E. (1969) Control of pasture Aedes mosquitoes by dripping larvicides into flowing water, with notes on residue in a pasture hatitat. Journal of Economic Entomology 62(2):365-370.
005000923	Mulla, M.S.; Darwazeh, H.A.; Peters, D.R. (1970) Mosquito control in sewage oxidation ponds with drip and pour-in larvicides. Mosquito News 30(3):456-460.
005000868	Mulla, M.S.; Khasawinah, A.M. (1969) Laboratory and field evaluation of larvicides against chironomid midges. Journal of Economic Entomology 62(1):37-41.
005020463	Mulla, M.S.; Majori, G.; Arata, A.A. (1979) Impact of biological and chemical mosquito control agents on nontarget biota in aquatic ecosystems. Pages 121-173, In Residue Reviews. Vol. 71. New York: Springer-Verlag.
005001105	Mulla, M.S.; Metcalf, R.L.; Geib, A.F. (1966) Laboratory and field evaluation of new mosquito larvicides. Mosquito News 26(2):236-242.
005000873	Mulla, M.S.; Norland, R.L.; Fanara, D.M.; Darwazeh, H.A.; McKean, D.W. (1971) Control of chironomid midges in recreational lakes. Journal of Economic Entomology 64(1):300-307.
005000838	Mulla, M.S.; Norland, R.L.; Westlake, W.E.; Dell, B.; St. Amant, J. (1973) Aquatic midge larvicides, their efficacy and residues in water, soil, and fish in a warm-water lake. Environmental Entomology 2(1):58-65.
005001024	Mulla, M.S.; St. Amant, J.; Anderson, L.D. (1967) Evaluation of organic pesticides for possible use as fish toxicants. Progressive Fish and Culturist 29(1):36-42.
005000813	Muller, R. (1970) Laboratory experiments on the control of cyclopstransmitting guinea worm. Bulletin of the World Health Organization 42(4):563-567.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000001325	Murphey, F.J.; Lake, R.W. (1966) The Toxicity of Mosquito Control Insecticides to Wildlife: Section 11-B. (Unpublished study received Nov 26, 1971 under 241-132; prepared by Univ. of Delaware, submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-S)
005000976	Murphy, S.D.; Cheever, K.L. (1971) Inhibition of rat tissue carboxylesterases and cholinesterase by Abate (O,O,O',O'-tetramethyl O,O'-thiodi-p-phenylene phosphorothioate). Toxicology and Applied Pharmacology 19(2):366-367.
005000800	Murphy, S.D.; Cheever, K.L. (1972) Carboxylesterase and cholinesterase inhibition in rats: Abate and interaction with malathion. Archives of Environmental Health 24(2):107-114.
005000751	Myers, C.M.; Gillies, P.A.; Frolli, R.F. (1969) Field tests of Abate and parathion granules in catch basins. Pages 61-63, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 37. Visalia, Calif.: CMCA.
005000844	Nagy, B.; Nadasy, M.; Kecskemeti, I. (1974) Preparation and examination of a new, highly potent larvicide. Hungarian Journal of Industrial Chemistry. Suppl 2(3):365-376.
005002559	Niazi, A.; Hat Hat, Y.A.K. (1973) Anti-larval activities in Iraq. Bulletin of Endemic Diseases 14(1/4):61-71.
000001373	Nielsen, R.S. (1965) Monthly Report: 52160 Lygus Control on Cotton. (Unpublished study received Jul 17, 1967 under 7G0566; submitted by American Cyanamid Co., Princeton, N.J.; CDL:090713-N)
000001372	Nielsen, R.S. (1966) Monthly Report: [52160 Lygus Control on Cotton]. (Unpublished study received Jul 17, 1967 under 7G0566; submitted by American Cyanamid Co., Princeton, N.J.; CDL:090713-M)
005021997	Nishiuchi, Y. (1979) Noyakuseizai no sushu tansuisan dobutsu ni taisuru dokusei--XXXXXVII. [Toxicity of pesticides to some freshwater organisms--LVII.] Suisan Zoshoku. [Aquiculture.] 27(1):36-41.
005001107	Novak, D. (1972) Granular Abate for control of mosquitoes larvae. Archives Roumaines de Pathologie Experimentale et de Microbiologie 31(4):579-581.
005000668	O'Brien, R.D. (1962) Organophosphorus insecticides. World Review of Pest Control 1(1):29-37.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000889	O'Neal, J.; Mangum, C.L.; Harris, T.M. (1975) Evaluation of insecticidal dips for effectiveness against Diaprepes Economic Entomology 68(5):650-652.
000001395	O'Neil, J.B. (1964) Malathion et al--Toxicity--Marine Animals. (Unpublished study received Jul 9, 1964 under unknown admin. no.; prepared in cooperation with Bureau of Marine Fisheries Laboratory, submitted by American Cyanamid Co., Princeton, N.J.; CDL:221977-A)
000001319	O'Neil, J.B. (1966) Insecticide Development: Abate--Toxicology--Aedes Program: Monthly Report, July 1966. (Unpublished study received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-M)
005000759	Oldham, M.; Lusk, E.E.; Womeldorf, D.J. (1972) Residual activity of various insecticides in treeholes. Pages 69-71, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 40. Visalia, Calif.: CMCA.
005000952	Oldham, M.L.; Lusk, E.E.; Womeldorf, D.J. (1977) Evaluation of serveral insecticides for the control of larval Aedes sierrensis (Ludlow). Mosquito News 37(2):218-221.
005003580	Olmi, M. (1977) Nuovo insetto nemico del riso in Italia: Donacia Chrysomelidae). [A new pest of rice in Italy: Donacia Chrysomelidae). Riso. [Rice.] 26(3):249-252.
000001298	Orloski, E.J. (1969) Abate® and Abate Sulfoxide Residues in Soil: Report No. C-208. (Unpublished study including report, received Nov 24, 1969 under 241-223; submitted by American Cyanamid Co., Princeton, N.J.; CDL:026905-H)
000001391	Orloski, E.J. (1971) Abate and Abate Sulfoxide Residues in Soil: Report No. C-208. Method dated Jun 17, 1969. (Unpublished study including letter dated Mar 30, 1971 from P.E. Gatterman to E.B. Bagan, received May 24, 1970 under 0F0913; prepared by American Cyanamid Co. in cooperation with Syracuse Univ. Research Corp., submitted by American Cyanamid Co., Princeton, N.J.; CDL:091568-AA)
000009241	Ota, A.K. (1969) Control of Platynota stultana, a tortricid leaf roller, in greenhouses. Journal of Economic Entomology 62(6): 1279-1282. (Also In unpublished submission received Aug 25, 1976 under 352-342; submitted by E.I. du Pont de Nemours & Co., Wilmington, Del.; CDL:227322-G)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005007458	Otsuki, A.; Takaku, T. (1979) Determination of an organophosphorus insecticide in water at parts per billion by reversed phase adsorption liquid chromatography. Analytical Chemistry 51(7):833-835.
005001010	Ott, D.E.; Gunther, F.A. (1966) Automated determination of orthophosphate: an application designed especially for the combusted product of organophosphorous pesticide residues. Bulletin of Environmental Contamination and Toxicology 1(3):90-96.
005014325	Ottinger, R.S.; Blumenthal, J.L.; Dal Porto, D.F.; Gruber, G.I.; Santy, M.J.; Shih, C.C. (1973) Recommended Methods of Reduction, Neutralization, Recovery or Disposal of Hazardous Waste. Volume XIV. Summary of Waste Origins, Forms, and Quantities. Cincinnati, Ohio: U.S. Environmental Protection Agency, National Environmental Research Center. (EPA-670/2-73-053-n; available from: NTIS, Springfield, VA; PB-224 593)
005000699	Palmer, J.S. (1968) Toxicologic effects of a phosphorothioic acid insecticide in cattle. Journal of the American Veterinary Medical Association 152(3):282-284.
005000948	Palmisano, C.T.; Steelman, C.D.; Schilling, P.E. (1976) Relative effects of insecticide usage in Louisiana mosquito control programs on the susceptibility of adult female Culex pipiens quinquetasciatus populations. Mosquito News 36(4):521-527.
000001369	Papke, C.C. (1966) Monthly Report: [Control of Lygus Bugs with 52160 on Cotton]. Summary of study 090713-K. (Unpublished study received Jul 17, 1967 under 7G0566; prepared by American Cyanamid Co. in cooperation with Univ. of Arizona Cotton Research Center, submitted by American Cynamid Co., Princeton, N.J.; CDL:090713-J)
005000781	Pasarela, N.R., inventor; American Cyanamid Co., assignee (1978) Akarizides Mittel auf Basis eines Polyvinylchloridharzes ais Traeger. [Acaricidal agent on a polyvinyl chloride carrier.] German (Fed. Rep.) offenlegungsschrift 27,54,700. Jun 15. 18 p. Int. Cl. 2* A61K31/325; Int. Cl.
005012518	Pasarela, N.R., inventor; American Cyanamid Co., assignee (1979) Acaricidal resin composition containing spiro[cyclopropane-1,-1'-indene]-2-carboxylic acid, 3,3-dimethyl-, alpha-cyano-m-phenoxybenzyl ester. U.S. patent 4,145,409. Mar 20. 6 p. Int. Cl. A 01K 27/00, A 01K 29/00, A 01M 1/20; U.S. Cl. 424/16,.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000007904	Pate, T.L.; Brazzel, J.R. (1966) Small Plot Insecticide Test at State College for Control of Boll Worms and Boll Weevils in Cotton during 1966. (Unpublished study received Apr 16, 1971 under 1G1144; prepared by Mississippi State College, Entomology Dept., submitted by E.I. du Pont de Nemours & Co., Inc., Wilmington, Del.; CDL:090931-R)
005000717	Patterson, R.S.; Wilson, F.L. (1966) Fogging and granule applications are teamed to control chironomid midges on Florida lake fronts. Pest Control 34(6):26,29-30,32.
005000660	Pelsue, F.W. (1971) Black flies in the Southeast Mosquito Abatement District. Pages 50, In Proceedings and Papers of the Thirty-ninth Annual Conference of the California Mosquito Control Association; Jan 25-27, 1971, Oakland, California. Visalia, Calif.: CMCA Press.
005000755	Pelsue, F.W.; McFarland, G.C. (1971) Laboratory and field studies of a new Chironomid species in the Southeast Mosquito Abatement District. Pages 74-79, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 39. Visalia, Calif.: CMCA.
005000662	Pelsue, F.W.; McFarland, G.C.; Beesley, C. (1974) Field evaluation of two insect growth regulators against Chironomid midges in water spreading basins. Pages 157-163, In Proceedings and Papers of the Forty-second Annual Conference of the California Mosquito Control Association; Feb 24-27, 1974, Anaheim, California. Edited by T.D. Mulhern. Visalia, Calif.: CMCA Press.
005000753	Pelsue, F.W.; McFarland, G.C.; Magy, H.I. (1970) Buffalo gnat (Simuliidae) control in the Southeast Mosquito Abatement District. Pages 102-104, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 38. Visalia, Calif.: CMCA.
005003577	Perlova, T.G.; Shvetsova-Shilovskaia, K.D.; Golovkin, G.V. (1976) Primesi v tekhnicheskom difose. [Impurities in commercial-grade diphos.] Khimiya v Sel'skom Khozyaistve.
000004969	Peterson, J.W. (1966) The Effects of Abate, an Experimental Mosquito Larvicide, on Fish, Wildlife and Marine Organisms. (Unpublished study received Mar 15, 1967 under unknown admin. no.; prepared by U.S. Fish and Wildlife Service, Div. of Wildlife Services, Pesticide Surveillance & Monitoring in cooperation with Univ. of Delaware, Dept. of Entomology and Applied Ecology, submitted by American Cyanamid Co., Princeton, N.J.; CDL: 222016-A)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005002856	Pfeifer, G. (1975) Foszforsaveszter tipusu inszekticidek molrefrakcioja. [Molar refraction of insecticides of phosphoric acid ester type.] Nehezvegyipari Kutato Intezet Kozlemenyei. Industry.] 5:97-101.
005002136	Phan-urai, P. (1975) Laboratory Evaluation of Abate Effectiveness in Constant Volume of Polluted Water. Geneva, Switzerland: World Health Organization. (WHO/VBC/75.511)
005000843	Phillips, D.J.H. (1978) Use of biological indicator organisms to quantitate organochlorine pollutants in aquatic environments - a review. Environmental Pollution 16(3):167-229.
005002887	Plato, C. (1972) Differential scanning calorimetry as a general method for determining purity and heat of fusion of high-purity organic chemicals: application to 64 compounds. Analytical Chemistry 44(8):1531-1534.
005002679	Popov, N.A.; Vershinin, P.V.; Ponomarenko, V.F. (1974) Perspektivnost' primeneniya preparata difos v meditsinskoi dezinsektsii. [Perspectives on the use of diphos in medical disinfestation.] Zhurnal Mikrobiologii, Epidemiologii i Immunobiologii. [Journal of Microbiology, Epidemiology and Immunobiology.] (5):127-128.
000001299	Post, A.; Castro, C. (1969) Determination of Abate in Bovine Muscle, Kidney and Liver. (Unpublished study including Determination of Abate in bovine fat, received Nov 24, 1969 under 241-223; prepared by Midwest Research Institute, submitted by American Cyanamid Co., Princeton, N.J.; CDL:026905-K)
000006597	Preiser, F.A.; Sawyer, R.L. (1965) Zolone EC--Potatoes: Final Report No. BB 65-131. (Unpublished study received Dec 7, 1972 under 3F1337; prepared in cooperation with Cornell Univ., submitted by Rhodia, Inc., New Brunswick, N.J.; CDL:094849-S)
005000968	Proctor, N.H.; Casida, J.E. (1975) Organophosphorus and methyl carbamate insecticide teratogenesis: diminished NAD in chicken embryos. Science 190:580-582.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005021730	Public Health Station of Chinwangtao (1978) Shuang liu lin de can xiao ce ding ji qi sha wen you chong xiao li de yan jiu. mosquito larva control.] K'un Ch'ung Hsueh Pao. Acta Entomological Sinica. [Chinese Journal of Entomology.] 21(1):49-54.
005003374	Quelennec, G. (1970) Essais sur le terrain de nouvelles formulations d'insecticides, OMS-187, OMS-786 et OMS-971, contre les larves de simulies. [Field trials of the new insecticide formulations, OMS-187, OMS-786 and OMS-971 against Simulium damnosum larvae.] Bulletin of the World Health Organization 43(2):313-316.
005000720	Quidet, P. (1975) Resultats des enquetes realisees en France en 1972, 1973, 1974 sur les causes de mortalites du gibier: influence des pesticides et evaluation des risques selon la nature des produits. [Results of investigations made in France in 1972, 1973, 1974 on the causes of mortlity in game species: influence of pesticides and evaluation of the risks with respect to the nature of the products.] Phytoma 269:26-32.
005003733	Radda, A.; Kunz, C.; Pretzmann, G. (1974) Ueber die Wirksamkeit von Abate gegen Ixodes ricinus. [The effectiveness of Abate against Ixodes ricinus.] Anzeiger fuer Schaedlingskunde, Pflanzen-und Umweltschutz 47(8):122-124.
005004486	Ragab, M.T.H. (1967) Direct fluorescent detection of organothiophosphorus pesticides and some of their sulfur-containing breakdown products after thin layer chromatography. Journal of the Association of Official Analytical Chemists 50(5):1088-1098.
005002338	Ragab, M.T.H. (1967) 4-(p-nitrobenzyl)pyridine as a spray reagent for organophosphorus pesticides and some of their breakdown products on thin-layer chromatograms. Bulletin of Environmental Contamination and Toxicology 2(5):279-288.
005000765	Ranasinghe, L.B.E.; Georghiou, G.P. (1976) An attempt to isolate organophosphorus resistance mechanisms in Culex pipiens 91-92, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 44. Visalia, Calif.: CMCA.
005020438	Ranasinghe, L.E.; Georghiou, G.P. (1979) Comparative modification of insecticide-resistance spectrum of Culex pipiens fatigens Wied. by selection with temephos and temephos/synergist combinations. Pesticide Science 10(6):502-508.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000811	Rangsriyam, Y.; Prownebon, S.; Hirakoso, S. (1968) Effects of insecticides on the feeding activity of the guppy, a mosquito-eating fish, in Thailand. Bulletin of the World Health Organization 39(6):977-980.
005009939	Rathburn, C.B., Jr. (1979) Insecticides for the control of mosquitoes and other Diptera. Mosquito News 39(1):58-63.
005010431	Redfern, R.E.; Walker, R.L.; Cantu, E. (1967) Promising Candidate Insecticides and Acaricides Evaluated at Brownsville, Tex., July 1963 to July 1966. Washington, D.C.: U.S. Department of Agriculture, Agricultural Research Service. (USDA report no. ARS 33-122)
005000640	Reese, C.D.; Becker, D.L. (1972) The Movement and Impact of Pesticides Used for Vector Control on the Aquatic Environment in the Northeastern United States. Environmental Protection Agency, Office of Water Programs, Applied Technology Division, Rural Wastes Branch. Cambridge, Mass.: Arthur D. Little. (Available from: GPO, Washington, D.C.: EP2.25:9; OWP Contract no. 68-01-0129; Pesticide study series-9)
005003350	Reish, D.J.; Kauwling, T.J.; Mearns, A.J.; Oshida, P.S.; Rossi, S.S.; Wilkes, F.G.; Ray, M.J. (1978) Marine and estuarine pollution. Journal of the Water Pollution Control Federation 50(6):1424-1469.
005000786	Rettich, F. (1976) Changes in susceptibility to temephos, pirimiphos-methyl, fenitrothion, and bromophos during preimaginal development of mosquitoes. Acta Entomologica Bohemoslovaca 73(6):382-387.
005000954	Rettich, F. (1977) The susceptibility of mosquito larvae to eighteen insecticides in Czechoslovakia. Mosquito News 37(2):252-257.
005008239	Rettich, F. (1979) Laboratory and field investigations in Czechoslovakia with fenitrothion, pirimiphos-methyl, temephos and other organophosphorous larvicides applied as sprays for control of Culex pipiens molestus Forskal and Aedes cantans Meigan. Mosquito News 39(2):320-328.
005003373	Rettich, F.; Privora, M. (1973) Freilandversuche mit dem Granulat "Abate" gegen Stechmueckenlarven in einem Auwald. (Diptera: Culicidae). [Field experiments with the granulate "Abate" against mosquito larvae in a lowland forest. (Diptera: Culicidae).] Zeitschrift fuer Angewandte Zoologie 60(3):293-298.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000653	Rice, R.E.; Reynolds, H.T.; Cudney, D.W. (1969) Organophosphorus resistance of cotton leaf perforator in areas infested by pink bollworm. California Agriculture 23(6):12-14.
005001112	Rice, R.E.; Reynolds, H.T.; Hannibal, R.M. (1969) Chemical control of pink bollworm in Imperial Valley. California Agriculture 23(5):19.
005000986	Rifaat, M.A.; Khalil, H.M.; Gad, A.M.; Sadek, S. (1974) Effect of sublethal concentrations of the insecticides DDT, Abate and Sevin applied to 3rd stage larvae of Anopheles pharoensis on malaria cycle in the adult mosquito. Journal of the Egyptian Public Health Association XLIX(6):329-340.
005000995	Rio, D.F. (1970) A laboratory study of the effects of Abate and malathion on the fiddler crab. Pages 99-102, In Proceedings of the Fifty-seventh Annual Meeting of the New Jersey Mosquito Extermination Association. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005000832	Rio, D.F. (1972) The survival and reproductive potential of Cyclops vernalis (Crustacea Copepoda) as affected by the organophosphorous mosquito larvicide Abate and three selected ecological factors. Dissertation Abstracts International B 32(9):5541.
005011387	Roberts, D.R.; Miller, T.A. (1971) Effects of Polymer Formulations of Dursban and Abate on Non-target Organism Populations. Edgewood Arsenal, Md.: U.S. Army Environmental Hygiene Agency. (U.S. Army Environmental Hygiene Agency entomological special study no. 31-004-71; available from: NTIS, Springfield, VA; AD-729 342)
005000650	Roberts, D.R.; Stoetzel, M.B.; Loy, V.A. (1969) Laboratory evaluation of two organophosphorus larvicides against pupae of Culex restuans Theobald and influence on adult longevity. Mosquito News 29(1):62-65.
005011388	Roberts, L.W.; Miller, T.A. (1970) Polymer Formulations of Dursban and Abate as Mosquito Larvicides. Edgewood Arsenal, Md.: U.S. Army Environmental Hygiene Agency. (U.S. Army Environmental Hygiene Agency entomological special study no. 31-006-71; available from: NTIS, Springfield, VA; AD-729 343)
005000775	Roberts, L.W.; Miller, T.A. (1970) Polymer Formulations of Dursban and Abate as Mosquito Larvicides, April-October, 1970. Edgewood Arsenal, Md.: U.S. Army Environmental Hygiene Agency. (Entomological special study no. 31-006-71; available from: NTIS, Springfield, Va.: AD-729 343)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005001774	Roberts, R.S. (1969) A review of pesticides used in environmental health activities. Journal of Environmental Health 31(6):560-567.
000001371	Roeuer, K. (1966) An Evaluation of American Cyanamid Compound 52160 for the Control of Lygus in Arizona: WESTAG 66015. (Unpublished study received Jul 17, 1967 under 7G0566; submitted by American Cyanamid Co., Princeton, N.J.; CDL:090713-L)
005001022	Rosen, J.D. (1972) Conversion of pesticides under environmental conditions. Pages 85-96, In Environmental Quality and Safety. Vol. I. Edited by F. Coulston and F. Korte. New York, N.Y.: Academic Press.
005000729	Rosen, J.D. (1972) The photochemistry of several pesticides. Pages 435-447, In Environmental Toxicology of Pesticides: Proceedings of a U.S.-Japanese Seminar; Oct 1971, Oiso, Japan. Edited by Fumio Matsumura, G. Mallory Bousch and Tomomasa Misato. New York: Academic Press.
005003645	Roslavtseva, G.B.; Ivanova, G.B.; Poliakova, V.K. (1973) Rezistentnosty komnatnykh mukh k benzofosfatu. [Resistance of houseflies to benzophosphate.] Khimiya v Sel'skom Khozyaistve.
005000727	Roslavtseva, S.; Ivanova, G. (1975) Specific features of resistance development of houseflies to organophosphate insecticides. Pages 447-449, In Environmental Quality and Safety Supplement. Vol. III: Pesticides. International Union of Pure and Applied Chemistry, Third International Congress; Jul 3-9, 1974, Helsinki, Finland. Edited by Frederick Coulston and Friedhelm Korte. Stuttgart, West Germany: George Thieme Publishers.
005002854	Roslavtseva, S.A.; Ivanova, G.B. (1974) Razvitie resistentnosti komnatnykh mukh k kumafosu v zavisimosti ot proiskhozhdeniya populyatsii. [Development of the resistance of houseflies to coumaphos in relation to the origin of the populations.] Khimiya v Sel'skom Khozyaistve. [Chemistry in Agriculture.] 12(4):283-284.
005003576	Roslavtseva, S.A.; Ivanova, G.B.; Poliakova, V.K.; Agashkova, T.M.; Skvortsova, I.V. (1975) O skorosti razvitiya i utraty rezistentnosti komnatnykh mukh k khlorofosu. [Rate of development and loss of resistance of houseflies to chlorophos.] Khimiya v Sel'skom Khozyaistve. [Chemistry in Agriculture.] 13(11):834-837.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005013684	Roslavtseva, S.A.; Ivanova, G.B.; Polyakova, V.K. (1973) Insektitisidy dlya bor'by s rezistentnymi k khlorofosu komnatnymi mukhami. [Insecticides for combating houseflies resistant to chlorophos.] Veterinariya (Moscow). [Veterinary Science.] (8):35.
005010983	Roslavtseva, S.A.; Ivanova, G.B.; Spirina, T.A.; Agashkova, T.M.; Skvortsova, I.V. (1979) Rezistentnost' komnatnykh mukh k Gardone. [Resistance of houseflies to Gardona.] Khimiya v Sel'skom Khozyaistve. [Chemistry in Agriculture.] 17(1):52-55.
005017548	Roslavtseva, S.A.; Ivanova, R.B. (1972) Razvitie rezistentnosti komnatnykh mukh k fozalonu. [Development of resistance to phosalone by the housefly.] Pages 176-178, In Problemy Parazitologii, Trudy Nauchnoi Konferentsii Parazitologov Ukrainskoi SSR, 7th. [Problems of Parasitology, Transactions of the 7th Scientific Conference of Parasi clogists of the Ukrainian SSR.]; Kiev, Russia. Vol. 2. Edited by A.P. Markevich. Kiev, USSR: Naukova Dumka.
005007005	Roslavtseva, S.A.; Spirina, T.A.; Agashkova, T.M.; Polyakova, V.K. (1978) Razvitie rezistentnosti komnatnykh mukh k ortenu. Sel'skom Khozyaistve. [Chemistry in Agriculture.] 16(1):51-54.
005010318	Ross, R.T.; Biros, F.J. (1970) Correlations between ?31*P n.m.r. chemical shifts and structures of some organophosphorus pesticides. Analytica Chimica Acta 52(1):139-141.
005000846	Roy, R.G.; Sitaraman, N.L.; Ghosh, R.B.; Samson, F.; Balakrishnan, R.; Chandrahasan, A.; Augustine, Y.; Pandy, A.; Ponnappan, C.V. (1978) Susceptibility of A. stephensi from some urban areas in Tamil Nadu and Karnataka to some commonly used insecticides. Indian Journal of Medical Research 67(6):947-952.
005000743	Ruber, E. (1971) Primary producers, pesticides and pyramids. Pages 162-174, In Proceedings of the Fifty-eighth Annual Meeting of the New Jersey Mosquito Extermination Association. Vol. 58. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
000001327	Ruber, E.; Baskar, J. (1968) Sensitivities of selected microcrustacea to eight mosquito toxicants. Pages 99-103, In Proceedings of Fifty-fifth Annual Meeting. (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-V)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005002681	Ruzimuradov, A. (1978) Organizatsiya meropriyatii protiv zoofil'nykh mukh. [Organization of safeguards against zoophilous flies.] Veterinariya (Moscow). [Veterinary Science.] (1):34-36.
005003737	Sacca, G.; Gandolfo, D.; Mastrilli, M.L.; Stella, E. (1971) Richerche di laboratorio e di campo per il controllo chimico di Culex pipiens L. con larvicidi. [Laboratory and field research on the chemical control of Culex pipiens L. with larvicides.] Parassitologia XIII(1/2):345-353.
005003376	Sacca, G.; Mastrilli, M.L. (1972) Tossicometria di alcuni esteri fosforici in Gambusia affinis holbrooki (Gir.) e in Culex in Gambusia affinis holbrooki (Gir.) and Culex pipiens L.] Parassitologia XIV(2/3):365-371.
005000984	Sadek, S.; Gad, A.M.; Rifaat, M.A.; Khalil, H.M. (1974) Effect of sublethal concentrations of the insecticides DDT, Abate and Sevin, applied to 3rd stage larvae of Culex pipiens molestus on certain biological aspects of the mosquito. Journal of the Egyptian Public Health Association XLIX(6):314-328.
000015099	Sakamoto, S.S.; Ewart, W.H.; Brawner, O.L.; et al. (1975) Orthene 75 S: Thrips Control in Citrus: General Summary of Data and Justification for Label Claims. (Unpublished study received Sep 30, 1975 under 239-2418; prepared in cooperation with Univ. of California--Riverside, San Joaquin Valley Agricultural Research Center, Dept. of Entomology and Univ. of Arizona, Yuma Experiment Station, Dept. of Entomology, submitted by Chevron Chemical Co., Richmond, Calif.; CDL:095911-R)
005000848	Samman, J.; Thomas, M.P. (1978) Changes in zooplankton populations in the White Volta with particular reference to the effect of Abate. International Journal of Environmental Studies 12(3):207-214.
005000847	Samman, J.; Thomas, M.P. (1978) Effect of an organophosphorus insecticide, Abate, used in the control of Simulium damnosum on nontarget benthic fauna. International Journal of Environmental Studies 12(2):141-144.
005020324	Samman, J.; Thomas, M.P. (1979) Observations on the effect of the aerial spraying of Abate, used in the control of Simulium larvae, on the drift rates of microcrustacea and Chaoboridae in the River Oti, Ghana. International Journal of Environmental Studies 14(1):43-47.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005005419	Sanchez, A.M.; Tonn, R.; Uribe, L.J.; Calheiros, L.B. (1978) Comparacion de la eficacia de varios metodos de aplicacion de insecticidas para el control o la erradicacion del Aedes different ways of applying insecticides to control or eradicate Aedes aegypti in Colombia.] Boletin de la Oficina Sanitaria Panamericana 84(1):24-37.
005009242	Sanders, H.O. (1969) Toxicity of Pesticides to the Crustacean Gammarus lacustris. Washington, D.C.: U.S. Bureau of Sport Fisheries and Wildlife. (U.S. Bureau of Sport Fisheries and Wildlife technical paper 25)
005010360	Sanders, H.O.; Cope, O.B. (1968) The relative toxicities of several pesticides to naiads of three species of stoneflies. Limnology and Oceanography 13(1):112-117.
005005683	Sandescu-Nosec, I.; Tacu, V.; Durbaca, S. (1975) Efficacite de certains insecticides organophosphores comme ovicides sur le groupe Maculipennis. [The efficiency of some organophosphorous insecticides as ovicides for the Maculipennis group.] Archives Roumaines de Pathologie Experimentale et de Microbiologie 34(3):271-276.
005000963	Santhoy, O.; Morallo-Rejesus, B. (1972) Toxicity of six organophosphorus insecticides to field-collected DDT-resistant strains of rice weevil, Sitophilus oryzae (L) and red flour beetle, Tribolium castaneum (Herbst). Phillippine Entomologist 2(4):283-290.
005000833	Sasaki, R.K. (1974) Organophosphorus insecticides and the fertilization and early embryonic development of Crassostrea viginica Gmelin: an unltra structural analysis. Dissertation Abstracts International B 34(10):4840-4841.
005000907	Sastry, S.C.; Jayakumar, K.; Lakshminarayana, V.; Rao, V.N.S. (1978) Abate - its value as a cyclopscide. Journal of Tropical Medicine and Hygiene 81(8):156-158.
005000851	Scanlon, J.E. (1967) Control of Aedes aegypti (Linn.) in Southeast Asia. Japanese Journal of Medical Science and Biology. Suppl 20:108-112.
000020560	Schafer, E.W. (1972) The acute oral toxicity of 369 pesticidal, pharmaceutical and other chemicals to wild birds. Toxicology and Applied Pharmacology 21(??):315-330. (Also In unpublished submission received Apr 25, 1978 under 476-2180; submitted by Stauffer Chemical Co., Richmond, Calif.; CDL:233577-C)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000744	Schmidt, R.F.; Sutherland, D.J.; Fitzpatrick, G. (1975) Abate-granular larviciding and its uniformity by helicopter. Pages 219-220, In Proceedings of the Sixty-second Annual Meeting of the New Jersey Mosquito Extermination Association. Vol. 62. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005020767	Schoettger, R.A. (1974) Toxicity of herbicides being investigated. Pages 1-5, In Annual Progress Report: 1973-1974, Fish-Pesticide Research Laboratory, Fish and Wildlife Service, Columbia, Missouri. Columbia, Mo.: Fish-Pesticide Research Laboratory.
005001017	Schoof, H.F. (1967) Insecticides for use against Aedes aegypti. Bulletin of the World Health Organization 36(4):618-622.
005019095	Schooley, D.A.; Quistad, G.B. (1979) High-pressure, high-resolution liquid chromatography and its application to pesticide analysis and biochemistry. Pages 1-113, In Progress in Drug Metabolism. Vol. 3. Edited by J.W. Bridges and L.F. Chasseaud. New York: John Wiley and Sons.
005001117	Schrader, G. (1965) The modification of biological activity by structural changes in certain organophosphorus compounds. World Review of Pest Control 4(4):140-149.
005000649	Schultz, H.A.; Webb, A.B. (1969) Laboratory bioassay of pesticide-impregnated rubber as a mosquito larvicide. Mosquito News 29(1):38-41.
005003372	Scirocchi, A.; D'Erme, A. (1977) Prove di laboratorio e di campo con alcuni insetticidi larvicidi per il controllo di Culex insecticides for the control of Culex pipiens L.] Rivista di Parassitologia. [Review of Parasitology.] XXXVIII(1):75-88.
005000897	Sechriest, R.E.; Sherrod, D.W. (1977) Pelleted bait for control of the black cutworm in corn. Journal of Economic Entomology 70(6):699-700.
005000324	See, G.G.; Yang, P.Y.; Kacholia, K.K. (1976) Research and Development of an Ozone Contractor System. Final Report. Washington, D.C.: U.S. Army Medical Research and Development Command. (Contract no. DAMD17-76-C-6041; available from NTIS, Springfield, Va;).
005000814	Self, L.S.; Tun, M.M. (1970) Summary of field trials in 1964-69 in Rangoon, Burma, of organophosphorus larvicides and oils against Culex pipiens fatigans larvae in polluted water. Bulletin of the World Health Organization 43(6):841-851.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005001111	Service, M.W. (1971) Conservation and the control of biting flies in temperate regions. Biological Conservation 3(2):113-122.
005020217	Sexton, J.D. (1978) The Study of Abate on Selected Non-Target Aquatic Organisms Under Laboratory and Field Conditions Minnesota. (Available from: University Microfilms, Ann Arbor, MI; 79-12080)
000001385	Shaffer, C.B. (1978) Addendum to Report of Two-Year Chronic Toxicity and Carcinogenesis Study of Temephos in the Rat. (Unpublished study received Apr 4, 1978 under 241-220; submitted by American Cyanamid Co., Princeton, N.J.; CDL:233402-A)
005000805	Shafik, M.T. (1968) A gas chromatographic method for the determination of low concentrations of Abate in water. Bulletin of Environmental Contamination and Toxicology 3(5):309-316.
005001015	Shafik, M.T.; Bradway, D.; Enos, H.F. (1971) A method for confirmation of organophosphorus compounds at the residue level. Bulletin of Environmental Contamination and Toxicology 6(1):55-66.
005016102	Sherma, J. (1973) Thin-layer chromatography: recent advances. Pages 3-87, In Thin-Layer and Liquid Chromatography and Analyses of Pesticides of International Importance. Edited by G. Zweig and J. Sherma. New York: Academic Press. (Analytical methods for pesticides and plant growth regulators, vol. VII)
005001026	Sherman, M.; Ross, E.; Chang, M.T.Y. (1964) Acute and subacute toxicity of several organophosphorus insecticides to chicks. Toxicology and Applied Pharmacology 6(2):147-153.
005019327	Siewierski, M. (1979) Chemistry and transformation of chemicals added to the environment [abstract no. 9.0089]. Toxicology Research Projects Directory. Section 1 4(9):12.
000012430	Simpson, R.G. (1964) Chemical Spray Trials. (Unpublished study received Nov 10, 1972 under 3F1323; prepared by Colorado State Univ., Dept. of Entomology, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:093552-F)
005003644	Sinegre, G.; Cousserans, J.; Vigo, G.; Crespo, O. (1971) Sensibilite des larves de Mansonia (C.) richiardii (Ficalbi) a quelques insecticides: effet de l'abate dans un biotope larvaire. [Sensitivity of Mansonia (C.) richiardii (Ficalbi) larvae to some insecticides: effect of abate in a larval biotope.] Cahiers ORSTOM, Serie Entomologie Medicale et Parasitologie IX(3):255-263.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005021621	Sinegre, G.; Gaven, B.; Jullien, J.L. (1977) Activite comparee de 31 insecticides sur des larves de Culex pipiens (L.) sensibles et resistantes au chlorpyrifos dans le midi de la France. (methoprene) in the south of France.] Parassitologia XIX(1/2):63-72.
005000970	Singh, K. I. (1973) Evaluation of insecticides against four strains of the house fly, Musca domestica L. (Diptera: Muscidae) from West Malaysia. Southeast Asian Journal of Tropical Medicine and Public Health 4(4):554-559.
005000746	Sjogren, R.D.; Mulla, M.S. (1968) Drip application of three organophosphorus insecticides for mosquito control. Pages 40, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 36. Visalia, Calif.: CMCA.
005000915	Sjogren, R.D.; Mulla, M.S. (1968) Drip application of three organophosphorus insecticides for mosquito control. Mosquito News 28(2):172-177.
005000908	Skrinde, R.T. (1969) A review of the literature of 1968 on wastewater and water pollution control. Analytical methods: organics. Journal of the Water Pollution Control Federation 41(6):875-888.
005000909	Skrinde, R.T. (1971) Addendum to review of 1970 literature. Analytical methods: organics. Journal of the Water Pollution Control Federation 43(7):1550-1557.
000013679	Sleesman, J.; Sakamoto, S.S.; Wells, A.; et al. (1977) [Evaluation of Various Insecticides on Vegetable Crops]: Report No. 31715. (Unpublished study including letter, report no. 36267, dated Sep 23, 1966 from W.H. Lange to Sid Sakamoto; letter, report no. 39337, dated Jan 4, 1974 from H.B. Green to O. Elmo Shipp and report nos. 31735, 36249, 36250, 36260 through 36263, 36265, 36266, 36268, 36269, 36279 through 36282, 36344, 37576, 38837, 39461, 40038, 40134, 40173, 40175, 40185, 40276, 41253, 42715, 42751, 46857, 51519, 55036, 55913, and 55926, received Jul 24, 1978 under NJ 78/14; prepared by Ohio Agricultural Research and Development Center, submitted by Mobay Chemical Corp., Agricultural Div., Kansas City, Mo.; CDL:235607-B)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000013675	Sleesman, J.; Wells, A.; Sigler, D.; et al. (1973) [Insect Control on Cabbage, Tomato, Pepper, Eggplant, Turnip, Lima Beans and Watermelon]. (Unpublished study including letter dated Sep 23, 1966 from W.H. Lange to Sid Sakamoto and report nos. 31715, 31726, 31733, 31723, 31735, 32153, 36249 through 36254, 36259 through 36270, 36274 through 36282, 36337, 36341, 36342 and 36344 through 36348, received Jul 24, 1973 under 4F1424; submitted by Mobay Chemical Corp., Agricultural Div., Kansas City, Mo.; CDL:094861-H)
000001309	Smith, C.N.; Roos, J. (1966) Results of Tests with Abate from American Cyanamid Company. (Unpublished study received Sep 1, 1972 under 241-230; prepared by U.S. Agricultural Research Service, Entomology Research Div., Insects Affecting Man and Animals Research Branch, submitted by American Cyanamid Co., Princeton, N.J.; CDL:002063-H)
000006333	Smyth, E.P.; Johansen, C. (1970) Final Report: Report No. 164670-015. (Unpublished study received Mar 24, 1976 under 359-620; prepared by Rhodia, Inc. in cooperation with Washington State Univ., submitted by Rhone-Poulenc, Inc., Monmouth Junction, N.J.; CDL:095197-AP)
005002919	Somermaa, K. (1970) Bekampningsforsok i laboratorium mot glasvingade angsstriten Javesella (Calligypona) pellucide (F.). [Control experiments in the laboratory against the Delphacia Javesella (Calligypona) pellucide (F.)]. Vaxtskyddsnotiser 34(4):59-64.
005008228	Spehar, R.L.; Holcombe, G.W.; Carlson, R.W.; Drummond, R.A.; Yount, J.D.; Pickering, Q.H. (1979) Effects of pollution on freshwater fish. Journal of the Water Pollution Control Federation 51(6):1616-1694.
005000703	St. John, L.E., Jr.; Lisk, D.J. (1968) Rapid, sensitive residue determination of organophosphorus insecticides by alkali thermionic gas chromatography of their methylated alkyl phosphate hydrolytic products. Journal of Agricultural and Food Chemistry 16(3):408-410.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005002858	Stan, H.J.; Abraham, B.; Jung, J.; Kellert, M.; Steinland, K. (1977) Nachweis von Organophosphorinsecticiden durch Gas-Chromatographie-Massenspektrometrie. [Detection of organophosphorus insecticides by gas chromatography/mass spectrometry.] Fresenius Zeitschrift fuer Analytische Chemie 287(4/5):271-285.
000001329	Steelman, C.D.; Colmer, A.R.; Cabes, L.; Barr, H.T.; Tower, B.A. (1966?) Relative toxicity of selected insecticides to bacterial populations in waste disposal lagoons. Journal of Economic Entomology 60(2):467-468. (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-X)
005000880	Steelman, C.D.; Poche, A.J. (1970) Insecticide-fertilizer mixtures for controlling the dark rice field mosquito. Journal of Economic Entomology 63(6):1986-1987.
005000734	Steinberg, M. (1973) Field studies of topically applied louse toxicants. Pages 181-186, In Proceedings of the International Symposium on the Control of Lice and Louse-born Diseases; Dec 4-6, 1972, Washington, D.C. Washington, D.C.: Pan American Health Organization. (Scientific publication no. 263)
00000 313	Steinberg, M.; Cole, M.M.; Miller, T.A.; Godke, R.A. (1972) Toxicological and entomological field evaluation of Mobam® and Abate® powders used as body louse toxicants (Ano plura: Pediculidae). Journal of Medical Entomology 9(1): 73-77. (Also In unpublished submission received Sep 1, 1972 under 241-230; submitted by American Cyanamid Co., Princeton, N.J.; CDL:002063-L)
005013883	Steinberg, M.; Rowe, S.E.; Boldt, R.E. (1970) Assessment of Relative Toxicity of Candidate Louse Toxicant ENT-27165 O,O,O',O'-Tetramethyl O,O'-Thiodi-P-Phenylene Phosphorothioate (Abate): Evaluation of Animal and Human Exposure Data and Recommendations Concerning a Field Test Program. Edgewood Arsenal, Md.: U.S. Army Environmental Hygiene Agency. (Available from: NTIS, Springfield, VA; AD-869 265)
000012511	Steinhauer, A.L.; Adler, V.E. (1964) Preliminary Summary of Results on Alfalfa Weevil Control--1964. (Unpublished study including letter dated Aug 9, 1965 from A.L. Steinhauer to Gentlemen, received Dec 28, 1967 under 8F0648; prepared by Univ. of Maryland, Dept. of Entomology in cooperation with U.S. Dept. of Agriculture, Alfalfa Weevil Investigations Laboratory, submitted by Geigy Chemical Corp., Ardsley, N.Y.; CDL:091129-S)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000001322	Stickel, W.H. (1967) Wildlife pesticides and mosquito control. Massachusetts Audobon 51(?/Spring):110-122. (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-P)
005005420	Straka, F. (1974) Vuzmozhnosti za borba sreshtu zelevata mukha s nyakoi granulirani preparati. [The possibilities of controlling Hylemya brassicae by some insecticides.] Rastitelna Zashtita. [Plant Protection.] 22(8):31-33.
005001275	Strong, R.G. (1970) Relative susceptibility of confused and red flour beetles to twelve organophosphorus insecticides, with notes on adequacy of the test method. Journal of Economic Entomology 63(1):258-263.
005001108	Strong, R.G.; Sbur, D.E. (1968) Evaluation of insecticides for control of stored-product insects. Journal of Economic Entomology 61(4):1034-1041.
005002677	Subra, R.; Bouchite, B.; Gayral, P. (1970) Evaluation a grande echelle du dursban et de l'abate pour le controle des larves de Culex pipiens fatigans Wiedemann, 1828, dans la ville de Bobo-Dioulasso (Haute-Volta). [Large-scale evaluation of dursban and abate to control the larvae of Culex pipiens (Upper-Volta).] Medecine Tropicale 30(3):393-402.
005004496	Subra, R.; Hebrard, G. (1973) Perspectives offertes par la lutte antilarvaire pour le controle de anopheles gambiae a Mayotte et la Grand Comore. (Archipel des Comores). [The outlook for combating larvae in the control of Anopheles gambiae at Mayotte and Grande Comore (Comoro Archipelago).] Pages 258, In Ninth International Congress on Tropical Medicine and Malaria; Oct 14-21, Athens,Greece. Vol. 1. Abstracts of Invited Papers. Edited by G.J. Papaevangelou. Athens, Greece: Congress Publication Committee. (Abstract no. 373)
005002912	Subra, R.; Hebrard, G.; Rabenirainy, L. (1973) Essai de lutte contre Anopheles gambiae, s.l. et Culex pipiens d'endemie filarienne, (Mayotte, Archipel des Comores). [Trial in the control of Anopheles gambiae s.l., and Culex pipiens filariasis, (Mayotte, Comores Archipelago).] Cahiers ORSTOM, Serie Entomologie Medicale et Parasitologie XI(4):225-231.
005000738	Sutherland, D.J. (1970) Newer chemicals for mosquito control. Pages 177-185, In Proceedings of the Fifty-seventh Annual Meeting of the New Jersey Mosquito Extermination Association. Vol. 57. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000935	Sutherland, D.J. (1971) Susceptibility of New Jersey mosquito larvae to DDT, Abate, and Baytex. Mosquito News 31(4):482-487.
005000939	Sutherland, D.J.; Schmidt, R.F.; Fitzpatrick, G. (1974) Distribution pattern of Abate granules applied by helicopter. Mosquito News 34(3):320-324.
005007498	Sutherland, D.J.; White, D.J. (1978) Distribution pattern of temephos granules applied by the Horn Seeder. Mosquito News 38(1):127-131.
005002913	Suzuki, T. (1968) [A note on a multiple-resistant or -tolerant colony of Culex pipiens to organophosphorus insecticides.] Eisei Dobutsu. [Sanitary Zoology.] 19(2):98-100.
000001250	Swabey, Y.H.; Schenk, C.F. (1966) Preliminary Report on the Use of Abate for Controlling Black Fly Larvae. (Unpublished study received Mar 15, 1967 under unknown admin. no.; prepared by Ontario Water Resources Commission, submitted by American Cyanamid Co., Princeton, N.J.; CDL:222016-B)
000001323	Swabey, Y.H.; Schenk, C.F.; Parker, G.L. (1967) Evaluation of two organophosphorus compounds as blackfly larvacides. Mosquito News 27(2):149-155. (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-Q)
000001390	Syracuse University Research Corporation (1969) Determination of Abate® & Abate Sulfoxide Residues in Soil. (Unpublished study received May 24, 1970 under 0F0913; submitted by American Cyanamid Co., Agricultural Div., Princeton, N.J.; CDL:091568-R)
005002915	Tabaru, Y. (1975) [Outbreak of chironomid midge in a polluted river and the chemical control of the larvae.] Eisei Dobutsu.
005003736	Tabaru, Y.; Ikeda, Y.; Hasegawa, M.; Hattori, K. (1973) [Studies on the control of the biting midges in Hokkaido: III. On the chemical control and vertical distribution of the larvae of Culicoides circumscriptus Kieffer (Diptera: Ceratopogonidae) in Saroma, shore of Lake Saroma, Hokkaido.] Eisei Dobutsu.
005002911	Tabaru, Y.; Ohkura, T., inventors; Sankyo Co., assignee (1977) Jan 12. 3 p. Int. Cl 2*. A 01N 9/36.
005000849	Tadano, T. (1970) Genetics of cross resistance to organophosphates, Abate, fenitrothion and malathion in larvae of Culex pipiens pallens Coquillet. Japanese Journal of Experimental Medicine 40(1):59-66.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005022039	Tanaka, J. (1978) Noyaku no "gyodokusei" ni tsuite. [On the "fish toxicity" of agricultural chemicals.] Seitai No Kagaku.
005000632	Taylor, R.T. (1968) Simulated field evaluation of promising insecticides for use against Aedes aegypti. Mosquito News 28(1):11-16.
005000651	Taylor, R.T.; Schoof, H.F. (1969) Field effectiveness of five organophosphorus compounds as thermal fogs. Mosquito News 29(1):7-8.
005000883	Taylor, R.T.; Schoof, H.F. (1971) Experimental field treatments with larvicides for control of Anopheles, Aedes, and Culex mosquitoes. Journal of Economic Entomology 64(5):1173-1176.
005001003	Thirugnanam, M.; Forgash, A.J. (1975) Environmental impact of mosquito pesticides - influence of Abate on the toxicity and acetylcholinesterase of killifishes. Pages 138-139, In Proceedings of the Sixty-second Annual Meeting of the New Jersey Mosquito Control Association; Mar 11-14, 1975, Atlantic City, New Jersey. New Brunswick, N.J.: New Jersey Mosquito Extermination Association.
005000842	Thirugnanam, M.; Forgash, A.J. (1975) Environmental impact of mosquito pesticides: influence of temefos on the brain acetylcholinesterase of killifish. Environmental Physiology and Biochemistry 5(6):451-459.
000001360	Thomas Ranch (1965) Relative Toxicity of Various Insecticide Sprays, Applied by Air, to a Number of Predatory Insects and Spiders in a Seed Alfafa Field: Table 7. (Unpublished study received Feb 3, 1967 under 7G0566; submitted by American Cyanamid Co., Princeton, N.J.; CDL:090712-R)
005018561	Thornburg, W. (1971) Pesticide residues. Analytical Chemistry 43(5):145-162.
005009700	Thornburg, W. (1973) Pesticide residues. Analytical Chemistry 45(5):151-167.
005017190	Thornburg, W.; Beckman, H. (1969) Pesticide residues. Analytical Chemistry 41(5):140-151.
005001668	Thornburg, W.; Beckman, H. (1969) Pesticide residues. Analytical Chemistry 41(5):140R-151R.
005000325	Tomlin, A.D. (1975) Toxicity of soil applications of insecticides to three species of springtails (Collembola) under laboratory conditions. The Canadian Entomologist 107(7):769-774.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005012234	Tomlin, A.D.; Forgash, A.J. (1972) Toxicity of insecticides to gypsy moth larvae. Journal of Economic Entomology 65(4):953-954.
000001324	Travis, B.V.; Dewey, J.E.; Pendleton, R.F. (1968) Comparative toxicity data on pesticides used for mosquito control. Pages 122-129, In Proceedings of the 55th Annual Meeting; March 1968, Atlantic City, N.J. Atlantic City, N.J.: New Jersey Mosquito Extermination Association. (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-R)
005001104	Travis, B.V.; Guttman, D. (1966) Additional tests with blackfly larvicides. Mosquito News 26(2):157-160.
005000735	Tripp, M.R. (1974) Effects of organophosphate pesticides on adult oysters Crassostrea virginica. Pages 225-236, In Pollution and Physiology of Marine Organisms. Edited by F. John Vernberg and Winona B. Vernberg. New York: Academic Press.
005006141	Tsin Wuang Tao City Revolution Committee Technical Group (1974) observations.] Chung-Hua i Hsueh Tsa Chih. [Chinese Medical Journal.] 54(1):47-49.
005012461	Tucker, R.K.; Haegele, M.A. (1971) Comparative acute oral toxicity of pesticides to six species of birds. Toxicology and Applied Pharmacology 20:57-65.
000001288	Turner, E.C., Jr. (1976?) Temephos Insecticide for Control of Lice on Beef Cattle. (Unpublished study received Jun 2, 1976 under 241-EX-77; prepared by Virginia Polytechnic Institute and State Univ., Dept. of Entomology, submitted by American Cyanamid Co., Princeton, N.J.; CDL:226521-D)
000011703	Tuttle, D.M. (1972) A Summary of Citrus Thrips. (Unpublished study received Jan 16, 1974 under unknown admin. no.; prepared by Univ. of Arizona, Entomology Dept., submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:122290-Q)
000012568	Tuttle, D.M.; Arvizo, G.L. (1964) Chemical Control of Insects and Mites on Alfalfa. (pp. 1,4-6 only; unpublished study received May 8, 1970 under 0F0892; prepared by Univ. of Arizona, Entomology Dept., submitted by Geigy Chemical Corp., Ardsley, N.Y.; CDL:091543-BF)
000012512	Tuttle, D.M.; Arvizo, G.L. (1965) Control of Alfalfa Insects. (Unpublished study received Dec 28, 1967 under 8F0648; prepared by Univ. of Arizona, Dept. of Entomology, submitted by Geigy Chemical Corp., Ardsley, N.Y.; CDL:091129-V)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000001284	Tuttle, D.M.; Arvizo, G.L. (1965) Control of Alfalfa Insects II. (Unpublished study received Feb 10, 1967 under unknown admin. no.; prepared by Univ. of Arizona--Yuma, submitted by American Cyanamid Co., Princeton, N.J.; CDL:122856-D)
000001283	Tuttle, D.M.; Arvizo, G.L. (1966) Control of Alfalfa Insects. (Unpublished study received Feb 10, 1967 under unknown admin. no.; prepared by Univ. of Arizona--Yuma, submitted by American Cyanamid Co., Princeton, N.J.; CDL:122856-C)
000001282	Tuttle, D.M.; Arvizo, G.L. (1966) Control of Alfalfa Insects with Various Insecticides. (Unpublished study received Feb 10, 1967 under unknown admin. no.; prepared by Univ. of Arizona--Yuma, submitted by American Cyanamid Co., Princeton, N.J.; CDL: 122856-B)
000012570	Tuttle, D.M.; Arvizo, G.L. (1966) Control of Alfalfa Insects with Various Insecticides. (Unpublished study received May 8, 1970 under 0F0892; prepared by Univ. of Arizona, submitted by Geigy Chemical Corp., Ardsley, N.Y.; CDL:091543-BI)
000001281	Tuttle, D.M.; Arvizo, G.L. (1966) Evaluation of Several Insecticides on Alfafa Insects. (Unpublished study received Feb 10, 1967 under unknown admin. no.; prepared by Univ. of Arizona--Yuma, submitted by American Cyanamid Co., Princeton, N.J.; CDL:122856-A)
000012571	Tuttle, D.M.; Arvizo, G.L. (1966) Evaluation of Several Insecticides on Alfalfa Insects. (Unpublished study received May 8, 1970 under 0F0892; prepared by Univ. of Arizona, submitted by Geigy Chemical Corp., Ardsley, N.Y.; CDL:091543-BJ)
000011688	Tuttle, D.M.; Arvizo, G.L. (1969) Evaluation of Chemicals for Control of Citrus Thrips on Lemon Trees. (Unpublished study received Jan 16, 1974 under unknown admin. no.; prepared by Univ. of Arizona, Dept. of Entomology, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:122290-B)
000011690	Tuttle, D.M.; Arvizo, G.L. (1971) Chemical Control of Citrus Thrips. (Unpublished study received Jan 16, 1974 under unknown admin. no.; prepared by Univ. of Arizona, Dept. of Entomology, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:122290-D)
000007895	Tuttle, D.M.; Arvizo, G.L. (1971) Chemical Control of Citrus Thrips. (Unpublished study received Mar 24, 1972 under 2F1254; prepared by Univ. of Arizona, Dept. of Entomology, submitted by E.I. du Pont de Nemours & Co., Inc., Wilmington, Del.; CDL: 095333-T)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000011693	Tuttle, D.M.; Arvizo, G.L. (1972) Control of Citrus Thrips on Grapefruit. (Unpublished study received Jan 16, 1974 under unknown admin. no.; prepared by Univ. of Arizona, Dept. of Entomology, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL: 122290-G)
000011694	Tuttle, D.M.; Arvizo, G.L. (1973) Insecticide Treatments for Control of Citrus Thrips on Grapefruit. (Unpublished study received Jan 16, 1974 under unknown admin. no.; prepared by Univ. of Arizona, Dept. of Entomology, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:122290-H)
005001034	U.N. Food and Agriculture Organization, Working Party of Experts on Pesticide Residues (1969) Pesticide Residues in Food; Report of the 1968 Joint Meeting of the FAO Working Party of Experts on Pesticide Residues and the WHO Expert Committee on Pesticide Residues; Dec 9-16, 1968, Geneva, Switzerland. Geneva, Switzerland: World Health Organization. (WHO technical report series no. 417; FAO agricultural report studies no. 78)
000013539	U.S. Fish and Wildlife Service (19??) Pesticide-Wildlife Studies, 1963: A Review of Fish and Wildlife Service Investigations during the Calendar Year. ?: U.S. Dept. of the Interior. (Circular 199; pp. 24,26,28; also In unpublished submission received Mar 4, 1966 under unknown admin. no.; submitted by Mobay Chemical Corp., Agricultural Div., Kansas City, Mo.; CDL:130685-O)
000001316	U.S. Fish and Wildlife Service (1971) Median lethal concentrations (LC_{50}'s) of pesticidal chemicals in 5-day diets of birds. By Bureau of Sport Fisheries and Wildlife, Patuxent Wildlife Research Center, Journal of Wildlife Management* (Oct): ? -? . (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-D)
000014011	U.S. Fish and Wildlife Service, Denver Wildlife Research Center (1966) 1966--Annual Progress Report, Denver Wildlife Research Center: Summary of Toxicity Tests, January 1 to November 21, 1966: Table 1. (Unpublished study received Jun 2, 1971 under 1F1062; submitted by Shell Chemical Co., Washington, D.C.; CDL: 091827-K)
000001406	U.S. Public Health Service (1966) Toxicology of Abate. By Comminicable Disease Center, Toxicology Section. (Unpublished study received Apr 13, 1966 under unknown admin. no.; CDL:129469-A)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005001114	Udomsakdi, S. (1973) Studies on hemorrhagic fever in Thailand 1958-1971: a review. Journal of the Medical Association of Thailand 56(1):40-66.
005002680	Umino, T.; Suzuki, T. (1969) [Effect of two new insecticides, Abate and Dursban, on larvae and adults of mosquitos.] Eisei Dobutsu. [Sanitary Zoology.] 20(2):95-99.
005002675	Vdovin, D.G.; Zhogolev, D.T. (1973) Chuvstvitel'nost' moskitov k insektitsidam abat i gardona. [Sensitivity of sand flies to the insecticides Abate and Gardona.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 42(3):352-353.
005000787	Verma, S.R.; Bhatnagar, M.C.; Dalela, R.C. (1978) Biocides in relation to water pollution. Part 2: bioassay studies of few biocides to a fresh water fish, Channa gachua. Acta Hydrochimica et Hydrobiologica 6(2):137-144.
005014582	von Ruemker, R.; Lawless, E.W.; Meiners, A.F.; Lawrence, K.A.; Kelso, G.L.; Horay, F. (1974) Production, Distribution, Use and Environmental Impact Potential of Selected Pesticides. Washington, D.C.: Council on Environmental Quality. (Available from: NTIS, Springfield, VA; PB-238 795)
000001251	Von Windeguth, D.L.; Patterson, R.S. (1965?) The Effects of Two Organic Phosphate Insecticides on Segments of the Aquatic Biota. (Unpublished study received Nov 29, 1965 under unknown admin. no.; prepared by Florida State Board of Health, submitted by American Cyanamid Co., Princeton, N.J.; CDL:222019-A)
000001320	Von Windeguth, D.L.; Patterson, R.S. (1966) The effects of two organic phosphate insecticides on segments of the aquatic biota. Mosquito News 26(3):377-380. (Also In unpublished submission received Nov 26, 1971 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-N)
005011209	Vrochinskii, K.K. (1975) Gerbitsidy kak potentsial'nyi faktor zagryazneniya vodoemov. [Herbicides as a potential factor of pollution in reservoirs.] Pages 11-15, In Mekhanizm Deistviya Gerbitsidov i Sinteticheskikh Regulyatorov Rosta Rastenii Ikh Sud'ba v Biosfere, Materialy Mezhdunarodnogo Simposiuma Stran-Chlenov SEV, 10th. [Mechanism of Action of Herbicides and Synthetic Plant Growth Regulators and Their Ultimate Fate in Biosphere, Materials of the International Symposium of Member Countries of the Council for Mutual Economic Assistance, 10th.]; Apr 15-17, 1975, Pushchino, USSR. Vol. 2. Edited by M.S. Sokolov. Pushchino, USSR: Akademiya Nauk SSSR, Nauchnyi Tsentr Biologicheskikh Issledovanii.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005003578	Vrochinskiy, K.K. (1973) Izuchenie migratsionnykh svoistv pestitsidov v modelnoi vodnoi ekosisteme. [Migratory properties of pesticides in a model water ecosystem.] Rybnoe Khozyaistvo (Moscow). [Fisheries.] 11:29-31.
005000932	Wall, W.J., Jr.; Marganian, V.M. (1973) Control of salt marsh Culicoides and Tabanus larvae in small plots with granular organophosphorus pesticides, and the direct effect on other fauna. Mosquito News 33(1):88-93.
000001229	Wang, G.T.; Colavita, J.H.; Phillips, A.A. (1973) Temephos: Safety of Temephos When Applied at Four Times the Recommended Level Once Weekly for Three Weeks: Project No.30-0960-3-755. (Unpublished study received Mar 6, 1974 under 241-234; submitted by American Cyanamid Co., Princeton, N.J.; CDL:026901-T)
000001231	Wang, G.T.; Colavita, J.H.; Phillips, A.A. (1974) Temephos: Safety of 2% Temephos Powder When Applied Topically on Cats at 80 mg Temephos/kg. Body Weight Once Weekly for Three Weeks: Project No. 30-0960-3-755. (Unpublished study received Mar 6, 1974 under 241-234; submitted by American Cyanamid Co., Princeton, N.J.; CDL:026901-V)
000001230	Wang, G.T.; Colavita, J.H.; Phillips, A.A. (1974) Temephos: Safety of 2% Temephos Powder When Applied Topically on Dogs at 20 mg Temephos/kg. Body Weight Five Days a Week for Three Weeks: Project No. 30-0960-3-755. (Unpublished study received Mar 6, 1974 under 241-234; submitted by American Cyanamid Co., Princeton, N.J.; CDL:026901-U)
005000804	Ward, D.V.; Howes, B.L. (1974) The effects of Abate, an organophosphorous insecticide, on marsh fiddler crab populations. Bulletin of Environmental Contamination and Toxicology 12(6):694-697.
005004596	Ward, D.V.; Howes, B.L.; Ludwig, D.F. (1976) Interactive effects of predation pressure and insecticide (temephos) toxicity on populations of the marsh fiddler crab Uca pugnax. Marine Biology 35:119-126.
005002138	Wattal, B.L.; Bhatnagar, V.N.; Joshi, G.C. (1975) Efficacy of different larvicides against Anopheles and Culicine mosquitoes in weekly application schedules. Journal of Communicable Diseases 7(4):358-363.
005000860	Watts, R.R.; Storherr, R.W. (1969) Gas chromatography of organophosphorus pesticides: retention times and response data on three columns. Journal of the Association of Official Analytical Chemists 52(3):513-521.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000989	Weaving, A.J.S. (1974) Preliminary screening tests of insecticides for contact residual effectiveness against the tobacco beetle Lasioderma serricorne (F.). Rhodesian Journal of Agricultural Research 12(2):177-193.
005000835	Webb, A.B. (1976) Residue evaluations of Abate briquettes for potential use as mosquito larvicides including related studies. Dissertation Abstracts International B 37(4):1544.
005020216	Webb, A.B. (1976) Residue Evaluations of Abate Briquettes for Potential Use as Mosquito Larvicides Including Related Studies (Available from: University Microfilms, Ann Arbor, MI; 76-22,892)
005000776	Weeks, M.H.; Boldt, R.E.; Steinberg, M. (1968) Hazards of Insecticides Formulated into Plastic Pellets O,O,O',O'-Tetramethyl, O,O'-Thiodi-p-Phenylene Phosphorothioate(Technical ENT-27165). Edgewood Arsenal, Md.: U.S. Army Environmental Hygiene Agency. (USAEHA project no. 33-20-68/69; available from: NTIS, Springfield, Va.: AD-846 601)
000001248	Weis, M.E. (1964) Monthly Report, May, 1964: E.I. 52,160 [for Mosquito Larvae Control]. (Unpublished study received May 27, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-AD)
000008944	Wells, A.; Cress, D. (1973) Asparagus Cutworm Research 1971-73. (Unpublished study received Jul 10, 1975 under 6F1654; prepared by Michigan State Univ., Dept. of Entomology, submitted by E.I. du Pont de Nemours & Co., Inc., Wilmington, Del.; CDL:094852-J)
000008466	Wells, A.; Grafius, E.; Carolus, B. (1970) Imported and Looper Control on Cabbage--1970: Table 1. (Unpublished study received Dec 8, 1976 under 352-362; submitted by E.I. du Pont de Nemours & Co., Wilmington, Del.; CDL:227248-E)
000008326	Wells, A.; Grafius, E.; Carolus, R. (1970) Snap Bean Insect Control 1970: Table 6. (Unpublished study including letter dated Jan 21, 1971 from W.D. Marks to Roy B. Boren, received Jun 29, 1976 under 201-347; prepared by Michigan State Univ., Dept. of Horticulture, submitted by Shell Chemical Co., Washington, D.C.; CDL:232406-T)
000008723	Wells, A.; Grafius, E; Carolus, R. (1970) Green Peach Aphid and Corn Borer Control on Peppers--1970: Table 2. (Unpublished study received Apr 13, 1971 under 352-342; submitted by E.I. du Pont de Nemours & Co., Wilmington, Del.; CDL:003023-C)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000722	Wenk, P.; Schulz-Key, H. (1974) Versuche zur quantitativen Wirkungsbestimmung von Larviziden gegen Simuliiden in Westafrica. [Experiments for qualitative evaluation of larvicides used for simulium control in West Africa.] Tropenmedizin und Parasitologie 25(4):381-394.
005003271	West, A.S. (1973) Recent Canadian developments in area chemical control of biting flies. Pages 15-18, In Biting Fly Control and Environmental Quality: Proceedings of a Symposium; May 16-18, 1972, University of Alberta, Edmonton, Alberta, Canada. Edited by A. Hudson. Edmonton, Alberta, Canada: University of Alberta. (Defence Research Board report DR 217)
000001243	Whalen, R.; Orloski, E.J. (1965) Residues of CL 52,160 in Oysters: Report No. C-72. (Unpublished study received May 27, 1965 under 241-132; submitted by American Cyanamid Co., Princeton, N.J.; CDL:001791-V)
005000867	Whitlaw, J.T., Jr.; Evans, E.S., Jr. (1968) Selected plastic formulations for use as mosquito larvicides. Journal of Economic Entomology 61(4):889-892.
005000878	Whitney, W.K.; Kenaga, E.E.; Hardy, J.L.; Doty, A.E. (1969) Rapid knockdown activity of Dowco 217, a new insecticide. Journal of Economic Entomology 62(3):567-568.
005011359	WHO Expert Committee on Insecticides (1973) Specifications for Pesticides Used in Public Health: Insecticides--Rodenticides--Molluscicides--Repellents--Methods. 4th ed. Geneva, Switzerland: World Health Organization.
005019466	WHO Expert Committee on Vector Biology and Control (1978) Chemistry and Specifications of Pesticides. Second Report of the WHO Expert Committee on Vector Biology and Control. Geneva, Switzerland: World Health Organization. (WHO technical report series no. 620)
005004227	WHO Scientific Group (1975) Chemical and Biochemical Methodology for the Assessment of Hazards of Pesticides for Man. Geneva, Switzerland: World Health Organization. (WHO technical report series no. 560)
005000665	Wicht, M.C., Jr.; Rodriguez, J.G. (1970) Integrated control of muscid flies in poultry houses using predator mites, selected pesticides and microbial agents. Journal of Medical Entomology 7(6):687-692.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000011924	Wilde, G. (1965) Potato Insecticide Trials. (Unpublished study received Nov 10, 1972 under 3F1323; prepared by Cornell Univ., Dept. of Entomology, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:093552-AF)
000012450	Wilde, G.; Semel, M. (1965) Control of Potato Insect Pests on Long Island, New York. (Unpublished study received Nov 10, 1972 under 3F1323; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:093552-AE)
005000381	Wilkinson, R.N.; Barnes, W.W.; Gillogly, A.R.; Minnemeyer, C.D. (1971) Field evaluation of slow-release mosquito larvicides. Journal of Economic Entomology 64(1):1-3.
000013568	Wilson, A.J., Jr.; Butler, P.A.; Lowe, J.I. (1964) Quarterly Project Progress Report: Effects of Pesticides. (Unpublished study received Jun 2, 1971 under 1F1062; prepared by U.S. Fish and Wildlife Service, Bureau of Commercial Fisheries, Biological Laboratory, submitted by Shell Chemical Co., Washington, D.C.; CDL:091826-P)
005000937	Wilton, D.P.; Fetzer, L.E., Jr.; Fay, R.W. (1973) Insecticide baits for Anopheline larvae. Mosquito News 33(2):198-203.
005000927	Windeguth, D.L. von; Eliason, D.A.; Schoof, H.F. (1971) Efficacy of carbaryl, propoxur, Abate, and methoxychlor as larvicides against field infestations of Aedes aegypti. Mosquito News 31(1):91-95.
005000712	Windeguth, D.L. von; Patterson, R.S. (1966) The effects of two organic phosphate insecticides on segments of the aquatic biota. Mosquito News 26(3):377-380.
005007868	Winner, R.A.; Steelman, C.D. (1978) Effects of selected insecticides on Romanomermis culicivorax, a mermithid nematode parasite of mosquito larvae. Mosquito News 38(4):546-553.
005009162	Winner, R.A.; Steelman, C.D.; Schilling, P.E. (1978) Effects of selected insecticides on Romanomermis culicivorax, a mermithid nematode parasite of mosquito larvae. Mosquito News 38(4):546-553.
005005436	Womeldorf, D.J. (1976) Role of the cooperative agreement between the California Department of Health and local vector control agencies in coordinating pesticide use. Pages 13-15, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association, Vol. 44. Visalia, Calif.: California Mosquito Control Association.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000820	Womeldorf, D.J.; Atkins, E.L.; Gillies, P.A. (1974) Honey bee hazards associated with some mosquito abatement aerial spray applications. California Vector Views 21(9):51-55.
005000748	Womeldorf, D.J.; Gillies, P.A.; White, K.E. (1968) Present status of insecticide resistance in Californian mosquito larvae. Pages 81-84, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association. Vol. 36. Visalia, Calif.: CMCA.
005000663	World Health Organization Expert Committee (1976) Epidemiology of Onchorceriasis. Geneva, Switzerland: World Health Organization. (WHO technical report series no.597)
005001007	World Health Organization Scientific Group (1975) Chemical and Biochemical Methodology for the Assessment of Hazards of Pesticides for Man: Report of a WHO Scientific Group. Geneva, Switzerland: World Health Organization. (WHO technical report series no.560)
005001006	World Health Organization. Expert Committee on Insecticides (1971) Chemistry and Specifications of Pesticides: Nineteenth Report of the WHO Expert Committee on Insecticides. Geneva, Switzerland: World Health Organization. (WHO technical report series no. 475)
005001028	World Health Organization, Expert Committee on Vector Biology and Control (1978) Chemistry and Specifications of Pesticides: Second Report of the WHO Expert Committee on Vector Biology and Control. Geneva, Switzerland: World Health Organization. (WHO technical report series no. 620)
005000772	World Health Organization, Division of Malaria and Other Parasitic Diseases (1973) Manual on Larval Control Operations in Malaria Programmes. Geneva, Switzerland: World Health Organization. (WHO offset publication no. 1)
005000325	World Health Organization, Expert Committee on Insecticides (1973) Specifications for Pesticides Used In Public Health. 4th ed. Geneva, Switzerland: WHO.
005001032	World Health Organization, Expert Committee on Malaria (1974) WHO Expert Committee on Malaria: Sixteenth Report. Geneva, Switzerland: World Health Organization. (WHO technical report series no. 549)
005000855	Wright, F.C.; Gilbert, B.N.; Riner, J.C. (1967) Gas chromatographic determination of Abate residues in water. Journal of Agricultural and Food Chemistry 15(6):1038-1039.

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
005000971	Yap, H.H.; Hanapi, S. (1976) Laboratory insecticide susceptibility tests against Mansonia larvae. Southeast Asian Journal of Tropical Medicine and Public Health 7(4):575-580.
005006245	Yasutomi, K. (1974) [Insecticide resistance in Culex pipiens pallens larval of Amagasaski City.] Bochu-Kagaku.
005005684	Zaini, M.A. (1969) Toxicity of various insecticides. Bulletin of Endemic Diseases 11(1/4):86-97,99.101,103.
005000872	Zeid, M.; Saad, A.A.; Ayad, A.M.; Tantawi, G.; Eldefrawi, M.E. (1968) Laboratory and field evaluation of insecticides against the Egyptian cotton leafworm. Journal of Economic Entomology 61(5):1183-1186.
000001321	Zimmerman, J.H. (1969) Toxicity of Paris Green, Methoxychlor and New Organophosphate Insecticides to Salt Marsh Killifish and Crustaceans: [Abate]. (Unpublished study received Nov 26, 1971 under 241-132; prepared by Univ. of Delaware, submitted by American Cyanamid Co., Princeton, N.J.; CDL:129448-O)
005000809	Ziv, M.; Brown, N.J.; Brown, A.W.A. (1969) Resistance potentialities of Aedes aegypti and Culex pipiens fatigans to organophosphorus and other insecticides. Bulletin of the World Health Organization 41(6):941-946.
000011154	Zoecon Corporation (1972?) 8-Day Subacute Toxicity Study--Bobwhite Quail--Altosid Technical--LC_{50} >10,000 PPM. (Unpublished study received Oct 26, 1977 under 20954-1; prepared in cooperation with Hazleton Laboratories, Inc.; CDL:232090-R)
000010674	Zoecon Corporation (1972*) 8-Day Subacute Toxicity Study--Bobwhite Quail--Altosid Technical--LC_{50} Greater than 10,000 PPM (Highest Dose Tested). (Unpublished study received Oct 10, 1974 under 20954-2; CDL:028329-R)
000011153	Zoecon Corporation (1972*) 8-Day Subacute Toxicity Study--Mallard Duck--Altosid Technical--LC_{50} >10,000 PPM. (Unpublished study received Oct 26, 1977 under 20954-1; prepared in cooperation with Hazleton Laboratories, Inc.; CDL.232090-Q)
000010673	Zoecon Corporation (1972*) 8-Day Subacute Toxicity Study--Mallard Duck--Altosid Technical--LC_{50} Greater than 10,000 PPM (Highest Dose Tested). (Unpublished study received Oct 10, 1974 under 20954-2; CDL:028329-Q)

Citations Examined and Judged to be Inappropriate For use in Developing the Standard

MRID	Citation
000010672	Zoecon Corporation (1973*) Acute Oral Toxicity Study--Mallard Duck--Altosid Technical--LD_{50} Greater than 2,000 Mg/Kg (Highest Dose Tested). (Unpublished study received Oct 10, 1974 under 20954-2; CDL.028329-P)
000011152	Zoecon Corporation (1973*) Acute Oral Toxicity Study--Mallard Ducks--Altosid Technical--LD_{50} >2,000 Mg/Kg. (Unpublished study received Oct 26, 1977 under 20954-1; prepared in cooperation with U.S. Bureau of Sport Fisheries and Wildlife; CDL: 232090-P)
000010401	Zoecon Corporation (1973*) Wildfowl Toxicity Studies. (Unpublished study received Jan 22, 1976 under 20954-EX-5; prepared in cooperation with U.S. Fish and Wildlife Service, Denver Wildlife Research Center and Hazleton Laboratories, Inc.; CDL.225689-F)
000010631	Zoecon Corporation (1976) Wildfowl Toxicity Studies. Summary of studies 231486-D through 231486-K. (Unpublished study received Feb 8, 1977 under 20954-1; CDL:231486-C)
005000659	Zurflueh, R.C. (1976) Phenylethers as insect growth regulators: laboratory and field experiments. Pages 61-74, In The Juvenile Hormones. Edited by L.I. Gilbert. New York: Plenum Press.
000001259	3M Company (1970?) [Chemical and Physical Properties of 3MCAP-1A]. (Unpublished study received Oct 16, 1970 under unknown admin. no.; 127730-A)
000001257	3M Company (1970?) [Efficacy Data on Field Tests of 3MCAP-1A]. (Unpublished study received Oct 16, 1970 under uknown admin. no.; CDL:127728-C)
000001258	3M Company (1970?) [Mechanism of Release: 3MCAP-1A]. (Unpublished study received Oct 16, 1970 under unknown admin. no.; CDL:127728-D)
000001256	3M Company (1970?) Toxicity Data of Abate--Reference 1. (Unpublished study received Oct 16, 1970 under unknown admin. no.; CDL: 27728-B)

Temephos Bibliography

Section 3

Standard Reference Material

The Federal Insecticide, Fungicide, and Rodenticide Act, as ameneded in 1978, 7tn U.S. Code, Chapter 135, 61 Statute 163.78 Statute 190.

U.S. Environmental Protection Agency (1978); Proposed Guidelines for Registering Pesticides in the United States. Federal Register, 43 (132) 29696.

U.S. Environmental Protection Agency (1978); Proposed Guidelines for Registering Pesticides in the United States; Hazard Evaluation Humans and Domestic Animals. Federal Register, 43 (163) 37336.

U.S. Environmental Protection Agency (1977); Regulations for the Enforcement of the Federal Insecticide, Fungicide, and Rodenticide Act, Title 40, Chapter 1, Part 162.

U.S. Environmental Protection Agency (in press) Proposed Guidelines for Registering Pesticides in the United States. Subparts G (Product Performance) and H (Label Development).

U.S. Environemental Protection Agency (1980); Proposed Guidelines for Registering Pesticides in the United States. Hazard Evaluation of Use of Pesticides for Nontarget Plants and Microorganisms. Federal Register, 45 (214) 72948.

U.S. Environmental Protection Agency (1975); Registration Reregistration ad Classification Procedures. Federal Register, 40 (129) 28242.

*U S GOVERNMENT PRINTING OFFICE: 1981 341-085/4469