

Air Quality Criteria for Ozone and Related Photochemical Oxidants (First External Review Draft)

Volume III of III

Air Quality Criteria for Ozone and Related Photochemical Oxidants

Volume III

National Center for Environmental Assessment-RTP Office
Office of Research and Development
U.S. Environmental Protection Agency
Research Triangle Park, NC

DISCLAIMER

This document is an external review draft for review purposes only and does not constitute U.S. Environmental Protection Agency policy. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

PREFACE

National Ambient Air Quality Standards (NAAQS) are promulgated by the United States Environmental Protection Agency (EPA) to meet requirements set forth in Sections 108 and 109 of the U.S. Clean Air Act (CAA). Sections 108 and 109 require the EPA Administrator (1) to list widespread air pollutants that reasonably may be expected to endanger public health or welfare; (2) to issue air quality criteria for them that assess the latest available scientific information on nature and effects of ambient exposure to them; (3) to set “primary” NAAQS to protect human health with adequate margin of safety and to set “secondary” NAAQS to protect against welfare effects (e.g., effects on vegetation, ecosystems, visibility, climate, manmade materials, etc); and (5) to periodically review and revise, as appropriate, the criteria and NAAQS for a given listed pollutant or class of pollutants.

In 1971, the U.S. Environmental Protection Agency (EPA) promulgated National Ambient Air Quality Standards (NAAQS) to protect the public health and welfare from adverse effects of photochemical oxidants. The EPA promulgates the NAAQS on the basis of scientific information contained in air quality criteria issued under Section 108 of the Clean Air Act. Following the review of criteria as contained in the EPA document, Air Quality Criteria for Ozone and other Photochemical Oxidants published in 1978, the chemical designation of the standards was changed from photochemical oxidants to ozone (O₃) in 1979 and a 1-hour O₃ NAAQS was set. The 1978 document focused primarily on the scientific air quality criteria for O₃ and, to a lesser extent, on those for other photochemical oxidants such as hydrogen peroxide and the peroxyacyl nitrates, as have subsequent revised versions of the ozone document.

To meet Clean Air Act requirements noted above for periodic review of criteria and NAAQS, the O₃ criteria document, *Air Quality Criteria for Ozone and Other Photochemical*

Oxidants, was next revised and then released in August 1986; and a supplement, *Summary of Selected New Information on Effects of Ozone on Health and Vegetation*, was issued in January 1992. These documents were the basis for a March 1993 decision by EPA that revision of the existing 1-h NAAQS for O₃ was not appropriate at that time. That decision, however, did not take into account some of the newer scientific data that became available after completion of the 1986 criteria document. Such literature was assessed in the next periodic revision of the O₃ air quality criteria document, which was completed in 1996 and provided scientific bases supporting the setting by EPA in 1997 of an 8-h O₃ NAAQS that is currently in force together with the 1-h O₃ standard.

The purpose of this revised air quality criteria document for O₃ and related photochemical oxidants is to critically evaluate and assess the latest scientific information published since that assessed in the above 1996 Ozone Air Quality Criteria Document (O₃ AQCD), with the main focus being on pertinent new information useful in evaluating health and environmental effects data associated with ambient air O₃ exposures. However, some other scientific data are also presented and evaluated in order to provide a better understanding of the nature, sources, distribution, measurement, and concentrations of O₃ and related photochemical oxidants and their precursors in the environment. The document assesses pertinent literature available through 2004.

The present draft document (dated January 2005) is being released for public comment and review by the Clean Air Scientific Advisory Committee (CASAC) to obtain comments on the organization and structure of the document, the issues addressed, the approaches employed in assessing and interpreting the newly available information on O₃ exposures and effects, and the key findings and conclusions arrived at as a consequence of this assessment. Public comments and recommendations will be taken into account making any appropriate further revisions to this document for incorporation into a Second External Review Draft. That draft will be released for further public comment and CASAC review before last revisions are made in response and incorporated into a final version to be completed by early 2006. Evaluations contained in the present document will be drawn on to provide inputs to associated PM Staff Paper analyses prepared by EPA's Office of Air Quality Planning and Standards (OAQPS) to pose options for consideration by the EPA Administrator with regard to proposal and, ultimately, promulgation of decisions on potential retention or revision, as appropriate, of the current O₃ NAAQS.

Preparation of this document was coordinated by staff of EPA's National Center for Environmental Assessment in Research Triangle Park (NCEA-RTP). NCEA-RTP scientific staff, together with experts from other EPA/ORD laboratories and academia, contributed to writing of document chapters. Earlier drafts of document materials were reviewed by non-EPA experts in peer consultation workshops held by EPA. The document describes the nature, sources, distribution, measurement, and concentrations of O₃ in outdoor (ambient) and indoor environments. It also evaluates the latest data on human exposures to ambient O₃ and consequent health effects in exposed human populations, to support decision making regarding the primary, health-related O₃ NAAQS. The document also evaluates ambient O₃ environmental effects on vegetation and ecosystems, man-made materials, and surface level solar UV radiation flux and global climate change, to support decision making on secondary O₃ NAAQS.

NCEA acknowledges the valuable contributions provided by authors, contributors, and reviewers and the diligence of its staff and contractors in the preparation of this draft document.

**Air Quality Criteria for Ozone and Related
Photochemical Oxidants
(First External Review Draft)**

VOLUME I

Executive Summary	E-1
1. INTRODUCTION	1-1
2. PHYSICS AND CHEMISTRY OF OZONE IN THE ATMOSPHERE	2-1
CHAPTER 2 ANNEX (ATMOSPHERIC PHYSICS/CHEMISTRY)	AX2-1
3. ENVIRONMENTAL CONCENTRATIONS, PATTERNS, AND EXPOSURE ESTIMATES	3-1
CHAPTER 3 ANNEX (AIR QUALITY AND EXPOSURE)	AX3-1

**Air Quality Criteria for Ozone and Related
Photochemical Oxidants
(First External Review Draft)**
(cont'd)

VOLUME II

4.	DOSIMETRY, SPECIES HOMOLOGY, SENSITIVITY, AND ANIMAL-TO-HUMAN EXTRAPOLATION	4-1
	CHAPTER 4 ANNEX (DOSIMETRY)	AX4-1
5.	TOXICOLOGICAL EFFECTS OF OZONE AND RELATED PHOTOCHEMICAL OXIDANTS IN LABORATORY ANIMALS AND IN VITRO TEST SYSTEMS	5-1
	CHAPTER 5 ANNEX (ANIMAL TOXICOLOGY)	AX5-1
6.	CONTROLLED HUMAN EXPOSURE STUDIES OF OZONE AND RELATED PHOTOCHEMICAL OXIDANTS	6-1
	CHAPTER 6 ANNEX (CONTROLLED HUMAN EXPOSURE)	AX6-1
7.	EPIDEMIOLOGICAL STUDIES OF HUMAN HEALTH EFFECTS ASSOCIATED WITH AMBIENT OZONE EXPOSURE	7-1
	CHAPTER 7 ANNEX (EPIDEMIOLOGY)	AX7-1
8.	INTEGRATIVE SYNTHESIS	8-1

**Air Quality Criteria for Ozone and Related
Photochemical Oxidants
(First External Review Draft)**
(cont'd)

VOLUME III

9.	ENVIRONMENTAL EFFECTS: OZONE EFFECTS ON VEGETATION AND ECOSYSTEMS	9-1
10.	TROPOSPHERIC OZONE EFFECTS ON UV-B FLUX AND CLIMATE CHANGE PROCESSES	10-1
11.	EFFECT OF OZONE ON MAN-MADE MATERIALS	11-1

Table of Contents

	<u>Page</u>
List of Tables	I-xiv
List of Figures	I-xvi
Authors, Contributors, and Reviewers	I-xviii
U.S. Environmental Protection Agency Project Team for Development of Air Quality Criteria for Ozone and Related Photochemical Oxidants	I-xix
U.S. Environmental Protection Agency Science Advisory Board (SAB) Staff Office Clean Air Scientific Advisory Committee (CASAC) Ozone Review Panel	I-xx
Abbreviations and Acronyms	I-xxi
 9. ENVIRONMENTAL EFFECTS: OZONE EFFECTS ON VEGETATION AND ECOSYSTEMS	 9-1
9.1 INTRODUCTION	9-1
9.2 METHODOLOGIES USED IN VEGETATION RESEARCH	9-2
9.2.1 Introduction	9-2
9.2.2 Methods Involving Experimental Exposures to O ₃	9-3
9.2.2.1 “Indoor”, Controlled Environment, and Greenhouse Chambers	9-3
9.2.2.2 Field Chambers	9-5
9.2.2.3 Plume Systems	9-7
9.2.2.4 Comparative Studies	9-10
9.2.2.5 Ozone Generation Systems	9-11
9.2.2.6 Experimental Exposure Protocols	9-11
9.2.3 Methods Involving Exposures to O ₃ in Ambient Air	9-12
9.2.3.1 Air-exclusion Systems	9-13
9.2.3.2 Natural Gradients	9-13
9.2.3.3 Use of Chemical Protectants	9-14
9.2.3.4 Biomonitoring	9-16
9.2.3.5 Calibrated Passive Monitors	9-24
9.2.4 Numerical/Statistical Methodologies	9-25
9.2.5 Improved Methods for Defining Exposure	9-27
9.3 SPECIES RESPONSE/MODE-OF-ACTION	9-28
9.3.1 Introduction	9-28
9.3.2 Mechanisms of Ozone-Induced Plant Alterations	9-31
9.3.2.1 Changes in Metabolic Processes	9-32
9.3.2.2 Modifications of Plant Physiological Processes	9-34
9.3.3 Ozone Uptake by Leaves	9-35
9.3.3.1 Possible Reactions Within the Leaf	9-43
9.3.3.2 Toxicants Within the Wall Space	9-46
9.3.3.3 Products of Ozone	9-48
9.3.3.4 Antioxidants Within the Apoplastic Space	9-55
9.3.4 Wounding and Pathogen Attack	9-64
9.3.4.1 Peroxidases	9-67
9.3.4.2 Jasmonic Acid and Salicylic Acid	9-69
9.3.4.3 Stress-Induced Alterations in Gene Expression	9-69

Table of Contents

(cont'd)

	<u>Page</u>
9.3.5	Primary Assimilation by Photosynthesis 9-73
9.3.5.1	Photooxidation: Light Reactions 9-73
9.3.6	Alteration of Rubisco by Ozone: Dark Reactions 9-74
9.3.7	Carbohydrate Transformations and Allocation 9-77
9.3.7.1	Lipid Synthesis 9-79
9.3.8	Role of Age and Size Influencing Response to Ozone 9-82
9.3.9	Summary 9-84
9.4	MODIFICATION OF FUNCTIONAL AND GROWTH RESPONSES 9-85
9.4.1	Introduction 9-85
9.4.2	Genetics 9-88
9.4.2.1	Genetic Basis of O ₃ Sensitivity 9-90
9.4.3	Environmental Biological Factors 9-91
9.4.3.1	Oxidant-Plant-Insect Interactions 9-92
9.4.3.2	Oxidant-Plant-Pathogen Interactions 9-95
9.4.3.3	Oxidant-Plant-Symbiont Interactions 9-100
9.4.3.4	Oxidant-Plant-Plant Interactions: Competition 9-101
9.4.4	Physical Factors 9-104
9.4.4.1	Light 9-104
9.4.4.2	Temperature 9-108
9.4.4.3	Humidity and Surface Wetness 9-111
9.4.4.4	Drought and Salinity 9-113
9.4.5	Nutritional Factors 9-116
9.4.6	Interactions with Other Pollutants 9-118
9.4.6.1	Oxidant Mixtures 9-119
9.4.6.2	Sulfur Dioxide 9-119
9.4.6.3	Nitrogen Oxides, Nitric Acid Vapor, and Ammonia 9-120
9.4.6.4	Hydrogen Fluoride and Other Gaseous Pollutants 9-123
9.4.6.5	Acid Deposition 9-123
9.4.6.6	Heavy Metals 9-125
9.4.6.7	Mixtures of Ozone with Two or More Pollutants 9-125
9.4.7	Interactions with Agricultural Chemicals 9-125
9.4.8	Factors Associated with Global Climate Change 9-126
9.4.8.1	Ozone-Carbon Dioxide-Temperature Interactions 9-128
9.4.8.2	Ozone-UV-B Interactions 9-151
9.4.8.3	Interactions of Ozone with Multiple Climate Change Factors 9-153
9.4.9	Summary - Environmental Factors 9-153
9.5	EFFECTS-BASED AIR QUALITY EXPOSURE- AND DOSE-RESPONSE INDICES 9-157
9.5.1	Introduction 9-157
9.5.2	Summary of Conclusions from the Previous Criteria Document 9-158
9.5.3	Use of Exposure Indices to Establish Exposure-Response Relationships 9-161

Table of Contents

(cont'd)

	<u>Page</u>
9.5.4	Identifying Exposure Components That Relate to Vegetation Effects 9-166
9.5.4.1	Role of Concentration 9-166
9.5.4.2	Role of Duration 9-169
9.5.4.3	Patterns of Exposure 9-169
9.5.4.4	Frequency of Occurrence of Peak Concentrations 9-172
9.5.4.5	Canopy Structure 9-173
9.5.4.6	Site and Climate Factors 9-174
9.5.4.7	Plant Defense Mechanism - Detoxification 9-174
9.5.5	Ozone Uptake or Effective Dose as an Index 9-174
9.5.5.1	Models of Stomatal Conductance 9-175
9.5.5.2	Nonlinear Response and Developing Flux Indices 9-178
9.5.5.3	Simulation Models 9-179
9.5.6	Summary 9-179
9.6	OZONE EXPOSURE-PLANT RESPONSE RELATIONSHIPS 9-181
9.6.1	Introduction 9-181
9.6.2	Summary of Key Findings/Conclusions from Previous Criteria Documents 9-182
9.6.3	Ozone Indices and Ambient Exposure 9-193
9.6.4	Effects of Ozone on Annual and Biennial Species 9-212
9.6.4.1	Effects on Growth, Biomass, and Yield of Individual Species 9-212
9.6.4.2	Effects on Plant Quality 9-217
9.6.4.3	Effects on Foliar Symptoms 9-219
9.6.4.4	Other Effects 9-220
9.6.4.5	Scaling Experimental Data to Field Conditions 9-221
9.6.4.6	European Critical Levels 9-224
9.6.4.7	Summary of Effects on Short-Lived Species 9-227
9.6.5	Effects of Ozone on Long-Lived (Perennial) Species 9-229
9.6.5.1	Herbaceous Perennial Species 9-229
9.6.5.2	Deciduous Woody Species 9-234
9.6.5.3	European Critical Levels 9-240
9.6.5.4	Summary of Effects on Deciduous Woody Species 9-240
9.6.5.5	Evergreen Woody Species 9-241
9.6.5.6	Summary of Effects on Evergreen Woody Species 9-244
9.6.5.7	Scaling Experimental Data to Mature Trees 9-245
9.6.6	Studies with the Chemical EDU 9-249
9.6.7	Summary 9-251
9.7	EFFECTS OF OZONE EXPOSURE ON NATURAL ECOSYSTEMS 9-254
9.7.1	Introduction 9-254
9.7.2	Case Studies 9-256
9.7.2.1	Valley of Mexico 9-256
9.7.2.2	San Bernardino Mountains 9-263
9.7.2.3	Sierra Nevada Mountains 9-268

Table of Contents

(cont'd)

	<u>Page</u>
9.7.2.4	Appalachian Mountains 9-270
9.7.2.5	Plantago Studies in the United Kingdom 9-271
9.7.2.6	Forest Health in the Carpathian Mountains 9-271
9.7.2.7	Field Exposure System (FACE), Rhinelander, Wisconsin 9-273
9.7.3	Landscape Condition 9-275
9.7.4	Biotic Condition 9-278
9.7.4.1	Ecosystems and Communities 9-278
9.7.4.2	Species and Populations 9-284
9.7.4.3	Organism Condition 9-287
9.7.5	Ecosystem, Chemical, and Physical Characteristics (water, soil) 9-295
9.7.5.1	Nutrient Concentrations, Trace Inorganic and Organic Chemicals 9-295
9.7.6	Ecological Processes 9-296
9.7.6.1	Energy Flow 9-296
9.7.6.2	Material Flow 9-298
9.7.7	Hydrological and Geomorphological 9-301
9.7.8	Natural Disturbance Regimes 9-301
9.7.9	Scaling to Ecosystem Levels 9-302
9.7.9.1	Scaling from Seedlings to Mature Trees 9-303
9.7.9.2	Surveys, Growth Correlations and Stand-Level Modeling 9-305
9.7.10	Summary of Ecological Effects of Ozone Exposure on Natural Ecosystems 9-312
9.8	ECONOMIC EVALUATION OF OZONE EFFECTS ON AGRICULTURE, FORESTRY AND NATURAL ECOSYSTEMS 9-315
9.8.1	Introduction 9-315
9.8.2	The Measurement of Economic Information 9-317
9.8.3	Understanding of Air Pollutants Effects on the Economic Valuation of Agriculture and Other Vegetation in the 1996 Criteria Document 9-318
9.8.3.1	Agriculture 9-319
9.8.3.2	Forests (Tree Species) and Natural Ecosystems 9-322
9.8.4	Studies Since 1996 of Ozone Exposure Effects on the Economic Value of Agriculture, Forests, and Ecosystems 9-323
9.8.5	Limitations of Scientific Studies and Economic Information 9-325
9.8.6	Conclusions 9-328
9.9	SUMMARY AND CONCLUSIONS FOR VEGETATION AND ECOSYSTEM EFFECTS 9-329
9.9.1	Introduction 9-329
9.9.2	Methodology 9-331
9.9.3	Mode-of-Action 9-332
9.9.4	Modification of Growth Response 9-334

Table of Contents

(cont'd)

	<u>Page</u>
9.9.5 Exposure Indices	9-337
9.9.6 Ozone Exposure-Plant Response Relationships	9-340
9.9.7 Ecosystem Effects	9-343
9.9.8 Economics	9-345
REFERENCES	9-347
 10. TROPOSPHERIC OZONE EFFECTS ON UV-B FLUX AND CLIMATE CHANGE PROCESSES	 10-1
10.1 INTRODUCTION	10-1
10.2 THE ROLE OF TROPOSPHERIC OZONE IN DETERMINING GROUND-LEVEL UV-B FLUX	 10-1
10.2.1 Factors Governing Ultraviolet Radiation Flux at the Earth's Surface	 10-2
10.2.1.1 UV Radiation: Wavelengths, Energies and Depth of Atmospheric Penetration	 10-2
10.2.1.2 Temporal Variations in Solar Flux	10-3
10.2.1.3 Atmospheric Radiative Interactions with Solar UV Radiation	 10-4
10.2.1.4 Modeling Surface UV-B Irradiance	10-10
10.2.2 Factors Governing Human Exposure to Ultraviolet Radiation	10-11
10.2.2.1 Outdoor Activities	10-12
10.2.2.2 Occupation	10-13
10.2.2.3 Age	10-13
10.2.2.4 Gender	10-14
10.2.2.5 Geography	10-15
10.2.2.6 Protective Behavior	10-15
10.2.2.7 Summary of Factors that Affect Human Exposures to Ultraviolet Radiation	 10-16
10.2.3 Factors Governing Human Health Effects due to Ultraviolet Radiation	 10-16
10.2.3.1 Erythema	10-17
10.2.3.2 Skin Cancer	10-19
10.2.3.3 Ultraviolet Radiation Exposure and the Incidence of Nonmelanoma Skin Cancers	 10-20
10.2.3.4 Ocular Effects of Ultraviolet Radiation Exposure	10-26
10.2.3.5 Ultraviolet Radiation and Immune System Suppression	 10-27
10.2.3.6 Protective Effects of Ultraviolet Radiation – Production of Vitamin D	 10-29
10.2.4 Summary and Conclusions for O ₃ Effects on UV-B Flux	10-30
10.3 TROPOSPHERIC OZONE AND CLIMATE CHANGE	10-31
10.3.1 The Projected Impacts of Global Climate Change	10-32
10.3.2 Solar Energy Transformation and the Components of the Earth's Climate System	 10-36

Table of Contents

(cont'd)

	<u>Page</u>
10.3.3 The Composition of the Atmosphere and the Earth's Radiative Equilibrium	10-37
10.3.3.1 Forcing of the Earth's Radiative Balance	10-39
10.3.4 Factors Affecting the Magnitude of Climate Forcing by Ozone	10-41
10.3.4.1 Global versus Regional Atmospheric Ozone Concentrations	10-41
10.3.4.2 Global Versus Regional Atmospheric Ozone Trends	10-42
10.3.4.3 The Sensitivity of Ozone-related Forcing Surface to Albedo	10-45
10.3.5 Estimated Forcing by Tropospheric Ozone	10-45
10.3.5.1 Direct Climate Forcing Due to Ozone	10-45
10.3.5.2 Indirect Forcing Due to Ozone	10-47
10.3.5.3 Predictions for Future Climate Forcing by Anthropogenic Ozone	10-48
10.3.6 Conclusion	10-48
REFERENCES	10-50
 11. EFFECT OF OZONE ON MAN-MADE MATERIALS	 11-1
11.11.1 Mechanisms of Ozone Damage and Exposure-Response	11-1
11.11.2 Textiles and Fabrics	11-3
11.11.3 Dyes, Pigments, and Inks	11-4
11.11.4 Artists' Pigments	11-5
11.11.5 Surface Coatings	11-8
REFERENCES	11-14

List of Tables

<u>Number</u>	<u>Page</u>
9-1	Advantages and Disadvantages of Protective Chemicals Used in Assessment of O ₃ Effects on Plants 9-17
9-2	Advantages and Disadvantages of Bioindicators Used to Study O ₃ Plant Effects 9-18
9-3	Advantages and Disadvantages of Cultivar Comparisons Used in Assessment of O ₃ Effects on Plants 9-22
9-4	Advantages and Disadvantages of Various Dendrochronological Techniques Used in Assessment of O ₃ Effects on Plants 9-24
9-5	Advantages and Disadvantages of Modeling Techniques Used in Assessment of O ₃ Effects on Plants 9-27
9-6	The Flow of Ozone into a Leaf and Possible Reactions 9-39
9-7	Some Rates of Reaction of Ozone With Critical Biochemicals 9-45
9-8	Superoxide Dismutase Isozymes and Isoforms 9-62
9-9	Gene Families and cDNA Clones Used as Probes for SAR (Ward et al., 1991) 9-66
9-10	Proteins Altered by Ozone as Measured by Molecular Biological Techniques as mRNA Level or Other Gene Activity Rather than Enzyme Activity 9-71
9-11	Interactions Involving O ₃ and Plant Pathogens 9-97
9-12	Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels 9-130
9-13	Summary of Ozone Exposure Indices Calculated for 3- or 5-Month Growing Seasons from 1982 to 1991 9-185
9-14	Ozone Exposure Levels (Using Various Indices) Estimated To Cause at Least 10% Crop Loss in 50 and 75% of Experimental Cases 9-187
9-15	SUM06 Levels Associated with 10 and 20% Total Biomass Loss for 50 and 75% of the Seedling Studies 9-190
9-16	Summary of Selected Studies of Ozone Effects on Annual Species 9-195
9-17	Summary of Selected Studies of the Effects of Ozone on Perennial Herbaceous Plants 9-199
9-18	Summary of Selected Studies of Ozone Effects on Deciduous Trees and Shrubs . . . 9-201

List of Tables
(cont'd)

<u>Number</u>		<u>Page</u>
9-19	Summary of Selected Studies of Ozone Effects on Evergreen Trees and Shrubs . . .	9-206
9-20	Ethylene Diurea Effects on Vegetation Responses to Ozone	9-210
9-21	Ozone Exposures at 35 Rural Sites in the Clean Air Status and Trends Network in the Central and Eastern United States From 1989 to 1995	9-228
9-22	Essential Ecological Attributes for Natural Ecosystems Affected by O ₃	9-257
9-23	Case Studies Demonstrating the Ecological Effects of O ₃	9-264
9-24	The Most Comprehensively Studied Effects of O ₃ on Natural Ecosystem is the San Bernardino Mountain Forest Ecosystem	9-266
9-25	Effects of Ozone, Ozone and N Deposition, and Ozone and Drought Stress on <i>Pinus ponderosa</i> and <i>Pinus jeffreyi</i> in the Sierra Nevada and the San Bernardino Mountains, California. Citations are Focused on Research Published Since U.S. EPA (1996)	9-267
9-26	Summary of Responses of <i>Populus tremuloides</i> to Elevated CO ₂ (+200 μmol mol ⁻¹), O ₃ (1.5 × ambient), or CO ₂ +O ₃ Compared with Control During Three Years of Treatments at the Aspen FACE Project (Modified from Karnosky et al. 2003a)	9-274
10-1	Examples of Impacts Resulting From Projected Changes in Extreme Climate Events	10-34
10-2	CTM Studies Assessed by the IPCC for its Estimate of the Change in Global, Total Column O ₃ Since the Pre-Industrial Era	10-44
10-3	Tropospheric O ₃ Change (ΔO ₃) in Dobson Units (DU) Since Pre-industrial Times, and the Accompanying net (SW plus LW) Radiative Forcings (Wm ⁻²), After Accounting for Stratospheric Temperature Adjustment (using the Fixed Dynamical Heating method)	10-46
11-1	Average 24-h Ozone Concentrations Producing the Highest Frequency of Cracks of a Certain Length in the Middle and Central Zones of the Rubber Test Strips	11-3
11-2	Cuprammonium Fluidity of Moist Cotton Cloth Exposed to 20 to 60 ppb Ozone	11-4
11-3	Color Change After 12 Weeks of Exposure to a Mixture of Photochemical Oxidants	11-11

List of Figures

<u>Number</u>		<u>Page</u>
9-1	Ozone uptake from the atmosphere	9-30
9-2	Absorption and transformation of O ₃ within the leaf	9-32
9-3	The uptake of O ₃ into the leaf. Each of the individual concentration layers of O ₃ represents a different process of movement and of plant/ microenvironmental interaction	9-36
9-4	The microarchitecture of a dicot leaf	9-37
9-5	The change in the O ₃ concentration inside a leaf with time	9-38
9-6	Possible transformations of O ₃ within a leaf	9-41
9-7	Possible reactions of O ₃ within water	9-41
9-8a,b	The Crigee mechanism of O ₃ attack of a double bond	9-44
9-9	Varied ESR radicals, trapped and not, generated by ozone under somewhat physiological conditions	9-47
9-10	Pathogen-Induced Hypersensitivity	9-50
9-11	The interaction of H ₂ O ₂ and Ca ²⁺ movements with ABA-induced stomatal closure	9-52
9-12	The reaction of ascorbate within the apoplasm of the cell wall and its ultimate reduction/oxidations	9-56
9-13	The pathway leading from phospholipids to jasmonic and traumatic acid	9-70
9-14	The production of Rubisco and its Calvin Cycle pathway reactions	9-75
9-15	Linkage of senescence with hypersensitivity reactions and first event of O ₃ attack of plants	9-82
9-16	Diagrammatic representation of several exposure indices, illustrating how they weight concentration and accumulate exposure	9-162
9-17	Trends in May-September 12-h SUM06, peak 1-h ozone concentration and number of daily exceedances of 95 ppb for Crestline in 1963-1999 in relation to trends in mean daily maximum temperature for Crestline and daily reactive organic gases (ROG) and oxides of nitrogen (NO _x) for San Bernardino county	9-168

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
9-18	Distribution of biomass loss predictions from Weibull and linear exposure-response models that relate biomass to O ₃ exposure	9-188
9-19	A conceptual diagram of processes and storage pools in sources and sinks that are affected by ozone exposure	9-276
9-20	Common anthropogenic stressors and the essential ecological attributes they affect	9-313
10-1	Extraterrestrial solar flux measured by the satellite UARS SOLSTICE instrument (dotted line)	10-3
10-2	Ozone column abundances from the years 1990 to 1992 for 0, 40, and 80° N as well as 80° S	10-6
10-3	The sensitivity of ground-level UV flux to a 1 DU change in total column O ₃ , under clear sky conditions, as a function of solar zenith angle (SZA)	10-10
10-4	Complexity of factors that determine human exposure to UV radiation	10-11
10-5	Estimated global mean radiative forcing exerted by gas and various particle phase species for the year 2000, relative to 1750	10-40
10-6	Mid-tropospheric O ₃ abundance (ppb) in northern midlatitudes (36 °N-59 °N) for the years 1970 to 1996	10-43
11-1	In-service fading of nylon 6 yarn inside house	11-6
11-2	In-service fading of nylon 6 yarn outside house	11-7
11-3	Observed color changes for natural colorant-on-paper systems during exposure to 0.40 ppm ozone at 25 °C ± 1 °C, 50% RH, in the absence of light	11-9
11-4	Observed color changes for natural colorant-on-site during exposure to 0.40 ppm ozone at 25 °C ± 1 °C, 50% RH, in the absence of light	11-10

Authors, Contributors, and Reviewers

To be inserted in Second External Review draft.

**U.S. ENVIRONMENTAL PROTECTION AGENCY
PROJECT TEAM FOR DEVELOPMENT OF AIR QUALITY CRITERIA
FOR OZONE**

To be inserted in Second External Review draft.

**U.S. ENVIRONMENTAL PROTECTION AGENCY
SCIENCE ADVISORY BOARD (SAB) STAFF OFFICE
CLEAN AIR SCIENTIFIC ADVISORY COMMITTEE (CASAC) OZONE
REVIEW PANEL***

To be inserted in Second External Review draft.

Abbreviations and Acronyms

AQCD	Air Quality Criteria Document
$C \times T$	Concentration times duration of exposure
CI	color index
CO ₂	carbon dioxide
COP	Conference of Parties
CTM	Chemistry Transport Model
DNA	Deoxyribonucleic acid
DNA	Deoxyribonucleic acid
EPA	U.S. Environmental Protection Agency
GSH	Glutathione
H ⁺	hydrogen ion
H ₂ O ₂	hydrogen peroxide
HFC	hydrofluorocarbon
HO	hydroxyl
HO ₂	hydroperoxyl; hydroperoxy
IPCC	Intergovernmental Panel on Climate Change
IR	infrared radiation
mRNA	Messenger ribonucleic acid
NHAPS	National Human Activity Pattern Survey
NMHC	nonmethane hydrocarbon
NO ₂	nitrogen dioxide
NO _x	nitrogen oxides
NRC	National Research Council
O(¹ D)	electronically excited oxygen atom
O ₃	ozone
OH	hydroxyl; hydroxy
PAN	Peroxyacetyl nitrate

RNA	Ribonucleic acid
RO ₂	organic peroxy; organic peroxy
SO ₂	sulfur dioxide
SOD	Superoxide dismutase
SUM00	Sum of all hourly average concentrations
SUM06	Seasonal sum of all hourly average concentrations ≥ 0.06 ppm
TAR	Third Assessment Report
UNEP	United Nations Environment Program
UV	Ultraviolet
UV-A	
UV-B	Ultraviolet radiation of wavelengths from 280 to 320 nm
UV-C	
W126	cumulative integrated exposure index with a sigmoidal weighting function
WMO	World Meteorological Organization
ZAPS	Zonal Air Pollution System

9. ENVIRONMENTAL EFFECTS: OZONE EFFECTS ON VEGETATION AND ECOSYSTEMS

9.1 INTRODUCTION

The preceding chapters of this document focused on discussion of: (a) background information regarding ozone (O₃)-related atmospheric chemistry, air quality and exposure aspects; and (b) dosimetric/health effects aspects, as well as the integrative synthesis of key information drawn from such chapters of most importance for EPA review of the primary O₃ NAAQS. On the other hand, this and the next two chapters assess available scientific information on O₃-related welfare effects of most pertinence for the review of secondary O₃ NAAQS. This includes discussion of three classes of environmental effects:

- O₃ effects on vegetation and ecosystem (Chapter 9);
- the role of tropospheric O₃ in climate change, including determining of ground-level UV-B flux (Chapter 10); and
- O₃ effects on manmade materials (Chapter 11).

As for the organizational structure of this chapter, after this opening Introduction, Section 9.2 highlights key features of methodologies used in vegetation research, followed by discussion of vegetation species response/mode-of-action aspects (Section 9.3) and factors that modify functional and growth responses (Section 9.4). Effects-based air quality exposure and dose-response indices are next discussed in Section 9.5, followed by assessment of O₃ exposure-plant response relationships (Section 9.6) and effects on natural ecosystems (Section 9.7). The economic evaluation of O₃ effects on agriculture, forestry, and natural ecosystems is then discussed in Section 9.8. Lastly, the chapter closes with Section 9.9, an overall summary of key findings and conclusions.

9.2 METHODOLOGIES USED IN VEGETATION RESEARCH

9.2.1 Introduction

The scale of investigations into the direct effects of O₃ on plants range from evaluation of sub-cellular to cellular, organismal, population, community, and ecosystem responses, each with its particular experimental methodologies and suite of specialized instrumentation, equipment, facilities, and experimental protocols. These investigations generate data; whereas other types of methodologies exist for the handling of data and statistical analysis as well as the utilization of data in developing the different exposure metrics or indices used to define exposure, quantitative exposure-response relationships, and computer simulation models of these. The objective of this section is not to provide an updated encyclopedia of all the methods that have been used, but rather to focus on approaches that have:

- (1) led to improved understanding of the quantitatively measurable growth and development responses of plants and plant communities to O₃, or
- (2) provided information about the extent and geographic distribution of the responses of herbaceous and woody plants, both cultivated and native, to ambient O₃ exposures.

The first objective is essential for determining dose-response functions used in the development of impact and risk assessments of the effects of O₃ and usually involves treating plants to a range of artificial O₃ exposures. The second objective is essential for determining the geographic distribution of the risk and usually involves plants subjected to ambient air O₃ exposures.

The types of methodologies used by biochemists, molecular biologists, or plant physiologists, whose interests lie in determining effects on specific constituents or in understanding the mode of action of O₃, are not discussed here. Methods used to characterize the O₃ content of ambient air and to define exposure and exposure-response relations are discussed in Sections 9.5 and 9.6, respectively.

The methodologies for exposure-response studies have used many different types of exposure facilities and protocols and have employed a range of statistical approaches to the analysis and interpretation of data. Most of the studies have been conducted using major agricultural crop species. The methodologies have improved over the years as a result of the development, availability, or application of new or improved instrumentation, physical systems, and numerical approaches to data analysis. Yet equally important to the roles played by these

advances has been the clearer understanding that has emerged from earlier work identifying the *type* of experimentation needed to achieve realistic assessments of the magnitude and extent of the impact of O₃ on vegetation of all types. As a result, significantly increased attention is now being paid to field observations of responses to ambient O₃ pollution, particularly to the responses of forest trees and native vegetation.

Other than in various exploratory studies that have used chamber-based steady-state exposure concentrations (so-called “square-wave” exposures), the trend in experimental exposure protocols has been to attempt to expose plants under conditions as natural as possible to temporal profiles that simulate the real-world, either by conducting experiments in the field or in elaborately controlled environment facilities that provide simulated field conditions.

Previous Air Quality Criteria Documents for Ozone and Other Photochemical Oxidants (U.S. Environmental Protection Agency, 1986; 1996) described the time-course for these methodological developments. Although this section provides a brief overview of the methodologies used in the past, and their limitations, it focuses mainly on those techniques that have come into prominence over the last decade. This has been aided considerably by several compilations of experimental methodologies and facilities, such as the earlier comprehensive review for the U.S. Environmental Protection Agency/National Acid Precipitation Assessment Program by Hogsett et al. (1987a; 1987b) and the more recent reviews by Manning and Krupa (1992), Musselman and Hale (1997) and Karnosky et al. (2001).

9.2.2 Methods Involving Experimental Exposures to O₃

9.2.2.1 “Indoor”, Controlled Environment, and Greenhouse Chambers

The earliest experimental investigations of the effects of O₃ on plants utilized simple glass or plastic-covered chambers, often located within greenhouses, into which a flow of O₃-enriched air or oxygen could be passed to provide the exposure. The types, shapes, styles, materials of construction, and locations of these chambers were as numerous as the different investigators and, in spite of providing little resemblance to real-world conditions, they yielded much of the basic information on the visible and physiological effects on plants. The construction and performance of more elaborate and better instrumented chambers dating back to the 1960s has been well-summarized in Hogsett et al. (1987a), including those installed in greenhouses (with or without some control of temperature and light intensity).

1 One greenhouse chamber approach that continues to yield useful information on the
2 relationships of O₃ uptake to both physiological and growth effects employs continuous stirred
3 tank reactors (CSTRs) first described by Heck et al. (1978). Although originally developed to
4 permit mass-balance studies of O₃ flux to plants, their use has more recently widened to include
5 short-term physiological and growth studies of O₃ × CO₂ interactions (e.g., Costa et al., 2001;
6 Heagle et al., 1994b; Loats and Rebbeck, 1999; Rao et al., 1995; Reinert and Ho, 1995; and
7 Reinert et al., 1997), and of surveys of native plant responses to O₃ (Orendovici et al., 2003).
8 In many cases, supplementary lighting and temperature control of the surrounding structure have
9 been used to control or modify the environmental conditions, e.g., Heagle et al. (1994a).

10 Many investigations have utilized commercially available controlled environment
11 chambers and walk-in rooms adapted to permit the introduction of a flow of O₃ into the
12 controlled air-volume. Such chambers continue to find use in genetic screening and in
13 physiological and biochemical studies aimed primarily at improving our understanding of mode
14 of action. For example, some of the ongoing studies of the O₃ responses of *Plantago major*
15 populations have been conducted in controlled environment chambers (Reiling and Davison,
16 1994; Whitfield et al., 1996b).

17 The environmental conditions provided by indoor chambers of any type will always
18 preclude the use of the information obtained with such chambers in predicting O₃ effects in the
19 natural environment, because the environmental conditions will always be measurably different
20 from field conditions. However, highly sophisticated controlled environment chambers such as
21 those described by Langebartels et al. (1997), which are subdivided into aerial and root
22 compartments, with dynamic control of light intensity and photoperiod, air and soil temperature,
23 humidity, soil moisture, wind speed, and exposure to O₃, may come close to simulating specific
24 natural conditions. Such chambers have provided meaningful insights into a wide array of early
25 biochemical responses of plants to O₃. They can minimize confounding factors that make indoor
26 chamber studies only rarely able to be extrapolated to field conditions, i.e., that shoots and roots
27 develop under different temperature regimes.

28 The applicability of the results of many chamber studies is also limited by their use of
29 container-grown plants. Several recent studies have raised serious questions as to the relevance
30 of pot-based studies to the true field situation (even when the studies are conducted in the field.)
31 Most of the questions have concerned studies of the effects of CO₂ enrichment, as discussed in

1 Section 9.4.7.1, but are also relevant to O₃ enrichment studies, as shown by Whitfield et al.
2 (1996a). Whitfield et al., reported significant interactive effects between O₃ and soil volume on
3 the growth of *Plantago major*. They noted that although container size may limit root and,
4 hence, plant growth, the reverse may also be true for single plants in large containers which do
5 not experience typical field competition for resources.

6 7 **9.2.2.2 Field Chambers**

8 Although closed field chambers have largely fallen out of favor in recent years, closed
9 “Solardome” field chambers (Lucas et al., 1987; Rafarel and Ashenden, 1991) have been used
10 recently in studies of O₃ × acid mist interactions (Ashenden et al., 1995; 1996).

11 Concern over the need to establish realistic plant-litter-soil relationships as a prerequisite to
12 studies of the effects of O₃ and CO₂ enrichment on ponderosa pine (*Pinus ponderosa*) seedlings
13 led Tingey et al. (1996) to develop closed, partially environmentally controlled, sun-lit chambers
14 (“terracosms”) incorporating 1 m-deep lysimeters containing forest soil that retained the
15 appropriate horizon structure.

16 In general, field chamber studies are dominated by the use of various versions of the open-
17 top chamber (OTC) design, first described by Heagle et al. (1973) and Mandl et al. in 1973.
18 Most chambers are ~3 m in diameter with 2.5 m high walls. Hogsett et al. (1987a) described in
19 detail many of the various modifications to the original OTC designs that appeared subsequently,
20 e.g., the use of larger chambers to permit exposing small trees (Kats et al., 1985); and grapevines
21 (Mandl et al., 1989), the addition of a conical baffle at the top to improve ventilation (Kats et al.,
22 1976), a frustrum at the top to reduce ambient air incursions, and a plastic rain-cap to exclude
23 precipitation (Hogsett et al., 1985). All of these modifications included the discharge of air via
24 ports in annular ducting or interiorly perforated double-layered walls at the base of the chambers
25 to provide turbulent mixing and upward mass flow of air.

26 Wiltshire et al. (1992) described a large open-top chamber suitable for small trees with
27 roll-up sides that permitted the trees to be readily subjected from time to time to episodic,
28 normal, “chamberless” environmental conditions. In the 6 m-high OTCs described by Seufert
29 and Arndt (1985) used with Norway spruce (*Picea abies*) trees, a second zone of annular
30 enrichment was also provided between 4 and 5 m. The use of OTCs was adopted for the large
31 European Stress Physiology and Climate Experiment on the effects of CO₂ and O₃ on spring

1 wheat (ESPACE-wheat), conducted in 1994-1996 at field sites in eight countries (Jäger et al.,
2 1999). However, typical European chambers have the introduction of O₃-enriched air at or
3 above canopy height. The relatively low costs of fabrication, operation and maintenance has
4 favored OTC use in field studies (Fangmeier et al., 1992; Musselman and Hale, 1997). The air
5 supplied to the chambers can be readily filtered through activated charcoal to reduce the O₃
6 concentration or it can be enriched with O₃ to provide a range of exposures.

7 All field chambers create internal environments that differ from ambient air, giving rise to
8 so-called “chamber effects” with the modification of microclimatic variables, including reduced
9 and uneven light intensity, uneven rainfall, constant wind speed, reduced dew formation, and
10 increased air temperatures (Fuhrer, 1994; Manning and Krupa, 1992). Several shortcomings of
11 the OTC design and operation relate to the means of introduction and mixing of enriched air to
12 produce a definable exposure. First, the plants are subjected to constant turbulence, which,
13 through increased uptake resulting from the consequently low boundary layer resistance to
14 diffusion, may lead to overestimation of the magnitude of real-world cause-effect relationships
15 (Krupa et al., 1995; Legge et al., 1995). Second, the introduction of the O₃-enriched air in the
16 lower part of chambers described by Heagle et al. (1973) and Mandl et al. (1973) results in a O₃
17 concentration gradient that decreases with increasing height, the converse of the situation
18 observed in ambient air, in which the O₃ concentration decreases markedly from above a plant
19 canopy to ground level (Grünhage and Jäger, 1994; Pleijel et al., 1995; Pleijel et al., 1996).
20 Concern that studies conducted in such OTCs may somewhat overestimate the effects of O₃ led
21 to the European design which provides a decreasing downward gradient. It seems unlikely that
22 the “chamber effects” produced by the two designs will be the same. These issues are discussed
23 more fully in Section 9.2.2.4.

24 It should also be noted that, although originally developed for exposing row crops in the
25 field, many recent studies employing OTCs have used potted plants in order to include or control
26 edaphic or nutritional factors or water relations within the experimental design. Therefore, the
27 same caveats as those discussed above (Section 9.2.2.1) with regard to extrapolating results of
28 pot studies to true field conditions apply to OTC studies.

29 The difficulties faced in the experimental exposure of forest trees to air pollutants in
30 chambers (e.g., Seufert and Arndt [1985]) led to the development of branch chambers such as
31 those described by Ennis et al. (1990), Houpis et al. (1991) and Teskey et al. (1991). These

chambers are essentially large cuvettes and, as noted by Musselman and Hale (1997), share many of the characteristics of CSTRs, i.e., transparent walls, internal fans, and inlet and outlet monitors to permit the determination of O₃ uptake, CO₂ exchange, and transpiration. Although they make it possible to expose whole branches to different O₃ regimes, the relevance of the data they yield in regard to the whole tree may be questioned. As noted by Saxe et al. (1998), the inevitable change in environmental conditions resulting from the isolation of the branch may cause different responses from those that would be obtained if the whole tree was subjected to the same environmental conditions.

9.2.2.3 Plume Systems

Plume systems are chamberless exposure facilities in which the atmosphere surrounding plants in the field is modified by the injection of pollutant gas into the air above or around them from multiple orifices that are spaced to permit diffusion and turbulence so as to establish relatively homogeneous conditions as the individual plumes disperse and mix with the ambient air. As pointed out by Manning and Krupa (1992), they can only be used to *increase* the O₃ levels in the ambient air. The volume of air to be modified is unconfined, and three approaches have been used to achieve desired pollutant concentrations in the air passing over the plants producing various systems that:

- (1) achieve a concentration gradient, in most instances dependent upon the direction of the prevailing wind;
- (2) achieve spatially uniform concentrations over a plot, dependent upon wind direction; and
- (3) seek to achieve spatially uniform concentrations over a plot, independent of wind speed and direction.

Gradient systems created by dispensing a pollutant gas into the air at canopy level from perforated horizontal pipes arranged at right angles to the prevailing wind were described for SO₂ studies in the early 1980s. A modified gradient system for O₃ was used by Bytnerowicz et al. (1988) to study effects on desert species, but there appear to have been no recent applications of the method. A gradient O₃-*exclusion* system is discussed in Section 9.2.3.1.

Systems designed to achieve spatially uniform pollutant levels by ensuring that the release of a pollutant is always on the upwind side of the study site were also originally described for

SO₂ studies, e.g. Greenwood et al. (1982). However, the adaptation of these concepts as introduced by McLeod et al. (1985) in constructing a large circular field site for exposing crops to SO₂ led to the subsequent development of both the large-scale O₃ and SO₂ fumigation system for forest trees in the United Kingdom in 1985 (the Liphook Forest Fumigation Project; McLeod et al., 1992), the smaller system for O₃ fumigation constructed at Kuopio, Finland in 1990 (Wulff et al., 1992), and the free-air carbon-dioxide enrichment (FACE) systems of gas dispersal over crops (Hendrey and Kimball, 1994) and forest trees (Hendrey et al., 1999). Although originally designed to provide chamberless field facilities for studying the CO₂ effects of climate change, large forest tree FACE systems have recently been adapted to include the dispensing of O₃ (Karnosky et al., 1999). Volk et al. (2003) have recently described a system for exposing grasslands that uses 7-m diameter plots. FACE systems discharge the pollutant gas (and/or CO₂) through spaced orifices along an annular ring (or torus) or at different heights on a ring of vertical pipes. Computer-controlled feedback from the monitoring of gas concentration regulates the feed rate of enriched air to the dispersion pipes. Feedback of wind speed and direction information ensures that the discharges only occur upwind of the treatment plots, and that discharge is restricted or closed down during periods of low wind speed or calm conditions. The diameter of the arrays and their heights in some FACE systems (25-30 m) requires large throughputs of enriched air per plot, particularly in forest tree systems. The cost of the throughputs tends to limit the number of enrichment treatments, although Hendrey et al. (1999) have argued that the cost on an enriched volume basis is comparable to that of chamber systems.

An alternative to the FACE system to free-air fumigation uses a horizontal grid system through which pollutant-air enriched is discharged over the canopies of plants in field plots. The original design, termed the Zonal Air Pollution System (ZAPS), was developed for studying the effects of SO₂ on native grasslands (Lee et al., 1975), and it was later modified by Runeckles et al. (1990) by randomly dividing each of three treatment plots into four sub-plots, each with different numbers of discharge orifices to provide various levels of O₃ enrichment. With the ZAPS system, changes in wind direction and speed result in varying degrees of carry-over from sub-plot to sub-plot, effectively resulting in twelve stochastically different seasonal exposures. The system was used for studies of growth effects on field crops and 2- to 4-year old Douglas fir (*Pseudotsuga menziesii*) saplings (Runeckles and Wright, 1996).

1 A larger ZAPS design was used by Wilbourn et al. (1995) on a grass (*Lolium perenne*)-
2 clover (*Trifolium repens*) mixture and by Ollerenshaw et al. (1999) on oilseed rape (*Brassica*
3 *napus*), whereby four replicate field plots were exposed to intermittent constant additions of O₃
4 to ambient air. A ZAPS design with eight spatially separated treatment plots was also developed
5 to obtain crop response data used in assessing crop losses in the Fraser Valley, British Columbia,
6 Canada (Runeckles and Bowen, 2000).

7 The FACE-type facility developed for the Kranzberg Ozone Fumigation Experiment in
8 Germany that began in 2000 (KROFEX; Werner and Fabian, 2002; Nunn et al., 2002) to study
9 the effects of O₃ on mature stands of beech (*Fagus sylvatica*) and spruce (*Picea abies*) trees is
10 more truly a zonal system that functions independently of wind direction. The enrichment of a
11 large volume of the ambient air immediately above the canopy takes place via orifices in vertical
12 tubes suspended from a horizontal grid supported above the canopy.

13 Recognizing the difficulties of modifying the aerial environments of large trees, Tjoelker
14 et al. (1994) devised a free-air system for exposing branches of sugar maple (*Acer saccharum*)
15 trees to O₃. Near the ends of up to 10 branches, enriched air was discharged through small holes
16 in 38-cm-diameter loops of 0.635-cm-OD teflon tubes positioned 20-30 cm below the terminal
17 foliage cluster.

18 Although plume systems make virtually none of the modifications to the physical
19 environment that are inevitable with chambers, their successful use depends on selecting the
20 appropriate numbers, sizes, and orientations of the discharge orifices to avoid “hot-spots”
21 resulting from the direct impingement of jets of pollutant-enriched air on plant foliage (Werner
22 and Fabian, 2002). However, because mixing is unassisted and completely dependent on wind
23 turbulence and diffusion, local gradients are inevitable even in large-scale FACE systems. Both
24 FACE and ZAPS systems have provisions for shutting down under low wind speed or calm
25 conditions and for an experimental area that is usually defined within a generous border, in order
26 to strive for homogeneity of the exposure concentrations within the treatment area. They are
27 also both dependent upon continuous computer-controlled feedback of the O₃ concentrations in
28 the mixed treated air and the meteorological conditions.

9.2.2.4 Comparative Studies

All experimental approaches to the exposure of plants to O₃ have shortcomings. The use of laboratory, greenhouse, or field chambers raises concerns for the roles of “chamber effects” on micrometeorology as well as the consistent turbulence over and within the plant canopy during chamber operation, in modifying plant responses.

With the widespread use of the OTC, numerous studies have demonstrated increased air temperatures and decreased light intensities as regularly noted features of their use. However, the question to be answered is whether or not these differences affect plant response to O₃. As noted in the 1996 criteria document (U.S. Environmental Protection Agency, 1996), evidence from the comparative studies of OTCs and from closed chamber and O₃-exclusion exposure systems on the growth of alfalfa (*Medicago sativa*) by Olszyk et al. (1986a) suggested that, since significant differences were found for fewer than 10% of the growth parameters measured, the responses were, in general, essentially the same regardless of exposure system used and “chamber effects” did not significantly affect response. In 1988, Heagle et al. (1988) concluded: “Although chamber effects on yield are common, there are no results showing that this will result in a changed yield response to O₃.” Several more recent studies have, however, indicated that the temperature effect alone may be sufficient to cause significant (though modest) shifts in quantitative responses to O₃, both positive and negative, as discussed below in Section 9.4.3.2. Still, it is not clear whether these effects are directly related to temperature or are the result of temperature interactions with other environmental variables. For example, Olszyk et al. (1992) undertook a 3-year study of the impact of O₃ on Valencia orange trees in large OTCs to determine if “insidious differences in microclimatic conditions could alter plant growth responses and susceptibility to pollutant stress.” Non-filtered chambers were found to have somewhat lower average O₃ concentrations than the ambient air, with fewer hourly exceedances of 100 ppb. In cool seasons, stomatal conductance was also lower, implying lower O₃ uptake. However, the cumulative fruit yields were doubled in the chamber trees even though photosynthetically active radiation was consistently reduced by about 19%, while leaf temperatures averaged more than 2°C higher. These data may be somewhat extreme, but they emphasize the need to avoid automatically assuming that OTCs yield response data that are all immediately relevant to the real world, particularly since, as in this study, no O₃ enrichment was involved as a complicating factor.

1 Plume systems avoid “chamber effects,” but because they rely solely upon diffusion and
2 natural turbulence to modify the ambient O₃ concentration, they may fail to achieve homogeneity
3 of the air to which the plants are exposed and may give rise to “hot spots” in which the enriched
4 air jets are inadequately diluted and impinge directly on foliage. A further deterrent to their
5 widespread use is the large-scale generation of O₃ needed, which has in most cases limited the
6 numbers of treatments that can be included in an experimental design. In spite of the various
7 advantages and disadvantages of the two field systems, it appears that no direct comparative
8 studies have been conducted.

9 10 **9.2.2.5 Ozone Generation Systems**

11 Two approaches have been used to generate the O₃ needed for enrichment purposes from
12 air or oxygen: (1) high voltage static discharge and (2) high intensity UV-irradiation. Using
13 gaseous oxygen as feedstock, both generate O₃-enriched oxygen, free from other impurities.
14 However, the use of high voltage discharge equipment with air as feedstock requires that the
15 output be scrubbed with water to remove appreciable amounts of the higher oxides of nitrogen
16 (especially N₂O₅, nitric acid vapor) that form concurrently with O₃ (Brown and Roberts, 1988;
17 Taylor et al., 1993).

18 19 **9.2.2.6 Experimental Exposure Protocols**

20 A few recent chamber studies of physiological or biochemical effects have continued to
21 use “square-wave” exposure profiles typified by a rapid rise to and falling off from a steady
22 target concentration. However, most recent experimentations into O₃ effects on plant growth and
23 development have employed either simulations of the diurnal ambient O₃ profile or
24 enhancement/reduction of the ambient O₃ concentrations.

25 Although Hogsett et al. (1985), Lefohn et al. (1986), and others have described the use in
26 controlled chambers of daily exposure profiles based on observed ambient O₃ profiles and such
27 profiles were used in the elaborately controlled chamber studies of Langbartels et al. (1997),
28 several recent chamber studies have used simpler computer-controlled half- or full-cosine wave
29 profiles to simulate the typical daily rise and fall in ambient O₃ levels (McKee et al., 1997a;
30 1997b; Mazarura, 1997).

1 The early studies with OTCs involved adding constant levels of O₃ to ambient air O₃
2 concentrations, but all recent studies have used enrichment delivery systems that maintain
3 proportionality to and track ambient O₃ concentrations to produce levels that more closely
4 resemble field observations. Both FACE and ZAPS studies have used proportional enrichment
5 to provide a range of treatments, although Wilbourn et al. (1995) and Ollerenshaw et al. (1999)
6 adjusted their systems manually to obtain a relatively constant target concentration during
7 exposure episodes.

9 **9.2.3 Methods Involving Exposures to O₃ in Ambient Air**

10 The experimental methods discussed above are largely aimed at developing quantitative
11 growth-response functions to permit the estimation of the effects of different ambient O₃
12 scenarios. Because such methodologies usually involve exposures to higher than ambient O₃
13 levels, the applicability of the functions obtained may, to some extent, be relevant only to
14 locations that are naturally subjected to high ambient O₃ levels. Furthermore, as pointed out in
15 Section 9.4, the response functions that they generate rarely incorporate other environmental,
16 genetic, and physiological factors, many of which can severely modify the magnitude of the
17 response to O₃. The consequences of ignoring such modifications have been well stated by De
18 Santis (1999). The European standard for protecting crops (based on the AOT40 index) was
19 derived from OTC studies of O₃-induced grain loss of wheat observed in experiments conducted
20 mostly in non-Mediterranean locations. However, the impact of ambient O₃ on wheat yields in
21 the Po Valley of northern Italy is much less than the devastatingly high loss (> 60%) suggested
22 by the seasonal exceedances of the standard. On a similar note, Manning (2003) has recently
23 urged the absolute necessity of seeking “ground truth” as verification of the nature and
24 magnitude of impacts on vegetation suggested by response functions using ambient O₃
25 monitoring data.

26 Such concerns clearly show that attention needs to be focused on incorporating
27 consideration of environmental and other factors into the response functions upon which
28 standards are based. This will require the development of improved simulation response models.
29 These concerns have also led to increasing attention being paid to seeking and developing
30 alternative approaches to the assessment of impact and the geographic extent of such impact,
31 approaches that are based on *in situ* exposures to ambient or sub-ambient O₃ levels. Although

one approach, the use of air exclusion systems, requires experimental facilities, the other approaches are generally based on simple field observations or measurements and, hence, can be undertaken on a wide geographic scale.

9.2.3.1 Air-exclusion Systems

The term, air-exclusion system, usually refers to chamberless field systems specifically designed to protect plants from exposure to polluted air by blowing filtered air through their canopies. Hogsett et al. (1987a; 1987b) describe several dedicated systems developed in the 1960s and 1970s, but there appear to have been no recent O₃-exclusion studies using systems specifically designed for the purpose since those described by Olszyk et al. (1986a; 1986b). Their system, a modification of the earlier system of Jones et al. (1977), consisted of perforated 31.8-cm OD inflatable polyethylene tubes laid between crop rows and supplied with charcoal-filtered air. By increasing the size of the orifices progressively in sections along the 9-m length of the tubes, an exclusion gradient was created with a progressive decrease in O₃ levels in the air surrounding the crop from one end of the system to the other. The system was used for studies on alfalfa (*Medicago sativa* L.) comparing plant response in OTCs, closed field chambers, the air-exclusion system, and ambient air plots (as discussed above in Section 9.2.2.4).

An air-exclusion component has also been part of the overall design of the many OTC experiments. Charcoal-filtered air or mixtures of charcoal-filtered and ambient air to chambers as part of the overall design.

9.2.3.2 Natural Gradients

Naturally occurring locational differences in ambient O₃ concentrations hold potential for the examination of plant response along a gradient of such concentrations. However, few such gradients can be found which meet the rigorous statistical requirements for comparable site characteristics such as soil type, temperature, rainfall, radiation, and aspect (Manning and Krupa, 1992), although with small plants, soil variability can be avoided by the use of potted plants.

Studies in the 1970s used the natural gradients occurring in Southern California to assess yield losses of alfalfa (*saliva*) and tomato (*Lycopersicon esculentum* L.) (Oshima et al., 1976; 1977). A transect study of the impact of O₃ on the growth of clover (*Trifolium repens* L.) and barley (*Hordeum vulgare* L.) in the United Kingdom was confounded by differences in the

concurrent gradients of SO₂ and NO₂ pollution (Ashmore et al., 1988). Studies of forest tree species in national parks in the eastern U.S. (Winner et al., 1989) revealed increasing gradients of O₃ and visible foliar injury with increased elevation.

9.2.3.3 Use of Chemical Protectants

The use of protective chemicals is a relatively inexpensive, promising alternative to experimental field exposures in chambers or free-air systems for determining plant response to O₃. Several chemical compounds (antioxidants, antisenescence agents, fungicides, pesticides, etc.) have been known for many years to provide plants some protection from photochemical oxidants, such as O₃ (Manning and Krupa, 1992). Most of these chemicals were originally used as a one-time application to reduce visible injury caused by acute O₃ exposures. The most widely used and popular of these has been ethylenediurea (EDU). Carnahan et al. (1978) reported that EDU protected pinto bean (*Phaseolus vulgaris*) from acute O₃ injury. After this initial investigation, EDU was shown to suppress visible O₃ injury on several species of plants under both controlled and field conditions (Brennan et al., 1987; Clarke et al., 1983). However, due to lack of a commercial market for this product, its commercial manufacture was largely discontinued. Other chemicals, including benomyl (Manning et al., 1974), carboxin (Rich et al., 1974), ascorbic acid (Dass and Weaver, 1968) and others (Manning and Krupa, 1992), exhibited some beneficial effects in reducing visible O₃ injury.

Despite their widespread use for O₃-effects studies, OTCs are not effective in determining plant responses in the field under truly natural conditions, as discussed above, and are not efficient or easily used in unmanaged ecosystems such as wilderness areas (Manning and Krupa, 1992). For these reasons, a renewed interest in using protective chemicals as research tools occurred in the early 1990s. These chemicals (mostly EDU) have recently been used in studies of different plant species, both in the United States (Bergweiler and Manning, 1999; Kuehler and Flagler, 1999) and in Europe (Bortier et al., 2001b; Pleijel et al., 1999; Wu and Tiedemann, 2002), to determine if ambient O₃ concentrations affect plant growth and productivity or are just exacerbating foliar injury. For example, Bortier et al. (2001a) injected seedlings of an O₃-sensitive poplar (*Populus nigra*) clone with EDU and measured growth over a 1-year period at a field site near Brussels, Belgium. Over the growing season, stem diameter increment was significantly higher (16%), biomass was increased (9%), and foliar O₃ symptoms were slightly

1 less for the EDU-treated seedlings. Ozone levels were reported to be low (AOT40 = 6170 ppb.h,
2 May-September) during the exposure period. In another study, Manning et al. (2003) applied
3 EDU (foliar spray) and sodium erythorbate (NaE) at various concentrations, biweekly for three
4 growing seasons to loblolly pine (*Pinus taeda*) at a field site in east Texas. After 3 years, the
5 trees were harvested and biomass measured. Neither EDU nor NaE prevented foliar O₃ injury,
6 but EDU applications at 450 ppm resulted in increases in stem diameter and height and total
7 above-ground biomass. These measures of growth also tended to slightly increase with
8 applications of NaE, but the effects were nonsignificant.

9 Several recent studies used EDU in assessing the response of several plant species to O₃ to
10 help validate the proposed critical level (AOT40 = 3000 ppb.h) for crop protection in Europe
11 (Ball et al., 1998; Ribas and Penuelas, 2000; Tonneijck and Van Dijk, 2002b; 2002a). EDU
12 appeared to provide protection from visible foliar injury, but the results regarding yield and
13 biomass reductions were mixed. In a 3-year study over 12 sites throughout Europe, Ball et al.
14 (1998) used the ratio of EDU-treated versus non-treated white clover (*Trifolium repens*)
15 biomass but did not find a significant relationship between biomass reductions and AOT40.
16 When other parameters, such as temperature and VPD were included in the model however,
17 they found a significant relationship ($r^2 = 0.79$) between biomass reductions and AOT40, and
18 the greatest biomass reductions occurred in areas with the highest levels of industrialization
19 (in Germany).

20 Tonneijck and Van Dijk (2002b) assessed the relationship of visible injury of subterranean
21 clover (*Trifolium subterraneum*) to ambient O₃ at four sites over three growing seasons in the
22 Netherlands, using EDU-treated and non-treated plants. Visible injury varied by site and year,
23 but was reduced to near zero by EDU treatment. However, no relationship indicative of a
24 protective effect of EDU with this plant species was observed for biomass. Tonneijck and
25 Van Dijk (2002a) also reported similar results with bean (*Phaseolus vulgaris*). Both EDU-
26 treated and non-treated plants were exposed to ambient O₃ at three locations in Spain over one
27 growing season (Ribas and Penuelas, 2000). Reductions in yield and biomass were correlated
28 with O₃ concentration and EDU provided some protective effect, although results varied by
29 location and with meteorological conditions.

30 The mechanisms by which protective chemicals, especially EDU, protect plants are poorly
31 understood. However, Wu and von Tiedemann (2002) reported that applications of two recently

developed fungicides (azoxystrobin and epoxiconazole) provided protection of spring barley to relatively high O₃ exposures (150-250 ppb, 5 days, 7h/d) and resulted in increases in leaf soluble protein content, as well as the activity of several antioxidative enzymes (e.g., superoxide dismutase, catalase, ascorbate-peroxidase, and glutathione reductase). In addition, the increase in these enzymes reduced superoxide levels in the leaves.

Before using these chemicals in a field setting, preliminary investigations should be done to evaluate the methods and timing of application, as well as proper application rates, so as to avoid any potential toxic effects (Manning, 2000; Manning and Krupa, 1992). Also, because certain plant species may be nonresponsive, careful pre-screening of various species to be tested is critical.

In summary, the use of protectants such as EDU (see Table 9-1) provides a ready means of demonstrating the local occurrence of adverse effects of O₃, both in terms of visible injury and growth reduction. However, even if accompanied by some type of O₃ and meteorological monitoring, their use has yet to give rise to a methodology for quantitatively assessing exposure response.

9.2.3.4 Biomonitoring

Bioindicators

The use of biological indicators to detect the presence of O₃ injury to plants is a longstanding and effective methodology (Chappelka and Samuelson, 1998; Manning and Krupa, 1992). A bioindicator can be defined as a vascular or non-vascular plant exhibiting a typical and verifiable response when exposed to a plant stress such as an air pollutant (Manning et al., 2003). To be considered a good indicator species, plants must:

- (1) exhibit a distinct, verified response;
- (2) have few or no confounding disease or pest problems; and
- (3) exhibit genetic stability.

Such sensitive plants can be used to detect the presence of a specific air pollutant such as O₃ in the ambient air at a specific location or region and, as a result of the magnitude of their response, provide unique information regarding specific ambient air quality. Bioindicators can be either introduced *sentinels*, such as the widely used tobacco (*Nicotiana tabacum*) variety

Table 9-1. Advantages and Disadvantages of Protective Chemicals Used in Assessment of O₃ Effects on Plants

Advantages
No chambers required. Plants exposed to ambient conditions of O ₃ , light, temperature, etc.
Can conduct studies “in situ.” Equipment needs are minimal. No “chamber effects”
A high degree of replication possible both within and among locations
Disadvantages
Exposure-response studies require inclusion of other methodologies (OTCs, etc.)
Need measurements of ambient O ₃ and other meteorological variables (temp, rainfall, etc)
Have to conduct preliminary toxicology studies to determine proper rate, timing etc.
Possible plant toxicity can result from repeated applications
Species response can vary; need to screen for proper species to use

Modified from Manning and Krupa (1992).

Bel W3, or *detectors*, which are sensitive native plant species. The approach is especially useful in areas where O₃ monitors are not operated (Manning et al., 2003). For example, in remote wilderness areas where instrument monitoring is generally not available, the use of bioindicator surveys in conjunction with the use of passive samplers (Krupa et al., 2001) is a particularly useful methodology (Manning et al., 2003). However, the method requires expertise or training in recognizing those signs and symptoms uniquely attributable to exposure to O₃ as well as their quantitative assessment.

Since the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) many new sensitive species have been identified from controlled exposure studies and verified in the field (Flagler, 1998; Innes et al., 2001). In addition, several new uses of this methodology have been demonstrated, including a national O₃ bioindicator network, studies in wilderness areas, and mature tree studies. Although it has been difficult to find robust relationships between the foliar injury symptoms caused by O₃ and effects on plant productivity or ecosystem function, visible injury correlations with growth responses have been reported (Table 9-2; Chappelka and Samuelson, 1998; Manning et al., 2003; Smith et al., 2003). One workshop on the utility of bioindicators of air pollutants led to a useful series of peer-reviewed publications in *Environmental Pollution* (Skelly, 2003).

**Table 9-2. Advantages and Disadvantages of Bioindicators Used to Study
O₃ Plant Effects**

Advantages	
No chambers required. Plants exposed to ambient conditions of O ₃ , light, temperature, etc.	
Relatively inexpensive. Equipment needs are minimal. No “chamber effects”	
A high degree of replication possible (sentinels) both within and among locations	
Disadvantages	
Individuals need to be trained and experienced in O ₃ symptom recognition	
Need adequate numbers of plants (detectors) to ensure valid results	
Need preliminary tests to insure a constant symptomatology of material used	
Use more than one indicator species (detector) per area if possible	
Quantify site characteristics (soils, light) that may influence symptom expression	
Need measurements of ambient O ₃ (active or passive) and other meteorological variables (temp, rainfall, etc)	
Need to ensure cultural (sentinels) practices (soil, irrigation, fertilization, etc.) are similar among sites	

National network

The U.S. Forest Service in cooperation with other federal and state agencies developed a network of O₃ bioindicators to detect the presence of O₃ in forested systems throughout the U.S. (Smith et al., 2003). This ongoing program was initiated in 1994; and 33 states currently participate. In a coordinated effort, a systematic grid system is used as the basis of plot selection, and field crews are trained to evaluate O₃ symptoms on sensitive plant species within the plots (Coulston et al., 2003; Smith et al., 2003).

The network has provided evidence of O₃ concentrations high enough to induce visible symptoms on sensitive vegetation. From repeated observations and measurements made over a number of years, specific patterns of areas experiencing visible O₃ injury symptoms can be identified. Coulston et al. (2003) used information gathered over a 6-year period (1994-1999) from the network to identify several species that were sensitive to O₃ over a regional scale

1 included sweetgum (*Liquidambar styraciflua*), loblolly pine (*Pinus Taeda*), and black cherry
2 (*Prunus serotina*).
3

4 ***Wilderness areas***

5 The use of bioindicator species as detectors has proven to be an effective technique for
6 deriving a relative estimate of O₃ injury in wilderness areas in both the US and Europe
7 (Chappelka et al., 1997; 2003; Manning et al., 2002). However, to be truly effective, these
8 regional and national bioindicator studies need the inclusion of air quality data and related
9 growth studies to determine effects on productivity and ecosystem function (Bytnerowicz et al.,
10 2002; Manning et al., 2003; Smith et al., 2003).

11 Chappelka et al. (1997; 2003) conducted surveys of foliar injury on several native plant
12 species throughout the Great Smoky Mountains National Park (GRSM), including black cherry
13 (*Prunus serotina*), tall milkweed (*Asclepias exaltata*), cutleaf coneflower (*Rudbeckia laciniata*)
14 and crownbeard (*Verbesina occidentalis*). Visible foliar symptoms were prevalent throughout
15 the Park, indicating that injury-producing O₃ levels were widespread in GRSM.

16 Manning et al. (2002) recently summarized a multi-year (1993-2000) bioindicator project
17 in the Carpathian Mountain range in eastern Europe. They evaluated numerous trees, shrubs,
18 forbs, and vines for possible symptoms of O₃ injury. Observations were made at plots located in
19 the vicinity of either active or passive O₃ monitors (Bytnerowicz et al., 2002). Approximately 30
20 species of native plants detectors were identified as possible bioindicators, the majority of which
21 (21) were shrubs (Manning et al., 2002). Based on these observations, it was concluded that O₃
22 concentrations were sufficiently high to impact ecosystems in the region. Similar investigations
23 regarding the sensitivity of native species have been conducted in Switzerland (Novak et al.,
24 2003) and Spain (Orendovici et al., 2003).
25

26 ***Mature tree detectors***

27 Many studies have reported visible injury of mature coniferous trees caused by O₃,
28 primarily in the western United States (Arbaugh et al., 1998) and, to a lesser extent, to mature
29 deciduous trees in eastern North America. In an effort to determine the extent and magnitude of
30 visible injury in mature tree canopies, Hildebrand et al. (1996) and Chappelka et al. (1999b)
31 conducted independent studies in the GRSM and the Shenandoah National Park (SHEN). The

1 species examined were sassafras (*Sassafras albidum*), black cherry, and yellow-poplar
2 (*Liriodendron tulipifera* L.) in GRSM and white ash (*Fraxinus americana* L.), black cherry, and
3 yellow-poplar in SHEN. Protocols were similar at both parks, and trees were located near O₃
4 monitors at three different areas in each park. Results from both studies indicated that symptoms
5 of O₃ injury were present in the trees and correlated with the amount of injury both spatially and
6 temporally. Ozone injury tended to be most severe at the highest elevation, except with yellow-
7 poplar.

8 Hildebrand et al. (1996) observed significant O₃ exposure-plant response relationships with
9 black cherry. The best relationships were found between foliar injury and the SUM06 and W126
10 exposure indices, indicating that higher O₃ concentrations were important in eliciting a response
11 in black cherry. No O₃ exposure-plant response relationships were found with any species tested
12 in GRSM (Chappelka et al., 1999b); but, when the data were combined for both parks, a
13 significant correlation $r = 0.72$) with black cherry was found for both SUM06 and W126, and
14 injury was the greatest ($r = 0.87$) at the higher elevations (Chappelka et al., 1999a).

15 Based on a study in which visible symptoms of O₃ injury were characterized for large,
16 mature yellow-poplar and black cherry trees in GRSM (Chappelka et al., 1999a), Somers et al.
17 (1998) compared radial growth differences among trees classified as sensitive or non-sensitive
18 based on the severity of visible foliar injury observed over a 3-year period (1991-1993).
19 Significantly more radial growth was observed over both a 5- and a 10-year period for the non-
20 sensitive compared to the sensitive trees. No significant relationship was found for black cherry
21 tree growth.

22 Vollenweider et al. (2003), using data collected from continuous forest inventory (CFI)
23 plots across Massachusetts, compared growth rates among either symptomatic or asymptomatic
24 mature black cherry trees. Of the 120 trees sampled in 1996, 47% exhibited visible foliar injury.
25 Using CFI data, growth rates were compared over a 31-year period. The growth rates for
26 symptomatic trees were reduced by 28% compared with the asymptomatic trees.

27 Because these studies (Somers et al., 1998; Vollenweider et al., 2003) were not controlled
28 studies and used a small sample of trees, they cannot validly be used to characterize cause and
29 effects related to the visible symptoms and radial growth they describe. However, the results
30 indicate the *possibility* that O₃ is correlated with growth losses in some sensitive genotypes,

1 illustrating the potential usefulness of this visible O₃ injury methodology in assessing effects on
2 the growth rates of mature deciduous trees.

4 ***Cultivar Comparisons***

5 The idea of using cultivars or isogenic lines of crop species that differed in O₃ sensitivity as
6 sentinels to determine the ambient effects of O₃ in the field was presented in the 1996 O₃ AQCD
7 (U.S. Environmental Protection Agency, 1996). The rationale was that comparing the ratio of
8 injury scores or some measure of growth between two different cultivars varying in O₃
9 sensitivity should be indicative of the relative amount of ambient stress to plants at a given
10 location. A sensitive:resistant ratio close to unity would indicate relatively low O₃
11 concentrations and a low ratio higher O₃ levels. Results from locations differing in O₃
12 concentrations could be evaluated to develop exposure-response models. The original protocol
13 was derived using two isogenic lines of white clover (*Trifolium repens*) differing in O₃
14 sensitivity (Heagle et al., 1994b; 1995).

15 This white clover model system has been used in several multi-location studies in the U.S.
16 (Heagle and Stefanski, 2000) and Europe (Ball et al., 2000; Bermejo et al., 2002; Mills et al.,
17 2000). Heagle and Stefanski (2000) compared results from eight sites over a 2-year period with
18 various exposure indices (SUM00, SUM06, W126 and others) to determine a best-fit regression.
19 They found that most of the indices performed similarly. The highest r² values (0.87-0.93) were
20 obtained using only the later harvests and a 6 h d⁻¹ index (1000-1600 h). Similar multiple-
21 comparison studies conducted in Europe using the AOT40 index (Ball et al., 2000; Mills et al.,
22 2000) yielded poorer r² values. Factors such as air temperature, NO_x (high levels at some sites)
23 and lower O₃ concentrations in Europe were suggested to account in part for the differences
24 between U.S. and Europe study results. Bermejo et al. (2002), in a study in Spain, improved the
25 model by comparing the biomass ratio of these white clover isolines to measures of O₃ uptake
26 (flux) rather than an exposure index (AOT40). Together, these studies indicate that systems
27 such as the white clover model can help reveal O₃ exposure-response relationships that can
28 provide valuable information regarding ambient O₃ conditions in a given location. Table 9-3
29 lists the advantages and disadvantages of the use of cultivar comparisons in assessing O₃ effects
30 of plants.

Table 9-3. Advantages and Disadvantages of Cultivar Comparisons Used in Assessment of O₃ Effects on Plants

Advantages
No chambers required. Plants exposed to ambient conditions of O ₃ , light, temperature, etc.
Relatively inexpensive. Equipment needs are minimal. No “chamber effects”
A high degree of replication possible both within and among locations
Can conduct studies “in situ”
Disadvantages
Preliminary tests to insure sensitivity and growth patterns of genotypes used are consistent are needed
Need measurements of ambient O ₃ and other meteorological variables (temp, rainfall, etc)
Have to ensure cultural practices (soil, irrigation, fertilization, etc.) are similar among sites
Closely monitor plants for presence of other factors that may cause a misinterpretation of results

Dendrochronological techniques

It has been difficult to determine whether O₃ significantly affects tree growth and productivity in the field, because O₃ concentrations are omnipresent and tree response to this pollutant is altered by many factors. The use of dendrochronological techniques to answer questions regarding ambient O₃ effects on forest growth and ecosystem function has recently emerged as a very useful biomonitoring methodology (Cook, 1990; McLaughlin et al., 2002). The technique is useful when either instrument or passive O₃ monitoring methods are used to determine ambient O₃ conditions.

Initial experiments were primarily correlative in nature and attempted to relate symptoms of visible injury with growth losses as revealed by tree ring analysis (Arbaugh et al., 1998; Benoit et al., 1983; Peterson et al., 1995; Somers et al., 1998; Swank and Vose, 1990). These studies evaluated radial growth patterns determined by cores removed from trees in the presence or absence of overt O₃ injury symptoms.

The method has also been adapted to better understand forest ecosystem function (McLaughlin and Downing, 1996; McLaughlin and Downing, 1995; Bartholomay et al., 1997; McLaughlin et al., 2003). The response of mature loblolly pine (*Pinus Taeda*) growing in

1 eastern Tennessee to ambient O₃ and moisture stress was evaluated by McLaughlin and Downing
2 (1996, 1995) over a 6-year period (1988 to 1983). They made radial growth measurements
3 from 12 to 37 times per year using dendrometer bands and determined relationships between O₃,
4 moisture stress, and radial growth. Exposures to O₃ concentrations $\geq 0.04 \mu\text{l L}^{-1}$ with high
5 temperatures and low soil moisture resulted in short-term depression in radial growth.
6 Reductions in growth were estimated to vary from 0 to 15% per year and averaged
7 approximately 5% per year.

8 Bartholomay et al. (1997) examined white pine (*Pinus strobus*) radial growth in eight
9 stands throughout Acadia National Park, Maine over a 10-year period from 1983 to 1992. They
10 related growth rates to several factors, including O₃ concentration. Ozone levels were negatively
11 correlated with radial growth in seven of the eight stands. Site characteristics were important in
12 the relationship: stands growing on shallow, poorly drained soils were most sensitive to O₃ in
13 the late portion of the growing season, possibly due to premature senescence of foliage.
14 However, litterfall measurements were not reported. Trees growing on better sites were more
15 sensitive to O₃ during the entire growing season, indicating the possibility of high O₃ uptake
16 rates throughout the growing season. Although these field studies (Bartholomay et al., 1997;
17 McLaughlin and Downing, 1996; McLaughlin and Downing, 1995) did not compare the firm O₃
18 effects on the two pine species, they indicate that potential interactions exist among O₃ and other
19 climatic and edaphic factors, such as temperature and soil moisture.

20 Using both automated and manual dendrometer bands, McLaughlin et al. (2002) examined
21 the growth response of yellow poplar (*Liriodendron tulipifera*) trees recently released from
22 competition. In addition to measuring growth, sap flow measurements were conducted and soil
23 moisture was measured in the vicinity of the trees. However, O₃ concentrations were low and
24 there were no negative O₃ effects observed in this 1-year study. Advantages and disadvantages
25 of dendrochronology techniques for evaluating whole-tree physiological responses for individual
26 trees and forest stands are listed in Table 9-4.

27 The use and evolution of various dendrochronological methods in the field of air pollution
28 effects research is reviewed in detail by McLaughlin et al. (2002). Automated dendrometer
29 bands provide a powerful tool for measuring radial growth responses of trees on an hourly or
30 daily basis. Diurnal patterns of growth can be related to water use and O₃ concentrations using
31

Table 9-4. Advantages and Disadvantages of Various Dendrochronological Techniques Used in Assessment of O₃ Effects on Plants

Advantages	
	Provide information regarding growth effects under ambient conditions
	Good historical information regarding O ₃ effects
	Can provide data on daily, and seasonal growth and O ₃ patterns and relate these to physiological function
	Provide information on forest function related to ambient O ₃ concentrations
	Can link data with process level growth models
Disadvantages	
	Results are generally correlative in nature with no true control
	Need background O ₃ and meteorological data (historical records)
	Need to account for other factors such as competition, in analyzing data
	Individuals need to be trained in counting growth rings
	Replication can be difficult (expensive and technological limitations)
	Complicated statistical analyses are sometimes required
	Can be expensive, especially if using automated growth (dendrometer) bands

time-series analyses. The major drawbacks of the method are that it is expensive and time consuming.

9.2.3.5 Calibrated Passive Monitors

Many studies have used passive monitors in the mapping of ambient O₃ concentrations, especially in remote areas (Cox and Malcolm, 1999; Grosjean et al., 1995; Krupa et al., 2001). Since they are cumulative recording devices, they cannot reveal short-term variations in O₃ concentration but only the total exposure over a given interval, usually 7 days. Thus, they produce a measurement that resembles the instrumentally derived exposure SUM00 index.

Runeckles and Bowen (2000) used the ZAPS system described in Section 9.2.2.3 to subject both crops and passive monitors (Williams, 1994) to a range of exposures. Passive monitors were also exposed at 16 agricultural field sites along a transect through the Fraser Valley, British

Columbia, Canada. Most field sites were downwind of the Greater Vancouver metropolitan area. All passive monitors were replaced at weekly intervals and the data from those in the ZAPS plots were “calibrated” to crop responses by means of Weibull exposure-response functions. Since the meteorological conditions throughout the valley were reasonably consistent from site to site, the use of these functions with data from the network passive monitors as inputs permitted the estimation of crop losses at the network sites. The overall method was thus a hybrid of several methodologies.

Although based on a single study, the use of passive monitors has potential for assessing crop losses at sites removed from locations with known ambient O₃ concentrations. Provided that the network and calibration sites have similar meteorological conditions, the method yields crop loss estimates that are responses to local ambient O₃ levels as influenced by local meteorological conditions

9.2.4 Numerical/Statistical Methodologies

Proper experimental design strategies including replication, randomization, and experimental protocols are paramount in O₃-effects research. These have been discussed in detail in previous O₃ AQCDs (U.S. Environmental Protection Agency, 1986, 1996), as have the different statistical analytical procedures used to determine the probable significance of results. However, new investigative approaches have demanded the adoption of new analytical methods. For example, the use of dendrochronological techniques has led to the use of time-series analysis (McLaughlin et al., 2003) and linear aggregate models (Cook, 1990), as reviewed by McLaughlin et al. (2002).

In spite of the rigors of the analyses, many differences occur in the published literature for almost any plant response to O₃ stress. Differences inevitably result from different researchers studying different locations, using different experimental methodologies and genetically different plant material even when using a common species. The techniques of meta-analysis can be used to consolidate and extract a summary of significant responses from a selection of such data.

Despite the differences in responses in the 53 primary studies used, a recent meta-analysis by Morgan et al. (2003) of the effects of O₃ on photosynthesis, growth, and yield of soybean (*Glycine max*) showed “overwhelming evidence for a significant decrease in photosynthesis, dry

1 matter production and yield ... across all the reported studies on effects of chronic O₃ treatment.”
2 The meta-analysis defined O₃ stress exposure to ~70 ppb for at least 7 days and found average
3 shoot biomass and seed yield decreases of 34% and 24%, respectively. Furthermore, although
4 other stress factors such as drought and UV-B did not affect the O₃ responses, elevated CO₂ was
5 found to significantly decrease O₃-induced losses.

6 The meta-analysis method clearly has the potential to consolidate and refine the
7 quantitative exposure-response models for many species. The majority of the reported growth
8 and physiological responses related to O₃ stress are for individual plants, primarily in various
9 types of exposure chambers. It is difficult to extrapolate these responses to stand/community,
10 ecosystem, or region-wide assessments, particularly in view of the importance of the significant
11 interactions that may occur between plant responses O₃ and other environmental stresses. Along
12 with the shift in effects research to a more ecological approach, these concerns necessitate a
13 move from simple regression analysis to more complex mathematical approaches to handle a
14 wider array of independent input variables than O₃ exposure alone. Other independent input
15 variables that must be accounted for include air and soil temperatures, soil moisture, relative
16 humidity, wind speed, and, particularly in the case of natural systems, biotic factors such as pests
17 and pathogens, plant density/spacing, and measures of plant competition.

18 Artificial neural network (ANN) methodology was used by Balls et al. (1995) for
19 “unraveling the complex interactions between microclimate, ozone dose, and ozone injury in
20 clover” and in the study with the protectant chemical EDU, discussed in Section 9.2.3.3 (Ball
21 et al., 1998). The multi-factor model for predicting the effects of ambient O₃ on clover
22 developed by Mills et al. (2000) utilized both ANN and multiple linear regression methods.

23 Models incorporating ANNs are of the “regression” type (Luxmoore, 1988) in contrast to
24 “mechanistic” or “phenomenological” models which have wider applicability. Process-level
25 models of either type have been developed at the organelle, individual plant (Constable and
26 Taylor, Jr., 1997; Weinstein et al., 1998), canopy (Amthor et al., 1994), and stand level (Ollinger
27 et al., 1997; Weinstein et al., 2001) and provide estimates of the rate of change of response
28 variables as affected by O₃ over time. However, as pointed out in the 1996 Criteria Document
29 (U.S. Environmental Protection Agency, 1996), mechanistic process models lack the precision of
30 regression models as well as their ability to estimate the likelihoods of responses. In their
31 extensive reviews, Kickert and Krupa (1991) and Kickert et al. (1999) summarized the

advantages and shortcomings of many different models and make the important point that most of the models that have been described provide *consequence* assessments that quantify the magnitudes of effects, but not *risk* assessments that quantify the likelihoods of such effects. Descriptions of several specific models are provided in other sections of this criteria document, and advantages and disadvantages of modeling techniques used in assessing O₃ effects on plants are summarized in Table 9-5.

Table 9-5. Advantages and Disadvantages of Modeling Techniques Used in Assessment of O₃ Effects on Plants

Advantages
<ul style="list-style-type: none"> • Provides an understanding of cause-effect relationships over time
Disadvantages
<ul style="list-style-type: none"> • Have to make assumptions based on scarcity of data • Most models are very complex and difficult to understand • Need to be evaluated for predictive validity

9.2.5 Improved Methods for Defining Exposure

Ambient air quality is defined in terms of the measured O₃ concentrations in the air at some standard elevation above ground level. Compilations of such concentration data have long been used as surrogates of the exposures to which plants are subjected. However, as long ago as 1965, field research provided evidence that plant response was a function, not of ambient O₃ concentration *per se*, but of the estimated flux of O₃ to the plant canopy (Mukammal, 1965). Subsequently, Runeckles (1974) introduced the term “effective dose” to define that part of the ambient exposure that was taken up by a plant. Fowler and Cape (1982) later referred it as “pollutant applied dose” (PAD), defined as the product of concentration, time and stomatal (or canopy) conductance, with units g m⁻². Such estimates of O₃ uptake or flux provide a more biologically relevant description of exposure than the simple product of concentration and time

1 alone, and formed the basis of Reich's 1983 "unifying theory" of plant response to O₃ (Reich,
2 1983).

3 However, it was not until the early 1990s that the inherent advantages of using O₃ flux
4 rather than O₃ concentration as a basis for determining response effects began to be widely
5 accepted, as demonstrated by the subsequent increase in publications involving flux
6 measurements and modeling (e.g., Fuhrer et al., 1997; Grünhage and Jäger, 2003; Grünhage
7 et al., 1993, 1997; Massman et al., 2000; Musselman and Massman, 1999; Pleijel, 1998). A key
8 requirement for flux determination is the measurement of stomatal or canopy conductances,
9 using established porometer/cuvette techniques or eddy correlation methods. The usefulness and
10 relevance of flux as a measure of exposure are discussed in detail in Section 9.5.

11 Efforts to develop models of O₃ deposition and stomatal uptake are currently under way
12 with a view to providing improved assessments of the risks to vegetation across Europe
13 (Emberson et al., 2000; Simpson et al., 2001; Simpson et al., 2003).

16 **9.3 SPECIES RESPONSE/MODE-OF-ACTION**

17 **9.3.1 Introduction**

18 The evaluation of O₃ risk to vegetation requires fundamental understanding of both the
19 functioning of the vegetation and how external environmental influences can alter that function.
20 For biological organisms subjected to atmospheric O₃, those alterations can be complex and
21 multiple. In addition, biological organisms have plasticity to external interactions due to their
22 complex internal, self-correcting systems, making the task of identifying their "correct"
23 functioning difficult. This section emphasizes reactions of O₃ with the cell and tissue rather than
24 the whole plant to describe the fundamental mechanisms known to govern the response of the
25 plant to O₃ exposure.

26 The many regulatory systems contained in leaves change both as a function of leaf
27 development and in response to various environmental stresses. Leaves function as the major
28 regulators of anatomical and morphological development of the shoot and control the allocation
29 of carbohydrates to the whole plant (Dickson and Isebrands, 1991). This section discusses the
30 movement of O₃ into plant leaves and their biochemical and physiological responses to O₃.

1 The 1996 criteria document (U.S. Environmental Protection Agency, 1996) assessed the
2 information available at that time concerning the biochemical and physiological responses to the
3 movement of O₃ into plant leaves. This information continues to be valid. Ozone uptake in a
4 plant canopy is a complex process involving adsorption to surfaces (leaves, stems, and soil) and
5 absorption into leaves (Figure 9-1). However, the initial biochemical changes that result within
6 leaf cells after the entry of O₃ and how these changes interact to produce plant responses remain
7 unclear. The response of vascular plants to O₃ may be viewed as the culmination of a sequence
8 of physical, biochemical, and physiological events. Only the O₃ that diffuses into a plant
9 through the stomata (which exert some control on O₃ uptake) to the active sites within a leaf
10 impairs plant processes or performance. An effect will occur only if sufficient amounts of O₃
11 reach sensitive cellular sites that are subject to the various physiological and biochemical
12 controls within the leaf cells. Ozone injury will not occur if (1) the rate and amount of O₃ uptake
13 is small enough for the plant to detoxify or metabolize O₃ or its metabolites, or if (2) the plant is
14 able to repair or compensate for the O₃ impacts (Tingey and Taylor, 1982; U.S. Environmental
15 Protection Agency, 1996). Therefore, a precondition for O₃ to affect plant function is that it
16 must enter the stomata and be absorbed into the water lining the mesophyll cell walls. The
17 response of each plant is determined by the amount of O₃ entering the leaves, which varies from
18 leaf to leaf.

19 Some potentially significant processes have been investigated since the 1996 criteria
20 document, especially detoxification and compensatory processes. The role of detoxification in
21 providing a level of resistance to O₃ has been investigated; however, it is still not clear as to what
22 extent detoxification can protect against O₃ injury. Data are needed especially on the potential
23 rates of antioxidant production and on the subcellular localization of the antioxidants. Potential
24 rates of antioxidant production are needed to assess whether they are sufficient to detoxify the O₃
25 as it enters the cell. The subcellular location(s) are needed to assess whether the antioxidants are
26 in cell wall or plasmalemma locations that permit contact with the O₃ before it has a chance to
27 damage subcellular systems. Various forms of compensation, especially the stimulation of new
28 leaf production and of higher photosynthetic performance of new leaves, have been reported.
29 Although these processes divert resources away from other sinks, these forms of compensation
30 may counteract the reduction in canopy carbon fixation caused by O₃. The quantitative
31 importance of these processes requires investigation.




Figure 9-1. Ozone uptake from the atmosphere. Ozone moves from the atmosphere above the canopy boundary layer into the canopy primarily by turbulent air flow. Canopy conductance, controlled by the complexity of the canopy architecture, is a measure of the ease with which gases move into the canopy. Within the canopy, O_3 is adsorbed onto surfaces as well as being absorbed into the foliage. Foliage absorption is controlled by two conductances, leaf boundary layer and stomatal, which together determine leaf conductance. The solid black arrows denote O_3 flow; dotted arrows indicate processes affecting uptake or response to O_3 . Boxes at the left with double borders are those processes described in the figure.

As a result of the research since the 1996 criteria document (U.S. Environmental Protection Agency, 1996), the way in which O_3 exposure reduces photosynthesis, especially its effects on the central carboxylating enzyme, Rubisco (ribulose-1,6- P_2 -carboxylase/oxygenase1), is better understood. The rate of leaf senescence has been shown to increase as a function of increasing O_3 exposure. The mechanism of the increased senescence is not known, and, hence, it deserves further study.

1 Finally, the role that changes in allocation of resources play in plant response to O₃ is now
2 better understood. Most studies have shown that O₃ decreases allocation of photosynthate to
3 roots. In some cases, allocation to leaf production has increased. Whether these changes are
4 driven entirely by changes in carbohydrate availability or are controlled by other factors (e.g.,
5 hormones) is not known. Physiological effects within the leaves inhibit photosynthesis, alter the
6 assimilation of photosynthate and shift its allocation patterns, and can lead to reduced biomass
7 production, growth, and yield (U.S. Environmental Protection Agency, 1986, 1996).

8 The major problem facing researchers trying to predict long-term O₃ effects on plants is
9 determining how plants integrate the responses to O₃ exposures into the overall naturally
10 occurring responses to environmental stressors. Little is now known about how plant responses
11 to O₃ exposures change with increasing age and size, but this information is crucial to predicting
12 the long-term consequence of O₃ exposure in forested ecosystems.

13 This section focuses on reactions of O₃ within cells and cellular tissue, in order to explain
14 known mechanisms that govern plant responses. The processes that occur at cell and tissue
15 levels within the leaf will be divided into several steps beginning with O₃ uptake and its initial
16 chemical transformations into a series of currently unknown, but suspected toxic, chemicals
17 (Figure 9-2). The discussion will then focus sequentially upon various cell regions, their general
18 physiology, and the changes that may occur within a plant after O₃ exposure. This is important
19 because the varying responses of the different plant species in a community ultimately lead to
20 an ecosystem response. Finally, a general summary is presented that discusses the known or
21 suspected changes that occur within the whole plant.

22 23 **9.3.2 Mechanisms of Ozone-Induced Plant Alterations**

24 Plant adaptations for surviving O₃ stress include exclusion or tolerance of it or its products
25 (Levitt, 1972; Tingey and Taylor, 1982). Ozone may be excluded from tissues or cells via
26 stomatal closure, by extracellular oxidants, or by membrane impermeability to O₃ or its products.
27 Past investigations of O₃ injury have indicated that physiological and metabolic changes occur
28 (see Heath, 1998; Reddy et al., 1993; Harris and Bailey-Serres, 1994). Many of these changes
29 are likely initiated via gene expression. During the last decade, our understanding of the cellular
30 processes within plants has increased. Although the fundamental hypotheses concerning




Figure 9-2. Absorption and transformation of O_3 within the leaf. The varied processes are broken down in to smaller mechanistic steps that lead from uptake of atmospheric O_3 into the alterations which may occur within the individual plant. Each plant responds to the O_3 level and therefore interacts with the total ecological setting to generate an ecosystem response due to the O_3 .

O_3 -induced changes in physiology have not changed, a more complete development of the theories is now nearing possibility.

9.3.2.1 Changes in Metabolic Processes: Current Theories

The current hypotheses regarding the biochemical response to O_3 fumigation revolve about injury and its prevention. These are well discussed by Pell et al. (1997) and are listed below in no order of importance. Although they are listed separately, some may be interlinked and related to each other.

- (1) Membrane Dysfunction. The membrane is altered by O₃, principally via protein changes not involving the lipid portions of the membrane (except at extremely high levels of O₃). These alterations involve increased permeability with perhaps lessened selectivity, declines in active transport, and changes in the trigger mechanisms of signal transduction pathways such that the signals are no longer suitable for the state of the cell. The cellular pools and transport systems of Ca²⁺/K⁺/H⁺ are the primary suspects.
- (2) Antioxidant Protectants. Varied antioxidants (both as metabolites and enzyme systems) can eliminate the oxidant or its products, if present at time of fumigation and in the sufficient abundance. However, oxidant entry that occurs rapidly can overwhelm the antioxidant response.
- (3) Stress Ethylene Interactions. Visible injury is caused by the interaction of O₃ with stress-induced ethylene, either by direct chemical transformation to a toxic product or by alteration of the biochemical relations at the ethylene binding site.
- (4) Impairment of Photosynthesis. A product of O₃ (and less probably, O₃ itself) enters the cell, causing a decline in the mRNA for Rubisco (especially the message RNA species of *rbcS* and *rbcL*) such that Rubisco levels slowly decline within the chloroplast, leading to a lowered rate of CO₂ fixation and productivity. This process is very similar to early senescence and may be linked to general senescence. Alternatively, a false signal is generated at the cell membrane which lowers the transcription of DNA to mRNA. Ozone alters the normal ionic and water relations of guard cells and subsidiary cells, causing the stomata to close and limit CO₂ fixation. In any case, the response of the stomata to the current environment does not promote efficient photosynthesis.
- (5) Translocation Disruption. One of the biochemical systems most sensitive to O₃ exposure is the translocation of sugars, such that even a mild exposure inhibits the translocation of carbohydrate (Grantz and Farrar, 1999, 2000).
- (6) General Impairment/Disruption of Varied Pathways of Metabolism. This is the oldest and most vague concept of how O₃ alters metabolism. It is based upon early work in which what enzymes and metabolites could be assayed were. Thus, these results were based upon what could be done, rather than a coherent hypothesis. The best examples are listed in Dugger and Ting (1970).

The latter two theories can be restated as a loss of productivity with three possible somewhat-independent causes — (a) a reduced production of the basic building blocks of growth and, hence, a slowing of growth in at least one organ; (b) a reduced ability to reproduce, leading to a decreased production of viable seeds or of fruits and nuts; and (c) a decreased ability to mount a defense against pathogens or insects, leading to weaker plants, which are more liable to be overcome by other stresses. It is important to separate out effects that may be detrimental or

1 disfiguring, such as the production of visible injury, but which have not been shown to lead
2 directly to a loss of productivity due to possible compensation by the remaining tissue. The loss
3 of some localized regions of photosynthesis tissue has not been shown to lead directly to a loss
4 of productivity.

6 **9.3.2.2 Modifications of Plant Physiological Processes**

7 The discussion that follows will focus on physiological processes; the species used to
8 develop an understanding of these processes are relatively unimportant. The study of any plant
9 that can help increase the biological understanding of the response of plants to O₃ exposures is of
10 use, regardless of how sensitive it is to O₃. Therefore, *Arabidopsis*, whose physiology and
11 genome continue to be studied and described by a large number of scientists is an appropriate
12 plant for studying O₃ injury. Though the responses of mature trees and understory plants are
13 critical to understanding plant interactions at an ecosystem level, the time required for trees to
14 reach maturity makes using them to study biological mechanisms a poor choice.

15 The high levels of O₃ used for some investigations do not automatically invalidate the
16 results obtained in those studies. Typically when a new hypothesis is being investigated,
17 extreme levels of the toxicant are used to clearly determine its effects. The older studies that
18 used concentrations as high 1 ppm, an extreme level, helped to define current studies. Later
19 experiments have used concentrations nearer ambient levels. Many of the current studies on
20 physiology use exposures between 0.15 and 0.25 ppm, which though higher than ambient levels
21 in some areas of the country, bypass confounding changes but allow for rapid experiments.

22 Three forms of air pollutant-induced injury patterns currently are known to exist: (a) acute
23 stress, generated by high atmospheric concentrations of pollutants for short periods of time;
24 (b) chronic stress, generated by lower concentrations of pollutants for long periods of time; and
25 (c) accelerated senescence, generated by very low concentrations of pollutants for very long
26 periods of time. At higher levels, distinct visible injury generally occurs due to cellular and
27 tissue death of regions of leaf mesophyll cells. This leads to a decline in the total area of
28 metabolically active tissue, with consequent loss of membrane integrity, loss of metabolites into
29 the extracellular tissue space, and formation of oxidative products. When no visible injury is
30 observed, lowered rates of photosynthesis or productivity are often used to index injury. Under
31 these conditions, metabolism is altered and the pool sizes of many metabolites are changed.

1 More importantly, the altered biochemical states within the tissue lead to the inability of the
2 plant to respond properly to existing environmental conditions and to other stressors (Manning
3 and Keane, 1988; Heath, 1988; 1994b; Koziol and Whatley, 1984; Schulte-Hosted et al., 1988).
4

5 **9.3.3 Ozone Uptake by Leaves**

6 Plants respond to O₃ similarly to other stressors on several levels: exclusion, tolerance, and
7 repair (Levitt, 1972; Tingey and Taylor, 1982). The response mechanism depends upon the O₃
8 concentration, environmental conditions, and the developmental and metabolic state of the plant
9 (Guzy and Heath, 1993). These responses are detrimental to plant productivity because they cost
10 the plant metabolic resources. In some cases, the stomata close under the O₃ exposure,
11 excluding the pollutant from the leaf interior and preventing injury. However, if this happens
12 too often, CO₂ fixation is also inhibited and plant productivity suffers.

13 Atmospheric O₃ does not cause injury, but rather it is the O₃ that enters the plant that
14 causes an effect (Tingey and Taylor, 1982). Three well-defined, sequential processes control the
15 movement of O₃ from the atmosphere into the sites of action within the leaf and must occur to
16 trigger O₃ stress (Heath, 1980). The processes are: (1) entry of O₃ into the leaf, (2) reactions of
17 O₃ and its possible reaction product(s) in the water phase at cell surfaces, and (3) movement of
18 O₃ reaction product(s) into the cell with enzymatic or chemical transformation of those products
19 in the cell.
20

21 *Process 1. Entry of O₃ into the leaf.* Often incorrectly, the external concentration of O₃ is
22 used to give an indication of “dose” (Heath, 1994a). Ozone-induced changes on the plants
23 cuticle are minimal, and O₃ does not penetrate the cuticle (Kerstiens and Lenzian, 1989) to
24 cause an effect. As O₃ has no easily measured isotope, virtually no measurements have been
25 done on an actual dose of O₃, i.e., the amount of O₃ which reacts with individual biochemicals in
26 the leaf. Yet the measurement of dose will be the amount of O₃ expected to penetrate into the
27 tissue through the stomata. Dose is expressed as a rate of delivery to a surface area (mol/m² s⁻¹).
28 Whether dose or total accumulation (mole/m², rate integrated over exposure time) is most critical
29 for the development of injury remains a major question.

30 Ozone uptake includes gaseous diffusion through the leaf boundary layer and stomata into
31 the substomatal cavity (Figure 9-3). Although the movement of pollutants through a boundary




Figure 9-3. The uptake of O₃ into the leaf. Each of the individual concentration layers of O₃ represents a different process of movement and of plant/microenvironmental interaction. This figure leads into Table 1, in which the amounts of O₃ along the pathway are calculated.

layer into the stomata region is known to be important, and even rate limiting in many cases of low wind velocity, its description has been defined from aeronautical concepts and usually relates to smooth surfaces that are not typical of leaf-surface morphology; however, it is nearly the only treatment available (Gates, 1968). Once through the boundary layer, the gas must enter the leaf through the stomata. The entry of gases into a leaf is dependent upon the physical and chemical processes of gas phase and surfaces and is a well defined path that approximately follows a linear flux law of:

$$j = g(C_o - C_i) \quad (9-1)$$

where the flux, j , into the internal space of a leaf is related to the conductance, g , through the boundary layer and stomata and the gradient of concentration of gas from the outside, C_o , inwards, C_i . This formulation has been used for years for both water and CO₂ (Figure 9-4), and for regions of varied CO₂ concentration that correspond to C_o (CO₂ of the atmospheric air, below the leaf proper) and C_i (CO₂ near the leaf's spongy mesophyll cells) (Farquar and Sharkey, 1982; Ball, 1987).

In the past, the internal concentration of O₃ has been assumed to be zero (Laisk et al., 1989), due to early studies that found that virtually no O₃ could pass through a leaf. That was expected because O₃ is extremely reactive with cellular biochemicals. If the assumption that the




Figure 9-4. The microarchitecture of a dicot leaf. While details among species vary, the general overview remains the same. Light that drives photosynthesis generally falls upon the upper (adaxial) leaf surface. Carbon dioxide and O₃ enters through the stomata on the lower (abaxial) leaf surface, while water vapor exits through the stomata (transpiration).

internal concentration zero is correct, then the effective delivery rate for O₃ is given as $g \times C_o$, with stomatal conductance being the major regulatory control (Taylor, Jr. et al., 1982; Amiro et al., 1984). However, a recent study by Moldau and Bichele (2002) indicated that the internal O₃ concentration may not be zero, as previous assumed. Moldau and Bichele (2002) permitted leaves of *Phaseolus vulgaris* L., which have stomata on both upper and lower leaf surfaces, to take up O₃ at a high rate for 3 to 5 min. Exposure of the lower leaf surface resulted in up to 5% of the O₃ that was taken up to be diffused through the leaf, emerging from the stomata on the upper surface. This suggested above-zero concentrations of O₃ in the intercellular leaf air spaces. The descriptive calculations and plots of Moldau and Bichele (2002) indicate that the rise in internal O₃ level (for both concentrations of external O₃) within the first few minutes of exposure is due to its reaction with an antioxidant, most probably absorbate, within the apoplastic space of the leaf (Figure 9-5). The rate of rise is probably due to more complete penetration of O₃ with a concurrent depletion of the external antioxidant. The rise peaks at about 2 min for 0.88 ppm and 3 min for 0.34 ppm and then falls to a lower level. This may be due to a




Figure 9-5. The change in the O₃ concentration inside a leaf with time. Data are from O₃ exposures at two different concentrations.

Source: Derived from data in Moldau and Bichele (2002).

replenishment of the antioxidant. The authors saw no injury to the plasmalemma (as measured by penetration of a dye) and no change in the stomatal conductance for the lower concentration of O₃ (Moldan and Bichele, 2002). The higher level (0.88 ppm) caused the plasmalemma of the mesophyll cells to pass a dye and a slight decline in stomatal conductance resulted at about 2.5 minutes. These data suggest that the antioxidant hypothesis is correct.

Gaseous pollutants flow from the substomatal cavity within the leaf into the cell, through the cell wall. It is suspected that the internal concentration of the pollutant is not uniform within the cavity. From within the wall, an equilibrium between the gas and aqueous phase must occur at the interface where the gaseous species dissolve into the water according to Henry's Law (Heath, 1980, 1987; Wellburn, 1990). It is important to understand exactly how much O₃ could move into the tissue of the leaves. Calculations in Table 9-6 give an indication of the amount of O₃ which may end up near the surface of cells within the leaf. The calculation is done for a standard temperature (25 °C), an ambient concentration of O₃ (0.10 ppm), and for nonspecific

Table 9-6. The Flow of Ozone into a Leaf and Possible Reactions

The level of O₃ in the atmosphere is chosen to be close to a standard and yet make calculations to other amounts easy. The same concept will be used for all standard parameters for these calculations.

DESCRIPTION

VALUES

The atmospheric level of O₃ is given as:

$$O_{3a} = 0.1 \text{ ppm}$$

For an air temperature of:

$$T_a = 25 \text{ }^{\circ}\text{C}$$

The perfect gas law ($pV = nRT$) is used to convert the O₃ level into standard mks. Further, the volume for a mole of gas ($V_o = 22400 \text{ m}^3$) will be used, from the perfect gas law with $R = 8.3144$

Thus, the concentration of O₃ within the atmosphere is:

$$C_{O_3} = \frac{O_{3a} \times 10^{-6} \cdot (T_a + 273.18)}{V_o \times 273.18}$$

$$C_{O_3} = 4.873 \times 10^{-12} \text{ moles/m}^3$$

The stomatal conductance of the gas must be chosen to be standard but adjustable. The number should be as large as typically measured, but allow for easy conversion, if necessary. For a stomatal conductance of:

$$gs_{wv} = 1 \text{ cm/s}$$

The amount of O₃ that will penetrate inside the leaf (for a typical concentration of nearly zero inside the leaf), is:

$$O_{3L} = \sqrt{\frac{18}{48} \frac{gs_{wv}}{100}} C_{O_3}$$

$$O_{3L} = 2.984 \times 10^{-14} \text{ mol/(m}^2 \text{ s)}$$

In terms of amount of water within the leaf we can assume that about 85% of the weight is water and the density of water is 1 g/ml. A typical leaf has a wet weight/area:

$$FW_L = 30 \text{ mg/cm}^2$$

Thus, the square surface area of the leaf will translate into water space (for concentration of chemicals), as:

$$Ar_L = \frac{FW_L \times 0.85}{100}$$

$$Ar_L = 0.255 \text{ L/m}^2$$

The maximum amount of toxic compound that will be generated, assuming all the O₃ is converted, is given below. Here the units of the leaf area weight are converted into the mks system and the water space units are converted into L, such that the concentrations calculated will be in mol/(L hr). The final units assume that the O₃ is present (and no back reactions occur) for one hour (short but typical units of exposure).

$$O_{3Lc} = (O_{3L} / Ar_L) \times 3600$$

$$O_{3Lc} = 4.21 \times 10^{-10} \text{ mol/(L hr)}$$

Table 9-6 (cont'd). The Flow of Ozone into a Leaf and Possible Reactions

Thus, the maximum amount of toxic chemicals generated per hour in a leaf would be:

$$O_{3Le} = 0.42 \text{ nmol/(L hr)}$$

Possible errors in these calculations (aside from the input numbers) are: (1) the O_3 within the leaf does not react uniformly within the leaf space, (2) the O_3 within the leaf does not totally convert to any one species, (3) varied products of O_3 react leading to innocuous chemicals, and (4) O_3 reactions can be catalytic and generate more by radical reaction cycling.

leaves. For example, 0.3 ppm would be the same general numbers but multiplied by 3. Similarly for more closed stomata, the value of 1.0 cm/s (equivalent to about 400 μmole^{-2} -leaf area s^{-1}) for a conductance would be reduced and the smaller values would lead to a smaller amount of O_3 moving into the tissue. Nonetheless, these values give some indication of what sort of chemical concentration can be expected. Under these conditions, a delivery rate of O_3 into the substomatal cavity near the spongy mesophyll tissue of about 0.42 nmol/(L hr) appears to be reasonable.

Process 2. Ozone diffuses into the leaf air spaces and reacts either with varied biochemical compounds (path 1) which are exposed to the air or is solubilized into the water lining the cell wall of the air spaces (path 2). As shown in Figure 9-6, each reaction has the possibility of transforming O_3 into another chemical species (a toxicant) which, in turn, may react with other chemical species and lead to a cascade of reactions.

Within the stomata, gases react with the water at the cell's surface and generate new species with the components within the cell wall region. The possible varied pathways are depicted in Figure 9-7. Although these chemical reaction are poorly understood, some of the fundamentals are known (Heath, 1987; 1988; Wellburn, 1990). Ozone reacts with organic molecules at the double bonds to form carbonyl groups and, under certain circumstances, generates peroxides, such as hydrogen peroxides (H_2O_2), superoxide (O_2^-) and its protonated form (HO_2^*), hydroxyl radical (HO^*), and peroxy radical (HO_2^*). Other chemicals present in the water phase can lead to many other oxygenated moities (Figure 9-6). Each of the steps are generally pH dependent (Walcek et al., 1997; Jans and Hoigne, 2000).




Figure 9-6. Possible transformations of O₃ within a leaf.




Figure 9-7. Possible reactions of O₃ within water. (a) Ozone reacts at the double bonds to form carbonyl groups. (b) Under certain circumstances, peroxides are generated.

1 Sulfhydryls are particularly easy targets, with the formation of disulfide bridges or sulfones
2 (Mudd and Kozlowski, 1975). In water, the reactions become more confusing, but some
3 products have been described by Heath and Castillo (Heath and Castillo, 1987), such as H_2O_2 ,
4 HO^\bullet , and O_2^- (Figure 9-7). Effective detoxification reactions can occur here via antioxidant
5 metabolites and enzymes, such as ascorbate, glutathione (GSH), and superoxide dismutase
6 (SOD), if they are present at high enough concentrations (Matters and Scandalios., 1987;
7 Castillo et al., 1987; Fong and Naider, 2002). If the levels are low, it is believed that stimulation
8 of their production is a response to O_3 , albeit a slow one (Harris and Bailey-Serres, 1994).
9 Certainly it is possible that chemical modification of wall-specific biochemicals (Castillo et al.,
10 1987), such as glucan synthase (Ordin et al., 1969) and diamine oxidase (Peters et al., 1988), is
11 possible.

12 Process 3. Movement of reaction product(s) into the cell and enzymatic or chemical
13 *transformations within the cell.* It is believed that the initial site of O_3 injury is near and within
14 the plasma membrane. Certainly, membrane functions, such as membrane fluidity (Pauls and
15 Thompson, 1980), permeability (Elkiey and Ormrod, 1979), K^+ -exchange via ATPase reactions
16 (Dominy and Heath, 1985) and Ca^{2+} exclusion (Castillo and Heath, 1990), are changed. The
17 similarity of wounding responses (Langebartels et al., 1991) and O_3 -induced membrane
18 disruption suggests the induction of normal wound-regulated genes (Mehlhorn et al., 1991;
19 Sandermann, Jr., 1998). This implies that O_3 can react with components of the cell wall
20 connected to the cytoplasm through the cell wall and membrane by membrane-specific proteins
21 not directly linked to transport.

22 Ozone is soluble in water and once having entered the aqueous phase, it can be rapidly
23 altered to form oxidative products that can diffuse more readily into and through the cell and
24 react with many biochemicals. Again, the presence of internal antioxidant would be critical to
25 reduce the concentration of most oxidants. A toxic product of O_3 may migrate through the
26 cytoplasm to react with photosynthetic processes, or a spurious signal generated at the membrane
27 may affect some control process or signal transduction pathway (Schraudner et al., 1998;
28 Overmyer et al., 2000, 2003; Rao et al., 2000a, 2002; Sandermann, 2000; Rao and Davis, 2001;
29 Leitao et al., 2003; Vahala et al., 2003; Booker et al., 2004).

9.3.3.1 Possible Reactions Within the Leaf

Ozone can react with many compounds within the substomatal cavity of the leaf¹ to produce a variety of oxidizing and toxic chemicals. Some of the possible reactions which will generate H_2O_2 , HO^\bullet , and SO_2^- , as well as charged O_3 intermediates are indicated in Figure 9-8. Many of these complex reactions have been studied within water solutions through research of O_3 induced water purification and are very dependent upon solutes present with the solutions, including H^+ (see von Gunten (Von Gunten, 2003)). An important point is that in alkaline media, O_3 forms H_2O_2 , but in acid media O_3 is relatively stable in the absence of free metal ions.


The rates of reaction of O_3 with several important compounds, including those with a double bond, the so-called Crigee Mechanism shown in Figure 9-8, can be calculated from the reaction coefficient as given by Atkinson (1990) (Table 9-7). The double bond of the ascorbate molecule is particularly sensitive to O_3 attack. An unstable ozonide product is formed and then accelerates the breakage of the double bond, leading to the formation of two products because of the ring formation of the ascorbate molecule. These products are relatively unstable and can lead to further reactions not shown in Figure 9-8. The rates of reactions can be calculated (Heath, 1987) from the concentrations of O_3 that should occur within the wall regions (Table 9-6). At a local concentration of $25\text{ }\mu\text{M}$ O_3 , it would take 5000 s (83 min) for all of the O_3 to react if there was no further flow of O_3 . Clearly, O_3 does not react rapidly with the compounds in Table 9-7 and, although some of the products would be formed through the Crigee Mechanism (see Figure 9-8a), they would be low in concentration². While other radicals, such as hydroxyl radical (see Figure 9-8b) can attack double bonds, the products differ. Of particular note for later discussion, is the reaction of O_3 with ascorbate (see Mudd, 1996; Figure 9-8b), which will cleave the double bond in the ring. Unfortunately, little work has been done to characterize possible products within the leaf (but see next section).

In a paper discussing the stability and reactivity of O_3 in the pulmonary air/tissue boundary, Pryor (1992) calculates that O_3 has a half-life of about $7 \times 10^{-8}\text{ s}$ in a bilayer.

¹The volume of the substomatal cavity (that are within the leaf immediately below the stomata) must be regarded as the region in which most O_3 reactions occur. That volume, at a relative humidity of near 100%, possesses many diverse surfaces with varied bonding, which could alter the fate of O_3 .

²For example, hydroxymethyl hydroperoxide would be expected to be formed by the reaction of O_3 with ethylene and its effects have been tested on peroxidases (Polle and Junkermann, 1994). Unfortunately, the concentration of required for inhibition is much higher than would be expected to be formed within the leaf.

a.



b.




Figure 9-8a,b. The Crigee mechanism of O_3 attack of a double bond. (a) The typical Crigee mechanism is shown in which several reactions paths from the initial product is shown. (b) Typical reaction of ascorbic acid with O_3 .

Source: Adapted from Mudd (1996).

Table 9-7. Some Rates of Reaction of Ozone With Critical Biochemicals

[a] Double bond reactions. The second column is taken from Atkinson (1990) and transformed into Column 3. Those rate coefficients are used to calculate the rate of reaction at a concentration of 10 ppm for the organic and 0.1 ppm for O₃ in the air stream within the leaf (localized concentration of about 25 mM, see Table 9-6).

Compound	$\times 10^{-18} \text{ cm}^3/\text{molecules s}^{-1}$	Rate coefficient (L/mole s ⁻¹)	Rate of reaction (M/s)
Ethane	1.7	1.02×10^3	4.3×10^{-11}
Propene	11.3	6.80×10^3	2.8×10^{-10}
1-butene	11	5.91×10^3	2.5×10^{-10}
trans-2 Butene	200	1.20×10^5	5.0×10^{-9}
α -pinene	85	5.12×10^4	2.1×10^{-9}

[b] Possible Oxidative Species. Another possibility is given by the reactions below from Walcek et al. (1997).

Reactions	Rate constants
(1) $\text{O}_3 + \text{OH}^- + \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}_2 + \text{O}_2 + \text{OH}^-$	$k_1 = 3.67 \times 10 \text{ mole}^{-1} \text{ L s}^{-1}$
(2) $\text{O}_3 + \text{O}_2^- \rightarrow \text{HO} \cdot + 2 \text{O}_2 + \text{OH}^-$	$k_2 = 1.26 \times 10^9 \text{ mole}^{-1} \text{ L s}^{-1}$
(3) $\text{O}_3 + \text{HO}_2^- \rightarrow \text{HO} \cdot + \text{O}_2^- + \text{O}_2$	$k_3 = 2.09 \times 10^6 \text{ mole}^{-1} \text{ L s}^{-1}$

[c] Possible Concentrations of Other Oxidative Species. Table from Heath (1987). Based upon 100 ppm O₃ in gas stream

Species	Concentration (M)		Molecules within wall	
	pH 7	pH 9		
Superoxide Radical (O ₂ [•])	8.75×10^{-15}	1×10^{-12}	5.5×10^{-6}	6.3×10^{-4}
Ozone Radical	4.16×10^{-15}	5×10^{-14}		
Protonated O ₃ radical (HO ₃ [•])	1.48×10^{-16}	1×10^{-18}		

Number of molecules within apoplastic space of (10⁻¹² L) at 0.1 ppm O₃.

1 However, the transit time through the lung lining fluid layer is about 2×10^{-6} s based upon a
2 reasonable estimate for the diffusion of O₃. This means that O₃ would suffer nearly 29 half-
3 lives³ in passage through the layer, which would reduce O₃ to about 3×10^{-9} of the original
4 concentration — zero for all practical aspects. In the same publication, Pryor points out that any
5 sulfhydryl or ascorbate would interact strongly with O₃, further reducing its net concentration.

³Here a half-life is the time that it takes the reactive species to travel a distance in which it loses 50% of its initial concentration. Therefore for a 29 half-life, the concentration has been reduced by 2⁻²⁹ or about a 10⁻⁹ decline.

1 The reactivity of cysteine is 10^9 , while the reactivity of tryptophan, methionine, polyunsaturated
2 fatty acids, and tyrosine is about 2×10^6 , and that of phenylalanine is only 10^3 . These numbers
3 are similar to what has been found for O_3 reactivity with amino acids and proteins in aqueous
4 solutions. In glycophorin (Banerjee and Mudd, 1992) and cytochrome C (Mudd et al., 1996,
5 1997b) in aqueous solutions, only the methionine was oxidized by O_3 , producing sulfoxide.
6 In other proteins lacking methionine, tryptophans were oxidized only if they were in an exposed
7 position on the surface of the proteins (Mudd et al., 1997b). Treatment of red blood cell ghosts
8 with O_3 , oxidized peripheral proteins of the plasma membrane before it oxidized lipids (Mudd
9 et al., 1997a).

11 9.3.3.2 Toxicants Within the Wall Space

12 While Mehlhorn et al. (1990) is often thought to have shown that free radicals were formed
13 in plant leaves under O_3 exposure, careful reading of that paper clearly shows that there was no
14 real evidence of free radicals induced by O_3 . Living tissues have many free radical signals,
15 making it difficult to observe changes in free radicals. Further, the work of Grimes et al. (1983)
16 has also been cited as showing the presence of free radicals in living tissues due to O_3 exposure;
17 however, no radical signals were found unless certain organic acids (e.g., caffeic acid) were
18 added to the tissue with the O_3 exposure. They used the radical trap TMPO (tetramethylphrrolise
19 1-oxide) which reacts with many types of free radicals to form a stable radical that can be used
20 to “trap” or increase the amount of radical present (see Figure 9-9a). Ozone would directly react
21 with this trap only if it were bubbled into the solution, not passed over the top of the solution.
22 In the presence of sorbitol or caffeic acid, the trap would indicate the presence of OH radical,
23 which would mean that $O_3 \rightarrow HO^\bullet$. Superoxide dismutase, catalase, or EDU had no effect upon
24 this signal, suggesting O_2^- and H_2O_2 were not involved in the above sequence. Both O_3 and O_3
25 plus caffeic acid had no effect upon the protoplasts’ intactness or viability. Thus, 10^{-5} M HO^\bullet
26 and/or 0.30 to 0.40 ppm O_3 did not react with the cell membrane. They found no signal in
27 normal cells after subjecting the leaf to O_3 and concluded that the radicals were produced via a
28 concerted mechanism with the acid. This does not fit with the mechanism postulated by
29 Mehlhorn et al. (1991), which involved a reaction of wound-induced ethylene and O_3 at the wall
30 level to generate some free radicals.




Figure 9-9. Varied ESR radicals, trapped and not, generated by ozone under somewhat physiological conditions. (a) The generation of a DMPO-trapped radical with caffeic acid in water solution (Grimes et al., 1983). (b) The generation of a DMPO-trapped radical within bean (*Phaseolus vulgaris* cv. Pinto) exposed to 0.10 ppm O_3 for 4 hours. The lower trace is the ESR signal produced with 0.3 ppm O_3 (Mehlhorn et al., 1990). (c) The EPR signal produced within a bluegrass leaf exposed to ppm of O_3 for 1 h (Runeckles, 1997). Although no trapping agent was used in this experiment, the signal is complex because of various free radicals normally present within the illuminated leaf.

The hypothesis that the production of wound-induced ethylene by O_3 exposure and its reaction with O_3 would result in the production of radicals was tested by Mehlhorn et al. (1990), using electron paramagnetic resonance spectroscopy. After 4 h of 300 ppb, an EPR signal of a compound was detected which resembled a butonyl radical (Figure 9-9b). Using 70 ppb, the signal was reduced by about one third of an ethyl radical⁴, leading to injury. However, the spraying of the plant with 1-aminoethoxyvinyl-Gly (AVG), which reduces the production of

⁴The reaction would be: $O_3 + H_2C = CH_2 \rightarrow$ varied C-1 compounds, due to double bond cleavage, at a rate constant of $1.7 \times 10^{-18} \text{ cm}^3/\text{molecule sec} = 1.02 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ (Atkinson R., 1990). This should be compared with a reaction of the hydroxyl radical with ethylene, which has a rate constant of $8.52 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, or $10^6 \times$ faster.

1 ethylene and visible injury, had no effect upon the EPR signal, suggesting that the radical is not
2 on a direct sequela to visible injury.

3 Runeckles and Vaartnou (1997; Figure 9-9c) discovered a signal by subtracting other EPR
4 signals of the leaf, which seemed to be due to an O₃ reaction with plant material, using 0.48 ppm
5 O₃. This difference signal looked very much like O₂⁻. At a lower concentration, they observed
6 that this signal still occurred but accumulated more slowly. Both bluegrass and ryegrass leaves
7 seemed to saturate after about 5 h of exposure at 22 to 28 units of signal, while radish leaves
8 reached a maximum of 7 units at 3 h and then declined. The problem, which is typical of any of
9 these methods, was that the detached leaf had to be rolled and placed into the EPR detection
10 cavity. Reichenauer et al. (1998) also detected an undefined free radical signal that seemed to be
11 related to a Mn(II) spectrum. The Nandu and Perlo cultivars of wheat were more sensitive to O₃
12 than Extradur (according to growth rate and closure of stomata under an O₃ exposure of 80 ppb
13 for 8 h/day, 7 days/week over 100 days), and these more sensitive cultivars had a greater, but
14 insignificant (P = 15%) EPR signal. Thus, data showing any production of a free radical must be
15 approached with some skepticism.

16 With an O₃ delivery rate of about 25 µM/h (Table 9-6), only 250 µM would be found after
17 a full day, if all of the O₃ were stable. While the use of free radical traps is the best method
18 available to observe any build-up of radicals, the traps are not as specific to individual radicals.
19 Currently, studies should be looking for hydroxyl radicals, superoxide, hydroperoxides, ethylene
20 radicals, and ascorbate radicals.

22 **9.3.3.3 Products of Ozone**

23 Ozone should reach a certain concentration in the substomatal cavity, which is dependent
24 upon its entry speed and its reactivity with the wall constituents. Once near the apoplastic space,
25 O₃ moves in two different pathways (Figure 9-6). It can react with constituents that are within
26 the wall as a gas in reaction 1A (path 1); or it can solubilize into a water space and travel to
27 another region within the water space and react through reaction 2A (path 2).

29 ***Hydrogen Peroxide***

30 Hydrogen peroxide, until recently was thought to be purely a toxic compound for cells.
31 However, it is now clear that it functions as a signaling molecule in plants and mediates

1 responses to abiotic and biotic stressors (Figure 9-10). Generation of H_2O_2 is increased in
2 response to various stressors, implicating it as key factor mediating the phenomena of
3 acclimation and cross tolerance, in which exposure to one stressor can induce tolerance of
4 subsequent exposure to the same or different stresses (Neill et al., 2002). The signaling response
5 to attack by invading pathogens using H_2O_2 has been described (Mehdy, 1994; Simon-Plas et al.,
6 1997). The reactions leading to hypersensitive cell death are caused by a pathogen recognition
7 step (Figure 9-10a), probably due to the plant cell wall releasing oligosaccharides in response to
8 the pathogen enzymatically breaking down the cell wall to penetrate it. A feed-forward step in
9 which H_2O_2 increases the level of benzoic acid leads to the activation of the hydroxylase step in
10 the production of salicylic acid and to a feed-back step in which the salicylic acid increases the
11 production of H_2O_2 (Leon et al., 1995).

12 An elicitor, e.g., a bacterial or fungal pathogen, induces a cascade of reactions within a cell
13 (Figure 9-10b). Some of the lipid reactions are thought to be due to the opening of the Ca^{2+}
14 channels and the alkalination of the cell wall region. The oxidative burst due to H_2O_2 production
15 is believed to lead to the transformation of a small population of lipids into jasmonic acid, which
16 is a secondary messenger.

17 Hydrogen peroxide also has an oxidative role in lignification (Schopfer, 1994). In the
18 interaction of lignification and the beginning processes of hypersensitivity, pectinase produced
19 by the pathogen disrupts pectin and dissolves the cell wall. Fragments of the dissolved cell wall
20 trigger an increase in the transcription of peroxidases within the remaining cell wall, leading to
21 lignification, which is a cross-linking of the cell wall that does not use pectin. This prevents
22 further pathogen disruption of the wall and reduces its further entry into the plant cell.

23 It is believed that the first species generated through a one-electron reduction of molecular
24 oxygen is SO_2^- . That generation is carried out using a cytochrome b_6 by the NAD(P)H oxidase
25 located on the cell membrane (Auh and Murphy, 1995). In the acid region of the cell wall, SO_2^-
26 is converted by a protonation and dismutation to H_2O_2 . The induced oxidative burst is believed
27 to play a role in stimulating the Cl^- and K^+ efflux, generating an alkalization of the
28 extracellular space (Cazale et al., 1998). In the wall region, H_2O_2 is not especially toxic, as no
29 necrosis was reported in tobacco when 500 mM peroxide was infiltrated into the leaf tissue.
30 However, the production of salicylic acid and benzoic 2-hydroxylase can be induced with only
31 30 and 0.3 mM H_2O_2 respectively, indicating some metabolic signaling (Leon et al., 1995).




Figure 9-10. Pathogen-Induced Hypersensitivity. (a) The reactions leading to hypersensitive cell death and the formation of a global response of salicylic acid. (b) The cascade of the elicitor-induced reactions within the cell.

- 1 On the other hand, 1M H₂O₂ infiltrated into soybean will generate lipid peroxidation after 1 h
- 2 with a peroxidation rate of 15 nmol/g-FW h (Degousee et al., 1994). Cells react to the system⁵

⁵Soybean suspension cells were inoculated with *Pseudomonas syringae* pv *syringae*, which generate an active oxygen response. Light emission by luminol, reacting with H₂O₂, was the assay for the peroxide.

1 and generate peroxide scavenging compounds within 1.5 to 2 hours, which appear to “mop up”
2 the excess H_2O_2 (Baker et al., 1995).

3 After O_3 exposure in birch, H_2O_2 has been found in the wall (Pellinen et al., 1999).
4 By using CeCl_2 as a cellular stain for H_2O_2 (as a cerium perhydroxide precipitate), Liu et al.
5 (1995) observed a gradual development of stain after 8 h of O_3 exposure (at 150 ppb). After 2 h
6 exposure, H_2O_2 stain was visible on the surfaces of both sets of mesophyll cells. Accumulation
7 of H_2O_2 stain continued for 16 h after exposure, suggesting a triggered-reaction rather than O_3
8 decomposition itself. H_2O_2 stain was present in the mitochondria, peroxisomes, and cytoplasm,
9 but not in the chloroplast. If methyl viologen (MV) was given to the leaves and then the leaves
10 were exposed to light, H_2O_2 stain could be observed within the chloroplast. This indicated that
11 the stain worked within the chloroplast if H_2O_2 were generated by the Mehler reaction
12 ($\text{MV} + \text{O}_2^-$). Thus, apparently, for birch, O_3 exposure does not generate excess H_2O_2 within the
13 chloroplast. Furthermore, these sets of experiments indicate that O_3 *per se* does not generate the
14 H_2O_2 , but rather triggers stress-related H_2O_2 formation similar to what occurs in a pathogen
15 attack (the Reactive Oxidative Species or ROS reaction).

16 The presence of higher than normal levels of H_2O_2 within the apoplastic space is a potential
17 trigger for the normal, well-studied pathogen defense pathway. Figure 9-10b depicts such a
18 pathway and suggests that all the events and activation of pathways/genes caused by pathogen
19 defense could be observed when plants are fumigated with O_3 . These events in Figure 9-10b will
20 be alluded to in later sections.

21 H_2O_2 has been linked to the hormone ABA-induced closure of the stomata by activating the
22 calcium influx in guard cells (Pell et al., 2000). The addition of H_2O_2 at a level of only 5 mM to
23 a guard cell preparation will cause a dramatic increase (ca. 9×) in electrical current at the
24 hyperpolarizing potential of -200 mV. Amounts as low 50 mM H_2O_2 will cause a less but still
25 sizable increase. Membrane stability is unaffected by the H_2O_2 and the activation of the channel
26 requires only about 2 to 3 minutes. Pell et al. (2000) also found that ABA induced the
27 production of H_2O_2 through ROS accumulation (also see Zhang et al., 2001).

28 Certain levels of ABA within the leaf lead to stomatal closure. The inactivation of a
29 phospho-tyrosine-specific protein phosphatase (ABI2) is an inhibitor of stomatal opening
30 induced by ABA but that enzyme is inhibited by H_2O_2 (Meinhard et al., 2002). This means that
31 H_2O_2 shifts the sensitivity of the stomatal opening to ABA (Figure 9-11), making the stomatal




Figure 9-11. The interaction of H_2O_2 and Ca^{2+} movements with ABA-induced stomatal closure. It is well known that certain levels of ABA within the leaf lead to closure of the stomata within the leaf. That level, however, can be shifted to make closure more or less sensitive to a given level of ABA. Recently it has been shown that H_2O_2 (externally or produced by the plant) within the cell wall region can shift that sensitivity. Here ABA stimulates the production of H_2O_2 , which in turn increases the rate of Ca^{2+} moving from the wall region into the cytoplasm. That shift in internal Ca^{2+} level increases the closure of the stomata. Hydrogen peroxide also blocks the activation of a polypeptide (ABI2) that inhibits stomatal closure seemingly induced by ABA.

Source: From Assmann (2003); Assmann and Wang (2001); and Zhang et al. (2001).

complex more sensitive to ABA. Thus, for a given level of ABA present in the guard cell complex due to environmental factors (e.g., low humidity, high air temperature, or low soil water potential), the generation of H_2O_2 would (by inhibiting ABI2) induce a closure of the stomata by increasing the sensitivity of the guard cells to ABA. In the past, it has been difficult to understand why O_3 would often decrease conductance in some cases, but not always (Heath, 1994b). This interaction between H_2O_2 and ABA could help understand this complexity.

Ethylene Reactivity

Ethylene (ET) is produced when plants are subjected to biotic stressors e.g., attacks by insects, fungi, and bacteria or abiotic stressors such as wounding or environmental stressors such as heat, cold, or oxidative stress and O_3 . If an O_3 stress has induced a wounding response with ET release, then ET within the substomatal cavity could react with O_3 , generating some relatively noxious chemicals (see Figure 9-6). The relationship between O_3 injury and wounding

1 is supported by the observation by Mehlhorn et al. (1991) that an inhibitor of ET formation,
2 AVG (an inhibitor of ACC synthase, a committed step to ET production), would block ET
3 formation and inhibit visible injury. Other studies with polyamine (which is closely linked to
4 ET production), including those of Ormrod and Beckerson (1986) who fed polyamines to the
5 transpirational stream and prevented visible injury, suggested a close involvement of both
6 pathways to the production of visible injury. Both the lack of ET production or an increased
7 level of polyamines slowed or prevented visible injury.

8 This concept was taken another step by Langerbartles (Langebartels et al., 1991). The
9 linkage to the pathogen wound responses and visible injury is well established (Sandermann,
10 1996). Sandermann (1998) used a system of Bell B and W3 tobacco, plants with differential O₃
11 sensitivities, in which the O₃ exposure level was chosen such that the sensitive cultivar was
12 injured, while the tolerant one was not. This led to a marvelous control which could be used to
13 their advantage. They followed a time sequence to show that the rise of varied systems followed
14 the same order as for a pathogen attack (Heath, 1994a).

15 More recent studies, however, indicate that O₃ responses resemble components of the
16 hypersensitive response (HR) observed in incompatible plant-pathogen interactions (Sanderman
17 et al., 1998). The similarity to the HR response may be related to the occurrence of ROS in the
18 apoplast. The O₃-derived ROS apparently trigger an oxidative burst in the affected cells by an as
19 yet unknown mechanism. An oxidative burst is similar to one of the earliest responses of plants
20 to microbial pathogens and is an integral component in HR-related cell death (Overmyer et al.,
21 2000).

22 In plants exposed to O₃, ET synthesis is a result of the specific ET induction of the genes
23 encoding 1-aminocyclopropane-1-carboxylase synthase (ACS), one of the fastest and most
24 obvious responses to O₃, which has been mechanistically linked to the regulation of O₃ lesion
25 formation. Biosynthesis of ET inhibited with ACS inhibitors significantly reduced the induction
26 of lesion formation in plant leaves exposed to O₃ concentration (Vahala et al., 2003; Melhorn
27 and Welburn, 1987; Mehlhorn et al., 1991). Ethylene biosynthesis correlates best with O₃
28 (Vahala et al., 2003; Overmyer, 2000). These data support the concept that elimination of ET
29 formation will prevent visible injury.
30

Ethylene-Interaction with Injury and Conductance

Increased ET production by plants exposed to O₃ stress was identified as a consistent marker for O₃ exposure decades ago (Tingey et al., 1976). They exposed more than 20 plant species and cultivars to O₃ to determine whether the production of O₃-induced stress-ET could be used to determine differences in plant sensitivity to O₃. Their studies suggested that increased production of stress-ET correlated well with the degree of foliar injury that developed within hours or days after O₃ exposure. The amount of ET released was exponentially related to the O₃ exposure. Furthermore, the amount of O₃-induced ET declined with repeated exposure, indicating an acclimatization to O₃. This acclimatization effect associated with repeated wounding has not yet been well described. The release of wound-induced ET is not linear with time, but declines after the initial response (Stan et al., 1981), as is also seen after O₃ exposure (Stan and Schicker, 1982). The stress-induced ET production correlates better with O₃ exposure level than with exposure duration. In other words, peaks of high O₃ (rather than accumulated dose) generate a higher rate of ET release, at least for a single O₃ exposure under an acute dose.

The production of ET after an O₃ exposure is thought to be a typical wounding response (Tingey et al., 1975). Prevention of ET release may prevent the formation of visible injury (Mehlhorn and Wellburn, 1987). However, the question arises as to whether this effect was limited to the prevention of visible injury or if the chemicals used to prevent ET release closed the stomata. Using *Glycine max* L., Taylor et al. (1988b) showed clearly that AVG did not necessarily close stomata nor inhibit carbon assimilation *per se*.

The correlation of ET release with O₃-induced visible injury was likewise shown in pea cultivars (Dijak and Ormrod, 1982). With O₃ exposure (generally 6 h at 0.3 ppm), the stomata closed by ~50% within 3 hours after a dose of 3×10^{-5} mol cm⁻² (with an average rate of 2×10^{-9} moles cm⁻² s⁻¹, as calculated from their data). Both sensitive and insensitive cultivars had a visible-linked-injury ET release, but sensitive cultivars scored higher both in visible injury and in ET release after a given exposure.

Gunderson and Taylor (1988, 1991) used exogenous ET to alter the gas exchange of *Glycine max* and found an exponential, but not simultaneous, decline of both stomatal conductance and carbon assimilation with ET. Interestingly, the exogenous ET caused a slight rise in difference of CO₂ within and without the leaf, indicating a lowering of internal CO₂, which was not observed in the experiments of Farage et al. (1991) for O₃ exposure. Ethylene

1 does inhibit both stomatal conductance and carbon assimilation to some extent (Taylor et al.,
2 1988b). Thus, one could postulate that O₃ generates a wounding response with a production of
3 ET, which would, in turn, generate the change in stomatal conductance and photosynthesis.
4 Clearly, these multiple events may have confounded some earlier studies.

6 **9.3.3.4 Antioxidants Within the Apoplastic Space**

7 The first line of defense against O₃ is a closure of the stomata to exclude its uptake. This is
8 counter-productive for efficient photosynthesis, but some amount of closure limits the rate of O₃
9 deposition into the leaf tissue to allow for a secondary line of defense to detoxify the O₃. The
10 secondary line of defense involves a range of antioxidants, which are highly reactive to the types
11 of chemicals that can be generated by O₃. Several antioxidant proteins are stimulated by O₃
12 in *Plumbagini folia*, including glutathione peroxidase (GSH-P_x), SOD, and catalase. The
13 timescales for changes in their levels vary: some rise rapidly, while others rise more slowly.
14 The pattern of changes in these particular proteins varies greatly among different species and
15 conditions.

17 ***Ascorbate Within the Cell Wall***

18 Most of the recent reports indicate that ascorbate within the cell wall is the real first line of
19 all defense. Ascorbate within the wall declines when the tissue is exposed to O₃ (Luwe et al.,
20 1993; Moldau, 1998; Turcsányi et al., 2000; Zheng et al., 2000). This decline appears to be
21 closely linked to the amount of O₃ penetrating the leaf tissue.

22 It has long been suspected that intracellular antioxidants play a role in preventing O₃-
23 induced injury to plant cells. Variation in the types of biochemical compounds present in the
24 apoplastic space can give rise to a multiplicity of reactions with O₃, but the predominant
25 biochemical species is ascorbate. Ascorbate is water-soluble, present in the solution where O₃
26 can dissolve, and is highly reactive. Unfortunately, a variety of antioxidants are found
27 throughout the cell and any measurements of one particular type within the total leaf tissue can
28 give misleading results. For example, ascorbate is present within the cell wall, within the
29 cytoplasm (Moldau, 1998; Burkey, 1999), and within the chloroplasts (Law et al., 1999); and
30 ascorbate can move between the cytoplasm and the cell wall with relative ease (Figure 9-12;
31 Bichelle et al, 2000). The total of all ascorbate pools are measured when the tissue is ground and




Figure 9-12. The reaction of ascorbate within the apoplasm of the cell wall and its ultimate reduction/oxidations.

- (a) Movements of reducing power (from Dietz, 1997).**
(b) The use of glutathione to maintain the level of ascorbate within the cell wall region (from Horemans et al., 2000).

1 assayed. If the cell wall ascorbate drops by 50% due to O₃ exposure but all other tissue
2 concentrations remain the same, the measurement of the total loss is dependent upon the amount
3 of ascorbate within the cell wall. Turcsányi et al. (2000) showed that, compared to the
4 concentration of apoplastic ascorbate, the rest of the cells contained about 38 times as much.
5 So a 50% loss of apoplastic ascorbate would be converted into only 2 to 3% loss of the total
6 ascorbate.

7 The ascorbate deficient *Arabidopsis thaliana* mutant has proven to be a powerful tool in
8 furthering the understanding of ascorbate biosynthesis in plants (Smirnoff et al., 2001). Three
9 classes of mutants were formed when *A. thaliana* seed was mutagenized with ethyl
10 methanesulfonate: (a) those deficient in SOD, (b) those that failed to accumulate more
11 antioxidant proteins upon increased O₃ exposure, and (c) those that were deficient (but not
12 depleted) in ascorbate. The low-ascorbate mutant type had 50 to 60% less ascorbate than the
13 wild type and displayed more foliar injury. This mutant is involved with the coding of the
14 GDP-D-Mannose pyrophosphorylase enzyme⁶ in the Smirnoff-Wheeler pathway for ascorbate
15 biosynthesis. Smirnoff et al. (2001) also suggested that other pathways can produce ascorbate
16 without relaying upon the pyrophosphorylase step, but most probably at a slower through-put
17 rate, because any fully ascorbate-deficient mutant would be lethal, perhaps because of ascorbate
18 use as a cofactor rather than its antioxidant properties.

19 The ascorbate peroxidase (APX, which uses ascorbate to detoxify peroxides) family
20 consists of at least five different isoforms, with isozymes in the apoplastic and cytosolic space.
21 Furthermore, most forms of ascorbate can move through the plasma membrane (Figure 9-12;
22 Bichele et al., 2000), making the levels of all forms of ascorbate interdependent and able to at
23 least partially influence each other. Dehydroascorbate (DHA) can be broken down into other
24 smaller fragments easily in vivo and represents a continuous loss of ascorbate from varied parts
25 of the cell if ascorbate is allowed to remain in the oxidized form in some regions. In fact, the
26 turnover rate in leaves is estimated to be from 2 to 13% / hour, depending upon species and
27 developmental age (Smirnoff et al., 2001). There are apparently three pathways for ascorbate

⁶ EC 2.7.7.13, Mannose-1 phosphate guanylyltransferase; mannose + GTP → GDP-mannose + ppi; this product leads into cell wall polysaccharide synthesis and protein glycosylation through GDP-galactose and GDP-fucose and ultimately, through Galactose, into ascorbate synthesis.

turnover (Figure 9-12a). The typical reaction is a reduction of DHA into ascorbate from which an oxidative step generates DHA. Pathway I requires a reductive step using NADH external to the plasma membrane generated from internal malate using a malate/oxaloacetate transporter. Pathway II uses a direct transporter of ascorbate/DHA. Pathway III moves the electron(s) required through a cytochrome b system, maintaining two separate pools of ascorbate/DHA within the cytoplasm and within the wall. Each of the pathways (Dietz, 1997) represented by Roman Numerals in the Figure 9-12, require only one NAD(P)H molecule to reduce the DHA molecule back to ascorbate. However, the transport properties and redox potential of the cell differ for each pathway. The efficiency of the reduction of DHA is dependent upon the redox coupling and the region in which the chemical species is located.

Turcsányi et al. (2000) exposed broad bean (*Vicia faba*) grown under two regimes in duplicate controlled chambers: charcoal/Purafil filtered air (CFA) or (CFA) plus 0.075 ppm O₃ for 7 h/day for 28 days (chronic exposure) or exposed to 0.150 ppm for 8 h (acute exposure). Responses of the two set of plants were similar except for stomatal conductance, which was 50% lower in the chronic exposure plants. Plants grown under acute exposures developed visible injury, while plants grown under chronic conditions developed no visible injury. Within an hour of the start of the acute exposure, the stomatal conductance was reduced by nearly 40% and assimilation was reduced by nearly 18% in the clean air plants; a reduction in conductance was only 21% and assimilation 16% in the plants subjected to chronic O₃ exposures. The assimilation was affected similarly in both cases, while the conductance showed less of a percentage drop in the chronic O₃-exposed plants, beginning at a lower O₃ level. The similarity of the assimilation indicated that the stomata were not limiting assimilation in either case before acute exposure. More to the point, the decline in ascorbate in the apoplastic space due to the O₃ exposure was "...more often than not, on the borderlines of statistical significance." However, a 30% decline in ascorbate after 4 hours of acute O₃ in both cases was observed. This lack of significance may be due to a relatively large standard error of the data, which in turn may be due to the difficulty of extracting and measuring ascorbate from the apoplastic space in quantitative terms.

1 The chemical reaction⁷ of ascorbate and O₃ is given by the molecular rate constant of
2 $4.8 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$. This is some 50,000× the rate constant for ET. Of course, it depends upon the
3 relative concentration of ascorbate and ET, but it is likely that ascorbate is in higher
4 concentration than ET. One would then expect that the rate reaction of ascorbate with O₃ would
5 greatly dominate any possible reaction of O₃ with ET. For a concentration of ascorbate in the
6 range of 1 mM and for an O₃ concentration of about 0.1 ppm or $4.2 \times 10^{-9} \text{ M}$, the detoxification
7 rate would be $4.8 \times 10^7 \times 10^{-3} \times 4.2 \times 10^{-9} \text{ M s}^{-1} = 2.0 \times 10^{-4} \text{ M s}^{-1}$. Turcsányi et al. (2000)
8 calculated an O₃ flux of about $1.6 \times 10^{-9} \text{ moles m}^{-2} \text{ sec}^{-1}$. With a wall thickness of $0.12 \times 10^{-6} \text{ m}$
9 and all the O₃ flux going into the wall region, this would give about $1.3 \times 10^{-2} \text{ mol m}^{-3} \text{ s}^{-1}$ or
10 $1.3 \times 10^{-5} \text{ M s}^{-1}$ flux, which is less than 10% the detoxification rate.

12 ***Glutathione***


13 Many of the initial studies of O₃ exposure used high concentrations and measured only the
14 total sulfhydryl contents of the tissues. For example, in some of the earlier work, exposures of
15 tobacco to 1 ppm O₃ for 30 min induced a 15% loss of the total sulfhydryls (0.74 μmole/g-FW;
16 Tomlinson and Rich, 1968). These results are similar to other studies at high O₃ levels (Dugger
17 and Ting, 1970). It is now suspected that severe injury in their studies resulted in a massive
18 collapse of the cells releasing most of their internal constituents. Much of the oxidation thus
19 observed may have been the result of chemical oxidations of the O₃ that subsequently entered the
20 damaged tissue. Even under milder conditions, changes in sulfhydryl components have still
21 been noted and any sulfhydryl on the surface of the cell would be at risk due to its high reactivity
22 with O₃ (Mudd et al., 1969; 1997a). For example, the level of sulfhydryl compounds within the
23 protein of isolated chloroplasts declined about 66% when the chloroplasts were subjected to O₃
24 (about 1 μmole O₃) exposure (Mudd et al., 1971).

25 At this stage, it is important to note that there are inherent problems with metabolic studies
26 of full tissues. The first is that most organs have several different types of tissues. For example,
27 leaves have, at the minimum, epidermal and vascular tissues and two types of mesophyll cells.
28 Each type of cell may be metabolizing quite differently, producing very different levels of

⁷These chemical rate constants are those constants within a bulk solution. In the apoplasm, the possibility exists for the chemicals to be preferentially oriented near a surface; so the constants may not be the same as for bulk solutions.

1 metabolites and enzymes. Furthermore, most pathways are well regulated and after any small
2 disruption, the pathway tends to return to near its former stability. Changes in the level of
3 enzymes are likewise difficult to measure. Many enzymes function below their maximum
4 activities. Their speeds of reactions are often increased through regulation, rather than through
5 the production of more enzyme.

6 Glutathione is a three-amino acid peptide, which has antioxidant properties due to its free
7 reducing sulfhydryl group (G-SH). Glutathione is generally kept in its reduced form by
8 glutathione reductase (GR) with the reaction:
9



10
11 GR has six isoforms⁸ within the chloroplast and six isoforms outside. The optimum activity
12 occurs at pH 7.8, suggesting it is located within the stroma of the chloroplast or the cytoplasm
13 rather than in the cell wall, which is at pH 4-5 (Madamanchi et al. (1992). Clearly, an increased
14 expression of GR (generated through transgenic implants) is important within the chloroplast to
15 prevent of some oxidations⁹ (Aono et al., 1995; Foyer et al., 1995).


16 17 *Catalase*

18 Catalase, even though it breaks down H₂O₂, does not appear to protect plants from O₃
19 exposures. Two principle reasons may cause this lack of reactivity: (1) catalase has a high K_m
20 for H₂O₂ and a low rate coefficient and (2) seems not to occur within the cell wall regions but
21 rather in the cytoplasm and peroxisomes (Buchanan et al., 2000). Only a few reports suggest
22 that catalase is increased by exposure to O₃ (Azevedo et al., 1998). Unfortunately H₂O₂ induced
23 by some forms of wounding in mesophyll cells can lead to induction of an increase in GSH and
24 the transient production of catalase (Vanacker et al., 2000). In general, it seems that catalase is

⁸An isoform is the same enzyme, with the same structure and perhaps within the same organelle, but its promoter region has different DNA codes. Thus, each protein segment is induced by different signals, and so its enzyme can be formed in response to different environments. This is in contrast to isozymes, which classically are similarly reacting, but structurally different, enzymes in different compartments.

⁹Typically this protection is observed in the paraquat sensitivity of plants. In this assay, added paraquat, the herbicide which intercepts electrons from the reducing end of photosystem I in the chloroplast, caused oxidations, chlorophyll loss, and death due to the buildup of superoxide and peroxides.

not really involved primarily in the defense of the cell due to O₃ attack but rather may be a secondary response. The reaction of catalase (Scandalios, 1993) is as follows:



Superoxide Dismutase

The varied compounds that O₃ can produce upon entering an aqueous solution are very similar to those involved in the HR when plants are infected by an avirulent pathogen (Figure 9-10). The sequence of the plant response to the pathogen is: (1) recognition of the gene products of the pathogen by the plant (elicitor), (2) generation of an immediate phytoresponse to attempt to localize the attack and its products, and (3) generation of a systemic acquired resistance (SAR) to subsequent attack by the pathogen. Inducible defense responses are phytoalexin synthesis and production of pathogenesis-related proteins (PR). One aspect of this total response is the production of O₂⁻ and H₂O₂ by the cell (Lamb and Dixon, 1997). The elicitor can generate a transient alkalinization of the apoplast, up to pH 7.2, caused by a lowering of the H⁺-pump rate and a increase in the H⁺-influx/K⁺-efflux exchange. Other effects include: a weak accumulation of transcripts for PAL (phenylalanine lyase); a larger and rapid induction of glutathione S-transferase, GSH-P_x; oxidative cross linking of cell wall proteins which is blocked by ascorbate acid; generation of localized apoptosis; and rapid influx of Ca²⁺, which activates apoptosis among other pathways (Lamb and Dixon, 1997). These effects seem to be very similar to those induced by O₃ exposure (Sandermann, 1996; 1998).

The putative antioxidant enzyme SOD (Equation 9-4 and Table 9-8) catalyzes the oxidoreductase reaction, which eliminates SO₂⁻ by dismutation (Bowler et al., 1992):



The number, as well as the activity, of isozymes of each type of SOD in Table 9-3 can vary with plant species. However, the isozymes that have been tabulated are Cu-Zn SODs, in cytosol and chloroplast; Fe-SOD, active in chloroplast stroma, and Mn-SOD, active in mitochondrial

Table 9-8. Superoxide Dismutase Isozymes and Isoforms

Reaction: $2\text{H}^+ + 2\text{O}_2^- \rightarrow \text{H}_2\text{O}_2 + \text{O}_2$			
Isozymes	M.W.	Isoforms	Cytolocation
Cu-Zn	20 kDa	<i>csd1</i>	
		<i>csd2</i>	Plastid
		<i>csd3</i>	Peroxisomal
Fe	23 kDa	<i>fsd1</i>	Mitochondrial
		<i>sd2</i>	
		<i>fsd3</i>	Plastid
Mn	23 kDa	<i>msd1</i>	Mitochondrial

matrix (Karpinski et al., 1993). In the experiment demonstrating the activation of varied SODs, there were three Cu-Zn SOD (*csd1*, *csd2*, *csd3*), three Fe-SODs (*fsd1*, *fsd2*, *fsd3*), and one Mn-SOD (*msd1*) (Kliebenstein et al., 1998). Ozone sensitivity was determined by exposure of plants to 8 h of 0.33 ppm of O₃. *csd1* induced by O₃ and UV-B was one of the earliest SOD increases and most pronounced responses for mRNA and protein. Also, some increase in *csd3* (thought to be peroxisomal) was induced when the plants were exposed to a high-intensity light pulse; *msd1* was unresponsive to the environmental stressors used here, including O₃; and *csd2* (thought to be chloroplast) showed little increase. The *fsd1* isozyme (present in the apoplasm) showed a slight decrease. On the other hand, an early report on snap beans in which the experimenters used EDU, N-[2-(2-oxo-1-imidazolidinyl)ethyl]-N-phenylurea (Carnahan et al., 1978; Kostka-Rick and Manning, 1993) to prevent visible injury by O₃, 4 h O₃ exposure at 0.45 ppm was correlated with an increase in general enzyme activity of SOD, i.e., the level rose nearly 2.5× in 2 weeks at a level of 50 mg EDU per pot (Lee and Bennett, 1982). It is believed that EDU may induce SOD, which then protects the plant. While gross assays of enzyme activity have not proven to be very useful in understanding the mechanism of O₃ action, in a well-crafted, long-term study involving ponderosa pine clones. Benes et al. (1995) stated that “changes in antioxidant enzyme activity were not a consistent response to the O₃ fumigation, but when observed, they occurred most often in the O₃-sensitive clone and in symptomatic,

1 fumigated branches...total (intra- and extracellular) activities of the antioxidant enzymes did not
2 appear to be good indicators of O₃ tolerance....”

3 Ozone exposure (70 ppb for 7 h/day for 14 to 42 days of exposure)¹⁰ caused an increase in
4 POD and a decline in SOD with no change in APX. No GSH was detected, but the
5 concentration of (ASC + DHA) was at 20 to 25 nmol/g-FW of extracellular fluid, compared with
6 2.4 to 3.0 mmol/g-FW of cell fluid. Glutathione within the cell was only 100 to 170 nmol/g-FW
7 of cell fluid. While these results are what one might expect for POD, the decline in SOD and
8 lack of change in APX are not what would be expected if protection was provided by SOD and
9 ascorbate. Yet as noted, because the rate of SOD reaction is many times higher than the rate of
10 O₃ entry, there may be no pressure to increase the SOD level.

11 Some protection against visible injury (induced by 59 ppb daily mean O₃ for 14 h/day for
12 7 days) was observed in genetically modified tobacco plants with excess chloroplast SOD¹¹ (2 to
13 4 times higher), but less protection was observed in plants that had an excess of mitochondrial
14 SOD (8× higher) (Van Camp et al., 1994). In all lines, the conductance of the leaves dropped
15 about 50%, compared with the unmodified plants. There was a correlation with age of leaf (less
16 injury in younger leaves) that corresponded to that found in spruce trees in which the amount of
17 SOD declined in relation to the longer that needles were held on the tree (Polle et al., 1989).
18 A slightly different study, however, found no O₃ protection with varied SOD within the needles
19 (Polle and Rennenberg, 1991). Interestingly, in maize, the synthesis of SOD (any form) was not
20 stimulated by O₃ exposure (at 0.50, or 0.75 ppm for 8 h, variable times thereafter) but was by
21 exposure to 90% O₃ (Matters and Scandalios, 1987). It may be that this high level of O₃ does not
22 affect the SOD, or perhaps it stimulates and degrades the enzyme simultaneously.

23 The conclusions to be drawn from these results are not obvious. There seems to be SOD (a
24 Cu-Zn form) present in the apoplastic space of some plants, but it does not necessarily rise with
25 O₃ exposure. Thus, either its concentration is sufficient to provide protection or it is not needed.
26 Over-expression of any SOD in other organelles may play a role, especially in the chloroplast

¹⁰At a level of 70 ppb, the concentration of O₃ in air was about 3.06×10^{-6} mol/m³, which with the conductance of 0.042 mol/m² s, gives a flux rate of O₃ of 1.27×10^{-8} mole/m² s. Converting the SOD rate of 23 units/g-FW into a SOD rate within the apoplastic space of 6.9×10^{-3} mol/m² s, or about 500,00 times the entry rate of O₃.

¹¹The SOD enzymes were from *Nicotiana plumbaginifolia* with appropriate transit sequence for targeting the correct organelle and expressed under control of cauliflower mosaic virus 35S promoter.

(Cu-Zn or Fe forms); but it may be playing a secondary role due to other effects of O₃ that generate conditions in which light can overload the chloroplast and generate detrimental circumstances, including the production of SO₂⁻. In addition, SOD is developmentally expressed in varied concentrations, so that long term exposure to O₃ may alter each leaf's developmental age and, in turn, alter what level of SOD is observed. In any case, SOD does not seem to be the primary antioxidant system to protect against O₃.

Changes to the Plasmalemma

Reports of “peroxidation” generally occur within unicellular organisms subjected to very high levels of O₃ (in *Chlorella* [Frederick and Heath, 1970]) and in *Euglena* [Chevrier et al., 1990]). Heath (1987) determined that by the time biochemical events were altered and MDA was produced in *Chlorella*, little permeability remained in the cells and most metabolic pathways were greatly disrupted by the subsequent loss of substrates. In fact, MDA production was concurrent with a high O₃ uptake by the cell, indicating a complete opening of the cell and associated with the concurrent inability to plate the cells on a glucose median (indicative of cell death). Heath reached the conclusion that no one had proven that lipid oxidation was in any way a part of the initial reactions of O₃ with the cell, a conclusion confirmed by Mudd et al. (1997a). An excellent review regarding the initial action induced by O₃ within a plant (Kangasjärvi et al., 1994) should be consulted. There is little data to show that lipids are attacked by O₃ in any living system that was not previously severely injured by O₃. Most of the data suggesting lipid attack by O₃ has been demonstrated in plants subjected to O₃ concentrations of 0.5 to 1.0 ppm for several hours, during which gross wilting of the plant tissues usually occurs, suggesting extreme water loss. It is not surprising that lipid and protein injury is observed under these conditions. While those reports were useful in the 1960s and 1970s, they are not especially insightful now when ambient levels of O₃ are rarely above 0.2 ppm.

9.3.4 Wounding and Pathogen Attack

The decline of an enzyme is more difficult to measure than the rise of a new enzyme; an increase from 0 to 2% may be within the precision of any assay, but a decrease from 100% to 98% is often masked by simple variation of the assay. Thus, measuring enzymes, which are in great abundance in pre-fumigated tissue, is a risky operation. On the other hand, if O₃ induces a

1 general physiological change that has characteristics similar to other well-studied, stress-induced
2 changes, then O₃ studies could “piggy-back” onto those studies to gain insight into the full scope
3 of metabolic alterations. It is now becoming clear that wounding and pathogen attack of plants
4 are similar to O₃-induced changes in plants, and a reasonable hypothesis is that O₃ must induce
5 one or more of the first steps seen in the wounding/pathogen-attack response.

6 Systemic acquired resistance (SAR) has been heavily investigated, and DNA probes have
7 existed for some time for a series of expressed genes (see Table 9-9). Several enzyme classes
8 are associated with O₃ injury, including glucanases and peroxidases and others, such as the PR
9 proteins and chitinases. Thus, strong evidence exists from enzyme function and genetic material
10 that O₃ induces an activation of a SAR-like response.

11 Mehdy (1994) described a model of how an elicitor produced by the pathogen attack
12 activates a G-protein, which opens the inward-flowing Ca²⁺ channel. The flow of Ca²⁺ into the
13 cytoplasm raises the internal level (at the μM level) and activates a protein kinase that increases
14 the activity of the plasma membrane NAD(P)H oxidase and generates O₂⁻. Superoxide
15 dismutase converts O₂⁻ into H₂O₂. Both O₂⁻ and H₂O₂ are responsible for the active oxygen
16 species response, which is believed to be a defense mechanism to kill the pathogen. In this
17 normal defensive reaction, a subsequent system induces either localized lipid peroxidation per se
18 or a membrane lipase to produce jasmonic acid or inositol triphosphate, which act as secondary
19 messages to activate the defense gene products.

20 Booker et al. (2004) found that G-proteins might be involved in the perception of O₃ in the
21 extracellular region using *A. thaliana* G-protein null mutants. The activation of a passive inward
22 flow of Ca²⁺, e.g., by an O₃-induced response, would serve the same function as activation of the
23 G-protein. Once the level of cytoplasmic Ca²⁺ rises, all else follows. It is suspected that
24 exposure of plants to O₃ does just that, as Castillo and Heath (1990) demonstrated — the in vivo
25 fumigation of bean plants both inhibits the outward-directed ATP-requiring Ca²⁺ pump and
26 increases the passive permeability of Ca²⁺. It was thought that the calcium transporter system
27 has a sensitive sulfhydryl group which, if oxidized, would alter normal Ca²⁺ movements. In
28 addition, Dominy and Heath (1985) observed that the K⁺-activated ATPase (believed to be
29 involved in K⁺ transport) was inactivated by in vivo exposure to O₃ and that inactivation was
30 traced to a sensitivity sulfhydryl. Mudd et al. (1996) argued that several amino acids are very
31 sensitive to O₃, including any with an exposed sulfhydryl. Thus, the O₃-induced change in Ca²⁺

Table 9-9. Gene Families and cDNA Clones Used as Probes for SAR (Ward et al., 1991)

Probe	Relevant Properties of Encoded Protein	Reference
PR-1	Acidic, extracellular; function unknown most abundant PR protein in tobacco; >90% identical to PR-1b and PR-1c	Payne et al. (1988b)
PR-2	Acidic, extracellular b-1,3-glucanase, >90% identical to PR-N and PR-O	Ward et al. (1991)
PR-3	Acidic, extracellular chitinase; also known as PR-O; >90% identical to PR-P	Payne et al. (1990a)
PR-4	Acidic, extracellular; unknown function; homologous to C-terminal domain of Win1 and Win2 of potato	Friedrich et al. (1991)
PR-5	Acidic, extracellular; homologous to thaumatin and bifunctional amylase/proteinase inhibitor of maize; also known as PR-R or PR-S	Payne et al. (1988a)
PR-1 basic	Basic isoform of acidic PR-1	Payne et al. (1989)
Basic class III chitinase	Homologous to cucumber chitinase (Metraux et al., 1989), structurally unrelated to PR-3	Lawton et al. (manuscript in preparation)
Acidic class III chitinase	Extracellular; approximately 60% identical to basic isoform	Lawton et al. (manuscript in preparation)
PR-O'	Acidic, extracellular b-1,3-glucanase; approximately 55% identical to PR-2 group	Payne et al. (1990b)
Basic, glucanase	Vacuolar; approximately 55% identical to PR-2 group and PR-O'	Shinshi et al. (1988)
Basic chitinase	Vacuolar; approximately 65% identical to PR-3 group	Shinshi et al. (1987)
SAR 8.2	Unknown function; cloned by \pm screen of cDNA library from secondary leaves of TMV-infected plants	Alexander et al. (manuscript in preparation)
Acidic peroxidase	Extracellular; lignin-forming	Lagrimini and Rothstein (1987)

permeability may be the trigger to most, if not all, the wounding responses. However, the difficult problem of proving that the cytoplasmic Ca^{2+} change is the first event in O_3 injury remains.

Some wound- and pathogen-induced genes that are activated or repressed in *Arabidopsis thaliana* are found with DNA arrays (Cheong et al., 2002). While these responses may not be uniform for all plants, they suggest the possibility of wide-ranging gene changes that may occur with a simple wound and that those changes are wide-ranging and diverse. As an example, these responses are related to hormonal responses that are related to jasmonic acid, ET, and auxin

pathways; signal transduction responses; and transcription factors for a variety of pathways.
The involvement of ET in wounding and pathogen attacks is discussed in Section 9.4.3.2.

9.3.4.1 Peroxidases

Increases in cytosolic and apoplastic peroxidase activity in response to O₃ are often observed, but the reasons and outcomes of these changes have yet to be fully explained. Increased activity is frequently correlated with O₃ injury. Dass and Weaver (1972) observed that increases in peroxidase after O₃ injury was similar to that observed for plant infection by a virus. Tingey et al. (1975) observed a 35% decrease in peroxidase activity immediately following O₃ exposure; however, within 24 to 48 h, activity had increased significantly and was above control level and remained there throughout the remainder of the study. Dijak and Ormrod (1982) also observed increases in peroxidase activity when two O₃-sensitive and two O₃-resistant varieties of garden peas (*Pisum sativum*) were exposed to O₃. Peroxidase activity was not related to cultivar sensitivity nor to visible injury. Unfortunately, there are many peroxidases (Birecka et al., 1976) therefore, any general increase is not specific. In ET-treated leaves, peroxidase reaction products were found between the plasma membrane and the cell wall, suggesting that ET itself could induce peroxidase activity (Abeles et al., 1989a,b; Birecka et al., 1976).

At the same time, others examined peroxidase reactions in general and found two types of peroxidases (designated as acidic or anionic and basic or cationic, EC 1.11.1.7, but also listed as EC 1.14.18.1). Many types of peroxidases are located in diverse organelles, and each seems to be activated by different conditions (e.g., pH for anionic and cation types and substrates such as guaiacol, syringaldazine, and ascorbate). Peroxidases belong to at least two groups, which catalyze two separate reactions: (1) the reaction of H₂O₂ with ascorbate to form DHA, discussed earlier (Thom and Maretzki, 1985), which is regenerated by plasma membrane electron transport using a dehydrogenase (Gross and Janse, 1977) that is now believed to be a malate/oxaloacetate shuttle through the membrane coupled to a NAD(P)H-cytochrome-b-reductoxidase; and (2) the reaction with coniferyl alcohol (from phenylalanine through phenylalanine ammonia lyase) to form lignin within the wall. The anionic peroxidase thought to be involved with lignification is within the cell wall (Buchanan et al., 2000; Taiz and Zeiger, 2002). Some basic peroxidases are maintained within the cell, while some are external to the cell. After wounding (Lagrimini and Rothstein, 1987; Gasper et al., 1985), some basic peroxidases can be activated by

1 processes leading to the synthesis of stress ET (Yang and Hoffman, 1984) and/or by excess Ca^{2+}
2 (Gasper et al., 1985). Elicitor treatment of plants change a series of peroxidases, some of which
3 are similar to those seen in O_3 -induced changes (see Table 9-9).

4 The formation of lignin is due to the phenylpropanoid metabolism (Buchanan et al., 2000).
5 Tyrosine and phenylalanine are converted to cinnamic and *p*-coumaric acid, which are in turn
6 converted to *p*-coumaryl, coniferyl, and sinapyl alcohols, and then into lignins. Hence, the
7 peroxidase activity is often measured by one of these substrates (Espelie et al., 1986; Gasper
8 et al., 1985). However, it is questionable whether apoplastic peroxidase activity is limiting for
9 lignification; laccases have a prominent role as well. Also, availability of monolignols is critical
10 for core lignin formation, and it is unclear whether levels of these metabolites change in
11 response to O_3 . Studies by Booker (Booker et al., 1991, 1996; Booker and Miller, 1998; Booker,
12 2000) indicated that O_3 did not increase core lignin concentrations in foliage of loblolly pine,
13 soybean, or cotton; although levels of phenolic polymers and cell wall-bound phenolics were
14 elevated in soybean. Increased phenolic polymers appear to be lignin in acid-insoluble lignin
15 assays and may well be responsible, along with polyphenol oxidase, for the stippling injury
16 observed in O_3 -treated plants. Cell wall function implies the transport of peroxidase molecules
17 out of the cell and, most likely, the regulation of their activities within the wall space. These
18 extracellular peroxidases may be observed by vacuum infiltration of buffer into leaf air spaces
19 and subsequent centrifugation of the tissues to remove the buffer with the apoplastic enzymes,
20 that wash out (Castillo and Greppin, 1986). However exposure to O_3 induces important changes
21 in the plant. For example, extracellular peroxidase activity in *Sedum album* leaves increased
22 nearly 3-fold over that in the control plants after 2-h exposure to 0.40 ppm (Castillo et al, 1984).
23 This O_3 -induced increase of extracellular peroxidase appears to be under the control of Ca^{2+}
24 (Castillo et al., 1984; Heath and Castillo, 1987). Initially, no effect on the anionic activity as
25 measured with syringaldazine (specific electron donor for lignifying peroxidases) was observed,
26 yet 21 hours later, the anionic peroxidase activity was increased, whereas the cationic (ascorbate
27 measured) peroxidase activity was decreased in O_3 -treated plants. This suggests an immediate
28 response (ascorbate peroxidase activation) and a secondary response that activates the lignifying
29 peroxidase via gene activation.

30 The rapid response of cationic peroxidase after O_3 exposure may not result from de novo
31 protein synthesis but from the secretion and direct activation by Ca^{2+} ions of enzyme molecules

1 already present in the tissue. Cationic peroxidases might attack the peroxides and, in this
2 manner, act as a detoxifying agent with ascorbate as the substrate in the apoplasm. The effect of
3 Ca^{2+} upon peroxidase activity is stronger at low H_2O_2 concentrations (Penel, 1986). Thus, one
4 can imagine that, when the H_2O_2 concentration is low, this peroxidase activation would have a
5 greater in vivo importance. Furthermore, the secretion of cationic peroxidases into the free
6 spaces as a result of O_3 treatment is accompanied by a simultaneous release of at least one of its
7 natural substrates (ascorbic acid); this cationic peroxidase exhibits a much higher affinity
8 towards ascorbate (up to 6-fold) than the anionic isozyme (Castillo and Greppin, 1986).

9 10 **9.3.4.2 Jasmonic Acid and Salicylic Acid**

11 Salicylic acid (SA) and jasmonic acid (JA) are considered to be regulators of the plant
12 defense response (Figure 9-13) (Buchanan et al., 2000). They tend to respond more slowly than
13 ET, causing widespread effects in the plant tissues. Both seem to be heavily involved in
14 responses of the plant to O_3 , once again linking the pathogen/wounding defense to O_3 -induced
15 injury; however, their roles are far from clear.

16 One of the lipoxidase isoforms is activated by pathogen infection (POTLX-3) within 6 h
17 and accumulates for a week (Kolomiets et al., 2000). This enzyme is the first stage of the JA
18 pathway which leads to 13-hydroperoxide linolenic acid (HPOT) which is converted either to
19 allene oxide through AOS or to C6 aldehydes through hydroperoxide lyase. These aldehydes act
20 as signaling agents via systemin (Sivasankar et al., 2000) or volatile odiferous compounds
21 (oxylipins) that have been implicated as antimicrobial toxins (Froehlich et al., 2001).
22 Interestingly, these compounds seem to target the chloroplast envelop where they interact with
23 its metabolism. As HPOT and AOs are both implicated in plant defense and are activated by O_3 ,
24 these interactions may be related to how chloroplast enzymes and their mRNAs are involved in
25 O_3 -induced injury.

26 27 **9.3.4.3 Stress-Induced Alterations in Gene Expression**

28 Early studies addressed the qualitative and quantitative effects of O_3 on protein metabolism
29 (Harris and Bailey-Serres, 1994). Subsequent reports suggested that the physiologic and
30 metabolic consequences of exposure to O_3 was, in part, mediated by increased gene expression.
31 A summary of the gene-linked changes in proteins induced by SAR may be seen in Table 9-9.




Figure 9-13. The pathway leading from phospholipids to jasmonic and traumatic acid. The role of lipoxygenase and the production of a hydroperoxyl moiety from the unsaturated fatty acid is clearly demonstrated. More importantly, several of the enzymes within this pathway have been shown to be activated by oxidative conditions including O₃ exposure. The production of both of these acid species could lead to a general global response of a whole plant to the O₃ exposure of a single leaf.

Source: Howe et al. (2000); Croft et al. (1993); Buchanan et al. (2000).

Of particular note are the productions of PR proteins, chitinase, glucanase, and acidic peroxidases that appear to be common markers used in many O₃ studies. A summary of varied proteins as measured by changes in the mRNA in *A. thaliana* induced by O₃ exposure is shown in Table 9-10. While studies on *Arabidopsis thaliana* required high concentrations of O₃ to produce a response, the levels reported in most of the studies did not induce visible injury. The types of messages induced included glutathione S-transferase, PAL, ACC synthase, SOD, and some PRs. Slower increases in messages are seen for other PR and SAR-senescence proteins. Declines in messages were observed for varied chloroplast enzymes, including those for Rubisco and chlorophyll binding proteins. A few new proteins were found — a casein kinase and three plasma membrane proteins. It is interesting to note that few messages for “new” proteins were generated by O₃ exposure.

Table 9-10. Proteins Altered by Ozone as Measured by Molecular Biological Techniques as mRNA Level or Other Gene Activity Rather than Enzyme Activity

Exposure	Physiological events	Identified proteins fast increase response	Slow increase response	Decline	Examined, but no change	Unknown proteins	Reference
150/300 ppb for 6 h daily	Leaf curling; reduced growth	GST, PAL	Pxase, SOD		CAT, LOX1		Sharma (1994)
300 ppb for 6 h daily	10 bands of 10 RNA					AtOZI1 >> casein kinase II	Sharma (1995)
200/500/1000 ppb for 2 h	Wilting (8 h); premature senescence					3 plasma membrane proteins: 75-, 45-, 35-kDa peptides	Tokarska-Schlattner (1997)
350 ppb for 1-6 h	Ethylene production; downward curvature; water logging	ACS-6			ACS-1, -2, -4, -5		Vahala (1998)
300 ppb for 6 h	Necrosis in NahG and Cvi-0 (accumulating SA), not in Col-0	Chl SOD, cytAPX, GST1	Chl GPX	Cab mRNA, cyto SOD, chl GR			Rao (1999)
150 ppm for 6 h/8 and 14 days	Downward rolling of leaf; early senescence		BCB, ERD1, SAG21	Cab, <i>rbcs</i>	Atgsr2, MT1, SAG 12, SAG 13, SAG 19, SAG 20		Miller (1999)
160 ppb for 3-72 h	Early senescence	GST1, VSP2;	MT1		CCH		Mira (2002)
(a) 250 ppb for 8 h; (b) 250 ppb for 2 h; (c) 175 ppb for 8 h/4 days	(a) little chlorosis or lesions; (c) growth retardation	GST Apx, CuZn-SOD	PAT1	Fe-SODI GR, cab, rbs			Conklin (1995)

Table 9-10 (cont'd). Proteins Altered by Ozone as Measured by Molecular Biological Techniques as mRNA Level or Other Gene Activity Rather than Enzyme Activity

Exposure	Physiological events	Identified proteins fast increase response	Slow increase response	Decline	Examined, but no change	Unknown proteins	Reference
200 ppb for 24 h		PR-1, PR-2a, PR-5 AtEDS1, AtGST1, AtGST2	PR-3b, PR-4		LOX2, AtOZI1, PAL, Lhcb, PAT1, HSP		Matsuyama (2002)
250 ppb for 6 h	Lesion initiated on margin and spread inward					rcd1, on chromosome 1, single Mendelian trait	Overmyer (2000)

Abbreviations used in Tables 5 and 6.

- GST = Glutathionine synthase

PAL = Phenylalanine ligase

PR-1 =

Pxase = Peroxidase

CAT = Catalase

LOX1 = Lipxygenase

ACS-6 = ACC synthatase

SOD = Superoxide dismutase

CuZn-SOD = cyto SOD

Fe-SODI = Chl SOD
- cyt APX = Ascorbate peroxidase

Chl GPX = Glutathione peroxidase

Cab mRNA = Chlorophyll a/b binding protein

chl GR = Gluthatione reductase

BCB =

ERD1 = Ethylene response

SAG21 = Senescence

rbcS = Rubisco small subunit

MT1 = Mitochondria

The working hypothesis is that O_3 , which is not eliminated by antioxidants in the cell wall, alters the properties of the plasma membrane. Specific polypeptides, indicative of these antioxidants, are induced. If specific receptor molecules or channels on the membrane are affected, the ionic balance within the cytoplasm is changed, leading to altered transcription or translation of the genes controlling those and other types of polypeptides. Once this membrane disruption occurs, the cell must mobilize repair systems to overcome the injury. Thus, carbon and energy sources once destined for productivity, must be used in repair processes. Some of these repairs are thought to result from the induction of specific genes. Photosynthesis is inhibited by direct inhibition of some of the enzymes, through byproducts of O_3 attack or by altered ionic balance. At the very least, the decrease in photosynthesis is a result of an O_3 -induced decrease in *rbcS* mRNA.

9.3.5 Primary Assimilation by Photosynthesis

9.3.5.1 Photooxidation: Light Reactions

Photooxidation refers to the oxidation of chlorophyll within the light reaction due to an imbalance between light absorption and the CO_2 use to produce carbohydrates. It was discovered in the 1920s and studied under the concept of chlorophyll bleaching and photo autooxidation (Asada, 1999; Rabinowitch, 1945). What generally occurs is that electron transfer from H_2O to NADPH declines, and a light reaction overload occurs. The slowdown of electron transfer may also be due to inhibition of the dark reactions, through the poor use of small molecular weight carbohydrates or a lowered amount of the fundamental substrate CO_2 . To counteract these detrimental reactions, a series of “antioxidant” reactions exist, which eliminate the buildup of oxidative intermediates.

A lowered CO_2 level, which can be caused by stomatal closure (Heath, 1996), blocks the use of reduced plastoquinone (PQH_2) in Photosystem II through NADP reduction in Photosystem I (Hankamer et al., 1997). The buildup of PQH_2 reduces the amount of Q_A , resulting in a buildup of $P_{680}^+|Pheo^-$ species (the primary photoact). The inability to reduce this radical leads to injury to the D_1 protein (32 kDa) and its fragmentation into 23-, 16-, and 10-kDa fragments (Hankamer et al., 1997). Ozone exposure of bean plants leads directly to the loss of this D_1 protein (Pino et al., 1995). The loss of D_1 stimulates the production of new D_1 (and its mRNA). Also, the production of the oxidized form of P_{680} (P_{680}^+) is harmful to the plant because

electron flow from water to P_{680}^{+} is limited, generating a P_{680}^T (the triplet form of P_{680}), which is highly oxidizing and can lead to dangerous reactions. One form of protection is the use of β -carotene to convert the triplet form back to its normal state; however, that reaction can lead to the loss of β -carotene. Without the protection of β -carotene, oxygen with oxidized products to produces singlet state of O_2 . This, in turn, can react with chlorophyll, leading to ring breakage that, in essence, leads to chlorosis. These types of reactions do not seem to occur often, but chlorosis is one form of visible injury, and loss of β -carotene has been reported. Farage et al. (1991) and Farage and Long (1999) studied these reactions in wheat and concluded that alterations to the dark reactions were much more common.

9.3.6 Alteration of Rubisco by Ozone: Dark Reactions

A large body of literature shows that O_3 exposure induces a decline in Rubisco (Pell et al., 1997). Treatment of a variety of plants with O_3 at near ambient levels results in a loss of Rubisco and of the mRNA coding for both subunits of Rubisco (*rbcS*, small and *rbcL*, large). Because Rubisco plays such an important role in the production of carbohydrates (Figure 9-14), any loss may have severe consequences for the plant's productivity.

The study by Noormets et al. (2001) used an exposure system of plants in a FACE exposure facility in which areas of ambient (daytime 360 ppm) and ambient with added CO_2 (560 ppm), with added O_3 (97.8 ppb), and with added CO_2 and O_3 , were used. Two clones of aspen (O_3 -tolerant and -sensitive) were used in all four cases. The study recognized that leaf age was important in the variation and tried to control for it by grouping the data by leaf plastochron index¹². These data, taken over many days using the LiCOR 6400, confirmed earlier observations that O_3 has the greatest effect on older leaves after causing a decline in assimilation and in conductance. They also confirmed that the internal CO_2 (calculated for within the leaf) is not affected by O_3 exposure. Higher levels of CO_2 increased the assimilation and lowered the conductance, maintaining the internal to external CO_2 ratio identical to that found with the ambient CO_2 level, corresponding to the theory of Farquhar et al. (1980). The level of Rubisco was not measured as frequently as the assimilation and conductance, but the significant (5%) increase in Rubisco was only observed in older leaves for both clones. More to the point was

¹²The plastochron index for the leaf measures the leaf age by degree of expansion rather than simply by chronological time.




Figure 9-14. The production of Rubisco and its Calvin Cycle pathway reactions. Two peptides are used to build Rubisco: *rbcS*, the small subunit produced by DNA within the nucleus; and *rbcL*, the large subunit produced by DNA within the chloroplast itself. Clearly both polypeptides must be closely regulated to produce the enzyme in a coherent manner. Furthermore, at least five isoforms of DNA can produce *rbcS*, each of which is regulated by a different promoter region.

that the stomatal limitation¹³ was not altered by O₃ exposure, with or without excess CO₂. It is critical to point out that mesophyll conductance is directly linked to internal CO₂ level¹⁴. So if C_i/C_o is constant and g_s declines, then g_m must likewise decline. If, as it is argued, Rubisco levels are constant or at least increasing, then a regeneration of RuBP must be the cause of the decline in g_m. Farquhar et al. (1980) were more concerned with high levels of CO₂ and had little

¹³The limitation was defined as the ratio of stomatal resistance to the total resistance (which included the operating point of the assimilation (A) versus internal CO₂ concentration (C_i) curve and the resistance of the boundary layer). The operating point of the curve was defined as the internal CO₂ level (which is calculated by the conductance and assimilation). The resistance of this operating point was calculated as the cotangent of the slope to the operating point. Unfortunately, the slope is not a dimensionless parameter but is rather moles of air per area of leaf ⁻²s of time and, thus, it is unclear whether the slope changes with added CO₂ and O₃.

¹⁴Respiration is generally small at saturating A and often is ignored. By transforming {A = g_s (C_o - C_i)} into {A = g_s C_o - g_m C_o = (g_s - g_m) C_o} where g_m = g_s (C_i/C_o) or the mesophyll conductance in earlier literature.

1 to say about O₃ exposure. One thing is certain however, the use of clones using different aged
2 leaves is a preferred method for this type of investigation.

3 The level of carbohydrate within the cell has an effect upon the amount of mRNA for
4 Rubisco (*rbcS*). Experiments by Krapp et al. (1993) indicated that a decline in carbohydrate
5 levels is probably due to the increased production of control metabolites, such as fructose
6 2,6-bisphosphate, which can shut down important sugar production pathways. This report also
7 leads to a measure of half-time for the decline in *rbcS* of about 2 days¹⁵ when 50 mM glucose is
8 added to a cell suspension of *Chenopodium*. Also, the carbohydrate level was increased by cold
9 girdling the petioles of tobacco and potato using intact plants. The levels of carbohydrate nearly
10 doubled in 5 days and the level of *rbcS* declined rapidly (reaching 25% after 12 h). A decline in
11 Rubisco followed, but more slowly (with an estimated half-time of about 108 h after a lag of at
12 least 12 hours). This, of course, is expected; the level of the enzyme would decline slowly with
13 a lag after a loss of the message.

14 A better estimation of the half-life of *rbcS* can be found in the Jiang et al. (1993) study of
15 the destabilization of the message by an antisense message. The wild type *rbcS* in tobacco had a
16 half-life of about 5 h compared to that in the mutant with the antisense. It was argued that the
17 antisense message increased the degradation of the normal *rbcS*. The estimated half-life of *rbcS*
18 under O₃ fumigation is about 1 hour (Pell et al., 1994). Although comparisons of these diverse
19 systems can not be easily made, the normal half-life of *rbcS* may be closer to 5 to 10 h; and O₃
20 fumigation does not simply stop the transcription of DNA, but rather it alters the rate of
21 degradation, either independently of, or simultaneously with, transcription.

22 Williams et al. (1994) developed a correlation between the levels of ABA after water stress
23 in *Arabidopsis thaliana* leaves and the loss of *rbcS*. Although their data were not quantitative, the
24 level of ABA had a half-time rise of about 1 to 2 h and the level of *rbcS* had a half-life decline of
25 about 2 to 4 hours. Their work suggests that drought stress may alter the CO₂ metabolism by
26 changing enzyme relationships much more than by merely closing the stomata. If an ABA rise
27 is lowering *rbcS*, *rbcS* may not be a good marker of O₃ fumigation except under highly
28 controlled conditions.

¹⁵The amount of Rubisco drops from an initial 0.12 to a final amount of 0.04 μ mole/g-FW s in 6 days.

9.3.7 Carbohydrate Transformations and Allocation

The question of whether translocation of the sugars out of the leaf is inhibited by O₃ exposure arises because productivity is often dramatically inhibited by O₃ fumigation. Though nearly 35 years have past since Dugger and Ting (1970) investigated the question of sugar transport within the leaf, the question has since been little studied. Translocation (Cooley and Manning, 1987) appears to be inhibited, because root functions are impaired by O₃ exposure. Many observed events suggest that while carbon assimilation within the leaf declines, translocation of carbon is inhibited even more so, because plant growth points are inhibited and root/shoot ratios are altered (Dugger and Ting, 1970; Gerant et al., 1996; Tjoelker et al., 1995).

Many of the experiments with O₃ fumigation indicate that O₃ exposure decreases the net growth or dry mass of the plant, but the mechanism is poorly understood. Generally the decrease in assimilation is much less than the decrease in growth, but not always. Under many conditions, the stomata will close partially, decreasing assimilation by a smaller factor. Only a long exposure, or high levels of exposure for a short time, generate enough decline in Rubisco to make the growth of the plant problematical. No convincing argument has linked the decrease in growth with a small decline in assimilation, either by a conductance- or Rubisco-limitation. Measures of assimilation with crops are frequently done on upper canopy leaves, which are the last leaves to exhibit O₃ injury, while leaves deeper in the canopy exhibit injury and early senescence. Crop root growth must be sensitive to these and other O₃ effects, because root biomass is often suppressed early by elevated O₃.

Volin et al. (1998) found O₃ exposures statistically decreased leaf area ratio, specific leaf area, leaf weight ratio, and root weight ratio in *Populus tremuloides* and two C3 grasses (*Agropyron smithii* and *Koeleria cristata*) but not in *Quercus rubra* and in the C4 grasses *Bouteloua curtipendula* and *Schizachyrium scoparium*. There was no statistically significant change in any species in leaf conductance (4% level decline in *K. cristata*) nor in assimilation (although there was a decline in assimilation at the 6% level for *P. tremuloides* and a decline at the 1% level in *B. curtipendula*). They also reported a correlation between growth decline and decreased stomatal conductance among all species.

Trade-offs are made by plants. Birch grown in highly fertilized conditions exhibited a greater leaf turnover when exposed to O₃, in that leaves not only formed faster but abscised faster, presumably due to early senescence; whereas birch grown under poorer fertilized

1 conditions retained their leaves longer and had a greater respiration rate within those leaves
2 (Maurer and Matyssek, 1997). Again, one must be careful in comparing short-term versus long-
3 term exposures. Grulke et al. (2001) observed that maximum concentrations of carbohydrates in
4 1-year old needles that had not abscised due to early senescence declined when subjected to
5 year-long exposures along an increasing pollution gradient. Furthermore, the monosaccharide
6 concentrations in fine roots (along with starch) were largely decreased, suggesting that needle
7 sugars were limiting, leading to root sugar limitations. However, determination of the total
8 productivity and detailed balance of carbohydrate was impossible, because these were older,
9 larger trees and the data were taken over a full growing season. For a shorter-term exposure of
10 9 days, Smeulders et al. (1995) observed that O₃ appeared to increase the retention of labeled
11 photosynthates within the needle, and, at higher exposures (400 versus 200 or 0 mg/m³), the total
12 starch within the needle decreased, suggesting that less carbohydrate was produced within the
13 cell or perhaps that it was in compounds not measured.

14 Studies with Pima cotton, aspen (*Populus* spp.) and bean seedlings (*Phaseolus vulgaris*)
15 indicate that acute O₃ exposures inhibit export of the current assimilate that provides
16 carbohydrates to the roots from source leaves of cotton as well as recent assimilate from the
17 older leaves of aspen and bean (Grantz and Yang, 2000). Grantz and Yuan (2000) attempted to
18 distinguish between potential mechanisms of O₃ phytotoxicity operating at the level of the whole
19 plant. Four hypothesis were tested by fumigating cotton: (1) O₃ exposure reduces leaf pools of
20 soluble sugars; (2) pruning leaf area and reducing source strength to match that of O₃-treated
21 plants reproduces O₃ effects; (3) pruning lower leaf area more closely reproduces O₃ effects than
22 pruning the upper leaf area; and (4) manipulating plant age and, thereby, plant size to match
23 O₃-treated plants reproduces O₃ effects. All were shown to be incorrect. Under each of the
24 above conditions, Grantz and Yang (2000) reduced the amount of foliage to match that caused
25 by O₃ injury. While the treatments reduced the biomass and leaf area, they did not alter biomass
26 allocation nor root function. They concluded that a simple loss of foliage does not induce the
27 changes in translocation to the roots to the same extent as does O₃ injury.

28 This finding by Grantz and Yang (2000) is important in that it suggests that O₃ can trigger
29 a plant-wide response that may be linked to alterations in signal transduction and the generation
30 of whole plant signals. Stitt (1996) suggested that "...allocation is regulated by long-distance
31 signals that act to influence growth of selected sinks and to modify the delivery of resources to

these sinks in parallel.” Cooley and Manning (1987), citing Mc Laughlin and McConathy (1983), suggested three possible ways that O₃ fumigation might alter translocation: (1) malfunction of the phloem loading process, (2) increased allocation to leaf injury repair, and (3) an altered balance between the leaf and sinks caused by reduced carbon fixation and a greater demand for assimilate in the leaf.

Ethylene has been shown to reverse this sugar inhibition of development and to be antagonistic to the ABA effect (Finkelstein and Gibson, 2002). However, these effects depend greatly upon the developmental stage of the plant. Thus, the balance of the effectors (sugars, ABA, and ET) may interact to generate the variation observed in the O₃-induced productivity decline. For example, O₃ fumigation can induce a shift in the carbon transfer between roots and shoot; and this shift can be amplified by mild drought (Gerant et al., 1996). Furthermore a regulation of source-sink relations with the defense responses induced by elicitors was observed by wounding the leaves of *Chenopodium rubrum*. Ethylene appears to be able to repress the expression of extracellular invertase, which is critical for control and down-loading of sucrose derived from the translocational stream (Roitsch, 1999; Lindow et al., 1996). In addition, the development of *Arabidopsis* at high concentrations of glucose or sucrose is arrested by increasing the ABA level (Coruzzi and Zhou, 2001).

Clearly more work is needed on the interactions between assimilation, translocation, and source/sink relations with O₃ exposure. In these interactions, one must be aware of the developmental age of the plants and their phytohormonal status.

9.3.7.1 Lipid Synthesis

Heath (1984) summarized several early reports of O₃-exposure induced lipid alterations. Most concerned the production of MDA as a measure of lipid oxidation as well as the loss of unsaturated fatty acids. However, a series of experiments by Sakaki and coworkers concentrated on one type of fumigation system and one metabolic pathway. This literature provides the best, most complete story with regard to lipid metabolism and O₃ fumigation and indicators that O₃ injures cellular membrane systems via lipid destruction.

Sakaki and coworkers used spinach, which is a sensitive plant but which has not been much evaluated with respect to O₃ fumigation. While the O₃ level was high (0.5 ppm), enough work has been done to be able to “tease apart” what is happening. The first paper showed that

1 chlorophyll bleaching does not begin until the plants have been exposed to O₃ for over 10 h,
2 whereas some MDA production begins with as little as 6 h exposure (Sakaki et al., 1983).
3 Consistent production of MDA, indicative of gross disruptions, occurred only after 8 h exposure
4 (Sakaki et al., 1985), within the timescale when chlorophyll and carotenoid levels began to
5 decline. Concurrently, the total fatty acid (FA) level decreased from ~481 to 358 nmol/cm² as
6 the MDA level increased from 0.6 to 2.4 nmol/cm², indicating FA peroxidation (Sakaki et al.,
7 1985).

8 Sakaki et al. (1983) also studied development of changes by cutting disks from exposed
9 leaves and floating them on water solutions for varied time periods (up to 24 h). This permitted
10 feeding experiments to be done easily, whereas the cutting gives rise to an additional wound
11 response and eliminates metabolite movement to and from other portions of the plant. The
12 floating experiments indicated that, after exposure, scavengers of singlet oxygen (¹O₂), such as
13 D₂O, and of hydroxyl radicals, such as benzoate and formate, have no effect on development of
14 the MDA response after 8 h of in vivo fumigation, while scavengers of (O₂⁻), such as tiron and
15 ascorbate, lowered the amount of MDA formed. By measuring metabolites immediately after
16 cessation of fumigation, they were able to show that ascorbate loss began with the onset of
17 fumigation, as did SOD loss. A lag time associated with the production of DHA suggested that
18 the reaction of ascorbate with fumigation did not immediately produce the oxidation product.
19 The first 4 h of exposure yielded 30 nmole/cm² of ascorbate loss with 5 nmole/cm² of DHA
20 production, whereas the second 4 h of exposure yielded 20 nmole/cm² of ascorbate loss with
21 20 nmole/cm² of DHA production.

22 Nouchi and Toyama (1988) exposed Japanese morning glory (*Ipomea nil*) and kidney bean
23 (*Phaseolus vulgaris*) to 0.15 ppm O₃ for 8 h. Under these conditions, little visible injury was
24 found with up to 4 h exposures, while injury increased by ~50% after 8 h of exposure. Morning
25 glory produced more MDA than kidney bean, which produced the same as the zero-time control.
26 Morning glory also demonstrated a slight (5%) drop in MDGD (monogalactosyldiacylglycerol),
27 with increases in PC (phosphatidylcholine), PG (phosphatidylglycerol), PI (phosphatidylinositol),
28 and PE (phosphatidylethanolamine) after 4 h. Twenty-four hours later, the drop in MDGD
29 (monogalactosyldiacylglycerol) was much larger and was thought to be related to an inhibition of
30 UDP-galactose 4-epimerase due to a rise in free fatty acids (FFAs) in the chloroplast. Note

1 that the two distinct timescales involved in O₃ fumigation, immediately post-fumigation and a
2 day or so later, allows for comparison after plant metabolism responds to the fumigation event.

3 The pathway for the formation of MGDG and DGDG (digalactosyldiacylglycerol) is located
4 on the chloroplast envelope. Diacylglycerol (DG) arrives from either the endomembrane system
5 or the stroma and the enzyme UDP-Galactose:1,2-diacylglycerol galactosyltransferase (UDGT)
6 forms MGDG with galactose from UDP-galactose. Sakaki et al. (1990) suggested that the
7 O₃-induced inhibition of UDGt was due to a release of FFAs from within the chloroplast. These
8 FFAs are inhibitory to UDGt, but not to GGGT, which is stimulated by high concentrations of
9 Mg²⁺ (Sakaki et al., 1990). The Sakaki et al. (1990) data indicate that the measured activities of
10 both enzymes isolated after fumigation has indicated that in vivo are not affected by O₃
11 fumigation in vivo. Both enzymes have sensitivity sulfhydryls, and both are located on the
12 envelope. Ozone, if it reaches those sulfhydryls, should inhibit these enzymes; yet inhibition
13 was not seen.

14 It has been thought for years that tocopherols functioned as antioxidants in biological
15 systems (Tappel, 1972). Hausladen et al. (1990) examined the role of antioxidants in red spruce
16 by following seasonal changes. They fit the level of tocopherol within the needles (/g-FW) to
17 the time of the year and found little change (fit as level = A + Bt + Ct²). From this empirical fit,
18 they found that the constant A was lower with higher levels of O₃. The seasonal variation
19 coefficients, B and C, were also lower, suggesting year long lows tocopherol levels. Variation
20 with the season is not particularly surprising, given that phytochrome action may be linked to
21 tocopherol biosynthesis (Lichtenthaler, 1977). Hansladen et al. (1990) reported a significant
22 (p < 0.05) trend in the difference between the high and low level of treatment; although there
23 was no discussion of why it occurred or what it meant in relation to metabolism. Their major
24 conclusion was that the antioxidant changes due to O₃ exposure may decrease cold hardiness.

25 Sterols, believed to act as membrane stabilizers, have been investigated by several groups
26 with mixed results. Tomlinson and Rich (1971), who exposed common bean at 0.25 ppm for
27 3 h, and Grunwald and Endress (1985), who exposed soybean at 0.07 ppm for 6 h for 48 days,
28 reported an increase in free sterols and a decline in esterified sterols. However, Trevanthen et al.
29 (1979) exposed tobacco at 0.3 ppm for 6 h and reported opposite results. None of these
30 investigators believed that O₃ had attacked the sterols directly, instead, they believed that these
31 changes involved metabolism and membrane stability. If O₃ induced a metabolic shift that

disturbed the polar lipid to sterol balance, membrane reactions to other stressors, such as cold tolerance, would certainly also be affected, perhaps detrimentally.

9.3.8 Role of Age and Size Influencing Response to Ozone

Clearly many changes occur with O₃ exposure can be observed within hours, or perhaps days, of the exposure. This document has argued that many of those events are connected with wounding and elicitor-induced changes in gene expression, but those are relatively fast acting changes (a timescale of tens of hours). Two other effects due to O₃ take longer to occur and tend to become most obvious under long periods of low-O₃ concentrations. These have been linked to senescence or some other physiological response very closely linked to senescence. These two responses, separated by a time sequence, are shown diagrammatically in Figure 9-15.




Figure 9-15. Linkage of senescence with hypersensitivity reactions and first event of O₃ attack of plants.

The understanding of how O₃ affects long-term growth and resistance to other biotic and abiotic insults in long-lived trees is unclear. Often, the conditions to which a tree is subjected to in one year will affect the response of that tree in the next year. This has been called “memory effect”, although the term “carry-over” is preferred. In other words, a condition in an earlier year sets the stage for a reaction in the next year; thereby giving a “cause-effect” scenario.

1 In perennial plant species, growth affected by a reduction in storage carbohydrates may
2 result in the limitation of growth the following year (carry-over effects). Carry-over effects have
3 been documented in the growth of tree seedlings (Hogsett et al., 1989; Sasek et al., 1991; Temple
4 et al., 1993; U.S. Environmental Protection Agency, 1996) and in roots (Andersen et al., 1991;
5 U.S. Environmental Protection Agency, 1996). Accumulation of the carry-over effects over time
6 will affect survival and reproduction. Data on the cumulative effects of multiple years of O₃
7 exposures have been, for the most part, the result of 2- to 3-year seedling studies. The difficulty
8 of experimentally exposing large trees to O₃ has lead to the tacit assumption that seedling
9 response to O₃ is a good predictor of large-tree response to O₃ (U.S. Environmental Protection
10 Agency, 1996).

11 The carry-over effects of O₃ exposures as observed in tree seedlings cited above by Hogsett
12 et al. (1989) have been termed “memory effects” by Langebartels et al. (1997) and proposed by
13 Schmieden and Wild (1995) to explain the sensitivity of spruce seedlings to frost in the winter
14 after having been exposed to O₃ during the previous summer. Norway spruce exposed to 80 ppb
15 for a whole growing season, demonstrated visible injury symptoms the following year when the
16 new needle flush appeared (Langebartels et al., 1997). Additional studies using Norway spruce
17 and Scots pine seedlings have showed similarly delayed responses following O₃ exposures.
18 Carry-over symptoms were noted to develop at different times of the year, depending on the
19 species of seedling exposed: in early spring for Norway spruce, and in early autumn for Scots
20 pine (Lange et al., 1989). Visible effects of O₃ exposures on spruce and pine may develop after a
21 substantial delay during the “sensitive” periods of the year when chlorophyll and needle loss
22 normally occur. Norway spruce and Scots pine differ in their sensitive periods because of the
23 different needle classes normally remaining on the tree (Langebartels et al., (1997).

24 Nutrient status of the tree during the over-wintering phase of its life (Schmieden and Wild,
25 1995) and chronic exposure to ambient O₃ (less severe with fewer peaks of very high levels)
26 induce (1) mineral nutrient deficiency; (2) alterations of normal metabolism, including
27 translocation and allocation of carbohydrates and probably nitrogen; and (3) disturbance of
28 normal transpiration and diurnal cycling, leading to water stress. This condition, termed
29 “Montane yellowing”, appears to be related to nutrient deficiencies rather than senescence
30 (although early loss of leaves and needles does occur). While generalized low nutrient
31 concentrations may not occur within the foliage, localized deficiencies may. However, they are

1 hard to observe or prove without a great deal of work involving all portions of a tree and without
2 a general hypothesis of what is occurring.

4 **9.3.9 Summary**

5 As the understanding of wounding responses of plants and more genome details and varied
6 plant mutants become available, the cellular and physiological responses of plants to O₃
7 exposures are slowly becoming clearer. However, more studies are needed on a larger variety of
8 species. Nevertheless, several key findings and conclusions can be highlighted:

- 9 (1) The entrance of O₃ into the leaf through the stomata remains the critical step in O₃
sensitivity. Not only does O₃ modify the opening of the stomata, usually closing it
partially, but O₃ also appears to alter the response of stomata to other stressful
situations, including a lowering of water potential and ABA responses. The
concentration of O₃ within the leaf is not the same as the external concentration due
to reactions within the leaf, but it is not “zero”.
- 10 (2) The initial reactions of O₃ within the leaf are still unclear, but the involvement of
H₂O₂ is clearly indicated. The detection of possible products by EPR spectroscopy
has progressed, but has not reached the point where any products can be identified.
Nonetheless, reaction of O₃ (or its product) with ascorbate and possibly other
antioxidants present in the apoplastic space of the mesophyll cells is clear and
serves to lower the amount of O₃ or product available to alter the plasma membrane
of the cells.
- 11 (3) The initial sites of membrane reactions seem to involve transport properties and,
possibly, the external signal transducer molecules. The alteration and mechanism of
the alteration of the varied carriers of K⁺ and Ca²⁺ is far from clear, but it would
seem that one of the primary triggers of O₃-induced cell responses is a change in
internal Ca²⁺ levels.
- 12 (4) The primary set of metabolic reactions that O₃ triggers now clearly includes those
typical of “wounding” responses generated by cutting of the leaf or by
pathogen/insect attack. Again, this seems to be due to a rise in cytoplasmic Ca²⁺
levels. Ethylene release and alteration of peroxidases and PAL activities, as well as
activation of many wound-derived genes, seem to be linked to some of the primary
reactions.
- 13 (5) The alteration of normal metabolism due to wounding has effects outside of the
cytoplasm. What effects are due to the “spreading of the problem” to other cellular
organelles is less clear. One of the secondary reactions is linked to an activation of
a senescence response. The loss of Rubisco and its messenger RNA is linked to an
early senescence or a speeding up of normal development leading to senescence.

The loss of photosynthetic capacity is linked to the lowered productivity of plants, and problems with efficient translocation are indicated, although photosynthesis and translocation still occur at a reasonable rate. The loss of productivity is not yet clearly explained.

It is important to note that the dramatic strides in understanding the genetic makeup of plants, gene control, and signal transduction/control over the last few years will likely accelerate in the future. That understanding will translate into better models of the hypotheses listed above and more detailed schemes of how O₃ alters much of basic plant metabolism. Thus, while understanding of how O₃ interacts with the plant at a cellular level has dramatically improved, the translation of those mechanisms into how O₃ is involved with altered cell metabolism, with whole plant productivity, and with other physiological facts remained to be more fully elucidated.

9.4 MODIFICATION OF FUNCTIONAL AND GROWTH RESPONSES

9.4.1 Introduction

The responses of plants to any air pollutant may be significantly influenced by a wide range of biological, chemical and physical factors. A plant's genetic make-up is an important inherent biological determinant of its response, but response can also be modified by other biological agents such as disease-causing organisms, insects and other pests, and by other higher plant species with which it may be competing for resources. Chemical factors that may influence response range from mineral nutrients obtained from the soil to other air pollutants and agricultural chemicals. Physical factors that may influence response include light, temperature and the availability of moisture, which are components of climate and climate change.

Some environmental factors are capable of being controlled, to some degree, by man, while others are not. The biological factors (pests, diseases, symbioses and competition) are partly controllable in agriculture but much less so (if at all) in natural ecosystems. It is possible to control agricultural soil fertility and the use of agricultural chemicals, as well as to exercise some control over the supply of water and airborne chemical factors. In contrast, the physical factors, i.e., light and temperature, are uncontrolled in the field even though they may be controllable in specialized situations such as greenhouses or shade houses.

1 The impacts of these various factors on plant response to O₃ and other oxidants were
2 extensively reviewed in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996).
3 It was noted in that document that, since any combination of these factors may come into play at
4 some time in a plant's life history, "response will be dictated by the plant's present and past
5 environmental milieu, which also includes the temporal pattern of exposure and the plant's stage
6 of development." That document also stressed that both the impact of environmental factors on
7 response to oxidants and the corollary effects of oxidants on responses to environmental factors
8 have to be considered in determining the impact of oxidants on vegetation in the field. The
9 variability observed in plant responses to defined exposures to O₃, particularly under field
10 conditions, is a consequence of the influences of genetics and the range of environmental
11 variables.

12 In view of the large number of factors to be considered and given that the purpose of this
13 document is to support the review of the O₃ NAAQS, including standards to protect vegetation,
14 this section focuses mainly on situations in which there is clear evidence that environmental
15 factors truly interact with oxidant effects, i.e., they magnify or diminish the impact of O₃ and are
16 *not merely additive* to it. Conversely, it will cover situations where O₃ acts synergistically or
17 antagonistically, but not additively, with effects induced by other factors. It will also emphasize
18 those interactions as a result of which overall plant growth and development, and yield are
19 adversely affected, rather than the details of interactions at the mechanistic level, unless the latter
20 are deemed to be essential to an understanding of larger scale effects.

21 To facilitate cross-reference, the present document uses essentially the same subsections as
22 in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996). Although light and
23 temperature are components of climate, they are initially reviewed as individual physical factors,
24 even though temperature effects are revisited to some extent in the discussion of interactions
25 with climate change.

26 Few studies reported since the 1996 document have systematically investigated
27 quantitative responses to O₃ concurrently with other variables. Although the 1996 document
28 cited almost 300 references pertaining to environmental interactions, and the present review cites
29 more than 350 new references, the bulk of the recently published work has continued to be
30 determined by the specific and frequently narrowly focused interests of individual researchers or
31 groups. Hence, the new findings are scattered and far from uniformly distributed among the

1 various sub-topics. In some instances there has been little or no research published that adds to
2 our understanding since the 1996 document. In such cases, the present review is therefore
3 restricted to summarizing the understanding that was current in 1996.

4 A few reviews have appeared since the early 1990s dealing with various environmental
5 interactions, and these are cited in relevant sections below. More general recent reviews are
6 those of Wellburn (1994); multi-authored volumes edited by Alscher and Wellburn (1994),
7 Yunus and Iqbal (1996), De Kok and Stulen (1998), and Bell and Treshow (2002); and reports
8 by the United Nations Environment Programme (UNEP) (1999) and the Intergovernmental
9 Panel on Climate Change (IPCC) (2001). Several biotic and abiotic interactions involving forest
10 trees are discussed in the review by Johnson et al. (1996b).

11 Although many reports have provided quantitative information on interactive effects, in
12 most cases the information is only descriptive of a specific situation involving only two or three
13 levels of a variable. While this may be adequate to provide statistical information about the
14 existence of interactions with environmental factors, it does not permit the development of
15 response surfaces or models to show the form that any influence of such factors might take on
16 O₃ exposure-response relationships or how O₃ might quantitatively influence responses to the
17 factors in question. This, together with the fragmented information available on the effects of
18 most factors, has contributed to the relative lack of development of simulation models of
19 oxidant-environmental factor interactions. Yet, as noted by Taylor et al. (1994), the large
20 number of variables constrains the assessment of pollution effects by experimentation alone.
21 The only alternative is to use mathematical models to attempt to predict the outcome of different
22 O₃ and environmental factor scenarios, building up their complexity in stages. The few models
23 thus far used to investigate O₃ stress have been adapted from existing process models of crop or
24 tree growth which include limited numbers of physical or chemical variables (such as
25 temperature, soil water stress, or nutrient deficiency). Taylor et al. (1994) provide a listing of
26 several simulation models developed for trees at the individual, stand, and regional levels, and
27 these and many other models have been critically reviewed by Kickert and Krupa (1991) and
28 Kickert et al. (1999). However, regardless of whether such models are descriptive/empirical or
29 process/mechanistic, their outputs will always be associated with varying degrees of uncertainty
30 and require validation against observable responses wherever possible. Kickert et al. (1999) also
31 point out that very few of the models that have been described provide *risk assessments* that

1 address likelihood, in contrast to *consequence assessments* that address the magnitudes of
2 effects. Thus, even though capable simulation models of plant response to O₃ involving
3 complex mixes of many biological, physical and chemical factors may be out of reach at the
4 present time, the use of newer mathematical approaches such as artificial neural networks
5 (ANNs) has enabled insightful analyses to be performed in several field studies involving
6 numerous micrometeorological and other environmental variables (e.g., Balls et al., 1996;
7 Mills et al., 2000).

8 Because the ensuing subsections deal with studies of O₃ interactions involving an
9 extremely wide array of biological, physical, and chemical factors in the plant's environment,
10 it is inevitable that many different exposure facilities and regimes have been used in these
11 studies. To provide specific information regarding the O₃ exposure concentrations, profiles,
12 hours and days of exposure (as well as the types of systems and facilities used for the exposures)
13 would add a wealth of detail that would do little to assist our understanding of the roles of
14 environment factors in modifying the impact of O₃ on vegetation or to facilitate our ability to
15 estimate the magnitudes of any such modifications. Thus, only experiments in which the
16 exposure levels and regimes were *within the bounds of ambient experience in North America*
17 are discussed in the ensuing subsections, regardless of the type of exposure profile used. The
18 cutoffs used have been: ~200 ppb for peak hourly concentrations or for short-term exposures;
19 ~100 ppb for daytime means involving prolonged exposures for several hours; or a doubling of
20 ambient levels, in cases in which enriched exposure levels were a function of ambient levels.
21 Actual details of the exposure regimes and conditions can, of course, be obtained from the
22 original references but are only stated here when any distinction needs to be made between the
23 effects of different exposure levels. Hence, it should be understood that ensuing statements such
24 as "... it was found that O₃ caused ..." should always be read as "... it was found that exposures to
25 O₃ (within the range of those that have been measured in ambient air) caused ..."

27 **9.4.2 Genetics**

28 The response of individual plants to O₃ is affected by several factors including the
29 environment in which it is growing, competition with neighbor plants, ontogeny, and genetics.
30 This section examines the role of genetics in plant response to O₃. In addition, major knowledge
31 gaps in the understanding of genetic aspects of O₃ response are pointed out.

1 It is well known that species vary greatly in their responsiveness to O₃ (Environmental
2 Protection Agency, 1996). This again has been recently demonstrated for grassland species
3 (Warwick and Taylor, 1995; Pleijel and Danielson, 1997; Grime et al., 1997; Bungener et al.,
4 1999a, b; Franzaring et al., 2000), wild herbaceous plants (Bergmann et al., 1999; Danielsson
5 et al., 1999; Nussbaum et al., 2001), agricultural crops (Renaud et al., 1997; Ollerenshaw et al.,
6 1999; Benton et al., 2000; Heagle and Stefanski, 2000; Fumagalli et al., 2001; Elagöz et al.,
7 2002; Nali et al., 2002; Köllner and Krause, 2003), horticultural shrubs and trees (Hormaza
8 et al., 1996; Findley et al., 1997), and forest trees (Pääkkönen et al., 1997; Volin et al., 1998; Pell
9 et al., 1999; Bortier et al., 2000; Postiglione et al., 2000; Landolt et al., 2000; Zhang et al., 2001;
10 Oksanen and Rousi, 2001; Matsumura, 2001; Guidi et al., 2001; Momen et al., 2002; Nali et al.,
11 2002; Saitanis and Karandinos, 2002). These studies have shown a wide range of responses to
12 O₃, from growth stimulation by a few species such as *Festuca ovina* L. (Pleijel and Danielson,
13 1997) and *Silene dioica* and *Chrysanthemum leucanthemum* (Bungener et al., 1999) to
14 significant growth reduction, depending on environmental conditions and exposure dose.

15 While determining the explanation for differences in species sensitivity to O₃ remains one
16 of the challenges facing plant biologists (Pell et al., 1999), a number of hypotheses have been
17 suggested. Reich (1987) proposed that variation in O₃ sensitivity could be explained by variation
18 in total uptake of the gas. Others have suggested that (1) fast-growing species are more sensitive
19 than slower growing ones (Bortier et al., 2000), (2) overall O₃ sensitivity may be closely linked
20 to root responses to O₃ (Warwick and Taylor, 1995) or (3) the relative ability of species to
21 detoxify O₃-generated reactive oxygen free radicals may determine O₃ sensitivity (Alscher et al.,
22 1997; Pell et al., 1999). Volin et al., (1998) suggest that the relative rate of stomatal
23 conductance and the photosynthesis rate at a given conductance both contribute strongly to
24 determining species sensitivity to O₃. Likely, there is more than one mechanism determining
25 sensitivity, even in a single species.

26 Within a given species, individual genotypes or populations can also vary significantly in
27 O₃ sensitivity (Environmental Protection Agency, 1996). For example, the intraspecific
28 variation in O₃ sensitivity for was a factor of two for *Phleum pratense* (Danielsson et al., 1999)
29 and *Trifolium repens* L. (Postiglione et al., 2000). A similar range of intraspecific variations in
30 O₃ responses was demonstrated for clonal differences in *Betula pendula* by Pääkkönen et al.
31 (1997) and *Prunus serotina* (Lee et al., 2002). These examples of wide ranges within species

responses suffice to show that caution should be taken when ranking species categorically as having an absolute degree of O₃ sensitivity (Davison and Barnes, 1998).

9.4.2.1 Genetic Basis of O₃ Sensitivity

Plant response to ozone is determined by genes that are directly related to oxidant stress and to an unknown number of genes that are not specifically related to oxidants. The latter includes genes that control leaf and cell wall thickness, stomatal conductance and the internal architecture of the air spaces. Although there is currently a great emphasis on individual antioxidants that can be manipulated by molecular methods, the challenge is to determine the relative contributions of all of the components to plant response and to understand the interplay between them. Recent studies using molecular biological tools are beginning to increase the understanding of O₃ toxicity and differences in O₃ sensitivity.

While much of the research in developing the understanding of O₃ responses has been correlative in nature, recent studies with transgenic plants have begun to positively verify the role of various genes and gene products in O₃ tolerance. Overexpressing MnSOD in chloroplasts increased O₃ tolerance in transgenic tobacco plants (Van Camp et al., 1994) provided the first definitive proof of antioxidants key role in O₃ tolerance. Subsequently, Broadbent et al. (1995) showed that overexpression of pea glutathione reductase simultaneously in both chloroplasts and mitochondria of transgenic tobacco enhanced O₃ tolerance. Similarly, increased O₃ tolerance to O₃-induced foliar necrosis was shown for transgenic tobacco plants overexpressing the cytosolic Cu/Zn-SOD gene (Pitcher and Zilinskas, 1996). Transgenic tobacco plants expressing antisense RNA for cytosolic ascorbate peroxidase, which reduces ascorbate peroxidase production, showed increased susceptibility to O₃ injury suggesting a key role in O₃ tolerance for the antioxidant ascorbate peroxidase (Örvar and Ellis, 1997).

The consensus among molecular studies of O₃ sensitivity is pointing to O₃ triggering salicylic acid, ethylene, and jasmonic acid and that the signaling of these molecules determines, in some cases, the O₃ susceptibility of plants (Overmyer et al., 2000; Rao and Davis, 1999; Rao et al., 2000; Langerbartels et al., 2002; Moeder et al., 2002; Tamaoki *et al.*, 2003; Vahala et al., 2003a, b). Increased levels of jasmonic acid production in O₃-tolerant compared to O₃-sensitive plants has been shown for *Arabidopsis* (Overmyer et al., 2000) and *Populus* (Koch et al., 1998, 2000). Blockage of ethylene production by using antisense methods with ACC synthase and

1 ACC oxidase suggest strongly that ethylene synthesis and perception are required for H₂O₂
2 production and cell death following O₃ exposure of *Lycopersicon esculentum* (Moeder et al.,
3 2002). Ethylene signaling may have multiple roles in O₃ tolerance determination as was
4 demonstrated recently by Vahala et al., (2003a, b) who found that in *Populus tremula* ×
5 *P. tremuloides* hybrid clones differing in O₃ sensitivity, ethylene accelerated leaf senescence in
6 sensitive plants under low O₃, but under acute O₃, ethylene seemed to be required for protection
7 from cell death.

8 While changing expressions of single antioxidant genes has proven very useful in
9 identifying possible mechanisms of O₃ sensitivity and tolerance (Kuzniak, 2002), it should be
10 noted that some studies of transgenic plants with enhanced antioxidant production have not
11 resulted in increased O₃ tolerance (Saji et al., 1997; Torsethaugen et al., 1997; Strohm et al.,
12 1999, 2002). Clearly, ethylene production plays a role in O₃ sensitivity but the role of various
13 antioxidants in O₃ tolerance regulation are yet to be fully elucidated (Wellburn and Wellburn,
14 1996). It is unlikely that single genes are responsible for O₃ tolerance responses, except in rare
15 exceptions (Engle and Gabelman, 1966). Regulation of stomatal opening and leaf structure
16 (Bennett et al., 1992; Elegoz and Manning, 2002) are also likely to play key roles in O₃ tolerance
17 in plants. Newly developing opportunities to examine simultaneous regulation of larger numbers
18 of genes will likely yield more clarification of genes controlling O₃ tolerance (Desikan et al.,
19 2001; Matsuyama et al., 2002).

20 Attempts to demonstrate conclusively changes in antioxidant and protective pigments for
21 O₃ sensitive and tolerant mature trees growing in the field have largely been unsuccessful (Tausz
22 et al., 1999a, b). However, evidence for antioxidant expression differences contributing to
23 differences in O₃ sensitivity of four-year-old *Populus tremuloides* trees has been found
24 (Wustman et al., 2001).

26 **9.4.3 Environmental Biological Factors**

27 The biological factors within the plant's environment that may influence its response to O₃
28 directly or may be influenced to the advantage or disadvantage of plants encompass insects and
29 other animal pests, diseases, weeds and other competing plant species. Although such
30 interactions are ecological in nature, those involving individual pests, plant pathogens, or weeds,

or agricultural crop or forest tree species are considered in this section. More complex ecological inter-species interactions are dealt with in Section 9.5.

The different types of biological factors are dealt with separately, as in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996). Still, it is important to recognize certain general features of relationships of plants with biological components of their environment:

- Successful infestation or infection involves complex interactions among the target or host species, the causal organism and environmental factors.
- Infestations and infections may co-occur.
- The successful development and spread of a pest, pathogen or weed require favorable environmental factors.
- Significant losses to crops and forest trees result from pests and pathogens.
- Significant losses to crops and seedling trees result from weed competition.

Ozone and other photochemical oxidants may influence the severity of a disease or infestation by a pest or weed, either by direct effects on the causal species, or indirectly by affecting the host, or both. In addition, the interaction between O₃, a plant and a pest, pathogen or weed may influence the response of the target host species to O₃. A perceptive overview of the possible interactions of O₃-exposure with insect pests and fungal diseases has been provided by Jones et al. (1994), based on a model system involving two insects and two pathogens affecting cottonwood (*Populus deltoides*). Their study also included effects on the decomposition of leaf litter.

In contrast to detrimental biological interactions, there are mutually beneficial relationships or symbioses involving higher plants and bacteria or fungi. These include (1) the nitrogen-fixing species *Rhizobium* and *Frankia* that nodulate the roots of legumes and alder, and (2) the mycorrhizae that infect the roots of many crop and tree species, all of which may be affected by exposure of the host plants to O₃.

In addition to the interactions involving animal pests, O₃ may also have indirect effects on higher herbivorous animals, e.g., livestock, due to O₃-induced changes in feed quality.

9.4.3.1 Oxidant-Plant-Insect Interactions

The 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) stressed the variability in the reported effects of O₃ on host plant-insect interactions. Since relatively few plant-insect

1 systems have been studied, few consistent patterns of response have emerged, as noted in other
2 reviews such as those of Colls and Unsworth (1992), Heliövaara and Väisänen (1993), Whittaker
3 (1994), Docherty et al. (1997) and, most recently, Flückiger et al. (2002).

4 None of the studies reported in the past decade have clarified the situation in terms of
5 clearly consistent effects. A 1997 review by Docherty et al. (1997), for example, examined
6 17 reports of studies of aphid species on a range of hosts and classed the O₃ effects on aphid
7 performance as: 35% positive; 41% negative; and 24% showing no significant effect.
8 A tabulation of 19 studies by Flückiger et al. (2002) gave the corresponding figures: 42%,
9 21%, and 37%.

10 Other recent studies with the aphids *Schizolachnus pineti* and *Cinara pinea* on Scots pine
11 (*Pinus sylvestris*) and *Cinara pilicornis* on Norway spruce (*Picea abies*) have also yielded
12 variable results, with them suggesting that O₃ enhances aphid density on pine and aphid
13 performance on spruce (Holopainen et al., 1997; Kainulainen et al., 2000a). In an earlier study
14 with *Schizolachnus pineti* on Scots pine, Kainulainen et al. (1994) had observed no significant
15 effects of O₃-treatment on aphid performance. However, more recent observations of long-term
16 effects on aphid populations on aspen (*Populus tremuloides*) exposed to O₃ in a FACE system
17 revealed that O₃ significantly increased aphid populations and decreased the populations of
18 predatory insects (Percy et al., 2002).

19 The observations of Brown et al. (1993) and Jackson (1995) led Whittaker (1994) and
20 Brown (1995) to suggest that aphid response was dependent on ambient temperature as well as
21 the dynamics of O₃ exposure; growth tended to be stimulated with maximum temperatures below
22 ~20 °C but was reduced at higher temperatures. The present situation with plant-aphid responses
23 therefore remains confused and, although numerous suggestions have been offered to explain
24 specific findings, they are difficult to assemble into a coherent picture.

25 Variability has also been found with the interactions involving chewing insects. For
26 example, Lindroth et al. (1993) reported a small negative O₃ effect (8% reduction) on the growth
27 of gypsy moth larvae (*Limantra dispar*) on hybrid poplar (*Populus tristis* × *balsamifera*) but no
28 effect when on sugar maple (*Acer saccharum*). Ozone resulted in reduced growth rate of the
29 larvae of the bug *Lygus rugulipennis* on Scots pine, but enhanced the growth of larvae of the
30 sawfly *Gilpinia pallida* (Manninen et al., 2000). Costa et al. (2001) observed no significant O₃

1 effects on the growth and fecundity of the Colorado potato beetle (*Leptinotarsa decemlineata*)
2 on potato in greenhouse and field experiments.

3 Fortin et al. (1997), in a two-year study of forest tent caterpillar (*Malacosoma disstria*) on
4 sugar maple, observed that O₃ exposure only increased the growth rate of female larvae in one
5 year; 4th- and 5th-instar larvae also showed a feeding preference for treated foliage in that year.
6 However, studies based on open-air exposures of aspen (*Populus tremuloides*) indicated
7 O₃-enhanced growth of *M. disstria* in terms of pupal weight (Percy et al., 2002) and larval
8 performance (Kopper and Lindroth, 2003). Jackson et al. (2000) observed inconsistency in
9 studies on the larva of the tobacco hornworm (*Manduca sexta*) on tobacco (*Nicotiana tabacum*).
10 In one year, feeding on O₃-treated foliage resulted in significantly greater larval weight, whereas
11 in a second year the increase was not statistically significant although survival was increased.
12 Also, oviposition by hornworm moths was increased if ambient O₃ levels were increased by 70%
13 but fell back to normal in ambient O₃ levels (Jackson et al., 1999).

14 Studies of the two-spotted spider mite (*Tetranychus urticae*) on white clover (*Trifolium*
15 *repens*) and peanut (*Arachis hypogaeae*) by Heagle et al. (1994) and Hummel et al. (1998)
16 showed that, on peanut and an O₃-sensitive clover clone, O₃-exposure stimulated mite
17 populations. The lack of significant effects on mites on the O₃-resistant clover clone suggests
18 that the responses were host-mediated.

19 With chewing insects and mites, there therefore appears to be a clearer indication of the
20 likelihood that increased insect performance will result from O₃-induced changes in the host
21 plant, but negative effects continue to be reported, indicating that the response is probably also
22 being determined in part by other environmental, genetic, or temporal variables.

23 Reported O₃-induced enhancement of attack by bark beetles (*Dendroctonus brevicornis*) on
24 Ponderosa pine (*Pinus ponderosa*) has been suggested by Dahlsten et al. (1997) to be due to
25 greater brood development on injured trees, possibly related to decreased numbers of predators
26 and parasitoids. This view gains some support from the observation that O₃ adversely affected
27 the searching behavior of a parasitoid, *Asobara tabida*, for larvae of *Drosophila subobscura*
28 which led to fewer parasitized hosts (Gate et al., 1995). Such observations reveal another level
29 of complexity in the O₃-plant-insect interrelationship: O₃ may reduce the effectiveness of the
30 natural control of insect pests. The phenomenon is probably related to effects on olfactory cues,

1 since it was shown by Arndt (1995) that O₃ can affect fly behavior by modifying the pheromones
2 that cause fly aggregation.

3 These reports focus on the direct or indirect effects on the insect or mite feeding on foliage
4 previously or currently exposed to O₃. They provide little if any information on effects on the
5 host plant other than qualitative references to the injury caused by the O₃-exposure. Enhanced
6 pest development will ultimately lead to increased adverse effects on the hosts in the long term,
7 but the only report of an O₃-plant-insect interaction directly affecting the host plant in the short
8 term still appears to be that of Rosen and Runeckles (1976). They found that infestation by the
9 greenhouse whitefly (*Trialeurodes vaporariorum*) sensitized bean plants (*Phaseolus vulgaris*) to
10 injury by otherwise non-injurious low levels of O₃, leading to premature senescence of the
11 leaves.

12 The overall picture regarding possible O₃ effects on plant-insect relations, therefore,
13 continues to be far from clear. Only a few of the very large number of such interactions that
14 may affect crops, forest trees and other natural vegetation have been studied. The trend
15 suggested in the 1996 criteria document that O₃ may enhance insect attack has received some
16 support from a few recent studies. However, the variability noted in most of the studies makes it
17 clear that we are still far from being able to predict the nature of any particular O₃-plant-insect
18 interaction or its magnitude or severity.

19 20 **9.4.3.2 Oxidant-Plant-Pathogen Interactions**

21 Plant diseases are caused by pathogenic organisms, e.g., fungi, bacteria, mycoplasmas,
22 viruses, and nematodes. Ozone impacts on disease are briefly discussed in earlier reviews by
23 Ayres (1991) and Colls and Unsworth (1992) and, more recently, by Flückiger et al. (2002).
24 Biotic interactions with forest trees have been reviewed by Chappelka and Samuelson (1998);
25 and Sandermann (1996) and Schraudner et al. (1996) have summarized molecular similarities
26 and interrelationships between necrotic O₃ injury to leaves and pathogen attack. A few recent
27 publications have added to our fragmented knowledge of O₃-plant-disease interactions and the
28 mechanisms involved, but there appear to have been no reports to date of studies involving
29 mycoplasmal diseases.

30 The 1996 criteria document (U.S. Environmental Protection Agency, 1996) noted the
31 concept put forward by Dowding, (1988) “that pathogens and pests which can benefit from

1 damaged host cells and from disordered transport mechanisms are enhanced by pollution insult
2 to their hosts, whereas those pathogens and other symbionts which require a healthy mature host
3 for successful invasion are depressed by pollutant stress to their host.” The pathogens of the first
4 type are mostly facultative, necrotrophic, fungal parasites, whereas the second type are largely
5 obligate biotrophic fungi, bacteria and viruses. Based on this distinction, the majority of the
6 cases cited in the 1996 document supported Dowding’s view, as have several more recent studies
7 summarized in Table 9-11. However, there are also some contradictions.

8 Most investigations have focused on the incidence and development of disease on plants
9 previously or concurrently exposed to O₃, rather than on the corollary effect of disease on the
10 response to O₃. In all of the studies of facultative pathogens and the nematode studies, exposure
11 to O₃ tended to result in increased disease severity through increased spore germination or
12 increased fungal growth and development, although in the case of grey mold (*Botrytis cinerea*)
13 on kidney bean (*Phaseolus vulgaris*) this was only observed on an O₃-sensitive cultivar
14 inoculated with conidia (Tonneijck, 1994). Using mycelial inoculation, O₃ reduced disease
15 development, but no satisfactory explanation was offered to account for the difference in
16 response. With leaf spot, *Marssonina tremulae*, on hybrid poplar (*Populus trichocarpa* ×
17 *balsamifera*), low level exposures to O₃ also increased disease (in agreement with theory) but
18 higher levels (200 ppb, 8h per day for 15 days) reduced conidial germination (Beare et al.,
19 1999b).

20 The situation with obligate biotrophic pathogens is less consistent. The effects on powdery
21 mildew (*Sphaerotheca fulginea*) on both bottle gourd (*Lagenaria siceraria*) and cucumber
22 (*Cucumis sativa*) resembled the situation with the necrotrophic poplar leaf spot disease
23 (*Marssonina tremulae*), since low O₃ exposures increased disease severity (in disagreement with
24 theory) although higher levels decreased it. The decreased infection in the pea-powdery mildew
25 (*Erysiphe polygoni*) situation agrees with theory, but the situations with leaf rust (*Melampsora*
26 sp.) on poplar (*Populus trichocarpa* × *balsamifera*) or aspen (*Populus tremuloides*) do not.
27 However, these reports are in contrast to earlier reports included in the 1996 criteria document
28 (U.S. Environmental Protection Agency, 1996) of observations with other species of *Erysiphe*
29 (Tiedemann et al., 1991) and *Melampsora* (Coleman et al., 1987). In contrast to the recent report
30 of a synergism with *Melampsora* on poplar, infections caused by the other biotrophs
31 (*Sphaerotheca*, *Erysiphe*, *Uromyces*) reduced the severity of injury caused by O₃ (in agreement

Table 9-11. Interactions Involving O₃ and Plant Pathogens

Host Plant	Pathogen	Effect of O ₃ on Disease	Effect of Disease on O ₃ Response	Reference
Obligate Biotrophs				
Bottle gourd (<i>Lagenaria siceraria</i>)	Powdery mildew (<i>Sphaerotheca fulginea</i>)	Increased in 50ppb O ₃ ; decreased in 100+ppb	Decreased; partial protection	Khan and Khan (1998a)
Cucumber (<i>Cucumis sativa</i>)	Powdery mildew (<i>Sphaerotheca fulginea</i> ;	Increased in 50ppb O ₃ ; decreased in 100+ppb	Synergistic increase in 50ppb O ₃ ; antagonistic decrease in 100+ppb; partial protection	Khan and Khan (1999)
Pea (<i>Pisum sativum</i>)	Powdery mildew (<i>Erysiphe polygoni</i>)	Decreased infection	Decreased; partial protection	Rusch and Laurence (1993)
Aspen (<i>Populus tremuloides</i>)	Leaf rust (<i>Melampsora medusae</i> f. sp. <i>tremuloidae</i>)	Increased severity	Not reported	Karnosky et al. (2002)
Hybrid poplar (<i>Populus trichocarpa</i> × <i>balsamifera</i>)	Leaf rust (<i>Melampsora larici-populina</i> or <i>M. allii-populina</i>)	Increased infection and severity	Increased sensitivity (synergistic)	Beare et al. (1999a)
Broad bean (<i>Vicia faba</i>)	Bean rust (<i>Uromyces viciae-fabae</i>)	Not reported	Decreased; partial protection	Lorenzini et al. (1994)
Facultative Necrotrophs				
Kidney bean (<i>Phaseolus vulgaris</i>)	Grey mold (<i>Botrytis cinerea</i>)	Increased from conidia on O ₃ -sensitive cultivar; decreased from mycelium	Not reported	Tonneijck (1994)
	Grey mold (<i>Botrytis cinerea</i>)	Increased infection	Not reported	Tonneijck and Leone (1993)
	White mold (<i>Sclerotinia sclerotiorum</i>)	Increased infection	Not reported	Tonneijck and Leone (1993)
Scots pine (<i>Pinus sylvestris</i>)	Annosus root and butt rot (<i>Heterobasidion annosum</i>)	Increased development*	Not reported	Bonello et al. (1993)

Table 9-11 (cont'd). Interactions Involving O₃ and Plant Pathogens

Host Plant	Pathogen	Effect of O ₃ on Disease	Effect of Disease on O ₃ Response	Reference
Facultative Necrotrophs (cont'd)				
Loblolly pine (<i>Pinus taeda</i>)	Pitch canker (<i>Fusarium subglutinans</i>)	Increased development	Increased sensitivity	Carey and Kelley (1994)
Hybrid poplar (<i>Populus deltoides</i> × <i>nigra</i>)	Canker (<i>Septoria musiva</i> [= <i>Mycosphaerella populinum</i>])	Increased incidence	Not reported	Woodbury et al. (1994)
Hybrid poplar (<i>Populus trichocarpa</i> × <i>balsamifera</i>)	Leaf spot (<i>Marssonina tremulae</i>)	Increased spore germination and lesion growth after 100ppb O ₃ (30 days); decreased germination after 200ppb (15 days)	Not reported	Beare et al. (1999b)
Wheat (<i>Triticum aestivum</i>)	Blotch (<i>Septoria nodorum</i>)	Increased infection	Not reported	Tiedemann and Firsching (1993)
	Tan spot (<i>Pyrenophora tritici-repentis</i>)	Increased infection of disease-susceptible genotypes	Not reported	Sah et al. (1993)
Nematodes				
Tomato (<i>Lycopersicon esculentum</i>)	Root-knot nematode (<i>Meloidogyne incognita</i>)	Increased development	Increased foliar injury; reduced plant growth (synergistic)	Khan and Khan (1997, 1998a)

* Increase completely countered by mycorrhizae (*Hebeloma crustuliniforme*).

1 with numerous earlier reports), but only at high O₃ exposures in the case of *Sphaerotheca* on
2 cucumber. At low exposure levels, the disease and O₃ acted synergistically. The only other
3 recent observation of such disease-related synergisms are the nematode-tomato reports of (Khan
4 and Khan, 1997, 1998b).

5 It is therefore clear that the type and magnitude of exposure to O₃ plays an important role
6 in determining both the response of the disease organism and of the host.

7 No recent studies involving interactions between O₃ and bacterial diseases appear to have
8 been reported. With regard to viruses, a laboratory study by Yalpani et al. (1994) added to
9 several reports of O₃ decreasing the severity of tobacco mosaic virus infection of tobacco,
10 *Nicotiana tabacum*; and Jimenez et al. (2001) reported that previous exposure to O₃ resulted in
11 increased adverse effects on tomato yield attributed to several virus diseases.

12 Similarities between the sensitivities of different cultivars or clones to O₃ and to specific
13 diseases have been noted. For example, Sah et al. (1993) found that the severity of injury caused
14 by tar spot and standard O₃ exposures of 12 wheat cultivars were closely correlated ($R^2 = 0.986$).
15 Such similarities appear to have a mechanistic basis, since several studies have noted similarities
16 in the molecular and biochemical changes that occur in plants infected with pathogens and plants
17 exposed to O₃. Schraudner et al. (1992), Ernst et al. (1992), Eckey-Kaltenbach et al. (1994a,b),
18 Yalpani et al. (1994) and Bahl et al. (1995) have presented evidence that exposures to O₃ result
19 in responses such as increased levels of salicylic acid, the signaling agent for increased induced
20 resistance to pathogens. This, in turn, leads to activation of the genes that encode defense
21 proteins, including the so-called pathogenesis-related proteins. The induction of such proteins
22 might account for the decreased infection with *Sphaerotheca* and *Melampsora* at higher O₃
23 exposures but does not account for increased infections seen at lower exposure levels. The issue
24 is discussed more fully by Sandermann (1996) and Schraudner et al. (1996). More recently
25 Sandermann has extended the theory relating O₃ and disease by suggesting that, because of O₃
26 “memory effects” in affected host plants that may persist over weeks or months, analysis for
27 various induced biomarkers of gene activation may provide a useful tool for improving our
28 ability to predict the outcome of O₃-plant-pathogen interactions (Sandermann, Jr., 2000).

29 There have been no reports of O₃ studies with mixed infections by pathogens, but the
30 complete suppression of *Heterobasidion* butt and root rot of Scots pine by the mycorrhizal

symbiont *Hebeloma crustuliniforme* indicates the possibility of interactions involving more than one fungus (see Section 9.4.4.3.3 below).

In summary, our understanding of oxidant-plant-disease interactions is far from complete. However, a combined tabulation of the evidence presented in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) and that noted in Table 9-11 leads to the following summary of O₃ effects on plant diseases and corollary effects of infection on plant response to O₃, as indicated by number of studies showing increases or decreases in disease or susceptibility:

For obligate biotrophic fungi, bacteria, nematodes:

O ₃ increased disease:	9	Increased susceptibility to O ₃ :	3.
-----------------------------------	---	--	----

O ₃ decreased disease:	15	Decreased susceptibility to O ₃ :	9.
-----------------------------------	----	--	----

For facultative necrotrophic fungi:

O ₃ increased disease:	25	Increased susceptibility to O ₃ :	2.
-----------------------------------	----	--	----

O ₃ decreased disease:	3	Decreased susceptibility to O ₃ :	4.
-----------------------------------	---	--	----

Thus, although O₃ may reduce the severity but not the incidence of some of the diseases caused by the obligate pathogens, the evidence overall indicates that with most diseases, their severity is more likely to be increased by O₃ than not. However, the actual consequences will be specific to the disease and level of exposure, and, most importantly, will be determined by environmental suitability and epidemiological requirements for disease to develop. Conversely, some evidence suggests that infection by obligate pathogens may confer some degree of “protection” against O₃, a dubious benefit from the plant’s point of view.

9.4.3.3 Oxidant-Plant-Symbiont Interactions

No further studies have appeared regarding O₃ effects on the important bacterial symbiont of legumes, *Rhizobium*, since those summarized in the 1996 document (U.S. Environmental Protection Agency, 1996). Hence, our present understanding is that, although relatively high levels of exposure (> 200 ppb) can result in severe (> 40%) reductions in nodulation (and therefore nitrogen-fixation) on soybean roots, lesser reductions in nitrogen-fixing may be caused by lower exposures. However, the data are inadequate to attempt to define any quantitative exposure-response relationships.

1 There have been a few recent reports on O₃-plant-mycorrhizae interrelationships. These
2 have mostly involved seedlings of coniferous tree species. A transient O₃-induced stimulation of
3 mycorrhiza on Scots pine roots reported by Kasurinen et al. (1999) was not observed in a later
4 study by Kainulainen et al. (2000b). Studies of the mycorrhiza *Paxillus involutus* on birch
5 (*Betula pendula*) seedlings showed that, although O₃ reduced mycorrhizal growth rate, it led to
6 greater extension growth which in turn resulted in greater mycorrhizal infection of neighboring
7 Aleppo pine (*Pinus halepensis*) seedlings (Kytöviita et al., 1999). However, O₃ reduced nitrogen
8 acquisition by *P. halepensis* from its mycorrhizal symbiont (Kytöviita et al., 2001). The
9 complex interrelationships that may occur in the rhizosphere were revealed by the observation
10 (Bonello et al., 1993) that the mycorrhiza *Hebeloma crustuliniforme* could overcome the
11 O₃-stimulated severity of root rot on Scots pine (*Pinus sylvestris*) caused by the fungus
12 *Heterobasidion annosum* (noted in Section 9.4.4.3.2).

13 In summary, the available evidence is far too fragmented and contradictory to permit
14 drawing any general conclusions about mycorrhizal impacts. The negative effects of O₃ on
15 mycorrhizae and their functioning that have been reported have not necessarily been found to
16 lead to deleterious effects on the growth of host plants. Thus, little has changed from 1991 when
17 Dighton and Jansen asked: “Atmospheric Pollutants and Ectomycorrhizae: More Questions
18 than Answers?” (Dighton and Jansen, 1991). Because of their important roles in ecosystems,
19 mycorrhizae are further discussed in Section 9.5 below.

21 **9.4.3.4 Oxidant-Plant-Plant Interactions: Competition**

22 Plant competition involves the ability of individual plants to acquire the environmental
23 resources needed for growth and development: light, water, nutrients and space. Intraspecific
24 competition involves individuals of the same species, typically in monocultural crop situations,
25 while interspecific competition refers to the interference exerted by individuals of different
26 species on each other when they are in a mixed culture.

27 In cropping situations, optimal cultural practices for row spacing and plant density/row
28 tend to balance the negative effects of intraspecific competition and the goal of maximum yield.
29 Although interspecific competition is agriculturally undesirable when it involves weak
30 infestations, the use of mixed plantings may be agriculturally deliberate, e.g., grass-legume

1 mixtures used for pasture or forage. In natural plant communities, monocultures are rare, and
2 complex interspecific competition is the norm.

3 Although weak competition is the largest global cause of crop losses, little is known about
4 the impact of O₃ on crop-weed interactions. The topic does not appear to have been investigated
5 in recent years. We can only speculate as to the possible consequences of O₃ exposure on weed
6 competition based on our limited understanding of the effects on a few, mostly two-component
7 mixtures of cultivated species.

8 The tendency for O₃-exposure to shift the biomass of grass-legume mixtures in favor of
9 grass species, reported in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) has
10 been confirmed by recent studies. In a ryegrass (*Lolium perenne*) + clover (*Trifolium repens*)
11 mixture grown in an open-air fumigation system, clover growth was impaired by extended
12 exposures to above-ambient O₃, leaving patches for weed invasion (Wilbourn et al., 1995).
13 An open-top chamber study by Nussbaum et al. (1995b) using the same species confirmed the
14 greater effect on clover but observed that the magnitude of the effect depended highly on the
15 pattern of O₃-exposures over extended growing periods. Low-level exposures shifted species
16 composition in favor of *Lolium*, but exposures to higher peak O₃ levels depressed total mixture
17 yield. With an alfalfa (*Medicago sativa*) + timothy (*Phleum pratense*) mixture, Johnson et al.
18 (1996a) noted that O₃ caused decreased alfalfa root growth and increased timothy shoot growth
19 and height. Nussbaum et al. (2000a) reported that, with increased exposure to O₃, well-watered
20 red clover (*Trifolium pratense*) plants suffered from increased competition from the grass,
21 *Trisetum flavescens*, but the O₃ exposure also negatively effected grass growth, depressing
22 overall total yield. However, a greater adverse effect on *Trisetum* resulted from O₃-induced
23 increased competition when grown with brown knapweed (*Centaurea jacea*), a weed species.

24 Andersen et al. (2001) demonstrated the potential for competition and O₃-exposure to work
25 together to affect the growth of tree seedlings. Ozone had no direct adverse effect on pine
26 growth in a 3-year study of ponderosa pine (*Pinus ponderosa*) seedlings grown in mesocosms
27 with three densities of blue wild-rye grass (*Elymus glaucus*), but the O₃-increased competitive
28 pressure of the grass caused a major reduction in pine growth.

29 Three studies have been reported on more complex plant associations. Ashmore and
30 Ainsworth (1995) studied mixed plantings of two grasses, *Agrostis capilaris* and *Festuca rubra*,

1 with two forbs¹⁶, *Trifolium repens* (a legume) and *Veronica chamaedrys*, exposed to O₃ in
2 open-top chambers (OTCs). The proportion of forbs, *Trifolium* in particular, declined, especially
3 when cut at biweekly intervals. In a related study, Ashmore et al. (1995) used artificial mixtures
4 of grasses and forbs and transplanted swards of native calcareous grassland species and found
5 that, regardless of whether total biomass was adversely affected by exposures to O₃, higher
6 exposures progressively shifted species composition, usually at the expense of the forb species.
7 The observed shifts in competitive balance in favor of grasses is consistent with observations
8 that many grass species are less sensitive to O₃ than forbs. However, as previously shown by
9 Evans and Ashmore (1992), knowledge of the relative sensitivities to O₃ of the component
10 species grown in isolation or in monoculture does not always predict the impact of O₃ on the
11 components in a mixed culture.

12 Barbo et al. (1998) exposed an early successional plant community to O₃ in OTCs for two
13 growing seasons. Ozone decreased community structure features such as height of canopy,
14 vertical canopy density (layers of foliage), and species diversity and evenness. Surprisingly,
15 blackberry (*Rubus cuneifolius*), a species considered to be O₃-sensitive, replaced sumac (*Rhus*
16 *copallina*) canopy dominance. Barbo et al. (2002) also demonstrated the role of competition in
17 determining the impact of O₃ on loblolly pine (*Pinus taeda*). They reported that the increased
18 growth of natural competitors in OTCs using charcoal-treated air to reduce the ambient O₃
19 concentrations resulted in decreased pine growth. They noted that this is contrary to the
20 frequently reported increased growth observed in reduced O₃ levels in the absence of
21 interspecific competition.

22 Competitively disadvantaged trees of four clones of aspen (*Populus tremuloides*) exposed
23 to 1.5× ambient O₃ in a FACE facility over 4 years were proportionately more adversely affected
24 by O₃ than competitively advantaged or neutral trees (McDonald et al., 2002). However, one
25 clone of the disadvantaged trees demonstrated enhanced growth.

26 In summary, our present knowledge of how O₃ may affect the competitive interspecific
27 plant-plant relationships typifying the agricultural and natural worlds is very limited. However,
28 as noted in the 1996 AQCD (U.S. Environmental Protection Agency, 1996), “the development
29 and use of field exposure systems have permitted many recent studies of crop species to be
30 conducted at normal planting densities and hence have incorporated intraspecific competition as

¹⁶Forb: any non-grassy herbaceous species on which animals feed.

an environmental factor.” Such facilities were used in most of the studies of interspecific competition discussed above. But we are still a long way from being able to bridge the gap between small model competing systems and the realities of natural ecosystem complexity.

9.4.4 Physical Factors

The physical features of a plant’s aerial and edaphic environments exercise numerous controls over its growth and development. Thus, many of their effects may be modified by exposure to atmospheric oxidants and, alternatively, plants may modify responses to such exposures. As in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996), this section focuses on the defining features of plant microclimate: light, temperature, relative humidity (HR, or saturation vapor pressure deficit), and the presence and availability of water, especially in the soil. Monteith and Elston (1993) suggested that light energy and mass of water should be viewed as climatic resources, and the other two elements (temperature and saturation vapor pressure deficit) as rate modifiers that determine how fast the resources are used. The modifications of plant response by physical environmental factors has recently been reviewed by Mills (2002).

Another physical feature of the microclimate, wind and air turbulence, which affects the thicknesses of the boundary layers over leaves and canopies and, hence, affects gas exchange rates (including the fluxes of O₃ and other oxidants into the leaves) is discussed elsewhere (Section 9.3).

Physical features of the environment are also important components of larger-scale regional and global climates. However, the following discussions are confined to issues related to individual factors at the plant level; meso-scale effects are reviewed in Section 9.4.4.8, which addresses the issues of climate change interactions.

9.4.4.1 Light

Plants are the primary producers of biomass on the planet through their ability to capture light energy (by the process of photosynthesis) and convert it to the many forms of chemical energy that sustain their own growth and that of secondary consumers and decomposers. Light *intensity* is critical since the availability of light energy (a resource, *sensu* [Monteith and Elston, 1993]) governs the rate at which photosynthesis can occur, while light *duration* (i.e.,

1 photoperiod) profoundly effects development in many species. Although light *quality* (i.e., the
2 distribution of incident wavelengths) may also affect some physiological plants processes, there
3 is no evidence to indicate that such effects are of relevance to concerns over oxidant pollution,
4 except at the short wavelengths of UV-B. This topic is discussed in the context of climate
5 change in Section 9.4.8, and as a stress factor per se affected by atmospheric O₃ in Chapter 8.
6 However, as noted above and in the 1996 O₃ AQCD (U.S. Environmental Protection Agency,
7 1996), none of the features is controllable in natural field situations. A brief discussion of light
8 intensity-O₃ interactions is included in the review by Chappelka and Samuelson (1998).

9 The conclusion in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) that
10 low light intensities and short photoperiods tended to increase susceptibility to foliar O₃-injury
11 may still be valid, but this may or may not translate into adverse effects on growth. For
12 example, Tjoelker et al. (1993) found that, when seedlings of sugar maple (*Acer saccharum*),
13 a shade-tolerant species, were grown in 7% full sunlight, O₃ reduced shoot and root growth, but
14 had no significant effect in 45% sunlight (a 6-fold increase). In contrast, the reverse was
15 observed with a shade-intolerant hybrid poplar (*Populus tristis* × *balsamifera*), with the greater
16 impact of O₃ occurring in the higher light intensity treatment.

17 The greater sensitivity of maple in low light has also been confirmed in other studies.
18 Tjoelker et al. (1995) noted a greater O₃-induced inhibition of photosynthetic CO₂ assimilation in
19 shaded leaves than in leaves in full sunlight. However, in the absence of differences in stomatal
20 conductance, the effect was considered to be independent of O₃ flux; it appeared to be a
21 consequence of reduced chlorophyll contents and quantum efficiencies induced by O₃.
22 In contrast, Bäck et al. (1999), who also observed a greater inhibition of net photosynthesis by
23 O₃ in shaded leaves, reported decreased stomatal conductance. Although reduced conductance
24 might suggest reduced O₃ flux and, therefore, decreased adverse effects, the authors concluded
25 that the effects of reduced conductance was offset by long-term changes in leaf structure, leading
26 to less densely packed mesophyll cells and greater internal air space within the leaves.
27 Morphological differences between lower and upper crown leaves of black cherry (*Prunus*
28 *serotina*) have been suggested as the basis for the greater O₃-susceptibility of the lower crown
29 leaves (Fredericksen et al., 1995). Bäck et al. (1999) also observed accelerated foliar senescence
30 induced by O₃ on shaded leaves, a response also noted by Topa et al. (2001). Sensitivity to O₃
31 was found to be increased in shade- but not sun-leaves of shade-tolerant red oak (*Quercus rubra*)

(Samuelson and Edwards, 1993). Similarly, Mortensen (1999) observed that seedlings of mountain birch (*Betula pubescens*) grown in 50% shade suffered greater foliar injury from O₃ than those grown in full sunlight.

Not all shade-intolerant species exhibit greater reductions in photosynthesis and growth due to O₃ when grown in full sunlight. Higher than ambient levels of O₃ failed to inhibit photosynthesis in leaves of shade-intolerant yellow poplar (*Liriodendron tulipifera*) grown in nearly full sunlight (Tjoelker and Luxmoore, 1991). Greater foliar injury in the lower, shaded leaves of shade-intolerant black cherry (*Prunus serotina*) trees and saplings, was attributed (Fredericksen et al., 1996a) to higher stomatal conductance and greater O₃ uptake relative to net photosynthetic rate. However, in a 3-year study of Norway spruce seedlings (*Picea abies*) in OTCs. Wallin et al. (1992) observed that photosynthetic efficiency was more adversely affected by O₃ in high than in low light.

The suggestion of greater sensitivity to O₃ of shade-tolerant species in low-light conditions and the greater sensitivity of shade-intolerant species in high light is somewhat of an oversimplification when dealing with mature trees, for which light intensity varies considerably within the canopy because of shading. Chappelka and Samuelson (1998) noted that the interaction between sensitivity to O₃ and the light environment in forest trees is further complicated by developmental stage, with seedlings, saplings, and mature trees frequently giving different results. Topa et al. (2001) also cautioned that O₃ effects on leaf-level photosynthesis may be poor predictors of the growth responses of sugar maple in different light environments.

In high-light intensities, many species exhibit some degree of photoinhibition of the photosynthetic process through the overloading of the mechanisms that protect the photosynthetic reaction centers in the chloroplasts. Guidi et al. (2000) reported complex interactions between high-light intensities (inducing photoinhibition) and O₃ exposures in kidney bean (*Phaseolus vulgaris*) with high intensities tending to enhance the detrimental effect of O₃ on photosynthesis. One of the studies in the extensive European Stress Physiology and Climate Experiment-wheat (ESPACE-wheat) program (Bender et al., 1999), conducted in 1994 and 1995, included an investigation of the effects of climatic variables on yield response to O₃ using two simulation models, AFRCWHEAT2-O₃ and LINTULCC (Ewart et al., 1999; Van Oijen and Ewart, 1999). Among the observed trends, it was noted that relative yield loss of wheat due to elevated O₃ tended to increase with light intensity. In contrast, Balls et al. (1996) used ANNS to

investigate microclimatic influences on injury caused by O₃ to clover (*Trifolium subterraneum*) and found that, especially at mid-range cumulative O₃ exposures (350 to 500 ppb-h), injury tended to decrease with increasing light intensity. Similar observations by Davison et al. (2003) of foliar injury to wild populations of cutleaf cone flower (*Rudbeckia laciniata*) exhibiting a range of PAR levels within their canopies led the authors to conclude that the variation in injury symptoms observed was “unlikely to be due to differences in ozone flux and more likely to be due to variation in light.” Antonielli et al. (1997) found evidence indicating that the high sensitivity of the bioindicator tobacco cultivar *Nicotiana tabacum* cv Bel-W3 is partly determined by its high photosynthetic electron transport rates at high-light intensities, which exceed the capabilities of the plant to dissipate energy and oxyradicals.

The 1996 O₃ AQCD referred to the important role of light in controlling stomatal opening and suggested that light duration (i.e., photoperiod) might dictate the actual uptake of O₃ to some degree. However, it should also be noted that Sild et al. (1999) found that clover plants could suffer foliar injury even if they were exposed to O₃ during the dark period of the day-night cycle, when stomatal conductance is at its lowest.

A possible indirect effect of light intensity was noted by Reiling and Davison (1992) in their study of the O₃-tolerance of *Plantago major* plants grown from seeds collected from populations at 28 different sites in Britain. Ozone-tolerance, defined in terms of plant growth, was found to be a function of both previous O₃-exposure history and hours of bright sunshine during the year before the seeds were collected. However, the authors cautioned that, since tropospheric O₃-formation is itself dependent upon irradiation, the observation does not necessarily imply a direct effect of light intensity on the plants’ response to O₃.

The only recent studies concerning interactions with light quality appear to be those involving O₃ and UV-B as a component of climate change. These are dealt with in Section 9.4.4.8.2. The effects of photoperiod on response to O₃ or the converse do not appear to have received any recent attention.

Although the intensity, quality, and duration of light are not controllable in the natural world, the interactions of O₃ with light intensity, in particular, clearly have relevance to the growth of shade-tolerant and shade-intolerant species in mixed forest stands. It appears that the nature of light intensity-O₃ interactions may depend upon the type of light environment to which the species are best adapted, with increased light intensity increasing the sensitivity of light-

1 tolerant and decreasing the sensitivity of shade-tolerant species to O₃. Although there is
2 certainly some evidence to the contrary, this hypothesis is a reasonable summation of current
3 understanding with regard to O₃-light intensity interactions.

4 5 **9.4.4.2 Temperature**

6 “Temperature determines the start and finish and rate and duration of organ growth and
7 development” (Lawlor, 1998). Such processes depend on fundamental physiological activities
8 that are mostly enzyme-mediated and whose kinetics are directly affected by temperature. Since
9 the processes of enzyme deactivation and protein denaturation also increase as temperatures rise,
10 each enzymatic process has a unique optimum temperature range for maximal function.
11 However, the optima for different processes within the plant vary appreciably and, hence, the
12 optimum temperature range for overall plant growth is one within which all of the individual
13 reactions and vital processes are *collectively functioning optimally*, not necessarily *maximally*.
14 Furthermore, individual features of plant development (e.g., shoot and root growth, flowering,
15 pollen tube growth, fruit set, seed development) have different specific optima, so that
16 differential responses to temperature occur, leading to temperature-induced developmental
17 changes. For example, despite increased assimilation, increased temperatures may result in
18 decreased grain yields of crops such as wheat, because the growing season is effectively
19 shortened by a more rapid onset of senescence (Van Oijen and Ewart, 1999).

20 Rowland-Bamford (2000) noted that a plant’s response to temperature changes will depend
21 upon whether it is growing at its near optimum temperature for growth or its near maximum
22 temperature, and whether any increase in mean temperature results in temperatures rising above
23 the threshold for beneficial responses. Impairment by O₃ of any process may be thought of as
24 being analogous to a downward shift below and away from the temperature optimum or an
25 upward shift above and away from the optimum. Since a temperature rise toward the optimum
26 would result in a rate increase, the combined effects of O₃ and such an increase might neutralize
27 each other, while the effects of O₃ and a decrease in temperature would likely be additively
28 negative. Above the optimum temperature, the situations would be reversed with the effects of
29 increased temperatures and O₃ being additively negative, and decreasing temperatures
30 counteracting any negative effect of O₃. Thus, it is difficult to generalize about the interactions
31 of temperature and O₃ on overall plant responses such as growth in which the different

1 temperature-rate relationships of different growth components are merged, because they depend
2 upon the relationship of any temperature changes to the optimum for a species.

3 Studies of the effects of temperature on the impact of O₃ have increased recently because
4 of an increased need to understand the consequences of global warming as a component of
5 climate change. Direct interactions of temperature with O₃ are reviewed here, but the issues are
6 addressed again in Section 9.4.8.1 in relation to changes in atmospheric CO₂ levels.

7 The 1996 criteria document (U.S. Environmental Protection Agency, 1996) stressed the
8 interdependence of the temperature within the tissues of the leaf (where the various temperature-
9 sensitive processes occur) on three distinct components: the ambient air temperature, the heating
10 effect of incident infrared radiation during the photoperiod, and the evaporative cooling effect
11 caused by transpirational loss of water. It also cautioned that, especially in experiments using
12 controlled environment chambers, the effects of temperature could well be confounded with
13 those of humidity/vapor pressure deficit (VPD). Temperature and VPD are strongly interrelated,
14 and VPD plays an important role in regulating stomatal transpiration. Because of the role that
15 evaporative cooling plays in determining internal leaf temperatures, any factor that causes
16 stomatal closure and reduced conductance inevitably leads to increased leaf temperatures. Such
17 interactions add to the difficulties in distinguishing the effects of temperature from those of other
18 factors, as actual leaf temperatures are rarely measured and reported.

19 Despite these caveats, there is some evidence that temperature per se influences plant
20 response to O₃. For example, in rapid-cycling Brassica (*Brassica rapa*) and radish (*Raphanus*
21 *sativus*), marked O₃-inhibitions of growth were observed at low root temperatures (13 °C) but
22 not at 18°C (Kleier et al., 1998, 2001). With regard to air temperature, this was included in the
23 range of micrometeorological variables studied in several recent extensive field studies and was
24 found to have a significant effect on response to O₃ in most cases. Ball et al. (1996) used ANNs
25 in an analysis of the growth of clover (*Trifolium subterraneum*) and concluded that light and
26 VPD had greater influences than temperature on the visible injury response to O₃. However, in
27 three studies with different cultivars of white clover (*Trifolium repens*), temperature was found
28 to be important to the growth response. Ball et al. (1998) exposed *T. repens* cv. Menna to
29 ambient O₃ in OTCs at 12 European sites at a range of latitudes and altitudes from 1994 to 1996.
30 The impact of O₃ on growth was determined as the ratio of growth with and without treatment
31 with the O₃-protectant, EDU (see Section 9.2). Artificial neural network analysis showed that O₃

1 exposure (measured as the AOT40 index, see Section 9.4.6), VPD, and temperature were
2 consistently the three most important variables governing response to O₃ over a range of
3 different ANN models. However, the authors did not describe the form of the O₃-response
4 relationship with temperature. Similar observations were reported by Ball et al. (2000) and Mills
5 et al. (2000) for O₃-sensitive and O₃-tolerant clones of *T. repens* cv. Regal, grown at 14 to
6 18 European locations from 1995 to 1998. In both studies, the impact of O₃ was measured as the
7 sensitive/tolerant growth (biomass) ratio. Although Ball et al. (2000) found temperature to be
8 less important than O₃ exposure and VPD, Mills et al. (2000) found temperature to be the most
9 important input variable after O₃ exposure (AOT40). In both cases, the adverse effect of O₃
10 increased with increasing temperature.

11 A study of black cherry (*Prunus serotina*) seedlings and mature trees in Pennsylvania,
12 using micrometeorological variables aimed to predict O₃ uptake, found temperature to be
13 unimportant (Fredericksen et al., 1996b), but in the study of populations of *Plantago major*
14 referred to in Section 9.4.4.1, Reiling and Davison (1992) noted a weak, positive correlation
15 between mean temperature at the collection site and O₃ tolerance (based on growth rate) of the
16 different populations. In contrast, Danielsson et al. (1999) collected genotypes of *Phleum*
17 *arvense* from a wide range of Nordic locations and found a positive effect of temperature on the
18 growth of genotypes from locations with higher summer temperatures, but sensitivity to O₃ did
19 not vary systematically with geographic location.

20 Van Oijen and Ewart (1999) studied the effects of climatic variables on the response to O₃
21 in the ESPACE-wheat program, based on two distinctive simulation models (AFRCWHEAT2-
22 O₃ and LINTULCC, [Ewart et al., 1999]) and noted that although the relative yield loss of wheat
23 due to elevated O₃ tended to increase with temperature, the effect was of minor significance.

24 In contrast to the variable results obtained in studies of the effects of temperature on
25 response to O₃, the corollary effect of O₃ exposure on subsequent sensitivity to low temperature
26 stress, noted in the 1996 criteria document, is well recognized. In reviewing low temperature-O₃
27 interactions, Colls and Unsworth (1992) noted that winter conditions produce three kinds of
28 stress: desiccation, chilling or freezing temperatures, and photooxidation of pigments. Of these,
29 they suggested that while the first two were important, the last may play a particularly significant
30 role because the “combination of high irradiance and low temperatures permits a buildup of free
31 radicals in leaf tissue, and these free radicals then attack chlorophyll.” Chappelka and Freer-

Smith (1995) suggested that the injury and losses to trees caused by this delayed impact of O₃ may be equally or more important than the direct impacts of O₃ on foliage of visible injury and necrosis, or the disruption of key physiological processes such as photosynthesis. In this context, the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) referred to the conceptual framework (Eamus and Murray, 1991), which is still valid: brief periods of mild temperatures in the severest winters result in dehardening; O₃ decreases frost hardiness per se, but also increases the predisposition to dehardening; dehardening places O₃-exposed trees at greater risk from subsequent low temperatures. However, no quantified models of these effects have yet appeared.

The 1996 criteria document also noted that O₃ adversely affects cold hardiness of herbaceous species. More recently, Foot et al. (1996, 1997) observed winter injury and decreased growth in low-growing perennial heather *Calluna vulgaris* exposed to O₃ (70 ppb, 8 h/day, 5 days/week for 6 months) during the winter (6.8 °C mean), but found no significant effects of the same exposures during the summer (12.3 °C mean). Although Potter et al. (1996) observed a similar situation with the moss *Polytrichum commune*, the reverse was found with the moss *Sphagnum recurvum*.

In summary, unequivocal evidence exists that O₃ causes sensitization to the adverse effect of low temperatures, but there is no clear pattern to evidence regarding the effects of temperature on O₃ response. The many contradictory responses to temperature and O₃ probably reflect our lack of detailed knowledge of the temperature optima for the different growth components of the studied species. The topic of temperature-oxidant interactions is revisited later in Section 9.4.4.8 in the context of global warming as a feature of climate change.

9.4.4.3 Humidity and Surface Wetness

The moisture content of the ambient air (or its VPD) is a rate modifier (*sensu* Monteith and Elston, 1993) and an environmental regulator of stomatal conductance. Both of the previous criteria documents (U.S. Environmental Protection Agency, 1986, 1996) concluded that the weight of evidence indicated that high RH (= low VPD) tended to increase the adverse effects of O₃, principally because the stomatal closure induced in most situations by O₃ is inhibited by high RH, leading to increased O₃ flux into the leaves.

Recent reports have confirmed this role of RH. The studies by Ball et al. (1995, 1996, 1998) showed that VPD was an important determinant of O₃-induced injury and reduced growth in two species of clover, *Trifolium repens* cv. Menna and *T. subterraneum*. However, Mills et al. (2000) found it to be unimportant in the case of *T. repens* cv. Regal. Such difference between cultivars is not unexpected since considerable differences also occur among species and genera. For example, Bungener et al. (1999a) studied 26 Swiss grassland species and found clear evidence that O₃ injury increased with decreased VPD (i.e., increased RH) in only eight species. However, the 1995 data from the European cooperative study of O₃ injury, which involved 28 sites in 15 countries and six crop species, led to the development of two 5-day critical-level scenarios involving O₃-exposure (calculated as the AOT40 index) and mean VPD (0930-1630h): 200 ppb-h at > 1.5 kPa, and 500 ppb-h at < 0.6 kPa (Benton et al., 2000).

With forest tree species, Fredericksen et al. (1996b) found significant correlations between stomatal conductance of black cherry (*Prunus serotina*) leaves and RH (+ve) and VPD (-ve), and studies on free-standing Norway spruce (*Picea abies*) and larch (*Larix decidua*) showed that although ambient VPD was highly positively correlated with ambient O₃ concentration, increased VPD caused stomatal closure, reducing O₃ uptake and impact (Wieser and Havranek, 1993, 1995).

Surface wetness may affect the response to O₃ through its direct effects on deposition to the surface and through changes in RH. Effects on the deposition of O₃ have been reviewed (Cape, 1996). A surface film of water on leaves was found to increase O₃ deposition in four studies involving field-grown grape (*Vitis vinifera*) (Grantz et al., 1995), red maple (*Acer rubrum*) (Fuentes and Gillespie, 1992), deciduous forest dominated by largetooth aspen (*Populus grandidentata*) and red maple (Fuentes et al., 1992), and clover-grass mixed pasture (*Trifolium pratense*, *Phleum pratense*, and *Festuca pratensis*) (Pleijel et al., 1995). In each case, the increased deposition could be attributed partly to an increased stomatal conductance through the abaxial (lower) surface and partly to uptake into the aqueous film on the adaxial (upper) surface. In contrast, decreased deposition was noted by Grantz et al. (1997) with field-grown cotton (*Gossypium hirsutum*). Since cotton is amphistomatous, with functional stomata on both leaf surfaces, it was suggested that, in this case, the water layer effectively sealed the adaxial surface stomata, more than offsetting any increase in conductivity of the stomata in the abaxial surface. However, none of the studies investigated the consequences of the differences in deposition.

1 Although it could be inferred that, with part of any increased deposition being the result of
2 increased O₃ flux into the leaves, there would be the likelihood of increased O₃ adverse effects,
3 as suggested by earlier studies (Elkiey and Ormrod, 1981) that, by misting bluegrass (*Poa*
4 *pratensis*) during exposure to O₃, injury was significantly increased.

5 To conclude, the effects of high RH (low VPD) and surface wetness have much in
6 common, as they both tend to enhance the uptake of O₃, largely through effects on stomata,
7 leading to increased impact.

9 **9.4.4.4 Drought and Salinity**

10 The 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) concluded that the
11 available evidence clearly indicated that exposure to drought conditions could reduce the adverse
12 effects of O₃ on the growth of herbaceous and woody plants, but it also noted that no quantitative
13 models of the O₃-soil moisture deficit (SMD) interaction had yet appeared in print.
14 Nevertheless, the “protective” effect was inconsistent, and only appeared when SMD was
15 accompanied by high evaporative demand. Since that time, further studies have confirmed the
16 interaction, and simulation models have begun to appear. Mills (2002) has recently provided a
17 brief review of the topic.

18 With regard to herbaceous species, Vozzo et al. (1995) observed less O₃-induced injury and
19 suppression of net photosynthesis and growth in water-deficient soybean (*Glycine max*) than in
20 well-watered plants. In several studies with wheat (*Triticum aestivum*), on the other hand,
21 although adverse effects of both O₃ and SMD were noted, they were consistently additive
22 (Bender et al., 1999; Fangmeier et al., 1994b, 1994a; Ommen et al., 1999).

23 In attempting to model the stomatal conductance of wheat in relation to O₃ and soil
24 moisture, Grütters et al. (1995) found that although O₃-induced stomatal closure was enhanced by
25 SMD, reducing O₃ uptake, the R² of the overall model was only 0.40, indicating that other
26 significant factors or relationships were involved.

27 With regard to native vegetation, Bungener et al. (1999a) used mixed plantings of 24 Swiss
28 grasses, herbs, and legumes and observed that, although O₃-drought interactions were species-
29 specific, they tended to reflect stomatal functioning. They found that SMD reduced O₃ injury in
30 two clovers (*Trifolium repens* and *T. pratensis*) and two grasses (*Trisetum flavescens* and
31 *Bromus erectus*), but noted no interactions in the other 20 species. With relative growth rate as

the measure of response to O₃, interactions with SMD were noted in only three species: *Trifolium repens* and two weedy herbs, *Knautia arvensis* and *Plantago lanceolata* (Bungener et al., 1999b). Although this variability in response among species was noted in the review by Davison and Barnes (1998), they also pointed out that in severely droughted regions of Europe, notably in Greece and Spain, O₃-induced injury and growth reductions were common on many (usually irrigated) crops, but there were virtually no records of injury symptoms in wild species.

Thus, the situation with herbaceous species is essentially unchanged from 1988 when Heagle et al. summarized the extensive NCLAN experiments that incorporated water stress as a variable: “SMD can reduce the response of crops to O₃ under some conditions but not under other conditions. Probably the occurrence of O₃ by SMD interactions was dependent on the degree of SMD-induced plant moisture stress.”

With regard to trees, O₃ interactions with soil water availability have been discussed in several recent reviews: Chappelka and Freer-Smith (1995), who focused on O₃-induced predisposition to drought stress; Johnson et al. (1996b); Chappelka and Samuelson (1998); and Skärby et al. (1998).

Several recent studies with conifers have yielded mixed results. No interactions with drought were observed by Broadmeadow and Jackson (2000) on Scots pine (*Pinus sylvestris*), by Karlsson et al. (2002) on Norway spruce (*Picea abies*), or by Pelloux et al. (2001) on Aleppo pine (*P. halepensis*). More recently, Le Thiec and Manninen (2003) reported that drought reduced O₃-induced growth suppression of Aleppo pine seedlings. Panek and Goldstein (2001) inferred less impact of O₃ on droughted Ponderosa pine (*P. ponderosa*), and Van den Driessche et al. (1994) reported that drought reduced injury and O₃-induced ethylene release by *Picea abies*. But Karlsson et al. (1997), in a comparative study of fast- and slow-growing clones of *P. abies*, only observed a drought-induced reduction of O₃-inhibited root growth in the fast-growing clone. In contrast, Grulke et al. (2002) noted a synergistic interaction between O₃ and drought stress on gross photosynthesis of *Pinus ponderosa*, and Wallin et al. (2002) reported a synergistic growth response of *Picea abies* in the third year of a 4-year study. A similar response was noted by Dixon et al. (1998) with the Istebna strain of *P. abies*.

With broad-leaved trees, studies of Durmast oak (*Quercus petraea*) (Broadmeadow et al., 1999; Broadmeadow and Jackson, 2000) and European ash (*Fraxinus excelsior*) (Broadmeadow and Jackson, 2000; Reiner et al., 1996) showed that drought provided partial protection against

1 O₃-induced growth reduction. Although European beech (*Fagus sylvatica*) is reportedly an
2 O₃-and drought-sensitive species, neither Pearson and Mansfield (1994) nor Broadmeadow et al.
3 (1999) observed any interactions between these stresses, while Dixon et al. (1998) observed
4 partial protection. Pääkönen et al. (1998) observed only additive effects in a sensitive clone of
5 birch (*Betula pendula*). However, the experiments of Schaub et al. (2003) and the survey by
6 Vollenweider et al. (2003) on black cherry (*Prunus serotina*) clearly indicate antagonism between
7 drought and O₃ stresses on this species.

8 With regard to the converse effect, in a critical review of the evidence for predisposition to
9 drought stress being caused by O₃, Maier-Maercker (1998) supported the hypothesis and
10 suggested that the effects were caused by the direct effects of O₃ on the walls of the stomatal
11 guard and subsidiary cells in the leaf epidermis, leading to stomatal dysfunction.

12 The Plant Growth Stress Model (PGSW) developed by Chen et al. (1994) is a physiology-
13 based process model which includes drought among several environmental variables.
14 Simulations for Ponderosa pine (*Pinus ponderosa*) incorporated antagonistic effects between O₃
15 and drought stresses, i.e. partial protection, although Karlsson et al. (2000) have since
16 emphasized that drought-induced “memory effects” should be considered when developing
17 simulation models incorporating stomatal conductance.

18 Retzlaff et al. (2000) used the single-tree model, TREEGRO, to simulate the combined
19 effects of O₃ and drought on white fir (*Abies concolor*). Although simulated reductions in
20 precipitation $\geq 25\%$ reduced growth, they also reduced O₃ uptake (and impact). But lesser
21 reductions in precipitation combined synergistically with O₃ stress to reduce growth, leading the
22 authors to conclude that moderate drought may not ameliorate the response of white fir to O₃.

23 On a much larger scale with a modified forest ecosystem model (PnEt-II) incorporating O₃-
24 response relationships for hardwood species, Ollinger et al. (1997) showed how predicted
25 changes in net primary production and mean wood production in the northeastern U.S. hardwood
26 forests due to O₃ would be reduced (but not countered or reversed) by drought stress, particularly
27 in the southern part of the region. This geographic distribution of the effect was substantiated by
28 the work of Lefohn et al. (1997) on the risk to forest trees in the southern Appalachian
29 Mountains, based on localized estimates of O₃ levels and SMD. The TREEGRO and ZELIG
30 models were combined by Laurence et al. (2001) to predict the impacts of O₃ and moisture (as
31 precipitation) on the growth of loblolly pine (*Pinus taeda*) and yellow poplar (*Liriodendron*

1 *tulipifera*). Based on O₃ and precipitation data from three sites in the eastern United States, the
2 six model regressions developed for the two species included both positive and negative
3 coefficients for O₃ exposure and precipitation as determinants of growth.

4 As noted in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996), the effects
5 of soil salinity are similar to those of SMD. In a study of rice (*Oryza sativa*) cultivars of
6 differing sensitivity to salinity, Welfare et al. (1996) noted that although both O₃ and salinity
7 reduced many features of growth additively, antagonistic interactions were only seen for leaf
8 length and potassium accumulation. Similarly, a recent study on chickpea (*Cicer arietinum*)
9 found no interactions with regard to most components of biomass accumulation (the effects
10 of O₃ and salinity were additive), but with root growth, salinity suppressed the adverse effects
11 of O₃.

12 In summary, the recently described interactions of O₃ and drought/salinity stresses are
13 consistent with the view that, in many species, drought/salinity reduces the impact of O₃ but O₃
14 increases sensitivity to drought stress, i.e., the type of response is determined by the sequence of
15 stresses. However, synergisms have also been observed and any antagonisms are species-
16 specific and unpredictable in the absence of experimental evidence. In no case has an
17 antagonism been found to provide complete protection.

18 19 **9.4.5 Nutritional Factors**

20 The 1996 criteria document (U.S. Environmental Protection Agency, 1996) noted that the
21 large number of macro- and micronutrients and the wide range of species had prevented
22 experimental investigation of all but a few cases of nutrient-O₃ interactions and most of these
23 concerned nitrogen (N) and crops or forest tree species. The document also provided a
24 comprehensive tabulation of the results of the relevant studies up to 1992.

25 The suboptimal supply of mineral nutrients to plants leads to various types of growth
26 reductions. The consequences of suboptimal nutrition might, therefore, be expected to have
27 some similarities to those of O₃ exposure. One might expect nutritional levels below the
28 optimum either to amplify any effects of O₃ or at least lead to additive responses. The difficulty
29 with this suggestion is that the available information has mostly been obtained from
30 experimentation conducted using two or more arbitrarily selected levels of fertility with little or
31 no regard to optima. Hence, it is not surprising that there have been contradictory reports, even

1 among studies with the same species or cultivars conducted by different workers at different
2 locations using different soils or soil mixes.

3 There appear to have been no recent studies on O₃ interactions with specific mineral
4 nutrients other than N. Hence, the previous conclusions are still valid, viz. that increasing levels
5 of the major elements potassium (K) and sulfur (S) usually reduce the impact of O₃, or,
6 deficiency increases susceptibility, whereas increased phosphorus (P) usually increases injury,
7 or, deficiency decreases susceptibility.

8 However, with N, a relationship to the optimum is usually demonstrable. Several earlier
9 studies of O₃ × N interactions reported that the adverse effects of O₃ on growth were greatest at
10 the optimum and decreased with increasing N-deficiency, a finding supported by the work on
11 aspen (*Populus tremuloides*) of Pell et al. (1995), who also confirmed that excess N decreased O₃
12 impact on growth. Similarly, the adverse effects of O₃ on growth rate in wheat (*Triticum*
13 *aestivum*) diminished with decreased N supply (Cardoso-Vilhena and Barnes, 2001). However,
14 the effects of N are far from consistent. For example, Greitner et al. (1994) reported that O₃ and
15 N-deficiency acted additively in aspen in reducing leaf surface area and rate of photosynthesis,
16 Bielenberg et al. (2001) reported that the rate of O₃-induced senescence was increased by
17 N-deficiency in hybrid poplar (*Populus trichocarpa* × *maximovizii*), and Pääkkönen and
18 Holopainen (1995) observed the least adverse effects of O₃ on birch (*Betula pendula*) at
19 optimum N-fertility levels. With cotton (*Gossypium hirsutum*), increased N-levels more than
20 overcame the adverse effect of O₃ on growth and boll yield (Heagle et al., 1999). In view of
21 these contradictions, one may conclude that other, unrecorded factors may have contributed to
22 the various findings. Thus, much remains unclear about O₃ × N-fertility interactions.

23 There have been two recent studies on the effects of overall soil fertility. Whitfield et al.
24 (1998) observed that low general fertility increased O₃ sensitivity in selections of *Plantago*
25 *major*. At the biochemical level, well-fertilized birch (*Betula pendula*) saplings were found to
26 be less adversely affected by O₃ than nutrient-stressed plants (Landolt et al., 1997).

27 TREEGRO model simulations of the growth of red spruce (*Picea rubra*) in conditions of
28 nutrient deficiency and O₃ stress showed that, in combination, the two stresses acted less than
29 additively (Weinstein and Yanai, 1994). Minimal amelioration by nutrient deficiency was
30 predicted with Ponderosa pine (*Pinus ponderosa*).

Plants may also obtain N and S from airborne sources such as NO_x , HNO_3 , NO_3^- , SO_2 , and SO_4^{2-} , although, depending upon their concentration, these may also be phytotoxic. In various parts of the world, the deposition of N and S in these forms contributes significantly to the levels of nutritionally available N and S in soils. Such depositions may, in turn, influence the impact of O_3 on sensitive species through their roles as nutrients independent of any interactions that may occur because of their acidic properties (see Section 9.4.6.5). For example, Takemoto et al. (2001) recently reviewed the situation in southern California's mixed conifer forests and noted that, where N-deposition is appreciable, its combination with O_3 is causing a shift in Ponderosa pine (*Pinus ponderosa*) biomass allocation towards that of deciduous trees, with increased needle drop so that only 1- and 2-year needle classes overwinter. Such changes are having significant consequences to the balance of the forest ecosystem and are discussed more fully in Section 9.5.

Of the micronutrient elements, only manganese (Mn) appears to have been studied recently. In beans (*Phaseolus vulgaris*) Mn-deficiency increased O_3 toxicity, despite causing reduced O_3 uptake (through decreased stomatal conductance) and inducing increased levels of Mn-SOD (Mehlhorn and Wenzel, 1996).

In view of the foregoing, it is impossible to generalize about the interactions of soil fertility with O_3 . While this is especially true of the interactions involving soil nitrogen, for which there is much conflicting evidence, the interactions with other nutrients need much more thorough investigation than has occurred to date, before any clear patterns become apparent.

9.4.6 Interactions with Other Pollutants

The ambient air may be polluted by gases other than O_3 and its photochemical oxidant relatives. In particular, industrial, domestic, and automobile emissions and accidents can lead to significant atmospheric concentrations of gases such as sulfur dioxide (SO_2), nitric oxide (NO), and nitrogen dioxide (NO_2), collectively referred to as NO_x , both locally and regionally. Local releases of gases such as hydrogen fluoride (HF), hydrogen chloride (HCl), and chlorine (Cl_2) may result from industrial emissions and accidents. Agricultural fertilizer and manure usage can lead to significant increases in ambient ammonia (NH_3). The sulfur and nitrogen oxides may undergo reactions in the atmosphere leading to the formation of sulphate (SO_4^{2-}) and nitrate (NO_3^-) ions and resultant acid deposition.

1 The 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) discounted much of
2 the early research on pollutant combinations, because of its lack of resemblance to the ambient
3 experience: the concentrations used were unrealistically high or the exposure regimes employed
4 almost invariably used gas mixtures, whereas Lefohn et al. (1987) showed that the co-occurrence
5 patterns of significant levels of O₃ with SO₂ or NO₂ in the United States were most frequently
6 sequential or partially sequential with overlap; only rarely were they entirely concurrent. On the
7 other hand, O₃ and peroxyacetylnitrate (PAN) frequently co-occur, as both form photochemically
8 under similar conditions.

9 To the list of reviews mentioned in the 1996 criteria document should be added the more
10 recent ones by Barnes and Wellburn (1998), Robinson et al. (1998), and Fangmeier et al. (2002),
11 which also explore some of the potential mechanisms underlying pollutant-pollutant interactions.
12

13 **9.4.6.1 Oxidant Mixtures**

14 In 1998, Barnes and Wellburn noted that virtually no information existed on the effects on
15 plants of concurrent exposures to O₃ and other components of photochemical oxidant other than
16 PAN. The situation has not changed since their review appeared, and the topic appears to have
17 attracted no research interest since before the 1996 O₃ AQCD (U.S. Environmental Protection
18 Agency, 1996). The continuing conclusion must, therefore, be that, from the limited information
19 available, the two gases appear to act antagonistically, with O₃ raising the threshold for the
20 visible injury response to PAN, and PAN reducing the harmful effects of O₃.
21

22 **9.4.6.2 Sulfur Dioxide**

23 In reviewing O₃ × SO₂ interactions, Barnes and Wellburn (1998) remarked: “The outcome
24 of exposure to this combination of pollutants has probably been the most studied, yet is one of
25 the least understood.” More recent studies have only added to the conflicts referred to in the
26 1996 criteria document (U.S. Environmental Protection Agency, 1996), rather than resolve them.
27 For example, Diaz et al. (1996) reported that, after a year of daily exposures of Aleppo pine
28 (*Pinus halepensis*) seedlings to 50 ppb O₃ and/or 40 ppb SO₂, the combination of pollutants
29 synergistically reduced shoot and root growth and impaired mycorrhizal colonization of the
30 roots. With tomato (*Lycopersicon esculentum*), on the other hand, effects on growth ranged
31 from synergistic at low exposures (50 ppb) to antagonistic at exposures of 200 ppb of each gas

(Khan and Khan, 1994). Although various physiological measurements were made in these and earlier studies, it has not been possible to determine any consistent mechanism or mechanisms that might account for the conflicting results.

Since the information available about $O_3 \times SO_2$ interactions appears to be highly dependent upon species, the type of response measured, and the experimental protocol used, it would still appear prudent to heed the statement of Heagle et al. (1988) in their summary of the studies undertaken in 12 field experiments over several years within the NCLAN program: “There were no cases where O_3 and SO_2 interactions *significantly* affected yield.” (emphasis added.)

9.4.6.3 Nitrogen Oxides, Nitric Acid Vapor, and Ammonia

The major oxides of nitrogen that occur in ambient air are nitrous oxide (N_2O), nitric oxide (NO) and nitrogen dioxide (NO_2), of which the latter two (conveniently symbolized as NO_x) are particularly important in connection with O_3 , because they are components of the reaction mix that leads to photochemical O_3 formation and because they can interact with O_3 -responses. Their reactions in the atmosphere can also lead to the occurrence of nitric acid vapor (HNO_3) in ambient air. The other major N-containing contaminant of ambient air in many parts of the world is ammonia (NH_3), largely released through agricultural practices.

Despite various combinations of O_3 and NO_x being probably the most common air pollutant combinations found in the field, Barnes and Wellburn (1998) noted that they have been little studied. Much early work with O_3 and NO_x focused on $O_3 \times NO_2$ interactions and can be discounted, because of the unrealistic concentrations employed and their use as mixtures rather than in types of sequences. The 1996 O_3 AQCD (U.S. Environmental Protection Agency, 1996) concluded that evidence from studies involving concurrent exposures to both O_3 and NO_x at realistic concentrations was so fragmented and varied that no firm conclusions could be drawn as to the likelihood and nature such interactions. However, the few recent investigations taken together with the earlier data are now beginning to reveal a pattern of response.

With regard to NO, Nussbaum et al. (1995a; 2000b) reported their findings with concurrent exposures to NO and O_3 and observed that, at low O_3 levels, NO tended to act similarly to O_3 by increasing the scale of responses such as growth reductions. However, in ambient air in which O_3 is a dominant factor, the effects of NO were usually found to be negligible due to low levels,

1 although the authors admitted that the effects observed were confounded by the inevitable
2 O₃-induced oxidation of NO to NO₂.

3 Two possible mechanisms whereby NO may influence plant response to O₃ are suggested
4 by recent biochemical studies. First, there is growing evidence for the role of NO as a signaling
5 agent in plants that can induce defense responses to a range of biotic and abiotic stressors
6 (Beligni and Lamattina, 2001; Neill et al., 2002). Second, a role for NO as an antioxidant
7 scavenger of reactive oxygen species has been demonstrated by Beligni and Lamattina (2002) in
8 potato (*Solanum tuberosum*) leaves and chloroplasts. However, both of these cases concern
9 endogenously synthesized NO, and it must be noted that in none of these or other reports of
10 studies of NO signaling have the authors considered the potential significance of exogenous NO
11 in ambient air.

12 An independent case for O₃ × NO interactions comes from Mills et al. (2000). The ANN
13 model developed to predict the O₃ effects on white clover (*Trifolium repens*) biomass based on
14 experiments at 18 locations throughout Europe found that the minimum daily NO concentration
15 (at 5 p.m.) was a significant contributor to adverse effects.

16 Turning to NO₂, Maggs and Ashmore (1998) found that, although concurrent but
17 intermittent exposures of Bismati rice (*Oryza sativa*) revealed no significant growth interactions,
18 NO₂ reduced the rate of O₃-induced senescence, an antagonistic response possibly related to
19 enhanced N-metabolism.

20 With regard to sequential exposures, two studies on gene activation in tobacco (*Nicotiana*
21 *tabacum*) revealed that NO₂ counteracted the effect of O₃ in reducing mRNA levels for three
22 genes encoding photosynthetic proteins (Bahl and Kahl, 1995) and tended to counteract the
23 O₃-induced enhancement of defense-protein gene activation (Bahl et al., 1995). However,
24 despite compelling evidence for significant interactive effects provided by earlier studies
25 (Bender et al., 1991; Goodyear and Ormrod, 1988; Runeckles and Palmer, 1987), the only recent
26 investigation of growth effects seems to have been that of Mazarura (1997) using sine-wave
27 exposure profiles. He found that although 4 weeks of twice daily 3-h exposures to NO₂ (120 ppb
28 peak concentrations) slightly stimulated growth of radish (*Raphanus sativa*) and while daily 6-h
29 exposures to O₃ (120 ppb peak concentration) did not significantly reduce growth, the daily
30 sequence, NO₂ - O₃ - NO₂, led to a 13% drop in dry matter production.

1 The combined evidence to date, therefore, suggests that, in leguminous species, the effects
2 of these sequences are antagonistic with NO₂ tending to reduce (or reverse) the negative effects
3 of O₃ on growth, while the effects are increased in other species. These conclusions differ from
4 those of Barnes and Wellburn (1998) who suggested that sequential exposures tended to result in
5 antagonistic effects (largely based on the summary by Bender and Weigel, 1992), whereas
6 simultaneous exposures were likely to lead to synergistic responses. With disagreements both
7 among the data and their interpretation, it is not possible to determine the circumstances under
8 which specific interactions of O₃ and NO₂ may occur, but there is no reason to doubt the validity
9 of the individual findings of each study. Far more systematic investigation is needed to clarify
10 the situation.

11 There appear to have been no studies of O₃ interacting with HNO₃ in the vapor phase.
12 However, in the southern California montane forests (Takemoto et al., 2001), in Sweden (Janson
13 and Granat, 1999), and elsewhere, significant amounts of N are deposited in this form because of
14 the vapor's high deposition velocity. As a consequence, although much of it ultimately reaches
15 the ground through leaching and leaf fall and enters the soil as nitrate (NO₃⁻), it may also be used
16 as a N source by the foliage itself. This nutritional role is independent of any contribution that
17 HNO₃ vapor may make to acidic deposition. Indirect interactions with the effects of O₃ through
18 N-deposition of NO_x, HNO₃, and NH₃ are related to the interactions of O₃ with N as a nutrient,
19 and have recently been examined in the review by Takemoto et al. (2001). The 1996 O₃ AQCD
20 (U.S. Environmental Protection Agency, 1996) stated that the evidence available at that time led
21 to estimates of total forest dry deposition, including HNO₃, ranging from 5.7 to 19.1 kg N ha⁻¹
22 year⁻¹ (Taylor, Jr. et al., 1988). However, Takemoto et al. (2001) pointed out that in parts of the
23 mid-elevation forests of southern California, dry deposition rates may reach more than 40 kg N
24 ha⁻¹ year⁻¹. As a result, some locations have seen the conversion from N-limited to N-saturated
25 forests. The concern for California's forests is well-stated by Takemoto et al.: "As potential
26 modifiers of long-term forest health, O₃ is a stressor and N deposition is an enhancer of
27 ponderosa/Jeffrey pine physiology and growth (Grulke and Balduman, 1999). The progression
28 toward a deciduous growth habit, higher shoot:root biomass ratios, increasing depths of litter,
29 tree densification, and elevated NO₃⁻ levels in soil and soil solution, all point to the replacement
30 of pine species with nitrophilous, shade- and O₃-tolerant tree species, such as fir and cedar
31 (Minnich et al., 1995) (Minnich, 1999)."

1 Few studies have been reported of interactions of O₃ with NH₃. The 1996 criteria
2 document made reference to the work on bean (*Phaseolus vulgaris*) by Tonneijck and van Dijk
3 (1994, 1998). Although NH₃ alone tended to increase growth and O₃ alone to inhibit it, one
4 interaction was noted (Tonneijck and van Dijk, 1994) on the number of injured leaves. Dueck
5 et al. (1998) studied the effects of O₃ and NH₃ on the growth and drought resistance of Scots
6 pine (*Pinus sylvestris*). Significant interactions were found for some growth features, but there
7 were no consistent patterns of the effects of NH₃ on O₃ response or vice versa. However, O₃ was
8 found to ameliorate the enhancement of drought stress caused by NH₃.

9 At this time there is insufficient information to offer any general conclusions about the
10 interactive effects of O₃ and NH₃.

11 12 **9.4.6.4 Hydrogen Fluoride and Other Gaseous Pollutants**

13 Although HF and other fluorides are important local air pollutants associated with
14 aluminum smelting and superphosphate fertilizer manufacture, no studies of possible interactions
15 with oxidants appear to have been reported since that of MacLean (1990). He found that HF
16 retarded the accelerated senescence and loss of chlorophyll resulting from O₃ exposure in corn
17 seedlings. However, such an isolated observation cannot be taken to indicate that HF can reduce
18 the impact of O₃ on other species or even that the effect would ultimately have led to an effect on
19 mature plants.

20 21 **9.4.6.5 Acid Deposition**

22 The deposition of acidic species onto vegetation may elicit direct effects on the foliage or
23 indirect effects via changes induced in the soil. The 1996 O₃ AQCD (U.S. Environmental
24 Protection Agency, 1996) included an extensive listing of investigations into the effects of O₃
25 and acid deposition (usually in the form of simulated acid rain, SAR) on plant growth and
26 physiology. The majority of studies found no effects of SAR or acid mists or fogs at pH values
27 greater than about 3.0 and no interactive effects with O₃. (In ambient air, pH values less than 3.0
28 have rarely been reported.) In the few reports in which significant interactions were found, most
29 were antagonistic and were explained as probably being the result of increased fertility due to
30 nitrate and sulfate supplied in the SAR.

1 Although numerous reviews have recently appeared (e.g., Bussotti and Ferretti, 1998;
2 Flückiger et al., 2002; Fox and Mickler, 1996; Nussbaum et al., 1999; and Sheppard and Cape,
3 1999), the shift in interest in air pollution effects away from acid deposition has resulted in little
4 new research having been reported over the past 10 or so years. In most of the reported studies,
5 no effects due to the O₃ exposures, the SAR treatments used, or their combinations were
6 observed, e.g., Baker et al. (1994) on loblolly pine (*Pinus taeda*); Laurence et al. (1997), and
7 Vann et al. (1995) on red spruce (*Picea rubens*); and Laurence et al. (1996) on sugar maple
8 (*Acer saccharum*). Branch chamber studies of 12-year old Ponderosa pine (*Pinus ponderosa*)
9 trees by Momen et al. (1997, 1999) revealed no O₃ effects or interactions. With red spruce
10 (*Picea rubens*), Sayre and Fahey (1999) noted no effects of O₃ on the foliar leaching of Ca or
11 Mg, which only became significant with SAR at pH 3.1. Izuta (1998) observed no interactions
12 with Nikko fir (*Abies homolepis*), although SAR at pH 4.0 reduced dry matter. Shan et al. (1996)
13 reported adverse effects of O₃ but none attributable to SAR on the growth of *Pinus armandi*.

14 With herbaceous species, Ashenden et al. (1995) noted significant antagonistic interactions
15 of O₃ and acid mist in white clover (*Trifolium repens*), in which the adverse effect of low pH was
16 countered by O₃. In contrast, Ashenden et al. (1996) found that, although pH 2.5 mist caused a
17 significant stimulation of the growth of ryegrass (*Lolium perenne*) attributed to a fertilizer effect,
18 and O₃ caused reduced growth, there was no interaction. Bentgrass (*Agrostis capillaris*) behaved
19 similarly.

20 A study by Bosley et al. (1998) on the germination of spores of the moss, *Polytrichum*
21 *commune*, and the ferns, *Athyrium felix-femina* and *Onoclea sensibilis*, revealed no effect of O₃
22 on moss spores, while SAR at pH < 4.0 was completely inhibitory. With the ferns, germination
23 was progressively reduced by both increased O₃ and acidity.

24 In summary, the few findings of interactions in these recent studies are consistent with the
25 previous conclusion regarding the likelihood of such interactions being antagonistic. However,
26 the interactions observed were in each case largely the result of the response to the lowest pH
27 used, which in several cases was below 3.0, and hence may not be relevant to most field
28 conditions.

9.4.6.6 Heavy Metals

Since there appears to have been no further research into the interactions of oxidants with heavy metal pollutants, our understanding is unchanged from at the time of the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996). As noted therein, the limited data available from early studies indicates varying degrees of enhancement of any adverse effects of O₃ but precludes the development of any response relationships.

9.4.6.7 Mixtures of Ozone with Two or More Pollutants

In many airsheds the mixtures that occur, both concurrently and over time, may involve three or more pollutants. Very little useful information exists on the effects of O₃ with multiple pollutants. As the 1996 criteria document and others have pointed out, most of the early studies on such combinations can be discounted because of their use of (1) high and environmentally irrelevant exposure concentrations and (2) unrealistic, repetitive exposure profiles (U.S. Environmental Protection Agency, 1996; Barnes and Wellburn, 1998).

The large investment in experimental facilities required to study these complex interactions is a major deterrent. So, although the topic has been included in several reviews that have appeared in the last decade, there appear to have been only two studies that have provided new information on the effects of O₃ in combination with more than one other pollutant stress. Ashenden et al. (1995, 1996) studied the effects of O₃ and/or (SO₂ + NO₂) with four acidities of SAR applied to each gas treatment, on white clover (*Trifolium repens*) and two pasture grasses (*Lolium perenne* and *Agrostis capillaris*). With each species, the antagonism reported for the O₃ × SAR interaction (Section 9.4.4.6.5) tended to be nullified by concurrent exposure to the other gases, while the combination of the three gaseous pollutants resulted in the most severe growth inhibition, regardless of the acidity of the SAR.

With such meager evidence, no clear conclusions can be drawn as to the ways in which the effects of multiple airborne stressors could influence or be influenced by O₃.

9.4.7 Interactions with Agricultural Chemicals

The review of interactions involving O₃, plants, and various agricultural chemicals presented in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) remains a valid assessment of our limited knowledge of these interrelationships. Our knowledge is largely based

on the protection against O₃ afforded to a range of crop species by applications of various chemicals, particularly fungicides, such as benomyl (benlate; methyl-1-[butylcarbamoyl]-2-benzimidazolecarbamate) and several carbamates and triazoles. A recent report has added azoxystrobin (AZO) and epoxyconazole (EPO) to the list (Wu and Tiedemann, 2002). Foliar sprays of either AZO or EPO provided 50 to 60% protection against O₃ injury to barley (*Hordeum vulgare*) leaves. Both had similar modes of action involving stimulation of the levels of antioxidant enzymes such as SOD, ascorbate peroxidase, guaiacol peroxidase, and catalase.

In contrast, applications of herbicides have yielded variable results ranging from increased sensitivity to protection from O₃; the nature of the effect is usually species- or cultivar-dependent. Although of less wide application, some plant growth retardants have also been found to provide protection, but no insecticide appears to have been clearly shown to have similar properties.

Despite the attraction of the use of permitted chemicals to provide crop protection, the statement in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) is still valid: “It is premature to recommend their use specifically for protecting crops from the adverse effects of O₃, rather than for their primary purpose.”

9.4.8 Factors Associated with Global Climate Change

During the last decade, interest in the effects of climatic change on vegetation has replaced concerns over the purported causes of forest decline and the effects of acidic deposition. Two specific components of climate change have been singled out as the foci of most of the research activity:

- the effects of increasing mean global CO₂ concentrations in the lower atmosphere, and
- the effects of increasing levels of surface-level irradiation by UV-B (the result of stratospheric O₃ depletion).

In spite of the crucial role of temperature as a climatic determinant (Monteith and Elston, 1993), the effects of increasing mean global temperatures and their interactions with increasing CO₂ levels in particular have received less attention.

All of the biotic and chemical interactions with oxidants discussed in the preceding sections may be modified by these climatic changes. However, research activities have largely

1 focused on the two-way $O_3 \times CO_2$ interaction. Little if any experimental evidence exists related
2 to three-way interactions such as $O_3 \times CO_2 \times$ disease or $O_3 \times CO_2 \times$ nutrient availability,
3 although such interactions cannot be predicted from the component two-way interactions.

4 Numerous reviews have appeared since the 1996 O_3 AQCD (U.S. Environmental
5 Protection Agency, 1996) dealing with the issues involved. General reviews include
6 publications of IPCC (1996); IPCC (2001); and UNEP (1993, 1999); the volume by Wellburn
7 (1994); the volumes edited by Alscher and Wellburn (1994), De Kok and Stulen (1998), Singh
8 (2000), and Yunus and Iqbal (1996); and papers by Idso and Idso (1994), Krupa and Groth
9 (2000), Luo et al. (1999), Polle and Pell (1999), Poorter and Pérez-Soba (2001), Runeckles
10 (2002), and Weubbles et al. (1999). Effects on agriculture and crop production, growth, and
11 metabolism have been reviewed by Groth and Krupa (2000), Rötter et al. (1999), and Schnug
12 (1998); effects on forests have been reviewed by Bortier et al. (2000); with focus on insect pests,
13 Docherty et al. (1997), Karnosky et al. (2001b,a), McLaughlin and Percy (1999), and Saxe et al.
14 (1998).

15 As background to the discussion of interactions with O_3 , it should be noted that the
16 increased levels of CO_2 experienced since the mid-18th century are such that, without abatement
17 of the rates of increase, increased levels of from 540 to 970 ppm have been projected by the year
18 2100 (IPCC, 2001). Such increases in the concentration of CO_2 , the principal GHG released into
19 the atmosphere, will inevitably lead to increased global mean temperatures, evidence for which
20 is already available from oceanic, icepack, and other records. The latest estimates of the global
21 warming are for an increase in the range 1.4 to 5.8 °C over this century, in contrast to the 0.6 °C
22 rise experienced since 1900 (IPCC, 2001). However, considerable uncertainty is associated with
23 such projections of future increases in global temperature.

24 The use of elevated CO_2 concentrations has been common practice for many years in the
25 production of many greenhouse crops. Much of our early knowledge of the effects of higher
26 than ambient CO_2 levels on plant growth derives from this application, coupled with research of
27 plant physiologists on how CO_2 concentrations affect the process of photosynthesis. Information
28 available about effects of increased CO_2 levels on photosynthesis and stomatal function, in
29 particular, has provided the underlying bases for numerous process models that simulate plant
30 growth under stress and in changed climates.

Although simple $O_3 \times$ temperature interactions were discussed in Section 9.4.4.2, the close linkage between global CO_2 levels and global mean temperatures in the context of climate change requires that an assessment of the interactive effects with O_3 should focus, as much as possible, on interactions involving all three factors.

9.4.8.1 Ozone-Carbon Dioxide-Temperature Interactions

Idso and Idso (1994) reviewed several hundred reports published between 1982 and 1994 on the effects of increased CO_2 on plant growth and net photosynthesis. Their survey covered a wide range of temperate and tropical, herbaceous and perennial species, including coniferous trees. They concluded that, for responses to a 300-ppm increase in CO_2 , somewhat less than a doubling of present-day levels, but somewhat greater than the 540 ppm lower limit suggested by the IPCC (2001), averaged across all species:

- light intensity had a negligible effect on net photosynthesis other than at limiting low intensities under which the CO_2 -driven enhancement was increased;
- increased temperature tended to increase the CO_2 -driven enhancement of dry matter accumulation (growth) and net photosynthesis;
- drought conditions tended to increase the CO_2 -driven enhancements of both growth and net photosynthesis, but increased salinity had little effect;
- mineral nutrient deficiency (especially of nitrogen) tended to increase the CO_2 -driven enhancement of growth; and
- in the presence of air pollutants (especially SO_2 and NO_x), the CO_2 -driven enhancement of net photosynthesis tended to be increased.

It should be noted that the statement that CO_2 -enhanced growth increased with temperature referred to total dry matter accumulation by the whole plant and not to the yield of grain, fruit, or seed. Unfortunately, despite the existence of several reports at the time, the summary of interactions with air pollutants contained only a single reference to O_3 , i.e., Pffirmann and Barnes (1993), who reported surprisingly that a doubling of CO_2 levels led to a 27% increase in dry weight of radish (*Raphanus sativus*) but that the combination with O_3 led to a 77% increase.

The more recent reviews by Rudorff et al. (2000) and Olszyk et al. (2000) have addressed $CO_2 \times O_3$ interactions in detail, with the latter focusing on the implications for ecosystems. They concluded that:

- the effects of both gases on stomatal closure were predominantly additive, with little evidence of interaction;
- increased photosynthesis resulting from elevated CO₂ may be canceled by exposures to high O₃ levels;
- foliar O₃ injury is reduced by elevated CO₂; and
- interactions between CO₂ and O₃ can affect storage carbohydrates, leaf free-radical metabolism, and carbon allocation to shoots and roots.

Olszyk et al. also made specific note of the relative lack of information on below-ground effects.

Much of the recently published information on the effects of increased CO₂ and O₃ levels is summarized in Table 9-12. Note that the table only lists the *directions* of O₃-induced effects and any modifications of these effects resulting from elevated CO₂, not their magnitudes. These directions are usually, but not necessarily, the same as the corollary effects of O₃ on CO₂-induced responses.

The bulk of the available evidence clearly shows that, under the various experimental conditions used (which almost exclusively employed abrupt or “step” increases in CO₂ concentration, as discussed below), increased CO₂ levels may protect plants from the adverse effects of O₃ on growth. This protection may be afforded in part by CO₂ acting together with O₃ in inducing stomatal closure, thereby reducing O₃ uptake, and in part by CO₂ reducing the negative effects of O₃ on Rubisco and its activity in CO₂-fixation. However, the situation with regard to the combined effects of O₃ and CO₂ on stomatal functioning is not clear-cut. In spite of the wealth of evidence supporting both CO₂-induced and O₃-induced short-term stomatal closure, studies over long-term exposures, especially with tree species, have revealed little effect of elevated CO₂ on stomatal conductance. Reverse effects of O₃ on stomata have also been noted, as a result of O₃-induced stomatal dysfunction after extended periods of exposure (Maier-Maercker, 1998).

At the mechanistic level, Rubisco plays a key role in CO₂-assimilation, and while both O₃ and elevated CO₂ per se can lead to reduced activity, CO₂ can also reverse the O₃-induced inhibition of Rubisco activity and photosynthesis (Table 9-12). However, in their review of the possible mechanisms involved, Polle and Pell (1999) cautioned that Rubisco should not be regarded “as a unique target for the interaction of the two gases.” But it is clear from the bulk of

Table 9-12. Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O₃ Effects: V, Decrease; ^, Increase; O, No Significant Effect. CO₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; O, No Significant Effect.					
Plant response	O ₃ Response ^a	CO ₂ Modification ^b	Species	Facility ^c	Reference
Biochemical/metabolic					
Ascorbate peroxidase	V	▼	Wheat (<i>Triticum aestivum</i>)	CSTR, P	Rao et al. (1995)
	V	O□	Sugar maple (<i>Acer saccharum</i>)	CEC, P	Niewiadomska et al. (1999)
	V	O□	Trembling aspen (<i>Populus tremuloides</i>)	FACE, G	Wustman et al. (2001)
Catalase	V	O□	Wheat (<i>T. aestivum</i>)	CEC, P	McKee et al. (1997b)
	^	O□		CEC, P	Niewiadomska et al. (1999)
Chlorophyll	V	▼	Wheat (<i>T. aestivum</i>)	OTC, G	Ommen et al. (1999); Donnelly et al. (2000)
	V	▼	Potato (<i>Solanum tuberosum</i>)	OTC, G	Donnelly et al. (2001a)
Glutathione reductase	V	▼	Wheat (<i>T. aestivum</i>)	CSTR, P	Rao et al. (1995)
	O	O□	Sugar maple (<i>A. saccharum</i>)	CEC, P	Niewiadomska et al. (1999)
	O	O□	Aspen (<i>P. tremuloides</i>)	FACE, G	Wustman et al. (2001)
Rubisco	V	▼	Soybean (<i>Glycine max</i>)	OTC, P	Reid et al. (1998)
	V	▼	Wheat (<i>T. aestivum</i>)	OTC, G	McKee et al. (2000)
	[^]	▼	Trembling aspen (<i>P. tremuloides</i>)	FACE, G	Noormets et al. (2001)
	V	▼	Sugar maple (<i>A. saccharum</i>)	CEC, P	Gaucher et al. (2003)

Table 9-12 (cont'd). Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O ₃ Effects: ▽, Decrease; ▲, Increase; ○, No Significant Effect. CO ₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; ○, No Significant Effect.					
Plant response	O ₃ Response ^a	CO ₂ Modification ^b	Species	Facility ^c	Reference
<i>Biochemical/metabolic (cont'd)</i>					
Rubisco activity	▽	▼	Soybean (<i>G. max</i>)	OTC, P	Reid et al. (1998)
	▽	▼	Wheat (<i>T. aestivum</i>)	CEC, P	McKee et al. (1995)
	▽	[▼]	Wheat (<i>T. aestivum</i>)	OTC, G	McKee et al. (2000)
	▽	▼	European beech (<i>Fagus sylvatica</i>)	CEC, P	Lütz et al. (2000)
Superoxide dismutase	○	○□	Wheat (<i>T. aestivum</i>)	CEC, P	McKee et al. (1997b)
	○	○□	Sugar maple (<i>A. saccharum</i>)	CEC, P	Niewiadomska et al. (1999)
Physiological					
Stomatal conductance	▽	▲	Radish (<i>Raphanus sativus</i>)	CEC, P	Barnes and Pfirrmann (1992)
	▽	▼	Soybean (<i>G. max</i>)	OTC, G	Mulchi et al. (1992)
	▽	▼	Bean (<i>Phaseolus vulgaris</i>)	OTC, P	Heagle et al. (2002)
	▽	[▲]	White clover (<i>Trifolium repens</i>) (O ₃ -sensitive)	CSTR, P	Heagle et al. (1993)
	▲ - ○□	▼ - ○□	White clover (<i>T. repens</i>) (O ₃ -tolerant)	CSTR, P	Heagle et al. (1993)
	▽	▲	Tomato (<i>Lycopersicon esculentum</i>)	CEC, P	Hao et al. (2000)

Table 9-12 (cont'd). Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O ₃ Effects: ▽, Decrease; ▲, Increase; ○, No Significant Effect. CO ₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; ○, No Significant Effect.					
Plant response	O ₃ response ^a	CO ₂ modification ^b	Species	Facility ^c	Reference
Physiological (cont'd)					
Stomatal conductance (cont'd)	▽	▲	Potato (<i>S. tuberosum</i>)	OTC, G	Finnan et al. (2002)
	▽	▲	Wheat (<i>T. aestivum</i>)	CEC, P	Balaguer et al. (1995) Barnes et al. (1995)
	[▽]	▲		CEC, P	McKee et al. (1995)
	▽	○□		OTC, G	Mulholland et al. (1997b)
	▽	○ - ▲		CEC, P	Donnelly et al. (1998)
	▲	○□		CEC, P	Tiedemann and Firsching (2000)
	▽	○□	<i>Agropyron smithii</i>	CEC, P	Volin et al. (1998)
	▽	○□	<i>Koeleria cristata</i>	CEC, P	Volin et al. (1998)
	○□	○□	<i>Bouteloua curtipendula</i>	CEC, P	Volin et al. (1998)
	○□	○□	<i>Schizachyrium scoparium</i>	CEC, P	Volin et al. (1998)
	○□	○□	Black cherry (<i>Prunus serotina</i>)	CSTR, P	Loats and Rebbeck (1999)
	○□	○□	Green ash (<i>Fraxinus pennsylvanica</i>)	CSTR, P	Loats and Rebbeck (1999)
	[▲]	▼	Yellow poplar (<i>Liriodendron tulipifera</i>)	CSTR, P	Loats and Rebbeck (1999)
	▽	▲	Trembling aspen (<i>P. tremuloides</i>)	CEC, P	Volin et al. (1998)

Table 9-12 (cont'd). Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O ₃ Effects: V, Decrease; ^, Increase; O, No Significant Effect. CO ₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; O, No Significant Effect.					
Plant response	O ₃ Response ^a	CO ₂ Codification ^b	Species	Facility ^c	Reference
Physiological (cont'd)					
Stomatal conductance (cont'd)	O - V	▼		FACE, G	Noormets et al. (2001)
	O□	O□	Red oak (<i>Quercus rubra</i>)	CEC, P	Volin et al. (1998)
	[^]	▼	Durmast oak (<i>Quercus petraea</i>)	CEC, P	Broadmeadow et al. (1999)
Photosynthesis	V	▼	Radish (<i>R. sativus</i>)	CEC, P	Barnes and Pfirmann (1992)
	V	▼	Soybean (<i>G. max</i>)	OTC, P	Booker et al. (1997)
	V	▼	Soybean (<i>G. max</i>)	OTC, G	Mulchi et al. (1992)
	V	▼	Bean (<i>P. vulgaris</i>)	OTC, P	Heagle et al. (2002)
	V	O□	Tomato (<i>L. esculentum</i>)	CEC, P	Hao et al. (2000)
	O□	O□	Potato (<i>S. tuberosum</i>)	OTC, G	Donnelly et al. (2001a)
	O□	[▼]		OTC, G	Lawson et al. (2001b)
	[V]	▼	Wheat (<i>T. aestivum</i>)	CEC, P	Barnes et al. (1995)
	V	▼	Wheat (<i>T. aestivum</i>)	OTC, G	Donnelly et al. (2000)
				OTC, P	Mulholland et al. (1997b)
				CEC, P	Reid and Fiscus (1998)
					Tiedemann and Firsching (2000)
	V	O□		CEC, P	Cardoso-Vilhena and Barnes (2001)

Table 9-12 (cont'd). Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O ₃ Effects: V, Decrease; Λ, Increase; O, No Significant Effect. CO ₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; O, No Significant Effect.					
Plant response	O ₃ Response ^a	CO ₂ Codification ^b	Species	Facility ^c	Reference
Physiological (cont'd)					
Photosynthesis (cont.)	O	O	<i>Agropyron smithii</i>	CEC, P	Volin et al. (1998)
	V	▼	<i>Koeleria cristata</i>	CEC, P	Volin et al. (1998)
	O - V	O	<i>Bouteloua curtipendula</i>	CEC, P	Volin et al. (1998)
	[V]	▼	<i>Schizachyrium scoparium</i>	CEC, P	Volin et al. (1998)
	V	▼	Ponderosa pine (<i>Pinus ponderosa</i>)	CEC, G	Olszyk et al. (2001)
	V	▼	Scots pine (<i>Pinus sylvestris</i>)	OTC, G	Kellomäki and Wang (1997b); Kellomäki and Wang (1997a)
	O	O	Black cherry (<i>P. serotina</i>)	CSTR, P	Loats and Rebbeck (1999)
	[V]	▼	Green ash (<i>F. pennsylvanica</i>)	CSTR, P	Loats and Rebbeck (1999)
	O	O	Yellow poplar (<i>L. tulipifera</i>)	CSTR, P	Loats and Rebbeck (1999)
	V	▼	Trembling aspen (<i>P. tremuloides</i>)	CEC, P	Volin et al. (1998)
	V	▼	European beech (<i>F. sylvatica</i>)	CEC, P	Grams et al. (1999)
	O	O	Red oak (<i>Q. rubra</i>)	CEC, P	Volin et al. (1998)
	V	▼	Sugar maple (<i>A. saccharum</i>)	CEC, P	Gaucher et al. (2003)

Table 9-12 (cont'd). Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O₃ Effects: V, Decrease; ^, Increase; O, No Significant Effect. CO₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; O, No Significant Effect.					
Plant response	O ₃ Response ^a	CO ₂ Codification ^b	Species	Facility ^c	Reference
Physiological (cont'd)					
Photorespiration	V	O□	Soybean (<i>G. max</i>)	OTC, P	Booker et al. (1997)
	V	▲	Wheat (<i>T. aestivum</i>)	CEC, P	McKee et al. (1997b)
Growth, Yield					
Total biomass	V	▼	Parsley (<i>Petroselinum sativum</i>)	CEC, P	Cardoso-Vilhena et al. (1998)
	V	▼	Bean (<i>P. vulgaris</i>)	CEC, P	Cardoso-Vilhena et al. (1998)
	V	▼		OTC, P	Heagle et al. (2002)
	V	▼	Soybean (<i>G. max</i>)	OTC, G CSTR, P	Mulchi et al. (1992) Reinert et al. (1997)
	[V]	O□	Alfalfa (<i>Medicago sativa</i>)	CEC, P	Johnson et al. (1996a)
	V	▼	White clover (<i>T. repens</i>) (O ₃ -sensitive)	CSTR, P	Heagle et al. (1993)
	O□	O□	White clover (<i>T. repens</i>) (O ₃ -tolerant)	CSTR, P	Heagle et al. (1993)
	V	▼	Tomato (<i>L. esculentum</i>)	CEC, P CSTR, P	Hao et al. (2000) Reinert and Ho (1995)
	O□	O□	Potato (<i>S. tuberosum</i>)	OTC, G	Donnelly et al. (2001b); Persson et al. (2003)

Table 9-12 (cont'd). Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O ₃ Effects: V, Decrease; ^, Increase; O, No Significant Effect. CO ₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; O, No Significant Effect.					
Plant response	O ₃ Response ^a	CO ₂ Codification ^b	Species	Facility ^c	Reference
Growth, Yield (cont'd)					
Total biomass (cont'd)	V	O		OTC, G	Lawson et al. (2001a)
	V	▼	Mustard (<i>Sinapis alba</i>)	CEC, P	Cardoso-Vilhena et al. (1998)
	V	▼	Plantain (<i>Plantago major</i>)	CEC, P	Cardoso-Vilhena et al. (1998)
	V	▼	Cotton (<i>Gossypium hirsutum</i>)	OTC, P	Booker, 2000) Heagle et al. (1999)
	V	▼	Wheat (<i>T. aestivum</i>)	CEC, P OTC, G OTC, P CEC, P OTC, G CSTR, P OTC, G	Cardoso-Vilhena et al. (1998) Fangmeier et al. (1996) Heagle et al. (2000) McKee et al. (1997a) Pleijel et al. (2000) Rao et al. (1995) Rudorff et al. (1996a)
	[V]	▼	Wheat (<i>T. aestivum</i>)	OTC, G	Bender et al. (1999); Mulholland et al. (1997a)
	V	O□	Wheat (<i>T. aestivum</i>)	CEC, P	Cardoso-Vilhena et al. (1998) Tiedemann and Firsching (2000)
	V	[▼]	Wheat (<i>T. aestivum</i>)	OTC, G	Ewart and Pleijel (1999)
	[V]	▼	Timothy (<i>Phleum pratense</i>)	CEC, P	Johnson et al. (1996a)
	V	▼	<i>Agropyron smithii</i>	CEC, P	Volin et al. (1998)
	V	▼	<i>Koeleria cristata</i>	CEC, P	Volin et al. (1998)

Table 9-12 (cont'd). Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O₃ Effects: V, Decrease; ^, Increase; O, No Significant Effect. CO₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; O, No Significant Effect.					
Plant response	O ₃ Response ^a	CO ₂ Codification ^b	Species	Facility ^c	Reference
Growth, Yield (cont'd)					
Total Biomass (cont'd)	V	▼	Corn (<i>Zea mays</i>)	OTC, G	Rudorff et al. (1996a)
	V	O□	<i>Bouteloua curtipendula</i>	CEC, P	Volin et al. (1998)
	O	O□	<i>Schizachyrium scoparium</i>	CEC, P	Volin et al. (1998)
	V	▼	Ponderosa pine (<i>P. ponderosa</i>)	CEC, G	Olszyk et al. (2001)
	V	▲	Birch (<i>Betula pendula</i>)	CEC, P	Kytöviita et al. (1999)
	[V]	O□	Black cherry <i>P. serotina</i>	CSTR, P	Loats and Rebbeck (1999)
	^	▼	Green ash (<i>F. pennsylvanica</i>)	CSTR, P	Loats and Rebbeck (1999)
	O	O□	European ash (<i>Fraxinus excelsior</i>)	OTC, G	Broadmeadow and Jackson (2000)
	[^]	▲	Yellow poplar (<i>L. tulipifera</i>)	CSTR, P	Loats and Rebbeck (1999)
	V	▼	Sugar maple (<i>A. saccharum</i>)	CEC, P	Gaucher et al. (2003)
	V	▼	Trembling aspen (<i>P. tremuloides</i>) (O ₃ -tolerant clone}	CEC, P OTC, P OTC, G	Volin et al. (1998) Dickson et al. (1998) Dickson et al. (2001)
	V	O□	(O ₃ -sensitive clone)	OTC, G	Dickson et al. (2001)
	O	O□	Red oak (<i>Q. rubra</i>)	CEC, P	Volin et al. (1998)
	V	▼	Durmast oak (<i>Q. petraea</i>)	CEC, P OTC, G	Broadmeadow et al. (1999) Broadmeadow and Jackson (2000)

Table 9-12 (cont'd). Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O ₃ Effects: V, Decrease; ^, Increase; O, No Significant Effect. CO ₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; O, No Significant Effect.					
Plant response	O ₃ Response ^a	CO ₂ Codification ^b	Species	Facility ^c	Reference
Growth, Yield (cont'd)					
Total Biomass (cont'd)	O	O□	Aleppo pine (<i>Pinus halepensis</i>)	CEC, P	Kytöviita et al. (1999)
	O	O□	Scots pine (<i>P. sylvestris</i>)	OTC, G	Broadmeadow and Jackson (2000)
Seed/grain/fruit/tuber yield	V	▼	Soybean (<i>G. max</i>)	OTC, P OTC, G OTC, G	Fiscus et al. (1997); Mulchi et al. (1992); Mulchi et al. (1995)
	V	▼	Bean (<i>P. vulgaris</i>)	OTC, P	Heagle et al. (2002)
	V	▼	Tomato (<i>L. esculentum</i>)	CSTR, P	Reinert and Ho (1995)
	V	▼	Potato (<i>S. tuberosum</i>)	OTC, G	Finnan et al. (2002)
	O	O		OTC, G	Persson et al. (2003)
	V	▼	Wheat (<i>T. aestivum</i>)	OTC, G OTC, G CEC, P OTC, G OTC, G OTC, G	Bender et al. (1999); Fangmeier et al. (1996); McKee et al. (1997a); Mulchi et al. (1995); Mulholland et al. (1998a) Rudorff et al. (1996b)
	[V]	▼	Wheat (<i>T. aestivum</i>) (cont.)	OTC, G	Fangmeier et al. (1996)
	V	[▼]		OTC, G	Mulholland et al. (1998b, 1998a)

Table 9-12 (cont'd). Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O ₃ Effects: ▽, Decrease; ▲, Increase; ○, No Significant Effect. CO ₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; ○, No Significant Effect.					
Plant response	O ₃ Response ^a	CO ₂ Codification ^b	Species	Facility ^c	Reference
Growth, Yield (cont'd)					
Relative growth rate	▽	▼	Wheat (<i>T. aestivum</i>)	CEC, P	Barnes et al. (1995)
	▽	▼		CEC, P	Cardoso-Vilhena and Barnes (2001)
	▽	▼	<i>Agropyron smithii</i>	CEC, P	Volin et al. (1998)
	▽	▼	<i>Koeleria cristata</i>	CEC, P	Volin et al. (1998)
	○	○□	<i>Bouteloua curtipendula</i>	CEC, P	Volin et al. (1998)
	○	○□	<i>Schizachyrium scoparium</i>	CEC, P	Volin et al. (1998)
	○	○□	European ash (<i>F. excelsior</i>)	OTC, G	Broadmeadow and Jackson (2000)
	▽	▼	Trembling aspen (<i>P. tremuloides</i>)	CEC, P	Volin et al. (1998)
	[▽]	▼	Red oak (<i>Q. rubra</i>)	CEC, P	Volin et al. (1998)
	▽	▼	Durmast oak (<i>Q. petraea</i>)	OTC, G	Broadmeadow and Jackson (2000)
	○	○□	Scots pine (<i>P. sylvestris</i>)	OTC, G	Broadmeadow and Jackson (2000)
Specific leaf area-SLA	▽	○□	Radish (<i>R. sativus</i>)	CEC, P	Barnes and Pfirrmann (1992)
	▽	▼	Soybean (<i>G. max</i>)	OTC, P	Reid et al. (1998)
	▽	○□		OTC, G	Mulchi et al. (1992)

Table 9-12 (cont'd). Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O ₃ Effects: ∨, Decrease; ∧, Increase; ○, No Significant Effect. CO ₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; ○, No Significant Effect.					
Plant response	O ₃ Response ^a	CO ₂ Codification ^b	Species	Facility ^c	Reference
Growth, Yield (cont'd)					
Specific leaf area-SLA (cont'd)	∨	▲	White clover (<i>T. repens</i>) (O ₃ -sensitive)	CSTR, P	Heagle et al. (1993)
	∧	▲	White clover (<i>T. repens</i>) (O ₃ -tolerant)	CSTR, P	Heagle et al. (1993)
	∧	▲	<i>Agropyron smithii</i>	CEC, P	Volin et al. (1998)
	∧	▲	<i>Koeleria cristata</i>	CEC, P	Volin et al. (1998)
	○	○	<i>Bouteloua curtipendula</i>	CEC, P	Volin et al. (1998)
	∧	○	<i>Schizachyrium scoparium</i>	CEC, P	Volin et al. (1998)
	∧	▲	Trembling aspen (<i>P. tremuloides</i>)	CEC, P	Volin et al. (1998)
	∧	▲	Red oak (<i>Q. rubra</i>)	CEC, P	Volin et al. (1998)
Root/shoot ratio	∨	▼	Radish (<i>R. sativus</i>)	CEC, P	Barnes and Pfirrmann (1992)
	∨	▲	Alfalfa (<i>M. sativa</i>)	CEC, P	Johnson et al. (1996a)
	∨	▼	White clover (<i>T. repens</i>) (O ₃ -sensitive)	CSTR, P	Heagle et al. (1993)
	○	○□	White clover (<i>T. repens</i>) (O ₃ -tolerant)	CSTR, P	Heagle et al. (1993)
	∧	▲	Wheat (<i>T. aestivum</i>)	CEC, P	McKee et al. (1997a)

Table 9-12 (cont'd). Effects of Increased Carbon Dioxide on Ozone-Induced Responses of Plants at the Metabolic, Physiological, and Whole-Plant Levels

O ₃ Effects: ▽, Decrease; ∧, Increase; ○, No Significant Effect. CO ₂ Effects: ▲, Additive/Synergistic; ▼, Antagonistic/Ameliorative; ○□, No Significant Effect.					
Plant response	O ₃ Response ^a	CO ₂ Codification ^b	Species	Facility ^c	Reference
Growth, Yield (cont'd)					
Root/shoot ratio (cont'd)	▽	▼	Timothy (<i>P. pratense</i>)	CEC, P	Johnson et al. (1996a)
	○	○□	Black cherry (<i>P. serotina</i>)	CSTR, P	Loats and Rebbeck (1999)
	○	○□	Green ash (<i>F. pennsylvanica</i>)	CSTR, P	Loats and Rebbeck (1999)
	○	○□	Yellow poplar (<i>L. tulipifera</i>)	CSTR, P	Loats and Rebbeck (1999)
	○	○□	Aspen (<i>P. tremuloides</i>)	OTC, G	Dickson et al. (2001)
Foliar injury	∧	▼	Potato (<i>S. tuberosum</i>)	OTC, G	Donnelly et al. (2001b); Persson et al. (2003)
	∧	▼	Bean (<i>P. vulgaris</i>)	OTC, P	Heagle et al. (2002)
	∧	▼	Cotton (<i>G. hirsutum</i>)	OTC, P	Heagle et al. (1999)
	∧	▼	Wheat (<i>T. aestivum</i>)	CEC, P OTC, G	Barnes et al. (1995) Mulholland et al. (1997a)
	∧	▼	Trembling aspen (<i>P. tremuloides</i>)	FACE, G	Karnosky et al. (1999) Wustman et al. (2001)
	∧	▼	European beech (<i>F. sylvatica</i>)	CEC, P	Grams et al. (1999)

O₃ m ≤ 0.15 ppm.

^b CO₂-modifications of O₃-effects resulting from ~2× present levels. (Trends are shown in brackets. Pronounced changes with ontogeny are, for example, indicated thus: ○□ ▼.)

^c Exposure facilities used: CEC: controlled environment chambers; CSTR: continuously stirred tank reactors (Heck et al. 1978); FACE: free air CO₂ enrichment facilities; OTC: open-top chambers. G: plants rooted in the ground; P: plants grown in pots. All species are C₃ except corn, *Bouteloua* and *Schizachyrium*.

the evidence in Table 9-12 that elevated CO₂ levels can ameliorate the inhibition of growth caused by O₃ in many species, although the precise balance among the mechanisms involved may well vary from species to species. Three important caveats must be raised with regard to the findings presented in Table 9-12:

- the applicability of results from experiments with an abrupt (step) increase in CO₂ level to understanding the consequences of the gradual increase in CO₂ predicted for the troposphere over the next hundred years;
- the validity of the findings in several long-term studies (particularly with tree species) conducted using potted plants, because of possible added stresses imposed on their root systems relative to trees growing in the field; and
- the relevance to understanding the effects of climate change of studies focused solely on CO₂ enrichment at current ambient conditions of temperature and precipitation patterns that provide no insights into possible interactive effects as these other climatic variables change concurrently with increasing CO₂ (IPCC, 2001).

The first caveat concerns the distinctly different natures of the exposures to O₃ and CO₂ experienced by plants in the field. Changes in the ambient concentrations of these gases have very different dynamics. In the context of climate change, CO₂ levels increase relatively slowly and may change little over several seasons of growth. On the other hand, O₃ presents a fluctuating stress with considerable hour-to-hour and day-to-day variability (Polle and Pell, 1999). Almost all of the evidence presented in Table 9-12 comes from experimentation involving plants grown from the outset in, or subjected to, an abrupt or step increase to a higher (more or less double), steady CO₂ concentration. In contrast, the O₃ exposure concentrations usually varied from day to day. Luo and Reynolds (1999), Hui et al. (2002), and Luo (2001) noted the difficulties in predicting the likely effects of a gradual CO₂ increase from experiments involving a step increase or those using a range of CO₂ concentrations. Indeed, although using the much accelerated time-scale of an 80-day growing season, Hui et al. (2002) clearly showed significant differences between the rates and magnitudes of various physiological and growth responses of plantain (*Plantago lanceolata*) to CO₂ between gradual and step increase treatments. The authors concluded that, even though there were major differences in most of the parameters studied between the gradual and step treatments, “the convergence of the measured parameters at the end of the experiment provides some encouragement for the applicability of

1 step-type experiments in the field; however, the study suggests caution in interpreting early
2 results from short-term studies.”

3 In long-term studies, the matter of photosynthetic acclimation to elevated CO₂ levels has to
4 be considered. Lawlor and Keys (1993) define *acclimation* in terms of long-term (days, weeks),
5 irreversible physiological changes, in contrast to *regulation*, which relates to more rapid
6 (minutes, hours), reversible changes. Each may be positive or negative, but many studies
7 indicate that, while positive acclimation to elevated CO₂ levels initially led to enhanced
8 photosynthesis and growth, negative acclimation ultimately ensued and reduced CO₂
9 assimilation and growth rates. However, the consensus from recent studies and reviews is that
10 such negative acclimation is most likely to occur in situations in which plants are grown under
11 some additional stress, induced, e.g., by limitations to growth posed by lack of resources such as
12 water or nutrients. The meta-analysis by Curtis (1996) revealed that slow or little negative
13 acclimation was noted in studies on unstressed tree species with unhindered opportunities for
14 root growth and development, a view originally suggested by Arp and Drake (1991) and largely
15 supported in the review by Eamus (1996). A nonwoody perennial, the rhizomatous wetland
16 sedge, *Scirpus olneyi*, grown in its natural environment with no edaphic limitations showed no
17 negative acclimation after 4 years; in fact, photosynthetic capacity increased by 31% (Arp and
18 Drake, 1991). No negative acclimation of well-watered, field-grown Ponderosa pine (*Pinus*
19 *ponderosa*) trees was observed by Tissue et al. (1999) after 6 years of growth at 2×-ambient CO₂
20 levels. Gifford and Morison (1993) have summarized the situation thus: “Where the aerial or
21 root environment for a plant is restricted (as with inter-plant competition, for example), positive
22 feedback is limited and adjustments to the changed resource input balance under high CO₂ can
23 include ‘down-regulation’ of leaf photosynthesis rate as an integral part of a positive growth
24 response.”

25 The influence of other environmental stress factors is borne out by several long-term tree
26 studies. After 3 years in 565 ppm CO₂ in the Duke Forest free-air CO₂ enrichment (FACE)
27 facility in North Carolina, maturing loblolly pine (*Pinus taeda*) trees showed only a marginal
28 CO₂-induced carbon gain if grown on a nutritionally moderate site, but zero gain if grown on a
29 nutritionally poor site (Oren et al., 2001). This is in sharp contrast to the substantially increased
30 initial growth rates in elevated CO₂ reported by DeLucia et al. (1999), but it is supported by the
31 observations of Tognetti et al. (2000) on five Mediterranean tree species growing for many years

1 adjacent to geothermal springs releasing CO₂ to provide ambient levels averaging 700 ppm.
2 No significant differences in radial growth of the oaks (*Quercus cerris*, *Q. ilex*, and
3 *Q. pubescens*), strawberry tree (*Arbutus Unedo*), and flowering ash (*Fraxinus Ornus*) could be
4 detected between trees at the naturally enriched site and those at a nearby site exposed to normal
5 ambient CO₂ (~350 ppm). The authors concluded that limited availability of water and nutrients
6 may have counteracted any positive effects of CO₂ on growth at the enriched site or that the trees
7 had acclimated to the higher CO₂ levels.

8 Because the ameliorative effects of CO₂ on responses to O₃ (Table 9-14) were reported
9 mostly in short-term studies involving an abrupt increase in CO₂ level, it is appropriate to ask
10 whether this amelioration is likely to persist to a time when the ambient CO₂ concentration is
11 relatively stable at such levels. Regardless of any negative acclimation due to resource
12 limitations that may occur in the interim, steadily rising CO₂ levels may well lead to natural
13 selection and genetic change. Nevertheless, it seems reasonable to expect that the amelioration
14 of O₃ impact at elevated CO₂ levels will be maintained in many situations, but the negative
15 acclimation that will probably occur in situations where other resources become limiting will
16 reduce the degree of protection.

17 Another caveat regarding the validity of some of the observations in Table 9-12 is related
18 to the matter of stress-induced negative acclimation to elevated CO₂ and concerns results
19 obtained using potted plants. Although much of the recent information on CO₂ effects has come
20 from experiments with plants rooted in the ground, more than half of the studies listed in
21 Table 9-12 used potted plants, whether in controlled environment and greenhouse chambers or in
22 OTCs. The recent meta-analysis of data on the effects of elevated CO₂ on soybean (*Glycine*
23 *max*) physiology and growth by Ainsworth et al. (2002) revealed a threefold smaller stimulation
24 of seed yield in pot-grown than in field-grown plants, even when large (> 9 L) pots were used.
25 Loats and Rebbeck (1999) noted that their inability to observe any CO₂-induced increases in the
26 root/shoot ratios in seedlings of three broad-leaved tree species (Table 9-13) may have resulted
27 from their use of potted plants. The use of potted plants was a confounding factor in the studies
28 of Taylor et al. (2001) of the differences in leaf growth of poplar (*Populus*) hybrids between
29 plants exposed to elevated CO₂ in controlled environment chambers (potted plants), OTCs or a
30 FACE facility. Eamus (1996) has suggested that any long-term observations of reduced stomatal
31 conductance are almost universally a consequence of pot-based trials at elevated CO₂ with

1 restricted root growth, in contrast to the lack of such a decline observed with trees rooted in the
2 ground. Although the majority of the cases cited in Table 9-12 indicate that O₃ and CO₂ act
3 additively or synergistically in causing stomatal closure, there are numerous exceptions.

4 Any reduction in stomatal aperture has consequences other than merely restricting O₃
5 uptake and the exchange of other gases. In particular, the rate of transpiration is reduced and,
6 while this tends to increase water-use efficiency, it may also lead to decreased mineral uptake,
7 which could adversely impact growth over extended periods. Furthermore, less transpiration
8 also means less evaporative cooling and an increase in leaf temperature, independent of any
9 change in mean air temperatures.

10 Hence, the final caveat regarding Table 9-12 concerns the interactions of O₃ and CO₂ with
11 other climatic variables, especially mean temperature. In light of the key role played by
12 temperature in regulating physiological processes and modifying plant response to increased
13 CO₂ levels (Long, 1991; Morison and Lawlor, 1999) and the knowledge that relatively modest
14 increases in temperature may lead to dramatic consequences in terms of plant development
15 (Lawlor, 1998), it is unfortunate that much of the large investment in time and resources spent
16 on recent studies of the effects of climate change on vegetation have gone into investigations
17 limited to increasing our knowledge of the effects of higher levels of CO₂ *at current ambient*
18 *temperatures*.

19 Some attention is now being paid to investigating the concurrent effects of CO₂ increases
20 and warming (recently reviewed by Rowland-Bamford, 2000 and Morison and Lawlor, 1999),
21 but the observed interactive effects on plant growth are inconsistent. For example, a FACE
22 study with ryegrass (*Lolium perenne*) showed that increased temperatures (provided by infrared
23 heaters) reduced the dry matter gain resulting from increased CO₂ levels (Nijs et al., 1996). The
24 field studies by Shaw et al. (2002) on a California annual grassland dominated by the grasses
25 *Avena barbata* and *Bromus hordeaceus* and the forbs *Geranium dissectum* and *Erodium botrys*
26 involved free-air increased CO₂ as well as increased temperature, precipitation, and N supply.
27 Not only did increased temperature reduce CO₂-stimulated net primary productivity (NPP), but
28 increased CO₂ itself, combined with other factors, was found to be able to cause reduced NPP.

29 There have been several investigations of effects on wheat (*Triticum aestivum*). Batts et al.
30 (1997) used plastic tunnels to create temperature gradients and maintain elevated CO₂ levels
31 over field-grown wheat and found that, in each of 4 years of study, a temperature rise of ~1.5 °C

consistently canceled the growth and yield increases caused by a doubling of the CO₂ level above ambient. Similar findings were reported by van Oijen et al. (1999) and van Oijen and Ewart (1999) in OTC field studies. Half of the chambers were cooled 1.6 to 2.4 °C below the uncooled chambers, to cancel out the normal temperature increase over ambient, due to the so-called “chamber effect” (usually a 1 to 3 °C increase above ambient temperature (Heagle et al., 1988). Although temperature had no effect on CO₂-enhanced assimilation rates, the CO₂-enhanced growth and grain yields observed in the cooled chambers were effectively canceled out in the warmer chambers. The authors attributed this effect to accelerated phenology, a shorter period for grain filling, and a lower leaf area index (LAI; total leaf area per unit ground area) in the warmer chambers. Wheeler et al. (1996) observed that the benefit to wheat of doubling the CO₂ level was offset by a mean seasonal increase of only 1 to 1.8 °C. With the continuing use of OTCs for field research, Runeckles (2002), has suggested that the temperature rise due to the chamber effect in OTCs should be exploited (and measured) as a means of exploring temperature × CO₂ as well as temperature × CO₂ × O₃ interactions.

An indirect affirmation of the importance of temperature as a component of climate change on wheat yield was provided by van Oijen and Ewart (1999), using two simulation models, AFRCWHEAT2-O₃ and LINTULCC (Ewart et al., 1999). They analyzed data from the ESPACE-wheat program, which involved 25 OTCs experiments in 1994, 1995, and 1996 at nine European locations (Jäger et al., 1999). Both models were able to predict control-treatment grain yields closely (5.5 ± 1.2 and 5.8 ± 1.2 t.ha⁻¹, respectively, versus the observed 5.9 ± 1.9 t.ha⁻¹), and both indicated a predominantly negative effect of temperature on the yield response to increased CO₂ (a 3 °C rise reduced the gain in yield from 30 to 14%). However, neither model had an R² > 0.3, indicating that the models included other sources of variability among the sites than the climatic factors. The multiple linear regression developed by Bender et al. (1999) based on the same data sets also included temperature as a highly significant covariant. Both studies are discussed more fully below.

Other studies, however, have found positive temperature-related growth effects, as suggested by the early Idso and Idso (1994) analysis. In an OTC study using the perennial grass *Festuca pratensis* in which a temperature increase of 3 °C above ambient was combined with CO₂-enrichment to 700 ppm, both CO₂ and temperature caused increases in total above-ground biomass (Hakala and Mela, 1996). Studies with potato (Cao et al., 1994) and soybean (Ziska and

Bunce, 1997) using potted plants in controlled environment chambers also showed temperature-enhanced increases in growth in enriched CO₂ atmospheres. Read and Morgan (1996) compared the effects of enriched CO₂ and temperature on two grasses: cool-season *Pascopyrum smithii* and warm-season *Bouteloua gracilis*. In the latter (a C₄ species), 750 ppm CO₂ resulted in increased dry matter production at daytime temperatures as high as 35 °C, but in *P. smithii* (a C₃ species), CO₂-stimulated growth was greatest at 20 °C. However, the stimulation was progressively attenuated by increased temperature, so that at 35 °C, growth in 750 ppm was only one third of that in 350 ppm CO₂ at 20 °C.

Although the picture we have of temperature × CO₂ interactions is inconsistent, Rowland-Bamford (2000) has provided persuasive evidence that the nature of response to temperature in the grain yield of crops with as different temperature optima as rice and wheat will depend upon whether the change is above or below the temperature optimum.

But what if we add O₃ as another variable? Unfortunately, there have been very few studies of the three-way interaction. With the information available on CO₂ × O₃ interactions (Table 9-12) and the limited information on temperature × O₃ interactions (discussed in Section 9.4.4.2) simulation modeling can attempt to provide estimates of O₃ × CO₂ × temperature effects, but experimental observation is still required to validate the models. The questions that need to be answered are: if increased temperature can offset the gains in productivity afforded by increased CO₂ in important species such as wheat, and increased CO₂ can offset the reductions in productivity caused by O₃, will increased temperature modify this protective effect. And if so, in what manner.

To date, the only information available appears to consist of the reports by van Oijen and Ewart (1999) and Bender et al. (1999) referred to above. In the former's simulation studies, the overall yield depression of wheat caused by O₃ was found to be 7 ± 4% for both AFRCWHEAT2-O₃ and LINTULCC models versus an observed 9 ± 11%. The enhancements due to CO₂ were predicted to be 24 ± 9% and 42 ± 11%, respectively, which straddled the observed 30 ± 22% gain. Based on the 13 experiments that included all four treatments (±O₃, ±CO₂), an actual 10% yield loss due to O₃ at ambient CO₂ levels was reduced to a 4% loss by the elevated CO₂. The AFRCWHEAT2-O₃ model predicted 7 and 4% losses, and LINTULCC model predicted 8 and 5% losses due to the O₃ and O₃ + CO₂ treatments. The actual and simulated yield increases in response to CO₂ increased further with increasing temperature, but

1 although temperature had no discernible effect on the observed depression of yield caused by O₃
2 alone, both models suggested that the yield reduction was diminished both by higher
3 temperatures and higher CO₂ levels.

4 The analysis of the ESPACE-wheat experiments by Bender et al. (1999) led to the
5 following multiple linear regression:

6
7
8
$$Y = 1004.6^{***} + 0.588^{***}[\text{CO}_2] - 1.908^{**}[\text{O}_3] - 31.230^{***}[T] + 7.309[I] - 1448.423^{***}[\text{H}_2\text{O}], \quad (9-5)$$

9
10

11 where Y = grain yield, g-m⁻²; [CO₂] = ppm CO₂; [O₃] = ppb O₃, 12-h mean; $[T]$ = °C; $[I]$ = light
12 intensity, MJ-m⁻²/day; and [H₂O] is a dummy variable: well watered = 1; limited water
13 supply = 2. (***, $p < 0.001$; **, $p < 0.01$; the coefficient for I was not significant.) With
14 $R^2 = 0.3983$, adjusted for 258 degrees of freedom, a large part of the variability was still
15 unaccounted for by the five variables. However, this analysis suggests that CO₂, O₃,
16 temperature, and water-status are important codeterminants of wheat yield but assumes no
17 interactions. Substitution in the model at summer light intensities and with well-watered plants
18 indicates that, at 20 °C, a doubling of CO₂ levels to 700 ppm alone, would lead to a 29.5%
19 increase in yield, while 50 ppb O₃ alone would decrease yield by 10.9%. With both gases at
20 those levels, the yield would only increase 20%, but with a concurrent temperature rise of 2 °C,
21 it would shrink to a 9.6% increase.

22 Both studies, therefore, indicate an amelioration of the effects of O₃ by CO₂, the magnitude
23 of which would be reduced at warmer temperatures. However, they relate to a single crop whose
24 response to CO₂ is temperature-sensitive. Information about other species in which the effects of
25 CO₂ and temperature are additive are limited. However, Wolf and Van Oijen (2003) recently
26 described a model (LPOTCO) simulating the effects of changes in climatic variables, CO₂ and
27 O₃ on tuber yield potential of irrigated potato (*Solanum tuberosum* cv. Bintje) over locations
28 within the European Union ranging from Finland to Italy. They noted that although increased
29 CO₂, O₃, and light intensity were predominant controlling factors, increased temperature also
30 influenced potential yields substantially, with increases in northern latitudes (attributed to a

1 longer growing season) but decreases in southern latitudes (attributed to decreased assimilate
2 production).

3 A clear understanding of the complex interactions of increased CO₂ and temperature with
4 O₃ must await further experimentation or simulations. However, it seems likely that any CO₂-
5 induced amelioration of the adverse effects of O₃ on aspects of growth other than seed or grain
6 yield may be lessened or increased by increased temperature, depending upon the temperature,
7 optima for the species, along the lines suggested by Rowland-Bamford (2000).

8 Other crop simulation models which incorporate O₃ and some of the various environmental
9 factors, including elements of climate change, have been reviewed by Kickert et al. (1999) and
10 Rötter and van de Geijn (1999). However, to date, the applications tend to have focused on
11 interactions of O₃ with factors such as soil moisture or nutrient availability.

12 With forest trees, the situation has the added complexity of a perennial growth form and
13 the inevitability, over time, of subjection to additional environmental stresses such as nutrient-
14 limitation. Here, too, although numerous models of tree growth have been described, there
15 appear to have been few applications to interactions of O₃ and factors of climate change.
16 Constable et al. (1996) used TREEGRO to model the growth of Ponderosa pine (*Pinus*
17 *ponderosa*) exposed to three O₃ levels (0.5×, 1.0×, and 2× ambient), two levels of CO₂ (ambient
18 and ambient + 200 ppm CO₂), and two temperature regimes (ambient and ambient + 4 °C). Plant
19 growth was predicted to be decreased 1, 19, and 39% by the three levels of O₃, respectively.
20 Increased CO₂ reduced the loss at the highest O₃ level to 7% but the combination of elevated
21 CO₂ with the higher temperature more than overcame the adverse effects of O₃, leading to a 4%
22 increase, largely attributed to increased fine root mass. The authors suggested that, in relation to
23 the baseline conditions used in the simulations (Corvallis, OR), higher concentrations of CO₂ and
24 O₃ and a warmer climate will have little impact on total-tree growth, but they noted the
25 importance of undertaking multiple stress studies in order to be able to make accurate forecasts
26 of the impact of such changes on forest trees.

27 More recently, Constable and Friend (2000) compared the capabilities of six published
28 process-based models (CARBON, ECOPHYS, PGSM, TRE-BGC, TREEGRO, and W91) for
29 simulating tree response to elevated CO₂, O₃, and temperature. They concluded that although
30 these models were capable integrators of the effects of various environmental factors on

individual processes such as photosynthesis, they were less reliable when extrapolating to growth.

Although most of the research emphasis has been on simple $\text{CO}_2 \times \text{O}_3$ interactions, a few isolated studies of interactions have involved O_3 , CO_2 , and biotic environmental factors. Heagle et al. (1994) observed that both O_3 and CO_2 tended to be additive in encouraging the growth of spider mite (*Tetranychus urticae*) populations on clover (*Trifolium repens*). Infection of wheat (*Triticum aestivum*) with leaf rust (*Puccinia recondita*) sensitized the plants to O_3 injury, but its severity was significantly reduced in elevated CO_2 (Tiedemann and Firsching, 2000). The effects of O_3 and CO_2 on mycorrhizal symbioses was studied by Kytöviita et al. (1999) who found that CO_2 did not ameliorate the adverse effects of O_3 on the root growth of Aleppo pine (*Pinus halepensis*) and birch (*Betula pendula*). In another study with Aleppo pine, Kytöviita et al. (2001) noted that both O_3 and elevated CO_2 reduced mycorrhiza-induced N-uptake by the roots. In Scots pine (*Pinus sylvestris*), Kasurinen et al. (1999) observed transient effects of elevated CO_2 and O_3 on root symbiosis, but none of the effects persisted over the 3 years of the study.

The soil water $\times \text{O}_3 \times \text{CO}_2$ interaction was experimentally investigated by Broadmeadow and Jackson (2000) in Durmast oak (*Quercus petraea*), European ash (*Fraxinus excelsior*), and Scots pine (*Pinus sylvestris*). No interactions were noted with ash and pine; but with oak, elevated CO_2 ameliorated and irrigation exacerbated the effects of O_3 , although the resultant effects were essentially additive.

Booker (2000) noted that soil nitrogen levels interacted only slightly with O_3 and CO_2 in determining the composition of cotton (*Gossypium hirsutum*) leaves and roots. Carbon dioxide reversed the inhibition of leaf growth caused by O_3 , but increased N-fertility tended to reduce this reversal.

Because of the small number of studies of possibly significant interactions of three or more environmental factors, it impossible to draw any sweeping conclusions as to how O_3 , in the context of global climate change, may affect relationships among plants and insects, diseases, and symbionts or among plants and nutrients or other air pollutants. The only interaction that has some degree of general support is the amelioration of adverse O_3 effects by elevated CO_2 .

9.4.8.2 Ozone-UV-B Interactions

As noted in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996), depletion of stratospheric O₃ by halofluorocarbons has resulted in increased intensities of ultraviolet-B (UV-B) radiation (280 to 320 nm wavelengths) at the earth's surface. The situation is discussed more fully in Chapter 10.

While stratospheric O₃ depletion may result in increased surface UV-B irradiation, absorption of UV-B is a property of the O₃ molecule regardless of its location; and surface UV-B flux is, therefore, also reduced by O₃ in the troposphere. Although only about 10% of the total atmospheric O₃ column occurs in the troposphere (Fishman et al., 1990), it contributes a disproportionately greater absorption effect than stratospheric O₃ because the UV radiation penetrating the troposphere becomes increasingly diffuse as it reaches the surface, with a consequent increase in mean path length (Brühl and Crutzen, 1989). Any benefits to vegetation from reduced ambient O₃ stress must, therefore, also be viewed in the context of possible adverse effects due to increased UV-B irradiation. There are, thus, two distinct types of possible interactions between surface level O₃ and UV-B radiation:

- direct interactions involving simultaneous, sequential, or mixed exposures to O₃ and UV-B stresses; and
- effects on responses to UV-B itself resulting from changes in radiation intensity caused by changes in surface level O₃ concentrations.

Only the first type of interaction is discussed here. The second type of interaction has broad implications for both health and welfare and focuses on the impacts of UV-B radiation per se. It is dealt with separately in Chapter 10, which includes a critical review of the experimental difficulties faced in undertaking meaningful plant research in order to reach a clear understanding of the effects of increased UV-B radiation, the evidence for both its adverse and beneficial effects on plants, and the potential for changes in ambient surface O₃ levels to modify these effects.

The most recent reviews specifically addressing the *combined* effects of tropospheric O₃ and UV-B on plants are by Runeckles and Krupa (1994) and Krupa and Jäger (1996), although the topic has also been included in several more general reviews of O₃ effects and factors of climate change such as those by Unsworth and Hogsett (1996), Krupa et al. (1998), Posthumus (1998), Groth and Krupa (2000), and Krupa and Groth (2000).

1 However, little new information has become available since Runeckles and Krupa (1994)
2 noted that the scanty knowledge of the effects of UV-B and O₃ combinations available at that
3 time was derived solely from studies of soybean (*Glycine max*). Miller et al. (1994) observed no
4 interaction and no effect of UV-B on yield, in contrast to a previous report by Teramura et al.
5 (1990) using the same cultivar, Essex. More recently, in a study of the saltmarsh grass *Elymus*
6 *athericus* subjected to reciprocal exposures to O₃ and UV-B, van de Staaij et al. (1997) observed
7 no interactive effects and no adverse effects of UV-B following 14-day exposures, even though
8 an earlier report showed that longer exposures to UV-B (65 days) could cause a 35% loss of
9 biomass (Van De Staaij et al., 1993). However, in a study in which ambient, high altitude UV-B
10 levels were compared with near zero levels, at ambient or 2×-ambient levels of O₃, interactions
11 involving the levels of antioxidants in Norway spruce (*Picea abies*) and Scots pine (*Pinus*
12 *sylvestris*) were reported by Baumbusch et al. (1998). Schnitzler et al. (1999) subsequently
13 reported that O₃-induced injury and adverse effects on photosynthesis were more pronounced
14 with near zero UV-B levels, indicating an amelioration of the O₃-response. A later study with
15 Scots pine (Zinser et al., 2000) revealed O₃ × UV-B interactions at the gene expression and
16 biochemical levels. In contrast, Ormrod et al. (1995) reported that UV-B predisposed
17 *Arabidopsis thaliana* to injurious growth effects of O₃.

18 At various organizational levels, Runeckles and Krupa (1994) identified several similarities
19 between plant response to O₃ and UV-B, and at the level of gene expression there have recently
20 been several reports of both similarities and distinctions. Willekens et al. (1994) reported similar
21 effects of O₃, UV-B, and SO₂ on the expression of antioxidant genes in *Nicotiana*
22 *plumbaginifolia*. In parsley (*Petroselinum crispum*), Eckey-Kaltenbach et al. (1994a) found that
23 O₃ was a cross-inducer for both the UV-B-induced enhanced biosynthesis of flavonoids and the
24 pathogen-induced furanocoumarin phytoalexins, in keeping with the previously observed
25 O₃-induction of fungal and viral defense reactions. In this regard, Yalpani et al. (1994) provided
26 evidence that, in tobacco (*Nicotiana tabacum*), O₃ and UV-B acted similarly in increasing
27 disease-resistance via a salicylate-mediated enhancement of defense proteins. However,
28 subsequent work with tobacco led Thalmair et al. (1996) to conclude that exposure to UV-B did
29 not lead to the accumulation of pathogenesis-related proteins. In Scots pine (*Pinus sylvestris*),
30 although O₃ is known to induce stilbene synthase and cinnamyl alcohol dehydrogenase, UV-B

was found to enhance the former but suppress the latter, revealing an interaction at the level of gene expression (Zinser et al., 2000).

In summary, the present base of information about possible interactions between increased UV-B radiation and O₃ is insufficient to draw any firm conclusions in terms of gross effects, but there is some evidence of similarities in the effects of O₃ and UV-B individually and of the mechanisms involved at the molecular level.

9.4.8.3 Interactions of Ozone with Multiple Climate Change Factors

Despite of the need for experimental investigations of three-way or more complex interactions among O₃, CO₂, UV-B, temperature, and other climate change factors, few studies have been reported, even without O₃ as a factor. In an isolated report, using tomato (*Lycopersicon esculentum*) seedlings, Hao et al. (2000) employed preexposure to UV-B (\pm CO₂ enrichment) followed by exposure to O₃ (\pm CO₂ enrichment). They observed that CO₂ more than overcame the inhibition of photosynthesis caused by O₃, but pretreatment with UV-B reduced the resultant increase.

In view of the unexpected observations made in their grassland study of the combined effects of CO₂, temperature, precipitation, and N-supply, Shaw et al. (2002) affirmed that: “Ecosystem responses to realistic combinations of global changes are not necessarily simple combinations of the individual factors.” The addition of O₃ to the list of variables results in further complexity.

Although computer simulation modeling may ultimately lead to improved understanding of these complex issues, to date, no such models appear to have been applied to these interactions, possibly because of the scarcity of experimental data for parameterization.

9.4.9 Summary - Environmental Factors

Although O₃ and other photochemical oxidants are phytotoxic, their actions on vegetation may be modified by a host of biotic and abiotic factors in the environment; conversely, they may modify plant response to these other factors. The extensive review of these biological, physical, and chemical factors conducted for the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) concluded with a statement that our understanding was too fragmented to permit drawing many general conclusions. With today’s increased awareness of the need for more complete

1 information on interactions, it is unfortunate that, in the interval since the 1996 criteria
2 document, rigorous, systematic investigations of interactions have been rare, and most of the
3 new information is as fragmented as before. This is inevitable partly in view of the vast scope of
4 the possible interactions between O₃ and other environmental variables. But it is also an
5 unavailable outcome of having the personal interests (and funding) of individual groups
6 determine the studies to be undertaken, instead of a coordinated program of research focused on
7 systematic investigations to improve our understanding our ability to assess the risks posed by
8 photochemical oxidants to cultivated and natural vegetation.

9 In the area of biotic interactions, new evidence with regard to insect pests and diseases has
10 done little to remove the uncertainties noted in the 1996 criteria document. Most of the large
11 number of such interactions that may affect crops, forest trees, and other natural vegetation have
12 yet to be studied. The trend suggested previously that O₃ increases the likelihood and success of
13 insect attack has received some support from recent studies, but only with respect to chewing
14 insects. With the economically important group of sucking insects such as the aphids, no clear
15 trends have been revealed by the latest studies. Hence, although it seems likely that some insect
16 problems could increase as a result of increased O₃ levels, we are still far from being able to
17 predict the nature of any particular O₃ plant insect interaction, its likelihood, or its severity.

18 The situation is a little clearer with respect to interactions involving facultative,
19 necrotrophic plant pathogens, with O₃ generally leading to increased disease. With obligate,
20 biotrophic fungal, bacterial, and nematode diseases, there are twice as many reports indicating
21 O₃-induced inhibitions than enhancements. The frequent reports that infection by obligate
22 biotrophs reduces the severity of O₃-induced foliar injury should not be interpreted as
23 “protection” because of the negative effects on the host plant of the disease per se. With obligate
24 biotrophs, the nature of any interaction with O₃ is probably dictated by the unique, highly
25 specific biochemical relationships between pathogen and host plant. At this time, therefore,
26 although some diseases may become more widespread or severe as a result of exposure to O₃, it
27 is still not possible to predict which diseases are likely to present the greatest risks to crops and
28 forests.

29 Several studies have indicated that the functioning of tree root symbioses with mycorrhizae
30 may be adversely affected by O₃, but there is also evidence that the presence of mycorrhizae may
31 overcome root diseases stimulated by O₃ and that O₃ may encourage the spread of mycorrhizae

1 to the roots of uninfected trees. The latest studies, therefore, present no clearer picture of the
2 likely nature of simple interactions of O₃ and root symbionts, but in view of the importance of
3 mycorrhizae as below-ground components of ecosystems, they are discussed more fully in
4 Section 9.5.

5 The few recent studies of the impact of O₃ on intraspecific plant competition have again
6 confirmed that grasses frequently show greater resilience than other types of plants. In grass-
7 legume pastures, the leguminous species suffer greater growth inhibition. And the suppression
8 of Ponderosa pine seedling growth by blue wild-rye grass was markedly increased by O₃.
9 However, we are far from being able to predict the outcome of the impact of O₃ on specific
10 competitive situations, such as successional plant communities or crop-weed interactions.

11 Light, a component of the plant's physical environment, is an essential "resource" whose
12 energy content drives photosynthesis and CO₂ assimilation. It has been suggested that increased
13 light intensity may increase the sensitivity to O₃ of light-tolerant species while decreasing that of
14 shade-tolerant species, but this appears to be an over-simplification with many exceptions.
15 Temperature affects the rates of all physiological processes based on enzyme-catalysis and
16 diffusion, and each process and overall growth (the integral of all processes) has a distinct
17 optimal temperature range. Although some recent field studies have indicated that O₃ impact
18 significantly increases with increased ambient temperature, other studies have revealed little
19 effect of temperature. But temperature is unquestionably an important variable affecting plant
20 response to O₃ in the presence of the elevated CO₂ levels contributing to global climate change
21 (see below). In contrast, evidence continues to accumulate to indicate that exposure to O₃
22 sensitizes plants to low temperature stress by reducing below-ground carbohydrate reserves,
23 possibly leading to responses in perennial species ranging from rapid demise to impaired growth
24 in subsequent seasons.

25 Although the RH of the ambient air has generally been found to increase the adverse
26 effects of O₃ by increasing stomatal conductance and thereby increasing O₃ flux, abundant
27 evidence indicates that the ready availability of soil moisture results in greater sensitivity to O₃.
28 The partial "protection" against the adverse effects of O₃ afforded by drought has been observed
29 in field experiments and modeled in computer simulations. There is also compelling evidence
30 that O₃ can predispose plants to drought stress. Hence, the response will depend to some extent
31 upon the sequence in which the stresses occur, but, even though the nature of the responses is

1 largely species-specific, successful applications of model simulations led to larger-scale
2 predictions of the consequences of $O_3 \times$ drought interactions. However, it must be recognized
3 that regardless of the interaction, the net result on growth in the short-term is negative, although
4 in the case of tree species, other responses such as increased water use efficiency could be a
5 benefit to long-term survival.

6 Mineral nutrients in the soil, other gaseous air pollutants, and agricultural chemicals
7 constitute chemical factors in the environment. The evidence regarding interactions with
8 specific nutrients is still contradictory. Some experimental evidence indicates that low general
9 fertility increases sensitivity to O_3 , while simulation modeling of trees suggests that nutrient
10 deficiency and O_3 act less than additively, but there are too many example of contrary trends to
11 permit any sweeping conclusions. Somewhat analogously with temperature, it appears that any
12 shift away from the nutritional optimum may lead to greater sensitivity, but the shift would have
13 to be substantial before a significant effect on response to O_3 was observed.

14 Interactions of O_3 with other air pollutants have received relatively little recent attention.
15 The situation with SO_2 remains inconsistent, but seems unlikely to pose any additional risk to
16 those related to the individual pollutants. With the NO and NO_2 , the situation is complicated by
17 their nutritional value as a N source. In leguminous species, it appears that NO_2 may reduce the
18 impact of O_3 on growth, with the reverse in other species, but the nature of the exposure pattern,
19 i.e., sequential or concurrent, also determines the outcome. Much more investigation is needed
20 before we will be able to predict the outcomes of different O_3 - NO - NO_2 scenarios. The latest
21 research into $O_3 \times$ acid rain interactions has confirmed that, at realistic acidities, significant
22 interactions are unlikely. A continuing lack of information precludes offering any
23 generalizations about interactive effects of O_3 with NH_3 , HF , or heavy metals. More evidence
24 has been reported that the application of fungicides affords some protective effects against O_3 .

25 Over the last decade, considerable emphasis has been placed on research into O_3
26 interactions with the components of global climate change: increased atmospheric CO_2 ,
27 increased mean global temperatures, and increased surface level UV-B radiation. However,
28 most of these studies have tended to regard increased CO_2 levels and increased mean
29 temperatures as unrelated phenomena. Experiments into the effects of doubled CO_2 levels at
30 today's mean ambient temperatures are not particularly helpful in trying to assess the impact of
31 *climate change* on responses to O_3 . To date, the limited experimental evidence and evidence

1 obtained by computer simulation suggest that in a 600+ ppm world, although the enriched CO₂
2 would more than offset the impact of O₃ on responses as varied as wheat yield or the growth of
3 young Ponderosa pine trees, the concurrent increase in temperature would reduce but probably
4 not eliminate the net gain. A similar decrease in the net gain resulting from the complete
5 reversal by CO₂ of the inhibition of photosynthesis caused by O₃ has been reported for increased
6 UV-B irradiation. However, these are preliminary results based on minimal data.

7 In conclusion, although the increased use of computer simulations may be important in
8 suggesting outcomes of the many complex interactions of O₃ and various combinations of
9 environmental factors, the results obtained will only be as reliable as the input data used for their
10 parameterization. The data needed for good simulations can only come from organized,
11 systematic study. For predicting the future, ignorance is as good as dependence on poor
12 simulations.

15 **9.5 EFFECTS-BASED AIR QUALITY EXPOSURE- AND** 16 **DOSE-RESPONSE INDICES**

17 **9.5.1 Introduction**

18 An index is needed that relates measured plant damage (i.e., growth) to ambient ozone
19 concentration over time. The index can provide both a consistency for reviewing exposure-
20 response effects in research, as well as a metric for developing a biologically-relevant air quality
21 standard that protects ecological resources. The quantifying function over some time frame has
22 frequently been referred to as “dose-response” and “exposure-response”. The distinction being
23 where the pollutant concentration is measured: “Dose” is the measure of the pollutant
24 concentration absorbed by the leaf over some time period, whereas “exposure” is the ambient air
25 concentration measured nearby the plant over some time period.

26 Plant ozone uptake from the ambient air (either rate or uptake or cumulative seasonal
27 uptake) is the ideal measure, because without ozone or its reactive product(s) reaching the target
28 tissue there is no effect. Uptake is controlled in part by stomata (see Section 9.3 for detailed
29 discussion). An uptake measure should integrate all those environmental factors that influence
30 stomatal conductance, e.g., temperature, humidity, soil water status. However, a direct measure
31 of the internal leaf concentration of ozone is technically difficult and thus uptake values are

generally obtained with models that require species- and site-specific variables. Because of this, a surrogate for uptake (i.e., exposure index) was sought early on using statistical summaries of ambient pollutant concentration over some integral of time (O’Gara, 1922; Lefohn and Benedict, 1982; Lee et al., 1988; U.S. Environmental Protection Agency, 1986, 1992, 1996).

An index of exposure that is biologically relevant must then consider those factors known to modify the plant response by altering ozone uptake (Hogsett et al., 1988; U.S. Environmental Protection Agency, 1996), including the temporal dynamics of exposure (e.g., concentration, frequency, duration), plant phenology (see Section 9.4), plant defense mechanisms (e.g., antioxidants) (see Section 9.3), and site climate and soil factors (e.g., temperature, vpd, soil moisture) (see Section 9.4). In using these indices to develop air quality standards, the needs of policy makers must also be considered, and those include simplicity and understandability (Fairley and Blanchard, 1991). The development of such indices continues to be a challenge.

9.5.2 Summary of Conclusions from the Previous Criteria Document

The 1996 AQCD (U.S. Environmental Protection Agency, 1996a) focused primarily on development of exposure indices, not flux, to quantify growth and yield effects in crops, perennials and trees (primarily seedlings), and not foliar injury. The testing of the adequacy of these indices to order the measured responses of growth and/or yield in crops and tree species, as seedlings, was accomplished through regression analyses of earlier exposure studies. No direct experimental testing of the range of indices has yet been accomplished. It was recognized that these indices were only surrogates for O₃ uptake or dose. Their development focused on consideration and inclusion of some, but not all, factors that affect O₃ uptake and expression of effects (e.g., Lee et al., 1988). The 1996 document (U.S. Environmental Protection Agency, 1996a) drew a number of conclusions that built on even earlier conclusions (U.S. Environmental Protection Agency, 1986, 1992). These conclusions are still valid today based on a review of research published since 1996.

Studies prior to 1996, and after, indicate that the components of exposure, including concentrations, temporal dynamics (e.g., time of day of peak events), frequency of occurrence, duration, and respite time, are integral to developing indices of exposure that relate to growth response. Evidence from the few direct experimental studies of varying exposure components indicate the importance of peak concentrations, occurrence, respite time and the importance of

1 cumulative the concentrations (Musselman et al., 1983, 1986, 1994; Hogsett et al., 1885; U.S.
2 Environmental Protection Agency, 1996a).

3 Exposure duration influences the degree of plant response. Single season, year-long or
4 multi-year experimental results indicate that greater yield losses occurred when plants were
5 exposed for the longer duration, and that a cumulative-type index was able to better describe the
6 relationship between exposure and yield. Those indices not considering duration, e.g., 7-h
7 seasonal mean concentration index, single peak event index, or the index that cumulates all
8 concentrations (i.e., SUM00), were unable to adequately describe the relationship. These single
9 event or mean-type indices do not consider the role of duration of exposure and focus either only
10 on the peak event or put too much focus on the lower hourly average concentrations (U.S.
11 Environmental Protection Agency, 1996a).

12 Higher hourly averaged concentrations had a greater affect on plant response. It was
13 concluded that cumulative indices that gave greater weight to higher concentrations relate well
14 with plant response (crops and tree seedlings) and order the treatment means in monotonically
15 decreasing fashion with increasing exposure, based on studies that apply two or more types of
16 exposure regimes with replicate studies of the same species. These indices include, among
17 others: SUM06, W126, AOT40 (U.S. Environmental Protection Agency, 1996a).

18 No studies before or after 1996, have enabled a discrimination among the various
19 weighted, cumulative indices. Various functional weighting approaches were used, including
20 allometric, sigmoid or threshold weighting, and compared for best statistical fit of the plant
21 growth or yield data; but no one functional weighting was favored. For use as an air quality
22 standard, however, the need for simplicity, understandability, and ease of monitoring, favored
23 the cumulative threshold-weighted index (SUM06) (U.S. Environmental Protection Agency,
24 1996a).

25 Since higher concentrations occur primarily during the daylight hours, those indices that
26 differentially weight higher concentrations give greater weight to daylight hour concentrations.
27 This is important since stomatal conductance is usually greatest during the daylight hours,
28 compared to concentrations at night when conductance is usually minimal. Peak concentrations,
29 however, do not occur throughout the day; thus the timing of the peak concentration and
30 maximum plant uptake is critical in determining plant response. An exposure index that
31 incorporated either the daily or seasonal temporal patterns of higher concentration occurrence

1 with the temporal pattern of individual species' stomatal conductance was not reported in the
2 prior 1996 review (U.S. Environmental Protection Agency, 1996).

3 The relative importance of cumulative peak concentrations (> 0.10 ppm) versus cumulative
4 mid-range concentrations (0.05 - 0.099 ppm) was questioned based on ambient field exposures of
5 sensitive species. Although controlled experiments provided important evidence that the higher
6 hourly average concentrations should be given greater weight than the mid-level values in
7 developing indices, there was concern that under ambient conditions in the field the higher
8 concentrations did not occur at the time of maximum plant uptake. This coincidence was
9 considered to be the critical factor in determining peak concentration impacts on plants. Based
10 on the evidence at that time, it was not possible to conclude whether the cumulative effects of
11 mid-range concentrations were of greater importance than those of peak hourly average
12 concentrations in determining plant response (U.S. Environmental Protection Agency, 1996a).
13 No direct experimental studies had addressed this question prior to 1996, nor have any since.

14 The composite exposure-response functions for crops and tree seedlings were derived from
15 single and multi-year exposure studies that used modified or simulated ambient exposure
16 profiles. These profiles were typified by episodic occurrence of a large number of high O_3
17 concentrations; and this type of pattern is prevalent in many but not all rural agricultural and
18 some forested areas in the United States. Selecting a concentration value from these crop and
19 seedling response models may result in an over or underestimation of growth effects if applied to
20 regions of the country where a different type of exposure pattern is prevalent (U.S.
21 Environmental Protection Agency, 1996a). A multi-component index was suggested that
22 combined the peak-weighted, cumulative index with the number of occurrences of hourly
23 averaged concentrations ≥ 0.10 ppm that might reduce the uncertainty associated with selecting
24 the exposure value for protection based on NCLAN type studies (Lefohn and Foley, 1992;
25 Musselman et al., 1994; U.S. Environmental Protection Agency, 1996a). No direct experimental
26 studies addressed this question prior to 1996, nor have any since.

27 Since 1996, additional research has focused on the time of day when the higher hourly
28 average concentrations occur, the time of day of maximum plant uptake, and the diurnal
29 variability of plant defense mechanisms, and various suggestions as to inclusion of these factors
30 in any one of the peak weighted cumulative exposure indices. A much broader literature has

focused since 1996 on developing indices using flux to relate plant response. These new developments are discussed in the sections that follow.

9.5.3 Use of Exposure Indices to Establish Exposure-Response Relationships

Mathematical approaches for summarizing ambient air quality information in biologically meaningful forms that can serve as surrogates for dose for O₃ vegetation effects assessment purposes have been explored for more than 80 years (O'Gara, 1922; U.S. Environmental Protection Agency, 1996). Several of the indices introduced have attempted to incorporate some of the biological, environmental, and exposure factors (directly or indirectly) that influence the magnitude of the biological response and contribute to observed variability (Hogsett et al., 1988). In the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996a), the exposure indices were arranged into 5 categories, including (1) One Event, (2) Mean, (3) Cumulative, (4) Concentration Weighted, and (5) Multi-component and were discussed in detail (Lee et al., 1988). Figure 9-16 illustrates how several of the indices weight concentration and accumulate exposure.

The indices were developed with knowledge of O₃ modes of action, as well as the role of the exposure components and their adequacy tested using earlier exposure studies, e.g., NCLAN (U.S. Environmental Protection Agency, 1996a; Hogsett et al., 1988). Various components of the exposure-response, including concentration, time of day, respite time, frequency of peak occurrence, plant phenology, predisposition, etc., were weighted with various functions and evaluated on their ability in ordering the regression of exposure vs. growth or yield response. The statistical evaluations for each of these indices were accomplished using growth/yield response data from many earlier exposure studies (e.g., NCLAN). This retrospective approach was necessary because there have been very few studies specifically designed to test the goodness of fit of the indices. The regression approach selects those indices that most properly order and space the treatment means to optimize the fit of a linear or curvi-linear model. This approach provides evidence for the best indices, albeit not as defensible as that from studies with experimental designs and analyses that focus on specific components of exposure.

Most of the early retrospective studies reporting regression approaches use data from the NCLAN program or from Corvallis, OR (Lee et al., 1987; Lee et al., 1988; Lefohn and Irving, 1988; Tingey et al., 1989) or use data collected in California (Musselman et al., 1988; U.S.




Figure 9-16. Diagrammatic representation of several exposure indices, illustrating how they weight concentration and accumulate exposure. (a) SUM06 – The upper graphic illustrates an episodic exposure profile; the shaded area under some of the peaks illustrates the concentrations greater than or equal to 0.06 ppm that are accumulated in the index. The insert shows the concentration weighting (0 or 1) function. The lower portion of the graphic illustrates how concentration is accumulated over the exposure period. (b) SIGMOID – The upper graphic illustrates an episodic exposure profile; the variable shaded area under the peaks illustrates the concentration dependent weights that are accumulated in the index. The insert shows the sigmoid concentration weighting function. The mid-point of the sigmoid weighting scheme was 0.062 ppm. The lower portion of the graphic illustrates how concentration is accumulated over the exposure period. (c) 2ndHDM and M-7 – The upper graphic illustrates an episodic exposure profile. The lower portion of the graphic illustrates that the 2ndHDM considers only a single exposure peak while the mean applies a constant exposure value over the exposure period.

Source: Tinge et al. (1991).

- 1 Environmental Protection Agency, 1986). These studies previously reviewed by EPA (U.S.
- 2 Environmental Protection Agency, 1992, 1996a) and were in general agreement and consistently
- 3 favored the use of cumulative peak-weighted exposure indices. Lee et al. (1987) suggested that
- 4 exposure indices that included all the data (24 h) performed better than those that used only 7 h
- 5 of data; this is consistent with the conclusions of Heagle et al. (1987) that plants receiving

1 exposures for an additional 5-h/day showed 10% greater yield loss than those exposed for
2 7-h/day. In a subsequent analysis using more of the NCLAN data, Lee et al. (1988) found the
3 “best” exposure index was a phenologically weighted, cumulative index, with sigmoid weighting
4 on concentration and a gamma weighting function as a surrogate for plant growth stage. This
5 index was the best statistical fit, but it depended upon more knowledge of species and site
6 conditions that made specification of weighting functions difficult for general use. This type of
7 multi-function index did not meet the other criteria for developing an air quality standard, that
8 being simplicity and understandability (Blanchard and Farley, 1999).

9 The next best fits were the several indices which only cumulated and weighted higher
10 concentrations (e.g., SUM06, SUM08, AOT40, W126). Amongst this group it was not possible
11 to distinguish a single best fit (Lee et al., 1988). Similarly Lefohn et al., (1992) reported that it
12 was not possible to differentiate among the SUM00, SUM06, SUM08, and W126 exposure
13 indices because the indices were highly correlated with one another in the experiment. Others
14 have reported similar results when attempting to identify optimum exposure indices (Musselman
15 et al., 1988).

16 Other factors, including predisposition time (Hogsett et al., 1988; McCool et al., 1988) and
17 crop development stage (Heagle et al., 1991; Tingey et al., 2002), contribute to variation in the
18 biological response and suggest the need for weighting O₃ concentrations to account for
19 predisposition time and phenology. However, the roles of predisposition and phenology in
20 influencing plant response vary considerably with species and environmental conditions, so that
21 specification of a weighting function for general use in characterizing plant exposure is not
22 possible at this time.

23 In similar retrospective analyses, using data from the European Open-Top Chamber
24 Program, Finnan et al. (1997) confirmed that cumulative exposure indices which emphasize
25 higher O₃ concentrations are best related to plant response and that cumulative exposure indices
26 which use weighting approaches provide a better fit (U.S. Environmental Protection Agency,
27 1996a).

28 The main conclusions from 1996 regarding a biologically-relevant index based on ambient
29 exposure still hold true today. No information has come forth in the interim that alters those
30 conclusions, and in fact, some recent studies have further substantiated them. These key
31 conclusions can be restated as follows:

- ozone effects are cumulative;
- peak concentrations appear to be more important than lower concentrations in eliciting a response;
- plant sensitivity to O₃ varies with time of day and crop development stage;
- exposure indices that accumulate the O₃ exposure and preferentially weight the peaks yield better statistical fits to response than do the mean and peak indices;
- to avoid possible overestimation of vegetation effects when using NCLAN-type yield data in exposure-response models, the cumulative exposure index might be combined with number of hours of high hourly average concentrations over the time period.

Following the 1996 review process (U.S. Environmental Protection Agency, 1996a,b), the EPA proposed an alternative form of the secondary NAAQS using a cumulative, concentration-weighted exposure index, the SUM06, to protect vegetation from damage (Federal Register, 1997). The EPA considered three specific concentration-weighted indices: the threshold weighted SUM06, the AOT40, and the sigmoid weighted W126 exposure index (U.S. Environmental Protection Agency, 1996b). Both indices performed equally well in predicting the exposure-response relationships observed in the crop and tree seedlings studies conducted during the prior 20 years (Heck and Cowling, 1997). In the absence of research results that differentiate the predictive power of these two forms, the EPA selected the SUM06 exposure index as the form for the proposed secondary standard, recognizing its simplicity, understandability, and ease of use (U.S. Environmental Protection Agency, 1996b). A 3-month, 12-hour SUM06 exposure index of 26.4 ppm hr was proposed (U.S. Environmental Protection Agency, 1996b). The value represented the concentration level that would protect 50% of the crop species from a 10% yield loss. The composite response model included 49 studies comprising 13 crop species and 5 locations across the U.S. where the crops are grown (U.S. Environmental Protection Agency, 1996a).

European scientists took a similar approach as the United States in developing indices describing growth and yield loss in crops and tree seedlings, using open-top chambers with modified ambient exposures, but many fewer crop species were employed in the European studies. The European countries were seeking a scientific basis for control strategies to reduce air pollution. They adopted the critical levels and loads approach, as per the UN Economic Commission for Europe (UNECE). A critical level was defined as “the concentration of

1 pollutant in the atmosphere above which direct adverse effects on receptors, such as plants,
2 ecosystems, or materials may occur according to present knowledge” (United Nations Economic
3 Commission for Europe, 1988). Critical levels are set to prevent long-term injury and damage to
4 the most sensitive elements of any ecosystem. They are used to map and identify areas in
5 Europe in which the levels are exceeded and that information is then used to plan optimized and
6 effect-based abatement strategies. As used by the UNECE, they are not air quality standards in
7 the US sense, but they have been used as targets for planning reductions in sulfur and nitrogen
8 emissions to protect ecological resources. The nature of the significant harmful effects is not
9 specified in the original definition, which provides for different levels for different types of
10 harmful effect (e.g., visible injury or loss of crop yield). There are also different levels for crops,
11 forests, and semi-natural vegetation. The caveat, “according to present knowledge,” is important
12 because critical levels are not rigid; they are revised periodically as new scientific information
13 becomes available. For example, the original critical level for O₃ specified concentrations for
14 three averaging times, but further research and debate led to the current critical levels being
15 stated as the cumulative exposure (concentration x hours) over a cut-off concentration of 40 ppb
16 (AOT40) (Fuhrer et al., 1997). The “Level I” critical level was used in the 1990s to map areas of
17 exceedance, but analyses of many exposure studies led to the conclusion that the simple,
18 exposure-based level leads to over-estimation of the effects in some regions and
19 under-estimation in others (Fuhrer et al., 1997; Kärenlampi and Skärby, 1996). The main
20 problem was that other environmental factors (vapor pressure deficit, water stress, temperature,
21 and light) and variation in canopy height altered O₃ uptake and its effects.

22 A decision was made to work towards a flux-based approach for the critical level, with the
23 goal of modeling O₃ flux-effect relationships for three vegetation types: crops, forests, and
24 semi-natural vegetation. Progress has been made in modeling flux (e.g., Grunhage and Jager,
25 2003; Ashmore et al., 2004a; Ashmore et al., 2004b) and the Mapping Manual is being revised
26 (Karlsson et al., 2003; Ashmore et al., 2004a; Ashmore et al., 2004b; Grennfelt, 2004). The
27 revisions may include a flux-based approach for 3 crops: wheat, potatoes, and cotton; but,
28 because of lack of flux data, a cumulative, concentration-based (AOTx) exposure index will
29 remain for most crops, and for forests and semi-natural herbaceous vegetation (Ashmore et al.,
30 2004; Jaeger and Grunhage, 2004).

9.5.4 Identifying Exposure Components That Relate to Vegetation Effects

The efficacy of exposure indices to predict biological response requires that researchers identify a relationship between exposure components and effects, as well as those environmental and site factors that control pollutant uptake by the plant. These relationships were identified and discussed in the 1996 review (U.S. Environmental Protection Agency, 1996a).

A significant, but in some instances, unquantified role was identified for: (1) concentration; (2) duration of exposure; (3) the diurnal and seasonal patterns of exposure, e.g., time of day of peak event, season of higher exposures, seasons of high precipitation and humidity, the frequency of occurrence of peak events and respite time (peak to valley ratios); (4) plant phenology; (5) plant canopy structure; (6) meteorological and site factors, e.g., light, humidity; and (7) plant defense mechanisms.

9.5.4.1 Role of Concentration

A significant role of higher concentrations was established earlier, based on several experimental studies (U.S. Environmental Protection Agency, 1996a). Recently Nussbaum et al., (1995) and Yun and Laurence (1999b) have added support for the important role that peak concentrations, as well as the pattern of occurrence, plays in plant response to O₃. Based on air quality data from 10 U.S. cities, three treatments of 4-week exposure to the same SUM06 value were constructed by Yun and Laurence (1999b). They used the regimes to explore effects of treatments with variable peak occurrence versus uniform peak occurrence during the exposure period. The authors reported that the peak exposures were important and that the same SUM06 resulted in very different foliar injury. Oksanen and Holopainen (2001) found that the peak concentrations and the shape of the O₃ exposure (i.e., duration of the event) were important in foliar injury of white birch saplings, but growth reductions were found to be more related to cumulative exposure. Nussbaum et al. (1995) also found peak concentrations and the pattern of occurrence to be critical in the measured response. The authors recommended that to describe the effect on total forage yield, peak concentrations > 0.11 ppm must be emphasized by using an AOT with higher threshold concentrations.

A greater role for higher concentrations affecting plant growth may be inferred based on recent air quality analyses for the Southern California area (Lee et al., 2003; Tingey et al., 2004). In the late 1960s and 1970s, extremely high O₃ concentrations had impacted the San Bernardino

1 National Forest. However, over the past 15 plus years, significant reductions in the O₃ exposure
2 have occurred in the San Bernardino National Forest (Lee et al., 2003; Lefohn and Shadwick,
3 2000; Davidson, 1993; Lloyd et al., 1989). An illustration of this improvement in air quality is
4 shown by the 37 year history of ozone air quality at a site in the San Bernardino Mountains
5 (Figure 9-17) (Lee et al., 2003). The O₃ exposure increased from 1963 to 1979 concurrent with
6 increased population and vehicular miles, followed by a decline to present mirroring decreases in
7 precursor emissions. The pattern in exposure was evident in various exposure indices including
8 the cumulative concentration weighted (SUM06), as well as maximum peak event (1-h peak),
9 and the number of days having hourly averaged O₃ concentrations \geq 95 ppb (i.e., the California
10 ozone standard). The number of days having hourly averaged O₃ concentrations \geq 95 ppb
11 declined significantly from 163 days in 1978 to 103 days in 1997. The changes in ambient
12 ozone air quality for the site were reflected in the changes in the frequency and magnitude of the
13 peak hourly concentration and the duration of the exposure (Figure 9-17). Considering the role
14 of exposure patterns in determining response, the seasonal and diurnal patterns in hourly O₃
15 concentration did not vary appreciably from year to year over the 37-year period (Lee et al.,
16 2003).

17 The inference for a role of higher concentrations comes from both results of ground
18 measures of tree conditions on established plots and from model simulations. Across a broad
19 area of the San Bernardino National Forest, the Forest Pest Management (FPM) method of injury
20 assessment indicated an improvement of crown condition from 1974 to 1988; and the area of
21 improvement in injury assessment is coincident with an improvement of O₃ air quality (Miller
22 and Rechel, 1999). A more recent analysis of forest changes in the San Bernardino National
23 Forest using an expanded network of monitoring sites has verified significant changes in growth,
24 mortality rates, basal area, and species composition throughout the area since 1974 (Arbaugh
25 et al., 2003). A model simulation of ponderosa pine growth over the 40 year period in the
26 San Bernardino showed a significant impact of ozone exposure on tree growth and indicates
27 improved growth with improving air quality. The improvement in growth was assigned to
28 improved air quality, but no distinction was made regarding the relative role of mid-range and
29 higher hourly concentrations, only that improved growth tracked both decreasing SUM06,
30 maximum peak concentration and number of days of hourly O₃ \geq 95 ppb (Tingey et al., 2004).




Figure 9-17. Trends in May-September 12-h SUM06, peak 1-h ozone concentration and number of daily exceedances of 95 ppb for Crestline in 1963-1999 in relation to trends in mean daily maximum temperature for Crestline and daily reactive organic gases (ROG) and oxides of nitrogen (NO_x) for San Bernardino county. Annual ROG and NO_x emissions data for San Bernardino county were obtained from Alexis et al. (2001) and the California Air Resource Board's emission inventory available at <http://www.arb.ca.gov/emisinv/emsmain/emsmain.htm>.

Source: Lee et al. (2003).

9.5.4.2 Role of Duration

Recent studies have called into question the period of time over which concentrations are accumulated and the form of the exposure index. Heagle and Stefanski (2000) reported that the form of the exposure index was important only for 24-h indices for which SUM00 provided the poorest fit. The authors reported that the SUM00, SUM06, W95 (Lefohn and Runeckles, 1987), W126, and AOT40 produced similarly good fits of the foliage biomass data for 6-, 5-, and 4-h midday accumulating periods. The study “pooled” data from San Bernardino (CA) and Riverside (CA) with data from Amherst (MA), Corvallis (OR), Kennedy Space Center (FL), Raleigh (NC), and Blacksburg (VA). Ozone exposures were much higher at the two California sites (indicated by high W126, SUM06, W95, and AOT40 values) in comparison to the other locations. Because of the pooling of the data, the large number of high hourly average O₃ concentrations that occurred at the California sites may have resulted in the exposure indices being highly correlated with one another and made it difficult to identify one optimal index.

In another study in California, Arbaugh et al. (1998) reported that the SUM00 exposure index performed better for describing visible injury than the SUM06, W126, number of hours greater than or equal to 0.08 ppm, and the number of days between measurement periods. These exposure indices were originally developed and tested using only growth/yield data, not foliar injury (U.S. Environmental Protection Agency, 1996a). This distinction is critical in comparing the efficacy of one index to another.

The SUM00 exposure index is a surrogate for a long-term average concentration and earlier growth response studies indicated that a long-term average was not adequate in predicting effects primarily because it gives equal weight to all concentrations (U.S. Environmental Protection Agency, 1996). However, for many locations in California, a large number of higher hourly average concentrations occur and the SUM00 could be highly correlated with the frequency of elevated hourly average concentrations and thus could be a good predictor of vegetation effects. In Section 9.3, it was noted that research results are mixed on what causes an effect in plants: an event or the cumulation of events.

9.5.4.3 Patterns of Exposure

A significant factor in developing exposure indices is the temporal patterns of ozone occurrence over a day, a month, a year, and seasonally overlaying the daily and seasonal

temporal patterns of those influential climatic and site factors. The coincidence of peak ozone and maximal stomatal conductance and detoxification processes are key to effecting plant growth response (Musselman and Minnick, 2000).

Daily Patterns

Recent experimental studies with tree species have demonstrated the de-coupling of ambient O₃ exposure, peak occurrence, and gas exchange, particularly in areas of drought (Panek, 2004). The coincidence of maximal conductance and occurrence of higher concentrations are relevant to the time of day that exposure is cumulated in those indices that cumulate and weight concentration.

The hours during the day over which ambient exposure is cumulated or effective dose is estimated is a surrogate for the coincidence of these patterns of conductance and O₃ occurrence. A 12-hr daylight period for cumulating exposure was proposed following the 1996 review (U.S. Environmental Protection Agency, 1996). An extensive review of the literature revealed that a large number of species had varying degrees of nocturnal stomatal conductance (Musselman and Minnick, 2000). Grulke et al. (2003) showed that the stomatal conductance at night for ponderosa pine in the San Bernardino (CA) National Forest ranged from one tenth to one fourth that of maximum daytime gas exchange. In June, at the high-elevation site, 11% of the total daily O₃ uptake of pole-sized trees occurred at night. In late summer, however, O₃ uptake at night was negligible. Birch seedlings exposed to O₃ at night show greater reductions in growth than those exposed to O₃ in daylight (Matyssek et al., 1995). *Brassica rapa* plants exposed to O₃ during the day or night, show little significant difference in the amounts of injury or reduced growth response to treatment; the conductance was 70 to 80% lower at night (Winner et al., 1989). Tissue biomass of ponderosa pine seedlings was significantly reduced when seedlings were exposed to either daytime or nighttime episodic profiles (Lee and Hogsett 1999). However, the biomass reductions were much greater with daytime peak concentrations than with nighttime peak concentrations.

Although conductance is lower at night than during the day for most plants, nocturnal conductance can result in some measurable O₃ flux into the plants and should be considered. Nocturnal O₃ flux also depends on the level of turbulence that intermittently occurs at night. In addition, plants can be more susceptible to O₃ exposure at night than during the daytime,

1 because of lower plant defenses at night (Musselman and Minnick, 2000). Massman (2004)
2 suggested that nocturnal stomatal O₃ uptake accounted for about 15% of the cumulative daily
3 effective O₃ dose that was related to predicted injury. Based on a review of the literature relating
4 to nocturnal stomatal conductance, Musselman and Minnick (2000) recommended that any O₃
5 exposure index used to relate air quality to plant response should use the 24-h cumulative
6 exposure period for both exposure-response and effective flux models.

7 In general, stomatal conductance needs to be taken into account in developing indices
8 (Panek et al., 2002). Stomatal conductances are linked to both diurnal and seasonal
9 meteorological activity and soil/site conditions (e.g., soil moisture). Daily patterns of
10 conductance are often highest in mid-morning, whereas higher ambient O₃ concentrations
11 generally occur in mid to late afternoon, when stomata are often partially closed and
12 conductances are lower. Total flux depends on atmospheric resistance and boundary layer
13 resistances, both of which exhibit variability throughout the day. Several recent studies have
14 suggested that ponderosa pine trees in the southern and northern Sierra Nevada Mountains may
15 not be as sensitive to high O₃ concentrations as to lower concentrations, due to reduced O₃
16 uptake during the period when the highest concentrations occur (Panek et al., 2002; Panek and
17 Goldstein 2001; Bauer et al., 2000; Arbaugh et al., 1998). Panek et al. (2002) compared direct
18 measurements of ozone flux into a canopy of ponderosa pine and demonstrated a lack of
19 correlation of daily patterns of conductance and ozone occurrence, especially in the late season
20 drought period; and they concluded that a consideration of climate or season was essential,
21 especially considering the role of soil moisture and conductance/uptake. In contrast, Grulke
22 et al. (2002) reported high conductance when O₃ concentrations were high in the same species,
23 but different growing site conditions. The uncoupling of conductance and higher ambient O₃
24 concentration would hold true for more mesic environments as well as xeric landscapes. The
25 longer term biological responses reported by Miller and Rechel (1999) for ponderosa pine in the
26 same region and the general reduction in recent years in ambient O₃ concentrations, suggest that
27 conductance alone may not be a sufficient indicator of potential vegetation injury or damage.

28 The generalized models of stomatal conductance may provide a means to link patterns of
29 O₃ occurrence with climatic and site factors that affect O₃ uptake to some degree, provided
30 conductance is modeled by regions of similar seasonal moisture and by similar canopy structure
31 (e.g., Emberson et al., 2000a,b; Grunhage et al., 2000; Massman, 2004)

Seasonal Patterns

Several of the recent studies measuring O₃ flux to pine canopies also reported on the importance of seasonal patterns in relating exposure to response (Panek, 2004). These seasonal patterns can be early vs late season occurrence of higher O₃ concentrations, reflecting climate and precursor availability. The patterns also reflect seasonal drought and the role of soil moisture plays in stomatal conductance and O₃ uptake. Recently, studies have looked directly at this linkage. Panek et al. (2002) compared direct measurements of ozone flux into a canopy of ponderosa pine with a number of exposure indices, demonstrated a lack of correlation especially in the late season drought period, and concluded that a consideration of climate was essential, especially soil moisture. They suggested that a better metric for a seasonally drought-stressed forests would be one that incorporates forest physiological activity, either through mechanistic modeling, by weighting ambient O₃ concentrations by stomatal conductance, or by weighting O₃ concentrations by site moisture conditions. Panek (2004) demonstrated a decoupling of O₃ exposure and uptake seasonally as well, via seasonal drought influence. Maximum O₃ uptake occurred at the beginning of the season and in the winter, whereas the pines were nearly dormant during August-September.

Using TREGRO, a process-based model, Tingey et al. (2004) simulated long-term growth of ponderosa pine over a 37-year period. The simulation showed high correlation between O₃ exposure and O₃-induced reductions in tree growth ($R^2 = 0.56$). The scatter about the line however indicates that other factors besides O₃ are required to describe the association between exposure and response. Incorporating annual temperature and precipitation increased the R^2 to 0.67. In keeping with the observations of Panek (2004) on the decoupling of peak O₃ occurrence and maximal conductance, the remaining unexplained variation is attributed to differences in timing of peak O₃ uptake and peak O₃ exposure over the years.

9.5.4.4 Frequency of Occurrence of Peak Concentrations

Several earlier studies demonstrated the greater effect of episodic occurrence of O₃ peaks compared to daily peak events (U.S. Environmental Protection Agency, 1996a); and, since the last review, a few studies have corroborated the importance of this pattern in growth response. Köllner and Krause (2003) reported that, under equal exposure conditions, the most pronounced effects on the yield of sugar beet and soybeans occurred with those regimes that emphasized the

episodic occurrence of peak events. Nussbaum et al. (1995) compared the effects of different patterns of peak occurrences with similar AOT40 values and reported a stronger effect on total forage yield from the episodic treatment.

9.5.4.5 Canopy Structure

Another factor important in either O₃ exposure or uptake, is how canopy structure affects O₃ concentration in and under forest canopies. There have been several comprehensive studies of O₃ concentrations under tree canopies (Enders, 1992; Fontan et al., 1992; Fredericksen et al., 1995; Joss and Graber, 1996; Kolb et al., 1997; Lorenzini and Nali, 1995; Neufeld et al., 1992; Samuelson and Kelly, 1997). In general they indicate a reduction in O₃ of ~20 to 40% below the canopy but above the shrub/herb layers. An essential component in the critical level AOT40 is the height of where the O₃ concentration is measured. The critical levels are related to the O₃ concentration measured at the top of the canopy, i.e., upper surface boundary of the (quasi-) laminar layer (Grunhage et al., 2003). This location is presumably more related to stomatal uptake. Weighting the O₃ concentration at this location takes into account stomatal opening and, if weighted with the Jarvis-Steward factors for radiation, temperature, and soil moisture, the “toxicologically” effective AOT40 is obtained (Grunhage et al., 2003).

In a study that considers those factors important in O₃ uptake that are also spatially distributed as a result of canopy structure, Davison et al. (2003) reported that the variation in visible injury in coneflower populations is unlikely to be due to differences in O₃ flux and more likely due to variation in PAR. At a height of 50 cm above ground, PAR was reduced by almost 90%, whereas the O₃ varied from about 15 to 90% of ambient. Ozone injury was not solely related to O₃ flux. Although there have been studies of the effects of different light levels on O₃ response, there have been few at the very low levels that occur in canopies of tall herbaceous stands or in the ground layer of forests. Davison et al. (2003) report that conductance was not related to diurnal changes in light. The O₃ levels were still about 90% of the O₃ concentration above the canopy when light was less than 5%. Light intensity dropped to 1.5% of open at 130 cm from the edge of the canopy, while O₃ dropped to only 42%. The study, although reporting on the adequacy of visible foliar injury as an indicator of O₃ effects, does suggest that consideration of other factors such as light are important in predicting response. How this may be included in developing exposure-response indices was not considered.

9.5.4.6 Site and Climate Factors

Soil moisture is a critical factor in controlling O₃ uptake through its effect on plant water status and stomatal conductance. In an attempt to relate uptake, soil moisture, and ambient air quality to identify areas of potential risk, available O₃ monitoring data for 1983 to 1990 were used, along with literature-based seedling exposure-response data from regions within the southern Appalachian Mountains that might have experienced O₃ exposures sufficient to inhibit growth (Lefohn et al., 1997). In a small number of areas within the region, O₃ exposures and soil moisture availability were sufficient to possibly result in growth losses in some sensitive species (e.g., black cherry). The conclusions were limited because of the interpolation of the O₃ exposures in many of the areas and the hydrologic index used might not reflect actual water stress.

9.5.4.7 Plant Defense Mechanism - Detoxification

The non-stomatal component of plant defenses are the most difficult to quantify, but some studies are available (Barnes et al., 2002; Plöchl et al., 2000; Chen et al., 1998; Massman and Grantz, 1995). Massman et al. (2000) developed a conceptual model of a dose-based index to determine plant injury response to O₃ that relates to the traditional exposure-based parameters. The index uses time-varying-weighted fluxes to account for the fact that flux is not necessarily correlated with plant injury or damage. Their model applies to plant foliar injury, and they suggest that application of flux-based models for determining plant damage (yield or biomass) will require a better understanding and quantification of the injury and damage relationship.

9.5.5 Ozone Uptake or Effective Dose as an Index

Developing an index that relates growth response to ambient exposure has been approached in the past through various weighting functions on those ambient exposure factors, including concentration, duration, and time of day for cumulating exposure. These indices have not incorporated factors of climate patterns and species- and site factors that control O₃ uptake via canopy and stomatal conductance. The other approach is basing the index on the O₃ concentration going into the leaf, or flux. This approach includes those factors controlling uptake through canopy and stomatal conductance and, by necessity, relies on models to predict flux and ultimately the “effective” flux: that concentration that reaches the target tissue to cause

an effect (Grunhage et al., 2004; Massman 2004; Massman et al., 2000) .Effective flux has been defined as the balance between the O₃ flux and the detoxification process (Dämmgen et al., 1993; Grunhage and Haenel, 1997; Musselman and Massman, 1999). The time-integrated effective flux is termed “effective dose”. The uptake mechanisms and the resistances in this process, including stomatal conductance and biochemical defense mechanisms, are discussed in the previous Section 9.3.

9.5.5.1 Models of Stomatal Conductance

Only a limited number of studies have measured O₃ concentration or its reaction products within the leaf. Otherwise the index of uptake is a modeled result that considers site, climatic, meteorological, and species-specific (e.g., detoxification reactions) factors. Models of O₃ conductance into plant tissue are available (Grunhage and Jager, 2003; Emberson et al., 2000a,b; Grunhage and Haenel, 1997; Grünhage et al., 1997; Massman, 1993; Wesely, 1989). The European Monitoring and Evaluation Program (EMEP) has an O₃ deposition model developed for application across Europe in conjunction with the EMEP photochemical model as a tool for the critical levels program. The model has been developed to estimate vegetation type-specific O₃ deposition and stomatal flux, calculated according to a standard 3-resistance formulation incorporating atmospheric, boundary layer, and stomatal resistances (Emberson 2000b). The model uses a multiplicative algorithm of the stomatal conductance of O₃ (Jarvis, 1976) and has been parameterized for 10 European tree species, 7 agricultural species, and 1 type of semi-natural vegetation. The model calculates conductance as a function of leaf phenology, temperature, photosynthetic flux density (PFD), vapor pressure deficit (VPD), and soil moisture (SMD). The environmental variables are site-specific (or regionally-specific). The most important factors limiting O₃ with this model were vapor pressure deficit (VPD), soil moisture deficit, and phenology (Emberson et al., 2000b). These factors demonstrate the critical linkage of high VPD and stomatal closure, which typically co-occurs with high O₃ concentrations.

A number of recent model-based studies relating flux and plant growth response have investigated several crop and forest tree species (Karlsson et al., 2004; Karlsson et al., 2004; Pleijel et al., 2004; Matyssek et al., 2004; Elvira et al., 2004; Soja et al., 2004; Weiser and Emberson, 2004; Touvinen et al., 2004; Altimir, et al., 2004; Gerosa et al., 2004; Mikkelsen, et al., 2004, Bassin, et al., 2004; Emberson et al., 2000). The studies have used earlier exposure

1 experiments as well as explicitly designed field studies but, to date, no abundance of data that
2 would provide the basis for a flux-based index.

3 The complexity of using flux as an index of O₃ exposure for response is shown in field
4 studies that measured O₃ flux into Norway spruce and cembra pine (Wieser et al., 2000). They
5 demonstrated that stomatal conductance was the main limiting factor for O₃ uptake and showed
6 the dependence of that measure on crown position, needle age, and altitude. Consideration of the
7 role of climate illustrates the importance of a flux measure. Pleijel et al., (2000) reported the
8 improved relationship of yield in spring and winter wheat grown in OTCs in many areas across
9 Europe when it was related to the cumulative stomatal O₃ uptake during the grain-filling period.
10 Compared to the AOT40, the cumulative uptake index estimated larger yield losses in the
11 relatively humid parts of western and northern Europe, while smaller yield loss was estimated
12 for the dry summer climates in south and central Europe.

13 Danielsson et al. (2003) compared the ability of two different stomatal models to relate
14 grain yield in field grown spring wheat to cumulated O₃ uptake and an exposure index of
15 AOT40, and found that the cumulated O₃ uptake determined with either model performed better
16 in relating exposure to yield than did the cumulative exposure index of AOT40.

17 Cumulative O₃ uptake was modeled for three deciduous and 2 coniferous species growing at
18 different sites and elevations and compared with exposure measure of AOT40 at these sites
19 (Matyssek et al., 2004). A general linearity was demonstrated between the two measures of O₃
20 exposure; and, at any given AOT40, there was a $25 \pm 11\%$ variation in CU. Although no
21 correlation of growth alterations was observed with either the exposure or the uptake measure,
22 the modeled cumulative uptake was able to describe the variation in tree size and site location
23 which makes for a better measure in risk assessment of O₃ (Matyssek et al., 2004). Karlsson
24 et al. (2004) compared the biomass-response relationship in young trees at seven experimental
25 sites across Europe using modeled cumulative O₃ uptake and AOT40. A weaker dose-response
26 relationships were reported for the cumulative uptake metric than the AOT40 (Karlsson et al.,
27 2004),

28 Concern about the complexity of the stomatal models and the data needed to model O₃
29 uptake, has led some researchers to offer modified accumulated exposure indices that consider
30 those meteorological factors controlling uptake (Karlsson et al., 2004; Gerosa et al., 2004). In a
31 study of subterranean clover in Austria, Belgium, and southern Sweden, Karlsson et al., (2004)

1 reported on the performance of a modified accumulated exposure over the threshold (mAOT)
2 which was based on solar radiation and vapor pressure deficit. This index improved the
3 relationship for observed visible injury. But if modeled uptake of O₃ was derived from a simple
4 stomatal conductance model considering solar radiation, VPD, and air temperature, then this
5 index gave an even greater improvement in the relationship to visible injury than did the ambient
6 exposure index of AOT40 (Karlsson et al., 2004B). The added value of the mAOT was
7 worthwhile and it had a lower degree of complexity and data requirements modeling O₃ uptake
8 with stomatal models. Based on a study of O₃ fluxes over a barley field in Italy, a similar
9 modified exposure index was reported and referred to as “effective exposure” (Gerosa et al.,
10 2004). Their approach was similar in its consideration of physiological aspects in conjunction
11 with monitored O₃ concentrations. It addressed the shortcoming of the data needs for modeled
12 O₃ uptake.

13 Models that partition O₃ uptake into stomatal and non-stomatal components are also now
14 available and predict a significant non-stomatal component in calculating O₃ flux (Altimir et al.,
15 2004; Mikkelsen et al., 2004; Nikolov and Zeller, 2003; Zeller and Nikolov, 2000; Bassin et al.,
16 2004; Nussbaum et al., 2003). Altimir et al. (2004) compared the relative contribution of
17 stomatal and non-stomatal sinks at the shoot level for Scots pine. Using the EMEP model with a
18 revised parameterization for Scots pine, they demonstrated that a major removal of O₃ was due
19 to the non-stomatal component; and when a non-stomatal term was introduced dependent on
20 ambient relative humidity, the non-stomatal contribution to the total conductance was about
21 50%. Zeller and Nikolov (2000) demonstrated a large non-stomatal O₃ uptake (41% of the total
22 annual flux) in subalpine fir at a site in southern Wyoming using the biophysical model
23 FORFLUX. In a 5 year study of measured O₃ flux to a Norway spruce canopy, Mikkelsen et al.
24 showed monthly patterns of non-stomatal and stomatal deposition as part of total deposition to
25 the canopy. Their study demonstrated that daily means of O₃ concentration and fluxes averaged
26 over 5 years correlate well, but the correlation is based on two different uncoupled processes
27 outside and inside the stomates. The destruction of O₃ in the canopy outside the stomates is
28 influenced by temperature, light and humidity, e.g., surface reactions, NO- and VOC-emissions;
29 and these same factors influence stomatal opening, e.g., mid-day and night closure.
30 Consequently, the diurnal O₃ concentration and O₃ flux do not correlate at all during the growing
31 season. The study estimated yearly stomatal uptake to be a minimum of 21% of total deposition

(i.e., non-stomatal as high as 80% of total). The stomatal uptake is highest in May-August (30-33%) and lowest in November-February (4-9%).

9.5.5.2 Nonlinear Response and Developing Flux Indices

If only O₃ flux were used as a metric to predict vegetation injury or damage, the prediction might be overestimated because of reported non-linear relationships between O₃ and plant response (Amiro et al., 1984; Amiro and Gillespie, 1985; Bennett, 1979; U.S. Environmental Protection Agency, 1978; 1986;1996). The non-linearity in the response surface suggested the existence of a biochemical threshold. More recently, nonlinear relationships between O₃ flux and yield were shown for potato (Pleijel et al.,2002) and spring wheat (Danielsson et al.,2003). The relationship between O₃ flux and potato yield suggested using an instantaneous flux threshold to overcome the nonlinear relationship (Pleijel et al., 2002). However, the authors did not report a substantial improvement in the mathematical fitting of the model when applying the threshold. Most of the flux was accumulated below 0.06 ppm. Danielsson et al. (2003) was able to show an improved relationship between O₃ uptake and yield of spring wheat using a threshold of 5 nmoles m⁻² sec⁻¹ (0.24 mg m⁻² sec⁻¹). These results suggest not all O₃ entering the stomata contribute to a reduction in yield, which depends to some degree on the amount of internal detoxification occurring with each particular species (see Section 9.3). The fact that the defense and repair mechanisms vary diurnally as well as seasonally may make it extremely difficult to apply a mathematically determined threshold to instantaneous flux measurements to calculate cumulative flux. The threshold models do not allow for the temporal (i.e., daily and seasonal) variability of defense mechanisms.

Musselman and Massman (1999) suggest that those species having high amounts of detoxification potential may show less of a relationship of O₃ stomatal uptake to plant response. There has been no direct experimental demonstration of this relationship, however. The cellular detoxification reactions and repair processes which both detoxify oxidants as well as play central role in the carbon economy of the plant are another level of resistance to O₃ reaching the target tissue (see Section 9.3). As indicated earlier, effects occur on vegetation when the amount of pollutant absorbed exceeds the ability of the plant to detoxify O₃ or to repair the initial impact. The magnitude of the response is determined by the amount of the pollutant reaching the target site and the ability of the plant to reestablish homeostatic equilibrium. Thus, one would expect

1 to observe a decoupling of O₃ uptake with vegetation effects, which would manifest itself as a
2 nonlinear relationship between O₃ flux and injury or damage.

3 Additional factors for inclusion in flux-based models to predict vegetation effects are the
4 defense and repair mechanisms. Specifically, the relationship between conductance, O₃
5 concentration, and defense/repair mechanisms needs to be included. Recently, Massman (2004)
6 described results that illustrate that the combination of conductance, O₃ concentration, and
7 diurnal variation of defense mechanisms show the daily maximum potential for plant injury
8 (based on effective dose) coincides with the daily peak in O₃ mixing ratio. Massman et al.
9 (2000) stress that the product of the overlapping mathematical relationships of conductance,
10 concentration, and defense mechanisms results in a much different picture of potential impact to
11 vegetation than just the use of conductance and concentration in predicting vegetation effects.
12

13 **9.5.5.3 Simulation Models**

14 Another approach for determining O₃ uptake and relating growth response to ambient O₃
15 exposure may be the use of physiologically-based simulation models. Several of these have
16 been used in various contexts, comparing O₃ response in a number of tree species with varying
17 climate and site factors (e.g., soil moisture) (e.g., Laurence et al., 2001; Ollinger et al., 1997;
18 Ollinger et al., 1998; Weinstein et al., 2001; Weinstein et al., 2004; Tingey et al., 2000; Tingey
19 et al., 2004). One of the important considerations in applying simulation modeling is to
20 carefully assess the uncertainties associated with the modeling predictions. Further efforts need
21 to be made to exercise the models so that they predict past growth losses associated with changes
22 in O₃ exposures that can be verified with on-the-ground surveys.
23

24 **9.5.6 Summary**

25 A few studies published since 1996 have substantiated earlier conclusions on the role of O₃
26 exposure components (including concentration, duration and exposure patterns) in describing
27 growth response to O₃ exposures. Recent studies of different exposure patterns have confirmed
28 earlier studies on the importance of higher concentrations and duration of exposure when
29 describing response. An inferred role of peak concentrations is possible from consideration of
30 improved air quality in regions like the San Bernardino Mountains in southern California.

1 Studies provide the basis for focusing on the higher O₃ concentrations, while including the lower
2 levels, when estimating the effects of emission reductions on vegetation.

3 A few studies have demonstrated the potential disconnection of peak events and maximal
4 stomatal conductance. In addition, a few studies have demonstrated the uptake of O₃ during
5 nighttime hours and suggested the need to cumulate O₃ exposure 24 h per day and not just during
6 daylight hours.

7 Several studies since 1996 have demonstrated another critical concern in developing an
8 index for exposure. The concern is that peak O₃ events and maximum stomatal conductance
9 may be temporally separate. This disconnection introduces uncertainty in assessing O₃ impact
10 when using the current ambient exposure based cumulative, concentration weighted indices.
11 If stomatal conductance is relatively low, as in the late afternoon in arid climates, and that is the
12 same time as the peak O₃ concentrations, then use of an exposure index that does not consider
13 this disconnect, will overestimate the effect of the exposure. This concern is especially apparent
14 when assessing the impact of O₃ across all the varied climatic regions of the United States or
15 Europe. Some studies use stomatal models to predict uptake or using process level models (e.g.,
16 TREGRO) to integrate those climate and site factors that drive this temporal pattern of stomatal
17 conductance and exposure, and thus reduce some of the uncertainty in regional to national
18 assessments of effects. These approaches however are still species dependent.

19 The results of these studies and reviews have indicated the need to continue to develop
20 indices that are more physiologically and meteorologically connected to the actual dose of O₃ the
21 plant receives. The cumulative concentration-weighted exposure indices are acknowledged
22 surrogates for effective dose that are simple conceptually and easy to measure. They do not fully
23 characterize the potential for plant uptake and resulting effects associated with O₃ because the
24 indices, being measures of ambient concentration, do not include the physical, biological, and
25 meteorological processes controlling the transfer of O₃ from the atmosphere through the leaf and
26 into the leaf interior. Use of such indices is especially limited in spatial risk characterizations
27 because of the lack of linkage of meteorology, species and site-specific factors influencing O₃
28 uptake. The flux-based approach should provide an opportunity to improve upon the
29 concentration-based (i.e., exposure indices) approach. A cautionary argument was advanced in a
30 few publications centered around the non-linear relationship between ozone uptake and plant
31 injury (not growth alterations) response. The concern was that not all O₃ stomatal uptake results

1 in a reduction in yield, which depends to some degree on the amount of internal detoxification
2 occurring with each particular species. Those species having high amounts of detoxification
3 potential may show less of a relationship between O₃ stomatal uptake and plant response.

4 The European approach and acceptance of flux-based critical values is a recognition of this
5 problem; and a concerted research effort is needed to develop the necessary experimental data
6 and modeling tools that will provide the scientific basis for such critical levels for O₃ (Grunhage
7 et al., 2004; Fuhrer et al., 1997; Grunhage and Jager, 1994).

8 At this time, based on the current state of knowledge, exposure indices that differentially
9 weight the higher hourly average O₃ concentrations, but include the mid-level values, represent
10 the best approach in the United States for relating vegetation effects to O₃ exposure.. A large
11 database exists that has been used for establishing exposure-response relationships. Such a
12 database does not yet exist for relating O₃ flux to growth response. Those pattern disconnects
13 between period of uptake and peak occurrence, as well as the potential for nocturnal uptake,
14 should be considered with some weighting functions in the currently used exposure indices.
15 Of particular consideration would be their inclusion in regional to national estimations of O₃
16 impacts on vegetation. Another useful approach to regional assessment for given specie(s) is
17 simulating growth effects with process-based models that account for seasonal climate and site
18 factors that control conductance

19 It is anticipated that, as the overlapping mathematical relationships of conductance,
20 concentration, and defense mechanisms are better defined, O₃ flux-based models may be able to
21 predict vegetation injury and/or damage at least for some categories of canopy-types with more
22 accuracy than the exposure-response models.

25 **9.6 OZONE EXPOSURE-PLANT RESPONSE RELATIONSHIPS**

26 **9.6.1 Introduction**

27 Ambient O₃ concentrations have long been known to cause visible symptoms, decreases in
28 photosynthetic rates, decreases in plant growth, and decreases in the yield of marketable organs
29 (U.S. Environmental Protection Agency, 1978, 1986, 1996). Yet, despite considerable research
30 in the U.S. and other countries during the past three decades, quantifying the effects of ambient
31 O₃ exposure on vegetation remains a challenge. Numerous studies have related O₃ exposure to

1 plant responses, with most effort focused on the yield of crops and the growth of tree seedlings.
2 Most experiments exposed individual plants grown in pots or soil under controlled conditions to
3 known concentrations of O₃ for a segment of daylight hours for some portion of the plant's life
4 span (Section 9.2). The response of a plant species or variety to O₃ exposure depends upon
5 many factors discussed in previous sections, including genetic characteristics (Section 9.4.2),
6 biochemical and physiological status (Section 9.4), and previous and current exposure to other
7 stressors (Sections 9.4, 9.5). Section 9.4 describes how O₃ moves from the atmosphere into the
8 leaf and the subsequent biochemical and physiological responses of plants. The current section
9 focuses on the quantitative responses of plants to seasonal or multiyear exposures to known
10 amounts of O₃. Quantitative responses include foliar symptoms and decreased growth of whole
11 plants or decreased harvestable portions of them. Because of the available information, most of
12 this section focuses on the response of individual plants, especially crop plants and tree
13 seedlings, with limited discussion of mixtures of herbaceous species. The responses of natural
14 ecosystems are discussed in Section 9.7.

15 This section will pay particular attention to studies conducted since the publication of the
16 1996 AQCD (U.S. Environmental Protection Agency, 1996). However, because much O₃
17 research was conducted prior to the 1996 AQCD, the present discussion of vegetation response
18 to O₃ exposure is largely based on the conclusions of the 1978, 1986, and 1996 criteria
19 documents (U.S. Environmental Protection Agency, 1978, 1986, 1996). To provide a context for
20 the discussion of recent research, the the key findings and conclusions of those three documents
21 are first summarized below.

22

23 **9.6.2 Summary of Key Findings/Conclusions from Previous Criteria** 24 **Documents**

25 Experimental data reviewed in the 1978 and 1986 criteria documents dealt primarily with
26 the effects of O₃ on agricultural crop species (U.S. Environmental Protection Agency, 1978,
27 1986). The chapter on vegetation effects in the 1978 document (U.S. Environmental Protection
28 Agency, 1978) emphasized foliar symptoms and growth effects, but not those effects that
29 affected yield, an emphasis dictated by the kind of data available at the time. The 1986
30 document reviewed a substantial new body of evidence based on open-top chamber (OTC)
31 experiments (see Section 9.2) showing that ambient O₃ exposures reduced the growth and yield

1 of herbaceous plants, again with a focus on major crop species. In the 1986 and 1996
2 documents, data were presented from regression studies conducted to develop exposure-response
3 functions for estimating yield loss of major crop species in different regions of the United States.
4 The 1996 document included results from additional herbaceous crop species as well as shrub
5 and tree species. For a number of tree species, biomass growth of seedlings was related to
6 growing season O₃ exposures to produce response functions for estimating O₃ exposures that
7 reduce growth by 10 or 30%. Also, in the 1986 and 1996 documents, data from studies using
8 ethylene diurea (EDU) as a protectant were reviewed. The 1978, 1986, and 1996 criteria
9 documents also reviewed data on the response to O₃ exposures of forest ecosystems in the
10 San Bernardino Mountains of southern California (U.S. Environmental Protection Agency, 1978,
11 1986, 1996). Because this region is exposed to high concentrations of O₃ and has shown
12 evidence of ecosystem-level changes, it remains an important study area (see Section 9.7).

13 Ozone can cause a range of effects, beginning with individual cells, leaves, and plants, and
14 proceeding to plant populations and communities. These effects may be classified as either
15 injury or damage. Injury encompasses all plant reactions, such as reversible changes in plant
16 metabolism (e.g., altered photosynthetic rate), altered plant quality, or reduced growth that does
17 not impair yield or the intended use or value of the plant (Guderian, 1977). In contrast, damage
18 includes all effects that reduce or impair the intended use or value of the plant. Damage includes
19 reductions in aesthetic values as well as losses in terms of weight, number, or size of the plant
20 part that is harvested (yield loss). Yield loss also may include changes in crop quality, i.e.,
21 physical appearance, chemical composition, or the ability to withstand storage. Losses in
22 aesthetic values are difficult to quantify. Although foliar symptoms cannot always be classified
23 as damage, their occurrence indicates that phytotoxic concentrations of O₃ are present, and,
24 therefore, studies should be conducted to assess the risk to vegetation.

25 Visible symptoms due to O₃ exposures reduce the market value of certain crops and
26 ornamentals for which leaves are the product, e.g., spinach, petunia, geranium, and poinsettia.
27 The concept of limiting values used to summarize foliar symptoms in the 1978 document (U.S.
28 Environmental Protection Agency, 1978) was also considered valid in the 1986 document (U.S.
29 Environmental Protection Agency, 1986). Jacobson (1977) developed limiting values by
30 assessing the available scientific literature and identifying the lowest exposure
31 concentration/duration reported to cause foliar symptoms in a variety of plant species.

1 A graphical analysis presented in those documents indicated that the limit for reduced plant
2 performance was an exposure to 50 ppb for several hours per day for more than 16 days.
3 Decreasing the exposure period to 10 days increased the concentration required to cause
4 symptoms to 100 ppb; and a short, 6-day exposure further increased the concentration to cause
5 symptoms to 300 ppb. These limiting values established in 1978 were still deemed appropriate
6 in the 1986 and 1996 criteria documents. Such foliar symptoms are caused by O₃ concentrations
7 that occur in the United States as shown in Table 9-13 (as adapted from U.S. Environmental
8 Protection Agency, 1996).

9 The 1986 document emphasized that, although foliar symptoms on vegetation are often an
10 early and obvious manifestation of O₃ exposure, O₃ effects are not limited to foliar symptoms.
11 Other effects include reduced growth of many organs (including roots), changes in crop quality,
12 and alterations in plant susceptibility to biotic stressors and sensitivity to abiotic stressors. The
13 1986 document also emphasized that O₃ exerts phytotoxic effects only if a sufficient amount of
14 O₃ reaches sensitive sites within the leaf (Section 9.3). Ozone injury will not occur if the rate of
15 O₃ uptake is low enough that the plant can detoxify or metabolize O₃ or its metabolites or if the
16 plant is able to repair or compensate for the effects (Tingey and Taylor, Jr., 1982; U.S.
17 Environmental Protection Agency, 1986). Cellular disturbances that are not repaired or
18 compensated for are ultimately expressed as foliar symptoms, reduced root growth, or reduced
19 yield of fruits or seeds.

20 Beginning in the 1986 document and continuing in the 1996 document, OTC studies were
21 reviewed that better quantified the relationship between O₃ exposure and effects on crop species,
22 with a focus on yield loss. These studies can be grouped into two types, depending on the
23 experimental design and statistical methods used to analyze the data: (1) studies that developed
24 predictive equations relating O₃ exposure to plant response, and (2) studies that compared the
25 effects of discrete treatment level(s) to a control. The advantage of the regression approach is
26 that exposure-response models can be used to interpolate results between treatment levels.

27 Discrete treatment experiments were designed to test whether specific O₃ treatments were
28 different from the control rather than to develop exposure-response equations, and the data were
29 analyzed using analyses of variance. When summarizing these studies using discrete treatment
30 levels, the lowest O₃ concentration that significantly reduced yield was determined from
31 analyses done by the original authors. Often, the lowest concentration used in the study was the

Table 9-13. Summary of Ozone Exposure Indices Calculated for 3- or 5-Month Growing Seasons from 1982 to 1991^a

<i>3-month growing season (June-August)</i>											
Year	No. of Sites ^b	HDM2 ^c ppm		M7 ppm		SUM00 ppm·h		SUM06 ppm·h		SIGMOID ppm·h	
		Mean	CV ^d	Mean	CV	Mean	CV	Mean	CV	Mean	CV
1982	99	0.114	23.7%	0	18.7%	82.9	19.1%	26.8	68.8%	26.3	56.7%
1983	102	0.125	24.9%	0	21.9%	86.1	22.1%	34.5	58.1%	33.0	52.3%
1984	104	0.117	24.6%	0	18.2%	84.1	19.9%	27.7	58.4%	27.4	47.9%
1985	117	0.117	24.6%	0	17.1%	84.6	18.0%	27.4	59.6%	27.4	47.6%
1986	123	0.115	21.8%	0	19.1%	85.3	18.0%	27.7	65.0%	27.7	51.8%
1987	121	0.119	22.9%	0	17.6%	86.9	17.3%	31.2	56.4%	30.4	46.8%
1988	139	0.129	21.3%	0	17.8%	97.6	19.6%	45.2	46.8%	42.9	42.4%
1989	171	0.105	23.1%	0	17.5%	86.4	19.9%	24.8	78.7%	25.8	59.4%
1990	188	0.105	21.6%	0	18.3%	85.7	21.0%	25.8	76.2%	26.6	59.2%
1991	199	0.106	22.0%	0	18.4%	87.7	21.3%	28.3	74.2%	28.9	59.5%
<i>Among Years</i>		0.113	11.1%	0	10.0%	87.0	9.9%	29.5	42.1%	29.4	31.0%
<i>5-month growing season (May-September)</i>											
Year	No. of Sites	M7 ppm		SUM00 ppm·h		SUM06 ppm·h		SIGMOID ppm·h			
		Mean	CV	Mean	CV	Mean	CV	Mean	CV		
1982	88	0.048	20.6%	122.9	22.3%	37.3	70.9%	37.1	57.8%		
1983	87	0.051	22.1%	129.6	24.4%	44.4	61.9%	43.8	52.7%		
1984	95	0.048	18.0%	126.2	19.1%	36.7	60.8%	37.6	46.9%		
1985	114	0.048	18.4%	124.5	19.4%	36.2	63.8%	37.0	50.3%		
1986	118	0.048	20.3%	123.3	21.4%	34.9	70.7%	35.6	55.7%		
1987	116	0.050	20.3%	128.7	20.4%	42.2	62.0%	41.8	50.3%		
1988	134	0.054	18.7%	141.7	22.0%	58.0	50.5%	55.6	45.0%		
1989	158	0.047	18.6%	127.8	22.5%	32.7	87.8%	35.2	64.1%		
1990	172	0.049	19.8%	129.4	22.7%	34.6	82.7%	37.0	62.1%		
1991	190	0.050	19.8%	130.6	23.6%	36.8	80.7%	38.8	62.9%		
<i>Among Years</i>		0.049	9.8%	129.0	9.9%	38.7	42.5%	39.6	29.8%		

^a Updated and additional years from data given in Table III of Tingey et al. (1991), where the spatial and temporal variation in ambient O₃ exposures is expressed in terms of several exposure indices.

^b Indicates the number of separate monitoring sites included in the analysis; fewer sites had 5 months of available data than had 3 months of available data.

^c The 2HDM index is calculated for sites with at least 3 months of available data. SUM00, SUM06, M7, SIGMOID, and 2HDM are the cumulative sum above 0.0 ppm, the cumulative sum above 0.06 ppm, the 7-h seasonal mean, the sigmoid weighted summed concentration, and the second highest daily maximum 1-h concentration, respectively.

^d CV = coefficient of variation.

Source: Table 5-30 from U.S. Environmental Protection Agency (1996) based on Tingey et al. (1991).

lowest concentration reported to reduce yield; hence, it was not always possible to estimate a no-effect exposure concentration. In general, the data indicated that 100 ppb O₃ (frequently the lowest concentration used in the studies) for a few hours per day for several days to several weeks usually caused significant yield reductions of 10 to 50%.

By the time the 1986 document was prepared, much new information concerning the effects of O₃ on the yield of crop plants had become available through EPA's NCLAN research program and through research funded by other agencies. The NCLAN project was initiated by EPA in 1980 primarily to improve estimates of yield loss under field conditions and to estimate the magnitude of crop losses caused by O₃ throughout the United States (Heck et al., 1982, 1991). The cultural conditions used in the NCLAN studies approximated typical agronomic practices. The primary objectives were:

- (1) to define relationships between yields of major agricultural crops and O₃ exposure as required to provide data necessary for economic assessments and development of O₃ NAAQS;
- (2) to assess the national economic consequences resulting from O₃ exposure of major agricultural crops; and
- (3) to advance understanding of cause-and-effect relationships that determine crop responses to pollutant exposures.

Using NCLAN data, the O₃ concentrations predicted to cause 10 or 30% yield loss were estimated using linear or Weibull response functions. The data in Table 9-14 are from the 1996 document and were based on yield-response functions for 38 species or cultivars developed from studies using OTCs of the type developed by Heagle et al. (1973) (see Section 9.2). Composite exposure-response functions for both crops and tree seedlings as a function of O₃ exposure expressed as SUM06 are shown in Figure 9-18. Review of these data indicate that 10% yield reductions could be predicted for more than 50% of experimental cases when: (1) 12-h SUM06 values exceeded 24.4 ppm·h, (2) SIGMOID values exceeded 21.5 ppm·h, or (3) 7-h seasonal mean concentrations were 50 ppb. The SIGMOID index is very similar to the W126 index (see Section 9.5 for further information about O₃ indices). Much lower values are required for each index to protect 75% of experimental cases (Table 9-15). Grain crops were generally found to be less sensitive than other crops. The data summarized in the 1996 criteria document also indicated that the variation in sensitivity within species may be as great as differences between species.

Table 9-14. Ozone Exposure Levels (Using Various Indices) Estimated To Cause at Least 10% Crop Loss in 50 and 75% of Experimental Cases^a

50th PERCENTILE^b	SUM06	SE^c	SIGMOID	SE	M7	SE	2HDM	SE
NCLAN Data (n = 49; wet and dry) ^d	24.4	3.4	21.5	2.0	0.049	0.003	0.094	0.006
NCLAN Data (n = 39; wet only)	22.3	1.0	19.4	2.3	0.046	0.003	0.090	0.010
NCLAN Data (n = 54; wet and dry) ^e	26.4	3.2	23.5	2.4	NA	NA	0.099	0.011
NCLAN Data (n = 42; wet only) ^e	23.4	3.1	22.9	4.7	NA	NA	0.089	0.008
NCLAN Data (n = 10; wet)	25.9	4.5	23.4	3.2	0.041	0.001	0.110	0.042
NCLAN Data (n = 10; dry)	45.7	23.3	40.6	0.1	0.059	0.014	0.119	0.017
Cotton Data (n = 5)	23.6	2.3	19.3	2.3	0.041	0.001	0.066	0.032
Soybean Data (n = 13)	26.2	5.4	22.6	3.6	0.044	0.005	0.085	0.013
Wheat Data (n = 6)	21.3	15.2	19.3	12.7	0.061	0.018	0.098	0.059
Cotton Data (n = 5) ^e	30.0	12.7	27.2	12.8	NA	NA	0.075	0.012
Soybean Data (n = 15) ^e	23.9	6.5	22.0	8.0	NA	NA	0.088	0.008
Wheat Data (n = 7) ^e	25.9	10.5	21.4	9.4	NA	NA	0.097	0.028
75th PERCENTILE^b								
NCLAN Data (n = 49; wet and dry)	14.2	4.2	11.9	5.6	0.040	0.007	0.051	0.010
NCLAN Data (n = 39; wet only)	14.3	2.7	12.6	2.3	0.039	0.005	0.056	0.006
NCLAN Data (n = 54; wet and dry) ^e	16.5	4.3	14.5	3.2	NA	NA	0.073	0.006
NCLAN Data (n = 42; wet only) ^e	17.2	3.0	14.7	2.4	NA	NA	0.070	0.006
NCLAN Data (n = 10; wet)	16.4	3.7	13.7	3.2	0.040	0.001	0.080	0.032
NCLAN Data (n = 10; dry)	24.0	0.8	22.3	0.1	0.053	0.022	0.093	0.003
Cotton Data (n = 5)	21.8	5.0	17.5	2.8	0.041	0.001	0.065	0.014
Soybean Data (n = 13)	14.2	0.1	12.4	0.1	0.041	0.006	0.069	0.004
Wheat Data (n = 6)	11.7	2.5	10.9	2.4	0.054	0.032	0.062	0.035
Cotton Data (n = 5) ^e	21.1	6.0	16.7	5.7	NA	NA	0.070	0.034
Soybean Data (n = 15) ^e	15.3	4.1	13.4	4.1	NA	NA	0.078	0.007
Wheat Data (n = 7) ^e	5.1	2.6	8.5	3.4	NA	NA	0.054	0.027

^aSee Appendix A for abbreviations and acronyms.

^bThe numbers in parentheses are the number of cases used in deriving the various exposure levels.

^cStandard error (SE).

^dNCLAN data refers to studies conducted as part of the NCLAN project. Wet and dry refer to watering regimes used in the studies, wet being well-watered, and dry meaning some level of drought stress was imposed.

^e24-h exposure statistics reported in Lee et al. (1994b). Relative yield loss for 2HDM is relative to yield at 40 ppb rather than 0 ppb as was used in Tingey et al. (1991).

Source: U.S. Environmental Protection Agency (1996), (modified from Tingey et al. (1991).




Figure 9-18. Distribution of biomass loss predictions from Weibull and linear exposure-response models that relate biomass to O₃ exposure. Exposure is characterized with the 24-h SUM06 statistic using data from (a) 31 crop studies from National Crop Loss Assessment Network (NCLAN) and (b) 26 tree seedling studies conducted at U.S. Environmental Protection Agency's Environmental Research Laboratory in Corvallis, OR; Smoky Mountains National Park, TN; Houghton, Michigan; and Delaware, Ohio. Separate regressions were calculated for studies with multiple harvests or cultivars, resulting in a total of 54 individual equations from the 31 NCLAN studies and 56 equations from the 26 seedling studies. Each equation was used to calculate the predicted relative yield or biomass loss at 10, 20, 30, 40, 50, and 60 ppm·h, and the distributions of the resulting loss were plotted. The solid line is the calculated Weibull fit at the 50th percentile.

Source: U.S. Environmental Protection Agency (1996); Hogsett et al. (1995).

1 The chemical protectant, ethylene diurea (EDU), was also used to provide estimates of
2 yield loss. The impact of O₃ on yield was determined by comparing the yield data from plots
3 treated with EDU versus untreated plots. Studies indicated that yields were reduced by 18 to
4 41% when daytime ambient O₃ concentrations exceeded 80 ppb for 5 to 18 days over the
5 growing season. For this approach to be credible, the effects of EDU itself on a particular
6 species must be preestablished under conditions without O₃ exposure (Kostka-Rick and
7 Manning, 1992).

8 The 1996 criteria document reviewed several experiments demonstrating that the seedlings
9 of some tree species such as poplars (*Populus*) and black cherry are as sensitive to O₃ as are
10 annual plants, in spite of the fact that trees are longer-lived and generally have lower rates of gas
11 exchange, and, therefore, a lower uptake of O₃. The 1996 document also reviewed data showing
12 that O₃ exposures that occur at present in the United States are sufficient to affect the growth of a
13 number of trees species. For example, exposure-response functions for 51 cases of tree seedling
14 responses to O₃, including 11 species representing deciduous and evergreen growth habits,
15 suggest that a SUM06 exposure for 5 months of 31.5 ppm·h would protect hardwoods from a
16 10% growth loss in 50% of the cases studied (Table 9-15). Similarly, a SUM06 exposure of
17 42.6 ppm·h should provide the same level of protection for evergreen seedlings. However, these
18 results do not take into the account the possibility of effects on growth in subsequent years.
19 Because multiple-year exposures may cause a cumulative effect on the growth of some trees
20 (Simini et al., 1992; Temple et al., 1992), it is likely that a number of species are currently being
21 affected even at ambient exposures (Table 9-14).

22 In 1986, the EPA (U.S. Environmental Protection Agency, 1986) established that 7-h per
23 day growing season mean exposures to O₃ concentrations above 50 ppb were likely to cause
24 measurable yield loss in agricultural crops. At that time, few conclusions could be drawn about
25 the response of deciduous or evergreen trees or shrubs, due to the lack of information about
26 response of such plants to season-long exposures to O₃ concentrations of 40 to 60 ppb and above.
27 However, the 1978 and 1986 criteria documents (U.S. Environmental Protection Agency, 1978,
28 1986) indicated that the limiting value for foliar symptoms on trees and shrubs was 60 to 100
29 ppb for 4 h. From 1986 to 1996, extensive research was conducted, establishing the sensitivity
30 of many tree species. Based on research published since the 1986 criteria document,, a number
31 of conclusions were drawn in 1996 AQCD (U.S. Environmental Protection Agency, 1996):

**Table 9-15. SUM06 Levels Associated with 10 and 20% Total Biomass Loss for
50 and 75% of the Seedling Studies
(The SUM06 value is adjusted to an exposure length of 92 days per year.)^a**

Weibull Equations (all 51 seedling studies):

$$50\text{th Percentile PRYL}^1 = 1 - \exp(-[\text{SUM06}/176.342]**1.34962)$$

$$75\text{th Percentile PRYL} = 1 - \exp(-[\text{SUM06}/104.281]**1.46719)$$

Weibull Equations (27 fast-growing seedling studies):

$$50\text{th Percentile PRYL} = 1 - \exp(-[\text{SUM06}/150.636]**1.43220)$$

$$75\text{th Percentile PRYL} = 1 - \exp(-[\text{SUM06}/89.983]**1.49261)$$

Weibull Equations (24 slow to moderate growing seedling studies):

$$50\text{th Percentile PRYL} = 1 - \exp(-[\text{SUM06}/190.900]**1.49986)$$

$$75\text{th Percentile PRYL} = 1 - \exp(-[\text{SUM06}/172.443]**1.14634)$$

Weibull Equations (28 deciduous seedling studies):

$$50\text{th Percentile PRYL} = 1 - \exp(-[\text{SUM06}/142.709]**1.48845)$$

$$75\text{th Percentile PRYL} = 1 - \exp(-[\text{SUM06}/87.724]**1.53324)$$

Weibull Equations (23 evergreen seedling studies):

$$50\text{th Percentile PRYL} = 1 - \exp(-[\text{SUM06}/262.911]**1.23673)$$

$$75\text{th Percentile PRYL} = 1 - \exp(-[\text{SUM06}/201.372]**1.01470)$$

**Levels Associated with Prevention of a 10 and 20% Total Biomass Loss
for 50 and 75% of the Seedlings**

All 51 Seedling Cases

		Percent of Seedlings	
		<i>50%</i>	<i>75%</i>
Relative	<i>10%</i>	33.3	22.5
Biomass Loss	<i>20%</i>	58.0	37.5

27 Fast-Growing Seedling Cases

		Percent of Seedlings	
		<i>50%</i>	<i>75%</i>
Relative	<i>10%</i>	31.3	19.4
Biomass Loss	<i>20%</i>	52.9	32.4

Table 9-15 (cont'd). SUM06 Levels Associated with 10 and 20% Total Biomass Loss for 50 and 75% of the Seedling Studies
(The SUM06 value is adjusted to an exposure length of 92 days per year.)^a

24 Slow-to-Moderate-Growth Seedling Cases

		Percent of Seedlings	
		50%	75%
Relative	10%	42.6	24.2
Biomass Loss	20%	70.2	46.6

28 Deciduous Seedling Cases

		Percent of Seedlings	
		50%	75%
Relative	10%	31.5	20.2
Biomass Loss	20%	52.1	33

23 Evergreen Seedling Cases

		Percent of Seedlings	
		50%	75%
Relative	10%	42.6	21.9
Biomass Loss	20%	78.2	45.9

^aSee Appendix [XXX] for abbreviations and acronyms.

^bPRYL = predicted relative yield (biomass) loss

Source: U.S. Environmental Protection Agency (1996), based on Hogsett et al. (1995).

- 1 (1) An analysis of 10 years of monitoring data from more than 80 to almost 200 non-urban sites in the United States established ambient 7-h growing season average concentrations of O₃ for 3 or 5 months of 51 to 60 ppb and 47 to 54 ppb, respectively. The SUM06 exposures ranged (a) from 24.8 to 45.2 ppm·h for 3 months and (b) from 32.7 to 58.0 ppm·h for 5 months (Tingey et al. (1991), Table 9-14).
- 2 (2) The results of OTC studies that compared yields at ambient O₃ exposures with those in filtered air and retrospective analyses of crop data (Table 9-14) established that ambient O₃ concentrations were sufficient to reduce the yield of major crops in the United States. Research results since 1978 did not invalidate EPA conclusions (U.S. Environmental Protection Agency, 1978, 1986) that foliar symptoms due to

O₃ exposures reduce the market value of certain crops and ornamentals where leaves are the product (such as spinach, petunia, geranium, and poinsettia) and that such damage occurs at ambient O₃ concentrations observed in the United States.

- (3) A 3-month SUM06 exposure of 24.4 ppm·h, corresponding to a 7-h mean of 49 ppb and a 2HDM of 94 ppb O₃ may prevent a 10% loss in 50% of the 49 experimental cases analyzed by Tingey et al. (1991). A 12-h growing season mean of 0.045 ppb should restrict yield losses to 10% in major crop species (Lesser et al., 1990).
- (4) Depending on duration, concentrations of O₃ and SUM06 exposures currently in the United States are sufficient to affect the growth of a number of tree species. Given the fact that multiple-year exposures may cause a cumulative effect on the growth of some trees (Simini et al., 1992; Temple et al., 1992), it is likely that a number of species currently are being impacted, even at ambient O₃ exposures (Table 9-14).
- (5) Exposure-response functions for 51 cases of seedling response to O₃ (Hogsett et al., 1995), including 11 species representing deciduous and evergreen growth habits, suggest that a SUM06 exposure for 5 months of 31.5 ppm·h would protect hardwoods from a 10% growth loss in 50% of the cases studied. A SUM06 exposure of 42.6 ppm·h should provide the same level of protection for evergreen seedlings. Note that these conclusions do not take into the account the possibility of effects on growth in subsequent years, an important consideration in the case of long-lived species.
- (6) Studies of the response of trees to O₃ have established that, in some cases (for instance, poplars and black cherry), trees are as sensitive to O₃ as are annual plants, in spite of the fact that trees are longer-lived and generally have lower gas exchange rates, and, therefore, lower O₃ uptake.
- (7) Use of the chemical protectant, EDU, is of value in estimating O₃-related losses in crop yield and tree growth, provided that care is exercised in establishing appropriate EDU dosages to protect the plants without affecting growth.

The major question to be addressed in the remainder of this section is whether new information supports or alters the 1996 criteria document conclusions summarized above. In particular, this section evaluates whether the response of plants to experimental treatments at or near O₃ concentrations characteristic of ambient levels in many areas of the United States (Table 9-14) can be compared to a control or reduced O₃ treatment to establish a potential adverse effect. Before evaluating new information from the literature on O₃ effects on vegetation, O₃ exposure indices used in O₃ studies and trends in O₃ exposure patterns during the past two decades are briefly reviewed.

9.6.3 Ozone Indices and Ambient Exposure

As recognized in both the 1986 and the 1996 criteria documents, the characterization and representation of the exposure of vegetation to O₃ is problematic, because the specific aspects of pollutant exposure that cause injury or damage are difficult to quantify. This issue is addressed in Section 9.5, and only a few points will be discussed here in order to provide a context for interpreting data on exposure-response relationships. The most important effects of O₃ on vegetation occur due to uptake of O₃ through stomata, with subsequent oxidative injury that appears to be rather nonspecific (Section 9.3). As has been discussed by numerous authors during the last three decades, from a toxicological and physiological view, it is much more realistic to relate effects to internal (absorbed) O₃ dose rather than to exposure near the leaf or canopy (Fowler and Cape, 1982; Fuhrer et al., 1992; Grünhage et al., 1993, 1999; Legge et al., 1995; Massman et al., 2000; Musselman and Massman, 1999; Pleijel et al., 1995; Runeckles, 1974; Taylor, Jr. et al., 1982; Tingey and Taylor, Jr., 1982) (see also Section 9.5). Theoretically, flux estimates should improve the assessment of O₃ effects, but despite recent attention to this topic, particularly in Europe, it remains difficult to estimate flux in the field outside of experimental sites where continuous measurements of wind speed and other environmental conditions are made. This topic is discussed further below in Section 9.6.4.5.

No simple exposure index can accurately represent all of the numerous factors operating at different timescales that affect O₃ flux into plants and subsequent plant response (Section 9.5). Indices of peaks, such as the 2HDM are not well suited for discerning exposure-response relationships, because they do not capture the effects of lower O₃ concentrations nor the cumulative effects of O₃ on vegetation (Heck and Cowling, 1997; U.S. Environmental Protection Agency, 1996). For this reason, peak indices have not been used in recent decades to develop exposure-response relationships for vegetation. Fortunately, other simple indices have shown substantial correlation with responses such as crop yield under experimental conditions. During the 1980s, the most commonly used indices for expressing O₃ exposure were 7-, 8-, or 12-h daytime average values over the duration of O₃ exposure, which was often 3 months or somewhat less for experimental studies with crops. These indices perform reasonably well for interpreting experimental data on the response of vegetation to ozone, particularly for individual experiments, although they do not explain all of the variation among experiments in retrospective analysis of multiple experiments (Lesser et al., 1990).

1 Since the 1980s, cumulative indices such as the SUM06, AOT40, or W126 that
2 preferentially weight higher concentrations have been used in conjunction with mean indices for
3 developing exposure-response relationships (Tables 9-15 and 9-16, and Figure 9-1). Such
4 indices are often more suitable than mean values because they are cumulative and because they
5 preferentially weight higher concentrations. Thus, these indices generally provide somewhat
6 better fits to experimental data than do mean indices, especially in retrospective analyses of
7 multiple experiments on multiple species (Lee et al., 1994a, 1994b; Lee and Hogsett, 1999;
8 Tingey et al., 1991). Unfortunately, no single index has been used consistently even in the
9 recent literature, making it difficult to compare results among and between experiments and with
10 ambient exposure data. However, Tables 9-14 and 9-17 provide summaries of ambient exposure
11 data for several indices that can be compared to the experimental results reviewed in the
12 remainder of this section. Of the cumulative indices that preferentially weight higher
13 concentrations, the SUM06 index has been used most commonly in the U.S. literature, and it was
14 selected in a meeting of scientific experts on O₃ effects on vegetation as suitable for a secondary
15 standard to protect vegetation (Heck and Cowling, 1997). However, it should be noted that the
16 W126 index has been selected for use in protecting vegetation in Class 1 areas (Federal Land
17 Managers' Air Quality Related Values Workgroup (FLAG), 2000). Even in recent studies, O₃
18 data are often presented using only a seasonal mean index value, and so mean values are
19 frequently presented in this section. Such reporting of mean indices should not be interpreted as
20 a preference for them, but rather as a limitation in the data reported in the literature. Additional
21 information about O₃ exposure for individual experiments, including the number and type of O₃
22 treatments (addition of a constant concentration of O₃ or an amount proportional to ambient
23 levels), and duration, are reported in Tables 9-16 through 9-19.

24 Since the 1996 document, the use of the AOT40 index has become quite common in
25 Europe for identifying and mapping areas of exceedance, but it has not been used much in the
26 United States. Thus, studies reporting O₃ exposure only as AOT40 values are presented in tables
27 summarizing effects on annual, herbaceous perennial, and woody vegetation. However, such
28 studies are not as commonly cited in the text of this section because AOT40 summary data on O₃
29 exposures in the United States are rarely available. This lack makes it difficult to compare
30 experimentally derived exposure-response data expressed as AOT40 to ambient U.S. O₃

Table 9-16. Summary of Selected Studies of Ozone Effects on Annual Species

Species	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (Decrease from lowest, %)	Reference
Bean, cv. Pros	OTC	The Netherlands	CF to CF75: 9-h mean = 3-70, AOT40 = 0 to 17.7 ppm•h	62 days	g=Green pod yield	29 at 9-h mean = 44 (AOT40 = 3.6 ppm•h)	Tonneijck and Van Dijk (1998)
Bean, cv. Lit	OTC	Germany	CF, NF, CF-1×, CF-2×: mean = 1, 14, 15 , 32	3 months	Pod yield	56 (CF, 2×)	Brunschon-Harti et al. (1995)
Corn	OTC	Beltsville, MD	CF, +40: 7-h mean = 20, 70	1 years	Grain yield	13	Mulchi et al. (1995) Rudorff et al. (1996c)
Cotton, cv. Deltapine	OTC	Raleigh, NC	CF, 1.5×: 12-h mean = 21, 71	1 years	Seed-cotton weight	22	Heagle et al. (1999)
Cotton, cv. Deltapine	OTC	Raleigh, NC	CF, NF, 1.5×, 12-h mean = 24, 51 , 78	1 years	Seed-cotton weight	21, 49 (NF, 1.5×)	Heagle et al. (1999)
Oat, cv. Vital	OTC	Ostad, Sweden	CF, NF: 7-h mean = 12, 27	1 years	Grain yield	+2 (n.s.)	Pleijel et al. (1994a)
Potato²	OTC	6 sites N. Europe	AOT40 = 6-27 ppm•h	2 years (1 year at 2 sites)	Tuber yield	4% average for all experiments	Craigon et al. (2002)
Rape, oilseed	Open Air	Northumberland, UK	AA, +O ₃ : 7-h mean for 17 days Aug.-Sept = 30 , 77, for 32 days in May-June = 31 , 80	17 days in fall, overwinter, 32 days in spring	Seed yield	14	Ollerenshaw et al. (1999)
Rice, cvs. Koshi-hikari, Nippon-bare	OTC	Japan	CF, 1×, 1.5×, 2×, 2.75×: 7-h mean = 13.5-93.4	3 years	Grain yield	3 to 10 at 40 ppb	Kobayashi et al. (1995)
Soybean	OTC	Beltsville, MD	CF, +40: 7-h mean = 25, 72	2 years	Seed yield	25	Mulchi et al. (1995)
Soybean, cv. Essex	OTC	Raleigh, NC	CF, 1.5×: 12-h mean for 3 years = 23, 82	3 years	Seed yield	41	Fiscus et al. (1997)

Table 9-16 (cont'd). Summary of Selected Studies of Ozone Effects on Annual Species

Species	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (Decrease from lowest, %)	Reference
Soybean, cvs. Forrest, Essex	OTC	Maryland	CF, +40: 7-h mean = 24 and 24, 63 and 62 for each year	2 years	Seed yield	10, 32 (2 cvs.)	Chernikova et al. (2000)
Soybean, cv. Essex	OTC	Raleigh, NC	CF, NF, 1.5×: 12-h mean = 20, 50 , 79	1 year	Seed yield	16, 37 (NF, 1.5×)	Heagle et al. (1998)
Soybean, cv. Essex	OTC	Raleigh, NC	CF, NF, 1.5×: 12-h mean = 18, 42 , 69	1 year	Seed yield	15, 40 (NF, 1.5×)	Heagle et al. (1998)
Soybean, cv. Holladay	OTC	Raleigh, NC	CF, NF, 1.5×: 12-h mean = 18, 42 , 69	1 year	Seed yield	22, 36 (NF, 1.5×)	Heagle et al. (1998)
Soybean, cv. NK-6955	OTC	Raleigh, NC	CF, NF, 1.5×: 12-h mean = 18, 42 , 69	1 year	Seed yield	+46, +4 (NF, 1.5×)	Heagle et al. (1998)
Soybean, 3 cvs.	OTC	Raleigh, NC	CF, NF, 1.5×: 12-h mean = 14, 36 , 64	3 months	Seed yield	At ambient = +14, 11, 16 for 3 cvs.	Miller et al. (1994)
Soybean, 3 cvs.	OTC	Raleigh, NC	CF, NF, 1.5×: 12-h mean = 24, 49 , 83	4 months	Seed yield	At ambient = 17, 13, 18 (3 cvs.)	Miller et al. (1994)
Soybean, cv. Essex	OTC	Raleigh, NC	CF, NF, 1.5×: 12-h mean = 20, 50 , 79	4 months	Seed yield	11, 22 (amb., 1.5×)	Miller et al. (1998)
Soybean, cvs, Essex, Forrest	OTC	Beltsville, MD	CF, NF+: 7-h mean = 24, 58	134 days	Seed yield	Essex = +11 (ns), Forrest = 21	Robinson and Britz (2000)
Timothy	OTC	Sweden	AOT40 = 10, 20, 340; 12-h mean = 20, 152	1 year	Biomass	58	Danielsson et al. (1999)
Watermelon	OTC	Spain	CF (O ₃ = 0), NF in 1988 AOT40 = 5.96 ppm•h, SUM06 = 0.29 ppm•h, in 1989 AOT40 = 18.92 ppm•h, SUM06 = 4.95 ppm•h	2 expts of 1 year	Fruit yield	19, 39 (2 expts)	Gimeno et al. (1999)

Table 9-16 (cont'd). Summary of Selected Studies of Ozone Effects on Annual Species

Species	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (Decrease from lowest, %)	Reference
Wheat¹, cv. Minaret	OTC	8 sites in N Europe	12-h mean (SD) low = 26.3 (12.2), 12-h mean (SD)-high = 51.4 (18.3) AOT40 mean (SD) low = 6.18 (8.54) ppm•h, AOT40 mean (SD) high = 28.23 (23.05) ppm•h	13 studies of 1 year each	Grain yield	13 (n.s.)	Bender et al. (1999) Hertstein et al. (1999)
Wheat¹	OTC	Sweden	AOT40 0 to 15 ppm•h	7 years	Grain yield	23 at AOT40 = 15 ppm•h	Danielsson et al. (2003)
Wheat, cv. promessa	OTC	SE Ireland	CF, +50: 12-h total = 5.6, 32.6 ppm•h	3 h/day, 5 d/week, 7 weeks	Grain yield	53	Finnan et al. (1996a)
Wheat, cv. promessa	OTC	SE Ireland	CF, +25: 12-h total = 6.2, 33.4 ppm•h	6 h/day, 5 d/week, 7 weeks	Grain yield	+17	Finnan et al. (1996a)
Wheat, cv. promessa	OTC	SE Ireland	CF, +25, +50: 12-h total = 6.7, 34, 34 ppm•h	+25 = 6 h/day, 5 d/week, +50 = 3 h/day, 5 d/week, both 7 weeks	Grain yield	Amb + 25 = 3 (n.s.); Amb + 50 = 17	Finnan et al. (1996a)
Wheat, cvs. Massey, Saluda	OTC	Beltsville, MD	CF, +40: 7-h mean = 19, 20 and 61, 65 (2 years)	2 years	Grain yield	20	Mulchi et al. (1995); Rudorff et al. (1996b)
Wheat, cv. Turbo	OTC	Germany	8-h mean = 5.9, 61.2, 92.5	1 year	Grain yield	14, 40 (mid, high O ₃)	Bender et al. (1994)
Wheat, cv. Turbo	OTC	Germany	8-h mean = 4.7, 86.4	1 year	Grain yield	20	Bender et al. (1994)
Wheat, cv. Turbo	OTC	Germany	7-h mean = 5, 41, 73	1 year	Grain yield	35	Fangmeier et al. (1994)

Table 9-16 (cont'd). Summary of Selected Studies of Ozone Effects on Annual Species

Species	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (Decrease from lowest, %)	Reference
Wheat, winter, 8 cvs.	OTC	Raleigh, NC	12-h mean = 27, 47, 90	2 months	Grain yield	5 (n.s.)	Heagle et al. (2000)
Wheat, winter, 8 cvs	OTC	Raleigh, NC	12-h mean = 22, 38, 74	2 months	Grain yield	16 (n.s.)	Heagle et al. (2000)
Wheat, cv. Drabant	OTC	Finland	1992: 12-h mean = 14, 30, 61; AOT40 = 16.3, 34.8, 54.6 ppm•h. 1993: 12-h mean = 9, 21, 45; AOT40 = 10.2, 24.8, 40.6 ppm•h	2 years	Grain yield	At highest O ₃ = 13 each year	Ojanpera et al. (1998)
Wheat, cv. Riband	Open Air	Northumberland, UK	AOT40 for Mar to Aug 93 = 3.5, 6.2 ppm•h	1 year, overwinter	Grain yield	13	Ollerenshaw and Lyons (1999)

¹ Values for ambient or NF treatments are indicated in bold.
² **Bold** indicates that multiple experiments (more than just 2 years at a single site) were included in the analysis.

Table 9-17. Summary of Selected Studies of the Effects of Ozone on Perennial Herbaceous Plants

Species	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (Decrease from lowest, %)	Reference
Alfalfa, cvs. Apica, Team	OTC	Quebec, Canada	12-h mean: 1991 = 6, 39, 49, 110; 1992 = 0, 34, 42, 94	3 months in each of 2 years	Biomass	For NF: Apica = 31, 21; Team = 14, 2 (n.s.)	Renaud et al. (1997)
Bahia grass	OTC	Auburn, AL	12-h mean = 22, 45, 91	24 weeks	Biomass at ambient O ₃ for 1st, 2nd cutting of early and late season plantings	34, 29 (n.s.), +6 (n.s.), 9 (n.s.)	Muntifering et al. (2000)
Bent grass (Capillaris sp.)	OTC	United Kingdom	AOT40 = 0.8-15.0 ppm•h	8 h/day for 3 months	Biomass, in competition with 3 other spp. Total biomass in uncut pots, aboveground biomass in cut pots (cut every 14 days).	8 (uncut), +18 (cut)	Ashmore and Ainsworth (1995)
Blackberry	Large OTC	Alabama	1994: AOT40 = 2-112 ppm•h, SUM06 = 1-162 ppm•h, 1995: AOT40 = 3-83 ppm•h, SUM06 = 0-132 ppm•h	7 months in 1994, 6 months in 1995	Percent canopy cover (grown in old field community), biomass ripe fruit number	+124 for cover, n.s. for biomass, 28% for ripe fruit number but sig. chamber effect.	Barbo et al. (1998) Chappelka (2002)
Clover, white	Ambient air	MA, OR, NC, CA (2 sites) and VA	SUM06 for 6-h/day = 10.2-39.4 ppm•h, AOT40 for 12-h/day = 0.6-50.1 ppm•h	2 growing seasons	Biomass ratio (sensitive/resistant)	4 at 6-h SUMO6 = 39.4 ppm•h; 12 h AOT40 = 50.1 ppm•h	Heagle and Stefanski (2000)
Clover, white	Ambient air	14 European sites	AOT40 for 28 d = 0-12 ppm•h	3 growing seasons	Biomass ratio sensitive/resistant)	5 at AOT40 for 28 days = 0.9-1.7 ppm•h	Mills et al. (2000)
Clover, white	OTC	United Kingdom	AOT40 = 0.8-15.0 ppm•h	8 h/day for 3 months	Biomass, in competition 3 other spp. Total biomass in uncut pots, aboveground biomass in cut pots (cut every 14 days).	18 (uncut), 40 (cut)	Ashmore and Ainsworth, (1995)
Clover, white and red	OTC	Switzerland	CF, NF, NF+, NF++: 12-h mean = 21, 39 , 47, 65	3.5 months/ year for 2 years	Biomass, in managed pasture	24, 26, 52	Fuhrer et al. (1994)
Clover, white, cv. Menna	OTC	Italy	CF, NF: AOT40 = 0.1; 8.9 ppm•h, 7-h mean = 24, 53	2 months	Biomass	20	Fumagalli et al. (1997)

Table 9-17 (cont'd). Summary of Selected Studies of the Effects of Ozone on Perennial Herbaceous Plants

Species	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (Decrease from lowest, %)	Reference
Fescue, red	OTC	United Kingdom	AOT40 = 0.8-15.0 ppm•h	8 h/day for 3 months	Biomass, in competition with 3 other spp. Total biomass in uncut pots, aboveground biomass in cut pots (cut every 14 days).	+ 30 (uncut), +13 (cut)	Ashmore and Ainsworth, (1995)
Lespediza, Sericea	OTC	Auburn, AL	CF, NF, 2×: 12-h mean = 23, 40 , 83, SUM06 = 0.2, 9.1 , 61.0, AOT40 = 0.6, 7.0 , 39.8	10 weeks	Biomass	n.s.	Powell et al. (2003)
Little bluestem	OTC	Auburn, AL	CF, NF, 2×: 12-h mean = 23, 40 , 83 ppb, SUM06 = 0.2, 9.1 , 61.0, AOT40 = 0.6, 7.0 , 39.8	10 weeks	Biomass	n.s.	Powell et al. (2003)
Phleum alpinum	OTC	Sweden	AOT40 = 0.01, 0.02, 0.34 ppm•h; 12-h mean = 20, 152	1 year	biomass	87	(Danielsson et al. (1999)
Raspberry	OTC	Ontario	1×, +12 +24: Ambient ≤ 4	1 day/week for 7 weeks	Fruit yield	No effect at +12, 52 at +24 in only one of two cultivars	Sullivan et al. (1994)
Speedwell, Germander	OTC	United Kingdom	AOT40 = 0.8-15.0 ppm•h	8 h/day for 3 months	Biomass, in competition with 3 other spp. Total biomass in uncut pots, aboveground biomass in cut pots (cut every 14 days).	14 (uncut), 26 (cut)	Ashmore and Ainsworth (1995)
Strawberry	OTC	United Kingdom	8-h mean = 27, 92; AOT40 for +O ₃ = 24.59 ppm•h	69 days	Fruit size, yield	Size = 14, yield = (n.s.)	Drogoudi and Ashmore (2000)
Sumac, winged	OTC	Alabama	SUM06 = 0 to 132 ppm•h	6 months	Percent canopy cover (grown in old field community)	95	(Barbo et al. (1998)
Timothy	OTC	Sweden	CF, NF, CF+: AOT40 = 0.0, 1.3 , 20.3 ppm•h; 12-h mean = 20, 68 , 152	1 year	Biomass	ns in NF, 58 in CF+	Danielsson et al. (1999)

¹ Values for ambient or NF treatments are indicated in bold.² **Bold** indicates that multiple experiments (more than just 2 years at a single site) were included in the analysis.

Table 9-18. Summary of Selected Studies of Ozone Effects on Deciduous Trees and Shrubs

Species	Age	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (Decrease from lowest, %)	Reference
Ash, European		OTC	Hampshire, UK	NF, NF+: Mean = 17.7 , 44.1; AOT40 for 24h = 1.9 , 59.9 ppm•h	3 years for day 100 to day 162	Growth and biomass of organs	n.s.	Broadmeadow and Jackson (2000)
Ash, European	Seedling	OTC	Switzerland	0.5×, 0.85×, 1×, 0.5×+30: AOT40 = 0.1, 3.4, 7.1 , 19.7 ppm•h	5 months	Biomass	26 at 1×, 50 at 0.5× + 30	Landolt et al. (2000)
Aspen	Cutting, Seedling	OTC	Michigan	CF, 1×, 2×: 3 months 7-h mean for 1990 = 7-69; for 1991 = 22-92	98 days	Total biomass	ns at 1×, 29 at 2× for all clones in each year	Karnosky et al. (1996)
Aspen	Cutting	OTC	Michigan	CF, 1×, 2×: SUM00 = 11, 58 , 71 ppm•h	98 days	Total biomass	25-38 at 1×	Dickson et al. (2001)
Aspen	Cutting	FACE	Wisconsin	ambient, +90 (exposure data not reported)	3 years	Volume (d2*h)	21 at +90	Isebrands et al. (2001)
Aspen	Cutting	Large OTC	New York	1×, 1.7×, 3×: SUM06 = 1 , 20, 62 ppm•h; 9-h mean = 40 , 74, 124	92 days	Shoot biomass	14, 25 for 2 clones at 1.7×	Yun and Laurence (1999a)
Aspen	First year	OTC	Pennsylvania	8-h mean = 39, 73	11 weeks	Biomass	14-30 for 3 of 6 N treatments	Pell et al. (1995)
Beech, European		OTC	Switzerland	0.5×, 0.85×, 1×, 0.5×+30: AOT40 = 0.1, 3.4, 7.1 , 19.7 ppm•h	5 months	Biomass	6 at 1×, 30 at 0.5×+30	Landolt et al. (2000)
Beech, European	Seedling	OTC	Belgium	CF, NF, +30: 8-h mean = 5. 29 , 33; AOT40 = 0. 4.06 , 8.88 ppm•h	23 April - 30 Sept	Growth	No effect	Bortier et al. (2000a)

Table 9-18 (cont'd). Summary of Selected Studies of Ozone Effects on Deciduous Trees and Shrubs

Species	Age	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (Decrease from lowest, %)	Reference
Beech, European	Seedling	Growth chamber	Belgium	CF, CF+40, CF+100: Sum0 = 0.48, 8.93, 25.14 ppm•h; AOT40 of NF+100 = 13.91 ppm•h; uptake = 159, 2965 7095 mol m ²	7 episodes of 5 days	Biomass, diameter	No effect	Bortier et al. (2001)
Beech, European	0-3 years	OTC	Switzerland	AOT40 for 24 h/days = 4-73 ppm•h	1-3 years	Total biomass	20 at AOT40 for 24 h = 32 ppm•h	Braun and Fluckiger (1995)
Beech, Japanese	4 years	Growth chamber	Japan	CF, +60 ppb for 7 h/day	156 days	Total biomass	19	Yonekura et al. (2001)
Birch, silver	Sapling	FACE	Finland	AOT40 = 1, 15 ppm•h; 7-h mean = 26, 40	5 years	Biomass	34 for root, ns for stem	Oksanen et al. (2001)
Birch, silver	Sapling	OTC	Sweden	NF, NF+, NF++, daylight mean 1997 = 29 , 37, 54; 1998 = 25 , 42, 71 ppb; AOT40 1997 = 2.4 , 6.9, 35.1; 1998 = 0.6 , 19.6, 74.7 ppm•h	2 years	Total biomass	Total biomass ns at NF+, 22 at NF++; root biomass 30 at NF++	Karlsson et al. (2003)
Birch, [<i>B. pubescens</i>]	Seedling	Chamber in glasshouse	Norway	AOT40 = 0.1, 2.5, 7.1, 7.4, 17.8, 19.8 ppm•h	40 days	Biomass	Sig. decrease in root at AOT40 = 2.5 ppm•h, shoot at 7.1 ppm•h	Mortensen (1998)
Cherry, black	2 years	OTC	Norris, TN	CF, 1×, 2×: 7-h mean = 21, 50 , 97	April to August	Biomass	No effect	Samuelson (1994)
Cherry, black	Seedling	OTC	GSMNP ¹	CF, 1×, 1.5×, 2×: SUM06 = 0-40.6 ppm•h, AOT40 = 0.03-28.3 ppm•h	76 days	Biomass	n.s. at 1x and 1.5×, 38 at 2×	Neufeld et al. (1995)

Table 9-18 (cont'd). Summary of Selected Studies of Ozone Effects on Deciduous Trees and Shrubs

Species	Age	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (Decrease from lowest, %)	Reference
Cherry, black	Seedling	OTC	GSMNP ¹	CF, 0.5×, 1×, 1.5×, 2×: SUM06 = 0-53.7 ppm•h; AOT40 = 0-40.4 ppm•h	140 days	Biomass	n.s. at 1× and 1.5×, 59 at 2×	Neufeld et al. (1995)
Cherry, black	1 year	OTC	Delaware, OH	CF, 0.5, 1, 1.5, 2×: SUM00 in 1990 = 17-107 ppm•h, in 1991 = 31-197 ppm•h	2 years (in 1990 for 3.5 months, 1991 for 4 months)	Total biomass	no effect at 1× and 1.5×, 32 at 2×	Rebbeck (1996)
Cherry, black	Seedling	OTC	Pennsylvania	CF, 0.75×, 0.97×: 7-h mean = 39 to 46 , SUM06 = 0- 10.34 ppm•h	3 years for 17 weeks	Total biomass	6 at 0.75×, 14 at 0.97×	Kouterick et al. (2000)
Cottonwood, Eastern	Cutting	Ambient, in buried pots with irrigation	In and within 100 km of New York City, NY	12 h mean = 23-49 ppb	2 months each year, 3 10-years experiments	Total biomass	33% decrease at 38 ppb compared to 23 ppb	Gregg et al. (2003)
Grape	3 years	OTC	Austria	CF, 1×, +30, +50: (AOT40 = 0-50 ppm•h	2 years (preflowering, past harvest)	Fruit yield	Calculated 10 at AOT40 = 27 ppm•h	Soja et al. (1997)
Oak	Seedling	OTC	Hampshire, UK	NF, NF+: Mean = 17.7 , 44, AOT40 for 24h = 1.9 , 59.9 ppm•h	3 years for day 100 to day 162	Biomass of organs	30 for total biomass	Broadmeado w and Jackson (2000)
Oak, red	Seedling	OTC	Norris, TN	SUM06 for 3 years = 0, 29, 326 ppm•h; SUMOO for 3 years = 147, 255 and 507 ppm•h	3 years	Total biomass	n.s.	Samuelson et al. (1996)

Table 9-18 (cont'd). Summary of Selected Studies of Ozone Effects on Deciduous Trees and Shrubs

Species	Age	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (Decrease from lowest, %)	Reference
Oak, red	30 years	OTC	Norris, TN	SUM06 for 3 years = 0, 29, 326 ppm•h; SUM00 for 3 years = 147, 255 and 507 ppm•h	3 years	Stem increment	n.s. despite 50% reduction in net photosynthesis	Samuelson et al. (1996)
Maple, red	2 years	OTC	Norris, TN	CF, 1×, 2× : 7-h mean = 21, 50 , 97 ppm•h	April to August	Biomass	No effect	Samuelson (1994)
Maple, sugar	1 year	OTC	Delaware, OH	CF, 0.5, 1.5, 2×: UM00 in 1990 = 17 to 107 ppm•h, in 1991 = 31 to 197 ppm•h	2 years (in 1990 for 3.5 months, 1991 for 4 months)	Total biomass	n.s., but linear trend	Rebbeck (1996)
Maple, sugar	Seedling	Large OTC	Ithaca, NY	CF, 1×, 1.5×, 2×: 3 years SUM00 = 148 to 591 ppm•h; daytime mean = 19.7 to 40.7	3 years for 134, 128, 109 days	Biomass	No effect	Laurence et al. (1996)
Maple, sugar	Seedling	Large OTC	Ithaca, NY	1×, 1.7×, 3×: 3 years 12-h mean = 38 , 69, 117	3 years for 109, 143, 116 days	Total biomass	For 1.7× and 3×: 21, 64 in low light, 26 and 41 in high light	Topa et al. (2001)
Plum, Casselman	Sapling	Large OTC	Fresno, CA	CF, 1×, +O ₃ : 12-h mean = 31, 48 , 91	4 years	Stem increment, fruit yield	Fruit yield 16 at 1×, stem +14 at +O ₃	Retzlaff et al. (1997)
Poplar, black	Seedling	OTC	Belgium	CF, NF, +30: 8-h mean = 5, 29 , 33; AOT40 = 0, 4 , 8.9 ppm•h	23 April - 30 Sept	Diameter, height	29 for diameter in NF+, no effect on height	Bortier et al. (2000b)

Table 9-18 (cont'd). Summary of Selected Studies of Ozone Effects on Deciduous Trees and Shrubs

Species	Age	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (Decrease from lowest, %)	Reference
Poplar, hybrid (P. tremuloides × P. tremula)	0 year	FACE	Finland	AOT40 = 0.07, 1.6 ppm•h; 7-h mean = 30, 38	2 months	Biomass, height	n.s. for biomass, 6 for height	Oksanen et al. (2001)
Poplar, hybrid	Cutting	OTC	Michigan	CF, CF+100: 12, 48 ppm•h	60 days	Total biomass	46 for average of 5 clones	Dickson et al. (1998)
Yellow-poplar	1 year	OTC	Delaware, OH	CF, 0.5, 1.5, 2×: SUM00 in 1990 = 17 to 107 ppm•h, in 1991 = 31 to 197 ppm•h	2 years (in 1990 for 3.5 months, 1991 for 4 months)	Total biomass	No effect	Rebbeck (1996)

¹ Values for ambient or NF treatments are indicated in bold.

² **Bold** indicates that multiple experiments (more than just 2 years at a single site) were included in the analysis.

Table 9-19. Summary of Selected Studies of Ozone Effects on Evergreen Trees and Shrubs

Species	Age	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (decrease from lowest, %)	Reference
Fir, Douglas	Seedling	Open air	British Columbia	12 trts: 12-h mean 1988 = 18-41; 1989 = 27-66	1988 = 92 days; 1989 = 101 days	Second flush biomass	Calculated 55 at highest exposure	Runeckles and Wright (1996)
Hemlock, eastern	Seedling	OTC	GSMNP ¹ , TN	CF to 2×: SUM06 = 0.2-108.1 ppm•h, AOT40 = 0.2-63.9 ppm•h	3 years	Biomass	No effect	Neufeld et al. (2000)
Pine, loblolly	12 weeks	OTC	Oak Ridge, TN	CF to 2×: 24-h summer = 74, 137, 169, 206, 284 ppm•h	3 months	Biomass	14 in 1× (avg for all families)	McLaughlin et al. (1994)
Pine, loblolly	1 year	OTC	Alabama	1994: AOT40 = 2-112 ppm•h, SUM06 = 10-162 ppm•h, 1995: AOT40 = 3-83 ppm•h, SUM06 = 0-132 ppm•h	2 years, April to October	Dry weight, height, diameter	n.s.	Barbo et al. (2002)
Pine, loblolly	3 years	OTC	Raleigh, NC	Ambient, CF, NF, 1.5×., 2.5×: 12-h mean = 54 , 29, 47, 76, 98	5 months	Height, diameter, needle length	No effect on stem height or diameter, decrease in needle length	Anttonen et al. (1996)
Pine, ponderosa	Seedling	OTC	Corvallis, OR	For CF 12-h SUM06 = 0 ppm•h; for +03 12-h SUM06 = 22, 27, 31 ppm•h for 3 years	3 years: 16 weeks, 16 weeks, 14 weeks	Total biomass	No effect without grass, 25 with grass present	Andersen et al. (2001)
Pine, ponderosa	39 to 45 years	Ambient gradient	CA	24-h mean for 3 weeks late July and early August for 1993 and 1994 = 70-90 ppb	Ambient gradient	Fine and medium root growth	85 at most polluted site.	Grulke et al. (1998)
Pine, ponderosa	Seedling	OTC	CA	CF, 1×, 2×: 24-h mean approx. 20, 60 , 120		Total biomass	n.s.	Takemoto et al. (1997)

Table 9-19 (cont'd). Summary of Selected Studies of Ozone Effects on Evergreen Trees and Shrubs

Species	Age	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (decrease from lowest, %)	Reference
Pine, Scots		OTC	Hampshire, UK	NF, NF+; Mean = 17.7 , 44.1; 24-h AOT40 = 1.9 , 59.9 ppm•h	3 years for 62 days	Total biomass	15	Broadmeadow and Jackson (2000)
Pine, Scots	3-6 years	Free air	Finland	Amb, +O ₃ ; AOT40 = 0-1 , 2-13 ppm•h	3 years	Biomass	No effect	Kainulainen et al. (2000)
Pine, Scots	Seedling	OTC	Switzerland	0.5×, 0.85×, 1×, 0.5×+30: AOT40 = 0.1, 3.4, 7.1 , 19.7 ppm•h	5 months	Biomass	14 at 1×, 22 at 0.5×+30	Landolt et al. (2000)
Pine, Scots	3 years	OTC	Finland	CF, 1×, +O ₃ ; 24 h AOT40 for 2 years = 0.5, 6 , 73 ppm•h	2 years (4 months each)	Biomass	No effect	Utriainen et al. (2000)
Pine, Scots	3 years	Free air	Finland	1×, +O ₃ ; 24 h AOT40 for 2 years = 2 , 37 ppm•h	3 years (3-4 months each)	Root and shoot biomass	32 only for root biomass in high N treatment	Utriainen and Holopainen (2001)
Pine, Table Mountain	Seedling	OTC	GSMNP ² , TN	CF to 2×: SUM06 = 0.2-116.4 ppm•h, AOT40 = 0.2-71.7 ppm•h	3 years	Biomass	Slight decrease in older needle mass only	(Neufeld et al. (2000)
Pine, Virginia	Seedling	OTC	GSMNP ² , TN	CF to 2×: SUM06 = 0.1-32.8, 47.9, 56.2 ppm•h; AOT40 = 0.1-19.3, 27.1, 34.4 ppm•h	1-2 years (3 expts)	Biomass	No effect	Neufeld et al. (2000)
Sequoia, giant	125 years	Branch chamber	California	0.25×, 1×, 2×, 3×: 24-h SUM00 approx. 10, 85 , 180, 560 ppm•h	61 days	Branch growth	No effect	Grulke et al. (1996)
Spruce, Norway	4-7 years	Open air	Finland	Amb, +O ₃ ; AOT40 = 0 1 , 2-13 ppm•h	3 years	Biomass	No effect	Kainulainen et al. (2000)

Table 9-19 (cont'd). Summary of Selected Studies of Ozone Effects on Evergreen Trees and Shrubs

Species	Age	Facility	Location	O ₃ Concentration (Units are ppb unless otherwise specified) ¹	Duration	Variable	Response (decrease from lowest, %)	Reference
Spruce, Norway	3-7 years	OTC	Sweden	CF, 1.5×: 12-h mean for 4 years = 12, 44; AOT40 = 2, 23 ppm•h	4 years	Total biomass	8	Karlsson et al. (2002)
Spruce, Norway	Seedling	OTC	Switzerland	0.5×, 0.85×, 1×, 0.5×+30: AOT40 = 0.1, 3.4, 7.1 , 19.7 ppm•h	5 months	Biomass	n.s.	Landolt et al. (2000)
Spruce, Norway	0-3 years	OTC	Switzerland	AOT40 for 24 h for 1 to 3 years = 22 to 63 ppm•h	1-3 years	Total biomass	n.s.	Braun and Fluckiger (1995)
Spruce, Norway	Seedling	OTC	Sweden	CF, 1×, 1.5×: AOT40 daylight for 4 years = 1, 16 , 79 ppm•h	4 years	Stem volume	No effect in final year	Wallin et al. (2002)
Spruce, red	Sapling	Large OTC	Ithaca, NY	CF, 1×, 1.5×, 2×: total for 4 years = 211 to 569 ppm•h; daytime mean = 21-71	4 years: 98-124 days/year	Biomass	No effect	Laurence et al. (1997)

¹ Values for ambient or NF treatments are indicated in bold.² Great Smoky Mountains National Park.

1 exposures. The development of critical levels in Europe has been based primarily on the AOT40
2 index, so this index is discussed in that context.

3 In addition to peak-weighting, there is also evidence that the timing of exposure during
4 plant growth is important. For example, the greatest effects on grain yield are due to exposure
5 during grain filling, rather than earlier or later in the growing season (Lee et al. 1988; Pleijel
6 et al., 1998; Soja et al., 2000; U.S. Environmental Protection Agency, 1996; Younglove et al.,
7 1994). The importance of respite times was also discussed in the previous criteria documents
8 (U.S. EPA, 1978, 1988, 1996) but remains difficult to quantify (Section 9.5). Even when some
9 of these aspects of O₃ exposure can be elucidated, it is difficult to apply this knowledge to
10 developing exposure-response relationships based on data in the scientific literature, because O₃
11 exposure is often reported only in the form of a summary index such as a 12- or 24-h mean,
12 SUM06, or AOT40.

13 Table 9-14 presents summaries of ambient O₃ exposure patterns in the United States for
14 1982 to 1981 for several indices including the 7-h mean and SUM06. More recent summaries
15 for the entire United States for these indices are not available, but Table 9-15 summarizes more
16 recent data for the central and eastern United States. As shown in Table 9-15, from 1989 to
17 1995, mean 12-h 3-month SUM06 values (in ppm-h) at 41 rural sites in the Clean Air Status and
18 Trends Network were 31.5 for the Midwest, 18.9 for the Upper Midwest, 33.2 for the Northeast,
19 13.2 for the Upper Northeast (NH, ME), 34.5 for the South-Central, and 19.2 for the Southern
20 Peripheral subregions (Baumgardner and Edgerton, 1998). These results are important because
21 these sites were selected to represent rural areas, while many other monitoring sites represent
22 urban or suburban areas. For these same subregions, W126 values ranged from 12.8 to
23 25.6 ppm-h. From 1989 to 1995, O₃ concentrations decreased about 5% for daily and 7% for
24 weekly values for most of these sites, after adjusting for meteorological conditions (Holland
25 et al., 1999). These trends were statistically significant at about 50% of the sites ($p \leq 0.05$).
26 However, because the trend analysis was intended to examine the efficacy of O₃ emissions
27 controls, the trends were adjusted for meteorological conditions. Thus, they do not reflect the
28 actual trends in O₃ exposure over time.

Table 9-20. Ethylene Diurea Effects on Vegetation Responses to Ozone

Species	Description	EDU application	Ozone exposure	Effects of EDU	Reference
Bean, cv. Lit	10-cm pots in OTCs in Germany	Soil drench 200 mL of 150 ppm solution per plant every 14 days	CF, NF, CF-1×, CF-2×; mean = 1, 14, 15, 32 ppb	O ₃ reduced pod, shoot, and root mass. EDU increased root, leaf, and shoot mass, but a significant interaction with O ₃ occurred only for root weight.	Brunschon-Harti et al. (1995)
Bean, cv. BBL-290	2 expts in 5.5 L pots in OTCs with 4 O ₃ treatments	Soil drench every 14 days, in expt 1 = 0.14, 28, 56, 120 mg/L potting medium; expt 2 = 0, 8, 16, 32 mg/L	2 expts with CF, NF and 2 constant additions of O ₃ . 7-h mean O ₃ (ppb) for Expt 1 = 34, 70, 95, 121; Expt 2 = 19, 42, 74, 106	Visible injury and reduced total biomass or yield, even in CF treatment. Within an O ₃ treatment, sometimes increased yield (Expt 2 only).	Miller et al. (1994)
Bean	Pots with potting mix at 3 locations in Spain (2 years at 1 site)	Soil drench of 200 mL of increasing concentrations of 100, 150, 200, 250 ppm every 14 days (4-10 mg l:1 soil)	AOT40 = 0.4-1.8 ppm-h	0 to 50% increase in pod mass, but did not restore yield at sites with higher O ₃ .	Ribas and Penuelas (2000)
Bean, cv. Lit	Pots with potting mix at 4 sites in the Netherlands	Soil drench of 200 mL of increasing concentrations of 100, 150, 200, 250 ppm every 14 days (4-10 mg l:1 soil)	AOT40 = 0.64-0.98, 7-h mean = 49-55 ppb	Average 20% yield increase at all sites.	Tonneijck and Van Dijk (1997)
Bean, cv. Lit	Pots with potting mix at 1 site in Belgium	Soil drench of 200 mL of increasing concentrations of 100, 150, 200, 250 ppm every 14 days (4-10 mg l:1 soil)	AOT40 = 0.81 ppm-h	16% yield increase.	Vandermeiren et al. (1995)
Clover, subterranean	Plants in 10-cm pots at 4 rural sites in the Netherlands for 3 years	100 ml of 150 ppm solution as soil drench every 14 days for 2 months	AOT40 = 0-0.56 ppm-h for 4-week periods	Injury, but not leaf biomass was affected by EDU and O ₃ exposure.	Tonneijck and Van Dijk (2002)

Table 9-20 (cont'd). Ethylene Diurea Effects on Vegetation Responses to Ozone

Species	Description	EDU application	Ozone exposure	Effects of EDU	Reference
Clover, white	15-cm pots in field, well watered, 12 locations throughout Europe, 3 years	100 ml of 150 ppm solution as soil drench every 14 days for 3 months	AOT40 (28 days) = 0-20 ppm-h	Change in biomass ratio, weak linear relationship ($r^2 = 0.16$) stronger relationship using ANN and climatic factors	Ball et al. (1998)
Clover, white, cv. Menna	2 expts, 10-cm pots in field in Italy, see also companion OTC expt	100 ml of 150 ppm solution as soil drench every 14 days for 2 months	AOT40 = 15.5, 12.1 ppm•h; 7-h mean = 69, 60	n.s.	Fumagalli et al. (1997)
Clover, white, cv. Menna	2 expts, 10-cm pots in OTCs in Denmark	Soil drench 100 mL of 150 ppm solution every 14 days	CF, NF, NF+ 25, NF+50 ppb, O ₃ exposure not reported	No effect of EDU despite highly significant effect of O ₃ on above-ground biomass	(Mortensen and Bastrup-Birk, 1996)
Poplar, hybrid	Stem injections, field, cuttings, 1 or 2 years	Approx. 125 or 250 mg/leaf (low, high EDU treatments) 5 times every 14 days	1991: 7-h mean = 56, AOT40 = 23; 1992: 7-h mean = 59, AOT40 = 27	No effect on biomass; 6%, 12% more severely O ₃ damaged leaves in high EDU for 2 years	Ainsworth et al. (1996)
Pine, loblolly	1 year old half-sib seedlings in field in TX for 3 years	150, 300, 450 ppm every 14 days	1995, 1996, 1997, no. h > 40 ppb = 1723, 2297, 2052; no. h > 60 ppb = 378, 584, 528; peak = 113, 102, 118	For EDU 450 trt, above-ground biomass increased approx 46% (n.s. in other treatments)	Manning et al. (2003)
Radish, cv. Cherry Belle	Plants in pots in potting mix exposed for 5 weeks in southern Sweden.	Soil drench containing 20 mg EDU applied 2 times, 14 days apart	24-h mean = 31 ppb, 7-h mean = 36 ppb, AOT40 = 1.3 ppm-h.	24% increase in hypocotyl mass, 18% increase in shoot mass	Pleijel et al. (1999b)

9.6.4 Effects of Ozone on Annual and Biennial Species

Much of the research on short-lived species during the last decade has been conducted in Europe. Several European studies have focused on wheat with an emphasis on developing critical levels, as discussed below in Section 9.6.4.6.

An extensive search of the literature was performed using several electronic databases to identify scientific articles containing quantitative information on both the amount of O₃ exposure and its effects on vegetation. Greater emphasis is placed on studies with longer duration with O₃ exposure concentrations and environmental conditions that were as similar as possible to ambient conditions. Many of the studies reviewed herein were conducted in OTCs. In the United States, nearly all of such studies have used the type of OTC developed by Heagle et al. (1973). For the few studies in the U.S. that used other types of OTCs, they are described briefly in the text. In Europe, a wide variety of styles of OTCs have been used. See Section 9.2 for further information about the use of OTCs. The emphasis in this subsection is on quantifying exposure-response relationships for annual plants, with a focus on the response of above-ground biomass and yield of species grown as crops or occurring as native or naturalized species in the United States. Emphasis is placed on studies not included in the 1996 AQCD (U.S. Environmental Protection Agency, 1996), including a few studies published prior to 1996. However, an attempt is made to compare the results of these recent studies of individual species to those reviewed in the 1996 AQCD.

9.6.4.1 Effects on Growth, Biomass, and Yield of Individual Species

Most research on the effects of O₃ on herbaceous species has evaluated growth, biomass, or yield of commercial portions of crop or forage species. It is well established that reproductive organs such as seeds may be particularly sensitive to injury or biomass reductions due to O₃, as reviewed recently by (Black et al. 2000}. As discussed in Section 9.4, numerous analyses of experiments conducted in OTCs and with naturally occurring gradients demonstrate that the effects of O₃ exposure vary depending on the growth stage of the plant. Plants grown for seed or grain are often most sensitive to exposure during the seed or grain-filling period (Lee et al., 1988; Pleijel et al., 1998; Soja et al., 2000; Younglove et al., 1994), whereas plants grown for biomass production, such as alfalfa, may be sensitive throughout the growth period (Younglove et al., 1994). However, because different species are sensitive during different periods of their

1 growth and, because planting or germination dates vary throughout large regions even for a
2 single species, no single phenological weighting scheme can appropriately and practically
3 represent all vegetation in all locations in the United States. For natural populations, reductions
4 in seed yield might be particularly important if subsequent seedling establishment is
5 compromised by O₃.

6 Green beans (cv. Pros) were grown in pots in OTCs in the Netherlands for 62 days and
7 exposed to 6 treatments consisting of constant O₃ additions to charcoal filtered (CF) chambers
8 (see Section 9.2) for 9 h/day (Tonneijck and Van Dijk, 1998). Bean yield response to O₃ was
9 nonlinear, with an apparent threshold near the CF30 (charcoal filtered with a constant addition of
10 30 ppb O₃) treatment with a 9-h mean O₃ concentration of 28 ppb and an AOT40 value of
11 0.1 ppm·h). Yield was reduced by 29% at a 9-h mean value of 44 ppb corresponding to an
12 AOT40 value of 3.6 ppm·h (Table 9-17). Beans were grown in pots in OTCs for 3 months with
13 the following O₃ treatments: CF, nonfiltered (NF), CF with O₃ added up to ambient, and CF
14 with 2×-ambient O₃ (Brunschon-Harti et al., 1995). Ozone reduced pod mass by 56% with a
15 mean concentration of 32 ppb in the 2× ambient treatment as compared with 1 ppb in the CF
16 treatment (daily averaging time not reported). A second treatment factor in this experiment was
17 addition of EDU, as discussed below under the heading “Studies Using Ethylene Diurea as a
18 Protectant”. These yield reductions are greater than those previously reported in four similar
19 studies summarized in the 1996 AQCD (Table 5-25 of U.S. Environmental Protection Agency,
20 1996). Greater sensitivity in the more recent experiments may be due to cultivar differences or
21 other differences in experimental protocols.

22 In a study with OTCs on silty loam soil in Beltsville, MD, corn yield was reduced by 13%
23 with exposure to a 7-h mean concentration of 70 ppb O₃ compared to a CF treatment with a 7-h
24 mean concentration of 20 ppb (Mulchi et al., 1995; Rudorff et al., 1996a; Rudorff et al., 1996c).
25 In this study, different amounts of O₃ were added above ambient levels for 5 days follows: 20,
26 30, 40, 50, 60 ppb, except that O₃ was not added to exceed a total concentration of 120 ppb
27 (Rudorff et al., 1996c, 1996a).

28 In two studies conducted in Raleigh, NC, cotton (cv. Deltapine 51) was grown in pots and
29 exposed to CF and 1.5× (nonfiltered, see Section 9.2) O₃ in one year, and CF, NF, and
30 1.5×-ambient O₃ in the second year, with ambient and elevated CO₂ concentrations (Heagle
31 et al., 1999; Table 9-17). In the first year, yield decreased by 22% with 1.5×-ambient O₃ (12-h

1 mean value of 71 ppb). In the second year, yield decreased by 21% and 49% with exposure to
2 ambient or 1.5× ambient O₃ (12-h mean values of 51 and 78 ppb, Table 9-17). Increased CO₂
3 levels prevented or reduced this yield suppression (Heagle et al., 1999). These yield reductions
4 are similar to those reported previously in four similar studies summarized in the 1996 AQCD
5 (Table 5-25 of U.S. Environmental Protection Agency, 1996).

6 In a study of oats in OTCs in southern Sweden, exposure to ambient (NF) O₃ did not affect
7 grain yield (Pleijel et al., 1994a). Ambient O₃ concentration expressed as a 7-h mean was
8 27 ppb, with only 1 h greater than 80 ppb and none above 90 ppb.

9 The interactive effects of elevated O₃ and CO₂ additions on potato yield (cv Bintje) were
10 studied in OTCs at 6 sites in northern Europe as part of the CHIP (Changing Climate and
11 Potential Impacts on Potato Yield and Quality) program (Craigon et al., 2002). Ozone was
12 added to a target daily average value of 60 ppb, and AOT40 values across all years and
13 experiments ranged from ~6 to 27 ppm·h. The O₃ treatment reduced total tuber yield an average
14 of 4.8% with elevated O₃ treatment across all experiments (Craigon et al., 2002). This total
15 effect was statistically significant even though the effects of individual experiments generally
16 were not (Craigon et al., 2002), due to the increased power of the pooled analysis. Several
17 publications report other aspects of the CHIP experiments or present results of individual
18 experiments (De Temmerman et al., 2002a, 2002b; Donnelly et al., 2001a, 2001b; Fangmeier
19 et al., 2002; Finnan et al., 2002; Hacour et al., 2002; Lawson et al., 2002; Pleijel et al., 2002;
20 Vandermeiren et al., 2002; Vorne et al., 2002).

21 The effect of an intermittent constant addition of O₃ using a free air exposure system in
22 Northumberland, UK was investigated with the oilseed rape cultivar “Eurol” (Ollerenshaw et al.,
23 1999). Ozone was added for 6 h/day for 17 days. The ambient treatment had a mean value of
24 30 ppb and the O₃ addition treatment had a mean of 77 ppb. After overwintering, O₃ was added
25 for 32 days for 7 h/day between May and June (mean values of 31 and 80 ppb). Yield was
26 reduced by 14% despite the lack of any foliar symptoms.

27 Field fumigation chambers ventilated with fans on both ends were used to assess effects of
28 five O₃ treatments on rice over 3 years in Japan (Kobayashi et al., 1994; 1995). All O₃
29 treatments used CF air, and O₃ was added to the 0.5, 1.0, 1.5, 2.0, or 2.75× ambient
30 concentration for 7 h/day. Based on a linear regression for the 3 years, yield decreased by 3 to
31 10% at a 7-h mean concentration of 40 ppb (Table 9-17). This decrease is greater than that

1 found for rice in earlier studies in California (Kats et al., 1985), although whether this difference
2 is due to differences in cultivars, experimental treatment, or environmental factors cannot be
3 determined.

4 During 3 years in Beltsville, MD, the soybean cultivars Essex and Forrest were exposed to
5 CF air and NF air in OTCs (Chernikova et al., 2000; Robinson and Britz, 2000) with O₃ added as
6 described for experiments with corn and wheat at Beltsville (Mulchi et al., 1995; Rudorff et al.,
7 1996c). During 1994 and 1995, as previously found for these cultivars, Essex was less sensitive
8 than Forrest, with yield decreases of 10% (n.s.: $p > 0.1$) compared to 32% for Forrest ($p < 0.01$)
9 (Chernikova et al., 2000). There was no evidence of water stress in this experiment. In 1997,
10 the two O₃ treatments were CF (7-h mean = 24 ppb) and NF with a constant addition of O₃ (7-h
11 mean = 58 ppb) (Robinson and Britz, 2000). The yield of Essex was not significantly affected,
12 while the yield of Forrest was decreased by 21% (Table 9-17).

13 In a study in Raleigh, NC the soybean cultivar Essex was grown in pots and exposed to CF
14 and 1.5× ambient O₃ concentrations during three growing seasons (Fiscus et al., 1997). Over the
15 3 years, exposure to an average 12-h mean O₃ concentration of 82 ppb reduced soybean yield by
16 41% (Table 9-17). In similar studies also in Raleigh, NC, Essex was exposed to CF, NF, and
17 1.5× ambient O₃ for two seasons (Heagle et al., 1998). Yield decreased by 16% and 15% in the
18 2 years by ambient O₃ (12-h mean values of 50 and 42 ppb), and decreased by 37 and 40% with
19 exposure to 1.5× ambient O₃ (12-h mean values of 79 and 69 ppb, Table 9-17). In this same
20 experiment in the second year, similar yield reductions were observed for the cultivar Holladay,
21 while the growth of cultivar NK-6955 was increased substantially by ambient O₃ exposure.
22 All three cultivars were grown in the same chambers in this experiment, and the authors
23 suggested that NK-6955 plants may have shaded the other cultivars to some extent.

24 In a 2-year study using OTCs in Raleigh, NC, the soybean cultivars Coker 6955, Essex,
25 and S53-34' were exposed CF, NF, and 1.5× ambient O₃ treatments (Miller et al., 1994).
26 Seasonal mean 12-h O₃ concentrations ranged from 14 to 83 ppb. As compared to the CF
27 treatment, ambient O₃ exposure (NF treatment) reduced seed yield by 11 to 18% except for
28 Coker 6955 in the first year (1989) which showed a yield increase of 14%. The 1.5× ambient O₃
29 treatment reduced yield by 32 to 56% in all cultivars in both years. In a similar subsequent
30 experiment with the cultivar Essex, exposure to a 12-h mean ambient O₃ concentration of 50 ppb

1 reduced yield by 11%, while exposure to 79 ppb reduced yield by 22% (Table 9-17) (Miller
2 et al., 1998).

3 These yield reductions for soybean are generally similar to those reported previously in
4 13 similar studies summarized in the 1996 AQCD (Table 5-23 of U.S. Environmental Protection
5 Agency, 1996).

6 A reanalysis of 7 years of data from OTC experiments with wheat in Ostad, Sweden
7 showed that relative yield linearly decreased with increasing O₃, with a maximum yield loss of
8 23% at an AOT40 value of 15 ppm·h (Danielsson et al., 2003). A very similar response was
9 found using the flux (stomatal conductance) model of Emberson et al. (2000b), and a similar
10 amount of the variance was explained by the flux model (for AOT40 model, $r^2 = 0.34$ and for the
11 Emberson flux model, $r^2 = 0.39$). A modified flux model developed and calibrated for this site
12 also had a similar linear response equation, but explained much more of the variance ($r^2 = 0.90$).

13 During the 1990s, a major European research program investigated the combined effects of
14 CO₂, O₃, and other physiological stresses on wheat (Bender et al., 1999; Hertstein et al., 1996,
15 1999; Jäger et al., 1999). The European Stress Physiology and Climate Experiment (“ESPACE-
16 wheat”) program included 13 experiments in OTCs at eight sites in northern Europe over 3
17 years. Low- and high-O₃ exposures in these experiments had the following values: 12-h mean
18 (SD) low = 26.3 ppb (12.2), 12-h mean (SD) high = 51.37 (18.3) ppb, AOT40 mean (SD) low =
19 6.2 (8.5) ppm-h, AOT40 mean (SD) high = 28.3 (23.0) ppm-h, as calculated from data presented
20 in Table 3 of Hertstein et al. (1999). An analysis of all 13 experiments showed that high O₃ at
21 ambient CO₂ reduced yield by 13% on average (Bender et al., 1999). However, this reduction
22 was not statistically significant based on an ANOVA, and the authors concluded that the wheat
23 cultivar Minaret may be relatively tolerant to O₃ (Bender et al., 1999). Results of some
24 individual studies within this program have been reported previously (Donnelly et al., 1999;
25 Fangmeier et al., 1996, 1999; Mulholland et al., 1997, 1998b, 1998a; Pleijel et al., 2000b).

26 In a study with OTCs on silty loam soil in Beltsville, MD, wheat yield was reduced by
27 20% on average over 2 years with 7-h mean concentrations of 61 and 65 ppb O₃ compared with
28 CF treatment with a 7-h mean concentration of 20 ppb (Mulchi et al., 1995; 1996a; Rudorff
29 et al., 1996c). In the above study, different amounts of O₃ were added above ambient (levels for
30 5 days) as follows: 20, 30, 40, 50, 60 ppb, except that O₃ was not added to exceed a total
31 concentration of 120 ppb (Rudorff et al., 1996c). Wheat grown in pots in OTCs was exposed to

elevated O₃ and water stress in Germany, and yield was decreased by 35% in the 2×-ambient treatment with a 7-h mean O₃ concentration of 71 ppb statistically significant effects were not seen in the 1×-ambient treatment (Fangmeier et al., 1994). In two studies conducted in Raleigh, NC, soft red winter wheat was grown in pots and exposed to CF, NF, and 1.5×-ambient O₃, with ambient and elevated CO₂ concentrations (Table 9-17) (Heagle et al., 2000). In the first experiment, eight cultivars were exposed to 12-h mean O₃ concentrations of 27, 47, and 90 ppb, and in the second experiment two of these cultivars were exposed to 22, 38, and 74 ppb. There was a trend toward decreased yield in both experiments, but these trends were not statistically significant. The wheat cultivar Drabant was exposed to CF, NF, and a constant addition of 35 ppb during 1992 and 1993 using the Heagle-type OTCs (Heagle et al., 1973) in Finland (Ojanpera et al., 1998). The following 12-h mean O₃ exposures were observed in 1992: 14, 30, 61 ppb. In 1993, the values were 9, 21, and 45 ppb, (see Table 9-17 for AOT40 values and other information). Yield was reduced 13% in each year by the added O₃ treatment.

The effect of an intermittent constant addition of O₃ using a free air exposure system was investigated with the winter wheat cultivar Riband in Northumberland, UK (Ollerenshaw and Lyons, 1999). Ozone exposures expressed as AOT40 values for September and October 1992 were 0.14 and 3.5 ppm·h; while for April to August 1993, values were 3.5 and 6.2 ppm·h. Yield was reduced by 13%.

These results provide an additional line of evidence supporting the OTC-studies that demonstrated yield reductions in wheat due to O₃ exposures that occur in the United States. These yield reductions for wheat are generally similar to those reported previously in 22 comparable studies summarized in the 1996 AQCD (Table 5-25 of U.S. Environmental Protection Agency, 1996).

9.6.4.2 Effects on Plant Quality

In addition to reductions in biomass or crop yield, O₃ may also reduce the quality or nutritive value of annual species. Many studies have shown effects of O₃ on various measures of plant organs that affect quality, with most studies focusing on characteristics important for food or fodder (U.S. Environmental Protection Agency, 1996).

The effect of a continuous intermittent addition of O₃ using a free air exposure system in Northumberland, UK was investigated with the oilseed rape cultivar Eurol as discussed above

(Ollerenshaw et al., 1999). Ozone exposures expressed as AOT40 values for August to October 1991 were 0.2 and 3.8 ppm·h and for June 1992 were 0.7 and 8.1 ppm·h. Yield quality measured as crude protein and oil content was decreased significantly. Because the price of the product is reduced in direct proportion to the oil content, such a decrease represents a substantial loss to growers (Ollerenshaw et al., 1999).

Two wheat cultivars, Massey and Saluda, were each grown for one year each in Beltsville, MD (Table 9-17) and exposed to either CF or an addition of 40 ppb for 7 h/day for 5 days/week (Mulchi et al., 1995, 1996a; Rudorff et al., 1996c). Milling and baking quality scores and flour protein were not significantly affected by elevated O₃ exposure, but the softness equivalent was increased slightly (2.4%) in both experiments (Rudorff et al., 1996b). The authors concluded that these changes, along with other slight changes due to an increased CO₂ treatment, suggested that O₃ and CO₂ had only minor effects on wheat grain quality. In wheat grown in Sweden, the harvest index was significantly decreased and the protein content increased due to exposure to a 12-h mean of 48 ppb (Gelang et al., 2000). In an analysis of 16 experiments conducted with spring wheat and either O₃ or CO₂ exposures in four Nordic countries, a negative linear relationship was found between grain yield and grain protein content ($y = -0.38x + 138.6$, expressed as percentages of the NF treatment (Pleijel et al., 1999a).

For three soybean cultivars grown in Raleigh, NC, O₃ significantly decreased oleic acid content, although the authors stated that the reduction was not large enough to be economically important (Heagle et al., 1998).

In a UK study, potato exposed during 1998 to an AOT40 value of 12.5 ppm·h in OTCs (in Nottingham) resulted in the paste from tubers being more viscous (Donnelly et al., 2001b). In this study, an AOT40 exposure of 27.11 ppm·h in 1999 caused starch granules to be less resistant to swelling, and total glycoalkaloid content was increased due to an increase in α-solanine (Donnelly et al., 2001b). Such increases in glycoalkaloid content have been observed previously in potato (Pell and Pearson, 1984) and may be important, because glycoalkaloids cause bitter flavors and, at higher concentrations, cause toxicity. The authors indicated that levels found in this study approached those that may cause bitterness, but not those of concern for toxicity (Donnelly et al., 2001b).

In the CHIP program the effects of O₃ were studied using OTCs at six sites in northern Europe, and yield decreases were observed as described above. The reducing sugar and starch

content of tubers decreased linearly due to O₃ exposure, while the ascorbic acid concentration increased linearly (Vorne et al., 2002). Compared to the CF treatment, exposure to an AOT40 value of 14 ppm·h decreased starch concentrations by 2%, decreased reducing sugar concentration by 30%, and increased ascorbic acid concentration by 20%. While the changes in reducing sugars and ascorbic acid increase tuber quality, the reduction in starch concentration decreases tuber quality.

In two 1-year studies using OTCs in commercial fields in Spain, the soluble solids content of watermelon was decreased 4 to 8% due to seasonal O₃ exposures as follows: AOT40 = 5.96 ppm·h and SUM06 = 0.295 ppm·h in one year; and AOT40 = 18.9, SUM06 = 4.95 in the second year (Gimeno et al., 1999).

9.6.4.3 Effects on Foliar Symptoms

For most annual crop species, the most important effects of O₃ are on yield of the commercially important part of the crop, expressed as the mass of the harvested portion. However, for some crops, foliar symptoms are important if they reduce the marketability of the crop. This is why efforts have been made to identify O₃ exposures associated with foliar symptoms. In Europe, Level I critical levels have been determined for such effects based on observations from experiments conducted in 15 countries under the auspices of the United Nations Economic Commission for Europe International Cooperative Programme on effects of air pollution and other stresses on crops and non-woody plants (UN/ECE ICP-Vegetation; formerly ICP-Crops), as well as on observations of symptoms in commercial fields from 1993 to 1996 (Benton et al., 1995, 2000). Because the occurrence of symptoms increased with greater humidity, these levels took into account the vapor pressure deficit (VPD). Two short-term critical levels were derived from 1995 data: an AOT40 value of 0.2 ppm·h over 5 days when mean VPD is below 1.5 KPa (0930 - 1630 h), and a value of 0.5 ppm·h when the mean VPD is above 1.5 Kpa (Benton et al., 1996). The 1996 data supported the critical levels in 83% of observations, although symptoms occurred on three occasions when the AOT40 was less than 0.05 ppm·h and the VPD was very low — less than 0.6 Kpa. The authors concluded that these critical levels are a good indicator of the likelihood of foliar symptoms, but that further refinement may be required, such as including factors that modify O₃ uptake by stomata.

1 In a study in Germany, 25 native herbaceous species were exposed to several square-wave
2 O₃ exposures in CF OTCs (Bergmann et al., 1999). Six of the 25 species showed O₃-specific
3 symptoms, and five species responded to single-day peaks of a single day. The most sensitive
4 species exhibiting O₃-specific symptoms were *Cirsium arvense* and *Sonchus asper*, which
5 responded to AOT40 values < 1.5 ppm·h (Bergmann et al., 1999).

6 In the United States, attention has been paid to foliar symptoms on annual plants in
7 previous criteria documents, and most recent literature has focused on other topics. However, a
8 study of O₃ effects was undertaken from 1990 to 1993 in Acadia National Park in Maine, which
9 experiences elevated O₃ exposures due to transport from urban areas located upwind (Kohut
10 et al., 2000). Because this study examined both herbaceous perennial species and woody
11 perennial species, the results are discussed in the perennial section below. Other studies and
12 biomonitoring programs in other U.S. National Parks and over larger areas are oriented primarily
13 towards forested ecosystems, so such studies are also discussed in the perennial subsection
14 below as well as in Section 9.7.

16 **9.6.4.4 Other Effects**

17 In addition to the effects seen on individual species grown in monocultures, effects may
18 occur due to competition when species are grown in mixtures, as reviewed by Davison and
19 Barnes (1998). Such effects have been investigated primarily in species grown for forage, which
20 often include perennial species; therefore, this topic is discussed in the herbaceous perennial
21 section, Section 9.6.5.1. Other stresses may interact with O₃, including pathogens and pests as
22 well as environmental conditions such as drought. These issues are discussed in Section 9.4.
23 Effects on reproduction are discussed briefly below.

24 Several studies during recent decades have demonstrated O₃ effects on different stages of
25 reproduction. Effects of O₃ have been observed on pollen germination, pollen tube growth,
26 fertilization, and abortion of reproductive structures as reviewed by Black et al. (2000). This
27 issue is not addressed here because reproductive effects will culminate for seed-bearing plants in
28 seed production, and the substantial body of evidence relating O₃ exposure and reduced seed
29 production is discussed above. However, one example of a native species will be presented
30 because of its implications for extrapolating exposure-response data to noncommercial species.
31 Spreading dogbane has been identified as a useful species for O₃ biomonitoring because of

O₃-induced diagnostic symptoms (Kohut et al., 2000). A study in Massachusetts found that exposure to O₃ in NF OTCs or ambient plots for 103 days produced significantly fewer flowers, and also that fewer of these flowers survived to produce mature fruits (Bergweiler and Manning, 1999). Because foliar symptoms were not common, the authors concluded they are not required for effects on reproduction to occur. Genotoxic effects and effects on population genetics are discussed in Section 9.4.

9.6.4.5 Scaling Experimental Data to Field Conditions

Substantial effort has been invested in the design of OTCs for assessing the effects of air pollutants on vegetation under near-ambient conditions. The design, construction, and performance of many types of chambers has been reviewed extensively (Hogsett et al., 1987a, 1987b). Despite such design efforts, the influence of experimental chambers on exposure-response functions has been debated for many years (e.g., Manning and Krupa, 1992) because several factors differ between OTC studies and actual fields. This issue is addressed in Section 9.2, and only a few comments about the implications of chamber artifacts for interpreting exposure-response relationships are presented here.

While it is clear that chambers can alter some aspects of plant growth, the more important issue is whether they alter the response of plants to O₃. A review of such chamber studies done in California found that plants responded similarly to O₃ whether OTCs, closed-top chambers, or air exclusion systems were used; differences were found for fewer than 10% of growth parameters (Olszyk et al., 1986). In a different review of literature about Heagle-type OTCs (Heagle et al., 1988), the authors concluded that “Although chamber effects on yield are common, there are no results showing that this will result in a changed yield response to O₃.” A more recent study of chamber effects examined the responses of tolerant and sensitive white clover clones to ambient O₃ in greenhouse, open-top, and ambient plots (Heagle et al., 1996). For individual harvests, O₃ reduced the forage weight of the sensitive clone 7 to 23% more in the greenhouse than in OTCs. However, the response in OTCs was the same as in ambient plots. Several studies have shown very similar yield response to O₃ for plants grown in pots or in the ground, suggesting that even such a significant change in environment does not alter the proportional response to O₃, at least as long as the plants are well watered (Heagle, 1979; Heagle et al., 1983).

1 Most experiments investigating the O₃ effects on annual vegetation provide adequate water
2 to avoid substantive drought stress. Because drought stress has generally been shown to reduce
3 the effect of O₃ on annual vegetation, such experiments may tend to overestimate O₃ effects on
4 crops and especially on unmanaged or seminatural vegetation.

5 As mentioned above, the use of O₃ flux rather than exposure is theoretically more realistic,
6 and such an approach would also address the vertical gradient issue (Section 9.2). Recently, a
7 number of investigators have suggested that modeling O₃ flux can improve estimates of O₃
8 effects on vegetation. Models of O₃ flux can reduce the variation in the response to O₃ that is
9 sometimes observed between years in an experiment (Emberson et al., 2000a, 2000b; Fuhrer
10 et al., 1992; Grünhage et al., 1993; Grünhage and Haenel, 1997; Pleijel et al., 2000a). In a study
11 of O₃ deposition to an oat crop in OTCs, O₃ flux in the chamber was estimated to be up to twice
12 that in an adjacent field based on a K-theory approach and measurements of stomatal
13 conductance and environmental conditions (Pleijel et al., 1994b). These measurements were
14 made for 2 hours on 5 days when the canopy was physiologically active and wind speeds were
15 moderate. However, the O₃ flux in a chamber without plants was nearly as high as that in the
16 open field. The authors conclude that O₃ uptake in the chamber was between 100 and 200% of
17 that in the field. These models of flux have a sound biological and meteorological basis and are
18 useful for interpreting experimental data. Flux models have been successfully applied at
19 intensive study sites with detailed site-specific data on stomatal conductance and
20 micrometeorological conditions (e.g., Grünhage et al., 1993b, 1994; Fredericksen et al., 1996).
21 Yet even at a single well-studied site, different methods can provide different estimates of O₃
22 flux. For example, at a site in a vineyard in California, an evapotranspiration-based method
23 overestimated the O₃ flux as compared to an eddy covariance approach by 20 to 26% (Massman
24 and Grantz, 1995). At a site in a nearby cotton field the evapotranspiration-based approach
25 overestimated the eddy-covariance method by 8 to 38%.

26 Interest has been increasing in recent years, particularly in Europe, in using mathematically
27 tractable flux models for O₃ assessments at the regional and national scale (Emberson et al.,
28 2000a, 2000b). However, methods for scaling site-specific models of O₃ flux to large areas
29 remains an active and challenging area of research (Massman et al., 1994; Massman and Ham,
30 1994; Wesely and Hicks, 2000). Reducing uncertainties in flux estimates for areas with diverse
31 surface or terrain conditions to within ±50% requires “very careful application of dry deposition

models, some model development, and support by experimental observations” (Wesely and Hicks, 2000). As an example, the annual average deposition velocity of O₃ among three nearby sites in similar vegetation was found to vary by $\pm 10\%$, presumably due to terrain (Brook et al., 1997). Moreover, the authors stated that the actual variation was even greater, because stomatal uptake was unrealistically assumed to be the same among all sites, and flux is strongly influenced by stomatal conductance (Brook et al., 1997). Stomatal conductance is affected by factors such as temperature, vapor pressure deficit, plant water status, and temperature. It is a challenging task to obtain such data for regional, national, or continental flux modeling (Pleijel, 1998). This topic is addressed further in Section 9.5.

If a flux approach is not used, another issue in scaling from experimental data to national assessments is the selection of an O₃-exposure index. As discussed in Section 9.5, there is no evidence to support a single value for defining a cutoff value for calculating a peak-weighted O₃ index. In Europe, it has become common practice to use a cutoff value of 40 ppb, as in the AOT40 index; while in the United States, a cutoff value of 60 ppb has been used, as in the SUM06 index. The W126 index has been promoted, because it has a continuous weighting function without an arbitrary cutoff value; but few exposure-response studies have used this index to date. There is evidence from some European studies that a cutoff value either lower (Pleijel et al., 1997) or higher (Finnan et al., 1996, 1997) may provide a better statistical fit to experimental data. However, the choice of the type of weighting depends on the type of response model that is fit. In an evaluation of numerous indices fit to seven studies of the response of spring wheat to O₃ in OTCs in Northern Europe, different types of indices performed better for linear response models than for Wiebull response models (Finnan et al., 1997). In this study, cumulative indices that give greater weight to higher O₃ concentrations performed best; and, while the AOT40 was not the best index, it did perform well, and slightly better than the SUM06 in conjunction with a linear model (Finnan et al., 1997). Although this section is focused on annual plants, the issue of cutoff values applies to perennial species as well; therefore, two examples of tree species will be discussed here. Aspen seedlings were exposed to O₃ in controlled environment chambers for 4 weeks, and a significantly greater degree of foliar symptoms occurred with a pattern of variable peak exposures compared to a constant peak exposure with the same total SUM06 exposure (Yun and Laurence, 1999b). However, a field experiment in large OTCs in Ithaca, NY with these two clones exposed to ambient, 1.7 \times , - and

1 3×-ambient O₃ concentrations (SUM06 = 1, 20, and 62 ppb) found SUM06 to be highly
2 correlated with foliar symptoms and biomass effects (Yun and Laurence, 1999a). When silver
3 birch was exposed to different patterns of O₃ with the same AOT40 value in growth chambers,
4 high peak concentrations and exposure shape were found to be important for symptom
5 production (Oksanen and Holopainen, 2001). However, growth reductions were best predicted
6 by total cumulative exposure.

7 An investigation of different exposure indices was conducted using data from experiments
8 conducted during 1993 at eight sites in the eastern United States in which O₃ sensitive and
9 O₃-tolerant white clover genotypes were grown using methods developed by Heagle et al.
10 (1994). A statistical approach based on profile likelihoods was used to estimate parameters in
11 generalized exposure indices similar to the SUM06 and AOT40 indices (Blankenship and
12 Stefanski, 2001). The results showed that for the SUMX family of indices, where X is a cutoff
13 value, hourly O₃ concentrations over ~71 ppb contribute the most to yield prediction. For the
14 AOTX family of indices, the parameter was 54.4. These values are similar to those used in the
15 SUM06 and AOT40 indices already in use. Furthermore, investigation of weighting for time of
16 day confirmed the importance of the mid-afternoon hours for this data set, unlike the results
17 found for wheat in Sweden (Danielsson et al., 2003; Pleijel et al., 2000a).

18 19 **9.6.4.6 European Critical Levels**

20 During the late 1980s and 1990s, substantial effort was expended in Europe to quantify the
21 effects of O₃ on crops and trees. A focus for this research effort was to develop and refine
22 “critical levels” for O₃ under the auspices of the United Nations Economic Commission for
23 Europe (UNECE) Convention on Long-Range Transboundary Air Pollution (Sanders et al.,
24 1995). A critical level was defined as “the concentration of pollutant in the atmosphere above
25 which direct adverse effects on receptors, such as plants, ecosystems or materials, may occur
26 according to present knowledge” (United Nations Economic Commission for Europe, 1988).
27 Areas where the levels were exceeded were identified to plan abatement strategies. Critical
28 levels are not air quality standards as are used in the United States for criteria pollutants, nor are
29 they legally enforced. However, they have been used successfully for planning reductions in
30 sulfur and nitrogen pollution. Critical levels for O₃ are intended to prevent long-term deleterious
31 effects on the most sensitive plant species under the most sensitive environmental conditions, but

1 not to quantify O₃ effects. The nature of the “significant harmful effects” is not specified in the
2 original definition, which allows different levels to be defined for different effects, such as foliar
3 symptoms or reduction in crop yield. Different levels also have been set for crops, forests and
4 seminatural vegetation. The caveat, “according to present knowledge” is important, because
5 critical levels are revised periodically as new scientific information becomes available.

6 Because these critical levels were designed for use in regional modeling and mapping, they
7 were required to be of a simple form (Sanders et al., 1995). During the early 1990s, the AOT40
8 index was selected as the form of the critical level, and Level I critical levels were determined
9 for arable crops and forestry. The AOT40 index is defined as the sum of the difference between
10 the hourly concentration (in ppb) and 40 ppb when the concentration exceeds 40 ppb for the
11 hours when global radiation exceeds 50 W m⁻² (Fuhrer et al., 1997). In regression analysis of
12 15 OTC studies of spring wheat, including one study from the United States and 14 from
13 locations ranging from southern Sweden to Switzerland, an AOT40 value of 5.7 ppm·h was
14 found to correspond to a 10% yield loss, and a value of 2.8 ppm·h corresponded to a 5% yield
15 loss (Fuhrer et al., 1997). Because a 4 to 5% yield loss could be detected with a confidence level
16 of 99%, a critical level of 3 ppm·h was selected an AOT40 value of in 1996 (Kärenlampi and
17 Skärby, 1996). This critical level is defined for a 3-month period calculated for daylight hours.
18 It was suggested that a 5-year average should be used to assess the exceedance of this critical
19 level, because this value was associated with a small yield decrement (Fuhrer et al., 1997). This
20 value is currently used for all crops, because it is the best-supported value and because the
21 limited data from other crop species do not provide strong evidence that a more stringent value is
22 required (Fuhrer et al., 1997).

23 The Level I critical levels were successfully used in the 1990s to map areas of exceedance,
24 but the research led to the conclusion that simple, exposure-based levels lead to overestimation
25 of the effects in some regions and underestimation in others. The main problem is that other
26 environmental factors (e.g., vapor pressure deficit, water stress, temperature, light) alter O₃
27 uptake and its effects. Therefore, the decision was made to work towards a flux-based approach,
28 aiming to be able to model O₃ flux-effect relationships for the three vegetation types (i.e., crops,
29 forests, seminatural vegetation). Progress has been made in modelling flux under experimental
30 conditions (Danielsson et al., 2003; Grünhage et al., 2001; Grünhage and Jäger, 2003; Pleijel

et al., 2000a), but because the UNECE meeting in November 2002 concluded that more research is needed, the subject is to be reassessed in 2 to 5 years' time.

There has been criticism that the Level I critical level for crops overestimates O₃ effects in the Mediterranean countries, because it was developed based on studies in Northern Europe (De Santis, 1999, 2000). However, there is evidence of substantial crop loss due to O₃ in some southern European counties, such as the Po valley in Northern Italy (Fumagalli et al., 2001). In these studies Heagle type OTCs were used. Losses in NF chambers as compared to CF chambers over several years at two sites ranged from 11.2 to 22.8% for barley and wheat, from 0.3 to 31.5% for other crop species, and from 4.1 to 19.8% for forage species (Fumagalli et al., 2001). Surprisingly, the least effect was observed for soybean, despite AOT40 values of 9.32 ppm·h, 3× the Level I critical level. Similarly, a review of studies in Northern Italy found that ambient O₃ episodes have been reported to cause foliar symptoms on 24 agricultural and horticultural crops in commercial fields (Fumagalli et al., 2001). Ambient O₃ has also been reported to cause yield losses in several crop species, although no data on O₃ exposure were presented by the reviewers (Fumagalli et al., 2001).

The Level I approach has also been criticized for focusing only on a single annual species (wheat) and a single woody perennial species (beech). However, this species focus is appropriate, because the goal was to determine an exposure-response relationship for a sensitive species based on available data. In support of standards in Germany, an effort was made to combine data from different species, and consisted of a meta-analysis of studies conducted in both closed chambers and OTCs (Grünhage et al., 2001). In this study, experiments published between 1989 and 1999 were analyzed if they met the following conditions: (1) a significant O₃ effect was determined (2) exposure conditions were well defined, (3) foliar symptoms, growth, or yield was measured, and (4) plant species were relevant to Europe (Grünhage et al., 2001). Despite the focus on European species, many of the species studies also occur in the United States. Separate regressions for herbaceous plants and for tree species were created as a function of duration of exposure at a given level of O₃ exposure at the top of the plant canopy. These regression equations, with confidence limits and with correction for the vertical gradient in O₃ from the top of the quasi-laminar boundary layer, can be used to define whether effects are unlikely (below the lower confidence limit), probable (between the confidence limits), or highly likely (above the upper confidence limit) to occur near a given O₃-monitoring station.

1 A further concern about the Level 1 approach is that foliar symptoms, rather than biomass,
2 may be an important endpoint, because foliar symptoms may be more sensitive than is biomass
3 increment (VanderHeyden et al., 2001). In an OTC study in southern Switzerland, it was shown
4 that a number of tree species show foliar symptoms at AOT40 values lower than the Level 1
5 value of 10 ppm·h (VanderHeyden et al., 2001).

6 7 **9.6.4.7 Summary of Effects on Short-Lived Species**

8 For annual vegetation, the data summarized in Table 9-17 show a range of growth and
9 yield responses both within species and among species. Nearly all of these data were derived
10 from studies in OTCs, with only two studies using open-air systems in the UK (Ollerenshaw
11 et al., 1999; Ollerenshaw and Lyons, 1999). It is difficult to compare studies that report O₃
12 exposure in different indices such as AOT40, SUM06, or 7-h or 12-h mean values. However,
13 when such comparisons can be made, the results of this recent body of research confirm earlier
14 results summarized in the 1995 AQCD (U.S. Environmental Protection Agency, 1996).

15 A summary of earlier literature concluded that a 7-h, 3-month mean of 49 ppb corresponding to a
16 SUM06 exposure of 24.4 ppm·h would cause 10% loss in 50% of 49 experimental cases (Tingey
17 et al., 1991). A similar study using a 24-h, rather than 7-h, averaging period found that a
18 SUM06 exposure of 26.4 ppb would cause 10% loss in 50% of 54 experimental cases (Lee et al.,
19 1994a, 1994b). Recent data summarized in Table 9-17 support this conclusion. These values
20 represent ambient exposure patterns that occur in some years over large portions of the United
21 States. Some annual species such as soybean are more sensitive, and greater losses may be
22 expected (Table 9-17). Thus, the recent scientific literature supports the conclusions of the 1996
23 AQCD that ambient O₃ concentrations are reducing the yield of major crops in the United States.

24 Much research in Europe has used the AOT40 exposure statistic, and substantial effort has
25 gone into developing Level-1 values for vegetation. Based on regression analysis of 15 OTC
26 studies of spring wheat, including one study from the United States and 14 from locations
27 ranging from southern Sweden to Switzerland, an AOT40 value of 5.7 ppm·h was found to
28 correspond to a 10% yield loss, and a value of 2.8 ppm·h corresponded to a 5% yield loss
29 (Fuhrer et al., 1997). Because a 4 to 5% decrease could be detected with a confidence level of
30 99%, a critical level of an AOT40 value of 3 ppm·h was selected in 1996 (Kärenlampi and
31 Skärby, 1996).

In addition to reductions in crop yield, O₃ may also reduce the quality or nutritive value of annual species. Many studies have shown effects of O₃ on various measures of plant organs that affect quality, with most studies focusing on characteristics important for food or fodder. These studies indicate that there may be economically important effects of ambient O₃ on the quality of crop and forage species. Previous criteria documents have concluded that foliar symptoms on marketable portions of crops and ornamental plants can occur with seasonal 7-h mean O₃ exposures of 40 to 100 ppb (U.S. Environmental Protection Agency, 1978, 1986, 1996). The recent scientific literature does not refute this conclusion.

The use of OTCs may reverse the usual vertical gradient in O₃ that occurs within a few meters above the ground surface (Section 9.2). Such a reversal suggests that OTC studies may overestimate, to some degree, the effects of an O₃ concentration measured several meters above the ground. However such considerations do not invalidate the conclusion of the 1996 AQCD (U.S. Environmental Protection Agency, 1996) that ambient (Table 9-14, Table 9-21) O₃ concentrations are sufficient to reduce the yield of major crops in the United States.

Table 9-21. Ozone Exposures at 35 Rural Sites in the Clean Air Status and Trends Network in the Central and Eastern United States From 1989 to 1995

Subregion	SUM06 12-h, 3-Month Mean	UM06 12-h, 3-Month SD	W126 3-Month Mean	W126 3-Month SD	Max. 8h > 80 ppb(n) Mean	Max. 8h > 80 ppb(n) SD
Midwest	31.5	10.2	25.1	7.7	13.8	10.6
Upper Midwest	18.9	8.5	16.0	5.9	5.6	5.6
Northeast	33.2	11.9	26.6	9.5	15.8	12.2
Upper Northeast	13.2	8.6	12.8	6.5	3.3	5.5
South Central	34.5	16.6	25.6	11.5	7.1	10.0
Southern Periphery	19.2	7.6	15.2	5.4	1.9	1.6

Units for SUM06 and W126 are ppm-h.
Source: Baumgardner and Edgerton (1998).

9.6.5 Effects of Ozone on Long-Lived (Perennial) Species

Although there has been considerable research in Europe on annual species during the past 10 years, much recent research in the United States has focused on perennial species. In Europe, and in a few studies in the United States, effects of O₃ on mixtures of annual and perennial herbaceous species have been investigated using growth chambers, greenhouses, and OTCs. Section 9.6.5.1 reviews such studies, with an emphasis on studies using OTCs.

9.6.5.1 Herbaceous Perennial Species

Two alfalfa cultivars were grown in pots and exposed to CF, NF, 1.5×-ambient and 2×-ambient O₃ concentrations in two 1-year studies in Quebec, Canada (Renaud et al., 1997). One cultivar, *Apica*, is commonly grown in the region, and another, *Team*, is normally grown farther south and is more tolerant to O₃. For *Apica* in both years and for *Team* in 1991, O₃ exposure caused a linear reduction in above-ground biomass. In the NF treatment, growth of *Apica* was decreased by 31 and 21% in the 2 years, while the growth of *Team* was reduced by 14% in 1991, but not reduced in 1992. The authors suggested that the differing effects on *Team* could be due to different progenies and propagation methods in the 2 years or to more rapid growth in 1991 along with higher O₃ peak values in 1991. In 1991, O₃ maxima exceeded 60 ppb in 15 days, whereas in 1992 there were only three such days. At the end of the growing season, total starch reserves in roots were decreased by O₃ due primarily to a decrease in root mass, which the authors suggested could accelerate decline in alfalfa yields. These yield reductions are generally similar to those reported previously in five similar studies summarized in the 1996 AQCD (Table 9-25 of U.S. Environmental Protection Agency, 1996).

A study in Alabama exposed early- and late-season-planted bahia grass (cultivar Pensacola) in OTCs to CF, NF or 2×-ambient O₃ treatments (Muntiferi et al., 2000). Ozone exposures expressed as 12-h mean values over the 24-week experiment were 22, 45, and 91 ppb, and the highest ambient O₃ concentrations were recorded in late June, late July, late August and mid-September at approximately 90 ppb. Above-ground biomass growth was reduced by the NF treatment for the first and second harvest by 34% and 29% for the early-season planting, but statistically significant effects were not observed in the late-season planting (Table 9-18). The 2×-ambient treatment did not cause further significant reductions in biomass. The authors suggested that the lack of a significant O₃ effect in the late planting may have been due to the

1 shorter total O₃ exposure time as well as to the lower O₃-exposure concentrations during the
2 weeks immediately preceding harvest. These results are important, because this is an
3 economically important species and because previous studies have focused on grass species that
4 use the C₃, rather than C₄, metabolic pathway.

5 An investigation of the use of different O₃ indices and averaging times on the correlation
6 with growth effects was undertaken with the NC clover system (Heagle et al., 1995). For 2 years
7 of data at six sites in Massachusetts, Oregon, North Carolina, California (2 sites), and Virginia,
8 averaging time was found to be more important than the choice of the type of index including
9 mean, SUM06, and AOT40 (Heagle and Stefanski, 2000). The best correlation between O₃
10 exposure and the ratio of sensitive to tolerant clover types was found for the 6-h period from
11 1000 to 1600 h. For this period, very similar r² values (0.91 to 0.94) were found for SUM06,
12 W126, and AOT40 (Heagle and Stefanski, 2000). For the above indices, a linear relationship
13 was found, with no effect in Corvallis, OR with exposure to a SUM06 value of 10.2 ppm·h and a
14 ratio of 0.53 (sensitive/tolerant) at San Bernardino with a SUM06 exposure of 39.4 ppm·h
15 (Heagle and Stefanski, 2000).

16 In a study of the biomass ratio of O₃-sensitive versus O₃-insensitive clover at 14 sites in
17 Europe during 1996 to 1998, a model that was developed using artificial neural network (ANN,
18 see Section 9.2) techniques had r² values for the training data of 0.84 and for unseen validation
19 data of 0.71 (Mills et al., 2000). The predictive factors in the model were AOT40, 24-h mean
20 O₃, daylight mean temperature, and 24-h mean temperature. This model was selected after a
21 thorough investigation of a number of models using many more or fewer parameters using both
22 ANN and multiple linear regression techniques. This model predicted that a 5% reduction in
23 biomass ratio was associated with AOT40 values in the range 0.9 to 1.7 ppm·h accumulated over
24 28 days, with plants being most sensitive under conditions of low NO_x, moderate temperature,
25 and high 24-h mean O₃ concentration.

26 Two experiments in Ontario investigated effects of square-wave additions of O₃ for 7 h on
27 1 day/week for 7 weeks on the growth and fruit production of two cultivars of primocane (first-
28 year fruiting) raspberry plants (Sullivan et al., 1994). Plants were grown in OTCs and exposed
29 to ambient O₃ (mean values of ≤ 4 ppb). In the both experiments, additions of 12 ppb O₃ on
30 1 day/week for 7 weeks did not significantly reduce fruit yield. In the second experiment,

1 additions of 24 ppb significantly reduced the fruit yield 52% below that of the ambient treatment
2 in one cultivar (Heritage), but not in another (Redwing).

3 Exposure to a square-wave 8-h mean O₃ concentration of 92 ppb for 62 days in an
4 experiment in OTCs the UK did not significantly reduce the total yield of strawberry fruits, but
5 did decrease the average size of the fruits by 14% (Drogoudi and Ashmore, 2000). This
6 contrasts with an increase in total yield (fruit weight) found in a previous study in California
7 (Takemoto et al., 1988).

8 When timothy was exposed in OTCs in Sweden to NF, CF, and CF+O₃ treatments, there
9 was no effect of a 12-h mean O₃ exposure of 68 ppb (NF treatment), but a 12-h mean exposure
10 of 152 ppb decreased yield by 58% (Danielsson et al., 1999). A similar lack of effect of
11 exposure to a 12-h mean O₃ exposure of 62 ppb was found in a previous study in the United
12 States (Kohut et al., 1988).

13 Although most investigations of O₃-response relationships focus on growth or yield of
14 marketable portions of plants, some studies also investigate effects on plant quality. In the study
15 of bahia grass in Alabama discussed above, in addition to the effects on yield, there were
16 significant effects on quality for ruminant nutrition (Muntifering et al., 2000). Concentrations of
17 neutral detergent fiber (NDF) were higher in primary-growth and regrowth forages from the
18 early-season planting when exposed to 2×-ambient O₃ than when exposed to the NF treatment.
19 The concentration of acid detergent fiber was higher in the 2×-ambient treatment than in NF
20 treatment regrowth, whereas acid detergent lignin concentration was higher in 2×-ambient than
21 in NF primary-growth forage. Crude protein concentrations were lower in CF-exposed than in
22 NF-exposed regrowth forage from the early planting and in CF- than in NF-exposed primary-
23 growth forage from the initial harvest of the late-season planting. No differences were observed
24 among treatments in concentrations of total phenolics in primary-growth or regrowth forages
25 from either planting, although concentrations of total phenolics tended to be higher in
26 CF-exposed than in NF-exposed primary-growth forage from the late-season planting. The
27 authors concluded that the alterations in quality of primary-growth and vegetative regrowth
28 forages were of sufficient magnitude to have nutritional and possibly economic implications to
29 their use for ruminant animal feed.

30 *Sericea lespedeza* and little bluestem were exposed to CF, NF, and 2×-ambient O₃ in OTCs
31 in Alabama for 10 weeks (Powell et al., 2003). Ozone treatments expressed as 12-h mean

1 concentrations were 23, 40, and 83 ppb and expressed as seasonal SUM06 values were 0.2, 9.1,
2 and 61.0 ppm·h. Although there were few statistically significant effects of O₃ on yield (the
3 yield of only the 2×-ambient compared to NF for *Sericea lespedeza* in the last of six harvests),
4 plant quality as feed for ruminants was reduced. The nutritive quality of *Sericea lespedeza* was
5 decreased by 7% and that of little bluestem by 2% as a result of increased cell wall constituents
6 and decreased in vitro digestibility.

7 For some annual species, particularly crops, the endpoint for an assessment of the risk of
8 O₃ exposure can be defined as yield or growth; e.g., production of grain. For plants grown in
9 mixtures such as hayfields, and natural or seminatural grasslands (including native
10 nonagricultural species), endpoints other than production of biomass may be important. Such
11 endpoints include biodiversity or species composition and measures of plant quality such as total
12 protein, and effects may result from competitive interactions among plants in mixed-species
13 communities. Most of the available data on non-crop herbaceous species are for grasslands.

14 In a study of two perennial grasses (bent grass and red fescue) and two forbs (white clover,
15 Germander speedwell) grown in pots in OTCs, O₃ effects differed among species and cutting
16 treatments (Ashmore and Ainsworth, 1995) (see also Table 9-18). Fescue biomass increased
17 with higher O₃ treatments both in pots that were not cut during the growing season (mid-June to
18 mid-September), and those that were cut every two weeks. However, bent grass biomass
19 decreased with higher O₃ exposure in the uncut treatment and increased in the cut treatment.
20 White clover and Germander speedwell biomass decreased substantially with higher O₃ exposure
21 with and without cutting, with greater decreases in the cut treatment. The authors cautioned that
22 the experiment did not replicate field circumstances. The plants were all cut to 1 cm above the
23 ground, which does not simulate grazing, and there may have been effects due to growing the
24 plants in pots. However, two key results of this study likely apply to mixtures of species
25 growing in hay or forage fields or seminatural and natural communities. First, O₃ exposure
26 increased the growth of O₃-tolerant species while exacerbating the growth decrease of O₃
27 sensitive species. Second, the total biomass of the mixed-species community was unaffected by
28 O₃ exposure due to the differential effects on O₃-sensitive and O₃-tolerant species.

29 In a 2-year study using OTCs placed over managed pasture in Switzerland, the above-
30 ground biomass of clover (red and white) was reduced linearly in response to increased O₃
31 exposure (Table 9-18) (Fuhrer et al., 1994). Exposure to a 12-h mean concentration of 39 ppb

O₃ reduced biomass by 24% as compared to the CF treatment with a 12-h mean concentration of 21 ppb O₃. There was a trend toward increased above-ground biomass of grasses (primarily orchard grass), but this trend was not statistically significant. As often found in other studies of mixtures of species, by O₃ exposure did not significantly affect total above-ground biomass O₃ exposure.

A field-grown grass/clover mixture was exposed to CF, NF, and two O₃ addition treatments for two growing seasons in OTCs in southern Sweden (Pleijel et al., 1996). The mixture consisted of 15% (by seed weight) red clover cv. Fanny, 60% timothy cv. Alexander, and 25% fescue cv. Svalofs Sena.. Ozone concentrations expressed as AOT40 ranged from 0 to approximately 47 ppm·h and expressed as 7-h mean from 11 to 62 ppb. Over this range, a slight, but statistically significant, linear decrease of 4% in total above-ground biomass was seen growth over six harvests. No significant decrease was seen in the proportion of clover, and the authors ascribed this lack of effect to the relatively higher O₃ sensitivity of timothy and lower sensitivity of this clover cultivar as compared to previously published results for other grass/clover mixtures (e.g., Fuhrer et al., 1994).

A mixture of species in an old farm field in Alabama was exposed to O₃ for two growing seasons in large OTCs (4.8 m high, and 4.5 m diameter); and a similar lack of effect of O₃ was found on total plant community growth measured as both canopy cover and vertical canopy density (Barbo et al., 1998). Of the 40 species in the plots, O₃ effects were examined only on the most common species: blackberry, broomsedge bluestem, bahia grass, *Panicum* spp., and winged sumac (second year only). Of these species, a 2×-ambient O₃ treatment increased the percent canopy cover of blackberry over 2 years by 124%, while that of winged sumac was decreased by 95% (Table 9-18). Blackberry showed no significant effect on biomass, but ripe fruit mass was decreased by 28% (Chappelka, 2002). However, there was a significant chamber effect for this latter response. Biomass was not reported for other species in this study due to a hurricane. Effects on loblolly pine grown in this experiment are discussed subsequently in Section 9.6.5.5.

In summary, results of studies on perennial herbaceous species conducted since the 1996 criteria document was prepared are presented in Table 9-18. As for single-season agricultural crops, yields of multiple-year forage crops are reduced at O₃ exposures that occur in some years over large areas of the United States (Table 9-14, Table 9-21). This result confirms that reported

1 in the 1996 AQCD (U.S. Environmental Protection Agency, 1996). When species are grown in
2 mixtures, O₃ exposure can increase the growth of O₃-tolerant species while exacerbating the
3 growth decrease of O₃-sensitive species (e.g., Ashmore and Ainsworth, 1995; Fuhrer et al.,
4 1994). Because of this competitive interaction, the total growth of the mixed-species community
5 may not be affected by O₃ exposure (Ashmore and Ainsworth, 1995; Barbo et al., 1998; Fuhrer
6 et al., 1994). However, in some cases mixtures of grasses and clover species have shown
7 significant decreases in total biomass growth in response to O₃ exposure in studies in the United
8 States (Heagle et al., 1989; Kohut et al., 1988) and in Sweden (Pleijel et al., 1996). In Europe,
9 a provisional critical level for perennials of an AOT40 value of 7 ppm·h over 6 months has been
10 proposed to protect sensitive plant species from the adverse effects of O₃.

12 **9.6.5.2 Deciduous Woody Species**

13 It is extremely difficult and costly to study entire mature trees under controlled conditions
14 such as those in OTCs, with the possible exception of some species managed for fruit or nut
15 production. For this reason, the great majority of investigations have been of seedlings in
16 growth chambers, greenhouses, or OTCs. A few investigations have been carried out on
17 saplings or more mature trees using free air exposure systems (Haeberle et al., 1999; Isebrands
18 et al., 2000, 2001; Werner and Fabian, 2002). Exposure-response functions based on 28
19 experimental cases of seedling response to O₃ suggest that a SUM06 exposure for 3 months of
20 31.5 ppm·h would protect hardwoods from a 10% growth loss in 50% of the cases (Table 9-19).
21 However, there is a substantial range in sensitivity among species. A risk analysis was
22 undertaken to predict tree biomass growth reductions due to O₃ based on exposure-response
23 equations for seedlings of individual species combined with the species' spatial distribution
24 across the eastern United States and interpolated O₃ exposure expressed as SUM06 (Hogsett
25 et al., 1997). The growth of sensitive species such as aspen and black cherry was predicted to be
26 reduced by at least 20% across 50% of their ranges in a high O₃ year and approximately 10% in a
27 lower-than-average O₃ year (Hogsett et al., 1997).

28 A few investigations reported since the last criteria document was prepared have examined
29 saplings or mature trees, notably of oak species in the southern Appalachian Mountains and pine
30 species in California. Most of these studies have been of natural (uncontrolled) O₃ exposures.
31 Additional studies have examined foliar symptoms on mature trees, and in recent years such

surveys have become more common and with greater attention to the standardization of methods and the use of reliable indicator species (Campbell et al., 2000; Smith et al., 2003). Previous criteria documents have noted the difficulty in relating foliar symptoms to effects on individual tree growth, stand growth, or ecosystem characteristics (U.S. Environmental Protection Agency, 1996). This difficulty still remains to the present day.

Some investigators have suggested that a comprehensive risk assessment of the effects of O₃ on mature tree species might best be accomplished by extrapolating measured effects of O₃ on seedlings to effects on forests using models based on tree physiology and forest stand dynamics (Fuhrer et al., 1997; Hogsett et al., 1997; Chappelka, 1998; Laurence et al., 2000, 2001). Several such efforts are discussed in Sections 9.4 and 9.7.

In this subsection, emphasis will be placed on experimental evidence of O₃ effects on the growth of woody species under controlled conditions with some information from observational studies under ambient conditions in forests. Experimental results are summarized for deciduous species in Table 9-20; and the species are discussed below in the order in which they appear in this table.

A series of studies out in Michigan and Wisconsin during the 1990s on clones of trembling aspen previously demonstrated that they differ in their O₃ sensitivity (Coleman et al., 1995b, 1995a, 1996; Dickson et al., 2001; Isebrands et al., 2000, 2001; Karnosky et al., 1996, 1998, 1999; King et al., 2001). Several of those studies were undertaken with plants in pots or in the ground in OTCs and additional studies were undertaken at three sites selected to differ primarily in O₃ exposure (Karnosky et al., 1999). An ongoing study was undertaken using a free air carbon dioxide and O₃ enrichment (FACE) facility in Rhinelander, WI (Isebrands et al., 2000, 2001). These studies showed that O₃-symptom expression was generally similar in OTCs and FACE and gradient sites, supporting the previously observed variation among aspen clones (Karnosky et al., 1999). In the Michigan OTC studies, O₃ decreased total plant biomass by an average of 18% in each of 2 years for three clones classified as high, intermediate and low in O₃ tolerance (clones 216, 271 and 259; Karnosky et al. [1996]). However, in the first experiment, an additional clone known to be O₃ insensitive showed no response (clone 253). In these studies, plants were grown in pots and exposed to CF, 1×-ambient, and 2×-ambient O₃ treatments in two separate experiments of 98 days each (additional treatments of 0.5×-ambient and 1.5×-ambient were used in the first experiment only (Karnosky et al., 1996). Ozone concentrations expressed

as 3-month, 7-h mean values for 1990 ranged from 7 to 69 ppb for the CF to 2×-ambient treatments; while for 1991, values ranged from 22 to 92 ppb.

The FACE study evaluated the effects of 3 years of exposure to combinations of elevated CO₂ and O₃ on growth responses in mixture of five trembling aspen clones (Isebrands et al., 2000, 2001). Height, diameter, and stem volume (diameter² × height) were decreased by elevated O₃. On average for all clones, stem volume was decreased by 20% over the 3 years in the elevated O₃ treatment as compared with the 1×-ambient treatment. However, one clone showed increased growth in response to O₃. Ozone concentrations were not reported. This FACE facility study is important, because it confirmed responses reported previously with these clones grown in pots or soil in OTCs, without the alterations of microclimate induced by chambers. Currently, this is the only U.S. study using this technology to have examined the effects of O₃ under these conditions. This study is also significant, because the elevated O₃-exposure pattern used was intended to reproduce the 6-year average pattern from Washtenaw County, Michigan (Karnosky et al., 1999).

Rooted cuttings of two aspen clones from Acadia National Park in Maine were exposed to 1×-ambient, 1.7×-ambient, and 3×-ambient O₃ concentrations in large OTCs in Ithaca, NY for much of one growing season (15 June to 15 September) (Yun and Laurence, 1999a). Both circular (4.7 m diameter, 3.7 m height) and rectangular (7.4 m × 2.75 m × 3.7 m height) chambers were used (Mandl et al., 1989). Exposure to 1.7×-ambient O₃ (SUM06 = 20 ppm·h, 9-h mean = 74 ppb) reduced shoot growth by 14 and 25% compared to ambient O₃ for the two clones (Yun and Laurence, 1999a). Total dry weight was reduced by 55 and 35% in the two clones by the 3×-ambient treatment (SUM06 = 62 ppm·h, 9-h mean = 124 ppb) compared to the ambient O₃ treatment.

When black poplar cuttings in OTCs in Belgium were exposed to 8-h mean O₃ concentrations of 5, 29, and 33 ppb, diameter growth decreased by 29% in the highest O₃ treatment, but height growth was unaffected (Bortier et al., 2000b). A 2-month study of hybrid poplar (*Populus tremuloides* × *P. tremula*) in a free air exposure system in Finland with 7-h mean O₃ concentrations of 30 and 38 ppb found a 6% decrease in height with no effect on biomass (Oksanen et al., 2001). Eastern cottonwood cuttings in pots buried in the ground with drip irrigation were exposed to ambient O₃ at several sites in and near New York City in three 2-month experiments during three summers (Gregg et al., 2003). Ozone concentrations were

1 lower at urban sites than at rural sites within 100 km of the urban sites. Total biomass growth
2 was greater in urban than rural sites, with a strong linear decrease in biomass with increasing O₃
3 across all sites and years ($r^2 = 93$). Total biomass decreased 33% with 12-h mean O₃ levels of
4 38 ppb compared to 23 ppb. Multiple regression analysis showed no significant temperature
5 effect on biomass; therefore, the authors suggested that O₃ exposures was the most likely
6 explanation for the reduced biomass in rural areas. The overall growth reductions and the
7 variation among genotypes seen on all of the above aspen and poplar studies is similar to those
8 previously reported in three OTCs studies summarized in the previous criteria document
9 (Table 9-26 of U.S. Environmental Protection Agency, 1996).

10 Black cherry seedlings grown in pots were exposed in OTCs in the Great Smoky Mountain
11 National Park in Tennessee to O₃ treatments ranging from CF to 2×-ambient in two experiments
12 during 1989 and 1992 (Neufeld et al., 1995). Ozone exposure, expressed as SUM06, ranged
13 from 0 to 40.6 ppm·h in 1989, and from 0 to 53.7 ppm·h in 1992. Corresponding AOT40 values
14 were 0.0 to 28.3 ppm·h in 1989, and 0 to 40.4 ppm·h in 1992. In 1989, total biomass was
15 decreased in the 1.5×-ambient treatment by 18% and in the 2×-ambient treatment by 38%.
16 In 1992, total biomass was decreased in the 1.5×-ambient treatment by 27%, and in the
17 2×-ambient treatment by 59%. In this study, SUM06 and AOT40 provided better fits than did
18 SUM00 with Weibull regression analyses to log-transformed biomass data. Although a Weibull
19 model was used, responses to O₃ expressed as SUM06 and AOT40 appeared to be linear. The O₃
20 exposures in the 1.5×-ambient and 2×-ambient treatments were reported to be similar to those
21 for a site near Charlotte, NC in a high-O₃ year (1988). In a 2-year experiment in OTCs in Ohio,
22 seedlings of black cherry, sugar maple, and yellow poplar were exposed to O₃ treatments with
23 SUM00 values ranging from 16 to 107 ppm·h in 1990 and 31 to 197 ppm·h in 1991 (Rebbeck,
24 1996). After two seasons of exposure, only black cherry showed a growth decrease: total
25 biomass was reduced by 32% in the 2×-ambient O₃ treatment compared to the CF treatment; root
26 biomass was decreased by 39%. These results contrast with those of a previous study with black
27 cherry seedlings in which significant biomass reductions with exposures up to 2×-ambient were
28 not observed (7-h mean = 21 to 97 ppb), perhaps because of the small sample size (3 seedlings
29 per chamber (Samuelson, 1994) in the earlier study.

30 A multiyear study of effects of O₃ on both seedling and mature (30-year-old) red oak trees
31 was conducted in Norris, TN in large OTCs with three replicates per O₃ treatment. Trees were

1 exposed for 3 years to CF, 1×-ambient, and 2×-ambient treatments, with the following
2 O₃ exposures: SUM06 for 3 years = 0, 29, 326 ppm·h; SUM00 for 3 years = 147, 255, and
3 507 ppm·h. The net photosynthetic rate in mature trees was reduced by 25% in the ambient
4 treatment and by 50% in the 2×-ambient treatment (Hanson et al., 1994; Samuelson and
5 Edwards, 1993; Wullschleger et al., 1996). Despite these large decreases, no significant effects
6 on stem increment at the base, stem increment in the canopy, or leaf mass were observed for the
7 mature trees (Samuelson et al., 1996). Similarly, seedling biomass was not significantly reduced
8 by O₃ exposure. The difficulty in replicating experiments with mature trees makes it difficult to
9 detect changes in growth or biomass. However, the mean values of the stem increment at the
10 base and within the canopy in the ambient treatment were larger than those in the CF treatment,
11 although those in the 2×-ambient treatment were lower. Therefore this study of mature trees
12 does not provide evidence that these ambient concentrations reduced aboveground tree growth,
13 even after 4-years exposure.

14 Sugar maple seedlings were exposed for 3 years to ambient, 1.7×-ambient, and 3×-ambient
15 O₃ treatments at both high-light (35% of ambient) and low-light levels (15% of ambient) (Topa
16 et al., 2001). This experiment was conducted in large OTCs near Ithaca, NY. Over the 3 years,
17 O₃ exposures expressed as SUM00 for the three treatments were 88, 126, and 185 ppm·h. After
18 3 years, total seedling biomass in the 3×-ambient treatment was reduced by 64% and 41% in the
19 low- and high-light treatments, respectively (compared to the 1×-ambient treatment). The larger
20 reduction of biomass under low-light conditions suggests that seedlings growing under closed
21 canopies may be substantially more sensitive to O₃ than are seedlings exposed to higher-light
22 levels in gaps or clearings. These results differ from other studies in which seedling biomass
23 was unaffected by exposure to SUM00 values of 304 ppm·h over 2 years (Rebbeck, 1996) or
24 591 ppm·h over 3 years (daytime mean of 40.7) (Laurence et al., 1996). However the latter two
25 studies used much higher light levels, which may have reduced the O₃ effect, based on the results
26 of Topa et al. (2001).

27 Although most studies demonstrate that O₃ decreases biomass growth, occasional results
28 indicate that O₃ can increase growth of some portions of woody perennials. When Casselman
29 plum trees near Fresno, CA were exposed to O₃ in large, rectangular OTCs to three O₃ treatments
30 (CH, 1×-ambient, and an above-ambient O₃ treatment) for 4 years (12-h mean = 31, 48, 91
31 ppm·h), stem increment increased 14% in the highest O₃ treatment compared to the CH

1 treatment; and this effect was statistically significant (Retzlaff et al., 1997). However, fruit yield
2 decreased in this treatment by 42% and also decreased by 16% in the 1×-ambient-O₃ treatment.
3 Root growth was not measured in this study. Hence, the increase in stem diameter may have
4 been at the expense of other organs. However, in a fifth year, all plants were exposed to
5 1×-ambient O₃, and there were no differences in fruit yield, suggesting that trees were able to
6 recover to some extent from the effects of O₃ exposure in prior years.

7 When yellow poplar seedlings were exposed to O₃ concentrations up to SUM00 values of
8 107 ppm·h in one year and 197 ppm·h in a second year, no effects on biomass were observed
9 (Rebbeck, 1996).

10 Many studies have demonstrated that root growth is more sensitive to O₃ exposure than is
11 stem growth. For example, in a study with black cherry seedlings exposed in OTCs in
12 Tennessee in 1989, root biomass in a 2×-ambient treatment was decreased by 42%, while stem
13 biomass was decreased by only 24%. However, in a second experiment in 1992, root and stem
14 growth reductions in the 2×-ambient treatment were similar (65% versus 60%) (Neufeld et al.,
15 1995). In Finland, reduced root growth was found for a number of clones of silver birch
16 (Oksanen and Saleem, 1999). After 5 years, root growth was decreased by 33%, but shoot
17 growth was not affected by O₃ exposures of a 7-h mean of 15 ppm·h over 5 years in a FACE
18 system (Oksanen, 2001). When first-year poplar seedlings (*P. tremuloides*) were exposed in
19 OTCs to two O₃ concentrations and six N concentrations, the root/shoot ratio was decreased soon
20 after exposure to O₃ in all N treatments, even though O₃ effects on total biomass were not
21 detected in the low-N and very high-N treatments (Pell et al., 1995). These results suggest that
22 effects on the root/shoot ratio occur before significant growth effects arise. In a series of OTC
23 experiments lasting 1 to 3 years at 3 different elevations in Switzerland, fine root growth in
24 European beech was found to be more sensitive to O₃ than was shoot or total biomass (Braun and
25 Fluckiger, 1995). Although the estimated effect of O₃ on fine root biomass was similar to that
26 for total biomass, fine root biomass was significantly decreased at AOT40 (24-h) values of only
27 3 ppm·h, while total biomass was not significantly decreased until AOT40 values reached 30 to
28 40 ppm·h.

9.6.5.3 European Critical Levels

In Europe, a Level I critical level has been set for forest trees based on OTC studies of saplings. This level is discussed here because it was based on a deciduous tree species. For consistency with the approach used for crops, an AOT40 index value was selected. A few studies have shown that O₃ can be taken up by tree species at nighttime, e.g., young birch trees (Matyssek et al., 1995). However, because most evidence suggests that O₃ deposition at nighttime is low (Coe et al., 1995; Rondon et al., 1993), a value for only daylight hours was selected in Europe (Fuhrer et al., 1997; Kärenlampi and Skärby, 1996). European beech was selected for development of a Level I critical level, because data from several studies were available for this species and because deciduous tree species were judged to be more sensitive to O₃ compared to evergreen tree species (Fuhrer et al., 1997; Kärenlampi and Skärby, 1996). A critical level was defined as an AOT40 value of 10 ppm·h for daylight hours for a 6-month growing season (Kärenlampi and Skärby, 1996). However, other studies have shown that other species such as silver birch may be more sensitive to O₃ than beech (Pääkkönen et al., 1996). As discussed for annual plants above (Section 9.6.4.6), Level I critical values are not designed for making quantitative estimates of the O₃ effects on vegetation at the regional scale. For long-lived perennials, additional problems complicate extrapolation. As discussed below (Section 9.6.5.7), considerable scaling is required to extrapolate from experiments conducted with tree seedlings to estimate effects on mature trees in forests. Because of these scaling issues, there is greater uncertainty in estimating effects on forest trees than on annual plants such as crops. While some information is available for addressing issues such as scaling from seedlings to mature trees and estimating O₃ uptake, this information may still be insufficient for developing a Level II approach that can provide quantitative estimates of forest growth losses due to O₃ exposure (Broadmeadow, 1998).

9.6.5.4 Summary of Effects on Deciduous Woody Species

Recent evidence from free air exposure systems and OTCs supports results observed previously in OTC studies (Table 9-16, Figure 9-1). Specifically, a series of studies undertaken using free air O₃ enrichment in Rhinelander, WI (Isebrands et al., 2000, 2001) showed that O₃-symptom expression was generally similar in OTCs, FACE, and ambient-O₃ gradient sites, supporting the previously observed variation among aspen clones using OTCs (Karnosky et al.,

1999). New evidence is also available comparing various aspects of O₃ sensitivity between seedlings and mature trees of some species, notably red oak. As has been observed in previous criteria documents, root growth is often found to be the most sensitive indicator in terms of biomass response to O₃.

Results since 1996 support the conclusions of the 1996 AQCD (U.S. Environmental Protection Agency, 1996) that individual deciduous trees are generally less sensitive to O₃ than are most annual plants, with the exception of a few genera such as *Populus*, which are highly sensitive. However, the data presented in Table 9-19 suggest that ambient exposures that occur in different regions of the United States can sometimes reduce the growth of seedlings of deciduous species. Results from multiple-year studies sometimes show a pattern of increasing effects in subsequent years. Although, in some cases, growth decreases due to O₃ become less significant or even disappear over time. While some mature trees show greater O₃ sensitivity in physiological parameters such as net photosynthetic rate compared to seedlings, these effects may not translate into measurable reductions in biomass growth. Because even multiple-year experiments do not expose trees to O₃ for more than a small fraction of their life span and because competition may, in some cases, exacerbate the effects of O₃ on individual species, determining the effects on mature trees remains a significant challenge. Effects on mature trees under natural conditions are discussed after the review of evergreen species below and more fully in Section 9.7, in the context of extrapolating from controlled studies to forest ecosystems.

9.6.5.5 Evergreen Woody Species

Most investigations have shown evergreen tree species to be less sensitive to O₃ compared to deciduous species (U.S. Environmental Protection Agency, 1996). For example, exposure-response functions based on 23 experimental cases of seedling response to O₃, suggest that a SUM06 exposure for 3 months of 42.6 ppm·h would protect evergreen species from a 10% growth loss in 50% of the cases (Table 9-16). For deciduous species, the corresponding SUM06 value was 31.5 ppm·h (Table 9-16). As another example, experiments in the Great Smoky Mountains National Park found black cherry seedlings to demonstrate substantial decreases in biomass, as discussed above and shown in Table 9-19 (Neufeld et al., 1995). However, exposure for up to three growing seasons did not decrease the biomass of eastern hemlock, Table

1 Mountain pine, and Virginia pine seedlings exposed to O₃ under similar conditions in this
2 location, as shown in Table 9-19 (Neufeld et al., 2000).

3 As for deciduous species, there is a substantial range in sensitivity among evergreen
4 species. As discussed above for deciduous species, a risk analysis was undertaken to predict tree
5 biomass growth reductions due to O₃ based on exposure-response equations for tree seedlings
6 combined with species distribution across the eastern United States and interpolated O₃ exposure
7 (Hogsett et al., 1997). While some species such as Virginia pine were predicted to be affected
8 only slightly even in a high O₃ year, the growth of sensitive evergreen species such white pine
9 was predicted to be reduced by 5% in a lower-than-average O₃ year and 10% in a high ozone
10 year across 50% of its range (Andersen et al., 1997). The remainder of this section discusses
11 experimental results for evergreen species in the order shown in Table 9-20.

12 Douglas fir seedlings were exposed to elevated O₃ concentrations in a free air zonal air
13 pollution system in British Columbia, Canada for two growing seasons with 12-h mean values in
14 1988 of 18 to 41 ppb and in 1989 of 27 to 66 ppb (Runeckles and Wright, 1996). Although
15 substantial variation was seen in effects among the different treatments, there was a significant
16 decrease in the growth of the second flush weight in the second year, with reductions of 55% at
17 the highest O₃ exposure, based on a linear regression. This result contrasts with the lack of
18 effect seen in a previous study with seedlings of this species grown in pots for 134 days and
19 exposed to 7-h mean O₃ concentrations up to 71 ppb (Table 9-30 in U.S. Environmental
20 Protection Agency, 1996).

21 First-year loblolly pine seedlings of 53 open-pollinated families were exposed to
22 1×-ambient O₃ in OTCs for a single growing season, and average growth volume was decreased
23 by 14% compared to a CF treatment (McLaughlin et al., 1994). The 1×-ambient O₃ exposure in
24 this study expressed as 24-h SUM00 was 137 ppm·h, and the CF treatment reduced O₃ by 47%.
25 In this study, the root/shoot ratio was decreased significantly in six of the nine families
26 examined. Exposure to O₃ with SUM06 values up to 162 ppm·h and 132 ppm·h in 2 successive
27 years in OTCs had no effect on seedlings grown in competition with various species of grasses
28 and forbs (Barbo et al., 2002). Exposure of 3-year-old seedlings to O₃ exposures of up to
29 2.5×-ambient (12-h mean of 98 ppb) also had no significant effect (Anttonen et al., 1996).
30 Information summarized in the 1996 AQCD (U.S. Environmental Protection Agency, 1996),
31 indicated that significant effects on seedling growth were observed in several studies of

1 seedlings exposed to elevated O₃ concentrations for one or more years. Several studies,
2 including that of McLaughlin et al. (1994), demonstrate considerable variation in O₃ sensitivity
3 among different genotypes of loblolly pine.

4 For ponderosa pine seedlings, the 1996 AQCD reviewed a number of studies with
5 exposures to elevated O₃ concentrations for one to three growing seasons (U.S. Environmental
6 Protection Agency, 1996). Recent similar studies support the earlier results (Table 9-20);
7 (Andersen et al., 2001; Takemoto et al., 1997). The 1996 criteria document also discussed at
8 some length the ongoing work examining effects of O₃ across a naturally-occurring O₃ gradient
9 in the San Bernardino Mountains in California. Since that time, much research on ponderosa
10 pine has focused on interactive effects of additional stresses such as nitrogen, and effects of O₃
11 on physiological parameters (Sections 9.4, 9.7). Effort has also focused on the effects of O₃ on
12 root growth because such effects could alter sensitivity to drought or nutrient stress. Ecosystem
13 level effects of ozone are discussed further in Section 9.7, but some information relevant to
14 exposure-response relationships is discussed below.

15 For several tree species, O₃ has been shown in experimental studies with seedlings to
16 reduce root growth more than shoot growth (U.S. Environmental Protection Agency, 1996).
17 Ponderosa pine has been shown to be sensitive to O₃, and studies with seedlings have shown
18 reduced root growth, decreases root/shoot ratios, and decreased allocation to roots (Andersen
19 et al., 1991; 1997; Andersen and Rygiewicz, 1995; Andersen and Scagel, 1997; Tingey et al.,
20 1976). More recently, data from a long-studied pollution gradient in the San Bernardino
21 Mountains of southern California suggests that O₃ substantially reduces root growth in natural
22 stands of ponderosa pine. Root growth in mature trees was decreased at least 87% in a high
23 pollution site as compared to a low pollution site (Grulke et al., 1998), and a similar pattern was
24 found in a separate study with whole tree harvest along this gradient (Grulke and Balduman,
25 1999). Because other potential influences on root growth, including shading by competing trees,
26 soil temperature, soil moisture, phenology, were not correlated with the observed pattern of
27 reduced root growth, the authors conclude that O₃ was the cause of the observed decline in root
28 growth. Further results of field investigations with ponderosa pine and other pine species native
29 to California are discussed below under the heading “Scaling experimental data on long-lived
30 species to field conditions” as well as in Section 9.7.

1 Table Mountain pine, Virginia pine, and eastern hemlock seedlings were exposed to
2 various levels of O₃ (CF to 2 × ambient) in OTCs for in a series of experiments two or three
3 years in Great Smoky Mountains National Park in Tennessee (Neufeld et al., 2000). There were
4 no statistically significant effects of O₃ exposure on stem or root biomass, and only slight effects
5 on the biomass of the oldest needles in Table Mountain pine in the 2× ambient treatment.

6 As reviewed in the 1996 criteria document, studies of the response of red spruce to
7 O₃ exposures generally have not found effects on growth of seedlings or saplings, even after
8 exposure to high concentrations (12-h mean of 90 ppb) for up to 4 years. A more recent report
9 confirms that this slow-growing species is O₃ insensitive for at least several years (Laurence
10 et al., 1997).

11 For perennial vegetation, cumulative effects over more than one growing season may be
12 important. For three-year-old Norway spruce in Sweden, exposure to elevated O₃ for three
13 growing seasons decreased total biomass by 18% and stem biomass by 28% (Karlsson et al.,
14 1995). However, after a fourth season of exposure, total biomass decreased significantly by only
15 8% (Karlsson et al., 2002). In this experiment, the O₃ exposures expressed as 12-h mean values
16 averaged over four growing seasons were 12 and 44 ppb for the CF and 1.5×-ambient treatments,
17 respectively; and AOT40 values were 2 and 23 ppm·h, respectively. Despite 4 years of
18 exposure, this experiment did not demonstrate a consistent trend in the O₃ effect on biomass to
19 indicate a significant carry-over effect.

21 **9.6.5.6 Summary of Effects on Evergreen Woody Species**

22 In summary, the O₃ sensitivity of different genotypes within species and between species
23 of evergreen vegetation varies widely. Based on studies with seedlings in OTCs, major species
24 in the United States are generally less sensitive than are most deciduous trees, and slower-
25 growing species are less sensitive than faster-growing ones. However, exposure to ambient O₃
26 may reduce the growth of seedlings of commonly occurring species. Because tree species are
27 long-lived, most experiments have only covered a very small portion of the life span of a tree,
28 making estimating of any effect on mature trees difficult. Considerations for scaling the results
29 of seedling studies to mature forest trees as well as additional information from field surveys and
30 studies of mature trees under natural conditions are discussed below and in Section 9.7.

9.6.5.7 Scaling Experimental Data to Mature Trees

As compared with annual crop species, it is much more difficult to define appropriate exposure-response relationships for tree species. For annual species, an experiment may cover the whole life span of the plant, but it is not feasible to provide controlled-exposure conditions for long-lived plants for any significant portion of their life spans. Most studies have used small seedlings because they are manageable under experimental conditions; but seedlings and mature trees may have different sensitivities to O₃. For perennial species, effects of O₃ may accumulate over more than 1 year, and may interact with other stresses such as drought stress over multiple growing seasons. As for annual species (Section 9.4.2), substantial variability occurs among evergreen genotypes and this variation may interact with other stress responses differently in different landscapes and regions. Despite these difficulties, investigators have addressed some of these issues since the 1996 AQCD (U.S. Environmental Protection Agency, 1996). New information is available on the response of mature evergreen trees to O₃ under field conditions, and models based on tree physiology and stand dynamics have been used to predict O₃ effects on forest stands and regions. The following issues are reviewed briefly below: (1) interaction of drought and O₃ stress, (2) scaling data from seedlings to mature-tree studies. Two additional scaling issues are addressed in Section 9.7: (1) scaling data to forest stands, and (2) scaling data to ecosystems and regions.

9.6.5.7.1 Interactive Effects of Drought and Ozone

Many interacting factors may influence the effect of O₃ on vegetation. For crop plants, environmental conditions are often managed such that nutrients and water are not strongly limiting; but for native vegetation, including most perennial species, such factors are likely to limit growth. The effects of interacting stresses on vegetation are reviewed in Section 9.4. However, because drought is common in many forests, and because there is substantial evidence that it alters the response of trees to O₃, it is discussed in this section in the context of determining exposure-response relationships for trees.

Controlled experiments with seedlings provide direct evidence that drought can reduce the impact of O₃. For example, for 3-year-old Norway spruce in Sweden, exposure to elevated O₃ for three growing seasons decreased total biomass by 18% and stem biomass by 28% (Karlsson et al., 1995). However, for droughted trees, both total and stem biomass decreased

only 5%, with a statistically significant interaction with O₃ for stem biomass. Yet after a fourth season of exposure, there was no longer any interaction between drought and O₃, while there was a significant decrease of 8% in the biomass when both drought and well-watered data were combined (Karlsson et al., 2002). In this study, seedlings were grown in sand in 120 L pots and for the drought treatment, water was withheld for 4 weeks during the first year and for 7 to 8 weeks during each of the last 3 years. In this experiment, the O₃ exposures expressed as 12 h seasonal daylight mean averaged over four growing seasons were 12 and 44 ppb for the CF and 1.5×-ambient treatments respectively. Over this period, the AOT40 values for the treatments averaged 2 and 23 ppm·h respectively. Despite 4 years of exposures, this experiment did not demonstrate a consistent trend in drought O₃ interactions. The difference in effects seen between the third and fourth season suggest that scaling drought-O₃ interactions from seedlings to mature trees may be difficult. However, evidence from biomonitoring surveys supports an interaction between drought and O₃ effects, at least for foliar symptoms. In systematic surveys of foliar symptoms on species selected as biomonitors throughout much of the eastern United States, symptoms were more common and more severe in areas with high O₃ concentrations (Smith et al., 2003). However, in very dry years, such as 1999, the occurrence and severity of symptoms was greatly reduced, even in areas with high ambient O₃ concentrations.

9.6.5.7.2 Scaling from Seedlings to Mature Trees

Because most experiments are conducted with seedlings, various methods are required to scale experimental data on seedlings to mature trees. An overview of physiological differences between young and old plants, and the consequences of these differences for O₃ sensitivity, is provided in Section 9.4.5.3. The discussion below focuses on information relevant to developing exposure-response relationships for mature trees. Information from a few experimental studies, as well as scaling efforts based on physiological characteristics incorporated into models, are discussed in Section 9.7.

Although most studies continue to examine the effects of O₃ on seedlings, during the 1990s some studies examined the effects of O₃ on the response of mature trees. Studies of mature trees demonstrate differences in some aspects of O₃ sensitivity between seedlings and mature trees. For some species, such as red oak, seedlings are less sensitive to O₃ than are mature trees (Hanson et al., 1994; Samuelson and Edwards, 1993; Wullschleger et al., 1996). Both red oak

seedlings and genetically related mature trees were exposed to CF, 1×-ambient, or 2×-ambient O₃ exposures in OTCs in Tennessee for two growing seasons (Hanson et al., 1994). Nine large chambers (4.6 × 8.2 m) were used to enclose individual mature trees and standard EPA-style OTCs were used for potted seedlings. Ozone exposures expressed as a 24-h SUM00 were 34, 79, and 147 ppm·h in 1992 and 37, 95, 188 ppm·h in 1993 for the sub-ambient, and 2×-ambient treatments. Mature trees had a greater light-saturated net photosynthetic rate and stomatal conductance compared to seedling foliage at physiological maturity. By the end of the growing season, exposure to 1×-ambient and 2×-ambient O₃ reduced the light-saturated net photosynthetic rate and stomatal conductance of mature trees by 25 and 50%, respectively, compared with the CF treatment (35 ppm·h). In seedlings, however, light-saturated net photosynthetic rate and stomatal conductance were less affected by O₃ exposure. The authors concluded that extrapolations of the results of seedling-exposure studies to foliar responses of mature forests without considering differences in foliar anatomy and stomatal response between juvenile and mature foliage may introduce large errors into projections of the O₃ responses of mature trees.

In a study of ponderosa pine in California, seedlings and branches of mature trees (in branch chambers) were exposed to O₃ concentrations of 0.5-, 1-, and 2×-ambient O₃ concentrations (Momen et al., 1997). Net photosynthetic rate of 1-year-old, but not current-year, foliage was reduced in mature trees while not significantly reduced in seedlings. This effect was not due to alteration of stomatal conductance by O₃. This result contrasts with those with earlier studies of red spruce (Rebbeck et al., 1993).

In contrast to the findings for red oak and ponderosa pine, giant sequoia seedlings had higher rates of stomatal conductance, CO₂-exchange rate, and dark respiration than did mature trees (Grulke and Miller, 1994). As compared to older trees, stomatal conductance was more than 7-fold greater in current-year and 4-fold greater in 2-year-old seedlings (Grulke and Miller, 1994). The authors concluded that giant sequoia seedlings are sensitive to atmospheric O₃ until ~5 years of age. Low conductance, high water use efficiency, and compact mesophyll all contribute to a natural O₃ tolerance, or to O₃ defense, or to both, in the foliage of older trees. Similarly, lower stomatal conductance was found in mature Norway spruce in Austria compared to seedlings grown with optimal water and nutrients in a growth chamber (Wieser, 1997). In this study, net photosynthetic rate was less sensitive to added O₃ in mature trees compared to

seedlings. In a related study, the average rate of O₃ uptake of 17-year-old trees ~0.6 nmol m⁻² s⁻¹, decreasing linearly in older trees, such that rates were only ~0.1 m⁻² s⁻¹ in 216-year-old trees (Wieser et al., 1999).

Based on a review of studies of stomatal conductance in both seedlings and mature trees, Samuelson and Kelly (2001) concluded that O₃ uptake in oak species, black cherry, sugar maple, and American beech averaged 47% lower in potted seedlings than in mature trees. For evergreen species, they concluded that O₃ uptake in seedlings averaged 26% higher than in mature trees. They also suggested that artifacts introduced by growth in pots confound these differences that exposure-response functions derived from seedlings grown in situ are more applicable to mature trees than are studies of seedlings grown in pots (Samuelson and Kelly, 2001).

As discussed above for annual vegetation, it has long been noted that internal O₃ dose is more appropriate than external O₃ exposure for assessing the effects of O₃ on vegetation, because effects occur primarily via the uptake of O₃ through the stomata (Section 9.3.2). However, external O₃ exposure sometimes has been shown to explain O₃ effects as well or better than calculated internal O₃ dose. For ponderosa pine, Grulke and others (2002) found little difference in the response of net photosynthetic rate and stomatal conductance to O₃ exposure as compared to calculated O₃ uptake; and estimated O₃ uptake by ponderosa pine and O₃ exposure at several sites were highly correlated (r² = 0.92). For red oak, Hanson and others (1994) found that SUM00 explained 83% of the variance in the response of light-saturated photosynthetic rate to O₃ levels, while estimated internal dose explained only 76% of the variance. In this same study, SUM06 explained only 49% of the variance. Due to genetic variation or other factors, individual mature trees will vary in their response to similar O₃ exposures. For example, in 125-year-old giant sequoia trees exposed to ~230 ppm·h of O₃ in branch chambers, O₃ uptake in one individual was ~5 mmol m⁻², while in another it was ~9.5 mmol m⁻² (Grulke et al., 1996).

Based on these results, stomatal conductance, O₃ uptake, and O₃ effects cannot be assumed to be equivalent in seedlings and mature trees. In general, mature deciduous trees are likely to be more sensitive to O₃ compared to seedlings, while mature evergreen trees are likely to be less sensitive than seedlings. However, even when differences in physiological traits occur, concomitant effects on stem growth may not be detected in the field. Additionally, complex interactions may occur between environmental conditions and O₃ responses; and artifacts may occur for seedling studies, especially for seedlings grown in pots. Section 9.7 further discusses

issues that must be addressed when scaling data from individual mature trees to forests and regions.

9.6.6 Studies with the Chemical EDU

The chemical EDU (*N*-[2-(2-oxo-1-imidagolidinyl)ethyl]-*N*-phenylurea) has been used with the goal of protecting plants from O₃ effects without controlling O₃ exposure (Table 9-21) (U.S. Environmental Protection Agency, 1986, 1996). The use of EDU has the potential to be a low-cost, practical method of evaluating ambient O₃ exposures on plants grown under natural conditions without the limitations imposed by methodologies such as OTCs (Section 9.2). EDU can be applied easily as a foliar spray or a soil drench and, with more effort, can be injected directly into plant stems. However, because EDU is phytotoxic, and may have effects on plants other than antioxidant protection, it is crucial that the correct dosage for protection from O₃, be determined without the direct effects of EDU. Unfortunately, although many studies with EDU have been conducted in recent decades, very few have used multiple EDU application levels along with multiple O₃ exposures to characterize the EDU system for a given plant species. Therefore, the text of this section focuses on how data from existing studies can be used for developing or validating exposure-response relationships, rather than reviewing results of all individual studies. Data from individual studies on O₃ exposure, EDU application rates, and the effects of EDU are presented in Table 9-21. In addition to EDU, sodium erythorbate has been used in a few studies as a protectant chemical. Since very few published studies have used sodium erythorbate and attempts to establish appropriate doses for individual species are even more limited, the use of this chemical is not reviewed here.

The phytotoxicity of EDU is well known, and the point has been made repeatedly that for a particular species or cultivar, tests under a range of environmental conditions and O₃ exposures must be made to establish the efficacy of EDU for quantifying O₃ effects (Heggestad, 1988; Kostka-Rick and Manning, 1992). A recent study has shown that even low concentrations of EDU (8 to 32 mg L⁻¹ soil), decreased bean yield under low O₃ exposure (7-h mean of 19 ppb) in CF OTCs (Miller et al., 1994). This study also demonstrated that phytotoxicity (both foliar symptoms and growth effects) can differ even in the same series of experiments, apparently due to changes in environmental conditions, and that EDU can suppress yield at application rates that do not always cause foliar symptoms (Miller et al., 1994). Finally, this study found that EDU

1 altered biomass partitioning by increasing vegetative growth and decreasing reproductive
2 growth. A study of bean grown in OTCs in Germany found that EDU treatment in CF OTCs
3 significantly increased yield, while EDU had no significant effect on yield in other O₃ treatments
4 (Brunschon-Harti et al., 1995). In this study, O₃ significantly reduced the mass of pods, shoots,
5 and roots. EDU increased root, leaf, and shoot mass across O₃ treatments. However, there was
6 only a significant interaction with O₃ for root mass. This study indicates that EDU can stimulate
7 above-ground growth and/or delay senescence regardless of O₃ treatment. Together, these
8 studies suggest that EDU has effects other than its antioxidant protection and phytotoxicity.

9 Recent studies have also shown that EDU does not always have greater effects at higher O₃
10 exposures. For bean grown in pots in studies in Spain and the Netherlands, EDU increased pod
11 yield (Ribas and Penuelas, 2000; Tonneijck and Van Dijk, 1997). However, this effect was not
12 greater at sites with higher O₃ exposure despite consistent experimental protocols at all sites,
13 including growing the same cultivar in pots with adequate water (Ribas and Penuelas, 2000;
14 Tonneijck and Van Dijk, 1997). Such results suggest that it may be difficult to quantify
15 ambient O₃ effects using EDU, because the amount of plant growth or yield expected at a low
16 (background) O₃ concentration cannot be inferred from EDU-treated plants grown at locations
17 with higher O₃ exposures.

18 Potted white clover exposed to ambient O₃ at 12 sites throughout Europe over
19 three growing seasons was evaluated in a meta-analysis (Ball et al., 1998). A soil drench of
20 100 mL EDU was applied every 2 weeks for 3 months per growing season. Only very weak
21 evidence of a linear relationship of the ratio of biomass in control versus EDU treated plants
22 across all sites ($r^2 = 0.16$) was seen. However, an artificial neural network (ANN) model
23 including VPD, temperature, longitude, year, and altitude explained much more of the variance
24 ($r^2 = 0.79$). The authors suggested that the greater sensitivity at certain sites in Germany may
25 have been due to occurrence of other pollutants. This meta-analysis indicates that EDU effects
26 may be influenced substantially by environmental factors.

27 In summary, the EDU method for assessing the impact of ambient O₃ exposures is
28 potentially useful, because it provides a separate line of evidence than other methods. However,
29 as has been pointed out by numerous authors, the system must be carefully characterized for
30 each species under different environmental conditions and different O₃ exposures.

31 Unfortunately, such characterization has so far been limited, although substantial progress has

1 been made for radish (Kostka-Rick et al., 1993; Kostka-Rick and Manning, 1992, 1993). Effects
2 of EDU can include phytotoxicity and alteration of biomass partitioning, along with differential
3 effects on the same species in different years or locations. Additionally, the degree of O₃
4 protection afforded by a certain application of EDU is difficult to quantify. Thus, it is difficult to
5 use data from existing EDU studies to develop exposure-response relationships or to quantify the
6 effects of ambient O₃ exposure. Despite these limitations, the EDU studies reviewed in previous
7 criteria documents (U.S. Environmental Protection Agency, 1986, 1996) and the more recent
8 studies summarized in Table 9-21 provide another line of evidence that ambient O₃ exposures
9 occurring in many regions of the United States may be reducing the growth of crops and trees.

11 **9.6.7 Summary**

12 Recently published data support the conclusions of previous criteria documents that there
13 is strong evidence that ambient concentrations of O₃ cause injury and damage to numerous
14 common and economically valuable plant species. For annual vegetation, the data summarized
15 in Table 9-17 show a range of growth and yield responses both within and among species.
16 Nearly all of these data were derived from studies in OTCs, with only two studies using open-air
17 systems in the UK (Ollerenshaw et al., 1999; Ollerenshaw and Lyons, 1999). It is difficult to
18 compare studies that report O₃ exposure using different indices, such as AOT40, SUM06, or 7-h
19 or 12-h mean values. However, when such comparisons can be made, the results of recent
20 research confirm earlier results summarized in the 1996 AQCD (U.S. Environmental Protection
21 Agency, 1996). A summary of earlier literature concluded that a 7-h, 3-month mean of 49 ppb
22 corresponding to a SUM06 exposure of 26 ppm·h would cause 10% loss in 50% of 49
23 experimental cases (Tingey et al., 1991). Recent data summarized in Table 9-17 support this
24 conclusion, and more generally indicate that ambient O₃ exposures can reduce the growth and
25 yield of annual species. Some annual species such as soybean are more sensitive, and greater
26 losses may be expected (Table 9-17). Thus the recent scientific literature supports the
27 conclusions of the 1996 AQCD (U.S. Environmental Protection Agency, 1996) that ambient O₃
28 concentrations are probably reducing the yield of major crops in the United States.

29 Much research in Europe has used the AOT40 exposure statistic, and substantial effort has
30 gone into developing Level-1 critical levels for vegetation using this index. Based on regression
31 analysis of 15 OTC studies of spring wheat including one study from the United States and

1 14 from locations ranging from southern Sweden to Switzerland, an AOT40 value of 5.7 ppm·h
2 was found to correspond to a 10% yield loss, and a value of 2.8 ppm·h corresponded to a 5%
3 yield loss (Fuhrer et al., 1997). Because a 4 to 5% decrease could be detected with a confidence
4 level of 99%, a critical level of an AOT40 value of 3 ppm·h was selected in 1996 (Kärenlampi
5 and Skärby, 1996).

6 In addition to likely reductions in crop yield, O₃ may also reduce the quality or nutritive
7 value of annual species. Many studies have shown effects of O₃ on various measures of plant
8 organs that affect quality, with most studies focusing on characteristics important for food or
9 fodder. These studies indicate that there may be economically important effects of ambient O₃
10 on the quality of crop and forage species. Previous criteria documents have concluded that
11 visible symptoms on marketable portions of crops and ornamental plants can occur with seasonal
12 7-h mean O₃ exposures of 40 to 100 ppb (U.S. Environmental Protection Agency, 1978, 1986,
13 1996). The recent scientific literature does not refute this conclusion.

14 The use of OTCs may reverse the usual vertical gradient in O₃ that occurs within a few
15 meters above the ground surface (Section 9.2). This reversal suggests that OTC studies may
16 somewhat overestimate the effects of an O₃ concentration measured several meters above the
17 ground. However such considerations do not invalidate the conclusion of the 1996 AQCD (U.S.
18 Environmental Protection Agency, 1996) that ambient O₃ exposures (Tables 9-14 and
19 Table 9-22) are sufficient to reduce the yield of major crops in the United States.

20 As for single-season agricultural crops, yields of multiple-year forage crops are reduced at
21 O₃ exposures that occur over large areas of the United States. This result is similar to that
22 reported in the 1996 AQCD (U.S. Environmental Protection Agency, 1996). When species are
23 grown in mixtures, O₃ exposure can increase the growth of O₃-tolerant species and exacerbate
24 the growth decrease of O₃-sensitive species (e.g., Ashmore and Ainsworth, 1995; Fuhrer et al.,
25 1994). Because of this competitive interaction, the total growth of the mixed-species community
26 may not be affected by O₃ exposure (Ashmore and Ainsworth, 1995; Barbo et al., 1998; Fuhrer
27 et al., 1994). However, in some cases, mixtures of grasses and clover species have shown
28 significant decreases in total biomass growth in response to O₃ exposure in studies in the United
29 States (Heagle et al., 1989; Kohut et al., 1988) and in Sweden (Pleijel et al., 1996). In Europe,
30 a provisional critical level for herbaceous perennials of an AOT40 value of 7 ppm·h over
31 6 months has been proposed to protect sensitive plant species from adverse effects of O₃.

1 For deciduous tree species, recent evidence from free air exposure systems and OTCs
2 supports results observed previously in OTC studies. For example, a series of studies
3 undertaken using free air O₃ enrichment in Rhinelander, WI (Isebrands et al., 2000, 2001)
4 demonstrated that O₃-symptom expression was generally similar in OTCs, FACE, and also sites
5 along an ambient O₃ gradient, supporting the previously observed variation among aspen clones
6 using OTCs (Karnosky et al., 1999). As has been observed in previous criteria documents, root
7 growth often is found to be the most sensitive biomass response indicator to O₃.

8 Results reported since 1996 support the conclusion of the 1996 AQCD (U.S.
9 Environmental Protection Agency, 1996) that deciduous trees are generally less sensitive to
10 O₃ than are most annual plants, with the exception of a few very sensitive genera such as
11 *Populus* and sensitive species such as black cherry. However, the data presented in Table 9-19
12 suggest that ambient O₃ exposures that occur in the United States can potentially reduce the
13 growth of seedlings of deciduous species. Results from multiple-year studies sometimes show a
14 pattern of increased effects in subsequent years. In some cases, however, growth decreases due
15 to O₃ may become less significant or even disappear over time. While some mature trees show
16 greater O₃ sensitivity in physiological parameters such as net photosynthetic rate than do
17 seedlings, these effects may not translate into measurable reductions in biomass growth.
18 However, because even multiple-year experiments do not expose trees to O₃ for more than a
19 small fraction of their life span, and because competition may in some cases exacerbate the
20 effects of O₃ on individual species, determining effects on mature trees remains a significant
21 challenge.

22 In Europe, a Level I critical level has been set for forest trees based on OTC studies of
23 European beech seedlings. A critical level was defined as an AOT40 value of 10 ppm·h for
24 daylight hours for a 6-month growing season (Kärenlampi and Skärby, 1996). However, other
25 studies show that some species such as silver birch may be more sensitive to O₃ compared to
26 beech (Pääkkönen et al., 1996).

27 For evergreen tree species, as for other tree species, the O₃ sensitivity of different
28 genotypes and different species varies widely. Based on studies with seedlings in OTCs, major
29 species in the United States are generally less sensitive than are most deciduous trees, and
30 slower- growing species are less sensitive than are faster-growing species. Interacting stresses
31 such as competition stress may increase the sensitivity of trees to O₃. As for deciduous species,

1 most experiments with evergreen species have only covered a small portion of the life span of a
2 tree and have been conducted with seedlings, making estimating effects on mature trees difficult.

3 For all types of perennial vegetation, cumulative effects over more than one growing
4 season may be important, furthermore, studies for only a single season may underestimate
5 effects. Mature trees may be more or less sensitive to O₃ than are seedlings, depending on the
6 species, but information on physiological traits may be used to predict some such differences.
7 In some cases, mature trees may be more sensitive to O₃ than seedlings due to differences in
8 their gas exchange rates, growth rates, greater cumulative exposures, or due to the interaction of
9 O₃ stress with other stresses.

12 **9.7 EFFECTS OF OZONE EXPOSURE ON NATURAL ECOSYSTEMS**

13 **9.7.1 Introduction**

14 The preceding section on species-level responses (9.6) provides a lead-in to address the
15 response of ecosystems to ozone (O₃). The conclusion of the 1996 O₃ AQCD was that aside
16 from the results from the San Bernardino Forest, there was no direct evidence that O₃ is altering
17 natural ecosystems in the United States. This conclusion is generally valid today, except that our
18 understanding of the effects of O₃ in the San Bernardino forest has been tempered by additional
19 understanding of the complicating role that N deposition plays in this system. Despite the lack
20 of any new, direct information linking O₃ with ecosystem changes, numerous publications since
21 1996 have highlighted ways in which O₃ may affect ecosystem structure and/or function. This
22 section addresses new and (where appropriate) older literature in order to illustrate possible
23 shifts in energy or material flow through ecosystems as a result of O₃ exposure.

24 An ecosystem is defined as comprising all of the organisms in a given area interacting with
25 the physical environment, so that a flow of energy leads to a clearly defined trophic structure,
26 biotic diversity, and cycling of materials between living and nonliving parts (Odum, 1963).
27 Individuals within a species and populations of species are the building blocks from which
28 communities and ecosystems are constructed. Classes of natural ecosystems are distinguished
29 by their dominant vegetation form, e.g., tundra, wetland, deciduous forest and conifer forest.
30 Boundaries of ecosystems are delineated when an integral unit is formed by the physical and

1 biological parts. Defined pathways for material transport and cycling, and for the flow of energy
2 are contained within the integrated unit.

3 Each level of organization within an ecosystem has functional and structural
4 characteristics. At the ecosystem level, functional characteristics include but are not limited to
5 energy flow; nutrient, hydrologic, and biogeochemical cycling; and maintenance of food chains.
6 The sum of the functions carried out by ecosystem components provides many benefits to
7 mankind as in the case of forest ecosystems (Smith, 1992). These include food, fiber production,
8 aesthetics, genetic diversity, and energy exchange.

9 Ecosystems are functionally highly integrated. Thus, changes in one part of an ecosystem,
10 such as the primary producer component, may have consequences for connected parts, such as
11 the consumer and decomposer components. For example, when needles are shed prematurely as
12 a result of O₃ exposure, successional development of phyllosphere fungi inhabiting the surface of
13 ponderosa pine needles may be truncated (Bruhn, 1980). In addition, decomposer populations in
14 the litter layer may be capable of higher rates of decomposition, because the younger age classes
15 of needles falling from O₃-damaged pines have higher nitrogen (N) content (Fenn and Dunn,
16 1989). Because ecological systems integrate the effects of many influences, the resulting effect
17 of O₃ exposure may depend on co-occurring influences that predispose an ecosystem to stress
18 (Colls and Unsworth, 1992). One important change in our thinking since the 1996 O₃ AQCD is
19 that at high levels of O₃ exposure that are known to result in detectable plant responses
20 (> 250 ppm h accumulated over a growing season), N deposition must also be considered as a
21 concurrent stressor. Changes in N cycling and compartmentalization in the ecosystem result
22 from both O₃ exposure and increased N deposition.

23 The vast majority of O₃ effects literature addresses individual species responses (see
24 Section 9.7.4.3), which was also true in 1996 (U.S. Environmental Protection Agency, 1996).
25 This section differs from the preceding one in that physiological stress of individual species is
26 considered only within the context of the natural ecosystem. Changes in function at the level of
27 the individual are propagated through higher levels of organization, resulting in changes in
28 ecosystem structure and function. However, since ecosystem-level responses result from the
29 interaction of organisms with one another and with their physical environment, it takes longer
30 for a change to develop to a level of prominence that can be identified and measured. The
31 paucity of scientific literature on O₃ effects at the ecosystem level is a result of both the

complexity of ecological systems, and long response times. In addition, “indirect” effects of O₃ on plants (e.g., affecting the plants' ability to integrate environmental stresses) may be more important than direct effects on photosynthesis and respiration at the leaf level (Johnson and Taylor, 1988).

A conceptual framework (see Table 9-22) suitable for organizing discussion of the effect of O₃ on ecosystems was developed by the EPA Science Advisory Board (Young and Sanzone, 2002). Their six essential ecological attributes (EEAs) include landscape condition, biotic condition, chemical and physical characteristics, ecological processes, hydrological and geomorphological processes, and natural disturbance regimes (listed as subsection headings below and in Table 9-22). The major ecological effects, and gaps in our knowledge of O₃ exposure effects at the ecosystem level are summarized at the end of this chapter. While the main focus is O₃ effects newly described since the last Ozone Criteria Document (EPA, 1996), many key historical papers are cited to demonstrate ecosystem response, particularly where they remain the only examples in the literature. Although the vast majority of published studies focus on individuals, five field examples and one FACE experiment have measured several ecosystem components simultaneously to better understand ecosystem response to O₃. We provide an overview of these six studies up-front because they provide a context for the subsequent discussion on possible ecosystem effects.

9.7.2 Case Studies

9.7.2.1 Valley of Mexico

The first evidence of air pollution impacts on vegetation in the Valley of Mexico (Mexico City Air Basin) were observations of foliar injury symptoms in bioindicator plants attributed to O₃, PAN, SO₂ and possibly other pollutants (de Bauer, 1972). Subsequently, O₃ injury to foliage and crowns of pine trees were reported in forests to the south and southwest of Mexico City (Krupa and de Bauer, 1976; de Bauer and Hernández-Tejeda, 1986). Ozone is considered to be the pollutant with the most severe impacts on vegetation within the Mexico City urban zone and in forests downwind of the city. *P. hartwegii* is the most O₃-sensitive pine species and is severely impacted by high O₃ exposures encountered to the south/southwest of the metropolitan area (Miller et al., 2002). The potential for O₃ injury is particularly high in this area, because O₃ levels are high during the summer rainy season when soil moisture availability and stomatal

Table 9-22. Essential Ecological Attributes for Natural Ecosystems Affected by O₃

Category	Species	Condition Measures	References
Landscape Condition			
• Habitat Types			
Biotic Condition			
• Ecosystems and Communities	Mixed conifer forest	Community composition, Stand structure	Miller et al. (1989); Miller and McBride (1999)
	Community Extent		
	Community Composition		
	<i>Pinus ponderosa</i>	Relative abundance	Miller (1973); Arbaugh et al. (2003)
	Grassland communities	Species composition	Ashmore and Ainsworth (1995); Ashmore et al. (1995)
	Coastal sage scrub	Species cover, richness, equitability	Westman (1979, 1981)
	Early successional plant community	Species richness, diversity, evenness	Barbo et al. (1998)
	<i>Populus tremuloides</i> & <i>Betula papyrifera</i>	Soil microbial community	Phillips et al. (2002)
	<i>Pinus ponderosa</i>	Soil microbial community	Scagel and Andersen (1997)
	<i>Pinus taeda</i>	Fungal morphotypes	Edwards and Kelly (1992); Qui et al. (1993)
Trophic Interactions	Insects		
	<i>Pinus ponderosa</i>	Bark beetle severity	Cobb et al. (1968)
	<i>Pinus ponderosa</i>	Bark beetle productivity and predator/parasitoid density	Dahlsten et al. (1997)
	<i>Populus tremuloides</i>	Blotch leaf miner performance	Kopper and Lindroth (2003a)
	<i>Populus tremuloides</i>	Aphid/natural enemy abundance	Percy et al. (2002)
Diseases	<i>Populus hybrids</i>	Septoria occurrence	Woodbury et al. (1994)
	<i>Populus hybrids</i>	Rust occurrence	Beare et al. (1999)
	<i>Populus tremuloides</i>	Rust occurrence	
	<i>Populus tremuloides</i>	Forest tent caterpillar/paratoid performance	Karnosky et al. (2002); Percy et al. (2002) Holton et al. (2003)
	<i>Picea abies</i> & <i>Picea sitchensis</i>	Needle fungi	Magan et al. (1995)
	<i>Pinus ponderosa</i>	Root disease x O ₃ interactions	Fenn et al. (1990)
	<i>Pinus taeda</i>	Canker dimensions	Carey and Kelly (1994)
	<i>Pinus sylvestris</i> / <i>mycorrhizae</i>	Disease susceptibility	Bonello et al. (1993)
Community Dynamics	<i>Pinus ponderosa</i> / <i>Abies concolor</i> / <i>calocedrus decurrens</i>	Abundance	Minnich et al. (1995)
	<i>Populus tremuloides</i>	Competitive status	McDonald et al. (2002)
	<i>Pinus ponderosa</i> / <i>Elymus glaucus</i>	O ₃ sensitivity	Andersen et al. (2001)
	<i>Pinus taeda</i> /diverse community	Tree growth	Barbo et al. (2002)

**Table 9-22 (cont'd). Essential Ecological Attributes for Natural Ecosystems
Affected by O₃**

Category	Species	Condition Measures	References
• Species and Populations			
Population Size	<i>Pinus strobus</i> <i>Pinus ponderosa</i>	Mortality Mortality	Karnosky (1981) Carroll et al. (2003)
Genetic Diversity/ Population Structure	<i>Lupinus bicolor</i> <i>Populus tremuloides</i> <i>Trifolium repens</i> <i>Plantago major</i>	% population sensitive % population sensitive % population sensitive % population sensitive	Dunn (1959) Berrang et al. (1986, 1989, 1991) Heagle et al. (1991) Reiling and Davison (1992a); Davison and Reiling (1995); Lyons et al. (1997)
Population Dynamics	<i>Trifolium repens</i> <i>Plantago major</i>	Adaptation Population changes over time	Heagle et al. (1991) Davison and Reiling (1995)
Organism Condition			
• Visible Symptoms			
	<i>Pinus ponderosa</i>	Foliar symptoms	Grulke and Lee (1997); Arbaugh et al. (1998); Salardino and Carroll (1998); Temple et al. (1992)
	<i>Pinus jeffreyi</i>	Foliar symptoms	Patterson and Rundel (1995); Salardino and Carroll (1998); Grulke et al. (2003b)
	<i>Prunus serotina</i>	Foliar symptoms	Fredericksen et al. (1995, 1996); Chappelka et al. (1997, 1999a,b); Hillebrand et al. (1996); Ghosh et al. (1998); Lee et al. (1999); Ferdinand et al. (2000, 2003);
	<i>Liriodendron tulipifera</i>	Foliar symptoms	Yuska et al. (2003) Somers et al. (1998); Hillebrand et al. (1996); Chappelka et al. (1999a)
	<i>Sassafras albidum</i> <i>Populus nigra</i> , <i>Fraxinus excelsior</i> & <i>Prunus avium</i>	Foliar symptoms Foliar symptoms	Chappelka et al. (1999a) Novak et al. (2003)
	<i>Fagus sylvatica</i>	Foliar symptoms	Gerosa et al. (2003); Vollenweider et al. (2003a)
	<i>Fraxinus americana</i>	Foliar symptoms	Schaub et al. (2003); Ferdinand et al. (2003)
	Grassland species	Foliar symptoms	Bungener et al. (1999a)
	Herbaceous species	Foliar symptoms	Bergmann et al. (1999)
	<i>Asclepias exaltata</i>	Foliar symptoms	Chappelka et al. (1997)
	<i>Rudbeckia laciniata</i> & <i>Verbesina occidentalis</i>	Foliar symptoms	Chappelka et al. (2003)
	<i>Asclepias incarnata</i>	Foliar symptoms	Orendovici et al. (2003)

**Table 9-22 (cont'd). Essential Ecological Attributes for Natural Ecosystems
Affected by O₃**

Category	Species	Condition Measures	References
• Physiological Status	<i>Pinus halepensis</i>	Allometry	Welburn and Wellburn (1994)
	<i>Populus tremuloides</i>	Crown architecture	Dickson et al. (2001)
	<i>Betula pendula</i>	Crown architecture	Kull et al. (2003)
	<i>Fagus sylvatica</i>	Crown architecture	Stribley and Ashmore (2002)
	<i>Populus tremuloides</i>	Root dry weight	Coleman et al. (1996)
	<i>Pinus ponderosa</i>	Root/shoot ratio	Grulke et al. (1998); Grulke and Balduman, (1999)
	<i>Festiva ovina</i>	Root/shoot ratio	Warwick and Taylor, (1995)
	<i>Betula pubescens</i>	Root/shoot ratio	Mortensen (1998)
	<i>Fagus sylvatica</i>	Root/shoot ratio	
	<i>Populus tremuloides</i> x <i>P. tremula</i>	Root/shoot ratio	Paludan-Müller et al. (1999); Landolt et al. (2000)
	<i>Prunus serotina</i>	Leaf area	Oksanen et al. (2001a)
	<i>Fraxinus excelsior</i>	Leaf area	
	<i>Populus tremuloides</i>	Leaf area index	
	<i>Pinus ponderosa</i>	Carbon allocation to mycorrhizae	Neufeld et al. (1995) Wiltshire et al. (1996)
	<i>Betula pendula</i>	Decreased winter bud formation	Karnosky et al. (2003a) Andersen and Rygielwicz (1995)
	<i>Betula pendula</i>	Delayed bud break	Oksanen (2003a,b)
	<i>Acer saccharum</i>	Early bud break	
			Prozherina et al. (2003) Bertrand et al. (1999)
• Reproductive Status	<i>Apocynun</i> <i>androsaemifolium</i>	Flowering time	Bergweiler and Manning, (1999)
	<i>Buddleia davidii</i>	Flowering time	Findley et al. (1997)
	<i>Rubus cuneifolius</i>	Pollen germination &	Chappelka (2002)
	<i>Plantago major</i> ,	Pollen tube elongation	Stewart (1998)
	<i>Fragaria x ananassa</i>	Fruit yield	
	<i>Plantago major</i>	Seed yield	Drogoudi and Ashmore (2000, 2001)
			Reiling and Davison (1992b); Pearson et al. (1996); Whitfield et al. (1997); Lyons and Barnes (1998)
	Understorey herbs	Seed yield	Harward and Treshow (1975)

**Table 9-22 (cont'd). Essential Ecological Attributes for Natural Ecosystems
Affected by O₃**

Category	Species	Condition Measures	References
Ecological Processes			
• Energy Flow			
Primary Production	<i>Pinus ponderosa</i>	Photosynthesis	Miller et al. (1969); Clark et al. (1995); Takemoto et al. (1997); Grulke et al. (2002b)
	<i>Pinus ponderosa</i> <i>Populus tremuloides</i>	Needle retention Photosynthesis	Temple et al. (1993) Coleman et al. (1995b); Noormets et al. (2001a,b); Sharma et al. (2003); Karnosky et al. (2003a) Oksanen (2003a,b)
	<i>Betula pendula</i> <i>Betula pendula</i>	Photosynthesis/conductance Stem respiration & radial growth	Matyssek et al. (2002) Kelting et al. (1995)
	<i>Quercus rubra</i>	Root turnover	Coleman et al. (1996)
	<i>Populus tremuloides</i>	Soil respiration	King et al. (2001)
	<i>Populus tremuloides</i>	Soil respiration	Andersen and Scagel (1997); Scagel and Andersen (1997), Andersen (2000)
	<i>Pinus ponderosa</i>	Soil respiration	Samuelson and Kelly (1996) Coleman et al. (1995b)
	<i>Quercus rubra</i>	Carbon partitioning & allocation	Andersen et al. (1997); Grulke et al. (1998); Grulke and Balduman (1999); Grulke et al. (2001)
	<i>Populus tremuloides</i> <i>Pinus ponderosa</i>	Carbon allocation Carbon allocation	
	<i>Betula pendula</i> <i>Betula pendula</i>	Carbon allocation Carbon allocation	Karlsson et al. (2003) Oksanen and Saleem, (1999); Salem et al. (2001)
	<i>Fragaria vesca</i> <i>Pinus taeda</i> <i>Lespedeza cuneata</i> & <i>Schizachyrium</i> <i>scoparium</i>	Carbon allocation Root respiration Yield	Manninen et al. (2003) Edwards (1991) Powell et al. (2003)
	<i>Liriodendron tulipifera</i> <i>Prunus serotina</i> <i>Pinus jeffreyi</i> <i>Pinus ponderosa</i> <i>Pinus strobus</i> <i>Pinus taeda</i>	Radial growth Radial growth Radial growth Radial growth (no effect) Radial growth Radial growth	Somers et al. (1998) Vollenweider et al. (2003b) Peterson et al. (1987) Peterson et al. (1993) Bartholomay et al. (1997) and Downing (1995, 1996) Braun et al. (1999)
	<i>Fagus sylvatica</i> <i>Picea abies</i> <i>Populus tremuloides</i> <i>Pinus ponderosa</i>	Stem volume Stem volume Volume growth Root growth	Wallin et al. (2002) Isebrands et al. (2001) Andersen et al. (1991)

**Table 9-22 (cont'd). Essential Ecological Attributes for Natural Ecosystems
Affected by O₃**

Category	Species	Condition Measures	References
Net Ecosystem Production	Northern hardwoods	NPP estimates	Laurence et al. (2000)
	Northern hardwoods	Biomass estimates	Hogsett et al. (1997)
Growth Efficiency	<i>Plantago major</i>	Relative growth rate	Reiling and Davison (1992a); Davison and Reiling (1995); Lyons et al. (1997); Davison and Barnes (1998)
	Grassland species	Relative growth rate	Bungener et al. (1999b)
	Native herbs	Relative growth rate	Warwick and Taylor (1995)
	Grasses and herbs	Relative growth rate	Pleijel and Danielsson (1997)
	<i>Populus tremuloides</i>	Relative growth rate	Yun and Laurence (1999)
	<i>Fagus sylvatica</i>	Relative growth rate	Bortier et al. (2000)
	<i>Picea abies</i>	Relative growth rate	Karlsson et al. (2002)
	<i>Prunus serotina</i>	Relative growth rate	Lee et al. (2002)
• Material Flow Organic Carbon Cycling	<i>Populus tremuloides</i> & <i>Betula papyrifera</i>	Altered foliar C:N ratio and N resorption efficiency	Lindroth et al. (2001)
	<i>Andropogon virginicus</i> & <i>Rubus cuneifolius</i>	Litter decomposition rate	Kim et al. (1998)
	<i>Liriodendron tulipera</i>	Litter decomposition rate	Scherzer and Rebbeck (1998)
	<i>Populus deltoides</i>	Litter decomposition rate	Findlay and Jones (1990)
	<i>Pinus ponderosa</i>	Litter decomposition rate	Fenn and Dunn (1989)
	<i>Pinus sylvestris</i>	Litter decomposition (no effect)	Kainulainen et al. (2003)
Nitrogen Cycling	<i>Pinus ponderosa</i>	Altered foliar N Foliar N & O ₃ exposure (no effects)	Momen and Helms (1996) Bytnerowicz et al. (1990)
	<i>Pinus taeda</i>		Mandersheid et al. (1992)
	<i>Prunus serotina</i> & <i>Liriodendron tulipifera</i>	Altered foliar N metabolism Altered foliar N	Boerner and Rebbeck (1995)
Other Nutrient Cycling	<i>Picea sitchensis</i> & <i>Pinus sylvestris</i>	Foliar leaching (no effect)	Skeffington and Sutherland (1995)
	<i>Pinus ponderosa</i>	Nutrient availability & O ₃	Bytnerowicz et al. (1990)
Hydrology and Geomorphology			
• Water Budget	<i>Picea rubens</i>	Water-use efficiency (no effect)	Laurence et al. (1997)
	<i>Pinus armandi</i>	Water-use efficiency	Shan et al. (1996)
	<i>Pinus jeffreyi</i>	Canopy transpiration	Grulke et al. (2003a)
	<i>Picea abies</i>	Transpiration, xylem sap flow	Maier-Maercker (1997)
	<i>Fraxinus excelsior</i>	Stem flow of water	Wiltshire et al. (1994)
	<i>Betula pendula</i>	Water-use efficiency	Maurer and Matysek (1997)
	<i>Populus</i> hybrids	Water-use efficiency	Reich and Lassoie (1984)

**Table 9-22 (cont'd). Essential Ecological Attributes for Natural Ecosystems
Affected by O₃**

Category	Species	Condition Measures	References
<i>Natural Disturbance Regimes</i>			
• Frequency	<i>Pinus ponderosa</i>	Frequency of fire	McBride and Laven (1976); Minnich et al. (1995); Miller and McBride (1999)
	<i>Pinus ponderosa</i>	Occurrence of bark beetle outbreaks	Pronos et al. (1999); Dahlsten et al. (1997)
• Intensity	<i>Picea sitchensis</i>	Winter damage	Lucas et al. (1988)
	<i>Pinus halepensis</i>	Reduced winter damage	Wellburn and Wellburn, (1994)
	<i>Picea rubens</i>	Freezing tolerance	Waite et al. (1994)
	<i>Fagus sylvatica</i>	Drought stress	Pearson and Mansfield (1993, 1994)
	<i>Picea abies</i>	Drought stress	Maier-Maercker (1998); Maier-Maercker and Koch (1992)
	<i>Pinus ponderosa</i>	Fire intensity	Miller and McBride (1999)
• Extent	<i>Pinus ponderosa</i>	Extent of bark beetle attack	Minnich et al. (1995)
• Duration	<i>Pinus ponderosa</i>	Duration of bark beetle attack	Minnich et al. (1995)

Source: Young and Sanzone (2002).

conductance are greatest and these factors enhance O₃ uptake and injury. Decline of *A. religiosa* (oyamel) in the Desierto de los Leones National Park is a well-known example of dramatic dieback and mortality of entire forest stands due primarily to air pollution stress (Alvarado-Rosales and Hernández-Tejeda, 2001). Other factors, such as a lack of stand thinning, also contribute to forest decline. Lead in automobile gasoline was phased out in 1990, and foliar concentrations of heavy metals in forest species are not now at phytotoxic levels (Fenn et al. 2001a). Sulfur dioxide concentrations decreased in the early 1990s as a result of regulatory mandates limiting their emissions. Sensitive plants in the NE and NW sectors of the Mexico City urban zone where concentrations are highest may still be impacted by exposure to ambient SO₂ levels. Deposition of ionic forms of N and S are high in forested areas southwest of Mexico City. The effects of these chronic nutrient inputs to the forest are only beginning to be investigated and understood.

1 The ecological perturbations caused by severe air pollution exposures in forests located
2 downwind of Mexico City are expected to continue for the near future (the next 5 to 10 years),
3 largely as a result of high O₃ concentrations, as well as of N oxides emissions. The longer-term
4 response is more uncertain, and depends largely on the effectiveness of regulatory emissions
5 control strategies. Currently, pollutant levels are declining. Forest responses to this trend will
6 depend on how long it takes to reduce levels sufficiently to allow sensitive species to recover.
7 Some of the change to the ecosystem is probably irreversible, such as the loss of lichen diversity
8 and of other O₃-sensitive species (Zambrano et al., 2002).

9 10 **9.7.2.2 San Bernardino Mountains**

11 The San Bernardino Mountains lie east of the Los Angeles Air Basin (California, USA),
12 and significant levels of pollution have been transported into the mountain range, including a
13 Class I Wilderness area. The effects of O₃ exposure on the mixed conifer forest of the
14 San Bernardino Mountains is perhaps the longest and most thoroughly documented O₃
15 ecological effects evaluation (Miller and McBride, 1999). In this classic case study linking
16 tropospheric O₃ exposure to damage to an entire forest ecosystem (U.S. Environmental
17 Protection Agency, 1996) (Table 9-23), Miller et al. (1963) first identified the unique foliar
18 chlorotic mottle that was occurring on two of the dominant tree species, *Pinus ponderosa* and
19 *P. jeffreyi*. Levels of O₃ averaging 100-120 ppb over 24 hours with 1-hour peaks well into the
20 200 ppb range were common in the region in the 1960s and 1970s (Miller and McBride, 1999).
21 Single hour peak values have declined in recent years due to heavily regulated pollution control,
22 but O₃ concentrations in the moderate range continue to rise and accumulate, increasing the
23 overall cumulative exposure (Arbaugh et al., 1998; Takemoto et al., 2001; Lee et al., 2003).

24 Since the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) was written, the
25 concurrent role of N deposition in modifying ecosystem response to O₃ exposure in the San
26 Bernardino Mountains has been further elucidated (Fenn et al., 1996, 2003; Bytnerowicz et al.,
27 1999; Takemoto et al., 2001; Bytnerowicz, 2002). Both O₃ exposure and N deposition reduce
28 foliar retention (Grulke and Balduman, 1999) and alter tissue chemistry of both needles and litter
29 (Poth and Fenn, 1998). In addition, confounding factors such as drought and fire suppression
30 add to the complexity of ecosystem response (Minnich et al., 1995; Takemoto et al., 2001;
31 Arbaugh et al., 2003). Extensive crown injury measurements have also been made, linking

Table 9-23. Case Studies Demonstrating the Ecological Effects of O₃

Study	Keystone Species	Study Type	Period Studied	Key Ecological Findings
Valley of Mexico	<i>Pinus hartwegii</i> , <i>Abies religiosa</i>	Field transects	35 yrs	<ul style="list-style-type: none"> • Significant foliar injury (de Bauer, 1972; Krupa and Bauer, 1976; Hernández-Tejeda, 1986) • Community composition changes (Alvarado-Rosales and Hernández-Tejeda, 1986) • Species richness changes (Zambrano et al., 2002)
San Bernardino Mountains	<i>Pinus ponderosa</i> , <i>P. jeffreyi</i>	Field transects	40 yrs	<ul style="list-style-type: none"> • Community composition changes (Miller, 1973; Minnich et al., 1995; Arbaugh et al., 2003) • Population changes (McBride and Laven, 1999) • O₃/pine/bark beetle interaction (Pronos et al. 1999) • Altered C flows (Grulke et al., 2002b; Grulke et al., 1998; Grulke and Balduman, 1999; Grulke et al., 2001; Arbaugh et al., 1999) • Interaction of O₃, drought, N deposition: (Fenn et al., 1996; Grulke, 1999; Takemoto et al., 2001) • Altered carbon cycling (Arbaugh et al., 1999)
Sierra Nevada Mountains	<i>Pinus ponderosa</i> , <i>P. jeffreyi</i>	Field	35 yrs	<ul style="list-style-type: none"> • Wide-scale nature of effects (Miller and Millecan, 1971; Miller et al., 1972) • Link to decreased growth (Peterson et al., 1987, 1991, 1995) • Quantification of O₃ flux (Bauer et al., 2000; Panek et al., 2002; Goldstein et al., 2003) • Cumulative O₃ effects (Takemoto et al., 1997) • Canopy level responses (Grulke et al., 2003a,b) • Population changes (Carroll et al., 2003)
Appalachian Mountains	<i>Fraxinus americana</i> , <i>Liriodendron tulipifera</i> , <i>Pinus strobus</i> , <i>Prunus serotina</i>	Field	25 yrs	<ul style="list-style-type: none"> • Link of visible symptoms to growth decreases (McLaughlin et al., 1982; Somers et al., 1998) • Wide-scale nature of effects (Chappelka et al., 1999a; Hildebrand et al., 1996)

Table 9-23 (cont'd). Case Studies Demonstrating the Ecological Effects of O₃

Study	Keystone Species	Study Type	Period Studied	Key Ecological Findings
Aspen FACE	<i>Acer saccharum</i> , <i>Betula papyrifera</i> , <i>Populus tremuloides</i>	Open-air O ₃ exposure	6 yrs	<ul style="list-style-type: none"> Competitive interactions (McDonald et al., 2002) O₃/aspen/rust interaction (Karnosky et al., 2002) Plant- insect interactions (Percy et al., 2002; Holton et al., 2003) C & N cycling (Lindroth et al., 2001; King et al., 2001) Moderation of CO₂ responses by O₃ (Isebrands et al., 2001; Wustman et al., 2001; McDonald et al., 2002; Karnosky et al., 2003a)
Plantago	<i>Plantago major</i>	Field	20 yrs	<ul style="list-style-type: none"> Population structure (Davison and Reiling, 1995; Lyons et al., 1997) O₃ resistance (Reiling and Davison, 1992a) Adaptation (Davison and Reiling, 1995)
Carpathian Mountains	<i>Pinus sylvestris</i> , <i>Picea abies</i>	Field	15 yrs	<ul style="list-style-type: none"> Significant foliar injury Community composition changes Species diversity changes

ambient O₃ exposure data to chlorotic mottle and fascicle retention (Arbaugh et al., 1998). Ozone exposure and N deposition reduce carbon allocation to stems and roots (Grulke et al., 1998; 2001), further predisposing trees to drought stress, windthrow, root diseases, and insect infestation (Takemoto et al., 2001). Increased mortality of susceptible tree species (*ponderosa* and *Jeffrey pine*, *Pinus ponderosa*, *P. Jeffreyi*) has shifted community composition towards white fir and incense cedar (*Abies concolor*, *Libocedrus decurrens*) and has altered forest stand structure (Miller et al., 1989). Ozone exposure is implicated in projected changes in stand composition (McBride and Laven, 1999) toward a predominance of oaks, rather than mixed conifer forests. Forest understory species have also been affected (Temple, 1999). These individual species responses collectively have affected trophic structure and food web dynamics (Dahlsten et al., 1997; Pronos et al., 1999), as well as C and N cycling (Arbaugh et al., 2003) (Table 9-24). Because of the high N deposition in the San Bernardino Mountains, it is difficult to separate out effects of only O₃ versus those due to the combined effects of O₃ and N deposition (Table 9-25).

Table 9-24. The Most Comprehensively Studied Effects of O₃ on Natural Ecosystem is the San Bernardino Mountain Forest Ecosystem. Citations Focus on Research Published Since U.S. EPA (1996).

Pollutant Occurrence	Reference
O ₃ exposures N deposition	Grulke et al. (1998); Fenn et al. (1996, 2000, 2003); Bytnerowicz et al. (1999)
Cellular, Biochemical	
Foliar pigments Antioxidants	Grulke and Lee (1997); Grulke (1999); Tausz et al. (1999a,b 2001); Tausz et al. (1999a,b,c, 2001)
Foliar Responses	
Foliar Symptoms	Grulke and Lee (1997); Arbaugh et al. (1998); Miller and Rechel (1999)
Gas Exchange	
Photosynthesis & Conductance O ₃ flux Foliar nutrients	Grulke (1999); Grulke and Retzlaff (2001); Grulke et al. (2002a,b)
Whole Organism	
Growth/Biomass	
• Aboveground	Grulke and Balduman (1999)
• Belowground	Grulke et al. (1998); Grulke and Balduman (1999)
• Root/shoot ratio	Grulke and Balduman (1999)
• Carbon allocation	Grulke et al. (2001)
• Crown vigor	Arbaugh et al. (1998); Miller and Rechel (1999)
Ecosystem	
Community dynamics/succession Simulations	McBride and Miller (1999); Arbaugh et al. (2002) Arbaugh et al. (1999)
Understory vegetation	Temple (1999)
Pest interactions	
• Bark beetle/predators	Dahlsten et al. (1997); Pronos et al. (1999)
• Disease occurrence	Miller and Rechel (1999); Pronos et al. (1999)
• Litter decomposition	
Disturbance	
• Bark beetle occurrence	Minnich et al. (1995); Minnich (1999); Miller and McBride (1999)
• Fire frequency	

Table 9-25. Effects of Ozone, Ozone and N Deposition, and Ozone and Drought Stress on *Pinus ponderosa* and *Pinus jeffreyi* in the Sierra Nevada and the San Bernardino Mountains, California. Citations are Focused on Research Published Since U.S. EPA (1996).

	O ₃	O ₃ + N deposition	O ₃ + Drought	References
<i>Foliar Biochemistry and Tissue Chemistry</i>				
Total ascorbate	d ¹	d	i	Tausz et al. (2001); Grulke et al. (2003b)
Dehydroascorbate	i	n.d.	d	Grulke et al. (2003b)
Total glutathione	d	i	d	Tausz et al. (2001)
Oxidized glutathione	i	i	d	Tausz et al. (2001)
á Carotenoids	i	n.d.	d	Grulke et al. (2003b)
Foliar nitrogen	d	i	d	Grulke and Lee (1997); Grulke et al. (1998); Poth and Fenn (1998)
C:N ratio of foliage ²	i	n.d.	d	Poth and Fenn (1998); Grulke et al. (2003b)
Starch	n.d.	d	i	Grulke et al. (2001)
Chlorophyll content	d	id	d	Takemoto et al. (1997); Grulke and Lee (1997; Grulke et al. (1998, 2003b); Grulke (1999); Tausz et al. (2001)
<i>Gas Exchange</i>				
A _{max} lower canopy	n.d.	i	d	Grulke (1999); Grulke and Retzlaff (2001); Grulke et al. (2002b); Panek (2004)
A _{max} whole canopy	d	n.d.	d	Panek and Goldstein (2001); Grulke et al. (2003b); Misson et al. (2004)
A _{max} seedlings	di	n.d.	n.d.	Grulke and Retzlaff (2001)
Stomatal limitation	á to ps	n.d.	i	Panek and Goldstein (2001)
Stomatal conductance	d	di	d	Grulke (1999; Grulke et al. (2003a); Panek (2004)
Foliar respiration	n.s.	i	d	Grulke et al. (2002a; Grulke (1999)
O ₃ flux	d	n.s.	d	Panek and Goldstein (2001); Panek et al. (2002; 2003); Grulke et al. (2002b; 2004)

Table 9-25 (cont'd). Effects of Ozone, Ozone and N Deposition, and Ozone and Drought Stress on *Pinus ponderosa* and *Pinus jeffreyi* in the Sierra Nevada and the San Bernardino Mountains, California. Citations are Focused on Research Published Since U.S. EPA (1996).

	O ₃	O ₃ + N deposition	O ₃ + Drought	References
<i>Growth and Productivity</i>				
Foliar biomass	n.d.	i	d	Grulke and Balduman (1999)
Height growth	n.d.	i	d	Grulke and Balduman (1999)
Bole diameter growth	d	i	d	Grulke and Balduman (1999)
Fine root biomass	d	d	i	Grulke et al. (1998)
Leaf Surfaces				
Stomatal occlusion	i	n.d.	n.d.	Bytnerowicz et al. (1999); Bytnerowicz and Turunen (1994)
Trophic Interactions				
Bark beetle	n.s.	i	i	Pronos et al. (1999)
<i>Ecosystem Level</i>				
Competitive indices	n.d.	d	i	Miller and Rechel (1999)

¹Responses are shown as significant increases (i), significant decreases (d), both significant decreases and increases reported (di), nonsignificant effects (n.s.), and no data (n.d.) compared to trees or seedlings at field sites with lower ozone, drought stress, or lack of significant N deposition (< 10 kg ha⁻¹ yr⁻¹). Frequently n.d. was used for lack of a control site without compounding high N deposition. Foliar analyses and leaf surface properties were largely determined from previous year needles. Gas exchange data were generally from previous year needles at peak growing season, prior to late summer drought (mid to late July).

²Abbreviations: C = carbon; N = nitrogen; A_{max} = maximum photosynthesis rate.

9.7.2.3 Sierra Nevada Mountains

The western slope of the Sierra Nevada Mountains in central and southern California has also been exposed to elevated O₃ for a long time, although the effects have been much less than those observed in the San Bernardino Mountains. Symptoms of O₃ injury have been found on ponderosa and Jeffrey pines in all of the Sierra Nevada National forests and parks (Carroll et al., 2003). First identified as a problem in the 1970s (Miller et al., 1972), elevated O₃ with daytime means of 60-80 ppb are common (Böhm et al., 1995; Bauer et al., 2000; Bytnerowicz et al., 2002a; Panek et al., 2002). The west-slope Sierra Nevada forests are also exposed to a wide range of additional gaseous and particulate pollutants, including various S and N compounds

(Bytnerowicz et al., 1999; Takemoto et al., 2001; Fenn et al., 2003), but at levels much lower than in the San Bernardino Mountains. Typical O₃-induced visible foliar symptoms, including chlorotic mottle, chlorophyll degradation, and premature senescence, are commonly found on sensitive genotypes of *ponderosa pine* (Peterson et al., 1991; Arbaugh et al., 1998; Staszak et al., 2003) and *Jeffery pine* (Peterson et al., 1987; Patterson and Rundel, 1995; Arbaugh et al., 1998; Grulke et al., 2003b). Other important conifers in the region, such as giant sequoia, appear to be relatively tolerant to O₃ (Grulke et al., 1996). The symptoms of foliar injury and growth reductions have been verified on seedlings in O₃ exposure chambers (Temple, 1988; Momen and Helms, 1996; Momen et al., 2002).

Ozone foliar injury of dominant pine species in the Sierra Nevada Mountains is correlated to decreased radial growth in both *ponderosa pine* (Peterson et al., 1991) and *Jeffreyi pine* (Peterson et al., 1987; Patterson and Rundel, 1995). Because of the large amount of intraspecific variation in O₃ sensitivity in these two species, O₃ exposure may be a selective agent (Patterson and Rundel, 1995), with differential mortality rates for sensitive individuals (Carroll et al., 2003). The region's forests may also be experiencing subtle changes in species composition and community structure (Patterson and Rundel, 1995; Takemoto et al., 2001).

Based on fire scar dating, reconstructions of stand age classes, historical records, and present stand structure, fire has been largely excluded in western forests for the last 75 to 100 years (Minnich, 1999; Minnich and Padgett, 2003). Fire exclusion has resulted in fewer, large, stand-replacing fires rather than a mosaic of smaller, lower intensity fires. The change in fire intensity may have selectively altered stand structure, fitness and competitiveness of component species, and their susceptibility to atmospheric pollutants and other stressors (Minnich, 1999). Short-lived (50-80 years) species such as knobcone and Coulter pine, which occur at the interface between the chaparral and the mixed conifer forest, may already have been selected for O₃ tolerance due to seedling establishment (the most sensitive tree age class in conifers) after large fires in the 1950's (Minnich, 1999). Strong measures to suppress fires have largely kept chaparral fires from invading the mixed conifer forests, and stand densification in the mixed conifer zone has increased. High stand density, in turn, may weaken the younger cohorts and increase sensitivity to atmospheric pollution (Minnich, 1999).

Other disturbances that play a potential role in sensitivity to atmospheric pollution include cycles of drought stress. Nearly every decade is marked by one or more years of very low

precipitation (Graumlich, 1993). During extended periods of drought, foliar injury is lower than in subsequent years with higher average precipitation (Carroll et al., 2003). In the first several years (1975-1977) of a Sierran-wide assessment of O₃ injury to pines, O₃ injury increased, because of greater water availability due to greater stomatal conductance and, presumably, greater O₃ uptake. Trees instrumented with monitors to directly measure canopy transpiration had 20% greater stomatal conductance in mesic microsites (riparian areas, mid-slope seeps) than trees in xeric microsites (rock outcrops) (Grulke et al., 2003a). Although the Sierra Nevada experienced a prolong drought between 1987 and 1993, it was less severe and O₃ injury did not significantly decrease (Carroll et al., 2003). The same plots showed only a slight increase in O₃ injury between 1993 and 2000. Drought stress, in general, can make trees more susceptible to insect and pathogen infestation. However, serious outbreaks of insect infestation are believed to be indicators, not a cause, of existing stress in the forest (Wickman, 1992).

9.7.2.4 Appalachian Mountains

The southern Appalachian Mountain region experiences some of the highest O₃ exposures of any natural areas in the eastern United States (Mueller, 1994; Hildebrand et al., 1996; Samuelson and Kelly, 1997; Chappelka et al., 1997). Since the region is the home of the Shenandoah and Great Smokey Mountains National Parks, which have Class I air quality designations by the 1977 Clean Air Act, there has been considerable study of the region's dominant forest species to determine O₃ effects. Visible foliar symptoms of O₃ have been found in natural ecosystems consisting of *Sassafras albidum* (Chappelka et al., 1999a), *Prunus serotina* (Hildebrand et al., 1996; Chappelka et al., 1997; 1999b; Samuelson and Kelly, 1997), *Liriodendron tulipifera* and *Fraxinus Americana* (Hildebrand et al., 1996; Chappelka et al., 1999a). Visible foliar symptoms induced by O₃ have been recreated on the same species in chamber studies (Duchelle et al., 1982; Chappelka et al., 1985; Chappelka and Chevone, 1986; Samuelson, 1994; Fredericksen et al., 1995). No response to O₃ exposure has been found for other hardwood trees, nor for the three conifer species tested (Neufeld et al., 2000).

Long term foliar injury symptoms have been correlated with decreased radial growth in tulip poplar and cherry (*Liriodendron tulipifera* and *Prunus serotina*; Somers et al. (1998) and with decreased biomass in cherry (Neufeld et al., 1995). Although climatic conditions (drought) largely explained radial growth reductions, O₃ exposure may have also contributed (McLaughlin

and Downing, 1996). Ozone exposure may also be affecting the understory vegetation in the region (Duchelle and Skelly, 1981; Duchelle et al., 1983; Chappelka et al., 1997; Chappelka et al., 2003; Davison et al., 2003) and community composition (Barbo et al., 1998), through impacts both on growth and reproduction (Chappelka, 2002). Foliar litter from trees exposed to elevated O₃ have lower decomposition rates (Kim et al. 1998). Other air pollutants are likely to be found in this ecosystem but not at such high deposition values found in the California studies. A decline in forest health in the northern Appalachians has been primarily attributed to the effects of acidic fog and rain on soil acidification, lower Ca²⁺ availability, reduction in fine root biomass, and modification of cuticular wax. However, fog- and O₃-exposed red spruce forests also show winter injury (Percy, 2002).

9.7.2.5 Plantago Studies in the United Kingdom

One of the most well-documented studies of population and community response to O₃ effects are the long-term studies of *Plantago major* in native plant communities in the United Kingdom (Reiling and Davison, 1992a; Davison and Reiling, 1995; Lyons et al., 1997). Sensitive populations of *P. major* had significant growth decreases in elevated O₃ (Reiling and Davison, 1992b and 1992c; Pearson et al., 1996; Whitfield et al., 1997) and reduced fitness as determined by decreased reproductive success (Reiling and Davison, 1992b; Pearson et al., 1996). While spatial comparisons of population responses to O₃ are complicated by other environmental factors, rapid changes in O₃ resistance were imposed by ambient levels and variations in O₃ exposure (Davison and Reiling, 1995). Molecular patterns of genetic variation suggest that change in O₃ resistance over time probably resulted from selection on genotypes already present in local populations, rather than through an influx of new *P. major* germplasm (Wolff et al., 2000). The highest correlations between O₃ resistance and ambient O₃ concentrations occurred at the site of seed collection (Lyons et al., 1997), rather than between O₃ resistance and other climatic variables, as was found for aspen (Berrang et al., 1991).

9.7.2.6 Forest Health in the Carpathian Mountains

The Carpathian Mountains cross five countries (the Czech Republic, the Slovak Republic, Poland, Romania, and the Ukraine) and contain many national parks and several biosphere reserves. The forests were largely cleared in the 15th century, and were then planted in Norway

1 spruce plantations. As elevation increases, beech (*Fagus sylvatica*) or beech-fir (*Abies alba*)
2 forests grade into Norway spruce (*Picea abies*) or spruce-fir forests. Near the treeline, Norway
3 spruce combines with dwarf mountain pine (*Pinus mugo*). Dwarf mountain pine forms an
4 almost pure stand just below alpine vegetation.

5 The forests of the Carpathian Mountains have been subjected to anthropogenic stressors
6 (shepherding, metal mining, wood harvest for structures and paper) for hundreds of years, as
7 described for the Tatra Mountains in the southern Carpathians (Wezyk and Guzik, 2004). The
8 Carpathians have been subjected to regional air pollution stresses since industrialization. Most
9 of the effects of air pollution on forest health degradation were due to (1) heavy metal
10 deposition, (2) soil acidification by acid deposition, and (3) subsequent pest outbreaks, the
11 combination of which led to the forest decline and dieback between 1970 and 1989 (Dunajski,
12 2004). By the 1980's, industrial pollutants such as SO₂ and heavy metals significantly declined,
13 but O₃ exposure has continued to increase (Bytnerowicz et al., 2004). Increased ownership and
14 use of private cars in Central Europe, as well as long-range transport of O₃ from western Europe,
15 are believed to be responsible for the continued increase in photooxidants. In 1995, drought
16 resulted in significant mortality, as well as an epidemic of bark beetle infestation in subsequent
17 years. A network of air quality monitoring sites was installed across Europe in the late 1980's, as
18 part of the International Cooperative Programme on Assessment and Monitoring of Air Pollutant
19 Effects on Forests (ICP Forests). Mean defoliation rates for six important forest species across
20 Europe have increased or remained unchanged from 1989 to 1999 (Percy, 2002). Ozone
21 concentrations experienced in the Tatra Mountains, especially along the southern slopes,
22 occasionally reach 190-200 ppb as two-week-long averages, with the highest values experienced
23 in early summer at elevations of 1700 to 2300 m (Bytnerowicz et al., 2004). In other parts of the
24 Carpathian Mountains, peak two-week averages of O₃ concentrations were lower, at 160 ppb
25 (Bytnerowicz et al., 2002). For all trees inventoried, about 13% exhibited greater than 25%
26 defoliation during 1997 to 2000. There was no difference between extent of damage for
27 broadleaves or conifers. Trees in Poland and the Czech Republic are the most affected by air
28 pollution, and the least damaged forests are in Romania (Badea et al., 2002).

29 The extent to which O₃ exposure affects forest health degradation, and slows forest
30 degradation, is still unknown in Europe. In many of the published studies, response to a known
31 O₃ gradient is largely confounded by other pollutants and/or climatic gradients (Bytnerowicz

et al., 1996; Szaro et al., 2002; Widadki et al., 2004). Current levels of ambient O₃ are believed to be high enough to reduce bole radial growth (Percy, 2004). Although average O₃ concentration alone was not related to bole growth, the peak hourly O₃ concentration was negatively correlated to growth (Muzika et al., 2002). Recent evidence indicates that canopy health of European white oak (*Quercus robur*), Norway spruce, maritime pine (*Pinus pinaster*), and beech has significantly declined (Huttunen et al., 2002). However, the canopy health of Scots pine (*Pinus sylvestris*) has improved. The network of air quality monitoring stations and forest plots is extensive and active. Subsequent correlative analyses including both meteorological and air quality attributes throughout the EU will help to determine the specific role of O₃ exposure in forest decline. Historical effects of anthropogenic disturbance may still be confounding.

9.7.2.7 Field Exposure System (FACE), Rhinelander, Wisconsin

The Aspen Free-Air CO₂ Enrichment (FACE) facility was designed to examine the effects of both elevated CO₂ and O₃ on aspen (*Populus tremuloides*), birch (*Betula papyfera*), and sugar maple (*Acer saccharum*) in a simple reconstructed plantation characteristic of Great Lakes Aspen-dominated forests (Karnosky et al., 1999, 2003a). Instead of using chambers to expose the plants to desired gas concentrations, the gas is piped up vertical delivery tubes in the open air. The vertical delivery pipes surround a 30-m diameter circular plot with five different aspen clones in half of the plot, one quarter of the plot planted in aspen and birch, and one quarter in aspen and maple. The O₃ treatment for the first five years was 1.5× ambient, with ambient O₃ exposures averaging 35 to 37 ppb (12 h daytime over the growing season) compared to elevated O₃ rings averaging 49 to 55 ppb for the same time period (Karnosky et al., 2003a).

Elevated CO₂, elevated O₃, and elevated CO₂ + O₃ have had effects on most system components being measured in the study (Table 9-26; Karnosky et al., 2003a). One interesting finding of the project has been the nearly complete offset by elevated O₃ of the enhancements induced by elevated atmospheric CO₂ for the pioneer keystone species *P. tremuloides* (Isebrands et al., 2001) and *B. papyrifera* (Percy et al., 2002), even though O₃ exposure alone did not always result in a significant response when compared to controls. They also found evidence that the effects on above- and below-ground growth and physiological processes have cascaded through the ecosystem, even affecting microbial communities (Larson et al., 2002; Phillips et al.,

Table 9-26. Summary of Responses of *Populus tremuloides* to Elevated CO₂ (+200 µmol mol⁻¹), O₃ (1.5 × ambient), or CO₂+O₃ Compared with Control During Three Years of Treatments at the Aspen FACE Project (Modified from Karnosky et al. 2003a)

	CO ₂	O ₃	CO ₂ + O ₃	Reference
<i>Foliar Gene Expression and Biochemistry</i>				
Rubisco; RbcS ² transcripts	d ¹	d	dd	Wustman et al. (2001); Noormets et al. (2001a)
PAL transcripts	d	i	d	Wustman et al. (2001)
Acc oxidase, catalase	d	i	d	Wustman et al. (2001)
Ascorbate peroxidase	d	n.s.	d	Wustman et al. (2001)
Glutathione reductase	d	i	d	Wustman et al. (2001)
Phenolic glycosides	i	d	n.s.	Lindroth et al. (2001); Kopper and Lindroth (2003a,b)
Tannins	n.s.	i	i	Lindroth et al. (2001); Kopper and Lindroth (2003a,b)
Foliar nitrogen	d	n.s.	d	Lindroth et al. (2001); Kopper and Lindroth (2003a,b)
C:N ratio of foliage	i	n.s.	ii	Lindroth et al. (2001)
Starch	d	d	n.s.	Wustman et al. (2001)
<i>Gas Exchange</i>				
A _{max} lower canopy	n.s.	dd	id	Takeuchi et al. (2001); Noormets et al. (2001a)
A _{max} whole canopy	ii	dd	n.s.	Noormets et al. (2001b); Sharma et al. (2003)
Stomatal limitation	d	n.s.	d	Noormets et al. (2001a)
Stomatal conductance	d	di	d	Noormets et al. (2001a)
Foliar respiration	n.s.	i	n.s.	Takeuchi et al. (2001); Noormets et al. (2001b)
Soil respiration	ii	d	n.s.	King et al. (2001)
Microbial respiration	ii	n.s.	n.s.	Phillips et al. (2002)
Stomatal density	n.s.	n.s.	n.s.	Percy et al. (2002)
Chlorophyll content	d	d	d	Wustman et al. (2001)
Chloroplast structure	i	d	d	Oksanen et al. (2001b); Takeuchi et al. (2001); Wustman et al. (2001)
O ₃ flux	d	ii	i	Noormets et al. (2001a)
<i>Growth and Productivity</i>				
Leaf thickness	i	n.s.	n.s.	Oksanen et al. (2001b)
Leaf size	i	d	d	Wustman et al. (2001)
Leaf area	i	d	n.s.	Noormets et al. (2001b)

2002). This study also confirmed earlier observations of changes in trophic interactions involving keystone tree species, as well as important insect pests and their natural enemies (Table 9-26) (Percy et al., 2002; Holton et al., 2003; Awmack et al., 2003).

9.7.3 Landscape Condition

In the SAB framework (Figure 9-19), landscape condition is assessed using the areal extent, composition of component landscape ecosystems or habitat types, and the pattern or structure of component ecosystems or habitat types (including biocorridors). To date, no publications exist on the impacts of O₃ exposure on landscape condition. The effects of O₃ exposure have only been reported at the community or stand level (see Biotic Conditions, below). The following is a description of current discussions by land stewards and of how difficult it will be to quantitatively assess the effect of O₃ exposure on landscape condition.

Landscapes are identified and preserved, such as National Parks, Class I Wilderness Areas, etc., so that they are protected from the effects of O₃ exposure by law. Efforts to determine whether landscapes have been affected by certain levels of exposure rely on valuation of landscape and ecosystem components. Several different approaches of valuation have been used, such as pathological (visible symptoms), biomass and allocation, and biogeochemical.

In the pathological approach, a “critical loads” concept is developed, with varying levels of impact viewed as acceptable, interim targets, or as unacceptable. As an example, land managers of Class I Wilderness Areas may consider a critical level acceptable if it resulted in no visible O₃ symptoms to sensitive species. In concrete terms, sensitive species may respond to peak O₃ exposures of 60 ppb (e.g., coneflower, *Rudbeckia laciniata*, in Great Smoky National Park; Davison et al., 2003), and so the critical exposure level would be < 60 ppb for any hourly valued during the growing season. An interim target would be that less than 5% of the sensitive plants would have visible symptoms of < 15% of the leaf surface. An unacceptable level of O₃ exposure would be any result more pronounced than the interim target. The advantage of the foliar injury approach is that large crews with relatively simple training can assess individual species within the landscape and “see” the effect of the oxidant exposure. There are several disadvantages, however. Some species (e.g., white fir) exhibit no foliar injury, but do have shifts in biomass allocation in response to oxidant exposure (Retzlaff et al., 2001). Other species have shown significant decreases in foliar injury due to needle loss, retranslocation of nutrients to




Figure 9-19. A conceptual diagram of processes and storage pools in sources and sinks that are affected by ozone exposure. A plus (+) denotes an increase in process rate or pool size, a minus (-) denotes a decrease in process rate or pool size, and a plus-minus (±) denotes that both increases and decreases have been reported in response to O₃. Primary effects in the shoots (1°) are distinguished from secondary effects in roots (2°) since the primary site of ozone action occurs in the leaves (from Andersen, 2003).

remaining foliage, with subsequent increased photosynthetic rate (Beyers et al., 1992).

In addition, the development of foliar symptoms within a species is related to sunlight and microclimate (Davison et al., 2003).

In the biomass approach, O₃ exposure resulting in a measureable decline in biomass (usually of a target, sensitive species) is used to evaluate landscape condition. The bulk of the information available is from seedling responses to controlled chamber exposures, reviewed in the previous section. Some information exists for species in natural environments, but teasing out concurrent stressors and finding adequate “controls” may be intractable. For example, in a

1 long-term gradient of O₃ exposure, N deposition, and drought, the site with the highest O₃
2 exposure had the greatest whole tree biomass (pole-sized trees) due to growth stimulation by N
3 deposition (Grulke and Balduman, 1999).

4 In the biogeochemical approach, changes in biogeochemical cycling are used to assess
5 landscape condition. Species sensitive to O₃ exposure have well known responses to O₃
6 exposure, including altered C allocation to below- and above-ground tissues, and altered rates of
7 leaf production, turnover, and decomposition. Changes in turnover rates of ephemeral tissues
8 (leaves, fine roots) also affect nutritional status of the remaining tissue. These shifts can affect
9 overall carbon and N loss from the ecosystem in terms of respired C, and leached aqueous
10 dissolved organic and inorganic C and N. Instability in C and N pools and dynamics can affect
11 landscape-level nutrient dynamics even without significant inputs of N deposition. The endpoint
12 assessment is based on changes in water quality from or in the landscape, correlated to a defined
13 oxidant exposure level. These approaches are linkable: visible injury at a particular level could
14 be related to reduction in photosynthate, which would reduce whole plant biomass (and carbon
15 dynamics). If O₃ sensitive species are dominant within the landscape, then changes in C and N
16 dynamics over time would be expected to alter biogeochemical cycles. Examples of forest types
17 that contain geographically extensive, O₃-sensitive species that could be used in assessing
18 landscape-level changes include ponderosa pine in the western United States, tulip poplar
19 (*Liriodendron tulipifera*) or loblolly pine (*Pinus taeda* L.) in the eastern U.S. deciduous forests,
20 and Norway spruce (*Picea abies* (L.) Karst) in the Carpathian Mountains of eastern Europe.

21 Water quantity may also be affected by O₃ exposure at the landscape level. Moderately
22 high O₃ exposure may affect the mechanism of stomatal opening (McAinsh et al., 2002),
23 resulting in sluggish stomatal opening and closing (Patterson and Rundel, 1989; Reich and
24 Lassoie, 1984). During moderately high O₃ exposure in a drought year, canopy transpiration was
25 greater for tulip poplar than on adjacent days with lower O₃ exposure. This could affect canopy
26 transpiration and landscape level water use (McLaughlin et al., 2004). Oxidant exposure (O₃ and
27 NO_x) may decrease the ability of exposed plants to close stomata at night (Grulke et al., 2004),
28 thus increasing water loss from the landscape. Ecosystem models should aid in interpreting
29 O₃-exposure effects at the landscape level.
30

9.7.4 Biotic Condition

9.7.4.1 Ecosystems and Communities

The SAB framework described by Young and Sanzone (2000) (Figure 9-20) identifies community extent, community composition, trophic structure, community dynamics, and physical structure as EEA's for assessing ecosystem health.

COMMUNITY EXTENT

Ecosystem function is dependent on areal extent, constituent species composition, trophic structure and its dynamics, and community physical structure. Genetic variation within species, and the dynamics of the interactions that exist among different species and their biotic and abiotic environment, are also involved (Agrawal and Agrawal, 2000). There are no reports of O₃ exposure altering community distribution or extent.

COMMUNITY COMPOSITION

Significant changes in plant community composition resulting directly from O₃ exposure has been demonstrated in two forested areas: the mixed conifer forest of the San Bernardino Mountains, CA and the mixed conifer forest of the Valley of Mexico near Mexico City. It is also likely that community composition has changed in response to O₃ exposure in the coniferous forests of the Carpathian Mountains, but this has not yet been definitively shown.

The first forest communities shown to be affected by O₃ were the *Pinus ponderosa*-dominated stands of the San Bernardino Mountains in southern California (Miller, 1973). Miller suggested that mixed forests of *P. ponderosa*, *Pinus jeffreyi* and *Abies concolor* were changing to predominantly *A. concolor* because of the greater sensitivity of the pines to O₃. Significantly greater mortality of young mature trees (50 to 99 years old) occurred in sites that also showed higher foliar injury relative to sites that showed slight foliar injury (McBride and Laven, 1999). For *P. ponderosa*, 33% of the trees in the high foliar injury sites died versus 7% of the trees in the low foliar injury sites over the decade-long census. In contrast, 24% of *Abies concolor* died in high foliar injury sites, whereas no trees died in slight injury sites. The authors suggested that certain age classes were especially sensitive to O₃ exposure because they are emerging into the canopy, where higher O₃ concentrations are encountered. Future projections based on past changes in community composition have been conducted for 2024 and 2074 (McBride and

1 Laven 1999). In their projections, the population of *Pinus ponderosa* nearly disappears in all
2 tree age classes, and the community is dominated by *Quercus kelloggii* in all tree age classes,
3 followed by *Calocedrus decurrens* and *Pinus lambertiana* by the year 2074. Their projections
4 do not account for potential changes in genetic structure of the more O₃-sensitive species.

5 In the Valley of Mexico, the closed forest structure changed to a woodland from high
6 pollutant exposure (Miller et al., 2002). Cryptogamic community diversity also significantly
7 declined in response to prolonged, extreme O₃ exposure (Zambrano et al., 2002). Together,
8 these two examples illustrate the potential for shifts in community composition in response to
9 O₃ stress.

11 **TROPHIC STRUCTURE**

12 *Above-Ground Structural*

13 One of the first reports of trophic level interactions in natural communities was the
14 O₃-induced predisposition of *Pinus ponderosa* to attack by bark beetles (Stark et al., 1968; Cobb
15 et al., 1968; Stark and Cobb, 1969). Trees exposed to oxidant injury had lower resin production,
16 flow, and exudation pressure. Also, several attributes associated with tree defense against beetle
17 attack were compromised by oxidant exposure including sapwood and phloem moisture content
18 and phloem thickness (Pronos et al., 1999). Another trophic level has been implicated, in that O₃
19 injured ponderosa pine had the same rate of bark beetle infection, but healthy trees had greater
20 numbers of bark beetle predators and parasitoids (Dahlsten et al., 1997). This suggests that O₃
21 damage rendered the pines inhospitable for the natural enemies of the bark beetles. Similar
22 findings were presented by Percy et al. (2002) for aphids whose abundance was increased in
23 young *Populus tremuloides* stands exposed to elevated O₃. In that study, the levels of natural
24 enemies of aphids (ladybirds, lacewings, spiders and parasitoids) were significantly decreased
25 under elevated O₃.

27 *Below-Ground*

28 Processing of plant-derived carbon compounds by soil organisms comprising the soil food
29 web is a fundamental property of a functional and stable below-ground ecosystem (de Ruiter
30 et al., 1998; Wolters, 1998). Soil food web organisms are responsible for recycling nutrients and
31 for development of soil properties such as porosity, aggregate structure, water-holding capacity

1 and cation exchange capacity. A shift in food web species diversity or functional complexity in
2 response to O₃ stress may alter ecosystem processes.

3 Evidence that soil organisms are affected by O₃ indicates the potential for changes in soil
4 food web structure and function. Since O₃ does not penetrate the soil beyond a few centimeters,
5 the proposed mechanism by which O₃ alters soil biota is through a change in carbon input to
6 soils (Andersen, 2003). Ozone can alter C inputs to soil and hence soil processes through four
7 different pathways: (1) leaf-litter quality and quantity (see Material Cycling, below), (2) carbon
8 allocation to roots (see Physiological Status, below), (3) interactions among root symbionts, and
9 (4) rhizodeposition. The complex nature of the effects of O₃ on trophic interactions and food
10 webs calls for additional basic research and modeling.

11 There have been no comprehensive studies on the effects of O₃ on structural components of
12 soil food webs, however studies have shown that O₃ affects free-living soil organisms of food
13 webs. In the few cases where soil microbial communities have been examined, O₃ has led to
14 changes in bacterial and fungal biomass, and in some cases changes in soil enzyme activity.
15 Phillips et al. (2002) examined the effects of elevated CO₂ and O₃ on C flow through
16 heterotrophic microbial communities in soils collected from a FACE study in Wisconsin. Ozone
17 decreased abundance of fungal phospholipid fatty acids in aspen and birch-aspen plots but had
18 few other direct effects on measured soil parameters. The greatest effect of O₃ was to eliminate
19 significant increases in microbial respiration resulting from elevated CO₂, suggesting an
20 important role for O₃ in altering C flow through soils. Shafer (1988) found that O₃ tended to
21 increase the number of fungal propagules and bacteria exhibiting phosphatase activity in the
22 rhizosphere of sorghum. Ozone in combination with simulated acid rain stimulated soil
23 arylsulfatase activity (Reddy et al., 1991). The response was observed at low concentrations, but
24 was reversed at high concentrations, suggesting a threshold level of O₃, possibly involving
25 different mechanisms. Ozone significantly decreased soil microbial biomass in the fall after one
26 season of exposure in a wheat (*Triticum aestivum*) and soybean (*Glycine max*) system (Islam
27 et al., 2000). Other studies have shown shifts in microbial and fungal biomass in response to O₃
28 stress, but responses were variable (Scagel and Andersen, 1997; Yoshida et al., 2001).

29 Decreased allocation to roots associated with O₃ exposure alters N fixation in legumes and
30 actinorrhizal species. Ozone exposure was found to decrease nodulation in a number of species
31 (Manning et al., 1971; Tingey and Blum, 1973). In alder (*Alnus serrulata*), host root cells of

1 nodules showed cytoplasmic breakdown and lacked organelles when seedlings were exposed to
2 27 d of O₃ (Greitner and Winner, 1989).

3 Ozone has been shown to affect mycorrhizal colonization (Ho and Trappe, 1981; McCool
4 et al., 1982; Simmons and Kelly 1989; Adams and O'Neill, 1991; Edwards and Kelly 1992;
5 Smith and Read, 1997). Although short-term in nature, several studies have found enhanced
6 mycorrhizal short-root formation under O₃ stress. White pine (*Pinus strobus*) (Stroo et al.,
7 1988), Norway spruce (Rantanen et al., 1994), Northern red oak (*Quercus rubra*) (Reich et al,
8 1985), Douglas-fir (*Pseudotsuga menziesii*) (Gorissen et al., 1991), European silver-fir (*Abies*
9 *alba*) (Wollmer and Kottke, 1990), and Scots pine (*P. sylvestris*) (Kasurinen et al., 1999) all
10 showed some increase in mycorrhizal presence when exposed to O₃. Others have shown
11 minimal or no effects of O₃ on mycorrhizas (Mahoney et al., 1985; Meier et al., 1990;
12 Kainulainen et al., 2000). Stroo et al. (1988) found that percent infection increased from 0.02 to
13 0.06 ppm O₃, then decreased from 0.06 to 0.14 ppm; the total number of short roots were
14 unaffected, however. In cases where stimulation was observed, the response was often noted
15 shortly after initiation of exposure, often at relatively low concentrations. Good examples of this
16 transitory response can be found in results with Norway spruce and Scots pine (Ratanen et al.,
17 1994; Kasurinen et al., 1999). In these studies, O₃ increased mycorrhizal short roots initially but
18 differences were not evident by the end of the experiment.

19 There is evidence which suggests that decreased below-ground allocation associated with
20 O₃ stress alters mycorrhizal host-symbiont compatibility. Edwards and Kelly (1992) found a
21 shift in fungal morphotypes present on loblolly pine roots, even though the number of
22 mycorrhizal short roots per gram fine root was not significantly affected by O₃. Qiu et al. (1993)
23 found increased numbers of morphotypes present on O₃-sensitive loblolly pine seedlings exposed
24 to O₃. Roth and Fahey (1998) found an interaction between O₃ and acid precipitation treatments
25 on the composition of fungi forming ectomycorrhizae on red spruce saplings, possibly driven by
26 nutrient availability. Carbohydrate requirements vary among fungal species (Bidartondo et al.,
27 2001), and O₃ may affect species composition by altering carbohydrate availability in roots.
28 A shift in species dominance could lead to a change in successional patterns of mycorrhizal
29 communities.

30 In the few studies that examined root exudation in response to O₃ exposure, O₃ was found
31 to alter the quantity and quality of root exudates. McCool and Menge (1983) found a significant

1 decrease in exudation of amino acids in tomato (*L. esculentum*) exposed to 300 ppb O₃. McCrady
2 and Andersen (2000) observed increased root exudation in nonmycorrhizal wheat (*Triticum*
3 *aestivum*). No apparent change in root exudation was found in labeling studies of ECM
4 ponderosa pine (Andersen and Rygiewicz, 1995). Inconsistency in the literature probably results
5 from species differences and experimental protocols, however these examples illustrate the
6 potential effects of O₃ on rhizosphere carbon flux.

7 Decreased carbon allocation to roots of O₃-exposed plants may reduce root longevity and
8 accelerate root turnover, increasing rhizodeposition of C and N. Fine root turnover decreased in
9 mature northern red oak exposed to elevated O₃ (seasonal exposure ranging from 152 to
10 189 ppm-h), whereas seedlings did not show any reduction in turnover (Kelting et al., 1995).
11 King et al. (2001) found a trend toward decreased live root biomass and increased dead root
12 biomass in aspen exposed to O₃ in a FACE study, suggesting possible changes in both
13 production and longevity.

14 Other studies also suggest that O₃ alters C flux to soils, resulting in changes in CO₂ efflux
15 from soils. Both root respiration and soil CO₂ efflux decreased from loblolly pine seedlings
16 exposed to O₃ (Edwards, 1991). Soil CO₂ efflux increased in response to O₃ in ponderosa pine
17 seedlings (Andersen and Scagel, 1997; Scagel and Andersen, 1997). No direct assessments of
18 hyphal growth and turnover in response to O₃ stress have been conducted. Ozone decreased C
19 allocation to extrametrical hyphae of a ponderosa pine mycorrhiza, which might be expected to
20 decrease growth and increase hyphal turnover (Andersen and Rygiewicz, 1995).

21 22 **COMMUNITY DYNAMICS, PHYSICAL STRUCTURE**

23 One of the best-documented examples of change in long-term forest community dynamics
24 of dominant overstory trees occurred in the San Bernardino Mountains between 1968 and 1974
25 (reported in Miller, 1973; Miller et al., 1989). Plots were recently re-inventoried ~25 years after
26 establishment (Arbaugh et al., 2003). Of the six codominant canopy species, white fir showed
27 the greatest change, increasing in both numbers and bole growth for a 286% change in basal
28 area/ha in the San Bernardino Mountains. Sugar pine basal area also increased significantly (by
29 334%), but this species represents only a small portion (1%) of the total basal area of the forest
30 sampled. The most sensitive species (Miller et al., 1983), ponderosa and Jeffrey pine, had the
31 lowest increase in basal area/ha (76 and 62%, respectively). These two species represented 72%

1 of the basal area/ha of all stands inventoried. Ponderosa pine had the greatest mortality rate of
2 all canopy species inventoried (46%), followed by white fir and black oak (35% and 33%),
3 Jeffrey pine (29%) and incense cedar and sugar pine (both 7%). In moist sites (at the western
4 end of the San Bernardino Mountains), there was significant recruitment of incense cedar, white
5 fir, and sugar pine. Only one study directly attributed tree mortality to O₃ exposure: it
6 accounted for 7% of mortality in the Sierra and Sequoia National Forests (Carroll et al., 2003).

7 Species diversity in the understory can be quite large, making studies of O₃ effects on
8 understory community dynamics very challenging. However, there have been some attempts to
9 quantify understory responses, ranging from describing relative sensitivity to their visible
10 symptoms (Treshow and Stewart, 1973; Temple, 1999) to very complex measures of community
11 structure and composition (Westman, 1979, 1981). The lowest percentage cover and lowest
12 species diversity in California coastal sage scrub was correlated with the highest O₃ exposures as
13 estimated by extrapolation from the closest air monitoring stations (Westman 1979). The
14 understory also has the potential to influence responses to O₃ of dominant keystone species, as
15 has been shown in controlled experiments with both *ponderosa pine* (Andersen et al., 2001) and
16 *loblolly pines* (Barbo et al., 2002). Barbo et al., (1998) exposed an early successional forest
17 community to ambient air, charcoal-filtered air, non-filtered air, and 2× ambient in the
18 Shenandoah National Park. They found changes in species performance, canopy structure,
19 species richness and diversity index consistent with the view that O₃ can induce a shift in
20 vegetation dominance and community structure.

21 There have been few studies evaluating the effect of O₃ exposure on the physical structure
22 of natural ecosystems. Despite an extensive array of allometric equations for conifers in the
23 west on United States (Ter-Mikaelian and Korzukhin, 1997), none appear to predict individual
24 tree shape in a site of moderate O₃ exposure, suggesting that O₃ may effect allometry (Grulke
25 et al., 2003a). Canopy structural changes are also implied by the measure of canopy
26 transparency used in the FHM assessment. The loss of epiphytic lichens within the canopy is a
27 clear example of plant community structural change occurring along an O₃ gradient (Nash and
28 Sigal, 1999; Zambrano et al., 2002).

29 As of yet, there have been no comprehensive studies on the effects of O₃ on structural or
30 functional components below-ground (Andersen, 2003). Phillips et al. (2002) found evidence
31 for changes in the bacterial and fungal biomass below *Populus tremuloides* and

1 *P. tremuloides/Betula papyrifera* stands exposed to elevated O₃. Subsequent study showed that
2 O₃ exposure decreased cellobiohydrolase activity in the soil microorganisms, driving the change
3 in the microbial community (Larson et al., 2002).
4

5 **9.7.4.2 Species and Populations**

6 Ozone can affect species and populations of species comprising ecosystems through
7 changes in population size, genetic diversity, population structure and/or dynamics, and habitat
8 suitability (Young and Sanzone, 2002). For example, if individuals of a species are lost due to
9 O₃ exposure, population size declines. Often very young (e.g., conifer seedlings, see Section
10 9.7.4.3 below) and old individuals differ in their sensitivity, so that population structure also will
11 be altered by O₃ exposure. If resource allocation to reproductive output is altered by O₃
12 exposure, population dynamics will be altered. Communities dominated by O₃ sensitive species
13 in the canopy or understory may be altered sufficiently for the habitat to become unsuitable for
14 other species. Genetic selection acts on the individual plant, which represents a certain
15 proportion of the populations' genetic variation. If an O₃ sensitive individual succumbs through
16 multiple stresses, including O₃ stress, the genetic variation represented in the population
17 generally declines, unless sensitive individuals have low inherent genetic variability (e.g.,
18 Staszak et al., 2004).

19 While the concept of natural selection induced by O₃ exposure and related changes in
20 natural plant communities has been around for a long time (Dunn, 1959; Miller et al., 1972), the
21 concept of evolution of O₃ tolerance is still not widely accepted. The unequivocal demonstration
22 that considerable genetic variation in O₃ resistance exists within and between plant populations,
23 and that ambient levels of O₃ may differentially affect fitness-related traits (i.e., growth, survival,
24 and fecundity), suggests that O₃ may potentially drive the natural selection for resistant
25 individuals. Dunn (1959) presented circumstantial evidence that ambient O₃ in the Los Angeles
26 area was high enough to drive the selection of O₃ resistant *Lupinus bicolor* genotypes. Since
27 Dunn's (1959) research on O₃-induced population changes, researchers have demonstrated
28 differences in O₃ tolerance among other plant populations. In the devastating forest decline
29 southeast of Mexico City, the remaining trees (primarily *Abies religiosa* in the "cemetery
30 forests") appear to be less affected by foliar injury than trees that were lost (Alvarado-Rosales
31 and Hernandez-Tejeda, 2002), despite continued, high O₃ exposures (Bravo-Alvarez and

1 Torres-Jardon, 2002). However, even the most convincing work in this field (Berrang et al.,
2 1986, 1989, 1991), with *Populus tremuloides*, where a strong correlation between visible foliar
3 injury after O₃ exposure and maximum O₃ concentration at the origin of the population was
4 shown (Berrang et al., 1991), a change in gene frequency at any one site over time has not yet
5 been demonstrated (Bell et al., 1991; Reiling and Davison, 1992b). Furthermore, the selection
6 intensity of O₃ has been questioned (Bell et al., 1991; Taylor and Pitelka, 1992; Taylor et al.,
7 1994) and the emergence of O₃ exposure since the 1950's as an environmental stressor may not
8 have been long enough to affect tree populations with long generation times (Barrett and Bush,
9 1991).

10 The loss of O₃-sensitive individuals results in natural selection favoring O₃-tolerant species
11 (Bradshaw and McNeilly, 1991). Increased levels of mortality of O₃-sensitive individuals have
12 occurred for *Pinus jeffreyi* and *P. ponderosa* exposed to ambient O₃ along the western slope of
13 the Sierra Nevadas (Peterson et al., 1987; Miller et al., 1998), for *Pinus strobus* exposed to
14 ambient O₃ in southern Wisconsin (Karnosky, 1981), and for *P. ponderosa* in the San Bernardino
15 Mountains (Carroll et al., 2003). In these examples, individuals that consistently had greater
16 foliar injury and lower needle retention were lost in repeated surveys. Ozone-induced loss of all
17 individuals except the most tolerant and breeding among the surviving individuals to yield more
18 more tolerant populations has not yet been demonstrated for plants exposed to O₃, except for the
19 relatively short term (2 years) adaptation exhibited in *Trifolium repens* (Heagle et al., 1991) and
20 *Plantago major* (Davison and Reiling, 1995). Heagle et al. (1991) were able to show the
21 adaptation of a *Trifolium repens* population to elevated O₃ in just two growing seasons.
22 Similarly, Davison and Reiling (1995) compared the O₃ resistance of *P. major* populations
23 grown from seed collected from the same sites over a period of increasing O₃. The two
24 independent populations studied exhibited increased O₃ resistance, consistent with the idea of
25 selection for O₃ tolerance. Using random amplified polymorphic DNA primers, this team also
26 showed that the later populations are subsets of the earlier ones, consistent with in-situ evolution
27 rather than with catastrophic loss and replacement of the populations (Wolff et al., 2000). The
28 problem is determining whether spatial patterns in O₃ resistance and changes in time are casually
29 related to O₃, because there were very strong correlations with other factors (Reiling and
30 Davison, 1992a; Davison and Barnes, 1998). The potential for evolution of O₃ resistance has
31 been clearly demonstrated by Whitfield et al. (1997) in their study of O₃ selection of common

1 plantain (*Plantago major* L.), where they showed that within a matter of a few generations, it
2 was possible to increase O₃ resistance in an initially O₃-sensitive population. Wild radish
3 (*Raphanus sativus* L.) developed O₃ resistance after only one generation of exposure to O₃
4 (Gillespie and Winner, 1989).

5 A third independent line of research suggesting O₃ may be affecting the genetic diversity of
6 wild plant populations was presented by Paludan-Müller et al. (1999) who showed that northwest
7 European provenances of European beech (*Fagus sylvatica* L.) were more sensitive to O₃ than
8 were southeast European provenances which had experienced higher O₃ levels. Recent research
9 on the genetic structure of 50-year-old *ponderosa pines* in the San Bernardino Mountains
10 suggests that distinct differences in frequency of some alleles and genotypes occurred, with the
11 O₃-tolerant trees being more heterozygous (Staszak et al., 2004). While both of these studies
12 were only correlational, they are consistent with previous studies of this type suggesting
13 O₃-induced population changes. Again, other environmental stressors besides O₃ exposure could
14 have been involved in effecting change within these populations.

15 Natural selection for O₃ tolerance can also be facilitated by reductions in fitness related to
16 lower seed yields of O₃-sensitive species or individuals. The impacts of O₃ on reproductive
17 development, recently reviewed by Black et al. (2000) can occur by influencing: (1) age of
18 flowering, particularly in long-lived trees that often have long juvenile periods of early growth
19 without flower and seed production; (2) flower bud initiation and development; (3) pollen
20 germination and pollen tube growth; and (4) seed, fruit, or cone yields and seed quality
21 (Table 9-24). In addition, vegetatively propagated species can have lower numbers of
22 propagules under elevated O₃ conditions (Table 9-24).

23 Several studies suggest that reproductive structures are clearly sensitive to O₃ and that O₃
24 can affect fitness of plants by affecting either the sporophytic or gametic generations. Decreased
25 numbers of flower spikes and seed capsules per plant were found for plantain growing under
26 elevated O₃ (Reiling and Davison, 1992b; Pearson et al., 1996; Lyons and Barnes, 1998).
27 Similar responses were seen for *Brassica campestris* L. plants exposed to a single dose of
28 100 ppb O₃ for 6 h. Stewart et al. (1996) and Bosac et al. (1998) reported an increase of flower
29 bud abortion for oilseed rape (*Brassica napus* L.) similarly exposed to a short duration of
30 elevated O₃. Floral initiation period can be delayed in O₃-sensitive plants, as was described for
31 dogbane (*Apocynum androsaemifolium*) grown under ambient O₃ in the eastern United States.

1 In one of the few comparisons of whole plant O₃ sensitivities with that of male gametophytes,
2 Hormaza et al. (1996) found a high correlation of relative O₃ sensitivity of pollen tube elongation
3 with that of O₃ effects on net photosynthesis and relative growth rates for 6 species of fruit trees.

4 Clearly, the concept of O₃-induced genetic change is an area that needs additional research
5 attention. Repeated collections over time from wild populations receiving high O₃ exposures to
6 examine population responses and relative sensitivity changes, the sampling of genetic diversity
7 along known O₃ gradients, and the use of modern biotechnological approaches to characterize
8 and quantify genetic diversity are useful approaches to test for O₃-induced impacts on diversity
9 in natural ecosystems.

11 **9.7.4.3 Organism Condition**

12 **PHYSIOLOGICAL STATUS**

13 The generalized effects of O₃ exposure on plants are well known, and have been reviewed
14 from several viewpoints over the last decade (De Kok and Tausz, 2001; Heath and Taylor, 1997;
15 Pell et al., 1997; Schraudner et al., 1997; Matyssek et al., 1995; Darrall 1989; Reich, 1987; U.S.
16 Environmental Protection Agency, 1996). The topic of individual species response and
17 modification of response by other factors has been addressed thoroughly in Sections 9.4.3 and
18 9.4.4 of this chapter. Here, physiological changes in response to O₃ that have been hypothesized
19 to lead to changes in ecosystem structure or function are highlighted.

21 *Above-Ground Responses*

22 The first critical step leading to O₃ response is uptake and movement of O₃ by the leaves,
23 leading to changes in C and nutrient relations that are thought to alter plant growth and
24 competitiveness (see Section 9.3). Ozone enters leaves through stomates, reacts with cell walls or
25 membranes, and starts a series of adverse reactions. Cuticular uptake of O₃ is believed to be
26 negligible (Kerstiens and Lendzian, 1989; Coe et al., 1995). Once inside the leaf, O₃ and its
27 byproducts lead to membrane disruption, chlorophyll breakdown, and decreased Rubisco levels
28 (Schweizer and Arndt, 1990). In turn, photosynthesis is decreased, as is stomatal conductance
29 (Weber et al., 1993). Also, O₃ often leads to increased maintenance respiration, decreased foliar
30 nutrient content, and imbalances in tissue nutrient content and retention. When photosynthetic
31 pigments have been damaged, the pigment must be fully broken down (and/or new N and Mg

1 must be taken up and transported to the leaf) for the pigment to be regenerated (Bjorkman and
2 Demmig-Adams, 1995). Ozone exposure alters within-plant priorities for resources: less C is
3 available for allocation to roots and spring regrowth, and less foliar biomass is retained. At the
4 whole-organism level, O₃ exposure decreases root mass (Grulke et al., 1998) and radial bole
5 growth (Peterson et al., 1991; Muzika et al., 2002) with little impact on height growth. Visible
6 symptoms of O₃ injury vary between species and genotypes but often include upper leaf surface
7 stipple, chlorotic mottle, or large bifacial blotches of necrotic tissue. Premature senescence is
8 typical of almost all O₃-induced foliar damage. All of these changes can alter the plant's ability
9 to function in a broader ecosystem context.

10 The underlying mechanisms of O₃-injury response in conifers, broadleaf deciduous trees,
11 and herbaceous species are assumed to be similar. However, several differences in long-lived
12 species are important at the ecosystem level. Most of the research on O₃ effects has been
13 conducted on herbaceous species (i.e., crops). Although a number of native herbaceous species
14 have been identified as O₃-sensitive, there are no published physiological studies on the effect of
15 O₃ exposure on herbaceous or shrub species in situ. In natural ecosystems, the majority of
16 species are not annuals, unless the system is highly disturbed. Nonetheless, response of crop
17 species to elevated O₃ may be used as an analog for native annual response: phenological
18 staging is accelerated (soybeans; Booker et al., 2004), thus "avoiding" additional O₃ exposure.

19 Conifers have roughly half the stomatal conductance of deciduous broadleaf trees (Reich,
20 1987), leading to proportionally less O₃ uptake at the same O₃ exposure level. Yet, except for
21 species of larch, individual conifer needles are longer-lived and active over a greater portion of
22 the year. Therefore, needle longevity can also work against the tree by increasing cumulative O₃
23 exposure and exposure to other stressors. Increased needle longevity is not always a
24 disadvantage, for example, conifers are physiologically active in early spring and late fall, during
25 times of lower oxidant concentrations. These periods can contribute significantly to a net
26 positive annual C balance and from the standpoint of nutrient storage, are important in reparation
27 responses to pollutants. Patterson and Rundel (1995) reported that Jeffrey pine had significant
28 stomatal opening (one third that of a typical summer day) in mid-winter with snow on the
29 ground. At least pole-sized and larger trees can mitigate reductions in C acquisition due to
30 oxidant exposure in the summer with C assimilation on favorable days in the winter. The
31 interaction of environmental factors, plant phenology (the timing of growth events; birth and

1 mortality of plant parts), physiological status (nutritional or moisture status; dormant or active
2 growth within the year), and tree age (interannual differences in resource acquisition and
3 allocation) all contribute to the complexity of long-lived species (and hence ecosystem) response
4 to O₃ exposure.

5 One widely observed response to O₃ exposure is premature leaf loss. As noted above,
6 premature leaf loss may reduce O₃ uptake during high-O₃ years, but it has several negative
7 consequences. Early leaf loss results in reduced C uptake through photosynthesis. Premature
8 needle loss also results in less N retranslocation compared to normally senescing leaves, which
9 reduces whole plant N balance (Fenn and Dunn, 1989) and carbohydrate availability for
10 overwinter storage (Grulke et al., 2001). Because of such effects accumulated over several years
11 of O₃ exposure, subsequent-year C and N reserves can be affected (Andersen et al., 1991).

12 Conversely, a series of drought years can decrease O₃ uptake, as well as reduce C and
13 nutrient acquisition, altering resource allocation to defenses (e.g., antioxidants) (Grulke et al.,
14 2003b) (or resins) against insect infestation, rendering the tree more susceptible to O₃ injury.
15 Conifers have thicker cuticles than either broadleaf deciduous or herbaceous species. Continued
16 O₃ exposure may compromise cuticular integrity (Percy et al., 1994). Once cuticular integrity is
17 breached, individual leaves (needles) are likely to be excised, thus contributing generally to
18 defoliation and reduced C acquisition.

19 With the exception of the extensive research conducted on mature tree response to O₃
20 exposure in California forests (Peterson et al., 1987, 1991, 1995, Arbaugh et al., 1998; Grulke
21 1999; Grulke and Balduman 1999; Grulke et al., 1996, 1998, 2001, 2002b, 2003a,b, 2004;
22 Weiser et al., 2002), the vast majority of studies of O₃ effects on forest trees have been
23 conducted on young seedlings (Chappelka and Samuelson, 1998) and little is known about
24 acclimation to O₃ (Skärby et al., 1998). Chamber exposure studies can be used to document
25 foliar symptoms and develop response variables for the whole plant. These response variables
26 can then be field tested on mature trees using correlative analyses (e.g., Grulke and Lee, 1997;
27 Grulke et al., 2003b). Without the initial work in chamber exposure studies, field responses to
28 O₃ exposure would be difficult to verify and distinguish from other concurrent stressors.

29 Predicting mature tree responses to O₃ solely from seedling response studies is complex,
30 because seedlings or saplings do not necessarily respond to O₃ in the same way as mature trees
31 (Norby et al., 1999; Karnosky, 2003). Ozone has been found to have stronger effects on leaf

function in younger rather than older trees (Kolb and Matyssek, 2001). Each component physiological attribute “matures” at a different rate. Gas exchange patterns differ between seedlings and mature trees. For example, leaf respiration of juvenile ponderosa pine was greater than that of mature trees (Momen et al., 1996). In the conifers tested, the highest gas exchange rates (and by inference stomatal uptake of O₃) are found in seedlings (e.g., in scions of red spruce, Rebbbeck and Jensen, 1993; giant sequoia, Grulke et al., 1994; ponderosa pine, Grulke and Retzlaff, 2001; and Norway spruce, Wieser et al., 2002). Patterns of biomass (Grulke and Balduman, 1999) and carbohydrate allocation (Grulke et al., 2001) differs between immature and mature trees. Pole-sized trees had greater reduction in root, foliar, and bole carbohydrate concentrations than did old growth trees. Antioxidant defenses vary with both tree age and needle age (Tegisher et al., 2002). Based on all attributes measured in both ponderosa pine and giant sequoia, the youngest tree age considered representative of mature trees was 20 years old (Grulke et al., 1996; Grulke and Retzlaff, 2001). In some broadleaf deciduous tree species, seedlings are more conservative, and mature trees have greater gas exchange rates, as is the case for *Quercus robur* (Edwards et al., 1994; Kelting et al., 1995; Samuelson and Kelly, 1996, 1997) and *Fagus sylvatica* (Braun et al., 1999). In another broadleaf deciduous tree species (cherry, *Prunus serotina*), gas-exchange rates of seedlings were faster, but total O₃ flux to leaves of seedlings was lower than that of mature trees due to differences in leaf ontogeny (Frederickson et al., 1995, 1996).

Nitrogen deposition modifies the effects of oxidant exposure through several offsetting physiological mechanisms (see Section 9.4.4). Nitrogen deposition, in wet or dry particulate form, ultimately increases site fertility, but increased soil N availability decreases C allocation to roots, further exacerbating the effects of O₃ exposure on roots (Grulke et al., 1998). Increased N availability also increases foliage turnover: fewer needle age classes are retained (Gower et al., 1993). Therefore, the combination of both increased N and O₃ exposure increases foliar turnover. Finally, N deposition and increased plant N nutrition can increase stomatal conductance, leading to increased O₃ uptake. Alternatively, increased N counteracts the effect of O₃ on photosynthesis by increasing photosynthetic pigments and enzymes. Nitrogen deposition may mitigate the degree of foliar injury from oxidant pollution via higher available N for reparation of photosynthetic pigments. Nitrogen amendments also modify the antioxidant defense system in complex ways (Polle, 1998).

Attributes of O₃ injury to trees (foliar injury, needle retention, and canopy transparency), as well as presence of pathogens and insect infestation, are routinely inventoried in established sample plots distributed on Federal lands across the United States (Forest Health Protection, USDA Forest Service). Foliar injury to several widespread, herbaceous species nationally recognized as sensitive (bioindicators) is also assessed (NPS, 2003). This assessment is part of a larger assessment of forest tree growth and dynamics (the Forest Inventory and Analysis Program; Smith, 2002; Smith et al., 2003). Risk of O₃ injury is then estimated for the dominant forest tree species in the sample plots. For example, 12% of sampled black cherry (*Prunus serotina*), 15% of loblolly pine (*Pinus taeda*), and 24% of sweetgum (*Liquidambar styraciflua*) were found to be in the highest risk category in the northeast and mid-Atlantic states (Coulston et al., 2003). In the Carpathian Mountains, 12 to 13% of all trees (broadleaf and coniferous) have greater than 26% crown defoliation (Badea et al., 2002). In general, broadleaves (primarily beech) trees were less affected (8 to 45%) than spruce (up to 37%) and fir (up to 50%) (Grodzinska et al., 2002). Ozone injury was directly correlated with cumulative O₃ exposure in the Sierra Nevada Mountains (Arbaugh et al., 1998); with the best correlation being found across sites where > 90% of the trees had O₃ injury. Although direct links of visible foliar symptoms induced by O₃ to adverse effects on biomass are not always found, visible foliar symptoms have been linked to decreased vegetative growth (Peterson et al., 1987; Karnosky et al., 1996; Somers et al., 1998), as well as reproductive function (Black et al., 2000; Chappelka, 2002).

Foliar O₃ injury has also been associated with adverse effects on competitive ability and survival in forest communities (Karnosky, 1981; McDonald et al., 2002; Karnosky et al., 2003b). Competition can alter organism condition and affect susceptibility to O₃. Ponderosa pine seedlings were more susceptible to O₃, as determined by decreased plant biomass, when grown in competition with blue wild-rye grass (Andersen et al. 2001). Similarly, the magnitude of O₃ effects on height and diameter growth depended on the competitive status of *Populus tremuloides* trees (McDonald et al., 2002). These studies show the importance of including competition as a concurrent stressor in assessing whole plant responses to O₃. The age of the community ("time since disturbance") may also affect the ability of individuals to effectively respond to O₃. Unfortunately, the vast majority of O₃ studies have been conducted on open-grown plants, often grown in pots where competition is absent both above and below ground.

Clearly, age-dependent O₃ responsiveness and juvenile-mature correlations remain important research questions in attempting to scale up to ecosystem level responses. Patterns of allocation between root, stem, and leaf differ between immature and mature trees. Tree architecture varies with tree age, and leaf area distribution in space and time may change in response to elevated O₃. All of these factors influence gas exchange in the canopy. Furthermore, there may be few generalities that can be made from seedling to mature tree response to O₃ within a plant functional group (Norby et al., 1999; Karnosky, 2003). Consequently, modeling ecosystem response is limited to either dealing with mono-specific plantations or assigning average responses to a mix of species.

Below-Ground Responses

The effect of O₃ on the soil ecosystem is thought to occur through physiological changes in the root and interactions with soil organisms (Andersen, 2003). Comparatively little is known about how changes in root growth and metabolism are translated through the soil food web, resulting in changes in soil and hence, ecosystem processes. An overview of physiological changes likely to lead to changes at the ecosystem level is provided below.

Ozone stress decreases carbon allocation to roots (Manning et al., 1971; McLaughlin and McConathy, 1983; McCool and Menge, 1983; Cooley and Manning, 1987; Gorissen and van Veen, 1988; Spence et al., 1990; Gorissen et al., 1994; Rennenberg et al., 1996; U.S. Environmental Protection Agency, 1996). Since roots are often dependent on current photosynthate for their structural development (van den Driessche, 1978; Ritchie and Dunlap, 1980; Marshall and Waring, 1985; van den Driessche, 1991), C-limiting stresses such as O₃ can have rapid and significant effects on root growth. In many cases, decreased allocation to roots in response to O₃ occurs quickly, with reductions in root growth occurring within one growing season (Gorissen and van Veen, 1988; Spence et al., 1990; Gorissen et al., 1991; Andersen and Rygielwicz, 1991, 1995; U.S. Environmental Protection Agency, 1996). Decreased C allocation below ground is often associated with decreased root-shoot ratio, but observed responses in root-shoot ratio are highly variable owing to several factors including intra- and interspecies variation, culture conditions, and ontogenetic drift (Reich, 2002). Root-shoot ratio is a point-in-time measurement that does not include C lost to exudation, respiration or turnover.

1 Therefore, biomass and ratios of biomass (such as root-shoot ratio) do not necessarily reveal
2 physiological changes in response to O₃ stress.

3 Decreased C acquisition leads to reduced carbohydrate levels and storage pools in
4 O₃-exposed plants (Tingey et al., 1976; Ito et al., 1985; Cooley and Manning, 1987; Rebbeck
5 et al., 1988; Gorissen et al., 1994; Andersen et al., 1997; McLaughlin et al., 1982). Although it
6 is difficult to quantify changes in the field, Grulke et al. (1998) found decreased medium and
7 fine root biomass with increased pollutant load across an O₃ gradient in southern California.
8 Coarse and fine root starch concentrations also were lowest in mature trees at the most polluted
9 site (Grulke et al., 2001). The effects of O₃ could not be completely separated from other known
10 stresses across the pollutant gradient, but it appeared that O₃ was an important factor in the
11 patterns observed.

12 Decreased storage pools can lead to carry-over effects on root growth that are compounded
13 over time. Decreased carbohydrate storage pools were associated with decreased root growth
14 during the spring following exposure to O₃, even in the absence of additional O₃ exposure
15 (Andersen et al., 1991, 1997). Decreased spring root growth was attributed to decreased stored
16 C reserves as well as to premature loss of older foliage age classes during the previous fall.
17 Aside from the loss of photosynthetic surface area associated with premature senescence, early
18 loss of foliage in the fall occurs when allocation to roots is at a maximum in many species
19 (Kozłowski and Pallardy, 1997). Older needle age classes preferentially allocate photosynthate
20 basipetally to stems and roots (Rangnekar et al., 1969; Gordon and Larson, 1970), and the loss of
21 older needles in the fall during allocation to root growth and storage, and in the spring during
22 periods of root growth, preferentially impacts roots and root processes.

23 Ozone has also been shown to affect root metabolism as evidenced by changes in root
24 respiration. Edwards (1991) found decreased root and soil CO₂ efflux during a 2-year exposure
25 of loblolly pine to O₃. Fine root respiration increased in mature red oak exposed to O₃, while
26 total soil CO₂ efflux increased in the spring and decreased in the summer and fall (Kelting et al.,
27 1995). The authors attributed increased root respiration to increased nutrient uptake in support
28 of increased demands in the shoot. Ozone decreased root system respiration in aspen after 12
29 weeks of exposure, but the decrease was closely associated with decreased root biomass and
30 probably not metabolic processes (Coleman et al., 1996). Whether other metabolic shifts occur
31 in the roots of plants exposed to O₃ needs to be examined.

Measurable effects on roots may occur before effects on shoots are observed because shoots have immediate access to C for repair and compensation whereas roots must compete with shoots for C. Mortensen (1998) found decreased root but not shoot growth in *Betula pubescens* at O₃ exposures of 42 nMol mol⁻¹L (applied 12 h d⁻¹), whereas both root and shoot growth were reduced at higher exposures. Chromosomal aberrations were found in root tips of Norway spruce exposed to O₃, even in the absence of biochemical changes in needles (Wonisch et al., 1998, 1999). Using relatively high O₃ concentrations (0.15 ppm O₃ 6 h d⁻¹), Hofstra et al. (1981) found metabolic changes in *Phaseolus vulgaris* root tips prior to the development of leaf injury. Morphological changes in root tips occurred within 2 to 3 days, and metabolism declined within 4 to 5 days of initiation of O₃ exposure.

Feedback signals from roots can influence the degree of O₃ response. Stolzy et al. (1964) exposed tomato roots (*Lycopersicon esculentum*) to periods of anoxic conditions and followed a change in leaf susceptibility to O₃. An exposure of roots to low oxygen conditions for 3 h did not alter photosynthesis, but foliar damage was decreased when the roots were subsequently exposed to O₃. In this case, a signal originating in the root appeared to alter leaf sensitivity to O₃, the signal possibly being hydraulic in nature and leading to decreased O₃ uptake.

SYMPTOMS OF DISEASE OR TRAUMA, SIGNS OF DISEASE

Although insects and diseases are dynamic components of forest ecosystems, trees can be especially susceptible to outbreaks due to the presence of multiple stressors such as drought and pollutant exposure. Ozone can have direct effects on insect or disease organisms, indirect effects on the insect or pathogen through changes to the host, and direct or indirect effects on natural enemies of the insect or pathogen (Pronos et al., 1999). A full discussion of O₃ effects on insect and pathogen interactions can be found in Section 9.4.

Although the multitude of interacting factors makes it difficult to identify causative factors in the field, some recent examples suggest a role for O₃ in the timing or magnitude of disease attacks in the field. After periods of drought stress (such as 1995 in central Europe), the incidence of bark beetle (*Ips spp.*) appears to increase. In 1998 and 1999, the mean daily capture of *Ips* was lowest in plots with low O₃ exposure; the converse was also true (Grodzki et al., 2002). Elevation confounded the relationships, but the differences in *Ips* frequency in relation to O₃ concentrations were highly significant at lower elevations. In the Valley of Mexico, a 1982

to 1983 drought was documented, but not described as precipitating a bark beetle attack in the early 1980's. However, a link between air pollutant exposure and bark beetle attacks were implicated, because attacked trees were already O₃-stressed at the time of the bark beetle attack (Alvarado-Rosales and Hernandez-Tejeda, 2002).

An early study showed that oxidant exposure predisposed *ponderosa pine* to the root pathogen *Fomes annosus* (James et al., 1980). Both root diseases (Pronos et al., 1999) and O₃ exposure (Grulke and Balduman, 1999) can each reduce root biomass, leading to increased drought stress, insect attack, and subsequent windthrow or death. Trees may take several years to die, and the patterns of precipitation and annual total rainfall interact to drive the level of drought stress experienced by the tree (Pronos et al., 1999). Additional research is necessary to fully understand the complex interactions occurring between O₃ stress and other biotic stresses.

9.7.5 Ecosystem, Chemical, and Physical Characteristics (water, soil)

9.7.5.1 Nutrient Concentrations, Trace Inorganic and Organic Chemicals

Ozone exposure reduces the nutritional content of tissues, as well as causing elemental imbalances. Foliar nutrient content may be too high (toxic) or too low (deficient), but the relative amounts and ratios among all nutrients can also result in imbalances.

Although N deposition and foliar N content increased with O₃ exposure in the San Bernardino Mountains, K, Mg, Fe, and Al were all also higher in ponderosa pine at sites more exposed to air pollution (Poth and Fenn, 1998). Trees with greater foliar injury (due to O₃ exposure) had higher current year needle concentrations of P, K, Zn, and Fe than trees at the same site that were less injured. In drought-stressed ponderosa pine with O₃ exposure, foliar N was also elevated and retained in the remaining needles (Temple et al., 1992). At a relatively clean site in the eastern San Bernardino Mountains, N, P, and K were efficiently readsorbed, but P remaining in the foliage was relatively high compared to defined thresholds. The fact that other elements were modified besides the N being deposited emphasizes the degree of chemical imbalance in the tissue. Foliar micronutrients were within the normal ranges reported for ponderosa pine (Poth and Fenn, 1998; Powers, 1981). Because both N deposition and O₃ exposure reduce root biomass, it was unlikely that the foliar nutrient content was higher due to greater uptake. Instead, it appears that retranslocation from senescing tissue was responsible. Across a pollution gradient in the Carpathian Mountains in eastern Europe, only S/N (expected

1 due to high S deposition) and Fe/Mn ratios were out of balance relative to established norms.
2 The S was relatively high due to SO₂ deposition, and the Fe was relatively high due to smelter
3 plumes. No imbalances could be directly attributed to O₃ exposure (Mankovska et al., 2004).
4

5 **9.7.6 Ecological Processes**

6 **9.7.6.1 Energy Flow**

7 All green plants generate and use energy-containing C compounds through the processes of
8 photosynthesis and respiration. Whole-plant C uptake is dependent on photosynthesis rates, leaf
9 area, and leaf phenology. The effects of O₃ at the site of action in the leaf are discussed in
10 Section 9.3. Here, the main focus is on whole-plant carbon dynamics resulting from changes in
11 C acquisition or use under O₃ stress.

12 In natural ecosystems, O₃ has been shown to depress photosynthesis in sensitive tree
13 species including *Pinus ponderosa* (Miller et al., 1969; Weber et al., 1993; Takemoto et al.,
14 1997; Grulke et al., 2002b) and *Populus tremuloides* (Coleman et al., 1995a; Yun and Laurence,
15 1999; Noormets et al., 2001a,b; Sharma et al., 2003). In a study of mature Jeffrey pine, trees in
16 mesic microsites had greater O₃ uptake over the growing season in comparison to trees in xeric
17 microsites (Grulke et al., 2003a) and greater O₃ uptake was correlated with lower mid canopy
18 needle retention, lower branch diameters, and lower foliar N content (Grulke et al., 2003a).
19 Chamber studies have also shown negative effects of O₃ on tree seedling canopy structure
20 (Dickson et al., 2001) and leaf area (Neufeld et al., 1995; Wiltshire et al., 1994). It is well
21 known from chamber and field studies that O₃ exposure is correlated with lower foliar retention
22 (Miller et al., 1963; 1972; Karnosky et al., 1996; Grulke and Lee, 1997; Pell et al., 1999; Topa
23 et al., 2001).

24 In contrast to the relatively consistent findings for photosynthesis, O₃ effects on respiration
25 have been more variable. Stem respiration was unaffected by O₃ exposure (Matyssek et al.,
26 2002), suggesting that construction costs of new stems are not affected by O₃. However, the bole
27 represents a relatively large storage pool of carbohydrates in mature trees, and the timing of
28 phenological events among individual trees may help to confound the ability to statistically
29 detect differences in stem respiration across pollutant gradients (Grulke et al., 2001). Below-
30 ground respiration has been found to both increase and decrease in response to O₃, depending on
31 the approach and timing of CO₂ measures (Coleman et al., 1996; Andersen and Scagel, 1997;

1 Scagel and Andersen, 1997; King et al., 2001). The decreased soil respiration is thought to be
2 due to reduced root growth under O₃ exposure, but could also be partially explained by
3 decreased microbial respiration in response to O₃. Additional research is necessary to identify
4 the role of O₃ in affecting root vs. heterotrophic respiration, particularly over long time intervals.

5 Carbohydrate availability and use influence the degree to which plants respond to O₃
6 exposure. A model simulation of the effect of O₃ exposure on bole growth of *Pinus ponderosa*
7 showed a 15% reduction in mass (Weber and Grulke, 1995), largely influenced by differences in
8 carbohydrates allocated and partitioned in repair processes elsewhere in the tree. Foliar
9 respiration is thought to increase under elevated O₃ as maintenance costs (energy needs) of
10 leaves damaged by O₃ are higher than normal (Grulke and Balduman, 1999; Noormets et al.,
11 2001b), but differences in foliar respiration are subtle and difficult to detect statistically. Foliar
12 carbohydrate studies also suggest that more C is used under O₃ stress for repair processes (Topa
13 et al., 2001; Grulke et al., 2001) which would result in increased respiration. Ozone exposure
14 also reduced enzymatic activities of carbohydrate metabolism related to the breakdown of
15 sucrose (Einig et al., 1997). Changes in soil respiration in response to O₃, even though O₃ does
16 not penetrate into the soil, illustrates the tight coupling of plant C balance and soil biota and
17 illustrates the potential role O₃ plays in altering ecosystem C balances (Andersen, 2000).

18 Ozone can affect plant allometry through changes in energy use, potentially affecting net
19 primary production (NPP) at larger scales. The net effect of O₃ impacts on photosynthesis and
20 respiration for sensitive components of natural ecosystems is that height growth (Isebrands et al.,
21 2001; Oksanen, 2003b); and radial growth (Peterson et al., 1987, 1991; Isebrands et al., 2001;
22 Terrazas and Bernal-Salazar, 2002; Oksanen, 2003b) can be negatively affected by O₃. This has
23 been extrapolated to decreased NPP (Hogsett et al., 1997; Laurence et al., 2000).

24 Energy flow in plant communities can be altered by O₃ through changes in C allocation.
25 It is well known that elevated O₃ affects C allocation to roots (Coleman et al., 1995b; Andersen
26 et al., 1997; Grulke et al., 1998; Grulke and Balduman, 1999; Grulke et al., 2001) by decreasing
27 or inhibiting phloem loading of carbohydrates (Grantz and Farrar, 1999; Landolt et al., 1997), or
28 of carbohydrate metabolism (Einig et al., 1997). This leads to depressed root growth (Andersen
29 et al., 1991; Coleman et al., 1996; Grulke et al., 1998) and the potential for plant communities to
30 have an increased susceptibility to drought through altered root-shoot balance. Furthermore, it
31 can negatively affect below ground food webs (Scagel and Andersen, 1997; Phillips et al., 2002).

Another energetically costly response to O₃ exposure is that the production of defense compounds, such as antioxidants, tend to increase under elevated O₃ conditions (Sheng et al., 1997; Tausz et al., 1999b; 2002). Antioxidants help the plant scavenge free radicals before they can cause damage to membranes or cell walls, but they demand C for production such that growth can be adversely affected. In mature Jeffrey pine, stomatal uptake of O₃ elicited one complex of antioxidant defenses in mesic microsites, while endogenously generated free radicals in the chloroplast elicited a second complex of antioxidant defenses in xeric microsites (Grulke et al., 2003b).

9.7.6.2 Material Flow

Plants as producers are responsible for using inorganic atmospheric C and reducing it into organic forms used by consumers, thus driving nutrient processes in ecosystems. Ozone has the potential to disrupt material flow through organic C cycling and changes in nutrient cycling, particularly N and P cycling. Although there is indirect evidence that O₃ is disrupting C and nutrient cycling at the ecosystem level, there is little direct evidence that O₃ alters nutrient processing at ecosystem scales.

The greatest annual nutrient and C input to ecosystems is from foliar and root turnover. Excision of plant parts and whole plant mortality are potentially much larger, but syncopated, ecosystem inputs. Ozone exposure alters C cycling in the ecosystem by affecting the within-plant C allocation and partitioning of dominant, O₃-sensitive plants, and through chemical composition and rate of decomposition of sloughed plant parts (roots, branches, leaves) (Figure 9-19).

In addition to O₃-induced changes in the quantity of C and nutrient inputs into ecosystems, O₃ also can alter the nutrient quality of inputs. Ozone exposure alters nutrient levels in the foliage (Boerner and Rebbeck, 1995; Lindroth et al., 2001; Fenn and Poth, 1998; Momen et al., 2002) and affects the C:N ratio (Andersen et al., 2001; Lindroth et al., 2001; Grulke and Lee, 1997; Grulke et al., 2003b). Concentrations of compounds such as tannins, lignin and phenolics (Kim et al., 1998; Findlay et al., 1996; Baumgarten et al., 2000; Saleem et al., 2001) are also affected by O₃ exposure, which in turn alters decomposability (Fenn and Dunn, 1989) and litter buildup in the ecosystem.

1 There are several possible pathways by which O₃ may affect litter quality and, hence, litter
2 decomposition, thus altering nutrient flow in ecosystems. These include altered C quality,
3 altered nutrient quality, and alteration of leaf surface organisms important in decomposition
4 pathways. For example, yellow poplar (*Liriodendron tulipifera*) and black cherry (*Prunus*
5 *serotina*) litter exposed to O₃ showed greater N loss during decomposition than charcoal filtered
6 controls, although mass loss did not vary among O₃ treatments (Boerner and Rebbeck, 1995).
7 Subsequent studies showed that although foliar N was not affected by O₃ exposure in yellow
8 poplar leaves, foliage decomposed more slowly (Scherzer et al., 1998). Other studies have also
9 shown a change in foliar N concentration in response to O₃ treatment, affecting the C/N ratio and
10 possible litter quality (Andersen et al., 2001).

11 In some cases, it appears that N remobilization from foliage into the plant is not complete
12 at the time of foliage abscission in O₃-exposed plants (Findlay and Jones, 1990; Stow et al.,
13 1992; Matyssek et al., 1993; Patterson and Rundel, 1995). Greater N content of senesced litter
14 could increase rates of decomposition. When O₃-exposed cottonwood (*Populus deltoides*) leaves
15 abscised at the same time as control leaves, they decomposed at similar rates; however,
16 prematurely senesced foliage from O₃-exposed cottonwood decomposed more slowly than
17 controls despite their higher N content (Findlay and Jones, 1990; Findlay et al., 1991). Higher N
18 in senesced leaves appeared to be related to organic complexes formed by bound phenolics in
19 O₃-exposed leaves, that made the litter less palatable to decomposers, thereby slowing
20 decomposition rates (Jones et al., 1994; Findlay et al., 1996). Increased phenolics also have
21 been found in European silver birch (*Betula pendula*) exposed to O₃ (Saleem et al., 2001).

22 Carbon quality in leaf litter also changes in O₃-exposed foliage. Compositional changes in
23 leaf structural characteristics, such as lignin content, would be expected to alter rates of litter
24 decomposition (Fogel and Cromak, 1977; Meentemeyer, 1978; Kim et al., 1998). Blackberry
25 (*Rubus cuneifolus*) litter exposed to elevated O₃ had greater permanganate lignin than control
26 treatments, which was inversely related to mass-loss rates in decomposition studies (Kim et al.,
27 1998).

28 Ozone may affect early stages of decomposition by altering populations of leaf surface
29 organisms before or after senescence. Magan et al. (1995) found a shift in phyllosphere fungi on
30 Scots pine (*Pinus sylvestris*), Sitka spruce (*Picea sitchensis*), and Norway spruce (*Picea abies*)
31 exposed to O₃, but the potential effect of these changes on subsequent litter decomposition was

1 uncertain. The slowest decomposition rates of pre-exposed blackberry leaves were found when
2 senesced foliage was exposed to O₃ during decomposition, suggesting a possible direct effect of
3 O₃ on microorganisms in decomposing litter (Kim et al., 1998). Whether O₃ concentrations at
4 the soil surface influence initial stages of litter decomposition remains to be addressed.

5 Ozone exposure also reduces nutritional content of foliage because of the degradation of
6 chlorophyll. Reconstruction of chlorophyll may be limited by nutritionally poor soils or low soil
7 moisture, as well as alteration of root uptake by O₃ exposure and other stressors. Foliar exposure
8 to O₃ may also increase leaching of nutrients (Kerstiens and Lendzian, 1989). Ozone exposure
9 promotes early senescence of foliage (Miller and Elderman, 1977; Heath and Taylor, 1997), with
10 higher nutrient content than if excised later in the growing season (Poth and Fenn, 1998).

11 Since O₃ can slow decomposition through changes in leaf quality and quantity, leaf litter
12 can accumulate (Fenn and Dunn, 1989). The accumulation of soil organic matter from increased
13 leaf litter, even in the absence of acidic deposition, can lower soil pH (Binkley, 1992). Lower
14 soil pH can promote loss of nutrients, particularly cations, from the system, further reducing
15 nutrient availability to the plant. Complex organic compounds in decomposing litter may also
16 tie up nutrients, rendering them less available to plants.

17 Other factors such as nutrient deposition also affect the degree to which O₃ effects C and
18 nutrient flow through ecosystems. In high-pollution sites, the effect of N deposition is difficult
19 to separate from O₃ exposure, and reviews of the effect of acidic (and N) deposition on
20 ecosystem nutrient dynamics are important to consider (see Binkley, 1992; Fenn et al., 1998;
21 2003). For example, at moderately high pollution sites, foliar content of N is higher than that at
22 lower pollution site, but so are P, Mg, and Fe contents (Poth and Fenn, 1998). Although
23 significant changes in foliar tissue chemistry have occurred in response to long-term pollutant
24 deposition in the Carpathians (Mankovska et al., 2002; Fenn et al., 2002), much of this response
25 is correlated to heavy metal and N and S deposition. At this point, the contribution of O₃
26 exposure alone cannot be isolated without careful between-site comparisons of plant response
27 and understanding the depositional velocities of constituent atmospheric species. The significant
28 effects of plant response along known pollution gradients are important to consider because
29 heavy metal and N and S deposition has significantly declined over the last decade, while O₃
30 exposures remain high and declines in forest health have been sustained.

1 Models provide a means to track material flows through ecosystems. Two biogeochemical
2 models were parameterized to capture long-term effects of O₃ exposure, N deposition, and
3 climate on a *ponderosa pine*-dominated site in the eastern San Bernardino Mountains.
4 Simulated O₃ exposure resulted in faster production and turnover of foliage and a shift in carbon
5 from the canopy (15% reduction) to the forest floor (increase of 50 to 60%) (Arbaugh et al.,
6 1999). When O₃ exposure was combined with that of N deposition, litter mass exponentially
7 increased.

8 The direct effect of O₃ exposure on below-ground nutrient dynamics and ecosystem
9 material flow is poorly understood. Additional research will be necessary to understand spatial
10 and temporal dynamics of nutrient and C flow in ecosystems and to separate the effects of O₃
11 from those attributable to N deposition.

12 13 **9.7.7 Hydrological and Geomorphological**

14 At present, there are no publications on the effects of O₃ exposure that are carried through
15 at the ecosystem level to changes in mass water flow, channel morphology, riparian habitat
16 complexity, or sediment movement. It is possible that processes occurring at smaller scales are
17 affecting geomorphological processes in ecosystems; however, difficulty in scaling these
18 responses spatially and temporally have made it difficult to show experimentally. It is possible
19 that O₃ exposure affects water quality through changes in energy and material flows, as
20 discussed previously.

21 22 **9.7.8 Natural Disturbance Regimes**

23 There has been little research on how natural disturbances interact with O₃ to affect
24 performance of plants, communities, and ecosystems. The frequency, intensity, extent, and
25 duration of natural disturbances are variable and unpredictable. However, there have been
26 enough ecophysiological studies to suggest that O₃ could predispose plant communities to
27 certain natural stresses, e.g., drought, stress, or extreme low temperature stress during the winter.

28 While several studies have shown that drought stress reduces O₃ uptake through stomatal
29 closure, evidence also suggests that O₃ can alter plant water use and susceptibility to drought.
30 In controlled studies, Reich and Lassoie (1984) showed that relatively low O₃ concentrations
31 could diminish stomatal control and alter water use efficiency. Ash trees (*Fraxinus excelsior*)

1 exposed to elevated O₃ had greater water use early in the growing season, but less water use late
2 in the growing season when exposed to elevated O₃ (Wiltshire et al., 1994). Under moderate
3 drought stress, *Picea abies* trees grown under elevated O₃ consumed water faster and showed
4 higher stomatal conductances than controls (Karlsson et al., 1995). Pearson and Mansfield
5 (1993) showed that successive O₃ episodes disrupted stomatal function, making *Fagus sylvatica*
6 seedlings more susceptible to drought. Previous year O₃ exposure was shown to have a
7 carryover effect in the following growing season for *Fagus sylvatica* (Pearson and Mansfield,
8 1994).

9 Few studies showing the effects of O₃ on water relations of field-grown trees are found in
10 the literature. However, Grulke et al. (2003a) examined the effects of O₃ on canopy transpiration
11 of *Pinus jeffreyi* from mesic and xeric microsites and found that trees from mesic sites had 20%
12 more O₃ uptake than those in the xeric sites. The authors also concluded that the mesic trees had
13 greater O₃ injury as evidenced by lower needle retention, whereas trees in xeric microsites had
14 greater chlorotic mottle. Chlorotic mottle induced by stomatal uptake of O₃ is indistinguishable
15 from that of endogenously produced oxidants resulting from partially closed stomata, a reduction
16 of CO₂ inside the leaf, and production of strong oxidizers within the chloroplast when excited
17 electrons are passed to O₂ instead of CO₂ under high light levels.

18 Trees living near the limits of their freezing-tolerance range may be especially susceptible
19 to predisposition of freezing injury by O₃ (Sheppard et al., 1989). However, *Pinus halepensis*
20 exposed to elevated O₃ had enhanced winter hardiness (Wellburn and Wellburn, 1994). As with
21 the seasonal carryover of drought susceptibility, the influence of elevated O₃ on freezing
22 tolerance is carried over from summer to winter. Such effects have been demonstrated for *Picea*
23 *sitchensis* (Lucas et al., 1988) and for *Picea rubens* (Waite et al., 1994). Sorting out the role of
24 elevated O₃ in contributing to frost or low-temperature damage in forests remains difficult due to
25 the presence of other factors that may affect senescence.

27 **9.7.9 Scaling to Ecosystem Levels**

28 The vast majority of literature describing O₃ effects comes from short-duration herbaceous
29 plant or tree seedling studies under controlled conditions. Scaling results from these studies
30 requires extrapolation over both space and time in order to understand the full extent of changes
31 in ecosystems. In addition to spatial, temporal, and age-related complexities, ecosystems are

composed of organisms whose lifetimes range from hours to centuries (Laurence and Andersen, 2003). Forested ecosystems are affected by environmental conditions such as water and nutrient availability, as well as by intra- and interspecies competition. Therefore, direct experimentation to determine the response of forested ecosystems is not simply a matter of determining the effect of O₃ on individual mature trees. In addition, even if an experiment can be conducted, extrapolation of the results across landscapes and regions remains challenging. Nonetheless, models provide a means to explore possible long-term changes and to identify important research uncertainties.

Approaches to scaling fall roughly into two categories: (1) process-based modeling to extrapolate physiological responses to O₃ based on seedling studies and (2) field assessments using surveys and growth correlations, often in association with stand-level models to address ecosystem complexity. Comparatively good information is available on process level effects of O₃ in seedlings and, therefore, some models offer the opportunity to use this information to scale O₃ effects at the stand and regional scales (Hogsett et al., 1997; Fuhrer et al., 1997; Chappelka and Samuelson, 1998; Laurence et al., 2000).

9.7.9.1 Scaling from Seedlings to Mature Trees

A number of investigators have used simulation models based on physiological processes to integrate available data and predict the effects of O₃ on mature trees. Such models predict tree growth by simulating fundamental mechanisms, rather than through a statistical analysis of empirical data. For instance, the process of photosynthesis is simulated based on environmental conditions and physiological characteristics, and then the fixed C is allocated to plant growth using principles of plant physiology. Models based on mechanisms should be applicable across wide areas if the important functional relationships are represented accurately in the models and if the environmental conditions are accurately identified. The ability of six models (TREGRO, CARBON, ECOPHYS, PGSM, TREE-BGC, and W91) to simulate the effects of climatic change and O₃ have been reviewed by Constable and Friend (2000). Of these models, only PGSM and TREGRO explicitly simulated the effects of O₃ on foliar processes.

The TREGRO model was used to simulate C allocation and tissue growth in seedlings and mature red oak trees based on the experimental data discussed above (Weinstein et al., 1998). For seedlings at 2×-ambient O₃, only the total nonstructural carbohydrate (TNC) storage pool

1 was predicted to be affected. For mature trees, large decreases were predicted for TNC, leaves,
2 stem, branch, and both fine and coarse roots. Most predicted effects in mature trees were
3 consistent with observations in the field, but the simulations overestimated the effect of
4 2×-ambient O₃ on root TNC and growth. The authors suggested that this discrepancy may
5 have been due to trees reducing respiration in response to O₃ stress, a response not simulated in
6 the model.

7 For *Abies concolor*, TREGRO was parameterized and simulated growth of a mature tree
8 for 3 years to test for effects of O₃ exposure and drought stress (Retzlaff et al., 2000).
9 Reductions in O₃ exposure-mediated carbon assimilation were translated to losses in whole tree
10 biomass that probably would not be detectable in the field. However, TNC levels in branch
11 tissue were simulated to be lowered by over 50%, and branch growth was reduced in a
12 moderately polluted site relative to a clean site. Low O₃ exposure (sufficient to decrease C
13 assimilation by 2.5%) and drought stress (25% reduction in annual precipitation, which is
14 common on a decadal scale) acted synergistically to reduce C gain of whole tree biomass of
15 *A. concolor*. Simulated results of the tests were comparable to effects found in OTCs for
16 seedlings and pole-sized trees in clean and moderately polluted sites.

17 Models such as TREGRO are usually parameterized from many different sources of data,
18 including chamber experiments and plantations, from seedlings to mature trees, making it
19 difficult to validate that they reproduce changes that occur as trees develop from seedlings to
20 maturity. To address this issue, physiological and growth data were collected from a natural
21 stand of *P. ponderosa* and used to parameterize the TREGRO model (Grulke and Retzlaff,
22 2001). Representative trees of each of five tree age classes were selected based on population
23 means of morphological, physiological, and nearest neighbor attributes. Seedlings were
24 observed to differ significantly from pole-sized and older trees in most physiological traits. The
25 changes in biomass with tree age predicted from the model closely matched those of trees in the
26 natural stand.

27 The PGSM model was used to simulate *Pinus ponderosa* seedling growth responses to O₃
28 exposure and drought stress (Chen et al., 1994). Drought stress was predicted to reduce the
29 effect of O₃ on growth, as was observed in the experimental data. The TREGRO model was
30 used to simulate responses of Pacific Coast and interior varieties of *P. ponderosa* to five
31 simulated O₃ exposures between subambient and 3×-ambient (Constable and Taylor, 1997).

1 Simulated growth of var. *ponderosa* was reduced more than that in var. *scopulorum* with all O₃
2 exposures. Drought was protective of O₃ exposure. Similar results were also found in a
3 relatively moist *ponderosa* pine plantation (Panek and Goldstein, 2001), whereas drought was
4 synergistically deleterious with cumulative O₃ exposure in a natural stand (Grulke et al., 2002b).

5 For *Pinus taeda*, the Plant Growth Stress Model (PGSM) was calibrated with seedling data
6 and then used to simulate the growth of mature trees over a 55 year period in the Duke Forest,
7 NC, using estimates of historical O₃ concentrations (Chen et al., 1998). Simulated stem diameter
8 and tree height were comparable to observed values. In another simulation using TREGRO,
9 loblolly pine was more sensitive (greatest reduction in C gain) to a peak O₃ episode in July
10 (Constable and Retzlaff 1997), whereas mature tulip poplar (*Liriodendron tulipifera*) was more
11 sensitive to a peak O₃ episode in August.

12 For *Populus tremuloides*, the ECOPHYS model was used to simulate the relative above-
13 ground growth response of an O₃-sensitive clone (259) exposed to square-wave variation in O₃
14 concentration (Martin et al., 2001). The model adequately simulated several effects of O₃,
15 including a greater effect on stem diameter than on stem height, earlier leaf abscission, and
16 reduced stem and leaf dry matter production at the end of the growing season. For *Acer*
17 *saccharum*, the TREGRO model was used to predict effects of a 10-year O₃ exposure on root and
18 stem growth of a simulated 160-year-old tree (Retzlaff et al., 1996). Twice-ambient O₃ exposure
19 (for Ithaca, NY) was predicted to deplete the TNC pools and reduce fine root production.

21 **9.7.9.2 Surveys, Growth Correlations and Stand-Level Modeling**

22 Stand-level studies have included surveys of O₃ symptoms, correlations of radial growth
23 with O₃ and other environmental factors, and regional scale modeling. In addition, open air O₃
24 exposure systems, such as those being used on *Populus tremuloides*-mixed stands in northern
25 Wisconsin (Karnosky et al., 2003a) and on *Fagus sylvatica* and *Picea abies* in Germany (Nunn
26 et al., 2002) offer an opportunity to examine larger plot sizes, older trees, and trees growing
27 under realistic competition. Plots along natural O₃ gradients, as have been used very effectively
28 in southern California forest studies (Miller and McBride, 1999) and with *P. tremuloides* stands
29 in the Great Lakes region (Karnosky et al., 1999), offer additional insights into ecosystem level
30 responses. Undoubtedly, however, simulation modeling will have to become an integral
31 component of research in order to predict adequately ecosystem responses to O₃ (Laurence and

Andersen, 2003). Results of these approaches are discussed below, organized into three U.S. regions: (1) northern states (including the upper Midwest and the Northeast), (2) southeastern states, and (3) western states (primarily California). A fourth section contains selected information from Europe.

Northern and Midwestern United States. In recent years, the USDA Forest Service has conducted systematic O₃ biomonitoring surveys in most north-central and northeastern states (Smith et al., 2003; Coulston et al., 2003). Plots are located on a systematic grid, and trained field crews evaluate up to 30 plants of up to six species that have foliar injury symptoms diagnostic of O₃ damage. For the United States as a whole, injury has been found more often in eastern than in interior or west-coast states. As expected, O₃ injury is more common and more severe in areas with higher O₃ concentrations. Of sampled *Prunus serotina* plots, ~12% were estimated to be at high risk for injury based on a injury index derived from the survey data (Coulston et al., 2003). *P. serotina* was estimated to be at risk for injury on the Allegheny Plateau and the Allegheny Mountains (in Pennsylvania, West Virginia, and Maryland), as well as in the coastal plain of Maryland and Virginia.

Ozone concentration, foliar injury, and physiological traits were measured on *P. serotina* trees of different sizes in Pennsylvania (Fredericksen et al., 1995). The proportion of foliage injured was 46% for seedlings, 15% for saplings, and 20% for canopy trees. Cumulative O₃ flux was the most useful O₃ metric for predicting injury. Injury was negatively correlated ($r^2 = 0.82$) with net photosynthetic rates, but was not related to stomatal conductance. *P. serotina* is discussed further below, because detailed surveys have been conducted in the Shenandoah and Great Smoky Mountains National Parks in the southeastern United States.

Over the past several decades, some surveys of *Pinus strobus* have reported significant associations between foliar injury and reduced growth (Anderson et al., 1988; Benoit et al., 1982). However, a review of 93 surveys conducted from 1900 through the late 1980s concluded that methodological problems were pervasive, including such issues as proximity to roads, lack of peer review, lack of random sampling, small sample sizes, and lack of quantitative methods to estimate severity (Bennett et al., 1994). Because of these problems, along with evidence of adequate growth rates for *P. strobus* regionally and contradictory evidence from numerous studies of symptom production in response to controlled O₃ exposure, these authors concluded

1 that there was no clear evidence of decline in *P. strobus* (Bennett et al., 1994). A more recent
2 study in Acadia National Park (ANP) in Maine found no association between O₃ exposure in
3 OTCs and symptom development in *P. strobus*, calling into question whether symptoms
4 previously ascribed to O₃ may be caused by some other stress (Kohut et al., 2000). However,
5 another ANP study found significant correlations between O₃ exposure and the radial growth of
6 trees during 10 years in 7 of 8 stands examined (102 trees total; Bartholomay et al., 1997).
7 Taken together, these results suggest that there may not an association between growth of *P.*
8 *strobus* trees and putative O₃ symptoms, but there may be an association between O₃ exposure
9 and radial growth of mature trees in the field.

10 The study of O₃ effects was undertaken from 1990 to 1993 in the ANP in Maine, because
11 this location experiences elevated O₃ exposures due to transport from urban areas located upwind
12 (Kohut et al., 2000). Thirty-two species of plants found in the park were propagated and
13 exposed to O₃ in OTCs. In addition, ambient O₃ concentrations were monitored at the study site
14 at 15 m above sea level and near the top of Cadillac Mountain at 470 m above sea level. At the
15 study site, the maximum 1-h O₃ concentration was 140 ppb, which occurred in both 1990 and
16 1991. Daytime 12-h O₃ concentrations were 35, 41, 36, and 37 ppb during the four years; and O₃
17 concentrations were consistently higher at the high-elevation site. Species showing foliar injury
18 at ambient O₃ concentrations included *Prunus serotina*, *Populus tremuloides*, *Fraxinus*
19 *americana*, *Pinus banksiana*, big-leaf aster, and spreading dogbane. Species showing foliar
20 injury at 1.5 × ambient O₃ concentrations included *Betula populifolia*, small sundrops, and
21 bunchberry. Species remaining uninjured at 2 × ambient O₃ concentrations included *Betula*
22 *papyrifera*, *Pinus strobus*, *Pinus rigida*, *Picea rubens*, *Thuja occidentalis*, *Quercus robur*,
23 Canada bluejoint grass, wild radish, and Canada mayflower. Because of their O₃ sensitivity and
24 diagnostic symptoms, big-leaf aster, spreading dogbane, *Populus tremuloides*, *Fraxinus*
25 *americana*, and *Prunus serotina* were recommended as bioindicators for the ANP.

26 The PnET-II model was applied to 64 locations across the northeastern United States to
27 simulate the effects of ambient O₃ on mature hardwood forests (Ollinger et al., 1997). In this
28 model, O₃ effects on each of several layers of the forest canopy were represented by a single
29 linear equation relating predicted O₃ uptake to decreased net photosynthetic rate. Wood growth
30 was predicted to decrease between 3 to 22%, with greatest reductions in southern portions of the
31 region where O₃ levels were highest and on soils with high water-holding capacity where

1 drought stress was absent. Little variation was predicted among years, because high O₃ often
2 coincided with hot, dry weather conditions that reduced predicted stomatal conductance and
3 O₃ uptake.

4 In order to estimate the impact of O₃ on forests, effects must be evaluated not only on
5 individuals, but also on mixtures of species and the composition of forest stands. The PnET
6 model described above evaluated the effects of O₃ on broad forest types (an evergreen/deciduous
7 mix), but did not address specific forest species composition. In order to address competition
8 among species, the TREGRO model was linked to the ZELIG forest stand growth model and a
9 geographic information system was created to predict the effects of O₃ across the north-central
10 and northeastern United States (Laurence et al., 2000). ZELIG is a gap-succession model used
11 to simulate succession in mixed stands typical of eastern and northern forests. Ambient O₃
12 generally caused a reduction of 2 to 4% in the growth of *Quercus robur* across the region during
13 the 100-year simulation. The response followed the pattern of O₃ exposure, with little effect in
14 the northwest part of the region, but with greater effect in southern locations. The O₃ response of
15 *Acer saccharum* to O₃ varied widely, but the overall growth response was always positive,
16 indicating that the evergreen/deciduous mix was able to take advantage of the decrease in the
17 growth of *Q. robur* and other species caused by O₃. In the northernmost part of the region,
18 *A. saccharum* growth increased by up to 3%, but in the southern part of the region, its growth
19 increased by up to 12%. The authors ascribed this enhanced growth to a combination of warmer
20 temperatures and reduction in the growth of *Prunus serotina*, a minor component of the
21 simulated stand that was very sensitive to O₃.

22
23 ***Southeast United States.*** In a survey of the Great Smoky Mountains National Park, foliar
24 injury attributed to O₃ was found on 47% of the more than 1,600 plants examined (Chappelka
25 et al., 1997). In subsequent surveys of injury in the park, injury was found on mature trees of the
26 following species: *Sassafras albidum*, *Prunus serotina*, and *Liriodendron tulipifera* (Chappelka
27 et al., 1999a,b). In a similar study in Shenandoah National Park, injury was found on *Fraxinus*
28 *americana*, *P. serotina*, and *L. tulipifera* (Hildebrand et al., 1996).

29 For *Prunus serotina* seedlings grown in soil in OTCs and exposed to relatively low
30 ambient levels of O₃ in Pennsylvania, there was no correspondence between visible foliar stipple,
31 leaf gas exchange, and seedling growth between two families previously shown to differ in O₃

symptoms (Kouterick et al., 2000). However, significant exposure-response relationships were found for foliar injury in the Great Smoky Mountains and Shenandoah National Parks. In each park, foliar injury was evaluated on mature *P. serotina* trees on three plots at different elevations near O₃ monitors during 1991 to 1993 (Hildebrand et al., 1996; Chappelka et al., 1999a,b). In 1991, incidence was 60% and 45% for the two parks and 33% in both parks during 1992 and 1993. Symptoms were greater at the highest elevations where O₃ concentrations were highest. In another study, radial growth rates were measured in 44 *P. serotina* trees ranging in age from 19 to 56 years old with and without O₃ symptoms, at three sites in the Great Smoky Mountains National Park. Trees with O₃ symptoms were compared to similar-sized trees with few symptoms. There was no evidence that trees with O₃ symptoms had lower growth rates ($p = 0.6$) (Somers et al., 1998).

In the Great Smoky Mountains National Park, radial growth rates were measured for 44 *L. tulipifera* trees ranging in age from 30 to 58 years old, with and without O₃ symptoms, at three sites at different elevations (Somers et al., 1998). Trees with O₃ symptoms averaged 30% lower growth rates over ten years ($p = 0.0005$). Seedlings of *Liriodendron tulipifera* were exposed for two seasons to 2×-ambient O₃ exposures in OTCs in Delaware, OH (seasonal SUM00 exposures of 107 and 197 ppmh) (Rebbeck, 1996). Foliar O₃ symptoms were observed, but growth was not reduced.

In order to evaluate the influence of interspecies competition on O₃ effects, the linked TREGRO and ZELIG modeling system was used to predict the effects of O₃ over 100 years on the basal area of species in a *Liriodendron tulipifera*-dominated forest in the Great Smoky Mountains National Park (Weinstein et al., 2001). Ambient O₃ was predicted to reduce the basal area of *L. tulipifera* by 10%, whereas a 1.5×-ambient exposure was predicted to cause a 30% reduction. Basal area of *Acer rubrum* and *Prunus serotina* was predicted to increase for some years, but then decrease by up to 30%, with few changes in the total basal area of all species by the end of the simulation.

In order to evaluate the influence of interspecies competition on O₃ effects, the linked TREGRO and ZELIG modeling system was used to predict the effects of O₃ on the basal area of *Pinus taeda* and *Liriodendron tulipifera* growth throughout their ranges (Laurence et al., 2003). The models were parameterized using biological and meteorological data from three sites in the southeastern United States (in Alabama, Louisiana, and North Carolina). Forest stand response

1 to five O₃-exposure regimes with annual SUM06 values ranging from 0 to 100 ppmh per year
2 was simulated for 100 years. The simulated basal area of the two species was generated within
3 the context of four other tree species common in southeastern forests. Basal area of *P. taeda* was
4 highly responsive to precipitation and O₃ exposure, with the greatest increases under high-
5 precipitation, low O₃-exposure scenarios and the greatest decreases under low-precipitation, high
6 O₃-exposure scenarios. The basal area of *L. tulipifera* did not significantly differ (+10%) from
7 simulations using a “base case” (ambient O₃, average precipitation).

8 Systematic biomonitoring surveys found that approximately 24% of sampled sweetgum
9 and 15% of sampled *Pinus taeda* plots were estimated to be at high risk for foliar injury on the
10 coastal plain of Maryland and Virginia (Smith et al., 2003; Coulston et al., 2003). In a study in
11 Tennessee, the effect of ambient (uncontrolled) O₃ on 28 mature canopy-dominant 50 to
12 90-year-old *P. taeda* trees in five stands was measured over a 5-year period (McLaughlin and
13 Downing, 1995, 1996). Of many O₃ metrics, a 3-day average of hourly O₃ values ≥ 40 ppb
14 (AOT40) was found to best explain short-term variation in stem expansion as measured with
15 dendrometer bands. Interactions between O₃, temperature, and drought stress (as indicated by
16 the weekly moisture stress index) accounted for 63% of the short-term variation in stem growth
17 rates. Because there are interactions among O₃, drought stress, and temperature that may differ
18 with the averaging time (days to years), this type of study cannot provide conclusive proof of
19 cause and effect (Reams et al., 1995). However, the results do suggest that the effects of O₃
20 measured on loblolly seedlings may also be occurring in mature trees in both wet and dry sites.
21 The magnitude of effects of O₃ on growth, including interactions with other variables in this
22 study, ranged from 0 to 15% over 5 years, with an average of 5.5%.

23
24 **Western United States.** The USDA Forest Service conducted O₃ biomonitoring surveys in
25 Washington, Oregon, and California during one year (1998), and in the Sierra Nevada and
26 Sequoia National Forest every other year for several decades (Campbell et al., 2000). Overall,
27 only one plot showed any symptoms of O₃ injury outside of the Sierra Nevada and Sequoia
28 National Forests. In the Sierra Nevada National Forest, between 30 and 40% of trees showed
29 injury from 1989 through 1997. In the Sequoia National Forest, between 40 and 50% of the
30 trees surveyed showed injury from 1990 through 1998.

1 For *Pinus ponderosa* along a well-studied gradient of O₃ exposure in the San Bernardino
2 mountains, chlorotic mottle was highest on foliage at the most polluted site, as has been found
3 previously (Grulke and Balduman, 1999). Based on whole-tree harvests, root biomass was
4 lowest at the most polluted sites, confirming previous studies with seedlings under controlled
5 conditions, as discussed above. Ozone responses in highly polluted environments such as
6 Southern California may not be predicted adequately by extrapolating effects from single-factor
7 experiments. Instead, combined approaches utilizing field experiments and modeling efforts
8 may be required to properly account for a combination of stressors including O₃, N deposition,
9 and drought. Furthermore, the available studies underscore the lack of correlation between O₃
10 symptoms and mature tree effects. If a field survey fails to find a correlation between mature
11 tree growth and O₃, this result may be due to the dominant effect of another factor such as N
12 deposition and may not be evidence that O₃ does not reduce the growth of mature trees.

13 The TREGRO model was used to evaluate how projected future temperature and CO₂
14 concentrations might affect the response of individual ponderosa pine to O₃ at seven sites in
15 California, Oregon, and Washington (Tingey et al., 2001). As expected, growth decreased with
16 increasing O₃ exposure. Differences in O₃ response among sites appeared to be due primarily to
17 differences in precipitation.

18 Often air quality standards do not translate directly into measureable improvements in tree
19 growth or productivity. To evaluate whether past improvements in air quality have improved
20 ponderosa pine growth, TREGRO was used to simulate growth at sites in the San Bernardino
21 Mountains in California (Tingey et al., 2004). Ozone and meteorological data from the past
22 37 years was used to run the simulations. Despite variation in precipitation and temperature,
23 O₃ was found to reduce simulated tree growth. The authors were able to simulate growth
24 improvements as air quality improved during the 1980s and 1990s, suggesting that
25 improvements in emission control strategies benefited ponderosa pine. The model simulations
26 were qualitatively consistent with improvements in canopy condition that were observed at sites
27 where O₃ reductions were the greatest.

28
29 ***Studies in Europe.*** In a 4-year study of *Fagus sylvatica* in Switzerland at 57 forest sites
30 ranging in age from 65 to 173 years, stem increment was found to decrease with increasing
31 maximum O₃ exposure (Braun et al., 1999). In this study, O₃ concentration was estimated by

interpolation among monitoring stations, and other site conditions such as soil water status and temperature were interpolated from weather stations. Other factors such as N deposition, tree diameter, and canopy dominance were also found to be significantly associated with stem increment. The maximum annual O₃ dose (expressed as AOT40) was found to be more strongly associated with decreased stem increment than was the average O₃ dose over the 4 years. A growth reduction of 22.5% (confidence interval 14.3 to 28.6%) was associated with each 10 ppmh increment of O₃ (expressed as AOT40). This decrease was steeper than the 6.1% growth reduction summarized previously from several OTC studies with *F. sylvatica* seedlings (Fuhrer et al., 1997). However, the authors suggest that this difference may be explained largely by the 4 years of exposure in their forest survey study as compared to the one-year exposures for seedlings. As with any forest survey, these results must be interpreted with caution because O₃ exposure was correlated with other variables, such as tree age and the deposition of NO₂ and SO₂.

9.7.10 Summary of Ecological Effects of Ozone Exposure on Natural Ecosystems

In this chapter, an effort has been made to discuss the adverse effects of O₃ on natural ecosystems within the context of the SAB framework for assessing and reporting ecological conditions (Young and Sanzone, 2002). Using this framework, there is evidence that tropospheric O₃ is an important stressor of natural ecosystems, with well-documented impacts on the biotic condition, ecological processes, and chemical/physical nature of natural ecosystems (Figure 9-20). In turn, the effects of O₃ on individual plants and processes are scaled up through the ecosystem affecting processes such as energy and material flow, intra and interspecies competition, and NPP. Thus, effects on individual keystone species and their associated microflora and fauna may cascade through the ecosystem to the landscape level. This suggests that by affecting water balance, cold hardiness, tolerance to wind, and by predisposing plants to insect and disease pests, O₃ may even influence the occurrence and impact of natural disturbance. Despite the probable occurrence of such effects, however, there are essentially no instances where highly integrated ecosystem-level studies have conclusively shown that O₃ is indeed altering ecosystem structure and/or function.




Figure 9-20. Common anthropogenic stressors and the essential ecological attributes they affect.

Source: Modified from Young and Sanzone (2002).

Systematic injury surveys demonstrate that foliar injury occurs on sensitive species in many regions of the United States. However, the frequent lack of correspondence between foliar symptoms and growth effects means that other methods must be used to estimate the regional effects of O₃ on tree growth rates. Investigations of the radial growth of mature trees in combination with data from many controlled studies with seedlings, as well as a few studies with

1 mature trees suggest that ambient O₃ is reducing the growth of mature trees in some locations.
2 Studies using models based on tree physiology and forest stand dynamics suggest that modest
3 effects of O₃ on growth may accumulate over time and may interact with other stresses. For
4 mixed-species stands, such models predict that overall stand growth rate is generally not likely to
5 be affected. However, competitive interactions among species may change as a result of growth
6 reductions of sensitive species. These results suggest that O₃ exposure over decades may be
7 altering the species composition of forests in some regions.

8 9 **RESEARCH NEEDS**

10 The knowledge base for examining the range of ecological effects of O₃ on natural
11 ecosystems is growing, but significant uncertainties remain regarding O₃ effects at the ecosystem
12 level. For example, there is a need for information on the following ecosystem-level responses:

- 13 • *Ecosystem Processes.* Little is known about the effects of O₃ on water, C, and nutrient cycling, particularly at the stand and community levels. Effects on belowground ecosystem processes in response to O₃ exposure alone and in combination with other stressors are critical to projections at the watershed and landscape scales. Little is yet known about the effects of O₃ on structural or functional components of soil food webs, or how these impacts could affect plant species diversity (Andersen, 2003).
- 14 • *Biodiversity and Genetic Diversity.* The study of genetic aspects of O₃ impacts on natural ecosystems has been largely correlational in nature and it remains to be shown conclusively whether O₃ affects biodiversity or genetic diversity (Pitelka, 1988; Winner et al., 1991; Davison and Barnes, 1998). Studies of competitive interactions under elevated O₃ levels are needed (Laurence and Andersen, 2003), and reexamination via new sampling of population studies to bring in a time component to previous studies showing spatial variability in population responses to O₃ are needed. These studies could be strengthened by modern molecular methods to quantify impacts on diversity.
- 15 • *Natural Ecosystem Interactions with the Atmosphere.* Little is known about feedbacks between O₃ and climate change on volatile organic compound (VOC) production, which in turn, could affect O₃ production (Fuentes et al., 2001). At moderate-to-high O₃ exposure sites, aberrations in stomatal behavior could significantly affect individual tree water balance of sensitive trees, and if the sensitive tree species is dominant, hydrologic balance at the watershed and landscape level could be affected. This has not been addressed in any model because O₃ exposure effects, if included in the modeling effort have assumed a linear relationship between assimilation and stomatal conductance.
- 16 • *Below-Ground Interactions.* While the negative effects of O₃ on below ground growth are well characterized, interactions of roots with the soil or microorganisms are not.

- 1 • *Other Interactive Effects.* Interaction studies with other components of global change (e.g., warming, increasing atmospheric CO₂, N deposition, etc.) or with various biotic stressors are needed to better predict complex interactions likely in the future (Laurence and Andersen, 2003). Whether O₃ will negate the positive effects of an elevated CO₂ environment on plant carbon and water balances is not yet known; nor is it known if these effects will scale up through the ecosystem. How might O₃ affect the progress of pest epidemics and insect outbreaks as concentrations increase is unclear (Skarby et al., 1998).
- 2 • *Reproduction Effects.* Information concerning the impact of O₃ on reproductive processes and reproductive development under realistic field or forest conditions are needed as well as examination of reproductive effects under interacting pollutants (Black et al., 2000).
- 3 • *Comparative Extrapolation.* The vast majority of O₃ studies of trees have been conducted with young, immature trees and in trees that have not yet formed a closed canopy. Questions remain as to the comparability of O₃ effects on juvenile and mature trees and on trees grown in the open versus those in a closed forest canopy in a competitive environment (Chappelka and Samuelson, 1998; Kolb and Matyssek, 2001; Samuelson and Kelly, 2001).
- 4 • *Scaling-Up Issues.* Scaling the effects of O₃ from the responses of single or a few plants to effects on communities and ecosystems is a complicated matter that will require a combination of manipulative experiments with model ecosystems, community and ecosystem studies along natural O₃ gradients, and extensive modeling efforts to project landscape-level, regional, national and international impacts of O₃. Linking these various studies via impacts on common research quantification across various scales using measures of such factors as leaf area index or spectral reflective data, which could eventually be remotely sensed (Kraft et al., 1996; Panek et al., 2003), would provide powerful new tools for ecologists.
- 5 • *Comparative Risk Assessment Methodologies.* Methodologies to determine the important values of services and benefits derived from natural ecosystems such that these could be used in comprehensive risk assessment for O₃ effects on natural ecosystems (Heck et al., 1998).

9.8 ECONOMIC EVALUATION OF OZONE EFFECTS ON AGRICULTURE, FORESTRY AND NATURAL ECOSYSTEMS

9.8.1 Introduction

The adverse consequences of ambient of air pollutant exposures on vegetation, ecosystems and components of the material environment have been documented since the beginning of the industrial revolution. Attempts to quantify the monetary damage and injury resulting from

1 tropospheric O₃ exposures to managed agriculture, forests, and natural ecosystems date back at
2 least to the 1950s.

3 Both methodological and data problems plagued early efforts to assess the monetary
4 damages of air pollution to crops and natural vegetation. Adams and Crocker (1989) discussed
5 the methodological issues, e.g., a lack of reliable data on effects from air pollutants on crop
6 yields or the failure to develop and apply appropriate economic models. Some of these problems
7 were remedied by the EPA's National Crop Loss Assessment Network (NCLAN) in the 1980s.
8 The EPA's NCLAN facilitated the performance of economic assessments by providing O₃-crop
9 yield data with which to estimate O₃-crop yield response functions (see Heagle [1988] for a
10 review of NCLAN procedures and findings). NCLAN also funded a series of economic
11 assessments that, along with subsequent economic assessments, documented substantial
12 economic damages to agriculture. (See Spash [1997] for a detailed review of economic
13 assessments, many of which used NCLAN data.)

14 Since the completion of the NCLAN program in the late 1980s, the number of economic
15 assessments of air pollution studies focusing on terrestrial ecosystems in general, and agriculture
16 in particular, has declined. For example, for the period of 1980 to 1990, 33 economic studies of
17 O₃ and other air pollutant effects on U.S. crops were published in peer-reviewed journal outlets
18 (Spash, 1997). However, in preparing this section of the current criteria document, only four
19 peer-reviewed economic assessments were found for the decade of 1991 to 2000 that addressed
20 vegetation in the United States. In addition, one peer-reviewed article (Kuik et al., 2000) was
21 found dealing with agriculture in the Netherlands. Recent interest in global climate change and
22 the potential effects of global warming on O₃ and other photochemical oxidants, has renewed
23 interest in the effects of air pollution on both managed and unmanaged terrestrial ecosystems
24 (Adams et al., 1998). In addition, concern is growing for regarding the effects of air pollutants
25 on natural ecosystems and on the services they provide (Daily, 1997). Unfortunately, this
26 interest has not yet translated into additional peer-reviewed publications addressing O₃ or other
27 air pollutants effects on ecosystems.

28 This section of the current criteria document first discusses the availability of economic
29 information and its usefulness in forming environmental policy. Next, economic assessments of
30 air pollution effects and findings from the 1996 AQCD (U.S. Environmental Protection Agency,
31 1996) are discussed, followed by a synthesis of the limited literature available since the 1996

AQCD with respect to O₃ effects on agriculture, forestry, and ecosystems. Finally, limitations and continuing uncertainties are reviewed. The most fundamental of these is the lack of measurements of the economic effects of air pollution on natural ecosystems. Other issues include the variability of performance in both managed and natural ecosystems under increased climatic and air pollution variability as well as the challenges related to spatial and temporal scales used in performing economic assessments. To date, this set of effects has been sparsely addressed.

9.8.2 The Measurement of Economic Information

Economic science is an exercise in deductive logic in which testable hypotheses about the behavior of economic agents (i.e., farmers, consumers, resource owners) and markets are deduced from a body of theory. That body of theory is based on a series of premises proposed by economists and philosophers dating back over two centuries to Adam Smith. These premises gradually evolved into a theoretical foundation primarily dealing with microeconomics, culminating in structural relationships that define the operation of markets. This foundation was first laid out in a comprehensive and rigorous fashion by Alfred Marshall in 1920. Paul Samuelson (1948) formalized these theoretical relationships, resulting in what is sometimes referred to as modern, or neoclassical, economics.

The insights gained from the theoretical foundation of economics helped shape the nature of applied economic, or policy-relevant research. An example of such applied research is when economists seek to measure the economic consequences of air pollution on agriculture. Such an application is described in Adams and Horst (2003), who provided a graphical representation of the measurement of the effects of air pollution on the well-being of producers and consumers. Economic theory is applied to real-world problems when the methods of economics and the need for data from other disciplines come into play. When measuring the economic effects of an environmental change, economists need an economic model or method that is theoretically consistent, i.e, defensible, as well as data with which to estimate both economic and environmental science relationships for use in the model. Among accepted economic assessment methods, the actual choice is frequently determined by the nature of the problem to be addressed. It should be noted that the choice of assessment method can affect the type of economic information that is obtained. Even with a given assessment method, results are sensitive to

specific data treatment or assumptions (Adams, 1999). For example, some methods only measure effects on a particular group e.g., farmers. Other methods may measure effects across several groups. Thus, one should not expect the magnitudes of damage or benefits to be identical across economic assessments. One should, however, expect that the direction of the effects will be similar.

Once it is established that an assessment meets basic economic criteria, e.g., including human behavioral responses, the selection of the specific economic assessment method is often a relatively minor issue in terms of estimating benefits of air pollution control (or disadvantages of increases in air pollution). Although results differ across approaches, the differences are largely attributable to specific features of the assessment (e.g., whether the natural science data include a particular effect or relationship, whether effects on consumers are measured, and so forth). The nature and quality of the air quality forecasts used in the assessments can greatly influence the sensitivity of the assessments (Adams and Crocker, 1989). This is particularly noticeable when dealing with forecasts of seasonal air pollution changes (Adams et al., 1988). From the standpoint of providing policy guidance, the differences in economic estimates attributable to the assessment methods are often swamped by uncertainty in the natural and physical science forecasts. This has also been noted in recent economic assessments of climate change (Adams et al., 1998). In many settings, the quality of economic assessments of air pollution is likely to be improved more by refining the physical and natural science data used in the assessments than by intensive efforts to fine-tune the assessment techniques (Adams, 1999).

9.8.3 Understanding of Air Pollutants Effects on the Economic Valuation of Agriculture and Other Vegetation in the 1996 Criteria Document

Evidence from the plant science literature cited in the 1996 O₃ AQCD (U.S. Environmental Protection Agency, 1996) is unambiguous with respect to the adverse effects of tropospheric O₃ on some types of vegetation. For example, the 1996 AQCD noted that findings from the EPA multiyear NCLAN program in the 1980s provided rigorous corroboration of at least two decades of previous research and a century of anecdotal observations that showed that O₃ at ambient levels caused physical damage to plants in general and to important crop species in particular. Specifically, NCLAN established that ambient O₃ levels resulted in statistically significant reductions in yields for some crop species (Heagle et al., 1988). The 1996 AQCD also assessed

1 the results of studies regarding O₃ effects on crops, forests, and natural vegetation in more detail.
2 More recent reviews, such as the comprehensive survey of the economic literature on
3 agricultural effects by Spash (1997) corroborate the synthesis of results reported in the 1996
4 AQCD.

5 The number and quality of assessments of the economic consequences of O₃ exposures on
6 vegetation reported in the 1996 AQCD are primarily a function of the state of evidence obtained
7 from scientific studies in each vegetation category. For example, the plant science evidence
8 reviewed in the 1996 AQCD concerning effects of O₃ exposures on agricultural crops was
9 reported to be more valid than for individual tree species or plant communities (ecosystems).
10 As a result, most economic assessments discussed in the 1996 AQCD focused on the data
11 obtained from studies of agricultural crops. The economic literature dealing with O₃ effects on
12 forest productivity in the 1996 AQCD is sparse. The few economic assessments of tree or forest
13 effects reported in the 1996 AQCD were confined to evaluations of assumed or hypothetical
14 changes in output, such as board feet of lumber (e.g., Haynes and Adams [1992]). As noted in
15 the 1996 AQCD, O₃ effects on ecosystems and their services had not been measured in any
16 systematic fashion and no peer-reviewed economic assessments were yet reported.

17 This section first briefly reviews economic assessments drawn from the review in the 1996
18 criteria document. This review is the benchmark against which recent articles are then discussed
19 in the subsequent section. As was the case in 1996, the discussion of economic valuation of
20 ecosystem effects is generally limited to conceptual and methodological issues, given the
21 continued lack of empirical analyses in this category.

23 **9.8.3.1 Agriculture**

24 In view of the importance of U.S. agriculture for both domestic and world consumption of
25 food and fiber, reductions in U.S. crop yields could adversely affect human welfare. The
26 plausibility of this premise has resulted in numerous attempts to assess, in monetary terms, the
27 losses from ambient O₃ exposures or the benefits of O₃ control, to agriculture. Twenty-three
28 assessments of the economic effects of O₃ exposures on agriculture were reviewed in the 1996
29 AQCD, highlighting key issues in the validity of these assessments (U.S. Environmental
30 Protection Agency, 1996). First, the evidence should reflect how crop yields will respond under
31 actual field conditions to O₃ exposures. Second, the air quality data used to frame current or

hypothetical effects of O₃ on crops should represent actual exposures sustained by crops at individual sites or production areas. Finally, the assessment methodology into which such data are entered should (1) capture the economic behavior of producers and consumers as they adjust to changes in crop yields and prices that may accompany changes in O₃ air quality; (2) accurately reflect institutional considerations, such as regulatory programs and income support policies (e.g., provisions of federal Farm Bill legislation), that may result in market distortions; and (3) use measures of well-being that are consistent with economic principles.

Assessments of O₃ damage to agricultural crops reported in the 1996 AQCD employed procedures for calculating economic losses that met the conditions described above. More specifically, the assessments provided 23 quantitative estimates of the economic consequences of exposures to O₃ and other air pollutants to agriculture that reflect producer-consumer decision-making processes, associated market adjustments, and some measure of distributional consequences between affected parties. Many of the economic assessments reviewed in previous O₃ documents also focused on O₃ effects in specific regions, primarily California and the Corn Belt (e.g., Garcia et al., 1986). This regional emphasis in the earlier literature may be attributed to the relative abundance of data on crop response and air quality for selected U.S. regions, as well as the importance of some agricultural regions (such as California) for the U.S. agricultural economy.

Two U.S. national studies described in previous O₃ criteria documents that are worthy of additional comment are Kopp et al. (1985) and Adams et al. (1986). These were judged to be “adequate” in terms of the three critical areas of data inputs in the 1996 AQCD. Together, it was argued, they provide a reasonably comprehensive estimate of the economic consequences of changes in ambient air O₃ levels on agriculture. Because of their central role in the 1996 criteria document, these two studies are reviewed briefly below.

The Kopp et al. (1985) and Adams et al. (1986) studies indicated that ambient levels of O₃ were imposing substantial economic costs of ~\$3.4 billion (in 2000 U.S. dollars) on agriculture. Both were judged to suffer from several sources of uncertainty, but the document concluded that these possible improvements in future assessments were not likely to greatly alter the range of agricultural benefit estimates arising from O₃ reductions for several reasons. First, the studies covered about 75 to 80 % of U.S. agricultural crops (by value). For inclusion of the other 20% to significantly change the estimates would require that their sensitivities to O₃ be much greater

1 than for the crops that have been included to date. Second, model-sensitivity analyses reported
2 in past studies indicate that changes in plant exposure response relationships must be substantial
3 to translate into major changes in economic benefits estimates. For example, it was assumed
4 unlikely that use of different exposure measures, or inclusion of interaction effects, would
5 greatly alter the magnitude of the economic estimates. Third, it was believed that countervailing
6 effects would mitigate against large swings in the estimates, e.g., longer O₃-exposure periods
7 may predict greater yield losses, but O₃-water stress interactions tend to reduce loss the
8 yield estimates.

9 Other national assessments reported in the 1996 AQCD provided general corroboration of
10 the results of Kopp et al. (1985) and Adams et al. (1986). An evaluation of these studies in
11 terms of the adequacy of information from plant exposure studies, and aerometric and economic
12 data was presented in the 1996 AQCD, along with estimates of benefits or damages associated
13 with changes in O₃. Most of the studies built on either Kopp et al. (1985) or Adams et al. (1986).
14 A relevant question was whether subsequent studies provided any “surprises” in terms of
15 magnitude of economic effects.

16 Common themes or findings from these and earlier O₃ and other air pollution studies were
17 summarized in two synthesis papers, Adams and Crocker (1989) and Segerson (1991). The
18 major conclusion is that the agricultural effects of tropospheric O₃ at ambient levels impose
19 economic costs to society or, conversely, that reductions in ambient O₃ should result in societal
20 benefits.

21 Several studies contained in the 1996 AQCD still are of interest. For example, one finding
22 pertains to the relationship between federal farm programs and air pollution regulations
23 (McGartland, 1987). In each case, the inclusion of farm programs in the economic models
24 resulted in modest reductions in the economic benefits of O₃ control due to increased farm
25 program costs. As Segerson (1987) noted, however, it is not clear that these increased costs
26 should be charged against the potential benefits of an O₃-regulatory standard, but rather,
27 considered as an additional cost associated with the inefficiencies of the farm program. It should
28 also be noted that the nature of federal farm programs was changed dramatically by Congress in
29 1996 in an attempt to reduce the federal government’s role in agriculture. Although more recent
30 federal legislation, such as the 2002 Farm Bill (U.S. Congress, 2002), appears to be restoring the
31 federal government’s role in the farming sector, this issue currently is not as important as

suggested by earlier studies, due to the declining reliance on deficiency payments to farmers, which tend to distort resource allocation.

Another national study (Adams et al., 1988) analyzed economic benefits under a regulatory alternative involving a seasonal (crop growing season) O₃-exposure index measured as a 12-h mean, instead of hourly levels or percent changes from ambient as reported in earlier studies. Specifically, a seasonal average of 50 ppb O₃ (measured as a 12-h seasonal mean) with a 95% compliance level, is reported in Adams et al. (1988). The result (a \$2.9 billion benefit in 2000 dollars) is similar to the assumed 25% reduction across all regions reported by Adams et al. (1986). At least one study also combined environmental stressors (e.g., O₃, UV-B radiation) in performing economic assessments. Adams and Rowe (1990), using the same model as Adams et al. (1986), reported that a 15% depletion of stratospheric O₃ (resulting in a 13% increase in tropospheric O₃) would cause an economic loss of ~\$1.4 billion in 2000 dollars attributed to the tropospheric O₃ increase. Reducing VOCs/NO_x motor vehicle emissions by 10% would result in a benefit of ~\$0.3 billion, while a complete elimination of motor vehicle emissions would yield a benefit of ~\$3.4 billion (1990 dollars). The range of these numbers is consistent with values reported in Adams et al. (1986), Kopp et al. (1985), and other national-level analyses, i.e., estimates of from \$1.0 to 2.0 billion for reductions in ambient O₃ of 25 to 40%.

9.8.3.2 Forests (Tree Species) and Natural Ecosystems

The long-term nature of air pollution effects on perennial species such as trees creates challenges to plant scientists in attempts to sort out the effects of specific individual stressors such as O₃ from among the many other potential causal factors (Skelly, 1988). It also creates problems in terms of measuring the impacts on direct economic value, such as reductions in board-feet of lumber produced per unit of time.

Most of the literature in the 1996 AQCD dealing with forest species reported the effects of O₃ exposures in terms of foliar injury (Davis and Skelly, 1992; Freer-Smith and Taylor, 1992; Simini et al., 1992; Taylor, Jr. and Hanson, 1992). This emphasis on foliar effects in the forest effects literature (rather than marketable yield) is similar to the state of science for agricultural crops prior to 1980. More recent studies address the effects of air pollutants on forest tree species diversity (Bringmark and Bringmark, 1995; Vacek et al., 1999; Weiner et al., 1997). However, such information is of limited use in economic assessments. The exception is in

1 measuring the economic value of aesthetic changes in a forest stock, where changes in species
2 composition may affect recreational values (Crocker, 1985).

3 The data concerning changes in marketed output such as board-feet of lumber or changes
4 in growth rates in managed forests, the affects on the growth of almond, peach, apricot, pear and
5 plum trees in orchards cited in the 1996 document (U.S. Environmental Protection Agency,
6 1996) have not been quantified. In addition, the economic impact of reductions in growth of
7 seedlings of evergreen trees, e.g., slash pine, presented in the same document have not been
8 valued. The few studies which attempted to measure economic losses arising from exposures to
9 O₃ or other pollutants circumvented the lack of plant science data by assuming (often arbitrary)
10 reductions in forest species growth or harvest rates (Adams, 1986; Callaway et al., 1986;
11 Crocker and Forster, 1986; Haynes and Adams, 1992). Although the economic estimates
12 reported in the 1996 AQCD are comparable to those reported for agriculture (e.g., \$2.6 billion
13 for eastern Canada forests, \$2.9 billion for eastern U.S. forests in 2000 dollars), the lack of yield
14 and/or growth effects data makes these studies only suggestive at best, of the economic
15 consequences of forest effects directly attributable to O₃ exposures questionable. Recent
16 developments in forestry economic modeling capabilities, in support of climate change research,
17 have enhanced the ability to measure the effects of environmental stressors on this sector
18 (Adams et al., 1996; McCarl et al., 1998). However, these models need data on changes in either
19 timber production or growth rates, both of which are lacking for forest species under alternative
20 O₃ levels.

22 **9.8.4 Studies Since 1996 of Ozone Exposure Effects on the Economic Value of** 23 **Agriculture, Forests, and Ecosystems**

24 Of the few current (post-1996) economic studies addressing agricultural effects, none offer
25 new insights of value in determining the economic cost of O₃ exposures. These post-1996
26 studies used variants of the economic methods from earlier assessments and measure yield
27 changes from response functions arising from the NCLAN or similar data. For example, Kim
28 et al. (1998) used a mathematical programming model of the San Joaquin Valley agricultural
29 sector in California, combined with crop yield response functions, to assess the economic effects
30 of O₃ on California crops. Their results showed net benefits from reductions in ambient O₃
31 levels, a finding consistent with all previous economic assessments. In another study,

1 Westenbarger and Frisvold (1994) measured agricultural exposures to O₃ (and acid precipitation)
2 in the United States. Though not an economic analyses of the costs of ambient exposures, they
3 identified areas of the United States of greatest potential economic damage based on the
4 interface between regional pollution levels and the value of crop production in each region.

5 A study by Murphy et al. (1999) of the economic effects of tropospheric O₃ on U.S.
6 agriculture is of note here, because it confirms the general magnitude of economic effects
7 reported by the two key studies performed a decade earlier (Kopp et al., 1985; Adams, 1986).
8 Specifically, Murphy et al. (1999) evaluated benefits to eight major crops associated with several
9 scenarios concerning the reduction or elimination of O₃ precursor emissions from motor vehicles
10 in the United States. Their analysis reported a \$2.8 to 5.8 billion (1990 dollars) benefit from
11 complete elimination of O₃ exposures from all sources, i.e., ambient O₃ reduced to a background
12 level assumed to be 0.025 to 0.027 ppm. While the analytical framework is similar to Adams
13 et al. (1986) in the use of NCLAN-based yield response functions and a mathematical
14 programming-based economic optimization model, the study is novel in its focus on the role of
15 motor vehicle emissions of VOCs/NO_x in total anthropogenic O₃ levels. The study is also
16 notable in its careful attention to federal farm program effects, particularly the deficiency
17 payment component.

18 In addition to these studies in peer-reviewed journals, a number of site-specific effects
19 studies have been performed, primarily by consulting companies for state public utility
20 commissions. Although perhaps of use to public utility commissioners concerned with effects
21 from single power plants or other localized sources, these regional studies generally contribute
22 little to the assessment of air pollution effects at the national level. Also, such reports are not
23 peer reviewed. Therefore, they are not discussed here.

24 There have been a number of recent studies of air pollutant effects on tree species in the
25 literature. Some have reported changes in total biomass and focused on European species
26 (Kurczynska et al., 1997). Other studies have assessed changes in composition of forest species
27 (biodiversity) or forest health due to exposure to air pollutants (Bringmark and Bringmark, 1995;
28 McLaughlin and Percy, 1999; Vacek et al., 1999). As noted previously, changes in forest
29 biomass and composition are more difficult to value than marketable products. However,
30 measures of forest composition or health have implications for an area of increasing policy
31 concern, that being the effect of air pollutants and other environmental stressors on unmanaged

(natural) ecosystems and the services they provide (Goulder and Kennedy, 1997; Pimentel et al., 1997). Considerable discussion has occurred among ecologists and economists as to the appropriate means for valuing these services (Anderson, 1990; Carpenter and Dixon, 1985; Common and Perrings, 1992). A number of conceptual articles have been published on this issue in both economic and ecological journals (Bergstrom, 1990; Castle, 1993; Pearce, 1993; Suter, II, 1990).

A continuing empirical challenge concerns the lack of information on how changes in biodiversity affect ecosystem performance resulting from O₃ stresses, and the problem of establishing economic values for such changes (Cairnes and Lackey, 1992; Norton, 1987; Pimm, 1997; Polasky, 2001; Randall, 1991). As noted in the 1996 AQCD, and more recently by Daily (1999, 2000) and Polasky (2001), there continues to be a lack of empirical studies that actually assess the economic value of changes in biodiversity or in service flows due to any environmental stressors. While some studies report monetary estimates, the estimates are generally for expository purposes and those would not be as defensible as the agricultural studies described earlier. For example, Costanza et al. (1997) assigned a value to the world's ecosystems, but the procedures used render this an exploratory study at best. As assessed by Polasky (2001), "In general, the field of valuation of ecosystem services is in its infancy." He attributed the lack of empirical studies due to both a lack of the understanding of ecology of ecosystem services and to the absence of reliable methods to estimate the value of these services.

In summary, the studies of crop and forest responses in the economic literature indicate that O₃ reduces crop yields and imposes economic costs. The economic literature also indicates that O₃ adversely influences the physiological performance of tree species and demonstrates, as expected, that changes in growth have economic consequences. However, the economic data and literature available on ecosystem effects of O₃ exposures are not sufficient to determine the economic costs.

9.8.5 Limitations of Scientific Studies and Economic Information

The 1996 O₃ AQCD discussed the need for additional research on both ecological functions and economic methodology in order to better understand the economic implications of air pollutants on ecosystem services. As noted by Daily (1999, 2000), Polasky (2001), and others, this research agenda continues to be important. Despite the large number of discussion

1 and survey articles published since 1996, there still are not sufficient data by which to estimate
2 confidently the magnitude of economic effects of O₃ to forests and natural ecosystems. Nor is it
3 apparent that ongoing research is adequate to answer this question in the next criteria document
4 cycle. Specifically, there does not appear to be any comprehensive, ecological studies underway
5 that attempt to measure changes in ecosystem outputs under alternative O₃ or other air pollutant
6 levels. Thus, in the near term, ecosystem services can only be discussed in qualitative terms.
7 However plausible the likelihood of economic damages to ecosystems, the available scientific
8 and economic information does not provide specific guidance on the magnitudes of these effects.

9 Beside the need to improve our understanding of the effects of O₃ exposures on natural
10 ecosystems, a number of areas of research could help assess the full economic consequences of
11 such pollutants on managed ecosystems. The first of these is the relationship between O₃
12 exposure levels and the variability of crop yields or changes in forest biodiversity. Most
13 assessments are based on the average or expected yield response of a crop to air pollution
14 exposure. However, the variability in yields (the spread or dispersion around the mean) appears
15 to be affected by the nature of plant exposure to pollutants (Hogsett et al., 1985; Musselman
16 et al., 1983). Plants exposed to the same mean dose but with different second moments of the
17 distribution of exposure may have different mean yields (Altshuller and Lefohn, 1996; Lefohn
18 and Benedict, 1982). In addition, the variability of the yield of the plant may also be increased
19 by greater variability in exposures (e.g., a higher frequency of extreme events). The economic
20 significance of higher yield varieties is such that variability may impose additional economic
21 costs, because most farmers have been reported to be averse to risk and prefer less variability for
22 a given yield (or profit). To date, no economic assessments of O₃ damages to agriculture or
23 vegetation in general include risk-averse behavior (studies cited here assumed risk neutrality).
24 To assess the economic consequences of a relationship between farmers' risk preferences and
25 O₃-induced changes in yield variability will require more information on the potential effects of
26 changes in O₃ on crop yield variability. While no economic studies were found on the effects of
27 O₃-induced changes in yield variability, a number of recent studies of climate change effects on
28 crop yields have indicated increased economic costs in the presence of increased climatic
29 variability (e.g., Mearns et al., 1997; Dixon and Segerson, 1999). Analogous economic costs
30 would be expected for changes in air pollution distributions; and these effects need to be
31 examined and quantified.

1 Another research area concerns the need improvement in our understanding of temporal
2 (dynamic) and spatial characteristics of O₃ exposures and their implications for crop yields,
3 production and producer profits. Most economic studies are static, in that they compare two
4 states of the world (e.g., economic activity at one O₃ level versus at an alternative level). In
5 addition, most national- level studies (the type needed to evaluate SNAAQS) display coarse
6 regional-level resolution in terms of crop response, O₃ exposure, and economic behavior. The
7 responses of producers to changes in yields due to changes in O₃ levels are generally assumed to
8 be similar over geographical areas up to several states in size. However, the changes between air
9 quality scenarios are more likely to be characterized by transient changes in exposure levels,
10 which means the producer responses are also likely to be gradual, rather than abrupt. Similarly,
11 the lack of finer scale (regional-level) data and modeling capabilities suggests that important
12 micro-level physical and economic effects are ignored. To what extent these abstractions
13 influence net economic effects is an empirical question. Research on these types of abstractions
14 and assumptions within other economic settings, such as climatic change, have shown that they
15 have implications for economic measurements (Adams, 2002).

16 Another issue is the natural or background level of O₃ (or other pollutants of interest)
17 assumed in economic studies. While many economic studies focus on changes in pollution
18 levels from current conditions, some studies have measured the economic damages between an
19 assumed, or pristine, level and current levels in agricultural areas. Such an analysis, it is
20 reasoned, will provide a measure of the net damages due to anthropogenic sources. The
21 challenge here is to have a correct measure of the background level of the pollutant. Recent
22 research by Lefohn et al. (2001) has suggested that background levels may be considerably
23 higher than assumed in some of the previous economic assessments reported in the 1996 AQCD
24 (25 to 30 ppb in most studies). For example, Lefohn et al. (2001) detected hourly readings of
25 from 40 to 80 ppb during winter and spring months in remote areas of the United States.
26 If background O₃ levels are in the range, then the economic damages estimated in studies with
27 lower background levels will be overstated. The issue of the background O₃ level is important to
28 all assessments of vegetative damages due to O₃.

29 In terms of expanding economic methods for future assessment, analysts should consider
30 using more “reduced form” estimation methods, particularly in situations where the availability
31 of dose-response functions is limited. This estimation approach is exemplified by Garcia et al.

(1986). Specifically, their econometric study of the impact of O₃ on producer profits used such a reduced-form approach. In this approach, farmer actions are modeled as a function of ambient O₃ levels (along with other explanatory variables) without the direct use of dose response functions. The advantage of this procedure is that one source of modeling uncertainty, the need for dose-response functions, including time-consuming crop experiments to generate data, is reduced. Actual responses of farmers' profits across air pollution gradients of ambient pollution are observed instead of hypothesized. Although this procedure has not been widely used in air pollution economic assessments, it has been used in a number of relatively recent climate change studies (e.g., Mendelsohn et al. [1994]). The reduced-form method suffers from the fact that if proposed O₃ levels are lower than those observed in the estimation sample, then the prediction accuracy of the method deteriorates. Also, some dose-response information is needed, if only to establish the plausibility of the economic estimates.

Another area that may improve economic assessments is incorporating consideration of livestock issues. To date, most agricultural economic assessments of O₃ impacts have ignored the livestock sector. Presumably this is because ambient O₃ levels do not noticeably affect meat yields. However, if feed prices and pasture conditions are affected by ambient O₃ levels, then a more accurate estimate of economic impacts would be forthcoming by including this link to livestock in the assessment. These types of feed production and feed price effects are included in the mathematical programming model used in Adams et al. (1986), but not in most other O₃ effects assessments. The significance of livestock-feed linkages are demonstrated in a recent study in the Netherlands by Kuik et al (2000). Using a mathematical programming model similar to that in Adams et al. (1986), Kuik et al. (2000) found that livestock effects were prominent, due mainly to improved pasture yields under reduced ambient O₃ levels.

9.8.6 Conclusions

Substantial progress has been made over the past two decades in our understanding of the effects of O₃ and other oxidants on vegetation, particularly for agriculturally important plant species. The physical and economic effects on agriculture are well documented and provide useful information for the setting of SNAQS. Effects on forests and natural ecosystems remain problematic, due to limitations in natural science data and economic methods. The problem is most acute for valuing natural ecosystem service flows.

1 The current limitations surrounding forests and natural ecosystems present a rich research
2 agenda. However, not all research needs are likely to lead to better policies. Thus, areas of
3 greatest potential value in terms of regional policymaking need to be prioritized. Such priority
4 setting can be assisted by sensitivity analyses with existing economic models. By measuring the
5 changes in economic effects arising from changes in key parameters, research data gaps most
6 likely to affect economic values can be identified.

9 **9.9 SUMMARY AND CONCLUSIONS FOR VEGETATION AND** 10 **ECOSYSTEM EFFECTS**

11 **9.9.1 Introduction**

12 A significant number of ozone-related studies were published between 1996 and 2004, and
13 they are reviewed in this document in the context of earlier publications reviewed in the previous
14 criteria documents (U.S. Environmental Protection Agency, 1978; 1986; 1992; 1996).

15 In general, there has been a shift away from chamber studies in favor of more field-based
16 approaches although chamber exposures still dominate the effects literature. Field-based
17 approaches include increased survey's of visible injury as well as free-air exposure systems that
18 eliminate some of the problems associated with chamber effects. Increased emphasis has also
19 been placed on quantifying aspects of ozone uptake to better link ambient exposure monitoring
20 with plant/tree response. Much of the progress in quantifying uptake has occurred in Europe
21 with application of models simulating uptake. Evaluation of this new information has added to
22 our knowledge, but has not fundamentally altered the conclusions of previous Ozone Criteria
23 Document (U.S. Environmental Protection Agency, 1996).

24 It is known that ozone is phytotoxic, and that toxicity occurs only if ozone or its reaction
25 products reach the target tissues in the plant cell. Recent studies have provided an increased
26 understanding of how ozone interacts with the plant at the cellular level. Progress has been
27 made on understanding the initial stages of ozone toxicity, which resemble a "wound" response
28 in plants. The alteration of normal metabolism due to a "wound" starts in the cytoplasm, and
29 'spreads' to other cellular organelles. One of the secondary reactions is linked to an activation of
30 a senescence response that results in loss of the critical enzymatic components of photosynthesis.
31 The loss of photosynthetic capacity is in turn linked to the lowered productivity of plants and

1 problems with efficient translocation of carbon. Although photosynthesis and translocation still
2 occurs at a reasonable rate, the loss of productivity, while better understood than in 1996, is still
3 not clearly explained. Dramatic strides in understanding the genetic make-up of plants, gene
4 control, and signal transduction/control over the last few years will accelerate in the future. This
5 understanding will translate into better models, more detailed schemes of how O₃ alters much of
6 the basic metabolism of plants, and how to construct an index that better quantifies exposure and
7 effect. However, the translation of those mechanisms into how O₃ is involved with the altered
8 cell metabolism and subsequent reductions in whole plant productivity and other physiological
9 facts has not yet been fully solved.

10 A number of biotic and abiotic factors are known, or suspected, to alter plant response to
11 ozone. New information published since 1996 does not significantly change the conclusions
12 arrived at in the previous criteria document regarding these multiple interactions. A major
13 concern from the last review was the potential, but unquantified, interaction of ozone and
14 diseases and insects. The nature of these interactions with O₃ are probably dictated by the
15 unique, highly specific biochemical relationships between pathogen and host plant. At this time,
16 therefore, although some diseases may become more widespread or severe as a result of
17 exposure to O₃, it is still not possible to predict which diseases are likely to present the greatest
18 risks to crops and forests.

19 Considerable emphasis during the last decade has been placed on research into O₃
20 interactions with the components of global climate change: increased atmospheric CO₂,
21 increased mean global temperatures, and increased surface level UV-B radiation. To date, the
22 limited experimental evidence and that obtained by computer simulation suggest that even
23 though an enriched CO₂ atmosphere (~600 ppm) would more than offset the impact of O₃ on
24 responses as varied as wheat yield or young Ponderosa growth, the concurrent increase in
25 temperature would reduce but probably not eliminate the net gain. A similar decrease in the net
26 gain resulting from the complete reversal by CO₂ of the inhibition of photosynthesis caused by
27 O₃ has been reported for increased UV-B irradiation. Overall, results are preliminary, based
28 upon minimal data, and do not allow adequate prediction of ozone effects under future climate
29 scenarios. Temperature is unquestionably also an important variable affecting plant response to
30 O₃ in the presence of the elevated CO₂ levels contributing to global climate change. In contrast,
31 evidence continues to accumulate to indicate that exposure to O₃ sensitizes plants to low

1 temperature stress, by reducing belowground carbohydrate reserves, which may lead to
2 responses in perennial species ranging from rapid demise to impaired growth in subsequent
3 seasons.

4 Efforts increased in the area of relating ozone flux (e.g., uptake) to response. Flux is
5 affected by a host of biotic and abiotic factors, including boundary layer conditions, stomatal
6 conductance, moisture conditions, solar irradiance and others. All of these factors provide a
7 more accurate estimation of flux for predicting plant, crop, or forest response. While this area of
8 research is promising, currently there is insufficient data linking flux with response across a
9 range of species or in different ecosystems. In addition, a flux-based standard would require
10 selection of a particular species or species assemblage across a meteorologically similar area in
11 order to adequately model the range of factors known to influence uptake. At this time, there is
12 insufficient information available to employ this approach nationally, but with continued
13 research may eventually provide an alternative index to the current exposure index.

15 **9.9.2 Methodology**

16 The majority of ozone effects studies are fumigation studies conducted in controlled
17 chambers, as noted in the previous document. The previous document noted that open-top
18 chambers (OTC's) represent the best technology for determination of crop yield to ozone at the
19 present time. While OTC's are still the best method for conducting controlled exposures of
20 varying length and frequency for exposure-response relationships, several new approaches have
21 been applied to ozone research, most notably free-air exposure or FACE systems. FACE
22 systems eliminate many of the concerns raised about closed or open-top chamber experiments,
23 including small plot size, altered microclimate within OTC's, and the effect of charcoal filtering
24 on overall air quality within OTC's. Although FACE systems have increased our understanding
25 in some areas, in most cases results from FACE systems have confirmed what we already
26 understood or hypothesized about how plants and plant assemblages respond to ozone.
27 In particular, studies with aspen showed that ozone symptom expression was generally similar in
28 OTC's, FACE, and also sites along an ambient ozone gradient, supporting the previously
29 observed variation among aspen clones using OTC's (Isebrands et al., 2000; 2001; Karnosky
30 et al., 1999). In addition, root growth is often found to be the most sensitive biomass response to
31 ozone, as reported in the previous document. Caution is still needed, however when making

quantitative extrapolations from OTC's, and free-air exposure systems will continue to provide valuable information in support of scaling efforts.

The lack of rural monitors continues to be a major problem in the characterization of ozone exposures in remote areas, and in linking effects with exposure in natural ecosystems. Since the last document, the use of passive samplers have expanded monitoring efforts to remote areas that were previously uncharacterized. This has greatly enhanced our ability to link ozone symptomology with elevated exposure in remote areas. Passive samplers do not capture temporal dynamics of exposure, however; therefore, passive samplers are not a substitute for active monitors when attempting to link dynamics of exposure with plant response, or when developing exposure/dose response relationships that are needed in the standard setting process.

Exclusion methods such as EDU are the least disruptive of ambient culture conditions in the field, as noted in the previous document. However, the level of protection afforded by EDU is site and species specific, and is subject to local meteorologic conditions. Because of the variability observed in the level of protection provided, and the fact that mechanisms of protection afforded by EDU and other exclusion methods are unknown, caution is needed in applying this approach to the study of ozone effects in the field.

9.9.3 Mode-of-Action

There are several steps in the process of ozone uptake and toxicity that are better understood than in 1996 and are important in developing more refined hypotheses on ozone uptake and mode of action on plants, and developing a flux-based index for use in quantifying response, and ultimately are relevant for developing an air quality standard. These are listed below:

- (1) The entrance of O₃ into the leaf through the stomata remains the critical step in O₃ sensitivity. Not only does O₃ modify the opening of the stomata, usually closing it partially, but O₃ also appears to alter the response of stomata to other stressful situations, including a lowering of water potential and ABA responses. The concentration of O₃ within the leaf is not the same as the external concentration due to reactions within the leaf but it is not "zero" (Moldau and Bichele, 2002).
- (2) The initial reactions of O₃ within the leaf are still unclear but the involvement of hydrogen peroxide is clearly indicated. The detection of possible products by EPR has progressed but is still not at the point where any products can be identified. Nonetheless, reaction of O₃ or its product with ascorbate and possibly other

antioxidants present in the apoplastic space of the mesophyll cells is clear and serves to lower the amount of O₃ or product available to alter the plasma membrane of the cells.

- 26 (3) The initial sites of membrane reactions seems to be at the transport properties and
possibly the external signal transducer molecules. The alteration and mechanism of
the alteration of the varied carriers of K⁺ and Ca²⁺ is far from clear but it would seem
that one of the primary triggers of O₃-induced cell responses is due to a change in
internal Ca²⁺ levels (Cazale et al., 1998).
- 27 (4) The primary set of metabolic reactions that O₃ triggers is now clearly the “wounding”
responses like that generated by cutting of the leaf or by pathogen/insect attack. This
seems to be due to a rise in cytoplasmic Ca²⁺. Ethylene release and alteration of
peroxidases and PAL activities as well as activation of many “wound”-derived genes
seem to be linked and some of the primary reactions to ozone exposure.
- 28 (5) The alteration of normal metabolism due to a “wound” has affects outside of the
cytoplasm. What effects are due to the “spreading of the problem” to other cellular
organelles is less clear. One of the secondary reactions is linked to an activation of a
senescence response. Certainly the loss of Rubisco and its messenger RNA is linked to
an early senescence or a speeding up of normal development leading to senescence.
Certainly the loss of photosynthetic capacity is linked to the lowered productivity of
plants and problems with efficient translocation are indicated. Although photosynthesis
and translocation still occurs in a reasonable rate, the loss of productivity is not
clearly explained.

29 The new information available on the mode of action of ozone is in part a result of
30 improved molecular tools for following rapid changes that occur within the leaf. Clearly there
31 are many changes that occur within hours or possibly days of the exposure to O₃. However,
32 there is another effect due to O₃ which takes longer to occur and tends to be most obvious under
33 exposure to low O₃ concentrations for long periods. These effects have been linked to
34 senescence or some physiological response very closely linked to senescence (i.e., translocation,
35 reabsorption, allocation of nutrients and carbon).

36 Langebartels et al. (1997) have discussed “memory” or “carry-over effects” within the
37 plant’s to explain sensitivity to frost in the winter after exposure to ozone in the summer. Others
38 have argued that it is nutrient status of the tree during the over-wintering phase of its life and
39 chronic exposure to ambient O₃ (less severe with fewer peaks of very high levels) which induces
40 (1) mineral nutrient deficiency; (2) alterations of normal metabolism, including translocation and
41 allocation of carbohydrates and probably nitrogen; (3) disturbance of normal transpiration and
42 diurnal cycling, leading to water stress (Schmieden and Wild, 1995). While general nutrient
43 concentrations within the foliage may not occur, it may be that localized deficiencies do. This is

1 hard to observe or prove without a great deal of work on all portions of a tree and without a
2 general hypothesis of what is occurring.

3 It is important to note that the dramatic strides in understanding the genetic make-up of
4 plants, gene control, and signal transduction/control over the last few years will only accelerate
5 in the future. That understanding will translate into better models of the hypotheses listed and
6 more detailed schemes of how O₃ alters much of the basic metabolism of plants. Thus overall,
7 understanding of how O₃ interacts with the plant at a cellular level has been likewise
8 dramatically improved. However, the translation of those mechanisms into how O₃ is involved
9 with the altered cell metabolism and subsequent reductions in whole plant productivity and other
10 physiological facts has not yet been fully solved. As the understanding of wounding responses
11 of plants and more genome details and varied plant mutants become available, the cellular and
12 physiological responses of plants to O₃ exposures is slowly becoming clearer. However, more
13 studies need to be made on a larger variety of species before this information can be
14 incorporated into indices of response and a protective standard.

16 **9.9.4 Modification of Growth Response**

17 There have been only a few studies reported since the 1996 document and none that
18 significantly change the conclusions of the previous criteria document with regard to factors
19 known to alter plant response to ozone.

20 In the area of biotic interactions, new evidence with regard to insect pests and diseases has
21 done little to remove the uncertainties noted in the previous criteria document (see Docherty
22 et al., 1997; and Fluckiger et al., 2002 for recent reviews). Most of the large number of such
23 interactions that may affect crops, forest trees and other natural vegetation have yet to be studied.
24 The trend suggested previously that O₃ increases the likelihood and success of insect attack has
25 received some support from recent studies, but only with respect to chewing insects. With the
26 economically important group of sucking insects such as the aphids, no clear trends have been
27 revealed by the latest studies. Hence, although it seems likely that some insect problems could
28 increase as a result of increased O₃ levels, we are still far from being able to predict the nature of
29 any particular O₃-plant-insect interaction, its likelihood or its severity.

30 The situation is a little clearer with respect to interactions involving facultative,
31 necrotrophic plant pathogens, with O₃ generally leading to increased disease. With obligate,

1 biotrophic fungal, bacterial and nematode diseases, however, there are twice as many reports
2 indicating O₃-induced inhibitions than enhancements. The frequent reports that infection by
3 obligate biotrophs reduces the severity of O₃-induced foliar injury does not result in true
4 “protection” since there are negative effects on the host plant of the disease *per se*. With
5 obligate biotrophs, the nature of any interaction with O₃ is probably dictated by the unique,
6 highly specific biochemical relationships between pathogen and host plant. At this time,
7 therefore, although some diseases may become more widespread or severe as a result of
8 exposure to O₃, it is still not possible to predict which diseases are likely to present the greatest
9 risks to crops and forests.

10 Recent studies have done little to improve our understanding of the nature of interactions
11 between O₃ and root symbionts. Several studies have indicated that the functioning of tree root
12 symbioses with mycorrhizae may be adversely affected by O₃, but there is also evidence that the
13 presence of mycorrhizae may overcome root diseases stimulated by O₃. There is also evidence
14 that O₃ may encourage the spread of mycorrhizae to the roots of uninfected trees.

15 The few recent studies of the impact of O₃ on intra-specific plant competition confirmed
16 that grasses frequently show greater resilience than other types of plants. In grass-legume
17 pastures, the leguminous species suffer greater growth inhibition (Johnson et al., 1996a;
18 Nussbaum et al., 2000a). And the suppression of Ponderosa pine seedling growth by blue wild-
19 rye grass was markedly increased by O₃ (Andersen et al., 2001). However, we are far from
20 being able to predict the outcome of the impact of O₃ on specific competitive situations such as
21 successional plant communities, or crop-weed interactions.

22 Although some recent field studies have indicated that O₃ impact significantly increases
23 with increased ambient temperature, other studies have revealed little effect of temperature.
24 Light, a component of the plant’s physical environment, is an essential ‘resource’ whose energy
25 content drives photosynthesis and CO₂ assimilation. It has been suggested that increased light
26 intensity may increase the sensitivity to O₃ of light-tolerant species, but decrease that of shade-
27 tolerant species, but this appears to be an over-simplification with many exceptions.
28 Temperature affects the rates of all physiological processes based on enzyme-catalysis and
29 diffusion, and each process and overall growth (the integral of all processes) has a distinct
30 optimal temperature range. It is important to note that a plants response to changes in
31 temperature will depend on whether it is growing near its optimum temperature for growth or

1 near its maximum temperature (Rowland-Bamford, 2000). But temperature is unquestionably an
2 important variable affecting plant response to O₃ in the presence of the elevated CO₂ levels
3 contributing to global climate change. In contrast, evidence continues to accumulate to indicate
4 that exposure to O₃ sensitizes plants to low temperature stress, by reducing belowground
5 carbohydrate reserves, which may lead to responses in perennial species ranging from rapid
6 demise to impaired growth in subsequent seasons (i.e., 'carry-over effects').

7 Although the relative humidity of the ambient air has generally been found to increase the
8 adverse effects of O₃ by increasing stomatal conductance and thereby increasing O₃ flux, there is
9 abundant evidence that the ready availability of soil moisture results in greater sensitivity to O₃.
10 The partial 'protection' against the adverse effects of O₃ afforded by drought has been observed
11 in field experiments and modeled in computer simulations (Broadmeadow and Jackson, 2000).
12 There is also compelling evidence that O₃ can predispose plants to drought stress (Maier-
13 Maercker, 1998). Hence the response will depend to some extent upon the sequence in which
14 the stresses occur, but, even though the nature of the responses is largely species-specific,
15 successful applications of model simulations have shown the way to larger scale predictions of
16 the consequences of O₃ × drought interactions. However, it must be recognized that regardless
17 of the interaction, the net result on growth in the short-term is negative, although in the case of
18 tree species, other responses such as increased water use efficiency could be a benefit to survival
19 in the long term.

20 Somewhat analogously with temperature, it appears that any shift away from the nutritional
21 optimum may lead to greater ozone sensitivity, but the shift would have to be substantial before
22 a significant effect on response to O₃ was observed. Mineral nutrients in the soil, other gaseous
23 air pollutants, and agricultural chemicals constitute chemical factors in the environment. The
24 evidence regarding interactions with specific nutrients is still contradictory: there is some
25 experimental evidence that low general fertility increases sensitivity to O₃, while simulation
26 modeling of trees suggests that nutrient deficiency and O₃ act less than additively, but there are
27 too many example of contrary trends to permit any sweeping conclusions.

28 Interactions of O₃ with other air pollutants have received relatively little recent attention
29 since 1996 (see Barnes and Wellburn, 1998; and Fangmeier et al., 2002 for recent reviews).
30 The situation with SO₂ remains inconsistent, but seems unlikely to pose any additional risk to
31 those related to the individual pollutants. With the NO and NO₂ the situation is complicated by

1 their nutritional value as a source of N. In leguminous species, it appears that NO₂ may reduce
2 the impact of O₃ on growth, with the reverse in other species, but the nature of the exposure
3 pattern, i.e., sequential or concurrent, also determines the outcome. Much more investigation is
4 needed before we will be able to predict the outcomes of different O₃-NO-NO₂ scenarios. The
5 latest research into O₃ × acid rain interactions has confirmed that, at realistic acidities, significant
6 interactions are unlikely. A continuing lack of information precludes offering any
7 generalizations about interactive effects of O₃ with NH₃, HF, or heavy metals. More evidence
8 has been reported for protective effects against O₃ afforded by the application of fungicides.

9 Considerable emphasis during the last decade has been placed on research into O₃
10 interactions with the components of global climate change: increased atmospheric CO₂,
11 increased mean global temperatures, and increased surface level UV-B radiation. However, it
12 must be noted that most of these studies have tended to regard increased CO₂ levels and
13 increased mean temperatures as unrelated phenomena. Experiments into the effects of doubled
14 CO₂ levels at today's mean ambient temperatures are of questionable value in trying to assess the
15 impact of *climate change* on responses to O₃. To date, the limited experimental evidence and
16 that obtained by computer simulation suggest that even though an enriched CO₂ atmosphere
17 (~600 ppm) would more than offset the impact of O₃ on responses as varied as wheat yield or
18 young Ponderosa growth, the concurrent increase in temperature would reduce but probably not
19 eliminate the net gain. A similar decrease in the net gain resulting from the complete reversal by
20 CO₂ of the inhibition of photosynthesis caused by O₃ has been reported for increased UV-B
21 irradiation. However, these are preliminary results based upon minimal data.

22 23 **9.9.5 Exposure Indices**

24 The previous Criteria Document (U.S. Environmental Protection Agency, 1996a) noted
25 that the complexities associated with uptake and ozone interactions with external physical and
26 internal genetic factors that influence plant response makes the development of an 'ideal'
27 exposure index that characterizes plant exposure and response extremely difficult. As a result, a
28 biologically relevant surrogate for uptake was recommended that related ambient exposure to
29 measured growth/yield response. Despite additional research linking estimates of flux with plant
30 response since 1996, there is still insufficient information to identify a flux-based model that
31 incorporates sufficient complexity across space and time to be non-site or non-species specific.

1 Therefore, based on the current state of knowledge, exposure indices that cumulate and
2 differentially weight the higher hourly average concentrations, but include the mid-level values,
3 still represents the best approach in the United States for relating vegetation effects to exposure.

4 The few studies published in this interim have substantiated earlier conclusions on the role
5 of exposure components, including concentration, duration and exposure patterns, in determining
6 the growth response of plants to ozone (Yun and Laurence, 1999; Oksanen and Holopainen,
7 2001). Recent studies using different exposure patterns have confirmed earlier studies on the
8 role of higher concentrations and exposure duration (Nussbaum et. al., 1995). A role for peak
9 concentrations is inferred based on improved air quality in regions like the San Bernardino
10 Mountains in southern California. The reductions of O₃ in the San Bernardino area are
11 associated with reductions in the higher hourly average concentrations, but not in the midrange
12 hourly average concentrations, which are increasing (Lee et al., 2003; Lefohn and Shadwick,
13 2000). General forest improvement is reported following a decrease of O₃ along a decreasing
14 gradient of exposure (Miller and Rechel, 1999; Arbaugh et al., 2003; Tingey et al., 2004). These
15 studies provide the basis for focusing on higher O₃ concentrations, while including the lower
16 levels, when estimating the effects of emission reductions on vegetation.

17 New studies have demonstrated the potential disconnection of peak events and maximal
18 stomatal conductance at xeric to mesic sites in California (Panek et al., 2002; Panek, 2004;
19 Grulke et al., 2002). In addition, a few studies have indicated the uptake of ozone during
20 nighttime hours is greater than previously thought (Grulke et al., 2003; Massman, 2004), and a
21 review of the literature suggests a large number of species exhibit some degree of conductance at
22 night (Musselman and Minnick, 2000). These studies suggest a reconsideration of the need to
23 cumulate exposure 24 hrs per day and not just during daylight hours in exposure index. This
24 lack of coincidence in temporal patterns of conductance and peak ambient concentrations
25 introduces uncertainty in assessing ozone's impact. The use of an exposure index that does not
26 consider regionally unique climate and site factors which modify stomatal conductance may, as a
27 result, under or over estimate growth effects. The shortcomings of an ambient exposure-based
28 index is especially apparent when assessing ozone's potential impact across broad climatic
29 regions of the U.S. or Europe.

30 It is apparent that additional research is needed to develop indices which are more
31 physiologically- and meteorologically-connected to the actual dose of ozone the plant receives.

1 The cumulative, concentration-weighted exposure indices are acknowledged surrogates for
2 effective dose that are conceptually simple and easy to measure. As discussed in the previous
3 Criteria Document, they do not fully characterize the potential for plant uptake and subsequent
4 response since they do not include the physical, biological, and meteorological processes
5 controlling the transfer of O₃ from the atmosphere through the leaf and into the leaf interior.

6 The flux-based approach should provide an opportunity to improve upon the
7 concentration-based (i.e., exposure indices) approach. A great deal of progress has occurred in
8 development and testing of stomatal models that may be generally applicable across certain
9 vegetation types (e.g., Matyssek et al., 2004; Grunhage and Jager 2003; Danielsson et al., 2003;
10 Emberson et al., 2000; Pleijel et al., 2000). While a flux-based approach is preferred, a
11 cautionary argument was advanced in a few publications based on the non-linear relationship
12 between ozone uptake and foliar injury (growth not assessed). The concern is that not all O₃
13 stomatal uptake results in a reduction in yield, which depends to some degree on the amount of
14 internal detoxification occurring with each particular species. Those species having high
15 amounts of detoxification potential may show little relationship between O₃ stomatal uptake and
16 plant response (Musselman and Massman, 1999).

17 The European approach and acceptance of flux-based critical values is in part a recognition
18 of the problems associated with ambient exposure-based indices. Research continues across
19 Europe to develop the necessary experimental database and modeling tools that will be required
20 to provide the scientific basis for a critical level for ozone (Grunhage et al., 2004; Fuhrer et al.,
21 1997; Grunhage and Jager, 1994).

22 Given the current state of knowledge and the best available data, exposure indices that
23 cumulate and differentially weight the higher hourly average concentrations, but include the
24 mid-level values, continue to offer the most defensible approach for vegetation protection.
25 A large database exists that has been used for establishing exposure-response relationships, and
26 at this time, such a database does not exist for relating ozone flux to growth response.

27 It is anticipated that as the overlapping relationships of conductance, concentration, and
28 defense mechanisms are better defined, the flux-based indices may be able to predict vegetation
29 injury and/or damage with more accuracy than the exposure-response models. However, it is
30 unclear that such is the case at this time. The translation of these research and assessment tools

to air quality standards has the additional need to be simple, understandable, and adaptive to a manageable monitoring program.

9.9.6 Ozone Exposure-Plant Response Relationships

Data published since 1996 continue to support the conclusions of previous criteria documents that there is strong evidence that ambient concentrations of O₃ cause foliar injury and growth and yield damage to numerous common and economically valuable plant and tree species. For annual vegetation, the data summarized in Table 4-19 show a range of growth and yield responses both within species and among species. Nearly all of these data were derived from studies in OTCs, with only two studies using open-air systems in the UK (Ollerenshaw et al., 1999; Ollerenshaw and Lyons, 1999). It continues to be difficult to compare studies that report O₃ exposure using different indices, such as AOT40, SUM06, or 7-h or 12-h mean values. However, when such comparisons can be made, the results of recent research confirm earlier results summarized in the previous criteria document (U.S. Environmental Protection Agency, 1996). A summary of earlier literature concluded that a 7-h 3-month mean of 49 ppb corresponding to a SUM06 exposure of 26 ppm·h, would cause 10% loss in 50% of 49 experimental cases (Tingey et al., 1991). Recent data summarized in Table 4-19 support this conclusion, and more generally indicate that ambient ozone exposures can reduce the growth and yield of annual species. Some annual species such as soybean are more sensitive, and greater losses may be expected (Table 4-19). Thus the recent scientific literature supports the conclusions of the previous criteria document that ambient O₃ concentrations are reducing the yield of major crops in the United States.

Much research in Europe has used the AOT40 exposure statistic, and substantial effort has gone into developing Level-1 critical levels for vegetation using this index. Based on regression analysis of 15 OTC studies of spring wheat including one study from the U.S. and 14 from locations ranging from southern Sweden to Switzerland, an AOT40 value of 5.7 ppm·h was found to correspond to a 10% yield loss, and a value of 2.8 ppm·h corresponded to a 5% yield loss (Fuhrer et al., 1997). Because a 4 to 5% decrease could be detected with a confidence level of 99%, a critical level of an AOT40 value of 3 ppm·h was selected in 1996 (Kärenlampi and Skärby, 1996).

1 In addition to reductions in crop yield, O₃ may also reduce the quality or nutritive value of
2 annual species. Many studies have shown effects of O₃ on various measures of plant organs that
3 affect quality, with most studies focusing on characteristics important for food or fodder. These
4 studies indicate that there may be economically important effects of ambient O₃ on the quality of
5 crop and forage species. Previous criteria documents have concluded that visible symptoms on
6 marketable portions of crops and ornamental plants can occur with seasonal 7-h mean O₃
7 exposures of 40 to 100 ppb (U.S. Environmental Protection Agency, 1978; 1986; 1996). The
8 recent scientific literature does not refute this conclusion.

9 The use of OTCs may reverse the usual vertical gradient in O₃ that occurs within a few
10 meters above the ground surface (Section 4.2). This reversal suggests that OTC studies may
11 overestimate to some degree the effects of an O₃ concentration measured several meters above
12 the ground. However such considerations do not invalidate the conclusion of the previous
13 criteria document that ambient O₃ exposures are sufficient to reduce the yield of major crops in
14 the United States.

15 As for single-season agricultural crops, yields of multiple-year forage crops are reduced at
16 ozone exposures that occur over large areas of the U.S. This result is similar to that reported in
17 the previous criteria document (U.S. Environmental Protection Agency, 1996). When species
18 are grown in mixtures, O₃ exposure can increase the growth of tolerant species while
19 exacerbating the growth decrease of O₃-sensitive species. Because of this competitive
20 interaction, the total growth of the mixed-species community may not be affected by O₃
21 exposure. However, in some cases mixtures of grasses and clover species have shown
22 significant decreases in total biomass growth in response to O₃ exposure in studies in the U.S.
23 and in Sweden. In Europe, a provisional critical level for herbaceous perennials of an AOT40
24 value of 7 ppm·h over six months has been proposed to protect sensitive plant species from
25 adverse effects of O₃.

26 For deciduous tree species, recent evidence from free air exposure systems and OTCs
27 supports results observed previously in OTC studies. For example, a series of studies was
28 undertaken using free air O₃ enrichment in Rhinelander, WI (Isebrands et al., 2000; 2001).
29 These studies showed that O₃ symptom expression was generally similar in OTCs, FACE, and
30 also sites along an ambient O₃ gradient, supporting the previously observed variation among

1 aspen clones using OTCs (Karnosky et al., 1999). As has been observed in previous criteria
2 documents, root growth often is found to be the most sensitive biomass response to O₃.

3 Results since 1996 support the conclusion of the previous criteria document (U.S.
4 Environmental Protection Agency, 1996) that deciduous trees are generally less sensitive to
5 O₃ than are most annual plants, with the exception of a few very sensitive genera such as
6 *Populus* and sensitive species such as black cherry. However, the data presented in Table 4-20
7 suggest that ambient exposures that occur in the U.S. can sometimes reduce the growth of
8 seedlings of deciduous species. Results from multiple-year studies sometimes show a pattern of
9 increased effects in subsequent years. In some cases however, growth decreases due to O₃ may
10 become less significant or even disappear over time. While some mature trees show greater O₃
11 sensitivity than do seedlings in physiological parameters such as net photosynthetic rate, these
12 effects may not translate into measurable reductions in biomass growth. However, because even
13 multiple-year experiments do not expose trees to O₃ for more than a small fraction of their life
14 span, and because competition may in some cases exacerbate the effects of O₃ on individual
15 species, determining effects on mature trees remains a significant challenge.

16 In Europe, a Level I critical level has been set for forest trees based on OTC studies of
17 European beech seedlings. A critical level was defined as an AOT40 value of 10 ppm·h for
18 daylight hours for a 6-month growing season (Kärenlampi and Skärby, 1996). However, other
19 studies show that other species such as silver birch may be more sensitive to O₃ than is beech
20 (Pääkkönen et al., 1996).

21 For evergreen tree species as for other tree species, the O₃ sensitivity of different genotypes
22 and different species varies widely. Based on studies with seedlings in OTCs, major species in
23 the US are generally less sensitive than are most deciduous trees, and slower growing species are
24 less sensitive than are faster growing species. There is evidence that interacting stresses such as
25 competition may increase the sensitivity of trees to O₃. As for deciduous species, most
26 experiments with evergreen species have only covered a very small portion of the life span of a
27 tree, and have been conducted with seedlings, so estimating effects on mature trees is difficult.

28 For all types of perennial vegetation, cumulative effects over more than one growing
29 season may be important, and studies for only a single season may underestimate effects.
30 Mature trees may be more or less sensitive to O₃ than are seedlings depending on the species,
31 but information on physiological traits can be used to predict such differences in some cases.

1 In some cases, mature trees may be more sensitive to O₃ than seedlings due to differences in gas
2 exchange rates, differences in growth rates, greater cumulative exposure, or the interaction of
3 ozone with other stresses.
4

5 **9.9.7 Ecosystem Effects**

6 There is evidence that tropospheric O₃ is an important stressor of ecosystems, with
7 documented impacts on the biotic condition, ecological processes, and chemical/physical nature
8 of natural ecosystems. In turn, the effects of O₃ on individual plants and processes are scaled up
9 through the ecosystem affecting processes such as energy and material flow, inter- and intra-
10 species competition, and NPP. Thus, effects on individual keystone species and their associated
11 microflora and fauna, which has been shown experimentally, may cascade through the
12 ecosystem to the landscape level, although it has not yet been demonstrated. By affecting water
13 balance, cold hardiness, tolerance to wind, and by predisposing plants to insect and disease pests,
14 O₃ may even impact the occurrence and impact of natural disturbance (e.g., fire, erosion).
15 Despite the probable occurrence of such effects, there are essentially no instances where
16 ecosystem level, highly integrated studies have conclusively shown that ozone is indeed altering
17 ecosystem structure and/or function.

18 Systematic injury surveys demonstrate that foliar injury occurs on sensitive species in
19 many regions of the USA. However, the frequent lack of correspondence between foliar
20 symptoms and growth effects means that other methods must be used to estimate the regional
21 effects of O₃ on tree growth rates. Investigations of the radial growth of mature trees in
22 combination with data from many controlled studies with seedlings and a few studies with
23 mature trees suggest that ambient O₃ is reducing the growth of mature trees in some locations.
24 Studies using models based on tree physiology and forest stand dynamics suggest that modest
25 effects of O₃ on growth may accumulate over time, and may interact with other stresses. For
26 mixed-species stands, such models predict that overall stand growth rate is generally not likely to
27 be affected. However, competitive interactions among species may change as a result of growth
28 reductions of sensitive species. These results suggest that O₃ exposure over decades may be
29 altering the species composition of forests in some regions.

30 Despite increased understanding of possible ecosystem effects of ozone, the data base
31 demonstrating and quantifying the degree to which O₃ is altering natural ecosystems is lacking.

1 Much of the speculation of ozone impact on ecosystems must be inferred from a number of case
2 studies of forest plot field-based data reporting on a number of different species. One means to
3 discuss our current knowledge is by listing the areas in which there is lack of knowledge. These
4 include:

5 *Ecosystem processes.* Very little is known about the effects of O₃ on water, carbon, and
6 nutrient cycling, particularly at the stand and community levels. Effects on belowground
7 ecosystem processes in response to O₃, exposure alone and in combination with other stressors
8 are critical to projections at the watershed and landscape scales. Little is yet known about the
9 effects of O₃ on structural or functional components of soil food webs or how these impacts
10 could affect plant species diversity (Andersen, 2003).

11 *Biodiversity and genetic diversity.* The study of genetic aspects of O₃ impacts on natural
12 ecosystems has been largely based on correlations and it remains to be shown conclusively
13 whether O₃ affects biodiversity or genetic diversity (Pitelka, 1988; Winner et al., 1991; Davison
14 & Barnes, 1998). Studies of competitive interactions under elevated O₃ are needed (Laurence &
15 Andersen, 2003), and re-examination via new sampling of population studies to bring in a time
16 component to previous studies showing spatial variability in population responses to O₃ are
17 needed. These studies could be strengthened by modern molecular methods to quantify impacts
18 on diversity.

19 *Natural ecosystem interactions with the atmosphere.* Little is known about feedbacks
20 between O₃ and climate change on production of volatile organic compounds, which in turn,
21 could affect O₃ production (Fuentes et al., 2001). At moderate to high O₃ exposure sites,
22 aberrations in stomatal behavior could significantly affect individual tree water balance of
23 sensitive trees, and if the sensitive tree species is dominant, hydrologic balance at the watershed
24 and landscape level could be affected. This has not been addressed in any model because O₃
25 exposure effects, if included at all in the modeling effort, have assumed a linear relationship
26 between assimilation and stomatal conductance. Interaction studies with other components of
27 global change (i.e., warming, increasing atmospheric CO₂, N deposition, etc.) or with various
28 biotic stressors are needed to better predict complex interactions likely in the future (Laurence &
29 Andersen, 2003). Whether O₃ will negate the positive effects of an elevated CO₂ environment
30 on plant carbon and water balances is not yet known nor is it known if these effects will scale-up

1 through the ecosystem. How O₃ affects the progress of pest epidemics and insect outbreaks as
2 concentrations increase is unclear (Skarby et al., 1998).

3 Information concerning the impact of O₃ on reproductive processes and reproductive
4 development under realistic field or forest conditions are needed as well as examination of
5 reproductive effects under interacting pollutants (Black et al., 2000).

6 *Scaling.* The vast majority of O₃ studies of trees have been conducted with young,
7 immature trees and in trees that have not yet formed a closed canopy. Questions remain as to the
8 comparability of O₃ effects on juvenile and mature trees and on trees grown in the open versus
9 those in a closed forest canopy in a competitive environment (Chappelka & Samuelson, 1998;
10 Kolb & Matyssek, 2001; Samuelson & Kelly, 2001). Scaling the effects of O₃ across spatial
11 scales is also difficult. Scaling responses of single or a few plants to effects on communities and
12 ecosystems are complicated matters that will require a combination of manipulative experiments
13 with model ecosystems, community and ecosystem studies along natural O₃ gradients, and
14 extensive modeling efforts to project landscape level, regional, national and international
15 impacts of O₃. Linking these various studies via impacts on common research quantification
16 across various scales using measures of such factors as leaf area index or spectral reflective data,
17 which could eventually be remotely sensed (Kraft et al., 1996; Panek et al., 2003), would provide
18 powerful new tools for ecologists.

19 *Identifying endpoints.* In general, methodologies to determine the important values of
20 services and benefits derived from natural ecosystems are lacking. Identifying and quantifying
21 factors that could be used in comprehensive risk assessment for O₃ effects on natural ecosystems
22 would increase societal awareness of the importance of protecting ecosystems (Heck et al.,
23 1998).

24 25 **9.9.8 Economics**

26 Substantial progress has been made over the past two decades in our understanding of the
27 effects of ozone and other oxidants on vegetation, particularly for agriculturally-important plant
28 species. The physical and economic effects on agriculture are well documented and provide
29 useful information for the consideration of establishing air quality standards for crops. Effects
30 on forests and natural ecosystems remain problematic, due to limitations in biological response

1 data and economic methods. The problem is even more acute for valuing natural ecosystem
2 service flows.

3 The current limitations surrounding forests and natural ecosystems present a rich research
4 agenda. However, not all research needs are likely to lead to better policies. Thus, areas of
5 greatest potential value in terms of regional policy making need to be prioritized. Such
6 priority-setting can be assisted by sensitivity analyses with existing economic models; by
7 measuring the changes in economic effects arising from changes in key parameters, it is possible
8 to identify those research data gaps most likely to affect economic values.

REFERENCES

- Abeles, F. B.; Biles, C. L.; Dunn, L. J. (1989a) Hormonal regulation and distribution of peroxidase isoenzymes in the cucurbitaceae. *Plant Physiol.* 91: 1609-1612.
- Abeles, F. B.; Hershberger, W. L.; Dunn, L. J. (1989b) Hormonal regulation, and intracellular localization of a 33-kD cationic peroxidase in excised cucumber cotyledons. *Plant Physiol.* 89: 664-668.
- Adams, R. M. (1986) Agriculture, forestry, and related benefits of air pollution control: a review and some observations. *Am. J. Agric. Econ.* 68: 464-472.
- Adams, R. M. (1999) On the search for the correct economic assessment method. An editorial comment. *Clim. Change* 41: 363-370.
- Adams, W. C. (2002) Comparison of chamber and face-mask 6.6-hour exposures to ozone on pulmonary function and symptoms responses. *Inhalation Toxicol.* 14: 745-764.
- Adams, R. M.; Crocker, T. D. (1989) The agricultural economics of environmental change: some lessons from air pollution. *J. Environ. Manage.* 28: 295-307.
- Adams, R. M.; Horst, R. L., Jr. (2003) Future directions in air quality research: economic issues. *Environ. Int.* 29: 289-302.
- Adams, R. M.; Rowe, R. D. (1990) The economic effects of stratospheric ozone depletion on U.S. agriculture: a preliminary assessment. *J. Environ. Manage.* 30: 321-335.
- Adams, R. M.; Hamilton, S. A.; McCarl, B. A. (1986) The benefits of pollution control: the case of ozone and U.S. agriculture. *Am. J. Agric. Econ.* 68: 886-893.
- Adams, R. M.; Glyer, J. D.; McCarl, B. A. (1988) The NCLAN economic assessment: approach, findings and implications. In: Heck, W. W.; Taylor, O. C.; Tingey, D. T., eds. *Assessment of crop loss from air pollutants: proceedings of an international conference; October 1987; Raleigh, NC.* New York, NY: Elsevier Applied Science; pp. 473-504.
- Adams, D. M.; Alig, R. J.; Callaway, J. M.; McCarl, B. A.; Winnett, S. M. (1996) The forest and agricultural sector optimization model (FASOM): model structure and policy application. Portland, OR: U.S. Department of Agriculture, Pacific Northwest Research Station, PNW-RP-495.
- Adams, R. M.; Hurd, B. H.; Lenhart, S.; Leary, N. (1998) Effects of global climate change on agriculture: an interpretative review. *Climate Research* 11: 19-30.
- Agrawal, M.; Agrawal, S. B. (2000) Effects of air pollution on plant diversity. In: Agrawal, S. B.; Agrawal, M., eds. *Environmental Pollution and Plant Responses.* Boca Raton, FL: Lewis Publishers; pp. 137-152.
- Ainsworth, N.; Fumagalli, I.; Giorcelli, A.; Mignanego, L.; Schenoe, G.; Vietto, L. (1996) Assessment of EDU stem injections as a technique to investigate the response of trees to ambient ozone in field conditions. *Agric. Ecosyst. Environ.* 59: 33-42.
- Ainsworth, E. A.; Davey, P. A.; Bernacchi, C. J.; Dermody, O. C.; Heaton, E. A.; Moore, D. J.; Morgan, P. B.; Naidu, S. L.; Yoora, H.-S.; Zhu, X.-G.; Curtis, P. S.; Long, S. P. (2002) A meta-analysis of elevated [CO₂] effects on soybean (*Glycine max*) physiology, growth and yield. *Global Change Biol.* 8: 695-709.
- Alscher, R. G.; Wellburn, A. R., eds. (1994) *Plant responses to the gaseous environment. Molecular, metabolic and physiological aspects.* London, UK: Chapman and Hall.
- Altshuller, A. P.; Lefohn, A. S. (1996) Background ozone in the planetary boundary layer over the United States. *J. Air Waste Manage. Assoc.* 46: 134-141.
- Amiro, B. D.; Gillespie, T. J. (1985) Leaf conductance response of *Phaseolus vulgaris* to ozone flux density. *Atmos. Environ.* 19: 807-810.
- Amiro, B. D.; Gillespie, T. J.; Thurtell, G. W. (1984) Injury response of *Phaseolus vulgaris* to ozone flux density. *Atmos. Environ.* 18: 1207-1215.
- Amthor, J. S.; Goulden, M. L.; Munger, J. W.; Wofsy, S. C. (1994) Testing a mechanistic model of forest-canopy mass and energy exchange using eddy correlation: carbon dioxide and ozone uptake by a mixed oak-maple stand. *Aust. J. Plant Physiol.* 21: 623-651.
- Anderson, E. (1990) The ethical limitations of the market. *Econ. Philos.* 6: 179-205.
- Andersen, C. P. (2000) Ozone stress and changes below-ground: linking root and soil processes. *Phyton* 40(Special issue 4): 7-12.
- Andersen, C. P. (2003) Source-sink balance and carbon allocation below ground in plants exposed to ozone. *New Phytol.* 157: 213-228.
- Andersen, C. P.; Rygielwicz, P. T. (1995) Allocation of carbon in mycorrhizal *Pinus ponderosa* seedlings exposed to ozone. *New Phytol.* 131: 471-480.

- 1 Andersen, C. P.; Scagel, C. F. (1997) Nutrient availability alters belowground respiration of ozone-exposed
2 ponderosa pine. *Tree Physiol.* 17: 377-387.
- 3 Anderson, R. L.; Brown, H. D.; Chevone, B. I.; McCartney, T. C. (1988) Occurrence of air pollution symptoms
4 (needle tip necrosis and chlorotic mottling) on eastern white pine in the southern Appalachian Mountains.
5 *Plant Dis.* 72: 130-132.
- 6 Andersen, C. P.; Hogsett, W. E.; Wessling, R.; Plocher, M. (1991) Ozone decreases spring root growth and root
7 carbohydrate content in ponderosa pine the year following exposure. *Can. J. For. Res.* 21: 1288-1291.
- 8 Andersen, C. P.; Wilson, R.; Plocher, M.; Hogsett, W. E. (1997) Carry-over effects of ozone on root growth and
9 carbohydrate concentrations of ponderosa pine seedlings. *Tree Physiol.* 17: 805-811.
- 10 Andersen, C. P.; Hogsett, W. E.; Plocher, M.; Rodecap, K.; Lee, E. H. (2001) Blue wild-rye grass competition
11 increases the effect of ozone on ponderosa pine seedlings. *Tree Physiol.* 21: 319-327.
- 12 Antonielli, M.; Pasqualini, S.; Ederli, L.; Batini, P.; Moscatello, S.; Loreto, F. (1997) Physiological characteristics of
13 tobacco cultivars with contrasting sensitivity to ozone. *Environ. Exp. Bot.* 38: 271-277.
- 14 Anttonen, S.; Sutinen, M. L.; Heagle, A. S. (1996) Ultrastructure and some plasma membrane characteristics of
15 ozone-exposed loblolly pine needles. *Physiol. Plant.* 98: 309-319.
- 16 Aono, M.; Saji, H.; Fujiyama, K.; Sugita, M.; Kondo, N.; Tanaka, K. (1995) Decrease in activity of glutathione
17 reductase enhances paraquat sensitivity in transgenic *Nicotiana tabacum*. *Plant Physiol.* 107: 645-648.
- 18 Arbaugh, M. J.; Miller, P. R.; Carroll, J. J.; Takemoto, B. L.; Proctor, T. (1998) Relationships of ozone exposure to
19 pine injury in the Sierra Nevada and San Bernardino Mountains of California, USA. *Environ. Pollut.*
20 101: 291-301.
- 21 Arbaugh, M.; Bytnerowicz, A.; Grulke, N.; Fenn, M.; Poth, M.; Temple, P.; Miller, P. (2003) Photochemical smog
22 effects in mixed conifer forests along a natural gradient of ozone and nitrogen deposition in the
23 San Bernardino Mountains. *Environ. Int.* 29: 401-406.
- 24 Ariens, E. J.; Simonis, A. M.; Offermeier, J. (1976) Introduction to general toxicology. New York, NY:
25 Academic Press.
- 26 Arndt, U. (1995) Air pollutants and pheromones - a problem? *Chemosphere* 30: 1023-1031.
- 27 Arp, W. J.; Drake, B. G. (1991) Increased photosynthetic capacity of *Scirpus olneyi* after 4 years of exposure to
28 elevated CO₂. *Plant Cell Environ.* 14: 1003-1006.
- 29 Asada, K. (1999) The water-water cycle in chloroplasts: scavenging of active oxygens and dissipation of excess
30 photons. *Ann. Rev. Plant Physiol. Mol. Biol.* 50: 601-639.
- 31 Ashenden, T. W.; Bell, S. A.; Rafarel, C. R. (1995) Responses of white clover to gaseous pollutants and acid mist:
32 implications for setting critical levels and loads. *New Phytol.* 130: 89-96.
- 33 Ashenden, T. W.; Bell, S. A.; Rafarel, C. R. (1996) Interactive effects of gaseous air pollutants and acid mist on two
34 major pasture grasses. *Agric. Ecosyst. Environ.* 57: 1-8.
- 35 Ashmore, M. R. (1984) Effects of ozone on vegetation in the United Kingdom. In: Grennfelt, P., ed. Proceedings of
36 the international workshop on the evaluation of the effects of photochemical oxidants on human health,
37 agricultural crops, forestry, materials and visibility; February-March; Göteborg, Sweden. Göteborg, Sweden:
38 Swedish Environmental Research Institute; pp. 92-104; report no. IVL-EM 1570.
- 39 Ashmore, M. R.; Ainsworth, N. (1995) The effects of ozone and cutting on the species composition of artificial
40 grassland communities. *Funct. Ecol.* 9: 708-712.
- 41 Ashmore, M.; Fuhrer, J. (2000) New directions: use and abuse of the AOT40 concept. *Atmos. Environ.*
42 34: 1157-1158.
- 43 Ashmore, M. R.; Bell, J. N. B.; Mimmack, A. (1988) Crop growth along a gradient of ambient air pollution.
44 *Environ. Pollut.* 53: 99-121.
- 45 Ashmore, M. R.; Thwaites, R. H.; Ainsworth, N.; Cousins, D. A.; Power, S. A.; Morton, A. J. (1995) Effects of
46 ozone on calcareous grassland communities. *Water Air Soil Pollut.* 85: 1527-1532.
- 47 Atkinson, R. (1990) Gas-phase tropospheric chemistry of organic compounds: a review. *Atmos. Environ. Part A*
48 24: 1-41.
- 49 Auh, C.-K.; Murphy, T. M. (1995) Plasma membrane redox enzyme is involved in the synthesis of O₂⁻ and H₂O₂ by
50 *Phytophthora elicitor*-stimulated rose cells. *Plant Physiol.* 107: 1241-1247.
- 51 Awmack, C. S.; Harrington, R.; Lindroth, R. L. (2004) Aphid individual performance may not predict population
52 responses to elevated CO₂ or O₃. *Global Change Biol.* 10: 1414-1423.
- 53 Ayres, P. G. (1991) Growth responses induced by pathogens and other stresses. In: Mooney, H. A.; Winner, W. E.;
54 Pell, E. J., eds. Responses of plants to multiple stresses. San Diego, CA: Academic Press; pp. 227-248.

- 1 Azevedo, R. A.; Alas, R. M.; Smith, R. J.; Lea, P. J. (1998) Response of antioxidant enzymes to transfer from
2 elevated carbon dioxide to air and ozone fumigation, in the leaves and roots of wild-type and a
3 catalase-deficient mutant of barley. *Physiol. Plant.* 104: 280-292.
- 4 Bäck, J.; Vanderklein, D. W.; Topa, M. A. (1999) Effects of elevated ozone on CO₂ uptake and leaf structure in sugar
5 maple under two light environments. *Plant Cell Environ.* 22: 137-147.
- 6 Bahl, A.; Kahl, G. (1995) Air pollutant stress changes the steady-state transcript levels of three photosynthesis genes.
7 *Environ. Pollut.* 88: 57-65.
- 8 Bahl, A.; Loitsch, S. M.; Kahl, G. (1995) Transcriptional activation of plant defence genes by short-term air pollutant
9 stress. *Environ. Pollut.* 89: 221-227.
- 10 Baker, T. R.; Allen, H. L.; Schoeneberger, M. M.; Kress, L. W. (1994) Nutritional response of loblolly pine exposed
11 to ozone and simulated acid rain. *Can. J. For. Res.* 24: 453-461.
- 12 Baker, C. J.; Harmon, G. L.; Glazener, J. A.; Orlandi, E. W. (1995) A noninvasive technique for monitoring
13 peroxidative and H₂O₂-scavenging activities during interactions between bacterial plant pathogens and
14 suspension cells. *Plant Physiol.* 108: 353-359.
- 15 Balaguer, L.; Barnes, J. D.; Panicucci, A.; Borland, A. M. (1995) Production and utilization of assimilates in wheat
16 (*Triticum aestivum* L.) leaves exposed to elevated O₃ and/or CO₂. *New Phytol.* 129: 557-568.
- 17 Ball, J. T. (1987) Calculations related to gas exchange. In: Zeiger, E.; Farquhar, G. D.; Cowan, I. R. *Stomatal*
18 *Function*. Stanford, CA: Stanford Press; pp. 445-476, pp. G-47.
- 19 Ball, G. R.; Benton, J.; Palmer-Brown, D.; Fuhrer, J.; Skärby, L.; Gimeno, B. S.; Mills, G. (1998) Identifying factors
20 which modify the effects of ambient ozone on white clover (*Trifolium repens*) in Europe. *Environ. Pollut.*
21 103: 7-16.
- 22 Ball, G. R.; Palmer-Brown, D.; Fuhrer, J.; Skärby, L.; Gimeno, B. S.; Mills, G. (2000) Identification of non-linear
23 influences on the seasonal ozone dose-response of sensitive and resistant clover clones using artificial neural
24 networks. *Ecol. Model.* 129: 153-168.
- 25 Balls, G. R.; Palmer-Brown, D.; Cobb, A. H.; Sanders, G. E. (1995) Towards unravelling the complex interactions
26 between microclimate, ozone dose, and ozone injury in clover. *Water Air Soil. Pollut.* 85: 1467-1472.
- 27 Balls, G. R.; Palmer-Brown, D.; Sanders, G. E. (1996) Investigating microclimatic influences on ozone injury in
28 clover (*Trifolium subterraneum*) using artificial neural networks. *New Phytol.* 132: 271-280.
- 29 Banerjee, S. K.; Mudd, J. B. (1992) Reaction of ozone with glycophorin in solution and in lipid vesicles.
30 *Arch. Biochem. Biophys.* 295: 84-89.
- 31 Barbo, D. N.; Chappelka, A. H.; Somers, G. L.; Miller-Goodman, M. S.; Stolte, K. (1998) Diversity of an early
32 successional plant community as influenced by ozone. *New Phytol.* 138: 653-662.
- 33 Barbo, D. N.; Chappelka, A. H.; Somers, G. L.; Miller-Goodman, M. S.; Stolte, K. (2002) Ozone impacts on loblolly
34 pine (*Pinus taeda* L.) grown in a competitive environment. *Environ. Pollut.* 116: 27-36.
- 35 Barnes, J. D.; Pfirrmann, T. (1992) The influence of CO₂ and O₃, singly and in combination, on gas exchange, growth
36 and nutrient status of radish (*Raphanus sativus* L.). *New Phytol.* 121: 403-412.
- 37 Barnes, J. D.; Wellburn, A. R. (1998) Air pollutant combinations. In: De Kok, L. J.; Stulen, I., eds. *Responses of*
38 *plant metabolism to air pollution and global change*. Leiden, The Netherlands: Backhuys Publishers;
39 pp. 147-164.
- 40 Barnes, J. D.; Ollerenshaw, J. H.; Whitfield, C. P. (1995) Effects of elevated CO₂ and/or O₃ on growth, development
41 and physiology of wheat (*Triticum aestivum* L.). *Global Change Biol.* 1: 129-142.
- 42 Barnes, J.; Zheng, Y.; Lyons, T. (2002) Plant resistance to ozone: the role of ascorbate. In: Omasa, K.; Saji, H.;
43 Youssefian, S.; Kondo, N., eds. *Air pollution and plant biotechnology - prospects for phytomonitoring and*
44 *phytoremediation*. Springer-Verlag; pp. 235-252.
- 45 Barrett, S. C. H.; Bush, E. J. (1991) Population processes in plants and the evolution of resistance to gaseous air
46 pollutants. In: Taylor, G. E., Jr.; Pitelka, L. F.; Clegg, M. T., eds. *Ecological genetics and air pollution*.
47 New York, NY: Springer-Verlag; pp. 137-165.
- 48 Bartholomay, G. A.; Eckert, R. T.; Smith, K. T. (1997) Reductions in tree-ring widths of white pine following ozone
49 exposure at Acadia National Park, Maine, USA. *Can. J. For. Res.* 27: 361-368.
- 50 Batts, G. R.; Morison, J. I. L.; Ellis, R. H.; Hadley, P.; Wheeler, T. R. (1997) Effects of CO₂ and temperature on
51 growth and yield of crops of winter wheat over four seasons. *Eur. J. Agron.* 7: 43-52.
- 52 Bauer, M. R.; Hultman, N. E.; Panek, J. A.; Goldstein, A. H. (2000) Ozone deposition to a ponderosa pine plantation
53 in the Sierra Nevada Mountains (CA): a comparison of two different climatic years. *J. Geophys. Res. [Atmos.]*
54 105: 22,123-22,136.

- 1 Baumbusch, L. O.; Eiblmeier, M.; Schnitzler, J.-P.; Heller, W.; Sandermann, H., Jr.; Polle, A. (1998) Interactive
2 effects of ozone and low UV-B radiation on antioxidants in spruce (*Picea abies*) and pine (*Pinus sylvestris*).
3 *Physiol. Plant.* 104: 248-254.
- 4 Baumgardner, R. E.; Edgerton, E. S. (1998) Rural ozone across the eastern United States: analysis of CASTNet data,
5 1988-1995. *J. Air Waste Manage. Assoc.* 48: 674-688.
- 6 Baumgarten, M.; Werner, H.; Häberle, K.-H.; Emberson, L. D.; Fabian, P.; Matyssek, R. (2000) Seasonal ozone
7 response of mature beech trees (*Fagus sylvatica*) at high altitude in the Bavarian Forest (Germany) in
8 comparison with young beech grown in the field and in phytotrons. *Environ. Pollut.* 109: 431-442.
- 9 Beare, J. A.; Archer, S. A.; Bell, J. N. B. (1999a) Effects of *Melampsora* leaf rust disease and chronic ozone exposure
10 on poplar. *Environ. Pollut.* 105: 419-426.
- 11 Beare, J. A.; Archer, S. A.; Bell, J. N. B. (1999b) *Marssonina* leafspot disease of poplar under elevated ozone:
12 pre-fumigated host and in vitro studies. *Environ. Pollut.* 105: 409-417.
- 13 Beligni, M. V.; Lamattina, L. (2001) Nitric oxide in plants: the history is just beginning. *Plant Cell Environ.*
14 24: 267-278.
- 15 Beligni, M.; Lamattina, L. (2002) Nitric oxide interferes with plant photo-oxidative stress by detoxifying reactive
16 oxygen species. *Plant Cell Environ.* 25: 737-748.
- 17 Bell, J. N. B.; Treshow, M., eds. (2002) Air pollution and plant life. 2nd ed. Chichester, United Kingdom: John Wiley
18 & Sons Ltd.
- 19 Bell, J. N. B.; Ashmore, M. R.; Wilson, G. B. (1991) Ecological genetics and chemical modifications of the
20 atmosphere. In: Taylor, G. E., Jr.; Pitelka, L. F.; Clegg, M. T., eds. *Ecological genetics and air pollution*.
21 New York, NY: Springer-Verlag; pp. 33-59.
- 22 Bender, J.; Weigel, H. J.; Wegner, U.; Jäger, H. J. (1994) Response of cellular antioxidants to ozone in wheat flag
23 leaves at different stages of plant development. *Environ. Pollut.* 84: 15-21.
- 24 Bender, J.; Herstein, U.; Black, C. R. (1999) Growth and yield responses of spring wheat to increasing carbon
25 dioxide, ozone and physiological stresses: a statistical analysis of 'ESPACE-wheat' results. *Eur. J. Agron.*
26 10: 185-195.
- 27 Benes, S. E.; Murphy, T. M.; Anderson, P. D.; Houpis, J. L. J. (1995) Relationship of antioxidant enzymes to ozone
28 tolerance in branches of mature ponderosa pine (*Pinus ponderosa*) trees exposed to long term, low
29 concentration, ozone fumigation and acid precipitation. *Physiol. Plant.* 94: 124-134.
- 30 Bennett, J. H. (1979) Foliar exchange of gases. In: Heck, W. W.; Krupa, S. V.; Linzon, S. N., eds. *Methodology for*
31 *the assessment of air pollution effects on vegetation: a handbook from a specialty conference; April 1978;*
32 *Minneapolis, MN. Pittsburgh, PA: Air Pollution Control Association; pp. 10-1 - 10-29.*
- 33 Bennett, J. P.; Anderson, R. L.; Mielke, M. L.; Ebersole, J. J. (1994) Foliar injury air pollution surveys of eastern
34 white-pine (*Pinus-strobus* L.) - a review. *Environ. Monit. Assess.* 30: 247-274.
- 35 Benoit, L. F.; Skelly, J. M.; Moore, L. D.; Dochinger, L. S. (1982) Radial growth reductions of *Pinus strobus* L.
36 correlated with foliar ozone sensitivity as an indicator of ozone-induced losses in eastern forests. *Can. J. For.*
37 *Res.* 12: 673-678.
- 38 Benoit, L. F.; Skelly, J. M.; Moore, L. D.; Dochinger, L. S. (1983) The influence of ozone on *Pinus strobus* L. pollen
39 germination. *Can. J. For. Res.* 13: 184-187.
- 40 Benton, J.; Fuhrer, J.; Gimeno, B. S.; Skärby, L.; Sanders, G. (1995) Results from the UN/ECE ICP-crops indicate the
41 extent of exceedance of the critical levels of ozone in Europe. *Water Air Soil Pollut.* 85: 1473-1478.
- 42 Benton, J.; Fuhrer, J.; Gimeno, B. S.; Skärby, L.; Palmer-Brown, D.; Ball, G.; Roadknight, C.; Sanders-Mills, G.
43 (1996) The critical level of ozone for visible injury on crops and natural vegetation (ICP-Crops). In:
44 Kärenlampi, L.; Skärby, L., eds. *Critical levels for ozone in Europe: testing and finalizing the concepts:*
45 *UN-ECE workshop report. UN-ECE Convention on Long-Range Transboundary Air Pollution; April; Kuopio,*
46 *Finland. Kuopio, Finland: University of Kuopio, Department of Ecology and Environmental Science;*
47 *pp. 44-57.*
- 48 Benton, J.; Fuhrer, J.; Gimeno, B. S.; Skärby, L.; Palmer-Brown, D.; Ball, G.; Roadknight, C.; Mills, G. (2000)
49 An international cooperative programme indicates the widespread occurrence of ozone injury on crops.
50 *Agric. Ecosyst. Environ.* 78: 19-30.
- 51 Berg, B.; Staaf, H. (1980) Decomposition rate and chemical changes of Scots pine needle litter. II. Influence of
52 chemical composition. *Ecol. Bull.* 32: 373-390.
- 53 Bergmann, E.; Bender, J.; Weigel, H.-J. (1999) Ozone threshold doses and exposure-response relationships for the
54 development of ozone injury symptoms in wild plant species. *New Phytol.* 144: 423-435.
- 55 Bergstrom, J. C. (1990) Concepts and measures of the economic value of environmental quality: a review. *J. Environ.*
56 *Manage.* 31: 215-228.

- 1 Bergweiler, C. J.; Manning, W. J. (1999) Inhibition of flowering and reproductive success in spreading dogbane
2 (*Apocynum androsaemifolium*) by exposure to ambient ozone. *Environ. Pollut.* 105: 333-339.
- 3 Bermejo, V.; Gimeno, B. S.; Granados, I.; Santamaria, J.; Irigoyen, J. J.; Bermejo, R.; Porcuna, J. L.; Mills, G. (2002)
4 Investigating indices to explain the impacts of ozone on the biomass of white clover (*Trifolium repens* L. cv.
5 Regal) at inland and coastal sites in Spain. *New Phytol.* 156: 43-55.
- 6 Berrang, P.; Karnosky, D. F.; Mickler, R. A.; Bennett, J. P. (1986) Natural selection for ozone tolerance in *Populus*
7 *tremuloides*. *Can. J. For. Res.* 16: 1214-1216.
- 8 Berrang, P.; Karnosky, D. F.; Bennett, J. P. (1989) Natural selection for ozone tolerance in *Populus tremuloides*: field
9 verification. *Can. J. For. Res.* 19: 519-522.
- 10 Bertrand, A.; Robitaille, G.; Nadeau, P.; Castonguay, Y. (1999) Influence of ozone on cold acclimation in sugar
11 maple seedlings. *Tree Physiol.* 19: 527-534.
- 12 Beyers, J. L.; Riechers, G. H.; Temple, P. J. (1992) Effects of long-term ozone exposure and drought on the
13 photosynthetic capacity of ponderosa pine (*Pinus ponderosa* Laws.). *New Phytol.* 122: 81-90.
- 14 Bichele, I.; Moldau, H.; Padu, E. (2000) Estimation of plasmalemma conductivity to ascorbic acid in intact leaves
15 exposed to ozone. *Physiol. Plant.* 108: 405-412.
- 16 Bielenberg, D. G.; Lynch, J. P.; Pell, E. J. (2001) A decline in nitrogen availability affects plant responses to ozone.
17 *New Phytol.* 151: 413-425.
- 18 Binkley, D. (1992) Sensitivity of forest soils in the western U.S. to acidic deposition. In: Olson, R. K.; Binkley, D.;
19 Böhm, M., eds. *The Response of Western Forests to Air Pollution*. New York, NY: Springer-Verlag;
20 pp. 153-182.
- 21 Birecka, H.; Catalfamo, J.; Urban, P. (1976) Cell isoperoxidases in sweet potato plants in relation to mechanical
22 injury and ethylene. *Plant Physiol.* 57: 74-79.
- 23 Black, V. J.; Black, C. R.; Roberts, J. A.; Stewart, C. A. (2000) Impact of ozone on the reproductive development of
24 plants. *New Phytol.* 147: 421-447.
- 25 Blankenship, E. E.; Stefanski, L. A. (2001) Statistical estimation of ozone exposure metrics. *Atmos. Environ.*
26 35: 4499-4510.
- 27 Boerner, R. E. J.; Rebbeck, J. (1995) Decomposition and nitrogen release from leaves of three hardwood species
28 grown under elevated O₃ and/or CO₂. *Plant Soil* 170: 149-157.
- 29 Böhm, M.; McCune, B.; Vandetta, T. (1995) Ozone regimes in or near forests of the western United States: I. regional
30 patterns. *J. Air Waste Manage. Assoc.* 45: 235-246.
- 31 Bonello, P.; Heller, W.; Sandermann, H., Jr. (1993) Ozone effects on root-disease susceptibility and defence
32 responses in mycorrhizal and non-mycorrhizal seedlings of Scots pine (*Pinus sylvestris* L.). *New Phytol.*
33 124: 653-663.
- 34 Booker, F. L. (2000) Influence of carbon dioxide enrichment, ozone and nitrogen fertilization on cotton
35 (*Gossypium hirsutum* L.) leaf and root composition. *Plant Cell Environ.* 23: 573-583.
- 36 Booker, F. L.; Reid, C. D.; Brunschön-Harti, S.; Fiscus, E. L.; Miller, J. E. (1997) Photosynthesis and
37 photorespiration in soybean [*Glycine max* (L.) Merr.] chronically exposed to elevated carbon dioxide and
38 ozone. *J. Exp. Bot.* 48: 1843-1852.
- 39 Bortier, K.; Ceulemans, R.; De Temmerman, L. (2000) Effects of tropospheric ozone on woody plants. In: Agrawal,
40 S. B.; Agrawal, M., eds. *Environmental pollution and plant responses*; pp. 153-182. Boca Raton, FL:
41 Lewis Publishers.
- 42 Bortier, K.; Ceulemans, R.; De Temmerman, L. (2000a) Effects of ozone exposure on growth and photosynthesis of
43 beech seedlings (*Fagus sylvatica*). *New Phytol.* 146: 271-280.
- 44 Bortier, K.; DeTemmerman, L.; Ceulemans, R. (2000b) Effects of ozone exposure in open-top chambers on poplar
45 (*Populus nigra*) and beech (*Fagus sylvatica*): a comparison. *Environ. Pollut.* 109: 509-516.
- 46 Bortier, K.; Vandermeiren, K.; De Temmerman, L.; Ceulemans, R. (2001a) Growth, photosynthesis and ozone uptake
47 of young beech (*Fagus sylvatica* L.) in response to different ozone exposures. *Trees-Struct. Funct.* 15: 75-82
- 48 Bortier, K.; Dekelver, G.; De Temmerman, L.; Ceulemans, R. (2001b) Stem injection of *Populus nigra* with EDU to
49 study ozone effects under field conditions. *Environ. Pollut.* 111: 199-208.
- 50 Bosac, C.; Black, V. J.; Roberts, J. A.; Black, C. R. (1998) The impact of ozone on seed yield and quality and
51 seedling vigour in oilseed rape (*Brassica napus* L.). *J. Plant Physiol.* 153: 127-134.
- 52 Bosley, A.; Peterson, R.; Rebbeck, J. (1998) The resistance of the moss *Polytrichum commune* to acute exposure of
53 simulated acid rain or ozone compared to two fern species: spore germination. *Bryologist* 101: 512-518.
- 54 Bowler, C.; Van Montagu, M.; Inzé, D. (1992) Superoxide dismutase and stress tolerance. *Ann. Rev. Plant Physiol.*
55 *Mol. Biol.* 43: 83-116.

- 1 Bradshaw, A. D.; McNeilly, T. (1991) Evolution in relation to environmental stress. In: Taylor, G. E., Jr.; Pitelka,
2 L. F.; Clegg, M. T., eds. Ecological genetics and air pollution. New York, NY: Springer-Verlag; pp. 11-31.
- 3 Braun, S.; Flückiger, W. (1995) Effects of ambient ozone on seedlings of *Fagus sylvatica* L. and *Picea abies* (L.)
4 Karst. New Phytol. 129: 33-44.
- 5 Braun, S.; Rihm, B.; Schindler, C.; Flückiger, W. (1999) Growth of mature beech in relation to ozone and nitrogen
6 deposition: an epidemiological approach. Water Air Soil Pollut. 116: 357-364.
- 7 Brennan, E. G.; Leone, I. A.; Clarke, B. B. (1987) EDU: a chemical for evaluating ozone foliar injury and yield
8 reduction in field-grown crops. Int. J. Trop. Plant Dis. 5: 35-42.
- 9 Bringmark, E.; Bringmark, L. (1995) Disappearance of spatial variability and structure in forest floors - a distinct
10 effect of air pollution? Water Air Soil Pollut. 85: 761-766.
- 11 Broadmeadow, M. (1998) Ozone and forest trees. New Phytol. 139: 123-125.
- 12 Broadmeadow, M. S. J.; Heath, J.; Randle, T. J. (1999) Environmental limitations to O₃ uptake - some key results
13 from young trees growing at elevated CO₂ concentrations. Water Air Soil Pollut. 116: 299-310.
- 14 Broadmeadow, M. S. J.; Jackson, S. B. (2000) Growth responses of *Quercus petraea*, *Fraxinus excelsior* and *Pinus*
15 *sylvestris* to elevated carbon dioxide, ozone and water supply. New Phytol. 146: 437-451.
- 16 Brook, J. R.; DiGiovanni, F.; Cakmak, S.; Meyers, T. P. (1997) Estimation of dry deposition velocity using inferential
17 models and site-specific meteorology—uncertainty due to siting of meteorological towers. Atmos. Environ.
18 31: 3911-3919.
- 19 Brown, V. C. (1995) Insect herbivores and gaseous air pollutants - current knowledge and predictions. In:
20 Harrington, R.; Stork, N. E., eds. Insects in a changing environment: 17th symposium of the Royal
21 Entomological Society. New York, NY: Academic Press; pp. 219-249.
- 22 Brown, K. A.; Roberts, T. M. (1988) Effects of ozone on foliar leaching in Norway spruce (*Picea abies* L. Karst):
23 confounding factors due to NO_x production during ozone generation. Environ. Pollut. 55: 55-73.
- 24 Brown, V. C.; Ashmore, M. R.; McNeill, S. (1993) Experimental investigations of the effects of air pollutants on
25 aphids on coniferous trees. Forstwiss. Centralbl. 112: 129-132.
- 26 Brunschön-Harti, S.; Fangmeier, A.; Jäger, H.-J. (1995) Influence of ozone and ethylenediurea (EDU) on growth and
27 yield of bean (*Phaseolus vulgaris* L.) in open-top field chambers. Environ. Pollut. 90: 89-94.
- 28 Buchanan, B. B.; Gruissem, W.; Jones, R. L., eds. (2000) Biochemistry and molecular biology of plants. Rockville,
29 MD: American Society of Plant Physiologists.
- 30 Bungener, P.; Balls, G. R.; Nussbaum, S.; Geissmann, M.; Grub, A.; Fuhrer, J. (1999a) Leaf injury characteristics of
31 grassland species exposed to ozone in relation to soil moisture condition and vapour pressure deficit.
32 New Phytol. 142: 271-282.
- 33 Bungener, P.; Nussbaum, S.; Grub, A.; Fuhrer, J. (1999b) Growth response of grassland species to ozone in relation
34 to soil moisture and plant strategy. New Phytol. 142: 283-293.
- 35 Burkey, K. O. (1999) Effects of ozone on apoplast/cytoplasm partitioning of ascorbic acid in snap bean. Physiol.
36 Plant. 107: 188-193.
- 37 Bussotti, F.; Ferretti, M. (1998) Air pollution, forest condition and forest decline in Southern Europe. Environ. Pollut.
38 101: 49-65.
- 39 Bytnerowicz, A. (2002) Physiological/ecological interactions between ozone and nitrogen deposition in forest
40 ecosystems. Phytol. 42: 13-28.
- 41 Bytnerowicz, A.; Turunen, M. (1994) Effects of ozone exposures on epicuticular wax of ponderosa pine needles. In:
42 Percy, K. E.; Cape, J. N.; Jagels, R.; Simpson, C. J., eds. Air Pollutants and the Leaf Cuticle. New York, NY:
43 Springer-Verlag; pp. 305-313. (NATO ASI Series; v. G 36).
- 44 Bytnerowicz, A.; Olszyk, D. M.; Fox, C. A.; Dawson, P. J.; Kats, G.; Morrison, C. L.; Wolf, J. (1988) Responses of
45 desert annual plants to ozone and water stress in an *in situ* experiment. JAPCA 38: 1145-1151.
- 46 Bytnerowicz, A.; Olszyk, D. M.; Huttunen, S.; Takemoto, B. K. (1989) Effects of photochemical smog on growth and
47 injury and gas exchange of pine seedlings. Can. J. Bot. 67: 2175-2181.
- 48 Bytnerowicz, A.; Poth, M.; Takemoto, B. K. (1990) Effects of photochemical smog and mineral nutrition on
49 ponderosa pine seedlings. Environ. Pollut. 67: 233-248.
- 50 Bytnerowicz, A.; Fenn, M. E.; Miller, P. R.; Arbaugh, M. J. (1999) Wet and dry pollutant deposition to the mixed
51 conifer forest. In: Miller, P. R.; McBride, J. R., eds. Oxidant Air Pollution Impacts in the Montane Forests of
52 Southern California: a Case Study of the San Bernardino Mountains. New York, NY: Springer; pp. 235-269.
53 (Ecological Series, no. 134).
54

- Bytnerowicz, A.; Godzik, B.; Frączek, W.; Grodzińska, K.; Krywult, M.; Badea, O.; Barančok, P.; Blum, O.; Černý, M.; Godzik, S.; Mankovska, B.; Manning, W.; Moravčík, P.; Musselman, R.; Oszlanyi, J.; Postelnicu, D.; Szdźuj, J.; Varšavova, M.; Zota, M. (2002) Distribution of ozone and other air pollutants in forests of the Carpathian Mountains in central Europe. *Environ. Pollut.* 116: 3-25.
- Bytnerowicz, A.; Tausz, M.; Alonso, R.; Jones, D.; Johnson, R.; Grulke, N. (2002) Summer-time distribution of air pollutants in Sequoia National Park, California. *Environ. Pollut.* 118: 187-203.
- Cairnes, M. S.; Lackey, R. T. (1992) Biodiversity and management of natural resources: the issues. *Fisheries* 17: 6-10.
- Callaway, J. M.; Darwin, R. F.; Nesse, R. J. (1986) Economic valuation of acidic deposition damages: preliminary results from the 1985 NAPAP assessment. In: *Proceedings of International Symposium on Acidic Precipitation*; September; Muskoka, Ontario, Canada. Richland, WA: U. S. Environmental Protection Agency; Pacific Northwest Laboratory. *Water Air Soil Pollut.* 31: 1019-1034.
- Campbell, S.; Temple, P.; Pronos, J.; Rochefort, R.; Andersen, C. (2000) Monitoring for ozone injury in west coast (Oregon, Washington, California) forests in 1998. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station; general technical report no. PNW-GTR-495. Available: <http://www.fs.fed.us/pnw/gtrs.htm> [11 April, 2003].
- Cao, W.; Tibbitts, T. W.; Wheeler, R. M. (1994) Carbon dioxide interactions with irradiance and temperature in potatoes. In: MacElroy, R. D. *Life sciences and space research XXV(3): natural and artificial ecosystems: proceedings of the topical meeting of the COSPAR Interdisciplinary Scientific Commission F (Meeting F4) of the COSPAR twenty-ninth plenary meeting*; August-September, 1992; Washington, DC. New York, NY: Committee on Space Research; pp. 243-250. (*Advances in space research*: v. 14, no. 11).
- Cape, J. N. (1996) Surface wetness and pollutant deposition. In: Kerstiens, G., ed. *Plant cuticles: an integrated functional approach*. Oxford, United Kingdom: BIOS Scientific Publishers; pp. 283-300. (*Environmental plant biology series*).
- Cardoso-Vilhena, J.; Barnes, J. (2001) Does nitrogen supply affect the response of wheat (*Triticum aestivum* cv. Hanno) to the combination of elevated CO₂ and O₃? *J. Exp. Bot.* 52: 1901-1911.
- Cardoso-Vilhena, J.; Balaguer, L.; Daymond, J.; Barnes, J. D. (1998) Amelioration of ozone damage by elevated CO₂. In: DeKok, L. J.; Stulen, I., eds. *Responses of plant metabolism to air pollution and global change*. Leiden, The Netherlands: Backhuys Publishers; pp. 281-282.
- Carey, W. A.; Kelley, W. D. (1994) Interaction of ozone exposure and *Fusarium subglutinans* inoculation on growth and disease development of loblolly pine seedlings. *Environ. Pollut.* 84: 35-43.
- Carey, W. A.; Kelley, W. D. (1994) Effects of ozone exposure and *Fusarium subglutinans* inoculation on growth and disease development of loblolly pine seedlings. *Environ. Pollut.* 84: 35-43.
- Carnahan, J. E.; Jenner, E. L.; Wat, E. K. W. (1978) Prevention of ozone injury to plants by a new protectant chemical. *Phytopathology* 68: 1225-1229.
- Carpenter, R. A.; Dixon, J. A. (1985) Ecology meets economics: a guide to sustainable development. *Environment* 27: 6-32.
- Carroll, J. J.; Miller, P. R.; Pronos, J. (2003) Historical perspectives on ambient ozone and its effects on the Sierra Nevada. In: Bytnerowicz, A.; Arbaugh, M. J.; Alonso, R., eds. *Ozone air pollution in the Sierra Nevada: distribution and effects on forests*; pp. 33-54. New York, NY: Elsevier. (*Developments in Environmental Science*: v. 2).
- Castillo, F. J.; Greppin, H. (1986) Balance between anionic and cationic extracellular peroxidase activities in *Sedum album* leaves after ozone exposure. Analysis by high-performance liquid chromatography. *Physiol. Plant.* 68: 201-208.
- Castillo, F. J.; Heath, R. L. (1990) CA²⁺ transport in membrane vesicles from pinto bean leaves and its alteration after ozone exposure. *Plant Physiol.* 94: 788-795.
- Castillo, F. J.; Penel, C.; Greppin, H. (1984) Peroxidase release induced by ozone in *Sedum album* leaves: involvement of Ca²⁺. *Plant Physiol.* 74: 846-851.
- Castillo, F. J.; Miller, P. R.; Greppin, H. (1987) Extracellular biochemical markers of photochemical oxidant air pollution damage to Norway spruce. *Experientia (Basel)* 43: 111-115.
- Castle, E. N. (1993) A pluralistic, pragmatic and evolutionary approach to natural resource management. *For. Ecol. Manage.* 56: 279-295.
- Cazalé, A.-C.; Rouet-Mayer, M.-A.; Barbier-Brygoo, H.; Mathieu, Y.; Laurière, C. (1998) Oxidative burst and hypoosmotic stress in tobacco cell suspensions. *Plant Physiol.* 116: 659-669.
- Chappelka, A. H. (2002) Reproductive development of blackberry (*Rubus cuneifolius*) as influenced by ozone. *New Phytol.* 155: 249-255.

- 1 Chappelka, A. H., III; Chevone, B. I. (1986) White ash seedling growth response to ozone and simulated acid rain.
2 Can. J. For. Res. 16: 786-790.
- 3 Chappelka, A. H.; Freer-Smith, P. H. (1995) Predisposition of trees by air pollutants to low temperatures and moisture
4 stress. Environ. Pollut. 87: 105-117.
- 5 Chappelka, A. H., III; Chevone, B. I.; Burk, T. E. (1985) Growth response of yellow-poplar (*Liriodendron tulipifera*
6 L.) seedlings to ozone, sulfur dioxide, and simulated acidic precipitation, alone and in combination. Environ.
7 Exp. Bot. 25: 233-244.
- 8 Chappelka, A.; Renfro, J.; Somers, G.; Nash, B. (1997) Evaluation of ozone injury on foliage of black cherry
9 (*Prunus serotina*) and tall milkweed (*Asclepias exaltata*) in Great Smoky Mountains National Park. Environ.
10 Pollut. 95: 13-18.
- 11 Chappelka, A. H.; Samuelson, L. J. (1998) Ambient ozone effects on forest trees of the eastern United States:
12 a review. New Phytol. 139: 91-108.
- 13 Chappelka, A.; Skelly, J.; Somers, G.; Renfro, J.; Hildebrand, E. (1999a) Mature black cherry used as a bioindicator
14 of ozone injury. Water Air Soil Pollut. 116: 261-266.
- 15 Chappelka, A.; Somers, G.; Renfro, J. (1999b) Visible ozone injury on forest trees in Great Smoky Mountains
16 National Park, USA. Water Air Soil Pollut. 116: 255-260.
- 17 Chappelka, A. H.; Neufeld, H. S.; Davison, A. W.; Somers, G. L.; Renfro, J. R. (2003) Ozone injury on cutleaf
18 coneflower (*Rudbeckia laciniata*) and crown-beard (*Verbesina occidentalis*) in Great Smoky Mountains
19 National Park. Environ. Pollut. 125: 53-60.
- 20 Chen, C. W.; Tsai, W. T.; Gomez, L. E. (1994) Modeling responses of Ponderosa pine to interacting stresses of ozone
21 and drought. For. Sci. 40: 267-288.
- 22 Chen, C. W.; Tsai, W. T.; Lucier, A. A. (1998) A model of air-tree-soil system for ozone impact analysis. Ecol.
23 Modell. 111: 207-222.
- 24 Cheong, Y. H.; Chang, H.-S.; Gupta, R.; Wang, X.; Zhu, T.; Luan, S. (2002) Transcriptional profiling reveals novel
25 interactions between wounding, pathogen, abiotic stress, and hormonal responses in Arabidopsis. Plant
26 Physiol. 129: 661-677.
- 27 Chernikova, T.; Robinson, J. M.; Lee, E. H.; Mulchi, C. L. (2000) Ozone tolerance and antioxidant enzyme activity in
28 soybean cultivars. Photosynth. Res. 64: 15-26.
- 29 Chevrier, N.; Chung, Y. S.; Sarhan, F. (1990) Oxidative damages and repair in *Euglena gracilis* exposed to ozone: II.
30 membrane permeability and uptake of metabolites. Plant Cell Physiol. 31: 987-992.
- 31 Clark, C. S.; Weber, J. A.; Lee, E. H.; Hogsett, W. E. (1995) Accentuation of gas exchange gradients in flushes of
32 ponderosa pine exposed to ozone. Tree Physiol. 15: 181-189.
- 33 Clarke, B. B.; Henninger, M. R.; Brennan, E. (1983) An assessment of potato losses caused by oxidant air pollution in
34 New Jersey. Phytopathology 73: 104-108.
- 35 Cobb, F. W., Jr.; Wood, D. L.; Stark, R. W.; Parmeter, J. R., Jr. (1968) Photochemical oxidant injury and bark beetle
36 (Coleoptera: Scolytidae) infestation of ponderosa pine. IV. Theory on the relationships between oxidant injury
37 and bark beetle infestation. Hilgardia 39: 141-152.
- 38 Coe, H.; Gallagher, M. W.; Choularton, T. W.; Dore, C. (1995) Canopy scale measurements of stomatal and cuticular
39 O₃ uptake by Sitka spruce. Atmos. Environ. 29: 1413-1423.
- 40 Coleman, J. S.; Jones, C. G.; Smith, W. H. (1987) The effect of ozone on cottonwood - leaf rust interactions:
41 independence of abiotic stress, genotype, and leaf ontogeny. Can. J. Bot. 65: 949-953.
- 42 Coleman, M. D.; Dickson, R. E.; Isebrands, J. G.; Karnosky, D. F. (1995a) Carbon allocation and partitioning in
43 aspen clones varying in sensitivity to tropospheric ozone. Tree Physiol. 15: 593-604.
- 44 Coleman, M. D.; Isebrands, J. G.; Dickson, R. E.; Karnosky, D. F. (1995b) Photosynthetic productivity of aspen
45 clones varying in sensitivity to tropospheric ozone. Tree Physiol. 15: 585-592.
- 46 Coleman, M. D.; Dickson, R. E.; Isebrands, J. G.; Karnosky, D. F. (1996) Root growth physiology of potted and
47 field-grown trembling aspen exposed to tropospheric ozone. Tree Physiol. 16: 145-152.
- 48 Colls, J. J.; Unsworth, M. H. (1992) Air pollution interactions with natural stressors. In: Barker, J. R.; Tingey, D. T.,
49 eds. Air pollution effects on biodiversity. New York, NY: Van Nostrand Reinhold; pp. 93-108.
- 50 Common, M.; Perrings, C. (1992) Towards an ecological economics of sustainability. Ecol. Econ. 6: 7-34.
- 51 Constable, J. V. H.; Friend, A. L. (2000) Suitability of process-based tree growth models for addressing tree response
52 to climate change. Environ. Pollut. 110: 47-59.
- 53 Constable, J. V. H.; Taylor, G. E., Jr. (1997) Modeling the effects of elevated tropospheric O₃ on two varieties of
54 *Pinus ponderosa*. Can. J. For. Res. 27: 527-537.
- 55 Constable, J. V. H.; Taylor, G. E., Jr.; Laurence, J. A.; Weber, J. A. (1996) Climatic change effects on the physiology
56 and growth of *Pinus ponderosa*: expectations from simulation modeling. Can. J. For. Res. 26: 1315-1325.

- 1 Cook, E. (1990) A conceptual linear aggregate model for tree rings. In: Cook, E. R.; Kairiukstis, L. A., eds. Methods
2 of Dendrochronology: applications in the environmental sciences. Dordrecht, The Netherlands: Kluwer
3 Academic Publishers, pp. 98-104.
- 4 Cooley, D. R.; Manning, W. J. (1987) The impact of ozone on assimilate partitioning in plants: a review. Environ.
5 Pollut. 47: 95-113.
- 6 Coruzzi, G. M.; Zhou, L. (2001) Carbon and nitrogen sensing and signaling in plants: emerging 'matrix effects'. Curr.
7 Opin. Plant Biol. 4: 247-253.
- 8 Costa, S. D.; Kennedy, G. G.; Heagle, A. S. (2001) Effect of host plant ozone stress on Colorado potato beetles.
9 Environ. Entomol. 30: 824-831.
- 10 Costanza, R.; d'Arge, R.; De Groot, R.; Farber, S.; Grasso, M.; Hannon, B.; Limburg, K.; Naeem, S.; O'Neill, R. V.;
11 Paruelo, J.; Raskin, R. G.; Sutton, P.; Van Den Belt, M. (1997) The value of the world's ecosystem services
12 and natural capital. Nature (London) 387: 253-259.
- 13
- 14 Coulston, J. W.; Smith, G. C.; Smith, W. D. (2003) Regional assessment of ozone sensitive tree species using
15 bioindicator plants. Environ. Monit. Assess. 83: 113-127.
- 16 Cox, R. M.; Malcolm, J. W. (1999) Passive ozone monitoring for forest health assessment. Water Air Soil Pollut. 116:
17 339-344.
- 18 Craigon, J.; Fangmeier, A.; Jones, M.; Donnelly, A.; Bindi, M.; De Temmerman, L.; Persson, K.; Ojanpera, K. (2002)
19 Growth and marketable-yield responses of potato to increased CO₂ and ozone. Eur. J. Agron. 17: 273-289.
- 20 Crocker, T. D. (1985) On the value of the condition of a forest stock. Land Econ. 61: 244-254.
- 21 Crocker, T. D.; Forster, B. A. (1986) Atmospheric deposition and forest decline. Water Air Soil Pollut. 31:
22 1007-1017.
- 23 Curtis, P. S. (1996) A meta-analysis of leaf gas exchange and nitrogen in trees grown under elevated carbon dioxide.
24 Plant Cell Environ. 19: 127-137.
- 25 Dahlsten, D. L.; Rowney, D. L.; Kickert, R. N. (1997) Effects of oxidant air pollutants on western pine beetle
26 (Coleoptera: Scolytidae) populations in southern California. Environ. Pollut. 96: 415-423.
- 27 Daily, G. C. (1999) Developing a scientific basis for managing Earth's life support systems. Conserv. Ecol. 3: 14.
28 Available: <http://www.ecologyandsociety.org/vol3/iss2/art14/> [21 April, 2004].
- 29 Daily, G. C. (2000) Ecological assessment (untitled article in this section). Issues Sci. Technol. 16: 6.
- 30 Dämmgen, U.; Grünhage, L.; Haenel, H.-D.; Jäger, H.-J. (1993) Climate and stress in ecotoxicology. A coherent
31 system of definitions and terms. Angew. Bot. 67: 157-162.
- 32 Danielsson, H.; Gelang, J.; Pleijel, H. (1999) Ozone sensitivity, growth and flower development in *Phleum* genotypes
33 of different geographic origin in the Nordic countries. Environ. Exp. Bot. 42: 41-49.
- 34 Danielsson, H.; Karlsson, G. P.; Karlsson, P. E.; Pleijel, J. (2003) Ozone uptake modelling and flux-response
35 relationships—an assessment of ozone-induced yield loss in spring wheat. Atmos. Environ. 37: 475-485.
- 36 Dass, H. C.; Weaver, G. M. (1968) Modification of ozone damage to *phaseolus vulgaris* by antioxidants, thiols and
37 sulfhydryl reagents. Can. J. Plant Sci. 48: 569-574.
- 38 Dass, H. C.; Weaver, G. M. (1972) Enzymatic changes in intact leaves of *Phaseolus vulgaris* following ozone
39 fumigation. Atmos. Environ. 6: 759-763.
- 40 Davidson, A. (1993) Update of ozone trends in California's South Coast Air Basin. Air Waste 43: 226-227.
- 41 Davis, D. D.; Skelly, J. M. (1992) Foliar sensitivity of eight eastern hardwood tree species to ozone. Water Air Soil
42 Pollut. 62: 269-277.
- 43 Davison, A. W.; Barnes, J. D. (1998) Effects of ozone on wild plants. New Phytol. 139: 135-151.
- 44 Davison, A. W.; Reiling, K. (1995) A rapid change in ozone resistance of *Plantago major* after summers with high
45 ozone concentrations. New Phytol. 131: 337-344.
- 46 Davison, A. W.; Neufeld, H. S.; Chappelka, A. H.; Wolff, K. W.; Finkelstein, P. L. (2003) Interpreting spatial
47 variation in ozone symptoms shown by cutleaf cone flower, *Rudbeckia laciniata* L. Environ. Pollut. 125:
48 61-70.
- 49 De Kok, L. J.; Stulen, I. (1998) Responses of plant metabolism to air pollution and global change. Leiden, The
50 Netherlands: Backhuys Publishers.
- 51 De Santis, F. (1999) New directions: will a new European vegetation ozone standard be fair to all European
52 countries? Atmos. Environ. 33: 3873-3874.
- 53 De Santis, F. (2000) Reply to 'Use and abuse of the AOT40 concept' by M. Ashmore and J. Fuhrer [letter]. Atmos.
54 Environ. 34: 1158-1159.
- 55 De Temmerman, L.; Hacour, A.; Guns, M. (2002a) Changing climate and potential impacts on potato yield and
56 quality 'CHIP': introduction, aims and methodology. Eur. J. Agron. 17: 233-242.

- 1 De Temmerman, L.; Wolf, J.; Colls, J.; Bindi, M.; Fangmeier, A.; Finnan, J.; Ojanperä, K.; Pleijel, H. (2002b) Effect
2 of climatic conditions on tuber yield (*Solanum tuberosum* L.) in the European 'CHIP' experiments. Eur. J.
3 Agron. 17: 243-255.
- 4 DeLucia, E. H.; Hamilton, J. G.; Naidu, S. L.; Thomas, R. B.; Andrews, J. A.; Finzi, A.; Lavine, M.; Matamala, R.;
5 Mohan, J. E.; Hendrey, G. R.; Schlesinger, W. H. (1999) Net primary production of a forest ecosystem with
6 experimental CO₂ enrichment. Science (Washington, DC) 284: 1177-1179.
- 7 Degoussée, N.; Triantaphyllides, C.; Montillet, J.-L. (1994) Involvement of oxidative processes in the signaling
8 mechanisms leading to the activation of glyceollin synthesis in soybean (*Glycine max*). Plant Physiol. 104:
9 945-952.
- 10 Devlin, W. S.; Gustine, D. L. (1992) Involvement of the oxidative burst in phytoalexin accumulation and the
11 hypersensitive reaction. Plant Physiol. 100: 1189-1195.
- 12 Diaz, G.; Barrantes, O.; Honrubia, M.; Gracia, C. (1996) Effect of ozone and sulphur dioxide on mycorrhizae of
13 *Pinus halepensis* Miller. Ann. Sci. For. 53: 849-856.
- 14 Dickson, R. E.; Coleman, M. D.; Reimenschneider, D. E.; Isebrands, J. G.; Hogan, G. D.; Karnosky, D. F. (1998)
15 Growth of five hybrid poplar genotypes exposed to interacting elevated CO₂ and O₃. Can. J. For. Res. 28:
16 1706-1716.
- 17 Dickson, R. E.; Coleman, M. D.; Pechter, P.; Karnosky, D. (2001) Growth and crown architecture of two aspen
18 genotypes exposed to interacting ozone and carbon dioxide. Environ. Pollut. 115: 319-334.
- 19 Dietz, K.-J. (1997) Functions and responses of the leaf apoplast under stress. Prog. Bot. 58: 221-254.
- 20 Dighton, J.; Jansen, A. E. (1991) Atmospheric pollutants and ectomycorrhizae: more questions than answers? Environ.
21 Pollut. 73: 179-204.
- 22 Dijak, M.; Ormrod, D. P. (1982) Some physiological and anatomical characteristics associated with differential ozone
23 sensitivity among pea cultivars. Environ. Exp. Bot. 22: 395-402.
- 24 Dixon, B. L.; Segerson, K. (1999) Impacts of increased climate variability on the profitability of Midwest agriculture.
25 J. Agric. Appl. Econ. 31: 537-548.
- 26 Dixon, M.; Le Thiec, D.; Garrec, J. P. (1998) Reactions of Norway spruce and beech trees to 2 years of ozone
27 exposure and episodic drought. Environ. Exp. Bot. 40: 77-91.
- 28 Docherty, M.; Salt, D. T.; Holopainen, J. K. (1997) The impacts of climate change and pollution on forest pests. In:
29 Watt, A. D.; Stork, N. E.; Hunter, M. D., eds. Forests and insects. New York, NY: Chapman and Hall; pp.
30 229-247.
- 31 Dockery, D. W.; Schwartz, J.; Spengler, J. D. (1992) Air pollution and daily mortality: associations with particulates
32 and acid aerosols. Environ. Res. 59: 362-373.
- 33 Dominy, P. J.; Health, R. L. (1985) Inhibition of the K⁺-stimulated ATPase of the plasmalemma of pinto bean leaves
34 by ozone. Plant Physiol. 77: 43-45.
- 35 Donnelly, A.; Jones, M. B.; Schneiders, B. J.; Burke, J. I. (1998) The interactive effects of CO₂, O₃ and nitrogen on the
36 photosynthetic response of juvenile wheat plants. In: De Kok, L. J.; Stulen, I., eds. Responses of plant
37 metabolism to air pollution and global change. Leiden, The Netherlands: Backhuys Publishers; pp. 291-293.
- 38 Donnelly, A.; Jones, M. B.; Burke, J. I.; Schnieders, B. (1999) Does elevated CO₂ protect grain yield of wheat from the
39 effects of ozone stress? Z. Naturforsch. C: J. Biosci. 54: 802-811.
- 40 Donnelly, A.; Jones, M. B.; Burke, J. I.; Schneiders, B. (2000) Elevated CO₂ provides protection from O₃ induced
41 photosynthetic damage and chlorophyll loss in flag leaves of spring wheat (*Triticum aestivum* L., cv.'Minaret').
42 Agric. Ecosyst. Environ. 80: 159-168.
- 43 Donnelly, A.; Craigon, J.; Black, C. R.; Colls, J. J.; Landon, G. (2001a) Elevated CO₂ increases biomass and tuber
44 yield in potato even at high ozone concentrations. New Phytol. 149: 265-274.
- 45 Donnelly, A.; Craigon, J.; Black, C. R.; Colls, J. J.; Landon, G. (2001) Does elevated CO₂ ameliorate the impact of O₃
46 on the chlorophyll content and photosynthesis in potato (*Solanum tuberosum*)? Physiol. Plant. 111: 501-511.
- 47 Donnelly, A.; Lawson, T.; Craigon, J.; Black, C. R.; Colls, J. J.; Landon, G. (2001b) Effects of elevated CO₂ and O₃ on
48 tuber quality in potato (*Solanum tuberosum* L.). Agric. Ecosyst. Environ. 87: 273-285.
- 49 Dowding, P. (1988) Air pollutant effects on plant pathogens. In: Schulte-Hostede, S.; Darrall, N. M.; Blank, L. W.;
50 Wellburn, A. R., eds. Air pollution and plant metabolism. London, United Kingdom: Elsevier Applied
51 Scientific; pp. 329-355.
- 52 Drogoudi, P. D.; Ashmore, M. R. (2000) Does elevated ozone have differing effects in flowering and deblossomed
53 strawberry? New Phytol. 147: 561-569.
- 54 Drogoudi, P. D.; Ashmore, M. (2001) ¹⁴C-allocation of flowering and deblossomed strawberry in response to elevated
55 ozone. New Phytol. 152: 455-461.

- 1 Duchelle, S. F.; Skelly, J. M. (1981) Response of common milkweed to oxidant air pollution in the Shenandoah
2 National Park in Virginia. *Plant Dis.* 65: 661-663.
- 3 Duchelle, S. F.; Skelly, J. M.; Chevone, B. I. (1982) Oxidant effects on forest tree seedling growth in the Appalachian
4 mountains. *Water Air Soil Pollut.* 18: 363-373.
- 5 Duchelle, S. F.; Skelly, J. M.; Sharick, T. L.; Chevone, B. I.; Yang, Y.-S.; Nellessen, J. E. (1983) Effects of ozone on
6 the productivity of natural vegetation in a high meadow of the Shenandoah National Park of Virginia. *J.*
7 *Environ. Manage.* 17: 299-308.
- 8 Dueck, Th. A.; Zuin, A.; Elderson, J. (1998) Influence of ammonia and ozone on growth and drought sensitivity of
9 *Pinus sylvestris*. *Atmos. Environ.* 32: 545-550.
- 10 Dunn, D. B. (1959) Some effects of air pollution on *Lupinus* in the Los Angeles area. *Ecology* 40: 621-625.
- 11 Eamus, D. (1996) Responses of field grown trees to CO₂ enrichment. *Commonw. For. Rev.* 75: 39-47.
- 12 Eamus, D.; Murray, M. (1991) Photosynthetic and stomatal conductance responses of Norway spruce and beech to
13 ozone, acid mist and frost - a conceptual model. *Environ. Pollut.* 72: 23-44.
- 14 Eckey-Kaltenbach, H.; Ernst, D.; Heller, W.; Sandermann, H., Jr. (1994) Biochemical responses to ozone. IV.
15 Cross-induction of defensive pathways in parsley (*Petroselinum crispum* L.) plants. *Plant Physiol.* 104: 67-74.
- 16 Eckey-Kaltenbach, H.; Grosskopf, E.; Sandermann, H., Jr.; Ernst, D. (1994) Induction of pathogen defence genes in
17 parsley (*Petroselinum crispum* L.) plants by ozone. *Proc. R. Soc. Edinburgh Sect. B: Biol. Sci.* 102: 63-74.
- 18 Edwards, N. T. (1991) Root and soil respiration responses to ozone in *Pinus taeda* L. seedlings. *New Phytol.* 118:
19 315-321.
- 20 Edwards, G. S.; Kelly, J. M. (1992) Ectomycorrhizal colonization of loblolly pine seedlings during three growing
21 seasons in response to ozone, acidic precipitation, and soil Mg status. *Environ. Pollut.* 76: 71-77.
- 22 Edwards, G. S.; Wullschlegel, S. D.; Kelly, J. M. (1994) Growth and physiology of northern red oak: preliminary
23 comparisons of mature tree and seedling responses to ozone. *Environ. Pollut.* 83: 215-221.
- 24 Einig, W.; Lauxmann, U.; Hauch, B.; Hampp, R.; Landolt, W.; Maurer, S.; Matyssek, R. (1997) Ozone-induced
25 accumulation of carbohydrates changes enzyme activities of carbohydrate metabolism in birch leaves. *New*
26 *Phytol.* 137: 673-680.
- 27 Elkies, T.; Ormrod, D. P. (1979) Leaf diffusion resistance responses of three petunia cultivars to ozone and/or sulfur
28 dioxide. *J. Air Pollut. Control Assoc.* 29: 622-625.
- 29 Elkies, T.; Ormrod, D. P. (1981) Sorption of O₃, SO₂, NO₂ or their mixture by nine *Poa pratensis* cultivars of differing
30 pollutant sensitivity. *Atmos. Environ.* 15: 1739-1743.
- 31 Emberson, L. D.; Wieser, G.; Ashmore, M. R. (2000a) Modelling of stomatal conductance and ozone flux of Norway
32 spruce: comparison with field data. *Environ. Pollut.* 109: 393-402.
- 33 Emberson, L.; Ashmore, M. R.; Cambridge, H. M.; Simpson, D.; Tuovinen, J.-P. (2000b) Modelling stomatal ozone
34 flux across Europe. *Environ. Pollut.* 109: 403-413.
- 35 Enders, G. (1992) Deposition of ozone to a mature spruce forest: measurements and comparison to models. *Environ.*
36 *Pollut.* 75: 61-67.
- 37 Ennis, C. A.; Lazrus, A. L.; Kok, G. L.; Zimmerman, P. R.; Monson, R. K. (1990) A branch chamber system and
38 techniques for simultaneous pollutant exposure experiments and gaseous flux determinations. *Tellus Ser. B* 42:
39 170-182.
- 40 Ernst, D.; Schraudner, M.; Langebartels, C.; Sandermann, H. (1992) Ozone-induced changes of messenger RNA levels
41 of beta-1,3-glucanase, chitinase and pathogenesis-related protein 1B in tobacco plants. *Plant. Mol. Biol.* 20:
42 673-682.
- 43 Espelie, K. E.; Franceschi, V. R.; Kolattukudy, P. E. (1986) Immunocytochemical localization and time course of
44 appearance of an anionic peroxidase associated with suberization in wound-healing potato tuber tissue. *Plant*
45 *Physiol.* 81: 487-492.
- 46 Evans, P. A.; Ashmore, M. R. (1992) The effects of ambient air on a semi-natural grassland community. *Agric.*
47 *Ecosyst. Environ.* 38: 91-97.
- 48 Ewart, F.; Pleijel, H. (1999) Phenological development, leaf emergence, tillering and leaf area index, and duration of
49 spring wheat across Europe in response to CO₂ and ozone. *Eur. J. Agron.* 10: 171-184.
- 50 Ewart, F.; Van Oijen, M.; Porter, J. R. (1999) Simulation of growth and development processes of spring wheat in
51 response to CO₂ and ozone for different sites and years in Europe using mechanistic crop simulation models.
52 *Eur. J. Agron.* 10: 231-247.
- 53 Fangmeier, A.; Stein, W.; Jäger, H.-J. (1992) Advantages of an open-top chamber plant exposure system to assess the
54 impact of atmospheric trace gases on vegetation. *Angew. Bot.* 66: 97-105.

- 1 Fangmeier, A.; Bockerhoff, U.; Grütters, U.; Jäger, H.-J. (1994a) Growth and yield responses of spring wheat
2 (*Triticum aestivum* L. cv. Turbo) grown in open-top chambers to ozone and water stress. *Environ. Pollut.* 83:
3 317-325.
- 4 Fangmeier, A.; Brunschön, S.; Jäger, H.-J. (1994b) Time course of oxidant stress biomarkers in flag leaves of wheat
5 exposed to ozone and drought stress. *New Phytol.* 126: 63-69.
- 6 Fangmeier, A.; Grütters, U.; Vermehren, B.; Jäger, H.-J. (1996) Responses of some cereal cultivars to CO₂ enrichment
7 and tropospheric ozone at different levels of nitrogen supply. *J. Appl. Bot.* 70: 12-18.
- 8 Fangmeier, A.; De Temmerman, L.; Mortensen, L.; Kemp, K.; Burke, J.; Mitchell, R.; van Oijen, M.; Weigel, H. J.
9 (1999) Effects on nutrients and on grain quality in spring wheat crops grown under elevated CO₂ concentrations
10 and stress conditions in the European, multiple-site experiment 'ESPACE-wheat'. *Eur. J. Agron.* 10: 215-229.
- 11 Fangmeier, A.; Bender, J.; Weigel, H.-J.; Jäger, H.-J. (2002) Effects of pollutant mixtures. In: Bell, J. N. B.; Treshow,
12 M., eds. *Air Pollution and Plant Life*. 2nd ed. Chichester, United Kingdom: John Wiley & Sons Ltd.; pp.
13 251-272.
- 14 Fangmeier, A.; De Temmerman, L.; Black, C.; Persson, K.; Vorne, V. (2002) Effects of elevated CO₂ and/or ozone on
15 nutrient concentrations and nutrient uptake of potatoes. *Eur. J. Agron.* 17: 353-368.
- 16 Farage, P. K.; Long, S. P. (1999) The effects of O₃ fumigation during leaf development on photosynthesis of wheat and
17 pea: an *in vivo* analysis. *Photosynth. Res.* 59: 1-7.
- 18 Farage, P. K.; Long, S. P.; Lechner, E. G.; Baker, N. R. (1991) The sequence of changes within the photosynthetic
19 apparatus of wheat following short-term exposure to ozone. *Plant Physiol.* 95: 529-535.
- 20 Farquhar, G. D.; Caemmerer, S. V.; Berry, J. A. (1980) A biochemical model of photosynthetic CO₂ assimilation in
21 leaves of C₃ species. *Planta* 149: 78-90.
- 22 Farquhar, G. D.; Sharkey, T. D. (1982) Stomatal conductance and photosynthesis. *Annu. Rev. Plant Physiol.* 33:
23 317-345.
- 24 Federal Land Manager's Air Quality Related Values Workgroup (FLAG). (2000) Phase I report. Denver, CO: National
25 Park Service, Air Resources Division. Available: <http://www2.nature.nps.gov/ard/flagfree/> [11 March, 2003].
- 26 Federal Register. (1997) National ambient air quality standards for ozone; final rule. *F. R.* (July 18) 62: 38856-38896.
- 27 Fenn, M. E.; Bytnerowicz, A. (1993) Dry deposition of nitrogen and sulfur to ponderosa and Jeffrey pine in the San
28 Bernardino National Forest in southern California. *Environ. Pollut.* 81: 277-285.
- 29 Fenn, M. E.; Dunn, P. H. (1989) Litter decomposition across an air-pollution gradient in the San Bernardino
30 Mountains. *Soil Sci. Soc. Am. J.* 53: 1560-1567.
- 31 Fenn, M. E.; Poth, M. A. (1998) Indicators of nitrogen status in California forests. U.S.D.A. Forest Service Gen. Tech.
32 Rep. PSW-GTR-166.
- 33 Fenn, M. E.; Dunn, R. H.; Wilborn, R. (1990) Black stain root disease in ozone-stressed ponderosa pine. *Plant Dis.* 74:
34 426-430.
- 35 Fenn, M. E.; Poth, M. A.; Johnson, D. W. (1996) Evidence for nitrogen saturation in the San Bernardino Mountains in
36 southern California. *For. Ecol. Manage.* 82: 211-230.
- 37 Fenn, M. E.; Poth, M. A.; Aber, J. D.; Baron, J. S.; Bormann, B. T.; Johnson, D. W.; Lemly, A. D.; McNulty, S. G.;
38 Ryan, D. F.; Stottliemyer, R. (1998) Nitrogen excess in North American ecosystems: predisposing factors,
39 ecosystem responses, and management strategies. *Ecol. Appl.* 8: 706-733.
- 40 Fenn, M. E.; Poth, M. A.; Schilling, S. L.; Grainger, D. B. (2000) Throughfall and fog deposition of nitrogen and
41 sulfur at an N-limited and N-saturated site in the San Bernardino Mountains, southern California. *Can. J. For.*
42 *Res.* 30: 1476-1488.
- 43 Fenn, M. E.; Poth, M. A.; Bytnerowicz, A.; Sickman, J. O.; Takemoto, B. K. (2003) Effects of ozone, nitrogen
44 deposition, and other stressors on montane ecosystems in the Sierra Nevada. In: Bytnerowicz, A.; Arbaugh, M.
45 J.; Alonso, R., eds. *Ozone air pollution in the Sierra Nevada: distribution and effects on forests*; pp. 111-155.
46 New York, NY: Elsevier. (Developments in Environmental Science: v. 2).
- 47 Fenn, M. E.; Haeuber, R.; Tonnesen, G. S.; Baron, J. S.; Grossman-Clarke, S.; Hope, D.; Jaffe, D. A.; Copeland, S.;
48 Geiser, L.; Rueth, H. M.; Sickman, J. O. (2003a) Nitrogen emissions, deposition, and monitoring in the western
49 United States. *BioScience* 53: 391-403.
- 50 Fenn, M. E.; Baron, J. S.; Allen, E. B.; Rueth, H. M.; Nydick, K. R.; Geiser, L.; Bowman, W. D.; Sickman, J. O.;
51 Meixner, T.; Johnson, D. W.; Neitlich, P. (2003b) Ecological effects of nitrogen deposition in the western
52 United States. *BioScience* 53: 404-420.
- 53 Ferdinand, J. A.; Fredericksen, T. S.; Kouteric, K. B.; Skelly, J. M. (2000) Leaf morphology and ozone sensitivity of
54 two open pollinated genotypes of black cherry (*Prunus serotina*) seedlings. *Environ. Pollut.* 108: 297-302.
- 55 Findlay, S.; Jones, C. G. (1990) Exposure of cottonwood plants to ozone alters subsequent decomposition. *Oecologia*
56 82: 248-250.

- Findlay, S.; Carreiro, M.; Krischik, V.; Jones, C. G. (1996) Effects of damage to living plants on leaf litter quality. *Ecol. Appl.* 6: 269-275.
- Findley, D. A.; Keever, G. J.; Chappelka, A. H.; Eakes, D. J.; Gillian, D. J. (1997) Differential responses of buddleia (*Buddleia davidii* Franch.) to ozone. *Environ. Pollut.* 98: 105-111.
- Finkelstein, R. R.; Gibson, S. I. (2002) ABA and sugar interactions regulating development: cross-talk or voices in a crowd? *Curr. Opin. Plant Biol.* 5: 26-32.
- Finnan, J. M.; Jones, M. B.; Burke, J. I. (1996) A time-concentration study on the effects of ozone on spring wheat (*Triticum aestivum* L.). 2. A comparison of indices. *Agric. Ecosyst. Environ.* 57: 169-177.
- Finnan, J. M.; Burke, J. L.; Jones, M. B. (1997) An evaluation of indices that describe the impact of ozone on the yield of spring wheat (*Triticum aestivum* L.). *Atmos. Environ.* 31: 2685-2693.
- Finnan, J. M.; Donnelly, A.; Burke, J. I.; Jones, M. B. (2002) The effects of elevated concentrations of carbon dioxide and ozone on potato (*Solanum tuberosum* L.) yield. *Agric. Ecosyst. Environ.* 88: 11-22.
- Fiscus, E. L.; Reid, C. D.; Miller, J. E.; Heagle, A. S. (1997) Elevated CO₂ reduces O₃ flux and O₃-induced yield losses in soybeans: possible implications for elevated CO₂ studies. *J. Exp. Bot.* 48: 307-313.
- Fishman, J.; Watson, C. E.; Larsen, J. C.; Logan, J. A. (1990) Distribution of tropospheric ozone determined from satellite data. *J. Geophys. Res. [Atmos.]* 95: 3599-3617.
- Flagler, R. B., ed. (1998) Recognition of air pollution injury to vegetation: a pictorial atlas. 2nd ed. Pittsburgh, PA: Air & Waste Management Association.
- Flückiger, W.; Braun, S.; Hiltbrunner, E. (2002) Effects of air pollutants on biotic stress. In: Bell, J. N. B.; Treshow, M., eds. *Air Pollution and Plant Life*. 2nd ed. Chichester, United Kingdom: John Wiley & Sons Ltd.; pp. 379-406.
- Fontan, J. A.; Minga, A.; Lopez, A.; Druilhet, A. (1992) Vertical ozone profiles in a pine forest. *Atmos. Environ. Part A* 26A: 863-869.
- Foot, J. P.; Caporn, S. J. M.; Lee, J. A.; Ashenden, T. W. (1996) The effect of long-term ozone fumigation on the growth, physiology and frost sensitivity of *Calluna vulgaris*. *New Phytol.* 133: 505-511.
- Foot, J. P.; Caporn, S. J. M.; Lee, J. A.; Ashenden, T. W. (1997) Evidence that ozone exposure increases the susceptibility of plants to natural frosting episodes. *New Phytol.* 135: 369-374.
- Fortin, M.; Maufette, Y.; Albert, P. J. (1997) The effects of ozone-exposed sugar maple seedlings on the biological performance and the feeding preference of the forest tent caterpillar (*Malacosoma disstria* Hbn.). *Environ. Pollut.* 97: 303-309.
- Fowler, D.; Cape, J. N. (1982) Air pollutants in agriculture and horticulture. In: Unsworth, M. H.; Ormrod, D. P. *Effects of gaseous air pollution in agriculture and horticulture*. London, United Kingdom: Butterworth Scientific; pp. 3-26.
- Fox, S.; Mickler, R. A., eds. (1996) *Impact of air pollutants on southern pine forests*. New York, NY: Springer-Verlag. (Lange, O. L.; Mooney, H. A., eds. *Ecological studies*: v. 118).
- Foyer, C. H.; Souriau, N.; Perret, S.; Lelandais, M.; Kunert, K.-J.; Pruvost, C.; Jouanin, L. (1995) Overexpression of glutathione reductase but not glutathione synthetase leads to increases in antioxidant capacity and resistance to photoinhibition in poplar trees. *Plant Physiol.* 109: 1047-1057.
- Frederick, P. E.; Heath, R. L. (1970) Ozone-induced fatty acid and viability changes in *Chlorella*. *Plant Physiol.* 55: 15-19.
- Fredericksen, T. S.; Joyce, B. J.; Skelly, J. M.; Steiner, K. C.; Kolb, T. E.; Kouterick, K. B.; Savage, J. E.; Snyder, K. R. (1995) Physiology, morphology, and ozone uptake of leaves of black cherry seedlings, saplings, and canopy trees. *Environ. Pollut.* 89: 273-283.
- Fredericksen, T. S.; Kolb, T. E.; Skelly, J. M.; Steiner, K. C.; Joyce, B. J.; Savage, J. E. (1996a) Light environment alters ozone uptake per net photosynthetic rate in black cherry trees. *Tree Physiol.* 16: 485-490.
- Fredericksen, T. S.; Skelly, J. M.; Snyder, K. R.; Steiner, K. C.; Kolb, T. E. (1996b) Predicting ozone uptake from meteorological and environmental variables. *J. Air Waste Manage. Assoc.* 46: 464-469.
- Freer-Smith, P. H.; Taylor, G. (1992) Comparative evaluation of the effects of gaseous pollutants, acidic deposition and mineral deficiencies on gas exchange of trees. *Agric. Ecosyst. Environ.* 42: 321-332.
- Froehlich, J. E.; Itoh, A.; Howe, G. A. (2001) Tomato allene oxide synthase and fatty acid hydroperoxide lyase, two cytochrome P450s involved in oxylipin metabolism, are targeted to different membranes of chloroplast envelope. *Plant Physiol.* 125: 306-317.
- Fuentes, J. D.; Gillespie, T. J. (1992) A gas exchange system to study the effects of leaf surface wetness on the deposition of ozone. *Atmos. Environ. Part A* 26: 1165-1173.
- Fuentes, J. D.; Gillespie, T. J.; den Hartog, G.; Neumann, H. H. (1992) Ozone deposition onto a deciduous forest during dry and wet conditions. *Agric. For. Meteorol.* 62: 1-18.

- 1 Fuentes, J. D.; Hayden, B. P.; Garstang, M.; Lerdau, M.; Fitzjarrald, D.; Baldocchi, D. D.; Monson, R.; Lamb, B.;
2 Geron, C. (2001) New directions: VOCs and biosphere-atmosphere feedbacks. *Atmos. Environ.* 35: 189-191.
- 3 Fuhrer, J. (1994) Effects of ozone on managed pasture. I. Effects of open-top chambers on microclimate, ozone flux,
4 and plant growth. *Environ. Pollut.* 86: 297-305.
- 5 Fuhrer, J. (1994) The critical level for ozone to protect agricultural crops - an assessment of data from European
6 open-top chamber experiments. In: Fuhrer, J.; Achermann, eds. *Critical levels for ozone: a UN-ECE workshop*
7 *report; FAC report no. 16.* Liebefeld-Bern, Switzerland: Swiss Federal Research Station for Agricultural
8 Chemistry and Environmental Hygiene; pp. 42-57.
- 9 Fuhrer, J. (1996) The critical level for effects of ozone on crops and the transfer to mapping. In: Kärenlampi, L.;
10 Skärby, L., eds. *Critical levels for ozone in Europe: testing and finalizing the concepts: UN-ECE workshop*
11 *report.* UN-ECE Convention on Long-Range Transboundary Air Pollution; April; Kuopio, Finland. Kuopio,
12 Finland: University of Kuopio, Department of Ecology and Environmental Science; pp. 27-43.
- 13 Fuhrer, J.; Grimm, A. G.; Tschannen, W.; Shariat-Madari, H. (1992) The response of spring wheat (*Triticum aestivum*
14 L.) to ozone at higher elevations: II. changes in yield, yield components and grain quality in response to ozone
15 flux. *New Phytol.* 121: 211-219.
- 16 Fuhrer, J.; Shariat-Madari, H.; Perler, R.; Tschannen, W.; Grub, A. (1994) Effects of ozone on managed pasture: II.
17 Yield, species composition, canopy structure, and forage quality. *Environ. Pollut.* 86: 307-314.
- 18 Fuhrer, J.; Skärby, L.; Ashmore, M. R. (1997) Critical levels for ozone effects on vegetation in Europe. *Environ.*
19 *Pollution* 97: 91-106.
- 20 Fumagalli, I.; Mignanego, L.; Violini, G. (1997) Effects of tropospheric ozone on white clover plants exposed in
21 open-top chambers or protected by the antioxidant ethylene-diurea (EDU). *Agronomie* 17: 271-281.
- 22 Fumagalli, I.; Ambrogi, R.; Mignanego, L. (2001) Yield responses of plants exposed to ambient ozone in the river Po
23 valley (Italy). *Agronomie* 21: 227-233.
- 24 Garcia, P.; Dixon, B.; Mjelde, J.; Adams, R. (1986) Measuring the benefits of environmental change using a duality
25 approach: the case of ozone and Illinois cash grain farms. *J. Environ. Econ. Manage.* 13: 69-80.
- 26 Gasper, T.; Penel, C.; Castillo, F. J.; Greppin, H. (1985) A two step control of basic and acidic peroxidases and its
27 significance for growth and development. *Physiol. Plant.* 64: 418-423.
- 28 Gate, I. M.; McNeill, S.; Ashmore, M. R. (1995) Effects of air pollution on the searching behaviour of an insect
29 parasitoid. *Water Air Soil Pollut.* 85: 1425-1430.
- 30 Gates, D. M. (1968) Transpiration and leaf temperature. *Annu. Rev. Plant Physiol.* 19: 211-238.
- 31 Gaucher, C.; Costanzo, N.; Afif, D.; Mauffette, Y.; Chevrier, N.; Dizengremel, P. (2003) The impact of elevated ozone
32 and carbon dioxide on young *Acer saccharum* seedlings. *Physiol. Plant.* 117: 392-402.
- 33 Gelang, J.; Pleijel, H.; Sild, E.; Danielsson, H.; Younis, S.; Selldén, G. (2000) Rate and duration of grain filling in
34 relation to flag leaf senescence and grain yield in spring wheat (*Triticum aestivum*) exposed to different
35 concentrations of ozone. *Physiol. Plant.* 110: 366-375.
- 36 Gerant, D.; Podor, M.; Grieu, P.; Afif, D.; Cornu, S.; Morabito, D.; Banvoy, J.; Robin, C.; Dizengremel, P. (1996)
37 Carbon metabolism enzyme activities and carbon partitioning in *Pinus halepensis* mill. Exposed to mild drought
38 and ozone. *J. Plant Physiol.* 148: 142-147.
- 39 Gerosa, G.; Marzuoli, R.; Bussotti, F.; Pancrazi, M.; Ballarin-Denti, A. (2003) Ozone sensitivity of *Fagus sylvatica*
40 and *Fraxinus excelsior* young trees in relation to leaf structure and foliar ozone uptake. *Environ. Pollut.* 125:
41 91-98.
- 42 Ghosh, S.; Skelly, J. M.; Innex, J. L.; Skelly, L. (1998) Temporal development of visual ozone injury on the foliage of
43 *Prunus serotina* - a statistical evaluation. *Environ. Pollut.* 192: 287-300.
- 44 Gifford, R. M.; Morison, J. I. L. (1993) Crop responses to the global increase in atmospheric carbon dioxide
45 concentration. In: Buxton, D. R., et al., eds. *International crop science I.* Madison, WI: Crop Science Society of
46 America; pp. 325-331.
- 47 Gimeno, B. S.; Bermejo, V.; Reinert, R. A.; Zheng, Y.; Barnes, J. D. (1999) Adverse effects of ambient ozone on
48 watermelon yield and physiology at a rural site in eastern Spain. *New Phytol.* 144: 245-260.
- 49 Goldstein, A. H.; Kurpius, M. R.; Panek, J. A. (2003) Ozone uptake by ponderosa pine in the Sierra Nevada: a
50 measurement perspective. In: Bytnerowicz, A.; Arbaugh, M. J.; Alonso, R., eds. *Ozone air pollution in the*
51 *Sierra Nevada: distribution and effects on forests*; pp. 83-109. New York, NY: Elsevier. (Developments in
52 *Environmental Science*: v. 2).
- 53 Goodyear, S. N.; Ormrod, D. P. (1988) Tomato response to concurrent and sequential NO₂ and O₃ exposures. *Environ.*
54 *Pollut.* 51: 315-326.

- Grams, T. E. E.; Anegg, S.; Haberle, K.-H.; Langebartels, C.; Matyssek, R. (1999) Interactions of chronic exposure to elevated CO₂ and O₃ levels in the photosynthetic light and dark reactions of European beech (*Fagus sylvatica*). *New Phytol.* 144: 95-107.
- Grantz, D. A.; Farrar, J. F. (1999) Acute exposure to ozone inhibits rapid carbon translocation from source leaves of Pima cotton. *J. Exp. Bot.* 50: 1253-1262.
- Grantz, D. A.; Yang, S. D. (1996) Effect of O₃ on hydraulic architecture in pima cotton. *Plant Physiol.* 112: 1649-1657.
- Grantz, D. A.; Yang, S. (2000) Ozone impacts on allometry and root hydraulic conductance are not mediated by source limitation nor developmental age. *J. Exp. Bot.* 51: 919-927.
- Grantz, D. A.; Zhang, X. J.; Massman, W. J.; Den Hartog, G.; Neumann, H. H.; Pederson, J. R. (1995) Effects of stomatal conductance and surface wetness on ozone deposition in field-grown grape. *Atmos. Environ.* 29: 3189-3198.
- Grantz, D. A.; Zhang, X. J.; Massman, W.; Delany, A.; Pederson, R. (1997) Ozone deposition to a cotton (*Gossypium hirsutum* L.) field: stomatal and surface wetness effects during the California Ozone Deposition experiment. *Agric. For. Meteorol.* 85: 19-31.
- Graumlich, L. J. (1993) A 1000-year record of temperature and precipitation in the Sierra Nevada. *Quaternary Res.* 39: 249-255.
- Greenwood, P.; Greenhalgh, A.; Baker, C.; Unsworth, M. (1982) A computer-controlled system for exposing field crops to gaseous air pollutants. *Atmos. Environ.* 16: 2261-2266.
- Gregg, J. W.; Jones, C. G.; Dawson, T. E. (2003) Urbanization effects on tree growth in the vicinity of New York City [letter]. *Nature (London, U.K.)* 424: 183-187.
- Greitner, C. S.; Pell, E. J.; Winner, W. E. (1994) Analysis of aspen foliage exposed to multiple stresses: ozone, nitrogen deficiency and drought. *New Phytol.* 127: 579-589.
- Grimes, H. D.; Perkins, K. K.; Boss, W. F. (1983) Ozone degrades into hydroxyl radical under physiological conditions. *Plant Physiol.* 72: 1016-1020.
- Grosjean, D.; Williams, E. L., II; Grosjean, E. (1995) Monitoring ambient ozone with a network of passive samplers: a feasibility study. *Environ. Pollut.* 88: 267-273.
- Gross, G. G.; Janse, C. (1977) Formation of NADH and hydrogen peroxide by cell wall-associated enzymes from forsythia xylem. *Z. Pflanzenphysiol.* 84: 447-452.
- Groth, J. V.; Krupa, S. V. (2000) Crop ecosystem responses to climatic change: interactive effects of ozone, ultraviolet-B radiation, sulphur dioxide and carbon dioxide on crops. In: Reddy, K. R.; Hodges, H. F., eds. *Climate change and global crop productivity*. New York, NY: CABI Publishing; pp. 387-405.
- Grulke, N. E. (1999) Physiological responses of ponderosa pine to gradients of environmental stressors. In: Miller, P. R.; McBride, J. R., eds. *Oxidant Air Pollution Impacts in the Montane Forests of Southern California*. New York, NY: Springer-Verlag; pp. 126-163. (Ecological Studies, v. 134).
- Grulke, N. E.; Balduman, L. (1999) Deciduous conifers: high N deposition and O₃ exposure effects on growth and biomass allocation in ponderosa pine. *Water Air Soil Pollut.* 116: 235-248.
- Grulke, N. E.; Lee, E. H. (1997) Assessing visible ozone-induced injury in ponderosa pine. *Can. J. For. Res.* 27: 1658-1668.
- Grulke, N. E.; Miller, P. R. (1994) Changes in gas-exchange characteristics during the life-span of giant sequoia: implications for response to current and future concentrations of atmospheric ozone. *Tree Physiol.* 14: 659-668.
- Grulke, N. E.; Retzlaff, W. A. (2001) Changes in physiological attributes of ponderosa pine from seedling to mature tree. *Tree Physiol.* 21: 275-286.
- Grulke, N. E.; Miller, P. R.; Scioli, D. (1996) Response of giant sequoia canopy foliage to elevated concentrations of atmospheric ozone. *Tree Physiol.* 16: 575-581.
- Grulke, N. E.; Andersen, C. P.; Fenn, M. E.; Miller, P. R. (1998) Ozone exposure and nitrogen deposition lowers root biomass of ponderosa pine in the San Bernardino Mountains, California. *Environ. Pollut.* 103: 63-73.
- Grulke, N. E.; Miller, P. R.; Leininger, T. D. (1998) Effect of ozone exposure on seasonal gas exchange of five western conifers. In: Bytnerowicz, A.; Arbaugh, M. J.; Schilling, S. L., eds. *Proceedings of the International Symposium on Air Pollution and Climate Change Effects on Forest Ecosystems*; February, 1996; Riverside, CA. Albany, CA: USDA Forest Service, Pacific Southwest General Technical Report no. PSW-GTR 166; pp. 229-238.
- Grulke, N. E.; Andersen, C. P.; Hogsett, W. E. (2001) Seasonal changes in above- and belowground carbohydrate concentrations of ponderosa pine along a pollution gradient. *Tree Physiol.* 21: 173-181.

- 1 Grulke, N. E.; Preisler, H. K.; Fan, C. C.; Retzlaff, W. A. (2002) A statistical approach to estimate O₃ uptake of
2 ponderosa pine in a Mediterranean climate. *Environ. Pollut.* 119: 163-175.
- 3 Grulke, N. E.; Preisler, H. K.; Rose, C.; Kirsch, J.; Balduman, L. (2002) O₃ uptake and drought stress effects on carbon
4 acquisition of ponderosa pine in natural stands. *New Phytol.* 154: 621-631.
- 5 Grulke, N. E.; Johnson, R.; Esperanza, A.; Jones, D.; Nguyen, T.; Posch, S.; Tausz, M. (2003a) Canopy transpiration
6 of Jeffrey pine in mesic and xeric microsites: O₃ uptake and injury response. *Trees (Berlin, Ger.)* 17: 292-298.
- 7 Grulke, N. E.; Johnson, R.; Monschein, S.; Nikolova, P.; Tausz, M. (2003b) Variation in morphological and
8 biochemical O₃ injury attributes of mature Jeffrey pine within canopies and between microsites. *Tree Physiol.*
9 23: 923-929.
- 10 Grulke, N. E.; Alonso, R.; Nguyen, T.; Cascio, C.; Dobrowolski, W. (2004) Stomata open at night in pole-sized and
11 mature ponderosa pine: implications for O₃ exposure metrics. *Tree Physiol.* 24: 1001-1010.
- 12 Grünhage, L.; Haenel, H.-D. (1997) PLATIN (PLant-ATmosphere-INteraction) I: a model of plant-atmosphere
13 interaction for estimating absorbed doses of gaseous air pollutants. *Environ. Pollut.* 98: 37-50.
- 14 Grünhage, L.; Jäger, H.-J. (1994) Influence of the atmospheric conductivity on the ozone exposure of plants under
15 ambient conditions: considerations for establishing ozone standards to protect vegetation. *Environ. Pollut.* 85:
16 125-129.
- 17 Grünhage, L.; Jäger, H.-J. (1996) Critical levels for ozone, ozone exposure potentials of the atmosphere or critical
18 absorbed doses for ozone: a general discussion. In: Kärenlampi, L.; Skärby, L., eds. Critical levels for ozone in
19 Europe: testing and finalizing the concepts: UN-ECE workshop report. UN-ECE Convention on Long-Range
20 Transboundary Air Pollution; April; Kuopio, Finland. Kuopio, Finland: University of Kuopio, Department of
21 Ecology and Environmental Science; pp. 151-168.
- 22 Grünhage, L.; Jäger, H.-J. (2003) From critical levels to critical loads for ozone: a discussion of a new experimental
23 and modelling approach for establishing flux-response relationships for agricultural crops and native plant
24 species. *Environ. Pollut.* 125: 99-110.
- 25 Grünhage, L.; Dämmgen, U.; Haenel, H.-D.; Jäger, H.-J.; Holl, A.; Schmitt, J.; Hanewald, K. (1993) A new potential
26 air-quality criterion derived from vertical flux densities of ozone and from plant response. *Angew. Bot.* 67:
27 9-13.
- 28 Grünhage, L.; Dämmgen, U.; Haenel, H.-D.; Jäger, H.-J. (1994) Response of a grassland ecosystem to air pollutants:
29 III--the chemical climate: vertical flux densities of gaseous species in the atmosphere near the ground. *Environ.*
30 *Pollut.* 85: 43-49.
- 31 Grünhage, L.; Jäger, H.-J.; Haenel, H.-D.; Hanewald, K.; Krupa, S. (1997) PLATIN (plant-atmosphere interaction) II:
32 co-occurrence of high ambient ozone concentrations and factors limiting plant absorbed dose. *Environ. Pollut.*
33 98: 51-60.
- 34 Grünhage, L.; Jäger, H.-J.; Haenel, H.-D.; Löpmeier, F.-J.; Hanewald, K. (1999) The European critical levels for
35 ozone: improving their usage. *Environ. Pollut.* 105: 163-173.
- 36 Grünhage, L.; Krause, G. H. M.; Köllner, B.; Bender, J.; Weigel, H.-J.; Jäger, H.-J.; Guderian, R. (2001) A new
37 flux-orientated concept to derive critical levels for ozone to protect vegetation. *Environ. Pollut.* 111: 355-362.
- 38 Grünhage, L.; Krupa, S. V.; Legge, A. H.; Jäger, H.-J. (2004) Ambient flux-based critical values of ozone for
39 protecting vegetation: differing spatial scales and uncertainties in risk assessment. *Atmos. Environ.* 38:
40 2433-2437.
- 41 Grunwald, C.; Endress, A. G. (1985) Foliar sterols in soybeans exposed to chronic levels of ozone. *Plant Physiol.* 77:
42 245-247.
- 43 Grütters, U.; Fangmeier, A.; Jäger, H.-J. (1995) Modelling stomatal responses of spring wheat (*Triticum aestivum* L.
44 cv. Turbo) to ozone and different levels of water supply. *Environ. Pollut.* 87: 141-149.
- 45 Guderian, R. (1977) Discussion of the suitability of plant responses as a basis for air pollution control measures. In:
46 Air pollution: phytotoxicity of acidic gases and its significance in air pollution control. Berlin, Federal Republic
47 of Germany: Springer Verlag; pp. 75-97. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S., eds. Ecological
48 studies: analysis and synthesis, v. 22).
- 49 Guidi, L.; Tonini, M.; Soldatini, G. F. (2000) Effects of high light and ozone fumigation on photosynthesis in
50 *Phaseolus vulgaris*. *Plant Physiol. Biochem.* 38: 717-725.
- 51 Guzy, M. R.; Heath R. L. (1993) Responses to ozone of varieties of common bean (*Phaseolus vulgaris* L.) *New*
52 *Phytol.* 124: 617-625.

- 1 Haber, F. (1924) Zur Geschichte des Gaskrieges [On the history of the gas war]. In: Fuenf Vortraege aus den Jahren
- 2 1920-1923 [Five lectures from the years 1920-1923]. Berlin, Germany: Verlag von Julius Springer; pp. 76-92.
- 3 Hacour, A.; Craigon, J.; Vandermeiren, K.; Ojanperä, K.; Pleijel, H.; Danielsson, H.; Høgy, P.; Finnán, J.; Bindi, M.
- 4 (2002) CO₂ and ozone effects on canopy development of potato crops across Europe. *Eur. J. Agron.* 17:
- 5 257-272.
- 6 Häberle, K. H.; Werner, H.; Fabian, P.; Pretzch, H.; Reiter, I.; Matyssek, R. (1999) 'Free-air' ozone fumigation of
- 7 mature forest trees: a concept for validating AOT40 under stand conditions. In: Fuhrer, J.; Achermann, B., eds.
- 8 Critical levels for ozone: level II: workshop under the Convention on Long-Range Transboundary Air Pollution
- 9 of the United Nations Economic Commission for Europe (UN/ECE); April; Gerzensee, Switzerland. Berne,
- 10 Switzerland: Bundesamt für Umwelt, Wald und Landschaft; pp. 133-137.
- 11 Hakala, K.; Mela, T. (1996) The effects of prolonged exposure to elevated temperatures and elevated CO₂ levels on the
- 12 growth, yield and dry matter partitioning of field-sown meadow fescue. *Agric. Food Sci. Finl.* 5: 285-298.
- 13 Hankamer, B.; Barber, J.; Boekema, E. J. (1997) Structure and membrane organization of photosystem II in green
- 14 plants. *Ann. Rev. Plant Physiol. Mol. Biol.* 48: 641-671.
- 15 Hanson, P. J.; Samuelson, L. J.; Wulschleger, S. D.; Tabberer, T. A.; Edwards, G. S. (1994) Seasonal patterns of
- 16 light-saturated photosynthesis and leaf conductance for mature and seedling *Quercus rubra* L. foliage:
- 17 differential sensitivity to ozone exposure. *Tree Physiol.* 14: 1351-1366.
- 18 Hao, X.; Hale, B. A.; Ormrod, D. P.; Papadopoulos, A. P. (2000) Effects of pre-exposure to ultraviolet-B radiation on
- 19 responses of tomato (*Lycopersicon esculentum* cv. New Yorker) to ozone in ambient and elevated carbon
- 20 dioxide. *Environ. Pollut.* 110: 217-224.
- 21 Harris, M. J.; Bailey-Serres, J. N. (1994) Ozone effects on gene expression and molecular approaches to breeding for
- 22 air pollution resistance. In: Basra, A. S., ed. Stress-induced gene expression in plants. Langhorne, PA: Harwood
- 23 Academic Publishers; pp. 185-207.
- 24 Harward, M.; Treshow, M. (1975) Impact of ozone on the growth and reproduction of understory plants in the Aspen
- 25 zone of western U.S.A. *Environ. Conserv.* 2: 17-23.
- 26 Hausladen, A.; Madamanchi, N. R.; Fellows, S.; Alscher, R. G.; Amundson, R. G. (1990) Seasonal changes in
- 27 antioxidants in red spruce as affected by ozone. *New Phytol.* 115: 447-458.
- 28 Haynes, R. W.; Adams, D. M. (1992) Assessing economic impacts of air pollution damage to U.S. forests. In: de
- 29 Steiguer, J. E., ed. The economic impact of air pollution on timber markets: studies from North America and
- 30 Europe. Asheville, NC: U.S. Department of Agriculture, Southeastern Forest Experiment Station; general
- 31 technical report no. SE-75.
- 32 Heagle, A. S. (1979) Effects of growth media, fertiliser rate and hour and season of exposure on sensitivity of four
- 33 soybean cultivars to ozone. *Environ. Pollut.* 18: 313-322.
- 34 Heagle, A. S.; Body, D. E.; Heck, W. W. (1973) An open-top field chamber to assess the impact of air pollution on
- 35 plants. *J. Environ. Qual.* 2: 365-368.
- 36 Heagle, A. S.; Body, D. E.; Neely, G. E. (1974) Injury and yield responses of soybean to chronic doses of ozone and
- 37 sulfur dioxide in the field. *Phytopathology* 64: 132-136.
- 38 Heagle, A. S.; Spencer, S.; Letchworth, M. B. (1979) Yield response of winter wheat to chronic doses of ozone. *Can. J.*
- 39 *Bot.* 57: 1999-2005.
- 40 Heagle, A. S.; Letchworth, M. B.; Mitchell, C. A. (1983) Effects of growth medium and fertilizer rate on the yield
- 41 response of soybeans exposed to chronic doses of ozone. *Phytopathology* 73: 134-139.
- 42 Heagle, A. S.; Heck, W. W.; Lesser, V. M.; Rawlings, J. O. (1987) Effects of daily ozone exposure duration and
- 43 concentration fluctuation on yield of tobacco. *Phytopathology* 77: 856-862.
- 44 Heagle, A. S.; Kress, L. W.; Temple, P. J.; Kohut, R. J.; Miller, J. E.; Heggstad, H. E. (1988) Factors influencing
- 45 ozone dose-yield response relationships in open-top field chamber studies. In: Heck, W. W.; Taylor, O. C.;
- 46 Tingey, D. T., eds. Assessment of crop loss from air pollutants: proceedings of an international conference;
- 47 October, 1987; Raleigh, NC. New York, NY: Elsevier Applied Science; pp. 141-179.
- 48 Heagle, A. S.; Rebbeck, J.; Shafer, S. R.; Blum, U.; Heck, W. W. (1989) Effects of long-term ozone exposure and soil
- 49 moisture deficit on growth of a ladino clover-tall fescue pasture. *Phytopathology* 79: 128-136.
- 50 Heagle, A. S.; McLaughlin, M. R.; Miller, J. E.; Joyner, R. L.; Spruill, S. E. (1991) Adaptation of a white clover
- 51 population to ozone stress. *New Phytol.* 119: 61-68.
- 52 Heagle, A. S.; Miller, J. E.; Rawlings, J. O.; Vozzo, S. F. (1991) Effect of growth stage on soybean response to chronic
- 53 ozone exposure. *J. Environ. Qual.* 20: 562-570.

- 1 Heagle, A. S.; Miller, J. E.; Sherrill, D. E.; Rawlings, J. O. (1993) Effects of ozone and carbon dioxide mixtures on
2 two clones of white clover. *New Phytol.* 123: 751-762.
- 3 Heagle, A. S.; Brandenburg, R. L.; Burns, J. C.; Miller, J. E. (1994) Ozone and carbon dioxide effects on spider mites
4 in white clover and peanut. *J. Environ. Qual.* 23: 1168-1176.
- 5 Heagle, A. S.; Miller, J. E.; Sherrill, D. E. (1994) A white clover system to estimate effects of tropospheric ozone on
6 plants. *J. Environ. Qual.* 23: 613-621.
- 7 Heagle, A. S.; Miller, J. E.; Chevone, B. I.; Dreschel, T. W.; Manning, W. J.; McCool, P. M.; Morrison, C. L.; Neely,
8 G. E.; Rebbeck, J. (1995) Response of a white clover indicator system to tropospheric ozone at eight locations
9 in the United States. *Water Air Soil Pollut.* 85: 1373-1378.
- 10 Heagle, A. S.; Reinert, R. A.; Miller, J. E. (1996) Response of white clover to ozone in different environments. *J.*
11 *Environ. Qual.* 25: 273-278.
- 12 Heagle, A. S.; Miller, J. E.; Pursley, W. A. (1998) Influence of ozone stress on soybean response to carbon dioxide
13 enrichment: III. Yield and seed quality. *Crop Sci.* 38: 128-134.
- 14 Heagle, A. S.; Miller, J. E.; Booker, F. L.; Pursley, W. A. (1999) Ozone stress, carbon dioxide enrichment, and
15 nitrogen fertility interactions in cotton. *Crop Sci.* 39: 731-741.
- 16 Heagle, A. S.; Miller, J. E.; Pursley, W. A. (2000) Growth and yield responses of winter wheat to mixtures of ozone
17 and carbon dioxide. *Crop Sci.* 40: 1656-1664.
- 18 Heagle, A. S.; Stefanski, L. A. (2000) Relationships between ambient ozone regimes and white clover forage
19 production using different ozone exposure indexes. *Atmos. Environ.* 34: 735-744.
- 20 Heagle, A. S.; Miller, J. E.; Burkey, K. O.; Eason, G.; Pursley, W. A. (2002) Growth and yield responses of snap bean
21 to mixtures of carbon dioxide and ozone. *J. Environ. Qual.* 31: 2008-2014.
- 22 Heath, R. L. (1980) Initial events in injury to plants by air pollutants. *Annu. Rev. Plant Physiol.* 31: 395-431.
- 23 Heath, R. L. (1984) Air pollutant effects on biochemicals derived from metabolism: organic, fatty and amino acids. In:
24 Koziol, M. J.; Whatley, F. R., eds. *Gaseous Air Pollutants and Plant Metabolism*. London, United Kingdom:
25 Butterworths; pp. 275-290.
- 26 Heath, R. L. (1987) The biochemistry of ozone attack on the plasma membrane of plant cells. *Recent Adv. Phytochem.*
27 21: 29-54.
- 28 Heath, R. L. (1988) Biochemical mechanisms of pollutant stress. In: Heck, W. W.; Taylor, O. C.; Tingey, D. T., eds.
29 *Assessment of crop loss from air pollutants: proceedings of an international conference; October, 1987; Raleigh,*
30 *NC. London, United Kingdom: Elsevier Applied Science; pp. 259-286.*
- 31 Heath, R. L. (1994a) Alterations of plant metabolism by ozone exposure. In: Alscher, R. G.; Wellburn, A. R., eds.
32 *Plant Responses to the Gaseous Environment: molecular, metabolic, and physiological aspects*. London, United
33 Kingdom: Chapman & Hall; pp. 121-146.
- 34 Heath, R. L. (1994b) Possible mechanisms for the inhibition of photosynthesis by ozone. *Photosynth. Res.* 39: 439-451.
- 35 Heath, R. L. (1996) The modification of photosynthetic capacity induced by ozone exposure. In: Baker, N. R., ed.
36 *Photosynthesis and the environment*. Dordrecht, The Netherlands: Kluwer Academic Publishers; pp. 409-433.
37 (Advances in Photosynthesis, v.5).
- 38 Heath, R. L.; Castillo, F. J. (1987) Membrane disturbances in response to air pollutants. In: Schulte-Hostede, S.;
39 Darrall, N. M.; Blank, L. W.; Wellburn, A. R., eds. *Air pollution and plant metabolism*. New York, NY:
40 Elsevier Applied Science; pp. 55-75.
- 41 Heath, R. L.; Taylor, G. E., Jr. (1997) Physiological processes and plant responses to ozone exposure. In: Sandermann,
42 H.; Wellburn, A. R.; Heath, R. L., eds. *Forest Decline and Ozone: a comparison of controlled chamber and field*
43 *experiments*. New York, NY: Springer-Verlag; pp. 317-368. (Ecological Studies, v. 127).
- 44 Heck, W. W.; Cowling, E. B. (1997) The need for a long term cumulative secondary ozone standard - an ecological
45 perspective. *EM (January)*: 23-33.
- 46 Heck, W. W.; Dunning, J. A.; Hindawi, I. J. (1966) Ozone: nonlinear relation of dose and injury in plants. *Science*
47 (Washington, DC) 151: 577-578.
- 48 Heck, W. W.; Philbeck, R. B.; Dunning, J. A. (1978) A continuous stirred tank reactor (CSTR) system for exposing
49 plants to gaseous air contaminants: principles, specifications, construction, and operation. *U.S. Agric. Res. Serv.*
50 *South. Reg. Rep. ARS-S-181*.
- 51 Heck, W. W.; Taylor, O. C.; Adams, R.; Bingham, G.; Miller, J.; Preston, E.; Weinstein, L. (1982) Assessment of crop
52 loss from ozone. *J. Air Pollut. Control Assoc.* 32: 353-361.

- 1 Heck, W. W.; Heagle, A. S.; Miller, J. E.; Rawlings, J. O. (1991) A national program (NCLAN) to assess the impact of
2 ozone on agricultural resources. In: Berglund, R. L.; Lawson, D. R.; McKee, D. J., eds. Tropospheric ozone and
3 the environment: papers from an international conference; March 1990; Los Angeles, CA. Pittsburgh, PA: Air
4 & Waste Management Association; pp. 225-254. (A&WMA transactions series no. TR-19).
- 5 Heck, W. W.; Furiness, C. S.; Cowling, E. B.; Sims, C. K. (1998) Effects of ozone on crop, forest, and natural
6 ecosystems: assessment of research needs. *EM* (October): 11-22.
- 7 Heggstad, H. E. (1988) Reduction in soybean seed yields by ozone air pollution? *JAPCA* 38: 1040-1041.
- 8 Heliövaara, K.; Väisänen, R. (1993) Insects and pollution. Boca Raton, FL: CRC Press.
- 9 Hendrey, G. R.; Kimball, B. A. (1994) The FACE program. *Agric. For. Meteorol.* 70: 3-14.
- 10 Hendrey, G. R.; Ellsworth, D. S.; Lewin, K. F.; Nagy, J. (1999) A free-air enrichment system for exposing tall forest
11 vegetation to elevated atmospheric CO₂. *Global Change Biol.* 5: 293-309.
- 12 Hertstein, U.; Fangmeier, A.; Jäger, H. J. (1996) ESPACE-wheat (European stress physiology and climate experiment
13 - Project 1: wheat): objectives, general approach, and first results. *J. Appl. Bot.* 70: 172-180.
- 14 Hertstein, U.; Colls, J.; Ewert, F.; Van Oijen, M. (1999) Climatic conditions and concentrations of carbon dioxide and
15 air pollutants during 'ESPACE--wheat' experiments. *Eur. J. Agron.* 10: 163-169.
- 16 Hildebrand, E.; Skelly, J. M.; Fredericksen, T. S. (1996) Foliar response of ozone-sensitive hardwood tree species from
17 1991 to 1993 in the Shenandoah National Park, Virginia. *Can. J. For. Res.* 26: 658-669.
- 18 Hogsett, W. E.; Plocher, M.; Wildman, V.; Tingey, D. T.; Bennett, J. P. (1985) Growth response of two varieties of
19 slash pine seedlings to chronic ozone exposures. *Can. J. Bot.* 63: 2369-2376.
- 20 Hogsett, W. E.; Tingey, D. T.; Holman, S. R. (1985) A programmable exposure control system for determination of the
21 effects of pollutant exposure regimes on plant growth. *Atmos. Environ.* 19: 1135-1145.
- 22 Hogsett, W. E.; Olszyk, D.; Ormrod, D. P.; Taylor, G. E., Jr.; Tingey, D. T. (1987a) Air pollution exposure systems
23 and experimental protocols: volume 1: a review and evaluation of performance. Corvallis, OR: U.S.
24 Environmental Protection Agency, Environmental Research Laboratory; EPA report no. EPA/600/3-87/037a.
25 Available from: NTIS, Springfield, VA; PB88-181680.
- 26 Hogsett, W. E.; Olszyk, D.; Ormrod, D. P.; Taylor, G. E., Jr.; Tingey, D. T. (1987b) Air pollution exposure systems
27 and experimental protocols: volume 2: description of facilities. Corvallis, OR: U.S. Environmental Protection
28 Agency, Environmental Research Laboratory; EPA report no. EPA/600/3-87/037b. Available from: NTIS,
29 Springfield, VA; PB88-181698.
- 30 Hogsett, W. E.; Tingey, D. T.; Lee, E. H. (1988) Ozone exposure indices: concepts for development and evaluation of
31 their use. In: Heck, W. W.; Taylor, O. C.; Tingey, D. T., eds. Assessment of crop loss from air pollutants:
32 proceedings of an international conference; October 1987; Raleigh, NC. New York, NY: Elsevier Applied
33 Science; pp. 107-138.
- 34 Hogsett, W. E.; Herstrom, A. A.; Laurence, J. A.; Lee, E. H.; Weber, J. E.; Tingey, D. T. (1995) Risk characterization
35 of tropospheric ozone to forests. In: Lee, S. D.; Schneider, T., eds. Comparative risk analysis and priority setting
36 for air pollution issues: proceedings of the 4th U.S.-Dutch international symposium; June 1993; Keystone, CO.
37 Pittsburgh, PA: Air & Waste Management Association; pp. 119-145. (AWMA publication no. VIP-43).
- 38 Hogsett, W. E.; Weber, J. E.; Tingey, D.; Herstrom, A.; Lee, E. H.; Laurence, J. A. (1996) An approach for
39 characterizing tropospheric ozone risk to forests. *Environ. Manage. (N. Y.)* 21: 105-120.
- 40 Hogsett, W. E.; Weber, J. E.; Tingey, D.; Herstrom, A.; Lee, E. H.; Laurence, J. A. (1997) Environmental auditing: an
41 approach for characterizing tropospheric ozone risk to forests. *Environ. Manage.* 21: 105-120.
- 42 Holland, D. M.; Saltzman, N.; Cox, L. H.; Nychka, D. (1999) Spatial prediction of sulfur dioxide in the eastern United
43 States. In: Gómez-Hernández, J.; Soares, A.; Froidevaux, R., eds. *geoENV II: geostatistics for environmental*
44 *applications*, proceedings of the second European conference; November 1998; Valencia, Spain. Dordrecht, The
45 Netherlands: Kluwer Academic Publishers; pp. 65-76. (Quantitative geology and geostatistics: v. 10).
- 46 Holopainen, J. K.; Kainulainen, P.; Oksanen, J. (1997) Growth and reproduction of aphids and levels of free amino
47 acids in Scots pine and Norway spruce in an open-air fumigation with ozone. *Global Change Biol.* 3: 139-147.
- 48 Holton, M. K.; Lindroth, R. L.; Nordheim, E. V. (2003) Foliar quality influences tree-herbivore-parasitoid interactions:
49 effects of elevated CO₂, O₃, and plant genotype. *Oecologia* 137: 233-244.
- 50 Horemans, N.; Foyer, C. H.; Asard, H. (2000) Transport and action of ascorbate at the plant plasma membrane. *Trends*
51 *Plant Sci.* 5: 263-267.
- 52 Hormaza, J. I.; Pinney, K.; Polito, V. S. (1996) Correlation in the tolerance to ozone between sporophytes and male
53 gametophytes of several fruit and tree nut species (Rosaceae). *Sex. Plant Reprod.* 9: 44-48.

- Houpis, J. L. J.; Costella, M. P.; Cowles, S. (1991) A branch exposure chamber for fumigating ponderosa pine to atmospheric pollution. *J. Environ. Qual.* 20: 467-474.
- Hui, D.; Sims, D. A.; Johnson, D. W.; Chang, W.; Luo, Y. (2002) Effects of gradual versus step increases in carbon dioxide on *Plantago* photosynthesis and growth in a microcosm study. *Environ. Exp. Bot.* 47: 51-66.
- Hummel, R. L.; Brandenburg, R. L.; Heagle, A. S.; Arellano, C. (1998) Effects of ozone on reproduction of two-spotted spider mite (Acari: Tetranychidae) on white clover. *Environ. Entomol.* 27: 388-394.
- Idso, K. E.; Idso, S. B. (1994) Plant responses to atmospheric CO₂ enrichment in the face of environmental constraints: a review of the past 10 years' research. *Agric. For. Meteorol.* 69: 153-203.
- Innes, J. L.; Skelly, J. M.; Schaub, M. (2001) Ozone and broadleaved species. A guide to the identification of ozone-induced foliar injury. Bern, Switzerland: Paul Haupt Publishers.
- Intergovernmental Panel on Climate Change (IPCC). (1996) Climate change 1995: the science of climate change: contribution of working group I to the second assessment of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom: Cambridge University Press.
- Intergovernmental Panel on Climate Change (IPCC). (1998) The regional impacts of climate change: an assessment of vulnerability, a special report of IPCC Working Group II. Watson, R. T.; Zinyowera, M. C.; Moss, R. H., eds. Cambridge, United Kingdom: Cambridge University Press.
- Isebrands, J. G.; Dickson, R. E.; Rebbeck, J.; Karnosky, D. F. (2000) Interacting effects of multiple stresses on growth and physiological processes in northern forest trees. In: Mickler, R. A.; Birsdey, R. A.; Hom, J., eds. Responses of northern U.S. forests to environmental change. New York, NY: Springer-Verlag; pp. 149-180. (Ecological studies: v. 139).
- Isebrands, J. G.; McDonald, E. P.; Kruger, E.; Hendrey, G.; Percy, K.; Pregitzer, K.; Sober, J.; Karnosky, D. F. (2001) Growth responses of *Populus tremuloides* clones to interacting carbon dioxide and tropospheric ozone. *Environ. Pollut.* 115: 359-371.
- Izuta, T. (1998) Ecophysiological responses of Japanese forest tree species to ozone, simulated acid rain and soil acidification. *J. Plant Res.* 111: 471-480.
- Jackson, G. E. (1995) The effect of ozone, nitrogen dioxide and nitric oxide fumigation of cereals on the rose grain aphid *Metopolophium dirhodum*. *Agric. Ecosyst. Environ.* 54: 187-194.
- Jackson, D. M.; Heagle, A. S.; Eckel, R. V. W. (1999) Ovipositional response of tobacco hornworm moths (Lepidoptera: Sphingidae) to tobacco plants grown under elevated levels of ozone. *Environ. Entomol.* 28: 566-571.
- Jackson, D. M.; Ruffy, T. W.; Heagle, A. S.; Severson, R. F.; Eckel, R. W. V. (2000) Survival and development of tobacco hornworm larvae on tobacco plants grown under elevated levels of ozone. *J. Chem. Ecol.* 26: 1-18.
- Jacobson, J. S. (1977) The effects of photochemical oxidants on vegetation. In: Ozon und Begleitsubstanzen im photochemischen Smog: das Kolloquium [Ozone and related substances in photochemical smog: the colloquium]; September 1976; Düsseldorf, Federal Republic of Germany. Düsseldorf, Federal Republic of Germany: VDI-Verlag GmbH; pp. 163-173. (VDI-Berichte nr. 270).
- Jäger, H. J.; Herstein, U.; Fangmeier, A. (1999) The European Stress Physiology and Climate Experiment - Project 1: wheat (ESPACE-wheat); introduction, aims and methodology. *Eur. J. Agron.* 10: 155-162.
- James, R. L.; Cobb, F. W., Jr.; Miller, P. R.; Parmeter, J. R., Jr. (1980) Effects of oxidant air pollution on susceptibility of pine roots to *Fomes annosus*. *Phytopathology* 70: 560-563.
- Janson, R.; Granat, L. (1999) A foliar rinse study of the dry deposition of nitric acid to a coniferous forest. *Agric. For. Meteorol.* 98-99: 683-696.
- Jarvis, P. G. (1976) The interpretation of the variations in leaf water potential and stomatal conductance found in canopies in the field. *Philos. Trans. R. Soc. B* 273: 593-610.
- Jiang, C.-Z.; Rodermel, S. R.; Shibles, R. M. (1993) Photosynthesis, rubisco activity and amount, and their regulation by transcription in senescing soybean leaves. *Plant Physiol.* 101: 105-112.
- Jimenez, A.; Calvo, E.; Martin, C.; Porcuna, J. L.; Sanz, M. J. (2001) Estudio de la interaccion entre el ozono y el desarrollo de las virosis en el cultivo del tomate. *Agricola Vergel* 231: 141-150.
- Johnson, B. G.; Hale, B. A.; Ormrod, D. P. (1996a) Carbon dioxide and ozone effects on growth of a legume-grass mixture. *J. Environ. Qual.* 25: 908-916.
- Johnson, J. D.; Chappelka, A. H.; Hain, F. P.; Heagle, A. S. (1996b) Interactive effects of air pollutants with abiotic and biotic factors on southern pine forests. In: Cox, S.; Mickler, R. A., eds. Impact of air pollutants on southern pine forests. New York, NY: Springer; pp. 281-312. (Ecological studies: v. 118).

- 1 Jones, H. G.; Lacasse, N. L.; Liggett, W. S.; Weatherford, F. (1977) Experimental air exclusion system for field studies
2 of SO₂ effects on crop productivity. Washington, DC: U.S. Environmental Protection Agency; report no.
3 EPA-600/7-77-122.
- 4 Jones, C. G.; Coleman, J. S.; Findlay, S. (1994) Effects of ozone on interactions between plants, consumers and
5 decomposers. In: Alscher, R. G.; Wellburn, A. R., eds. Plant responses to the gaseous environment. Molecular,
6 metabolic and physiological aspects. London, United Kingdom: Chapman and Hall; pp. 339-363.
- 7 Joss, U.; Graber, W. K. (1996) Profiles and simulated exchange of H₂O, O₃, NO₂ between the atmosphere and the
8 HartX Scots pine plantation. Theor. Appl. Climatol. 53: 157-172.
- 9 Kainulainen, P.; Holopainen, J. K.; Hyttinen, H.; Oksanen, J. (1994) Effect of ozone on the biochemistry and aphid
10 infestation of Scots pine. Phytochemistry 35: 39-42.
- 11 Kainulainen, P.; Holopainen, J. K.; Holopainen, T. (2000a) Combined effects of ozone and nitrogen on secondary
12 compounds, amino acids, and aphid performance in Scots pine. J. Environ. Qual. 29: 334-342.
- 13 Kainulainen, P.; Utriainen, J.; Holopainen, J. K.; Oksanen, J.; Holopainen, T. (2000b) Influence of elevated ozone and
14 limited nitrogen availability on conifer seedlings in an open-air fumigation system: effects on growth, nutrient
15 content, mycorrhiza, needle ultrastructure, starch and secondary compounds. Global Change Biol. 6: 345-355.
- 16 Kainulainen, P.; Holopainen, T.; Holopainen, J. K. (2003) Decomposition of secondary compounds from needle litter
17 of Scots pine grown under elevated CO₂ and O₃. Glob. Change Biol. 9: 295-304.
- 18 Kangasjärvi, J.; Talvinen, J.; Utriainen, M.; Karjalainen, R. (1994) Plant defense systems induced by ozone. Plant Cell
19 Environ. 17: 783-794.
- 20 Kärenlampi, L.; Skärby, L. (1996) Critical levels for ozone in Europe: testing and finalizing the concepts. UN-ECE
21 workshop report. In: Proceedings of UN-ECE convention on long-range transboundary air pollution workshop;
22 April; Kuopio, Finland. Kupio, Finland: University of Kuopio, Department of Ecology and Environmental
23 Science.
- 24 Karlsson, P. E.; Medin, E.-L.; Wickström, H.; Selldén, G.; Wallin, G.; Ottosson, S.; Skärby, L. (1995) Ozone and
25 drought stress: interactive effects on the growth and physiology of Norway spruce (*Picea abies* (L.) Karst.).
26 Water Air Soil Pollut. 85: 1325-1330.
- 27 Karlsson, P. E.; Medin, E. L.; Wallin, G.; Selldén, G.; Skärby, L. (1997) Effects of ozone and drought stress on the
28 physiology and growth of two clones of Norway spruce (*Picea abies*). New Phytol. 136: 265-275.
- 29 Karlsson, P. E.; Pleijel, H.; Karlsson, G. P.; Medin, E. L.; Skärby, L. (2000) Simulations of stomatal conductance and
30 ozone uptake to Norway spruce saplings in open-top chambers. Environ. Pollut. 109: 443-451.
- 31 Karlsson, P. E.; Medin, E. L.; Selldén, G.; Wallin, G.; Ottosson, S.; Pleijel, H.; Skärby, L. (2002) Impact of ozone and
32 reduced water supply on the biomass accumulation of Norway spruce saplings. Environ. Pollut. 19: 237-244.
- 33 Karlsson, P. E.; Selldén, G.; Pleijel, H. (2003) Establishing ozone critical levels II. UNECE workshop report.
34 Gothenburg, Sweden: IVL Swedish Environmental Research Institute; IVL report B 1523 A,B,C. Available:
35 <http://www.ozoneworkshop.ivl.se/wsreport.htm> [7 July, 2004].
- 36 Karlsson, G. P.; Karlsson, P. E.; Danielsson, H.; Pleijel, H. (2003) Clover as a tool for bioindication of phytotoxic
37 ozone—5 years of experience from southern Sweden—consequences for the short-term critical levels. Sci. Total
38 Environ. 301: 205-213.
- 39 Karlsson, P. E.; Uddling, J.; Skärby, L.; Wallin, G.; Selldén, G. (2003) Impact of ozone on the growth of birch (*Betula*
40 *pendula*) saplings. Environ. Pollut. 124: 485-495.
- 41 Karnosky, D. F. (1981) Changes in eastern white pine stands related to air pollution stress. Mitt. Forstl.
42 Bundesversuchsanst. Wien 137: 41-45.
- 43 Karnosky, D. F. (2003) Impacts of elevated atmospheric CO₂ on forest trees and forest ecosystems: knowledge gaps.
44 Environ. Int. 29: 161-169.
- 45 Karnosky, D. F.; Gagnon, Z. E.; Dickson, R. E.; Coleman, M. D.; Lee, E. H.; Isebrands, J. G. (1996) Changes in
46 growth, leaf abscission, biomass associated with seasonal tropospheric ozone exposures of *Populus tremuloides*
47 clones and seedlings. Can. J. For. Res. 26: 23-37.
- 48 Karnosky, D. F.; Podila, G. K.; Gagnon, Z.; Pechter, P.; Akkapeddi, A.; Coleman, M.; Dickson, R. E.; Isebrands, J. G.
49 (1998) Genetic control of responses to interacting O₃ and CO₂ in *Populus tremuloides*. Chemosphere 36:
50 807-812.
- 51 Karnosky, D. F.; Mankovska, B.; Percy, K.; Dickson, R. E.; Podila, G. K.; Sober, J.; Noormets, A.; Hendrey, G.;
52 Coleman, M. D.; Kubiske, M.; Pregitzer, K. S.; Isebrands, J. G. (1999) Effects of tropospheric ozone on

- trembling aspen and interaction with CO₂: results from an O₃-gradient and a FACE experiment. *Water Air Soil Pollut.* 116: 311-322.
- Karnosky, D. F.; Ceulemans, R.; Scarascia-Mugnozza, G. E.; Ines, J. L., eds. (2001) The impact of carbon dioxide and other greenhouse gases on forest ecosystems: report no. 3 of the IUFRO Task Force on Environmental Change. Wallingford, United Kingdom: CABI Publishing. (IUFRO research series no. 8). Available: <http://www.cabi-publishing.org/Bookshop/Readingroom/0851995519.asp> [19 August, 2003].
- Karnosky, D. F.; Gielen, B.; Ceulemans, R.; Schlesinger, W. H.; Norby, R. J.; Oksanen, E.; Matyssek, R.; Hendrey, G. R. (2001a) FACE systems for studying the impacts of greenhouse gases on forest ecosystems. In: Karnosky, D. F.; Ceulemans, R.; Scarascia-Mugnozza, G. E.; Innes, J. L. eds. The impact of carbon dioxide and other greenhouse gases on forest ecosystems; report no. 3 of the IUFRO Task Force on Environmental Change. Wallingford, UK: CABI Publishers, pp. 297-324. Available: <http://www.cabi-publishing.org/Bookshop/Readingroom/0851995519.asp> [16 April, 2004].
- Karnosky, D. F.; Oksanen, E.; Dickson, R. E.; Isebrands, J. G. (2001b) Impact of interacting greenhouse gases on forest ecosystems. In: Karnosky, D. F.; Ceulemans, R.; Scarascia-Mugnozza, G. E.; Innes, J. L., eds. The impact of carbon dioxide and other greenhouse gases of forest ecosystems: report no. 3 of the IUFRO Task Force on Environmental Change. Wallingford, United Kingdom: CABI Publishing; pp. 253-267. (IUFRO research series no. 8).
- Karnosky, D. F.; Percy, K. E.; Xiang, B.; Callan, B.; Noormets, A.; Mankovska, B.; Hopkin, A.; Sober, J.; Jones, W.; Dickson, R. E.; Isebrands, J. G. (2002) Interacting elevated CO₂ and tropospheric O₃ predisposes aspen (*Populus tremuloides* Michx.) to infection by rust (*Melampsora medusae* f. sp. *tremuloidae*). *Global Change Biol.* 8: 329-338.
- Karnosky, D. F.; Zak, D. R.; Pregitzer, K. S.; Awmack, C. S.; Bockheim, J. G.; Dickson, R. E.; Hendrey, G. R.; Host, G. E.; King, J. S.; Kopper, B. J.; Kruger, E. L.; Kubiske, M. E.; Lindroth, R. L.; Mattson, W. J.; McDonald, E. P.; Noormets, A.; Oksanen, E.; Parsons, W. F. J.; Percy, K. E.; Podila, G. K.; Riemenschneider, D. E.; Sharma, P.; Thakur, R.; Söber, A.; Söber, J.; Jones, W. S.; Anttonen, S.; Vapaavuori, E.; Mankovska, B.; Heilman, W.; Isebrands, J. G. (2003a) Tropospheric O₃ moderates responses of temperate hardwood forests to elevated CO₂: A synthesis of molecular to ecosystem results from the Aspen FACE project. *Funct. Ecol.* 17: 289-304.
- Karnosky, D. F.; Percy, K.; Mankovska, B.; Pritchard, T.; Noormets, A.; Dickson, R. E.; Jepsen, E.; Isebrands, J. G. (2003b) Ozone affects the fitness of trembling aspen. In: Karnosky, D. F.; Percy, K. E.; Chappelka, A. H.; Simpson, C. J.; Pikkariainen, J., eds. Air pollution, global change and forests in the new millenium. Oxford, United Kingdom: Elsevier Science. (Developments in Environmental Science, 3).
- Karpinski, S.; Wingsle, G.; Karpinska, B.; Hällgren, J.-E. (1993) Molecular response to photooxidative stress in *Pinus sylvestris* (L.): II. Differential expression of Cu-Zn superoxide dismutases and glutathione reductase. *Plant Physiol.* 103: 1385-1391.
- Kasurinen, A. E.; Helmisaari, H.-S.; Holopainen, T. (1999) The influence of elevated CO₂ and O₃ on fine roots and mycorrhizas of naturally growing young Scots pine trees during three exposure years. *Global Change Biol.* 5: 771-780.
- Kats, G.; Thompson, C. R.; Kuby, W. C. (1976) Improved ventilation of open top greenhouses. *J. Air Pollut. Control Assoc.* 26: 1089-1090.
- Kats, G.; Dawson, P. J.; Bytnerowicz, A.; Wolf, J. W.; Thompson, C. R.; Olszyk, D. M. (1985) Effects of ozone or sulfur dioxide on growth and yield of rice. *Agric. Ecosyst. Environ.* 14: 103-117.
- Kellomäki, S.; Wang, K.-Y. (1997a) Effects of elevated O₃ and CO₂ concentrations on photosynthesis and stomatal conductance in Scots pine. *Plant Cell Environ.* 20: 995-1006.
- Kellomäki, S.; Wang, K.-Y. (1997b) Effects of elevated O₃ and CO₂ on chlorophyll fluorescence and gas exchange in Scots pine during the third growing season. *Environ. Pollut.* 97: 17-27.
- Kelting, D. L.; Burger, J. A.; Edwards, G. S. (1995) The effects of ozone on the root dynamics of seedlings and mature red oak (*Quercus rubra* L.). *For. Ecol. Manage.* 79: 197-206.
- Kerstiens, G.; Lendzian, K. J. (1989) Interaction between ozone and plant cuticles: I. ozone deposition and permeability. *New Phytol.* 112: 13-19.
- Khan, M. R.; Khan, M. W. (1994) Single and interactive effects of O₃ and SO₂ on tomato. *Environ. Exp. Bot.* 34: 461-469.
- Khan, M. R.; Khan, M. W. (1997) Effects of the root-knot nematode, *Meloidogyne incognita*, on the sensitivity of tomato to sulfur dioxide and ozone. *Environ. Exp. Bot.* 38: 117-130.

- 1 Khan, M. R.; Khan, M. W. (1998a) Interactive effects of ozone and powdery mildew (*Sphaerotheca fuliginea*) on
2 bottle gourd (*Lagenaria siceraria*). Agric. Ecosyst. Environ. 70: 109-118.
- 3 Khan, M. R.; Khan, M. W. (1998b) Interactive effects of ozone and root-knot nematode on tomato. Agric. Ecosyst.
4 Environ. 70: 97-107.
- 5 Khan, M. R.; Khan, M. W. (1999) Effects of intermittent ozone exposures on powdery mildew of cucumber. Environ.
6 Exp. Bot. 42: 163-171.
- 7 Kickert, R. N.; Krupa, S. V. (1991) Modeling plant response to tropospheric ozone: a critical review. Environ. Pollut.
8 70: 271-383.
- 9 Kickert, R. N.; Tonella, G.; Simonov, A.; Krupa, S. V. (1999) Predictive modeling of effects under global change.
10 Environ. Pollut. 100: 87-132.
- 11 Kieliszewska-Rokicka, B.; Rudawska, M.; Leski, T. (1998) Ectomycorrhizae of young and mature Scots pine trees in
12 industrial regions in Poland. In: Bytnerowicz, A.; Arbaugh, M. J.; Schilling, S. L., eds. Proceedings of the
13 International Symposium on Air Pollution and Climate Change Effects on Forest Ecosystems; February, 1996;
14 Riverside, CA. Albany, CA: USDA Forest Service, Pacific Southwest General Technical Report no. PSW-GTR
15 166; pp. 205-214.
- 16 Kim, H. J.; Helfand, G. E.; Howitt, R. E. (1998) An economic analysis of ozone control in California's San Joaquin
17 Valley. J. Agric. Resource Econ. 23: 55-70.
- 18 Kim, J. S.; Chappelka, A. H.; Miller-Goodman, M. S. (1998) Decomposition of blackberry and broomsedge bluestem
19 as influenced by ozone. J. Environ. Qual. 27: 953-960.
- 20 King, J. S.; Pregitzer, K. S.; Zak, D. R.; Sober, J.; Isebrands, J. G.; Dickson, R. E.; Hendrey, G. R.; Karnosky, D. F.
21 (2001) Fine-root biomass and fluxes of soil carbon in young stands of paper birch and trembling aspen as
22 affected by elevated atmospheric CO₂ and tropospheric O₃. Oecologia 128: 237-250.
- 23 Kleier, C.; Farnsworth, B.; Winner, W. (1998) Biomass, reproductive output, and physiological responses of
24 rapid-cycling Brassica (*Brassica rapa*) to ozone and modified root temperature. New Phytol. 139: 657-664.
- 25 Kleier, C.; Farnsworth, B.; Winner, W. (2001) Photosynthesis and biomass allocation of radish cv. "Cherry Belle" in
26 response to root temperature and ozone. Environ. Pollut. 111: 127-133.
- 27 Kliebenstein, D. J.; Monde, R.-A.; Last, R. L. (1998) Superoxide dismutase in Arabidopsis: an eclectic enzyme family
28 with disparate regulation and protein localization. Plant Physiol. 118: 637-650.
- 29 Kobayashi, K.; Okada, M.; Nouchi, I. (1994) A chamber system for exposing rice (*Oryza sativa* L.) to ozone in a
30 paddy field. New Phytol. 126: 317-325.
- 31 Kobayashi, K.; Okada, M.; Nouchi, I. (1995) Effects of ozone on dry matter partitioning and yield of Japanese
32 cultivars of rice (*Oryza sativa* L.). Agric. Ecosyst. Environ. 53: 109-122.
- 33 Kohut, R. J.; Laurence, J. A.; Amundson, R. G. (1988) Effects of ozone and sulfur dioxide on yield of red clover and
34 timothy. J. Environ. Qual. 17: 580-585.
- 35 Kohut, R. J.; Laurence, J. A.; King, P.; Raba, R. (1997) Identification of bioindicator species for ozone and assessment
36 of the responses to ozone of native vegetation at Acadia National Park. Denver CO: National Park Service; Air
37 Resources Division; NPS technical report Acadia D-175.
- 38 Kolb, T. E.; Matyssek, R. (2001) Limitations and perspectives about scaling ozone impacts in trees. Environ. Pollut.
39 115: 373-393.
- 40 Kolb, T. E.; Fredericksen, T. S.; Steiner, K. C.; Skelly, J. M. (1997) Issues in scaling tree size and age responses to
41 ozone: a review. Environ. Pollut. 98: 195-208.
- 42 Köllner, B.; Krause, G. H. M. (2003) Effects of two different ozone exposure regimes on chlorophyll and sucrose
43 content of leaves and yield parameters of sugar beet (*Beta Vulgaris* L.) and rape (*Brassica Napus* L.). Water Air
44 Soil Pollut. 144: 317-332.
- 45 Kolomiets, M. V.; Chen, H.; Gladon, R. J.; Braun, E. J.; Hannapel, D. J. (2000) A leaf lipoxygenase of potato induced
46 specifically by pathogen infection. Plant Physiol. 124: 1121-1130.
- 47 Kopp, R. J.; Vaughn, W. J.; Hazilla, M.; Carson, R. (1985) Implications of environmental policy for U.S. agriculture:
48 the case of ambient ozone standards. J. Environ. Manage. 20: 321-331.
- 49 Kopper, B. J.; Lindroth, R. L. (2003) Responses of trembling aspen (*Populus tremuloides*) phytochemistry and aspen
50 blotch leafminer (*Phyllonorycter tremuloidiella*) performance to elevated levels of atmospheric CO₂ and O₃.
51 Agric. For. Entomol. 5: 17-26.
- 52 Kopper, B. J.; Lindroth, R. L. (2003) Effects of elevated carbon dioxide and ozone on the phytochemistry of aspen and
53 performance of an herbivore. Oecologia 134: 95-103.

- 1 Kostka-Rick, R.; Manning, W. J. (1992) Partitioning of biomass and carbohydrates in field-grown radish under
2 ambient concentrations of ozone, and treated with the anti-ozonant ethylene-diurea (EDU). *New Phytol.* 121:
3 187-200.
- 4 Kostka-Rick, R.; Manning, W. J. (1993) Dose-response studies with the antiozonant ethylenediurea (EDU), applied as
5 a soil drench to two growth substrates, on greenhouse-grown varieties of *Phaseolus vulgaris* L. *Environ. Pollut.*
6 82: 63-72.
- 7 Kostka-Rick, R.; Manning, W. J. (1993) Dynamics of growth and biomass partitioning in field-grown bush bean
8 (*Phaseolus vulgaris* L.), treated with the antiozonant ethylenediurea (EDU). *Agric. Ecosyst. Environ.* 47:
9 195-214.
- 10 Kostka-Rick, R.; Manning, W. J.; Buonaccorsi, J. P. (1993) Dynamics of biomass partitioning in field-grown radish
11 varieties, treated with ethylenediurea. *Environ. Pollut.* 80: 133-145.
- 12 Kouterick, K. B.; Skelly, J. M.; Fredericksen, T. S.; Steiner, K. C.; Kolb, T. E.; Ferdinand, J. A. (2000) Foliar injury,
13 leaf gas exchange and biomass responses of black cherry (*Prunus serotina* Ehrh.) half-sibling families to ozone
14 exposure. *Environ. Pollut.* 107: 117-126.
- 15 Koziol, M. J.; Whitley, F. R. (1984) Gaseous air pollutants and plant metabolism. London, United Kingdom:
16 Butterworths Scientific.
- 17 Kozlowski, T. T.; Kramer, P. J.; Pallardy, S. G. (1991) The physiological ecology of woody plants. San Diego, CA:
18 Academic Press, Inc.; pp. 338-375. (Mooney, H. A., ed. *Physiological ecology: a series of monographs, texts,*
19 *and treatises*).
- 20 Kraft, M.; Weigel, H.-J.; Mejer, G.-J.; Brandes, F. (1996) Reflectance measurements of leaves for detecting visible and
21 non-visible ozone damage to crops. *J. Plant Physiol.* 148: 148-154.
- 22 Krapp, A.; Hofmann, B.; Schäfer, C.; Stitt, M. (1993) Regulation of the expression of *rbcS* and other photosynthetic
23 genes by carbohydrates: a mechanism for the "sink regulation" of photosynthesis? *Plant J.* 3: 817-828.
- 24 Kress, L. W.; Miller, J. E.; Smith, H. J. (1985) Impact of ozone on winter wheat yield. *Environ. Exp. Bot.* 25: 211-228.
- 25 Krupa, S. V.; Nosal, M.; Legge, A. H. (1994) Ambient ozone and crop loss: establishing a cause-effect relationship.
26 *Environ. Pollut.* 83: 269-276.
- 27 Krupa, S. V.; Grünhage, L.; Jäger, H.-J.; Nosal, M.; Manning, W. J.; Legge, A. H.; Hanewald, K. (1995) Ambient
28 ozone (O₃) and adverse crop response: a unified view of cause and effect. *Environ. Pollut.* 87: 119-126.
- 29 Krupa, S. V.; Jäger, H.-J. (1996) Adverse effects of elevated levels of ultraviolet (UV)-B radiation and ozone (O₃) on
30 crop growth and productivity. In: Bazzaz, F.; Sombroek, W., eds. *Global climate change and agricultural*
31 *production*. Chichester, United Kingdom: John Wiley & Sons; pp. 141-169. Available:
32 <http://www.fao.org/docrep/W5183E/w5183e00.htm> [28 August, 2003].
- 33 Krupa, S. V.; Kickert, R. N.; Jäger, H.-J. (1998) Elevated ultraviolet (UV)-B radiation and agriculture. New York, NY:
34 Springer-Verlag Environmental Intelligence Unit.
- 35 Krupa, S. V.; Groth, J. V. (2000) Global climate change and crop responses: uncertainties associated with the current
36 methodologies. In: Agrawal, S. B.; Agrawal, M., eds. *Environmental pollution and plant responses*. Boca Raton,
37 FL: Lewis Publishers; pp. 1-18.
- 38 Krupa, S. V.; Legge, A. H. (2000) Passive sampling of ambient, gaseous air pollutants: an assessment from an
39 ecological perspective. *Environ. Pollut.* 107: 31-45.
- 40 Krupa, S. V.; Nosal, M.; Peterson, D. L. (2001) Use of passive ozone O₃ samplers in vegetation effects assessment.
41 *Environ. Pollut.* 112: 303-309.
- 42 Krupa, S. V.; Nosal, M.; Ferdinand, J. A.; Stevenson, R. E.; Skelly, J. M. (2003) A multi-variate statistical model
43 integrating passive sampler and meteorology data to predict the frequency distributions of hourly ambient ozone
44 (O₃) concentrations. *Environ. Pollut.* 124: 173-178.
- 45 Kuehler, E. A.; Flagler, R. B. (1999) The effects of sodium erythorbate and ethylenediurea on photosynthetic function
46 of ozone-exposed loblolly pine seedlings. *Environ. Pollut.* 105: 25-35.
- 47 Kuik, O. J.; Helming, J. F. M.; Dorland, C.; Spaninks, F. A. (2000) The economic benefits to agriculture of a reduction
48 of low-level ozone pollution in The Netherlands. *Eur. Rev. Agric Econ.* 27: 75-90.
- 49 Kull, O.; Tulva, I.; Vapaavuori, E. (2003) Influence of elevated CO₂ and O₃ on *Betula pendula* Roth crown structure.
50 *Ann. Bot.* 91: 559-569.
- 51 Kurezyńska, E. U.; Dmuchowski, W.; Wloch, W.; Bytnerowicz, A. (1997) The influence of air pollutants on needles
52 and stems of scots pine (*Pinus Sylvestris* L.) trees. *Environ. Pollut.* 98: 325-334.
- 53 Kurpius, M. R.; McKay, J.; Goldstein, A. H. (2002) Annual ozone deposition to a Sierra Nevada ponderosa pine
54 plantation. *Atmos. Environ.* 36: 4503-4515.

- 1 Kytöviita, M.-M.; Pelloux, J.; Fontaine, V.; Botton, B.; Dizengremel, P. (1999) Elevated CO₂ does not ameliorate
2 effects of ozone on carbon allocation in *Pinus halepensis* and *Betula pendula* in symbiosis with *Paxillus*
3 *involutus*. *Physiol. Plant.* 106: 370-377.
- 4 Kytöviita, M.-M.; Le Thiec, D.; Dizengremel, P. (2001) Elevated CO₂ and ozone reduce nitrogen acquisition by *Pinus*
5 *halepensis* from its mycorrhizal symbiont. *Physiol. Plant.* 111: 305-312.
- 6 Lagrimini, L. M.; Rothstein, S. (1987) Tissue specificity of tobacco peroxidase isozymes and their induction by
7 wounding and tobacco mosaic virus infection. *Plant Physiol.* 84: 438-442.
- 8 Laisk, A.; Kull, O.; Moldau, H. (1989) Ozone concentration in leaf intercellular air spaces is close to zero. *Plant*
9 *Physiol.* 90: 1163-1167.
- 10 Lamb, C.; Dixon, R. A. (1997) The oxidative burst in plant disease resistance. *Ann. Rev. Plant Physiol. Mol. Biol.* 48:
11 251-275.
- 12 Landolt, W.; Günthardt-Goerg, M. S.; Pfenninger, I.; Einig, W.; Hampp, R.; Maurer, S.; Matyssek, R. (1997) Effect of
13 fertilization on ozone-induced changes in the metabolism of birch (*Betula pendula*) leaves. *New Phytol.* 137:
14 389-397.
- 15 Landolt, W.; Bühlmann, U.; Bleuler, P.; Bucher, J. B. (2000) Ozone exposure-response relationships for biomass and
16 root/shoot ratio of beech (*Fagus sylvatica*), ash (*Fraxinus excelsior*), Norway spruce (*Picea abies*) and scots
17 pine (*Pinus sylvestris*). *Environ. Pollut.* 109: 473-478.
- 18 Langebartels, C.; Kerner, K.; Leonardi, S.; Schraudner, M.; Trost, M.; Heller, W.; Sandermann, H., Jr. (1991)
19 Biochemical plant responses to ozone: I. differential induction of polyamine and ethylene biosynthesis in
20 tobacco. *Plant Physiol.* 95: 882-889.
- 21 Langebartels, C.; Ernest, D.; Heller, W.; Lutz, C.; Payer, H.-D.; Sandermann, J. (1997) Ozone responses of trees:
22 results from controlled chamber exposures at the GSF phytotron. In: Sandermann, J. H.; Wellburn, A. R.; Heath,
23 R. L., eds. *Forest decline and ozone: a comparison of controlled chamber and field experiments*. Berlin,
24 Germany: Springer-Verlag; pp. 163-200. (Ecological studies: v. 127).
- 25 Larsen, R. I.; Heagle, A. S.; Heck, W. W. (1983) An air quality data analysis system for interrelating effects, standards,
26 and needed source reductions: part 7. an O₃-SO₂ leaf injury mathematical model. *J. Air Pollut. Control Assoc.*
27 33: 198-207.
- 28 Larsen, R. I.; Heck, W. W. (1984) An air quality data analysis system for interrelating effects, standards, and needed
29 source reductions: part 8. an effective mean O₃ crop reduction mathematical model. *J. Air Pollut. Control Assoc.*
30 34: 1023-1034.
- 31 Larson, T. V.; Vong, R. J. (1990) A theoretical investigation of the pressure and temperature dependence of
32 atmospheric ozone deposition to trees. *Environ. Pollut.* 67: 179-189.
- 33 Larson, J. L.; Zak, D. R.; Sinsabaugh, R. L. (2002) Extracellular enzyme activity beneath temperate trees growing
34 under elevated carbon dioxide and ozone. *Soil Sci. Soc. Am. J.* 66: 1848-1856.
- 35 Laurence, J. A.; Kohut, R. J.; Amundson, R. G.; Weinstein, D. A.; MacLean, D. C. (1996) Response of sugar maple to
36 multiple year exposure to ozone and simulated acidic precipitation. *Environ. Pollut.* 92: 119-126.
- 37 Laurence, J. A.; Amundson, R. G.; Kohut, R. J.; Weinstein, D. A. (1997) Growth and water use of red spruce (*Picea*
38 *rubens* Sarg.) exposed to ozone and simulated acidic precipitation for four growing seasons. *For. Sci.* 43:
39 355-361.
- 40 Laurence, J. A.; Retzlaff, W. A.; Kern, J. S.; Lee, E. H.; Hogsett, W. E.; Weinstein, D. A. (2001) Predicting the
41 regional impact of ozone and precipitation on the growth of loblolly pine and yellow poplar using linked
42 TREGRO and ZELG models. *For. Ecol. Manage.* 146: 247-263.
- 43 Law, M. Y.; Charles, S. A.; Halliwell, B. (1983) Glutathione and ascorbic acid in spinach (*Spinacia oleracea*)
44 chloroplasts: the effect of hydrogen peroxide and of Paraquat. *Biochem. J.* 210: 899-903.
- 45 Lawlor, D. W. (1998) Plant responses to global change: temperature and drought stress. In: De Kok, L. J.; Stulen, I.,
46 eds. *Responses of plant metabolism to air pollution and global change*. Leiden, The Netherlands: Backhuys
47 Publishers; pp. 193-207.
- 48 Lawlor, D. W.; Keys, A. J. (1993) Understanding photosynthetic adaptation to changing climate. In: Fowden, L.;
49 Mansfield, T.; Stoddart, J., eds. *Plant adaptation to environmental stress*; London, United Kingdom: Chapman
50 and Hall; pp. 85-106.
- 51 Lawson, T.; Craigan, J.; Black, C. R.; Colls, J. J.; Tulloch, A.-M.; Landon, G. (2001a) Effects of elevated carbon
52 dioxide and ozone on the growth and yield of potatoes (*Solanum tuberosum*) grown in open-top chambers.
53 *Environ. Pollut.* 111: 479-491.

- 1 Lawson, T.; Craigon, J.; Tulloch, A.-M.; Black, C. R.; Colls, J. J.; Landon, G. (2001b) Photosynthetic responses to
2 elevated CO₂ and O₃ in field-grown potato (*Solanum tuberosum*). J. Plant Physiol. 158: 309-323.
- 3 Lawson, T.; Craigon, J.; Black, C. R.; Colls, J. J.; Landon, G.; Weyers, J. D. B. (2002) Impact of elevated CO₂ and O₃
4 on gas exchange parameters and epidermal characteristics in potato (*Solanum tuberosum* L.). J. Exp. Bot. 53:
5 737-746.
- 6 Le Thiec, D.; Manninen, S. (2003) Ozone and water deficit reduced growth of Aleppo pine seedlings. Plant Physiol.
7 Biochem. 41: 55-63.
- 8 Lee, E. H.; Bennett, J. H. (1982) Superoxide dismutase: a possible protective enzyme against ozone injury in snap
9 beans (*Phaseolus vulgaris* L.). Plant Physiol. 69: 1444-1449.
- 10 Lee, E. H.; Hogsett, W. E. (1999) Role of concentrations and time of day in developing ozone exposure indices for a
11 secondary standard. J. Air Waste Manage. Assoc. 49: 669-681.
- 12 Lee, J. J.; Lewis, R. A.; Body, D. E. (1975) A field experimental system for the evaluation of the bioenvironmental
13 effects of sulfur dioxide. In: Proceedings of the Fort Union Coal Field Symposium; April; Billings, MT.
14 Billings, MT: Montana Academy of Science, v. 5, pp. 608-620.
- 15 Lee, E. H.; Tingey, D. T.; Hogsett, W. E. (1987) Selection of the best exposure-response model using various 7-hour
16 ozone exposure statistics. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air
17 Quality Planning and Standards.
- 18 Lee, E. H.; Tingey, D. T.; Hogsett, W. E. (1988) Evaluation of ozone exposure indices in exposure-response modeling.
19 Environ. Pollut. 53: 43-62.
- 20 Lee, E. H.; Tingey, D. T.; Hogsett, W. E. (1989) Interrelation of experimental exposure and ambient air quality data
21 for comparison of ozone exposure indices and estimating agricultural losses. Corvallis, OR: U.S. Environmental
22 Protection Agency, Environmental Research Laboratory; EPA report no. EPA-600/3-89-047. Available from:
23 NTIS, Springfield, VA; PB89-195036.
- 24 Lee, E. H.; Hogsett, W. E.; Tingey, D. T. (1994a) Alternative attainment criteria for a secondary federal standard for
25 ozone. In: Nriagu, J. O.; Simmons, M. S., eds. Environmental oxidants. New York, NY: John Wiley & Sons;
26 pp. 549-584. (Nriagu, J. O., ed. Advances in environmental sciences and technology series: v. 28).
- 27 Lee, E. H.; Hogsett, W. E.; Tingey, D. T. (1994b) Attainment and effects issues regarding alternative secondary ozone
28 air quality standards. J. Environ. Qual. 23: 1129-1140.
- 29 Lee, J. C.; Skelly, J. M.; Steiner, K. C.; Zhang, J. W.; Savage, J. E. (1999) Foliar response of black cherry (*Prunus*
30 *serotina*) clones to ambient ozone exposure in central Pennsylvania. Environ. Pollut. 105: 325-331.
- 31 Lee, J. C.; Steiner, K. C.; Zhang, J. W.; Skelly, J. M. (2002) Heritability of ozone sensitivity in open-pollinated
32 families of black cherry (*Prunus serotina* Ehrh.). For. Sci. 48: 111-117.
- 33 Lee, E. H.; Tingey, D. T.; Hogsett, W. E.; Laurence, J. A. (2003) History of tropospheric ozone for the San Bernardino
34 Mountains of southern California, 1963-1999. Atmos. Environ. 37: 2705-2717.
- 35 Lefohn, A. S. (1984) A comparison of ambient ozone exposures for selected nonurban sites. Presented at: 77th annual
36 meeting of the Air Pollution Control Association; June; San Francisco, CA. Pittsburgh, PA: Air Pollution
37 Control Association; paper no. 84-104.1.
- 38 Lefohn, A. S.; Benedict, H. M. (1982) Development of a mathematical index that describes ozone concentration,
39 frequency and duration. Atmos. Environ. 16: 2529-2532.
- 40 Lefohn, A. S.; Foley, J. K. (1992) NCLAN results and their application to the standard-setting process: protecting
41 vegetation from surface ozone exposures. J. Air Waste Manage. Assoc. 42: 1046-1052.
- 42 Lefohn, A. S.; Irving, P. M. (1988) Characterizing ambient ozone exposure regimes in agricultural areas. Presented at:
43 81st annual meeting of the Air Pollution Control Association; June; Dallas, TX. Pittsburgh, PA: Air Pollution
44 Control Association; 88-69.1.
- 45 Lefohn, A. S.; Jones, C. K. (1986) The characterization of ozone and sulfur dioxide air quality data for assessing
46 possible vegetation effects. J. Air Pollut. Control Assoc. 36: 1123-1129.
- 47 Lefohn, A. S.; Runeckles, V. C. (1987) Establishing standards to protect vegetation - ozone exposure/dose
48 considerations. Atmos. Environ. 21: 561-568.
- 49 Lefohn, A. S.; Shadwick, D. S. (2000) Differences in trending estimates in the United States using several ozone
50 metrics. Presented at: 93rd annual meeting of the Air & Waste Management Association; June; Salt Lake City,
51 UT. Pittsburgh, PA: Air & Waste Management Association.
- 52 Lefohn, A. S.; Tingey, D. T. (1985) Comments on "Injury response of *Phaseolus vulgaris* to ozone flux density."
53 Atmos. Environ. 19: 206-207.

- 1 Lefohn, A. S.; Hogsett, W. E.; Tingey, D. T. (1986) A method for developing ozone exposures that mimic ambient
2 conditions in agricultural areas. *Atmos. Environ.* 20: 361-366.
- 3 Lefohn, A. S.; Davis, C. E.; Jones, C. K.; Tingey, D. T.; Hogsett, W. E. (1987) Co-occurrence patterns of gaseous air
4 pollutant pairs at different minimum concentrations in the United States. *Atmos. Environ.* 21: 2435-2444.
- 5 Lefohn, A. S.; Laurence, J. A.; Kohut, R. J. (1988a) A comparison of indices that describe the relationship between
6 exposure to ozone and reduction in the yield of agricultural crops. *Atmos. Environ.* 22: 1229-1240.
- 7 Lefohn, A. S.; Laurence, J. A.; Kohut, R. J. (1988b) A comparison of indices that describe the relationship between
8 exposure to ozone and reduction in the yield of agricultural crops [response to comments by V. C. Runeckles].
9 *Atmos. Environ.* 22: 1242-1243.
- 10 Lefohn, A. S.; Runeckles, V. C.; Krupa, S. V.; Shadwick, D. S. (1989) Important considerations for establishing a
11 secondary ozone standard to protect vegetation. *JAPCA* 39: 1039-1045.
- 12 Lefohn, A. S.; Shadwick, D. S.; Mohnen, V. A. (1990) The characterization of ozone concentrations at a select set of
13 high-elevation sites in the eastern United States. *Environ. Pollut.* 67: 147-178.
- 14 Lefohn, A. S.; Shadwick, D. S.; Somerville, M. C.; Chappelka, A. H.; Lockaby, B. G.; Meldahl, R. S. (1992) The
15 characterization and comparison of ozone exposure indices used in assessing the response of loblolly pine to
16 ozone. *Atmos. Environ. Part A* 26: 287-298.
- 17 Lefohn, A. S.; Jackson, W.; Shadwick, D. S.; Knudsen, H. P. (1997) Effect of surface ozone exposures on vegetation
18 grown in the southern Appalachian Mountains: identification of possible areas of concern. *Atmos. Environ.* 31:
19 1695-1708.
- 20 Lefohn, A. S.; Shadwick, D. S.; Ziman, S. D. (1998) The difficult challenge of attaining EPA's new ozone standard.
21 *Environ. Sci. Technol.* 32: 276A-282A.
- 22 Lefohn, A. S.; Oltmans, S. J.; Dann, T.; Singh, H. B. (2001) Present-day variability of background ozone in the lower
23 troposphere. *J. Geophys. Res. [Atmos.]* 106: 9945-9958.
- 24 Legge, A. H.; Grünhage, L.; Nosal, M.; Jäger, H.-J.; Krupa, S. V. (1995) Ambient ozone and adverse crop response: an
25 evaluation of North American and European data as they relate to exposure indices and critical levels. *Angew.*
26 *Bot.* 69: 192-205.
- 27 Lenzian, K. J.; Kerstiens, G. (1991) Sorption and transport of gases and vapors in plant cuticles. *Rev. Environ.*
28 *Contam. Toxicol.* 121: 65-128.
- 29 León, J.; Lawton, M. A.; Raskin, I. (1995) Hydrogen peroxide synthesis stimulates salicylic acid biosynthesis in
30 tobacco. *Plant Physiol.* 108: 1673-1683.
- 31 Lesser, V. M.; Rawlings, J. O.; Spruill, S. E.; Somerville, M. C. (1990) Ozone effects on agricultural crops: statistical
32 methodologies and estimated dose-response relationships. *Crop Sci.* 30: 148-155.
- 33 Levitt, J. (1972) Responses of plants to environmental stress. New York, NY: Academic Press.
- 34 Lichtenthaler, H. K. (1977) Regulation of prenylquinone synthesis in higher plants. In: Tevini, M.; Lichtenthaler, H.
35 K., eds. *Lipids and lipid polymers in higher plants*. New York, NY: Springer-Verlag; pp. 231-258.
- 36 Lindroth, R. L.; Reich, P. B.; Tjoelker, M. G.; Volin, J. C.; Oleksyn, J. (1993) Light environment alters response to
37 ozone stress in seedlings of *Acer saccharum* March, and hybrid *Populus* L. III. Consequences for performance
38 of gypsy moth. *New Phytol.* 124: 647-654.
- 39 Lindroth, R. L.; Kopper, B. J.; Parsons, W. F. J.; Bockheim, J. G.; Karnosky, D. F.; Hendrey, G. R.; Pregitzer, K. S.;
40 Isebrands, J. G.; Sober, J. (2001) Consequences of elevated carbon dioxide and ozone for foliar chemical
41 composition and dynamics in trembling aspen (*Populus tremuloides*) and paper birch (*Betula papyrifera*).
42 *Environ. Pollut.* 115: 395-404.
- 43 Liu, L.; Eriksson, K.-E. L.; Dean, J. F. D. (1995) Localization of hydrogen peroxide production in *Pisum sativum* L.
44 using epi-polarization microscopy to follow cerium perhydroxide disposition. *Plant Physiol.* 107: 501-506.
- 45 Lloyd, A. C.; Lents, J. M.; Green, C.; Nemeth, P. (1989) Air quality management in Los Angeles: perspectives on past
46 and future emission control strategies. *JAPCA* 39: 696-703.
- 47 Loats, K. V.; Rebbeck, J. (1999) Interactive effects of ozone and elevated carbon dioxide on the growth and
48 physiology of black cherry, green ash, and yellow poplar seedlings. *Environ. Pollut.* 106: 237-248.
- 49 Logan, J. A. (1989) Ozone in rural areas of the United States. *J. Geophys. Res. [Atmos.]* 94: 8511-8532.
- 50 Long, S. P. (1991) Modification of the response of photosynthetic productivity to rising temperature by atmospheric
51 CO₂ concentrations: has its importance been underestimated? *Plant Cell Environ.* 14: 729-739.
- 52 Longstreth, J. D.; De Gruijl; Kripke, M. L.; Takizawa, Y.; van der Leun, J. C. (1995) Effects of increased solar
53 ultraviolet radiation on human health. *Ambio* 24: 153-165.

- 1 Lorenzini, G.; Nali, C. (1995) Analysis of vertical ozone and nitrogen oxides profiles in a *Prunus cerasifera* canopy.
2 Int. J. Biometeorol. 39: 1-4.
- 3 Lorenzini, G.; Medeghini Bonatti, P.; Nali, C.; Baroni Fornasiero, R. (1994) The protective effect of rust infection
4 against ozone, sulphur dioxide and paraquat toxicity symptoms in broad bean. Physiol. Mol. Plant Pathol. 45:
5 263-279.
- 6 Lucas, P. W.; Cottam, D. A.; Mansfield, T. A. (1987) A large-scale fumigation system for investigating interactions
7 between air pollution and cold stress on plants. Environ. Pollut. 43: 15-28.
- 8 Lucas, P. W.; Cottam, D. A.; Sheppard, L. J.; Francis, B. J. (1988) Growth responses and delayed winter hardening in
9 Sitka spruce following summer exposure to ozone. New Phytol. 108: 495-504.
- 10 Luo, Y. (2001) Transient ecosystem response to free-air CO₂ enrichment (FACE): experimental evidence and methods
11 of analysis. New Phytol. 152: 3-8.
- 12 Luo, Y.; Reynolds, J. F. (1999) Validity of extrapolating field CO₂ experiments to predict carbon sequestration in
13 natural ecosystems. Ecology 80: 1568-1583.
- 14 Luo, Y.; Canadell, J.; Mooney, H. A. (1999) Interactive effects of carbon dioxide and environmental stress on plants
15 and ecosystems: a synthesis. In: Luo, Y.; Mooney, H. A., eds. Carbon dioxide and environmental stress. San
16 Diego, CA: Academic Press; pp. 393-408.
- 17 Lutz, C.; Anegg, S.; Gerant, D.; Alaoui-Sossé, B.; Gérard, J.; Dizengremel, P. (2000) Beech trees exposed to high CO₂
18 and to simulated summer ozone levels: effects on photosynthesis, chloroplast components and leaf enzyme
19 activity. Physiol. Plant. 109: 252-259.
- 20 Luwe, M. W. F.; Takahama, U.; Heber, U. (1993) Role of ascorbate in detoxifying ozone in the apoplast of spinach
21 (*Spinacia oleracea* L.) leaves. Plant Physiol. 101: 969-976.
- 22 Luxmoore, R. J. (1988) Assessing the mechanisms of crop loss from air pollutants with process models. In: Heck, W.
23 W.; Taylor, O. C.; Tingey, D. T., eds. Assessment of crop loss from air pollutants: proceedings of an
24 international conference; October 1987; Raleigh, NC. Essex, United Kingdom: Elsevier Science Publishers,
25 Ltd.; pp. 417-444.
- 26 Lyons, T. M.; Barnes, J. D. (1998) Influence of plant age on ozone resistance in *Plantago major*. New Phytol. 138:
27 83-89.
- 28 Lyons, T. M.; Barnes, J. D.; Davison, A. W. (1997) Relationships between ozone resistance and climate in European
29 populations of *Plantago major*. New Phytol. 136: 503-510.
- 30 Lyons, T.; Ollerenshaw, J. H.; Barnes, J. D. (1999) Impacts of ozone on *Plantago major*: apoplastic and symplastic
31 antioxidant status. New Phytol. 141: 253-263.
- 32 MacLean, D. C. (1990) Joint action of ozone and hydrogen fluoride on foliar senescence in maize. Environ. Pollut. 63:
33 283-292.
- 34 Madamanchi, N. R.; Anderson, J. V.; Alscher, R. G.; Cramer, C. L.; Hess, J. L. (1992) Purification of multiple forms
35 of glutathione reductase from pea (*Pisum sativum* L.) seedlings and enzyme levels in ozone-fumigated pea
36 leaves. Plant Physiol. 100: 138-145.
- 37 Magan, N.; Kirkwood, I. A.; McLeod, A. R.; Smith, M. K. (1995) Effect of open-air fumigation with sulphur dioxide
38 and ozone on phyllosphere and endophytic fungi of conifer needles. Plant Cell Environ. 18: 291-302.
- 39 Maggs, R.; Ashmore, M. R. (1998) Growth and yield responses of Pakistan rice (*Oryza sativa* L.) cultivars to O₃ and
40 NO₂. Environ. Pollut. 103: 159-170.
- 41 Maier-Maercker, U. (1997) Experiments on the water balance of individual attached twigs of *Picea abies* (L.) Karst. in
42 pure and ozone-enriched air. Trees 11: 229-239.
- 43 Maier-Maercker, U. (1998) Predisposition of trees to drought stress by ozone. Tree Physiol. 19: 71-78.
- 44 Maier-Maercker, U.; Koch, W. (1992) The effect of air pollution on the mechanism of stomatal control. Trees (Berlin)
45 7: 12-25.
- 46 Manderscheid, R.; Jäger, H.-J.; Kress, L. W. (1992) Effects of ozone on foliar nitrogen metabolism of *Pinus taeda* L.
47 and implications for carbohydrate metabolism. New Phytol. 121: 623-633.
- 48 Mandl, R. H.; Weinstein, L. H.; McCune, D. C.; Keveny, M. (1973) A cylindrical, open-top chamber for the exposure
49 of plants to air pollutants in the field. J. Environ. Qual. 2: 371-376.
- 50 Mandl, R. H.; Laurence, J. A.; Kohut, R. J. (1989) Development and testing of open-top chambers for exposing large,
51 perennial plants to air pollutants. J. Environ. Qual. 18: 534-540.
- 52 Manninen, A.-M.; Holopainen, T.; Lyytikäinen-Saarenmaa, P.; Holopainen, J. K. (2000) The role of low-level ozone
53 exposure and mycorrhizas in chemical quality and insect herbivore performance on Scots pine seedlings. Global
54 Change Biol. 6: 111-121.

- 1 Manninen, S.; Siivonen, N.; Timonen, U.; Huttunen, S. (2003) Differences in ozone response between two Finnish
2 wild strawberry populations. *Environ. Exp. Bot.* 49: 29-39.
- 3 Manning, W. J. (2000) Use of protective chemicals to assess the effects of ambient ozone on plants. In: Agrawal, S. B.;
4 Agrawal, M., eds. *Environmental pollution and plant responses*. Boca Raton, FL: Lewis Publishers; pp.
5 247-258.
- 6 Manning, W. J. (2003) Detecting plant effects is necessary to give biological significance to ambient ozone monitoring
7 data and predictive ozone standards. *Environ. Pollut.* 126: 375-379.
- 8 Manning, W. J.; Keane, K. D. (1988) Effects of air pollutants on interactions between plants, insects, and pathogens.
9 In: Heck, W. W.; Taylor, O. C.; Tingey, D. T., eds. *Assessment of crop loss from air pollutants: proceedings of*
10 *an international conference; October 1987; Raleigh, NC. Essex, United Kingdom: Elsevier Science Publishers,*
11 *Ltd.; pp. 365-386.*
- 12 Manning, W. J.; Krupa, S. V. (1992) Experimental methodology for studying the effects of ozone on crops and trees.
13 In: Lefohn, A. S., ed. *Surface level ozone exposures and their effects on vegetation*. Chelsea, MI: Lewis
14 Publishers, Inc.; pp. 93-156.
- 15 Manning, W. J.; Feder, W. A.; Vardaro, P. M. (1974) Suppression of oxidant injury by benomyl: effects on yields of
16 bean cultivars in the field. *J. Environ. Qual.* 3: 1-3.
- 17 Manning, W. J.; Godzik, B.; Musselman, R. (2002) Potential bioindicator plant species for ambient ozone in forested
18 mountain areas of central Europe. *Environ. Pollut.* 119: 283-290.
- 19 Manning, W. J.; Flagler, R. B.; Frenkel, M. A. (2003) Assessing plant response to ambient ozone: growth of
20 ozone-sensitive loblolly pine seedlings treated with ethylenediurea or sodium erythorbate. *Environ. Pollut.* 126:
21 73-81.
- 22 Martin, M. J.; Host, G. E.; Lenz, K. E.; Isebrands, J. G. (2001) Simulating the growth response of aspen to elevated
23 ozone: a mechanistic approach to scaling a leaf-level model of ozone effects on photosynthesis to a complex
24 canopy architecture. *Environ. Pollut.* 115: 425-436.
- 25 Massman, W. J. (1993) Partitioning ozone fluxes to sparse grass and soils and the inferred resistances to dry
26 deposition. *Atmos. Environ. Part A* 27A: 167-174.
- 27 Massman, W. J. (2004) Toward an ozone standard to protect vegetation based on effective dose: a review of deposition
28 resistances and a possible metric. *Atmos. Environ.* 38: 2323-2337.
- 29 Massman, W. J.; Grantz, D. A. (1995) Estimating canopy conductance to ozone uptake from observations of
30 evapotranspiration at the canopy scale and at the leaf scale. *Global Change Biol.* 1: 183-198.
- 31 Massman, W. J.; Ham, J. M. (1994) An evaluation of a surface-energy balance method for partitioning ET data into
32 plant and soil components for a surface with partial canopy cover. *Agric. For. Meteorol.* 67: 253-267.
- 33 Massman, W. J.; Pederson, J.; Delany, A.; Grantz, D.; Denhartog, G.; Neumann, H. H.; Oncley, S. P.; Pearson, R.;
34 Shaw, R. H. (1994) An evaluation of the regional acid deposition model surface module for ozone uptake at 3
35 sites in the San-Joaquin valley of California. *J. Geophys. Res. [Atmos.]* 99: 8281-8294.
- 36 Massman, W. J.; Musselman, R. C.; Lefohn, A. S. (2000) A conceptual ozone dose-response model to develop a
37 standard to protect vegetation. *Atmos. Environ.* 34: 745-759.
- 38 Matters, G. L.; Scandalios, J. G. (1987) Synthesis of isozymes of superoxide dismutase in maize leaves in response to
39 O₃, SO₂, and elevated O₂. *J. Exp. Bot.* 38: 842-852.
- 40 Matyssek, R.; Günthardt-Goerg, M. S.; Maurer, S.; Keller, T. (1995) Nighttime exposure to ozone reduces whole-plant
41 production in *Betula pendula*. *Tree Physiol.* 15: 159-165.
- 42 Matyssek, R.; Günthardt-Goerg, M. S.; Maurer, S.; Christ, R. (2002) Tissue structure and respiration of stems of
43 *Betula pendula* under contrasting ozone exposure and nutrition. *Trees* 16: 375-385.
- 44 Maurer, S.; Matyssek, R. (1997) Nutrition and the ozone sensitivity of birch (*Betula pendula*) II. Carbon balance,
45 water-use efficiency and nutritional status of the whole plant. *Trees* 12: 11-20.
- 46 Mazarura, U. (1997) Effects of fluctuating levels of ozone and nitrogen dioxide alone and in sequence on plant growth
47 and photosynthesis [dissertation]. Vancouver, British Columbia, Canada: University of British Columbia.
- 48 McBride, J. R.; Laven, R. D. (1976) Scars as an indicator of fire frequency in the San Bernardino Mountains,
49 California. *J. For.* 74: 439-442.
- 50 McBride, J. R.; Laven, R. D. (1999) Impact of oxidant air pollutants on forest succession in the mixed conifer forests
51 of the San Bernardino Mountains. In: Miller, P. R.; McBride, J. E., eds. *Oxidant air pollution impacts in the*
52 *montane forests of southern California: a case study of the San Bernardino Mountains*. New York, NY:
53 Springer; pp. 338-352. (Ecological Studies, v. 134).

- 1 McBride, J. R.; Semion, V. P.; Miller, P. R. (1975) Impact of air pollution on the growth of Ponderosa pine. *Calif.*
2 *Agric.* 29: 8-9.
- 3 McCarl, B. A.; Alig, R.; Adams, D. M. (1998) Impacts of incorporating land exchanges between forestry and
4 agriculture in sector models. *J. Agric. Appl. Econ.* 30: 389-401.
- 5 McCool, P. M.; Musselman, R. C.; Younglove, T.; Teso, R. R. (1988) Response of kidney bean to sequential ozone
6 exposures. *Environ. Exp. Bot.* 28: 307-313.
- 7 McDonald, E. P.; Kruger, E. L.; Riemenschneider, D. E.; Isebrands, J. G. (2002) Competitive status influences
8 tree-growth responses to elevated CO₂ and O₃ in aggrading aspen stands. *Funct. Ecol.* 16: 792-801.
- 9 McGartland, A. M. (1987) The implications of ambient ozone standards for U.S. agriculture: a comment and some
10 further evidence. *J. Environ. Manage.* 24: 139-146.
- 11 McKee, I. F.; Farage, P. K.; Long, S. P. (1995) The interactive effects of elevated CO₂ and O₃ concentration on
12 photosynthesis in spring wheat. *Photosyn. Res.* 45: 111-119.
- 13 McKee, I. F.; Bullimore, J. F.; Long, S. P. (1997a) Will elevated CO₂ concentrations protect the yield of wheat from O₃
14 damage? *Plant Cell Environ.* 20: 77-84.
- 15 McKee, I. F.; Eiblmeier, M.; Polle, A. (1997b) Enhanced ozone-tolerance in wheat grown at an elevated CO₂
16 concentration: ozone exclusion and detoxification. *New Phytol.* 137: 275-284.
- 17 McKee, I. F.; Mulholland, B. J.; Craigon, J.; Black, C. R.; Long, S. P. (2000) Elevated concentrations of atmospheric
18 CO₂ protect against and compensate for O₃ damage to photosynthetic tissues of field-grown wheat. *New Phytol.*
19 146: 427-435.
- 20 McLaughlin, S. B.; Downing, D. J. (1995) Interactive effects of ambient ozone and climate measured on growth of
21 mature forest trees. *Nature (London)* 374: 252-254.
- 22 McLaughlin, S. B.; Downing, D. J. (1996) Interactive effects of ambient ozone and climate measured on growth of
23 mature loblolly pine trees. *Can. J. For. Res.* 670-681.
- 24 McLaughlin, S.; Percy, K. (1999) Forest health in North America: some perspectives on actual and potential roles of
25 climate and air pollution. *Water Air Soil Pollut.* 116: 151-197.
- 26 McLaughlin, S. B.; McConathy, R. K.; Duvick, D.; Mann, L. K. (1982) Effects of chronic air pollution stress on
27 photosynthesis, carbon allocation, and growth of white pine trees. *For. Sci.* 28: 60-70.
- 28 McLaughlin, S. B.; Layton, P. A.; Adams, M. B.; Edwards, N. T.; Hanson, P. J.; O'Neill, E. G.; Roy, W. K. (1994)
29 Growth-responses of 53 open-pollinated loblolly-pine families to ozone and acid-rain. *J. Environ. Qual.* 23:
30 247-257.
- 31 McLaughlin, S. B.; Shortle, W. C.; Smith, K. T. (2002) Dendroecological applications in air pollution and
32 environmental chemistry: research needs. *Dendrochronologia* 20: 133-157.
- 33 McLaughlin, S. B.; Wullschlegel, S. D.; Nosal, M. (2003) Diurnal and seasonal changes in stem increment and water
34 use by yellow poplar trees in response to environmental stress. *Tree Physiol.* 23: 1125-1136.
- 35 McLeod, A. R.; Fackrell, J. E.; Alexander, K. (1985) Open-air fumigation of field crops: criteria and design for a new
36 experimental system. *Atmos. Environ.* 19: 1639-1649.
- 37 McLeod, A. R.; Shaw, P. J. A.; Holland, M. R. (1992) The Liphook Forest Fumigation Project: studies of sulphur
38 dioxide and ozone effects on coniferous trees. *For. Ecol. Manage.* 51: 121-127.
- 39 Mearns, L. O.; Rosenzweig, C.; Goldberg, R. (1997) Mean and variance change in climate scenarios: methods,
40 agricultural applications and measures of uncertainty. *Clim. Change* 35: 367-396.
- 41 Mehdy, M. C. (1994) Active oxygen species in plant defense against pathogens. *Plant Physiol.* 105: 467-472.
- 42 Mehlhorn, H.; Wellburn, A. R. (1987) Stress ethylene formation determines plant sensitivity to ozone. *Nature*
43 (London) 327: 417-418.
- 44 Mehlhorn, H.; Wenzel, A. A. (1996) Manganese deficiency enhances ozone toxicity in bush beans (*Phaseolus vulgaris*
45 L. cv. Saxa). *J. Plant Physiol.* 148: 155-159.
- 46 Mehlhorn, H.; Tabner, B. J.; Wellburn, A. R. (1990) Electron spin resonance evidence for the formation of free
47 radicals in plants exposed to ozone. *Physiol. Plant.* 79: 377-383.
- 48 Mehlhorn, H.; O'Shea, J. M.; Wellburn, A. R. (1991) Atmospheric ozone interacts with stress ethylene formation by
49 plants to cause visible plant injury. *J. Exp. Bot.* 42: 17-24.
- 50 Meinhard, M.; Rodriguez, P. L.; Grill, E. (2002) The sensitivity of ABI2 to hydrogen peroxide links the abscisic
51 acid-response regulator to redox signalling. *Planta* 214: 775-782.
- 52 Mendelsohn, R.; Nordhaus, W.; Shaw, D. (1994) The impact of global warming on agriculture: a Ricardian analysis.
53 *Am. Econ. Rev.* 84: 753-771.

- 1 Miller, P. L. (1973) Oxidant-induced community change in a mixed conifer forest. In: Naegele, J. A., ed. Air pollution
2 damage to vegetation. Washington, DC: American Chemical Society; pp. 101-117. (Advances in chemistry
3 series: v. 122).
- 4 Miller, P. R.; Elderman, M. J. (1977) Photochemical oxidant air pollutant effects on a mixed conifer forest ecosystem:
5 a progress report, 1976. Corvallis, OR: U.S. Environmental Protection Agency, Corvallis Environmental
6 Research Laboratory; EPA report no. EPA-600/3-77-104. Available from: NTIS, Springfield, VA; PB-274531.
- 7 Miller, P. R.; McBride, J. R. (1999) Assessment of ecological risks and implications for policy and management in the
8 San Bernardino Mountains. In: Miller, P. R.; McBride, J. R., eds. Oxidant air pollution impacts in the Montane
9 forests of southern California: a case study of the San Bernardino Mountains. New York, NY: Springer; pp.
10 397-404. (Ecological Studies, v. 134).
- 11 Miller, P. R.; Millecan, A. A. (1971) Extent of oxidant air pollution damage to some pines and other conifers in
12 California. Plant Dis. Rep. 55: 555-559.
- 13 Miller, P. R.; Rechel, J. (1999) Temporal changes in crown condition indices, needle litterfall, and collateral needle
14 injuries of Ponderosa and Jeffrey pines. In: Miller, P. R.; McBride, J. R., eds. Oxidant air pollution impacts in
15 the Montane forests of southern California: a case study of the San Bernardino Mountains. New York, NY:
16 Springer; pp. 164-178.
- 17 Miller, P. R.; Parmeter, J. R., Jr.; Taylor, O. C.; Cardiff, E. A. (1963) Ozone injury to the foliage of *Pinus ponderosa*.
18 Phytopathology 53: 1072-1076.
- 19 Miller, P. R.; Parmeter, J. R., Jr.; Flick, B. H.; Martinez, C. W. (1969) Ozone dosage response of ponderosa pine
20 seedlings. J. Air Pollut. Control Assoc. 19: 435-438.
- 21 Miller, P. R.; McCutchan, M. H.; Milligan, H. P. (1972) Oxidant air pollution in the Central Valley, Sierra Nevada
22 foothills, and Mineral King Valley of California. Atmos. Environ. 6: 623-633.
- 23 Miller, P. R.; McBride, J. R.; Schilling, S. L.; Gomez, A. P. (1989) Trend of ozone damage to conifer forests between
24 1974 and 1988 in the San Bernardino Mountains of southern California. In: Olson, R. K.; Lefohn, A. S., eds.
25 Effects of air pollution on western forests [an A&WMA symposium; June; Anaheim, CA]. Pittsburgh, PA: Air
26 and Waste Management Association; pp. 309-323. (APCA transactions series: no. 16).
- 27 Miller, J. E.; Booker, F. L.; Fiscus, E. L.; Heagle, A. S.; Pursley, W. A.; Vozzo, S. F.; Heck, W. W. (1994)
28 Ultraviolet-B radiation and ozone effects on growth, yield, and photosynthesis of soybean. J. Environ. Qual. 23:
29 83-91.
- 30 Miller, J. E.; Heagle, A. S.; Pursley, W. A. (1998) Influence of ozone stress on soybean response to carbon dioxide
31 enrichment: II. Biomass and development. Crop Sci. 38: 122-128.
- 32 Miller, P. R.; Bytnerowicz, A.; Fenn, M.; Poth, M.; Temple, P.; Schilling, S.; Jones, D.; Johnson, D.; Chow, J.;
33 Watson, J. (1998) Multidisciplinary study of ozone, acidic deposition and climatic effects on a mixed conifer
34 forest in California, USA. Chemosphere 36: 1001-1006.
- 35 Mills, G. (2002) Modification of plant response by environmental conditions. In: Bell, J. N. B.; Treshow, M., eds. Air
36 Pollution and Plant Life. 2nd ed. Chichester, United Kingdom: John Wiley & Sons Ltd.; pp. 343-358.
- 37 Mills, G.; Ball, G.; Hayes, F.; Fuhrer, J.; Skarby, L.; Gimeno, B.; De Temmerman, L.; Heagle, A.; Members of the ICP
38 Vegetation programme. (2000) Development of a multi-factor model for predicting the effects of ambient ozone
39 on the biomass of white clover. Environ. Pollut. 109: 533-542.
- 40 Minnich, R. A. (1999) Vegetation, fire regimes, and forest dynamics. In: Oxidant air pollution impacts in the montane
41 forests southern California: a case study of the San Bernardino mountains. New York, NY: Springer; pp. 44-80.
42 (Ecological studies: v. 134).
- 43 Minnich, R. A.; Padgett, P. E. (2003) Geology, climate, and vegetation of the Sierra Nevada and the mixed-conifer
44 zone: an introduction to the ecosystem. In: Bytnerowicz, A.; Arbaugh, M. J.; Alonso, R., eds. Ozone Air
45 Pollution in the Sierra Nevada: Distribution and Effects on Forests. New York, NY: Elsevier; pp. 1-31.
46 (Developments in Environmental Science, v. 2).
- 47 Minnich, R. A.; Barbour, M. G.; Burk, J. H.; Fernau, R. F. (1995) Sixty years of change in California conifer forests of
48 the San Bernardino mountains. Conserv. Biol. 9: 902-914.
- 49 Moldau, H. (1998) Hierarchy of ozone scavenging reactions in the plant cell wall. Physiol. Plant. 104: 617-622.
- 50 Moldau, H.; Bichele, I. (2002) Plasmalemma protection by the apoplast as assessed from above zero ozone
51 concentration in leaf intercellular air spaces. Planta 214: 484-487.
- 52 Momen, B.; Helms, J. A. (1996) Effects of simulated acid rain and ozone on foliar chemistry of field-grown *Pinus*
53 *ponderosa* seedlings and mature trees. Environ. Pollut. 91: 105-111.

- 1 Momen, B.; Helms, J. A.; Criddle, R. S. (1996) Foliar metabolic heat rate of seedlings and mature trees of *Pinus*
- 2 *ponderosa* exposed to simulated acid rain and ozone. *Plant Cell Environ.* 19: 747-753.
- 3 Momen, B.; Anderson, P. D.; Helms, J. A.; Houpis, J. L. J. (1997) Acid rain and ozone effects on gas exchange of
- 4 *Pinus ponderosa*: a comparison between trees and seedlings. *Int. J. Plant Sci.* 158: 617-621.
- 5 Momen, B.; Anderson, P. D.; Helms, J. A. (1999) Temperature dependency of acid-rain effect on photosynthesis of
- 6 *Pinus ponderosa*. *For. Ecol. Manage.* 113: 223-230.
- 7 Momen, B.; Anderson, P. D.; Houpis, J. L. J.; Helms, J. A. (2002) Growth of ponderosa pine seedlings as affected by
- 8 air pollution. *Atmos. Environ.* 36: 1875-1882.
- 9 Monteith, J. L.; Elston, J. (1993) Climatic constraints on crop production. In: Fowden, L.; Mansfield, T.; Stoddart, J.,
- 10 eds. *Plant adaptation to environmental stress*. London, United Kingdom: Chapman and Hall; pp. 3-18.
- 11 Morgan, P. B.; Ainsworth, E. A.; Long, S. P. (2003) How does elevated ozone impact soybean? A meta-analysis of
- 12 photosynthesis, growth and yield. *Plant Cell Environ.* 26: 1317-1328.
- 13 Morison, J. I. L.; Lawlor, D. W. (1999) Interactions between increasing CO₂ concentration and temperature on plant
- 14 growth. *Plant Cell Environ.* 22: 659-682.
- 15 Mortensen, L. M. (1998) Growth responses of seedlings of six *Betula pubescens* Ehrh. Provenances to six ozone
- 16 exposure regimes. *Scand. J. For. Res.* 13: 189-196.
- 17 Mortensen, L. M. (1999) Foliar injuries caused by ozone in *Betula pubescens* Ehrh. and *Phleum pratense* L. as
- 18 influenced by climatic conditions before and during O₃ exposure. *Acta Agric. Scand. B* 49: 44-49.
- 19 Mortensen, L.; Bastrup-Birk, A. (1996) Critical levels for O₃ for biomass production in white clover. In: Karenlampi,
- 20 L.; Skärby, L., eds. *Critical levels for ozone in Europe: testing and finalizing the concepts: UN-ECE workshop*
- 21 *report*. UN-ECE Convention on Long-Range Transboundary Air Pollution; April; Kuopio, Finland. Kuopio,
- 22 Finland: University of Kuopio, Department of Ecology and Environmental Science; pp. 285-289.
- 23 Mudd, J. B. (1996) Biochemical basis for the toxicity of ozone. In: Yunus, M.; Iqbal, M., eds. *Plant response to air*
- 24 *pollution*. New York, NY: John Wiley & Sons; pp. 267-283.
- 25 Mudd, J. B.; Kozłowski, T. T., eds. (1975) *Responses of plants to air pollution*. New York, NY: Academic Press.
- 26 (Physiological ecology, a series of monographs, texts, and treatises).
- 27 Mudd, J. B.; Leavitt, R.; Ongun, A.; McManus, T. T. (1969) Reaction of ozone with amino acids and proteins. *Atmos.*
- 28 *Environ.* 3: 669-681.
- 29 Mudd, J. B.; McManus, T. T.; Ongun, A. (1971) Inhibition of lipid metabolism in chloroplasts by ozone. In: England,
- 30 H. M.; Beery, W. T., eds. *Proceedings of the second clean air congress*. New York, NY: Academic Press; pp.
- 31 256-260.
- 32 Mudd, J. B.; Dawson, P. J.; Adams, J. R.; Wingo, J.; Santrock, J. (1996) Reaction of ozone with enzymes of
- 33 erythrocyte membranes. *Arch. Biochem. Biophys.* 335: 145-151.
- 34 Mudd, J. B.; Dawson, P. J.; Santrock, J. (1997a) Ozone does not react with human erythrocyte membrane lipids. *Arch.*
- 35 *Biochem. Biophys.* 341: 251-258.
- 36 Mudd, J. B.; Dawson, P. J.; Tseng, S.; Liu, F.-P. (1997b) Reaction of ozone with protein tryptophans: band III, serum
- 37 albumin, and cytochrome C. *Arch. Biochem. Biophys.* 338: 143-149.
- 38 Mueller, S. F. (1994) Characterization of ambient ozone levels in the Great Smoky Mountains National Park. *J. Appl.*
- 39 *Meteorol.* 33: 465-472.
- 40 Mukammal, E. I. (1965) Ozone as a cause of tobacco injury. *Agric. Meteorol.* 2: 145-165.
- 41 Mulchi, C. L.; Slaughter, L.; Saleem, M.; Lee, E. H.; Pausch, R.; Rowland, R. (1992) Growth and physiological
- 42 characteristics of soybean in open-top chambers in response to ozone and increased atmospheric CO₂. *Agric.*
- 43 *Ecosyst. Environ.* 38: 107-118.
- 44 Mulchi, C.; Rudorff, B.; Lee, E.; Rowland, R.; Pausch, R. (1995) Morphological responses among crop species to
- 45 full-season exposures to enhanced concentrations of atmospheric CO₂ and O₃. *Water Air Soil Pollut.* 85:
- 46 1379-1386.
- 47 Mulholland, B.; Craigon, J.; Black, C.; Atherton, J.; Landon, G. (1997a) Effects of elevated carbon dioxide and ozone
- 48 on the growth and yield of spring wheat (*Triticum aestivum* L.). *J. Exper. Bot.* 48: 113-122.
- 49 Mulholland, B. J.; Craigon, J.; Black, C. R.; Colls, J. J.; Atherton, J.; Landon, G. (1997b) Impact of elevated
- 50 atmospheric CO₂ and O₃ on gas exchange and chlorophyll content in spring wheat (*Triticum aestivum* L.). *J.*
- 51 *Exp. Bot.* 48: 1853-1863.
- 52 Mulholland, B. J.; Craigon, J.; Black, C. R.; Colls, J. J.; Atherton, J.; Landon, G. (1998a) Effects of elevated CO₂ and
- 53 O₃ on the rate and duration of grain growth and harvest index in spring wheat (*Triticum aestivum* L.). *Global*
- 54 *Change Biol.* 4: 627-635.

- 1 Mulholland, B. J.; Craigon, J.; Black, C. R.; Colls, J. J.; Atherton, J.; Landon, G. (1998b) Growth, light interception
2 and yield responses of spring wheat (*Triticum aestivum* L.) grown under elevated CO₂ and O₃ in open-top
3 chambers. *Global Change Biol.* 4: 121-130.
- 4 Muntifering, R. B.; Crosby, D. D.; Powell, M. C.; Chappelka, A. H. (2000) Yield and quality characteristics of
5 bahiagrass (*Paspalum notatum*) exposed to ground-level ozone. *Anim. Feed Sci. Technol.* 84: 243-256.
- 6 Murphy, J. J.; Deluki, M. A.; McCubbin, D. R.; Kim, H. J. (1999) The cost of crop damage caused by ozone air
7 pollution from motor vehicles. *J. Environ. Manage.* 55: 273-289.
- 8 Musselman, R. C.; Hale, B. A. (1997) Methods for controlled and field ozone exposures of forest tree species in North
9 America. In: Sandermann, H.; Wellburn, A. R.; Health, R. L., eds. *Forest decline and ozone*. Berlin, Germany:
10 Springer-Verlag; pp. 277-315. (Ecological Studies, v.127).
- 11 Musselman, R. C.; Massman, W. J. (1999) Ozone flux to vegetation and its relationship to plant response and ambient
12 air quality standards. *Atmos. Environ.* 33: 65-73.
- 13 Musselman, R. C.; Minnick, T. J. (2000) Nocturnal stomatal conductance and ambient air quality standards for ozone.
14 *Atmos. Environ.* 34: 719-733.
- 15 Musselman, R. C.; Oshima, R. J.; Gallavan, R. E. (1983) Significance of pollutant concentration distribution in the
16 response of 'red kidney' beans to ozone. *J. Am. Soc. Hortic. Sci.* 108: 347-351.
- 17 Musselman, R. C.; Huerta, A. J.; McCool, P. M.; Oshima, R. J. (1986) Response of beans to simulated ambient and
18 uniform ozone distributions with equal peak concentration. *J. Am. Soc. Hortic. Sci.* 111: 470-473.
- 19 Musselman, R. C.; McCool, P. M.; Younglove, T. (1988) Selecting ozone exposure statistics for determining crop
20 yield loss from air pollutants. *Environ. Pollut.* 53: 63-78.
- 21 Musselman, R. C.; Younglove, T.; McCool, P. M. (1994) Response of *Phaseolus vulgaris* L. to differing ozone
22 regimes having identical total exposure. *Atmos. Environ.* 28: 2727-2731.
- 23 Neill, S. J.; Desikan, R.; Clarke, A.; Hurst, R. D.; Hancock, J. T. (2002) Hydrogen peroxide and nitric oxide as
24 signalling molecules in plants. *J. Exper. Bot.* 53: 1237-1247.
- 25 Neill, S.; Desikan, R.; Hancock, J. (2002) Hydrogen peroxide signalling. *Curr. Opin. Plant Biol.* 5: 388-395.
- 26 Neufeld, H. S.; Lee, E. H.; Renfro, J. R.; Hacker, W. D. (2000) Seedling insensitivity to ozone for three conifer species
27 native to Great Smoky Mountains National Park. *Environ. Pollut.* 108: 141-151.
- 28 Neufeld, H. S.; Renfro, J. R.; Hacker, W. D.; Silsbee, D. (1992) Ozone in Great Smoky Mountains National Park:
29 dynamics and effects on plants. In: Berglund, R. L., ed. *Tropospheric ozone and the environment II: effects,*
30 *modeling and control [papers from an international specialty conference];* November; Atlanta, GA. Pittsburgh,
31 PA: Air & Waste Management Association; pp. 594-617. (A&WMA transactions series no. 20).
- 32 Neufeld, H. S.; Lee, E. H.; Renfro, J. R.; Hacker, W. D.; Yu, B.-H. (1995) Sensitivity of seedlings of black cherry
33 (*Prunus serotina* Ehrh.) to ozone in Great Smoky Mountains National Park. I. Exposure-response curves for
34 biomass. *New Phytol.* 130: 447-459.
- 35 Niewiadomska, E.; Gaucher-Veilleux, C.; Chevrier, N.; Mauffette, Y.; Dizengremel, P. (1999) Elevated CO₂ does not
36 provide protection against ozone considering the activity of several antioxidant enzymes in the leaves of sugar
37 maple. *J. Plant Physiol.* 155: 70-77.
- 38 Nijs, I.; Teughels, H.; Blum, H.; Hendrey, G.; Impens, I. (1996) Simulation of climate change with infrared heaters
39 reduces the productivity of *Lolium perenne* L. in summer. *Environ. Exp. Bot.* 36: 271-280.
- 40 Nikolov, N.; Zeller, K. (2003) Modeling coupled interactions of carbon, water, and ozone exchange between terrestrial
41 ecosystems and the atmosphere. I. Model description. *Environ. Pollut.* 124: 231-246.
- 42 Noormets, A.; McDonald, E. P.; Dickson, R. E.; Kruger, E. L.; Sôber, A.; Isebrands, J. G.; Karnosky, D. F. (2001) The
43 effect of elevated carbon dioxide and ozone on leaf- and branch-level photosynthesis and potential plant-level
44 carbon gain in aspen. *Trees* 15: 262-270.
- 45 Noormets, A.; Sôber, A.; Pell, E. J.; Dickson, R. E.; Posila, G. K.; Sôber, J.; Isebrands, J. G.; Karnosky, D. F. (2001)
46 Stomatal and non-stomatal limitation to photosynthesis in two trembling aspen (*Populus tremuloides* Michx.)
47 clones exposed to elevated CO₂ and/or O₃. *Plant Cell Environ.* 24: 327-336.
- 48 Norby, R. J.; Wullschleger, S. D.; Gunderson, C. A.; Johnson, D. W.; Ceulemans, R. (1999) Tree responses to rising
49 CO₂ in field experiments: implications for the future forest. *Plant Cell Environ.* 22: 683-714.
- 50 Norton, B. G. (1987) *Why preserve natural variety?* Princeton, NJ: Princeton University Press.
- 51 Nouchi, I.; Aoki, K. (1979) Morning glory as a photochemical oxidant indicator. *Environ. Pollut.* 18: 289-303.
- 52 Nouchi, I.; Toyama, S. (1988) Effects of ozone and peroxyacetyl nitrate on polar lipids and fatty acids in leaves of
53 morning glory and kidney bean. *Plant Physiol.* 87: 638-646.

- 1 Novak, K.; Skelly, J. M.; Schaub, M.; Krauchi, N.; Hug, C.; Landolt, W.; Bleuter, P. (2003) Ozone air pollution and
2 foliar injury development on native plants of Switzerland. *Environ. Pollut.* 125: 41-52.
- 3 Nunn, A. J.; Reiter, I. M.; Haberle, K.-H.; Werner, H.; Langebartels, C.; Sandermann, H.; Heerdt, C.; Fabian, P.;
4 Matyssek, R. (2002) "Free-air" ozone canopy fumigation in an old-growth mixed forest: concept and
5 observations in beech. *Phyton* 42: 105-119.
- 6 Nussbaum, S.; Geissman, M.; Fuhrer, J. (1995) Effects of nitric oxide and ozone on spring wheat (*Triticum aestivum*).
7 *Water Air Soil Pollut.* 85: 1449-1454.
- 8 Nussbaum, S.; Geissmann, M.; Fuhrer, J. (1995) Ozone exposure-response relationships for mixtures of perennial
9 ryegrass and white clover depend on ozone exposure patterns. *Atmos. Environ.* 29: 989-995.
- 10 Nussbaum, S.; Ammann, M.; Fuhrer, J. (1999) Foliar absorption and use of airborne oxidized nitrogen by terrestrial
11 plants. In: Srivastava, H. S.; Singh, R. P., eds. *Nitrogen Nutrition and Plant Growth*; pp. 101-172. New Delhi,
12 India: Oxford & IBH Publishing Co. Pvt. Ltd.
- 13 Nussbaum, S.; Bungener, P.; Geissmann, M.; Fuhrer, J. (2000) Plant-plant interactions and soil moisture might be
14 important in determining ozone impacts on grasslands. *New Phytol.* 147: 327-335.
- 15 Nussbaum, S.; Geissman, M.; Saurer, M.; Siegwolf, R.; Fuhrer, J. (2000) Ozone and low concentrations of nitric oxide
16 have similar effects on carbon isotope discrimination and gas exchange in leaves of wheat (*Triticum aestivum*
17 L.). *J. Plant Physiol.* 156: 741-745.
- 18 O'Gara, P. J. (1922) Sulfur dioxide and fume problems and their solution. (presented at the 14th semiannual meeting of
19 the American Institute of Chemical Engineers, Niagara Falls, Canada, June 19-22, 1922). Presented at: 14th
20 semiannual meeting of the American Institute of Chemical Engineers; June; Niagara Falls, Canada. *Proceedings*
21 *and papers of the meeting summarized in J. Ind. Eng. Chem.* 14: 744-745 by J. C. Olsen.
- 22 Ojanperä K.; Pätsikkä, E.; Ylärinta, T. (1998) Effects of low ozone exposure of spring wheat on net CO₂ uptake,
23 Rubisco, leaf senescence and grain filling. *New Phytol.* 138: 451-460.
- 24 Oksanen, E. J. (2001) Increasing tropospheric ozone level reduced birch (*Betula pendula*) dry mass within a five years
25 period. *Water Air Soil Pollut.* 130: 947-952.
- 26 Oksanen, E. (2003a) Physiological responses of birch (*Betula pendula*) to ozone: a comparison between
27 open-soil-grown trees exposed for six growing seasons and potted seedlings exposed for one season. *Tree*
28 *Physiol.* 23: 603-614.
- 29 Oksanen, E. (2003b) Responses of selected birch (*Betula pendula* Roth) clones to ozone change over time. *Plant Cell*
30 *Environ.* 26: 875-886.
- 31 Oksanen, E.; Holopainen, T. (2001) Responses of two birch (*Betula pendula* Roth) clones to different ozone profiles
32 with similar AOT40 exposure. *Atmos. Environ.* 35: 5245-5254.
- 33 Oksanen, E.; Saleem, A. (1999) Ozone exposure results in various carry-over effects and prolonged reduction in
34 biomass in birch (*Betula pendula* Roth). *Plant Cell Environ.* 22: 1401-1411.
- 35 Oksanen, E.; Amores, A.; Kokko, H.; Santamaria, J. M.; Kärenlampi, L. (2001a) Genotypic variation in growth and
36 physiological responses of Finnish hybrid aspen (*Populus tremuloides* x *P. tremula*) to elevated tropospheric
37 ozone concentration. *Tree Physiol.* 21: 1171-1181.
- 38 Oksanen, E.; Söber, J.; Karnosky, D. F. (2001b) Interactions of elevated CO₂ and ozone in leaf morphology of aspen
39 (*Populus tremuloides*) and birch (*Betula papyrifera*) in aspen FACE experiment. *Environ. Pollut.* 115: 437-446.
- 40 Ollerenshaw, J. H.; Lyons, T. (1999) Impacts of ozone on the growth and yield of field-grown winter wheat. *Environ.*
41 *Pollut.* 106: 67-72.
- 42 Ollerenshaw, J. H.; Lyons, T.; Barnes, J. D. (1999) Impacts of ozone on the growth and yield of field-grown winter
43 oilseed rape. *Environ. Pollut.* 104: 53-59.
- 44 Ollinger, S. V.; Aber, J. D.; Reich, P. B. (1997) Simulating ozone effects on forest productivity: interactions among
45 leaf-, canopy-, and stand-level processes. *Ecol. Appl.* 7: 1237-1251.
- 46 Ollinger, S. V.; Aber, J. D.; Federer, C. A. (1998) Estimating regional forest productivity and water yield using an
47 ecosystem model linked to a GIS. *Landscape Ecol.* 13: 323-334.
- 48 Olszyk, D. M.; Bytnerowicz, A.; Kats, G.; Dawson, P. J.; Wolf, J.; Thompson, C. R. (1986a) Crop effects from air
49 pollutants in air exclusion systems vs. field chambers. *J. Environ. Qual.* 15: 417-422.
- 50 Olszyk, D. M.; Kats, G.; Dawson, P. J.; Bytnerowicz, A.; Wolf, J.; Thompson, C. R. (1986b) Characteristics of air
51 exclusion systems vs. chambers for field air pollution studies. *J. Environ. Qual.* 15: 326-334.
- 52 Olszyk, D. M.; Takemoto, B. K.; Kats, G.; Dawson, P. J.; Morrison, C. L.; Preston, J. W.; Thompson, C. R. (1992)
53 Effects of open-top chambers on 'Valencia' orange trees. *J. Environ. Qual.* 21: 128-134.

- 1 Olszyk, D. M.; Tingey, D. T.; Watrud, L.; Seidler, R.; Andersen, C. (2000) Interactive effects of O₃ and CO₂:
2 implications for terrestrial ecosystems. In: Singh, S. N., ed. Trace gas emissions and plants. Dordrecht, The
3 Netherlands: Kluwer Academic Publishers; pp. 97-136.
- 4 Olszyk, D. M.; Johnson, M. G.; Phillips, D. L.; Seidler, R. J.; Tingey, D. T.; Watrud, L. S. (2001) Interactive effects of
5 CO₂ and O₃ on a ponderosa pine/litter/soil mesocosm. Environ. Pollut. 115: 447-462.
- 6 Ommen, O. E.; Donnelly, A.; Vanhoutvin, S.; Van Oijen, M.; Manderscheid, R. (1999) Chlorophyll content of spring
7 wheat flag leaves grown under elevated CO₂ concentrations and other environmental stresses within the
8 'ESPACE-wheat' project. Eur. J. Agron. 10: 197-203.
- 9 Ordin, L.; Hall, M. A.; Kindinger, J. I. (1969) Oxidant induced inhibition of enzymes involved in cell wall
10 polysaccharide synthesis. Arch. Environ. Health 18: 623-626.
- 11 Oren, R.; Ellsworth, D. S.; Johnsen, K. H.; Phillips, N.; Ewers, B. E.; Maier, C.; Schäfer, K. V. R.; McCarthy, H.;
12 Hendrey, G.; McNulty, S. G.; Katul, G. G. (2001) Soil fertility limits carbon sequestration by forest ecosystems
13 in a CO₂-enriched atmosphere. Nature (London) 411: 469-472.
- 14 Orendovici, T.; Skelly, J. M.; Ferdinand, J. A.; Savage, J. E.; Sanz, M.-J.; Smith, G. C. (2003) Response of native
15 plants of northeastern United States and southern Spain to ozone exposures; determining exposure/response
16 relationships. Environ. Pollut. 125: 31-40.
- 17 Ormrod, D. P.; Beckerson, D. W. (1986) Polyamines as antiozonants for tomato. HortScience 21: 1070-1071.
- 18 Ormrod, D. P.; Landry, L. G.; Conklin, P. L. (1995) Short-term UV-B radiation and ozone exposure effects on
19 aromatic secondary metabolite accumulation and shoot growth of flavonoid-deficient *Arabidopsis* mutants.
20 Physiol. Plant. 93: 602-610.
- 21 Oshima, R. J. (1975) Development of a system for evaluating and reporting economic crop losses caused by air
22 pollution in California. III. Ozone dosage-crop loss conversion function--alfalfa, sweet corn. IIIA. Procedures
23 for production, ozone effects on alfalfa and sweet corn, and evaluation of these systems. Sacramento, CA:
24 California Air Resources Board; report no. ARB-R-3-690-75-35. Available from: NTIS, Springfield, VA;
25 PB-272994.
- 26 Oshima, R. J.; Poe, M. P.; Braegelmann, P. K.; Baldwin, D. W.; Van Way, V. (1976) Ozone dosage-crop loss function
27 for alfalfa: a standardized method for assessing crop losses from air pollutants. J. Air Pollut. Control Assoc. 26:
28 861-865.
- 29 Oshima, R. J.; Braegelmann, P. K.; Baldwin, D. W.; Van Way, V.; Taylor, O. C. (1977) Responses of five cultivars of
30 fresh market tomato to ozone: a contrast of cultivar screening with foliar injury and yield. J. Am. Soc. Hortic.
31 Sci. 102: 286-289.
- 32 Pääkkönen, E.; Holopainen, T. (1995) Influence of nitrogen supply on the response of clones of birch (*Betula pendula*
33 Roth) to ozone. New Phytol. 129: 595-603.
- 34 Pääkkönen, E.; Holopainen, T.; Kärenlampi, L. (1996) Relationships between open-field ozone exposures and growth
35 and senescence of birch (*Betula pendula* and *Betula pubescens*). In: Kärenlampi, L.; Skärby, L., eds. Critical
36 levels for ozone in Europe: testing and finalizing the concepts: UN-ECE workshop report. UN-ECE Convention
37 on Long-Range Transboundary Air Pollution; April; Kuopio, Finland. Kuopio, Finland: University of Kuopio,
38 Department of Ecology and Environmental Science; pp. 298-302.
- 39 Pääkkönen, E.; Günthardt-Georg, M. S.; Holopainen, T. (1998) Responses of leaf processes in a sensitive birch (*Betula*
40 *pendula* Roth) clone to ozone combined with drought. Ann. Bot. 82: 49-59.
- 41 Paludan-Müller, G.; Saxe, H.; Leverenz, J. W. (1999) Responses to ozone in 12 provenances of European beech
42 (*Fagus sylvatica*): genotypic variation and chamber effects on photosynthesis and dry-matter partitioning. New
43 Phytol. 144: 261-273.
- 44 Panek, J. A.; Goldstein, A. H. (2001) Responses of stomatal conductance to drought in ponderosa pine: implications
45 for carbon and ozone uptake. Tree Physiol. 21: 337-344.
- 46 Panek, J.; Kurpius, M. R.; Goldstein, A. H. (2002) An evaluation of ozone exposure metrics for a seasonally
47 drought-stressed ponderosa pine ecosystem. Environ. Pollut. 117: 93-100.
- 48 Panek, J. A.; Baldocchi, D. D.; Goldstein, A. H. (2003) The need for spatially and functionally integrated models of
49 ozone deposition to Sierra Nevada forests. In: Bytnerowicz, A.; Arbaugh, M. J.; Alonso, R., eds. Ozone Air
50 Pollution in the Sierra Nevada: Distribution and Effects on Forests. New York, NY: Elsevier; pp. 325-357.
51 (Developments in Environmental Science, v. 2).
- 52 Patterson, M. T.; Rundel, P. W. (1995) Stand characteristics of ozone-stressed populations of *Pinus jeffreyi* (Pinaceae):
53 extent, development, and physiological consequences of visible injury. Am. J. Bot. 82: 150-158.

- Pauls, K. P.; Thompson, J. E. (1980) In vitro simulation of senescence-related membrane damage by ozone-induced lipid peroxidation. *Nature (London)* 283: 504-506.
- Pearce, D. W. (1993) Economic values and the natural world. London, United Kingdom: Earthscan Publications, Ltd.
- Pearson, M.; Mansfield, T. A. (1993) Interacting effects of ozone and water stress on the stomatal resistance of beech (*Fagus sylvatica* L.). *New Phytol.* 123: 351-358.
- Pearson, M.; Mansfield, T. A. (1994) Effects of exposure to ozone and water stress on the following season's growth of beech (*Fagus sylvatica* L.). *New Phytol.* 126: 511-515.
- Pearson, S.; Davison, A. W.; Reiling, K.; Ashenden, T.; Ollerenshaw, J. H. (1996) The effects of different ozone exposures on three contrasting populations of *Plantago major*. *New Phytol.* 132: 493-502.
- Pel, Z.-M.; Murata, Y.; Benning, G.; Thomine, S.; Klüsener, B.; Allen, G. J.; Grill, E.; Schroeder, J. I. (2000) Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling in guard cells. *Nature (London)* 406: 731-734.
- Pell, E. J.; Pearson, N. S. (1984) Ozone-induced reduction in quantity and quality of two potato cultivars. *Environ. Pollut. Ser. A* 35: 345-352.
- Pell, E. J.; Eckardt, N.; Enyedi, A. J. (1992) Timing of ozone stress and resulting status of ribulose biphosphate carboxylase/oxygenase and associated net photosynthesis. *New Phytol.* 120: 397-405.
- Pell, E. J.; Eckardt, N. A.; Glick, R. E. (1994) Biochemical and molecular basis for impairment of photosynthetic potential. *Photosynth. Res.* 39: 453-462.
- Pell, E. J.; Sinn, J. P.; Johansen, C. V. (1995) Nitrogen supply as a limiting factor determining the sensitivity of *Populus tremuloides* Michx. to ozone stress. *New Phytol.* 130: 437-446.
- Pell, E. J.; Schlagnhauser, C. D.; Arteca, R. N. (1997) Ozone-induced oxidative stress: mechanisms of action and reaction. *Physiol. Plant.* 100: 264-273.
- Pell, E. J.; Sinn, J. P.; Brendley, B. W.; Samuelson, L.; Vinten-Johansen, C.; Tien, M.; Skillman, J. (1999) Differential response of four tree species to ozone-induced acceleration of foliar senescence. *Plant Cell Environ.* 22: 779-790.
- Pellinen, R.; Palva, T.; Kangasj vi, J. (1999) Subcellular localization of ozone-induced hydrogen peroxide production in birch (*Betula pendula*) leaf cells. *Plant. J.* 20: 349-356.
- Pelloux, J.; Jolivet, Y.; Fontaine, V.; Banvoy, J.; Dizengremel, P. (2001) Changes in Rubisco and Rubisco activase gene expression and polypeptide content in *Pinus halepensis* M. subjected to ozone and drought. *Plant Cell Environ.* 24: 123-131.
- Penel, C. (1986) The role of calcium in the control of peroxidase activity. In: Greppin, H.; Penel, C.; Gasper, T., eds. *Molecular and physiological aspects of plant peroxidases*. Geneva, Switzerland: Universite de Genieve, Centre de Botanique; pp. 155-164.
- Percy, K. E.; Awmack, C. S.; Lindroth, R. L.; Kubiske, M. E.; Kopper, B. J.; Isebrands, J. G.; Pregitzer, K. S.; Hendry, G. R.; Dickson, R. E.; Zak, D. R.; Oksanen, E.; Sober, J.; Harrington, R.; Karnosky, D. F. (2002) Altered performance of forest pests under atmospheres enriched with CO₂ and O₃. *Nature (London)* 420: 403-407.
- Percy, K. E.; Karnosky, D. F.; Mankovska, B. (2003) Interactive O₃ and CO₂ effects on leaf surface physicochemical characteristics in three aspen clones: two-year results from the Aspen FACE experiment. In: Karnosky, D. F.; Percy, K. E.; Chappelka, A. H.; Simpson, C. J.; Pikkariainen, J., eds. *Air pollution, global change and forests in the new millenium*. New York, NY: Elsevier Science. (Developments in Environmental Science, 3).
- Persson, K.; Danielsson, H.; Seld n, G.; Pleijel, H. (2003) The effects of tropospheric ozone and elevated carbon dioxide on potato (*Solanum tuberosum* L. cv. Bintje) growth and yield. *Sci. Total Environ.* 310: 191-201.
- Peters, J. L.; Castillo, F. J.; Heath, R. L. (1988) Alteration of extracellular enzymes in pinto bean leaves upon exposure to air pollutants, ozone and sulfur dioxide. *Plant Physiol.* 89: 159-164.
- Peterson, D. L.; Arbaugh, M. J.; Wakefield, V. A.; Miller, P. R. (1987) Evidence of growth reduction in ozone-injured Jeffrey pine (*Pinus jeffreyi* Grev. and Balf.) in Sequoia and Kings Canyon National Parks. *JAPCA* 37: 906-912.
- Peterson, D. L.; Arbaugh, M. J.; Robinson, L. J. (1991) Regional growth changes in ozone-stressed ponderosa pine (*Pinus ponderosa*) in the Sierra Nevada, California, USA. *Holocene* 1: 50-61.
- Peterson, D. L.; Silsbee, D. G.; Poth, M.; Arbaugh, M. J.; Biles, F. E. (1995) Growth response of bigcone Douglas fir (*Pseudotsuga macrocarpa*) to long-term ozone exposure in southern California. *J. Air Waste Manage. Assoc.* 45: 36-45.
- Pfaffmann, T.; Barnes, J. D. (1993) Interactive effects of CO₂ and O₃ on growth of radish (*Raphanus sativus* L.). *Vegetatio* 104-105: 464-466.
- Phillips, R. L.; Zak, D. R.; Holmes, W. E.; White, D. C. (2002) Microbial community composition and function beneath temperate trees exposed to elevated atmospheric carbon dioxide and ozone. *Oecologia* 131: 236-244.

- 1 Pimm, S. L. (1997) The value of everything. *Nature* (London) 387: 231-232.
- 2 Pino, M. E.; Mudd, J. B.; Bailey-Serres, J. (1995) Ozone-induced alterations in the accumulation of newly synthesized
- 3 proteins in leaves of maize. *Plant Physiol.* 108: 777-785.
- 4 Pitcher, L. H.; Zilinskas, B. A. (1996) Overexpression of copper/zinc superoxide dismutase in the cytosol of transgenic
- 5 tobacco confers partial resistance to ozone-induced foliar necrosis. *Plant Physiol.* 110: 583-588.
- 6 Pitcher, L. H.; Brennan, E.; Zilinskas, B. A. (1992) The antiozonant ethylenediurea does not act via superoxide
- 7 dismutase induction in bean. *Plant Physiol.* 99: 1388-1392.
- 8 Pitelka, L. F. (1988) Evolutionary responses of plants to anthropogenic pollutants. *Trends in Ecology and Evolution* 3:
- 9 233-236.
- 10 Pleijel, H. (1998) A suggestion of a simple transfer function for the use of ozone monitoring data in dose-response
- 11 relationships obtained using open-top chambers. *Water Air Soil Pollut.* 102: 61-74.
- 12 Pleijel, H.; Danielsson, H. (1997) Growth of 27 herbs and grasses in relation to ozone exposure and plant strategy. *New*
- 13 *Phytol.* 135: 361-367.
- 14 Pleijel, H.; Skarby, L.; Ojanpera, K.; Selldén, G. (1994) Exposure of oats, *avena sativa* L., to filtered and unfiltered air
- 15 in open-top chambers: Effects on grain yield and quality. *Environ. Pollut.* 86: 129-134.
- 16 Pleijel, H.; Wallin, G.; Karlsson, P. E.; Skärby, L.; Selldén, G. (1994) Ozone deposition to an oat crop (*Avena sativa* L.)
- 17 grown in open-top chambers and in the ambient air. *Atmos. Environ.* 28: 1971-1979.
- 18 Pleijel, H.; Wallin, G.; Karlsson, P.; Skärby, L.; Selldén, G. (1995) Gradients of ozone at a forest site and over a field
- 19 crop - consequences for the AOT40 concept of critical level. *Water Air Soil Pollut.* 85: 2033-2038.
- 20 Pleijel, H.; Karlsson, G. P.; Danielsson, H.; Selldén, G. (1995) Surface wetness enhances ozone deposition to a pasture
- 21 canopy. *Atmos. Environ.* 29: 3391-3393.
- 22 Pleijel, H.; Karlsson, G. P.; Sild, E.; Danielsson, H.; Skärby, L.; Selldén, G. (1996) Exposure of a grass-clover mixture
- 23 to ozone in open-top chambers—effects on yield, quality and botanical composition *Agric. Ecosyst. Environ.* 59:
- 24 55-62.
- 25 Pleijel, H.; Ojanperä, K.; Mortensen, L. (1997) Effects of tropospheric ozone on the yield and grain protein content of
- 26 spring wheat (*Triticum aestivum* L.) in the nordic countries. *Acta Agric. Scand. Sect. B* 47: 20-25.
- 27 Pleijel, H.; Danielsson, H.; Gelang, J.; Sild, E.; Selldén, G. (1998) Growth stage dependence of the grain yield response
- 28 to ozone in spring wheat (*Triticum aestivum* L.). *Agric. Ecosyst. Environ.* 70: 61-68.
- 29 Pleijel, H.; Mortensen, L.; Fuhrer, J.; Ojanperä, K.; Danielsson, H. (1999) Grain protein accumulation in relation to
- 30 grain yield of spring wheat (*Triticum aestivum* L.) grown in open-top chambers with different concentrations of
- 31 ozone, carbon dioxide and water availability. *Agric. Ecosyst. Environ.* 72: 265-270.
- 32 Pleijel, H.; Norberg, P. A.; Selldén, G.; Skärby, L. (1999) Tropospheric ozone decreases biomass production in radish
- 33 plants (*Raphanus sativus*) grown in rural south-west Sweden. *Environ. Pollut.* 106: 143-147.
- 34 Pleijel, H.; Danielsson, H.; Karlsson, G. P.; Gelang, J.; Karlsson, P. E.; Selldén, G. (2000) An ozone flux-response
- 35 relationship for wheat. *Environ. Pollut.* 109: 453-462.
- 36 Pleijel, H.; Gelang, J.; Sild, E.; Danielsson, H.; Younis, S.; Karlsson, P.-E.; Wallin, G.; Skärby, L.; Selldén, G. (2000)
- 37 Effects of elevated carbon dioxide, ozone and water availability on spring wheat growth and yield. *Physiol. Plant.*
- 38 108: 61-70.
- 39 Pleijel, H.; Danielsson, H.; Vandermeiren, K.; Blum, C.; Colls, J.; Ojanperä, K. (2002) Stomatal conductance and ozone
- 40 exposure in relation to potato tuber yield—results from the European CHIP programme. *Eur. J. Agron.* 17:
- 41 303-317.
- 42 Plöchl, M.; Lyons, T.; Ollerenshaw, J.; Barnes, J. (2000) Simulating ozone detoxification in the leaf apoplast through
- 43 the direct reaction with ascorbate. *Planta* 210: 454-467.
- 44 Polasky, S., ed. (2001) The economics of biodiversity conservation. Burlington, VT: Ashgate Publishers.
- 45 Polle, A.; Pell, E. J. (1999) Role of carbon dioxide in modifying the plant response to ozone. In: Luo, Y.; Mooney, H.
- 46 A., eds. Carbon dioxide and environmental stress. San Diego, CA: Academic Press; pp. 193-213.
- 47 Polle, A.; Rennenberg, H. (1991) Superoxide dismutase activity in needles of Scots pine and Norway spruce under field
- 48 and changer conditions: lack of ozone effects. *New Phytol.* 117: 335-343.
- 49 Polle, A.; Krings, B.; Rennenberg, H. (1989) Superoxide dismutase activity in needles of Norwegian spruce trees (*Picea*
- 50 *abies* L.) *Plant Physiol.* 90: 1310-1315.
- 51 Poorter, H.; Pérez-Soba, M. (2001) The growth response of plants to elevated CO₂ under non-optimal environmental
- 52 conditions. *Oecologia* 129: 1-20.

- 1 Posthumus, A. C. (1998) Air pollution and global change: significance and perspectives. In: De Kok, L. J.; Stulen, I.,
2 eds. Responses of plant metabolism to air pollution and global change. Leiden, The Netherlands: Backhuys
3 Publishers; pp. 3-14.
- 4 Poth, M. A.; Fenn, M. E. (1998) Mature ponderosa pine nutrient use and allocation responses to air pollution. In:
5 Bytnerowicz, A.; Arbaugh, M. J.; Schilling, S. L. Proceedings of the International Symposium on Air Pollution
6 and Climate Change Effects on Forest Ecosystems; February, 1996; Riverside, CA. Albany, CA: USDA Forest
7 Service, Pacific Southwest General Technical Report no. PSW-GTR 166; pp. 239-247.
- 8 Potter, L.; Foot, J. P.; Caporn, S. J. M.; Lee, J. A. (1996) The effects of long-term elevated ozone concentrations on the
9 growth and photosynthesis of *Sphagnum recurvum* and *Polytrichum commune*. New Phytol. 134: 649-656.
- 10 Powell, M. C.; Muntifering, R. B.; Lin, J. C.; Chappelka, A. H. (2003) Yield and nutritive quality of sericea lespedeza
11 (*Lespedeza cuneata*) and little bluestem (*Schizachyrium scoparium*) exposed to ground-level ozone. Environ.
12 Pollut. 122: 313-322.
- 13 Pronos, J.; Merrill, L.; Dahlsten, D. (1999) Insects and pathogens in a pollution-stressed forest. In: Miller, P. R.;
14 McBride, J. R., eds. Oxidant air pollution impacts in the montane forests of southern California. New York, NY:
15 Springer, pp. 317-337.
- 16 Prozherina, N.; Freiwald, V.; Rousi, M.; Oksanen, E. (2003) Interactive effect of springtime frost and elevated ozone on
17 early growth, foliar injuries and leaf structure of birch (*Betula pendula*). New Phytol. 159: 623-636.
- 18 Pryor, W. A. (1992) How far does ozone penetrate into the pulmonary air/tissue boundary before it reacts? Free Radic.
19 Biol. Med. 12: 83-88.
- 20 Qui, Z.; Chappelka, A. H.; Somers, G. L.; Lockaby, B. G.; Meldahl, R. S. (1993) Effects of ozone and simulated acidic
21 precipitation on ectomycorrhizal formation on loblolly pine seedlings. Environ. Exp. Bot. 33: 423-431.
- 22 Rabinowitch, E. I. (1945) Photosynthesis and related processes. Vol. 1: chemistry of photosynthesis, chemosynthesis
23 and related processes in vitro and in vivo. New York, NY: Interscience Publishing, Inc.
- 24 Rafarel, C. R.; Ashenden, T. W. (1991) A facility for the large-scale exposure of plants to gaseous atmospheric
25 pollutants. New Phytol. 117: 345-349.
- 26 Randall, A. (1991) The value of biodiversity. Ambio 20: 64-68.
- 27 Rao, M.; Davis, K. R. (1999) Ozone-induced cell death occurs via two distinct mechanisms in *Arabidopsis*: the role of
28 salicylic acid. Plant J. 17: 603-614.
- 29 Rao, M. V.; Hale, B. A.; Ormrod, D. P. (1995) Amelioration of ozone-induced oxidative damage in wheat plants grown
30 under high carbon dioxide. Role of antioxidant enzymes. Plant Physiol. 109: 421-432.
- 31 Rawlings, J. O.; Lesser, V. M.; Heagle, A. S.; Heck, W. W. (1988) Alternative ozone dose metrics to characterize ozone
32 impact on crop yield loss. J. Environ. Qual. 17: 285-291.
- 33 Read, J. J.; Morgan, J. A. (1996) Growth and partitioning in *Pascopyrum smithii* (C3) and *Bouteloua gracilis* (C4) as
34 influenced by carbon dioxide and temperature. Ann. Bot. (London, UK) 77: 487-496.
- 35 Reams, G. A.; Van Deusen, P. C.; Lucier, A. A. (1995) Ambient ozone and loblolly pines. Nature (London) 378:
36 449-450.
- 37 Rebbeck, J. (1996) Chronic ozone effects on three northeastern hardwood species: growth and biomass. Can. J. For.
38 Res. 26: 1788-1798.
- 39 Rebbeck, J.; Jensen, K. F.; Greenwood, M. S. (1993) Ozone effects on grafted mature and juvenile red spruce:
40 photosynthesis, stomatal conductance, and chlorophyll concentration. Can. J. For. Res. 23: 450-456.
- 41 Reddy, G. N.; Arteca, R. N.; Dai, Y.-R.; Flores, H. E.; Negm, F. B.; Pell, E. J. (1993) Changes in ethylene and
42 polyamines in relation to mRNA levels of the large and small subunits of ribulose biphosphate
43 carboxylase/oxygenase in ozone-stressed potato foliage. Plant Cell Environ. 16: 819-826.
- 44 Reich, P. B. (1983) Effects of low concentrations of O₃ on net photosynthesis, dark respiration, and chlorophyll contents
45 in aging hybrid poplar leaves. Plant Physiol. 73: 291-296.
- 46 Reich, P. B.; Lassoie, J. P. (1984) Effects of low level O₃ exposure on leaf diffusive conductance and water-use
47 efficiency in hybrid poplar. Plant Cell Environ. 7: 661-668.
- 48 Reichenauer, T. G.; Goodman, B. A.; Kostecki, P.; Soja, G. (1998) Ozone sensitivity in *Triticum durum* and *T. aestivum*
49 with respect to leaf injury, photosynthetic activity and free radical content. Physiol. Plant. 104: 681-686.
- 50 Reid, C. D.; Fiscus, E. L. (1998) Effects of elevated [CO₂] and/or ozone on limitations to CO₂ assimilation in soybean
51 (*Glycine max*). J. Exp. Bot. 49: 885-895.
- 52 Reid, C. D.; Fiscus, E. L.; Burkey, K. O. (1998) Combined effects of chronic ozone and elevated CO₂ on rubisco
53 activity and leaf components in soybean (*Glycine max*). J. Exp. Bot. 49: 1999-2011.

- 1 Reiling, K.; Davison, A. W. (1992a) Spatial variation in ozone resistance of British populations of *Plantago major* L.
2 New Phytol. 122: 699-708.
- 3 Reiling, K.; Davison, A. W. (1992b) Effects of a short ozone exposure given at different stages in the development of
4 *Plantago major* L. New Phytol. 121: 643-647.
- 5 Reiling, K.; Davison, A. W. (1992c) The response of native, herbaceous species to ozone: growth and fluorescence
6 screening. New Phytol. 120: 29-37.
- 7 Reiling, K.; Davison, A. W. (1994) Effects of exposure to ozone at different stages in the development of *Plantago*
8 *major* L. on chlorophyll fluorescence and gas exchange. New Phytol. 128: 509-514.
- 9 Reimer, J.; Whittaker, J. B. (1989) Air pollution and insect herbivores: observed interactions and possible mechanisms.
10 In: Bernays, E. A., ed. Insect-plant interactions, v. 1. Boca Raton, FL: CRC Press, pp. 73-105.
- 11 Reiner, S.; Wiltshire, J. J. J.; Wright, C. J.; Colls, J. J. (1996) The impact of ozone and drought on the water relations of
12 ash trees (*Fraxinus excelsior* L.). J. Plant Physiol. 148: 166-171.
- 13 Reinert, R. A.; Ho, M. C. (1995) Vegetative growth of soybean as affected by elevated carbon dioxide and ozone.
14 Environ. Pollut. 89: 89-96.
- 15 Reinert, R. A.; Eason, G.; Barton, J. (1997) Growth and fruiting of tomato as influenced by elevated carbon dioxide and
16 ozone. New Phytol. 137: 411-420.
- 17 Renaud, J. P.; Allard, G.; Mauffette, Y. (1997) Effects of ozone on yield, growth, and root starch concentrations of two
18 alfalfa (*Medicago sativa* L.) cultivars. Environ. Pollut. 95: 273-281.
- 19 Retzlaff, W. A.; Weinstein, D. A.; Laurence, J. A.; Gollands, B. (1996) Simulated root dynamics of a 160-year-old sugar
20 maple (*Acer saccharum* Marsh.) tree with and without ozone exposure using the TREGRO model. Tree Physiol.
21 16: 915-921.
- 22 Retzlaff, W. A.; Williams, L. E.; DeJong, T. M. (1997) Growth and yield response of commercial bearing-age
23 'Casselman' plum trees to various ozone partial pressures. J. Environ. Qual. 26: 858-865.
- 24 Retzlaff, W. A.; Arthur, M. A.; Grulke, N. E.; Weinstein, D. A.; Gollands, B. (2000) Use of a single-tree simulation
25 model to predict effects of ozone and drought on growth of a white fir tree. Tree Physiol. 20: 195-202.
- 26 Ribas, A.; Peñuelas, J. (2000) Effects of ethylene diurea as a protective antiozonant on beans (*Phaseolus vulgaris* cv lit)
27 exposed to different tropospheric ozone doses in Catalonia (NE Spain). Water Air Soil Pollut. 117: 263-271.
- 28 Rich, S.; Ames, R.; Zukel, J. W. (1974) 1,4-oxanthin derivatives protect plants against ozone. Plant Dis. Rep. 58:
29 162-164.
- 30 Robinson, J. M.; Britz, S. J. (2000) Tolerance of a field grown soybean cultivar to elevated ozone level is concurrent with
31 higher leaflet ascorbic acid level, higher ascorbate-dehydroascorbate redox status, and long term photosynthetic
32 productivity. Photosynth. Res. 64: 77-87.
- 33 Robinson, M. F.; Heath, J.; Mansfield, T. A. (1998) Disturbance in stomatal behaviour caused by air pollutants. J. Exp.
34 Bot. 49: 461-469.
- 35 Roitsch, T. (1999) Source-sink regulation by sugar and stress. Curr. Opin. Plant Biol. 2: 198-206.
- 36 Rondón, A.; Johansson, C.; Granat, L. (1993) Dry deposition of nitrogen-dioxide and ozone to coniferous forests. J.
37 Geophys. Res. [Atmos.] 98(D3): 5159-5172.
- 38 Rosen, P. M.; Runeckles, V. C. (1976) Interaction of ozone and greenhouse whitefly in plant injury. Environ. Conserv. 3:
39 70-71.
- 40 Rötter, R.; Van De Geijn, S. C. (1999) Climate change effects on plant growth, crop yield and livestock. Clim. Change
41 43: 651-681.
- 42 Rowland-Bamford, A. J. (2000) Plant responses to changing carbon dioxide and temperature. In: Singh, S. N., ed. Trace
43 gas emissions and plants. Dordrecht, The Netherlands: Kluwer Academic Publishers; pp. 63-74.
- 44 Rudorff, B. F. T.; Mulchi, C. L.; Daughtry, C. S. T.; Lee, E. H. (1996) Growth, radiation use efficiency, and canopy
45 reflectance of wheat and corn grown under elevated ozone and carbon dioxide atmospheres. Remote Sens.
46 Environ. 55: 163-173.
- 47 Rudorff, B. F. T.; Mulchi, C. L.; Fenny, P.; Lee, E. H.; Rowland, R. (1996) Wheat grain quality under enhanced
48 tropospheric CO₂ and O₃ concentrations. J. Environ. Qual. 25: 1384-1388.
- 49 Rudorff, B. F. T.; Mulchi, C. L.; Lee, E. H.; Rowland, R.; Pausch, R. (1996) Effects of enhanced O₃ and CO₂ enrichment
50 on plant characteristics in wheat and corn. Environ. Pollut. 94: 53-60.
- 51 Rudorff, B. F. T.; Mulchi, C. L.; Lee, E. H. (2000) Plant responses to elevated CO₂ and interactions with O₃. In: Singh, S.
52 N., ed. Trace gas emissions and plants. Dordrecht, The Netherlands: Kluwer Academic Publishers; pp. 155-179.
- 53 Runeckles, V. C. (1974) Dosage of air pollutants and damage to vegetation. Environ. Conserv. 1: 305-308.

- 1 Runeckles, V. C. (2002) Air pollution and climate change. In: Bell, J. N. B.; Treshow, M., eds. Air Pollution and Plant
2 Life. 2nd ed. Chichester, United Kingdom: John Wiley & Sons Ltd.; pp. 431-454.
- 3 Runeckles, V. C.; Bowen, P. A. (2000) The use of calibrated passive monitors to assess crop loss due to ozone in rural
4 locations. In: Agrawal, S. B.; Agrawal, M., eds. Environmental pollution and plant responses. Boca Raton, FL:
5 Lewis Publishers; pp. 307-316.
- 6 Runeckles, V. C.; Krupa, S. V. (1994) The impact of UV-B radiation and ozone on terrestrial vegetation. Environ. Pollut.
7 83: 191-213.
- 8 Runeckles, V. C.; Palmer, K. (1987) Pretreatment with nitrogen dioxide modifies plant response to ozone. Atmos.
9 Environ. 21: 717-719.
- 10 Runeckles, V. C.; Rosen, P. M. (1974) Effects of pretreatment with low ozone concentrations on ozone injury to bean
11 and mint. Can. J. Bot. 52: 2607-2610.
- 12 Runeckles, V. C.; Vaartnou, M. (1997) EPR evidence for superoxide anion formation in leaves during exposure to low
13 levels of ozone. Plant Cell Environ. 20: 306-314.
- 14 Runeckles, V. C.; Wright, E. F. (1996) Delayed impact of chronic ozone stress on young Douglas-fir grown under field
15 conditions. Can. J. For. Res. 26: 629-638.
- 16 Runeckles, V. C.; Wright, E. F.; White, D. (1990) A chamberless field exposure system for determining the effects of
17 gaseous air pollutants on crop growth and yield. Environ. Pollut. 63: 61-77.
- 18 Rusch, H.; Laurence, J. A. (1993) Interactive effects of ozone and powdery mildew on pea seedlings. Phytopathology 83:
19 1258-1263.
- 20 Sah, D. N.; Von Tiedemann, A.; Fehrmann, H. (1993) Effects of ozone exposure on resistance of wheat genotypes to
21 *Phrenophora tritici-repentis*. J. Phytopathol. 138: 249-256.
- 22 Sakaki, T.; Kondo, N.; Sugahara, K. (1983) Breakdown of photosynthetic pigments and lipids in spinach leaves with
23 ozone fumigation: role of active oxygens. Physiol. Plant. 59: 28-34.
- 24 Sakaki, T.; Ohnishi, J.-I.; Kondo, N.; Yamada, M. (1985) Polar and neutral lipid changes in spinach leaves with ozone
25 fumigation: triacylglycerol synthesis from polar lipids. Plant Cell Physiol. 26: 253-262.
- 26 Sakaki, T.; Saito, K.; Kawaguchi, A.; Kondo, N.; Yamada, M. (1990) Conversion of monogalactosylidacylglycerol to
27 triacylglycerols in ozone-fumigated spinach leaves. Plant Physiol. 94: 766-772.
- 28 Salardino, D. H.; Carroll, J. J. (1998) Correlation between ozone exposure and visible foliar injury in ponderosa and
29 Jeffrey pines. Atmos. Environ. 32: 3001-3010.
- 30 Saleem, A.; Loponen, J.; Pihlaja, K.; Oksanen, E. (2001) Effects of long-term open-field ozone exposure on leaf
31 phenolics of European silver birch (*Betula pendula* Roth). J. Chem. Ecol. 27: 1049-1062.
- 32 Salisbury, F. B.; Ross, C. (1969) Plant physiology. Belmont, CA: Wadsworth Publishing Company, Inc.
- 33 Samuelson, L. J. (1994) Ozone-exposure responses of black cherry and red maple seedlings. Environ. Exp. Bot. 34:
34 355-362.
- 35 Samuelson, L. (2001) Relating ozone uptake in forest trees to risk assessment. Human Ecol. Risk Assess. 7: 1161-1167.
- 36 Samuelson, L. J.; Edwards, G. S. (1993) A comparison of sensitivity to ozone in seedlings and trees of *Quercus rubra* L.
37 New Phytol. 125: 373-379.
- 38 Samuelson, L. J.; Kelly, J. M. (1996) Carbon partitioning and allocation in northern red oak seedlings and mature trees in
39 response to ozone. Tree Physiol. 16: 853-858.
- 40 Samuelson, L. J.; Kelly, J. M. (1997) Ozone uptake in *Prunus serotina*, *Acer rubrum* and *Quercus rubra* forest trees of
41 different sizes. New Phytol. 136: 255-264.
- 42 Samuelson, L.; Kelly, J. M. (2001) Scaling ozone effects from seedlings to forest trees. Tansley review no. 21. New
43 Phytol. 149: 21-41.
- 44 Samuelson, L. J.; Kelly, J. M.; Mays, P. A.; Edwards, G. S. (1996) Growth and nutrition of *Quercus rubra* L. seedlings
45 and mature trees after three seasons of ozone exposure. Environ. Pollut. 91: 317-323.
- 46 Sandermann, H., Jr. (1996) Ozone and plant health. Annu. Rev. Phytopathol. 34: 347-366.
- 47 Sandermann, H., Jr. (1998) Ozone: an air pollutant acting as a plant-signaling molecule. Naturwissenschaften 85:
48 369-375.
- 49 Sandermann, H., Jr. (2000) Ozone/biotic disease interactions: molecular biomarkers as a new experimental tool. Environ.
50 Pollut. 108: 327-332.
- 51 Sanders, G. E.; Skärby, L.; Ashmore, M. R.; Fuhrer, J. (1995) Establishing critical levels to the effects of air pollution on
52 vegetation. Water Air Soil Pollut. 85: 189-200.
- 53 Saxe, H.; Ellsworth, D. S.; Heath, J. (1998) Tree and forest functioning in an enriched CO₂ atmosphere. New Phytol.
54 139: 395-436.

- 1 Sayre, R. G.; Fahey, T. J. (1999) Effects of rainfall acidity and ozone on foliar leaching in red spruce *Picea rubens*. Can.- 2 J. For. Res. 29: 486-496.
- 3 Scagel, C. F.; Andersen, C. P. (1997) Seasonal changes in root and soil respiration of ozone-exposed ponderosa pine
4 (*Pinus ponderosa*) grown in different substrates. New Phytol. 136: 627-643.
- 5 Scandalios, J. G. (1993) Oxygen stress and superoxide dismutases. Plant Physiol. 101: 7-12.
- 6 Schaub, M.; Skelly, J. M.; Steiner, K. C.; Davis, D. D.; Pennypacker, S. P.; Zhang, J.; Ferdinand, J. A.; Savage, J. E.;
7 Stevenson, R. E. (2003) Physiological and foliar injury responses of *Prunus serotina*, *Fraxinus americana*, and
8 *Acer rubrum* seedlings to varying soil moisture and ozone. Environ. Pollut. 124: 307-320.
- 9 Scherzer, A. J.; Rebbeck, J.; Boerner, R. E. J. (1998) Folia nitrogen dynamics and decomposition of yellow-poplar and
10 eastern white pine during four seasons of exposure to elevated ozone and carbon dioxide. For. Ecol. Manage. 109:
11 355-366.
- 12 Schmieden, U.; Wild, A. (1995) The contribution of ozone to forest decline. Physiol. Plant. 94: 371-378.
- 13 Schnitzler, J.-P.; Langebartels, C.; Heller, W.; Liu, J.; Lippert, M.; Dohring, T.; Bahnweg, G.; Sandermann, H. (1999)
14 Ameliorating effect of UV-B radiation on the response of Norway spruce and Scots pine to ambient ozone
15 concentrations. Global Change Biol. 5: 83-94.
- 16 Schnug, E. (1998) Response of plant metabolism to air pollution and global change - impact on agriculture. In: De Kok,
17 L. J.; Stulen, I., eds. Responses of plant metabolism to air pollution and global change. Leiden, The Netherlands:
18 Backhuys Publishers; pp. 15-22.
- 19 Schopfer, P. (1994) Histochemical demonstration and localization of H₂O₂ in organs of higher plants by tissue printing
20 on nitrocellulose paper. Plant Physiol. 104: 1269-1275.
- 21 Schraudner, M.; Ernst, D.; Langebartels, C.; Sandermann, H., Jr. (1992) Biochemical plant responses to ozone. III.
22 Activation of the defense-related proteins β -1,3-glucanase and chitinase in tobacco leaves. Plant Physiol. 99:
23 1321-1328.
- 24 Schraudner, M.; Langebartels, C.; Sandermann, H., Jr. (1996) Plant defence systems and ozone. Biochem. Soc. Trans.
25 24: 456-461.
- 26 Schulte-Hosted, S.; Darrall, N. M.; Blank, L. W.; Wellburn, A. R. (1988) Air pollution and plant metabolism. New York,
27 NY: Elsevier Applied Science.
- 28 Segerson, K. (1987) Impacts of ozone and acid rain: discussion. Am. J. Agric. Econ. 69: 970-971.
- 29 Segerson, K. (1991) Air pollution and agriculture: a review and evaluation of policy interactions. In: Just, R. E.;
30 Bockstael, N., eds. Commodity and resource policies in agricultural systems. Berlin, Germany: Springer-Verlag;
31 pp. 349-367.
- 32 Seufert, G.; Arndt, U. (1985) Open-top-Kammern als teil eines konzepts zur ökosystemaren untersuchung der neuartigen
33 waldschaden. Allg. Forstz. 40: 13-18.
- 34 Shan, Y.; Feng, Z.; Izuta, T.; Aoki, M.; Totsuka, T. (1996) The individual and combined effects of ozone and simulated
35 acid rain on growth, gas exchange and water-use efficiency of *Pinus armandi* Franch. Environ. Pollut. 91:
36 355-361.
- 37 Sharma, P.; Sôber, A.; Sôber, J.; Podila, G. K.; Kubiske, M. E.; Mattson, W. J.; Isebrands, J. G.; Karnosky, D. F. (2003)
38 Moderation of [CO₂]-induced gas exchange responses by elevated tropospheric O₃ in trembling aspen and sugar
39 maple. Ekologia (Journal of Ecology) (Bratislava): in press.
- 40 Shaw, M. R.; Zavaleta, E. S.; Chiariello, N. R.; Cleland, E. E.; Mooney, H. A.; Field, C. B. (2002) Grassland responses
41 to global environmental changes suppressed by elevated CO₂. Science (Washington, DC, U.S.) 298: 1987-1990.
- 42 Sheng, Y.; Podila, G. K.; Karnosky, D. F. (1997) Differences in O₃-induced superoxide dismutase and glutathione
43 antioxidant expression in O₃ tolerant and sensitive trembling aspen (*Populus tremuloides* Michx.) clones. For.
44 Gen. 4: 25-33.
- 45 Sheppard, L. J.; Cape, J. N. (1999) Forest growth responses to the pollution climate of the 21st century: proceedings of
46 the 18th IUFRO workshop on air pollution stress; 1998; Edinburgh, Scotland. Water Air Soil Pollut. 116(1-2
47 special issue).
- 48 Sheppard, L. J.; Smith, R. I.; Cannell, M. G. R. (1989) Frost hardiness of *Picea rubens* growing in spruce decline regions
49 of the Appalachians. Tree Physiol. 5: 25-37.
- 50 Showman, R. E. (1991) A comparison of ozone injury to vegetation during moist and drought years. J. Air Waste
51 Manage. Assoc. 41: 63-64.
- 52 Sild, E.; Pleijel, H.; Selldén, G. (1999) Response of subterranean clover (*Trifolium subterraneum*) to ozone in relation to
53 plant age and light conditions during exposure. New Phytol. 143: 315-321.

- 1 Simini, M.; Skelly, J. M.; Davis, D. D.; Savage, J. E.; Comrie, A. C. (1992) Sensitivity of four hardwood species to
2 ambient ozone in north central Pennsylvania. *Can. J. For. Res.* 22: 1789-1799.
- 3 Simon-Plas, F.; Rustérucci, C.; Milat, M.-L.; Humbert, C.; Montillet, J.-L.; Blein, J.-P. (1997) Active oxygen species
4 production in tobacco cells elicited by cryptogein. *Plant Cell Environ.* 20: 1573-1579.
- 5 Simpson, D.; Tuovinen, J.-P.; Emberson, L.; Ashmore, M.R. (2001) Characteristics of an ozone deposition module.
6 *Water Air Soil Pollut. FOCUS* 1: 253-262.
- 7 Simpson, D.; Tuovinen, J.-P.; Emberson, L.; Ashmore, M.R. (2003) Characteristics of an ozone deposition module II:
8 Sensitivity analysis. *Water Air Soil Pollut.* 143: 123-137.
- 9 Singh, S. N., ed. (2000) Trace gas emissions and plants. Dordrecht, The Netherlands: Kluwer Academic Publishers.
- 10 Sivasankar, S.; Sheldrick, B.; Rothstein, S. J. (2000) Expression of allene oxide synthase determines defense gene
11 activation in tomato. *Plant Physiol.* 122: 1335-1342.
- 12 Skärby, L.; Ro-Poulsen, H.; Wellburn, F. A. M.; Sheppard, L. J. (1998) Impacts of ozone on forests: a European
13 perspective. *New Phytol.* 139: 109-122.
- 14 Skeffington, R. A.; Sutherland, P. M. (1995) The effects of SO₂ and O₃ fumigation on acid deposition and foliar leaching
15 in the Liphook forest fumigation experiment. *Plant Cell Environ.* 18: 247-261.
- 16 Skelly, J. M. (1988) Forest decline versus tree decline the pathological considerations. *Environ. Monit. Assess.* 12:
17 23-27.
- 18 Skelly, J. M. (2000) Tropospheric ozone and its importance to forests and natural plant communities of the northeastern
19 United States. *Northeast. Nat.* 7: 221-236.
- 20 Skelly, J. M. (2003) Native plants as bioindicators of air pollutants: contributed papers to a symposium held in
21 conjunction with the 34th air pollution workshop. *Environ. Pollut.* 125: 1-2.
- 22 Smeulders, S. M.; Gorissen, A.; Joosten, N. N.; Van Veen, J. A. (1995) Effects of short-term ozone exposure on the
23 carbon economy of mature and juvenile Douglas firs [*Pseudotsuga menziesii* (Mirb.) Franco]. *New Phytol.* 129:
24 45-53.
- 25 Smirnoff, N.; Conklin, P. L.; Loewus, F. A. (2001) Biosynthesis of ascorbate acid in plants: a renaissance. *Annu. Rev.*
26 *Plant Physiol. Mol. Biol.* 52: 437-467.
- 27 Smith, W. B. (2002) Forest inventory and analysis: a national inventory and monitoring program. *Environ. Pollut.* 116:
28 S233-S242.
- 29 Smith, G.; Greenhalgh, B.; Brennan, E.; Justin, J. (1987) Soybean yield in New Jersey relative to ozone pollution and
30 antioxidant application. *Plant Dis.* 71: 121-125.
- 31 Smith, G.; Coulston, J.; Jepsen, E.; Prichard, T. (2003) A national ozone biomonitoring program - results from field
32 surveys of ozone sensitive plants in northeastern forests (1994-2000). *Environ. Monit. Assess.* 87: 271-291.
- 33 Soja, G.; Eid, M.; Gangl, H.; Redl, H. (1997) Ozone sensitivity of grapevine (*Vitis vinifera* L.): evidence for a memory
34 effect in a perennial crop plant? *Phyton (Austria)* 37: 265-270.
- 35 Soja, G.; Barnes, J. D.; Posch, M.; Vandermeiren, K.; Pleijel, H.; Mills, G. (2000) Phenological weighting of ozone
36 exposures in the calculation of critical levels for wheat, bean and plantain. *Environ. Pollut.* 109: 517-524.
- 37 Somers, G. L.; Chappelka, A. H.; Rosseau, P.; Renfro, J. R. (1998) Empirical evidence of growth decline related to
38 visible ozone injury. *For. Ecol. Manage.* 104: 129-137.
- 39 Spash, C. L. (1997) Assessing the economic benefits to agriculture from air pollution control. *J. Econ. Surv.* 11: 47-70.
- 40 Stan, H.-J.; Schicker, S. (1982) Effect of repetitive ozone treatment on bean plants - stress ethylene production and leaf
41 necrosis. *Atmos. Environ.* 16: 2267-2270.
- 42 Stan, H.-J.; Schicker, S.; Kassner, H. (1981) Stress ethylene evolution of bean plants - a parameter indicating ozone
43 pollution. *Atmos. Environ.* 15: 391-395.
- 44 Stark, R. W.; Cobb, F. W., Jr. (1969) Smog injury, root diseases and bark beetle damage in ponderosa pine. *Calif. Agric.*
45 23(9): 13-15.
- 46 Stark, R. W.; Miller, P. R.; Cobb, F. W., Jr.; Wood, D. L.; Parmeter, J. R., Jr. (1968) Photochemical oxidant injury and
47 bark beetle (Coleoptera: Scolytidae) infestation of ponderosa pine. I. Incidence of bark beetle infestation in injured
48 trees. *Hilgardia* 39: 121-126.
- 49 Staszak, J.; Grulke, N. E.; Prus-Glowacki, W. (2004) Genetic differences of *Pinus ponderosa* (Dougl. Ex Laws) trees
50 tolerant and sensitive to ozone. *Water Air Soil Pollut.* 153: 3-14.
- 51 Stewart, C. A. (1998) Impact of ozone on the reproductive biology of *Brassica campestris* L. and *Plantago major* L
52 [dissertation]. Leicestershire, United Kingdom: University of Loughborough, Department of Geology.
- 53 Stewart, C. A.; Black, V. J.; Black, C. R.; Roberts, J. A. (1996) Direct effects of ozone on the reproductive development
54 of *Brassica species*. *J. Plant Physiol.* 148: 172-178.

- 1 Stitt, M. (1996) Plasmodesmata play an essential role in sucrose export from leaves: a step towards an integration of
- 2 metabolic biochemistry and cell biology. *Plant Cell* 8: 565-571.
- 3 Stribley, G. H.; Ashmore, M. R. (2002) Quantitative changes in twig growth pattern of young woodland beech (*Fagus*
- 4 *sylvatica* L.) in relation to climate and ozone pollution over 10 years. *For. Ecol. Manage.* 157: 191-204.
- 5 Sullivan, J. A.; Hale, B. A.; Ormrod, D. P. (1994) Impact of seasonal ozone exposure on yield and vegetative growth of
- 6 primocane-fruiting raspberry. *HortScience* 29: 1059-1061.
- 7 Suter, G. W., II. (1990) Endpoints for regional ecological risk assessments. *Environ. Manage.* (N. Y.) 14: 9-23.
- 8 Swank, W. T.; Vose, J. M. (1990) Watershed-scale responses to ozone events in a *Pinus strobus* L. plantation. *Water Air*
- 9 *Soil Pollut.* 54: 119-133.
- 10 Taiz, L.; Zeiger, E. (2002) *Plant physiology*. 3rd ed. Sunderland, MA: Sinauer Associates, Publishers.
- 11 Takemoto, B. K.; Olszyk, D. M.; Johnson, A. G.; Parada, C. R. (1988) Yield responses of field-grown crops to acidic fog
- 12 and ambient ozone. *J. Environ. Qual.* 17: 192-197.
- 13 Takemoto, B. K.; Bytnerowicz, A.; Dawson, P. J.; Morrison, C. L.; Temple, P. J. (1997) Effects of ozone on *Pinus*
- 14 *ponderosa* seedlings: comparison of responses in the first and second growing seasons of exposure. *Can. J. For.*
- 15 *Res.* 27: 23-30.
- 16 Takemoto, B. K.; Bytnerowicz, A.; Fenn, M. E. (2001) Current and future effects of ozone and atmospheric nitrogen
- 17 deposition on California's mixed conifer forests. *For. Ecol. Manage.* 144: 159-173.
- 18 Takeuchi, Y.; Kubiske, M. E.; Isebrands, J. G.; Pregitzer, K. S.; Hendrey, G.; Karnosky, D. F. (2001) Photosynthesis,
- 19 light and nitrogen relationships in a young deciduous forest canopy under open-air CO₂ enrichment. *Plant Cell*
- 20 *Environ.* 24: 1257-1268.
- 21 Tappel, A. L. (1972) Vitamin E and free radical peroxidation of lipids. In: Nair, P. P.; Kayden, H. J., eds. *Vitamin E and*
- 22 *its role in cellular metabolism*. New York, NY: New York Academy of Sciences; pp. 12-28. (*Annals of the New*
- 23 *York Academy of Sciences*: v. 203).
- 24 Tausz, M.; Bytnerowicz, A.; Arbaugh, M. J.; Weidner, W.; Grill, D. (1999a) Antioxidants and protective pigments of
- 25 *Pinus ponderosa* needles at gradients of natural stresses and ozone in the San Bernardino Mountains in California.
- 26 *Free Rad. Res.* 31(suppl.): S113-S120.
- 27 Tausz, M.; Bytnerowicz, A.; Weidner, W.; Arbaugh, M. J.; Padgett, P.; Grill, D. (1999b) Pigments and phytoprotection
- 28 in needles of *Pinus ponderosa* trees with and without symptoms of ozone injury. *Phyton* (Horn, Austria) 39:
- 29 219-224.
- 30 Tausz, M.; Bytnerowicz, A.; Weidner, W.; Arbaugh, M. J.; Padgett, P.; Grill, D. (1999c) Changes in free-radical
- 31 scavengers describe the susceptibility of *Pinus ponderosa* to ozone in southern Californian forests. *Water Air Soil*
- 32 *Pollut.* 116: 249-254.
- 33 Tausz, M.; Bytnerowicz, A.; Arbaugh, M. J.; Wonisch, A.; Grill, D. (2001) Multivariate patterns of biochemical
- 34 responses of *Pinus ponderosa* trees at field plots in the San Bernardino Mountains, southern California. *Tree*
- 35 *Physiol.* 21: 329-336.
- 36 Tausz, M.; Herbinger, K.; Posch, S.; Grulke, N. E. (2002) Antioxidant status of *Pinus jeffreyi* needles from mesic and
- 37 xeric microsites in early and late summer. *Phyton* (Horn, Austria) 42: 201-207.
- 38 Taylor, G. E., Jr.; Hanson, P. J. (1992) Forest trees and tropospheric ozone: role of canopy deposition and leaf uptake in
- 39 developing exposure-response relationships. *Agric. Ecosyst. Environ.* 42: 255-273.
- 40 Taylor, G. E., Jr.; Pitelka, L. F. (1992) Genetic diversity of plant populations and the role of air pollution. In: Barker, J.
- 41 R.; Tingey, D. T., eds. *Air pollution effects on biodiversity*. New York, NY: Van Nostrand Reinhold; pp. 111-130.
- 42 Taylor, G. E., Jr.; McLaughlin, S. B.; Shriner, D. S. (1982) Effective pollutant dose. In: Unsworth, M. H.; Ormrod, D. P.,
- 43 eds. *Effects of gaseous air pollution in agriculture and horticulture*. London, United Kingdom: Butterworth
- 44 Scientific; pp. 458-460.
- 45 Taylor, G. E., Jr.; Tingey, D. T.; Ratsch, H. C. (1982) Ozone flux in *Glycine max* (L.) Merr.: sites of regulation and
- 46 relationship to leaf injury. *Oecologia* 53: 179-186.
- 47 Taylor, G. E., Jr.; Hanson, P. J.; Baldocchi, D. D. (1988a) Pollutant deposition to individual leaves and plant canopies:
- 48 sites of regulation and relationship to injury. In: Heck, W. W.; Taylor, O. C.; Tingey, D. T., eds. *Assessment of*
- 49 *crop loss from air pollutants*. New York, NY: Elsevier Applied Science; pp. 227-257.
- 50 Taylor, G. E., Jr.; Ross-Todd, B. M.; Gunderson, C. A. (1988b) Action of ozone on foliar gas exchange in *Glycine max*
- 51 L. Merr.: a potential role for endogenous stress ethylene. *New Phytol.* 110: 301-307.
- 52 Taylor, G. E.; Owens, J. G.; Grizzard, T.; Selvidge, W. J. (1993) Atmosphere x canopy interactions of nitric acid vapor in
- 53 loblolly-pine grown in open-top chambers. *J. Environ. Qual.* 22: 70-80.

- 1 Taylor, G. E., Jr.; Johnson, D. W.; Andersen, C. P. (1994) Air pollution and forest ecosystems: a regional to global
2 perspective. *Ecol. Appl.* 4: 662-689.
- 3 Taylor, G.; Ceulemans, R.; Ferris, R.; Gardner, S. D. L.; Shao, B. Y. (2001) Increased leaf area expansion of hybrid
4 poplar in elevated CO₂. From controlled environments to open-top chambers and to FACE. *Environ. Pollut.* 115:
5 463-472.
- 6 Temple, P. J. (1988) Injury and growth of Jeffrey pine and giant sequoia in response to ozone and acidic mist. *Environ.*
7 *Exp. Bot.* 28: 323-333.
- 8 Temple, P. J. (1999) Effects of ozone on understory vegetation in the mixed conifer forest. In: Miller, P. R.; McBride, J.
9 R., eds. *Oxidant air pollution impacts in the montane forests of southern California: a case study of the San*
10 *Bernardino Mountains*. New York, NY: Springer; pp. 208-222. (Ecological studies: v. 134).
- 11 Temple, P. J.; Miller, P. R. (1994) Foliar ozone injury and radial growth of ponderosa pine. *Can. J. For. Res.* 24:
12 1877-1882.
- 13 Temple, P. J.; Riechers, G. H.; Miller, P. R. (1992) Foliar injury responses of ponderosa pine seedlings to ozone, wet and
14 dry acidic deposition, and drought. *Environ. Exp. Bot.* 32: 101-113.
- 15 Temple, P. J.; Riechers, G. H.; Miller, P. R.; Lennox, R. W. (1993) Growth responses of ponderosa pine to long-term
16 exposure to ozone, wet and dry acidic deposition, and drought. *Can. J. For. Res.* 23: 59-66.
- 17 Teramura, A. H.; Sullivan J. H.; Lydon, J. (1990) Effects of UV-b radiation on soybean yield and seed quality: a 6-year
18 field study. *Physiol. Plant.* 80: 5-11.
- 19 Teskey, R. O.; Dougherty, P. M.; Wiselogle, A. E. (1991) Design and performance of branch chambers suitable for
20 long-term ozone fumigation of foliage in large trees. *J. Environ. Qual.* 20: 591-595.
- 21 Thalmair, M.; Bauw, G.; Thiel, S.; Döhning, T.; Langebartels, C.; Sandermann, H., Jr. (1996) Ozone and ultraviolet B
22 effects on the defense-related proteins β -1,3-glucanase and chitinase in tobacco. *J. Plant Physiol.* 148: 222-228.
- 23 Thom, M.; Maretzki, A. (1985) Evidence for a plasmalemma redox system in sugarcane. *Plant Physiol.* 77: 873-876.
- 24 Tiedemann, A. V.; Firsching, K. H. (1993) Effects of ozone exposure and leaf age of wheat on infection processes of
25 *Septoria nodorum* Berk.. *Plant Pathol.* 42: 287-293.
- 26 Tiedemann, A. V.; Firsching, K. H. (2000) Interactive effects of elevated ozone and carbon dioxide on growth and yield
27 of leaf rust-infected versus non-infected wheat. *Environ. Pollut.* 108: 357-363.
- 28 Tiedemann, A. V.; Weigel, H.-J.; Jäger, H.-J. (1991) Effects of open-top chamber fumigations with ozone on three
29 fungal leaf diseases of wheat and the mycoflora of the phyllosphere. *Environ. Pollut.* 72: 205-224.
- 30 Tingey, D. T. (1984) The effects of ozone on plants in the United States. In: Grennfelt, P., ed. *Proceedings of the*
31 *international workshop on the evaluation of the effects of photochemical oxidants on human health, agricultural*
32 *crops, forestry, materials and visibility; February-March; Göteborg, Sweden*. Göteborg, Sweden: Swedish
33 *Environmental Research Institute*; pp. 60-75; report no. IVL-EM 1570.
- 34 Tingey, D. T.; Andersen, C. P. (1991) The physiological basis of differential plant sensitivity to changes in atmospheric
35 quality. In: Taylor, G. E., Jr.; Pitelka, L. F.; Clegg, M. T., eds. *Ecological genetics and air pollution*. New York,
36 NY: Springer-Verlag.
- 37 Tingey, D. T.; Olszyk, D. M. (1985) Intraspecific variability in metabolic responses to SO₂. In: Winner, W. E.; Mooney,
38 H. A.; Goldstein, R. A., eds. *Sulfur dioxide and vegetation: physiology, ecology, and policy issues*. Stanford, CA:
39 *Stanford University Press*; pp. 178-205.
- 40 Tingey, D. T.; Taylor, G. E., Jr. (1982) Variation in plant response to ozone: a conceptual model of physiological events.
41 In: Unsworth, M. H.; Ormrod, D. P., eds. *Effects of gaseous air pollution in agriculture and horticulture*. London,
42 *United Kingdom: Butterworth Scientific*; pp. 113-138.
- 43 Tingey, D. T.; Fites, R. C.; Wickliff, C. (1975) Activity changes in selected enzymes from soybean leaves following
44 ozone exposure. *Physiol. Plant.* 33: 316-320.
- 45 Tingey, D. T.; Standley, C.; Field, R. W. (1976a) Stress ethylene evolution: a measure of ozone effects on plants. *Atmos.*
46 *Environ.* 10: 969-974.
- 47 Tingey, D. T.; Wilhour, R. G.; Standley, C. (1976b) The effect of chronic ozone exposures on the metabolite content of
48 ponderosa pine seedlings. *For. Sci.* 22: 234-241.
- 49 Tingey, D. T.; Hogsett, W. E.; Lee, E. H. (1989) Analysis of crop loss for alternative ozone exposure indices. In:
50 Schneider, T.; Lee, S. D.; Wolters, G. J. R.; Grant, L. D., eds. *Atmospheric ozone research and its policy*
51 *implications: proceedings of the 3rd US-Dutch international symposium; May 1988; Nijmegen, The Netherlands*.
52 *Amsterdam, The Netherlands: Elsevier Science Publishers*; pp. 219-227. (Studies in environmental science 35).
- 53 Tingey, D. T.; Hogsett, W. E.; Lee, E. H.; Herstrom, A. A.; Azevedo, S. H. (1991) An evaluation of various alternative
54 ambient ozone standards based on crop yield loss data. In: Berglund, R. L.; Lawson, D. R.; McKee, D. J., eds.

- Tropospheric ozone and the environment: papers from an international conference; March 1990; Los Angeles, CA. Pittsburgh, PA: Air & Waste Management Association; pp. 272-288. (A&WMA transactions series no. TR-19).
- Tingey, D. T.; McVeety, B. D.; Waschmann, R.; Johnson, M. G.; Phillips, D. L.; Rygiewicz, P. T.; Olszyk, D. M. (1996) A versatile sun-lit controlled-environment facility for studying plant and soil processes. *J. Environ. Qual.* 25: 614-625.
- Tingey, D. T.; Laurence, J. A.; Weber, J. A.; Greene, J.; Hogsett, W. E.; Brown, S.; Lee, E. H. (2001) Elevated CO₂ and temperature alter the response of *Pinus ponderosa* to ozone: A simulation analysis. *Ecol. Appl.* 11: 1412-1424.
- Tingey, D. T.; Rodecap, K. D.; Lee, E. H.; Hogsett, W. E.; Gregg, J. W. (2002) Pod development increases the ozone sensitivity of *Phaseolus vulgaris*. *Water Air Soil Pollut.* 139: 325-341.
- Tingey, D. T.; Hogsett, W. E.; Lee, E. H.; Laurence, J. A. (2004) Stricter ozone ambient air quality standard has beneficial effect on Ponderosa pine in California. *Environ. Manage.* 34: 397-405.
- Tissue, D. T.; Griffin, K. L.; Ball, T. (1999) Photosynthetic adjustment in field-grown ponderosa pine trees after six years of exposure to elevated CO₂. *Tree Physiol.* 19: 221-228.
- Tjoelker, M. G.; Luxmoore, R. J. (1991) Soil nitrogen and chronic ozone stress influence physiology, growth and nutrient status of *Pinus taeda* L. and *Liriodendron tulipifera* L. seedlings. *New Phytol.* 119: 69-81.
- Tjoelker, M. G.; Volin, J. C.; Oleksyn, J.; Reich, P. B. (1993) Light environment alters response to ozone stress in seedlings of *Acer saccharum* Marsh. and hybrid *Populus* L. I. *In situ* net photosynthesis, dark respiration and growth. *New Phytol.* 124: 627-636.
- Tjoelker, M. G.; Volin, J. C.; Oleksyn, J.; Reich, P. B. (1994) An open-air system for exposing forest-canopy branches to ozone pollution. *Plant Cell Environ.* 17: 211-218.
- Tjoelker, M. G.; Volin, J. C.; Oleksyn, J.; Reich, P. B. (1995) Interaction of ozone pollution and light effects on photosynthesis in a forest canopy experiment. *Plant Cell Environ.* 18: 895-905.
- Tognetti, R.; Cherubini, P.; Innes, J. L. (2000) Comparative stem-growth rates of Mediterranean trees under background and naturally enhanced ambient CO₂ levels. *New Phytol.* 146: 59-74.
- Tomlinson, H.; Rich, S. (1968) The ozone resistance of leaves as related to their sulfhydryl and adenosine triphosphate content. *Phytopathology* 58: 808-810.
- Tomlinson, H.; Rich, S. (1971) Effect of ozone on sterols and sterol derivatives in bean leaves. *Phytopathology* 61: 1404-1405.
- Tonneijck, A. E. G. (1994) Effects of various ozone exposures on the susceptibility of bean leaves (*Phaseolus vulgaris* L.) to *Botrytis cinerea*. *Environ. Pollut.* 85: 59-65.
- Tonneijck, A. E. G.; Leone, G. (1993) Changes in susceptibility of bean leaves (*Phaseolus vulgaris*) to *Sclerotinia sclerotiorum* and *Botrytis cinerea* by pre-inoculative ozone exposures. *Neth. J. Plant Pathol.* 99: 313-322.
- Tonneijck, A. E. G.; Van Dijk, C. J. (1994) Combined effects of ozone and ammonia on growth and yield of bean (*Phaseolus vulgaris* L. cv. Pros). In: Fuhrer, J.; Achermann, B., eds. Critical levels for ozone: a UN-ECE workshop report; November 1993; Bern, Switzerland. Liebefeld-Bern, Switzerland: Swiss Federal Research Station for Agricultural Chemistry and Environmental Hygiene; pp. 282-286. (Schriftenreihe der FAC Liebefeld no. 16).
- Tonneijck, A. E. G.; Van Dijk, C. J. (1997) Effects of ambient ozone on injury and yield of *Phaseolus vulgaris* at four rural sites in The Netherlands as assessed by using ethylenediurea (EDU). *New Phytol.* 135: 93-100.
- Tonneijck, A. E. G.; Van Dijk, C. J. (1998) Responses of bean (*Phaseolus vulgaris* L. cv. Pros) to chronic ozone exposure at two levels of atmospheric ammonia. *Environ. Pollut.* 99: 45-51.
- Tonneijck, A. E. G.; Van Dijk, C. J. (2002a) Assessing effects of ambient ozone on injury and yield of bean with ethylenediurea (EDU): three years of plant monitoring at four sites in The Netherlands. *Environ. Monit. Assess.* 77: 1-10.
- Tonneijck, A. E. G.; Van Dijk, C. J. (2002b) Injury and growth response of subterranean clover to ambient ozone as assessed by using ethylenediurea (EDU): three years of plant monitoring at four sites in The Netherlands. *Environ. Exper. Bot.* 48: 33-41.
- Topa, M. A.; Vanderklein, D. W.; Corbin, A. (2001) Effects of elevated ozone and low light on diurnal and seasonal carbon gain in sugar maple. *Plant Cell Environ.* 24: 663-677.
- Treshow, M.; Stewart, D. (1973) Ozone sensitivity of plants in natural communities. *Biol. Conserv.* 5: 209-214.
- Tuovinen, J.-P. (2002) Assessing vegetation exposure to ozone: is it possible to estimate AOT40 by passive sampling. *Environ. Pollut.* 119: 203-214.

- 1 U.S. Congress. (2002) Food security and rural investment act of 2002, PL 107-171, May 13, 2002. Washington, DC:
2 U.S. Government Printing Office.
- 3 U.S. Environmental Protection Agency. (1986) Air quality criteria for ozone and other photochemical oxidants. Research
4 Triangle Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment
5 Office; report nos. EPA-600/8-84-020aF-eF. 5v. Available from: NTIS, Springfield, VA; PB87-142949.
- 6 U.S. Environmental Protection Agency. (1992) Summary of selected new information on effects of ozone on health and
7 vegetation: supplement to 1986 air quality criteria for ozone and other photochemical oxidants. Research Triangle
8 Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; report
9 no. EPA/600/8-88/105F. Available from: NTIS, Springfield, VA; PB92-235670.
- 10 U.S. Environmental Protection Agency. (1993) Air quality criteria for oxides of nitrogen. Research Triangle Park, NC:
11 Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; report nos.
12 EPA/600/8-91/049aF-cF. 3v. Available from: NTIS, Springfield, VA; PB95-124533, PB95-124525, and
13 PB95-124517.
- 14 U.S. Environmental Protection Agency. (1996a) Air quality criteria for ozone and related photochemical oxidants.
15 Research Triangle Park, NC: Office of Research and Development; report nos. EPA/600/AP-93/004aF-cF. 3v.
16 Available from: NTIS, Springfield, VA; PB96-185582, PB96-185590, and PB96-185608. Available:
17 <http://cfpub2.epa.gov/ncea/>.
- 18 U.S. Environmental Protection Agency. (1996b) Review of national ambient air quality standards for ozone: assessment
19 of scientific and technical information. OAQPS staff paper. Research Triangle Park, NC: Office of Air Quality
20 Planning and Standards; report no. EPA/452/R-96/007. Available from: NTIS, Springfield, VA; PB96-203435.
21 Available: http://www.epa.gov/ttn/naaqs/standards/ozone/s_o3_pr_sp.html (9 September 2003).
- 22 U.S. Environmental Protection Agency. (2003) Technology Transfer Network: Air Quality System (AQS). Washington,
23 DC: Office of Air and Radiation. Available: <http://www.epa.gov/ttn/airs/airsaqs/> [1 April, 2004].
- 24 United Nations Economic Commission for Europe, UNECE. (1988) ECE critical levels workshop; March; Bad
25 Harzburg, Germany [final report]. Geneva, Switzerland: United Nations Economic Commission for Europe.
- 26 United Nations Environment Programme (UNEP). (1993) Environmental data report 1993-94. Oxford, United Kingdom:
27 Blackwell Publishers.
- 28 United Nations Environment Programme (UNEP). (1999) Geo-2000: global environmental outlook 2000. London,
29 United Kingdom: Earthscan Publishers. Available: <http://www.unep.org/geo2000/english/index.htm> (26 July
30 2002).
- 31 Unsworth, M. H.; Hogsett, W. E. (1996) Combined effects of changing CO₂, temperature, UV-B radiation and O₃ on
32 crop growth. In: Bazzaz, F.; Sombroek, W., eds. Global climate change and agricultural production. Chichester,
33 United Kingdom: John Wiley & Sons; pp. 171-197. Available:
34 <http://www.fao.org/docrep/W5183E/w5183e00.htm> [28 August, 2003].
- 35 Unsworth, M. H.; Heagle, A. S.; Heck, W. W. (1984a) Gas exchange in open-top field chambers - I. measurement and
36 analysis of atmospheric resistances to gas exchange. Atmos. Environ. 18: 373-380.
- 37 Unsworth, M. H.; Heagle, A. S.; Heck, W. W. (1984b) Gas exchange in open-top field chambers - II. resistances to
38 ozone uptake by soybeans. Atmos. Environ. 18: 381-385.
- 39 Utriainen, J.; Holopainen, T. (2001) Nitrogen availability modifies the ozone responses of Scots pine seedlings exposed
40 in an open-field system. Tree Physiol. 21: 1205-1213.
- 41 Utriainen, J.; Janhunen, S.; Helmisaari, H. S.; Holopainen, T. (2000) Biomass allocation, needle structural characteristics
42 and nutrient composition in Scots pine seedlings exposed to elevated CO₂ and O₃ concentrations. Trees (Berlin,
43 Ger.) 14: 475-484.
- 44 Vacek, S.; Bastl, M.; Leps, J. (1999) Vegetation changes in forests of the Krkonose Mountains over a period of air
45 pollution stress (1980-1995). Plant Ecol. 143: 1-11.
- 46 Van Camp, W.; Willekens, H.; Bowler, C.; Van Montagu, M.; Inzé, D.; Reupold-Popp, P.; Sandermann, H., Jr.;
47 Langebartels, C. (1994) Elevated levels of superoxide dismutase protect transgenic plants against ozone damage.
48 Biotechnology 12: 165-168.
- 49 Van De Staaij, J. W. M.; Lenssen, G. M.; Stroetenga, M.; Rozema, J. (1993) The combined effects of elevated CO₂ levels
50 and UV-B radiation on growth characteristics of *Elymus athericus* (= *E. pycnanthus*). Vegetatio 104/105: 433-439.
- 51 Van De Staaij, J. W. M.; Tonneijck, A. E. G.; Rozema, J. (1997) The effect of reciprocal treatments with ozone and
52 ultraviolet-B radiation on photosynthesis and growth of perennial grass *Elymus athericus*. Environ. Pollut. 97:
53 281-286.

- 1 Van Den Driessche, R.; Langebartels, C. (1994) Foliar symptoms, ethylene biosynthesis and water use of young Norway
2 spruce (*Picea abies* (L.) Karst.) exposed to drought and ozone. *Water Air Soil Pollut.* 78: 153-168.
- 3 Van Oijen, M.; Ewart, F. (1999) The effects of climatic variation in Europe on the yield response of spring wheat cv.
4 Minaret to elevated CO₂ and O₃: an analysis of open-top chamber experiments by means of two crop growth
5 simulation models. *Eur. J. Agron.* 10: 249-264.
- 6 Van Oijen, M.; Schapendonk, A. H. C. M.; Jansen, M. J. H.; Pot, C. S.; Maciorowski, R. (1999) Do open-top chambers
7 overestimate the effects of rising CO₂ on plants? An analysis using spring wheat. *Glob. Change Biol.* 5: 411-421.
- 8 Vanacker, H.; Carver, T. L. W.; Foyer, C. H. (2000) Early H₂O₂ accumulation in mesophyll cells leads to induction of
9 glutathione during the hyper-sensitive response in the barley-powdery mildew interaction. *Plant Physiol.* 123:
10 1289-1300.
- 11 VanderHeyden, D.; Skelly, J.; Innes, J.; Hug, C.; Zhang, J.; Landolt, W.; Bleuler, P. (2001) Ozone exposure thresholds
12 and foliar injury on forest plants in Switzerland. *Environ. Pollut.* 111: 321-331.
- 13 Vandermeiren, K.; De Temmerman, L.; Hookham, N. (1995) Ozone sensitivity of *Phaseolus vulgaris* in relation to
14 cultivar differences, growth stage and growing conditions. *Water Air Soil Pollut.* 85: 1455-1460.
- 15 Vandermeiren, K.; Black, C.; Lawson, T.; Casanova, M. A.; Ojanperä, K. (2002) Photosynthetic and stomatal responses
16 of potatoes grown under elevated CO₂ and/or O₃—results from the European CHIP-programme. *Eur. J. Agron.* 17:
17 337-352.
- 18 Vann, D. R.; Strimbeck, G. R.; Johnson, A. H. (1995) Effects of mist acidity and ambient ozone removal on montane red
19 spruce. *Tree Physiol.* 15: 639-647.
- 20 Volin, J. C.; Reich, P. B.; Givnish, T. J. (1998) Elevated carbon dioxide ameliorates the effects of ozone on
21 photosynthesis and growth: species respond similarly regardless of photosynthetic pathway or plant functional
22 group. *New Phytol.* 138: 315-325.
- 23 Volk, M.; Geissmann, M.; Blatter, A.; Contat, F.; Fuhrer, J. (2003) Design and performance of a free-air exposure system
24 to study long-term effects of ozone on grasslands. *Atmos. Environ.* 37: 1341-1350.
- 25 Vollenweider, P.; Woodcock, H.; Kelty, M. J.; Hofer, R.-M. (2003a) Reduction of stem growth and site dependency of
26 leaf injury in Massachusetts black cherries exhibiting ozone symptoms. *Environ. Pollut.* 125: 467-480.
- 27 Vollenweider, P.; Ottiger, M.; Günthardt-Goerg, M. S. (2003b) Validation of leaf ozone symptoms in natural vegetation
28 using microscopical methods. *Environ. Pollut.* 124: 101-118.
- 29 Von Gunten, U. (2003) Ozonation of drinking water: part I. Oxidation kinetics and product formation. *Water Res.* 37:
30 1443-1467.
- 31 Vorne, V.; Ojanperä, K.; De Temmerman, L.; Bindi, M.; Högy, P.; Jones, M. B.; Lawson, T.; Persson, K. (2002) Effects
32 of elevated carbon dioxide and ozone on potato tuber quality in the European multiple-site experiment
33 'CHIP-project'. *Eur. J. Agron.* 17: 369-381.
- 34 Vozzo, S. F.; Miller, J. E.; Pursley, W. A.; Heagle, A. S. (1995) Effects of ozone and water deficit on field-grown
35 soybean: I. Leaf gas exchange. *J. Environ. Qual.* 24: 663-670.
- 36 Waite, C. E.; DeHayes, D. H.; Rebbeck, J.; Schier, G. A.; Johnson, A. H. (1994) The influence of elevated ozone on
37 freezing tolerance of red spruce seedlings. *New Phytol.* 126: 327-335.
- 38 Walcek, C. J.; Yuan, H.-H.; Stockwell, W. R. (1997) The influence of aqueous-phase chemical reactions on ozone
39 formation in polluted and nonpolluted clouds. *Atmos. Environ.* 31: 1221-1237.
- 40 Wallin, G.; Skärby, L.; Selldén, G. (1992) Long-term exposure of Norway spruce, *Picea abies* (L.) Karst., to ozone in
41 open-top chambers. 3. Effects on the light response of net photosynthesis in shoots of different ages. *New Phytol.*
42 121: 387-394.
- 43 Wallin, G.; Karlsson, P. E.; Selldén, G.; Ottosson, S.; Medin, E.-L.; Pleijel, H.; Skärby, L. (2002) Impact of four years
44 exposure to different levels of ozone, phosphorus and drought on chlorophyll, mineral nutrients, and stem volume
45 of Norway spruce, *Picea abies*. *Physiol. Plant.* 114: 192-206.
- 46 Wang, D.; Hinckley, T. M.; Cumming, A. B.; Braatne, J. (1995) A comparison of measured and modeled ozone uptake
47 into plant leaves. *Environ. Pollut.* 89: 247-254.
- 48 Ward, E. R.; Uknes, S. J.; Williams, S. C.; Dincher, S. S.; Wiederhold, D. L.; Alexander, D. C.; Ahi-Goy, P.; Métraux,
49 J.-P.; Ryals, J. A. (1991) Coordinate gene activity in response to agents that induce systemic acquired resistance.
50 *Plant Cell* 3: 1084-1094.
- 51 Warwick, K. R.; Taylor, G. (1995) Contrasting effects of tropospheric ozone on five native herbs which coexist in
52 calcareous grassland. *Global Change Biol.* 1: 143-151.
- 53 Weber, J.; Grulke, N. E. (1995) Effects of air pollution on stem growth. In: Gartner, B., ed. *Stems and trunks in plant*
54 *form and function*. San Diego, CA: Academic Press; pp. 343-364.

- 1 Weber, J. A.; Clark, C. S.; Hogsett, W. E. (1993) Analysis of the relationships among O₃ uptake, conductance, and
2 photosynthesis in needles of *Pinus ponderosa*. *Tree Physiol.* 13: 157-172.
- 3 Weiner, J.; Fredro-Boniecki, S.; Reed, D.; Maclean, A.; Strong, M. (1997) NiepoLomice forest-A GIS analysis of
4 ecosystem response to industrial pollution. *Environ. Pollut.* 3: 381-388.
- 5 Weinstein, D. A.; Yanai, R. D. (1994) Integrating the effects of simultaneous multiple stresses on plants using the
6 simulation model TREGRO. *J. Environ. Qual.* 23: 418-428.
- 7 Weinstein, D. A.; Samuelson, L. J.; Arthur, M. A. (1998) Comparison of the response of red oak (*Quercus rubra*)
8 seedlings and mature trees to ozone exposure using simulation modeling. *Environ. Pollut.* 102: 307-320.
- 9 Weinstein, D. A.; Gollands, B.; Retzlaff, W. A. (2001) The effects of ozone on a lower slope forest of the Great Smoky
10 Mountain National Park: simulations linking an individual tree model to a stand model. *For. Sci.* 47: 29-42.
- 11 Weinstein, D. A.; Woodbury, P. B.; Gollands, B.; King, P.; Lepak, L.; Pendleton, D. (2002) Assessment of effects of
12 ozone on forest resources in the southern Appalachian mountains. Asheville, NC: Southern Appalachian
13 Mountains Initiative. Available: <http://cycas.cornell.edu/ebp/pmg.html> [19 March, 2003].
- 14 Welfare, K.; Flowers, T. J.; Taylor, G.; Yeo, A. R. (1996) Additive and antagonistic effects of ozone and salinity on the
15 growth, ion contents and gas exchange of five varieties of rice (*Oryza sativa* L.). *Environ. Pollut.* 92: 257-266.
- 16 Wellburn, A. R. (1990) Why are atmospheric oxides of nitrogen usually phytotoxic and not alternative fertilizers? *New*
17 *Phytol.* 115: 395-429.
- 18 Wellburn, A. (1994) Air pollution and climate change: the biological impact. Harlow, Essex, United Kingdom: Longman
19 Scientific & Technical.
- 20 Wellburn, F. A. M.; Wellburn, A. R. (1994) Atmospheric ozone affects carbohydrate allocation and winter hardiness of
21 *Pinus halepensis* (Mill.) *J. Exp. Bot.* 274: 607-614.
- 22 Werner, H.; Fabian, P. (2002) Free-air fumigation of mature trees. A novel system for controlled ozone enrichment in
23 grown-up beech and spruce canopies. *Environ. Sci. Pollut. Res.* 9: 117-121.
- 24 Wesely, M. L. (1989) Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical
25 models. *Atmos. Environ.* 23: 1293-1304.
- 26 Westenbarger, A. D.; Frisvold, G. B. (1994) Agricultural exposure to ozone and acid precipitation. *Atmos. Environ.* 28:
27 2895-2907.
- 28 Westman, W. E. (1979) Oxidant effects on Californian coastal sage scrub. *Science* (Washington, DC) 205: 1001-1003.
- 29 Westman, W. E. (1981) Factors influencing the distribution of species of Californian coastal sage scrub. *Ecology* 62:
30 439-455.
- 31 Weubbles, D. J.; Jain, A.; Edmonds, J.; Harvey, D.; Hayhoe, K. (1999) Global change: state of the science. *Environ.*
32 *Pollut.* 100: 57-86.
- 33 Wheeler, T. R.; Hong, T. D.; Ellis, R. H.; Batts, G. R.; Morrison, J. I. L.; Hadley, P. (1996) The duration and rate of grain
34 growth, and harvest index, of wheat (*Triticum aestivum* L.) in response to temperature and CO₂. *J. Exper. Bot.* 47:
35 623-630.
- 36 Whitfield, C. P.; Davison, A. W.; Ashenden, T. W. (1996a) Interactive effects of ozone and soil volume on *Plantago*
37 *major*. *New Phytol.* 134: 287-294.
- 38 Whitfield, R. G.; Biller, W. F.; Jusko, M. J.; Keisler, J. M. (1996b) A probabilistic assessment of health risks associated
39 with short-term exposure to tropospheric ozone. Argonne, IL: Argonne National Laboratory, Decision and
40 Information Sciences Division; report no. ANL/DIS-3.
- 41 Whitfield, C. P.; Davison, A. W.; Ashenden, T. W. (1997) Artificial selection and heritability of ozone resistance in two
42 populations of *Plantago major*. *New Phytol.* 137: 645-655.
- 43 Whitfield, C. P.; Davison, A. W.; Ashenden, T. W. (1998) The effects of nutrient limitation on the response of *Plantago*
44 *major* to ozone. *New Phytol.* 140: 219-230.
- 45 Whittaker, J. B. (1994) Interactions between insects and air pollutants. In: Alscher, R. G.; Wellburn, A. R., eds. *Plant*
46 *responses to the gaseous environment. Molecular, metabolic and physiological aspects.* London, United Kingdom:
47 Chapman and Hall; pp. 365-384.
- 48 Wickman, B. E. (1992) Forest Health in the Blue Mountains: the influence of insects and diseases. Portland OR: U.S.
49 Department of Agriculture, Forest Service, Pacific Northwest Research Station; general technical report
50 PNW-GTR-295. (Quigley, T. M., ed. *Forest health in the Blue Mountains: science perspectives*).
- 51 Wieser, G. (1997) Ozone impact on photosynthetic capacity of mature and young Norway spruce (*Picea abies* (L.)
52 Karst.). External versus internal exposure. *Phyton* (Austria) 37: 297-302.
- 53 Wieser, G.; Havranek, W. M. (1993) Ozone uptake in the sun and shade crown of spruce: quantifying the physiological
54 effects of ozone exposure. *Trees* (Berlin, Ger.) 7: 227-232.

- 1 Wieser, G.; Havranek, W. M. (1995) Environmental control of ozone uptake in *Larix decidua* Mill.: a comparison
2 between different altitudes. *Tree Physiol.* 15: 253-258.
- 3 Wieser, G.; Häsler, R.; Götz, B.; Koch, W.; Havranek, W. M. (1999) Seasonal ozone uptake of mature evergreen
4 conifers at different altitudes. *Phyton (Austria)* 39: 233-240.
- 5 Wieser, G.; Tegischer, K.; Tausz, M.; Häberle, K.-H.; Grams, T. E. E.; Matyssek, R. (2002) Age effects on Norway
6 spruce (*Picea abies*) susceptibility to ozone uptake: a novel approach relating stress avoidance to defense. *Tree*
7 *Physiol.* 22: 583-590.
- 8 Wilbourn, S.; Davison, A. W.; Ollerenshaw, J. H. (1995) The use of an unenclosed field fumigation system to determine
9 the effects of elevated ozone on a grass-clover mixture. *New Phytol.* 129: 23-32.
- 10 Willekens, H.; Van Camp, M.; Van Montagu, M.; Inze, D.; Langebartels, C.; Sandermann, H., Jr. (1994) Ozone, sulfur
11 dioxide and ultraviolet B have similar effects on mRNA accumulation of antioxidant genes in *Nicotiana*
12 *plumbaginifolia* L. *Plant Physiol.* 106:1007-1014.
- 13 Williams, T. P. W. (1994) Use of passive monitors to assess plant bioindicators for ground-level ozone. New Brunswick,
14 Canada: University of New Brunswick.
- 15 Williams, J.; Bulman, M. P.; Neill, S. J. (1994) Wilt-induced aba biosynthesis, gene expression and down regulation of
16 *rbcs* mRNA levels in *Arabidopsis thaliana*. *Physiol. Plant.* 91: 177-182.
- 17 Wiltshire, J. J. J.; Wright, C. J.; Unsworth, M. H. (1992) A new method for exposing mature trees to ozone episodes. *For.*
18 *Ecol. Manage.* 51: 115-120.
- 19 Wiltshire, J. J. J.; Unsworth, M. H.; Wright, C. J. (1994) Seasonal changes in water use of ash trees exposed to ozone
20 episodes. *New Phytol.* 127: 349-354.
- 21 Wiltshire, J. J. J.; Wright, C. J.; Colls, J. J.; Craigon, J.; Unsworth, M. H. (1996) Some foliar characteristics of ash trees
22 (*Fraxinus excelsior*) exposed to ozone episodes. *New Phytol.* 134: 623-630.
- 23 Winner, W. E.; Lefohn, A. S.; Cotter, I. S.; Greitner, C. S.; Nellessen, J.; McEvoy, L. R., Jr.; Olson, R. L.; Atkinson, C.
24 J.; Moore, L. D. (1989) Plant responses to elevational gradients of O₃ exposures in Virginia. *Proc. Natl. Acad. Sci.*
25 *U. S. A.* 86: 8828-8832.
- 26 Winner, W. E.; Coleman, J. S.; Gillespie, C.; Mooney, H. A.; Pell, E. J. (1991) Consequences of evolving resistance to air
27 pollutants. In: Taylor, G. E.; Pitelka, L. F.; Clegg, M. T., eds. *Ecological genetics and air pollution*. New York,
28 NY: Springer-Verlag; pp. 177-202.
- 29 Wolf, J.; Van Oijen, M. (2003) Model simulation of effects of changes in climate and atmospheric CO₂ and O₃ on tuber
30 yield potential of potato (cv. Bintje) in the European Union. *Agric. Ecosyst. Environ.* 94: 141-157.
- 31 Wolff, K.; Morgan-Richards, M.; Davison, A. W. (2000) Patterns of molecular genetic variation in *Plantago major* and
32 *P. intermedia* in relation to ozone resistance. *New Phytol.* 145: 501-509.
- 33 Wolters, J. H. B.; Martens, M. J. M. (1987) Effects of air pollutants on pollen. *Bot. Rev.* 53: 372-414.
- 34 Woodbury, P. B.; Laurence, J. A.; Hudler, G. W. (1994) Chronic ozone exposure increases the susceptibility of hybrid
35 *Populus* to disease caused by *Septoria musiva*. *Environ. Pollut.* 86: 109-114.
- 36 Wu, Y.-X.; Tiedemann, A. V. (2002) Impact of fungicides on active oxygen species and antioxidant enzymes in spring
37 barley (*Hordeum vulgare* L.) exposed to ozone. *Environ. Pollut.* 116: 37-47.
- 38 Wulff, A.; Hänninen, O.; Tuomainen, A.; Kärenlampi, L. (1992) A method for open-air exposure of plants to ozone.
39 *Ann. Bot. Fennici.* 29: 253-262.
- 40 Wullschlegel, S. D.; Hanson, P. J.; Edwards, G. S. (1996) Growth and maintenance respiration in leaves of northern red
41 oak seedlings and mature trees after 3 years of ozone exposure. *Plant Cell Environ.* 19: 577-584.
- 42 Wustman, B. A.; Oksanen, E.; Karnosky, D. F.; Noormets, A.; Isebrands, J. G.; Pregitzer, K. S.; Hendrey, G. R.; Sober,
43 J.; Podila, G. K. (2001) Effects of elevated CO₂ and O₃ on aspen clones varying in O₃ sensitivity: can CO₂
44 ameliorate the harmful effects of O₃? *Environ. Pollut.* 115: 473-481.
- 45 Yalpani, N.; Enyedi, A. J.; Leön, J.; Raskin, I. (1994) Ultraviolet light and ozone stimulate accumulation of salicylic
46 acid, pathogenesis-related proteins and virus resistance in tobacco. *Planta* 193: 372-376.
- 47 Yonekura, T.; Dokiya, Y.; Fukami, M.; Izuta, T. (2001) Effects of ozone and/or soil water stress on growth and
48 photosynthesis of *Fagus crenata* seedlings. *Water Air Soil Pollut.* 130: 965-970.
- 49 Young, T. F.; Sanzone, S., eds. (2002) A framework for assessing and reporting on ecological condition: an SAB report.
50 Washington, DC: U.S. Environmental Protection Agency, Science Advisory Board; report no.
51 EPA-SAB-EPEC-02-009. Available: <http://www.epa.gov/sab/pdf/epec02009.pdf> [9 December, 2003].
- 52 Younglove, T.; McCool, P. M.; Musselman, R. C.; Kahl, M. E. (1994) Growth-stage dependent crop yield response to
53 ozone exposure. *Environ. Pollut.* 86: 287-295.

- 1 Yun, S.-C.; Laurence, J. A. (1999a) The response of clones of *Populus tremuloides* differing in sensitivity to ozone in the
2 field. New Phytol. 141: 411-421.
- 3 Yun, S.-C.; Laurence, J. A. (1999b) The response of sensitive and tolerant clones of *Populus tremuloides* to dynamic
4 ozone exposure under controlled environmental conditions. New Phytol. 143: 305-313.
- 5 Yunus, M.; Iqbal, M. (1996) Plant response to air pollution. Chichester, United Kingdom: John Wiley & Sons Ltd.
- 6 Yuska, D. E.; Skelly, J. M.; Ferdinand, J. A.; Stevenson, R. E.; Savage, J. E.; Mulki, J.; Hines, A. (2003) Use of
7 bioindicators and passive sampling devices to evaluate ambient ozone concentrations in north central
8 Pennsylvania. Environ. Pollut. 125: 71-80.
- 9 Zeller, K.; Nikolov, N. (2000) Quantifying simultaneous fluxes of ozone, carbon dioxide and water vapor above a
10 subalpine forest ecosystem. Environ. Pollut. 107: 1-20.
- 11 Zhang, X.; Zhang, L.; Dong, F.; Gao, J.; Galbraith, D. W.; Song, C.-P. (2001) Hydrogen peroxide is involved in abscisic
12 acid-induced stomatal closure in *Vicia faba*. Plant Physiol. 126: 1438-1448.
- 13 Zheng, Y.; Lyons, T.; Ollerenshaw, J. H.; Barnes, J. D. (2000) Ascorbate in the leaf apoplast is a factor mediating ozone
14 resistance in *Plantago major*. Plant Physiol. Biochem. (Paris, Fr.) 38: 403-411.
- 15 Zinser, C.; Jungblut, T.; Heller, W.; Seidlitz, H. K.; Schnitzler, J.-P.; Ernst, D.; Sandermann, H., Jr. (2000) The effect of
16 ozone in Scots pine (*Pinus sylvestris* L.): gene expression, biochemical changes and interactions with UV-B
17 radiation. Plant Cell Environ. 23: 975-982.
- 18 Ziska, L.; Bunce, J. A. (1997) The role of temperature in determining the stimulation of CO₂ assimilation at elevated
19 carbon dioxide concentrations in soybean seedlings. Physiol. Plant. 100: 126-132.
- 20

10. TROPOSPHERIC OZONE EFFECTS ON UV-B FLUX AND CLIMATE CHANGE PROCESSES

10.1 INTRODUCTION

In addition to exerting direct effects on human health and vegetation/ecosystems, tropospheric ozone (O₃) is involved in determining ground-level flux of solar ultraviolet (UV) radiation and, also, in other processes that alter the earth's radiative balance and contribute to resulting climate change. This chapter first discusses the role of tropospheric ozone in determining surface-level UV flux and, then, secondly, discusses tropospheric O₃ involvement in global climate change.

10.2 THE ROLE OF TROPOSPHERIC OZONE IN DETERMINING GROUND-LEVEL UV-B FLUX

Stratospheric O₃ plays a crucial role in reducing the exposure of living organisms to solar ultraviolet radiation. The stratosphere, the atmospheric layer immediately above the troposphere, is roughly 50 km thick and possesses 90% or more of the atmosphere's O₃. Ozone depletion due to the release of long-lived anthropogenic chlorinated and fluorinated hydrocarbons was discovered over the course of the 1970s and early 1980s and led to establishment of an international treaty for the protection of stratospheric O₃, the 1987 Montreal Protocol on Substances that Deplete the Ozone Layer.

Active research has been underway into the effects of increased UV radiation on ecosystems and on human health since the discovery of the seasonal polar "Ozone Hole" and the declining stratospheric O₃ concentrations in the midlatitudes. An outcome of this effort is a body of literature that also describes the effects of tropospheric pollutants, PM, and O₃, on ground-level UV radiation flux. The Montreal Protocol requires routine review of the latest scientific information available on the status of the O₃ layer and of UV radiation levels at the earth's surface. The World Meteorology Organization (WMO) and UN Environmental Program (UNEP) are responsible for assessing the state of the science regarding the O₃ layer and for

reporting on trends in surface UV radiation levels. The latest WMO/UNEP assessment was published in 2003 (WMO/UNEP, 2002).

This section describes the current level of scientific understanding of the factors influencing the flux of UV radiation at the earth's surface, including geophysical factors, tropospheric O₃, PM, and cloud cover, with reference to the WMO/UNEP assessment and the peer-reviewed literature. Factors influencing the degree of human exposure to UV-B and the resulting health effects are also discussed.

10.2.1 Factors Governing Ultraviolet Radiation Flux at the Earth's Surface

10.2.1.1 UV Radiation: Wavelengths, Energies and Depth of Atmospheric Penetration

Designations for portions of the electromagnetic spectrum have evolved over time and are usually associated with general function or effects caused by photons in a given wavelength range. Gamma rays, at or below 0.1 nm in wavelength, are especially damaging high energy photons emitted during radioactive decay and by stellar activity. Radiowaves, wavelengths greater than 10⁸ nm, are very low in energy and function as carriers for broadcast communications. The energy possessed by any photon is inversely proportional to its wavelength.

The wavelengths ranging between 50 and 400 nm in length are denoted "ultraviolet." Solar radiation of wavelengths < 280 nm, including UV-C (200 to 280 nm), is almost entirely blocked by Earth's upper atmosphere due to photoionization and photo-dissociation processes. Figure 10-1 shows a comparison between solar flux above the atmosphere and the flux at the earth's surface. The flux of solar radiation between 280 and 320 nm (UV-B) is absorbed or scattered in part within the atmosphere, while radiation between 320 and 400 nm (UV-A) can be scattered but are not absorbed by gases to any meaningful degree. UV-B and UV-A photons contain the necessary levels of energy needed to break (photolyze) chemical bonds. Both UV-A and UV-B are associated with human health- and ecosystem-damaging effects. Because UV-B is more energetic, it is potentially capable of producing more biological damage than UV-A. UV-A comprises 90% of the UV radiation that reaches the earth's surface, yet is 1000 times less damaging to keratinocytes (skin cells), than UV-B (Hildesheim, 2004).




Figure 10-1. Extraterrestrial solar flux measured by the satellite UARS SOLSTICE instrument (dotted line). The dashed line represents calculated atmospheric transmittance and the solid line is the calculated absolute flux of UV radiation for a solar zenith angle of 50deg, total column O3 of 275 DU, and a surface reflectivity of 8%. The fine structure on the surface flux trace results from Fraunhofer lines (absorption specific wavelengths within the solar atmosphere).

Source: Krotkov et al. (1998).

10.2.1.2 Temporal Variations in Solar Flux

The magnitude of the solar radiation flux entering the atmosphere is dependent upon long-term solar activity, sunspot cycle (11 years), solar rotation (27 days) and the position of the earth in its orbit around the sun. A variety of changes in solar irradiance can be found in the historical data, beginning in 1700 and leading up to the present. Fligge and Solanki (2000) concluded that solar irradiation changes on time scales of days to centuries can be attributed to variations in solar magnetic features. Since the last Maunder minimum in 1700, solar irradiance has increased very slightly, at approximately 3.0% for wavelengths < 300 nm and at ~1.3% for the UV-B and UV-A range. Including visible wavelengths, Fligge and Solanki (2000) estimated that the overall increase in solar irradiance was ~0.3%. Rozema et al. (2001) pointed out that any

1 increase in wavelengths < 300 nm (UV-C) would initiate additional O₃ formation in the
2 stratosphere, therefore increasing its UV radiation absorptive capacity.

3 Solar rotation and sunspot activity have the greatest effect on the radiation flux that
4 originates in the highest levels of the solar atmosphere. The amplitude of the associated cyclical
5 changes in solar shortwave radiation flux follows an inverse relationship between the photon's
6 wavelength and the solar altitude at which it was emitted. The maximum level of radiation
7 (solar-max) differs from the minimum (solar-min) by as much as 10% for wavelengths near
8 160 nm. This peak-to-trough difference declines to around 1% for 300 nm (UV-B range)
9 (Salby, 1996).

10 The combined effects of the earth's obliquity (the angle of the earth's axis of rotation with
11 respect to the plane of its orbit around the sun) and its precession (the rotation of the earth's axis
12 with respect to a perpendicular line through the plane of its solar orbit) yield variations of up to
13 30% in total summertime solar flux, depending on latitude (Hartmann, 1994).

14 15 *Zenith Angle: Latitude, Season and Time of Day*

16 The sun's relative elevation is measured with respect to the vertical and is known as its
17 "zenith angle." This angle varies hourly, seasonally, and with latitude. Daily and seasonal
18 changes in solar zenith angle result in the largest changes in the magnitude of solar radiation
19 flux, with higher zenith angles corresponding to lower solar flux. The largest natural fluxes
20 occur in the tropical regions, where solar noon occurs at a zenith angle at or near 0°. Seasonal
21 variation in solar flux ranges from small changes at the equator to very large changes at high
22 latitudes. Daily variations in solar flux, from sunrise to sunset, show added wavelength
23 dependence as a function of zenith angle, because transmission of some wavelengths are
24 sensitive to atmospheric pathlength due to scattering and absorption processes. These processes
25 will be discussed further below.

26 27 **10.2.1.3 Atmospheric Radiative Interactions with Solar UV Radiation**

28 *Radiative Interactions in the Stratosphere*

29 The stratosphere contains 90% or more of the total column density of O₃, the principle gas
30 phase absorber of UV-B. Ozone interacts with UV radiation by scattering the photon or
31 absorbing and transforming its energy. Upon absorbing a UV photon, O₃ may photodissociate,
32 or become electronically- and vibrationally-excited.

Photoabsorption by O₃ occurs with very high efficiency. After electronically-excited O₃ (O₃^{*}) is formed, it will either dissociate into ground-state oxygen, O₂, and an electronically excited oxygen radical, O(¹D), or a collision between the O₃ molecule with another gas molecule (M) may occur (See Reactions 1 and 2). Intermolecular collisions degrade the excess electronic energy of the O₃ molecule by transferring it to other molecules as vibrational, rotational, and/or translational energies, which warm the atmosphere. The electronically excited oxygen radical can then react with H₂O to form two hydroxyl radicals, OH (Reaction 3). See Chapter 2 for further discussion of odd oxygen and HO_x photochemistry.



Either of these photochemical processes transforms the energy of the UV photon into a form lacking the potential for human health or ecosystems damage.

The WMO/UNEP (2002) scientific assessment reported that global average total column O₃ had declined by 3% from pre-1980 levels, due to the presence of anthropogenic ozone-depleting substances in the atmosphere. Ozone depletion has a strong latitude and seasonal dependence. The seasonality of total O₃ changes differ between the Northern and Southern Hemispheres. In the northern midlatitudes, total column O₃ declined by ~4% during the winter/spring seasons and by approximately half that amount in the summer/fall of the 1997-2001 time period, relative to pre-1980 total column O₃ levels. In southern midlatitudes, total column O₃ declined ~6% during all seasons.

The concentration of O₃ in a vertical column extending from the earth's surface is expressed in Dobson Units (DU), corresponding to the column height in hundredths of a millimeter of O₃ at standard temperature and pressure (273 K and 1 atmosphere) (Wayne, 2000). One DU = 2.587 × 10¹⁵ molecules of O₃/cm². The total O₃ column effectively prevents any UV-C from reaching the surface, reduces the penetration of UV-B to the surface but does little to

attenuate the intensity of UV-A except at the shorter wavelengths close to the cut off for UV-B. Cutchis (1974) calculated that with overhead sun a 10% decrease in the O₃ column would lead to 20, 250, and 500% increases in flux at 305, 290, and 287 nm, respectively, values that have been supported by ground observations in Toronto, ON (49° N; Kerr and McElroy, 1993). Rapid changes of this magnitude appear to happen naturally. As seen in data collected by the Total Ozone Mapping Satellite (TOMS) (Figure 10-2), the total O₃ column undergoes wide natural variation on short timescales (Cockell, 2001).




Figure 10-2. Ozone column abundances from the years 1990 to 1992 for 0, 40, and 80° N as well as 80° S. The data for 80° S are incomplete, but the graph shows the effects of the Antarctic O₃ hole on total column abundances at this latitude. The data for the Northern Hemisphere illustrate the natural variations in the O₃ column over time. The data are taken from the TOMS (Total Ozone Monitoring Satellite) data set (1979 to 1993).

Source: Cockell (2001).

1 Nacreous and polar stratospheric clouds scatter radiation and aerosols, such as those
2 injected into the stratosphere by explosive volcanic eruptions, both absorb and scatter radiation.
3 Relative to the troposphere, the stratosphere is low in atmospheric pressure. Stratospheric clouds
4 and aerosols are also more dispersed than in the troposphere. Consequently, UV radiation can
5 traverse the stratosphere with a substantially lower probability of encountering a gas molecule or
6 cloud or aerosol particle than it would in the troposphere. In the radiative transfer literature, the
7 stratosphere is described as a “single scattering” regime for UV radiation, and UV that has
8 penetrated the stratosphere is referred to as “direct beam UV.” The troposphere, due to its high
9 gas and particle concentrations is referred to as a “multiple scattering” regime.

11 ***Radiative Interactions in the Troposphere: Solar Irradiance vs. Actinic Flux***

12 The troposphere contains $\leq 10\%$ of the total column O_3 but $\sim 78\%$ of the total atmospheric
13 mass including clouds, gas- and particle-phase radiation scatterers and absorbers, making it a
14 “multiple scattering” regime for UV radiation. These scattering processes increase the mean-
15 free path a photon must travel before reaching the surface, transforming the direct beam UV
16 solar irradiance that has penetrated the stratosphere into diffuse, or actinic, UV (Brühl and
17 Crutzen, 1989).

18 Atmospheric scattering processes, such as Rayleigh and Mie scattering, are particle-size
19 dependent. In Rayleigh scattering, gas molecules that are smaller than the wavelength of the
20 incident photon isotropically deflect incoming photons. With Mie scattering, aerosol and cloud
21 droplets scatter incoming radiation with forward- and back-scattering tendencies. Actinic flux,
22 especially at the earth’s surface, is directly proportional to surface albedo (Wendisch and Mayer,
23 2003). Surface albedo is very strongly wavelength dependent. For example, fresh and wet snow
24 reflects 60 to 90% of violet light, while soil and grass surfaces reflect $< 5\%$ of incident violet
25 light (Xenopoulos and Schindler, 2001). Wendisch and Mayer (2003) found, in their in situ
26 measurement and modeling study of the vertical distribution of solar irradiance, that surface
27 albedos must be measured in order to accurately simulate solar flux, due to possibly large
28 variations in albedo within a given surface type. Snow cover, even many kilometers from
29 measurement sites is known to increase detected UV irradiances. Complicated interactions
30 result when radiation is scattered by snow (or other bright surfaces) and backscattered or
31 absorbed by atmospheric particles and clouds in the same vicinity (WMO/UNEP, 2002).

Variation in Solar Flux with Altitude

Solar flux increases with altitude above sea level, due to the decreased presence of clouds and declining concentrations of scattering and absorbing atmospheric pollutants. Rayleigh scattering, discussed below, also lessens with decreasing atmospheric pressure. A number of measurements of UV radiation have been taken as a function of altitude and are reviewed by Xenopoulos and Schindler (2001). Increases in flux as a function of altitude are given as percent irradiance enhancement per 1000 m relative to sea level. The effect can range from 9 to 24% /1000 m as function of the altitude at which the measurement was taken (Xenopoulos and Schindler, 2001). The effect corresponds to the relative pathlength traveled by the solar photon: flux is strongest when the photon is not impeded by atmospheric scattering or absorbing agents. Similarly, this effect is seen as a function of solar zenith angle, i.e., flux is at its maximum when the atmospheric depth through which the photon must pass is at its shallowest.

Clouds

In principle, clouds have the largest influence on surface level UV irradiance, but their effects are difficult to quantify. The depth and composition of a cloud determines, in part, the amount and wavelengths of radiation that it will scatter or absorb. Geometry is an especially important factor, as scattered or broken clouds may enhance, rather than reduce, surface flux. For example, if the sun is not fully blocked by a cloud, the reduction in irradiance may be small (WMO/UNEP, 2002). Quantifying the effect of clouds on surface UV flux, therefore, requires information on cloud composition, geometry, altitude, and the position of the sun relative to the cloud and the underlying surface, as a function of time. Provided that all of this detailed information is available, a three dimensional model is then required to calculate surface-level reductions or enhancements in UV flux.

Particulate Matter

On a zonally averaged basis, PM does not contribute significantly to lower tropospheric absorption of UV radiation. However, in urban areas or other areas subject to high smog levels (areas of significant biomass combustion), PM may be the most important determinant of ground-level erythemal UV flux, second only to cloud cover (U.S. Environmental Protection Agency, 2004; WMO/UNEP, 2003). Model-to-measurement comparisons of ground-level flux

for Greece and Toronto have shown 20 and 5-10% reductions, respectively (McKenzie et al., 2001). Increases over the past 20 to 30 years in combustion-associated PM and black carbon may account for the inability to detect a surface trend in UV-B radiation caused by a known decrease in stratospheric O₃ over the Northern Hemisphere (Barnard et al., 2003).

Gases

In the upper troposphere, the UV-absorbing gases O₃ and, of lesser importance, formaldehyde (CH₂O) and SO₂ are vented or diffuse from the surface. Stratospheric intrusions extrude O₃-rich air into the troposphere where it mixes, increasing regional background O₃ levels (see Chapter 3). Tropospheric O₃ data are typically expressed on a concentration basis, e.g., parts per billion by volume (ppbv), where 1 ppbv tropospheric O₃ ≡ 0.65 DU (IPCC, 2001). The mean values of O₃ in the free troposphere reported in the literature range from ~50 to ~80 ppbv, with higher values occurring at the tropopause. For example, a series of ozonesonde soundings over France during the period from 1976 to 1995 showed an O₃ increase from 48.9 ppbv in the 2.5 to 3.5 km layer to 56.5 ppbv in the 6.5 to 7.5 km layer, although the data revealed no significant increasing trend over time (Ancellet and Beekmann, 1997).

Close to the surface in polluted urban settings, photochemistry produces a diurnal rise and fall in O₃ and PM concentrations. Temperature inversions tend to prevent the upward mixing and dilution of ground-level O₃, while trapping primary and secondary PM. A recent study of the concurrence of O₃ and PM is provided by Koloutsou-Vakakis et al. (2001). No measurement technique is currently available that can distinguish between absorption of incident UV radiation by O₃ versus absorption by PM.

UV absorption by gases becomes important under aerosol- and cloud-free conditions. Figure 10-3 shows a calculation by Krotkov et al. (1998) of the sensitivity, as a function of wavelength, of ground-level UV flux to a 1-DU decrease in total column O₃ under cloud- and aerosol-free conditions. A 1991 to 1992 study in Chicago in which ambient O₃, broadband UV irradiance and total sunlight were monitored (Frederick et al., 1993), produced a statistically significant negative correlation between the UV irradiance and ambient O₃ when the atmosphere was relatively free of clouds and haze. Although they estimated that a 10-ppbv reduction in O₃ was associated with a 1.3% increase in erythemally-weighted UV-B, they cautioned that this figure had a comparatively large uncertainty (±1.2%).




Figure 10-3. The sensitivity of ground-level UV flux to a 1 DU change in total column O_3 , under clear sky conditions, as a function of solar zenith angle (SZA).

Source: Krotkov et al. (1998).

10.2.1.4 Modeling Surface UV-B Irradiance

The WMO/UNEP (2003) stated that, in principle, if the spatial distribution of all UV absorbers and scatterers were fully known, the wavelength and angular distribution of the UV irradiance at the earth's surface could be determined with model calculations. However, the poorly known and complicated distribution of the primary components (clouds, particles, O_3 , and surface albedo) makes detailed predictions extremely difficult.

Each of the factors described above must be integrated with human behaviors that also influence potential exposure to UV-B radiation in order to calculate the human health risks associated with small changes in total column O_3 density. These human factors and the related health consequences are described in detail below.

10.2.2 Factors Governing Human Exposure to Ultraviolet Radiation

Multiple, complex factors determine the solar flux of UV radiation at the ground level, as discussed in Section 10.2.1. Figure 10-4 illustrates some of these geophysical/atmospheric factors including stratospheric and tropospheric O₃, clouds, aerosols, and Rayleigh scattering. An estimate of the impact of changes in surface-level O₃ on UV-related human health effects requires information about human physical, behavioral, and demographic factors associated with exposure to UV radiation. In this section, the various factors that govern the degree of human exposure to UV radiation are examined.




Figure 10-4. Complexity of factors that determine human exposure to UV radiation. In addition to the geophysical/atmospheric factors (e.g., stratospheric and tropospheric O₃, clouds, aerosols, and Rayleigh scattering) that affect the solar flux of UV radiation at surface level, there are human physical, behavioral and demographic factors that influence human exposure to UV radiation.

10.2.2.1 Outdoor Activities

Exposure to solar UV radiation is related to one predominating factor – time spent in outdoor microenvironments during daylight hours. A large U.S. study was conducted using the EPA National Human Activity Pattern Survey (NHAPS) to assess UV radiation dose in Americans (Godar, 2001; Godar et al., 2001, 2003). The EPA NHAPS recorded the activity profiles of 9,386 Americans (age 0 to 60+ years) over a 24-month period to assess their exposure to various environmental pollutants, plus UV radiation. This study indicated that there was a strong seasonal preference for outdoor activities, with people spending the most time outdoors during the summer, followed by spring, fall, and, lastly, winter (Godar et al., 2001). Because the solar erythema (i.e., skin reddening) UV radiation dose is also highest during the summer, the estimated UV radiation dose of Americans was more than ten-fold greater in the summer compared to the winter season (Godar et al., 2001). Note that the error associated with estimating UV radiation dose from exposure surveys and one EPA UV-monitoring site located at each quadrant of the U.S. will be high.

Vacationing at the beach during the summer was associated with higher UV radiation exposures (Godar et al., 2001; Thieden et al., 2001). Even after accounting for sunscreen use at the beach, the erythema UV radiation doses were more than 40% higher during a three-week beach vacation compared to a three-week stay at home (Godar et al., 2001). Danish children and adolescents were found to receive > 50% of their annual UV radiation dose while vacationing at European beaches (Thieden et al., 2004a). The high UV radiation dose received at beaches is due to increased time spent outdoors during daylight hours and increased risk behavior, namely sunbathing. Sunbathing also was associated with increased annual UV radiation dose in the Canadian National Survey on Sun Exposure and Protective Behaviours (Shoveller et al., 1998). Among the 3,449 adults (age 25+ years) who completed the telephone household survey, 21% stated that they spent time actively sunbathing. In a Danish study with 164 participants, all children (age 1 to 12 years) and teenagers (age 13 to 19 years) and 94% of adults (age 20 to 76 years) had days with risk behavior (Thieden et al., 2004b). Teenagers had the highest annual UV radiation doses, as compared to children and adults, a finding likely attributable to their having the highest number of risk-behavior days. Among teenagers, 76% of their UV radiation dose during the measurement period was received on risk-behavior days, as determined using personal electronic UV dosimeters and exposure diaries (Thieden et al., 2004b). In addition to

vacationing and sunbathing, participation in outdoor sports (e.g., basketball, soccer, golfing, swimming, cycling) also significantly increased UV radiation exposure (Moehrle, 2001; Moehrle et al., 2000; Moise et al., 1999; Thieden et al., 2004a,b).

10.2.2.2 Occupation

Of the various factors that affect human exposure to UV radiation, occupation is also important. Approximately 5% of the American workforce perform work in outdoor microenvironments, as determined by the EPA NHAPS (Godar et al., 2001). American indoor workers, on average, spend ~10% of their day outdoors and are exposed to ~30% of the total ground-level UV flux, as measured by the EPA UV-monitoring program, during this time period (Godar et al., 2001). In comparison to indoor or home workers, outdoor workers are exposed to much higher levels of UV radiation (Kimlin et al., 1998a; Thieden et al., 2004a), frequently at levels that are above current exposure limits set by the International Commission on Non-Ionizing Radiation Protection (ICNIRP, 2004). For example, Thieden et al. (2004a) observed that the annual UV radiation dose, estimated using personal electronic UV dosimeters and exposure diaries, ~70% higher for gardeners than indoor workers. The gardeners received the majority (55%) of their UV radiation dose on working days (Thieden et al., 2004a). Another study found that outdoor workers received three to four times the annual UV radiation exposure of indoor workers (Diffey, 1990). At-risk working populations include farmers (Airey et al., 1997; Schenker et al., 2002), fisherman (Rosenthal et al., 1988), landscapers (Rosenthal et al., 1988), building and construction workers (Gies and Wright, 2003), physical education teachers (Vishvakarman et al., 2001), mail delivery personnel (Vishvakarman et al., 2001), and various other workers who spend the majority of their day in outdoor microenvironments during peak UV radiation hours.

10.2.2.3 Age

Age-related differences in UV radiation dose was examined in a U.S. study using the EPA NHAPS (Godar, 2001; Godar et al., 2001). The average UV radiation dose among American children (age < 12 years) was estimated to be similar to that of adults (age 20+ years), but ~20% higher than that of adolescents (age 13 to 19 years) (Godar, 2001). During the summer, higher UV radiation doses were observed in children < 5 years of age and men 40+ years of age (Goder,

2001). In a large Canadian survey, 89% of children (age < 12 years) had 30 minutes or more of daily UV exposure, compared to 51% for both adults (age 25+ years) and youth (age 15 to 24 years) (Lovato et al., 1998a, 1998b; Shoveller et al., 1998). In an English study (Diffey et al., 1996), UV radiation exposure was estimated in 180 children (age 9 to 10 years) and adolescents (age 14 to 15 years) using personal film badges and exposure records. Once again, children were found to have received higher UV radiation exposure compared to adolescents (Diffey et al., 1996). However, in contrast to the results from the studies discussed above, a Danish study observed that the annual UV radiation dose in teenagers (age 13 to 19 years) was ~20% higher compared to children (age 1 to 12 years) (Thieden et al., 2004b). This increase in UV radiation dose in teenagers was attributed to their increased risk-behavior days. The time profiles of daily UV radiation exposure among grade 8 students was assessed in an Australian study using a routinely operating UV-Biometer and questionnaires (Moise et al., 1999). The results indicated that up to 47% of the daily UV radiation dose fell within the time periods when students were outdoors during school hours, sitting under shaded structures during lunch breaks and participating in routine outdoors or sports activities (Moise et al., 1999).

Two studies examined lifetime UV radiation exposure among persons in the U.S. (Godar et al., 2001) and Denmark (Thieden et al., 2004b). Both studies observed that while there are slight differences in UV radiation dose by age, generally people receive fairly consistent UV doses at different age intervals throughout their lives.

10.2.2.4 Gender

Studies have indicated that females generally spend less time outdoors and, consequently, have lower UV radiation exposure compared to males (Gies et al., 1998; Godar et al., 2001; Shoveller et al., 1998). The U.S. study by Godar et al. (2001) observed that even though both males and females had somewhat consistent erythematous UV radiation doses throughout their lives, males received consistently higher UV doses compared to females at all age groups. Among all Americans, the lowest exposure to UV radiation was received in females during their child-raising years (age 22 to 40 years) (Godar et al., 2001). The highest exposure was observed in males aged 41 to 59 years in the U.S. study (Godar et al., 2001), while a similar Canadian survey found that younger adult males had the greatest exposures to UV radiation (Shoveller et al., 1998).

10.2.2.5 Geography

In the U.S. study by Godar et al. (2001), erythema UV radiation doses were examined in persons living in northern and southern regions. Northerners and southerners were found to spend an equal amount of time outdoors; however, the higher solar flux at lower latitudes significantly increased the annual UV radiation dose for southerners (Godar et al., 2001). The annual UV radiation doses in southerners were 25 and 40% higher in females and males, respectively, compared to northerners (Godar et al., 2001). Other studies also have shown that altitude and latitude influence personal exposure to UV radiation (Kimlin et al., 1998b; Rigel et al., 1999).

10.2.2.6 Protective Behavior

Protective behaviors such as using sunscreen (e.g., Nole and Johnson, 2004), wearing protective clothing (e.g., Rosenthal et al., 1988; Sarkar, 2004; Wong et al., 1996), and spending time in microenvironments with shaded surfaces (Moise et al., 1999; Parisi et al., 1999) have been shown to reduce exposure to UV radiation. In one study, the use of sunscreen was associated with extended intentional UV radiation exposure (Autier et al., 1999); however, a follow-up study indicated that sunscreen use increased duration of exposures to doses of UV radiation that were below the threshold level for erythema (Autier et al., 2000).

In a national study of U.S. youths aged 11 to 18 years, the most prevalent protective behavior was sunscreen use (39.2%) followed by use of a baseball hat (4.5%) (Davis et al., 2002). There were significant differences in the use of sunscreen by age group and gender, with the younger age group (age 11 to 13 years) and girls having greater likelihood (47.4 and 48.4%, respectively) of using sunscreen (Davis et al., 2002). The Canadian National Survey on Sun Exposure and Protective Behaviours observed that less than half of the adults (age 25+ years, n = 3,449) surveyed took adequate protective actions (Shoveller et al., 1998). Once again, children (age < 12 years, n = 1,051) were most protected from exposure to UV radiation, with 76% using sunscreen and 36% avoiding the sun, as reported by their parents (Lovato et al., 1998a). However, the protection level was still not adequate, as indicated by the high 45% rate of erythema in children. Among Canadian youth (age 15 to 24 years, n = 574), protective actions from UV radiation exposure included wearing a hat (38%) and seeking shade and avoiding the sun between the peak hours of 11:00 a.m. to 4:00 p.m. (26%) (Lovato et al., 1998b).

The lowest prevalence of protective behavior among the youth was likely responsible for the highest proportion of erythema (68%) experienced in this age group. A Danish study observed that both children and teenagers applied sunscreen on more days than adults, but teenagers had the most days with erythema, due to their increased risk behavior (Thieden et al., 2004b). A survey in Switzerland of 1,285 individuals, including children and parents, indicated that sunscreen use was the protective action most commonly used, but only at the beach and not in routine daily exposure (Berret et al., 2002). Protective clothing and avoiding the sun were not highly used among these individuals.

10.2.2.7 Summary of Factors that Affect Human Exposures to Ultraviolet Radiation

The factors that potentially influence UV radiation doses were discussed in the previous sections and include choice of leisure activities, occupation, age, gender, geography, and protective behavior. Results from the various studies indicate that specific populations may be at risk for higher exposures to UV radiation. Of particular concern are the following potentially susceptible populations:

- Individuals who engage in high-risk behavior, viz., sunbathing;
- Individuals who participate in outdoor sports and activities, including professional athletes;
- Individuals who work in outdoor microenvironments with inadequate shade, e.g., farmers, fishermen, gardeners, landscapers, building and construction workers;
- Young children; and
- Individuals living in geographic areas with higher solar flux (i.e., lower latitudes [e.g., Honolulu, HI] and higher altitudes [e.g., Denver, CO]).

10.2.3 Factors Governing Human Health Effects due to Ultraviolet Radiation

Ultraviolet radiation occupies a specific region of the electromagnetic spectrum of wavelengths and can be further subdivided into three parts, UV-A (320 to 400 nm), UV-B (280 to 320 nm), and UV-C (200 to 280 nm). Most of the health risks associated with UV radiation exposure are wavelength dependent. Wavelengths < 180 nm are of little practical biological significance as they are readily absorbed in the air (ICNIRP, 2004). Until 1980, it was generally thought that wavelengths < 315 nm were responsible for the most significant

adverse UV radiation health effects; however, recent studies have found that the longer wavelengths in the UV-A range also might produce adverse responses at substantially higher doses (ICNIRP, 2004).

Action spectra of a given biological response to UV radiation across its spectral range are used to estimate exposure by weighting individual wavelength intensities according to the associated response. The overall effectiveness of the incident flux at inducing the biological response of interest is computed by means of the relationship

$$\text{effective irradiance} = \int_{\lambda} I_{\lambda} E_{\lambda} d\lambda \quad (10-4)$$

where I_{λ} and E_{λ} are, respectively, the irradiance and its relative effectiveness at wavelength λ .

UV-A and UV-B radiations differ in their abilities to initiate DNA damage, cell signaling pathways, and immune alterations. In this section, the various adverse health effects associated with acute and chronic UV radiation exposure will be discussed.

10.2.3.1 Erythema

Association Between Ultraviolet Radiation Exposure and Erythema

The most conspicuous and well-recognized acute response to UV radiation is erythema, or the reddening of the skin, which is likely caused by direct damage to DNA by UV-B and UV-A radiation (Matsumura and Ananthaswamy, 2004). Indirect oxidative damage also might occur at longer wavelengths (Matsumura and Ananthaswamy, 2004). Skin type appears to play a large role in the sensitivity to UV radiation-induced erythema. The Fitzpatrick classifications for skin types are: (1) skin type I – individuals with extremely sensitive skin that sunburns easily and severely, and is not likely to tan (e.g., very fair skin, blue eyes, freckles); (2) skin type II – individuals with very sensitive skin that usually sunburns easily and severely, and tans minimally (e.g., fair skin, red or blond hair, blue, hazel or brown eyes); (3) skin type III – individuals with sensitive skin that sunburns moderately and tans slowly (e.g., white skin, dark hair); (4) skin type IV – individuals with moderately sensitive skin that sunburns minimally and usually tans well (e.g., white or light brown skin, dark hair, dark eyes); (5) skin type V – individuals with minimally sensitive skin that rarely sunburns and tans deeply (e.g., brown

1 skin); and (6) skin type VI – individuals with nonsensitive skin that never sunburns and tans
2 profusely (e.g., dark skin). A study by Harrison and Young (2002) found that the perceptible
3 minimal erythema dose was approximately two-fold greater among individuals with skin type
4 IV compared to skin type I, although there was considerable overlap in the minimal erythema
5 dose among the four skin types. Waterston et al. (2004) further observed that within an
6 individual, erythema response differed by body site (e.g., abdomen, chest, front upper arm, back
7 of thigh). These differences were likely attributable to body site-specific variations in melanin
8 pigmentation.

9 Kollias et al. (2001) investigated the change in erythema response following a previous
10 exposure to UV radiation. Body sites that received a second exposure to UV radiation always
11 showed a reduced erythema response compared to body sites with a single exposure, especially
12 when the first exposure was at levels greater than the minimal erythema dose. The suppression
13 of erythema was more pronounced when the second exposure was given 48 hours after the first.
14 However, Kaidbey and Kligman (1981) found that multiple exposures to subthreshold doses of
15 UV radiation at 24-hour intervals were found to lower the minimal erythema dose. Henrisken
16 et al. (2004) observed that the change in threshold depended on skin type. In 49 healthy
17 volunteers with skin types II, III, and IV, just perceptible erythema 24 hours post-exposure was
18 chosen as the minimal erythema dose. After four days of repeated UV radiation, the minimal
19 erythema dose was lowered by 40 to 50% in darker-skinned persons. However, among fair-
20 skinned individuals, there was no change in the erythema threshold dose with repeated exposure
21 to UV radiation.

22 A reference erythema action spectrum was adopted by the Commission Internationale de
23 l'Eclairage (International Commission on Illumination, CIE) in 1987 (McKinlay and Diffey,
24 1987). The CIE erythema action spectrum indicates that UV-B radiation is orders of magnitude
25 more effective per unit dose than UV-A radiation. However, a follow-up study by Diffey (1994)
26 that compared the observed and predicted minimal erythema doses found that the erythema
27 sensitivity of skin at longer UV wavelengths (> 350 nm) was greater than predicted from the CIE
28 reference action spectrum.

Risk of Erythema from Changes in Tropospheric Ozone Levels

There is no literature on the risk of erythema associated with changes in tropospheric or ground-level O₃ levels. However, one study conducted a risk assessment of the effects of stratospheric O₃ depletion on the risk of erythema (Longstreth et al., 1998). Stratospheric O₃ depletion will result in only a slight increase influence rate of UV-A as O₃ only absorbs a very little part of this UV spectrum. However, the ground-level UV-B flux will likely increase as O₃ absorbs radiation in that wavelength range with high efficiency.

The risk analysis by Longstreth et al. (1998) concluded that erythema will not appreciably increase with depletion of the O₃ layer. This is due to the powerful adaptation of the skin to different levels of UV radiation, as evidenced by its ability to cope with changes in UV radiation by season (van der Leun and de Gruijl, 1993). Gradual exposure to increasing UV radiation from the winter to summer leads to decreased sensitivity of the skin. In midlatitudes, the UV-B radiation in the summer is ten-fold greater than in the winter. In contrast, the steady depletion of the O₃ layer has been estimated to likely lead to an ~20% increase in UV-B in 10 years (Longstreth et al., 1998). Such a comparatively small increase in UV radiation throughout the years, therefore, would not seem likely to significantly increase the risk of erythema. Given that the stratospheric O₃ depletion is estimated as not being likely to affect the risk of erythema, one can conclude that changes in ground-level O₃ (which only constitutes no more than 10% of total atmospheric O₃) is not likely to result in increased risk.

10.2.3.2 Skin Cancer

According to the Skin Cancer Foundation, one in six Americans will develop skin cancer during their lifetime (Gloster and Brodland, 1996). The three main forms of skin cancer include basal cell carcinoma and squamous cell carcinoma, which are both nonmelanoma skin cancers, and malignant melanoma. Nonmelanoma skin cancers constitute more than one-third of all cancers in the U.S. and ~90% of all skin cancers, with basal cell carcinoma being approximately four times as common as squamous cell carcinoma (Diepgen and Mahler, 2002; ICNIRP, 2004). The incidence of malignant melanoma (3 to 4% of all cancers) is much lower compared to nonmelanoma skin cancers, but melanoma has great metastatic potential and accounts for the majority of skin cancer deaths (Jemal et al., 2004).

Exposure to UV radiation is considered to be a major risk factor for all three forms of skin cancer (Gloster and Brodland, 1996; Diepgen and Mahler, 2002; IARC, 1992). Ultraviolet radiation is especially effective in inducing genetic mutations and acts as both a tumor initiator and promoter. Keratinocytes have evolved DNA repair mechanisms that correct the damage induced by UV, but mutations do occur, leading to skin cancers that are appearing with increasing frequency (Hildesheim and Fornace, 2004). The relationship between skin cancer and chronic exposure to UV radiation is further explored below, followed by discussion of the influence of O₃ on the incidence of skin cancer.

10.2.3.3 Ultraviolet Radiation Exposure and the Incidence of Nonmelanoma Skin Cancers

The incidence of all three types of cancers has been shown to rise with increasing UV radiation concentrations across the U.S. (de Gruijl, 1999); however, the most convincing evidence for a causal relationship exists between UV radiation and squamous cell carcinoma. Squamous cell carcinoma occurs almost exclusively on skin that is regularly exposed to the sun, such as the face, neck, arms, and hands. The incidence is higher among whites in areas of lower latitudes, where solar flux is greater (Kricker et al., 1994). The risk of squamous cell carcinoma was shown to increase with life-long accumulated exposure to UV radiation in one cross-sectional study (Vitasa et al., 1990), but was found to be associated only with the exposure ten years prior to diagnosis in a case-control study (Gallagher et al., 1995a). One of the major concerns with both types of studies is the potential for recall bias in reporting their UV radiation exposure as individuals are already aware of their disease status.

Ultraviolet radiation also has been linked to basal cell carcinoma. Basal cell carcinoma is common on the face and neck (80-90%) but rarely occurs on the back of the hands (de Gruijl, 1999). While cumulative UV radiation exposure was not associated with risk of basal cell carcinoma (Vitasa et al., 1990), increased risk was observed in individuals with increased recreational UV radiation exposure in adolescence and childhood (age < 19 years) and individuals with a history of severe erythema in childhood (Gallagher et al., 1995b). Once again, consideration must be given to potential recall bias in assessing these results. Thus, there is suggestive evidence that UV radiation also plays a role in the development of basal cell carcinoma, but the etiologic mechanisms for squamous cell carcinoma and basal cell carcinoma likely differ. In an Australian study conducted in a subtropical community, having fair skin,

1 a history of repeated sunburns, and nonmalignant solar skin damage diagnosed by dermatologists
2 were strongly associated with both types of nonmelanoma skin cancer (Green et al., 1996).
3 Outdoor occupation was not associated with nonmelanoma skin cancer, which was likely due to
4 significant self-selection. Individuals with fair or medium complexions and a tendency to
5 sunburn, though they accounted for more than 80% of the community study sample, were
6 systematically underrepresented among outdoor workers (Green et al., 1996). This selection bias
7 may partly explain the lack of consistent quantitative evidence of a causal link between UV
8 radiation and skin cancer in humans.

9 De Gruijl et al. (1993) assessed the action spectrum for nonmelanoma skin cancers using
10 hairless albino mice. Human data are not available to examine the wavelength dependence of
11 the carcinogenicity of UV radiation. After adjusting for species differences, the Skin Cancer
12 Utrecht-Philadelphia action spectrum indicated the highest effectiveness in the UV-B range with
13 a maximum at 293 nm, which dropped to 10^{-4} of this maximum at the UV-A range above
14 340 nm (de Gruijl et al., 1993). The mutations commonly present in the *p53* tumor suppressor
15 gene in individuals with squamous cell carcinoma and basal cell carcinoma are called the
16 “signature” mutations of UV-B radiation (de Gruijl, 2002). UV-B radiation is highly mutagenic
17 due to the fact that DNA is a chromophore for UV-B, but not for UV-A, radiation (Ichihashi
18 et al., 2003). However, other studies have found that UV-A radiation, in addition to UV-B
19 radiation, can induce DNA damage (Persson et al., 2002; Ruenger et al., 2000). DNA damage by
20 UV-A is mediated by reactive oxygen species, thus is indistinguishable from damage caused by
21 other agents that generate reactive oxygen species (de Gruijl, 2002). Epidemiologic evidence of
22 a carcinogenic effect of UV-A was found in a study of psoriasis patients receiving oral psoralen
23 and UV-A radiation treatment (Stern et al., 1998). High-dose exposure to oral psoralen and
24 UV-A radiation was associated with a persistent, dose-related increase in the risk of squamous
25 cell cancer, but with much less effect on the risk of basal cell cancer. Therefore, although UV-B
26 radiation has long been considered the main culprit for nonmelanoma skin cancer, limited
27 evidence suggest that UV-A radiation may also play a role.

28 Susceptible populations for nonmelanoma skin cancers include individuals with reduced
29 capacity for nucleotide excision repair, the primary repair mechanism for UV radiation-induced
30 DNA lesions (Ichihashi et al., 2003). At particular risk are individuals with xeroderma
31 pigmentosum, as they have defective nucleotide excision repair in all tissues (Kraemer, 1997;

1 Sarasin, 1999). Skin type also largely affects susceptibility to skin cancer. Of the six skin
2 phenotypes, the most sensitive individuals are those with skin types I and II, who have a fair
3 complexion, blue or green eyes, and red or blond hair (Diepgen and Mahler, 2002). These
4 individuals tend to sunburn easily, tan poorly, and freckle with sun exposure. A history of
5 repeated sunburns also appears to increase the risk of both cancers, while sunburns during
6 childhood are more associated with increased basal cell carcinoma (Gallagher et al., 1995b;
7 Green et al., 1996).

9 ***Ultraviolet Radiation and the Incidence of Cutaneous Malignant Melanoma***

10 From 1973 to 1994, the incidence rate of melanoma increased 120.5% along with an
11 increased mortality rate of 38.9% among whites in the U.S. (Hall et al., 1999). The ICNIRP
12 (2004) states that during the past 40 years or so, each decade has seen a two-fold increase in the
13 incidence of malignant melanoma in white populations, with increased incidence observed more
14 prominently in individuals living in lower latitudes. Cutaneous malignant melanoma has
15 multifactorial etiology, with environmental, genetic, and host factors (Lens and Dawes, 2004).
16 While the major environmental factor of melanoma has been identified as UV radiation
17 exposure, the risk of melanoma appears to depend on the interaction between the nature of the
18 exposure and skin type (Lens and Dawes, 2004).

19 Fears et al. (2002) examined the association between invasive cutaneous melanoma and
20 UV radiation in non-Hispanic whites using a case-control study design. Lifetime residential
21 history was coupled with mid-range UV-B radiation flux measurements to reduce exposure
22 misclassification and recall bias. A 10% increase in the average annual UV-B flux was
23 significantly associated with a 19% increase in individual odds for melanoma in men and a
24 16% increase in women. Whiteman et al. (2001) conducted a systematic review of studies that
25 examined the association between childhood UV radiation exposure and risk of melanoma.
26 Researchers found that ecological studies assessing ambient sun exposure consistently reported
27 higher risks of melanoma among people who resided in an environment with high UV radiation
28 during their childhood (Whiteman et al., 2001). The lack of consistency among the case-control
29 studies was likely due to the varying methods used to assess UV radiation dose.

30 While the evidence is generally suggestive of a causal relationship between UV radiation
31 and malignant melanoma, possibly conflicting data has been observed. For example, the highest

1 occurrence of malignant melanoma is on men's backs and women's legs, areas that do not have
2 prolonged exposure to the sun (Rivers, 2004). This indicates that malignant melanoma tends to
3 occur in sites of intermittent, intense sun exposure (trunk and legs), rather than in areas of
4 cumulative sun damage (head, neck, and arms), in contrast to nonmelanoma skin cancers
5 (Swetter, 2003).

6 The available data conflict with regard to the relative importance of UV-A versus UV-B in
7 inducing melanomas. UV-A has a much higher flux rate at the earth's surface, as it is not
8 absorbed by O₃ and it is able to penetrate more deeply into the skin surface due to its longer
9 wavelength. However, UV-B, as mentioned earlier, is much more energetic and, therefore, more
10 effective in photochemically altering DNA. The individual roles of UV-B and UV-A in the
11 development of cutaneous malignant melanoma have been examined in several studies.
12 A case-control study of 571 patients and 913 matched controls observed an elevated odds ratio
13 for development of malignant melanoma in individuals who regularly used tanning beds, which
14 typically are UV-A sources (Westerdahl et al., 2000). In a study by Setlow et al. (1993),
15 an action spectrum using the tropical fish *Xiphophorus* indicated that UV-A range wavelengths
16 were especially important in malignant melanoma induction. However, an action spectrum
17 using the opossum *Monodelphis domestica* found that the potency of UV-A for melanoma
18 induction was extremely low compared to that of UV-B (Robinson et al., 2000). A recent study
19 by De Fabo et al. (2004) examined the differences in wavelength effectiveness using a
20 hepatocyte growth factor/scatter factor-transgenic mouse model. The epidermal tissue of these
21 transgenic mice behaves very similarly to the human epidermis in response to UV exposure.
22 Given the absence of a mammalian melanoma action spectrum, the standardized CIE erythema
23 action spectrum was used to deliver identical erythemally effective doses. Only UV-B radiation
24 was found to initiate mammalian cutaneous malignant melanoma. UV-A radiation, even at
25 doses considered physiologically relevant, were ineffective at inducing melanoma (De Fabo
26 et al., 2004). Overall, current evidence suggests that UV-B, and not UV-A, is the primary risk
27 factor for malignant melanoma (ICNIRP, 2004).

28 The populations susceptible for malignant melanoma are similar to those for nonmelanoma
29 skin cancers. Once again, individuals with xeroderma pigmentosum or a reduced capacity of
30 nucleotide excision repair are at increased risk (Tomescu et al., 2001; Wei et al., 2003).
31 Individuals with skin types I and II, or the fair-skin phenotype (blue or green eyes; blond or red

1 hair; skin that freckles, sunburns easily, and does not tan), have increased susceptibility to
2 malignant melanoma (Evans et al., 1988; Swetter, 2003; Veierød et al., 2003). However, the
3 incidence of melanoma was also positively associated with UV radiation in Hispanics and blacks
4 (Hu et al., 2004). Although the incidence of melanoma is much lower in Hispanics and blacks
5 compared to whites, melanomas in these populations are more likely to metastasize and have a
6 poorer prognosis (Black et al., 1987; Bellows et al., 2001). Among children, malignant
7 melanoma appears to have similar epidemiologic characteristics to the adult form of the disease
8 (Whiteman et al., 1997). Individuals with intermittent, intense sun exposure, particularly during
9 childhood, were found to have increased risk of melanoma (Whiteman et al., 2001), in contrast
10 to the association between cumulative exposure and increased risk of squamous cell carcinoma.
11 One study found that a personal history of nonmelanoma skin cancer or precancer, higher
12 socioeconomic status, and increased numbers of nevocytic nevi also were associated with
13 increased incidence of melanoma (Evans et al., 1988).

15 ***Effect of Changes in Tropospheric Ozone Levels on Skin Cancer Incidence***

16 The current evidence strongly suggests a causal link between exposure to UV radiation and
17 the incidence of both nonmelanoma and melanoma skin cancer. Genetic factors, including skin
18 phenotype and ability to repair DNA, affect an individual's susceptibility to skin cancer.
19 Quantifying the relationship between UV radiation and skin cancer is complicated by the
20 uncertainties involved in the selection of an action spectrum and appropriate characterization of
21 dose (e.g., peak or cumulative levels of exposure, childhood or lifetime exposures). In addition,
22 there are multiple complexities in attempting to quantify the effect of tropospheric O₃ levels on
23 UV-radiation exposure, as described in Section 9.11.2. There is an absence of published studies
24 that critically examine any increased incidence of skin cancer that may be attributable to
25 decreased tropospheric O₃ exposures (which reflects the significant challenges in determining
26 ground-level O₃-related changes in UV radiation exposure).

27 Several studies have examined the potential effect of stratospheric O₃ depletion on the
28 incidence of skin cancer (de Gruijl, 1995; Longstreth et al., 1995; Madronich and de Gruijl,
29 1993; Slaper et al., 1996). Stratospheric O₃ depletion is likely to increase the ground-level UV-B
30 flux, as O₃ absorbs radiation in that wavelength range with high efficiency. Because UV-B
31 radiation is primarily implicated in the induction of skin cancer, especially among persons with

1 skin phenotypes I and II, there is concern that the depletion of the O₃ layer will result in
2 significantly increased incidence of skin cancers.

3 Estimation of the increased risk in melanoma associated with O₃ depletion cannot be done
4 adequately due to the lack of a mammalian action spectrum for melanoma. In addition, the
5 complexity of the UV-related induction mechanism of melanoma adds an additional layer of
6 uncertainty to the calculations. The excess risk in nonmelanoma skin cancers associated with a
7 decrease in stratospheric O₃ was estimated using the Skin Cancer Utrecht-Philadelphia action
8 spectrum based on hairless albino mice (Longstreth et al., 1995). Quantification of how much
9 more UV radiation would reach ground level with each percentage decrease in O₃ required
10 several assumptions: (1) annual doses are an appropriate measure; (2) personal doses are
11 proportional to ambient doses; and, most notably, (3) each percentage decrease in O₃ is
12 associated with a 1.2% increase in UV radiation. Next, the relationship between UV radiation
13 and nonmelanoma skin cancer incidence was determined. Each percent increase in annual UV
14 radiation dose was estimated to cause a 2.5% increase in squamous cell carcinoma and 1.4%
15 increase in basal cell carcinoma over a human lifetime. Incorporating all these factors,
16 Longstreth et al. (1995) calculated that a sustained 10% decrease in stratospheric O₃
17 concentration would result in 250,000 additional nonmelanoma skin cancer cases per year.
18 Madronich and de Gruijl (1993) noted that the largest percent of O₃-induced nonmelanoma skin
19 cancer increases would be at high latitudes, where baseline incidence of skin cancer is usually
20 small. Assuming a phaseout of primary O₃-depleting substances by 1996, as established by the
21 Copenhagen Amendments in 1992, Slaper et al. (1996) estimated that the number of excess
22 nonmelanoma skin cancers in the U.S. caused by O₃ depletion would exceed 33,000 per year
23 (or ~7 per 100,000) around the year 2050.

24 However, estimating the increase in nonmelanoma skin cancer incidence attributable to the
25 depletion of the stratospheric O₃ layer is marred by uncertainty. The following statement by
26 Madronich and de Gruijl (1994) describes the uncertainty of estimating the effect of
27 stratospheric O₃ depletion on the incidence of skin cancer:

28 Extrapolating trends and effects of UV into the future is very hypothetical due to
29 uncertainties that arise from atmospheric chemistry, epidemiology, and related
30 disciplines. The values that we calculated are one plausible measure of the
31 magnitude of the O₃-UV effects....The timescales for atmospheric change and skin-
32 cancer development are still far from certain: O₃ reductions are expected to continue
33 well into next century, and the time between UV exposure and development of skin
34 cancer is essentially unknown. . .
35

Therefore, much caution is necessary when assessing and interpreting the quantitative results of excess nonmelanoma skin cancer incidence due to stratospheric O₃ depletion. Note that although the effect of changes in ground-level or tropospheric O₃ concentrations on skin cancer incidence has not been assessed, it would be expected to be much less compared to the effect from the depletion of the stratospheric O₃ layer, given that tropospheric O₃ makes up $\leq 10\%$ of the total atmospheric O₃.

10.2.3.4 Ocular Effects of Ultraviolet Radiation Exposure

Ultraviolet Radiation Exposure and Risk of Ocular Damage

Ocular damage from UV radiation exposures includes effects on the cornea, lens, iris, and associated epithelial and conjunctival tissues. Absorption of UV radiation differs by wavelength, with short wavelengths (< 300 nm) being almost completely absorbed by the cornea, whereas longer wavelengths are transmitted through the cornea and absorbed by the lens (McCarty and Taylor, 2002). The most common acute ocular effect of environmental UV exposure is photokeratitis, also known as snowblindness, caused by absorption of short wavelength UV radiation by the cornea. The action spectrum indicated that maximum sensitivity of the human eye was found to occur at 270 nm (ICNIRP, 2004; Pitts, 1993). The threshold for photokeratitis in humans varied from 4 to 14 mJ/cm² for wavelengths 220 to 310 nm. For UV-A radiation, levels exceeding 10 J/cm² were necessary for corneal injury.

Exposure to longer wavelengths has been shown to cause both transient and permanent opacities of the lens, or cataracts. There is extensive toxicologic and epidemiologic evidence supporting the causal association between UV radiation and cataracts (Hockwin et al., 1999; McCarty and Taylor, 2002). Ultraviolet radiation-induced cataracts are hypothesized to be caused by oxidative stress leading to increased reactive species in the lens, which then causes damage to lens DNA and cross-linking of proteins. Exposure time to low-dose UV radiation was found to strongly influence cataract formation (Ayala et al., 2000). An action spectrum determined using young female rats indicated that the rat lens was most sensitive to 300 nm, correcting for corneal transmittance (Merriam et al., 2000). Oriowo et al. (2001) examined the action spectrum for cataract formation using whole cultured lens from pigs. As pigs lens are similar in shape and size to the human lens, some inferences may be made. Results indicated that the 270 to 315 nm waveband was most effective in producing UV-induced cataracts in vitro.

1 However, the threshold values varied widely within that range, from 0.02 J/cm² for 285 nm to
2 0.74 J/cm² for 315 nm (Oriowo et al., 2001). At wavelengths > 325 nm, the threshold levels
3 were orders of magnitude larger, with a minimum threshold value of 18.7 J/cm². A U.S. study of
4 838 watermen found that UV-B radiation was significantly associated with cortical, but not
5 nuclear, cataract formation (Taylor et al., 1988). No association was observed between cataracts
6 and UV-A radiation in this outdoor-working population.

7 8 ***Risk of Ocular Damage from Changes in Tropospheric Ozone Levels***

9 Cataracts are the most common cause of blindness in the world. McCarty et al. (2000)
10 calculated that ocular UV radiation exposure accounted for 10% of the cortical cataracts in an
11 Australian cohort of 4,744 individuals from both urban and rural areas. A study by Javitt and
12 Taylor (1994-1995) found that the probability of cataract surgery in the U.S. increased by 3% for
13 each 1E decrease in latitude. These results suggest that depletion of the stratospheric O₃ layer
14 may increase UV radiation-induced cataract formation. After assuming a certain wavelength
15 dependency along with some additional assumptions, every 1% decrease in the stratospheric O₃
16 layer was estimated to be associated with a 0.3 to 0.6% increase in cataracts (Longstreth et al.,
17 1995). Longstreth et al. (1995) noted that this estimate has a high degree of uncertainty due to
18 inadequate information on the action spectrum and dose-response relationships. Quantitative
19 estimates have not been possible for photokeratitis, pterygium, or other UV-related ocular effects
20 due to lack of epidemiologic and experimental data. As is the case for all of the other UV-
21 related health outcomes, there is no published information on the potential effects of decreased
22 tropospheric O₃ concentrations on cataract formation due to any changes in surface-level UV
23 flux resulting from decreases in tropospheric O₃.

24 25 **10.2.3.5 Ultraviolet Radiation and Immune System Suppression**

26 Experimental studies have suggested that exposure to UV radiation may suppress local and
27 systemic immune responses to a variety of antigens (Clydesdale et al., 2001; Garssen and van
28 Loveren, 2001; Selgrade et al., 1997). In rodent models, these effects have been shown to
29 worsen the course and outcome of some infectious diseases and cancers (Granstein and Matsui,
30 2004; Norval et al., 1999). Granstein and Matsui (2004) stated that exposure to UV-B radiation
31 caused immunosuppression in mice ultimately by releasing cytokines that prevent antigen-

1 presenting cells from performing their normal functions and causing direct damage to epidermal
2 Langerhans cells. Noonan et al. (2003) investigated UV skin cancer induction in two strains of
3 reciprocal F1 hybrid mice and found that genetically determined differences in susceptibility to
4 UV-induced immunosuppression was a risk factor for skin cancer. At high-UV radiation doses,
5 mice with greater susceptibility to immune suppression had a larger proportion of skin tumors
6 compared to those with low susceptibility (Noonan et al., 2003).

7 While results from animal models are supportive of an association between UV radiation
8 and local and systemic immunosuppression, evidence for UV-induced immunosuppression in
9 humans is limited. There is some evidence that UV radiation has indirect involvement in viral
10 oncogenesis through the human papillomavirus (Pfister, 2003). Additional evidence of
11 UV-related immunosuppression comes from an epidemiologic study of 919 patients with rare
12 autoimmune muscle diseases from 15 cities on four continents with variable UV radiation
13 intensity (Okada et al., 2003). Ultraviolet radiation was strongly associated with the prevalence
14 of dermatomyositis, an autoimmune disease distinguished by the presence of photosensitive
15 pathognomonic rashes (Okada et al., 2003). In patients with the human immunodeficiency virus,
16 UV-B radiation lead to activation of the virus in their skin through the release of cytoplasmic
17 nuclear factor kappa B (Breuher-McHam et al., 2001). In a study by Selgrade et al. (2001),
18 UV-induced immunosuppression was examined in 185 subjects with different skin
19 pigmentations. To assess immune suppression, dinitrochlorobenzene was applied to irradiated
20 buttock skin 72 hours after irradiation. Differences in sensitivity were not related to skin type
21 based on the Fitzpatrick classification or minimal erythral dose (Selgrade et al, 2001).
22 However, erythral reactivity, assessed by the steepness of the erythral dose-response curve,
23 was shown to be significantly associated with UV-induced immunosuppression. Only subjects
24 with steep erythral responses, which included individuals with skin types I through V, showed
25 a dose-response relationship between UV exposure and immune suppression (Selgrade et al,
26 2001).

27 Most action spectrum investigations have concluded that immunosuppression is caused
28 most effectively by the UV-B waveband (Garssen and van Loveren, 2001). The effects of UV-A
29 on local and systemic immunosuppression have been unclear and inconsistent. There is some
30 evidence that high doses of UV-A is protective of immunosuppression induced by UV-B
31 exposure (Halliday et al., 2004). Given the variety of outcomes of immune suppression and

possible mechanisms of effect, little detailed information exists on UV radiation action spectrums and dose-response relationships. The available data are insufficient to develop quantitative risk estimates for UV radiation-induced immunosuppression in humans.

10.2.3.6 Protective Effects of Ultraviolet Radiation – Production of Vitamin D

Most humans depend on sun exposure to satisfy their requirements for vitamin D (Holick, 2004). UV-B photons are absorbed by 7-dehydrocholesterol in the skin, leading to its transformation to previtamin D₃, which is rapidly converted to vitamin D₃. Vitamin D₃ is metabolized in the liver, then in the kidney to its biologically active form. Vitamin D deficiency is known to cause metabolic bone disease among children and adults, but also may increase the risk of many common chronic diseases, including type I diabetes mellitus and rheumatoid arthritis (Holick, 2004). Vitamin D also is capable of inhibiting the growth of various cancer cells in cell cultures. Thus, vitamin D deficiency may lead to increased incidence of carcinomas in various organs (Studzinski and Moore, 1995).

Large geographical gradients in mortality rates for a number of cancers in the U.S. are not explained by dietary or other risk factors; therefore, it has been hypothesized that some carcinomas are due to insufficient UV-B radiation. For example, published literature indicates that solar UV-B radiation, by increasing production of vitamin D, is associated with reduced risk of cancer of the breast (Freedman et al., 2002; Garland et al., 1990; Gorham et al., 1990; Grant, 2002a; John et al., 1999), colon (Freedman et al., 2002), ovary (Freedman et al., 2002; Lefkowitz and Garland, 1994), and prostate (Freedman et al., 2002; Hanchette and Schwartz, 1992), as well as non-Hodgkin lymphoma (Hartge et al., 1996). Eight other cancers, including bladder, esophageal, kidney, lung, pancreatic, rectal, stomach, and corpus uteri, have been found to exhibit an inverse correlation between mortality rates and UV-B radiation (Grant, 2002b).

Using UV-B data from July 1992 and U.S. cancer mortality rates from 1970 to 1994, premature cancer deaths attributable to reduced UV-B exposure were analyzed in an ecologic study (Grant, 2002b). The annual number of premature deaths from various cancers due to lower UV-B exposures was 21,700 for white Americans; 1,400 for black Americans; and 500 for Asian Americans and other minorities. Uncertainty in the estimation of UV-B exposure limits the confidence for the estimates of excess cancer deaths attributable to reduced exposure. No study has assessed the decreased risk of cancer mortality resulting from increased UV

radiation attributable to decreased tropospheric O₃ levels, but the change in risk is expected to be unappreciable.

In establishing guidelines on limits of exposure to UV radiation, the ICNIRP agreed that some low-level exposure to UV radiation benefits health (ICNIRP, 2004). However, the adverse health effects necessitated the development of exposure limits for UV radiation. The ICNIRP recognized the challenge in establishing exposure limits that would achieve a realistic balance between beneficial and adverse health effects.

10.2.4 Summary and Conclusions for O₃ Effects on UV-B Flux

Increased solar radiation, especially UV-B radiation, can be detrimental as well as beneficial to the exposed human population. Analogously, both positive and negative effects of UV-B radiation would be expected on plant and animal biota, and on man-made materials (e.g., Van der Leun et al., 1998; U.S. Environmental Protection Agency, 2002). Other environmental factors include the daily nonlinearity of UV-B flux, nonlinear absorption by O₃, and absorption and scattering of UV-B radiation by both stratospheric and tropospheric processes, including ground-level environmental pollutants such as PM and O₃. Surface flux variability of UV-B is partially dependent upon cloud and tree cover, latitude, time of year, and time of day. Given the magnitude of this natural variability, it would be extremely difficult with current instrumentation and atmospheric models to measure and/or estimate a change in the ground-level UV-B flux and to attribute it to small reductions in tropospheric O₃ and/or PM.

Of equal or even greater importance are changes in human habits of daily activities, recreation, dress, and skin care. Little is known about the impact of variability in these human factors that can modify individual exposure to UV radiation. Most information about potential UV exposure to the human population relies on demographic information. There is some information available on activity patterns (e.g., EPA NHAPS) and protective behaviors (e.g., Canadian National Survey on Sun Exposure and Protective Behaviors), but the data are inadequate to accurately assess UV radiation exposure in the population. The uncertainties in characterizing UV radiation exposure limit any UV-related health risk assessment.

Another difficulty in examining health risks of UV radiation exposure in the population is due to the lack of detailed information regarding the relevant type (e.g., peak or cumulative) and time period (e.g., childhood or lifetime) of exposure, wavelength dependency of biological

1 responses, and interindividual variability in UV resistance. For example, exposure to solar UV-
2 B radiation appears to be the most important environmental risk factor for basal and squamous
3 cell carcinomas and cutaneous malignant melanoma in fair-skinned individuals. Originally, total
4 cumulative exposure to UV-B was believed to be the most important environmental factor in
5 determining risk for these tumors. New information now suggests that only squamous cell
6 carcinoma risk is related to total accumulated exposure. For basal cell carcinoma and for
7 malignant melanoma, new information suggests that increases in risk are linked to early
8 exposures (before about age 15), particularly those leading to severe sunburns. There is also
9 controversy regarding the effect of UV-A on health outcomes. Though most studies have found
10 that UV-B radiation is highly mutagenic, some evidence indicates that UV-A radiation, at high
11 doses, may also be carcinogenic.

12 Susceptibility to the various UV radiation-induced health effects, generally, appears to be
13 related to skin type. Numerous studies have shown that individuals with Fitzpatrick skin types I
14 and II (i.e., fair-skinned phenotypes) are at increased risk for erythema and skin cancer.
15 In addition, individuals with a reduced ability to repair DNA have increased susceptibility to
16 both nonmelanoma and melanoma skin cancers. For UV-related immunosuppression, however,
17 it appears that erythema reactivity, rather than skin type, is a marker for susceptibility.

18 There is some evidence that increased UV radiation exposures resulting from depletion of
19 the stratospheric O₃ layer will lead to increases in skin cancer and cataracts. However, the
20 numerous assumptions necessary to calculate these quantitative risk estimates lead to high levels
21 of uncertainty. No study has yet examined the benefits of increased UV radiation exposure due
22 to enhanced production of vitamin D. With the currently available information, the effect of
23 changes in tropospheric O₃ on UV-induced health outcomes cannot be critically assessed.

26 **10.3 TROPOSPHERIC OZONE AND CLIMATE CHANGE**

27 Water vapor, CO₂, O₃, N₂O, CH₄, CFCs, and other polyatomic gases present in the earth's
28 troposphere, trap infrared radiation emitted by the earth's surface, leading to surface warming.
29 This phenomenon is widely known as the "Greenhouse Effect" (Arrhenius, 1896), and the gases
30 involved are known as "greenhouse gases" (GHGs). The term used for the role a particular
31 atmospheric component, or any other component of the greater climate system, plays in altering

1 the earth's radiative balance is "forcing." In the past decade, the global atmospheric sciences
2 and climate communities have made significant progress in determining the specific role O₃
3 plays in forcing climate.

4 The Intergovernmental Panel on Climate Change (IPCC) was founded in 1988 by the
5 World Meteorological Society (WMO) and the United Nations Environmental Program (UNEP)
6 to support the work of the Conference of Parties (COP) to the United Nations Framework
7 Convention on Climate Change (UNFCCC). Drawing from the global climate and atmospheric
8 sciences community for its authors and reviewers, the IPCC produces reports containing
9 thorough assessments of the available, peer-reviewed science regarding the physical climate
10 system, past and present climate, and evidence of human-induced climate change. This section
11 will summarize the reviews of the available information on the forcing properties of tropospheric
12 O₃ provided by IPCC Third Assessment Report (IPCC, 2001a) will be given, and will describe
13 some of the more recent developments on the subject.

14 The projected effects of global climate change will be briefly explained to provide the
15 context within which O₃ serves as a regional, and possibly global, anthropogenic pollutant.
16 The concept of climate forcing is also explained, along with the factors that influence the extent
17 of climate forcing by ozone. The section concludes with a summary of the various estimates that
18 have been placed on the globally-averaged forcing due to ozone.

20 **10.3.1 The Projected Impacts of Global Climate Change**

21 The study of the atmospheric processes involved in global climate change and its potential
22 consequences for human health and global ecosystems is an area of active research. The IPCC
23 Third Assessment Report (TAR) is the most thorough evaluation of current scientific
24 understanding of climate change available at this time. In addition to the first and second IPCC
25 assessments in 1990 and 1995, along with other IPCC reports, earlier assessments included those
26 conducted by the United Nations Environment Program (UNEP, 1986), the World
27 Meteorological Organization (WMO, 1988), the U.S. Environmental Protection Agency (1987),
28 and others (e.g., Patz et al., 2000a,b). The reader is referred to these documents for a complete
29 discussion of climate change science. An abbreviated list of the IPCC conclusions, to date, and a
30 short discussion of the potential impacts of climate change on human health and welfare is

provided here to serve as the context for the discussion of the role of the increasing tropospheric O₃ concentration in climate change.

According to various historic and modern measurement records, atmospheric GHG concentrations have increased dramatically in the past century, and have been attributed to human activities. The IPCC TAR describes the scientific theory and evidence tying increases in GHGs to human activities (IPCC 2001a).

An increasing body of geophysical observations shows that the earth is warming and that other climate changes are underway. These observations include the global surface temperature record assembled since the year 1860, the satellite temperature record begun in 1979, recorded changes in snow and ice cover since the 1950s, sea level measurements taken throughout the 20th century, and sea surface temperature observations recorded since the 1950s. Other evidence includes a marked increase over the past 100 years in the frequency, intensity, and persistence of the zonal atmospheric circulation shifts known as the El Nino-Southern Oscillation (ENSO). ENSO events occur when the tropical Pacific Ocean has accumulated a large, localized mass of warm water that interrupts cold surface currents along South America, altering precipitation and temperature patterns in the tropics, subtropics and the midlatitudes.

IPCC (1998, 2001a) reports also describe the results of general circulation model (GCM) studies that predict that human activities will alter the climate system in a manner that will likely lead to marked global and regional changes in temperature, precipitation and other climate properties. These changes are expected to increase the global mean sea level, as well as increase the number of extreme weather events including floods and droughts, increased wind speeds and precipitation intensity of tropical cyclones, and changes in soil moisture. These predicted changes can be expected to directly impact human health, ecosystems, and global economic sectors, e.g., hydrology and water resources, food and fiber production (IPCC, 1998, 2001b). Table 10-1 summarizes these projected impacts.

Wide variations in the course and net impacts of climate change in different geographic areas are expected. In general, the projected changes will result in additional stresses on natural ecosystems and human societal systems already impacted by increasing resource demands, unsustainable resource management practices, and pollution. Some of the predicted changes include alterations in ecological balances; in the availability of adequate food, water, clean air;

Table 10-1. Examples of Impacts Resulting From Projected Changes in Extreme Climate Events

Projected changes during the 21st Century in Extreme Climate Phenomena and their Likelihood ^a	Representative Examples of Projected Impacts ^b (<i>all high confidence of occurrence in some areas^c</i>)
<i>Simple Extremes</i>	
Higher maximum temperatures; more hot days and heat waves ^d over nearly all land areas (<i>very likely^a</i>)	<ul style="list-style-type: none"> • Increased incidence of death and serious illness in older age groups and urban poor • Increased heat stress in livestock and wildlife • Shift in tourist destinations • Increased risk of damage to a number of crops • Increased electric cooling demand and reduced energy supply reliability
Higher (increasing) minimum temperatures; fewer cold days, frost days, and cold waves ^d over nearly all land areas (<i>very likely^a</i>)	<ul style="list-style-type: none"> • Decreased cold-related human morbidity and mortality • Decreased risk of damage to a number of crops, and increased risk to others • Extended range and activity of some pest and disease vectors • Reduced heating energy demand
More intense precipitation events (<i>very likely^a</i> over many years)	<ul style="list-style-type: none"> • Increased flood, landslide, avalanche, and mudslide damage • Increased soil erosion • Increased flood runoff could increase recharge of some floodplain aquifers • Increased pressure on government and private flood insurance systems and disaster relief
<i>Complex Extremes</i>	
Increased summer drying over most midlatitude continental interiors and associated risk of drought (<i>likely^a</i>)	<ul style="list-style-type: none"> • Decreased crop yields • Increased damage to building foundations caused by ground shrinkage • Decreased water resource quantity and quality • Increased risk of forest fire
Increase in tropical cyclone peak wind intensities, mean and peak precipitation intensities (<i>likely^a</i> over some areas) ^e	<ul style="list-style-type: none"> • Increased risk to human life, risk of infections, disease epidemics, and many other risks • Increased coastal erosion and damage to coastal buildings and infrastructure • Increased damage to coastal ecosystems such as coral reefs and mangroves
Intensified droughts and floods associated with El Niño events in many different regions (<i>likely^a</i>) (see also under droughts and intense precipitation events)	<ul style="list-style-type: none"> • Decreased agricultural and rangeland productivity in drought- and flood-prone regions • Decreased hydro-power potential in drought-prone regions
Increased Asian summer monsoon precipitation variability (<i>likely^a</i>)	<ul style="list-style-type: none"> • Increased flood and drought magnitude and damages in temperate and tropical Asia

Table 10-1 (cont'd). Examples of Impacts Resulting From Projected Changes in Extreme Climate Events

Projected changes during the 21st Century in Extreme Climate Phenomena and their Likelihood ^a	Representative Examples of Projected Impacts ^b (<i>all high confidence of occurrence in some areas^c</i>)
Increased intensity of midlatitude storms (little agreement between current models) ^d	<ul style="list-style-type: none"> • Increased risks to human life and health • Increased property and infrastructure losses • Increased damage to coastal ecosystems

^aLikelihood refers to judgmental estimates of confidence used by TAR WGI: *very likely* (90-99% chance); *likely* (66-90% chance). Unless otherwise stated, information on climate phenomena is taken from the Summary for Policymakers, TAR WGI. TAR WGI = Third Assessment Report of Working Group 1 (IPCC, 2001a).

^bThese impacts can be lessened by appropriate response measures.

^cHigh confidence refers to probabilities between 67 and 95%.

^dInformation from TAR WGI, Technical Summary.

^eChanges in regional distribution of tropical cyclones are possible but have not been established.

Source: IPCC (2001b).

and in human health and safety. Poorer nations can be expected to suffer the most, given their limited adaptive capabilities.

Although many regions are predicted to experience severe, possibly irreversible, adverse effects due to climate change, beneficial changes may also take place. For example, certain regions may benefit from warmer temperatures or increased CO₂ fertilization, e.g., U.S. West Coast coniferous forests, some Western rangelands. Specific benefits may include reduced energy costs for heating, reduced road salting and snow-clearance costs, longer open-water seasons in northern channels and ports, and improved agricultural opportunities in the northern latitudes, as well as the Western interior and coast. For further details about the projected effects of climate change on a U.S.-regional scale, the reader is also referred to several regionally-focused reports (MARAT, 2000; Yarnal et al., 2000; NERAG, 2001; GLRAG, 2000), as well as a report on potential impacts upon human health due to climate change (Bernard et al., 2001). The IPCC report, "The Regional Impacts of Climate Change," (IPCC, 1998) describes the projected effects of human-induced climate change on the different regions of the globe, including Africa, the Arctic and Antarctic, the Middle East and arid Asia, Australasia, Europe, Latin America, North America, the small island nations, temperate Asia, and tropical Asia. While climate models are successful in simulating present annual mean global climate and the

seasonal cycles on continental scales, they have been less successful on regional scales. Clouds and humidity, essential factors in defining local and regional (sub-grid scale) climate, are significantly uncertain (IPCC, 2001a). Due to modeling uncertainties, both in reproducing regional climate and in predicting the future economic activity that will govern future GHG emissions, the projected impacts discussed above are also uncertain.

Findings from the United States Global Change Research Program (USGCRP) (NAST, 2000) report and related reports illustrate the considerable uncertainties and difficulties in projecting likely climate change impacts on regional or local scales. The USGCRP findings also reflect the mixed nature of projected potential climate change impacts (i.e., combinations of mostly deleterious, but other possibly beneficial, effects) for U.S. regions and their variation across different regions. Difficulties in projecting region-specific climate change impacts are complicated by the need to evaluate the potential effects of regional- or local-scale changes in key air pollutants not only on global-scale temperature trends, but also regional- or local-scale temperature and precipitation patterns.

10.3.2 Solar Energy Transformation and the Components of the Earth's Climate System

Mass, in any form, has the capacity to interact with solar and terrestrial radiation, but the manner in which mass interacts with radiation is governed by its particular physical form and/or molecular properties. The most interesting example of radiative properties as a function of physical form is water. In its gas phase, water is the most important GHG present in the climate system, due to its ability to absorb long-wave terrestrial radiation. Conversely, in its frozen form as snow or sea ice, the most important role for water in the climate system is scattering ultraviolet and visible solar radiation back to space, i.e., decreasing the earth's net solar radiation receipts by increasing the earth's reflective properties (albedo). In its liquid aerosol form as clouds, water scatters radiation. In its bulk liquid form as ocean water, it absorbs terrestrial radiation, and represents the earth's most important reservoir of heat energy due to its mass. Molecular properties become important in comparing the radiative absorptive capacity of different gases. In the case of gases, the atomic composition and molecular structure determine the wavelengths of light that a gas molecule can absorb. Ozone and O₂, provide one example of the importance of molecular structure in determining absorption capacity. These molecules are

composed solely of atomic oxygen atoms, but their bond structures are distinct. Ozone, because of its three-atom, bent molecular structure, has the capacity to absorb terrestrial (infrared) wavelengths, making it a GHG. Tropospheric O₃, like stratospheric O₃, also has the capacity to absorb ultraviolet radiation of 320 nm and shorter, increasing the energy-absorbing capacity of the troposphere. Molecular oxygen, O₂, due to its structure, is limited to absorbing very short-wave ultraviolet light – and does so at altitudes too high to influence the climate system significantly.

Each component of the climate system plays a role in absorbing, transforming, storing, dispersing, or scattering solar radiation. Weather is a consequence of the transformation and dispersion of terrestrial radiation. The term, “weather,” refers to the condition of the earth’s atmosphere at a given time and place. It is defined by the air temperature, air pressure, humidity, clouds, precipitation, visibility, and wind speed. The “climate” for a given place on the earth’s surface is a long-term average of these elements of weather, accounting for daily and seasonal weather events. The frequency of extreme weather events is used to distinguish among climates that have similar averages (Ahrens, 1994).

The earth’s capacity for retaining heat is increased by the transformation of its oxygen, for example, into O₃ by way of air pollution chemistry. Climate components, including GHGs, land, oceans, sea ice, land ice and snow, atmospheric particles, vegetation, clouds, etc., all contribute to the earth’s heat capacity with respect to solar energy. Changes in the properties (or mass) of these components will “force” the climate system in one direction or the other, i.e., warmer versus cooler. Climate forcing is further described, below.

10.3.3 The Composition of the Atmosphere and the Earth’s Radiative Equilibrium

The Greenhouse Effect is the term given to the decreased rate of reemission of absorbed solar energy due to the heat-retaining properties of the Earth’s atmosphere. According to simple radiative transfer theory, at thermal equilibrium, the earth’s temperature should be near – 15 °C. This is the temperature of a theoretical “black body” that is receiving and then reemitting 342.5 Wm⁻², i.e., the globally-averaged amount of full-spectrum solar energy absorbed, and then reemitted by the earth as infrared terrestrial radiation, per square meter. In fact, satellite observations well above the atmosphere indicate that the earth’s average *planetary* temperature

1 is remarkably close to its theoretical black body value at -18°C , a temperature at which liquid
2 water ordinarily does not exist.

3 At its *surface*, however, the earth's average temperature is $+15^{\circ}\text{C}$. The $+33^{\circ}\text{C}$
4 temperature differential between the Earth's planetary and surface temperatures is due to the
5 presence of infrared radiation-absorbing components in the atmosphere, such as water vapor,
6 CO_2 , CH_4 , several other trace gases, and some types of particles and clouds.

7 The atmosphere, when cloud free, is largely transparent in the solar wavelength range.
8 A small fraction of this radiation is absorbed and reemitted as black body radiation by dark
9 atmospheric particles (IPCC, 2001a). The majority of clouds and particles offset, in part, the
10 greenhouse effect by increasing the earth's albedo and, therefore, decreasing the overall amount
11 of solar radiation absorbed by the earth system.

12 Ozone, SO_2 and NO_2 also absorb ultraviolet and near ultraviolet wavelengths, in addition to
13 infrared wavelengths. Once absorbed by a gas molecule, the energy introduced by a photon may
14 induce a photochemical reaction with the residual energy thermally exciting (heating) the
15 products of the reaction. Alternatively, the energy introduced into the molecule by the photon
16 may be dispersed amongst neighboring molecules via intermolecular collisions, or reemitted in
17 part as a lower energy (IR) photon.

18 Radiation from the sun or the earth's surface that is absorbed by gases and particles is
19 reemitted isotropically, i.e., it is equally likely to be emitted in all directions. Therefore, to a
20 first approximation, half of the radiation trapped by the earth's atmosphere is reflected back to
21 its surface. A portion of this radiation is transformed into the heat energy that drives the
22 atmospheric processes that form the basis of weather and climate. Radiation which is not
23 absorbed by gases and aerosols reaches the earth's surface where it is scattered (reflected) or
24 absorbed, depending on the surface albedo.

25 Successful modeling of the earth's climate and, therefore, the assessment of the extent of
26 human-induced climate change and development of appropriate policy depends on the quality of
27 available information on the relative efficiencies, amounts, and spatial and temporal distributions
28 of the various radiatively active components of the atmosphere that absorb and/or reflect solar
29 and terrestrial radiation, along with all the other non-atmospheric components of the earth
30 system.

10.3.3.1 Forcing of the Earth's Radiative Balance

As mentioned earlier, the commonly used measure of the relative influence of a given component of the climate system on the earth's radiative balance is its radiative forcing (IPCC, 2001a; Houghton et al., 1990). Radiative forcing, in Wm^{-2} , is a quantity that was developed by the climate modeling community as a first order-only means of estimating relative effects of individual anthropogenic and natural processes on the energy balance within the climate system. Discussions within the climate community are underway regarding a metric to replace forcing as the standard measure of climate impact – one that will account for more of the factors that determine the effectiveness of a given atmospheric component in this capacity. However, forcing remains the current standard (National Research Council, 2005).

When the effect of a particular component of the climate system is to reduce the amount of solar energy absorbed, usually by increasing the earth's albedo, this component is said to provide a “negative” forcing, measured in Wm^{-2} . The convention assigns a positive value to the forcing induced by climate system components that enhance the greenhouse effect, or otherwise act to increase the heat absorbing capacity of the earth system. Purely reflective atmospheric aerosol, clouds, white roof tops, snow-covered land surfaces, and dense sea ice provide a negative forcing, while highly absorbing dark-colored atmospheric aerosols, GHGs, and increases, due to the melting of sea ice sheets, in dark ocean surfaces positively force the climate system.

Global and regional climate are roughly defined by the balance between the large number of positive and negative forcings induced by the many different components of the earth system. However, the earth system responds to these forcings in a complex way, due to feedback mechanisms that are theorized but very difficult to resolve at the quantitative level.

A simple example would be the positive feedback associated with melting sea ice. As sea ice melts with increasing surface temperatures, the dark ocean surface is revealed which is more efficient at absorbing infrared radiation, further increasing the rate of warming. The role of feedbacks in determining the sensitivity of climate to changes in radiative forcing is described in detail in the IPCC TAR (IPCC, 2001a). In the absence of complete, quantitative, information about climate feedbacks the use of radiative forcing values for the many components of the climate system remains the primary method for estimating their relative importance in climate change.

The IPCC has reported estimated values for forcing by individual radiatively active gases, and particle-phase components of the atmosphere, derived primarily through expert judgment incorporating the results of peer-reviewed modeling studies. The forcing estimates, shown in Figure 10-5, are global averages attributed to known greenhouse gases, including O₃; particles; anthropogenic cirrus clouds; land-use change; and natural solar flux variations. Uncertainty ranges are assigned to reflect the range of modeled values reported in those studies. The current estimate of forcing due to long-lived, well-mixed, greenhouse gases accumulated in the atmosphere from the preindustrial era (ca., 1750) through the year 2000 is + 2.4 Wm⁻² ± 10% (IPCC, 2001a). An indication of the level of confidence in each of these estimates is given along the bottom of this figure, again reflecting the expert judgment of the IPCC.




Figure 10-5. Estimated global mean radiative forcing exerted by gas and various particle phase species for the year 2000, relative to 1750.

Source: IPCC (2001a).

1 The IPCC reported a global average forcing value of $0.35 \pm 0.15 \text{ Wm}^{-2}$ for tropospheric O_3 ,
2 based on model calculations constrained by climatological observations. The considerations and
3 studies used to estimate this value will be outlined below.

5 **10.3.4 Factors Affecting the Magnitude of Climate Forcing by Ozone**

6 Tropospheric O_3 is estimated to have provided the third largest increase in direct forcing
7 since pre-industrial times. It may also play a role in indirect forcing through its participation in
8 the oxidative removal of other radiatively active trace gases, such as CH_4 and the HCFCs. Given
9 its relatively short atmospheric lifetime, the distribution of tropospheric O_3 is highly variable in
10 geographic extent and time. The direct and indirect forcing it imposes on the climate system,
11 therefore, depends upon its geospatial and temporal distribution, but also depends upon the
12 albedo of the underlying surface. These several variables introduce substantial uncertainty into
13 tropospheric ozone forcing estimates.

15 **10.3.4.1 Global versus Regional Atmospheric Ozone Concentrations**

16 Ozone reacts photochemically at time-scales generally shorter than those for large-scale
17 mixing processes in the atmosphere. Concentrated O_3 plumes evolve downwind of strong
18 sources of its precursor pollutants: reactive nitrogen, CO, and non-methane hydrocarbons
19 (NMHCs). The most important of these sources are midlatitude industrialized areas and tropical
20 biomass burning. When viewed from above the atmosphere by satellite-borne spectrometers, O_3
21 enhancements appear as relatively localized air masses or regional-scale plumes, usually
22 originating from industrialized areas or areas in which active biomass burning is underway. The
23 IPCC (2001a) describes the efforts of several research teams who have analyzed data supplied by
24 the satellite-borne Total Ozone Mapping Spectrometer (TOMS) and other remote-sensing
25 instruments to map the global distribution of tropospheric O_3 and to attempt to identify processes
26 that influence the global tropospheric O_3 budget (IPCC, 2001a). More recently, coincident
27 observations of total O_3 by TOMS and the Solar Backscattered UV (SBUV) instrument were
28 used by Fishman et al. (2003) to construct well-resolved spatial and temporal maps of the
29 regional distribution of tropospheric O_3 . Their results were consistent with those reported by
30 others, but with higher regional-scale resolution. They reported large O_3 enhancements in the
31 southern tropics in austral Spring, and in the northern temperate latitudes in the summer. The

regional nature of high O₃ concentrations was clearly visible in northeastern India, the eastern United States, eastern China, and west and southern Africa, each coincident with high population densities. Fishman et al. (2003) noted, as have the other groups cited above, significant interannual variability in the concentrations observed over these regions. *In situ* measurements of tropospheric O₃ concentrations range from 10 ppb over remote oceans, to 100 ppb in both the upper troposphere and in plumes downwind from polluted metropolitan regions (IPCC, 2001a). Ground-level concentrations in urban areas are often higher than 100 ppb. In the southern hemisphere, one of the largest sources of O₃ precursors is biomass burning. Biomass burning elevates O₃ on large spatial scales, particularly in the tropical Atlantic west of the coast of Africa and in Indonesia.

Current estimates place the global burden of tropospheric O₃ at a highly uncertain 370 Tg, equivalent to an average column density of 34 Dobson Units (1 DU = 2.687×10^{16} molecules/cm⁻²) or a mean concentration of 50 ppb (IPCC, 2001a). Accounting for differences in levels of industrialization between the hemispheres, the average column burden in the Northern Hemisphere is estimated to be 36 DU, with the Southern Hemisphere estimated to average 32 DU. Due to its rapid photochemistry, individual surface measurements of tropospheric O₃ cannot capture large scale concentrations, nor will it represent the concentration at higher altitudes. Dense surface and vertical measurements (ozonesondes) would be required to supplement available output from remote sensing instruments to provide the complete set of observations needed to derive a credible global O₃ budget. Such a measurement program appears, at present, to be impractical.

Little historical data exists that might be used to estimate the global ozone burden prior to industrialization. Although a few late 19th century measurements suggest that O₃ has more than doubled in Europe during the 20th century. The insufficient data record on pre-industrial tropospheric O₃ distributions introduces a major uncertainty in the estimation of the ozone-induced forcing (IPCC, 2001a).

10.3.4.2 Global Versus Regional Atmospheric Ozone Trends

For the northern hemisphere, weekly continuous data since 1970 are available from only nine stations in the latitude range 36°N to 59°N (IPCC, 2001a). Available tropospheric O₃ measurements do not reveal a clear trend in concentration, while trends in the stratosphere are

more readily identified. Different trends are seen at different locations for different periods, consistent with regional changes in pollutant emission, especially NO_x. Logan et al. (1999) analyzed the composite record of mid-tropospheric ozone abundance from the nine station network. A plot of data is shown in Figure 10-6. While no clear trend appeared for 1980 through 1996, the second half of this record (about 57 ppb) is clearly greater than the first half (about 53 ppb). The trend may be consistent with changes in regional NO_x emission rates due to pollution reduction efforts in developed countries and increasing emissions in rapidly growing economies in Asia.




Figure 10-6. Mid-tropospheric O₃ abundance (ppb) in northern midlatitudes (36 °N-59 °N) for the years 1970 to 1996. Observations between 630 and 400 hPa are averaged from nine ozone sonde stations (four in North America, three in Europe, two in Japan), following the data analysis of Logan et al. (1999). Values are derived from the residuals of the trend fit with the trend added back to allow for discontinuities in the instruments. Monthly data (points) are shown with a smoothed 12-month-running mean (line).

Source: IPCC (2001).

It must be noted that the measurements shown in Figure 10-6 are for surface concentrations, only. Many fewer locations have measured changes in the concentrations of O₃ as a function of altitude. Fewer still are locations that have collected and maintained data

records prior to 1970. The absence of historical data on the vertical distribution of O₃ adds to the difficulty in estimating historical atmospheric burdens and trends in O₃-related climate forcing.

The IPCC (2001a) surveyed the results of published chemistry transport model (CTM) modelling studies, listed in Table 10-2, that estimated the global average increase since the pre-industrial era in total column O₃. Model estimates ranged from +7 to +12 DU. On the basis of these estimates, available measurements and other analyses, the IPCC estimated that total column O₃ has increased by 9 DU, with a 67% confidence range of +6 to +13 DU. In some of the modelling studies, emissions scenarios were used that predict a further increase in column O₃ due to growing emissions of O₃ precursors.

Table 10-2. CTM Studies Assessed by the IPCC for its Estimate of the Change in Global, Total Column O₃ Since the Pre-Industrial Era

Estimated Change in Column Ozone in DU	Model Used	Authors (Publication Date)
7.9	GFDL	Haywood et al. (1998)
8.9	MOZART-1	Hauglustaine et al. (1998)
8.4	NCAR/2D	Kiehl et al. (1999)
9.5	GFDL-scaled	Levy et al. (1997)
12	Harvard/GISS	Mickley et al. (1999)
7.2	ECHAM4,	Roelofs et al. (1997)
8.7	UKMO,	Stevenson et al. (2000)
9.6	UIO	Berntsen et al. (1999)
8	MOGUNTIA	VanDorland et al. (1997)

Source: IPCC (2001).

Chapters 2 and 3 of this document provide an expanded discussion of the issues associated with determining the global tropospheric O₃ background.

10.3.4.3 The Sensitivity of Ozone-related Forcing Surface to Albedo

The characteristics of the surface underlying an O₃ enhancement play a large role in the O₃ forcing effect. Highly reflective, warm surfaces, such as light-colored deserts, scatter solar ultraviolet (UV) radiation; absorb, and then emit infrared terrestrial radiation. Both forms of radiation can be trapped, transformed, and/or reemitted back to the surface by tropospheric O₃. Highly reflective, cold, surfaces will scatter more radiation while emitting less terrestrial infrared radiation. Dark, warm, surfaces such as tropical ocean hot spots predominantly emit terrestrial radiation. Ozone will trap heat at differing efficiencies as a consequence of the amount and type of radiation reflected or emitted from the surface underneath it. Studies by two groups, Hauglustaine et al. (1998) and Mickley et al. (1999), have shown that industrial pollution that has been transported to the Arctic induces a high, regional O₃-related forcing due to the highly reflective underlying ice and snow surface.

10.3.5 Estimated Forcing by Tropospheric Ozone

10.3.5.1 Direct Climate Forcing Due to Ozone

The inhomogeneous distribution of O₃ within the troposphere coupled with the large uncertainty in the global O₃ budget significantly complicates the matter of estimating the global average direct forcing due to O₃. The IPCC TAR (2001a) lists the results of several modeling studies that estimated the annual change in the relative forcing by O₃ since pre-industrial times. It was noted that the differences amongst the estimates were most likely due to differences in predicted O₃ chemistry, including the emissions inventories used, and the chemical process and transport mechanisms incorporated into the models, rather than by factors relating to radiative transfer. The IPCC intercomparison of the models and their results indicated that the uncertainties in estimated forcings due to O₃ have reduced since the IPPC Second Assessment Report (1996)

The O₃-related forcings estimated by studies considered by the IPCC (2001a) are listed in Table 10-3. Ten of the listed estimates are based on global chemistry/transport models calculations. One study was constrained by a climatology derived from observations. Given the differences in calculated total column O₃ amongst the models, a normalized forcing (Wm⁻² per Dobson Unit of tropospheric O₃ change) is listed in addition to the absolute forcing (Wm⁻²)

Table 10-3. Tropospheric O₃ Change (ΔO_3) in Dobson Units (DU) Since Pre-industrial Times, and the Accompanying net (SW plus LW) Radiative Forcings (Wm^{-2}), After Accounting for Stratospheric Temperature Adjustment (using the Fixed Dynamical Heating method). Estimates are Taken From the Published Literature. Normalized Forcings (norm.) Refer to Radiative Forcing per O₃ Change (Wm^{-2} per DU)

Estimated Global Average Forcing Due to Tropospheric Ozone					
Reference	Clear sky conditions			Total sky conditions	
	ΔO_3	Net	Net (norm.)	Net	Net (norm.)
Berntsen et al. (1997) – [Reading model]	7.600	0.310	0.041	0.280	0.037
Stevenson et al. (1998)	8.700	0.391	0.045	0.289	0.033
Berntsen et al. (1997) – [Oslo model]	7.600	0.390	0.051	0.310	0.041
Haywood et al. (1998a)	7.900	0.380	0.048	0.310	0.039
Kiehl et al. (1999)	8.400	0.379	0.045	0.320	0.038
Berntsen et al. (2000)	9.600	0.428	0.045	0.342	0.036
Brasseur et al. (1998)	—	—	—	0.370	—
van Dorland et al. (1997)	8.070	0.443	0.055	0.380	0.047
Roelofs et al. (1997)	7.200	0.397	0.055	0.404	0.056
Lelieveld and Dentener (2000)	—	—	—	0.420	—
Hauglustaine et al. (1998)	8.940	0.511	0.057	0.426	0.048
Mean	8.224	0.403	0.049	0.343	0.042

Source: IPCC (2001).

estimated by each model. Both clear sky (cloud-free) and total sky (including clouds) forcing estimates are listed.

The largest O₃-related forcings coincide with strongest sources of tropospheric ozone, the models predict that occur in the northern midlatitude regions (40° to 50° N), reaching as much as 1 Wm^{-2} in the summer and in the tropics, related to biomass burning. In general, the estimates are comparable in magnitude and show similarity in geographic distribution. For total sky conditions, the range in globally and annual averaged tropospheric O₃ forcing from all of these models is from 0.28 to 0.43 Wm^{-2} , while the normalized forcing is 0.033 to 0.056 Wm^{-2} per DU. As expected, they are opposite in sign to the forcing estimated for sulphate aerosols, which

scatter radiation. The range in normalized forcings emphasizes the differences in assumptions used by the different models. The tropospheric O₃ forcing constrained by the observational climatology is 0.32 Wm⁻² for globally averaged, total sky conditions. As shown in Figure 10-1, the IPCC (2001) concluded that 0.35 ± 0.15 Wm⁻² represents the most likely value for annually and globally-averaged forcing by tropospheric O₃.

10.3.5.2 Indirect Forcing Due to Ozone

Ozone has an indirect climate forcing effect due to its role in the oxidative removal of other reactive GHGs, including CH₄, hydrofluorocarbons (HFCs) and other reactive non-methane hydrocarbons (NMHCs). The primary actor in this effect is a second generation product of the photolysis of O₃, the hydroxyl radical (OH). Hydroxyl radical is produced by way of a pair of reactions that start with the photodissociation of O₃ by solar UV.



Reactions with OH are the primary removal mechanism for CH₄ and NMHCs as well as the pollutants NO_x and CO. Methane and CO have especially high abundances in the global atmosphere. OH is estimated to react with these two gases within one second of its formation. In addition to CH₄, NO_x, CO and the NMHCs, OH concentrations are controlled by local concentrations of H₂O (humidity) and the intensity of solar UV. Different atmospheric concentrations of the required precursors suggest that pre-industrial OH concentrations are likely to have been different from present-day concentrations, but there is no consensus on the magnitude of this difference. Observations of global atmospheric concentrations of chloroform (CH₃CCl₃), a well-mixed tropospheric species that reacts with OH, have been used to estimate OH abundances. Independent studies have shown overlapping trends for the period 1978 to 1994, but none outside the given uncertainty ranges (0.5 ± 0.6%/yr) (Prinn et al., 1995; Krol et al., 1998). The IPCC (2001) reported a range of +5% to -20% for predicted changes in global OH abundances.

Given the difficulty in estimating global OH abundances in the past, present and future, estimates of indirect forcing due to O₃ have been difficult to obtain and are highly uncertain. Attempts have been made to account for OH feedbacks in estimating the global warming potential of CH₄.

10.3.5.3 Predictions for Future Climate Forcing by Anthropogenic Ozone

The rate of increase in surface O₃ in Europe and North America, since 1980, appears to be slowing, likely due to control measures intended to improve urban air quality. Not surprisingly, CTM modeling attempts to predict future precursor emissions and resulting O₃ abundances indicate that the largest future O₃-related forcings will be related to population growth and economic development in Asia (van Dorland et al., 1997; Brasseur et al., 1998). The results of these modeling studies suggested that a higher globally averaged total radiative forcing due to O₃ from pre-industrial times to 2050 of 0.66 Wm⁻² and 0.63 Wm⁻². Chalita et al. (1996) predicted a globally averaged radiative forcing from pre-industrial times to 2050 of 0.43 Wm⁻². Stevenson et al. (1998) predicted an O₃-related forcing of 0.48 Wm⁻² in 2100. All of these predictions must be viewed with much caution given the considerable uncertainties associated with such estimates.

10.3.6 Conclusion

The general consensus within the atmospheric sciences community, as represented by the United Nations Intergovernmental Panel on Climate Change (IPCC), is that human activities have a discernable effect on the earth's climate. However, quantifying the extent of human-induced forcing on climate requires detailed information about human-induced change on the components of the earth system that govern climate. Troposphere ozone is a well-known GHG, but information regarding its historical trends in concentration, its current and future atmospheric burden, and other critical details needed for estimating its direct and indirect forcing effects on the climate system are highly uncertain.

The IPCC has estimated that the globally averaged forcing due to O₃ is approximately 0.35 ± 0.15 Wm⁻². The role of O₃ in climate is likely to be much more pronounced adjacent to the sources of its chemical precursors, consistent with satellite observations of high regional scale column densities near large urban areas and large-scale biomass burning activity.

1 Modeling studies evaluated by the IPCC have estimated that regional scale forcing due to O₃ can
2 approach 1 Wm⁻², or two-thirds the forcing estimated for CO₂. However, more reliable
3 estimates of the overall importance of forcing due to tropospheric O₃ await further advances in
4 monitoring and chemical transport modeling.
5

REFERENCES

- Ahrens, D. C. (1994) *Meteorology today: an introduction to weather, climate and the environment*. Minneapolis, MN: West Publishing Co., p. 17.
- Airey, D. K.; Wong, J. C.; Fleming, R. A.; Meldrum, L. R. (1997) An estimate of the UV-B exposure for outdoor workers during a south-east Queensland summer. *Health Phys.* 72: 544-549.
- Ancellet, G.; Beekmann, M. (1997) Evidence for changes in the ozone concentrations in the free troposphere over southern France from 1976 to 1995. *Atmos. Environ.* 31: 2835-2851.
- Autier, P.; Doré, J. F.; Négrier, S.; Liénard, D.; Panizzon, R.; Lejeune, F. J.; Guggisberg, D.; Eggermont, A. M. (1999) Sunscreen use and duration of sun exposure: a double-blind, randomized trial. *J. Natl. Cancer Inst.* 91: 1304-1309.
- Autier, P.; Doré, J.-F.; Reis, A. C.; Grivegnée, A.; Ollivaud, L.; Truchetet, F.; Chamoun, E.; Rotmensz, N.; Severi, G.; Césarini, J. P.; EORTC Melanoma Co-operative Group. (2000) Sunscreen use and intentional exposure to ultraviolet A and B radiation: a double blind randomized trial using personal dosimeters. *Br. J. Cancer* 83: 1243-1248.
- Ayala, M. N.; Michael, R.; Soderberg, P. G. (2000) Influence of exposure time for UV radiation-induced cataract. *Invest. Ophthalmol. Visual Sci.* 41: 3539-3543.
- Barnard, W. F.; Saxena, V. K.; Wenny, B. N.; DeLuisi, J. J. (2003) Daily surface UV exposure and its relationship to surface pollutant measurements. *J. Air Waste Manage. Assoc.* 53: 237-245.
- Bellows, C. F.; Belafsky, P.; Fortgang, I. S.; Beech, D. J. (2001) Melanoma in African-Americans: trends in biological behavior and clinical characteristics over two decades. *J. Surg. Oncol.* 78: 10-16.
- Berntsen, T. K.; Isaksen, I. S. A. (1999) Effects of lightning and convection on changes in tropospheric ozone due to NO_x emissions from aircraft. *Tellus* 51B: 766-788.
- Berntsen, T. K.; Isaksen, I. S. A.; Myhre, G.; Fuglestad, J. S.; Stordal, F.; Larsen, T. A.; Freckleton, R. S.; Shine, K. P. (1997) Effects of anthropogenic emissions on tropospheric ozone and its radiative forcing. *J. Geophys. Res. [Atmos.]* 102: 28,101-28,126.
- Berntsen, T. K.; Myhre, G.; Stordal, F.; Isaksen, I. S. A. (2000) Time evolution of tropospheric ozone and its radiative forcing. *J. Geophys. Res. [Atmos.]* 105: 8915-8930.
- Berret, J.; Liardet, S.; Scaletta, C.; Panizzon, R.; Hohlfeld, P.; Applegate, L. A. (2002) Use of sunscreens in families living in Switzerland. *Dermatology* 204: 202-208.
- Black, W. C.; Goldhahn, R. T., Jr.; Wiggins, C. (1987) Melanoma within a southwestern Hispanic population. *Arch. Dermatol.* 123: 1331-1334.
- Brasseur, G. P.; Kiehl, J. T.; Müller, J.-F.; Schneider, T.; Granier, C.; Tie, X.-X.; Hauglustaine, D. (1998) Past and future changes in global tropospheric ozone: impact on radiative forcing. *Geophys. Res. Lett.* 25: 3807-3810.
- Breuer-McHam, J.; Simpson, E.; Dougherty, I.; Bonkobara, M.; Ariizumi, K.; Lewis, D. E.; Dawson, D. B.; Duvic, M.; Cruz, P. D., Jr. (2001) Activation of HIV in human skin by ultraviolet B radiation and its inhibition by NFκB blocking agents. *Photochem. Photobiol.* 74: 805-810.
- Brühl, C.; Crutzen, P. J. (1988) On the disproportionate role of tropospheric ozone as a filter against solar UV-B radiation. *Geophys. Res. Lett.* 16: 703-706.
- Chalita, S.; Hauglustaine, D. A.; Le Treut, H.; Müller, J. F.; Penkett, S., eds. (1996) Radiative forcing due to increased tropospheric ozone concentrations. *Atmos. Environ.* 30: 1641-1646.
- Clydesdale, G. J.; Dandie, G. W.; Muller, H. K. (2001) Ultraviolet light induced injury: immunological and inflammatory effects. *Immunol. Cell Biol.* 79: 547-568.
- Cockell, C. S. (2001) A photobiological history of Earth. In: Cockell, C. S.; Blaustein, A. R., eds. *UV radiation flux into ecosystems*. New York, NY: Springer-Verlag.
- Cutchis, P. (1974) Stratospheric ozone depletion and solar ultraviolet radiation on Earth. *Science (Washington, DC)* 184: 13-19.
- Davis, K. J.; Cokkinides, V. E.; Weinstock, M. A.; O'Connell, M. C.; Wingo, P. A. (2002) Summer sunburn and sun exposure among US youths ages 11 to 18: national prevalence and associated factors. *Pediatrics* 110: 27-35.
- De Fabo, E. C.; Noonan, F. P.; Fears, T.; Merlino, G. (2004) Ultraviolet B but not ultraviolet A radiation initiates melanoma. *Cancer Res.* 64: 6372-6376.
- De Gruijl, F. R. (1995) Action spectrum for photocarcinogenesis. *Recent Results Cancer Res.* 139: 21-30.
- De Gruijl, F. R. (1999) Skin cancer and solar UV radiation. *Eur. J. Cancer* 35: 2003-2009.
- De Gruijl, F. R. (2002) Photocarcinogenesis: UVA vs. UVB radiation. *Skin Pharmacol. Appl. Skin Physiol.* 15: 316-320.

- 1 De Gruijl, F. R.; Henricus, J. C. M.; Sterenborg, H.; Forbes, P. D.; Davies, R. E.; Cole, C.; Kelfkens, G.;
2 Van Weelden, H.; Slaper, H.; Van der Leun, J. C. (1993) Wavelength dependence of skin cancer induction by
3 ultraviolet irradiation of albino hairless mice. *Cancer Res.* 53: 53-60.
- 4 Diepgen, T. L.; Mahler, V. (2002) The epidemiology of skin cancer. *Br. J. Dermatol.* 146(suppl. 61): 1-6.
- 5 Diffey, B. L. (1990) Human exposure to ultraviolet radiation. *Semin. Dermatol.* 9: 2-10.
- 6 Diffey, B. L. (1994) Observed and predicted minimal erythema doses: a comparative study. *Photochem. Photobiol.*
7 60: 380-382.
- 8 Diffey, B. L.; Gibson, C. J.; Haylock, R.; McKinlay, A. F. (1996) Outdoor ultraviolet exposure of children and
9 adolescents. *Br. J. Dermatol.* 134: 1030-1034.
- 10 Evans, R. D.; Kopf, A. W.; Lew, R. A.; Rigel, D. S.; Bart, R. S.; Friedman, R. J.; Rivers, J. K. (1988) Risk factors
11 for the development of malignant melanoma--I: review of case-control studies. *J. Dermatol. Surg. Oncol.*
12 14: 393-408.
- 13 Evans, W. F. J.; Ferguson, C. R.; Puckrin, E. (2004) Radiative forcing by tropospheric ozone. *Bull. Am. Meteorol.*
14 *Soc.* 85: 347.
- 15 Fears, T. R.; Bird, C. C.; Guerry, D., IV; Sagebiel, R. W.; Gail, M. H.; Elder, D. E.; Halpern, A.; Holly, E. A.;
16 Hartge, P.; Tucker M. A. (2002) Average midrange ultraviolet radiation flux and time outdoors predict
17 melanoma risk. *Cancer Res.* 62: 3992-3996.
- 18 Fiore, A. M.; Jacob, D. J.; Field, B. D.; Streets, D. G.; Fernandes, S. D.; Jang, C. (2002) Linking ozone pollution and
19 climate change: the case for controlling methane. *Geophys. Res. Lett.* 29(19): 10.1029/2002GL015601.
- 20 Fishman, J.; Wozniak, A. E.; Creilson, J. K. (2003) Global distribution of tropospheric ozone from satellite
21 measurements using the empirically corrected tropospheric ozone residual technique: identification of the
22 regional aspects of air pollution. *Atmos. Chem. Phys.* 3: 893-907.
- 23 Fligge, M.; Solanki, S. K. (2000) Reconstruction of past solar irradiance. *Space Sci. Rev.* 94: 127-138.
- 24 Frederick, J. E.; Koob, A. E.; Weatherhead, E. C. (1993) Empirical studies of tropospheric transmission in the
25 ultraviolet: roadband measurements. *J. Appl. Meteorol.* 32: 1883-1892.
- 26 Freedman, D. M.; Dosemeci, M.; McGlynn, K. (2002) Sunlight and mortality from breast, ovarian, colon, prostate,
27 and non-melanoma skin cancer: a composite death certificate based case-control study. *Occup. Environ. Med.*
28 59: 257-262.
- 29 Gallagher, R. P.; Hill, G. B.; Bajdik, C. D.; Coldman, A. J.; Fincham, S.; McLean, D. I.; Threlfall, W. J. (1995a)
30 Sunlight exposure, pigmentation factors and risk of nonmelanocytic skin cancer. II. Squamous cell carcinoma.
31 *Arch. Dermatol.* 131: 164-169.
- 32 Gallagher, R. P.; Hill, G. B.; Bajdik, C. D.; Fincham, S.; Coldman, A. J.; McLean, D. I.; Threlfall, W. J. (1995b)
33 Sunlight exposure, pigmentary factors, and risk of nonmelanocytic skin cancer. I. Basal cell carcinoma.
34 *Arch. Dermatol.* 131: 157-163.
- 35 Garland, F. C.; Garland, C. F.; Gorham, E. D.; Young, J. F. (1990) Geographic variation in breast cancer mortality in
36 the United States: a hypothesis involving exposure to solar radiation. *Prev. Med.* 19: 614-622.
- 37 Garsen, J.; Van Loveren, H. (2001) Effects of ultraviolet exposure on the immune system. *Crit. Rev. Immunol.*
38 21: 359-397.
- 39 Gauss, M.; Myhre, G.; Pitari, G.; Prather, M. J.; Isaksen, I. S. A.; Berntsen, T. K.; Brasseur, G. P.; Dentener, F. J.;
40 Derwent, R. G.; Hauglustaine, D. A.; Horowitz, L. W.; Jacob, D. J.; Johnson, M.; Law, K. S.; Mickley, L. J.;
41 Müller, J.-F.; Plantevin, P. H.; Pyle, J. A.; Rogers, H. L.; Stevenson, D. S.; Sundet, J. K.; Van Weele, M.;
42 Wild, O. (2003) Radiative forcing in the 21st century due to ozone changes in the troposphere and the lower
43 stratosphere. *J. Geophys. Res. (Atmos.)* 108(D9): 10.1029/2002JD002624.
- 44 Gies, P.; Wright, J. (2003) Measured solar ultraviolet radiation exposures of outdoor workers in Queensland in the
45 building and construction industry. *Photochem. Photobiol.* 68: 78-83.
- 46 Gies, P.; Roy, C.; Toomey, S.; MacLennan, R.; Watson, M. (1998) Solar UVR exposures of primary school children
47 at three locations in Queensland. *Photochem. Photobiol.* 68: 78-83.
- 48 Gloster, H. M., Jr.; Brodland, D. G. (1996) The epidemiology of skin cancer. *Dermatol. Surg.* 22: 217-226.
- 49 Godar, D. E. (2001) UV doses of American children and adolescents. *Photochem. Photobiol.* 74: 787-793.
- 50 Godar, D. E.; Wengraitis, S. P.; Shreffler, J.; Sliney, D. H. (2001) UV doses of Americans. *Photochem. Photobiol.*
51 73: 621-629.
- 52 Godar, D. E.; Urbach, F.; Gasparro, F. P.; Van Der Leun, J. C. (2003) UV doses of young adults. *Photochem.*
53 *Photobiol.* 77: 453-457.
- 54 Gorham, E. D.; Garland, F. C.; Garland, C. F. (1990) Sunlight and breast cancer incidence in the USSR. *Int. J.*
55 *Epidemiol.* 19: 820-824.
- 56 Granstein, R. D.; Matsui, M. S. (2004) UV radiation-induced immunosuppression and skin cancer. *Cutis*
57 74(suppl. 5): 4-9.

- 1 Grant, W. B. (2002a) An estimate of premature cancer mortality in the U.S. due to inadequate doses of solar
2 ultraviolet-B radiation. *Cancer* 94: 1867-1875.
- 3 Grant, W. B. (2002b) An ecologic study of dietary and solar ultraviolet-B links to breast carcinoma mortality rates.
4 *Cancer* 94: 272-281.
- 5 Great Lakes Regional Assessment Group (GLRAG). (2000) Preparing for a changing climate: the potential
6 consequences of climate variability and change. Great Lakes overview. Washington, DC: U.S. Environmental
7 Protection Agency; Office of Research and Development; U.S. Global Change Research Program (USGCRP)
8 and Ann Arbor, MI: University of Michigan; Atmospheric, Oceanic and Space Sciences Department.
9 Available: <http://www.geo.msu.edu/glra/assessment/assessment.html> [17 April 2002].
- 10 Green, A.; Battistutta, D.; Hart, V.; Leslie, D.; Weedon, D. (1996) Skin cancer in a subtropical Australian
11 population: incidence and lack of association with occupation. The Nambour Study Group. *Am. J. Epidemiol.*
12 144: 1034-1040.
- 13 Hall, H. I.; Miller, D. R.; Rogers, J. D.; Bewerse, B. J. (1999) Update on the incidence and mortality from melanoma
14 in the United States. *J. Am. Acad. Dermatol.* 40: 35-42.
- 15 Halliday, G. M.; Byrne, S. N.; Kuchel, J. M.; Poon, T. S.; Barnetson, R. S. (2004) The suppression of immunity by
16 ultraviolet radiation: UVA, nitric oxide and DNA damage. *Photochem. Photobiol. Sci.* 3: 736-740.
- 17 Hanchette, C. L.; Schwartz, G. G. (1992) Geographic patterns of prostate cancer mortality. *Cancer* 70: 2861-2869.
- 18 Harrison, G. I.; Young, A. R. (2002) Ultraviolet radiation-induced erythema in human skin. *Methods* (San Diego,
19 CA, U.S.) 28: 14-19.
- 20 Hartge, P.; Devesa, S. S.; Grauman, D.; Fears, T. R.; Fraumeni, J. F., Jr. (1996) Non-Hodgkin's lymphoma and
21 sunlight. *J. Natl. Cancer Inst.* 88: 298-300.
- 22 Hartmann, D. L. (1994) Global physical climatology. New York, NY: Academic Press. (International geophysics:
23 v. 56).
- 24 Hauglustaine, D. A.; Brasseur, G. P.; Walters, S.; Rasch, P. J.; Müller, J.-F.; Emmons, L. K.; Carroll, M. A. (1998)
25 MOZART, a global chemical transport model for ozone and related chemical tracers 2. model results and
26 evaluation. *J. Geophys. Res. (Atmos.)* 103: 28291-28335.
- 27 Haywood, J. M.; Schwarzkopf, M. D.; Ramaswamy, V. (1998) Estimates of radiative forcing due to modeled
28 increases in tropospheric ozone. *J. Geophys. Res. [Atmos.]* 103: 16,999-17,007.
- 29 Henriksen, M.; Na, R.; Agren, M. S.; Wulf, H. C. (2004) Minimal erythema dose after umltiple UV exposures
30 depends on pre-exposure skin pigmentation. *Photodermatol. Photoimmunol. Photomed.* 20: 163-169.
- 31 Hildesheim, J.; Fornace, A. J., Jr. (2004) The dark side of light: the damaging effects of UV rays and the protective
32 efforts of MAP kinase signaling in the epidermis. *DNA Repair* 3: 567-580.
- 33 Hockwin, O.; Kojima, M.; Sakamoto, Y.; Wegener, A.; Shui, Y. B.; Sasaki, K. (1999) UV damage to the eye lens:
34 further results from animal model studies: a review. *J. Epidemiol.* 9(suppl. 6): S39-S47.
- 35 Holick, M. F. (2004) Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and
36 cardiovascular disease. *Am. J. Clin. Nutr.* 80(6 suppl.): 1678S-1688S.
- 37 Hu, S.; Ma, F.; Collado-Mesa, F.; Kirsner, R. S. (2004) UV radiation, latitude, and melanoma in US Hispanics and
38 blacks. *Arch. Dermatol.* 140: 819-824.
- 39 Ichihashi, M.; Ueda, M.; Budyanto, A.; Bito, T.; Oka, M.; Fukunaga, M.; Tsuru, K.; Horikawa, T. (2003)
40 UV-induced skin damage. *Toxicology* 189: 21-39.
- 41 Intergovernmental Panel on Climate Change. (1996) Climate change 1995: the science of climate change. Summary
42 for policymakers and technical summary of the working group I report. Presented at: the fifth session of the
43 IPCC Working Group I; November; Madrid, Spain. Geneva, Switzerland: World Meteorological Association.
- 44 Intergovernmental Panel on Climate Change (IPCC). (2001a) Climate change 2001: the scientific basis. Contribution
45 of working group I to the third assessment report of the Intergovernmental Panel on Climate Change.
46 Cambridge, United Kingdom: Cambridge University Press.
- 47 Intergovernmental Panel on Climate Change (IPCC). (2001b) Climate change 2001: impacts, adaptation, and
48 vulnerability. Contribution of working group II to the third assessment report of the Intergovernmental Panel
49 on Climate Change. Cambridge, United Kingdom: Cambridge University Press.
- 50 International Agency for Research on Cancer. (1992) Solar and ultraviolet radiation. Lyon, France: International
51 Agency for Research on Cancer. (IARC monographs on the evaluation of carcinogenic risks to humans:
52 v. 55).
- 53 International Commission on Non-Ionizing Radiation Protection. (2004) Guidelines on limits of exposure to
54 ultraviolet radiation of wavelengths between 180 nm and 400 nm (incoherent optical radiation). *Health Phys.*
55 87: 171-186.
- 56 Javitt, J. C.; Taylor, H. R. (1994-95) Cataract and latitude. *Doc. Ophthalmol.* 88: 307-325.

- 1 Jemal, A.; Tiwari, R. C.; Murray, T.; Ghafoor, A.; Samuels, A.; Ward, E.; Feuer, E. J.; Thun, M. J. (2004) Cancer
2 statistics, 2004. *CA-Cancer J. Clin.* 54: 8-29.
- 3 John, E. M.; Schwartz, G. G.; Dreon, D. M.; Koo, J. (1999) Vitamin D and breast cancer risk: the NHANES I
4 Epidemiologic Follow-up Study, 1971-1975 to 1992. *Cancer Epidemiol. Biomarkers Prev.* 8: 399-406.
- 5 Kaidbey, K. H.; Kligman, A. M. (1981) Cumulative effects from repeated exposures to ultraviolet radiation.
6 *J. Invest. Dermatol.* 76: 352-355.
- 7 Kerr, J. B.; McElroy, C. T. (1993) Evidence for large upward trends of ultraviolet-B radiation linked to ozone
8 depletion. *Science (Washington, DC)* 262: 1032-1034.
- 9 Kiehl, J. T.; Schneider, T. L.; Portmann, R. W.; Solomon, S. (1999) Climate forcing due to tropospheric and
10 stratospheric ozone. *J. Geophys. Res. [Atmos.]* 104: 31,239-31,254.
- 11 Kimlin, M. G.; Parisi, A. V.; Wong, J. C. (1998) Quantification of personal solar UV exposure of outdoor workers,
12 indoor workers and adolescents at two locations in Southeast Queensland. *Photodermatol. Photoimmunol.*
13 *Photomed.* 14: 7-11.
- 14 Kimlin, M. G.; Wong, J. C.; Parisi, A. V. (1998) Simultaneous comparison of the personal UV exposure of two
15 human groups at different altitudes. *Health Phys.* 74: 429-434.
- 16 Kollias, N.; Stamatas, G. N.; Youn, J. I. (2001) Suppression of UVB-induced cutaneous erythema by a previous
17 UVB exposure. *Photochem. Photobiol.* 74: 471-476.
- 18 Koloutsou-Vakakis, S.; Helmis, C. G.; Assimakopoulos, V.; Güsten, H. (2001) Middle and lower troposphere
19 aerosol characteristics and ozone concentrations over northwestern Greece during STAAARTE 1997.
20 *Atmos. Environ.* 35: 1517-1526.
- 21 Kraemer, K. H. (1997) Sunlight and skin cancer: another link revealed. *Proc. Natl. Acad. Sci. U. S. A.* 94: 11-14.
- 22 Krickler, A.; Armstrong, B. K.; English, D. R. (1994) Sun exposure and non-melanocytic skin cancer. *Cancer Causes*
23 *Control* 5: 367-392.
- 24 Krol, M.; Van Leeuwen, P. J.; Lelieveld, J. (1998) Global OH trend inferred from methylchloroform measurements.
25 *J. Geophys. Res. [Atmos.]* 103: 10,697-10,711.
- 26 Krotkov, N. A.; Bhartia, P. K.; Herman, J. R.; Fioletov, V.; Kerr, J. (1998) Satellite estimation of spectral surface
27 UV irradiance in the presence of tropospheric aerosols 1. cloud-free case. *J. Geophys. Res. (Atmos.)*
28 103: 8779-8793.
- 29 Lefkowitz, E. S.; Garland, C. F. (1994) Sunlight, vitamin D, and ovarian cancer mortality rates in U.S. women.
30 *Int. J. Epidemiol.* 23: 1133-1136.
- 31 Lelieveld, J.; Dentener, F. J. (2000) What controls tropospheric ozone? *J. Geophys. Res. [Atmos.]* 105: 3531-3551.
- 32 Lens, M. B.; Dawes, M. (2004) Global perspectives of contemporary epidemiological trends of cutaneous malignant
33 melanoma. *Br. J. Dermatol.* 150: 179-185.
- 34 Levy, H., II; Kasibhatla, P. S.; Moxim, W. J.; Klonecki, A. A.; Hirsch, A. I.; Oltmans, S. J.; Chameides, W. L.
35 (1997) The global impact of human activity on tropospheric ozone. *Geophys. Res. Lett.* 24: 791-794.
- 36 Liao, H.; Seinfeld, J. H.; Adams, P. J.; Mickley, L. J. (2004) Global radiative forcing of coupled tropospheric ozone
37 and aerosols in a unified general circulation model. *J. Geophys. Res. [Atmos.]* 109(D16):
38 10.1029/2003JD004456.
- 39 Logan, J. A. (1999) An analysis of ozonesonde data for the troposphere: recommendations for testing 3-D models
40 and development of a gridded climatology for tropospheric ozone. *J. Geophys. Res.* 104: 16,115-16,149.
- 41 Longstreth, J.; De Gruijl, F. R.; Kripke, M. L.; Abseck, S.; Arnold, F.; Slaper, H. I.; Velders, G.; Takizawa, Y.;
42 Van Der Leun, J. C. (1998) Health risks. *J. Photochem. Photobiol. B* 46: 20-39.
- 43 Lovato, C. Y.; Shoveller, J. A.; Peters, L.; Rivers, J. K. (1998a) Canadian National Survey on Sun Exposure &
44 Protective Behaviours: parent's reports on children. *Cancer Prev. Control.* 2: 123-128.
- 45 Lovato, C. Y.; Shoveller, J. A.; Peters, L.; Rivers, J. K. (1998b) Canadian National Survey on Sun Exposure &
46 Protective Behaviours: youth at leisure. *Cancer Prev. Control* 2: 117-122.
- 47 Madronich, S.; De Gruijl, F. (1993) Skin cancer and UV radiation. *Nature (London, UK)* 366(6450): 23.
- 48 Madronich, S.; De Gruijl, F. (1994) Reply to "Krickler, A.; Armstrong, B. K.; McMichael, A. J. (1994) Skin cancer
49 and ultraviolet. *Nature (London, UK)* 368: 594." *Nature (London, UK)* 368(6472): 594.
- 50 Matsumura, Y.; Ananthaswamy, H. N. (2004) Toxic effects of ultraviolet radiation on the skin. *Toxicol. Appl.*
51 *Pharmacol.* 195: 298-308.
- 52 McCarty, C. A.; Nanjan, M. B.; Taylor, H. R. (2000) Attributable risk estimates for cataract to prioritize medical and
53 public health action. *Invest. Ophthalmol. Vis. Sci.* 41: 3720-3725.
- 54 McCarty, C. A.; Taylor, H. R. (2002) A review of the epidemiologic evidence linking ultraviolet radiation and
55 cataracts. *Dev. Ophthalmol.* 35: 21-31.

- 1 McKenzie, R.; Smale, D.; Bodeker, G.; Claude, H. (2003) Ozone profile differences between Europe and New
2 Zealand: effects on surface UV irradiance and its estimation from satellite sensors. *J. Geophys. Res. [Atmos.]*
3 108(D6): 10.1029/2002JD002770.
- 4 McKinlay, A. F.; Diffey, B. L. (1987) A reference action spectrum for ultraviolet induced erythema in human skin.
5 In: Passchier, W. F.; Bosnijakovic, B. F. M., eds. *Human exposure to ultraviolet radiation: risks and*
6 *regulations*. Amsterdam, The Netherlands: Elsevier Science Publishers; pp. 83-87.
- 7 Merriam, J. C.; Lofgren, S.; Michael, R.; Soderberg, P.; Dillon, J.; Zheng, L.; Ayala, M. (2000) An action spectrum
8 for UV-B radiation and the rat lens. *Invest. Ophthalmol. Vis. Sci.* 41: 2642-2647.
- 9 Mickley, L. J.; Murti, P. P.; Jacob, D. J.; Logan, J. A.; Koch, D. M.; Rind, D. (1999) Radiative forcing from
10 tropospheric ozone calculated with a unified chemistry-climate model. *J. Geophys. Res. [Atmos.]*
11 104: 30,153-30,172.
- 12 Mickley, L. J.; Jacob, D. J.; Field, B. D.; Rind, D. (2004) Climate response to the increase in tropospheric ozone
13 since preindustrial times: a comparison between ozone and equivalent CO₂ forcings. *J. Geophys. Res.*
14 *[Atmos.]* 109(D5): 10.1029/2003JD003653.
- 15 Mid-Atlantic Regional Assessment (MARA) Team. (2000) Preparing for a changing climate: the potential
16 consequences of climate variability and change. Mid-Atlantic overview. Washington, DC: U.S.
17 Environmental Protection Agency; Office of Research and Development; U.S. Global Change Research
18 Program (USGCRP) and University Park, PA: Pennsylvania State University. Available:
19 <http://www.usgcrp.gov/usgcrp/nacc/midatlantic.htm> [17 April 2002].
- 20 Moehrle, M. (2001) Ultraviolet exposure in the Ironman triathlon. *Med. Sci. Sports Exercise* 33: 1385-1386.
- 21 Moehrle, M.; Heinrich, L.; Schmid, A.; Garbe, C. (2000) Extreme UV exposure of professional cyclists.
22 *Dermatology* 201: 44-45.
- 23 Moise, A. F.; Buttner, P. G.; Harrison, S. L. (1999) Sun exposure at school. *Photochem. Photobiol.* 70: 269-274.
- 24 National Assessment Synthesis Team (NAST). (2000) Climate change impacts on the United States: the potential
25 consequences of climate variability and change. Overview. Washington, DC: U.S. Global Change Research
26 Program. Available at: <http://www.usgcrp.gov/usgcrp/Library/nationalassessment/overview.htm>
27 (17 January 2003).
- 28 National Research Council. (2005) Radiative forcing of climate change: expanding the concept and addressing
29 uncertainties. Washington, DC: The National Academies Press.
- 30 New England Regional Assessment Group (NERAG). (2001) Preparing for a changing climate: the potential
31 consequences of climate variability and change. New England regional overview. Washington, DC: U.S.
32 Environmental Protection Agency; Office of Research and Development; U.S. Global Change Research
33 Program (USGCRP) and Durham, NH: University of New Hampshire; Institute for the Study of Earth,
34 Oceans, and Space. Available: <http://www.necci.sr.unh.edu/2001-NERA-report.html> [17 April 2002].
- 35 Nole, G.; Johnson, A. W. (2004) An analysis of cumulative lifetime solar ultraviolet radiation exposure and the
36 benefits of daily sun protection. *Dermatol. Ther.* 17(suppl. 1): 57-62.
- 37 Noonan, F. P.; Muller, H. K.; Fears, T. R.; Kusewitt, D. F.; Johnson, T. M.; De Fabo, E. D. (2003) Mice with
38 genetically determined high susceptibility to ultraviolet (UV)-induced immunosuppression show enhanced
39 UV carcinogenesis. *J. Invest. Dermatol.* 121: 1175-1181.
- 40 Norval, M.; Garssen, J.; Van Loveren, H.; El-Ghorr, A. A. (1999) UV-induced changes in the immune response to
41 microbial infections in human subjects and animal models. *J. Epidemiol.* 9(suppl. 6): S84-S92.
- 42 Okada, S.; Weatherhead, E.; Targoff, I. N.; Wesley, R.; Miller, F. W.; International Myositis Collaborative Study
43 Group. (2003) Global surface ultraviolet radiation intensity may modulate the clinical and immunologic
44 expression of autoimmune muscle disease. *Arthritis Rheum.* 48: 2285-2293.
- 45 Oriowo, O. M.; Cullen, A. P.; Chou, B. R.; Sivak, J. G. (2001) Action spectrum and recovery for in vitro
46 UV-induced cataract using whole lenses. *Invest. Ophthalmol. Vis. Sci.* 42: 2596-2602.
- 47 Parisi, A. V.; Willey, A.; Kimlin, M. G.; Wong, J. C. (1999) Penetration of solar erythemal UV radiation in the
48 shade of two common Australian trees. *Health Phys.* 76: 682-686.
- 49 Patz, J. A.; Engelberg, D.; Last, J. (2000a) The effects of changing weather on public health. *Annu. Rev. Public*
50 *Health* 21: 271-307.

- 1 Patz, J. A.; McGeehin, M. A.; Bernard, S. M.; Ebi, K. L.; Epstein, P. R.; Grambsch, A.; Gubler, D. J.; Reiter, P.;
2 Romieu, I.; Rose, J. B.; Samet, J. M.; Trtanj, J. (2000b) The potential health impacts of climate variability and
3 change for the United States: executive summary of the report of the health sector of the U.S. national
4 assessment. *Environ. Health Perspect.* 108: 367-376.
- 5 Persson, A. E.; Edstrom, D. W.; Backvall, H.; Lundeberg, J.; Ponten, F.; Ros, A. M.; Williams, C. (2002) The
6 mutagenic effect of ultraviolet-A1 on human skin demonstrated by sequencing the p53 gene in single
7 keratinocytes. *Photodermatol. Photoimmunol. Photomed.* 18: 287-293.
- 8 Pfister, H. (2003) Human papillomavirus and skin cancer. *J. Natl. Cancer Inst. Monogr.* 31: 52-56.
- 9 Pitts, D. G. (1993) Ocular effects of radiant energy. In: Pitts, D. G.; Kleinstein, R. N., eds. *Environmental Vision*.
10 Stoneham, MA: Butterworth-Heinemann, pp. 151-220.
- 11 Prinn, R. G.; Weiss, R. F.; Miller, B. R.; Huang, J.; Alyea, F. N.; Cunnold, D. M.; Fraser, P. J.; Hartley, D. E.;
12 Simmonds, P. G. (1995) Atmospheric trends and lifetime of CH₃CCl₃ and global OH concentrations. *Science*
13 (Washington, DC) 269: 187-192.
- 14 Rigel, D. S.; Rigel, E. G.; Rigel, A. C. (1999) Effects of altitude and latitude on ambient UVB radiation. *J. Am.*
15 *Acad. Dermatol.* 40: 114-116.
- 16 Rivers, J. K. (2004) Is there more than one road to melanoma? *Lancet* 363: 728-730.
- 17 Robinson, E. S.; Hill, R. H., Jr.; Kripke, M. L.; Setlow, R. B. (2000) The Monodelphis melanoma model: initial
18 report on large ultraviolet A exposures of suckling young. *Photochem. Photobiol.* 71: 743-746.
- 19 Roelofs, G. J.; Lelieveld, J.; Van Dorland, R. (1997) A three-dimensional chemistry/general circulation model
20 simulation of anthropogenically derived ozone in the troposphere and its radiative climate forcing.
21 *J. Geophys. Res. [Atmos.]* 102: 23,389-23,401.
- 22 Rosenthal, F. S.; Phoon, C.; Bakalian, A. E.; Taylor, H. R. (1988) The ocular dose of ultraviolet radiation to outdoor
23 workers. *Invest. Ophthalmol. Vis. Sci.* 29: 649-656.
- 24 Rozema, J.; Manetas, Y.; Bjorn, L. O., eds. (2001) Responses of plants to UV-B radiation. Dordrecht, The
25 Netherlands: Kluwer Academic. [Advances in Vegetation Science, v. 18].
- 26 Runger, T. M.; Moller, K.; Jung, T.; Dekant, B. (2000) DNA damage formation, DNA repair, and survival after
27 exposure of DNA repair-proficient and nucleotide excision repair-deficient human lymphoblasts to UVA1
28 and UVB. *Int. J. Radiat. Biol.* 76: 789-797.
- 29 Salby, M. L. (1996) Fundamentals of atmospheric physics. New York, NY: Academic Press. (The International
30 Geophysics Series, Monographs and Textbooks, v. 61).
- 31 Sarasin, A. (1999) The molecular pathways of ultraviolet-induced carcinogenesis. *Mutat. Res.* 428: 5-10.
- 32 Sarkar, A. K. (2004) An evaluation of UV protection imparted by cotton fabrics dyed with natural colorants. *BMC*
33 *Dermatol.* 4: 15.
- 34 Schenker, M. B.; Orenstein, M. R.; Samuels, S. J. (2002) Use of protective equipment among California farmers.
35 *Am. J. Ind. Med.* 42: 455-464.
- 36 Selgrade, M. K.; Repacholi, M. H.; Koren, H. S. (1997) Ultraviolet radiation-induced immune modulation: potential
37 consequences for infectious, allergic, and autoimmune disease. *Environ. Health Perspect.* 105: 332-334.
- 38 Selgrade, M. K.; Smith, M. V.; Oberhelman-Bragg, L. J.; LeVee, G. J.; Koren, H. S.; Cooper, K. D. (2001) Dose
39 response for UV-induced immune suppression in people of color: differences based on erythral reactivity
40 rather than skin pigmentation. *Photochem. Photobiol.* 74: 88-95.
- 41 Setlow, R. B.; Grist, E.; Thompson, K.; Woodhead, A. D. (1993) Wavelengths effective in induction of malignant
42 melanoma. *Proc. Natl. Acad. Sci. U. S. A.* 90: 6666-6670.
- 43 Shindell, D. T.; Faluvegi, G.; Bell, N. (2003) Preindustrial-to-present-day radiative forcing by tropospheric ozone
44 from improved simulations with the GISS chemistry-climate GCM. *Atmos. Chem. Phys.* 3: 1675-1702.
- 45 Shoveller, J. A.; Lovato, C. Y.; Peters, L.; Rivers, J. K. (1998) Canadian National Survey on Sun Exposure &
46 Protective Behaviours: adults at leisure. *Cancer Prev. Control* 2: 111-116.
- 47 Slaper, H.; Velders, G. J. M.; Daniel, J. S.; de Gruijl, F. R.; Van der Leun, J. C. (1996) Estimates of ozone depletion
48 and skin cancer incidence to examine the Vienna Convention achievements. *Nature (London, U.K.)*
49 384: 256-258.
- 50 Stern, R. S.; Liebman, E. J.; Vakeva, L. (1998) Oral psoralen and ultraviolet-A light (PUVA) treatment of psoriasis
51 and persistent risk of nonmelanoma skin cancer. PUVA Follow-up Study. *J. Natl. Cancer Inst.* 90: 1278-1284.
- 52 Stevenson, D. S.; Johnson, C. E.; Collins, W. J.; Derwent, R. G.; Shine, K. P.; Edwards, J. M. (1998) Evolution of
53 tropospheric ozone radiative forcing. *Geophys. Res. Lett.* 25: 3819-3822.
- 54 Stevenson, D. S.; Johnson, C. E.; Collins, W. J.; Derwent, R. G.; Edwards, J. M. (2000) Future estimates of
55 tropospheric ozone radiative forcing and methane turnover - the impact of climate change. *Geophys. Res.*
56 *Lett.* 27: 2073-2076.
- 57 Studzinski, G. P.; Moore, D. C. (1995) Sunlight--can it prevent as well as cause cancer? *Cancer Res.* 55: 4014-4022.

- Swetter, S. M. (2003) Dermatological perspectives of malignant melanoma. *Surg. Clin. N. Am.* 83: 77-95.
- Taylor, H. R.; West, S. K.; Rosenthal, F. S.; Munoz, B.; Newland, H. S.; Abbey, H.; Emmett, E. A. (1988) Effect of ultraviolet radiation on cataract formation. *N. Engl. J. Med.* 319: 1429-1433.
- Thieden, E.; Agren, M. S.; Wulf, H. C. (2001) Solar UVR exposures of indoor workers in a working and a holiday period assessed by personal dosimeters and sun exposure diaries. *Photodermatol. Photoimmunol. Photomed.* 17: 249-255.
- Thieden, E.; Philipsen, P. A.; Heydenreich, J.; Wulf, H. C. (2004a) UV radiation exposure related to age, sex, occupation, and sun behavior based on time-stamped personal dosimeter readings. *Arch. Dermatol.* 140: 197-203.
- Thieden, E.; Philipsen, P. A.; Sandby-Moller, J.; Heydenreich, J.; Wulf, H. C. (2004b) Proportion of lifetime UV dose received by children, teenagers and adults based on time-stamped personal dosimetry. *J. Invest. Dermatol.* 123: 1147-1150.
- Tomescu, D.; Kavanagh, G.; Ha, T.; Campbell, H.; Melton, D. W. (2001) Nucleotide excision repair gene *XPD* polymorphisms and genetic predisposition to melanoma. *Carcinogenesis* 22: 403-408.
- U.S. Environmental Protection Agency. (1987) Assessing the risks of trace gases that can modify the stratosphere, v. 1-8. Washington, DC: Office of Air and Radiation; report no. EPA 400/1-87/001A-H.
- U.S. Environmental Protection Agency. (2004) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment; report no. EPA/600/P-99/002aF-bF. 2v. Available: <http://cfpub.epa.gov/ncea/> [9 November, 2004].
- United Nations Environment Programme (UNEP). (1986) Report of the international conference on the assessment of the role of carbon dioxide and of other greenhouse gases in climate variations and associated impacts; October 1985; Villach, Austria. Geneva, Switzerland: World Meteorological Organization; WMO no. 661.
- Van Der Leun, J. C.; Takiyawa, Y.; Longstreth, J. D. (1989) Human health. In: Van der Leun, J. C.; Trevini, M.; Worress, R. C., eds. *Environmental Effects Panel Report*. Nairobi, Kenya: United Nations Environment Programme, pp. 11-24. Available: <http://www.ciesin.org/docs/001-538/001-538.html>.
- Van Dorland, R.; Dentener, F. J.; Lelieveld, J. (1997) Radiative forcing due to tropospheric ozone and sulfate aerosols. *J. Geophys. Res. [Atmos.]* 102: 28,079-28,100.
- Veierød, M. B.; Weiderpass, E.; Thörn, M.; Hansson, J.; Lund, E.; Armstrong, B.; Adami, H.-O. (2003) A prospective study of pigmentation, sun exposure, and risk of cutaneous malignant melanoma in women. *J. Natl. Cancer Inst.* 95: 1530-1538.
- Vishvakarman, D.; Wong, J. C.; Boreham, B. W. (2001) Annual occupational exposure to ultraviolet radiation in central Queensland. *Health Phys.* 81: 536-544.
- Vitasa, B. C.; Taylor, H. R.; Strickland, P. T.; Rosenthal, F. S.; West, S.; Abbey, H.; Ng, S. K.; Munoz, B.; Emmett, E. A. (1990) Association on nonmelanoma skin cancer and actinic keratosis with cumulative solar ultraviolet exposure in Maryland watermen. *Cancer* 65: 2811-2817.
- Waterston, K.; Naysmith, L.; Rees, J. L. (2004) Physiological variation in the erythral response to ultraviolet radiation and photoadaptation. *J. Invest. Dermatol.* 123: 958-964.
- Wayne, R. P. (2000) *Chemistry of Atmospheres: an introduction to the chemistry of the atmospheres of Earth, the planets, and their satellites*. 3rd ed. New York, NY: Oxford University Press, Inc.
- Wei, Q.; Lee, J. E.; Gershenwald, J. E.; Ross, M. I.; Mansfield, P. F.; Strom, S. S.; Wang, L. E.; Guo, Z.; Qiao, Y.; Amos, C. I.; Spitz, M. R.; Duvic, M. (2003) Repair of UV light-induced DNA damage and risk of cutaneous malignant melanoma. *J. Natl. Cancer Inst.* 95: 308-315.
- Wendisch, M.; Mayer, B. (2003) Vertical distribution of spectral solar irradiance in the cloudless sky: a case study. *Geophys. Res. Lett.* 30: 10.1029/2002GL016529.
- Westerdahl, J.; Ingvar, C.; Måsbäck, A.; Jonsson, N.; Olsson, H. (2000) Risk of cutaneous malignant melanoma in relation to use of sunbeds: further evidence for UV-A carcinogenicity. *Br. J. Cancer* 82: 1593-1599.
- Whiteman, D. C.; Valery, P.; McWhirter, W.; Green, A. C. (1997) Risk factors for childhood melanoma in Queensland, Australia. *Int. J. Cancer* 70: 26-31.
- Whiteman, D. C.; Whiteman, C. A.; Green, A. C. (2001) Childhood sun exposure as a risk factor for melanoma: a systematic review of epidemiologic studies. *Cancer Causes Control* 12: 69-82.
- Wigley, T. M. L.; Smith, S. J.; Prather, M. J. (2002) Radiative forcing due to reactive gas emissions. *J. Clim.* 15: 2690-2696.
- Wong, J. C.; Airey, D. K.; Fleming, R. A. (1996) Annual reduction of solar UV exposure to the facial area of outdoor workers in southeast Queensland by wearing a hat. *Photodermatol. Photoimmunol. Photomed.* 12: 131-135.
- Woodward, A.; Boffetta, P. (1997) Environmental exposure, social class, and cancer risk. In: Kogevinas, M.; Pearce, N.; Susser, M.; Boffetta, P. *Social inequalities and cancer*. Lyon, France: International Agency for Research on Cancer; pp. 361-367. (IARC scientific publications: v. 138).

- 1 World Meteorological Organization. (1988) Developing policies for responding to climatic change: a summary of
2 the discussions and recommendations of workshops; September-October 1987; Villach, Austria; and
3 November 1987; Bellagio, Austria. Geneva, Switzerland: World Meteorological Organization; report no.
4 WMO/TD; no. 225. [World Climate Impact Programme series report no. WCIP-1].
- 5 World Meteorological Organization (WMO). (2003) Scientific assessment of ozone depletion: 2003. Geneva,
6 Switzerland: United Nations Environment Programme. [Global Ozone and Monitoring Project; report no. 47].
- 7 Xenopoulos, M. A.; Schindler, D. W. (2001) Physical factors determining ultraviolet radiation flux into ecosystems.
8 In: Cockell, C.; Blaustein, A. R. Ecosystems, evolution, and ultraviolet radiation. New York, NY: Springer;
9 pp. 36-62.
- 10 Yarnal, B.; Kalkstein, L. S.; Scheraga, J. D. (2000) Mid-Atlantic regional assessment of climate change impacts.
11 Clim. Res. (CR Special 7) 14: 153-269.
- 12 Zeng, G.; Pyle, J. A. (2003) Changes in tropospheric ozone between 2000 and 2100 modeled in a chemistry-climate
13 model. Geophys. Res. Lett. 30: 10.1029/2002GL016708.
- 14
15

11. EFFECT OF OZONE ON MAN-MADE MATERIALS

Ozone and other photochemical oxidants react with many of the economically important man-made materials, decreasing their useful life and aesthetic appearance. Some of the materials known to be damaged by ozone include elastomers, fibers and dyes, and paints. This section will provide a brief discussion on the effects of ozone on man-made materials including the damage mechanisms and, where possible, concentration-response relationships. Since only limited information has been published on the effects of ozone on materials, this section will provide a summary of information presented in the previous ozone criteria document (U.S. Environmental Protection Agency, 1996) and a more detailed discussion of studies published since publication of the previous ozone criteria document. The reader is referred to the previous ozone criteria document for a more detailed discussion of the earlier studies.

11.11.1 Mechanisms of Ozone Damage and Exposure-Response

Elastomer Cracking

The elastomeric compounds, natural rubber and synthetic polymers and copolymers of butadiene, isoprene, and styrene, are particularly susceptible to even low levels of ozone. Elastomeric compounds are long chain unsaturated organic molecules. Ozone damages these compounds by breaking the molecular chain at the carbon-carbon double bond; a chain of three oxygen atoms is added directly across the double bond, forming a five-membered ring structure (Mueller and Stickney, 1970). The change in structure promotes the characteristic cracking of stressed/stretched rubber called “weathering.” A 5% tensile strain will produce cracks on the surface of the rubber that increase in number with increased stress/stretching. The rate of crack growth is dependent on the degree of stress, the type of rubber compound, ozone concentration, time of exposure, ozone velocity, and temperature (Bradley and Haagen-Smit, 1951; Lake and Mente, 1992) (Gent and McGrath, 1965). Once cracked, there is further ozone penetration and in additional cracking and eventually mechanical weakening or stress relaxation (U.S. Environmental Protection Agency, 1996). Razumovskii et al. (1988) demonstrated the effect of ozone on stress relaxation of polyisoprene vulcanizates. A decrease in stress (stress relaxation) was caused by ozone-induced cracks in exposed elastomers resulting in irreversible changes in the elastomer dimensions and decreased tensile strength. A description of findings of earlier

1 studies appear in the 1996 Air Quality Criteria Document for ozone (U.S. Environmental
2 Protection Agency, 1996).

3 To counteract the effect of ozone on elastomers, antiozonants and wax have been added to
4 the elastomeric formulations during processing. An antiozonant is an additive used to protect a
5 polymer against the effects of ozone-induced degradation and, hence, used mainly in diene
6 rubbers. Antiozonant protection works either by providing a physical barrier to ozone
7 penetration forming a thin surface film of an ozone-resisting wax or by chemically reacting with
8 ozone or polymer ozonolysis products, as do aromatic diamines such as p-phenylene diamine
9 derivatives. The antiozonant diffuses to the surface of the elastomeric material where it reacts
10 with ozone faster than ozone reacts to break the molecular chain and the carbon-carbon double
11 bond, or the antiozonant diffuses to the surface of the material but is not reactive with ozone and
12 serves as a protective coating against ozone attack. The antiozonant may also serve to scavenge
13 ozone while also providing protective film against ozone attack (Andries et al., 1979; Lattimer
14 et al., 1984).

15 Most of the studies on ozone effects on elastomers were designed to evaluate the
16 effectiveness of antiozonants in counteracting the rubber cracking produced by ozone exposure.
17 Consequently, many of the studies were conducted using ozone concentrations higher than those
18 typically found in the ambient air. Natural rubber strips exposed to high concentrations of ozone
19 (20,000 ppm) under stressed conditions cracked almost instantaneously and were broken within
20 1 sec. When the ozone concentration was lowered (0.02 to 0.46 ppm), the time to required to
21 produce cracks in the exposed rubber material was increased (Bradley and Haagen-Smit, 1951).
22 Lake and Mente (1992) studied the effect of temperature on ozone-induced elastomer cracking
23 and antiozonant protection on natural rubber, epoxidised natural rubber, and two acrylonitrile-
24 butadiene copolymers under constant strain. Temperatures ranged from -20° C to +70° C. The
25 elastomers were exposed to 0.05 to 1,000 ppm ozone for 16 h. Ozone cracking decreased at
26 lower ambient temperatures, however, diffusing of both chemical and wax antiozonants also
27 were slowed at the lower temperatures. Cracking was slightly increased at the higher
28 temperatures but the antiozonants offered more protection.

29 Serrano et al. (1993) evaluated the appropriateness of using ozone-induced elastomer
30 cracking to estimated the ambient ozone concentrations. Two vulcanized natural rubber
31 compounds were exposed for 24 h to varying ozone concentrations under stressed conditions.

Ozone concentrations were 60, 80, 90, 100, and 120 ppb for durations of 2, 4, or 6 h. The 24 h average ozone concentrations ranged from 31 to 57.5 ppb. There was a clear relationship between the 24-h average ozone concentration and the distribution of crack length frequencies on the rubber surface. Table 11-1 gives the average 24-h ozone concentration and lengths for two vulcanized natural rubber strips.

Table 11-1. Average 24-h Ozone Concentrations Producing the Highest Frequency of Cracks of a Certain Length in the Middle and Central Zones of the Rubber Test Strips

Crack Length (mm)	1% Antiozonant 4010NA #		0.5% Antiozonant 4010NA	
	Middle Zones	Central Zones	Middle Zones	Central Zones
0.05 - 0.10	37.5	37.5	40.0	42.5
0.10 - 0.15	45.0	48.0	48.0	53.0
0.15 - 0.20	48.0	≥ 57.5	≥ 57.5	≥ 57.5
0.20 - 0.40	≥ 57.5	≥ 57.5	≥ 57.5	≥ 57.5

Ozone concentrations given in ppb.

Adapted from Serrano et al. (1993).

11.11.2 Textiles and Fabrics

Ozone can damage textiles and fabrics by methods similar to those associated with elastomers. Generally, synthetic fibers are less affected by ozone than natural fibers, however, ozone contribution to the degradation of textiles and fabrics is not considered significant (U.S. Environmental Protection Agency, 1996). A study reported by Bogaty et al. (1952) showed that ozone effects moistened cloth more than dry cloth. Scoured cotton duck cloth and commercially bleached cotton print cloth were exposed to 20 to 60 ppb for 1,200 h (50 days). The rate of deterioration was measured by the changes in cuprammonium fluidity values and the fabric breaking strength. At the end of the 1,200-h exposure, there was a 20% loss in breaking strength. Table 11-2 list the changes in cuprammonium fluidity values for both fabrics.

**Table 11-2. Cuprammonium Fluidity of Moist Cotton Cloth Exposed
to 20 to 60 ppb Ozone**

	Duration of Exposure (h)	Cuprammonium Fluidity (rhes)
Duck Cloth	0	2.6
	200	2.8
	680	4.0
	960	6.8
	1200	9.5
Bleached Print Cloth	0	8.2
	200	8.7
	510	9.4
	650	12.0
	865	12.7
	1500	16.5

Adapted from Bogaty et al. (1952).

11.11.3 Dyes, Pigments, and Inks

Ozone fading of textile dyes is diffusion-controlled; the rate of fading is controlled by the diffusion of the dye to the fiber surface. Many textile dyes react with ozone; however, the rate and severity of the ozone attack is influenced by the chemical nature of the textile fiber and the manner in which the dye is applied. Ozone molecules break the aromatic ring portion of the dye molecule, oxidizing the dye (U.S. Environmental Protection Agency, 1996). In case of aromatic azo dyes, ozone attacks the aromatic rings and electron rich nitrogen atoms (Matsui et al., 1988). Grosjean et al. (1987; 1988a,b) proposed a mechanism of reaction of ozone and indigo, thioindigo, and dibromoindigo, alazarin, and curcumin dyes under dark conditions. Ozone attaches to the dye molecule at the unsaturated carbon = carbon bond. An ozone adduct is formed (1,2,3-trioxolane), followed by scission of the carbon–carbon bond and the subsequent formation of the corresponding Criegee biradical. A similar mechanism was proposed for the reaction of ozone with triphenylmethane colorant Basic Violet 14. Ozone attacked Basic Violet 14 at the carbon=carbon unsaturated bond and at the carbon–nitrogen unsaturated bond under dark conditions. Other members of the group of triphenylmethane colorants with unsaturated carbon–carbon bonds also are expected to be subject to ozone fading. Triphenylmethane colorants that are expected to be ozone-fugitive include the amino-substituted cationic dyes

(Malachite Green, Brilliant Green, Crystal Violet, Pararosaniline Chloride, Methyl Green, and others) (Grosjean et al., 1989).

An indication that ozone caused textile dye fading was first reported by Salvin and Walker (1955). The researchers found that the fading was primarily the result of the destruction of the blue dye molecule. Drapes made of acetate, Arnel, and Dacron and dyed with anthraquinone blue dye exhibited a decrease in shade that was not accompanied by the characteristic reddening caused by NO_x . Figures 11-1 and 11-2 demonstrate the effect of ozone exposure on nylon 6 yarn colored with several blue dyes. Nylon samples inside the home were located on a wall away from sunlight. Outside nylon samples were placed on a covered patio or under the eaves of the house to minimize exposure to sunlight and rain. Ozone concentrations ranged from 2 to 5 ppb outside and 0 to 2 ppb inside. The percent change in dye color was determined monthly by extraction and analysis of the remaining dye or by instrumental measurement of the color change (Haylock and Rush, 1978).

11.11.4 Artists' Pigments

Several artists' pigments are sensitive to fading and oxidation by ozone when exposed to concentrations found in urban areas (Shaver et al., 1983; Drisko et al., 1985; Whitmore et al., 1987; Whitmore and Cass, 1988; Grosjean et al., 1993). The organic pigments that are ozone fugitive include alizarin red pigments containing lakes of the polycyclic aromatic compound 1,2-dihydroxyanthraquinone, blue-violet pigments containing substituted triphenylmethane lakes, indigo, and yellow coloring agents containing polyfunctional, polyunsaturated compounds such as curcumin (Grosjean et al., 1987). Because of the potential of ozone to damage works of art, recommended limits on ozone concentrations in museums, libraries, and archives are relatively low, ranging from 0.013 to 0.01 ppm.

Experimental studies demonstrate a concentration \times time ($C \times T$) relationship between ozone concentration and exposure time and pigment fading. Cass et al. (1991) summarized some of the earlier research on the effects of ozone on artists' pigments. In studies evaluating the effect of ozone on organic and inorganic watercolors and traditional organic pigments only the traditional organic pigments showed measurable fading from ozone exposure. Of the inorganic pigments tested, only the arsenic sulfides showed ozone-related changes. The pigments were exposed to 0.3 to 0.4 ppm ozone for 3 mo in the absence of light, at 22 °C and




Figure 11-1. In-service fading of nylon 6 yarn inside house. ● = C.I. Disperse Blue 3; O = C.I. Basic Blue 22; ▲ = C.I. Acid Blue 27; x = C.I. Disperse Blue 56; Δ = C.I. Acid Blue 232.

Source: Haylock and Rush (1978).

50% RH. The authors equated this exposure to a C × T of 6 to 8 years inside a Los Angeles museum with air conditioning but without a pollutant removal system.

Whitmore and Cass (1988) studied the effect of ozone on traditional Japanese colorants. Most of these compounds are insoluble metal salts that are stable in light and air. Suspensions or solutions of the colorants were airbrushed on hot-pressed watercolor paper or silk cloths. A sample of Japanese woodblock print also was included in the analysis. Samples were exposed to 0.4 ppm ozone at 22 °C, 50% relative humidity, in the absence of light for 12 wk. Changes in reflectance spectra were used to evaluate the effect of ozone exposure on colorant fading. Among the colorants tested on paper, curmin, indigo, madder lake, and lac lake were the most sensitive to ozone exposure. Gamboge was relatively insensitive to ozone.




Figure 11-2. In-service fading of nylon 6 yarn outside house. ● = C.I. Disperse Blue 3; O = C.I. Basic Blue 22; ▲ = C.I. Acid Blue 27; x = C.I. Disperse Blue 56; △ = C.I. Acid Blue 232.

Source: Haylock and Rush (1978).

The blue and green areas of the sample from the woodblock print was very reactive due to the indigo dye ozone sensitivity. The other colorants, red, yellow, and purple, showed very little sensitivity to ozone. The textiles dyes that reacted with ozone were indigo, alone or in combination with several yellow dyes.

Ye et al. (2000) reported the rate of ozone fading of traditional Chinese plant dyes. Twelve different colorants were applied to watercolor paper and silk and exposed to 0.4 ppm ozone at 25° C, at 50% RH, in the absence of light for 22 wks. Dye fading was greater when the colorant was applied to the watercolor paper compared to the silk cloth due to the darker initial depth of the shade, the greater saturation of the colorant throughout the cloth. Tumeric, gromwell, and violet on paper was particularly reactive. Tangerine peel was moderately reactive and sappan wood, dalbergia wood, Chinese gall, indigo, and Chinese yellow cork tree were slightly reactive

1 to ozone. Black tea was not reactive to ozone. The colorants on silk samples showing color
2 changes were gromwell, sappan wood, gardenia, tumeric, and violet. Figures 11-3 and 11-4
3 demonstrate the color change of the various colorants on watercolor paper and silk.

4 Artists' pigments also have exhibited fading when exposed to a mixture of photochemical
5 oxidants. Grosjean et al. (1993) exposed 35 artists' pigments to a mixture of photochemical
6 oxidants consisting of ozone, nitrogen dioxide (NO₂), and peroxyacetyl nitrate (PAN) for
7 12 wks. Weekly average photochemical concentrations were 200 ppb for ozone, 56 ± 12 to
8 99 ± 24 for NO₂, and 11 ± 3 to 18 ± 2 for PAN. All exposures were carried out at room
9 temperature in the absence of light. To determine the effect of humidity on pigment fading, the
10 relative humidity was increased from 46% after 8 weeks of exposure to 83% for a 2 week period
11 and then returned to 46% for the remainder of the exposure.

12 Table 11-3 lists the artists' pigment and degree of fading. Eleven of the pigments
13 exhibited negligible color change, 12 had small color changes, 3 had modest color changes, and
14 9 exhibited substantial color changes. Fading of Disperse Blue 3 and Reactive Blue 2 were
15 likely the result of NO₂ exposure, the fading of triphenylmethanes is consistent with exposure to
16 nitric acid formed under high humidity conditions. Fading of the indigos was dominated by
17 ozone exposure and curcumin was faded by all of the photochemicals studied. Increasing the
18 relative humidity resulted in a substantial color change for all of the pigments, with the
19 exception of curcumin and indigo.

21 **11.11.5 Surface Coatings**

22 Ozone will act to erode some surface coatings (paints, varnishes, and lacquers). However,
23 many of the available studies on ozone degradation of surface coatings do not separate the
24 effects of ozone from other pollutants or environmental factors such as weather, humidity, and
25 temperature. Campbell et al. (1974) attempted to demonstrate an ozone related effect on oil
26 house paint, acrylic latex coating, alkyd industrial maintenance coating, urea alkyd coil coating,
27 and nitrocellulose/acrylic automotive paint. Painted test panels were exposed to 100 and
28 1,000 ppb ozone in a xenon arc accelerated weathering chamber for up to 1,000 h. Using weight
29 loss as a measure of ozone-induced erosion the researchers concluded that all of the paints tested
30 suffered degradation in the presence of ozone and that the automotive finish suffered the most





Figure 11-3. Observed color changes for natural colorant-on-paper systems during exposure to 0.40 ppm ozone at 25 °C ± 1 °C, 50% RH, in the absence of light.

Source: Ye et al. (2000).





Figure 11-4. Observed color changes for natural colorant-on-site during exposure to 0.40 ppm ozone at 25 °C ± 1 °C, 50% RH, in the absence of light.

Source: Ye et al. (2000).

Table 11-3. Color Change After 12 Weeks of Exposure to a Mixture of Photochemical Oxidants

Colorant*	Color Change (ΔE units)[†]	Chemical Functionality or Chemical Composition
Acid Red 37 (17045) [‡]	11.7 ± 0.5	Aminophenyl-substituted azo dye, sulfonate salt
Acid Yellow 65 [‡]	1.8 ± 0.5	Nitro- and phenyl-substituted azo dye, sulfonate salt
Alizarin Carmine	1.8 ± 0.2	Alizarin lake
Alizarin Crimson (Pigment Red 83)	1.4 ± 0.2	Alizarin lake
Aurora Yellow (77199)	0.5 ± 0.1	Cadmium sulfide
Basic Fuschin (42510) [‡]	33.4 ± 3.0	Amino-substituted triphenylmethane
Brilliant Green (42040) [‡]	20.6 ± 2.1	Amino-substituted triphenylmethane
Brown Madder	1.7 ± 0.1	Alizarin lake
Cadmium Yellow (77199)	0.4 ± 0.1	Cadmium sulfide
Carmine	1.8 ± 0.2	Lake of cochineal (substituted anthraquinone)
Chrome Yellow (77600) [‡]	1.7 ± 1.2	Lead chromate
Copper phthalocyanine (Pigment Blue 15)	1.0 ± 0.1	Copper phthalocyanine
Crimson Lake	3.5 ± 0.3	Alizarin lake
Curcumin (Natural Yellow 3)	15.2 ± 2.6	1,7 bis (4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione
Disperse Blue 3	10.8 ± 0.1	Amino-substituted anthraquinone
French Ultramarine Blue	0.8 ± 0.3	
Gamboge (Natural Yellow 24)	0.4 ± 0.1	Gambogic acid
Hooker's Green Light	1.5 ± 0.4	Chlorinated copper phthalocyanine plus ferrous beta naphthol derivative
Indigo (a formulation)	1.1 ± 0.1	Alizarin lake plus lampblack plus copper phthalocyanine
Indigo carmine [‡]	14.0 ± 1.9	5,5-indigo disulfonic acid, sodium salt
Indigo (73000) [‡]	64.1 ± 4.5	
Mauve	3.6 ± 0.5	Lake of triphenyl methane (basic fuschin) plus copper phthalocyanine
New Gamboge	0.9 ± 0.1	Arylamide yellow (CI 11680) plus toluidine red

Table 11-3 (cont'd). Color Change After 12 Weeks of Exposure to a Mixture of Photochemical Oxidants

Colorant*	Color Change (ΔE units)[†]	Chemical Functionality or Chemical Composition
Pararosaniline base (42500) [‡]	25.6 ± 4.7	Amino-substituted triphenylmethane
Payne's Grey	1.0 ± 0.1	Alizarin lake plus prussian blue plus lampblack plus ultramarine blue
Permanent Magenta	1.1 ± 0.1	Quinacridone
Permanent Rose	2.0 ± 0.1	Quinacridone
Prussian Blue	0.7 ± 0.2 1.6 ± 0.3	Ferric ferrocyanide
Prussian Green	0.9 ± 0.2	Arylamide yellow plus prussian blue
Purple Lake	2.3 ± 0.3	Alizarin lake
Reactive Blue 2 (61211) [‡]	14.4 ± 1.1	Amino-substituted anthraquinone, sulfonate salt
Rose Carthane (12467)	0.8 ± 0.2	Arylamide (Pigment Red 10) plus xanthene (Pigment Red 90)
Rose Doré	2.0 ± 0.2	Quinacridone plus Yellow 3
Thioindigo Violet (73312) [‡]	1.9 ± 1.2	Chlorinated thioindigo
Winsor Yellow (11680)	0.5 ± 0.2	Arylamide yellow

* On watercolor paper unless otherwise indicated. Color Index (CI) names or CI numbers are given in parentheses.

[†] Mean ± one standard deviation for triplicate samples calculated from the parameters L^* , a^* , and b^* measured with the color analyzer.

[‡] On Whatman 41 paper.

Source: Grosjean et al. (1993)

ozone-induce degradation. When ozone degradation was measured using scanning electron microscopy, the oil house paint and latex coating samples showed erosion above that seen with clean air but only at the highest exposure level. No effects were noted for the automotive paint. The other painted surfaces were not evaluated.

Spence et al. (1975) studied the effect of air pollutants and relative humidity on oil based house paint, acrylic latex house paint, acrylic coil coating, and vinyl coil coating under laboratory conditions. Test panels were exposed in weathering chambers equipped with a xenon are light for simulating sunlight to low and high levels of ozone (0.08 and 0.5 ppm), sulfur

1 dioxide (0.03 and 0.5 ppm), and nitrogen dioxide (0.05 and 0.5 ppm) and relative humidity
2 (50 and 90%). Samples were exposed for a total of 1000 h. The exposure cycle consisted of
3 20 min of dew and 20 min of light. The effects of the exposure on the painted surfaces were
4 measured by weight loss and loss in film thickness. The acrylic coil coating had the lowest
5 erosion rate of the surface coatings tested. However, ozone was the only pollutant that had a
6 significant effect on the surface erosion. Sulfur dioxide and relative humidity were significant
7 factors in the erosion of oil base house paints and vinyl coil coating. The findings for acrylic
8 latex house paint were not reported.

REFERENCES

- Andries, J. C.; Rhee, C. K.; Smith, R. W.; Ross, D. B.; Diem, H. E. (1979) A surface study of ozone attack and antiozonant protection of carbon black loaded natural rubber compounds. *Rubber Chem. Technol.* 52: 823-837.
- Bogaty, H.; Campbell, K. S.; Appel, W. D. (1952) The oxidation of cellulose by ozone in small concentrations. *Text. Res. J.* 22: 81-83.
- Bradley, C. E.; Haagen-Smit, A. J. (1951) The application of rubber in the quantitative determination of ozone. *Rubber Chem. Technol.* 24: 750-755.
- Campbell, G. G.; Schurr, G. G.; Slawikowski, D. E.; Spence, J. W. (1974) Assessing air pollution damage to coatings. *J. Paint Technol.* 46: 59-71.
- Cass, G. R. (1991) Deterioration of materials due to ozone exposure: current problems and future research needs. In: Berglund, R. L.; Lawson, D. R.; McKee, D. J., eds. *Tropospheric ozone and the environment: papers from an international conference; March 1990; Los Angeles, CA. Pittsburgh, PA: Air & Waste Management Association; pp. 311-320. (A&WMA transactions series no. TR-19).*
- Drisko, K.; Cass, G. R.; Whitmore, P. M.; Druzik, J. R. (1985) Fading of artists' pigments due to atmospheric ozone. In: Vendl, A.; Pichler, B.; Weber, J.; Banik, G., eds. *Wiener Berichte über Naturwissenschaft in der Kunst: Doppelband 2/3. Vienna, Austria: Verlag ORAC; pp. 66-87.*
- Gent, A. N.; McGrath, J. E. (1965) Effect of temperature on ozone cracking of rubbers. *J. Polymer Sci. A* 3: 1473-1482.
- Grosjean, D.; Whitmore, P. M.; De Moor, C. P.; Cass, G. R.; Druzik, J. R. (1987) Fading of alizarin and related artists' pigments by atmospheric ozone: reaction products and mechanisms. *Environ. Sci. Technol.* 21: 635-643.
- Grosjean, D.; Whitmore, P. M.; Cass, G. R.; Druzik, J. R. (1988a) Ozone fading of natural organic colorants: mechanisms and products of the reaction of ozone with indigos. *Environ. Sci. Technol.* 22: 292-298.
- Grosjean, D.; Whitmore, P. M.; De Moor, C. P.; Cass, G. R.; Druzik, J. R. (1988b) Ozone fading of organic colorants: products and mechanism of the reaction of ozone with curcumin. *Environ. Sci. Technol.* 22: 1357-1361.
- Grosjean, D.; Whitmore, P. M.; Cass, G. R.; Druzik, J. R. (1989) Ozone fading of triphenylmethane colorants: reaction products and mechanisms. *Environ. Sci. Technol.* 23: 1164-1167.
- Grosjean, D.; Grosjean, E.; Williams, E. L., II. (1993) Fading of artists' colorants by a mixture of photochemical oxidants. *Atmos. Environ. Part A* 27: 765-772.
- Haylock, J. C.; Rush, J. L. (1978) Studies on the ozone fading of anthraquinone dyes on nylon fibers. Part II: In-service performance. *Text. Res. J.* 48: 143-149.
- Lake, G. J.; Mente, P. G. (1992) Ozone cracking and protection of elastomers at high and low temperatures. *J. Nat. Rubber Res.* 7: 1-13.
- Lattimer, R. P.; Layer, R. W.; Rhee, C. K. (1984) Mechanisms of antiozonant protection: antiozonant-rubber reactions during ozone exposure. In: 32nd annual conference on mass spectrometry and allied topics. May/June; San Antonio, TX. Bethesda, MD: American Society for Mass Spectrometry; pp. 357-358.
- Matsui, M.; Koike, T.; Shibata, K. (1988) Ozone fading of phenolphthalein and aurin. *J. Soc. Dyers Colour* 104: 482-486.
- Mueller, W. J.; Stickney, P. B. (1970) A survey and economic assessment of the effects of air pollution on elastomers: final report. Columbus, OH: Battelle Memorial Institute, Columbus Laboratories; National Air Pollution Control Administration contract no. CPA-22-69-146.
- Razumovskii, S. D.; Podmasteriev, V. V.; Zaikov, G. E. (1988) Kinetics and mechanism of stress relaxation of polyisoprene vulcanizates under ozone ageing. *Polym. Degrad. Stab.* 20: 37-47.
- Salvin, V. S.; Walker, R. A. (1955) Service fading of disperse dyestuffs by chemical agents other than the oxides of nitrogen. *Text. Res. J.* 25: 571-585.
- Serrano, E.; Castro, M.; Macías, A. (1993) An improved direct method of rubber cracking analysis for estimating 24-hour ozone levels. *Atmos. Environ. Part A* 27: 431-442.
- Shaver, C. L.; Cass, G. R.; Druzik, J. R. (1983) Ozone and the deterioration of works of art. *Environ. Sci. Technol.* 17: 748-752.
- Spence, J. W.; Haynie, F.; Upham, J. B. (1975) Effects of gaseous pollutants on paints: a chamber study. *J. Paint Technol.* 47: 57-63.

- 1 U.S. Environmental Protection Agency. (1996) Air quality criteria for ozone and related photochemical oxidants.
2 Research Triangle Park, NC: Office of Research and Development; report nos. EPA/600/AP-93/004aF-cF. 3v.
3 Available from: NTIS, Springfield, VA; PB96-185582, PB96-185590, and PB96-185608. Available:
4 <http://cfpub2.epa.gov/ncea/>.
- 5 Whitmore, P. M.; Cass, G. R. (1988) The ozone fading of traditional Japanese colorants. *Stud. Conserv.* 33: 29-40.
6 Whitmore, P. M.; Cass, G. R.; Druzik, J. R. (1987) The ozone fading of traditional natural organic colorants on
7 paper. *J. Am. Inst. Conserv.* 26: 45-58.
- 8 Ye, Y.; Salmon, L. G.; Cass, G. R. (2000) The ozone fading of traditional Chinese plant dyes. *J. Am. Inst. Conserv.*
9 39: 245-257.

10