




Contaminant Trends In Lake Trout (*Salvelinus namaycush*) From the Upper Great Lakes



EPA 905/3-85-001
and
GLFL no. 639

CONTAMINANT TRENDS
IN
LAKE TROUT (Salvelinus namaycush)
OF THE
UPPER GREAT LAKES

by

David S. DeVault
U.S. Environmental Protection Agency
Great Lakes National Program Office

Wayne A. Willford and Robert J. Hesselberg
U.S. Fish and Wildlife Service
Great Lakes Fishery Laboratory
1451 Green Road
Ann Arbor, Michigan 48105

For

U.S. Environmental Protection Agency
Great Lakes National Program Office
536 South Clark Street
Chicago, Illinois 60605

DISCLAIMER

"This report has completed the EPA peer and publications review process and is approved for publication as an EPA document. Mention of trade names or commercial products does not constitute endorsement or recommendation by EPA."

FOREWORD

The Great Lakes National Program Office (GLNPO) of the U.S. Environmental Protection Agency was established in Region V, Chicago to focus attention on the significant and complex natural resource represented by the Great Lakes.

GLNPO implements a multi-media environmental management program drawing on a wide range of expertise represented by universities, private firms, State, Federal and Canadian governmental agencies and the International Joint Commission. The goal of the GLNPO program is to develop programs, practices and technology necessary for a better understanding of the Great Lakes system and to eliminate or reduce to the maximum extent practicable the discharge of pollutants into the Great Lakes system. The GLNPO also coordinates U.S. actions in fulfillment of the Agreement between Canada and the United States of America on Great Lakes Water Quality of 1978.

This study was carried out under a Cooperative Agreement with U.S. Fish and Wildlife Service, Great Lakes Fishery Laboratory, Ann Arbor, Michigan.

ABSTRACT

Contaminant body burdens in lake trout from the Upper Great Lakes have been monitored since 1970 on Lake Michigan and since 1977 and 1978 on Lakes Superior and Huron by USEPA, Great Lakes National Program Office and USFWS, Great Lakes Fishery Laboratory.

Analysis of the Lake Michigan data shows that mean PCB concentrations declined from a maximum of 22.91 mg/kg in 1974 to 5.63 in 1982. Mean total DDT concentrations declined from 19.19 mg/kg in 1970 to 2.74 mg/kg in 1982. The decline in both contaminants closely followed first order loss kinetics. If the current declines continue, PCB concentrations will decline to the USFDA tolerance of 2.0 mg/kg in 1988. Mean total DDT concentrations will fall to the IJC objective of 1.0 mg/kg by 1991. Mean dieldrin concentrations increased significantly from 0.20 mg/kg in 1971 to 0.58 mg/kg in 1979 before declining to 0.21 mg/kg in 1982. The decline from 1979-1982 followed first order loss kinetics. As this decline is not reflected in other species (bloater chubs, smelt) it will require additional years of monitoring to determine if the decline in dieldrin concentrations between 1979 and 1982 truly represents a declining trend.

Contaminants in lake trout from Lake Superior and Lake Huron generally declined over the study period. The only statistically significant trend other than in Lake Michigan was for total DDT which declined significantly in Lake Superior lake trout. Large data variance and the short time frame covered (1977-1982) interfered with detection of trends on Lakes Superior and Huron.

CONTENTS

Foreword.....ii
Abstract.....iii
Tables and Figures.....v
Introduction.....1
Methods.....2
Results.....8
Discussion and Conclusions.....19
References.....21

Table and Figures

Table 1 - Sample Collection Data.....	6
Table 2 - PCB and Pesticide Detection Limits.....	7
Table 3 - Mean Contaminant Concentrations in Lake Trout from Lake Michigan.....	9
Table 4 - Mean Contaminant Concentrations in Lake Trout from Lake Huron.....	16
Table 5 - Mean Contaminant Concentrations in Lake Trout from Lake Superior.....	17

Figures

Figure 1 - Lake Trout Collection Sites.....	3
Figure 2 - PCB Concentrations in Lake Michigan Lake Trout.....	11
Figure 3 - Total DDT Concentrations in Lake Michigan Lake Trout.....	12
Figure 4 - Dieldrin Concentrations in Lake Michigan Lake Trout.....	13
Figure 5 - Oxychlordanes Concentrations in Lake Michigan Lake Trout.....	14
Figure 6 - Total DDT Concentrations in Lake Superior Lake Trout.....	18

INTRODUCTION

The presence of toxic organic contaminants, many of which bioaccumulate in fish, is one of the most serious environmental problems in the Great Lakes. During the past decade, contaminants such as PCB, DDT and dieldrin have been found in Great Lakes fish at concentrations exceeding U.S. Food and Drug Administration action levels. Bans and restrictions on the use of these compounds have been implemented in an effort to control these problems.

In response to its responsibilities under the U.S. and Canadian Great Lakes Water Quality Agreements of 1972 and 1978, the International Joint Commission (IJC) has assisted in the development and implementation of a coordinated surveillance and monitoring program "to assess compliance with pollution control requirements and achievement of objectives, to provide information for measuring loads and whole lake response to control measures, and to identify emerging problems." As part of the coordinated surveillance program, the U.S. Environmental Protection Agency's, Great Lakes National Program Office (GLNPO) and U.S. Fish and Wildlife Service (USFWS), Great Lakes Fishery Laboratory have cooperated in a program to monitor contaminant trends in fish collected from each of the Great Lakes and Lake St. Clair. In this first report resulting from the Cooperative Program we present the results of the analysis of lake trout (Salvelinus namaycush) collected in Lakes Michigan, Superior, and Huron in 1977-1982.

METHODS

Lake Trout were collected by gillnet from Lakes Michigan, Huron, and Superior (Figure 1) in the fall of the year by USFWS personnel. The fish (60 per collection site) were put into plastic bags, frozen whole, and transported to the USFWS, Great Lakes Fishery Laboratory in Ann Arbor, Michigan where they were sorted into small (300-450 mm), medium (451-650 mm), and large (>650 mm) size categories (20 fish per category) for each collection site and year. Within each size category the fish were grouped into four composite samples consisting of five fish each and homogenized in a Hobart™ vertical chopper/ mixer. Subsamples of the homogenized tissue were weighed and stored frozen at <-30°C in solvent-washed glass jars with foil-lined screw caps until analyzed by the USEPA Central Regional Laboratory (CRL), Chicago, Illinois.

At CRL, the samples were thawed, dried with Na₂SO₄, and lipids and contaminants were column extracted with two 50-ml aliquots of 10% ethyl acetate in petroleum ether (Hesselberg and Johnson 1972). Lipids were separated from the extract by gel permeation chromatography (Stalling et al. 1972). Pesticides and PCB were separated on a column of silica gel (Snyder and Reinert 1971). Final separation and quantitation of pesticides and PCBs was by a Varian 3700 dual column gas chromatograph with Ni⁶³ electron capture detectors (CRL 1982) under the following operating conditions:

	Column A	Column B
Column length	2.4mx2mm i.d.	2.4mx2mm i.d.
Column materials	4% SE-30/6% SP2401 on Supelcoport 100-120 mesh	3% OV-101 on Supelcoport 80-100 mesh
Injection temperature	270°C	270°C
Oven temperature	200°C	200°C
Detector temperature	320°C	320°C
N ₂ carrier gas flow	ca. 30 ml/min	ca 30 ml/min
Range (amps/mv)	10-10 amp	10-10 amp

All results were computed on a mg/kg wet weight basis, and were not corrected for extraction or recovery efficiency.




Figure 1. Lake trout collection sites

As initially designed, the sampling program was intended to compare contaminant levels in fish, both temporally and spatially, in the three size categories of fish collected from each location. The results were to be compared by using analysis of covariance techniques. This approach requires that a significant correlation exist between contaminant concentrations and the size of fish. If differences are to be tested for, the regressions of contaminant levels and fish size must also have similar slopes. The data for 1977-1979 from Lakes Michigan, Huron, and Superior were initially analyzed by using this approach. Since data for three years, 13 organic contaminants and three sites were included in the analysis, 117 regressions between contaminant concentration and fish size were possible. Of these, 63 were significant at the 95% confidence level. Among the significant regressions, several exhibited significant ($p < 0.05$) differences in slopes, either between years or collection sites. Thus, less than half of the available data met the requirements for the test. We therefore, decided that although the analysis of covariance technique may be worthy of further evaluation, the use of mean statistics with specific size ranges of fish was the most appropriate approach for analyzing available data and determining differences in contaminant concentrations between sites and within sites over time.

To achieve a comparable size range of fish from 1977 through 1981 and into future years (current collections consist of fish 600-700 mm long) we used the following procedure. The mean length of all medium and large size group composite samples from each site and year was calculated. When this mean was below 620 mm or above 640 mm, the composite sample with the smallest or largest length was dropped and the mean recalculated. This procedure was repeated until we arrived at a mean length within the 620 mm to 640 mm range. The composite samples

thus selected were then treated as environmental replicates and mean statistics were used to compare contaminant concentrations. Table 1 presents the fish size and lipid data for the samples thus developed.

For purposes of statistical analyses and graphical display, we calculated concentrations of contaminants detected, but below quantitation limits, as 1/2 of the limits given in Table 2.

TABLE 1

SAMPLE COLLECTION DATA

Lake and Year	N	Length (mm) Mean SE	Weight (g) Mean SE	Lipid % Mean SE
Lake Michigan*				
Saugatuck				
1970	18	613 (9)	2084 (149)	25.63 (1.69)
1971	20	579 (8)	2314 (125)	18.03 (0.68)
1972	9	648 (15)	2576 (178)	18.52 (1.81)
1973	30	602 (10)	2353 (120)	16.00 (0.70)
1974	30	616 (10)	2516 (130)	16.50 (0.56)
1975	29	613 (8)	2581 (92)	19.10 (0.55)
1976	30	606 (9)	2337 (103)	16.80 (0.49)
1977	3	636 (53)	2888 (681)	16.59 (0.88)
1978	8	631 (27)	2447 (311)	17.50 (0.76)
1979	6	630 (40)	2844 (461)	16.78 (1.10)
1980	6	637 (18)	2843 (296)	17.68 (0.74)
1981	3	620 (37)	2646 (497)	17.96 (0.88)
1982	10	612 (4)	2466 (47)	16.53 (0.33)
Lake Huron				
Rockport				
1977			NO SAMPLE	
1978	6	637 (18)	2522 (210)	19.19 (0.65)
1979	5	640 (21)	3017 (308)	21.30 (0.78)
1980	6	633 (18)	2674 (230)	19.03 (1.10)
1981	6	628 (28)	2775 (343)	18.88 (0.64)
1982	10	623 (7)	2527 (84)	18.10 (0.71)
Lake Superior				
Apostle Island				
1977	7	640 (27)	2927 (425)	16.46 (1.80)
1978	7	638 (27)	3080 (400)	21.14 (1.20)
1979	6	630 (24)	2797 (312)	21.80 (1.80)
1980	7	636 (24)	2868 (360)	15.10 (1.30)
1981	5	634 (29)	2762 (386)	15.92 (0.93)
1982	10	619 (7)	2479 (103)	15.67 (0.43)

* Data for 1970-76 from the U.S. Fish and Wildlife Service, Great Lakes Fishery Laboratory, collected before the Cooperative Program began, and are based on analyses of individual, whole fish.

Table 2
PCB and Pesticide Detection Limits

Compound	Quantitation Limits mg/kg
o,p'-DDE	0.002
p,p'-DDE	0.002
o,p'-DDD	0.005
p,p'-DDD	0.005
o,p'-DDT	0.020
p,p'-DDT	0.002
Dieldrin	0.002
Total Aroclor (as 1254)	0.030 - 0.050
Heptachlor	0.001
Oxychlorane	0.005

RESULTS

PCBs were the dominate contaminant at all sites reflecting its once wide spread use (Durfee 1976), environmental persistence, and tendency to bioaccumulate (EPA 1983). The continued presence of DDT residues illustrates the duration of contaminant problems long after measures to ban manufacture and use are inacted. Lake Michigan lake trout generally contained the highest contaminant concentrations and Lake Superior the lowest. Clark et al. (1984) found a similar pattern in the relative contaminant levels in coho salmon. Studies by Frank et al. (1980) and Frank et al. (1981) found the sediment concentrations of PCB, chlordane plus heptachlor and DDT to be similar in Lake Michigan and Huron and substantially lower in Lake Superior

Lake Michigan

Table 3 presents the annual means and 95 percent confidence intervals for contaminants in lake trout off saugatuck, Michigan in 1970-1982. Before 1977 individual fish were analyzed by the Great Lakes Fishery Laboratory by the analytical methodology described in Willford et al. (1976). Since 1977 five fish composite samples were analyzed by USEPA.

Mean PCB concentrations (mg/kg) increased from 12.86 in 1972 to 22.91 in 1974, and the declined to 6.49 in 1981.

Mean PCB concentrations from 1974 - 1981 declined, generally following first order loss rate kinetics in the form of the following equation:

$$C_t = C_0 e^{-kT} \quad (1)$$

where C_t = concentration at time, C_0 = initial concentration, K = rate constant, and T = Time

Table 3. Mean Contaminant Concentrations in lake trout from Lake Michigan off Saugatuck, mg/kg wet weight (95% confidence intervals in parenthesis)

Year	PCB*	DDT	Oxychlorane	Dieldrin	Number of Samples
1970	NA	19.19 (15.90, 22.46)	NA	0.27 (0.22, 0.32)	18
1971	NA	13.00 (11.24, 14.76)	NA	0.20 (0.17, 0.23)	20
1972	12.86 (8.11, 17.61)	11.31 (8.05, 14.57)	NA	0.20 (0.14, 0.26)	9
1973	18.93 (16.85, 21.01)	9.96 (8.60, 11.32)	NA	0.27 (0.24, 0.30)	30
1974	22.91 (19.18, 26.64)	8.42 (6.68, 10.16)	NA	0.30 (0.27, 0.33)	30
1975	22.28 (19.38, 25.18)	7.50 (6.25, 8.75)	NA	0.35 (0.32, 0.38)	29
1976	18.68 (16.02, 21.34)	5.65 (4.61, 6.69)	NA	0.30 (0.28, 0.32)	30
1977	11.58 (-2.30, 25.4)	6.34 (-0.40, 13.10)	0.230 (-0.71, 0.531)	0.40 (0.35, 0.45)	3
1978	8.18 (5.50, 10.90)	4.58 (2.32, 6.83)	0.175 (0.072, 0.278)	0.44 (0.28, 0.61)	8
1979	8.82 (5.20, 12.50)	6.91 (2.80, 11.00)	0.240 (0.136, 0.344)	0.58 (0.25, 0.91)	6
1980	9.93 (2.00, 17.80)	4.74 (1.90, 7.50)	0.160 (0.039, 0.280)	0.335 (0.20, 0.47)	6
1981	6.49 (4.51, 8.46)	3.22 (2.11, 4.34)	0.061 (0.049, 0.073)	0.26 (0.24, 0.28)	3
1982	5.63 (5.36, 5.89)	2.74 (2.34, 3.14)	0.075 (0.057, 0.093)	0.21 (0.16, 0.28)	10

NA = Not Analyzed

1977-1982 and beginning in 1977 only Aroclor 1254.

Figure 2 illustrates the loss trajectory based on the least squares fit ($r^2 = 89.6$, $P < 0.01$) to the mean yearly PCB concentrations. The rate constant is $-0.18/\text{yr}$.

Mean total DDT concentrations (mg/kg) declined from 19.19 in 1970 to 2.74 in 1982. The observed decline in mean total DDT concentrations also followed first order loss rate kinetics (equation 1). Figure 3 illustrates the loss trajectory from 1970 to 1982. The rate constant is $-0.13/\text{yr}$ with $r^2 = 90.9$ ($P < 0.01$).

Mean dieldrin concentrations (mg/kg) decreased from 0.27 in 1970 to 0.20 in 1971, then increased to 0.58 in 1979 after which concentrations declined to 0.22 in 1982. The increase in mean concentrations between 1972 and 1979 was highly significant ($P < 0.01$, $R^2 = 86.6$) following the linear equation:

$$C_t = 0.045 (Y) - 3.02 \quad (2)$$

Where C_t = mean dieldrin concentration (mg/kg) in year t and Y = year (1900's) from 72-79

From 1979 to 1982 mean concentrations declined, following first order loss kinetics (equation 1), with a loss rate of $-0.32/\text{yr}$. The curve on Figure 4 illustrates the loss trajectory based on a least squares fit to the data ($r^2 = 93.2$, $P < 0.05$).

The chlordane metabolite, oxychlordane, increased from 1977 through 1979 and decreased from 1979 through 1982. The decrease (Figure 5) in mean concentration from 1979 to 1982 was significant ($P < 0.05$, $R^2 = 85.1$) following the linear equation.

$$C_t = 0.059 (y) + 4.92 \quad (3)$$

Where C_t = mean oxychlordane concentration (mg/kg) in year t and y = year (1900's) from 79 to 82.




Figure 2. PCB Trends in Lake Michigan Lake Trout




Figure 3. Total DDT trends in Lake Michigan lake trout.
12




Figure 4. Dieldrin trends in Lake Michigan lake trout.




Figure 5. Oxychlordane trends in Lake Michigan lake trout.

Lake Huron

Unlike Lake Michigan, where comparable trend monitoring began as early as 1970, the Lake Huron data base began in 1978 (Table 4). Despite general declines in the mean concentrations of PCB and DDT from 1979 to 1982 no statistically significant ($P < 0.05$) trends were observed for any of the monitored compounds. Annual mean total DDT and PCB concentrations exceeded the IJC (1978) objectives (1.0 mg/kg and 0.1 mg/kg, respectively) in each year of the study.

Lake Superior

Contaminant concentrations in Lake Superior lake trout are given in Table 5. The relatively low levels of PCB, dieldrin, and oxychlordan did not exhibit trends nor statistically significant ($P < 0.05$) differences over the study period.

Total DDT concentrations in Lake Superior lake trout (Figure 6) declined significantly ($P < 0.05$, $r^2 = 78.9$) from 1977 to 1982, following the equation:

$$C_t = -0.162 (y) + 13.7 \quad (4)$$

Where C_t = mean total DDT concentration (mg/kg) in year t and y = year (1900s) from 77 to 82.

The annual mean PCB concentration exceeded the IJC (1978) objective in each year of the study. The IJC (1978) total DDT objective was exceeded by the mean concentrations in 1977 through 1979, but not from 1980 to 1982.

Table 4
 Results of Contaminant Monitoring of Lake Trout
 From Rockport, Lake Huron
 Mean (95% Confidence Interval)
 All data mg/kg wet weight

Year	PCB*	DDT	Oxychlordan	Dieldrin	Number of Samples
1978	2.92 (2.32, 3.52)	2.19 (2.05, 2.33)	0.038 (0.033, 0.043)	0.16 (0.14, 0.17)	6
1979	3.66 (1.57, 5.75)	2.85 (1.94, 3.76)	0.060 (0.024, 0.097)	0.19 (0.14, 0.24)	5
1980	3.44 (2.46, 4.42)	1.53 (1.10, 1.96)	0.056 (0.039, 0.073)	0.13 (0.10, 0.15)	6
1981	3.51 (2.31, 4.72)	1.79 (1.15, 2.43)	0.058 (0.040, 0.076)	0.14 (0.12, 0.16)	6
1982	2.10 (1.72, 2.49)	1.31 (1.13, 1.49)	0.059 (0.044, 0.074)	0.15 (0.11, 0.18)	10

*PCB quantitated as Aroclor 1254

Table 5
 Results of Contaminant Monitoring of Lake Trout
 From the Apostle Island, Lake Superior
 Mean (95% Confidence Interval)
 All Data mg/kg Wet Weight

Year	PCB*	DDT	Oxychlordan	Dieldrin	Number of Samples
1977	1.87 (0.95, 2.80)	1.2 (0.76, 1.80)	0.120 (0.059, 0.180)	0.05 (0.03, 0.06)	7
1978	0.93 (0.65, 1.21)	1.02 (0.66, 1.39)	0.402 (0.110, 0.700)	0.06 (0.05, 0.07)	7
1979	0.88 (0.59, 1.17)	1.095 (0.61, 1.57)	0.136 (0.102, 0.375)	0.07 (0.05, 0.08)	6
1980	1.89 (0.41, 3.36)	0.62 (0.13, 1.11)	0.026 (0.018, 0.035)	0.045 (0.02, 0.06)	7
1981	1.40 (0.37, 2.44)	0.81 (-0.09, 1.71)	0.056 (0.001, 0.111)	0.05 (0.03, 0.07)	5
1982	0.48 (0.36, 0.60)	0.288 (0.214, 0.47)	0.041 (0.036, 0.047)	0.05 (0.04, 0.05)	10

*PCB quantitated as Aroclor 1254




Figure 6. Total DDT trends in Lake Superior lake trout.

Discussion and Conclusions

Of the data sets for the three upper lake monitoring sites reported here, only that for Lake Michigan, was sufficient to allow rigorous statistical evaluation of trends with time. This is principally because FWS had collected data for 6 years before the current U.S. EPA/U.S. FWS cooperative program was begun in 1977. As a result of these extra years of data, we were able to demonstrate that mean PCB and mean total DDT concentrations in lake trout off Saugatuk declined in a manner that approximated first order loss kinetics. Rodgers and Swain (1983) reported that the decline of PCB concentrations in bloaters (*Coregonus hoyi*) from Lake Michigan (off Saugatuck) from 1972 to 1980 also approximated first order loss kinetics with a rate constant of $-0.12/\text{yr}$. They used the trend data for bloaters to hindcast loads and then used their loading estimates to forecast concentrations in other fish species. They estimated that PCB concentrations in Lake Michigan trout would reach 2 mg/kg in 1987. By solving equation (1) for t , we calculate that PCB concentrations will drop to 2 mg/kg in 1988. This estimate is in excellent agreement with the projections of Rodgers and Swain (1983), particularly when the difference in approach and the uncertainties of such projections are considered. These projections presuppose that PCB loading into Lake Michigan will continue to decrease at a rate at least equal to that before 1982. As atmospheric deposition now contributes will over 50 percent of the total load (Murphy and Rzesutko 1977), this decrease may not be affected.

A similar projection of total DDT concentration suggests that mean concentrations will reach the IJC objective of 1.0 mg/kg in about 1991.

Despite cancellation of most uses of aldrin and dieldrin in 1974 (EPA 1980), mean concentrations in lake trout in Lake Michigan off Saugatuck increased from 0.27 mg/kg in 1970 to 0.58 mg/kg in 1979. From 1979 to 1982 mean dieldrin concentrations declined in a manner approximating first order loss kinetics. However, other species such as bloater chubs and rainbow smelt have not exhibited the decline observed in lake trout. Dieldrin concentrations in bloaters in Lake Michigan increased from a mean of 0.27 (mg/kg) in 1970 to 0.43 in 1982 (Willford 1982), while rainbow smelt remained relatively constant (range 0.046 - 0.071 mg/kg) from 1977 to 1982 (GLNPO-unpublished data). Elevated dieldrin concentrations resulted in seizure of commercial catches of Lake Michigan bloaters by the U.S. Food and Drug Administration in 1984 (David Crockett, U.S. Food and Drug Administration, Chicago, Illinois, personal communications, 1984). Therefore, while the apparent decline in lake trout is encouraging, it will require additional years of monitoring to test this trend.

With the exception of DDT which declined significantly in Lake Superior lake trout from 1977 to 1982, no significant changes in contaminant concentrations were observed in lake trout from Lake Huron or Lake Superior. The combination of limited data sets, large data variance, and relatively lower contaminant concentrations interfered with the detection of trends. The ineffectiveness of the study design used from 1977 to 1981 (sampling three size ranges of fish), and the resulting requirements to select fish over a large size range and greatly reduce the number of yearly samples, was partly responsible for the large variance associated with the 1977 to 1981 data. Although the covariance approach will be further evaluated, the monitoring protocol used since 1982 for this cooperative program utilizes mean statistics on results from 10 composite samples of 5 fish each, all of which are 600 to 700 mm long. The reduced variance under this more stringent sampling protocol will greatly enhance the likelihood of detecting significant differences between any two sites or years when contaminant concentrations are changing.

References

- Clark, J.R., D.DeVault, R.J. Bowden, and J.A. Weishaar. 1984. Contaminant analysis of fillets from Great Lakes coho salmon, 1980. *J. Great Lakes Res.* 10:38-47.
- Central Regional Laboratory (CRL). 1982. Standard operating procedure for the analysis of contaminants in fish for the open lake program Great Lakes fish contaminant dynamics project. USEPA. Chicago, Illinois.
- DeVault, D.S., and J.A. Weishaar. 1984. Contaminant analysis of 1982 fall run coho salmon. U.S. Environmental Protection Agency, Great Lakes National Program Office. EPA 905/3-84-004.
- Durfee, R.L. 1976. Production and usage of PCBs in the United States. In: *Proceedings of the National Conference on Polychlorinated Biphenyls*. U.S. Environmental Protection Agency, Office of Toxic Substances, Washington, D.C. pp 103-107, EPA-560/6-75-004.
- EPA. 1980. Ambient water quality criteria for dieldrin. U.S. Environmental Protection Agency, Office of Water Regulations and Standards, Washington, D.C. EPA 440/5-80-019.
- EPA. 1983. Environmental transport and transformation of polychlorinated biphenyls. U.S. Environmental Protection Agency, Office of Pesticides and Toxic Substances, Washington, D.C. EPA 560/5-83-025.
- Frank R., R.L. Thomas, H.E. Braun, J.Rasper, and R. Dawson. 1980. Organochlorine insecticides and PCB in surficial sediments of Lake Superior (1973). *J. Great Lakes Res.* 6:113-120.
- Frank R., R.L. Thomas, H.E. Braun, D.L. Gross and T.T. Davies. 1981. Organochlorine insecticides and PCB in surficial sediments of Lake Michigan (1975). *J. Great Lakes Res.* 7:42-50.
- Hesselberg R.J., and J.L. Johnson. 1972. Column extraction of pesticides from fish, fish food, and mud. *Bull. Environ. Contam Toxicol.* 7:115-120.
- International Joint Commission. 1978. Status Report on persistent-toxic pollutants in the Lake Ontario Basin. Great Lakes Water Quality Board, Windsor, Ontario, Canada.
- Murphy, T.J. and C.P. Rzeszutko. 1977. Precipitation inputs of PCBs to Lake Michigan. *J. Great Lakes Res.* 3:305-312. *
- Rodgers, P.W., and W.R. Swain. 1983. Analysis of polychlorinated biphenyl (PCB) loading trends in Lake Michigan. *J. Great Lakes Res.* 9:548-558.

- Snyder, D. and R. Reinert. 1971. Rapid separation of polychlorinated biphenyls from DDT and its analogues on silica gel. Bull. Environ. Contam. and Toxicol. 6:385-390.
- Stalling, D.L., R.C. Tindle, and J.L. Johnson. 1972. Cleanup of pesticide and polychlorinated biphenyl residues in fish extracts by gel permeation chromatography J. Assoc. Off. Anal. Chem. 55:32-38.
- Willford, A., J. Hesselberg, and W. Nicholson. 1976. Trends of polychlorinated biphenyls in three Lake Michigan fishes. In: Proceedings of the National Conference on Polychlorinated Biphenyls. U.S. Environmental Protection Agency, Office of Toxic Substances, Washington, D.C. pp 177-181, EPA-560/6-75-004.
- Willford, A. 1982. Evaluating the presence and effects of contaminants in fish of the Great Lakes. Minutes of the Great Lakes Fishery Commission Annual Meeting, Green Bay, Wisconsin, June 9-10, 1982, Agenda Item 14.C., Appendix XXVI:273-284.
- Zar, J. 1974. Biostatistical Analysis. Prentice Hall. Inc., Englewood Cliffs, N.J.

TECHNICAL REPORT DATA (Please read Instructions on the reverse before completing)		
1. REPORT NO. EPA-905/3-85-001	2.	3. RECIPIENT'S ACCESSION NO.
4. TITLE AND SUBTITLE Contaminant Trends In Lake Trout (Salvelinus namaycush) of the Upper Great Lakes	5. REPORT DATE April 1985	6. PERFORMING ORGANIZATION CODE 5GL
	8. PERFORMING ORGANIZATION REPORT NO.	
7. AUTHOR(S) David S. DeVault	10. PROGRAM ELEMENT NO.	
9. PERFORMING ORGANIZATION NAME AND ADDRESS U.S. Fish and Wildlife Service Great Lakes Fishery Laboratory 1451 Green Road Ann Arbor, Michigan 48105	11. CONTRACT/GRANT NO.	
	13. TYPE OF REPORT AND PERIOD COVERED Fish Monitoring 1977-1982	
12. SPONSORING AGENCY NAME AND ADDRESS U.S. Environmental Protection Agency Great Lakes National Program Office 536 South Clark Street, Room 958 Chicago, Illinois 60605	14. SPONSORING AGENCY CODE Great Lakes National Program Office-USEPA, Region V	
	15. SUPPLEMENTARY NOTES	
16. ABSTRACT Contaminant body burdens in lake trout from the Upper Great Lakes have been monitored since 1970 on Lake Michigan and since 1977 and 1978 on Lakes Superior and Huron by USEPA, Great Lakes National Program Office and USFWS, Great Lakes Fishery Laboratory. Analysis of the Lake Michigan data shows that mean PCB concentrations declined from maximum of 22.91 mg/kg in 1974 to 5.63 in 1982. Mean total DDT concentrations declined from 19.19 mg/kg in 1970 to 2.74 mg/kg in 1982. The decline in both contaminants closely followed first order loss kinetics. If the current declines continue, PCB concentrations will decline to the USFDA tolerance of 2.0 mg/kg in 1988. Mean total DDT concentrations will fall to the IJC objective of 1.0 mg/kg by 1991. Mean dieldrin concentrations increased significantly from 0.20 mg/kg in 1971 to 0.58 mg/kg in 1979 before declining to 0.21 mg/kg in 1982. The decline from 1979-1982 followed first order loss kinetics. As this decline is not reflected in other species (bloaters, chubs, smelt) it will require additional years of monitoring to determine if the decline in dieldrin concentrations between 1979 and 1982 truly represents a declining trend.		
17. KEY WORDS AND DOCUMENT ANALYSIS		
a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
Contaminants Monitoring Toxic organic Pesticides Lipids Lake Trout		
18. DISTRIBUTION STATEMENT Document is available to the public through the National Technical Information Service, Springfield, VA 22151	19. SECURITY CLASS (This Report)	21. NO. OF PAGES 30
	20. SECURITY CLASS (This page)	22. PRICE