

Environmental Protection  
Agency

Office of Emergency and  
Remedial Response  
Washington DC 20460

Office of Research and Development  
Office of Health and Environmental  
Assessment  
Environmental Criteria and  
Assessment Office  
Cincinnati OH 45268


---

Superfund

---



HEALTH EFFECTS ASSESSMENT  
FOR DDT



HEALTH EFFECTS ASSESSMENT  
FOR DDT

U.S. Environmental Protection Agency  
Office of Research and Development  
Office of Health and Environmental Assessment  
Environmental Criteria and Assessment Office  
Cincinnati, OH 45268

U.S. Environmental Protection Agency  
Office of Emergency and Remedial Response  
Office of Solid Waste and Emergency Response  
Washington, DC 20460

## DISCLAIMER

This report has been funded wholly or in part by the United States Environmental Protection Agency under Contract No. 68-03-3112 to Syracuse Research Corporation. It has been subject to the Agency's peer and administrative review, and it has been approved for publication as an EPA document. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

## PREFACE

This report summarizes and evaluates information relevant to a preliminary interim assessment of adverse health effects associated with DDT. All estimates of acceptable intakes and carcinogenic potency presented in this document should be considered as preliminary and reflect limited resources allocated to this project. Pertinent toxicologic and environmental data were located through on-line literature searches of the Chemical Abstracts, TOXLINE, CANCERLINE and the CHEMFATE/DATALOG data bases. The basic literature searched supporting this document is current up to September, 1984. Secondary sources of information have also been relied upon in the preparation of this report and represent large-scale health assessment efforts that entail extensive peer and Agency review. The following Office of Health and Environmental Assessment (OHEA) sources have been extensively utilized:

U.S. EPA. 1980a. Hazard Assessment Report on DDT, DDD, DDE. Environmental Criteria and Assessment Office, Cincinnati, OH. Internal report.

U.S. EPA. 1980b. Ambient Water Quality Criteria for DDT. Environmental Criteria and Assessment Office, Cincinnati, OH. EPA 440/5-80-038. NTIS PB 81-117491.

U.S. EPA. 1985. The Carcinogen Assessment Group's Evaluation of the Carcinogenicity of Dicofof (Kelthane), DDT, DDE and DDD (TDE). Prepared by the Carcinogen Assessment Group, OHEA, Washington, DC for the Hazard Evaluation Division Office of Pesticide Programs. EPA 600/6-85-002X. Internal Review Draft.

The intent in these assessments is to suggest acceptable exposure levels whenever sufficient data were available. Values were not derived or larger uncertainty factors were employed when the variable data were limited in scope tending to generate conservative (i.e., protective) estimates. Nevertheless, the interim values presented reflect the relative degree of hazard associated with exposure or risk to the chemical(s) addressed.

Whenever possible, two categories of values have been estimated for systemic toxicants (toxicants for which cancer is not the endpoint of concern). The first, the AIS or acceptable intake subchronic, is an estimate of an exposure level that would not be expected to cause adverse effects when exposure occurs during a limited time interval (i.e., for an interval that does not constitute a significant portion of the lifespan). This type of exposure estimate has not been extensively used or rigorously defined, as previous risk assessment efforts have been primarily directed towards exposures from toxicants in ambient air or water where lifetime exposure is assumed. Animal data used for AIS estimates generally include exposures with durations of 30-90 days. Subchronic human data are rarely available. Reported exposures are usually from chronic occupational exposure situations or from reports of acute accidental exposure.

The AIC, acceptable intake chronic, is similar in concept to the ADI (acceptable daily intake). It is an estimate of an exposure level that would not be expected to cause adverse effects when exposure occurs for a significant portion of the lifespan [see U.S. EPA (1980c) for a discussion of this concept]. The AIC is route specific and estimates acceptable exposure for a given route with the implicit assumption that exposure by other routes is insignificant.

Composite scores (CSs) for noncarcinogens have also been calculated where data permitted. These values are used for ranking reportable quantities; the methodology for their development is explained in U.S. EPA (1983).

For compounds for which there is sufficient evidence of carcinogenicity, AIS and AIC values are not derived. For a discussion of risk assessment methodology for carcinogens refer to U.S. EPA (1980c). Since cancer is a process that is not characterized by a threshold, any exposure contributes an increment of risk. Consequently, derivation of AIS and AIC values would be inappropriate. For carcinogens, q<sub>1</sub>\*s have been computed based on oral and inhalation data if available.

## ABSTRACT

In order to place the risk assessment evaluation in proper context, refer to the preface of this document. The preface outlines limitations applicable to all documents of this series as well as the appropriate interpretation and use of the quantitative estimates presented.

U.S. EPA (1985) estimated a human  $q_1^*$  of  $0.34 \text{ (mg/kg/day)}^{-1}$  for orally administered DDT based on the geometric average of potencies from six separate studies.

Data are not available for the evaluation of the potential carcinogenicity of DDT following inhalation exposure.

## ACKNOWLEDGEMENTS

The initial draft of this report was prepared by Syracuse Research Corporation under Contract No. 68-03-3112 for EPA's Environmental Criteria and Assessment Office, Cincinnati, OH. Dr. Christopher DeRosa and Karen Blackburn were the Technical Project Monitors and Helen Ball was the Project Officer. The final documents in this series were prepared for the Office of Emergency and Remedial Response, Washington, DC.

Scientists from the following U.S. EPA offices provided review comments for this document series:

Environmental Criteria and Assessment Office, Cincinnati, OH  
Carcinogen Assessment Group  
Office of Air Quality Planning and Standards  
Office of Solid Waste  
Office of Toxic Substances  
Office of Drinking Water

Editorial review for the document series was provided by:

Judith Olsen and Erma Durden  
Environmental Criteria and Assessment Office  
Cincinnati, OH

Technical support services for the document series was provided by:

Bette Zwyer, Pat Daunt, Karen Mann and Jacky Bohanon  
Environmental Criteria and Assessment Office  
Cincinnati, OH

# TABLE OF CONTENTS

	<u>Page</u>
1. ENVIRONMENTAL CHEMISTRY AND FATE. . . . .	1
2. ABSORPTION FACTORS IN HUMANS AND EXPERIMENTAL ANIMALS . . . . .	2
2.1. ORAL . . . . .	2
2.2. INHALATION . . . . .	2
3. TOXICITY IN HUMANS AND EXPERIMENTAL ANIMALS . . . . .	3
3.1. SUBCHRONIC . . . . .	3
3.1.1. Oral. . . . .	3
3.1.2. Inhalation. . . . .	3
3.2. CHRONIC. . . . .	3
3.2.1. Oral. . . . .	3
3.2.2. Inhalation. . . . .	7
3.3. TERATOGENICITY AND OTHER REPRODUCTIVE EFFECTS. . . . .	7
3.3.1. Oral. . . . .	7
3.3.2. Inhalation. . . . .	8
3.4. TOXICANT INTERACTIONS. . . . .	8
4. CARCINOGENICITY . . . . .	11
4.1. HUMAN DATA . . . . .	11
4.1.1. Oral. . . . .	11
4.1.2. Inhalation. . . . .	11
4.2. BIOASSAYS. . . . .	11
4.2.1. Oral. . . . .	11
4.2.2. Inhalation. . . . .	17
4.3. OTHER RELEVANT DATA. . . . .	17
4.4. WEIGHT OF EVIDENCE . . . . .	18
5. REGULATORY STANDARDS AND CRITERIA . . . . .	19



# TABLE OF CONTENTS (cont.)

	<u>Page</u>
6. RISK ASSESSMENT . . . . .	20
6.1. ACCEPTABLE INTAKE SUBCHRONIC (AIS) . . . . .	20
6.2. ACCEPTABLE INTAKE CHRONIC (AIC). . . . .	20
6.3. CARCINOGENIC POTENCY (q <sub>1</sub> *) . . . . .	20
6.3.1. Oral. . . . .	20
6.3.2. Inhalation. . . . .	26
7. REFERENCES. . . . .	31
APPENDIX: Summary Table for DDT. . . . .	43

# LIST OF TABLES

<u>No.</u>	<u>Title</u>	<u>Page</u>
3-1	Subchronic Oral Toxicity of DDT in Rats . . . . .	4
3-2	Chronic Oral Toxicity of DDT. . . . .	5
3-3	Reproductive Studies. . . . .	9
4-1	Oral Carcinogenicity of DDT . . . . .	12
6-1	Incidence of the Most Commonly Occurring Malignant Tumors in Each of Five Generations of BALB/c Mice Fed DDT. . . . .	21
6-2	Incidence of Benign Liver Tumors in Each of Six Generations of CF-1 Mice Fed DDT. . . . .	23
6-3	Incidence of Benign Liver Tumors in BALB/C Mice Fed DDT During a 2-Generation Experiment. . . . .	25
6-4	Incidence of Liver Tumors (Benign and Malignant) in CF-1 Mice Fed DDT for a Single Generation. . . . .	27
6-5	Incidence of Benign Liver Tumors in CF-1 Mice Fed DDT for 15 or 30 Weeks and then Sacrificed at 65, 95 and 120 Weeks. . . . .	28
6-6	Incidence of Benign Liver Tumors in Rats Fed DDT. . . . .	29
6-7	Summary of Quantitative Potency Estimates for DDT . . . . .	30

## LIST OF ABBREVIATIONS

ADI	Acceptable daily intake
AIC	Acceptable intake chronic
AIS	Acceptable intake subchronic
BCF	Bioconcentration factor
bw	Body weight
CAS	Chemical Abstract Service
CS	Composite score
ppm	Parts per million
SER	Smooth endoplasmic reticulum
STEL	Short-term exposure limit
TLV	Threshold limit value
TWA	Time-weighted average

## 1. ENVIRONMENTAL CHEMISTRY AND FATE

The physical and chemical properties and environmental fate of DDT (CAS No. 50-29-3 for p,p'-DDT and 789-02-6 for o,p'-DDT) are as follows:

Chemical class:	pesticide (Callahan et al., 1979)
Molecular weight:	354.5 (Callahan et al., 1979)
Vapor pressure at 20°C:	1.5x10 <sup>-7</sup> mm Hg for p,p'-DDT 5.5x10 <sup>-6</sup> mm Hg for o,p'-DDT (Callahan, et al., 1979)
Water solubility at 25°C:	1.2-25 µg/l for p,p'-DDT 26-85 µg/l for o,p'-DDT (Callahan et al., 1979)
Log octanol/water partition coefficient:	3.98-6.19 (Callahan et al., 1979)
BCF:	10 <sup>3</sup> to 10 <sup>6</sup> (Callahan et al., 1979)
Half-life in Water:	56-110 days in lake water (Zoeteman et al., 1980)
Soil:	3-15 years (IARC, 1974)

The mobility of DDT in soils has been studied by various authors (U.S. EPA, 1980a) and has been reported to be extremely slow. Therefore, the leaching of DDT from soil is expected to be very slow, particularly in soil with high organic carbon content. Nevertheless, leaching of DDT from soil into groundwater at a frequency of 8-9% has been reported (Page, 1981).

The half-life of DDT in the atmosphere is uncertain. DDT is probably lost from the atmosphere by rainout, fallout or photochemical degradation (Spencer, 1975). The photolytic half-life for the transformation of DDT to DDE and DDD in the atmosphere can be estimated to be ~17 days (Crosby and Moilanen, 1977). The fact that DDT has been found to participate in long distance aerial transport (Callahan et al., 1979) indicates that it has a long half-life in the atmosphere.

## 2. ABSORPTION FACTORS IN HUMANS AND EXPERIMENTAL ANIMALS

### 2.1. ORAL

Jensen et al. (1957) reported that 95% of the ingested DDT in rats is absorbed from the gastrointestinal tract. Over 65% of the labeled DDT was recovered in the bile collected from rats for 9 days following DDT ingestion (Jensen et al., 1957). More DDT is absorbed in rats when it is dissolved in digestible vegetable oils than when it is suspended in water or dissolved in indigestible mineral oils (Keller and Yeary, 1980).

In humans, the absorption of DDT follows the same pattern as the absorption of dietary fat, that is, absorption is relatively slow; however, absorption appears to be complete after 24 hours (Morgan and Roan, 1977).

### 2.2. INHALATION

Pertinent data regarding the absorption of DDT by inhalation could not be located in the available literature.

### 3. TOXICITY IN HUMANS AND EXPERIMENTAL ANIMALS

#### 3.1. SUBCHRONIC

3.1.1. Oral. Laug et al. (1950) exposed rats to different levels of DDT (1-50 ppm) in the food and examined the histology of the liver 15-27 weeks after the start of treatment. No effects were reported at an exposure of 1 ppm; however, at all the higher exposure levels (5-50 ppm), hepatic cell hypertrophy appeared to be dose-related and minimal alterations in liver histology were reported at an exposure of 5 ppm (Laug et al., 1950). Hart and Fouts (1965) reported increased liver microsomal enzyme activity in rats at 5 and 50 ppm. Ortega et al. (1956a,b) reported ultrastructural changes at 5 and 15 ppm in livers of male rats (Table 3-1).

3.1.2. Inhalation. Pertinent data regarding the inhalation of DDT could not be located in the available literature.

#### 3.2. CHRONIC

3.2.1. Oral. Exposure of rats to 200 ppm (10 mg/kg/day) DDT in the food for 2 years resulted in increased liver weight in the females (Fitzhugh and Nelson, 1947; Fitzhugh, 1948). At higher doses (400-800 ppm), the liver alterations became more severe and were present in both sexes. Liver lesions were present at all dose levels tested, the lowest of which was 10 ppm (0.5 mg/kg/day) (Fitzhugh and Nelson, 1947; Fitzhugh, 1948) (Table 3-2). In chronic exposure studies with DDT, effects appeared to be most severe in offspring exposed to DDT for their entire lifespan, especially during in utero development and nursing. This is demonstrated in the study of Treon and Cleveland (1955) in which male rats exposed to 25 ppm (1.25 mg/kg/day) DDT in the food for 2 years and 3 generations had increased liver weights. In mice, lactation was decreased and the mortality of the offspring increased at an exposure level of 100 ppm (13.0 mg/kg/day) for 6 generations (Keplinger et al., 1968). The incidence of tumors was increased at exposure levels of 50

TABLE 3-1

## Subchronic Oral Toxicity of DDT in Rats

Isomer	Dose*		Duration of Exposure	Sex	Number Used	Effects	Reference
	ppm	mg/kg/day					
NR	50	2.5	15-27 weeks	M/F	75/75	Hepatic cell hypertrophy was definitely present at dose levels of 50 and 10 ppm, and minimally present at 5 ppm. No effects were reported at 1 ppm.	Laug et al., 1950
	10	0.5					
	5	0.25					
	1	0.05					
NR	350	17.0	33-60 weeks	NR	NR	No histopathologic alteration in the exposed as compared with the control group.	Cameron and Cheng, 1951
NR	50	2.5	3 months	NR	NR	Liver microsomal enzyme activity was increased at both exposure levels.	Hart and Fouts, 1965
	5	0.25					
NR	15	0.75	2-18 months	NR	NR	Males had histopathologic alterations (proliferation of SER and concentric membrane arrays) in the liver at both exposure levels.	Ortega et al., 1956a,b
	5	0.25					
NR	0.2	0.01	1-13 weeks	NR	NR	Liver microsomal enzyme activity was not increased.	Kinoshita et al., 1966

\*The fraction of body weight consumed per day in food, called the food factor, is 0.13 for mice and 0.05 for rats. Exposures given in ppm of food are converted to mg/kg/day by multiplying the ppm exposure level by the food factor.

NR = Not reported

TABLE 3-2

## Chronic Oral Toxicity of DDT

Isomer <sup>a</sup>	Dose <sup>b</sup>		Duration of Exposure	Species	Sex	Number Used	Effects	Reference
	ppm	mg/kg/day						
Technical	800	40	2 years	rat/ Osborne- Mendel	M/F	24/24	At 800 ppm there were increased mortality and liver lesions. Increased liver weight occurred at 200 ppm for females and at higher doses for both males and females. No second generation offspring survived in animals exposed to 600 ppm. A reduction in survival of preweanling rats occurred at DDT levels of >50 ppm. Liver lesions were present at all dose levels, but no reproductive effects were reported at exposures of 10 ppm.	Fitzhugh and Nelson, 1947; Fitzhugh, 1948
	600	30			M/F	24/24		
	400	20			M/F	24/12		
	200	10			M/F	24/12		
	100	5			M	12		
	50	2.5			NR	NR		
	10	0.5			NR	NR		
	0				M/F	24/12		
NR	25	1.25	2 years/ 3 generations	rat	M/F	40/40	Males had increased liver weights at an exposure level of 25 ppm. Although size of the litter and the number of pregnancies remained unaffected by DDT, the mortality of the offspring was increased at all exposure levels.	Treon and Cleveland, 1955
	12.5	0.625			M/F	40/40		
	2.5	0.125			M/F	40/40		
	0				M/F	40/40		
Technical	200	10	2 generations	rat/ Sprague- Dawley	NR	NR	Increased liver weight, but no reproductive disturbance, was seen in exposed animals.	Ottoboni, 1969
Technical	250	32.5	2 generations	mouse	NR	NR	There was an increased incidence of liver tumors at all exposure levels tested. At an exposure of 250 ppm, there was increased mortality in the neonates, and reduced lifespan. Tremors and convulsions occurred in some animals.	Tomatis et al., 1972
	50	6.5						
	10	1.3						
Technical	250	32.5	4 generations	mouse/ BALB/c	NR	NR	Although no effects on reproduction were reported, females suffered reduced lifespans in addition to gastrointestinal bleeding, convulsion and increased incidence of tumors.	Terracini et al., 1973a,b
	20	2.6	2 generations	mouse/ BALB/c	NR	NR		



TABLE 3-2 (cont.)

Isomer <sup>a</sup>	Dose <sup>b</sup>		Duration of Exposure	Species	Sex	Number Used	Effects	Reference
	ppm	mg/kg/day						
p,p'-DDT	100 50	13.0 6.5	2 years	mouse	NR	NR	The incidence of liver tumors was increased at both exposure levels.	Thorpe and Walker, 1973; Walker et al., 1972
Technical p,p'-DDT	200 200	8.0 8.0	3.5-7.5 years	monkey	NR	NR	No observed effects reported.	Durham et al., 1963
Technical	3200 2000 400 0	128 80 16	39-40 months	dog	M/F M/F M/F M/F	7/7 2/2 1/1 1/1	No effects were observed at 400 ppm, but liver damage occurred at 2000 ppm and was more severe at 3200 ppm.	Lehman, 1952, 1965
p,p'-DDT	1000 500 0	80 40	44/48 weeks	hamster	M/F M/F M/F	25-30/ 25-30/ 25-30/ 79	Increased mortality as well as nervousness and convulsions were observed in exposed animals.	Agthe et al., 1970
Technical	1000 500 250	80 40 20	18 months	hamster	NR NR NR	NR NR NR	Increased liver weight and enzyme (glucose-6-phosphate dehydrogenase) activity and decreased lifespan were reported.	Graillet et al., 1975
NR	500 250 125	40 20 10	lifespan	hamster	NR	NR	No observed effects on growth or survival.	Cabral and Shubik, 1977

<sup>a</sup>Technical DDT consists of ~77.1% p,p'-DDT, 14.9% o,p'-DDT, 0.3% p,p'-DDD, 0.1% o,p'-DDD, 4.0% p,p'-DDE, 0.1% o,p'-DDE and 3.5% unidentified compounds (U.S. EPA, 1980a).

<sup>b</sup>The fraction of body weight consumed per day in food, called the food factor, is 0.13 for mice and 0.05 for rats. Exposures given in ppm of food are converted to mg/kg/day by multiplying the ppm exposure level by the food factor.

NR = Not reported

ppm (6.5 mg/kg/day) for 2 years (Thorpe and Walker, 1973; Walker et al., 1972) and at 20 ppm (2.6 mg/kg/day) in a 2-generation study (Terracini et al., 1973a,b). The lifespan of the females that suffered from gastrointestinal bleeding and convulsions was decreased at an exposure level of 20 ppm (2.6 mg/kg/day) for 2-4 generations (Terracini et al., 1973a,b). Durham et al. (1963) reported that there was no observed effect in monkeys exposed to 200 ppm (8.0 mg/kg/day) DDT in the food for 3.5-7.5 years; however, the offspring of dogs exposed to 1 ppm (0.04 mg/kg/day) DDT in the food for 3 generations had consistently increased liver weights (Ottoboni et al., 1977). Hamsters were exposed to much higher levels of DDT, ranging from 125 ppm (10 mg/kg/day) to 1000 ppm (80 mg/kg/day). At an exposure level of 500 ppm (40 mg/kg/day), Agthe et al. (1970) reported nervousness, convulsions and increased mortality in exposed animals; Graillot et al. (1975) reported increased liver weights; and Cabral and Shubik (1977) reported that there was no effect on the growth or survival of exposed hamsters.

3.2.2. Inhalation. Pertinent data regarding the chronic toxicity of inhaled DDT could not be located in the available literature.

### 3.3. TERATOGENICITY AND OTHER REPRODUCTIVE EFFECTS

3.3.1. Oral. When pregnant mice were exposed to 1 mg/kg bw on days 10, 12 and 17 of gestation, morphologic changes were seen in the gonads and the fertility of the female offspring was reduced (McLachlan and Dixon, 1972). Although no teratogenesis was observed, 2.5 mg/kg given daily to pregnant mice was significantly embryotoxic, as well as blastotoxic and fetotoxic (Schmidt, 1973). Similarly in rabbits, increased resorption, premature delivery and reduced fetal growth, but no teratogenicity, were caused by exposure to 50 mg/kg DDT on days 7, 8 and 9 of pregnancy. When rats were exposed to 7 ppm DDT in the diet for 60 days before mating and for the duration of pregnancy, fertility was decreased and the number of resorptions increased, but there was

no evidence of teratogenicity (Green, 1969) (Table 3-3). Although there have been no reports of any teratogenic effects of DDT, with the possible exception of ringtail occurring in rats exposed to 200 ppm DDT in the diet (Ottoboni, 1969), DDT has consistently caused a decrease in reproductive capacity in mice (Keplingler et al., 1968; McLachlan and Dixon, 1972; Schmidt, 1973), rats (Fitzhugh and Nelson, 1947; Fitzhugh, 1948; Treon and Cleveland, 1955; Clement and Okey, 1974; Jonsson et al., 1975) and dogs (Deichmann et al., 1971; Deichmann and MacDonald, 1971). Some studies reported no observed effects on reproduction in rats (Ware and Good, 1967; Duby et al., 1971) and dogs (Ottoboni et al., 1977) (see Tables 3-2 and 3-3).

3.3.2. Inhalation. Pertinent data regarding the teratogenicity of inhaled DDT were not located in the available literature.

#### 3.4. TOXICANT INTERACTIONS

Although there is widespread concern about the synergistic interaction of DDT with other potentially hazardous chemicals, the interaction of DDT with only a few chemicals has been studied. Weisburger and Weisburger (1968) reported that 10 mg/day of DDT significantly increased the incidence of hepatomas in rats, caused by the ingestion of 1 mg/day N-fluorenylacetamide (2-AAF). They postulated that DDT exerted its influence by stimulating the hepatic mixed function oxidase system (Weisburger and Weisburger, 1968). On the other hand, after evaluating the interaction of DDT, aramite, methoxychlor, thiouracil and aldrin, Deichmann et al. (1967) concluded that the chemicals did not act synergistically, and, in fact, may have interacted antagonistically. Although there was no increase in the overall incidence of liver tumors when mice were exposed to 100 ppm DDT and 5 ppm dieldrin simultaneously, the histological characteristics of the liver tumors were more malignant after the mice were exposed to DDT and dieldrin simultaneously (Walker et al., 1972).

TABLE 3-3  
Reproductive Studies

Isomer <sup>a</sup>	Dose <sup>b</sup>		Duration of Exposure	Species	Sex	Number Used	Effects	Reference
	ppm	mg/kg/day						
Technical	7	0.9	120 days	mouse/ BALB/c and CFW	NR	NR	No observed effect on reproduction was reported.	Ware and Good, 1967
p,p'-DDT	15	0.75	2 generations	rat	NR	NR	No effects on reproduction were reported.	Duby et al., 1971
o,p'-DDT	15	0.75			NR	NR		
Technical	15	0.75						
NR	1	0.05	175 days		NR	NR		
p,p'-DDT	500	25	6 months	rat	NR	NR	No effects were seen at the 20 ppm dose level. Growth of offspring was severely depressed at the 200 ppm dose level, and all offspring died within 10 days of birth when the animals were exposed to 500 ppm p,p'-DDT.	Clement and Okey, 1974
	200	10			NR	NR		
	20	1			NR	NR		
					NR	NR		
o,p'-DDT	1000	50	6 months	rat	NR	NR	The only effects on reproduction were seen at the 1000 ppm dose levels, which caused decreased fertility and growth in exposed animals.	Clement and Okey, 1974
	200	10			NR	NR		
	20	1			NR	NR		
NR	150	7.5	8-36 weeks	rat	NR	NR	An exposure level of 150 ppm caused reproductive failure. At 75 ppm, the number of pups/litter was not changed, but there was a decreased number of pregnant females.	Jonsson et al., 1975
	75	3.75			NR	NR		
NR	7	0.35	2 generations	rat	NR	131	None of the rats in the second generation were able to get pregnant. There was decreased fertility and increased mortality in the first generation.	Green, 1969
	0				NR	396		
Technical	200	10	2 generations	rat/ Sprague-Dawley	NR	NR	No reproductive disturbance was seen in exposed animals.	Ottoboni, 1969
NR	25	1.25	2 years/ 3 generations	rat	M/F	40/40	Although size of the litter and the number of pregnancies remained unaffected by DDT, the mortality of the offspring was increased at all exposure levels.	Treon and Cleveland, 1955
	12.5	0.625			M/F	40/40		
	2.5	0.125			M/F	40/40		
	0				M/F	40/40		

TABLE 3-3 (cont.)

Isomer <sup>a</sup>	Dose <sup>b</sup>		Duration of Exposure	Species	Sex	Number Used	Effects	Reference
	ppm	mg/kg/day						
Technical	250	32.5	2 generations	mouse	NR	NR	At an exposure of 250 ppm, there was increased mortality in the neonates, and reduced lifespan. Tremors and convulsions occurred in some animals.	Tomatis et al., 1972
	50	6.5						
	10	1.3						
Technical	250	32.5	4 generations	mouse/ BALB/c	NR	NR	No effects on reproduction were reported.	Terracini et al., 1973a,b
	20	2.6	2 generations		NR	NR		
NR	250	32.5	6 generations	mouse	NR	NR	No reproductive effects were seen at exposure levels of 25 ppm. At an exposure level of 100 ppm lactation was reduced and mortality of the offspring increased. The severity of the effects produced at 100 ppm were greatly increased at 250 ppm.	Keplinger et al., 1968
	100	13.0			NR	NR		
	25	3.25			NR	NR		
p,p'-DDT		12	5 times/week, 14 months	dog	M/F	4/3	Decreased male libido and delayed female estrus were reported, as well as infertility, reduced lactation and mammary gland development, and increased maternal and fetal mortality.	Deichmann et al., 1971; Deichmann and MacDonald, 1971
Technical	10	0.4	3 generations	dog	NR	NR	Except that estrus occurred earlier in exposed dogs than in controls, there were no observed effects on reproduction. However, liver weights were consistently higher in pups exposed to DDT.	Ottoboni et al., 1977
	5	0.2						
	1	0.04						

<sup>a</sup>Technical DDT consists of ~77.1% p,p'-DDT, 14.9% o,p'-DDT, 0.3% p,p'-DDD, 0.1% o,p'-DDD, 4.0% p,p'-DDE, 0.1% o,p'-DDE and 3.5% unidentified compounds (U.S. EPA, 1980a).

<sup>b</sup>The fraction of body weight consumed per day in food, called the food factor, is 0.13 for mice, 0.05 for rats and 0.04 for dogs. Exposures given in ppm of food are converted to mg/kg/day by multiplying the ppm exposure level by the food factor.

NR = Not reported

## 4. CARCINOGENICITY

### 4.1. HUMAN DATA

4.1.1. Oral. A group of prison volunteers ingested daily doses of DDT (3.5 mg/man/day or 35 mg/man/day) for 21.5 months (Hayes et al., 1971). No ill effects ascribed to DDT ingestion were reported 4-5 years after the start of the experiment.

4.1.2. Inhalation. Human data regarding the carcinogenicity of DDT has been collected from occupational exposure, which occurs chiefly through inhalation. In occupational exposures where doses ranged from an estimated 10-40 mg/man/day for 1-8 years (Ortelee, 1958) and 3-18 mg/man/day for an average of 15 years (range, 11-19 years) (Laws et al., 1967), no increased incidence of cancer was reported.

### 4.2. BIOASSAYS

4.2.1. Oral. The carcinogenicity of DDT has been tested extensively in mice, rats and hamsters. An NCI study, reported by Innes et al. (1969), found increased incidence of hepatomas in male and female mice exposed to DDT beginning at 7 days of age. At first the DDT was introduced by gavage; after weaning at 4 weeks the DDT was incorporated in the diet. In addition to liver tumors, females had an increased incidence of lymphomas (Innes et al., 1969) (Table 4-1).

Tarjan and Kemeny (1969) studied five generations of BALB/c mice exposed to DDT at a level of 2.8-3.0 ppm in the diet. A variety of tumors accounted for the increased incidence of cancer in exposed (28.7%) as compared with control (3.2%) groups. Lung carcinoma, which occurred in 116 of the 196 animals with tumors, was the predominant type. Leukemia was found in 23 females and 21 males of the 196 animals that developed tumors. Tumors were also observed in the liver, kidney, spleen, ovary and other organs. The

TABLE 4-1  
Oral Carcinogenicity of DDT

Vehicle	Dose	Length of Treatment	Length of Experiment	Species	Sex	Number Tested	Effects	Reference
NR	0 ppm 3.0 ppm	11fespan	5 generations	mice/ BALB/c	M/F M/F	406 683	Combined tumor incidence across 5 generations: lung tumors (16.9% treated, 1.2% control), lymphomas (4.8% treated, 1.0% control) and leukemias (12.4% treated, 2.5% control) occurred in the mice exposed to DDT.	Tarjan and Kemeny, 1969
Gelatin/ diet	control 46.4 mg/kg/day, days 7-28/ 18.2 mg/kg/day (140 ppm), 4-81 weeks	81 weeks	NR	mice/X <sup>a</sup>	M/F M/F	79/87 18/18	About 30% of the females died during treatment. Hepatomas occurred in 11/18 male and 4/18 female mice exposed to DDT, whereas only 8/79 male and 0/87 female controls had hepatomas.	Innes et al., 1969
Gelatin/ diet	control 46.4 mg/kg/day, days 7-28/ 18.2 mg/kg/day (140 ppm), 4-81 weeks	81 weeks	NR	mice/Y <sup>b</sup>	M/F M/F	90/82 18/18	About 30% of the females died during treatment. Hepatomas occurred in 7/18 male and 1/18 female mice exposed to DDT, whereas only 5/90 male and 1/82 female controls had hepatomas.	Innes et al., 1969
NR	0 ppm <sup>c</sup> (0.26 mg/kg/day) 2 ppm (1.3 mg/kg/day) 10 ppm (6.5 mg/kg/day) 50 ppm (32.5 mg/kg/day) 250 ppm	11fespan	2 generations	mouse/ CF1	M/F M/F M/F M/F	113/111 124/105 104/124 127/104 103/90	The incidence of hepatomas was significantly increased (p<0.01) in both males and females exposed to 250 ppm DDT. When the animals were aged past 60 weeks, the increased incidence of liver tumors was significant (p<0.01) at all dose levels in males, but not females.	Tomatis et al., 1972
NR	0 ppm (6.5 mg/kg/day) 50 ppm (13.0 mg/kg/day) 100 ppm	112 weeks	NR	mouse/ CF1	M/F M/F M/F	47/47 32/30 32/32	Liver tumors in mice on diets containing DDT at a dose of 0, 50 or 100 ppm occurred in 13, 37 and 53% of the males, and 17, 50 and 76% of the females, respectively.	Walker et al., 1972
NR	0 ppm 100 ppm	110 weeks	NR	mouse/ CF1	M/F M/F	45/44 30/30	Liver tumors in mice on diets containing 0 or 100 ppm DDT occurred in 24 and 80% of the males and 23 and 87% of the females, respectively. Malignant liver tumors occurred in 4.4 and 30% of the males and 0 and 40% of the females.	Thorpe and Walker, 1983

TABLE 4-1 (cont.)

Vehicle	Dose	Length of Treatment	Length of Experiment	Species	Sex	Number Tested	Effects	Reference
NR	0 ppm (0.26 mg/kg) 2 ppm (1.5 mg/kg) 10 ppm (6.5 mg/kg) 50 ppm (32.5 mg/kg) 250 ppm	lifetime	6 generations	mouse/ CF1	M/F M/F M/F M/F M/F	328/340 354/339 362/355 383/328 350/293	Cumulative incidence of hepatomas males: 29.5, 50.5, 50.0, 55.9, 86.0% for 0, 2, 10, 50 and 250 ppm, respectively. Cumulative hepatomas females: 4.7, 3.5, 9.0, 13.1, 65.5% for 0, 2, 10, 50 and 250 ppm, respectively.	Turusov et al., 1973
NR	0 ppm (0.26 mg/kg/day) 2 ppm (2.6 mg/kg/day) 20 ppm (32.5 mg/kg/day) 250 ppm	2 generations	lifespan	mouse/ CF1	M/F M/F M/F M/F	107/131 112/136 106/128 106/121	Only liver tumors occurred more frequently in treated as compared with control animals. Male mortality caused by DDT toxicity was increased only in the group receiving a dose of 250 ppm. In males, those on diets of 0, 2, 20 and 250 ppm of DDT had incidences of liver tumors combined across 2 generations of 2/107, 3/112, 1/106, 15/106. For females, the corresponding incidences were 0/131, 0/136, 1/128 and 71/121.	Terracini et al., 1973a,b
NR	0 ppm (32.5 mg/kg/day) 250 ppm	15-30 weeks	65, 95 and 120 weeks	mouse/ CF1	M/F M/F	481 713	Incidence of hepatomas in males treated for 15 weeks or 30 weeks with DDT and killed 65, 95 and 120 weeks after the start of the experiment was 13/60, 25/60, 25/60 and 38/60, 41/60 and 37/60, respectively. Corresponding values for controls, were 12/70, 24/83 and 33/98. When females were treated for 15 weeks or 30 weeks with DDT and killed at 65, 95 and 120 weeks after the start of the experiment, the incidence of hepatomas was 3/60, 11/60, 5/60 and 4/54, 11/65 and 11/54, respectively. Corresponding control values for females were 0/69, 0/72 and 1/90.	Tomatis et al., 1974
NR	0 ppm <sup>d</sup> (3.3 mg/kg/day) 22 ppm (6.6 mg/kg/day) 44 ppm 0 ppm (13.05 mg/kg/day) 87 ppm (26.25 mg/kg/day) 175 ppm	78 weeks	93 weeks	mouse/ B6C3F1	M M M F F F	20 ~50 ~50 20 ~50 ~50	Mortality was significantly increased in treated females and the increase appeared dose-related. No other toxic or carcinogenic effects were seen.	NCI, 1978



TABLE 4-1 (cont.)

Vehicle	Dose	Length of Treatment	Length of Experiment	Species	Sex	Number Tested	Effects	Reference
011	0 ppm 5 mg/kg/day (100 ppm) 10 mg/kg/day (200 ppm) 20 mg/kg/day (400 ppm) 30 mg/kg/day (600 ppm) 40 mg/kg/day (800 ppm)	18 months	NR	rats/ Osborne- Mendel	M/F M M/F M/F M/F	24/12 12 24/12 24/12 36/24	The authors concluded that there was an increased incidence of liver tumor formation, but did not relate the effect to dose levels or sex, and did not provide any statistical analysis. Treatment began at weaning.	Fitzhugh and Nelson, 1947
NR	0 ppm 80 ppm 200 ppm	2 years	NR	rats	M/F M/F M/F	30/30 30/30 30/30	No difference in tumor incidence between treated and control groups, except for bronchogenic carcinomas which occurred in 2/60 controls, 8/60 rats fed DDT at 80 ppm and 0/60 rats fed 200 ppm.	Deichmann et al., 1967; Radomski et al., 1965
NR	0 ppm (25 mg/kg/day) 500 ppm	lifetime	lifetime	rat/ Wistar	M/F M/F	35/32 27/28	Increased incidence of liver tumors were seen in males (9/27, 35%) and females (15/28, 56%). No liver tumors were seen in controls.	Rossi et al., 1977
NR	15 mg/rat, 5 days/week by gavage (612.24 ppm)	1 year	18 months	rats/ Fischer	M/F	15/15	No liver tumors were seen.	Weisburger and Weisburger, 1968
NR	0 ppm (16.05 mg/kg/day) 321 ppm (32.1 mg/kg/day) 642 ppm 0 ppm (10.05 mg/kg/day) 210 ppm (21.0 mg/kg/day) 420 ppm	78 weeks	111 weeks	rat/ Osborne- Mendel	M M M F F F	20 ~50 ~50 20 ~50 ~50	No increase in tumor incidence or signs of toxicity were noted for DDT.	NCI, 1978
NR	0 ppm (6.25 mg/kg/day) 125 ppm (12.5 mg/kg/day) 250 ppm (25 mg/kg/day) 500 ppm	lifetime	lifetime	rats/MRC Portion	M/F M/F M/F M/F	38/38 30/30 30/30 38/38	The incidence of hepatomas in males was 1/38, 0/30, 1/30 and 2/38 for the 0, 125, 250 and 500 ppm groups, respectively. Corresponding incidences for females were 0/38, 2/30, 4/30 and 7/30.	Cabral et al., 1982b
NR	0 ppm (10 mg/kg/day) 125 ppm (20 mg/kg/day) 250 ppm (40 mg/kg/day) 500 ppm	NR	NR	hamster	M/F M/F M/F M/F	40/39 30/28 31/28 39/40	No liver tumors were seen in hamsters at any feeding level. There was a significant trend in total number of tumor-bearing females: 7.5, 16.6, 25.8 and 28.2% for 0, 125, 250 and 500 ppm, respectively.	Cabral et al., 1982a

TABLE 4-1 (cont.)

Vehicle	Dose	Length of Treatment	Length of Experiment	Species	Sex	Number Tested	Effects	Reference
NR	0 ppm (80 mg/kg/day) 1000 ppm	NR	NR	hamster	M/F M/F	31/42 35/36	An increased incidence of liver tumors was not seen. Incidences of adrenal adenomas were significantly elevated in females, but not males. Incidence for males: 0 ppm, 8/31; 1000 ppm, 14/35 and for females: 0 ppm, 2/42; 1000 ppm, 10/36.	Rossi et al., 1983

<sup>a</sup>C57BL/6 females mated with C3H/Anf males

<sup>b</sup>C57BL/6 females mated with AKR males

<sup>c</sup>The fraction of body weight consumed per day in food, called the food factor, is 0.13 for mice and 0.05 for rats. Exposures given in ppm of food are converted to mg/kg/day by multiplying the ppm exposure level by the food factor.

<sup>d</sup>Doses given are the TWA dose computed by NCI.

authors emphasized that the transplacental and translactational exposure to DDT that occurred during the experiment may have had an important, but unquantified, effect upon the results.

Under the auspices of IARC, Tomatis et al. (1972), Terracini et al. (1973a,b), Turusov et al. (1973) and Shabad et al. (1973) investigated DDT carcinogenicity in mice. Several multigeneration studies were conducted in which mice were exposed continuously to DDT, including transplacentally and translactationally. Tomatis et al. (1972) found that mice of both sexes exposed to 250 ppm DDT had a significantly higher incidence of hepatomas. When the mice exposed to lower doses of DDT (2-50 ppm) were allowed to age past 60 weeks, the males, but not the females, had a significantly higher incidence of hepatomas. In another experiment, Tomatis et al. (1974) demonstrated that hepatoma formation was dose-dependent and that exposure to DDT early in life may result in tumor formation in the aging animal even after removal of the source of DDT. Terracini et al. (1973a) found that the incidence of hepatomas was increased in BALB/c mice exposed to DDT. In addition, the mice exposed transplacentally and translactationally had an even higher incidence of liver tumors than the mice exposed after weaning only.

Walker et al. (1972) and Thorpe and Walker (1973) reported an increased incidence of liver tumors in CF1 mice exposed to DDT in their food for 2 years. They divided the liver lesions into two histological types. Subsequent analysis by Reuber (1974,1976) classified the lesions as hepatocellular carcinomas and preneoplastic lesions.

The NCI (1978) bioassay reported no carcinogenic effect from DDT exposure, although the mortality of the female mice was significantly increased in a dose-related manner. The dose levels, however, were low in comparison with other bioassays.

In rats exposed to 100-800 ppm DDT in the diet for 18 months, the dose-related changes included hypertrophy of centrilobular hepatic cells, hyalinization of the cytoplasm and focal necrosis (Fitzhugh and Nelson, 1947). More than half (111 out of 192) of the rats died during the experiment. Of the 81 that remained alive at the end of the treatment period, 4 had liver carcinomas and 11 had preneoplastic hepatic lesions. No liver pathology was observed in the control animals (Fitzhugh and Nelson, 1947). Rossi et al. (1977) reported an increased incidence of liver tumors in rats exposed to 500 ppm DDT in their diets. Other studies in rats on the carcinogenicity of DDT in doses ranging from 210-642 ppm (NCI, 1978; Weisburger and Weisburger, 1968; Deichmann et al., 1967; Radomski et al., 1965) reported no increased incidence of liver tumors.

None of the bioassays in hamsters revealed an increased incidence of tumors in DDT exposed animals (Agthe et al., 1970; Cabral and Shubik, 1977; Graillot et al., 1975), and Graillot et al. (1975) reported a marked dose-related decrease in the incidence of lymphosarcoma in the DDT exposed animals as compared with controls.

4.2.2. Inhalation. Pertinent data regarding the carcinogenicity of inhaled DDT could not be located in the available literature.

#### 4.3. OTHER RELEVANT DATA

No increase in the number of reverse mutants was caused by DDT in Salmonella typhimurium (strains TA1535, TA1536, TA1537 and TA1538) or Escherichia coli (strains B/r try, WP2, WP2 try and hcr), regardless of the presence of rat liver microsomes (Shirasu et al., 1976; Marshall et al., 1976). DDT caused no significant increase in recombination mutants in Bacillus subtilis (Shirasu et al., 1976), in recessive lethal mutants in Neurospora crassa (Clark, 1974; Luers, 1953) or in mutation frequency in Salmonella typhimurium

(Buselmaier et al., 1972). Chromosomal aberrations have been reported in cultivated rat-kangaroo cells (Palmer et al., 1972) and human lymphocytes (Lessa et al., 1976), but not in rat cells (Legator et al., 1973) and in mouse bone marrow cells in vivo (Markaryan, 1966). An increase in dominant lethal mutations was caused by DDT in Drosophila melanogaster (Clark, 1974), the Swiss mouse (Clark, 1974) and the rat (Palmer et al., 1972), but not in the CF1 mouse (Wallace et al., 1976) or the ICR/Ha mouse (Epstein et al., 1972).

#### 4.4. WEIGHT OF EVIDENCE

DDT has been associated with liver tumors in mice (Innes et al., 1969; Tomatis et al., 1972; Terracini et al., 1973a,b) and rats (Rossi et al., 1977), and lymphomas and pulmonary tumors (Tarjan and Kemeny, 1969) in mice. Based on inadequate evidence for carcinogenicity in humans and sufficient evidence of carcinogenicity in animals in spite of inadequate evidence for carcinogenicity in short-term tests (i.e., mutagenicity), DDT is most appropriately classified following the scheme proposed by the Carcinogen Assessment Group of the U.S. EPA (Federal Register, 1984) as a Group B2 - Probable Human Carcinogen.

## 5. REGULATORY STANDARDS AND CRITERIA

The WHO (1971) recommended a maximum interim ADI in food of 0.005 mg/kg/bw for DDT. The ACGIH (1983) recommended a TLV-TWA of 1 mg/m<sup>3</sup> and a TLV-STEL of 3 mg/m<sup>3</sup>, based on analogy with lindane, which was judged to be twice as toxic as DDT (ACGIH, 1980).

## 6. RISK ASSESSMENT

### 6.1. ACCEPTABLE INTAKE SUBCHRONIC (AIS)

DDT is known to be an animal carcinogen and data are sufficient for computing a  $q_1^*$ . Therefore, it is inappropriate to calculate an oral or inhalation AIS for DDT.

### 6.2. ACCEPTABLE INTAKE CHRONIC (AIC)

DDT is a chemical known to be an animal carcinogen and for which data are sufficient for computing a  $q_1^*$ . Therefore, it is inappropriate to calculate an oral or inhalation AIC for DDT.

### 6.3. CARCINOGENIC POTENCY ( $q_1^*$ )

6.3.1. Oral. U.S. EPA (1985) has developed a quantitative estimate of the carcinogenic potency of DDT. The studies selected for quantitative evaluation were: Tarjan and Kemeny (1969), Turusov et al. (1973), Terracini et al. (1973), Thorpe and Walker (1973), Tomatis and Turusov (1975), Cabral et al. (1982b) and Rossi et al. (1977).

The data used for estimation of a  $q_1^*$  from the Tarjan and Kemeny (1969) study as shown in U.S. EPA (1985) are presented in Table 6-1. Group sizes were considered inadequate for evaluation in the  $F_1$  and  $F_2$  generations. A potency estimate of  $7.27 \text{ (mg/kg/day)}^{-1}$  was estimated based on the geometric mean of the potencies for generations  $F_3$ - $F_5$  for lung cancers and leukemia.

The data used for estimation of a  $q_1^*$  from the Turusov et al. (1973) study as shown in U.S. EPA (1985) are presented in Table 6-2. A geometric mean of the potencies across six generations for liver tumors of  $0.80 \text{ (mg/kg/day)}^{-1}$  for males and  $0.42 \text{ (mg/kg/day)}^{-1}$  for females was estimated.

The data used for estimation of a  $q_1^*$  from the Terracini et al. (1973) study as shown in U.S. EPA (1985) are presented in Table 6-3. A geometric mean potency ( $q_1^*$ ) of  $0.082 \text{ (mg/kg/day)}^{-1}$  for liver tumors was calculated across sexes and two generations.

TABLE 6-1

Incidence of the Most Commonly Occurring Malignant Tumors in Each of Five Generations  
of BALB/c Mice Fed DDT<sup>a</sup>

Site/Dose Group	Incidence by Generation <sup>b</sup> (combined male and female) (%)				
	F <sub>1</sub>	F <sub>2</sub>	F <sub>3</sub>	F <sub>4</sub>	F <sub>5</sub>
<u>Lung (carcinomas)</u>					
Control	0/3 (00.0)	0/39 (00.0)	3/51 (5.9)	0/144 (00.0)	2/169 (1.2)
3 ppm DDT <sup>c</sup>	2/10 (20.0)	10/35 (28.5)	13/69 (18.8)	41/264 (15.5)	50/305 (16.4)
Significance <sup>d</sup>	--	p=0.001	p=0.007	p<0.002	p<0.001
q <sub>1</sub> * <sup>e</sup>	--	18.78	9.09	7.45	7.37
q <sub>1</sub> * <sup>f</sup>	--	17.20	9.95	7.16	7.68



TABLE 6-1 (cont.)

Site/Dose Group	Incidence by Generation <sup>b</sup> (combined male and female) (%)				
	F <sub>1</sub>	F <sub>2</sub>	F <sub>3</sub>	F <sub>4</sub>	F <sub>5</sub>
<u>Leukemia</u>					
Control	2/3 (66.6)	1/39 (2.6)	0/51 (00.0)	3/144 (2.1)	4/169 (2.4)
3 ppm DDT <sup>c</sup>	4/10 (40.0)	2/35 (5.7)	11/69 (15.9)	35/264 (13.2)	33/305 (10.8)
Significance <sup>d</sup>	--	p=0.924	p=0.008	p<0.001	p=0.002
q <sub>1</sub> <sup>*e</sup>	--	4.67	9.48	5.79	4.50
q <sub>1</sub> <sup>*f</sup>	--	5.01	8.98	6.22	4.83

-22-

<sup>a</sup>Source: U.S. EPA, 1985<sup>b</sup>Number of animals with tumors/number of animals examined (percent). The F<sub>1</sub> generation contained too few effective animals for reliable statistical analysis.<sup>c</sup>The human equivalent doses are calculated by multiplying the ppm values by 0.13 and then by the cube root of 0.030/70. 3 ppm DDT to BALB/c mice = 0.029 mg/kg bw/day for humans. The DDT was given to the mice for lifetime via the diet every day, so no time correction is necessary.<sup>d</sup>Beneath each dose group incidence is the p value for the comparison of the dose group incidence with that of the control group. The F<sub>1</sub> generation was not analyzed.<sup>e</sup>The q<sub>1</sub>'s were calculated using the human equivalent dose. The index values assume that DDT contamination in the control diets was zero.<sup>f</sup>The q<sub>1</sub>\* were calculated using a level of 0.20 ppm DDT<sub>r</sub> (combined DDT-related residues) in the control feed, as reported by the authors.

TABLE 6-2

Incidence of Benign Liver Tumors in Each of Six Generations of CF-1 Mice Fed DDT<sup>a</sup>

Sex/Dose Group	Benign Liver Tumor Incidence by Generation (%) <sup>b</sup>					
	Parental	F <sub>1</sub>	F <sub>2</sub>	F <sub>3</sub>	F <sub>4</sub>	F <sub>5</sub>
<u>Males</u>						
Control	14/60 (24)	13/60 (21)	20/60 (34)	21/60 (35)	16/60 (26)	23/60 (39)
2 ppm <sup>c</sup>	26/60 (44)	29/60 (48)	38/60 (63)	30/60 (50)	34/60 (57)	25/60 (42)
10 ppm	32/60 (53)	28/60 (47)	33/60 (55)	36/60 (60)	24/60 (40)	26/60 (44)
50 ppm	27/60 (45)	35/60 (58)	41/60 (69)	36/60 (60)	32/60 (53)	28/60 (47)
250 ppm	46/60 (76)	51/60 (85)	53/60 (89)	53/60 (89)	57/60 (95)	48/60 (80)
q <sub>1</sub> * <sup>d</sup>	0.572	0.873	0.935	0.878	1.096	0.598

TABLE 6-2 (cont.)

Sex/Dose Group	Benign Liver Tumor Incidence by Generation (%) <sup>b</sup>					
	Parental	F <sub>1</sub>	F <sub>2</sub>	F <sub>3</sub>	F <sub>4</sub>	F <sub>5</sub>
<u>Females</u>						
Control	3/60 (5)	2/60 (3)	1/60 (2)	2/60 (3)	4/60 (7)	5/60 (8)
2 ppm <sup>c</sup>	3/60 (5)	1/60 (2)	3/60 (5)	5/60 (9)	0/60 (0)	0/60 (0)
10 ppm	2/60 (3)	8/60 (13)	8/60 (13)	3/60 (5)	5/60 (8)	6/60 (10)
50 ppm	8/60 (13)	7/60 (12)	8/60 (13)	9/60 (15)	10/60 (16)	7/60 (11)
250 ppm	37/60 (61)	43/60 (71)	31/60 (52)	40/60 (67)	48/60 (80)	38/60 (64)
q <sub>1</sub> * <sup>d</sup>	0.372	0.471	0.369	0.434	0.526	0.370

<sup>a</sup>Source: U.S. EPA, 1985

<sup>b</sup>Number of animals with tumors/number of animals examined (percent). The effective number of animals was given by Turusov et al. (1973) as 50-60; 60 has been used for every group because the exact number was not given.

<sup>c</sup>The human equivalent doses are calculated by multiplying the ppm values by 0.13 and then by the cube root of 0.030/70 (=0.0753949). No adjustment for time was made because these were lifetime tests and CF-1 mice were fed DDT continuously during that time. For example, human equivalent doses are: 2 ppm = 0.0196, 10 ppm = 0.0980, 50 ppm = 0.4900 and 250 ppm = 2.45 mg/kg bw/day.

<sup>d</sup>The q<sub>1</sub>\*s of the upper-bound limits in units of (mg/kg bw/day of dietary exposure)<sup>-1</sup> were calculated using the multistage model.

TABLE 6-3

Incidence of Benign Liver Tumors in BALB/c Mice Fed DDT During  
a 2-Generation Experiment<sup>a,b</sup>

Dose Group	Incidence of Benign Liver Tumors by Generation <sup>c</sup>		
	Males	Females	
	Parental + F <sub>1</sub>	Parental	F <sub>1</sub>
0 ppm	2/107 (1.9)	0/62 (0)	0/69 (0)
Trend <sup>d</sup>	p<0.001	p<0.001	p<0.001
2 ppm	3/112 (2.7)	0/63 (0)	0/73 (0)
20 ppm	1/106 (0.9)	1/61 (1.6)	0/67 (0)
250 ppm	15/106 (14.2)	28/63 (44.4)	43/58 (74.1)
q <sub>1</sub> * <sup>e</sup>	0.074	0.080	0.094
High dose q <sub>1</sub> *	0.086	0.324	0.718

<sup>a</sup>Source: U.S. EPA, 1985

<sup>b</sup>Number of animals with tumors/number of animals examined (percent). Malignant tumors were not observed in liver.

<sup>c</sup>The numbers in the groups of males were reduced by fighting, so the 2 generations of males were pooled. Each high-dose group shown is statistically different from its control group (p<0.001). Other pairwise tests were not significant.

<sup>d</sup>Beneath the control incidence is the p value for positive trend in incidence over the dose levels.

<sup>e</sup>The q<sub>1</sub>\*s were calculated using the human equivalent dose. The "high-dose q<sub>1</sub>\*" is the result of using only the controls and the high-dose groups in the calculations. The human equivalent doses are calculated by multiplying the ppm values by 0.13 and then by the cube root of 0.030/70 (=0.0753949). For example, 250 ppm = 2.45 mg/kg/day for humans.

The data used for estimation of a  $q_1^*$  from the Thorpe and Walker (1983) study as presented in U.S. EPA (1985) are shown in Table 6-4.  $q_1^*$  values for the incidence of malignant liver tumors in males and females were calculated as  $0.52 \text{ (mg/kg/day)}^{-1}$  and  $0.81 \text{ (mg/kg/day)}^{-1}$ , respectively.

The data used for estimation of a  $q_1^*$  from the Tomatis and Turusov (1975) study as presented in U.S. EPA (1985) are shown in Table 6-5.  $q_1^*$  values for benign liver tumors in males and females were  $1.04 \text{ (mg/kg/day)}^{-1}$  and  $0.49 \text{ (mg/kg/day)}^{-1}$ , respectively.

The data used for estimation of a  $q_1^*$  from the Cabral et al. (1982b) study are shown in Table 6-6. A  $q_1$  of  $0.084 \text{ (mg/kg/day)}^{-1}$  was calculated for liver tumors in female rats. The data used for  $q_1^*$  estimation from Rossi et al. (1977) are also shown in Table 6-6.  $q_1^*$  values for liver tumors in male and female rats of  $0.16$  and  $0.27 \text{ (mg/kg/day)}^{-1}$ , respectively, were calculated.

Table 6-7 presents a summary of the calculated potency estimates. Of these values, the  $q_1^*$  based on Tarjan and Kemeny (1969) was eliminated based on the Dixon statistical criterion for rejecting outliers ( $p=0.01$ ) and the additional considerations that the study was from an unaudited laboratory and that the feed was contaminated with DDT. A geometric average of the values from the remaining six studies resulted in a final  $q_1^*$  estimate for DDT of  $0.34 \text{ (mg/kg/day)}^{-1}$ :  $(0.80 \times 0.42 \times 0.082 \times 0.52 \times 0.81 \times 1.04 \times 0.49 \times 0.084 \times 0.16 \times 0.27)^{1/10} = 0.34$ .

6.3.2. Inhalation. Pertinent data regarding the carcinogenicity of inhaled DDT could not be located in the available literature.

TABLE 6-4

Incidence of Liver Tumors (Benign and Malignant) in CF-1 Mice  
Fed DDT for a Single Generation<sup>a</sup>

Dose Group	Incidence of Benign Liver Tumors <sup>b</sup>	Incidence of Malignant Liver Tumors <sup>b</sup>
<u>Males</u>		
Controls	11/45 (24%)	2/45 (4.4%)
100 ppm	23/30 (80%)	9/30 (30%)
q1*	ND	0.52
<u>Females</u>		
Controls	10/44 (23%)	0/44 (0%)
100 ppm	26/30 (87%)	12/30 (40%)
q1*	ND	0.81

<sup>a</sup>Source: U.S. EPA, 1985

<sup>b</sup>Benign liver tumors in this study were referred to as "type a" and malignant liver tumors as "type b."

ND = Not determined

TABLE 6-5

Incidence of Benign Liver Tumors in CF-1 Mice Fed DDT for 15 or 30 Weeks and then Sacrificed at 65, 95 and 120 Weeks<sup>a,b,c</sup>

Dose Group	Males at Week			Females at Week		
	65	95	120	65	95	120
0 ppm <sup>d</sup>	12/70 (17)	24/83 (29)	33/98 (34)	0/69 (0)	0/72 (0)	1/90 (1)
250 ppm for 15 weeks <sup>e</sup>	13/60 (22) p=0.142	25/60 (42) p=0.040	25/60 (42) p=0.080	3/60 (5) p=0.097	11/60 (14) p<0.001	5/50 (10) p=0.034
250 ppm for 30 weeks <sup>e</sup>	38/60 (63) p<0.001	41/60 (68) p<0.001	37/60 (62) p<0.001	4/54 (7) p=0.034	11/55 (20) p<0.001	11/54 (20) p<0.001
<u>q<sub>1</sub>*<sup>f</sup> values</u>						
all q <sub>1</sub> * <sup>f</sup>	0.36	1.04	0.84	0.19	0.49	0.35
30 week	1.38	1.43	1.06	0.19	0.79	0.43

<sup>a</sup>Source: U.S. EPA, 1985

<sup>b</sup>Number of animals with tumors/number of animals examined (percent).

<sup>c</sup>Some groups were exposed for 15 weeks; other groups were exposed for 30 weeks. All groups were sacrificed serially at 30, 65, 95 and 120 weeks.

<sup>d</sup>The human equivalent dose for 1 ppm for 15 weeks is 0.4084 mg/kg/day and for 30 weeks is 0.8168 mg/kg/day. The human equivalent doses are calculated by multiplying the ppm values by 0.13 and then by the cube root of 0.030/70 (=0.0753949). Adjustments for time consist of multiplying the 15-week dose by 15/90 and the 30-week exposure by 30/90.

<sup>e</sup>Beneath each dosed group incidence is the p value for comparison of the incidence in the dose group with that in the control group.

<sup>f</sup>The q<sub>1</sub>\*s were calculated based on the human equivalent dose shown in footnote d. The term "all q<sub>1</sub>\*" indicates that the dosed groups and the control group were used in the calculation. The "30 week" row contains the results of using only the 30-week exposure cancer data with the control cancer data.

TABLE 6-6

Incidence of Benign Liver Tumors in Rats Fed DDT<sup>a,b</sup>

Dose Group <sup>e</sup>	Cabral et al., 1982 <sup>b,c</sup>		Rossi et al., 1977 <sup>d</sup>	
	Males	Females	Males	Females
0 ppm Trend <sup>f</sup>	1/38 (0) NS	0/38 (0) p=0.003	0/35 (0) --	0/32 (0) --
125 ppm	0.30 (0) NS	2/30 (6.7) NS	--	--
250 ppm	1/30 (3.3) NS	4/30 (13.3) p=0.033	--	--
500 ppm	2/38 (5.3) NS	7/38 (18.4) p=0.005	9/27 (33.3) p<0.001	15/28 (53.6) p<0.001
q <sub>1</sub> * <sup>g</sup>	ND <sup>h</sup>	0.084	0.16	0.27

<sup>a</sup>Source: U.S. EPA, 1985

<sup>b</sup>Number of animals with tumor/number of animals examined (percent).

<sup>c</sup>These were Portion (Wistar derived) rats.

<sup>d</sup>These were Wistar rats.

<sup>e</sup>The human equivalent doses are calculated by multiplying the ppm values by 0.0085499, which is 0.05 mg/kg/day (for rats) multiplied by the cube root of 0.350/70 (=0.0753949). No adjustment for time was made because rats were fed continuously for a lifetime.

<sup>f</sup>Beneath the control group incidence is the p value for a positive trend of incidences as the dose increases, when the p values is less than p=0.05, otherwise NS (not significant). Beneath each dosed group incidence is the p value for the comparison of the incidence in the dosed group with its control group when it is less than p=0.05, otherwise NS.

<sup>g</sup>The q<sub>1</sub>\*s were calculated using the human equivalent dose. For example, 500 ppm = 4.275 mg/kg/day for humans.

<sup>h</sup>Not calculated due to lack of statistical increase in hepatomas.

NS = Not significant; ND = not determined



TABLE 6-7  
Summary of Quantitative Potency Estimates for DDT\*

Species	Tumor Site	q <sub>1</sub> * (mg/kg/day) <sup>-1</sup>		Reference
		Males	Females	
Mouse	lung/leukemia	7.27 (combined)		Tarjan and Kemeny, 1969
Mouse	liver	0.80	0.42	Turusov et al., 1973
Mouse	liver	0.082 (combined)		Terracini et al., 1973
Mouse	liver	0.52	0.81	Thorpe and Walker, 1983
Mouse	liver	1.04	0.49	Tomatis and Turusov, 1975
Rat	liver		0.084	Cabral et al., 1982b
Rat	liver	0.16	0.27	Rossi et al., 1977

\*Source: Adapted from U.S. EPA, 1985

## 7. REFERENCES

ACGIH (American Conference of Governmental Industrial Hygienists). 1980. Documentation of the Threshold Limit Values for Substances in Workroom Air, 4th ed. with supplements through 1981. Cincinnati, OH. p. 117-118.

ACGIH (American Conference of Governmental Industrial Hygienists). 1983. Threshold Limit Values for Chemical Substances and Physical Agents in the Workroom Environment with Intended Changes for 1983-1984. Cincinnati, OH. p. 16.

Agthe, C., H. Garcia, P. Shubik, L. Tomatis and E. Wenyon. 1970. Study of the potential carcinogenicity of DDT in the Syrian golden hamster (34740). Proc. Soc. Exp. Med. 134: 113-116. (Cited in IARC, 1974; NIOSH, 1978)

Buselmaier, W. and G. Rohrborn and C. Propping. 1972. Mutagenitäts-Untersuchungen mit Pestiziden im Host-mediated assay und mit den Dominanten Letaltest an der Maus. Biol. Zentrabl. 91: 307-325. (Ger.) (Cited in NIOSH, 1978)

Cabral, J.R.P. and P. Shubik. 1977. Lack of carcinogenicity of DDT in hamsters. Fed. Proc. 36: 1086. (Cited in NIOSH, 1978)

Cabral, J.R.P., R.K. Hall, L. Rossi, S.A. Bronczyk and K.P. Shubik. 1982a. Lack of carcinogenicity of DDT in hamsters. Tumori. 68: 5-10. (Cited in U.S. EPA, 1985)

Cabral, J.R.P., R.K. Hall, L. Rossi, S.A. Bronczyk and K.P. Shubik. 1982b. Effects of long-term intake of DDT on rats. *Tumori*. 68: 11-17. (Cited in U.S. EPA, 1985)

Callahan, M.A., M.W. Slimak, N.W. Gabel, et al. 1979. Water-Related Environmental Fate of 129 Priority Pollutants, Vol. I. U.S. EPA, Office of Water Planning and Standards, Office of Water and Waste Management, Washington, DC. EPA 440/4-79-029.

Cameron, G.R. and K.K. Cheng. 1951. Failure of oral DDT to induce toxic changes in rats. *Br. Med. J.* 2: 819-821. (Cited in NIOSH, 1978)

Clark, J.M. 1974. Mutagenicity of DDT in mice, Drosophila melanogaster and Neurospora crassa. *Aust. J. Biol.* 27: 427-440. (Cited in NIOSH, 1978)

Clement, J.G. and A.B. Okey. 1974. Reproduction in female rats born to DDT-treated parents. *Bull. Environ. Contam. Toxicol.* 12: 373-377. (Cited in NIOSH, 1978)

Crosby, D.G. and K.W. Moilanen. 1977. Vapor-phase photodecomposition of DDT. *Chemosphere*. 6: 167-172.

Deichmann, W.B. and W.E. MacDonald. 1971. Organochlorine pesticides and human health. *Food Cosmet. Toxicol.* 9: 91-103. (Cited in NIOSH, 1978)

Deichmann, W.B., M. Keplinger, F. Sala and E. Glass. 1967. Synergism among oral carcinogens: IV. The simultaneous feeding of four tumorigens to rats. *Toxicol. Appl. Pharmacol.* 11: 88-103. (Cited in NIOSH, 1978; U.S. EPA, 1980a)

Deichmann, W.B., W.E. MacDonald, A.G. Beasley and D. Cubit. 1971. Subnormal reproduction in beagle dogs induced by DDT and aldrin. *Ind. Med.* 40: 10-20. (Cited in NIOSH, 1978)

Duby, R.F., H.F. Travis and C.E. Terrill. 1971. Uterotrophic activity of DDT in rats and mink and its influence on reproduction in the rat. *Toxicol. Appl. Pharmacol.* 8: 348-355. (Cited in NIOSH, 1978)

Durham, W.F., P. Ortega and W.J. Hayes, Jr. 1963. The effect of various dietary levels of DDT on liver function, cell morphology, and DDT storage in the rhesus monkey. *Arch. Int. Pharmacodyn. Ther.* 141: 111-129. (Cited in NIOSH, 1978)

Epstein, S.S., E. Arnold, J. Andrea, W. Bass and Y. Bishop. 1972. Detection of chemical mutagens by the dominant lethal assay in the mouse. *Toxicol. Appl. Pharmacol.* 23: 288-325. (Cited in NIOSH, 1978)

Federal Register. 1984. Environmental Protection Agency. Proposed guidelines for carcinogenic risk assessment. 49 FR 46294-46299.

Fitzhugh, O.G. 1948. Use of DDT insecticides on food products. *Ind. Eng. Chem.* 40: 704-705. (Cited in NIOSH, 1978)

Fitzhugh, O.G. and A.A. Nelson. 1947. The chronic oral toxicity of DDT (2,2-bis(p-chlorophenyl)-1,1,1-trichloroethane). *J. Pharmacol. Exp. Ther.* 89: 1830. (Cited in NIOSH, 1978)

Grailliot, C., J.C. Gak, C. Lacret, R. Truhaut, D. Fournier and C. Martin. 1975. Recherchers sur les modalites et les mecanismes d'action toxique des insecticides organochlores: II. Etude chez le hamster des effect de toxicite a long terme du DDT. Eur. J. Toxicol. 8: 353-359. (Fre.) (Cited in NIOSH, 1978)

Green, V.A. 1969. Effects of pesticides on rat and chick embryo. Proc. Conf. Trace Substances. Environ. Health. 2: 183-209. (Cited in NIOSH, 1978)

Hart, L.G. and J.R. Fouts. 1965. Further studies on the stimulations of hepatic microsomal drug metabolizing enzymes by DDT and its analogs. Naunyn-Schmiedebergs Arch. Path. Pharmacol. 249: 496-500. (Cited in NIOSH, 1978)

Hayes, W.J., Jr., W.E. Dale, and C.I. Pirkle. 1971. Evidence of safety of long-term, high, oral doses of DDT for man. Arch. Environ. Health. 22: 119-136. (Cited in NIOSH, 1978)

Innes, J.R.M., et al. 1969. Bioassay of pesticides and industrial chemicals for tumorigenicity in mice: A preliminary note. J. Natl. Cancer. Inst. 42: 1101. (Cited in U.S. EPA, 1980a)

IARC (International Agency for Research on Cancer). 1974. DDT. In: Some Organochlorine Pesticides. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man. WHO, IARC, Lyon, France. Vol. 5, p. 83-124.

Jensen, J.A., et al. 1957. DDT metabolites in feces and bile of rats. J. Agric. Food Chem. 5: 919. (Cited in U.S. EPA, 1980a)

Jönsson, H.T., Jr., J.E. Keil, R.G. Gaddy, C.B. Loadholt, G.R. Hennigar and E.M. Walker, Jr. 1975. Prolonged ingestion of commercial DDT and PCB: Effects on progesterone levels and reproduction in the mature female rat. Arch. Environ. Contam. Toxicol. 3: 479-490. (Cited in NIOSH, 1978)

Keller, W.C. and R.A. Yeary. 1980. A comparison of the effects of mineral oil, vegetable oil and sodium sulfate on the intestinal absorption of DDT in rodents. Clin. Toxicol. 16(2): 223. (Cited in U.S. EPA, 1980a)

Keplinger, M.L., W.B. Deichmann and F. Sala. 1968. Effects of combinations of pesticides on reproduction in mice. In: Pesticides Symposia, W.B. Deichmann, Ed. Halos and Associates, Inc., Miami, FL. p. 125-138. (Cited in NIOSH, 1978)

Kinoshita, F.K., J.P. Frawley and K.P. Dubois. 1966. Quantitative measurement of induction of hepatic microsomal enzymes by various dietary levels of DDT and toxaphene in rats. Toxicol. Appl. Pharmacol. 9: 505-513. (Cited in NIOSH, 1978)

Laug, E.P., A.A. Nelson, O.G. Fithugh and F.M. Kunze. 1950. Liver cell alteration and DDT storage in the fat of the rat induced by dietary levels of 1-50 ppm DDT. J. Pharmacol. Exp. Ther. 98: 268-273. (Cited in NIOSH, 1978)

Laws, E.R., A. Curley and F.J. Biros. 1967. Men with intensive occupational exposure to DDT. Arch. Environ. Health. 15: 766-775. (Cited in NIOSH, 1978)

Legator, M.S., K.A. Palmer and I-D. Adler. 1973. A collaborative study of in vivo cytogenetic analysis. Toxicol. Appl. Pharmacol. 24: 337-370. (Cited in NIOSH, 1978)

Lehman, A.J. 1952. Chemicals in foods -- A Report to the Association of Food and Drug Officials on Current Developments: II. Pesticides; III. Subacute and chronic toxicity. Quart. Bull. Assoc. Food Drug Officials USA. 16: 47. (Cited in IARC, 1974; NIOSH, 1978)

Lehman, A.L., Ed. 1965. DDT [a mixture of 1,1,1-trichloro-2,2-bis(p-chlorophenyl)ethane and 1,1,1-trichloro-2-(0-chlorophenyl)-2-(p-chlorophenyl)-ethane]. In: Summaries of Pesticide Toxicity, Food and Drug Administration, U.S. DHEW, Washington, DC. U.S. GPO. p. 16-17. (Cited in IARC, 1974; NIOSH, 1978)

Lessa, J.M.M., W. Becak, M.N. Rabello, C.A.B. Pereira and M.T. Ungaro. 1976. Cytogenetic study of DDT on human lymphocytes in vitro. Mutat. Res. 40: 131-138. (Cited in NIOSH, 1978)

Luers, H. 1953. Untersuchung zur Frage der Mutagenitat des Kontaktinsektizids DDT an Drosophilla melangaster. Naturwiss. 40: 293. (Ger.) (Cited in NIOSH, 1978)

Markaryan, D.S. 1966. Cytogenetic effect of some chlororganic insecticides on the nuclei of mouse bone-marrow cells. Genetika. 1: 132-137. (Cited in NIOSH, 1978)

Marshall, T.C., H. Dorough and H.E. Swim. 1976. Screening of pesticides for mutagenic potential using Salmonella typhimurium mutants. J. Agric. Food Chem. 24: 560-563. (Cited in NIOSH, 1978)

McLachlan, J.A. and R.L. Dixon. 1972. Gonadal function in mice exposed prenatally to p,p'-DDT. Toxicol. Appl. Pharmacol. 22: 327. (Cited in NIOSH, 1978)

Morgan, D.P. and C.C. Roan. 1977. The metabolism of DDT in man. Essays Toxicol. 5: 139. (Cited in U.S. EPA, 1980a)

NCI (National Cancer Institute). 1978. Bioassays of DDT, DDE and p,p'-DDE for possible carcinogenicity. CAS No. 50-29-3, 72-54-8, 72-55-9, NCI-CG-TR-1321. U.S. DHEW. (Cited in U.S. EPA, 1980a)

NIOSH (National Institute for Occupational Safety and Health). 1978. Special Occupational Hazard Review: DDT. U.S. DHEW (NIOSH) Publ. No. 78-200.

Ortega, P., W.J. Hayes, Jr. and W.F. Durham. 1956a. Pathologic changes in the liver of rats after feeding low levels of various insecticides. Am. Med. Assoc. Arch. Pathol. 64: 614. (Cited in NIOSH, 1978)

Ortega, P., W.J. Hayes, Jr., W.F. Durham and A. Mattson. 1956b. DDT in the diet of the rat: Its effect on DDT storage, liver function and cell morphology. Publ. Health Monogr. No. 43. (Cited in NIOSH, 1978)



Ortlee, M.F. 1958. Study of men with prolonged intensive occupational exposure to DDT. Arch. Ind. Health. 18: 433-439. (Cited in NIOSH, 1978)

Ottoboni, A. 1969. Effect of DDT on reproduction in the rat. Toxicol. Appl. Pharmacol. 14: 74-81. (Cited in NIOSH, 1978)

Ottoboni, A., G.D. Bissell and A.C. Hexter. 1977. Effects of DDT reproduction in multiple generations of beagle dogs. Arch. Environ. Contam. Toxicol. 6: 83-101. (Cited in NIOSH, 1978)

Page, G.W. 1981. Comparison of groundwater and surface water for patterns and levels of contamination by toxic substances. Environ. Sci. Technol. 15: 1475-1481.

Palmer, K.A., S. Green and M.S. Legator. 1972. Cytogenic effects DDT and derivatives of DDT in a cultured mammalian cell line. Toxicol. Appl. Pharmacol. 22: 355-364. (Cited in NIOSH, 1978)

Radomski, J.L., W.B. Deichmann, W.E. MacDonald and E.M. Glass. 1965. Synergism among oral carcinogens. I. Results of the simultaneous feeding of four tumorigens to rats. Toxicol. Appl. Pharmacol. 7: 652-656. (Cited in NIOSH, 1978; U.S. EPA, 1980a)

Reuber, M.D. 1974. Statement of testimony at public hearings on cancellation of registrations of aldrin/dieldrin (EPA Exhibit 42). U.S. EPA, Washington, DC. (Cited in NIOSH, 1978)

Reuber, M.D. 1976. Histopathology of carcinomas of the liver in mice ingesting dieldrin or aldrin. Tumori. 62: 463-472. (Cited in NIOSH, 1978)

Rossi, L., M. Ravera, G. Repetti and L. Santi. 1977. Long-term administration of DDT or phenobarbital-Na in Wistar rats. Int. J. Cancer. 19: 179-185. (Cited in NIOSH, 1978)

Rossi, L., O. Barbieri, M. Sanguineti, J.R.P. Cabral, P. Bruzzi and L. Santi. 1983. Carcinogenicity study with technical-grade DDT and DDE in hamsters. Cancer Res. 43: 776-781. (Cited in U.S. EPA, 1985)

Schmidt, R. 1973. Effect of 1,1,1-trichloro-2,2-bis(p-chlorophenyl)-ethane (DDT) on the prenatal development of the mouse (under consideration of distribution of tritium-labeled and carbon-14-labeled DDT in pregnant mice). (Ger.) Biol. Rundsch. 11: 316-317. (Cited in NIOSH, 1978)

Shabad, L., T.S. Kolesnichenko and T.V. Nikonova. 1973. Transplacental and combined long-term effect of DDT in five generations of A-strain mice. Int. J. Cancer. 11: 688-693. (Cited in U.S. EPA, 1980a)

Shirasu, Y., M. Moriya, K. Kato, A. Furuhashi and T. Kada. 1976. Mutagenicity screening of pesticides in the microbial system. Mutat. Res. 40: 19-30. (Cited in NIOSH, 1978)

Spencer, W.W. 1975. Movement of DDT and its derivatives into the atmosphere. Res. Rev. 59: 91-117.

Tarjan, R. and T. Kemeny. 1969. Multigeneration studies on DDT in mice. Food Cosmet. Toxicol. 7: 215-222. (Cited in NIOSH, 1978; U.S. EPA, 1980a)

Terracini, B., M.C. Testa, J.R. Cabral and N. Day. 1973a. The effects of long-term feeding of DDT to BALB/c mice. Int. J. Cancer. 11: 747-764. (Cited in NIOSH, 1978; U.S. EPA, 1980a)

Terracini, B., R.J. Cabral and M.C. Testa. 1973b. A multigeneration study of the effects of continuous administration of DDT to BALB/c mice. In: Pesticides and the Environment: A Continuing Controversy, W.B. Deichmann, Ed. Intercontinental Medical Book Corp., NY. p. 77-86. (Cited in NIOSH, 1978)

Thorpe, E. and A.I.T. Walker. 1973. The toxicology of dieldrin (HEOD): II. Comparative long-term oral toxicity studies in mice with dieldrin, DDT, phenobarbitone, beta-BHC, and gamma-BHC. Food Cosmet. Toxicol. 11: 433-442. (Cited in NIOSH, 1978)

Tomatis, L., V. Turusov, N. Day and R.T. Charles. 1972. The effect of long-term exposure to DDT on CF-1 mice. Int. J. Cancer. 10: 489-506. (Cited in U.S. EPA, 1980a)

Tomatis, L., B. Turusov, N. Day and R.T. Charles. 1972. The effect of long-term exposure to DDT on CF-1 mice. Int. J. Cancer. 10: 489-506. (Cited in NIOSH, 1978; U.S. EPA, 1980a)

Tomatis, L., V. Turusov, R.T. Charles, M. Boiocchi and E. Gati. 1974. Liver tumors in CF-1 mice exposed for limited periods to technical DDT. Z. Krebsforsch. 82: 25-35. (Cited in NIOSH, 1978)

Treon, J.F. and F.P. Cleveland. 1955. Toxicity of certain chlorinated hydrocarbon insecticides for laboratory animals, with special reference to aldrin and dieldrin. Agric. Food Chem. 3: 402-408. (Cited in NIOSH, 1978; IARC, 1974)

Turusov, V.S., N.E. Day, L. Tomatis, E. Gati and R.T. Charles. 1973. Tumors in CF-1 mice exposed for six consecutive generations to DDT. J. Natl. Cancer Inst. 51: 983-995. (Cited in U.S. EPA, 1980a)

U.S. EPA. 1980a. Hazard Assessment Report on DDT, DDD, DDE. Environmental Criteria and Assessment Office, Cincinnati, OH. Internal report.

U.S. EPA. 1980b. Ambient Water Quality Criteria for DDT. Environmental Criteria and Assessment Office, Cincinnati, OH. EPA 440/5-80-038. NTIS PB 81-117491.

U.S. EPA. 1980c. Guidelines and Methodology Used in the Preparation of Health Effects Assessment Chapters of the Consent Decree Water Quality Criteria. Federal Register. 45: 79347-79357.

U.S. EPA. 1983. Methodology and Guidelines for Reportable Quantity Determinations Based on Chronic Toxicity Data. Prepared by the Environmental Criteria and Assessment Office, Cincinnati, OH, OHEA for the Office of Solid Waste and Emergency Response, Washington, DC.

U.S. EPA. 1985. The Carcinogen Assessment Group's Evaluation of the Carcinogenicity of Dicofol (Kelthane), DDT, DDE and DDD (TDE). Prepared by the Carcinogen Assessment Group, OHEA, Washington, DC for the Hazard Evaluation Division Office of Pesticide Programs. EPA 600/6-85-002X. Internal Review Draft.

Walker, A.I.T., E. Thorpe and D.E. Stevenson. 1972. The toxicology of dieldrin (HEOD): I. Long-term oral toxicity studies in mice. Food Cosmet. Toxicol. 11: 415-432. (Cited in NIOSH, 1978; U.S. EPA, 1980a)

Wallace, M.E., P. Knights and A.O. Dye. 1976. Pilot study of the mutagenicity of DDT in mice. Environ. Pollut. 11: 217-222. (Cited in NIOSH, 1978)

Ware, G.W. and E.E. Good. 1967. Effects of insecticides on reproduction in the laboratory mouse: II. Mirex, Telodrin and DDT. Toxicol. Appl. Pharmacol. 10: 54-61. (Cited in NIOSH, 1978)

Weisburger, J.H. and E.K. Weisburger. 1968. Food additives and chemical carcinogens: On the concept of zero tolerance. Food Cosmet. Toxicol. 6: 235. (Cited in U.S. EPA, 1980a)

WHO (World Health Organization). 1971. International Standards for Drinking Water, 3rd ed. Geneva. (Cited in U.S. EPA, 1980a)

Zoeteman, B.C.J., K. Harmsen, J.B.H.J. Linders, C.F.H. Morra and W. Slooff. 1980. Persistent organic pollutants in river water and groundwater of The Netherlands. Chemosphere. 9: 231-249.

APPENDIX  
Summary Table for DDT

	Species	Experimental Dose/Exposure	Effect	q <sub>1</sub> <sup>*</sup>	Reference
Inhalation					
				ND	
				ND	
				ND	
Oral					
				ND	
				ND	
	mice and rats	combined data from six studies	increased incidence of liver tumors	0.34 (mg/kg/day) <sup>-1</sup>	U.S. EPA, 1985

ND = Not derived