

Research and Development



Health Assessment Document for Vinylidene Chloride

Review Draft

(Do Not
Cite or Quote)

NOTICE

This document is a preliminary draft. It has not been formally released by EPA and should not at this stage be construed to represent Agency policy. It is being circulated for comment on its technical accuracy and policy implications.



**Review Draft
EPA-600/8-83-031A
October 1983**

Health Assessment Document for Vinylidene Chloride

NOTICE

This document is a preliminary draft. It has not been formally released by the U.S. Environmental Protection Agency and should not at this stage be construed to represent Agency policy. It is being circulated for comment on its technical accuracy and policy implications.

**U.S. Environmental Protection Agency
Region V Library
200 South Dearborn Street
Chicago, Illinois 60604**

**U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Research and Development
Office of Health and Environmental Assessment
Environmental Criteria and Assessment Office
Research Triangle Park, North Carolina 27711**

DISCLAIMER

The report is an external draft for review purposes only and does not constitute Agency Policy. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

U.S. Environmental Protection Agency

PREFACE

The Office of Health and Environmental Assessment has prepared this health assessment to serve as a "source document" for EPA use. The health assessment document was originally developed for use by the Office of Air Quality Planning and Standards to support decision-making regarding possible regulation of vinylidene chloride as a hazardous air pollutant. However, the scope of this document has since been expanded to address multimedia aspects.

In the development of the assessment document, the scientific literature has been inventoried, key studies have been evaluated and summary/conclusions have been prepared so that the chemical's toxicity and related characteristics are qualitatively identified. Observed effect levels and other measures of dose-response relationships are discussed, where appropriate, so that the nature of the adverse health responses are placed in perspective with observed environmental levels.

The EPA Office of Health and Environmental Assessment (OHEA) is responsible for the preparation of this health assessment document. The OHEA Environmental Criteria and Assessment Office (ECAO/RTP) had overall responsibility for coordination and direction of the document and production effort (Dr. Robert M. Bruce, Project Manager). The chapters addressing physical and chemical properties, sampling and analysis, air quality and biological effects in animals and man were all originally written by Syracuse Research Corporation. The principal authors of these chapters are listed below.

Dr. Dipak K. Basu
Life and Environmental Sciences Division
Syracuse Research Corporation
Syracuse, NY

Mr. John Becker
Life and Environmental Sciences Division
Syracuse Research Corporation
Syracuse, NY

Dr. Joan T. Colman
Life and Environmental Sciences Division
Syracuse Research Corporation
Syracuse, NY

Dr. Michael W. Neal
Life and Environmental Sciences Division
Syracuse Research Corporation
Syracuse, NY

Dr. Joseph Santodonato
Life and Environmental Sciences Division
Syracuse Research Corporation
Syracuse, NY

Dr. Richard Sugatt
Life and Environmental Sciences Division
Syracuse Research Corporation
Syracuse, NY

The OHEA Carcinogen Assessment Group (CAG) was responsible for reviewing and updating the sections on carcinogenicity that were originally written by

Syracuse Research Corporation. Participating members of the CAG are listed below (principal authors of the present carcinogenicity sections are designated by asterisk).

Roy Albert, M.D. (Chairman)
Elizabeth L. Anderson, Ph.D.
Larry D. Anderson*
Steven Bayard, Ph.D.*
David L. Bayliss, M.S.*
Chao W. Chen, Ph.D.
Herman J. Gibb, M.S., M.P.H.
Bernard H. Haberman, D.V.M., M.S.
Charalingayya B. Hiremath, Ph.D.
Robert McGaughy, Ph.D.
Dharm V. Singh, D.V.M., Ph.D.
Todd W. Thorslund, Sc.D.

The following individuals within EPA provided peer-review of this draft or earlier drafts of this document:

Larry Anderson, Ph.D.
Office of Health and Environmental Assessment
Carcinogen Assessment Group

H. Matthew Bills
Acting Director for Monitoring
Systems and Quality Assurance
Office of Research and Development

Robert M. Bruce, Ph.D.
Office of Health and Environmental Assessment
Environmental Criteria and Assessment Office

John B. Clements, Ph.D.
Environmental Monitoring and Support Laboratory, RTP
Office of Research and Development

Larry Cupitt, Ph.D.
Environmental Res. Lab. (RTP)
Office of Research and Development

Paul E. des Rosiers
Industrial and Extractive Processes Division
Office of Environmental Engineering and Technologies

Linda Erdreich, Ph.D.
Office of Health and Environmental Assessment
Environmental Criteria and Assessment Office

James W. Falco, Ph.D.
Office of Health and Environmental Assessment
Exposure Assessment Group

Bruce Gay, Ph.D.
Environmental Sciences Res. Lab. (RTP)
Office of Research and Development

Lester D. Grant, Ph.D.
Office of Health and Environmental Assessment
Environmental Criteria and Assessment Office

Larry L. Hall, Ph.D.
Health Effects Res. Lab. (RTP)

Gary E. Hatch, Ph.D.
Health Effects Research Laboratory
Office of Research and Development

Donna Kuroda
Office of Health and Environmental Assessment
Reproductive Effects Assessment Group

James J. Lichtenberg
Environmental Monitoring and Support Lab. (Cin.)
Office of Research and Development

Steve Lutkenhoff
Office of Health and Environmental Assessment
Environmental Criteria and Assessment Office

Robert McGaughy, Ph.D.
Office of Health and Environmental Assessment
Carcinogen Assessment Group

Raymond Merrill, Ph.D.
Industrial Environmental Res. Lab. (RTP)
Office of Research and Development

Joseph Padgett
Office of Air Quality Planning and Standards
Strategies and Air Standards Division

Nancy Pate, D.V.M.
Office of Air Quality Planning and Standards
Strategies and Air Standards Division

Shabeg Sandhu, Ph.D.
Health Effects Research Lab.
Office of Research and Development

Robert D. Schonbroo, Ph.D.
Environmental Monitoring Systems Laboratory
Las Vegas, Nevada
Office of Research and Development

Jerry F. Stara, D.V.M.
Office of Health and Environmental Assessment
Environmental Criteria and Assessment Office

Bruce Tichenor, Ph.D.
Industrial Environmental Research Laboratory (RTP)
Office of Research and Development

Peter Voytek, Ph.D.
Office of Health and Environmental Assessment
Reproductive Effects Assessment Group

Herbert Wiser, Ph.D.
Senior Science Advisor
Office of Research and Development

Consultants and/or Reviewers

Julian B. Andelman, Ph.D.
Professor of Chemistry
Graduate School of Public Health
University of Pittsburgh
Pittsburgh, PA

I.W.F. Davidson, Ph.D.
Professor of Pharmacology
Bowman Gray School of Medicine
Wake Forest University

Rudolph J. Jaeger, Ph.D.
Consulting Toxicologist
7 Bogert Place
Westwood, NJ

Edmond J. LaVoie, Ph.D.
Head, Metabolic Biochemistry Section
Naylor Dana Institute for Disease Prevention
American Health Foundation
Valhalla, NY

Richard R. Monson, M.D., S.C.D.
Assoc. Prof. of Epidemiology
School of Public Health
Harvard University

TABLE OF CONTENTS

	<u>Page</u>
1. SUMMARY AND CONCLUSIONS.....	1-1
2. INTRODUCTION.....	2-1
3. PHYSICAL AND CHEMICAL PROPERTIES.....	3-1
3.1 SYNONYMS AND TRADE NAMES.....	3-1
3.2 STRUCTURAL AND MOLECULAR FORMULAE AND MOLECULAR WEIGHT.....	3-1
3.3 PHYSICAL PROPERTIES.....	3-1
3.3.1 Description.....	3-1
3.3.2 Boiling Point.....	3-1
3.3.3 Freezing Point.....	3-1
3.3.4 Density	3-2
3.3.5 Refractive Index	3-2
3.3.6 Solubility	3-2
3.3.7 Volatility	3-2
3.3.8 Volatility from Water	3-2
3.3.9 Hazard Parameters	3-3
3.3.10 Dielectric Constant	3-3
3.3.11 Thermodynamic Data	3-3
3.3.12 Viscosity	3-3
3.3.13 Conversion Factors at 25°C and 760 nm pressure	3-4
3.4 STORAGE AND TRANSPORTATION OF THE MONOMER	3-4
3.5 CHARACTERISTICS OF THE COMMERCIAL PRODUCT	3-5
3.6 PURIFICATION OF THE MONOMER	3-5
3.7 CHEMICAL REACTIVITY	3-5
4. SAMPLING AND ANALYTICAL METHODS.....	4-1
4.1 AIR.....	4-1
4.1.1 Air Sampling.....	4-1
4.1.2 Storage of Samples Collected in Solid Sorbent Tubes....	4-7
4.1.3 Analytical Methods.....	4-7
4.2 WATER.....	4-13
4.2.1 Water Sampling.....	4-13
4.2.2 Analytical Methods.....	4-14
4.3 SOIL AND DISPOSAL SITE SAMPLES.....	4-17
4.3.1 Sampling.....	4-17
4.3.2 Analysis.....	4-18

TABLE OF CONTENTS (cont.)

	<u>Page</u>
4.4 MONOMER CONTENT IN POLYMERS AND MONOMER MIGRATION INTO FOOD-SIMULATING SOLVENTS.....	4-19
4.4.1 Sample Collection.....	4-19
4.4.2 Analysis.....	4-19
4.5 ANALYSIS OF FOODS AND OTHER BIOLOGICAL SAMPLES.....	4-21
5. SOURCES IN THE ENVIRONMENT.....	5-1
5.1 SOURCES OF AIR POLLUTION.....	5-1
5.1.1 Manufacture of Vinylidene Chloride Monomer.....	5-2
5.1.2 Manufacture of Polymers Containing Polyvinylidene Chloride.....	5-3
5.1.3 Processing of Fabrication of Polymers.....	5-8
5.1.4 Storage, Handling and Transportation of the Monomer.....	5-8
5.1.5 Chemical Intermediate Production.....	5-15
5.1.6 Incineration of Polymers Containing Polyvinylidene Chloride.....	5-15
5.2 SOURCES IN WATER.....	5-16
5.3 SOURCES IN OIL.....	5-16
5.4 SOURCES IN FOOD.....	5-17
5.5 SUMMARY OF ENVIRONMENTAL LOSSES.....	5-17
6. ENVIRONMENTAL FATE, TRANSPORT, AND DISTRIBUTION.....	6-1
6.1 ATMOSPHERIC FATE, TRANSPORT, AND DISTRIBUTION.....	6-1
6.1.1 Reaction with Atmospheric Radicals and Ozone.....	6-1
6.1.2 Atmospheric Photochemical Reactions.....	6-3
6.1.3 Atmospheric Physical Processes.....	6-4
6.2 AQUATIC FATE, TRANSPORT, AND BIOACCUMULATION.....	6-4
6.2.1 Fate and Transport in Water.....	6-5
6.2.2 Bioaccumulation of Vinylidene Chloride in Aquatic Organisms.....	6-8
6.3 FATE, PERSISTENCE, AND TRANSPORT IN SOIL.....	6-8
7. ENVIRONMENTAL LEVELS AND EXPOSURE.....	7-1
7.1 AIR.....	7-1
7.1.1 Environmental Levels.....	7-1
7.1.2 Exposure.....	7-9

TABLE OF CONTENTS (cont.)

	<u>Page</u>
7.2 WATER.....	7-11
7.2.1 Environmental Levels.....	7-11
7.2.2 Exposure.....	7-17
7.3 SOIL.....	7-17
7.4 FOODS.....	7-18
8. BIOLOGICAL EFFECTS ON PLANTS AND MICROORGANISMS.....	8-1
9. BIOLOGICAL EFFECTS ON AQUATIC ORGANISMS.....	9-1
9.1 ACUTE TOXICITY.....	9-1
9.1.1 Freshwater Fish.....	9-1
9.1.2 Freshwater Invertebrates.....	9-1
9.1.3 Marine Fish.....	9-3
9.1.4 Marine Invertebrates.....	9-3
9.2 SUBACUTE TOXICITY.....	9-3
10. BIOLOGICAL EFFECTS.....	10-1
10.1 PHARMACOKINETICS.....	10-1
10.1.1 Absorption and Distribution.....	10-1
10.1.2 Metabolism.....	10-5
10.1.3 Excretion.....	10-27
10.1.4 Summary of Pharmacokinetics.....	10-29
10.2 ACUTE, SUBACUTE, AND CHRONIC TOXICITY.....	10-31
10.2.1 Acute Exposure.....	10-31
10.2.2 Subacute and Chronic Exposure.....	10-48
10.2.3 Summary of Toxicity.....	10-58
10.3 TERATOGENICITY AND REPRODUCTIVE TOXICITY.....	10-59
10.4 MUTAGENICITY.....	10-69
10.4.1 Mutagenicity in Bacteria.....	10-69
10.4.2 Mutagenicity in Plants and Yeast.....	10-74
10.4.3 Mutagenicity in Cultured Mammalian Cells.....	10-75
10.4.4 Mutagenicity <u>In Vivo</u>	10-75
10.4.5 <u>In Vivo</u> DNA Repair.....	10-76
10.5 CARCINOGENICITY.....	10-78
10.5.1 Animal Studies.....	10-78
10.5.2 Epidemiologic Studies.....	10-118
10.5.3 Quantitative Estimation.....	10-120
11. REFERENCES.....	11-1

LIST OF TABLES

<u>No.</u>	<u>Title</u>	<u>Page</u>
4-1	Breakthrough Volume of Vinylidene Chloride as a Function of Flow Rate, Concentration, and Relative Humidity.....	4-5
5-1	Monomer Production Facilities in the United States and their Capacities.....	5-3
5-2	Air Emissions of Vinylidene Chloride During Monomer Manufacture.....	5-4
5-3	Yearly Consumption of Vinylidene Chloride in Different Industries.....	5-5
5-4	Polymerization Processes, Products, and Their Applications.....	5-7
5-5	Polymerization Sites and Type of Polymer Produced.....	5-9
5-6	Estimated Annual Emissions of Vinylidene Chloride from Polymer Synthesis.....	5-10
5-7	Major Polyvinylidene Chloride Processors.....	5-11
5-8	Manufacturers of Polyvinylidene Chloride-Coated Cellophane.....	5-12
5-9	Major Extruders of Polyvinylidene Chloride Film.....	5-13
5-10	Estimated Emissions from Polyvinylidene Chloride Processing.....	5-14
5-11	Summary of Estimated Environmental Losses of Vinylidene Chloride.....	5-18
7-1	Vinylidene Chloride Concentrations in the Ambient Air Throughout the Continental U.S.....	7-4
7-2	Mean and Median Vinylidene Chloride Concentrations in U.S. Ambient Air of Different Site Types.....	7-8
7-3	Estimated Population Residing Near Plant Producing or Fabricating Monomers and Polymers of Vinylidene Chloride.....	7-10

LIST OF TABLES (cont.)

<u>No.</u>	<u>Title</u>	<u>Page</u>
7-4	Industrial Occurrence of Vinylidene Chloride in Raw Wastewater.....	7-13
7-5	Industrial Occurrence of Vinylidene Chloride in Treated Wastewater.....	7-14
7-6	Vinylidene Chloride Concentration in a Few Waters Near Industrial Sites.....	7-15
7-7	Analysis of Food-Packaging Films for Vinylidene Chloride.....	7-19
7-8	Migration of Vinylidene Chloride from Saran Films to Heptane, Corn Oil, and Water at 49 °C.....	7-20
9-1	Acute Toxicity Values for Marine and Freshwater Fish and Invertebrates Exposed to Vinylidene Chloride.....	9-2
10-1	End-Exposure Body Burdens and Disposition of ¹⁴ C Activity in Rats 72 Hours Following Exposure to 10 or 200 ppm of ¹⁴ C-Vinylidene Chloride for 6 hours.....	10-4
10-2	End-Exposure Body Burdens and Disposition of ¹⁴ C Activity in Rats and Mice 72 Hours Following Inhalation Exposure to 10 ppm ¹⁴ C-Vinylidene Chloride for 6 Hours.....	10-14
10-3	Covalently Bound ¹⁴ C-Activity in Tissues 72 Hours Following Inhalation Exposure to 10 ppm ¹⁴ C Vinylidene Chloride for 6 Hours.....	10-15
10-4	Relative Proportions of ¹⁴ C Urinary Metabolites After Intragastric Administration of 350 mg/kg 1- ¹⁴ C-Vinylidene Chloride or 50 mg/kg ¹⁴ C Monochloroacetic Acid to Male Rats.....	10-19
10-5	Relative Proportions of ¹⁴ C Excretory Products After Intragastric Administration of 50 mg/kg of 1- ¹⁴ C-Vinylidene Chloride to Male Rats or Mice.....	10-20
10-6	Metabolism of ¹⁴ C-Vinylidene Chloride and Covalent Binding of ¹⁴ C Activity to Rat Hepatic Tissue After Intragastric Dose of ¹⁴ C-Vinylidene Chloride.....	10-23

LIST OF TABLES (cont.)

<u>No.</u>	<u>Title</u>	<u>Page</u>
10-7	Metabolism of ^{14}C -Vinylidene Chloride and Covalent Binding of ^{14}C Activity to Rat Hepatic Tissue After Inhalation Exposure to 10 or 200 ppm of ^{14}C -Vinylidene Chloride.....	10-24
10-8	Excretion of Radioactivity by Male Rats Given 0.5 mg/kg or 350 mg/kg ^{14}C -Vinylidene Chloride Intragastrically, Intravenously, or Intraperitoneally.....	10-28
10-9	Half-Lives of Excretion of ^{14}C -Vinylidene Chloride, $^{14}\text{CO}_2$, and Radiolabeled Urinary Metabolites of ^{14}C -Vinylidene Chloride.....	10-30
10-10	Influence of 24-Hour Fasting on the Effect of Oral Administration of Vinylidene Chloride (400 mg/kg) in Corn Oil on Plasma Urea Nitrogen Concentration in Male Rats.....	10-37
10-11	Relationships of Plasma Indicators of Kidney Damage to Dose 24 Hours After Oral Administration of Vinylidene Chloride to Male Rats.....	10-38
10-12	Comparison of Prevalence of Histopathologic Effects of Oral Administration of Vinylidene Chloride (400 mg/kg) in Male and Female Rat Kidneys.....	10-40
10-13	Toxicity of 60 ppm Vinylidene Chloride in Male Mice and Rats.....	10-49
10-14	Effect on Experimental Animals of Long Term Inhalation of Vinylidene Chloride.....	10-51
10-15	Pathologic Effects of Long-Term Ingestion of Vinylidene Chloride Incorporated in the Drinking Water of Sprague-Dawley Rats.....	10-57
10-16	Exposure Levels and Duration of Exposure to Vinylidene Chloride During Gestation in Mice.....	10-61
10-17	Number of Animals and Exposure Levels Used to Study the Teratogenicity of Vinylidene Chloride.....	10-63
10-18	Incidence of Fetal Alterations Among Rats Exposed to Vinylidene Chloride By Inhalation or By Ingestion.....	10-65

LIST OF TABLES (cont.)

<u>No.</u>	<u>Title</u>	<u>Page</u>
10-19	Incidence of Fetal Malformation Among Litters of Rabbits Exposed to Vinylidene Chloride.....	10-66
10-20	Mouse Tissue Mediated Mutagenicity of Vinylidene Chloride in <u>S. typhimurium</u>	10-73
10-21	Results of Carcinogenicity Bioassays of Vinylidene Chloride.....	10-79
10-22	Gas Chromatography Analysis of Vinylidene Chloride.....	10-81
10-23	Experiment DT401: Exposure by Inhalation to Vinylidene Chloride (VDC) in Air at 150, 100, 50, 25, 10 ppm, 4 Hours Daily, 4-5 Days Weekly, For 52 Weeks.....	10-81
10-24	Exposure By Inhalation to Vinylidene Chloride (VDC) in Air at 200, 100, 50, 25, 10 ppm, 4 Hours Daily, 4-5 Days Weekly, For 52 weeks.....	10-82
10-25	Experiment BT405: Exposure By Inhalation to Vinylidene Chloride (VDC) in Air at 25 ppm, 4 Hours Daily, 4-5 Days Weekly, For 52 Weeks.....	10-83
10-26	Experiment BT403: Exposure By Ingestion (Stomach Tube) to Vinylidene Chloride in Olive Oil at 20, 10, 5 mg/kg Body Weight, Once Daily 4-5 Days Weekly, For 52 Weeks.....	10-84
	Experiment BT404: Exposure by Ingestion (Stomach Tube) to Vinylidene Chloride in Olive Oil at 0.5 mg/kg Body Weight, Once Daily, 4-5 Days Weekly, for 52 Weeks	10-84
10-27	Experiment BT401: Exposure By Inhalation to Vinylidene Chloride (VDC) in Air at 150, 100, 50, 25, 10 ppm, 4 Hours Daily, 4-5 Days For 52 Weeks. Results After 137 Weeks (End of Experiment).....	10-86
10-28	Experiment BT402: Exposure By Inhalation to Vinylidene Chloride (VDC) in Air at 200, 100, 50, 25, 10 ppm, 4 Hours Daily, 4-5 Days For 52 Weeks. Results After 121 Weeks (End of Experiment).....	10-88
10-29	Statistical Analyses of Survival, Mammary Carcinoma Incidence and Pulmonary Adenoma Incidence For Male And Female Swiss Mice in a Carcinogenicity Study of Vinylidene Chloride.....	10-91
10-30	Tumor Incidence in Rats and Mice Exposed to Vinylidene Chloride.....	10-95

LIST OF TABLES (cont.)

<u>No.</u>	<u>Titles</u>	<u>Page</u>
10-31	Tumor Incidence in Rats and Mice Exposed to Vinylidene Chloride.....	10-97
10-32	Vinylidene Chloride: A Chronic Inhalation Toxicity and Oncogenicity Study in Rats. Cumulative Percent Mortality for Male Rats.....	10-99
10-33	Vinylidene Chloride: A Chronic Inhalation Toxicity and Oncogenicity Study in Rats. Cumulative Percent Mortality for Female Rats.....	10-100
10-34	Vinylidene Chloride: A Chronic Inhalation Toxicity and Oncogenicity Study in Rats. Mean Body Weight for Male Rats.....	10-101
10-35	Vinylidene Chloride: A Chronic Inhalation Toxicity and Oncogenicity Study in Rats. Mean Body Weight for Female Rats.....	10-102
10-36	Vinylidene Chloride: A Chronic Inhalation Toxicity and Oncogenicity Study in Rats. Histopathologic Diagnosis and Number of Tumors in Female Rats.....	10-104
10-37	Representative Tissue Specimens Obtained at Necropsy From All Animals.....	10-106
10-38	Tumor Incidence Following Ingestion of VDC.....	10-108
10-39	Tissues Examined For Histologic Changes in the NCI Bioassay.....	10-109
10-40	Mean Body Weight Change (Relative to Controls) of Mice Administered Vinylidene Chloride by Gavage.....	10-110
10-41	Tumors With Increased Incidence in Rats and Mice, as Indicated by the Fisher Exact Test or the Cochran-Armitage Test for Linear Trend.....	10-112
10-42	Tumor Incidence in Female BDIV Rats Treated with VDC on Day 17 of Pregnancy, in Their Progeny Treated Weekly for Life and Controls.....	10-114
10-43	Estimated Cumulative Dose, Duration of Exposure and Date of First Exposure Among 138 Individuals Exposed to Vinylidene Chloride.....	10-120

LIST OF TABLES (cont.)

<u>No.</u>	<u>Title</u>	<u>Page</u>
10-44	Data From Maltoni Inhalation Study on Male Swiss Mice.....	10-132
10-45	Relative Carcinogenic Potencies Among 53 Chemicals Evaluated by the Carcinogen Assessment Group as Suspect Human Carcinogens.....	10-135


LIST OF FIGURES

<u>No.</u>	<u>Title</u>	<u>Page</u>
4-1	Desorption Efficiency vs. Amount Found by GC Analysis.....	4-12
10-1	Metabolic Pathways for Vinylidene Chloride.....	10-6
10-2	Comparisons of Observed LT50 Data for Vinylidene Chloride with Theoretical Curves Predicted for Two Different Mechanisms of Toxicity.....	10-10
10-3	Dose-Response Relationship for Hepatic Glutathione (GSH) Levels, Total Metabolism of ¹⁴ C-Vinylidene Chloride (VDC), and Covalent Binding of Radioactivity to Hepatic Macromolecules.....	10-26
10-4	Dose-Mortality Curves for Administration of Single, Oral Doses of Vinylidene Chloride Dissolved in Corn Oil to Fasted Male Rats of Various Sizes.....	10-44
10-5	Effect of Increasing Concentration of Vinylidene Chloride on Mortality in Mature Male Rats.....	10-47
10-6	Histogram Representing the Frequency Distribution of the Potency Indices of 53 Suspect Carcinogens Evaluated by the Carcinogen Assessment Group.....	10-134

1. SUMMARY AND CONCLUSIONS

Vinylidene chloride is a highly reactive, flammable, clear, colorless liquid that can produce complex peroxides in the absence of chemical inhibitors (Hushon and Kornrieck, 1978; Wessling and Edwards, 1970). The peroxides are violently explosive, and formaldehyde, phosgene, and hydrochloric acid are produced as decomposition products. Vinylidene chloride has a boiling point of 31.6°C at 760 mm Hg and a vapor pressure of 600 mm Hg at 25°C. The solubility of vinylidene chloride in water is 2250 mg/l at 25°C, and the density of the liquid is 1.2132 g/cm³ (20°C). Vinylidene chloride vapor is 3.34 times as dense as air.

Synonyms for vinylidene chloride are 1,1-dichloroethene, 1,1-DCE, and 1,1-dichloroethylene. Vinylidene chloride has a molecular weight of 96.95 and a molecular formula of C₂H₂Cl₂. The structural formula is given below.



Vinylidene chloride monomer production capacity in the United States is approximately 178 million pounds per year (Neufeld et al., 1977; Anonymous, 1978). Virtually all of the vinylidene chloride produced is used in the production of copolymers with vinyl chloride or acrylonitrile (Hawley, 1977). A small percentage (4%) of vinylidene chloride production is used as chemical intermediates (Neufeld et al., 1977).

The two primary methods that have been used in recent years for the sampling and analysis of vinylidene chloride in ambient air are the freeze-trap method and the sorption onto Tenax-GC method with subsequent analysis of the desorbed vinylidene chloride by high resolution gas chromatography with either flame ionization, electron capture, electrical conductivity or mass spectrometric

detectors. Both the freeze-trap and Tenax-GC methods of sample collection have some disadvantages. For example, the freeze-trap method using liquid oxygen (Singh et al., 1979) is a cumbersome method both for sample collection and transportation, and the Tenax-GC method (Pellizzari and Bunch, 1979) may suffer from serious problems of artifact formation.

The two methods commonly used for the analysis of vinylidene chloride in grab aqueous samples are the static head-space method and the dynamic purge-trap method. However, for aqueous samples containing very low levels of vinylidene chloride (e.g., potable water), the dynamic purge-trap is more suitable than the static head-space method because of the higher sensitivity of the former. Although gas chromatography with either flame ionization, electron capture, electrical conductivity or mass spectrometric detectors has been used for the final quantification of vinylidene chloride by both methods, the electrical conductivity detector is preferable to other detectors because of its greater sensitivity and selectivity. The mass spectrometric method is usually used as a confirmatory technique.

Vinylidene chloride in soil samples has been analyzed by solvent extraction in sealed vials with subsequent quantification by GC-FID, and using mass spectrometry as the confirmatory technique (DeLeon et al., 1980). The analysis of vinylidene chloride in food wrapping materials, foods, and biological tissues has been performed either by the static head-space method or by the dynamic purge-trap method in a manner similar to that employed for aqueous samples.

Because of the high volatility of vinylidene chloride, it is lost to the atmosphere during industrial manufacturing of the monomer and polymer, and during storage and handling. The total emission of vinylidene chloride to all media from these facilities has been estimated to be 1,300,400 pounds per year. Moreover, vinylidene chloride originally in aqueous solution is likely to

contribute to air contamination as a result of its high volatility from water (Dilling, 1977).

Under atmospheric smog conditions, the half-life of vinylidene chloride in air has been determined to be 5 to 12 hours (Dilling et al., 1976). In the absence of smog conditions, vinylidene chloride may persist in the atmosphere with a half-life of approximately 2 days (Cupitt, 1980). Volatilization from aquatic media is probably the most significant fate-determining process for vinylidene chloride, although the role of biodegradation still remains uncertain. The half-lives of volatilization of vinylidene chloride from pond, river, and lake water have been estimated to be 6.1 days, 1.2 days, and 4.2 days, respectively. The fate of vinylidene chloride in soils has not been evaluated with certainty. However, it has been concluded from the limited data that both volatilization and leaching may play significant roles in determining the fate of this chemical in soils.

The median ambient air level of vinylidene chloride in urban/suburban areas of the U.S. was estimated to be 20 ng/m^3 (averaging time ranged from 1 hour to 1091 days). However, the median concentration value is substantially higher ($14 \text{ } \mu\text{g/m}^3$) for ambient air in the vicinity of point sources of emission. The estimated daily vinylidene chloride intake from ambient air in urban/suburban areas through inhalation is $0.4 \text{ } \mu\text{g}$. However, the daily inhalation exposure from ambient air may be as high as 0.3 mg in the immediate vicinity of point sources. Vinylidene chloride has been detected in approximately 3% of the total drinking water supplies in the U.S. at an estimated mean concentration of $0.3 \text{ } \mu\text{g/l}$ and a concentration range of 0.2 to $0.5 \text{ } \mu\text{g/l}$. For the majority of the U.S. population, the daily exposure to vinylidene chloride from ingestion of drinking water has been estimated to be less than $0.6 \text{ } \mu\text{g}$, although the maximum daily exposure in

certain communities could exceed 1 μ g. Because of the paucity of data, no estimate of the dietary intake of vinylidene chloride in the U.S. can be made at the present time.

Vinylidene chloride is acutely toxic to aquatic animals at exposure concentrations in the milligrams per liter range. The lowest concentration reported to be acutely toxic to an aquatic organism is 2.4 mg/l (U.S. EPA, 1978). Reported acute and subchronic LC₅₀ values ranged between 11.6 and 250 mg/l for aquatic animals. Vinylidene chloride was not acutely toxic to aquatic algae at a concentration of 712 to 798 mg/l. Vinylidene chloride was, however, toxic to yeast. No information was found concerning the toxicity of vinylidene chloride to domestic animals and non-aquatic wildlife.

Vinylidene chloride is readily absorbed by mammals following oral or inhalation exposure. Vinylidene chloride is metabolized in the liver with a number of possible reactive intermediates, including an epoxide, being formed. These reactive intermediate metabolites can react with macromolecules; this is a characteristic of many chemical carcinogens. The metabolites of vinylidene chloride produce toxic lesions in the liver and kidneys, with inhibitors of metabolism providing protection from vinylidene chloride toxicity. The hepatotoxic effect can be extensive and histological effects can be noted within 2 hours after the onset of exposure. The acute hepatotoxicity of vinylidene chloride was shown to be greater than that of any other chloroethylene. The liver and kidneys remain the target organs for toxic effects regardless of the route of administration (administration vehicle may influence vinylidene chloride metabolism and therefore affect the extent of toxicity) or whether acute, subacute, or chronic exposure occurs.

Vinylidene chloride is mutagenic in bacterial assay systems with positive results being reported in the Ames assay using S. typhimurium and in a reverse

mutation assay using E. coli. In both assays metabolic activation was employed. In the Ames assay, the efficiency of activation depended on both the species and the tissue from which the enzyme system was prepared. There is also some evidence that vinylidene chloride reacts with the DNA of mouse kidney in vivo resulting in low levels of unscheduled DNA synthesis. Other assays, such as the mutation in vitro of cultured Chinese hamster V79 cells and the in vivo dominant lethal assay, have not been able to demonstrate an interaction of vinylidene chloride with genetic material. Vinylidene chloride has not been adequately tested to determine whether the positive mutagenic response observed in bacteria is relevant to nonbacterial systems as well.

Vinylidene chloride has been described as a possible weak teratogen in a study using rats and mice (Short et al., 1977c). In this study, the levels of vinylidene chloride used were toxic to the dams which confounded the interpretation of the results. It was clear, however, that exposure of pregnant rats and mice to high levels of vinylidene chloride could cause fetotoxicity and adversely affect the outcome of pregnancy. Preliminary results from another study (Murray et al., 1978) indicated that vinylidene chloride was not a teratogen in rats or rabbits. The effect of vinylidene chloride on reproduction and fetal and neonatal development of rats was tested in a three generation study with six litter groups being produced (Nitschke et al., 1980). Some toxic effects were seen in adults, but reproductive capacities were not altered and embryological and neonatal development were not adversely affected.

A single epidemiological investigation (Ott et al., 1976) has been performed on a cohort of 138 workers exposed primarily to vinylidene chloride. From the data presented, the length and extent of exposure could not be determined for each individual in the study. Thus, it is possible that the population examined in this study was not adequate. For this cohort there was no

significant difference in the results of clinical tests when compared to matched controls.

There have been eight cancer bioassay studies (Table 10-21) in which vinylidene chloride was administered to rats, mice, or hamsters by either the inhalation or the oral route. Among these studies, only the interim report of Maltoni et al. (1977) indicated a possible positive carcinogenic response.

Maltoni et al. (1977) exposed rats by inhalation to 10, 25, 50, and 100 ppm, mice to 10 and 25 ppm, and hamsters to 25 ppm vinylidene chloride for 4 hours/day, 5 days/week for 12 months. After approximately 80 weeks, preliminary findings of this study were reported on the animals that had died or had palpable tumors. During this observation period, hamsters developed no tumors. Rats showed a possible treatment-related incidence of mammary carcinomas, although the significance of these results was uncertain due to the lack of appropriate control data presented in the interim report. In male mice, kidney carcinomas were observed at the 25 ppm exposure level, while no similar tumors were seen in the control or exposed female mice or in the male mice of the 10 ppm exposure group. The authors assert that these tumors were directly related to the vinylidene chloride treatment; however, it is also possible that these tumors arose as a result of the non-specific facilitation of tumor development as a result of the severe chronic toxicity of vinylidene chloride in male mice.

In a personal communication (Norris, 1982) following the completion of this study, Maltoni stated that the number of mammary carcinomas was not significantly increased in rats. The kidney tumor incidence was considered to be significantly elevated; however, Maltoni said that he has not been able to reproduce these results in other strains of mice. The other seven chronic toxicity studies, including data from an NCI/NTP (1982) bioassay, have failed to demonstrate a significant carcinogenic response with vinylidene chloride in rats or mice.

Thus only limited data suggest that vinylidene chloride may be an animal carcinogen. Vinylidene chloride has been shown to be a mutagen in the Ames assay in the presence of a metabolic activation system, and vinylidene chloride has produced kidney tumors in male Swiss mice, which appears to be a strain-specific response (Norris, 1982; Maltoni, 1977). There is some suggestion that the kidney tumors were produced by a non-specific mechanism as a result of severe kidney toxicity and accompanying compensatory growth. Other bioassays, however, have not been able to demonstrate that vinylidene chloride is a carcinogen in rats, hamsters, or other strains of mice. Although the only available epidemiological study also showed no difference between the exposed and control groups, the study population was small and the extent of individual exposure unknown. Applying the International Agency for Research on Cancer (IARC) criteria for animal studies, the level of currently available evidence for vinylidene chloride carcinogenicity would be regarded as limited and insufficient to provide a firm conclusion regarding its carcinogenic potential in humans.

2. INTRODUCTION

EPA's Office of Research and Development has prepared this health assessment to serve as a "source document" for Agency use. This health assessment was originally developed for use by the Office of Air Quality Planning and Standards to support decision-making regarding possible regulations of vinylidene chloride under the Clean Air Act. However, the scope of this document has been expanded to address vinylidene chloride in relation to sectors of the environment outside of air. It is expected that this document will be able to serve the information needs of many government agencies and private groups that may be involved in decision-making activities related to vinylidene chloride.

3. PHYSICAL AND CHEMICAL PROPERTIES

The data and the discussion presented below have been obtained from Hushon and Kornreich (1978) and Wessling and Edwards (1970), except in cases where the reference is specifically cited.

3.1 SYNONYMS AND TRADE NAMES



Chemical Abstracts Name: 1,1-dichloroethene

CAS No.: 75-35-4

RTECS No.: KV92750

Synonyms: VDC, 1,1-dichloroethene (1,1-DCE), 1,1-dichloroethylene, vinylidene chloride monomer, vinylidene dichloride

3.2 STRUCTURAL AND MOLECULAR FORMULAE AND MOLECULAR WEIGHT



Molecular Weight: 96.95

3.3 PHYSICAL PROPERTIES

3.3.1 Description

Vinylidene chloride is a clear, colorless liquid with a characteristic "sweet" odor.

3.3.2 Boiling Point

31.56°C (at 760 mm Hg)

3.3.3 Freezing Point

-122.5°C

3.3.4 Density

Liquid density at 0°C: 1.2517 g/cm³

Liquid density at 20°C: 1.2132 g/cm³

Vapor density: 3.34 (air = 1)

Density in saturated air: 2.8 (air = 1)

Percent in saturated air: 78%

3.3.5 Refractive Index

n_D^{20} : 1.42468

3.3.6 Solubility

Solubility of vinylidene chloride in water: 2250 mg/l (at 25°C) (Delassus and Schmidt, 1981)

Solubility of water in vinylidene chloride: 0.35 g/l (at 25°C)

Vinylidene chloride is soluble in acetone, ethanol, benzene, diethyl ether, and chloroform.

3.3.7 Volatility

Vapor Pressure (P) in mm Hg: $\log P_{\text{mm}} = 6.9820 - \frac{1104.29}{t + 237.7}$

(where t = temperature in °C)

Therefore, P = 600 mm Hg at 25°C

3.3.8 Volatility from water

Calculated Henry's law constant (unitless) at 25°C: 7.8 (Dilling, 1977).

Half-life ($t_{1/2}$) for evaporation from water at 25°C for a solution depth of 6.5 cm (Dilling, 1977):

Calculated: 20.1 min

Experimental: 27.2 min

The theoretical equation of Dilling (1977) was derived for a specific hydrodynamic regime and may not be applicable for all waters.

3.3.9 Hazard Parameters

Flash point (tag open-cup): 3°F

Flash point (tag closed-cup): -2°F

Autoignition temperature: 1031°F

Explosive limit with air at 28°C

lower: 7.3% by volume

upper: 16.0% by volume

3.3.10 Dielectric Constant

4.67 (at 16°C)

3.3.11 Thermodynamic Data

Latent heat of vaporization at 25°C: 6328 cal/mol

Latent heat of vaporization at boiling point: 6257 cal/mol

Latent heat of fusion: 1557 cal/mol

Specific Heat: 0.275 cal/g

Heat of combustion: 261.93 kcal/mol

Heat of polymerization: -18.0 kcal/mol

Heat of formation (liquid monomer): -6 kcal/mol

Heat of formation (gaseous monomer): 0.3 kcal/mol

Heat capacity at 25.15°C (liquid monomer): 26.745 cal/mol/degree

Heat capacity at 25.15°C (ideal gas): 16.04 cal/mol/degree

Critical parameters:

Temperature (T_c): 222°C

Volume (V_c): 2.19 cm³/mol

Pressure (P_c): 51.3 atm

3.3.12 Viscosity

0.3302 cP (centipoise) (at 20°C)

3.3.13 Conversion Factors at 25°C and 760 mm pressure:

1 ppm \equiv 3.97 mg/m³ (air)

3.4 STORAGE AND TRANSPORTATION OF THE MONOMER

Without added inhibitors, vinylidene chloride in the presence of air forms a complex peroxide compound at temperatures as low as -40°C. The peroxide is violently explosive. Decomposition products of vinylidene chloride peroxides are formaldehyde, phosgene, and hydrochloric acid. Since the peroxide is a polymerization initiator, formation of insoluble polymer in stored vinylidene chloride monomer may be an indication of peroxide formation. The peroxides are adsorbed on the precipitated polymer. Any dry composition containing more than about 15% peroxide detonates from a slight mechanical shock or from heat; therefore, the separation of the polymer from the monomer by filtration, evaporation, or drying may result in explosion.

Inhibitors are used to prevent formation of peroxides in the monomer. The most common practice to inhibit peroxide formation is the addition of 200 ppm of the monomethyl ether of hydroquinone (MEHQ). Other inhibitors include alkylamines, phenols, and organic sulfur derivatives. Inhibited vinylidene chloride is shipped under a nitrogen blanket to avoid contact with air and to prevent the formation of peroxides. In the absence of water, light, and excessive heat, inhibited vinylidene chloride under a blanket of nitrogen can be stored indefinitely.

To retard polymerization, uninhibited vinylidene chloride should be kept away from light and at a temperature below -10°C. Uninhibited vinylidene chloride can be stored in mild steel, stainless steel, or nickel containers. Contact with copper and aluminum should be avoided because the acetylinic impurities in vinylidene chloride may form copper acetylides and aluminum acetylides, and because aluminum reacts with vinylidene chloride to form

aluminum chloroalkyls. Both of these classes of compounds are extremely reactive and potentially hazardous.

3.5 CHARACTERISTICS OF THE COMMERCIAL PRODUCT

A typical analysis of commercial-grade vinylidene chloride monomer (excluding inhibitors) is as follows: vinylidene chloride 99.8%; trans-1,2-dichloroethylene, 900 ppm; vinyl chloride, 800 ppm; 1,1,1-trichloroethane, 150 ppm; cis-1,2-dichloroethylene, 10 ppm; and 1,1-dichloroethane, ethylene chloride, and trichloroethylene--each less than 10 ppm.

3.6 PURIFICATION OF THE MONOMER

Vinylidene chloride forms an azeotrope with 6% methanol. Distillation of the azeotrope, followed by extraction of the methanol with water, is a method of purification of vinylidene chloride.


3.7 CHEMICAL REACTIVITY

Vinylidene chloride, like other olefins, is expected to undergo addition reactions. An example of this addition reaction with HCl is as follows:



This reaction is used commercially for the production of 1,1,1-trichloroethane (methyl chloroform).

The addition reaction of vinylidene chloride with bromine is shown below:



The reaction product from bromination of vinylidene chloride was used as a confirmatory method for the identification and quantification of the parent compound (Going and Spigarelli, 1977). Other reactions of vinylidene chloride with thiolate (McKinney et al., 1959) and cysteinate (Francis and Leitch, 1957) are also known.

In the presence of polymerization initiators, vinylidene chloride undergoes self-polymerization to form homopolymers, or polymerizes with other compounds--

namely, vinyl chloride, alkyl acrylates, and acrylonitrile--to yield copolymers. Copolymers containing various amounts of vinylidene chloride are used as packaging films, carpet backing materials, and modacrylic fibers. When compared with that of other monomers, the reactivity of vinylidene chloride towards polymerization is considered to be average.

4. SAMPLING AND ANALYTICAL METHODS

Several methods are available for the measurement of vinylidene chloride in air, water, soil and disposal site samples, and polymers, and as migrated monomer into food-simulating solvents. The analysis of vinylidene chloride in air is more frequently studied because atmospheric air is probably the most prevalent medium of vinylidene chloride contamination. Because of its high volatility, vinylidene chloride lost during its industrial production and processing primarily enters the atmosphere. Moreover, vinylidene chloride originally present in aqueous media is likely to contribute significantly to air contamination as a result of its high volatility from water.

The sampling and analysis of vinylidene chloride in different media are discussed individually in the following sections.

4.1 AIR

4.1.1 Air Sampling

Three general methods have been used in the past for the collection of vinylidene chloride in air samples: grab sampling, cold-trapping, and trapping in sorbent.

Grab samples can be collected in vacuum bottles or in evacuated syringes (Irish, 1967, cited in Hushon and Kornreich, 1978). The difficulties with this method of collection are twofold. First, since it does not allow sample preconcentration, the detection limit of the method is too high to permit detection of most ambient air levels of vinylidene chloride. Second, the inherent limitation of any grab sampling method is that samples collected by this method will not permit the determination of time-weighted average concentrations.

The inability to collect samples representative of time-weighted average concentrations with the grab sampling method was overcome by the use of a

personal sampler that used the critical orifice concept (Williams et al., 1976). The sampler consisted of an evacuated container (<1 torr) equipped with a critical orifice (2 to 5 μ m diameter). In such a sampler, the gas flow into the container remains constant as long as a critical flow pattern exists. In order for this to occur, the internal pressure of the container must remain less than half the external atmospheric pressure during sampling. Thus, the length of sampling time depends on both the size of the orifice and the volume of the sample container. For a container of 100 ml volume equipped with a 2 μ m orifice, the critical flow will exist for 22 hours. The use of a critical orifice for time averaged sampling has been used in source sampling for many years. The major problem that occurs is blockage of the orifice by particulate matter.

In the cold-trap method, air containing vinylidene chloride is pumped through a cold-trap maintained at a suitable temperature (Irish, 1967, cited in Hushon and Kornreich, 1978). In order to avoid sample loss, the temperature of the cold-trap must be maintained so that the trapped compound has negligible vapor pressure under the applied pressure. The cold-trap method was used by Singh et al. (1981a,b) for the measurement of ambient air concentrations of vinylidene chloride. In this method, sample preconcentration was conducted with a 4-inch long stainless steel tubing packed with 100/120 mesh glass beads and maintained at liquid O₂ temperature.

The collection of atmospheric vinylidene chloride by sorbent-trapping has been used by several investigators. The advantage of the sorbent-trapping method is that it permits collection of a large volume of air and it concentrates the compound of interest in a small volume of the sorbent. Both liquid and solid sorbents have been used. Gronsborg (1975) bubbled air through pyridine to collect vinylidene chloride. Guillemin et al. (1979) suggested bubbling air through an impinger containing chilled water for the collection of halogenated

ethylenes; however, the method was not applied for sampling air and the efficiency of collection was not established. The disadvantage of handling the impingers in the field and transporting them to the laboratory renders this method unattractive.

The trapping of vinylidene chloride in a solid sorbent is a more suitable method for its collection from air samples. Although silica gel was used in the past (Irish, 1967, cited in Hushon and Kornreich, 1978), sorption on activated carbon has been used extensively in recent years for the collection of vinylidene chloride in air. Many investigators (Russell, 1975; Severs and Skory, 1975; Foerst, 1979) used this method for the monitoring of the higher levels of vinylidene chloride present in workplace atmospheres. In only one investigation (Going and Spigarelli, 1977) was this method utilized for monitoring ambient air levels of vinylidene chloride.

The adsorption capacity of vinylidene chloride on activated carbon depends on the characteristics of the carbon (source, particle size, nature of pretreatment) and the amount used. Generally, for short-term sampling (less than 8 hours) sample tubes containing smaller amounts of carbon are used than for long-term sampling (about 24 hours). For example, an increase in the amount of a specific activated charcoal (PCB 12 x 30) from 150 mg to 600 mg allowed an increase in collection time from 10 to 15 minutes to at least 75 minutes at a flow rate of 1 liter/minute (Severs and Skory, 1975). The effect of adsorption capacity on the nature of activated carbon was also demonstrated by these investigators. Segmentation of the carbon tubes into a front and back section is usually employed for determination of the breakthrough limit during sample collection. If the back section retains more than a certain predetermined percentage of vinylidene chloride, it is taken as an indication that the breakthrough limit has been exceeded.

The sampling unit usually consists of the following components assembled in the order given: probe, charcoal tube, critical orifice, and a battery-operated pump. The tube containing the charcoal is held vertically at a height of 4 to 6 feet from the ground and air is drawn through it at a constant flow rate for a specified length of time. At the end of sample collection period, the tube is closed with caps and shipped to the laboratory for analysis.

For a known weight and nature of activated carbon, the breakthrough characteristics during sample collection are dependent on: (1) concentration of vinylidene chloride in air, (2) flow rate of air through the carbon bed, (3) relative humidity of air, and (4) presence of other contaminants in the air. The influence of the first three factors on the breakthrough volume has been thoroughly studied by Foerst (1979). This investigator sampled vinylidene chloride through a 100 mg front section and 50 mg back section of MSA-8 charcoal tubes, and monitored the vinylidene chloride concentration by gas chromatography-flame ionization detector (GC-FID). The results of his experiments showing the dependence of breakthrough volume on the three parameters are given in Table 4-1. It is evident from Table 4-1 that the flow rate and concentration of vinylidene chloride have relatively less effect on both the breakthrough volume and loading at breakthrough than do the changes in relative humidity.

The effect of loading on breakthrough volume of Carbosieve B was also studied by Russell (1975). This investigator stated that, "the quantity of the compound of interest collected affects the breakthrough volume very slightly. A pollutant present at 20 ppm will break through at a slightly smaller volume than when at 1 ppm." Relative humidity, however, has a more profound effect on the breakthrough characteristics. A change in relative humidity from 17 to 95% resulted in an almost 90% decrease in both the breakthrough volume and loading capacity. It should be noted that the findings of Russell (1975) regarding the

TABLE 4-1

Breakthrough Volume of Vinylidene Chloride as a Function
of Flow Rate, Concentration, and Relative Humidity^a

Concentration (mg/m ³)	Flow Rate (l/min)	Relative Humidity (%)	Breakthrough ^b Volume (l)	Loading at Breakthrough (μg)
Effect of Flow Rate				
13.5	1.00	87	8.9	121
13.4	0.20	85	6.9	93
Effect of Concentration				
114	1.00	95	6.0	684
13.5	1.00	87	8.9	121
Effect of Humidity				
114	0.74	17	56.0	6375
114	1.00	95	6.0	684

^aSource: Foerst, 1979

^bVolume at which the effluent from the charcoal tubes is equal to 5% of the concentration being sampled.

effect of relative humidity on breakthrough volume are consistent with those of Foerst (1979) as shown in Table 4-1.

It is interesting to point out that the breakthrough studies done by Going and Spigarelli (1977) employed air with a relative humidity of 39 to 42% for sampling vinylidene chloride through a 2 g bed of Fisher activated charcoal. They concluded that the maximum loading before breakthrough occurs at approximately 2.5% of the weight of total charcoal bed and that, for vinylidene chloride concentrations much lower than 10 ppm (as expected in ambient air), sampling can be performed at 1 liter/minute for 24 hours without exceeding the breakthrough characteristics. While this may be true at 39 to 42% relative humidity, the results of Foerst (1979) indicate that the volume and loading characteristics will drastically change at considerably higher relative humidity. Although the relative humidity during the collection of samples was not recorded by Going and Spigarelli (1977), light rain was falling during the collection of some samples, which indicates that the relative humidity during field sampling was much higher than 39 to 42%. The field samples collected by Going and Spigarelli (1977) at high relative humidity at a sampling rate of 1 liter/minute for 24 hours might have exceeded the breakthrough characteristics; consequently, some of the results reported by these investigators may be erroneous.

The possibility of using other sorbents for the preconcentration of vinylidene chloride during atmospheric sample collection was studied by Pellizzari and Bunch (1979). They evaluated three sorbents, namely, charcoal, XAD-2, and Tenax GC for the evaluation. It was determined by these investigators that while charcoal and XAD-2 may form traces of halogenated hydrocarbons by in situ reactions of Cl_2 , olefins, and other pollutants present in the atmosphere, the possibility of artifact formation was minimal with Tenax GC. Therefore, Tenax GC sorbent was used by these investigators for the collection of vinylidene chloride

and other volatile organics in the atmosphere. The performance characteristics of Tenax GC as a sorbent for the collection of organics in ambient air were also evaluated by Brown and Purnell (1979). It should be mentioned that although Pellizzari and Bunch (1979) reported minimal artifact formation with Tenax GC, Singh et al. (1979) observed "serious" artifact formation problems with this adsorbent. In addition, Singh et al. (1979) encountered "serious difficulties" in quantitatively adsorbing and desorbing a few specific species with this sorbent.

4.1.2 Storage of Samples Collected in Solid Sorbent Tubes

The stability of vinylidene chloride collected in charcoal-packed tubes was studied by a number of investigators (Severs and Skory, 1975; Going and Spigarelli, 1977; Foerst, 1979). No loss of vinylidene chloride was detected when the samples in charcoal-packed glass collecting tubes were stored in a refrigerator (2 to 4°C) for 16 to 21 days (Going and Spigarelli, 1977; Foerst, 1979). Similarly, Tenax-GC cartridges stored in cooled culture tubes showed high recoveries with halogenated aliphatic compounds (Pellizzari, 1982; Krost et al., 1982).

4.1.3 Analytical Methods

Vinylidene chloride collected by the grab method can be analyzed by three different methods: gas chromatography with various detectors, Infra-red (IR) analysis, and Laser Stark Spectrometry.

In the combined gas chromatography-mass spectrometric (GS-MS) method applied by Grimsrud and Rasmussen (1975), 20 ml samples were flushed onto a SCOT OV-101 chromatographic column temperature-programmed from -60°C to 100°C. The effluents from the GC were passed to a mass spectrometer operated in the single-ion mode. The identity of vinylidene chloride was established on the basis of chromatographic retention time and by monitoring ions of masses 61 and 96 in the

mass spectrum. The detection limit of the method was 5 ppt; however, the chromatographic conditions employed could not separate vinylidene chloride from its isomers cis- and trans-1,2-dichloroethylene.

A portable IR analyzer was used by Foerst (1979) for continuously monitoring vinylidene chloride produced in a dynamic flow vapor generator. A Wilks Miran IA IR analyzer operated in the absorbance mode at a wavelength of 9.1 μm , with a cell path length of 20.25 meters, and a slit width of 2 mm was used. The response time for vinylidene chloride was 4 seconds. This method, however, is unsuitable for monitoring vinylidene chloride in atmospheric air for two reasons. First, the concentration of vinylidene chloride in atmospheric air (on the order of ppt) is far below the detection limit of the IR analyzer. Second, some of the other halohydrocarbons present in atmospheric air samples will produce interference in the vinylidene chloride determinations.

Laser Stark Spectrometry with a 9.6 μm line from a CO_2 IR laser can be used for the determination of vinylidene chloride; however, the detection limit of this method with a 40 cm Stark cell is in the ppm range (Sweger and Travis, 1979). A multipass Stark cell could result in a significant increase in vinylidene chloride detection sensitivity but is not yet available. Other disadvantages of this method include interferences from other halohydrocarbons such as Freons, methyl chloride, methyl fluoride, and vinyl chloride (Sweger and Travis, 1979).

Both the IR and Laser Stark Spectrometry methods have the inherent capability for continuous monitoring of vinylidene chloride levels in the workplace where the concentration of vinylidene chloride is considerably higher than that of atmospheric air and where few interfering substances may be present.

A modified commercial nitrogen oxide analyzer with an ethylene supply system substituted for the ozone source may also be used for the monitoring of compounds containing unsaturated double bonds (Hilborn et al., 1976). This

instrument, which utilizes the principle of chemiluminescence, may be useful for monitoring total unsaturated hydrocarbons in the atmosphere (including vinylidene chloride), but is not suitable for monitoring individual unsaturated hydrocarbons when they are present in a mixture. In addition to detecting olefinic compounds, the instrument also responds to compounds containing sulfide and amine groups (Hilborn et al., 1976).

In one of the more recent methods for determining vinylidene chloride in the ambient air, Singh et al. (1981a,b) collected the samples by cryogenic cooling (see Section 4.1.1). With this method, the desorption of chemicals from the cooling trap was accomplished by maintaining the trap at boiling water temperature and purging it with carrier gas. The desorbed vinylidene chloride was analyzed by a long (33 foot) packed column and electron capture detector.

When vinylidene chloride was collected in an impinger containing pyridine, a colorimetric method was used for its quantification (Gronsborg, 1975). The pyridine solution containing vinylidene chloride was heated with NaOH in a sealed container and then coupled with barbituric acid or $C_6H_5NH_2$ in acetic acid to form a cyanine dye. The dye color intensity was measured with a spectrophotometer. Neither vinyl chloride nor acrylonitrile interfered with vinylidene chloride determination. The detection limit of the method (10 mg/m^3 of air), however, renders it unsuitable for monitoring vinylidene chloride in ambient air.

Vinylidene chloride collected by bubbling air through water in an impinger can be analyzed by injecting the aqueous solution onto a steam-modified gas-solid chromatograph (Guillemin et al., 1979). The details of this technique are given in Subsection 4.2.2. A Spherosil XOA-400 column operated at an isothermal temperature of 128°C with a mobile phase containing a mixture of 50 ml/minute N_2 gas and 16 ml/minute steam has been successfully used for the separation and quantification (by flame ionization detectors) of the haloethylenes including

vinylidene chloride (Guillemin et al., 1979). By using a 200 μ l aqueous solution injection with this method, 20 ppb vinylidene chloride in water can easily be detected.

Vinylidene chloride collected by adsorption on solid sorbents is analyzed by a two-step method. The first step in the analysis is the desorption of vinylidene chloride from the sorbent. The second step is the separation, identification, and quantification of the desorbed vinylidene chloride, usually by gas chromatography employing a flame ionization detector.

Thermal elution and solvent extraction are two commonly used methods for the desorption of vinylidene chloride from solid sorbents. The thermal desorption technique was used by several authors (Pellizzari and Bunch, 1979; Bozzelli et al., 1980). In one thermal desorption method, the sampling tube was reversed (the end where air entered during sampling was attached to the injection port) to allow backflushing of the sorbed compounds into the injection port of a GC. With a 2 x 0.25 inch tube packed with Carbosieve B, Russell (1975) determined the recovery of vinylidene chloride to be 100% ($\pm 3\%$ standard deviation) when the tube was heated at 270°C for 5 minutes at a nitrogen flow of 20 ml/minute. When the dimension of the collection tube was increased to 5.5 by 0.25 inches, however, Severs and Skory (1975) demonstrated that 10 liters of N₂ gas were required to pass through a heated (430°C) PCB 12 x 30 activated carbon collection tube to desorb vinylidene chloride. The desorption efficiency under the above conditions was determined to be 88%.

The sensitivity of the activated carbon adsorption and thermal desorption technique is approximately 1 ppb for a 1-liter air sample using flame ionization detection (Russell, 1975). The high temperature required for desorption, however, may cause some organic compounds to chemically react on the catalytic surface of the sorption tube, thereby altering their chemical composition. For

example, methyl bromide appears to exchange a Br atom for a Cl atom (present as an impurity in the adsorbent) during thermal desorption from Carbosieve B. Thus, peaks from both methyl bromide and methyl chloride appear in the chromatogram (Russell, 1975).

In some of the more recent methods that used Tenax GC as the sorbent, the thermally desorbed vinylidene chloride was either collected in stainless steel tubes under pressure (Bozzelli et al., 1980) or the desorbed gas purged with helium was collected into a liquid nitrogen-cooled nickel capillary trap (Pellizzari and Bunch, 1979). In both cases, the final separation and analysis of vinylidene chloride was done by high resolution gas chromatography with flame ionization detectors.

Desorption by solvent extraction has been used by a few investigators. In this technique, the adsorbent is added to 1 to 10 ml CS_2 either at room temperature in a crimp-sealed vial (Foerst, 1979), or cooled in wet ice (Severs and Skory, 1975), or in dry ice (Going and Spigarelli, 1977). The effect of CS_2 both at 30°C and at -78°C showed almost the same recovery for desorption times ranging from 15 to 90 minutes. The CS_2 desorption efficiency was determined to be 100% by Severs and Skory (1975) and by Going and Spigarelli (1977). Foerst (1979), however, reported that the desorption efficiency was dependent on loading level. There was a trend towards lower desorption efficiencies at lower loadings. This dependence of desorption efficiency on loading level is shown in Figure 4-1. The stability of dilute vinylidene chloride solutions in CS_2 was studied by Going and Spigarelli (1977). About 20% loss of vinylidene chloride was observed whether the sample was stored at -18°C (freezer), 4°C, or 25°C for 7 days. The loss was much less (about 6%) when the solution was stored for only 4 days at the same three temperatures. Therefore, the desorbed vinylidene chloride in CS_2 should be analyzed within 4 days, regardless of the temperature of storage.




Figure 4-1. Desorption Efficiency vs. Amount Found by GC Analysis (Foerst, 1979)

The second step in the analysis of vinylidene chloride by the solvent extraction method involves injection of the CS_2 solution into a gas chromatograph equipped with a flame ionization detector (FID). The choice of FID is based upon its compatibility with CS_2 ; that is, CS_2 has very little response with FID. A number of columns including 10% TCEP, OV-225, 3% Dexsil, Tenax GC, Porapak N, and Chromosorb 102 were tested (Foerst, 1979). None of these columns were able to separate vinylidene chloride, vinyl chloride, and CS_2 . The most suitable column was found to be Durapak OPN on Porasil C (Foerst, 1979; Going and Spigarelli, 1977). Limited success was attained with 20% DC200 (Severs and Skory, 1975), but the separation of vinylidene chloride from CS_2 solvent was less than ideal with this packing material.

The confirmation for vinylidene chloride quantified by FID was made both by mass spectrometry and by bromination of vinylidene chloride (Going and Spigarelli, 1977). The bromination was performed by reacting vinylidene chloride with a 500-fold molar excess of Br_2 for 30 minutes. The excess Br_2 was removed by 0.5 M NaOH solution. The brominated vinylidene chloride was dried over anhydrous Na_2SO_4 and analyzed on a 1.5% OV-101 column equipped with an FID.

4.2 WATER

4.2.1 Water Sampling

Grab water samples should be collected in narrow-mouth, glass, screw-cap bottles with Teflon-faced, silicone rubber, septa cap liners (Federal Register, 1979; Going and Spigarelli, 1977; Bellar and Lichtenberg, 1979; Bellar et al., 1979). To avoid vinylidene chloride loss, the bottles should be filled to overflow with water and should be sealed immediately without leaving any head-space. If the sample contains free or combined chlorine, sodium thiosulfate (10 mg/40 ml for 5 ppm Cl_2) should be added to the water at the time of sample collection (Federal Register, 1979; Bellar et al., 1979). If left unneutralized,

Cl_2 will react with organics in water and will increase the levels of chlorinated hydrocarbons (Bellar et al., 1974, 1979). The samples collected this way should be shipped in ice and refrigerated during storage. A sample stored at 4°C was found to be stable for at least 9 days (Going and Spigarelli, 1977) and samples similarly stored for up to 27 days were stable (Bellar and Lichtenberg, 1981). It was suggested that the analysis of the stored samples be performed within 14 days (Federal Register, 1979).

4.2.2 Analytical Methods

Vinylidene chloride in water samples has been analyzed by three different techniques: direct head-space analysis, direct injection on a steam-modified gas-solid chromatograph, and by the purge-trap technique.

The direct head-space analysis studied by Piet et al. (1978) has the advantage that no preconcentration and prehandling of the samples are required; therefore, systematic error is reduced. The additional advantages of the method are its good GC separability due to the absence of solvents and its low detection limit. The method can be applied for tap water, ground water, and surface waters. In this method, 100 ml N_2 is bubbled through 1 liter of water at 30°C and the N_2 containing the head-space vinylidene chloride is trapped between two columns of water in a special apparatus (Piet et al., 1978). Up to 500 μl of the head-space gas could be injected into a gas chromatograph equipped with an electron capture detector. The GC column used was a wide-bore glass capillary coated with OV-225 and operated isothermally at 22°C. The detection limit for a 100 μl injection was determined to be 2 $\mu\text{g/l}$.

Castello et al. (1982) also used the head-space technique for the determination of halocarbons in drinking water. In their technique, a 5 ml aliquot of water was placed in a closed vial, the sample was thermostatically held at 30°C for 1 hour in a water bath, and 200 μl of the head-space gas was injected into the

GC column attached to an electron capture detector. The authors did not provide the detection limit for vinylidene chloride by this method.

The steam-modified gas-solid chromatography technique, which utilizes a composite of inert carrier gas and a controlled amount of steam as the mobile phase, a porous packing material as the stationary phase, and a flame ionization detector, appears to be a powerful tool for the analysis of aqueous samples (Guillemin et al., 1979). For the analysis of aqueous samples of haloethylenes including vinylidene chloride, a Spherosil XOA-400 column operated isothermally at 128°C and a mobile phase containing a mixture of 50 ml/minute N₂ and 16 ml/minute steam were successfully used. One advantage of this method is that it allows for direct aqueous injections, thereby eliminating time-consuming and loss-incurring pretreatment of samples. The other advantage of this technique over the traditional GC technique is that large amounts of aqueous samples can be injected onto the column without causing any trouble or artifact on the baseline stability, even at maximum sensitivity of the detector amplifier (Guillemin et al., 1979). By using a 200 µl injection with this method, 20 ppb vinylidene chloride in aqueous samples can easily be detected. The method may not be applicable to polluted water samples, however, because of GC separation problems arising from co-contaminants and rapid deterioration of the GC column due to sorption of impurities.

The purge-trap technique is the most widely applied method for the analysis of vinylidene chloride in aqueous samples (Bellar et al., 1979; Bellar and Lichtenberg, 1974, 1979; Going and Spigarelli, 1977; Dowty et al., 1975). This method is applicable for vinylidene chloride analysis in a diversity of media including drinking water, on one extreme, and in raw undiluted sewage on the other (Bellar et al., 1979). This technique has been adopted as the recommended method for the analysis of vinylidene chloride in water samples (Federal

Register, 1979). In the purge-trap technique, the water sample is purged with an inert gas, such as N₂ or He, at room temperature. The purged vinylidene chloride is adsorbed on a trapping column. The trapped vinylidene chloride is thermally desorbed and passes through an analytical column.

The optimum conditions for purging were determined to be either 38 ml/minute for 10 minutes (Going and Spigarelli, 1977) or 40 ml/minute for 11 to 15 minutes (Bellar and Lichtenberg, 1979). Increasing the purge time at the specified flow rate of inert gas resulted in a decrease in vinylidene chloride recovery (Going and Spigarelli, 1977). The purged vinylidene chloride was trapped in a Tenax GC precolumn (Going and Spigarelli, 1977; Bellar et al., 1979). The thermal desorption was performed in the backflush mode for 4 minutes at a temperature of 180°C with an inert gas flow rate of 20-60 ml/minute (Bellar et al., 1979). The average recovery of vinylidene chloride determined for the entire method by multiple analyses over 21 days using repeated 5 ml aliquots of 2 µg/l aqueous solution was determined to be 88% by this purge-trap thermal desorption technique (Bellar and Lichtenberg, 1979). The sensitivity of this technique is dependent on the method of detection and the volume of water used for purging. However, the method is very sensitive because the entire amount of purged vinylidene chloride is used for a single analysis. A method detection limit (MDL) of 0.003 µg/l was reported in EPA method 502.1 (Bellar and Lichtenberg, 1981).

Analytical columns packed with four kinds of materials--0.2% Carbowax 1500 on 80/100 mesh Carbopack-C (Bellar and Lichtenberg, 1979; Going and Spigarelli, 1977), 1% SP-1000 on 60/80 mesh Carbopack-B (Federal Register, 1979), n-octane on 100/120 mesh Porasil C (Federal Register, 1979), and 10% SF-96 and 1% Igepal CO 880 capillary column (Dowty et al., 1975)--were found to be best suited for the separation of vinylidene chloride from co-contaminants. Proper temperature programming was found to be necessary for successful separation.

Both flame ionization (Dowty et al., 1975; Going and Spigarelli, 1977) and halogen specific (Bellar et al., 1979; Federal Register, 1979) detectors were used for the quantification of vinylidene chloride. The flame ionization detector is somewhat unsuitable, however, because the calibration procedure needed with the purge-trap method requires the preparation of aqueous standards by mixing reagent-free water with aliquots of standard solution in water-miscible solvents, such as methanol. Consequently, the aqueous standards prepared in this manner contain relatively large quantities of the miscible organic solvent, relative to the compound of interest. While these organic solvents are purged with relatively low efficiency, their relatively high concentration may lead to significant interferences with the FID. Therefore, the halogen specific detector is more suitable for the purge-trap technique.

The confirmation of vinylidene chloride quantified by GC-FID or GC-Hall detector was made by the GC-MS technique (Bellar et al., 1979; Going and Spigarelli, 1977; Dowty et al., 1975; Federal Register, 1979). Mass spectrometry, either in the specific ion mode (SIM) or in the mass scan mode, was used as the confirmatory technique. The specific ion mode, however, showed better sensitivity of detection (Bellar et al., 1979).

The purge-trap technique described in the preceding discussions, or slightly modified methods, have been used by several investigators recently for the determination of vinylidene chloride in aqueous samples (Pereira and Hughes, 1980); Otson et al., 1982a,b). The precision of the purge-trap method using GC/MS also has been determined recently by Olynyk et al. (1981).

4.3 SOIL AND DISPOSAL SITE SAMPLES

4.3.1 Sampling

DeLeon et al. (1980) reported taking samples from 30-foot vertical borings using the split-spoon method. The samples were placed in glass jars,

sealed with Teflon-lined screw caps, and shipped in containers maintained at temperatures between 6°C and 10°C. Upon arrival at the laboratory, the samples were transferred to a freezer, where they were kept at -20°C until analyzed. Speis (1980) collected sediment with a methanol rinsed stainless steel spatula in 40 ml vials sealed with teflon baked silicon septa. The vials were filled to capacity to minimize headspace and stored up to 1 week at wet ice temperatures.

4.3.2 Analysis

DeLeon et al. (1980) solvent extracted the samples in sealed vials with either n-hexane (for FID quantification) or n-hexadecane (for MS confirmation) by mixing them with a vortex for 30 seconds. Tribromomethane (bromoform) was added as an internal standard for the samples analyzed by FID. The solvent extracts were quantified for vinylidene chloride by the GC-FID method and by GC coupled with MS for confirmation. A temperature-programmed capillary column coated with SE-52 was employed as the analytical column. This column (27 meters long) was found to be suitable for the separation of vinylidene chloride from its isomers cis- and trans-1,2-dichloroethylene. The recoveries of vinylidene chloride by this method were determined at different levels of contamination. At vinylidene chloride levels of 10 µg/g, 100 µg/g, and 300 µg/g, the percent recoveries were determined to be 110.5 ± 3.5 (standard deviation), 86.9 ± 8.3 , and 80.2 ± 3.7 , respectively. The detection limit of the method was determined to be 10 µg/g.

Speis (1980) transferred a weighed portion of sediment into a Hypo vial, sealed the vial, and injected internal standards. The vials were then placed in a sand bath at 110°C and purged with an inert gas at 40 ml/min by means of a sharpened stainless steel tube inserted through the septum to the bottom of the vial. The outlet was passed through a water trap to a Tenax silica gel trap. Desorption and analysis were virtually identical to EPA method 624. Detection

limits were reported to be 5 µg/kg and recoveries were 67% for a 10 µg spike, 90% for a 20 µg spike, 81.1% for a 40 µg spike and 84% for an 80 µg spike.

4.4 MONOMER CONTENT IN POLYMERS AND MONOMER MIGRATION INTO FOOD-SIMULATING SOLVENTS

Saran film is a copolymer of vinyl chloride (15-20%) and vinylidene chloride (80-85%). It is marketed primarily as household film (0.5 mil thickness) and as industrial film (2.0 mil thickness). The monomers in saran films used for food packaging applications have the potential for migration into foods. Therefore it is important to determine the monomer content in the polymer and the extent of the monomer's migration into food-simulating solvents.

4.4.1 Sample Collection

The saran films required for analysis of residual monomer or of its migration into food-simulating solvents can be obtained from the manufacturer or can be purchased directly at retail stores.

4.4.2 Analysis

The analysis of saran films for the monomer concentration has been accomplished by two methods. In one method (Birkel et al., 1977; Hollifield and McNeal, 1978), the film was dissolved in tetrahydrofuran (THF) in a teflon-lined screw cap glass bottle at 70°C. The dissolved polymer containing the monomer was directly injected into a gas chromatograph equipped with an electron capture detector for quantification. The analytical column used was packed either with Chromosorb-102 (Birkel et al., 1977) or Chromosorb-104 (Hollifield and McNeal, 1978). When THF spiked with vinylidene chloride was used, this method yielded recoveries of 96-102% through the quantification steps (Birkel et al., 1977). When a Chromosorb-104 column was used, the detection limit of the method was determined to be about 1 ppm vinylidene chloride in the polymer (Hollifield and McNeal, 1978). The confirmatory technique used with gas-solid chromatography-

electron capture detectors (GSC-ECD) was GC-MS in the specific ion mode (SIM). The disadvantage of this method is that the injections of polymer solutions leave polymer residue in the glass liner of the injection port. If the injection port liner is not cleaned frequently, the polymers may enter the column and alter its performance.

This disadvantage in the analysis of vinylidene chloride monomer by solvent dissolution of the polymer can be overcome by analyzing the monomer content of the polymer by head-space technique. The additional advantage of the head-space technique is that the method is free from solvent interference such as that encountered with the solvent dissolution method. The head-space technique was used by Going and Spigarelli (1977). In this method, the polymer was crimp-sealed in a vial (teflon-lined cap) and the vial was heated. The optimum conditions for heating were determined to be 100°C for 90 minutes. At a higher temperature (150°C), vinylidene chloride was found to decompose; at lower temperatures (50°C and 75°C), vinylidene chloride was not released from the polymer. A 100 µl head-space gas was injected into a Durapak OPN on a 80/100 mesh Porasil C GC column and the separated vinylidene chloride was quantified by FID. This technique, however, has its own disadvantage. The calibration of the method requires standard polymer samples that have known vinylidene chloride concentration. In the absence of such a standard, it becomes impossible to determine the recovery of vinylidene chloride from the thermal release procedure used in the head-space technique.

In order to determine the extent of the migration of vinylidene chloride, the analysis of food-simulating solvents in contact with polymers was performed by Hollifield and McNeal (1978). They used three food-simulating solvents--corn oil, heptane, and water. Strips of saran wraps were placed in 400 ml food-simulating solvents in a sealed can and stored at 49°C; this temperature is

suggested in the Code of Federal Regulations (CFR, 1977, cited in Hollifield and McNeal, 1978) as the most severe temperature condition to which the food packaging material is expected to be subjected. After the desired storage time, 20 ml of the solvent was placed in screw cap bottles sealed with teflon-lined septa. The bottles were heated to 90°C for 30 minutes or more and 1-2 ml of the head-space gas was withdrawn for analysis. For canned water samples, the head-space gas at 25°C was analyzed. The quantitative analysis was done by a GC with a Chromosorb-104 column and an electron capture detector. A gas chromatograph equipped with the same column and coupled with a mass spectrometer operated in the selective ion mode was used as a confirmatory analysis. The recovery of vinylidene chloride after spiking the migration system solution and analyses by the head-space gas-solid chromatography technique was found to be quantitative (95 to 138%). The vinylidene chloride detection limit was determined to be 5-10 ppb in the food-simulating solvents (Hollifield and McNeal, 1978).

A similar head-space technique was employed by Gilbert et al. (1980) for the determination of vinylidene chloride monomer in packaging films. In this method, the films cut into narrow strips were put into hypovials and sealed with rubber septa. The hypovials were heated to 120°C by an air-circulating fan oven for 30 minutes and 1 ml of the head-space gas was removed for analysis by GC with an electron capture detector.

4.5 ANALYSIS OF FOODS AND OTHER BIOLOGICAL SAMPLES

The analysis of vinylidene chloride in food products in contact with polyvinylidene chloride wrappings was performed by Gilbert et al. (1980) by a head-space method similar to that described for wrapping materials. For cooked meat and cheese products, the samples were ground under liquid nitrogen to a fine powder prior to their placements into hypovials. The hypovials were heated to 60°C in a waterbath for a minimum of 90 minutes prior to analysis.

The analysis of vinylidene chloride in fish samples and in body tissues was performed by the purge-trap method similar to that described for aqueous samples (Lin et al., 1982; Easley et al., 1981). Lin et al. (1982) used computerized GC-MS for the determination of vinylidene chloride in fish, and Easley et al. (1981) used a GC equipped with a Hall electrolytic conductivity detector for vinylidene chloride determination in body tissues. The approximate limit of detection for vinylidene chloride in spiked fish samples was determined to be 10 µg/kg by Easley et al. (1981). Lin et al. (1982) determined the absolute detection limit of 50 µg/kg for the analysis of vinylidene chloride in body tissues by their method.

Hiatt (1981, 1983) analyzed fish tissue for volatile compounds including vinylidene chloride by vacuum distillation. Whole fish, stored in dry ice, were homogenized in a food cutter with added liquid nitrogen to prevent loss of volatiles. A 10 g fish sample in the sample flask was then heated to 50°C at 360 torr while immersed in an ultrasonic cleaner bath. Distillation was continued for 15 min.; the volatiles were trapped at liquid nitrogen temperature. The contents of the trap were transferred to a DB5 coated fused-silica capillary column (FSCC) (30 m x 0.32 mm, 0.1 µm film thickness) at -99°C, which was then programmed up to 180°C. Analysis was performed by mass spectrometry. Recovery was reported to be $74 \pm 8\%$ as opposed to $59 \pm 19\%$ for purge and trap of a diluted sample. FSCC chromatography gave better resolution than packed columns for late eluting peaks.

5. SOURCES IN THE ENVIRONMENT

The purpose of this document is to present available information relevant to human health effects that could be caused by this substance.

Any information regarding sources, emissions, ambient air concentrations, and public exposure has been included only to give the reader a preliminary indication of the potential presence of this substance in the ambient air. While the available information is presented as accurately as possible, it is acknowledged to be limited and dependent in many instances on assumption rather than specific data. This information is not intended, nor should it be used, to support any conclusions regarding risks to public health.

If a review of the health information indicates that the Agency should consider regulatory action for this substance, a considerable effort will be undertaken to obtain appropriate information regarding sources, emissions, and ambient air concentrations. Such data will provide additional information for drawing regulatory conclusions regarding the extent and significance of public exposure to this substance.

Vinylidene chloride is a man-made chemical. To date, there is no evidence that any natural process produces this chemical. The sources of vinylidene chloride in four different environmental media--air, water, soil, and foods--are discussed below.

5.1 SOURCES OF AIR POLLUTION



The following operations contribute to the presence of vinylidene chloride in the air:

- (1) Manufacture of vinylidene chloride monomer,
- (2) Manufacture of polymers containing polyvinylidene chloride,

- (3) Processing or fabrication of polyvinylidene homopolymers and copolymers,
- (4) Storage, handling, and transportation of the monomer,
- (5) Chemical intermediate production, and
- (6) Incineration of polymers containing polyvinylidene chloride.

5.1.1 Manufacture of Vinylidene Chloride Monomer

Vinylidene chloride is manufactured industrially either as a coproduct of ethylene dichloride, produced from the chlorination of ethane or ethylene, or by the dehydrochlorination of 1,1,2-trichloroethane. The chemical reactions for the processes are the following:



The current domestic manufacturers of vinylidene chloride monomer and their estimated annual capacity are given in Table 5-1.

The Lake Charles facility of PPG Industries, which used 130 million pounds of vinylidene chloride annually for captive consumption in the manufacture of 1,1,1-trichloroethane, now produces 1,1,1-trichloroethane by a route that does not utilize vinylidene chloride (PPG Industries verbal communication, 1980). The Lake Charles facility of PPG continues to produce vinylidene chloride, however, for sale in the merchant market.

It is reported that the Dow facility at Freeport produces about two-thirds of its total annual capacity (Neufeld et al., 1977). Dow captively consumes about 85% of the total vinylidene chloride it manufactures annually. The balance is shipped to its other polymer-producing facilities or sold to the merchant market.

Vinylidene chloride is emitted to the air during the manufacture of the monomer. The air emission factors, that is, the fraction of vinylidene chloride lost to the air during its manufacture, can be estimated from material balance.

TABLE 5-1

Monomer Production Facilities in the United States and Their Capacities^a

Manufacturer	Location	Estimated Capacity (million lb/yr)
PPG Industries	Lake Charles, LA	78 ^b
Dow Chemical	Freeport, TX and Plaquemine, LA	95-100

^aSource: Neufeld et al., 1977

^bFigure obtained from Anonymous, 1978

These air emission factors for vinylidene chloride manufacturing processes are given by A.D. Little (1976, cited in Going and Spigarelli, 1977). Based on these values and the estimated production (estimated as 80% of capacity) of vinylidene chloride monomer, the annual emission rates derived are shown in Table 5-2.

The estimated emission of 450,000 pounds/year is considerably lower than the figure given by other authors (Hushon and Kornreich, 1978). The difference is due mainly to PPG Industries' no longer producing 1,1,1-trichloroethane from vinylidene chloride. However, PPG still produces vinylidene chloride for sale to polymer producing facilities.

5.1.2 Manufacture of Polymers Containing Polyvinylidene Chloride

The two primary uses of vinylidene chloride are in the manufacture of polymers and chemical intermediates. The breakdown of vinylidene chloride consumption in different industrial uses is given in Table 5-3.

TABLE 5-2

Air Emissions of Vinylidene Chloride During Monomer Manufacture

Producer	Location	Air Emission Factor	
		(emission in lb/100 lb monomer produced)	Annual Emission (thousands of lb/yr)
PPG Industries	Lake Charles, LA	0.10	62
Dow Chemical	Freeport, TX and Plaquemine, LA	0.48	384

TABLE 5-3

Yearly Consumption of Vinylidene Chloride in Different Industries^a

Industry	Amount Consumed (million lb/yr)	Percent of Total
Polymers		
Domestic Use	108-115	81%
Export	20	15%
Chemical Intermediate	5	4%
TOTAL	133-140	100%

^aSource: Neufeld et al., 1977

It can be seen from Table 5-3 that except for a small application as a chemical intermediate, most of the vinylidene chloride is consumed domestically for the production of polymers. Because of its excellent fire and barrier resistance properties, vinylidene chloride polymer has been used in the manufacture of copolymers with other monomers. In general, copolymers that contain a high percentage of vinylidene chloride (70-95%) are used for the barrier resistance characteristic, and copolymers with a lower percentage of vinylidene chloride (10-40%) are used for fire retardant properties. The three different polymerization processes used industrially for the production of polymeric materials and the uses of these polymers are shown in Table 5-4.

Emulsion latexes used as barrier coatings usually contain 70 to 95% vinylidene chloride copolymerized with a variety of other monomers such as vinyl chloride, acrylic acid, and acrylonitrile. These copolymers have a unique and excellent barrier resistance to gases, water vapor, organic vapors, and odors. The speciality latexes obtained by copolymerization of 10 to 40% vinylidene chloride and butadiene-styrene have excellent fire retardant properties and are used as carpet backing materials.

The solid resins obtained by converting emulsion or suspension latexes to the powdered resin may contain different amounts of vinylidene chloride. As in the case of latexes, the characteristics of the copolymer are strongly influenced by the vinylidene chloride content. Solid resins are used for coating cellophane and extrusion into film or for manufacturing multilayer laminates.

Solution polymerization of 10 to 30% vinylidene chloride with acrylonitrile produces modacrylic fibers. The modacrylics are produced solely for the flame retardant characteristics they impart to sleepwear fabric, drapery fabric, and automobile upholstery.

TABLE 5-4

Polymerization Processes, Products, and Their Applications^a

Process	Product	Application
Emulsion Polymerization	Emulsion Latex	Barrier coating
	Speciality Latex	Carpet backing
Suspension Polymerization	Resin	Barrier coating on multilayer laminate
		Molded plastics
		Extruding films and pipes
Solution Polymerization	Fiber	Modacrylic fibers

^aSource: Neufeld et al., 1977

The major plant site for vinylidene chloride polymerization and the type of polymer produced by these facilities are shown in Table 5-5.

The estimated annual emissions of vinylidene chloride from polymer synthesis are given in Table 5-6.

5.1.3 Processing or Fabrication of Polymers

The fabrication processes include coating of paper, glassine, paper products, cellophane, and plastic films. Fabrication processes also include extrusion of films and monofilament fibers and molding operations.

The major polyvinylidene chloride latex users for miscellaneous fabrication processes are given in Tables 5-7, 5-8, and 5-9.

The estimated annual emission rate of vinylidene chloride in the processing of polyvinylidene chloride polymers is given in Table 5-10.

5.1.4 Storage, Handling, and Transportation of the Monomer

Vinylidene chloride monomer is shipped from the production sites to the user site in bulk railroad tank cars or tank trucks. A small unspecified quantity is shipped in drums (Neufeld et al., 1977). Approximately 90 million pounds of vinylidene chloride is transported annually in various ways. Vinylidene chloride emissions can be expected to occur during storage, transfer, and filling operations. It has been estimated that of the total vinylidene chloride loss during its manufacture, subsequent polymerization, and processing, 25% takes place during storage, transfer, and handling and 75% takes place during processing (Neufeld et al., 1977). Since only half of the total vinylidene chloride manufactured is transported to other locations (the other half is captively used), this half is solely responsible for losses related to storage, transportation, and handling. The total vinylidene chloride loss during manufacture, subsequent polymerization, and processing adds up to 1,160,000

TABLE 5-5

Polymerization Sites and Type of Polymer Produced^a

Manufacturer	Location	Polymer Type	Primary Usage
Dow Chemical	Midland, MI	Suspension latex	Barrier coating
		Emulsion latex	Barrier coating
		Solid resin	Barrier coating
		Speciality latex	Carpet backing
W.R. Grace	Owensburg, KY	Emulsion latex	Barrier coating
		Speciality latex	Carpet backing
Morton Chemicals	Ringwood, IL	Emulsion latex	Barrier coating
A.E. Staley	Lemont, IL	Emulsion latex	Barrier coating
Rohm and Haas	Knoxville, TN	Emulsion latex	Barrier coating
GAF	Chattanooga, TN	Emulsion latex	Barrier coating
		Speciality latex	Carpet backing
Reichhold Chemicals	Cheswold, DE	Emulsion latex	Barrier coating
		Speciality latex	Carpet backing
National Starch	Meridosia, IL	Emulsion latex	Barrier coating
		Speciality latex	Carpet backing
Eastman	Kingsport, TN	Modacrylics	Fiber
Monsanto	Decatur, AL	Modacrylics	Fiber
American Cyanamid	Pensacola, FL	Modacrylics	Fiber
DuPont	Circleville, OH	Solid resin	Barrier coating

^aSource: Neufeld et al., 1977

TABLE 5-6

Estimated Annual Emissions of Vinylidene Chloride from Polymer Synthesis^a

Polymer	Emission Factor (lb/100 lb vinylidene chloride polymerized)	Annual Emissions (thousands of lb/yr)
Barrier-coating latex	0.60	120
Latex for miscellaneous coatings	1	150
Synthetic fibers	1	160
Resin for cellophane coating	0.7	182
Extrusion resin (emulsion)	0.13	27
Extrusion resin (suspension)	0.26	40
TOTAL		679

^aSource: A.D. Little, Inc., 1976

TABLE 5-7

Major Polyvinylidene Chloride Processors^a

Company	Plant Location
Barrier-Coating on Paper and Glassine	
Deerfield-Reed Corporation	Clifton, NJ
American Bag and Paper	Philadelphia, PA
Consolidated Paper	Wisconsin Falls, WI
Crown Zellerbach	Portland, OR
Daniels	Rhineland, WI
Diversa-Pax	St. Petersburg, FL
Dixico	Dallas, TX
DuPont	Circleville, OH
Milprint	Milwaukee, WI
Philip Morris	Nicholasville, KY
Rexham	Memphis, TN
St. Regis	Rhineland, WI
Thilmany	Kaukauna, WI
Chase Bag Company	Hudson Falls, NY
Print Pak	Atlanta, GA
Barrier-Coating on Paperboard	
Gordon Carton	Baltimore, MD
Green Bay Packaging	Green Bay, WI
Interstate Folding Box	Middletown, OH
Michigan Carton Company	Battle Creek, MI
Olin Kraft	W. Monroe, LA
Zumbril	Cincinnati, OH
Barrier-Coating on Plastic	
DuPont	Circleville, OH
Cryovac	Simpsonville, SC
Minnesota Mining and Manufacturing	Irvington, NJ
	Decatur, AL
Allied Chemical Corporation	Pittsville, PA
Hercules	Convington, VA
Milprint	Milwaukee, WI
Cryovac	Simpsonville, SC
Bemis	New London, WI
American Can Company	Neenah, WI
Standard Packaging	Clifton, NJ
Sealed Air Corporation	Fairlawn, NJ

^aSource: Neufeld et al., 1977

TABLE 5-8

Manufacturers of Polyvinylidene Chloride-Coated Cellophane^a

Manufacturer	Share of Market	Location
DuPont	40%	Richmond, VA Clinton, IA Tecumseh, KS
FMC	35%	Fredericksburg, VA
Olin	25%	Pisgah Forrest, NC Covington, VA

^aSource: Neufeld et al., 1977; A.D. Little, Inc., 1976

TABLE 5-9
Major Extruders of Polyvinylidene Chloride Film^a

Manufacturer	Plant Location
Cryovac	Simpsonville, SC Cedar Rapids, IA Camarillo, CA Iowa Park, TX
Oscar Mayer	Madison, WI Chicago, IL Davenport, IA Philadelphia, PA Nashville, TN Vernon, CA Sherman, TX
Dow Chemical	Midland, MI
American Can Company	Cleveland, OH
Union Carbide Corporation	Centerville, IA
Amtech, Inc. ^b	Odenton, MO

^aSource: Neufeld et al., 1977; A.D. Little, Inc., 1976

^bThis plant extrudes monofilament, which is used primarily as filter cloth in the chemical industry.

TABLE 5-10

Estimated Emissions from Polyvinylidene Chloride Processing^a

Process	Emission Factor (lb/100 lb Polymer Used)	Annual Vinylidene Chloride Emissions (lb/yr)
Coating cellophane	0.06	1,560
Coating plastics, paper and glassine	0.03	16,380
Extrusion	0.001	430
Miscellaneous coating	0.04	12,000
TOTAL		30,370

^aSource: A.D. Little, Inc., 1976

pounds per year. Half of this figure has been used in the following calculation to determine the losses during storage, transportation, and handling:

$$1,160,000 \times 0.5 \times 25\% = x$$

$$x = 145,000 \text{ lb/yr}$$

This annual loss of 145,000 pounds during storage, handling, and transportation does not include any additional losses due to accidental spillage during transportation.

5.1.5 Chemical Intermediate Production

The primary chemical intermediate that is manufactured from vinylidene chloride is chloroacetyl chloride. Since PPG Industries, the sole manufacturer of 1,1,1-trichloroethane from vinylidene chloride, no longer uses this process for 1,1,1-trichloroethane manufacture, no vinylidene chloride emissions can be expected from this intermediate. Chloroacetyl chloride is captively used by Dow Chemical Company and no losses of vinylidene chloride to the environment are reported during its manufacture (Neufeld et al., 1977). The purified chloroacetyl chloride used as an intermediate for further chemical reaction is reported to be free of vinylidene chloride contamination (Neufeld et al., 1977). Thus, its usage cannot be a source of vinylidene chloride in air.

5.1.6 Incineration of Polymers Containing Polyvinylidene Chloride

Over 95% of vinylidene chloride polymers are used for packaging, in the manufacture of textile fibers, or as a flame-retardant carpet backing (Neufeld et al., 1977). All these consumable items are disposed of at the end of their useful life. The ultimate disposal operation is either incineration or placement in a solid waste landfill. If incineration is used as the disposal process, it may contribute to small amounts of vinylidene chloride in air. The residual monomer in the polymer may be released into the air during heating. The decomposition of polymer to monomer is not likely to occur. The products of thermal

degradation will depend primarily on the heating temperature. Thermal degradation of polyvinylidene chloride has been studied in detail by Wessling and Edwards (1970). These investigators found that polyvinylidene chloride begins to decompose at about 125°C. At very high temperatures, graphitization of carbonaceous residue takes place. On pyrolysis, polyvinylidene chloride yields 70% hydrogen chloride, 26% carbonaceous ash, and 4% volatile hydrocarbons consisting of benzene, chlorobenzene, and three dichlorobenzenes (O'Mara et al., 1973). The thermal decomposition is also catalyzed by various metal salts. The mechanism of thermal decomposition of polyvinylidene chloride was also studied by Ballistreri et al. (1981). Currently an estimate of the amount of vinylidene chloride released to the air from incineration is not available. However, incineration is not expected to be a significant source of vinylidene chloride in air.

5.2 SOURCES IN WATER

The only industrial operation that produces liquid effluents containing vinylidene chloride monomer is reported to be the polymerization operation (Neufeld et al., 1977). The two main sources of vinylidene chloride loss during polymerization operations are waste polymer sludges arising from reactor cleaning during normal operation or from products containing bad batches, and wastewater from condensation of gases during stripping operations. Neufeld et al. (1977) estimate that approximately 4100 pounds per year of vinylidene chloride loss can be attributed to this source.

A small but unknown amount of vinylidene chloride may originate in water from the cleaning of rugs backed with speciality latexes.

5.3 SOURCES IN SOIL

The disposal of consumer products that contain processed vinylidene chloride polymers (1 to 3 ppm residual vinylidene chloride monomer) by municipal

solid waste disposal operations may result in soil contamination. Neufeld et al. (1977) estimate that the maximum potential quantity of vinylidene chloride monomer in solid wastes would be 180 pounds per year.

5.4 SOURCES IN FOODS

The three potential sources of vinylidene chloride contamination in foods are as follows: (1) migration of monomer from the polymeric wrapping materials to foods, (2) contaminated drinking water, (3) edible aquatic foods that have bioconcentrated vinylidene chloride from polluted water. All these are secondary sources, however, whose primary sources (i.e., polymers and water) have already been discussed.

5.5 SUMMARY OF ENVIRONMENTAL LOSSES

This section presents estimates of the amount of vinylidene chloride monomer lost to the atmosphere, to water streams, or to soils. The losses are summarized in Table 5-11.

TABLE 5-11

Summary of Estimated Environmental Losses of Vinylidene Chloride

Source	Vinylidene Chloride Contamination (lb/yr)		
	Air	Water	Soil
Monomer Synthesis	446,000	0	0
Polymer Synthesis	679,000	4100	0
Polymer Fabrication	30,400	0	0
Storage, Handling, and Transportation	145,000	0 ^a	0 ^a
Chemical Intermediate Production	0	0	0
Disposal of Polymers	0 ^b	0	180
TOTAL	1,300,400	4100	180

^aSome water and soil contamination may arise from this source.

^bIncineration of polymers may contaminate air with an undeterminable amount.

6. ENVIRONMENTAL FATE, TRANSPORT, AND DISTRIBUTION

6.1 ATMOSPHERIC FATE, TRANSPORT, AND DISTRIBUTION

6.1.1 Reaction with Atmospheric Radicals and Ozone

Very little information is available on this subject. Upon entering the atmosphere, vinylidene chloride, like other chloro-olefin vapors, may interact with $O\cdot$, $OH\cdot$, and $RO_2\cdot$ radicals, and with ozone already present in the atmosphere. In addition, chloro-olefins in the atmosphere will generate chlorine atoms, which may interact with the parent molecules. The gas phase oxidation of halo-olefins at room temperature initiated by Cl atoms, $O(^3P)$ radicals, and O_3 has been reviewed (Sanhueza et al., 1976; Heicklen et al., 1975). In the presence of O_2 , the oxidation of vinylidene chloride by chlorine atoms produced 98% chloroacetyl chloride, and about 2% phosgene and CO as the reaction products. The reaction of vinylidene chloride with $O(^3P)$ in the presence of O_2 produced CO, chloroacetyl chloride, polymer, and another unidentified compound. The rate of $O(^3P)$ attack on vinylidene chloride is the same as the rate of $O(^3P)$ attack on C_2H_4 (Sanhueza, 1976; Heicklen et al., 1975); however, the relative reactivity of $O(^3P)$ with vinylidene chloride (1.0) in the gas phase at room temperature is higher than the reactivity of Cl atoms with vinylidene chloride (0.72) under the same conditions (Heicklen et al., 1975). Based on a reaction rate constant of $7.8 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ sec}^{-1}$ (Graedel, 1978) for $O(^3P)$ reaction with vinylidene chloride (same as C_2H_4 reaction rate) and the concentration of $O(^3P)$ in the ambient atmosphere as $5 \times 10^4 \text{ molecules cm}^{-3}$ (Cupitt, 1980), the half-life for this reaction can be estimated to be approximately 206 days.

In the absence of O_2 , $O(^3P)$ forms the same reaction products it produces in the presence of O_2 ; however, the quantum yield for carbon monoxide formation is

reduced from 0.78 in the presence of O_2 to 0.35 in the absence of O_2 (Heicklen et al., 1975).

The main products of O_3 interaction with vinylidene chloride in the vapor phase at 25°C are phosgene and formaldehyde (Sanhueza et al., 1976; Heicklen et al., 1975; Brown et al., 1975). The formation of chloroacetyl chloride, CO, CO_2 , HCl, and possibly water has also been reported (Hull et al., 1973). The yield of phosgene is always either comparable to or greater than the yield of formic acid (Heicklen et al., 1975). The O_3 reaction in the presence of O_2 is, however, too slow to be important in the urban atmosphere (Sanhueza et al., 1976; Heicklen et al., 1975). Based on a reaction rate constant of $4 \times 10^{-20} \text{ cm}^3 \text{ molecule}^{-1} \text{ sec}^{-1}$ (Cupitt, 1980) and the concentration of ozone in the ambient atmosphere at $1 \times 10^{12} \text{ molecule cm}^{-3}$ (Cupitt, 1980), the half-life for ozone reaction with vinylidene chloride in the atmosphere can be calculated to be approximately 201 days.

The interaction of RO_2 and OH radicals with vinylidene chloride was reported by Brown et al. (1975). The formation of phosgene and formaldehyde was reported with both RO_2 and OH radicals. The rate constant for the oxidation with OH radicals was reported to be $4 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ sec}^{-1}$ (Cupitt, 1980). If the concentration of OH radicals in the ambient atmosphere is assumed to be $10^{-6} \text{ radicals cm}^{-3}$ (Cupitt, 1980), the half-life for this reaction can be calculated to be 2 days. The half-life for peroxy radicals reaction with vinylidene chloride in the atmosphere has been reported to be about 22 years by Brown et al. (1975).

It can be concluded from the above discussion that the most significant oxidation reaction of vinylidene chloride in the ambient atmosphere is its reaction with hydroxyl radicals. The half-life for this reaction has been estimated to be about 2 days.

6.1.2 Atmospheric Photochemical Reactions

Gay et al. (1976) studied the photoreaction of vinylidene chloride in air in the presence of nitrogen dioxide with ultraviolet light. At a concentration of 4.85 ppm and in the presence of 2.26 ppm NO_2 , vinylidene chloride was found to decompose rapidly--83% decomposition had taken place within 140 minutes. The reaction products identified were HCOOH , HCl , CO , HCHO , O_3 , COCl_2 , chloroacetyl chloride, formyl chloride, and nitric acid.

The photolysis of vinylidene chloride under simulated atmospheric conditions was also studied by Dilling et al. (1976). In the presence of 5 ppm NO and at a relative humidity of 35%, 10 ppm vinylidene underwent photolysis at wavelengths greater than 290 nm, and disappeared with a half-life of 2.1 hours. Addition of cyclohexane was found to retard the photodecomposition rate of vinylidene chloride. Although vinylidene chloride was not tested, reaction of a closely related halo-olefin (trichloroethene) showed similar reactivity with ozone and NO_2 as with NO .

The above experiment of Dilling et al. (1976) was conducted at ultraviolet light intensity of about 2.6 times that of natural sunlight at noon on a summer day in Freeport, TX. Therefore, the estimated half-life of vinylidene chloride under bright sunlight and in the presence of NO is approximately 5 to 12 hours (Dilling et al., 1976). In the absence of NO , the half-life for photodecomposition of vinylidene chloride under bright sunlight may increase five-fold. This estimate is based on the experimental half-life of a closely-related halo-olefin (trichloroethene) in the presence and absence of NO as determined by Dilling et al. (1976). Therefore, it is estimated that vinylidene chloride will photodecompose under bright sunlight with a half-life of one to three days. Pearson and McConnell (1975), on the other hand, exposed quartz flasks containing halogenated aliphatic compounds in air to incident radiations from the sun and

estimated the tropospheric photolytic half-life of 56 days for vinylidene chloride. Although the experiments of Pearson and McConnell (1975) were not adequately controlled in terms of air composition, temperature, and light intensity, it is difficult to explain the large discrepancy in the photolytic half-life of vinylidene chloride between this investigation and the work of Dilling et al. (1976). That the photolysis of vinylidene chloride in the atmosphere is a possibility has also been indicated (but no quantitative rate data given) by Cupitt (1980). In conclusion, it can be predicted that photolysis of vinylidene chloride in the atmosphere may be competitive with its reaction involving hydroxyl radicals. The half-life of vinylidene chloride in air under atmospheric smog conditions (air containing NO_x , O_3 , etc.) may be further reduced due to its greater photochemical reactivity in the presence of NO_x and O_3 .

6.1.3 Atmospheric Physical Processes

Recently, the atmospheric fate of vinylidene chloride under physical and chemical removal processes was studied by Cupitt (1980). The atmospheric half-life of vinylidene chloride removal by rain droplets was estimated to be 110,000 years. The lifetime of vinylidene chloride removal by adsorption onto aerosol particles was estimated to be 5.3×10^{10} days. Therefore, the possibility of vinylidene chloride removal by these two physical removal processes is remote.

6.2 AQUATIC FATE, TRANSPORT, AND BIOACCUMULATION

Very little experimental data pertaining to the various fate processes of this chemical in aquatic media and its bioaccumulation potential in aquatic organisms are available in the literature. In the following discussion the experimental data whenever available have been used to predict the fate of vinylidene chloride in aquatic media. In the absence of such data, the prediction of the fate has been based on calculated or estimated data using empirical or theoretical methods. Extreme caution should be exercised in

interpreting the fate of a chemical based solely on calculated or estimated data. This reservation concerning the estimated data is justified by the large discrepancies observed in a few instances between the experimental and estimated data where both data are available.

6.2.1 Fate and Transport in Water

The photolysis and hydrolysis of vinylidene chloride in natural aquatic media are probably not significant processes (Mabey et al., 1981). No experimental data on the possibility of oxidation of vinylidene chloride by singlet oxygen (O_2) and peroxy radicals present in aquatic media could be found. However, based on the experimental data from closely related analogues (e.g., trichloroethene and tetrachloroethene), Callahan et al. (1979) concluded that vinylidene chloride is "more amenable" to oxidation in aquatic media than both trichloroethene and tetrachloroethene. From the calculated (based on structure/activity relationship) rate constant values of less than $10^8 \text{ M}^{-1} \text{ hr}^{-1}$ and $3 \text{ M}^{-1} \text{ hr}^{-1}$ (Mabey et al., 1981) for singlet oxygen and peroxy radical reactions, respectively, with vinylidene chloride in aquatic media and the estimated concentrations of 10^{-12} M and 10^{-9} M for singlet oxygen and peroxy radicals in aquatic media (Mill and Mabey, 1980), the oxidation of vinylidene chloride by either process can be calculated to be environmentally insignificant.

The biodegradability of vinylidene chloride was studied by Tabak et al. (1981) by the static-culture flask-screening procedure with settled domestic wastewater as microbial inoculum. These authors reported a significant degradation of vinylidene chloride with the unacclimated sludge. At 5 mg/l initial concentration, 78% of the compound biodegraded in 7 days of incubation at 25°C. The biodegradation was only 45% when the initial concentration was 10 mg/l and the other conditions were the same. Similarly, it has been reported by

Patterson and Kodukala (1981) that activated sludge treatment of a municipal wastewater at an influent vinylidene chloride level of 0.04 mg/l resulted in 97% removal of the compound. It can be concluded from these experimental results that vinylidene chloride may undergo biodegradation in ambient aquatic media. However, until the role of biodegradation of this compound in natural aquatic media is defined, it is not possible to predict the significance of this process in determining its fate in aquatic media. It should be mentioned that on the basis of limited experimental evidence on related chloroaliphatics, Callahan et al. (1979) have concluded that vinylidene chloride probably biodegrades in aquatic media at a slow rate.

The transport of vinylidene chloride from aquatic media to the atmosphere through volatilization appears to be its primary transport process (Callahan et al., 1979). The evaporation characteristics of vinylidene chloride in water were studied by Dilling (1977). When a 1 mg/l aqueous solution of 6.5 cm depth was stirred at 200 rpm in still air (<0.2 mph air current) at 25°C, the half-life for evaporation was found to be 27.2 minutes. The measured half-life for evaporation of vinylidene chloride from dilute aqueous solution was reasonably close to the calculated value of 20.1 minutes (Dilling, 1977) based on liquid and vapor-phase mass transfer coefficients for exchange between the water body and the atmosphere. However, this agreement is somewhat fortuitous since the calculated value depends on the exchange constants, whereas the experimental values depend on the stirring rate, air current, and other factors (Dilling, 1977).

The half-life for the evaporation of vinylidene chloride from pond, river, and lake waters can also be calculated from the reaeration rate constant of this compound. Assuming the reaeration rate ratios (K_v^c/K_v^o) of 0.601 (calculated from diffusion coefficients) (Mabey et al., 1981) for vinylidene chloride, and the values for oxygen reaeration rate constants of 0.19 day^{-1} , 0.96 day^{-1} , and

0.24 day⁻¹ for pond, river, and lake waters, respectively (Mabey et al., 1981), the half-lives for evaporation can be calculated to be 6.1 days, 1.2 days, and 4.2 days from pond, river, and lake waters, respectively.

No specific information pertaining to the sorption of vinylidene chloride onto sediments could be found. However, based on the experimental data from related analogues (e.g., trichloroethene and tetrachloroethene), Callahan et al. (1979) concluded that the sorption of vinylidene chloride onto sediments is probably an insignificant process. The significance of sorption of this compound onto sediments can also be predicted from the correlation equation of Schwarzenbach and Westall (1981). Assuming the octanol/water partition coefficient value of 69 for vinylidene chloride (Mabey et al., 1981), the log K_{oc} value ($K_{oc} = \frac{\text{sorption coefficient} \times 100}{\% \text{ organic carbon}}$) in sediments containing 0.5% organic carbon can be calculated to be 1.81 from the equation of Scharzenbach and Westall (1981). This log K_{oc} value is somewhat lower than the experimental log K_{oc} value of 1.92 for benzene (Schwarzenbach and Westall, 1981). Therefore, the rate of sorption of vinylidene chloride onto sediments is expected to be less than benzene sorption onto sediments. In other words, the sorption of vinylidene chloride onto sediments containing low organic carbon is not likely to be significant but may increase as the organic carbon content in the sediment increases.

It can be concluded from the above discussions that volatilization from aquatic media is probably the most significant process for vinylidene chloride. The role of biodegradation in determining the fate of this compound in natural aquatic media has not been definitely determined. All other aquatic processes are probably insignificant in determining the fate of vinylidene chloride in natural aquatic media.

6.2.2 Bioaccumulation of Vinylidene Chloride in Aquatic Organisms

No experimental data are available for the bioaccumulation factor (BCF) for vinylidene chloride in any kind of fish or other edible aquatic organisms. A number of theoretical correlation equations are available that relate the BCF to either the octanol-water partition coefficient or the water solubility of the chemical. These equations are given below:

$$\text{Log BCF} = 0.76 \log K_{ow} - 0.23 \text{ (Veith et al., 1979)}$$

$$\text{Log BCF} = 0.542 \log K_{ow} + 0.124 \text{ (Neely et al., 1974)}$$

$$\text{Log BCF} = -0.508 \log S + 3.41 \text{ (Chiou et al., 1977)}$$

where K_{ow} = partition coefficient of the chemical between octanol-water,
and

S = water solubility of the chemical expressed in $\mu\text{mol/l}$.

The above equation of Veith et al. (1979) is applicable for the whole fish, whereas the equations of Neely et al. (1974) and Chiou et al. (1977) are applicable for fish muscle only.

If the values of $\log K_{ow}$ and S for vinylidene chloride are assumed to be 1.48 (Tute, 1971) and $2.19 \times 10^3 \mu\text{mol/l}$ (Hushon and Kornreich, 1978), respectively, the theoretical values for BCF can be calculated to be 7.8 (equation of Veith et al., 1979) for the whole fish and 8.5 (equation of Neely et al., 1974) or 51 (equation of Chiou et al., 1977) for fish muscle. The relatively low value for BCF indicates that vinylidene chloride will probably not bioaccumulate in fish to any significant extent.

6.3 FATE, PERSISTENCE, AND TRANSPORT IN SOIL

No information pertaining specifically to the fate, persistence, and transport of vinylidene chloride in soils could be found. However, the fate and transport processes of this compound can be partially predicted from the estimated $\log K_{oc}$ value of 1.81 in soils containing 0.5% organic carbon. This

relatively low $\log K_{oc}$ value, coupled with a solubility of 2250 mg/l and a vapor pressure of 600 mm of mercury at 25°C would indicate that both volatilization and leaching may play significant roles in determining the fate of this chemical in soils. In fact, the detection of this compound in several groundwaters (see Section 7.2) is indicative of the leaching of vinylidene chloride from soils. Unless more experimental or calculated data are available, no prediction regarding the microbial degradability and chemical oxidation of this chemical in soils can be made, although Kobayashi and Rittmann (1982) have reported the biodegradation of vinylidene chloride by soil bacteria.

7. ENVIRONMENTAL LEVELS AND EXPOSURE

When an environmental pollutant is suspected of being a potential health hazard, it becomes necessary to determine the level of that pollutant in the environment and its possible human intake from various environmental media. The four environmental media of interest in this regard are air, water, soil, and foods. The vinylidene chloride levels from each individual medium and the possible human exposure therefrom are discussed in the following subsections.

The purpose of this document is to present available information relevant to human health effects that could be caused by this substance. Any information regarding sources, emissions, ambient air concentrations, and public exposure has been included only to give the reader a preliminary indication of the potential presence of this substance in the ambient air. While the available information is presented as accurately as possible, it is acknowledged to be limited and dependent in many instances on assumption rather than specific data. This information is not intended, nor should it be used, to support any conclusions regarding risks to public health.

If a review of the health information indicates that the Agency should consider regulatory action for this substance, a considerable effort will be undertaken to obtain appropriate information regarding sources, emissions, and ambient air concentrations. Such data will provide additional information for drawing regulatory conclusions regarding the extent and significance of public exposure to this substance.

7.1 AIR

7.1.1 Environmental Levels

The concentration of vinylidene chloride in air can be determined by monitoring its level in the atmosphere. In the absence of monitored data, the

concentration can be theoretically estimated by dispersion modelling based on the amount of vinylidene chloride losses in air from each plant. Such theoretical modelling, which is generally recognized to be accurate within a factor of two in predicting the concentration of a pollutant in the vicinity of a plant, has been used for estimating the population exposure to other atmospheric pollutants, such as acrylonitrile (Suta, 1979). A vinylidene chloride model based on analogy to 1,1,1-trichloroethane has been developed to estimate potential human exposure (Wapora, 1982). Using dispersion modelling, Hushon and Kornreich (1978) estimated the vinylidene chloride concentration levels at a single distance from three vinylidene chloride monomer production sites in the United States using atmospheric dispersion modeling. The 10-minute peak exposures of an individual 500 m downwind from the three plants were estimated to be 0.87 ppm, 0.44 ppm, and 0.52 ppm for the Dow plant at Freeport, the Dow plant at Plaquemine, and the PPG plant at Lake Charles, respectively. The corresponding 24-hour peak concentrations at the same three locations were estimated to be 0.52 ppm, 0.26 ppm, and 0.32 ppm. In order to estimate these values, it was assumed that these plants have certain yearly emission rates of vinylidene chloride (Hushon and Kornreich, 1978), the wind speed was 6 m/s, the height of the plants was 12.3 m, and vinylidene chloride monomer would be non-reactive in the atmosphere. The last assumption may have lead to erroneous estimates of vinylidene chloride concentration at longer distances. The half-life of vinylidene chloride under simulated smog conditions was estimated to be 5 to 12 hours (Dilling et al., 1976) and approximately 2 days under atmospheric hydroxyl reactions (Cupitt, 1980). This relatively rapid atmospheric oxidation may not have a significant effect on ambient air concentration at short distances from the source, but may have a dramatic effect at longer distances. Thus, any dispersion modelling for estimation of vinylidene chloride concentrations at various

distances from the source should include the effect of disappearance of vinylidene chloride due to atmospheric oxidation.

One of the earlier attempts to experimentally determine atmospheric levels of halocarbons including vinylidene chloride was made by Grimsrud and Rasmussen (1975) in rural Pullman, WA, during the period of December 1974 to February 1975. They used a GC-MS method for the quantification of vinylidene chloride. Because of the limitations of the analytical methodology (inadequate GC column), however, they could not separate vinylidene chloride from its isomers cis- and trans-1,2-dichloroethylene, and reported the combined concentration of the three isomers to be <5 ppt.

The monitoring of vinylidene chloride near industrial sites was conducted by Research Triangle Institute (Research Triangle Institute, 1977a,b, cited in Hushon and Kornreich, 1978) during the period 1976 to 1977. Vinylidene chloride concentrations ranging from 9 ppt to 249 ppt were detected near Dow plant sites, one at Freeport, TX, and four at Plaquemine, LA. Seven other Dow sites at Plaquemine, LA, showed no detectable trace of vinylidene chloride.

The more systematic environmental monitoring for vinylidene chloride near industrial sites was performed by Going and Spigarelli (1977) during the period of January 1977 to April 1977. These investigators selected sampling points near six industrial plants representing manufacturers of vinylidene chloride monomers and polymers and fabricators of the polymers. The analyses of the samples were performed by gas chromatography-flame ionization detectors (GC-FID) and gas chromatography-mass spectrometry (GC-MS).

The concentrations of vinylidene chloride in the ambient air throughout the U.S. have been measured by several other investigators. The ambient atmospheric levels of vinylidene chloride throughout the continental U.S. as determined by these and other investigators are shown in Table 7-1. It should be noted that

TABLE 7-1

Vinylidene Chloride Concentrations in the Ambient Air Throughout the Continental U.S.

Location	Sampling Date (duration)	Mean Conc. (ppt)	Std. ^a Dev. (ppt)	Data ^b Quality	Reference
Aldine, TX	1977 (88 days)	36	62	4	Pellizzari et al., 1979
Baton Rouge, LA	1977 (78 days)	ND	0	4	Pellizzari et al., 1979
Batsto, NJ	1979 (306 days)	ND	0	4	Bozzelli et al., 1980
Beaumont, TX (grab sample)	1980	12,000	NR	4	Wallace, 1981
Bridgeport, NJ	1977 (1 hour)	ND	0	4	Pellizzari and Bunch, 1979
Bristol, PA	1977 (1 hour)	ND	0	4	Pellizzari and Bunch, 1979
Camden, NJ	1979 (198 days)	ND	0	4	Bozzelli et al., 1980
Chapel Hill, NC (grab sample)	1980	2,800	NR	4	Wallace, 1981
Charleston, WV	1977 (53 days)	ND	0	4	Pellizzari, 1978b
Denver, CO	1980 (10 days)	6.4	21	3	Singh et al., 1981a
Edison, NJ	1976 (98 days)	ND	0	4	Pellizzari et al., 1979
Elizabeth, NJ	1979 (347 days)	ND	0	4	Bozzelli et al., 1980
Freeport, TX	1976 (2 hours)	130	NR	4	Pellizzari et al., 1979

TABLE 7-1 (cont.)

Location	Sampling Date (duration)	Mean Conc. (ppt)	Std. ^a Dev. (ppt)	Data ^b Quality	Reference
Front Royal, VA	1977 (29 days)	49	53	4	Pellizzari, 1978b
Geismar, LA (grab sample)	1977	5	NR	4	Pellizzari, 1978a
Houston, TX	1977 (1091 days)	16	15	3	Singh et al., 1981a
	1977	ND	0	4	Pellizzari et al., 1979
Institute, WV	1977 (1 day)	ND	0	4	Pellizzari et al., 1978b
Lake Charles, LA	1977 (1 day)	6,700	73	2	Going and Spigarelli, 1977
Liberty Mound, OK	1977 (42 days)	ND	0	4	Pellizzari, 1978b
Marcus Hook, PA	1977 (1 hour)	ND	0	4	Pellizzari and Bunch, 1979
Midland, MI	1977 (1 day)	6,100	6	2	Going and Spigarelli, 1977
N. Philadelphia, PA	1977 (1 hour)	ND	0	4	Pellizzari and Bunch, 1979
Newark, NJ	1979 (342 days)	ND	0	4	Bozzelli et al., 1980
Nitro, WV	1977 (51 days)	ND	0	4	Pellizzari, 1978b
Plaquemine, LA	1977 (29 days)	1,100	69	2	Going and Spigarelli, 1977
	1977	30	NR	4	Pellizzari et al., 1979
Riverside, CA	1980 (10 hours)	6.5	3	3	Singh et al., 1981a

TABLE 7-1 (cont.)

Location	Sampling Date (duration)	Mean Conc. (ppt)	Std. ^a Dev. (ppt)	Data ^b Quality	Reference
Rutherford, NJ	1979 (348 days)	ND	0	4	Bozzelli et al., 1980
S. Charleston, WV	1977 (51 days)	ND	0	4	Pellizzari, 1978b
S. Amboy, NJ	1977 (336 days)	ND	0	4	Bozzelli et al., 1980
St. Louis, MO	1980 (9 days)	3.6	1.7	3	Singh et al., 1981a
St. Albans, WV	1977 (28 days)	ND	0	4	Pellizzari, 1978b
Tulsa, OK	1977 (72 days)	ND	0	4	Pellizzari, 1978b
Vera, OK	1977 (3 hours)	ND	0	4	Pellizzari, 1978b
W. Belle, WV	1977 (51 days)	ND	0	4	Pellizzari, 1978b

^aStandard deviation

^bData quality; 1: excellent, 2: good, 3: acceptable, 4: questionable
ND = Not Detected; NR = Not Reported

the data presented in Table 7-1 represent vinylidene chloride levels in air samples for various durations, and from different types of sites, namely, rural/remote areas, urban/suburban areas, and source areas. An assessment of the quality of data presented in Table 7-1 was made by Brodzinsky and Singh (1982) on the basis of the error limits of the data. Some of the investigators did not provide adequate data to calculate the error limit. In these cases, the quality codes were assigned on the basis of expert judgement of Brodzinsky and Singh (1982). The quality codes of 1, 2, 3, and 4 signify excellent, good, acceptable and questionable data, respectively. The mean concentrations of vinylidene chloride in the ambient air samples collected in 1979 from Los Angeles, CA, Phoenix, AZ, and Oakland, CA, were measured to be 4.9 ppt, 29.8 ppt, and 12.6 ppt, respectively (Singh et al., 1981b). These values have not been presented in Table 7-1 because no assessment of the quality of these data was made by Brodzinsky and Singh (1982).

The mean ambient air concentrations of vinylidene chloride, the averaging times (duration), the standard deviations, and the data qualities for all the data categorized by the types of sites are given in Table 7-1. Data with assessed qualities of excellent, good, and acceptable are presented in Table 7-2 in parentheses. Additionally, the median vinylidene chloride levels in air samples from different site types for which the determined data have been assessed to be acceptable or better also are shown in Table 7-2. It is obvious from Table 7-2 that the median concentration of vinylidene chloride in U.S. urban/suburban ambient air is approximately 5 ppt. However, this value is substantially higher (3600 ppt) for ambient air samples collected from vinylidene chloride source areas.

TABLE 7-2

Mean and Median Vinylidene Chloride Concentrations in U.S. Ambient Air
of Different Site Types^a

Site Type	No. of Samples ^b	Mean Conc. (ppt)	Std. Dev.	Median Conc. ^c (ppt)	Data Quality
Grand totals	767 (339)	260 (170)	1500 (930)	5	3.5 (3)
Rural/remote	2 (0)	ND (0)	0 (0)	0	4.0 (0)
Urban/suburban	665 (325)	220 (8)	1500 (14)	5	3.5 (3)
Source areas	100 (14)	540 (3800)	1700 (2800)	3600	3.7 (2)

^aSource: Brodzinsky and Singh, 1982; averaging times ranged from 1 hour to 1091 days.

^bThe values in parentheses refer only to data with quality of 3 or lower.

^cThe median concentrations refer only to data with quality of 3 or lower.

7.1.2 Exposure

The Office of Air Quality Planning and Standard (OAQPS), U.S. EPA, Research Triangle Park, NC, has prepared a document estimating human exposure to vinylidene chloride in the atmosphere around plants producing the monomer and the polymer. The ranges of measured concentrations associated with the monomer and polymer plants were 90-100 $\mu\text{g}/\text{m}^3$ and 25-50 $\mu\text{g}/\text{m}^3$, respectively. For a detailed discussion, the readers are referred to the OAQPS document (Wapora, 1982). Landau and Manos (1976) estimated the number of Americans who reside within 5 miles of 38 major plants in the United States that produce monomers and polymers and fabricate polymers. The estimated values are given in Table 7-3. The values are only population estimates and are not meant to indicate the absolute number of people exposed to vinylidene chloride.

The size of the population residing near plant sites as given in Table 7-3 may have changed slightly due to shut down of a few plants or addition of new plants and due to some population shifts. It should be noted that the number of people likely to be exposed to vinylidene chloride is not shown in Table 7-3.

The estimation of these values would require knowledge of the chemical's transport characteristics. Theoretically, dispersion modelling could provide an estimate of vinylidene chloride concentrations at different distances from the plant sites; however, such data are not presently available.

Experimental data could provide a solution to this problem. Going and Spigarelli (1977) detected vinylidene chloride at a distance of 1.5 miles from a vinylidene chloride monomer production facility. Their study was not designed to provide vinylidene chloride levels at different distances from the source. Therefore, the study of Going and Spigarelli (1977) cannot be used to estimate vinylidene chloride exposure.

TABLE 7-3

Estimated Population Residing Near Plant Producing or
Fabricating Monomers and Polymers of Vinylidene Chloride
(Landau and Manos, 1976)

Perimeter (miles)	Producing or Process Plants			Total
	Monomer	Polymer	Fabrication	
0-1/2	315	11,114	37,310	48,739
1/2-1	1,035	34,091	100,974	136,100
1-3	17,356	221,966	1,064,994	1,304,316
3-5	70,230	337,547	1,676,463	2,084,240
TOTAL	88,936	604,718	2,879,741	3,573,395

With the median ambient levels (averaged over 1 hour up to 1091 days) given in Table 7-1, and assuming 20 m^3 of air is inhaled per day by an individual, the estimated daily vinylidene chloride intake from urban/suburban areas through inhalation is approximately $0.4 \text{ } \mu\text{g}$. Pellizzari et al. (1979) estimated the daily average vinylidene chloride intake through inhalation of ambient air from the Baton Rouge, Geismen, and Plaquemine, LA, area and the Houston, Deer Park, and Pasadena, TX, area to be $0.7 \text{ } \mu\text{g}$ and $1.6 \text{ } \mu\text{g}$, respectively. However, the daily inhalation exposure may be as high as 0.3 mg near the immediate vicinity of source areas if a median vinylidene chloride level of 3600 ppt (see Table 7-2) is employed to calculate the exposure value.

7.2 WATER

7.2.1 Environmental Levels

The presence of vinylidene chloride in water samples was reported as early as 1975. Monitoring investigations performed prior to or during 1976 have reported the presence of vinylidene chloride in a wide range of water samples ranging from industrial effluents to drinking waters. Some of the earlier attempts at analysis, however, could not separate vinylidene chloride from its isomers due to inadequate GC column selection. The presence of unseparated vinylidene chloride and its isomers has been reported in effluents from the textile industry, effluents from the latex industry, effluents from sewage treatment plants, raw surface waters, finished drinking waters, and even in well water (Schackelford and Keith, 1976).

The presence of vinylidene chloride alone, separated from its isomers, has been reported four times in industrial effluents, two times in raw surface waters, seven times in finished drinking water, and one time in well water (Shackelford and Keith, 1976). No quantitative values for vinylidene chloride concentrations are available from most of these earlier reports, however,

because the vinylidene chloride concentrations were very low and mass spectrometry was used for its identification and not for quantification. The levels of vinylidene chloride in industrial raw wastewaters and treated wastewaters have been determined recently and are shown in Table 7-4 and Table 7-5. These tables also indicate the expected vinylidene chloride loading in surface waters due to the discharge of the wastewaters into these water bodies. According to these tables, nonferrous metals manufacturing industries are likely to contribute to the maximum vinylidene chloride loading into the water bodies.

In a recent publication, Keith and Telliard (1979) reported the approach used by the U.S. EPA in establishing a priority pollutant list for toxic chemicals. Industrial effluents were screened by GC-MS for various pollutants, including vinylidene chloride. Among the over 2,000 samples representing 17 different industrial categories that were analyzed up until August 1978, 7.7% qualitatively showed the presence of vinylidene chloride.

The levels of vinylidene chloride in surface waters around a few industrial sites were determined by Going and Spigarelli (1977). The samples were analyzed by a combination of GC-FID and GC-MS. Table 7-6 lists the concentrations of vinylidene chloride in these water samples.

The levels of vinylidene chloride in U.S. raw surface and ground waters used for the abstraction of drinking waters and in potable drinking waters were monitored as part of the National Organic Monitoring Survey (NOMS) (U.S. EPA, 1977) and National Organics Reconnaissance Survey (NORS) (Coleman et al., 1976). The results of these two surveys were combined with four other surveys conducted for the U.S. EPA (namely, surveys by Stanford Research Institute, Office of Toxic Substances (OTS), Annapolis Field Study and U.S. EPA Region V Survey) by Coniglio et al. (1980) to derive statistics on the levels of volatile organics in raw and finished waters in the U.S. None of the surface waters from the 105 cities

TABLE 7-4
Industrial Occurrence of Vinylidene Chloride in Raw Wastewater^{a,b}

Industry	Number of Samples	Number of Detections	Concentration, $\mu\text{g/l}$			Loading, kg/d^g Mean
			Minimum	Maximum	Mean	
Auto and Other Laundries (c)	5	3	2.0	23	7.5	0.0014
Coal Mining (d)	47	3	3	3	3	0.011
Aluminum Forming	NA	ND				
Battery Manufacturing	27	8	ND	10	10	0.0008
Coil Coating	30	2	ND	36	10	0.0011
Metal Finishing (d)	95	16	2	10,000	760	0.61
Porcelain Enameling	1	ND				
Nonferrous Metals Manufacturing	66	9 (e)	ND	6,100	200	1.6
Organic Chemicals Manufacturing/ Plastics	NA	22	NA	NA	200	NA
Paint and Ink Formulation (c)	36	8	NA	620	78	0.00031
Soap and Detergent Manufacturing (c)	3	3	11	25	18	0.0016
Steam Electric Power Plants (f)	11	ND				
Textile Mills (d)	50	1		<5		<0.0035

^aSource: U.S. EPA, 1981

^bThe pollutant was not detected during screening analyses for the following industries: Leather Tanning and Finishing, Foundries, Explosives Manufacture, Gum and Wood Chemicals, Pharmaceutical Manufacturing, Nonferrous Metals Manufacturing, Pulp and Paperboard Mills, Rubber Processing, Timber Products Processing.

^cScreening data

^dScreening and verification data

^eDetections >10 $\mu\text{g/l}$

^fVerification data plus surveillance and analyses

^gPollutant loadings determined by multiplying mean pollutant concentration by industry wastewater discharges; where mean is not available one-half the reported maximum was utilized.

NA = Not Available; ND = Not Detected

TABLE 7-5

Industrial Occurrence of Vinylidene Chloride in Treated Wastewater^{a,b}

Industry	Number of Samples	Number of Detections	Concentration, $\mu\text{g/l}$			Loading, kg/d^g Mean
			Minimum	Maximum	Mean	
Auto and Other Laundries (a)	1	ND				
Coal Mining (b)	51	3	3	3	3	0.0056
Coil Coating	5	ND				
Explosives Manufacture	NA	NA	NA	NA	NA	NA
Pharmaceutical Manufacturing	2	2	10	10	10	0.0091
Nonferrous Metals Manufacturing	70	10 (c)	ND	4,100	120	0.96
Organic Chemicals Manufacturing/ Plastics	NA	15	NA	NA	6.8	NA
Paint and Ink Formulations (a)	24	4	NA	44	19	0.000076
Soap and Detergent Manufacturing	NA	NA	NA	NA	NA	NA
Steam Electric Power Plants (d)	12	1		10		0.015

^aSource: U.S. EPA, 1981^bThe pollutant was not detected during screening analyses for the following industries: Leather Tanning and Finishing, Aluminum Forming, Battery Manufacturing, Foundries, Porcelain Enameling, Gum and Wood Chemicals, Rubber Processing, Timber Products Processing.^cScreening data^dScreening and verification data^eDetections $>10 \mu\text{g/l}$ ^fVerification data plus surveillance and analyses^gPollutant loading determined by multiplying mean pollutant concentration by industry wastewater discharges; where mean is not available, one-half the reported maximum was utilized.

NA = Not Available; ND = Not Detected

TABLE 7-6

Vinylidene Chloride Concentration in a Few Waters Near Industrial Sites^a

Sample Source _b	Industrial Site	Industrial Process	Frequency of Detection ^c	Concentration	
				High (µg/l)	Low (µg/l)
Mississippi River	Dow Chemical, Plaquemine, LA	Monomer producer	1/2	0.2	ND
PPG Canal	PPG Industries Lake Charles, LA	Monomer and methyl chloroform producer ^d	1/2	550	ND
Ohio River	W.R. Grace, Owensboro, KY	Polymer producer	0/2	ND	ND
Holston River	Eastman, Kingsport, TN	Modacrylic producer	2/4	1	ND
Tittabawassee River	Dow Chemical, Midland, MI	Polymer producer and extruder	1/3	<1	ND

^aSource: Going and Spigarelli, 1977^bThe samples were collected at different points upstream and downstream from the discharge outfall.^cThe first figure indicates the number of samples that showed presence of compound and the second figure indicates the total number of samples analyzed.^dThis plant no longer uses vinylidene chloride to produce methyl chloroform.

ND = Not Detected

monitored in this survey showed any detectable level of vinylidene chloride. Only two finished waters abstracted from surface waters from 103 cities showed the presence of this compound at concentrations of 0.2 µg/l and 0.51 µg/l. Since the compound was not present in the raw water, it is likely that the treatment process is the source of vinylidene chloride in these treated waters. Going and Spigarelli (1977) also monitored the treated surface waters from Cincinnati, OH, Philadelphia, PA, and Lawrence, MA, for vinylidene chloride. Neither the pre-chlorinated nor the post-chlorinated treated water from any of these cities showed any detectable level of vinylidene chloride. Recently, Pellizzari et al. (1979) reported the mean concentration of vinylidene chloride in the drinking water from New Orleans and Baton Rouge, LA, areas to be 0.2 µg/l.

A survey of raw and treated groundwaters from 13 U.S. cities was also conducted by Coniglio et al. (1980). Raw groundwaters from two cities both showed detectable quantities of vinylidene chloride at a level of 0.5 µg/l. Vinylidene chloride was detected in treated groundwater from only one city at a concentration of 0.2 µg/l. Going and Spigarelli (1977) also detected 0.059 µg/l and 0.045 µg/l of vinylidene chloride in pre-chlorinated and post-chlorinated treated groundwater, respectively, from Miami, FL. Eight state agencies (e.g., AL, FL, KY, ME, MA, NC, SC, and TN) also monitored the levels of dichloroethylenes (three unseparated isomers) in their well water. Of the total of 781 samples tested, 23% showed detectable levels of dichloroethylenes with a maximum level of 860 µg/l (Coniglio et al., 1980). The levels of vinylidene chloride in 378 groundwater samples from urban, suburban, rural, and dump site areas in New Jersey were monitored by Page (1981). Approximately 44% of the groundwaters showed detectable levels of vinylidene chloride. Although the highest concentration of vinylidene chloride detected in a groundwater, probably from a dump

site (author did not specify the sites) was 17.3 mg/l, the median concentration of vinylidene chloride in all samples was below the detection limit (10 µg/l).

The levels of vinylidene chloride in raw and treated (both ground and surface waters) from 30 Canadian potable water treatment facilities serving about 5.5 million consumers across Canada were monitored by Otson et al. (1982a,b). Vinylidene chloride was not detected in any of the raw water samples, and only one finished water sample collected during the summer months (Aug.-Sept.) showed a mean concentration of less than 1 µg/l and a maximum concentration of 20 µg/l for vinylidene chloride.

7.2.2 Exposure

It can be concluded from the discussions in Section 7.2.1 that about 3% of the total drinking water supplies in the U.S. contain vinylidene chloride at an estimated mean concentration of 0.3 µg/l. Assuming that an individual consumes 2 liters of water per day, the daily exposure to vinylidene chloride for these consumers (3% water supplies) would be approximately 0.6 µg. For the majority of the U.S. population, the daily exposure to vinylidene chloride from the consumption of drinking water would be lower than 0.6 µg. Therefore, the daily exposure of vinylidene chloride to the general population in the U.S. from the consumption of drinking water would be <0.6 µg, although the maximum daily exposure in certain communities could exceed 1 µg (see Section 7.2.1).

7.3 SOIL

DeLeon et al. (1980) analyzed over 100 soil samples within and around a few chemical waste disposal sites for 11 chlorinated hydrocarbons including vinylidene chloride. The analytical method used was gas chromatography-electron capture detector (GC-ECD) for quantification and gas chromatography-mass spectrometry (GC-MS) for confirmation. One of the three samples collected from a midwestern chemical disposal site (name not mentioned) showed 21.9 µg/g (dry

weight) of vinylidene chloride, while none was detected in the other two samples. The detection limit of the method was given as 10 $\mu\text{g/g}$.

7.4 FOODS

Human exposure to vinylidene chloride from foods can occur in two ways: from consumption of food contaminated by polyvinylidene wrappings, and from consumption of edible aquatic foods that have bioconcentrated vinylidene chloride from water. Vinylidene chloride content both in food-packaging materials and in the foods themselves have been determined. The extent of migration of residual vinylidene chloride from polyvinylidene packaging materials into foods is customarily tested with food-simulating solvents rather than with foods themselves (Hollifield and McNeal, 1978). The three food-simulating solvents, which represented aqueous, oily, and fatty foods, were water, corn oil, and heptane, respectively (Hollifield and McNeal, 1978).

The results of analysis of vinylidene chloride content in saran and meat packaging films as determined by Going and Spigarelli (1977) are shown in Table 7-7.

Hollifield and McNeal (1978) analyzed five commercial food-packaging saran films of 0.5 mil thickness and determined that the residual monomer concentration varied between 1.6 $\mu\text{g/g}$ and 8.1 $\mu\text{g/g}$.

The extent of migration of vinylidene chloride from saran films to food-simulating solvents at 49°C is shown in Table 7-8.

It can be concluded from Table 7-8 that the migration of the residual monomer from the 0.5 mil film was essentially complete in corn oil and heptane in well under 24 hours. It leveled off at about 60% in water after 24 hours.

It is difficult to estimate the extent of human exposure to vinylidene chloride resulting from consumption of food contaminated by polyvinylidene packaging materials. The exposure to vinylidene chloride from consuming

TABLE 7-7

Analysis of Food-Packaging Films for Vinylidene Chloride^a

Sample	Concentration ($\mu\text{g/g}$)
Saran Wrap	41-58 ^b
Saran Wrap	4.9
Oscar Mayer Film A	ND
Oscar Mayer Film B	ND
Oscar Mayer Film C	ND

^aSource: Going and Spigarelli, 1977^bThis saran wrap, presumably produced in 1974, contained higher amounts of vinylidene chlorideND = Not Detected. The detection limit was 0.005 $\mu\text{g/g}$.

TABLE 7-8

Migration of Vinylidene Chloride from Saran Films to Heptane,
Corn Oil, and Water at 49°C^a

Film Thickness	Residue Monomer Concentration μg/g	Time of Contact (hours)	Concentration Found in Solvent (ppb)	Percent of Total Extracted
Heptane				
0.5 mil	8.1	0.5	34	76%
		1.0	37	88%
		2.0	41	94%
		2.5	44	106%
Corn Oil				
0.5 mil	8.1	1	18	41%
		3	34	76%
		6	40	94%
		24	39	94%
		48	41	94%
Water				
0.5 mil	8.1	4	26	59%
		24	27	65%
		312	24	59%
		696	24	59%

^aSource: Hollifield and McNeal, 1978

contaminated edible aquatic foods is equally difficult to estimate, even when the bioaccumulation factor is available. To calculate body burden, a knowledge of the concentration of vinylidene chloride in the water from which the aquatic organism originated is necessary.

8. BIOLOGICAL EFFECTS ON PLANTS AND MICROORGANISMS

The 96-hour EC_{50} (median effective concentration) values for the freshwater alga Selenastrum capricornutum and the marine alga Skeletonema costatum were found to be greater than 798 or 712 mg vinylidene chloride/l, respectively (U.S. EPA, 1978). These were the highest concentrations tested. The reported "no effect concentration" was <80 mg/l for Selenastrum capricornutum and 712 mg/l for Skeletonema costatum. The criteria for toxic effect in these bioassays were reduction in cell number or chlorophyll a content in algae cultures grown for 96 hours.

No studies were found in the literature searched on the biological effects of vinylidene chloride on microorganisms.

9. BIOLOGICAL EFFECTS ON AQUATIC ORGANISMS

9.1 ACUTE TOXICITY

The acute toxicity of vinylidene chloride has been determined with several species of freshwater and marine fish and invertebrates. Acute toxicity values are presented in Table 9-1.

9.1.1 Freshwater Fish

The 96-hour LC_{50} (95% confidence interval) for bluegill sunfish, Lepomis macrochirus, tested under static exposure conditions was 73.9 (56.8-90.7) mg/l (U.S. EPA, 1978; Buccafusco et al., 1981). This value is the same as the 24-hour LC_{50} (Table 9-1), indicating that all mortalities during the test occurred within the first day of exposure. The "no effect" concentration was 32.0 mg/l. Dawson et al. (1977) reported a much higher 96-hour LC_{50} for bluegill sunfish (220 mg/l). The chemical used was identified as "vinylidene chlorine" and presumable was vinylidene chloride. This test was conducted under static exposure conditions. Dill et al. (n.d., cited in U.S. EPA, 1979) determined the 96-hour LC_{50} for fathead minnows, Pimephales promelas, under static and flow-through exposure conditions. The values were 169 mg/l and 108 mg/l, respectively. The different results indicate that the flow-through technique gave a higher estimate of the acute toxicity of vinylidene chloride than did the static exposure technique.

9.1.2 Freshwater Invertebrates

The only freshwater invertebrate for which vinylidene chloride toxicity data are available is the water flea, Daphnia magna. The 24-hour and 48-hour static LC_{50} (95% confidence interval) values were 98 (71-130) and 79 (62-110) mg/l, respectively (LeBlanc, 1980). The "no discernible effect" concentration was <2.4 mg/l, which was the lowest concentration tested. In another study (Dill

TABLE 9-1
Acute Toxicity Values for Marine and Freshwater Fish and Invertebrates Exposed to Vinylidene Chloride

Species	Test Type	Water Type	LC50 (mg/l) ^a				No Effect Concentration (mg/l)	References
			24-h	48-h	72-h	96-h		
FISH								
Bluegill Sunfish (<u>Lepomis macrochirus</u>)	ST	FW	73.9 (56.8-90.7)	73.9 (56.8-90.7)	73.9 (56.8-90.7)	73.9	32.0	U.S. EPA, 1978 Buccafusco et al., 1981 Dawson et al., 1977
	ST	FW	--	--	--	220	--	
Fathead Minnows (<u>Pimephales promelas</u>)	ST	FW	--	--	--	169		Dill et al., n.d. Dill et al., n.d.
	ST	FW	--	--	--	108	--	
Sheepshead Minnow (<u>Cyprinodon variegatus</u>)	ST	SW	249 (198-338)	249 (198-338)	249 (198-338)	249 (198-338)	79.8	U.S. EPA, 1978 Heitmuller et al., 1981
Tidewater Silversides (<u>Menidia beryllina</u>)	ST	SW	--	--	--	250	--	Dawson et al., 1977
INVERTEBRATES								
Water Fleas (<u>Daphnia magna</u>)	ST	FW	--	11.6 ^b	--	--	--	Dill et al., n.d.
Water Fleas (<u>Daphnia magna</u>)	ST	FW	98 (71-130)	79 (62-110)	--	--	<2.4	LeBlanc, 1980
Mysid Shrimp (<u>Mysidopsis bahia</u>)	ST	SW	>798	>798	>798	224 (106-434)	14.2	U.S. EPA, 1978

^aNumbers in parentheses represent the 95% confidence interval

^bMedian effective concentration (EC₅₀)

ST = static exposure; FT = flow-through exposure; FW = freshwater; SW = sea water

et al., n.d., cited in U.S. EPA, 1979), the 48-hour static EC₅₀ (median effective concentration) was 11.6 mg/l, which is much lower than the previously cited LC₅₀ values. The difference may be due to the different toxicity endpoints measured (effective concentration vs. lethal concentration) or some other factor(s).

9.1.3 Marine Fish

The 96-hour LC₅₀ (95% confidence interval) for sheepshead minnows, Cyprinodon variegatus, tested under static exposure conditions, was 249 (198-338) mg/l (U.S. EPA, 1978; Heitmuller et al., 1981). As occurred with bluegill sunfish, the 96-hour LC₅₀ was the same as the 24-hour LC₅₀ (Table 9-1). The "no effect" concentration was 79.8 mg/l. The only other study with marine fish was by Dawson et al. (1977), who reported a 96-hour LC₅₀ value of 250 mg/l for tidewater silversides, Menidia beryllina, tested under static exposure conditions. This value is nearly identical to the 96-hour LC₅₀ for sheepshead minnows.

9.1.4 Marine Invertebrates

The mysid shrimp, Mysidopsis bahia, is the only species of marine invertebrate that has been tested for vinylidene chloride toxicity. The 96-hour LC₅₀ value (95% confidence interval) for this species was 224 (106-434) mg/l. The LC₅₀ value at 24, 48, and 72 hours could not be determined precisely, but was greater than 798 mg/l for each time period. The "no effect" concentration was 14.2 mg/l. The 96-hour LC₅₀ value for mysid shrimp was similar to the 96-hour LC₅₀ values for sheepshead minnows and tidewater silversides.

9.2 SUBACUTE TOXICITY

The only information regarding the subacute effects of vinylidene chloride on aquatic organisms concerns one species of freshwater fish.

Dill et al. (n.d., cited in U.S. EPA, 1979) determined the 13-day LC_{50} for fathead minnows under flow-through conditions. This value, 29 mg/l, was considerable lower than the equivalent 96-hour LC_{50} value (108 mg/l).

No adverse effects were observed in a fathead minnow embryo-larval test at 2.8 mg/l, which was the highest concentration tested (U.S. EPA, 1978). In this test, fathead minnow eggs were exposed under flow-through conditions to a series of vinylidene chloride concentrations through the hatching and early larval life stages.

Most of the previously cited aquatic toxicity tests were conducted under static exposure conditions. As far as can be determined, the LC_{50} values obtained from these tests were calculated on the basis of initial nominal concentrations, rather than actual measured concentrations.

Due to the high volatility of vinylidene chloride (Section 3), it is probable that test organisms in static toxicity tests would be exposed to rapidly decreasing concentrations of vinylidene chloride. This rapid loss may account for the observations that the static LC_{50} values for bluegill sunfish and sheepshead minnow (U.S. EPA, 1978) did not decrease between 24 and 96 hours of exposure and that the static 96-hour LC_{50} for fathead minnows was higher than the flow-through 96-hour LC_{50} with the same species (Dill et al., cited in U.S. EPA, 1979). Although the static tests with bluegills and sheepshead minnows produced apparent lethal thresholds by 24 hours of exposure, Dill et al. (n.d., cited in U.S. EPA, 1979) showed that flow-through exposure with fathead minnows resulted in sufficient additional mortality between 4 and 13 days of exposure to lower the LC_{50} from 108 to 29 mg/l. Although static tests may provide some useful information concerning the potential ecological effects of vinylidene chloride in spill

situations, the flow-through results indicate that static tests may underestimate the effects of acute and chronic exposure to constant vinylidene chloride concentrations.

In summary, the reported acute LC_{50} or EC_{50} values for four fish species and two invertebrate species ranged between 11.6 and 250 mg/l. The acute static "no effect" concentrations reported in U.S. EPA (1978) decreased in this order: sheepshead minnow (79.8 mg/l) > bluegill sunfish (32.0 mg/l) > mysid shrimp (14.2) > Daphnia magna (>2.4 mg/l, the lowest concentration tested). However, insufficient species have been tested to definitely establish whether or not there may be significant differences in sensitivity to vinylidene chloride between marine and freshwater organisms, or between fish and invertebrates. Of the species tested, the most sensitive species appears to be Daphnia magna, which was adversely affected at 2.4 mg/l, the lowest concentration tested with that species in the U.S. EPA (1978) study. This concentration is lower than the concentration (2.8 mg/l) reported in the same study to cause no adverse effects on fathead minnows in a flow-through embryo-larval test.

Additional flow-through chronic or subchronic testing with Daphnia magna and other freshwater and marine species would be required to estimate more accurately the range of species sensitivity and to determine the maximum acceptable concentrations of vinylidene chloride compatible with protection of aquatic life.

10. BIOLOGICAL EFFECTS

10.1 PHARMACOKINETICS

The available data characterize the disposition and metabolism of vinylidene chloride in experimental animals when administered as a single dose or exposure. The effects of age, sex, route, species and fasting on disposition and metabolism are also characterized. Information on the disposition and metabolism of vinylidene chloride during subchronic or chronic exposure, however, is lacking.

10.1.1 Absorption and Distribution

Studies with experimental animals indicate that vinylidene chloride is readily absorbed and rapidly distributed in the body. Systemic absorption after intragastric administration of ^{14}C -vinylidene chloride to male or female rats appears to be rapid and fairly complete based on the rapid appearance of labeled vinylidene chloride in the expired air (McKenna et al., 1978b; Reichert et al., 1979) and the extensive pulmonary and urinary excretion of radioactivity (Jones and Hathway, 1978a; McKenna et al., 1978b; Reichert et al., 1979). Intraperitoneal administration also resulted in extensive pulmonary and urinary excretion (Jones and Hathway, 1978a).

Anderson et al. (1979a) studied the rate of uptake of vinylidene chloride vapor by fasted male rats. Fasting (for 18 hours) has been shown to deplete the glutathione content of the liver, and to increase the sensitivity of rats to the toxicity of vinylidene chloride; these observations are consistent with the role of glutathione in the detoxification of vinylidene chloride (Section 10.1.2.2.1). Throughout Section 10.1, animals can be assumed to be fed unless fasting is specified. Uptake was determined from disappearance of vinylidene chloride from a closed chamber as a function of time, corrected for nonspecific

loss. Uptake could be resolved into a rapid and a slow phase. The rapid phase was not affected by pretreatment of the rats with inhibitors of microsomal metabolism and could therefore be separated from the slow phase, in which the rate was markedly reduced by these inhibitors (the slow phase is discussed in Section 10.1.2.1). The rapid phase was a first-order process with a rate constant of 2.2 hr^{-1} (half-life = 0.315 hr). The rate constant did not vary significantly with initial concentration of vinylidene chloride. The magnitude of the rapid component was proportional to the initial concentration of vinylidene chloride and the weight of rats in the chamber. The concentration of vinylidene chloride in the body as a function of concentration in the chamber at equilibrium (whole body solubility coefficient) was 0.0163 mg/kg body weight/ppm chamber, equivalent to a body/gas distribution coefficient of 4.04. The rapid phase was essentially complete in 60 to 80 minutes. On the basis of these characteristics, Andersen et al. (1979a) concluded that the rapid phase represented whole body equilibration.

Jaeger et al. (1977a) investigated the distribution of radioactivity in various tissues 30 minutes after cessation of a 2-hour exposure of fed and fasted male rats to ^{14}C -vinylidene chloride. Exposure occurred in a closed chamber; the initial concentration of vinylidene chloride was 2000 ppm. Although there was no significant difference between fed and fasted rats in the overall rate of uptake of vinylidene chloride, the tissues and serum of fasted rats contained more radioactivity than those of fed rats. The highest concentration of radioactivity was found in kidney, and the next highest in liver. Spleen contained an intermediate level of radioactivity; heart and brain contained the least. In both fed and fasted rats, most of the radioactivity in kidney was trichloroacetic acid (TCA)-soluble. Approximately one-third of the radioactivity in liver was

TCA-precipitable, indicating that it was covalently bound to macromolecules. Liver contained more TCA-precipitable radioactivity than did kidney.

McKenna et al. (1978a) from the Dow Chemical Company investigated the fate of ^{14}C -vinylidene chloride in fed and fasted (deprived of food for 18 hours) male rats 72 hours after a 6-hour inhalation exposure to 10 or 200 ppm. As presented in Table 10-1, fasted rats had lower body burdens of ^{14}C than did fed rats. Although most of the body burdens had been excreted within 72 hours, a small but significant amount remained in the carcasses. After the 200 ppm exposure, more radioactivity was retained in the carcasses of fasted rats than in those of fed rats. All of the radioactivity found in urine, feces, and tissues was nonvolatile and therefore was considered to represent nonvolatile metabolites of vinylidene chloride. Seventy-two hours after exposure to either 10 or 200 ppm vinylidene chloride, concentrations of radioactivity were highest in kidney and liver and decreased in the following order: kidney > liver > lung > skin > plasma > fat, muscle, and remaining carcass. At both exposure levels, concentrations of radioactivity remaining in the tissues of fasted rats were greater than those in the tissues of fed rats when the data were normalized to account for the lower end-exposure body burdens of fasted rats.

In similar experiments with intragastric administration of 1 or 50 mg ^{14}C -vinylidene chloride to fed and fasted male rats, McKenna et al. (1978b) found at 72 hours after dosing that concentrations of radioactivity were highest in liver and next highest in kidney. Levels of radioactivity in lung were not as high as noted in the inhalation experiment previously described (McKenna et al., 1978a). Jones and Hathway (1978a) reported that the whole animal autoradiography of young male rats after intragastric administration of ^{14}C -vinylidene chloride revealed high concentrations of ^{14}C in the kidneys and liver 30 minutes after dosing, followed at 1 hour by a more general distribution of radioactivity

Table 10-1

End-Exposure Body Burdens and Disposition of ^{14}C Activity in Rats 72 Hours
Following Exposure to 10 or 200 ppm of ^{14}C -Vinylidene Chloride for 6 Hours (McKenna et al., 1978a)

	10 ppm		200 ppm	
	Fasted rats	Fed rats	Fasted rats	Fed rats
Body burden ^a				
Total milligram equivalents of ^{14}C -vinylidene chloride/rat	0.589 \pm 0.19 ^b	0.748 \pm 0.041	9.57 \pm 0.08 ^b	11.41 \pm 0.64
10-4 Milligram equivalents of ^{14}C -vinylidene chloride/kg	2.30 \pm 0.05	2.89 \pm 0.12	35.39 \pm 0.30 ^b	44.53 \pm 0.64
Percentage body burden				
Expired vinylidene chloride	1.60 \pm 0.19	1.63 \pm 0.07	8.36 \pm 0.27 ^b	4.17 \pm 0.62 ^c
$^{14}\text{CO}_2$	8.27 \pm 0.80	8.74 \pm 1.86	7.24 \pm 0.51	8.22 \pm 0.54
Urine	78.19 \pm 1.42	74.72 \pm 1.65	70.41 \pm 1.86	74.66 \pm 2.62
Feces	6.75 \pm 0.56 ^b	9.73 \pm 0.05	2.72 \pm 0.41	6.39 \pm 1.50
Tissues and carcass	3.91 \pm 0.05	3.45 \pm 0.30	7.45 \pm 1.01 ^b	4.26 \pm 0.09 ^c
Skin	1.37 \pm 0.25	1.30 \pm 0.09	3.07 \pm 1.0	1.92 \pm 0.29
Cage wash	0.27 \pm 0.11	0.44 \pm 0.14	0.76 \pm 0.10 ^b	0.34 \pm 0.02

^aThe values are given as $\bar{x} \pm \text{SE}$ (n = four per group).

^bSignificantly different from fed animals of the same exposure group, $p < 0.05$, Student's t test.

^cSignificantly different from fed animals of the 10 ppm exposure group, $p < 0.05$, Student's t test.

throughout the soft tissues. According to the authors, liver and kidney retained radioactivity longer than did other tissues.

10.1.2 Metabolism

The metabolism of vinylidene chloride by experimental animals has been extensively studied. A metabolic scheme that integrates the data from several laboratories is presented in Figure 10-1. In this scheme, vinylidene chloride is metabolized to reactive intermediates, 1,1-dichloroethylene oxide, chloroacetyl chloride, and monochloroacetic acid. These reactive intermediates are detoxified by conjugation with glutathione (GSH), ultimately leading to the formation of mercapturic acids (derivatives of N-acetyl cysteine) or degradation to various thioglycolic acids. A minor pathway for detoxification of monochloroacetic acid involves further oxidation to CO_2 . 1,1-Dichloroethylene oxide, chloroacetyl chloride, and monochloroacetic acid can react with (i.e., alkylate) cellular macromolecules; this type of reaction could be involved in the production of toxic effects.

From an analysis of the literature on the metabolism of chlorinated ethylenes, Lowe et al. (1983) have concluded that the extent of metabolism of these compounds generally decreases with increasing degree of chlorine substitution. Thus the extent of metabolism appears to be: vinyl chloride \approx vinylidene chloride > trans or cis-dichloroethylene > trichloroethylene > tetrachloroethylene. These conclusions were drawn from a comparison of the results of a number of oral and inhalation studies, taking into account species and dose differences.

10.1.2.1 Metabolism to Reactive Intermediates

Vinylidene chloride is thought to be metabolized by the microsomal mixed-function oxidase system to an epoxide (oxirane), 1,1-dichloroethylene oxide (Bonse and Henschler, 1976; Liebman and Ortiz, 1977; McKenna et al., 1977).




Figure 10-1 Metabolic Pathways for Vinylidene Chloride. Identified metabolites are underlined. Identification was usually by gas chromatography-mass spectrometry. R' is considered to be OH; R is unknown. (Adapted from Jones and Hathway, 1978a, 1978b; Reichert et al., 1979; Leibman and Ortiz, 1977)

According to Bonse and Henschler (1976), 1,1-dichloroethylene oxide is extremely unstable and could not be isolated during attempts at chemical synthesis, although its expected rearrangement product, chloroacetyl chloride, was formed after chemical oxidation of vinylidene chloride with m-chloroperbenzoic acid. Metabolism of vinylidene chloride to this epoxide has not been directly demonstrated, but rather is inferred by analogy with the metabolism of other chlorinated ethylenes (Leibman and Ortiz, 1977; Bonse and Henschler, 1976; Uehleke et al., 1976) and from the observed metabolism of vinylidene chloride to monochloroacetic acid.

The metabolism of vinylidene chloride by hepatic cytochrome P-450 in microsomes from male Long-Evans rats was studied in vitro by Costa and Ivanetich (1982) in an attempt to identify metabolites and detoxification pathways. Vinylidene chloride was shown to be metabolized to dichloroacetaldehyde and to monochloroacetic acid by hepatic microsomes, after binding to cytochrome P-450 in the presence of an NADPH-generating system. Certain forms of cytochrome P-450 appear to play minor roles in the metabolism of vinylidene chloride (such as the phenobarbital-inducible form) while other forms may be more efficient.

Radioactively-labeled monochloroacetic acid has been identified in the urine of rats after intragastric administration in corn oil of ^{14}C -vinylidene chloride (Jones and Hathway, 1978a), in rat livers perfused with ^{14}C -vinylidene chloride (Reichert and Bashti, 1976), and after incubation of ^{14}C -vinylidene chloride with 9000 x g supernatants from rat liver homogenates (Leibman and Ortiz, 1977). Monochloroacetic acid is the expected product of rearrangement of the epoxide to chloroacetyl chloride and subsequent hydrolysis (Bonse and Henschler, 1976; Leibman and Ortiz, 1977). Alternatively, monochloroacetic acid could result from metabolism of the epoxide by epoxide hydrase to the diol shown

in Figure 10-1, followed by rearrangement to chloroacetyl chloride and hydrolysis to monochloroacetic acid (Leibman and Ortiz, 1977).

The relative importance of the diol pathway is difficult to assess. Jones and Hathway (1978b) suggested that conversion to the diol is not important because CO_2 , a possible degradation product, is found only in small amounts in vivo. It is not known, however, whether the diol is metabolized to CO_2 to a significant extent. Leibman and Ortiz (1977) concluded that diol formation is relatively unimportant in the metabolism of vinylidene chloride to monochloroacetic acid because, 1,1,1-trichloropropene-2,3-oxide and cyclohexene oxide (inhibitors of epoxide hydrase) stimulated, rather than inhibited, the formation of monochloroacetic acid in incubations of vinylidene chloride with 9000 x g supernatants from rat liver homogenates. These two inhibitors react with glutathione (Oesch and Daly, 1972). Thus the results of Leibman and Ortiz (1977) could be explained by the depletion of glutathione, thereby resulting in the accumulation of monochloroacetic acid. Furthermore, according to Oesch (1972), there are two types of epoxide hydrase activity--one is tightly coupled with mixed-function oxidase activity, whereas the other is not. The coupled form is thought to be more important in detoxifying epoxides formed by mixed function oxidases and is relatively resistant to inhibitors (Oesch, 1972).

Andersen et al. (1979a) investigated the metabolism of vinylidene chloride by studying the rate of uptake of vinylidene chloride vapor by fasted male rats. Uptake was determined from the disappearance of vinylidene chloride as a function of time (usually plotted on linear coordinates) from a closed chamber, corrected for nonspecific loss. The uptake of vinylidene chloride by fasted male rats was biphasic; the initial rapid phase represented tissue equilibration (Section 10.1.1) and the slow phase represented metabolism. The slow phase of uptake was relatively independent of concentration at higher exposure levels and dependent

on concentration at lower exposures. A series of uptake experiments using different initial concentrations of vinylidene chloride were performed. The instantaneous rates of uptake were determined from tangents to the curves at 60 to 80 minutes of exposure, so that the contribution from the rapid phase was negligible. Instantaneous rates plotted against actual chamber concentration yielded a rectangular hyperbola typical of Michaelis-Menten kinetics, as would be expected if the slow phase represented metabolism. V_{\max} , the maximum velocity at saturating exposure levels, was 132 ppm/kg/hr (equivalent to 15.87 mg of vinylidene chloride metabolized/kg/hr). K_m , the exposure level that produced half the maximum velocity, was 335 ppm (equivalent to a body burden of 5.43 mg/kg).

At an initial concentration of vinylidene chloride of 300 ppm in air (below the K_m), the rate constant of the slow phase was reduced 84, 92, and 65% by pretreatment with pyrazole, carbon tetrachloride, and aminotriazole, respectively, but was unaffected by pretreatment with SKF 525A or 2,3-epoxypropan-1-ol. These compounds, with the exception of 2,3-epoxypropan-1-ol, are inhibitors of microsomal metabolism; all have been shown to affect the acute toxicity of vinylidene chloride (Section 10.2) (Andersen et al., 1980). At high initial concentrations of vinylidene chloride, the V_{\max} was unaffected by pretreatment with phenobarbital or by using fed rather than fasted rats, and was similar for immature and mature rats (Andersen et al., 1979a).

As determined by Andersen et al. (1979b), the acute toxicity of inhaled vinylidene chloride was relatively independent of concentration and dependent on duration of exposure at concentrations above 200 ppm (Section 10.2). In an attempt to correlate metabolism with toxicity, Andersen et al. (1979a) calculated the two theoretical curves shown in Figure 10-2. The solid line is based on the assumption that the product of exposure concentration times the duration




Figure 10-2 Comparisons of Observed LT50 Data for Vinylidene Chloride with Theoretical Curves Predicted for Two Different Mechanisms of Toxicity. As indicated by the symbol in upper right, data are given as LT50 and its 95% confidence interval. (Andersen et al., 1979a)

of exposure required to produce a given effect is constant. The dashed line is based on the assumption that the product of the rate of metabolism (at each exposure concentration) times the duration of exposure required to produce a given effect is constant. The experimentally determined LT50 (time required to produce 50% mortality rate) of 4.1 hours at a 200 ppm exposure concentration was used to calculate the constant. The experimentally determined LT50 values (Andersen et al., 1979b) fit the theoretical curve based on metabolism more closely than they fit the theoretical curve based on exposure concentration, suggesting that toxicity is a function of the amount of metabolite formed rather than the concentration of vinylidene chloride (Andersen et al., 1979a).

Using an experimental protocol similar to that of Andersen et al. (1979a), Filser and Bolt (1979) investigated the kinetics of metabolism of vinylidene chloride in the male rat. Filser and Bolt (1979) focused on strict zero-order and first-order kinetics, whereas Andersen et al. (1979a) focused on Michaelis-Menten kinetics. Both groups of investigators, however, demonstrated the saturable dose-dependent nature of vinylidene chloride metabolism. As determined by Filser and Bolt (1979); the V_{\max} (velocity for zero-order metabolism) was 100 $\mu\text{mol/kg/hr}$ (9.7 mg/kg/hr).

Reichert and Henschler (1978) reported that pretreatment of female rats with an inducer of mixed-function oxidases, DDT, increased the uptake of vinylidene chloride 35%, while direct addition of inhibitors (pyrazole and ethanol) of mixed-function oxidases decreased the uptake of vinylidene chloride 35 to 40% in isolated, perfused rat livers. Pretreatment with phenobarbital, or direct addition of SKF 525A, 6-nitro-1,2,3-benzothiadiazole, and 5,6-dimethyl-1,2,3-benzothiadiazole to the perfusate had little effect on uptake when low concentrations of vinylidene chloride were used. The uptake of vinylidene chloride, measured as the difference between prehepatic and posthepatic concentrations of

vinylidene chloride under steady state conditions was taken to be equivalent to metabolism. The authors stated that with a high concentration of vinylidene chloride in the perfusate, pretreatment with phenobarbital elevated the uptake of vinylidene chloride by approximately 30%. The results of Reichert and Henschler (1978) for perfused livers of female rats are in general agreement with the results of Andersen et al. (1979a) for intact fasted male rats except for the stimulation of uptake (metabolism) by phenobarbital pretreatment noted by Reichert and Henschler (1978). Pretreatment with ethanol for a 3 week period enhanced the activity of rat liver drug-metabolizing enzymes six-fold in their ability to convert vinylidene chloride (Sato et al., 1980). The authors discussed the increased rate of vinylidene chloride metabolism and noted that this may increase the toxicity of the compound if metabolites are the actual causative agents.

A comparison of sex, age, and species differences in metabolism and toxicity suggests that vinylidene chloride is metabolized to a toxic intermediate by mixed-function oxidases. In general, microsomal oxidation of compounds was found to be greater in male rats than in female rats (Kato, 1974). Male rats were much more susceptible to the hepatotoxic effects of vinylidene chloride than were female rats (Andersen and Jenkins, 1977). Microsomal oxidase activity reached a maximum in the rat at 30 to 40 days of age (Kato et al., 1964), an age range which corresponded to the size (100-150 g) of rats most susceptible to the toxicity of vinylidene chloride in the report of Andersen and Jenkins (1977). All of the studies reviewed in Section 10.1 of this report, however, were performed on mature (180-350 g) rats and mature mice unless otherwise specified.

A greater percentage of an orally administered dose (50 mg/kg) of ^{14}C -vinylidene chloride was metabolized by mice than by rats, based on quantitation of excreted radiolabeled metabolites and unchanged vinylidene chloride (Jones

and Hathway, 1978b). Similarly, as presented in Table 10-2, mice metabolized more vinylidene chloride per kg body weight than did rats after inhaling 10 ppm ¹⁴C-vinylidene chloride in air for 6 hours (McKenna et al., 1977). The overall pattern of excretion for mice and rats was similar. The concentration of covalently bound radioactivity in liver and kidney was much higher in mice than in rats, as shown in Table 10-3. Binding was measured as radioactivity that was TCA-precipitable and 80% methanol-insoluble and thus represented radioactivity covalently bound to protein and nucleic acid. As discussed in Section 10.4.1, the 9000 x g supernatant from mouse liver was more active than the 9000 x g supernatant from rat liver in metabolizing vinylidene chloride to mutagenic substances. These observations of increased metabolism and covalent binding in mice versus rats parallel the greater sensitivity of mice to the acute lethality, hepatotoxicity, and renal toxicity of vinylidene chloride (Short et al., 1977a,b). A brief review of the metabolism of vinylidene chloride by rats and mice is presented by Cooper (1980).

Hypoxia (7% oxygen), which inhibits mixed-function oxidase activity, decreased the hepatotoxicity of inhaled vinylidene chloride to fasted male rats (Jaeger, 1978). Disulfuram, which has been reported to inhibit hepatic mixed-function oxidases (Zemaitis and Green, 1976), protected male mice from the acute lethal and hepatotoxic effects of inhaled vinylidene chloride (Short et al., 1977a,b). Measured at 4 and 24 hours after intraperitoneal administration of ¹⁴C-vinylidene chloride, covalent binding of radioactivity to protein in liver and kidney was significantly reduced in mice fed disulfuram before and during exposure (Short et al., 1977a,b). Additional studies dealing with the relationship between the induction and inhibition of microsomal mixed function oxidases and the toxicity of vinylidene chloride are discussed in Section 10.2.

Table 10-2
End-Exposure Body Burdens and Disposition of ^{14}C -Activity in
Rats and Mice, 72 Hours Following Inhalation Exposure to
10 ppm ^{14}C -Vinylidene Chloride for 6 Hours^a

	Mice ^d	Rats ^e
Body burden, mg-Eq ^{14}C -vinylidene chloride/kg	5.30 \pm 0.75	2.89 \pm 0.24
Total metabolized vinylidene chloride mg-Eq ^{14}C -vinylidene chloride/kg ^c	5.27 \pm 0.74	2.84 \pm 0.26
Percentage body burden ^b		
Expired vinylidene chloride	0.65 \pm 0.07	1.63 \pm 0.14
Expired $^{14}\text{CO}_2$	4.64 \pm 0.17	8.74 \pm 3.72
Urine	80.83 \pm 1.68	74.72 \pm 2.30
Feces	6.58 \pm 0.81	9.73 \pm 0.10
Carcass	5.46 \pm 0.41	4.75 \pm 0.78
Cage Wash	1.83 \pm 0.84	0.44 \pm 0.28

^aSource: McKenna et al., 1977

^b $\bar{X} \pm \text{SE}$, n = 4.

^cCalculated from the total ^{14}C -activity recovered (end-exposure body burden) minus the ^{14}C -vinylidene chloride exhaled from each rat

^dMice were males of the Ha(ICR) strain.

^eRats were males of the Sprague-Dawley strain.

Table 10-3

Covalently Bound ^{14}C -Activity in Tissues 72 Hours
 Following Inhalation Exposure to 10 ppm
 ^{14}C -Vinylidene Chloride for 6 Hours (McKenna et al., 1977)

	^{14}C -vinylidene chloride, $\mu\text{g-Eq/g protein}$ ($\bar{X} \pm \text{SE}$, n = 4)	
	Liver	Kidney
Mice	22.29 \pm 3.77	79.55 \pm 19.11
Rats	5.28 \pm 0.14	13.14 \pm 1.15

The identification of methylthio-acetylaminoethanol as a urinary metabolite of vinylidene chloride suggested the alkylation of lipids by a reactive intermediate (Reichert et al., 1979; Henschler and Hoos, 1981). The pathway of formation of methylthio-acetylaminoethanol has not yet been clarified. Reichert et al. (1979) hypothesized that the first step could be the reaction of chloroacetyl chloride with phosphatidyl ethanolamine (a constituent of lipid membranes). This postulated pathway is illustrated in Figure 10-1.

10.1.2.2 Detoxification Pathways

10.1.2.2.1 Conjugation with Glutathione

Conjugation of vinylidene chloride or its metabolites with glutathione is indicated because administration of ^{14}C -vinylidene chloride (intragastrically or by inhalation) to rats and mice produced several S-containing radioactive metabolites. The major urinary metabolites of vinylidene chloride in rats have been identified as thiodiglycolic acid (thiodiacetic acid) (Jones and Hathway, 1978a,b; Reichert et al., 1979; McKenna et al., 1977, 1978a,b) and mercapturic acids: an N-acetyl-S-cysteinyl acetyl derivative shown in Figure 10-1 (Jones and Hathway, 1978a,b), N-acetyl-S-(carboxymethyl)cysteine (Reichert et al., 1979), and N-acetyl-S-(2-hydroxyethyl)cysteine (McKenna et al., 1977, 1978a,b). N-acetyl-S-(2-hydroxyethyl)cysteine is not shown as part of the metabolic scheme in Figure 10-1 because its relationship to other metabolites of vinylidene chloride is unclear. The major urinary metabolites of vinylidene chloride in mice were dithioglycolic acid and an N-acetyl-S-cysteinyl acetyl derivative shown in Figure 10-1 (Jones and Hathway, 1978b).

Vinylidene chloride itself does not appear to react with glutathione. McKenna et al. (1977) stated that they were unable to demonstrate conjugation of vinylidene chloride with glutathione either nonenzymatically or in the presence of the soluble fraction (which contains glutathione S-transferases) of a rat

liver homogenate. Conjugation with glutathione required the presence of a microsomal enzyme system, suggesting that vinylidene chloride was metabolized to a reactive intermediate before conjugation with glutathione. Unfortunately, no data or experimental details were presented.

Additional evidence that the S-containing metabolites (mercapturic and thioglycolic acids) of vinylidene chloride arise from conjugation with glutathione is as follows. The origin of the cysteine moiety in mercapturic acid derivatives of many xenobiotics has been shown to be glutathione (Chasseaud, 1973). In vivo experiments with unlabeled vinylidene chloride, in which the cysteine-cystine pools were labeled with ^{14}C , resulted in the production of labeled thiodiglycolic acid (Jones and Hathway, 1978a). Thus, at least part of the carbon skeleton of thiodiglycolic acid must have been derived from cysteine.

Administration of vinylidene chloride to male or female rats in vivo, or during perfusion of isolated rat livers, depleted the liver glutathione content (Jaeger et al., 1974; Reichert et al., 1978; McKenna et al., 1977; McKenna, 1979). Depletion of glutathione was maximal at about 4 hours after oral administration of vinylidene chloride to female rats and was dependent on the dose administered (Reichert et al., 1978). Depletion of glutathione after inhalation exposure of male rats was similarly dose dependent (McKenna et al., 1977; McKenna, 1979).

Reichert et al. (1978) attempted to demonstrate a correlation between the glutathione content of isolated, perfused livers from female rats and the rate of metabolism of vinylidene chloride measured as uptake (see Section 10.1.2.1, Reichert and Henschler, 1978). When the glutathione content was decreased 80% by the prior addition of diethyl maleate to the perfusate, an 18% decrease in vinylidene chloride metabolism was measured. A less pronounced decrease in glutathione content, produced by prior fasting of the rats, was not associated

with a decrease in vinylidene chloride metabolism; a slight increase (10%) was observed; neither change in metabolism was statistically significant. The authors concluded that the glutathione content of liver becomes a limiting factor in the metabolism of vinylidene chloride only when the glutathione content has been severely depleted.

Evidence for the metabolic pathways shown in Figure 10-1 that involve conjugation with glutathione has been obtained from the analysis of the urinary metabolites resulting from the administration of ^{14}C -labeled vinylidene chloride, monochloroacetic acid, and other metabolic intermediates to rats and mice. As shown in Table 10-4, the N-acetyl-S-cysteinyl acetyl derivative was detected only when rats were given ^{14}C -vinylidene chloride and not when they were given ^{14}C -monochloroacetic acid. The N-acetyl-S-cysteinyl acetyl derivative must therefore have come from conjugation of glutathione with a metabolite arising prior to the formation of monochloroacetic acid (Jones and Hathway, 1978a).

The administration of either ^{14}C -vinylidene chloride or ^{14}C -monochloroacetic acid gave rise to ^{14}C -labeled S-(carboxymethyl)cysteine, N-acetyl-S-(carboxymethyl)cysteine, thiodiglycolic acid, thioglycolic acid, and dithioglycolic acid (Tables 10-4 and 10-5, Jones and Hathway, 1978a). These results suggest that monochloroacetic acid is a key intermediate in the metabolism of vinylidene chloride and the S-(carboxymethyl)glutathione can be metabolized to thioglycolic acids or to a mercapturic acid.

S-(Carboxymethyl)cysteine was metabolized to thiodiglycolic acid and to CO_2 (Green and Hathway, 1977; Yllner, 1971; Jones and Hathway, 1978b). Because labeled CO_2 was detected only when the cysteine moiety of S-(carboxymethyl)-cysteine was uniformly labeled with ^{14}C (Jones and Hathway, 1978b) but not when the carboxymethyl moiety was labeled (Yllner, 1971), CO_2 must have been produced

Table 10-4

Relative Proportions of ^{14}C Urinary Metabolites After Intragastric Administration of 350 mg/kg 1- ^{14}C -Vinylidene Chloride or 50 mg/kg ^{14}C -Monochloroacetic Acid to Male Rats
(Adapted from Jones and Hathway, 1978a; Hathway, 1977*)

Vinylidene Chloride and Chloroacetic Acid Metabolites	% of Urinary Radioactivity	
	Vinylidene Chloride	Chloroacetic Acid
Thiodiglycolic acid	37.0	90.0
N-acetyl-S-cysteinyl-acetyl derivative	48.0	0
Dithioglycolic acid	5.0	3.0
Thioglycolic acid	3.0	3.0
Chloroacetic acid	3.0	0
Urea	0.5	0.5
S-(Carboxymethyl)cysteine	0	2.0
N-acetyl-S-(carboxymethyl)cysteine	--	2.0*

Table 10-5

Relative Proportions of ^{14}C Excretory Products After Intragastric
Administration of 50 mg/kg of 1- ^{14}C -Vinylidene Chloride to
Male Rats or Mice (Jones and Hathway, 1978b)

^{14}C Excretory Products	^{14}C Expressed at Percent of Dose	
	Mice	Rats
Unchanged vinylidene chloride (pulmonary excretion)	6	28
Carbon dioxide (pulmonary excretion)	3	3.5
Chloroacetic acid	0	1
Thiodiglycolic acid	3	22
Thioglycolic acid	5	3
Dithioglycolic acid	23	5
Thioglycolyloxalic acid	3	2
N-Acetyl-S-cysteiny acetyl (derivative)	50	28
N-Acetyl-S-(2-carboxymethyl) cysteine	4	0
Urea	3	3.5

from the decarboxylation of cysteine. ^{14}C -Thiodiglycolic acid was converted to thioglycolic acid, dithioglycolic acid, and thioglycolyloxalic acid, indicating that these metabolites arise from β -thionase hydrolysis of thiodiglycolic acid (Jones and Hathway, 1978b), shown in Figure 10-1. As can be seen from the relative proportions of thioglycolic acids in Table 10-5, β -thionase conversion occurred to a greater extent in mice than in rats (Jones and Hathway, 1978b).

Results of experiments with rats suggest that conjugation with glutathione serves to detoxify reactive metabolites of vinylidene chloride. The hepatotoxicity of vinylidene chloride in vivo and in the isolated perfused liver was greater if vinylidene chloride was administered when glutathione levels in the liver had been diminished by fasting (Jaeger et al., 1974, 1975a; McKenna et al., 1978a), diurnal variation (Jaeger et al., 1973a, 1975a), by pretreatment with diethyl maleate (Jaeger et al., 1974, 1975a; Andersen et al., 1980), or by pretreatment with thyroxine (Jaeger et al., 1977b). The degree to which pretreatment with various epoxides or diethyl maleate exacerbated the acute toxicity of vinylidene chloride was related to the degree to which these compounds depleted glutathione (Andersen et al., 1980). Increasing the levels of glutathione in the liver by chemical or surgical thyroidectomy protected against the toxicity of vinylidene chloride (Jaeger et al., 1977b). Hypoxia, which inhibited mixed-function oxidase activity, protected against the hepatotoxicity of vinylidene chloride and concomitantly prevented depletion of liver glutathione (Jaeger, 1978).

Additional evidence for the role of glutathione in detoxification can be obtained from studies of covalent binding. Fasting and, hence, a low level of liver glutathione were associated with decreased total metabolism of ^{14}C -vinylidene chloride, increased covalent binding of radioactivity to liver macromolecules, and centrilobular hepatic necrosis in male rats (McKenna et al., 1978a,b).

Data for intragastric and inhalation administration are shown in Tables 10-6 and 10-7, respectively. Binding was measured 72 hours after exposure, as radioactivity that was TCA-precipitable and 80% methanol-insoluble, and thus represents radioactivity covalently bound to protein and nucleic acid. At the higher exposure levels, fasted male rats had significantly greater concentrations of bound radioactivity in their livers when the data were normalized to account for differences in metabolism.

Jaeger et al. (1977a) studied the subcellular distribution of free and bound radioactivity in livers from fed and fasted male rats exposed to ^{14}C -vinylidene chloride. The analyses were performed 30 minutes after the cessation of a 2-hour inhalation exposure to vinylidene chloride (initial concentration, 2000 ppm) in a closed chamber. Significantly more radioactivity was found in all subcellular fractions from fasted animals than was found in those from fed animals. In livers from both fed and fasted rats, the radioactivity in the mitochondrial and microsomal fractions was largely TCA-precipitable; radioactivity in the cytoplasmic fractions was largely TCA-soluble. Because substantial amounts of the TCA-precipitable radioactivity of mitochondria and microsomes were soluble in chloroform, the authors suggested that binding to lipid had occurred. As discussed in Section 10.1.2.1, methylthio-acetylaminoethanol, a metabolite of vinylidene chloride, is thought to be a product of the alkylation of lipids by a reactive intermediate of vinylidene chloride (Reichert et al., 1979; Henschler and Hoos, 1981). The amounts (per mg protein) of radioactivity presumably covalently bound to protein and nucleic acid (TCA-precipitable, chloroform-insoluble material) and to lipids (TCA-precipitable, chloroform-soluble material) were greater in mitochondrial and microsomal fractions from fasted rats than in the corresponding fractions from fed rats. Similarly, fasted rats

Table 10-6

Metabolism of ^{14}C -Vinylidene Chloride and Covalent Binding of ^{14}C Activity
to Rat Hepatic Tissue after Intragastric Dose of ^{14}C -Vinylidene Chloride^a (McKenna et al., 1978b)

Dose	n	A	B	B/A ^b
		Metabolized vinylidene chloride (mg Eq of ¹⁴ C- vinylidene chloride/kg)	Covalently bound vinylidene chloride (μg Eq of ¹⁴ C- vinylidene chloride/g of liver protein)	
1 mg/kg				
Fed rats	4	0.956 ± 0.136 ^c	0.854 ± 0.003	0.89 ± 0.36
Fasted rats	4	0.924 ± 0.032	0.861 ± 0.022	0.93 ± 0.04
50 mg/kg				
Fed rats	3	39.08 ± 3.58	37.67 ± 3.16	0.96 ± 0.12
Fasted rats	3	31.86 ± 1.81 ^d	44.45 ± 3.21	1.40 ± 0.08 ^d

^aAll values represent the $\bar{X} \pm \text{SE}$ for the number of rats indicated in the table.

^bCovalent binding data normalized to account for differences in metabolized vinylidene chloride.

^cCalculated from the total recovery of $^{14}\text{CO}_2$ plus nonvolatile radioactivity from each animal.

^dSignificant fed vs. fasted difference, $p < 0.05$, Student's t test.

Table 10-7

Metabolism of ^{14}C -Vinylidene Chloride and Covalent Binding of ^{14}C Activity to Rat Hepatic Tissue after Inhalation Exposure to 10 or 200 ppm of ^{14}C -Vinylidene Chloride (McKenna et al., 1978a)

Exposure concentration and pretreatment	Body burden (mg of ^{14}C -vinylidene chloride/kg)	A	B	B/A ^a
		Metabolized ^{14}C -vinylidene chloride (mg Eq/kg)	Microgram equivalents of ^{14}C -vinylidene chloride bound per gram of liver protein	
10-24 10 ppm				
Fed	2.89 ± 0.12^b	2.84 ± 0.13^c	2.49 ± 0.17	0.88 ± 0.08
Fasted	2.30 ± 0.05	2.26 ± 0.06	2.47 ± 0.29	1.10 ± 0.15
200 ppm				
Fed	44.53 ± 6.05^d	42.73 ± 3.18^d	64.18 ± 7.97	1.49 ± 0.10
Fasted	35.93 ± 0.30^d	32.92 ± 0.32^d	79.46 ± 4.90	2.42 ± 0.17^d

^aCovalent binding data normalized to account for differences in metabolized vinylidene chloride.

^bAll values represented the mean \pm SE for four rats.

^cCalculated from the total ^{14}C activity recovered (end-exposure body burden) minus the ^{14}C -vinylidene chloride exhaled from each rat.

^dSignificantly different from fed rats exposed to the same concentration of vinylidene chloride, $p < 0.05$, Student's t test.

had greater amounts (per mg protein) of radioactivity bound to protein and nucleic acid of the cytoplasmic fraction than did fed rats.

The relationship among glutathione depletion, total metabolism, and covalent binding to macromolecules is not clear, as shown in Figure 10-3. Immediately after a 6-hour inhalation exposure to ^{14}C -vinylidene chloride, male rats were sacrificed and hepatic glutathione and covalent binding of radioactivity to protein and nucleic acids were measured. The total amount of vinylidene chloride metabolized was apparently measured, as indicated in the legend to Table 10-7. Glutathione depletion and metabolism of vinylidene chloride displayed a similar, non-linear, dose dependence, whereas covalent binding increased linearly with increased exposure concentrations (McKenna et al., 1977; Dedrick, 1979; McKenna, 1979).

Suggestive evidence for an enhancement of hepatic lipid peroxidation in vitro after vinylidene chloride administration in vivo has been obtained in rats under conditions designed to overwhelm the GSH conjugating capacity of the liver: i.e., the rats were pretreated with phenobarbital to induce mixed-function oxidases and were then treated with a very high dose of vinylidene chloride (500 mg/kg) (Siegers et al., 1982). This dose is higher than those used by other investigators to produce hepatic damage in noninduced rats (Section 10.2.1). In general, however, the hepatotoxic effects of vinylidene chloride are not thought to be due to a lipoperoxidative mechanism of action (Section 10.2.1).

10.1.2.2.2 Minor Oxidative Pathway for Monochloroacetic Acid

Yllner (1971) demonstrated the oxidative degradation of ^{14}C -monochloroacetic acid to glycolic acid, oxalic acid, and CO_2 by mice after intraperitoneal administration. S-(Carboxymethyl-1,2- ^{14}C)cysteine was not metabolized to appreciable amounts of labeled CO_2 and hence is not an intermediate in the metabolism of monochloroacetic acid to CO_2 (Yllner, 1971). Trace amounts of

A28264 U



(McKenna *et al.*, 1977)(McKenna *et al.*, 1977, replotted by Dedrick, 1979)

Figure 10-3 Dose-Response Relationship for Hepatic Glutathione (GSH) Levels, Total Metabolism of ^{14}C -Vinylidene Chloride (VDC), and Covalent Binding of Radioactivity to Hepatic Macromolecules.

labeled oxalic acid have been detected in experiments with ^{14}C -monochloroacetic acid in rats (Jones and Hathway, 1978a). Small amounts of labeled CO_2 and urea were identified as metabolites of ^{14}C -vinylidene chloride (see Table 10-5) (Jones and Hathway, 1978b). Thus it appears that the metabolism of monochloroacetic acid to CO_2 is a minor pathway in the metabolism of vinylidene chloride.

10.1.3 Excretion

Vinylidene chloride and its metabolites are excreted fairly rapidly by experimental animals. The influence of route of administration and size of dose on the pattern of excretion of vinylidene chloride and its metabolites by male rats is shown in Table 10-8 (Jones and Hathway, 1978a). According to Jones and Hathway (1978a), 60% of the 0.5 mg/kg intravenous dose of ^{14}C -vinylidene chloride was expired unchanged within 5 minutes of injection; 80% was expired unchanged within 1 hour. These results indicated an efficient arterial-alveolar transfer of vinylidene chloride. With intragastric or intraperitoneal administration of 0.5 mg/kg of ^{14}C -vinylidene chloride, most of the radioactivity was excreted in the urine within the first 24 hours, whereas with 350 mg/kg, most of the radioactivity was expired as unchanged vinylidene chloride. The percentage of radioactivity expired as CO_2 decreased with the higher dose of vinylidene chloride. These changes in the pattern of excretion were attributed by Jones and Hathway (1978a) to saturable drug metabolism and to the efficient transfer of vinylidene chloride from systemic blood to the alveoli, leaving a relatively low concentration of vinylidene chloride available for metabolism in subsequent passes through the liver.

Similar dose-related shifts in the relative proportion of radioactivity excreted by various routes have been reported by Reichert et al. (1979) for intragastric administration of ^{14}C -vinylidene chloride (0.5, 5.0, and 50 mg/kg) to female rats and by McKenna et al. (1978b) for intragastric administration of 1

Table 10-8

Excretion of Radioactivity by Male Rats Given 0.5 mg/kg or 350 mg/kg ^{14}C -Vinylidene Chloride Intragastrically, Intravenously, or Intraperitoneally (Jones and Hathway, 1978a)

Size of Dose	Time (h)	Radioactivity Excreted (Percent of Dose) ^a											
		Intragastric				Intravenous				Intraperitoneal			
		Exhaled Air		Urine	Feces	Exhaled Air		Urine	Feces	Exhaled Air		Urine	Feces
		Vinylidene Chloride	Carbon Dioxide			Vinylidene Chloride	Carbon Dioxide			Vinylidene Chloride	Carbon Dioxide		
0.5 mg/kg	0-24	0.6±0.2	3.9±0.7	71.3±1.2	5.1±0.7	80.0±4.0	3.5±0.6	14.4±3.6	0.3	11.4±2.8	2.6±0.7	65.8±2.5	14.2±4.4
	24-48	0.06	0.5±0.4	5.3±0.9	2.7±0.9	0	0	0.7±0.2	0.1	0.2	0.5±0.1	2.0±0.1	1.6±1.1
	48-72	0.08	0.5±0.2	3.6±1.7	0.6±0.2	0	0	0	0	0.1	0.5±0.2	1.2±0.3	0.4
	Total	0.7±0.1	4.8±1.3	80.2±1.4	8.3±0.1	80.0±4.0	3.5±4.0	15.0±3.9	0.4	11.7±2.8	3.6±0.7	69.0±2.8	16.2±4.5
150 mg/kg	0-24	62.4±4.3	0.3	17.6±4.4	0.4					90.5±2.9	0.7±0.4	7.1±2.1	0.5±0.2
	24-48	4.8±2.8	0.4	10.0±4.0	0.5±0.3					0.6±0.3	0.5±0.2	0.3	0.1
	48-72	0.1	0.3	1.9±1.0	0.4					0	0.1	0.3	0.1
	Total	67.3±4.3	29.5±6.7	1.3±0.4						91.1±3.2	1.3±0.6	7.7±2.1	0.7±0.3

^aValues shown are the Mean ± Standard Deviation of those means (n=four per group).

and 50 mg/kg ^{14}C -vinylidene chloride to fed and fasted male rats. Fed and fasted male rats that inhaled 200 ppm ^{14}C -vinylidene chloride exhaled a greater percentage of their body burden as unchanged vinylidene chloride than did rats exposed to 10 ppm, but this was the only significant dose-related change in excretion (see Table 10-1) (McKenna et al., 1978a). After intragastric administration of 50 mg/kg ^{14}C -vinylidene chloride, fasted male rats exhaled a greater percentage of the dose as unchanged vinylidene chloride and a lesser percentage in the urine than did fed male rats (McKenna et al., 1978b). No such difference was apparent with a dose of 1 mg/kg. These results indicate a reduced capacity for metabolism of vinylidene chloride in fasted animals. Fed/fasted and species differences in excretion that have already been noted in Section 10.1.2 will not be discussed here.

Following intragastric administration of 350 mg/kg ^{14}C -vinylidene chloride, urinary excretion of radioactivity in intact male rats was approximately equal to the sum of the separate urinary and biliary excretions in male rats with biliary fistulae (Jones and Hathway, 1978a). This finding suggests that the origin of part of the urinary radioactivity was via the enterohepatic cycle.

Reported half lives of excretion for different nutritional states (fed versus fasted) and dose levels are compiled in Table 10-9. The only difference between fed and fasted rats was that fed rats excreted urinary metabolites more rapidly during the first 48 hours after administration of 200 ppm than did fasted rats (McKenna et al., 1978b). Changes in half-lives with increasing doses are inconsistent. It is not known whether the inconsistencies could be due to normal experimental variation, to an effect on more than one process, or to possible toxic effects.

10.1.4. Summary of Pharmacokinetics. Vinylidene chloride is readily absorbed by mammals following oral or inhalation exposure. Judging from excretion in the

Table 10-9
Half-Lives of Excretion of ^{14}C -Vinylidene Chloride, $^{14}\text{CO}_2$, and
Radiolabeled Urinary Metabolites of ^{14}C -Vinylidene Chloride

Excretion	Dose		Sex	Rates	Half-lives		Reference
	Route	Amount		Nutritional Status	Rapid Phase	Slow Phase	
<u>Pulmonary</u>							
Vinylidene chloride	intragastric	1 mg/kg	male	fed and fasted*	25 min	117 min	McKenna et al., 1978b
		50 mg/kg	male	fed and fasted*	21 min	66 min	
	intragastric	0.5 mg/kg	female	fed	43 min	---	Reichert et al., 1979
		5.0 mg/kg	female	fed	18 min		
		50.0 mg/kg	female	fed	27 min	381 min	
	inhalation	10 ppm	male	fed and fasted*	20 min	217 min	McKenna et al., 1978a
200 ppm		male	fed and fasted*	21 min	133 min		
Carbon Dioxide	intragastric	0.5 mg/kg	female	fed	4 hr 30 min	35 hr	Reichert et al., 1979
		5.0 mg/kg	female	fed	4 hr 45 min	24 hr	
		50.0 mg/kg	female	fed	2 hr 45 min	20 hr 45 min	
<u>Urinary</u>							
Metabolites	intragastric	1 mg/kg	male	fed and fasted*	6 hr	16.8 hr	McKenna et al., 1978b
		50 mg/kg	male	fed and fasted*	6 hr	16.8 hr	
	intragastric	0.5 mg/kg	female	fed	4 hr 30 min	18 hr 30 min	Reichert et al., 1979
		5.0 mg/kg	female	fed	3 hr 45 min	14 hr	
		50.0 mg/kg	female	fed	5 hr 15 min	25 hr	
	inhalation	10 ppm	male	fed and fasted*	3.1 hr	19.3 hr	McKenna et al., 1978a
		200 ppm	male	fed	4.5 hr	1 hr	
		200 ppm	male	fasted	3.8 hr	23.9 hr	

*One group fed and another group fasted

exhaled air and urine, most of an orally administered dose is absorbed through the gastrointestinal tract. Data for retention (as a percent of the inhaled concentration) during inhalation exposure were not encountered. Vinylidene chloride is metabolized in the liver with a number of possible reactive intermediates, including an epoxide, being formed. These reactive intermediates may react with macromolecules, producing toxic effects, and are detoxified primarily by conjugation with glutathione. Excretion of metabolites and parent compound occurs primarily via the urine and the exhaled air, with greater percentages of the dose being exhaled as unchanged vinylidene chloride at high doses/exposures. Vinylidene chloride does not appear to be stored or accumulated in the tissues.

10.2 ACUTE, SUBACUTE, AND CHRONIC TOXICITY

10.2.1 Acute Exposure

As a class, the halogenated olefins and the chloroethylenes in particular show many similarities in their biologic effects. Anesthesia, hepatotoxicity, and nephrotoxicity are commonly measured indices of acute exposure in experimental animals, although considerable differences exist in relative potency. Factors that influence the toxicity of chloroethylenes include inherent chemical reactivity, species, sex, diet, and exposure to exogenous chemicals that modify drug-metabolizing enzyme activity. With the chloroethylenes (e.g., vinylidene chloride, trichloroethylene, vinyl chloride, perchloroethylene), toxic effects appear to be mediated by the metabolic formation of a critical intermediate, which is probably an epoxide (Andersen and Jenkins, 1977; Andersen et al., 1979b; Reynolds and Moslen, 1977; Jaeger, 1977).

Carpenter and coworkers (1949) published the first report on the toxicity of vinylidene chloride. They indicated that a single 4-hour inhalation exposure to 32,000 ppm was lethal to 2, 3, or 4 rats out of a group of 6 (exact number that died was not specified) over a 14-day observation period. More subjective

information on the acute inhalation toxicity of vinylidene chloride was published in 1962 by Irish (1962) at the Dow Chemical Company. He reported that exposures to 4000 ppm could rapidly produce stupor and unconsciousness, but noted that complete recovery from the anesthetic effect was probable if duration of exposure was brief. No-effect levels for experimental animals were estimated at 1000 ppm for up to 1 hour and 200 ppm for up to 8 hours. These values are in reasonable agreement with the data of Siegel et al. (1971), indicating that the 4-hour LC50 for vinylidene chloride in rats is about 6350 ppm.

Over the past decade, several groups of investigators have explored the biochemical mechanism of vinylidene chloride-induced hepatotoxicity in rats. These studies were prompted by early observations that the toxicity of vinylidene chloride was similar to that of carbon tetrachloride, a well documented hepatotoxin (Jenkins et al., 1972). Initial studies involving the administration of single oral doses (500 mg/kg) to rats established that vinylidene chloride produced biochemical changes that were qualitatively similar to those produced by carbon tetrachloride (Jenkins et al., 1972). These included depressed liver glucose-6-phosphatase activity and increased activities of liver alkaline phosphatase and tyrosine transaminase, and of plasma alkaline phosphatase and alanine aminotransferase. In addition, vinylidene chloride displayed greater potency than did carbon tetrachloride in its effect on several of these biochemical parameters. In contrast to carbon tetrachloride, however, the biochemical alterations induced by vinylidene chloride were less pronounced in female rats and were reduced by pretreatment of animals with phenobarbital, an inducer of hepatic drug-metabolizing enzymes. On the other hand, Carlson and Fuller (1972) demonstrated that enzyme inducers (phenobarbital and 3-methylcholanthrene) as well as enzyme inhibitors (SKF 525A and Lilly 18947) increased the lethality of vinylidene chloride by inhalation in rats, whereas the opposite

effect was seen in rats receiving carbon tetrachloride. These results strongly suggested a mechanism of toxic action for vinylidene chloride that differed from that for carbon tetrachloride. Confirmation of this belief was provided by Jaeger and coworkers (1973a,b), when they demonstrated that the hepatotoxic effects of vinylidene chloride are not due to a lipoperoxidative mechanism of action, as is the case with carbon tetrachloride.

Further studies on the hepatotoxicity of vinylidene chloride have focused on the relationship of metabolism to acute toxicity, and the influence of nutritional status, age, and sex on mortality. Jaeger and coworkers (1973a) observed that the acute lethality of vinylidene chloride administered by inhalation in male rats was greater during the period from 6:00 p.m. to 10:00 p.m. than during the period 6:00 a.m. to 10:00 a.m. This pattern of sensitivity to vinylidene chloride was correlated with the diurnal variation in hepatic glutathione concentration, which is highest during the day and lowest during the night. Thus, the hypothesis was offered that a glutathione-dependent pathway mediates the detoxification of vinylidene chloride, a mechanism which is known to be important in the detoxification of certain carcinogenic chemical intermediates (e.g., epoxides) (Jaeger, 1979). The importance of adequate hepatic glutathione concentrations in protecting against acute intoxication with vinylidene chloride has subsequently been confirmed in studies using fasted animals that have reduced hepatic glutathione levels (Jaeger et al., 1974; Andersen and Jenkins, 1977). In male rats fasted for 18 hours, the 24-hour LC50 following a 4-hour inhalation exposure to vinylidene chloride was 600 ppm, whereas in fed animals an estimated LC50 of 15,000 ppm was obtained (Jaeger et al., 1974). The minimum lethal concentration of vinylidene chloride was 200 ppm in fasted animals and 10,000 ppm for fed animals. The functional state of the thyroid gland, which apparently played a role in the regulation of glutathione levels in the liver, also

influenced the hepatotoxicity of vinylidene chloride (Jaeger et al., 1977b; Szabo et al., 1977). Chemical or surgical thyroidectomy was associated with an increase in hepatic glutathione, and a concomitant decrease in the mortality and hepatic necrosis caused by a 4-hour inhalation exposure of fasted male rats to 2000 ppm vinylidene chloride.

The studies of Jaeger (1977) and his coworkers (Jaeger et al., 1977a,b) led to the hypothesis that the mechanism of toxicity for vinylidene chloride involved damage to hepatic cellular mitochondria. This postulation was supported by evidence showing biochemical alterations following exposure of fasted male rats to vinylidene chloride (200-500 ppm for 1-24 hours); these alterations were indicative of an inhibition of the tricarboxylic acid cycle. Among these changes was a significant elevation of hepatic citric acid concentration in fasted (i.e., glutathione deficient) rats after 12 hours of exposure to 250 ppm vinylidene chloride in air. Thus, the authors suggested that vinylidene chloride may induce mitochondria-specific injury via the metabolic formation of monochloroacetic acid, which may subsequently give rise to monochlorocitric acid.

Studies conducted by Reynolds et al. (1975) and Reynolds and Moslen (1977) supported the theory that vinylidene chloride is a unique hepatotoxin in terms of its rapid action and its effect on mitochondria. The acute hepatotoxicity of chloroethylenes was shown to be, in decreasing order: vinylidene chloride > trichloroethylene > vinyl chloride > perchloroethylene. Exposure of rats to vinylidene chloride by inhalation (200 ppm, 4 hours) produced an abrupt hemorrhagic centrilobular necrosis. Parenchymal cell injury is characterized by retracted cell borders with pericellular spaces forming that may contain red blood cells and fibrin. Nuclear changes include the loss of perinucleolar chromatin and the aggregation of chromatin along the nuclear perimeter against the nuclear envelope. Within 2 hours after onset of exposure, swollen and

ruptured mitochondria were observed in hepatic parenchymal cells, whereas the rough and the smooth endoplasmic reticulum appeared normal. In contrast, trichloroethylene, carbon tetrachloride, vinyl chloride, and perchloroethylene all produced damage to the endoplasmic reticulum.

Reynolds et al. (1980) associated the biochemical changes in the liver, such as sodium, potassium, calcium and glutathione (GSH) levels, with subsequent histological changes in rats exposed to 200 ppm vinylidene chloride for 1 to 4 hours. By the end of the first hour of exposure, sorbital dehydrogenase (SDH) activity was elevated and hepatic Na levels increased while GSH decreased. This trend continued for the second hour of exposure. By the third and fourth hour SDH, as well as serum alanine aminotransferase activity, and sodium and calcium levels were greatly increased above controls. These increases remained high for 12 hours post-exposure. Histological injury included nuclear changes and centrilobular necrosis. Ion changes were attributed to changes in the membrane ion pumps, either directly or due to an adenosine triphosphate (ATP) deficiency. Decreased levels of GSH were attributed to a conjugation reaction in the metabolism of vinylidene chloride. Enzymatic activity increased during detoxification (Reynolds et al., 1980).

Similarly, Chieco et al. (1982) combined histological and chemical analyses to study the hepatotoxic effect of acute exposure to vinylidene chloride in rats. Histochemical testing was conducted at 1, 2, 4, or 6 hours after oral doses of 50, 100, 150, or 200 mg vinylidene chloride. The degree of liver or hepatocellular damage was assessed by histology and by measurements of liver ion levels (Na, Ca and K) and serum transaminase activity. Injury was noted in the form of increased transaminase activity, decreased membrane adenosine triphosphatase activity, decreased succinate dehydrogenase activity (on the inner mitochondrial membrane) and corresponding changes in the plasma and mitochondrial

membrane histology. The investigators (Chieco et al., 1982) concluded that membranous organelles and membrane bound enzymes were the primary sites of hepatocellular damage by vinylidene chloride. These changes in turn resulted in secondary alterations in glutathione levels, cellular ion imbalances, and cytoplasmic changes (Chieco et al., 1982).

Recently, vinylidene chloride has been shown to inhibit the activity of rat liver microsomal calcium pumps in the presence of a NADPH-generating system. The calcium pumps were not inhibited when vinylidene chloride was applied in vitro in the absence of a NADPH-generating system (Moore, 1980; Ray and Moore, 1982). Moore (1982) also reported inhibition of the liver endoplasmic reticulum calcium pump activity by vinylidene chloride. The concomitant rise in liver calcium levels occurred soon after vinylidene chloride administration and was attributed to the inactivity of the calcium pumps. Calcium released from the endoplasmic reticulum may serve as a trigger for the influx of extracellular calcium and, ultimately, cytotoxicity (Moore, 1982).

Information regarding the acute toxicity of vinylidene chloride is not restricted entirely to its effect on the liver. Jenkins and Andersen (1978) recently reported a nephrotoxic action of vinylidene chloride following oral administration to rats. Biochemical indices of kidney damage (plasma urea nitrogen, plasma creatinine) were elevated in fasted male rats given a single oral 400 mg/kg dose of vinylidene chloride (Table 10-10). Conversely, rats that were not fasted were protected from the nephrotoxic effect. At doses below 400 mg/kg, no significant alterations were noted in plasma indicators of kidney damage (Table 10-11). In addition, female rats were much less sensitive to the effects of vinylidene chloride on the kidney. In both male and female rats, however, histological evidence of renal damage was obtained at various times following the administration of a single oral dose of 400 mg/kg vinylidene

Table 10-10
Influence of 24-Hour Fasting on the Effect of Oral
Administration of Vinylidene Chloride (400 mg/kg)
in Corn Oil on Plasma Urea Nitrogen Concentration
in Male Rats (341 ± 7 g)^a

Status	Challenge	Hours After Challenge	Urea Nitrogen (mg/100 ml)
Fasted	Corn Oil	24	21 ± 1^b
		48	20 ± 1
		72	22 ± 1
Fasted	Vinylidene chloride	24	51 ± 16^c
		48	82 ± 39^c
		72	50 ± 3^c
Fed	Corn Oil	24	20 ± 2
		48	21 ± 1
		72	22 ± 1
Fed	Vinylidene chloride	24	19 ± 2
		48	12 ± 2^d
		72	15 ± 1^c

^aSource: Jenkins and Andersen, 1978

^bAll data are expressed as mean \pm SE of groups of 3 to 6 rats.

^cSignificantly higher ($p < 0.05$) than fasted, corn oil-challenged controls at the same time after challenge (method of analysis not mentioned).

^dSignificantly lower ($p < 0.05$) than fed, corn oil-challenged controls at the same time after challenge.

Table 10-11

Relationships of Plasma Indicators of Kidney Damage to Dose 24 Hours
After Oral Administration of Vinylidene Chloride to Male Rats
(289 \pm 9 g)^a

Dose (mg/kg)	Urea Nitrogen (mg/100 ml)	Creatinine (mg/100 ml)
0	20 \pm 2 ^b	0.76 \pm 0.06
50	23 \pm 1	0.61 \pm 0.11
100	26 \pm 1	0.62 \pm 0.06
200	32 \pm 7	0.80 \pm 0.06
400	80 \pm 1 ^c	2.33 \pm 0.05 ^c
800	108 \pm 18 ^c	ND ^d

^aSource: Jenkins and Andersen, 1978

^bAll data are expressed as mean \pm SE of groups of two to six rats.

^cSignificantly different from zero-dose control ($p < 0.05$); however, the method of analysis was not mentioned.

^dNot done

chloride (Table 10-12). Since female rats showed the same, if not greater, histologic damage to the kidney as did males, it was suggested that biochemical parameters may not be sensitive indicators of kidney damage in females.

A single report has been published (Siletnik and Carlson, 1974) concerning the cardiac sensitizing effects of vinylidene chloride by acute inhalation exposure in rats. Cardiac arrhythmias could not be produced by a single 4 µg/kg injection of epinephrine to male rats; however, when rats inhaled 25,600 ppm vinylidene chloride for 80 minutes, doses of epinephrine as low as 0.5 µg/kg could elicit serious cardiac arrhythmias, which could be further enhanced by pretreatment of rats with phenobarbital (50 mg/kg i.p. for 4 days).

Pulmonary damage has been observed in mice after either oral or intraperitoneal administration of vinylidene chloride (Forkert and Reynolds, 1982; Krijgsheld et al., 1983). At 6 hours after administration, a single oral dose of 100 mg/kg produced degeneration of the endoplasmic reticulum of the Clara cells of the bronchiolar epithelium; a single oral dose of 200 mg/kg resulted in necrosis of both ciliated and Clara cells, with exfoliation of the bronchiolar epithelium (Forkert and Reynolds, 1982). Hypoxia (measured as a decrease in oxygen partial pressure of arterial blood) was observed 24 hours after administration of the higher dose, but the animals recovered and the bronchiolar epithelium had regained a normal appearance by 7 days. Clinical chemistry indices of liver damage (SGOT and SGPT levels) were greatly elevated even at the 100 mg/kg dose, 24 hours after administration.

Twenty-four hours after intraperitoneal injection of mice with vinylidene chloride at a dose of 125 mg/kg, microscopic examination of the lungs revealed necrosis and sloughing of the Clara cells lining the bronchioles (Krygsheld et al., 1983). Pulmonary cytochrome P-450 levels and related monooxygenase activities were decreased; the authors suggested that these decreases may have been a

Table 10-12

Comparison of Prevalence of Histopathologic Effects of Oral Administration of Vinylidene Chloride (400 mg/kg) in Male and Female Rat Kidneys (Jenkins and Andersen, 1978)

Time in hours after Vinylidene Chloride Treatment	Sex	Prevalence of Effect					
		Normal	Chronic Inflammation ^a	Pigment	Dilatation ^c	Tubular Necrosis ^d	Vacuolization ^e
10-40	0	M	6/8 ^f	1/8	1/8	--	--
		F	2/4	2/4	--	--	--
	2	M	7/8	--	--	--	1/8
		F	1/4	3/4	--	--	--
	4	M	8/8	--	--	--	--
		F	1/4	3/4	--	--	--
	8	M	4/4	--	--	--	--
		F	--	4/4	--	--	--
	24	M	1/2	--	1/2	--	--
		F	1/4	3/4	1/4	--	--
	48	M	1/3	--	2/3	1/3	--
		F	--	2/4	--	3/4	1/4
	72	M	--	--	2/3	1/3	--
		F	--	--	3/4	4/4	1/4

Table 10-12 (cont.)

Time in hours after Vinylidene Chloride Treatment	Sex	Prevalence of Effect					
		Normal	Chronic Inflammation ^a	Pigment	Dilatation ^c	Tubular Necrosis ^d	Vacuolization ^e
10-41	96	M	--	--	2/3	1/2	--
		F	1/4	--	2/4	1/4	1/4
	120	M	--	--	2/2	1/2	--
		F	--	--	2/4	1/4	1/4
	144	M	--	--	2/2	1/2	--
		F	--	--	3/4	4/4	--

^aFocal collections of mononuclear cells were present.

^bFew to several tubules contained blue-black amorphous material.

^cFocally, the tubules were distended and lined by basophilic regenerated epithelium.

^dThe tubular epithelium, chiefly proximal, showed eosinophilic coagulative necrosis with loss of cellular detail and inflammatory cell infiltration.

^eThe lining epithelium of occasional tubules contained cytoplasmic vacuoles.

^fNumber of tissues in which effect was seen per number of tissues examined.

reflection of the destruction of the Clara cells, which are known to contain relatively high levels of cytochrome P-450. The livers and kidneys of the treated mice were relatively unaffected, as judged by microscopic appearance, cytochrome P-450 content and monooxygenase activities.

10.2.1.1 Mechanisms of Toxicity


The acute toxicity of vinylidene chloride is altered by several factors in addition to the effects on sensitivity produced by fasting. A series of time-course and dose-response studies that explored mechanisms of toxicity for vinylidene chloride have recently been published.

Chieco et al. (1981) examined the effect of the administration vehicle on the toxicity and biological fate of vinylidene chloride in fasted and fed male Sprague-Dawley rats. Vinylidene chloride was given at 200 mg/kg orally in corn oil, mineral oil, or aqueous Tween-80 (0.5%; Sigma). Measurements of exhaled vinylidene chloride (unchanged) were taken at 15 minute intervals for 5 hours and rats were sacrificed at 6 hours for biochemical analyses as an assessment of hepatic injury. The vehicle did not affect the overall amount of vinylidene chloride exhaled, which ranged from 37 to 51% of the administered dose. The nature of the administration vehicle did, however, alter the rate at which vinylidene chloride was absorbed and exhaled. Aqueous Tween-80, which is readily absorbed, allowed the vinylidene chloride to be readily absorbed and exhaled in a shorter period of time. Corn oil was intermediate, while the poorly digested and absorbed mineral oil prolonged absorption and exhalation times in fasted and fed rats. Hepatic injury, measured as elevated glutamic oxalacetic transaminase (GOT) and glutamic pyruvic transaminase (GPT) serum levels, was greatest in fasted rats treated with vinylidene chloride in mineral oil or corn oil (up to 150-fold increase in GOT and GPT levels). Fasted rats treated with vinylidene chloride in Tween-80 were moderately affected (15 to 18 times increase in GOT and

GPT levels). In fed rats treated with vinylidene chloride in corn or mineral oil, enzymatic activity was slightly elevated (2 to 5 times) above the enzymatic activity in fed rats, while GOT was not different from control levels. Chieco et al. (1981) in connection with results of this study suggested that decreased hepatic injury in fed animals is due to the ability of these animals to detoxify vinylidene chloride over a longer period of time than fasted animals.

Histologically, fasted rats fed vinylidene chloride in corn or mineral oil showed massive hepatic necrosis, which was consistent with biochemical changes. Only scattered necrotic hepatocytes were seen in fasted animals given vinylidene chloride in aqueous Tween-80. Fasted animals given vinylidene chloride in corn oil or mineral oil also had granular "heme" casts in Henle's loop of the kidney. Concomitant high levels of free plasma hemoglobin and "pink urine" were noted in these animals. No pathological abnormalities were observed in the heart, lungs, spleen, adrenals, or duodenum in vinylidene chloride-treated animals (Chieco et al., 1981).

Andersen and Jenkins (1977) observed that body size or the age of the animal had a dramatic effect on the toxicity of vinylidene chloride when a single oral dose was given to fasted male rats. Dose-response curves for vinylidene chloride-induced mortality were constructed for animals of three different sizes (Figure 10-4). Mortality among large animals (395 ± 11 g) increased linearly from 0 to 100% at doses between 800 and 2000 mg/kg; however, minimum lethal doses were found to be as low as 50 mg/kg. In medium-sized rats (224 ± 1 g), doses between 50 and 800 mg/kg produced mortality which varied from 10 to 33%. The most aberrant results were obtained in small rats (73 ± 1 g). Mortality increased to 100% at a dose of 300 mg/kg, but then decreased with increases in dose up to 800 mg/kg. The LD_{50} values calculated from the data presented in



- Mortality in groups of 6 rats for 30 rats weighing 395 ± 11 g.
- Mortality in groups of rats for 40 weanling rats weighing 73 ± 1 g.
- △ Mortality in groups of 10 rats for 40 rats weighing 224 ± 1 g.

Figure 10-4 Dose-Mortality Curves for Administration of Single, Oral Doses of Vinylidene Chloride Dissolved in Corn Oil to Fasted Male Rats of Various Sizes (Andersen and Jenkins, 1977a)

Figure 10-4 would appear to be widely variable as shown by dependence on body size.

In an attempt to explain the effect of body size on acute mortality, Andersen and Jenkins (1977) examined the relationship between hepatotoxicity (indicated by plasma enzyme levels) and body size following the administration of a single oral 50 mg/kg dose of vinylidene chloride in corn oil. They found that both mortality within the first 24 hours and plasma enzyme activities after 24 hours were greatest among rats weighing between 130 and 160 g. In animals of this size, the estimated LD₅₀ was less than 50 mg/kg, whereas in large rats, the LD₅₀ would be 30 times greater. The unusual dose-mortality curve for immature rats (see Figure 10-4) was partially explained by the fact that increases in plasma transaminase activities paralleled mortality, and the retention of orally administered vinylidene chloride reached a maximum at a total dose of 100 mg/kg and then leveled off.

In comparison to fasted male rats, female animals were much less susceptible to the effects of vinylidene chloride. Plasma transaminase levels were not altered by a 50 mg/kg dose, regardless of body size. In female rats, the threshold dose for expression of acute toxic effects (plasma transaminase elevation) was about 100 mg/kg. Based on their results and on analogy to other halogenated hydrocarbons, Andersen and Jenkins (1977) postulated that vinylidene chloride is metabolized by a saturable microsomal enzyme-mediated reaction to a toxic intermediate that is probably an epoxide. A second enzyme-mediated reaction that converts the toxic intermediate to a less toxic form was also proposed. In young rats, this detoxification mechanism may be absent or present at only low levels, thus accounting for increased susceptibility.

In subsequent studies, Andersen and coworkers (1978, 1979b) examined in greater detail the relationship between vinylidene chloride metabolism and

expression of toxic effects. Chemical modifiers of microsomal enzyme activity were administered to fasted male rats followed by a single oral dose of vinylidene chloride (Andersen et al., 1978). Phenobarbital pretreatment, which causes microsomal enzyme induction, protected rats against the lethal effects of vinylidene chloride. This protection by phenobarbital was opposite to that observed by Carlson and Fuller (1972) following inhalation exposure of rats. This protective action increased with increasing body size. Conversely, the microsomal enzyme inhibitor SKF 525A greatly enhanced the lethality of vinylidene chloride in large rats (261 ± 2 g) but had little influence in immature rats (88 ± 2 g). In addition, pretreatment of rats with the metabolic inhibitors pyrazole, aminotriazole, and carbon tetrachloride protected animals of all sizes against the lethal action of vinylidene chloride. These results supported the argument that two sequential enzyme-mediated reactions are involved in the activation of vinylidene chloride. The first reaction, which is inhibited by pyrazole, aminotriazole, and carbon tetrachloride, leads to formation of a toxic intermediate. The second reaction, which is affected by phenobarbital and SKF 525A pretreatment, is a detoxification step that apparently exists at greater levels in mature animals.

Subsequent studies by Andersen and coworkers (1979b) demonstrated that mortality induced in rats by inhalation of vinylidene chloride failed to follow a strict concentration x time relationship. Instead, mortality was more dependent on time of exposure than on concentration once a threshold level (about 200 ppm) had been reached (Figure 10-5). These data supported the involvement of a saturable enzymatic reaction in the production of a metabolite that is responsible for the expression of vinylidene chloride-induced toxicity. The authors appropriately noted that, under conditions of exposure that produce enzyme




Figure 10-5 Effect of Increasing Concentration of Vinylidene Chloride on Mortality in Mature Male Rats (Anderson et al., 1978) (Each point represents percentage mortality of a group of 6 animals exposed for 4 hours.)

saturation, misleading estimates of the LC_{50} (inhalation) or LD_{50} (oral and parenteral) could result.

Studies by Short and coworkers (1977a,b) with rats and mice support the conclusions reached by Andersen et al. (1978, 1979b). Non-fasted animals (groups of 10) inhaled vinylidene chloride for 22 to 23 hours per day for up to 7 days. Short et al. (1977a,b) found that male mice were more sensitive than were male rats to lethal effects and hepatotoxic effects (measured by serum glutamic-oxaloacetic transaminase and serum glutamic-pyruvic transaminase activities) of vinylidene chloride (Table 10-13). Histopathologic examination revealed hepatic and renal damage among male mice exposed to 15, 30, and 60 ppm vinylidene chloride (1-5 days); hepatic damage only was seen in male rats exposed at 60 ppm for 3 days. When mice were given disulfuram (0.10% in feed 23 days before and during exposure), or thiram (0.10% in feed 3 days before and during exposure), the acute lethality of vinylidene chloride was reduced. The authors observed that these compounds are known to protect against drug-induced toxicity, and may act by decreasing metabolic activation or increasing detoxification or both.

10.2.2 Subacute and Chronic Exposure

10.2.2.1 Inhalation Studies

Several groups of investigators have characterized the toxicity of vinylidene chloride by repeated administration to experimental animals. Both ingestion and inhalation have been employed as routes of exposure. These studies established that, in general, the target organs affected by chronic exposures (i.e., liver and kidneys) are the same as those affected by acute administration of vinylidene chloride.

The first published report regarding the chronic toxicity of vinylidene chloride was a brief summary of inhalation studies conducted by the Dow Chemical Company (Irish, 1962). Unspecified animals exposed to 100 ppm and 50 ppm vinyli-

Table 10-13

Toxicity of 60 ppm Vinylidene Chloride in Male Mice and Rats
(Short et al., 1977b)

Species	Days Exposed	SGOT (IU/l) ^a	SGPT (IU/l) ^a	Ratio Dead/Exposed
mouse	1	1946 \pm 270	3045 \pm 209	2/10
	2	751 \pm 150	1112 \pm 226	8/10
rat	1	74 \pm 6	44 \pm 7	0/10
	2	263 \pm 33	198 \pm 29	0/10

^aMean \pm SE for 2 to 5 determinations

SGOT = Serum glutamic-oxaloacetic transaminase; SGPT = Serum glutamic-pyruvic transaminase

dene chloride (5 days per week, 8 hours per day for several months) developed liver and kidney damage. Minimal lesions were also observed in the liver and kidney of animals exposed at concentrations of 25 ppm.

Prendergast and coworkers (1967) published the first detailed account of the chronic toxicity of vinylidene chloride. Groups of rats, rabbits, dogs, and monkeys were exposed in one of two ways: 30 exposures, 8 hours per day, 5 days per week to a concentration of 395 mg/m^3 (100 ppm) or 90 days of continuous exposure to a concentration of 189 mg/m^3 (48 ppm), 101 mg/m^3 (26 ppm), 61 mg/m^3 (16 ppm), or 20 mg/m^3 (5 ppm). The results of these studies are summarized in Table 10-14. The most prominent effects observed included hepatic damage in dogs, monkeys, and rats exposed to 189 mg/m^3 ; this damage consisted of fatty metamorphosis, focal necrosis, hemosiderin deposition, lymphocytic infiltration, bile duct proliferation, fibrosis, and pseudo-lobule formation. Rats also showed nuclear hypertrophy of the renal tubular epithelium when exposed to vinylidene chloride at 189 mg/m^3 .

Gage (1970) conducted limited studies in rats exposed to vinylidene chloride by inhalation that support the previous observations of Prendergast and coworkers (1967). Four male and 4 female rats received twenty 6-hour exposures to vinylidene chloride at concentrations of 500 or 200 ppm. At the higher concentration, retarded weight gain and nasal irritation were noted. Liver cell degeneration was observed upon autopsy. Inhalation of 200 ppm vinylidene chloride produced slight nasal irritation but no significant findings at autopsy; however, the group size was very small.

Lee and coworkers (1977) exposed CD-1 mice (36 males and 36 females) to 55 ppm of vinylidene chloride for 6 hours per day, 5 days per week, for up to 12 months. Two male mice died on the 13th day of exposure, and revealed acute toxic hepatitis and tubular necrosis of the renal cortex. Hepatic hemangiosarcomas were

Table 10-14

Effect on Experimental Animals of Long Term Inhalation of Vinylidene Chloride
(data from Prendergast et al., 1967)

Concentration	Schedule	Species	Mortality	Significant Findings
100 ppm (395 \pm 32 mg/m ³)	30 exposures, 8 hr/day, 5 days/week	rat	0/15	None.
		guinea pig	0/15	None.
		rabbits	0/3	Weight loss in treated animals.
		dog	0/2	None.
		monkey	0/3	Weight loss in treated animals.
48 ppm (189 \pm 6.2 mg/m ³)	90 days, 24 hr/day	rat	0/15	Animals gained less weight than controls. Hepatic lesions (see text). Renal lesions (see text).
		guinea pig	7/15	Mortality occurred between day 4 and day 9 of exposure. Slight elevation of liver alkaline phosphatase activity and serum glutamic-pyruvic transaminase activity.
		dog	0/2	Animals lost weight. Hepatic lesions (see text). One dog developed an adrenal cortical adenoma.
		monkey	3/9	Mortality occurred on days 26, 60 and 64. Animals lost weight. Hepatic lesions (see text).

Table 10-14 (cont.)

Concentration	Schedule	Species	Mortality	Significant Findings
26 ppm (101 \pm 4.4 mg/m ³)	90 days, 24 hr/day	rat	0/15	None.
		guinea pig	3/15	Mortality occurred between day 3 and day 5 of exposure.
		rabbit	0/3	Animals lost weight.
		dog	0/2	Animals lost weight.
		monkey	2/3	Mortality occurred between day 3 and day 6 of exposure.
16 ppm (61 \pm 5.7 mg/m ³)	90 days 24 hr/day	rat	0/15	Animals gained less weight than controls.
		guinea pig	3/15	Mortality occurred on day 3 and day 4.
		dog	0/2	None.
		monkey	0/9	Animals lost weight.
5 ppm (20 \pm 2.1 mg/m ³)	90 days 24 hr/day	rat	2/45	Animals gained less weight than controls.
		guinea pig	2/45	None.
		dog	0/6	Animals lost weight.
		monkey	1/21	None.
Control		rat	7/304	
		guinea pig	2/314	
		rabbit	2/48	
		dog	0/34	
		monkey	1/57	

observed in three exposed mice, and various hepatic lesions that included enlarged and basophilic hepatocytes, enlarged nuclei with eosinophilic inclusions, mitotic figures or polyploidy, microfoci of mononuclear cells, focal degeneration, and necrosis were generally observed (specific incidences were not given). Hepatic hemangiosarcomas were not observed in any of 72 control mice. Bronchiolo-alveolar adenomas were observed in six exposed mice and one control mouse.

Dow Chemical Company performed an inhalation toxicity study with vinylidene chloride and interim results were published by Rampy and coworkers (1977). In a 90-day study, groups of 40 Sprague-Dawley rats (20 male and 20 female) were exposed for 6 hours per day, 5 days per week, to vinylidene chloride at a concentration of 25 or 75 ppm. Eight animals of each sex were sacrificed after 30 days, and the remainder killed after 90 days. Among animals exposed at both concentrations, increased cytoplasmic vacuolation of hepatocytes was seen 30 days or longer after initial exposure. No other remarkable effects were attributed to the vinylidene chloride exposure.

In another study, Rampy and coworkers (1977) exposed Sprague-Dawley rats for 18 months (6 hours/day, 5 days/week) to vinylidene chloride at several concentrations in air. For the first 5 weeks, rats (10⁴ males and 10⁴ females) were exposed to levels of 10 or 40 ppm, after which the concentrations were raised to 25 and 75 ppm, respectively, for the remainder of the 18-month exposure period. Animals were maintained for an additional 6 months after termination of exposure. Although male rats gained weight at a slightly slower rate than controls, no clinical evidence of overt toxicity was seen during the study. Both male and female rats sacrificed at an interim of 1 year, however, showed hepatic lesions consisting of increased cytoplasmic vacuolation of hepatocytes. In

addition, higher kidney weights were seen in female rats exposed at both treatment levels of vinylidene chloride.

The final data of this 2-year inhalation study (interim results reported by Rampy et al., 1977) were recently compiled and reported by McKenna et al. (1982). Data were reported on hematologic parameters, urinalysis, clinical chemistry, cytogenic changes in the bone marrow, body weight, major organ weights, and histopathology following necropsy on all animals (either at interim periods of 1, 6, 12, or 18 months or at the termination of the experiment at 734 to 736 days). Statistical evaluations of data were made using analysis of variance and Dunnett's test (body weight, hematology, urinalysis, clinical chemistry, and organ weight data) or by the Fischer exact test (mortality rates, gross and microscopic pathologies, and tumor incidence). The level of significance was chosen as $P \leq 0.05$.

Mortality figures for male rats exposed to vinylidene chloride at either dose level did not reveal significant increases. Females exposed at 25 ppm vinylidene chloride had increased mortality rates when compared to controls from the 15th through 22nd months. Statistically significant increased mortality rates in females exposed at 75 ppm vinylidene chloride were noted during months 15, 17, and 21 of the study. The earlier onset of mammary tumors and higher incidences of pneumonia in treated groups may have contributed to these higher mortality rates.

No consistent dose-related decrease in mean body weight for male rats exposed to 25 or 75 ppm vinylidene chloride was observed. A general trend of decreased body weight was noted in exposed males when compared to control males but increased mean body weights were noted in the exposed groups sporadically throughout the study. Generally, females exposed to either 25 or 75 ppm vinyli-

dene chloride had higher mean body weights for the first 6 months but were comparable to controls throughout the remainder of the study.

No significant dose-related effects in hematological, urinalysis, clinical chemistry, or cytogenetic parameters were reported in any exposure group. The only significant changes in organ weight were a decreased mean liver weight in males exposed to 25 or 75 ppm vinylidene chloride at the 1-year interim sacrifice and an increased mean kidney weight in both exposure groups of female rats at 1 year. Both organ weight changes were statistically significant when compared to controls and considered to result from vinylidene chloride exposure (McKenna et al., 1982).

Many gross and microscopic lesions were seen in control and vinylidene chloride-exposed rats; most were spontaneously occurring, age-related changes common to Sprague-Dawley rats. The incidence of lesions from chronic murine pneumonia in exposed male and female rats was elevated, but lower levels in controls were attributed to the physical separation of exposed and control groups which deterred the spread of pneumonia. Chronic renal disease was the predominant cause of death in male rats of the study. Females had a high incidence of subcutaneous tumors of the mammary gland region which was frequently the cause of death. There was a statistically significant increase in the incidence of hepatocellular fatty change in female rats exposed to 75 ppm during the 18 month exposure period. These changes were reversed during the 6 month post-exposure period, however. Females treated at 25 ppm showed a similar but non-significant trend. This reversible and non-progressive liver change was noted in males at the 6 and 12 months interim sacrifice but not at 18 months.

The incidence of pituitary adenomas was decreased in both dose groups of male rats and the 25 ppm vinylidene chloride dose group of females. There were fewer pancreatic islet cell adenomas and thyroid adenocarcinoma in male rats

exposed to 75 ppm vinylidene chloride. Mammary adenocarcinomas were statistically increased in female rats treated at 25 ppm vinylidene chloride. Subcutaneous or mammary tumors were more prevalent in 75 ppm vinylidene chloride treated females than in controls. The total number of rats with primary neoplasms was similar for both sexes at all dose groups, including controls. The total number of rats with primary neoplasms was similar (ranging from 84 to 86) for all 6 dose/sex groups. Summarizing the tumor incidence data, the authors reported that none of the statistically significant differences were considered to be direct effects of vinylidene chloride exposure.

10.2.2.2 Ingestion Studies

A single long-term study has been conducted with vinylidene chloride administered in the drinking water of rats. This study was conducted by the Dow Chemical Company for the Manufacturing Chemists' Association (Humiston et al., 1978). Groups of 96 Sprague Dawley rats (48 males and 48 females) were exposed for 2 years to vinylidene chloride incorporated into the drinking water at nominal concentrations of 50, 100, and 200 ppm. These dose levels corresponded to approximate daily intakes of vinylidene chloride in the range of 5-12, 8-20, and 16-40 mg/kg at the 50, 100, and 200 ppm concentrations, respectively. In comparison to control animals, vinylidene chloride-treated rats displayed no significant or consistent differences in general appearance, body weight, food consumption, water consumption, hematologic values, urinalysis, clinical chemistry values, or organ weights. Gross and histopathologic examination of tissues from treated rats, however, revealed a number of statistically significant lesions. These are partially summarized in Table 10-15. The authors considered the most important lesions to be the hepatocellular fatty change and periportal hepatocellular hypertrophy which occurred in male rats at the 200 ppm

Table 10-15

Pathologic Effects of Long-Term Ingestion of Vinylidene Chloride
Incorporated in the Drinking Water of Sprague-Dawley Rats
(data from Humiston et al., 1978)

Effect	Dose Level					
	50 ppm		100 ppm		200 ppm	
	M	F	M	F	M	F
Increased incidence of intra-abdominal fluid or blood in the abdominal cavity			X			
Increased incidence in the total number of rats with hepatocellular fatty change or fatty degeneration				X	X	X
Increased incidence of hepatocellular fatty change with location in lobule not specified		X		X		X
Increased incidence in periportal hepatocellular fatty change						X
Increased incidence of periportal hepatocellular hypertrophy		X		X	X	X
Increased incidence of hepatic centrilobular atrophy						X
Increased incidence of mammary gland fibroadenomas/adenofibromas		X				

M = male; F = female

dose level and in females at all dose levels. The authors did not observe any hepatocellular necrosis that could be considered treatment-related.

Results of this long-term, oral study with rats were subsequently published in the open literature by Quast et al. (1983), along with results of a 97-day oral study with dogs; the dog study was also conducted by the Dow Chemical Company. Groups of four male and four female beagle dogs were administered vinylidene chloride in doses of 0, 6.25, 12.5 or 25 mg/kg daily for 97 days. The chemical or vehicle (peanut oil) was administered orally in a capsule once a day. As compared with controls, the groups of treated dogs had no differences in general appearance or demeanor, body weight, food consumption, hematologic and clinical chemistry values, urinalysis, organ weights and gross and microscopic appearance of the tissues.

10.2.3 Summary of Toxicity

The biologic activity of vinylidene chloride has been thoroughly studied in experimental mammals; no information is available concerning the effects of vinylidene chloride on domestic animals or wildlife.

The toxicity of vinylidene chloride varies with the age, sex, and species of the animal exposed. In the most studied animal, rat, inhalation exposure to 32,000 to 6,350 ppm has proven to be fatal. In rats fasted prior to exposure, the LC_{50} value was as low as 600 ppm. For oral administration of vinylidene chloride, however, the determination of an LD_{50} value is difficult as a result of the relatively flat dose mortality curve which possesses an extended plateau region. The target organs affected by acute exposure to vinylidene chloride are the liver and kidneys with evidence from studies of the liver to indicate that mitochondria damage occurs. Both liver and kidney damage are also observed following subchronic and chronic exposure to vinylidene chloride by either inhalation or ingestion. Continuous inhalation exposure of rats to 5 ppm of vinylidene

chloride has been shown to adversely affect body weight gain while ingestion of vinylidene chloride by rats at 5 to 12 mg/kg/day produced fatty livers. A no-observed effect level (NOEL) for inhalation exposure (<5 ppm) or ingestion (<5 mg/kg/day) has not been demonstrated.

10.3 TERATOGENICITY AND REPRODUCTIVE TOXICITY

Short et al. (1977c) performed an extensive investigation of the toxicity during gestation of inhaled vinylidene chloride in mice and rats (approximately 20 animals in the exposed groups and 60 in the control groups). This study examined toxicity to the dams, toxicity to the fetus, and teratogenic and behavioral effects in the pup. In the initial part of the experiment, rats (18 to 20 per group) were exposed to 15, 57, 300, and 449 ppm vinylidene chloride, and mice to 15, 30, 57, 144, and 300 ppm vinylidene chloride for 23 hours per day from day 6 to 16 of gestation. In rats, appreciable deaths (over 25%) of the dams occurred in the 300 to 449 ppm group, and maternal well being, as measured by food consumption and weight gain, was adversely affected in the 15 ppm and above groups. Similar results were observed in mice, with no pregnant mice surviving at the 144 or 300 ppm levels, and food consumption and weight gain being reduced in all groups except the 15 ppm animals. Early resorptions were common in exposed animals, with 100% resorption in mice exposed to 30 and 57 ppm vinylidene chloride (resorption in the 15 ppm groups was comparable to controls), and 49 and 64% resorption in rats exposed to 57 and 449 ppm vinylidene chloride, respectively. Dams that survived the exposure at other concentration levels had resorption rates comparable to those of controls. There were no increases in external gross abnormalities noted in surviving pups in any of the exposed groups of rats or mice; however, fetotoxicity was observed as soft tissue anomalies in rats (none were observed in mice) and some skeletal ossification problems in all groups of mice and rats. The authors concluded that these anomalies could

not be evaluated in regard to teratogenic effects, since there were overt signs of toxicity (reduced weight gain and food consumption) due to the treatment in the dams. Abnormalities at a similar incidence to those observed in the treated animals were also observed in a group of animals on a food-restricted diet.

The study was continued using only mice, with a group of 10 to 24 dams exposed to a variety of concentrations of vinylidene chloride for different periods of time during gestation (Table 10-16), in an attempt to alleviate maternal toxicity as usually indicated by decreased maternal weight gain. By shortening the exposure periods, the incidence of resorption was lessened; however, treatment-related weight loss in the dams was still evident. It was interesting to note that exposures starting on days 10 and 12 and ending on day 15 were associated with a higher resorption rate than that observed when exposures were started on day 8 of gestation. This may have occurred because of some adaptation of the animals as a result of the longer exposure prior to a critical period during the latter part of gestation. Pups from vinylidene chloride-exposed dams again had a variety of soft tissue and skeletal anomalies, but these occurred only when maternal welfare had been adversely affected by the exposure to vinylidene chloride. In addition, the behavioral studies demonstrated no major adverse effect on pups exposed in utero to vinylidene chloride. These tests included: surface righting, pivoting, auditory startle, bar holding, righting in air, visual placing, swimming ability, physical maturation, and activity test. The authors concluded that vinylidene chloride was possibly a weak teratogen based on the increase in soft tissue anomalies although these anomalies were not statistically significant. However, a conclusion of teratogenicity is weakened by additional effects on the pups as a result of maternal toxicity at nearly all concentrations of vinylidene chloride used.

Table 10-16

Exposure Levels and Duration of Exposure to Vinylidene Chloride
During Gestation in Mice (Short et al., 1977c)

Vinylidene Chloride ppm	Days of Gestation											
	6	7	8	9	10	11	12	13	14	15	16	17
54	6-----									15		
74			8-----							15		
54			8-----							15		
41			8-----							15		
54					10-----					15		
54							12-----			15		
112	6-----			9								
81	6-----			9								
56	6-----			9								
112				9-----			12					
81				9-----			12					
56				9-----			12					
112							12-----			15		
81							12-----			15		
56							12-----			15		
112										15-----	17	
81										15-----	17	
56										15-----	17	

----- = Duration of Exposure

Murray et al. (1979) exposed rabbits to vinylidene chloride by inhalation, and rats by inhalation and by incorporating the compound into the drinking water (Table 10-17). Exposure occurred during days 6-15 or 6-18 of gestation for rats and rabbits, respectively. Inhalation exposure was for 7 hours/day to vinylidene chloride which was 99.5% pure and contained the polymerization inhibitor monomethylether of hydroquinone (MEHQ) at a level of 200 to 400 ppm. An exposure level of 200 ppm was employed in the drinking water study using redistilled vinylidene chloride containing only 1 to 5 ppm of residual MEHQ. On the termination of the observation period, on days 21 and 29 of gestation for rats and rabbits, respectively, the dams were killed and examined for changes in liver weight, and in non-pregnant animals for signs of conception. The fetuses were measured, examined for external anomalies, and one-third of the animals were dissected for soft-tissue alteration, while all fetuses were examined for skeletal alterations.

In the inhalation study in rats, maternal weight gain was decreased in the 80 and 160 ppm groups during the exposure period and increased during the post-exposure period, while no changes were observed in the 20 ppm groups or the animals in the drinking water study. Food consumption followed similar trends. Only in dams exposed to 160 ppm was there an increase in the liver to body weight ratio, although the absolute liver weight was similar to control animals. Exposure to vinylidene chloride did not affect the outcome of pregnancy as indicated by number of implants, live fetuses, or resorption rates, nor were there changes in fetal sex ratios or fetal body weights. There was a significant increase in the crown-rump length in fetuses of rats receiving vinylidene chloride in the drinking water. Although no increase in major malformations (taken individually or collectively) was present, the fetuses of animals exposed in the 80 and 160 ppm groups showed an increased incidence of minor skeletal alterations (Table

Table 10-17

Number of Animals and Exposure Levels Used to Study
the Teratogenicity of Vinylidene Chloride*

Route	Concentration	Species	No. of Animals
inhalation	160	rats	30
	160	rabbits	18
	80	rats	30
	80	rabbits	22
	20	rats	44
	0	rats	20-47/group
	0	rabbits	16
oral	200	rats	26
	0	rats	24

*Source: Murray et al. (1979)

10-18). The most common skeletal alteration was wavy ribs, which was observed to be more severe at the higher exposure level. Wavy ribs and delayed ossification were interpreted as fetotoxic and embryotoxic manifestations of maternal toxicity at the higher exposure levels. Rats receiving vinylidene chloride in the drinking water were estimated (using data from metabolism studies) to have received the equivalent of a 7-hour inhalation exposure to 120 ppm of vinylidene chloride. It was speculated that a lack of toxic effects in the rats maintained on drinking water containing 200 ppm vinylidene chloride as compared to the high dose inhalation group may have resulted from diurnal variation in metabolism, although other mechanisms are also possible.

In rabbits exposed to 160 ppm of vinylidene chloride, there was a decrease in body weight gain during exposure and an increase after termination of exposure. Only dams exposed at 80 ppm had an increase in liver-to-body-weight-ratio. There was also an increase in the resorption rate in the 160 ppm group, with resorption associated with animals having the greatest weight loss. Similar to the study in rats, exposure to vinylidene chloride produced no increase in major malformations, although there was an increase in minor skeletal alterations in the 160 ppm group (Table 10-19). Again, the fetal anomalies observed in rabbits were attributed to maternal toxicity rather than to a teratogenic effect of vinylidene chloride.

The effect of vinylidene chloride on reproduction and its teratogenic potential when ingested in the drinking water was tested in a three generation study in Sprague-Dawley rats (Nitschke et al., 1980; Nitschke et al., 1983). The parental generation (F_0), consisting of 10 male and 20 females in each of three treatment groups, received 50, 100, or 200 ppm vinylidene chloride. No vinylidene chloride was administered to a control group of 15 males and 30 females. Equivalent doses for males ingesting 50, 100, or 200 ppm vinylidene chloride

TABLE 10-18

Incidence of Fetal Alterations Among Rats Exposed to Vinylidene Chloride
By Inhalation or by Ingestion^a

	Route of exposure ^b (ppm vinylidene chloride)						Drinking Water	
	Inhalation							
	0	20	0	80	0	160	0	200
	No. fetuses examined/No. litters examined							
External and skeletal examination	470/40	462/40	230/20	292/27	209/17	291/26	250/24	290/25
Soft tissue examination	162/40	158/40	77/20	103/27	70/17	100/26	86/24	100/25
Bones of the skull	308/39	304/39	153/20	189/26	139/17	191/26	164/23	190/24
External examination	No. fetuses (litters) affected							
Multiple defects ^c	1(1)	0	0	0	1(1)	0	0	0
Omphalocele	0	0	0	1(1)	0	0	0	0
Narrow head, point snout	0	0	0	1(1)	0	0	0	0
Soft tissue examination								
Diaphragmatic hernia	0	0	1(1)	0	0	0	0	0
Ectopic ovaries (only)	1(1)	0	0	0	0	0	0	0
Skeletal examination								
Missing one thoracic and one lumbar vertebrae	0	0	0	0	2(1)	14(4)	0	0
Extra thoracic vertebrae and pair of ribs	0	0	0	0	1(1)	0	0	0
Vertebrae, delayed ossification of centra of cervical vertebrae ^d	34(17)	58(19)	24(14)	47(17)	30(13)	107(20) ^e	46(16)	72(18)
Wavy ribs ^d								
Type I ^f	5(5)	10(7)	0	9(8) ^e	2(1)	9(5)	0	3(2)
Type II ^f	4(4)	3(2)	0	2(2)	1(1)	13(9) ^e	1(1)	0
Total	9(7)	13(7)	0	11(9) ^e	3(1)	22(10) ^e	1(1)	3(2)
Skull, delayed ossification ^d	42(17)	36(18)	13(7)	38(13) ^e	24(8)	59(20) ^e	19(10)	32(11)
Total malformed fetuses	2(2)	0	1(1)	2(2)	4(2)	14(4)	0	0

^aSource: Murray et al., 1979

^bRats were exposed to 20, 80, or 160 ppm vinylidene chloride for 7 hours/day or were given drinking water containing 200 ppm of the compound from days 6 to 15 of gestation.

^cEach of these fetuses had: short trunk, hypoplastic tail, ectopic ovaries, missing all lumbar and sacral vertebrae, and missing ribs.

^dThis alteration was considered to be a skeletal variant and was not included in the calculation of the total malformed fetuses.

^eSignificantly different from control value by a modified Wilcoxon test, $p < 0.05$

^fType I, gentle wave; Type II, callous or "U"-shaped bend in rib

TABLE 10-19

Incidence of Fetal Malformation Among Litters of Rabbits
Exposed to Vinylidene Chloride^a

	Vinylidene Chloride ^b (ppm)			
	0	80	0	160
	No. fetuses/No. litters examined			
External and skeletal examination	116/14	155/18	111/13	91/12
Soft tissue examination	44/14	56/18	42/13	37/12
External examination	No. fetuses (litters) affected			
No malformation observed				
Soft tissue examination				
Dilated cerebral ventricles	0	1(1)	0	0
Thinning of ventricular wall of heart	0	0	0	1(1)
Skeletal examination				
Missing ribs	0	0	1(1)	0
Extra vertebrae	0	1(1)	1(1)	0
Hemivertebrae and fused ribs	0	1(1)	0	1(1)
Total malformed fetuses	0	3(3)	2(2)	2(2)

^aSource: Murray et al., 1979

^bRabbits were exposed to 80 or 160 ppm vinylidene chloride for 7 hr/day from days 6 to 15 of gestation. No value differed significantly from the control value by a modified Wilcoxon test, $p < 0.05$.

continuously in the drinking water were 6, 10, or 19 mg/kg/day, and 8, 13, or 26 mg/kg/day, respectively. Following 100 days of exposure to vinylidene chloride, all groups of males and females (f_0) were mated (within dose groups), resulting in the first filial (f_{1A}) generation. Due to the low fertility rate during this initial breeding, the f_0 rats were re-mated after a period of no less than 10 days after weaning of the f_{1A} , which produced the f_{1B} offspring. The f_0 generation rats were exposed to vinylidene chloride at the three respective doses for a 2-year toxicity study (Humiston et al., 1975). The f_{1A} and f_{1B} litters of all dose groups were examined for reproductive indices (number of litter/number of dams; survival of pups at 1, 7, 14, and 21 days). The animals to be raised as the f_1 generation adults were randomly selected from the f_{1B} generation and continued to be exposed to vinylidene chloride. All other f_{1A} and f_{1B} rats were sacrificed at 21-24 days of age for internal and external examinations.

At ≈ 110 days of age, the f_1 adults were mated, producing the f_2 progeny. The f_2 litters were examined similar to the previous litters, and randomly selected males and females were maintained for f_2 adults. The remaining f_2 weanlings were terminated and examined for external and internal aberrations. Adult f_2 rats were mated at ≈ 110 days of age to produce the f_{3A} generation. Survival of the f_{3A} litters in all groups receiving vinylidene chloride was decreased so the f_2 adults were re-mated after the f_{3A} litters were weaned. To determine if the decreased survival in the dose groups was due to prenatal or postnatal exposure to vinylidene chloride, one-third of the f_{3B} litters born to 200 ppm vinylidene chloride-treated dams were exchanged with litters of untreated dams. No dose-related decreased survival was noted in any groups of the f_{3B} litter. At least 10 days after the f_{3B} litters were weaned, the f_2 parents were mated again to produce the f_{3C} litters to determine survival rates of this group. A selected number of male and female rats from each dose group of

the f_{3B} litters were raised for 185-213 days as f_3 adults. All remaining weanling rats of the f_{3A} , f_{3B} , and f_{3C} litters were sacrificed at 21-24 days of age for external and internal examination (Nitschke et al., 1980).

The water containing the vinylidene chloride was available continuously throughout this study to all treated rats. The first generation rats (f_0) were treated during their maturation, during mating periods, throughout gestation and lactation for females, and were subsequently used in a 2-year toxicity study. Pups were exposed prenatally and throughout their life until their sacrifice, or longer if they were selected as breeders of that generation (Nitschke et al., 1980).

The results of this study indicate few, if any, dose-related effects on reproduction or the development of progeny in rats treated with vinylidene chloride in the drinking water at the dose levels tested (Nitschke et al., 1980). A slight decrease in the fertility (number of litters/number of mated dams) of the f_0 rats was noted in controls and the two highest dose groups (120 and 200 ppm) when compared to historical controls of Sprague-Dawley rats. Survival of f_2 progeny was decreased in all treated groups but this response was not dose-related since animals treated at 100 ppm vinylidene chloride showed the greatest mortality, while groups treated at 50 and 200 ppm were similar. Decreased survival during the lactation period in the f_{3A} generation did appear dose-related, but survival in the f_{3B} litters of dams given 100 and 200 ppm vinylidene chloride was higher than controls. It was not possible to ascertain the cause, whether it be prenatal or postnatal exposure to vinylidene chloride, of the decreased survival in the f_{3A} generation mice. No decrease in any group of the f_{3B} , including those cross-fostered (control group dams rearing 200 ppm progeny and vice versa), was noted. The decreased survival of the f_{3A} generation was

negated since the similarly treated f_{3C} generation showed no dose-related, decreased survival.

Fetal survival, litter size, and growth rates of the neonates showed no consistent dose-related effects for any generation during this study. At necropsy, histopathological examination revealed hepatocellular fatty change and pronounced hepatic lobular patterns in adult f_1 rats treated with 100 or 200 ppm vinylidene chloride. Fatty changes of the hepatocytes were also noted in adult f_2 females. Liver degeneration was noted in f_1 adults treated with 100 or 200 ppm vinylidene chloride and at all dose levels in f_2 adults. No dose-related malformations or changes in organ weights were revealed after necropsy of weanlings in any dose group or generation (Nitschke et al., 1980).

In summary, the three generation study by Nitschke et al. (1980) indicated that vinylidene chloride in drinking water of rats at 50, 100, or 200 ppm did not adversely affect reproduction or neonatal development. Neonatal survival was decreased in the f_{3A} litters, but contradictory results were shown in the f_{3B} and f_{3C} litters. Histopathological examination revealed mild dose-related hepatotoxic effects in adult rats treated with vinylidene chloride, but no dose-related changes or malformations were seen in neonates. Reproductive capacities in adults did not appear altered by vinylidene chloride exposure (Nitschke et al., 1980).

10.4 Mutagenicity

10.4.1. Mutagenicity in Bacteria

A variety of assay methods have been used to assess the mutagenic potential of vinylidene chloride. These include the standard Salmonella typhimurium plate incorporation assay of Ames, the liquid suspension assay, the host mediated assay, as well as exposure of the bacteria to an atmosphere containing vinylidene chloride. Due to the volatile nature of vinylidene chloride, the most commonly

used assay method involves the exposure of the tester bacteria (in combination with a mammalian metabolic activating system, if used) to a defined atmosphere of vinylidene chloride in a desiccator. Following the exposure period, the bacteria are incubated in petri dishes to allow the growth of mutant colonies.

Bartsch et al. (1975) exposed bacteria of S. typhimurium strains TA100 and TA1530 to an atmosphere of 0.2, 2.0, or 20% vinylidene chloride (v/v) in air for 4 hours. The concentration of vinylidene chloride in the agar after 2 hours of exposure was shown by gas chromatography to be 3.3×10^{-4} , 3.3×10^{-3} , and 3.3×10^{-2} M, respectively, with no further increases in the vinylidene chloride concentration for up to 7 hours of exposure. In the presence of a metabolic activation system prepared from the livers of mice pretreated with phenobarbital (0.1% in the drinking water for 7 days), both TA100 and TA1530 showed increases in the number of revertant colonies. The maximum mutation rate occurred in TA100 with 2.0% vinylidene chloride (≈ 11 -fold); it was suggested that the higher concentrations of vinylidene chloride caused inactivation of the activating enzymes. Vinylidene chloride has recently been shown to inhibit the activity of microsomal cytochrome P-450 in vitro (Poptawski-Tabarelli and Uehleke, 1982). The same group of workers (Bartsch et al., 1975) have repeatedly demonstrated the sensitivity of TA100 to the mutagenic effect of vinylidene chloride in a study characterizing the stability of the metabolic activation system (Malaveille et al., 1977). It was shown that these results were not due to the stabilizer 4-methoxyphenol present in the vinylidene chloride in these assays.

In more limited studies, other investigators have demonstrated the mutagenic potential of vinylidene chloride following vapor phase exposure of S. typhimurium. In an investigation of suspected mutagenic compounds in drinking water, Simmon (Simmon et al., 1977; Simmon, 1978) demonstrated a six-fold increase (125 in controls to 650 in tested) in revertant colony number using

tester strain TA100 in the presence of rat liver homogenate following exposure for 9 hours to a 5% (v/v) atmosphere of vinylidene chloride. Also, Baden et al. (1976) used a 3% atmosphere (exposure period of 8 hours) and Waskell (1978) used a 5% atmosphere (exposure period of 4 hours) of vinylidene chloride as a positive control in a study assessing the mutagenic potential of volatile compounds. TA100 and TA98 showed a four-fold and 2.5-fold increase, respectively, compared to controls in revertant colony number in the presence of a metabolic activation system from Aroclor 1254-pretreated female BDV1 rats. In the absence of metabolic activation, the number of revertants did not vary significantly from the spontaneous controls.

The mutagenicity of vinylidene chloride (99.9% pure) has also been demonstrated when the bacteria (E. coli or S. typhimurium) were exposed to the compound in sealed tubes in liquid media (liquid suspension assay). Greim et al. (1975) used a strain of E. coli K12 sensitive to back mutation (gal^+ , arg^+ , and nad^+) and one forward mutation system which leads to resistance to 5-methyl-DL-tryptophan. Only the reverse mutation in the Arg^+ strain showed an increased number of revertant colonies (2.3-fold over control) following exposure of the bacteria to 2.5 mM vinylidene chloride for 2 hours. A liver homogenate from phenobarbital-pretreated mice was employed for metabolic activation. Baden and coworkers (Baden et al., 1976, 1977, 1978) have used the liquid suspension assay with S. typhimurium to assess the mutagenicity of volatile anesthetics, and have used vinylidene chloride as a positive control for these assays. Both tester strain TA1535 and TA100 showed positive responses in the presence of a mammalian metabolic activating system after a 1-hour exposure to vinylidene chloride at a concentration that made the head space gas 3% vinylidene chloride.

An abstract (Cerna and Kypenova, 1977) reported vinylidene chloride to be mutagenic in the host-mediated assay with S. typhimurium strains TA1950, TA1951,

and TA1952. Vinylidene chloride was administered to ICR mice at doses equal to the LD₅₀ and one-half the LD₅₀ dose. There were fewer revertant colonies at the higher exposure level. This assay has been demonstrated to have a poor correlation with known genotoxic compounds (Simmon et al., 1979).

Many compounds (pro-mutagens) are inactive until metabolized by mammalian enzyme systems to ultimate mutagens. Metabolism occurs in many cases via the mixed-function oxidases associated with the microsomal fraction of a tissue homogenate. Vinylidene chloride requires metabolic activation in order to produce a mutagenic response in bacterial in vitro assays. In a single report in which vinylidene chloride was not mutagenic (using DNA repair-deficient B. subtilis and a comparison of zones of inhibition with the wild type and the standard Ames plate incorporation assay), a metabolic activation system was not employed (Laumbach et al., 1977).

Livers from a variety of species have been used to metabolically activate vinylidene chloride. These include livers from mice, rats, and humans (Baden et al., 1976; 1977). Of these species, the 9000 x g supernatant (S9) containing microsomal and soluble enzymes from the livers of mice was most effective, particularly when the enzyme systems were induced by pretreatment of the animals with phenobarbital or Aroclor 1254. Liver biopsy samples from 4 humans were only 11, 16, 17, and 38% as effective in activating vinylidene chloride as was mouse liver, as indicated by the relative numbers of revertant colonies in S. typhimurium strains TA100 or TA1530 (Bartsch, 1976; Bartsch et al., 1979). Bartsch et al. (1975) have demonstrated that S9 prepared from the liver, kidney, or lung of OF-1 male mice was effective in activating vinylidene chloride, although the lung possessed only marginal activity (Table 10-20). The ability to metabolically activate vinylidene chloride may have wide tissue and species

Table 10-20

Mouse Tissue Mediated Mutagenicity of Vinylidene Chloride in S. typhimurium
(modified from Bartsch et al., 1975)

Experiment Number	Species	Phenobarbitone Pretreatment	Tissue (9000 x g) supernatant) ^a	Cofactors ^b	2% Vinylidene Chloride in Air, his ⁺ Revertant per plate ^c	20% Vinylidene Chloride in Air, his ⁺ Revertants per plate ^c
1	OF-1 mouse/male	Yes	liver	+	500 ± 23	330 ± 29
2	OF-1 mouse/male	Yes	liver	-	23 ± 10	7 ± 5
3	OF-1 mouse/male	No	liver	+	330 ± 49	435 ± 46
4	OF-1 mouse/male	No	liver	-	16 ± 4	1 ± 3
5	OF-1 mouse/male	Yes	kidney	+	147 ± 15	173 ± 5
6	OF-1 mouse/male	Yes	kidney	-	31 ± 7	17 ± 2
7	OF-1 mouse/male	No	kidney	+	67 ± 2	125 ± 5
8	OF-1 mouse/male	No	kidney	-	20 ± 3	16 ± 1
9	OF-1 mouse/male	Yes	lung	+	34 ± 4	48 ± 5
10	OF-1 mouse/male	Yes	lung	-	5 ± 4	10 ± 1
11	OF-1 mouse/male	No	lung	+	21 ± 5	37 ± 3
12	OF-1 mouse/male	No	lung	-	6 ± 9	14 ± 8

^aEquivalent to 38 mg wet tissue per plate

^bNADP⁺ (2.0 μmol per plate) and glucose-6-phosphate (2.5 μmol per plate)

^cMean values ± S.E. from 1-4 experiments, each using pooled tissue from 4 mice. The number of spontaneous mutations per plate (49 ± 2) has been subtracted from each value.

distribution, but it is apparent that there is considerable variability in the efficiency of activation by different S9 preparations.

There has been speculation as to the nature of the ultimate mutagen formed during metabolic activation of vinylidene chloride. Barstch et al. (1975) have suggested that an alkylating intermediate is formed, since mutations occur in S. typhimurium strain TA1530--which is reported by Ames et al. (1973) to be specifically reverted by monofunctional alkylating agents. Nevertheless, an alkylating compound was not identified in the trapping agent 4-(4'-nitrobenz)-pyridine when a gas that contained vinylidene chloride was passed through a mouse S9 activating system, even though this system provided activated metabolites that mutated S. typhimurium strain TA100 (Barbin et al., 1978). It has also been suggested that an unstable oxirane, possibly 1,1-dichloroethylene oxide, may be formed during metabolism and that this metabolite plays a significant role in the mutagenic activity of vinylidene chloride (Greim et al., 1975; Bonse et al., 1975; Bartsch et al., 1979). Presently, however, there is little evidence identifying the exact metabolic intermediate or intermediates that constitute the ultimate mutagenic form of vinylidene chloride (see Section 10.1.2.1).

10.4.2 Mutagenicity in Plants and Yeast

The plant Tradescantia has been used as a detector of gaseous mutagens in the laboratory and in tests conducted in the field. This assay system involves a phenotypic change in flower color from blue to pink as a measure of mutagenic pollutants in the atmosphere. The plants are highly sensitive and produce an effect from contaminated ambient air. Collection and concentration of the gaseous contaminants is not necessary. A mutation (or loss) of a dominant gene (for blue color) results in the expression of recessive pink flower pigmentation. In this assay system, vinylidene chloride exposure for 24 hours at 22 ppm was sufficient to induce the mutagenic response. A maximum concentration at which no

effects was observed after a 6-hour exposure was 1288 ppm (Van't Hof and Schairer, 1982).

Bronzetti et al. (1981) reported that vinylidene chloride was mutagenic with D7 yeast (Saccharomyces cerevisiae) only in the presence of a mammalian activation system. When a mouse hepatic supernatant was included, vinylidene chloride (above 20 mM) was effective in increasing point mutations and gene conversions in a dose-related response. In an intrasanguineous host-mediated assay, vinylidene chloride was mutagenic to yeast (injected into mice) which were removed from the liver and kidneys after both acute (400 mg/kg by gavage to mice) and subacute (100 mg/kg/day, 5 times/week for 23 administrations) vinylidene chloride exposure.

10.4.3 Mutagenicity in Cultured Mammalian Cells

Drevon and Kuroki (1979) exposed V79 Chinese hamster cells in a desiccator to an atmosphere of 2 or 10% vinylidene chloride for 5 hours. Following exposure, the cells were assayed for viability and mutations to 8-azaguanine, or ouabain resistance. Dose-dependent toxicity was observed in cells exposed to vinylidene chloride in the presence of a 15,000 x g supernatant from rat liver. Cells exposed to vinylidene chloride in the presence of a 15,000 x g supernatant from mouse liver or in the absence of a metabolic activating system showed no apparent toxic effect. Both rats and mice were pretreated with 1.0% phenobarbital in drinking water for 7 days prior to sacrifice. There was no indication under any of the experimental conditions of an increase in 8-azaguanine or ouabain resistance.

10.4.4 Mutagenicity In Vivo

Dominant lethal studies with vinylidene chloride have been performed in male CD-1 mice (Anderson et al., 1977) and in CD rats (Short et al., 1977d). Male mice (20 animals/group) were exposed to 10 and 30 ppm of vinylidene chloride for

6 hours per day for 5 days. One group was exposed to 50 ppm for 6 hours per day for 2 days; however, the poor survival of these animals precluded any assessment of mutagenic damage. The male rats (11 animals) were exposed to 55 ppm vinylidene chloride for 6 hours per day, 5 days per week for 11 weeks. At the end of these exposure periods, the animals were mated with nonexposed virgin females. In neither rats nor mice did the treatment result in an excess of preimplantation or postimplantation losses. In the studies performed by Andersen et al. (1977), a positive control of cyclophosphamide was included to assure the sensitivity of this strain of mouse in the dominant lethal assay.

10.4.5 In Vivo DNA Repair

Reitz et al. (1980) used CD-1 male mice and Sprague-Dawley male rats to study the effect of vinylidene chloride on DNA synthesis and repair in the liver and kidneys. The mice were exposed at 10 ppm for 6 hours. Total DNA synthesis was measured by ^3H -thymidine incorporation into DNA 48 hours after exposure. DNA repair synthesis was measured immediately after exposure to vinylidene chloride by ^3H -thymidine incorporation into DNA in the presence of hydroxyurea. Hydroxyurea inhibits replicative DNA synthesis, but allows DNA repair synthesis to occur. The ability of vinylidene chloride to alkylate DNA was also examined by exposing the animals to ^{14}C -vinylidene chloride followed by isolation and analysis of the DNA from the liver and kidneys. Vinylidene chloride was most efficient at alkylating the DNA in mouse kidney, with 30 alkylations/ 10^6 nucleotides in animals exposed to 50 ppm vinylidene chloride and 11 alkylations/ 10^6 nucleotides in animals exposed to 10 ppm. In mouse liver, the values were 6.1 and 0.94 alkylations/ 10^6 nucleotides in the 50 and 10 ppm exposure groups, respectively. In rats, the rate of alkylation was 2 and 0.87 alkylations/ 10^6 nucleotides in the kidney and liver, respectively. These alkylation rates were considerably less than the 3000 to 4000 per 10^6 nucleotides observed in rats with

the potent carcinogen dimethylnitrosamine (DMN). The elimination of alkylated-nucleotides was rapid, with a biphasic elimination pattern. The first phase occurred during the first 8 hours and the second phase continued for 192 hours.

The pattern of DNA repair synthesis was similar to that of alkylation. Some repair synthesis was noted in both kidney and liver; however, only the synthesis in the kidneys of mice exposed to the highest concentration was significantly elevated over control levels. Although repair synthesis was increased only marginally following vinylidene chloride treatment, total DNA synthesis was increased markedly in the kidney of mice. This indicates that vinylidene chloride caused cell damage and compensatory growth, but did not interact with DNA directly and cause repair synthesis. The liver carcinogen DMN caused little increase in total DNA synthesis, while dramatically increasing DNA repair synthesis. The authors concluded that the kidney tumors in mice reported by Maltoni et al. (1977) were not the direct result of a genetic effect of vinylidene chloride, but may have been the result of an epigenetic effect following the kidney toxicity.

10.5 CARCINOGENICITY

10.5.1 Animal Studies

There have been a number of laboratory investigations of the carcinogenic potential of vinylidene chloride (VDC). These studies have been performed using rats, mice, and hamsters, with VDC administered by inhalation, gavage, incorporation into drinking water, subcutaneous injection, and topical applications. These results are summarized in Table 10-21, and have also been summarized by Chu and Milman (1981). Of the studies performed, only the results reported by Maltoni et al. (1977, 1980) indicated a positive carcinogenic effect following VDC treatment.

Maltoni et al. (1980, 1977) exposed Sprague-Dawley rats, Swiss mice, and Chinese hamsters to VDC (99.95% pure, Table 10-22) in an inhalation chamber for 4 hours daily, 4 to 5 days a week for 12 months. The exposure concentration used depended on the species' susceptibility to VDC's toxic action; the highest tolerated dose in mice and rats was used as the maximum concentration. The exposure level for hamsters was not selected on the basis of achieving a maximum tolerated dose. Rats, initially 16 weeks old, were exposed to VDC at 150, 100, 50, 25, and 10 ppm (Table 10-23), while mice, initially 16 weeks or 9 weeks old, were chronically exposed to VDC at 25 and 10 ppm (Table 10-24), and hamsters, initially 28 weeks old, to a concentration of VDC at 25 ppm (Table 10-25). In addition, rats, initially 9 weeks old, were treated by gavage with 20, 10, 5, and 0.5 mg/kg/day of VDC dissolved in olive oil (Table 10-26). Body weights were recorded every 2 weeks during treatment and monthly thereafter. At the time of their 1980 report, the studies were completed (137 weeks for the inhalation study in rats, 147 weeks for the

TABLE 10-21. RESULTS OF CARCINOGENICITY BIOASSAYS OF VINYLIDENE CHLORIDE

Species	Dose	Route of administration	Total duration of observation	Findings	Reference
Sprague-Dawley rats	10, 25, 50, 100, 150 ppm 4 to 5 days/week for 12 months	Inhalation 4 hr/day	137 weeks	Statistically significant increase in total mammary tumors, but not carcinomas alone, only at 10 and 100 ppm, no dose response	Maltoni et al., 1980, 1977
Swiss mice	10, 25 ppm 4 to 5 days/week for 12 months	Inhalation 4 hr/day	121 weeks	Kidney carcinomas at 25 ppm in males (none in controls) Statistically significant increase in mammary carcinomas in females, no dose response.	Maltoni et al., 1980, 1977
Chinese hamsters	25 ppm 4 to 5 days/week for 12 months	Inhalation 4 hr/day	157 weeks	No statistically significant increase	Maltoni et al., 1980, 1977
Sprague-Dawley rats	20, 10, 5, 0.5 mg/kg for 12 months	Gavage daily	147 weeks	No statistically significant increase	Maltoni et al., 1980, 1977
Wistar rats	200 ppm for 6 months, followed by 100 ppm for 6 months, 5 days/week	Inhalation 4 hr/day	Lifetime	No statistically significant increase	Viola and Caputo, 1977
Sprague-Dawley rats	100, 75 ppm 5 days/week for 12 months	Inhalation 4 hr/day	Lifetime	No statistically significant increase	Viola and Caputo, 1977
CD-1 mice	55 ppm 5 days/week	Inhalation 6 hr/day	12 months	No statistically significant increase	Lee et al., 1978
CD rats	55 ppm 5 days/week for 12 months	Inhalation 6 hr/day	12 months	No statistically significant increase	Lee et al., 1978
Sprague-Dawley rats	25, 75 ppm for 24 months	Inhalation	104 weeks	No statistically significant increase	McKenna et al., 1982
Sprague-Dawley rats	50, 100, 200 ppm in drinking water	Ingestion	104 weeks	No statistically significant increase	Quast et al., 1983

TABLE 10-21. (continued)

Species	Dose	Route of administration	Total duration of observation	Findings	Reference
Fischer 344 rats	5 ml/kg of a 1000 or 200 ppm solution	Gavage 5 days/week	103 weeks	No statistically significant increase	NCI/NTP, 1981
B6C3F1 mice	10 ml/kg of a 1000 or 200 ppm solution	Gavage 5 days/week	103 weeks	No statistically significant increase	NCI/NTP, 1981
CD mice	55 ppm, 5 days/week 1, 3, or 6 months	Inhalation 6 hr/day	13, 15, or 18 months	No statistically significant increase	Hong et al., 1981
CD rats	55 ppm, 5 days/week 1, 3, 6, or 10 months	Inhalation 6 hr/day	13, 15, 18, or 22 months	No statistically significant increase	Hong et al., 1981
Ha:ICR Swiss	121 mg/mouse	Skin application, 3 times/week	Lifetime	No tumors	Van Duuren et al., 1980
Ha:ICR Swiss	2 mg/mouse	Subcutaneous injection once/week	Lifetime	No tumors at site of injection or in other organs	Van Duuren et al., 1980
Sprague-Dawley rats	10, 25, 50, 100, 150 ppm	Inhalation 4 to 5 days/week	52 weeks	No brain tumors	Maltoni et. al., 1982
Sprague-Dawley rats	0.5, 5, 10, 20 mg/kg/day	Gavage, 5 days/week	52-59 weeks	No brain tumors	Maltoni et al., 1982

TABLE 10-22. GAS CHROMATOGRAPHY ANALYSIS OF VINYLIDENE CHLORIDE

Compound	Amount
Vinylidene chloride (1,1-Dichloroethylene)	999.5 g/kg
1,2-Dichloroethylene trans	0.4 g/kg
Acetone	0.1 g/kg
Methylene chloride	0.05 g/kg
Monochloroacetylene and Dichloroacetylene	0.02 g/kg
Paramethoxyphenol (as stabilizer)	200 ppm

Source: Maltoni et al., 1977

TABLE 10-23. EXPERIMENT DT401: EXPOSURE BY INHALATION TO VINYLIDENE CHLORIDE (VDC) IN AIR AT 150, 100, 50, 25, 10 PPM, 4 HOURS DAILY, 4-5 DAYS WEEKLY, FOR 52 WEEKS

GROUP NOS.	CONCENTRATION	ANIMALS (Sprague-Dawley rats, 16 weeks old at start)		
		Males	Females	Total
I	200 (a) - 150 ppm	60	60	120
II	100 ppm	30	30	60
III	50 ppm	30	30	60
IV	25 ppm	30	30	60
V	10 ppm	30	30	60
VI	No treatment	100	100	200
Total		250	280	560

(a) Two treatments only because of the high toxicity of this dose level.

Source: Maltoni et al.; 1980.

TABLE 10-24. EXPOSURE BY INHALATION TO VINYLIDENE CHLORIDE (VDC) IN AIR
AT 200, 100, 50, 25, 10 PPM, 4 HOURS DAILY,
4-5 DAYS WEEKLY, FOR 52 WEEKS

GROUP NOS.	TREATMENT		ANIMALS Swiss mice 16 weeks old (groups I, II, III, IV, V, VI) and 9 weeks old (groups IV bis and VII)		
	Concentration	Length	Males	Females	Total
I	200 ppm	2 days (d)	60	60	120
II	100 ppm	2 days (d)	30	30	60
III	50 ppm	1 week (d)	30	30	60
IV	25 ppm	52 weeks	30	30	60
IV bis (a)	25 ppm	52 weeks	120	120	240
V	10 ppm	52 weeks	30	30	60
VI	No treatment (Controls)(b)		100	100	200
VII	No treatment (Controls)(c)		90	90	180
Total			490	490	980

- (a) The treatment started two weeks later than in other groups.
 (b) Controls to the groups I, II, III, IV, V.
 (c) Controls to the group IV bis.
 (d) The treatment was interrupted because of the high toxic effects and high mortality.

Source: Maltoni et al., 1980.

TABLE 10-25. EXPERIMENT BT405: EXPOSURE BY INHALATION TO VINYLIDENE CHLORIDE (VDC) IN AIR AT 25 PPM, 4 HOURS DAILY, 4-5 DAYS WEEKLY, FOR 52 WEEKS

GROUP NOS.	CONCENTRATION	ANIMALS (Chinese hamsters, 28 weeks old at start)		
		Males	Females	Total
I	25 ppm	30	30	60
II	No treatment (Controls)	18	17	35
Total		48	47	95

Source: Maltoni et al., 1980.

TABLE 10-26. EXPERIMENT BT403: EXPOSURE BY INGESTION (STOMACH TUBE) TO VINYLIDENE CHLORIDE IN OLIVE OIL AT 20, 10, 5 MG/KG BODY WEIGHT, ONCE DAILY 4-5 DAYS WEEKLY, FOR 52 WEEKS

GROUP NOS.	CONCENTRATION	ANIMALS (Sprague-Dawley rats, 9 weeks old at start)		
		Males	Females	Total
I	20 mg/kg	50	50	100
II	10 mg/kg	50	50	100
III	5 mg/kg	50	50	100
IV	None (olive oil alone) (Controls)	100	100	200
Total		250	250	500

EXPERIMENT BT404: EXPOSURE BY INGESTION (STOMACH TUBE) TO VINYLIDENE CHLORIDE IN OLIVE OIL AT 0.5 MG/KG BODY WEIGHT, ONCE DAILY, 4-5 DAYS WEEKLY, FOR 52 WEEKS

GROUP NOS.	CONCENTRATION	ANIMALS (Sprague-Dawley rats, 9 weeks old at start)		
		Males	Females	Total
I	0.5 mg/kg	50	50	100
II	None (olive oil alone) (Controls)	82	77	159
Total		132	127	259

Source: Maltoni et al., 1980.

ingestion study in rats, 121 weeks for the inhalation study in mice, and 157 weeks for the inhalation study in hamsters. Animals were allowed to survive until spontaneous death. Each animal was necropsied, and tissues and organs, as well as tumors and lesions, were examined histopathologically.

No treatment-related effect on body weights in rats was evident. Body weight data for mice and hamsters were not reported, but no effect attributable to treatment was noted.

There were no significant ($P < 0.05$) differences in tumor development between control and treated hamsters.

A number of neoplastic lesions were observed in control and treated rats, with no treatment- or dose-related effect in evidence from either inhalation exposure or gavage administration. The 1977 report by Maltoni et al. presented interim (82 weeks) results showing an increased incidence of mammary tumors in treated rats compared to controls in the inhalation study; however, at that time, a clear dose-related effect was not evident, spontaneous mammary tumor incidence in female controls was 32%, and histopathologic diagnosis of the mammary tumors was ongoing. The final mammary tumor data for female rats in this study, as described in the Maltoni et al. (1980) report and Table 10-27 herein, show statistically significant ($P < 0.05$) increases in the total number of tumor-bearing animals in the 10 ppm and 100 ppm groups compared to controls, and in the number of animals with fibromas and fibroadenomas in each treatment group compared to controls. However, there was no clear dose-related increase in mammary tumor incidence, latency time for mammary tumor formation was similar among all groups, there was a high (61%) incidence of spontaneously formed mammary tumors in controls, and mammary carcinoma

TABLE 10-27. EXPERIMENT BT401: EXPOSURE BY INHALATION TO VINYLIDENE CHLORIDE (VDC) IN AIR AT 150, 100, 50, 25, 10 PPM, 4 HOURS DAILY, 4-5 DAYS FOR 52 WEEKS. RESULTS AFTER 137 WEEKS (END OF EXPERIMENT).

DISTRIBUTION OF THE DIFFERENT TYPES OF MAMMARY TUMORS

MAMMARY TUMORS (b)																	
GROUP NOS.	CONCENTRATIONS	ANIMALS (Sprague-Dawley rats, 16 weeks old at start)				Total No.	% (c)	Average latency time (weeks) (d)	No. of tumors/tumor-bearing animals	Total No.	% (e)	Histologically Examined					
		Sex	No. at start	Corrected number (a)	Histotype												
					Fibromas and fibroadenomas							Carcinomas					
					No.							% (f)	Average latency time (weeks) (d)	No.	% (f)	Average latency time (weeks) (d)	
I	150 ppm	M	60	60	8	13.3	97±14	1.0	8	100.0	6	75.0	109± 8	1	12.5	26	
		F	60	60	44	73.3	82± 3	1.5	43	97.7	38*	88.4	83± 3	9	20.9	78± 8	
		M and F	120	120	52	43.3	82± 3	1.4	51	98.1	44	86.3	86± 3	10	19.6	73± 8	
II	100 ppm	M	30	30	5	16.7	104± 9	1.0	5	100.0	5	100.0	104± 9	0	-	-	
		F	30	30	25*	83.3	82± 4	1.7	23	92.0	21*	91.3	83± 5	3	13.0	102±10	
		M and F	60	60	30	50.0	85± 4	1.6	28	93.3	26	92.8	87± 4	3	10.7	102±10	
III	50 ppm	M	30	30	7	23.3	106± 5	1.0	7	100.0	7	100.0	106± 5	0	-	-	
		F	30	30	23	76.7	79± 4	1.9	22	95.6	21*	95.4	82± 4	1	4.5	68	
		M and F	60	60	30	50.0	86± 4	1.7	29	96.7	28	96.5	88± 4	1	3.4	68	
IV	25 ppm	M	30	28	4	14.3	103±10	1.0	4	100.0	4	100.0	103±10	0	-	-	
		F	30	30	21	70.0	86± 4	1.6	20	95.2	20*	100.0	87± 4	4	20.0	82±10	
		M and F	60	58	25	43.1	88± 4	1.5	24	96.0	24	100.0	90± 4	4	16.7	82±10	
V	10 ppm	M	30	29	3	10.3	81±23	1.0	3	100.0	3	100.0	81±23	0	-	-	
		F	30	30	28*	93.3	83± 4	1.6	24	85.7	24*	100.0	85± 4	5	20.8	90±14	
		M and F	60	59	31	52.5	81± 4	1.5	27	87.1	27	100.0	85± 4	5	18.5	90±14	
VI	No treatment (Controls)	M	100	87	11	12.6	115± 6	1.0	11	100.0	11	100.0	115± 6	0	-	-	
		F	100	99	61	61.6	87± 2	1.5	56	91.8	44	78.6	88± 3	16	28.6	95± 5	
		M and F	200	186	72	38.7	91± 3	1.4	67	93.0	55	82.1	93± 3	16	23.9	95± 5	
Total			560	543													

(a) Alive animals after 10 weeks, when the first tumor (a leukemia) was observed.

(b) Two or more tumors of the same and/or different types (fibroadenomas, carcinomas, sarcomas, carcinosarcomas) may be present in the same animals. A carcinosarcoma was found in one male in the 150 ppm group, and no animals were observed to have sarcomas.

(c) The percentages refer to the corrected numbers.

(d) Average age at the onset of the first mammary tumor per animal, detected at the periodic control or at autopsy.

(e) The percentages refer to total numbers of animals bearing mammary tumors.

(f) The percentages refer to total numbers of animals bearing mammary tumors, histologically examined.

* Statistically significant increase compared to control by chi-square test (P<0.05). Comparisons are made between numbers with tumors/corrected numbers.

incidence in treated groups was not significantly ($P < 0.05$) different from and was actually consistently less than that of controls. Hence, the evidence for a carcinogenic effect of inhaled vinylidene chloride in female Sprague-Dawley rats, as mammary tumors, in this study would appear to be inconclusive.

In male Swiss mice, kidney adenocarcinomas were observed following inhalation exposure to VDC at 25 ppm (Table 10-28). In female Swiss mice, mammary carcinoma incidence in both the 10 ppm and combined 25 ppm groups compared to combined controls was significantly ($P < 0.01$) increased, and a small incidence of mammary carcinomas was evident in females exposed to VDC at 50, 100, or 200 ppm for 1 week or less (Table 10-28). The incidence of pulmonary adenomas was significantly ($P < 0.01$) increased in male and female mice exposed to VDC at 10 and 25 ppm compared to controls (Table 10-28); however, pulmonary carcinomas were not found in any of the mice.

Maltoni et al. (1980) concluded that the kidney tumors in male mice developed in response to VDC treatment, particularly since no corresponding spontaneous tumors were noted in the 190 control male mice. Maltoni (1977) also concluded that the observed kidney tumors in Swiss mice were a strain-specific phenomenon. It has been noted that male mice of several strains are particularly sensitive to the toxic effects of VDC, and that a direct relationship may exist between the degree of toxicity and the carcinogenic effect of VDC (Maltoni et al. 1977; Maltoni 1977; Henck et al. 1980). These authors postulated that a metabolite is responsible for both effects of VDC. Maltoni et al. (1977) noted that degeneration and necrotic changes in kidneys, especially in the tubular region, in male Swiss mice which died from exposure to 200 ppm VDC, 4 hours daily, for 2 days were observed as

TABLE 10-28. EXPERIMENT BT402: EXPOSURE BY INHALATION TO VINYLIDENE CHLORIDE (VDC) IN AIR AT 200, 100, 50, 25, 10 PPM, 4 HOURS DAILY, 4-5 DAYS FOR 52 WEEKS. RESULTS AFTER 121 WEEKS (END OF EXPERIMENT).

DISTRIBUTION OF THE DIFFERENT TYPES OF MAMMARY TUMORS

GROUPS NOS.	TREATMENT		ANIMALS (Swiss mice 16 weeks old (Grs I,II, III,IV,V,VI) and 9 weeks old (Grs IV bis, VII) at start)		ANIMALS WITH TUMORS											
	CONCEN- TRATIONS	LENGTH			Kidney adenocarcinomas				Mammary Tumors ^c				Pulmonary adenomas ^d			
					Corrected number (a)	No.	%	Average la- tency time (weeks)(b)	Corrected number (a)	No.	%	Average la- tency time (weeks)(b)	Corrected number (a)	No.	%	Average la- tency time (weeks)(b)
			Sex	No. at start												
I	200 ppm	2 days	M	60	1	0	-	-	6	0	-	-	5	0	-	-
			F	60	28	0	-	-	53	1	1.9	87	46	1	2.2	57
			M and F	120	29	0	-	-	59	1	1.7	87	53	1	1.9	57
II	100 ppm	2 days	M	30	12	0	-	-	21	0	-	-	18	2	11.1	62± 7
			F	30	13	0	-	-	28	3	10.7	46± 3	26	2	7.7	53± 2
			M and F	60	25	0	-	-	49	3	6.1	46± 5	44	4	9.1	58± 4
III	50 ppm	1 week	M	30	17	1	5.9	64	27	0	-	-	26	1	3.8	62
			F	30	14	0	-	-	28	2	7.1	39±13	27	3	11.1	80± 8
			M and F	60	31	1	3.2	64	55	2	3.6	39±13	53	4	7.5	75± 7
IV	25 ppm	52 weeks	M	30	21	3 ^e	14.3	71± 5	29	0 ^f	-	-	28	7 ^g	25.0	73± 6
			F	30	26	0	-	-	30	4 ^f	13.3	68±11	29	7 ^g	24.1	85± 6
			M and F	60	47	3	6.4	71± 5	59	4	6.8	68±11	57	14	24.6	30± 4
IV bis	25 ppm	52 weeks	M	120	98	25 ^e	25.5	75± 2	117	1 ^f	0.8	46	113	16 ^g	14.2	77± 3
			F	120	112	1	0.9	77	118	12 ^f	10.2	69± 4	118	11 ^g	9.3	78± 6
			M and F	240	210	26	12.4	75± 2	235	13	5.5	67± 4	231	27	11.7	77± 3
V	10 ppm	52 weeks	M	30	25	0	-	-	30	0 ^f	-	-	28	11 ^g	39.3	71± 5
			F	30	26	0	-	-	30	6 ^f	20.0	63± 5	30	3 ^g	16.0	68± 4
			M and F	60	51	0	-	-	60	6	10.0	63± 5	58	14	24.1	70± 4
VI	No treatment (Controls)		M	100	56	0	-	-	92	1	1.1	25	80	3	3.7	66± 7
			F	100	73	0	-	-	97	2	2.1	49± 7	92	4	4.3	56± 4
			M and F	200	129	0	-	-	189	3	1.6	41± 9	172	7	4.1	60± 4
VII	No treatment (Controls)		M	90	70	0	-	-	80	0	-	-	73	3	4.1	56±11
			F	90	85	0	-	-	88	1	1.1	83	86	2	2.3	75±12
			M and F	180	155	0	-	-	168	1	0.6	83	159	5	3.1	64± 8

(a) Alive animals when the first tumor was observed: kidney adenocarcinoma, 55 weeks; mammary tumor, 27 weeks; pulmonary adenoma, 36 weeks. The percentages refer to the corrected numbers.

(b) Average time from the start of the experiment to the detection (at the periodic control or at autopsy).

(c) All mammary tumors in females were histologically diagnosed as carcinomas.

(d) Some pulmonary adenomas were cellular atypias.

(e) $P < 0.01$, combined 25 ppm (28/119) males vs. combined control males (0/196) by chi-square test. Based on corrected numbers.

(f) $P < 0.01$ Combined control females (3/185) vs. 10 ppm females (6/30) and vs. combined 25 ppm females (16/148). Based on corrected numbers.

(g) $P < 0.01$. Combined control males (6/153) vs. 10 ppm males (11/28) and vs. combined 25 ppm males (29/294). Also, combined control females (6/178) vs. 10 ppm females (3/30) and vs. combined 25 ppm females (18/147).

a similar effect in the kidneys of mice which developed kidney adenocarcinomas in the lifetime inhalation study.

The significant ($P < 0.01$) increase in mammary carcinomas in treated female Swiss mice would indicate evidence for the carcinogenicity of VDC in these animals. However, Maltoni et al. (1980) concluded that a direct relationship between induction of mammary tumors, as well as pulmonary adenomas, in mice and vinylidene chloride exposure remains open and needs further clarification because of reasons that include the following, as quoted from the Maltoni et al. (1980) paper:

- 1) "There are some significant higher rates of mammary and pulmonary tumors in VDC exposed groups when compared to the controls;"
- 2) "When the incidence of these tumors is adjusted for survival rate, the relevance of the difference between treated and control groups is reduced;"
- 3) "No dose-response relationship could be calculated either during the total time of the experiment or in different time intervals;" and
- 4) "Overall there was a clear fluctuation and imbalance of this trend in the different groups."

The authors (1980) stated that survival of mice exposed to 10 and 25 ppm VDC was higher than survival of the control groups. Numbers of mice surviving

at 27, 36, and 55 weeks are given in Table 10-28, herein, and Maltoni et al. (1977) observed an interim survival for female mice of: (1) 47% in the 25 ppm group (IV), 20% in the 10 ppm group (V), and 21% in the matched control group (VI) when these groups were 98 weeks old, and (2) 40% in the 25 ppm (Group IV bis) and 40% in the matched control group (VII) when these groups were 91 weeks old. Maltoni et al. (1980) presented survival data and their statistical analyses of pulmonary adenoma and mammary carcinoma data in mice to support their conclusions quoted above. Their analyses include data presented in Table 10-29 which were obtained from an appendix in the Maltoni et al. (1980) report. The authors noted that "with one exception" slightly higher mean survival times for the treatment groups were not significantly ($P < 0.05$) different from those of their matched controls by a rank test of Krauth. The significant ($P < 0.05$) differences shown in Table 10-29 for survival include comparisons of Groups IV and V with Group VII, which was not the matched control group for these two treatment groups, and Group IV bis with Group IV, which were the two groups exposed to 25 ppm VDC; hence, the authors (1980) indicated that "with one exception" significant differences in mean survival were found only between groups which began their exposure at different ages.

Statistically significant ($P < 0.01$) differences in tumor incidences between control and treated mice shown in Table 10-28 were calculated using the chi-square test by combining control and 25 ppm groups since, outside of differences in age at the start of the study and the two week difference in the starting times of the study (Table 10-24), similar tumor incidences for the separate control and 25 ppm groups were found with the same protocol. Maltoni et al. (1980) found statistically significant ($P < 0.05$) increases in

TABLE 10-29. STATISTICAL ANALYSES OF SURVIVAL, MAMMARY CARCINOMA INCIDENCE AND PULMONARY ADENOMA INCIDENCE FOR MALE AND FEMALE SWISS MICE IN A CARCINOGENICITY STUDY OF VINYLIDENE CHLORIDE (MALTONI ET AL. 1980)

Mean and Standard Deviation (in weeks) of lifetime
(Test of Krauth, one-sided)

Group ^a	Male			Female		
	Animals at start	Mean	Lifetime Stand. Dev.	Animals at start	Mean	Lifetime Stand. Dev.
VII	90	70.7 C	22.58	90	81.0 C	18.51
VI	100	75.4 C	22.08	100	83.1 C	20.32
V	30	83.7 x *	15.77	30	87.5 *	12.97
IV bis	120	75.0	15.85	120	84.3	17.41
IV	30	79.6 *	17.03	30	90.0 x *	18.40

Tumor incidence for MA and PA at the end of experiment; N.TU (respectively N.MA, N.PA) is the number of animals with tumors (respectively MA, PA) (Exact Fisher Test, one-sided)

Group	Male				Female			
	Animals at start	N.TU ⁶	N.MA ^c	N.PA ^d	Animals at start	N.TU	N.MA	N.PA
VI	100	7	0	3	100	15	2	4
V	30	11 x	0	11 x	30	15 x	6 x	3
IV	30	11 x	0	7 x	30	17 x	4 *	7 x
VII	90	7	0	3	90	12	1	2
IV bis	120	43 x	1	16 x	120	40 x	12 x	11 *

Tumor incidence for MA and PA at the end of experiment; N.MA (respectively N.PA) is the number of animals with MA (respectively PA) (Logrank test)

Group	Male			Female		
	Animals at start	N.MA	N.PA	Animals at start	N.MA	N.PA
VI	100	0	3	100	2	4
V	30	0	11 x	30	6 x	3
IV	30	0	7 x	30	4	7 *
VII	90	0	3	90	1	2
IV bis	120	1	16*	120	12 *	11

^aVII - Matched control for Group IV bis, VI - Matched control for Groups V and IV, V - 10 ppm VDC exposure, IV bis and IV - 25 ppm VDC exposure.

^ccontrol

* $P \leq 0.05$

^x $P < 0.01$

^bN.TU -- Number of animals with tumors

^cN.MA -- Number of animals with mammary tumors

^dN.PA -- Number of animals with pulmonary tumors

mammary carcinoma and pulmonary adenoma incidences in treated mice compared to matched control mice by the Fisher exact test (Table 10-29); however, their analysis of these data by the Log Rank test, to take into account survival patterns, indicated a lower level of statistical significance for these differences between control and treated mice. A dose-related increase in tumor incidence in treated mice was not clearly apparent from the data in Table 10-29.

In summary, the studies by Maltoni et al. (1977, 1980) show significant ($P < 0.01$) increases in the incidence of kidney adenocarcinomas in male Swiss mice and mammary carcinomas in female Swiss mice chronically exposed to VDC. However, the authors concluded that the relationship between VDC exposure and mammary carcinoma induction in female mice remains open as discussed above. There was a significant ($P < 0.05$) increase in the incidence of fibromas and fibroadenomas in each treated group of female Sprague-Dawley rats compared to controls in the inhalation study; however, there were no significant ($P < 0.05$) increases in mammary carcinoma incidence between treated and control rats, and there was no clear dose-response for induction of mammary tumor in treated rats.

A broader evaluation of the carcinogenicity of VDC by Maltoni et al. (1977, 1980) might have been possible if exposure periods longer than 12 months had been used and if, at the beginning of treatment, younger animals, such as weanlings, had been evaluated to cover the portion of their lifespans during growth to adulthood. It does not appear that doses as high as those maximally tolerated were used in the inhalation study in hamsters and the gavage study in rats, in that neither toxicologic effects from VDC treatment nor an attempt to select a maximally tolerated dose is indicated in the report on these studies.

Viola and Caputo (1977) exposed male and female Wistar strain rats initially 2 months old in an inhalation chamber for 4 hours per day, 5 days per week, to VDC at 200 ppm (99.8% pure) for 5 months, followed by an additional 7 months of reduced exposure to VDC at 100 ppm to avoid toxicity. All animals were observed until death (22 to 24 months). Thirty untreated controls of each sex, 23 treated females, and 51 treated males were examined for tumors. During the exposure period, a rapidly growing mass appeared in the external ear duct of many of the animals, but on biopsy examination, only signs of an inflammatory reaction were present with no indications of dermal or epidermal neoplasia. Tumors of the abdominal cavity were present at the same incidence in both experimental (23%) and control (25%) rats. Although histology of the tumors was described, there was no mention of which organs, if any, were routinely evaluated histologically for neoplastic or pre-neoplastic changes. Furthermore, the 12-month exposure period was shorter than potential lifetime exposures for laboratory rats. In a second study using male and female Sprague-Dawley rats exposed to VDC at 100 and 75 ppm (started in the summer of 1975) for an unspecified duration, there was again no increased tumor incidence in the experimental animals as compared with the controls. There were 30 rats in each control and 100 ppm group, and 21 females and 16 males in the 75 ppm group examined for tumors. At the time of the report, only gross tumors were enumerated; results from microscopic examination of the tissues from the Sprague-Dawley rats were to be reported later.

Lee et al. (1978), in a study primarily concerned with chronic inhalation exposure of CD-1 mice and CD rats (Charles River), initially about 2 months old, to vinyl chloride (99.8% pure), exposed a group of rats and mice (36

males and 36 females) to a single 55 ppm concentration of VDC. Concurrent control groups exposed to inhalation of air alone consisted of 36 males and 36 females. The exposure was for 6 hours daily, 5 days per week for 12 months, at which time all surviving animals were terminated. Following sacrifice, the brain, pituitary, thyroid, respiratory tract, alimentary canal, urogenital organs, thymus, heart, liver, pancreas, spleen, mesenteric lymph nodes, and other organs with gross pathological lesions were histologically examined. The incidence of tumors observed in both rats and mice is given in Table 10-30. There was no statistically significant ($P < 0.05$, Fisher exact test) increase in tumors at any of the sites examined as compared with control animals. The short duration (12 months) of this study may have precluded observing tumors that have a long latency period.

In a follow-up study, Lee and coworkers (Hong et al., 1981) exposed 8 to 12 CD-1 mice of each sex and 4-16 CD rats of each sex, initially 2 months old, to VDC vapors (99% pure) at a concentration of 55 ppm. The mice were exposed 6 hours/day, 5 days/week for a period of 1, 3, or 6 months (12 mice of each sex were treated for 6 months), while the rats were exposed on the same schedule for 1, 3, 6, or 10 months (16 rats of each sex were treated for 10 months). Concurrent controls exposed to inhalation of air alone consisted of 28 mice and 16 rats of each sex. All animals were observed for an additional period of 12 months following exposure. Histologic examinations were performed as described above in Lee et al. (1978). Focusing on males and females, respectively, in the groups with the longest durations of exposure, 18 and 21% of control mice, 50 and 42% of treated mice, 38 and 44% of control rats, and 79 and 56% of treated rats died before terminal sacrifice. Although tumors were observed in some exposed and control animals, the incidence of

TABLE 10-30. TUMOR INCIDENCE IN RATS AND MICE EXPOSED TO VINYLIDENE CHLORIDE

Species	Sex	Tumor type	No. tumors/No. ^a exposed animals (55 ppm vinylidene chloride)	No. tumors/No. control animals
CD-1 mice	M	Bronchioloalveolar adenoma	6/35	1/26
CD-1 mice	F	Bronchioloalveolar adenoma	0/35	0/36
CD-1 mice	M	Hemangiosarcoma of the liver	2/35	0/26
CD-1 mice	F	Hemangiosarcoma of the liver	1/35	0/36
CD-1 mice	M	Heptoma	2/35	0/26
CD-1 mice	F	Heptoma	1/35	0/36
CD rats	M	Hemangiosarcoma of any organ except liver	2/36	0/35
CD rats	F	Hemangiosarcoma of any organ except liver	0/35	0/35

^a None of the tumor increases were statistically significant when compared to controls.

Source: Lee et al., 1978.

these tumors was not significantly ($P < 0.05$) increased over control levels (Table 10-31). The small number of animals in each group weakens the ability of this study to detect a tumorigenic response, and the exposure durations were less than potential lifetime exposures. A broader evaluation could have been made with additional exposure levels in each study.

Two studies have been performed for the Manufacturing Chemists Association to assess the effects of chronic inhalation and ingestion (in drinking water) of VDC. The studies were conducted in two phases, the first for 90 days and the second for 2 years. Norris (1977) presented preliminary results of these studies with a detailed final report of the drinking water study presented by Humiston et al. (1978) and Quast et al. (1983). An additional report of these two studies to the open literature was provided by Rampy et al. (1977), and the final results of the inhalation study have been presented by McKenna et al. (1982).

In the inhalation study (Rampy et al., 1977; McKenna et al., 1982), Sprague-Dawley rats, with initial body weights of 250 to 300 g for males and 190 to 230 g for females, were exposed to VDC ($\geq 99\%$ pure) for 6 hours per day, 5 days per week for a period of 18 months, and animals that survived the treatment were observed for an additional 6 months. A total of 86 males and 86 females per group were used for each exposure group plus a concurrent untreated control group. The animals were initially exposed to VDC at 10 or 40 ppm; however, after 5 weeks of exposure, an interim kill revealed no effects of exposure, and the concentrations were raised to 25 and 75 ppm, with these concentrations being used for the remainder of the experiment. Analytical and nominal atmospheric concentrations of VDC in inhalation

TABLE 10-31. TUMOR INCIDENCE IN RATS AND MICE EXPOSED TO VINYLIDENE CHLORIDE

Species	Sex	Tumor type	No. tumors/No. exposed animals ^a (55 ppm vinylidene chloride)	No. tumors/No. control animals ^a
CD-1 mice	M	Hemangiosarcoma of the liver	0/28	0/60
CD-1 mice	F	Hemangiosarcoma of the liver	0/28	1/60
CD-1 mice	M	Hepatocellular tumor	4/28	10/60
CD-1 mice	F	Hepatocellular tumor	0/28	1/60
CD-1 mice	M	Hemangiosarcoma--various organs	1/28	0/60
CD-1 mice	M	Bronchioloalveolar tumor	4/28	8/60
CD-1 mice	F	Bronchioloalveolar tumor	1/28	8/60
CD rats	M	Hepatocellular carcinoma	0/34	1/36
CD rats	M	Hemangiosarcoma of the liver	1/34	0/36
CD rats	F	Mammary fibroadenoma	5/36	5/36
CD rats	F	Mammary adenocarcinoma/carcinoma	0/36	1/36

^aCombined incidences for all exposure periods.

Source: Hong et al., 1981.

chambers were equivalent. The overall evaluation of toxicity included survival, body weights, organ weights, hematology, urinalysis, clinical chemistry, cytogenetics, gross pathology, and histopathology.

There were no overt signs of toxicity shown by the overall evaluation in the treated groups, and a significant ($P < 0.05$) increase in mortality was observed only in female rats exposed to 25 ppm during months 15 to 23 and 75 ppm at months 15, 17, and 21 (Tables 10-32 and 10-33). However, increases in mortality in treated groups might have been influenced by histopathologically diagnosed increases in chronic murine pneumonia incidence (number of rats with pneumonia/cumulative number of animals examined) as follows: males - 0/86 control, 26/85 low dose, 24/86 high dose; females - 9/84 control, 21/86 low dose, 22/84 high dose. Although control cages were in the inhalation chamber room during exposures, control animals were not placed in the inhalation chambers. Decreased body weights in both treated groups of male rats compared to controls are indicated in Table 10-34, but the relationship to a direct effect of VDC appears rather uncertain in that a consistent slight but significant ($P < 0.05$) decrease is evident only in the low-dose group of males, no clear dose response is evident, and information on food consumption could have shown whether this could have affected the body weight trends shown. Body weights between control and treated female rats were similar (Table 10-35).

Treatment-related induction of non-neoplastic lesions was not apparent, with the possible exception of ovarian cysts in high-dose females [6/84 in controls; 8/86 in the low-dose group; 14/84, ($P < 0.05$) in the high-dose group]. An increase in midzonal fatty changes in the livers of treated female rats was

TABLE 10-32. VINYLIDENE CHLORIDE: A CHRONIC INHALATION TOXICITY AND ONCOGENICITY STUDY IN RATS

CUMULATIVE PERCENT MORTALITY FOR MALE RATS

Months on Study	Exposure level		
	Control number dead (% dead)	25 ppm number dead (% dead)	75 ppm number dead (% dead)
Number of rats alive on day 0 ^a	86	85	86
1	1 (1)	3 (4)	1 (1)
2	1 (1)	3 (4)	1 (1)
3	1 (1)	3 (4)	2 (2)
4	1 (1)	4 (5)	3 (4)
5	1 (1)	4 (5)	3 (4)
6	3 (4)	4 (5)	4 (5)
7	4 (5)	4 (5)	4 (5)
8	5 (6)	5 (6)	5 (6)
9	5 (6)	5 (6)	5 (6)
10	5 (6)	6 (7)	7 (8)
11	6 (7)	6 (7)	8 (9)
12	8 (9)	7 (8)	8 (9)
13	9 (11)	8 (9)	9 (11)
14	10 (12)	8 (9)	10 (12)
15	14 (16)	8 (9)	12 (14)
16	19 (22)	17 (20)	12 (14)
17	23 (27)	18 (21)	16 (19)
18	27 (31)	25 (29)	27 (31)
19	32 (37)	36 (42)	37 (43)
20	39 (45)	47 (55)	47 (55)
21	49 (57)	50 (59)	56 (65)
22	54 (63)	58 (68)	66* (77)
23	63 (73)	65 (77)	71 (83)
24	72 (84)	72 (85)	74 (86)
25	73 (85)	72 (85)	78 (91)
Terminals Kill	13	13	8
30-Day Interim Kill	4	4	4
6-Month Interim Kill	5	5	5
12-Month Interim Kill	5	5	5
26 Weeks Cytogenetic Kill ^b	4	3	4
Total Rats in Study	104	103 ^c	104

^a Excludes those rats in the interim kills and the cytogenetic kill.

^b Four rats per exposure level were added one month after start of study.

^c Includes one rat designated for cytogenetics study but not used due to death six weeks prior to cytogenetic kill.

* Statistically different from control data when analyzed using Fisher's Exact Probability test, $P < 0.05$.

Source: McKenna et al., 1982.

TABLE 10-33. VINYLIDENE CHLORIDE: A CHRONIC INHALATION TOXICITY AND ONCOGENICITY STUDY IN RATS

CUMULATIVE PERCENT MORTALITY FOR FEMALE RATS

Months on study	Exposure level					
	Control		25 ppm		75 ppm	
Number of rats alive on day 0 ^a	number dead (% dead)		number dead (% dead)		number dead (% dead)	
	84		86		84	
1	0	(0)	0	(0)	0	(0)
2	0	(0)	0	(0)	0	(0)
3	0	(0)	0	(0)	0	(0)
4	1	(1)	0	(0)	0	(0)
5	1	(1)	0	(0)	0	(0)
6	1	(1)	1	(1)	1	(1)
7	1	(1)	1	(1)	1	(1)
8	1	(1)	1	(1)	1	(1)
9	1	(1)	3	(4)	1	(1)
10	1	(1)	3	(4)	1	(1)
11	2	(2)	6	(7)	1	(1)
12	2	(2)	6	(7)	5	(6)
13	4	(5)	9	(11)	6	(7)
14	4	(5)	10	(12)	7	(8)
15	5	(6)	18*	(21)	13*	(16)
16	8	(10)	19*	(22)	16	(19)
17	9	(11)	19*	(22)	18*	(21)
18	16	(19)	29*	(34)	20	(24)
19	23	(27)	37*	(43)	29	(35)
20	30	(36)	49*	(57)	41	(49)
21	39	(46)	55*	(64)	51*	(61)
22	46	(55)	59*	(69)	54	(64)
23	56	(67)	65	(76)	61	(73)
24	64	(76)	72	(84)	68	(81)
25	65	(77)	75	(87)	68	(81)
Terminal Kill	19		11		16	
30-Day Interim Kill	4		4		4	
6-Month Interim Kill	5		5		5	
12-Month Interim Kill	5		5		5	
26 Weeks Cytogenetic Kill ^b	4		4		4	
Total Rats in Study	102		104		102	

^a Excludes those rats in the interim kills and the cytogenetic kill.

^b Four rats per exposure level were added one month after start of study.

* Statistically different from control data when analyzed using Fisher's Exact Probability test, $P < 0.05$.

Source: McKenna et al., 1982.

TABLE 10-34. VINYLIDENE CHLORIDE: A CHRONIC INHALATION TOXICITY
AND ONCOGENICITY STUDY IN RATS

MEAN BODY WEIGHT FOR MALE RATS

Study Time		Controls	15 ppm	75 ppm
Month	Day			
0	0	275.82	276.05	282.56*
	8	319.74	305.84*	328.20*
	13	346.77	340.06*	353.56*
	20	378.09	363.06*	375.61
1	26	399.47	385.93*	396.02
	39	428.59	412.48*	423.88
2	52	448.09	431.80*	447.24
3	79	489.09	468.71*	488.15
4	113	518.47	500.00*	518.94
5	140	536.55	512.86*	539.54
6	174	561.78	528.74*	549.27
	201	579.24	528.15*	556.47*
	209	572.32	528.50*	552.79*
	212	601.29	563.12*	557.76*
7	219	565.01	535.02*	558.60
8	233	579.07	550.81*	559.31*
9	260	598.12	555.11*	565.36*
10	287	603.05	569.36*	572.19*
11	321	615.80	586.04*	588.57*
12	348	639.17	593.14*	586.36*
13	375	637.45	611.82*	625.81
14	409	638.81	613.47	639.74
15	436	638.46	623.26	645.81
16	463	631.71	618.00	635.24
17	497	659.41	614.94*	628.62*
18	524	627.18	585.60*	618.39
19	551	606.18	596.84	583.75
20	578	609.37	575.20	585.84
21	612	607.78	582.11	563.10*
22	646	597.50	569.04	541.85
23	677	574.96	554.65	536.13
24	704	587.27	549.69	529.55

* Significantly different from control mean, $p < .05$.

Source: McKenna et al., 1982.

TABLE 10-35. VINYLIDENE CHLORIDE: A CHRONIC INHALATION TOXICITY
AND ONCOGENICITY STUDY IN RATS

MEAN BODY WEIGHT FOR FEMALE RATS

Study Time		Controls	25 ppm	75 ppm
Month	Day			
0	0	210.24	212.58	203.22*
	8	230.20	221.38*	241.57*
	13	238.48	243.40*	247.40*
	20	252.87	246.59*	255.68
1	26	259.52	258.60	264.34
	39	269.89	277.79*	280.79*
2	52	278.78	280.84	289.26*
3	79	294.14	301.51*	308.79*
4	113	311.28	317.23	325.10*
5	140	322.03	329.17	335.96*
6	174	332.67	342.63*	342.12*
	201	336.18	331.11	339.93
	209	337.11	334.58	336.71
	212	332.63	335.27	343.73*
7	219	331.73	335.80	340.29
8	233	338.73	338.93	340.23
9	260	339.60	348.01	348.26
10	287	343.67	362.06*	356.57
11	321	356.89	370.98*	368.90*
12	348	367.94	377.14	363.14
13	375	376.74	375.25	387.52
14	409	379.60	386.23	397.57*
15	436	386.38	391.36	402.62
16	463	390.53	397.31	397.10
17	497	406.30	411.39	419.64
18	524	396.26	419.26	426.13
19	551	395.10	429.48	421.21
20	578	414.86	405.68	413.13
21	612	443.06	411.72	448.65
22	646	412.23	435.19	457.27
23	677	420.33	460.71	469.91
24	704	427.00	462.67	438.62

* Significantly different from control mean, $p < .05$.

Source: McKenna et al., 1982.

evident [1/84 in controls; 6/86, ($P>0.05$) in the low-dose group; 8/84, ($P<0.05$) in the high-dose group] which may be treatment-related. However, these changes were mainly evident in females sacrificed at the end of exposure at 8 months, and only 1 high-dose female had this midzonal fatty change in liver during the 6-month post-exposure period. Furthermore, the total numbers of females with any type of fatty change in liver were as follows: 15/84 controls, 13/86 low-dose rats, and 16/84 high-dose rats.

VDC exposure did not result in significant ($P<0.05$) increases in tumor incidence in this study, except for the mammary tumor data discussed below. Mammary tumor data are presented in Table 10-36 for comparison with mammary tumor data on female Sprague-Dawley rats exposed to VDC in the inhalation study by Maltoni et al. (1980; Table 10-27). The incidence of adenocarcinoma without metastasis was significantly ($P<0.05$) increased in low-dose females compared to controls; however, mammary adenocarcinoma incidence was not significantly ($P<0.05$) increased in high-dose females, and adenocarcinoma without metastasis was found in another control female, which causes the adenocarcinoma incidence in low-dose females to lose significance ($P>0.05$). The significant increase ($P<0.05$) in total mammary tumor incidence in high-dose females compared to control females includes animals with both benign and malignant tumors and according to the authors, total mammary tumor incidence in the control group and in each treatment group is within the historical control range for the Sprague-Dawley strain of rats at the investigating laboratory. Thus, demonstration of an induction of mammary tumor formation in female Sprague-Dawley rats under the conditions of the study by McKenna et al. (1982) does not seem apparent. Overt toxicity of VDC in Sprague-Dawley rats does not appear evident, and an attempt to use higher

TABLE 10-36. VINYLIDENE CHLORIDE: A CHRONIC INHALATION TOXICITY AND ONCOGENICITY STUDY IN RATS

HISTOPATHOLOGIC DIAGNOSIS AND NUMBER OF TUMORS IN FEMALE RATS

		0-6 Months	7-12 Months	13-18 Months	19-24 Months	Terminal Kill	Cumulative Results
Number of rats necropsied during the time period indicated	Control	1	1	14	49	19	85
	25 ppm	1	5	23	46	11	86
	75 ppm	1	4	15	48	16	85
MAMMARY GLAND							
Fibroadenoma/adenofibroma S-25	Control	0	0	17/10	78/39	37/15	132/64 ^b
	25 ppm	0	2/2	40/20	76/38	16/8	134/68 ^b
	75 ppm	1/1	6/4	17/14	100/40	33/15	157/74 ^b
Fibroma S-26	Control	0	0	0	0	0	0
	25 ppm	0	0	0	0	0	0
	75 ppm	0	0	0	1/1	1/1	2/2 ^b
Adenocarcinoma with pulmonary metastasis S-40	Control	0	0	0	1/1	0	1/1 ^b
	25 ppm	0	0	0	0	0	0
	75 ppm	0	0	0	0	0	0
Adenocarcinoma without metastasis S-27	Control	0	0	0	1/1	0	1/1 ^b
	25 ppm	0	1/1	2/2	1/1	4/3	8/7 ^{a,b}
	75 ppm	0	0	2/2	0	2/2	4/4 ^b
Leiomyosarcoma without metastasis S-41	Control	0	0	0	0	0	0
	25 ppm	0	0	0	1/1	0	1/1 ^b
	75 ppm	0	0	0	0	0	0
Carcinosarcoma of the mammary gland S-43	Control	0	0	0	0	1/1	1/1 ^b
	25 ppm	0	0	0	0	0	0
	75 ppm	0	0	0	0	0	0
Adenofibroma based on gross examination only S-44	Control	0	0	0	0	0	0
	25 ppm	0	0	0	0	0	0
	75 ppm	0	0	0	1/1	0	1/1 ^b
Total number of animals with a subcutaneous and/or mammary gland tumor ^c S-02, S-05, S-25, S-26, S-27, S-40, S-41, S-43, S-44	Control	0	0	10	39	16	65
	25 ppm	0	3	20	38	9	70
	75 ppm	1	4	14	41	15	75 ^a
Number of rats with a malignant tumor in mammary region or skin ^c S-05, S-27, S-40, S-41, S-43	Control	0	0	0	2	2	4
	25 ppm	0	1	2	2	3	8
	75 ppm	0	0	2	0	2	4

^aSignificantly different from control data when analyzed using Fisher's Exact Probability Test, $P < 0.05$.^bNumber of tumors/number of animals with tumors.^cA subcutaneous fibroma was found in one female in the 75 ppm group, and a subcutaneous fibrosarcoma without metastasis was observed in one female in the control group.

Source: McKenna et al., 1982.

exposure levels in this study might have provided a broader evaluation for carcinogenicity.

In the reports of Quast et al. (1983) and Humiston et al. (1978), male and female Sprague-Dawley rats, initially 6 to 7 weeks old, were exposed to VDC at 50, 100, and 200 ppm (99.5% pure) in drinking water for two years. Because of the volatile nature of VDC, the water was made up at higher concentrations, and the above figures reflect the average concentrations during 24-hour periods. The actual VDC levels measured in the drinking water were 68 ± 21 , 99 ± 22 , and 206 ± 33 ppm (mean \pm S.D.). When taking into account the water consumption of the rats, the time-weighted average dose was equivalent to 7, 10, or 20, and 9, 14, or 30 mg/kg/day in males and females, respectively. The VDC used in this study was distilled prior to making up the test water in order to reduce to 1 to 5 ppm the concentration of the inhibitor monomethyl ether of hydroquinone (this inhibitor would normally be at low levels in copolymers used for food packaging applications). Each treatment group consisted of 48 rats of each sex, and an untreated control group consisted of 80 males and 80 females. Animals were evaluated for survival, body weight, food and water consumption, hematology, clinical chemistry, urinalysis, gross pathology, and histopathology. The animals were observed until morbidity, or 24 months, at which time surviving animals were sacrificed. The specific tissues examined are presented in Table 10-37. There was no evidence of statistically significant ($P < 0.05$) treatment-related effects of VDC on the toxicologic endpoints evaluated in treated animals as compared to controls. At least 50% of each group survived for approximately as long as 20 months.

TABLE 10-37. REPRESENTATIVE TISSUE SPECIMENS OBTAINED AT NECROPSY
FROM ALL ANIMALS

esophagus	peripheral nerve
salivary glands	trachea
stomach	lungs (bronchi)
large intestine	nasal turbinates
pancreas	sternum and sternal bone marrow ^b
liver	spleen
kidneys	mediastinal lymphoid tissue ^a
urinary bladder	(thymus, mediastinal lymph nodes)
prostate	lymph nodes (mesenteric)
accessory sex glands	heart
epididymides	aorta
testes	skeletal muscle
ovaries	adrenal glands
uterus	thyroid gland
mammary tissue (females)	parathyroid gland ^a
brain (cerebrum, cerebellum with brain stem)	adipose tissue
pituitary gland	skin
spinal cord	any gross lesion or mass
	eyes

^aThese tissues were evaluated histologically only to the extent that they were included in the routine sections of adjacent larger organs.

^bSternum and enclosed bone marrow were to be evaluated histologically only if indicated by abnormal findings in the hematological studies.

^cEyes were saved in formalin if the rat died or was culled from the study. If the rat was from the terminal kill, a portion of these were fixed in Zenker's fixative and the remainder in formalin. Only selected animals had their eyes microscopically examined.

Source: Humiston et al., 1978

A number of neoplastic lesions were observed in both control and experimental animals. The total tumor incidence is given in Table 10-38. The tumor frequency at specific sites, as reported by Quast et al. (1983), did not show a treatment-related effect of VDC. The only statistically ($P < 0.05$) significant increase in a specific neoplasm was in female rats exposed to VDC at 50 ppm. These animals had an increased incidence of mammary gland fibroadenomas/adenofibromas (40/48 50 ppm females vs. 53/80 control females). This tumor was not considered by the authors to be treatment-related, since there was no dose relationship between tumor incidence, i.e., 36/48 100 ppm females and 35/48 200 ppm females, and exposure to VDC. Also, the spontaneous incidence of mammary tumors was high in this strain of rats, and the tumor incidence observed in the 50 ppm females was within the limits for historical controls at the testing laboratory. Four control, 1 low-dose, and 2 mid-dose females had mammary carcinomas. No treatment-related effects of VDC on non-neoplastic lesions were evident, except for minimal fatty change and swelling in the livers of high-dose males and females in all treatment groups. The dose of VDC ingested in the drinking water appeared low enough to produce no overt toxic effects. The use of higher exposure levels might have provided a stronger evaluation of the carcinogenic potential of VDC; however, the authors indicated that 200 ppm, the maximum concentration of VDC in water used in this study, was the highest concentration possible given the solubility of VDC (2.25 g/l at 25°C).

The National Cancer Institute/National Toxicology Program (NCI/NTP, 1982) has prepared a report on a cancer bioassay of VDC (99% pure) performed on Fischer 344 rats and B6C3F1 mice, initially 9 weeks old. Male and female rats (50 animals of each sex) received VDC, dissolved in corn oil, by gavage at

TABLE 10-38. TUMOR INCIDENCE FOLLOWING INGESTION OF VDC

Concentration of VDC water (ppm)	Total number of neoplasms		Total number of rats in group		Average number of neoplasms/number of rats	
	Male	Female	Male	Female	Male	Female
0	100	187	80	80	1.3	2.3
50	53	132	48	48	1.1	2.8
100	43	120	48	48	0.9	2.5
200	53	124	47	48	1.1	2.6

Source: Humiston et al., 1978.

doses of 1 and 5 mg/kg, while male and female mice (50 animals of each sex) received doses of 2 and 10 mg/kg. The animals were treated 5 days/week for a total of 104 weeks. Animals were observed for gross signs of toxicity during the exposure period, as indicated by food consumption patterns, body weight gain, and mortality. The treated and control animals were killed after 104 weeks of exposure or when moribund, and the tissues indicated in Table 10-39 were examined histologically for both tumorigenic and non-tumorigenic pathology.

The survival (greater than 50% for the whole study for all groups) and weight gain of rats were unaffected by treatment with VDC; however, 12 control and 10 low-dose male animals were accidentally killed as a result of a 5-hour exposure to 37°C on week 82 of the study. Including the male rats accidentally killed, survival of rats at terminal sacrifice was as follows: 20/50 vehicle control males, 24/50 low dose males, 37/50 high dose males, 27/50 vehicle control females, 28/50 low dose females, and 29/50 high dose females. In mice, only survival in the low-dose female group was adversely affected by treatment. However, at least 64% of the mice in all groups survived the study (33/50 vehicle control males, 35/50 low dose males, 36/50 high dose males,

TABLE 10-39. TISSUES EXAMINED^a FOR HISTOLOGIC CHANGES IN THE NCI BIOASSAY

skin	liver
lungs and bronchi	pancreas
trachea	stomach
bone and bone marrow	small intestine
spleen	large intestine
lymph nodes	kidney
heart	urinary bladder
salivary gland	pituitary
adrenal	mammary gland
thyroid	prostate
parathyroid	brain
seminal vesicles (male)	uterus (female)
testis (male)	ovary (female)
thymus	larynx
esophagus	

^aThese tissues were examined in all animals, except where advanced autolysis or cannibalism prevented meaningful evaluation.

Source: NCI/NTP, 1981.

40/50 vehicle control females, 32/50 low dose females, and 42/50 high dose females). Body weight gain was slightly depressed in both dose groups of male mice and in female mice of the low-dose group as compared to either the corresponding high-dose or control animals (Table 10-40); however, a dose-response was not evident, and body weight decreases were greater in low-dose than in high-dose mice. Dose-related non-neoplastic lesions consisted of liver necrosis in mice (male controls, 1/46, 2%; low-dose, 3/46, 7%; high-dose, 7/49, 14%; female controls, 0/47, 0%; low-dose, 4/49, 8%; high-dose, 1/49, 2%) and inflammation of the kidney in rats (male controls, 26/49, 53%; low-dose, 24/48, 50%; high-dose, 43/48, 90%; female controls, 3/49, 6%; low-dose, 6/49, 12%; high-dose, 9/49, 20%). It is noted in the NTP (1982) report that chronic nephritis is a common lesion in aging rats. Other histopathologic changes of a non-tumorigenic type were observed randomly in control and treated animals, and were not considered treatment-related.

TABLE 10-40. MEAN BODY WEIGHT CHANGE (RELATIVE TO CONTROLS) OF MICE
ADMINISTERED VINYLIDENE CHLORIDE BY GAVAGE

		Mean body weight change grams			Weight change rela- tive to controls (a) (percent)	
Week No.		Control	Low dose	High dose	Low dose	High dose
Male mice	0	28 ^(b)	24 ^(b)	25 ^(b)		
	1	1	2	2	+100	+100
	20	13	11	13	-15	0
	40	19	17	19	-20	0
	60	22	21	23	-5	+5
	80	21	20	22	-5	+5
	100	20	21	21	+5	+5
Female mice	0	19 ^(b)	18 ^(b)	18 ^(b)		
	1	2	2	3	0	+50
	20	9	8	9	-11	0
	40	12	12	13	0	+8
	60	17	15	18	-12	+6
	80	19	18	14	-5	+35
	100	24	21	24	-13	0

(a) Weight change relative to controls =

$$\frac{\text{Weight change (dosed group)} - \text{Weight change (control group)}}{\text{Weight change (control group)}} \times 100$$

^bInitial weight

Source: NCI/NTP, 1981.

In rats and mice, increased tumor incidences were observed in a number of organs (Table 10-41). While unadjusted analyses of these data suggested that some responses were marginally statistically significant at the $P = .05$ level, such analyses do not take into account the multiple organs compared. As indicated by NCI/NTP (1980) and discussed in greater detail by Gart et al. (1979), it is more accurate to account for multiplicity of comparisons by the Bonferroni correction when using the Fisher exact test, and to account for differing patterns of mortality among the dose groups by time-adjusted analyses. Making these corrections, the only statistically significant ($P < 0.05$) increased response was for lymphomas in the low-dose group female mice compared to the matched controls using the Fisher exact test. Furthermore, only lymphomas represent a statistical increase in malignant tumors in Table 10-41. However, a similar increase in the high-dose group was not noted, and the Cochran-Armitage test for linear trends was not significant. Since the matched controls had a low incidence (4%) of lymphomas when compared to the incidence (9.8%) in historical controls, this effect was not considered to be treatment-related. Thus, the conclusion in the NTP (1981) report is that VDC was not carcinogenic to F344 rats or B6C3F1 mice under the conditions of this assay. Survival and body weight data suggest that higher doses of VDC could have been given to rats and mice to more strongly challenge these animals for carcinogenicity.

VDC was tested for its potential as a brain carcinogen on the basis that vinyl chloride causes neuroblastomas in animal studies and is associated with brain neoplasms in exposed humans (Maltoni et al., 1982). No increase in the incidence of gliomas, meningiomas, or ependymomas was reported among Sprague-Dawley rats exposed by inhalation to VDC at 10, 25, 50, 100, or 150 ppm for

TABLE 10-41. TUMORS WITH INCREASED INCIDENCE IN RATS AND MICE, AS INDICATED BY THE FISHER EXACT TEST OR THE COCHRAN-ARMITAGE TEST FOR LINEAR TREND

Species	Tumor	Control	Incidence	
			Low Dose	High Dose
rat (male)	adrenal, pheochromocytomas	6/50 (12%) (P=0.01)*	5/48 (10%)	13/47 (28%) (P=0.045)**
rat (male)	pancreatic islet-cell, adenomas/carcinoma (a)	4/49 (8%) (P=0.025)*	1/47 (2%)	8/48 (17%)
rat (male)	testes, interstitial-cell	43/50 (86%) (P=0.013)*	39/47 (83%)	47/48 (98%) (P=0.034)**
rat (female)	pituitary, adenomas	16/48 (33%) (P=0.017)	20/49 (41%)	24/43 (56%) (P=0.026)**
rat (male)	subcutaneous, fibromas	0/50 (0%) (P=0.024)*	1/48 (2%)	4/48 (8%)
mice (female)	lymphomas	2/48 (4%)	9/48 (18%) (P=0.028)**	6/50 (12%)

(a) All pancreatic tumors were carcinomas, except for adenomas in two high-dose males. Carcinomas alone were not significant by the Cochran-Armitage test.

* Significant for linear trend by Cochran-Armitage test ($P < 0.05$).

**Significant by Fisher exact test ($P < 0.05$).

Source: NCI/NTP, 1982.

4 or 5 days weekly for 52 weeks. Male and female rats exposed by ingestion (gavage) to 0.5, 5, 10, or 20 mg/kg of VDC for 5 days weekly for 52-59 weeks did not show an increased incidence of brain tumors when compared to controls.

Ponomarev and Tomatis (1980) assessed the carcinogenic potential of VDC in BDIV rats exposed to the compound starting in utero and continuing to 120 weeks of age. Pregnant rats (24 animals) received a single 150 mg/kg dose of VDC (99% pure) by gavage in corn oil on day 17 of gestation. Following birth,

the pups (89 males and 90 females) received VDC by gavage at a dose of 50 mg/kg weekly for the duration of the study. The animals were killed at the termination of the study or when moribund, and major organs and gross lesions were examined histologically.

There was no treatment-related effect on the reproductive success of the dams or on the survival or weight gain of the offspring of rats treated with VDC. Non-tumorigenic lesions in the liver, consisting of degeneration of the parenchymal cells, liver necrosis, and hemorrhage, were observed in treated animals moribund after 80 weeks of treatment or killed at termination. In cases of early death (30 weeks of treatment), lung and kidney congestion were also observed in treated rats. The total tumor incidence in either the dams or the offspring was not significantly increased by treatment (Table 10-42); however, there was an increased incidence of meningiomas in male rats, although this was also not significantly different from control levels. Although slight differences in tumor incidence were observed following VDC treatment, this study did not demonstrate a statistically significant effect of VDC treatment on the induction of tumors in either the dams or their offspring. Although liver lesions were seen, it appears that higher doses could possibly have been used, since no effect was observed on survival and body weight.

Skin application has been used to assay VDC for tumor initiation and complete carcinogen action in lifetime studies using the two-stage tumorigenesis model in female Ha:ICR Swiss mice (Van Duuren et al., 1979). In two-stage tumorigenesis, an initiator agent is applied in a single dose to the skin of a mouse. The initiator does not produce tumors at the applied

Table 10-42.

Tumor incidence in female BDIV rats treated with VDC on day 17 of pregnancy,
in their progeny treated weekly for life and controls (Ponomarev and Tomatis, 1980)

Group	Effective number of rats ^a	Tumor bearing rats		Number of tumors		Animals with more than one tumor	
		n	%	Total	per rat	n	%
Females given VDC	23	11	47.8	14	0.6	3	13
Progeny treated weekly with VDC							
Males	81	31	38.3	35	0.4	4	4.9
Females	90	53	66.3	64	0.8	11	13.8
Females given olive oil	14	5	35.7	7	0.5	2	14.3
Progeny treated weekly with olive oil							
Males	49	16	32.7	16	0.3	-	-
Females	47	24	51.1	29	0.6	5	10.6

Table 10-42. (cont.)

Group	Distribution of Tumors																Liver hyperpl. nodules	
	meninges		oral cavity		stomach		liver		soft tissue		mammary gland		ovary		other			
	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Females given VDC	-	-	2	8.7	1	4.3	-	-	-	-	8	34.8	2	8.7	1 ^b	4.3	2	8.7
Progeny treated weekly with VDC																		
Males	6	7.4	5	6.2	1	1.2	1	1.2	9	11.1	1	1.2	-	-	12 ^c	14.8	2	2.5
Females	-	-	1	1.3	2	2.5	3	3.8	7	8.8	39	48.8	6	7.5	6 ^d	7.5	6	7.5
Females given olive oil	-	-	1	7.1	-	-	-	-	1	7.1	4	28.6	-	-	1 ^e	7.1	-	-
Progeny treated weekly with olive oil																		
Males	1	2.0	2	4.1	-	-	-	-	4	8.2	-	-	-	-	9 ^f	18.4	-	-
Females	-	-	1	2.1	1	2.1	-	-	-	-	22	46.8	3	6.4	2 ^g	4.3	-	-

^aThe percentages and the number of tumors per rat are expressed in relation to the effective number of rats.

^bSurvivors at the time the first tumors were observed.

^cUrinary bladder papilloma; 1 lymphoma, 4 pituitary adenomas; 3 adrenal cortical adenomas; 1 spleen haemangioma; 1 lung sarcoma, pleomorphic; 1 skin squamous cell carcinomas; 1 seminoma.

^d1 salivary gland carcinoma; 1 salivary gland adenoma; 1 lymphoma; 1 pituitary adenoma; 1 rectal adenomatous polyp; 1 uterine adenoma

^eAdrenal cortical adenoma

^f1 osteosarcoma; 1 mediastinal sarcoma; 1 lung epidermoid carcinoma; 2 lymphomas; 1 spleen haemangioma; 2 pituitary adenomas; 1 adrenal cortical adenoma

^g1 lymphoma; 1 uterine adenoma

concentration, but predisposes the skin so that later repeated applications of a promotor (an agent that by itself will not produce tumors) will cause the formation of tumors. A complete carcinogen is one which, if applied in sufficient concentrations, can produce tumors by itself. When VDC was applied 3 times each week at a dose of 121 mg/mouse to the shaved backs of 30 mice, no tumors were observed; however, when a single dose of 121 mg/mouse of VDC was applied to the skin of 30 mice, followed by repeated application of phorbol myristate acetate (PMA) as the promotor, a significant increase ($P < 0.005$) in skin papillomas (8 mice with papillomas/9 total papillomas) was observed as compared to controls. No skin papillomas were observed in 30 controls treated with acetone, the dosing vehicle, and 100 untreated controls, whereas 9 mice (10 total papillomas) of 120 mice given 0.0025 mg PMA and 6 mice (7 total papillomas) of 90 mice given 0.005 mg PMA on the same schedule as that for treated mice were found. Squamous cell carcinomas on the skin in the initiation-promotion study were observed on one mouse in the VDC group, one mouse in the low-dose PMA group, and 2 mice in the high-dose PMA group. A positive control group of 30 mice in the initiation-promotion study given 0.02 mg/mouse of 7,12-dimethylbenz[a]anthracene as the initiating agent developed papillomas in 29 mice (317 total papillomas) and local squamous cell carcinomas in 14 mice. In the same study, VDC was also tested for complete carcinogenic action in 30 female Swiss ICR/Ha mice by repeated subcutaneous injection once a week at a dose of 2 mg/mouse. Following treatment for 548 days, no tumors were observed at the site of injection or at sites distant from the site of injection. Although this study indicates that VDC is a tumor initiator in mouse skin, it is not clear how this relates to the processes of complete chemical carcinogenesis in other organs, or if tumor initiation as observed here is a phenomenon solely restricted to the skin of mice. The

relevance of positive results in this tumor initiation study with regard to the assessment of human health effects is not clear, particularly since complete carcinogenic activity could not be demonstrated. Maximally tolerated doses, estimated from preliminary short-term tests, were used in the tests for carcinogenicity performed by Van Duuren et al. (1979).

In summary, the evidence supporting the carcinogenicity of VDC is limited. It has been shown that VDC is a mutagen in bacterial assay systems, and Van Duuren et al. (1979) have demonstrated that VDC acts as a tumor initiator in mouse skin; however, of the three animal species used to assess the carcinogenicity of this compound--rats, mice, and hamsters--only one strain of mice developed a unique tumor type (kidney carcinomas) following exposure to VDC. Maltoni (1977) and Norris (1982) suggested that the observed increase in kidney tumors may be a species- and strain-specific effect, and Maltoni et al. (1980) and Maltoni (1977) obtained experimental evidence for a direct relationship between the sensitivity to acute toxic effects of VDC and carcinogenic responses in the species and strains of rats and mice evaluated for VDC carcinogenicity in their studies. Norris (1982) reviewed some of the pharmacokinetic, toxicological, and carcinogenic data on the effects of VDC in animals, and emphasized the unique susceptibility of Swiss mice to the toxic and tumorigenic effects of this compound. A statistically significant increase in mammary carcinomas was found in female Swiss mice by Maltoni et al. (1982), but these investigators concluded that this evidence was not conclusive due to reasons discussed in their report.

There have been a number of other carcinogenicity bioassays of VDC in which no statistically significant increase in tumor incidence was observed

(Table 10-21). From these studies, it could be concluded that VDC has not been found to be carcinogenic in rats or hamsters, nor in mice when administered by the oral or dermal route. The only published inhalation studies in mice, except for the positive bioassay of Maltoni et al. (1977), used either a shorter exposure period (Hong et al., 1981) or a shorter observation period (Lee et al., 1978), although the level of exposure in both studies was higher (55 ppm, 6 hours/day) than the high-dose group (25 ppm for 4 hours/day) used by Maltoni et al. (1977). Furthermore, higher doses possibly could have been tested in several of the negative studies. Nonetheless, in view of the present evidence for VDC carcinogenicity found in one mouse strain in the inhalation study by Maltoni et al. (1980, 1977), the available evidence regarding the carcinogenicity of VDC in animals is thus seen to be limited and not sufficient for a firm conclusion on the potential carcinogenicity of this compound in humans.

10.5.2 Epidemiologic Studies

Adequate data regarding the carcinogenic potential of VDC in humans are lacking. One study, that of Ott et al. (1976) investigated 138 Dow Chemical Company workers exposed primarily to VDC, and gave mortality data and the results of health examinations. The authors noted that vinyl chloride is a common contaminant of VDC monomer and that liquid VDC contained less than 0.2% by weight of vinyl chloride in recent years prior to publication of their study. In this study, time-weighted average (TWA) exposures were estimated based on job descriptions and industrial hygiene surveys. Based on TWA exposure estimates, each job initially fell into one of four categories: <10 ppm, 10 to 24 ppm, 25 to 49 ppm, and 50+ ppm. However, since operators in

fiber production were the only workers with exposures above 25 ppm TWA, only three concentration categories were used to calculate cumulative dose: <10 ppm (5 ppm used for calculation); 10-24 ppm (17 ppm used for calculation); \geq 25 ppm (43 ppm used for calculation). Cumulative career exposures were estimated by multiplying the TWA times the duration, in months, of potential exposure (Table 10-43). As indicated in the table, the size of the population having a lengthy duration of exposure or a long latency period since the initial time of exposure was small. Moreover, it cannot be determined from the data presented in the study report whether those individuals for whom the greatest time had elapsed since initial exposure were also among the most heavily exposed subjects. Thus, it is possible that for detection of long-latency diseases such as cancer, the population examined in this study may not be adequate.

Cohort mortality was compared with U.S. white male mortality for 1942, 1947, 1952, 1957, 1962, 1967, and 1971 by the indirect method. The most recent health inventory for the cohort was compared with matched controls. For the total cohort of 138 persons, 5 deaths were recorded. One death of an individual with a 745 ppm cumulative dose was attributed to malignancy (respiratory cancer); however, his smoking history was not known. The expected respiratory cancer death rate was 0.3 for the total cohort; 0.2 for the total cohort 15+ years after first exposure to VDC; and 0.2 for the cohort with 500+ ppm months of exposure. An examination of company health inventory records on this same cohort revealed no statistically significant ($P < 0.05$) clinical difference between the VDC-exposed group and matched (for age and smoking) controls. Ott et al. (1976) mentioned that the cohort exposed to VDC was also exposed to copolymers other than vinyl chloride, and that individuals

TABLE 10-43. ESTIMATED CUMULATIVE DOSE, DURATION OF EXPOSURE AND DATE OF FIRST EXPOSURE AMONG 138 INDIVIDUALS EXPOSED TO VINYLIDENE CHLORIDE

Exposure measures	Total population
Estimated career dosage (TWA x months of exposure)	
<500 ppm months	50
500-999 ppm months	28
1000-1999 ppm months	28
2000+ ppm months	32
Duration of exposure	
<12 months	35
12-59 months	43
60-119 months	35
120+ months	25
Date of first exposure	
1940-1949	9
1950-1959	74
1960-1969	55

^aStatus as of January 1974.

Source: Ott et al., 1976.

in the matched control population might have been exposed to a number of other chemicals.

10.5.3 Quantitative Estimation

This quantitative section deals with the unit risk for VDC in air, and the potency of VDC relative to other carcinogens that the CAG has evaluated. The unit risk estimate for an air pollutant is defined as the lifetime cancer risk occurring in a hypothetical population in which all individuals are exposed continuously from birth throughout their lifetimes to a concentration of $1 \mu\text{g}/\text{m}^3$ of the agent in the air they breathe. Unit risk estimates are used

for two purposes: 1) to compare the carcinogenic potency of several agents with each other, and 2) to give a crude indication of the population risk which might be associated with air or water exposure to these agents, if the actual exposures are known.

10.5.3.1 Procedures for Determination of Unit Risk

The data used for the quantitative estimation of unit risk for VDC were taken from a lifetime animal study. In animal studies it is assumed, unless evidence exists to the contrary, that if a carcinogenic response occurred at the dose levels used in the study, then responses would also occur at all lower doses, at an incidence determined by an extrapolation model. However, there is no solid scientific basis for any mathematical extrapolation model that relates carcinogen exposure to cancer risks at the extremely low concentrations that must be dealt with in evaluating environmental hazards. For practical reasons, such low levels of risk cannot be measured directly either by animal experiments or by epidemiologic studies. It is necessary, therefore, to depend on current knowledge of the mechanisms of carcinogenesis for guidance as to the correct risk model to use.

At the present time, the dominant view is that most cancer-causing agents also cause irreversible damage to DNA. This position is reflected by the fact that a very large proportion of agents that cause cancer are also mutagenic. There is reason to expect that the quantal type of biological response, which is characteristic of mutagenesis, is associated with a linear non-threshold dose-response relationship. Indeed, there is substantial evidence from mutagenicity studies with both ionizing radiation and a wide

variety of chemicals that this type of dose-response model is the appropriate one to use. This is particularly true at the lower end of the dose-response curve; at higher doses, there can be an upward curvature, probably reflecting the effects of multistage processes on the mutagenic response. The linear non-threshold dose-response relationship is also consistent with the relatively few epidemiologic studies of cancer responses to specific agents that contain enough information to make the evaluation possible (e.g., radiation-induced leukemia, breast and thyroid cancer, skin cancer induced by arsenic in drinking water, liver cancer induced by aflatoxins in the diet). There is also some evidence from animal experiments that is consistent with the linear non-threshold model (e.g., the initiation stage of the two-stage carcinogenesis model in rat liver and mouse skin).

Because its scientific basis, although limited, is the best of any of the current mathematical extrapolation models, the linear non-threshold model has been adopted as the primary basis for risk extrapolation to low levels of the dose-response relationship. The risk estimates made with this model should be regarded as conservative, representing the most plausible upper limit for the risk; i.e., the true risk is not likely to be higher than the estimate, but it could be lower.

The mathematical formulation chosen to describe the linear non-threshold dose-response relationship at low doses is the linearized multistage model. This model employs enough arbitrary constants to be able to fit almost any monotonically increasing dose-response data, and it incorporates a procedure for estimating the largest possible linear slope (in the 95% confidence limit sense) at low extrapolated doses that is consistent with the data at all dose levels of the experiment.

10.5.3.2 Description of the Low-Dose Animal Extrapolation Model

Let $P(d)$ represent the lifetime risk (probability) of cancer at dose d . The multistage model has the form

$$P(d) = 1 - \exp [-q_0 + q_1 d + q_1 d^2 + \dots + q_k d^k]$$

where

$$q_i \geq 0, \quad i = 0, 1, 2, \dots, k$$

Equivalently,

$$P_t(d) = 1 - \exp [q_1 d + q_2 d^2 + \dots + q_k d^k]$$

where

$$P_t(d) = \frac{P(d) - P(0)}{1 - P(0)}$$

is the extra risk over background rate at dose d .

The point estimate of the coefficients q_i , $i = 0, 1, 2, \dots, k$, and consequently, the extra risk function, $P_t(d)$, at any given dose d , is calculated by maximizing the likelihood function of the data.

The point estimate and the 95% upper confidence limit of the extra risk, $P_t(d)$, are calculated by using the computer program GLOBAL 79, developed by Crump and Watson (1979). At low doses, upper 95% confidence limits on the extra risk, and lower 95% confidence limits on the dose producing a given risk, are determined from a 95% upper confidence limit, q_1^* , on parameter q_1 . Whenever $q_1 > 0$, at low doses the extra risk $P_t(d)$ has approximately the form $P_t(d) = q_1^* \times d$. Therefore, $q_1^* \times d$ is a 95% upper confidence limit on the extra risk and R/q_1^* is a 95% lower confidence limit on the dose, producing an extra risk of R . Let L_0 be the maximum value of the log-likelihood function. The upper limit, q_1^* , is calculated by increasing q_1 to a value q_1^* such that

when the log-likelihood is remaximized subject to this fixed value q_1^* for the linear coefficient, the resulting maximum value of the log-likelihood L_1 satisfies the equation

$$2 (L_0 - L_1) = 2.70554$$

where 2.70554 is the cumulative 90% point of the chi-square distribution with one degree of freedom, which corresponds to a 95% upper limit (one-sided).

This approach of computing the upper confidence limit for the extra risk, $P_t(d)$, is an improvement on the Crump et al. (1977) model. The upper confidence limit for the extra risk calculated at low doses is always linear. This is conceptually consistent with the linear non-threshold concept discussed earlier. The slope, q_1^* , is taken as an upper bound of the potency of the chemical in inducing cancer at low doses. (In the section calculating the risk estimates, $P_t(d)$ will be abbreviated as P.)

In fitting the dose-response model, the number of terms in the polynomial is chosen equal to $(h-1)$, where h is the number of dose groups in the experiment, including the control group.

Whenever the multistage model does not fit the data sufficiently well, data at the highest dose is deleted and the model is refit to the rest of the data. This is continued until an acceptable fit to the data is obtained. To determine whether or not a fit is acceptable, the chi-square statistic

$$\chi^2 = \sum_{i=1}^h \frac{(X_i - N_i P_i)^2}{N_i P_i (1 - P_i)}$$

is calculated where N_i is the number of animals in the i^{th} dose group, X_i is the number of animals in the i^{th} dose group, with a tumor response, P_i is the probability of a response in the i^{th} dose group estimated by fitting the multistage model to the data, and h is the number of remaining groups. The fit is determined to be unacceptable whenever X^2 is larger than the cumulative 99% point of the chi-square distribution with f degrees of freedom, where f equals the number of dose groups minus the number of non-zero multistage coefficients.

10.5.3.3 Calculation of Human Equivalent Dosages from Animal Data

Following the suggestion of Mantel and Schneiderman (1975), we assume that mg/surface area/day is an equivalent dose between species. Since, to a close approximation, the surface area is proportional to the $2/3$ power of the weight, as would be the case for a perfect sphere, the exposure in mg/day per $2/3$ power of the weight is also considered to be equivalent exposure. In an animal experiment this equivalent dose is computed in the following manner:

Let

L_e = duration of experiment

l_e = duration of exposure

m = average dose per day in mg during administration of the agent
(i.e., during l_e), and

W = average weight of the experimental animal.

Then, the lifetime average exposure is

$$d = \frac{l_e \times m}{L_e \times W^{2/3}}$$

Inhalation

When exposure is via inhalation, the calculation of dose can be considered for two cases where 1) the carcinogenic agent is either a completely water-soluble gas or an aerosol and is absorbed proportionally to the amount of air breathed in, and 2) where the carcinogen is a poorly water-soluble gas which reaches an equilibrium between the air breathed and the body compartments. After equilibrium is reached, the rate of absorption of these agents is expected to be proportional to the metabolic rate, which in turn is proportional to the rate of oxygen consumption, which in turn is a function of surface area.

Case 1

Agents that are in the form of particulate matter or virtually completely absorbed gases, such as sulfur dioxide, can reasonably be expected to be absorbed proportionally to the breathing rate. In this case the exposure in mg/day may be expressed as

$$m = I \times v \times r,$$

where I = inhalation rate per day in m^3 , v = mg/m^3 of the agent in air, and r = the absorption fraction.

The inhalation rates, I, for various species can be calculated from the observations (FASEB, 1974) that 25 g mice breathe 34.5 liters/day and 113 g rats breathe 105 liters/day. For mice and rats of other weights, W (in kilograms), the surface area proportionality can be used to find breathing rates in m³/day, as follows:

$$\text{For mice, } I = 0.0345(W/0.025)^{2/3} \text{ m}^3/\text{day}$$

$$\text{For rats, } I = 0.105 (W/0.113)^{2/3} \text{ m}^3/\text{day}.$$

For humans, the values of 20 m³/day* is adopted as a standard breathing rate (ICRP, 1977).

The equivalent exposure in mg/W^{2/3} for these agents can be derived from the air intake data in a way analogous to the food intake data. The empirical factors for the air intake per kg per day, i = I/W, based upon the previously stated relationships, are tabulated as follows:

Species	W	i = I/W
Man	70	0.29
Rats	0.35	0.64
Mice	0.03	1.3.

Therefore, for particulates or completely absorbed gases, the equivalent exposure in mg/W^{2/3} is

$$d = \frac{m}{W^{2/3}} = \frac{Ivr}{W^{2/3}} = \frac{iWvr}{W^{2/3}} = iW^{1/3}vr$$

*From "Recommendation of the International Commission on Radiological Protection, p. 3⁹. The average breathing rate is 10⁷ cm³ per 8-hr. workday and 2 x 20⁷ cm³ in 24 hrs.

In the absence of experimental information or a sound theoretical argument to the contrary, the fraction absorbed, r , is assumed to be the same for all species.

Case 2

The dose in mg/day of partially soluble vapors is proportional to the O_2 consumption, which in turn is proportional to $W^{2/3}$ and is also proportional to the solubility of the gas in body fluids, which can be expressed as an absorption coefficient, r , for the gas. Therefore, expressing the O_2 consumption as $O_2 = k W^{2/3}$, where k is a constant independent of species, it follows that

$$m = k W^{2/3} \times v \times r$$

or

$$d = \frac{m}{W^{2/3}} = kvr$$

As with Case 1, in the absence of experimental information or a sound theoretical argument to the contrary, the absorption fraction, r , is assumed to be the same for all species. Therefore, for these substances a certain concentration in ppm or $\mu\text{g}/\text{m}^3$ in experimental animals is equivalent to the same concentration in humans. This is supported by the observation that the minimum alveolar concentration necessary to produce a given "stage" of anesthesia is similar in man or animals (Dripps et al. 1977). When the animals are exposed via the oral route and human exposure is via inhalation or vice versa, the assumption is made, unless there is pharmacokinetic evidence to the contrary, that absorption is equal by either exposure route.

10.5.3.4 Calculation of the Unit Risk from Animal Studies

The 95% upper limit risk associated with $d \text{ mg/kg}^{2/3}/\text{day}$ is obtained from GLOBAL 79, and for most cases of interest to risk assessment, can be adequately approximated by $P(d) = 1 - \exp(-q_1^* d)$. A "unit risk" in units X is simply the risk corresponding to an exposure of $X = 1$. This value is estimated simply by finding the number of $\text{mg/kg}^{2/3}/\text{day}$ corresponding to one unit of X and substituting this value into the above relationship. Thus, for example, if X is in units of $\mu\text{g}/\text{m}^3$ in the air, we have that for case 1, $d = 0.29 \times 70^{1/3} \times 10^3 \text{ mg/kg}^{2/3}/\text{day}$, and for case 2, $d = 1$, when $\mu\text{g}/\text{m}^3$ is the unit used to compute parameters in animal experiments.

If exposures are given in terms of ppm in air, the following calculation may be used:

$$1 \text{ ppm} = 1.2 \times \frac{\text{molecular weight (gas) mg/m}^3}{\text{molecular weight (air)}}$$

Note that an equivalent method of calculating unit risk would be to use mg/kg for the animal exposures, and then to increase the j th polynomial coefficient by

$$(W_h/W_a)^{j/3} \quad j = 1, 2, \dots, k$$

and use mg/kg equivalents for the unit risk values.

10.5.3.5 Interpretation of Quantitative Estimates

For several reasons, the unit risk estimate based on animal bioassays is only an approximate indication of the absolute risk in populations exposed to

known carcinogen concentrations. First, there are important species differences in uptake, metabolism, and organ distribution of carcinogens, as well as species differences in target site susceptibility, immunological responses, hormone function, dietary factors, and disease. Second, the concept of equivalent doses for humans compared to animals on a mg/surface area basis is virtually without experimental verification as far as carcinogenic response is concerned. Finally, human populations are variable with respect to genetic constitution and diet, living environment, activity patterns, and other cultural factors.

The unit risk estimate can give a rough indication of the relative potency of a given agent compared with other carcinogens. The comparative potency of different agents is more reliable when the comparison is based on studies in the same test species, strain, and sex, and by the same route of exposure, preferably inhalation.

The quantitative aspect of carcinogen risk assessment is included here because it may be of use in the regulatory decision-making process, e.g., in setting regulatory priorities, evaluating the adequacy of technology-based controls, etc. However, it should be recognized that with present technology, only imprecise estimations are possible concerning cancer risks to humans at low levels of exposure. At best, the linear extrapolation model used here provides a rough but plausible estimate of the upper limit of risk, and while the true risk is not likely to be more than the estimated risk, it could be considerably lower. The risk estimates presented in subsequent sections should not be regarded, therefore, as accurate representations of the true cancer risks even when the exposures are accurately defined. The estimates

presented may, however, be factored into regulatory decisions to the extent that the concept of upper risk limits is found to be useful.

10.5.3.6 Alternative Methodological Approaches

The methods used by the CAG for quantitative assessment are consistently conservative; that is, they tend toward high estimates of risk. The use of the linear non-threshold extrapolation model tends to contribute to this conservatism. Other extrapolation models are available that would give lower risk estimates. These alternative models have not been used by the CAG in the following analysis because the limited available data show a positive response at only one dose level. Thus only the background and linear parameters have any meaning.

10.5.3.7 Estimation of Unit Risk Using the Maltoni Inhalation Study

The only animal study showing a significant increase in tumors is the inhalation study of Maltoni et al. (1980), in which male Swiss mice developed kidney adenocarcinomas. The long-term (52 weeks) exposure data from this study are presented in Table 10-44. The two control groups were combined, as well as the two groups treated at 25 ppm, since the responses within treatment groups were not statistically different. Lacking information on early mortality and scheduled sacrifice, the number surviving to the time of the first kidney adenocarcinoma is used as the denominator.

Since VDC is only slightly soluble in water, it is considered a partially soluble vapor in which the inhalation dose is proportional to O₂ consumption.

TABLE 10-44. DATA FROM MALTONI INHALATION STUDY ON MALE SWISS MICE

Dose ppm daily for 12 months	Lifetime continuous equivalent ppm	Number of animals at start	Kidney adenocarcinomas/ number surviving 55 weeks from start
0	0	100	0/56*
0	0	90	0/70*
10	0.54	30	0/25
25	1.34	30	3/21(14.3%)**
25	1.34	120	25/98(25.5%)**

* Groups were combined since responses were not statistically different.

**These groups were combined, since responses using the same dose level were not statistically different.

Furthermore, evidence by Dallas et al. (1983) indicates that the small uncharged lipid-soluble molecule is easily absorbed across the lung membranes into the systemic circulation, and that at low doses all of the chemical presented to the animal will be retained until near-equilibrium is reached. This behavior is considered similar to that of a water-insoluble anesthetic gas. (As discussed in the methodology section on equivalent inhalation dose (Case 2), exposure with these types of compounds to concentrations in ppm or $\mu\text{g}/\text{m}^3$ is considered equivalent between animals and humans).

The lifetime continuous equivalent exposure is determined by dividing total dose by total lifetime. In this experiment, inhalation exposure was 4 hours/day, 4-5 days/week for one year. On a continuous basis, this is

$$10 \text{ ppm} \times 4/24 \times 4.5/7 \times 1/2 = 0.54.$$

The calculation of the upper confidence level of risk by the multistage model yields a slope factor of

$$q_1^* = 1.7 \times 10^{-1} \text{ (ppm)}^{-1}.$$

In terms of risk, a lifetime exposure to VDC at 1 ppm corresponds to a lifetime risk of induced cancer of

$$P = 1 - \exp(-1.7 \times 10^{-1}) = 0.16.$$

To express the unit risk in terms of $\mu\text{g}/\text{m}^3$, the following conversion is used:

$$\begin{aligned} 1 \mu\text{g}/\text{m}^3 &= 1(\mu\text{g}/\text{m}^3) \times 10^{-3} (\text{m}^3/\text{l}) / (1.2(\text{g}/\text{l}) \times \text{MW}(\text{VDC})/\text{MW}(\text{air})) \\ &= 10^{-3} / (1.2 \times 97/28.8) \mu\text{g}/\text{g} = 2.5 \times 10^{-4} \text{ ppm}. \end{aligned}$$

For a lifetime continuous exposure to 1 $\mu\text{g}/\text{m}^3$ of VDC, the corresponding estimate is

$$P = 1 - \exp(-1.7 \times 10^{-1} \times 2.5 \times 10^{-4}) = 4.2 \times 10^{-5}.$$

10.5.3.8 Relative Potency

One of the uses of the concept of unit risk is to compare the relative potency of carcinogens. To estimate the relative potency on a per mole basis, the unit risk slope factor is multiplied by the molecular weight, and the resulting number is expressed in terms of $(\text{mMol}/\text{kg}/\text{day})^{-1}$. This is called the relative potency index.

Figure (10-6) is a histogram representing the frequency distribution of potency indices of 53 chemicals evaluated by the CAG as suspect carcinogens. The actual data summarized by the histogram are presented in Table (10-45).




Figure 10-6. Histogram representing the frequency distribution of the potency indices of 53 suspect carcinogens evaluated by the Carcinogen Assessment Group.

TABLE 10-45. RELATIVE CARCINOGENIC POTENCIES AMONG 53 CHEMICALS EVALUATED
BY THE CARCINOGEN ASSESSMENT GROUP AS SUSPECT HUMAN CARCINOGENS ^{1,2,3}

Compounds	Slope (mg/kg/day) ⁻¹	Molecular Weight	Potency Index	Order of Magnitude (log ₁₀) Index
Acrylonitrile	0.24(W)	53.1	1x10 ⁺¹	+1
Aflatoxin B ₁	2924	312.3	9x10 ⁺⁵	+6
Aldrin	11.4	369.4	4x10 ⁺³	+4
Allyl Chloride	1.19x10 ⁻²	76.5	9x10 ⁻¹	0
Arsenic	15(H)	149.8	2x10 ⁺³	+3
B[a]P	11.5	252.3	3x10 ⁺³	+3
Benzene	5.2x10 ⁻² (W)	78	4x10 ⁰	+1
Benzidine	234(W)	184.2	4x10 ⁺⁴	+5
Beryllium	4.86	9	4x10 ⁺¹	+2
Cadmium	6.65(W)	112.4	7x10 ⁺²	+3
Carbon Tetrachloride	1.30x10 ⁻¹	153.8	2x10 ⁺¹	+1
Chlordane	1.61	409.8	7x10 ⁺²	+3
Chlorinated Ethanes				
1,2-dichloroethane	6.90x10 ⁻²	98.9	7x10 ⁰	+1
Hexachloroethane	1.42x10 ⁻²	236.7	3x10 ⁰	0
1,1,2,2-tetrachloroethane	0.20	167.9	3x10 ⁺¹	+1
1,1,1-trichloroethane	1.6x10 ⁻³	133.4	2x10 ⁻¹	-1
1,1,2-trichloroethane	5.73x10 ⁻²	133.4	8x10 ⁰	+1
Chloroform	7x10 ⁻²	119.4	8x10 ⁰	+1
Chromium	41	104	4x10 ⁺³	+4
DDT	8.42	354.5	3x10 ⁺³	+3
Dichlorobenzidine	1.69	253.1	4x10 ⁺²	+3
1,1-dichloroethylene	1.47x10 ⁻¹ (I)	97	1x10 ⁺¹	+1

TABLE 10-45. (continued)

Compounds	Slope (mg/kg/day) ⁻¹	Molecular Weight	Potency Index	Order of Magnitude (log ₁₀) Index
Dieldrin	30.4	380.9	1x10 ⁺⁴	+4
Dinitrotoluene	0.31	182	6x10 ⁺¹	+2
Diphenylhydrazine	0.77	180	1x10 ⁺²	+2
Epichlorohydrin	9.9x10 ⁻³	92.5	9x10 ⁻¹	0
Bis(2-chloroethyl)ether	1.14	143	2x10 ⁺²	+2
Bis(chloromethyl)ether	9300(I)	115	1x10 ⁺⁶	+6
Ethylene Dibromide (EDB)	8.51	187.9	2x10 ⁺³	+3
Ethylene Oxide	0.63(I)	44.0	3x10 ⁺¹	+1
Formaldehyde	2.14x10 ⁻² (I)	30	6x10 ⁻¹	0
Heptachlor	3.37	373.3	1x10 ⁺³	+3
Hexachlorobenzene	1.67	284.4	5x10 ⁺²	+3
Hexachlorobutadiene	7.75x10 ⁻²	261	2x10 ⁺¹	+1
Hexachlorocyclohexane				
technical grade	4.75	290.9	1x10 ⁺³	+3
alpha isomer	11.12	290.9	3x10 ⁺³	+3
beta isomer	1.84	290.9	5x10 ⁺²	+3
gamma isomer	1.33	290.9	4x10 ⁺²	+3
Nickel	1.15(W)	58.7	7x10 ⁺¹	+2
Nitrosamines				
Dimethylnitrosamine	25.9(not by q _f ⁺)	74.1	2x10 ⁺³	+3
Diethylnitrosamine	43.5(not by q _f ⁺)	102.1	4x10 ⁺³	+4
Dibutylnitrosamine	5.43	158.2	9x10 ⁺²	+3
N-nitrosopyrrolidine	2.13	100.2	2x10 ⁺²	+2
N-nitroso-N-ethylurea	32.9	117.1	4x10 ⁺³	+4

TABLE 10-45. (continued)

Compounds	Slope (mg/kg/day) ⁻¹	Molecular Weight	Potency Index	Order of Magnitude (log ₁₀) Index
N-nitroso-N-methylurea	302.6	103.1	3x10 ⁺⁴	+4
N-nitroso-diphenylamine	4.92x10 ⁻³	198	1x10 ⁰	0
PCBs	4.34	324	1x10 ⁺³	+3
Phenols				
2,4,6-trichlorophenol	1.99x10 ⁻²	197.4	4x10 ⁰	+1
Tetrachlorodioxin	4.25x10 ⁵	322	1x10 ⁺⁸	+8
Tetrachloroethylene	5.31x10 ⁻²	165.8	9x10 ⁰	+1
Toxaphene	1.13	414	5x10 ⁺²	+3
Trichloroethylene	1.26x10 ⁻²	131.4	2x10 ⁰	0
Vinyl Chloride	1.75x10 ⁻² (I)	62.5	1x10 ⁰	0

Remarks:

1. Animal slopes are 95% upper-limit slopes based on the linearized multistage model. They are calculated based on animal oral studies, except for those indicated by I (animal inhalation), W (human occupational exposure), and H (human drinking water exposure). Human slopes are point estimates, based on the linear nonthreshold model.
2. The potency index is a rounded-off slope in (mMol/kg/day)⁻¹ and is calculated by multiplying the slopes in (mg/kg/day)⁻¹ by the molecular weight of the compound.
3. Not all the carcinogenic potencies presented in this table represent the same degree of certainty. All are subject to change as new evidence becomes available.

Where human data were available for a compound, they were used to calculate the index. Where no human data were available, animal oral studies and animal inhalation studies were used, in that order. Animal oral studies were selected over animal inhalation studies because most of the chemicals have been tested with animal oral studies, thus allowing potency comparisons by route.

The potency index for VDC based on kidney adenocarcinomas in the Maltoni (1980) inhalation bioassay is $1.4 \times 10^{+1} \text{ (mMol/kg/day)}^{-1}$. This is derived as follows: the slope estimate from the Maltoni study, $4.2 \times 10^{-5} (\mu\text{g}/\text{m}^3)^{-1}$, is first converted to units of $(\text{mg}/\text{kg}/\text{day})^{-1}$, assuming a breathing rate of 20 m^3 of air per day and a 70-kg person.

$$4.2 \times 10^{-5} (\mu\text{g}/\text{m}^3)^{-1} \times \frac{1 \text{ day}}{20 \text{ m}^3} \times \frac{1 \mu\text{g}}{10^{-3} \text{ mg}} \times 70 \text{ kg} \\ = 1.47 \times 10^{-1} (\text{mg}/\text{kg}/\text{day})^{-1}.$$

Multiplying by the molecular weight of 97 gives a potency index of $1.4 \times 10^{+1}$. Rounding off to the nearest order of magnitude gives a value of 10^{+1} , which is the scale presented on the horizontal axis of Figure 10-1. The index of $1.4 \times 10^{+1}$ lies at the bottom of the third quartile of the 53 suspect carcinogens.

Ranking of the relative potency indices is subject to the uncertainty of comparing estimates of potency of different chemicals based on different routes of exposure to different species using studies of different quality. Furthermore, all the indices are based on estimates of low-dose risk using linear extrapolation from the observational range. Thus, these indices are not valid to compare potencies in the experimental or observational range if linearity does not exist there.

SUMMARY AND CONCLUSIONS

QUALITATIVE SUMMARY

The carcinogenicity of VDC was evaluated by inhalation exposure of both sexes of Swiss mice and Sprague-Dawley rats to VDC concentrations as high as those maximally tolerated (25 ppm for mice and 150 ppm for rats) for 12 months followed by lifetime observation as well as in Chinese hamsters exposed to 25 ppm VDC (less than a maximally tolerated dose) for 12 months followed by lifetime observation (Maltoni et al., 1980). A statistically significant increase in kidney adenocarcinomas was found in male Swiss mice, and, although a statistically significant increase in mammary carcinomas in treated female Swiss mice was evident, the investigators concluded that a direct relationship between this response in female mice and VDC treatment remains open, largely due to stated results of their statistical analysis indicating a weaker response when the data are adjusted for survival and lack of a dose-related response over the course of the study. A carcinogenic effect of VDC in Sprague-Dawley rats and Chinese hamsters was not apparent. The 12-month exposure period used in this study was below lifetime exposures.

Additional inhalation exposure studies of VDC in animals have been done (Table 10-21). Exposure of male and female Sprague-Dawley rats to 25 ppm and 75 ppm VDC for 18 months followed by 6 months of observation did not show a carcinogenic effect (McKenna et al., 1982); however, the exposure levels used did not appear to be overtly toxic, and use of additional higher exposure levels might have provided a broader evaluation for carcinogenicity.

No statistically significant increase in tumor formation in treated animals was shown in any of the following separate studies of exposure to VDC:

(1) Male and female Wistar rats exposed to 200 ppm for 5 months and 100 ppm for 7 months followed by observation for an additional 12 months (Viola and Caputo, 1977);

(2) Male and female Sprague-Dawley rats exposed to 75 ppm and 100 ppm for an unspecified duration (Viola and Caputo, 1977);

(3) CD-1 mice and CD rats exposed to 55 ppm and observed for a period of 12 months (Lee et al., 1978);

(4) CD-1 mice exposed for 55 ppm for as long as 6 months and observed for 12 additional months (Hong et al., 1981); and

(5) CD rats exposed to 55 ppm for as long as 10 months and observed for an additional 12 months (Hong et al., 1981).

The known exposure periods of 12 months or less used in these studies were shorter than potential lifetime exposure periods, and use of more than one exposure level in the studies with CD-1 mice and CD rats could have provided a stronger evaluation of dose-response.

Several carcinogenicity studies in which VDC was administered orally to experimental animals have been reported as negative. These studies include:

(1) Gavage administration of VDC in olive oil at doses as high as 20 mg/kg to male and female Sprague-Dawley rats for 12 months, followed by lifetime observation (Maltoni et al., 1980);

(2) Administration of VDC in drinking water at levels of maximum water solubility (200 ppm) to male and female Sprague-Dawley rats for 2 years (Quast et al., 1983; Humiston, et al., 1978).

(3) Gavage administration of VDC in corn oil to male and female Fischer 344 rats at 1 and 5 mg/kg and male and female B6C3F1 mice at 2 and 10 mg/kg for 2 years (NCI/NTP, 1981);

(4) Gavage administration of 150 mg/kg of VDC in corn oil to pregnant BDIV rats on day 17 of gestation, followed by weekly gavage doses of 50 mg/kg of VDC to the pups for their lifetimes following birth (Ponomarev and Tomatis, 1980).

The 12-month exposure in Sprague-Dawley rats given VDC by gavage was below the potential lifetime exposure, and a maximally tolerated dose does not appear to have been selected. No clear effect on survival and body weight in Fischer 344 and B6C3F1 mice given VDC in corn oil suggests that higher doses could have been tested to challenge the animals more strongly for carcinogenicity. A broader evaluation of VDC carcinogenicity in BDIV rats might have been possible with the addition of more than the one dose level used in the study, and with more than one treatment each week.

VDC was not carcinogenic in female ICR/Ha mice when applied to the skin at 121 mg/mouse three times per week, or when injected subcutaneously at 2 mg/mouse once weekly at maximum tolerated doses in lifetime studies (Van Duuren et al., 1980). When a single application of 121 mg/mouse of VDC was followed by repeated application of the tumor-promoting agent phorbol myristate acetate, skin papillomas resulted in ICR/Ha mice.

One epidemiologic study on a population of 138 workers (Ott et al., 1976) showed no carcinogenic effects attributable to VDC. At the time the study was published in 1976, the dates of first exposures for 55 workers were between 1960 and 1969, thus indicating that nearly 40% of the workers had no more than 16 years latency since first exposure and that continued follow-up could strengthen the evaluation for carcinogenicity.

Quantitative Summary

Only the Swiss mice inhalation study of Maltoni et al. (1980) provides sufficient evidence for a quantitative cancer risk estimate for VDC. Based on kidney adenocarcinomas in male mice, the 95% upper-limit unit risk estimate for additional cancer from a lifetime of continuous exposure to $1\mu\text{g (VDC)}/\text{m}^3$ (air) is 4.2×10^{-5} . The relative potency of VDC, expressed in molar units, is $1.4 \times 10^{+1} (\text{mMol}/\text{kg}/\text{day})^{-1}$. Among 53 chemicals which the CAG has evaluated as suspect carcinogens, VDC ranks at the bottom of the third quartile.

Conclusions

Evidence for the carcinogenicity of VDC in one strain of mice was found in an inhalation study (Maltoni et al., 1980). Kidney adenocarcinomas were diagnosed in treated male Swiss mice, and, although a statistically significant excess of mammary carcinomas was evident in female Swiss mice, the study investigators concluded that a direct correlation between this response in female mice and VDC treatment could not be made. As discussed herein, VDC has demonstrated mutagenicity in bacterial assay systems with metabolic activation and can be metabolized to intermediates capable of reacting with cellular

macromolecules. A 6-hour exposure of male CD-1 mice and male Sprague-Dawley rats to 10 or 50 ppm VDC was reported to produce minimal DNA alkylation and DNA repair synthesis in liver and kidney, and increased DNA replication only in mouse kidney.

Applying the International Agency for Cancer Research (IARC) approach for classifying carcinogenic agents, this level of evidence can be considered to be limited and not sufficient to make a firm conclusion regarding the carcinogenicity of VDC in experimental animals. Applying IARC criteria for evidence of human carcinogenicity, currently available data could also be considered inadequate for making a judgment on the carcinogenicity of VDC in humans. Based on the overall evidence, VDC would be, according to the IARC method, a Group 3 chemical, by definition according to the IARC method, cannot be classified as to its carcinogenicity to humans.

The kidney adenocarcinomas in male Swiss mice, however, provide data for estimating an upper limit of potential human risk. This 95% upper-limit risk is 4.2×10^{-5} for a lifetime continuous exposure to $1\mu\text{g (VDC)}/\text{m}^3$ air. Among 53 chemicals which the CAG has evaluated as suspect carcinogens, VDC would rank at the bottom of the third quartile.

11. REFERENCES

A.D. Little, Inc. (1976). Vinylidene Chloride Monomer Emissions from the Monomer, Polymer, and Polymer Processing Industries. Prepared by A.D. Little, Inc. under Contract No. 68-02-1332. U.S. Environmental Protection Agency. (Cited in Going and Spigarelli, 1977).

Ames, B.N.; Durston, W.E.; Yamasaki, E.; and Lee, F.D. (1973). Carcinogens are mutagens: A simple test system combining liver homogenates for activation and bacteria for detection. *Proc. Nat. Acad. Sci.* 70:2281-2285.

Andersen, M.E.; Gargas, M.L.; Jones, R.A.; and Jenkins, L.J., Jr. (1979a). The use of inhalation techniques to assess the kinetic constants of 1,1-dichloroethylene metabolism. *Toxicol. Appl. Pharmacol.* 47(2):395-409.

Andersen, M.E.; French, J.E.; Gargas, M.L.; Jones, R.A.; and Jenkins, L.J., Jr. (1979b). Saturable metabolism and the acute toxicity of 1,1-dichloroethylene. *Toxicol. Appl. Pharmacol.* 47(2):385-393.

Andersen, M.E., and Jenkins, L.J., Jr. (1977). Oral toxicity of 1,1-dichloroethylene in the rat: Effects of sex, age, and fasting. *Environ. Health Perspect.* 21:157-163.

Andersen, M.E.; Jones, R.A.; and Jenkins, L.J., Jr. (1978). The acute toxicity of single, oral doses of 1,1-dichloroethylene in the fasted, male rat: Effect of induction and inhibition of microsomal enzyme activities on mortality. Toxicol. Appl. Pharmacol. 46(1):227-234.

Andersen, M.E.; Thomas, O.E.; Gargas, M.L.; Jones, R.E.; and Jenkins, L.G., Jr. (1980). The significance of multiple detoxification pathways for reactive metabolites in the toxicity of 1,1-dichloroethylene. Toxicol. Appl. Pharmacol. 52:422-432.

Anderson, D.; Hodge, M.C.E.; and Purchase, I.F.H. (1977). Dominant lethal studies with the halogenated olefins vinyl chloride and vinylidene chloride in male CD-1 mice. Environ. Health Perspect. 21:71-78.

Anonymous (1978). Aliphatic organics: Vinylidene chloride. Chemical Marketing Reporter, February 27, 1978, p. 13.

Baden, J.M.; Brinkenhoff, M.; Wharton, R.S.; Hitt, B.A.; Simmon, V.F.; and Mazze, R.I. (1976). Mutagenicity of volatile anesthetics: halothane. Anesthesiology 45(3):311-318.

Baden, J.M.; Kelley, M.; Simmon, V.F.; Rice, S.A.; and Mazze, R.I. (1978). Fluroxene mutagenicity. Mutat. Res. 58:183-191.

Baden, J.M.; Kelley, M.; Wharton, R.S.; Hitt, B.A.; Simmon, V.F.; and Mazze, R.I. (1977). Mutagenicity of halogenated ether anesthetics. Anesthesiology 46:346-350.

Ballistreri, A.; Foti, S.; Maravigna, P.; Montaudo, G.; and Scamporrino, E. (1981). Mechanism of thermal decomposition of poly(vinylidene chloride), Polymer. 22:131-132.

Barbin, A.; Planche, G.; Croisy, A.; Malaveille, C.; and Bartsch, H. (1978). Detection of electrophilic metabolites of halogenated olefins with 4-(4-nitrobenzyl) pyridine or with Salmonella typhimurium. Mutat. Res. 53:150.

Bartsch, H.; Malaveille, C.; Barbin, A.; and Planche, G. (1979). Mutagenic and alkylating metabolite of haloethylenes, chlorobutadienes, and dichlorobutenes produced by rodent or human liver tissues. Evidence for oxirane formation by P-450 linked microsomal oxygenase. Arch. Toxicol. 41:249-277.

Bartsch, H. (1976). Mutagenicity tests in chemical carcinogenesis. In: Environmental Pollution and Carcinogenic Risks, IARC Scientific Publications No. 13. INSERM Symposia Series. Lyon, France: International Agency for Research on Cancer, Vol. 52, pp. 229-240.

Bartsch, H.; Malaveille, C.; Montesano, R.; and Tomatis, L. (1975). Tissue mediated mutagenicity of vinylidene chloride and 2-chlorobutadiene in Salmonella typhimurium. Nature (London) 225(5510):641-643.

Bellar, T.A.; Budde, W.L.; and Eichelberger, J.W. (1979). The identification and measurement of volatile organic compounds in aqueous environmental samples. In 94th ACS Symposium Series: Monitoring Toxic Substances, p. 49-62.

Bellar, T.A. and Lichtenberg, J.J. (1979). Semiautomated Headspace Analysis of Drinking Waters and Industrial Waters for Purgeable Volatile Organic Compounds. Measurement of Organic Pollutants in Water and Wastewater, ASTM STP 686, C.E. Van Hall, Ed. American Society for Testing and Materials, p. 108-129.

Bellar, T.A.; and Lichtenberg, J.J. (1974). Determining volatile organics at microgram-per-liter levels by gas chromatography. J. Am. Water Works Assoc. 66(12): 739-744.

Bellar, T.A.; and Lichtenberg, J.J. (1981). The Determination of Halogenated Chemicals in Water by the Purge and Trap Method. Method 502.1. U.S. EPA No. 600/4-81-059.

Bellar, T.A.; Lichtenberg, J.J.; and Kroner, R.C. (1974). The occurrence of organohalides in chlorinated drinking waters. J. Am. Water Works Assoc. 66(12): 703-706.

Birkel, T.J.; Roach, J.A.G.; and Sphon, J.A. (1977). Determination of vinylidene chloride in saran films by electron capture gas-solid chromatography and confirmation by mass spectrometry. J. Assoc. Off. Anal. Chem. 60(5):1210-1213.

Bonse, G.; Urban, T.; Reichert, D.; Henschler, D. (1975). Chemical reactivity, metabolic oxirane formation and biological reactivity of chlorinated ethylenes in the isolated perfused rat liver preparation. Biochem. Pharmacol. 24:1829-1834.

Bonse, G., and Henschler, D. (1976). Chemical reactivity, biotransformation, and toxicity of polychlorinated aliphatic compounds. CRC Critical Rev. Toxicol. 4:395-409.

Bozzelli, J.W.; Kebbekus, B.; and Greenburg, A. (1980). Analysis of Selected Toxic and Carcinogenic Substances in Ambient Air in New Jersey. Prepared for New Jersey Dept. of Environmental Protection; Prepared by New Jersey Institute of Technology, Newark, New Jersey.

Brodzinsky, R., and Singh, H.B. (1982). Volatile Organic Chemicals in the Atmosphere: An Assessment of Available Data. Final report on contract No. 68-02-3452, Environ. Sciences Res. Lab, Office of Res. and Dev., U.S. EPA, Research Triangle Park, N.C.

Bronzetti, G.; Bauer, C.; Corsi, C.; Leporini, C.; Nieri, R.; Del Carratore, R. (1981). Genetic activity of vinylidene chloride in yeast. Mutat. Res. 89: 179-185.

Brown, R.H., and Purnell, C.J., (1979). Collection and analysis of trace organic vapor pollutants in ambient atmospheres. The performance of a Tenax-G adsorbent tube. J. Chromatography, 178:79-90.

Brown, S.L.; Chan, F.Y.; Jones, J.L.; Liu, D.H.; McCaleb, K.E.; Mill, T.; Sapio, K.N.; and Schendel, D.E. (1975). Research Program on Hazard Priority Ranking of Manufactured Chemicals. Phase II: Final Report (Chemicals 1-20), Report No. NSF-RA-E-75-190A. Prepared by Stanford Research Institute, Menlo Park, CA, for Research Applied to National Needs (RANN), National Science Foundation, Washington, DC. Available from: National Technical Information Service, Springfield, VA (NTIS PB-263-161).

Buccafusco, R.J.; Ells, S.J.; LeBlanc, G.A., (1981). Acute toxicity of priority pollutants to bluegill (*Lepomis macrochirus*). Bull. Environ. Contam. Toxicol. 26(4):446-452.

Callahan, M.A. et al., (1979). Water-related Environmental Fate of 129 Priority Pollutants. Vol. 2, U.S. EPA Report No. 440(4-79-029), of Water and Waste Management, U.S. EPA, Washington, D.C.

Carlson, G.P., and Fuller, G.C. (1972). Interaction of modifiers of hepatic microsomal drug metabolism and the inhalation toxicity of 1,1-dichloroethylene. Res. Commun. Chem. Pathol. Pharmacol. 4(3):553-560.

Carpenter, C.P.; Smyth, H.P., Jr.; and Pozzani, D.C. (1949). The assay of acute vapor toxicity, and the grading and interpretation of results on ninety-six chemical compounds. J. Ind. Hyg. Toxicol. 31(6):343-346.

Castello, G.; Gerbino, T.C., and Kanitz, S. (1982). Gas chromatographic analysis of halocarbons in drinking water by headspace extraction and mixed column separation. J. Chromatography. 247:263-272.

Cerna, M., and Kypenova, H. (1977). Mutagenic activity of chloroethylenes analyzed by sceening system tests. Mutat. Res. 46(3)214-215.

CFR (Code of Federal Regulations) (1977). Title 21, 175.300(d). (Cited in Hollifield and McNeal, 1978).

Chasseaud, L.F. (1973). The nature and distribution of enzymes catalyzing the conjugation of glutathione with foreign compounds. Drug Metab. Rev. 2:185-220.

Chieco, P.; Moslen, M.T.; Reynolds, E.S. (1982). Histochemical evidence that plasma and mitochondria membranes are primary foci of hepatocellular injury caused by 1,1-dichloroethylene. Lab. Invest. 46:413-421.

Chieco, P.; Moslen, M.T.; Reynolds, E.S. (1981). Effect of administrative vehicle on oral 1,1-dichloroethylene toxicity. Toxicol. Appl. Pharmacol. 57:146-155.

Chiou, C.T.; Freed, V.H.; Schmedding, D.W.; and Kohnert, R.L. (1977). Partition coefficient and bioaccumulation of selected organic chemicals. Environ. Sci. Technol. 11:475-478.

Chu, K.C., and Milman, H.A. (1981). Review of Experimental Carcinogenesis by Compounds Related to Vinyl Chloride. Environ. Health Perspect. 41:211-220.

Coleman, W.E.; Lingy, R.D.; Melton, R.G.; and Kopfler, F.C. (1976). The occurrence of volatile organics in five drinking water supplies using gas chromatography/mass spectrometry. In: Identification and Analysis of Organic Pollutants in Water. Keith, L.H., editor. Ann Arbor, MI: Ann Arbor Science Publishers, Inc., pp. 305-327.

Coniglio, W.A.; Miller, K., and MacKeever, D. (1980). The Occurrence of Volatile Organics in Drinking Water. Briefing for U.S. EPA Deputy Assistant Administration for Drinking Water. Prepared by Criteria and Standards Division, Science and Technology Branch, U.S. EPA, Washington, D.C.

Cooper, P. (1980). The varying fate of vinylidene chloride. Food Cosmet. Toxicol. 18:435-436.

Costa, A.K.; Ivanetich, K.M. (1982). Vinylidene chloride: its metabolism by hepatic microsomal cytochrome P-450 in vitro. Biochem. Pharmacol. 31:2083-2092.

Cupitt, L.T., (1980). Fate of Toxic and Hazardous Materials in the Air Environment, EPA Report No. EPA-600/83-80-084, NTIS No. PB 80-221948. Prepared by Environmental Sciences Research Lab., Research Triangle Park, N.C.

Dawson, G.W.; Jennings, A.L.; Drozdowski, D.; and Rider, E. (1977). The acute toxicity of 47 industrial chemicals to fresh and saltwater fishes. J. Hazard. Mater. 1(1975/77):303-318.

Dedrick, R.L. (1979). Pharmacokinetics of vinylidene chloride in the rat. Environ. Health Perspect. 28:311-313.

DeLassus, P.T., and Schmidt, D.D. (1981). Solubilities of vinyl chloride and vinylidene chloride in water. J. Chem. Eng. Data. 26:274-276.

DeLeon, I.R.; Maberry, M.A.; Overton, E.B.; Raschke, C.K.; Remele, P.C.; Steele, C.F.; Warren, W.L.; and Laseter, J.L. (1980). Rapid gas chromatographic method for the determination of volatile and semi-volatile organochlorine compounds in soil and chemical waste disposal site samples. J. Chromatogr. Sci. 18:85-88.

Dill, D.E. et al. (n.d.). Toxicity of 1,1-dichloroethylene (vinylidene chloride) to aquatic organisms. Dow Chemical Co. (Manuscript.) (Cited in U.S. EPA, 1979).

Dilling, W.L. (1977). Interphase transfer processes. II. Evaporation rates of chloromethanes, ethanes, ethylenes, propanes, and propylenes from dilute aqueous solutions: Comparison with theoretical predictions. Environ. Sci. Technol. 11(4):405-409.

Dilling, W.L.; Bredeweg, C.J.; and Tefertiller, N.B. (1976). Organic photochemistry: Simulated atmospheric photodecomposition rates of methylene chloride, 1,1-trichloroethane, trichloroethylene, tetrachloroethylene and other compounds. Environ. Sci. Technol. 10(4):351-356.

Dilling, W.L.; Tefertiller, N.B.; and Kallos, G.J. (1975). Evaporation rates and reactivities of methylene chloride, chloroform, 1,1,1-trichloroethane, trichloroethylene, tetrachloroethylene, and other chlorinated compounds in dilute aqueous solutions. Environ. Sci. Technol. 9(9):833-838.

Dowty, B.J.; Carlisle, D.R.; and Laseter, J.L. (1975). New Orleans drinking water sources tested by gas chromatography-mass spectrometry. Environ. Sci. Technol. 9:762-765.

Drevon, C., and Kuroki, T. (1979). Mutagenicity of vinyl chloride, vinylidene chloride and chloroprene in V79 Chinese-hamster cells. Mutat. Res. 67(20):173-182.

Easley, D.M.; Kleopfer, R.D., and Carasea, A.M. (1981). Gas chromatographic-mass Spectrometric determination of volatile organic compounds in fish. J. Assoc. Off. Anal. Chem. 64:653-656.

Federal Register (1979). Method 601, purgeable halocarbons. 44(233):69468-69473.

Filser, J.G., and Bolt, H.M. (1979). Pharmacokinetics of halogenated ethylenes in rats. Arch. Toxicol. 42:123-136.

Foerst, D. (1979). Sampling and analytical method for vinylidene chloride in air. Amer. Ind. Hyg. Assoc. J. 40(10):888-893.

Forkert, P.G.; and Reynolds, E.S. 1982. 1,1-Dichloroethylene-induced pulmonary injury. Exp. Lung. Res. 3(1): 57-68.

Francis, J.R., and Leitch, L.C. (1957). Organic deuterium compounds. XVII. Synthesis of deuteriated methyl chloroform. Can. J. Chem. 35:348-50. Taken from: Chem. Abst. 51:13750d, 1957.

Gage, J.C. (1970). The subacute inhalation toxicity of 109 industrial chemicals. Brit. J. Ind. Med. 27(1):1-18.

Gart, J.J.; Chu, K.C., and Tarone, R.E. (1979). Statistical Issues in Interpretation of Chronic Bioassay Tests for Carcinogenicity. J. Natl. Cancer Inst. 62(4):957-974.

Gay, B.W.; Hanst, P.L.; Bufalini, J.J.; and Noonan, C.D. (1976). Atmospheric oxidation of chlorinated ethylenes. Environ. Sci. Technol. 10(1):58-67.

Gilbert, J.; Shepherd, M.J.; Startin, J.R.; and McWeeny, D.J. (1980). Gas chromatographic determination of vinylidene chloride monomer in packaging films and in foods. J. Chromatography. 197:71-78.

Going, J.E., and Spigarelli, J.L. (1977). Environmental Monitoring Near Industrial Sites: Vinylidene Chloride, Publication No. EPA 560/6-77-026. Prepared by Midwest Research Institute, Kansas City, MO, under Contract No. 68-01-4115. U.S. Environmental Protection Agency, Office of Toxic Substances, Washington, D.C.

Graedel, T.E. (1978). Chemical Compounds in the Atmosphere. Academic Press, New York. pp. 85.

Green, T., and Hathway, D.E. (1977). The chemistry and biogenesis of the S-containing metabolites of vinyl chloride. Chem.-Biol. Interact. 17:137-150

Greim, H.; Bonse, G.; Radwan, Z.; Reichert, D.; and Henschler, D. (1975). Mutagenicity in vitro and potential carcinogenicity of chlorinated ethylenes as a function of metabolic oxirane formation. *Biochem. Pharmacol.* 24(21):2013-2017.

Grimsrud, E.P., and Rasmussen, R.A. (1975). Survey and analysis of halocarbons in the atmosphere by gas chromatography-mass spectrometry. *Atmos. Environ.* 9(11):1014-017.

Gronsborg, E. Sh. (1975). Determination of vinylidene chloride in the air. *Gig. Sanit.* 7:77-79. (In Russ.) Taken from: *Chem. Abst.* 84:21558n, 1976.

Guillemin, C.L.; Martinez, F.; and Thiault, S. (1979). Steam-modified gas-solid-chromatography: A complementary technique for organic pollutant survey. *J. Chromtogr. Sci.* 17:677-681.

Hathway, D.E. (1977). Comparative mammalian metabolism of vinyl chloride and vinylidene chloride in relation to oncogenic potential. *Environ. Health Perspect.* 21:55-59.

Heicklen, J.P.; Sanhueza, E.; Hisatsune, I.E.; Jayanty, R.K.M.; Simonatis, R.; Hull, L.A.; Blume, C.W.; and Mathias, E. (1975). The Oxidation of Halocarbons, Publication No. EPA 650/3-75-008. Prepared by Center for Air Environment Studies, Pennsylvania State University, University Park, PA, under Grant No. 800949 U.S. Environmental Protection Agency, Research Triangle Park, NC. Taken from: *Chem. Abst.* 84:155023t, 1976.

Heitmuller, P.T.; Hollister, T.A; Parrish, P.R. (1981). Acute toxicity of 54 industrial chemicals to sheepshead minnows (*Cyprinodon variegatus*). Bull. Environ. Contam. Toxicol. 27:596-604.

Henck, J.W.; Quast, J.F.; Rampy, L.W., and Norris, J.M. (1980). 10-Day Toxicity Study on Inhaled Vinylidene chloride in four strains of laboratory mice. Report of the Dow Chemical Company, Cited in Norris , 1982.

Henschler, D; and Hoos, R. (1981). Metabolic activation and deactivation mechanisms of di-, tri- and tetrachloroethylenes. Adv. Exp. Med. Biol. 136(A): 659-666.

Hiatt, M.H. (1981). Analysis of fish and sediment for volatile priority pollutants. Anal. Chem. 53: 1541-1543.

Hiatt, M.H. (1983). Determination of volatile organic compounds in fish samples by vacuum distillation and fasted silica capillary gas chromatography/mass spectrometry. Anal. Chem. 55: 506-516.

Hilborn, J.C.; Findlay, W.J.; and Quickert, N. (1976). The application of chemiluminescence to the measurement of reactive hydrocarbons in ambient air. Proc. Int. Conf. Environ. Sensing Assess., Vol. 2, pp. 24-2 to 24-5.

Hollifield, H.C., and McNeal, T. (1978). Gas-solid chromatographic determination of vinylidene chloride in saran film and three food-simulating solvents. J. Assoc. Off. Anal. Chem. 61(3):537-544.

Hong, C.B.; Winston, J.M.; Thornburg, L.P.; Lee, C.C.; and Woods, J.S. (1981). Follow-up study on the carcinogenicity of vinyl chloride and vinylidene chloride in rats and mice. Tumor incidence and mortality subsequent to exposure. J. Toxicol. Environ. Health 7:909-924.

Hull, L.A.; Hisatsume, I.C.; and Heicklen, J. (1973). The reaction of O_3 with CCl_2CH_2 . Can. J. Chem. 51:1504-1510.

Humiston, C.G.; Quast, J.F.; Wade, C.E.; Ballard, J.; Beyer, J.E.; and Lisowe, R.W. (1978). Results of a Two-Year Toxicity and Oncogenicity Study with Vinylidene Chloride Incorporated in the Drinking Water of Rats, MCA Report No. VCD 1.3-Tox-Orl-Dow. Toxicology Research Laboratory Health and Environmental Research, Dow Chemical U.S.A., Midland, MI.

Hushon, J., and Kornreich, M. (1978). Air pollution Assessment of Vinylidene Chloride, Publication No. EPA 450/3-78-015. Prepared by the Metrek Division of the Mitre Corp., McLean, VA, under Contract No. 68-02-1495. U.S. Environmental Protection Agency, Research Triangle Park, NC. Available from: National Technical Information Service, Springfield, VA (NTIS PB-280-624).

Irish, D.D. (1962). Aliphatic halogenated hydrocarbons. In: Toxicology. Vol. 2 of Industrial Hygiene and Toxicology, 2nd ed. Patty, F.A., editor. New York: John Wiley and Sons, Inc., pp. 1305-1308.

Irish, D.D. (1967). Vinylidene Chloride. In: Toxicology. Vol. 2 of Industrial Hygiene and Toxicology, 2nd ed. Patty, F.A., editor. New York: Interscience Publishers, pp. 1245-1248. (Cited in Hushon and Kornreich, 1978).

Jaeger, R.J. (1977). Effect of 1,1-dichloroethylene exposure on hepatic mitochondria. Res. Commun. Chem. Pathol. Pharmacol. 18(1):83-94.

Jaeger, R.J. (1978). Effect of hypoxia on the toxicity of 1,1-dichloroethylene or acrylonitrile. Toxicol. Appl. Pharmacol. 45(1):257.

Jaeger, R.J. (1979). Time-related variation of non-protein sulfhydryl concentrations in rat tissues and human blood. Int. Arch. Occup. Environ. Health 42:141-148.

Jaeger, R.J.; Conolly, R.B.; and Murphy, S.D. (1973a). Diurnal variation of hepatic glutathione concentration and its correlation with 1,1-dichloroethylene inhalation toxicity in rats. Res. Commun. Chem. Pathol. Pharmacol. 6(2):465-471.

Jaeger, R.J.; Trabulus, M.J.; and Murphy, S.D. (1973b). Biochemical effects of 1,1-dichloroethylene in rats. Dissociation of its hepatotoxicity from a lipoperoxidative mechanism. Toxicol. Appl. Pharmacol. 24(3):457-467.

Jaeger, R.J.; Conolly, R.B.; and Murphy, S.D. (1974). Effect of 18-hour fast and glutathione depletion on 1,1-dichloroethylene-induced hepatotoxicity and lethality in rats. Exp. Mol. Pathol. 20(2):187-198.

Jaeger, R.J.; Conolly, R.B.; Reynolds E.S.; and Murphy, S.D. (1975a). Biochemical toxicology of unsaturated halogenated monomers. Environ. Health Perspect. 11:121-128.

Jaeger, R.J.; Conolly, R.B.; and Murphy, S.D. (1975b). Short-term inhalation toxicity of halogenated hydrocarbons: Effects on fasting rats. Arch. Environ. Health 30(1):26-31.

Jaeger, R.J.; Shoner, L.G.; and Coffman, L. (1977a). 1,1-dichloroethylene hepatotoxicity: Proposed mechanism of action and distribution and binding of carbon-14 radioactivity following inhalation exposure in rats. Environ. Health Perspect. 21:113-119.

Jaeger, R.J.; Szabo, S.; and Coffman, L.J. (1977b). 1,1-dichloroethylene hepatotoxicity: Effect of altered thyroid function and evidence for the subcellular site of injury. J. Toxicol. Environ. Health 3(3):545-555.

Jenkins, L.J., Jr., and Andersen, M.E. (1978). 1,1-dichloroethylene nephrotoxicity in the rat. Toxicol. Appl. Pharmacol. 46(1):131-142.

Jenkins, L.J., Jr.; Trabulus, M.J.; and Murphy, S.D. (1972). Biochemical effects of 1,1-dichloroethylene in rats. Comparison with carbon tetrachloride and 1,2-dichloroethylene. Toxicol. Appl. Pharmacol. 23(3):501-510.

John, J.A.; Murray, F.J.; Smith, F.A.; and Schwetz, B.A. (1978). Teratologic evaluation of vinyl chloride, vinylidene chloride, and styrene in laboratory animals. Teratology 17(2):48A.

Jones, B.K., and Hathway, D.E. (1978a). The biological fate of vinylidene chloride in rats. Chem.-Biol. Interact. 20(1):27-41.

Jones, B.K., and Hathway, D.E. (1978b). Differences in metabolism of vinylidene chloride between mice and rats. *Brit. J. Cancer* 37(3):411-417.

Kato, R. (1974). Sex-related differences in drug metabolism. *Drug Metab. Rev.* 3:1-32.

Kato, R.; Vassanelli, P.; Frontino, G.; and Chiesara, E. (1964). Variation in the activity of liver microsomal drug-metabolising enzymes in rats in relation to the age. *Biochem. Pharmacol.* 13:1037-1052.

Keith, L.H., and Telliard, W.A. (1979). Priority pollutants. I. A perspective view. *Environ. Sci. Technol.* 13(4):416-423.

Kobayashi, H., and Rittman, B.E. (1982). Microbial removal of hazardous organic compounds. *Environ. Sci. Technol.* 16:170A-182A.

Krijgsheld, K.R.; Lowe, M.C.; Mimnaugh, E.G.; Trush, M.A.; Ginsburg, E.; and Gram, T.E. (1983). Lung-selective impairment of cytochrome P-450-dependent monooxygenases and cellular injury by 1,1-dichloroethylene in mice. *Biochem. Biophys. Res. Commun.* 110(2): 675-681.

Krost, K.J.; Pellizzari, E.D.; Walburn, S.G.; and Hubbard, S.A. (1982). Collection and analysis of hazardous organic emissions. *Anal. Chem.* 54: 810-817.

Landau, E., and Manos, N.E. (1976). Epidemiology Studies. Task IV: Estimation of Population Residing Near Vinylidene Chloride Plants, Publication No. EPA 560/6-76-022. Prepared by American Public Health Association, Washington, D.C., under Contract No. 68-01-2490. U.S. Environmental Protection Agency, Office of Toxic Substances, Washington, D.C. Available from: National Technical Information Service, Springfield, V.A. (NTIS PB-257-490).

Laumbach, A.D.; Lee, S.; Wong, J.; and Streips, U.N. (1977). Studies on the mutagenicity of vinyl chloride metabolites and related chemicals. Prev. Detect. Cancer Proc. Int. Symp. 1:155-170.

LeBlanc, G.A. (1980). Acute toxicity of priority pollutants to water flea (Daphia magna). Bull. Environ. Contam. Toxicol. 24:684-691.

Lee, C.C.; Bhandari, J.C.; Winston, J.M.; House, W.B.; Dixon, R.L.; and Woods, J.S. (1978). Carcinogenicity of vinyl chloride and vinylidene chloride. J. Toxicol. Environ. Health. 4(1):15-30.

Lee, C.C.; Bhandari, J.C.; Winston, J.M.; House, W.B.; Peters, P.J.; Dixon, R.L.; and Woods, J.S. (1977). Inhalation Toxicity of Vinyl Chloride and Vinylidene Chloride. Health Perspect. 21:25-32.

Leibman, K.C., and Ortiz, E. (1977). Metabolism of halogenated ethylenes. Environ. Health Perspect. 21:91-97.

Lin, S.N.; Fu, F.W.Y.; Bruckner, J.V., and Feldman, S. (1982). Quantitation of 1,1- and 1,2-dichloroethylene in body tissues by purge-and-trap gas chromatography. J. Chromatography 244:311-320.

Loew, G.H.; Kurkjian, E; and Rebagliati, M. (1983). Metabolism and relative carcinogenic potency of chloroethylenes: a quantum chemical structure-activity study. 43(1): 33-66.

Mabey, W.R.; Smith, J.H.; Podoll, R.T.; Johnson, H.L.; Mill, T.; Chou, T.W.; Gates, J.; Partridge, I.W., and Vandenburg, D. (1981). Aquatic Fate Process Data for Organic Priority Pollutants, EPA Report No. 440/4-81-014. Monitoring and Data Support Div, Office of Water Regulations and Standards, Washington, D.C.

Malaveille, C.; Planche, G.; and Bartsch, H. (1977). Factors for efficiency of the Salmonella microsome mutagenicity assay. Chem.-Biol. Interact. 17(2):129-136.

Maltoni, C. (1977). Recent findings on the carcinogenicity of chlorinated olefins. Environ. Health Perspect. 21:1-5.

Maltoni, C.; Ciliberti, A.; Carretti, D. (1982). Experimental contributions in identifying brain potential carcinogens in the petrochemical industry. Ann. N.Y. Acad. Sci. 381:216-249.

Maltoni, C.; Cotti, G.; Morisi, L.; and Chieco, P. (1977). Carcinogenicity bioassays of vinylidene chloride. Research plan and early results. Med. Lav. 68(4):241-262.

McKenna, M.J. (1979). Pharmacokinetics of vinylidene chloride in the rat. Reply to comments. Environ. Health Perspect. 28:313.

McKenna, M.J.; Quast, J.F.; Yakel, H.O.; Balmer, M.F.; Rampy, L.W. (1982). Vinylidene Chloride: A Chronic Inhalation Toxicity and Oncogenicity Study in Rats. Toxicology Research Laboratory, Health and Environmental Sciences, Dow Chemical USA, Midland, Michigan.

McKenna, M.J.; Watanabe, P.G.; and Gehring, P.G. (1977). Pharmacokinetics of vinylidene chloride in the rat. Environ. Health. Perspect. 21:99-105.

McKenna, M.J.; Zemple, J.A.; Madrid, E.O.; and Gehring, P.J. (1978a). The pharmacokinetics of [^{14}C] -vinylidene chloride in rats following inhalation exposure. Toxicol. Appl. Pharmacol. 45(2):599-610.

McKenna, M.J.; Zemple, J.A.; Madrid, E.O.; Braun, W.H.; and Gehring, P.J. (1978b). Metabolism and pharmacokinetic profile of vinylidene chloride in rats following oral administration. Toxicol. Appl. Pharmacol. 45(3):821-835.

McKinney, L.L.; Eldridge, A.C.; and Cowan, J.C. (1959). Cysteine thioethers from chloroethylenes. J. Amer. Chem. Soc. 81:1423-1427.

Mill, T., and Mabey, W.R. (1980). Test Protocols for Evaluating the Fate of Organic Chemicals in Air and Water. Final Report. EPA contract No. 68-03-2227, Environmental Res. Lab, Office of Res. and Dev., U.S. EPA, Athens, GA.

Moore, L. (1980). Inhibition of liver-microsome calcium pump by in vivo administration of carbon tetrachloride, chloroform, and 1,1-dichloroethylene (vinylidene chloride). *Biochem. Pharmacol.* 29(18):2505-2511.

Moore, L. (1982). 1,1-Dichloroethylene inhibition of liver endoplasmic reticulum calcium pump function. *Biochem. Pharmacol.* 31(7):1463-1465.

Murray, F.; Nitschke, K.; Ramey, L.; and Schwetz, B. (1979). Embryotoxicity and fetotoxicity of inhaled or ingested vinylidene chloride in rats and rabbits. *Toxicol. Appl. Pharmacol.* 49:189-202.

Murray, F.J.; Nitschke, K.D.; Smith, F.A.; John, J.A.; Humiston, C.G.; Quast, J.F. and Schwetz, B.A. (1978). Teratology and reproduction studies of vinylidene chloride in rats and rabbits. *Toxicol. Appl. Pharmacol.* 45(1):341 (abst. no. 285).

NCI/NTP (National Cancer Institute/National Toxicology Program) (1980). Experimental Design Status Report/Individual Animal Pathology Report for Vinylidene Chloride.

NCI/NTP (National Cancer Institute/National Toxicology Program) (1982). NTP Technical Report on the Carcinogenesis Bioassay of Vinylidene Chloride in F344/N Rats and B6C3F1/Mice (Gavage Study). NTP No. 80-82. NIH Pub No. 82-1784. NTP Research Triangle Park, N.C. and Bethesda, MD. U.S. Department of Health and Human Services, Public Health Services, National Institute of Health.

Neely, W.B.; Branson, D.R.; and Blau, G.E. (1974). Partition coefficient to measure bioconcentration potential of organic chemicals in fish. Environ. Sci. Technol. 8(13):1113-1115.

Neufeld, M.L.; Sittenfield, M.; Wolk, K.F.; and Boyd, R.E. (1977). Market Input/Output Studies. Task I: Vinylidene Chloride, Publication No. EPA 560/6-77-003. Prepared by Auerbach Associates, Inc., Philadelphia, PA, under Contract No. 68-01-1996. U.S. Environmental Protection Agency, Office of Toxic Substances, Washington, D.C. Available from: National Technical Information Service, Springfield, VA (NTIS PB-273-205).

Nitschke, K.D.; Schwetz, B.A.; Humiston, C.G.; Murray, F.J.; Quast, J.F.; Smith, F.A.; Ioset, H.D.; John, J.A.; Crawford, A.A.; and Schwetz, R.W. (1980). A Multiple Generation Reproduction Study in Rats Maintained on Drinking Water Containing Vinylidene Chloride. Toxicology Research Laboratory, Health and Environmental Research, Dow Chemical USA, Midland, Michigan.

Nitschke, K.D.; Smith, F.A.; Quast, J.F.; Norris, J.M.; and Schwetz, B.A. (1983). A three-generation rat reproductive toxicity study of vinylidene chloride in the drinking water. Fundamental and Applied Toxicology. 3: 75-79.

Norris, J.M. (1977). Toxicological and pharmacokinetic studies on inhaled and ingested vinylidene chloride in laboratory animals. Pap. Synth. Conf. (in preparation), pp.45-50. Taken from: Chem. Abst. 88:45825f, 1978.

Norris, J.M. (1982). Interpretive Review of the Animal Toxicological, Pharmacokinetic/Metabolism, Biomolecular and In-Vitro Mutagenicity Studies on Vinylidene Chloride and the Significance of the findings for Man. Health and Environmental Sciences, The Dow Chemical Company, Midland, Michigan.

Oesch, R. (1972). Mammalian epoxide hydrases: Inducible enxymes catalyzing the inactivation of carcinogenic and cytotoxic metabolites derived from aromatic and olefinic compounds. *Xenobiotica* 3:305-340.

Oesch, F., and Daly, J. (1972). Conversion of naphthalene to transnaphthalene dihydrodiol: Evidence for the presence of a coupled aryl monooxygenase epoxide hydrase system in hepatic microsomes. *Biochem. Biophys. Res. Commun.* 46:1713-1720.

Olynyk, P.; Budde, W.L.; and Eichelberger, J.W. (1981). Simultaneous qualitative and quantitative analysis. I. Precision study of compounds amenable to the inert gas-purge and trap method. *J. Chromatography. Sci.* 19:377-382.

O'Mara, M.M.; Ward, W.; Knechtges, D.P.; and Meyer, R.J. (1973). Fire retardation of polyvinyl chloride and related polymers. In: *Flame Retardancy of Polymeric Materials*. Kuryla, W.C., and Paper, A.J., editors. New York: Marcel Dekker, Inc., Vol. 1, pp. 195-275.

Otson, R.; Williams, D.T., and Biggs, D.C. (1982a). Relationships between raw water quality, treatment, and occurrence of organics in Canadian potable water. *Bull. Environ. Contam. Toxicol.* 28:396-403.

Otson, R.; Williams, D.T., and Bothwell, P.D. (1982b). Volatile organic compounds in Water at Thirty Canadian potable water treatment facilities. J. Assoc. Off. Anal. Chem. 65:1370-1374.

Ott, M.G.; Fishbeck, W.A.; Townsend, J.C.; and Schneider, E.J. (1976). A health study of employees exposed to vinylidene chloride. J. Occup. Med. 18(11):735-738.

Page, W.G. (1981). Comparison of groundwater and surface water for patterns and levels of contamination by toxic substances. Environ. Sci. Technol. 15:1475-1481.

Patterson, J.W., and Kodukala, P.S. (1981). Biodegradation of hazardous organic pollutants. CEP, April 1981. pp. 48-55.

Pearson, C.R., and McConnell, G. (1975). Chlorinated C₁ and C₂ hydrocarbons in the marine environment. Proc. Roy. Soc. London, B. 189:305-322.

Pellizzari, E.D., and Bunch, J.E. (1979). Ambient Air Carcinogenic vapors, Improved Sampling and Analytical Techniques and Field Studies, EPA Report No. EPA-600/2-79-081. Prepared for Environmental Sciences Research Lab., Office of Research and Development, U.S. EPA, Research Triangle Park, N.C.

Pellizzari, E.D.; Erickson, M.D., and Zweidinger, R.A. (1979). Formulation of a Preliminary Assessment of Halogenated Organic Compounds in man and Environmental Media, EPA Report No. EPA/560/13-79/006, NTIS PB No. 80-112170. Prepared by Research Triangle Institute, Research Triangle Park, N.C.

Pellizzari, E.D. (1978a). Measurement of Carcinogenic Vapors in Ambient Atmospheres. NTIS PB-283023, EPA Report No. 600/7-78-062, prepared for U.S. EPA, by Research Triangle Institute, Research Triangle Park, N.C.

Pellizzari, E.D. (1978b). Quantification of Chlorinated Hydrocarbons in Previously Collected Air Samples. NTIS PB-289804, U.S. EPA Report No. 450/3-78-112. Prepared by Research Triangle Institute, Research Triangle Park, N.C.

Pereira, W.E., and Hughes, B.A. (1980). Determination of selected volatile organic priority pollutants in water by computerized gas chromatography-quadrupole mass spectrometry. J. Am. Water Works Assoc. 72:220-230.

Piet, G.J.; Slingerland, P.; de Grunt, F.E.; v.d. Heuvel, M.P.M.; and Zoeteman, B.C.J. (1978). Determination of very volatile halogenated organic compounds in water by direct head-space analysis. Anal. Lett. A11(5):437-448.

Ponomarev, V., and Tomatis, L. (1980). Long-term testing of vinylidene chloride and chloroprene for carcinogenicity in rats. Oncology 37:136-141.

Poplawski-Tabarelli, S., and Uehleke, H. (1982). Inhibition of microsomal drug oxidations by aliphatic halohydrocarbons: correlation with vapor pressure. Xenobiotica. 12(1):55-61.

PPG Industries (1980). Verbal communication between Syracuse Research Corp. and a PPG plant employee. April 11, 1980.

Prendergast, J.A.; Jones, R.A.; Jenkins, L.J., Jr.; and Siegel, J. (1967). Effects on experimental animals of long-term inhalation of trichloroethylene, carbon tetrachloride, 1,1,1-trichloroethane, dichlorodifluoromethane, and 1,1-dichloroethylene. *Toxicol. Appl. Pharmacol.* 10(2):270-289.

Quast, J.F.; Humiston, C.G.; Wade, C.E. et al. (1983). A chronic toxicity and oncogenicity study in rats and subchronic toxicity study in dogs on ingested vinylidene chloride. *Fundam. Appl. Toxicol.* 3(1): 55-62.

Rampy, L.W.; Quast, J.F.; Humiston, C.G.; Balmer, M.F.; Schwetz, B.A. (1977). Interim results of two-year toxicological studies in rats of vinylidene chloride incorporated in the drinking water or administered by repeated inhalation. *Environ. Health Perspect.* 21:33-43.

Ray, P., and Moore, L. (1982). 1,1-Dichloroethylene inhibition of liver microsomal calcium pump in vitro. *Arch. Biochem. Biophys.* 218(1):26-30.

Reichert, D., and Bashti, N. (1976). Metabolism and disposition of 1,1-dichloroethylene in the isolated blood perfused liver of the rat. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 293(suppl.):R64.

Reichert, D., and Henschler, D. (1978). Uptake and hepatotoxicity of 1,1-dichloroethylene by the isolated blood-perfused rat liver. *Int. Arch. Occup. Environ. Health* 41(3):169-178.

Reichert, D., Werner, H.W., and Henschler, D. (1978). Role of liver glutathione in 1,1-dichloroethylene metabolism and hepatotoxicity in intact rats and isolated perfused rat liver. Arch. Toxicol. 41(3):169-178.

Reichert, D., Werner, H.W., Metzler, M., and Henschler, D. (1979). Molecular mechanism of 1,1-dichloroethylene toxicity: Excreted metabolites reveal different pathways of reactive intermediates. Arch. Toxicol. 42(3):159-169.

Reitz, R.H., Watanabe, P.G., McKenna, M.J., Quast, J.F., and Gehring, P.J. (1980). Effects of vinylidene chloride on DNA synthesis and DNA repair in the rat and mouse: A comparative study with dimethylnitrosamine. Toxicol. Appl. Pharmacol. 52: 357-370.

Research Triangle Institute. (1977a). Identification and Analysis of Ambient Air Pollutants Using the Combined Techniques of Gas Chromatography and Mass Spectrometry. Quarterly Report 1, April 7, 1977. Research Triangle Park, NC. (Cited in Hushon and Kornreich, 1978).

Research Triangle Institute. (1977b). Development of Techniques for Measuring Ambient Atmospheric Carcinogenic vapors. Monthly Progress Report 8, April 6, 1977. Research Triangle Park, NC. (Cited in Hushon and Kornreich, 1978).

Reynolds, E.S., Moslen, M.T., Boor, P.J., and Jaeger, R.J. (1980). 1,1-Dichloroethylene hepatotoxicity. Time course of GSH changes and biochemical aberrations. Am. J. Pathol. 101(2):331-343.

Reynolds, E.S., and Moslen, M.T. (1977). Damage to hepatic cellular membranes by chlorinated olefins with emphasis on synergism and antagonism. Environ. Health Perspect. 21:137-147.

Reynolds, E.S., Moslen, M.T., Szabo, S., Jaeger, R.J., and Murphy, S.D. (1975). Hepatotoxicity of vinyl chloride and 1,1-dichloroethylene: Role of mixed function oxidase system. Am. J. Pathol. 81:219.

Russell, J.W. (1975). Analysis of air pollutants using sampling tubes and gas chromatography. Environ. Sci. Technol. 9(13):1175-1178.

Sanhueza, E., Hisatsune, I.C., and Heicklen, J. (1976). Oxidation of haloethylenes. Chem. Rev. 76:801-826.

Sato, A., Nakajima, T., and Koyama, Y. (1980). Effects of chronic ethanol consumption on hepatic metabolism of aromatic and chlorinated hydrocarbons in rats. Br. J. Ind. Med. 37(4):382-386.

Schwarzenbach, R.P., and Westall, J. 1981. Transport of non-polar organic compounds from surface water to groundwater. Laboratory sorption studies. Environ. Sci. Technol. 15:1360-1367.

Severs, L.W., and Skory, L.K. (1975). Monitoring personnel exposure to vinyl chloride, vinylidene chloride and methyl chloride in an industrial work environment. Amer. Ind. Hyg. Assoc. J. 36(9):669-691.

Shackelford, W.M., and Keith, L.H. (1976). Frequency of organic compounds identified in water. Publication No. EPA-600/4-76-062. U.S. Environmental Protection Agency, Office of Research and Development, Environmental Research Laboratory, Athens, GA.

Short, R.D., Winston, J.M., Minor, J.L., Hong, C.D., Seifter, J., and Lee, C.C. (1977a). Toxicity of vinylidene chloride in mice and rats and its alterations by various treatments. J. Toxicol. Environ. Health. 3:(5-6):913-921.

Short, R.D., Winston, J.M., Minor, J.L. Seifter, J. and Lee, C.C. (1977b). Effect of various treatments on toxicity of inhaled vinylidene chloride. Environ. Health Perspect. 21:125-129.

Short, R.D., Jr., Minor, J.L., Winston, J.M., Ferguson, B., and Unger, T. (1977c). Toxicity Studies of Selected Chemicals. Task II. The Developmental Toxicity of Vinylidene Chloride Inhaled by Rats and Mice During Gestation. Publication No. EPA-560/6-77-022. Prepared by Midwest Research Institute, Kansas City, MO, under Contract No. 68-01-3242. U.S. Environmental Protection Agency, Office of Toxic Substances, Washington, DC.

Short, R.D. Minor, J.L., Winston, J.M., and Lee, C.C. (1977d). A dominant lethal study in male rats after repeated exposures to vinyl chloride or vinylidene chloride. J. Toxicol. Environ. Health. 3:5-6.

Siegel, J., Jones, R.A., Coon, R.A., and Lyon, J.P. (1971). Effects on experimental animals of acute repeated and continuous inhalation exposures to dichloroacetylene mixtures. Toxicol. Appl. Pharmacol. 18(1):168-174.

Siegers, C.P.; Huebscher, W.; and Younes, M. 1982. Glutathione S-transferase and GSH peroxidase activities during the state of GSH depletion leading to lipid peroxidation in rat liver. Res. Commun. Chem. Pathol. Pharmacol. 37(2): 163-169.

Siletnik, L.M., and Carlson, G.P. (1974). Cardiac sensitizing effects of 1,1-dichloroethylene: Enhancement by phenobarbital pretreatment. Arch. Int. Pharmacodyn. 210(2):359-364.

Simmon, V.F. (1978). Structural correlations of carcinogenic and mutagenic alkyl halides. In: Structural Correlations between Carcinogenicity and Mutagenicity, DHEW (ISS-FDA) Publication No. 78-1046, pp. 163-171.

Simmon, V.F., Kauhanen, K., and Tardiff, R.C. (1977). Mutagenic activity of chemicals identified in drinking water. In: Progress in Genetic Toxicology.

Scott, D., Bridges, B.A., and Sobels, F.H., editors. Proceedings of the Second International Conference on Environmental Mutagens, Edinburgh, July 11-15, 1977. Amsterdam, The Netherlands: Elsevier/North-Holland Biomedical Press, pp. 249-258.

Simmon, V.F., Rosenkranz, H.S., Zeiger, E., and Poirier, L.A. (1979). Mutagenic activity of chemical carcinogens and related compounds in the intraperitoneal host-mediated Assay. J. Natl. Cancer Inst. 62:911-918.

Singh, H.B., Salas, L.J., Smith, A., and Shigeishi, H. (1979). Atmospheric measurements of selected toxic organic chemicals. Report on EPA Grant No. 805990. Prepared for Environmental Sciences Research Lab. U.S. EPA, Research Triangle Park, NC.

Singh, H.B., Salas, L.J., Smith, A., Stiles, R., and Shigeishi, H. (1981a). Atmospheric measurements of selected hazardous chemicals. NTIS PB 81-200628. Prepared for Environmental Sciences Research Lab., Office of Research Development. U.S. EPA, Research Triangle Park, NC.

Singh, H.B., Salas, L.J., Smith, A.J., and Shigeiski, H. (1981b). Measurements of some potentially hazardous organic chemicals in urban environments. Atmos. Environ. 15:601-612.

Suta, B.E. (1979). Assessment of Human Exposures to Atmospheric Acrylonitrile. Prepared by Stanford Research Institute, Menlo Park, CA, under Contract No. 68-02-2835. U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards, Research Triangle Park, NC.

Sweger, D.M., and Travis, J.C. (1979). An application of infrared lasers to the selective detection of trace organic gases. Appl. Spectrosc. 33(1):46-51.

Szabo, S.; Jaeger, R.J.; Moslen, T.M.; and Reynolds, E.S. (1977). Modification of 1,1-dichloroethylene hepatotoxicity by hypothyroidism. Toxicol. Appl. Pharmacol. 42(2):367-376.

Tabak, H.H., Quave, S.A., Mashni, C.I., and Barth, E.F. (1981). Biodegradability studied with organic priority pollutant compounds. J. Water Pollut. Control Fed. 53:1503-1518.

Tute, M.S. (1971). Principles and practice of Hansch analysis: A guide to structure-activity correlation for the medicinal chemist. Adv. Drug Res. 6:1-77.

Uehleke, H., Poplawski, S., Bonse, G., and Henschel, D. (1976). Spectral evidence for 2,2,3-trichloro-oxirane formation during microsomal trichloroethylene oxidation. Naunyn-Schmiedeberg's Arch. Pharmacol. 293(suppl.):R64.

U.S. EPA (U.S. Environmental Protection Agency) (1977). The National Organic Monitoring Survey. Technical Support Division, Office of Water Supply, U.S. EPA.

U.S. EPA (U.S. Environmental Protection Agency) (1978). In-Depth Studies on Health and Environmental Impacts of Selected Water Pollutants. Contract No. 68-01-4646. U.S. EPA. (Results summary obtained from Mr. John Carroll, Project Monitor, U.S. EPA, Washington, DC.)

U.S. EPA (U.S. Environmental Protection Agency) (1979). Ambient Water Quality Criteria Document for Dichloroethylenes. Draft. U.S. EPA, Criteria and Standards Division, Office of Water Planning and Standards, Washington, DC. Available from: National Technical Information Service, Springfield, VA (NTIS PB 292-430).

U.S. EPA (U.S. Environmental Protection Agency) (1981). Treatability Manual. Volume I. Treatability Data. U.S. EPA Report No. 600/2-82-001a, Office of Research and Development, U.S. EPA, Washington, DC.

Van Duuren, B.L., Goldschmidt, B.M. Loewengert, G., Smith, A.C., Melchienne, S., Seidman, L., Roth, D. (1979). Carcinogenicity of halogenated olefinic and aliphatic hydrocarbons in mice. J. National Cancer Inst. 63:1433-1439.

Van't Hof, J., and Schairer, L.A. (1982). Tradescantia assay system for gaseous mutagens. A report of the U.S. Environmental Protection Agency Gene-Tox Program. Mutat. Res. 99(3):303-315.

Veith, G.D., DeFoe, D.L., and Bergstedt, B.V. (1979). Measuring and estimating the bioconcentration factor of chemicals in fish. J. Fish Res. Board Can. 36:1040-1048.

Viola, P.L., and Caputo, A. (1977). Carcinogenicity studies on vinylidene chloride. Environ. Health Perspect. 21:45-47.

Wallace, L. (1981). Measurements of Volatile Organic Compounds in Breathing-Zone Air, Drinking Water, and Exhaled Breath. (Preliminary draft).

Wapora, Inc. (1982). Human Exposure to Atmospheric Concentrations of Vinylidene Chloride. Final Report. Project 507-13. Prepared for the U.S. EPA, Research Triangle Park, NC.

Waskell, L. (1978). Study of the mutagenicity of anesthetics and their metabolites. *Mutat. Res.* 57:141-153.

Wessling, R., and Edwards, F.C. (1970). Poly(vinylidene chloride). In: Kirk-Othmer Encyclopedia of Chemical technology, 2nd ed. Standen, A., editor. New York: John Wiley and Sons, Inc., Vol. 21, pp. 275-303.

Williams, F.W., Stone, J.P., and Eaton, H.G. (1976). Personal atmospheric gas sampler using the critical orifice concept. *Anal. Chem.* 48(2):442-445.

Yllner, S. (1971). Metabolism of chloroacetate-1-¹⁴C in the mouse. *Acta Pharmacol. Toxicol.* 30:69-80.

Zemaitis, M.A., and Green, F.E. (1976). Impairment of hepatic microsomal drug metabolism in the rat during daily disulfiram administration. *Biochem. Pharmacol.* 25:1355.

U.S. Environmental Protection Agency
Region V Library
200 South Dearborn Street
Chicago, Illinois 60604