



## *Project Summary*

# Sulfur Dioxide Oxidation Reactions in Aqueous Solutions

L. R. Martin, D. E. Damschen, and H. S. Judeikis

The kinetics and mechanisms of more than one hundred reactions involving the aqueous phase oxidation of sulfur dioxide and nitrogen oxides in mixed catalyst/oxidant systems at pH = 0 to 3 were investigated.

The systems investigated involve different redox reaction types:  $\text{SO}_2/\text{O}_2$ ;  $\text{SO}_2/\text{NO}_x$ ;  $\text{SO}_2/\text{H}_2\text{O}_2$ ;  $\text{SO}_2/\text{O}_3$ ;  $\text{NO}_x/\text{O}_3$ ;  $\text{NO}_x/\text{H}_2\text{O}_2$ . The  $\text{SO}_2/\text{O}_2$  system demonstrated little reactivity unless a catalyst was present. The most effective metal catalysts were  $\text{Fe}^{+++}$  and  $\text{Mn}^{++}$  with a complex synergism between Fe and Mn.

In the  $\text{SO}_2/\text{NO}_x$  system the nitrite ion caused a rapid reaction producing  $\text{N}_2\text{O}$ . The reaction did show slight inhibition by toluene and hexene, but was unaffected by metal ion catalysts.

The  $\text{SO}_2/\text{H}_2\text{O}_2$  reaction in water is extremely fast and insensitive to catalysis or inhibition.

The  $\text{SO}_2/\text{O}_3$  reaction is also rapid and similarly insensitive to catalysis or inhibition.

Nitrite oxidation by both ozone and hydrogen peroxide is rapid. The pH dependence of these two reactions was determined.

*This Project Summary was developed by EPA's Environmental Sciences Research Laboratory, Research Triangle Park, NC, to announce key findings of the research project that is fully documented in a separate report of the same title (see Project Report ordering information at back).*

### Introduction

Our understanding of the atmospheric chemistry of sulfur and nitrogen oxides

is important for establishing control strategies both for the primary pollutants, sulfur dioxide and nitric oxide, and for their aerosol formation products, sulfates and nitrates. Research in the past ten years has identified the importance of several gas phase reactions leading to sulfate or nitrate formation. It is only recently that liquid droplet phase mechanisms have been addressed as a possible explanation for the wintertime presence of sulfate and nitrate in regions where photochemical oxidant potential is obviously low.

This project has addressed, in a systematic fashion, over one hundred aqueous phase reactions of potential significance in sulfate and nitrate formation. Environmentally important metal catalysts, commonly found oxidants ( $\text{O}_2$ ,  $\text{O}_3$ ,  $\text{H}_2\text{O}_2$ ) in the ambient air, and potential inhibitors or accelerators of the main reactions,  $\text{SO}_2 \rightarrow \text{SO}_4^-$  and  $\text{NO}_x \rightarrow \text{NO}_3^-$  were studied for their reaction rates.

### Procedure

The apparatus used consisted of an ultraviolet light source, a Durrum D-103 stopped flow system, monochromator (PAR 1208) and two detection systems interfaced to a minicomputer. The detection systems used, depending on reactions conditions, were (1) a PAR 1205D Silicon Intensified Target Detector linked to a PAR 1205A Optical Multichannel Analyzer and (2) an RCA IP28 phototube maintained at -650v by a Fluke 405 Power Supply. The tube is linked to a Biomation 802 Transient Recorder to record waveshape during selected periods of time. The minicomputer used was

a Cromemco Z2 which utilizes BASIC language.

Fresh aqueous solutions were prepared daily using 18 megohm resistivity water obtained from a Milli-Q water purification system. HCl was used to maintain a pH of less than 4. The pH was measured before each run using an Orion 701A digital ionalyzer. The water for the SO<sub>2</sub> solutions was deoxygenated by passing a steady flow of nitrogen through it. For the ozone reactions an Orec Model 03V5-0 ozonator was used to prepare 1x10<sup>-4</sup>M solutions. The reductant is placed on one side of the stopped flow system while the oxidant plus catalysts or inhibitors is placed on the other. The capability to detect 1x10<sup>-5</sup>M of the absorbing species existed; usually solutions were prepared at concentrations of 1x10<sup>-3</sup>M to 1x10<sup>-4</sup>M to obtain a better signal to noise ratio.

## Results

The following rate expressions have been determined during the course of this project:

- (1) For the SO<sub>2</sub> + O<sub>2</sub> reaction at pH = 2

$$d(S_{VI})/dt = 1970 (Mn^{++}) + 82(Fe^{+++}) \\ \left[ 1 + \frac{2.67 \times 10^3 (Mn^{++})^{1.5}}{6.31 \times 10^{-6} + (Fe^{+++})} \right] (S_{IV})$$

The reaction is zeroth order in O<sub>2</sub>.

- (2) For the SO<sub>2</sub> + NO<sub>2</sub><sup>-</sup> reaction:

$$d(S_{VI})/dt = 142(H^+)^{1/2} (HNO_2 + NO_2^-) \\ (SO_2 \cdot Aq + HSO_3^-)$$

- (3) For the SO<sub>2</sub> + H<sub>2</sub>O<sub>2</sub> reaction:

$$d(S_{VI})/dt = \frac{7.2 \times 10^4 (H_2O_2)(SO_2 \cdot Aq)}{0.1 + (H^+)} \\ = \frac{4.5 \times 10^6 (HSO_3^-)(H_2O_2)(H^+)}{0.1 + H^+}$$

- (4) For the SO<sub>2</sub> + O<sub>3</sub> reaction:

$$d(S_{VI})/dt = \frac{1.9 \times 10^4 (SO_2 \cdot Aq + HSO_3^-)(O_3)}{(H^+)^{1/2}}$$

- (5) For the NO<sub>2</sub><sup>-</sup> + O<sub>3</sub> reactions:

$$\frac{-d(O_3)}{dt} = \frac{200(O_3)(HNO_2 + NO_2^-)}{(H^+)}$$

- (6) For the NO<sub>2</sub><sup>-</sup> + H<sub>2</sub>O<sub>2</sub> reaction:

$$\frac{-d(N_{III})}{dt} = 4.6 \times 10^3 (N_{III})(H_2O_2)(H^+)$$

## Conclusions

Of the six reaction rate expressions listed in the Results section, the SO<sub>2</sub> + H<sub>2</sub>O<sub>2</sub> reaction appears the most likely to be significant in the atmosphere when acidic aerosols are present, but any of the six can become important under certain conditions. For example, the Fe/Mn synergistic effect may be the predominant pathway to sulfate formation in a coastal urban environment during wintertime conditions.

As previously stated the SO<sub>2</sub> + H<sub>2</sub>O<sub>2</sub> appears to be the most significant of those studied. At low pH (0 to 3) the reaction is faster than one might expect from a straight line extrapolation of the higher pH data obtained by others. The peroxide reaction was found to be insensitive to catalysis or inhibition by Fe<sup>+++</sup>, Mn<sup>++</sup>, toluene, pinene, hexene or hydroquinone.

The temperature dependence of the SO<sub>2</sub> + H<sub>2</sub>O<sub>2</sub> reaction was determined and the conclusion was that the relatively low activation energy for the kinetics leads to a net increase in the atmospheric rate as the temperature is lowered because of the increasing Henry's Law constants for SO<sub>2</sub> and H<sub>2</sub>O.

Estimates of the atmospheric oxidation rate for 1 ppb H<sub>2</sub>O<sub>2</sub> in the gas phase and in a typical cloud are on the order of 2% min<sup>-1</sup> for SO<sub>2</sub>, which is higher than any reported rates, and as high as the rate allowed by the heterogeneous reactivity of SO<sub>2</sub> with droplet surfaces.

*L. R. Martin, D. E. Damschen, and H. S. Judeikis are with Aerospace Corporation, El Segundo, CA 90245.*

*H. Barnes is the EPA Project Officer (see below).*

*The complete report, entitled "Sulfur Dioxide Oxidation Reactions in Aqueous Solution," (Order No. PB 81-196 834; Cost: \$6.50, subject to change) will be available only from:*

*National Technical Information Service  
5285 Port Royal Road  
Springfield, VA 22161  
Telephone: 703-487-4650*

*The EPA Project Officer can be contacted at:  
Environmental Sciences Research Laboratory  
U.S. Environmental Protection Agency  
Research Triangle Park, NC 27711*

United States  
Environmental Protection  
Agency

Center for Environmental Research  
Information  
Cincinnati OH 45268

Postage and  
Fees Paid  
Environmental  
Protection  
Agency  
EPA 335



Official Business  
Penalty for Private Use \$300

PS 0000329  
U S ENVIR PROTECTION AGENCY  
REGION 5 LIBRARY  
230 S DEARBORN STREET  
CHICAGO IL 60604