



## Project Summary

# Modeling of SO<sub>2</sub> Oxidation in Smog

David F. Miller

Smog chamber experiments were conducted to investigate the kinetics of the free radical reactions of SO<sub>2</sub> in smog and the transformation of SO<sub>2</sub> to sulfate under simulated urban and rural atmospheric conditions. Rate constants were derived for three reactions: SO<sub>2</sub> + HO + M → SULFATE (60); SO<sub>2</sub> + HO<sub>2</sub> → SULFATE (61); and SO<sub>2</sub> + CH<sub>3</sub>O<sub>2</sub> → SULFATE (64); where k<sub>60</sub> = 1600, k<sub>61</sub> < 0.2, and k<sub>64</sub> < 0.8 ppm<sup>-1</sup> min<sup>-1</sup>, respectively. Oxidation of SO<sub>2</sub> by the unstable intermediate HO led exclusively to particulate sulfate. However, even under favorable NO<sub>x</sub> conditions, particulate nitrate was not a product. Hydrocarbon mixtures typical of urban environments promoted SO<sub>2</sub> oxidation faster than hydrocarbons selected to represent rural conditions. Maximum SO<sub>2</sub> oxidation rates occurred during periods of peak NO<sub>2</sub> formation. Oxidation rates of SO<sub>2</sub> in aged smog were 25 to 35% of the maximum rates.

Model calculations indicated that the SO<sub>2</sub> + HO + M → SULFATE reaction dominated sulfate production from SO<sub>2</sub> in polluted air. The model also indicated that SO<sub>2</sub> oxidation is heavily influenced by solar radiation intensity and pollutant conditions. For clear-sky and a variety of HC-NO<sub>x</sub> conditions, maximum SO<sub>2</sub> conversion rates ranged from 3.7 to 7.4%/h.

Models used to simulate free radical reactions in power plant plumes showed that SO<sub>2</sub> oxidation is strongly dependent on plume dispersion rates, as well as on the same parameters governing SO<sub>2</sub> oxidation in ambient

air. Although exceptions do occur for certain HC-NO<sub>x</sub> conditions, SO<sub>2</sub> oxidation rates in plumes are generally bound by the ambient rates.

*This Project Summary was developed by EPA's Environmental Sciences Research Laboratory, Research Triangle Park, NC, to announce key findings of the research project that is fully documented in a separate report of the same title (see Project Report ordering information at back).*

### Introduction

The widespread distribution of sulfate aerosols in the atmosphere and their associated effects on visibility and precipitation pH (acid rain) are serious environmental concerns in the United States. The Environmental Sciences Research Laboratory of the U.S. Environmental Protection Agency is working to formulate strategies for the effective control of sulfates. One step toward achieving such control is the development of models to predict regional production of sulfate by SO<sub>2</sub> sources. The study summarized here was conducted by Battelle Columbus Laboratories to broaden the information base available on SO<sub>2</sub> reactions and transformations and to provide the Environmental Sciences Research Laboratory with the predictive models necessary for sulfate control.

Oxidation of SO<sub>2</sub> in the atmosphere is considered the primary source of sulfate particulates. The chemical and microphysical processes involved in the gas-to-particulate transformations of oxidation, however, are highly complex.

Despite numerous research efforts to define these processes more clearly, many essential details pertaining to reaction rates and mechanisms are missing. To date, kinetic and mechanistic studies of the elementary gas phase reactions of SO<sub>2</sub> in air have yielded only that all known reactions involving electronically excited states of SO<sub>2</sub> are insignificantly slow in the troposphere and that HO, HO<sub>2</sub>, and RO<sub>2</sub> are the only tropospheric species capable of oxidizing SO<sub>2</sub> at substantial rates.

Numerous studies have presented evidence that submicron-radii sulfates are predominantly (NH<sub>4</sub>)<sub>x</sub>H<sub>2-x</sub>SO<sub>4</sub> with associated water. However, although microphysical aspects of H<sub>2</sub>SO<sub>4</sub> particle formation are well understood, much uncertainty surrounds defining new particle nucleation. Some investigators have concluded that SO<sub>2</sub> oxidation in the gas phase is the rate limiting step for the formulation and growth of secondary H<sub>2</sub>SO<sub>4</sub> particles in the atmosphere. One objective of this study was to examine these rate limiting reactions.

Finally, based on the new findings and on additional smog chamber data obtained with rural and urban hydrocarbon systems, a photochemical smog model was applied to simulate gas phase SO<sub>2</sub> oxidation rates for various atmospheric conditions.

## Procedure

Experiments to investigate the kinetics of the free radical reactions of SO<sub>2</sub> in smog and to obtain data on SO<sub>2</sub> transformation to sulfate under various atmospheric conditions were conducted in Battelle's 17.4 m<sup>3</sup> smog chamber. Rate constants were derived for the following reactions: SO<sub>2</sub> + HO + M → SULFATE (60); SO<sub>2</sub> + HO<sub>2</sub> → SULFATE (61); and SO<sub>2</sub> + CH<sub>3</sub>O<sub>2</sub> → SULFATE (64). Pseudo second-order rate constants (ppm<sup>-1</sup> min<sup>-1</sup>) were k<sub>60</sub> = 1600, k<sub>61</sub> < 0.2, and k<sub>64</sub> < 0.8. The values for k<sub>61</sub> and k<sub>64</sub> were only upper limit estimates; the rates pertained to the overall conversion of SO<sub>2</sub> to particulate sulfate.

Although the mechanisms of SO<sub>2</sub> conversion were not investigated, some work was done to determine if NO and/or NO<sub>2</sub> form stable aerosol products with the intermediates created after HO addition to SO<sub>2</sub>.

## Methods for Kinetic Studies

The smog chamber facility has a surface:volume ratio of 0.8 m<sup>-1</sup>; the surface is polished aluminum (~80%)

and FEP Teflon™ (~20%). Irradiation was provided by a bank of 95 fluorescent blacklamps and 15 fluorescent sunlamps. Combined intensity of the lamps yielded a NO<sub>2</sub> photodissociation rate of approximately 0.14 min<sup>-1</sup>.

Background air for the chamber was passed through a purification system for dehumidification and filtering. SO<sub>2</sub> and NO<sub>2</sub> concentrations were usually below the detection levels of 1 ppb and 2 ppb, respectively. Background NO concentrations ranged from 1 to 2 ppb. Background total hydrocarbon ranged between 1.9 and 2.5 ppmC (of which most was CH<sub>4</sub>). Gas chromatographic analyses of background air showed no detectable olefins or aromatics in the C<sub>2</sub>-C<sub>9</sub> range when nonmethane hydrocarbon was 0.1 to 0.3 ppmC. When relatively small concentrations of nitrous acid (a source of HO radicals and NO<sub>x</sub>) were added to the background air under high molecular weight hydrocarbon contamination, irradiation resulted in peak O<sub>3</sub> concentrations in the 8 to 12 ppb range.

The reactivity of this amount of hydrocarbon contamination in terms of O<sub>3</sub> production corresponds to the reactivity of about 4 to 6 ppm CO. Thus, intentional addition of 4.5 ppmC hydrocarbons and/or large quantities of CH<sub>4</sub> and CO clearly overwhelmed any effects due to background HC contaminants. In addition, intentional introduction of nitrous acid into the air for most of the experiments was thought to overcome any spurious surface contaminant effects which might have led to the generation of free radicals at the outset of irradiation.

Gas phase analyses included measurement of PAN, C<sub>1</sub>-C<sub>9</sub> hydrocarbons, CO, O<sub>3</sub>, H<sub>2</sub>O<sub>2</sub>, HNO<sub>2</sub>, NO<sub>2</sub>, NO<sub>x</sub>, NO and SO<sub>2</sub>. Initial measurements of SO<sub>2</sub> were taken with a Meloy 285 flame photometer. Upon irradiation, the instrument was used to monitor sulfate aerosol. The photometer was intermittently operated in the total sulfur mode (SO<sub>2</sub> plus sulfate) to check on the SO<sub>2</sub> concentration.

Gas calibrations were performed routinely. The chemiluminescence analyzer used to monitor NO, HNO<sub>2</sub>, and NO<sub>2</sub> was spanned before almost every experiment. *In situ* O<sub>3</sub> titrations were conducted in the chamber to check on the span of the O<sub>3</sub> analyzer and the efficiency of the NO<sub>x</sub> converters. After an experiment (lamps off), excess NO was injected into the chamber to more than titrate the remaining O<sub>3</sub>.

Condensed phase analysis concentrated on the measurement of sulfate aerosol. The primary method for monitoring sulfate was based on continuous sampling by a flame photometric detector (FPD). Checks were made on this method by taking simultaneous large volume collections on quartz filters and comparing the results. Using this tremendous difference in diffusivity between SO<sub>2</sub> and H<sub>2</sub>SO<sub>4</sub> aerosols, a diffusion-denuder tube was designed to remove virtually all of the SO<sub>2</sub> in the sample entering the FPD, while allowing most of the particulate sulfur to pass to the detector. Optimization between maximum SO<sub>2</sub> removal and minimum H<sub>2</sub>SO<sub>4</sub> loss was determined experimentally by generating independent known concentrations of SO<sub>2</sub> and H<sub>2</sub>SO<sub>4</sub> aerosol in the smog chamber. The optimization work resulted in a sample flow of 200 ml/min through a 35 cm lead acetate diffusion tube, with >99.5% removal of SO<sub>2</sub> and 35% loss of H<sub>2</sub>SO<sub>4</sub> aerosol.

## Kinetic Data

The objective of the first portion of this project was to investigate the kinetic (and mechanisms where feasible) of SO<sub>2</sub> oxidation by free radicals; namely HO, HO<sub>2</sub>, and CH<sub>3</sub>O<sub>2</sub>. The approach taken to study these reactions was to generate a variety of distributions between HO, HO<sub>2</sub>, and CH<sub>3</sub>O<sub>2</sub> in a smog chamber containing SO<sub>2</sub>. This work began with a series of smog chamber experiments using an inorganic smog system (HNO<sub>2</sub>/NO<sub>x</sub>/CO/SO<sub>2</sub>). By selecting the initial concentrations of system components, it was possible to affect a range of HO:HO<sub>2</sub> ratios and thereby observe the relative rates of SO<sub>2</sub> oxidation attributable to the respective radicals.

To obtain estimates of the free radical concentrations, and hence to calculate absolute rate constants, the experimental data were fitted with a chemical kinetics model. Throughout the study the SO<sub>2</sub> oxidation rates related to the subject radicals were determined solely in terms of sulfate aerosol product. Any sulfur products remaining in the gas phase were not determined. As such the rate constants obtained represent rate limiting kinetics for conversion of SO<sub>2</sub> to sulfate and not the upper limit rates for SO<sub>2</sub> removal. In general, good material balances were observed between gas-phase and condensed-phase sulfur, but once the reactions began

was not certain that the gas-phase sulfur was exclusively  $\text{SO}_2$ .

In the inorganic smog system,  $\text{NO}_x$  (i.e.,  $\text{HNO}_2$ ,  $\text{NO}$ , and  $\text{NO}_2$ ) governed the free radical concentrations because their reaction rates with the radicals were typically orders of magnitude larger than those of the corresponding reactions with  $\text{SO}_2$ . Thus, before attempting to interpret experimental data pertaining to  $\text{SO}_2$  oxidation, the  $\text{NO}_x$  chemistry as it occurred in the smog chamber had to be investigated. Substantial concentrations of  $\text{HNO}_2$  were used as a radical source in many of the experiments.

Three initial experiments were conducted to determine the reaction rates for  $\text{HNO}_2$  under experimental conditions and to account for the large  $\text{NO}_x$  losses observed in most experiments. The first experiment showed that the  $\text{NO}$  concentration was largely a function of  $\text{HNO}_2$ 's photolysis rate and the  $\text{HO} + \text{NO}$  recombination rate. Material balances between the model and experimental data were reasonably good, although the model slightly underestimated  $\text{NO}_x$  losses.

The next experiment, conducted to investigate further the smog chamber reactions involving  $\text{HNO}_2$  and the formation of  $\text{HNO}_3$ , investigated two routes for  $\text{HNO}_3$  formation in the gas phase:  $\text{HNO}_2 + \text{O}_3 \rightarrow \text{HNO}_3 + \text{O}_2$  and  $\text{N}_2\text{O}_5 + \text{H}_2\text{O} \rightarrow 2\text{HNO}_3$ . The results suggested that neither reaction produced appreciable amounts. The experiments also provided no conclusive data on the  $\text{HNO}_2$  photolysis rate in the smog chamber. However the choice of  $k = 0.28 \times 10^{-1} \text{min}^{-1}$  seemed to fit these and other experimental data as well as alternative values.

The last of these three experiments attempted to measure the reactions involving  $\text{N}_2\text{O}_5$  formation. Thermal decomposition of  $\text{N}_2\text{O}_5$  apparently limited its concentrations in the troposphere. The slow rate found for homogeneous conversion to  $\text{HNO}_3$  coupled to the strong temperature dependence for  $\text{N}_2\text{O}_5$  decomposition suggested that  $\text{N}_2\text{O}_5$  persists for substantial periods in the colder regions of the lower atmosphere and that clouds and perhaps moist aerosols may be the dominant sink for  $\text{N}_2\text{O}_5$ .

A series of experiments was also conducted in the inorganic smog system to study the conversion of  $\text{SO}_2$  to sulfate via hydroxyl radicals.  $\text{SO}_2$  was added to air mixtures of  $\text{HNO}_2$  and  $\text{NO}_x$  in the chamber. Sulfate aerosol formation was

observed immediately after the lamps were turned on, and the rate of sulfate formation was generally found to maximize early in the irradiation period and then gradually diminish.

The eight experiments were simulated by kinetic modeling in order to estimate the  $\text{HO}$  profiles and assess the overall conversion rate for  $\text{SO}_2$ , as initiated by  $\text{HO}$  radicals.

Some investigators had suggested that  $\text{NO}_x$  could have various roles in the transformation process and even become incorporated in the condensed phase as the reactions reached completion. To investigate these hypotheses, various amounts of  $\text{NO}$  or  $\text{NO}_2$  were added to some of the mixtures, in addition to the amounts produced while charging  $\text{HNO}_2$  to the chamber.

The model gave satisfactory fits to the experimental data, with only one experiment in the series excepted. The model tended to underestimate  $\text{NO}_2$  conversion and sulfate aerosol concentrations, particularly during the latter half of the experimental periods. The discrepancies rarely exceeded 20% of the experimental values, however.

The data supported a rate constant of  $1.6 \times 10^3 \text{ppm}^{-1} \text{min}^{-1}$  for the conversion of  $\text{SO}_2$  to sulfate aerosol, as initiated by the  $\text{HO}$  radical.

The last series of experiments in the inorganic smog system was conducted to determine the kinetics of  $\text{SO}_2$  oxidation by hydroperoxy radicals. To investigate the reactions in the smog chamber, relatively large concentrations of  $\text{CO}$  were added to mixtures of  $\text{NO}_x/\text{HNO}_2/\text{SO}_2$ .  $\text{CO}$  was found to give a negative interference to the analyses of  $\text{SO}_2$  and sulfate aerosols with the FPD. This interference severely limited the range of useful  $\text{CO}$  concentration.

Based on the data from this group of experiments, the selection of an upper limit for the  $\text{SO}_2 + \text{HO}_2 \rightarrow \text{SULFATE}$  reaction was arbitrary. Thus for subsequent modeling,  $0.2 \text{ppm}^{-1} \text{min}^{-1}$  was chosen as a certain upper limit, although the true rate is likely to be much lower.

In the organic smog system, experimenters investigated the methylperoxy- $\text{SO}_2$  reaction beginning with the same approach used for the  $\text{HO}_2$ - $\text{SO}_2$  kinetics studies. The approach consisted of irradiating high concentrations of  $\text{CH}_4$  with controlled  $\text{HNO}_2/\text{NO}_x/\text{SO}_2$  mixtures in the smog chamber. The reaction of principal concern was  $\text{CH}_3\text{O}_2 + \text{SO}_2 \rightarrow \text{CH}_3\text{O} + \text{SULFATE}$ .

Two initial experiments were undertaken to determine the overall rate of

the above reaction based on sulfate aerosol formation in an atmosphere of moderate humidity. Methane was used as the methylperoxy source in the experiments. Although results were inconclusive, an upper limit estimate of  $0.8 \text{ppm}^{-1} \text{min}^{-1}$  was obtained for the conversion of  $\text{SO}_2$  to sulfate via the methylperoxy radical. The mechanism of the conversion was not investigated. Because of the inconclusive results obtained with the  $\text{CH}_4/\text{HNO}_2/\text{NO}_x/\text{SO}_2$  smog system, additional experiments were conducted using azomethane as the methylperoxy source. However, these experiments too were quite limited.

### **Role of $\text{NO}_x$ in $\text{SO}_2$ Oxidation Via $\text{HO}$ Radicals**

Experiments conducted with the  $\text{HNO}_2/\text{NO}_x/\text{SO}_2$  smog system served to investigate the possibility for particulate nitrate formation. At the conclusion of an experiment, sample air was drawn through acid washed and neutralized quartz filters and analyzed for  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ , and  $\text{NH}_4^+$ . To avoid  $\text{HNO}_2$  or  $\text{HNO}_3$  volatilization losses, filter samples were immediately halved; one-half was stored in a desiccatory containing  $\text{NH}_3$ , and the other was refrigerated without treatment.

The analytical results from nine experiments showed no substantial differences in either  $\text{SO}_4^{2-}$  or  $\text{NO}_3^-$  for the  $\text{NH}_3$  treated samples versus the untreated samples. Although ammoniated samples contained somewhat higher  $\text{NH}_4^+$  concentrations, nearly all samples were deficient in  $\text{NH}_3$  when compared to the expected stoichiometry for  $(\text{NH}_4)_2\text{SO}_4$  and  $\text{NH}_4\text{NO}_3$ .

In every case, the particulate nitrate concentrations were very low compared to the sulfate concentrations. The highest particulate nitrate concentrations were recorded for the two experiments without  $\text{SO}_2$ . Thus a substantial amount of nitrate was not incorporated in the particulate phase under the conditions of these experiments. While the results do not rule out the possibility that  $\text{HOSO}_2\text{O}$  interacts with  $\text{NO}_2$  to produce  $\text{H}_2\text{SO}_4$  and  $\text{HNO}_3$ , they do indicate that nitrate (if formed via this route) is not maintained in the aerosol phase, even under the presumably stabilizing influence of  $\text{NH}_3$ .

### **$\text{SO}_2$ Oxidation in Simulated Urban and Rural Atmospheres**

In the second portion of this study, a series of smog chamber experiments

was conducted to investigate some of the factors affecting SO<sub>2</sub> oxidation under simulated atmospheric conditions and to develop data needed to evaluate photochemical models of SO<sub>2</sub> oxidation in polluted air.

Experiments were conducted using hydrocarbon mixtures selected to represent urban and rural conditions. For each experiment, the initial non-methane hydrocarbon concentration was about 4.5 ppmC, and the initial NO<sub>x</sub> concentration was 0.3 ppm. Apart from the control experiments where no SO<sub>2</sub> was introduced, 0.6 ppm SO<sub>2</sub> was injected into the chamber at either 1 or 5 h intervals after the irradiation began. Replicate experiments were also conducted in which 0.15 ppm NO was injected, together with 0.6 ppm SO<sub>2</sub>. This approach was designed to simulate conditions in which SO<sub>2</sub> is emitted into polluted air from point sources. The experimenters attempted to determine whether or not SO<sub>2</sub> oxidation rates depended substantially on the interval of the photochemical smog period, and to what extent the presence of NO (NO is generally emitted along with SO<sub>2</sub>) affected the oxidation rates.

Analytical methods for determining SO<sub>2</sub> and sulfate aerosols were identical to those used in the kinetic experiments. For this experimental series, two Meloy sulfur analyzers were used to monitor SO<sub>2</sub> and particulate sulfate simultaneously.

During the period of these experiments, Battelle's large smog chamber was lined with 5 mil FEP Teflon™ filter. The light intensity corresponding to *k* was 0.17 min<sup>-1</sup>. Gas chromatographic analyses for hydrocarbons were obtained initially and at irradiation intervals of 1, 3, 5, and 7 h. PAN concentrations were obtained hourly.

For both urban and rural hydrocarbon mixtures, the rates of SO<sub>2</sub> oxidation were greatest near the periods of peak NO<sub>2</sub> formation, and gradually diminished thereafter. After 24 h of irradiation, the rate of SO<sub>2</sub> oxidation was about one-fourth as large as the maximum rate observed with the urban hydrocarbon mix: for the rural hydrocarbon mix, the rate of SO<sub>2</sub> oxidation at 24 h was about one-third the maximum rate.

The addition of NO with SO<sub>2</sub> (at SO<sub>2</sub>:NO ratios of 4:1) at different intervals of the irradiation period had mixed results. When added at 5 h, NO had small positive effects on the rates of

SO<sub>2</sub> oxidation, but when added at 1 h, NO reduced the rates of SO<sub>2</sub> oxidation.

### **Kinetic Model Application**

One objective of this portion of the study was to determine if a relatively simple model of atmospheric chemistry could suffice to represent SO<sub>2</sub> oxidation in polluted air. The smog chamber data previously presented and some field data from the Midwest Interstate Sulfur Transformation and Transport (MISTT) project provided the basis for applying the model.

In most cases, the predictions for SO<sub>2</sub> conversion to sulfate were reasonably good; within 20% for the inorganic smog systems and within 25% for the HC-NO<sub>x</sub> systems. The model tended to underestimate SO<sub>2</sub> conversion, and the hydrocarbon systems when the model estimates were only one-half the actual rates. For the smog conditions that were modeled, it was estimated that at least 75% and sometimes greater than 90% of the SO<sub>2</sub> oxidation was initiated by reactions of SO<sub>2</sub> with HO radicals.

Although several shortcomings of the lumped model were found during its application to these data, it was still of interest to adapt the model to estimate diurnal patterns of SO<sub>2</sub> oxidation in comparison with field data. The objective of this exercise was to estimate the rate of SO<sub>2</sub> oxidation for power plant plumes resident in polluted atmospheres, as a function of solar radiation, plume emission periods, and plume dispersion rates.

According to the modeling results, the production of HO and the oxidation of SO<sub>2</sub> depended strongly on solar radiation intensity. Reduced radiation resulted in disproportionately lower rates of SO<sub>2</sub> oxidation. For reduced radiation conditions, the maximum rates of SO<sub>2</sub> oxidation were predicted to occur near mid-afternoon, while for clear sky conditions, the maximum oxidation rates occurred between 11:00 a.m. and noon. The rates of SO<sub>2</sub> oxidation were shown to depend on ambient HC and NO<sub>x</sub> conditions. For the range of conditions modeled, the maximum rate of SO<sub>2</sub> oxidation (clear sky) varied from 3.7 to 7.4%/h.

Plume dispersion rates affected the induction period of NO and SO<sub>2</sub> oxidation. Variations in solar radiation intensity and thus the diurnal period of the power plant emissions strongly affected SO<sub>2</sub> oxidation rates. For well-dispersed plumes and clear sky condi-

tions, maximum SO<sub>2</sub> oxidation rate (ranging from 2.7 to 6.9%/h) were estimated to occur during the noon to early afternoon periods. The patterns of SO<sub>2</sub> oxidation predicted by the model agreed well with the data for the Labadie plume as reported in the MISTT project.

For relatively low HC:NO<sub>x</sub> ratios for ambient air, SO<sub>2</sub> oxidation rates computed for plumes were bound by the ambient rates of SO<sub>2</sub> oxidation. At high HC:NO<sub>x</sub> ratios, the rates of SO<sub>2</sub> oxidation in plumes were predicted to exceed the rates in ambient air. Under such conditions, the model also predicted an excess (or bulge) in the plume O concentrations. These features have been observed in actual plume studies

### **Conclusions and Recommendations**

Regarding reaction 60 (SO<sub>2</sub> + OH + M → SULFATE), it appears that all of the SO<sub>2</sub> oxidized by this route results in the formation of sulfate particles. According to the analyses for particulate nitrate, they did not form stable aerosol products with the intermediates created after HO addition to SO<sub>2</sub>.

In the course of the experiment with inorganic smog systems (HNO<sub>2</sub>/NO<sub>x</sub>/SO<sub>2</sub>), additional kinetic information was obtained, particularly as it pertained to the experimental system. An upper limit rate of 3 × 10<sup>-7</sup> ppm<sup>-1</sup>min<sup>-1</sup> was derived for the homogeneous reaction between dinitrogen pentoxide and water vapor to yield nitric acid (N<sub>2</sub>O<sub>5</sub> + H<sub>2</sub>O → HNO<sub>3</sub>). It was also determined that ozone and nitrous acid do not react at a significant rate to produce nitric acid. No heterogeneous reaction between SO<sub>2</sub> and nitrous acid in the presence of sulfuric acid aerosols could be detected.

Although the average reactivity for the urban and rural hydrocarbon mixtures, based on rate constants of individual hydrocarbons with HO radicals, was nearly equal, there were marked differences in several important smog parameters. With the urban hydrocarbon mixture, the rates of NO oxidation and xylene disappearance and the maximum rates of SO<sub>2</sub> oxidation were about twice as great as for the rural hydrocarbon mixture. The formation of PAN and other products of NO<sub>2</sub> oxidation were also greater for the experiments with urban hydrocarbon.

Since SO<sub>2</sub> oxidation is dominated by the reaction with hydroxyl radicals, and since this initial reaction is the rate-

limiting step in SO<sub>2</sub>'s conversion to sulfate, it follows that relatively simple kinetic models which adequately describe the variations in atmospheric HO should suffice to estimate gas phase SO<sub>2</sub> transformation rates.

Experimental and theoretical simulations of urban and rural atmospheres showed that the rates of SO<sub>2</sub> oxidation are strongly coupled to solar radiation intensity and pollution conditions. Although high concentrations of NO generally suppressed or delayed HO production and SO<sub>2</sub> oxidation, NO<sub>2</sub> inhibited SO<sub>2</sub> oxidation by scavenging HO. Aldehydes, CO, and possibly some hydrocarbons tended to inhibit SO<sub>2</sub> oxidation by converting HO into products which were less efficient in oxidizing SO<sub>2</sub>. In general, reactive hydrocarbons served to promote SO<sub>2</sub> oxidation, and the stoichiometry or net flux of HO affected by hydrocarbon degradation appeared to be key to the acceleration of SO<sub>2</sub> oxidation in polluted air.

SO<sub>2</sub> oxidation rates in moderately polluted air or in well-aged air that was once polluted were found for the most part to be 60 to 80% lower than the rates in freshly polluted air.

Modeling results indicated that SO<sub>2</sub> oxidation rates in power plant plumes are highly dependent on plume dispersion rates and are coupled to the same parameters governing SO<sub>2</sub> oxidation in the ambient air. For well-dispersed plumes, most rapid SO<sub>2</sub> conversion was predicted to occur from noon through early afternoon. The accompaniment of NO in plumes generally suppressed HO and the rate of SO<sub>2</sub> oxidation. However, under certain circumstances where high concentrations of reactive HC were mixed into a rapidly-dispersing plume, the rate of SO<sub>2</sub> oxidation was expected to exceed the oxidation rate in ambient air. Under such circumstances, an excess of ozone was also predicted in the plume.

This study pointed out the need for further study in the area of sulfate aerosol formation. In particular, since the overall conversion rate for the SO<sub>2</sub> + HO<sub>2</sub> → SULFATE reaction was considerably less in this study than the literature value, it is recommended that additional high-pressure measurements of the reaction be made. Any such effort should include the determination of reaction products.

*David F. Miller is with Battelle-Columbus Laboratories, Columbus, OH 43201. Joseph J. Bufalini is the EPA Project Officer (see below).*

*The complete report, entitled "Modeling of SO<sub>2</sub> Oxidation in Smog," (Order No. PB 82-101 932; Cost: \$13.50, subject to change) will be available only from:*

*National Technical Information Service*

*5285 Port Royal Road*

*Springfield, VA 22161*

*Telephone: 703-487-4650*

*The EPA Project Officer can be contacted at:*

*Environmental Sciences Research Laboratory*

*U.S. Environmental Protection Agency*

*Research Triangle Park, NC 27711*

United States  
Environmental Protection  
Agency

Center for Environmental Research  
Information  
Cincinnati OH 45268

Postage and  
Fees Paid  
Environmental  
Protection  
Agency  
EPA 335



---

Official Business  
Penalty for Private Use \$300

• PS 0000329 •  
U S ENVIR PROTECTION AGENCY  
REGION 5 LIBRARY  
230 S DEARBORN STREET  
CHICAGO IL 60604  
•