


7281  
no. 343

905R76010

**U.S. ENVIRONMENTAL PROTECTION AGENCY  
NATIONAL EUTROPHICATION SURVEY  
WORKING PAPER SERIES**



REPORT  
ON  
VERSAILLES LAKE  
RIPLEY COUNTY  
INDIANA  
EPA REGION V  
WORKING PAPER No. 343

**CORVALLIS ENVIRONMENTAL RESEARCH LABORATORY - CORVALLIS, OREGON  
and  
ENVIRONMENTAL MONITORING & SUPPORT LABORATORY - LAS VEGAS, NEVADA**

## CONTENTS

	<u>Page</u>
Foreword	ii
List of Indiana Study Lakes	iv
Lake and Drainage Area Map	v, vi
 <u>Sections</u>	
I. Conclusions	1
II. Lake and Drainage Basin Characteristics	4
III. Lake Water Quality Summary	5
IV. Nutrient Loadings	9
V. Literature Reviewed	14
VI. Appendices	15

## F O R E W O R D

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nationwide threat of accelerated eutrophication to fresh water lakes and reservoirs.

### OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

### ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

- a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.
- b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.
- c. With such a transformation, an assessment of the potential for eutrophication control can be made.

### LAKE ANALYSIS

In this report, the first stage of evaluation of lake and watershed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning {§303(e)}, water quality criteria/standards review {§303(c)}, clean lakes {§314(a,b)}, and water quality monitoring {§106 and §305(b)} activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

ACKNOWLEDGMENT

The staff of the National Eutrophication Survey (Office of Research & Development, U. S. Environmental Protection Agency) expresses sincere appreciation to the Indiana State Board of Health for professional involvement, to the Indiana National Guard for conducting the tributary sampling phase of the Survey, and to those Indiana wastewater treatment plant operators who provided effluent samples and flow data.

The staff of the Division of Water Pollution Control, Indiana State Board of Health, provided invaluable lake documentation and counsel during the Survey, reviewed the preliminary reports, and provided critiques most useful in the preparation of this Working Paper series.


Major General Alfred F. Ahner, Adjutant General of Indiana, and Project Officers Lt. Colonel Charles B. Roberts (Retired) and Colonel Robert L. Sharp, who directed the volunteer efforts of the Indiana National Guardsmen, are also gratefully acknowledged for their assistance to the Survey.


## NATIONAL EUTROPHICATION SURVEY

## STUDY LAKES

STATE OF INDIANA

<u>LAKE NAME</u>	<u>COUNTY</u>
Bass	Starke
Cataract	Owen, Putnam
Crooked	Steuben
Dallas	LaGrange
Geist	Hamilton, Marion
Hamilton	Steuben
Hovey	Posey
James	Kosciusko
James	Steuben
Long	Steuben
Marsh	Steuben
Mississinewa	Grant, Miami, Wabash
Maxinkuckee	Marshall
Monroe	Brown, Monroe
Morse	Hamilton
Olin	LaGrange
Oliver	LaGrange
Pigeon	Steuben
Sylvan	Noble
Tippecanoe	Kosciusko
Versailles	Ripley
Wawassee	Kosciusko
Webster	Kosciusko
Westler	LaGrange
Whitewater	Union
Winona	Kosciusko
Witmer	LaGrange





VERSAILLES LAKE

STORET NO. 1850

I. CONCLUSIONS

A. Trophic Condition:

Survey data indicate that Versailles Lake is eutrophic. It ranked twenty-third in overall trophic quality when the 27 Indiana lakes sampled in 1973 were compared using a combination of six parameters\*. Twenty-four of the lakes had less median total phosphorus, 22 had less median dissolved phosphorus, 18 had less median inorganic nitrogen, 19 had less mean chlorophyll a, and 25 had greater mean Secchi disc transparency. Near-depletion of dissolved oxygen with depth occurred at sampling station 3 in August and October.

Survey limnologists reported rooted aquatic vegetation along about 20% of the shoreline in August and October and some algae along the shore near station 1 in October.

B. Rate-Limiting Nutrient:

The algal assay results indicate that Versailles Lake was phosphorus limited at the time the sample was collected (04/29/73). The lake data indicate phosphorus limitation at the other sampling times as well.

C. Nutrient Controllability:

1. Point sources--The phosphorus contributions of known point sources amounted to 42.9% of the total reaching Versailles Lake

\* See Appendix A.

during the sampling year. The communities of Ba  
Osgood (2.7%), and Napoleon (0.5%) were the sour  
loads. However, the non-point phosphorus export  
Creek was substantially higher than would be exp  
12). It is possible that additional point sour  
ville or Osgood municipal areas contributed sig  
phosphorus load, the load from Napoleon may hav  
or both.

The present phosphorus loading of 44.42 g/r  
that proposed by Vollenweider (Vollenweider an  
a eutrophic loading (see page 13). However, t  
retention time of the lake probably is quite s  
and Vollenweider's model may not apply.

A 100% reduction in the phosphorus loads f  
sources would lower the overall loading rate t  
Because Versailles Lake is phosphorus limited,  
inputs should be minimized to the greatest pr  
slow the aging of this water body.

2. Non-point sources--About 57% of the t  
to Versailles Lake came from non-point source  
year. Laughery Creek contributed 55.4%; howe  
above, the phosphorus export rate of 58 kg/kr  
higher than the rates of other tributaries i

This may have been due in part to unknown point sources rather than to non-point source inputs. Other contributing gaged tributaries were Cedar Creek, 0.7%; Unnamed Stream C-1, 0.5%; and Falling Timber Creek, 0.3%. The ungaged tributaries were estimated to have contributed 0.2% of the overall phosphorus load.

## II. LAKE AND DRAINAGE BASIN CHARACTERISTICS<sup>†</sup>

### A. Lake Morphometry<sup>††</sup>:

1. Surface area: 0.93 kilometers<sup>2</sup>.
2. Mean depth: Unknown.
3. Maximum depth: 9.1 meters.
4. Volume: Unknown.
5. Mean hydraulic retention time: Unknown but probably is quite short; if the mean depth is equal to the maximum depth, the retention time would only be 24 days.

### B. Tributary and Outlet: (See Appendix C for flow data)

#### 1. Tributaries -

<u>Name</u>	<u>Drainage area (km<sup>2</sup>)*</u>	<u>Mean flow (m<sup>3</sup>/sec)*</u>
Laughery Creek	396.5	3.78
Falling Timber Creek	15.5	0.14
Unnamed Stream C-1	7.8	0.07
Cedar Creek	11.9	0.11
Minor tributaries & immediate drainage -	<u>4.1</u>	<u>0.02</u>
Totals	435.8	4.12

#### 2. Outlet -

Laughery Creek	436.7**	4.12
----------------	---------	------

### C. Precipitation\*\*\*:

1. Year of sampling: 128.3 centimeters.
2. Mean annual: 105.4 centimeters.

<sup>†</sup> Table of metric conversions--Appendix B.

<sup>††</sup> Winters, 1975.

<sup>\*</sup> For limits of accuracy, see Working Paper No. 175, "...Survey Methods, 1973-1976".

<sup>\*\*</sup> Includes area of lake.

<sup>\*\*\*</sup> See Working Paper No. 175.

### III. LAKE WATER QUALITY SUMMARY

Versailles Lake was sampled three times during the open-water season of 1973 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from a number of depths at three stations on the lake (see map, page v). During each visit, a single depth-integrated (4.6 m or near bottom to surface) sample was composited from the stations for phytoplankton identification and enumeration; and during the first visit, a single 18.9-liter depth-integrated sample was composited for algal assays. Also each time, a depth-integrated sample was collected from each of the stations for chlorophyll a analysis. The maximum depths sampled were 0.9 meters at station 1, 1.5 meters at station 2, and 4.6 meters at station 3.

The sampling results are presented in full in Appendix D and are summarized in the following table.

A. SUMMARY OF PHYSICAL AND CHEMICAL CHARACTERISTICS FOR VERSAILLES LAKE  
STORE CODE 1850

PARAMETER	1ST SAMPLING ( 4/29/73)			2ND SAMPLING ( 8/ 2/73)			3RD SAMPLING (10/10/73)		
	3 SITES			3 SITES			3 SITES		
	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN
TEMP (C)	13.5 - 14.3	14.0	14.0	22.8 - 24.9	24.4	24.7	21.2 - 22.4	21.7	21.6
DISS OXY (MG/L)	6.5 - 8.8	7.7	7.7	0.5 - 8.6	5.7	6.8	0.5 - 9.2	4.8	4.8
CNDCTVY (MICROM)	300. - 350.	324.	330.	245. - 324.	279.	271.	309. - 323.	315.	315.
pH (STAND UNITS)	7.8 - 8.0	7.9	7.9	7.3 - 8.4	7.9	8.2	7.2 - 8.2	7.7	7.8
TOT ALK (MG/L)	118. - 139.	127.	126.	98. - 128.	110.	109.	136. - 146.	140.	139.
TOT P (MG/L)	0.173 - 0.230	0.194	0.193	0.087 - 0.180	0.118	0.115	0.089 - 0.139	0.108	0.108
ORTHO P (MG/L)	0.054 - 0.070	0.063	0.064	0.010 - 0.041	0.017	0.011	0.010 - 0.019	0.014	0.013
NO2+NO3 (MG/L)	1.000 - 1.300	1.129	1.100	0.750 - 0.850	0.802	0.810	0.030 - 0.050	0.042	0.040
AMMONIA (MG/L)	0.100 - 0.180	0.137	0.120	0.060 - 0.390	0.174	0.100	0.050 - 0.800	0.326	0.220
KJEL N (MG/L)	0.700 - 1.200	0.914	0.900	1.100 - 1.600	1.320	1.200	1.300 - 2.000	1.600	1.500
INORG N (MG/L)	1.160 - 1.420	1.266	1.220	0.830 - 1.140	0.976	0.930	0.080 - 0.840	0.368	0.260
TOTAL N (MG/L)	1.700 - 2.500	2.043	1.900	1.910 - 2.450	2.122	1.970	1.340 - 2.050	1.642	1.540
CHLRPYL A (UG/L)	3.2 - 19.8	11.0	9.9	4.9 - 64.7	32.9	29.1	22.6 - 41.3	31.4	30.2
SECCHI (METERS)	0.2 - 0.3	0.2	0.2	0.2 - 0.6	0.4	0.5	0.5 - 0.8	0.6	0.7

## B. Biological characteristics:

## 1. Phytoplankton -

<u>Sampling Date</u>	<u>Dominant Genera</u>	<u>Algal Units per ml</u>
04/29/73	1. <u>Cryptomonas sp.</u> 2. <u>Oscillatoria sp.</u> 3. Centric diatoms 4. <u>Synedra sp.</u> 5. <u>Navicula sp.</u>	1,292 775 52 34 <u>17</u>
		Total 2,170
08/02/73	1. <u>Stephanodiscus sp.</u> 2. <u>Chroomonas sp.</u> 3. Flagellates 4. <u>Chlorogonium sp.</u> 5. <u>Melosira sp.</u> Other genera	4,946 1,371 979 539 196 <u>1,763</u>
		Total 9,794
10/10/73	1. <u>Oscillatoria sp.</u> 2. <u>Melosira sp.</u> 3. Centric diatoms 4. <u>Merismopedia sp.</u> 5. <u>Cryptomonas sp.</u> Other genera	10,228 3,133 2,580 1,843 1,659 <u>4,607</u>
		Total 24,050

## 2. Chlorophyll a -

<u>Sampling Date</u>	<u>Station Number</u>	<u>Chlorophyll a (<math>\mu\text{g/l}</math>)</u>
04/29/73	1	3.2
	2	9.9
	3	19.8
08/02/73	1	64.7
	2	29.1
	3	4.9
10/10/73	1	30.2
	2	41.3
	3	22.6

## C. Limiting Nutrient Study:

## 1. Autoclaved, filtered, and nutrient spiked -

<u>Spike (mg/l)</u>	<u>Ortho P Conc. (mg/l)</u>	<u>Inorganic N Conc. (mg/l)</u>	<u>Maximum yield (mg/l-dry wt.)</u>
Control	0.050	1.280	19.7
0.050 P	0.100	1.280	22.6
0.050 P + 1.0 N	0.100	2.280	29.0
1.0 N	0.050	2.280	21.8

## 2. Filtered, and nutrient spiked -

<u>Spike (mg/l)</u>	<u>Ortho P Conc. (mg/l)</u>	<u>Inorganic N Conc. (mg/l)</u>	<u>Maximum yield (mg/l-dry wt.)</u>
Control	0.035	1.185	10.9
0.050 P	0.085	1.185	27.5
0.050 P + 1.0 N	0.085	2.185	30.3
1.0 N	0.035	2.185	13.2

## 3. Discussion -

The control yields of the assay alga, Selenastrum capricornutum, indicate that the potential primary productivity of Versailles Lake was high at the time the sample was collected (04/29/73). Both the autoclaved-filtered and filtered-only assays showed significant increases in yield with the addition of orthophosphorus alone, although smaller increases than would be expected. Note that the addition of nitrogen alone did not result in significantly increased yields as compared to the control; hence, phosphorus limitation is indicated.

The lake data indicate phosphorus limitation at all sampling times; i.e., the mean inorganic nitrogen/orthophosphorus ratios were 20/1 or greater.

IV. NUTRIENT LOADINGS  
(See Appendix E for data)

For the determination of nutrient loadings, the Indiana National Guard collected monthly near-surface grab samples from each of the tributary sites indicated on the map (page vi), except for the high runoff months of January and February when two samples were collected. Sampling was begun in June, 1973, and was completed in May, 1974.

Through an interagency agreement, stream flow estimates for the year of sampling and a "normalized" or average year were provided by the Indiana District Office of the U.S. Geological Survey for the tributary sites nearest the lake.

In this report, nutrient loads for sampled tributaries were determined by using a modification of a U.S. Geological Survey computer program for calculating stream loadings\*. Nutrient loads shown are those measured minus point-source loads, if any.

Nutrient loads for unsampled "minor tributaries and immediate drainage" ("ZZ" of U.S.G.S.) were estimated using the means of the nutrient loads, in kg/km<sup>2</sup>/year, at stations B-1, C-1, and D-1 and multiplying the means by the ZZ area in km<sup>2</sup>.

The operators of the Osgood and Batesville wastewater treatment plants provided monthly effluent samples and corresponding flow data. The community of Napoleon did not participate in the Survey; nutrient loads from this source were estimated at 0.567 kg P and 3.401 kg N/capita/year, and flows were estimated at 0.3785 m<sup>3</sup>/capita/day.

\* See Working Paper No. 175.

All estimated phosphorus loads were reduced by 50% to adjust for a phosphate detergent limitation in effect in Indiana since January, 1972.

A. Waste Sources:

1. Known municipal<sup>†</sup> -

<u>Name</u>	<u>Pop. Served</u>	<u>Treatment</u>	<u>Mean Flow (m<sup>3</sup>/d)</u>	<u>Receiving Water</u>
Batesville #1		tr. filter	1,097.8	Little Laughery Creek
Batesville #2	} 4,000	act. sludge	1,047.3	Little Laughery Creek
Osgood	1,500	tr. filter + P-removal*	662.4	Branch of Plum Creek
Napoleon	400**	stab. pond	151.4	Laughery Creek

2. Known industrial - None

<sup>†</sup> Treatment plant questionnaires.

\* Begun in November, 1974.

\*\* Anonymous, 1971.

## B. Annual Total Phosphorus Loading - Average Year:

## 1. Inputs -

<u>Source</u>	<u>kg P/ yr</u>	<u>% of total</u>
a. Tributaries (non-point load) -		
Laughery Creek	22,875	55.4
Falling Timber Creek	140	0.3
Unnamed Stream C-1	195	0.5
Cedar Creek	285	0.7
b. Minor tributaries & immediate drainage (non-point load) -	80	0.2
c. Known municipal STP's -		
Batesville #1	9,920	24.0
Batesville #2	6,475	15.7
Osgood	1,100	2.7
Napoleon	225	0.5
d. Septic tanks* -	<5	<0.1
e. Known industrial - None	-	-
f. Direct precipitation** -	<u>15</u>	<u>&lt;0.1</u>
Total	41,315	100.0

## 2. Outputs -

Lake outlet - Laughery Creek 28,700

3. Net annual P accumulation - 12,615 kg.

\* Estimate based on two camps and one park; see Working Paper No. 175.

\*\* See Working Paper No. 175.

## C. Annual Total Nitrogen Loading - Average Year:

## 1. Inputs -

<u>Source</u>	<u>kg N/ yr</u>	<u>% of total</u>
a. Tributaries (non-point load) -		
Laughery Creek	302,320	89.9
Falling Timber Creek	4,785	1.4
Unnamed Stream C-1	3,690	1.1
Cedar Creek	7,445	2.2
b. Minor tributaries & immediate drainage (non-point load) -	1,925	0.6
c. Known municipal STP's -		
Batesville #1	5,430	1.6
Batesville #2	4,215	1.3
Osgood	3,875	1.2
Napoleon	1,360	0.4
d. Septic tanks* -	175	<0.1
e. Known industrial - None	-	-
f. Direct precipitation** -	<u>1,005</u>	<u>0.3</u>
Total	336,225	100.0

## 2. Outputs -

Lake outlet - Laughery Creek 353,070

3. Net annual N loss - 16,845 kg.

## D. Mean Annual Non-point Nutrient Export by Subdrainage Area:

<u>Tributary</u>	<u>kg P/km<sup>2</sup>/yr</u>	<u>kg N/km<sup>2</sup>/yr</u>
Laughery Creek	58	762
Falling Timber Creek	9	309
Unnamed Stream C-1	25	473
Cedar Creek	24	626

\* Estimate based on two camps and one park; see Working Paper No. 175.

\*\* See Working Paper No. 175.

E. Yearly Loads:

In the following table, the existing phosphorus loadings are compared to those proposed by Vollenweider (Vollenweider and Dillon, 1974). Essentially, his "dangerous" loading is one at which the receiving water would become eutrophic or remain eutrophic; his "permissible" loading is that which would result in the receiving water remaining oligotrophic or becoming oligotrophic if morphometry permitted. A mesotrophic loading would be considered one between "dangerous" and "permissible".

Note that Vollenweider's model may not be applicable to water bodies with short hydraulic retention times.

	Total Phosphorus Total	Total Phosphorus Accumulated	Total Nitrogen Total	Total Nitrogen Accumulated
grams/m <sup>2</sup> /yr	44.42	13.56	361.5	loss*

Vollenweider phosphorus loadings  
(g/m<sup>2</sup>/yr) based on surface area and mean  
outflow of Versailles Lake:

"Dangerous" (eutrophic loading)	2.20
"Permissible" (oligotrophic loading)	1.10

\* There was an apparent nitrogen loss during the sampling year. This may have been due to nitrogen fixation in the lake, solubilization of previously sedimented nitrogen, recharge with nitrogen-rich ground water, unknown and unsampled point sources discharging directly to the lake, or (probably) insufficient sampling in relation to the short hydraulic retention time of the lake (see page 4). Whatever the cause, a similar nitrogen loss has occurred at Shagawa Lake, Minnesota, which has been intensively studied by EPA's former National Eutrophication and Lake Restoration Branch (Malueg et al., 1975).

## V. LITERATURE REVIEWED

Anonymous, 1971. Inventory of municipal waste facilities. EPA Publ. OWP-1, vol. 5, Wash., DC.

Crayer, John W., 1973. Treatment plant questionnaire (Osgood STP). Osgood Sanitation Dept., Osgood.

Flodder, Art, 1973. Treatment plant questionnaires (Batesville STP's). Batesville.

Malueg, Kenneth W., D. Phillips Larsen, Donald W. Schults, and Howard T. Mercier; 1975. A six-year water, phosphorus, and nitrogen budget for Shagawa Lake, Minnesota. Jour. Environ. Qual., vol. 4, no. 2, pp. 236-242.

Vollenweider, R. A., and P. J. Dillon, 1974. The application of the phosphorus loading concept to eutrophication research. Natl. Res. Council of Canada Publ. No. 13690, Canada Centre for Inland Waters, Burlington, Ontario.

Winters, John, 1975. Personal communication (lake morphometry). IN Div. of Water Poll. Contr., Indianapolis.

VI. APPENDICES

APPENDIX A

LAKE RANKINGS

## LAKE DATA TO BE USED IN RANKINGS

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500-MEAN SEC	MEAN CHLORA	15-MIN DU	DISS ORTHO P	MEDIAN P
1805	CATARACT LAKE	0.058	1.660	466.667	10.744	15.000	0.013	
1811	GEIST RESERVOIR	0.074	1.080	472.500	45.950	11.600	0.009	
1817	JAMES LAKE	0.024	1.030	434.000	11.533	15.000	0.008	
1827	MISSISSINNEWA RESERVOIR	0.107	2.400	473.444	15.778	15.000	0.029	
1828	MONROE RESERVOIR	0.025	0.325	436.823	6.947	15.000	0.007	
1829	MORSE RESERVOIR	0.084	3.325	473.222	56.167	15.000	0.009	
1836	WAHSEE LAKE	0.012	0.210	364.500	5.000	14.600	0.003	
1837	WEBSTER LAKE	0.025	0.790	431.000	11.500	15.000	0.005	
1839	WHITEWATER LAKE	0.084	1.620	470.167	33.083	15.000	0.012	
1840	WINONA LAKE	0.035	1.250	444.667	11.211	15.000	0.011	
1841	WESTLER LAKE	0.035	0.860	427.125	10.712	15.000	0.013	
1842	WITMER LAKE	0.035	0.300	440.333	11.917	15.000	0.011	
1843	LAKE MAXINKUCKEE	0.050	0.220	400.400	5.483	15.000	0.003	
1844	TIPPECANOE LAKE	0.019	0.195	391.500	6.050	15.000	0.005	
1845	DALLAS LAKE	0.029	0.830	413.333	10.067	15.000	0.014	
1846	OLIN LAKE	0.012	1.460	403.333	4.867	14.900	0.003	
1847	OLIVER LAKE	0.009	0.920	392.000	3.767	14.800	0.004	
1848	SYLVAN LAKE	0.170	0.130	469.833	47.480	14.800	0.017	
1849	HOVEY LAKE	0.062	1.050	489.333	84.267	7.600	0.024	
1850	VERSAILLES LAKE	0.139	1.090	482.000	25.078	14.500	0.019	
1851	BASS LAKE	0.040	0.250	471.375	29.367	7.000	0.012	
1852	CROOKED LAKE	0.019	0.120	410.111	5.578	15.000	0.005	
1853	LAKE JAMES	0.016	0.190	352.444	4.856	15.000	0.005	
1854	LONG LAKE	0.204	1.920	442.667	16.100	15.000	0.150	
1855	PIGEON LAKE	0.058	1.945	442.667	11.900	15.000	0.015	
1856	MARSH LAKE	0.093	0.270	451.333	34.467	15.000	0.055	
1857	HAMILTON LAKE	0.033	0.720	413.167	17.450	15.000	0.018	

## PERCENT OF LAKES WITH HIGHER VALUES (NUMBER OF LAKES WITH HIGHER VALUES)

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500-MEAN SEC	MEAN CHLORA	15-MIN DO	MEDIAN DISS ORTHO P	INDEX NO
1805	CATARACT LAKE	37 ( 9)	15 ( 4)	31 ( 8)	62 ( 16)	35 ( 0)	37 ( 9)	217
1811	GEIST RESERVOIR	27 ( 7)	35 ( 9)	15 ( 4)	12 ( 3)	92 ( 24)	62 ( 16)	243
1817	JAMES LAKE	73 ( 19)	42 ( 11)	58 ( 15)	50 ( 13)	35 ( 0)	65 ( 17)	323
1827	MISSISSINNEWA RESERVOIR	12 ( 3)	4 ( 1)	8 ( 2)	38 ( 10)	35 ( 0)	8 ( 2)	105
1828	MONROE RESERVOIR	67 ( 17)	69 ( 18)	54 ( 14)	73 ( 19)	35 ( 0)	69 ( 18)	367
1829	MORSE RESERVOIR	23 ( 6)	0 ( 0)	12 ( 3)	4 ( 1)	35 ( 0)	58 ( 15)	132
1836	WAWASEE LAKE	94 ( 24)	85 ( 22)	96 ( 25)	88 ( 23)	85 ( 22)	98 ( 25)	546
1837	WEBSTER LAKE	67 ( 17)	62 ( 16)	62 ( 16)	54 ( 14)	35 ( 0)	81 ( 21)	361
1839	WHITEWATER LAKE	19 ( 5)	19 ( 5)	23 ( 6)	19 ( 5)	35 ( 0)	42 ( 11)	157
1840	WINONA LAKE	50 ( 12)	27 ( 7)	38 ( 10)	58 ( 15)	35 ( 0)	52 ( 13)	260
1841	WESTLER LAKE	50 ( 12)	54 ( 14)	65 ( 17)	65 ( 17)	35 ( 0)	37 ( 9)	306
1842	WITMER LAKE	50 ( 12)	50 ( 13)	50 ( 13)	42 ( 11)	35 ( 0)	52 ( 13)	279
1843	LAKE MAXINKUCKEE	77 ( 20)	81 ( 21)	85 ( 22)	85 ( 22)	35 ( 0)	98 ( 25)	461
1844	TIPPECANOE LAKE	85 ( 22)	88 ( 23)	92 ( 24)	77 ( 20)	35 ( 0)	85 ( 22)	462
1845	DALLAS LAKE	62 ( 16)	58 ( 15)	69 ( 18)	69 ( 18)	35 ( 0)	31 ( 8)	324
1846	OLIN LAKE	94 ( 24)	23 ( 6)	81 ( 21)	92 ( 24)	73 ( 19)	92 ( 24)	455
1847	OLIVER LAKE	100 ( 26)	46 ( 12)	88 ( 23)	100 ( 26)	79 ( 20)	88 ( 23)	501
1848	SYLVAN LAKE	4 ( 1)	96 ( 25)	27 ( 7)	8 ( 2)	79 ( 20)	23 ( 6)	237
1849	HOVEY LAKE	31 ( 8)	38 ( 10)	0 ( 0)	0 ( 0)	96 ( 25)	12 ( 3)	177
1850	VERSAILLES LAKE	8 ( 2)	31 ( 8)	4 ( 1)	27 ( 7)	88 ( 23)	15 ( 4)	173
1851	BASS LAKE	42 ( 11)	77 ( 20)	19 ( 5)	23 ( 6)	100 ( 26)	46 ( 12)	307
1852	CROOKED LAKE	81 ( 21)	100 ( 26)	77 ( 20)	81 ( 21)	35 ( 0)	75 ( 19)	449
1853	LAKE JAMES	88 ( 23)	92 ( 24)	100 ( 26)	96 ( 25)	35 ( 0)	75 ( 19)	486
1854	LONG LAKE	0 ( 0)	12 ( 3)	44 ( 11)	35 ( 9)	35 ( 0)	0 ( 0)	126
1855	PIGEON LAKE	37 ( 9)	8 ( 2)	44 ( 11)	46 ( 12)	35 ( 0)	27 ( 7)	197
1856	MARSH LAKE	15 ( 4)	73 ( 19)	35 ( 9)	15 ( 4)	35 ( 0)	4 ( 1)	177
1857	HAMILTON LAKE	58 ( 15)	65 ( 17)	73 ( 19)	31 ( 8)	35 ( 0)	19 ( 5)	281

## LAKES RANKED BY INDEX NOS.

RANK	LAKE CODE	LAKE NAME	INDEX NO
1	1836	WANASEE LAKE	546
2	1847	OLIVER LAKE	501
3	1853	LAKE JAMES	486
4	1844	TIPPECANOE LAKE	462
5	1843	LAKE MAXINKUCKEE	461
6	1846	OLIN LAKE	455
7	1852	CROOKED LAKE	449
8	1828	MONROE RESERVOIR	367
9	1837	WEBSTER LAKE	361
10	1845	DALLAS LAKE	324
11	1817	JAMES LAKE	323
12	1851	BASS LAKE	307
13	1841	WESTLER LAKE	306
14	1857	HAMILTON LAKE	281
15	1842	WITHER LAKE	279
16	1840	WINONA LAKE	260
17	1811	GEIST RESERVOIR	243
18	1848	SYLVAN LAKE	237
19	1805	CATARACT LAKE	217
20	1855	PIGEON LAKE	197
21	1856	MARSH LAKE	177
22	1849	HOVEY LAKE	177
23	1850	VERSAILLES LAKE	173
24	1839	WHITEWATER LAKE	157
25	1829	MORSE RESERVOIR	132
26	1854	LONG LAKE	126
27	1827	MISSISSINNEWA RESERVOIR	105

## **APPENDIX B**

### **CONVERSION FACTORS**

## CONVERSION FACTORS

Hectares x 2.471 = acres

Kilometers x 0.6214 = miles

Meters x 3.281 = feet

Cubic meters x  $8.107 \times 10^{-4}$  = acre/feet

Square kilometers x 0.3861 = square miles

Cubic meters/sec x 35.315 = cubic feet/sec

Centimeters x 0.3937 = inches

Kilograms x 2.205 = pounds

Kilograms/square kilometer x 5.711 = lbs/square mile

## APPENDIX C

### TRIBUTARY FLOW DATA

SUBMITTAL FORM INFORMATION FOR INDIANA

03/29/16

VERBALS | 185

TOTAL DRAINAGE AREA OF LAKE (SQ KM) 434.9

TRANSITORY STATE INFORMATION FOR INDIANA

1/30/20

SUB-DRAINAGE AREA (SQ KM)	TRIBUTARY	NORMALIZED FLOWS (CMS)						NOV	DEC	MEAN
		JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	
436.7	1850A1	5.72	6.37	8.69	8.30	5.47	4.05	2.53	1.11	0.95
396.5	1850A2	5.18	5.78	7.90	7.53	4.96	3.68	2.29	1.00	0.86
15.5	1850B1	0.178	0.221	0.320	0.297	0.190	0.147	0.088	0.026	0.027
7.0	1850C1	0.088	0.110	0.161	0.150	0.093	0.074	0.042	0.012	0.013
11.9	1850D1	0.136	0.170	0.246	0.227	0.144	0.113	0.065	0.020	0.020
2.2	1850Z2	0.023	0.031	0.045	0.042	0.026	0.021	0.012	0.003	0.003

### TOTAL DRAINAGE AREA OR SUM OF SUB-DRAINAGE AREAS

TOTAL DRAINAGE AREA OF LAKE = 434.9  
SUM OF SUB-DRAINAGE AREAS = 433.9

7	73	13	0.094	0.094
6	73	13	0.283	0.164
9	73	13	0.054	0.034
10	73	13	0.595	0.283
11	73	13	6.201	1.614
12	73	13	4.927	6.881
	12	120	17	16.282
1	74	13	6.003	1.869
2	74	13	6.003	20

TRIBUTARY FLOW INFORMATION FOR INDIANA							03/29/76
LAKE CODE	1850	VERSAILLES	MEAN MONTHLY FLOWS AND DAILY FLOWS (CMS)				
TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW
1850B1	6	73	0.283	9	0.042		
	7	73	0.481	13	0.014		
	8	73	0.006	17	0.003		
	9	73	0.001	17	0.000		
	10	73	0.014	15	0.003		
	11	73	0.218	16	0.045		
	12	73	0.176	21	0.023		
	1	74	0.425	17	0.566	29	0.453
	2	74	0.210	15	0.054	20	0.255
	3	74	0.283	15	0.198		
	4	74	0.510	19	0.045		
	5	74	0.283	26	0.031		
1850C1	6	73	0.139	9	0.028		
	7	73	0.238	13	0.011		
	8	73	0.006	17	0.003		
	9	73	0.001	17	0.001		
	10	73	0.011	15	0.003		
	11	73	0.110	16	0.028		
	12	73	0.088	21	0.125		
	1	74	0.215	17	0.283	29	0.227
	2	74	0.108	15	0.034	20	0.127
	3	74	0.150	15	0.113		
	4	74	0.255	19	0.028		
	5	74	0.167	26	0.020		
1850D1	6	73	0.042	9	0.008		
	7	73	0.071	13	0.003		
	8	73	0.002	17	0.001		
	9	73	0.000	17	0.000		
	10	73	0.003	15	0.001		
	11	73	0.034	16	0.008		
	12	73	0.025	21	0.040		
	1	74	0.065	17	0.088	29	0.068
	2	74	0.031	15	0.011	20	0.037
	3	74	0.042	15	0.034		
	4	74	0.076	19	0.008		
	5	74	0.042	26	0.006		
1850Z2	6	73	0.040	9	0.008		
	7	73	0.068	13	0.003		
	8	73	0.001	17	0.001		
	9	73	0.000	17	0.000		
	10	73	0.003	15	0.001		
	11	73	0.031	16	0.008		
	12	73	0.025	21	0.034		
	1	74	0.059	17	0.082	29	0.062
	2	74	0.031	15	0.008	20	0.037
	3	74	0.042	15	0.031		
	4	74	0.071	19	0.008		
	5	74	0.040	26	0.008		

**APPENDIX D**

**PHYSICAL and CHEMICAL DATA**

STORED RETRIEVAL DATE 76/03/30

185001  
39 04 52.0 085 14 05.0 3  
VERSAILLES LAKE  
18137 INDIANA

052192

DATE FROM TO	TIME OF DAY	DEPTH FEET	WATER TEMP CENT	00010 DO MG/L	000300 TRANSI SECCHI INCHES	00077 CNDCTVY FIELD MICROMHO	00094 NH3-N TOTAL MG/L	11EPALES 0007 FEET			2111202 DEPTH CLASS 00		
								TALK CACO3 SU	NH3-N TOTAL MG/L	TOT KJEL N MG/L	N-TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	
73/04/29	13 45	0000	14.3		8	330	7.90	126	0.120	1.100	1.100		
73/08/02	13 45	0003	14.0	8.8		310	8.00	128	0.100	0.700	1.100		
73/10/10	12 00	0000	24.9	8.6	20	285	8.20	109	0.100	1.500	0.830		
	10 00	0000	21.6	8.6	28	309	7.80	136	0.070	2.000	0.010		
											0.012		

DATE FROM TO	TIME OF DAY	DEPTH FEET	PHOS-TOT MG/L P	00665 CHLRPHYL A UG/L	32217 3.2	32217 CHLRPHYL A UG/L		
						0.197 0.176	0.119 0.109	64.7 30.2
73/04/29	13 45	0000						
73/08/02	13 45	0003						
73/10/10	12 00	0000						
	10 00	0000						

STORRET RETRIEVAL DATE 76/03/30

185002  
39 05 15.0 085 14 35.0 3  
VERSAILLES LAKE  
18137 INDIANA

052192

11EPALES				2111202			
0007 FEET				DEPTH	CLASS	00	
DATE	TIME	DEPTH	00010	00077	00094	00410	00625
FROM OF TO	DAY	FEET	WATER TEMP CENT	TRANS SECCHI INCHES	CONDUTCTV FIELD MICROMHO	PH TALK CACO3 MG/L	NH3-N TOT KJEL TOTAL N MG/L
73/04/29	13	15 0000	14.0		10	330	8.00
	13	15 0005	13.9	8.8		350	8.00
73/08/02	12	15 0000	24.5	5.6		324	7.60
73/10/10	10	10 0000	22.4	9.2	18	319	6.26
DATE	TIME	DEPTH	00665	32217	CHLRPHYL A UG/L		
FROM OF TO	DAY	FEET	PHOS-TOT				
73/04/29	13	15 0000	0.189				
	13	15 0005	0.203				
73/08/02	12	15 0000	0.115				
73/10/10	10	10 0000	0.108				
				139	0.800	1.300	0.040
						1.200	0.070
						0.110	0.066
						1.28	0.012
						0.240	0.013
						1.39	0.013

STORED RETRIEVAL DATE 76/03/30

185003 39 04 32.0 085 14 48.0 3  
VERSAILLES LAKE  
10137 INDIANA

052192

11EPALES				2111202			
DATE		TIME	DEPTH	0019 FEET		DEPTH	CLASS
FROM TO	OF DAY	FEET	CENT	PH	TALK	KJEL	00
73/04/29	14	00	0000	00000	00000	00000	00
	14	00	0003	14.1	118	1.000	00000
	14	00	0015	13.9	0.160	0.900	00000
	14	00	0030	13.5	0.170	0.700	00058
73/08/02	12	25	0000	6.6	300	1.00	00054
	12	25	0005	6.5	330	0.800	0.054
	12	25	0015	24.9	7.80	0.180	0.054
	12	25	0020	24.7	271	0.12	0.054
	12	25	0030	8.0	8.40	0.060	0.011
	12	25	0040	24.7	270	0.20	0.011
	12	25	0050	22.8	0.20	0.060	0.011
	12	25	0100	0.5	245	7.30	0.810
	12	25	0115	22.8	98	0.390	0.041
73/10/10	10	25	0000	22.0	315	7.90	0.750
	10	25	0005	21.4	315	7.30	0.030
	10	25	0015	1.0	315	1.43	0.040
	10	25	0020	0.5	323	7.20	0.014
	10	25	0030	21.2	323	146	0.050
	10	25	0040	21.4	323	146	0.019

DATE FROM TO	TIME OF DAY	DEPTH FEET	PHOS-TOT MG/L	CHLORPHYL A UG/L	
73/04/29	14 00	0 000	0.173	19.8	32217
	14 00	0 003	0.193		
	14 00	0 015	0.230		
73/08/02	12 25	0 000	0.091	4.9	
	12 25	0 005	0.087		
73/10/10	12 25	0 015	0.180	22.6	
	10 25	0 000	0.094		
	10 25	0 005	0.089		
	10 25	0 012	0.139		

APPENDIX E

TRIBUTARY DATA

STORET RETRIEVAL DATE 76/03/30

DATE FROM TO	TIME OF DAY	DEPTH FEET	NO2&NO3 N-TOTAL MG/L	TOT KJEL N MG/L	NH3-N TOTAL MG/L	PHOS-DIS ORTHO MG/L P	PHOS-TOT MG/L P
73/06/09	10 00		0.294	1.600	0.019	0.023	0.145
73/07/13	09 40		0.820	1.980	0.115	0.023	0.115
73/08/17	11 12		0.810	1.100	0.240	0.110	0.180
73/09/17	09 25		0.760	1.800	0.550	0.260	0.410
73/10/15	10 50		0.410	1.500	0.460	0.180	0.260
73/11/16	10 55		0.384	0.850	0.044	0.044	0.110
73/12/21	13 30		1.760	0.800	0.144	0.184	0.250
74/01/17	15 00		1.900	0.900	0.144	0.132	0.300
74/01/29	10 05		1.440	1.500	0.065	0.085	0.320
74/02/15	10 00		1.700	1.000	0.105	0.115	0.210
74/02/20	11 10		1.760	1.700	0.095	0.115	0.320
74/03/15	10 00		1.520	1.300	0.075	0.100	0.175
74/04/19	10 30		0.970	0.900	0.050	0.025	0.050
74/05/26	13 50		1.600	1.600	0.035	0.010	0.185

1850A1  
39 04 05.0 085 14 16.0 4

LAUGHERY CREEK  
18 7.5 MILAN  
O/VERSAILLES  
COVERED BRDG E OF VERSAILLES  
NIEPALES 052192  
0000 FEET DEPTH CLASS 00

STORRET RETRIEVAL DATE 76/03/30

1850A2		39 06 13.0 085 14 16.0 4	
		LAUGHERY CREEK	
		18 7.5 MILAN	
		1/VERSAILLS	
		BANK AT END OF SEC RD	
		052192	
		1IEPALES	
		2111204	
		0000 FEET DEPTH CLASS 00	
DATE	TIME	DEPTH	NO2&NO3
FROM TO	OF DAY	FEET	N-TOTAL MG/L
73/06/09	10 10	1.700	00625 TOT KJEL
73/07/13	09 15	0.810	00630 N TOTAL
73/08/17	10 50	0.013	00630 MG/L
73/09/17	08 55	0.010K	00625 NH3-N
73/10/15	10 20	0.013	00625 TOTAL
73/11/16	10 30	0.384	00610 MG/L
74/01/17	14 30	4.500	00671 PHOS-UIS
74/01/29	09 30	0.920	00671 ORTHO
74/02/15	09 25	2.000	00665 MG/L P
74/02/20	10 40	2.000	00665 PHOS-TUT
74/03/15	09 30	1.800	
74/04/19	10 10	0.720	
74/05/26	13 30	1.500	

K VALUE KNOWN TO BE  
LESS THAN INDICATED

STORED RETRIEVAL DATE 76/03/30

DATE FROM TO		TIME OF DAY	DEPTH FEET	N02&N03 N-TOTAL MG/L	TOT KJEL N MG/L	NH3-N TOTAL MG/L	00610 PHOS-DIS ORTHO MG/L P	00671 PHOS-TOT MG/L P	00665
73/06/09	11	05		0.016	0.665	0.029	0.010	0.015	
73/07/13	10	05		0.058	0.480	0.038	0.010	0.017	
73/08/17	12	30		0.010K	0.880	0.033	0.005K	0.019	
73/09/17	09	50		0.011	0.460	0.069	0.014	0.080	
73/10/15	11	00		0.035	0.550	0.024	0.026	0.055	
73/11/16	11	10		0.022	0.250	0.017	0.016	0.025	
74/01/17	15	30		2.300	0.400	0.028	0.020	0.055	
74/01/29	10	20		0.616	0.300	0.020	0.010	0.020	
74/02/15	10	20		0.690	0.600	0.015	0.010	0.030	
74/02/20	11	25		0.690	1.300	0.115	0.015	0.045	
74/03/15	10	10		0.368	0.500	0.015	0.005	0.012	
74/04/19	10	45		0.068	0.450	0.030	0.005K	0.005	
74/05/26	14	10		0.032	0.300	0.025	0.020	0.020	

K VALUE KNOWN TO BE  
LESS THAN INDICATED

STORET RETRIEVAL DATE 76/03/30

DATE FROM TO	TIME OF DAY	DEPTH FEET	NO2&NO3 N-TOTAL MG/L	TOT KJEL N MG/L	00630		00625		00610		00671		00665	
					M6/L	P	M6/L	P	M6/L	P	M6/L	P	M6/L	P
73/06/09	11	35		0.086	0.820		0.056	0.023	0.045					
73/07/13	10	30		0.073	0.520		0.008	0.022	0.065					
73/08/17	12	00		0.048	0.250		0.019	0.036	0.050					
73/09/17	10	30		0.056	0.800		0.039	0.028	0.040					
73/10/15	11	55		0.026	3.150		0.105	0.033	0.080					
73/11/16	12	00		0.240	1.200		0.029	0.020	0.195					
74/01/17	14	15		1.260	0.500		0.088	0.076	0.135					
74/01/29	11	15		1.700	0.600		0.015	0.090	0.230					
74/02/15	11	05		0.830	1.300		0.085	0.015	0.060					
74/02/20				0.810	2.200		0.280	0.010	0.065					
74/03/15	10	45		0.368	0.300		0.015	0.015	0.035					
74/04/19	11	40		0.940	1.000		0.015	0.025	0.080					
74/05/26	14	35		0.088	0.200		0.010	0.015	0.035					

1850CL  
 39 04 44.0 085 15 05.0 4  
 UNNAMED STREAM  
 18 7.5 VERSAILLES  
 T/VERSAILLES 051892  
 BANK NEAR MOUTH WALK FROM END OF SEC RD  
 11EPALES 2111204  
 0000 FEET DEPTH CLASS 00

STORET RETRIEVAL DATE 76/03/30

DATE FROM TO	TIME OF DAY	DEPTH FEET	NO2&N03 N-TOTAL MG/L	TOT KJEL N MG/L	NH3-N TOTAL MG/L	PHOS-DIS ORTHO MG/L P	PHOS-TOT MG/L P
73/06/09	11	45	0.090	0.960	0.040	0.034	0.065
73/07/13	11	00	0.180	0.820	0.054	0.037	0.050
73/08/17	11	50	0.110	0.355	0.014	0.029	0.035
73/09/17	10	10	0.115	0.920	0.038	0.028	0.040
73/10/15	11	40	0.084	0.400	0.025	0.033	0.045
73/11/16	11	50	0.240	1.200	0.035	0.012	0.092
74/01/17	15	20	0.750	0.300	0.072	0.044	0.075
74/01/29	11	00	0.810	0.100	0.005	0.010	0.051
74/02/15	11	20	1.260	1.100	0.075	0.060	0.145
74/02/20			1.440	1.900	0.210	0.077	0.195
74/03/15	10	30	1.400	0.800	0.040	0.085	0.160
74/04/19	11	20	0.552	1.100	0.045	0.015	0.050
74/05/26	14	20	0.128	0.250	0.025	0.015	0.037

185001  
39 04 53.0 085 15 02.0 4

CEDAR CREEK

18 7.5 VERSAILLES

T/VERSAILLES  
BANK NEAR MOUTH WALK FROM END HASSMER RD  
11EPALES 051892

2111204  
0000 FEET DEPTH CLASS 00

STORED RETRIEVAL DATE 76/03/30

DATE FROM TO	TIME OF DAY	DEPTH FEET	NO2&NO3 N-TOTAL	TOT KJEL N MG/L	NH3-N TOTAL MG/L	PHOS-DIS ORTHO MG/L P	00671 00610 PHOS-TOT INST MGD	00665 50051 FLOW RATE INST MGD	50053 CONDUIT FLOW-MGD MONTHLY
73/06/23	10	30	0.130	10.000	2.700	4.500	5.200	0.200	
73/09/18	11	00	3.635	17.600	5.230	6.360	8.200		
74/01/24	14	00	2.880	21.000	0.150	1.440	3.700	0.150	
74/05/05	12	00	0.840	23.000	3.000	2.100	4.300		

1850XA  
39 07 52.0 085 16 50.0 4  
OSGOOD

18 7.5 OSGOOD  
T/VERSAILLES LAKE 052192  
NO NAME DITCH/LAUGHERY CREEK  
11EPALES 2141204  
0000 FEET DEPTH CLASS 00

TF1850XA  
P001500\*



STORET RETRIEVAL DATE 76/03/30

					1850AB	TF1850AB	
DATE	TIME	DEPTH	NO2&N03	TOT KJEL	00610	00665	P004000
FROM TO	OF DAY	N-TOTAL FEET	N	TOTAL MG/L	PHOS-DIS ORTHO MG/L P	PHOS-TOT RATE MG/L P	
73/06/08	07 00	8.100	3.500	0.160	20.100	23.000	0.380
CP(T)-							0.350
73/06/08	17 00	6.3/07/09	07 00	4.500	6.400	4.200	6.000
73/07/05	17 00	7.3/08/08	07 00	5.900	8.750	0.980	29.000
CP(T)-							0.360
73/08/08	17 00	73/09/10	07 00	9.200	10.000	1.040	15.200
CP(T)-							0.250
73/09/10	17 00	73/10/03	07 00	11.800	7.150	1.600	11.600
CP(T)-							0.210
73/10/03	17 00	73/11/08	07 00	5.600	13.000	2.940	32.000
CP(T)-							0.200
73/11/08	17 00	73/12/06	07 00	6.000	8.200	0.350	36.000
CP(T)-							0.260
73/12/06	17 00	74/01/09	07 00	3.300	9.700	1.560	25.200
CP(T)-							0.220
74/01/09	17 00	74/02/14	07 00	5.300	9.100	1.300	27.500
CP(T)-							0.220
74/02/14	17 00	74/02/14	17 00	5.040	7.700	0.330	20.500
CP(T)-							0.260
74/03/07	17 00	74/04/04	07 00	5.880	1.400	0.130	15.000
CP(T)-							0.320
74/04/04	17 00	74/05/07	07 00	6.700	6.900	0.150	24.000
CP(T)-							0.380
74/05/07	17 00						

1850AB  
39 18 30.0 085 13 00.0 4  
BATESVILLE  
18 RIPLEY CO MAP  
VERSAILLES LAKE 051392  
LITTLE LAUGHERY CREEK  
11EPALES  
0000 FEET DEPTH CLASS 00

U.S. Environmental Protection Agency,  
Region 5, Library (PL-12J)  
77 West Jackson Boulevard, 12th Floor  
Chicago, IL 60604-3590