

United States  
Environmental Protection  
Agency

Environmental Sciences Research  
Laboratory  
Research Triangle Park NC 27711

EPA-600 4-80 006  
January 1980

Research and Development



# Regional Air Pollution Study

## Gas Chromatography Laboratory Operation



## RESEARCH REPORTING SERIES

Research reports of the Office of Research and Development, U S Environmental Protection Agency, have been grouped into nine series. These nine broad categories were established to facilitate further development and application of environmental technology. Elimination of traditional grouping was consciously planned to foster technology transfer and a maximum interface in related fields. The nine series are

- 1 Environmental Health Effects Research
- 2 Environmental Protection Technology
- 3 Ecological Research
- 4 Environmental Monitoring
- 5 Socioeconomic Environmental Studies
- 6 Scientific and Technical Assessment Reports (STAR)
- 7 Interagency Energy-Environment Research and Development
- 8 "Special" Reports
- 9 Miscellaneous Reports

This report has been assigned to the ENVIRONMENTAL MONITORING series. This series describes research conducted to develop new or improved methods and instrumentation for the identification and quantification of environmental pollutants at the lowest conceivably significant concentrations. It also includes studies to determine the ambient concentrations of pollutants in the environment and/or the variance of pollutants as a function of time or meteorological factors.

EPA-600/4-80-006  
January 1980

REGIONAL AIR POLLUTION STUDY  
Gas Chromatography Laboratory Operation

by

G. Cardwell  
Rockwell International  
Environmental Monitoring & Services Center  
Environmental & Energy Systems Division  
11640 Administration Drive  
Creve Coeur, MO 63141

Contract No. 68-02-2093  
Task Order 113

Project Officer

Stanley Kopczynski  
Atmospheric Chemistry and Physics Division  
Environmental Sciences Research Laboratory  
Research Triangle Park, North Carolina 27711

ENVIRONMENTAL SCIENCES RESEARCH LABORATORY  
OFFICE OF RESEARCH AND DEVELOPMENT  
U.S. ENVIRONMENTAL PROTECTION AGENCY  
RESEARCH TRIANGLE PARK, NORTH CAROLINA 27711

## DISCLAIMER

This report has been reviewed by the Environmental Sciences Research Laboratory, U.S. Environmental Protection Agency, and approved for publication. Approval does not signify that the contents necessarily reflect the views and policies of the U.S. Environmental Protection Agency, nor does mention of trade names or commercial products constitute endorsement or recommendation for use.

## ABSTRACT

A gas chromatography laboratory was set up to analyze air samples collected in 100-liter Teflon bags and 8-liter stainless steel tanks. Samples were analyzed for total hydrocarbons (THC), methane, and CO on a Beckman Model 6800 gas chromatograph, and for C<sub>2</sub>-C<sub>10</sub> hydrocarbons on a Perkin-Elmer Model 900 gas chromatograph equipped with dual columns, connected to a PEP-1 data system. A phenyl isocyanate column resolved the C<sub>2</sub>-C<sub>5</sub> compounds and a support-coated-open-tubular squalane column was used for C<sub>6</sub>-C<sub>10</sub> compounds. Experiments were performed to establish optimum temperature programming, flow rates, and column lengths to yield good compound resolution within reasonable elution times.

A total of 455 samples including replicates, were analyzed during the summer and fall, 1976. Of these samples, 292 were collected at 12 of the Regional Air Monitoring System (RAMS) sites to yield data on spatial and temporal distributions of hydrocarbons. All were analyzed for THC, methane, CO, and C<sub>2</sub>-C<sub>5</sub> hydrocarbons. In addition, early morning (0600-0800) samples were analyzed for C<sub>6</sub>-C<sub>10</sub> compounds. Portions of these data are included in this report. Sampling results indicate excessive concentration of ethylene within some of the RAMS stations, probably from the ozone monitoring system used at all RAMS sites. Some samples collected during roadway studies were re-analyzed on subsequent days; results showed good reproducibility using the Teflon bags. System reproducibility from quality control checks was good; with analyses of standards indicating deviations generally less than 5 percent.

All data, including the sums of paraffins, olefins, aromatics and total non-methane hydrocarbons are stored in the RAPS Data Bank at Research Triangle Park, North Carolina.

## CONTENTS

Abstract . . . . .	iii
Figures. . . . .	vii
Tables . . . . .	viii
1.0 Introduction. . . . .	1
2.0 Summary . . . . .	2
3.0 Development of the Squalane Column. . . . .	4
3.1 Literature Search. . . . .	4
3.2 Development of the 60.96 and 30.48 Meter Columns . . . . .	6
3.2.1 Preliminary Experiments . . . . .	6
3.2.2 C <sub>2</sub> -C <sub>5</sub> Analysis on the Phenyl Isocyanate (Dura Pak) Column . . . . .	7
3.2.3 Use of Dual Columns . . . . .	7
3.2.4 Response Factors for Hydrocarbons C <sub>6</sub> -C <sub>10</sub> . . . . .	12
4.0 Analysis of RAMS Stations Samples During the Summer and Fall Intensives of 1976. . . . .	16
4.1 Selection of Sampling Bags and Tanks . . . . .	16
4.2 Ambient Air Sampling . . . . .	19
4.2.1 Bag Preparation and Sample Collection . . . . .	19
4.3 Ambient Air Sampling . . . . .	22
4.4 Data Validation, Interpretation and Distribution . . . . .	22
5.0 Special Studies . . . . .	27
5.1 Collection and Analyses of Roadway Samples . . . . .	27
5.2 Ethylene Studies . . . . .	29
5.3 Special Audits . . . . .	29
5.3.1 Total Hydrocarbon, Methane and Carbon Monoxide Audits . . . . .	31
5.3.2 Isobutane, Normal Hexane and Synthetic Bag Mixture Audits . . . . .	33
5.4 Special Studies at RAMS 103 and 107. . . . .	33
5.5 Data Validation, Interpretation and Distribution . . . . .	34

## CONTENTS (continued)

6.0	Quality Control . . . . .	.38
6.1	Standards for Beckman 6800 . . . . .	.38
6.2	Standards for Perkin Elmer 900 . . . . .	.38
6.3	Daily Standards . . . . .	.38
7.0	Data Processing . . . . .	.42
7.1	Data Tabulation. . . . .	.42
7.2	Keypunching and Processing . . . . .	.42
	Literature Cited . . . . .	.44
	Appendices	
A.	Operation of Beckman 6800 Gas Chromatograph . . . . .	.45
B.	Operation of the Perkin Elmer 900 Gas Chromatograph . . . . .	.52
C.	Data from Summer and Fall Intensives. . . . .	.65
D.	Special Studies . . . . .	117

## FIGURES

<u>Number</u>		<u>Page</u>
1	Squalane Column Analysis of Atmospheric Sample for C <sub>2</sub> - C <sub>10</sub> Hydrocarbons . . . . .	5
2	Laboratory Air - Phenyl Isocyanate Column 10/14/76 (New Method). . . . .	8
3	Sample of Ambient Air on the Squalane Column (New Method) . .	9
4	Hydrocarbons C <sub>2</sub> - C <sub>5</sub> on Dura Pak, 3 Minutes at 0°, Programmed at 6°/Minute (Old Method). . . . .	10
5	Atmospheric Organics Analysis System - P.E. 900 . . . . .	11
6	Quantitative Study of Hydrocarbons C <sub>5</sub> - C <sub>10</sub> 6/17/76 . . . .	13
7	Bag 17 Analyzed on the Phenyl Isocyanate Column 6/9/76. . .	17
8	Bag 17 Analyzed on Squalane Column 6/19/76. . . . .	18
9	Background of Stainless Steel Tank. . . . .	20
10	RAMS Bag Sampling System Flow Diagram . . . . .	21
11	Five Point Calibration of Beckman Model 6800 Gas Chromatograph . . . . .	32



## TABLES

<u>Number</u>		<u>Page</u>
1	Experimentally Determined Response Factors for C <sub>2</sub> - C <sub>10</sub> Hydrocarbons . . . . .	14
2	Estimated Response Factors for C <sub>6</sub> - C <sub>10</sub> Hydrocarbons. . . .	15
3	Ambient Air Samples Analyzed in the Gas Chromatography Laboratory 1976 . . . . .	23
4	Summary of Summer and Fall Intensives 1976. . . . .	25
5	Average Carbon Monoxide to Acetylene Ratios . . . . .	26
6	Carbon Monoxide to Acetylene Ratios of Roadway Samples. . .	28
7	Storage of Roadside Samples in Teflon Bags. . . . .	30
8	Carbon Monoxide Audits. . . . .	31
9	Methane Audits. . . . .	31
10	Isobutane, N-Hexane and Special Bag Audits. . . . .	33
11	Quantitative Studies at RAMS 103 and 107. . . . .	35
12	Propane Standard. . . . .	40
13	Toluene Standard. . . . .	41

## 1.0 INTRODUCTION

The Regional Air Pollution Study (RAPS) is directed toward a quantitative understanding of urban pollution, including the monitoring of ambient levels of pollution, the gathering of micro-meteorological data and of a comprehensive emission inventory.

The gas chromatography laboratory was established to support these investigations. In particular it was charged with the collection and analysis of samples of hydrocarbons  $C_1 - C_{10}$ , carbon monoxide and tracer gases in ambient air. In order to find a more accurate and selective method for identifying hydrocarbons  $C_2 - C_{10}$ , modifications were made on the Perkin Elmer 900 Gas Chromatograph such that detailed analyses of these hydrocarbons in the parts-per-billion range could be performed. Samples were run and the results were tabulated and classified.

## 2.0 SUMMARY

The gas chromatography laboratory modified the analysis of  $C_2 - C_{10}$  hydrocarbons from the methods described in the final reports of Task Orders 53 and 103. A Perkin Elmer (P.E.) Model 900 gas chromatograph equipped with dual columns and a PEP-1 data system was used. A phenyl isocyanate (Dura Pak) column gave good resolution for  $C_2 - C_5$  compounds and a Support-Coated-Open-Tubular (SCOT) squalane column was used to analyze hydrocarbons  $C_6 - C_{10}$ . The modifications included shortening the SCOT squalane column from 30.48 meters to 15.24 meters and raising the initial temperature from  $-3^\circ\text{C}$  to  $25^\circ\text{C}$ . These changes permitted the accurate and methodical identification of hydrocarbons  $C_6 - C_{10}$  on the squalane column with an analysis time of 55 minutes. As result of these modifications the precision of the analyses done on the squalene column was improved as indicated by the reproducibility of the analysis of a toulene standard over a three month interval.

A total of 455 samples of ambient air (including triplicate analyses) was analyzed during the summer and fall intensives and for special studies during this task order period. These samples were collected from RAMS stations and roadside locations and were analyzed for total hydrocarbons (THC), carbon monoxide (CO) and methane ( $\text{CH}_4$ ) on the Beckman Model 6800 gas chromatograph and hydrocarbons  $C_2 - C_{10}$  on the P.E. 900 gas chromatograph.

Two hundred and ninety-two ambient air samples from twelve RAMS sites were analyzed during the summer and fall intensives. All of the samples were analyzed for THC,  $\text{CH}_4$ , and CO (Beckman 6800) and  $C_2 - C_5$  hydrocarbons (P.E. 900). The 0600-0800 (CST) samples were analyzed for THC,  $\text{CH}_4$  and CO and  $C_2 - C_{10}$  hydrocarbons (P.E. 900). Tables containing a portion of these data are in Appendix C.

High concentrations of ethylene at RAMS sites 101 and 124 led to a special study of ethylene concentration. The samples were collected in 109 litre Teflon bags (5 mil thickness) and 8 litre stainless steel tanks, in

and around the stations as requested by the EPA Task Coordinator. The results of this special study indicate that in certain cases the values reported for ethylene may be biased by fugitive emission of ethylene from the ozone monitor and associated hardware found in each of the RAMS stations.

Roadway samples were analyzed by the gas chromatography laboratory for a special study conducted by the EPA Task Coordinator. Some of the samples were reanalyzed on subsequent days. Relatively good reproducibility was obtained using the 5 mil Teflon bags.

To insure quality control, standards were run daily and records maintained of the areas and peak heights. The triplicate analyses of the first bag sample each day showed a standard deviation of less than 2.0 ppbC for the task order period.

All of these data, including the sums of the paraffins, olefins, aromatics and total non-methane hydrocarbons, were processed and prepared for entry into the RAPS Computer Data Bank, Research Triangle Park (RTP), North Carolina.

### 3.0 DEVELOPMENT OF THE SQUALANE COLUMN

In the past, the gas chromatography laboratory experienced some difficulty in the analysis of  $C_5 - C_{10}$  compounds on the squalane column. During Task Orders 53 and 103 periods the initial 60.96 meters x 0.05 cm squalane SCOT column was reduced to 30.48 meters because of excessive column bleed (1, 2). A Carbowax 20M-TPA on Chromosorb W-AW post column was added during Task Order 53 but was replaced with glass beads during Task Order 103 because of the problems incurred with the Carbowax column. A potassium carbonate ( $K_2CO_3$ ) pretrap was added during Task Order 53 in order to minimize the effect of water and polar compounds in the squalane system. Figure 1 is an example of hydrocarbons  $C_2 - C_{10}$  analyzed during both task order periods. Because of the length of time required for analysis and the broadening of the peaks,  $C_9 - C_{10}$  were not really identified. Alternate methods of analysis were explored.

The literature was searched and the problem examined in a systematic manner with the aim of developing a method of analysis that: (1) provided a reliable basis for identification of hydrocarbons on the squalane SCOT column; (2) could be easily incorporated in the equipment of the laboratory; and (3) preferably employed only one column for the analyses of compounds  $C_2 - C_{10}$ .

#### 3.1 LITERATURE SEARCH

Extensive work on the squalane SCOT column has been reported in the literature. W. O. McReynolds (3) and others described the analyses of hydrocarbons  $C_2 - C_{10}$  on 60.96 and 30.48 meters x 0.05 cm columns. The problem encountered was the length of time required to complete the analyses since in some cases five or more hours were required. L. S. Ettre (4) described the analysis of  $C_1 - C_9$  hydrocarbons on a 30.48 meter x 0.05 cm column requiring only 35 minutes. The method, although attractive, had some shortcomings: it did not identify all the compounds in which the gas




FIGURE 1. SQUALANE COLUMN ANALYSIS OF ATMOSPHERIC SAMPLE FOR  $C_2 - C_{10}$  HYDROCARBONS

chromatography laboratory was interested, e.g. ethyl benzene, the xylenes and decanes; therefore, modifications had to be made.

### 3.2 DEVELOPMENT OF THE 60.96 AND 30.48 METER COLUMNS

#### 3.2.1 Preliminary Experiments

Two 100 litre Teflon bags were filled with 50 litres of Scott-Marrin ultrapure air by using a mass flow meter which was corrected for standard temperature and pressure. Bag 1 was injected with 10 ml of each of the following gases using the appropriate size precision sampling syringe: methane ( $\text{CH}_4$ ), ethane ( $\text{C}_2\text{H}_6$ ), ethylene ( $\text{C}_2\text{H}_4$ ), propane ( $\text{C}_3\text{H}_8$ ), acetylene ( $\text{C}_2\text{H}_2$ ), isobutane (I- $\text{C}_4\text{H}_{10}$ ), normal butane (N- $\text{C}_4\text{H}_{10}$ ); and 0.05 ml each of the following liquids: isopentane (I- $\text{C}_5\text{H}_{12}$ ), normal pentane (N- $\text{C}_5\text{H}_{12}$ ), hexane ( $\text{C}_6\text{H}_{14}$ ), heptane ( $\text{C}_7\text{H}_{16}$ ), toluene ( $\text{C}_7\text{H}_8$ ), octane ( $\text{C}_8\text{H}_{18}$ ), ethyl benzene ( $\text{C}_8\text{H}_{10}$ ), meta xylene ( $\text{C}_8\text{H}_{10}$ ), ortho xylene ( $\text{C}_8\text{H}_{10}$ ), nonane ( $\text{C}_9\text{H}_{20}$ ), mesitylene ( $\text{C}_9\text{H}_{12}$ ) and decane ( $\text{C}_{10}\text{H}_{22}$ ). After equilibration, one milliliter of this mixture was injected into Bag 2 (dilution bag). The diluted contents of Bag 2 were analyzed on the 60.96 and 30.48 meters x 0.05 cm squalane columns.

The 60.96 and 30.48 meters squalane columns provided acceptable separation for hydrocarbons  $\text{C}_1 - \text{C}_7$ ; but, the heavier hydrocarbons ( $\text{C}_8 - \text{C}_{10}$ ) took an exceptionally long time to elute. The physical conditions (flow rate and temperature) were the same as described in the final report for Task Orders 53 and 103. In an attempt to overcome the elution problem with the heavier hydrocarbons, the initial oven temperature was varied from  $-3^\circ\text{C}$  to  $0^\circ$ ,  $3^\circ$ ,  $15^\circ$  and  $25^\circ\text{C}$  and the final temperature from  $75^\circ\text{C}$  to  $90^\circ\text{C}$  and  $115^\circ\text{C}$ . The carrier gas flow rate was increased from 12 cc/min. to 15 cc/min. The results of these changes were not satisfactory; an increase in temperature and carrier gas flow rate yielded no data on  $\text{C}_1 - \text{C}_6$  hydrocarbons (they eluted in the initial 15 seconds) and the higher temperatures yielded significant shifts in the baseline of the chromatograph at  $90^\circ\text{C}$  and above. These changes had little or no effect on the heavier hydrocarbons.

The 15.24 meter squalane SCOT column was then employed and was found suitable for the analyses of  $\text{C}_6 - \text{C}_{10}$  hydrocarbons. It was found that hydrocarbons  $\text{C}_1 - \text{C}_5$  elute too fast (within the first two minutes) to be

properly identified. The hydrocarbons were analyzed under the same physical conditions (temperatures and flow rates) as described for the 60.96 and 30.48 meters columns. The initial temperature was increased to 25°C and the final temperature increased to 90°C; the carrier gas flow rate remained at 12 cc/min. These conditions proved ideal for the analysis of hexane, which eluted at 3.6 minutes, through normal butyl benzene, which eluted at 55 minutes. It was then decided to continue to employ two columns in the P.E. 900 in the analyses of hydrocarbons  $C_2 - C_{10}$ . Figures 2 and 3 show typical examples of ambient air analyzed with the two columns employed in the P.E. 900 gas chromatograph.

### 3.2.2 $C_2 - C_5$ Analysis on the Phenyl Isocyanate (Dura Pak) Column

During the preceding task order period (103)  $C_2 - C_5$  hydrocarbons were analyzed on the 1.83 meter x 0.63 cm ID phenyl isocyanate (Dura Pak) column where the initial oven temperature was held at 0° for 3 minutes then programmed at a rate of 6°/minute until normal pentane ( $N-C_5H_{12}$ ), the last compound analyzed on the column, eluted. The whole analysis took 13 minutes to complete.


With the modified procedure, initial oven temperature at 25°C for 8 minutes, the programmed at a rate of 2°/minute and employing the Dura Pak column, normal pentane elutes in 10 minutes. Figures 2 and 4 show that the change in environment had no adverse effect on the  $C_2 - C_5$  analysis.

### 3.2.3 Use of Dual Columns

The analysis of hydrocarbons  $C_2 - C_{10}$  was accomplished by using two columns connected to two flame ionization detectors in the Perkin Elmer Model 900 gas chromatograph (Figure 5 shows the schematic diagram of the analysis system). Hydrocarbons  $C_2 - C_5$  were analyzed on a 1.83 meter x 0.63 cm ID phenyl isocyanate (Dura Pak) column that was conditioned by heating it at 115°C with helium flowing through over a 24 hour period. Hydrocarbons  $C_6 - C_{10}$  were analyzed on a 15.24 meter x 0.05 cm ID squalane SCOT column that was conditioned at 115°C with helium flowing through over a 60 hour period. Both columns shared the same oven which was held at 25°C for eight minutes, then programmed at a rate of 2°C per minute until the



FIGURE 2. LABORATORY AIR - PHENYL ISOCYANATE  
COLUMN 10/14/76 (NEW METHOD)






FIGURE 3. SAMPLE OF AMBIENT AIR ON THE SQUALANE COLUMN  
(NEW METHOD)




FIGURE 4. HYDROCARBONS  $C_2 - C_5$  ON DURA PAK , 3 MINUTES AT  $0^\circ$ ,  
PROGRAMMED AT  $6^\circ/\text{MINUTE}$  (Old Method)




FIGURE 5. ATMOSPHERIC ORGANICS ANALYSIS SYSTEM - P.E. 900

system reached 90°C, where it was held for the remainder of the analysis. The total analysis times were 10 minutes for hydrocarbons C<sub>2</sub> - C<sub>5</sub> on the Dura Pak column and 55 minutes for hydrocarbons C<sub>6</sub> - C<sub>10</sub> on the squalane column.

### 3.2.4 Response Factors For Hydrocarbons C<sub>6</sub> - C<sub>10</sub>

In order to obtain accurate response factors (RF) for C<sub>6</sub> - C<sub>10</sub> hydrocarbons for the squalane column, two synthetic mixtures of the various compounds were carefully prepared, using the procedure described in Section 3.2.1, then analyzed on the P.E. 900 gas chromatograph. The concentrations of the liquid compounds were computed in the following manner.

Example: Toluene

$$\text{Bag-1} \quad \frac{22.4 \times \frac{295^\circ}{273^\circ} \text{ l/mole} \times \frac{1 \text{ mole}}{92.15 \text{ gm}} \times 0.8669 \frac{\text{gm}}{\text{ml}} \times 0.05 \text{ ml}}{50 \text{ litres of ultrapure air}} = 228 \text{ ppm}$$

$$\text{Bag-2} \quad \frac{1 \text{ ml of contents of Bag 1} \times 228 \text{ ppm}}{50 \text{ litres of ultrapure air}} \times 7 \text{ carbon number} = 31.9 \text{ ppbC}$$

The PEP-1 computer was programmed in such a manner (area normalization) that only the areas of the selected compounds were computed. The response factors for the hydrocarbons were calculated in the following manner.

Example: Toluene

$$\text{RF} = \frac{\text{concentration}}{\text{area} \times 5}$$

$$\text{RF} = \frac{31.9}{2.044 \times 5}$$

$$\text{RF} = 3.12$$

Figure 6 shows an example of a chromatograph that was used in calculating the response factors for hydrocarbons C<sub>6</sub> - C<sub>10</sub>. It was observed that although equivalent amounts of each hydrocarbon were in the final mixture, the detector of the P.E. 900 indicated varying integrated peak areas. Table 1 shows experimentally determined response factors for C<sub>2</sub> - C<sub>10</sub> hydrocarbons. In instances where the pure compound could not be obtained for analysis, the response factors were estimated (Table 2).




FIGURE 6. QUANTITATIVE STUDY OF HYDROCARBONS  $C_5$  -  $C_{10}$  6/17/76

TABLE 1. EXPERIMENTALLY DETERMINED RESPONSE FACTORS  
FOR C<sub>2</sub> - C<sub>10</sub> HYDROCARBONS

<u>COMPOUNDS</u>	<u>RESPONSE FACTOR (RF)</u>
Ethane	3.91
Ethylene	3.85
Propane	4.12
Acetylene	4.84
Isobutane	4.50
N-Butane	4.16
Propylene	4.36
Iso-Pentane	4.11
N-Pentane	4.10
N-Hexane	4.24
Trans 2-Hexene	3.72
Cis 2-Hexene	3.72
Cyclobexane	3.74
N-Heptane	2.98
Toluene	3.12
Octane	3.26
Ethyl Benzene	3.39
Methyl Xylene	3.61
Ortho Xylene	3.67
Nonane	2.99
N-Propyl Benzene	3.94
Mesitylene	3.88
N-Decane	4.34
N-Butyl Benzene	4.16

TABLE 2. ESTIMATED RESPONSE FACTORS FOR C<sub>6</sub> - C<sub>10</sub> HYDROCARBONS

<u>COMPOUNDS</u>	<u>RESPONSE FACTOR (RF)</u>
Olefins C <sub>6</sub> to C <sub>10</sub>	3.70
Paraffins C <sub>6</sub> to C <sub>10</sub>	3.50
Alkyl Aromatics	3.80



#### 4.0 ANALYSIS OF RAMS STATIONS SAMPLES DURING THE SUMMER AND FALL INTENSIVES OF 1976

During the period of performance of Task Order 113, the gas chromatography laboratory analyzed atmospheric samples from various RAMS stations. Sampling commenced on 23 June 1976 with a collection of two or four samples per site, five sites per day, two days per week. The choice of sites and times of sampling were determined by the EPA Task Coordinator. The sampling period ended on 18 November 1976.

##### 4.1 SELECTION OF SAMPLING BAGS AND TANKS

Ambient air samples were collected in Teflon bags and stainless steel tanks. The 100 litre Teflon bags (5 mil thickness) were checked for leaks by filling the bags with approximately 60 litres of helium, then going over them thoroughly (especially the seams and Teflon fittings) with a Gow Mac helium leak detector. The detector was zeroed, then the detector's nozzle was passed over the bag. If the gauge remained at zero, the bag was leak free; but, if the gauge moved from zero, the bag contained a leak. Leaks that occurred around the fittings were repaired by tightening them. Bags with leaks in their seams were set aside to be repaired at RTP in Durham, N.C. The "leak free" bags, filled to capacity with helium, were left on the shelves overnight, then again checked for leaks. These bags were purged three times with helium then filled with approximately 60 litres of ultrapure air and again left on the shelves overnight. The bags were analyzed on the Beckman 6800 for THC, CH<sub>4</sub> and CO and the P.E. 900 for C<sub>2</sub> - C<sub>10</sub> hydrocarbons. Bags possessing more than 2 ppbC of hydrocarbons were put aside; those having less than 2 ppbC hydrocarbon contamination were placed into circulation to be used in the RAMS sampling system. A supply of at least 25 "good bags" were kept in the system at all times. Bags were replaced in the system as needed. Figures 7 and 8 show the background chromatographs of a typical good bag.




FIGURE 7. BAG 17 ANALYZED ON THE PHENYL ISOCYANATE COLUMN 6/9/76




FIGURE 8. BAG 17 ANALYZED ON SQUALANE COLUMN 6/19/76

During the 1976 fall intensive and special studies periods, four 8 litre stainless steel tanks were introduced into the sampling system. The tanks were fitted with a vacuum gauge on the rear and a two way needle valve on the front. The tanks were purged with helium, pressurized to approximately 30 psi with ultrapure air, and left to stand overnight in order to ascertain the amount of leakage and the degree of hydrocarbon contamination. The contents of the tank were analyzed on the P.E. 900 gas chromatograph the following day. The air in the tanks was removed by attaching a regulator with a quick connect fitting to the two-way valve. The valve was opened all the way so that fractionating of the gases would not occur and the regulator was partially opened. The quick connect fitting was joined to its counterpart on the P.E. 900 in injection mode (see Figure 5). Three of the four tanks were found suitable for sampling. Figure 9 shows an example of a chromatogram of a hydrocarbon-free stainless steel tank.

## 4.2 AMBIENT AIR SAMPLING

The gas chromatography laboratory was responsible for the bag preparation, placing bags at the RAMS stations, sample transportation, sample analyses, data validation and computer tape and printout distribution of the data. These were accomplished as follows:

### 4.2.1 Bag Preparation and Sample Collection

In order for the Teflon bags to be in place at the RAMS stations at 0600 hours CST, they were purged three times with helium and stored in a box at 0400 on the sampling day. The bags were transported to the various sites in a van and placed in the bag boxes at the RAMS stations. Figure 10 shows the sampling system and the flow diagram of the system at each site. A glass wool filter was used for the removal of ozone and particulants.

One bag was placed at each site in each bag box and upon command from the central computer at 0600 CST, the port between 1 and 2 would open on box 1 (see Figure 10) and air would flow through for two hours at an approximate rate of 833 cc/minute. (This rate varied somewhat from station to station owing to the idiosyncrasy of the equipment at that station.) The port between 1 and 2 would open and ambient air would flow through 2




FIGURE 9. BACKGROUND OF STAINLESS STEEL TANK




FIGURE 10. RAMS BAG SAMPLING SYSTEM FLOW DIAGRAM

to 1, then into the bag. Although port 3 was closed, the vacuum pump continued to evacuate the air in the air box; this caused the ambient air to fill the bag.

After two hours, a command from central would shut off the air flow on box 1 and turn on the air flow of box 2 and the procedure would be repeated.

The 0600 ambient air samples were removed and an empty bag put in its place in order to collect a sample at 1000-1200 hours. This procedure continued until the final samples were collected at all stations at 1200-1400 hours. The ambient air samples, protected from direct sunlight, were transported back to the laboratory for analyses.

#### 4.3 AMBIENT AIR SAMPLING

Ambient air samples, which were collected in the 100 litre Teflon bags at the RAMS stations over a two hour period of time, were analyzed for THC, CH<sub>4</sub> and CO on the Beckman 6800 and for hydrocarbons C<sub>2</sub> - C<sub>10</sub> on the Perkin Elmer 900 (procedures are described in Appendices A and B). Initially all bag samples were analyzed for all compounds but this proved too difficult due to the late arrivals of the 0600-0800 and 0800-1000 samples at the laboratory (13:00-14:00 hours), the number of samples (maximum of 16) and the time required to complete the squalane analysis (1.5 hours including printout and turnaround times). The following changes were made by the EPA Task Coordinator: ambient air samples that were collected at 0600-0800 hours were analyzed for THC, CO and C<sub>1</sub> - C<sub>10</sub> and subsequent bags were analyzed for THC, CO and C<sub>1</sub> - C<sub>5</sub>. Table 3 shows the number of ambient air samples analyzed by the gas chromatography laboratory during this Task Order period. The analyses commenced immediately upon arrival of the bags at the laboratory and continued until all analyses were complete for all samples, which required the operation of two shifts at the laboratory.

#### 4.4 DATA VALIDATION, INTERPRETATION AND DISTRIBUTION

Each chromatogram of ambient air samples of the summer and fall intensive was visually analyzed by the EPA Task Coordinator and laboratory personnel working together. At the conclusion of the Intensive and data validation, the arithmetic mean and standard deviation of the concentrations of all time segments were calculated for total hydrocarbons (THC), methane (CH<sub>4</sub>),

TABLE 3. AMBIENT AIR SAMPLES ANALYZED IN THE  
GAS CHROMATOGRAPHY LABORATORY 1976

<u>MONTH</u>	<u>NUMBER</u>
June	20
July	98
August	96
September	9
October	0
November	<u>69</u>
TOTAL	292



carbon monoxide (CO) and acetylene ( $C_2H_2$ ). Table 4 shows data on a station by station basis. The sums of the parafins, olefins, aromatics and total non-methane hydrocarbons per sample were calculated and recorded. The validated data, on one copy of a 600 foot, 9 track, 800 BPI, odd parity magnetic data tape, along with two copies of the printout were given to the EPA Task Coordinator. Appendix C contains some of the results of the data collected during the summer and fall intensives.

The carbon monoxide to acetylene ratios were computed from the arithmetic mean for each station where acetylene was consistently analyzed. These data are shown in Table 5. It should be noted that acetylene was not present in sufficient quantities to be analyzed at all times for many of the RAMS sites.

TABLE 4. SUMMARY OF SUMMER AND FALL INTENSIVES 1976\*

RAMS STATIONS	THC		CH <sub>4</sub>		CO		C <sub>2</sub> H <sub>2</sub>	
	$\bar{X}$	S <sub>x</sub>	$\bar{X}$	S <sub>x</sub>	$\bar{X}$	S <sub>x</sub>	$\bar{X}$	S <sub>x</sub>
Summer 101	3283	1137	1880	193	1101	530	12	8
Summer 102**	2660	306	1972	97	935	214		
Summer 103	2328	424	1860	273	763	412	4	2
Summer 108	2223	270	1872	98	603	388		
Summer 113	1874	78	1912	82	957	167		
Summer 114	2613	912	1851	308	599	302	7	9
Summer 115	2427	908	1788	147	498	213	3	2
Summer 118	2791	1310	1770	133	479	301	3	2
Summer 121	1960	311	1872	44	364	183		
Summer 122	2939	1771	1864	212	607	240	5	2
Summer 124	2264	382	1931	402	579	214	3	1
Fall 101	2906	788	1734	240	1815	1322	33	26
Fall 103	2676	817	1739	367	1073	844	15	14
Fall 114	2451	1299	1541	581	1187	1392	19	22
Fall 115	2321	417	1637	225	666	388	9	7

$\bar{X}$  Mean  
S<sub>x</sub> Standard Deviation

\* Concentrations in ppbC

\*\* Values artificially high due to small amount of data

TABLE 5. AVERAGE CARBON MONOXIDE TO ACETYLENE RATIOS

<u>RAMS STATION</u>	<u>RATIO OF CO TO C<sub>2</sub>H<sub>2</sub></u>
Summer 101	96
Summer 103	177
Summer 114	84
Summer 115	143
Summer 118	142
Summer 122	127
Summer 124	196
	$\bar{X} = 138 \quad S_x = 40$
Fall 101	56
Fall 103	72
Fall 114	62
Fall 115	71
	$\bar{X} = 65 \quad S_x = 8$

## 5.0 SPECIAL STUDIES

Special studies and audits were carried out by the gas chromatography laboratory, as requested by the EPA Task Coordinator. These included road-way sampling, ethylene concentration verifications and audits on various gas samples.

### 5.1 COLLECTION AND ANALYSES OF ROADWAY SAMPLES

The EPA Task Coordinator requested and collected roadway samples for the gas chromatography laboratory to analyze THC, CO and  $C_1 - C_{10}$  hydrocarbons. A mobile air monitoring laboratory was used as the vehicle from which the samples were collected from heavily traveled thoroughfares during the morning rush hours. The bags used as containers for ambient air samples were purged three times with helium, filled with approximately 60 litres of helium, then left with the EPA Task Coordinator the day prior to sampling. Prior to sampling, a metal bellows pump was used to purge the bag of the helium, fill the bag with approximately 60 litres of ambient air, then purge the bag again. The 50-60 litre ambient air sample for analysis was collected by using the metal bellows pump with a Teflon sample line (~ 1.2 m long) extended out of the driver's side of the mobile laboratory for periods of 5-10 minutes. The samples were transported back to the laboratory for analysis. These roadway samples were collected mainly from I-270, I-40, Olive Road, Manchester Road and Babler Park. Detailed analytical results are contained in Appendix D.

The carbon monoxide to acetylene ratios were computed for each sample collected. The arithmetic mean of the ratio, 68.5, (Table 6) is similar to the carbon monoxide to acetylene ratio ( $63.4 \pm 6.1$ ) computed for automobile emissions in the Lincoln Tunnel by EPA researchers (5).

TABLE 6. CARBON MONOXIDE TO ACETYLENE RATIOS OF ROADWAY SAMPLES

<u>Location</u>	<u>Date</u>	<u>Ratio of CO to C<sub>2</sub>H<sub>2</sub></u>
Manchester Road	9/22/76	70
I-40	9/22/76	78
I-270	9/22/76	60
N on I-270 (I-40 to Olive)*	9/27/76	(41)
W on Manchester	9/27/76	55
W on Olive	9/27/76	76
N on I-270 Olive to Page	9/27/76	83
W on Manchester (I-270 to Weidman Road)	10/29/76	60
W on Manchester Weidman Road to Ries Rd.	10/29/76	61
S on I-270 (Manchester to I-44)	10/29/76	67
N on I-270 (I-44 to Manchester)	10/29/76	75
		$\bar{X} = 68.5 \quad S_x = 9.35$
Babler Park**	9/22/76	96
Olive to Eatherton (west of airport)**	9/22/76	61

\* Extremely high acetylene - not used in determining  $\bar{X}$  and  $S_x$

\*\*Rural Samples

In order to ascertain the quality of data obtained from ambient air samples stored in Teflon bags over a period of time, the EPA Task Coordinator requested that some of the roadway samples be analyzed on the day collected and again 48 and 168 hours later. Table 7 shows samples collected from Manchester Road, I-40 and I-270 at the requested times.

## 5.2 ETHYLENE STUDIES

Ethylene studies were carried out at RAMS stations 101 and 124 on 20, 22, 28 and 31 October 1976 and 1 and 4 November 1976 because of periodic abnormally high concentrations of that compound found in these stations during the summer intensive. Ethylene was of particular interest because of its use with the ozone analyzers at the RAMS stations. For these studies, the ambient air samples were collected in 100 litre Teflon bags and in 8 litre stainless steel tanks. The samples were taken from the following locations on various days:

- A. From RAMS station bag box with a sample taken outside simultaneously, 10-17 meters upwind and 2-3 meters above the ground using a metal bellows pump.
- B. Inside the station between the pump box and bag box.
- C. From the outlet of the center most blower of the pump box.
- D. From the outlet of the vacuum pump in pump box.

The bags and tanks were immediately transported back to the laboratory and analyzed on the phenyl isocyanate column of the Perkin Elmer Model 900 gas chromatograph for  $C_2$ - $C_5$  hydrocarbons. Detailed analytical results are contained in Appendix D.

## 5.3 SPECIAL AUDITS

At the request of the EPA Task Coordinator, the cylinders MM11435, MM11437, MM11436 (from Scott-Marrin) and cylinder number FF3753 (from AIRCO Industrial Co.) were analyzed for CO, THC and  $CH_4$  and RSG-80-4802 and RSG-80-7384 (from AIRCO Industrial Co.) for isobutane and normal hexane respectively. The audits were done in the following manner.

TABLE 7. STORAGE OF ROADSIDE SAMPLES IN TEFLON BAGS\*

	MANCHESTER ROAD			INTERSTATE 40		INTERSTATE I-270		
	†C9/22/76							
	§A9/22/76	A9/24/76	A9/28/76	A9/22/76	A9/24/76	A9/22/76	A9/24/76	A9/28/76
<u>Dura Pak</u>								
C <sub>2</sub> H <sub>6</sub>	22.3	22.2	24.0	18.9	19.1	42.0	47.1	44.9
C <sub>2</sub> H <sub>4</sub>	157.1	148.6	158.5	127.3	121.0	253.0	280.1	254.4
C <sub>3</sub> H <sub>8</sub>	27.6	26.7	29.0	14.4	14.2	54.8	61.5	56.8
C <sub>2</sub> H <sub>2</sub>	111.5	99.5	105.4	83.1	78.0	190.1	208.2	185.0
IsoC <sub>4</sub> H <sub>10</sub>	15.2	14.5	15.8	11.3	11.1	19.8	24.2	22.0
N-C <sub>4</sub> H <sub>10</sub>	91.0	88.0	94.0	68.0	66.0	131.0	149.0	137.0
C <sub>3</sub> H <sub>6</sub>	73.7	69.8	74.2	60.3	57.3	112.7	125.0	113.5
IsoC <sub>5</sub> H <sub>12</sub>	125.0	119.0	130.0	84.0	81.0	172.0	191.0	179.0
N-C <sub>5</sub> H <sub>12</sub>	68.0	64.0	72.0	41.0	38.0	86.0	85.0	94.0
<u>Squalane</u>								
N-C <sub>6</sub> H <sub>14</sub>	46.0	41.0		64.0	27.0	38.0	87.0	
N-C <sub>7</sub> H <sub>16</sub>	15.0	15.0		32.0	14.0	51.0	28.0	
Toluene	72.0	71.0		165.0	67.0	239.0	140.0	
N-C <sub>8</sub> H <sub>18</sub>	12.0	14.0		16.0	12.0	45.0	25.0	
E-C <sub>6</sub> H <sub>5</sub>	30.0	36.0		68.0	37.0	107.0	66.0	
M-XYL	77.0	93.0		136.0	90.0	277.0	131.0	
O-XYL	30.0	36.0		65.0	32.0	91.0	56.0	
N-C <sub>9</sub> H <sub>20</sub>	4.0	4.0		4.0	1.0	21.0	10.0	
N-P-C <sub>6</sub> H <sub>5</sub>	5.0	8.0		6.0	2.0	18.0	10.0	
N-C <sub>10</sub> H <sub>22</sub>	16.0	9.0			5.0	41.0	25.0	

\* Concentration in ppbC

† C - Collected

§ A - Analyzed

### 5.3.1 Total Hydrocarbon, Methane and Carbon Monoxide Audits

A five point calibration (Figure 11) was performed on the Beckman 6800 as described in Appendix A. The tanks to be audited were analyzed on the gas chromatograph and their concentrations determined from the 5-point calibration curve.

The results of the audit on 10/21/76 are shown in Tables 8 and 9.

TABLE 8. CARBON MONOXIDE AUDITS

Tank #	Manufacturer's Conc. in ppm	Rockwell Conc. in ppm	% Difference Rockwell vs. Manufacturer	EPA Conc. in ppm	% Difference Rockwell vs. EPA
MM11435	15.35	16.24	5.5	15.56	4.2
MM11436	8.00	8.31	3.7	8.17	1.7
MM11437	5.02	5.23	4.0	5.16	1.3
*FF3753		21.62		20.32	6.0
L-2359	5.06	5.33	5.1	5.29	0.7
L-1749	5.11			5.34	

\*Tank prepared by EPA at RTP

TABLE 9. METHANE AUDITS

Tank #	Manufacturer's Conc. in ppm	Rockwell Conc. in ppm	% Difference Rockwell vs. Manufacturer	EPA Conc. in ppm	% Difference Rockwell vs. EPA
MM11435	8.04	7.81	2.9	8.35	6.5
MM11436	4.98	4.83	3.0	5.23	7.6
MM11437	2.02	1.95	3.5	2.08	6.3
*FF3753		5.85		6.33	7.6
L-1749	1.99	1.81	9.0	2.09	13.4
L-2359	1.95	1.82	6.7	2.20	17.3

\*Tank prepared by EPA at RTP






FIGURE 11. FIVE POINT CALIBRATION OF BECKMAN  
MODEL 6800 GAS CHROMATOGRAPH

### 5.3.2 Isobutane, Normal Hexane and Synthetic Bag Mixture Audits

Audits were performed on AIRCO cylinders RGS-80-7384 and RGS-80-4802 containing isobutane and normal hexane respectively in nitrogen and on a synthetic mixture of carbon monoxide and methane in ultrapure air. In order to perform these audits, primary standards were prepared of  $I-C_4H_{10}$  and  $N-C_6H_{14}$  as described in Appendix B and a 5-point calibration was performed to facilitate the analysis of the synthetic bag mixture of CO and  $CH_4$  which was prepared by the EPA Task Coordinator. The results are given in Table 10.

TABLE 10. ISOBUTANE, N-HEXANE AND SPECIAL BAG AUDITS

	AIRCO RGS-80-7384	AIRCO RGS-80-4802
	Isobutane	Hexane
	Conc. in ppm	Conc. in ppm
Manufacturer	57.0	50.0
Rockwell	62.1	36.5
EPA	58.4	47.0

The results of the special bag samples are:

	Measured	Value of EPA Std.
$CH_4$	4.5	4.59
CO	8.9	9.32

### 5.4 SPECIAL STUDIES AT RAMS 103 AND 107

Special studies were conducted at RAMS stations 103 and 107 in order to compare concentrations of compounds collected simultaneously in Teflon bags and stainless steel tanks; determine contamination at stations by collecting samples upwind; and to ascertain the accuracy of the data obtained from the gas chromatography laboratory. The samples were analyzed using the Dura Pak Column.

The bag and tank samples were collected by the same methods described in sections 4.2.1 and 4.1 respectively. The actual collection time varied from sample to sample, but bag and tank samples that were collected simultaneously

were collected over the same time interval. Tank samples were collected upwind of the stations.

Table 11 shows the results of replicate analyses of the samples, a comparison of duplicate bag samples, and a comparison of station and upwind samples.

#### 5.5 DATA VALIDATION, INTERPRETATION AND DISTRIBUTION

Each chromatogram of ambient air samples collected for the roadway, ethylene and special studies at RAMS 103 and 107 was visually analyzed by the EPA Task Coordinator and laboratory personnel jointly. The validated data on one copy of a 6800 foot, 9 track, 800 BPI, odd parity magnetic data tape, along with 2 copies of the printout were given to the EPA Task Coordinator. These data contain the sums of the paraffins, olefins, aromatics and total non-methane hydrocarbons per sample and are furnished in Appendix D of this text.

TABLE 11. QUANTITATIVE STUDIES AT RAMS 103 and 107

## A. Reproducibility of Analyses

RAMS 103

Sample 103-1

Collection Time and Date: 7:15 - 9:05, November 24, 1976

Dura Pak Column. Concentrations in ppbC

Compound	First Analysis	Repeat Analysis	Duplicate Analysis	Average	Standard Deviation
C <sub>2</sub> H <sub>6</sub>	20.0	19.7	20.2	20.0	0.25
C <sub>2</sub> H <sub>4</sub>	48.7	48.3	49.6	48.9	0.67
C <sub>3</sub> H <sub>8</sub>	56.4	55.4	57.3	56.4	0.95
C <sub>2</sub> H <sub>2</sub>	29.5	26.6	28.4	28.2	1.46
I-C <sub>4</sub> H <sub>10</sub>	16.1	16.6	17.0	16.6	0.45
N-C <sub>4</sub> H <sub>10</sub>	53.1	54.5	55.2	54.3	1.07
C <sub>3</sub> H <sub>6</sub>	9.6	11.3	12.8	11.2	1.60
I-C <sub>5</sub> H <sub>12</sub>	43.7	44.8	45.5	44.7	0.90
N-C <sub>5</sub> H <sub>12</sub>	15.1	17.2	15.7	16.0	1.08

RAMS 107

Sample 107-1

Collection Time and Date: 7:22 - 9:12, November 30, 1976

Dura Pak Column. Concentrations in ppbC

Compound	First Analysis	Repeat Analysis	Duplicate Analysis	Average	Standard Deviation
C <sub>2</sub> H <sub>6</sub>	14.7	14.5	14.8	14.7	0.17
C <sub>2</sub> H <sub>4</sub>	47.3	46.9	47.6	47.3	0.30
C <sub>3</sub> H <sub>8</sub>	12.2	11.7	12.1	12.0	0.29
C <sub>2</sub> H <sub>2</sub>	37.3	38.8	43.1	39.7	3.01
I-C <sub>4</sub> H <sub>10</sub>	9.2	8.9	9.2	9.1	0.17
N-C <sub>4</sub> H <sub>10</sub>	41.4	40.8	42.3	41.5	0.75
C <sub>3</sub> H <sub>6</sub>	13.8	12.9	14.2	13.7	0.71
I-C <sub>5</sub> H <sub>12</sub>	40.5	40.6	40.6	40.6	0.06
N-C <sub>5</sub> H <sub>12</sub>	16.2	15.7	18.3	16.8	1.40

(continued)

TABLE 11 (continued)

## B. Duplication of Samples

RAMS 107

Samples 107-1, 107-2

Collection Time and Date: 7:22 - 9:12, November 24, 1976

Dura Pak Column. Concentrations in ppbC

<u>Compound</u>	<u>Bag #1</u>	<u>Bag #2</u>	<u>Average</u>	<u>Standard Deviation</u>
C <sub>2</sub> H <sub>6</sub>	29.3	29.8	29.6	0.35
C <sub>2</sub> H <sub>4</sub>	76.2	74.8	75.5	0.99
C <sub>3</sub> H <sub>8</sub>	25.3	25.7	25.5	0.28
C <sub>2</sub> H <sub>2</sub>	72.3	72.6	72.5	0.21
I-C <sub>4</sub> H <sub>10</sub>	23.5	20.6	22.1	2.05
N-C <sub>4</sub> H <sub>10</sub>	94.0	90.4	94.2	2.54
C <sub>3</sub> H <sub>6</sub>	27.5	28.4	28.0	0.64
I-C <sub>5</sub> H <sub>12</sub>	84.9	82.3	83.6	1.83
N-C <sub>5</sub> H <sub>12</sub>	35.2	34.9	35.1	0.21

RAMS 107

Samples 107-1, 107-2

Collection Time and Date: 7:22 - 9:12, November 30, 1976

Dura Pak Column. Concentrations in ppbC

<u>Compound</u>	<u>Bag #1</u>	<u>Bag #2</u>	<u>Average</u>	<u>Standard Deviation</u>
C <sub>2</sub> H <sub>6</sub>	14.8	14.7	14.8	0.07
C <sub>2</sub> H <sub>4</sub>	47.3	56.1	51.7	6.22
C <sub>3</sub> H <sub>8</sub>	12.2	12.9	12.6	0.49
C <sub>2</sub> H <sub>2</sub>	37.3	38.1	37.7	0.56
I-C <sub>4</sub> H <sub>10</sub>	9.2	8.8	9.0	0.28
N-C <sub>4</sub> H <sub>10</sub>	41.4	41.1	41.3	0.21
C <sub>3</sub> H <sub>6</sub>	13.8	13.4	13.6	0.28
I-C <sub>5</sub> H <sub>12</sub>	40.6	40.1	40.4	0.35
N-C <sub>5</sub> H <sub>12</sub>	16.2	15.9	16.1	0.21

(continued)

TABLE 11 (continued)

## C. Comparison of Upwind and Station Data

RAMS 103

Samples 103-1 (average), 103 Tank

Collection Time and Date: 7:15 - 9:05, November 23, 1976

Upwind Sample Collected in Stainless Steel Tank

Dura Pak Column. Concentrations in ppbC

Compound	Station Sample (average)	Upwind Sample (S.S. Tank)	% $\Delta$
C <sub>2</sub> H <sub>6</sub>	20.0	19.5	2.5
C <sub>2</sub> H <sub>4</sub>	48.9	27.5	43.7
C <sub>3</sub> H <sub>8</sub>	56.4	52.3	7.3
C <sub>2</sub> H <sub>2</sub>	28.2	26.7	5.3
I-C <sub>4</sub> H <sub>10</sub>	16.6	14.1	15.1
N-C <sub>4</sub> H <sub>10</sub>	54.3	47.0	13.4
C <sub>3</sub> H <sub>6</sub>	11.2	11.1	0.9
I-C <sub>5</sub> H <sub>12</sub>	44.7	41.7	6.7
N-C <sub>5</sub> H <sub>12</sub>	16.0	15.0	6.2

RAMS 107

Samples 107-1 (average), 107 Tank

Collection Time and Date: 7:22 - 9:12, November 24 and 30, 1976

Upwind Sample Collected in Stainless Steel Tank

Dura Pak Column. Concentrations in ppbC

Compound	11/24/76			11/30/76		
	Station Sample	Upwind Sample	% $\Delta$	Station Sample	Upwind Sample	% $\Delta$
C <sub>2</sub> H <sub>6</sub>	29.6	30.2	-2.0	14.7	15.1	-2.7
C <sub>2</sub> H <sub>4</sub>	75.5	63.8	15.5	50.2	33.7	32.8
C <sub>3</sub> H <sub>8</sub>	25.5	25.1	1.6	12.3	12.4	-0.8
C <sub>2</sub> H <sub>2</sub>	72.5	76.2	-5.1	39.3	38.8	1.3
I-C <sub>4</sub> H <sub>10</sub>	22.1	19.4	12.2	9.0	10.7	-1.9
N-C <sub>4</sub> H <sub>10</sub>	94.2	85.2	9.6	41.4	42.3	-2.2
C <sub>3</sub> H <sub>6</sub>	28.0	27.3	2.5	13.6	12.3	9.6
I-C <sub>5</sub> H <sub>12</sub>	83.6	76.7	8.3	40.5	41.7	-3.0
N-C <sub>5</sub> H <sub>12</sub>	35.1	31.8	9.4	16.6	15.0	9.6

## 6.0 QUALITY CONTROL

To insure accurate results for all analyses done in the laboratory, quality control procedures were employed. The instruments were standardized using standard gases from Scott-Marrin for the Beckman 6800 and accurately prepared laboratory standards for the P.E. 900. (See Appendix B, Sec. 1.3) Records of area sizes and peak heights were maintained and checked daily, duplicate and repeat analyses were performed on the samples that were collected.

The first daily sample was analyzed three times in order to insure that the Perkin Elmer 900 was functioning properly. The first repeat analysis came directly after the original analysis, to provide a check on reproducibility of the machine; the second duplicate analyses were performed at the end of the day in order to ascertain whether the gas chromatograph maintained its stability. These duplications were usually within  $\pm 5\%$  of each other; if not, the samples were reanalyzed.

### 6.1 STANDARDS FOR BECKMAN 6800

Scott-Marrin gases were chosen as standards because of the good reputation of the company and the availability of the THC, CH<sub>4</sub>, and CO gases in stable all-aluminum cylinders. EPA Quality Assurance Branch at the Research Triangle Park provided the calibrations used to determine the concentrations of the gases used as primary and secondary standards.

### 6.2 STANDARDS FOR PERKIN ELMER 900

Appendix B describes the methods employed to prepare standards. Propane was used on the phenyl isocyanate column and toluene on the squalane column in the P.E. 900 gas chromatograph. The gases were stored in stainless steel cylinders in order to insure their stability.

### 6.3 DAILY STANDARDS

A tank containing laboratory air was used as a secondary standard.

Its composition was checked against the primary standards. The tank was refilled periodically because of the size of the cylinder and the volume of gas used per analysis, reanalyzed, then standardized against the primary standards.

Tables 12 and 13 show that from 20 July 1976 through 30 November 1976, the standard deviation of the analysis of propane, in two tank standards, were 0.08 ppbC and 1.07 ppbC with a standard error of 0.02 ppbC and 0.32 ppbC, respectively. For the same period, toluene exhibits a standard deviation of 0.58 ppbC and a standard error of 0.15 ppbC.

The comparison of areas of the standards was used as a quality control criterion as well as to detect the presence of leaks in the system. This approach provided a high level of confidence for the work done over this task order period.



TABLE 12. PROPANE STANDARD\*

DATE	AREA	DATE	AREA
7/20/76	14.7193	8/24/76	2.2904
7/22/76	13.7971	8/26/76	2.3351
7/27/76	16.7500	8/31/76	2.3192
7/29/76	16.7449	9/2/76	2.3775
8/3/76	16.7033	9/22/76	2.3649
8/4/76	17.0707	9/24/76	2.4480
8/5/76	16.9472	9/27/76	2.3218
8/10/76	17.1686	9/28/76	2.2637
8/12/76	16.7462	10/20/76	2.1580
8/17/76	16.5964	10/22/76	2.1924
8/19/76	16.7571	10/28/76	2.2105
	$\bar{X} = 16.36$	10/29/76	2.2111
	$S_x = 1.07$	11/1/76	2.1606
	$S_x^2 = 0.32$	11/2/76	2.2017
		11/3/76	2.1917
		11/4/76	2.2104
		11/8/76	2.2781
		11/10/76	2.2583
		11/12/76	2.2524
		11/16/76	2.1953
		11/18/76	2.2167
		11/24/76	2.1489
		11/30/76	2.2672
			$\bar{X} = 2.26$
			$S_x = 0.08$
			$S_x^2 = 0.02$

\* These areas are directly proportional to the response factors and the concentrations. Areas in arbitrary units.

TABLE 13. TOLUENE STANDARD\*

DATE	AREA	DATE	AREA
7/20/76	0.9344	8/26/76	12.9900
7/22/76	1.0363	8/31/76	13.2486
7/27/76	1.8308	9/2/76	13.0400
8/3/76	1.4842	9/22/76	13.0912
8/5/76	1.6171	9/24/76	13.1302
8/12/76	0.9089	9/27/76	13.2902
8/17/76	1.4664	10/29/76	13.1078
8/19/76	0.8082	11/8/76	12.6534
	$\bar{X} = 1.26$	11/10/76	12.6560
	$S_x = 0.38$	11/12/76	12.5260
	$S_{\bar{x}} = 0.14$	11/16/76	12.1836
		11/18/76	13.0540
		11/29/76	14.1280
		11/30/76	14.3846
			$\bar{X} = 13.1100$
			$S_x = 0.58$
			$S_{\bar{x}} = 0.15$

\* These areas are directly proportional to the response factors and the concentrations. Areas in arbitrary units.

## 7.0 DATA PROCESSING

Data generated by the RAPS Gas Chromatography Laboratory were processed and submitted to the RAPS Computer Data Bank, Research Triangle Park (RTP), North Carolina. Data processing from analysis to submission to the data bank was performed as described in the following sections.

### 7.1 DATA TABULATION

It was planned to perform approximately thirty-two analyses per week for up to sixty components. The data were initially recorded in the form of strip chart chromatograms, punched tape and/or teletype printouts. Next, the data were given a first quality review by visually inspecting the data for general chromatographic form. Quantitative values for each component were then established. Following review and approval, the data were tabulated on a special preprinted form for keypunching.

### 7.2 KEYPUNCHING AND PROCESSING

At the end of the analysis period, the data forms were presented to the EPA Task Coordinator for review and approval and subsequently keypunched and keypunch validated. Key punching errors were corrected by computer operators at the RAMS Computer Facility.

Data processing entailed checking the cards for index number consistency, as provided for by the form, and then producing a triple copy printout of labelling information: the name, code number, concentration (PPB), ratio relative to CO, and flags if the concentration or ratio was outside an upper and lower set of limits provided by EPA. Four quantities aggregated by the software were treated as components in all respects: sum of nonmethane paraffins, olefins, aromatics, and nonmethane hydrocarbons. Validation of the data was carried out by visual inspection and comparison of the data with the chromatogram and original tabulated data. Also, special attention was directed to flagged data for validity and proper annotation.

Upon completion of data validation, three copies of a 600 foot, 9 track, 800 BPI, odd-parity magnetic data tape were prepared. One copy was sent to RTP, along with a copy of the printout. The other copy of the tape and a printout was delivered to the EPA Task Order Coordinator (St. Louis) and a third copy of the printout and tape retained by the RAMS Central Computer Facility.

#### LITERATURE CITED

1. Mindrup, R. F., RAPS Final Report, Task Order 53, pg. 5-10.
2. Seeger, G. and Hern, D., RAPS Final Report, Task Order 103, pg. 20-30.
3. McReynolds, W. O., Gas Chromatograph Retention Data, pg. 22-26, 154-155.
4. Ettre, L. S., Open Tubular Columns, Perkin Elmer 1973, pg. 32-33.
5. Lonneman, et al, Environmental Science and Technology, Vl. 8, pg. 231.

APPENDIX A  
OPERATION OF BECKMAN 6800 GAS CHROMATOGRAPH

## CONTENTS

A.0	Total Hydrocarbons, Carbon Monoxide and Methane Analyses. . . . .	.48
A.1	Reproducibility of the Beckman 6800. . . . .	.48
A.2	Linearity of Detector Response . . . . .	.48
A.3	Primary Standard . . . . .	.49
A.4	Secondary Standards. . . . .	.49
A.5	Calculations . . . . .	.49

TABLES

<u>Number</u>		<u>Page</u>
A-1	Tank L-2327 - Tank Standard for 6800 . . . . .	51



## A.0 TOTAL HYDROCARCONS, CARBON MONOXIDE AND METHANE ANALYSES

Analyses of CO, THC and CH<sub>4</sub> were done on a Beckman Model 6800 Gas Chromatograph in the following manner:

The recorder was turned on, set at 10 mv range and zeroed with the zero volt button. The auto-zero switch was actuated in order to zero the electrometer of the chromatograph. The pump on the back of the chromatograph was turned on and a sample bag was connected to commence analyses.

To analyze total hydrocarbons, the attenuation was normally set at 4 and the range at 10 and the valve B toggle switch was actuated for approximately 15-20 seconds. The recorder was allowed to return to zero, aided with the auto-zero toggle switch. The attenuation was changed from 4 to 1 for the methane and carbon monoxide analyses and the valve A toggle switch actuated for 45 seconds.

### A.1 REPRODUCIBILITY OF THE BECKMAN 6800

Reproducibility of the results of the analyses of THC, CH<sub>4</sub> and CO was checked by running duplicate analyses; that is, the first sample and the standard were rerun after the analysis of all other samples. During the course of this task order, these duplicates agreed closely with the initial analysis (+5%).

### A.2 LINEARITY OF DETECTOR RESPONSE

Five point calibrations were carried out in duplicate to insure that the detector response was linear with respect to sample concentration. The tests were performed for THC, CH<sub>4</sub> and CO. These checks were determined from synthetic bag mixtures prepared in the laboratory. The procedure was as follows:

Five bags were prepared. The bags were labeled and filled with 100 liters of zero air using a mass flow meter. The appropriate amounts of CH<sub>4</sub> and CO were injected into each bag using a gas-tight calibrated syringe and allowed to diffuse throughout the bag. The bags were then analyzed. The make-up of the mixtures is shown below:

<u>BAG #</u>	<u>ZERO AIR</u>	<u>CH<sub>4</sub></u>	<u>CO</u>	<u>CONC.</u>
1	100 litres	0.1 ml	0.1 ml	1 ppm
2	100 litres	0.2 ml	0.2 ml	2 ppm
3	100 litres	0.3 ml	0.3 ml	3 ppm
4	100 litres	0.4 ml	0.4 ml	4 ppm
5	100 litres	0.5 ml	0.5 ml	5 ppm

### A.3 PRIMARY STANDARD

A cylinder containing small amounts of methane and carbon monoxide in air (Scott-Marrin L 1749) was analyzed by the EPA at Research Triangle Park, North Carolina. Its concentrations was computed to be 5.33 ppm CO and 1.99 ppm CH<sub>4</sub>. This cylinder served as the primary standard for the Beckman 6800 and was used to calibrate other tanks.

### A.4 SECONDARY STANDARDS

Two other cylinders (Scott-Marrin L 2327 and L 2359) served as secondary standards for the Beckman 6800 during this task order period. When analyzed against the primary standard Scott-Marrin L 1749, their concentrations were computed to be 5.28 ppm CO, 4.38 ppm CH<sub>4</sub> and THC for Scott L 2327 and 5.29 ppm CO and ppm CH<sub>4</sub> and 2.20 ppm CH<sub>4</sub> and THC for Scott L 2349. The secondary standards were checked against the primary ones once a month.

### A.5 CALCULATIONS

Calculations of the concentrations of THC, CH<sub>4</sub> and CO in various samples are straight forward and were done by hand. When analyses were to be performed, the secondary standard was analyzed; that is, its peak height was measured. Since its concentration was known, it was possible to compute the concentration of unknown once its peak height was measured:

$$\frac{\text{Standard concentration in ppm}}{\text{Peak height of standard in mm}} \times \text{peak height of unknown in mm} = \text{concentration of unknown}$$

Initially, a sample of standard gas was injected into a Teflon bag, then analyzed on the Beckman 6800. It was observed that the peak height varied as much as 42.6% for THC, 9.7% for CH<sub>4</sub> and 58% for CO. These variations were unacceptable; therefore, another method was devised by which the standard gas was passed directly into the Beckman 6800. With this procedure, maximum peak height variation for THC was 8.4%, CH<sub>4</sub> was 14.9%, and CO was 17.9%. The standard deviations were computed to be 4.88 mm (3.8%) for THC, 7.25 mm (6.5%) for CH<sub>4</sub> and 9.89 mm (9.2%) for CO as shown in Table A-1.

TABLE A-1. TANK L-2327 - TANK STANDARD FOR 6800

DATE	THC	CH <sub>4</sub>	CO
6/23/76	129.5*	105.0	102.0
6/28/76	127.0	103.5	100.5
7/06/76	127.0	106.5	92.0
7/08/76	142.5	104.0	98.5
7/13/76	132.0	102.0	85.5
7/15/76	115.5	97.0	86.8
7/20/76	130.0	106.0	103.8
7/22/76	129.5	112.0	110.7
7/27/76	122.6	117.3	117.0
7/29/76 am	124.8	106.6	96.7
7/29/76 pm	124.7	121.8	122.4
8/03/76 am	126.4	115.1	112.8
8/03/76 pm	129.1	115.0	109.0
8/03/76	125.8	116.5	114.6
8/05/76	126.4	116.9	116.5
8/10/76 am	132.9	102.3	106.5
8/10/76 pm	130.1	112.5	106.9
8/12/76 am	128.4	120.9	121.0
8/12/76 pm	127.1	118.3	119.1
8/24/76	127.0	117.2	113.3
8/26/76 am	125.1	117.5	114.7
8/26/76 pm	125.1	118.7	115.8
	$\bar{X} = 127.7$	$\bar{X} = 111.48$	$\bar{X} = 108.00$
	$S_x = 4.88$	$S_x = 7.25$	$S_x = 9.89$
	$S^2_x = 1.04$	$S^2_x = 1.54$	$S^2_x = 2.11$

\* Response in mm

APPENDIX B  
OPERATION OF THE PERKIN ELMER 900 GAS CHROMATOGRAPH

## CONTENTS

B.0	C <sub>2</sub> - C <sub>10</sub> Analysis . . . . .	56
B.1	General Procedures . . . . .	56
B.2	Reproducibility of the Phenyl Isocyanate Column. . . . .	57
B.3	Linearity of Detector Response of Phenyl Isocyanate Column .	57
B.4	PEP-1 Computer Calculations for P.E. 900 . . . . .	62
	B.4.1 Reference Peak(s) . . . . .	62
	B.4.2 Relative Retention Times. . . . .	62
	B.4.3 Response Factor . . . . .	63
	B.4.4 Response Ractor Update. . . . .	64
B.5	Calculations Used to Determine Standard Concentrations . . .	64
B.6	Primary Standards (Propane and Toluene) P.E. 900 . . . . .	64
B.7	Secondary Standards (Propane and Toluene) P.E. 900 . . . . .	64

## FIGURES

<u>Number</u>		<u>Page</u>
B-1	Five Point Calibration of Propane on Perkin Elmer 900 Gas Chromatograph. . . . .	59
B-2	Five Point Calibration of Toluene on Perkin Elmer 900 Gas Chromatograph. . . . .	60

## TABLES

<u>Number</u>		<u>Page</u>
B-1	Five Point Calibration of Propane . . . . .	61
B-2	Five Point Calibration of Toluene . . . . .	61



## B.0 C<sub>2</sub> - C<sub>10</sub> HYDROCARBON ANALYSIS

The RAHS bag samples were analyzed for C<sub>2</sub>-C<sub>10</sub> hydrocarbons on the PE 900 gas chromatograph. The ambient air samples were analyzed and the concentrations of the compounds determined by comparing the samples to the standard. One sample was analyzed in duplicate and in some cases triplicate to insure the reproducibility of the entire set of samples and the precision and accuracy of the gas chromatograph.

### B.1 GENERAL PROCEDURES

The hydrogen and air sources were turned on and the flame ionization detectors lighted. The oven was cooled from 90°C (overnight and weekend temperature) to 25°C by turning on the liquid nitrogen. The recorders were turned on, checked for positive response, then zeroed; the vacuum line was turned on. Directives of identification and sampling time were entered on the teletype to the PEP-1 integrator.

The stainless steel concentration trap was placed in a Dewar flask which contained liquid oxygen. The appropriate vacuum line was connected to the vacuum source and the bag or tank connected to the inlet line. The vacuum source was opened and a small sample was drawn through the inlet line to purge the line of any air from a previous sample. The inlet valve was switched to the trapping mode and a measured volume of the sample was cold-trapped into the liquid oxygen cooled concentration trap.

After the sample was trapped, the ready light on the PEP-1 interface was actuated, the valve switched to the injection mode and a Dewar flask of hot water applied to the concentration trap. The "ready light" on the gas chromatograph was immediately pressed to initiate the temperature programming. The chromatograph was programmed to hold 25°C for 8 minutes. The temperature was then raised at a rate of 2°C per minute to 90°C, where it was held for the remainder of the run. The valve was returned from the injection mode to the backflush (neutral) position after 2 minutes. The analyses times were 10

minutes for the phenyl isocyanate column ( $C_2-C_5$ ) and 55 minutes for the squalane column ( $C_5-C_{10}$ ).

Upon completion of the analyses, the PEP-1 integrator tabulated the area measured, the concentration and the identification of peaks. The reset button on the gas chromatograph was pressed and the above procedure was repeated. The analyses and turnaround time for trapping sample, analysis and computer printout times for  $C_2-C_5$  were approximately 0.5 hours on the phenyl isocyanate column and 1.35 hours for  $C_5-C_{10}$  on the squalane column. The files of the memory bank of the PEP-1 computer were normally erased at the end of the working day.

## B.2 REPRODUCIBILITY OF THE PHENYL ISOCYANATE COLUMN

To ascertain and maintain reproducibility of this column, the propane standard and triplicate sample analyses were used as indicators. Any major change in the area size of the propane standard or major deviation from the concentrations of nine compounds analyzed on the column signaled problems in the system. The magnetic tapes and the computer printouts demonstrate the reproducibility of the triplicate analyses done on the first daily bag samples. The reproducibility of the daily duplicate and triplicate analyses were generally within  $\pm 5\%$ .

## B.3 LINEARITY OF DETECTOR RESPONSE OF PHENYL ISOCYANATE COLUMN

Five point calibrations were carried out on the P.E. 900 to insure that the detector response was linear with respect to sample concentration. Five mil (0.005 inch) Teflon bags were filled with ultrapure air and injected with accurately measured quantities of propane and toluene. After allowing half an hour for diffusion, samples were withdrawn from this bag ("A") and injected into five other bags, each of which had been filled with 50 l of ultrapure air according to the schedule shown on the following page:

<u>Bag #</u>	<u>Zero Air</u>	<u>C<sub>3</sub>H<sub>8</sub> (99% purity)</u>	<u>Conc.</u>	<u>Toluene (99% purity)</u>	<u>Conc.</u>
A	50 liters	5 ml of propane	100 ppm	0.05 ml toluene	226.9 ppm
1	50 litres	1 ml from Bag A	6 ppbC	1 ml from Bag A	31.9 ppbC
2	50 litres	2 ml from Bag A	12 ppbC	2 ml from Bag A	63.8 ppbC
3	50 litres	3 ml from Bag A	18 ppbC	3 ml from Bag A	95.7 ppbC
4	50 litres	4 ml from Bag A	24 ppbC	4 ml from Bag A	127.6 ppbC
5	50 litres	5 ml from Bag A	30 ppbC	5 ml from Bag A	159.5 ppbC

The bags were analyzed in descending order, then reanalyzed in the same order for duplicate analyses. Figures B-1 and B-2 and Tables B-1 and B-2 are examples of the 5 point calibrations of propane and toluene respectively.




FIGURE B-1. FIVE POINT CALIBRATION OF PROPANE ON PERKIN ELMER 900 GAS CHROMATOGRAPH




FIGURE B-2. FIVE POINT CALIBRATION OF TOLUENE ON PERKIN ELMER 900 GAS CHROMATOGRAPH

---

TABLE B-1. FIVE POINT CALIBRATION OF PROPANE

---

Bag A: 5.0 ml gas 50L of ultrapure air

Bag 1 to 5: 50 litres of ultrapure air each

Number	Amount Injected From Bag A	AREA			Conc.
		Run - 1	Run - 2	Average	
1	1.0 ml	0.0642	0.0641	0.0641	6 ppbC
2	2.0 ml	0.1068	0.1061	0.1064	12 ppbC
3	3.0 ml	0.1631	0.1652	0.1641	18 ppbC
4	4.0 ml	0.2288	0.2297	0.2292	24 ppbC
5	5.0 ml	0.3080	0.3082	0.3081	30 ppbC

---



---

TABLE B-2. FIVE POINT CALIBRATION OF TOLUENE

---

Bag A: 0.0500 ml liquid 50L of ultrapure air

Bag 1 to 5: 50 litres of ultrapure air each

Number	Amount Injected From Bag A	AREA			Conc.
		Run - 1	Run - 2	Average	
1	1.0 ml	3.5380	--	3.5380	31.9 ppbC
2	2.0 ml	4.4350	4.4036	4.4193	63.8 ppbC
3	3.0 ml	6.6937	6.4601	6.5769	95.7 ppbC
4	4.0 ml	8.5107	8.1254	8.3181	127.6 ppbC
5	5.0 ml	10.9798	10.7468	10.8633	159.5 ppbC

---

#### B.4 PEP-1 COMPUTER CALCULATIONS FOR P.E. 900

The PEP-1 computer was programmed in such a manner that it could detect, file, store, compute the concentration of selected compounds and print out pertinent data of samples that were analyzed on the P.E. 900.

In order to accurately compute the concentrations of the label compounds, the following information was input to the computer:

- 1) reference peak(s)
- 2) relative retention times for all compounds
- 3) response factor for all compounds
- 4) response factor update

##### B.1.4.1 Reference Peak(s)

N-Butane and toluene were chosen as the reference peaks for the phenyl isocyanate and squalane columns respectively because both compounds occur abundantly in ambient air samples and no large peaks appear nearby, making identification easy. When writing the programs for the phenyl isocyanate and squalane columns, n-butane was assigned a reference number of one (1) and toluene was assigned a reference number of five (5). Along with the reference numbers, time spans for the elutions of the two compounds were provided.

##### B.4.2 Relative Retention Times

Relative retention times may be obtained two ways: manual computation or the PEP-1 system. The G. C. Laboratory employs both methods; the procedure and results of both are identical. As an example, the calculation of RRT of  $C_2H_6$  is as follows:

- X = Relative retention time of n-butane = 1.0.  
T = The elution time in minutes of n-butane = 2.5 minutes.  
 $X^1$  = The unknown relative retention time of  $C_2H_6$ .  
 $T^1$  = The known elution time of the  $C_2H_6$  = 0.87 minutes.

$$\frac{X}{T} = \frac{X^1}{T^1}$$

$$\frac{1}{2.5} \times .87 = X^1$$

$$X^1 = .348$$

It should be noted here that once the reference peak has been designated and its elution time standardized, the PEP-1 automatically assigned the relative retention times to all other compounds occurring in each sample. The same procedure was used for toluene.

#### B.4.3 Response Factor

To obtain response factors (RF) for hydrocarbons  $C_2 - C_{10}$  a primary standard was made up and analyzed in the laboratory. The PEP-1 was programmed in such a manner that only the areas of the selected compounds were computed. After manually calculating the concentrations of the hydrocarbons and obtaining the various areas from the computer, it was possible to calculate the response factors for each compound.

The PEP-1 used the following formula:

Area x response factor x 5 = concentration

To calculate the response factors of the hydrocarbons:

$$RF = \frac{\text{Concentration}}{\text{Area} \times 5}$$

Example: Toluene

$$RF = \frac{31.9}{2.0440 \times 5}$$

$$RF = 3.12$$

Response factors for all selected  $C_2 - C_{10}$  hydrocarbons were determined in the above fashion.

After all of the calculations are done and the computer has been properly programmed, the PEP-1 uses; area x response factor x 5 to compute the concentrations of unknowns against the concentration of a standard.



#### B.4.4 Response Factor Update

Once the concentration of the standard was calculated and an acceptable range ( $\pm 3\%$  for propane and  $\pm 3\%$  for toluene) for the area size of each compound had been established, the response factor update feature of the PEP-1 data system was used to correct the small changes that occur in the detector from day to day. The daily computed concentrations for the propane and toluene standards were compared to the calculated value for these standards; the relative change was fed to the computer and the response factor for each compound that is analyzed is multiplied by that factor. During this task order period, the detectors showed little or no change. The changes that occurred were random with a standard deviation of less than  $\pm 3\%$ .

#### B.5 CALCULATIONS USED TO DETERMINE STANDARD CONCENTRATIONS

The preparation of primary standards and the secondary standard was done at the same time. Only one secondary standard was needed because ambient air contains the propane and toluene that was used by the Gas Chromatography Laboratory on the phenyl isocyanate and squalane columns, respectively. The PEP-1 computer system uses the area x response factor x 5 method for computing the concentrations of compounds.

#### B.6 PRIMARY STANDARDS (Propane and Toluene) P.E. 900

The primary standards for the P.E. 900 were synthetically prepared in the laboratory by using the same reagents and equipment described in Section B.3 of this appendix. It should be noted here that only one dilution bag was used.

#### B.7 SECONDARY STANDARDS (Propane and Toluene) P.E. 900

Since the hydrocarbons to be analyzed occurred in laboratory air, secondary standards were prepared by simply storing a pressurized sample of ambient air in an 8 litre stainless steel tank, equipped with the appropriate valves. A sample of the secondary standard was then analyzed and the concentration of propane and toluene computed.

APPENDIX C  
DATA FROM SUMMER AND FALL INTENSIVES

## CONTENTS

C.0 Introduction . . . . .	69
----------------------------	----

## FIGURES

<u>Number</u>		<u>Page</u>
C-1	Listing of all compounds recorded on magnetic tapes . . . .	70

## TABLES

<u>Number</u>		<u>Page</u>
C-1	RAMS Station 101 "Summer Intensive" . . . . .	73
C-2	RAMS Station 102 "Summer Intensive" . . . . .	81
C-3	RAMS Station 103 "Summer Intensive" . . . . .	82
C-4	RAMS Station 108 "Summer Intensive" . . . . .	88
C-5	RAMS Station 113 "Summer Intensive" . . . . .	89
C-6	RAMS Station 114 "Summer Intensive" . . . . .	90
C-7	RAMS Station 115 "Summer Intensive" . . . . .	95
C-8	RAMS Station 118 "Summer Intensive" . . . . .	97
C-9	RAMS Station 121 "Summer Intensive" . . . . .	99
C-10	RAMS Station 122 "Summer Intensive" . . . . .	100
C-11	RAMS Station 124 "Summer Intensive" . . . . .	103
C-12	RAMS Station 101 "Fall Intensive" . . . . .	105
C-13	RAMS Station 103 "Fall Intensive" . . . . .	108
C-14	RAMS Station 114 "Fall Intensive" . . . . .	111
C-15	RAMS Station 115 "Fall Intensive" . . . . .	114

## C.0 INTRODUCTION

Appendix C contains data obtained from samplings at RAMS stations. The results are in table form in order to provide an overview of the data from RAMS stations. The arithmetic mean ( $\bar{X}$ ) and standard deviation ( $S_x$ ) were calculated for total hydrocarbon (THC), methane ( $\text{CH}_4$ ), carbon monoxide (CO) and acetylene ( $\text{C}_2\text{H}_2$ ) for each table so that comparisons could be made between and among the RAMS stations. Not all compounds are listed in these tables. The magnetic tape of data delivered to the EPA contains the concentration for all compounds as well as sums for olefins, paraffins, etc. The following marks and abbreviations were used: "Trace" indicates less than 1 ppbC in the squalane analyses, "BDL" below detection limits, "NM" not measurable (because of interference, etc.). All concentrations in the following tables are in ppbC and the time of samples is Central Daylight Time (CDT) from April 25 to October 30 and Central Standard Time (CST) after October 30.

Figure C-1 shows the header for that tape and lists the various compounds which are reported.

001TOTAL SULFUR  
 002HYDROGEN SULFIDE  
 003SULFUR DIOXIDE  
 004METHYL MERCAPTAN  
 005NITROGEN OXIDES (NOX)  
 006SULFUR HEXAFLUORIDE  
 007TRICHLORO-FLUORO METHANE  
 008DICHLORO-DIFLUORO METHANE  
 009TOTAL ORGANICS  
 010METHANE  
 011CARBON MONOXIDE  
 012ETHYLENE  
 013ETHANE  
 014ACETYLENE  
 015PROPANE  
 016PROPYLENE  
 017ISOBUTANE  
 018ISOBUTYLENE + BUTENE-1  
 019N-BUTANE  
 020T-2-BUTENE  
 021C-2-BUTENE  
 0223-M-1 BUTENE  
 023ISOPENTANE  
 024PENTENE-1  
 0252-M-1 BUTENE  
 026N-PENTANE  
 027T-2-PENTENE  
 028C-2-PENTENE  
 0292-M-2 BUTENE  
 0302,2-DM-BUTANE  
 031CYCLOPENTANE  
 0323-M-1-PENTENE + 4-M-1-PENTENE  
 0334-M-C-2-PENTENE  
 034CYCLOPENTANE  
 0352,3-DM-BUTANE + 4-M-T-2 PENTENE  
 0362-M-PENTANE  
 0372-M-1-PENTENE  
 0383-M-PENTANE + HEXENE-1 + 2-E-1-BUTENE  
 039T-3-HEXENE  
 0402-M-2-PENTENE + 3-M-CYCLOPENTANE  
 0413-M-C-2 PENTENE  
 042N-HEXANE  
 043T-2-HEXENE  
 044C-2-HEXENE  
 0453-M-T-2-PENTENE  
 046METHYLCYCLOPENTANE + 3,3-DM-1-PENTENE  
 047BENZENE  
 0482,4-DM-PENTENE  
 0492,2,3-TM-BUTANE  
 0502,4-DM-1-PENTENE  
 0511-METHYLCYCLOPENTENE + 2-M-C-3-HEXENE  
 0522,4-DM-2-PENTENE + 3-E-1-PENTENE + 3-M-1-HEXENE  
 0532-M-T-3-HEXENE + 5-M-1-HEXENE  
 054CYCLOHEXANE + 4-M-C-2-HEXENE

(continued)

FIGURE C-1. LISTING OF ALL COMPOUNDS RECORDED ON MAGNETIC TAPES

0554-M-1-HEXENE + 4-M-T-2-HEXENE  
 0563-M-2-E-1-BUTENE + 5-M-T-2-HEXENE  
 057CYCLOHEXENE  
 0582-M-HEXENE + 5-M-C-2-HEXENE  
 0592,3-DM-PENTANE + 1,1-DM-CYCLOPENTANE  
 0603-M-HEXANE  
 0611-C-3-DM-CYCLOPENTANE + 2-M-1-HEXANE  
 0621-T-3-DM-CYCLOPENTANE + 1-HEPTENE + 2-E-1-PENTANE  
 0633-E-PENTANE + 3-M-T-2-HEXENE  
 0642,2,4-TM-PENTANE + T-3-HEPTENE  
 065C-3-HEPTENE  
 0663-M-C-3-HEXENE + 2-M-2-HEXENE + 3-M-T-3-HEXENE  
 0673-E-2-PENTENE  
 068N-HEPTANE  
 0692,3-DH-2-PENTENE + C-2-HEPTENE  
 0701-C-2-DM-CYCLOPENTANE  
 071M-CYCLOHEXANE + 2,2-DM-HEXANE + 1,1,3-TM-CYCLOPENTANE  
 0724-M-CYCLOHEXENE  
 0732,5-DM-HEXANE  
 074ETHYLCYCLOPENTANE  
 0752,4-DM-HEXANE  
 0762,2,3-TM-PENTANE  
 0771-T-2-C-4-TM-CYCLOPENTANE  
 078TOLUENE  
 0792,3,4-TM-PENTANE  
 0802,3,3-TM-PENTANE  
 0812,3-DM-HEXANE + 2-M-3-E-PENTANE  
 0822-M-HEPTANE  
 0834-M-HEPTANE  
 0843,4-DH-HEXANE + 1-C-2-T-4-TM-CYCLOPENTANE  
 0853-M-HEPTANE + 3-M-3-E-PENTANE  
 0862,2,5-TM-HEXANE + 1-C-2-C-4-TM-CYCLOPENTANE  
 0871-T-4-DM-CYCLOHEXANE  
 0881-M-T-3-E-CYCLOPENTANE  
 0892,2,4-TM-HEXANE  
 090CYCLOHEPTANE + 1-M-1-E-CYCLOPENTANE  
 0911-T-2-DM-CYCLOHEXANE + 1-C-2-C-3-TM-CYCLOPENTANE  
 092N-OCTANE  
 0931-T-3-DM-CYCLOHEXANE  
 0942,4,4-TM-HEXANE  
 0952,3,5-TM-HEXANE  
 0962,2-DM-HEPTANE  
 0972,4-DM-HEPTANE + 2,2,3-TM-HEXANE  
 0982,2-DM-5-PENTANE + 2-M-4-E-HEXANE  
 0992,6-DM-HEPTANE + 1-C-2-DM-CYCLOHEXANE  
 100N-PROPYLCYCLOPENTANE  
 101ETHYLCYCLOHEXANE  
 1022,5-DM-HEPTANE + 3,5-DM-HEPTANE  
 103ETHYLBENZENE  
 1043,3-DM-HEPTANE  
 1052,3,3-TM-HEPTANE  
 106P-XYLENE  
 107M-XYLENE  
 1084-M-OCTANE

(continued)

FIGURE C-1 (continued)



1092-M-OCTANE  
1103-E-HEPTANE  
1113-M-OCTANE  
1120-XYLENE  
1132,2,4-TM-HEPTANE  
1142,2,5-TM-HEPTANE + 2,2,5-TM-HEPTANE  
1152,5,5ITM-HEPTANE + 2,4,4ITM-HEPTANE  
116N-NONANE  
117N-PROPYLBENZENE  
1182,2,3,3-TM-HEXANE  
1191-M-2-E-BENZENE  
1201,3,5-TM-BENZENE  
121TERT-BUTYLBENZENE  
1221,2,4-TM-BENZENE  
123SEC-BUTYLBENZENE + ISOBUTYLBENZENE  
124N-DECANE  
1251,2,3-TM-BENZENE + 1-M-4-ISOPROPYLBENZENE  
126N-BUTYLBENZENE  
127PARAFFINS  
128OLEFINS  
129AROMATICS  
130TOTAL NON-METHANE HYDROCARBONS

FIGURE C-1 (continued)

TABLE C-1. RAMS STATION 101

<u>SUMMER INTENSIVE</u>							
	Mean				Standard Deviation		
THC	3720 ppbC				494 ppbC		
CH <sub>4</sub>	2016 ppbC				262 ppbC		
CO	1621 ppbC				634 ppbC		
C <sub>2</sub> H <sub>2</sub>	19.8 ppbC				11 ppbC		
Date	6/23/76 7-9	6/23/76 9-11	6/23/76 11-13	6/23/76 13-15	7/13/76 9-11	7/15/76 7-9	7/15/76 9-11
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>					8.7	8.0	5.9
C <sub>2</sub> H <sub>4</sub>					996.5	967.5	18.8
C <sub>3</sub> H <sub>8</sub>					22.8	8.2	5.3
C <sub>2</sub> H <sub>2</sub>					8.7	8.6	5.0
IsoC <sub>4</sub> H <sub>10</sub>					12.6	4.1	4.5
N-C <sub>4</sub> H <sub>10</sub>					62.6	17.7	17.3
C <sub>3</sub> H <sub>6</sub>					2.4	3.0	BDL*
IsoC <sub>5</sub> H <sub>12</sub>					55.0	22.0	14.0
N-C <sub>5</sub> H <sub>12</sub>					14.0	9.0	7.0
<u>Beckman 6800</u>							
THC	2690	2600	3550	2540	3850	3640	2730
CH <sub>4</sub>	1960	1940	1900	1900	2060	2030	2090
CO	1520	1010	1050	1220	2210	1280	1460

\* Below detection limits

(continued)

TABLE C-1 (continued)

Date	7/15/76 11-13	7/15/76 13-15	7/20/76 7-9	7/20/76 9-11	7/20/76 11-13	7/20/76 13-15	7/22/76 7-9
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	5.3	4.1	8.4	8.0	4.9	4.5	12.6
C <sub>2</sub> H <sub>4</sub>	1055.6	12.7	1345.9	30.0	1253.4	11.7	1553.2
C <sub>3</sub> H <sub>8</sub>	6.0	3.4	10.4	9.7	5.2	4.1	16.7
C <sub>2</sub> H <sub>2</sub>	4.5	6.3	10.9	5.5	4.8	2.1	19.3
IsoC <sub>4</sub> H <sub>10</sub>	3.5	4.1	7.2	6.4	5.4	4.1	15.1
N-C <sub>4</sub> H <sub>10</sub> '	12.6	15.6	32.7	24.5	32.3	21.3	72.9
C <sub>3</sub> H <sub>6</sub>	1.3	1.0	6.3	3.0	BDL*	BDL*	11.5
IsoC <sub>5</sub> H <sub>12</sub>	14.0	13.0	36.0	25.0	26.0	18.0	73.0
N-C <sub>5</sub> H <sub>12</sub>	7.0	4.6	18.0	8.0	8.0	6.0	28.5
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>			12				27
N-C <sub>7</sub> H <sub>16</sub>			5				9
Toluene			50				62
N-C <sub>8</sub> H <sub>18</sub>			15				5
E-C <sub>6</sub> H <sub>5</sub>			18				25
M-XYL			88				67
O-XYL			19				22
N-C <sub>9</sub> H <sub>20</sub>			40				2
N-P-C <sub>6</sub> H <sub>5</sub>			1				3
N-C <sub>10</sub> H <sub>22</sub>			189				5
<u>Beckman 6800</u>							
THC	3680	2470	3690	7950	3560	2270	3960
CH <sub>4</sub>	1960	1870	1920	1800	1780	1720	1920
CO	730	700	1320	926	712	610	2120

\* Below detection limits

(continued)

TABLE C-1 (continued)

Date	7/22/76 9-11	7/22/76 13-15	7/27/76 7-9	7/27/76 9-11	7/27/76 11-13	7/27/76 13-15	7/29/76 7-9
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	9.4	5.8	14.5	10.4	12.3	8.7	11.3
C <sub>2</sub> H <sub>4</sub>	33.5	21.9	1240.8	30.1	1137.4	14.3	1044.0
C <sub>3</sub> H <sub>8</sub>	9.1	5.5	23.4	18.6	12.0	9.0	12.0
C <sub>2</sub> H <sub>2</sub>	10.8	4.2	17.3	11.9	11.2	8.5	12.4
IsoC <sub>4</sub> H <sub>10</sub>	6.6	6.2	13.8	12.4	5.0	4.4	4.4
N-C <sub>4</sub> H <sub>10</sub>	27.3	28.9	45.6	36.9	22.1	23.3	24.2
C <sub>3</sub> H <sub>6</sub>	3.6	BDL*	34.0	43.0	66.0	25.0	15.0
IsoC <sub>5</sub> H <sub>12</sub>	28.0	24.0	43.0	30.0	22.0	21.0	26.0
N-C <sub>5</sub> H <sub>12</sub>	19.0	10.0	17.0	11.0	10.0	6.0	9.9
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>			10				13
N-C <sub>7</sub> H <sub>16</sub>			4				2
Toluene			48				117
N-C <sub>8</sub> H <sub>18</sub>			3				4
E-C <sub>6</sub> H <sub>5</sub>			13				30
M-XYL			40				73
O-XYL			11				24
N-C <sub>9</sub> H <sub>20</sub>			2				3
N-P-C <sub>6</sub> H <sub>5</sub>			1				2
N-C <sub>10</sub> H <sub>22</sub>							3
<u>Beckman 6800</u>							
THC	2370	2520	4010	2600	4860	2260	3630
CH <sub>4</sub>	1870	1800	1630	1480	2110	1710	2080
CO	1060	525	1130	796	758	781	1190

\* Below detection limits

(continued)

TABLE C-1 (continued)

Date	7/29/76 9-11	7/29/76 11-13	7/29/76 13-15	8/3/76 7-9	8/3/76 9-11	8/3/76 11-13	8/3/76 13-15
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	11.7	12.2	14.1	9.8	6.6	5.0	6.0
C <sub>2</sub> H <sub>4</sub>	26.9	1402.6	41.3	1345.6	22.9	1151.5	20.7
C <sub>3</sub> H <sub>8</sub>	16.5	19.2	20.8	17.7	11.4	7.2	8.6
C <sub>2</sub> H <sub>2</sub>	12.7	13.4	12.9	10.7	5.6	3.7	4.7
IsoC <sub>4</sub> H <sub>10</sub>	8.5	10.4	10.3	4.2	4.2	2.7	6.1
N-C <sub>4</sub> H <sub>10</sub>	43.3	63.6	53.0	22.7	15.1	10.6	22.2
C <sub>3</sub> H <sub>6</sub>	11.5	14.0	8.0	6.8	3.1	1.5	1.3
IsoC <sub>5</sub> H <sub>12</sub>	38.0	50.0	42.0	23.0	10.0	9.2	19.0
N-C <sub>5</sub> H <sub>12</sub>	15.0	18.0	18.0	12.0	5.0	4.3	9.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>				8			
N-C <sub>7</sub> H <sub>16</sub>				2			
Toluene				81			
N-C <sub>8</sub> H <sub>18</sub>				4			
E-C <sub>6</sub> H <sub>5</sub>				9			
M-XYL				24			
O-XYL				8			
N-C <sub>9</sub> H <sub>20</sub>				1			
N-P-C <sub>6</sub> H <sub>5</sub>				BDL*			
N-C <sub>10</sub> H <sub>22</sub>				2			
<u>Beckman 6800</u>							
THC	3320	2960	6440	4060	2470	3180	2240
CH <sub>4</sub>	2000	1740	1710	2090	1920	1680	1720
CO	1250	1070	1120	1040	754	585	645

\* Below detection limits

(continued)

TABLE C-1 (continued)

Date	8/5/76 7-9	8/5/76 9-11	8/5/76 11-13	8/5/76 13-15	8/12/76 7-9	8/12/76 9-11	8/12/76 11-13
<u>Dura Pak</u>							
$C_2H_6$	9.0	7.0	6.1	6.0	8.6	8.9	8.3
$C_2H_4$	1321.0	14.9	15.8	19.9	24.9	1864.0	43.8
$C_3H_8$	11.2	7.3	5.5	4.9	19.4	18.4	7.7
$C_2H_2$	14.4	8.3	6.5	6.2	29.7	10.3	7.4
$IsoC_4H_{10}$	8.2	5.0	6.7	5.1	14.0	14.0	5.5
$N-C_4H_{10}$	48.3	26.7	41.0	28.0	70.6	47.2	24.9
$C_3H_6$	5.1	2.5	1.0	1.1	7.3	3.4	1.8
$IsoC_5H_{12}$	49.0	29.0	30.0	25.0	54.0	30.0	23.0
$N-C_5H_{12}$	21.0	11.0	13.0	11.0	28.0	13.0	10.0
<u>Squalane</u>							
$N-C_6H_{14}$	9				23		
$N-C_7H_{16}$	3				7		
Toluene	32				73		
$N-C_8H_{18}$	2				8		
$E-C_6H_5$	9				55		
M-XYL	32				199		
O-XYL	12				47		
$N-C_9H_{20}$	1				8		
$N-P-C_6H_5$	1				3		
$N-C_{10}H_{22}$	3				20		
<u>Beckman 6800</u>							
THC	3810	2350	2300	2200	3110	3790	2390
$CH_4$	1820	1730	1720	1720	1820	1690	1740
CO	1220	811	770	635	1810	850	630

(continued)

TABLE C-1 (continued)

Date	8/17/76 7-9	8/17/76 9-11	8/17/76 11-13	8/19/76 7-9 Repeat Analysis	8/19/76 7-9	8/19/76 11-13
<u>Dura Pak</u>						
C <sub>2</sub> H <sub>6</sub>	14.1	7.9	5.6	19.5	19.2	8.2
C <sub>2</sub> H <sub>4</sub>	158.1	83.1	79.4	108.3	108.1	30.8
C <sub>3</sub> H <sub>8</sub>	21.0	11.7	6.6	32.8	32.9	9.0
C <sub>2</sub> H <sub>2</sub>	21.6	13.5	6.6	19.6	18.5	7.5
IsoC <sub>4</sub> H <sub>10</sub>	10.7	7.2	3.0	9.1	9.0	4.8
N-C <sub>4</sub> H <sub>10</sub>	43.8	30.4	14.2	37.7	37.4	21.2
C <sub>3</sub> H <sub>6</sub>	11.2	8.4	2.3	11.0	11.1	2.3
IsoC <sub>5</sub> H <sub>12</sub>	40.0	29.0	13.0	31.0	35.0	19.0
N-C <sub>5</sub> H <sub>12</sub>	19.0	13.0	6.0	25.0	21.0	10.0
<u>Squalane</u>						
N-C <sub>6</sub> H <sub>14</sub>	18				13	
N-C <sub>7</sub> H <sub>16</sub>	8				8	
Toluene	48				57	
N-C <sub>8</sub> H <sub>18</sub>	5				6	
E-C <sub>6</sub> H <sub>5</sub>	14				25	
M-XYL	39				42	
O-XYL	8				13	
N-C <sub>9</sub> H <sub>20</sub>	5				4	
N-P-C <sub>6</sub> H <sub>5</sub>	2				2	
N-C <sub>10</sub> H <sub>22</sub>	25				19	

(continued)

TABLE C-1 (continued)

Date	8/24/76 7-9	8/24/76 9-11	8/24/76 11-13	8/24/76 13-15	8/26/76 7-9	8/26/76 9-11	8/26/76 11-13
<u>Dura Pak</u>							
$C_2H_6$	30.2	10.0	9.0	7.0	18.6	12.3	10.3
$C_2H_4$	65.5	35.4	103.5	17.3	89.3	117.4	31.1
$C_3H_8$	70.8	13.4	9.6	9.3	23.9	14.4	12.9
$C_2H_2$	23.3	10.9	9.9	8.0	48.1	20.9	14.0
$Isoc_4H_{10}$	88.6	8.3	6.6	5.0	27.2	14.7	10.9
$N-C_4H_{10}$	240.2	35.2	41.3	29.8	144.3	98.9	61.8
$C_3H_6$	34.8	3.5	2.0	0.8	19.8	7.9	3.7
$Isoc_5H_{12}$	163.0	24.0	25.0	19.0	147.0	80.0	49.0
$N-C_5H_{12}$	98.0	14.0	14.0	10.0	61.0	33.0	21.0
<u>Squalane</u>							
$N-C_6H_{14}$					26		
$N-C_7H_{16}$					8		
Toluene					59		
$N-C_8H_{18}$					6		
$E-C_6H_5$					23		
M-XYL					66		
O-XYL					23		
$N-C_9H_{20}$					5		
$N-P-C_6H_5$					4		
$N-C_{10}H_{22}$					14		
<u>Beckman 6800</u>							
THC	4570	2750	2490	2060	3750	3200	2760
$CH_4$	2600	1960	1880	1740	2310	1940	1960
CO	2020	780	860	460	3180	1520	1200

(continued)



Table 1 - continued

Date	8/26/76	3/10/78
	13-15	1-3

Surf Pak

C <sub>2</sub> H <sub>6</sub>	9.9	4.3
C <sub>2</sub> H <sub>4</sub>	112.2	11.4
C <sub>2</sub> H <sub>2</sub>	9.4	1.1
C <sub>2</sub> H <sub>2</sub>	10.3	1.4
Isoc <sub>4</sub> H <sub>10</sub>	11.9	2.6
N <sub>2</sub> C <sub>4</sub> H <sub>10</sub>	29.1	1.1
C <sub>3</sub> H <sub>8</sub>	3.4	1
Isoc <sub>5</sub> H <sub>12</sub>	43.0	29.0
N <sub>2</sub> C <sub>5</sub> H <sub>12</sub>	1.0	1

Brockman 6800

CH <sub>4</sub>	3290	100
CH <sub>4</sub>	1840	1200
CO	1010	100

TABLE C-2. RAMS STATION 102

<u>SUMMER INTENSIVE</u>				
	Mean		Standard Deviation	
THC	2660 ppbC		306 ppbC	
CH <sub>4</sub>	1972 ppbC		97 ppbC	
CO	935 ppbC		214 ppbC	

Date	6/23/76 7-9	6/23/76 9-11	6/23/76 11-13	6/23/76 13-15
<u>Beckman 6800</u>				
THC	3010	2820	2440	2370
CH <sub>4</sub>	1970	2110	1900	1910
CO	1100	1140	750	750

TABLE C-3. RAMS STATION 103

SUMMER INTENSIVE

	Mean	Standard Deviation
THC	2327 ppbC	424 ppbC
CH <sub>4</sub>	1860 ppbC	273 ppbC
CO	763 ppbC	412 ppbC
C <sub>2</sub> H <sub>2</sub>	4 ppbC	2 ppbC

Date	6/28/76 7-9	6/28/76 9-11	6/28/76 11-13	6/28/76 13-15	6/30/76 7-9	6/30/76 9-11	6/30/76 11-13
<u>Beckman 6800</u>							
THC	3670	2530	2280	2350	2210	2070	2140
CH <sub>4</sub>	3200	1920	1900	1850	1900	1880	1820
CO	1400	870	880	750	740	630	660

(continued)

TABLE C-3 (continued)

Date	6/30/76 13-15	7/13/76 11-13	7/13/76 13-15	7/15/76 7-9	7/15/76 9-11	7/20/76 7-9	7/20/76 9-11
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>		5.8	7.5	8.6	5.7	6.9	7.2
C <sub>2</sub> H <sub>4</sub>		26.2	14.3	23.8	1128.0	26.8	224.2
C <sub>3</sub> H <sub>8</sub>		10.9	13.4	7.6	5.9	9.0	8.6
C <sub>2</sub> H <sub>2</sub>		3.3	4.7	5.0	2.1	5.6	3.7
IsoC <sub>4</sub> H <sub>10</sub>		3.4	7.3	4.2	2.0	6.5	3.7
N-C <sub>4</sub> H <sub>10</sub>		12.7	31.8	18.2	8.5	42.8	17.5
C <sub>3</sub> H <sub>6</sub>		BDL*	Trace	BDL*	BDL*	3.7	BDL*
IsoC <sub>5</sub> H <sub>12</sub>		9.0	22.0	14.0	6.5	27.0	12.0
N-C <sub>5</sub> H <sub>12</sub>		1.0	9.0	4.5	1.4	8.0	4.8
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>						NM**	
N-C <sub>7</sub> H <sub>16</sub>						2	
Toluene						49	
N-C <sub>8</sub> H <sub>18</sub>						2	
E-C <sub>6</sub> H <sub>5</sub>						31	
M-XYL						138	
O-XYL						27	
N-C <sub>9</sub> H <sub>20</sub>						2	
N-P-C <sub>6</sub> H <sub>5</sub>						BDL*	
N-C <sub>10</sub> H <sub>22</sub>						36	
<u>Beckman 6800</u>							
THC	2090	2090	2240	2470	3570	2730	2650
CH <sub>4</sub>	1840	1910	1870	2030	1990	1880	1780
CO	370	610	800	1220	840	1170	990

\* Below detection limits

\*\* Not measurable (because of interference etc.)

(continued)

TABLE C-3 (continued)

Date	7/20/76 11-13	7/20/76 13-15	7/22/76 7-9	7/22/76 9-11	7/22/76 13-15	7/27/76 7-9	7/27/76 9-11
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	4.8	3.9	10.7	8.1	5.2	11.8	6.9
C <sub>2</sub> H <sub>4</sub>	38.9	13.6	69.0	32.8	26.5	20.1	19.9
C <sub>3</sub> H <sub>8</sub>	4.9	3.9	34.8	16.5	18.0	37.9	11.5
C <sub>2</sub> H <sub>2</sub>	1.5	1.5	9.1	5.5	2.2	6.6	4.3
IsoC <sub>4</sub> H <sub>10</sub>	3.7	3.1	23.1	4.5	2.6	15.6	6.7
N-C <sub>4</sub> H <sub>10</sub>	15.8	13.6	62.9	18.4	13.1	37.8	19.1
C <sub>3</sub> H <sub>6</sub>	BDL*	BDL*	5.9	2.8	BDL*	49.8	40.4
IsoC <sub>5</sub> H <sub>12</sub>	10.0	8.0	38.0	15.0	5.0	26.0	15.0
N-C <sub>5</sub> H <sub>12</sub>	4.8	3.0	15.0	6.0	4.0	13.0	6.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>			25			11	
N-C <sub>7</sub> H <sub>16</sub>			211			3	
Toluene			586			16	
N-C <sub>8</sub> H <sub>18</sub>			1008			2	
E-C <sub>6</sub> H <sub>5</sub>			1262			14	
M-XYL			1695			39	
O-XYL			3423			10	
N-C <sub>9</sub> H <sub>20</sub>			1426			2	
N-P-C <sub>6</sub> H <sub>5</sub>			6			2	
N-C <sub>10</sub> H <sub>22</sub>			24			BDL*	
<u>Beckman 6800</u>							
THC	2340	2160	1390	2310	1980	2500	2180
CH <sub>4</sub>	1760	1720	2000	1800	1760	1520	1470
CH	380	430	2510	1050	440	550	410

\* Below detection limits

(continued)

TABLE C-3 (continued)

Date			Special Bags				
	7/27/76 11-13	7/27/76 13-15	7/29/76 0800- 0815	7/29/76 1515- 1530	7/29/76 7-9	7/29/76 9-11	7/29/76 11-13
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	7.1	6.5	55.4	55.3	9.7	10.1	12.0
C <sub>2</sub> H <sub>4</sub>	15.2	13.7	68.6	61.4	24.7	27.6	48.8
C <sub>3</sub> H <sub>8</sub>	13.0	8.9	19.2	18.9	19.3	18.0	19.8
C <sub>2</sub> H <sub>2</sub>	4.0	3.0	+ 7.9	+ 7.0	7.6	7.0	8.5
IsoC <sub>4</sub> H <sub>10</sub>	2.8	2.3	11.2	9.5	5.6	5.2	8.0
N-C <sub>4</sub> H <sub>10</sub>	9.8	7.6	117.7	98.9	19.2	18.1	30.9
C <sub>3</sub> H <sub>6</sub>	35.6	20.4	8.9	1.8	12.2	12.7	10.0
IsoC <sub>5</sub> H <sub>12</sub>	7.8	4.0	88.0	64.0	19.0	18.0	26.0
N-C <sub>5</sub> H <sub>12</sub>	3.0	1.0	15.0	14.0	7.3	7.7	11.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>					5		
N-C <sub>7</sub> H <sub>16</sub>					1		
Toluene					27		
N-C <sub>8</sub> H <sub>18</sub>					2		
E-C <sub>6</sub> H <sub>5</sub>					10		
M-XYL					29		
O-XYL					8		
N-C <sub>9</sub> H <sub>20</sub>					2		
N-P-C <sub>6</sub> H <sub>5</sub>					2		
N-C <sub>10</sub> H <sub>22</sub>					2		
<u>Beckman 6800</u>							
THC	2690	2090	+ 2650	2790	2190	2220	2420
CH <sub>4</sub>	1740	1680	1970	1760	1880	1920	1710
CO	410	390	2070	2230	810	840	811

+Values are not included in calculations

(continued)

TABLE C-3 (continued)

Date	8/3/76 7-9	8/3/76 9-11	8/3/76 11-13	8/3/76 13-15	8/5/76 7-9	8/5/76 9-11	8/5/76 11-13
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	11.2	4.7	4.0	5.9	6.8	8.4	6.0
C <sub>2</sub> H <sub>4</sub>	37.3	18.3	8.4	19.5	8.4	29.4	17.9
C <sub>3</sub> H <sub>8</sub>	24.3	7.2	4.9	7.5	31.0	17.6	12.4
C <sub>2</sub> H <sub>2</sub>	4.1	2.9	2.2	2.7	3.3	5.8	3.4
IsoC <sub>4</sub> H <sub>10</sub>	3.5	2.5	2.3	4.8	15.8	8.0	6.6
N-C <sub>4</sub> H <sub>10</sub>	11.5	9.5	7.2	4.8	47.6	59.2	22.4
C <sub>3</sub> H <sub>6</sub>	5.1	11.8	2.2	4.0	2.0	2.0	1.0
IsoC <sub>5</sub> H <sub>12</sub>	10.0	8.0	3.0	6.0	40.0	32.0	17.0
N-C <sub>5</sub> H <sub>12</sub>	4.0	3.0	NM**	3.0	9.0	10.0	8.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>	3				38		
N-C <sub>7</sub> H <sub>16</sub>	1				1		
Toluene	8				16		
N-C <sub>8</sub> H <sub>18</sub>	BDL*				3		
E-C <sub>6</sub> H <sub>5</sub>	4				7		
M-XYL	13				29		
O-XYL	2				5		
N-C <sub>9</sub> H <sub>20</sub>	Trace				1		
N-P-C <sub>6</sub> H <sub>5</sub>	Trace				1		
N-C <sub>10</sub> H <sub>22</sub>	1				3		
<u>Beckman 6800</u>							
THC	2270	1940	1950	2100	2360	2360	2360
CH <sub>4</sub>	1920	1750	1830	1710	1800	1770	1720
CO	570	530	450	560	670	820	660

\* Below detection limits

\*\* Not measurable (because of interference etc.)

(continued)

TABLE C-3 (continued)

Date	8/5/76 13-15
<u>Dura Pak</u>	
$C_2H_6$	5.3
$C_2H_4$	19.2
$C_3H_8$	8.5
$C_2H_2$	2.5
$IsoC_4H_{10}$	4.9
$N-C_4H_{10}$	26.2
$C_3H_6$	1.0
$IsoC_5H_{12}$	18.0
$N-C_5H_{12}$	6.0



TABLE C-4. RAMS STATION 108

<u>SUMMER INTENSIVE</u>					
	Mean			Standard Deviation	
THC	2223 ppbC			270 ppbC	
CH <sub>4</sub>	1872 ppbC			98 ppbC	
CO	603 ppbC			388 ppbC	

Date	6/23/76 7-9	6/23/76 11-13	6/23/76 13-15	6/28/76 9-11	6/28/76 11-13
<u>Beckman 6800</u>					
THC	2870	2260	2210	2260	2210
CH <sub>4</sub>	2130	1840	1840	1840	1840
CO	1480	680	520	680	560

Date	6/30/76 7-9	6/30/76 9-11	6/30/76 11-13	6/30/76 13-15
<u>Beckman 6800</u>				
THC	2190	2100	1980	1930
CH <sub>4</sub>	1880	1820	1840	1820
CO	410	740	180	180

TABLE C-5. RAMS STATION 113

<u>SUMMER INTENSIVE</u>			
	Mean		Standard Deviation
THC	1874 ppbC		78 ppbC
CH <sub>4</sub>	1912 ppbC		82 ppbC
CO	957 ppbC		167 ppbC

---

Date	7/6/76 7-9	7/6/76 9-11	7/6/76 13-15
<u>Beckman 6800</u>			
THC	1963	1816	1843
CH <sub>4</sub>	1995	1912	1830
CO	1150	861	861

TABLE C-6. RAMS STATION 114

<u>SUMMER INTENSIVE</u>							
	Mean				Standard Deviation		
THC	2613 ppbC				912 ppbC		
CH <sub>4</sub>	1851 ppbC				310 ppbC		
CO	599 ppbC				302 ppbC		
C <sub>2</sub> H <sub>2</sub>	7.1 ppbC				8.9 ppbC		
Date	6/23/76 9-11	6/23/76 11-13	6/23/76 13-15	7/15/76 9-11	7/15/76 11-13	7/15/76 13-15	7/20/76 9-11
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>				4.8	4.6	2.4	5.7
C <sub>2</sub> H <sub>4</sub>				16.4	21.8	10.9	20.5
C <sub>3</sub> H <sub>8</sub>				4.8	4.8	2.8	6.3
C <sub>2</sub> H <sub>2</sub>				1.7	1.4	0.7	3.0†
IsoC <sub>4</sub> H <sub>10</sub>				1.0	1.0	1.0	3.6
N-C <sub>4</sub> H <sub>10</sub>				6.1	6.6	4.4	15.4
C <sub>3</sub> H <sub>6</sub>				BDL*	BDL*	0.5	BDL*
IsoC <sub>5</sub> H <sub>12</sub>				5.0	4.0	2.0	10.5
N-C <sub>5</sub> H <sub>12</sub>				1.0	1.0	1.0	2.3
<u>Beckman 6800</u>							
THC	2200	2750	2160	2450	2330	2350	2490
CH <sub>4</sub>	1820	1840	1840	1960	1920	1650	1760
CO	520	560	540	550	400	330	460

\* Below detection limits

(continued)

TABLE C-6 (continued)

Date	7/20/76 11-13	7/20/76 13-15	7/22/76 9-11	7/22/76 11-13	7/22/76 13-15	7/27/76 9-11	7/27/76 11-13
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	5.0	3.9	8.2	7.5	5.5	7.5	6.6
C <sub>2</sub> H <sub>4</sub>	32.6	9.7	15.5	41.9	10.8	19.2	27.8
C <sub>3</sub> H <sub>8</sub>	4.9	3.7	8.9	7.6	4.9	14.7	10.1
C <sub>2</sub> H <sub>2</sub>	2.0	2.0	4.7	2.9	1.0	3.7	2.6
IsoC <sub>4</sub> H <sub>10</sub>	3.0	1.8	4.1	2.6	2.1	8.6	3.3
N-C <sub>4</sub> H <sub>10</sub>	12.8	9.8	14.2	12.4	11.2	19.0	9.9
C <sub>3</sub> H <sub>6</sub>	BDL*	BDL*	0.7	BDL*	BDL*	31.0	51.7
IsoC <sub>5</sub> H <sub>12</sub>	10.0	6.8	14.0	10.0	9.0	11.0	4.0
N-C <sub>5</sub> H <sub>12</sub>	3.0	3.0	5.0	4.0	3.0	6.0	6.0
<u>Beckman 6800</u>							
THC	2650	2240	2080	2040	1950	2090	2470
CH <sub>4</sub>	1740	1610	1820	1800	1760	1390	1660
CO	410	480	660	410	480	340	340
Date	7/27/76 13-15	7/29/77 11-13	8/3/76 11-13	8/3/76 13-15	8/5/76 9-11	8/5/76 11-13	8/5/76 13-15
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	6.8	14.9	7.7	5.5	13.3	5.7	4.6
C <sub>2</sub> H <sub>4</sub>	39.9	29.6	113.3	17.0	52.9	102.5	10.5
C <sub>3</sub> H <sub>8</sub>	8.8	25.8	18.6	11.2	12.1	5.5	3.5
C <sub>2</sub> H <sub>2</sub>	2.0	5.6	2.8	2.4	11.5	5.3	2.7
IsoC <sub>4</sub> H <sub>10</sub>	2.5	14.3	16.2	13.1	7.4	3.8	1.8
N-C <sub>4</sub> H <sub>10</sub>	8.4	35.7	48.1	35.2	34.9	17.3	8.5
C <sub>3</sub> H <sub>6</sub>	35.0	13.0	24.5	10.3	4.5	6.5	BDL*
IsoC <sub>5</sub> H <sub>12</sub>	2.0	25.0	27.0	35.0	31.0	13.0	6.0
N-C <sub>5</sub> H <sub>12</sub>	3.0	11.0	14.0	11.0	15.0	7.0	3.0
<u>Beckman 6800</u>							
THC		2300	2450	3870	2110	2740	2120
CH <sub>4</sub>		1680	1790	2230	1710	1720	1670
CO		510	340	650	840	840	730

\* Below detection limits

(continued)

TABLE C-6 (continued)

Date	8/10/76 9-11	8/10/76 11-13	8/10/76 13-15	8/12/76 7-9	8/12/76 9-11	8/12/76 13-15	8/17/76 7-9
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	5.5	51.9	4.2	37.6	6.7	5.4	6.2
C <sub>2</sub> H <sub>4</sub>	19.6	26.6	10.2	20.5	20.8	11.5	26.1
C <sub>3</sub> H <sub>8</sub>	6.1	26.9	7.5	19.8	7.4	5.7	15.1
C <sub>2</sub> H <sub>2</sub>	5.7	49.4	3.8	18.1	6.7	21.7	5.3
IsoC <sub>4</sub> H <sub>10</sub>	4.0	21.6	4.6	9.1	3.0	1.0	7.1
N-C <sub>4</sub> H <sub>10</sub>	18.7	56.3	19.8	29.0	12.5	4.7	27.4
C <sub>3</sub> H <sub>6</sub>	0.9	13.0	BDL*	6.0	1.5	BDL*	2.2
IsoC <sub>5</sub> H <sub>12</sub>	15.0	59.0	14.0	26.0	11.0	3.0	13.0
N-C <sub>5</sub> H <sub>12</sub>	6.0	53.0	6.0	39.0	5.0	2.0	9.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>				9			9
N-C <sub>7</sub> H <sub>16</sub>				14			3
Toluene				45			14
N-C <sub>8</sub> H <sub>18</sub>				3			1
E-C <sub>6</sub> H <sub>5</sub>				30			5
M-XYL				89			19
O-XYL				24			5
N-C <sub>9</sub> H <sub>20</sub>				3			1
N-P-C <sub>6</sub> H <sub>5</sub>				1			NM**
N-C <sub>10</sub> H <sub>22</sub>				8.			1
<u>Beckman 6800</u>							
THC				2370	2110	2090	
CH <sub>4</sub>				1880	1690	1670	
CO				590	580	290	

\* Below detection limits

\*\* Not measurable (because of interference etc.)

(continued)

TABLE C-6 (continued)

Date	8/17/76 9-11	8/17/76 11-13	8/18/76 13-15	8/19/76 7-9	8/19/76 9-11	8/19/76 11-13	8/19/76 13-15
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	6.5	5.2	4.8	15.2	13.1	6.1	5.8
C <sub>2</sub> H <sub>4</sub>	30.8	14.9	13.2	19.6	42.7	14.3	15.0
C <sub>3</sub> H <sub>8</sub>	13.0	8.4	6.6	55.5	53.5	6.1	8.8
C <sub>2</sub> H <sub>2</sub>	4.2	5.0	6.3	4.9	5.8	3.7	2.2
IsoC <sub>4</sub> H <sub>10</sub>	4.7	4.3	2.5	44.3	11.1	2.3	2.1
N-C <sub>4</sub> H <sub>10</sub>	15.8	20.1	11.3	131.1	46.5	8.1	8.0
C <sub>3</sub> H <sub>6</sub>	2.1	1.3	1.0	BDL*	BDL*	BDL*	BDL*
IsoC <sub>5</sub> H <sub>12</sub>	11.0	15.0	10.0	65.0	26.0	5.0	5.0
N-C <sub>5</sub> H <sub>12</sub>	5.0	6.0	3.0	42.0	11.0	3.0	3.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>				24			
N-C <sub>7</sub> H <sub>16</sub>				17			
Toluene				21			
N-C <sub>8</sub> H <sub>18</sub>				9			
E-C <sub>6</sub> H <sub>5</sub>				12			
M-XYL				30			
O-XYL				10			
N-C <sub>9</sub> H <sub>20</sub>				2			
N-P-C <sub>6</sub> H <sub>5</sub>				1			
N-C <sub>10</sub> H <sub>22</sub>				5			

\* Below detection limits

(continued)

TABLE C-6 (continued)

Date	8/26/76 7-9	8/26/76 9-11	8/31/76 7-9	8/31/76 9-11	9/2/76 7-9	9/2/76 9-11	9/2/76 13-15
<u>Dura Pak</u>							
$C_2H_6$	15.8	14.2	57.3	13.7	15.4	7.6	4.6
$C_2H_4$	35.4	37.0	24.9	55.0	18.2	25.3	31.6
$C_3H_8$	19.5	15.9	40.1	35.6	51.1	24.4	5.1
$C_2H_2$	10.1	15.3	21.5	16.3	3.2	2.0	2.3
$IsoC_4H_{10}$	8.3	7.8	38.0	23.8	73.9	20.1	2.0
$N-C_4H_{10}$	30.0	37.0	173.3	61.1	210.2	77.6	7.5
$C_3H_6$	3.9	4.7	10.7	6.1	19.7	73.9	0.7
$IsoC_5H_{12}$	25.0	32.0	193.0	52.0	95.0	45.0	8.0
$N-C_5H_{12}$	11.0	14.0	98.0	25.0	69.0	19.0	2.0
<u>Squalane</u>							
$N-C_6H_{14}$	11		15		36		
$N-C_7H_{16}$	2		5		15		
Toluene	23		56		21		
$N-C_8H_{18}$	2		4		2		
$E-C_6H_5$	11		52		17		
M-XYL	32		131		55		
O-XYL	9		37		14		
$N-C_9H_{20}$	1		2		2		
$N-P-C_6H_5$	1		11		2		
$N-C_{10}H_{22}$	8		7		1		
<u>Beckman 6800</u>							
THC	2850	2910	6140	3000	2800	2020	1780
$CH_4$	2290	2200	3100	2220	1690	1720	1690
CO	830	1200	1330	1540	450	350	410

TABLE C-7. RAMS STATION 115

SUMMER INTENSIVE							
	Mean				Standard Deviation		
THC	2427 ppbC				908 ppbC		
CH <sub>4</sub>	1788 ppbC				147 ppbC		
CO	498 ppbC				213 ppbC		
C <sub>2</sub> H <sub>2</sub>	3 ppbC				2 ppbC		
Date	6/28/76 9-11	6/28/76 11-13	6/28/76 13-15	6/30/76 9-11	6/30/76 11-13	6/30/76 13-15	
<u>Beckman 6800</u>							
THC	2660	2490	2550	2310	2330	2120	
CH <sub>4</sub>	1880	1880	1900	1840	1800	1840	
CO	550	650	430	630	260	260	
Date	7/13/76 13-15	7/15/76 9-11	7/15/76 11-13	7/15/76 13-15	7/20/76 9-11	7/20/76 11-13	7/20/76 13-15
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	7.6	4.7	3.4	2.5	5.7	4.8	4.2
C <sub>2</sub> H <sub>4</sub>	20.6	33.9	28.5	36.2	34.0	52.8	31.2
C <sub>3</sub> H <sub>8</sub>	14.8	4.6	3.1	2.1	6.6	5.0	4.3
C <sub>2</sub> H <sub>2</sub>	3.0	1.7	2.2	0.9	2.5	2.3	2.4
IsoC <sub>4</sub> H <sub>10</sub>	4.9	1.5	1.4	0.9	3.1	2.0	2.3
N-C <sub>4</sub> H <sub>10</sub>	10.9	5.6	5.3	4.2	12.9	8.4	9.6
C <sub>3</sub> H <sub>6</sub>	2.0	BDL*	BDL*	BDL*	BDL*	BDL*	BDL*
IsoC <sub>5</sub> H <sub>12</sub>	8.0	3.0	2.0	3.0	5.0	7.0	8.0
N-C <sub>5</sub> H <sub>12</sub>	2.9	1.0	BDL*	BDL*	3.0	2.0	2.0
<u>Beckman 6800</u>							
THC	2160	2520	2260	2140	2220	2170	2090
CH <sub>4</sub>	1870	1930	1870	1830	1760	1720	1720
CO	860	490	330	610	690	410	410

\* Below detection limits

(continued)



TABLE C-7 (continued)

Date	7/22/76 9-11	7/22/76 13-15	7/27/76 9-11	7/27/76 11-13	7/27/76 13-15	7/29/76 9-11	7/29/76 11-13
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	9.9	6.0	6.6	6.8	5.7	13.9	11.0
C <sub>2</sub> H <sub>4</sub>	38.8	24.2	33.2	39.9	17.6	19.8	25.4
C <sub>3</sub> H <sub>8</sub>	14.6	5.2	9.0	8.8	5.1	22.8	16.2
C <sub>2</sub> H <sub>2</sub>	6.3	1.3	3.1	2.0	1.5	6.9	3.9
IsoC <sub>4</sub> H <sub>10</sub>	7.0	2.1	4.8	2.5	1.2	9.4	7.8
N-C <sub>4</sub> H <sub>10</sub>	24.1	10.6	12.1	8.4	4.8	27.9	22.5
C <sub>3</sub> H <sub>6</sub>	BDL*	BDL*	44.9	35.0	19.3	14.2	11.0
IsoC <sub>5</sub> H <sub>12</sub>	19.0	9.0	6.0	2.0	2.0	26.0	15.0
N-C <sub>5</sub> H <sub>12</sub>	9.0	2.0	2.0	4.0	BDL*	10.0	7.0
<u>Beckman 6800</u>							
THC	2210	2010	3830	2460	2080	2030	2860
CH <sub>4</sub>	1920	1770	1350	1610	1580	2060	1650
CO	160	380	280	330	330	950	450

Date	7/29/76 13-15	8/3/76 9-11	8/3/76 11-13	8/3/76 13-15	8/5/76 9-11
<u>Dura Pak</u>					
C <sub>2</sub> H <sub>6</sub>	11.3	3.2	8.0	2.9	7.4
C <sub>2</sub> H <sub>4</sub>	12.9	21.6	36.0	17.0	27.8
C <sub>3</sub> H <sub>8</sub>	16.9	3.1	5.2	1.9	10.1
C <sub>2</sub> H <sub>2</sub>	4.5	2.6	10.4	1.9	6.8
IsoC <sub>4</sub> H <sub>10</sub>	8.0	2.1	6.0	2.0	4.0
N-C <sub>4</sub> H <sub>10</sub>	23.7	10.9	37.1	7.3	22.6
C <sub>3</sub> H <sub>6</sub>	8.1	14.4	2890.9	6.7	19.2
IsoC <sub>5</sub> H <sub>12</sub>	14.0	10.0	28.0	3.0	18.0
N-C <sub>5</sub> H <sub>12</sub>	7.0	3.0	6.0	2.0	9.0
<u>Beckman 6800</u>					
THC	1060	1980	6200	2000	2230
CH <sub>4</sub>	1670	1690	1980	1800	1790
CO	440	700	350	560	940

\* Below detection limits

TABLE C-8. RAMS STATION 118

SUMMER INTENSIVE							
	Mean				Standard Deviation		
THC	2791 ppbC				1311 ppbC		
CH <sub>4</sub>	1770 ppbC				133 ppbC		
CO	479 ppbC				301 ppbC		
C <sub>2</sub> H <sub>2</sub>	3.4 ppbC				2.3 ppbC		
Date	7/13/76 13-15	7/15/76 7-9	7/15/76 13-15	7/20/76 7-9	7/22/76 7-9	7/22/76 13-15	7/27/76 7-9
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	6.2	5.0	2.1	4.8	6.6	4.6	6.3
C <sub>2</sub> H <sub>4</sub>	41.4	15.0	7.9	24.8	9.3	13.5	18.2
C <sub>3</sub> H <sub>8</sub>	27.6	6.8	2.0	4.6	8.3	3.9	7.7
C <sub>2</sub> H <sub>2</sub>	8.5	2.7	0.6	1.5	1.1	0.2	5.6
IsoC <sub>4</sub> H <sub>10</sub>	2.4	1.2	0.5	1.1	2.4	1.2	3.3
N-C <sub>4</sub> H <sub>10</sub>	6.6	7.9	2.4	6.4	9.7	3.9	14.6
C <sub>3</sub> H <sub>6</sub>	BDL	BDL	BDL	BDL	BDL	0.7	12.7
IsoC <sub>5</sub> H <sub>12</sub>	3.0	5.0	1.0	2.0	4.0	2.0	8.0
N-C <sub>5</sub> H <sub>12</sub>	1.0	2.0			14.0	1.0	2.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>				NM			2
N-C <sub>7</sub> H <sub>16</sub>				8			1
Toluene				26			4
N-C <sub>8</sub> H <sub>18</sub>				23			1
E-C <sub>6</sub> H <sub>5</sub>				19			5
M-XYL				84			18
O-XYL				21			6
N-C <sub>9</sub> H <sub>20</sub>				37			Trace
N-P-C <sub>6</sub> H <sub>5</sub>				1			Trace
N-C <sub>10</sub> H <sub>22</sub>				126			1
<u>Beckman 6800</u>							
THC	3850	2330	2280	2340		2670	2080
CH <sub>4</sub>	1870	1900	1870	1820		1740	1480
CO	1380	580	210	356		191	380

TABLE C-8 (continued)

Date	7/27/76 13-15	7/29/76 7-9	7/29/76 13-15	8/03/76 7-9	8/03/76 13-15	8/05/76 7-9	8/05/76 13-15
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	7.8	6.4	8.4	6.2	3.8	5.3	4.2
C <sub>2</sub> H <sub>4</sub>	4.3	25.2	22.7	32.5	11.2	27.2	12.6
C <sub>3</sub> H <sub>8</sub>	5.5	7.6	10.2	22.7	7.0	4.7	3.0
C <sub>2</sub> H <sub>2</sub>	4.1	4.7	5.0	5.2	2.8	1.9	1.3
IsoC <sub>4</sub> H <sub>10</sub>	1.8	2.8	3.1	6.5	3.5	1.8	0.8
N-C <sub>4</sub> H <sub>10</sub>	7.8	15.3	15.8	20.0	13.5	5.6	3.2
C <sub>3</sub> H <sub>6</sub>	10.0	24.4	19.4	92.0	BDL*	BDL*	BDL*
IsoC <sub>5</sub> H <sub>12</sub>	6.0	11.0	11.0	12.0	9.0	3.0	2.0
N-C <sub>5</sub> H <sub>12</sub>	1.0	2.0	3.0	9.0	4.0	5.2	1.2
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>		2		5		1	
N-C <sub>7</sub> H <sub>16</sub>		1		2		Trace	
Toluene		13		6		3	
N-C <sub>8</sub> H <sub>18</sub>		2		2		1	
E-C <sub>6</sub> H <sub>5</sub>		10		3		3	
M-XYL		43		11		17	
O-XYL		9		4		2	
N-C <sub>9</sub> H <sub>20</sub>		1		1		Trace	
N-P-C <sub>6</sub> H <sub>5</sub>		1		Trace		BDL*	
N-C <sub>10</sub> H <sub>22</sub>		21		2		1	
<u>Beckman 6800</u>							
THC	6070	2030	2610	2280	2130	2060	2100
CH <sub>4</sub>	1990	1780	1610	1810	1750	1720	1670
CO	460	490	430	530	590	410	230

\* Below detection limits

TABLE C-9. RAMS STATION 121

<u>SUMMER INTENSIVE</u>					
	Mean			Standard Deviation	
THC	1960 ppbC			311 ppbC	
CH <sub>4</sub>	1872 ppbC			43 ppbC	
CO	364 ppbC			183 ppbC	
Date	7/6/76 9-11	7/6/76 11-13	7/6/76 13-15	7/8/76 9-11	7/8/76 13-15
<u>Beckman 6800</u>					
THC	1790	1880	2500	1910	1720
CH <sub>4</sub>	1870	1930	1830	1900	1830
CO	660	370	717	350	270

TABLE C-10. RAMS STATION 122

SUMMER INTENSIVE						
	Mean			Standard Deviation		
THC	2939 ppbC			1771 ppbC		
CH <sub>4</sub>	1864 ppbC			212 ppbC		
CO	607 ppbC			240 ppbC		
C <sub>2</sub> H <sub>2</sub>	5 ppbC			3 ppbC		
Date	8/10/76 13-15	8/12/76 7-9	8/12/76 13-15	8/17/76 7-9	8/17/76 13-15	8/19/76 13-15
<u>Dura Pak</u>						
C <sub>2</sub> H <sub>6</sub>	4.5	12.3	7.4	4.2	2.9	4.9
C <sub>2</sub> H <sub>4</sub>	6.4	15.4	16.1	6.6	5.3	6.5
C <sub>3</sub> H <sub>8</sub>	6.0	26.1	6.8	8.5	3.4	4.3
C <sub>2</sub> H <sub>2</sub>	5.0	8.4	4.4	4.1	2.0	2.4
IsoC <sub>4</sub> H <sub>10</sub>	3.3	20.4	2.3	2.3	0.9	1.3
N-C <sub>4</sub> H <sub>10</sub>	15.8	47.2	7.8	9.1	3.4	4.2
C <sub>3</sub> H <sub>6</sub>	BDL*	4.9	1.0	1.1	BDL*	BDL*
IsoC <sub>5</sub> H <sub>12</sub>	12.0	36.0	7.0	6.0	1.9	2.0
N-C <sub>5</sub> H <sub>12</sub>	4.0	18.0	3.0	2.0	0.8	2.0
<u>Squalane</u>						
N-C <sub>6</sub> H <sub>14</sub>				2		
N-C <sub>7</sub> H <sub>16</sub>				3		
Toluene				12		
N-C <sub>8</sub> H <sub>18</sub>				Trace		
E-C <sub>6</sub> H <sub>5</sub>				8		
M-XYL				28		
O-XYL				5		
N-C <sub>9</sub> H <sub>20</sub>				Trace		
N-P-C <sub>6</sub> H <sub>5</sub>				1		
N-C <sub>10</sub> H <sub>22</sub>				2		
<u>Beckman 6800</u>						
THC	2070	2580	1950			
CH <sub>4</sub>	1810	1840	1670			
CO	450	650	300			

\* Below detection limits

(continued)

TABLE C-10 (continued)

Date	8/24/76 7-9	8/24/76 13-15	8/26/76 7-9	8/26/76 13-15	8/31/76 7-9	8/31/76 13-15	9/2/76 7-9
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	7.5	5.8	9.4	7.7	18.1	8.4	0.7
C <sub>2</sub> H <sub>4</sub>	18.9	8.1	43.3	25.6	16.8	5.7	3.1
C <sub>3</sub> H <sub>8</sub>	10.1	4.6	10.0	7.4	29.1	28.2	4.4
C <sub>2</sub> H <sub>2</sub>	3.9	3.0	10.0	3.9	10.6	5.3	1.3
IsoC <sub>4</sub> H <sub>10</sub>	2.0	1.7	7.1	2.1	12.4	23.4	1.0
N-C <sub>4</sub> H <sub>10</sub>	7.8	4.1	31.2	6.7	43.1	76.3	4.2
C <sub>3</sub> H <sub>6</sub>	0.7	BDL*	1.3	0.4	5.3	1.1	0.1
IsoC <sub>5</sub> H <sub>12</sub>	5.0	3.0	22.0	5.0	37.0	46.0	2.0
N-C <sub>5</sub> H <sub>12</sub>	3.0	2.0	13.0	2.0	17.0	27.0	7.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>			6		10		5
N-C <sub>7</sub> H <sub>16</sub>			2		2		5
Toluene			18		24		4
N-C <sub>8</sub> H <sub>18</sub>			1		2		1
E-C <sub>6</sub> H <sub>5</sub>			16		12		5
M-XYL			48		31		16
O-XYL			14		10		4
N-C <sub>9</sub> H <sub>20</sub>			1		1		1
N-P-C <sub>6</sub> H <sub>5</sub>			2		1		1
N-C <sub>10</sub> H <sub>22</sub>			8		5		2
<u>Beckman 6800</u>							
THC	2270	2080	2520	2210	2750	2160	1600
CH <sub>4</sub>	1890	1690	1900	1800	2390	1800	1650
CO	460	400	760	490	970	620	310

\* Below detection limits

(continued)

TABLE C-10 (continued)

Date	9/2/76 13-15
------	-----------------

Dura Pak

$C_2H_6$	4.7
$C_2H_4$	5.1
$C_3H_8$	5.2
$C_2H_2$	1.9
$IsoC_4H_{10}$	1.9
$N-C_4H_{10}$	5.8
$C_3H_6$	
$IsoC_5H_{12}$	3.0
$N-C_5H_{12}$	2.0

TABLE C-11. RAMS STATION 124

<u>SUMMER INTENSIVE</u>							
	Mean				Standard Deviation		
THC	2264 ppbC				382 ppbC		
CH <sub>4</sub>	1931 ppbC				402 ppbC		
CO	579 ppbC				214 ppbC		
C <sub>2</sub> H <sub>2</sub>	3 ppbC				2 ppbC		
Date	8/12/76 7-9	8/12/76 13-15	8/17/76 11-13	8/24/76 7-9	8/26/76 7-9	8/26/76 13-15	8/31/76 7-9
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	4.8	7.6	8.4	10.4	6.0	7.2	1.8
C <sub>2</sub> H <sub>4</sub>	255.8	40.0	112.9	54.1	151.7	113.5	38.5
C <sub>3</sub> H <sub>8</sub>	5.5	6.9	15.6	14.8	5.9	5.9	5.7
C <sub>2</sub> H <sub>2</sub>	2.5	2.6	2.6	7.1	2.6	1.7	3.5
IsoC <sub>4</sub> H <sub>10</sub>		1.8	3.1	5.2	1.1	1.7	1.5
N-C <sub>4</sub> H <sub>10</sub>	4.4	4.3	11.7	30.8	4.5	6.2	8.4
C <sub>3</sub> H <sub>6</sub>				1.9	0.9	0.5	1.6
IsoC <sub>5</sub> H <sub>12</sub>	3.0	3.0	4.0	14.0	3.0	5.0	5.0
N-C <sub>5</sub> H <sub>12</sub>	2.0	2.0	3.0	6.0	3.0	4.0	3.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>							
N-C <sub>7</sub> H <sub>16</sub>							
Toluene							
N-C <sub>8</sub> H <sub>18</sub>							
E-C <sub>6</sub> H <sub>5</sub>							
M-XYL							
O-XYL							
N-C <sub>9</sub> H <sub>20</sub>							
N-P-C <sub>6</sub> H <sub>5</sub>							
N-C <sub>10</sub> H <sub>22</sub>							
<u>Beckman 6800</u>							
THC	2380	1950		2270	2630	2210	
CH <sub>4</sub>	1720	1670		1890	1900	1930	
CO	290	300		460	460	300	

(continued)



TABLE C-11 (continued)

Date	8/31/76 13-15	9/2/76 7-9	9/2/76 13-15
<u>Dura Pak</u>			
$C_2H_6$	6.3	6.9	5.4
$C_2H_4$	278.6	111.0	180.8
$C_3H_8$	4.3	13.1	7.7
$C_2H_2$	2.9	2.4	1.6
$IsoC_4H_{10}$	2.0	3.2	2.4
$N-C_4H_{10}$	6.9	10.5	7.1
$C_3H_6$	0.7	0.9	0.5
$IsoC_5H_{12}$	5.0	5.0	5.0
$N-C_5H_{12}$	3.0	3.0	3.0
<u>Squalane</u>			
$N-C_6H_{14}$			
$N-C_7H_{16}$			
Toluene			
$N-C_8H_{18}$			
$E-C_6H_5$			
M-XYL			
O-XYL			
$N-C_9H_{20}$			
$N-P-C_6H_5$			
$N-C_{10}H_{22}$			
<u>Beckman 6800</u>			
THC	2900	1910	1850
$CH_4$	2140	1750	1690
CO	620	280	220

TABLE C-12. RAMS STATION 101

	FALL INTENSIVE				Standard Deviation		
	Mean						
THC	2906 ppbC				789 ppbC		
CH <sub>4</sub>	1734 ppbC				240 ppbC		
CO	1815 ppbC				1322 ppbC		
C <sub>2</sub> H <sub>2</sub>	33 ppbC				26 ppbC		
Date	11/8/76 13-15	11/10/76 7-9	11/10/76 9-11	11/10/76 11-13	11/10/76 13-15	11/12/76 9-11	11/12/76 13:30-15
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	18.0	11.9	10.4	7.9	6.4	9.2	7.6
C <sub>2</sub> H <sub>4</sub>	46.4	24.3	20.4	10.7	7.7	21.1	16.5
C <sub>3</sub> H <sub>8</sub>	11.4	6.8	8.8	5.2	4.9	9.0	6.6
C <sub>2</sub> H <sub>2</sub>	40.3	25.8	13.6	5.3	5.7	18.2	10.2
IsoC <sub>4</sub> H <sub>10</sub>	11.8	6.7	5.0	3.1	3.3	4.9	3.1
N-C <sub>4</sub> H <sub>10</sub>	47.5	29.6	22.8	15.4	16.0	20.9	12.8
C <sub>3</sub> H <sub>6</sub>	12.4	7.0	6.0	3.4	2.1	4.1	1.5
IsoC <sub>5</sub> H <sub>12</sub>	44.0	28.0	18.0	9.0	10.0	14.0	9.0
N-C <sub>5</sub> H <sub>12</sub>	22.0	13.0	9.0	6.0	5.0	6.0	5.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>		NM				7	
N-C <sub>7</sub> H <sub>16</sub>		3				4	
Toluene		38				42	
N-C <sub>8</sub> H <sub>18</sub>		14				3	
E-C <sub>6</sub> H <sub>5</sub>		11				15	
M-XYL		27				37	
O-XYL		9				10	
N-C <sub>9</sub> H <sub>20</sub>		43				2	
N-P-C <sub>6</sub> H <sub>5</sub>		2				1	
N-C <sub>10</sub> H <sub>22</sub>		370				4	
<u>Beckman 6800</u>							
THC		2690				2330	1990
CH <sub>4</sub>		2040				1580	1470
CO		2115				740	250

(continued)

TABLE C-12 (continued)

Date	11/16/76 7-9	11/16/76 9-11	11/16/76 11-13	11/16/76 13-15	11/18/76 7-9	11/18/76 9-11	11/18/76 11-13
<u>Dura Pak</u>							
$C_2H_6$	40.0	33.5	30.0	10.7	75.6	66.4	23.8
$C_2H_4$	87.9	185.5	120.3	19.7	57.4	40.5	28.7
$C_3H_8$	37.0	43.5	31.3	14.8	75.5	73.4	27.0
$C_2H_2$	71.2	62.8	87.2	14.2	60.1	42.0	16.7
$IsoC_4H_{10}$	23.0	33.4	47.6	8.4	32.0	27.8	21.0
$N-C_4H_{10}$	101.3	160.8	187.2	30.8	134.2	103.2	95.0
$C_3H_6$	26.0	24.5	57.5	5.8	22.2	15.1	8.8
$IsoC_5H_{12}$	92.0	108.0	161.0	23.0	99.0	71.0	61.0
$N-C_5H_{12}$	38.0	46.0	76.0	11.0	54.0	37.0	28.0
<u>Squalane</u>							
$N-C_6H_{14}$	29				31		
$N-C_7H_{16}$	13				5		
Toluene	87				49		
$N-C_8H_{18}$	10				15		
$E-C_6H_5$	31				16		
M-XYL	68				40		
O-XYL	28				17		
$N-C_9H_{20}$	3				35		
$N-P-C_6H_5$	4				3		
$N-C_{10}H_{22}$	15				10		
<u>Beckman 6800</u>							
THC	3900	4020		2310	3860	3300	2540
$CH_4$	2120	1980		1490	1970	1830	1470
CO	3660	3810		710	3100	2250	900

(continued)

TABLE C-12 (continued)

Date	11/18/76 13-15
------	-------------------

Dura Pak

C <sub>2</sub> H <sub>6</sub>	46.2
C <sub>2</sub> H <sub>4</sub>	20.2
C <sub>3</sub> H <sub>8</sub>	53.3
C <sub>2</sub> H <sub>2</sub>	17.2
IsoC <sub>4</sub> H <sub>10</sub>	22.5
N-C <sub>4</sub> H <sub>10</sub>	99.4
C <sub>3</sub> H <sub>6</sub>	9.5
IsoC <sub>5</sub> H <sub>12</sub>	58.0
N-C <sub>5</sub> H <sub>12</sub>	28.0

Beckman 6800

THC	2670
CH <sub>4</sub>	1610
CO	900

TABLE C-13. RAMS STATION 103

FALL INTENSIVE							
	Mean				Standard Deviation		
THC	2676 ppbC				817 ppbC		
CH <sub>4</sub>	1739 ppbC				367 ppbC		
CO	1073 ppbC				844 ppbC		
C <sub>2</sub> H <sub>2</sub>	15 ppbC				14 ppbC		
Date	11/8/76 7-9	11/8/76 11-13	11/8/76 13-15	11/10/76 7-9	11/10/76 9-11	11/10/76 11-13	11/10/76 13-15
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	19.1	5.0	45.2	9.7	6.4	9.0	7.8
C <sub>2</sub> H <sub>4</sub>	47.2	15.4	9.4	31.0	20.1	15.7	16.0
C <sub>3</sub> H <sub>8</sub>	13.5	4.8	65.7	7.6	7.8	6.0	6.6
C <sub>2</sub> H <sub>2</sub>	26.0	3.9	7.5	17.7	8.2	5.0	6.6
IsoC <sub>4</sub> H <sub>10</sub>	6.8	2.8	17.7	4.5	3.2	4.9	2.8
N-C <sub>4</sub> H <sub>10</sub>	31.5	9.1	63.8	22.2	12.0	43.0	18.4
C <sub>3</sub> H <sub>6</sub>	7.7	2.5	2.4	6.1	3.0	4.3	3.9
IsoC <sub>5</sub> H <sub>12</sub>	30.0	6.0	30.0	17.0	10.0	19.0	7.0
N-C <sub>5</sub> H <sub>12</sub>	14.0	3.0	10.0	7.0	4.0	7.0	5.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>	11			16			
N-C <sub>7</sub> H <sub>16</sub>	3			2			
Toluene	90			41			
N-C <sub>8</sub> H <sub>18</sub>	14			2			
E-C <sub>6</sub> H <sub>5</sub>	21			9			
M-XYL	37			22			
O-XYL	15			9			
N-C <sub>9</sub> H <sub>20</sub>	38			2			
N-P-C <sub>6</sub> H <sub>5</sub>	3			1			
N-C <sub>10</sub> H <sub>22</sub>	258			5			
<u>Beckman 6800</u>							
THC	3066	1785	1850				
CH <sub>4</sub>	2270	1683	1767				
CO	1560	281	336				

(continued)

TABLE C-13 (continued)

Date	11/12/76 9-11	11/12/76 1130 - 1330	11/12/76 1330 - 1530	11/16/76 7-9	11/16/76 9-11	11/16/76 11-13	11/16/76 13-15
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	7.0	6.7	6.3	42.1	30.4	9.0	11.4
C <sub>2</sub> H <sub>4</sub>	11.9	9.2	9.2	121.6	54.7	7.8	13.2
C <sub>3</sub> H <sub>8</sub>	7.7	6.5	6.4	223.8	105.7	18.7	34.1
C <sub>2</sub> H <sub>2</sub>	7.9	4.5	4.8	51.0	39.0	6.0	8.4
IsoC <sub>4</sub> H <sub>10</sub>	2.7	2.9	2.1	65.0	30.7	5.7	8.8
N-C <sub>4</sub> H <sub>10</sub>	11.8	11.5	7.6	174.9	148.2	24.9	43.8
C <sub>3</sub> H <sub>6</sub>	2.2	1.5	1.2	24.6	14.4	1.5	2.4
IsoC <sub>5</sub> H <sub>12</sub>	11.0	9.1	6.4	118.0	75.0	11.0	17.0
N-C <sub>5</sub> H <sub>12</sub>	1.0	1.2	1.0	43.0	31.0	4.0	6.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>	2			48			
N-C <sub>7</sub> H <sub>16</sub>	1			6			
Toluene	6			40			
N-C <sub>8</sub> H <sub>18</sub>	1			5			
E-C <sub>6</sub> H <sub>5</sub>	4			21			
M-XYL	10			48			
O-XYL	4			16			
N-C <sub>9</sub> H <sub>20</sub>				3			
N-P-C <sub>6</sub> H <sub>5</sub>	1			3			
N-C <sub>10</sub> H <sub>22</sub>	1			11			
<u>Beckman 6800</u>							
THC	2010	1990	2960	4470	3360	2090	2280
CH <sub>4</sub>	1500	1480	1470	2550	1890	1440	1480
CO	490	250	280	2640	2160	410	530

(continued)

TABLE C-13 (continued)

Date	11/18/76 7-9	11/18/76 9-11
<u>Dura Pak</u>		
$C_2H_6$	58.6	60.6
$C_2H_4$	23.2	28.3
$C_3H_8$	69.1	84.8
$C_5H_{12}$	15.9	22.7
$IsoC_4H_{10}$	31.6	28.5
$N-C_4H_{10}$	197.3	139.8
$C_3H_6$	6.3	7.8
$IsoC_5H_{12}$	57.0	55.0
$N-C_5H_{12}$	30.0	28.0
<u>Squalane</u>		
$N-C_6H_{14}$	6	
$N-C_7H_{16}$	2	
Toluene	24	
$N-C_8H_{18}$	2	
$E-C_6H_5$	4	
M-XYL	16	
O-XYL	4	
$N-C_9H_{20}$	2	
$N-P-C_6H_5$	1	
$N-C_{10}H_{22}$	5	
<u>Beckman 6800</u>		
THC	2920	2990
$CH_4$	1740	1740
CO	1350	1690

TABLE C-14. RAMS STATION 114

FALL INTENSIVE							
	Mean			Standard Deviation			
THC	2451 ppbC			1299 ppbC			
CH <sub>4</sub>	1541 ppbC			581 ppbC			
CO	1187 ppbC			1392 ppbC			
C <sub>2</sub> H <sub>2</sub>	19 ppbC			22 ppbC			
Date	11/8/76 7-9	11/8/76 13-15	11/10/76 7-9	11/10/76 7-9 Tank (Upwind)	11/10/76 9-11 Tank (Upwind)	11/10/76 11-13	11/10/76 13-15
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	11.7	6.7	9.4	6.9	5.1	9.9	4.9
C <sub>2</sub> H <sub>4</sub>	9.6	10.7	16.6	7.8	3.6	5.6	7.0
C <sub>3</sub> H <sub>8</sub>	8.0	11.2	7.9	6.4	7.2	8.1	4.7
C <sub>2</sub> H <sub>2</sub>	9.0	8.0	15.1	9.2	2.3	1.7	1.6
IsoC <sub>4</sub> H <sub>10</sub>	5.0	6.1	4.4	3.4	1.1	2.4	4.0
N-C <sub>4</sub> H <sub>10</sub>	12.2	29.6	16.4	12.2	3.8	11.1	16.8
C <sub>3</sub> H <sub>6</sub>	3.9	2.3	4.1	2.9	1.9	0.9	
IsoC <sub>5</sub> H <sub>12</sub>	10.0	17.0	15.0	10.0	2.0	11.0	9.0
N-C <sub>5</sub> H <sub>12</sub>	10.0	8.0	6.0	4.0	2.0	6.0	2.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>	8		6	3			
N-C <sub>7</sub> H <sub>16</sub>	3		2	1			
Toluene	19		13	9			
N-C <sub>8</sub> H <sub>18</sub>	1		1	1			
E-C <sub>6</sub> H <sub>5</sub>	7		5	9			
M-XYL	15		16	26			
O-XYL	6		8	10			
N-C <sub>9</sub> H <sub>20</sub>	1		BDL*	1			
N-P-C <sub>6</sub> H <sub>5</sub>	1		1	1			
N-C <sub>10</sub> H <sub>22</sub>	2		1	2			
<u>Beckman 6800</u>							
THC	2654	2254					
CH <sub>4</sub>	2140	1863					
CO	1330	536					

\* Below detection limits

(continued)



TABLE C-14 (continued)

Date	11/12/76 7-9	11/12/76 0930 - 1130	11/12/76 1130 - 1330	11/12/76 1330 - 1530	Tank 11/12/76 7-9 (Upwind)	Tank 11/12/76 9-11 (Upwind)	11/16/76 7-9
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	7.8	6.2	5.7	5.4	8.2	6.5	37.0
C <sub>2</sub> H <sub>4</sub>	11.0	8.3	19.5	4.9	5.3	3.3	94.2
C <sub>3</sub> H <sub>8</sub>	9.6	6.8	3.4	5.3	10.3	7.4	39.4
C <sub>2</sub> H <sub>2</sub>	7.4	3.9	28.2	9.3	8.2	4.8	72.8
IsoC <sub>4</sub> H <sub>10</sub>	1.7	1.2	2.8	1.7	1.8	1.4	78.0
N-C <sub>4</sub> H <sub>10</sub>	7.8	6.2	7.0	5.0	5.9	4.3	235.9
C <sub>3</sub> H <sub>6</sub>	1.6	1.1	1.1	1.1	2.6	1.9	36.1
IsoC <sub>5</sub> H <sub>12</sub>	4.0	4.0	5.0	4.0	4.0	3.0	156.0
N-C <sub>5</sub> H <sub>12</sub>	4.0	2.0	3.0	2.0	3.0	2.0	71.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>	NM**	10			2	5	
N-C <sub>7</sub> H <sub>16</sub>	1				1		
Toluene	4	3			5	3	
N-C <sub>8</sub> H <sub>18</sub>	1	1			1	1	
E-C <sub>6</sub> H <sub>5</sub>	3	1			7	5	
M-XYL	11	4			23	18	
O-XYL	3	1			9	5	
N-C <sub>9</sub> H <sub>20</sub>	Trace	Trace			BDL*	BDL*	
N-P-C <sub>6</sub> H <sub>5</sub>	Trace	Trace			1	BDL*	
N-C <sub>10</sub> H <sub>22</sub>	1	1			2	1	
<u>Beckman 6800</u>							
THC	2030	1890	2060	1970			5170
CH <sub>4</sub>	1540	1470	1460	1450			1880
CO	350	210	170	220			4010

\* Below detection limits

\*\* Not measurable (because of interference etc.)

(continued)

TABLE C-14 (continued)

Date	11/16/76 9-11	11/16/76 13-15	11/18/76 7-9	Tank 11/18/76 7-9 (Upwind)
<u>Dura Pak</u>				
C <sub>2</sub> H <sub>6</sub>	40.3	6.1	66.2	63.0
C <sub>2</sub> H <sub>2</sub>	72.9	38.4	80.2	23.9
C <sub>3</sub> H <sub>8</sub>		5.3	69.6	66.4
C <sub>2</sub> H <sub>2</sub>	69.1	16.7	37.1	32.1
IsoC <sub>4</sub> H <sub>10</sub>	25.3	6.8	19.6	17.3
N-C <sub>4</sub> H <sub>10</sub>	108.9	24.2	67.2	57.1
C <sub>3</sub> H <sub>6</sub>	23.0	7.1	8.7	10.0
IsoC <sub>5</sub> H <sub>12</sub>	84.0	18.0	43.0	34.0
N-C <sub>5</sub> H <sub>12</sub>	39.0	10.0	25.0	19.0
<u>Squalane</u>				
N-C <sub>6</sub> H <sub>14</sub>			24	13
N-C <sub>7</sub> H <sub>16</sub>	9		4	4
Toluene	57		39	37
N-C <sub>8</sub> H <sub>18</sub>	6		3	2
E-C <sub>6</sub> H <sub>5</sub>	17		12	8
M-XYL	43		27	23
O-XYL	17		12	8
N-C <sub>9</sub> H <sub>20</sub>	4		2	1
N-P-C <sub>6</sub> H <sub>5</sub>	4		1	1
N-C <sub>10</sub> H <sub>22</sub>	10		NM*	2
<u>Beckman 6800</u>				
THC	4010		2990	
CH <sub>4</sub>	2190		1770	
CO	3280		1550	

\* Not measurable (because of interference etc.)

TABLE C-15. RAMS STATION 115

	FALL INTENSIVE						
	Mean				Standard Deviation		
THC	2321 ppbC				417 ppbC		
CH <sub>4</sub>	1637 ppbC				225 ppbC		
CO	666 ppbC				388 ppbC		
C <sub>2</sub> H <sub>2</sub>	9 ppbC				7 ppbC		
Date	11/8/76 7-9:45	11/8/76 13-15	11/10/76 7-9	11/10/76 9-11	11/10/76 11-13	11/10/76 13-15	11/12/76 9:30 - 11:30
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	11.7	5.4	8.4	6.4	4.8	5.7	13.0
C <sub>2</sub> H <sub>4</sub>	40.7	27.6	33.4	32.5	42.6	45.2	28.1
C <sub>3</sub> H <sub>8</sub>	20.5	5.4	10.3	7.0	5.3	5.8	63.7
C <sub>2</sub> H <sub>2</sub>	14.9	3.8	7.3	2.1	2.7	2.5	7.4
IsoC <sub>4</sub> H <sub>10</sub>	8.6	1.7	5.4	3.2	1.8	2.2	28.5
N-C <sub>4</sub> H <sub>10</sub>	24.9	8.1	15.1	8.9	7.9	12.9	53.1
C <sub>3</sub> H <sub>6</sub>	5.7	1.2	3.2	0.9	BDL*	BDL*	17.5
IsoC <sub>5</sub> H <sub>12</sub>	19.0	6.0	11.0	6.0	2.0	8.0	33.0
N-C <sub>5</sub> H <sub>12</sub>	9.0	3.0	4.0	3.0	2.0	3.0	14.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>	8		17				14
N-C <sub>7</sub> H <sub>14</sub>	2		1				2
Toluene	12		9				9
N-C <sub>8</sub> H <sub>18</sub>	1		1				2
E-C <sub>6</sub> H <sub>5</sub>	3		5				4
M-XYL	12		17				9
O-XYL	5		6				3
N-C <sub>9</sub> H <sub>20</sub>	1		1				1
N-P-C <sub>6</sub> H <sub>5</sub>	1		1				TRACE
N-C <sub>10</sub> H <sub>22</sub>	4		5				2
<u>Beckman 6800</u>							
THC	2285		1967				2360
CH <sub>4</sub>	2152		1841				1500
CO	768		350				350

\* Below detection limits

(continued)

TABLE C-15 (continued)

Date	11/12/76 1130- 1330	11/12/76 1330- 1530	11/16/76 7-9	11/16/76 9-11	11/16/76 11-13	11/16/76 13-15	11/18/76 7-9
<u>Dura Pak</u>							
C <sub>2</sub> H <sub>6</sub>	8.3	8.1	23.9	15.0	9.2	10.6	65.8
C <sub>2</sub> H <sub>4</sub>	24.4	60.0	54.2	30.9	30.1	32.1	37.9
C <sub>3</sub> H <sub>8</sub>	16.4	19.9	48.5	26.9	20.6	19.6	91.0
C <sub>2</sub> H <sub>2</sub>	6.2	6.7	24.5	13.8	5.8	8.6	17.2
IsoC <sub>4</sub> H <sub>10</sub>	14.5	19.0	11.6	8.0	6.0	5.5	24.0
N-C <sub>4</sub> H <sub>10</sub>	20.9	30.9	41.4	32.5	21.0	19.4	66.9
C <sub>3</sub> H <sub>6</sub>	4.1	5.8	10.6	5.0	1.2	2.1	12.2
IsoC <sub>5</sub> H <sub>12</sub>	13.0	21.0	32.0	16.0	12.0	13.0	33.0
N-C <sub>5</sub> H <sub>12</sub>	5.0	8.0	16.0	9.0	4.0	5.0	19.0
<u>Squalane</u>							
N-C <sub>6</sub> H <sub>14</sub>			8				8
N-C <sub>7</sub> H <sub>16</sub>			3				3
Toluene			15				14
N-C <sub>8</sub> H <sub>18</sub>			3				2
E-C <sub>6</sub> H <sub>5</sub>			6				6
M-XYL			18				15
O-XYL			8				5
N-C <sub>9</sub> H <sub>20</sub>			1				1
N-P-H <sub>5</sub>			1				1
N-C <sub>10</sub> H <sub>22</sub>			4				3
<u>Beckman 6800</u>							
THC	2000		2620	2500	2100	2150	2720
CH <sub>4</sub>	1450		1710	1580	1450	1500	1770
CO	300		1070	840	430	470	980

(continued)

TABLE C-15 (continued)

Date	11/18/76 9-11	11/18/76 11-13	Bag Box 1 11/18/76 13-15	Tank 11/18/76 13-15 (Upwind)	Bag Box 2 11/18/76 13-15
<u>Dura Pak</u>					
$C_2H_6$	65.1	48.4	28.1	27.9	27.8
$C_2H_4$	74.7	65.6	35.1	6.9	19.6
$C_3H_8$	91.2	65.7	35.2	35.5	35.2
$C_2H_2$	26.1	11.1	5.7	5.7	6.7
$IsoC_4H_{10}$	24.3	16.5	11.0	10.6	10.8
$N-C_4H_{10}$	85.5	55.8	40.2	36.9	36.7
$C_3H_6$	7.0	4.5	1.3	2.6	1.0
$IsoC_5H_{12}$	46.0	27.0	21.0	18.0	21.0
$N-C_5H_{12}$	27.0	14.0	10.0	8.0	6.0
<u>Squalane</u>					
$N-C_6H_{14}$				30	21
$N-C_7H_{16}$				1	1
Toluene				10	12
$N-C_8H_{18}$				1	1
$E-C_6H_5$				5	6
M-XYL				17	19
O-XYL				5	5
$N-C_9H_{22}$				1	1
$N-P-C_6H_5$				1	1
$N-C_{10}H_{22}$				2	2
<u>Beckman 6800</u>					
THC	3020	2580	2350		2270
$CH_4$	1830	1710	1580		1560
CO	1610	690	450		410

APPENDIX D  
SPECIAL STUDIES

## CONTENTS

	<u>Page</u>
1.0 INTRODUCTION . . . . .	120

## TABLES

<u>Number</u>		<u>Page</u>
D-1	Roadway Samples Collected and Analyzed 9/22/76 . . . . .	121
D-2	Roadway Samples Collected 9/22/76 Analyzed 9/24/76 . . . . .	123
D-3	Roadway Samples Collected and Analyzed 9/27/76 . . . . .	124
D-4	Roadway Samples Collected and Analyzed 10/29/76. . . . .	125
D-5	C <sub>2</sub> H <sub>4</sub> Bag Samples at Station 101 Collected and Analyzed 10/22/76. . . . .	127
D-6	Ethylene Bag Sampling at Station 101 (with sampling pump). .	128
D-7	Ethylene Bag Sampling at Station 124 (with sampling pump). .	129
D-8	Ethylene Bag Sampling at Station 124 (with sampling pump). .	130



## D.0 INTRODUCTION

Appendix D contains the results obtained from special studies in table form in order to provide an overview of the data.

TABLE D-1. ROADWAY SAMPLES COLLECTED AND ANALYZED 9/22/76 \*

	Babler Park # 5	Babler Park Repeat	Manchester Road	Ultrapure Air	Interstate 40
<u>Dura Pak</u>					
C <sub>2</sub> H <sub>6</sub>	4.1	3.7	22.3	0.3	18.9
C <sub>2</sub> H <sub>4</sub>	5.2	5.4	157.1	2.2	127.3
C <sub>3</sub> H <sub>8</sub>	5.7	5.3	27.6	0.3	14.4
C <sub>2</sub> H <sub>2</sub>	3.0	3.2	111.5	0.9†	83.2
IsoC <sub>4</sub> H <sub>10</sub>	0.8	0.8	15.2	BDL**	11.3
N-C <sub>4</sub> H <sub>10</sub>	3.4	3.3	84.1	1.2	62.7
C <sub>3</sub> H <sub>6</sub>	1.8	1.4	73.7	0.7	60.3
IsoC <sub>5</sub> H <sub>12</sub>	1.7	1.8	125.0	0.6	84.0
N-C <sub>5</sub> H <sub>12</sub>	1.5	1.5	68.0	BDL**	41.0
<u>Squalane</u>					
N-C <sub>6</sub> H <sub>14</sub>			46.0	NM†	64
N-C <sub>7</sub> H <sub>16</sub>			15.0	NM†	32
Toluene	6		72.0	9	165
N-C <sub>8</sub> H <sub>18</sub>	1		12.0	Trace	16
E-C <sub>6</sub> H <sub>5</sub>	9		30	10	68
M-XYL	30		78	33	136
O-XYL	4		30	6	65
N-C <sub>9</sub> H <sub>20</sub>	Trace		4	Trace	4
N-P-C <sub>6</sub> H <sub>5</sub>	BDL**		5	1	6
N-C <sub>10</sub> H <sub>22</sub>	NM†		16	1	NM†
<u>Beckman 6800</u>					
THC	1930		4010	210	3450
CH <sub>4</sub>	1850		2270		1960
CO	290		7850	240	6510

\* Concentrations in ppbC

\*\* Below detection limits

† Not measurable (because of interference etc.)

(continued)

TABLE D-1 (continued)\*

	Interstate 270	Babler Duplicate
<u>Dura Pak</u>		
$C_2H_6$	42.0	4.6
$C_2H_4$	253.0	5.6
$C_3H_8$	54.9	5.6
$C_2H_2$	190.1	3.3
$IsoC_4H_{10}$	19.8	0.8
$N-C_4H_{10}$	121.4	5.3
$C_3H_6$	112.7	1.8
$IsoC_5H_{12}$	159.1	2.9
$N-C_5H_{12}$	79.7	1.5
<u>Squalane</u>		
$N-C_6H_{14}$	38	
$N-C_7H_{16}$	51	
Toluene	239	
$N-C_8H_{18}$	45	
$E-C_6H_5$	107	
M-XYL	277	
O-XYL	91	
$N-C_9H_{20}$	21	
$N-P-C_6H_5$	18	
$N-C_{10}H_{22}$	41	
<u>Beckman 6800</u>		
THC	6250	
$CH_4$	3280	
CO	11360	

\* Concentrations in ppbC

TABLE D-2. ROADWAY SAMPLES COLLECTED 9/22/76 ANALYZED 9/24/76\*

	I-270	I-40	Manchester Road	Ultrapure Air	I-270 Duplicate
<u>Dura Pak</u>					
C <sub>2</sub> H <sub>6</sub>	47.1	19.1	22.2	0.4	40.8
C <sub>2</sub> H <sub>4</sub>	280.1	121.0	148.6	2.8	242.6
C <sub>3</sub> H <sub>8</sub>	61.5	14.2	26.7	0.4	53.1
C <sub>2</sub> H <sub>2</sub>	208.2	78.0	99.5	0.6	184.8
IsoC <sub>4</sub> H <sub>10</sub>	24.2	11.1	14.5	0.2	20.3
N-C <sub>4</sub> H <sub>10</sub>	138.3	61.3	80.6	1.4	118.5
C <sub>3</sub> H <sub>6</sub>	125.0	57.3	76.0	0.9+	108.6
IsoC <sub>5</sub> H <sub>12</sub>	191.0	81.0	119.0	0.7	165.0
N-C <sub>5</sub> H <sub>12</sub>	85.0	38.0	64.0	0.7	77.0
<u>Squalane</u>					
N-C <sub>6</sub> H <sub>14</sub>	89	27	41	1	
N-C <sub>7</sub> H <sub>16</sub>	28	14	15	1	
Toluene	140	67	73	12	
N-C <sub>8</sub> H <sub>18</sub>	25	12	14	2	
E-C <sub>6</sub> H <sub>5</sub>	66	37	36	16	
M-XYL	131	90	92	53	
O-XYL	56	32	36	10	
N-C <sub>9</sub> H <sub>20</sub>	10	1	4	NM +	
N-P-C <sub>6</sub> H <sub>5</sub>	10	2	5	NM +	
N-C <sub>10</sub> H <sub>22</sub>	25	5	9	NM +	

\* Concentrations in ppbC

+ Not Measurable (because of interference etc.)

TABLE D-3. ROADWAY SAMPLES COLLECTED AND ANALYZED 9/27/76 \*

	Bag 3 No. on 270	Bag 3 N. on 270 Repeat	Bag 1 West on Manchester	Bag 4 West on Olive	Bag 5 N. on 270 (Olive-Page)	Bag 1 (Dupl) West on Manchester
<u>Dura Pak</u>						
C <sub>2</sub> H <sub>6</sub>	38.6	38.4	16.1	17.1	21.4	16.0
C <sub>2</sub> H <sub>4</sub>	306.6	305.3	94.0	72.9	115.9	93.9
C <sub>3</sub> H <sub>8</sub>	16.6	16.6	12.0	12.4	14.1	12.2
C <sub>2</sub> H <sub>2</sub>	355.4	361.0	105.5	68.7	67.1	109.6
IsoC <sub>4</sub> H <sub>10</sub>	33.5	33.0	12.0	8.4	13.3	12.0
N-C <sub>4</sub> H <sub>10</sub>	180.3	179.1	57.6	32.0	60.8	57.5
C <sub>3</sub> H <sub>6</sub>	141.2	141.4	39.1	26.4	49.0	39.2
IsoC <sub>5</sub> H <sub>12</sub>	136.0	243.0	75.0	38.0	73.5	60.0
N-C <sub>5</sub> H <sub>12</sub>	135.0	134.0	42.0	19.8	43.4	36.4
<u>Squalane</u>						
N-C <sub>6</sub> H <sub>14</sub>	91		45	30	27	
N-C <sub>7</sub> H <sub>16</sub>	31		11	6	11	
Toluene	126		44	30	51	
N-C <sub>8</sub> H <sub>18</sub>	21		52	6	10	
E-C <sub>6</sub> H <sub>5</sub>	55		93	14	22	
M-XYL	123		88	40	98	
O-XYL	61		27	14	22	
N-C <sub>9</sub> H <sub>20</sub>	6		135	2	3	
N-P-C <sub>6</sub> H <sub>5</sub>	22		894	6	5	
N-C <sub>10</sub> H <sub>22</sub>	16		4331	3	5	
<u>Beckman 6800</u>						
THC	6000		9000	2900	3500	10020
CH <sub>4</sub>	1910		1800	1850	1940	1840
CO	14600		5850	5190	5540	5890

\* Concentrations in ppbC

TABLE D-4. ROADWAY SAMPLES COLLECTED AND ANALYZED 10/29/76\*

	W. on Manchester I-270 to Weidman	W. on Manchester Weidman to Ries	S. on I-270 Manchester to I-44
<u>Dura Pak</u>			
C <sub>2</sub> H <sub>6</sub>	36.1	53.6	33.0
C <sub>2</sub> H <sub>4</sub>	215.1	259.1	207.9
C <sub>3</sub> H <sub>8</sub>	47.0	40.5	48.5
C <sub>2</sub> H <sub>2</sub>	205.4	257.8	134.7
Iso-C <sub>4</sub> H <sub>10</sub>	32.0	33.0	19.9
N-C <sub>4</sub> H <sub>10</sub>	153.3	194.3	114.8
C <sub>3</sub> H <sub>6</sub>	95.6	114.1	93.7
IsoC <sub>5</sub> H <sub>12</sub>	145.2	182.6	116.7
N-C <sub>5</sub> H <sub>12</sub>	61.4	75.8	62.9
<u>Squalane</u>			
N-C <sub>6</sub> H <sub>14</sub>	38	74	44
N-C <sub>7</sub> H <sub>16</sub>	19	25	20
Toluene	94	119	83
N-C <sub>8</sub> H <sub>18</sub>	12	20	20
E-C <sub>6</sub> H <sub>5</sub>	38	41	24
M-XYL	90	129	74
O-XYL	47	52	41
N-C <sub>9</sub> H <sub>20</sub>	5	8	31
N-P-C <sub>6</sub> H <sub>5</sub>	7	12	6
N-C <sub>10</sub> H <sub>22</sub>	9	15	NM**
<u>Beckman 6800</u>			
THC	5230	6410	5000
CH <sub>4</sub>	2350	2690	2090
CO	12360	15660	8970

\* Concentration in ppbC

(continued)

\*\* Not measurable (because of interference etc.)

TABLE D-4 (continued)\*

	W. on Olive I-44 to Manchester	W. on Olive I-270 to Eatherton	Dupl. Analysis W. on Manchester I-270 to Weidman
<u>Dura Pak</u>			
$C_2H_6$	29.8	5.3	37.3
$C_2H_4$	206.0	6.6	220.4
$C_3H_8$	42.3	11.8	47.8
$C_2H_2$	127.4	5.8	208.4
Iso- $C_4H_{10}$	17.1	2.2	33.0
N- $C_4H_{10}$	95.5	7.7	156.8
$C_3H_6$	95.9	2.4	95.2
Iso- $C_5H_{12}$	99.5	9.8	148.9
N- $C_5H_{12}$	51.0	2.2	61.0
<u>Squalane</u>			
N- $C_6H_{14}$	28	BDL	38
N- $C_7H_{16}$	15	1	21
Toluene	68	5	99
N- $C_8H_{18}$	10	2	13
E- $C_6H_5$	21	3	32
M-XYL	62	5	88
O-XYL	34	5	39
N- $C_9H_{20}$	4	1	5
N-P- $C_6H_5$	9	NM**	3
N- $C_{10}H_{22}$	14	NM**	8
<u>Beckman 6800</u>			
THC	4290	2030	- -
$CH_4$	1990	1720	- -
CO	9580	350	- -

\* Concentration in ppbC

\*\* Not measurable (because of interference etc.)

TABLE D-5. C<sub>2</sub>H<sub>4</sub> BAG SAMPLES AT STATION 101 COLLECTED & ANALYZED 10/22/76\*

	Dura Pak								
	C <sub>2</sub> H <sub>6</sub>	C <sub>2</sub> H <sub>4</sub>	C <sub>3</sub> H <sub>8</sub>	C <sub>2</sub> H <sub>2</sub>	ISO-C <sub>4</sub> H <sub>10</sub>	N-C <sub>4</sub> H <sub>10</sub>	C <sub>3</sub> H <sub>6</sub>	ISO-C <sub>5</sub> H <sub>12</sub>	N-C <sub>5</sub> H <sub>10</sub>
BAG BOX 7-8	27.7	70.7	17.6	62.1	23.8	117.5	26.3	97.0	51.0
BAG BOX (REPEAT) 7-8	27.7	70.7	17.5	60.8	22.8	116.7	24.5	96.0	53.0
AUTO (UPWIND) 7-8	29.7	61.1	26.0	68.4	79.2	485.8	26.8	231.0	103.0
BAG BOX 8-9	24.0	71.6	19.8	57.3	28.6	143.0	23.4	107.0	52.0
AUTO (UPWIND) 8-9	21.9	53.8	19.1	53.7	21.9	98.4	25.3	91.0	48.0
Between PUMP BOX and BAG BOX 9-10	12.7	1740.5	18.0	21.5	17.5	134.7	8.4	58.0	29.0
CENTERMOST BLOWER OUTLET 10-11	11.0	9199.4	34.4	8.2	12.0	55.9	5.0	38.0	12.0
INOPERATIVE BLOWER OUTLET 11-12	8.2	92.9	16.0	10.1	7.6	27.7	3.6	21.0	10.0
OUTLET OF VACUUM PUMP 12-13	22.6	9042.8	32.9	11.6	7.7	32.2	17.5	24.0	12.0
PUMP BOX INTERIOR 13-14	9.5	1486.1	20.1	12.0	12.5	105.3	6.6	39.0	14.0
BAG BOX (DUPLICATE) 7-8	28.2	72.5	17.2	62.7	22.5	116.0	25.7	98.0	54.0

\* Concentrations in ppbC



TABLE D-6. ETHYLENE BAG SAMPLING AT STATION 101 (WITH AIR SAMPLING PUMP)\*

	C <sub>2</sub> H <sub>6</sub>	C <sub>2</sub> H <sub>4</sub>	C <sub>3</sub> H <sub>8</sub>	C <sub>2</sub> H <sub>2</sub>	ISO-C <sub>4</sub> H <sub>10</sub>	N-C <sub>4</sub> H <sub>10</sub>	C <sub>3</sub> H <sub>6</sub>	ISO-C <sub>5</sub> H <sub>12</sub>	N-C <sub>5</sub> H <sub>12</sub>
BAG 1 9-9:15	22.1	36.1	31.2	33.5	27.9	430.2	15.0	84.0	25.0
BAG 1 9-9:15	22.1	35.9	31.2	33.5	27.9	427.4	15.0	82.0	25.0
TANK 1 9-9:15	20.3	271.0	30.4	31.3	27.1	418.8	13.0	73.0	26.0
BAG 2 10:10-10:15	13.7	28.9	20.0	18.5	21.6	207.7	8.0	65.0	25.0
TANK 2 10:10-10:15	13.2	21.9	20.4	19.2	20.8	223.4	8.0	59.0	30.0
BAG 3 10:45-11	10.1	108.2	12.6	12.5	11.7	65.7	5.0	40.0	14.0
TANK 3 10:45-11	9.5	13.0	12.0	12.4	11.7	64.5	5.0	39.0	18.0
BAG BOX (NO PUMP) 9-11	16.6	1765.4	24.9	29.3	23.5	259.5	11.0	75.0	24.0
BAG 1 9-9:15 Duplicate	21.6	36.4	31.3	34.3	27.6	431.5	14.0	77.0	30.0

\* Concentrations in ppbC Collected and Analyzed 11/2/76

TABLE D-7. ETHYLENE BAG SAMPLING AT STATION 124 (WITH AIR SAMPLING PUMP)\*

	Dura Pak									
	C <sub>2</sub> H <sub>6</sub>	C <sub>2</sub> H <sub>4</sub>	C <sub>3</sub> H <sub>8</sub>	C <sub>2</sub> H <sub>2</sub>	ISO-C <sub>4</sub> H <sub>10</sub>	N-C <sub>4</sub> H <sub>10</sub>	C <sub>3</sub> H <sub>6</sub>	ISO-C <sub>5</sub> H <sub>12</sub>	N-C <sub>5</sub> H <sub>10</sub>	
BAG 2 10-10:30	4.3	7.5	4.8	1.6	0.8	4.4	0.7	4.0	2.0	
BAG 2 (REPEAT) 10-10:30	4.5	7.5	4.8	1.9	1.1	4.5	0.7	4.0	2.0	
TANK 2 10-10:30	4.5	6.2	5.6	2.6	2.0	4.0	1.1	5.0	3.0	
BAG 3 10:45-11:10	4.5	7.9	4.4	1.5	0.9	3.3	0.7	3.0	2.0	
TANK 3 10:45-11:10	4.5	4.5	4.0	1.9	0.8	2.3	0.9	2.0	2.0	
BAG 4 (BAG BOX - NO PUMP) 10-11:15	4.7	21.4	4.8	1.2	1.3	7.2	1.0	5.0	2.0	
TANK 1 (OUTSIDE AIR) 9:30-9:45	4.4	2.2	5.8	2.0	2.0	15.2	1.1	4.0	3.0	
BAG 5 (STATION AIR)	5.8	2716.8	7.2	1.9	1.6	6.0	1.1	2.0	2.0	
BAG 2 (DJPLICATE) 10-10:30	4.3	7.3	4.9	1.6	0.6	4.2	0.7	4.0	2.0	

\* Concentrations in ppbC

Collected and Analyzed on 11/3/76

TABLE D-8. ETHYLENE BAG SAMPLING AT STATION 124 (WITH AIR SAMPLING PUMP)\*

	Dura Pak												
	C <sub>2</sub> H <sub>6</sub>	C <sub>2</sub> H <sub>4</sub>	C <sub>3</sub> H <sub>8</sub>	C <sub>2</sub> H <sub>2</sub>	ISO-C <sub>4</sub> H <sub>10</sub>	N-C <sub>4</sub> H <sub>10</sub>	C <sub>3</sub> H <sub>6</sub>	ISO-C <sub>5</sub> H <sub>12</sub>	N-C <sub>5</sub> H <sub>12</sub>				
BAG 2 9:55-10:10	5.4	11.4	4.7	4.9	2.2	8.8	2.0	9.0	3.0				
BAG 2 (REPEAT) 9:55-10:10	6.1	11.3	4.7	3.8	1.5	8.5	1.0	8.0	3.0				
TANK 2 9:55-10:10	5.2	8.1	4.3	4.5	1.4	5.4	1.0	4.0	2.0				
BAG 3 10:45-11	4.7	9.4	3.8	3.3	1.7	7.1	3.0	4.0	3.0				
TANK 3 10:45-11	4.7	8.6	3.6	2.8	1.1	4.1	1.0	3.0	3.0				
BAG 4 (STATION AIR -1) 10:20-10:45	5.2	2013.2	7.8	4.1	1.9	8.8	2.0	6.0					
TANK 1 (AIR UPWIND OF STATION)	5.5	4.2	5.4	3.5	4.1	34.2	2.0	16.0	3.0				
BAG 5 (BAG BOX-NO PUMP) 9:30-11	5.3	13.8	4.7	4.5	1.6	6.9	2.0	4.0	2.0				
BAG 6 (STATION AIR -2 LEE SIDE OF STATION)	4.6	923.5	5.3	3.5	1.4	5.7	1.0	3.0	2.0				
TANK 4 (AIR UPWIND OF STATION)	4.7	3.3	4.2	3.0	2.2	19.1	2.0	11.0	4.0				
BAG 2 (DUPLICATE) 9:55-10:10	5.3	11.4	4.6	4.3	1.7	8.1	1.0	5.0	3.0				

\* Concentrations in ppbC

Collected and Analyzed 11/4/76

<b>TECHNICAL REPORT DATA</b> <i>(Please read Instructions on the reverse before completing)</i>		
1. REPORT NO. EPA-600/4-80-006	2.	3. RECIPIENT'S ACCESSION NO.
4. TITLE AND SUBTITLE REGIONAL AIR POLLUTION STUDY Gas Chromatography Laboratory Operation		5. REPORT DATE January 1980
7. AUTHOR(S) G. Cardwell		6. PERFORMING ORGANIZATION CODE
9. PERFORMING ORGANIZATION NAME AND ADDRESS Rockwell International Environmental Monitoring & Services Center 11640 Administration Drive Creve Coeur, MO 63141		8. PERFORMING ORGANIZATION REPORT NO.
12. SPONSORING AGENCY NAME AND ADDRESS Environmental Sciences Research Laboratory - RTP, N.C. Office of Research and Development U.S. Environmental Protection Agency Research Triangle Park, N.C. 27711		10. PROGRAM ELEMENT NO. 1AA603 AA-126 (FY-79)
		11. CONTRACT/GRANT NO. 68-02-2093 Task Order 113
		13. TYPE OF REPORT AND PERIOD COVERED Final
		14. SPONSORING AGENCY CODE EPA/600/09
15. SUPPLEMENTARY NOTES		
16. ABSTRACT  A gas chromatography laboratory was set up to analyze air samples collected in Teflon bags and stainless steel tanks. Samples were analyzed for total hydrocarbons, methane, CO, and C <sub>2</sub> -C <sub>10</sub> hydrocarbons. A total of 455 samples, including replicates, were analyzed during the summer and fall of 1976. Many samples were collected at 12 of the Regional Air Monitoring Systems (RAMS) sites to yield data on spatial and temporal distributions of hydrocarbons. Additional sampling was performed to study ethylene contamination in and around RAMS stations. Roadway samples were collected to determine the composition of freshly emitted vehicular pollution. Quality control audits indicated good system performance during the study. Replicate samples indicated good reproducibility for samples stored for as long as six days in the Teflon bags.  All data, including sums of paraffins, olefins, aromatic, and total non-methane hydrocarbons are stored in the RAPS Data Bank at Research Triangle Park, North Carolina.		
17. KEY WORDS AND DOCUMENT ANALYSIS		
a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
*Air pollution *Hydrocarbons *Carbon monoxide *Gas chromatography *Chemical laboratories	Regional Air Pollution Study St. Louis, MO	13B 07C 07B 07D 14D
18. DISTRIBUTION STATEMENT  RELEASE TO PUBLIC	19. SECURITY CLASS (This Report) UNCLASSIFIED	21. NO. OF PAGES 139
	20. SECURITY CLASS (This page) UNCLASSIFIED	22. PRICE