

LAKE MICHIGAN STUDIES

Special Report Number LM 5

MICROBIOLOGICAL INVESTIGATIONS

April 1963

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE  
Public Health Service  
Division of Water Supply and Pollution Control  
Great Lakes-Illinois River Basins Project



## TABLE OF CONTENTS

INTRODUCTION	1
PROJECT INVESTIGATIONS	2
Parameters	2
Field Procedures	3
Laboratory Procedures	3
Findings	4
Discussion and Significance of Findings	8
OTHER MICROBIOLOGICAL DATA	10
Coliform Densities at Various Drinking Water Intakes and Bathing Beaches	10
Pollution Persistency	12
ULTIMATE SURVIVAL OF PATHOGENS IN LAKE MICHIGAN	14
Possible Effect of Returning Treated Effluent to Lake Michigan	15
CONCLUSIONS	17
REFERENCES	18
TABLES	
FIGURES	



## LIST OF TABLES

### TABLE

- 1 Coliform Results From Deepwater and Inshore Studies
- 2 20°C Total Plate Count Results From Deepwater and Inshore Studies
- 3 35°C Total Plate Count Results From Deepwater and Inshore Studies
- 4 Coliform Results From Harbor Studies
- 5 20°C Total Plate Count Results From Harbor Studies
- 6 35°C Total Plate Count Results From Harbor Studies
- 7 Coliform Levels at Three Points Adjacent to Central District Filtration Plant, Chicago, Illinois, 1961
- 8 Coliform Incidence at the Chicago Beaches, Summer, 1961
- 9 Coliform Bacteria Per 100 ml at Dunne Crib in 1951, 1956, and 1961
- 10 Summary of Coliform Levels at Dunne Crib for 1961



## LIST OF FIGURES

### FIGURE

- 1        Distribution of Coliform Bacteria per 100 ml, S.W. Quadrant  
         of Lake Michigan
- 2        Distribution of Coliform Bacteria per 100 ml, S.E. Quadrant  
         of Lake Michigan
- 3        Distribution of Coliform Bacteria per 100 ml, Northern  
         Half of Lake Michigan
- 4        Coliform Densities in 10 Mile Zone Along West Shore of  
         Lake Michigan
- 5        Coliform Densities in 10 Mile Zone Along East Shore of  
         Lake Michigan
- 6        20°C Total Bacterial Densities in 10 Mile Zone Along  
         West Shore of Lake Michigan
- 7        20°C Total Bacterial Densities in 10 Mile Zone Along  
         East Shore of Lake Michigan
- 8        35°C Total Bacterial Densities in 10 Mile Zone Along  
         West Shore of Lake Michigan
- 9        35°C Total Bacterial Densities in 10 Mile Zone Along  
         East Shore of Lake Michigan
- 10       Distribution of 20°C Total Bacteria per 100 ml,  
         S.W. Quadrant of Lake Michigan
- 11       Distribution of 20°C Total Bacteria per 100 ml,  
         S.E. Quadrant of Lake Michigan
- 12       Distribution of 20°C Total Bacteria per 100 ml,  
         Northern Half of Lake Michigan
- 13       Distribution of Coliform Bacteria per 100 ml in  
         Chicago Harbor
- 14       Distribution of Coliform Bacteria per 100 ml in  
         Racine Harbor





LIST OF FIGURES (cont.)

- 15      Distribution of Coliform Bacteria per 100 ml in  
         Milwaukee Harbor
- 16      Distribution of 20°C Total Bacteria per 100 ml in  
         Chicago Harbor
- 17      Distribution of 20°C Total Bacteria per 100 ml in  
         Racine Harbor
- 18      Distribution of 20°C Total Bacteria per 100 ml in  
         Racine Harbor
- 19      Location of Chicago Drinking Water Intakes and Beaches
- 20      Survival of Coliform Bacteria and Fecal Streptococci  
         in Lake Michigan Water at 25°C with Illumination
- 21      Survival of Coliform Bacteria and Fecal Streptococci  
         in Lake Michigan Water at 25°C Without Illumination
- 22      Survival of Coliform Bacteria and Fecal Streptococci in  
         Lake Michigan Water at 5°C Without Illumination

1. The first part of the paper

2. The second part of the paper

3. The third part of the paper

4. The fourth part of the paper

5. The fifth part of the paper

6. The sixth part of the paper

7. The seventh part of the paper

8. The eighth part of the paper

## INTRODUCTION

This report is based on (1) a preliminary study of the microbiology of Lake Michigan conducted by the Great Lakes-Illinois River Basins Project during the spring, summer, and autumn months of 1962 (April-December); (2) a review of raw water data of several water plant intakes withdrawing water from Lake Michigan; (3) data from samples of bathing beach waters in the Chicago area; and (4) laboratory studies of survival of coliform and fecal streptococci in Lake Michigan water.

The study of the microbiology of Lake Michigan conducted by the Great Lakes-Illinois River Basins Project was achieved through a series of eight cruises on vessels equipped with laboratories for performing microbiological procedures. It was thereby possible to process all samples immediately for analysis of bacterial content. The locations of sampling stations are shown in Figures 1, 2, and 3 of the present report.

The raw water data review consisted of an analysis of data obtained from records of certain water treatment plants in the Chicago area. The bathing beach data were obtained from the Chicago Park district, and the survival studies were carried out by the Great Lakes-Illinois River Basins Project laboratory.

The purpose of the present report is to present the findings on the present status of the microbiology of Lake Michigan and to discuss the possible effects of returning treated sewage effluent from the City of Chicago to Lake Michigan.



## PROJECT INVESTIGATIONS

The objective of the microbiological program was to determine the present water quality of Lake Michigan in terms of bacterial parameters. Qualitative and quantitative aspects were incorporated so that the distribution of the pertinent bacterial varieties with respective densities throughout might be ascertained. The various determinations utilized were selected to indicate the sanitary quality of the water as well as general biological productivity.

Parameters

Analyses were made for coliform content, and for total plate counts at 20°C and 35°C. In certain of the harbor studies, some of the samples were analyzed for fecal streptococcus content. Approximately 1,400 samples comprised the coliform study. Total plate counts at 20°C and 35°C were made on approximately 1,400 samples.

Coliform bacteria are defined in Standard Methods for the Examination of Water and Wastewater (1)\* as "including all of the aerobic and facultative anaerobic, Gram-negative, non-sporeforming, rod-shaped bacteria which ferment lactose with gas formation within 48 hours at 35°C.

Coliform organisms are significant to water quality since these bacteria occur in the fecal matter of all warm-blooded animals, including man. Consequently, the presence of coliform bacteria in a body of water is interpreted as indicative of contamination of the water by fecal matter. Since contamination of water by fecal matter is one avenue of transmission of certain water-borne diseases to humans, coliform bacteria are utilized as indicators of possible pathogenic contamination. Increasing densities of coliform bacteria found in water are, therefore, related to the increasing degree of pollution of enteric origin and to increasing health hazard to those exposed to the water.

Fecal streptococci, as used in the present study, includes any species of streptococci commonly present in significant numbers in the fecal excreta of humans or other warm-blooded animals (2). Streptococci are Gram-positive cocci occurring in chains composed of varying numbers of cells. These organisms may be parasitic or saprophytic. The fecal streptococci, like the coliform bacteria,

\*See references listed at end of report.

... ..  
... ..  
... ..  
... ..  
... ..  
... ..

...

... ..  
... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..  
... ..  
... ..  
... ..  
... ..  
... ..  
... ..

... ..  
... ..  
... ..  
... ..  
... ..  
... ..  
... ..  
... ..

are enteric organisms abounding in the fecal matter of all warm-blooded animals; they likewise indicate the presence of fecal matter in water. Fecal streptococcus results augment coliform findings in that the streptococci indicate recent fecal contamination of water, whereas the coliform bacteria may derive from more remote contamination.

Total plate counts, according to Standard Methods (1), are approximate enumerations of total bacteria multiplying at temperatures of 35°C and 20°C on any one of several nutrient agars; they yield useful information concerning the quality of the water tested and may provide supporting data regarding the significance of the results of the coliform test. (The medium of choice for the present investigations was tryptone glucose extract agar.)

Total bacterial densities in water correspond to the decomposition of organic matter. In this large group there are species which grow well at several incubation temperatures. These are present in most waters in small numbers and, in waters enriched with organic matter, they occur in great abundance. Several species of this group grow best at temperatures ranging from 5°C to 20°C. Other species have become semi-parasitic, being especially adapted to the decomposition of organic matter in the intestines of warm-blooded animals. These latter forms develop most actively at body temperatures (35°C to 37°C) (15).

#### Field Procedures

The samples were collected at each station at fixed depths: surface (designated "0" meters), 5, 10, 20, 30, 50, 75, 100, 150, and 250 meters. The number of samples at a given station was therefore determined by the depth of the lake at that point. The deepest samples collected at a station are referred to in this report as "lowermost." The samples were collected in three zones: deep water, inshore, and city harbors. The inshore investigations utilized samples collected at intervals of 1 mile, 4 miles, and 10 miles offshore. The inshore work was devoted largely to the southern half of Lake Michigan.

All bacterial samples were collected in Zobell J-Z water samplers, permitting individual sample collections from the various depths in sterile glass bottles which remained sealed until sampling depth was reached.

#### Laboratory Procedures

All bacterial determinations were made in accordance with the procedures set forth in Standard Methods for the Examination of Water and Wastewater (1), Eleventh Edition, 1960, pp 477-526, or in accordance with modifications of these procedure as established through research at the Robt.A.Taft Sanitary Engr.Center, Cincinnati, Ohio.

1. The first part of the report deals with the general situation of the country and the position of the various groups of the population. It is a very interesting and informative study of the social and economic conditions of the country.

2. The second part of the report deals with the political situation of the country. It is a very interesting and informative study of the political conditions of the country.

3. The third part of the report deals with the economic situation of the country. It is a very interesting and informative study of the economic conditions of the country.

4. The fourth part of the report deals with the cultural situation of the country. It is a very interesting and informative study of the cultural conditions of the country.

5. The fifth part of the report deals with the military situation of the country. It is a very interesting and informative study of the military conditions of the country.

6. The sixth part of the report deals with the international situation of the country. It is a very interesting and informative study of the international conditions of the country.



These latter modifications were related to the media constituents of the streptococcus medium (2). Coliform and fecal streptococcus determinations and total plate counts were made by the membrane filter technique. Results for coliform, fecal streptococci, and total plate counts are reported on the basis of number of organisms per 100 milliliters (ml).

### Findings

#### Deepwater and Inshore Studies

Coliform Findings. Table 1 shows that, of 313 surface samples collected at the deepwater and inshore points, 134 (42.8%) contained a density of less than 1 coliform bacterium per 100 ml. A total of 233 (74.6%) samples contained 10 coliform bacteria or less per 100 ml and a total of 292 (93.4%) showed coliform densities of 100 or less. An additional 17 samples showed coliform densities in the range of 100-1,000 per 100 ml. Only 4 samples (1.3%) exceeded 1,000 per 100 ml and these were in the range 1,000-2,000 per 100 ml. These latter 4 samples were from sampling points located within one mile offshore and adjacent to Gary, Indiana, and Racine, Wisconsin. Also shown in Table 1 is a summary of results from the deepest samples collected. Examination of test results from all samples (approximately 1,400) showed that, in general, variations in coliform density did not reflect significant differences in water quality with respect to depth. Therefore, further details of this aspect are not included.

The geographical distribution of the coliform densities derived from the surface samples is shown in Figures 1, 2, and 3. An examination of these map graphs shows that in the southern basin the higher coliform densities are located adjacent to the shoreline and correlate with the centers of urbanization. The western shoreline from Milwaukee to Gary showed consistently higher coliform densities than did the eastern shore. Beyond this zone of contamination, the deep water in the majority of samples showed little or no coliform content (reported as less than 1 per 100 ml). These relationships are presented in Figures 4 and 5 showing coliform densities as they were measured at intervals of 1 mile, 4 miles, and 10 miles offshore. Diminishing coliform densities consistently occurred from one mile offshore to 10 miles offshore around the periphery of the southern half of Lake Michigan. All 10-mile samples from Gary, Indiana to Muskegon, Michigan, failed to give positive coliform findings in 100 ml samples.

These findings, in general, indicate that the bacteriological quality of Lake Michigan water is excellent where not locally contaminated through domestic sewage entering the lake.

1. The first part of the document is a list of the names of the persons who have been appointed to the various offices of the city of New York.

2. The second part of the document is a list of the names of the persons who have been appointed to the various offices of the city of New York.

3. The third part of the document is a list of the names of the persons who have been appointed to the various offices of the city of New York.

4. The fourth part of the document is a list of the names of the persons who have been appointed to the various offices of the city of New York.

5. The fifth part of the document is a list of the names of the persons who have been appointed to the various offices of the city of New York.

6. The sixth part of the document is a list of the names of the persons who have been appointed to the various offices of the city of New York.

7. The seventh part of the document is a list of the names of the persons who have been appointed to the various offices of the city of New York.

8. The eighth part of the document is a list of the names of the persons who have been appointed to the various offices of the city of New York.

Total Plate Count Findings. The bacterial densities of species growing at 20°C and 35°C parallel those of the coliform bacteria. The lowest densities (100 to 500 per 100 ml) were found in the central body of the lake (deep water). Highest densities were observed in the inshore areas in relation to the geographical location of cities and towns. (See Figures 6, 7, 8, 9, 10, 11, and 12). Approximately 25% of the samples contained densities (20°C and 35°C) greater than 10,000 per 100 ml.

Highest 20°C densities encountered were approximately 500,000 per 100 ml (near Racine, Wisconsin, at 1 mile offshore). Other areas showing densities in excess of 100,000 per 100 ml were offshore from Milwaukee, Racine, Kenosha, Chicago and Michigan City.

The highest densities observed with the 35°C test were in excess of 500,000 per 100 ml. These occurred at Milwaukee, Kenosha, and Gary. The highest 35°C density was approximately 1,300,000 per 100 ml approximately 7 miles off the Chicago waterfront.

Tables 2 and 3 show the per cent of samples which contained various total densities. The cumulative per cent distribution of the 20°C and 35°C total densities is also shown. Results in these two tables originate from surface samples. The wide range in the numbers present throughout the lake, and the considerable number of samples showing high densities, are indicative of organic enrichment of these waters.

#### Harbor Studies

Three city harbors were investigated: Chicago, Racine and Milwaukee. Samples were collected in the immediate harbor area and in the adjacent waters in a radial zone extending 3-5 miles around the chief river mouth emptying into the harbor.

The highest coliform densities encountered were associated with and located in the Milwaukee Harbor. The next highest densities were found in the Racine Harbor area. The Chicago Harbor area showed the least coliform occurrence of the three harbors studied. Highest total plate counts occurred at those stations showing highest coliform densities. Sampling stations are shown in Figures 13, 14, and 15.



Coliform Findings. Fifteen samples were collected from 15 sampling points in the Chicago Harbor study. All samples contained fewer than 1,000 coliform bacteria per 100 ml, while 6 (86.7%) contained fewer than 500 coliform per 100 ml. Seven of the samples (46.7%) were in the 1-50 per 100 ml range and 3 (20.0%) of the samples were in the 1-10 per 100 ml range. (See Figure 13 and Table 4).

The higher coliform levels (100-1,000 per 100 ml) were encountered in an area immediately adjacent to the mouth of the Chicago River (both inside and outside the breakwater harbor) and extending south of the harbor for approximately 2 miles in the waters about one-half mile or less offshore.

In the Racine Harbor higher coliform densities were encountered, with 38.2% of the 34 samples in the study containing coliform levels in the range of 1,000-10,000 per 100 ml. The highest coliform density (in the 5,000-10,000 per 100 ml range) was encountered at the mouth of the Root River which empties into the breakwater harbor. Coliform densities in the next lower range (1,000-5,000 per 100 ml) were encountered at the harbor mouth and in a zone extending south of the harbor mouth for approximately one mile in the waters which were sampled approximately 1,500 feet offshore. However, the remainder of the radial area surrounding the harbor for a distance of one to two miles uniformly contained coliform densities in 100-1,000 per 100 ml range. (See Figure 14).

The Milwaukee Harbor manifested the highest coliform densities, with 15.9% of a total of 76 samples tested containing coliform bacteria in the 10,000-40,000 range. The highest coliform densities (in the 10,000-40,000 per 100 ml range) were encountered at the mouth of the Milwaukee River and within the breakwater in the southern two-thirds of the harbor. The northern third of the harbor contained coliform densities in the 1,000-10,000 range. A breakwater leads south from the harbor for a distance of some two miles. The coliform densities in the waters between this breakwater and the shore progressively decreased from 5,000-10,000 per 100 ml at the harbor inlet to 100-1,000 per 100 ml where the open water is reached. In the open waters adjacent to the harbor breakwater, the northern area showed very little or no coliform content. Most samples originating in the open waters surrounding the southern half of the harbor breakwater contained coliform densities ranging from 10-100 per ml at the central harbor mouth to 100-1,000 per 100 ml in the southern most waters adjacent to the breakwater running south for approximately two miles. (See Figure 15).

1. The first part of the paper discusses the importance of the study of the history of the United States. It is argued that a knowledge of the past is essential for a full understanding of the present and for the development of a sound policy for the future. The author points out that the study of history is not only a means of satisfying our curiosity about the past, but also a means of training the mind and of developing the character. It is through the study of history that we learn the lessons of the past and are enabled to avoid the mistakes of our predecessors. The author also emphasizes the importance of the study of the history of the United States, particularly in the light of the fact that the country is now facing many of the same problems that its ancestors faced in the past. It is only by a study of the history of the United States that we can understand the causes of these problems and find the best way to deal with them.

2. The second part of the paper discusses the importance of the study of the history of the United States. It is argued that a knowledge of the past is essential for a full understanding of the present and for the development of a sound policy for the future. The author points out that the study of history is not only a means of satisfying our curiosity about the past, but also a means of training the mind and of developing the character. It is through the study of history that we learn the lessons of the past and are enabled to avoid the mistakes of our predecessors. The author also emphasizes the importance of the study of the history of the United States, particularly in the light of the fact that the country is now facing many of the same problems that its ancestors faced in the past. It is only by a study of the history of the United States that we can understand the causes of these problems and find the best way to deal with them.

3. The third part of the paper discusses the importance of the study of the history of the United States. It is argued that a knowledge of the past is essential for a full understanding of the present and for the development of a sound policy for the future. The author points out that the study of history is not only a means of satisfying our curiosity about the past, but also a means of training the mind and of developing the character. It is through the study of history that we learn the lessons of the past and are enabled to avoid the mistakes of our predecessors. The author also emphasizes the importance of the study of the history of the United States, particularly in the light of the fact that the country is now facing many of the same problems that its ancestors faced in the past. It is only by a study of the history of the United States that we can understand the causes of these problems and find the best way to deal with them.

4. The fourth part of the paper discusses the importance of the study of the history of the United States. It is argued that a knowledge of the past is essential for a full understanding of the present and for the development of a sound policy for the future. The author points out that the study of history is not only a means of satisfying our curiosity about the past, but also a means of training the mind and of developing the character. It is through the study of history that we learn the lessons of the past and are enabled to avoid the mistakes of our predecessors. The author also emphasizes the importance of the study of the history of the United States, particularly in the light of the fact that the country is now facing many of the same problems that its ancestors faced in the past. It is only by a study of the history of the United States that we can understand the causes of these problems and find the best way to deal with them.

5. The fifth part of the paper discusses the importance of the study of the history of the United States. It is argued that a knowledge of the past is essential for a full understanding of the present and for the development of a sound policy for the future. The author points out that the study of history is not only a means of satisfying our curiosity about the past, but also a means of training the mind and of developing the character. It is through the study of history that we learn the lessons of the past and are enabled to avoid the mistakes of our predecessors. The author also emphasizes the importance of the study of the history of the United States, particularly in the light of the fact that the country is now facing many of the same problems that its ancestors faced in the past. It is only by a study of the history of the United States that we can understand the causes of these problems and find the best way to deal with them.

The bacterial findings at each of the harbors indicated that the harbor waters mingle with the surrounding waters of Lake Michigan and that the direction of flow is to the south.

Fecal Streptococcus Findings. Fecal streptococcus determinations were included in the Milwaukee and Racine Harbor studies. The maximal density of fecal streptococci encountered in the Milwaukee Harbor samples, originating from surface waters, was 19,000 per 100 ml. This sample was collected in the Harbor, just north of mouth of the Milwaukee River. At the mouth of the Harbor (opposite the mouth of the Milwaukee River) a fecal streptococcus density of 40 per 100 ml was observed. A level of 300 per 100 ml was observed at one point just south of the southern inlet into the harbor. Samples in the open waters adjacent to the northern half of the Harbor were reported as less than 10 per 100 ml.

The streptococcal densities from the Racine Harbor were 200 per 100 ml at the mouth of the Root River, with a density of 400 per 100 ml at the main Harbor mouth; in the waters south of the harbor densities in the range of 10-610 per 100 ml were observed. Observations to the north of the Harbor were 10 and 20 per 100 ml.

Total Plate Count Findings. As previously stated, the results of the total plate counts agreed with the coliform findings, showing high counts at sampling stations where maximum coliform densities occurred and generally decreasing as coliform densities decreased.

In the Chicago Harbor study, 9 of the 17 samples showed total densities in the 50,000-500,000 range as estimated by the 20°C total plate count. The 35°C total plate count showed similar total densities, with 9 of the 14 samples tested falling into the 50,000-500,000 per 100 ml range. The highest densities (300,000-500,000 per 100 ml) occurred in the waters immediately outside the breakwater and in the area extending approximately two miles to the south and one-half mile or less offshore. (See Tables 5 and 6). The distribution of 20°C total bacterial densities in the Chicago Harbor are shown in Figure 16.

In the Racine Harbor the total plate counts at 20°C gave rise to estimations of higher densities than did the 35°C plate count. 93.7% of 32 samples contained total densities in the 100,000-1 million range as estimated by the 20°C total plate count, and 89.3% of the 28 samples tested at 35°C contained densities in the 10,000-500,000 range. Densities of approximately one million (20°C) were encountered at the Root River mouth and throughout the harbor. (Figure 17). The entire one-to-two-mile radial zone around this harbor contained total 20°C densities of approximately 200,000 per 100 ml.





The Milwaukee Harbor showed the highest total bacterial densities of the three harbors herein discussed. Of 75 samples, 32 (42.7%) contained densities in the 50,000 to less than 1 million range as estimated by the 20°C total plate count (Figure 18) and 30 (40.0%) of 75 samples were in the same range as estimated by the 35°C total plate count. Densities of 1,300,000 to 1,500,000 per 100 ml (20°C) were encountered at the mouth of the Milwaukee River and in the central one-third of the Harbor. The corresponding 35°C density at the river's mouth was 2,200,000 per 100 ml. At the extreme northern end of the harbor the total densities were 180,000 and 140,000 (20° and 35°C, respectively). At the southern extremity of the harbor the respective densities were 170,000 and 130,000. In the breakwater zone south of the harbor the 20°C densities progressively diminished from 260,000 per 100 ml to 48,000 (at open water). The 35°C densities diminished from 210,000 to 24,000 throughout the same area. In the open waters adjacent to the northern portion of the harbor much lower densities were often observed (from 10,000 to only a few hundred per 100 ml on both 20° and 35°C tests). The densities at the central harbor mouth were 13,000 and 7,200 (20° and 35°C, respectively). This latter situation may indicate prevailing inflow of Lake Michigan water at the harbor mouth. In the open waters along the breakwater extending south of the harbor, 20°C densities were fairly constant at approximately 30,000 per 100 ml. The 35°C densities in the same area were somewhat lower (28,000 to 9,700). (See Tables 5 and 6).

#### Discussion and Significance of Findings

From the above findings, it is apparent that the bacterial quality of Lake Michigan water is generally good in deep water and is more degraded in varying degrees along the shoreline and in city harbors. The coliform findings indicate the presence of pollution of fecal origin in these shoreline and harbor areas, showing increased densities with respect to sampling stations located in these areas.

At a distance of 1 mile offshore along the western shore from Milwaukee, Wisconsin, to Gary, Indiana, the coliform levels were frequently in the 100-1,000 per 100 ml range, representing substantial pollution in the 1 mile zone. The highest coliform densities were encountered in water contiguous to the towns of Milwaukee, and Racine, Wisconsin; Chicago, Illinois; and Hammond, and Gary, Indiana. The water quality was much less degraded in the northern half of Lake Michigan.



The coliform and fecal streptococcus findings in the Milwaukee Harbor indicate the presence of gross pollution. The same may be said of the Racine Harbor as well as of the waters adjacent to Whiting and Gary, and Hammond, Indiana, although the pollution encountered was not as intense as that of the Milwaukee Harbor.



## OTHER MICROBIOLOGICAL DATA

Coliform Densities at Various Drinking Water Intakes and Bathing Beaches

In addition to the data collected by the Great Lakes-Illinois River Basins Project, daily water sampling results at the various city water intakes as well as routine sampling of bathing beaches, gives much additional information on prevailing quality as well as the occurrence of sudden and localized deteriorating influences. Tables 7 and 8 are included to show the presence of coliform densities occurring at substantially higher levels than encountered in the Great Lakes-Illinois River Basins Project cruises. Additional microbiological aspects pertinent to the evaluation of the present and future quality of the waters of Lake Michigan are to be found in the discussion below.

From the records of the water treatment plants at Chicago and Evanston, Illinois (3, 4), it is indicated that the water of Lake Michigan is of good bacterial quality as long as pollution from domestic sewage inflows is not present. Bacteriological samples collected at the water intakes at the Chicago and Evanston water treatment plants often have shown coliform density of less than 2.2 per 100 ml. At times, however, evidence of pollution of varying degrees is encountered at the water intakes (located one to four miles offshore), as judged by coliform content. (See Figure 19 for location of the Chicago water intakes). Such increase in pollution is reported to be encountered at the Dunne Crib when the prevailing winds are southerly. The pollution accruing from the cities located at the southern tip of the lake, as well as polluted water from the Calumeg-Sag Channel and other points flowing into Lake Michigan during heavy rainfalls, thus is introduced into the water intake of the City of Chicago's South District Filtration Plant (Dunne Crib). This polluted water often bears marked phenolic tastes and odors believed to originate in the industrial and other wastes introduced into the lake along with the domestic sewage cited above.

Table 9, comprised of data from the official records of the City of Chicago, summarizes variations in prevailing water quality in terms of the coliform content encountered at Dunne Crib for the years 1951, 1956, and 1961, as examples typifying conditions in past records. The total number of days when the coliform density (MPN per 100 ml) was 2.2 or less per 100 ml were: 1951, 186; 1956, 90; and 1961, 174. Days when the coliform density was 50 or greater were as follows: 1951, 11 days; 1956, 48 days; and 1961, 54 days. The maximum coliform densities encountered in 1951, 1956, and 1961 were 374, 1100, and 3000 per 100 ml, respectively.



The maximum coliform density reported at the Evanston water treatment plant for 1956 was 23,000. This density was encountered on the following dates: January 10, 12, 13, 17; September--1 day; November 24, 25, 29; and December 15, 21, and 25 (4).

Inspection of coliform levels at these two locations, separated by 20 miles, but both located on the west shore of Lake Michigan, near heavily populated metropolitan areas, reveals great variation and lack of homogeneity in the distribution of coliform bacteria present in Lake Michigan water, particularly when a density of 23,000 per 100 ml prevailed for more than one day consecutively at Evanston and nothing comparable was detected simultaneously at the Dunne Crib.

Earlier records show that heavily contaminated water would reach and travel past Dunne Crib as in the years 1940, 1941, 1942, when isolated coliform levels of 300,000 were detected. On October 19, 1940, a maximum of 300,000 coliform per 100 ml was detected in the Dunne Crib intake, while an adjacent water intake located only 300 feet closer to shore and sampled within 15 minutes, showed a level of 2400 coliform per 100 ml (3). Pollution in this order of magnitude was known to originate in cities located at the southern tip of Lake Michigan which contributed large amounts of domestic and industrial wastes to Lake Michigan. Improvements in these local conditions resulted in sharply reduced maximum coliform levels reaching the Dunne Crib in subsequent years. It was this degraded water quality in the southern end of Lake Michigan that determined the need for and location of the City of Chicago's first water filtration plant (the South District Filtration Plant, put into service in 1946) (6, p.15).

Table 7 presents data revealing the coliform densities that prevailed during the months of June-November, 1961, inclusive, at three locations adjacent to the north side of the City of Chicago's Central District Filtration Plant, currently under construction (3). It will be noted that the levels of coliform at all points exceeded the recommended maximum of 50 to 100 coliform bacteria per 100 ml for source water to be used for public water supplies where chlorination is the only treatment provided (7, p.11). The maximum MPN per 100 ml reported from these points was 24,000.

The range of commonly accepted standards established by States for water to be used for swimming and other recreational uses is 50 to 2400 coliform bacteria per 100 ml (14). Table 8 presents the coliform levels occurring at each of the beaches located on the Chicago lakefront during the summer of 1961 and is based on samples collected during the months of May through September at each sampling point (5). The southern-most beaches show days of gross pollution





with Most Probable Numbers (MPN) greater than 110,000. Three points on the Calumet Beach showed MPN's per 100 ml in excess of 1,000 in 70% of the samples collected. Beaches at Hammond and Whiting, Indiana showed 72% of samples in excess of 1,000. The range for the other 24 beaches was 19-60% wherein MPN's per 100 ml were detected in excess of 1,000. The highest MPN per 100 ml detected was 240,000 occurring at 12th Street Beach on September 15, 1961 in the wake of the polluted water backflowing through the locks into Lake Michigan (see discussion below).

#### Pollution Persistency

The material and data presented above demonstrate the lack of uniformity in the quality of Lake Michigan water, wherein local areas of the lake show excessive pollution, as measured by the coliform MPN, and other areas, either adjacent or more removed, may show quite low coliform content. Polluted slugs of water entering the lake are not immediately dispersed and tend to maintain their identity for days. The latter condition has been demonstrated when the locks on the Chicago River have been opened to alleviate flood conditions in the Upper Illinois Waterway.

The locks at the Chicago River were opened to release flood waters on September 14, 1961. At the same time flood water was released from the Upper Illinois Waterway through the locks at the upper end of the North Shore Channel at Wilmette, Illinois. At the southern end of Lake Michigan, massive amounts of flood water entered the lake from the Calumet River when the direction of flow in the river was reversed in response to the heavy rainfall. A survey to determine the location, migration, and persistency of the flood water introduced into Lake Michigan was undertaken by the Great Lakes-Illinois River Basins Project (8). The discharge from the Chicago River was more intensively studied than the other two massive discharges. Testing and identification of the polluted water was maintained in terms of coliform and fecal streptococcus levels (membrane filter determinations). Coliform densities per 100 ml in the magnitude of 180,000-150,000 were encountered 1 3/8 and 7/8 miles, respectively, offshore opposite the mouth of the Chicago River, the first day, with slight diminution on the second day, and a maximum of 11,000 appearing on the third day. On the third day a density of 320,000 was encountered to the north near Wilson Avenue Crib (water intake). This body of polluted water may have originated from the flood water released concurrently on September 14, 1961, from the North Shore Channel at Wilmette.

1. The first part of the document discusses the importance of maintaining accurate records of all transactions and activities. It emphasizes that this is essential for ensuring transparency and accountability in the organization's operations.

2. The second part of the document outlines the various methods and tools used to collect and analyze data. It highlights the need for a systematic approach to data collection and the importance of using reliable sources of information.

3. The third part of the document describes the process of identifying and addressing potential risks and challenges. It stresses the importance of proactive risk management and the need to develop effective strategies to mitigate potential threats.

4. The fourth part of the document discusses the role of communication and collaboration in achieving the organization's goals. It emphasizes the importance of clear communication and the need for all team members to work together effectively.

5. The fifth part of the document outlines the various metrics and indicators used to measure the organization's performance. It highlights the need for a balanced scorecard approach that takes into account both financial and non-financial factors.

6. The sixth part of the document describes the process of reviewing and evaluating the organization's progress. It stresses the importance of regular reviews and the need to make adjustments as needed to ensure that the organization is on track to achieve its goals.

7. The seventh part of the document discusses the importance of continuous improvement and the need to seek out new opportunities for growth and innovation. It emphasizes the importance of a culture of learning and the need to embrace change.

8. The eighth part of the document outlines the various challenges and obstacles that the organization may face. It highlights the need for a strong leadership team and the importance of having a clear vision and mission statement.

9. The ninth part of the document discusses the importance of maintaining a strong relationship with stakeholders and the need to communicate effectively with all parties involved. It emphasizes the importance of transparency and the need to be open to feedback.

10. The tenth part of the document outlines the various steps and actions that the organization needs to take to achieve its goals. It stresses the importance of a clear plan and the need to monitor progress closely.

The bacteriological findings of this investigation indicated that the polluted flood water tended to maintain its identity for at least 3 days and was kept fairly close to its point of introduction (around the mouth of the Chicago River) by the prevailing winds. A tongue of polluted water appeared to extend some 4 miles in a southerly direction from the point of heaviest concentration of pollution.

The normal quality of Lake Michigan water in the Chicago area was apparently influenced by the heavy rainfalls and subsequent run-off which occurred during September, 1961. The total precipitation recorded at Midway Airport (14.17") was the heaviest for any month of record at this station (1928-1961). The Chicago Avenue and Wilson Avenue Control Stations of the Chicago water department as well as Dunne Crib, reported sustained periods of high chlorine demand occurring throughout the remainder of September, October, and November. Table 10 summarizes the change in coliform content of water taken in at Dunne Crib throughout the year of 1961, with the exception of December. It is apparent that higher levels of coliform bacteria were encountered in September, October, and November at greater frequency than at any other period of the year.



## ULTIMATE SURVIVAL OF PATHOGENS IN LAKE MICHIGAN

The question of ultimate survival of bacteria, including pathogens, artificially introduced into Lake Michigan (through sewage effluents, flood water overflow from polluted sewers, etc.) is one of great importance and is of particular significance in relation to public health and safety. Pathogenic forms such as typhoid organisms have been known to survive winter conditions in northern locations, travel downstream following thaws and subsequently precipitate epidemics among water users (9). Pathogenic enteroviruses have been shown to be much more resistant to chlorination than are vegetative bacterial cells, and may be presumed to be persistent in nature until proven otherwise (10, 16, 17). Other enteric parasites (such as Endamoeba histolytica and nematode eggs) are known to survive outside the human intestinal tract and to be more resistant to chlorination than the sewage indicator organisms, the coliform group (10).

Little information is available on the survival of any of these forms, including the coliform bacteria in Lake Michigan. The very nature of the problem and the peculiarities of microbial survival and reproduction render field investigations extremely difficult since no known method is available on a practical basis to follow a given inoculum of bacteria through the many depths and currents of a body of water like Lake Michigan. Moreover, the introduction of pathogenic forms into a body of water used as a source for public drinking water for study purposes is not tenable. Laboratory studies cannot easily duplicate the conditions found in nature. Nonetheless, such studies do provide useful information in considerations related to survival of enteric microorganisms in nature. While many laboratory studies relating to the survival of pathogenic forms in either laboratory or natural conditions can be cited, one survival study on Shigella sonnei seems particularly noteworthy. In this study S. sonnei remained viable when stored in tap water for 211 days (11). In another investigation Salmonella typhosa was found to survive in impounded surface water up to 26 days (12).

A survival study using Lake Michigan water was conducted in the laboratories of the Great Lakes-Illinois River Basins Project wherein the survival of coliform and fecal streptococci was investigated. In this experiment, Lake Michigan water was collected 20 miles offshore. Initial coliform measurements on the water as collected were less than one coliform cell per 100 ml, with no fecal streptococci detected. This water was seeded with untreated sewage collected from the West-Southwest Treatment works and transferred to sterile gallon jugs. These containers were stored at the following temperatures and conditions:

1. The first part of the document discusses the importance of maintaining accurate records of all transactions and the role of the accounting department in ensuring the integrity of the financial statements. It also highlights the need for regular audits and the importance of transparency in financial reporting.

2. The second part of the document focuses on the implementation of internal controls to prevent fraud and ensure the accuracy of financial data. It outlines the key components of a robust internal control system, including segregation of duties, authorization procedures, and regular monitoring and evaluation.

3. The third part of the document addresses the challenges faced by organizations in managing their financial resources effectively. It discusses the importance of budgeting, forecasting, and cost management, and provides practical advice on how to overcome common financial management challenges.

4. The fourth part of the document explores the role of technology in modern accounting and finance. It discusses the benefits of using accounting software and the importance of staying up-to-date with the latest technological advancements in the field.

5. The fifth part of the document concludes by emphasizing the importance of continuous learning and professional development for accounting and finance professionals. It encourages individuals to stay current in their knowledge and skills to meet the evolving demands of the industry.

- (1) 25°C with room illumination during working hours
- (2) 25°C in the dark, and
- (3) 5°C in the dark.

Under each of these test conditions, the fecal streptococci showed less survival capacity than did the coliform bacteria. Initial streptococcus levels were in the ranges of 680,000 to 1,000,000 cells/100 ml. At 25°C with light, no streptococci were detected at the end of 5 days. In the dark at 25°C, 2 per 100 ml were measured at the end of 15 days, and at 5°C in the dark, 340 per 100 ml were viable at the end of 15 days. Determinations for both the streptococci and coliform were via direct colony count using membrane filter techniques.

Initial levels of coliform were approximately 500,000 in each of the test containers. At the end of 15 days the following levels were detected as still viable: (1) 25°C with light, 3400 per 100 ml; (2) 25°C dark, 7400 per 100 ml, and (3) 5°C dark, 11,000 per 100 ml. Under the conditions of (2) the coliform bacteria underwent an initial increase of 30% at the end of the first day and under the conditions of (3) a 50% increase by the end of the fifth day. Following these peaks, the numbers decreased without interruption through the 15-day test period. These results are set forth in Figures 20, 21, and 22. While this study was a small investigation, and exploratory in nature, its immediate significance lies in the fact that bacteria from Chicago sewage wastes were combined with Lake Michigan water on a test basis. The results wherein coliform survived at 5°C through 15 days (and would, no doubt, have been detected as viable for a more extended period had the test not been terminated at the end of 15 days) are of particular importance if the same or greater survival rate should prevail from sewage bacteria introduced into Lake Michigan. It has been observed in some areas (13) that residual bacteria remaining viable in partially chlorinated sewage treatment plant effluent may multiply in great numbers in the waters receiving the effluent. This rapid multiplication occurs since competitors for the food supply (other bacteria and other microbial forms) and predators are killed off, leaving the residual bacteria surviving uninhibited in an environment rich in the nutrients upon which they thrive. Currents and wave distribution could, under certain conditions, carry such contamination to the vicinity of water intakes and bathing beaches within a few days at most.

#### Possible Effect of Returning Treated Effluent to Lake Michigan

The possible effect of returning sewage effluent produced by the MSD plants to Lake Michigan is complex, with manifold ramifications. The factors relating to the fate or survival of pathogens introduced into the lake are largely unknown. Distribution and survival of living pathogens, both surface-wise and depth-wise, cannot, at the present time, be intelligently discussed since little information is available relating to both the functioning of the biological entities in question and the physical and limnological characteristics of Lake Michigan.

1. The first part of the document discusses the importance of maintaining accurate records of all transactions and activities. It emphasizes that proper record-keeping is essential for transparency and accountability, particularly in financial matters. The text suggests that organizations should implement robust systems to track every aspect of their operations, from procurement to sales.

2. The second part of the document addresses the challenges of managing a large and diverse workforce. It highlights the need for effective communication and collaboration across different departments and regions. The author suggests that regular meetings and open lines of communication are crucial for ensuring that everyone is on the same page and working towards common goals.

3. The third part of the document focuses on the importance of continuous learning and development. It argues that in a rapidly changing business environment, employees must be equipped with the latest skills and knowledge. The text recommends investing in training programs and encouraging a culture of lifelong learning.

4. The fourth part of the document discusses the role of technology in modern business operations. It notes that while technology offers many advantages, it also presents challenges, such as data security and integration. The author suggests that organizations should carefully evaluate their technology needs and implement solutions that enhance efficiency without compromising security.

5. The fifth part of the document touches upon the importance of maintaining a strong corporate culture. It suggests that a clear set of values and a strong sense of purpose can help guide decision-making and foster a positive work environment. The text encourages leaders to model the behaviors they expect from their employees.

6. The sixth part of the document discusses the importance of financial management. It suggests that organizations should regularly review their financial statements and budgets to ensure they are on track. The text also mentions the importance of maintaining good relationships with creditors and investors.

7. The seventh part of the document discusses the importance of legal compliance. It suggests that organizations should stay up-to-date on relevant laws and regulations to avoid potential legal issues. The text also mentions the importance of having a clear policy on data privacy and protection.

8. The eighth part of the document discusses the importance of customer satisfaction. It suggests that organizations should regularly gather feedback from their customers and use it to improve their products and services. The text also mentions the importance of providing excellent customer service.

9. The ninth part of the document discusses the importance of environmental sustainability. It suggests that organizations should consider the environmental impact of their operations and take steps to reduce their carbon footprint. The text also mentions the importance of using sustainable materials and practices.

10. The tenth part of the document discusses the importance of social responsibility. It suggests that organizations should consider the social impact of their operations and take steps to support the communities in which they operate. The text also mentions the importance of being transparent about their social and environmental performance.



It can be proposed, however, on the basis of available information that the addition of sewage effluent from the MSD plants would contribute to the deterioration of the microbiological quality of Lake Michigan water. It has been observed that under certain conditions polluted water may maintain its identity and move in discrete masses within the lake. It is reasonable to assume that drinking water intakes and bathing beaches would, at times of varying frequency, be influenced by water of objectionable quality. On many days in the bathing season of 1961 the present water quality at several of the Chicago beaches had very high coliform levels. Any addition to the prevailing levels of pollution could be expected to seriously menace this use of Lake Michigan. The same observations would apply to the lake as a source of public water supplies.

1. The first part of the document is a letter from the author to the reader, explaining the purpose of the study and the methods used. The author states that the study is a qualitative investigation into the experiences of young people who have been involved in the criminal justice system. The methods used are semi-structured interviews and focus groups. The author also mentions that the study is part of a larger project funded by the Home Office.

2. The second part of the document is a literature review, which discusses the existing research on young people and the criminal justice system. The author identifies a gap in the literature, which is the lack of research on the experiences of young people who have been involved in the criminal justice system. The author also discusses the importance of understanding the experiences of young people in order to develop effective interventions.

3. The third part of the document is the methodology, which describes the research design and the data collection methods. The author explains that the study is a qualitative investigation and that the data collection methods are semi-structured interviews and focus groups. The author also discusses the ethical considerations of the study and the steps taken to ensure that the study is conducted in a ethical and responsible manner.

4. The fourth part of the document is the results, which presents the findings of the study. The author discusses the experiences of young people who have been involved in the criminal justice system and the factors that influence their experiences. The author also discusses the implications of the findings for practice and policy.

5. The fifth part of the document is the conclusion, which summarizes the findings of the study and the implications for practice and policy. The author states that the study has identified a gap in the literature and that the findings have implications for practice and policy. The author also states that the study has contributed to the understanding of the experiences of young people who have been involved in the criminal justice system.

## CONCLUSIONS

1. The coliform concentrations in certain areas of Lake Michigan water at times exceed the level regarded as the safe limit for water used as a source of public water supply where chlorination is the only treatment provided.
2. The most recent records of the quality of Lake Michigan water at points up to two and one-half miles offshore in the vicinity of Chicago bathing beaches show that coliform concentrations are often in excess of acceptable limits for swimming and other recreational uses.
3. Coliform densities are so high in city harbors and adjacent areas studied, that a grave threat to health must be presumed to exist through contact with these waters.
4. The fecal streptococcus levels, as well as the coliform bacteria, indicate gross pollution in the harbors studied.
5. Lowest total bacterial densities were observed in the central body of lake water with increasing densities occurring in relation to the placement of cities and towns. Increasing total bacterial densities usually indicate deterioration in the sanitary quality of water through the presence of pollutants contributing to the enrichment of the water.
6. Masses of bacterially-polluted water introduced into Lake Michigan may maintain their identity for several days.
7. It is known that microbial forms pathogenic to man survive in natural bodies of water for varying periods; some forms may be able to survive longer than the indicator organisms commonly used, i.e., the coliform bacteria.
8. The return to Lake Michigan of sizable quantities of treated waste waters, now being discharged to the Illinois River, would increase the densities of coliform bacteria and other microbial forms in the vicinity of the outlet. This would increase the potential hazards to public health in the use of waters in the area, and heighten the need for protective measures - including, but not limited to: complete treatment of waste waters before discharge to the Lake and complete treatment of municipal supply taken from the Lake.

# THEORY OF THE EARTH

The theory of the earth is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the processes which have shaped the earth and its various parts.

The theory of the earth is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the processes which have shaped the earth and its various parts.

The theory of the earth is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the processes which have shaped the earth and its various parts.

The theory of the earth is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the processes which have shaped the earth and its various parts.

The theory of the earth is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the processes which have shaped the earth and its various parts.

The theory of the earth is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the processes which have shaped the earth and its various parts.

The theory of the earth is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the processes which have shaped the earth and its various parts.

The theory of the earth is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the processes which have shaped the earth and its various parts.

## REFERENCES

1. Standard Methods for the Examination of Water and Wastewater. Eleventh Edition. American Public Health Association, Inc. New York, N.Y. (1960).
2. Recent Developments in Water Bacteriology. U.S. Department of Health, Education, and Welfare; Public Health Service; Division of Water Supply and Pollution Control; Robert A. Taft Sanitary Engineering Center, Cincinnati, Ohio. (1961).
3. City of Chicago Water Department. Unpublished Water Quality Data Records.
4. City of Evanston Water Department. Unpublished Water Quality Data Records.
5. Chicago Park District. Unpublished Water Quality Data Records.
6. U.S. Public Health Service. The Chicago Cook County Health Survey. Columbia University Press. New York. (1949).
7. U.S. Public Health Service. Manual of Recommended Water Sanitation Practice. Public Health Service Publication No. 525. (1958).
8. U.S. Public Health Service. Movements in Lake Michigan of Water Discharged 14 September 1961 from The Chicago Sanitary Canal System, U.S. Exhibit No. 4, Chicago Diversion Case, Mimeo, Unpublished (1961).
9. Healy, W.A. and Grossman, R.P. Water-Borne Typhoid Fever Epidemic at Keene, New Hampshire. J. New England Waterworks Association, 75:38 (1961).
10. Kabler, P.W.; Clark, H.F.; and Clarke, N.A. Pathogenic Microorganisms and Water-Borne Disease. Proceedings Rudolfs Research Conference: Public Health Hazards of Microbial Pollution of Water. Dept. of Sanitation, College of Agriculture, Rutgers University. (1961). pp. 9-56.
11. Kraus, P., and Weber, G. Investigations on the Viability of Pathogenic Organisms in Water Supplies and Surface Water. Zbl. Bakt. I, 171:509 (1958).
12. Shrewsbury, J.F.D., and Barson, G.J. A Note on the Absolute Viability in Water of *S. Typhi* and the Dysentery Bacilli. Brit. Med. Jour., 1:954 (1952).

1. The first part of the paper is devoted to the study of the properties of the function  $f(x)$  defined by the equation

$$f(x) = \int_0^x \frac{1}{1+t^2} dt, \quad (1)$$

where  $x$  is a real number. It is well known that the function  $f(x)$  is increasing and concave down.

2. In the second part, we shall study the properties of the function  $g(x)$  defined by the equation

$$g(x) = \int_0^x \frac{t}{1+t^2} dt, \quad (2)$$

where  $x$  is a real number. It is well known that the function  $g(x)$  is increasing and concave up.

3. In the third part, we shall study the properties of the function  $h(x)$  defined by the equation

$$h(x) = \int_0^x \frac{t^2}{1+t^2} dt, \quad (3)$$

where  $x$  is a real number. It is well known that the function  $h(x)$  is increasing and concave down.

4. In the fourth part, we shall study the properties of the function  $k(x)$  defined by the equation

$$k(x) = \int_0^x \frac{t^3}{1+t^2} dt, \quad (4)$$

## REFERENCES (cont.)

13. Rudolfs, W., and Gehm, H.W. Multiplication of Total Bacteria and B. coli after Sewage Chlorination. Sewage Works Journal 7:991 (1935).
14. State Water Pollution Control Board. Water Quality Criteria. Sacramento, California. 1957.
15. Prescott, S.C.; Winslow, C.A.; and McCrady, M. Water Bacteriology. John Wiley and Sons, Inc. New York. (1950).
16. Rhodes, A.J.; Clark, E.M.; Knowles, D.S. and Goodfellow, A.M. Prolonged Survival of Human Poliomyelitis Virus in Experimentally Infected River Water. Canad. Jour. Pub. Health, 41:146 (1950).
17. Gilcreas, F.W. and Kelly, S.M. Relationship of Coliform-Organism Test to Enteric Virus Pollution. Jour. Amer. Wat. Wks. Assn. 47:683 (1955).

1. The first part of the document discusses the importance of maintaining accurate records of all transactions and the role of the accounting department in ensuring the integrity of the financial statements. It also highlights the need for transparency and accountability in the reporting process.

2. The second part of the document outlines the various methods used to collect and analyze data, including surveys, interviews, and focus groups. It emphasizes the importance of using a mix of qualitative and quantitative techniques to gain a comprehensive understanding of the research topic.

3. The third part of the document presents the results of the study, which show a significant positive correlation between the variables being investigated. The findings suggest that the proposed intervention could have a beneficial impact on the outcome being measured.

4. The final part of the document discusses the implications of the study and provides recommendations for future research. It suggests that further exploration is needed to understand the underlying mechanisms and to test the findings in a larger, more diverse sample.



TABLE 1

## COLIFORM RESULTS FROM DEEPPWATER AND INSHORE STUDIES

No. per 100 ml	Surface Samples			Lowermost Samples		
	No. of Samples	Per Cent	Cumulative Totals	No. of Samples	Per Cent	Cumulative Totals
< 1	134	42.8	134	156	56.3	156
1-10	99	31.8	233	75	27.1	231
11-50	47	15.0	280	28	10.1	259
51-100	12	3.8	292	4	1.4	263
110-200	8	2.5	300	7	2.5	270
210-300	4	1.3	304	2	0.7	272
310-400	1	0.3	305	0	0	-
410-500	1	0.3	306	1	0.4	273
510-600	2	0.6	308	3	1.1	276
610-700	0	0	-	0	0	-
710-800	1	0.3	309	0	0	-
810-900	0	0	-	0	0	-
910-1,000	0	0	-	0	0	-
1,100-2,000	4	1.3	313	0	0	-
2,000-2,500	0	0	-	1	0.4	277
Total	313	100.00	-	277	100.0	-

100.0



TABLE 2  
20°C TOTAL PLATE COUNT RESULTS FROM DEEPWATER AND INSHORE STUDIES

No. per 100 ml	Surface Samples			Lowermost Samples		
	No. of Samples	Per Cent	Cumulative Totals	Per Cent	No. of Samples	Cumulative Totals
100-490	17	6.3	17	6.3	13	13
500-990	18	6.6	35	12.9	17	30
1,000-4,900	109	40.2	144	53.1	85	115
5,000-9,900	48	17.7	192	70.8	59	174
10,000-49,000	46	16.9	238	87.7	25	199
50,000-99,000	12	4.5	250	92.2	15	214
100,000-490,000	21	7.8	271	100.0	21	235
500,000-990,000	0	0	-	100.0	2	237
Total	271	100.0			237	100.0



TABLE 3

## 35°C TOTAL PLATE COUNT RESULTS FROM DEEPWATER AND INSHORE STUDIES

No. per 100 ml	Surface Samples			Lowermost Samples		
	No. of Samples	Per Cent	Cumulative Totals	Per Cent	No. of Samples	Cumulative Totals
10-90	8	2.6	8	2.6	0	0
100-490	37	11.9	45	14.5	26	26
500-990	34	10.9	79	25.4	19	45
1,000-4,900	99	31.8	178	57.2	82	127
5,000-9,000	53	17.0	231	74.2	32	159
10,000-49,000	37	11.9	268	86.1	21	180
50,000-99,000	19	6.1	287	92.2	13	193
100,000-490,000	21	6.9	308	99.1	16	209
500,000-990,000	1	0.3	309	99.4	2	211
1 M+	2	0.6	311	100.0	1	212
Total	311	100.0			212	100.0

1. The first part of the paper is devoted to a discussion of the general principles of the theory of the structure of the atom. It is shown that the structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the uncertainty of the position and momentum of the particles.

2. In the second part of the paper, the author discusses the problem of the structure of the nucleus. It is shown that the structure of the nucleus is determined by the laws of quantum mechanics, which are based on the principle of the uncertainty of the position and momentum of the particles.

3. In the third part of the paper, the author discusses the problem of the structure of the molecule. It is shown that the structure of the molecule is determined by the laws of quantum mechanics, which are based on the principle of the uncertainty of the position and momentum of the particles.

4. In the fourth part of the paper, the author discusses the problem of the structure of the crystal. It is shown that the structure of the crystal is determined by the laws of quantum mechanics, which are based on the principle of the uncertainty of the position and momentum of the particles.

5. In the fifth part of the paper, the author discusses the problem of the structure of the solid. It is shown that the structure of the solid is determined by the laws of quantum mechanics, which are based on the principle of the uncertainty of the position and momentum of the particles.

TABLE 4  
COLIFORM RESULTS FROM HARBOR STUDIES (SURFACE SAMPLES ONLY)

No. per 100 ml	No. Stations Sampled	No. Samples	Per Cent	Cumulative Totals	Cumulative Per Cent
<u>Chicago Harbor</u>					
1-10		3	20.0	3	20.0
11-50		4	26.7	7	46.7
100-490		6	40.0	13	86.7
500-990		2	13.3	15	100.0
Total	15	15	100.0		
<u>Racine Harbor</u>					
10-50		3	8.8	3	8.8
50-100		2	5.9	5	14.7
100-490		11	32.4	16	47.1
500-990		5	14.7	21	61.8
1000-5000		12	35.3	33	97.1
5000-10,000		1	2.9	34	100.0
Total	34	34	100.0		
<u>Milwaukee Harbor</u>					
1		9	11.8	9	11.8
1-10		5	6.5	14	18.3
10-50		8	10.5	22	28.8
50-100		4	5.3	26	34.1
100-490		18	23.7	44	57.8
500-990		3	3.9	47	61.7
1000-5000		10	13.2	57	74.9
5000-10,000		7	9.2	64	84.1
10,000-20,000		4	5.3	68	89.4
20,000-30,000		4	5.3	72	94.7
30,000-40,000		4	5.3	76	100.0
Total	76	76	100.0		

1875

1875

1875

1875

1875

1875

1875

1875

1875

1875



TABLE 5

## 20°C TOTAL PLATE COUNT RESULTS FROM HARBOR STUDIES (SURFACE SAMPLES ONLY)

No. per 100 ml	No. Stations Sampled	No. Samples	Per Cent	Cumulative Totals	Cumulative Per Cent
<u>Chicago Harbor</u>					
500-990		1	5.8	1	5.8
1000-4900		4	23.5	5	29.3
5000-9900		3	17.8	8	47.1
50,000-99,000		1	5.8	9	52.9
100,000-490,000		8	47.1	17	100.0
Total	17	17	100.0		
<u>Racine Harbor</u>					
5000-9900		1	3.1	1	3.1
10,000-49,000		1	3.1	2	6.2
100,000-490,000		23	71.9	25	78.1
500,000-990,000		5	15.6	30	93.7
1 million $\neq$		2	6.3	32	100.0
Total	32	32	100.0		
<u>Milwaukee Harbor</u>					
500-990		1	1.3	1	1.3
1000-4900		9	12.0	10	13.3
5000-9900		12	16.0	22	29.3
10,000-49,000		21	28.0	43	57.3
50,000-99,000		8	10.7	51	68.0
100,000-490,000		14	18.7	65	86.7
500,000-1 million		4	5.3	69	92.0
1 million $\neq$		6	8.0	75	100.0
Total	75	75	100.0		



TABLE 6  
35°C TOTAL PLATE COUNT RESULTS FROM HARBOR STUDIES (SURFACE SAMPLES ONLY)

No. per 100 ml	No. Stations Sampled	No. Samples	Per Cent	Cumulative Totals	Cumulative Per Cent
		<u>Chicago Harbor</u>			
500-990		2	14.3	2	14.3
1,000-4,900		1	7.1	3	21.4
10,000-49,000		2	14.3	5	35.7
50,000-99,000		5	35.7	10	71.4
100,000-490,000		3	21.5	13	92.9
500,000		1	7.1	14	100.0
Total	14	14	100.0		
		<u>Racine Harbor</u>			
1,000-4,900		2	7.1	2	7.1
5,000-9,900		1	3.6	3	10.7
10,000-49,000		9	32.1	12	42.8
50,000-99,000		4	14.3	16	57.1
100,000-490,000		12	42.9	28	100.0
Total	28	28	100.0		
		<u>Milwaukee Harbor</u>			
100-490		4	5.3	4	5.3
500-990		3	4.0	7	9.3
1,000-4,900		15	20.0	22	29.3
5,000-9,900		8	10.7	30	40.0
10,000-49,000		15	20.0	45	60.0
50,000-99,000		5	6.6	50	66.6
100,000-490,000		17	22.7	67	89.3
1 million		8	10.7	75	100.0
Total	75	75	100.0		



TABLE 7  
COLIFORM LEVELS AT THREE POINTS  
ADJACENT TO CENTRAL DISTRICT FILTRATION PLANT, CHICAGO, ILLINOIS, 1961\*  
MPN per 100/ml

	Distance from Shore	June	July	August	September	October	November
East End	App. 1 mile	Max. 2,400 Av. 738	24,000 4,114	780 198	9,200** 1,629	790 400	1,300 285
Center	App. $\frac{1}{2}$ mile	Max. 5,400 Av. 1,500	13,000 2,486	780 165	490 187	1,300 339	16,000 2,104
West End	Near Shore	Max. 3,300 Av. 1,273	4,900 1,613	790 235	220 138	1,100 393	2,400 487

\* Data Supplied by: Department of Water and Sewers, City of Chicago

\*\* September 14, 1961

TABLE 8

## COLIFORM INCIDENCE AT THE CHICAGO BEACHES, SUMMER, 1961\*

BEACH	SAMPLING POINT	COLIFORM MPN/100 ml		
		MAX	No. days at Max.	No. days ≥ 1000
1. Juneway Terrace	301	1,500	1	5
2. Rogers Ave.	302	12,000	1	10
3. Howard Street	303	11,000	1	9
4. Sherwin Ave.	304	110,000	2	8
5. Rogers Park	305	110,000	1	8
6. Farwell Beach	306	46,000	1	9
7. Pratt Blvd.	307	4,600	1	6
8. Columbia Ave.	308	11,000	1	5
9. North Shore Ave.	309	15,000	1	5
10. Albion Ave.	310	11,000	1	6
11. Granville Ave.	311	7,500	1	10
12. Thorndale Ave.	312	21,000	1	7
13. Hollywood Ave.	313	7,500	1	9
14. Foster Ave.	314	24,000**	1	11
15. Montrose Beach	315	24,000**	1	16
16. North Ave. (North)	316	15,000	3	8
17. North Ave. (South)	317	110,000	1	10
18. Oak Street	318	15,000	1	12
19. 12th Street	319	240,000**	1	17
20. 31st Street	320	46,000	1	10
21. 49th Street	321	1,100	1	5
22. 57th Street	322	4,600	2	12
23. Jackson Park	323	46,000	1	12
24. 67th Street	324	4,600	1	9
25. Rainbow (North)	325	110,000	1	10
26. Rainbow (South)	326	11,000	1	10
27. Calumet (Outer)	327	46,000	1	15
28. Calumet (North)	328	110,000	1	18
29. Calumet (South)	329	110,000	1	22
30. Hammond	330	110,000	2	20
31. Whiting	331	110,000	2	19

\*\*9-15-61

\*Data supplied by the Park District, City of Chicago



TABLE 9

COLIFORM BACTERIA Per 100 ml at DUNNE CRIB in 1951, 1956, and 1961 (Standard MPN)\*

	1951		1956		1961	
	< 2.2	> 50	< 2.2	> 50	< 2.2	> 50
No. of days	186	11	90	48	174	54
Highest Daily maximum		374		1100		3000

\* Data supplied by: Dept. of Water and Sewers, City of Chicago



TABLE 10

Summary of Coliform Levels at Dunne Crib  
for 1961 (MPN/100 ml)

	Number of Samples Having an MPN of		Maximum MPN During Month
	< 2.2	> 50	
JANUARY	24	0	10.0
FEBRUARY	23	0	6.9
MARCH	10	3	120.0
APRIL	19	3	88.0
MAY	16	8	510.0
JUNE	18	0	33.0
JULY	5	1	120.0
AUGUST	13	0	44.0
SEPTEMBER	2	9	450.0
OCTOBER	1	18	1800.0
NOVEMBER	0	14	3000.0
DECEMBER			

Data Supplied by: Department of Water and Sewers, City of  
Chicago

