

EPA/625/3-91/019A
SAB/SAP Review Draft
February 1991

ALPHA-2U-GLOBULIN: ASSOCIATION WITH CHEMICALLY-INDUCED RENAL TOXICITY AND NEOPLASIA IN THE MALE RAT

Prepared for the
Risk Assessment Forum
U.S. Environmental Protection Agency
Washington, DC

Authors

Karl Baetcke
Imogene Sevin Rodgers
Letitia Tahan

Gordon Hard
Robert McGaughy

Technical Panel

Karl P. Baetcke, Ph.D.*
Letitia Tahan, M.S.*
Marion Copley, D.V.M.

Julie Du, Ph.D.
Robert McGaughy, Ph.D.
William PePelko, Ph.D.

*Technical Panel Co-Chairman

Project Officer: Imogene Sevin Rodgers, Ph.D.

Consultant: Gordon C. Hard, B.V.Sc., Ph.D., D.Sc., F.R.C. Path.,
F.R.C.V.S., F.A. Tox. Sci.

U.S. ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

DRAFT -- DO NOT QUOTE OR CITE

This document is a draft for review purposes only and does not constitute Agency policy. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

TABLE OF CONTENTS

PREFACE	vii
GLOSSARY	viii
EXTERNAL PEER REVIEWERS	ix
EXECUTIVE SUMMARY	1
 I. Introduction	 9
 PART 1 - NEPHROTOXICITY	 17
II. Hyaline droplets and alpha-2u-globulin; physiology and biochemistry	17
A. Filtration, reabsorption, and catabolism of low-molecular-weight proteins by the kidney	17
B. Hyaline droplets in renal tubules	20
C. Factors affecting kidney accumulation of low-molecular-weight proteins	23
D. The alpha-2u-globulin superfamily of proteins	25
E. Characteristics of alpha-2u-globulin	29
F. Sex and species comparison of urinary protein content of the lipocalin superfamily	31
G. Noncovalent binding to alpha-2u-globulin and its homologues	33
H. Catabolism of alpha-2u-globulin complexed with CIGA	38
I. Structure-activity relationships for CIGA	41
 III. Alpha-2u-globulin nephropathy	 43
A. Pathologic features of alpha-2u-globulin nephropathy	44
B. Rat urine chemistry and CIGA	48

C. Species variation in the renal response to CIGA	50
D. Factors affecting the expression of alpha-2u-globulin nephropathy	53
E. Chronic progressive nephropathy	56
F. Renal toxicity observed in chronic bioassays of chemicals that induced kidney tumors in rats	59
 PART 2 - CARCINOGENICITY	 66
IV. Pathologic features of renal carcinogenesis induced by classical carcinogens	66
A. Early nephrotoxicity	67
B. Karyomegaly	68
C. Tubule cell hyperplasia	68
D. Adenoma	69
E. Adenocarcinomas and carcinomas	70
F. Tumor progression	70
G. Site of origin of renal tubule tumors	71
V. Neoplastic and pre-neoplastic lesions observed in the 2-year bioassays	71
A. Generic considerations	80
B. Renal tumor incidence	82
C. Histogenesis of renal tumors	83
D. Renal tumor latency and progression	84
E. Induction of other tumor types	84
VI. Additional evidence concerning the renal carcinogenicity of CIGA	86
A. Genetic toxicology studies	86
B. Initiation-promotion data	91

VII.	Comparison of CIGA with classical renal carcinogens	94
VIII.	Evidence concerning human kidney cancer	97
	A. Morphology and histogenesis	97
	B. Incidence and mortality	98
	C. Environmental and lifestyle factors	101
	D. Occupational factors	102
	E. Renal cancer and hydrocarbon, solvent or petroleum product exposure	103
IX.	Evidence for dose- and time-dependent progression from early to late lesions	107
	A. Association between CIGA, hyaline droplet formation and alpha-2u-globulin accumulation	107
	B. Association between hyaline droplet formation, cell necrosis, and tubule cell regeneration	109
	C. Progression to cast formation, tubule dilation and mineralization	115
	D. Association between CIGA and chronic progressive nephropathy	119
	E. Evidence concerning progression from nephrotoxicity to renal neoplasia	120
	 PART 3 - EVALUATION OF THE HYPOTHESIS	123
X.	Summary of the evidence on the renal effects of CIGA	123
	A. Association between alpha-2u-globulin and nephrotoxicity	123
	B. Association between nephrotoxicity and renal cancer	125
	C. Information reducing confidence in the conclusion that the alpha-2u-globulin response is specific to the male rat	126
XI.	Conclusions	128
XII.	Research needs	134

PART 4 - POLICY

137

REFERENCES

138

APPENDICES

161

PREFACE

The U.S. Environmental Protection Agency (EPA) Risk Assessment Forum was established to promote scientific consensus on risk assessment issues and to ensure that this consensus is incorporated into appropriate risk assessment guidance. To accomplish this, the Risk Assessment Forum assembles experts from throughout the EPA in a formal process to study and report on these issues from an Agency-wide perspective.

For major risk assessment activities, the Risk Assessment Forum has established Technical Panels to conduct scientific review and analysis. Members are chosen to assure that necessary technical expertise is available. Outside experts may be invited to participate as consultants or, if appropriate, as Technical Panel members.

The use of male rat kidney tumors in risk assessment has been the subject of much recent discussion. For a certain group of chemicals, investigators have reported renal tubule tumor formation in male rats as the sequela of renal toxicity commencing with an excessive accumulation of the protein, alpha-2u-globulin (α_{2u} -g), in renal tubules. Renal tubule tumor formation with protein accumulation has not been observed in female rats or other tested species, most notably the mouse. The NCI Black Reiter rat, which does not produce α_{2u} -g, also fails to show a proliferative response in the kidney or evidence of a promotional effect when exposed to chemicals that induce protein droplet accumulation in male rats of other strains; its response has not been tested in a conventional two year animal bioassay. Some scientists apply the observations seen in animals to conclude that any renal tubule tumor in male rats observed in connection with α_{2u} -g accumulation is a species-specific effect inapplicable to human risk assessment. Other scientists argue that more information on humans is needed and that all male rat kidney tumors should continue to be considered as relevant to human risk as other tumors.

Because the question is relevant in assessing risk for a number of chemicals of interest to EPA, the Risk Assessment Forum established a Technical Panel to assemble and evaluate the current evidence and to develop science policy recommendations for Agency-wide use. This document is the product of that effort.

The literature review supporting this document is current as of February 25, 1991.

NOTE: Except for SAB/SAP review, the scientific analysis in this report is complete; editorial review is incomplete. Accordingly, this draft is being submitted simultaneously to the SAB for scientific peer review, and to technical editors for final editing, formatting, and reference review.

GLOSSARY

AAT	aspartate aminotransferase
ABS	chromosome aberrations in CHO cells
α_{2u} -g	alpha-2u-globulin
CHO	Chinese hamster ovary
CI	confidence interval
CIGA	<u>C</u> hemical(s) <u>I</u> nducing alpha-2u- <u>G</u> lobulin <u>A</u> ccumulation
CPN	chronic progressive nephropathy
1,2-DCB	1,2-dichlorobenzene
1,4-DCB	1,4-dichlorobenzene
DEN	diethylnitrosamine
DMN	dimethylnitrosamine
EHEN	N-ethyl-N-hydroxyethylnitrosamine
FBPA	N-4'-(fluoro-4-biphenyl)acetamide
IRDC	International Research and Development Corporation
MLA	TK-gene mutation assay in L5178Y cells
MTD	maximum tolerated dose
MUP	mouse major urinary protein
NAG	N-acetyl- β -glucosaminidase
NBR	NCI Black-Reiter rat
NTP	National Toxicology Program
NCI	National Cancer Institute
OR	odds ratio
RR	relative risk
P1	first convoluted segment of proximal tubule
P2	second convoluted segment of proximal tubule
P3	pars recta of proximal tubule
SAL	salmonella
SCE	sister chromatid exchange
SEER	Surveillance, Epidemiology and End Results Program of NCI
SLRL	sex-linked recessive lethal
TFT	trifluorothymidine
TK	thymidine-kinase
TMP	2,2,4-trimethylpentane
TMPOH	2,4,4-trimethylpentanol
UDS	unscheduled DNA synthesis

EXTERNAL PEER REVIEWERS

This draft report was evaluated at a two-day Peer Review Workshop sponsored by the U.S. EPA Risk Assessment Forum. The meeting, held in Gaithersburg, Maryland, on November 13 and 14, 1990, was chaired by Richard Griesemer, director of the Division of Toxicology Research and Testing, National Toxicology Program (NTP). A separate report of this meeting will be available as EPA publication no. [____]. In addition to plenary sessions on each day, workgroups were asked to address specific issues on four topics, nephropathy and biochemistry, cancer, criteria for evaluating renal carcinogens, and risk characterization.

The Nephropathy and Biochemistry Workgroup was chaired by Michael Olson of General Motors Research Laboratories. Other participants in that group included: Carl Potter (Risk Reduction Engineering Laboratory, Cincinnati) and James McKinney (Health Effects Research Laboratory, Research Triangle Park) of EPA, Benjamin Trump of the University of Maryland, and Dennis Lynch of the Division of Biological and Behavioral Sciences, National Institute for Occupational Safety and Health.

The Cancer Workgroup was chaired by John Ashby of the Central Toxicology Laboratory at International Chemical Industries, Ltd. Other members included: R. Daniel Benz of the Center for Food Safety and Applied Nutrition at the Food and Drug Administration, James Popp, head of the Department of Experimental Pathology and Toxicology at Chemical Industries Institute of Toxicology, Michael Elwell of NTP, and Joseph McLaughlin and Jerrold Ward of the National Cancer Institute.

The Criteria Workgroup was composed of pathologists who had specific research experience either in examining the lesions hypothesized to be associated with alpha-2u-globulin accumulation or in renal carcinogenesis. This group was chaired by Gordon Hard, of the Medical Research Council. Other members included: William Busey of Experimental Pathologies Laboratories, Scot Eustis of the NTP, Lois Lehman-McKeeman of Procter and Gamble Company, and James Swenberg of the University of North Carolina.

Participation in the Risk Characterization Workgroup was limited to government officials except for the chair, Norbert Page of Page Associates. Others were: William Farland and Penny Fenner-Crisp of EPA, Deborah Barsotti of the Division of Toxicology at the Agency for Toxic Substances Disease Registry, Murray Cohn of the Directorate for Health Sciences at Consumer Product Safety Commission, Elizabeth Grossman of the Office of Risk Assessment at the Occupational Safety and Health Administration, and Lauren Zeiss of the State of California Health Department.

Drafts prepared in advance of the Peer Review Workshop were reviewed and commented on by the following external reviewers.

Carl Alden (first and second drafts)
The Procter and Gamble Company
Miami Valley Laboratories
Cincinnati, Ohio

William Busey (first draft)
Experimental Pathology Laboratories
Herndon, VA

Gordon Hard (first draft)
Medical Research Council Toxicology Unit
Great Britain

Michael Lipsky (first draft)
University of Maryland
School of Medicine

Michael Olson (first and second drafts)
General Motors Research Laboratories
Warren, Michigan

James Popp (second draft)
Department of Experimental Pathology and Toxicology
Chemical Industries Institute for Toxicology
Research Triangle Park, North Carolina

James Swenberg (first draft)
University of North Carolina
Department of Pathology
Chapel Hill, North Carolina

The Technical Panel and the Risk Assessment Forum also acknowledge with appreciation the special contributions of Lawrence Valcovic, who prepared the section on mutagenicity, Joseph McLaughlin and Cheryl Siegel Scott, who greatly assisted in the preparation of the epidemiology section, and Ila Cote, Margaret M.L. Chu, and Richard N. Hill for their thoughtful comments.

EXECUTIVE SUMMARY

A Technical Panel of the U.S. Environmental Protection Agency's (EPA) Risk Assessment Forum advises EPA risk assessors against using information on renal tubule tumors or nephrotoxicity that is associated with alpha-2u-globulin (α_{2u} -g) accumulation in hyaline droplets in male rats to assess human risk. The scientific information reviewed by the Technical Panel provides reasonable evidence to suggest that the acute and chronic renal effects observed in male rats from chemically-induced α_{2u} -g accumulation are unlikely to occur in the absence of α_{2u} -g, or a protein with a structurally similar binding domain, in the large quantities typically seen in the male rat. Thus, if a chemical induces α_{2u} -g accumulation in hyaline droplets, the associated nephropathy observed in male rats may not be an appropriate endpoint for assessing noncancer risk in humans. Likewise, a carcinogenic response in the male rat kidney attributable to a process involving α_{2u} -g accumulation in the renal proximal tubule may not be an appropriate endpoint for assessing carcinogenic risk to humans.

The policy set out in this report provides guidance on determining when it is reasonable to presume that renal tumors in male rats result from a process involving α_{2u} -g accumulation and on selecting appropriate procedures for estimating risks to humans under such circumstances. It also defines situations that suggest different approaches and calls for research to clarify questions raised because of the existence of human proteins that may be structurally similar to α_{2u} -g.

In the male rat, the production of renal tumors by chemicals inducing α -2u-globulin accumulation (CIGA) is preceded by the renal lesions ascribed to α_{2u} -g-associated nephropathy. The involvement of hyaline droplet accumulation in the early nephrotoxicity associated with CIGA is a major difference from the sequence seen for classical carcinogens. The pathologic changes that precede the proliferative sequence for classical renal carcinogens also include a form of early nephrotoxicity, but no apparent hyaline droplet accumulation.

Investigations performed in multiple laboratories over the last decade have demonstrated a consistent association between hyaline droplets containing α_{2u} -g and production of certain lesions in the male rat kidney. These renal lesions are not found in mice, female rats, or other laboratory species tested. The histopathological sequence in the male rat consists of the following:

- (1) an excessive accumulation of hyaline droplets containing α_{2u} -g in renal proximal tubules;
- (2) subsequent cytotoxicity and single cell necrosis of the tubule epithelium;
- (3) sustained regenerative tubule cell proliferation, providing exposure continues;
- (4) development of intralumenal granular casts from sloughed cell debris associated with tubule dilation and papillary mineralization;
- (5) foci of tubule hyperplasia in the convoluted proximal tubules; and finally,
- (6) renal tubule tumors.

Biochemical studies with model compounds show that CIGA or

their metabolites bind specifically, but reversibly, to male rat α_{2u} -g. The resulting α_{2u} -g-CIGA complex appears to be more resistant to hydrolytic degradation by lysosomal enzymes than native, unbound α_{2u} -g. Inhibition of the catabolism of α_{2u} -g, a protein only slowly hydrolyzed by renal lysosomal enzymes under normal physiological conditions, provides a plausible basis for the initial stage of protein overload in the nephropathy sequence.

It is instructive to compare CIGA renal carcinogens with other renal carcinogens. Several genotoxic chemicals recognized as classical inducers of rodent kidney tumors have been used to study the pathogenesis of renal tubule cancer in laboratory animals. In general, these prototypic renal carcinogens produce tumors in both males and females. Although the wide range of chemicals represented suggests multiple mechanisms of action, many of the classical renal carcinogens or their active metabolites are electrophilic species able to bind covalently to macromolecules and likely to form DNA adducts in the kidney. In contrast, CIGA renal carcinogens are not known to react with DNA and are generally negative in short-term tests for genotoxicity. CIGA renal carcinogens also interact with α_{2u} -g in a reversible and noncovalent manner.

CIGA produced minimal changes in urine chemistry and very little or no glomerular dysfunction in male rats. The mild tubule toxicity of CIGA, in contrast to the obvious urinary changes induced by renal toxins such as mercuric chloride or hexachlorobutadiene, is characteristic of CIGA and is consistent

with the notion that CIGA do not bind covalently to α_{2u} -g.

Classical renal carcinogens, such as certain nitrosamines, induce renal tubule cancer in rats or mice with high incidence, minimal duration of exposure, and clear dose-response relationships. There is usually no absolute sex-specificity, although males and females may be susceptible to different degrees. In contrast, the renal tumors produced by the eight model carcinogens examined in this report tended not to be life-threatening, occurred late in life usually being found at terminal sacrifice, and were frequently microscopic. Even though the maximum tolerated dose was exceeded for some of the eight model carcinogens, the renal tumor incidence rate, adjusted for intercurrent mortality, was never greater than 28%. An increase in renal tubule tumors was not found in mice or female rats exposed to these chemicals. Initiation/promotion studies with gasoline, trimethylpentane (TMP), and d-limonene in Fischer rats showed that these CIGA promoted atypical tubule cell hyperplasia and/or renal tubule tumors in males but not in females. In contrast, d-limonene did not promote these lesions in males of the NCI Black-Reiter (NBR) strain in the same initiation/promotion model. Such differences in potency and species-, strain-, and sex-susceptibility suggest that CIGA renal carcinogens act via different mechanisms than classical renal carcinogens.

Renal tubule tumors produced by CIGA carcinogens also have features in common with other renal tubule tumors observed in the male rat. For renal carcinogens, in general, there is a continuum

of chemically induced steps from atypical hyperplasia through microscopic adenomas to macroscopic adenocarcinomas or carcinomas. Renal tubule tumors induced by the eight model carcinogens are morphologically indistinguishable from those induced by classical carcinogens. Likewise, the sequence of development of CIGA carcinogen-induced renal tumors from tubule cell hyperplasia to carcinoma appears identical. Furthermore, none of these chemically-induced tumors can be differentiated from spontaneous tumors.

All eight of the model carcinogens examined in this report were also capable of producing renal tubule hyperplasia in male rats. In general, this hyperplasia became more severe with increasing dose. The occurrence of these preneoplastic lesions together with the neoplastic lesions provides indirect evidence of progression that is in accord with generally accepted views on renal tubule tumor formation.

Dose- and time-related associations between the administration of CIGA to male rats and the various histological stages have been observed. These relationships were demonstrated between CIGA administration for both hyaline droplet formation and $\alpha_2\mu$ -g accumulation. Although the relationships between increased hyaline droplets and cell necrosis or between cell necrosis and cell regeneration have not been quantified, a correlation between hyaline droplet response and the number of cells excreted in the urine has been observed for CIGA. Dose-response relationships between hyaline droplet accumulation and proximal tubule cell

proliferation have been shown for TMP and unleaded gasoline. Clear dose-response relationships were demonstrated between linear mineralization in the renal medulla and incidence of renal tubule neoplasia in male rats in several bioassays. A recent study of d-limonene demonstrated a relationship between severity of nephropathy and renal tubule cancer in male rats.

The Technical Panel is not aware of any epidemiological study that has been designed or conducted specifically to examine the applicability of the CIGA hypothesis in humans. Several epidemiologic studies were reviewed for this report, but they are of limited value for this analysis because they involved exposure to complex blends, such as gasoline, or otherwise involved multiple exposures to both CIGA and non-CIGA. In addition, these studies were of limited statistical power and were not able to account for possibly confounding factors, such as smoking or obesity, which are known to influence renal cell cancer rates. In a few studies, slight increases in risk of renal cell cancer have been observed; however, it is difficult to identify specifically the agent responsible for the increased risk. These studies, therefore, are considered inadequate for purposes of hazard identification.

Low-molecular-weight proteins that probably have a three dimensional structure similar to α_{2u} -g have been identified in mice and other species, including humans. In vitro studies have shown that the active metabolite of TMP forms complexes with some of these proteins. Other in vitro studies indicate, however, that reversible binding does not necessarily increase resistance to

hydrolytic degradation, a feature apparently required for hyaline droplet formation.

Extensive studies in mice, whose urine contains large amounts of mouse major urinary proteins (MUP), have found no evidence of renal lesions similar to those associated with the α_{2u} -g syndrome. Thus, the presence of a structurally-related protein, even in large quantities in the urine, does not imply that another species will respond in a manner similar to the male rat.

The form of α_{2u} -g which originates in the liver of the male rat is not detected in the female rat. Like the mouse, the female rat shows no evidence of an α_{2u} -g-like nephropathy when exposed to CIGA. In cases where nephrotoxicity was observed in mice or female rats, it was less severe and qualitatively different from that in male rats and did not involve the spectrum of discrete lesions associated with α_{2u} -g accumulation in the male rat.

Specialized studies of rats, such as those involving immature, aged and castrated male rats, males of the NCI Black Reiter (NBR) strain (which does not synthesize α_{2u} -g in the liver), and injection of male rats with estrogen and female rats with α_{2u} -g, show that development of the early features of the specific nephropathy syndrome is dependent on the presence of α_{2u} -g. Very limited information in dogs, hamsters, guinea pigs, and monkeys also supports this statement. These studies further support the hypothesis that this α_{2u} -g-related nephropathy occurs specifically in the male rat.

In summation, the reversible binding of the compound to α_{2u} -

g, which results in a shift in balance between reabsorption and hydrolysis and the accumulation of α_{2u} -g in hyaline droplets in the P2 segment of the renal tubule provides a plausible explanation for the initial steps in a sequence of events leading to the formation of renal tubule tumors in the male rat. A sustained protein overload would result in single cell necrosis in the tubule epithelium and increased cell regeneration. This increased proliferative response caused by chemically-induced cytotoxicity may be a plausible reason for the development of renal tubule tumors in male rats, and renal tubule tumors produced in male rats where there was CIGA-induced α_{2u} -g nephropathy should be distinguished from other renal tumors wherever possible for purposes of their use in human risk assessment.

I. INTRODUCTION

For most hazardous chemicals, adequate human data are not available, and risk analyses must rely on information from laboratory studies of rats or mice. The inference that the results of animal experiments can be applied to humans is a fundamental principle of all toxicologic research. This paper deals with a specific case, however, where the male rat seems to respond in a different manner than other laboratory species. The possibility of a unique response in the rat among laboratory animals raises questions about the applicability of certain rat data to other species, including humans. This document provides guidance for the assessment of such information.

A variety of organic chemicals have produced specific renal lesions in male rats, in the form of a hyaline droplet nephropathy accompanied by accumulation of the protein, alpha-2u-globulin (α_{2u} -g) (reviewed in HEI, 1985, 1988). Among the chemicals tested are paraffins (Halder et al., 1984; Phillips and Cockrell, 1984), decalin (decahydronaphthalene) (Alden et al., 1984; Kanerva et al., 1987a), petroleum-based and synthetic fuels (MacNaughton and Uddin, 1984), military aviation propellants (Bruner, 1984), and 2,2,4-trimethylpentane (TMP) (Halder et al., 1985). As seen in Table 1, which lists a sampling of chemicals that have been tested, many are of major regulatory and commercial interest. For example, isophorone is a chemical intermediate of major industrial importance. Aviation and automotive fuels fit into the category, as does the natural food product, d-limonene, found in citrus oils.

TABLE 1. SOME EXAMPLES OF ORGANIC CHEMICALS THAT PRODUCE RENAL INJURY IN MALE RATS CHARACTERIZED BY HYALINE DROPLET ACCUMULATION BUT NOT IN FEMALE RATS OR OTHER SPECIES

CHEMICAL	SPECIES TESTED	RENAL TOXICITY	REFERENCE
Decalin	Rats (m/f) Mice (m/f) Dogs (m/f) Guinea pigs (m/f)	+/- -/- -/- -/-	Alden et al. (1985) USEPA (1987)
Dimethyl methyl phosphonate	Rats (m/f) Mice (m/f)	+/- -/-	NTP-TR-323 (1987b)
Isophorone	Rats (m/f) Mice (m/f)	+/- -/-	NTP-TR-291 (1986a)
JP-5 shale-derived jet fuel	Rats (m/f) Mice (m/f) Dogs (m/f)	+/- -/- -/-	MacNaughton & Uddin (1984)
JP-4 jet fuel	Rats (m/f) Mice (m/f) Dogs (m/f)	+/- -/- -/-	MacNaughton & Uddin (1984)
d-Limonene	Rats (m/f) Mice (m/f) Dogs (m)	+/- -/- -	NTP-TR-347 (1990) Webb et al. (1991)
Methyl isobutyl ketone	Rats (m/f) Mice (m/f) Dogs (m) Monkeys (m)	+/- -/- - -	Alden et al. (1984) Phillips et al. (1987)
Pentachloroethane	Rats (m/f) Mice (m/f)	+/- -/-	NTP-TR-232 (1983)
Unleaded gasoline	Rats (m/f) Mice (m/f)	+/- -/-	USEPA (1987)

m = male
f = female
+ = positive
- = negative

This analysis focuses on model compounds having both an adequate animal carcinogenesis bioassay and information on $\alpha_2\text{u}$ -g or hyaline droplet accumulation in the male rat. These substances are seven chemicals, 1,4-dichlorobenzene (1,4-DCB), dimethyl methyl phosphonate, hexachloroethane, isophorone, d-limonene, pentachloroethane, tetrachloroethylene, and a mixture, unleaded gasoline. These eight substances are compared and contrasted with two related non- $\alpha_2\text{u}$ -g-inducers, chlorothalonil and trichloroethylene. The analysis also relies on research studies on two other model compounds, decalin and 2,2,4-trimethylpentane (TMP), which have extensive information on $\alpha_2\text{u}$ -g nephropathy but no chronic bioassay data. More limited data on 22 additional substances is also discussed where appropriate.

Among the eight model chemicals tested in chronic animal bioassays, all invoked a specific type of protein droplet nephropathy in male rats and all also produced renal tumors in male rats but not in other species tested. It has been proposed that such renal tumors are the end product in the following sequence of functional changes in the epithelial cells of proximal tubules (UAREP, 1983; Alden et al., 1984; Halder et al., 1984; HEI, 1988; Swenberg et al., 1989a).

- Excessive accumulation of hyaline droplets in proximal tubules, representing lysosomal overload, leads to tubule cell degeneration, cell loss, and regenerative cellular proliferation.
- Cell debris in the form of granular casts accumulates at the "corticomedullary" junction with associated dilation of the affected tubule segment and more distally, mineralization of tubules within the renal medulla.

- Single cell necrosis accompanied by compensatory cell proliferation and exacerbation of the chronic progressive nephropathy (CPN) characteristically found in aging rats occurs.
- Renal tubule hyperplasia and neoplasia develop subsequently.

According to this hypothesis, the increased proliferative response caused by the chemically-induced cytotoxicity results in clonal expansion of spontaneously initiated renal tubule cells and increased incidence of renal tumor formation (Trump et al., 1984b; Alden, 1989; Swenberg et al., 1989a). This line of reasoning leads supporters of the hypothesis to conclude that the acute and chronic renal effects induced in male rats by these chemicals will be unlikely to occur in any species not producing α_{2u} -g or a very closely related protein in the large quantities typically seen in the male rat (Alden 1989; Borghoff et al., 1990; Green et al., 1990; Olson et al., 1990; Flamm and Lehman-McKeeman, 1991).

This report examines the hypothesis that the male rat is predisposed to the nephrotoxic effects induced by certain classes of chemicals, such as volatile light hydrocarbons and organohalides. It also examines data that support or contradict the concept that the renal tumors produced in male rats by these chemicals are causally related to the nephrotoxicity. Based on the Risk Assessment Forum's conclusions regarding these data, the document proposes a uniform approach for EPA to use in risk assessments dealing with this spectrum of lesions and category of chemicals.

Information for this Risk Assessment Forum report was obtained

initially from a 1988 review entitled "Evaluation of Data Concerning the Relationships among Chemically-induced Renal Alpha_{2u}Globulin or Hyaline Droplet Accumulation, Nephropathy, and Renal Neoplasia" prepared for the Office of Toxic Substances by Dr. William Richards of Dynamac Corporation, Rockville, Maryland. Additional information considered in this report includes recent comprehensive reviews of the subject, comments from peer reviewers, and other original work, especially publications subsequent to the 1988 Dynamac review.

The document has four parts. Following this brief introduction, Part 1 addresses the characteristics of hyaline droplets and the protein, α_{2u} -g, and the nephropathy associated with α_{2u} -g accumulation (Sections II and III).

Part 2 (Sections IV-IX) presents data on the carcinogenic potential of CIGA in the male rat. Section IV describes the preneoplastic and neoplastic lesions produced by classical renal carcinogens. Section V considers generic factors relevant to all studies of potential renal carcinogenicity in laboratory animals and then analyzes and discusses data on the renal lesions observed in 2-year bioassays with chemicals causing the hyaline droplet nephropathy. Section VI examines additional information that assists in defining renal carcinogens as CIGA, in particular genotoxicity and initiation-promotion data. In Section VII, CIGA are compared with classical renal carcinogens, while Section VIII considers the human evidence for kidney cancer, its histogenesis and epidemiology. Section IX examines evidence for dose- and time-

dependent progression of the lesions hypothesized to lead to this nephropathy.

Part 3 evaluates the evidence considered in Parts 1 and 2 with regard to the hypothesis that α_{2u} -g accumulation in the kidney is an initial step in a succession of histopathologic events that may culminate in renal tubule tumor formation in male rats. This part also lists priorities for future research.

Part 4 comprises the Agency policy statement regarding approaches to risk assessment for this category of chemicals.

For clarity throughout the review, nomenclature is standardized, and abbreviations are used for frequently repeated terms. Insofar as hyaline droplet represents a morphological entity requiring only light microscopy for identification, this term will be used in preference to the synonymous protein droplet¹. The designation, alpha-2u-globulin (α_{2u} -g) nephropathy is used to connote the full sequence of pathologic lesions from hyaline droplet formation to restorative hyperplasia and medullary mineralization. Toxic tubular nephropathy is a non-specific term commonly used in rodent bioassay reports to describe various forms of nephrotoxicity induced by chemicals, including the specific lesions of α_{2u} -g nephropathy. The spontaneous age-related syndrome of rat kidney disease otherwise known in the literature as old rat

¹Hyaline droplets refer to spherical inclusions in the cytoplasm which are homogeneous and eosinophilic, representing overdistended phagolysosomes. They may contain various macromolecules including α -2u-globulin. The morphology of droplets containing different proteins may be identical and therefore immunocytochemistry is required for precise definition of contents.

nephropathy, chronic nephrosis, glomerulosclerosis, and progressive glomerulonephrosis, is standardized according to Barthold (1979) as chronic progressive nephropathy (CPN). The term lipocalin is used according to the terminology of Pervaiz and Brew (1987) to describe the superfamily of low-molecular-weight proteins which appear to transport lipophilic substances.

In rats, the proximal tubule of the nephron is divisible morphologically into three parts (see Figure 1). The first segment is in continuity with the parietal epithelium of Bowman's capsule surrounding the glomerular tuft. Together, the first and second segments represent the convoluted portion of the proximal tubule and are situated wholly in the cortex, the outermost zone of the rat kidney. The third segment is the straight portion of the proximal tubule (pars recta) comprising the outer stripe of the outer medulla but also the medullary rays arising in the cortex. The abbreviations P1, P2, P3 are used conventionally to denote these three segments. The term renal tubule tumor describes neoplasms of the renal cortical tubule epithelium comprising collectively adenoma, adenocarcinoma and carcinoma according to standardized nomenclature determined by the Society of Toxicologic Pathologists (Hard et al., 1991). The same neoplasms are referred to as renal cell tumors in humans, in keeping with the general literature (Bannayan and Lamm, 1980).




FIGURE 1. DIAGRAM OF ZONATION AND TUBULE SEGMENTATION IN RAT KIDNEY.

G: Glomerulus; P₁: First segment of proximal convoluted tubule; P₂: Second segment of proximal convoluted tubule; P₃: Pars recta of proximal tubule.
(Adapted from Bachmann et al., 1986)

PART 1. NEPHROTOXICITY

II. HYALINE DROPLETS AND ALPHA-2u-GLOBULIN; PHYSIOLOGY AND BIOCHEMISTRY

Information on the renal processing of low-molecular-weight proteins, sex and species differences in urinary proteins, and the characteristics of α_{2u} -g provides an explanatory basis for the accumulation of α_{2u} -g in hyaline droplets in the male rat following exposure to CIGA. It is pertinent, therefore, to examine the physiological and biochemical characteristics of α_{2u} -g and related proteins, particularly those that occur in humans, before exploring the possible associations between α_{2u} -g accumulation, renal toxicity and renal tumor formation and their relevance to human risk assessment.

A. Filtration, reabsorption, and catabolism of low-molecular-weight proteins by the kidney

The mammalian kidney has a major role in maintaining the plasma concentrations of circulating low-molecular-weight proteins at their normally low, physiological levels. Thus, low-molecular-weight proteins are continually removed from the plasma by glomerular filtration followed by reabsorption and catabolism in the proximal tubules (Maack et al., 1985), or excretion. Figure 2 is a schematic representation of the cellular uptake and disposition of filtered proteins by the renal tubule.




FIGURE 2: Schematic representation of endocytic uptake of filtered proteins. Filtered proteins are adsorbed to endocytic sites at the luminal membrane and segregated in endocytic vacuoles (EV). These EV migrate to the cell interior, where they fuse with lysosomes (L) to form secondary lysosomes (SL) or phagolysosome where digestion of the protein takes place. The products of hydrolysis (amino acids) permeate the SL membrane, cross the contraluminal cell membrane, and return to the circulation. (Adapted from Kaysen et al., 1986).

The normal renal glomerulus freely passes proteins with a molecular weight of less than 20,000 daltons, including peptides such as insulin, lysozyme, rat growth hormone, myoglobin, and cytochrome C (Maack et al., 1985). For larger proteins like the albumins and globulins, which have a far greater plasma concentration and much lower filtration rate than low-molecular-weight proteins, the kidney has no regulating role in plasma protein concentration.

Reabsorption of filtered protein occurs predominantly in the convoluted part of the proximal tubule and to a lesser extent in the pars recta cells. Tubular absorption of a protein is a complex process initiated by binding of the protein to the microvilli of the proximal tubule epithelium. This is followed by migration to the base of the microvilli and adsorptive endocytosis whereby invagination of the surface membrane internalizes the protein (Kaysen et al., 1986). While reabsorption was once considered largely non-selective, high capacity, low affinity transport (Maack et al., 1985), from recent work it now appears that interaction between the protein and the brush border membrane is the step at which a degree of selectivity in the absorption process occurs (Kaysen et al., 1986).

Within proximal tubule cells, endocytic vesicles fuse to form endocytic vacuoles which in turn coalesce with lysosomes derived from the Golgi apparatus, forming secondary lysosomes. The hydrolysis of proteins by protease enzymes takes place within the secondary lysosomes. The lysosomal enzymes of renal cortical

tubules include two major classes of acid proteinases - cysteine proteinases (cathepsin B, H and L) and an aspartic acid proteinase, cathepsin D (Lehman-McKeeman et al., 1990a). Lehman-McKeeman et al. (1990a) have shown that both of these endopeptidase classes contribute to the degradation of α_{2u} -g.

Lysosomes have a large, but not unlimited, capacity to cope with increased amounts of hydrolyzable proteins, but the proteins differ in susceptibility to hydrolysis. Protein half-lives, which are indices of their catabolism by proteases in the kidney, depend on specific molecular determinants in the protein. The primary amino acid sequence may be one important factor in determining protein half-lives (Dice, 1987). The plasma half-lives of many low-molecular-weight proteins are measured typically in minutes (Maack et al., 1985). Alpha-2u-globulin, with a half-life measured in hours (Geertzen et al., 1973), is one of the exceptions.

Whether or not low-molecular-weight proteins like α_{2u} -g accumulate in kidney tubules depends on the balance between the rate of reabsorption by epithelium and the rate of hydrolysis in the cells. Based on the information presented below, it is believed that exposure to CIGA results in a shift of this balance in male rats.

B. Hyaline droplets in renal tubules

The product of protein reabsorption and accumulation in renal tubule cells is visualized by light microscopy as hyaline droplets. Small protein reabsorption droplets of uniform size are a

constitutive feature of normal mature male rats being particularly evident in the P2 segment of proximal tubules (Logothetopoulos and Weinbren, 1955; Maunsbach, 1966a; Goldsworthy et al., 1988a). Ultrastructurally, hyaline droplets are abnormally large, dense, secondary lysosomes (also termed phagolysosomes), representing fusion of endocytic vacuoles with primary lysosomes. Some hyaline droplets show crystalloid changes by electron microscopy which are not observed in the lysosomes of female rats (Maunsbach, 1966b). Crystalline formation in the normal male rat is believed to indicate the presence of a poorly catabolized protein in pure solution (Pesce et al., 1980), presumably α_{2u} -g in the kidney lysosomes.

Hyaline droplets in the proximal tubules of normal male rats contain α_{2u} -g (Alden et al., 1984; Garg et al., 1987; Goldsworthy et al., 1988), and their occurrence appears to parallel the variable synthesis of this protein. Thus, hyaline droplets become apparent in male rats at the time of puberty, but they decline progressively with increasing age after 18 months (Logothetopoulos and Weinbren, 1955; Murty et al., 1988). In female rats, protein droplets in proximal tubules are either absent, or considerably less frequent than in males, and they do not contain α_{2u} -g (Logothetopoulos and Weinbren, 1955; Maunsbach, 1966a; Goldsworthy et al., 1988a; Burnett et al., 1989). Hyaline droplets are substantially reduced in castrated male rats (Logothetopoulos and Weinbren, 1955).

Because an abnormal increase in hyaline droplets has more than

one etiology and can be associated with the accumulation of different proteins, it is necessary to apply special diagnostic methods such as immunohistochemical staining to make the association between chemical exposure and pathologic accumulation of α_{2u} -g.

Abnormal accumulation of hyaline droplets in rodent kidney is seen in certain disease processes. Both male and female rats with histiocytic sarcoma show hyaline droplet accumulation in the proximal tubules, indistinguishable from the CIGA-induced lesion. The accumulating protein in these tumor-bearing animals has been identified as lysozyme (Hard and Snowdon, 1991). Similarly, in male and female mice with histiocytic tumors, abnormal accumulation of lysozyme-containing hyaline droplets sometimes occurs in proximal tubules (Hard and Snowdon, 1991).

In humans, the Bence-Jones proteins, a class of light chain immunoglobulins, are produced in large amounts in multiple myeloma patients (Pirani et al., 1983). In cases of mononuclear cell leukemia, lysozyme is produced (Muggia et al., 1969). The kidney injury seen with these neoplastic diseases has been described as similar to that produced by administration of decalin to male rats (Alden, 1986), including protein droplet accumulation in renal tubules (Oliver and MacDowell, 1958; Pirani et al, 1983; Pruzansky and Platts, 1970). Patients with epidemic hemorrhagic fever, infused with large amounts of concentrated human serum albumin as a therapeutic procedure for shock have also developed a comparable form of hyaline droplet accumulation (Oliver and MacDowell, 1958).

C. Factors affecting kidney accumulation of low-molecular-weight proteins

Protein accumulation in the proximal tubule can reach pathological levels resulting in excessive hyaline droplet formation for several reasons: (1) the rate of protein delivery to the tubule cells is abnormally high, (2) the proteins delivered are difficult to hydrolyze, or (3) the lysosomal hydrolysis capacity is sufficiently reduced.

The rate of protein delivery to the tubule can be abnormally high under conditions when the capillary wall of the glomerulus fails to provide the normal filtration barrier. This happens, for example where there is immunological, inflammatory, or toxic disease in the glomerulus or when the permselectivity barrier is overloaded by filterable proteins (Kaysen et al., 1986).

The increased urinary excretion of low-molecular weight proteins seen in diseases, such as multiple myeloma in humans or histiocytic sarcoma in rats, is primarily the result of an increase in plasma concentration caused by overproduction of specific small proteins (Maack et al., 1985). Lysozyme (histiocytic sarcoma) and light chain immunoglobulins (multiple myeloma) are proteins also relatively resistant to hydrolysis (Maack et al., 1985). This suggests a combination of features (1) and (2) as an etiologic factor in the accumulation of protein observed in rats with histiocytic sarcoma (lysozyme) and in human patients with multiple myeloma (light chain immunoglobulins). The combination of difficult hydrolysis of the protein, as suggested by its long half life, coupled with high rate of protein delivery to tubule cells

in the sexually mature male rat also appears to be a factor in the accumulation of α_{2u} -g in the renal tubules of male rats.

The process of protein hydrolysis can be reduced or inhibited when lysosomes are unable to maintain the low pH required for hydrolytic enzyme function. Inhibition of the metabolically driven hydrogen ion pump, by metabolic poisons or the presence of a weak base in tubule lysosomes, alters the pH and results in the accumulation of proteins (Maack et al., 1985). In the presence of a reduced lysosomal hydrolysis capacity, the most hydrolytically resistant proteins, like α_{2u} -g, tend to accumulate first. Testosterone is known to have a suppressive effect on the activity of some major proteolytic enzymes in the male rat kidney (Kugler and Vornberger, 1986). Consequently, the lysosomal protease activity in male proximal tubules is lower than those of females (Jedrzejewski and Kugler, 1982; Kugler and Vornberger, 1986) implying that the male rat could be intrinsically more prone to protein overload in the renal tubules than the female rat.

Reduction of the hydrolytic capacity of renal lysosomes and increased resistance of protein to hydrolysis can both be affected by exogenous chemicals. Although CIGA may not compromise kidney lysosomal enzyme activity per se (Murty et al., 1988; Lehman-McKeeman et al., 1990), any chemically-induced impediment to α_{2u} -g digestibility caused by CIGA would be further superimposed on the factors considered above that alone can cause excessive protein accumulation in renal tubules.

D. The alpha-2u-globulin superfamily of proteins

Alpha 2u-globulin is a member of a large superfamily of low molecular weight proteins. The complete amino acid sequence of α_{2u} -g was first deduced by Unterman et al. (1981). Even though, with the exception of α_{2u} -g and mouse major urinary protein(s) (MUP), the sequence homology between any pair of proteins in this superfamily is small, about 20 percent, statistical analysis shows that the proteins are related evolutionarily (Akerstrom and Logdberg, 1990).

Of the approximately 20 proteins now considered to be potential members of the superfamily (Akerstrom and Logdberg, 1990), the three dimensional structure is known for only three, retinol-binding protein, β -lactoglobulin, and insecticyanin (Sawyer, 1987). The central core of these three proteins is composed of eight strands with a β -barrel structure forming a hydrophobic pocket that appears to enclose the ligand (Papiz et al., 1986; Sawyer, 1987). This structure has been described as resembling a coffee filter paper (Akerstrom and Logdberg, 1990). In addition to the β -structural motif, one helical rod and several other structural elements appear to be conserved among the proteins. Protein folding patterns tend to be highly conserved in homologous proteins even though they may diverge considerably in structure and function, suggesting that other members of the superfamily, including α_{2u} -g, possess a similar three dimensional structure.

The only member of the protein superfamily with a clearly

defined physiological function is retinol-binding protein. More circumstantial evidence suggests that the superfamily members serve as carriers of lipophilic molecules (Pervaiz and Brew, 1987). The mode of binding in which the lipid ligand is enclosed within the β -barrel impressed Pervaiz and Brew as not unlike the role of the calyx to a flower. On this basis, they suggested the illustrative name, lipocalins, for the superfamily of proteins.

Table 2 illustrates the information available on several members of the lipocalin superfamily, which includes α_{2u} -g, retinol-binding protein, apolipoprotein D, α_1 -acid glycoprotein and α_1 -microglobulin of humans, bovine β -lactoglobulin and pyrazine-binding protein (i.e., odorant-binding protein), rat odorant-binding protein and major urinary protein(s) (MUP) of mice. Some of the members of the lipocalin superfamily, such as retinol-binding protein, β -lactoglobulin, and α_1 -microglobulin have been identified in many species, and their properties appear to be species independent, suggesting that they share a common vital function (Akerstrom and Logdberg, 1990). Others, such as α_{2u} -g and MUP seem to be species-specific.

TABLE 2. SUPERFAMILY OF LIPOPHILIC LIGAND-BINDING CARRIER PROTEINS ¹

SPECIES	PROTEIN	TISSUE OR BODY FLUID	MOLECULAR WEIGHT	NO. OF AMINO ACIDS
Human	α_1 -Acid glycoprotein	Plasma	18,944 ²	167
	Apolipoprotein D	Plasma	19,300	169
	Pregnancy-associated endometrial α_2 -globulin	Placenta	25,000	Not known
	Protein HC; α_1 microglobulin	Plasma, urine, spinal fluid	20,619	182
	Retinol-binding protein	Liver	22,868	199
Cow	β -lactoglobulin	Milk	18,281	162
	Pyrazine-binding protein	Nasal epithelium	19,000	Not known ³
Rat	α_{2u} -globulin	Primarily male liver, urine	18,709	162
	Androgen-dependent secretory protein	Epididymis	18,500	184
	Odorant-binding protein	Nasal epithelium	18,091	172
	Fatty-acid-binding protein	Liver	14,000	Not known ⁴
Mouse	Major urinary protein	Liver (both sexes), urine	18,730	162
Chick	Purpurin	Retina	21,924	196
Frog	Bowman's gland protein	Olfactory epithelium	20,300	182
Insect ⁵	Insecticyanin	Hemolymph	21,382	189

¹ Adapted from Pevsner et al., 1988, except as noted.

² In rat (Pervaiz and Brew, 1987)

³ Cavaggioni, et al., 1987

⁴ Kimura, et al., 1989

⁵ Tobacco hornworm

Several functions have been suggested for α_{2u} -g. Cavaggioni et al. (1987) speculated that α_{2u} -g may serve to transfer odorants such as ethereal lipid pheromones from male rat urine to the air for attracting females. Glandular tissue production of α_{2u} -g helps support these speculations (Murthy et al., 1987; Mancini et al., 1989). In addition, α_{2u} -g has been identified as a fatty acid-binding protein of the kidney (Kimura et al., 1989) and may serve to transport fatty acid, an important energy source in kidney, within renal epithelial cells. Brooks (1987) found a protein structurally related to α_{2u} -g which is synthesized and secreted by the rat epididymis under the influence of androgenic hormones. He speculated that the function of these proteins may be to carry retinoids within the lumen of the male reproductive tract.

Other members of the lipocalin superfamily, such as retinol-binding protein, apolipoprotein D, β -lactoglobulin, and α_1 -acid glycoprotein, function in the transport of lipids between cells and across hydrophilic barriers (Pevsner et al., 1988). The lipids bound by the proteins differ considerably in structure and range from odorants in rat nasal epithelium to human cholesterol and retinol (vitamin A). It is not yet clear how selective these proteins are for specific ligands or whether a given protein might bind a wide spectrum of small hydrophobic molecules. Both cases might occur since retinol-binding protein is quite specific for retinol, whereas odorant-binding proteins may have a broad specificity (Godovac-Zimmermann, 1988).

Cavaggioni et al. (1990) reported substantial differences in

the binding affinities of α_{2u} -g, MUP and pyrazine-binding protein isolated from calf nasal mucosa for a series of odorants. MUP bound only one of these chemicals; pyrazine-binding protein bound six; and α_{2u} -g bound twelve. The best ligands for each of the three proteins were chemically unrelated; close structural analogs of the best ligands were also only weak ligands, much weaker than structurally unrelated chemicals. This study suggests that structure-activity and binding affinities for one lipocalin are poor predictors for other members of the superfamily.

E. Characteristics of alpha-2u-globulin

Alpha-2u-globulin was first characterized in male rat urine (Roy and Neuhaus, 1967). All isoforms of α_{2u} -g are anionic at neutral pH although they have varying isoelectric points. The molecular weight of α_{2u} -g has been reported to be 18,000-20,000 daltons. In all known rat strains, except for the NCI Black-Reiter (NBR) rat (Chatterjee et al., 1989), the major urinary source of α_{2u} -g is the liver where α_{2u} -g mRNA constitutes approximately 1% of the hepatic mRNA population (Sippel et al., 1976; Kurtz and Feigelson, 1978). The hepatic isoforms of α_{2u} -g may vary throughout the lifetime (Roy et al., 1983). Synthesis of the protein in rat liver is under multihormonal control, particularly androgen, but also glucocorticoids, thyroid hormones, insulin and growth hormone (Feigelson and Kurtz, 1977; Roy and Chatterjee, 1983). These hormones appear to act by regulating the steady-state level of α_{2u} -g mRNA (Kurtz and Feigelson, 1977). Neither α_{2u} -g nor its corresponding mRNA are detectable in the livers of

sexually intact female rats (Sippel et al., 1975, 1976; MacInnes et al., 1986). However, a very low background level of the mRNA has been indicated in the ovariectomized female rat (Chatterjee et al., 1979), and ovariectomy in concert with androgen treatment induces a parallel increase in α_{2u} -g and its mRNA in female rat liver (Roy and Neuhaus, 1967; Sippel et al., 1975).

Although plasma and urinary α_{2u} -g derives predominantly from the liver in male rats, high levels of α_{2u} -g and its mRNA are also present in the preputial gland of both male and female rats, and neither castration nor ovariectomy significantly alter the preputial concentration of this protein and its mRNA (Murty et al., 1988). Alpha-2u-globulin mRNA has also been detected in the female mammary gland during pregnancy, and in the submaxillary, lacrymal, Meibomian, and perianal glands of rats of both sexes (MacInnes et al., 1986; Mancini et al., 1989). The female forms of α_{2u} -g show distinct differences from male rat α_{2u} -g suggesting that they are encoded by different genes (Vandoren et al., 1983).

Low levels of α_{2u} -g first become detectable in the male rat liver under the stimulus of testosterone at 35-40 days, reaching maximum adult levels by 60-80 days (Roy et al., 1983; MacInnes et al., 1986; Motwani et al., 1984). At some stage after 100-150 days of age, due to the development of hepatic insensitivity to androgen during senescence, hepatic synthesis of α_{2u} -g falls gradually to 50 percent of peak levels in 600 day old male rats, beyond 750 days of age, becoming undetectable (Roy et al., 1983; Motwani et al., 1984). Renal cortical tissue content (Murty et al., 1988) and

urinary excretion (Neuhaus and Flory, 1978; Motwani et al., 1984) of α_{2u} -g reflect the same age related trends as synthesis in the liver.

In the mature male rat, approximately 50 mg of α_{2u} -g is filtered per day, 40% of the filtered protein being excreted in the urine and 60% undergoing reabsorption and catabolism (Neuhaus et al., 1981; Caudill et al., 1991). It is catabolized slowly relative to most other proteins in the glomerular filtrate, the half-life in plasma or kidney cytosol or lysosomal preparations being 5-8 hours (Geertzen et al., 1973; Ekstrom, 1983; Lehman-McKeeman et al., 1990a). In vitro studies indicate that α_{2u} -g is more resistant to lysosomal enzyme digestion than bovine β -lactoglobulin and lysozyme (Charbonneau et al., 1988). In another study comparing members of the protein superfamily, α_{2u} -g and α_1 -acid glycoprotein were the most resistant to proteinase K digestion while retinol-binding protein and β -lactoglobulin were 1000- to 100,000-fold more easily hydrolysed (Borghoff et al., 1990). These data indicate that α_{2u} -g may be more likely to accumulate in the kidney than most other members of the superfamily if shifts in the balance between reabsorption and hydrolysis occur.

F. Sex and species comparison of urinary protein content of lipocalin superfamily

Relative to the female rat, and other species including humans, the normal mature male rat is physiologically proteinuric. This is due to the amount of α_{2u} -g in male rat urine, 1.36-8.64 mg/day/gm kidney (Neuhaus and Lerseth, 1979), which is 100 to 300 times more than observed in female rat urine (Shapiro and

Sachchidananda, 1982; Vandoren et al., 1983). The mouse can also be described as physiologically proteinuric because of a high urinary content of MUP (Thung, 1962). MUP shows the greatest similarity to α_{2u} -g in the lipocalin superfamily, sharing 90% amino acid sequence homology (Dolan et al., 1982). Representing a group of proteins encoded by a multigene family, MUP is synthesized in the liver of mice of both sexes but at rates four to five times greater in males than females (Hastie et al., 1979; Roy and Chatterjee, 1983). Daily urinary excretion of MUP varies considerably among strains (Szoka and Paigen, 1978). In the B6C3F1 strain, males have been shown to excrete 14.9 mg of MUP/day in the urine, and females, 2.1 mg/day (Lehman-McKeeman et al., 1990b). Adjusted for body weight, a male B6C3F1 mouse therefore excretes approximately 600 mg/kg/day of MUP, some 12 fold higher than α_{2u} -g urinary excretion by the male rat. Unlike the rat, however, where 60% of filtered α_{2u} -g is reabsorbed by the kidney, MUP is not reabsorbed in the mouse and appears to be totally excreted (Caudill et al., 1991).

In contrast, normal human urine contains relatively little protein, only 1% of the total concentration present in mature male rat urine (Olson et al., 1990). Human urinary proteins are predominantly high-molecular-weight species with only minor components weighing less than 66,000 daltons. Within the low-molecular-weight fraction, trace amounts of proteins represent the lipocalin superfamily, but none appear to share molecular weight identity with α_{2u} -g. The urinary excretion of retinol-binding

protein, α_1 -acid glycoprotein and α_1 -microglobulin has been measured at 0.0001-0.0007, 0.0006-0.002, and 0.02-0.05 mg/day/gm kidney, respectively (Berggard, 1970; Peterson and Berggard, 1971; Ekstrom and Berggard, 1977). Thus, the urinary excretion of α_{2u} -g in the male rat is approximately two orders of magnitude greater than the human urinary content of the three superfamily proteins combined.

Recently, a sex-dependent protein of unknown origin and function, termed urine protein 1, was identified in normal human urine (Bernard et al., 1989, 1990). The molecular features of protein 1 are similar to α_{2u} -g as it has a molecular weight of approximately 21,000 daltons and an isoelectric point around 4.8 (Bernard et al., 1990). As its amino acids have not been sequenced, it cannot be placed in the lipocalin superfamily. Protein 1 occurs in both sexes from an early age, but increases substantially in males after puberty, reaching up to a fifty-fold difference over females during late adolescence. A five-fold male to female differential persists through adulthood. Average urinary concentrations of protein 1 have been determined as 108 and 3.2 μ g/liter respectively for males and females aged 15 to 20 years, and 24.7 and 5.8 μ g/liter for males and females in the 20 to 60 year age-range (Bernard et al., 1989). Such levels of protein 1 in human male urine, however, are calculated as four to five orders of magnitude less than α_{2u} -g concentrations in the urine of male rats (Bernard et al., 1990).

G. Noncovalent binding to alpha-2u-globulin and its homologues

It has been suggested that CIGA bind reversibly and

noncovalently to α_{2u} -g in the male rat, forming a resultant complex that is even more poorly digested than α_{2u} -g (Swenberg et al., 1989a).

In a few instances, the specific chemical entity complexed with α_{2u} -g has been identified. TMP, a branched chain aliphatic hydrocarbon present in gasoline (Halder et al., 1985) was the first model CIGA to be studied in this manner. When [14-C]TMP was administered in a single oral dose to male or female rats, radioactivity was retained in the kidneys of males, but not of females (Kloss et al., 1985; Charbonneau et al., 1987). The major metabolite of TMP detected in the kidneys of male rats was identified as 2,4,4-trimethyl-2-pentanol (TMPOH) (Charbonneau et al., 1987). In a separate report, it was demonstrated that TMPOH is the only compound that binds to α_{2u} -g whenever TMP is administered to the male rat (Lock et al., 1987a). TMPOH was not detected in the kidney tissue of the female rats, which excreted more conjugated TMPOH (glucuronides and sulfates) than the males (Charbonneau et al., 1987). Later studies confirmed, as suspected, that the TMPOH- α_{2u} -g complex is cleared slowly from male rat kidney (Swenberg, 1989b).

It has been shown since for d-limonene, that the metabolite interacting predominantly with α_{2u} -g is d-limonene-1,2-oxide although there is also some binding to the parent material (Lehman-McKeeman et al., 1989). For isophorone, the bound material is the parent compound (Strasser et al., 1988). Following exposure of the male rat to 1,4-DCB, both the parent chemical and the metabolite,

2,5-dichlorophenol, bind reversibly to α_{2u} -g (Charbonneau et al., 1989).

The nature of the association of CIGA with α_{2u} -g was explored initially by Lock et al. (1987a) who dosed sexually mature Fischer 344 rats with [3-H]TMP, killed them 8-72 hours later, and homogenized the kidneys. Cytosol, obtained by centrifugation of the homogenate at 116,000 g, was applied to a Sephadex G-75 column. For the males, 26% of the cytosol radiolabel (15% of all radiolabel in the kidney) eluted in the fraction containing α_{2u} -g. Approximately 19% of the radiolabel present in male rat kidney cytosol was nondialyzable following overnight equilibrium dialysis against phosphate buffer. Chromatography of the dialyzed cytosol showed that the nondialyzable radiolabeled material coeluted with the peak containing α_{2u} -g. When 0.1% sodium dodecyl sulfate, a detergent which affects the secondary and tertiary structure of proteins, was added to the dialysis buffer, there was a significant loss of binding. These results suggest a reversible binding between TMP metabolite and the protein fraction containing α_{2u} -g (Lock et al., 1987a). The reversibility of the chemical binding with α_{2u} -g, whether parent compound or metabolite, has been confirmed with isophorone (Strasser et al., 1988), 1,4-DCB (Charbonneau et al., 1989), and d-limonene (Lehman-McKeeman et al., 1989).

In the d-limonene study (Lehman-McKeeman et al., 1989), the amount of radioactivity observed in the kidneys of Sprague Dawley rats 24 hours after oral administration of [14-C]d-limonene was

about 2.5 times higher in the males than in the females. Equilibrium dialysis in the presence or absence of sodium dodecyl sulfate indicated that approximately 40% of the radioactive material retained in the male rat kidney was associated with proteins in a reversible manner. Gel filtration high performance liquid chromatography (HPLC), reverse phase HPLC, and amino acid sequencing demonstrated that this radioactive material was associated with α_2 -g. No d-limonene or d-limonene metabolite was seen to coelute with female rat kidney proteins.

Reversible binding generally implies a dissociable chemical-protein interaction in which the free chemical can be liberated from the protein without having produced molecular damage. In contrast, in covalent binding a reactive chemical species, usually an electrophile, reacts with nucleophilic centers in target molecules comprising enzymes, other proteins, nucleic acids or lipids. CIGA appear to differ from many known chemical toxins, nephrotoxins included, which bind covalently and irreversibly to proteins and/or DNA and through this process cause cellular injury.

A DNA binding study with F344 rats and B6C3F1 mice of both sexes was performed using [1,3,5,- 14 C]-isophorone (Thier, et al., 1990). Twenty-four hours after the animals were administered a 500 mg dose by gavage, liver and kidneys were processed for determination of DNA binding. Neither isophorone nor its metabolites showed covalent binding to DNA. In addition, metabolically formed degradation products were not incorporated into the DNA by de novo synthesis of DNA from labelled fragments

of the xenobiotic.

The non-CIGA, 1,2-dichlorobenzene (1,2-DCB), unlike its closely related isomer, 1,4-DCB, binds to α_{2u} -g and other proteins in the kidney cytosol without inducing an increase in hyaline droplets. The binding of 1,2-DCB to α_{2u} -g was less reversible than it was for the hyaline droplet inducer, 1,4-DCB. This finding is consistent with the more severe and different nephrotoxicity seen for 1,2-DCB compared with 1,4-DCB these two compounds (NTP, 1987a; Charbonneau et al., 1989).

Gas chromatographic analysis in experiments with liver microsomes have shown that mice are able to oxidize d-limonene to cis-d-limonene-1,2 oxide, as in the rat, although some quantitative and qualitative species differences were noted (Lehman-McKeeman, 1990b). However, equilibrium saturation binding studies then demonstrated a lack of any interaction between d-limonene or its metabolites and MUP in male or female mice (Lehman-McKeeman, 1990b; Caudill et al., 1990). These results add further support to the specificity of the interaction between CIGA and α_{2u} -g.

The capacity of CIGA for association with other low-molecular-weight proteins, some of which are found in humans, that share some homology with α_{2u} -g has been investigated in in vitro assays. For example, the alcohol metabolite of TMP, TMPOH, which binds reversibly to α_{2u} -g in vitro, also binds reversibly to three other members of the superfamily, ie., retinol-binding protein, α_1 -acid glycoprotein, and β -lactoglobulin (Borghoff et al., 1988). It did not bind to the β_2 -microglobulin or lysozyme, low-molecular-weight

proteins that are not members of the superfamily.

When [H3]-retinol was administered to male rats, retinol-derived radioactivity coeluted with the protein fraction in cytosol containing α_{2u} -g. However, retinol did not produce hyaline droplet or α_{2u} -g accumulation (Borghoff et al., 1989). In vitro studies on the binding affinities of retinol and several CIGA for α_{2u} -g show that retinol can compete with CIGA for binding to α_{2u} -g (Borghoff et al., 1991). These studies suggest that hyaline droplet accumulation may not depend on how strongly a chemical binds to α_{2u} -g, but on whether the chemical causes a conformational change in the protein (Borghoff et al., 1990) which inhibits protein catabolism.

Binding affinities measured in in vitro studies generally have not correlated well with the efficacy of chemicals for causing hyaline droplet accumulation. Other factors affecting the development of hyaline droplet accumulation are the protein concentration in the tubule lumen, the rate of breakdown of the protein-hydrocarbon complexes in the tubule cells, the death of cells resulting from abnormal accumulation of hyaline droplets, and the subsequent appearance of cell debris in the lumen of tubule cells. These factors are discussed in the following sections.

H. Catabolism of alpha-2u-globulin complexed with CIGA

Reduced renal lysosomal catabolism of the CIGA- α_{2u} -g complex leads to its accumulation in the cells of the proximal renal tubule, causing lysosomal protein overload and individual cell death (Swenberg et al., 1989a). Figure 3 illustrates this proposed




FIGURE 3: Schematic representation of the uptake and fate of alpha-2u-globulin complexed with a CIGA in hydrocarbon nephrotoxicity.

sequence of events.

Lysosomal degradation of α_{2u} -g bound to CIGA has been studied by measuring the digestion rate of the protein recovered from treated male rat kidney (Charbonneau et al., 1988) or of purified urine-derived protein conjugated with CIGA in vitro (Lehman-McKeeman et al, 1990a). Charbonneau et al. (1988) found that both a mixture of standard protease enzymes of non-rat origin or proteinase K digested α_{2u} -g from rats treated with TMP at a much slower rate than α_{2u} -g from untreated rats.

Using an in vitro incubation system with renal cortex lysosomes prepared from male rats, Lehman-McKeeman et al. (1990b) demonstrated that the reversible binding of three other CIGA or their metabolites impaired the degradation of α_{2u} -g by one-third. Under the experimental conditions employed, this was equivalent to an extension of the apparent half-life of α_{2u} -g from 6.67 to 10 hours. The study is particularly interesting in that it shows reversible binding of a CIGA to α_{2u} -g does not necessarily alter the rate of protein degradation but that this may be a function of a metabolite. Thus, d-limonene and 1,4-DCB did not impair hydrolysis of α_{2u} -g but their respective bound metabolites, d-limonene-1,2-oxide and 2,5-dichlorophenol, did. With isophorone, however, it was the parent compound alone which produced the effect. This apparent need to reduce protein degradation might offer an explanation to describe why chemicals, such as retinol, have been shown to bind to α_{2u} -g, without producing hyaline droplet accumulation.

Other evidence (Olson et al, 1988) shows that administration to male rats of leupeptin (an inhibitor of the lysosomal peptidase cathepsin B), causes a rapid α_{2u} -g accumulation in the kidney, indistinguishable from that induced by TMP and gasoline. These various observations provide evidence that CIGA-induced hyaline droplet accumulation may result from a reduced protein degradation rate either by 1) making the protein harder to digest or 2) inhibiting enzymatic components of the proteolytic process. Studies by Charbonneau et al. (1988) and Lehman-McKeeman et al. (1990a) support the former by indicating that the TMP metabolite-protein complex is more resistant to hydrolysis than free α_{2u} -g. Furthermore, Murty et al. (1988) found that unleaded gasoline was not associated with a reduction, but rather an increase, in rat kidney lysosomal proteolytic enzyme activity.

I. Structure-Activity Relationships for CIGA.

An ability to predict those chemicals that will induce accumulation of α_{2u} -g in the male rat through structural relationships would be clearly advantageous. The fact that relatively minor metabolites such as d-limonene 1,2-epoxide can account for the majority of the association with α_{2u} -g, however, restricts the present utility of structure activity calculations as a predictive tool. Nevertheless, some associations have been observed. Lehman-McKeeman et al. (1990a) noted that retarded degradation of α_{2u} -g correlates with the presence on the active CIGA or metabolite of an oxygen function of one type or another, i.e., a hydroxyl group for TMPOH and 2,5-dichlorophenol, an epoxide

for d-limonene-1,2-oxide, and a ketone function for isophorone.

Another recent study employed a quantitative approach to determine the structural features necessary to induce excessive hyaline droplet activity in male rats (Bomhard et al., 1990). Based on data for a number of light hydrocarbons, Bomhard et al. surmised that an n-octanol water partition coefficient above 3.5 and the presence of an isopentyl structural moiety are associated with increased hyaline droplet formation in male rats. A binding site model for aliphatics was derived from this information. The model was then generalized to include cycloaliphatics by substituting the requirement for an isopentyl structure with a requirement for the presence of at least one tertiary carbon atom. Using this binding site model, Bomhard et al. predicted the hyaline-droplet inducing activity of 18 previously untested hydrocarbons. These chemicals were then tested for ability to induce hyaline droplet accumulation in adult male Wistar rats. Although the binding site model was based on the structure of the parent compound and did not allow for active metabolites, the results in the rats were described as being in good agreement with the predictions.

Borghoff et al. (1991) determined the apparent binding affinity to α_{2u} -g for a number of chemicals associated with α_{2u} -g-nephropathy and measured their ability to compete with TMPOH. Using molecular modeling and information on the most active compounds, these investigators concluded that the presence of an electronegative atom for hydrogen bonding is a critical factor in

determining binding affinity. Lipophilicity also seemed crucial for hydrophobic interactions, but the presence of an electronegative atom was necessary for greater activity. Steric volume was also considered to play an essential role in binding activity.

The conclusions of Borghoff et al. (1991) are consistent with the notion that α_{2u} -g is capable of transporting lipophilic compounds within a binding site pocket of specific dimensions. Since binding affinity does not correlate well with hyaline droplet formation, however, the ability of this structural feature to serve as a predictive tool would appear limited.

III. ALPHA-2U-GLOBULIN NEPHROPATHY

Substances reported to induce increased formation of hyaline droplets in proximal tubule cells of male rats are listed in Appendix 1, along with available information on whether the accumulating protein is α_{2u} -g. The nephrotoxicity that can ensue from hyaline droplet accumulation is novel because it is associated with excessive α_{2u} -g accumulation. This α_{2u} -g accumulation is believed to initiate a sequence of events resulting in chronic proliferation of tubule epithelium, as well as an exacerbation of CPN. Because α_{2u} -g is a male rat-specific protein, nephropathy induced by accumulation of α_{2u} -g would not be expected to occur in female rats, mice of either sex, or other species.

The proposed sequence of histopathological changes is based mainly on research studies with four model substances, unleaded gasoline and TMP (Short et al., 1986; 1987; 1989a), decalin (Alden

et al., 1984; Kanerva et al., 1987a,b,c; Stone et al., 1987) and d-limonene (Kanerva et al., 1987b; Webb et al., 1989). For even these four substances, not all of the individual lesions in the proposed progression have been shown to belong to a sequence of interrelated events. Specific information pertaining to lesion nature and sequence is lacking for many of the hyaline-droplet inducers listed in Appendix 1.

Much of the information useful for defining the pathologic sequelae to α_{2u} -g accumulation does not require chronic exposure. Accumulation of α_{2u} -g is visible within a matter of days and the response to chronic administration of CIGA might even diminish since α_{2u} -g levels decline in aging male rats (Murty et al., 1988). The nephrotoxicity associated with α_{2u} -g accumulation might also be influenced by age. Certainly, the age-related progression of CPN obscures the lesions directly related to CIGA administration, making evaluation of the chronic sequence of lesions especially difficult.

A. Pathologic features of alpha-2u-globulin nephropathy

Renal lesions that have been associated with α_{2u} -g nephropathy are listed in Table 3. The first morphological manifestation of α_{2u} -g nephropathy is the rapid accumulation of hyaline droplets in proximal tubule cells, developing within the first 24 hours after dosing (Webb et al., 1989).

The droplets stain positively with Mallory's Heidenhain stain but are negative for periodic acid Schiff, indicating their protein composition (Alden et al., 1984). Mallory's Heidenhain stain is

**TABLE 3. SUMMARY OF THE HISTOPATHOLOGY AND LESION PROGRESSION
REPORTED IN ALPHA-2U-GLOBULIN-ASSOCIATED NEPHROTOXICITY**

1. Excessive accumulation of hyaline droplets in the P2 segment of the proximal tubule region of kidney occurs after 1 or 2 days. This is reversible within 3 days to 2 weeks after exposure ceases.
2. Evidence of single cell necrosis in P2 segment epithelium and exfoliation after 5 days of continuous exposure.
3. Accumulation of granular casts formed from the cellular debris and subsequent tubule dilation, at the junction of the P3 segment and the thin loop of Henle, following 20 to 40 days of continuous exposure. Granular casts have been observed at 3 to 13 weeks after commencing exposure and sometimes beyond, up to two years. *
4. Increase in cell proliferation within the P2 segment following 3 weeks of continuous exposure, remaining elevated above normal at 48 weeks of exposure.
5. Linear mineralization of tubules within the renal papilla, appearing between 3 and 12 months after a 28-day exposure, and sometimes observed at the end of a two year study. *
6. Hyperplasia of the renal pelvic urothelium observed around 1 year. *
7. Exacerbation of the spontaneous chronic progressive nephropathy syndrome common in aging rats. *
8. Formation of occasional hyperplastic foci within cortical epithelium at chronic time-points.

*Indirect consequence of progression of lesions.

therefore more useful than conventional hematoxylin and eosin for visualizing and quantitating the droplets. As they represent lysosome-derived entities, the droplets are strongly autofluorescent (yellow) in paraffin sections under ultraviolet illumination (unpublished observations, G.C. Hard). In plastic-embedded tissue, hyaline droplets can be visualized easily with Lee's methylene blue basic fuschin (Short et al., 1986).

Excessive hyaline droplet formation occurs primarily in cells of the P2 segment, but small increases in the number of hyaline droplets may also be seen in P1 and P3 (Short et al., 1987). By light microscopic immunohistochemistry, α_{2u} -g has been clearly and specifically localized to the hyaline droplets within proximal tubules (Burnett et al., 1989). Ultrastructurally, the hyaline droplets are enlarged secondary lysosomes partially composed of α_{2u} -g (Garg et al., 1989a). Many are polyangular or irregular in shape, containing a condensed crystalline core suggestive of aggregated protein in pure form. Although the α_{2u} -g-associated hyaline droplet accumulation persists during chronic exposure, the severity becomes less with increasing duration of exposure beyond about three weeks (Short et al., 1989a). This apparent waning of the response with continued exposure could be related to declining α_{2u} -g production by the male rat beginning at some stage after 100-150 days of age (Roy et al., 1983; Motwani et al., 1984).

With continued exposure, the initial accumulation of α_{2u} -g-containing hyaline droplets may be followed by a sequence of interrelated pathological events. (1) Scattered single cell

necrosis occurs predominantly in the P2 segment cells (Short et al., 1987) with subsequent exfoliation of these degenerate cells and cell fragments laden with crystalloid phagolysosomes into the tubule lumen. With decalin, a minimal degree of cell degeneration/necrosis was reported to be present in the proximal convoluted tubules after 5 days of exposure, becoming maximal at 19 days, but reverting to the minimal level after 31 days of exposure (Kanerva et al., 1987a). Occasional exfoliation of droplet-affected cells was observed after 48 weeks of exposure to unleaded gasoline or TMP (Short et al., 1989a), indicating sustained single cell loss while exposure to CIGA continues.

(2) Epithelial cell proliferation primarily involving the P2 segment occurs as a regenerative response to cell damage and loss. This can be seen as increased numbers of mitotic figures or demonstrated by labeling techniques for DNA-synthetic activity. Increased proliferative activity has been recorded after only three weeks of petroleum hydrocarbon exposure (Short et al., 1987) but it persisted during 48 weeks of chronic exposure (Short et al., 1989a).

(3) Granular casts composed of sloughed cell debris accumulate at the junction between the P3 segment of the proximal tubule and the descending thin loop of Henle, that is, at the junction between the inner and outer stripes of outer medulla, with consequent tubule dilation at this part of the nephron (Alden et al., 1984). This can occur as early as two to three weeks after initial exposure (Alden et al., 1984; Kanerva et al., 1987a). As

well as comprising recognizable cell debris, the granular casts stain positively for α_{2u} -g (unpublished observations, R.J. Foster, Central Toxicology Laboratory, ICI, Macclesfield) indicating probable derivation of the debris from cells which had accumulated this protein. Granular cast formation appears to be associated with higher doses of compound rather than with the lowest doses that can induce increased hyaline droplet accumulation. An absence of casts after treatment might therefore reflect a dose-related decrease in the severity of cell necrosis and exfoliation (Short et al., 1986, 1987).

(4) At chronic timepoints, linear mineralization develops in the renal papilla, outlining affected medullary tubules, along with hyperplasia of the pelvic urothelium (Alden, 1989). The mineralization appears to form within the loops of Henle and has been identified as calcium hydroxyapatite (Trump et al., 1984b). The relationship between papillary mineralization and the proximal tubule lesion remains undetermined but the medullary lesion is presumed to represent mineralized remnants of debris from disintegrating granular casts that lodge in the prebend segments of Henle's loop (Bruner, 1984; Alden, 1989). In turn, urothelial hyperplasia, which mainly affects the surface of the renal papilla, may be a response of the renal pelvis lining to papillary mineralization (Bruner, 1984; Alden, 1991).

B. Rat urine chemistry and CIGA

Several studies have examined renal function in rats treated with CIGA and subsequently developing α_{2u} -g nephropathy. Two days

of treatment with TMP resulted in mild urinary increase in the lysosomal enzyme N-acetyl- β -glucosaminidase (NAG) and alkaline phosphatase, a decrease in creatinine, and mild increase in urinary cell debris. Other parameters, aspartate aminotransferase (AAT), urine osmolality and volume, were not affected (Fowlie et al., 1987). A single, oral dose of TMP had no effect on renal function (Stonard et al., 1986). In a 14-day study with decalin, of six urinary enzymes tested, only AAT, lactate dehydrogenase and NAG were altered (increases) at days 21 and/or 28 (Evans et al., 1986). Similar results were obtained for levamisole except that AAT remained normal (Evans et al., 1988). During prolonged treatment with C₁₀-C₁₁ isoparaffinic solvent, up to 8 weeks, the only urinary changes observed were mild elevation of glucose and albumin, slightly decreased concentrating power and osmolality, and epithelial cell debris in the urine. There was no alteration in urinary β_2 -microglobulin content (Phillips and Egan, 1984).

Taken together, these studies suggest that CIGA produce minimal changes in urinary chemistry and very little or no glomerular dysfunction or damage in the days following their administration. The minor alterations seen in urine composition following administration of CIGA suggest also that hyaline droplet accumulation is not related to increased passage of serum proteins by the glomerulus. The mild tubule toxicity identified by clinical chemistry is a characteristic of CIGA, which contrasts with the obvious urinary changes associated with the nephrotoxicity induced by such classical renal toxins as mercuric chloride,

hexachlorobutadiene, aminoglycosides and papillotoxic agents (Stonard, 1987).

C. Species variation in the renal response to CIGA

The male-specific effects of hyaline droplet inducers have been demonstrated over a range of rat strains including Fischer 344, Sprague-Dawley, Buffalo and Brown Norway rats (Ridder et al., 1990). Hyaline droplet accumulation or the spectrum of lesions comprising $\alpha_2\mu$ -g nephropathy have not been observed in female rats, or mice of either sex, following treatment with these chemicals (Alden et al., 1984; Swenberg et al., 1989a). In addition to these studies, other hyaline-droplet inducers have been tested for toxicity in hamsters (jet fuels), guinea pigs (decalin), dogs (decalin, jet fuels, d-limonene and methyl isobutyl ketone) and monkeys (gasoline and methyl isobutyl ketone). No renal pathology was demonstrated in these species at doses known to cause nephropathy in male rats (Alden et al., 1984; Kuna and Ulrich, 1984; MacFarland, 1984; MacNaughton and Uddin, 1984; Phillips et al., 1987) except for one report of minor changes in dogs treated for 6 months with d-limonene (Tsuji et al., 1975). In this chronic study, an increased incidence of proteinaceous casts was observed in male and female beagles, but no tubule epithelium changes, tubule lumen dilation or mineralization. However, Webb et al. (1990) were unable to demonstrate any renal pathology in dogs after 6 months of d-limonene treatment at comparable dose-levels.

Knowledge concerning the renal effects of CIGA in humans is hampered by the lack of data on specific chemicals in this

category, and the limitations imposed by a multiplicity of types of occupational and non-occupational exposures. Case studies have reported a link between chronic renal disease with gasoline, solvents, jet and diesel fuels including rare cases of acute tubular necrosis (proximal and distal tubule epithelium) following severe exposure to petroleum distillates (e.g. Barrientos et al., 1977; Crisp et al., 1979). Case reports cannot be used to establish a causal relationship but may serve to initiate formal epidemiologic investigation (Churchill et al., 1983).

Epidemiological studies concerning non-neoplastic kidney disease and occupational exposure to hydrocarbons and solvents have been conducted only since 1975 (Reviewed by Askergren, 1986; Daniell et al., 1988; Phillips et al., 1988). A majority of these studies have indicated an association between glomerulonephritis and exposure to hydrocarbons, especially organic solvents or gasoline. Some have suggested a positive association between the presence of glomerular disease and duration and severity of occupational exposure to hydrocarbon solvents, including tetrachloroethylene which is a CIGA in male rats (Kluwe et al., 1984). However, many of the earlier studies are considered to be methodologically limited (Churchill et al., 1983; Askergren, 1986; Phillips et al., 1988). Their major shortcomings have been heterogeneous case definition, use of inappropriate control groups or non-blinded interviewers, and failure to consider recall bias or to adequately define hydrocarbon exposure (Phillips et al., 1988).

More recently, Steenland et al. (1990), investigating specific

occupational exposures associated with end-stage renal disease in male workers, found elevated risks for solvents used as cleaning agents or degreasers (odds ratio (OR) 2.5; 95% confidence interval (CI) 1.56-3.95) but not for exposure to gasoline and diesel fuel (OR 0.98; 95% CI 0.49-1.06) or motor and fuel oil (OR 1.13; 95% CI 0.69-1.84). Harrington et al. (1989) found no association (OR 1.0; 95% CI 0.16-6.3) between occupational exposure to inorganic solvents and glomerulonephritis, but the authors also concluded that the statistical power of this case-referent study was not sufficient to detect other than large risk estimates.

The glomerulonephritis reported in the positive epidemiologic studies has involved thickening of glomerular basement membranes or deposition of antibodies against glomerular basement membrane, a mild degree of albuminuria, and sometimes tubule atrophy and tubular basement membrane thickening (Kluwe et al., 1984; Phillips et al., 1988).

Levamisole, a drug used as an anthelmintic, in cancer chemotherapy, and in the treatment of rheumatoid arthritis in humans, falls into the CIGA category because it induces both hyaline droplet and α_{2u} -g accumulation in male rats (Read et al., 1988). Based on an absence of elevated levels of urinary NAG in patients receiving 150 mg levamisole per day for 26 weeks, there is little evidence to indicate that this compound is nephrotoxic in humans (Dieppe et al., 1978). Since urinary NAG is only slightly elevated in male rats exposed to CIGA, however, urine chemistry may not be a good biological monitor of the type of nephrotoxicity

associated with CIGA.

In a study of 16 females exposed to tetrachloroethylene from their employment in dry-cleaning shops an average of 11 years (range 1-25 years), Vyskocil et al. (1990) found no evidence of renal damage except for an increase in lysozyme in the urine. Although a high concentration of lysozyme in the urine can be a measure of decreased tubular reabsorption, the authors discounted this explanation because there was no statistically significant increase in urinary excretion of β_2 -microglobulin, lactate dehydrogenase, or glucose, other markers of tubular dysfunction.

The evidence regarding renal injury in humans from chronic organic chemical exposure is inadequate to demonstrate whether or not CIGA exposure can affect the human renal tubule cell. Existing reports imply that, if the association is real, it is the glomerulus that is pathologically involved. However, this may simply reflect study designs which concentrated on detection of glomerular effects. Since the injury to the rat tubule cells is relatively mild, insensitive tests, such as urine chemistry, which are generally used for evaluating humans might be inadequate to detect changes.

D. Factors affecting the expression of alpha-2u-globulin nephropathy

Various conditions, including age, hormone manipulation and genetics, have the potential for altering the expression of CIGA-induced α_{2u} -g nephropathy. Experimental studies have investigated the influence of these factors on CIGA nephrotoxicity as well as determining the effects of α_{2u} -g in female rats.

1. Age-related effects

As discussed earlier, the hepatic synthesis and urinary excretion of α_{2u} -g in the male rat are highly age-dependent, with prepubertal and aged animals showing negligible amounts of this protein (Neuhaus and Flory, 1978; Roy et al., 1983). Accordingly, administration of either decalin to immature male rats (Alden et al., 1984) or unleaded gasoline to aged, 26 month old, male rats (Murty et al., 1988) failed to produce renal cortical α_{2u} -g accumulation or an increase in hyaline droplets.

2. Effect of hormone manipulation

As α_{2u} -g synthesis is primarily under androgenic control, the effects of castration, which depresses hepatic synthesis of α_{2u} -g (Roy and Neuhaus, 1967), were explored by Hobson et al. (1986) using TMP. Although a significant increase in hyaline droplet formation was observed in both castrated and uncastrated male F344 rats exposed to a single oral dose of TMP, the severity of the lesion was less in the former. Thus, castration diminished but did not abolish the TMP-induced nephrotoxicity.

Estrogen is known to inhibit the hepatic synthesis of α_{2u} -g in the rat (Roy et al., 1975). Garg and coworkers (1988) used estradiol administration to study the influence of inhibition of new synthesis of α_{2u} -g on recovery from CIGA-induced renal tubule changes. Commencing treatment on the ninth and final day of unleaded gasoline exposure, estradiol reduced renal cortical α_{2u} -g content by 25%, 41% and 52% on post-exposure days 3, 6, and 9 respectively, compared to rats receiving no hormone treatment. At

the same time, hyaline droplet removal appeared to be accelerated in rats treated conjointly with hormone. Hyaline droplet number and size (qualitative observations) in hormone-treated rats approached control levels at 3 days post-exposure, compared with up to 9 days for complete resolution in unleaded gasoline-exposed rats not receiving estradiol.

In a subsequent study, Garg et al. (1989b) demonstrated that pretreatment of mature male rats with subcutaneous injections of estradiol for 10 days before gasoline exposure completely inhibited the renal accumulation of α_{2u} -g and hyaline droplets normally induced by gasoline.

3. Genetic variants

The NBR rat is a strain that appears to have a tissue- and gene-specific regulatory defect involving α_{2u} -g. This rat has no detectable levels of hepatic α_{2u} -g mRNA in either sex and, therefore, is unable to synthesize α_{2u} -g in the liver although high constitutive levels of the mRNA are present in the preputial gland (Chatterjee et al., 1989). Under exposure conditions that produce α_{2u} -g nephropathy in Fischer 344 rats, d-limonene, TMP, isophorone, and 1,4-DCB did not induce any detectable α_{2u} -g accumulation, hyaline droplets or other lesions in the male NBR rat (Dietrich and Swenberg, 1990a). Identical results were obtained for decalin (Ridder et al., 1990) and lindane (Dietrich and Swenberg, 1990b).

4. Alpha-2u-globulin infusion in female rats

Ridder et al. (1990) intraperitoneally administered α_{2u} -g (purified from mature male rat urine) at hourly intervals to

decalin-treated female Sprague-Dawley rats for a total of 8 injections and examined kidney samples for hyaline droplets and α_{2u} -g one hour after the last protein injection (9 hours after decalin treatment). Although droplet formation was not evident in kidney sections from the α_{2u} -g-infused female rats stained with Mallory's Heidenhain, hyaline droplet and α_{2u} -g accumulation were clearly demonstrated in females exposed to both hydrocarbon and male urinary protein. By means of two-dimensional gel electrophoresis, the investigators showed slight, but apparent, renal cortical accumulation of α_{2u} -g in the infused females. Accumulation of the protein greatly increased in females that were both infused with α_{2u} -g and decalin-treated.

These various studies indicate a direct dependence of CIGA-induced renal lesion expression on the presence of α_{2u} -g.

E. Chronic progressive nephropathy

Rats are particularly predisposed to an age-related spontaneous nephropathy, CPN, that is more severe in males than in females and that affects certain strains more than others. CPN is more common in Sprague-Dawley and Fischer 344 rats than the Wistar strain (Gray, 1986) and it is also common in the Osborne-Mendel rat (Goodman et al., 1980). The etiology of CPN is not known but the severity of the syndrome is influenced by a number of factors, particularly dietary manipulation affecting protein content, or caloric intake (Masoro and Yu, 1989).

Exacerbated CPN, involving enhanced severity and earlier onset of the disease, is generally observed after chronic administration

of CIGA to male rats (Trump et al., 1984b). It has been stated that exacerbated CPN is one component (together with hyaline droplet accumulation and granular cast formation) of a triad of lesions that specifies the nephropathic response to CIGA (Kanerva et al., 1987a; Webb et al., 1989). Exacerbated CPN is usually recognized after months of continuous treatment (Trump et al., 1984b; Short et al., 1989a) although Alden et al. (1984) reported early signs after 2-3 weeks with decalin. These authors (Alden et al., 1984) consider that exacerbated CPN develops as a tertiary response to nephron obstruction caused by the CIGA-induced granular casts.

The pathologic features of CPN (listed in Table 4) include certain lesions that are also found in α_{2u} -g nephropathy, as well as lesions that are distinctive. Single cell necrosis, regenerating basophilic tubules and focal hyperplasia of proximal tubule epithelium are common to spontaneous CPN and to α_{2u} -g nephropathy (UAREP, 1983). CPN is characterized by certain lesions which are not components of α_{2u} -g nephropathy, including conspicuous thickening of tubule and glomerular basement membranes, hyaline casts consisting of homogeneous, proteinaceous material (distinct from granular casts containing cellular debris), interstitial mononuclear cell infiltration, fibrosis, tubule atrophy and sclerotic glomeruli. Conversely, early and late stages of α_{2u} -g nephropathy exhibit a number of characteristics unlike CPN, such as hyaline droplet accumulation associated with α_{2u} -g in the P2 segment, granular casts at the corticomedullary

TABLE 4. SUMMMARY OF THE HISTOPATHOLOGY OF SPONTANEOUS
CHRONIC PROGRESSIVE NEPHROPATHY OF AGING RATS

1. Thickening of tubular and glomerular basement membranes.
2. Basophilic segments of proximal tubules with sporadic mitoses indicative of tubule cell proliferation.
3. Tubular hyaline casts of proteinaceous material originating in the more distal portion of the nephron, mainly in the medulla, and later plugging a considerable length of the tubule.
4. Focal interstitial aggregations of mononuclear inflammatory cells within areas of affected tubules.
5. Glomerular hyalinization and sclerosis.
6. Interstitial fibrosis and scarring.
7. Tubular atrophy involving segments of proximal tubule.
8. Chronically in advanced cases, occasional hyperplastic foci in affected tubules.
9. In some advanced cases, accumulation of protein droplets in sporadic proximal tubules.

junction, and linear mineralization in the papilla (Trump et al., 1984b). In very advanced cases of spontaneous CPN, sporadic tubules may contain excessive numbers of hyaline droplets similar in appearance to those induced by CIGA. However, these do not show immunochemical evidence of α_{2u} -g (unpublished observations, G.C. Hard). The urine and serum chemistry of advanced CPN also differs from α_{2u} -g nephropathy. Albuminuria, hypoalbuminemia, and hypocholesterolemia typify CPN, with increases in serum creatinine and urea nitrogen levels in end-stage disease (Barthold, 1979).

F. Renal toxicity observed in chronic bioassays of chemicals that induced kidney tumors in rats

For the purpose of the current review, bioassays were identified and the data examined on seven chemicals tested for chronic toxicity and carcinogenicity by the National Toxicology Program (NTP) or the National Cancer Institute (NCI). All seven produced accumulation of hyaline droplets, nephropathy, and kidney tumors in male rats. These model compounds are d-limonene, dimethyl methylphosphonate, hexachloroethane, 1,4-DCB, tetrachloroethylene, pentachloroethane, and isophorone¹. Information on unleaded gasoline (tested at International Research and Development Corporation (IRDC) for the American Petroleum Institute), which is a mixture regarded as a CIGA, was also

¹Several of these seven chemicals cannot be described as true "CIGA carcinogens" since the accumulating protein in the hyaline droplets has not been confirmed to be α_{2u} -g. They are occasionally described as "potential CIGA" or "potential CIGA carcinogens" for purposes of developing the discussion on cancer. This should not be construed to mean that all seven chemicals fit the Policy Statement developed in Part IV of this document.

examined. The two non-CIGA, trichloroethylene and chlorothalonil are included for comparative purposes. Although extensive acute and subchronic studies have been performed on two other chemicals (decalin and TMP), both of which cause the sequence of nephropathy in male rat kidney beginning with α_{2u} -g accumulation, carcinogenicity bioassay data are not available for these compounds.

Trichloroethylene, which was tested by NTP, induces kidney tumors in male rats only (NTP, 1988a) but does not cause an accumulation of hyaline droplets or an increase in α_{2u} -g levels (Goldsworthy et al, 1988a). There is also some evidence that trichloroethylene metabolites bind covalently to renal macromolecules (Bruckner et al, 1989). Consequently, this compound is not considered to be a CIGA.

Chlorothalonil, a fungicide tested on separate occasions by industry and a government agency, induced renal tubule tumors in male and female rats and in male mice (NCI, 1978a). It also induced hyaline droplet accumulation in proximal convoluted tubules of male rats (USEPA, 1988), but these may not become apparent during the first few weeks of treatment (Killeen et al, 1990). Electron microscopic studies of male rat kidney following subchronic chlorothalonil exposure revealed angular membrane-bound lysosomes containing crystalline structures similar to those observed in α_{2u} -g nephropathy (personal communication, William M. Busey and James C. Killeen). However, α_{2u} -g has not been detected in the renal tubules of chlorothalonil-exposed rats (Swenberg,

1989a). The progression of chlorothalonil nephrotoxicity involves initially, vacuolar degeneration of proximal tubule epithelium followed 4 weeks later by tubule cell hypertrophy, hyperplasia, and tubule dilation (Killeen et al, 1990). Therefore, this compound appears not to produce the same spectrum or sequence of lesions induced by CIGA. Furthermore, chlorothalonil has been shown to interact with cellular macromolecules including histones and thiol proteins, possibly through covalent binding of a metabolite with sulfhydryl groups (Rosanoff and Siegel, 1981). Chlorothalonil also induces overt renal dysfunction in both sexes of rats. At doses from 40 mg/kg/day, blood urea nitrogen and creatinine were increased while circulating levels of glucose and albumin were decreased (U.S. EPA, 1988). For these various reasons, chlorothalonil is not considered a member of the CIGA class.

A summary of the non-neoplastic and preneoplastic kidney effects observed in male rats after administration of the ten selected chemicals is presented in Table 5. Non-neoplastic and preneoplastic lesions reported in female rats and mice of both sexes are summarized in Table 6. The data in these two Tables were extracted from the NTP Technical Reports (see Appendix 2) and other relevant literature.

TABLE 5. SUMMARY OF DATA ON NON-NEOPLASTIC AND PRENEOPLASTIC KIDNEY LESIONS IN MALE RATS ASSOCIATED WITH EIGHT MODEL COMPOUNDS THAT INDUCED RENAL TUMOURS IN 2-YEAR BIOASSAYS

CHEMICAL	<u>Toxic Nephropathy</u>			Cast Formation	<u>Mineralization^a</u>			Karyomegaly	<u>Hyperplasia</u>	
	Hyaline Droplets	Dose-Response	Increased Severity		Present	Dose-Response	Present		Present	Dose-Response
d-limonene	+	+	+	+	+	+	+	N.R.	+	+
Dimethyl methyl-phosphonate	+	+	+	+	+	+	+	N.R.	+	+
Pentachloroethane	+	+	N.R.	+	+	+	+	N.R.	+	N.R.
Isophorone	+	- (slight)	+	+	N.R.	N.R.	+	N.R.	+	+
1,4-Dichlorobenzene	+	+	+	+	+	+	+	N.R.	+	+
Tetrachloroethylene	+	+	N.R.	+	N.R.	N.R.	+	+	+	+
Hexachloroethane	+	+	+	+	+	+	+	N.R.	+	+
Unleaded gasoline	+	+	+	+	+	+	^b	+/-	^b	N.R.

+ Positive
- Negative

N.R. = Not reported

a, localized to renal papilla

b, data from research studies with unleaded gasoline.

TABLE 6. SUMMARY OF DATA FROM 2-YEAR BIOASSAYS ON NON-NEOPLASTIC AND PRE-NEOPLASTIC KIDNEY LESIONS IN MICE AND FEMALE RATS EXPOSED TO EIGHT MODEL COMPOUNDS THAT INDUCED RENAL TUMORS IN MALE RATS

CHEMICAL	Hyaline Droplets	Toxic Nephropathy	Cast Formation	Mineralization	Karyomegaly	Hyperplasia
d-limonene	-	-	-	-	N.R.	-
Dimethyl methyl- phosphonate	-	-	-	-	N.R.	-
Pentachloroethane	-	-	-	-	N.R.	-
Isophorone	-	-	-	-	N.R.	-
1,4-Dichlorobenzene	-	+ (male mice and female rats)	-	+ (female rats)a	N.R.	-
Tetrachloroethylene	-	N.R.	+ (mice)	-	+ (female rats and mice)b	+ (male mice)b
Hexachloroethane	-	+ (female rats and mice)	+ (mice - hyaline)	+ mice (Ca deposition)	N.R.	-
Unleaded Gasoline	-	-	-	-	+	-

a, located to renal papilla

b, data from research studies

In male rats, renal tubule cell hyperplasia was reported in the 2-year bioassays for the 7 renal carcinogens tested by NTP. Although not reported in the bioassay for unleaded gasoline, this lesion was observed in later research studies with the mixture (Short et al., 1989b). None of the eight bioassayed chemicals produced tubule cell hyperplasia in female rats, although this lesion was reported in male mice exposed to tetrachloroethylene. In male rats, renal changes described as "toxic tubular nephropathy" (encompassing degeneration of tubule epithelium, necrosis, epithelial cell regeneration, and cast formation) were seen following administration of all 8 of the renal carcinogens (Table 6). Some aspect of toxic tubular nephropathy was also observed in female rats or mice administered hexachloroethane, 1,4-DCB, or tetrachloroethylene (Table 6). For example, calcium deposition or mineralization was seen after administration of hexachloroethane to mice or 1,4-DCB to female rats. Cast formation was reported in mice following administration of hexachloroethane and tetrachloroethylene.

Several difficulties arise in the interpretation and utilization of the bioassay-derived data when mouse and female rat lesions are considered. The nature of casts (granular vs. hyaline) is not always described, and for mineral deposits, the site (papillary vs. corticomedullary) and form (linear vs. globular) may not be specified. The range of lesions encompassed by the term "toxic nephropathy" is not always defined, and there is sometimes no clear distinction from CPN. Nevertheless, it appears from the

data that female rats and mice do not develop as broad a spectrum of nephrotoxic lesions as those proposed to be associated with α_{2u} -g nephropathy and renal tumor formation in the male rat. Furthermore, where nephrotoxicity was reported in both male and female rats, the males had more lesions and the female response never demonstrated the characteristics seen in the male response to CIGA. Therefore, the lesions caused by CIGA seem to be both qualitatively and quantitatively different for male rats compared to mice and female rats.

PART 2. CARCINOGENICITY

The second major part of this document describes information from NTP (or NCI) assays for renal neoplasia induced by chemicals that produced hyaline droplets and/or accumulation of $\alpha_2\mu$ -g and compares and contrasts this information with the kidney tumors induced by classical renal carcinogens. In addition, other information, such as mutagenic activity and tumor-promoting ability, which help to define a CIGA carcinogen or point to possible mechanism of action, are evaluated.

Epidemiological studies of human renal cell cancer are reviewed for consistency with the hypothesis that CIGA-induced renal cancer in male rats is an inappropriate endpoint for assessing human risk. Implicit in this evaluation is a presumption of male rat-to-human tumor site concordance, a supposition EPA generally does not make. In this special case, however, the hypothesized mechanism being examined depends on the accumulation of low-molecular-weight protein in the renal tubule, regardless of species. Hence, the predicted target site in humans, as in the rats, would be the renal tubule.

IV. PATHOLOGIC FEATURES OF RENAL CARCINOGENESIS INDUCED BY CLASSICAL CARCINOGENS

Among the many chemicals recognized as inducers of rodent cancer, several have been used as model kidney carcinogens for studying the pathogenesis of renal tubule tumors in rats. These are dimethylnitrosamine (DMN), diethylnitrosamine (DEN), N-nitrosomorpholine, N-ethyl-N-hydroxyethylnitrosamine (EHEN), lead

acetate, N-(4'-fluoro-4-biphenyl)acetamide (FBPA), and aflatoxin B₁ (Hard, 1990). In the mouse, certain nitrosamines, streptozotocin and ochratoxin A are strong inducers of renal tubule tumors, while the classical renal carcinogen in hamsters is diethylstilbestrol (Hard, 1987). In general, these prototypic renal carcinogens are active in both males and females.

Studies on the pathogenesis of renal tubule tumor formation using model carcinogens in rats demonstrate that a continuum of chemically-induced steps leads from atypical hyperplasia in tubules (also termed hyperplastic tubules), to tubule dysplasia and atypical cell foci through microscopic adenomas, to macroscopic adenocarcinomas or carcinomas (Hard, 1987; Lipsky and Trump, 1988).

In addition there are invariably pathologic changes which precede the proliferative sequence of preneoplastic and neoplastic lesions including a period of early nephrotoxicity and, often, karyomegaly. These various lesions are described below in chronological sequence.

A. Early nephrotoxicity

Acute toxic changes occur in the proximal tubules shortly after the administration of classical renal carcinogens. They include mild lipid droplet accumulation and scattered single cell necrosis (Hard, 1987). Depending on the carcinogen used, this early damage can be observed in different segments of the renal tubule. For instance, with DMN it is localized to the P2 segment (Hard et al, 1984) and with FBPA, to the P3 segment (Dees et al, 1980a,b).

Detailed histological and/or ultrastructural observation shows that hyaline droplet accumulation is not induced by DMN (Hard and Butler, 1971; Hard et al, 1984) or DEN (G.C. Hard, unpublished observations); nor has it been described in studies using other carcinogens, such as FBPA (Dees et al, 1980a,b), as models for renal carcinogenesis.

More is known about DMN than other classical renal carcinogens concerning molecular interactions during the time that acute toxic changes are seen in the proximal tubules. DNA adduct formation in rat renal tissue occurs rapidly following a single administration of DMN. O^6 -Methylguanine formed in the renal cortex (Fan et al, 1989) persists at least 4 days post-injection (Nicolli et al, 1975), which is consistent with the notion that methylation of the O^6 position of guanine in DNA is the most likely initiating event (Pegg, 1984).

B. Karyomegaly

Conspicuous nuclear enlargement, indicative of increased chromosome number without completion of mitosis (Jackson, 1974), may occur in scattered proximal tubule cells during the weeks preceding development of carcinogen-induced proliferative foci. Although karyomegaly is produced by many, but perhaps not all renal carcinogens, there is no evidence that these cells participate in the initial formation of proliferative foci. Hence karyomegaly is not regarded as a preneoplastic lesion (Dees et al, 1980a; Hard, 1987; Lipsky and Trump, 1988).

C. Tubule cell hyperplasia

Tubule cell hyperplasia leads to the appearance of tubules

with proliferating epithelium, usually multilayered, that partially or completely fills the tubular lumen. Although luminal dilation may be pronounced (sometimes to cystic proportions), the structure of the individual tubule remains intact with a confluent basal lamina. Affected cells may be eosinophilic, basophilic or pale-staining and often with vesicular nuclei and prominent nucleoli. Mitotic figures are variable. As a preneoplastic lesion, the hyperplastic tubule is usually associated with some degree of cellular atypia (dysplasia) in the form of cell pleomorphism and increased nuclear to cytoplasmic area ratio (Hard, 1987; Lipsky and Trump, 1988). Preneoplastic tubule hyperplasia is generally considered to be distinguishable from the background tubular regeneration that is a component of spontaneous CPN (Lipsky and Trump, 1988; NTP, 1988a).

D. Adenoma

Adenomas are small neoplastic foci representing epithelial cell proliferation beyond the well-defined structure of individual tubules. These lesions are solid or cystic in form and the cellular morphology and architectural appearance is similar to that of adenocarcinomas, which are described below, particularly the well-differentiated variants. Whereas adenomas and hyperplastic tubules can be differentiated on the basis of finite structure, the distinction between adenomas and adenocarcinomas/carcinomas is an arbitrary one based on size. Neoplasms in the rat kidney parenchyma less than approximately 0.5 cm tend to lack significant vascularization, hemorrhage and degeneration, although there may be

single cell necrosis, mitosis and cell pleomorphism (Hard, 1990).

E. Adenocarcinomas and carcinomas

Renal tubule tumors comprise histological variants based on staining characteristics and architectural organization. In the rat, renal tubule tumors consist mainly of lightly basophilic, granular and/or clear cells organized in tubular, lobular, solid or papillary patterns. Glandular differentiation as opposed to solid sheets of cells distinguishes adenocarcinomas from carcinomas. Increased cellular pleomorphism tends to correlate with a decreasing degree of tubular differentiation and anaplastic variants occur occasionally.

Cells within adenocarcinomas maintain many of the light and electron microscopic characteristics of proximal tubule epithelium, in particular, microvilli resembling brush border, basement membrane and cytoplasmic vesicles. Brush border may occur inappropriately between adjacent cells, along any cell border, or as intracellular profiles. Adenocarcinomas/carcinomas are well vascularized and usually display areas of hemorrhage and degeneration (UAREP, 1983; Lipsky and Trump, 1988; Hard, 1990).

F. Tumor progression

Renal tubule tumors of the rat are slowly growing neoplasms usually requiring about 40 weeks to become clinically palpable in most experimental systems (Hard, 1987). They can grow to large dimensions, several centimeters in diameter.

Unlike their spontaneously occurring human counterparts, renal tubule tumors induced in rats by chemical carcinogens metastasize

infrequently (Lipsky and Trump, 1988). However, effective life-span in chronic-exposure regimens may be a limiting factor. Single-dose studies with DMN, which maximize the life-span following tumor initiation, have demonstrated a link between survival period, tumor size, and incidence of metastasis in renal carcinogenesis (Hard, 1984). For example, rats that survived at least 1.5 years after dosing with DMN showed a high rate of metastasis, approximately 50%, whenever epithelial tumor dimensions exceeded 2.4 cm. These data confirm the malignant potential of renal tubule tumors induced in the rat by a classical carcinogen.

G. Site of origin of renal tubule tumors

The precise location within the nephron from which experimental renal tubule tumors arise varies with the carcinogen, and correlates with the site of the induced early nephrotoxicity. Thus, the P3 segment is the site of origin for FBPA-induced tumors (Dees et al, 1980a,b), while DMN tumors arise from the convoluted segments of proximal tubules, probably P2 (Hard, 1990). Lead-acetate and DEN-induced tumors appear to originate in both P₂ and P₃ segments (Nogueira, 1987).

V. NEOPLASTIC AND PRE-NEOPLASTIC LESIONS OBSERVED IN THE 2-YEAR BIOASSAYS

Data for all reported renal tubule tumors and tubule hyperplasia in male rats from the 2-year bioassays on the eight model chemicals are summarized in Table 7. Information on tumors at non-renal sites with a statistically significant increase are also mentioned. Table 7-a provides similar information for trichloroethylene and cholerothalonil.

In addition to the specific results obtained from individual bioassays, there are considerations generic to all bioassays conducted by the NTP. For example, the NTP position with regard to evaluation of rare tumors and the use of historical controls influences NTP interpretation of the evidence for carcinogenicity of CIGA (Haseman et al., 1984). Likewise, survival rates influence the ability to analyze information from animal bioassays. These generic issues are explored before describing the results of individual studies.

TABLE 7. INCIDENCES OF RENAL TUBULE PRENEOPLASIA AND NEOPLASIA IN RATS TAKEN FROM 2-YEAR BIOASSAYS ON EIGHT MODEL COMPOUNDS

Chemical	Strain	Sex	Changes	Doses (mg/kg/day)		
				0	150	300
1,4-Dichloro- benzene (NTP-TR-319) 1987a Gavage	F344	M	Survival (%)	77	69	43
			Hyperplasia (%)	0	2	18
			<u>Adenomas</u>			
			Incidence	0/50	0/50	1/50
			Adj. Rate (%)	0	0	4
			<u>Adenocarcinomas</u>			
			Incidence	1/50	3/50	7/50
			Adj. Rate (%)	3	9	26
			<u>Combined</u>			
			Incidence	1/50	3/50	8/50
			Adj. Rate (%)	3	9	28
<u>Other Tumors:</u> Hepatocellular tumors in mice						
Chemical	Strain	Sex	Changes	Doses (mg/kg/day)		
				0	500	1000
Dimethyl methyl phosphonate (NTP-TR-323) 1987b Gavage	F344	M	Survival (%)	56	34	19
			Hyperplasia (%)	0	16	18
			<u>Adenomas</u>		None	
			<u>Adenocarcinomas</u>			
			Incidence	0/50	2/50	3/49
			Adj. Rate (%)	0	9	19
			<u>Other Tumors:</u> Mononuclear cell leukemia; transitional cell papillomas			

continued

TABLE 7. INCIDENCES OF RENAL TUBULE PRENEOPLASIA AND NEOPLASIA
IN RATS TAKEN FROM 2-YEAR BIOASSAYS ON EIGHT MODEL COMPOUNDS
(continued)

Chemical	Strain	Sex	Changes	Doses (mg/kg/day)			
				0	0	212	423
Hexachloro- ethane (NTP-TR-68) NCI 1978b	Osborne- Mendel	M	Survival (%)	56	65	20	18
			Hyperplasia (%)	Not Reported			
			<u>Adenomas</u>				
			Incidence	0/20	0/18	4/37	0/29
			Adj. Rate	0	0	11	0
Gavage			<u>Carcinoma</u>	None			

Other Tumors: Hepatocellular tumors in mice

Chemical	Strain	Sex	Changes	Doses (mg/kg/day)		
				0	10	20
Hexachloro- ethane (NTP-TR-361) 1989	F344	M	Survival (%)	62	58	52
			Hyperplasia (%)	4	8	22
			<u>Adenomas</u>			
			Incidence	1/50	2/50	4/50
			Adj. Rate (%)	3	6	15
Gavage			<u>Adenocarcinomas</u>			
			Incidence	0/50	0/50	3/50
			Adj. Rate (%)	0	0	9
			<u>Combined</u>			
			Incidence	1/50	2/50	7/50
			Adj. Rate (%)	3	6	24

Other Tumors: Marginal increase in pheochromocytomas in M rats

continued

**TABLE 7. INCIDENCES OF RENAL TUBULE PRENEOPLASIA AND NEOPLASIA
IN RATS TAKEN FROM 2-YEAR BIOASSAYS ON EIGHT MODEL COMPOUNDS
(continued)**

Chemical	Strain	Sex	Changes	Doses (mg/kg/day)		
				0	250	500
Isophorone (NTP-TR-291) 1986a	F344	M	Survival (%)	66	66	28
			Hyperplasia (%)	0	2	8
			<u>Adenomas</u>			
Gavage			Incidence	0/50	0/50	2/50
			Adj. Rate (%)	0	0	8
	<u>Adenocarcinomas</u>					
			Incidence	0/50	3/50	1/50
			Adj. Rate (%)	0	9	4
<u>Combined</u>						
			Incidence	0/50	3/50	3/50
			Adj. Rate (%)	0	9	12
<u>Other Tumors:</u> Preputial gland tumors in male rats; hepatocellular tumors, mesenchymal tumors & malignant lymphomas in male mice						

Chemical	Strain	Sex	Changes	Doses (mg/kg/day)		
				0	75	150
d-Limonene (NTP-TR-347) 1990	F344	M	Survival (%)	60	68	69
			Hyperplasia (%)	0	4	7
			<u>Adenomas</u>			
Gavage			Incidence	0/50	4/50	8/50
			Adj. Rate (%)	0	12	19
	<u>Adenocarcinomas</u>					
			Incidence	0/50	4/50	3/50
			Adj. Rate (%)	0	12	7
<u>Combined</u>						
			Incidence	0/50	8/50	11/50
			Adj. Rate (%)	0	23	25
<u>Other Tumors:</u> None in mice or rats						
continued						

continued

TABLE 7. INCIDENCES OF RENAL TUBULE PRENEOPLASIA AND NEOPLASIA IN RATS TAKEN FROM 2-YEAR BIOASSAYS ON EIGHT MODEL COMPOUNDS (continued)

Chemical	Strain	Sex	Changes	Doses (mg/kg/day)		
				0	75	150
Pentachloro-ethane	F344	M	Survival (%)	82	68	52
			Hyperplasia (%)	0	0	2
(NTP-TR-232) 1983			<u>Adenomas</u>			
			Incidence	0/50	1/49	4/50
			Adj. Rate (%)	0	3	14
Gavage			<u>Adenocarcinomas</u>			
			Incidence	1/50	1/49	0/50
			Adj. Rate (%)	2	3	0
			<u>Combined</u>			
			Incidence	1/50	2/49	4/50
			Adj. Rate (%)	2	6	14

Other Tumors: Hepatocellular tumors in mice

Chemical	Strain	Sex	Changes	Doses (ppm)		
				0	200	400
Tetrachloro-ethylene	F344	M	Survival (%)	48	40	24
(NTP-TR-311) 1986b			Hyperplasia (%)	0	6	10
			<u>Adenomas</u>			
Inhalation			Incidence	1/49	3/49	2/50
			Adj. Rate (%)	4	11	11
			<u>Adenocarcinomas</u>			
			Incidence	0/49	0/49	2/50
			Adj. Rate (%)	0	0	11
			<u>Combined</u>			
			Incidence	1/49	3/49	4/50
			Adj. Rate (%)	4	11	22

Other Tumors: Leukemia in rats; hepatocellular tumors in mice
continued

**TABLE 7. INCIDENCES OF RENAL TUBULE PRENEOPLASIA AND NEOPLASIA
IN RATS TAKEN FROM 2-YEAR BIOASSAYS ON EIGHT MODEL COMPOUNDS
(continued)**

Mixture	Strain	Sex	Changes	Doses (ppm)			
				0	67	292	2056
Unleaded gasoline (US EPA) 1987	F344	M	Survival (%)	Not affected			
			Hyperplasia (%)				
			<u>Adenomas</u>				
			Incidence	0/49	1/59	2/56	1/45
			Adj. Rate (%)	0	2	4	2
			<u>Carcinomas</u>				
			Incidence	0/49	1/59	2/56	6/45
			Adj. Rate (%)	0	2	4	14
			<u>Combined</u>				
			Incidence	0/49	1/59	5/56	7/45
			Adj. Rate (%)	0	2	9	16

Other Tumors: Hepatocellular tumors in F mice

TABLE 7a. INCIDENCES OF RENAL TUBULE PRENEOPLASIA AND NEOPLASIA IN RATS TAKEN FROM 2-YEAR BIOASSAYS ON CHLOROTHALONIL AND TRICHLOROETHYLENE

Chemical	Strain	Sex	Changes	Doses (ppm)		
				0	5063	10126
Chlorotha- lonil (NTP-TR-41) NCI 1978a	Osborne- Mendel	M	Survival (%)	82	40	40
			Hyperplasia (%)		none	
			<u>Adenomas</u>			
			Incidence	0/10	2/46	1/49
			Rate (%)	0	4	2
			<u>Carcinomas</u>			
			Incidence	0/10	1/46	3/49
			Rate (%)	0	2	6
			<u>Combined</u>			
			Incidence	0/10	3/46	4/49
			Rate (%)	0	6	8
<hr/>						
		F	Survival (%)	50	62	72
			Hyperplasia (%)		none	
			<u>Adenomas</u>			
			Incidence	0/10	0/48	3/50
			Rate (%)	0	0	6
			<u>Carcinomas</u>			
			Incidence	0/10	1/48	2/50
			Rate (%)	0	2	4
			<u>Combined</u>			
			Incidence	0/10	1/48	5/50
			Rate (%)	0	2	10

Other Tumors: none

(continued)

**TABLE 7a. INCIDENCES OF RENAL CELL PRENEOPLASIA AND NEOPLASIA
IN RATS TAKEN FROM 2-YEAR BIOASSAYS ON CHLOROTHALONIL AND
TRICHLOROETHYLENE**
(continued)

Chemical	Strain	Sex	Changes	Doses (mg/kg/day)			
				0	0	500	1000
Trichloro- ethylene (NTP-TR-273) 1988a Gavage	Osborne Mendel	M	Survival (%)	42	44	34	30
			Hyperplasia (%)	0	0	10	6
			<u>Adenomas</u>				
			Incidence	0/50	0/50	6/50	1/50
			Adj. Rate (%)	0	0	32	6
			<u>Carcinomas</u>				
			Incidence	0/50	0/50	0/50	1/50
			Adj. Rate (%)	0	0	0	6
			<u>Combined</u>				
			Incidence	0/50	0/50	6/50	2/50
			Adj. Rate (%)	0	0	32	11

Tumors in Other Strains: 2-4% Renal tubule tumors in 3 other strains.

Other Tumors: Malignant mesothelioma in M rats; hepatocellular tumors in male and female mice and lymphoma in F mice

A. Generic Considerations

Renal tubule tumors are neoplasms with a low background incidence in laboratory animals including the rat strains used in the chronic bioassays on CIGA, namely Fischer 344 and Osborne-Mendel. The overall historical incidence of these tumors in male Fischer 344 rats is considered by the NTP to be 0.5% based on data reported on 1,943 animals which served as vehicle controls in studies involving administration of chemicals via corn oil gavage (NTP, 1990). In a larger historical control data-base, involving 2,320 male and 2,370 female Fischer 344 rats used as untreated controls in NTP two-year bioassays, the incidence was 0.35% for males and 0.17% for females suggesting a male predilection for renal tubule tumors (Solleveld et al., 1984). This is supported by spontaneous renal tubule tumor incidence rates recorded for Osborne-Mendel rats used as controls in the NCI Carcinogenesis Testing Program (Goodman et al., 1980). In 975 males and 970 females the incidence was 0.3% and 0% respectively. Because of the infrequency of renal tubule tumors, even marginal increases in their incidence in treated animals (statistically significant when compared to historical rather than concurrent controls) is regarded by the NTP as biologically significant and attributable to compound administration (Haseman et al., 1984; NTP, 1989).

In the 2-year studies with the eight selected renal carcinogens, the observed incidences of renal tumors for individual chemically-dosed groups were less than 25%, and no higher than 16% for most. Because of the low background rate in both concurrent

and historical controls, however, development of renal tubule tumors at these incidences was ascribed to an effect of the chemical.

The NTP bioassays provide little insight into the histogenesis of the renal tumors as they were designed and performed with the prime objective of determining the presence or absence of carcinogenic activity of the test chemical. Although an industry-sponsored study of unleaded gasoline included interim sacrifices, even this bioassay did not incorporate serial sacrifices designed to provide information on the site of origin or histogenesis of tumors.

Survival rates in high dose male rats were poor in several of the NTP bioassays, which complicates interpretation of the data. The high mortality rate observed in some of these studies cannot be attributed to the renal tumors (Hoel et al., 1988). In fact poor survival rates usually indicated excessive toxicity. For the 1,4-DCB bioassay, survival of the high-dose males, 40% at termination, became significantly lower than that of vehicle controls after week 97 (NTP, 1987a). Nearly all deaths were non-accidental. A similar situation pertains to isophorone where only 28% of high dose males survived to termination (NTP, 1986a).

The decreased survival rates suggest that a maximum tolerated dose (MTD) was exceeded since the early deaths could not be attributed to tumors. Administration of a chemical at dose-levels exceeding an MTD may alter responses that would be seen at lower dose-levels (OSTP, 1985). However, exceeding an MTD, by itself, is

not compelling evidence that tumors are produced only when detoxification mechanisms are overwhelmed. In fact, survival of male rats in low-dose groups administered isophorone, 1,4-DCB, hexachloroethane and tetrachloroethane was equivalent to that of the concurrent control groups and renal tumor incidence was elevated in these animals. Survival was excellent for all dose groups of male rats administered d-limonene or unleaded gasoline. However, it is difficult to compare tumor incidences among studies with marked differences in survival rates, especially when there is the potential for development of slow-growing tumors, such as renal neoplasms.

B. Renal tumor incidence

Among the eight model carcinogens, the overall unadjusted incidence rates for renal tubule tumors (adenomas and adenocarcinomas/ carcinomas combined) in male rats ranged from 3% to 11% at low-dose levels and 0% to 22% at the high dose. The highest unadjusted incidence (22%) was associated with d-limonene. For the remainder of the chemicals, incidences of renal tumors were 16% or less. When adjusted for intercurrent mortality, the incidence rates for combined renal tumors ranged from 0% to 28% with 1,4-DCB highest (Table 7).

For all of the eight model carcinogens, and also for trichloroethylene and chlorothalonil, the increase in the incidences of renal tubule tumors, where adjusted for intercurrent mortality, was dose-related. Because the incidence of renal tubule tumors was low and there were confounding factors such as toxicity

occurring at all dose-levels in most studies, it is not possible from the NTP bioassay data to determine if there was a relationship between increasing dose and percentage of tumors classified as adenocarcinomas rather than adenomas. In its 1986 Cancer Guidelines, EPA discussed its strategy for analyzing combinations of benign and malignant tumors (U.S. EPA, 1986). In general, the Agency stated that it would consider the combination of benign and malignant tumors to be scientifically defensible if the benign tumors have the potential to progress to the associated malignancies of the same histogenic origin. The weight-of-evidence that a chemical is potentially carcinogenic for humans would increase when there is a dose-related increase in the proportion of tumors that are malignant. Conversely, if only benign tumors were observed, this would constitute less evidence of human cancer potential. Since the distinction between adenomas and adenocarcinomas for renal tubule tumors in rats is rather arbitrary, based mainly on size, these general principles cannot be rigidly applied.

C. Histogenesis of renal tumors

As previously indicated, NTP bioassays are designed to determine whether or not a chemical is a carcinogen. They are not designed with the intent of providing information to evaluate the developmental stages of renal neoplasia. Although renal tubule hyperplasia was reported in the male rat for seven of the eight bioassays and incidences of this lesion generally increased with increasing dose, further insight with respect to histogenesis into

possible interrelationships between hyperplasia, adenomas, and adenocarcinomas is not possible because of the low overall frequency of these lesions. The occurrence together of pre-neoplastic and neoplastic lesions in most studies with the eight chemicals does provide indirect evidence of progression from tubule cell hyperplasia via adenomas to adenocarcinomas. In studies with d-limonene (NTP, 1990) and hexachloroethane (NTP, 1989), these lesions were stated to be part of a continuous morphologic spectrum. This accords with the generally accepted view on renal tubule tumor formation and progression (Lipsky and Trump, 1988; Hard, 1990).

D. Renal tumor latency and progression

Renal tubule tumors produced by administration of CIGA appear to be late developing neoplasms. Times at which such tumors were first observed in bioassays of the eight model carcinogens usually exceeded 18 months. In general, the first renal tumor observed in each of the bioassays occurred about 5-10 weeks earlier in the high-dose than in low-dose animals. Because renal tubule tumors are not immediately life-threatening, they were usually detected in bioassays at terminal sacrifice or at death of the animal from other causes. Out of the eight bioassays, there was only one case of renal tumor metastasis, occurring in the high-dose group of hexachloroethane (NTP, 1989).

E. Induction of other tumor types

Six of the eight model compounds produced liver tumors in male and/or female mice but not in male or female rats. These chemicals

were hexachloroethane, unleaded gasoline, isophorone, 1,4-DCB, pentachloroethane, and tetrachloroethylene. A different mechanism, independent of hyaline droplet accumulation, may be involved in the production of liver tumors by these six chemicals. Some authors suggest a mechanism involving peroxisome proliferation to account for the production of such liver tumors (Elcombe et al., 1985; Goldsworthy and Popp, 1987).

An alternative explanation for the liver tumors is that both CIGA-induced liver and kidney tumors are produced by a common mechanism (direct or indirect) not involving α_{2u} -g. Available data do not tend to support this hypothesis, although a recent inhalation toxicity study of 1,4-DCB illustrates other types of data needed before these questions can be resolved. In this study, significantly higher levels of 1,4-DCB were found in the kidneys of male rats and in the livers of female rats following exposure at 500 ppm for 24 hours (Umemura et al., 1990). Although the Umemura study may simply demonstrate reaction of 1,4-DCB with α_{2u} -g, it may also indicate metabolic differences among species and sexes that influence the effective doses delivered to the tumor sites.

Primary tumors were not consistently produced in rats or mice at organ sites other than the liver following administration of the eight chemicals. The production of tumors at other sites, however, raises the possibility that other mechanisms could also be contributing to the overall kidney tumor incidence in male rats. This possibility has been suggested for perchloroethylene (Green et al., 1990; Dekant et al., 1989). Dekant and colleagues have

proposed a mechanism involving hepatic glutathione S-conjugate formation and, ultimately, bioactivation by renal cysteine conjugate β -lyase in the nephrotoxic and carcinogenic response to halogenated alkenes, including perchloroethylene, although they also do not rule out a role for α -_{2u}-g-induced nephrotoxicity. Within this context, it is noteworthy that in the tetrachloroethylene bioassay a renal tubule adenocarcinoma was observed in a single low dose male mouse, clearly a statistically nonsignificant event, but less readily regarded as biologically irrelevant.

VI. ADDITIONAL EVIDENCE CONCERNING THE RENAL CARCINOGENICITY OF CIGA

Key evidence relevant to providing information on carcinogenic mechanisms can also be derived from short term tests, such as assays for gene mutations and DNA damage, and from studies testing the tumor-promoting effects of CIGA.

A. Genetic toxicology studies

The available genotoxicity data for the eight model carcinogens and for trichloroethylene and chloroethalonil are summarized in Table 8. The four assays listed in the table (Salmonella (SAL), chromosome aberrations in Chinese hamster ovary cells (ABS), sister chromatid exchange in Chinese hamster ovary cells (SCE), and thymidine-kinase (TK)-gene mutations in L5178Y cells (MLA)) are the only ones with enough common data for comparative purposes. It is not coincidental that these are the assays employed by the NTP. Consequently, this analysis of

genotoxicity data was limited, for the main part, to the 10 chemicals with bioassay data. Data from *Drosophila* tests conducted by the NTP (Yoon et al., 1985) and in human lymphoblasts (Richardson et al., 1986) are also cited in Table 8 when available.

All eight renal carcinogens selected as potential CIGA have been tested for chromosome aberrations in Chinese hamster ovary (CHO) cells (Galloway et al., 1987a) and in Salmonella (Haworth et al., 1983; Mortelmans et al., 1986; Ashby and Tennant, 1988; NTP, 1987). All results were negative both in the absence and presence of exogenous activation provided by S9 extracts from rat liver. Two presumed intermediate metabolites of the CIGA, d-limonene, (the 1,2- and 8,9-epoxides) were also tested in Salmonella with and without induced S9, and no increase in revertants was observed (Watabe et al., 1981). Several chemicals have tested positive, at least under some conditions, for sister chromatid exchange in CHO cells (Galloway et al., 1987a) and in the mouse lymphoma TK gene mutation assay (McGregor et al., 1988). Four of the eight potential CIGA and both the non-CIGAs were positive. Richardson et al. (1986) reported negative results for unleaded gasoline and its known CIGA component, TMP, in assays for TK-gene mutations and SCE in the TK6 human lymphoblast cell line. A cursory appraisal of only positive and negative responses leads to the conclusions that there is significant heterogeneity and the CIGA groups are not distinguishable from non-CIGA by their genotoxic activity. Upon more detailed analysis, it becomes apparent that the majority of the positive responses of the eight model carcinogens selected as

TABLE 8. SUMMARY OF GENOTOXICITY DATA FOR 10 SELECTED MALE RAT KIDNEY CARCINOGENS

AGENT	SAL	ABS	SCE	MLA	COMMENTS
Chloroethalonil	-	+	+	+	ABS and MLA positive without S9; MLA not tested with S9. Negative in Drosophila SLRL.
1,4-Dichlorobenzene	-	-	-	E	MLA with S9 there was a marginal positive result in one of three experiments. Negative in <u>in vivo</u> chromosome aberrations, micronuclei, and dominant lethals.
Dimethyl methylphosphonate	-	E	+	+	MLA and SCE results positive without S9; MLA not tested with S9; SCE negative with S9. ABS negative in two labs, both +/-S9 (NTP) but positive reported by Aerospace Med. (-S9). Drosophila SLRL positive, but translocations negative. Dominant lethal positive in both rats and mice.
Hexachloroethane	-	-	+		SCE reproducible positive only with S9. No data for MLA.
Isophorone	-	-	+	±	SCE only positive with cell cycle delay without S9; MLA replicated positive without S9, not tested with S9 in NTP studies. CMA reported negatives for hepatocyte UDS, mouse micronuclei, and MLA (both +/- S9).
D-Limonene	-	-	-	-	Clear negative in all NTP studies.
Pentachloroethane	-	-	+	+	MLA and SCE positive without S9 (reproduced); SCE negative with S9. Negative in rat <u>in vivo</u> kidney UDS assay.
Tetrachloroethylene	-	-	-	-	In NTP studies all clear negatives both with and without S9; Also Drosophila SLRL negative. Negative in rat <u>in vivo</u> kidney UDS assay. Recent positive in Salmonella TA100 with GSH and kidney microsomes.

(continued)

AGENT	SAL	ABS	SCE	MLA	COMMENTS
Trichloroethylene	-	-	+	+	Unpublished studies negative in SAL, MLA as well as in dominant lethal and bone marrow cytogenetic studies in mice. MLA on various catalytic fractions gave mixed results. Positive UDS in rat, mouse, and human hepatocytes. In vivo studies with gavage were positive in mouse, but not rat liver. Kidney UDS in rats negative (both gavage and inhalation).
Unleaded gasoline	-	-	-	+/-	In NTP studies both positive responses were only with S9. Other studies confirm negative in bacteria. In yeast, positives have been reported for mitotic recombination and both positive and negative responses for gene mutations. Negative in rat in vivo kidney UDS assay.

SAL = Salmonella; ABS = Chromosome aberrations in CHO cells; SCE = Sister chromatid exchange in CHO cells; MLA = Thymidine-kinase (TK)-gene mutation assay in L5178Y cells; GSH = Glutathione; TFT = Trifluorothymidine; SLRC = Sex-linked recessive lethal; UDS = unscheduled DNA synthesis.

hyaline droplet inducers were observed in the absence, but not in the presence, of exogenous S9 activation and at concentrations greater than 100 µg/ml.

Dimethyl methylphosphonate appears to present a unique genotoxicity profile among the eight model carcinogens. Because dimethyl methylphosphonate has high water solubility and low toxicity, in vitro assays have employed very high concentrations of dimethyl methylphosphonate, as high as 30 mg/ml. Galloway et al. (1987b) reported that at least some of the observed in vitro mutagenic activity seen for dimethyl methylphosphonate occurred at levels that decreased cell growth and greatly increased the osmotic strength. Similar levels of osmolality and chromosome aberrations were observed, for example, with 160 mM of potassium chloride and 30 mg/ml of dimethyl methylphosphonate. The SCE increases observed for dimethyl methylphosphonate, however, occurred at concentrations causing only slight increases in osmolality.

Of particular relevance to this report are those studies in which rodent kidney or kidney extracts are combined with a genotoxic endpoint. Loury et al. (1987) reported that unleaded gasoline was negative in an in vivo/ in vitro kidney UDS assay indicative of DNA damage and repair. Similar results were reported for pentachloroethane and tetrachloroethylene by Goldsworthy et al. (1988b). However, both studies reported significant elevation of replicative DNA synthesis in kidneys of male rats treated with these compounds.

Recently, Vamvakas et al. (1989) reported clear dose-related

positive results in Salmonella TA100 with tetrachloroethylene in the presence of glutathione and rat kidney microsomes. The glutathione conjugate S-(1,2,2-trichlorovinyl)glutathione was also mutagenic in the presence of kidney microsomes and the activity was reduced in the presence of a β -lyase inhibitor. The importance of these findings in the formation of the kidney tumors of male rats exposed to tetrachloroethylene is yet unclear, but similar studies with other kidney carcinogens seem to be in order before direct interaction with DNA can be excluded.

In summary, the preponderance of available data suggest that the CIGA group possess little, if any, genotoxic activity. However, the dearth of data in the kidney or with glutathione conjugates for these chemicals precludes closure on the question.

B. Initiation-promotion studies

The multistage concept of carcinogenesis, involving in its simplest form an irreversible initiation phase followed by a stage of tumor promotion (Pitot, 1982), implies that chemicals may play a role in assisting, as well as directly causing, cancer formation. There have been two research studies testing the potential of CIGA for promoting or cocarcinogenic activity in an established initiation-promotion model of renal carcinogenesis.

Using 2 weeks exposure to 170 ppm of EHEN in the drinking water as the initiating agent, the first initiation-promotion experiment of Short et al. (1989b) included both sexes of Fischer 344 rats, multiple dose-levels of the two test compounds (unleaded gasoline and TMP), short-term versus long-term promotion exposures,

and a sequence-reversal study to discriminate any cocarcinogenic from promotional effects. The test compounds were unleaded gasoline (3 inhalation concentration-levels of 10, 70 and 300 ppm), and TMP (one oral dose-level of 50 ppm). Treatment groups, comprised of approximately 30 animals, included a control, 2 promotion controls, an EHEN initiation control, reverse-sequence initiation control, initiation-promotion group with a promotion phase of 24 weeks, initiation-promotion group with a promotion phase of 59 weeks, and a reverse-sequence test group where 24 weeks of exposure to unleaded gasoline or TMP preceded the 2-week period of EHEN administration. All animals were killed at 65 to 67 weeks after the commencement of the experiment. The results were assessed in terms of the incidence of foci of tubule hyperplasia (called atypical cell foci by the authors) and renal tubule tumors.

Dose-related increases in hyperplastic foci were observed in male rats promoted with unleaded gasoline or TMP for both the short- and long-term promotion periods. A significant linear trend in the incidence of renal tubule tumors with increasing gasoline dose was also observed in male rats promoted with unleaded gasoline for 24 weeks but not for 59 weeks. The latter discrepancy reflects an experimental design weakness in the study, namely underestimation of an optimal initiating dose of EHEN, which resulted in a very low basal incidence of renal tumors. Nevertheless, the results with the single dose-level of TMP, and the absence of renal tumors in any negative control group, supported the observed trends with unleaded gasoline.

In the sequence-reversal study, there was no increase in renal tumors although the incidence of hyperplastic foci was significantly elevated for both compounds. Foci of CPN were also scored in these various groups with an increase upon CIGA exposure apparent in male rats. However, no correlation of incidence of CPN lesions with numbers of hyperplastic foci or incidence of renal tubule tumors was found.

On the basis of the results, the authors' conclusions that unleaded gasoline and TMP have promoting activity for renal tubule tumors in the male rat, rather than acting as cocarcinogens, appear reasonable. Furthermore, there was no elevation of either hyperplastic foci or renal tumors in female rats in the study, emphasizing once again, the male-specificity of the renal response to CIGA.

A second initiation-promotion assay using the same EHEN model was conducted with d-limonene (Deitrich and Swenberg, 1991). This study specifically addressed the comparison of responses between the male Fischer 344 rat and the α_{2u} -g-deficient NBR strain. The initiating dose of EHEN was 500 ppm administered in the drinking water for two weeks, followed by d-limonene by daily gavage at 150 mg/kg for 30 weeks. An initiation control (EHEN), promotion control (d-limonene), and a vehicle control was included for both strains. In the Fischer rats administered EHEN and d-limonene, atypical tubule cell hyperplasia and renal tubule adenomas were increased ten-fold as compared to the EHEN control group. In contrast, no tumors were observed in any of the NBR groups. Such

negative results in the NBR rat strongly suggest a clear dependence on α_{2u} -g for the promoting activity of d-limonene.

The promotional effect of unleaded gasoline, TMP, and d-limonene may be occurring through the influence of sustained tubule cell proliferation which has been demonstrated with these same compounds (Short et al., 1989a; Dietrich and Swenberg, 1991). The extent of cell proliferation is regarded as an important factor in chemical carcinogenesis (Grisham et al., 1983; Cohen and Ellwein, 1990) and stimulation of cell turnover is one of the key mechanisms believed to operate in tumor promotion (Farber, 1988).

VI. COMPARISON OF CIGA WITH CLASSICAL RENAL CARCINOGENS

In general, classical renal carcinogens or their active metabolites are electrophilic species binding covalently to macromolecules and forming, in particular, DNA adducts (Hard, 1987; Lipsky and Trump, 1988; Alden, 1991). Such DNA reactivity is putatively the mechanistic basis of renal carcinogenesis induced by these chemicals. For example, carcinogenic nitrosamines can form various alkylation products in DNA, including O⁶-alkylguanine which is a promutagenic lesion (Pegg, 1984). Accordingly, classical renal carcinogens are usually positive in short-term mutagenicity assays.

In contrast, CIGA are not known to react with DNA and are generally negative in short-term tests for genotoxicity. As described previously (Section IIG) CIGA binding to α_{2u} -g is reversible and not covalent in nature.

Classical renal carcinogens can induce renal tubule cancer in

rats or mice in high incidence, with minimal duration of exposure, clear dose-response relationships, and with decreased latent period of development (Hard, 1987; Alden, 1991). Tumor frequencies are often over 50% and up to 100%, much higher than the low incidences (2-28% adjusted) recorded for CIGA. Some genotoxic carcinogens, e.g. DMN, DEN and streptozotocin, are highly effective by single dose. Unlike CIGA-induced renal carcinogenesis, there is usually no absolute sex-specificity, with males and females both susceptible, but sometimes to varying degree. These differences in potency and species- and sex-susceptibility, suggest that classical renal carcinogens and CIGA act via different mechanisms in kidney carcinogenesis.

The lack of involvement of hyaline droplet accumulation in the early nephrotoxicity associated with classical carcinogens (definite with DMN and DEN and apparent with the others) is a major difference from the sequence of early pathological events induced by CIGA in the male rat.

Pathology reports indicate that renal tubule tumors induced by CIGA are morphologically indistinguishable from spontaneous tumors or those induced by classical carcinogens, with both granular and clear cell types occurring. Likewise, despite differences in toxicity observed, the sequence of development of CIGA-induced renal tumors from tubule hyperplasia to carcinoma appears identical. However, some evidence from the bioassays suggests that the CIGA tumors may, in general, have a smaller size, probably because of the difference in potency between these chemicals and

classical carcinogens, affecting the latent period of tumor development.

As with classical carcinogens, metastases have been rarely reported for renal tubule tumors related to treatment by chemicals inducing hyaline droplets and/or α_{2u} -g. The one case of metastasis noted with hexachloroethane suggests, however, that a malignant potential exists for such neoplasms.

Although the specific site of origin for the renal tubule tumors produced by CIGA is not known, the P2 region of the proximal tubule as the primary site would be consistent with existing information. Based on studies with classical carcinogens this does not represent an unusual location.

VIII. EVIDENCE CONCERNING HUMAN KIDNEY CANCER

Although not one of the most common neoplasms in the United States, renal cell adenocarcinoma/carcinoma is regarded as an important human cancer. This is because the disease is unpredictable and a significant proportion of patients, approximately one third, have distant metastasis at the time of diagnosis (Bennington and Beckwith, 1975; NCI, 1987). The mortality rate in these cases is high, and overall, the survival rate for patients with renal cell cancer is 48% (Devesa et al., 1990). In addition, the etiology of kidney cancer in humans is poorly understood.

A. Morphology and histogenesis

Human renal cell tumors, which are morphologically similar to those of rodents, are classified according to cell type and cellular arrangement. Thus, two main cell forms are recognized, granular and clear, and the usual patterns of organization are tubular, solid, papillary and cystic. Individual tumors may show an admixture of patterns and of cell types. Infrequently, renal cell carcinoma presents as a sarcomatoid form composed of spindle cells (Bennington and Beckwith, 1975; Bannayam and Lamm, 1980; Tannenbaum, 1983).

It is generally accepted that the origin of renal cell carcinoma is the proximal tubule, based on both immunological study (Wallace and Nairn, 1972) and ultrastructural features (Tannenbaum, 1971; Bennington and Beckwith, 1975). Electron microscopy reveals many similarities between the tumor cells and normal proximal

tubule epithelium, including brush border elements, membrane-associated vesicles, and basilar infoldings of the plasma membrane (Tannenbaum, 1971). Ultrastructurally, the amount of intracellular lipid, particulate glycogen, and organelles distinguishes clear from granular cells.

It is widely considered that human renal adenomas represent small adenocarcinomas or carcinomas as there are no microscopic, histochemical or immunologic features which discriminate them, other than size, and this is not an absolute biologic parameter (Bennington and Beckwith, 1975; Ritchie and Chisholm, 1983; Tannenbaum, 1983). Adenomas are therefore considered part of an evolutionary continuum from hyperplasia, through adenoma, to adenocarcinoma/carcinoma, as in rodents. As a general observation, there is a direct relationship between tumor size and frequency of metastasis (Bell, 1950; Hellsten et al., 1981; Ritchie and Chisholm, 1983).

B. Incidence and mortality

Kidney cancer statistics are usually reported in a form which encompasses all types of malignant cancer affecting kidney, renal pelvis, and sometimes ureter and urethra. Renal cell cancer rarely occurs under the age of 40 years (McLaughlin and Schuman, 1983; Asal et al., 1988) and represents about 70% of all kidney tumors in adults (Devesa et al., 1990). Kidney cancer statistics, therefore, provide an approximation only of renal cell tumor prevalence.

The number of new cases of kidney and urinary tract cancer (excluding bladder) estimated for 1989 in the U.S. is 23,100 with

a mortality estimate of 10,000 deaths (Silverberg and Lubera, 1989). These figures represent approximately 2% of both new cancer cases at all sites and total cancer deaths. The age-adjusted incidence rates in the U.S. for the period between 1975-1985 obtained from the NCI Surveillance, Epidemiology and End Results Program (SEER) data for renal cell cancer are 8.4 per 100,000 for males and 3.7 per 100,000 for females, with no difference among racial groups (Devesa et al., 1990). Most studies indicate a consistent male to female ratio of 2:1 for the incidence of renal cell tumors (Asal et al., 1988; Devesa et al., 1990).

In considering renal cell tumors specifically, the highest rates internationally have been reported from Iceland and other Scandinavian countries. Renal cell carcinoma is the fifth most common malignant tumor of males in Iceland although it ranks only tenth in females (Thorhallson and Tulinius, 1981). The lowest rates for renal cancer are recorded in Africa, Asia and South America (McLaughlin and Schuman, 1983). Within the U.S., mortality surveys indicate that the North Central region and some areas in the Northeast have the highest incidence rate for renal cell carcinoma (Pickle et al., 1987). It has been suggested that the clustering in the North Central region may be partially explained by the predominantly German and Scandinavian origin of the area's population (McLaughlin et al., 1984). Several studies have reported that the urban rates for renal cell tumor incidence are higher than for rural areas, but the correlation is considered to be weak (Newsom and Vugrin, 1987).

In contrast to the relatively low incidence and mortality figures for malignant kidney and related tumors provided by cancer statistics data, the occurrence of renal cell adenomas at autopsy is common. The reported incidence has ranged from 15% (Bannayam and Lamm, 1980) up to 25%, the latter for males over the age of 50 (Reese and Winstanley, 1958). These findings have led to speculation that a proportion of adenomas may reach a limit of growth and/or remain quiescent (Bannayam and Lamm, 1980; Warter 1983).

Over the period 1950-1985, the U.S. Cancer Statistics data indicate an increase of 82% in the incidence of kidney and renal pelvis cancer combined (NCI, 1987). For renal cell cancer alone, the increase among whites was estimated at about 30% between 1969-1971 and 1983-1985 representing an average annual percent change in incidence of 2.0 for males and 1.8 for females (Devesa et al., 1990). Data from Cancer Registries in Scotland between 1967 and 1979 also indicate an increase of approximately 37% in the incidence of renal cell carcinoma for males, although no overall increase in females (Ritchie and Chisholm, 1983). Despite an improvement in mortality rates since 1950 compared to incidence rates (NCI, 1987), the relative 5-year survival rates, which are close to 50%, have not altered since the early 1970's (Silverberg and Lubera, 1989), suggesting little improvement in treatment over the past two decades. On the other hand, diagnostic detection measures have improved dramatically during this time which may explain, at least in part, the observed increase in renal cancer

incidence (Higginson et al., 1984; NCI, 1989).

Renal cell carcinoma has been diagnosed with increasing frequency in patients with chronic renal failure (Hughson et al., 1986; Newsom and Vugrin, 1987). In particular, this appears to reflect an association with the development of acquired renal cystic disease which frequently occurs in patients on long-term hemodialysis. The incidence of renal cell carcinoma in patients with acquired cystic disease has been estimated as approximately 6% (Hughson et al., 1986). Thus, current data suggest that a growing population of humans receiving maintenance dialysis may be at risk for developing renal cell tumors.

C. Environmental and lifestyle factors

Potential etiological associations between renal cell cancer and exogenous and endogenous environmental factors, lifestyle and occupation, have been sought in cohort and case-control studies. Of all the environmental and lifestyle factors investigated, tobacco use in the form of cigarette, cigar or pipe smoking has been the one most consistently associated with renal cell carcinoma (Dayal and Kinman, 1983; McLaughlin and Schuman, 1983; Yu et al., 1986; Asal et al., 1988; Brownson, 1988). Although a few studies have failed to identify a statistical association between smoking and renal cell cancer, it has been estimated that 30% of renal cell carcinomas in males and 24% in females may be attributable to cigarette smoking (McLaughlin, et al., 1984) and that there is evidence for a moderate dose-response (McLaughlin and Schuman, 1983). One study has also linked use of chewing tobacco with renal

cell carcinoma in males (Goodman et al., 1986) and another has associated smoking with renal adenoma (Bennington et al., 1968).

Other possible risk factors which have been reported include coffee and tea consumption, artificial sweeteners, high body mass index (maintained from 20 years of age), high dietary animal protein and fat, lower educational levels, long-term analgesic use, and diuretics (reviewed in Dayal and Kinman, 1983; McLaughlin and Schuman, 1983; McLaughlin, 1984; McLaughlin et al., 1984; Goodman et al., 1986; Yu et al., 1986; Asal et al., 1988; McCredie et al., 1988). Of these, the evidence is least consistent for beverage consumption, artificial sweeteners, other dietary factors, and socioeconomic status, and strongest for high body mass index and drug use (phenacetin and diuretics).

D. Occupational factors

Although a number of epidemiological studies have reported some association between occupation and renal cancer, clear occupational determinants have yet to be demonstrated and it is considered that much epidemiological research is needed to further define and quantify potential risks (McLaughlin and Schuman, 1983). Occupational exposures in North America where at least one study has reported an association with increased kidney cancer rates include asbestos (Selikoff et al., 1979; Smith et al., 1989), coke-oven emissions in the steel industry (Redmond et al., 1972), printing press chemicals (Paganini-Hill et al., 1980), laundry- and dry-cleaning agents (Blair et al., 1979; Katz and Jowett, 1981; Duh and Asal, 1984), exhaust fumes in truck drivers (Brownson, 1988),

petroleum, tar and pitch products (Thomas et al., 1980; Hanis et al., 1982; Wen et al., 1983; McLaughlin et al., 1984; Savitz and Moure, 1984; Kadamani et al., 1989) and aviation and jet fuels (Siemiatycki et al., 1987). In these studies, information on smoking history was rarely available, so that its possible influence could not be determined.

A study of renal cancer by occupation in Sweden, where the incidence rates are higher than in the U.S., did not detect increased risk for hearth and furnace workers in the steel industry, printing workers, laundry-and dry-cleaners, or workers in petroleum refineries and gasoline stations (McLaughlin et al., 1987). Instead, the Swedish study reported an increase in incidence of renal cell cancer among health care professionals.

E. Renal cancer and hydrocarbon, solvent or petroleum product exposure

Several of the occupations listed above involve exposure to certain classes of chemicals that may fall into the CIGA category. Besides CIGA, however, non-CIGA compounds are also present making it difficult to attribute elevations in risk with a unique exposure (e.g., CIGA). In a recent population-based case-control study, Kadamani et al. (1989) reported a weak positive association (OR 1.6; 95% CI 0.7-3.6) between renal cell carcinoma and high occupational exposure to hydrocarbons in males but not in females (OR 0.8; 95% CI 0.3-2.3). The authors reported a dose-response relationship for the older age groups and for workers with the greatest duration of exposure.

The synthetic solvents that have been widely used in dry-

cleaning include one chemical shown in rodent tests to be a CIGA, namely tetrachloroethylene, as well as Stoddard and 140F solvents which are mixtures of hydrocarbons including straight and branched chain paraffins. Three studies analyzing proportional mortality data on laundry- and dry-cleaning workers in several U.S. states reported elevated risks for kidney cancer (Blair et al., 1979; Duh and Asal, 1984; Katz and Jowett, 1981). More recently, however, a better designed cohort mortality study on a larger population of dry cleaning workers by Blair et al. (1990) revealed no excess kidney cancer (Standardized Mortality Ratio (SMR) 0.5; 95% CI 0.1-1.8). In considering occupational exposure to solvents as a general chemical category, Harrington et al. (1989) found no relationship with renal cancer (OR 1.0; 95% CI 0.2-4.9) although the statistical power of this study was acknowledged by the authors as sufficient to identify only large risk estimates.

Siemiatycki et al. (1987) conducted a population-based case-referent study in Montreal on cancer associations with exposure to 12 petroleum-derived liquids. These various mixtures included automotive and aviation gasolines, and distillate jet fuel. Aviation gasoline differs in composition from the automotive counterpart by its high content of alkylate naphthas, constituted mainly of branched alkanes (Siemiatycki et al., 1987). No statistically significant risk of renal cancer was found with exposure to automotive gasoline (OR 1.2; 90% CI 0.8-1.6). Statistically significant elevations, however, were noted at the 90% confidence level with exposure to aviation gasoline (OR 2.6;

90% CI 1.2-5.8) and to jet fuel (OR 2.5; 90% CI 1.1-5.4). Six of the seven cases with exposure to aviation gasoline also had exposure to jet fuel, making it difficult to distinguish a unique exposure. In depth analyses of the two associations using logistic regression methods indicated, however, a greater role for aviation gasoline than for jet fuel.

Wong and Raabe (1989) conducted a quantitative meta-analysis by cancer site of petroleum industry employees from the U.S., Canada, United Kingdom, Europe, Australia and Japan, critically reviewing almost 100 published and unpublished epidemiological reports. Standardized mortality ratios observed for kidney cancer in the industry as a whole were similar to those for the general population. Results from refinery studies ranged from non-significant deficits to non-significant excesses. However, the possibility of an elevated kidney cancer risk was raised for one specific group within the industry. Drivers among British distribution workers showed borderline significance for excess kidney cancer mortality. These authors concluded that additional data, particularly involving exposure to downstream gasoline, are needed to resolve the issue. In a large population-based case-control study adjusted for the confounding factors of age and cigarette smoking, no overall association (OR 1.0; 95% CI 0.7-1.4) was observed between risk for renal cell cancer and employment in a range of occupations with potential for exposure to petroleum products (McLaughlin et al., 1985). There was, however, a small excess risk among gasoline station attendants (OR 1.2; 95% CI 0.6-

2.3) which increased with duration of employment, although individual point estimates and tests for trends were not statistically significant. A case-control study on a combined cohort of approximately 100,000 male refinery workers from five petroleum companies, sponsored by the American Petroleum Institute (Poole et al., 1990), suggested increases in kidney cancer risk for laborers (Relative Risk (RR) 1.9; 95% CI 1.0-3.9), workers in receipt, storage and movements (RR 2.5; 95% CI 0.9-6.6), and refinery unit cleaners (RR 2.3; 95% CI 0.5-9.9) when compared with a reference group of office workers, professionals and technicians. In this study there were 102 kidney cancer cases among 18,323 deaths.

In evaluating unleaded gasoline, 55 relevant studies were reviewed by USEPA (1987) to determine whether there was any epidemiologic evidence for an association between gasoline exposure and cancer risk. The evidence for drawing causal inferences between unleaded gasoline and cancer was considered inadequate under the EPA guidelines for epidemiologic evidence. As Enterline and Viren (1985) have emphasized in their review on the epidemiology of renal cancer and gasoline exposure, most of the studies have not been designed or analyzed with a specific hypothesis associating gasoline exposure and renal cancer in mind. The cohort studies of petroleum workers do not lend themselves for a comparison since they shed no light on gasoline exposure, per se. Exposures in these studies have been varied, and the only common element is the place of work. Thus, who in the cohort had the

exposure of interest, i.e. gasoline, cannot be identified.

As a general conclusion from the foregoing, small risks cannot be excluded for specific job categories, but the association between human kidney cancer and exposure to petroleum distillates, if there is one, does not suggest high risks for the types of exposures that have occurred in the past.

IX. EVIDENCE FOR DOSE- AND TIME-DEPENDENT PROGRESSION FROM EARLY TO LATE LESIONS

An important aspect for examining the hypothesis that renal tumor formation is directly associated with accumulation of α_{2u} -g in the male rat kidney is a demonstration of the progression of lesions proposed to culminate in neoplasia. For some of the steps, clear dose-response relationships have been shown. As the information presented below shows, however, data demonstrating the existence of other steps in the proposed progression are extremely limited, hindering the ability to reach judgments on the nature of the association.

Evaluation of the events leading to neoplasia is further complicated by the low incidence of renal tumors induced by the CIGA studied. Such information makes it difficult to identify possible relationships between the induced nephropathy and renal carcinogenesis.

A. Association between CIGA, hyaline droplet formation, and α -2u-globulin accumulation

Dose-dependent relationships have been demonstrated between the administration of d-limonene (Lehman-McKeeman et al., 1989) and gabapentin (Dominick, et al., 1990) and excessive formation of

hyaline droplets, and between unleaded gasoline or TMP and α_{2u} -g accumulation (Olson et al., 1987; Charbonneau et al., 1987). In the d-limonene study, hyaline droplets were graded on a scale of 0-12 according to size, eosinophilic intensity, and the number of tubules loaded with droplets. The droplet scores for d-limonene doses of 0, 0.1, 0.3, 1.0 and 3.0 mmol/kg were, control to high dose, 3, 4.5, ca.7, 8 and 10 (Lehman-McKeeman et al., 1989). The dose-response relationship with α_{2u} -g accumulation is exemplified by measurements following administration of TMP, which, given at single doses of 0.044, 0.440, and 4.000 mmol/kg, induced α_{2u} -g concentrations in rat kidney tissue at 24 hours of 10.3, 17.3 and 28.1 mg/g wet weight, respectively, against a control value of 9.5 mg/g wet weight (Charbonneau et al., 1987). With orally administered gasoline, the α_{2u} -g concentrations were dose-responsive only in the range of 0.04 to 1.00 ml/kg (Olson et al., 1987).

In a special NTP study, male and female F344 rats were exposed to d-limonene by gavage for 14 days over a 21-day period (NTP, 1990). The α_{2u} -g content, quantitated with an ELISA test in kidney homogenates, increased significantly in dosed male rats relative to vehicle controls. At 75 mg/kg, the low dose employed for male rats in the 2-year bioassay, α_{2u} -g levels were approximately double those in controls. In females, increasing the dose as high as 1,200 mg/kg had no measurable effect on α_{2u} -g levels in the kidney. Although microscopic examination of kidney sections stained with hematoxylin and eosin showed no visible differences between dose

and vehicle control male rats, in plastic embedded sections stained with Lee's methylene blue basic fuchsin, differences in the distribution, amount, and shape of intracytoplasmic granules in the proximal tubules were detected.

In contrast to the 21-day follow-up study, the 13 week range-finding study conducted before the d-limonene bioassay failed to detect an accumulation of hyaline droplets. The NTP report (1990) acknowledged that this failure might have been related to the fact that several days passed between the time the chemical was last administered and the time the animals were killed for histological examination. Other studies have shown that renal α_{2u} -g concentrations decline rapidly, reaching pre-exposure levels by the third day after treatment, although hyaline droplets, being structural entities, require up to 9 days for complete resolution (Garg et al., 1988). This suggests that the interval between the time the chemical was last administered and the time the animals were killed for histological examination is critical to finding hyaline droplets and probably accounts for discrepancies found among some studies.

These various observations, along with the results of α_{2u} -g localization studies and binding studies considered earlier, support a causal association between the administration of CIGA and α_{2u} -g accumulation in hyaline droplets.

B. Association between hyaline droplet formation, cell necrosis, and tubule cell regeneration.

Hyaline droplet accumulation, single cell necrosis and cell proliferation occur predominantly in the P2 segment of the proximal

tubule following CIGA administration (Short et al., 1987; 1989a; 1989b). Although single cell necrosis has been clearly demonstrated in association with cellular hyaline droplet accumulation (Kanerva et al., 1987a; Short et al., 1987), there are no dose-response studies quantitating the relationship between increased hyaline droplets and cell necrosis in histological sections, or between cell necrosis and cell regeneration. However, Alden (1991) has shown a correlation between the hyaline droplet response, increased mitotic index in proximal convoluted tubules, and elevation of the number of cells excreted hourly in the urine (an index of exfoliated necrotic tubule cells), using two dose-levels of d-limonene given orally for 3 weeks.

Dose-response relationships between hyaline droplet accumulation and proximal tubule cell proliferation have been observed. Short and coworkers exposed male rats for 3 weeks to TMP (oral) or unleaded gasoline (inhalation) and then measured [^3H]-thymidine labeling indices (1987). The extent and severity of hyaline droplet accumulation paralleled the extent and localization of cell proliferation in proximal tubule cells, and both parameters were increased in dose-dependent fashion (Figures 4 and 5). In an extended study of the same compounds, Short et al. (1989a) observed 6- to 11-fold increases in labeling indices in the P2 segment of the rat kidney after the rats received 3, 10 and 22 weeks of exposure to 300 ppm unleaded gasoline or 50 ppm TMP. These labeling indices remained 4- to 6-fold higher than control values during the 48th week of exposure.




FIGURE 4: Dose-response relationship between renal hyaline droplet accumulation (D) and [3H]-thymidine labeling index (L) of proximal tubule P2 cells in male F-344 rats gavaged with TMP for 5 consecutive days per week for 3 weeks. Seven-day osmotic minipump implanted on twelfth day after start of dosing. Rats killed and fixed on 22nd day. (Adapted from Short et al., 1987).




FIGURE 5: Effect of 0-2,000 ppm unleaded gasoline on continuous uptake of $[^3\text{H}]\text{TdR}$ by P1, P2, and P3 segments of the proximal tubule epithelium. Each test point is the mean value determined from 3 rats. Dosage is presented on a log scale. (Adapted from Short et al., 1987)

In contrast, Viau et al. (1986) did not observe a sustained regenerative response in the kidneys of male rats exposed to an isoparaffinic solvent consisting of saturated C₁₀-C₁₂ aliphatic hydrocarbons beyond 5.5 weeks. Labeling indices in the cortex of treated rats at 46 and 68 weeks were no different from the controls. This apparent discrepancy with the gasoline and TMP results undoubtedly reflects differences in the technique of radioactive labeling. Viau et al. (1986) used a single injection of tritiated-thymidine 1 hour prior to sacrifice whereas Short et al. (1989a) labeled continuously by subcutaneous osmotic minipump infusion over a 7-day period, the preferred method for cell populations with a low cell turnover, thereby increasing the amount of radiolabel incorporated into renal tissue.

In recovery studies with unleaded gasoline and TMP, Short and coworkers (1989a) showed that neither increased hyaline droplets nor cell proliferation were observable 7 days after discontinuing the 3-week exposures, indicating complete recovery. However, after 10 and 22 week periods of exposure, recovery was only partial, labeling indices remaining nearly three times above controls following 10 days in a gasoline-or TMP-free environment. Thus, proximal tubule cell proliferation is a persistent phenomenon in chronic exposure to CIGA, becoming less amenable to recovery with increasing duration of exposure.

Furthermore, in promotion studies with d-limonene, cell proliferation, assessed by bromodeoxyuridine labeling via subcutaneous osmotic minipump implants, was not induced beyond background by d-limonene after 5 or 30 weeks of exposure in the α_{2u} -g-deficient NBR rat, compared to a five-fold increase in the tubule cell labeling of d-limonene-promoted Fischer rats initiated with EHEN (Dietrich and Swenberg, 1991). This result suggests that the sustained proliferative response induced by a CIGA is dependent on the α_{2u} -g syndrome.

Thus, the sequence of events following CIGA administration involves lysosomal overload, cell necrosis, and cell replication. All three of these occur in the same segment of the nephron in conventional strains of rats, but none occur in the NBR rat. Whereas these events are temporally correlated, it is not yet clear whether the lysosomal overload causes necrosis or whether necrosis can be dissociated from replication. These questions need further

investigation and hypothesis development in order to establish mechanisms of action.

C. Progression to cast formation, tubule dilation and mineralization

Since few chronic studies incorporated serial sacrifices, it is difficult to assess the time-dependence of the development and progression of the sequential lesions proposed to be associated with α_{2u} -g nephropathy.

Granular cast formation was recorded exclusively in male rats for most of the selected chemicals evaluated in 13-week toxicity studies by the NTP and sometimes in the 2-year bioassays. In another study, Viau et al. (1986) exposed male rats to C_{10} - C_{12} aliphatic hydrocarbons by inhalation for 5.5, 46, or 68 weeks and found granular casts at the earliest time-point, but they were absent at the later time-points. One explanation for these results is that certain lesions in the sequence are transitory in nature. Granular casts, for example, are assumed to be linked to the active hyaline droplet overload. Once α_{2u} -g levels become low because of age, after approximately 18 months, the number of new hyaline droplets being formed should become minimal. A second explanation is that subtle changes such as granular cast formation and the associated tubule dilation can be obscured by the development of CPN in later stages.

Tubule dilation is presumed to follow obstruction of the nephron by the accumulation of granular casts composed of sloughed epithelial cell debris in the tubule lumen. Figure 6 shows one example of the interrelationships observed between




FIGURE 6: Time-sequence for the development of hyaline droplets (o), regenerating tubule epithelium (+), and tubule dilation (*), in male F-344 rats administered 2 g/kg unleaded gasoline daily by gavage for a 28-day period. (Adapted from Thomas et al., 1985).

hyaline droplet formation, epithelial cell proliferation, and tubule dilation. In this study, male rats were administered unleaded gasoline (2 g/kg/day) for a period of 28 days and examined at 5 interim time-points (Thomas et al., 1985). An initial accumulation of hyaline droplets, commencing on the first day of exposure and persisting throughout, was followed at 14, 21 and 28 days, by increases in epithelial cell proliferation and tubule dilation associated with luminal accumulation of granular debris.

Linear mineralization in the renal papilla of male rats has been consistently observed in a number of NTP and other 2-year bioassays with potential CIGA carcinogens but not in the 13-week toxicity studies. Clear dose-response relationships were demonstrated for 1,4-DCB (NTP, 1987a), JP-4 mixed distillate (MacNaughton and Uddin, 1984), and unleaded gasoline (USEPA, 1987). In the 2-year unleaded gasoline study there were interim sacrifices at 3, 6, 12 and 18 months permitting quantitative observation on the incidence of mineralization (USEPA, 1987). Although this lesion was termed pelvic rather than medullary mineralization in the original report from the IRDC, it was qualified as referring to material located within tubules of the renal pelvis, thus conforming to the medullary site seen with other CIGA. Table 9 presents a summary of these data which shows a clear dose-related progression in the incidence of mineralization from 6 months up to, and including, the 2-year sacrifice. Parallel dose-response increases have been demonstrated for medullary mineralization and urothelial hyperplasia with JP-5 jet fuels, Diesel Fuel Marine and

decalin (Bruner 1986), supporting the notion that the pelvic hyperplasia is a urothelial response to mineralization in the papilla.

TABLE 9. INCIDENCE OF MEDULLARY MINERALIZATION IN MALE RATS DURING INHALATION EXPOSURE TO UNLEADED GASOLINE

Observation Time-Points (months)	Exposure levels of U.G. Vapor (ppm)			
	0	67	292	2056
3	0	0	0	0
6	0	0	0	20*
12	0	0	20	80
18	0	0	20	80
24	0	5	63	91

*The incidence figures are percent of animals affected.

The data was taken from USEPA, 1987.

D. Association between CIGA and chronic progressive nephropathy

Although exacerbation of spontaneous CPN by CIGA has been noted in many studies, quantitation of this response has been attempted on few occasions. Short et al. (1989a) compared the number of CPN foci per kidney section in male rats at three dose-levels of unleaded gasoline exposure and two chronic time-points, with control specimens. For a daily dose range of 0, 10, 70 and 300 ppm unleaded gasoline, the numbers of foci observed at 22 weeks of exposure were 0.4, 0, 1.0 and 6.3 respectively, and at 48 weeks of exposure, 5.0, 4.0, 10.0 and 9.0. This study therefore supports the conclusion that there is an earlier onset of CPN, demonstrable by 5 months, and a higher incidence of disease with the middle and high doses of unleaded gasoline in male rats.

In the NTP bioassay of d-limonene, (NTP, 1990), treated male rats showed a spectrum of compound-related kidney lesions, including exacerbation of CPN, mineralization in the renal medulla, hyperplasia of the epithelium lining the renal papilla, and proliferative lesions of the renal tubule epithelium. The severity of CPN was graded as "not present, minimal, mild, moderate, or marked." The mean value increased with increasing d-limonene dose from 1.5 in vehicle controls to 1.8 and 2.2 in animals dosed at 75 and 150 mg/kg, respectively.

As CPN is exacerbated by CIGA administration, and CPN-affected tubules have a high cell turnover rate, it has been suggested that CPN may play a role in renal tumor production following $\alpha_2\text{u}$ -g nephropathy because enhanced regeneration is considered a risk

factor for carcinogenesis (Trump et al., 1984a; Short et al., 1989b). There is no firm evidence available to date that substantiates or disproves a link between CPN and renal tubule tumor induction. Nevertheless, in a specialized initiation-promotion study with unleaded gasoline and TMP where the authors quantified foci of CPN, some adenomas were described as arising within foci of CPN (Short et al., 1989b).

E. Evidence concerning progression from nephrotoxicity to renal neoplasia

For the eight selected carcinogens examined in this report, there was an overall pattern indicative of dose-related increases in the incidences of toxic nephropathy, hyperplasia, and renal tubule tumors in male rats. For two CIGA, unleaded gasoline and TMP, dose related increases in renal tubule proliferation were sustained throughout chronic administration. It is believed that the likelihood of producing a cancerous cell is increased, not only if there is a probability of a genetic transition, but also if the rate of cell replication is increased (Cohen and Ellwein, 1990; Deal et al., 1989). Thus, a finding that a sustained state of cell turnover in the target cell population is a mechanistic link between α_{2u} -g nephropathy and renal neoplasia should be considered a plausible, but unproven, description of the observed results.

The hyperplastic tubules and adenomas produced by CIGA carcinogens appear to arise from the cortex, which includes the P2 segment of the proximal tubule, the main site of cellular injury in α_{2u} -g nephropathy, providing further support for their linkage. Furthermore, Goldsworthy et al. (1988a) have shown an increase in

cell replication rates specifically in the histologically damaged P2 segments after tetrachloroethylene or pentachloroethane exposure in male rats. Under the same conditions, cell replication did not differ from controls in female rats given these chemicals nor in rats of both sexes treated with a non-CIGA, trichloroethylene.

Recent studies of the promotion potential of d-limonene, TMP, and gasoline also provide convincing evidence to support a linkage between α_{2u} -g nephropathy and renal tubule neoplasia. Dietrich and Swenberg (1991) demonstrated that d-limonene promoted renal tubule tumors in male F-344 rats, an animal that produces α_{2u} -g. In addition, there was a five-fold increase of P2-labeling index in the F-344 rats treated with d-limonene. In contrast, no response was recorded for proliferation, hyperplasia, or renal tubule adenomas in the NBR rat, an α_{2u} -g-deficient animal which does not develop the characteristic nephropathy. These results substantiate those of an earlier study where dose-related increases in atypical cell foci were observed in male rats promoted with unleaded gasoline or TMP for 24 or 60 weeks (Short et al., 1989b). In that study, there was a significant linear trend in incidence of renal tubule tumors in the male rat promoted with unleaded gasoline for 24 weeks. In contrast, none of these changes was observed in similarly treated female rats.

Finally, the nephrotoxicity seen in male rats in the selected two year bioassays of renal tubule carcinogens was characteristic of that proposed to result from cell damage caused by α_{2u} -g accumulation. In contrast, whenever nephrotoxicity was observed in

female rats or mice of either sex, ie. for hexachloroethane, tetrachloroethylene, and 1,4-DCB, the lesions were not characteristic of CIGA and probably were a response caused by an independent mechanism.

PART 3 - EVALUATION OF THE HYPOTHESIS

X. SUMMARY OF THE EVIDENCE ON THE RENAL EFFECTS OF CIGA

Several lines of evidence establish an association between exposure of the male rat to chemicals that induce α_{2u} -g accumulation (CIGA) and nephrotoxicity, and strongly support an association between this nephrotoxicity and renal tubule tumors.

A. Association between α_{2u} -g accumulation and nephropathy

The information that supports an association between α_{2u} -g accumulation and male rat-specific renal toxicity following CIGA administration is summarized below.

(1) Thirty-two organic compounds including fuels, solvents, and other chemicals (listed in Appendix 1), examined in this report, have been shown to induce an excessive accumulation of hyaline droplets in the renal proximal tubule epithelium of male rats. The results in female rats for many of these compounds and the results in mice for about half were also examined, with no finding of hyaline droplet accumulation. However, hyaline droplet accumulation per se is not necessarily diagnostic of a CIGA until proven to represent α_{2u} -g accumulation. Of the 32 substances, the presence of α_{2u} -g has been confirmed in male rats for 17.

(2) There is convincing evidence that the excessive accumulation of hyaline droplets is followed sequentially by tubule epithelial cell necrosis, cast formation and other aspects of α_{2u} -g nephropathy in the male rat. Five of the 32 hyaline-droplet inducers were tested in species other than the mouse or the rat,

although possibly not as rigorously. Characteristic lesions were observed in the male rat kidney for these five substances, but there was no apparent nephrotoxic response in the female rat or any other species tested, which included mice (all 5 substances), hamsters (jet fuels), guinea pigs (decalin), dogs (jet fuels, decalin, d-limonene, and methyl isobutyl ketone), and monkeys (methyl isobutyl ketone and gasoline).

(3) The increase in hyaline droplets, tubule dilation caused by granular cast formation, tubule cell proliferation, and medullary mineralization is dose dependent as shown by research studies conducted to date with four model CIGA (decalin, d-limonene, unleaded gasoline, and TMP).

(4) In general, the chronic administration of CIGA to male rats and the ensuing nephrotoxicity enhanced the age-related renal degenerative process by exacerbating spontaneous CPN.

(5) Specialized studies involving rats of varying age, castrated or estrogen-treated rats, the NBR strain, and α_{2u} -g-treated female rats have shown that development of the early features of α_{2u} -g nephropathy is dependent on the presence of α_{2u} -g formed in the liver.

(6) For three of the eight model carcinogens (hexachloroethane, tetrachloroethylene, and 1,4-DCB), renal toxicity was observed in chronic studies of female rats or mice, but the renal toxicity appeared to be less severe and qualitatively different, not involving the same spectrum of discrete lesions associated with α_{2u} -g nephropathy.

(7) CIGA bind reversibly to α_{2u} -g as a target molecule, and the renal accumulation of α_{2u} -g and hyaline droplet formation may be explained by chemical inhibition of α_{2u} -g catabolism after reabsorption of the complex by the proximal tubule.

(8) TMPOH, the active metabolite of TMP can form in vitro complexes with retinol-binding protein and α_1 -acid glycoprotein, members of the lipocalin superfamily found in humans. In vivo data on retinol and α_{2u} -g, however, demonstrate that such an association does not necessarily lead to α_{2u} -g accumulation or hyaline droplet formation.

B. Association between alpha-2u-globulin nephropathy and renal cancer

Based on information from the rodent bioassays examined in this report and additional key data, features of renal tumors occurring subsequent to the development of nephropathy in the male rat can be identified.

(1) The eight model carcinogens produced hyperplasia, adenomas, and adenocarcinomas in the renal tubule of the male rat.

(2) All eight that produced renal tumors in male rats also produced nephrotoxicity.

(3) In general, the nephrotoxicity that preceded renal tumor formation in male rats was characteristic of the form associated with α_{2u} -g as distinguished from other forms of toxicity associated with non-CIGA renal toxicants.

(4) The incidence of renal tumors produced in the male rat by the eight model carcinogens was relatively low. These tumors were morphologically indistinguishable from renal tubule neoplasia that

occurs rarely, but spontaneously, in male and female rats.

(5) The renal tumors produced by the eight model carcinogens occurred late usually being found at sacrifice, metastasized rarely, and were not life-threatening.

(6) For d-limonene, the one CIGA studied in an initiation-promotion in male rats of the NBR and another strain, α_{2u} -g accumulation was necessary for promotion of male rat renal tubule tumors initiated by EHEN.

(7) CIGA appear to be non-genotoxic or only marginally so and may, therefore, not depend on direct genetic injury as the mechanism for tumor induction.

(8) Trichloroethylene, a compound structurally similar to hexachloroethane and tetrachloroethylene produced renal tumors apparently by mechanisms, such as covalent binding to DNA, which do not appear applicable to the CIGA hypothesis.

C. Information reducing confidence in the conclusion that the α_{2u} -g response is specific to the male rat.

Although the evidence available to date supports the hypothesized association between α_{2u} -accumulation and renal tubule tumors in the male rat, confidence in this assertion would be improved if the same results were found in an expanded database. In addition, the paucity of data on the lipocalin superfamily, in general, leaves several questions unanswered regarding the specificity of the response to the male rat.

(1) Pathological accumulation of hyaline droplets is a reaction to excessive protein load not exclusively related to α_{2u} -g accumulation. Although there are 32 hyaline droplet-inducing

compounds identified in Appendix 1 of this report, the accumulating protein responsible for hyaline droplet formation has not been identified for about half of these compounds.

(2) Data sufficient to demonstrate interdependence of the lesions in the proposed pathological sequence from hyaline droplet accumulation to chronic toxicity exist for only a few substances. Data to define dose-response relationships for tubule cell necrosis and its association with cell proliferation are even more limited, as is dose-related information on increased cell proliferation rates over chronic exposure periods.

(3) Hexachloroethane, tetrachloroethylene, and 1,4-DCB produced renal toxicity in female rats or mice indicating that some CIGA may have additional effects on rodent kidney not limited to the α_{2u} -g-induced sequence of lesions.

(4) Information on a possible association between renal cell tumors and CIGA exposure in humans is inconclusive since exposures in the reviewed epidemiologic studies have been to both CIGA and non-CIGA compounds.

(5) Information on the in vivo binding of CIGA with other lipocalins in the α_{2u} -g superfamily of proteins is too limited to demonstrate conclusively that toxicity in humans does not occur via this mechanism.

(6) Although there are major quantitative and qualitative differences between male rats and humans in the amounts of protein excreted in urine, little is known concerning the relative quantities of low-molecular-weight proteins that are normally

filtered by the human glomerulus and reabsorbed by the renal tubules for catabolism.

(7) The mechanism whereby α_{2u} -g accumulation leads to cell death has not been established.

The scientific data summarized above were used to draw conclusions with regard to the role of α_{2u} -g accumulation and hyaline droplet formation in producing male rat-specific nephropathy and renal tubule neoplasia and to determine the relevance of this information to assessing human risk.

XI. CONCLUSIONS

The available information on CIGA-associated renal tubule carcinogenesis in the male rat can be described by a suggested sequence of critical molecular and cellular events. According to this description, the reaction of a lipophilic compound with the low-molecular-weight protein, α_{2u} -g, appears to lead to the formation of a complex which is more resistant to lysosomal degradation than the unreacted protein. This results in a shift in balance between reabsorption and hydrolysis leading to an abnormal accumulation of the protein in the P2 segment of the renal tubule of male rats. If exposure ceases after a short time period, recovery is complete. Continued exposure, however, results in a nephrotoxic response that is less readily reversible and a sustained increase in cell turnover, enhancing the chance that lesions occurring in the kidney may be replicated rather than repaired.

Because there are substantial data gaps, especially with regard to the expected response in humans and the critical linkages between single cell necrosis and increased cell turnover, and tubule hyperplasia and renal tubule cancer, this α_{2u} -g syndrome should be considered a satisfactory working hypothesis but not a proven mechanism of action to describe renal tubule cancer in male rats exposed to CIGA. As such, it provides an empirical description of a series of observed events in laboratory animals which could be modified or expanded upon as additional information becomes available.

Despite these limitations and the fact that α_{2u} -g accumulation also exacerbates CPN, chemically induced α_{2u} -g-associated nephropathy in the male rat can be distinguished histopathologically from other chemically-induced nephrotoxicities and also from CPN. Excessive hyaline droplet formation is the earliest morphologic manifestation and an important characteristic, although a chemical can be described as a CIGA with certainty only when there is a positive identification of α_{2u} -g in the hyaline droplets. Other observable characteristics indicative of possible CIGA-induced nephrotoxicity include single cell necrosis of the tubule epithelium, granular casts at the corticomedullary junction caused by sloughing of necrotic cells, mitotic figures indicative of regeneration or increased cell turnover, and often medullary mineralization.

The hepatic synthesis of the lipocalin, α_{2u} -g, is not known to occur normally in any species other than the male rat. Alpha-

α_{2u} -globulin-induced nephropathy is also a distinct entity specific to the male rat among the laboratory species and genders tested to date. The characteristic nephropathy has been found only when α_{2u} -g formed in the liver is present. Thus, female rats do not develop hyaline droplets when exposed to CIGA unless they have been administered α_{2u} -g isolated from male rat urine. NBR rats which do not carry the mRNA for liver α_{2u} -g and castrated male rats also respond differently from conventional male rats. Of the other species, the mouse is the most thoroughly tested. Although the mouse produces large amounts of a structurally similar lipocalin, MUP, this protein is not known to bind with CIGA; it is not reabsorbed from the urine; and the mouse does not develop kidney tumors or the characteristic nephropathy seen in male rats. Limited testing in dogs, hamsters, guinea pigs, and monkeys has not shown hyaline droplet accumulation or nephropathy in these species, further suggesting that the α_{2u} -g syndrome occurs specifically in the male rat.

With regard to the potential for a chemical to produce renal tubule neoplasia in the male rat, there are common characteristics among the substances evaluated in this report. First, these compounds (and their CIGA-binding metabolites) possess little or no mutagenic activity in standard batteries of tests, they are lipophiles and not electrophilic substances, and they do not appear to bind covalently to DNA. Second, the nephrotoxic response characteristic of CIGA always preceded renal tumor formation in the male rat, a finding not characteristic of classical renal

carcinogens. Third, for all eight model compounds examined in this report, additional sexes/strains were tested, and the increased incidence of renal tumors was found only in the male rat.

The manner in which the human male responds to CIGA has not been tested directly although there are human proteins that, like α_{2u} -g, are members of the lipocalin superfamily. Human urine contains small amounts of a sex-linked urinary protein. Epidemiological studies have focused on glomerulonephritis or renal cancer and organic chemical exposure, in general, and not on renal tubule damage and CIGA exposure, and they do not yield results useful for testing the hypothesized mechanism in humans. Protein overload can result in formation of hyaline droplets in human kidneys, although there is no evidence that this response has occurred from lipocalin accumulation in the human kidney. While it is not possible to resolve the issue of how the human renal tubule responds to CIGA exposure from the available data, the uniqueness of the male rat response among the tested laboratory species and the high doses needed to produce an effect, even in the male rat, suggest that this reaction would not occur in humans, especially under typical conditions of exposure.

Several factors complicate the analysis of data on the renal effects of CIGA. Unbound moieties, either the CIGA or its metabolites that do not bind to α_{2u} -g, can exist in the kidney along with the protein-bound material. The potential toxicities of these unbound moieties to the kidney need to be taken into account. For example, perchloroethylene, in addition to showing α_{2u} -g

nephropathy, displays evidence of renal toxicity typical of chlorinated hydrocarbons. This example demonstrates how other mechanisms may play some role in the observed results. Since not all CIGA present in the male rat kidney is protein-bound, the possibility that the toxicity of the moieties not bound to α_{2u} -g may also play some role should be kept in mind when evaluating CIGA for purposes of human risk assessment.

At present, there is insufficient information on CIGA and their metabolites to confidently predict activity on the basis of structural analogy. Recent research on structural correlations suggests the presence of an electronegative atom for hydrogen bonding, lipophilicity, and steric volume are important considerations. Conformational changes or other structural alterations to the protein may also be necessary since binding of the compound in the protein pocket, alone, appears to be an insufficient condition to cause reduced digestibility of the protein.

Evidence of dose-responsiveness between CIGA administration and the degree of hyaline droplet or α_{2u} -g formation has been demonstrated in several studies. These findings are frequently based on subjective histopathological criteria, however, limiting their usefulness for making quantitative judgments about the relative hazard potential of different chemicals.

It is also important to recognize that for various reasons (eg., doses administered too low, animals killed before the latency period of these slow growing tumors is attained, number of specimens and histological sections insufficient, competing

toxicity in kidney or other organs), the entire pathological sequence culminating in renal tubule neoplasia may not be demonstrated in all cases of CIGA administration. Thus, not all CIGA would be expected to demonstrate renal tubule neoplasia in the male rat in a 2-year animal bioassay. Such a finding would not negate the applicability of the hypothesized CIGA syndrome to the evaluation of nephropathy data.

Based on the cancer bioassays and other laboratory data, an increased proliferative response caused by chemically-induced cytotoxicity appears to play a role in the development of renal tubule tumors seen in male rats. Among the laboratory animals tested to date, this response to CIGA administration seems to be specific to the male rat. These conclusions can probably be extended to analysis of human hazard potential, especially whenever human exposure to CIGA is not excessively high for sustained periods of time, when short-term tests for genotoxicity of the compound are negative, when the nephrotoxic response and increased cell turnover characteristic of CIGA have been demonstrated in the male rat, and other species/sex combinations were tested but renal tubule tumors were observed only in male rats.

XII. RESEARCH NEEDS

Certain studies, suggested to fill key data gaps, are listed below. There has been no attempt to outline all the possible avenues for research on CIGA and on lipocalins, since a vast array of useful experiments could be envisioned. Instead, recommended studies would greatly improve the data base on these chemicals,

provide needed information to answer questions of human relevance, and set up a framework for improving the testing of chemicals that are potentially male rat renal tubule tumorigens. These research needs are listed as follows.

- (1) Extend studies in humans, wherever possible, to determine directly the effects of hydrocarbon and solvent exposure, focusing on specific jobs known to have relatively pure CIGA exposure. Any human pathology found should be compared with α_{2u} -g nephropathy in the male rat, and urine should be examined for the presence of cells and casts since this noninvasive technique is readily applied to humans.
- (2) Examine human subpopulations that excrete abnormal amounts of low-molecular weight protein in the urine to determine if they are at risk of renal disease or renal cell cancer.
- (3) Examine the binding of CIGA to lipocalins, such as retinol-binding protein, α_1 -acid glycoprotein, and urine protein 1, to be followed with a determination of those complexes that have a slower degradation rate as a result of binding.
- (4) Thoroughly characterize potential protein droplet nephrotoxicity resulting from administration of known CIGA (eg., d-limonene, TMP) to additional species (eg., dog, hamster, rabbit, guinea pig, and especially non-human primate).
- (5) Further characterize the kidney response to CIGA and non-CIGA renal carcinogens in the NBR rat which appears not to synthesize α_{2u} -g. These studies should verify in a two year chronic bioassay that the NBR rat kidney is responsive to classical renal carcinogens already tested in other strains, and they should evaluate the suitability of this strain of rat as a test species. If the NBR rat meets these two criteria, the possibility of employing a separate test group, consisting of male NBR rats, should be considered for conventional bioassays whenever it is suspected that the α_{2u} -g syndrome would influence the results.
- (6) Develop a standard short-term protocol (eg. the 2-week study) to look for the presence of hyaline droplets in the male rat kidney before potential nephrotoxins are placed on chronic study. If hyaline droplets are discovered, this information should be taken into account in designing the chronic study to ensure that the maximum information is attained during the study.

- (7) Serial-sacrifice studies of CIGA and non-CIGA renal carcinogens designed to determine if a distinctly different progression from α_{2u} -g nephropathy to tumor formation can be seen for the CIGA. Studies should involve chronic exposures, examine the histogenesis of the renal cell tumors, and include "stop" experiments and time-dependent appearance of tumor markers.
- (8) Dose-response studies designed to quantitate the relationship between increased hyaline droplets and cell necrosis and between cell necrosis and cell regeneration. In addition, the possibility of additional steps in the progression that might further define the expression of cancer in the male rat and the cause of cell death should be explored.
- (9) Metabolism and disposition studies of CIGA in other species, compared with male rats, to determine the causative chemical for the nephropathy, and to clarify sites of biotransformation and deposition and fate of these compounds.

Additional work, not as critical as the above, but which would also assist in understanding this disease process includes the following.

- (1) Identification of the accumulating material contained in hyaline droplets of proximal tubules for chemicals that are apparent, but unverified CIGA, and 2-year bioassays for decalin and TMP.
- (2) In vitro assays using rodent kidney extracts to more specifically determine mutagenic potential of CIGA (or active metabolites).
- (3) Studies on the genesis of CPN and its relationship to α_{2u} -g nephropathy as well as the possible role of CPN as a co-carcinogenic factor for renal tumor induction.
- (4) More information on the renal catabolism of α_{2u} -g and the rate and efficiency of protease-mediated hydrolysis in control and CIGA-treated rats.
- (5) Studies on the binding relationships between CIGA and α_{2u} -g (e.g. affinity, concentration ranges, binding effectors) and determination of the site at which binding of CIGA to α_{2u} -g occurs (eg., liver, plasma, or urine) to investigate the hypothesis that the protein-CIGA complex is only formed at high concentrations of the chemical.
- (6) Determination of the reasons that the amount of low molecular

weight protein in the human urine is much less than it is in male rats.

P A R T 4. P O L I C Y

THIS PART WILL BE DISTRIBUTED LATER UNDER SEPARATE COVER.

REFERENCES

- Akerstrom B, Logdberg L: 1990. An intriguing member of the lipocalin protein family: α_1 -microglobulin. *TIBS* 15: 240-243.
- Alden CL: 1986. A review of unique male rat hydrocarbon nephropathy. *Toxicol. Pathol.* 14: 109-111.
- Alden CL: 1989. Male rat specific α_{2u} globulin nephropathy and renal tumorigenesis. In: Bach PH, Lock EA (eds.), *Nephrotoxicity. In Vitro to In Vivo. Animals to Man*. Plenum Press, New York, pp. 535-541.
- Alden CL, Chestnut MH, LeBoeuf RA, Burns JL, Eurell TE, Poiley JA, Raineri R: 1991. Exploration of the tumorigenic potential of a unique male rat protein - α_{2u} globulin. *Toxicologist* 11: 137 (abstract).
- Alden CL, Kanerva RL, Ridder G, Stone LC: 1984. The pathogenesis of the nephrotoxicity of volatile hydrocarbons in the male rat. In: Mehlman MA, Hemstreet GP, Thorpe JJ, Weaver NK (eds.), *Advances in Modern Environmental Toxicology*. Vol. VII. Renal Effects of Petroleum Hydrocarbons. Princeton Scientific Publishers, Inc., Princeton, New Jersey, pp. 107-120.
- Alden CL, Ridder G, Stone L, Kanerva RL: 1985. Pathology of petrochemical fuels in male rats. Acute toxicity. In: Bach PH, Lock EA (eds.), *Renal Heterogeneity and Target Cell Toxicity*. John Wiley and Sons, New York, pp. 461-472.
- Asal NR, Risser DR, Kadamani S, Geyer JR, Lee, ET, Cherng N: 1988. Risk factors in renal cell carcinoma: I. Methodology, demographics, tobacco, beverage use, and obesity. *Cancer Detection Prevention* 11: 359-377.
- Ashby J, Tennant RW: 1988. Chemical structure. Salmonella mutagenicity and extent of carcinogenicity as indicators of genotoxic carcinogenesis among 222 chemicals tested in rodents by the U.S. NCI/NTP. *Mutat. Res.* 204: 17-115.
- Askergren A: 1986. Solvents and the kidney. *Progr. Clin. Biol. Res.* 220: 155-167.
- Bachmann S, Sakai T, Kriz W: 1986. Nephron and collecting duct structure in the kidney, rat. In: Jones TC, Mohr U, Hunt RD (eds.), *Urinary System. ILSI Monographs on Pathology of Laboratory Animals*. Springer-Verlag, New York, pp.3-24.
- Bannayam GA, Lamm DL: 1980. Renal cell tumors. *Pathol. Ann.* 15: part 2, 271-308.
- Barnett JW, Johnson DE, Boothe AD, Johannsen FR: 1987. Protein

accumulation as the initial event in tricyanohexane induced nephropathy in male rats. Toxicologist 7: 27 (abstract).

Barrientos A, Ortuno MT, Morales JM, Martinez Tello F, Rodicio JL: 1977. Acute renal failure after use of diesel fuel as shampoo. Arch. Intern. Med. 137: 1217.

Barthold SW: 1979. Chronic progressive nephropathy in aging rats. Toxicologic Pathol. 7: 1-6.

Bennington JL, Beckwith JB: 1975. Tumors of the Kidney, Renal Pelvis, and Ureter. Atlas of Tumor Pathology. Second Series, Fascicle 12. Armed Forces Institute of Pathology, Washington, DC, pp. 93-199.

Bennington JL, Ferguson BL, Campbell PB: 1968. Epidemiologic studies of carcinoma of the kidney. II. Association of renal adenoma with smoking. Cancer 22: 821-823.

Bell ET: 1950. Renal Diseases. Lea and Febiger, Philadelphia, p. 428.

Berggard I: 1970. Plasma proteins in normal human urine. In: Manuel Y, Revillard JP, Betuel H (eds.), Proteins in normal and pathological urine. S. Karger, Basel, pp. 7-19.

Bernard AM, Lauwerys RR, Noel A, Vandeleene B, Lambert A: 1989. Urine protein 1: a sex dependent marker of tubular or glomerular dysfunction. Clin. Chem. 35: 2141-2142.

Bernard A, Lauwerys R, Noel A, Vandeleene B, Lambert A: 1990. Determination by latex immunoassay of protein 1 in normal and pathological urine. Submitted to Clin Chem.

Blair A, Decoufle P, Grauman D: 1979. Causes of death among laundry and dry cleaning workers. Am. J. Public Health 69: 508-511.

Blair A, Stewart PA, Tolbert PE, Grauman D, Moran FX, Vaught J, Rayner J: 1990. Cancer and other causes of death among a cohort of dry cleaners. Brit. J. Industr. Med. 47: 162-168.

Bomhard E, Luckhaus G, Voight W-H, Loeser E: 1988. Induction of light hydrocarbon nephropathy by p-dichlorobenzene. Arch. Toxicol. 61: 433-439.

Bomhard E, Marsmann M, Ruhl-Fehlert CH, Zyweitz A: 1990. Relationships between structure and induction of hyaline droplet accumulation in the renal cortex of male rats by aliphatic and alicyclic hydrocarbons. Arch. Toxicol. 64: 530-538.

Borghoff SJ, Miller AB, Bowen JP, Swenberg JA: 1991.

Characteristics of chemical binding to α_{2u} -globulin in vitro: evaluating structure-activity relationships. Toxicol. Appl. Pharmacol. 107, 228-238.

Borghoff SJ, Short BG, Swenberg JA: 1990. Biochemical mechanisms and pathobiology of α_{2u} -globulin nephropathy. Ann. Rev. Pharmacol. Toxicol. 30: 349-367.

Borghoff SJ, Strasser J, Charbonneau M, Swenberg JA: 1988. Analysis of 2,4,4-trimethyl-2-pentanol (TMP-OH) binding to male rat kidney α_{2u} -globulin (α_{2u}) and other proteins. Toxicologist 8: 135 (abstract).

Borghoff SJ, Upton PB, Swenberg JA: 1989. Characteristics of 2,4,4-trimethyl-2-pentanol (TMP-OH) binding to α_{2u} -globulin and other compounds that cause protein droplet nephropathy. Toxicologist 9: 79 (abstract).

Brooks DE: 1987. The major androgen-regulated secretory proteins of the rat epididymis bear sequence homology with members of the α_{2u} -globulin superfamily. Biochem. Internat. 14: 235-240.

Brownson RC: 1988. A case-control study of renal cell carcinoma in relation to occupation, smoking, and alcohol consumption. Arch. Environ. Health. 43: 238-241.

Bruner RH: 1984. Pathologic findings in laboratory animals exposed to hydrocarbon fuels of military interest. In: Mehlman MA, Hemstreet GP, Thorpe JJ, Weaver NK (eds.), Advances in Modern Environmental Toxicology. Vol. VII. Renal Effects of Petroleum Hydrocarbons. Princeton Scientific Publishers, Inc., Princeton, New Jersey, pp. 133-140.

Bruner RH: 1986. Chronic consequence of α_{2u} -globulin nephropathy: review. Proceedings, Toxicology Forum, Aspen, Colorado, July 15, pp. 161-166.

Bruckner JV, Davis BD, Blancato JN: 1989. Metabolism, toxicity, and carcinogenicity of trichloroethylene. CRC Critical Rev. Toxicol. 20: 31-50.

Burnett VL, Short BG, Swenberg JA: 1989. Localization of α_{2u} -globulin within protein droplets of male rat kidney: immunohistochemistry using perfusion-fixed, GMA-embedded tissue sections. J. Histochem. Cytochem. 37: 813-818.

Caudill D, Asquith TN, Lehman-McKeeman LD: 1991. Biochemical differences between α_{2u} -globulin (α_{2u}) and mouse urinary protein (MUP): explanation of murine resistance to develop hyaline droplet nephropathy (HDN). Toxicologist 11: in press (Abstract).

Cavaggioni A, Findlay JBC, Tirindelli R: 1990: Ligand binding

characteristics of homologous rat and mouse urinary proteins and pyrazine-binding protein of calf. Comp. Biochem. Physiol. 96B: 513-520.

Cavaggioni A, Sorbi RT, Keen JN, Pappin DJC, Findlay JBC: 1987. Homology between the pyrazine-binding protein from nasal mucosa and major urinary proteins. FEBS Lett. 212: 225-228.

Charbonneau M, Lock EA, Strasser J, Cox MG, Turner MJ, Bus JS: 1987. 2,2,4-Trimethylpentane-induced nephrotoxicity. I. Metabolic disposition of TMP in male and female Fischer 344 rats. Toxicol. Appl. Pharmacol. 91: 171-181.

Charbonneau M, Strasser J, Borghoff SJ, Swenberg JA: 1988. In vitro hydrolysis of [¹⁴C]- α_2 u-globulin (α_2 u) isolated from male rat kidney. Toxicologist 8: 135 (abstract).

Charbonneau M, Strasser J, Lock EA, Turner MJ, Swenberg JA: 1989. Involvement of reversible binding to α_2 u-globulin in 1,4-dichlorobenzene-induced nephrotoxicity. Toxicol. Appl. Pharmacol. 99: 122-132.

Chatterjee B, Demyan WF, Song CS, Garg BD, Roy AK: 1989. Loss of androgenic induction of α_2 u-globulin gene family in the liver of NIH Black rats. Endocrinology 125: 1385-1388.

Chatterjee B, Hopkins J, Dutchak D, Roy AK: 1979. Superinduction of α_2 u-globulin by actinomycin D: evidence for drug-mediated increase in α_2 u mRNA. Proc. Natl. Acad. Sci. 76: 1833-1837.

Churchill DN, Fine A, Gault MH: 1983. Association between hydrocarbon exposure and glomerulonephritis. An appraisal of the evidence. Nephron 33: 169-172.

Cohen SM, Ellwein LB: 1990. Cell proliferation in carcinogenesis. Science 249: 1007-1011.

Cohen C, McCue PA, Derosé PB: 1988. Histogenesis of renal cell carcinoma and renal oncocytoma. An immunohistochemical study. Cancer 62: 1946-1951.

Crisp AJ, Bhalla AK, Hoffbrand BI: 1979. Acute tubular necrosis after exposure to diesel oil. Brit. Med. J. 2: 177.

Daniell WE, Couser WG, Rosenstock L: 1988. Occupational solvent exposure and glomerulonephritis. A case report and review of the literature. J. Am. Med. Assoc. 259: 2280-2283.

Dayal H, Kinman J: 1983. Epidemiology of kidney cancer. Semin. Oncol. 10: 366-377.

Deal FH, Richardson FC, Swenberg, JA: 1989. Dose response of

hepatocyte replication in rats following continuous exposure to diethylnitrosamine. Cancer Res. 49: 6985-6988.

Dees JH, Heatfield BM, Reuber MD, Trump BF: 1980a. Adenocarcinoma of the kidney. III. Histogenesis of renal adenocarcinomas in rats by N-(4'-fluoro-4-biphenyl)acetamide. J. Natl. Cancer Inst. 64: 1537-1545.

Dees JH, Heatfield BM, Trump BF: 1980b. Adenocarcinoma of the kidney. IV. Electronmicroscopic study of the development of renal adenocarcinomas induced in rats by N-(4'-fluoro-4-biphenyl)acetamide. J. Natl. Cancer Inst. 64: 1547-1562.

Dekant W, Vamvakas S, Anders MW: 1989. Bioactivation of nephrotoxic haloalkanes by glutathione conjugation -- formation of toxic and mutagenic intermediates by cysteine conjugate β -lyase. Drug Metab. Rev. 20: 43-83.

Devesa SS, Silverman DT, McLaughlin JK, Brown CC, Connelly RR, Fraumeni JF: 1990. Comparison of the descriptive epidemiology of urinary tract cancers. Cancer Causes and Control 1: 133-141.

Dice JF: 1987. Molecular determinants of protein half-lives in eukaryotic cells. FASEB J. 1: 349-357.

Dieppe PA, Doyle DV, Burry HC: 1978. Renal damage during treatment with antirheumatic drugs. Brit. Med. J. 2: 664.

Dietrich DR, Swenberg JA: 1990a. NBR male rats fail to develop renal disease following exposure to agents that induce alpha-2u-globulin (A2u) nephropathy. Toxicologist 10: 266 (abstract).

Dietrich DR, Swenberg JA: 1990b. Lindane induces nephropathy and renal accumulation of α 2u-globulin in male but not in female Fischer 344 rats or male NBR rats. Toxicol. Lett. 53: 179-181.

Dietrich DR, Swenberg JA: 1990. NBR male rats fail to develop renal disease following exposure to agents that induce alpha-2u-globulin (A2u) nephropathy. Toxicologist 10: 266 (abstract).

Dietrich DR, Swenberg JA: 1991. Alpha $_{2u}$ -globulin is necessary for d-limonene promotion of male rat kidney tumors. Toxicologist 11: In press (abstract).

Dodd DE, Losco PE, Troup CM, Pritts IM, Tyler TR: 1987. Hyaline droplet nephrosis in male Fischer 344 rats following inhalation of diisobutyl ketone. Toxicol. Indust. Health 3: 443-457.

Dolan KP, Unterman R, McLaughlin M, Nakhasi HL, Lynch KR, Feigelson P: 1982. The structure and expression of very closely related members of the α_{2u} -globulin gene family. J. Biol. Chem. 257: 13527-13534.

Dominick MA, Robertson DG, Sifler MR, Susick RL, Bobrowski WF: 1990. Absence of nephrocarcinogenesis in male rats administered gabapentin, a compound that causes alpha-2u-globulin nephropathy. Proceedings of the American College of Veterinary Pathology, December 9-14, 1990, p. 45 (abstract).

Duh R-W, Asal NR: 1984. Mortality among laundry and dry cleaning workers in Oklahoma. Am. J. Public Health 74: 1278-1280.

Ekstrom B, Berggard I: 1977. Human α_1 -microglobulin: purification procedure, chemical and physicochemical properties. J. Biol. Chem. 252: 8048-8057.

Ekstrom RC: 1983. Characterization and metabolism of α_{2u} -globulin - a male sex-dependent protein of the rat. Ph.D. thesis, University of Wisconsin-Madison.

Elcombe CR, Rose MS, Pratt IS: 1985. Biochemical, histological, and ultrastructural changes in rat and mouse liver following the administration of trichloroethylene: possible relevance to species differences in hepatocarcinogenicity. Toxicol. Appl. Pharmacol. 79: 365-376.

Enterline PE, Viren J: 1985. Epidemiologic evidence for an association between gasoline and kidney cancer. Environ. Health Persp. 62: 303-312.

Evans GO, Goodwin DA, Parsons CE, Read NG: 1988. The effects of levamisole on urinary enzyme measurements and proximal tubule cell inclusions in male rats. Brit. J. Exp. Pathol. 69: 301-308.

Evans GO, Morgan RJI: 1986. Urinary enzyme measurements in male rats after oral administration of decalin. Human Toxicol. 5: 120 (abstract).

Fan CY, Butler WH, O'Connor PJ: 1989. Cell and tissue specific localization of O^6 -methylguanine in the DNA of rats given N-nitrosodimethylamine: effects of protein deficient and normal diets. Carcinogenesis 10: 1967-1970.

Farber E: 1988. Cancer development and its natural history. A cancer prevention perspective. Cancer 62: 1676-1679.

Feigelson P, Kurtz DT: 1977. Hormonal modulation of specific messenger RNA species in normal and neoplastic rat liver. Adv. Enzymol. 47: 275-312.

Flamm WG, Lehman-McKeeman LD: 1991. The human relevance of the renal tumor-inducing potential of d-limonene in male rats: implications for risk assessment. Regulatory Toxicol. Pharmacol. In press.

Fowlie AJ, Grasso P, Bridges JW: 1987. Renal and hepatic lesions induced by 2,2,4-trimethylpentane. J. Appl. Toxicol. 7: 335-341.

Galloway SM, Armstrong MJ, Reuben C, Colman S, Brown B, Cannon C, Bloom AD, Nakamura F, Ahmed M, Duk S, Rimpo J, Margolin BH, Resnick MA, Anderson B, Zeiger E: 1987a. Chromosome aberrations and sister chromatid exchanges in Chinese hamster ovary cells--evaluations of 108 chemicals. Environ. Mol. Mutagen. 10(Suppl. 10): 1-175.

Galloway SM, Deasy DA, Bean CL, Kraynak AR, Armstrong MJ, Bradley MO: 1987b. Effects of high osmotic strength on chromosome aberrations, sister-chromatid exchanges and DNA strand breaks, and the relation to toxicity. Mutat. Res. 189: 15-25.

Garg BD, Olson MJ, Demyan WF, Roy AK: 1988. Rapid post-exposure decay of α_{2u} -globulin and hyaline droplets in the kidneys of gasoline-treated male rats. J. Toxicol. Env. Health 24: 145-160.

Garg BD, Olson MJ, Li LC: 1989a. Phagolysosomal alterations induced by unleaded gasoline in epithelial cells of the proximal convoluted tubules of male rats: effect of dose and treatment duration. J. Toxicol. Env. Health 26: 103-116.

Garg BD, Olson MJ, Li LC, Mancini MA, Roy AK: 1987. Immunoelectron microscopic localization of α_{2u} -globulin in the male rat kidney after gasoline exposure. In: Bailey GW (ed.), Proceedings of the 45th Annual Meeting of the Electron Microscope Society of America. San Francisco, California. pp. 872-873.

Garg BD, Olson MJ, Li LC, Mancini MA, Roy AK: 1989b. Estradiol pretreatment of male rats inhibits gasoline-induced renal hyaline droplet and alpha-2u-globulin accumulation. Research Publication, General Motors Research Laboratories. GMR-6557. Dated March 1989.

Gaworski CL, Haun CC, MacEwen JD, Vernot EH, Bruner RH, Amster RL, Cowan MJ, Jr.: 1985a. A 90-day vapor inhalation toxicity study of decalin. Fund. Appl. Toxicol. 5: 785-793.

Gaworski CL, MacEwen JD, Vernot EH, Bruner RH, Cowan MJ: 1984. Comparison of the subchronic inhalation toxicity of petroleum and oil shale JP-5 jet fuels. In: Mehlman MA, Hemstreet GP, Thorpe JJ, Weaver NK (eds.), Advances in Modern Environmental Toxicology. Vol. VII. Renal Effects of Petroleum Hydrocarbons. Princeton Scientific Publishers, Inc., Princeton. pp. 3-47.

Gaworski CL, MacEwen JD, Vernot EH, Haun CC, Leahy HF, Bruner RH, Hall A, Latendresse JR, Mattie DR, Pitts LL: 1985b. Evaluation of 90-day inhalation toxicity of petroleum and oil shale diesel fuel marine (DFM). Harry G. Armstrong Aerospace Medical Research Laboratory, AAMRL-TR-85-074, NMRI 85-57, December 1985.

Geertzen HGM, van den Ouderaa FJG, Kassenaar AAH: 1973. Isolation

and metabolism of male sex-dependent urinary protein from rats. *Acta Endocrinologica* 72: 197-208.

Goldsworthy TL, Lyght O, Burnett VL, Popp JA: 1988a. Potential role of α 2u-globulin, protein droplet accumulation, and cell replication in the renal carcinogenicity of rats exposed to trichloroethylene, perchloroethylene, and pentachloroethane. *Toxicol. Appl. Pharmacol.* 96: 367-379.

Goldsworthy TL, Popp JA: 1987. Chlorinated hydrocarbon-induced peroxisomal enzyme activity in relation to species and organ carcinogenicity. *Toxicol. Appl. Pharmacol.* 88: 225-233.

Goldsworthy TL, Smith-Oliver T, Loury DJ, Popp JA, Butterworth BE: 1988b. Assessment of chlorinated hydrocarbon-induced genotoxicity and cell replication in rat kidney cells. *Environ. Mol. Mutagen.* 11(Suppl. 11): 39.

Goodman DG, Ward JM, Squire RA, Paxton MB, Reichardt WD, Chu KC, Linhart MS: 1980. Neoplastic and non-neoplastic lesions in aging Osborne-Mendel rats. *Toxicol. Appl. Pharmacol.* 55: 433-447.

Goodman MT, Morgenstern H, Wynder EL: 1986. A case-control study of factors affecting the development of renal cell cancer. *Am. J. Epidemiol.* 124: 926-941.

Godovac-Zimmermann J: The structural motif of β -lactoglobulin and retinol-binding protein: a basic framework for binding and transport of small hydrophobic molecules? 1988. *TIBS* 13: 64-66.

Gray JE: 1986. Chronic progressive nephrosis, rat. In: Jones TC, Mohr U, Hunt RD (eds.), *Urinary System. ILSI Monographs on Pathology of Laboratory Animals.* Springer-Verlag, New York, pp. 174-179.

Grisham JW, Kaufmann WK, Kaufmann DG: 1983. The cell cycle and chemical carcinogenesis. *Survey Synth. Pathol. Res.* 1: 49-66.

Green T, Odum J, Nash JA, Foster JR: 1990. Perchloroethylene-induced rat kidney tumors: an investigation of the mechanisms involved and their relevance to humans. *Toxicol. Appl. Pharmacol.* 102: 77-89.

Halder CA, Holdsworth CE, Cockrell BY, Piccirillo VJ: 1985. Hydrocarbon nephropathy in male rats: identification of the nephrotoxic components of unleaded gasoline. *Toxicol. Indust. Health* 1: 67-87.

Halder CA, Warne TM, Hatoum NS: 1984. Renal toxicity of gasoline and related petroleum naphthas in male rats. In: Mehlman MA, Hemstreet GP, Thorpe JJ, Weaver NK (eds.), *Advances in Modern Environmental Toxicology. Vol. VII. Renal Effects of Petroleum*

Hydrocarbons. Princeton Scientific Publishers, Inc., Princeton, New Jersey, pp. 73-88.

Hanis NM, Holmes TM, Shallenberger LG, Jones KE: 1982. Epidemiologic study of refinery and chemical plant workers. J. Occup. Med. 24: 203-212.

Hard GC: 1984. High frequency, single-dose model of renal adenoma/carcinoma induction using dimethylnitrosamine in Crl:(W)BR rats. Carcinogenesis 5: 1047-1050.

Hard, GC: 1987. Chemically induced epithelial tumors and carcinogenesis of the renal parenchyma. In: Bach PH, Lock EA (eds.) Nephrotoxicity in the Experimental and the Clinical Situation, Part I. Martinus Nijhoff Publishers, Lancaster, pp. 211-250.

Hard, GC: 1990. Tumors of the kidney, renal pelvis and ureter. In: Turusov VS, Mohr U (eds.), Pathology of Tumors in Laboratory Animals. Vol. 1 - Tumors of the Rat. Second Edition. International Agency for Research on Cancer, Lyon. IARC Scientific Publications No.99 (in press).

Hard GC, Alden CL, Stula EF, Trump BF: 1991. Proliferative lesions of the kidney in rats. Society of Toxicologic Pathologists. Standard System of Nomenclature and Diagnostic Pathology. In press.

Hard GC, Butler WH: 1971. Morphogenesis of epithelial neoplasms induced in the rat kidney by dimethylnitrosamine. Cancer Res. 31: 1496-1505.

Hard GC, Mackay RL, Kochhar OS: 1984. Electron microscopic determination of the sequence of acute tubular and vascular injury induced in the rat kidney by a carcinogenic dose of dimethylnitrosamine. Lab. Invest. 50: 659-672.

Hard GC, Snowden RT: 1991. Hyaline droplet accumulation in rodent kidney proximal tubules: an association with histiocytic sarcoma. Toxicol. Pathol. (submitted).

Harrington JM, Whitby H, Gray CN, Reid FJ, Aw TC, Waterhouse JA: 1989. Renal disease and occupational exposure to organic solvents: a case-referent approach. Brit. J. Industr. Med. 46: 643-650.

Haseman JK, Huff J, Boorman GA: 1984. Use of historical control data in carcinogenicity studies in rodents. Toxicologic Pathol. 12: 126-135.

Hastie ND, Held WA, Toole JJ: 1979. Multiple genes coding for the androgen-regulated major urinary proteins of the mouse. Cell 17: 449-457.

Haworth S, Lawlor T, Mortelmans K, Speck W, Zeiger E: 1983. Salmonella mutagenicity test results for 250 chemicals. Environ. Mutagen. 5(Suppl. 1): 1-142.

HEI: 1985. Gasoline Vapor Exposure and Human Cancer: Evaluation of Existing Scientific Information and Recommendations for Further Research. Health Effects Institute, Report of the Institute's Health Review Committee, September 1985.

HEI: 1988. An Update on Gasoline Vapor Exposure and Human Cancer: An Evaluation of Scientific Information Published Between 1985 and 1987. Health Effects Institute, Report of the Institute's Health Review Committee, January 6, 1988.

Hellsten S, Berge T, Wehlin L: 1981. Unrecognised renal cell carcinoma. Clinical and pathological aspects. Scand. J. Urol. Nephrol. 8: 273-278.

Higginson J, Muir CS, Buffler PA: 1984. The epidemiology of renal carcinoma in humans with a note on the effect of exposure to gasoline. In: Mehlmán MA, Hemstreet GP, Thorpe JJ, Weaver NK (eds.), Advances in Modern Environmental Toxicology. Vol. VII. Renal Effects of Petroleum Hydrocarbons. Princeton Scientific Publishers, Inc., Princeton, New Jersey, pp. 203-226.

Hobson DW, Mattie DR, Bruner RH, Uddin DE, Eurell T, Olson CT: 1986. Effect of castration on the development of 2,2,4-trimethylpentane-induced hydrocarbon nephropathy in the male F-344 rat. Toxicologist 6: 173 (abstract).

Hoel DG, Haseman JK, Hogan MD, Huff J, McConnell EE: 1988. The impact of toxicity on carcinogenicity studies: implications for risk assessment. Carcinogenesis 9: 2045-2052.

Hughson MD, Buchwold D, Fox M: 1986. Renal neoplasia and acquired cystic kidney disease in patients receiving long term dialysis. Arch. Pathol. Lab. Med. 110: 592-600.

Jackson MR: 1974. The nature of dimethylnitrosamine induced enlargement of rat hepatocyte nuclei. J. Pathol. 113: 197-208.

Jedrzejewski K, Kugler P: 1982. Peptidases in the kidney and urine of rats after castration. Histochemistry 74: 63-84.

Johnson DE: 1987. Protein induced nephropathy in male rats. In: Holcenberg JS, Winkelhake JL (eds.), The Pharmacology and Toxicology of Proteins. Alan R. Liss Inc., New York, pp.165-171.

Kadamani S, Asal NR, Nelson RY: 1989. Occupational hydrocarbon exposure and risk of renal cell carcinoma. Am. J. Industr. Health 15: 131-141.

- Kanerva RL, McCracken MS, Alden CL, Stone LC: 1987a. Morphogenesis of decalin-induced renal alterations in the male rat. *Fd. Chem. Toxicol.* 25: 53-61.
- Kanerva RL, Ridder GM, Lefever FR, Alden CL: 1987b. Comparison of short-term renal effects due to oral administration of decalin or d-limonene in young adult male Fischer-344 rats. *Fd. Chem. Toxicol.* 25: 345-353.
- Kanerva RL, Ridder GM, Stone LC, Alden CL: 1987c. Characterization of spontaneous and decalin-induced hyaline droplets in kidneys of adult male rats. *Fd. Chem. Toxicol.* 25: 63-82.
- Katz GV, Renner ER, Terharr CJ: 1986. Subchronic inhalation toxicity of methyl isoamyl ketone in rats. *Fund. Appl. Toxicol.* 76: 498-505.
- Katz RM, Jowett D: 1981. Female laundry and dry cleaning workers in Wisconsin: a mortality analysis. *Am. J. Public Health*, 71: 305-307.
- Kaysen GA, Myers BD, Couser WG, Rabkin R, Felts JM: 1986. Biology of Disease. Mechanisms and consequences of proteinuria. *Lab. Invest.* 54: 479-498.
- Killeen JC, Wilson NH, Ford WH, Siou G, Busey WM, Eilrich GL: 1990. Progression of renal and forestomach effects following administration of chlorothalonil to rats. *Toxicologist* 10: 269 (abstract).
- Kimura H, Odani S; Arakawa M, Ono T. 1989. Kidney fatty acid-binding protein. *FEBS Letters* 246: 102-104.
- Kloss MW, Cox MG, Norton RM, Swenberg JA, Bus JA: 1985. Sex-dependent differences in the disposition of [¹⁴C-5]2,2,4-trimethylpentane in Fischer 344 rats. In: Bach PH, Lock EA (eds.), *Renal Heterogeneity and Target Cell Toxicity*. Wiley, Chichester, pp. 489-492.
- Kluwe WM, Abdo KM, Huff J: 1984. Chronic kidney disease and organic chemical exposures: evaluations of causal relationships in humans and experimental animals. *Fund. Appl. Toxicol.* 4: 889-901.
- Kugler P, Vornberger: 1986. Renal cathepsin-B activities in rats after castration and treatment with sex hormones. *Histochem.* 85: 157-161.
- Kuna RA, Ulrich CE: 1984. Subchronic inhalation toxicity of two motor fuels. *J. Am. Coll. Toxicol.* 3: 217-229.
- Kurtz DT, Feigelson P: 1977. Multihormonal induction of hepatic α_2 -globulin mRNA as measured by hybridization to complementary

- DNA. Proc. Natl. Acad. Sci. 74: 4791-4795.
- Lehman-McKeeman LD, Rivera-Torres MI, Caudill D: 1990a. Lysosomal degradation of α_2 u-globulin and α_2 u-globulin-xenobiotic conjugates. Toxicol. Appl. Pharmacol. 103: 539-548.
- Lehman-McKeeman LD, Caudill D, Takigiku R, Schneider RE, Young JA: 1990b. Comparative disposition of d-limonene in rats and mice: relevance to male-rat-specific nephrotoxicity. Toxicol. Lett. 53: 193-195.
- Lehman-McKeeman LD, Rodriguez PA, Caudill D, Fey ML, Eddy CL, Asquith TN: 1990. Male rat-specific nephrotoxicity resulting from exposure to 3,5,5-trimethyl hexanoic acid (TMHA). Toxicologist 10: 178. (Abstract).
- Lehman-McKeeman LD, Rodriguez PA, Takigiku R, Caudill D, Fey ML: 1989. d-Limonene-induced male rat-specific nephrotoxicity: evaluation of the association between d-limonene and α_2 u-globulin. Toxicol. Appl. Pharmacol. 99: 250-259.
- Lipsky MM, Trump BF: 1988. Chemically induced renal epithelial neoplasia in experimental animals. Intl. Rev. Exptl. Pathol. 30: 357-383.
- Lock EA, Charbonneau M, Strasser J, Swenberg JA, Bus JS: 1987a. 2,2,4-Trimethylpentane-induced nephrotoxicity. II. The reversible binding of a TMP metabolite to a renal protein fraction containing α_2 u-globulin. Toxicol. Appl. Pharmacol. 91: 182-192.
- Lock EA, Stonard MD, Elcombe CR: 1987b. The induction of W and β -oxidation of fatty acids and effect on α_2 u-globulin content in the liver and kidney of rats administered 2,2,4-trimethylpentane. Xenobiotica 17: 513-522.
- Logothetopoulos J, Weinbren K: 1955. Naturally occurring protein droplets in the proximal tubule of the rat's kidney. Brit. J. Exptl. Pathol. 36: 402-406.
- Loury DJ, Smith-Oliver T, Butterworth BE: 1987. Assessment of unscheduled and replicative DNA synthesis in rat kidney cells exposed in vitro or in vivo to unleaded gasoline. Toxicol. Appl. Pharmacol. 87: 127-140.
- Maack T, Park CH, Camargo MJF: 1985. Renal filtration, transport, and metabolism of proteins. In: Seldin DW, Giebisch G (eds.), The Kidney: Physiology and Pathophysiology. Raven Press, New York, pp. 1773-1803.
- MacFarland HN: 1984. Xenobiotic induced kidney lesions: hydrocarbons. The 90-day and 2-year gasoline studies. In: Mehlman MA, Hemstreet GP, Thorpe JJ, Weaver NK (eds.), Advances in Modern

Environmental Toxicology. Vol. VII. Renal Effects of Petroleum Hydrocarbons. Princeton Scientific Publishers, Inc., Princeton, New Jersey, pp. 51-56.

MacInnes JJ, Nozik ES, Kurtz DT: 1986. Tissue-specific expression of the rat α_{2u} globulin gene family. Molec. Cell. Biol. 6: 3563-3567.

MacNaughton MG, Uddin DE: 1984. Toxicology of mixed distillate and high-energy synthetic fuels. In: Mehlman MA, Hemstreet GP, Thorpe JJ, Weaver NK (eds.), Advances in Modern Environmental Toxicology. Vol. VII. Renal Effects of Petroleum Hydrocarbons. Princeton Scientific Publishers, Inc., Princeton, New Jersey, pp. 121-132.

Mancini M, Majumdar D, Chatterjee B, Roy AK. 1989. α_{2u} globulin in modified sebaceous glands with pheromonal functions--localization of the protein and its mRNA in preputial, Meibomian, and perianal glands. J. Histochem. Cytochem. 37: 149-157.

Masoro EJ, Yu BP: 1989. Diet and nephropathy. Lab. Invest. 60: 165-167.

Mattie DR, Bruner RH, Hoeflich TJ, Kendrick JM: 1988. Effects of decalin and JP-10 on the function and morphology of male rat kidneys. Toxicologist 8: 187 (abstract).

Maunsbach AB: 1966a. Observations on the segmentation of the proximal tubule in the rat kidney. Comparison of results from phase contrast, fluorescence and electron microscopy. J. Ultrastruct. Res. 16: 239-258.

Maunsbach AB: 1966b. Electron microscopic observations of cytoplasmic bodies with crystalline patterns in rat kidney proximal tubule cells. J. Ultrastruct. Res. 14: 167-189.

McCredie M, Ford JM, Stewart JH: 1988. Risk factors for cancer of the renal parenchyma. Int. J. Cancer 42: 13-16.

McGregor DB, Brown A, Cattnach P, Edwards IA, McBride D, Riach V, Caspary WJ: 1988. Responses of the L5178Y tk /tk mouse lymphoma cell forward mutation assay. III. 72 coded chemicals. Environ. Mol. Mutagen. 12; 85-154.

McLaughlin JK: 1984. Risk factors from a population-based case-control study of renal cancer. In: Mehlman MA, Hemstreet GP, Thorpe JJ, Weaver NK (eds.), Advances in Modern Environmental Toxicology. Vol. VII. Renal Effects of Petroleum Hydrocarbons. Princeton Scientific Publishers, Inc., Princeton, New Jersey, pp. 227-244.

McLaughlin JK, Blot WJ, Mehl ES, Stewart PA, Venable FS, Fraumeni JF: 1985. Petroleum-related employment and renal cell cancer. J. Occup. Med. 27: 672-674.

McLaughlin JK, Malaker HSR, Stone BJ, Weiner JA, Malaker BK, Ericsson JLE, Blot WJ, Fraumeni JF: 1987. Occupational risks for renal cancer in Sweden. *Brit. J. Industr. Med.* 44: 119-123.

McLaughlin JK, Mandel JS, Blot WJ, Schuman LM, Mehl ES, Fraumeni JF: 1984. A population-based case-control study of renal cell carcinoma. *J. Natl. Cancer Inst.* 72: 275-284.

McLaughlin JK, Schuman LM: 1983. Epidemiology of renal cell carcinoma. *Rev. Cancer Epidemiol.* 2: 170-210.

Mortelmans K, Haworth S, Lawlor T, Speck W, Tainer B, Zeiger E: 1986. Salmonella mutagenicity tests. II. Results from the testing of 270 chemicals. *Environ. Mutagen.* 8 (Suppl. 7): 1-119.

Motwani NM, Caron D, Demuan WF, Chatterjee B, Hunter S, Poulik MD, Roy AK: 1984. Monoclonal antibodies to α_{2u} -globulin and immunocytofluorometric analysis of α_{2u} -globulin-synthesizing hepatocytes during androgenic induction and aging. *J. Biol. Chem.* 259: 3653-3657.

Muggia FM, Heinemann HO, Farhangi M, Osserman EF: 1969. Lysozymuria and renal tubular dysfunction in monocytic and myelomonocytic leukemia. *Am. J. Med.* 47: 351-366.

Murty CVR, Olson MJ, Garg BD, Roy AK: 1988. Hydrocarbon-induced hyaline droplet nephropathy in male rats during senescence. *Toxicol. Appl. Pharmacol.* 96: 380-392.

Murty CVR, Sarkar FH, Mancini MA, Roy AK: 1987. Sex-independent synthesis of α -2 μ -globulin and its messenger ribonucleic acid in the rat preputial gland--biochemical and immunocytochemical analysis. *Endocrinology* 121: 1000-1005.

NCI: 1978a. Bioassay of chlorothalonil for possible carcinogenicity. Carcinogenesis Testing Program, National Cancer Institute. DHEW Publication No. (NIH) 78-841 (NTP-TR-41).

NCI: 1978b. Bioassay of hexachloroethane for possible carcinogenicity. Carcinogenesis Testing Program, National Cancer Institute. CAS No. 67-72-1. DHEW Publication No. (NIH) 78-1318. (NTP-TR-68).

NCI: 1987. Annual Cancer Statistics Review. National Cancer Institute, US DHHS. NIH Publication No. 88-2789.

Neuhaus OW, Flory W: 1978. Age-dependent changes in the excretion of urinary proteins by the rat. *Nephron* 22: 570-576.

Neuhaus OW, Lerseth DS: 1979. Dietary control of the renal reabsorption and excretion of α_{2u} -globulin. *Kidney Intl.* 16: 409-415.

Neuhaus OW, Flory W, Biswas N, Hollerman CE: 1981. Urinary excretion of α 2u-globulin and albumin by adult male rats following treatment with nephrotoxic agents. Nephron 28: 133-140.

Newsom GD, Vugrin D: 1987. Etiologic factors in renal cell adenocarcinoma. Semin. Nephrol. 7: 109-116.

Nicoll JW, Swann PF, Pegg AE: 1975. Effect of dimethylnitrosamine on persistence of methylated guanines in rat liver and kidney DNA. Nature 254: 261-262.

Nogueira E: 1987. Rat renal carcinogenesis after chronic simultaneous exposure to lead acetate and N-nitrosodiethylamine. Virchows Arch. (Cell Pathol.) 53: 365-374.

National Research Council (NRC). 1983. Risk assessment in the Federal government: managing the process. Washington, D.C.: National Academy Press.

NTP: 1983. Carcinogenesis bioassay of pentachloroethane (CAS No. 76-01-7) in F344/N rats and B6C3F1 mice (gavage study). NTP-81-34. NIH Publication No. 83-1788. NTP-TR-232. Dated April 1983.

NTP: 1986a. Toxicology and carcinogenesis studies of isophorone (CAS No. 78-59-1) in F344/N rats and B6C3F1 mice (gavage studies). NIH Publication No. 86-2547. NTP-TR-291, dated January 1986.

NTP: 1986b. Toxicology and carcinogenesis of tetrachloroethylene (perchloroethylene) (CAS No. 127-18-4) in F344/N rats and B6C3F1 mice (inhalation studies). NIH Publication No. 86-2567. NTP-TR-311, dated August 1986.

NTP: 1987a. Toxicology and carcinogenesis studies of 1,4-dichlorobenzene (CAS No. 106-46-7) in F344/N rats and B6C3F1 mice (gavage studies). NIH Publication No. 87-2575. NTP-TR-319, dated January 1987.

NTP: 1987b. Toxicology and carcinogenesis studies of dimethyl methylphosphonate (CAS No. 756-79-6) in F344/N rats and B6C3F1 mice (gavage studies). NIH Publication No. 87-2579. NTP-TR-323, dated November 1987.

NTP: 1988. Toxicology and carcinogenesis studies of trichloroethylene (CAS No. 79-01-6) in four strains of rats (ACI, August, Marshall, Osborne-Mendel) (gavage studies). NIH Publication No. 88-2529. NTP-TR-273, dated April 1988.

NTP: 1989. Toxicology and carcinogenesis studies of hexachloroethane (CAS No. 67-72-1) in F344/N rats (gavage studies). NIH Publication No. 89-2816. NTP-TR-361, dated August 1989.

NTP: 1990. Toxicology and carcinogenesis studies of d-limonene (CAS No. 5989-27-5) in F344/N rats and B6C3F1 mice (gavage

studies). NIH Publication No. 88-2902. NTP-TR-347, dated January 1990.

Oliver J, MacDowell M: 1958. Cellular mechanisms of protein metabolism in the nephron. VII. The characteristics and significance of the protein absorption droplets (hyaline droplets) in epidemic hemorrhagic fever and other renal diseases. J. Exptl. Med. 107: 731-754.

Olson MJ, Garg BD, Murty CVR, Roy AK: 1987. Accumulation of α_{2u} -globulin in the renal proximal tubules of male rats exposed to unleaded gasoline. Toxicol. Appl. Pharmacol. 90: 43-51.

Olson MJ, Johnson JT, Reidy CA: 1990. A comparison of male rat and human urinary proteins: implications for human resistance to hyaline droplet nephropathy. Toxicol. Appl. Pharmacol. 102: 524-536.

Olson MJ, Mancini MA, Garg BD, Roy AK: 1988. Leupeptin-mediated alteration of renal phagolysosomes: similarity to hyaline droplet nephropathy of male rats exposed to unleaded gasoline. Toxicol. Lett. 41: 245-254.

OSTP: 1985. Chemical Carcinogens; A Review of the Science and Its Associated Principles. Office of Science and Technology Policy, February 1985. Federal Register, March 14, 1985, pp. 10371-10442.

Paganini-Hill A, Glazer E, Henderson BE, Ross RK: 1980. Cause-specific mortality among newspaper web pressmen. J. Occup. Med. 22: 542-544.

Papiz MZ, Sawyer L, Eliopoulos EE, North ACT, Findlay JBC, Sivaprasadarao R, Jones TA, Newcomer ME, Kraulis PJ: 1986. The structure of β -lactoglobulin and its similarity to plasma retinol-binding protein. Nature 324: 383-385.

Parker GA, Bogo V, Young RW: 1981. Acute toxicity of conventional versus shale-derived JP5 jet fuel: light microscopic, hematologic, and serum chemistry studies. Toxicol. Appl. Pharmacol. 57: 302-317.

Pegg AE: 1984. Methylation of the O^6 -position of guanine in DNA is the most likely initiating event in carcinogenesis by methylating agents. Cancer Investig. 2: 223-231.

Pervaiz S, Brew K: 1987. Homology and structure-function correlations between α^1 -acid glycoprotein and serum retinol-binding protein and its relatives. FASEB J. 1: 209-214.

Pesce AJ, Clyne DH, Pollak VE, Kant SF, Foulkes EC, Selenke WM: 1980. Renal tubular interactions of proteins. Clin. Biochem. 13: 209-215.

- Peterson PA, Berggard I: 1971. Isolation and properties of a human retinol-transporting protein. *J. Biol. Chem.* 246: 25-33.
- Pevsner J, Reed RR, Feinstein PG, Snyder SH: 1988. Molecular cloning of odorant-binding protein: member of a ligand carrier family. *Science* 241: 336-339.
- Phillips RD, Cockrell BY: 1984. Kidney structural changes in rats following inhalation exposure to C₁₀-C₁₁ isoparaffinic solvent. *Toxicology* 3: 261-273.
- Phillips RD, Egan GF: 1984. Effect of C₁₀-C₁₁ isoparaffinic solvent on kidney function in Fischer 344 rats during eight weeks of inhalation. *Toxicol. Appl. Pharmacol.* 73: 500-510.
- Phillips RD, Moran EJ, Dodd DE, Fowler EH, Kary CD, O'Donoghue J: 1987. A 14-week vapor inhalation toxicity study of methyl isobutyl ketone. *Fund. Appl. Toxicol.* 9: 380-388.
- Phillips SC, Petrone RL, Hemstreet GP: 1988. A review of the non-neoplastic kidney effects of hydrocarbon exposure in humans. *Occupat. Med: State of the Art Reviews* 3: 495-509.
- Pickle LW, Mason TJ, Howard N, Hoover R, Fraumeni JF: 1987. Atlas of U.S. cancer mortality among whites: 1950-1980. Washington, D.C. U.S. Department of Health and Human Services Publication 46 (NIH 87-2900).
- Pirani CL, Silva FG, Appel GB: 1983. Tubulo-interstitial disease in multiple myeloma and other nonrenal neoplasias. In: Cotran RS, Brenner BM, Stein JH (eds.), *Tubulo-Interstitial Nephropathies*. Churchill Livingstone, New York. pp. 287-334.
- Pitot HC: 1982. The natural history of neoplastic development: the relation of experimental models to human cancer. *Cancer* 49: 1206-1211.
- Poole C, Satterfield MH, Levin L, Rothman KJ, Dreyer NA: 1990. A case-control study of kidney cancer among petroleum refinery workers. Final Report to American Petroleum Institute, Washington D.C. by Epidemiology Resources Inc. Chestnut Hill, Massachusetts. Dated January 2nd, 1990.
- Pruzanski W, Platts ME: 1970. Serum and urinary proteins, lysozyme (muramidase), and renal dysfunction in mono- and myelomonocytic leukemia. *J. Clin. Invest.* 49: 1694-1707.
- Read NG, Astbury PJ, Morgan RJI, Parsons DN, Port CJ: 1988. Induction and exacerbation of hyaline droplet formation in the proximal tubular cells of the kidneys from male rats receiving a variety of pharmacological agents. *Toxicology* 52: 81-101.

Redmond CK, Ciocco A, Lloyd W, Ruch HW: 1972. Long-term mortality study of steelworkers. VI. Mortality from malignant neoplasms among coke oven workers. J. Occup. Med. 14: 621-629.

Reese AJM, Winstanley DP: 1958. The small tumour-like lesions of the kidney. Brit. J. Cancer 12: 507-516.

Richardson KA, Wilmer JL, Smith-Simpson D, Skopek TR: 1986. Assessment of the genotoxic potential of unleaded gasoline and 2,2,4-trimethylpentane in human lymphoblasts in vitro. Toxicol. Appl. Pharmacol. 82: 316-322.

Ridder G, Von Barger E, Parker R, Alden C: 1987. A comparative study of the acute effects of d-limonene and decalin on male rat nephropathy. Toxicologist 7: 238 (abstract).

Ridder GM, Von Barger EC, Parker RD, Alden CL: 1988. Spontaneous and induced accumulation of α_2 u globulin in the kidney cortex of rats and mice. Toxicologist 8: 89 (abstract).

Ridder GN, Von Barger EC, Alden CL, Parker RD: 1990. Increased hyaline droplet formation in male rats exposed to decalin is dependent on the presence of α_2 u globulin. Fund. Appl. Toxicol. 15: 732-743.

Ritchie AWS, Chisholm GD: 1983. The natural history of renal carcinoma. Semin. Oncol. 10: 390-400.

Rosanoff KA, Siegel MR: 1981. Mechanism of action and fate of the fungicide chlorothalonil (2,4,5,6-tetrachloroisophthalonitrile) in biological systems 3. Interaction with mammalian DNA, histones, and isolated rat liver nuclei. Pesticide Biochem. Physiol. 16: 120-128.

Roy AK, Neuhaus OW: 1967. Androgenic control of a sex-dependent protein in the rat. Nature 214: 618-620.

Roy AK, Chatterjee B: 1983. Sexual dimorphism in the liver. Ann. Rev. Physiol. 45: 37-50.

Roy AK, McMinn DM, Biswas NM: 1975. Estrogenic inhibition of the hepatic synthesis of α_2 u-globulin in the rat. Endocrinology 97: 1501-1508.

Roy AK, Nath TS, Motwani NM, Chatterjee B: 1983. Age-dependent regulation of the polymorphic forms of α_2 u-globulin. J. Biol. Chem. 258: 10123-10127.

Savitz DA, Moure R: 1984. Cancer risk among oil refinery workers: a review of epidemiological studies. J. Occup. Med. 26: 662-670.

Sawyer L: 1987. Protein structure - One fold among many. Nature

327, 659.

Selikoff IJ, Hammond EC, Seidman M: 1979. Mortality experience of insulation workers in the United States and Canada, 1943-76. Ann. N.Y. Acad. Sci. 330: 91-116.

Serve MP, Olson CT, Llewellyn BM, Bruner RH, Yu KO, Hobson DW: 1988. The metabolism and nephrotoxicity of tetralin in Fischer 344 rats. Toxicologist 8: 180 (abstract).

Shapiro LE, Sachchidananda J: 1982. Regulation of proteins by thyroid hormone and glucocorticoid: the responses of hepatic α_{2u} -globulin and pituitary growth hormone differ in adult male hypothyroid rats. Endocrinology 111: 653-660.

Short BG, Burnett VL, Swenberg JA: 1986. Histopathology and cell proliferation induced by 2,2,4-trimethylpentane in the male rat kidney. Toxicologic Pathol. 14: 194-203.

Short BG, Burnett VL, Cox MG, Bus JS, Swenberg JA: 1987. Site-specific renal cytotoxicity and cell proliferation in male rats exposed to petroleum hydrocarbons. Lab. Invest. 57: 564-577.

Short BG, Burnett VL, Swenberg JA: 1989a. Elevated proliferation of proximal tubule cells and localization of accumulated α_{2u} -globulin in F344 rats during chronic exposure to unleaded gasoline or 2,2,4-trimethylpentane. Toxicol. Appl. Pharmacol. 101: 414-431.

Short BG, Steinhagen WH, Swenberg JA: 1989b. Promoting effects of unleaded gasoline and 2,2,4-trimethylpentane on the development of atypical cell foci and renal tubular cell tumors in rats exposed to N-ethyl-N-hydroxyethylnitrosamine. Cancer Res. 49: 6369-6378.
Siemiątycki J, Dewar R, Nadon L, Gerin M, Richardson L, Wacholder S: 1987. Associations between several sites of cancer and twelve petroleum-derived liquids. Scand. J. Work Environ. Health 13: 493-504.

Sippel AE, Feigelson P, Roy AK: 1975. Hormonal regulation of the hepatic messenger RNA levels for α_{2u} globulin. Biochemistry 14: 825-829.

Sippel AE, Kurtz DT, Morris HP, Feigelson P: 1976. Comparison of in vivo translational rates and messenger RNA levels of α_{2u} -globulin in rat liver and Morris hepatoma 5123D. Cancer Res. 36: 3588-3593.

Silverberg E, Lubera JA: 1989. Cancer statistics. 1989. CA. 39: 3-20.

Smith AH, Shearn VI, Wood R: 1989. Asbestos and kidney cancer: the evidence supports a causal association. Am. J. Industr. Med. 16:

159-166.

Solleveld HA, Haseman JK, McConnell EE: 1984. Natural history of body weight gain, survival, and neoplasia in the F344 rat. J. Natl. Cancer Inst. 72: 929-940.

Steenland NK, Thun MJ, Ferguson CW, Port FK: 1990. Occupational and other exposures associated with male end-stage renal disease: a case/control study. Am. J. Public Hlth. 80: 153-157.

Stonard MD: 1987. Proteins, enzymes and cells in urine as indicators of the site of renal damage. In: Bach PH, Lock EA (eds.), Nephrotoxicity in the Experimental and Clinical Situation. Part 2. Martinus Nijhoff, Lancaster, pp. 563-592.

Stonard MD, Foster JR, Phillips PGN, Simpson MG, Lock EA: 1985. Hyaline droplet formation in rat kidney induced by 2,2,4-trimethylpentane. In: Bach PH, Lock EA, (eds.), Renal Heterogeneity and Target Cell Toxicity. John Wiley and Sons, Chichester, pp. 485-488.

Stonard MD, Phillips PGN, Foster JR, Simpson MG, Lock EA: 1986. α_{2u} -Globulin: measurement in rat kidney following administration of 2,2,4-trimethylpentane. Toxicology 41: 161-168.

Stone LC, Kanerva RL, Burns JL, Alden CL: 1987. Decalin-induced nephrotoxicity: light and electronmicroscopic examination of the effects of oral dosing on the development of kidney lesions in the rat. Fd. Chem. Toxicol. 25: 43-52.

Strasser J, Charbonneau M, Borghoff SJ, Turner MJ, Swenberg JA: 1988. Renal protein droplet formation in male Fischer 344 rats after isophorone (IPH) treatment. Toxicologist 8: 136 (abstract).

Swenberg JA: 1989b. Lecture sponsored by International Life Sciences Institute, presented at Brookings Institution, Washington D.C., December 13.

Swenberg JA, Short B, Borghoff S, Strasser J, Charbonneau M: 1989a. The comparative pathobiology of α_{2u} -globulin nephropathy. Toxicol. Appl. Pharmacol. 97: 35-46.

Szoka PR, Paigen K: 1978. Regulation of mouse major urinary protein production by the MUP-A gene. Genetics 90: 597-612.

Tannenbaum M: 1971. Ultrastructural pathology of human renal cell tumors. Pathol. Ann. 6: 249-277.

Tannenbaum M: 1983. Surgical and histopathology of renal tumors. Sem. Oncol. 10: 385-389.

Thier R, Peter H, Wiegand HJ, Bolt HM: 1990. DNA binding study of

isophorone in rats and mice. Arch. Toxicol. 64: 684-685.

Thomas FB, Halder CA, Holdsworth CE, Cockrell BY: 1985. Hydrocarbon nephropathy in male rats. Temporal and morphologic characterization of the renal lesions. In: Bach PH, Lock EA (eds.), Renal Heterogeneity and Target Cell Toxicity. John Wiley and Sons, New York, pp. 477-480.

Thomas TL, Decoufle P, Moure-Eraso R: 1980. Mortality among workers employed in petroleum refining and petrochemical plants. J. Occup. Med. 22: 97-103.

Thorhallson P, Tulinius H: 1981. Tumors in Iceland. 3. Malignant tumors of kidney. A histological classification. Acta Path. Microbiol. Scand. Sect. A 89: 403-410.

Thung PJ: 1962. Physiological proteinuria in mice. Acta Physiol. Pharmacol. Neerl. 10: 248-261.

Trump BF, Jones TW, Heatfield BM: 1984a. The biology of the kidney. In: Mehlman MA, Hemstreet GP, Thorpe JJ, Weaver NK (eds.), Advances in Modern Environmental Toxicology, Vol. VII, Renal Effects of Petroleum Hydrocarbons. Princeton Scientific Publishers, Inc., Princeton, pp. 27-49.

Trump, BF, Lipsky MM, Jones TW, Heatfield BM, Higginson J, Endicott K, Hess HB: 1984b. An evaluation of the significance of experimental hydrocarbon toxicity to man. In: Mehlman MA, Hemstreet GP, Thorpe JJ, Weaver NK (eds.), Advances in Modern Environmental Toxicology, Vol. VII, Renal Effects of Petroleum Hydrocarbons. Princeton Scientific Publishers, Inc., Princeton, pp. 273-288.

Tsuji M, Fujisaki Y, Arikawa Y, Masuda S, Tanaka T, Sato K, Noda K, Ide H, Kikuchi M: 1975. Studies on d-limonene, as gallstone solubilizer (IV). Chronic toxicity in dogs (in Japanese). Pharmacometrics 9: 775-808.

UAREP: 1983. Hydrocarbon toxicity: acute, subchronic, and chronic effects in relation to unleaded gasoline exposure of rodents with comments on the significance to human health. Universities Associated for Research and Education in Pathology, Inc. Contract No. PS-6 UAREP (504-2) with the American Petroleum Institute. December 1983.

Umemura T, Takada K, Ogawa Y, Kamata E, Saito M, Kurokawa Y: 1990. Sex difference in inhalation toxicity of p-dichlorobenzene (p-DCB) in rats. Tox. Lett. 52: 209-214.

Unterman RD, Lynch KR, Nakhasi HL, Dolan KP, Hamilton JW, Cohn DV, Feigelson P. 1981. Cloning and sequence of several α_2 -globulin cDNAs. Proc. Natl. Acad. Sci. U.S.A. 78, 3478-3482.

USEPA: 1985. Health Assessment Document for Tetrachloroethylene (Perchloroethylene). Office of Health and Environmental Assessment, U.S. Environmental Protection Agency, July 1985. pp. 8-23, 8-24.

USEPA: 1987. Evaluation of the carcinogenicity of unleaded gasoline. Carcinogen Assessment Group, United States Environmental Protection Agency, September 1987.

USEPA: 1988. Chlorothalonil. Health advisory. Office of Drinking Water, U.S. Environmental Protection Agency, June 1988.

USEPA: 1991. Report of the Peer Review Workshop on the Association of Alpha-2u-G Accumulation with Renal Effects in the Male Rat. In preparation.

Vamvakas S, Herkenhoff M, Dekant W, Henschler D: 1989. Mutagenicity of tetrachloroethylene in the Ames test--metabolic activation by conjugation with glutathione. J. Biochem. Toxicology 4: 21-27.

Vandoren G, Mertens B, Heyns W, Van Baelen H, Rombauts W, Verhoeven G: 1983. Different forms of α_{2u} -globulin in male and female rat urine. Eur. J. Biochem. 134: 175-181.

Viau C, Bernard A, Gueret F, Maldague P, Gengoux P, Lauwerys R: 1986. Isoparaffinic solvent-induced nephrotoxicity in the rat. Toxicology 38: 227-240.

Wallace AC, Nairn RC: 1972. Renal tubular antigens in kidney tumors. Cancer 29: 977-981.

Warter HL: 1983. Recent progress in the pathological anatomy of cancers of the kidney. Progr. Surg. (Basel) 17: 32-57.

Watabe T, Hiratsuka A, Oxawa N, Isobe M: 1981. A comparative study on the metabolism of d-limonene and 4-vinylcyclohex-1-ene by hepatic microsomes. Xenobiotica 11: 333-344.

Webb DR, Kanerva RL, Hysell DK, Alden CL, Lehman-McKeeman LD: 1990. Assessment of the subchronic oral toxicity of d-limonene in dogs. Fd. Chem. Toxicol. 28: In press.

Webb DR, Ridder GM, Alden CL: 1989. Acute and subchronic nephrotoxicity of d-limonene in Fischer 344 rats. Fd. Chem. Toxicol. 27: 639-649.

Wen CP, Tsai SP, McLellan WA, Gibson RL: 1983. Long-term mortality study of oil refinery workers: mortality of hourly and salaried workers. Am. J. Epidemiol. 118: 526-542.


Wong O, Raabe GK: 1989. Critical review of cancer epidemiology in petroleum industry employees, with a quantitative meta-analysis by

cancer site. Am. J. Industr. Med. 15: 283-310.

Yoon JS, Mason JM, Valencia R, Woodruff EC, Zimmering S: 1985. Chemical mutagenesis testing in Drosophila. IV. results of 45 coded compounds tested for the National Toxicology Program, Environ. Mutagen. 7: 349-367.

Yu MC, Mack TM, Hanisch R, Cicioni C, Henderson BE: 1986. Cigarette smoking, obesity, diuretic use, and coffee consumption as risk factors for renal cell carcinoma. J. Natl. Cancer Inst. 77: 351-356.

A P P E N D I X



APPENDIX 1. SUBSTANCES OR CHEMICALS THAT INDUCE THE ACCUMULATION OF HYALINE DROPLETS IN RENAL PROXIMAL TUBULE EPITHELIAL CELLS AND/OR ELEVATED LEVELS OF RENAL ALPHA-2U-GLOBULIN

Substance/Chemical	Evidence for exacerbation of hyaline droplets in renal proximal tubule cells			Evidence for increased renal alpha-2u-globulin levels		
	Males	Females	References	Males	Females	References
Unleaded gasoline	+	-	Halder et al. (1984) Thomas et al. (1985) Olson et al. (1987)	+	NR	Olson et al. (1987) Garg et al. (1988)
Murty et al. (1988)			Garg et al. (1988) Murty et al. (1988)			
2,2,4-Trimethyl pentane	+	-	Stonard et al. (1985, 1986) Short et al. (1986, 1987)	+	-	Stonard et al. (1986) Charbonneau et al. (1987) Lock et al. (1987b)
JP-5 jet fuel (mixed distillate hydrocarbons)	+	-	Parker et al. (1981) Bruner (1984) Gaworski et al. (1984) MacNaughton and Uddin (1984)	NR	NR	
JP-4 jet fuel (mixed distillate hydrocarbons)	+	-	Bruner (1984) MacNaughton and Uddin (1984)	NR	NR	
Diesel fuel, marine	+	-	Bruner (1984) Gaworski et al. (1985b)	NR	NR	
JP-10 synthetic jet fuel + (exohexahydro-4,7-methanoindan)	+	NR	MacNaughton and Uddin (1984) Mattie et al. (1988)	NR	NR	

+ Positive
- Negative
N.R. = Not reported

APPENDIX 1. SUBSTANCES OR CHEMICALS THAT INDUCE THE ACCUMULATION OF HYALINE DROPLETS IN RENAL
(cont'd) PROXIMAL TUBULE EPITHELIAL CELLS AND/OR ELEVATED LEVELS OF RENAL ALPHA-2U-GLOBULIN

Substance/Chemical	Evidence for exacerbation of hyaline droplets in renal proximal tubule cells			Evidence for increased renal alpha-2u-globulin levels		
	Males	Females	References	Males	Females	References
RJ-5 synthetic jet fuel (hydrogenated dimers of norbornadiene)	+	-	MacNaughton and Uddin (1984)	NR	NR	
JP-7 distillate jet fuel	+	-	Bruner (1990, unpublished data) Alden (1989)	NR	NR	
JP-15 distillate jet fuel	+	-	Bruner (1990, unpublished data) Alden (1989)	NR	NR	
Stoddard solvent	+	-	Phillips and Cockrell (1984)	NR	NR	
C ₁₀ - C ₁₂ isoparaffinic solvent (saturated aliphatic hydrocarbons)	+	-	Phillips and Cockrell (1984) Viau et al. (1986)	+	-	Viau et al. (1986)
Decalin	+	-	Alden et al. (1984, 1985) Bruner (1984) Gaworski et al. (1985a) Kanerva et al. (1987a) Stone et al. (1987)	+	-	Alden et al. (1984, 1985) Kanerva et al. (1987b)
Tetralin	+	NR	Serve et al. (1988)	NR	NR	
d-Limonene	+	-	Ridder et al. (1987) Kanerva et al. (1987a) NTP (1990) Lehman-McKeeman et al. (1989) Webb et al. (1989)	+	-	Ridder et al. (1988) Lehman-McKeeman et al. (1989) Webb et al. (1989)

APPENDIX 1. SUBSTANCES OR CHEMICALS THAT INDUCE THE ACCUMULATION OF HYALINE DROPLETS IN RENAL
(cont'd) PROXIMAL TUBULE EPITHELIAL CELLS AND/OR ELEVATED LEVELS OF RENAL ALPHA-2U-GLOBULIN

Substance/Chemical	Evidence for exacerbation of hyaline droplets in renal proximal tubule cells			Evidence for increased renal alpha-2u-globulin levels		
	Males	Females	References	Males	Females	References
1,4-Dichlorobenzene	+	-	NTP (1987a) Bonhard et al. (1988) Charbonneau et al. (1989)	+	NR	Charbonneau et al. (1989)
Tetrachloroethylene (Perchloroethylene)	+	-	Goldsworthy et al. (1988) Green et al. (1990)	+	-	Goldsworthy et al. (1989)
Pentachloroethane	+	-	Goldsworthy et al. (1988)	+	-	Goldsworthy et al. (1988)
Hexachloroethane	+	-	NTP (1989)	NR	NR	
Isophorone	+	NR	Strasser et al. (1988)	+	NR	Strasser et al, 1988
Lindane	+	-	Dietrich and Suenberg (1990)	+	-	Dietrich and Suenberg (1990)
Dimethyl methyl- phosphonate	+	-	NTP (1987b)	NR	NR	
Methyl isobutyl ketone	+	-	Phillips et al. (1987)	NR	NR	
Methyl isoamyl ketone	+	-	Katz et al. (1986)	NR	NR	
Diisobutyl ketone	+	-	Dodd et al. (1987)	NR	NR	
BW540C (3-methylamino-1- (3-trifluoromethylphenyl)- 2-pyrazoline)	+	-	Read et al. (1988)	+	NR	Read et al. (1988)
BW58C (mixture of isomeric <u>cis</u> and <u>trans</u> forms of 2-(4'-t-butylcyclohexyl)-3- hydroxy-1-4-naphthoquinone)	+	-	Read et al. (1988)	+	NR	Read et al. (1988)

APPENDIX 1. SUBSTANCES OR CHEMICALS THAT INDUCE THE ACCUMULATION OF HYALINE DROPLETS IN RENAL PROXIMAL TUBULE EPITHELIAL CELLS AND/OR ELEVATED LEVELS OF RENAL ALPHA-2U-GLOBULIN (cont'd)

Substance/Chemical	Evidence for exacerbation of hyaline droplets in renal proximal tubule cells			Evidence for increased renal alpha-2u-globulin levels		
	Males	Females	References	Males	Females	References
Levamisole (levoisomer of 2,3,5,6-tetrahydro-6-phenylimidazo-(2,1-b) thiazole)	+	-	Read et al. (1988)	+	NR	Read et al. (1988)
Gabapentin	+	+	Dominick et al. (1990)	+	+	Dominick et al. (1990)
3,5,5-Trimethyl-hexanoic acid derivatives	+	NR	Lehman-McKeeman et al. (1990)	+	NR	Lehman-McKeeman et al. (1990)
Tridecyl acetate	+	NR	Daugherty et al. (1990)	+	NR	Daugherty et al. (1990)
Isopropylcyclohexane	+	NR	Henningsen et al. (1988)	+	NR	Henningsen et al. (1988)
1,3,6-Tricyanohexane	+	-	Barnett et al. (1987) Johnson (1987)	NR	NR	

APPENDIX 2. NON-NEOPLASTIC FINDINGS REPORTED IN RODENT 2-YEAR STUDIES ON TEN SELECTED SUBSTANCES (continued)

CHEMICAL	NON-NEOPLASTIC LESIONS
<p>Tetrachloroethylene species: F344 rats B6C3F1 mice route: inhalation ref.: NTP-TR-311, 1986b</p>	<p>Both male and female rats exhibited renal tubule cell karyomegaly. Karyomegaly also occurred in mice. Cast formation noted in male and female mice. Tubule cell hyperplasia seen in male rats and male mice.</p>
<p>Unleaded gasoline species: F344 rat route: inhalation ref.: USEPA, 1987</p>	<p>Dose-related kidney lesions at 3-6 months in males included focal tubule basophilia and tubular casts at the corticomedullary junction. Interrelated increase in the incidence of renal pelvis mineralization was also reported at 12 months, 18 months, and terminal sacrifice. Progressive glomerulonephrosis was reported in one high-dose male at 12 months; the incidence was higher at 18 months but was dose-related; at the final sacrifice, nearly all male rats exhibited this lesion (MacFarland et al., 1984). Karyomegaly, i.e. very large nuclei within cells of tubule epithelium, first noted at 12 months; at 18 months, more numerous karyomegalic cells were observed in treated males, particularly in the high-dose group (UAREP, 1983).</p>
<p>Chloroethalonil species: Osborne-Mendel rats B6C3F1 mice route: diet ref.: NTP-TR-41, NCI, 1978a</p>	<p>No non-neoplastic renal lesions were reported in the NTP bioassays.</p>
<p>Trichloroethylene (cont.) species: ACI rats August rats Marshall rats Osborne-Mendel rats route: gavage ref.: NTP-TR-273, 1988a</p>	<p>Cytomegaly noted in males and females of all strains. Toxic nephropathy increased in both sexes of all strains. Calcification was produced in kidneys of ACI male and female rats.</p>

APPENDIX 2. NON-NEOPLASTIC FINDINGS REPORTED IN RODENT 2-YEAR STUDIES ON TEN SELECTED SUBSTANCES
THAT PRODUCE RENAL TUBULE CELL TUMORS IN RATS

CHEMICAL

NON-NEOPLASTIC LESIONS

1,4-Dichlorobenzene
species: F344 rats
B6C3F1 mice
route: gavage
ref.: NTP-TR-319, 1987a

Renal damage was not observed in female rats or mice in 13-week studies. Cell degeneration or necrosis of tubule epithelium was observed in male rats in a 13-week study. In 2-year studies, the severity of nephropathy was greater in male rats than female rats. Nephropathy characterized by degeneration and regeneration of renal tubule epithelium, tubule dilation with attenuation and atrophy of the epithelium, granular casts in tubules of the outer stripe of the medulla, thickening of basement membranes, and minimal accumulation of interstitial collagen. Renal tubular regeneration was noted in female mice. Nephropathy increased with dose in male mice and female rats.

Dimethyl methyl phosphonate
species: F344 rats
B6C3F1 mice
route: gavage
ref.: NTP-TR-323, 1987b

Kidneys of dosed male rats but not dosed male mice had varying degrees of tubule cell regeneration, hyaline droplet degeneration, and cellular infiltration (13 wks). At 2-years, the average severity of nephropathy increased and calcification was observed in the collecting tubules of the renal pelvis of male rats. The nephropathy was characterized by degeneration of the tubule epithelium, tubule dilation with attenuation and atrophy of the epithelium, granular casts in the tubules of the outer stripe of the outer medulla, thickening of basement membranes, minimal to mild accumulation of interstitial collagen, and minimal to mild inflammatory cell infiltrates. The increase in severity of nephropathy was ranked, control to high dose, 1.9, 2.5, 2.8, on scale of 1 to 4.


Hexachloroethane


species: Osborne Mendel rat
B6C3F1 mice
route: gavage
ref.: NTP-TR-68, NCI,
1978b

Toxic tubule nephropathy was observed in all groups of treated rats and in male and female mice. The nephropathy in rats was characterized by degeneration, necrosis, and the presence of large hyperchromatic regenerating epithelial cells. Overlying the tubule lesions were chronic interstitial fibrosis and nephritis, focal pyelonephritis, tubular ectasia, cast formation, and focal glomerulosclerosis. In mice, nephropathy was characterized by degeneration of affected tubules containing hyaline casts. The kidney often showed infiltration of inflammatory cells, fibrosis, and calcium deposition. The incidences of toxic nephropathy were higher in mice than in rats.

APPENDIX 2. NON-NEOPLASTIC FINDINGS REPORTED IN RODENT 2-YEAR STUDIES ON 10 SELECTED SUBSTANCES (continued)

CHEMICAL	NON-NEOPLASTIC LESIONS
<p>Hexachloroethane (cont.) species: F344/N rats route: gavage ref.: NTP-TR-361, 1989</p>	<p>Nephropathy observed in nearly all males; overall average severity mild in vehicle controls and mild to moderate in dosed males. Incidence and severity of nephropathy was increased in dosed females relative to controls. Nephropathy in each sex consisted of tubule cell degeneration, regeneration and dilation, atrophy, glomerulosclerosis, interstitial fibrosis, and chronic inflammation. Linear mineralization of the renal papillae showed dose-related increase in male rats. Hyperplasia of pelvic transitional epithelium was increased. Incidences of mineralization and pelvic epithelial hyperplasia were not increased in females.</p>
<p>Isophorone species: F344 rat B6C3F1 mouse route: gavage ref.: NTP-TR-291, 1986a</p>	<p>Tubular cell mineralization was increased in dosed male rats but not in dosed female rats. This lesion was characterized by basophilic aggregates of mineral most often found in the medullary collecting ducts and occurred coincidentally with lesions of chronic nephropathy. The incidence of nephropathy was similar in dosed and vehicle control male rats, the severity was greater in low dose males. Hyperplasia of the renal pelvis was observed in dosed male rats but not in vehicle controls.</p>
<p>d-Limonene species: F344 rats B6C3F1 mice route: gavage ref.: NTP-TR-347, 1990</p>	<p>Increased severity of nephropathy in male rats at 13 weeks characterized by degeneration of epithelial cells in convoluted tubules, granular casts in the outer medulla, and epithelial regeneration. No lesions in female rats. Increase in severity of nephropathy and linear deposits of mineral in the renal medulla and papilla in male rats in 2-year studies.</p>
<p>Pentachloroethane species: F344 rats B6C3F1 mice route: gavage ref.: NTP-TR-232, 1983</p>	<p>Chronic, diffuse inflammation, distinguishable from nephropathy seen in aging F344 rats, found in male rats in a significant dose-related increase. Interstitial fibrosis and tubule dilation more severe than in old-age nephropathy. Mineralization of the renal papilla observed at increased incidences in dosed male rats. Some dilated tubules with giant cells and casts were observed. No indication of renal toxicity was reported for female rats or mice.</p>





United States
Environmental Protection
Agency

Center for Environmental Research
Information
Cincinnati OH 45268

Official Business
Penalty for Private Use, \$300

BULK RATE
POSTAGE & FEES PAID
EPA
PERMIT NO. G-35

Please make all necessary changes on the above label,
detach or copy, and return to the address in the upper
left-hand corner.

If you do not wish to receive these reports CHECK HERE ☐
detach, or copy this cover, and return to the address in the
upper left-hand corner.

EPA/625/3-91/019A