WAT
EPAX
9205-
0094
c.2
EPA
500
ECAO-
CIN-
442
c.2
#25902604

United States
Environmental Protection
Agency

ECAO-CIN-442
May, 1991
Revised January 1992

EPA

Research and Development

DRINKING WATER CRITERIA DOCUMENT
FOR CYANIDE

Prepared for

HEALTH AND ECOLOGICAL CRITERIA DIVISION
OFFICE OF SCIENCE AND TECHNOLOGY
OFFICE OF WATER

Prepared by

Environmental Criteria and Assessment Office
Office of Health and Environmental Assessment
U.S. Environmental Protection Agency
Cincinnati, OH 45268

U.S. EPA Headquarters Library
Mail code ~~3201~~ 3404T
1200 Pennsylvania Avenue NW
Washington DC 20460

JUN 18 1992

JUL 2 1992

HEADQUARTERS LIBRARY
ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

DISCLAIMER

This document has been reviewed in accordance with U.S. Environmental Protection Agency policy and approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

FOREWORD

Section 1412 (b)(3)(A) of the Safe Drinking Water Act, as amended in 1986, requires the Administrator of the Environmental Protection Agency to publish maximum contaminant level goals (MCLGs) and promulgate National Primary Drinking Water Regulations for each contaminant, which, in the judgment of the Administrator, may have an adverse effect on public health and which is known or anticipated to occur in public water systems. The MCLG is nonenforceable and is set at a level at which no known or anticipated adverse health effects in humans occur and which allows for an adequate margin of safety. Factors considered in setting the MCLG include health effects data and sources of exposure other than drinking water.

This document provides the health effects basis to be considered in establishing the MCLG. To achieve this objective, data on pharmacokinetics, human exposure, acute and chronic toxicity to animals and humans, epidemiology and mechanisms of toxicity are evaluated. Specific emphasis is placed on literature data providing dose-response information. Thus, while the literature search and evaluation performed in support of this document has been comprehensive, only the reports considered most pertinent in the derivation of the MCLG are cited in the document. The comprehensive literature data base in support of this document includes information published up to March 1987; however, more recent data may have been added during the review process. Health effects and toxicokinetic data provided in this document are limited to HCN and free CN (CN^-).

When adequate health effects data exist, Health Advisory values for less than lifetime exposures (1-day, 10-day and longer-term, ~10% of an individual's lifetime) are included in this document. These values are not used in setting the MCLG, but serve as informal guidance to municipalities and other organizations when emergency spills or contamination situations occur.

Tudor Davies
Office of Science and
Technology

James Elder, Director
Office of Groundwater
and Drinking Water

DOCUMENT DEVELOPMENT

Linda R. Papa, M.S., Document Manager
Environmental Criteria and Assessment Office, Cincinnati
U.S. Environmental Protection Agency

Helen H. Ball, M.S., Project Officer
Environmental Criteria and Assessment Office, Cincinnati
U.S. Environmental Protection Agency

Authors

Dipak K. Basu, Ph.D.
Michael W. Neal, Ph.D.
Sharon B. Wilbur, M.S.
Syracuse Research Corporation
Syracuse, New York
(EPA Contract #68-03-3112)

Editorial Reviewers
Judith Olsen, B.A.
Environmental Criteria and
Assessment Office, Cincinnati
U.S. Environmental Protection Agency

Scientific Reviewers

Cynthia Sonich-Mullin M.S.
Michael L. Dourson, Ph.D.
Environmental Criteria and
Assessment Office, Cincinnati
U.S. Environmental Protection Agency

Ernest C. Foulkes, Ph.D.
Kettering Laboratory
Department of Environmental Health
University of Cincinnati
Cincinnati, Ohio
(EPA Contract #68-03-3234,
Eastern Research Group)
Nancy H. Chiu, Ph.D.
James J. Murphy, Ph.D.
Edward V. Ohanian, Ph.D.
Office of Science and Technology
U.S. Environmental Protection Agency

Julio A. Salinas, Ph.D.
1372 High Street
Westwood, MA
(EPA Contract #68-03-3234,
Eastern Research Group)

Document Preparation

Technical Support Services Staff, Environmental Criteria and Assessment Office, Cincinnati

TABLE OF CONTENTS

		Page
I.	SUMMARY	I-1
II.	PHYSICAL AND CHEMICAL	II-1
	STRUCTURE AND IDENTIFICATION	II-1
	PHYSICAL AND CHEMICAL PROPERTIES	II-1
	ENVIRONMENTAL FATE IN AQUATIC MEDIA	II-5
	SUMMARY	II-8
III.	TOXICOKINETICS	III-1
	ABSORPTION	III-1
	DISTRIBUTION	III-4
	METABOLISM	III-10
	EXCRETION	III-17
	SUMMARY	III-19
IV.	HUMAN EXPOSURE	IV-1
V.	HEALTH EFFECTS IN ANIMALS	V-1
	GENERAL TOXICITY	V-1
	Acute Exposure	V-2
	Subchronic Exposure	V-15
	Chronic Exposure	V-24
	TARGET ORGAN TOXICITY	V-26
	Central Nervous System	V-26
	Heart	V-28
	Thyroid	V-28
	OTHER EFFECTS	V-28
	Carcinogenicity	V-28
	Mutagenicity	V-29
	Teratogenicity and Other Reproductive Effects	V-30
	SUMMARY	V-35
VI.	HEALTH EFFECTS IN HUMANS	VI-1
	ACUTE EXPOSURE	VI-1
	Oral	VI-1
	Inhalation	VI-4
	Dermal	VI-7

TABLE OF CONTENTS (cont.)

	Page
SUBCHRONIC ANC CHRONIC EXPOSURE	VI-7
Oral	VI-7
Inhalation	VI-8
Occupational Exposure	VI-9
Epidemiologic Studies	VI-10
HIGH RISK SUBPOPULATIONS	VI-13
SUMMARY	VI-13
VII. MECHANISMS OF TOXICITY	VII-1
ACUTE	VII-1
CHRONIC	VII-2
SYNERGISM	VII-4
ANTAGONISM	VII-4
SUMMARY	VII-6
VIII. QUANTIFICATION OF TOXICOLOGIC EFFECTS	VIII-1
INTRODUCTION	VIII-1
NONCARCINOGENIC EFFECTS	VIII-6
Short-Term Exposure	VIII-7
Long-Term Exposure	VIII-14
QUANTIFICATION OF NONCARCINOGENIC EFFECTS	VIII-16
Derivation of 1-Day HA	VIII-16
Derivation of 10-Day HA	VIII-17
Derivation of Longer-Term HA	VIII-18
Assessment of Lifetime Exposure and Derivation of a DWEL	VIII-18
CARCINOGENIC EFFECTS	VIII-20
QUANTIFICATION OF CARCINOGENIC EFFECTS	VIII-20
EXISTING GUIDELINES, RECOMMENDATIONS AND STANDARDS	VIII-21
SPECIAL GROUPS AT RISK	VIII-22
SUMMARY	VIII-23
IX. REFERENCES	IX-1

LIST OF TABLES

No.	Title	Page
II-1	Structure and Identification Symbols of Selected Cyanides	II-2
II-2	Physical and Chemical Properties of the Selected Cyanides	II-3
III-1	Mean Levels and Ranges of Cyanide Ion Concentration in Human Organs in Cases of Fatal Poisoning	III-6
III-2	Cyanide Levels in Human Tissues and Fluids After Fatal Cyanide Poisoning	III-7
III-3	Cyanide Concentrations in Various Organs of Rats Treated with NaCN Orally (A) or HCN by Inhalation (B)	III-8
III-4	Comparison of Cyanide Concentrations in Tissues from Rabbits Killed by HCN with Concentrations in Tissues from Rabbits Killed with KCN	III-11
III-5	Rhodanese Activity in Tissues of the Dog, Rhesus Monkey, Rabbit and Rat (mg CN^- converted to CNS/g of tissue after 15 minutes)	III-13
V-1	Single-Dose LD_{50} Values for Cyanides	V-4
V-2	Acute Toxicity of Cyanides in Laboratory Animals	V-7
V-3	Sensitivity of Various Species to Inhalation Exposures of HCN	V-16
V-4	Effects of Subchronic Exposure of Laboratory Animals to Cyanide	V-17
V-5	Effects of Chronic Exposure of Laboratory Animals to Cyanide	V-25
V-6	Fetotoxic and Teratogenic Effects of NaCN in Hamsters	V-31
VI-1	Fatal Oral Doses of Cyanide Compounds	VI-2
VI-2	Reported (Estimated) Human Responses to Various Concentrations of HCN Vapors	VI-5
VI-3	Summary of Epidemiologic Studies of Cyanide Exposure	VI-11
VIII-1	Acute LD_{50} Values for Cyanides	VIII-8
VIII-2	Summary of Quantification of Toxicologic Effects	VIII-24

LIST OF ABBREVIATIONS

CNS	Central Nervous System
CuCN	Copper cyanide
DWEL	Drinking water equivalent level
EEG	Electroencephalogram
EKG	Electrocardiogram
GI	Gastrointestinal
HA	Health Advisory
HCN	Hydrogen cyanide (hydrocyanic acid)
i.p.	Intraperitoneal
i.m.	Intramuscular
KCN	Potassium cyanide
LD_{50}	Dose lethal to 50% of recepients
LC_{50}	Concentration lethal to 50% of recepients
LOAEL	Lowest-observed-adverse-effect level
MF	Modifying Factor
NaCN	Sodium cyanide
NOAEL	No-observed-adverse-effect level
RfD	Reference Dose
s.c.	Subcutaneous
TSH	Thyroid stimulating hormone
UF	Uncertainty Factor
UV	Ultraviolet

I. SUMMARY

Cyanides are a group of organic and inorganic compounds that contain the cyano (CN) radical. Free cyanide is defined as the sum of cyanide present as HCN and cyanide ion as CN^-. In water, hydrogen cyanide and the alkali-metal cyanides are very soluble and dissociate into their respective anions and cations in water and, except for HCN, are only slightly soluble in organic solvents. The alkaline earth metal cyanides, in general, are not very soluble in water. HCN is volatile, and in aqueous solution is a very weak acid. At pH below 8, most of HCN in an aqueous media is present in the nondissociated form. As the pH increases, HCN will dissociate into CN^-. CN^- in water may form simple or complex cyanides depending on whether there is an excess of cyanide or metal present in the aqueous media. The metal cyanides can be oxidized to isocyanate and ultimately to CO_2 and N_2 in the presence of strong oxidizing agents.

The fate of cyanides in the aquatic media may vary widely. Hydrogen cyanide and the most common alkali-metal cyanides may be lost from aquatic media primarily through the volatilization process. Some of these cyanides may also be lost through microbial degradation in aquatic media and sorption to particulate matters in water. However, both these processes are less significant than volatilization.

The sparingly soluble metal cyanides, such as CuCN in acidic waters (pH <7) may form some HCN that may subsequently volatilize from water. But the predominant fate of these compounds is sedimentation and microbial degradation.

The fate of water soluble complex metal cyanides, such as ferrocyanide and ferricyanides, with respect to biodegradation is not known with certainty. In the absence of destabilizing factors in water (high temperature, sunlight, extreme pH conditions), these complexes are expected to have long lifetimes and can be transported in the aquatic media.

The simple metal cyanides and HCN do not bioaccumulate in aquatic organisms. The water soluble complex metal cyanides may bioaccumulate to some extent in aquatic organisms although the bioaccumulation factors for such compounds are not known.

Cyanide is readily absorbed from the lungs, the GI tract and the skin. Inhalation exposure to HCN provides the most rapid route of entry, resulting in the most rapid onset of toxic effects. Cyanide enters erythrocytes and is found in the blood at low levels in normal humans. Transplacental transfer can also occur. Cyanide is detoxified by an intramitochondrial enzyme, rhodanese, which catalyzes the transfer of sulfur from a donor to cyanide to form the less toxic thiocyanate. Thiocyanate is excreted in the urine. Rhodanese is widely distributed throughout the body; the highest levels are found in the liver. Minor detoxification pathways include spontaneous reaction with cystine to form 2-imino-4-thiazolidine-carboxylic acid, and the reaction with hydroxycobalamin (vitamin B_{12}) to form cyanocobalamin. Both products are excreted in urine. The major route of cyanide elimination is via thiocyanate in the urine, although some cyanide can enter 1-carbon-compound metabolic pathways and be eliminated as CO_2 in expired air. A small amount of unchanged HCN is eliminated in expired air.

Signs of acute poisoning by cyanide include rapid breathing, gasping, tremors, convulsions and death. The severity and rapidity of onset of effects depend on route, dose, duration of exposure and compound administered. Inhalation exposure results in the most rapid absorption and appearance of toxic signs. Oral exposure to cyanide salts results in slower GI absorption, passage to the liver and faster detoxification. The acute oral (gavage) LD_{50} for cyanide is 4 mg CN^-/kg bw in rats and 3.4 mg CN^-/kg in mice. The LC_{50} for inhaled HCN by mice is 184 mg/m³. Species differ with respect to sensitivity with the dog being the most sensitive. Cyanide is less toxic when administered subchronically and chronically in the diet. Subchronic and chronic subcutaneous administration of cyanide to rats results in histopathologic damage to the brain and spinal cord. Subchronic oral exposure to cyanide in capsules has resulted in histopathologic lesions in the CNS of dogs. The results of chronic oral administration reveal no evidence of carcinogenicity. Negative results for KCN in the reverse mutation assay were obtained in five strains of Salmonella typhimurium with or without metabolic activation. A marginally positive response was found for HCN gas in strain TA100 of S. typhimurium. Cyanide was negative in a modified rec assay in Bacillus subtilis. Severe teratogenic effects were observed following the administration of high doses of NaCN by subcutaneously implanted osmotic minipumps in hamsters.

Ingestion of cyanide by humans at doses of 0.5-3.5 mg/kg bw was found to be fatal following acute axposure. Death, which is preceded by hyperventilation, vomiting, unconsciousness, convulsions, rapid heart rate, gasping and vascular collapse, occurs within 20 minutes of ingestion. A no-effect level for ingestion by humans is 0.06 mg CN^-/kg bw. The

estimated 1-minute LC_{50} for inhalation of HCN by humans is 3763 mg/m³. Inhalation of ~2000 mg/m³ result in dyspnea, followed by rapid breathing, then apnea, gasping and death within minutes. Many people are unable to smell cyanide, which has the odor of bitter almond, due to a sex-linked recessive gene. Sublethal concentrations produce dizziness, headache, confusion, nausea and numbness. Chronic oral exposure to HCN, KCN or NaCN in humans has not been described. Ingestion of cyanogenic plants, in combination with dietary deficiencies in protein, vitamin B_{12} and riboflavin, may result in neuropathies. Ingestion of cyanogenic chemicals combined with iodine deficiency may be associated with the etiology of goiter and cretinism. Other disorders associated with defective cyanide metabolism include tobacco amblyopia, retrobulbar neuritis and Leber's hereditary optic atrophy. Smoking during pregnancy may result in birth of low-weight infants. This may be partly due to the cyanide content of tobacco smoke. Case studies and epidemiologic studies of workers occupationally exposed to cyanide describe effects such as headache, dizziness, nausea and thyroid enlargement.

Cyanide exerts its toxic effects by reacting with ferric iron (Fe^{+++}) in cytochrome oxidase, the enzyme that catalyzes the terminal step in the electron transport chain, thereby preventing utilization of oxygen by cells. The CNS and the heart are particularly sensitive to this histotoxic hypoxia. If the exposure to cyanide is high enough, the detoxification mechanism via rhodanese may be overloaded and toxic signs occur. Chronic or repeated exposure to high doses may result in repeated hypoxic insult to the CNS and degenerative changes in the CNS of rats and dogs.

Synergistic effects may occur if cyanide exposure is accompanied by exposure to other inhibitors of cytochrome oxidase, such as azide and sulfide. Treatments for cyanide poisoning are based on the generation or administration of compounds that can compete with cytochrome oxidase for cyanide. Sodium nitrite can generate methemoglobin that binds cyanide. Any compound that can act as a sulfur donor for rhodanese, such as sodium thiosulfate, can be an effective antidote. Cobalt-containing compounds can compete with cytochrome oxidase for cyanide and are also effective antidotes.

The available data are insufficient to develop a 1-day, 10-day or longer-term HAs for cyanide. It is recommended that the DWEL of 0.7 mg/ℓ be adopted for the adult longer-term HA and the adjusted DWEL of 0.2 mg/ℓ be adopted for the child 1-day, 10-day and longer-term HAs. The available human and animal cancer studies are inadequate to determine the carcinogenic potential of cyanide and hence cyanide is accorded a Group D weight of evidence according to EPA's cancer risk assessment procedures.

II. PHYSICAL AND CHEMICAL PROPERTIES

Structure and Identification

Cyanides are a group of organic and inorganic compounds that contain the cyano (CN) radical. Free cyanide is defined as the sum of cyanide present as HCN and as CN^-. A large number of inorganic cyanides are used industrially, both as simple salts or as complex cyanides. The usefulness of cyanides stems from the tendency of these compounds to form strong complexes with most metals. Organic cyanides known as nitriles can dissociate to yield CN^- or HCN. Compounds such as acrylonitrile and adiponitrile are the main cyanides produced in the United States (Towill et al., 1978). In this section, only a few widely used industrial inorganic cyanides will be discussed. The structure, molecular formula, molecular weight, Chemical Abstract Services (CAS) Registry number and Registry of Toxic Effects of Chemical Substances (RTECS) number of selected cyanides are given in Table II-1.

Physical and Chemical Properties

Physical and chemical properties (Weast, 1980; Towill et al., 1978) of selected cyanides are given in Table II-2.

Hydrogen cyanide is a colorless, flammable liquid or gas that is miscible with ethanol and water but is only slightly soluble in ether (Towill et al., 1978). Liquid HCN is unstable and can polymerize violently in the absence of stabilizers. In aqueous solutions, UV light may induce polymerization of HCN. Among the many polymerized products of HCN are the trimer, tetramer and other high molecular weight polymers (Cotton and Wilkinson, 1980).

TABLE II-1

Structure and Identification Symbols of Selected Cyanides*

Compound	Molecular Formula	Molecular Weight	CAS No.	RTECS No.
Hydrogen cyanide; hydrocyanic acid; prussic acid; formo nitrile	HCN	27.03	74-90-8	MW6825000
Sodium cyanide	NaCN	49.01	143-33-9	VZ7525000
Potassium cyanide	KCN	65.12	151-50-8	TS8750000
Copper (1) cyanide; cupricin; cuprous cyanide	CuCN	89.56	544-92-3	GL7150000
Potassium ferricyanide; Tripotassium hexa-cyanoferrate	$K_3Fe(CN)_6$	326.27	13746-66-2	LJ8225000

*Source: NIOSH, 1981

TABLE II-2

Physical and Chemical Properties of the Selected Cyanides[a]

Compound	Physical State and Description	Melting Point (°C)	Boiling Point (°C)	Density (g/cm³)	Vapor Pressure (mm Hg)	Water Solubility (g/100 mℓ)	K_{ow}[b]	Conversion Factor (1 ppm equivalent of mg/m³)
HCN	colorless gas or liquid with bitter almond odor	-13.24	25.70	0.688 at 20°C (liquid)	807.2 at 27.2°C	completely miscible	0.66[c]	1.123
NaCN	colorless crystalline solid	563.7	1496	1.60-1.62[d]	0.76 at 800°C	48 at 10°C	-0.44[e]	2.037
KCN	colorless crystalline solid	634.5	NA	1.553 at 20°C	NA	71.6 at 25°C	NA	2.707
CuCN	white crystalline solid	473 (in N_2)	decomposes	2.92[d]	NA	insoluble[f]	NA	3.722
$K_3Fe(CN)_6$	red crystalline solid	decomposes	NA	1.85 at 25°C	NA	33 at 4°C	NA	13.561

[a]Adapted from Weast (1980) and Towill et al. (1978) unless otherwise stated.

[b]K_{ow} is the octanol/water partition coefficient

[c]Calculated average of values given by Leo et al. (1971)

[d]The temperature at which this value was measured is not available.

[e]Calculated by the method of Leo et al. (1971) from the value given by Hansch and Leo (1979)

[f]The exact solubility data is not available.

NA = Not available

In aqueous solution, HCN is a very weak acid having a pK_a (dissociation constant) value of 9.21 at 25°C (Cotton and Wilkinson, 1980). The relationship of pH to percent undissociated HCN in aqueous solutions at 25°C is shown below (Callahan et al., 1979):

pH	% undissociated HCN
<7	>99
8	93.3
9	58
10	13

Therefore, at pH below 8, most of HCN in aqueous media is present in the undissociated form.

Simple cyanides are represented by the formula $A(CN)_x$, where A is an alkali (sodium, potassium) or a metal. The alkali-metal cyanides, NaCN and KCN, are less soluble in ethanol and methanol than in water (Weast, 1980). Since these compounds are ionic, their solubility in less polar and nonpolar organic solvents may be even lower than in alcohols. Both NaCN and KCN are strongly hydrolyzed in aqueous media, producing basic solutions. In the presence of strong mineral acids, the salts will liberate HCN from aqueous solutions (Towill et al., 1978).

Many simple metal cyanides (CaCn, AgCN) are sparingly soluble or almost insoluble. Copper (I) cyanide has a solubility product of only 3.2×10^{-20} in aqueous solution at ambient temperature (NIOSH, 1976); it is soluble in HCl and also in KCN and NH_4OH, with the formation of complexes (Cotton and Wilkinson, 1980). In acidic solution, CuCN will form HCN, which may be released from the aqueous phase. At a pH of 3, 78% of the cyanide may be

removed from a 52 ppm CuCN solution in 30 minutes by the application of vacuum (Watson, 1973).

Simple metal cyanides can form a variety of highly soluble, complex metal cyanides in the presence of alkali cyanides: $A_yM(CN)_x$. In this formula, A represents the alkali, M the heavy metal (ferrous and ferric iron, cadmium, copper, nickel, silver, zinc, or others). Initial dissociation of each of these soluble, alkali-metallic, complex cyanides yields an anion that is the radical $M(CN)_x^{-y}$. This may dissociate further, depending on several factors, with the liberation of CN and consequent formation of HCN. The degree of dissociation of the various metallocyanide complexes at equilibrium, which may not be attained for a long time, increases with decreased concentration and decreased pH, and is inversely related to their highly variable stability. The zinc- and cadmium-cyanide complexes are dissociated almost totally in very dilute solutions; thus these complexes can result in acute toxicity to fish at ordinary pH. In equally dilute solutions there is much less dissociation for the nickel-cyanide complexes and more stable cyanide complexes formed with copper and silver. Acute toxicity (to fish) of dilute solutions containing copper- or silver-cyanide complex anions can be mainly due to the toxicity of the complex ions, and not HCN. The iron-cyanide complex ions are very stable.

Potassium ferricyanide is less soluble in ethanol than in water (Towill et al., 1978). The aqueous solution is unstable and decomposes slowly on standing. This compound normally does not produce HCN or cyanide ions in water, but may produce HCN when exposed to sunlight or UV radiation (Towill et al., 1978).

The metal cyanides can be oxidized to isocyanate and ultimately to CO_2 and N_2 in the presence of strong oxidizing agents such as Cl_2 gas, O_3 gas, catalytic O_2 and peroxides. In fact, such processes, in addition to electrolytic decomposition, ionizing radiation, heat, activated carbon adsorption, ion floatation and liquid-liquid extraction are suitable for the treatment of cyanide wastes (Watson, 1973).

Environmental Fate in Aquatic Media

The three likely chemical processes that may cause loss of simple cyanides in aquatic media are oxidation, hydrolysis and photolysis. Cyanides are known to be oxidized to isocyanates by strong oxidizing agents. The isocyanates may then hydrolyze to amine and CO_2 (Towill et al., 1978). Whether such oxidation and subsequent hydrolysis of isocyanates will significantly occur in natural waters known to contain peroxy radicals has not yet been determined.

The hydrolysis of alkali-metal cyanides proceeds rapidly in aquatic media, with production of HCN and alkali-metal hydroxides. The HCN produced may undergo further hydrolysis according to the following reaction (Callahan et al., 1979):

$$HCN \rightleftharpoons HCN: \xrightleftharpoons{H_2O} H\text{-}CO\text{-}NH_2 \xrightleftharpoons{H_2O} NH_4^+ + HCO^-$$

The reaction rate for the alkaline and acidic hydrolysis of HCN at the normal pH region (6-9) encountered in natural aquatic media is too slow (Callahan et al., 1979) to be significant in determining the fate of cyanides in aquatic media.

The photolysis of HCN and cyanide ions in aqueous solution was studied by Frank and Bard (1977). These compounds were found to be very resistant to photolysis with available sunlight. However, in the presence of TiO_2 powder, more than 99% of the 26 mg CN^-/ℓ of solution was oxidized in 2 days with sunlight irradiation (Frank and Bard, 1977), presumably through a heterogeneous photocatalytic oxidation process. However, such photocatalytic oxidation may not be very significant in natural aquatic systems because of significant light reduction at increasingly greater depths below the surface.

The photodecomposition of aqueous ferrocyanide and ferricyanide solution in sunlight, with the resultant production of HCN, has been observed (Callahan et al., 1979). A 5-hour sunlight exposure of 100 mg/ℓ potassium ferrocyanide produced cyanide ions at a concentration of 6 mg/ℓ (Callahan et al., 1979). While it is known that such compounds will photolytically produce HCN in sunlight, the importance of this photodecomposition process in determining the fate of ferrocyanide and ferricyanide in aquatic media cannot be determined unless the rate constants for these processes are known.

Several investigators have demonstrated that cyanides in aquatic media can be biodegraded at low concentrations by both single and mixed microorganisms (Callahan et al., 1979). Both aerobic and anaerobic microbial degradation of cyanide during sewage treatment plant operations have also been demonstrated (Callahan et al., 1979). It is evident from the literature that cyanides at low concentrations can be biodegraded in natural surface waters. However, additional data are needed to assess the relative importance of this process in determining the fate of aquatic cyanides. The

studies by Raef et al. (1977a,b) indicate that volatilization of HCN from aquatic media may be a far more significant process for cyanides than biodegradation.

The two physical processes that contribute to the loss of cyanides from aquatic media are volatilization and sorption. In most natural waters, cyanides will be present both as CN^- and HCN. As has been shown in the Physical and Chemical Properties Section, the percent of HCN in aqueous solution increases with decreasing pH of the solution.

Since the pH of most natural waters ranges between 6 and 9, a large percent of dissociated cyanide anion will be present in the form of HCN. Hydrogen cyanide being extremely volatile, the undissociated HCN is likely to volatilize from aquatic media. It has been shown that the half-life of HCN evaporation from solutions at concentrations of 25-200 μg/ℓ ranges from 22-110 hours in the laboratory. When the experiments were performed outdoors at moderate wind-speed, the rate of HCN loss increased by a factor of 2-2.5 (Callahan et al., 1979).

Hydrogen cyanide and alkali-metal cyanides are not likely to be strongly sorbed onto sediments because of high water solubilities. It was shown by Raef et al. (1977b), in an aerobic microbial biodegradation study of a cyanide, that the biological solids removed very little cyanide from solution through adsorption. From this result and the results of adsorption studies of other investigators (Callahan et al., 1979), it appears that sorption may not be an important reaction of water soluble cyanides in aquatic media.

There are no data available to indicate bioconcentration of cyanide in aquatic organisms. Using the equation of Veith et al. (1979) for the bioconcentration factor (BCF) of a chemical in whole fish (log BCF = 0.76 log K_{ow}) and the value of K_{ow} given in Table II-2, BCF values of 1.9 and 0.27 can be calculated for HCN and NaCN, respectively. According to U.S. EPA (1980), cyanides are not bioaccumulated in aquatic organisms, and the concentration of cyanides in tissues is considered the same as the concentration of cyanide in the surrounding media. However, there is some evidence of bioaccumulation of metal cyanide complexes in fish, although the actual bioaccumulation factors are not known. It is difficult to assess the environmental significance of bioaccumulation of cyanide complexes since they are far less toxic than soluble HCN, NaCN and KCN.

Summary

Hydrogen cyanide and the alkali-metal cyanides are very soluble in water and except for HCN are not very soluble in organic solvents. The alkaline-earth metal cyanides, in general, are not very soluble in water. Hydrogen cyanide is highly volatile, and in aqueous solution is a very weak acid. The alkali-metal salts of cyanides are easily hydrolyzed in an aqueous solution. The metal cyanides can be oxidized to isocyanate and ultimately to CO_2 and N_2 in the presence of strong oxidizing agents. Some of the complex metal cyanides, such as the ferrocyanides and ferricyanides, liberate HCN when the solutions are exposed to sunlight.

The fate of cyanides in aquatic media may vary widely. Hydrogen cyanide and the most common alkali-metal cyanides may be lost from aquatic media primarily through volatilization. Some of these cyanides may also be lost

through microbial degradation in aquatic media and sorption to particulate matters in water. However, both these processes are less significant than volatilization.

The sparingly soluble metal cyanides, such as CuCN, may form some HCN in acidic waters (pH <7) that may subsequently volatilize. However, the predominant fate of these compounds is sedimentation and microbial degradation.

The water-soluble complex metal cyanides, such as ferrocyanide and ferricyanide, may undergo some photodecomposition. The fate of these compounds with respect to biodegradation is not known with certainty. In the absence of destabilizing factors in water (high temperature, sunlight, extreme pH conditions), these complexes are expected to have long lifetimes and they can be transported in the aquatic media.

The simple metal cyanides and HCN do not bioaccumulate in aquatic organisms. The water-soluble complex metal cyanides, on the other hand, may bioaccumulate to some extent in aquatic organisms, although the bioaccumulation factors for such compounds are not known.

III. TOXICOKINETICS

Absorption

The cyanides are rapidly absorbed by animals and humans, whether exposure occurs by inhalation of HCN gas, by ingestion of the cyanide salts KCN or NaCN, or by dermal exposure to HCN gas or to aqueous solutions of HCN, KCN and NaCN.

Gettler and Baine (1938) reported some early observations of quantitative absorption data in dogs and humans. Three dogs received single gavage doses of KCN equivalent to 1.57, 4.42 and 8.40 mg HCN/kg bw corresponding to amounts of 20, 50 and 100 mg HCN, respectively. The dogs died within 155, 21 and 8 minutes, respectively, after dosing. At necropsy, the amount of HCN remaining in the GI tract was measured. The difference between the amount administered and the amount remaining in the GI tract was considered to represent the amount absorbed. According to this definition, the low-dose dog absorbed 14.4 mg of the administered 20 mg or 1.13 mg/kg bw (72%). The dog that received 50 mg HCN absorbed 12 mg or 1.06 mg/kg bw (24%). The dog that received 100 mg HCN absorbed 16.6 mg or 1.39 mg/kg bw (16.6%). In 3 cases of fatal human poisoning, Gettler and Baine (1938) estimated the total absorbed dose of cyanide as 228.1 (15.7%), 101.0 (18.7%) and 59.9 (19.5%) mg HCN. The actual time of death for these cases was unknown. In a fourth case in which the time of death occurred 3 hours after the ingestion of 29.8 mg HCN, the amount of HCN absorbed was estimated to be 24.4 mg (81.9%).

Liebowitz and Schwartz (1948) reported a case of attempted suicide in which an 80 kg man had ingested an estimated 3-5 g (37.5-62.5 mg/kg bw) KCN and recovered. He vomited 0.5 hour following ingestion, and gastric lavage

was performed immediately on admission to a hospital. Two hours after ingestion the concentration of HCN in blood was 200 mg/ℓ, giving an estimated total of 1200 mg HCN in blood (assuming a volume of 6 ℓ of blood) and 2400 mg HCN in the whole body. These HCN concentrations correspond to 192 mg CN^-/ℓ blood, and an estimated total of 1152 mg CN^- in blood and 2304 mg CN^- in the whole body.

Absorption of HCN across the GI mucosa proceeds rapidly because it is a weak acid with a pK_a of 9.2. The acidic environment in the stomach favors the nonionized form of HCN and, hence, facilitates the absorption. The physiologic pH of body tissues also favors the undissociated species (U.S. EPA, 1980; Callahan et al., 1979).

Quantitative data on absorption of HCN by inhalation in dogs was reported by Gettler and Baine (1938). The HCN gas was generated into an inhalation apparatus from a tube containing a solution of hydrocyanic acid. Exhaled cyanide was bubbled through a solution of sodium hydroxide. The amount of absorbed HCN was estimated by subtracting the amount of cyanide remaining in the apparatus from the initial amount in the tube. Although the original amount of HCN in the tube was not reported, one dog was calculated to have absorbed 16.0 mg (1.55 mg/kg bw) and the other dog 10.1 mg (1.11 mg/kg bw). These doses were fatal to the dogs in 15 and 10 minutes, respectively.

Absorption of low levels of cyanide by humans following inhalation can be inferred from data on blood levels of thiocyanate in cigarette smokers compared with nonsmokers (Wilson and Matthews, 1966). Although plasma levels of cyanide did not differ significantly between the two groups,

smokers had significantly ($p<0.001$) higher levels of thiocyanate, presumably formed from metabolism of inhaled cyanide in cigarette smoke, than did non-smokers.

The air concentrations of HCN resulting in certain blood levels in humans were tabulated by Knowles and Bain (1968). At >300 ppm (330 mg/m³), blood levels reached >10 mg/ℓ. Concentrations of >200 ppm (220 mg/m³), >110 ppm (111 mg/m³) and >50 ppm (55 mg/m³) in air resulted in blood levels of 8-10, 3-8 and 2-4 mg/ℓ, respectively. It was emphasized that wide variations exist, and the values were presented as examples. Humans retained ~60% of HCN in the lungs, following the inhalation of HCN (0.0005-0.02 mg/ℓ) through normal breathing by mouth (Landahl and Herrmann, 1950).

Cyanides can also be absorbed by the dermal route. Walton and Witherspoon (1926) held a test tube of 1-inch diameter containing liquid HCN against the shaved bellies of guinea pigs and observed signs of toxicity. Dogs that were exposed to HCN gas in a chamber that allowed their heads to be excluded from exposure also developed toxic signs.

Drinker (1932) reported that three men working in an atmosphere of 2% (v/v) (20,000 ppm or 22,000 mg/m³) HCN gas for 8-10 minutes developed symptoms of dizziness, weakness and throbbing pulse, despite the fact that they were wearing gas masks. Potter (1950) described a case of an HCN process worker who entered the chamber wearing a gas mask and protective clothing. While attempting to take a sample of hydrocyanic acid, he removed one glove, and some of the acid ran over his hand. Within 5 minutes, he became dizzy, had difficulty breathing and became unconscious.

Hydrogen cyanide, either liquid or gas, is more readily absorbed dermally than are the cyanide salts (Wolfsie and Shaffer, 1959), although cases of toxicity in humans have resulted from dermal exposure to solutions of KCN (NIOSH, 1976). Since HCN can be generated from solutions of KCN and NaCN, some absorption may be attributed to inhalation exposure (see Chapter II).

Distribution

Once cyanide is absorbed it is rapidly distributed by the blood throughout the body. Within the blood, a greater concentration of cyanide was found in erythrocytes than in plasma when HCN was administered (McMillan and Svoboda, 1982); however, this was not the case when KCN was administered (Farooqui and Ahmed, 1982). Farooqui and Ahmed (1982) studied the incorporation of radioactivity in the erythrocytes of rats treated orally with $K^{14}CN$ (5 mg/kg). Levels of radioactivity declined rapidly from whole blood and plasma with a small increase in erythrocytes over 24 hours. Following hemolysis, the majority of the radioactivity (94.32%) in the erythrocytes was found in the hemolysate rather than the membranes. The heme fraction contained 70% of the radioactivity while 14-25% and 5-10% were found in the globin and cell membrane, respectively. Since cyanide reacts readily with iron in the ferric state (Fe^{+++}) (Hartung, 1982; NIOSH, 1976), the accumulation of cyanide within erythrocytes is mainly due to the binding of the cyanide ion to Fe^{+++} in methemoglobin to form the nontoxic complex cyanomethemoglobin (Chen and Rose, 1952).

Feldstein and Klendshoj (1954) reported blood levels of cyanide in normal humans that ranged from 0-0.107 μg/mℓ with an average of 0.048 μg/mℓ.

Tissue levels of cyanide in humans fatally poisoned with cyanide have been obtained at autopsy (Ansell and Lewis, 1970; Gettler and Baine, 1938; Finck, 1969). In examining the data, two issues should be noted. First, when the concentration found equals zero, a loss of HCN from the sample must be suspected. This could occur by volatilization between the time of death and the time of assay, or by metabolism of CN^- to SCN^- (which was not assayed). Secondly, the time between exposure and death depends on the dose of cyanide, which in these cases was unknown. If death occurred very rapidly, it is possible that the tissue distribution of cyanide was not complete and, therefore, tissue levels of cyanide could be irrelevant. Ansell and Lewis (1970) tabulated the mean levels of cyanide found in postmortem samples of tissues (Table III-1). The highest concentrations of cyanide were found in the spleen and blood. Gettler and Baine (1938), however, found high cyanide levels in brains and livers of three humans who had ingested fatal doses of cyanide. The importance of analyzing many tissues to determine the route of exposure was emphasized by Finck (1969). Table III-2 presents the distribution of cyanide in tissues of three fatally poisoned humans. For case one, no lung tissue had been submitted to the toxicologist for analysis. Based on low levels of cyanide in the gastric contents, cyanide inhalation may be suspected. Cases two and three, however, demonstrated high levels in stomach contents following ingestion.

The differences in cyanide distribution following the oral and inhalation exposures in rats were examined by Yamamoto et al. (1982). Rats were treated by gavage with NaCN to give a dose of 7 and 21 mg CN^-/kg bw. Another group of rats inhaled HCN at concentrations that averaged 356 and 1180 ppm (392 and 1298 mg/m^3, respectively). The results are presented in Table III-3. Since little difference in distribution was observed with

TABLE III-1

Mean Levels and Ranges of Cyanide Ion Concentration in Human Organs in Cases of Fatal Poisoning[a,b]

Tissue	No. of Cases with Quantitative Levels Available	Mean (mg %)	Range (mg %)
Blood	58	2.39	0-5.3
Brain	34	1.20	0-19.9
Liver	48	1.62	0-25.0
Kidney	34	0.61	0-2.8
Spleen	22	3.77	0-37.5
Stomach Contents	49	160.0	0.2-2800
Urine	17	0.08	0-0.96

[a]Source: Adapted from Ansell and Lewis, 1970

[b]Note: for variable routes of exposure

TABLE III-2

Cyanide Levels in Human Tissues and Fluids After Fatal Cyanide Poisoning[a]

Sample	Cyanide Content (mg/100 g or mg/100 mℓ)		
	Case 1[b]	Case 2[c]	Case 3[c]
Gastric contents	0.03	15	20
Lung	NA	0.90	0.70
Blood	0.50	0.75	0.80
Liver	0.03	0.40	0.50
Kidney	0.11	0.35	0.40
Muscle	NA	0.30	NA
Brain	0.07	0.25	0.06
Urine	0.20	0.20	NA
Fat	NA	0.20	NA

[a]Source: Adapted from Finck, 1969

[b]Death from inhalation of cyanide gas

[c]Death from ingestion of cyanide salt

NA = Not analyzed

TABLE III-3

Cyanide Concentrations in Various Organs of Rats Treated with NaCN Orally (A) or HCN by Inhalation (B)[a]

Organs[c]	(A) Dose[b]			(B) Concentration		
	7 mg/kg (n=10)	21 mg/kg (n=9)	Combined[d] (n=19)	392 mg/m³ (n=12)	1298 mg/m³ (n=12)	Combined[d] (n=24)
Blood	5.01 ± 1.61	4.79 ± 2.04	4.91 ± 1.78	2.91 ±0.41	3.11 ± 0.51	3.01 ± 0.46
Liver	7.53 ± 1.45 (161 ± 50)[e]	10.40 ± 2.17 (230 ± 47)	8.89 ± 2.30 (194 ± 59)	2.19 ± 0.77 (76 ± 29)	2.11 ± 1.10 (68 ± 32)	2.15 ± 0.93 (72 ± 30)
Lungs	5.44 ± 2.27 (114 ± 46)	6.30 ± 2.19 (146 ± 66)	5.85 ± 2.22 (129 ± 57)	4.78 ± 1.61 (170 ± 69)	4.08 ± 2.24 (141 ± 104)	4.42 ± 2.02 (156 ± 88)
Spleen	2.48 ± 0.98 (52 ± 22)	1.76 ± 0.50 (36 ± 12)	2.09 ± 0.88 (45 ± 19)	0.58 ± 0.28 (20 ± 9)	0.79 ± 0.81 (28 ± 24)	0.68 ± 0.60 (24 ± 18)
Brain	1.31 ± 0.47 (27 ± 8)	1.76 ± 0.70 (40 ± 8)	1.52 ± 0.43 (33 ± 10)	1.34 ± 0.54 (45 ± 14)	1.56 ± 0.38 (50 ± 10)	1.45 ± 0.47 (48 ± 12)
Time to death (minutes)	10.3 ± 4.3	3.3 ± 2.1	NA	9.6 ± 2.3	5.4 ± 0.9	NA

[a]Source: Yamamoto et al., 1982

[b]The dose is expressed as CN^-

[c]Organ levels are µg/g wet weight or µg/mL blood

[d]Data from the two dose groups were combined

[e]The number in parentheses are the organ levels of CN expressed as mean % of the blood levels

NA - Not available

respect to dose or concentrations, the authors combined the data. Although these combined data are provided in Table III-3, one must examine the appropriateness of doing so. The results of time to death indicate that animals receiving 21 mg/kg bw or 1298 mg/m³ died within significantly less time (average 3.3 and 5.4 minutes, respectively) than those receiving 7 mg/kg bw or 392 mg/m³ (average time to death 10.3 and 9.6 minutes, respectively). This suggests that a uniform body distribution of cyanide may not have been reached at the time of death of the animals receiving the higher dose. In order to compare the distributions for the two routes of administration, the organ levels of cyanide were examined and expressed as percentage of the respective blood levels. Following oral administration, cyanide levels in the liver were higher than those in the blood, and were dependent on the dose ingested. No differences were observed in the lungs. Following inhalation exposure, cyanide levels in both the liver and lungs did not differ from that observed in the blood and were unrelated to the concentration in the air. In the spleen, cyanide levels were higher following oral administration than inhalation regardless of the dose or concentration. These results suggest that body distribution of cyanide depends primarily on the route of administration.

The pattern of distribution of cyanide may also depend on the type of compound administered. Ballantyne et al. (1972) compared the tissue distribution of cyanide from KCN and HCN in rabbits following intramuscular injection. For either case, the administered dose of cyanide was 8 mg/kg. Levels of cyanide were measured in tissues of two groups of rabbits (6/group) at 0.5 hours after death. In another two groups of rabbits, the organs were perfused with saline in order to measure the actual tissue levels without interference from blood cyanide levels in the tissues. The

results are presented in Table III-4. The higher levels of cyanide in blood and tissues following HCN rather than KCN administration were presumably due to the more rapid absorption and distribution of HCN. High levels of cyanide were found in perfused brain, spinal cord and liver.

Cyanide does not accumulate in blood and tissues following chronic exposure. Howard and Hanzal (1955) treated rats with dietary concentrations of HCN of 100 and 300 ppm (mg/kg diet) for 2 years. At the end of the 2-year period, virtually no cyanide was found in plasma or kidneys. Low levels were found in erythrocytes (mean of 1.97 μg/100 mℓ). Increased levels of thiocyanate, the less toxic primary metabolite of cyanide (see Metabolism Section) were found in plasma (1123 μg/100 mℓ), erythrocytes (246 μg/100 mℓ), liver (665 μg/100 g) and kidney (1188 μg/100 g).

Transplacental transfer of cyanide presumably occurs. Doherty et al. (1982) reported that administration of solutions of NaCN to pregnant hamsters by subcutaneous infusion using osmotic minipumps resulted in malformed fetuses (see Teratogenicity and Other Reproductive Effects Section in Chapter V); this finding, however, does not prove occurrence of transplacental transfer.

Metabolism

The major metabolite of cyanide is the less toxic thiocyanate (U.S. EPA, 1980; Ansell and Lewis, 1970; Williams, 1959; de Duve et al., 1955), formed by the transfer of sulfur from a sulfur donor to the cyanide ion by the

TABLE III-4

Comparison of Cyanide Concentrations in Tissues from Rabbits Killed by HCN with Concentrations in Tissues from Rabbits Killed with KCN[a,b]

Tissue	Cyanide concentration mean ± standard error[c]		p[d]
	HCN	KCN	
	Containing blood		
Skeletal muscle	35.0 ± 5.2	29.6 ± 2.4	<0.5
Kidney	74.7 ± 10.3	52.0 ± 11.0	<0.1
Liver	148.7 ± 32.3	82.0 ± 8.0	<0.1
Spinal cord	48.5 ± 4.9	36.8 ± 3.5	<0.1
Brain	145.3 ± 37.2	106.5 ± 12.4	<0.1
Whole blood	685.0 ± 83.0	453.0 ± 34.0	<0.05
Serum	275.0 ± 18.0	161.0 ± 21.0	<0.005
	Perfused with saline		
Skeletal muscle	9.3 ± 2.7	7.8 ± 2.4	<0.7
Kidney	11.0 ± 4.3	2.3 ± 1.1	<0.1
Liver	43.7 ± 13.5	6.5 ± 0.8	<0.025
Spinal cord	49.8 ± 14.7	22.5 ± 3.8	<0.2
Brain	289.0 ± 67.7	98.0 ± 5.0	<0.02
Whole blood	761.0 ± 129.0	438.0 ± 8.0	<0.05
Serum	261.0 ± 48.0	134.0 ± 8.0	<0.05

[a]Source: Ballantyne et al., 1972

[b]For both compounds the dose was equivalent to 8 mg CN^-/kg bw

[c]Concentrations expressed as µg CN^-/100 g tissue wet weight or µg CN/100 mℓ blood or serum

[d]Significance of difference in cyanide concentrations between animals killed with HCN and those killed with KCN

enzymatic action of rhodanese (Lang, 1933). Rhodanese is the trivial name for thiosulfate: cyanide sulfurtransferase, EC 2.8.1.1. or 3-mercapto-pyruvate:cyanide sulfurtransferase EC 2.8.1.2. (Pettigrew and Fell, 1973). The species and tissue distribution of rhodanese were investigated by Himwich and Saunders (1948); the results are presented in Table III-5. The activity of the enzyme was highly variable among species and tissues. In the dog, the highest activity of rhodanese was found in the adrenal gland, ~2.5 times greater than the activity in liver. Monkeys, rabbits and rats had highest activity in liver and kidney, with relatively low levels of rhodanese in adrenals. It should be noted that the enzyme activity in the livers of monkeys, rabbits and rats was ~10-20 times higher than the activity in the liver of dogs, and that total activity in the other species was higher than that in dogs. Dogs are particularly sensitive to the acute effects of cyanide (see General Toxicity Section in Chapter V). Similar activities of the enzyme among the species were found for brain, testes, lungs, spleen and muscle. The ubiquitous nature of the enzyme was demonstrated in the dog for which low levels of rhodanese activity were also measured in erythrocytes, anterior pituitary, thyroid, pancreas, lymph nodes, salivary glands, optic nerve, eye, intestine and heart.

The activities of different compounds as sulfur-donors have been studied. Himwich and Saunders (1948) found that sodium thiosulfate (100% relative activity) was the best sulfur donor when compared with sodium sulfide (4%), thiourea (4.5%), α-naphthylthiourea (4.6%), thiouracil (1%) and cysteine (1%). No activity was measured when thiodiglycol, diphenyl sulfide or diphenyldisulfide were tested. Sorbo (1953) found that p-toluene thiosulfonate was ~4.5 times more active as a sulfur donor than thiosulfate, while ethyl thiosulfate, ethyl xanthate, diethyldithiocarbamate and

TABLE III-5

Rhodanese Activity in Tissues of the Dog, Rhesus Monkey, Rabbit and Rat (mg CN^- converted to CNS/g of tissue after 15 minutes)[a]

Tissue	Dog		Rhesus Monkey		Rabbit		Rat	
	Range[b]	Number of Observations	Range[b]	Number of Observations	Range	Number of Observations	Range	Number of Observations
Suprarenals								
whole	2.14-3.60 (5.46, 4.50)	6	0.14-1.35	3	1.24-3.94	2	0.27-0.41	2
cortex	2.06-5.62	2	ND	NA	ND	NA	ND	NA
medulla	0.27-1.12	2	ND	NA	ND	NA	ND	NA
Liver	0.78-1.46 (4.91, 6.28)	7	10.98-15.16 (5.98)	4	7.98-18.92	9	14.24-28.38	9
Brain								
cortex	0.34-0.92	7	0.27	1	1.41-1.44	2	0.70-0.72	2
caudate nucleus	0.27-1.06	7	0.34-0.50	2	0.13-0.18	2	ND	NA
midbrain	0.52-1.35	6	0.22-0.80	2	1.17-1.39	2	0.73-1.13	2
cerebellum	0.21-1.22	7	0.33	1	0.63-1.24	2	ND	NA
medulla	0.38-1.52	7	0.49-0.85	2	0.91	1	ND	NA
Spinal cord								
cervical	0.15-1.08	7	0.56-0.57	2	0.89-0.90	2	0.16-0.18	2
lumbar	0.12-0.84	4	0.20-0.52	2	0.35-1.74	2	0.23-0.27	2
sacral	0.16-1.41	4	0.23-0.28	2	0.59-1.10	3	0.56-0.74	2
Heart	0.11-0.14	6	0.48-0.82	3	ND	NA	ND	NA
Kidney	0.42-0.74	6	2.46-3.58	4	6.20-7.69	3	10.44-11.08	2
Testes	0.32-0.41	5	0.38-0.46	3	0.32-0.36	2	1.24-1.61	2
Epididymis	0.29	1	ND	NA	ND	NA	ND	NA
Ovaries	0.42	1	ND	NA	0.30	1	ND	NA
Lung	0.16-0.17	3	0.11-0.21	2	0.40	1	ND	NA
Spleen	0.10-0.14	2	0.12-0.34	2	0.20	1	ND	NA
Muscle	0.03-0.19	6	0.23-0.57	3	0.18	1	ND	NA
Intestine								
duodenum	0.05-0.11	3	ND	NA	ND	NA	ND	NA
jejunum	0.04	1	ND	NA	ND	NA	ND	NA

TABLE III-5 (cont.)

Tissue	Dog		Rhesus Monkey		Rabbit		Rat	
	Range[b]	Number of Observations	Range[b]	Number of Observations	Range	Number of Observations	Range	Number of Observations
Eye	0.02	1	ND	NA	ND	NA	ND	NA
Optic nerve	0.35	1	ND	NA	ND	NA	ND	NA
Salivary gland, parotid	0.05-0.36	3	0.99	1	ND	NA	ND	NA
Lymph node	0.08-0.13	2	ND	NA	ND	NA	ND	NA
Pancreas	0.14-0.28	4	0.12-0.44	2	ND	NA	ND	NA
Thyroid	0.05-0.94	3	ND	NA	ND	NA	ND	NA
Anterior pituitary	0.26	1	ND	NA	ND	NA	ND	NA
Whole blood	0.01-0.02	2	ND	NA	ND	NA	ND	NA
Erythrocytes	0.01-0.02	2	ND	NA	ND	NA	ND	NA
Plasma	0.01	1	ND	NA	ND	NA	ND	NA

[a]Source: Himwich and Saunders, 1948

[b]Figures in parentheses are single observations falling outside the normal range.

ND - Not determined; NA - Not applicable

hydrosulfite were relatively inactive. The structural requirements, therefore, were for a free sulfur to be adjacent to another sulfur atom in the molecule. Since the greatest activity of rhodanese is found in mitochondria (de Duve et al., 1955), and rhodanese is widely distributed among tissues (Himwich and Saunders, 1948), the rate limiting factor for cyanide detoxification to thiocyanate is the intracellular, and especially intramitochondrial, availability of an appropriate endogenous sulfur donor. The nature of the endogenous sulfur donor is unknown (U.S. EPA, 1980). Westly (1983) hypothesized that the sulfone carrier is albumin. Radioisotope studies have indicated that albumin does interact with elemental sulfur and that this complex can react with cyanide.

Pettigrew and Fell (1973) investigated cyanide-thiocyanate interconversion by administering KCN i.p. to Wistar rats fed either a normal diet or a vitamin B_{12}-deficient diet. Both groups received a daily dose of 9.2 μmole (0.6 mg) CN^-/100 g bw, 5 days/week for 3 weeks. This dose approximates the minimum dose that has previously been shown (Lessell, 1971) to produce neurologic disturbances in rats. Among the vitamin B_{12}-deficient rats, the mean whole blood CN^- concentrations rose 6-fold while the mean plasma thiocyanate concentration rose 8-fold. Among the normally fed rats, blood was sampled at various intervals following the injection. The whole blood CN^- concentration rose quickly reaching a maximum ~20 minutes post-injection and then fell rapidly to minimal levels ~2 hours postinjection. Conversely, the plasma thiocyanate levels increased slowly, reaching a maximum between 1 and 4 hours postinjection after which it decreased gradually.

A detoxification rate of 0.076 mg/kg/minute was determined in guinea pigs during continuous intravenous infusion of cyanide (Lendle, 1964). The estimated detoxification rate in humans has been estimated as 0.017 mg/kg/minute (McNamara, 1976). This estimate was based on a study in men injected intravenously with HCN.

The overall rate of conversion of cyanide to thiocyanate will depend upon the rate of conversion of thiocyanate to cyanide. The rhodanese reaction was shown to be nonreversible; however, conversion of thiocyanate to cyanide was found to be mediated by a different enzyme, thiocyanate oxidase, which has been found in erythrocytes of humans, dogs, rabbits and rats (Goldstein and Rieders, 1951, 1953). A minor activity of the glutathione-S-transferases in converting organic thiocyanates to cyanide was reported by Habig et al. (1975).

The overall rate of detoxification of cyanide will also depend upon the contribution of minor pathways. Cyanide can react spontaneously with cystine to yield 2-aminothiazoline-4-carboxylic acid, which tautomerizes to 2-imino-4-thiazolidinecarboxylic acid (Wood and Cooley, 1956). Rats pretreated with ^{35}S-cystine excreted 16% of a subcutaneous dose of NaCN as ^{35}S-labeled 2-imino-4-thiazolidine-carboxylic acid in the urine. Thiocyanate in urine accounted for 80% of the cyanide dose.

Another minor route of cyanide detoxification is via the reaction of cyanide with the B_{12} vitamin, hydroxocobalamin, to yield cyanocobalamin, a complex that is an essential element, is nontoxic, and is excreted in urine (Brink et al., 1950; Boxer and Rickards, 1951) and bile (Herbert, 1975). A metabolic pathway involving the oxidation of cyanide to CO_2 and formate

was studied by Boxer and Rickards (1952b). A dog injected with ~0.1 mg $^{14}CN^-$/kg, once an hour for 5 hours, excreted 0.007% of the dose as formate and 92% as thiocyanate over 2 days. Rats similarly treated with $^{14}CN^-$ (215 µg/rat/injection), excreted 1.7% of the dose in the expired air. Of this amount of radioactivity, 10% was $H^{14}CN$ and 90% was $^{14}CO_2$. Thus, carbon derived from cyanate can enter one-carbon-compound metabolic pathways. The various pathways of cyanide biotransformation are summarized in Figure III-1.

Excretion

By far the major route of cyanide elimination from the body is via urinary excretion of thiocyanate. Rats eliminated 80% of subcutaneously-injected cyanide as thiocyanate in the urine, while 16% was eliminated as urinary 2-imino-4-thiazolidine-carboxylic acid (Wood and Cooley, 1956). A man who had ingested 3-5 g KCN (1.2 g HCN was estimated in blood, assuming 6 ℓ blood, 2 hours later) eliminated a total of 237.14 mg of thiocyanate in 72-hour urine (Liebowitz and Schwartz, 1948). Maliszewski and Bass (1955) found increased levels of thiocyanate in the urine, saliva and perspiration of cigarette smokers compared with nonsmokers. Okoh (1983) reported the pattern of elimination of ^{14}C in rats maintained on a diet of KCN (~2 mg CN^-/rat/day) for 6 weeks, and then injected subcutaneously with $Na^{14}CN$. Over 57% of the administered radioactivity was excreted in the 24-hour urine compared with 4.9% in expired air and 1.7% in feces. Thiocyanate accounted for 78.8% and cyanide for 1.3% of the urinary radioactivity. Of the radioactivity in expired air, 89.5% was $^{14}CO_2$ and 9.4% was $^{14}CN^-$. In a similar study, Okoh and Pitt (1982) also found radioactivity excreted in feces following intraperitoneal injection of $Na^{14}CN$. This finding, along with relatively high levels of radioactivity in the stomach



FIGURE III-1

Fate of Cyanide in the Body

Source: Ansell and Lewis, 1970

wall and stomach contents compared with other tissues, suggested that GI recirculation was occurring. Approximately 80% of the radioactivity in the stomach contents was in the form of thiocyanate.

The finding of low levels of unchanged HCN in expired air of rats and humans was reported by Boxer and Rickards (1952a). Guinea pigs eliminated 1-2% of intravenously-infused HCN as unchanged gas in the expired air (Friedberg and Schwarzkopf, 1969). Ohsy et al. (1987) found that >50% of an administerd dose of cyanide was excreted in the urine of hens as thiocyanate within 6 hours. Little cyanide was excreted directly and the rate of thiocyanate excretion closely paralleled plasma thiocyanate concentrations. As mentioned previously in the Metabolism Section, cyanocobalamin is excreted in urine (Boxer and Rickards, 1951). Additionally, in humans, 3-8 μg of cyanocobalamin are excreted into the GI tract, mainly in the bile. All but 1 μg of this excreted cyanocobalamin is reabsorbed by the intrinsic factor mechanism at the level of the ileum, thus allowing effective enterohepatic recirculation of vitamin B_{12} (Herbert, 1975).

Summary

Cyanides are readily absorbed from the lungs, the GI tract and skin (U.S. EPA, 1980). Following uptake by blood, cyanide is rapidly distributed throughout the body. Cyanide accumulates within erythrocytes (Farooqui and Ahmed, 1982; McMillan and Svoboda, 1982) where it combines with Fe^{+++} of methemoglobin (Chen and Rose, 1952) and the heme moiety of hemoglobin (Farooqui and Ahmed, 1982). Cyanide can be detected at low levels in blood of normal humans (Feldstein and Klendshoj, 1954). Relatively high levels of cyanide have been found in spleen, brain and liver at autopsy of humans fatally poisoned with cyanide. Distribution patterns differ with respect to

route of exposure (Finck, 1969; Yamamoto et al., 1982) and compound (Ballantyne et al., 1972). Cyanide and thiocyanate accumulate in blood and tissues following chronic exposure (Howard and Hanzal, 1955). The major route of metabolism and detoxification of cyanide is via rhodanese (thiosulfate:cyanide sulfurtransferase), which catalyzes the transfer of sulfur from thiosulfate to cyanide to yield thiocyanate (Lang, 1933; Sorbo, 1953). Rhodanese is widely distributed and the activity is highly variable among tissues and species; high levels are found in the liver and kidneys of monkeys, rabbits and rats and in the adrenals of dogs (Himwich and Saunders, 1948). The greatest activity of rhodanese is located in mitochondria (de Duve et al., 1955), and the enzyme is specific for a sulfur donor with a free sulfur atom adjacent to another sulfur atom in the molecule (Sorbo, 1953). The rate of detoxification in guinea pigs has been estimated at 0.076 mg/kg/minute (Lendle, 1964) and in humans as 0.017 mg/kg/minute (McNamara, 1976). Minor detoxification pathways include the spontaneous reaction of cyanide with cystine to form 2-imino-4-thiazolidinecarboxylic acid (Wood and Cooley, 1956), and the reaction of cyanide with hydroxocobalamin (vitamin B_{12a}) to yield cyanocobalamin (vitamin B_{12}) (Brink et al., 1950; Boxer and Rickards, 1951). Both of these compounds are excreted in urine. Cyanide can also be oxidized to CO_2 and formate (Boxer and Rickards, 1952b). The major route of cyanide elimination is via urinary excretion of thiocyanate (Wood and Cooley, 1956; Okoh, 1983). Small percentages of radioactivity derived from $Na^{14}CN$ administered intraperitoneally to rats were excreted in expired air and feces (Okoh, 1983). The fecal excretion, coupled with the detection of radioactivity in the stomach wall and contents following intraperitoneal injection of ^{14}CN, suggested GI recirculation (Okoh and Pitt, 1982).

IV. HUMAN EXPOSURE

This chapter will be submitted by the Science and Technology Branch, Criteria and Standards Division, Office of Drinking Water.

IV. HUMAN EXPOSURE

Humans may be exposed to chemicals such as cyanide from a variety of sources, including drinking water, food, ambient air, occupational settings and consumer products. This analysis of human exposure to cyanide is limited to drinking water, food and ambient air because those media are considered to be sources common to all individuals. Even in limiting the analysis to these three sources, it must be recognized that individual exposure will vary widely based on many personal choices and on several factors over which there is little control. Where one lives, works and travels, what one eats, and physiologic characteristics related to age, sex and health status can all profoundly affect daily exposure and intake. Individuals living in the same neighborhood or even in the same household can experience vastly different exposure patterns.

In the Exposure Estimation Section of this chapter, available information is presented on the range of human exposure and intake for cyanide from drinking water, food and ambient air for the 70 kg adult male. It is not possible to provide an estimate of the number of individuals experiencing specific combined exposures from those three sources.

Exposure Estimation

Drinking Water. Essentially no information was obtained indicating the occurrence of cyanide in drinking water supplies. Only one of the Federal surveys [the 1969 Community Water Survey (CWSS)] included cyanide as an analyte.

The CWSS (McCabe et al., 1970) examined 120 surface water supplies, 613 groundwater supplies, 152 supplies with mixed water or purchased sources and 84 special systems (trailer parks, tourist and other institutions) in the United States in 1969. Samples were analyzed for various chemicals including cyanide. A total of 2595 distribution water samples were collected. The U.S. Public Health Service mandatory drinking water limit for cyanide in effect at the time of this study was 0.01 mg/ℓ. Cyanide concentrations in all samples were below this limit. The maximum cyanide concentration reported was 0.008 mg/ℓ.

According to U.S. EPA (1980), cyanide is an uncommon pollutant in U.S. water supplies. Levels in excess of the U.S. Public Health Service mandatory limit (McCabe et al., 1970) were not documented in U.S. EPA (1980). Apparently, the general recognition of the high toxicity of cyanide has made its removal standard practice in industry. As reported in U.S. EPA (1980), accidental perturbations have resulted in fish kills, livestock death and other environmental damage. Approximately 1500 drums (30 and 55 gallon) containing cyanides disposed near Byron, IL caused significant environmental damage and livestock death. Surface water runoff from the area was reported to contain up to 365 mg/ℓ of cyanide (Towill et al., 1978).

Given the lack of information on cyanide in public drinking water supplies, human intake of cyanide from drinking water could not be evaluated.

<u>Diet</u>. Although it is uncommon to find cyanide in foods in the United States, certain nitriles naturally occur in plants such as soybeans, lima beans and cassava (U.S. EPA, 1980; Honig et al., 1983). Honig et al. (1983)

found levels of cyanide <0.1 mg/kg in raw soybean meal and <1.5 mg/kg in soybean product samples tested under standard conditions. Honig and coworkers maintained that, according to the Food Protein Council (1978), consumption of edible soy protein products in the United States is >1 billion pounds annually, which is equivalent to <3 g of protein/person/day. The authors, therefore, concluded that soybean cyanide levels do not appear to be of nutritional significance.

Although the lima bean has been studied more than any other cyanogenetic plant food consumed by man (Leiner, 1966), none of these studies have focused on cyanide levels in lima beans consumed in the United States. Similarly, the cassava plant is used to produce tapioca, a food infrequently eaten in the United States; no studies have been conducted in the United States on cyanide levels in cassavas (Honig et al., 1983).

No data are available from the Food and Drug Administration (FDA) on the occurrence of cyanide in foods nor is any information available on the daily dietary intake of cyanide.

Air. No information is available on the levels of cyanide in ambient air.

Summary

Insufficient information was available to assess human exposure to cyanide or to determine the relative source contribution to total exposure by the three common media (drinking water, food and ambient air).

References

Food Protein Council. 1978. Soy protein improving our food system. Food Protein Council, Washington, DC. (Cited in Honig et al., 1983)

Honig, D.H., M.E. Hockridge, R.M. Gould and J.J. Rackis. 1983. Determination of cyanide in soybeans and soybean products. J. Agric. Food Chem. 31(2): 272-275.

Leiner, I.E. 1966. Cyanogenetic glycosides. In: Toxicants occurring naturally in foods. Food Protection Committee, Food and Nutrition Board, Natl. Acad. Sci., Washington, DC. p. 58-61.

McCabe, L.J., J.M. Symons, R.D. Lee and G.G. Robeck. 1970. Survey of community water supply systems. J. Am. Water Works Assoc. 62: 670-687.

Towill, L.E., et al. 1978. Reviews of the environmental effects of pollutants: V. Cyanide. U.S. EPA. (Cited in U.S. EPA, 1980)

U.S. EPA. 1980. Ambient Water Quality Criteria for Cyanides. Environmental Criteria and Assessment Office, Cincinnati, OH. EPA-440/5-80-037. NTIS PB 81-117483.

HEALTH EFFECTS IN ANIMALS

General Toxicity

There are numerous studies on the toxic effects of the cyanides administered to animals by a variety of routes. In the current discussion, studies employing the oral route of administration are emphasized. No attempt was made to review all the studies employing other routes of exposure; however, several representative reports are included for the sake of completeness.

An important consideration in evaluating oral toxicity of cyanides is not only the total amount administered but also the rate of its absorption. This follows from the fact that the liver is the major site of cyanide detoxification; like any other enzymatically catalyzed reaction this detoxification reaches a maximum rate in the presence of excess substrate. If cyanide absorption proceeds too fast, the capacity of the liver to form thiocyanate upon first pass of mesenteric blood through the organ may be exceeded. In contrast, slow absorption of the same total oral load of the poison may allow complete metabolism by the liver. For the most part, cyanide is readily absorbed from the GI tract, especially since at physiologic pH it is present mostly in the highly diffusible nonionized form. However, the rate of absorption may be influenced by factors such as the composition and volume of the intestinal contents and by the rate of peristalsis.

An additional factor, which occurs during cyanide feeding studies, is the likely loss of this substance by volatilization of HCN before the food is consumed. Even if attempts are made to chemically determine total cyanide loss, average concentrations of cyanide in food cannot be computed unless it is known whether such loss follows zero or first-order kinetics.

Two important conclusions follow from these considerations: use of feeding studies as the basis for the setting of water standards introduces an uncertainty in the case of cyanide, which must be accounted for. Secondly, toxicity of cyanide may differ greatly depending upon the route of administration; after inhalation the poison reaches the liver at a rate quite different from that following ingestion. Use of data other than oral for setting permissible drinking water standards for cyanide introduces additional uncertainties.

Acute Exposure. The lethal effects of cyanide exposure by any route are well known; however, the severity of effects and the time course depend upon the route, the dose, the duration and the compound administered. By far, inhalation of the HCN gas results in the most rapid absorption and, hence, the most rapidly appearing signs of toxicity. Gastrointestinal absorption of the salts, NaCN and KCN, is slower and results in passage through the portal system into the liver. Rhodanese, the enzyme(s) responsible for converting the cyanide ion into thiocyanate, is found in tissues throughout the body, but the major site in most species is the liver. Therefore, first passage of cyanide into the liver following GI absorption results in a greater degree of detoxification than if cyanide is absorbed from the lungs. Cyanide exerts its toxic effect by reacting with the ferric ion (Fe^{+++}) in mitochondrial cytochrome oxidase, thereby interfering with cellular respiration (see Chapter VII).

Single-dose LD_{50} values for cyanides are presented in Table V-1. For the sake of comparison, all doses are represented as the dose of the administered compound, as well as the equivalent dose of cyanide ion. Ballantyne et al. (1971, 1972) demonstrated that intramuscular administration of HCN to rabbits resulted in lower LD_{50}s than did intramuscular administration of KCN to rabbits with the LD_{50} for HCN in female rabbits being significantly lower; however, when expressed as dose of cyanide ion, the LD_{50}s were approximately equivalent. Oral LD_{50} values are available for NaCN and KCN. For NaCN, a human oral LD_{50} of 2.86 mg NaCN/kg bw (1.52 mg CN^-/kg bw) and a rat oral LD_{50} of 6.44 mg NaCN/kg bw (3.41 mg CN^-/kg bw) are reported (NIOSH, 1976). For KCN, oral LD_{50}s are reported as follows: 10 mg KCN/kg bw (4 mg CN^-/kg bw) for rats (Gaines, 1969; Hayes, 1967), 8.5 mg KCN/kg bw (3.4 mg CN^-/kg bw) for mice (Sheehy and Way, 1968), 5 mg KCN/kg bw (2 mg CN^-/kg bw) for rabbits (NIOSH, 1976) and 2.86 mg KCN/kg bw (1.14 mg CN^-/kg bw) for humans (NIOSH, 1976). Note that in the case of cyanide, such determinations differ from the classical determination of an LD_{50} (i.e., counting of deaths during 14 days after dose administration) since cyanide is extremely acutely toxic. For the LD_{50} determination in mice, the animals were observed for 24 hours because many mice that appeared to be moribund recovered. The LD_{50} for intraperitoneal NaCN in mice (Kruszyna et al., 1982) is essentially the same as that for oral KCN in mice (Sheehy and Way, 1968), when expressed in terms of cyanide ion (3.2 vs. 3.4). KCN and NaCN have been administered intraperitoneally or subcutaneously to mice and dogs in order to determine the effects of antidotal treatment on the LD_{50} (Kruszyna et al., 1982; Way et al., 1966, 1972; Isom and Way, 1973; Chen and Rose, 1952). Streicher (1951) demonstrated that an increase in temperature decreased the toxicity of KCN. The LC_{50} for inhalation (not

TABLE V-1

Single-Dose LD_{50} Values for Cyanides

Route	Administered Compound	Species/Strain	Age/Weight (g)	Sex/No.	Vehicle	LD_{50} (mg/kg bw)	LD_{50} (mg CN^-/kg bw)	Comment	Reference
Oral	KCN	rat/Sherman	adult/NR	M/50	water	10 (8.7-11.5)	4 (3.5-4.6)	The rats died within 1 hour. The LD_1 was 5.9 mg KCN/kg bw (2.4 mg CN^-/kg).	Hayes, 1967; Gaines, 1969
Oral	KCN	mouse/Swiss-Webster	NR/NR	NR/20/group	water	10.8 (10.0-11.7)	4.3 (4.0-4.7)	The compound was administered in a volume corresponding to 5% bw.	Ferguson, 1962
Oral	KCN	mouse/Swiss-Webster	NR/NR	NR/20/group	water	15.8 (14.6-17.1)	6.3 (5.8-6.8)	The volume corresponded to 1% bw.	Ferguson, 1962
Oral	KCN	mouse/Swiss-Webster	NR/18-20	M/10/group	NR	8.5 (8.1-9.0)	3.4 (3.2-3.6)	The LD_{50} is for mice dying up to 24 hours after dose. Signs of toxicity appeared in 30-90 seconds. After a short period of convulsions, mice were moribund; many at doses below LD_{50} appeared dead but recovered.	Sheehy and May, 1968
Ocular	HCN	rabbit/NR	NR/NR	NR/NR	see comment	1.05	1.01	It was stated that the toxicity of NaCN was not different whether applied in solution, as a solid or mixed with an inert powder. Deaths occurred in 3-12 minutes.	Ballantyne, 1983
	KCN					7.87	3.15		
	NaCN					5.05	2.68		
i.p.	NaCN	mouse/CD-1	NR/25-30	M/NR	water	6.0 (6.0-6.4)	3.2 (3.2-3.4)	Pretreatment of mice with hydroxylamine, dimethylaminophenol or sodium nitrite increased the LD_{50} of NaCN.	Kruszyna et al., 1982
s.c.	KCN	mouse/Swiss-Webster	NR/18-23	M/10/group	saline	8.5 (7.7-9.4)	3.4 (3.1-3.8)	LD_{50} increased by pretreatment with sodium nitrite and sodium thiosulfate.	May et al., 1966
s.c.	KCN	mouse/Swiss-Webster	NR/18-23	M/10/group	saline	12.0 (10.8-13.3)	4.8 (4.3-5.3)	LD_{50} increased by pretreatment with cobaltous chloride alone or in combination with sodium nitrite and/or sodium thiosulfate.	Isom and May, 1973

10/06/90

TABLE V-1 (cont.)

Route	Administered Compound	Species/Strain	Age/Weight (g)	Sex/No.	Vehicle	LD_{50} (mg/kg bw)	LD_{50} (mg CN^-/kg bw)	Comment	Reference
s.c.	KCN	mouse/Swiss-Webster	NR/18-23	M/10/ group	saline	11.8 (10.6-13.2)	4.7 (4.2-5.3)	LD_{50} increased by pretreatment with sodium nitrite and sodium thiosulfate, enhanced by O_2.	Way et al., 1972
s.c.	KCN	mouse/NR	NR/18-32	F/10/ group	saline	6.02 ± 0.33	2.41 ± 0.12	At 23-25°C	Streicher, 1951
						2.86 ± 0.16	1.14 ± 0.06	At 4°C	
s.c.	NaCN	dog/NR	NR/NR	NR/16	NR	5.36 ± 0.28	2.84 ± 0.15	LD_{50} increased by immediate post-treatment administration of sodium thiosulfate, amyl nitrite, sodium nitrite and combinations of antidotes.	Chen and Rose, 1952
i.m.	HCN	rabbit/NR	adult/NR	M/10-15/ group	NR	1.50 (1.27-1.80)	1.44 (1.22-1.73)	Increased rate of breathing, vociferation, uncoordinated movement of head, ataxia, tremors, respiratory and cardiac arrest.	Ballantyne et al., 1971, 1972
				F/10-15/ group		0.95 (0.81-1.11)	0.91 (0.78-1.07)		
i.m.	KCN	rabbit/NR	adult/NR	M/10-15/ group	NR	3.06 (2.61-3.63)	1.22 (1.04-1.45)	Same as above, the rabbits lived longer following KCN rather than HCN administration.	Ballantyne et al., 1971, 1972
				F/10-15/ group		3.27 (2.70-4.08)	1.31 (1.08-1.63)		

NR - Not reported

s.c. = subcutaneous; i.p. = intraperitoneal; i.m. = intramuscular

included in Table V-1) of HCN in Swiss-Webster mice for a 30-minute exposure and a 10-minute recovery period was determined to be 166 ppm (182.6 mg HCN/m³) (Matijak-Schaper and Alarie, 1982).

Table V-2 is a summary of studies of acute toxicity other than LD_{50} determinations. For the purposes of this review, durations ranging from a single exposure up to 14 daily doses are considered to be acute. Time to death depends upon dose and rapidity of absorption. Three dogs receiving oral doses of KCN of 3.8, 10.7 and 20.2 mg KCN/kg bw (1.5, 4.3 and 8.1 mg CN^-/kg bw, respectively) died in 155, 21 and 8 minutes, respectively (Gettler and Baine, 1938). Oral doses of 10 and 15 mg KCN/kg bw (4 and 6 mg CN^-/kg bw, respectively) were fatal to rats and mice, respectively; the incidence of mortality depended on the amount administered (Ferguson, 1962). Basu (1983) found that an oral dose of 8 mg KCN/kg bw (3.2 mg CN^-/kg bw) resulted in no signs of toxicity in five guinea pigs, and slight tremors with complete recovery in three guinea pigs. This dose is approximately equal to the LD_{50} in mice (see Table V-1). Kreutler et al. (1978) studied the effect of KCN administered in the diet of rats in relation to protein content and iodine deficiency. The rats tolerated a much higher dose of cyanide (80 mg CN^-/kg bw/day) when it was mixed in the diet than when administered by gavage. All of the rats on the low protein diet lost body weight, regardless of other treatments. KCN-treated rats on a low protein diet deficient in iodine had increased thyroid weight, increased thyroid-to-body weight ratio and increased levels of plasma thyroid stimulating hormone (TSH). Addition of iodine to the diet protected the rats from the effects on the thyroid. Rats on high protein diets gained

TABLE V-2

Acute Toxicity of Cyanides in Laboratory Animals

Route	Administered Compound	Species/ Strain	Age/ Weight (g)	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/Concentration as CN^-	Duration	Effects	Reference
Oral	KCN	dog/NR	NR/ 12.7 kg	NR/3	NR	3.8 mg/kg bw	1.5 mg/kg bw	single dose	death in 155 minutes	Gettler and Baine, 1938
			NR/ 11.3 kg			10.7 mg/kg bw	4.3 mg/kg bw	single dose	death in 21 minutes	
			NR/ 11.9 kg			20.2 mg/kg bw	8.1 mg/kg bw	single dose	death in 8 minutes	
Oral	KCN	rat/ Sprague-Dawley	adult/NR	NR/20/ group	water	10 mg/kg bw	4.0 mg/kg bw	single dose	The compound was administered in volumes corresponding to % bw. At 1.35%, 10/20 rats died; at 2.5%, 14/20 died; at 5%, 19/20 died.	Ferguson, 1962
Oral	KCN	mouse/ Swiss-Webster	NR/NR	NR/20/ group	water	15 mg/kg bw	6.0 mg/kg bw	single dose	When administered in a volume corresponding to 1.25% bw, 7/20 died; at 2.5%, 15/20 died; at 5%, 19/20 died.	Ferguson, 1962
Oral	KCN	guinea pig/ Duncan-Hartley	NR/ 200-250	NR/8	sucrose	8 mg/kg bw	3.2 mg/kg bw	single dose	5 of the guinea pigs had no signs of toxicity after 30 min., 3 had slight tremors but recovered completely in 15 minutes.	Basu, 1983
Oral	KCN	rat/ Holtzman	NR/ 90-100 g	M/24	2% casein diet	0	0	14 days	Mean loss of bw = 30%	Kreutler et al., 1978
				M/14	2% casein diet	2000 mg/kg diet[a] (200 mg/kg bw/day)	800 mg/kg diet[a] (80 mg/kg bw/day)	14 days	Mean loss of bw = 31%; significantly increased mean absolute thyroid weight (~2x); mean thyroid-to-bw ratio (~2.5x) and plasma TSH (27x) compared with controls ($p<0.05$).	

TABLE V-2 (cont.)

Route	Administered Compound	Species/ Strain	Age/ Weight (g)	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/Concentration as CN^-	Duration	Effects	Reference
Oral				M/14	2% casein diet + KI	2000 mg/kg diet (200 mg/kg bw/day)	800 mg/kg diet (80 mg/kg bw/day)	14 days	Mean loss of bw = 33%; mean absolute thyroid weight and mean plasma TSH were not significantly different from control. Thyroid-to-bw ratio was significantly decreased compared to control (~25% decrease), and compared with group without added KI to diet (~3.2x) ($p<0.05$).	Kreutler et al., 1978
				M/11	20% casein diet	0	0	14 days	Mean gain of bw = 37%. No significant change in absolute thyroid weight, thyroid-to-bw ratio or plasma TSH compared with 2% casein diet control.	
				M/10	20% casein diet	2000 mg/kg diet (200 mg/kg bw/day)	800 mg/kg diet (80 mg/kg bw/day)	14 days	Mean gain of bw = 41%. No significant change in absolute thyroid weight, thyroid-to-bw ratio, and plasma TSH compared with 20% casein diet control.	
Inhalation	HCN	dog/NR	NR/ T0.3 kg	NR/1	air	1.55 mg/kg bw[b]	1.49 mg/kg bw[c]	15 minutes	death	Gettler and Baine, 1938
			NR/ 9.1 kg	NR/1	air	1.11 mg/kg bw[b]	1.07 mg/kg bw[c]	10 minutes	death	
Inhalation	HCN	dog/ mongrels	adult/NR	NR/6	air	620 mg/m³	595 mg/m³	2.0 minutes	Died in ~16 hours, apneic	Haymaker et al., 1952
						590 mg/m³	566 mg/m³	2.0 minutes	Died in ~16 hours, apneic and comatose before death.	
						700 mg/m³	672 mg/m³	1.75 minutes	Died in 20 hours, apneic and comatose before death.	
						700 mg/m³	672 mg/m³	1.75 minutes	Prior to sacrifice at 24 hours, dog became comatose but recovered full consciousness.	

TABLE V-2 (cont.)

Route	Administered Compound	Species/ Strain	Age/ Weight (g)	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/Concentration as CN^-	Duration	Effects	Reference
Inhalation						165 mg/m³	158 mg/m³	10.0 minutes	Prior to sacrifice at 26 hours, dog became comatose but recovered; this dog received one ampule of amyl nitrate after gasing.	Haymaker et al., 1952
						690 mg/m³	662 mg/m³	2.0 minutes	Prior to sacrifice at 28 hours, dog became comatose but recovered. 4 of the dogs had convulsive seizures (individuals not specified). Histopathologic damage to gray matter of brain included necrosis of cerebral cortex, caudate nucleus and putamen, substantia nigra, globus pallidus, pulvinar of the thalamus and cerebellar cortex.	
Inhalation	HCN	rat/F344	NR/200-300 g	M/6/ group	air	176.5 mg/m³	169.9 mg/m³	30 minutes	All animals died either within the 30-minute exposure period or during the 24-hour post-exposure observation period.	Levin et al., 1987

TABLE V-2 (cont.)

Route	Administered Compound	Species/ Strain	Age/ Weight (g)	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/Concentration as CN^-	Duration	Effects	Reference
Inhalation	HCN	rat/Long-Evans	adult/ 350	M/4/ group	air	0, 220 mg/m³ (exposed 5 times)	0, 211 mg/m³ (exposed 5 times)	one 12.5-minute exposure; 5 exposures with 4-day intervals between exposures	Increased activity of cardiac specific creatine phosphokinase ~2-fold over control, was seen after each exposure but the increases at each exposure were not influenced by previous exposure; increased numbers of ectopic heart beats induced by norepinephrine ~8x above control; no consistent increase in incidence of histopathologic damage to myocardium. Pretreatment of rats with chlorpromazine-thiosulfate reduced the increase in plasma cardiac-specific creatine kinase activity and the number of ectopic beats induced by norepinephrine but increased the incidence of myocardial lesions.	O'Flaherty and Thomas, 1982
Inhalation	HCN	monkey/ cynomolgus	NR/3-4 kg	NR/7	air	110-172 mg/m³	106-165 mg/m³	30 minutes	Displayed hyperventilation, loss of consciousness, bradycardia with arrhythmias and T wave abnormalities followed by rapid recovery.	Purser et al., 1984

TABLE V-2 (cont.)

Route	Administered Compound	Species/ Strain	Age/ Weight (g)	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/Concentration as CN^-	Duration	Effects	Reference
Dermal	HCN	dog/NR	NR/NR	NR/11	air	6500-16,900 mg/m³	6240-16,224 mg/m³	47-180 minutes	The heads of the dogs were excluded from the exposure chamber. No deaths were observed at 6500-10,900 mg HCN/m³ in 30 or 60 minutes. Respiration decreased and was labored and irregluar, face and throat twitches observed, deaths followed. Lungs were bright pink and collapsed.	Walton and Witherspoon, 1926
Dermal	HCN	guinea pig/NR	NR/NR	NR/8/ group	none	not quantified	NA	~8 minutes	The guinea pigs were prevented from inhaling vapors. A tube filled with liquid HCN was pressed against the abdomen. Signs of toxicity: rapid respiration, muscle twitches and convulsions began at 5-7 minutes, death at 7-8 minutes.	Walton and Witherspoon, 1926

TABLE V-2 (cont.)

Route	Administered Compound	Species/ Strain	Age/ Weight (g)	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/Concentration as CN^-	Duration	Effects	Reference
i.p.	KCN	mouse/ Swiss-Webster	NR/NR	M/NR	water	0, 1, 2, 3, 4, 5 or 6 mg/kg bw	0, 0.4, 0.8, 1.2, 1.6, 2.0 or 2.4 mg/kg bw	single injection	Effects were rapid breathing, agitation, loss of coordination and convulsions. These signs were scarcely observed at 1 and 2 mg KCN/kg. At 3, 4 and 5 mg KCN/kg, there was gasping, irregular breathing and convulsions, beginning at 2-3 minutes. At 6 mg/kg, severe convulsions, cessation of breathing, death of 20% of the mice. At doses >6 mg KCN/kg, all died in 4 minutes. Changes corresponded to degree of inhibition of cytochrome oxidase.	Isom et al., 1982
s.c.	NaCN	rat/ Wistar	adult/ 200-220	M/5-6/ group	saline	3 x 0 and 3 x 0.04 mg/kg bw	0.043 x 0 and 3 x 0.02 mg/kg bw	3 injections 10 minutes apart	The treated rats survived 5 hours; there was a significant increase in acetylcholinesterase activity ($p<0.05$) in cerebral cortex (57%), hippocampus (28%) and midbrain (18%) compared with controls ($p<0.05$).	Owasoyo and Iramain, 1980
s.c.	KCN	mouse/ Webster	adult/ 25-30	M/4-9/ group	saline	10, 12 or 15 mg/kg bw	4.0, 5 or 6 mg/kg bw	single injection	Significantly increased whole brain calcium cyanide-induced tremors, severity and dose response paralleled increase brain calcium levels.	Johnson et al., 1986

TABLE V-2 (cont.)

Route	Administered Compound	Species/ Strain	Age/ Weight (g)	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/Concentration as CN^-	Duration	Effects	Reference
i.v.	NaCN	dog/ mongrels	NR/15 kg	NR/NR	saline	1 mg/kg bw	0.5 mg/kg bw	single injection	EEG: lower brain electrical activity for 6 minutes, then depressed wave amplitude. EKG: sinus pause, bradycardia (slow heart rate), elevated T or biphasic T wave. Initial decrease followed by marked increase in systolic, diastolic and venous blood pressure for 10 minutes, returned to pre-NaCN level in 20-30 minutes. There was hyper- followed by hypoventilation.	Burrows et al., 1973

[a]Confusion in reference in which KCN was the compound administered as described in text, and HCN was the compound referred to in table.

[b]Dose in mg/kg bw was experimentally determined.

[c]Dose in mg/kg/day was calculated from dietary concentration on the assumption that a young growing rat consumes a quantity of food equal to 10% of its bw/day.

NR = Not reported

TSH = Thyroid stimulating hormone

KI = Potassium iodide

body weight regardless of treatment. Although these diets were not supplemented by iodine, no effects on the thyroid were observed from cyanide exposure. Ingestion of cyanogen glycoside-containing foods, along with nutritional deficiencies, is suspected of having a goitrogenic effect in humans (Ermans et al., 1972).

Effects of acute inhalation exposure of dogs and rats to HCN are also summarized in Table V-2. A dog receiving a bolus of 3.8 mg KCN/kg bw (1.5 mg CN^-/kg bw) by gavage died in 155 minutes, while a dog inhaling a total of 1.55 mg/kg bw (1.5 mg CN^-/kg bw) died in 15 minutes (Gettler and Baine, 1938). Correlation of duration of exposure and concentration with lethal effect is difficult because of the wide range of sensitivity among species and the small number of animals exposed/group. This phenomenon was demonstrated in dogs by Haymaker et al. (1952) (see Table V-2). Histopathologic lesions to the gray matter of the brain were observed when the dogs were necropsied. Moss et al. (1951) reported a minimum lethal concentration of 55 mg HCN/m³ for $\leq$20 minutes in female rats. Rats tolerated repeated acute inhalation exposures (one 12.5-minute exposure every 4 days) to HCN of 220 mg/m³ (O'Flaherty and Thomas, 1982). During the experimental period, the rats had increased activities of cardiac specific creatine phosphokinase (an index of myocardial damage) and increased numbers of ectopic heart beats in response to norepinephrine. The rats were killed and necropsied within 2 weeks of the last exposure. No consistent increase in incidence of lesions to the myocardium was observed.

Barcroft (1931) compared the sensitivities of a variety of species to HCN. Animals were exposed by inhalation to 1000 mg/m³ HCN and the time at

which death occurred was recorded. The results are presented in Table V-3. The species were also compared with respect to the highest concentration that the animals could breathe "indefinitely." It was not clear as to how this determination was made, but the reported concentrations ranged from 100-400 mg HCN/m³. The order of sensitivities to these nonlethal concentrations, from the most sensitive to the least sensitive species was dog = rat > mouse > rabbit = monkey = cat > goat > guinea pig.

Toxic effects of cyanides administered by other routes are described in Table V-2. Dermal absorption of HCN gas resulted in death in dogs and guinea pigs (Walton and Witherspoon, 1926). A dose-response for toxic effects of KCN administered intraperitoneally to mice was defined by Isom et al. (1982). Such signs as rapid breathing, agitation, loss of coordination and convulsions were rare at doses of 1 and 2 mg KCN/kg bw (0.4 and 0.8 mg CN^-/kg bw, respectively). At 3, 4 and 5 mg KCN/kg bw (1.2, 1.6 and 2.0 mg CN^-/kg bw, respectively), these effects were observed within 2-3 minutes. At 6 mg KCN/kg bw (2.4 mg CN^-/kg bw), there was 20% mortality in 4 minutes. Subcutaneous administration of NaCN to rats resulted in significantly increased acetylcholinesterase activity in the cerebral cortex, hippocampus and midbrain compared with controls (Owasoyo and Iramain, 1980). Intravenous administration of 1 mg NaCN/kg bw (0.5 mg CN^-/kg bw) to dogs resulted in EEG, EKG and blood pressure changes (Burrows et al., 1973).

Subchronic Exposure. A summary of the effects following subchronic exposure of animals to cyanides was provided in Table V-4. The intermittent ingestion of low doses over a day appears to allow for sufficient detoxification to account for the sublethal effects. Rats tolerated 25

TABLE V-3

Sensitivity of Various Species to Inhalation Exposures of HCN*

Species (number/species not given)	Lethal time of exposure to a Concentration of 1000 mg HCN/m³ (minutes)
Dog	0.8
Mouse	1.0
Cat	1.0
Rabbit	1.0
Rat	2.0
Guinea pig	2.0
Goat	3.0
Monkey	3.5

*Source: Barcroft, 1931

TABLE V 4

Effects of Subchronic Exposure of Laboratory Animals to Cyanide

Route	Administered Compound	Species/ Strain	Age/ Weight	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/concentration as CN^-	Duration	Effects	Reference
Oral	KCN	rat/Sprague-Dawley	NR/70 g	M/7/ group	drinking water	0 and 200 mg/L[a] (30 mg/kg bw/day)	0 and 80 mg/mL[a] (12 mg/kg bw/day)	21 days	Increased liver weight of treated animals (15.8 g) compared with controls (13.5 g), $p<0.05$. No effect on bw gain.	Palmer and Olson, 1979
Oral	KCN	rat/Sprague-Dawley	NR/70 g	M/7/ group	diet	0 and 200 mg/kg[b] diet (20 mg/kg bw/day)	0 and 80 mg/kg[b] diet (8 mg/kg bw/day)	21 days	No increase in liver weight of treated animals (13.7 g) compared with controls (13.5 g). No effect on bw gain.	Palmer and Olson, 1979
Oral	KCN	rat/Sherman	adult/NR	M/NR	diet	0, 10 mg/kg bw/day	0, 4 mg/kg bw/day	25 days	No mortality; this dose is equal to the 1-day oral LD_{50} of KCN in the same strain of rats; however, the rats tolerated the dose when mixed in feed.	Hayes, 1967
Oral	KCN	rat/Sherman	adult/NR	M/NR	diet	0-250 mg/kg bw/day	1-100 mg/kg bw/day	90 days	No mortality. The rats tolerated the doses, which were up to 25x the acute oral LD_{50} when mixed in the diet.	Hayes, 1967
Oral	KCN	rat/Sprague-Dawley	weanling/ NR	M+F/8/ group	diet	0 and 1875 mg/kg diet	0 and 750 mg/kg diet	56 days	No significant difference in food consumption, bw gain, food efficiency, protein efficiency ratio and liver and kidney to bw ratio.	Iewe and Maner, 1982
Oral	KCN	rat/ Chiceiomys Waterhouse (African giant rat)	weanling/ ~90 g	M/6/ group /2/	diet	0 and 2500[c] mg/kg diet (90 mg/kg bw/day)	0 and 1000[c] mg/kg diet (36 mg/kg bw/day)	84 days	Slightly reduced feed consumption and bw gain of treated animals compared with controls. No histopathologic changes in thyroid, kidney, liver or spleen.	Iewe, 1982

TABLE V-4 (cont.)

Route	Administered Compound	Species/ Strain	Age/ Weight	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/concentration as CN^-	Duration	Effects	Reference
Oral	KCN	pig/ Yorkshire	weanling/ NR	M + f	diet	0 and 1250[c] mg/kg diet	0 and 500[c] mg/kg diet	56 days	A small but significant reduction in average food consumption ($p<0.05$), but no significant difference in bw gain between treated and control pigs. No significant difference in food efficiency, protein efficiency ratio, organ to bw ratios of thyroid, liver, kidney, spleen or heart. No histopathologic changes.	Tewe and Maner, 1980
Oral	NaCN	dog/beagle	6.5 months/ Mean-7.1 kg at start and mean 8.1 kg at end of test	1M, 1F control, 3M, 2F treated	diet	0 and 2500[d] mg/kg diet (6 mg/kg bw/day)	0 and 1000[d] mg/kg diet (3 mg/kg bw/day)	30-32 days	No effect on food consumption or bw gain, no effect on hemoglobin concentration, hematocrit or differential white cell count. Complete autopsies and comprehensive histologic examinations revealed no effect on organ to bw ratios and no treatment-related gross or histopathologic lesions.	American Cyanamid Co., 1959
Oral	NaCN	dog/NR	NR/NR	NR/1 control, 3 treated	capsule	0, 0.5, 1.0 up to 2 x 2.0 mg/kg bw/day (6 mg/kg bw day)	0, 0.27, 0.53 up to 2 x 1.1 mg/kg bw/day (3 mg/kg bw/day)	15 months	At >1.0 mg NaCN/kg bw, acute signs of toxicity were observed immediately after dosing, but recovery occurred in <0.5 hours. No pathologic changes in organs or metabolic systems, but erythrocyte count was increased and serum albumin decreased. Degenerative changes in ganglion cells of CNS and especially Purkinje cells of cerebellum (necrosis, reduced RNA content, inflammation) in all treated dogs.	Herlting et al., 1960

TABLE V-4 (cont.)

Route	Administered Compound	Species/ Strain	Age/ Weight	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/concentration as CN^-	Duration	Effects	Reference
Inhalation	HCN	rabbit/ Danish	NR/2.0-2.5 kg	M/22/ group	air	0, 0.55 mg/m³	0, 0.53 mg/m³	28 days	No significant difference between control and treated animals with respect to ultrastructural changes in myocardium.	Hugod, 1981
s.c.	NaCN	rat/Charles River	adult/ 200-350 g	M/221	saline	0 and TWA[e] doses of 0.20, 1.16, 2.48 and 3.24 mg/kg bw/day	0 and TWA[e] doses of 0.11, 0.61, 1.31 and 1.72 mg/kg/day	3 days a week for 3 months (90 days)	There was high mortality in ≥0.61 mg/kg treated groups: 159/208 treated rats died before the end of the experiment; 107 died in the first 3 weeks. In 15 minutes after each injection most rats became comatose, but recovered. Many appeared blind and ataxic before death and had necrotic lesions of corpus callosum and optic nerve on autopsy. Severely affected rats had focal constriction of optic nerve. A no-effect level was 0.20 mg/kg bw/day TWA dose of NaCN (0.11 mg CN^-/kg bw/day) for 90 days.	Lessell, 1971
s.c.	KCN	rat/Wistar	adult/NR	NR/3/ group	NR	0 and 0.5 mg/rat (1.43 mg/kg bw/[f] injection)	0.2 mg/rat (0.57 mg/kg bw/[f] injection)	once weekly for 22 weeks	Treated rats had degenerating cells in cortex, degenerating Purkinje cells in cerebellum and hippocampus; degenerating neurons in brain; pallor in myelin of corpus callosum; decreased vitamin B_{12} levels in liver.	Smith et al., 1963

TABLE V 4 (cont.)

Route	Administered Compound	Species/ Strain	Age/ Weight	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/concentration as CN^-	Duration	Effects	Reference
s.c.	NaCN	baboon/ *Papio cynocephalus* and *P. anubis*	9 months/ 3.7 kg	M/7 M/10	phosphate buffer and adrenalin	0 1 mg/kg bw, 5 days/week	0 0.4 mg/kg bw, 5 days/week	42 months	Rise in hemoglobin ($p<0.01$), low mean erythrocyte volume ($p<0.01$), decreased mean erythrocyte hemoglobin content ($p<0.01$) from pre-test determination.	Crampton et al., 1979
i.m.	KCN	monkey/ rhesus	NR/2.5-5.0 kg	NR/25	water	TWA dose[e] 1.9-38 mg/kg/day	TWA 0.76-15.2[e] mg/kg bw/day	17-103 days	All treatments were fatal in <2 hours to 13 days after the last dose; 18 monkeys had lesions in white matter (cell proliferation, demyelination necrosis), 13 had lesions of gray matter (minor damage and necrosis). It was not possible to correlate dose with effect.	Hurst, 1940

[a]The dose in mg/kg bw/day was calculated from the concentration in drinking water, assuming that a young rat consumes an amount of water equal to ~15% of its bw per day.

[b]The dose in mg/kg bw/day was calculated from the concentration in food, assuming that a young growing rat consumes an amount of food equal to ~10% of its bw per day.

[c]The dose in mg/kg bw/day was calculated from food consumption and body weight information supplied in the report.

[d]The doses were calculated from data supplied in the method of meal preparation on the quantity of NaCN (45 mg) mixed with each meal. The dogs consumed all the meals, and the average weight of the dogs during the test was 7.6 kg. The concentration of NaCN in the meal was 500 mg/kg meal.

[e]TWA (time-weighted average) doses were calculated. In the study, the doses were raised throughout the experiment; the investigator supplied cumulative doses, and these cumulative doses were divided by 90 days to give the TWA doses.

[f]The dose in mg/kg bw was calculated by assuming the the average weight of a rat is 0.35 kg.

daily doses of KCN, each of which were equal to the acute oral LD_{50} (10 mg KCN/kg bw or 4 mg CN^-/kg bw), when the chemical was mixed in the diet (Hayes, 1967). In a second experiment, 90 daily doses that were equal to 25 times the LD_{50} did not result in mortality when administered in the diet (Hayes, 1967). Rats appeared to tolerate a higher oral dose of KCN (~30 mg KCN/kg bw/day or 12 mg CN^-/kg bw/day for 21 days) that was administered in drinking water (Palmer and Olson, 1979), than when KCN was administered in a bolus by gavage with water as the vehicle (Hayes, 1967; Gaines, 1969). The rats that had received KCN in the drinking water had significantly increased liver weights when compared with controls (Palmer and Olson, 1979). Rats receiving KCN in the diet for 21 days (20 mg KCN/kg bw/day or 8 mg CN^-/kg bw/day) did not have increased liver weights, but the doses were lower than when KCN was added to the drinking water. Other parameters were not examined (Palmer and Olson, 1979).

Tewe (1982) studied the effects of dietary KCN, administered for 84 days, in the African giant rat (8/group). The concentration in the diet (2500 mg KCN/kg diet) was approximately equivalent to 90 mg KCN/kg bw/day (36 mg CN^-/kg bw/day). The effects in the African giant rat were slightly reduced food consumption, slightly reduced body weight gain and an increase in serum urea. The differences from control were significant only for serum urea. The brain was not examined histologically, although it is a target organ for subcutaneously administered cyanide in rats (Smith et al., 1963; Lessell, 1971).

Tewe and Maner (1980, 1982) conducted studies in Yorkshire pigs and Sprague-Dawley rats to determine the interrelationships of cyanide, dietary

protein and iodine on the physiologic performance. Only those portions of the reports where pigs and rats were maintained on standard diets with or without KCN are discussed here. There were no significant differences between treated rats (750 ppm CN^- in diet for 56 days) and controls when compared for food consumption, body weight gain, food efficiency (food intake/body weight gain), protein efficiency ratio (body weight gain/protein intake) and liver and kidney weight-to-body weight ratios (Tewe and Maner, 1982). Histologic examinations were not performed. Treated pigs (500 ppm CN^-) had slightly reduced food consumption; however, body weight gain was not affected (Tewe and Maner, 1980). Histologic examination of the pigs revealed no pathologic changes in thyroid, hypophysis, stomach, liver, cardiovascular tissue, spleen, tonsils, thymus, intestinal mesentery, kidney, eye, brain, spinal cord, neural ganglia or bone. Organ-to-body weight ratios of thyroid, liver, kidney, spleen and heart were not different from controls.

The American Cyanamid Co. (1959) conducted a study in beagles on the effects of NaCN mixed with their food. The dogs consumed 3 mg CN^-/kg bw/day for 30-32 days. There were no clinical signs of toxicity or effects on food consumption, body weight gain or hematologic parameters. Histologic examination was extensive and included tongue, tonsils, cervical lymph node, salivary gland, thyroid, trachea, myocardium, aorta, lung, thymus, liver, gall bladder, pancreas, esophagus, stomach, duodenum, jejunum, ileum, colon, cecum, rectum, mesenteric lymph node, spleen, adrenal, kidney, ureter, urinary bladder, skeletal muscle, cerebellum, cerebrum, spinal cord, pituitary, testis, prostate, ovary, uterus and vagina. No treatment-related abnormalities were found. There were no differences in organ-to-body weight

ratios of submaxillary glands, thyroid, heart, lung, liver, adrenal, pancreas, spleen, kidney, brain, pituitary, testis or ovary. The dose had been chosen on the basis of the concentration of NaCN in the food that the dogs would accept and eat completely. Dogs receiving doses of >1.0 mg NaCN/kg bw/day (0.5 mg CN^-/kg bw/day) for up to 15 months had signs of intoxication immediately after each dosing, but they recovered in <0.5 hour (Hertting et al., 1960). The NaCN was administered in a capsule. Degenerative changes in ganglion cells of the CNS were observed in all three treated dogs. The lowest exposure was 0.27 mg CN^-/kg bw/day.

Effects of subchronic exposure to cyanide by other routes are also included in Table V-4. Rabbits exposed to 0.55 mg HCN/m^3 air for 28 days had no treatment-related changes in the ultrastructure of the myocardium (Hugod, 1981). Rats treated subcutaneously 3 days/week for 90 days with NaCN in TWA (time-weighted average) doses of 0.61, 1.31 and 1.72 mg CN^-/kg bw/day had high rates of mortality and histopathologic damage to the brain (Lessell, 1971). The damage consisted of necrotic lesions in the corpus callosum and the optic nerve. A no-effect level was a TWA dose of 0.11 mg CN^-/kg bw/day for 90 days. Smith et al. (1963) also found histopathologic damage to the brains of rats treated once weekly for 22 weeks with subcutaneous injections of 0.57 mg CN^-/kg bw. Hurst (1940) found necrotic lesions in white and gray matter of the brains of rhesus monkeys treated with lethal intramuscular TWA doses of KCN (0.76-15.2 mg CN^-/kg bw/day) for 17-103 days. Baboons treated subcutaneously for 42 months with 1 mg NaCN/kg bw (0.4 mg CN^-/kg bw) 5 days/week, had increased hemoglobin and decreased MCU and MCHC values compared with controls (Crampton et al., 1979). There was no effect on body weight gain, and the treated animals

appeared to be in good health throughout the experiment. Animal studies have shown that repeated injections of cyanide can cause CNS damage, particularly to the white matter (Levine, 1967).

Chronic Exposure. Only two studies were found that provided information on the effects of long-term exposure of animals to cyanide. Details of the studies are presented in Table V-5.

In the study of Howard and Hanzal (1955), groups of 10 male and 10 female Carworth Farms rats were maintained for 104 weeks on diets that had been fumigated with HCN. It was necessary to measure the loss of HCN due to evaporation from the chow and to prepare fresh rations every other day to nominal concentrations of 100 and 300 mg HCN/kg diet. Results of the analysis of residues over the 2-year duration of the study indicated an average drop in the dietary concentration of HCN over 2 days from 100 mg/kg diet to 51.9 mg/kg diet (48% loss) for the low-dose level and from 300 mg/kg diet to 80.1 mg/kg diet (73% loss) for the high-dose level. Thus, the average concentrations are 76 mg/kg diet for the low-dose level and 190 mg/kg diet for the high-dose level.

During the 2 years of this study, the growth curves for all groups did not vary as a result of treatment. Food consumption, hematologic values and survival were also similar in all groups. At termination of the study, the only pathologic lesions observed were those associated with aging. The only effects of treatment were increased CN^- level in the red blood cells, and increased thiocyanate levels in the plasma, red blood cells, liver and kidneys of animals from both treatment groups.

TABLE V-5

Effects of Chronic Exposure of Laboratory Animals to Cyanide

Administered Compound	Species/ Strain	Age/ Weight	Sex/No.	Vehicle	Dose/Concentration of Compound	Dose/Concentration as CN^-	Duration	Effects	Reference
HCN	rat/ Carworth Farms	weanling/ ~57 g	M/10/ group	diet	0, 76 and 190[a] mg/kg diet (0, 3.3 and 8.1 mg/kg bw/day)	0, 73 and 183[a] mg/kg diet (0, 3.6 and 7.5 mg/kg bw/day)	104 weeks	No effect on growth rate, no signs of toxicity, no histopathologic changes in heart, lungs, liver, spleen, stomach, small and large intestines, kidney, adrenals, thyroid, testes, uterus, ovary and the cerebrum and cerebellum of the brain.	Howard and Hanzal, 1955
			F/10/ group		(0, 4.5 and 10.8 mg/kg bw/day)	(0, 4.6 and 10.8 mg/kg bw/day)			
KCN	rat/NR	weanling/ 43 g	M/10/ group	diet	0, 1500 mg/kg[b] diet (75 mg/kg bw/day)	0, 600 mg/kg[b] diet 30 mg/kg bw/day)	11.5 months	No deaths or clinical signs. Significantly (no p values) reduced bw gain, increased excretion of thiocyanate at 4 months and at at 11 months, decreased plasma thyroxine levels, decreased secretion rates at 4 months. No effect on plasma thyroxine at 11 months, no definitive lesion to sciatic, thyroid, optic or CNS tissue. Vacuoles and accumulation of fluid in spinal cord white matter with moderate primary myelin degeneration were observed in treated rats. These changes were more pronounced in KCN-treated rats that were maintained on methionine and vitamin B_{12} deficient diets.	Philbrick et al., 1979

[a]The doses in mg/kg bw/day were calculated using data on food consumption and bw supplied in the report.

[b]The dose in mg/kg bw/day was calculated assuming that a rat consumes an amount of feed equal to 5% of its body weight/day.

NR = Not reported

Philbrick et al. (1979) mixed KCN in the diet at a concentration of 1500 mg/kg diet and maintained groups of 10 male weanling rats (strain not specified) on this diet for 11.5 months. When expressed in terms of cyanide, this dietary concentration is higher than the concentration of cyanide in the food of the rats in the Howard and Hanzal (1955) study. Positive controls received a diet containing 10% casein supplemented with 0.3% DL-methionine, KI and vitamin B_{12}. Two groups received this diet plus either 1500 ppm KCN or 2240 KSCN. Negative controls received the 10% casein diet supplemented with methionamine but with no added KI or vitamin B_{12}. Two additional groups received this negative control diet plus either 1500 ppm KCN or 2240 ppm KSCN. Thus, of those two groups receiving KCN, one received a complete diet whereas the other was restricted in iodine and vitamin B_{12}. No deaths occurred nor were there any gross signs of toxicity. Effects were reduced body weight gain, decreased plasma thyroxine levels and decreased rate of thyroxine secretion. The effects appeared greater in the animals on the deficient diet, although the difference was not always significant. There were no definitive histopathologic lesions in the optic or CNS tissue, thyroid or sciatic tissue; however, vacuolization and myelin degeneration were observed in spinal cord sections.

Target Organ Toxicity

Central Nervous System. Several studies discussed in the General Toxicity Section described histopathologic damage to brain tissues. Haymaker et al. (1952) found lesions in the gray matter of brains of dogs 28 hours after fatal inhalation exposure to 690 mg/m³ HCN for 2 minutes. The damage included necrosis of the cerebral cortex, caudate nucleus and putamen, substantia nigra, globus pallidus, pulvinar of the thalamus and

cerebellum cortex. Dogs given subchronic oral doses of ≥1.0 mg/kg/day NaCN showed degenerative changes in ganglion cells, especially Purkinje cells of the cerebellum (Hertting et al., 1960). EEG changes such as low electrical activity and depressed wave amplitude were observed in dogs given a single dose of 1 mg NaCN/kg bw intravenously (Burrows et al., 1973).

Smith et al. (1963) found that subcutaneous administration of 1.43 mg KCN/kg bw to rats once weekly for 22 weeks resulted in degeneration of pyramidal cells in the cerebral cortex, degeneration of the cerebellum (Purkinje cells) and hippocampus, degenerating neurons in the brain and pallor of myelin in the corpus callosum. Lessell (1971) described necrotic lesions in the corpus callosum and optic nerves of rats treated subcutaneously with ≥1.16 mg NaCN/kg/day. Focal constriction of the optic nerve was observed in severely affected rats. In a subsequent study, in which the doses of NaCN were adjusted in order to keep rats in a coma for 225-260 minutes, there were severe ultrastructural pathologic changes in the optic nerve, particularly in the retrobulbar zone (Lessell and Kuwabara, 1974). The optic nerve lesions included edema, enlarged astrocytes, clumping of myelin, nerve fiber degeneration and necrotic and swollen axons that had few microtubules. Owasoyo and Iramain (1980) found increased activity of acetylcholinesterase in the cerebral cortex, hippocampus and midbrain of rats given subcutaneous doses of NaCN. Ferraro (1933) studied the CNS histopathology of cats and monkeys given increasing doses of KCN. The damage included demyelination in white matter of the frontal, occipital, parietal and temporal lobes, the corpus callosum, the cerebellum, the spinal cord and optic nerve. Degeneration of axis cylinder, gliosis, hypertrophy and vacuolization of oligodendroglia were also observed. Philbrick et al.

(1979) reported vacuolization and myelin degeneration in the white matter of the spinal cord in rats fed diets containing 75 mg KCN/kg bw (30 mg CN/kg bw).

Heart. O'Flaherty and Thomas (1982) found that inhalation of HCN by rats resulted in increased activity of plasma cardiac specific creatine phosphokinase, an indicator of myocardial damage, although there was no consistent increase in the incidence of lesions to the myocardium. Hugod (1981) also failed to detect changes in the ultrastructure of the myocardium of rabbits after HCN inhalation. Purser et al. (1984) exposed cynomolgus monkeys to concentrations of HCN ranging from 100-172 ppm and observed bradycardia with arrhythmias and T-wave abnormalities, followed by rapid recovery. EKG changes observed following intravenous injections of NaCN in dogs included sinus pause, bradycardia and elevated or biphasic T waves (Burrows et al., 1973). There was also a marked increase in systolic, diastolic and venous blood pressures.

Thyroid. Kreutler et al. (1978) administered KCN orally (2000 ppm in the diet) for 14 days. Decreases in thyroid weights and plasma levels of TSH in rats were inhibited by addition of iodine to the diet, or by providing adequate levels of protein in the diet. However, Philbrick et al. (1979) reported decreased thyroxine levels and decreased rates of thyroxine secretion in rats treated chronically with standard diets containing KCN.

Other Effects

Carcinogenicity. Pertinent data regarding the carcinogenicity of HCN, KCN and NaCN were not located in the available literature; HCN and KCN are

not scheduled for testing by the National Toxicology Program (NTP, 1991). Short-term toxicity studies for NaCN have been completed and are currently undergoing review (NTP, 1991). Perry (1935) found that inhalation exposure of rats to hydrogen cyanide retarded the growth of implanted Jensen rat sarcomas, but effective concentrations were very near the lethal concentrations. Bown et al. (1960) found that intraperitoneal injections of cyanide prolonged the lives of mice with transplanted Ehrlich ascites tumors and Sarcoma 180. Human patients with advanced cancers in the pelvic region tolerated pelvic perfusions of cyanide without evidence of toxicity. Other studies onthe anti-tumor activity of cyanide and the controversial cyanogenic glycoside, amygdalin (laetrile), which releases cyandie on enzymatic hydrolysis, have been reviewed elsewhere (Towill et al., 1978).

Mutagenicity. Most of the assays of cyanides for mutagenicity and effects on DNA synthesis have been negaitve. De Flora (1981) and DeFlora et al. (1984) found that KCN was negative for reverse mutation in *Salmonella typhimurium* strains TA1535, TA1537, TA1538, TA97, TA98, TA100 and TA102, using both the spot test and plate incorporation application technique, up to a dose of $3x10^{-3}$ nmoles/plate. This was the highest dose tested due to bacterial toxicity. Addition of the S-9 mix, prepared from livers of Aroclor-induced rats, had no effect on the response. Kushi et al. (1983) reported that HCN gas was marginally mutagenic to *S. typhimurium* strain TA100 in doses $\leq$5 mg/plate. Addition of S-9 mix decreased the mutagenic response. Negative results were obtained in strain TA98. DeFlora et al. (1984) also found that KCN was negative for inducing DNA damage in repair-deficient *Escherichia coli* strains WP67, CM871 and WP2. Karube et al. (1981) tested cyanide, a known inhibitor of cellular respiration (see

Chapter VII), by a method employing the rec assay in Bacillus subtilis strains M45 Rec^- and H17 Rec^+. The technique involved the use of two microbial electrodes consisting of immobilized bacteria on oxygen electrodes to measure the preferential death of the Rec^- strain in response to decreased respiration by the DNA-damaged bacteria. If the test chemical damaged the DNA, then the Rec^+ strain could repair the damage, and the rate of current increase at the Rec^+ electrode would be slower than at the electrode of the repair-deficient Rec^- strain. Cyanide increased the rate of current at both electrodes equally (dose-related), since the inhibition of cytochrome oxidase, rather than DNA damage, was responsible for the decrease in cellular respiration (see Chapter VII).

Teratogenicity and Other Reproductive Effects. Doherty et al. (1982) studied the teratogenicity of NaCN in pregnant Golden Syrian hamsters. On day 6 of gestation, osmotic minipumps were implanted subcutaneously at the back of the necks of 5-7 animals/group. The pumps delivered doses of NaCN of 0, 0.126, 0.1275 or 0.1295 mmoles/kg bw/hour (0, 148.2, 149.9 or 152.3 mg NaCN/kg bw/day or 0, 78.5, 79.4 or 80.7 mg CN^-/kg bw/day, respectively) from days 6-9 of gestation, at which time the pumps were removed. The hamsters were killed on day 11 of gestation. The osmotic minipump method of administration was chosen in order to approximate the slow release of cyanide from nitriles during in vivo metabolism. It was believed that the teratogenic effect of nitriles and cyanogenic glycosides, e.g., laetrile (Willhite et al., 1981; Willhite, 1982, 1983) was due to in vivo release of cyanide. The results are presented in Table V-6. Severe teratogenic effects were observed at all three doses. In addition, mild maternal toxicity was observed at 148.2 and 149.9 mg/kg/day, and more severe effects were seen at 152.3 mg/kg day. Toxic signs included shortness of breath,

TABLE V-6

Fetotoxic and Teratogenic Effects of NaCN in Hamsters[a]

	Treatment[b]			
		NaCN (mg/kg bw/day)[c]		
	Distilled Water (1.7 µℓ/hour)	148.2	149.9	152.3
Litters	6	5	6	6
Maternal bw change g ± SD	4 ± 5	-16 ± 9	-13 ± 10	-16 ± 12
Litters completely absorbed	0	2	3	5
Total resorptions/ total implantations (%)	8/83 (10)	44/70 (63)[d]	60/83 (72)[d]	72/87 (83)[d]
Number litters[e] affected (%)	3 (50)	2 (67)	3 (100)	1 (100)
Total malformed/ total number live fetuses (%)	4/75 (5)	16/26 (62)[d]	10/23 (43)[d]	1/15 (7)[f]

TABLE V-6 (cont.)

	Treatment[b]			
	Distilled Water (1.7 µℓ/hour)	NaCN (mg/kg bw/day)[c]		
		148.2	149.9	152.3
Abnormalities in live fetuses				
neural tube[g]	0	16	6	1
heart	0	0	4	0
limb or tail	3	0	4	0
other[h]	1	0	1	0
Crown-rump length, mm ± SD	8.19 ± 0.54	3.33 ± 0.29[d]	6.61 ± 1.64[d]	NR

[a]Source: Doherty et al., 1982

[b]The NaCN was delivered by osmotic minipumps implanted subcutaneously.

[c]The doses correspond to rates of 0.126, 0.1275 and 0.1295 mmole/kg bw/hour.

[d]Significantly different from control ($p<0.05$).

[e]Litters affected of those litters with one or more live fetuses.

[f]Not included in statistical analysis, too few fetuses.

[g]Includes exencephaly, encephalocele and nonclosure.

[h]Includes microphthalmia and one fetus with a small caudal half.

NR = Not reported

incoordination, reduced body temperature and loss of body weight. Preliminary range-finding experiments revealed that a dose of 0.126 mmole/kg/hour (148.0 mg/kg/day) resulted in no fetal abnormalities, while a dose of 0.133 mmole/kg/hour (156.4 mg/kg/day) resulted in 100% resorptions and death to some of the dams. In experiments with pair-fed controls, it was shown that the loss of body weight, in itself, was not related to the fetal abnormalities. No significant relationship between maternal toxicity and the incidence of fetal malformations was found by analysis of variance of the transformed data ($p>0.05$). When thiosulfate was co-administered with cyanide, no teratogenic effect was found. The investigators stated that the results do not imply a dose-response relationship because the variability in certified pump rates of individual minipumps is larger than the percentage difference among doses. Furthermore, the pumping rates cannot be verified *in vivo*.

Tewe and Maner (1981a) studied the effect of KCN added at a concentration of 500 ppm as the cyanide ion (500 mg CN^-/kg diet or 1250 mg KCN/kg diet) to a basal diet containing low-HCN cassava (21 mg HCN/kg) meal on the reproductive performance of female Wistar rats. Control rats were fed the basal diet (which contained 12 mg HCN/kg diet) in quantities that were equal to the diet consumed by treated rats. The KCN-treated rats were fed the test diet from 19.7±1.8 days before pregnancy, through lactation and during the post-weaning period. The controls were fed the control diet from 16.3±1.1 days before pregnancy and through post-weaning. Male rats did not receive the diets. Prior to parturition, the pregnant rats were separated, so that half the treated dams would remain on the test diet, the other half on the basal diet. Controls were separated similarly. At weaning, two rats

from each litter were selected and assigned at random to control and treatment groups for a 28-day post-weaning period. No significant differences were found among the various treated and control groups with respect to weight gain during gestation, litter size, birth weight of pups, feed consumption and body weight change during lactation, weights of maternal liver and kidney, weanling weights or mortality of offspring. The offspring that were continued on the diets during the post-weaning period consumed significantly less food and grew at a significantly slower rate than the basal diet offspring, regardless of previous cyanide exposure (in utero and/or in milk and/or in diet). The protein efficiency ratio of rats exposed in utero and fed cyanide during the post-weaning phase was significantly reduced compared with basal diet rats. Assuming that a weanling rat weighs approximately 45 g and consumes food equivalent to 10% of its body weight, then the rats fed diets with added KCN consumed approximately 50 mg CN^-/kg/day.

In a similar study, Tewe and Maner (1981b) studied the effect of cyanide on the reproductive performance of pregnant Yorkshire pigs. Cassava (containing a low level of HCN) was supplemented with protein, minerals and vitamin mix to which was added KCN. The groups, consisting of six pigs each, received the basal diet of low cyanide (30.3 mg CN^-/kg diet) or the basal diet plus cyanide, which provided total CN^- levels of 276.6 and 520.7 mg CN^-/kg diet. The concentrations as mg CN^-/kg diet were calculated by the investigators. The diets were started on the day after breeding and continued until parturition. Two pregnant pigs/group were killed on day 110 of gestation; the remaining pigs were fed their respective diets until parturition and then fed standard diets (no cyanide) during the

56 days of lactation. The piglets were not fed any cyanide. No differences were found with respect to litter size, litter size at weaning, birth weight of piglets, daily feed intake of sows or piglets. In the pigs that were killed on day 110 of gestation, no significant differences were found among groups for body weight, body weight gain during gestation, organ-to-body weight ratios of thyroid, spleen, liver, kidney and heart, number of fetuses/litter, weight of fetuses, weight of fetal liver or kidney. The thyroid-to-body weight ratios of fetuses from sows receiving 276.6 mg CN^-/kg diet were significantly reduced ($p<0.05$) compared with the low-cyanide group (30.3 mg CN^-/kg diet), but the high-cyanide (520.7 mg CN^-/kg diet) group was not different from the low-cyanide group. Fetal spleen-to-body weight ratio was significantly reduced ($p<0.05$) in the high-cyanide group compared with both other groups. The fetal heart-to-body weight ratio of the high-cyanide group was significantly reduced ($p<0.05$) compared with the low-cyanide group. Histopathologic examination of two killed pigs/group on day 110 of gestation indicated hyperplasia of kidney glomerular cells in one sow from each of the low- and medium-cyanide groups and in both sows of the high-cyanide group. Both high-cyanide sows had an accumulation of colloid and follicular cells of low height in the thyroid. No difference in hypophysis, adrenal, pancreas, tongue, esophagus, stomach, liver, cardiovascular tissues, lymphoreticular system of the spleen, tonsils, thymus, intestinal mesenteries, eye, brain or spinal cord were observed between the treatment groups.

<u>Summary</u>

Signs of acute poisoning by cyanide include rapid breathing, gasping, tremors, convulsions and death. The speed of onset and the severity of the

effects depend upon the route, the dose, the duration of exposure and the compound administered. Inhalation of HCN results in the fastest absorption and appearance of toxic signs. Gastrointestinal absorption of the cyanide salts results in passage into the liver, the major site of detoxification. Cyanide exposure results in death because the cyanide ion reacts with cytochrome oxidase, blocking cellular respiration. If enough cyanide is detoxified before death occurs, the animal may recover.

The acute oral LD_{50} for KCN was 10 mg/kg bw (4 mg CN^-/kg bw) in rats (Gaines, 1969; Hayes, 1967) and 8.5 mg KCN/kg bw (3.4 mg CN^-/kg bw) in mice (Sheehy and Way, 1968). Intraperitoneal injection of NaCN in mice results in a similar LD_{50} as orally administered KCN when expressed as cyanide (3.2 mg CN^-/kg bw) (Kruszyna et al., 1982). The dose-response for intraperitoneally administered KCN in mice indicated that 1 or 2 mg KCN/kg bw (0.4 or 0.8 mg CN^-/kg bw) had minimal or no effects, while 3-5 mg KCN/kg bw (1.2-2.0 mg CN^-/kg bw) resulted in signs of toxicity (convulsions, agitation) (Isom et al., 1982). A dose of 6 mg KCN/kg bw (2.4 mg CN^-/kg bw) resulted in 20% mortality. Doses that are fatal to one species may be harmless to others. An oral dose of 3.8 mg KCN/kg bw (1.5 mg CN^-/kg bw) was fatal to a dog in 155 minutes (Gettler and Baine, 1938) but a higher dose of 8 mg KCN/kg bw (3.2 mg CN^-/kg bw), equal to the LD_{50} in mice, had only minimal effects on guinea pigs (Basu, 1983). Rats tolerated much higher doses of cyanide (80 mg CN^-/kg bw/day) when mixed in the diet (Kreutler et al., 1978) than when administered by gavage at 4.0 mg CN^-/kg bw (Ferguson, 1962).

The LC_{50} in mice for inhaled HCN was 184 mg/m³ for 30 minutes (Matjak-Schaper and Alarie, 1982). Correlation of concentration and duration of exposure with effect was difficult due to the high variability in response even within the same species, in which some animals recovered from doses that were fatal to others (Haymaker et al., 1952). When the concentration of HCN was kept at 1000 mg/m³, the time it took for different species to die decreased in the order: monkey > goat > guinea pig = rat > rabbit = cat = mouse > dog (Barcroft, 1931). Therefore, monkeys were least sensitive; dogs were most sensitive.

Animals tolerated higher doses of cyanide when administered in the diet or in the drinking water during subchronic exposure (Hayes, 1967; Palmer and Olson, 1979). Rats receiving ~12 mg CN^-/kg bw/day in drinking water had significantly increased liver weights compared with controls, while rats receiving ~8 mg CN^-/kg bw/day in the diet did not (Palmer and Olson, 1979). No clinical signs of toxicity, effects on body weight, hematology and no histopathologic lesions were found in beagles that consumed 3 mg CN^-/kg bw/day (calculated from body weight data and amount of CN consumed) in the diet for 30 days (American Cyanamid Co., 1959). However, dogs receiving $\geq$0.27 mg CN^-/kg bw/day, administered in a capsule for 15 months, had degenerative changes in ganglion cells of the CNS (Hertting et al., 1960). Subchronic exposure to cyanide by subcutaneous and intramuscular routes has resulted in histopathologic lesions in the brain and spinal cord of rats and monkeys (Lessell, 1971; Smith et al., 1963; Hurst, 1940) and changes in hematologic parameters in baboons (Crampton et al., 1979). Chronic dietary exposure to cyanides (~3.2 and 4.3 mg CN^-/kg bw/day) resulted in no effects clinically or histologically (Howard and Hanzal,

1955), but daily doses of ~30 mg CN^-/kg bw/day in the diet of rats resulted in changes in thyroid homeostasis, reduced body weight gain and moderate demyelination in the spinal cord (Philbrick et al., 1979).

Pertinent data regarding the carcinogenicity of cyanides were not located in the available literature. Cyanide may have antitumor activity (Perry, 1935; Brown et al., 1960). Negative results for KCN were obtained for mutagenicity in Salmonella typhimurium strains TA1535, TA1537, TA1538, TA97, TA98, TA100 and TA102, with and without metabolic activation (De Flora, 1981; DeFlora et al., 1984); although in a study by Kushi et al. (1983) mutagenic activity of HCN gas was found in S. typhimurium strain TA100. Cyanide was negative in a modified rec assay in Bacillus subtilis (Karube et al., 1981). NaCN induced severe teratogenic effects in hamsters when administered by subcutaneously implanted osmotic minipumps that delivered cyanide at a rate of 3.3-3.4 mg CN^-/kg bw/hour (79.2-81.6 mg CN^-/kg bw/day) from days 6-9 of gestation (Doherty et al., 1982). There were no effects of dietary cyanide in a concentration of 500 mg CN^-/kg diet on the reproductive performance of pregnant rats fed KCN throughout gestation and lactation (Tewe and Maner, 1981a). Offspring that were continued on the test diet after weaning consumed less food and grew at a significantly reduced rate compared with control offspring. The effects on the reproductive performance of pigs maintained on diets containing cyanide (30.3, 276.6 and 520.7 mg CN^-/kg diet) throughout gestation and lactation included reduced piglet organ-to-body weight ratios of the thyroid in the medium-dose group relative to the low-dose group, of the spleen in the high-dose group relative to the medium- and low-dose groups and of the heart in the high-dose group relative to the low-dose group of fetuses (Tewe and

Maner, 1981b). Treated sows had hyperplasia of kidney glomerular cells and accumulation of colloid and morphologic changes in follicular cells of the thyroid.

VI. HEALTH EFFECTS IN HUMANS

Acute Exposure

The acute effects of cyanide exposure in humans have been well documented. Several reviews have described the effects of ingestion and inhalation of HCN, KCN and NaCN (DiPalma, 1971, Gosselin et al., 1976; NIOSH, 1976; Hartung, 1982). In addition, systemic toxicity following dermal exposure has been described.

Oral. Acute exposure to cyanide by the oral route has usually occurred from suicide attempts (NIOSH, 1976). Reports of fatal oral doses are in good agreement. These data are summarized in Table VI-1. Gettler and St. George (1934) estimated a lethal dose of cyanide, as hydrocyanic acid (HCN) to be 50 mg (0.71 mg/kg bw for a 70 kg human). Gettler and Baine (1938) reported that the minimum lethal oral dose of HCN was 0.5 mg/kg bw. A range of doses of 0.5-3.5 mg HCN/kg bw taken orally was quoted by Ermans et al. (1972). This range includes the 60-90 mg (0.86-1.29 mg/kg bw) range cited by Gosselin et al. (1976) equivalent to ~1 teaspoon of a 2% hydrocyanic acid solution. Ingestion of 50-100 mg (0.71-1.42 mg/kg bw) of NaCN or KCN can result in immediate collapse and respiratory arrest (Hartung, 1982). The minimum lethal dose of cyanide salts was quoted by DiPalma (1971) as 200 mg for adults (2.9 mg/kg bw), although suicide attempts usually involved ingestion of ≥1-6 g corresponding to "a spoonful." When this amount of cyanide is ingested, ~3.5 mg/kg bw had been absorbed before death resulted. Gastrointestinal absorption of inorganic cyanide salts is slower than pulmonary absorption of HCN, and the onset and severity of symptoms are delayed and diminished. The time-course of responses has been described by DiPalma (1971). Within 1-5 minutes after

TABLE VI-1

Fatal Oral Doses of Cyanide Compounds

Compound	Dose		Reference
	mg	mg CN^-/kg bw	
HCN	50	0.71	Gettler and St. George, 1934
HCN	--	0.5	Gettler and Baine, 1938
CN salts	200	2.9	DiPalma, 1971
HCN	--	0.5-3.5	Ermans et al., 1972
HCN	60-90	0.86-1.29	Gosselin et al., 1976
NaCN or KCN	50-100	0.71-1.42	Hartung, 1982

ingestion, chemoreceptor stimulation results in hyperventilation, the stomach lining becomes irritated and vomiting may occur. At 5-20 minutes, unconsciousness, convulsions, muscular contraction of the jaw, flushed and dry skin, rapid and irregular pulse and gasping occur. Vascular collapse, dilation of the pupils and cyanosis follow. Bodansky and Levy (1923) administered KCNS or KCN in a gelatin capsule on two consecutive days to 25 human subjects. On the first day, the subjects received 15 mg KCNS, and on the second day, 10 mg KCN. Assuming the average weight of a human to be ~70 kg, these amounts represent doses of 0.14 mg KCN/kg bw and 0.21 mg KCNS/kg bw (both equivalent to ~0.06 mg CN^-/kg bw). All subjects effectively detoxified these doses, as determined by thiocyanate level measurements.

Liebowitz and Schwartz (1948) reported a case of a 60-year-old, 80 kg man who had attempted suicide by ingesting an estimated 3-5 g of KCN. At one-half hour after ingestion, he vomited. One-half hour later, on admission to the hospital, he was comatose, and gastric lavage was performed. One-half hour after lavage, he regained consciousness; 8 hours after admission he had short-lived lingering effects of nausea and weakness. Subsequently he recovered fully. The blood level of cyanide at 2 hours after admission was 0.2 mg HCN/mℓ. Assuming a blood volume of 6 ℓ, the investigators calculated that 1.2 g (15 mg/kg bw) of HCN was present in the circulating blood. This absorbed dose is at least 5 times the lethal dose of ingested KCN, and yet, the patient recovered. No apparent reason for his recovery was discovered; however, recovery from ingestion of cyanide salts is not unprecedented. Liebowitz and Schwartz (1948) cited 22 cases of recovery. Utti et al. (1985) reported a case in which an 18-year-old man ingested 975-1300 mg KCN (13.9-18.5 mg KCN/kg bw), was treated and survived.

Shortly after recovery, the man developed severe Parkinsonian syndrome and died 19 months after the cyanide ingestion. The clinicopathologic findings indicated that the Parkinsonism was the result of cyanide poisonings.

Inhalation. Inhalation of HCN gas results in the most rapid onset of poisoning, producing almost immediate collapse, respiratory arrest and death within minutes (DiPalma, 1971). Although cyanide has a characteristic odor of bitter almond, in human, the ability to smell cyanide is controlled by a sex-linked recessive gene (Kirk and Stenhouse, 1953). NIOSH (1976) compiled data from human case reports and animal studies to estimate concentration-response effects for humans (Table VI-2). These data indicated that air-borne concentrations of HCN of 99-528 mg/m³ are fatal in 30-60 minutes. However, some reports indicated no injury at concentrations as high as 550 mg/m³ for 1 minute, while others reported that 297 mg/m³ was immediately fatal. Discrepancies may be due to individual variations and, in some cases, the problems inherent in extrapolating data from animal experiments in order to predict human responses.

Barcroft (1931) placed a man (70 kg) and a dog (12 kg) in an inhalation chamber and exposed them simultaneously to HCN at concentrations of 550-688 mg/m³. After 50 seconds, the dog became unsteady; at 1.25 minutes, the dog collapsed and was unconscious; at 1.5 minutes, the dog had tetanic convulsions. One second later (91 seconds of exposure), the man emerged from the chamber, and put on a respirator, but felt no symptoms. Two seconds later, the dog was believed to be dead (it recovered, and was walking around the next morning). Within a few minutes after exposure, the man was briefly nauseated and had difficulty concentrating.

TABLE VI-2

Reported (Estimated) Human Responses to Various Concentrations of HCN Vapors[a]

Reponses	Concentration (mg/m^3)[b]
Fatal in 6-8 minutes	297
Fatal after 30 minutes	243-528
Fatal after 10 minutes	199
Fatal after 30 minutes	149
Fatal after 30-60 minutes	121-149
Fatal after 30-60 minutes	110-264
Fatal after 60 minutes	99
Tolerated for 30-60 minutes without immediate or after effects	50-59
Complaints of headache, nausea, vomiting, cardiac symptoms	50
Minimal symptoms after several hours of exposure	22-44
Effects after several hours of exposure	20-40
No observed effect	0-19; mean 5.4
Fatigue, headache, body weakness, tremor, pain, nausea	5.5-14.3
Headache, weakness, changes in taste and smell, throat irritation, nausea, effort dyspnea, enlarged thyroids, changes in blood chemistry	4.6-13.6; mean, 9.1
Increased thiocyanate excretion in urine, but to a lesser extent than in cigarette smokers; no other effects noted	2.2-8.8; mean 5.5
Slight decrease in leukocytic activity of cytochrome oxidase, peroxidase and succinate dehydrogenase after an average of 5.4 years of exposure	0.25

TABLE VI-2 (cont.)

Reponses	Concentration (mg/m^3)[b]
No effects	0.11-0.99
No symptoms after 6 hours	20-40
No serious consequences in 1 minute	550
No injury in 1 minute	550
No injury in 1.5 minutes	413
Nausea and difficulty concentrating after 91-second exposure	550-688

[a]Source: NIOSH, 1976

[b]The concentrations in mg/m^3 were calculated from concentrations expressed in ppm by multiplying by 1.1.

According to DiPalma (1971), acute exposure of humans to inhalation of ~2200 mg HCN/m³ results first in shortness of breath, local irritation to the throat and a feeling of warmth. Following a period of rapid breathing and sometimes an outcry, apnea, gasping, collapse, convulsions and often cardiac arrest occur within minutes. It was estimated that rapid death results from an absorbed dose of as little as 0.7 mg HCN/kg bw. At lower airborne concentrations (~5-50 mg/m³) (see Table VI-2), symptoms include dizziness, numbness, weakness, headache, nausea and vomiting, confusion, rapid breathing and rapid pulse (NIOSH, 1976; Hartung, 1982). Wexler et al. (1947) found that intravenous injection of 0.11-0.2 mg NaCN/kg bw (roughly equivalent to 0.06-0.11 mg HCN/kg bw absorbed following inhalation, as estimated by ratio of molecular weights) in humans resulted in sinus pause and irregularly slowed heart rate, followed by an increased pulse above control.

Dermal. Absorption of HCN or solutions of cyanide salts through the skin may result in effects similar to the sublethal effects of inhalation exposure (Potter, 1950). Local effects to the skin such as dermatitis and rash may also result (Drinker, 1932; NIOSH, 1976). An LD_{50} for dermal absorption of HCN, which is much slower than pulmonary absorption, was estimated as 100 mg/kg bw (DiPalma, 1971).

Subchronic and Chronic Exposure

Oral. Pertinent data regarding chronic oral exposure of humans to hydrocyanic acid, KCN and NaCN were not located in the available literature. There is a body of literature on the etiology of thyroid disorders and neuropathies characterized by optic atrophy, nerve deafness and spinal

ataxia in people living in certain tropical areas of Africa, where the staple diet consists largely of cassava (Monekosso and Wilson, 1966; Osuntokun, 1968, 1972; Osuntokun et al., 1969, 1970; Makene and Wilson, 1972; Ermans et al., 1972; Delange and Ermans, 1971). Cassava is a root food that contains a high level of the cyanogenic glycoside, linamarin. Linamarin releases cyanide on metabolism or acid hydrolysis in vivo; therefore, exposure is not to cyanide per se. Tropical neuropathies have also been associated with such factors as infections, protein deficiency, riboflavin deficiency, vitamin B_{12} (cobalamin) deficiency or defective B_{12} metabolism (Osuntokun, 1968; Osuntokun et al., 1969; Makene and Wilson, 1972). Ingestion of cassava, in combination with iodine deficiency, has been associated with high incidences of goiter and cretinism in Zaire (Delange and Ermans, 1971; Ermans et al., 1972).

Inhalation.

Cigarette Smoke -- Inhalation exposure to cyanide in tobacco smoke by heavy smokers has been associated with the condition of tobacco amblyopia, characterized by a loss of visual acuity; Leber's hereditary optic atrophy, a similarly manifested disorder; and with retrobulbar neuritis complicating pernicious anemia and optic atrophy (Wokes, 1958; Pettigrew and Fell, 1972, 1973; Wilson and Matthews, 1966; Foulds et al., 1968; Wilson, 1983). These disorders appear to involve defective metabolism of cyanide to thiocyanate, as well as a deficiency in vitamin B_{12} or defective vitamin B_{12} metabolism.

Smoking during pregnancy is associated with a higher risk of giving birth to low body weight infants and of perinatal death (Andrews, 1973).

Andrews (1973) compared plasma thiocyanate levels in maternal and cord blood taken at delivery from 21 women who had smoked during pregnancy and from 22 women who had not. For both maternal and cord blood, smokers (>10 cigarettes/day) had significantly ($p<0.001$) higher levels of thiocyanate than did nonsmokers. Although no differences were found for infant weight, duration of gestation or onset of labor, group sizes were small (21 smokers, 22 nonsmokers). This study does demonstrate, however, the potential for intra-uterine exposure. The use of low-tar, low-nicotine or filter cigarettes does not reduce the HCN concentration in cigarette smoke (Way, 1984).

Occupational Exposure.

Case Studies -- Chronic exposure to the cyanides by the inhalation and dermal routes involves occupational exposure of persons engaged in case hardening and polishing of metals and photographic materials workers. Sandberg (1967) described a case of a goldsmith apprentice. In the course of his work, the man cleaned goldware 5-10 times daily for 4 years. The cleaning solution was prepared by adding 15 g KCN to a liter of water, heating to boiling and adding 50 mℓ of hydrogen peroxide. This procedure caused spattering on the skin. His symptoms included headaches, listlessness, partial paralysis of his left arm and left leg, a grayish pallor and partial loss of vision in the left eye. Hardy et al. (1950) described a case hardener who had been consistently exposed to HCN in his work. His symptoms included headaches, episodes of dizziness, confusion, muscular weakness, poor vision, slurred speech, abdominal cramps, tremor, body rash and an enlarged thyroid. In another case, a man who had been frequently exposed to cyanide for 5 months while working as a machinist developed a goiter. The thyroid enlargements in these two cases were presumed to be a

result of the action of thiocyanate formed upon metabolism of cyanide. Many additional cases of chronic occupational exposure to cyanide of individuals who developed similar conditions have been reviewed by NIOSH (1976).

Epidemiologic Studies. NIOSH (1976) reviewed several epidemiologic studies that have been published in various countries around the world (Saia et al., 1970; Carmelo, 1955; Dinca et al., 1972; and Radojicic, 1973), including one conducted in Egypt (El Ghawabi et al., 1975) and one in India (Chandra et al., 1980). These studies are summarized in Table VI-3. Typical symptoms and manifestations of cyanide poisoning were observed in each study, with the exception of the study by Dinca et al. (1972), which studied enzyme activity only. NIOSH (1976) questioned the validity of the techniques in measuring enzyme activity. Blood smears were stained for detection of enzyme granules in neutrophilic leukocytes. In the study of Carmelo (1955), the original exposed cohort had consisted of 600 fumigators who had experienced acute episodes of intoxication. Their symptoms were typical. In the study of El Ghawabi et al. (1975), the study groups had been instructed to avoid foods containing cyanogenic and thiocyanogenic glucosides (e.g., cabbage-like vegetables) (Ermans et al., 1972). The increased mean uptake of ^{131}I by the exposed workers, who had not been at work for 2 days prior to the determination, was explained as a response to acute cyanide withdrawal. Chandra et al. (1980) also instructed the subjects to avoid cyanogenic foods. In addition, data from smokers and nonsmokers were analyzed separately. Smokers showed higher levels of cyanide and thiocyanate in their blood and urine.

TABLE VI-3

Summary of Epidemiologic Studies of Cyanide Exposure*

Study Population	Control Population	Duration of Exposure	Level of Exposure	Effects	Reference
36 male electroplaters from 3 factories in Egypt Age: 30-50+ years All were non-smokers	20 normal male workers of same age and socio-econimic status, with no exposure to cyanide All were non-smokers	5-15 years	Mean concentration of HCN in breathing zone was 7.1, 8.9 and 11.4 mg/m³ for the 3 factories	Of the 36 exposed men, the following incidences were recorded: headache-29, weakness-28, changes in smell and taste-28, giddiness-20, throat irritation-16, vomiting-16, difficulty breathing-16, lacrimation-9, precordial pain-7, excessive salivation-3, difficulty focusing eyes-3, psychosis (confusion, hallucination, slurred speech)-2. The incidence of these symptoms were much lower in controls. 20 of the exposed men had mild to moderate thyroid enlargement. The exposed workers had significantly higher uptake of ^{131}I by the thyroid 4 and 24 hours after administration than the controls ($p<0.001$). The exposed group had significantly higher hemoglobin and lymphocyte counts ($p<0.001$).	El Ghawabi et al., 1975
23 (of 40 male workers) were available for study from a factory engaged in electroplating and casehardening in India Age: 23-40 years	20 male workers of the same age and socio-economic status with no exposure to cyanide or other hazardous chemicals	NR The study was conducted during months of May and June	Airborne concentration in workroom was 0.2-0.8 mg/m³, mean, 0.45 mg/m³ Airborne concentration in breathing zone was 0.1-0.2 mg/m³, mean, 0.15 mg/m³	Exposed workers had significantly higher levels of cyanide and thiocyanate in blood and urine than the control group. The exposed workers complained of typical symptoms	Chandra et al., 1980
21 workers of a galvanizing department who worked near cyanide vats and 19 wire hardeners and zinc platers	22 men who washed the wire in caustic soda	NR	NR	The exposed group had twice the incidence of insomnia, disturbed sleep, tremors, dermatitis (thought to be due to poor work habits), nosebleed and vertigo. The control group had relatively high incidences of headache, dizziness, buzzing in ears, irritation of eyes, cold sweat, poor digestion, abdominal pain, muscle weakness and shortness of breath (they worked with caustic soda).	Sala et al., 1970

TABLE VI-3 (cont.)

Study Population	Control Population	Duration of Exposure	Level of Exposure	Effects	Reference
13 HCN fumigators who had suffered acute episodes of HCN poisoning with loss of consciousness	4 HCN fumigators who did not report acute episodes of of HCN poisoning	1-27 years	NR	The 13 intoxicated men had high incidences of nervous disorders, such as, disturbance of equilibrium, vertigo, rapid eye movement, moebius sign (inability to maintain conversion of eyes in exopthalmic goiter). Precordial pain was a complaint of 9 of the 13 men, 11 had EKG abnormalities, 11 had hypertropic gastritis. None of the 4 controls had stomach or intestinal disorders.	Carmelo, 1955
31 men, 12 women in Romania who worked in metal galvanizing with chronic exposure to airborne HCN Age: 18-49 years Mean - 36 years	no details of control group	0.25-16 years mean: 5.4 years	Average exposure for last 5 years: 0.26 mg/m³	Significantly decreased activity of cytochrome oxidase compared with controls. Considerably reduced activity of peroxidase and succinodehydrogenase.	Dinca et al., 1972
43 electroplaters and annealers in Yugoslavia exposed to cyanide	20 non-exposed workers	NR	7-11 mg/m³	Most of the exposed workers had symptoms of fatigue, headache, weakness, tremor of hands and feet, pain and nausea. Exposed workers had higher urinary thiocyanate levels than non-exposed workers.	Radojicic, 1973

*The studies of Sala et al., 1970; Carmelo, 1955; Dinca et al., 1972 and Radojicic, 1973 were cited in NIOSH, 1976.

NR - Not reported

High Risk Subpopulations

Pregnant women who smoke may increase the susceptibility of their infants to toxic effects of cyanide. Smoking during pregnancy has been linked to low-weight of infants and perinatal death, and higher plasma thiocyanate levels were found in cord blood of smokers than nonsmokers (Andrews, 1973).

A metabolic disturbance in the conversion of cyanide to thiocyanate and vitamin B_{12} deficiency have been associated with such conditions as tobacco amblyopia, Leber's hereditary optic atrophy and retrobulbar neuritis in people who are exposed to excess cyanide in tobacco smoke (Wilson, 1983). Excess exposure to cyanide and iodine deficiencies has been associated with thyroid abnormalities, such as goiter and cretinism (Delange and Ermans, 1971; Ermans et al., 1972). Neuropathies have been associated with increased cyanide exposure due to chronic dietary intake of cassava as a staple food, in combination with protein deficiency, riboflavin and vitamin B_{12} deficiencies and infections (Osuntokun et al., 1969; Osuntokun, 1968; Makene and Wilson, 1972). Therefore, individuals with a metabolic defect in the rhodanese system (the enzyme that transfers sulfur to cyanide, see Metabolism Section in Chapter III), a vitamin B_{12} deficiency or defective B_{12} metabolism or iodine deficiency, and fetuses exposed *in utero*, are at a higher risk to the toxic effects of excessive cyanide intake than the general population.

Summary

Acute oral exposure to cyanide has usually occurred from suicide attempts. Fatal orally absorbed doses range from 0.5-3.5 mg HCN/kg bw

(Gettler and St. George, 1934; Gettler and Baine, 1938; Ermans et al., 1972; Gosselin et al., 1976). The minimum lethal dose of orally ingested cyanide salts is ~2.9 mg/kg bw (DiPalma, 1971). When 1-6 g ("a spoonful") is ingested, ~3.5 mg/kg bw has been absorbed by the time of death. Within 20 minutes of exposure to fatal doses, events progress from hyperventilation, vomiting, unconsciousness, convulsions, rapid and irregular heart rate, gasping, vascular collapse and cyanosis, to death. Several cases of recovery have been reported after ingestion of cyanide at doses that have proved fatal in other cases (Liebowitz and Schwartz, 1948; Utti et al., 1985).

Inhalation of HCN is the most rapid route of fatal exposure and poisoning. Acute exposure of humans to ~2200 mg HCN/m³ results in shortness of breath, irritation to the throat and warmth followed by rapid breathing, an outcry, apnea, gasping, collapse, convulsions and death in minutes (DiPalma, 1971). As little as 0.7 mg HCN/kg bw absorbed after inhalation may result in rapid death. Sublethal concentrations (5-50 mg/m³) result in dizziness, confusion, headache, nausea, numbness and hyperventilation (NIOSH, 1976). Acute dermal exposure produces similar effects, although much more slowly (Potter, 1950; DiPalma, 1971).

Chronic oral exposure to HCN, KCN or NaCN has not been described. People living in tropical areas of Africa consume cassava as a staple food, which contains high levels of linamarin, a cyanogenic glycoside that releases cyanide on metabolism or acid hydrolysis <u>in vivo</u> (Osuntokun et al., 1969; Osuntokun, 1968; Makene and Wilson, 1972). These people have high incidences of neuropathies associated with cyanide, vitamin B_{12} deficiency, protein deficiency and infections. Ingestion of cassava, together

with iodine deficiency, has been associated with the etiology of goiter and cretinism (Ermans et al., 1972, Delange and Ermans, 1971).

Chronic exposure to cyanide has been associated with tobacco amblyopia in heavy cigarette smokers, Leber's hereditary optic atrophy and retrobulbar neuritis (Wilson, 1983). A defect in cyanide metabolism and vitamin B_{12} deficiency appear to be involved in these disorders. Smoking during pregnancy may result in low birth-weight infants (Andrews, 1973).

Case studies and epidemiologic studies of metal case hardeners and HCN fumigators occupationally exposed to cyanide fumes, describe effects in workers typical of sublethal cyanide poisoning, including headache, dizziness and thyroid enlargement (NIOSH, 1976).

VII. MECHANISMS OF TOXICITY

Acute

The mechanism by which cyanide exerts its acute toxic effects is well understood and has been reviewed in numerous reports. The following discussion is based on reviews by DiPalma (1971), Gosselin et al. (1976), U.S. EPA (1980), Solomonson (1981), Hartung (1982) and Way (1984).

The cyanide ion can be formed *in vivo* by dissociation of HCN, KCN, NaCN and any other cyanogenic compounds. The ion has a high affinity for many biologically active metal ions. It forms a stable complex with iron in the ferric state (Fe^{+++}). Cytochrome oxidase, the terminal enzyme in the mitochondrial electron transport chain, is inhibited by HCN complexing with the heavy metal ions contained in the enzyme (Way, 1984). The inactivated enzyme is unable to catalyze the reaction:

$$Fe^{2+}\text{-cytochrome} + 0.5\ O \xrightarrow[\text{oxidase}]{\text{cytochrome}} Fe^{3+}\text{-cytochrome} + 0.5O^{2-}$$

resulting in inhibition of cellular respiration. The inhibition results in a histotoxic hypoxia since cells are unable to utilize oxygen. Oxidative phosphorylation is unable to proceed, and there is a shift to anaerobic metabolism with resultant accumulation of lactate. The cells cannot use oxygen delivered by oxyhemoglobin, thus, venous blood appears bright red since hemoglobin continues to carry oxygen. Since CNS and the heart are particularly sensitive to hypoxia, toxic effects are manifested. The time course for these events is very rapid and death can ensue in minutes (DiPalma, 1971).

Cyanide is capable of complexing with many enzymes and compounds that contain Fe^{+++} including catalase, peroxidase and methemoglobin. Cyanide can also form complexes with compounds that contain metals other than iron such as hydroxycobalamin (vitamin B_{12a}), phosphatase, tyrosinase, ascorbic acid-oxidase, xanthine oxidase and succinic dehydrogenase (DiPalma, 1971). However, the reaction with cytochrome oxidase is the most important for the onset of the acute toxic effects.

As discussed in Chapter III, cyanide is detoxified by the action of the enzyme rhodanese (Lang, 1933), the trivial name for cyanide sulfurtransferase. The enzyme transfers sulfur from a sulfur donor to cyanide to form the less toxic thiocyanate, which is excreted in the urine (Williams, 1959). The major site of rhodanese activity in most species (rat, rabbit, monkey) is the liver (Himwich and Saunders, 1948), with particular localization in mitochondria (de Duve et al., 1955). The rate-limiting factor in cyanide detoxication, therefore, is the availability of an intramitochondrial sulfur donor.

Chronic

The effects of chronic low-level exposure to cyanide are neurologic in nature. Repeated insults to neurologic tissue resulting in histotoxic hypoxia may be responsible for the demyelination and necrosis of many tissues of the CNS, including the optic nerve (Ferraro, 1933; Lessell, 1971; Lessell and Kuwabara, 1974; Smith et al., 1963). Wilson (1983) has reviewed the role of cyanide in human diseases. Tobacco amblyopia, associated with heavy smoking and vitamin B_{12} depletion, and retrobulbar neuritis may be the result of abnormal cyanide and vitamin B_{12} metabolism (Wilson, 1983;

Wokes, 1958). These conditions are effectively treated with hydroxycobalamin (vitamin B_{12}), which can react with cyanide to form cyanocobalamin (Wilson et al., 1971). If there were a defect in cyanide conversion to thiocyanate, or insufficient vitamin B_{12} to detoxify the accumulated cyanide, neurologic disorders may occur. Similar interrelationships may be involved in the etiology of such disorders as Leber's hereditary optic atrophy (Wilson, 1983). Neuropathies and amblyopias among people living in the tropical areas may also involve vitamin B_{12} deficiency. This is especially true in Africa, where cassava, a root food rich in the cyanogenic glycosides, linamarin and lotaustralin, is a dietary staple. In these disorders, many complicating factors, such as protein and other vitamin deficiencies, may also be involved (Wilson, 1983). The interrelationships between cyanide, thiocyanate, vitamin B_{12} and inherent defects in the metabolic control of the conversions are not completely understood. One suggestion was that cyanide, formed <u>in vivo</u> from thiocyanate by the enzymatic action of thiocyanate oxidase in red blood cells (Goldstein and Rieders, 1953) may reach levels that are chronically toxic if there is a deficiency in vitamin B_{12}. Therefore, vitamin B_{12} may play a role in the regulation of cyanide metabolism. Increased blood levels of thiocyanate resulting from ingestion of cyanogenic foods, in combination with iodine deficiency, may play a role in the etiology of goiter and cretinism among inhabitants of the tropics (Ermans et al., 1972; Delange and Ermans, 1971). Thiocyanate is known competitively to inhibit the uptake of iodide by the thyroid.

Synergism

Other chemicals known to inhibit cytochrome oxidase are sulfide and azide (Smith and Gosselin, 1979; Smith et al., 1977), and they may as such have a synergistic or additional effect on cyanide toxicity. Sulfide is a more potent inhibitor of cytochrome oxidase than is cyanide, and may act by the same mechanism (Nicholls, 1975; Smith et al., 1977). Basu (1983) found that guinea pigs pretreated with ascorbate prior to oral administration of KCN developed a 100% incidence of severe tremor, ataxia, muscle twitches, paralysis and convulsion compared with only slight tremors in 38% of the animals treated only with KCN. It was postulated that ascorbate ties up cysteine so that it cannot act as a sulfur donor for the enzymatic action of rhodanese to convert cyanide to thiocyanate. Administration of cysteine had a protective effect on the enhanced toxicity of ascorbate plus cyanide. In a recent paper, Levin et al. (1987) examined the effects of exposure to a variety of gases including HCN and low oxygen levels that could be produced by fires. Thirty-minute studies on the lethal effects of carbon monoxide and HCN indicated they act in an additive fashion. Exposure to 5% CO reduced the LC_{50} value for HCN exposure in rats from 110 ppm to 75 ppm.

Antagonism

Since cyanide acts by inhibiting cytochrome oxidase, thereby inhibiting the cellular utilization of oxygen, oxygen is not an antagonist of cyanide toxicity (U.S. EPA, 1980). Since cyanide can bind to Fe^{+++}, methemoglobin, which contains the ferric ion, can compete with cytochrome oxidase for cyanide, forming cyanomethemoglobin (Hartung, 1982; Way 1981, 1984). The use of sodium nitrite, hydroxylamine, amyl nitrite and other compounds capable of generating methemoglobin as antidotes to acute cyanide poisoning

has its basis in this reaction (Smith and Olson, 1973; Way, 1981, 1984). Methylene blue has been used as an antidote in the past (Trautman, 1930), and its use may still be advocated by some physicians; however, it does not form methemoglobin as readily as the nitrites (Way, 1981).

Although oxygen in itself is not an antagonist, numerous studies have found that oxygen enhances the antidotal efficacy of the combination of sodium thiosulfate (a sulfur donor for the rhodanese reaction) and sodium nitrite (a methemoglobin generator) (Way et al., 1966; Burrows et al., 1973; Isom and Way, 1974; Sheehy and Way, 1968). Schwartz et al. (1979) found that pretreatment of mice with pyruvate had a protective effect against cyanide poisoning. Pyruvate enhanced the antidotal effect of sodium thiosulfate, but not of sodium nitrite. When used in combination with sodium thiosulfate and sodium nitrite, maximum antidotal efficacy was observed.

Administration of compounds that can serve as sulfur donors for rhodanese may be a useful antidotal approach. Sorbo (1953) compared the effectiveness of many sulfur containing compounds and concluded that the structural requirements were for a free sulfur atom to be adjacent to another sulfur atom in the molecule. Frankenberg (1980) injected mice intravenously with bovine liver rhodanese and a sulfur donor, following the administration of cyanide and found the enzyme to have an antidotal effect on cyanide poisoning. Disulfonic acid (DIDS) has been found to reduce cyanide-induced contractions <u>in vitro</u> by blocking ionic channel mechanisms that facilitate the entry of cyanide into vascular smooth muscle cells (Robinson et al., 1985).

Cobalt and cobalt containing compounds, such as vitamin B_{12} (hydroxocobalamin) (Mushett et al., 1952), cobalt-EDTA (Davison, 1969; Friedberg and Schwarzkopf, 1969) and cobaltous histidine (Friedberg and Schwarzkopf, 1969) have been used as effective antidotes to cyanide toxicity because of the affinity of cyanide for cobalt.

Summary

Cyanide exerts its acute toxic effect by forming a complex with Fe^{+++} in cytochrome oxidase, the terminal enzyme in the mitochondrial electron transport chain, thereby inhibiting utilization of oxygen by cells (DiPalma, 1971). The inhibition results in histotoxic hypoxia, preventing oxidative phosphorylation. There is a shift to anaerobic metabolism. The CNS and the heart are particularly sensitive to hypoxia, and toxicity is manifested. Cyanide can complex with other compounds that contain metals, but the reaction with cytochrome oxidase is the most important, toxicologically. Cyanide is detoxified by the enzymatic activity of rhodanese, which transfers sulfur from a donor molecule to cyanide to form thiocyanate.

Chronic cyanide exposure may contribute to the etiology of neurologic and optical disorders (tobacco amblyopia, tropical neuropathy, Leber's hereditary optic atrophy), since repeated insults of hypoxia to neurologic tissue results in demyelination and necrosis of many tissues of the CNS, including the optic nerve in animals (Ferraro, 1933; Lessell, 1971; Lessell and Kuwabara, 1974; Smith et al., 1963). Defects in cyanide metabolism, along with nutritional deficiencies of protein and vitamin B_{12}, are believed to be involved (Wilson, 1983). The goitrogenic effect of cyanide may be due to thiocyanate, which inhibits the uptake of iodine by the thyroid (Ermans et al., 1972).

Synergistic effects may occur if cyanide exposure is accompanied by exposures to other substances known to inhibit cytochrome oxidase, such as sulfide and azide (Smith and Gosselin, 1979). Ascorbate potentiated the toxicity of cyanide in guinea pigs, perhaps by reacting with cysteine and thereby preventing cysteine from acting as a sulfur donor for rhodanese (Basu, 1983).

Cyanide poisoning can be antagonized by methemoglobin, or any compound that can generate methemoglobin in vivo, such as sodium nitrite (Way, 1981, 1984). Oxygen enhances the antidotal efficacy of sodium thiosulfate and sodium nitrite (Way, 1981), as does pyruvate (Schwartz et al., 1979). Cyanide probably reacts with the carbonyl group of pyruvate to yield a cyanohydrin. Any compound that can act as a sulfur donor for rhodanese would be a cyanide antagonist (Sorbo, 1953), and treatment with rhodanese itself was an effective antidote (Frankenberg, 1980). Cobalt containing compounds have also been used as effective antidotes to cyanide poisoning (Mushett et al., 1952; Davison, 1969; Friedberg and Schwarzkopf, 1969).

VIII. QUANTIFICATION OF TOXICOLOGIC EFFECTS

Introduction

The quantification of toxicologic effects of a chemical consists of separate assessments of noncarcinogenic and carcinogenic health effects. Chemicals that do not produce carcinogenic effects are believed to have a threshold dose below which no adverse, noncarcinogenic health effects occur, while carcinogens are assumed to act without a threshold.

In the quantification of noncarcinogenic effects, a Reference Dose (RfD), [formerly termed the Acceptable Daily Intake (ADI)] is calculated. The RfD is an estimate (with uncertainty spanning perhaps an order of magnitude) of a daily exposure to the human population (including sensitive subgroups) that is likely to be without an appreciable risk of deleterious health effects during a lifetime. The RfD is derived from a no-observed-adverse-effect level (NOAEL), or lowest-observed-adverse-effect level (LOAEL), identified from a subchronic or chronic study, and divided by an uncertainty factor(s) times a modifying factor. The RfD is calculated as follows:

$$\text{RfD} = \frac{\text{(NOAEL or LOAEL)}}{\text{[Uncertainty Factor(s) x Modifying Factor]}} = ____ \text{ mg/kg bw/day}$$

Selection of the uncertainty factor (UF) to be employed in the calculation of the RfD is based upon professional judgment, while considering the entire data base of toxicologic effects for the chemical. In order to ensure that uncertainty factors are selected and applied in a consistent manner, the U.S. EPA (1991) employs a modification to the

guidelines proposed by the National Academy of Sciences (NAS, 1977, 1980) as follows:

Standard Uncertainty Factors (UFs)

- Use a 10-fold factor when extrapolating from valid experimental results from studies using prolonged exposure to average healthy humans. This factor is intended to account for the variation in sensitivity among the members of the human population. [10H]

- Use an additional 10-fold factor when extrapolating from valid results of long-term studies on experimental animals when results of studies of human exposure are not available or are inadequate. This factor is intended to account for the uncertainty in extrapolating animal data to the case of humans. [10A]

- Use an additional 10-fold factor when extrapolating from less than chronic results on experimental animals when there is no useful long-term human data. This factor is intended to account for the uncertainty in extrapolating from less than chronic NOAELs to chronic NOAELs. [10S]

- Use an additional 10-fold factor when deriving an RfD from a LOAEL instead of a NOAEL. This factor is intended to account for the uncertainty in extrapolating from LOAELs to NOAELs. [10L]

Modifying Factor (MF)

- Use professional judgment to determine another uncertainty factor (MF) that is greater than zero and less than or equal to 10. The magnitude of the MF depends upon the professional assessment of scientific uncertainties of the study and data base not explicitly treated above, e.g., the completeness of the overall data base and the number of species tested. The default value for the MF is 1.

The uncertainty factor used for a specific risk assessment is based principally upon scientific judgment rather than scientific fact and accounts for possible intra- and interspecies differences. Additional considerations not incorporated in the NAS/ODW guidelines for selection of an uncertainty factor include the use of a less than lifetime study for deriving an RfD, the significance of the adverse health effects and the counterbalancing of beneficial effects.

From the RfD, a Drinking Water Equivalent Level (DWEL) can be calculated. The DWEL represents a medium specific (i.e., drinking water) lifetime exposure at which adverse, noncarcinogenic health effects are not anticipated to occur. The DWEL assumes 100% exposure from drinking water. The DWEL provides the noncarcinogenic health effects basis for establishing a drinking water standard. For ingestion data, the DWEL is derived as follows:

$$\text{DWEL} = \frac{(\text{RfD}) \times (\text{Body weight in kg})}{\text{Drinking Water Volume in } \ell/\text{day}} = ____ \text{ mg}/\ell$$

where:

Body weight = assumed to be 70 kg for an adult
Drinking water volume = assumed to be 2 ℓ/day for an adult

In addition to the RfD and the DWEL, Health Advisories (HAs) for exposures of shorter duration (1-day, 10-day and longer-term) are determined. The HA values are used as informal guidance to municipalities and other organizations when emergency spills or contamination situations occur. The HAs are calculated using an equation similar to the RfD and DWEL; however, the NOAELs or LOAELs are identified from acute or subchronic studies. The HAs are derived as follows:

$$\text{HA} = \frac{(\text{NOAEL or LOAEL}) \times (\text{bw})}{(\text{UF}) \times (___ \ \ell/\text{day})} = ____ \text{ mg}/\ell$$

Using the above equation, the following drinking water HAs are developed for noncarcinogenic effects:

1. 1-day HA for a 10 kg child ingesting 1 ℓ water per day.
2. 10-day HA for a 10 kg child ingesting 1 ℓ water per day.
3. Longer-term HA for a 10 kg child ingesting 1 ℓ water per day.
4. Longer-term HA for a 70 kg adult ingesting 2 ℓ water per day.

The 1-day HA calculated for a 10 kg child assumes a single acute exposure to the chemical and is generally derived from a study of <7 days duration. The 10-day HA assumes a limited exposure period of 1-2 weeks and is generally derived from a study of <30 days duration. The longer-term HA is derived for both the 10 kg child and a 70 kg adult and assumes an exposure period of ~7 years (or 10% of an individual's lifetime). The longer-term HA is generally derived from a study of subchronic duration (exposure for 10% of animal's lifetime).

The U.S. EPA categorizes the carcinogenic potential of a chemical, based on the overall weight-of-evidence, according to the following scheme:

> Group A: Human Carcinogen. Sufficient evidence exists from epidemiology studies to support a causal association between exposure to the chemical and human cancer.
>
> Group B: Probable Human Carcinogen. Sufficient evidence of carcinogenicity in animals with limited (Group B1) or inadequate (Group B2) evidence in humans.
>
> Group C: Possible Human Carcinogen. Limited evidence of carcinogenicity in animals in the absence of human data.
>
> Group D: Not Classified as to Human Carcinogenicity. Inadequate human and animal evidence of carcinogenicity or for which no data are available.
>
> Group E: Evidence of Noncarcinogenicity for Humans. No evidence of carcinogenicity in at least two adequate animal tests in different species or in both adequate epidemiologic and animal studies.

If toxicologic evidence leads to the classification of the contaminant as a known, probable or possible human carcinogen, mathematical models are used to calculate the estimated excess cancer risk associated with the ingestion of the contaminant in drinking water. The data used in these

estimates usually come from lifetime exposure studies using animals. In order to predict the risk for humans from animal data, animal doses must be converted to equivalent human doses. This conversion includes correction for noncontinuous exposure, less than lifetime studies and for differences in size. The factor that compensates for the size difference is the cube root of the ratio of the animal and human body weights. It is assumed that the average adult human body weight is 70 kg and that the average water consumption of an adult human is 2 ℓ of water per day.

For contaminants with a carcinogenic potential, chemical levels are correlated with a carcinogenic risk estimate by employing a cancer potency (unit risk) value together with the assumption for lifetime exposure from ingestion of water. The cancer unit risk is usually derived from a linearized multistage model with a 95% upper confidence limit providing a low dose estimate; that is, the true risk to humans, while not identifiable, is not likely to exceed the upper limit estimate and, in fact, may be lower. Excess cancer risk estimates may also be calculated using other models such as the one-hit, Weibull, logit and probit. There is little basis in the current understanding of the biological mechanisms involved in cancer to suggest that any one of these models is able to predict risk more accurately than any other. Because each model is based upon differing assumptions, the estimates derived for each model can differ by several orders of magnitude.

The scientific data base used to calculate and support the setting of cancer risk rate levels has an inherent uncertainty that is due to the systematic and random errors in scientific measurement. In most cases, only studies using experimental animals have been performed. Thus, there is

uncertainty when the data are extrapolated to humans. When developing cancer risk rate levels, several other areas of uncertainty exist, such as the incomplete knowledge concerning the health effects of contaminants in drinking water, the impact of the experimental animal's age, sex and species, the nature of the target organ system(s) examined and the actual rate of exposure of the internal targets in experimental animals or humans. Dose-response data usually are available only for high levels of exposure and not for the lower levels of exposure closer to where a standard may be set. When there is exposure to more than one contaminant, additional uncertainty results from a lack of information about possible synergistic or antagonistic effects.

Noncarcinogenic Effects

Cyanides are readily absorbed from the lungs, the GI tract and the skin by animals and humans. Inhalation exposure to HCN provides the most rapid rate of entry, resulting in the most rapid onset of toxic effects. Absorption also occurs rapidly by ingestion of the cyanide salts, KCN and NaCN, by dermal exposure to HCN, aqueous solutions of HCN (hydrocyanic acid), KCN and NaCN.

Several studies reported quantitative absorption data in animals and humans (Gettler and Baine, 1938; Liebowitz and Schwartz, 1948; Lendle, 1964; McNamara, 1976). These studies suggest that the absolute amount absorbed was relatively constant and not related to the time between administration and death.

The main site of cyanide detoxification is in the liver via the enzymatic action of rhodanese (the detoxifying enzyme in the liver) to form the less toxic thiocyanate. Differences among species with respect to distribution of rhodanese may account in part for their differences in sensitivity to cyanide. Dogs were found to have very low levels of rhodanese. Monkeys, rabbits and rats had hepatic levels ~10-18 times higher than dogs (Himwich and Saunders, 1948). Total levels of the enzyme in all tissues, except the adrenals, were lower in dogs than in the other species. Levels of rhodanese in other species, including humans, have not been reported.

The major route of cyanide elimination from the body is via urinary excretion of thiocyanate. Rats eliminated 80% of subcutaneously-injected cyanide as thiocyanate in the urine, while 15% was eliminated as urinary 2-imino-4-thiazolidine carboxylic acid (Wood and Cooley, 1956). A man who had ingested 3-5 g (at least 1.2 g HCN was present in blood 3 hours later) eliminated a total of 237 mg thiocyanate in 72-hour urine (Liebowitz and Schwartz, 1948).

<u>Short-Term Exposure</u>. In animals, acute LD_{50} values for cyanide range from ~0.91 mg CN^-/kg by intramuscular administration in rabbits, to 6.3 mg CN^-/kg by oral ingestion in mice. Acute LD_{50} values are summarized in Table VIII-1. For comparison purposes all doses are presented as the dose of the administered compound as well as the equivalent dose of cyanide ion (CN^-).

A dermal LD_{50} for HCN in humans has been estimated to be 100 mg/kg (DiPalma, 1971).

TABLE VIII-1

Acute LD_{50} Values for Cyanides

Route	Administered Compound	Species/Strain	Age/Weight (g)	Sex/No.	Vehicle	LD_{50} (mg/kg bw)	LD_{50} (mg CN^-/kg bw)	Comment	Reference
Oral	KCN	rat/Sherman	adult/NR	M/50	water	10	4.0	The rats died within 1 hour. LD_1 was 5.9 mg KCN/kg bw (2.4 mg CN^-/kg bw).	Hayes, 1967; Gaines, 1969
Oral	KCN	mouse/Swiss-Webster	NR/NR	NR/20/group	water	10.8	4.3	The compound was administered in a volume corresponding to 5% bw.	Ferguson, 1962
Oral	KCN	mouse/swiss Webster	NR/NR	NR/20/group	water	15.8	6.3	The volume corresponded to 1% bw.	Ferguson, 1962
Oral	KCN	mouse/Swiss-Webster	NR/18-20	M/10/group	NR	8.5	3.4	The LD_{50} is for mice dying up to 24 hours after dose. Signs of toxicity appeared in 30-90 seconds. After a short period of convulsions, mice were moribund; many at doses below LD_{50} appeared dead but recovered.	Sheehy and Way, 1968
Ocular	HCN	rabbit/NR	NR/NR	NR/NR	see comment	1.05	1.01	It was stated that the toxicity of NaCN was not different whether applied in solution, as a solid or mixed with an inert powder. Deaths occurred in 3-12 minutes.	Ballantyne, 1983
	KCN					7.87	3.15		
	NaCN					5.05	2.68		
i.p.	NaCN	mouse/CD-1	NR/25-30	M/NR	water	6.0	3.2	Pretreatment of mice with hydroxylamine, dimethylaminophenol or sodium nitrite increased the LD_{50} of NaCN.	Kruszyna et al., 1982
s.c.	KCN	mouse/Swiss-Webster	NR/18-22	M/10/group	saline	8.5	3.4	LD_{50} increased by pretreatment with sodium nitrite and sodium thiosulfate.	Way et al., 1966

TABLE VIII-1 (cont.)

Route	Administered Compound	Species/Strain	Age/Weight (g)	Sex/No.	Vehicle	LD_{50} (mg/kg bw)	LD_{50} (mg CN^-/kg bw)	Comment	Reference
s.c.	KCN	mouse/Swiss-Webster	NR/18-23	M/10/ group	saline	12.0	4.8	LD_{50} increased by pretreatment with cobaltous chloride alone or in combination with sodium nitrite and/or sodium thiosulfate.	Isom and May, 1973
s.c.	KCN	mouse/Swiss-Webster	NR/18-23	M/10/ group	saline	11.8	4.7	LD_{50} increased by pretreatment with sodium nitrite and sodium thiosulfate, enhanced by O_2.	May et al., 1972
s.c.	KCN	mouse/NR	NR/18-32	F/10/ group	saline	6.02 2.86	2.41 1.14	At 23-25°C At 4°C	Streicher, 1951
s.c.	NaCN	dog/NR	NR/NR	NR/16	NR	5.36	2.84	LD_{50} increased by immediate post-treatment administration of sodium thiosulfate, amyl nitrite, sodium nitrite and combinations of antidotes.	Chen and Rose, 1952
i.m.	HCN	rabbit/NR	adult/NR	M/10-15/ group	water	1.50	1.44	Increased rate of breathing, vociferation, uncoordinated movement of head, ataxia, tremors, respiratory and cardiac arrest.	Ballantyne et al., 1971, 1972
				F/10-15/ group		0.95	0.91		
i.m.	KCN	rabbit/NR	adult/NR	M/10-15/ group	water	3.06	1.22	Same as above, the rabbits lived longer following KCN rather than HCN administration.	Ballantyne et al., 1971, 1972
				F/10-15/ group		3.27	1.31		

NR = Not reported

i.p. = intraperitoneal; s.c. = subcutaneous; i.m. = intramuscular

Most of the studies on the health effects of oral exposure to cyanides in animals have been conducted using KCN, although NaCN has also been administered in some studies. Few studies on acute oral toxicity of cyanides providing dose-response data were encountered. Oral (via gastric intubation) doses of KCN equivalent to 4.0 and 6.0 mg CN^-/kg were fatal to rats and mice, respectively (Ferguson, 1962). Increasing the volume of water in which the dose was administered had the effect of increasing the incidence of death in the animals. Data are not available that indicate an effect of vehicle volume at sublethal doses.

Of 8 guinea pigs administered a single oral dose of KCN equivalent to 3.2 mg CN^-/kg, 5 had no signs of toxicity; the other 3 had slight tremors from which they recovered within 15 minutes (Basu, 1983). It should be noted that this low effect level in guinea pigs is equal to the LD_{50} in mice.

Rats can tolerate higher doses of cyanide when administered in the diet than when administered orally by gavage. The GI absorption following intermittent ingestion of low doses of cyanide over a day may allow for sufficient detoxication to occur (see Chapter V). Sherman rats survived 25 daily doses of 4 mg CN^-/kg bw when given as KCN mixed with the food (Hayes, 1967). This daily dose is equal to the acute oral LD_{50} in aqueous vehicle in the same strain of rats.

Male Sprague-Dawley rats (70 g, 7 rats/group) tolerated KCN at a level of 0.08 mg CN^-/mℓ when KCN was dissolved in the drinking water for 21 days (Palmer and Olson, 1979). This dose can be calculated as 12 mg CN^-/kg/day by assuming that a young growing rat consumes an amount of

water equal to ~15% of its body weight each day. The ingestion of water by rats is intermittent, thus GI absorption is intermittent, as in the case of dietary doses. There was no effect on body weight gain, but there was a significant increase in the average liver weight when compared with controls ($p<0.05$). When the KCN was mixed in the food at a concentration equivalent to 80 mg CN^-/kg feed, a daily dose of 8 mg CN^-/kg bw (calculated by assuming that a young growing rat, in a subchronic study, consumes an amount of food equal to ~10% of its body weight per day), did not result in increased liver weight. Potassium cyanide in the diet may have been more slowly absorbed than KCN in water due to physical interference with absorption or possible influence of chemical interactions of cyanide with components of the food. The Sprague-Dawley rats used in this study (Palmer and Olson, 1979) were not examined histologically, but African giant rats (Cricetomys gambianus, Waterhouse), maintained on diets containing 1000 mg CN^-/kg diet (daily dose of 36 mg CN^-/kg bw) for 84 days did not have histopathologic lesions in the thymus, kidney, liver and spleen (Tewe, 1982). Thus, a daily dose of 8 mg CN^-/kg bw administered orally in food (Palmer and Olson, 1979) is a NOAEL for increased liver weight in rats.

In a dietary study conducted by the American Cyanamid Co. (1959), three male and two female beagle dogs received NaCN mixed with food for 30-32 days. One male and one female dog served as controls. Each meal, which the dogs consumed completely, contained 24 mg CN^-. The average weight of the dogs was 7.6 kg; therefore, the daily dose was ~3 mg CN^-/kg bw. There were no effects during the dietary study on body weight, hemoglobin content, hematocrit or differential white cell count. No treatment-related histopathologic lesions or organ-to-body weight ratio changes were observed during exhaustive histologic examinations. A single gavage dose at this

level in dogs resulted in death in another study (Gettler and Baine, 1938). Dogs receiving 1.5, 4.3 and 8.1 mg CN^-/kg bw as single oral doses of KCN died in 155, 21 and 8 minutes, respectively. In the American Cyanamid study the dogs tolerated the higher dose because absorption may have been slowed due to the presence of food in the stomach.

Isom et al. (1982) studied the dose-response relationship of intraperitoneally administered KCN in male Swiss-Webster mice (4/group). The dose-dependent toxic signs included agitation, uncoordinated movement, gasping, irregular breathing, convulsion, respiratory arrest and death. At doses of KCN equivalent to 0.4 and 0.8 mg CN^-/kg bw, these signs were "barely detectable.". At doses of 1.2, 1.6 and 2.0 mg CN^-/kg bw, the mice started gasping, and had irregular respiration and convulsions within 2-3 minutes of cyanide administration. At 2.4 mg CN^-/kg bw, the toxic signs were more severe and 20% of the mice died. At doses >2.4 mg CN^-/kg bw (not specified), all the mice died within 4 minutes. Assuming that "barely detectable" means that the mice were slightly agitated and breathed slightly more rapidly, 0.8 mg CN^-/kg bw may represent a LOAEL and 0.4 mg CN^-/kg bw probably represents a NOAEL for intraperitoneally administered cyanide in mice. Because cyanide is rapidly absorbed from the GI tract and is rapidly detoxified on first passage through the liver, the effects of intraperitoneal administration may well be equivalent to the effects of oral administration. Data shown in Table VIII-1 indicate that the intraperitoneal LD_{50} and the oral LD_{50} in mice (3.2 and 3.4 mg CN^-/kg bw, respectively) are essentially equivalent.

The administration of NaCN by osmotic minipumps, implanted subcutaneously in pregnant hamsters on days 6-9 of gestation, resulted in severe teratogenic effects (Doherty et al., 1982). The pumps delivered NaCN at rates of ~0.1275 mmole NaCN/kg bw or a daily dose of 150 mg NaCN/kg bw (80 mg CN^-/kg bw). This study has little relevance to the risk assessment due to the high dose of cyanide and the method of administration, which was chosen in order to mimic the slow release of cyanide from nitriles during in vivo metabolism.

In an acute study, a man and a dog were exposed to HCN gas (between 500 and 625 ppm) simultaneously in an inhalation chamber (Barcroft, 1931). The man mimicked the movements of the dog, to approximate the same degree of activity, until the dog collapsed and went into convulsion. The man then exited the chamber with no symptoms except mild nausea and a sense of confusion. The dog was removed from the chamber and was believed to be dead, but subsequently recovered.

Information on the acute effects of ingestion of cyanides in humans has been obtained from case reports of suicide attempts. For HCN taken orally, absorbed doses of 0.5-3.5 mg HCN/kg bw (0.48-3.4 mg CN^-/kg bw) are fatal (Gettler and St. George, 1934; Gosselin et al., 1976; Ermans et al., 1972). Lethal doses of NaCN or KCN were reported to range from 50-200 mg or 0.7-2.9 mg/kg bw for a 70 kg human (~0.3-1.5 mg CN^-/kg bw) (Hartung, 1982; DiPalma, 1971). However, a case was reported in which a man recovered following ingestion of 3-5 g of KCN (~15 mg CN^-/kg bw was absorbed) (Liebowitz and Schwartz, 1948).

Bodansky and Levy (1923) administered cyanide orally in a capsule to 25 human subjects on 2 consecutive days. The capsules contained 15 mg of KCNS on the first day and 10 mg KCN on the second day. Assuming that the average body weight of humans is 70 kg, the doses were 0.21 mg KCNS and 0.14 mg KCN/kg bw or 0.06 mg CN^-/kg bw. The subjects effectively detoxified these doses as determined from measurements of thiocyanate levels in saliva. The subjects did not complain of any symptoms. While a daily dose of 0.06 mg CN^-/kg bw thus exerted no acute oral effect in humans, it does not necessarily represent the maximum no-effect level.

Long-Term Exposure. In a 2-year dietary study, Carworth Farms rats (10 animals/sex/group) were administered diets fumigated with HCN (Howard and Hanzal, 1955). Diets contained 0, 100 or 300 mg HCN/kg. Results of the residue analyses (done to measure evaporation loss) indicated that in 2 days the dietary concentration of HCN fell, on an average, from 100 to 51.9 mg/kg diet for the low-dose level and from 300 to 80.1 mg/kg diet for the high-dose level. The average daily low and high concentrations were ~76 and 190 mg HCN/kg diet (73 and 183 mg CN^-/kg diet), repectively. From data on reported food consumption and body weight, the estimated daily doses were 3.6 and 4.6 mg CN^-/kg bw for the low-dose males and females, respectively, and 7.5 and 10.8 mg CN^-/kg bw for the high-dose males and females, respectively. There were no treatment-related effects on growth rate, no gross signs of toxicity, nor any effects on organ-to-body weight ratios for liver, kidney, spleen, brain, heart, adrenals or gonads. There were no histopathologic lesions of the heart, lung, liver, spleen, stomach, intestines, kidney, adrenal, testes, uterus, ovary, cerebrum or cerebellum. The number of animals histologically examined was not specified, but it was stated as a

"representative" number. Thus, the daily doses of 3.6 and 7.5 mg CN^-/kg bw (for males) and 4.6 and 10.8 mg CN^-/kg bw (for females) appear to be no-adverse-effect levels for 2-year cyanide feeding to rats.

Philbrick et al. (1979) fed KCN in the diet to a group of 10 male rats (strain not specified) for 11.5 months at a concentration of 1500 mg KCN/kg diet (600 mg CN^-/kg diet). Another group of 10 rats served as the control. Assuming that a rat in a long-term feeding study consumes a quantity of food equal to ~5% of its body weight, the dietary level corresponds to a daily dose of ~30 mg CN^-/kg bw. When compared with controls, the treated rats had a 40% reduction in mean body weight gain, 53% decrease in plasma thyroxine levels and 68% decrease in thyroxine secretion rates. There were no definitive histopathologic lesions to thyroid, sciatic, optic or other neural tissue. Mild degenerative changes in the myelin of the spinal cord were observed. Thus, a daily dose of 30 mg CN^-/kg bw was a LOAEL for dietary cyanide in rats.

Hertting et al. (1960) treated three dogs orally with NaCN in a gelatin capsule for up to 14.5 months. There was one control dog. Treatments were as follows: dog I received a daily dose of 0.27 mg CN^-/kg bw for 13.5 months; dog II received a daily dose of 0.53 mg CN^-/kg bw for 16 weeks and then a daily dose of 2.2 mg CN^-/kg bw for ~10.5 months longer; dog III received a daily dose of 1.1 mg CN^-/kg bw for 14.5 months. When daily doses >0.53 mg CN^-/kg bw were administered, there were signs of acute intoxication immediately after dosing; however, recovery occurred in <0.5 hours. In all treated dogs, histologic examination revealed degeneration of CNS ganglion cells and necrosis and inflammation of Purkinje cells of the

cerebellum. Thus, a no-effect level in dogs cannot be identified. The finding of histologic damage to the CNS of dogs following subchronic oral dosing with low levels of cyanide may reflect the greater sensitivity of dogs to cyanide poisoning. Dogs have a decreased ability to detoxify cyanide since levels of rhodanese are very low in the dog, especially in the liver, when compared with rats, rabbits and monkeys (Himwich and Saunders, 1948).

No information was available on chronic oral exposure of humans to HCN, KCN or NaCN. Chronic ingestion of cyanogenic plants, such as cassava, a root food that forms the staple diet of people living in certain tropical areas of Africa, has been associated with the etiology of neuropathies and thyroid disorders (Osuntokun, 1972; Delange and Ermans, 1971). Exposure, however, was to linamarin and lotaustralin, cyanogenic glycosides contained in cassava, in addition to cyanide per se. Other complicating factors, such as vitamin B_{12}, protein and iodine deficiencies also played a role (Wilson, 1983).

Quantification of Noncarcinogenic Effects

Derivation of 1-Day HA. No suitable study was identified from which to calculate a 1-day health advisory. The results of Bodansky and Levy (1923) refer to one arbitrarily chosen low dose of cyanide and provide little indication of permissible loads. Indeed, if these results were used, the resultant 1-day HA would actually fall below those criteria estimated from the more extensive results for more prolonged exposure. Thus, it is recommended that the 1-day HA be set at the same level as the 10-day HA.

Derivation of 10-Day HA. The study by Palmer and Olson (1979) could be used for the derivation of the 10-day HA. Using this data, the 10-day HA for a 10 kg child could be calculated as follows:

$$\text{10-day HA} = \frac{(8 \text{ mg CN}^-/\text{kg bw/day})\ (10 \text{ kg})}{(1\ \ell/\text{day})\ (100)\ (5)}$$

$$= 0.16 \text{ mg CN}^-/\ell \text{ (rounded 0.2 mg CN}^-/\ell)$$

where:

8 mg CN^-/kg bw/day	=	NOAEL for effects on liver weight or body weight in rats following a 21 day exposure via food (Palmer and Olson, 1979)
10 kg	=	assumed weight of a child
1 ℓ/day	=	assumed water consumption by a child
100	=	uncertainty factor. An uncertainty factor of 100 is selected based upon U.S. EPA (1991) and NAS/ODW guidelines in which a NOAEL from an animal study is used.
5	=	modifying factor. A modifying factor of 5 is selected because of the possible problems associated with the use of a dietary study to estimate a drinking water criterion.

However, it is recommended that the DWEL of 0.7 mg/ℓ be used for the child 1-day and 10-day HAs. The NOAEL of 10.8 mg/kg/day used for the DWEL comes from a study where the animals were exposed for 2 years; the 8 mg/kg/day NOAEL is from a study where animals were exposed for only 21 days. Therefore, because there were no adverse effects when the exposure was for a longer time at higher levels, the DWEL of 0.7 mg/ℓ would be protective for the child 1-day and 10-day exposures also. The corresponding 1-day and 10-day HAs for a 10 kg child consuming 1 ℓ of water would be 0.2 mg/ℓ.

Derivation of Longer-Term HA. The available data are insufficient to develop longer-term HAs for cyanide. It is recommended that the DWEL of 0.7 mg/ℓ be used as the longer-term HA for the 70 kg adult and the modified DWEL of 0.2 mg/ℓ (adjusted for a 10 kg child) be used for the longer-term HA for a 10 kg child.

Assessment of Lifetime Exposure and Derivation of a DWEL. The study by Howard and Hanzal (1955) has been selected to serve as the basis for an RfD and lifetime DWEL because it is the only long-term study for which a NOAEL was identified. In this study female rats (Cartworth Farm) exposed to 4.6 or 10.8 mg CN/kg bw for 2 years experienced no adverse effects. An RfD of 0.02 mg/kg/day was verified (verification date 03/23/88) by the Agency-wide RfD Workgroup using the 10.8 mg/kg bw dose as a NOAEL (U.S. EPA, 1991).

In deriving daily doses, certain assumptions and decisions are made as follows:

1. The HCN concentrations reported by the authors are converted to CN^- concentrations on a molecular weight basis.

2. The body weight data presented by the authors are used rather than employing the assumption that a rat weighs an average of 350 g over its lifespan (45 FR 79352, November 28, 1980). The average body weights for all exposure groups are estimated from the growth charts using linear interpolation to estimate missing data points.

3. The food consumption data presented by the authors are used rather than assuming an average food consumption for all dose groups.

4. Average CN^- concentrations over the life of the experiment are estimated from the data presented by the authors and by assuming that the rate of loss of HCN can be described by a first-order process.

5. It is inappropriate to use a net absorption coefficient to account for presumed differences in the absorption of CN^- in food vs. drinking water. However, there is a given measure of uncertainty associated with the use of a dietary study to estimate a drinking water criterion, thus, an additional 5-fold uncertainty factor is used.

Thus, the data are:

Dose Group	Average Concentration (mg CN^-/kg diet)	Average Food Consumption (av. g/rat/day)	Body Weight (g)
Low male	73	19.46	390
High male	183	18.50	394
Low female	73	14.69	232
High female	183	17.24	255

Average daily doses are calculated as follows:

low dose male:
(73 mg CN^-/kg diet x 0.01946 kg)/0.390 kg = 3.6 mg CN^-/kg bw/day

high dose male:
(183 mg CN^-/kg diet x 0.01850 kg)/0.394 kg = 7.5 mg CN^-/kg bw/day

low dose female:
(73 mg CN^-/kg diet x 0.01469 kg)/0.232 kg = 4.6 mg CN^-/kg bw/day

high dose female:
(183 mg CN^-/kg diet x 0.01724 kg)/0.255 kg = 10.8 mg CN^-/kg bw/day

<u>Step 1 - RfD Derivation</u>

$$RfD = \frac{(10.8 \text{ mg CN /kg bw})}{(100)\ (5)}$$

= 0.022 mg/kg/day (rounded to 0.02 mg/kg/day)

where: 10.8 mg CN^-/kg bw = NOAEL for absence of clinical and histologic effects in a 2-year dietary study in rats (Howard and Hanzal, 1955).

100	= uncertainty factor. An uncertainty factor of 100 is selected based upon U.S. EPA (1991) and NAS/ODW guidelines in which a NOAEL from an animal study is used.
5	= modifying factor. A modifying factor of 5 is selected due to possible problems associated with the use of a dietary study to estimate a drinking water criterion.

Step 2 - DWEL Derivation

$$DWEL = \frac{0.02\ mg/kg/day \times 70\ kg}{2\ \ell/day} = 0.7\ mg/\ell\ (700\ \mu g/\ell)$$

where:

0.02 mg/kg/day = RfD

2 ℓ/day = assumed water consumption by an adult

70 kg = assumed body weight of an adult

This DWEL calculation assumes 100% of the human exposure derives from drinking water. The DWEL may be modified upon the availability of relative source contribution data providing human exposure estimates from food, air and possibly the occupational environment. The ultimate goal is to establish a DWEL so that human exposure from all sources does not exceed the RfD.

Carcinogenic Effects

Potassium cyanide was negative for reverse mutation in five strains of *Salmonella typhimurium*, with and without metabolic activation (De Flora, 1981). A marginally positive response was obtained in strain TA100 with HCN gas (Kushi et al., 1983). Cyanide was also negative in a modified rec assay in *Bacillus subtilis* (Karube et al., 1981).

Quantification of Carcinogenic Effects

Studies regarding the carcinogenicity of cyanide were not located in the available literature. The lack of information concerning the carcinogenic potential of cyanide precludes any further analysis of carcinogenic potential. The International Agency for Research on Cancer (IARC) has not evaluated the carcinogenic potential of cyanide (WHO, 1982). Applying the criteria described in U.S. EPA (1986) Guidelines for Carcinogen Risk Assessment, cyanide has been classified in Group D: not classifiable in this case due to a lack of bioassay studies. This category is for agents with inadequate or insufficient human and animal studies for carcinogenicity.

Existing Guidelines, Recommendations and Standards

The ambient water quality criterion has been proposed at 3.77 mg CN^-/ℓ assuming that a 70 kg human consumes 2 ℓ of water and 6.5 g of fish per day with a bioconcentration factor of 1.0 (U.S. EPA, 1982).

The U.S. Public Health Service (1962) recommended that concentrations of cyanide in water supplies not exceed 0.2 mg CN^-/ℓ in order to protect human health. The value appears to be calculated from the TLV, which gives a water level of ~19 mg/ℓ followed by the application of a 100-fold uncertainty factor. This uncertainty factor was used due to the steep dose-effect relationship observed for cyanide when body levels exceed detoxification capacity. The U.S. Public Health Service (1962) also recommended that concentrations in drinking water be kept below 0.01 mg CN^-/ℓ since this level or lower can be achieved by proper treatment.

A TLV for alkali cyanides in workroom air of 5 mg CN^-/m^3 is recommended by the ACGIH (1980) based upon irritation to the respiratory system as well as protection from the effects of chronic exposure. The TLV of 5 mg CN^-/m^3 was also recommended by NIOSH (1976) and adopted by OSHA (1981).

Special Group at Risk

The only special considerations identified for cyanide exposure were the potential effects to high risk subpopulations. Infants and fetuses may be at higher risk to cyanide. Pregnant hamsters exposed to high doses ($\geq$78 mg/kg) of cyanide experienced severe teratogenic effects (Doherty et al., 1982). Andrews (1973) found that pregnant women who smoke have higher levels of plasma thiocyanate in blood than nonsmokers. It has been suggested that cyanide exposure may result in low body weight infants.

A metabolic disturbance in the conversion of cyanide to thiocyanate and vitamin B_{12} deficiency has been associated with such conditions as tobacco amblyopia and Leber's hereditary optic atrophy in persons who are exposed to excess cyanide in tobacco smoke (Wilson, 1983). Iodine deficiency, along with excess exposure to cyanide, may be involved in the etiology of such thyroid disorders as goiter and cretinism (Delange and Ermans, 1971; Ermans et al., 1972). Protein deficiencies, vitamin B_{12} and riboflavin deficiencies may subject people in the tropics who eat cassava, a cyanogenic plant, to increased risks of tropical neuropathies (Osuntokun, 1972; Osuntokun et al., 1969; Makene and Wilson, 1972). Therefore, individuals with a metabolic defect in the rhodanese system, a vitamin B_{12} deficiency or defective B_{12} metabolism, iodine deficiency, protein deficiency as well as fetuses exposed *in utero*, are at a higher risk to the toxic effects of cyanide exposure than the general population.

Additional groups at a higher risk may be strict vegetarians who may become vitamin B_{12} deficient and people unable to smell cyanide.

Summary

The recommended values for the 1-day and 10-day HAs for the child and longer-term HAs for both adults and children and the DWEL are summarized in Table VIII-2.

TABLE VIII-2

Summary of Quantification of Toxicologic Effects

	Drinking Water Concentration (mg CN^-/ℓ)		
	10 kg Child	70 kg Adult	Reference[a]
1-day Health Advisory	b/		--
10-day Health Advisory	0.2[c]		Howard and Hanzal, 1955
DWEL	d/	0.7	Howard and Hanzal, 1955

[a]Reference for NOAEL used in calculating HA or RfD values

[b]No suitable study identified; it is recommended that the 1-day HA be set at the same levels as the 10-day HA.

[c]Calculated from the NOAEL used for the DWEL.

[d]Not calculated because the child will not remain in the high risk population for a lifetime.

IX. REFERENCES

ACGIH (American Conference of Governmental Industrial Hygienists). 1980. Documentation of the threshold limit values for substances in workroom air, 4th ed., with supplements through 1981. Cincinnati, OH. p. 109-110.

American Cyanamid Co. 1959. Report on sodium cyanide: 30-day repeated feedings to dogs. Central Med. Dept. Report Number 59-14.

Andrews, J. 1973. Thiocyanate and smoking in pregnancy. J. Obstet. Gynaecol. Br. Common. 80: 810-814.

Ansell, M. and F.S. Lewis. 1970. A review of cyanide concentrations found in human organs: A survey of literature concerning cyanide metabolism, "normal," non-fatal and fatal body cyanide levels. J. Forensic Med. 17: 148-155.

Ballantyne, B. 1983. Acute systemic toxicity of cyanides by topical application to the eye. J. Toxicol. 2: 119-129.

Ballantyne, B., J. Bright, D.W. Swanston and P. Williams. 1971. Toxicity of cyanides given by intramuscular injection. Br. J. Pharmacol. 41: 423-424.

Ballantyne, B., J. Bright, D.W. Swanston and P. Williams. 1972. Toxicity and distribution of free cyanides given intramuscularly. Med. Sci. Law. 12: 209-219.

Barcroft, J. 1931. The toxicity of atmospheres containing hydrocyanic acid gas. J. Hyg. 31: 1-34.

Basu, T.K. 1983. High-dose ascorbic acid decreases detoxification of cyanide derived from amygdalin (laetrile): studies in guinea pigs. Can. J. Physiol. Pharmacol. 61(11): 1426-1430.

Bodansky, M. and M.D. Levy. 1923. Some factors influencing the detoxification of cyanides in health and disease. Arch. Int. Med. 31: 373-389.

Boxer, G.E. and J.C. Rickards. 1951. Chemical determination of vitamin B_{12}. IV. Assay of vitamin B_{12} in multivitamin preparations and biological materials. Arch. Biochem. 30: 392-401.

Boxer, G.E. and J.C. Rickards. 1952a. Determination of traces of hydrogen cyanide in respiratory air. Arch. Biochem. Biophys. 39: 287-291.

Boxer, G.E. and J.C. Rickards. 1952b. Studies on the metabolism of the carbon of cyanide and thiocyanate. Arch. Biochem. 39: 7-26.

Brink, N.G., F.A. Kuehl and K. Folkers. 1950. Vitamin B_{12}: The identification of vitamin B_{12} as a cyano-cobal coordination complex. Science. 112: 354.

Brown, W.E., C.D. Wood and A.N. Smith. 1960. Sodium cyanide as a cancer chemotherapeutic agent. Am. J. Obstet. Gynecol. 80: 907-918.

Burrows, G.E., D.H.W. Liu and J.L. Way. 1973. Effect of oxygen on cyanide intoxication: V. Physiologic effects. J. Pharmacol. Exp. Ther. 184: 739-748.

Callahan, M.A., M.W. Slimak, N.W. Gabel, et al. 1979. Water-Related Environmental Fate of 129 Priority Pollutants, Vol. 1. Office of Water Planning and Standards, Office of Water and Waste Management, U.S. EPA, Washington, DC. EPA-440/4-79/029a.

Carmelo, S. 1955. New contributions to the study of subacute-chronic hydrocyanic acid intoxication in man. Rass Med. Ind. 24: 254-271. (Cited in NIOSH, 1976)

Chandra, H., B.N. Gupta, S.K. Bhargava, S.H. Clerk and P.N. Mahendra. 1980. Chronic cyanide exposure - a biological and industrial hygiene study. J. Anal. Toxicol. 4: 161-165.

Chen, K.K. and C.L. Rose. 1952. Nitrite and thiosulfate therapy in cyanide poisoning. J. Am. Med. Assoc. 149: 113-119.

Cotton, F.A. and G. Wilkinson. 1980. Advanced Inorganic Chemistry, 4th ed. John Wiley and Sons, NY. p. 1396

Crampton, R.E., I.F. Gaunt, R. Harris et al. 1979. Effect of low cobalamin diet and chronic cyanide toxicity in baboons. Toxicology. 12: 221-234.

Davison, V. 1969. Cyanide poisoning. Kelocyanor -- A new treatment as seen at Imperial Chemical Industries, Billingham. Occup. Health. 21: 306-308.

de Duve, C., B.C. Pressman, R. Gianetto, R. Wattiaux and F. Appelmans. 1955. Tissue fractionation studies: 6. Intracellular distribution patterns of enzymes in rat-liver tissue. Biochem. J. 60: 604-617.

De Flora, S. 1981. Study of 106 organic and inorganic compounds in the Salmonella/microsome test. Carcinogenesis. 2(4): 283-298.

De Flora, S., A. Camoirano, P. Zanacchi and C. Bennicelli. 1984. Mutagenicity testing with TA97 and TA102 of 30 DNA-damaging compounds, negative with other Salmonella strains. Mutat. Res. 134(2-3): 159-165.

Delange, F. and A.M. Ermans. 1971. Role of a dietary goitrogen in the etiology of endemic goiter on Idjwi Island. Am. J. Clin. Nutr. 24: 1354-1360.

Dinca, C. L. Pod and I. Galetariu. 1972. Considerations on leukocytic oxidative enzyme changes in subjects exposed to prolonged action of cyanhydric acid in industry. Med. Int. 24: 1385-1392. (Cited in NIOSH, 1976)

DiPalma, J.R., Ed. 1971. Noxious gases and vapors: I. Carbon monoxide, cyanides, methemoglobin and sulfhemoglobin. In: Drill's Pharmacology in Medicine. McGraw-Hill Book Co., NY. p. 1189-1205.

Doherty, P.A., V.H. Ferm and R.P. Smith. 1982. Congenital malformations induced by infusion of sodium cyanide in the golden hamster. Toxicol. Appl. Pharmacol. 64: 456-464.

Drinker, P. 1932. Hydrocyanic acid gas poisoning by absorption through the skin. J. Ind. Hyg. 14: 1-2.

El Ghawabi, S.H., M.A. Gaafar, A.A. El-Saharti, S.H. Almed, K.K. Malash and R. Fares. 1975. Chronic cyanide exposure: a clinical, radioisotope and laboratory study. Br. J. Ind. Med. 32: 215-219.

Ermans, A.M., F. Delange, M. Van der Velden and J. Kinthaert. 1972. Possible role of cyanide and thiocyanate in the etiology of endemic cretinism. Adv. Exp. Med. Biol. 30: 455-486.

Farooqui, M.Y.H. and A.E. Ahmed. 1982. Molecular interaction of acrylonitrile and potassium cyanide with rat blood. Chem. Biol. Interact. 38: 145-159.

Feldstein, M. and N.C. Klendshoj. 1954. The determination of cyanide in biologic fluids by microdiffusion analysis. J. Lab Clin. Med. 44: 166-170.

Ferguson, H.C. 1962. Dilution of dose and acute oral toxicity. Toxicol. Appl. Pharmacol. 4: 759-762.

Ferraro, A. 1933. Experimental toxic encephalomyelopathy. Psychiatr. Q. 7: 267-283.

Finck, P.A. 1969. Postmortem distribution studies of cyanide. Med. Am. DC. 38: 357-358.

Foulds, W.S., J.S. Cant, I.A. Chisholm, J. Bronte-Stewart and J. Wilson. 1968. Hydroxocobalamin in the treatment of leber's hereditary optic atrophy. Lancet. 1: 896-897.

Frank, S.N. and A-J. Bard. 1977. Heterogeneous photocatalytic oxidation of cyanide ion in agueous solutions of TiO_2 powder. J. Am. Chem. Soc. 99: 303-304.

Frankenberg, L. 1980. Enzyme therapy in cyanide poisoning: effect of rhodanese and sulfur compounds. Arch. Toxicol. 45(4): 315-323.

Friedberg, K.D. and H.A. Schwarzkopf. 1969. Blausaureexhalation bei der cyanidvergiftung (The exhalation of hydrocyanic acid in cyanide poisoning). Arch. Toxicol. 24: 235-248.

Gaines, T.B. 1969. Acute toxicity of pesticides. Toxicol. Appl. Pharmacol. 14: 515-534.

Gettler, A.O. and J.O. Baine. 1938. The toxicology of cyanide. Am. J. Med. Sci. 195: 182-198.

Gettler, A.O. and A.V. St. George. 1934. Cyanide poisoning. J. Clin. Pathol. 4: 429-437.

Goldstein, F. and F. Rieders. 1951. Formation of cyanide in dog and man following administration of thiocyanate. Am. J. Physiol. 167: 47-51.

Goldstein, F. and F. Rieders. 1953. Conversion of thiocyanate to cyanide by an erythrocytic enzyme. Am. J. Physiol. 173: 287-280.

Gosselin, R.E., H.C. Hodge, R.P. Smith and M.N. Gleason. 1976. Clinical Toxicology of Commercial Products, 4th ed. Williams and Wilkins Co., Baltimore, MD. p. 105-112.

Habig, W.H., J.H. Keen and W.B. Jakoby. 1975. Glutathione s-transferase in the formation of cyanide from organic thiocyanates and as an organic nitrate reductase. Biochem. Biophys Res. Commun. 2: 501-506.

Hansch, C. and A.J. Leo. 1979. In: Substituent Constants for Correlation Analysis in Chemistry and Biology. John Wiley and Sons, NY. p. 172.

Hardy, H.L. W.M. Jeffries, M.M. Wesserman and W.R. Waddell. 1950. Thiocyanate effect following industrial cyanide exposure -- report of two cases. N. Engl. J. Med. 242: 968-972.

Hartung, R. 1982. Cyanides and Nitriles. In: Patty's Industrial Hygiene and Toxicology, 3rd revised ed., Vol. 2C, G.D. Clayton and F.E. Clayton, Ed. John Wiley and Sons, Inc., NY. p. 4845-4900.

Hayes, W.T. 1967. The 90-dose LD_{50} and a chronicity factor as measures of toxicity. Toxicol. Appl. Pharmacol. 11: 327-335.

Haymaker, W., A.M. Ginzler and R.L. Ferguson. 1952. Residual neuropathological effects of cyanide poisoning -- a study of central nervous system of 23 dogs exposed to cyanide compounds. Mil. Surg. 111: 231-246.

Herbert, V. 1975. Drugs effective in megaloblastic anemias. In: The Pharmacological Basis of Therapeutics, 5th ed., L.S. Goodman and A. Gilman, Ed. McMillan Publishing Co., New York. p. 1332.

Hertting, G., O. Kraupp, E. Schnetz and S. Wieketich. 1960. Untersuchungen uber die Folgen einer chronischen Verabreichung akut toxischer Dosen von Natriumcyanid an Hunden. Acta. Pharmacol. Toxicol. 17: 27-43. (Ger.)

Himwich, W.A. and J.P. Saunders. 1948. Enzymatic conversion of cyanide to thiocyanate. Am. J. Physiol. 153: 348-354.

Howard, J.W. and R.F. Hanzal. 1955. Chronic toxicity for rats of food treated with hydrogen cyanide. J. Agric. Food Chem. 3: 325-329.

Hugod, Carl. 1981. Myocardial morphology in rabbits exposed to various gasphase constituents of tobacco smoke - an ultrastructural study. Atherosclerosis. 40: 181-190.

Hurst, E.W. 1940. Experimental demyelination of the central nervous system: 1. The encephalopathy produced by potassium cyanide. Aust. J. Exp. Biol. Med. Sci. 18: 201-223.

Isom, G. and J.L. Way. 1973. Cyanide intoxication: Protection with cobaltous chloride. Toxicol. Appl. Pharmacol. 24: 449-456.

Isom, G.E. and J.L. Way. 1974. Effect of oxygen on cyanide intoxication: VI. Reactivation of cyanide-inhibited glucose metabolism. J. Pharmacol. Exp. Ther. 189: 235-243.

Isom, G.E., G.E. Burrows and J.L. Way. 1982. Effect of oxygen on the antagonism of cyanide intoxication-cytochrome oxidase, in vivo. Toxicol. Appl. Pharmacol. 65(2): 250-256.

Johnson, J.D., T.L. Meisenheimer and G.E. Isom. 1986. Cyanide induced neurotoxicity: Role of neuronal calcium. Toxicol. Appl. Pharmacol. 84(3): 464-469.

Karube, I, T. Matsunaga, T. Nakahara, S. Suzuki and T. Kada. 1981. Preliminary screening of mutagens with a microbial sensor. Anal. Chem. 53(7): 1024-1026.

Kirk, R.L. and N.S. Stenhouse. 1953. Ability to smell solutions of potassium cyanide. Nature. 171: 698-699.

Knowles, E.L. and J.T.B. Bain. 1968. Medical cover required in large scale production of cyanides and hydrocyanic acid. Chem. Ind. 8: 232-235.

Kreutler, P.A., V. Varbanov, W. Goodman, G. Olaya and J.B. Stanbury. 1978. Interactions of protein deficiency, cyanide and thiocyanate on thyroid function in neonatal and adult rats. Am. J. Clin. Nutrit. 31: 282-289.

Kruszyna, R., H. Kruszyna and R.P. Smith. 1982. Comparison of hydroxylamine, 4-dimethylaminophenol and nitrite protection against cyanide poisoning in mice. Arch. Toxicol. 49: 191-202.

Kushi, A, T. Matsumoto and D. Yoshida. 1983. Mutagen from the gaseous phase of protein pyrolyzate. Agric. Biol. Chem. 47(9): 1979-1982.

Landahl, H.D. and R.G. Herrmann. 1950. Retention of vapors and gases in the human nose and lung. Arch. Ind. Hyg. Occup. Med. 1: 36-45.

Lang, K. 1933. Die Rhodanbildung im Tierkorper. Biochem. Zeitschrift. 259: 243-256. (Ger.)

Lendle, L. 1964. Wirkunsbedingungen von blausaure und schwefelwasserstoff und moglichkreiten der vergiftungsbehandlung. Jap. J. Pharmacol. 14: 215-224. (Ger.)

Leo, A., C. Hansch and D.A. Elkins. 1971. Partition coefficients and their uses. Chem. Rev. 71(6): 525-616.

Lessell, S. 1971. Experimental cyanide optic neuropathy. Arch. Ophthalmol. 86: 194-204.

Lessell. S. and T. Kuwabara. 1974. Fine structure of experimental cyanide optic neuropathy. Invest. Ophthalmol. 13: 748-756.

Levin, B.C., M. Paabo, J.L. Gurman and S.E. Harris. 1987. Effects of exposure to single or multiple combinations of the predominant toxic gases and low oxygen atmospheres produced in fires. Fund. Appl. Toxicol. 9(2): 236-250.

Liebowitz, D. and H. Schwartz. 1948. Cyanide poisoning: Report of a case with recovery. Am. J. Clin. Pathol. 18: 965-970.

Makene, W.J. and J. Wilson. 1972. Biochemical studies in Tanzanian patients with ataxic tropical neuropathy. J. Neurol. Neurosurg. Psychiatry. 35: 31-33.

Maliszewski, T.F. and D.E. Bass. 1955. 'True' and 'apparent' thiocyanate in body fluids of smokers and nonsmokers. J. Appl. Physiol. 8: 289-291.

Matijak-Schaper, M. and Y. Alarie. 1982. Toxicity of carbon monoxide, hydrogen cyanide and low oxygen. J. Combust. Toxicol. 9(2): 21-61.

McMillan, D.E. and A.C. Svoboda, IV. 1982. The role of erythrocytes in cyanide detoxification. J. Pharmacol. Exp. Ther. 221(1): 37-42.

McNamara, B.P. 1976. Estimates of the toxicity of hydrocyanic acid vapors in man. Edgewood Arsenal Technical Report EB-TR-76026, Dept. of the Army, August, 1976. (Cited in NIOSH, 1976)

Monekosso, G.L. and J. Wilson. 1966. Plasma thiocyanate and vitamin B_{12} in Nigerian patients with degenerative neurological disease. Lancet. 1: 1062-1064.

Moss, R.H., C.F. Jackson and J. Seiberlich. 1951. Toxicity of carbon monoxide and hydrogen cyanide gas mixtures -- a preliminary report. Arch. Ind. Hyg. Occup. Med. 4: 53-64.

Mushett, C.W., K.L. Kelley, G.E. Boxer and J.C. Rickards. 1952 Antidotal efficacy of vitamin $B_{12}a$ (hydroxo-cobalamin) in experimental cyanide poisoning. Proc. Soc. Exp. Biol. Med. 81: 234-237.

NAS (National Academy of Sciences). 1977. Drinking Water and Health. Vol. I, p. 19-63.

NAS (National Academy of Sciences). 1980. Drinking Water and Health. Vol. III, p. 25-67.

Nicholls, P. 1975. The effect of sulphide on cytochrome aa_3: isosteric and allosteric shifts of the reduced α peak. Biochem. Biophys. Acta. 396: 24-35.

NIOSH (National Institute for Occupational Safety and Health). 1976. Criteria for a Recommended Standard. Occupational Exposure to Hydrogen Cyanide and Cyanide Salts (NaCN, KCN and $Ca(CN)_2$). NIOSH Publ. No. 77-108. Dept. Health Educ. Welfare. U.S. Gov. Printing Off., Washington, DC.

NIOSH (National Institute for Occupational Safety and Health). 1981. Occupational Guildelines for Cyanide. NIOSH Publ. No. 81-123. Dept. of Human Health Services, Washington, DC.

NTP (National Toxicology Program). 1991. Toxicology Research and Testing Program. Management Status Report, dated 04/03/91.

O'Flaherty, J.E. and W.C. Thomas. 1982. The cardiotoxicity of hydrogen cyanide as a component of polymer pyrolysis smokes. Toxicol. Appl. Pharmacol. 63(3): 373-381.

Ohsy, S. Jalaludin, R.H. Davis and A.H. Sykes. 1987. Detoxification of cyanide in the chicken by conversion to thiocyanate. Com. Biochem. Physiol. 86(1): 129-133.

Okoh, P.N. 1983. Excretion of ^{14}C-labeled cyanide in rats exposed to chronic intake of potassium cyanide. Toxicol. Appl. Pharmacol. 70(2): 335-339.

Okoh, P.N. and G.A.J. Pitt. 1982. The metabolism of cyanide and the gastrointestinal circulation of the resulting thiocyanate under conditions of chronic cyanide intake in the rat. Can. J. Physiol. Pharmacol. 60(3): 381-386.

OSHA (Occupational Safety and Health Administration). 1981. General Industry OSHA Safety and Health Standards (29CFR 1910). OSHA 2206. U.S. Dept. Labor, Washington, DC. p. 633.

Osuntokun, B.O. 1968. An ataxic neuropathy in Nigeria. Brain. 91: 215-248.

Osuntokun, B.O. 1972. Chronic cyanide neurotoxicity and neuropathy in Nigerians. Plant Foods Hum. Nutr. 2: 215-266.

Osuntokun, B.O., G.L. Monekosso and J. Wilson. 1969. Relationship of a degenerative tropical neuropathy to diet, report of a field study. Br. Med. J. 1: 547-550.

Osuntokun, B.O., A. Aladetoyinbo and A.O. G. Adeuja. 1970. Free-cyanide levels in tropical atoxic neuropathy. Lancet. 2: 372-373.

Owasoyo, J.O. and C.A. Iramain. 1980. Acetylcholinesterase activity in rat brain: effect of acute cyanide intoxication. Toxicol. Lett. 6(1): 1-3.

Palmer, I. S. and O.E. Olson. 1979. Partial prevention by cyanide of selenium poisoning in rats. Biochem. Biophys. Res. Commun. 90(4): 1379-1386.

Perry, I.H. 1935. The effect of prolonged cyanide treatment on body and tumor growth in rats. Am. J. Cancer. 25: 592-598.

Pettigrew, A.R. and G.S. Fell. 1972. Simplified colorimetric determination of thiocyanate in biological fluids and its application to investigation of toxic amblyopias. Clin. Chem. 18: 996-1000.

Pettigrew, A.R. and G.S. Fell. 1973. Microdiffusion method for estimation of cyanide in whole blood and its application to the study of conversion of cyanide to thiocyanate. Clin. Chem. 19: 466-471.

Philbrick, D.J., J.B. Hopkins, D.C. Hill, J.C. Alexander and R.G. Thomson. 1979. Effects of prolonged cyanide and thiocyanate feeding in rats. J. Toxicol. Environ. Health. 5: 579-592.

Potter, A.L. 1950. The successful treatment of two recent cases of cyanide poisoning. Br. J. Ind. Med. 7: 125-130.

Purser, D.A., P. Grimshaw and K.R. Berrill. 1984. Intoxication by cyanide in fires: A study in monkeys using polyacrylonitrile. Arch. Environ. Health. 39(6): 394-400.

Radojicic, B. 1973. Determining thiocyanate in urine of workers exposed to cyanides. Arch. Hig. Rada. 24: 227-232. (Cited in NIOSH, 1976)

Raef, S.F, W.G. Characklis, M.A. Kessick and C.H. Ward. 1977a. Fate of cyanide and related compounds in aerobic microbial systems. II. Microbial degradation. Water Res. 11: 485-492.

Raef, S.F. W.G. Characklis, M.A. Kessick and C.H. Ward. 1977b. Fate of cyanide and related compounds in aerobic microbial systems. I. Chemical reaction with substrate and physical removal. Water Res. 11: 477-483.

Robinson, C.P., S.I. Baskin and D.R. Franz. 1985. The mechanism of action of cyanide on the rabbit aorta. J. Appl. Toxicol. 5(6): 372-377.

Saia, B., E. DeRosa, L. Galzigna. 1970. Remarks on the chronic poisoning from cyanide. Med. Lav. 61: 580-586. (Cited in NIOSH, 1976)

Sandberg, C.G. 1967. A case of chronic poisoning with potassium cyanide. Acta. Med. Scand. 181: 233-236.

Schwartz, C., R.L. Morgan, L. Ronald, L.M. Way and J.L. Way. 1979. Antagonism of cyanide intoxication with sodium pyruvate. Toxicol. Appl. Pharmacol. 50(3): 437-441.

Sheehy M. and J.L. Way. 1968. Effect of oxygen on cyanide intoxication III. Mithridate. J. Pharmacol. Exp. Ther. 161: 163-168.

Smith, R.P. and R.E. Gosselin. 1979. Hydrogen sulfide poisoning. J. Occup. Med. 21: 93-97.

Smith, R.P. and M.V. Olson. 1973. Drug-induced methemoglobinemia. Sem. Hematol. 10: 253-268.

Smith, A.D.M., S. Duckett and A.H. Waters. 1963. Neuropathological changes in chronic cyanide intoxication. Nature. 200: 179-181.

Smith, L., H. Kruszyna and R.P. Smith. 1977. The effect of methemoglobin on the inhibition of cytochrome c oxidase by cyanide, sulfide or azide. Biochem. Pharmacol. 26: 2247-2250.

Solomonson, L.P. 1981. Cyanide as a metabolic inhibitor. In: Cyanide in Biology, B. Vennesland, E.E. Conn and C. Knowles, Ed. Academic Press, London, England. p. 11-28.

Towill, L.E., J.S. Arury, B.L. Whitfield, E.B. Lewis, E.L. Galyan and A.S. Hammons. 1978. Reviews of the Environmental Effects of Pollutants: V. Cyanide. Health Effects Research Laboratory, Office of Research and Development, U.S. EPA, Cincinnati, OH. EPA-600/1-78-027. NTIS PB 289920.

Trautman, J.A. 1930. Methylene blue in the treatment of HCN gas poisoning. Public Health Report. 48: 1443-1447.

U.S. EPA. 1980. Ambient Water Quality Criteria Document for Cyanide. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC. EPA-440/5-80-037. NTIS PB 81-117483.

U.S. EPA. 1982. Errata for Ambient Water Quality Criteria Documents. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH for the Office of Water Regulations and Standards, Washington, DC.

U.S EPA. 1986. Guidelines for Carcinogen Risk Assessment. Federal Register. 51(185): 33992-34003.

U.S. EPA. 1991. Integrated Risk Information System (IRIS). Online. Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH.

U.S. Public Health Service. 1962. Drinking water standards. U.S. Gov. Printing Off., Washington, D.C. PHS Publ. No. 956.

Sorbo, B.H. 1953. On the substrate specificity of rhodanese. Acta Chem. Scand. 7: 32-37.

Streicher, E. 1951. Toxicity of colchicine, di-isopropyl fluorophosphate, intocostrin and potassium cyanide in mice at 4°C. Proc. Soc. Exp. Biol. Med. 76: 536-538.

Tewe, O.O. 1982. Effect of dietary cyanide on the performance, metabolism and pathology of the African rat (Cricetomys gambianus Waterhouse). Nutr. Rep. Int. 26(4): 529-536.

Tewe, O.O. and J.H. Maner. 1980. Cyanide, protein and iodine interactions in the performance, metabolism and pathology of pigs. Res. Vet. Sci. 29(3): 271-276.

Tewe, O.O., J.H. Maner. 1981a. Long-term and carry-over effect of dietary inorganic cyanide (KCN) in the life cycle performance and metabolism of rats. Toxicol. Appl. Pharmacol. 58(1): 1-7.

Tewe, O.O. and J.H. Maner. 1981b. Performance and pathophysiological changes in pregnant pigs fed cassava diets containing different levels of cyanide. Res. Vet. Sci. 30(2): 147-151.

Tewe, O.O. and J.H. Maner. 1982. Cyanide, protein and iodine interaction in the physiology and metabolism of rats. Food Chem. 9(3): 195-204.

Weast, R.C., Ed. 1980. CRC Handbook of Chemistry and Physics, 61st ed. CRC Press, Inc., Boca Raton, FL. p. B-98, B-133, B-147.

Westley, J. 1983. Sulfur-transferases. Fund. Appl. Toxicol. 3: 377-382.

Wexler, J., J.L. Whittenberger and P.R. Dumke. 1947. The effect of cyanide on the electrocardiogram of man. Am. Heart J. 34: 163-173.

WHO (World Health Organization). 1982. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Chemicals, industry process and industries associated with cancer in humans. International Agency for Research on Cancer Monographs. Vol. 1 to 29(Suppl. 4). Geneva.

Willhite, C.C. 1982. Congenital malformations induced by laetrile. Science. 215(4539): 1513-1515.

Willhite, C.C. 1983. Development toxicology of acetonitrile in the Syrian golden hamster. Teratology. 27(3): 313-325.

Willhite, C.C., V.H. Ferm and R.P. Smith. 1981. Teratogenic effects of aliphatic nitriles. Teratology. 23: 317-323.

Williams, R.T. 1959. Detoxication Mechanisms, 2nd ed. John Wiley and Sons, Inc., NY.

Wilson, J. 1983. Cyanide in human disease: A review of clinical and laboratory evidence. Fund. Appl. Toxicol. 3(5): 397-399.

Utti, R.J., A.H. Rajput, E.M. Ashenhurst and B. Rozdilsky. 1985. Cyanide-induced Parkinsonism: A clinicopathologic report. Neurology. 35(6): 291-295.

Veith, G.D., D.L. DeFoe and B.V. Bergstedt. 1979. Measuring and estimating the bioconcentration factor of chemicals in fish. J. Fish. Res. Board Can. 36: 1040-1048.

Walton, D.C. and M.G. Witherspoon. 1926. Skin absorption of certain gases. J. Pharmacol. Exp. Ther. 26: 315-324.

Watson, M.R. 1973. Cyanide Removal from Water. In: Pollution Technology Review, No. 5: Pollution Control in Metal Finishing. Noyes Data Corp., Park Ridge, NJ. p. 147-178.

Way, J.L. 1981. Pharmacologic aspects of cyanide and its antagonism. In: Cyanide in Biology, B. Vennesland, E.E. Conn and C. Knowles, Ed. Academic, London, England. p. 29-49.

Way, J.L. 1984. Cyanide intoxication and its mechanism of antagonism. Ann. Rev. Pharmacol. Toxicol. 24: 451-481.

Way, J.L., S.L. Gibbon and M. Sheehy. 1966. Cyanide intoxication: protection with oxygen. Sciences. 152: 210-211.

Way, J.L., E. End, M.H. Sheehy et al. 1972. Effect of oxygen on cyanide intoxication: IV. Hyperbaric oxygen. Toxicol. Appl. Pharmacol. 22: 415-421.

Wilson, J. and D.M. Matthews. 1966. Metabolic interrelationships between cyanide, thiocyanate and vitamin B_{12} in smokers and non-smokers. Clin. Sci. 31: 1-7.

Wilson, J., J.C. Linnell and D.M. Matthews. 1971. Plasma-cobalamins in neuro-ophthalmological diseases. Lancet. 1: 259-261.

Wokes, F. 1958. Tobacco amblyopia. Lancet. 2: 526-527.

Wolfsie, J.H. and C.B. Shaffer. 1959. Hydrogen cyanide: Hazards, toxicology, prevention and management of poisoning. J. Occup. Med. 1: 281-288.

Wood, J.L. and S.L. Cooley. 1956. Detoxication of cyanide by cystine. J. Biol. Chem. 218: 449-457.

Yamamoto, K., Y. Yamamoto, H. Hattori and T. Samori. 1982. Effects of routes of administration on the cyanide concentration distribution in the various organs of cyanide-intoxicated rats. Tohoku J. Exp. Med. 137(1): 73-78.

U.S. EPA Headquarters Library
Mail code 3201 3404T
1200 Pennsylvania Avenue NW
Washington DC 20460