




**Annual Water Sampling  
and Analysis at the  
Salmon Test Site Area,  
Lamar County, Mississippi  
April 2001**



# **Annual Water Sampling and Analysis at the Salmon Test Site Area**

**Lamar County, Mississippi  
April 2001**

by

Max G. Davis

Radiation and Indoor Environments National Laboratory  
Office of Radiation and Indoor Air  
U.S. Environmental Protection Agency  
P.O. Box 98517  
Las Vegas, NV 89193-8517

Prepared for the U.S. Department of Energy  
under Interagency Agreement  
DE-AI08-96NV11969

RADIATION AND INDOOR ENVIRONMENTS NATIONAL LABORATORY  
U.S. ENVIRONMENTAL PROTECTION AGENCY  
LAS VEGAS, NV 89193-8517

## **NOTICE**

The information in this document has been funded wholly or in part by the United States Environmental Protection Agency (EPA) through Interagency Agreement (IAG) DE-AI 08-96NV11969 from the United States Department of Energy (DOE). It has been subjected to the Agency's peer and administrative review, and it has been approved for publication as an EPA document. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

## ABSTRACT

In 1964 and 1966, nuclear explosives were detonated approximately 2,700 feet (823 m) underground in the Salmon Test Site Area located in Lamar County, Mississippi. Drilling and clean-up activities have resulted in tritium contamination in close proximity to the surface ground zero. The Long-Term Hydrological Monitoring Program (LTHMP), directed by the EPA, conducts annual water sampling on and around the Salmon Test Site Area and monitors radiological sampling equipment on the site throughout the year.

In this report the 2001 annual water sampling at the Salmon Site is described, and the analytical results of the collected samples are given. The highest tritium concentration onsite was  $2.67 \times 10^4$  pCi/L in water from one of the new wells added in 1997, see Appendix B, page 16. No radioactivity attributable to the test site was found in any offsite water sample. The highest tritium concentration offsite was  $9.8 \pm 3.4$  pCi/L at the Bill Ray Anderson residence.

All samples, with the exception of HMH-1 through HMH-16 and the 15 new wells added in 1998 (see Appendix B), were analyzed for the presence of gamma-ray emitting radionuclides. None were detected above the minimum detectable concentration (MDC).

This page is left blank intentionally.

## CONTENTS

|  | Page |
|--|------|
| Notice .....   | ii   |
| Abstract .....   | iii  |
| Figures .....  | vi   |
| Acronyms and Abbreviations .....                                   | vii  |
| Acknowledgments .....  | viii |
| Introduction .....   | 1    |
| History .....  | 1    |
| Historical Monitoring Results .....                                | 1    |
| Sampling Collection .....  | 4    |
| Sample Analysis .....  | 7    |
| Water Analysis Results .....                                       | 10   |
| Conclusions .....  | 11   |
| References .....   | 12   |
| Glossary of Terms .....  | 13   |
| Appendices   |      |
| A. Summary of Analytical Procedures .....                          | 15   |
| B. Tritium Results for Water Samples Collected in April 2001 ..... | 16   |

## FIGURES

| Figure Number | Page  |
|---------------|---|
| 1             | General site location of Project Salmon Test Site Area . . . . . 2  |
| 2             | Topographic map of the Salmon Test Site Area showing the Surface<br>Ground Zero and outline of Test Area at 2,700 feet below land surface . . . . . 3 |
| 3             | Test cavity and aquifers . . . . . 4  |
| 4             | Tritium concentration vs. sampling year for HMH-1 (depth = 10 ft) . . . . . 5   |
| 5             | Tritium concentration vs. sampling year for HMH-2 (depth = 10 ft) . . . . . 5   |
| 6             | Tritium concentration vs. sampling year for HMH-5 (depth = 10 ft) . . . . . 6   |
| 7             | Tritium concentration vs. sampling year for HM-S (depth = 30 ft) . . . . . 6  |
| 8             | Tritium concentration vs. sampling year for HM-L (depth = 200 ft) . . . . . 7   |
| 9             | Locations on the Salmon Test Site Area sampled in 2001 . . . . . 8  |
| 10            | Offsite locations sampled in 2001 . . . . . 9   |

## ACRONYMS AND ABBREVIATIONS

|                |   |
|----------------|---|
| DCG            | Derived Concentration Guide   |
| DOE            | U.S. Department of Energy   |
| EPA            | U. S. Environmental Protection Agency   |
| g              | gram  |
| HpGe           | high purity germanium gamma detector  |
| IAG            | Interagency Agreement   |
| keV            | kilo electron volts = thousand electron volts   |
| kg             | kilogram, 1000 grams  |
| kt             | kiloton (TNT equivalent)  |
| L              | liter   |
| LTHMP          | Long Term Hydrological Monitoring Program   |
| m              | meter   |
| MDC            | minimum detectable concentration  |
| MeV            | one million electron volts  |
| min            | minute  |
| mL             | milliliter = one thousandth of a liter  |
| ORIA           | Office of Radiation and Indoor Air  |
| pCi/L          | picocuries per liter = $10^{-12}$ curies per liter = 1/1,000,000,000,000 curies per liter |
| PHS            | U.S. Public Health Service  |
| R&IE           | Radiation and Indoor Environments National Laboratory                                     |
| SGZ            | surface ground zero   |
| USGS           | U.S. Geological Survey  |
| $^3\text{H}$   | Tritium   |
| $^3\text{H}^+$ | Enriched Tritium  |
| HMH            | Hydrological Monitoring Hole (1-16)   |
| HM-L, HM-L2    | Hydrological Monitoring Well Local Aquifer  |
| HM-S           | Hydrological Monitoring Well - Surficial Aquifer  |
| HM-1           | Hydrological Monitoring Well - Aquifer 1  |
| HM-2a          | Hydrological Monitoring Well - Aquifer 2a   |
| HM-2b          | Hydrological Monitoring Well - Aquifer 2b   |
| HM-3           | Hydrological Monitoring Well - Aquifer 3  |
| HT-2c          | Hydrological Test Hole  |
| HT-4           | Hydrological Test Hole  |
| HT-5           | Hydrological Test Hole  |
| REECo          | Reynolds Electrical & Engineering Co  |
| SA Wells       | Source Area Wells   |



## **ACKNOWLEDGMENTS**

External peer review was provided by Vernon Hodges, PhD, Chemistry, University Nevada Las Vegas. In addition, the author would like to acknowledge Ken Giles, Dennis Farmer, Richard Flotard, Pat Honsa, Rose Houston, and the staff of the Hydrological Monitoring Group, EPA, for their dedication to quality and their tireless work in the execution of the sampling effort. The author would also like to thank Terry Mouck for her outstanding support in preparing the graphics and for providing the desktop publishing in the production of this report.

---

## INTRODUCTION

Under an Interagency Agreement (IAG) with the DOE, the EPA's Radiation and Indoor Environments National Laboratory (R&IE) located in Las Vegas, NV, conducts sampling to measure radioactivity in water sources near the sites of underground nuclear explosions. The results provide assurance that radioactive materials from the tests have not migrated into drinking water supplies. This report presents the results for samples collected under EPA's LTHMP in April 2001, on and around the Salmon Test Site Area, Lamar County, Mississippi.

### History

Project Dribble, consisting of two nuclear explosions, and Project Miracle Play, consisting of two non-nuclear gas explosions, were conducted in the Salmon Test Site Area, near Baxterville, Lamar County, Mississippi, between 1964 and 1970. The general area is depicted in Figure 1. The Salmon Test Site Area (Figure 2) contains approximately 1,470 acres located in Sections 11, 12, 13, and 14, Township 2 North, Range 16 West.

| Test Date | Name        | Type    | Yield (kt) |
|-----------|-------------|---------|------------|
| 10-22-64  | Salmon      | Nuclear | 5.3        |
| 12-03-66  | Sterling    | Nuclear | 0.38       |
| 02-02-69  | Diode Tube  | Gas     | 0.32       |
| 04-19-70  | Humid Water | Gas     | 0.32       |

These tests were part of the Vela Uniform Program of the U.S. Atomic Energy Commission (a predecessor agency of the DOE). The purpose was to measure and evaluate the phenomena of seismic waves that


are induced from the explosions as compared to those that occur naturally from earthquakes.

The first explosion, the Salmon Event, created a cavity in the salt dome underlying the test area. The top of the cavity is 1,160 feet (360 m) below the top of the salt dome which lies 1,500 feet (460 m) below the land surface (Figure 3). The Salmon detonation cavity was subsequently used to contain the next three explosions.


Following each detonation, the surrounding area was closely monitored by the U.S. Public Health Service (PHS). Radiological monitoring became the responsibility of the EPA at its inception in 1970, and after the second site cleanup operation in 1971-72, the LTHMP was instituted. In this program, all potable aquifers, several wells, public water supplies, and some surface waters in the vicinity of the Salmon Test Site are sampled and analyzed to determine the presence of tritium and other radioactive contaminants.

### Historical Monitoring Results


The disposal of drilling mud and fluids near the surface ground zero (SGZ) is responsible for tritium ( $^3\text{H}$ ) contamination of the soil zone and underlying shallow aquifer. These waters lie at depths of 4 to 10 feet (1.3 to 3 m) and 30 feet (9 m), respectively, and are not potable. Tritium contamination is also present in the potable water of the local aquifer which lies at about 200 feet (62 m). The observed  $^3\text{H}$  concentration in the local aquifer is well below the 20,000 pCi/L guideline specified in the National Primary Drinking Water Regulations (40CFR141), and is thought to be



**Figure 1** General site location of Project Salmon Test Site Area.



**Figure 2** Topographic map of the Salmon Test Site Area showing the Surface Ground Zero and outline of Test Area at 2,700 feet below land surface.




**Figure 3** Test cavity and aquifers.


due to drilling activities at the site (Fenske and Humphrey, 1980;. Fordham and Fenske, 1985).

Of the 55 wells sampled on the Salmon Site, five regularly have tritium values above those expected in surface water values. These include three soil zone wells near the SGZ (wells HMH-1, HMH-2, and HMH-5), well HM-S in the shallow aquifer, and well HM-L in the


underlying local aquifer. Plots of tritium concentration vs. time for these wells are shown in Figures 4 to 8. The solid line in the graph represents the normal radioactive decay of tritium. Surface water collected from the Half Moon Creek overflow pond adjacent to the SGZ area, has tritium values above background as does the REECop Pit drainage area which was used for the disposal of drilling mud.




**Figure 4** Tritium concentration vs. sampling year for HMH-1 (depth = 10 ft).




**Figure 5** Tritium concentration vs. sampling year for HMH-2 (depth = 10 ft).



**Figure 6** Tritium concentration vs. sampling year for HMH-5 (depth = 10 ft).



**Figure 7** Tritium concentration vs. sampling year for HM-S (depth = 30 ft).



**Figure 8** Tritium concentration vs. sampling year for HM-L (depth = 200 ft).


## Sample Collection

According to standard operating procedures agreed to by DOE (U.S. DOE 1981), the HMH wells on the test site were first sampled, pumped-dry, and sampled on the following day as were the 15 new wells added in 1997. The 15 new wells added in 1998 were purged of three volumes of water and then sampled. Wells HM-1, HM-2A, HM-2B, HM-3, and HM-L, which lie adjacent to SGZ, were first sampled and then pumped steadily while further samples were taken at 30 min intervals until the pH and conductivity of the water stabilized. A final sample was taken from each well 30 min after stability was reached. Water samples were taken from sources near the SGZ area (i.e., Half Moon Creek, Half Moon Creek Overflow, and the Pond west of SGZ) before and after the


pumping operations to identify any resulting changes in tritium concentration. Well HM-L2 was first sampled and then pumped for one hour before a second sample was taken. All other water supplies were sampled only once.

For wells with operating pumps, the samples were collected at the nearest convenient outlet. If the well has no pump, a truck-mounted sampling unit was used. With this unit it is possible to collect three-liter samples from wells as deep as 1800 meters (5,900 ft.). The pH, conductivity, water temperature, and sampling depth was measured and recorded as each sample was collected. Disposal of water from wells HM-3, SA-1-1H, and SA1-7H, were contained in a Frac Tank. The water was then transported to the Columbia Waste Water Plant for disposal.





**Figure 9** Locations on the Salmon Test Site Area sampled in 2001.



**Figure 10. Offsite locations sampled in 2001**

The locations of all sampling sites are shown in Figures 9–10. Sampling also included three locations in Columbia, Mississippi (not shown). The sampling results are discussed in the following sections.

## Sample Analysis

Radiochemical procedures used to analyze the samples collected for this report are described in Johns et al. (1979) and summarized in Appendix A. These include standard methods to identify natural and man-made gamma-emitting radionuclides, tritium, plutonium, strontium, and uranium in water samples.

Two types of tritium analyses were performed: conventional and electrolytic enrichment. The enrichment method lowers the minimum detectable concentration (MDC) from approximately 300 pCi/L to about 5 pCi/L. An upper limit of activity of 700–800 pCi/L has been established for the tritium enrichment method because sample cross-contamination becomes a problem at higher levels.

In late 1995, it was decided that a maximum of 25 percent of all samples collected would be analyzed by the low-level enrichment method. This decision was based on the time required for analysis, budgetary constraints, and an assessment of past results. Under the current sampling and analysis protocol for the site, all samples are initially screened for tritium activity by the conventional method and selected samples enriched. At this time, only sampling locations that are in position to show migration are selected for enrichment.

Sufficient sample is collected from new sampling locations to perform all routine analyses and a full-suite of other radio-

chemical determinations including assays for strontium-90, plutonium, and uranium.

## Water Analysis Results

Gamma-ray spectral analysis results indicated that no man-made gamma-emitting radionuclides were detected in any onsite or offsite samples. Tritium concentrations above normal background values were not detected in any offsite samples. Long-term decreasing trends in tritium concentrations are evident for onsite locations that have shown detectable tritium activity since monitoring began under the LTHMP (wells HMH-1, HMH-2, HMH-5, HM-S, and HM-L, depicted in Figures 4 thru 8).

Only three wells were above the MDC in the samples collected from the offsite sampling locations. Tritium activity in the offsite samples ranged from less than the MDC to 9.8 pCi/L (~0.5 Bq/L), 0.05 percent of the DCG. These results do not exceed the natural tritium activity expected in rain water in this area.

Due to the high rainfall in the area, the sampling procedure for selected onsite wells is modified as follows: after collection of an initial sample, the well is purged, and a second sample taken after the well refills. The second sample is representative of water that has infiltrated through the soil zone, whereas the first sample may represent a mixture of direct rainwater influx at top of the well and infiltrated or soil zone water.

In April 2001, a total of 21 sampling locations onsite were not sampled by order of DOE reducing the number of location onsite from 61 to 40. Of the 40 locations sampled onsite, 33 sites were sampled twice (pre-and post-sampling), 25 yielded tritium activities greater

than the MDC in either the first or second sample. Of these, 9 yielded results higher than normal background (approximately 25 - 40 pCi/L [1 - 1.5 Bq/L]) as shown in Appendix B. The locations where the highest tritium activities were measured generally correspond to areas of known contamination.

In 1997, an additional 15 shallow wells were added to the annual sampling, which range in depth of 195' to 2100', increasing the total sampling locations onsite to 61. Five of the previous locations regularly have tritium values above those expected in surface water samples; of the 15 new wells, tritium values were all below the MDC. The water in these wells is not accessible to the public, nor suitable for drinking due to its brackishness.

No radioactive materials from the Salmon Test Site Area were detected in any water samples collected offsite. The tritium concentrations of water samples collected onsite and offsite are consistent with those of past studies at the Salmon Site. The highest tritium concentration found in water collected in the offsite area was  $9.8 \pm 3.4$  pCi/L, which is typical of background tritium levels, and is 0.059 percent of the National Interim Primary Drinking Water Regulations (40CFR141).

The highest tritium concentration found onsite was 26,700 pCi/L. This was detected in a water sample collected from Well SA1-1H which is a shallow well (40') near SGZ. The water from this well is not available to the public nor is it potable.

The tritium concentrations, except for Well SA1-1H, were all well below the 20,000 pCi/L level defined in the EPA Drinking Water Regulations (40CFR141).

All samples were analyzed for presence of gamma-ray emitting radionuclides with the exception of HMH-1 through HMH-16. None was detected above the MDC (see Appendix B on page 16).

## REFERENCES

Code of Federal Regulations, Vol. 41, title 40, Part 141, July 9, 1976, National Interim Primary Drinking Water Regulations.

A Guide for Environmental Radiological Surveillance at U.S. Dept. of Energy Installations, July 1981, Office of Operational Safety Report. Las Vegas, NV: U.S. Department of Energy; DOE/EP-0023.

Fenske, P. R.; Humphrey, T. M., Jr. The Tatum Dome Project Lamar County, Mississippi. Las Vegas, NV: U.S. Department of Energy, Nevada Operations Office; NVO-225; 1980.

Fordham, J. W. Fenske, P. R. Tatum Dome Field Study Report and Monitoring Data Analysis, Las Vegas, NV: U.S. Department of Energy, Nevada Operations Office; DOE/NV/10384-03; 1985.

Johns, F., et al. 1979. Radiochemical and Analytical Procedures for Analysis of Environmental Samples. Las Vegas, NV: U.S. Environmental Protection Agency; EMSL-LV-0539-17-1979.

## GLOSSARY OF TERMS

### Background Radiation

The radiation in man's environment, including cosmic rays and radiation from naturally-occurring and man-made radioactive elements, both outside and inside the bodies of humans and animals. The usually quoted average individual exposure from background radiation is 125 millirem per year in mid-latitudes at sea level (Shein & Terplak, 1984).

### Curie (Ci)

The basic unit used to describe the rate of radioactive disintegration. The curie is equal to 37 billion disintegrations per second, which is the equivalent of 1 gram of radium. Named for Marie and Pierre Curie who discovered radium in 1898. One microcurie (pCi) is one millionth of a Ci.

### Isotope

Atoms of the same element with different numbers of neutrons in the nuclei. Thus  $^{12}\text{C}$ ,  $^{13}\text{C}$ , and  $^{14}\text{C}$  are isotopes of the element carbon, the numbers denoting the approximate atomic weights. Isotopes have very nearly the same chemical properties, but often different physical properties (for example  $^{12}\text{C}$  and  $^{13}\text{C}$  are stable,  $^{14}\text{C}$  is radioactive).

### Enrichment Method

A method of electrolytic concentration that increases the sensitivity of the analysis of tritium in water. This method is used by R&IE in selected samples if the tritium concentration is less than 700 pCi/L.

### Minimum Detectable Concentration (MDC)

The smallest amount of radioactivity that can be reliably detected with a probability of Type I and Type II errors at 5 percent each (DOE 1981).

### Offsite

Areas exclusive of the immediate Salmon Test Site Area.

### Onsite

Refers to the immediate vicinity of the Salmon Test Site Area.

### Shallow ground water

Water found near the soil surface, caused by precipitation infiltration of the soil. This shallow ground water is not an aquifer.

## **GLOSSARY OF TERMS**

**(Continued)**

### **Surficial Aquifer**

The ground water layer located closest to the surface, generally at a depth of approximately 30 feet at SGZ.

### **Tritium**

A radioactive isotope of hydrogen that decays by beta emission. Its half-life is about 12.5 years.

### **Pre Sample**

First sample taken from wells onsite (before pumping).

### **Post Sample**

Last sample taken from wells onsite (after recharge).

### **Type I Error**

The statistical error of accepting the presence of radioactivity when none is present. Sometimes called alpha error.

### **Type II Error**

The statistical error of failing to recognize the presence of radioactivity when it is present. Sometimes called beta error.

## APPENDIX A

### Summary of Analytical Procedures

| Type of Analysis           | Analytical Equipment  | Counting Period (Min) | Analytical Procedures  | Sample Size | Approximate Detection Limit <sup>a</sup>   |
|----------------------------|---|-----------------------|--|-------------|--|
| HpGe Gamma <sup>b</sup>    | HpGe detector calibrated at 0.5 keV/channel (0.04 to 2 MeV range). Individual detector efficiencies ranging from 15 to 35%. | 150                   | Radionuclide concentration quantified from gamma spectral data by online computer program. | 3.5L        | Varies with radionuclides and detector used, normally counted to a MDC of approx. 5 pCi/L for Cs-137 |
| <sup>3</sup> H             | Automatic liquid scintillation counter  | 300                   | Sample prepared by distillation.   | 5 to 10 mL  | 300 to 700 pCi/L   |
| <sup>3</sup> H+ Enrichment | Automatic liquid scintillation counter  | 300                   | Sample concentrated by electrolysis followed by distillation.                              | 5 mL        | 5 pCi/L  |

<sup>a</sup> The detection limit is defined as the smallest amount of radioactivity that can be reliably detected, i.e., probability of Type I and Type II error at 5 percent each (DOE 1981).

<sup>b</sup> Gamma spectrometry using a high purity intrinsic germanium (HpGe) detector.

### Typical MDA Values for Gamma Spectroscopy (100 minute count time)

|           |           |         |          |
|-----------|-----------|---------|----------|
| Geometry* | Marinelli | Model   | 430G     |
| Matrix    | Water     | Density | 1.0 g/ml |
| Volume    | 3.5 liter | Units   | pCi/L    |
| Isotope   | MDA       | Isotope | MDA      |
|           |           | Ru-106  | 4.76E+01 |
| Be-7      | 4.56E+01  | Sn-113  | 8.32E+00 |
| K-40      | 4.92E+01  | Sb-125  | 1.65E+01 |
| Cr-51     | 5.88E+01  | I-131   | 8.28E+00 |
| Mn-54     | 4.55E+01  | Ba-133  | 9.16E+00 |
| Co-57     | 9.65E+00  | Cs-134  | 6.12E+00 |
| Co-58     | 4.71E+00  | Cs-137  | 6.43E+00 |
| Fe-59     | 1.07E+01  | Ce-144  | 7.59E+01 |
| Co-60     | 5.38E+00  | Eu-152  | 2.86E+01 |
| Zn-65     | 1.24E+01  | Ra-226  | 1.58E+01 |
| Nb-95     | 5.64E+00  | U-235   | 1.01E+02 |
| Zr-95     | 9.06E+00  | Am-241  | 6.60E+01 |

#### Disclaimer

The MDA's provided are for background matrix samples presumed to contain no known analytes and no decay time. All MDA's provided here are for one specific \*Germanium detector and the geometry of interest. The MDA's in no way should be used as a source of reference for determining MDA's for any other type of detector. All gamma spectroscopy MDA's will vary with different types of shielding, geometries, counting times, and decay time of sample.



**APPENDIX B****TRITIUM RESULTS FOR WATER SAMPLES COLLECTED IN APRIL 2001**

| Sample<br>Location           | Collection<br>Date | Enriched<br>Tritium<br>pCi/L $\pm$ 2 SD | (MDC) | Tritium<br>pCi/L $\pm$ 2 SD | (MDC) | Comments             | Gamma<br>Spectrometry <sup>(b)</sup><br>(MDC) |
|------------------------------|--------------------|---|-------|-----------------------------|-------|----------------------|---|
| <b>Baxterville, MS</b>       |                    |   |       |                             |       |                      |   |
| Anderson, Billy Ray          | 4-16               | 7.2 $\pm$ 3.8                           | (6.0) |                             |       |                      | ND (5.0)                                      |
| Anderson Pond                | 4-16               | 9.8 $\pm$ 3.4                           | (5.0) |                             |       |                      | ND (4.0)                                      |
| Vines, James                 | 4-16               |   |       | -28 $\pm$ 136               | (225) |                      | ND (5.0)                                      |
| Anderson, Rhonda             | 4-16               |   |       | 29.5 $\pm$ 137              | (225) |                      | ND (4.6)                                      |
| Anderson, Robert Lowell, Jr. | 4-17               |   |       | -85 $\pm$ 135               | (225) |                      | ND (4.6)                                      |
| Anderson, Robert Lee, Jr.    | 4-17               |   |       | 26 $\pm$ 137                | (225) |                      | ND (4.9)                                      |
| Anderson, Tony               | 4-16               |   |       |                             |       | No sample city water |   |
| Burge, Joe                   | 4-17               |   |       | -63 $\pm$ 136               | (225) |                      | ND (4.9)                                      |
| Half Moon Creek              | Pre 4-16           | 1.3 $\pm$ 3.4                           | (5.0) |                             |       |                      | ND (4.6)                                      |
|                              | Post 4-17          | 10.3 $\pm$ 2.2                          | (4.5) |                             |       |                      | ND (4.6)                                      |
| Half Moon Creek<br>Overflow  | Pre 4-16           | 214 $\pm$ 5.7                           | (5.2) |                             |       |                      | ND (4.6)                                      |
|                              | Post 4-17          | 222 $\pm$ 5.7                           | (5.0) |                             |       |                      | ND (4.8)                                      |
| Hibley, Billy                | 4-16               |   |       | 33 $\pm$ 137                | (225) |                      | ND (4.9)                                      |
| Napier, Denice               | 4-17               | 11 $\pm$ 3.6                            | (5.6) |                             |       |                      | ND (5.0)                                      |
| Lee, Perry T., Jr.           | 4-16               |   |       | -42 $\pm$ 136               | (225) |                      | ND (5.0)                                      |
| Little Creek #1              | 4-16               |   |       | -70.5 $\pm$ 136             | (225) |                      | ND (5.0)                                      |
| Lower Little Creek #2        | 4-16               |   |       | -13.4 $\pm$ 137             | (225) |                      | ND (1.5)                                      |
| Mills, Roy                   | 4-16               |   |       |                             |       | No sample city water |   |

<sup>(a)</sup> Indicates results are less than MDC

<sup>(b)</sup> No gamma radionuclides detected above MDC

ND Non-detected, represents <sup>137</sup>Cs (pCi/L)

**APPENDIX B (Continued)****TRITIUM RESULTS FOR WATER SAMPLES COLLECTED IN APRIL 2001**

| Sample Location               | Collection Date | Enriched Tritium<br>pCi/L $\pm$ 2 SD (MDC) | Tritium<br>pCi/L $\pm$ 2 SD (MDC)    | Comments             | Gamma Spectrometry <sup>(b)</sup><br>(MDC) |
|-------------------------------|-----------------|--|--------------------------------------|----------------------|--|
| <b>Baxterville, MS (Cont)</b> |                 |  |                                      |                      |  |
| Mills, AC                     | 4-16            |  |                                      | No sample city water |  |
| Nobles Pond                   | 4-17            |  | 33 $\pm$ 137 (225)                   |                      | ND (4.6)                                   |
| Noble, Evelyn                 | 4-17            |  | -17 $\pm$ 136 (225)                  |                      | ND (4.7)                                   |
| Pond West of GZ               | Pre 4-16        | 4.9 $\pm$ 3.4 (5.4)                        |                                      |                      | ND (4.3)                                   |
|                               | Post 4-17       | 9.6 $\pm$ 3.6 (5.6)                        |                                      |                      | ND (4.5)                                   |
| REECo Pit Drainage-A          | 4-16            |  | 142 $\pm$ 202 (225)                  |                      | ND (4.6)                                   |
| REECo Pit Drainage-B          | 4-16            |  | 25.9 $\pm$ 123 (225)                 |                      | ND (4.9)                                   |
| REECo Pit Drainage-C          | 4-16            | 22.4 $\pm$ 3.8 (5.5)                       |                                      |                      | ND (4.7)                                   |
| Saucier, Dennis               | 4-16            | 14 $\pm$ 4.0 (6.3)                         |                                      |                      | ND (4.6)                                   |
| Well Ascot 2                  | 4-16            |  |                                      | Not sampled per DOE  |  |
| Baxterville Well City         | 4-17            |  | 4.5 $\pm$ 137 (225)                  |                      | ND (1.5)                                   |
| Well E-7                      | 4-20            |  | -70 $\pm$ 199 (329)                  |                      | ND (4.8)                                   |
| Well HM-1                     | Pre 4-16        |  | 24 $\pm$ 200 <sup>(a)</sup> (329)    |                      | ND (4.9)                                   |
|                               | 1st 30 Min 4-16 |  | 98 $\pm$ 199 <sup>(a)</sup> (329)    |                      | ND (4.9)                                   |
|                               | 2nd 30 Min 4-16 |  | -4.2 $\pm$ 200 <sup>(a)</sup> (320)  |                      | ND (4.8)                                   |
|                               | Post 4-16       |  | -2.4 $\pm$ 200 <sup>(a)</sup> (329)  |                      | ND (4.3)                                   |
| Well HM-2A                    | Pre 4-16        |  | -18.6 $\pm$ 200 <sup>(a)</sup> (329) |                      | ND (5.0)                                   |
|                               | 1st 30 Min 4-16 |  | -4.3 $\pm$ 200 <sup>(a)</sup> (329)  |                      | ND (4.9)                                   |
|                               | 2nd 30 Min 4-16 |  | 30.6 $\pm$ 200 <sup>(a)</sup> (329)  |                      | ND (5.0)                                   |
|                               | 3rd 30 Min 4-16 |  | 50 $\pm$ 199 <sup>(a)</sup> (329)    |                      | ND (5.0)                                   |
|                               | Post 4-16       |  | 27 $\pm$ 200 <sup>(a)</sup> (329)    |                      | ND (4.9)                                   |
| Well HM-2B                    | Pre 4-16        |  | 210 $\pm$ 202 <sup>(a)</sup> (529)   |                      | ND (5.0)                                   |
|                               | 1st 30 Min 4-16 | -0.39 $\pm$ 3.1 (5.1)                      |                                      |                      | ND (1.9)                                   |
|                               | Post 4-16       | -3.2 $\pm$ 3.2 (5.0)                       |                                      |                      | ND (4.0)                                   |

<sup>(a)</sup> Indicates results are less than MDC<sup>(b)</sup> No gamma radionuclides detected above MDCND Non-detected, MDC for gamma represents <sup>137</sup>Cs (pCi/L)

**APPENDIX B (Continued)****TRITIUM RESULTS FOR WATER SAMPLES COLLECTED IN APRIL 2001**

| Sample Location                |            | Collection Date 2001 | Enriched Tritium<br>pCi/L $\pm$ 2 SD (MDC) | Tritium<br>pCi/L $\pm$ 2 SD (MDC) | Comments          | Gamma Spectrometry <sup>(b)</sup><br>(MDC) |
|--------------------------------|------------|----------------------|--|-----------------------------------|-------------------|--|
| <b>Baxterville, MS (cont.)</b> |            |                      |  |                                   |                   |  |
| Well HM-3                      | Pre        | 4-16                 | 1.5 $\pm$ 3.7                              | (6.1)                             |                   | ND (4.6)                                   |
|                                | 1st 30 Min | 4-16                 | -55 $\pm$ 3.8                              | (6.0)                             |                   | ND (4.6)                                   |
| Well HM3                       | 2nd 30 Min | 4-16                 | -50 $\pm$ 3.3 <sup>(a)</sup>               | (5.5)                             |                   | ND (4.9)                                   |
|                                | 3rd 30 Min | 4-16                 | 3.8 $\pm$ 3.7 <sup>(a)</sup>               | (6.0)                             |                   | ND (4.7)                                   |
|                                | Post       | 4-16                 | -48 $\pm$ 3.3 <sup>(a)</sup>               | (5.4)                             |                   | ND (4.5)                                   |
| Well HM-L                      | Pre        | 4-16                 | 67 $\pm$ 8.9                               | (5.0)                             |                   | ND (4.0)                                   |
|                                | 1st 30 Min | 4-16                 | 4.0 $\pm$ 7.0 <sup>(a)</sup>               | (5.0)                             |                   | ND (4.5)                                   |
|                                | 2nd 30 Min | 4-16                 | 347 $\pm$ 7.0                              | (5.0)                             |                   | ND (5.0)                                   |
|                                | 3rd 30 Min | 4-16                 | 358 $\pm$ 7.0                              | (5.0)                             |                   | ND (5.0)                                   |
|                                | 4th 30 Min | 4-16                 | 2.0 $\pm$ 6.0 <sup>(a)</sup>               | (5.0)                             |                   | ND (5.0)                                   |
|                                | 5th 30 Min | 4-16                 | 4.0 $\pm$ 7.0 <sup>(a)</sup>               | (5.0)                             |                   | ND (5.0)                                   |
|                                | Post       | 4-16                 | 4.0 $\pm$ 7.0 <sup>(a)</sup>               | (5.0)                             |                   | ND (5.0)                                   |
| Well HM-L2                     | Pre        | 4-17                 | -1.6 $\pm$ 3.5 <sup>(a)</sup>              | (5.7)                             |                   | ND (4.7)                                   |
|                                | Post       | 4-17                 | -1.3 $\pm$ 3.4 <sup>(a)</sup>              | (5.7)                             |                   | ND (1.9)                                   |
| Well HM-S                      | Pre        | 4-16                 |  | 1710 $\pm$ 216 <sup>(a)</sup>     | (329)             | ND (4.8)                                   |
|                                | Post       | 4-17                 |  | 1620 $\pm$ 212 <sup>(a)</sup>     | (329)             | ND (5.0)                                   |
| Well HMH-1                     | Pre        | 4-16                 |  | 182 $\pm$ 202                     | (329)             |  |
|                                | Post       | 4-17                 |  | 476 $\pm$ 204                     | (329)             |  |
| Well HMH-2                     | Pre        | 4-16                 |  | 35.0 $\pm$ 203                    | (329)             |  |
|                                | Post       | 4-17                 |  | 216 $\pm$ 202 <sup>(a)</sup>      | (329)             |  |
| Well HMH-3                     | Pre        | 4-16                 |  | -133 $\pm$ 199 <sup>(a)</sup>     | (329)             |  |
|                                | Post       | 4-17                 |  | 122 $\pm$ 201 <sup>(a)</sup>      | (329)             |  |
| Well HMH-4                     | Pre        | 4-16                 |  |                                   | No Sample per DOE |  |
|                                | Post       | 4-16                 |  |                                   | No Sample per DOE |  |
| Well HMH-5                     | Pre        | 4-16                 |  | 373 $\pm$ 203                     | (329)             |  |
|                                | Post       | 4-17                 |  | 708 $\pm$ 207                     | (329)             |  |
| Well HMH-6                     | Pre        | 4-16                 |  |                                   | No Sample per DOE |  |
|                                | Post       | 4-16                 |  |                                   | No Sample per DOE |  |

<sup>(a)</sup> Indicates results are less than MDC<sup>(b)</sup> No gamma radionuclides detected above MDCND Non-detected, MDC for gamma represents <sup>137</sup>Cs (pCi/L)

**APPENDIX B (Continued)****TRITIUM RESULTS FOR WATER SAMPLES COLLECTED IN APRIL 2001**

| Sample<br>Location             |      | Collection<br>Date<br>2001 | Enriched<br>Tritium<br>pCi/L $\pm$ 2 SD (MDC) | Tritium<br>pCi/L $\pm$ 2 SD (MDC)   | Comments | Gamma<br>Spectrometry <sup>(b)</sup><br>(MDC) |
|--------------------------------|------|----------------------------|---|-------------------------------------|----------|---|
| <b>Baxterville, MS (cont.)</b> |      |                            |   |                                     |          |   |
| Well HMH-7                     | Pre  | 4-16                       |   |                                     |          | No Sample per DOE                             |
|                                | Post | 4-16                       |   |                                     |          | No Sample per DOE                             |
| Well HMH-8                     | Pre  | 4-16                       |   |                                     |          | No Sample per DOE                             |
|                                | Post | 4-16                       |   |                                     |          | No Sample per DOE                             |
| Well HMH-9                     | Pre  | 4-16                       |   | -7 $\pm$ 200 <sup>(a)</sup> (329)   |          |   |
|                                | Post | 4-17                       |   | -139 $\pm$ 199 <sup>(a)</sup> (329) |          |   |
| Well HMH-10                    | Pre  | 4-16                       |   | 304 $\pm$ 203 <sup>(a)</sup> (329)  |          |   |
|                                | Post | 4-17                       |   | 327 $\pm$ 203 <sup>(a)</sup> (329)  |          |   |
| Well HMH-11                    | Pre  | 4-16                       |   | 47 $\pm$ 200 <sup>(a)</sup> (329)   |          |   |
|                                | Post | 4-17                       |   | 25 $\pm$ 127 <sup>(a)</sup> (329)   |          |   |
| Well HMH-12                    | Pre  | 4-16                       |   |                                     |          | No Sample per DOE                             |
|                                | Post | 4-16                       |   |                                     |          | No Sample per DOE                             |
| Well HMH-13                    | Pre  | 4-16                       |   |                                     |          | No Sample per DOE                             |
|                                | Post | 4-16                       |   |                                     |          | No Sample per DOE                             |
| Well HMH-14                    | Pre  | 4-16                       |   |                                     |          | No Sample per DOE                             |
|                                | Post | 4-16                       |   |                                     |          | No Sample per DOE                             |
| Well HMH-15                    | Pre  | 4-16                       |   |                                     |          | No Sample per DOE                             |
|                                | Post | 4-16                       |   |                                     |          | No Sample per DOE                             |
| Well HMH-16                    | Pre  | 4-16                       | 10 $\pm$ 3.5 (5.5)                            |                                     |          |   |
|                                | Post | 4-17                       | 11 $\pm$ 3.7 (6.0)                            |                                     |          |   |
| SA1-1H                         | Pre  | 4-17                       |   | 23,200 $\pm$ 415 (202)              |          | ND (4.9)                                      |
|                                | Post | 4-18                       |   | 26,700 $\pm$ 328 (202)              |          | ND (4.7)                                      |
| SA1-2H                         | Pre  | 4-17                       |   | 2520 $\pm$ 172 (202)                |          | ND (4.8)                                      |
|                                | Post | 4-18                       |   | 2280 $\pm$ 168 (202)                |          | ND (4.3)                                      |
| SA1-3H                         | Pre  | 4-17                       | 886 $\pm$ 103 (5.1)                           |                                     |          | ND (4.9)                                      |
|                                | Post | 4-18                       | 603 $\pm$ 8.5 (4.9)                           |                                     |          | ND (4.9)                                      |

<sup>(a)</sup> Indicates results are less than MDC<sup>(b)</sup> No gamma radionuclides detected above MDCND Non-detected, MDC for gamma represents <sup>137</sup>Cs (pCi/L)

**APPENDIX B (Continued)****TRITIUM RESULTS FOR WATER SAMPLES COLLECTED IN APRIL 2001**

| Sample Location                |      | Collection Date | Enriched Tritium<br>pCi/L $\pm$ 2 SD (MDC) | Tritium<br>pCi/L $\pm$ 2 SD (MDC)   | Comments | Gamma Spectrometry <sup>(b)</sup><br>(MDC) |
|--------------------------------|------|-----------------|--|-------------------------------------|----------|--|
| <b>Baxterville, MS (cont.)</b> |      |                 |  |                                     |          |  |
| SA1-4H                         | Pre  | 4-17            |  | 290 $\pm$ 130 (202)                 |          | ND (4.5)                                   |
|                                | Post | 4-18            |  | 229 $\pm$ 128 (202)                 |          | ND (4.8)                                   |
| SA1-5H                         | Pre  | 4-17            |  | 891 $\pm$ 142 (202)                 |          | ND (4.7)                                   |
|                                | Post | 4-18            |  | 879 $\pm$ 142 (202)                 |          | ND (4.7)                                   |
| SA1-6H                         | Pre  | 4-17            |  | 33,000 $\pm$ 453 (202)              |          | ND (5.0)                                   |
|                                | Post | 4-18            |  | 52 $\pm$ 124 (202)                  |          | ND (4.3)                                   |
| SA1-7H                         | Pre  | 4-16            |  | 47 $\pm$ 124 <sup>(a)</sup> (202)   |          | ND (4.7)                                   |
|                                | Post | 4-17            |  | 57 $\pm$ 124 <sup>(a)</sup> (202)   |          | ND (4.6)                                   |
| SA1-8-L                        |      | 4-18            |  | 17.0 $\pm$ 123 <sup>(a)</sup> (202) |          | ND (1.5)                                   |
| SA1-9-2A                       |      | 4-19            | 4.3 $\pm$ 3.1 <sup>(a)</sup> (5.0)         |                                     |          | ND (4.8)                                   |
| SA1-10-2B                      |      | 4-19            | 2.8 $\pm$ 3.6 <sup>(a)</sup> (5.8)         |                                     |          | ND (4.9)                                   |
| SA1-11-3                       |      | 4-19            |  | -17 $\pm$ 122 <sup>(a)</sup> (202)  |          | ND (4.9)                                   |
| SA2-1-L                        |      | 4-18            |  | 4.7 $\pm$ 123 <sup>(a)</sup> (202)  |          | ND (5.0)                                   |
| SA2-2-L                        |      | 4-16            |  | No sample per DOE                   |          |  |
| SA2-3-L                        |      | 4-18            | 5.7 $\pm$ 3.8 <sup>(a)</sup> (6.2)         |                                     |          | ND (4.9)                                   |
| SA2-4-L                        |      | 4-18            |  | 69 $\pm$ 125 <sup>(a)</sup> (202)   |          | ND (4.6)                                   |
| SA2-5-L                        |      | 4-18            |  | 112<br>ND (5.0)                     |          | $\pm$ 125 <sup>(a)</sup> (202)             |
| SA3-1M                         | Pre  | 4-18            |  | 112                                 |          | $\pm$ 125 <sup>(a)</sup> (202)             |
|                                | Post | 4-19            |  | 82                                  |          | ND (4.8)<br>$\pm$ 125 <sup>(a)</sup> (202) |
| SA3-3M                         | Pre  | 4-18            | 15 $\pm$ 3.2 (4.9)                         |                                     |          | ND (4.8)                                   |
|                                | Post | 4-19            | 14 $\pm$ 3.4 (5.2)                         |                                     |          | ND (5.0)<br>ND (5.0)                       |

<sup>(a)</sup> Indicates results are less than MDC<sup>(b)</sup> No gamma radionuclides detected above MDCND Non -detected, MDC for gamma represents <sup>137</sup>Cs (pCi/L)

**APPENDIX B (Continued)****TRITIUM RESULTS FOR WATER SAMPLES COLLECTED IN APRIL 2001**

| Sample Location                |      | Collection Date 2001 | Enriched Tritium<br>pCi/L $\pm$ 2 SD (MDC) | Tritium<br>pCi/L $\pm$ 2 SD (MDC)   | Comments          | Gamma Spectrometry <sup>(b)</sup><br>(MDC) |
|--------------------------------|------|----------------------|--|-------------------------------------|-------------------|--|
| <b>Baxterville, MS (cont.)</b> |      |                      |  |                                     |                   |  |
| SA3-4H                         | Pre  | 4-17                 | 16.4 $\pm$ 3.4 (5.0)                       |                                     |                   | ND (5.0)                                   |
|                                | Post | 4-18                 | 16.6 $\pm$ 3.3 (5.0)                       |                                     |                   | ND (5.0)                                   |
| SA3-5-H                        | Pre  | 4-17                 |  | 73.5 $\pm$ 125 <sup>(a)</sup> (202) |                   | ND (5.0)                                   |
|                                | Post | 4-18                 |  | 56 $\pm$ 124 <sup>(a)</sup> (202)   |                   | ND (4.9)                                   |
| SA3-8-1                        |      | 4-20                 | .60 $\pm$ 3.0 (4.9)                        |                                     |                   | ND (4.9)                                   |
| SA3-10-2B                      |      | 4-20                 |  | 86 $\pm$ 125 <sup>(a)</sup> (202)   |                   | ND (4.6)                                   |
| SA3-11-3                       |      | 4-20                 |  | -8.6 $\pm$ 123 <sup>(a)</sup> (202) |                   | ND (1.6 )                                  |
| SA4-1M                         | Pre  | 4-17                 | 7.4 $\pm$ 3.3 (5.0)                        |                                     |                   | ND (4.9)                                   |
|                                | Post | 4-18                 | 5.7 $\pm$ 3.0 (5.0)                        |                                     |                   | ND (4.7)                                   |
| SA5-1M                         | Pre  | 4-18                 | 7.8 $\pm$ 3.5 <sup>(a)</sup> (5.0)         |                                     |                   | ND (4.6)                                   |
|                                | Post | 4-19                 | 9.2 $\pm$ 3.3 <sup>(a)</sup> (5.0)         |                                     |                   | ND (4.6)                                   |
| SA5-2M                         | Pre  | 4-18                 | 13.7 $\pm$ 3.3 (5.0)                       |                                     |                   | ND (5.0)                                   |
|                                | Post | 4-19                 | 12.7 $\pm$ 3.5 (5.4)                       |                                     |                   | ND (4.8)                                   |
| SA5-3M                         | Pre  | 4-18                 |  | 82 $\pm$ 125 <sup>(a)</sup> (202)   |                   | ND (4.6)                                   |
|                                | Post | 4-19                 |  | 86.5 $\pm$ 125 <sup>(a)</sup> (202) |                   | ND (4.6)                                   |
| SA5-4-4                        |      | 4-20                 | 2.4 $\pm$ 3.2 <sup>(a)</sup> (5.3)         |                                     |                   | ND (4.8)                                   |
| SA5-5-4                        |      | 4-20                 | 1.84 $\pm$ 3.5 <sup>(a)</sup> (5.6)        |                                     |                   | ND (4.8)                                   |
| Well HT-2C                     |      | 4-16                 |  |                                     | No Sample per DOE |  |
| Well HT-4                      |      | 4-16                 |  |                                     | No Sample per DOE |  |
| Well HT-5                      |      | 4-16                 |  |                                     | No Sample per DOE |  |

<sup>(a)</sup> Indicates results are less than MDC<sup>(b)</sup> No gamma radionuclides detected above MDCND Non -detected, MDC for gamma represents <sup>137</sup>Cs (pCi/L)

**APPENDIX B (Continued)****TRITIUM RESULTS FOR WATER SAMPLES COLLECTED IN APRIL 2001**

| Sample<br>Location      | Collection<br>Date<br>2001 | Enriched<br>Tritium<br>pCi/L $\pm$ 2 SD (MDC) | Tritium<br>pCi/L $\pm$ 2 SD (MDC)    | Comments            | Gamma<br>Spectrometry <sup>(b)</sup><br>(MDC) |
|-------------------------|----------------------------|---|--------------------------------------|---------------------|---|
| <b>Columbia, MS</b>     |                            |   |                                      |                     |   |
| Well 46-003-5-6-7       | 4-17                       |   | -11 $\pm$ 136 <sup>(a)</sup> (225)   |                     | ND (1.6)                                      |
| <b>Lumberton, MS</b>    |                            |   |                                      |                     |   |
| Anderson, Arleene       | 4-18                       |   | -45.6 $\pm$ 136 <sup>(a)</sup> (225) |                     | ND (4.9)                                      |
| James Lowe              | 4-14                       |   | 40 $\pm$ 137 <sup>(a)</sup> (225)    |                     | ND (4.6)                                      |
| Crawfish Pond           | 4-16                       |   | -38 $\pm$ 125 <sup>(a)</sup> (225)   |                     | ND (4.7)                                      |
| Powell, Shannon         | 4-14                       |   | 104 $\pm$ 139 <sup>(a)</sup> (225)   |                     | ND (4.4)                                      |
| Smith, Howard Pond      | 4-16                       |   | -42 $\pm$ 136 <sup>(a)</sup> (225)   |                     | ND (4.6)                                      |
| Thompson, Mike          | 4-16                       |   | -57 $\pm$ 136 <sup>(a)</sup> (225)   |                     | ND (1.8)                                      |
| Thompson, Roswell       | 4-16                       |   | -24 $\pm$ 136 <sup>(a)</sup> (225)   |                     | ND (4.4)                                      |
| Well 2 City             | 4-17                       | -2.5 $\pm$ 3.5 <sup>(a)</sup> (5.9)           |                                      |                     | ND (4.2)                                      |
| Burge, Willie           | 4-17                       |   | 121 $\pm$ 135 <sup>(a)</sup> (225)   |                     | ND (1.4)                                      |
| Kiffe, Richie & Patsy   | 4-16                       |   | -6.3 $\pm$ 137 <sup>(a)</sup> (225)  |                     | ND (5.0)                                      |
| <b>Purvis, MS</b>       |                            |   |                                      |                     |   |
| City Supply Purvis      | 4-17                       | .89 $\pm$ 3.7 <sup>(a)</sup> (6.1)            |                                      |                     | ND (1.9)                                      |
| Rain Sample IT Compound | 4-28                       |   |                                      | No sample - no rain |   |

<sup>(a)</sup> Indicates results are less than MDC<sup>(b)</sup> No gamma radionuclides detected above MDCND Non -detected, MDC for gamma represents <sup>137</sup>Cs (pCi/L)