

Pesticide



N,N-diethyl-m-toluamide (Deet) Pesticide Registration Standard



DEET

Pesticide Registration Standard

Joseph Panetta
Owen Beeder
Lynn F. Brown
Tonda Hicks
Richard Johnson
Jeff Kempter
Cathy Kessler
Gary Otakie
Wendy Weinstein

William Boodee
Roger Gardner
Franklin D.R. Gee
Carolyn Gregorio
Raymond Kent
John Leitzke
Linda Priddy
Arthur Schlosser

Team Leader (SPRD)
Project Manager (SPRD)
Project Manager (SPRD)
Secretary (SPRD)
Project Manager (SPRD)
Section Head (SPRD)
Student Assistant (SPRD)
Project Manager (SPRD)
Student Assistant (SPRD)

Chemist (HED)
Toxicologist (HED)
Product Manager (RD)
Toxicologist (HED)
Residue Chemist (HED)
Wildlife Biologist (HED)
Entomologist (BFSD)
Environmental Chemist (HED)

December 1980

Office of Pesticides and Toxic Substances

Environmental Protection Agency
401 M Street, SW
Washington, DC 20460

TABLE OF CONTENTS

I. How to Register Under a Registration Standard	1
II. Regulatory Position	7
A. Introduction	7
B. Background	7
C. Regulatory Decision	7
D. Terms and Conditions of Registration	9
III. Product Chemistry	
A. Introduction	24
B. Disciplinary Review	24
1. Chemistry Profile	
2. Generic Data Gaps	
3. Required Labeling	
C. Topical Discussions	25
IV. Environmental Fate	
A. Disciplinary Review	30
1. Use Profile	
2. Environmental Fate Profile	
3. Exposure Profile	
4. Data Gaps	
5. Required Labeling	
B. Topical Discussions	34
V. Toxicology	
A. Disciplinary Review	36
1. Toxicology Profile	
2. Human Hazard Assessment	
3. Data Gaps	
4. Required Labeling	
B. Topical Discussions	40

VI. Ecological Effects	63
A. Disciplinary Review	63
1. Ecological Effects Profile and Hazard Assessment	
2. Data Gaps	
3. Required Labeling	
B. Topical Discussions	63
VII. Efficacy	65
A. Introduction	65
B. Efficacy Assessment	65
C. Data Gaps	71
D. Labeling	72
VIII. Case Bibliography	81
Guide to the Use of the Bibliography	81

I. HOW TO REGISTER UNDER A REGISTRATION STANDARD

Organization of the Standard

Purpose of the Standard

Requirement to Re-register Under the Standard

"Product-Specific" Data and "Generic" Data

Data Compensation Requirements under FIFRA 3(c) (1) (D)

Amendments to the Standard

Organization of the Standard

This first chapter explains the purpose of a Registration Standard and summarizes the legal principles involved in registering or re-registering under a Standard. The second chapter sets forth the requirements that must be met to obtain or retain registration for products covered by this particular Registration Standard. In the remaining chapters, the Agency reviews the available data by scientific discipline, discusses the Agency's concerns with the identified potential hazards, and logically develops the conditions and requirements that would reduce those hazards to acceptable levels.

Purpose of the Standard

Section 3 of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) provides that "no person in any State may distribute, sell, offer for sale, hold for sale, ship, deliver for shipment, or receive (and having so received) deliver or offer to deliver, to any person any pesticide which is not registered with the Administrator [of EPA]." To approve the registration of a pesticide, the Administrator must find, pursuant to Section 3(c) (5) that:

- "(A) its composition is such as to warrant the proposed claims for it;
- (B) its labeling and other material required to be submitted comply with the requirements of this Act;
- (C) it will perform its intended function without unreasonable adverse effect on the environment; and
- (D) when used in accordance with widespread and commonly recognized practice it will not generally cause unreasonable adverse effects on the environment."

In making these findings, the Agency reviews a wide range of data which registrants are required to submit, and assesses the risks and benefits associated with the use of the proposed pesticide. But the established approach to making these findings has been found to be defective on two counts:

First, EPA and its predecessor agency, the United States Department of Agriculture (USDA), routinely reviewed registration applications on a 'product-by-product' basis, evaluating each product-specific application independently. In the review of products containing similar components, there was little opportunity for a retrospective review of the full range of pertinent data available in Agency files and in the public literature. Thus the 'product-by-product' approach was often inefficient and sometimes resulted in inconsistent or incomplete regulatory judgments. Second, over the years, as a result of continuing advances in scientific knowledge, methodology, and policy, the data base for many pesticides came to be considered inadequate by current scientific and regulatory standards. Given the long history of pesticide regulation in several agencies, it is even likely that materials may have been lost from the data files. When EPA issued new requirements for registration in 1975 (40 CFR 10, 1978 and 43 FR 37336, August 2, 1978), many products that had already been registered for years were being sold and used without the same assurances of human and environmental safety as was being required for new products. Because

of this inconsistency, Congress directed EPA to re-register all previously registered products, so as to bring their registrations and their data bases into compliance with current requirements [See FIFRA Section 3(g)].

Facing the enormous job of re-reviewing and calling-in new data for the approximately 35,000 current registrations, and realizing the inefficiencies of the 'product-by-product' approach, the Agency decided that a new, more effective method of review was needed.

A new review procedure has been developed. Under it, EPA publishes documents called Registration Standards, each of which discusses a particular pesticide active ingredient. Each Registration Standard summarizes all the data available to the Agency on a particular active ingredient and its current uses, and sets forth the Agency's comprehensive position on the conditions and requirements for registration of all existing and future products which contain that active ingredient. These conditions and requirements, all of which must be met to obtain or retain full registration or re-registration under Section 3(c) (5) of FIFRA, include the submission of needed scientific data which the Agency does not now have, compliance with standards of toxicity, composition, labeling, and packaging, and satisfaction of the data compensation provisions of FIFRA Section 3(c) (1) (D).

The Standard will also serve as a tool for product classification. As part of the registration of a pesticide product, EPA may classify each product for "general use", or "restricted use" when some special regulatory restriction is needed to ensure against unreasonable adverse effects to man or the environment. Many such risks of unreasonable adverse effects can be lessened if expressly-designed label precautions are strictly followed. Thus the special regulatory restriction for a "restricted use" pesticide usually requires that an applicator be certified by the state or Federal government as being competent to use pesticides safely, responsibly, and in accordance with label directions. A restricted-use pesticide can have other regulatory restrictions [40 CFR 162.11(c) (5)] instead of, or in addition to, the certified applicator requirement. These other regulatory restrictions may include such actions as seasonal or regional limitations on use, or a requirement for the monitoring of residue levels after use. A pesticide classified for "general use," or not classified at all, is available for use by any individual who is in compliance with state or local regulations. The Registration Standard review compares information about potential adverse effects of specific uses of the pesticide with risk criteria listed in 40 CFR 162.11(c), and thereby determines whether a product needs to be classified for "restricted use." If the Standard does classify a pesticide for "restricted use," this determination is stated in the Chapter 2.

Requirement to Re-register Under the Standard

FIFRA Section 3(g), as amended in 1978, directs EPA to re-register all currently registered products as expeditiously as possible. Congress also agreed that re-registration should be accomplished by the use of Registration Standards.

Each registrant of a currently registered product to which this Standard applies, and who wishes to continue to sell or distribute his product in commerce, must apply for re-registration. Each application must contain proposed labeling that complies with this Standard.

EPA will issue a notice of intent to cancel the registration of any currently registered product to which this Standard applies if the registrant fails to comply with the procedures for re-registration set forth in the Guidance Package which accompanies this Standard.

"Product-Specific" Data and "Generic" Data

In the course of developing this Standard, EPA has determined the types of data needed for evaluation of the properties and effects of products to which the Standard applies, in the disciplinary areas of Product Chemistry, Environmental Fate, Toxicology, Residue Chemistry, and Ecological Effects. These determinations are based primarily on the data Guidelines proposed in 1978 (43 FR 29686, July 10, 1978, and 43 FR 37336, August 2, 1978), as applied to the use patterns of the products to which this Standard applies. Where it appeared that data from a normally applicable Guidelines requirement was actually unnecessary to evaluate these products, the Standard indicates that the requirement has been waived. On the other hand, in some cases studies not required by the Guidelines may be needed because of the particular composition or use pattern of products the Standard covers; if so, the Standard explains the Agency's reasoning. Data guidelines have not yet been proposed for the Residue Chemistry discipline, but the requirements for such data have been in effect for some time and are, the Agency believes, relatively familiar to registrants. Data which we have found are needed to evaluate the registrability of some products covered by the Standard may not be needed for the evaluation of other products, depending upon the composition, formulation type, and intended uses of the product in question. The Standard states which data requirements apply to which product categories (see the second chapter). The various kinds of data normally required for registration of a pesticide product can be divided into two basic groups:

- (A) data that is "product-specific," i.e., data that relates only to the properties or effects of a product with a particular composition (or a group of products with closely similar composition); and
- (B) "generic" data that pertains to the properties or effects of a particular ingredient, and thus is relevant to an evaluation of the risks and benefits of all products containing that ingredient (or all such products having a certain use pattern), regardless of any such product's unique composition.

The Agency requires certain "product-specific" data for each product to characterize the product's particular composition and physical/chemical properties (Product Chemistry), and to characterize the product's acute toxicity (which is a function of its total composition). The applicant for registration or re-registration of any product, whether it is a manufacturing-use or end-use product, and without regard to its intended use pattern, must submit or cite enough of this kind of data to allow EPA to evaluate the product. For such purposes, "product-specific" data on any product other than the applicant's is irrelevant, unless the other product is closely similar in composition to the applicant's. (Where it has been found practicable to group similar products for purposes of evaluation, with a single set of tests, of all products in the group, the Standard so indicates.) "Product-specific" data on the efficacy of particular end-use products is also required where the exact formulation may affect efficacy and where failure of efficacy could cause public health problems.

All other data needed to evaluate pesticide products concerns the properties or effects of a particular ingredient of products (normally a pesticidally active ingredient, but in some cases a pesticidally inactive, or "inert," ingredient). Some data in this "generic" category are required to evaluate the properties and effects of all products containing the ingredient (e.g., the acute LD-50 of the active ingredient in its technical or purer grade; see proposed 40 CFR 163.81-1(a), 43 FR 37355].

Other "generic" data are required to evaluate all products which both contain a particular ingredient and are intended for certain uses (see proposed 40 CFR 163.82-1, 43 FR 37363, which requires subchronic oral testing of the active ingredient with respect to certain use patterns only). Where a particular data requirement is use-pattern dependent, it will apply to each end-use product which is to be labeled for that use pattern (except where such end-use product is formulated from a registered manufacturing-use product permitting such formulations) and to each manufacturing-use product with labeling that allows it to be used to make end-use products with that use pattern. Thus, for example, a subchronic oral dosing study is needed to evaluate the safety of any manufacturing-use product that legally could be used to make an end-use, food-crop pesticide. But if an end-use product's label specified it was for use only in ways that involved no food/feed exposure and no repeated human exposure, the subchronic oral dosing study would not be required to evaluate the product's safety; and if a manufacturing-use product's label states that the product is for use only in making end-use products not involving food/feed use or repeated human exposure, that subchronic oral study would not be relevant to the evaluation of the manufacturing-use product either.

If a registrant of a currently registered manufacturing-use or end-use product wishes to avoid the costs of data compensation [under FIFRA Section 3(c) (1) (D)] or data generation [under Section 3(c) (2) (B)] for "generic" data that is required only with respect to some use patterns, he may elect to delete those use patterns from his labeling at the time he re-registers his product. An applicant for registration of a new product under this Standard may similarly request approval for only certain use patterns.

Data Compensation Requirements under FIFRA 3(c) (1) (D)

Under FIFRA Section 3(C) (1) (D), an applicant for registration, re-registration, or amended registration must offer to pay compensation for certain existing data the Agency has used in developing the Registration Standard. The data for which compensation must be offered is all data which is described by the following criteria:

- (1) the data were first submitted to EPA (or to its predecessor agencies, USDA or FDA), on or after January 1, 1970;
- (2) the data were submitted to EPA (or USDA or FDA) by some other applicant or registrant in support of an application for an experimental use permit, an amendment adding a new use to a registration, or for re-registration, or to support or maintain in effect an existing registration;
- (3) the data are relevant to the Agency's decision to register or re-register the applicant's product under the Registration Standard, taking into account the applicant's product's composition and intended use pattern(s);
- (4) the data are determined by EPA to be valid and usable in reaching regulatory conclusions; and

- (5) the data are not those for which the applicant has been exempted by FIFRA Section 3(c) (2) (D) from the duty to offer to pay compensation. (This exemption applies to the "generic" data concerning the safety of an active ingredient of the applicant's product, not to "product specific" data. The exemption is available only to applicants who propose to purchase a registered pesticide from another producer in order to formulate such purchased pesticide into an end-use product.)

An applicant for re-registration of an already registered product under this Standard, or for registration of a new product under this Standard, accordingly must determine which of the data used by EPA in developing the Standard must be the subject of an offer to pay compensation, and must submit with his application the appropriate statements showing his compliance with FIFRA Section 3(C) (1) (D).

An applicant would never be required to offer to pay for "product-specific" data submitted by another firm. In most cases, data which are specific to another firm's product will not be useful to the Agency in determining whether the applicant's product is registrable. There may be cases, however, where because of close similarities between the compositions of two or more products, another firm's data may suffice to allow EPA to evaluate some or all of the "product specific" aspects of the applicant's product. In such a case, the applicant may choose to cite that data instead of submitting data from tests on his own product. If he chooses that option, he would have to comply with the offer-to-pay requirements of Section 3(C) (1) (D) for the data.

Each applicant for registration or re-registration of a manufacturing-use or end-use product who is not exempted by FIFRA Section 3(c) (2) (D), must comply with the Section 3(c) (1) (D) requirements with respect to each item of "generic" data that relates to his product's intended uses.

A detailed description of the procedures an applicant must follow in applying for re-registration (or new registration) under this Standard is found in the Guidance Package for this Standard.

Obtaining Data to Fill "Data Gaps"; FIFRA 3(c) (2) (B)

Some of the kinds of data EPA needs for its evaluation of the properties and effects of products to which this Standard applies have never been submitted to the Agency (or, if submitted, have been found to have deficiencies rendering them inadequate) and have not been located in the literature search that EPA conducted as part of preparing this Standard. Such instances of missing but required data are referred to in the Standard as "data gaps".

FIFRA Section 3(c) (2) (B), added to FIFRA by the Congress in 1978, authorizes EPA to require registrants to whom a data requirement applies to generate (or otherwise produce) data to fill such "gaps" and submit these data to EPA. EPA must allow a reasonable period of time for this to be accomplished. If a registrant fails to take appropriate and timely steps to fill the data gaps identified by a Section 3(c) (2) (B) order, his product's registration may be suspended until the data are submitted. A mechanism is provided whereby two or more registrants may agree to share in the costs of producing data for which they are both responsible.

The Standard lists the "generic" data gaps and notes the classes of products to which these data gaps pertain in its summary second chapter. The Standard also points out that to be registrable under the Standard, a product must be supported by certain required "product-specific" data.

As part of the process of re-registering currently registered products, EPA will issue Section 3(c) (2) (B) directives requiring the registrants to take appropriate steps to fill all identified data gaps — whether the data in question is "product-specific" or "generic" — in accordance with a schedule.

Persons who wish to obtain registrations for new products under this Standard will be required to submit (or cite) sufficient "product-specific" data before their applications are approved. Upon registration, they will be required under Section 3(c)(2)(B) to take appropriate steps to submit data needed to fill "generic" data gaps. (We expect they will respond to this requirement by entering into cost-sharing agreements with other registrants who previously have been told they must furnish the data.) The Guidance Package for this Standard details the steps that must be taken by registrants to comply with Section 3(c)(2)(B).

Amendments to the Standard

Applications for registration which propose uses or formulations that are not presently covered by the Standard, or which present product compositions, product chemistry data, hazard data, toxicity levels, or labeling that do not meet the requirements of the Standard, will automatically be considered by the Agency to be requests for amendments to the Standard. In response to such applications, the Agency may request additional data to support the proposed amendment to the Standard, or may deny the application for registration on the grounds that the proposed product would cause unreasonable adverse effects to the environment. In the former case, when additional data have been satisfactorily supplied, and providing that the data do not indicate the potential for unreasonable adverse effects, the Agency will then amend the Standard to cover the new registration.

Each Registration Standard is based upon all data and information available to the Agency's reviewers on a particular date prior to the publication date. This "cut-off" date is stated at the beginning of the second chapter. Any subsequent data submissions and any approved amendments will be incorporated into the Registration Standard by means of addenda, copies of which are available for inspection at EPA in Washington, D.C., and may be requested from the Agency. When all the present "data gaps" have been filled and the submitted data have been reviewed, the Agency will revise the Registration Standard. Thereafter, when the Agency determines that the internally maintained addenda have significantly altered the conditions for registration under the Standard, the document will be updated and re-issued for publication.

While the Registration Standard discusses only the uses and hazards of products containing the designated active ingredient(s), the Agency is also concerned with the potential hazards of some inert ingredients and impurities. Independent of the development of any one Standard, the Agency has initiated the evaluation of some inert pesticide ingredients. Where the Agency has identified inert ingredients of concern in a specific product to which the Standard applies, these ingredients will be pointed out in Chapter 2.

II. REGULATORY POSITION

A. Introduction

This chapter is the central part of the Registration Standard. It presents the Agency's regulatory position based on an evaluation of all registered products containing Deet as the sole active ingredient. After briefly discussing background information on the regulatory history, uses and production of Deet, this chapter explains the Agency's major concerns about the toxicology of Deet, the regulatory actions which will be pursued, and the criteria by which applications for registration of Deet products will be approved. Thus, this chapter contains all of the Agency's requirements or continued registration of Deet products and new product registrations that are covered by the Standard. Detailed analyses of the data upon which this regulatory position is based are presented in each of the disciplinary chapters, III through VII.

B. Background

"Deet" is the common name for N,N-diethyl-m-toluamide, a multi-purpose insect repellent registered for direct application to human skin, clothing, household pets, tents and bedrolls and screens. Deet is a rather unique pesticide, because it is applied directly to the human body for purposes of repelling insects. It was developed and patented by the U.S. Army in 1946 for use by military personnel in insect-infested areas. Because Deet was recognized as one of the few products effective against mosquitoes and biting flies, it was registered for "domestic use" (use by the general public in the U.S.) in 1957. Subsequently, all Deet-containing products have been registered without any restrictions as to the amount or frequency of application to the body.

Deet products currently available to the public are marketed in a variety of liquids, foams, lotions, sprays, and impregnated materials. Formulations registered for direct application to human skin contain from 11.27% to 99.9% Deet as the active ingredient.

C. Regulatory Decision

A review of the very limited data available as of October 24, 1980 indicates that the criteria for unreasonable adverse effects have not been met for products containing Deet (40 CFR 162.11(a)). However, the data do suggest that more complete testing must be required to satisfy specific health effects concerns revealed during the review of Deet.

First, the Deet data base is very incomplete, particularly in the area of chronic effects studies. Further, the available chronic toxicity data were inadequate to assess the long-term effects of Deet, i.e. oncogenicity, teratology, reproduction and mutagenicity. Several supplementary studies (dominant lethal mutagenicity, oral teratology, subchronic inhalation) indicate possible reproductive toxicity. Although the data are only suggestive it is imperative to conduct the necessary studies to resolve the possibility of adverse effects.

A second major concern is that because Deet is applied directly to the human body, an individual's exposure can be very high, especially when applications are repeated daily or even more frequently over a period of time. The available data indicate that about 38% of the population uses an insect repellent. Of this group, many use Deet only occasionally. However, some people apply large doses daily (up to 117.0 mg/kg/day) and repeatedly throughout part of the year. Exposures are even higher for those involved in outdoor work including military personnel (10.7 mg/kg/day up to 60 times per year.)

This profile of widespread and often repeated use creates a high potential for unreasonable adverse effects, especially considering the absence of many chronic studies and considering the questions raised above by existing studies.

Of additional concern is that several acute eye irritation studies demonstrate that certain Deet products are severe eye irritants. Eye irritation studies using technical grade (99.9%) Deet demonstrate that it produces temporary corneal opacity (clouding of the eye) in rabbits. Based on these results, the Agency determines that products containing 99.9% Deet are not registerable for domestic use (see 40 CFR 162.11 (c) (2) (i) (A)). Because the Agency has no eye irritation studies for products containing from 30% to 99.9% Deet, these data must be provided as soon as possible to determine if public health risks exist from use of these products.

Further, an Agency review of Deet confidential Statements of Formula shows that many registrants have not submitted revised Statements demonstrating that Freon propellants have been removed. The Agency reminds registrants that Freons have been banned for use in Deet pressurized liquids because they cause a depletion of the ozone layer (43 FR 11318). Products containing Freons are therefore not registerable under this Standard.

Finally, toxicologic concerns have emerged as to the health effects of pesticides in children. These concerns justify the need to study the dermal absorption rate of Deet in laboratory animals to determine if there is greater absorption through the skin of children than adults.

Based on all these concerns, the Agency has concluded that the following regulatory actions must be initiated immediately:

1. All basic chronic and acute toxicology data for technical and formulated Deet, as well as additional special studies must be submitted. These studies are listed in Tables 1 and 2; protocols for the special studies are available from the Agency. They must be submitted according to the stringent, accelerated schedule shown in the tables.
2. The Agency will prohibit registration or continued registration formulated Deet product which demonstrates acute eye irritation effects of corneal opacity, or eye irritation persisting for 7 days. These products currently include formulations containing 99.9% Deet. The Agency will not change or reduce the requirements for eye irritation testing of Deet, because of its intended use directly on human skin.

3. Deet products are registerable only if all of the terms and conditions of this Standard are met. These conditions are presented in the remainder of this chapter. Products not brought into compliance with the requirements of this Standard will be subject to suspension and/or cancellation. New products registered under this Standard are subject to the same conditions.

4. This Standard will be effective immediately. It will be appropriately revised after all required data have been reviewed and evaluated by the Agency. The Standard may be amended to add certain new uses and formulations not currently registered.

D. Terms and Conditions of Registration

Applicants for registration or continued registration of Deet products subject to this Standard must comply with all terms and conditions described in this Standard, including commitment to fill data gaps according to the schedule specified in Tables 1 and 2. When applicable (see Chapter 1), applicants must offer to pay compensation to the extent required by Sections 3(c) (1) (D) and 3(c) (2) (D) of the Federal Insecticide, Fungicide and Rodenticide Act [FIFRA] as amended, 7 U.S.C. 136 (c) (1) (D) and 136 (c) (2) (D). The following registrants have submitted data in support of Deet registrations, and have not waived their rights to compensation for these data: Cutter Laboratories; Rhone-Poulenc Co.; Charles Pfizer and Company, Inc.; McLaughlin Gormley King Co.; S.C. Johnson and Sons, Inc.; Straight Arrow Co and; Tillar Enterprises.

Following is a summary of the terms and conditions of this standard, including the areas of product composition, acute toxicity, use sites, data requirements and labeling.

Product Composition Standard

a. Technical Deet

To be registered under this Standard, technical Deet must comply with the product composition standard developed in the Product Chemistry chapter. Technical Deet must therefore contain a minimum of 95% active ingredient, with appropriate certification of upper limits of unintentional ingredients as defined in Chapter 3.

b. Formulated Deet

The following types of formulated Deet, with less than 95% active ingredient, and acceptable acute toxicity ratings are acceptable under this Standard:

Ready-to-Use Solution
Pressurized Liquid
Impregnated Material

Applicants for registration of these types of formulations must certify upper limits of impurities as described in Chapter 3, as well as upper and lower limits of inert ingredients.

The Agency finds the inert ingredient, Freon, indentified through a review of Confidential Statements of Formula, unacceptable for use in Deet formulations because of the potential to cause depletion of the ozone layer of the atmosphere (43 FR 11318).

Acute Toxicity Limits

1. Technical Deet

Because technical Deet products are only for manufacturing use, there are no established acute toxicity limits for technical Deet.

2. Formulated Deet

To be registered for domestic use under this Standard, in accordance with FIFRA Section 3(d) (1) (C) (i), formulated (see Table 24) Deet products must have ratings no higher than Toxicity Category III or IV for each of the following acute effects:

- Acute Oral Toxicity
- Acute Dermal Toxicity
- Acute Inhalation Toxicity
- Primary Dermal Irritation

Additionally, formulated Deet products must not demonstrate eye irritation persisting for seven days or any corneal opacity in test animals.

Use Patterns

1. Technical Deet

To be registered under this Standard, technical Deet products may be repackaged or formulated only into end-use, insect repellents for domestic use.

2. Formulated Deet

To be registered under this Standard, formulated Deet products shall be labelled for use only as insect repellents for application to human skin, household pets, clothing, bedding, tents, and screens. Application to food or livestock shall be prohibited by label direction.

Data Requirements

Applicants for registration of technical or formulated Deet products must cite or submit the data listed in Tables 1 and 2, respectively, by specified due dates. Data in this Standard that satisfy registration requirements may be cited, if the applicant establishes that the proposed product is substantially similar (as determined by percentage of active and inert ingredients, manufacturing impurities and different uses) to another product for which the Agency has received acceptable data. If data in this Standard are cited, compensation must first be offered to the submitter(s) of the data as explained in Chapter 1. The Agency will examine both active and inert ingredients to determine if products are similar. In front of each requirement in Tables 1

and 2 is listed the section of the Proposed Guidelines (43 FR 29696, July 10, 1978; and 43 FR 37336, August 22, 1978) which describe that type of data and when it is required. Applicants must submit, solely or through joint agreement, all information in the tables identified as data gaps.

Applicants will not be required to submit residue chemistry data because Deet has no food uses and is not expected to occur indirectly as a food residue.

Required Labeling

All technical and formulated Deet products must bear appropriate labeling as specified in the Guidance Package which accompanies this Standard.

1. Technical Deet

All technical Deet products must list on the label the intended end-use of formulated products produced from the technical products. Therefore, in addition to basic labeling, all technical Deet product labels must bear the following statement:

"For Formulation into End-Use Insect Repellent Products Intended Only for Domestic, Nonfood Use."

2. Formulated Deet

In addition to the basic labeling requirements, to reduce the possibility of inadvertent application to food and livestock, all formulated Deet products must bear the following statement:

"Do Not Apply Near Food or to Livestock"

All formulated Deet products must bear an appropriate statement of practical first-aid treatment in case of accidental ingestion or eye contact.

For formulated Deet products which present label claims that the product is effective against insects which might affect public health, the Agency will require registrants to submit data to support these claims. All labels which claim that Deet repels the following pests, must be supported by company data or published literature:

- biting flies (black fly, sandfly, horsefly and Ceratopogonid species only)
- chiggers
- deerflies
- fleas
- leaches
- mosquitoes
- stableflies
- ticks

Data required by the Agency in support of Deet's efficacy are listed in Table 3. These data are explained in detail in Chapter VII.

PRODUCT-SPECIFIC DATA REQUIREMENTS FOR TECHNICAL PRODUCTS

Product-Chemistry

Table 1

Guideline Citation	Name of Test	Composition Characteristics	Does EPA Have Data To Satisfy This Requirement ?	Bibliographic Citation	Must Data Be Submitted Under FIFRA 3(c) (2) (B) ?	Time Allowed Before Submission
163.61-3	Product identity and disclosure of ingredients	Minimum 95% active ingredient	No	-	Yes	6 months
-4	Description of manufacturing process	"	Yes--incomplete	Pfizer, Inc., 1976 MRID #00001036; Bogard, T., 1976 MRID #GS0002014	Yes	24 months
-5	Discussion on formation of unintentional ingredients	"	No	-	Yes	24 months
-6	Declaration and certification of ingredients limits	"	No	-	Yes	6(24mos.)*
-7	Product analytical methods and data	"	No	-	Yes	6 months
163.61-8(1)	Color	"	Yes	Hercules Powder Co, 1957 MRID #00001025	No	-
-8(2)	Odor	"	Yes	S.C. Johnson & Son, Inc., 1977 MRID #00001100	No	-
-8(4)	Solubility	"	No	-	-	6 months
-8(5)	Stability	"	Yes	S.C. Johnson & Son, Inc., 1977 MRID #00001100	No	-
-8(6)	Octanol/Water partition coefficient	"	No	-	Yes	6
-8(7)	Physical State	"	Yes	Hercules Inc., 1977 MRID # 00001060	No	-

* Any impurities at 0.1% or less-Information need not be submitted until 24 mos.

PRODUCT-SPECIFIC DATA REQUIREMENTS | TECHNICAL PRODUCTS

Product Chemistry

Table 1 Cont'd

Guideline Citation	Name of Test	Composition Characteristics	Does EPA Have Data To Satisfy This Requirement ?	Bibliographic Citation	Must Data Be Submitted Under FIFRA 3(c)(2)(B) ?	Time Allowed Before Submission
-8(8)	Specific gravity	Minimum 95% active ingredient	Yes	Hercules Inc., 1977 MRID # 00001008	No	-
-8(9)	Boiling point	"	Yes	S.C. Johnson & Son, Inc., 1977 MRID #00001100	No	-
-8(10)	Vapor pressure	"	Yes	Blaine et al., 1974 MRID # 05002553	No	-
-8(11)	pH	"	No	-	Yes	6 months
-8(12)	Storage stability	"	No	-	Yes	6 months
-8(13)	Flammability	"	Yes	S.C. Johnson & Son, Inc., 1977 MRID # 00001100 Hercules, Inc., 1972 MRID #00001152	No	-
-8(14)	Oxidizing or reducing action	"	No	-	Yes	6 months
-8(15)	Explosiveness	"	No	-	Yes	6 months
-8(17)	Viscosity coefficient	"	Yes	S.C. Johnson & Son, Inc., 1977 MRID #00001100 Hercules, Inc., 1972 MRID # 00001152	No	-
-8(18)	Corrosion characteristics	"	Yes	Hercules Powder Co., 1957 MRID # 00001025	No	-

GENERIC DATA REQUIREMENTS FOR TECHNICAL PRODUCTS

Environmental Fate

Table 1 cont'd.

Guidelines Citation	Name of Test	Composition	Does EPA Have Data To Satisfy This Requirement?	Bibliographic Citation	Must Data Be Submitted Under FIFRA 3(c) (2) (B)?	Time Allowed Before Submission
163.62-7(c)	Hydrolysis	Minimum 95% active ingredient	No	-	Yes	6 months
163.62-8(g)	Activated sludge metabolism	"	No	-	Yes	6 months
See Chapter 4	Human use and exposure data	"	No	-	Yes	12 months

GENERIC DATA REQUIREMENTS FOR PESTICIDAL PRODUCTS

Toxicology

Table 1 Cont'd

Guideline Citation	Name of Test	Composition Characteristics	Does EPA Have Data to Satisfy This Requirement ?	Bibliographic Citations	Must Data Be Submitted Under FIFRA 3 (c) (2) (B) ?	Time Allowed Before Submission
163.82-3	Subchronic 90-Day Dermal Toxicity	Minimum 95% Active Ingredient	No	-	Yes	12 months
163.82-4	Subchronic Inhalation Toxicity	"	Yes	U.S. Army, 1980 MRID # GS0002032	No	-
163.83-1	Chronic Dermal Toxicity	"	No	-	Yes	4 1/2 years
163.83-2	Dermal Oncogenicity (Mouse)	"	No	-	Yes	4 1/2 years
163.83-3	Dermal Teratogenicity (Rat)	"	No	-	Yes	1 year
163.83-4	Dermal Reproduction (Rat)	"	No	-	Yes	3 1/2 years
163.84-2 through -4	Mutagenicity	"	No	-	Yes	12 months
163.85-1	Metabolism (Identification of Metabolites)	"	No	-	Yes	18 months
Protocols will be furnished by the Agency	Age-Related Dermal Absorption Studies (Rats)	"	No (Special Testing)	-	Yes	18 months

GENERIC DATA REQUIREMENTS FOR TECHNICAL PRODUCTS

Toxicology

Table 1 Cont'd

Guideline Citation	Name of Test	Composition Characteristics	Does EPA Have Data to Satisfy This Requirement ?	Bibliographic Citations	Must Data Be Submitted Under FIFRA 3 (c) (2) (B)?	Time Allowed Before Submission
163.81-1	Acute Oral Toxicity	Minimum 95% Active Ingredient	Yes	See Table 8	No	-
163.81-2	Acute Dermal Toxicity	"	Yes	See Table 11	No	-
163.81-3	Acute Inhalation Toxicity	"	Yes	U.S. Army 1979 MRID #GS0002034	No	-
163.81-4	Primary Eye Irritation	"	Yes	U.S. Army 1979 MRID #GS0002025	No	-
163.81-5	Primary Dermal Irritation	"	Yes	See Table 15	No	-
163.81-6	Dermal Sensitization	"	Yes	U.S. Army 1979 MRID #GS0002026	No	-

GENERIC DATA REQUIREMENTS FOR PESTICIDAL PRODUCTS

Ecological Effects

Table 1 Cont'd

Guideline Citation	Name of Test	Composition	Does EPA Have Data to Satisfy this Requirement?	Bibliographic Citation	Must Data Be Submitted Under FIFRA 3(c) (2) (B)	Months Allowed Before Submission
163.71-1	Avian single-dose oral LD 50 (wild waterfowl, preferably mallard duck)	Minimum 95% active ingredient	No	-	Yes	6 months
163.72-1	Fish acute LC 50 (coldwater species)	"	Yes	McCann, J.A. (1972) MRID # 00001026	No	6 months
163.72-1	Fish acute LC 50 (warmwater species)	"	No	-	Yes	6 months
163.72-2	Acute toxicity to aquatic invertebrates	"	No	-	Yes	6 months

PRODUCT-SPECIFIC DATA REQUIREMENTS FOR FORMULATED PRODUCTS

Product Chemistry

Table 2

Guideline Citation	Name of Test	Composition Characteristics	Does EPA Have Data To Satisfy This Requirement ?	Bibliographic Citation	Must Data Be Submitted Under FIFRA 3(c) (2) (B) ?	Time Allowed Before Submission
163.61-3	Product identity and disclosure of ingredients	Any percentage active ingredient	No	-	Yes	6 months
-4	Description of manufacturing process	"	No	-	Yes	24 months
-5	Discussion on formation of unintentional ingredients	"	No	-	Yes	24 months
-6	Declaration and certification of ingredients limits	"	No	-	Yes	6 (24mos.)*
-7	Product analytical methods and data	"	No	-	Yes	6 months
163.61-8(1)	Color	"	No	-	Yes	6 months
-8(2)	Odor	"	No	-	Yes	6 months
-8(8)	Density or specific gravity	"	No	-	Yes	6 months
-8(9)	Boiling point	"	No	-	Yes	6 months
-8(10)	Vapor pressure	"	No	-	Yes	6 months
-8(11)	pH (Manufacturing-use product and the technical)	"	No	-	Yes	6 months
-8(12)	Storage stability	"	No	-	Yes	6 months
-8(13)	Flammability	"	No	-	Yes	6 months
-8(17)	Viscosity	"	No	-	Yes	6 months
-8(18)	Corrosion characteristics	"	No	-	Yes	6 months

* For impurities at 0.1% or less, information need not be submitted until 24 mos.

PRODUCT-SPECIFIC DATA REQUIREMENTS FOR FORMULATED PRODUCTS

Environmental Fate

Table 2 cont'd

Guideline Citation	Name of Test	Composition Characteristics	Does EPA Have Data to Satisfy This Requirement ?	Bibliographic Citations	Must Data Be Submitted Under FIFRA 3(c) (2) (B)?	Time Allowed Before Submission
163.62-7 (c)	Hydrolysis	Any percentage active ingredient	No	-	Yes	6 months

PRODUCT - SPECIFIC DATA REQUIREMENTS FOR FORMULATED PRODUCTS

Toxicology

Table 2 cont'd

Guideline Citation	Name of Test	Composition Characteristics	Does EPA Have Data to Satisfy This Requirement ?	Bibliographic Citations	Must Data Be Submitted Under FIFRA 3(c) (2) (B) ?	Time Allowed Before Submission
		1. All Ready-to-use				
163.81-1	Acute Oral Toxicity	"	Yes	Warf 1976 MRID 00001101	No	-
163.81-2	Acute Dermal Toxicity	"	Yes	Davidson 1968 MRID # 00001139	No	-
163.81-3	Acute Inhalation Toxicity	"	Yes *	-	No	-
163.81-4	Primary Eye Irritation (Rabbit)	"	+ Some (Incomplete)	Davidson 1968 MRID # 00001139 Warf 1976 MRID # 00001101	Yes	6 months
163.81-5	Primary Dermal Irritation (Rabbit)	"	+ Some (Incomplete)	Davidson 1969 MRID # 00001139 Warf 1976 MRID 00001101	Yes	6 months
163.81-6	Dermal Sensitization (Guinea Pig)	"	+ Some (Incomplete)	Ambrose 1959 MRID # 00001051	Yes	9 months
		2. All Pressurized Liquid				
163.81-1	Acute Oral Toxicity	"	Yes	Warf 1975a MRID 00001085 Warf 1975b MRID 00001086 Howard 1971 00001080	No	-
163.81-2	Acute Dermal Toxicity	"	Yes*	-	No	-
163.81-3	Acute Inhalation Toxicity	"	Yes	Warf 1975 MRID 00001086	No	-
163.81-4	Primary Eye Irritation (Rabbit)	"	Some	Warf, 1975b MRID # 00001085 Durlow & Woodward 197b MRID 00001082	Yes	6 months

* This information has been extrapolated from other tests, and is adequate without further testing.

+ See Chapter 5, Topical Discussions.

PRODUCT - SPECIFIC DATA REQUIREMENTS FOR FORMULATED PRODUCTS

Toxicology

Table 2 cont'd

Guideline Citation	Name of Test	Composition Characteristics	Does EPA Have Data to Satisfy this Requirement	Bibliographic Citations	Must Data Be Submitted Under FIFRA 3(c) (2) (B)?	Time Allowed Before Submission
163.81-5	Primary Dermal Irritation (Rabbit)	Pressurized Liquid	Some	Johnson 1972 MRID # 00001073 Ware 1975a MRID # 00001085 Ware 1975b MRID # 00001086 Durlco & Woodland 9171b MRID # 00001081	Yes	6 months
163.81-6	Dermal Sensitization (Guinea Pig)	Pressurized Liquid	No	-	Yes	9 months
		3. All Impregnated Materials				
163.81-1	Acute Oral Toxicity	"	Yes *	-	No	-
163.81-2	Acute Dermal Toxicity	"	Yes *	-	No	-
163.81-3	Acute Inhalation Toxicity	"	Yes *	-	No	-
163.81-4	Primary Eye Irritation (Rabbit)	"	+ Some (Incomplete)	-	Yes	6 months
163.81-5	Primary Dermal Irritation (Rabbit)	"	No	-	Yes	6 months
163.81-6	Dermal Sensitization (Guinea Pig)	"	No	-	Yes	9 months

* This information has been extrapolated from other tests, and is adequate without further testing.

+ See Chapter 5, Topical Discussions.

PRODUCT-SPECIFIC DATA REQUIREMENTS FOR FORMULATED PRODUCTS

Efficacy

Table 3

Target Pest	Acceptable Use Sites (No Data Required)	Use Sites for which Data are Required	Must Data be Submitted to Support Duration Claims?
Biting Flies (black flies, sand-flies and Ceratopogonidae);	Skin Clothing Tents and Bedrolls Screens	Outdoor Mist	Yes
horseflies	None	Skin, Clothing, Tents and Bedrolls, Screens	Yes
Chiggers	Skin	*Clothing *Tents and Bedrolls Outdoor mist	Yes
Deerflies	Skin Clothing Tents and Bedrolls	Outdoor Mist	Yes
Fleas	Skin Clothing Tents and Bedrolls	Outdoor Mist	Yes
Leeches	Skin Clothing Tents and Bedrolls	Outdoor Mist	Yes
Mosquitoes	Skin Clothing Tents and Bedrolls Screens	Outdoor Mist	Up to 2 hrs. may be claimed without data for acceptable sites.
Stable flies	Skin Clothing Tents and Bedrolls	Outdoor Mist	Up to 2 hrs. may be claimed without data for acceptable sites.

PRODUCT-SPECIFIC DATA REQUIREMENTS FOR FORMULATED PRODUCTS

Efficacy

Table 3 Con't

Target Pest	Acceptable Use Sites (No Data Required)	Use Sites for which Data are Required	Must Data be Submitted to Support Duration Claims?
Ticks	None	Skin Clothing Tents and Bedrolls Outdoor Mist	Yes

* Submit calculations which demonstrate that the product will deposit a minimum of 1 gm. AI/sq. ft., of treated material when used properly.

III. PRODUCT CHEMISTRY

A. Introduction

FIFRA 3(c) (2) (A) requires the Agency to establish guidelines for registering pesticides in the United States. The Agency requires registrants to provide quantitative data on all added ingredients, active and inert, which are equal to or greater than 0.1% of the product by weight.

To establish the composition of products proposed for registration, the Agency requires data and information not only on the manufacturing and formulation processes, but also a discussion on the formation of manufacturing impurities and other product ingredients, intentional and unintentional. Further, to assure that the composition of the product as marketed will not vary from the composition evaluated at the time of registration, applicants are required to submit a statement certifying upper and lower composition limits for the added ingredients, and the upper limits only for some unintentional ingredients. Subpart D (43 FR 29696, July 10, 1978) suggests specific precision limits for ingredients based on the percentage of ingredient and the standard deviation of the analytic method.

In addition to the data on product composition, the Agency also requires data to establish the physical and chemical properties of both the pesticide active ingredient and its formulations. For example, data are needed concerning the identity and physical state of the active ingredient (e.g., melting and boiling point data, ambient vapor pressure and solubility). Data are also required on the properties of the formulated product to establish labeling cautions (e.g., flammability, corrosivity or storage stability). The Agency uses these data to characterize each pesticide and to determine its environmental and health hazards.

B. Disciplinary Review

1. Chemistry Profile

Deet, (N,N-diethyl-m-toluamide), is an all-purpose individual insect repellent which contains a minimum of 95% of the meta isomer, the most effective form of diethyl toluamide, as a technical active ingredient.

Technical Deet is a nearly colorless liquid with a faint characteristic odor. It is relatively stable, highly hygroscopic and sensitive to light. Technical Deet is practically insoluble in water and glycerin but miscible with several organic solvents. It has a specific gravity of 0.990-1.000 at 25°C, a boiling point of 111°C at 1 mm Hg, and a vapor pressure of 1.67×10^{-3} mm Hg at 25°C.

Technical Deet, as it is considered under this standard, is a "manufacturing-use product" which is intended for (re)formulation or repackaging for end use as a "formulated product." End use products are formulated as solutions, lotions, gels, aerosol sprays, sticks and impregnated towlettes. Although several multiple active ingredient products containing Deet are currently registered, this Registration Standard covers only those products which contain Deet as the single active ingredient. The amount of Deet in these products ranges from 11.27% - 100% of the product composition by weight.

2. Generic Data Gaps

All of the generic data needed to evaluate the continued registerability of products to which this Standard applies are listed in Tables 1 and 2, Chapter 2.

3. Required Labeling

All technical and formulated Deet products must bear appropriate product chemistry labeling as specified in the Guidance Package which accompanies this Standard.

C. Topical Discussions

In accordance with each of the topical discussions listed below, a detailed explanation of the minimum data that the Agency requires in order to adequately assess a pesticide's product chemistry can be found in the "Proposed Guidelines for Registering Pesticides" of July 10, 1978 (43 FR Part 163.61-2).

<u>Data Requirement</u>	<u>Guidelines Section</u>
<u>Technical Deet</u>	
Chemical Identity	163.61-3
Manufacturing Process	163.61-4
Discussion of Formation of Pesticide Products	163.61-5
Percentages of Components in Unintentional Ingredients	163.61-6
Product Analytical Methods and Data	163.61-7
Physical/Chemical Properties-	
Color	163.61-8(c)1
Odor	163.61-8(c)2
Solubility (in quantitative terms)	163.61-8(c)4
Stability	163.61-8(c)5
Octanol/Water Partition Coefficient	163.61-8(c)6
Physical State	163.61-8(c)7
Specific Gravity	163.61-8(c)8
Boiling Point	163.61-8(c)9
Vapor Pressure	163.61-8(c)10
Viscosity	163.61-8(c)17
Corrosion Characteristics	163.61-8(c)18
pH Measurement	163.61-8(c)11
Storage stability	163.61-8(c)12
Flammability (flash point by closed cup method)	163.61-8(c)13
<u>b. Formulated Deet</u>	
Chemical Identity	163.61-3
Manufacturing Process	163.61-4
Discussion on Formation of Unintentional Ingredients	163.61-5
Percentages of Components in Pesticide Products	163.61-6
Product Analytical Methods and Data	163.61-7
Physical/Chemical Properties-	
Color	163.61-8(c)1
Odor	163.61-8(c)2

Physical State	163.61-8(c)7
Density or specific gravity	163.61-8(c)8
Boiling point	163.61-8(c)9
Vapor pressure	163.61-8(c)10
pH Measurement	163.61-8(c)11
Storage stability	163.61-8(c)12
Flammability (flashpoint, flame extension)	163.61-8(c)13
Viscosity	163.61-8(c)17
Corrosion characteristics	163.61-8(c)18


a. Technical Deet

(1) Chemical Identity

"Deet" is the acceptable common name of the Entomological Society of America for N,N-diethyl-m-toluamide. The American National Standards Institute does not recognize Deet as the common name for N,N-diethyl-m-toluamide.

The name "Deet" will be routinely used in this Registration Standard in lieu of the more complex chemical name or of its trade names. Trade names for Deet include: Metadelphene[®] (Hercules, Inc.), OFF[®] (S.C. Johnson & Son, Inc.), and MGK Diethyltoluamide (McLaughlin-Gormley-King Co.).

Deet is a diethyltoluamide consisting primarily of 95% of the meta isomer; it is an all-purpose individual insect repellent. Its molecular configuration is shown below:



(2) Manufacturing Process

In general, the open literature describes four methods of synthesizing N,N-diethyl-toluides:

- acid-chloridizing m-toluic acid and then reacting with diethylamine in the presence of a catalyst in an organic solvent;
- reacting m-toluic acid and diethylamine with an appropriate catalyst in a continuous vapor-phase;
- reacting m-toluic acid and diethylamine in an anhydrous acetic acid medium; and
- reacting m-toluic acid with diethylamine in the presence of N-mono-ethyl-m-toluamide under pressure and raised temperature.

In addition, there are several other means of producing N,N-diethyl-m-toluamide in a laboratory; however, these procedures are not economical for continuous large scale industrial operation.

No data were submitted concerning the specific procedures, equipment and manufacturing conditions required for commercial production of the chemical.

(3) Discussion on Formation of Unintentional Ingredients

The chemical reactions employed in the synthesis and purification of any active ingredient might cause potentially harmful impurities to be produced. The presence of manufacturing impurities is dependent upon the nature of the manufacturing process used.

Data considered as "Confidential Business Information" concerning the formation of each substance that might reasonably be identified as present in technical Deet were reported by McLaughlin, Gormley, King Company and Pfizer, Inc. The data fulfill requirements for these particular registrants. A data gap exists for other registrants.

(4) Percentages of Components in Pesticide Products

Technical Deet contains a minimum of 95% of the active ingredient, N,N-diethyl-m-toluamide. Depending upon the manufacturing process used, up to about 5% of technical impurities (including moisture) can be expected.

(5) Product Analytical Methods and Data

Although some of the manufacturing impurities of technical Deet have been identified by its manufacturers (McLaughlin, Gormley, King, Company, 1976, MRID # GS0002014); Pfizer Inc., 1976, 00001036), the methods that were used in the analyses were not described.

(6) Physical/Chemical Properties

Color: Not darker than Hazen number 100 (Hercules Powder Co., 1957 MRID #00001025).

Odor: Nearly odorless, or has a characteristic faint odor depending on the purity of technical Deet and the kind of manufacturing process used (S.C. Johnson & Son, Inc., 1977 MRID # 00001100).

Solubility: The solubility of technical Deet was reported as follows (Cutter Laboratories, 1975 MRID#00001008):

- practically insoluble in water
- practically insoluble in glycerin
- miscible with ethanol
- miscible with ether
- miscible with isopropanol
- miscible with chloroform
- miscible with carbon disulfide

The proposed guidelines state that solubility, preferably in g/100 ml of solvent at 20°C (68°F), or reported in other terms such as ppm (mg/kg) is required for the technical chemical.

Stability: Technical Deet is a relatively stable compound. It is highly hygroscopic and sensitive to light. When stored in a closed container protected from light it is regarded as being stable for at least one year (S.C. Johnson & Son, Inc., 1977 MRID #00001100).

Octanol/Water Partition Coefficient: No coefficient has been reported.

Physical State: Technical Deet is a liquid (Hercules, Inc., 1977 MRID #00001069).

Specific Gravity: The specific gravity of technical Deet ranges from 0.990 - 1.000 at 25°C (Hercules, Inc., 1977 MRID #0001068).

Boiling Point: At 1 mm Hg. the boiling point of technical Deet is 111°C (S.C. Johnson & Son, Inc. MRID #00001100; Blaine, R.L. et al., 1974 MRID #05002553).

Vapor Pressure: 1.67×10^{-3} mm Hg at 25°C (Blaine, R.L. et al., 1974 MRID #05002553).

pH: none reported

Storage Stability: none reported

Flammability: The flashpoint (open cup) for technical Deet ranges from 150°C - 155°C (Hercules, Inc., 1972 MRID # 00001152; S.C. Johnson & Son, Inc., 1977 MRID# 00001100).

Viscosity: The viscosity for technical Deet was reported as 13.3 centipoises at 30°C (Hercules, Inc., 1972 MRID # 00001152; S.C. Johnson & Son, Inc., 1977 MRID# 00001100).

Corrosion Characteristics: Technical Deet has no corrosive action on most metals; however, metal containers may be corroded due to the presence of one or more of the other ingredients in the formulated product. To prevent the possibility of corrosion, a three-quarter or one-pound tin plate is recommended. Because excessive moisture can cause corrosion of pressurized containers, Deet is made available almost entirely free of water. As a hygroscopic agent, Deet should be protected from possible moisture pick-up in handling (Hercules Powder Co., 1957 MRID #00001025).

Submittal of Samples: Applicants for registration or reregistration will be notified at the time of application with regard to the submission of samples.

b. Formulated Deet

(1) Chemical Identity

There are no data available on the chemical identity of Deet formulations. This constitutes a data gap.

(2) Manufacturing Process

There are no data available on the manufacturing process of Deet formulations. This constitutes a data gap.

(3) Discussion on Formulation of Unintentional Ingredients

No data have been submitted. This constitutes a data gap.

(4) Percentages of Components in Pesticides Products

Data on currently registered formulated Deet products are listed below.

<u>Type of Formulation</u>	<u>% Active Ingredient</u>
Solution, Ready-to-Use	15.0 - 99.9%
Pressurized Liquid	12.5 - 75.0%
Impregnated Material	11.7 - 38.8 %

The compositional data as to inert ingredients will not be discussed because of the extensive number of Deet products. The data were given full consideration by the Agency in developing this Registration Standard.

(5) Product Analytical Methods and Data

No method has been submitted for determining or measuring any of the impurities in Deet products; however, the literature indicated the identification and measurement of these impurities by thin layer and gas chromatography (Voronkina, T.M., et al., 1971, 05000597).

Methods for detecting and measuring the active ingredient Deet in registered products have been submitted (Sarmienta and Beroza, 1975, 05000048; Cutter Laboratories, 1975, 00001008; Pfizer, Inc, 1976, 00001036).

(6) Physical/Chemical Properties

No data pertaining to the physical/chemical properties of Deet products were submitted. Accordingly, none of the minimum data requirements listed at the beginning of the Topical Discussions are satisfied and this constitutes a data gap.

IV. ENVIRONMENTAL FATE

A. Disciplinary Review

1. Use Profile

The major use site for Deet is human skin. Formulations registered for such use are labeled to warn against contact with eyes or lips; pressurized liquids are not to be sprayed directly into the face.

Deet is also registered for application to household pets. Labels advise against contact with eyes, lips and open wounds.

Clothing, tents and bedrolls may be treated with Deet formulations registered for these uses. Labels advise against application to rayon, spandex, dynel and verel fabrics.

Other registered uses for Deet are application to screens and screen doors. Labels on formulations intended for these uses caution against application to plastics, painted surfaces and other finishes.

Although all of the above uses can occur outdoors, residues occurring in the environment after application will be considered insignificant for purposes of this Standard. Likewise, while Deet can be applied around food in picnic and camping areas, residues on food items are considered insignificant for purposes of this Standard.

2. Environmental Fate Profile

Registered outdoor uses of Deet are not expected to result in the introduction of significant amounts of this pesticide into the environment. However, technical Deet can potentially enter the aquatic environment through indirect or accidental discharge into lakes, streams or wastewater treatment systems. The data necessary, but currently not available, for an evaluation of technical Deet's effect on the aquatic environment are hydrolysis and activated sludge metabolism studies; formulated Deet can also potentially enter wastewater treatment systems and for this reason hydrolysis studies are necessary. The only portion of an environmental fate profile necessary to this Standard is an evaluation of these effects.

3. Exposure Profile

a. Technical Deet

For persons involved in the manufacture, handling, storage or shipment of technical Deet, this Standard will not include an assessment of hazard because the responsibility of establishing standards for such exposure falls within the jurisdiction of the Occupational Safety and Health Administration.

b. Formulated Deet

Ready-to-use solutions, pressurized liquids and impregnated materials containing Deet are registered for use on all exposed human skin areas. It is estimated that this use accounts for 99% of the single active ingredient Deet produced (Hales, Y. and Radtke, 1980 MRID #GS0002001).

In 1978, an estimated 38% of the total U S population used an insect repellent, although not all of these products contained Deet (S.C. Johnson and Co. 1979 MRID #GS0002002). Highest use of insect repellent was reported in the Southwest (49%) and lowest on the Pacific Coast (27%). Of the various products used (Deet containing and otherwise), 73% were pressurized liquids, 19% were ready-to-use solutions, and 8% were impregnated materials. Deet concentrations in the pressurized liquid products (the most widely used) ranged from 15 to 20%. Based on these data, it is estimated that 22% of the general population is exposed to pressurized liquid products that contain 15 to 20% Deet as an active ingredient. Remaining usage is divided among other types of products and pressurized formulations.

An Insecticide User Profile prepared by S.C. Johnson Co, Inc. in 1975 provided most of the available data on exposure of the general population to Deet (MRID #GS0002003). The study is not considered to be fully representative of the U.S. population because respondents were all S.C. Johnson employees. Nevertheless, these data are useful in the absence of other information about the extent of use of Deet by the general population.

Based on the S.C. Johnson Co. data, average usage of Deet products by the general population will result in an average daily exposure to active ingredient ranging from 0.944 g (for a 20% a.i. product) to 3.5 g (for a 75% product). This is equivalent to 13.5 - 50.6 mg/kg/day for a 70 kg man and 15.7-59 mg/kg/day for a 60 kg woman. The Johnson data also estimated usage by 90% or less of the general population to be 1.06g and 3.98g per day for 20% and 75% products, respectively. This is equivalent to 15.0 - 56.8 mg/kg/day for a 70 kg man and 17.7-66.3 mg/kg/day for a 60 kg woman.

Additional data provided by S.C. Johnson for a 15% pressurized liquid formulation related grams of product applied on the users forearms to the number of users applying a specific amount of the product. These data, based on a survey of 71 people, indicate that 90% of those interviewed used 1.4 grams of product or less per forearm and 99% used 2.9 grams or less per forearm. From these data the Agency calculated, through extrapolation, that 99% of users are exposed to 11.0 g/day or less of formulation applied to all exposed skin. This is equivalent to a daily exposure of 1.65 grams of Deet or 23.5 mg/kg/day for a 70 kg man using a 15% formulation. If a 75% Deet product were used at the 11.0 g/day rate assumed above, the exposure would 8.23 g/day of active ingredient or 117.6 mg/kg/day for a 70 kg man.

Thus, based on the S.C. Johnson survey, 1% of users may be exposed to greater than 1.65 g/day assuming use of a 15% product. In the case of a 75% product used at the rate of 11 g/day, exposure would be 5 times that of the 15% product, or 8.25 g/day/ of active ingredient.

The Agency recognizes that Deet products composed solely of technical Deet are also available for use by the general population. Although no data are available on use rates of these products, the Agency concludes that exposures will be proportionately higher than those presented above if the products are applied with the same frequency.

It must be noted that all of the above calculations reflect Deet exposure at the surface of the skin only. Data revealed by Agency toxicologists, and explained in detail in Chapter V, show that 10% of Deet applied directly to the

body is absorbed through the skin. All of the available exposure data on the general population and the Agency's estimates based on these data are summarized in Table 4.

Table 4
Estimated Exposure to Deet

Formulation (% Deet)	Average Daily Exposure ^{1/} gram/day mg/kg/day		Maximum use by 90% of Subjects ^{2/} gram/day mg/kg/day		Maximum use by 99% of Subject (Based on 11.0 gr product/day) grams/day mg/kg/day	
15%	ND	ND	ND	ND	1.65	23.40
20%	0.944	13.50 * 15.70	1.06	15.00 * 17.70	ND	ND
30%	1.42	20.23	1.59	22.70	ND	ND
50%	2.36	33.70	2.65	37.80	ND	ND
75%	3.54	50.60 * 59.00	3.98	56.8 * 66.3	+8.23	+117.6

^{1/} Based on use of 16-20 g. of formulation per 4-week period (S.C. Johnson & Son, Inc. 1979 MRID # GS0002002).

^{2/} Based on use of 48 g. of formulation per 4-week period (S.C. Johnson & Son, Inc. 1979 MRID #GS0002002).

* Figures assume a 60 Kg woman. All other figures assume a 70 Kg man.

Figures assume a 60 Kg woman. All other figures assume a 70 Kg man.

+ Based on extrapolation from data on application to forearm only.

ND = No data available

Several occupational groups are expected to experience exposures higher than those of the general population. Military personnel have been identified as a high-exposure group based on an estimated usage of 1 ml of a 75% formulation approximately 60 times per year. This is equivalent to a total annual exposure of 43 grams of active ingredient to each of approximately 628,000 exposed individuals, or 10.7 mg/kg/day (Bales, Y and H.E. Radtke 1980, MRID #GS0002001).

Other high exposure groups include persons engaged in such outdoor occupations such as forestry, research, fishing, lumbering, park and refuge maintenance, outdoor sports and gardening. The only quantitative exposure data submitted

relate to a research biologist in the Florida Everglades, who reported using 2 one-ounce bottles of 28.74% Deet per week on the skin, and 2 two-ounce cans of 71.25% Deet per week on clothing (Mazzotti, F. 1980 MRID #GS0002004).

Applications were made 4 days per week from May to October. Based on these data, the Agency estimates that exposure is 4.25 g/day or 60.7 mg/kg/day, assuming a body weight of 70 Kg. Application to clothing, also based on a 4-day week, would be 20.2 g/day; an undetermined amount of the Deet applied to clothing would be transferred to the skin. These exposure estimates are almost three times as great as the maximum daily exposure to 90% of the general population reported by S.C. Johnson Co. for a 30% active ingredient product (63.4 mg/kg/day vs. 22.7 mg/kg/day).

In conclusion, an average exposure to Deet active ingredient is probably in the range of 10-20 mg/kg/day, with 90% of the population exposed to less than about 35 mg/kg/day. It has been estimated that about 1% of users apply 11.0 gm or more per day to exposed skin areas. Thus, use of a 20% formulation would result in an exposure of 31 mg/kg/day; for a 75% formulation, 117.6 mg/kg/day; and for a 100% active product, 157 mg/kg/day. The above figures for exposure are only estimates based on limited data for the numbers of grams of products applied by users. Few data were available on the actual formulations used, although pressurized liquid products containing 15-20% active ingredient are claimed to be the most popular. The Agency acknowledges that the above estimates are based on limited data taken from a small sample group; however, they represent all the data currently available.

Occupational exposures could not be fully evaluated, although available data indicate that military personnel may experience an annual exposure of up to 10.7 mg/kg/day and a research biologist in the Florida Everglades is exposed to 63.4 mg/kg/day. Although these exposure levels to military personnel appear to be about the same as those of the general population, they occur with greater frequency throughout the year; thus, the total annual exposures are likely to be much greater.

Before a complete exposure profile is prepared, additional human exposure data will be required, under Data Gaps below.

4. Data Gaps

To support the registration of Deet products, it is necessary to submit the following data, which are explained in detail in the "Proposed Guidelines for Registering Pesticides in the United States" (43 FR 29696, July 10, 1978).

a. Technical Deet

Hydrolysis
Activated Sludge

Guidelines Section

163.62-7 (a)
163.62-8 (g)

b. Formulated Deet

Hydrolysis

163.62.7 (g)

Human Exposure Data

Special Requirements
(FIFRA Sec 3(c) (2) (B))

1. Statistical data giving the percentage of total U.S. population exposed to Deet products, broken down by sex, age group (adults, teenagers, children, and pregnant women). Data on pregnant women are requested as a result of questions of possible embryotoxicity (See Chapter V);

2. Statistical data giving the geographical breakdown of the use frequency and amounts of Deet products (i.e., the Southeast, Gulf States and Pacific Northwest);

3. Data estimating the expected "average" or "normal" maximum exposure per day to an individual using Deet products. The data must encompass 90 to 99% of the total population exposed to such products. In addition, worst case and occupational exposure situations must be identified, including situations that could be reasonably expected to occur, such as application to large areas of the skin, frequent applications per day, etc. These estimates should be given for each of the population groups specified above;

4. Identification of the types of Deet products most commonly used and the percent active ingredient contained in each product.

5. Required Labeling

There are no environmental fate labeling requirements for technical or formulated Deet.

B. Topical Discussions

Corresponding to each of the Topical Discussions listed below is the number of the section in the "Proposed Guidelines for Registering Pesticides in the United States" (43 FR 29696, July 10, 1978) which explains the minimum data that the Agency requires to adequately assess the environmental fate of Deet. Special data requirements are also specified.

1. Technical Deet

<u>Data Requirement</u>	<u>Guidelines Section</u>
Physico-Chemical Transformation (Hydrolysis)	163.62-7 (a)
Metabolism (Activated Sludge)	163.62-8 (g)

a. Hydrolysis

Technical Deet could potentially enter natural water via industrial discharge, and as a result of disposal and cleanup of containers and equipment. Hydrolysis data are therefore required on technical Deet to support the registration of technical Deet. No data on the hydrolysis of Deet are available; this constitutes a data gap.

b. Activated Sludge Metabolism

For the assessment of Deet's potential effect on the wastewater treatment process through indirect discharge into treatment systems, an activated sludge metabolism study is required. No data on activated sludge metabolism of technical Deet are available; this constitutes a data gap.

2. Formulated Deet

<u>Data Requirement</u>	<u>Guidelines Section</u>
Physico-Chemical Transformation (Hydrolysis)	163.62-7 (a)
Human Exposure	Special requirement FIFRA Sec. 3(c) (2) (B)

Hydrolysis

Data are required for the same reasons as those listed under part (a) of this section. A data gap exists for formulated Deet.

V. TOXICOLOGY

A. Disciplinary Review

1. Toxicology Profile

a. Technical Deet

Sufficient data were available to assess the acute toxicity of technical Deet. The oral LD₅₀ in rats (male rats = 2.43 ml/kg; female rats = 1.78 ml/kg) indicates a potentially low acute oral toxicity in humans. The dermal LD₅₀ in rabbits (3.18 ml/kg or greater) indicates a potentially low dermal toxicity in humans. Likewise, the inhalation LC₅₀ in rats (5.95 mg/l) indicates a low acute inhalation toxicity in humans.

Some information was provided on the irritation and sensitization potential of technical Deet. In an eye irritation study conducted on rabbits, 0.1 ml of technical Deet induced marked transitory irritation and opacity. This study strongly suggests that technical Deet is potentially a severe eye irritant. Dermal irritation and sensitization studies conducted on rabbits and guinea pigs, respectively, demonstrated that technical Deet is not a potential skin irritant or sensitizer. No adequate subchronic toxicity data were available. In several inadequate subchronic (90-day) dermal studies, technical Deet was administered to rabbits. Conflicting results and inappropriate reporting preclude the use of these studies to assess the subchronic dermal toxicity potential; additional testing is required. However, an adequate subchronic inhalation study conducted on rats showed no abnormal behavioral signs or pathology at 7.5 mg/l; the lowest effect level was 15.0 mg/l, the effect being abnormal sperm morphology. Additional special testing was conducted on sperm count, spermhead, morphology and sperm viability in rats dermally administered doses of 0, 100, 300, or 1000 mg/kg/day of Deet 5 days each week for 9 weeks. There were no compound-related effects at any dose.

Inadequate chronic toxicity data were available to assess long-term effects of technical Deet. Supplementary studies¹ conducted on mice and rabbits demonstrated no oncogenic potential; however, inappropriate testing protocol precludes the use of these studies and additional data are required. In addition, conflicting teratogenicity studies were performed. An adequate dermal teratology test conducted using rabbits showed no effect at the highest dose of 5000 mg/kg/day. However, an oral teratology test conducted on rats resulted in a possible embryotoxic effect at 80 mg/kg/day; this test is not adequate to satisfy the data requirement due to inappropriate testing protocols and is therefore considered supplementary. An additional dermal teratology test is required in the same species to clarify questions raised by the first study. No adequate data were available to assess the reproduction toxicity potential of technical Deet. Mutagenicity testing is incomplete. The available mutagenicity data demonstrate that technical Deet does not induce reverse mutations in *S. typhimurium*. No dominant lethal effects at the dose tested (600 mg/kg in male mice) were indicated in a supplementary study (deemed supplementary because the maximum tolerated dose was not used). However, the

¹ / For the purposes of this Standard, a supplementary study is one that does not meet required testing standards but does provide some relevant information.

dominant lethal study showed a reduction of implants in pregnant females, suggesting that further reproduction testing is necessary. In addition, further mutagenicity testing is required to satisfy data requirements.

The available metabolism data were incomplete, but animal data suggest that Deet is absorbed (rabbit is 36%; male rat is 43%; female rat is 32%; humans is 8-10%), rapidly excreted (rats = 68% in 24 hrs) and not bioaccumulated. However, no metabolite identification data are available and additional metabolism testing is required.

b. Formulated Deet

(1) Ready-To-Use (RTU)

Based on available data (see Topical Discussions for details), statements of formula and the intended pattern of use for Deet, the acute toxicity (oral, dermal, inhalation), irritation (eye and dermal) and dermal sensitization potentials are as indicated in Table 5.

Table 5

Summary of Acute Toxicity Data on Ready-To-Use Formulations

Toxicity Testing	Existing RTU Products			
	15-25%	40-55%	75%	100%
Acute Oral	Low	Low 1/	Low 1/	Low
Acute Dermal	Low 1/	Low 1/	Low 1/	Low
Acute Inhalation	Very low 1/	Very Low 1/	Very Low 1/	Very Low
Primary Eye	Low	Data Gap 2/	Data Gap 2/	Severe eye irritation
Primary Dermal	Very Low	Data Gap 2/	Data Gap 2/	Very low
Primary Sensitization	Data gap 2/	Data Gap 2/	Data Gap 2/	No Sensitization

- 1/ Based on data extrapolation
 2/ Testing is required

(2) Pressurized Liquid (PrL)

Based on the available data (See Topical Discussions for details), statements of formula, and the intended pattern of Use for Deet, the acute toxicity (oral, dermal, inhalation), irritation (eye and dermal) and dermal sensitization potentials are indicated in the Table 6.

Table 6

Summary of Acute Toxicity Data on Pressurized Liquid Formulations

Toxicity Testing	Existing PrL Products 1/	
	12-30%	75%
Acute Oral	Low	Low ^{2/}
Acute Dermal	Low	Low ^{2/}
Acute Inhalation	Very Low ^{2/}	Very Low ^{2/}
Primary Eye	Low	Data Gap ^{3/}
Primary Skin	Very Low	Data Gap ^{3/}
Primary Sensitization	Data Gap ^{2/}	Data Gap ^{3/}

1/ Those existing PrL products which do not contain freon propellants.

2/ Based on data extrapolation

3/ Testing is required

(3) Impregnated Material (IMM)

Based on available data (see Topical Discussions for details), statements of formula, and the intended pattern of use for Deet, the acute toxicity (oral, dermal, inhalation), irritation (eye and dermal) and dermal sensitization potentials are as indicated in Table 7.

Table 7

Summary of Acute Toxicity Data on Impregnated Material Formulations

Toxicity Testing	Registered IMM Products	
	11-27%	33%
Acute Oral (LD ₅₀)	Low ^{1/}	Low ^{1/}
Acute Dermal (LD ₅₀)	Low ^{1/}	Low ^{1/}
Acute Inhalation (LC ₅₀)	Very Low ^{1/}	Very Low ^{1/}
Primary Eye	Data Gap ^{2/}	Data Gap ^{2/}
Primary Dermal	Data Gap ^{2/}	Data Gap ^{2/}
Primary Dermal Sensitization	Data Gap ^{2/}	Data Gap ^{2/}

1/ Based on data extrapolation

2/ Testing is required

2. Toxicology Hazard Assessment

a. Technical Deet

The information available to assess potential hazard as a result of chronic exposure is incomplete and inconclusive (see Topical Discussions and the Toxicology Profile for details). Therefore, the oncogenic, teratologic, mutagenic and reproduction effects resulting from chronic exposure to formulated Deet, based on testing of technical Deet, cannot be adequately evaluated. Several supplemental studies (dominant lethal mutagenicity, teratology, and subchronic inhalation toxicity) indicate possible reproductive toxicity. Although the data are only suggestive, it is imperative to conduct the necessary studies to resolve the questions of adverse effects.

b. Formulated Deet

(1) Ready-To-Use

Because of the intended pattern of use for RTU Deet products, the likely routes of exposure are dermal and occasional accidental oral, eye, and inhalation. The existing RTU products have a low acute oral and inhalation toxicity potential; therefore, single accidental exposures are not expected to pose a hazard. Accidental eye exposure is potentially serious because corneal opacity has been noted in a technical product; therefore, extreme caution should be used when applying RTU products which contain more than 30% Deet. The information currently available to assess the primary dermal toxicity and dermal sensitization of RTU formulations is incomplete. Likewise, the data for chronic exposure is incomplete and therefore a chronic toxicity risk cannot be determined at this time.

(2) Pressurized Liquid (PrL)

Due to the intended pattern of use for PrL Deet products, dermal, and occasional accidental oral, eye, and inhalation exposure are the likely routes of exposure. However, the existing products have a low acute oral and inhalation toxicity potential; therefore, accidental exposures are not expected to pose a hazard. Accidental eye exposure is potentially serious as corneal opacity has been noted in a technical Deet product; therefore, extreme caution should be used when applying RTU products. The information currently available to assess the primary dermal toxicity and dermal sensitization is incomplete. Likewise, the data for chronic exposure to technical Deet is incomplete and therefore a chronic toxicity risk cannot be determined at this time.

(3) Impregnated Material (ImM)

Due to the intended pattern of use for ImM Deet products, dermal and occasional accidental oral, eye, and inhalation are the likely exposure routes. However, the existing products have a low acute oral and inhalation toxicity potential; therefore, these single accidental exposures are not expected to pose a hazard. Accidental eye exposure is potentially serious because corneal opacity has been noted in a technical Deet product. The information currently available to assess the primary dermal toxicity and dermal sensitization is incomplete. Likewise, the data for chronic exposure is incomplete; therefore, a chronic toxicity risk cannot be determined at this time.

3. Generic Toxicology Data Gaps

Technical Deet

Guidelines Section:

Subchronic 90-day dermal (rabbits)	163.82-3
2-yr Dermal exposure combined with oncogenicity (rat)	163.83-2
Dermal Oncogenicity (mouse)	163.83-2
Dermal Teratogenicity (rat)	163.83-3
Dermal Reproduction (rat)	163.83-4
Mutagenicity	163.84-2
	163.84-3
	163.84-4
Metabolism (identification of metabolites)	163.85-1

Formulated Deet

Ready-To-Use (RTU)

Primary Eye Irritation (40-55% and 75%) (rabbit)	163.81-4
Primary Dermal Irritation (40-55% and 75%) (rabbits)	163.81-5
Dermal Sensitization (15-25%; 40-55%; 75%) (Guinea Pig)	163.81-6

Pressurized Liquid (PrL)

Primary Eye Irritation (75%) (rabbit)	163.81.4
Primary Dermal Irritation (75%) (rabbit)	163.81-5
Dermal Sensitization (12-30%; 75%) (Guinea Pig)	163.81-6

Impregnated Material (Imm)

Primary Eye Irritation (11.27% and 33%) (rabbit)	163.81.4
Primary Dermal Irritation (11.27% and 33%) (rabbit)	163.81-5
Dermal Sensitization (11.27% and 33%) (Guinea Pig)	163.81-6

4. Required Labeling

All technical and formulated Deet products must bear appropriate acute toxicity labeling and statements of practical treatment as specified in the Guidance Package which accompanies this Standard.

B. Topical Discussions

1. Acute Oral Toxicity (163.81-1)

The minimum data requirement for testing acute oral toxicity (LD_{50}) is one test for the technical formulation and one test for each registered product, preferably using the laboratory rat.

a. Technical

Adequate acute oral toxicity studies were conducted as indicated in Tables 8 and 9.

Table 8
Acute Oral Toxicity of Technical Deet

Animal	%-DEET	LD ₅₀	Toxicity Category	Reference
Rat (M)	90-100%	2.43 ml/kg	III	Ambrose, 1959 MRID #00001051
Rat (F)	90-100%	1.78 ml/kg	III	Ambrose, 1959 MRID #00001051
Rat (M)	Unspecified	2.68 ml/kg	III	Carpenter, et al., 1974 MRID #05000243
Rat (M)	95%	3.29G/kg	III	U.S. Army, 1979 MRID #GS0002030
Rat (F)	95%	2.42G/kg	III	U.S. Army, 1979 MRID #GS0002030
Rat (M)	95% (in corn oil)	3.16G/kg	III	U.S. Army, 1979 MRID #GS0002030
Rat (F)	95% (in corn oil)	2.17G/kg	III	U.S. Army, 1979 MRID #GS0002030

The above information is sufficient to satisfy the data requirement for acute oral toxicity for technical DEET. The data indicate that technical Deet should be assigned to Toxicity Category III for acute oral toxicity, which corresponds to a low potential acute oral hazard.

In additional testing using the ortho(o)-isomer and the para(p)-isomer, which occur as impurities in Deet, the acute oral toxicity in rats was as follows:

Table 9
Acute Oral Toxicity of o- and p- Isomers of Deet

Animal	%-Deet	LD ₅₀	Toxicity Category	Reference
Rat (M/F)	95% o-Deet	1.21 g/kg	III	Ambrose & Yost, 1965 MRID #00001102
Rat (M/F)	95% p-Deet	2.3 g/kg	III	Ambrose & Yost, 1965 MRID #00001102

The data indicate that o-Deet and p-Deet should be assigned to Toxicity Category III for acute oral toxicity, which corresponds to a potentially low acute oral hazard.

b. Formulated Deet

(1) Ready-To-Use (RTU)

The acute oral LD₅₀ in the rat is 2.10 ml/kg (Warf, 1976 MRID #00001101) for an 18% RTU product. The data indicate that the 18% RTU product should be assigned to Toxicity Category III for acute oral Toxicity.

Because oral toxicity studies place both technical deet and the 18% RTU product in Toxicity Category III, and because the inert ingredients in RTU Deet formulations are not expected to increase the acute oral toxicity potential, all existing RTU concentrations are placed in Toxicity Category III. Toxicity Category III corresponds to a low acute oral toxicity potential.

(2) Pressurized Liquid (PrL)

Adequate acute oral toxicity studies were conducted as reported Table 10.

Table 10
Acute Oral Toxicity of PrL Deet

Animal	%m-DEET	LD ₅₀	Toxicity Category	Reference
Rat (M)	15%	5-10 ml/kg	III	Warf, 1975a, 1975b
Rat (F)	15%	2-10 ml/kg	III	Warf, 1975a, 1975b MRID #00001085, #00001086
Rat (M)	30%	2.61 ml/kg	III	Howard, 1971
Rat (F)	30%	2.3 g/kg	III	Howard, 1971 MRID #00001080

The table shows that PrL products at 15% and 30% concentrations are in Toxicity Category III. Because technical Deet and the 15% and 30% PrL formulations are in Toxicity Category III, and because the inert ingredients in the PrL formulations are not expected to increase the acute oral toxicity potential, existing PrL products between 12% and 75% concentrations, except those that contain freon propellants, are classified in Toxicity Category III. Toxicity Category III corresponds to a low acute oral toxicity potential.

NOTE: Freon is regarded as a major hazard to the environment and its use in consumer products has been prohibited by the Agency. For all existing products which contain freon propellants the registrant must submit formulation statements with replacement propellants. The Agency will evaluate the new submissions on a case by case basis.

(3) Impregnated Materials (ImM)

No data were available for the assessment of the acute oral toxicity of ImM products. Because technical Deet is placed in Toxicity Category III) and because the inert ingredients in ImM products are not expected to increase the

acute oral toxicity potential, all existing IMM products between 11% and 33% concentrations are placed in Toxicity Category III. Toxicity Category III corresponds to a low acute oral toxicity potential.

2. Acute Dermal Toxicity (163.81-2)

The minimum data requirement for testing acute dermal toxicity (LD₅₀) is one test for the technical product and one test for each registered formulation, preferably using the albino rabbit.

a. Technical

Adequate acute dermal toxicity studies were conducted as indicated in Table 11.

Table 11

Animal	%-DEET	LD ₅₀	Toxicity Category	Reference
Rabbit	90-100%	greater than 4 ml/kg	III	Ambrose, 1959 MRID #00001051
Rabbit	unspecified	3.18 ml/kg	III	Carpenter, 1974 MRID #05000243
Rabbit	95%	4.28 g/kg	III	U.S. Army, 1979 MRID #GS0002026

The data indicate that technical Deet should be assigned to Toxicity Category III for acute toxicity, which indicates a low potential acute dermal hazard.

The above data are sufficient to satisfy the data requirement for acute dermal toxicity for technical Deet.

b. Formulated Deet

(1) Ready-To-Use (RTU)

The acute dermal LD₅₀ in the rabbit (abraded and unabraded skin) is greater than 4 g/kg for a 15% RTU product (Davidow, 1960 MRID #00001139). The data indicate that the 15% product should be assigned Toxicity Category III for acute dermal toxicity.

Because acute dermal toxicity studies place both technical deet and the 15% RTU product in Toxicity Category III, and because the inert ingredients in RTU Deet formulations are not expected to increase the acute dermal toxicity potential, all existing RTU products between 15% and 75% concentrations are placed in Toxicity Category III. Toxicity Category III corresponds to a low acute dermal toxicity potential.

(2) Pressurized Liquid (PrL)

No data were available to assess the acute dermal toxicity of the existing PrL product. Because technical Deet is placed in Toxicity Category III, and because the inert ingredients in PrL products are not expected to increase the acute dermal toxicity potential, all existing PrL products between 12% and 75%

concentrations, except those that incorporate from propellants, are classified in Toxicity Category III. Toxicity Category III corresponds to a low acute dermal toxicity potential.

(3) Impregnated Materials (ImM)

No data were available to assess the acute dermal toxicity of the existing ImM products. Because technical Deet is placed in Toxicity Category III, and because the inert ingredients in ImM products are not expected to increase the acute dermal toxicity potential, all existing ImM products between 11% and 33% are classified in Toxicity Category III. Toxicity Category III corresponds to a low acute dermal toxicity potential.

3. Acute Inhalation Toxicity (163.81-3)

The minimum data requirement for testing acute inhalation toxicity (LC_{50}) is one test on the technical chemical and on each manufacturing use and formulated product, preferably using the laboratory rat.

An acute inhalation toxicity (LC_{50}) test is required for each formulation that causes a respirable vapor, or for which 20% or more of the aerodynamic equivalent is composed of particles not larger than 10 microns.

a. Technical

A supplemental acute inhalation study was conducted on rats (Ambrose, 1959 MRID #0000105). The animals were exposed to an unspecified amount of technical Deet (85% A.I.) aerosol for 6 hours. Toxic signs observed were slight bloody discharge around the eyes and nose immediately after exposure; all rats appeared normal at 24 hours. All rats survived the exposure; histopathologic examination was negative. This study does not meet the data requirement for acute inhalation because the dose was not reported and therefore is considered supplementary.

An adequate acute inhalation study was conducted on male and female rats (U.S. Army, 1979 MRID #GS0002034). The animals were exposed to 0, 3.70, 4.26, 5.19, 5.48 or 5.95 mg/l of technical Deet for 4 hours. The LC_{50} was calculated to be 5.95 mg/l which corresponds to Toxicity Category IV, indicating a very low acute inhalation toxicity potential.

The above data are sufficient to satisfy the acute inhalation toxicity data requirement for technical DEET.

b. Formulated

(1) Ready-To-Use (RTU)

No data were available to assess the acute inhalation toxicity of registered RTU products. Because technical Deet is placed in Toxicity Category IV, and because the inert ingredients in RTU products are not expected to increase the acute inhalation toxicity potential, all existing RTU products between 15% and 75% concentrations are placed in Toxicity Category IV. Toxicity Category IV corresponds to a very low acute inhalation toxicity potential.

(2) Pressurized Liquids (PrL)

The available data were invalid to assess the acute inhalation toxicity of the 15% PrL product. Because technical Deet is placed in Toxicity Category IV, and because the inert ingredients in PrL products are not expected to increase the acute inhalation toxicity potential, all existing PrL products between 12% and 75% concentrations, except those that incorporate freon propellants, are placed in Toxicity Category IV. Toxicity Category IV, corresponding to a very low acute inhalation Toxicity potential.

(3) Impregnated Material (ImM)

No data were available to assess the acute inhalation potential of existing ImM products. Because technical Deet is placed in Toxicity Category IV, and because the inert ingredients in ImM products are not expected to increase the acute inhalation toxicity potential, all existing ImM products between 11% and 33% concentrations are placed in Toxicity Category IV. Toxicity Category IV corresponds to a very low acute inhalation toxicity potential.

4. Primary Eye Irritation (163.81-4)

The minimum data requirement for primary eye irritation is one test for the manufacturing-use product and one test for each registered product, preferably using the albino rabbit.

a. Technical/Manufacturing-Use Product

A primary eye irritation study was conducted on rabbits (Ambrose, 1959 MRID #00001051) using one drop (0.04 mg) of Deet. The Deet produced moderate edema and erythema and some "cloudiness" in 3 rabbits for 72 hours; after 5 days all eyes were normal. This study is not sufficient to meet data requirements for eye irritation because the scoring system used was not identified and the individual scores were not reported. It is therefore considered supplementary. However, this study does suggest that technical Deet is a potentially severe eye irritant in rabbits.

An adequate eye irritation study was conducted on rabbits (U.S. Army, 1979 MRID # GS0002025). A dose of 0.1 ml of technical Deet was instilled into the lower conjunction of the rabbit eye and evaluated by the method of Draize (1959). The following results were reported:

Table 12

Results of U.S. Army (1979) Eye Irritation Study

Structure	Mean Draize Score			
	24 hrs.	48 hrs.	72 hrs.	7 days
Cornea	0.5	0.2	0.0	0.0
Iris	0.2	0.0	0.0	0.0
Conjunctivae	2.7	1.0	0.4	0.0

This study indicates that technical Deet induced marked transitory eye irritation and transitory corneal opacification. This information indicates

that technical Deet is irritating to the rabbit eye due to the transitory corneal opacity noted. This information is sufficient to satisfy the data requirement for primary eye irritation for technical Deet.

b. Formulated Deet

(1) Ready-To-Use (RTU)

Two adequate primary eye irritation studies were conducted on existing RTU products as shown in Table 13.

Table 13

Primary Eye Irritation of RTU Deet

Animal	%m-DEET	Dose	Results	Reference
Rabbit ^{1/}	15%	100 mg	Mild irritation at 48 hrs but absent at 72 hrs; no corneal opacity.	Davidow 1960, MRID #00001139
Rabbit ^{1/}	15%	Oil	Mild irritation at 48 hrs. but absent at 72 hrs; no corneal opacity	Warf 1976, MRID #00001101

^{1/} Washed and Unwashed Eyes

The above data are sufficient to classify registered 15% and 18% RTU products as mild, transitory eye irritants. However, due to the reported results on technical Deet, each of the existing RTU products between 20% and 75% must be tested for eye irritation potential.

(2) Pressurized Liquid (PrL)

Adequate primary eye irritation studies were conducted on existing PrL products as shown in Table 14.

Table 14

Primary Eye Irritation of PrL Deet

Animal	%DEET	Dose	Results	Reference
Rabbit ^{1/}	15%	1 second spray	Mild irritation at 72 hrs but absent at 4 days, no corneal opacification	WARF 1975b MRID #00001085
Rabbit ^{1/}	15%	0.1 ml condensate	Mild irritation at 72 hrs, but absent at 4 days; no corneal opacity	WARF 1975b MRID #00001085 Woodward
Rabbit ^{1/}	30%	0.1 ml condensate	Mild irritation at 48 hrs; but absent at 72 hrs; no corneal opacity	Durloo and Woodward 1971b MRID #00001082
Rabbit ^{1/}	30%	2 second spray	No irritation at 48 hrs; no corneal opacity	Durloo and Woodward 1971b MRID # 00001082

1/ Washed and Unwashed Eyes

2/ Unwashed Eyes

The above data are sufficient to classify the tested 15% and 30% PrL products as mild, transitory eye irritants.

However, due to the reported results on technical Deet and the possible eye irritating inert ingredients in some products, each of the existing products must be tested for eye irritation potential.

(3) Impregnated Material (IMM)

No data were available for the assessment of the eye irritation potential of registered IMM products. However, due to the reported results on technical Deet all IMM products must be tested for eye irritation potential.

5. Primary Dermal Irritation (163.81-5)

The minimum data requirement for primary dermal irritation is one test for the manufacturing-use product and one test for each registered product, preferably using the albino rabbit.

a. Technical/Manufacturing-Use Product.

Adequate primary dermal irritation studies were conducted as indicated in 15.

Table 15

Primary Dermal Irritation of Technical Deet

Animal	%m-DEET	Irritation Signs	Toxicity Category	Reference
Rabbit	90-100%	48 hrs - slight edema 48 hrs - slight erythema 7 days-clear	IV	Philipps 1972, MRID # 05000242
Rabbit ^{1/}	95%	24 hrs - slight erythema 48 hrs - slight erythema 7 days - clear	IV	U.S. Army, 1979 MRID # GS00020269

1/ Abraded and unabraded skin

These studies show that technical Deet is mildly irritating to rabbit skin and corresponds to Toxicity Category IV, indicating a very low primary dermal irritation potential. This information is sufficient to satisfy the data requirement for primary skin irritation for technical Deet.

b. Formulated

(1) Ready-To-Use (RTU)

Adequate primary dermal irritation studies using rabbits were conducted on the registered RTU products shown in Table 16.

Table 16

Primary Dermal Irritation of RTU Deet

% m-DEET	Results	Toxicity Category	Reference
15%	No irritation at 72 hrs. using abraded and nonabraded skin.	IV	Davidow 1960 MRID # 00001139
18%	Mild irritation at 72 hrs. using abraded and nonabraded skin.	IV	WARF 1976 MRID # 00001101

As per the available data on the 15% and 18% RTU products (Toxicity Category IV) and technical Deet (Toxicity Category IV), products between 15% and 25% may be classified as Toxicity Category IV, corresponding to a very low primary

dermal irritation potential. However, due to the intended use pattern of Deet and the lack of available data, all products which contain greater than 25% Deet must be tested for primary dermal irritation potential.

(2) Pressurized Liquids (PrL)

The following adequate dermal irritation studies using rabbits were conducted on registered PrL products:

Table 17

Primary Dermal Irritation of PrL Products

% m-DEET	Results	Toxicity Category	Reference
15%-15.45%	No irritation at 72 hrs. using abraded and nonabraded skin.	IV	Johnson, 1972; MRID # 00001073 WARF, 1975a; MRID # 00001085 WARF, 1975b MRID # 00001086
30%	No irritation at 72 hrs. using abraded and nonabraded skin.	IV	Durloo and Woodward, 1971a MRID #00001081

The data show that 15% and 30% PrL products are placed in Toxicity Category IV. Because technical Deet and the 15% and 30% PrL products are placed in Toxicity Category IV, and because the inert ingredients in PrL products are not expected to increase the primary dermal irritation potential, all existing PrL products between 12% and 30% concentrations, that do not incorporate from propellants, are placed in Toxicity Category IV. Toxicity Category IV corresponds to a very low primary dermal irritation potential.

(3) Impregnated Material (ImM)

No data were available for the assessment of the primary dermal irritation potential of the registered ImM products. Due to the intended use pattern of Deet and the lack of available data, all ImM products must be tested for primary dermal irritation potential.

6. Dermal Sensitization (163.81-6)

The minimum data requirement for dermal sensitization is a test for the technical product and each registered formulation preferably using the guinea pig.

a. Technical/Manufacturing-Use

An adequate dermal sensitization study was conducted on guinea pigs (U.S. Army 1979 MRID GS0002026). The animals received 10 applications of 0.1% technical Deet followed by an insult application after a 2 week rest. No sensitization reaction was noted. These data indicate that technical Deet is not a skin sensitizer.

b. Formulated

(1) Ready-To-Use (RTU)

A supplementary dermal sensitization study was conducted on guinea pigs (Ambrose, 1959 MRID #0001051). The animals received 10 applications of 1 ml of 10% Deet in isopropanol on the depilated flank. Fifteen days following the last application, the animals received a single insult application. Slight dermal irritation was seen after the 3rd or 4th application; no other reactions were noted. This data suggests that 10% Deet in isopropanol is not a skin sensitizer.

Due to the intended use pattern of Deet and the limited available data, all RTU products must be tested for dermal sensitization.

(2) Pressurized Liquid (PrL)

No data were available for the assessment of the dermal sensitization potential of any PrL product. Due to the intended use pattern of Deet, each of the registered PrL products between 12% and 75% and all new products must be tested for dermal sensitization potential.

(3) Impregnated Materials (ImM)

No data were available for the assessment of the dermal sensitization potential of any impregnated materials product. Due to the use pattern of Deet, each of the existing ImM products and all new products must be tested for dermal sensitization potential.

7. Subchronic Oral Toxicity

The minimum data requirement for testing subchronic oral toxicity is testing in two mammalian species, preferably the dog and rat, using the technical product. Subchronic oral testing is not required for Deet because a significant exposure via the oral route is unlikely. However, the following data were available:

A supplementary 200 day feeding study was conducted on rats (Ambrose, 1959 MRID #00001051). Groups of rats (10 male, 10 female) were fed 0, 0.01, 0.05, 0.1, 0.5, or 1% m-Deet (equivalent to 4.5-10.0, 23.5-45.6, 48.0-92.2, 267.0-474.0, and 531.0-1055.0 mg/kg/day, respectively) in the diet. Male and female rats in the 1% group showed lower weight gains than controls. Average daily food consumption, red cell and plasma cholinesterase values, and hemoglobin counts did not differ from control values for any treatment group.

Control and treated animals had comparable blood cholinesterase values, gross pathology and histopathology. However, there was a statistically significant increase in the organ-to-body weight ratio in the testes, kidneys and liver as indicated in Table 18.

Table 18

Organ-to-Body Weight Ratios in Ambrose, 1959 Study

% Deet	Body weight (gm)		Organ to body weight ratios					
	Male	Female	Testes**		Kidneys		Liver	
			Male	Female	Male	Females	Males	Females
0	405	232	0.71+0.03	0.55+0.01	0.57+0.01	2.5+0.07	2.7+0.15	
0.01	408	242	0.76+0.03	0.57+0.02	0.53+0.06	2.7+0.10	2.5+0.25	
0.05	389	237	0.75+0.01	0.59+0.03	0.57+0.02	2.6+0.09	2.6+0.23	
0.10	386	229	0.81+0.02	0.57+0.02	0.57+0.01	2.6+0.10	2.3+0.06	
0.50	385	224	0.86+0.05*	0.99+0.03	0.59+0.02	3.0+0.11*	3.0+0.03	
1.00	366*	209*	0.87+0.02*	0.62+0.02*	0.64+0.03*	3.1+0.14*	3.6+0.12*	

*Significant at 5% level.

**The increase in body-weight ratio in the testes suggest a dose-response relationship.

Another supplementary subchronic oral study using technical Deet in dogs was reviewed (Woodward, 1959 MRID # 00001029). Groups of two male and two female beagles were given 85% m-Deet with 10% other isomers) orally by capsule at 0, 0.1, or 0.3 ml/kg/day for 13 weeks. After each daily dose, dogs at the 0.3 ml/kg/day dose level showed signs of slight-to-moderate central nervous system excitation consisting of tremors and hyperactivity. Emesis occurred "from time to time." Animals treated at 0.1 ml/kg/day showed only slight hyperactivity. Hemograms taken throughout the study failed to reveal any consistent changes. At necropsy, there was no evidence of gross pathology associated with the treatment. Organ weights were within normal limits for all animals. Histological studies of all tissues examined (liver, kidney, heart, spleen, ovary, uterus, testis, and adrenal) were within normal limits.

8. Subchronic 21-Day Dermal Toxicity (163.82-2)

The minimum data requirement for subchronic 21-day dermal toxicity is one test for the technical chemical, preferably using the albino rabbit. However, a subchronic 21-day dermal test is not required for Deet because the intended use of Deet is purposeful application to the skin and the need for a 90-day dermal toxicity test precludes the requirement for a 21-day dermal toxicity test.

9. Subchronic 90-Day Dermal Toxicity (163.82-3)

The minimum data requirement for subchronic 90-day dermal toxicity is one test for the technical chemical, preferably using the albino rabbit. The subchronic 90-day dermal toxicity is required because pesticidal use of Deet involves purposeful application to the skin.

Two supplementary subchronic dermal toxicity tests were conducted on rabbits (Ambrose, 1959 MRID #00001051 and Woodard, 1959 MRID # 00001029). These were as follows:

1. The application of Deet (85-100% m-Deet) to intact rabbit skin for 90 days at a dose of 1 ml/kg/day produced moderate skin irritation but no effect on weight, behavior, or mortality (Ambrose, 1959 MRID #00001051). Although these data suggest no systemic effects of technical Deet, the lack of

pathologic examination and reporting of clinical parameters precludes the use of this study.

2. In the other study, rabbits (six per dose) dermally dosed the 0, 0.75, 1.50 and 3.0 ml/kg/day of Deet (85% m-Deet, 10% Deet isomers) for 90 days showed mild dermal irritation (0.75 ml/kg/day) to severe dermal irritation (3.0 ml/kg/day) (Woodard, 1959 MRID # 00001029). In addition, increased kidney weights and marked histopathological changes were noted in the intermediate dose animals (1.5 ml/kg/day). Two rabbits dosed at 1.5 ml/kg/day died, and all the rabbits at the high dose (3.0 ml/kg/day) dies. However, the lack of clinical data (e.g. growth curves, blood chemistries, etc.) and the lack of data on individual animal preclude the use of this study.

The available information is not sufficient to satisfy the requirement on subchronic 90-day dermal toxicity; additional testing is required.

10. Subchronic Inhalation Toxicity (163.82-4)

The minimum data requirement for subchronic inhalation is one test for the technical formulation, preferably using the laboratory rat. The registered end uses of Deet could result in repeated inhalation exposure.

A supplementary subchronic inhalation study was conducted on rats (Ambrose, 1959 MRID #00001051). Rats exposed to air "saturated" with Deet in an inhalation chamber for 8 hours a day, 5 days a week, for 7 weeks showed no effect on behavior, mortality, or histopathology. Although the data suggest that no adverse effects resulted from subchronic inhalation exposure, this study is not sufficient to meet data requirements for subchronic inhalation because actual doses were not reported, and incomplete pathology was reported.

An adequate subchronic inhalation study was conducted on rats (male and female) and Beagle dogs (male and female) (U.S. Army, 1979 MRID #GS 0002034). The animals were exposed to aerosol concentrations of 0 (room air), 2.5, 7.5, or 15.0 mg/l of Deet for 6 hours per day, 5 days per week for 13 weeks. The lowest effect level (LEL) for rats is 15.0 mg/l (transient red exudate around the eyes and nose were noted); the no effect level is 7.5 mg/l. In addition, an increase in sperm with abnormal morphology was noted in rats at the 15.0 mg/l dose. However, this effect is evidence that mutagenicity and reproductive testing are necessary.

11. Chronic Feeding/Dermal Toxicity (163.83-1)

The minimum data requirement for chronic feeding/dermal toxicity is one test for the technical chemical, preferably using the laboratory rat. A chronic dermal study is required because pesticidal use of Deet could result in application to the skin for significant periods of time. The chronic dermal toxicity study would be conducted as a part of a chronic feeding study.

12. Oncogenicity (163.83-2)

The minimum data requirement for oncogenicity is testing in two mammalian species, preferably the rat and mouse, using the technical chemical.

A screening bioassay was conducted on rabbits and mice (Stenbaeck, 1977 MRID# 05000045). Groups of 50 females mice or 5 rabbits (sex distribution unspecified) received dermal applications of 10%, 50% or 100% Deet in ethanol.

The positive control used was DMBA. This study resulted in no statistically significant results, and is also not adequate to satisfy testing requirements for the following reasons:

The mouse experiment was conducted on only female mice with limited histological examinations performed (only grossly observed lesions were examined microscopically and the extent of the examination was not described); no data were provided on individual animals; no clinical data were reported; and there was; no indication if the maximum tolerated dose was administered.

The Rabbit study was conducted using an Insufficient number of animals; no data were provided on individuals animals; no pathology report was provided for tumor types; no clinical data were reported; there was no; no indication if the maximum tolerated dose was administered; and there were early and abrupt deaths (40% at the 10% and 100% dose levels at 80 weeks; 100% at 10%, 50% and 100% dose levels at 90 weeks), causing concern about the conduct of the study.

Additional oncogenic testing is required in two mammalian species.

13. Teratology (163.83-3)

The minimum data requirement for teratology is testing in two mammalian species using the technical chemical.

An adequate dermal teratology study (U.S. Army, 1980 MRID #GS0002036) was conducted on rabbits. Groups of 20 pregnant rabbits received daily dermal applications of 0, 50, 100, 500, 1000, or 5000 mg Deet/kg/day in ethanol (vehicle control) on shaved backs from day 0 through day 29 of gestation. Precautions were taken to prevent the animals from licking treated skin and skin was not washed between treatments. There were no significant differences between control and treated animals with respect to the fertility index, number of implantations per animal, or number of fetuses per animal. In addition, treatment did not change fetal weight, fetal length or placental weights and no increases in the incidence of skeletal or soft tissue anomalies were observed in treated groups when compared with untreated controls. This study demonstrated that Deet has no teratogenic or embryotoxic effects in rabbits exposed dermally to technical Deet.

An additional supplementary teratology study (Grahwitt, 1977 MRID # 00001063) was conducted on rats. Groups of 20 pregnant rats were daily administered 10 ml of peanut oil containing 0, 8, 20 or 80 mg/kg/Deet by gavage from day 5 through day 15 of gestation. No significant differences were reported between control and treated mothers with respect to fertility, fetuses per litter, fetal weight or fetal survival. However, the study did show decreases in number of implantation sites per dam and number of fetuses per animal. In addition, a related increase was observed in the number of resorptions per dam (see summary table below).

Table 19
Summary of Results
in
U.S Army (1980) Teratology Study

Parameter	Control	8mg/kg	20mg/kg	80mg/kg
Implantations/dam	10.65	10.00	10.21	9.10
Fetuses/dam	10.45	9.25	9.37	7.70
% Live fetuses	100	100	99.44	100
Fetuses/litter	10.45	9.74	9.89	9.10
Resorption/dam	0.20	0.75	0.84	1.43

These summary data are somewhat misleading because two animals in the high dose group had low body weights and had 8 resorption sites each. A third animal, which had 5 resorption sites, delivered viable fetuses and its body weight was comparable to control animals. These 3 animals represent 75% of the total number of resorptions for this group. In addition, the number of implantations per animal in the high dose group was lower than that in control animal (9.10 and 10.65, respectively). These results are not compound related because Deet dosing began 3 days after implantation normally occurs in rats.

These data suggest the possibility of impaired maternal health prior to the administration of Deet; therefore, the significance of the suggested embryotoxicity at 80 mg/kg/day cannot be determined. However, the suggested embryotoxic effect is evidence that additional teratology testing is required in the same species.

Another teratogenicity study was reviewed and deemed insufficient to assess potential teratogenicity (Gleiberman, 1975 MRID #05000007). There are insufficient details reported to assess the protocol and no characterization of the chemical administered was reported in the study.

14. Reproduction (163.83-4)

The minimum data requirement for measuring reproductive indices is one test in the laboratory rat lasting two generations using the technical formulation.

An inadequate study was provided for assessment of the reproductive toxicity of Deet (Gleiberman, S., 1976 MRID # 05000008). There are insufficient details reported to assess the validity of the study and no characterization of the chemical administered was reported.

Although this study is not suitable to meet data requirements for reproductive testing it does suggest potential sperm abnormality. The authors report that by "the special function-morphological examination of spermatozoa, both motility time and the number of pathological forms (deformation, absence of tail or head, thickening of middle piece, etc.) differed reliably from similar indices in control animals." Without additional specific information the Agency cannot interpret the significance or the validity of this finding.

Additional reproductive testing is required.

15. Mutagenicity (163.84-1 through-4)

Although the Agency's mutagenic testing requirements are not final, refer to the "Proposed Guidelines" (43 FR No. 163, August 22, 1978) for information concerning the types of studies the Agency is considering.

The following studies are representative and are likely to be required:

- 1) Microbial point mutation
- 2) Mammalian in vitro point mutation
- 3) In vitro or in vivo cytogenetics or one of the following: heritable translocation or dominant lethal
- 4) Primary DNA damage, e.g. sister chromatid exchange or inscheduled DNA synthesis.

Mutagenic studies are required because pesticidal use of Deet involves purposeful application to the skin.

Several supplementary mutagenicity studies were reviewed for this Standard. In one study (Litton Bionetics, Inc. 1977 MRID#GS0002036), Deet did not induce reverse mutations in S-9 activated or inactivated *S. typhimurium* TA-1535, 1537, 1538, 1598 and 15100, and did not induce gene conversions in unactivated *S. cerevisia* D4. No conclusion could be drawn regarding the S-9 activation system in *S. cerevisia* D4 in the same study due to uncertainties in the preincubation suspension technique.

A supplementary dominant lethal study was conducted using three groups of ten male ICR/Ha Swiss mice (U.S. Army, 1979, GS0002021). The concurrent control group was administered a single dose by gavage of 5 mg/kg corn oil, the positive control group was administered 10 mg/kg TEM, and the experimental group was administered 600 mg/kg Deet. The study resulted in no dominant lethal effects, however, a statistically significant ($p < 0.02$) reduction of a plants was noted in pregnant females. This effect is evidence that a reproductive study is necessary. Although this study was generally conducted adequately, the maximum tolerated dose was not administered; therefore, the study is not sufficient to meet the data requirement for mutagenicity.

16. Metabolism in Laboratory Animals (163.85-1)

The minimum data requirement for metabolism is a single dose using the analytically pure grade of the active ingredient in the radioactively labelled form. A metabolism study is required using technical Deet because oncogenicity and chronic studies are required.

Metabolism studies showing excretion levels and absorption rates are summarized in Table 20. Tissue distribution studies are summarized in Table 21 and several studies conducted during pregnancy are summarized in Table 22.

The studies on animals suggest that m-Deet is rapidly absorbed and excreted in the urine. Distribution studies show the liver, kidney, bladder and lacrimal glands are the preferred organs, however excretion rates are so high (98%) that there is no evidence of bio accumulation. In addition, there is suggestive evidence that m-Deet does cross the placenta in mice (Bloomquist, 1975 MRID # 05000001) and rabbits (U.S. Army, 1979 GS0002023); however, there is no accumulation of Deet in any fetal organ indicating rapid fetal excretion.

Additional data is needed to identify metabolites of Deet and their potential bioaccumulation.

Table 20
Metabolism in Laboratory Animals

Animal	Dose	Route	% Absorbed	Excretion (Where ex- creted/Time)	Reference
Guinea Pigs	50 ₄ mg/kg C ¹⁴ Deet	Dermal	20-50%	80% urine/24hrs. 1% feces/8 days	Schmidt, (1959) MRID
Guinea Pigs	51 ₄ mg/kg C ¹⁴ Deet	Dermal	NS	70-80% urine/24 1% feces/un- specified	Kingscote, (1959) MRID #05000271
Mice	15 ₄ mg/kg C ¹⁴ Deet	Dermal	NS	13% urine/2, 5hrs. 34% urine/48hrs.	Bloomquist (1977) MRID #05000002
Mice	50 ₄ µCi /10gm C ¹⁴ Deet	I.V.	NS	95.6% urine/8hrs. 97.3% urine/40 hrs. 0.6% feces/48hrs.	Bloomquist (1975) MRID 05000001
Rats	100 ₄ mg/kg C ¹⁴ Deet	Dermal	NS	42% urine/6hrs. 68% urine/24hrs.	Lure et al. (1978) MRID #05000020
Rat/Male	NS	Dermal	43%	NS	U.S. Army (1969) MRID #GS0002034
Rat/Female	NS	Dermal	32%	NS	U.S. Army (1979) MRID #GS0002034
Rabbit	NS	Dermal	36%	NS	U.S. Army (1979) MRID #GS0002024
Dog	NS	Dermal	30%	NS	U.S. Army (1979) MRID #GS0002034

NS = Not specified

Table 21

Tissue Distribution Studies

Animal	Dose	Route	Tissue Distribution/Time After Dosing	Reference
Mice	15mg/kg	Dermal	1) Lacrimal Gland, liver, Kidney nasal mucosa/2 hrs. 2) Skin at application area, traces in Bladder/5 days 3) Skin at application area/36 days.	Bloomquist, (1977) MRID #05000002
Mice	5u Ci/ 10 gm	I.V.	1) Lacrimal gland, liver, kidney, small intestine (high level); Adipose, central nervous system (low level)/5 min. 2) Lacrimal gland, renal marrow, nasal mucosa, bladder, small intestine/1 hr. 3) Lacrimal gland, liver, nasal mucosa, bladder, intestine (low level)/24 hrs. 4) No trace of radioactivity at 4 days.	Bloomquist, (1975) MRID #05000001

Table 22

Tissue Distribution Studies
in Pregnant Animals

Animal	Gestation Time	Dose	Route	Time/Result	Reference
Mice	"Advances Dermal Pregnancy"	5uCi/ 10G	I-V	(1) 20 minutes/fetal bone marrow, urinary bladder, gastric mucosa lens showed "some accumulation"; average low concentration in the fetus was "very low." (2) 4 hours/no accumulation in any fetal organ.	Bloomquist (1977) MRID #05000002 Bloomquist, (1975) MRID #05000001
Rabbit	Day-1 through Day-29	?	Dermal	No detectable radioactivity above background was observed in any fetus.	U.S. Army (1979) MRID # GS0002034
Rabbit	Day-15	?	Dermal	Radioactivity was "lower" in fetuses than comparable maternal specimens	U.S. Army (1979) MRID #GS0002034

17. Clinical Trials —Metabolism and Pharmacodynamics

Although comprehensive metabolic and pharmacodynamic studies of the absorption, fate and excretion of Deet and its metabolites in human beings are not available, a few studies have measured rates of evaporation, dermal absorption, and urinary excretion.

Spencer et al. (1979, MRID #05007487) investigated the rate of evaporation of Deet after dermal application. The average total loss from human skin by evaporation, wiping, and stripping of the stratum corneum was 49% of the applied dose. This also suggests that some of the applied Deet that was not recovered penetrated the skin.

Feldman and Maibach (1970, MRID # 05003588) investigated the dermal absorption and excretion of radiolabeled Deet in four human volunteers. A total of 52 ug of Deet was applied once to the ventral forearm. The total average urinary excretion at 24 hours was 13.3% of the dose, at 48 hours it was 15.3%, and at 120 hours it was 16.7%.

Blomquist and Thorsell (1977, MRID # 05000002) applied 250 uCi of ¹⁴C-Deet in 0.03 ml (0.12 mg/kg of body weight) of a 20% ethanol solution to the forearm of one woman volunteer. The experiment was performed twice, but the time interval between the two experiments was not specified. The amount of radioactivity recovered in the urine during 48 hours was 5.5% of the first dose applied and 3.8% of the second. Washing of the treated skin area 8 hours after each application resulted in the recovery of 8% of the first dose and 15% of the second.

Markina and Yatsenko (1971, MRID # 05000599) investigated the dermal absorption of different formulations of Deet (20-40% active ingredient) in 15 persons and found that only trace amounts of Deet were detected in the urine when silicone or cellulose acetobutyrate additives were added to the Deet cream or lotion. No toxicity was reported after 30 days of daily application of the lotion. Medical examination revealed no effects.

These studies suggest that the dermal absorption of Deet is rapid but that excretion rates are variable. The number of subjects in these studies, however, is too small to allow more than a tentative conclusion regarding excretion rates. Pharmacodynamic studies also suggest that a considerable amount (15% or more) of the applied dose remains on the skin for at least 8 hours after application.

18. Dermal Irritation and Sensitization in Humans

Technical Deet (purity unspecified) in ethanol was not irritating to human skin when applied as a single dose (Phillips et al., 1972 MRID # 05000242). When applied to occluded human skin daily for 21 consecutive days as 1, 10, 20, 30, 60, or 80% ethanol solutions, mean scores over the 21-day period on a scale of 0-4 were 0.43, 0.13-1.13, 2.9, 2.4, 2.3, and 1.0, respectively. No irritation was observed when 100% technical grade Deet or a 50% ethanol solution was applied daily for 21 consecutive days to unoccluded human skin.

Ambrose (1959 MRID #00001051) applied a 50% solution of Deet to the arms and faces of five human volunteers once a day for 5 days. No irritation of the

arms was observed. However, after the third application some desquamation around the nose was apparent in all subjects. No further symptoms were noted after application of the solution was discontinued.

Lamberg and Mulrennan (1969 MRID # 05002308) applied patches moistened with 75% Deet (as a liquid in 25% ethanol or as a spray in 25% dichlorodifluoromethane) to the antecubital fossae of 77 men. This resulted in burns or in skin reactions of either bullar or erosions resembling abrasions in 48% of the men. These reactions led to necrosis and scarring in some of the subjects. However, when patches were applied to the upper inner part of the arm of 62 of these men, there were no skin reactions.

Vos (1972 MRID #00001093) patch-tested 52 adults with 15% Deet liquid three times a week for 9 weeks. After a 2-week rest period, a challenge patch was applied. Because only one of the subject developed a slight erythema after removal of the challenge patch, it was concluded that Deet is not a skin sensitizer.

Blau and Kanof (1961 MRID #00001182) applied blotting paper impregnated with a spray containing 7.5% Deet and 1.5% isobutyl salicyl cinnamate (Revlon Sun Bath) to the backs of 402 women. After 48 hours, the patches were removed and the skin was observed for signs of irritation. Of the 402 subjects, 9 showed a "minimal plus-minus" reaction; the rest were negative.

These studies show that Deet is not usually a dermal irritant or a skin sensitizer. However, in selected individuals and under special circumstances Deet can cause extensive skin reactions.

19. Case Reports

Maibach and Johnson (1975 05000021) reported that a 35-year-old woman developed red raised lesions after application of an insect repellent containing an unspecified amount of Deet. She was given open patch tests for sensitivity to several insecticides but only pure samples of Deet produced a wheal and flare reaction, which is considered contact urticaria or immediate-type hypersensitivity. Further investigation revealed that this response could be passively transferred, indicating that the contact urticaria to Deet was caused by an immunologic response.

Gryboski et al. (1961 MRID #05000328) described the case of a 3.5-year-old girl who, after exposure over a 2-week period to 180 ml of 15% Deet spray, experienced a toxic encephalopathy including disorientation, staggering gait, slurred speech, and episodes of crying out, stiffening into a sitting position, extending the extremities, flexing the fingers, and dorsiflexing the toes. She recovered after 4 days, and since that time she has reportedly been well. All clinical examinations to discover other etiologic factors, including family history, virology, bacteriology, and lead testing, proved to be negative; however, tests to confirm that Deet was the causative agent in this episode were not performed.

Zadjoff (1979 MRID #05007489) reported two additional cases of toxic encephalopathy in children after exposure to Deet. A 5-year-old girl sprayed nightly for 3 months with Mylol (containing 10% Deet) experienced a toxic encephalopathy, including slurred speech ataxia, and generalized convulsions. She died 24 days after hospitalization. The brain showed generalized edema with congestion of the meninges, and the cerebral blood vessels showed swelling

of endothelial cells. In another case, an 18-month-old girl ingested an unknown amount of Mylert liquid containing 10% deet. The next day she became irritable, displayed bizarre movements, and had depressed muscle stretch reflexes. She improved somewhat during 6 weeks of hospitalization. After 6 weeks, her head control was not yet normal, and her tendon reflexes remained depressed, but the case was not followed up.

Other reports of toxicity related to Deet exposure are discussed in Rabinovich (1966 MRID #05000693) and the Pesticide Incident Monitoring System Report Number 121 (PIMS 1978).

Rabinovich (1966 MRID #05000693) conducted a field trial on 600 lumbermen who used a 60% Deet solution and reported occasional contact dermatitis, aggravation of preexisting acne, conjunctivitis, and burning eyes.

Pesticide Incident Monitoring System Report Number 121 (PIMS 1978) lists 45 cases of Deet exposure found in the files of the Pesticide Incident Monitoring System for the period covering 1966 to December 1978. For 29 of these cases, data on the incident and sequelae were insufficient to draw any conclusions relating to the toxicity of Deet. The remaining cases included 10 incidents of temporary eye irritation; 6 cases of skin reactions involving blisters, rashes, irritation, and hives; and 1 case of a generalized allergic response requiring hospitalization.

The cited literature indicates that the general toxicity of Deet is low; that local skin reactions are infrequent, but can be severe depending on the area of application and on individual response; and the eye irritation, although slight and temporary, occurs frequently. There is inadequate information to judge potential effects in persons with preexisting allergic conditions or to determine if other populations, for example, children and pregnant women are at risk.

20. Emergency Treatment

No information has been submitted on emergency treatment of Deet intoxication.

21. Special Testing—Sperm Count, Spermhead Morphology, Sperm Viability

A valid morphology, viability and sperm count was conducted in rats (Litton Bionetics 1980 MRID #GS0002036). Groups of 80 male sprague-Dawley rats were dermally administered doses 0, 100, 300 or 1000 mg/kg/day of Deet for 5 days each week for 9 weeks.

There were no significant differences between treated and control rats with respect to mean body weights and food consumption (no statistical tests were considered necessary by the authors). Histological examination of the testes of rats in the high dose and control groups revealed no lesions nor the occurrence of compound related hypospermatogenesis. The appearance of liver and kidneys in treated rats was normal at necropsy. The only organ weights affected were those of liver and kidneys in the high dose rats at the first kill (no effect on testes weight was as shown in Table 23).

Table 23

Organ Weight Results in Litton (1980) Sperm Study

Dose (mg/kg)	Mean liver wt. (g)	% Body wt.	Mean Kidney wt. (g)	% body wt.
0	19.2550	4.4521	3.5100	0.8154
1000	19.8400	4.7427	4.2050	1.0062*

* significantly different from controls
(P is less than 0.05) using the
students t-test; no data from low
and misdosed groups reported.

There were no compound-related effects on sperm count, sperm head morphology, or sperm viability of treated male rats at repeated dermal doses up to 1000 mg/kg/day.

22. Age-Related Dermal Absorption Studies

The Agency is concerned with the possible age related differences in dermal absorption of Deet because Deet is used on children. Therefore, a dermal absorption study is required in weanling and young adult rats. Rats are chosen as the experimental animal because most of the toxicology and metabolism data will have been performed in rats, thus permitting correlation of data.

Table 24
Toxicity Category Indicators

Hazard Indicators	Toxicity Categories			
	I	II	III	IV
Oral LD ₅₀	Up to and including 50 mg/kg	From 50-500 mg/kg	From 500-5000 mg/kg	Greater than 5000 mg/kg
Inhalation LD ₅₀	Up to and including 0.2 mg/liter	From 0.2-2 mg/liter	From 2-20 mg/liter	Greater than 20 mg/liter
Dermal LD ₅₀	Up to and including 200 mg/kg	From 200-2000 mg/kg	From 2,000-20,000 mg/kg	Greater than 20,000 mg/kg
Skin Effects	Corrosive	Severe irritation at 72 hours	Moderate irritation at 72 hours	Mild or slight irritation at 72 hours or no effects

Table 24 A
Eye Irritation Toxicity
Indicators for Formulated Deet Products

Acceptable for Domestic Use	Not Acceptable for Domestic Use
No corneal opacity; irritation reversible within 7 days	Corrosive; any corneal opacity; irritation persisting for 7 days

VI. ECOLOGICAL EFFECTS

A. Disciplinary Review

1. Ecological Effects Profile and Hazard Assessment

Subpart E, "Hazard Evaluation: Wildlife and Aquatic Organisms," of the "Proposed Guidelines, Registration of Pesticides in the United States", published in the Federal Register on July 10, 1978, describes the fish and wildlife data required by the Agency to assess the hazards of pesticides to nontarget organisms and to provide for adequate precautionary labeling (43 FR 132, Part 163.70 through .72).

Formulated Deet is neither intended for application to outdoor sites, nor registered for such use; consequently, formulated Deet will not be reviewed in this Chapter. However, an understanding of technical Deet's toxicity to fish, aquatic invertebrates and waterfowl is fundamental to the development of appropriate clean-up measures in the event of a chemical spill into lakes and streams. The Agency's concern in this regard is supported by data indicating that a high volume of technical Deet is manufactured in the U.S.; manufacturers are located in diverse areas of the country; and there are a large number of formulators (over 100). The ecological effects review in this Chapter relates only to fish, aquatic invertebrates and waterfowl, and not to upland game birds or other non-aquatic wildlife, because the latter are not expected to experience effects of aquatic spills.

In general, the available data were insufficient to support an assessment of technical Deet's ecological effects. The only valid study, (McCann, 1972, MRID # 00001026) provided data which demonstrate that technical Deet has a slight acute toxicity to coldwater fish.

2. Data Gaps

The following gaps in the Ecological Effects data base must be filled to adequately support the continued registration of technical Deet. After each requirement is listed the section in the Proposed Guidelines of July 10, 1978 (43 FR 132, Part 163.70-.72) which describes the test in detail. These data requirements are also listed in Table 1, Chapter 2.

1. Avian single-dose LD₅₀ (wild waterfowl, preferably mallard duck) 163.71-1
2. Fish 96-hour LC₅₀ for warmwater species 163.72-2
3. Aquatic invertebrate acute LC₅₀ 163.72-2

3. Required Labeling

There are no ecological effects labeling requirements for technical or formulated Deet.

B. Topical Discussions

The data requirements listed below correspond to the sections in the "Proposed Guidelines for Registration of Pesticides in the United States", published in the Federal Register of July 10, 1978, which explain the data the Agency requires to adequately assess the hazards of technical Deet to fish and wildlife.

Data Requirements

Guidelines Section

Birds (wild waterfowl only)
Fish (cold and warmwater species)
Aquatic Invertebrates

163.71-1
163.72-1
163.72-2

Birds (wild waterfowl)

No valid studies on the effects of single oral doses of Deet to wild waterfowl were submitted.

Because data were not available, no conclusion can be drawn about the toxicity of Deet to wild waterfowl; this constitutes a data gap.

Fish

One valid study concerning the acute toxicity of technical Deet (95% a.i.) to fish (McCann, 1972, MRID #00001026) provided data on coldwater species. McCann reported a 24-hour LC-50 of 125 ppm and a 96-hour LC-50 of 172 ppm for rainbow trout. These results are sufficient to characterize Deet as slightly toxic to coldwater fish.

No data concerning the acute toxicity of technical Deet to warmwater fish were submitted; this constitutes a data gap.

Aquatic Invertebrates

No studies concerning the acute toxicity of Deet to aquatic invertebrates were submitted; this constitutes a data gap.

VII. EFFICACY

A. Introduction

This chapter deals with the efficacy of Deet when utilized as the single active ingredient in an insect repellent product. The review of Deet data and the identification of data gaps is limited to efficacy data only as it relates to public health applications. The Agency has provided for the waiver of efficacy data submission as a part of the registration process in all other instances (44 FR 2793; May 11, 1979).

The pest species found on registered Deet labels that fall within the Agency's area of public health concern are as follows:

- Bitting flies (black fly, sand fly, horse fly and Ceratopogonid species)
- Chiggers
- Deerflies
- Fleas
- Leeches
- Mosquitoes
- Stableflies
- Ticks

The Agency reviewed all pertinent studies. Many papers, while providing valid and useful data, were not considered because the data related to formulations containing more than one active ingredient. The Agency did not reject any data on the basis of test methodology, but considered that all data would contribute to the assessment of efficacy (See Appendix I.).

B. Efficacy Assessment

1. Factors Influencing Efficacy

Several factors appear to influence the efficacy of Deet. Major factors are environmental conditions, extent of absorption and elimination, type of repellent formulation, and avidity of the test species. One or more of these conditions always influences the duration of Deet's protection.

Environmental conditions, such as temperature and wind velocity, alter Deet activity. Deet efficacy is reduced by increased perspiration rate, although not by elevated temperature.

Although data generated by Smith (Smith et al., 1963, MRID# 05000300) were inconsistent, they suggest that perspiring might reduce Deet protection time against the mosquito, Aedes aegypti. Wind velocity also influences Deet protection time. Khan (Khan et al. 1973 MRID 05000193) demonstrated that wind velocities of 192 meters per minute decreased efficacy on the order of 66% when compared to data collected in a room with "normal" air exchange rates.

The duration of repellent activity is influenced by washoff, abrasion from the treated surface, and absorption. Water, from either rainfall or perspiration, decreases the efficacy of Deet. Gouch (Gouch et al., 1971, MRID# 05000214) reported that netting treated with 0.25 gram active ingredient of Deet per gram of netting was effective in repelling 90% of a population of Aedes taeniorhynchus mosquitoes for a period of 54 days. Approximately 0.3 inches of precipitation, however, reduced the effectiveness to less than 90% when

measured over the same period. Similarly, Schiefer (Schiefer et al., 1976, MRID 05000223) noted a reduction in the overall efficacy in response to rainfall. Deet's effectiveness has been reported to be reduced by loss through cutaneous absorption (Maibach et al. 1974, MRID 0500116). Since Deet is highly lipophilic, it is reasonable to assume that loss by absorption would be accelerated when lotions, creams, and stick-type formulations are applied. Much data has been reviewed about the formulation dependency of Deet activity; the Agency has considered only the effect of the range of percent active ingredient in its target pest assessment.

Given the absence of data relating to the upper limit of Deet efficacy based on percent active ingredient, the Agency will not prescribe the maximum level of Deet allowable in any formulation. However, the Agency will consider toxicology hazards of all formulations, balancing the risks against the benefits of any formulation for which a suspected hazard exists.

Although overall efficacy of Deet can also be seriously reduced by inadvertent abrasion from the skin, there are data that suggest the Deet can protect surrounding untreated skin up to 4-8 cm. from the treated area (Kahn and Maibach 1972, MRID 00001163).

2. Use Sites

a. Human Skin

The data clearly indicate that Deet products which contain greater than 10% active ingredient are efficacious for use on human skin. Any performance claims made must fall within the general parameters prescribed in each of the pest-specific discussions in Part 3 of this section.

b. Clothing

To repel fleas and chiggers, the product must be applied to clothing in order to work satisfactorily. In the case of other pests, (e.g. black flies, mosquitoes, etc.), clothing treatment may be advisable. The Agency, therefore, recognizes clothing treatment efficacy. The Agency also recognizes the potential utility of specialty formulations for this use. The Agency will consider the specialty products supported under this standard. Any performance claims made must fall within the general parameters prescribed in each of the pest-specific discussions under Target Pests below.

c. Tents and Bedrolls

A limited number of Deet products bear claims for treatment of tents and bedrolls. The Agency has very little data directly related to this use (e.g., netting and general cloth treatment). However, through extrapolation from these data, the Agency considers applications for use on tents and bedrolls to be supported under this standard. Claims made for such applications must fall within the general parameters prescribed in each of the following pest-specific discussions.

d. Treatment of Screens in Domestic Dwellings

As in the preceding case, a limited number of Deet formulations bear claims for the treatment of screens. Again, while no data directly related to screens are available, the Agency will extrapolate from the data available for

netting treatment. Screen treatments may not be claimed for all pests, however. Claims for screen application may only be made for mosquitoes, ceratopogonid species and black flies. Any claims made for these pests must fall within the parameters prescribed in each of the following pest-specific discussions.

e. Outdoor Mist Application

A single registered Deet product is claimed to be efficacious for outdoor mist application on and around lawn chairs, picnic tables, boats and other areas where repellency might be desired. The Agency has been unable to relate these sites or this method of application to available data. Since available data show that Deet is effective only as much as 4 to 8 centimeters out from a treated surface, (Kahn and Maibach 1972, MRID 00001163), the Agency contends that such applications are ineffective. These claims will not be supported under this standard. Registrants desiring to make these claims will be required to submit supporting efficacy data.

3. Target Pests

a. Black flies

Deet, applied in formulations where it is the sole active ingredient, has been demonstrated effective in repelling several Simuliidae species for periods of 2.6 to 10 hours. Formulations for which acceptable data have been reviewed range from 15 to 65% active ingredient. Although the available data are inconclusive in establishing the low dose range at which Deet might prove effective, they do indicate a distinct dose related response. A 15% Deet formulation, applied at 1 gram per 645 cm² was reported to effectively repel S. venustum for a period of 2.6 hours (S.C. Johnson and Son, Inc. 1974, MRID 00001095). Formulations of 50 and 60% were effective in extending the protection time to 10 and 10.6 hours respectively (Garnett and Lomax 1973, MRID 00001137).

b. Ceratopogonidae (Biting midges, sandflies)

Formulations containing a minimum of 15% Deet active ingredient, have been shown to be effective against the Ceratopogonidae. The dose-effect relationship is more obscure for this target pest, probably as a result of the somewhat smaller data base. Deet formulations ranging from 15 to 50% have been shown to protect against Ceratopogonidae species for periods of 3.0 to 7.0 hours.

c. Horse flies

There is an absence of data relating to the efficacy of Deet on horse fly species. Claims for horse flies, therefore, are not supported by this Standard. Any registrant desiring to add claims for horse flies must submit appropriate efficacy data.

d. Chiggers

All available data on Deet efficacy against chiggers were derived from laboratory studies involving cloth treatment. Although all the studies were laboratory tests, the results were so consistent that the Agency considers the data base adequate. Deet activity against chiggers primarily results

from contact toxicity rather than repellency. Studies indicate that Deet, applied to cloth test surfaces, often exhibits toxic action in less than 60 seconds after application (USDA 1959, MRID 00001165); (Gertler 1962, MRID 00001018). In tests conducted against Trombicula splendons, one of the two most frequently encountered chiggers species within the United States, cloth treated at 1 gram per square foot remained effective for a period of 15 days (Kochhar 1974, MRID 05000150); (Gilbert 1957, MRID 05000237). Trombicula alfreddugesi, was also killed at an application rate of 2 gm/ft² (Gerberg 1966, MRID 0001171).

The data base for technical Deet is adequate, even when restricted only to proof of Deet's toxicity to chigger species. It is also clear that Deet-impregnated cloth can remain effective over a period of several days.

e. Deerflies

The data reviewed have indicated that the deerfly is the least susceptible of all dipteran species to Deet products. Formulations ranging from 12.5 to 75% have been evaluated. The duration of acceptable repellency has ranged from 1 to 8 hours, with an average of 1 to 2 hours. There are distinct differences occurring between deerfly species. Schreck (Schreck et al. 1976 MRID 05002304) noted differences in Deet protection time when tested against Chrysops atlanticus and C. flavidus. C. flavidus appeared more responsive to Deet; a 12.5% formulation afforded 135 minutes of protection. C. atlanticus however, was repelled for only 13 to 32 minutes by a formulation containing 50% active Deet. Similar results were obtained in an additional study utilizing C. atlanticus as the test subject (S.C. Johnson and Son, Inc. 1974 MRID 00001095). In this study, Deet provided adequate repellent activity for 36 and 27 minutes for skin-applied formulations containing 10 and 15% Deet respectively. Available data establish that the Deet repels deerflies. The data also indicate that protection time varies considerably due to sensitivity of different species.

f. Fleas

Few studies are available on the efficacy of Deet products on fleas. Those studies which have been located and reviewed deal primarily with Xenopsylla and Ceratophyllus species. In laboratory studies (Kasafutdiner 1971, MRID 05000077), Deet applied to cloth at 40 grams of active ingredient per square meter provided 100% repellency against Ceratophyllus tesquorum for a period of 30 days. In field tests with humans (Fristane et al. 1970 MRID 05000589), a 40% Deet formulation applied to footwear and overalls at a rate of 80 grams of active ingredient per square meter afforded protection for 1-2 months against Xenopsylla gerbilli fleas. Dremova (Dremova et al. 1977, MRID 05000086) conducted laboratory trials and found that Deet treated cloth (20 grams active ingredient per square meter) provided over 90% repellency of Xenopsylla cheopis for a period of 15 days. In another laboratory study, Zolotarev and Stavroskaya (Zolotarev and Stavroskaya 1960, MRID 05004738) found the p-isomer of Deet to be the most effective in repelling a mixed population of fleas. The m-isomer, while somewhat less effective than the p-isomer, was effective for periods of 2 to 35 days; the variation in the protection period being largely a function of application rate. Despite the relatively small data base, the available data are adequate in establishing Deet repellency toward fleas for products containing 10% or more active ingredient of Deet. There is only a slight variation between the different species' response to Deet. There are no data on formulations containing 10% active ingredient or less. The Zolotarev

paper clearly establishes a dose/response relationship.

g. Leeches

Field and laboratory data indicate that Deet provides short-term protection against Haemadipsa and Hirudo leeches. Saxena et al. (Saxena, et al. 1969, MRID 05003639) conducted field tests to determine the efficacy of Deet against the terrestrial leech Haemadipsa sylvestris. Deet was reported effective against the test species for at least 2 days after treatment when applied to boots (6 milliliters) and anklets (4 milliliters). Four milliliters of Deet applied to the forearms, lower portion of the legs, neck and face of test subjects provided protection against the leech for at least 8 hours after treatment.

In tests conducted with aquatic leeches, Hirudo spp., Deet efficacy was marginal. Laboratory trials conducted with a 33% active ingredient preparation produced 30 minutes repellency. Field trials with the same formulation, however, provided only 12 minutes of protection. Additional field trials conducted with 45-50% Deet in varying inert formulations yielded protection times ranging from 1 to 50 minutes (Keegan et al. 1964, MRID 05002310). In a separate study, Keegan and Weaver (Keegan and Weaver 1964, MRID 00001035) evaluated 75% lotions and pressurized sprays containing 12.75% of Deet active ingredient. A 50% reduction in the number of leeches on a test subject was observed after five minute exposure periods. Neither formulation remained effective for a second five minute period.

The data indicate that formulations containing at least 50% Deet active ingredient adequately repels terrestrial leeches. With respect to aquatic leeches, the data are inadequate to support label claims.

h. Mosquitoes

(1) Laboratory Studies

Most of the laboratory studies reviewed evaluated the efficacy of Deet against Aedes aegypti using the "first confirmed bite" method. Formulations ranging from 6.25 to 100% active ingredient of Deet afforded protection for periods from 2 to 14 hours. The wide variation in response resulted from a variety of factors. These factors may have included subject susceptibility, individual differences in laboratory-reared populations, time of day of the test, and other undescribed environmental conditions. The data further indicate that Deet can provide at least 3 hours of protection against mosquitoes following direct skin applications of formulations containing 10% or greater of the single active ingredient. Laboratory tests conducted to determine the minimum effective dose indicate that at least 0.02 to 0.20 milligram of Deet per square centimeter of skin must be applied to protect the forearms of human subjects exposed to caged A. aegypti.

(2) Field Studies

Numerous studies have been conducted to evaluate the effectiveness of Deet against mosquitoes. The data demonstrate that Deet is effective in repelling a wide spectrum of mosquito species, i.e., Aedes, Culex, Mansonia, Anopheles, and Psorophora. The protection times afforded by various dose levels ranged from 1 to 24 hours. Following exposure to 100% active ingredient Deet, maximum protection times of 20 hours were obtained against Culex pipiens and mixed

populations comprised primarily of Aedes and Mansonia species. Formulations ranging from 10 to 50% afforded adequate protection from both Aedes and Culex mosquitoes.

Both the field and laboratory data currently available clearly indicate that Deet is effective in repelling mosquitoes.

i. Stableflies

A relatively large number of papers dealing with Deet efficacy against the stablefly (Stomoxys calcitrans) are available. They provided a clear indication that Deet is effective in protecting human subjects from stablefly biting. A distinct dose/response relationship exists. The minimum protection time afforded a human test subject was 2.7 hours (S.C. Johnson and Son, Inc. 1974, MRID 00001095). This protection time resulted from a 1 gram per 645 cm² application of a 10% single active Deet formulation, the lowest percent active ingredient formulation for which data appear available. The maximum protection time indicated by the data was 8 hours, afforded by a 25% Deet active ingredient and ethanol formulation (Gilbert et al. 1957, MRID 05000236). Although formulation strengths up to 50% have been tested, the 8 hour duration reported by Gilbert was the maximum recorded for all studies. It can not be inferred, however, that 25% is the optimum Deet concentration for protection against stableflies, since other factors (wind, precipitation, etc.) also affect the product's effectiveness.

The data currently available clearly indicate that Deet is effective in repelling stableflies.

j. Ticks

Wide variations in response levels have been noted in data on Deet applied to ticks. The variation appears relative to species, sex and life stage (Simironva and Dremova 1971, MRID 05002661) (Novak 1973, MRID 05000290). Even in studies conducted on the same subjects, considerable variation in test results are reported. In studies involving the Lone Star Tick, Amblyomma americanum, Gilbert (Gilbert 1957, MRID 0500237) reported marginal repellency when Deet was applied to cloth test surfaces at the rate of 2 grams active ingredient per square foot. Eight hour repellency was recorded at 52% active ingredient. In a field study; however, Gerberg (Gerberg 1966, MRID 00001171) reported adequate repellency for 7 hours utilizing the same 2 gram active ingredient per square foot application rate. Similarly, studies conducted on the Brown Dog Tick, Rhiphicephalus sanguineus, provided divergent results. In laboratory trials, Deet-treated cardboard (70% m-isomer at 5 milliliters per square inches) yielded virtually no repellent activity (Kochhar 1974, MRID 05000150). A 25% Deet cream formulation, however, provided 21 to 47 hours of protection when applied at a rate of 0.018 - 0.028 milligrams per cm² to the shaved back of rabbits.

Although the Agency has reviewed papers dealing with a wide range of tick species, most available data are from foreign sources and involve species not found in the United States.

Among the species of concern to the Agency, Demacentor andersoni, D. variabilis, Ixodes scapularis, and I. pacificus do not appear within the literature. Given the absence of data relating to the majority of the tick species and the inconclusive nature of the overall data base, claims for tick

repellency are not adequately substantiated. Claims for ticks, therefore, are not supported by this Standard. Any registrant desiring to retain or add claims for ticks must submit the appropriate efficacy data.

C. Data Gaps

1. Use Sites

Registrants desiring to make claims of Deet efficacy for outdoor mist application will be required to submit supporting data.

2. Target Pests

a. Ceratopogonidae

The Agency will require formulation-specific data in each instance where a specific duration claim is requested.

b. Fleas

The Agency will require formulation-specific data in each instance where a specific duration claim is requested.

c. Chiggers

Virtually no data are available related to formulated products. Rather than require formulation-specific data for repellancy, at the time of product reregistration, registrants claiming on the label, "kills chiggers", must submit the calculations that demonstrate that the product will satisfactorily deposit a minimum of 1 gram of Deet active ingredient per square foot of treated cloth, when used in accordance with label directions. Since the Agency will not prescribe acceptable duration claims, whenever a claim of product duration is requested, formulation-specific data must be developed and submitted for Agency evaluation.

d. Leeches

Registrants desiring claims for aquatic leeches will be required to submit data derived from testing of the specific formulation proposed for (re)registration.

e. Mosquitoes

Registrants desiring duration claims of more than two hours will be required to submit formulation specific data.

f. Stableflies

Deet products, of any formulation type and containing 10% or more active ingredient, may bear claims for stablefly repellency. Sufficient data are available to allow for a general, limited duration claim. Utilizing the previous logic for mosquitoes, products containing 10% or greater concentration of Deet active ingredient may claim "repels stableflies for up to two hours." Registrant who wish to put this claim on their labels must also include a brief statement to the effect that the duration of protection may vary with the individual and that re-application may be required. Registrants desiring duration claims for periods longer than two hours will be required to submit

formulation-specific data.

D. Labeling

1. Use Sites

a. Human Skin, Clothing, Tents and Bedrolls

Claims may be made for biting flies, chiggers, deer flies, mosquitoes, fleas and stableflies.

b. Treatment of Screens in Domestic Dwellings

Claims for screen application may only be made for mosquitoes, ceratopogonid species, and black flies.

2. Target Pests

a. Black flies

Data currently available clearly indicate that Deet is effective in repelling black flies. The data also indicate efficacy of various doses and formulations. The Agency, therefore, will set limits on product labels. The lower limit of formulations tested was 15%, so the Agency will not accept efficacy claims for products containing less than 15% Deet as the sole active ingredient. Because of the limits of the data reviewed, the Agency cannot prescribe claims of product duration for all formulations, strengths and types. The Agency will require formulation-specific data in cases where producers wish to make a claim.

b. Deerflies

Given the extreme variation in protection times for different species, the Agency will not accept duration claims for deerflies. Duration claims for deerfly protection, therefore, will be refused for Deet products. Deet formulations containing 10% or more active ingredient however, may bear a limited, general claim for deer fly repellency.

c. Leeches

Because there are no terrestrial leeches in the U.S., the Agency does not believe that the public interest would be served by permitting these claims on consumer product labeling. In the past, there have been occasions where these products were required for use by American military forces. The Agency is prepared to permit the labeling of products for terrestrial leeches with the qualification that all such products bear the statement "For Use in Tropical Areas Where Pest Occurs."

d. Mosquitoes

Deet products of any formulation or type, containing 10% or more active ingredient, may bear claims for mosquito repellency. With regard to duration claims, the Agency will not prescribe claims for all variations of formulation strengths and types. There do appear, however, sufficient data to allow for a general, limited duration claim. Products containing 10% or greater concentrations of Deet may claim "repels mosquitoes for up to two hours."

Registrants desiring to put this claim on their labels must also include a brief statement to the effect that the duration of protection may vary with the individual and that re-application maybe required.

APPENDIX I
SUPPORTING COMPANY EFFICACY DATA

CITATION	Blackflies	Ceratopogonid species	Chiggers	Deerflies	Fleas	Houseflies	Leeches	Mosquitoes	Stableflies	Ticks
Allenbaugh (1965) 00001003				X	X			X	X	X
Altman (1969) 00001172								X		
Eastman (1969) 00001169	X			X				X		
Gerberg (1966) 00001171	X	X	X		X			X		X
Gerberg (1973) 00001065								X		
Gertler et al. (1962) 00001018			X					X		X
Gilbert et al. (1957) 00001021		X		X				X	X	
Gilbert et al. (1957) 00001022			X		X			X		X
Gilbert et al. (1965) 00001033						X				

APPENDIX I

SUPPORTING COMPANY EFFICACY DATA

CITATION	Blackflies	Ceratopogonid species	Chiggers	Deerflies	Fleas	Houseflies	Leeches	Mosquitoes	Stableflies	Ticks
Gilbert et.al. (1970) 00001162								X		
Gilbert et.al. (1970) 00001167								X	X	
Gouck et.al. (1967) 00001032							X			
Granett (1969) 00001109				X				X	X	
Granett et.al. (1971) 00001138	X							X		
Granett et.al. (1973) 00001137	X									
Granett et.al. (1973) 00001127	X									
Grothaus et.al. (1975) 00001038	X	X		X				X		
Haynes (1969) 00004864				X				X	X	

APPENDIX I

SUPPORTING COMPANY EFFICACY DATA

[illegible]

APPENDIX I

SUPPORTING COMPANY EFFICACY DATA

CITATION	Blackflies	Ceratopogonid species	Chiggers	Deerflies	Fleas	Houseflies	Leeches	Mosquitoes	Stableflies	Ticks
Lahr (1970) 00004870								X		
Lahr (1972) 00001124	X			X				X	X	
Lomax (1968) 00001110	X	X		X				X	X	
Lomax (1968) 00004865	X	X		X				X	X	
Lomax (1969) 00001118	X	X		X				X	X	
Lomax (1970) 00001115								X	X	
Lomax (1970) 00001117								X	X	
Mace (1966) 00001084								X		
McAndless (1974) 00001037				X				X		

APPENDIX I

SUPPORTING COMPANY EFFICACY DATA

CITATION	Blackflies	Ceratopogonid species	Chiggers	Deerflies	Fleas	Houseflies	Leeches	Mosquitoes	Stableflies	Ticks
MGK Co. (1957) 00001055		X	X	X	X			X	X	X
MGK Co. (1974) 00001053								X		
Miller (1955) 00001000									X	
Mitchell (1971) 00001129						X				
Neumiller (1972) 00001072								X	X	
Pierce (1958) 00001104			X					X	X	
S.C.Johnson (1964) 00001168								X	X	
S.C.Johnson (1969) 00001170								X	X	
S.C.Johnson (1974) 00001095	X			X				X	X	

APPENDIX I

SUPPORTING COMPANY EFFICACY DATA

CITATION	Blackflies	Ceratopogonid species	Chiggers	Deerflies	Fleas	Houseflies	Leeches	Mosquitoes	Stableflies	Ticks
S.C.Johnson (1974) 00001097								X	X	
S.C.Johnson (1977) 00001103	X	X		X				X	X	
Schley et.al. (1974) 00001052								X		
Smith (1957) 00001113			X		X			X		X
Smith (1970) 00001166								X		
Smith et.al. (1957) 00001010								X		
Smith et.al. (1959) 00001164			X					X		X
Smith et.al. (1961) 00001184								X		
Union Carbide (1970) 00001116								X	X	

APPENDIX I

SUPPORTING COMPANY EFFICACY DATA

[illegible]

VIII. CASE BIBLIOGRAPHY

Guide to Use of This Bibliography

1. Content of Bibliography. This bibliography contains citations of all the studies reviewed by EPA in arriving at the positions and conclusions stated elsewhere in this standard. The bibliography is divided into 3 sections: (1) citations that contributed information useful to the review of the chemical and considered to be part of the data base supporting registrations under the standard; (2) citations examined and judged to be inappropriate for use in developing the standard; and (3) standard reference material. Primary sources for studies in this bibliography have been the body of data submitted to EPA and its predecessor agencies in support of past regulatory decisions, and the published technical literature.
2. Units of Entry. The unit of entry in this bibliography is called a "study". In the case of published materials, this corresponds closely to an article. In the case of unpublished materials submitted to the Agency, the Agency has sought to identify documents at a level parallel to a published article from within the typically larger volumes in which they were submitted. The resulting "studies" generally have a distinct title (or at least a single subject), can stand alone for purposes of review, and can be described with a conventional bibliographic citation. The Agency has attempted also to unite basic documents and commentaries upon them, treating them as a single study.
3. Identification of Entries. The entries in this bibliography are sorted by author, date of the document, and title. Each entry bears, to the left of the citation proper, an eight-digit numeric identifier. This number is unique to the citations, and should be used at any time specific reference is required. This number is called the "Master Record Identifier", or "MRID". It is not related to the six-digit "Accession Number" which has been used to identify volumes of submitted data; see paragraph 4(d)(4) below for a further explanation. In a few cases, entries added to the bibliography late in the review may be preceded by a nine-character temporary identifier. This is also to be used whenever a specific reference is needed.
4. Form of the Entry. In addition to the Master Record Identifier (MRID), each entry consists of a bibliographic citation containing standard elements followed, in the case of materials submitted to EPA, by a description of the earliest known submission. The bibliographic conventions used reflect the standards of the American National Standards Institute (ANSI), expanded to provide for certain special needs. Some explanatory notes of specific elements follow:
 - a. Author. Whenever the Agency could confidently identify one, the Agency has chosen to show a personal author. When no individual was identified, the Agency has shown an identifiable laboratory or testing facility as author. As a last resort, the Agency has shown the first known submitter as author.

- b. Document Date. When the date appears as four digits with no question marks, the Agency took it directly from the document. When a four-digit date is followed by a question mark, the bibliographer deduced the date from evidence in the document. When the date appears as (19??), the Agency was unable to determine or estimate the date of the document.
- c. Title. This is the third element in the citation. In some cases it has been necessary for Agency bibliographers to create or enhance a document title. Any such editorial insertions are contained between square brackets.
- d. Trailing Parentheses. For studies submitted to us in the past, the trailing parentheses include (in addition to any self-explanatory text) the following elements describing the earliest known submission:
 - (1) Submission Date. Immediately following the word 'received' appears the date of the earliest known submission.
 - (2) Administrative Number. The next element, immediately following the word 'under', is the registration number, experimental permit number, petition number, or other administrative number associated with the earliest known submission.
 - (3) Submitter. The third element is the submitter, following the the phrase 'submitted by'. When authorship is defaulted to the submitter, this element is omitted.
 - (4) Volume Identification. The final element in the trailing parenthesis identifies the EPA accession number of the volume in which the original submission of the study appears. The six-digit accession number follows the symbol 'CDL', standing for "Company Data Library". This accession number is in turn followed by an alphabetic suffix which shows the relative position of the study within the volume. For example, within accession number 123456, the first study would be 123456-A; the second, 123456-B; the 26th, 123456-Z; and the 27th, 123456-AA.

Section I

Farm Chemical Handbook. (1979) Meister publishing. Willoughby. Ohio.

The Federal Insecticide, Fungicide, and Rodenticide Act, as amended in 1978,
7th U.S. Code, Chapter 135, 61 Statute 163.78 Statute 190.

Pesticide Process Encyclopedia, 1977 Noyes Data Corp., Park Ridge, New
Jersey.

Pesticide Index: Basic information on the chemicals used as active
components of pesticides.; Martin, H., and Worthington, C.R., eds., 1977.
5th ed., British Corp. Protection Council, Worcestershire, England.

U.S. Environmental Protection Agency (1978a); Proposed Guidelines for Regis-
tering Pesticides in the United States. Federal Register, 43 (132) 29696.

U.S. Environmental Protection Agency (1978b); Proposed Guidelines for Regis-
tering Pesticides in the United States; Hazard Evaluation: Humans and
Domestic Animals. Federal Register, 43(163) 37336.

U.S. Environmental Protection Agency (1980); Regulations for the Enforcement of
the Federal Insecticide, Fungicide, and Rodenticide Act, Title 40, Chapter
1, Part 162.

U.S. Environmental Protection Agency (in press) Proposed Guidelines for Regis-
tering Pesticides in the United States. Subparts G (Product Performance)
H (Label Development).

Section II

OFFICE OF PESTICIDE PROGRAMS REGISTRATION STANDARD BIBLIOGRAPHY

Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

PRODUCT CHEMISTRY

MRID	CITATION
05002553	Blaine, R.L.; Levy, P.F. (1974) The use of thermal evolution analysis (TEA) for the determination of vapor pressure of agricultural chemicals.
GS0002014	Bogard, T. 1976. Information on manufacturing process of deet. (Unpublished data submitted by MGK Co., June 1, 1976.)
00001008	Cutter Laboratories (1975) Diethyltoluamide: Analytical Test Methods: [Gas chromatography dated Sep 1973 and IR spectroscopic determination dated Feb 1975]. (Unpublished study received Jun 11, 1975 under 121-17; CDL:106619-A)
05000251	Ellis, L.C.; Kise, M.A., inventors; Virginia Chemicals, Inc., assignee (1972) Verfahren zur Herstellung von N,N-Diethyl-m-Toluamid in Waebrieger Atzalkaliloesung. [N,N-diethyl-m-toluamide] GDY 2, 153, 307. May 4. 13p. Int. Cl-2 AOIH 9/20
00001152	Hercules Incorporated (1972) Properties: Physical and Chemical: [Metadelphene]: AP-105B. (Incomplete study, p. 2; unpublished study received Jun 12, 1972 under 11525-12; submitted by Peterson Puritan, Inc., Danville, Ill.; CDL*004536-E)
00001068	Hercules Incorporated (1977) Metadelphene(R): The All-Purpose Insect Repellent: Bulletin AP-105B. (Unpublished study received Jul 7, 1969 under 1516-52; submitted by Curtis Laboratories, Inc., Kansas City, Mo.; CDL:024967-B)
00001025	Hercules Powder Company (1957) Meta Delphene "The Complete Insect Repellent." Wilmington, Del., Hercules Powder Co., Inc. (Bulletin No. 213; Also In unpublished submission received Aug 13, 1957 under unknown admin. no.; submitted by Hercules Powder Co., Wilmington, Del.; CDL:108471-A)
00001036	Pfizer, Incorporated (1976) Product Chemistry for Technical Chemical: Deet Insect Repellent. (Unpublished study that includes manufacturing process and analytical methods, received Sep 13, 1976 under 1007-27; submitted by Pfizer, Inc., New York, N.Y.; CDL:226112-A)
05000048	Sarmiento, R.; Beroza, M. (1975) Analysis of the insect repellent deet and its isomers by gas chromatography on graphitized carbon black. Journal of Economic Entomology 68(2): 258-260

00001100

S.C. Johnson & Son, Incorporated (1977) Chemical and Physical Properties [of Diethyltoluamide]: Summary. (Unpublished study received Mar 21, 1978 under 4822-160; CDL:233261-S)

05000597

Voronkina, T.M.; Latyshev, V.I.; Zhukova, E.V.; Yudina, I.P.; Sakodinskii, K.I. (1971) Khromatografiya v opredelenii nekotorykh repellentov i ikh primesei [chromatography in the determination of some (insect) repellents and their impurities] Trudy, Vsesoyuznyi Nauchno-Issledovatel'skiy Institut Veterinarnoi Sanitaii 39:189-192

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

ENVIRONMENTAL FATE

MRID	CITATION
GS0002001	Hales, Y. and H.E. Radtke Jr. Use Data Related to Exposure for Deet. GEOMET Technologies, Inc., under Contract #68-01-5155
GS0002002	S.C. Johnson & Son, Inc. 1979. Consumer Exposure & Use Rates of Insect Repellents. (This study includes confidential business information.)
GS0002003	S.C. Johnson & Son, Inc. 1980. DEET Consumer Use Information. (This package contains studies using confidential business information.)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

TOXICOLOGY

MRID	CITATION
00001051	Ambrose, A.M., Huffman, D.K., and Salamone, R.T. 1959 Pharmacologic and toxicologic studies on N,N-diethyltoluamide: I. N,N-Diethyl-m-toluamide. Toxicol. Appl. Pharmacol. 1(1):97- 115. (MRID:05000240. 17A-0036) (Also in unpublished submission received Oct. 22, 1974 under 1021-1323; prepared by U.S. Army Environmental Health Lab., Toxicology Div., submitted by McLaughlin Gormley King Co., Minneapolis, Minn; CDI:028339-B)
00001102	Ambrose, A.M., Yost, D.H. 1965 Pharmacologic and toxicologic studies on N,N-diethyltoluamide: II. N,N-Diethyl-o-toluamide and N,N-Diethyl-p-toluamide. Toxicol. Appl. Pharmacol. 7(6): 772-780 (MRID: 00001102, 17A-0023) (Also in unpublished submission received Mar 21, 1978 under 4822-160; prepared by U.S. Army Environmental Health Agency, Toxicology Div., in cooperation with U.S. Army Medical Research Laboratory, Pathology Div., submitted by S.C. Johnson & Son, Inc., Racine, Wis.; CDI:233262-C)
GS0002016	Beliles, R.P., and HORN, H.J. 1967. Acute Oral Toxicity to Rats of 2370D-89: Repeated Dermal Application to Rabbits and Single Application to the Eyes of Rabbits. (Unpublished study that includes tables 1-21 and appended letter, received Jul 10, 1976, under 4822-10; prepared by Woodard Research Corp., submitted by S.C. Johnson and Son, Inc., Racine, Wis.; CDL:233491-D)
GS0002004	Blau, S.J., and KANOF, N. 1961. CDK 79/2-88/1: Patch Tests: DEET. (Unpublished study that includes letter, received December 19, 1961, under 6650-1; submitted by Revlon, Inc., Bronx, N.Y.; CDL:229503-B)
05000001	Blomquist, L., Stroman, L., and Thorsell, W. 1975, Distribution and fate of the insect repellent 14C-N,N-diethyl-m-toluamide in the animal body: I. Distribution and excretion after injection into mice. Acta Pharmacol. Toxicol. 37:121-133 (MRID: 05000001, 17A-0033)
05000002	Blomquist, L., and Thorsell, W. 1977, Distribution and fate of insect repellent 14C-N,N-diethyl-m-toluamide in the animal body: II. Distribution and excretion after cutaneous application. Acta Pharmacol. Toxicol. 41:235-243 (MRID: 05000002. 17A-0038)
05000243	Carpenter, C.P., Weil, C.S. and Smyth, H.F., Jr. 1974, Range- finding toxicity data: List VIII. Toxicol. Appl. Pharmacol. 28:313-319 (MRID: 0500243. 17A-0024)

- 00001139 Davidow, B. 1960, Lee's Sports' Balm: Acute Oral Toxicity, Primary Irritation, Acute Dermal Toxicity, Eye Irritation. (Unpublished study that includes tables 1-5, received June 14, 1960, under 7165-1; prepared by New Drug Institute, submitted by William W. Lee and Co., Inc., Watervliet, N.Y.; CDL:050672-A) (MRID: 00001139, 174-0021c)
- GS0002005 Durloo, R.S., and Woodard, M.W. 1971a. Safety Evaluation by Repeated Dermal Application to Rabbits: Formulation 3369D138 and Formulation 3369D139. (Unpublished study that includes tables A-1-A-10, received September 9, 1971, under 4822-114; prepared by Woodard Research Corp., submitted by S.C. Johnson and Son, Inc., Racine, Wis.; CDL:133553-C)
- GS0002006 Durloo, R.S., and Woodard, M.W. 1971b. Safety Evaluation by Test for Irritation in Rabbits: Formulation 3369D135-1. (Unpublished study that includes tables A-1-A-15 and toxicology data, received September 9, 1971, under 4822-RRU (114); prepared by Woodard Research Corp., submitted by S.C. Johnson and Son, Inc., Racine, Wis.; CDL:133553-B)
- GS0002007 Evers, W.D. 1969. The Dermal Toxicity of Insect Repellent Gel D-554-9: Report No. WDE 69:11. (Unpublished study received February 12, 1969, under 134-48; submitted by Hess and Clark, Research Department, Division of Richardson Merrell, Inc., Ashland, Ohio; CDL:000602-B)
- 05003588 Feldmann, R.J., and Maibach, H.I. 1970. Absorption of some organic compounds through the skin in man. J. Invest. Dermatol. 54-399-404 (MRID: 05003588, 17A-0005)
- 05000134 Ficsor, G., Nii, L.O.; and Piccolo, G.M. 1972. Survey of pesticides for mutagenicity by the bacterial-plate assay method. E.M.S. Newsletter 6:6-8 (MRID: 05000134, 17A-0026)
- 05000061 Gleiberman, S.E., and Volkova, A.P. 1974. [Comparative toxicity of diethyltoluamide synthesized on petroleum and coal metaxylol.] NTIS AD/A-000 176 (Translated from Russian) (MRID: 05000061, 17A-0008)
- 05000007 Gleiberman, S.E., Volkova, A.P., Nikolaev, G.M., and Zhukova, E.V. 1975. [Embryotoxocity of the insect repellent diethyltoluamide.] Farmakol. Toksikol. 38:202-205 (Translated from Russian) (MRID: 05000007, 17A-0027)
- 05000008 Gleiberman, S.E., Volkova, A.P., Nikolaev, G.M., and Zhukova, E.V. 1976. [Study of the remote results of the use of repellents: I. Experimental study of the effects remote results of long-term exposure to the repellent diethyltoluamide (DEET).] Med. Parazitol. (Mosk.) 45:65-69 (Translated from Russian) (MRID: 05000008, 17A-0010)

- 00001063 Grahwit, G., Sterner, W., and Chibanguza, G. 1977. Effect of "MGK Diethyltoluamide" on the Embryonic Development of Rats after Oral Application. (Unpublished study that includes appended letter, received May 23, 1977, under 1021-891; prepared by International Bio-Research, Inc., submitted by McLaughlin Gormley King Co., Minneapolis, Minn.; CDL:230396-A) (MRID: 00001063, 17A-0028)
- 05000328 Gryboski, J., Weinstein, D., and Ordway, N.K. 1961. Toxic encephalopathy apparently related to the use of an insect repellent. N. Engl. J. Med. 264:289-291 (MRID: 05000328, 17A-0029)
- GS0002008 Horn, H.J. 1968. Repeated Dermal Application to Rabbits 2693D102. (Unpublished study received Jul 10, 1976, under 4822-10; prepared by Woodard Research Corp., submitted by S.C. Johnson and Son, Inc., Racine, Wis.; CDL:233491-E)
- 00001080 Howard, D.J.; Woodard, M.W. (1971) Acute Oral Toxicity to Rats: Formulation 3369D135-1. (Unpublished study that includes toxicology data, received Sept 9, 1971 under 4822-114; prepared by Woodard Research Corp., submitted by S.C. Johnson & Sons, Inc., Racine, Wis.; CDL:133553-A)
- 00001073 Johnston, C.D. (1972) Safety Evaluation by Acute-Dermal Irritation Test in the Rabbit: 3560D105-1 and 3560D138-2. (Unpublished study that includes tables and formulations, received Jun 12, 1972 under 4822-119; prepared by Woodard Research Corp. submitted by S.C. Johnson & Son, Inc., Racine, Wis.; CDL:007514-B)
- 00001039 Jolly, E.R.; Sloughfy, C.A. (1974) Irritation and Sensitization Potential of Insect Repellent: Final Report: IC-1433. (Unpublished study received Feb 16, 1977 under 1021-567; prepared by Biometric Testing, Inc., submitted by McLaughlin Gormley King co., Minneapolis, Minn.; CDL:228664-A)
- 05000271 Kingscote, A.A., and Allen, J.R. 1959. The fate of Insect Repellents Applied to Skin. (Unpublished study received August 4, 1960, under DRB Extra-Mural) (MRID: 05000271, 17A-0030)
- 05002807 Kurasova, V.V., and Leschev, V.V. 1969. [Toxicity of benzimine, diethyltoluamide, and carboxide for rabbits and cattle during their external application.] Tr. Vsesoy. Nauch. Iasled, Inst. Vet, Sanit, 33:235-245 (Translated from Russian) (MRID: 05002807, 17A-0011)
- 050023008 Lamberg, S.I., and Mulrennan, J.A., Jr. 1969, Bullous reaction to diethyltoluamide (DET): Resembling a blistering insect eruption,... Arch. Dermatol. 100:582-586 (MRID: 050023008, 17A-0012)
- GS0002036 Litton Bionetics, Inc. 1977. Mutagenicity Evaluation of Diethyltoluamide technical. Final report submitted to the U.S. Army, Project #20838.

- GS0002009 Lukes, T.H., and PAA, H. 1975. 90-Day Subacute Dermal Toxicity Study with 3989BN69-1 in Albino Rabbits: IBT No. 601-06046. (Unpublished study that includes tables I-XLIX, received July 10, 1976, under 4322-10; prepared by Industrial Bio-Test Laboratories, Inc., submitted by S.C. Johnson and Son, Inc., Racine, Wis.; CDL:233491-J)
- 0500020 Lure, A.A., Gleiberman, S.E., and Tsizin, Y.S. 1978. [Pharmacokinetics of the repellent N,N-diethyl-3-methyl-benzamide.] Med. Parazit. (Mosk.) 47:72-77 (Translated from Russian) (MRID: 0500020, 17A-0000)
- 05000021 Maibach, H.I., and Johnson, H.L. 1975. Contact urticaria syndrome: Contact urticaria to diethyltoluamide—immediate-type hypersensitivity. Arch. Dermatol. 111:726-730 (MRID: 05000021, 17A-0031)
- 05000599 Markina, V.V. and Yatsenko, V.K. 1971. [A comparative evaluation of the absorbability of the repellents diethyltoluamide, benzimine, and carboxide in the form of lotions and creams] Zdravookhr. Turkmen. 15:41-44 (Translated from Russian) (MRID: 05000599, 17A-0014)
- GS0002010 Pesticide Incident Monitoring System (PIMS). 1978. Summary of Reported Pesticide Incidents Involving DEET. Report No. 121. Human Effects Monitoring Branch, Office of Pesticide Programs, U.S. Environmental Protection Agency, Washington, D.C.
- 05000242 Phillips, L., Jr., Steinberg, M., Maibach, H.I., and Akers, W.A. 1972. A comparison of rabbit and human skin response to certain irritants. Toxicol. Appl. Pharmacol. 21:359-382 (MRID: 05000242, 17A-0032)
- 05000693 Rabinovich, M.V. 1966. [The effect of diethyltoluamide on the human skin and mucous membranes of the eyes.] Zh. Mikrobiol. Epidemiol. Immunobiol. 43:109-112 (Translated from Russian) (MRID: 05000693, 17A-0016)
- 05000174 Schmidt, D.H., Acree, F., Jr., and Bowman, M.C. 1959. Fate of C¹⁴ diethyltoluamide applied to guinea pigs. J. Econ. Entomol. 52:928-930 (MRID: 05000174, 17A-0017)
- GS0002011 Seater, S., and Horn, H.J. 1967. Repeated Dermal Application to Rabbits and Single Application to the Eyes of Rabbits of 2370D89, 2539D91-1, and 2539D91-3. (Unpublished study that includes tables, received July 10, 1976, under 4822010; prepared by Woodard Research Corp., submitted by S.C. Johnson and Son, Inc., Racine, Wis.; CDL:233491-F)
- 05007487 Spencer, T.S., Hill, J.A., Feldman, R.J., and Maibach, H.I. 1979. Evaporation of diethyltoluamide from human skin in vivo and in vitro. J. Invest. Dermatol. 72:317-319 (MRID: 05007487, 17A-0034)

- 05000045 Stenbaeck, P. (1977) Local and systemic effect of commonly used agents: Lifetime studies of 16 compounds in mice and rabbits. *Acta Pharmacologica et Toxicologica* 41(5):417-431
- 05001247 Swentzel, K.C. 1977. Investigation of N,N-Diethyl-m-toluamide (M-DET) for Dominant Lethal Effects in the Mouse. Aberdeen Proving Ground, Md.; U.S. Army Environmental Hygiene Agency (USAEHA Study No. 51-0034-78, available from NTIS, Springfield, Va.; AD-A508 414/4GA) (MRID: 05001247, 17A-0019)
- GS0002021 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special study No. 75-51-0034,80. Appendix I: Investigation of N,N-Diethyl-M-toluamide (M-DET) for Dominant Lethal Effects (17D-0008)
- GS0002022 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix J: Mutagenicity Evaluation of Diethyltoluamide Technical Grade LB-468-77 Lot No. 7141 (17D-0009)
- GS0002023 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix P: Prenatal Toxicity in Rabbits (17D-0015)
- GS0002024 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix S: Determination of Metabolites of M-DET in Urine of Rabbits (17D-0018)
- GS0002025 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix B: Protocol-Primary Eye Irritation (17D-0001)
- GS0002026 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix A: Protocol-Primary Skin Irritation (17D-0000).
- GS0002027 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study 75-51-0034,80. Appendix C: Primary Eye Irritation (17D-0002)
- GS0002028 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix D: Photochemical Skin Irritation-Evaluation Protocol (17D-0003)

- GS0002029 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix E: Protocol LD₅₀ Determinations (Dermal) (17D-0004)
- GS0002030 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix F: Acute Oral LD₅₀ Determinations (17D-0005)
- GS0002031 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix H: Protocol-Guinea Pig Sensitization Test (17D-0007)
- GS0002032 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix K: Protocol-Saturated Vapor Inhalation Toxicity Study (17D-0010)
- GS0002033 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix L: Protocol-Acute Aerosol Inhalation Toxicity (LC₅₀) (17D-0011)
- GS0002034 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix M: Inhalation Toxicities of N,N-diethyl-meta-toluamide (M-DET) (17D-0012)
- GS0002035 U.S. Army Environmental Hygiene Agency. 1979. Preliminary assessment of relative toxicity of insect repellent N,N-diethyl-meta-toluamide. Special Study No. 75-51-0034,80. Appendix Q: Dermal Penetration of 14C-labeled M-DET (17D-0016)
- GS0002036 U.S. Army Environmental Hygiene Agency. 1980. Effect of Dermal Application of N,N-Diethyl-Meta-Toluamide (M-DET) on the Embryonic Development of Rabbits. Study No. 75-81-0034-80.
- 00001093 Vos, B.J. (1972) Insect Repellent Off. Repeated Insult Human Patch Test. (Unpublished study that includes table, received July 10, 1976 under 4822-10; prepared by Woodard Research Corp., submitted by S.C. Johnson & Son, Inc., Racine, Wis.; CDL:233491-K)
- 00001099 WARF Institute, Incorporated. 1974. Spray in Eye Irritation of 4071 D9-3: WARF No. 4044151. (Unpublished study received Aug 7, 1974 under 4822-138; submitted by S.C. Johnson & Son, Inc., Racine, Wis.; CDL:140048-B). File No. 00001099 (AA0085)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

EFFICACY

MRID	CITATION
00001003	Allenbaugh, R.B. (1965) Review of Literature on N,N-Diethyl-m-toluamide as an Insect Repellent. (Unpublished study received Feb 12, 1969 under 134-38; submitted by Hess and Clark, Research Department, Division of Richardson-Merrell, Inc., Ashland, OH; CIL:000602-C)
00001172	Altman, R.M. (1969) Repellent tests against <u>Anopheles albimanus</u> Wiedemann in the Panama Canal Zone. Mosquito News 29 (1): 110-112. (Also In unpublished study received Jun 23, 1969 under 4822-EX-8; submitted by S.C. Johnson and Son, Inc., Racine, WI; CIL:127306-D)
05000167	Altman, R.M.; Smith, C.N. (1955) Investigations of repellents for protection against mosquitoes in Alaska, 1953. Journal of Economic Entomology 48(1):67-72
05002317	Axtell, R.C. (1967) Evaluations of repellents for <u>Hippelates</u> eye gnats. Journal of Economic Entomology 60(1):176-180.
05000211	Bar-Zeev, M.; Ben-Tamar, D. (1968) Effectiveness of repellents on cloth as determined by oviposition of <u>Aedes aegypti</u> . L. Mosquito News 28(3):396-403.
05002987	Bar-Zeev, M.; Ben-Tamar, D. (1971) Evaluation of Mosquito repellents. Mosquito News 31(1):56-61.
05000216	Bar-Zeev, M.; Ben-Tamar, D.; Gothilf, S. (1974) Field evaluation of repellents against mosquitoes in Israel. Mosquito News 34(2):199-203.
05000201	Bar-Zeev, M.; Gothilf, S. (1972) Laboratory evaluation of flea repellents. Journal of Medical Entomology 9(3): 215-218.
05000173	Bar-Zeev, M.; Schmidt, C.H. (1959) Action of a repellent as indicated by a radioactive tracer. Journal of Economic Entomology 52(2):268-269.
05000172	Bar-Zeev, M.; Smith, C.N. (1959) Action of repellents on mosquitoes feeding through treated membranes or on treated blood. Journal of Economic Entomology 52(2):263-267.
05000284	Berry, R.; Joseph, S.R.; Langford, G.S. (1965) The question of area mosquito repellents. Pages 190-193, In New Brunswick, N.J.: New Jersey Mosquito Extermination Association, New Brunswick, NJ.

- 05002313 Boparai, M.S.; Varma, R.N. (1971) Field evaluation of diethyltoluamide (deet) as a mosquito repellent. Indian Journal of Medical Research 59(3):504-509.
- 05000093 Bukshytov, V.I. (1964) Comparative tests of repellents for protection of bighorn cattle from pests. Trudy, Vsesoyuznyi Nauchno-Issledovatel'skii Institute Veterinarnoi Sanitarii. Transactopms. [All-Union Scientific Research Institute of Veterinary Science.] 24:44-50.
- 05006505 Burton, D.J. (1969) Intrinsic mosquito repellency values of some chemical compounds. American Perfumer and Cosmetics 84(4):41-42,44.
- 05000101 Carrillo, A.L. (1972) Barrier coatings for skin to bind Deet. Springfield, VA: NTIS (NTIS AD-763 217).
- 05000184 Catts, E.P. (1968) Deet-impregnated net shirt repels biting flies. Journal of Economic Entomology 61(6):1765-1766.
- 05000066 Cherepanov, A.I.; Gomoionova, N.P. (1963) Primenenie setok, propitanny'kh dietilamidom metatoluilovoi kisloty, pri [Use of netting impregnated with metatoluic acid diethylamide for individual protection of humans from horseflies and mosquitoes.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasite Diseases] 32(3):341-343.
- 05003637 Cherepanov, A.I.; Gomoionova, N.P.; Bogomiakova, N.P.; Dariichuk, Z.S. (1963) Novoe sredstvo boryby s gnusom. [A new means of gnat control.] Lesnaya Promyshlennost. [Forest Industry.] 6:12.
- 05000132 Clyde, D.F.; Kingazi, H. (1957) Diethyl toluamide as a mosquito repellent. East African Medical Journal 34(5):185-189.
- 05000592 Davydova, Z.V.; Rezaev, N.I.; Makolkin, I.A. (1970) Issledovanie spektrov Kombinatsionnogo rasseyaniya repellentov. [An investigation of the Raman spectra of repellents.] Vestnik Moskovskogo Universiteta Biologiya, Pochvovedenie. [Journal of Moscow University, Biology and Soil Science.] 25(1):42-48.
- 05002556 Drake, N.L.; Eaker, C.M.; Kilmer, G.W.; Melamed, S.; Shenk, W.J.; Weaver, W.E. (1945) The Preparation of Some Compounds for Testing as Insect Repellents. Washington, D.C.: U.S. Department of Commerce, Office of Technical Services. (PB Report No. 27409. issued by the Office of Scientific Research and Development, National Defense Research Committee, Division 9, as report OSRD No. 6370; National Research Council, Insect Control Committee Report No. 153)

- GS0002012 WARF Institute, Inc. 1975a. Acute Oral LD₅₀ (Male and Female), Skin and Eye Irritation, Acute Inhalation LD₅₀ of 423SD111-1: WARF No. 5030701. (Unpublished study that includes tables I-II, received July 10, 1976, under 4822-10; submitted by S.C. Johnson and Son, Inc., Racine, Wis.; CDL:233491-C)
- 00001085 WARF Institute, Inc. 1975b. Acute Oral LD₅₀ (Male and Female), Skin and Eye Irritation, Eye Irritation (Direct Spray) of 3872D102: WARF No. 4122920. (Unpublished study that includes tables I-II, received July 10, 1976, under 4822-10; submitted by S.C. Johnson & Son, Inc., Racine Wis.; CDL:233491-B). (MRID: 00001085, 17A-0022c)
- 00001101 WARF Institute, Incorporated. (1976). Summary of Toxicology Data on Insect Repellent 431SD81-2. (Unpublished study that includes WARF No. 5120760, received Mar 21, 1978 under 4822-160; CDL:233262-A) File No. 00001101 (AA0100).
- 05000047 Wong, M.H.; Yew, D.T. (1978) Dermotoxicity of the mosquito repellent related to rabbit ears. Acta Anatomica 100 (1):129-131
- GS0002013 Woodard, G. 1959. Meta Delphene: Toxicity by Repeated Oral and Dermal Administration. (Unpublished study that includes charts 1-101, tables 1-125, plates 1-201, reports with summaries on rabbits and dogs, received October 1959, under unknown administration number; prepared by Woodard Research, submitted by Hercules Powder Co., Agricultural Chemicals Division, Wilmington, Del.; CDL:133545-A)
- 00001043 Zabel, L. (1976) Acute Aerosol Inhalation Toxicity Study in Rats: TL-1480, Code No. 369-76 Intermediate 2007: IBT No. 3562-09154. (Unpublished study received Sept 27, 1976 under 1021-567; prepared by Industrial Bio-Test Laboratories, Inc., submitted by McLaughlin Gormley King Co., Minneapolis, Minn.; CDL:226115-B)
- 05007489 Zadjkoff, C.M. 1979. Toxic encephalopathy associated with use of insect repellent. J. Pediatr., 95:140-142 (MRID: 05007489, 17A-0035)
- 05004736 Zakamardin, I.A. 1969, [Toxicity of diethyltoluamide for laboratory and agricultural animals.] Scientific papers of the Kazan Veterinary Institute 105:275-280 (Translated from Russian) (MRID: 05004736, 17A-0020)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

ECOLOGICAL EFFECTS

MRID

CITATION

00001026

McCann, J.A. 1972. Fish Toxicity Laboratory Report: Toxicity of Hercules Metadelphene to rainbow trout Test No. 483. Prepared by Animal Biology Laboratory, EPA; submitted by Hercules, Inc. (Acc. No. 129780).

- 00001171 Gerberg, E.J. (1966) Field and laboratory repellency tests with 2,2,4-trimethyl-1,3-pentanediol (TMPD). Journal of Economic Entomology 59(4):872-875. (Also in unpublished study received Jun 23, 1969 under 4822-EX-8; prepared by Insect Control and Research, Inc., submitted by S.C. Johnson and Son, Inc., Racine, WI; CDL:127306-C)
- 00001065 Gerberg, E.J. (1973) Mosquito Protection Time Test With 25% Deet vs. 15% Deet. (Unpublished study received Aug 27, 1973 under 3282-38; prepared by Insect Control and Research, Inc., submitted by D-Con Co., Inc., New York, NY; CDL:016035-A)
- 00001018 Gertler, S.I.; Gouck, H.K.; Gilbert, I.H. (1962) N-alkyl toluamides in cloth as repellents for mosquitoes, ticks and chiggers. Journal of Economic Entomology 53(4):451-452. (Also in unpublished study received May 7, 1970 under 891-37; submitted by Hercules, Inc., Agricultural Chemicals, Wilmington, DE; CDL:005111-D)
- 05000170 Gilbert, I.H.; (1957) Evaluation of repellents against mosquitoes and deer flies in Oregon. Journal of Economic Entomology 50(1):46-48.
- 05000141 Gilbert, I.H.; Gouck, H.K. (1955) Evaluation of repellents against mosquitoes in Panama. Florida Entomologist XXXVIII(4):153-163.
- 00001167 Gilbert, I.H.; Gouck, H.K.; Schreck, C.E. (1970) Comparison of four cinchoninates, an oxetanone, and two standard skin repellents against Aedes aegypti and Stomoxys calcitrans. The Florida Entomologist 53(2):89-92. (Also in unpublished study received Jan 11, 1978 under 4822-10; prepared by USDA, Entomology Research Division, Agricultural Research Service, submitted by S.C. Johnson and Son, Inc., Racine, WI; CDL:233318-M)
- 05000169 Gilbert, I.H.; Gouck, H.K.; Smith, C.N. (1955) New mosquito repellents. Journal of Economic Entomology 48(6):741-743.
- 00001022 Gilbert, I.H.; Gouck, H.K.; Smith, C.N. (1957) Diethyltoluamide: New insect repellents: Part II—Clothing treatments. Soap and Chemical Specialties 33(6):95-109. (Also in unpublished study received Oct 17, 1957 under 891-12; prepared by USDA Agricultural Research Service, Entomology Research Division, submitted by Hercules, Inc., Agricultural Chemicals, Wilmington, DE; CDL:022000-B)
- 05000222 Gabel, M.L.; Spencer, T.S.; Akers, W.A. (1976) Evaporation rates and protection times of mosquito repellents. Mosquito News 36(2):141-146.
- 00001171 Gerberg, E.J. (1966) Field and laboratory repellency tests with 2,2,4-trimethyl-1,3-pentanediol (TMPD). Journal of Economic Entomology 59(4):872-875. (Also in unpublished study received Jun 23, 1969 under 4822-EX-8; prepared by Insect Control and Research, Inc., submitted by S.C. Johnson and Son, Inc., Racine, WI; CDL:127306-C).

- 00001065 Gerberg, E.J. (1973) Mosquito Protection Time Test With 25% Deet vs. 15% Deet. (Unpublished study received Aug 27, 1973 under 3282-38; prepared by Insect Control and Research, Inc., submitted by D-Con-Co., Inc., New York, NY; CDL:016035-A)..
- 00001018 Gertler, S.I.; Gouck, H.K.; Gilbert, I.H. (1962) N-alkyl toluamides in cloth as repellents for mosquitoes, ticks, and chiggers. Journal of Economic Entomology 53(4):451-452. (Also In unpublished study received May 7, 1970 under 891-33; submitted by Hercules, Inc., Agricultural Chemicals, Wilmington, DE; CDL:005111-D).
- 05000170 Gilbert, I.H. (1957) Evaluation of repellents against mosquitoes and deer flies in Oregon. Journal of Economic Entomology 50(1):46-48.
- 05000141 Gilbert, I.H.; Gouck, H.K. (1955) Evaluation of repellents against mosquitoes in Panama. Florida Entomologist XXXVIII(4):153-163.
- 00001167 Gilbert, I.H.; Gouck, H.K.; Schreck, C.E. (1970) Comparison of four cinchoninates, an oxetanone, and two standard skin repellents against Aedes aegypti and Stomoxys calcitrans. The Florida Entomologist 53(2):89-92. (Also In unpublished study received Jan 11, 1978 under 4822-10; prepared by USDA, Entomology Research Division, Agricultural Research Service, submitted by S.C. Johnson and Son, Inc., Racine, WI: (CDL:233318-M).
- 05000169 Gilbert, I.H.; Gouck, H.K.; Smith, C.N. (1955) New mosquito repellents. Journal of Economic Entomology 48(6):741-743.
- 00001022 Gilbert, I.H.; Gouck, H.K.; Smith, C.N. (1957) Diethyltoluamide: New insect repellents: Part II-Clothing treatments. Soap and Chemical Specialties 33(6):95-109. (Also In unpublished study received Oct 17, 1975 under 891-12; prepared by USDA Agricultural Research Service, Entomology Research Division, submitted by Hercules, Inc., Agricultural Chemicals, Wilmington, DE; CDL:022000-8).
- 05000237 Gilbert, I.H.; Gouck, H.K.; Smith, C.N. (1957) Diethyltoluamide: New insect repellent, Part II-Clothing treatments. Soap and Chemical Specialties 33(6):95,97,99,109.
- 00001021 Gilbert, I.H.; Gouck, H.K.; Smith, C.N. (1957) New insect repellent. Soap and Chemical Specialties 33(5):115-133. (Also In unpublished study received Oct 17, 1957 under 891-12; prepared by USDA, Agricultural Research Service, Entomology Research Division, submitted by Hercules, Inc., Agricultural Chemicals, Wilmington, DE; CDL:022000-A)
- 05000236 Gilbert, I.H.; Gouck, H.K.; Smith, C.N. (1957) New insect repellent. Soap and Chemical Specialties 33(5):115-117,129,131,133.

- 05000073 Dremova, V.P. (1967) Effektivnost' primeneniya nekotorykh perellentov dlya zashchity rabochikhneftyanikov ot napadeniya gnusa v usloviyakh Tyumenskoy Oblasti. [Efficiency of use of some repellents for protection of oil-industry workers from attacks of blood-sucking Diptera in the Tyumen region.] Meditsinskaya Parazitologiya Parazitarnye Bolezni. [Medical Parasitology and Parasite Diseases.] 36(1):60-66.
- 05000089 Dremova, V.P.; Belan, A.A.; Smirnova, S.N.; Yantsen, M.M. (1968) Repellentnaya aktivnost' nekotorykh preparatov po ottnosheniyu kleshcham Ixodes persulcatus P. Sch. i Dermacentor pictus Herm. [Effect of some repellents on Ixodes persulcatus and Dermacentor pictus Herm.] Parazitologiya. [Parasitology.] 2(5):430-432.
- 05000083 Dremova, V.P.; Gleiberman, S.Ye.; Markina, V.V.; Babenko, Z.I.; Tsetlin, V.M.; Smirnova, S.N.; Nefedova, L.F.; Zvyagintseva, repelliennykh preparatov na osnove N,N-diethylamida fenoksiuksusnoi kisloty (P-203). [Entomological and toxicological evaluation of repellent preparations based on phenoxyacetic acid diethylamide (R-203).] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 44(6):699-703.
- 05000086 Dremova V.P.; Gleiberman, S.E.; Naumov, Y.A.; Zvyagintseva, T.V.; Terekhova, A.I.; Markina, V.V.; Smirnova, S.N.; Isakova, A.P.; Kazitskaya, I.I.; Yaroslavskaya, L.A. (1977) Rezultaty izucheniya i perspektivy primeneniya repellenta N,N-diethyl-2,5-dimetilbenzamida (DEKSA). [The results of a study of the repellent N,N-diethyl-2,5-dimethylbenzamide (DEXA) and the prospects for its use.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 46(2):216-221.
- 05000081 Dremova, V.P.; Gleiberman, S.E.; Smirnova, S.N.; Voronkina, T.V.; Tsetlin, V.M.; Markina, V.V. (1974) Issledovanie los'ona "deta-1", ego repellentnoi effektivnosti i toksichnosti dlya teplokrovnykh. [Lotion "deta"-1, its repellent effectiveness, and toxicity to warm blooded animals.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 43(1):67-73.
- 05002662 Dremova, V.P.; Gleiberman, S.E.; Smirnova, S.N.; Voronkina, T.V.; Tsetlin, V.M.; Markina, V.V. (1974) Issledovanie los'ona "Deta-1", teplokrovnykh. [Study of the lotion "DETA-1", its repellent effectiveness and toxicity in warm-blooded animals.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 43(1):67-73.
- 05002846 Dremova, V.P.; Markina, V.V.; Smirnova, S.N.; Volkova, T.V. (1973) Effektivnost' primeneniya repellentov v tundre Krasnoyarskogo Kraya. [Effectiveness of application of repellents in the tundra of the Krasnoyarsk territory.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 42(1):21-27.

- 05000153 Dremova, V.P.; Markina, V.V.; Zvyagintzeva, T.V. (1976) Methodology for the primary selection of insect repellents in field conditions. *International Pest Control* 18(5):12-14.
- 05000151 Dremova, V.P.; Smirnova, S.N. (1970) Effects of repellents on hard (Ixodidae) and soft (Argasidae) ticks. *International Pest Control* 12(3):10-14.
- 05000055 Dremova, V.P.; Tsetlin, V.M.; Zhuk, E.B.; Yankovskis, E.Y.; Gorbatkova, V.V. (1969) Activity of repellents employed in aerosol form. *Journal of Hygiene, Epidemiology, Microbiology and Immunology* 13(1):64-72.
- 05000582 Dubitskii, A.M. (1966)) polozhitel'noi reaktsii Komarov; mokretsov na repellenty. [Positive reaction of mosquitoes and biting midges to repellents.] *Izvestiya Akademii Nauk Kazakhskoi SSR, Seriya Biologicheskaya*. [Proceedings of the Academy of Sciences of the Kazakh SSR, Biological Series.] 1:53-56.
- 05000197 Dukes, J.C.; Rodriguez, J.G. (1976) A bioassay for host-seeking responses of tick nymphs (Ixodidae). *Journal of the Kansas Entomological Society* 49(4):562-566.
- 00001169 Eastman Chemical Products, Incorporated (1969) Technical Data: Repellent-Grade TMPD (2,2,4-Trimethyl-1,3-Pentanediol): TDS No.X-185. (Unpublished study received Jun 23, 1969 under 48220EXOX0185; submitted by S.C. Johnson and Son, Inc., Racine, WI; CDL:127306-A).
- 05000124 Elliott, R. (1964) Studies on the kinetic response of mosquitoes to chemicals. *Bulletin of the World Health Organization* 31(5):657-667.
- 05000590 Fedder, M.L. (1960) Laboratornoe izuerie insektorepellentnykh svoistv novykh otpugivayushchikh preparatov. [Laboratory study of insect-repellent properties of new repellent preparations.] *Trudy Tsentral' nogo Nauchno-Issledovatel'skogo Dezinfektsionnogo Instituta*. [Transaction of the Central Scientific Research Institute of Disinfection.] 13:278-286.
- 05000069 Fedeer, M.L.; Tsetlin, V.M.; Grikitys, E.Ya. (1964) Opyt primeneniya diefiltoluamida v aerosol'nykh ballonakh. [Experiment on the use of diethyltoluamide in aerosol cylinders.] *Meditinskaya Parazitichlogiya i Parazitarnye Bolezni*. [Medical Parasitology and Parasitic Diseases.] 33(1):61-63.
- 05000006 Frommer, R.L.; Carestia, R.R.; Vavra, R.W., Jr. (1975) Field evaluation of deet treated mesh jacket against black flies (Simuliidae). *Journal of Medical Entomology* 12(5):558-561.
- 05000222 Gabel, M.L.; Spencer, T.S.; Akers, W.A. (1976) Evaporation rates and protection times of mosquito repellents. *Mosquito News* 36(2):141-146.

- 00001033 Gilbert, I.H.; Keegan, H.L.; Eaton, R.J.; Isarangura, V.; Sirivorasarn, P. (1965) [Testing repellents against leeches in Rangong, Thailand.] Quarterly Report of Entomological Research by the USDA on Insects of Military Importance, Quarter Ending Dec 31, 1965: 585-586. (Also In unpublished study received Aug 10, 1971 under 901-337; prepared by U.S. Agricultural Research Service, Insects Affecting Man and Animals Research Branch in cooperation with U.S. Army and Royal Thai Army Medical Department, submitted by Airosol Co., Inc., Neodesha, KS; CDL:133092-C)
- 05000187 Gilbert, I.H.; Scanlon, J.E.; Bailey, D.L. (1970) Repellents against mosquitoes in Thailand. *Journal of Economic Entomology* 63(4):1207-1209.
- 00001162 Gilbert, I.H.; Smith, N.; Gouck, H.K. (1970) Skin repellents against *Aedes aegypti*. *Journal of Economic Entomology* 63(4):1203-1204. (Also In unpublished study received Jan 11, 1978 under 4822-10; prepared by U.S. Agricultural Research Service, Entomology Research Div., submitted by S.C. Johnson and Son, Inc., Racine, WI; CDL:233318-C)
- 05002847 Gleiberman, S.E.; Dremova, V.P.; Smirnova, S.N.; Durdyev, D.; Artykov, A.; Gyshikov, S. (1970) Izuchenie toksichnosti i repellentnoi aktivnosti dietiltolylamida, sintezirovannogo naosnove neftyanogo meta-ksilola, v usloviyakh turkmenii.. Study of the toxicity and repellent activity of diethyltoluamide synthesized from petroleum meta-xylene in the Turkmen SSR. *Zdravookhranenie Turkmenistana*. [Public Health of Turkmenistan.] (6):29-31.
- 05000217 Gorham, J.R. (1974) Tests of mosquito repellents in Alaska. *Mosquito News* 34(4):409-415.
- 05000168 Gouck, H.K.; Gilbert, I.H. (1955) Field Tests with new tick repellents in 1954. *Journal of Economic Entomology* 48(5):499-550.
- 05000214 Gouck, H.K.; Godwin, D.R.; Posey, K.; Schreck, C.E.; Weidhaas, D.E. (1971) Resistance to aging and rain of repellent-treated netting used against salt-marsh mosquitoes in the field. *Mosquito News* 31(1):95-99.
- 00001032 Gouck, H.K.; Taylor, J.D., Jr.; Barnhart, C.S. (1967) Screening of repellents and rearing methods for water leeches. *Journal of Economic Entomology* 60(4):959-961. (Also In unpublished study received Aug 10, 1971 under 901-337; prepared by U.S. Agricultural Research Service, Entomology Research Div., submitted by Airosol Co., Inc., Neodesha, KS; CDL:133092-A)
- 05000281 Granett, P. (1960) Use of an animal membrane in the evaluation of repellents against the stable fly (*Stomoxys calcitrans*). *Journal of Economic Entomology* 53(3):432-435.

- 00001109 Granett, P. (1969) Entomological Performance Data: Some of the Advantages of the New "6-12" Water Spray. (Unpublished study that includes summary tables of laboratory and field insect repellent efficacy tests, received Sep 18, 1969 under 3282-46, prepared by Rutgers University, submitted by Union Carbide Corp., Tarrytown, NY; CDL:007759-A)
- 0001138 Granett, P.; Lomax, J.L. (1971) Summary of Rutgers University Efficacy Data-June, 1971: "6-12" Brand Super Insect Repellent Stick Formula No. 1481-141B Stick vs. 50% Deet (95%) Liquid Formula No. 1481-141A: Exhibit I. (Unpublished study that includes tables of efficacy tests, received 1972? under 5769-Q; prepared by Rutgers University, submitted by Union Carbide Corp., Tarrytown, NY; CDL:228420-A)
- 00001137 Granett, P.; Lomax, J. (1973) [Summarized results from field tests in Maine.] (Unpublished study received Aug 24, 1973 under 3282-56; prepared by Rutgers University, Department of Entomology, submitted by Union Carbide Corp., Tarrytown, NY; CDL:009768-A)
- 00001127 Granett, P.; Lomax, J. (1973) Summarized Results from Rutgers University: Field Tests on Insect Repellent Sprays. (Unpublished study received Aug 24, 1973 under 5769-97; prepared by Rutgers University, Department of Entomology, submitted by Union Carbide Corp., New York, NY; CDL:003305-A)
- 05000209 Grothaus, R.H.; Adams, J.F. (1972) An innovation in mosquito-borne disease protection. Military Medicine 137(5):181-184.
- 05000107 Grothaus, R.H.; Gouck, H.K.; Weidhaas, D.E.; Jackson, S.C. (1974) Wide-mesh netting, and improved method of protection against blood-feeding Diptera. American Journal of Tropical Medicine and Hygiene 23(3):533-537.
- 00001038 Grothaus, R.H.; Haskins, J.R.; Bruner, L.L. (1975) Insect Repellent Jackets: A Protective Item for Use Against Blood-Feeding Diptera. Interim Report. By U.S. Navy Department, Bureau of Medicine and Surgery. Camp LeJeune, N.C.; U.S. Naval Medical Field Research Laboratory. (Report 17, Vol. XXV, No. 7; MF51.524.009-8008.17; also In unpublished study received Apr 22, 1977 under 901-37; prepared by U.S. Naval Medical Field Research Laboratory, submitted by Aerosol Co., Inc., Neodesha, KS; CDL:229634-A)
- 05000221 Grothaus, R.H.; Haskins, J.R.; Schreck, C.E.; Gouck, H.K. (1976) Insect repellent jacket: Status, value and potential. Mosquito News 36(1):49-152.
- 05000200 Grothaus, R.H.; Hirst, J.M.; Gouck, H.K.; Weidhaas, D.E. (1972) Field tests with repellent-treated wide-mesh netting against mixed mosquito populations. Journal of Medical Entomology 9(2): 149-152.

- 05000010 Grothaus, R.H.; Reed, J.T.; Passingham, L.H. (1976) Field evaluation of arthropod repellent netting as a ground cloth against ticks. *American Journal of Tropical Medicine and Hygiene* 25(5):747-750.
- 05000207 Gualtieri, F.; Tsakotellis, P.; Skinner, W.; Johnson, H.; Skidmore, D.; Maibach, H. (1973) Topical mosquito repellents. VI. Sulforamides and quinoline-4-carboxylic acid derivatives. *Journal of Pharmaceutical Sciences* 62(5):849-851.
- 05006357 Hadani, A.; Ziv, M.; Rechav, Y. (1977) A laboratory study of tick repellents. *Entomologia Experimentalis et Applicata* 22(1):53-59.
- 05000091 Hasegawa, M.; Hattori, K. (1970) Studies on the control of the biting midges Culicoides circumscriptus on the shore of Lake Saroma, Hokkaido. Hokkaidoritsu Eisei Kenkyosho Ho. [Report of the Hokkaido Institute of Public Health.] 20:91-95.
- 00004864 Haynes, H.L. (1969) 3. Entomological Performance Data. (Unpublished study received Sept 18, 1969 under 3282-45. prepared by Union Carbide Consumer Products Co. in cooperation with Rutgers, University, submitted by D-Con Co., Inc., New York, NY; CDL:022574-A)
- 05006137 Haynes, H.L.; Lahr, P.H., inventors; Union Carbide Corp., assignee (1974) Synergistic insect repellent compositions containing N,N-diethyl meta toluamide and 2-ethyl-1,3-hexanediol. Canadian patent 960140. Dec 31. 22 p.
- 00001068 Hercules Incorporated (1977) Metadelphene: The All-Purpose Insect Repellent: Bulletin AP-105B. (Unpublished study received Jul 7, 1969 under 1516-52; submitted by Curtis Laboratories, Inc., Kansas City, MO; CDL:024967-B)
- 00001025 Hercules Powder Company (1957) Meta Delphene: The Complete Insect Repellent. Bulletin No. 213. Wilmington: Hercules Powder Company. (Also in unpublished study received Aug 13, 1957 under ?; submitted by Hercules Powder Co., Wilmington, DE; CDL:108471-A)
- 00001105 Hercules Powder Company (1961) Metadelphene: The All-Purpose Insect Repellent. Wilmington, DE: Hercules Powder Company. (Number NS-292; Also in unpublished submission received Mar 21, 1978 under 4822-160; submitted by S.C. Johnson and Son, Inc., Racine, WI; CDL:233;263-H)
- 05008237 Hill, J.A.; Robinson, P.B.; McVey, D.I.; Akers, W.A.; Reifenrath, W.G. (1979) Evaluation of mosquito repellents on the hairless dog. *Mosquito News* 39(2):307-310.
- 05002327 Husbands, R.C.; Lewallen, L.L. (1976) Laboratory and field test on deet repellency to mosquitoes as a means of protecting squabs from fowl pox. Pages 100-102, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association, 44th.

- 05000064 Ivanova, L.V.; Stavrovskaya, V.I. (1960) Predvaritel'nye dannye polevogo ispitaniya dietiltoluamidov. [Field trials of diethyltoluamides.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 29:564-570.
- 05002306 Ivanova, L.V.; Stavrovskaya, V.I. (1960) Predvaritel'nye dannye polevogo ispytaniya novogo repellenta piperidilamida m-toluilovoy kisloty. [Preliminary results of field experiments with a new repellent, N-(m-toluoyl)piperidine.] Meditsinskaya Parazitologiya i Parazitarnye Bolenzi. [Medical Parasitology and Parasite Diseases.] 35(3):320-323.
- 0500015 Johnson, H.; DeGraw, J.; Engstrom, J. (1975) Topical mosquito repellents VII: Akyl triethylene glycol monoethers. Journal of Pharmaceutical Sciences 64(4):693-695.
- 05000198 Johnson, H.L.; Skinner, W.A.; Skidmore, D.; Maibach, H.I. (1968) Topical mosquito repellents. II. Repellent potency and duration off ring-substituted N,N-dialkyl-and amino-alkylbenzamides. Journal of Medical Chemistry 11:1265-1268.
- 05002555 Kappor, D.; Paul, C.F.; Perti, S.L. (1972) Evaluation of certain insecticides and repellents against ticks. Defence Science Journal 22(3):185-190.
- 05002665 Kashafutdinov, G.A. (1969) Znachenie pola nasekomykh i rastvoritelei pri sravnitel'noi otsenke repellentov. [Significance of the sex of insects and the effect of solvents during a comparative evaluation of repellents.] Uchenye Zapiski Kazanskogo Veterinarnogo Instituta. [Scientific papers of the Kazan Veterinary Institute.] 105:267-269.
- 05000077 Kashafutdinov, G.A.; Il'ina, N.A. (1971) Repellent activity of cyclohexenyl ethers of amidophosphoric acid. Meditsinskaya Parazitologiya Parazitarnye Bolezni 40(1):36-39.
- 05001824 Kazhdan, V.B.; Naumov, Y.A.; Dunaeva, I.D.; Moskvicheva, M.V. (1972) Synthesis and repellent properties of N-alkylanilides and esters of butanesulfonic acid. Khimiko-Farmatsevticheskii Zhurnal. [Chemical-Pharmaceutical Journal.] 6(2):6-10.
- 5002309 Keegan, H.L.; Weaver, R.E. (1964) Studies of Taiwan leeches. II: Field tests of effectiveness of insect repellents against aquatic leeches at Cha'o Chow, Pingtung, Taiwan. Bulletin of the Institute of the Zoological Academy of Sinica 3(2):83-92.
- 00001034 Keegan, H.L.; Weaver, R.E. (1964) Studies of Taiwan Leeches: 2: Field Tests of Effectiveness of Insect Repellents Against Aquatic Leeches at Cha'o Chow, Pingtung, Taiwan. 406th Medical Laboratory Research Report. (Also In unpublished study received Jul 15, 1974 under 901-37; prepared by U.S. Army Medical Command, Department of Entomology, 406th Medical Laboratory, submitted by Aerosol Co., Inc., Neodesha, KS; CIL:221875-A)

- 00001035 Keegan, H.L.; Weaver, R.E.; Fleshman, P.; Zarem, M. (1964) Studies of Taiwan Leeches. 3. Further Tests of Repellents Against Aquatic, Blood-sucking Leeches. 406th Medical Laboratory Research Report. (Also In unpublished study received ? under 901-37; prepared by U.S. Army Medical Command, Department of Entomology, 406th Medical Laboratory, submitted by Aerosol Co., Inc., Neodesha, KS; CDL:221876-A)
- 05002310 Keegan, H.L.; Weaver, R.E.; Fleshman, P.; Zarem, M. (1964) Studies of Taiwan leeches. III. Further tests of repellents against aquatic, blood-sucking leeches. Bulletin of the Institute of the Zoological Academy of Sinica 3(2):93-106.
- 0500127 Kellogg, F.E.; Burton, D.J.; Wright, R.H. (1968) Measuring mosquito repellency. Canadian Entomologist 100(7):763-768.
- 00001163 Kahn, A.A.; Maibach, H.I. (1972) A study of insect repellents: 1: Effect on the flight and approach by Aedes aegypti. Journal of Economic Entomology 65(5):1318-1321. (Also In unpublished study received Jan 11, 1978 under 4822-10; prepared by University of California School of Medicine, Department of Dermatology, submitted by S.C. Johnson and Son, Inc., Racine, WI; CDL:233318-D)
- 05000193 Kahn, A.A.; Maibach, H.I.; Skidmore, D.L. (1973) Insect repellents. 2. Effect of temperature on protection time. Journal of Economic Entomology 6(2):437-438.
- 05000195 Kahn, A.A.; Maibach, H.I.; Skidmore, D.L. (1975) Insect repellent: Effect of mosquito and repellent-related factors on protection time. Journal of Economic Entomology 68(1):43-45.
- 05000226 Kahn, A.A.; Maibach, H.I.; Skidmore, D.L. (1977) Increased abrasion and wash resistance of repellents with addition of polymers. Mosquito News 37(1):123-126.
- 05000084 Kharitonova, S.I. (1975) O prolongiruyushchem deistvii fiksatorov parfyumerii i etilstellyulosey v zhidkikh repellentnykh sostavakh. [Prolonging effect of perfume fixatives and ethylcellulose in liquid repellent compositions.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 44(6):704-707.
- 05000299 King, W.V., (1954) Chemicals Evaluated as Insecticides and Repellents at Orlando, FL. By U.S. Department of Agriculture, Agricultural Research Service, Entomology Research Branch. Washington, D.C.: U.S. Government Printing Office. (Agricultural Handbook No. 69)
- 05000150 Kochhar, R.K.; Dixit, R.S.; Somaya, C.I. (1974) A critical analysis of deet as a repellent against arthropods of public health importance and water leeches. Indian Journal of Medical Research 62(1):125-133.

- 05000075 Koshkina, I.V.; Kharitonova, S.I. (1970) Ethyl cellulose as an agent prolonging the action of repellents, part 2. Meditsinskaya Parazitologiya Parazitarnye Bolezni 39(2):224-227.
- 05002302 Koshkina, I.V.; Kharitonova, S.I. (1976) Vliyanie fiksatorov i polimerov na dlitel'nost' otpugivayushchego deistviya repellentov na tkani. [The effect of fixatives and polymers on the duration of the protective action on fabric. Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 45(4):446-449.
- 05001823 Krasovskaya, S.B.; Virnik, A.D.; Dremova, V.P.; Rogovin, Z.A. (1972) Izvestiya Vysshikh Uchebnykh Zavedenii, Tekhnologiya Tekstil'noi Promyshlennosti. Technology of the Textile Industry USSR. 72(1):84-88. [Insect-repelling fibrous materials.] [Technology of the Textile Industry USSR.] 72(1)-:84-88.
- 05000229 Kuraishy, N.A.; Talibi, S.A.; Debnath, R.K. (1962) Report on a mosquito repellent cream Flypel with metatoluamide as active ingredient. Pakistan Journal of Health XII(1):22-26.
- 00004869 Lahr, P.H. (1970) Letter sent to H.L. Haynes dated June 10, 1970. [Additional data concerning the synergism of mixtures of ethylhexanediol and diethyltoluamide.] (Unpublished study received Jul 1, 1970 under 3282-48; prepared by Union Carbide Corp., submitted by D-Con Co., Inc., New York, NY; CDL:007765-C)
- 00004868 Lahr, P.H. (1970) Letter sent to H.L. Haynes dated Jun 8, 1970. [Summary of data concerning the efficacy and synergism of mixtures of ethylhexanediol and diethyltoluamide. Includes appendix with tables from the Rutgers Report-May, 1970.] (In unpublished study received Jul 1, 1970 under 3283-48; prepared by Rutgers University, submitted by D-Con Co., Inc., New York, NY; CDL:007765-B)
- 00004870 Lahr, P.H. (1970) Letter sent to H.L. Haynes dated Jun 10, 1970. [Summary of data to show graphically the improvement due the admixture with diethyltoluamide. Includes appendix with tables from the Rutgers Report done by J.L. Lomax.] (In unpublished study received Jul 1, 1970 under 3282-48; prepared by Rutgers University, submitted by D-Con Co., Inc., New York, NY; CDL:007765-D)
- 00001108 Lahr, P.H. (1970) Synergistic Mixtures of 2-Ethyl-1,3-hexanediol and M-Diethyltoluamide. Tarrytown Technical Memorandum TTM-81. (Unpublished study that includes summary tables and graphs from data done by J.L. Lomax, received Mar 24, 1970 under 5769-27; prepared by Rutgers University, submitted by Union Carbide Corp., New York, NY; CDL:007758-D)

- 00001124 Lahr, P.H. (1972) Performance of "6-12 Plus" Spray Formula vs. 15 Percent Deet Based on Rutgers Laboratory an Field Test for 1968 through 1971. Tarrytown Technical Memorandum TTM-102. (Unpublished study that includes report by J.L.Lomax entitled "Rutgers Field Testing of Insect Repellents from 1968 to 1972," received Jun 27, 1972 under 3282-48; prepared by Rutgers University, submitted by Union Carbide Corp., Tarrytown, NY; CDL:007768-A)
- 05002319 Langford, G.S.; Joseph, S.R.; Berry, R.. (1966) Some observations on mosquito repellents. Mosquito News 26(3):399-404.
- 05004984 Lindsay, I.S.; McAndless, J.M. (1978) Permethrin-treated jackets versus repellent-treated jackets and hoods for personal protection against black flies and mosquitoes. Mosquito News 38(3):350-356.
- 00001110 Lomax, J.L. (1968) Annual Report for 1968. [Report concerns the efficacy of pest repellents in laboratory and field tests.] (Unpublished study received Sep 18, 1969 under 3282-46; prepared by Union Carbide Consumer Products Co. in cooperation with Rutgers University, Department of Entomology and Economic Zoology, submitted by Union Carbide Corp., Tarrytown, NY; CDL:007759-B)
- 00004865 Lomax, J.L. (1968) Annual Report for 1968. [Report concerns the efficacy of pest repellents in laboratory and field tests.] (Unpublished study received Sep 18, 1969 under 3282-45; prepared by Union Carbide Consumer Products Co. in cooperation with Rutgers University, Department of Entomology and Economic Zoology, submitted by D-Con Co., Inc., New York, NY; CDL:022574-B)
- 00001118 Lomax, J.L. (1969) Summary and Discussion of All Duration Field Test Results for Summer 1969. (Unpublished study received Apr 30, 1970 under 4769-48; prepared by Rutgers University, College of Agriculture, Department of Entomology submitted by Union Carbide Corp., New York, NY; CDL:007762-F)
- 00001115 Lomax, J.L. (1970) Summary and Discussion of Repellent Test Results: Duration Type in the Laboratory. (Unpublished study received May 14, 1970 under 5769-44; prepared by Rutgers University, College of Agriculture, Department of Entomology, submitted by Union Carbide Corp., New York, NY; CDL:007761-E)
- 00001117 Lomax, J.L. (1970) Summary and Discussion of Repellent Test Results: Duration Type in the Laboratory. (Unpublished study received Apr 30, 1970 under 5769-48; prepared by Rutgers University, College of Agriculture, Department of Entomology, submitted by Union Carbide Corp., New York, NY; CDL:007762-E)

- 00001084 Mace, E.F. (1966) [Efficacy tests on Off! and 2370D52-2.] (Unpublished study received Oct 8, 1966 under 4822-10; prepared by S.C. Johnson and Son, Inc., Research and Development Division, Product Research Department, submitted by S.C. Johnson and Son, Inc., Racine, WI; CDL:009371-A)
- 05000116 Maibach, H.I.; Kahn, A.A.; Akers, W. (1974) Use of Insect repellents for maximum efficacy. Archives of Dermatology 109(1): 32-35.
- 05000070 Mal'tsev, M.I.; Katkov, V.M.; Achilov, R. (1964) Results of tests of some repellents under natural conditions of Turkmenia. Meditsinskaya Parazitologiya i Parazitarnye Bolezni 33(5):613-614.
- 05000065 Mandel'baum, Y.A.; Saf'yanova, V.M. (1960) Dietilamid metatoluilovci kisloty-effektivnyi repellent proti krovososushchikh masekomykh i kleshchei. [Diethylamide of metatoluyil acid-an effective repellent against blood-sucking insects and ticks.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 29(5):570-575.
- 05000583 Markina, V.V.; Dremova, V.P. (1975) Prolongation of the protective action of repellents applied to fabric. Khimiko Farmatsevticheskii matsevticheskii Zhurnal. [Chemical-Pharmaceutical Journal.] 9(1):24-27.
- 05003370 Markina, V.V.; Dremova, V.P.; Kamennov, N.A. (1970) Nekotoryedannye o dinamike udaleniya DETA s obrabotannoi poverkhnosti. [Dynamics of removal of diethyltoluamide from treated surface.] Masloboino-Zhirovaya Promyshlennost. [Oil and Fat Industry.] 36(2):30-32.
- 05000062 Markina, V.V.; Dremova, V.P.; Vashkov, V.I.; Kamennov, N.A. (1971) Dinamika ispareniya i vsasyvaemosti los'onov i kremov na osnove repellentov deta, benzimina i karboksida. [Dynamics of evaporation and absorption of repellent lotions and creams based on DETA, benzimine and carboxide.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 40(6):690-695.
- 00001037 McAndless, J.M. (1974) Personal Protection Against Biting Flies: The Relative Effectiveness of Topical Repellent and Repellent-treated Overjackets, Collars and Wristbands. By the Department of National Defense Canada, Research and Development Branch. Ottawa, Canada: Defense Research Establishment Ottawa. (Technical Note No. 74-28; Project No. 79-03-05; Also In unpublished study received Apr 27, 1977 under 901-37; prepared by NBC Defense Div., Materials Section, submitted by Aerosol Co., Inc., Neodesha, KS; CDL:229633-A)

- 05000024 McAndless, J.M.; Lindsay, I.S. (1977) Pyrethroid-treated jackets versus repellent-treated jackets and hoods for personal protection against biting flies. Ottawa, Ontario, Canada: Defense Research Establishment Ottawa. (Available from: NTIS, Springfield, VA; AD-A047 324/9ST)
- 05000202 McDonald, J.L.; Grothaus, R.H. (1973) Field studies using wide mesh mosquito bed nets in Taiwan and Indonesia. *Journal of Medical Entomology* 10(3):299.
- 05000194 McGovern, T.P.; Gouck, H.R.; Burden, G.S.; Sarmiento, R.; Beroza, M.; Schmidt, C.H. (1974) N,N-substituted n-alkanesulfonamides as repellents for the yellowfever mosquito and the German cockroach. *Journal of Economic Entomology* 67(1):71-73.
- 00001055 McLaughlin Gormley King Company (1957) Recommendations for the Use of Diethyltoluamide by the Armed Forces. (Unpublished study received Oct 22, 1974 under 1021-1323; submitted by ?; CDL:028339-I)
- 00001053 McLaughlin Gormley King Company (1974) Personal Repellency, Aedes aegypti. MGK File No. D-1030-74. (Unpublished study received Oct 22, 1974 under 1021-1323; prepared by MGK Research and Development Department, Biological Testing Laboratory, submitted by ?; CDL:028339-D)
- 00001000 Miller, W.V. (1955) Insect Repellents for Canines. (Unpublished study received Mar 2, 1956 under 59-82; submitted by Burroughs Wellcome Co., Research Triangle Park, NC; CDL:000126-A)
- 05001270 Mironov, V.P.; Krashkevich, K.V.; Krivtsova, Y.N.; Kul'kova, T.A.; Roginskaya, Y.Y. (1961) Laboratory investigations of the effect of certain repellents on mature Dermacentor pictus Herm ticks. *Vestnik Moskovskogo Universiteta*. [Moscow University Herald.] XVI (1):26-31.
- 00001129 Mitchell, R.B. (1971) "6-12 Plus" Insect Repellent Spray with Knockdown Power: Efficacy Data. (Unpublished study that includes Tarrytown Technical Memorandum TTM-95 and tables, received Oct 13, 1971 under 5769;86; prepared by Union Carbide Corp. in cooperation with Crop Protection Institute; submitted by Union Carbide Corp., Tarrytown, NY; CDL:007772-A)
- 05002307 Moore, J.B. (1962) Insect repellent formulations for personal use. *Aerosol Age* 7(12):43,46,123,128.
- 05000220 Mulrennan, J.A.; Lewis, L.A.; Grothaus, R.H. (1975) Field test with repellent treated wide-mesh netted jackets against the valley Black Gnat, Leptoconops carteri. *Mosquito News* 35(2):228-229.

- 00001072 Neumiller, P.J. (1972) [Efficacy of Deet in Combination with Other Repellents.] (Project 573; unpublished study that includes discussion by E.F. Mace and data summaries with accompanying tables by P.J. Neumiller, received Jun 12, 1972 under 4822-119; prepared by S.C. Johnson and Son, Inc., submitted by S.C. Johnson and Son, Inc., Racine, WI; CDL:007514-A)
- 05000290 Novak, D. (1973) Preliminary laboratory tests with tick repellents. Pages 685-686, In Proceedings of the International Congress of Acarology, 3rd; Prague; 1971.
- 05000060 Pervomaiskii, G.S.; Shustrov, A.K. (1969) Repellentnoe deystvie dets na komarov i kleshchey. [Repellent effect of dets on mosquitoes and ticks.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 38(1):52-55.
- 05000235 Pierce, H.F. (1958) Diethyltoluamide in aerosol repellents. Soap and Chemical Specialties 34(6):80-81,83-85-86, 191.
- 00001104 Pierce, H.F. (1958) Diethyltoluamide in aerosol repellents. Soap and Chemical Specialties Jun: 43-48. Also In unpublished study received Mar 21, 1978 under 4822-160; prepared by Hercules Powder Co.; submitted by S.C. Johnson and Son, Inc., Racine, WI; CDL:233263-F)
- 05000269 Pierce H.F.(1958) The use of diethyltoluamide insect repellent in aerosols. Page 47-50, In Proceedings of the Mid-Year Meeting, Chemical Specialties Manufacturers Association, No. 44, Dec 9-12, 1957. New York NY: Chemical Specialties Manufacturers Association.
- 05002659 Polyakov, V.A. (1977) Zashchita severnykh oleney ot gnusa. [Protection of reindeer from biting flies.] Veterinariya (Moscow). [Veterinary Science.] 7:37-38.
- 05000026 Potapov, A.A.; Tsizin, Y.S.; Kharitonova, S.I.; Vladimirova, V.V.; Sazonova, E.V. (1977) Repellentsalfetki dlya zashchity ot gnusa. [Repellent napkins for protection against blood-sucking Diptera.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 4:434-439.
- 05000079 Potapov, A.A.; Vorobieva, Z.G.; Vladimirova, V.V.; Kharitonova, S.I.; Koshkina, I.V. (1972) N,N-diethylamide of benzoic acid (R-2) as a repellent for protection against biting Diptera. Meditsinskaya Parazitologiya Parazitarnye Bolezni 41(2):185-189.
- 05000027 Potapov, A.A.; Zharova, A.N.; Vladimirova, V.V.; Kornienko, N.I. [Resistance of Repellents to washing off with water after application to fabric and skin (laboratory trials).] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 47(1):81-85.

- 05000190 Quintana, R.P.; Mui, P.T.; Fisher, R.G.; Schreck, C.; Gouck, H.K. (1972) Dihydroxyacetone monoester mosquito repellents. Effect of branching, cyclization, and unsaturation upon repellent efficacy. *Journal of Economic Entomology* 65(1):66-69.
- 05000189 Ramachandran, P.K.; Koshy, T.; Sastry, K.G.K.; Singh, S.P.; Srinivasan, M.N.; Ganguly, S.K. (1971) Leech repellents. *Journal of Economic Entomology* 64(5):1293-1294.
- 05000196 Rietschel, R.L.; Spencer, T.S. (1975) Correlation between mosquito repellent protection time and insensible water loss from the skin. *Journal of Investigative Dermatology* 65(4):385-387.
- 05002333 Rutledge, L.C.; Moussa, M.A.; Lowe, C.A.; Sofield, R.K. (1978) Comparative sensitivity of mosquito species and strains to the repellent diethyltoluamide. *Journal of Medical Entomology* 14(5):536-541.
- 05003639 Saxena, B.N.; Khalsa, H.G.; Pillai, K.R.M. (1969) Evaluation of repellents against land leeches--Part II. Defence Science Journal 19(2):93-96.
- 00001168 S.C. Johnson and Son, Incorporated (1964) [Repellency Tests (Duration, Abrasion and Water Resistance) Off! Foam] (Unpublished study that includes bioassay reports with accompanying tables, received Mar 10, 1964 under 4822-51; prepared by S.C. Johnson and Son, Inc., submitted by ?; CDL:009364-A)
- 00001170 S.C. Johnson and Son, Incorporated (1969) [Repellency Tests with Deet as Single Active and in Combination with Other active Ingredients.] (Unpublished study that includes bioassay reports with accompanying discussion and tables, received Jun 23, 1969 under 4822-EXP-8G; prepared by S.C. Johnson and Son, Inc., submitted by ?; CDL:127306-B)
- 00001095 S.C. Johnson and Son, Incorporated (1974) Fly and Mosquito Repellent 3936D84-2. (Unpublished study received Aug 7, 1974 under 4822-137; prepared by S.C. Johnson and Son, Inc., Product Research Department, submitted by ?; CDL:022152-A)
- 00001097 S.C. Johnson and Son, Incorporated (1974) Fly and Mosquito Repellent 4071D9-3. Project 752. (Unpublished study received Aug 7, 1974 under 4822-138; prepared by S.C. Johnson and Son, Inc., submitted by ?; CDL:102574-A)
- 00001103 S.C. Johnson and Son, Incorporated (1977) Efficacy Data Summary. (Unpublished study that includes summaries of laboratory and field data with accompanying tables, received Mar 21, 1978 under 4822-106; prepared by S.C. Johnson and Son, Inc., submitted by ?; CDL:233263-A)

- 05000613; Schiefer, B.A.; Carestia, R.R.; Vavra, R.W., Jr.; Frommer, R.I. (1977) Development of Ground and Aerial Adult Control Measures for Biting Diptera without Using Persistent Pesticides. By U.S. Army Environmental Sciences Branch, Regional Division North, Radiation and Entomological Sciences Branch, Fort George G. Meade, MD. Aberdeen Proving Ground, MD.: U.S. Army Environmental Hygiene Agency. (Report. No: 61-0614-77 Contract No. 3A06 1102-B-7P01)
- 05000223 Schiefer, B.A.; Vavra, R.W., Jr.; Frommer, R.I.; Gerber, E.J. (1976) Field evaluation of several repellents against black flies (Diptera, Simuliidae). Mosquito News 36(3):242-247.
- 00001052 Schley, G.; Preiss, F.; Mitchell, K.; Bollenback, W. (1974) [Mosquito Repellency. MGK File No. D-1036-74.] (Unpublished study received Oct 22, 1974 under 1021-1323; prepared by McLaughlin Gormley King Co., submitted by McLaughlin Gormley King Co., Minneapolis, MN; CDL:028339-C)
- 05000009 Schmidt, M.L. (1977) Relative effectiveness of repellents against Simulium damnosum (Diptera: Simuliidae) and Glossina morsitans (Diptera: Glossinidae) in Ethiopia. Journal of Medical Entomology 14(3):276-278.
- 05000199 Schmidt, M.L.; Schmidt, J.R. (1969) Relative effectiveness of chemical repellents against Phlebotomus papatasi (Scopoli). Journal of Medical entomology 6 (1):79-80.
- 005000113 Schreck, C.E. (1977) Techniques for the evaluation of insect repellents: A critical review. Annual Review of Entomology 22:101-119.
- 05000188 Schreck, C.E.; Gilbert, I.H.; Weidhaas, D.E.; Posey, K.H. (1970) Spatial action of mosquito repellents. Journal of Economic Entomology 63(5):1576-1578.
- 05000033 Schreck, C.E.; McGovern, T.P.; Smith, N. (1978) Repellency of selected esters and amides of four alicyclic acids against the stable fly, Stomoxys calcitrans (Diptera: Muscidae). Journal of Medical Entomology 14(5):589-591.
- 05000030 Schreck, C.E.; Posey, K; Smith D. (1977) Repellent Activity of Compounds Submitted by Walter Reed Army Institute of Research. Part 1. Protection Time and Minimum Effective Dosage Against Aedes aegypti Mosquitoes. By U.S. Department of Agriculture, Agricultural Research Service in cooperation with Walter Reed Army Institute of Research. Washington, DC: United States Department of Agriculture. (USDA, ARS Technical Bulletin No. 1549).
- 05000031 Schreck, C.E.; Smith, N. (1977) Repellency of cyclohexamethylene carbamide and N,N-diethyl-m-toluamide against Aedes aegypti and Ae. taeniorhynchus. Mosquito News 37(3):525-5527.

- 05002304 Schreck, C.E.; Smith, N.; Gouck, H.K. (1976) Repellency of N,N-diethyl-M-Toluamide (deet) and 2 hydroxyethyl cyclohexanecarboxylate against the deer flies Chrysops atlanticus Pechuman and Chrysops flavidus Wiedemann. Journal of Medical Entomology 13(1):115-118.
- 0500096 Semenov, P.V.; Gomoionova, N.P.; Korolev, A.P. (1968) Ispytaniya dietiltoluidina (deta) dlya zashchity zhivotnykh ot slepnei. [A test of diethyltoluamide (deta) in protecting animals from the gadfly.] Veterinaria. [Veterinary Science.] 45(6):87-89.
- 05000292 Shimmin, R.; Bayles, S.; Spencer, T.; Akers, W.; Grothaus, R. (1974) Four-site method for mosquito repellent field trials Pages 24-27, In Proceedings and Papers of the Annual Conference of the California Mosquito Control Association, Inc. (42nd) and the Annual Meeting of the American Mosquito Control Association Inc.(30th). Visalia, Calif.: California Mosquito Control Association.
- 05000224 Sholdt, L.L.; Holloway, M.L.; Grothaus, R.H.; Schreck, C.E. (1976) Dwelling space repellents: Effect upon behavioral responses of mosquitoes in Ethiopia. Mosquito News 36(3):327-331.
- 05000036 Sholdt, L.L.; Holloway, M.L.; Chandler, J.A.; Fontaine, R.E.; Elsen, A.V. (1977) Dwelling space repellents: Their use on military tentage against mosquitoes in Kenya, East Africa. Journal of Medical Entomology 14(2):252-253.
- 05000039 Skinner, W.A.; Crawford, H.T.; Skidmore, D.; Maibach, H.I. (1977) Topical mosquito repellents X: 2-oxazolidones. Journal of Pharmaceutical Sciences 66(4):587-589.
- 05000038 Skinner, W.A.; Crawford, H.T.; Tong, H.; Skidmore, D.; Maibach, H.I. (1976) Topical mosquito repellents IX: Quinolines, isoquinolines, and quinoxalines. Journal of Pharmaceutical Sciences 65(9):1404-1407.
- 05000054 Smatov, Z.S.; Kozlova, R.G. (1976) Repellentnaya aktivnost antikomarina v usloviakh Kazakhstana. [The repellent activity of Antikomarin in the conditions of Kazakhstan.] Izvestiya Akademii Nauk Kazakhskoi SSR Seriya Biologicheskaya. [Proceeding of the Academy of Sciences of the Kazakh SSR, Biological Series. 3:32-35.
- 05000090 Smirnova, S.N.; Dremova, V.P. (1971) Chuvstvitel'nost' k repellentam u iksodovykh kleshchei rodov Rhipicephalus i Hyalomma. [Sensitivity of the ixodid tick genera Rhipicephalus and Hyalomma to repellents.] Parazitologiya. Parasitology. 5(4):357-360.

- 05002661 Smirnova, S.N.; Dremova, V.P. (1971) *Izmenenie uroviya Chuvstvitel'nosti k repellentam u iksodovykh kleschei v protsesse metamorfoza.* [Change of susceptibility level to repellents in hard ticks during metamorphosis. Pages 145-149, In] *Khemoretseptsiya Nasekomykh Materialy k Vsesoyuznomu Simpoziumu po Khemoretseptsii Nasekomykh*, 1st; Sep 8-10, Vilnyus, USSR. [Insect Chemoreception, Materials for All-Union Symposium on Insect Chemoreception, 1st. Edited by A. Skirkevicius.] Vilnyus, USSR
- 05000076 Smirnova, S.N.; Dremova, V.P. (1971) The sensitivity to repellents of *Alectorobius tholozani papillipes* (Birula) at different stages of development. *Meditinskaya Parazitologiya Parazitarnye Bolezni* 40(1):45-49.
- 00001113 Smith, C.N. (1957) Insect Repellents. Pages 1-8, In 44th Annual Meeting Proceedings of the Chemical Specialties Manufacturing Association, Inc.; Dec 1957, Hollywood, FL. New York, NY: The Chemical Specialties Manufacturers Association, Inc. (Also In unpublished submission received Feb 26, 1970 under 3282-446; prepared by USDA Agricultural Research Service, Entomology Research Division, submitted by Union Carbide Corp., New York, NY; CDL:007760-C).
- 05000270 Smith, C.N. (1958) Insect repellents. Pages 96-104, In Proceedings of the Annual Meeting, Chemical Specialties Manufacturers Association, No. 44; Dec 9-12, 1957. New York, NY: Chemical Specialties Manufacturers Association.
- 05000276 Smith, C.N. (1962) Factors affecting the protection obtained with insect repellents. Pages 482-485, In Transactions of The International Congress of Entomology, 11th Congress, vol. 2, sect. 7/14.
- 05000052 Sova, C.R. (1965) Initial studies on mosquito repellents. *Farmakoterap Zpravy* 11(1):89-92.
- 05000041 Spencer, T.S.; Akers, W.A. (1976) Field trials of mosquito repellents in Florida, Alaska, and California. Pages 102-105, in Proceedings and papers of the Forty-fourth Annual Conference of the California Mosquito Control Association Inc.; Jan 25-28, 1976. Visalia, Calif.: California Mosquito Control Association. (Available from: NTIS, Springfield, VA; AD-A051 081).
- 05000042 Spencer, T.S.; Bayles, S.F.; Shimmin, R.K.; Gabel, M.I.; Akers, W.A. (1975) Interactions between mosquito repellents and human skin. Pages 1-14, In Proceedings of the Ninth Army Science Conference; June 1975; West Point, NY (Available from NTIS, Springfield, VA; AD-A050-958/8ST).

- 05000043 Spencer, T.S.; Hill, J.A.; Akers, W.A.; Bjorkland, G. (1977) Studies of repellent formulations with N,N-diethyl-m-toluamide. Pages 121-123, In Proceedings and Papers of the Forty-fifth Annual Conference of the California Mosquito Vector Control Association, Inc.; Feb 13-16, 1977. Visalia, Calif.: California Mosquito and Vector Control Association.
- 05000040 Spencer, T.S.; Zeller, K.L.; Brodel, C.F.; Akers, W.A. (1977) Analysis of four-site method for testing mosquito repellents. Pages 119-120, In Proceedings and Papers of the Forty-Sixth Annual Conference of the California Mosquito Vector Control Association, Inc.; Feb 13-16, 1977. Visalia, Calif.; California Mosquito and Vector Control Association.
- 05000119 Varma, R.N. (1971) Evaluation of repellents against mosquitoes. Armed Forces Medical Journal, India 27(2):164-168.
- 05000067 Violovich, N.A. (1960) Repellent Properties of diethylamide of metatoluic acid in regard to horseflies. Meditsinskaya Parazitologiya i Parazitarnye Bolezni 32(5):539-542.
- 05000085 Vladimirova, V.V.; Potapov, A.A (1977) Rezultaty ispytaniya golovnykh nakidok iz raznykh tkanei, propitannykh repellentami, dlya zashchity ot komorov i moshek. [The results of tests of different materials, impregnated with repellents, for protection against mosquitoes and simuliids.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 46(1):60-63.
- 05000233 Whittemore, F.W., Jr.; Keegan, H.L.; Hedeon, R.A.; Fowler, H.W., Jr. (1961) Relative effectiveness of diethyltoluamide and M-2020 against Aedes scapularis (Rondani). Public Health Reports 76(2):179-182.
- 05000275 Wilson, H.G.; LaBrecque, G.C.; Gahan, J.B. (1961) Laboratory tests of selected house fly repellents. Florida Entomologist 44(3):123-124.
- 05006242 Yanovich, G.I. (1962) Dietiltoluamid—effektivnoe sredstvo protiv gnusa. Diethyltoluamide—effective agent against gnats. Zhivotnovodstvo. Animal Husbandry. (7):84-85.
- 05000058 Yanovich, G.I. (1967) Primenenie dietiltoluamid dlya zashchity zhivotnykh ot komarov i slepney. [Use of diethyltoluamide to protect animals from mosquitoes and horseflies.] Khimiya v Sel'skom Khozyaistve. [Chemistry in Agriculture.] 5(9):691-692.
- 05000291 Yeoman, G.H.; Lindberg, U.H. (1971) Insect repellency of certain alkyl-substituted cinchoninic acid esters. Pages 293-295, In Proceedings of the International Congress of Entomology, 13th, vol. 2. Vet. Res. Stn., Wheathampstead, Engl.
- 05000268 Yeoman, G.H.; Warren, B.C. (1970) Repellents for Stomoxys calcitrans, the stable fly: techniques and a comparative laboratory assessment of butyl methylcinchoninate. Bulletin of Entomological Research 59(4):563-577.

- 05005681 Zakamyrdin, I.A. (1973) K voprosu razrabotki khimicheskogo metoda zashchity sel'skokhoziaystvennykh zhivotnykh ot gnusa. [Development of a chemical method of protecting farm animals from blood-sucking flies.] Uchenye Zapiski Kazanskogo Veterinarnogo Instituta. [Scientific Papers of the Kazan Veterinary Institute.] 115:244-249.
- 05004980 Zaugg, J.L. (1978) Field tests with deet-treated wide-mesh netting against mixed hematophagous fly populations. Mosquito News 38(4):559-562.
- 05000071 Zhukova, L.I. (1965) Productional trials for some repellents against gnats and Ixodes persulcatus ticks. Meditsinskaya Parazitologiya i Parazitarnye Bolezni 34(5):540-543.
- 05004737 Zolotarev, E.K.; Kalakutskaya, T.V. (1960) Issledovanie repellentov. IX. Diethyltoluamidy. Sravnitel'noe izuchenie repellentnosti O-, M-, i P-izomerov dlya kleshei i komarov. [Study of repellents. IX. Diethyltoluamides. A comparative study of the repellency of O-, M-, and P-isomers for ticks and mosquitoes.] Vestnik Moskovskogo Universiteta, Biologiya, Pochvovedenie. [Journal of Moscow University, Biology and Soil Science.] 15(3):18-21.
- 05004738 Zolotarev, E.K.; Stavrovskaya, V.I. (1960) Issledovanie repellentov. X. Diethyltoluamidy. Sravnitel'noe izuchenie repellentnosti orto-, meta-, i paraizomerov dlya blokh. [An investigation of repellents: X. Diethyltoluamides. A comparative study of the repellent effect of ortho-, meta-, and para-isomers on fleas.] Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 29(5):559-563.

Section III

OFFICE OF PESTICIDE PROGRAMS REGISTRATION STANDARD BIBLIOGRAPHY

Examined and Judged to be Citations Inappropriate For Use in
Developing the Standard

CASE	GS0002	N,N-Diethyl-meta-toluamide and other isomer	9/8/80	Page
MRID	CITATION			
05002316	Acree, F., Jr.; Beroza, M. (1972) Gas chromatography of insect repellents. <i>Journal of Economic Entomology</i> 55(1):128-130.			
05000179	Acree, F., Jr.; Beroza, M. (1971) Quantitative gas chromatography of isomers of insect repellent N, N-diethyltoluamide. <i>Journal of Economic Entomology</i> 55(5):619-622.			
05002329	Adkins, H.; Cope, A.C.; Fuson, R.C.; Koelsch, C.F.; Shriner, R.L.; Smith, L.I.; Wilds, A.L. (1945) Some Compounds Submitted for Testing as Insect Repellents. Washington, D.C.: U.S. Department of Commerce, Office of Technical Services. (PB report no. 2/410, issued by the National Defense Research Committee of the Office of Scientific Research and Development as OSRD no. 6371).			
00001146	Aero Mist, Incorporated (1972) Flame Extension Test Results: Misty Super Insect Repellent. (Unpublished study received Aug 11, 1972 under 10807-28; CDL:222071-A)			
05000111	All, J.N.; Benjamin, D.M. (1975) A feeding preference bioassay for <u>Neodiprion</u> saw flies on jack pine <u>Pinus banksiana</u> . <i>Annals of the Entomological Society of America</i> 68(3) :553-556.			
05000303	Almazova, V.V.; Bataev, P.S.; Stavrovskaya, V.I.; Akseenke, G.R.; Bezzubova, V.P.; Vorob'eva, Z.G.; Gladkikh, V.F.; Zhukova, L.I.; Zueva, I.K.; Korogodina, Yu.V. (1961) Polevye isytaniya novykh repellentov. [Field tests of new repellents.] <i>Meditinskaya Parazitologiya i Parazitarnye Bolezni</i> . [Medical Parasitology and Parasitic Diseases.] 30(4) :457-464.			
00001154	Anon. (1965) Personal insect repellents: You can select a very efficient product if you know what to look for on the labels. <i>Consumer Reports</i> . Jul: 352-355. (Also In unpublished study received Jan 2, 1974 under 1266-152; submitted by Malter International Corp., New Orleans, LA; CDL:024809-A)			
00001067	Anon. (1968) Insect repellents. <i>U.S. Navy Medical News Letter</i> 52 (6): ?. (Also In unpublished study received Jul 7, 1969 under 1516-52; submitted by Curtis Laboratories, Inc., Kansas City, MO; CDL:024967-A)			

- 05002321 Anon. (1972) CSMA Chemical analysis test methods. Soap, Cosmetics, Chemical Specialties 48(4):152, 155-163
- 05000105 Atkins, E.L.; Kellum, D.; Neuman, K.J. (1977) Repellent additives to reduce pesticide hazards to honey bees. American Bee Journal 117(7) :438-439,457.
- 05000604 Atkins, E.L.; MacDonald, R.L.; Greywood-Hale, E.A. (1975). Repellent additives to reduce pesticide hazards to honey bees: field tests. Environmental Entomology 4(2) :207-210.
- 05000158 Atkins, E.L.; MacDonald, R.L.; McGovern, T.P.; Beroza, M.; Greywood-Hale, E.A. (1975) Repellent additives to reduce pesticide hazards to honeybees: Laboratory testing. Journal of Apicultural Research 14(2) :85-97.
- 00001042 Baker, R.G. 1976. Acute Toxicity Studies with Pyroicide Intermediate 5734: IBT No. 8530-09175. (Unpublished study received September 27, 1976, under 1021-567; prepared by Industrial Bio-Test Laboratories, Inc., submitted by McLaughlin Gormley King Co., Minneapolis, Minn.; CDL:226225-A) (MRID: C0001042. 17A-0037c)
- 05000121 Bar-Zeev, M. (1962) A rapid method for screening and evaluating mosquito repellents. Bulletin of Entomological Research 53 (3) :521-528.
- 05002330 Bartlett, P.D.; Dauben, H.J., Jr.; Hammond, G.S.; McKusick, B.C.; Mueller, G.P.; Packer, C.K.; Ross, S.D.; Schneider, A.; Siegel, S.; Woods, G.F. (1945) The Preparation of Some Organic Compounds for Testing as Insect Repellents. Washington, D.C.: U.S. Department of Commerce, Office of Technical Services. (U.S. Department of Commerce, Office of Technical Services PB report no. 27406 issued by the Office of Scientific Research and Development, National Defense Research Committee as report OSRD no. 6367; National Research Council, Insect Control Committee report no. 155)
- 050002399 Bataev, P.S.; Gridnev, V.I.; Darydova, Z.V.; Makolkin, I.A. (1968) Inftrakrasnye spektry poloshcheniya nekotorykh repellentov [Infrared absorption spectra of some repellents] Trudy, Moskovskii Institut Narod Nogo Khozyaistva. [Transactions, Moscow Institute of the National Economy.] 46:53-61
- 05000261 Bataev, P.S.; Stavrovskaya, V.I.; Luchkovskaya, O.N, inventors; Institute of Medical Parasitology and Tropical Medicine, assignee (1966) Repellents: N-Substituted amides of aromatic and aliphatic-aromatic acids. SU 184,060. Jul 9, 2p. Cl. 451. 9/02
- 05000302 Bataev, P.S.; Iudin, L.G. (1960) Repellentny moshchnoe sredstvo bor'by s gnyssom. [Repellents are a powerful means of control for blood-sucking Diptera.] Priroda (Moscow) 7:82-84.

- 00001121 Beam, F.; Granett, P. (1965) Repellency Tests June-September 1965. (Unpublished study that includes Sections II, III and IV and accompanying tables, received Dec 3, 1966 under 5769-44; prepared by Union Carbide Consumer Products Co. in cooperation with Rutgers State University, submitted by Union Carbide Corp., Tarrytown, NY; CDL:024243-A)
- 05004226 Beckerman, S.C. 1960 An unusual case of aplastic anemia. J. Maine Med. Assoc. 51:53-54 (MRID: 05004226, 17A-0001)
- 05000175 Bigley, W.S.; Roth, A.R.; Eddy, G.W. (1960) Laboratory and field tests against mite and lice attacking poultry. Journal of Economic Entomology 53:12-14.
- 05002553 Blaine, R.L.; Levy, P.F. (1974) The use of thermal evolution analysis (TEA) for the determination of vapor pressure of agricultural chemicals. Analytical Calorimetry 3:185-198
- 05000159 Bowman, M.C.; Beroza, M. (1975) Extraction p-values of pesticides and related compounds in six binary solvent systems. Journal of the Association of Official Analytical Chemists 48(5):943-953
- 05000145 Brady, U.E. (1969) Response of male Sitotroga to the sex pheromones of Sitotroga and Pectinophora females and to Propylure and deet. Deet Analysis in Sitotroga females. Journal of the Georgia Entomological Society 4(1):11-4.
- 00001001 Burkhart, D.M. (1969) Efficacy of Diethyltoluamide as a Fly Repellent. Research Report No. 69:1. Research Book No. 538. (Unpublished study received Feb 12, 1969 under 134-48; prepared by Hess and Clark, Research Department, Division of Richardson-Merrell, Inc., submitted by Hess and Clark, Research Department, Division of Richardson-Merrell, Inc., Ashland, OH; CDL:000602-A)
- 05002323 Camin, J.H.; Clarke, G.K.; Goodson, L.H.; Shuyler, H.R. (1964) Control of the snake mite, Ophionyssus natricis in captive reptile collections. Zoologica 49(2):65-79.
- 00001006 Campbell, L.A. (1970) Repel-A-Gel Stability and Package Compatibility: Research Report No. JAC 70:17. (Unpublished study received Jul 6, 1970 under 134-48 submitted by Hess Clark Research Dept. Div. of Richardson-Merrell, Inc., Ashland, Ohio, CDL:00604-A)
- 00001178 Charles Pfizer & Company, Incorporated (1963) Deet (N, N-Diethyl-m-Toluamide, Pfizer): The Broad-Spectrum Insect Repellent: Data Sheet No. 614. (Unpublished study received Jan 24, 1966 under unknown admin. no; submitted by Diversey Corp., Chicago, III.; CDL:123935-A)

- 00001147 Charles Pfizer and Company, Incorporated (1968) Deet (N,N-diethyl-m-Toluamide, Pfizer): The Broad-Spectrum Insect Repellent: Data Sheet No. 614. (Unpublished study received Apr 2, 1974 under 34281-1; submitted by Tillar Enterprises, Los Angeles, Calif.; CDL:026356-A)
- 05000109 Clark, W.H. (1959) Infrared analysis of the isomers of N,N-diethyltoluamide. *Analytical Chemistry* 31:197-199
- 05000098 Corba, J. (1970) Pouzitie repelentov na ochranu oci havaedzieho dobytku pred medzihostitel'mi telazii. [Repellents for the protection of the eyes of cattle against the intermediate hosts of *Thelazia* spp.] *Veterinarsky Casopis*. [Veterinary Journal.] 13(2):59-63.
- 00004863 Cutter, R.E.; Morison, C.R.W. (1961) A Comparison of the Effect of 6-12 and Deet Insect Repellents Upon Common Surface Coatings. (Unpublished study received Jan 8, 1962 under 3282-45; prepared by Union Carbide Chemical Co., submitted by D-Con Co., Inc., New York, NY: CDL:022564-B)
- 05000301 Cutting, W.C. (1967) Insecticides, repellents and rodenticides. Pages 132-140, In *Handbook of Pharmacology*, 3rd Edition. New York: Appleton-Century-Crofts.
- 05000256 Daniher, F.A., Zaslowsky, J.A., inventors; CPC International, Inc., assignee (1974) Preparation of amides. U.S. patent 3,856,791. Dec 24, 8p.Int. Cl. C07d 87/28, C07d 87/36
- 05000191 Davis, E.E.; Rebert, C.S. (1972) Elements of olfactory receptor coding in the yellowfever mosquito. *Journal of Economic Entomology* 65(4):1058-1061.
- 05005641 Davis, E.E.; Sokolove, P.G. (1976) Lactic acid-sensitive receptors on the antennae of the mosquito, *Aedes aegypti*. *Journal of Comparative Physiology* 105(1):43-54.
- 05002556 Drake, N.L.; Baker, C.M.; Kilmer, G.W.; Melamed, S.; Shenk, W.J.; Weaver, W.E. (1945) The Preparation of Some Compounds for Testing as Insect Repellents. Washington, D.C.: U.S. Department of Commerce, Office of Technical Services. (PB report no. 27409, issued by the Office of Scientific Research and Development, National Defense Research Committee as report OSRD no. 6370; National Research Council, Insect Control Committee report no. 153)
- 05000005 Draughon, F.A.; Ayres, J.C. (1978) Effect of selected pesticides on growth and citrinin production by *penicillium citrinum*. *Journal of Food Sciences* 43(2):576-578, 583.
- 05000078 Dremova, V.P. (1971) Methods for testing repellents on fleas in laboratory conditions. *Meditinskaya Parazitologiya Parazitarnye Bolezni* 40(5):624-625.

- 05000591 Dremova, V.P.; Belan, A.A.; Sytova, N.A.; Reshitova, B.K.; Bodnyuk, N.G.; Agafonova, G.V. (1970) Sravnitel'noe akarorepellentnoe deistvie n ekotorykh preparatov po otnosheniyu k Ixodes persulcatus laboratornykh i polevykh usloviyakh [Comparative acarorepellent action of preparation in relation to Ixodes persulcatus under laboratory and field conditions.] Trudy Tsentral'nogo Nauchno-Issledcuatel'skogo Dezinfedtsionnogo Instituta. [Transactions of the Central Scientific Research Institute of Disinfection.] 19:350-358.
- 05001099 Dremova, V.P.; Markina, V.V. (1975) Zur Frage der minimalen effektiven Dosierungen der Repellents [Minimal effective dosage of repellents] Angewandte Parasitologie 16(3):177-181.
- 05000598 Dremova, V.P.; Markina, V.V.; Akyev, S.A.; Karakhanov, A.G.; Tsetlin, V.M.; Smirnova, S.N. (1971) Ot individualynoi zashchite naseleniya ot gnusa v usloviyakh Turkmenskoi SSSR [Individual protection of the population from mosquitoes in the Turkmen SSR] Zdravookhranenie Turkmenistana 15(3):38-40.
- 05000154 Dremova, V.P.; Zvyagintzeva, T.V.; Markina, V.V. (1977) The sensitivity of repellents of the mosquitoes Anopheles pulcherrimus Theob. International Pest Control 19(3):18-19.
- 05000083 Dremova, V.P., Gleiberman, S.E., Markina, V.V., Babenko, Z.I., Tsetlin, V.M., Smirnova, S.N., Nefedova, L.P., and Zvyagintseva, T.V. 1975. [Entomological and toxicological evaluation of repellent preparations based on diethylamide of phenoxyacetic acid (R-203).] Med. Parazitoll. (Mosk.) 44:699-703 (Translated from Russian) (MRID: 05000083, 17A-0002)
- 05000081 Dremova, V.P., Gleiberman, S.E., Smirnova, S.N., Veronkina, T.V., Tsetlin V.M., Markina, V.V., 1974. [Study of the lotion "DETA-1", its repellent effectiveness and toxicity in warm-blooded animals.] Med. Parazitoll. (Mosk.) 43:67-73 (Translated from Russian) (MRID: 05002662, 17A-0003)
- 05002660 Dubitskii, A.M. (1964) Ispytani ye zashchitnogo deystviya repellentov protiv komarov v usloviyakh pustynnoy zony Kazakhstana [Tests on the protective effect of repellents against mosquitoes under conditions of the desert zone of Kazakhstan.] Izvestiya Akademii Nauk Kazakhskoi SSR, Seriya Biologicheskaya. [Proceedings of the Academy of Sciences of the Kazakh SSR, Biological Series.] 2:72-74.
- 05000285 Eldridge, B.F. (1973) Repellents and impregnants for the control of body lice. Scientific Publication, Pan American Health Organization 263:177-178.

- 05000305 Elizarov, Y.A. (1961) Effects of repellants on physiological characteristics of the chemoreceptors of tick Ixodes persulcatus p. Sch.: An English translation. Moscow Universitet Vestnik Seniyab Biologiya Pochvovedenie. 6(4):45-50.
- 05000588 Elizarov, Y.A. (1964) [Investigation of chemoreception in insects and mites [I] Electrical activity of chemoreceptors of the Haller's organ under the action of repellents] Nauchnye Doklady Vysshei Shkoly, Biologicheskie Nauki. [Scientific Reports of Higher Schools, Biological Sciences. 2:55-59.]
- 05002764 Elizarov, Y.A.; Sinitsina, E.E. (1971) Zlektrofiziologicheskoe issledovanie kontaktnykh khemoretseptornykh sensill osennei zhigalki Stomoxys calcitrans pri deistvii nekotorykh insektorepellentov [Electrophysiological investigation of Stomoxys calcitrans contact chemosensory sensilla to stimulation of some insect repellents Pages 95-104, In Khemoretseptiva Nasekomykh, Materialy k Vsesoyuznomu Simpoziumu po Khemoretseptsii Nasekomykh, Materialy k Vsesoyuznomu Simpoziumu po Khemoretseptsii Nasekomykh, 1st; Sep 8-10, Vilnyus, USSR. [Insect Chemoreception, Materials for the All-Union Symposium on Insect Chemoreception, 1st.] Edited by A. Skirkevicius. Vilnyus, USSR: All-Union Symposium on Insect Chemoreception.
- 05000251 Ellis, L.C.; Kise, M.A., inventors; Virginia Chemicals, Inc., assignee (1972) Verfahren zur Herstellung von N,N-Diaethyl-m-Toluamid in waehriger Atzalkalioesung. [N,N-diethyl-m-toluamide.] GDY 2, 153,307. May 4, 13p. Int. Cl² A0 1H 9/20
- 05000117 Falk, M.S. 1976. Contact urticaria syndrome (letter). Arch. Dermatol. 112:559 (MRID: 05000117, 17A-0025)
- 00001007 Farnam Companies, Incorporated (1968) Repellency Tests: Farnam Wipe-On Gnat and Mosquito Repellent. (Unpublished study received Oct 21, 1968 under 270-49; submitted by ?; CDL:-001925-A)
- 00001070 Fisher, A.M.; Birdsall, C.M. (1972) Flash Point on Insect Repellent Liquid #1659-142. (Unpublished study received May 23, 1972 under unknown admin. no.; submitted by Union Carbide Corp., Agricultural Products, Washington, D.C.; DCL: 11587-A)
- 00001064 Fletcher, D. (1977) 8-Day dietary LC₅₀ study with diethyl toluamide in mallard ducklings: IBT No. 8580-10301. Prepared by Industrial Bio-Test Labs., Inc., submitted by McLaughlin Gormley King Co. (Acc. No. 233280)

- 00001180 Fluno, J.A.; Weidhaas, D.E. (1968) Saving your skin: Insect Repellents. Pages 293-296, In Yearbook of Agriculture, 1968. By U.S. Department of Agriculture. Washington, D.C.: U.S. Government Printing Office. (Yearbook Separate No. 3605: Also In unpublished study received Jun 10, 1971 under 3282-48 and 3282-51; prepared by U.S. Agricultural Research Service, Entomology Research Division, Insects Affecting Man and Animals Research Branch, submitted by Union Carbide Corp., Tarrytown, NY; CDL:007766-B)
- 00004897 Fultz, T.O., Jr. Sandfly Efficacy Report (Comparative Evaluations of Three Repellents. Unpublished study received Feb 21, 1978 under 40195-1; submitted by Reaenge Insect Repellent Corporation., Savannah, GA; CDL:232893-A)
- 05002765 Garnham, P.C.C. (1949) Modern concepts in malaria control. Royal Sanitary Institute Journal 69:617-625.
- 05000185 Garson, L.R.; Buckner, J.H.; Schreck, C.E.; Weidhaas, D.E.; Gilbert, I.H. (1970) Insect repellency of N,N-diethyl-m-toluidine-a nitrogen isotere of deet. Journal of Economic Entomology 63(4) :116-1117.
- 00001071 Gilbert, E.L.R. (1957) Entomological Test Report. Paradichlorobenzene and diethyl toluamide. (Misc. No. 1695.) (Unpublished study received Sep 16, 1957 under ?; prepared by Idea of Asco Electronic Co., submitted by Idea of Asco Electronics Co., Andover, MA; CDL:228307-A)
- 05000157 Gilbert, I.H. (1966) Evaluation and use of mosquito repellents. Journal of the American Medical Association 196(3):163-165.
- 0500287 Gleiberman, S.E., Dremova, V.P., Smirnova, S.N., Durdyev, D., Artydov, A., and Gyshikov, S. 1970. [Study of the toxicity and repellent activity of diethyltoluamide synthesized from petroleum meta-zylene in the Turkmen SSR]. Zdravookhr. Turkmen. 6:29-31 (MRID: 0500287, 17A-0009)
- 05000063 Gleiberman, S.E., and Voronkina, T.M. 1972. [Absorption of diethyltoluamide repellent.] Med. Parazitol. (Mosk.) 41:189-197 (Translated from Russian) MRID: 05000063, 17A-0007)
- 05003638 Gleiberman, S.E., Voronkina, T.M., Latyshev, V.I., and Tsetlin, V.M. 1974. [An experimental study of the resorptive-excretory properties of some repellents.] Med. Parazitol. (Mosk.) 43:446-455 (MRID: 05003638, 17A-0006)
- 05002326 Goodhue, L.D.; Howell, D.E. (1960) Repellents and attractants in pest control operations. Pest Control 28(8):44,46,48,50.
- 05000137 Gothilf, S.; Bar-Zeev, M. (1972) Effect of vapor repellents on the electrical response of insect antennae. Experientia 28(5):601-603.

- 05000274 Gouck, H.K. (1956) Recent developments with a new mosquito repellent. Florida Anti-Mosquito Association, Report of the Annual Meeting 27:124.
- 00001141 Gouck, H.K. (1966) Protection from ticks, fleas, chiggers, and leeches. Archives of Dermatology 93: 112-113. (Also in unpublished study received Jul 22, 1966 under 9110-1; prepared by U.S. Agricultural Research Service, Entomology Research Div., submitted by Paul B. Elder Co., Bryan, OH; CDL: 220305-A)
- 05000114 Gouck, H.K. (1966) Protection from ticks, fleas, chiggers, and leeches. Archives of Dermatology 93(1):112-113.
- 05000280 Gouck, H.K.; Bowman, M.C. (1959) Effect of repellents on the evolution of carbon dioxide and moisture from human arms. Journal of Economic Entomology 52(6):1257-1259.
- 05000213 Gouck, H.K.; Moussa, M.A. (1969) Field tests with bed nets treated with repellents to prevent mosquito bites. Mosquito News 29(2):263-265.
- 05000282 Granett, P. (1960) Insect Repellents for Personal Use. New Brunswick: New Jersey Agricultural Experimental Station. (N.J. A.E.S. circular no. 592).
- 05000288 Griffiths, D.C. (1970) Laboratory test of feeding deterrents, and baits against wireworms (Agriotes species). Pages 398-404, In Proceedings of the British Insecticide and Fungicides Conference, 5th; 1969. Worcester, England: A.W. Pillitt.
- 05006136 Griffiths, D.C.; Jeffs, K.A.; Scott, G.C.; Maskell, P.E.; Mathias, P.L.; Roberts, P.F. (1974) Repellents, synthetic pyrethroids, and organophosphorus compounds as seed dressings against wheat bulb fly Leptohylemyia coarctata (Fall). Plant Pathology 23(2):51-60.
- 00001009 Gunderson, G.R. (1969) Repel-A-Gel Irritation Test on Horses. Research Report No. GRG 69:73. (Unpublished study received Feb 12, 1969 under 134-48; prepared by Hess and Clark, Research Dept., Div. of Richardson-Merrell, Inc., submitted by Hess and Clark, Research Dept., Div. of Richardson-Merrell, Inc., Ashland, OH; CDL:133550-B)
- 05002325 Hall, S.A.; Green, N.; Beroza, M. (1957) Insect repellents and attractants. Journal of Agricultural and Food Chemistry 5(9):663-667, 669.
- 050002014 Hardwicke Chemical Co. (1976) Deet. Confidential Company Study; submitted by Hardwicke Chemical Company, Elgin, S.C. #accession 226379
- 05000012 Hasegawa, Y., Shiota, N.; Yamazaki, Y., inventors; Mitsubishi Gas Chemical Co., Inc., assignee (1977) [N,N-diethyltoluamides.] Japanese ko kai 77 125,142. Oct 20, 4p

- 050002015 Hasegawa, Y.; Shirota, N.; Yamazaki, Y., inventors; Mitsubishi Gas Chemical Co., Inc., assignee (1977) [N,N-diethyltoluamides.] Japanese Kokai 77 125,142. Oct 20, 4 p
- 00001122 Haynes, H.L. (1972) Supporting Data for "6-12 Plus" Brand Insect Repellent Spray. (Unpublished study that includes tables and figures, received Jun 27, 1972 under 3282-48; submitted by Union Carbide Corp., New York, NY; CDL:007767-A)
- 05000203 Heller-Haupt, A.; Rusvine, J.R. (1974) Tests of acaricides against house dust mites. Journal of Medical Entomology 11(5):551-558.
- 00001068 Hercules Incorporated (1977) Metadelphene: The All-Purpose Insect Repellent: Bulletin AP-105B. (Unpublished study received Jul 7, 1969 under 1516-52; submitted by Curts Laboratories, Inc., Kansas City, Mo., CDL:024967-B)
- 00001152 Hercules Incorporated (1972) Properties: Physical and Chemical: [Metadelphene]: AP-105B. (Incomplete study, p. 2; unpublished study received Jun 12, 1972 under 11515-12; submitted by Peterson Puritan, Inc., Danville, Ill.; CDL:004536-E)
- 00001025 Hercules Powder Company (1975) Meta Delphene: "The Complete Insect Repellent." Wilmington, Del., Hercules Powder Co., Inc. (Bulletin No. 213; Also In unpublished submission received Aug. 13, 1957 under unknown admin. no., submitted by Hercules Powder Co., Wilmington, Del.; CDL: 108471-A)
- 05000208 Iglitsina, L.S.; Fedder, M.L. (1959) sintez dietiltoluamida-novogo repellenta. [Synthesis of diethyltoluamide—a new repellent.] Meditsinskaya Promyshlennost SSSR. [Medical Industry of the USSR.] 13(8):20-22
- 05000304 Ikeda, Y. (1959) Some simplified methods for the evaluation of the effectiveness of fly repellents in laboratory and outdoors. Botyu Kagaku. [Scientific Pest Control.] 24(4):175-181.
- 05000050 Ikeda, Y.; Kondo, M. (1961) Efficiency of certain repellents to female German cockroaches, Blattella germanica. Botyu Kagaku 26(4):112-114.
- 05000014 Ito, M.; Imaseki, K.; Hirano, S.; Amano, K., inventors; Tsumura Juntendo Co., Ltd., assignee (1977) [Diethyltoluamide insect repellent] Japanese patent specification 77 110,824. Sep 17. 1p
- 05000257 Ito, M.; Naito, T., inventors; Showa Denko K.K, Japan, assignee (1976) Process for preparation of N,N-dialkyltoluamide. U.S. patent 3,947, 51114. Mar 30, 4p

- 05000032 Ivanov, V.I. (1975) Antropofilii oten'ei Krovososki Lipoptena cervi L. (Diptera, Hippoboscidae). [Anthropophilicity in Lipoptena cervi L. (Diptera, Hippoboscidae). Meditsinskaya Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 44(4):491-495.
- 05000112 Jacobson, M. (1966) Chemical insect attractants and repellents. Annual Review of Entomology 11:403-422.
- 05002312 Jacobson, M.; Jones, W.A. (1974) Attraction of the male pink bollworm moth under laboratory and field conditions. Environmental Letters 6(4):297-301.
- 05000181 Johnson, H.L.; Skinner, W.A.; Maibach, H.I.; Pearson, T.P. (1967) Repellent activity and physical properties of ring-substituted N,N-diethylbenzamides. Journal of Economic Entomology 60(1):173-176.
- 00001083 Johnson, S.C. & Sons, Incorporated (1976) Toxicology Data Applicable to Hydrocarbon Propellant Off: Pressurized. (Unpublished study received Jul 10, 1976 under 4822-10; CDL:233491-A)
- 00001100 Johnson, S.C. & Sons, Incorporated (1977) Chemical and Physical Properties [of Diethyltoluamide]: Summary. (Unpublished study received Mar 21, 1978 under 4822-160; CDL:233261-A)
- 05002788 Kashin, P.; Kardatzke, M.L. (1975) Diurnal rhythm, age, and other variables affecting yellowfever mosquito avidity and the laboratory assay of repellents. Journal of Economic Entomology 68(6):766-768.
- 05000103 Kettle, D.S. (1969) The ecology and control of blood-sucking ceratopogonids. Acta Tropica 26(3):235-248.
- 05000100 Khan, A.A. (1965) Effects of repellents on mosquito behavior. Quaestiones Entomologicae 1(1):1-35.
- 05000218 Khan, A.A.; Maibach, H.I.; Skidmore, D.L. (1975) Addition of perfume fixatives to mosquito repellents to increase protection time. Mosquito News 35(1):23-26.
- 05000017 Klier, M.; Hamburg, A.G. (1976) Neue Insektenabwehrmittel - am Stickstoff disubstituierte beta-Alaninderivate. [New insect repellents - derivatives of N-disubstituted beta-alanine. Journal of the Society of Cosmetic Chemists 27(3):141-153.
- 05000296 Knipling, E.F. (1960) Use of insecticides, attractants and repellents. Washington D.C.: USDA. (U.S. Department of Agriculture, Agricultural Research Service).

- 00001074 Knott, W.B.; Johnston, C.D. (1972) Acute Oral Toxicity to Rats: 3560D105-5. (Unpublished study that includes formulations, received June 12, 1972 under 4822-119; prepared by Woodard Research Corp., submitted by S.C. Johnson & Son, Inc., Racine, Wis.; CDL:007514-C)
- 05000601 Kolomiets, A.F.; Bliznyuk N,K.; Strel'tsov, R.V.; Savenkov, N.F.; Kalutskii, L.A. (1969) Synthesis of carboxylic acid amides. A translation of Zhurnal Prikladnoi Khimii 42(4):963-966
- 05000161 Krejci, M.; Vespalec, R.; Sirec, M. (1972) Sampling system and analysis of basic components of sprays by liquid chromatography. Journal of Chromatography 65(1):333-340
- 05002671 Kunzmann, T. (1961) Insektenabwehrende und hautschuetzende Prepare in Aerosolform. [Insect repellent and sunscreen preparations in aerosol form.] Seifen, Oele, Fetten Wachse 87(8):179
- 05002336 Lal, H. (1964) Development of an Orally Effective Insect Repellent. Washington, D.C.: U.S. Army Medical Research and Development Command, Office of the Surgeon General. (IIT Research Institute, Chicago, IL report no. IITRI-C222-9; available from: NTIS, Springfield, VA; AD-603 114)
- 05000294 Lal, H.; Ginocchio, S.; Hawrylewicz, E.J. (1963) Procedure for bioassaying mosquito repellents in laboratory animals. Proceedings of the Society for Experimental Biology and Medicine 113(3):770-772.
- 05000600 Leitman, Y. I.; Perzner, M.S. (1963) Development of methods for producing the repellent diethyltoluamide. A translation of: Zhurnal Prikladnoi Khimii 36(3):632-639
- 05000058 Leshchev, V.V. 1970. [Stability of the repellents benzimine, carboxide, and N,N-diethyl-m-toluamide (DETA) used to protect animals.] Khim. Sel: Khoz. 8:376-379 (Translated from Russian) (MRID: 05000058, 17A-0013)
- 05000059 Leshchev, V.V. (1970) Ustoichivost' repellentor benzimina, Karboksida i dietiltoluamido, primaya emykh dlya zashchity zhivotnykh. [Stability of the repellents benzimine, carboxide, and N,N-diethyl-m-toluamide (DETA) used to protect animals.] Khimiya v Sel'skom Khozyaistve. [Chemistry in Agriculture.] 8(5):376-379
- 05000102 Lindberg, U.H.; Ulff, B.; Yeoman, G. (1968) Cinchoninic and 3-alkylcinchoninic acid alkyl esters. A new class of insect repellents. Acta Pharmaceutica Suecica 5(5):441-448.

- 00001128 Lomax, J.L.; Gränett, P. (1971) Current Laboratory procedures for the development of improved insect repellents at Rutgers-the State University. Pages 41-47, In Proceedings of the 58th Annual Meeting of the New Jersey Mosquito Extermination Association; Mar 17-19, 1971, Atlantic City, NJ. (Also In unpublished study received Aug 24, 1973 under 5769-97; prepared by Rutgers University, Department of Entomology, submitted by Union Carbide Corp., New York, NY; CDL:003305-B)
- 05002609 Lust, S. (1960) Ueber Wirkung and biologische Pruefung von Repellentien) [The effect and biological testing of repellents] Parfuemerie Kosmetik 41(8):304-306.
- 05000022 Maibach, H.I.; Akers, W.A.; Johnson, H.L. (1974) Insects. Topical insect repellents. Clinical Pharmacology and Therapeutics 16(511):970-973.
- 05000587 Mandel'baum, Y.A.; Lomakina, V.I.; Mel'nikov, N.N. (1961) Amidy Kislota kak otpugivayushchie sredstva (repellent) [Amides of acids as repellents.] Moskv Nauk Inst Udobr Insektofungitsidam Sborn Rabot 171:151-158.
- 05000584 Mandel'baum, Y.A.; Khcheyan, K.E. (1961) Effektivnyy Repellent-diethylamid metatoluilovoy kistoty preparat Deta. [An effective repellent—the diethylamide of m-toluic acid (DETA).] Khimicheskaya Promshlennost SSR. [Chemical Industry of the USSR.] 10:686-690
- 05000238 Markarian, H.; Enos, H.F.; Pratt, J.J., Jr. (1965) Analysis of diethyltoluamide. Soap and Chemical Specialties 41(10):94, 96-97, 220
- 05000023 Massie, F.S. (1974) Papular urticaria: etiology, diagnosis and management. Cutis 13(6):980-986.
- 00001173 Mastri, C. (1969) Acute Oral Toxicity Study on 3071D14-2 in Albino Rats: Report to S.C. Johnson & Son, Inc.: [Deet]: IBT No. A7340. (Unpublished study received Jun 23, 1969 under 4822-Ex-8; prepared by Industrial Bio-Test Laboratories, Inc., submitted by S.C. Johnson & Son, Inc., Racine, Wis.; CDL:127306-E)
- 00001176 Mastri, C. (1969) Addendum Report to S.C. Johnson and Son, Inc.: 20-Day Subacute Dermal Toxicity Study on Two Formulations of Insect Repellent in Albino Rabbits: [DEET]: IBT No. A7241. (Unpublished study received Jun 23, 1969 under 4822-Ex-8; prepared by Industrial Bio-Test Laboratories, Inc., submitted by S.C. Johnson & Son, Inc., Racine, Wis.; CDL:127306-H). File No. 00001176 (AA0104)
- 00001174 Mastri, C. (1969) Eye Irritation Test on 3071D15-2 in Albino Rabbits: Report to S.C. Johnson and Son, Inc., submitted by S.C. Johnson & Son, Inc., Racine, Wis.; CDL:127306-F)

- C0001175 Mastri, C. (1969) 20-Day Subacute Dermal Toxicity Study on two Formulations of Insect Repellent in Albino Rabbits: [DEET]: Report to S.C. Johnson and Son, Inc.: IBT No. A7241. (Unpublished study received Jun 23, 1969 under 4822-EX-8; prepared by Industrial Bio-Test Laboratories Inc., submitted by S.C. Johnson & Son, Inc., Racine, Wis.; CDL: 127306-G). File No. 00001175 (AA0075)
- 05001271 McAndless, J.M., Underwood, G.P. (1976) Personal Protection Against Biting Flies: The Relative Effectiveness of Over-jackets Treated with Various Insect Repellents. Ottawa, Ontario, Canada: Department of Defence. (Defence Research Establishment technical note no. 76-28: available from: NTIS, Springfield, VA; AD A034660)
- 00001026 McCann, J.A. (1972) Fish Toxicity Laboratory report: Toxicity of Hercules Metadelphene to rainbow trout: Test No. 483. Prepared by Animal Biology Laboratory, EPA; submitted by Hercules, Inc., (Acc. No. 129780)
- 00001186 McLaughlin Gormley King Company (1958) (Testimonial Letters: Deet.) (Unpublished study that includes 9 letters, received Feb 1956 under 1021-511; submitted by ?; CDL:005214-A)
- 00001187 McLaughlin Gormley King Company (1958) (Testimonial Letters: Deet.) (Unpublished study that includes 5 letters, received Aug 5, 1958 under 1021-512; CDL:005216-A)
- 00001057 McLaughlin Gormley King Company (1965) MGK (R) Repellents for personal Use. (Unpublished study received Feb 15, 1965 under unknown admin. no.; CDL: 105970-B)
- 00001061 McLaughlin Gormley King Company (1973) Repellency. MGK File No. D-1017-73. (Unpublished study received Feb 27, 1974 under 1021-1312; prepared by McLaughlin Gormley King Company, MGK Research and Development Department, Biological Testing Laboratory, submitted by ?; CDL:221198-A)
- 00001144 Med-Pak Corporation (1970) Deet (N, N-Diethyl-M-Toluamide). (Incomplete study, pp.2-5; unpublished study received May 27, 1970 under 10823-1; CDL:226620-A)
- 05000215 Michael, A.G., Grant, G.S. (1974) Toxicity of the repellent DEET (N,N-diethyl-meta-toluamide) to Gambusia affinis (Baird and Girard). Mosquito News 34(1): 32-34
- 00001004 Miller, R.L. (1969) Effect of Repel-A-Gel on Horses. Research Report No. RLM 69:68. (Unpublished study received Oct 30, 1969 under 134-48; prepared by Hess and Clark, Research Dept., Div. of Richardson-Merrell, Inc., submitted by Hess and Clark, Research Dept., Div. of Richardson-Merrell, Inc., Ashland, OH; CDL:000603-A)
- 05000051 Navr'atil, K. (1966) Dieltamid. (ointment). Cesk Farm 15(8):453.

- 05002303 Neumark, S.; Jacobson, M.; Teich, I. (1974) Field evaluation of propylure, hexalure, and their formulations with deet, dodecyl acetate, and an antioxidant as attractants for male pink bollworm moths. *Environmental Letters* 7(2):21-30.
- 05000315 Newland, G.C.; Greear, G.R.; Tamblyn, J.W., inventors; Eastman Kodak Co., assignee (1966) Opaque polyethylene film containing pro-oxidants. British patent specification 1,052,998. Dec 30, 5p. Int. cl.-2 C 08f 29/02.
- 05000691 Nudel'Man, Z.N. (1955) Repellents. *Priroda* (Moscow). [Nature.] 44(10):86-88.
- 05000314 Norton, R.V., inventor; Sun Ventures, Inc., assignee (1976) Preparation of alkylcarboxamides. U.S. patent 3,985,805. Oct 12, 4p. Int. Cl² c 07 102/08; C 07C 103/76
- 05000687 Ota, Y.; Tateishi, M. (1963) [The control of tyroglyphid mites by chemicals] *Igaku to Seibutsugaku*. [Medicine and Biology] 66(3):103106.
- 05000108 Palmer, J.S. (1969) Toxicologic effects of aerosols of N,N-diethyl-m-toluamide (deet) applied on skin of horses. *American Journal of Veterinary Research* 30(11): 1929-1932
- 05002332 Phalen, W.L., Jr., inventors; Hercules Powder Co., assignee (1961) N,N-diethyltoluamides. U.S. patent 2,991,310. Jul 4. 3p
- 05000312 Phalen, W.L., Jr., inventors; Hercules Powder Co., assignee (1961) Recovery of N,N-diethyltoluamide. U.S. patent 2,995,604. Aug 8, 4p
- 05000088 Pokrovskaya, E.I.; Bataev, P.S.; Ryabykh, I.V. (1952) Ispytanie novykh preparatov, otpugivayushchikh komarov v prirodnykh usloviyakh voronezhskoi oblasti. [Testing new preparations repelling mosquitoes under natural conditions in the Voronezh Region.] *Nauchnye Doklady Vysshei Shkoly, Biologicheskie Nauki*. [Scientific Reports of Higher Schools, Biological Science.] 3:23-26.
- 05000695 Pokrovskaya, E.I.; Ryabykh, L.V.; Bataev, P.S. (1961) Predvaritelnye polevye ispytaniya novykh preparatov otpugivayushchikh komarov, v prirodnykh usloviyakh voronezhskoy oblasti [Preliminary field trial of new mosquito repellents] *Meditinskaya Parazitologiya i Parazitarnye Bolezni*. [Medical Parasitology and Parasitic Diseases. 30(6):723-726.
- 05000097 Polyakov, V.A. (1975) The effectiveness of treatments applied to reindeer against blood-sucking insects. *Veterinariya* 5:42-44.
- 05000063 Potapov, A.A.; Vladimirova, V.V. (1963) The effect of repellents on some horsefly species at different air temperatures (field trials with Skufiin's traps). *Meditinskaya Parazitologiya i Parazitarnye Bolezni* 32(5):542-546.

- 05000080 Potapov, A.A.; Vladimirova, V.V.; Sazonova, E.V.; Bogdanova, B.I., (1973) [Comparative tests of dimethyl phthalate, diethyl phthalate, and dibutyl phthalate as repellents. Meditsinskaya-Parazitologiya i Parazitarnye Bolezni. [Medical Parasitology and Parasitic Diseases.] 42(6):692-697.
- 05000026 Potapov, A.A., Tsizin, Y.S., Kharitonova, S.I., Vladimirova, V.V., and Sazonova, E.V. 1977. [Repellent napkins for protection against blood-sucking Diptera.] Med. Parazitol. (Mosk.) 4:434-439 (Translated from Russian) (MRID: 05000026, 17A-0015)
- 05000148 Raha, C.R. (1960) Mosquito repellent preparation from xylene. Indian Journal of Applied Chemistry 23(1): 65-66
- 00001062 Rausina, G. (1977) Four-day static aquatic toxicity studies with diethyl toluamide in rainbow trout and bluegills: IBT No. 8560-10299; Prepared by Industrial Bio-Test Labs., Inc.; submitted by McLaughlin Gormley King Co. (Acc. No. 230395)
- 05000028 Rustamov, B.R.; Tadzhieva, V.S. (1976) Problema Bor'by s komarami i profilaktika malyarii na territorii Arnasaiskogo ponizheniya v Uzbekskoi SSR. [Problems of mosquito control and prophylaxis against malaria in the Arnasai depression in the Uzbek SSR. Meditsinskii Zhurnal Uzbekistana. [Medical Journal of Uzbekistan] 2:46-50.
- 00001075 S.C. Johnson and Son, Incorporated (1972) Repellent Lotion 3727D50. Project 573. (Unpublished study received Sep 18, 1972 under 4822-125; prepared by S.C. Johnson and Sons, Inc., Product Research Dept., submitted by ?; CDL:007521-A)
- 00001076 S.C. Johnson and Son, Incorporated (1973) Insect Repellent Gel (3936D1-1). (Unpublished study received Jul 19, 1973 under 4822-129; prepared by S.C. Johnson and Son, Inc., Product Research Dept., submitted by ?; CDL:007525-A)
- 00001094 S.C. Johnson and Son, Incorporated (1974) Insect Repellent Towellettes 3936D74-1. (Unpublished study received Nov 4, 1974 under 4822-142; prepared by S.C. Johnson and Son Inc., Product Research Department, submitted by ?; CDL:000918-A)
- 00001155 S.C. Johnson and Son, Incorporated (1977) Mosquito Repellency (Aedes aegypti). (Project Assignment A-5589; unpublished study received Jun 13, 1977 under 4822-119; prepared by S.C. Johnson and Son., Inc., Research and Development Div., Biology Research Dept., submitted by ?; CDL:230624-A)
- 00001156 S.C. Johnson and Son, Incorporated (1977) Stable Fly Repellency (Stomoxys calcitrans). (Project Assignment A-5589; unpublished study received Jun 13, 1977 under 4822-119; prepared by S.C. Johnson and Son., Inc., Research and Development Div., Biology Research Dept., submitted by ?; CDL:230624-B)

- 05002557 Saf'yanova, V.M. (1960) Ob ispol'zovanii kolokola A.S. Monchad-
skogo pri polevom ispytanii deystviya repellentov na krovo-
sosushchikh Dvukrylykh [Use of the A.S. Monchadsky bell in
field tests of the effect of repellents against blood-sucking
Diptera] Zoologicheskii Zhurnal. [Journal of Zoology.]
XXXIX(8) :1169-1173.
- 05000690 Saf'yanova, V.M. (1963) Results of testing individual bed-nets
impregnated with repellents for protection against sand-
flies. Meditsinskaya Parazitologiya i Parazitarnye Bolezni.
[Medical Parasitology and Parasitic Diseases.] 35(5):549-
551.
- 05000602 Saf'yanova, V.M. (1963) Rezulytaty polevogo ispytaniya nekotorykh
repellentov v otnosheniia moskitov (podsemeistvo Phleb-
otominae) [Results of field tests of some repellents
against sand flies belonging to the subfamily Phleboto-
minae Zoologicheskii Zhurnal 42(3):461-462.
- 00001142 Samuel J. Milazzo Manufacturing Company (1970) Qik Joe Mosquito
Chaser: Effectiveness Report. (Unpublished study received
Feb 18, 1970 under 8218-2; prepared by Samuel J. Milazzo
Manufacturing Company, submitted by ?; CDL:227966-A)
- 05000610 Saruto, K.; Gono T.; Nakajima. G.; Yashitomi Pharamaceutical
Industries, Ltd., inventors; (1966) [N,N-dialkylaryamides.]
Japanese patent specification 20,822. Dec 5, 3p
- 05002065 Schmidt, C.H.; Bowman, M.C.; Acree, F., Jr. (1958) Ultraviolet
determination of the insect repellent diethyltoluamide. Jour-
nal of Economic Entomology 51(5): 694-697
- 05002305 Schwartz, R.D.; Mathews, R.G.; Pedro, N.A. (1977) High-tempera-
ture gas-liquid chromatography with a dendritic salt sup-
port. Journal of Chromatography 142:103-107
- 05000029 Schreck, C.E.; Weidhaas, D.E.; Smith, N. (1977) Evaluation of
electronic sound producing devices against Aedes taenior
hynchus and Aedes-sollicitans. Mosquito News 37(3):529-531.
- 05000263 Shambaugh, G.P.; Brown, R.P.; Pratt, J.J., Jr. (1957) Repellents
for biting arthropods. Pages 277-303, In Advances in
Post Control Research. New York: New York Interscience.
- 05000133 Sholdt, L.L.; Grothaus, R.E.; Schreck, C.B.; Gouck, H.K. (1975)
Field studies using repellent-treated wide-mesh net jackets
against Glossina morsitans in Ethiopia. East African
Medical Journal 52(5):277-283.
- 05000595 Shrunderenko, G.V.; Burykina, A.P.; Kharitonova, S.I. (1976)
Effektivnostv repellentnykh salfetok v polevykh usloviyukh)
[Effectiveness of repellent handkerchiefs under field
conditions] Voenno Meditsinskii Zhurnal. [Journal of Mili-
tary Medicine.] (6):76.

- 05000049 Sixl, W. von; Stueznier, D. (1975) Autan als Zeckenabwehrmittel.
[Autan as a product to control ticks.] Angewandte Parasitologie 16(2):106-108.
- 05002337 Skinner, W.A. (1956) Attractiveness and Repellency of Man to Mosquito Bites. Menlo Park, Calif.: Stanford Research Institute. (Available from: NTIS, Springfield, VA; AD-571)
- 05000037 Skinner, W.A.; Tong, H.C.; Johnson, H.; Parkhurst, R.M.; Thomas, D.; Spencer, T.; Akers, W.; Skidmore, D.; Maibach, H. (1977) Influence of human skin surface lipids on protection time of topical mosquito repellent. Journal of Pharmaceutical Sciences 66(12):1764-1766.
- 00001020 Smith, C.N. (1965) Insect repellents: Their present usefulness and future development. Pages 507-509, In Section 8: Insecticides and Toxicology: (Proceedings of the) XIIth International Congress of Entomology; 1964 (1965), London, N.P. (Also in unpublished study received May 7, 1970 under 891-37; prepared by U.S. Agricultural Research Service, Entomology Research Div., submitted by Hercules, Inc., Agricultural Chemicals, Wilmington, DE; CDL:005111-G)
- 05000277 Smith, C.N.; Gilbert, I.H.; Gouck, H.K. (1962) Factors in the attraction of mosquitoes to hosts, and their relation to protection with repellents. Pages 82-86, In Transactions of the International Congress of Entomology, 11th Congress, vol. 3, supp. 1/2, 5/7.
- 05000300 Smith, C.N.; Gilbert, I.H.; Gouck, H.K.; Bowman, M.C.; Acree, P., Jr.; Schmidt, C.H. (1963) Factors affecting the protection period of mosquito repellents. Pages 1-36, In U.S. Department of Agriculture Technical Bulletin, No. 1285.
- 00001031 Smith, C.N.; Gilbert, I.H.; Gouck, H.K. (1960) Use of Insect Repellents. ARS-33-26. (Unpublished study received 1960 under ?; prepared by U.S. Agricultural Research Service, Entomology Research Div., submitted by Union Carbide Corp., Agricultural Products, Washington, D.C.; CDL:117588-A)
- 00001183 Smith, W.W. (1960) Exposure Test: Inspection Report. Report 1-Final. (Efficacy tests on CDK 47/1, Normal Sun Bath and CZ8-11/1 when applied to normal human skin subjected to sun-shine.) (Unpublished study received Dec 19, 1961 under 6650-1; prepared by South Florida Test Service, Inc., submitted by Revlon, Inc., Bronx, NY; CDL:229503-C)
- 05000603 Spencer, T.S.; Shimmin, R.K.; Schoeppner, B.F. (1975) Field test of repellents against the valley black gnat, Leptoconops carteri Hoffman (Diptera: Ceratopogonidae). California Vector Views 22(1):5-7.
- 00004233 Standard Seed and Feed company. 1950. Pharmacological and Rodenticide Laboratory Report. (Unpublished study received on unknown data under unknown administration number; CDL: 133493-A) (MRID: 00004233, 17A-0018c)

- 05000094 Starcheus, A.P. (1963) Dietiltoluamid dlya zashchity zhyvotnykh ot gnusa. [Diethyltoluamide for protection of animals against gnats.] Veterinariya. [Veterinary Science.] 40(6):78-79
- 05000095 Starcheus, A.P. (1964) Zashchita zhyvotnykh of gnusa. [Protection of livestock against blood-sucking insects.] Veterinariya [Veterinary Science.] 41(6):109-111.
- 05000589 Starkov, A.V.; Katumina, V.I.; Stomakhina, Z.P.; Latyshev, V.I. (1967) Preparation of diethyltoluamide by a transamidation method. A translation of: Khimko Farmatsevticheskii Zhurnal]. [Chemical Pharmaceutical Journal] (ii):23-25
- 00001145 Straight Arrow Incorporated (1971) "1971" Study Report on Formula to be Marketed as "Jungle Jel". (Unpublished study that includes appended incomplete Hercules Agricultural Chemicals report AP-105B, received May 16, 1972 under 11314-1; CDL:-013082-A)
- 05002664 Ternovskaia, N.T.; Kazinik, E.M.; Pastushenko, V.G.; Samokhina, L.A. (1973) Kromatograficheskoe operdelenie diztiltoluamide v repellentnykh prepartakh na ego osnove [Chromatographic determination of diethyltoluamide in its repellent preparations] sbornik Trudov po Bytovio Khimii. [Chemistry.] 1:105-109
- 00001005 Thompson, W.S. (1969) Effect of Repel-A-Gel on Horses. Research Report No. WST 69:24. (Unpublished study received Oct 30, 1969 under 134-48; prepared by Hess and Clark, Research Dept., Div. of Richardson-Merrell, Inc., submitted by Hess and Clark, Research Dept., Div. of Richardson-Merrell, Inc., Ashland, OH; CDL:000603-B)
- 05002315 Touhey, J.G.; Bray, D.F. (1961) Certain compounds as feeding deterrents against the smaller European elm bark beetle, Scolytus multistriatus. Journal of Economic Entomology 54(2):293-296.
- 05000606 Traub, R.; Elisberg, B.L. (1962) Comparative efficacy of diethyltoluamide skin-application repellent (deet) and M-1960 clothing impregnant against mosquitoes in the nipah palm-mangrove swamps in Malaya. Pacific Insects 4(2): 314-318.
- 05000605 Traub, R.; Elisberg, B.L. (1962) Field tests on diethyltoluamide (deet), a highly effective repellent against mosquitoes in the nipah palm-mangrove swamps in Malaya. Pacific Insects 4(2):303-313.
- 05000589 Tristan, D.F.; Rudenchik, Y.V.; Chekalin, V.B.; Vishnyakova, I.K.; Dzhumabekov, K.D. (1970) O pyt primeneniya repellentov protiv blokh i kleshchei [Use of flea and tick repellents] Problemy Osobo Opasnykh Infeksii. [Problems of Highly Dangerous Infections.] (5):124-127.

05000206 Tsakotellis, P.; Johnson, H.L.; Skinner, W.A.; Skidmore, D.; Maibach, H.I. (1971) Topical mosquito repellents. III. Carboxamide acetals and ketals and related carbonyl addition derivatives. Journal of Pharmaceutical Sciences 60:84-89.

05000262 Tsetlin, V.M.; Dremova, V.P.; Zhuk, E.B.; Pumpurs, A.I., inventors; Central Scientific-Research Disinfectant Institute, assignee (1968) Charge for aerosol repellent containers. SU 227,013. Sep 26, lp. Cl. 451, 9/00

00001054 U.S. Department of Agriculture. (1957) A New Repellent, Diethyltoluamide: Correspondence Aid No.6, revised. (Unpublished study received Oct 22, 1974 under 1021-1323; prepared by Agricultural Research Service, Entomology Research Div., submitted by McLaughlin Gormley King Co., Minneapolis, Minn.; CDL:028339-H)

05000147 U.S. Department of Agriculture. (1976) Controlling Chiggers. Washington, D.C.: U.S.D.A. (USDA home garden bulletin, no. 137).

00001019 U.S. Department of Agriculture. (?) Protecting Yourself From Insects in Recreational Areas. Pages 11-14, In Home and Garden Bulletin No. 4. Washington, D.C.: U.S. Department of Agriculture. (Also in unpublished study received May 7, 1970 under 891-37; submitted by Hercules, Inc., Agricultural Chemicals Wilmington, DE; CDL:005111-F)