U.S. ENVIRONMENTAL PROTECTION AGENCY NATIONAL EUTROPHICATION SURVEY WORKING PAPER SERIES



REPORT
ON
SCHROON LAKE
ESSEX AND WARREN COUNTIES
NEW YORK
EPA REGION II
WORKING PAPER No. 169

PACIFIC NORTHWEST ENVIRONMENTAL RESEARCH LABORATORY

An Associate Laboratory of the NATIONAL ENVIRONMENTAL RESEARCH CENTER - CORVALLIS, OREGON and

NATIONAL ENVIRONMENTAL RESEARCH CENTER - LAS VEGAS, NEVADA

REPORT
ON
SCHROON LAKE
ESSEX AND WARREN COUNTIES
NEW YORK
EPA REGION II
WORKING PAPER No. 169

WITH THE COOPERATION OF THE

NEW YORK STATE DEPARTMENT OF ENVIRONMENTAL CONSERVATION

AND THE

NEW YORK NATIONAL GUARD

NOVEMBER, 1974

CONTENTS

	Page
Foreword	ii
List of New York Study Lakes	iv
Lake and Drainage Area Map	V
Sections	
I. Conclusions	1
II. Lake and Drainage Basin Characteristics	3
III. Lake Water Quality Summary	4
IV. Nutrient Loadings	9
V. Literature Reviewed	14
VI. Appendices	15

FOREWORD

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nation-wide threat of accelerated eutrophication to fresh water lakes and reservoirs.

OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

- a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.
- b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.
- c. With such a transformation, an assessment of the potential for eutrophication control can be made.

LAKE ANALYSIS

In this report, the first stage of evaluation of lake and watershed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning [§303(e)], water quality criteria/standards review [§303(c)], clean lakes [§314(a,b)], and water quality monitoring [§106 and §305(b)] activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

ACKNOWLEDGMENT

The staff of the National Eutrophication Survey (Office of Research & Development, U. S. Environmental Protection Agency) expresses sincere appreciation to the New York Department of Environmental Conservation for professional involvement and to the New York National Guard for conducting the tributary sampling phase of the Survey.

Henry L. Diamond, Commissioner of the New York Department of Environmental Conservation, and Leo J. Hetling, Director, and Italo G. Carcich, Senior Sanitary Engineer, Environmental Quality Research, Department of Environmental Conservation, provided invaluable lake documentation and counsel during the Survey.

Major General John C. Baker, the Adjutant General of New York, and Project Officer Lieutenant Colonel Fred Peters, who directed the volunteer efforts of the New York National Guardsmen, are also gratefully acknowledged for their assistance to the Survey.

NATIONAL EUTROPHICATION SURVEY

STUDY LAKES

STATE OF NEW YORK

LAKE NAME

Allegheny Reservoir

Black Canandaigua Cannonsville Carry Falls Cassadaga Cayuga Champlain

Chautauqua Conesus Cross Goodyear Huntington Keuka Long

Lower St. Regis

Otter Owasco

Raquette Pond

Round

Sacandaga Res.

Saratoga Schroon Seneca Swan

Swinging Bridge Res.

COUNTY

Cattaraugas, NY; McLean,

Warren, PA
St. Lawrence
Ontario
Delaware
St. Lawrence
Chautauqua

Seneca, Tompkins

Clinton, Essex, NY; Addison, Chittenden, Franklin, VT

Chautauqua Livingston

Cayuga, Onondaga

Otsego Sullivan Ontario Hamilton Franklin Cayuga Cayuga Franklin Saratoga

Fulton, Saratoga

Saratoga Essex, Warren

Seneca, Schyler, Yates

Sullivan Sullivan

SCHROON LAKE

STORET NO. 3634

CONCLUSIONS

A. Trophic Condition:

Survey data indicate that Schroon Lake is oligotrophic. Of the 26 New York lakes studied, none had lower mean total phosphorus, only one had less mean inorganic nitrogen, and none had less mean chlorophyll a.

Survey limnologists noted that Schroon Lake exhibited high clarity and was clear of algal blooms at all sampling times.

B. Rate-Limiting Nutrient:

Algal assay results show that Schroon Lake was phosphorus limited at the time the assay sample was collected. The lake data also indicate phosphorus limitation at all sampling times (N/P ratios were greater than 36/1, and phosphorus limitation would be expected).

C. Nutrient Controllability:

1. Point sources--It is estimated that point sources accounted for only 3% of the total phosphorus load to Schroon Lake during the sampling year. The calculated <u>input</u> phosphorus loading rate was less than that proposed by Vollenweider (in press) as "permissible" (i.e., an oligotrophic rate--see page 13); and it might be concluded that point-source control of phosphorus would have little

effect on the trophic condition of the lake. However, during the sampling year some 17,800 pounds more phosphorus and 260,500 pounds more nitrogen were measured leaving the lake than can be accounted for by all measured and estimated inputs. While nitrogen loss can occur under certain conditions, with losses of both primary nutrients at the indicated magnitude, it is most probable that unknown and unmeasured direct-discharge nutrient inputs were the cause of the imbalance.

Now, if the amount of phosphorus measured leaving the lake was equivalent to the sum of the actual inputs, then Schroon Lake received a phosphorus load at a rate of 7.8 lbs/acre/yr or 0.88 g/ $\rm m^2/yr$. This rate is well above the oligotrophic rate of 0.56 g/ $\rm m^2/yr$ and, if continued, would affect the existing trophic condition of the lake.

It is apparent that a more detailed study of Schroon Lake nutrient sources is needed to provide a basis for assessment of nutrient controllability.

2. Non-point sources (see page 13)--The phosphorus exports of the Schroon Lake tributaries were relatively low and compare well with the exports of unimpacted tributaries studied elsewhere in New York (e.g., Sacandaga Reservoir tributaries*).

In all, it is estimated that non-point sources contributed 97% of the total phosphorus load to Schroon Lake during the sampling year.

^{*} Working Paper No. 167.

II. LAKE AND DRAINAGE BASIN CHARACTERISTICS

- A. Lake Morphometry[†]:
 - 1. Surface area: 4,128 acres.
 - 2. Mean depth: 47 feet.
 - 3. Maximum depth: 152 feet.
 - 4. Volume: 194,016 acre/feet.
 - 5. Mean hydraulic retention time: 153 days.
- B. Tributary and Outlet: (See Appendix A for flow data)

1. Tributaries -

	Name	Drainage area*	Mean flow*
	Schroon River Rogers Brook Horseshoe Pond Brook Unnamed Brook (F-1) Mill Brook Minor tributaries &	24.7 mi ²	257.6 cfs 13.0 cfs 3.7 cfs 3.7 cfs 36.8 cfs
	immediate drainage -	214.2 mi ²	328.1 cfs
	Totals	425.6 mi ²	642.9 cfs
2.	Outlet -		
	Schroon River	432.0 mi ² **	640.8 cfs

C. Precipitation***:

- 1. Year of sampling: 38.5 inches.
- 2. Mean annual: 37.4 inches.

[†] Aulenbach, 1973.

^{*} Drainage areas are accurate within $\pm 5\%$, except for small basins ($\pm 10\%$); mean daily flows are accurate within ± 5 to 25%; and normalized mean monthly flows are accurate within $\pm 15\%$.

^{**} Includes area of lake.

^{***} See Working Paper No. 1, "Survey Methods".

III. LAKE WATER QUALITY SUMMARY

Schroon Lake was sampled three times during the open-water season of 1972 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from two stations on the lake and from a number of depths at each station (see map, page v). During each visit a single depth-integrated (15 feet or near bottom to surface) sample was collected from the two stations for phytoplankton identification and enumeration; and during the last visit, a single fivegallon depth-integrated sample was collected for algal assays. Also each time, a depth-integrated sample was collected from each of the stations for chlorophyll \underline{a} analysis. The maximum depths sampled were 41 feet at station 1 and 32 feet at station 2.

The results obtained are presented in full in Appendix B, and the data for the fall sampling period, when the lake was essentially well-mixed, are summarized below. Note, however, the Secchi disc summary is based on all values.

For differences in the various parameters at the other sampling times, refer to Appendix B.

A. Physical and chemical characteristics:

FALL VALUES

(10/10/72)

Parameter	<u>Minimum</u>	<u>Mean</u>	<u>Median</u>	Maximum
Temperature (Cent.) Dissolved oxygen (mg/l) Conductivity (µmhos) pH (units) Alkalinity (mg/l) Total P (mg/l) Dissolved P (mg/l) NO ₂ + NO ₃ (mg/l) Ammonia (mg/l)	7.8 7.1 57 6.6 10.0 0.003 0.001 0.080 0.030	12.4 8.6 · 59 7.1 10.0 0.004 0.003 0.114 0.046	13.5 9.0 59 7.1 10.0 0.004 0.003 0.090 0.040	13.8 9.1 60 7.2 11.0 0.007 0.005 0.220 0.070
Secchi disc (inches)	118	ALL VALUES	<u>S</u> 142	180

B. Biological characteristics:

Phytoplankton -

Sampling Date		inant era	Number per ml
06/01/72	1. 2. 3. 4. 5.	Dinobryon Cyclotella Fragilaria Achnanthes Mallomonas Other genera	219 132 70 27 14 55
		Total	517
07/25/72	1. 2. 3. 4. 5.	Microcystis Chroococcus Flagellates Dinobryon Merismopedia Other genera	76 60 45 34 31 157
		Total	403
10/10/72	1. 2. 3. 4. 5.	Dinobryon Flagellates Aphanocapsa Fragilaria Tabellaria Other genera	163 139 127 87 78 223
		Total	817

2. Chlorophyll \underline{a} - (Because of instrumentation problems during the 1972 sampling, the following values may be in error by plus or minus 20 percent.)

Sampling Date	Station Number	Chlorophyll <u>a</u> (µg/l)
06/01/72	01 02	1.9 1.9
07/25/72	01 02	1.8 2.9
10/10/72	01 02	2.1 1.9

C. Limiting Nutrient Study:

1. Autoclaved, filtered, and nutrient spiked -

Spike (mg/l)	Ortho P Conc. (mg/1)	Inorganic N Conc. (mg/l)	Maximum yield (mg/l-dry wt.)	N/P <u>Ratio</u>
Control 0.010 P 0.020 P 0.050 P 0.050 P + 5.0 N 0.050 P + 10.0 N 10.0 N	0.004 0.014 0.024 0.054 0.054 0.054 0.004	0.062 0.062 0.062 0.062 5.062 10.062	0.1 0.3 0.5 0.4 13.9 14.3 0.1	16/1 4/1

2. Discussion -

The control yield of the assay alga, <u>Selenastrum capri-cornutum</u>, indicates that the potential primary productivity of Schroon Lake was very low at the time the assay sample was collected as were the levels of primary nutrients (there was some loss of inorganic nitrogen in the assay sample from the time of collection until the assay was begun; however, the results would not have differed significantly had the loss not occurred).

The N/P ratio of the control sample, and the lack of response to the nitrogen only spike, show that the control sample was phosphorus limited. However, the first phosphorus spike of only 10 µg/l more than tripled the orthophosphorus concentration in the sample and nitrogen became limiting, so there was no significant yield response to increments of phosphorus (note the N/P ratio after the first addition of phosphorus). Because of the very low levels of both nutrients, the sample essentially was co-limited, and the only significant yield response occurred when both nitrogen and phosphorus were added.

The lake data indicate phosphorus limitation at all sampling times; i.e., the N/P ratios were 36/1 or greater, and phosphorus limitation would be expected.

IV. NUTRIENT LOADINGS (See Appendix C for data)

For the determination of nutrient loadings, the New York National Guard collected monthly near-surface grab samples from each of the tributary sites indicated on the map (page v), except for the high runoff months of April and May when two samples were collected. Sampling began in November, 1972, and was completed in October, 1973.

Through an interagency agreement, stream flow estimates for the year of sampling and a "normalized" or average year were provided by the New York District Office of the U.S. Geological Survey for the tributary sites nearest the lake.

Except for Rogers Brook, nutrient loads for sampled tributaries were determined by using a modification of a U.S. Geological Survey computer program for calculating stream loadings[†]. Because of the possible effect of septic tank discharges on the nutrient concentrations measured in Rogers Brook at station D-1, the nutrient loads for this stream were calculated using the mean concentrations at station D-2 and the mean flow at station D-1.

Nutrient loadings for unsampled "minor tributaries and immediate drainage" ("ZZ" of U.S.G.S.) were estimated by using the means of the nutrient loads, in $lbs/mi^2/year$, at stations E-1, F-1, and G-1 and multiplying the means by the ZZ area in mi^2 .

⁺ See Working Paper No. 1.

During the sampling year, the City of Schroon Lake was served by septic tanks, and estimates of nutrient loadings are based on a population served of 150*. It is assumed that all of these nutrients reached the lake. A new wastewater treatment facility was expected to begin operation by the summer of 1973**.

A. Waste Sources:

Known municipal* -

Name Pop.		Treatment	Mean	Receiving		
Served			<u>Flow (mgd</u>)	<u>Water</u>		
Schroon Lake	150	Septic tanks	?	Rogers Brook		

Known industrial - None

^{*} Anonymous, 1971. ** Aulenbach, 1973.

B. Annual Total Phosphorus Loading - Average Year:

1. Inputs -

Sou	<u>rce</u>	lbs P/ yr	% of total
a.	Tributaries (non-point load)	-	
	Schroon River Rogers Brook Horseshoe Pond Brook Unnamed Brook (F-1) Mill Brook	4,980 380 80 70 910	34.5 2.6 0.6 0.5 6.3
b.	Minor tributaries & immediate drainage (non-point load) -		48.1
с.	Known municipal - None	-	-
d.	Septic tanks* -	430	3.0
e.	Known industrial - None	-	-
f.	Direct precipitation** -	640	4.4
	Total	14,420	100.0
0ut	puts -		
Lak	e outlet - Schroon River	32,290	

^{2.}

3. Net annual P loss - 17,870 pounds

^{*} Estimate based on that portion of the City of Schroon Lake population served by septic tanks (150) and 627 shoreline dwellings (U.S.G.S. map, 1953); see Working Paper No. 1.

** See Working Paper No. 1.

C. Annual Total Nitrogen Loading - Average Year:

1. Inputs -

Sou	rce	lbs N/ yr	% of total
a.	Tributaries (non-point load) -	
	Schroon River Rogers Brook Horseshoe Pond Brook Unnamed Brook (F-1) Mill Brook	556,640 28,410 6,090 5,930 91,470	41.3 1.6 0.5 0.4 6.8
b.	Minor tributaries & immedia drainage (non-point load)		45.3
c.	Known municipal - None	-	-
d.	Septic tanks* -	16,150	1.2
e.	Known industrial - None	-	-
f.	Direct precipitation** -	39,770	2.9
	Total	1,354,930	100.0
Out	puts -		

2.

Lake outlet - Schroon River 1,615,440

3. Net annual N loss - 260,510 pounds

^{*} Estimate based on that portion of the City of Schroon Lake population served by septic tanks (150) and 627 shoreline dwellings (U.S.G.S. map, 1953); see Working Paper No. 1.
** See Working Paper No. 1.

D. Mean Annual Non-point Nutrient Export by Subdrainage Area:

<u>Tributary</u>	<u>lbs P/mi²/yr</u>	lbs N/mi ² /yr
Schroon River	29	3,218
Rogers Brook	44	3,266
Horseshoe Pond Brook	32	2,456
Unnamed Brook (F-1)	28	2,391
Mill Brook	37	3,703

E. Yearly Loading Rates:

In the following table, the existing phosphorus loading rates are compared to those proposed by Vollenweider (in press). Essentially, his "dangerous" rate is the rate at which the receiving waters would become eutrophic or remain eutrophic; his "permissible" rate is that which would result in the receiving water remaining oligotrophic or becoming oligotrophic if morphometry permitted. A mesotrophic rate would be considered one between "dangerous and "permissible".

	Tota	1 Phosphorus	Total Nitrogen		
Units	Total	Accumulated	Total	Accumulated	
lbs/acre/yr	3.5	loss*	328.2	loss*	
lbs/acre/yr grams/m²/yr	0.39	-	36.8	-	

Vollenweider loading rates for phosphorus (g/m²/yr) based on mean depth and mean hydraulic retention time of Schroon Lake:

"Dangerous" (eutrophic rate) 1.12
"Permissible" (oligotrophic rate) 0.56

^{*} An apparent loss of some 17,870 lbs of total phosphorus and 260,510 lbs of total nitrogen occurred during the sampling year. While nitrogen loss can occur with certain conditions, losses of both nutrients at that magnitude makes it almost certain that unknown and unmeasured nutrient inputs were involved.

V. LITERATURE REVIEWED

- Anonymous, 1971. Inventory of municipal waste facilities. Publ. No. OWP-1, Vol. 2, EPA, Washington, D.C.
- Aulenbach, Donald B., 1973. National Eutrophication Survey questionnaire. Rensselaer Polytechnic Inst., Troy.
- Vollenweider, Richard A. (in press). Input-output models. Schweiz. A. Hydrol.

VII. APPENDICES

APPENDIX A

TRIBUTARY FLOW DATA

LAKE CODE 3634 SCHROON LAKE

TOTAL DRAINAGE AREA OF LAKE 432.00

Su	JB-DRAINAGE						NOR	MAL I ZED	FLOWS					
TRIBUTARY	AREA	MAL	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
3634A1	173.00	181.00	170.00	319.00	888 -00	509.00	244.00	119.00	70.90	79.20	119.00	191.00	204.00	257.55
3634A2	432.00	459.00	390.00	810.00	2180.00	1290.00	599.00	301.00	180.00	195.00	302.00	470.00	518.00	640.82
3634D1	8.72	9.11	8.58	16.10	44.80	25.70	12.30	5.97	3.57	3.99	5.99	9,63	10.30	12.99
3634E1	2.48	2.59	2.44	4.57	12.70	7.30	3.49	1.70	1.02	1.14	1.70	2.74	2.92	3.69
3634F1	2.48	2.59	2.44	4.57	12.70	7.30	3.49	1.70	1.02	1.14	1.70	2.74	2.92	3.69
363461	24.70	25.80	24.30	45.50	127.00	. 72.70	34.80	16.90	10.10	11.30	17.10	27.30	29.10	36.78
363477	550.65	230.00	217.00	406.00	1130.00	649.00	311.00	151.00	90.50	101.00	152.00	244.00	260.00	328.08

			_	
21	IMN	l A	D	v

TOTAL DRAINAGE AREA OF LAKE	=	432.00	TOTAL FLOW IN =	7722.15
SUM OF SUB-DRAINAGE AREAS	=	432.00	TOTAL FLOW OUT =	7694.00

MFAN	MONTHI Y	FLOWS	ΔND	DATLY	FLOWS

TRIBUTARY	нтиом	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
3634A1	11	72	276.00	5	908.00				
	12	72	218.00	5 3	450.00				
	1	73	348.00	7	600.00				
	2	73	315.00	25	200.00				
	2	73	845.00						
	4	73	884.00	8	830.00	23	1440.00		
	4 5	73	540.00	6	452.00	26	655.00		
	6	73	247.00	9	312.00		•		
	7	73	168.00	10	190.00				
	8	73	62.70	3	206.00	31	72.50		
	8 9	73	72.70						
	10	73	45.00						
3634A2	11	72	1190.00						
	12	72	711.00						
	1	73	1010.00	7	882.00	28	1170.00		
	2	73	684.00	25	574.00				
	3	73	2010.00						
		73	2070.00	8	3020.00	23	1620.00		
	4 5	73	1520.00	6	1110.00	26	2080.00		
	6	73	1050.00	9	1400.00				
	7	73	565.00	10	733.00				
	8	73	217.00	3	296.00	31	170.00		
	9	73	214.00	29	252.00				
	10	73	208.00						

LAKE CODE 3634

SCHROON LAKE

MEAN MONTHLY FLOWS AND DAILY FLOWS

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
3634D1	11	72	13.90	5	57.10				
	12	72	1.00	3	28.20				
	1	73	17.50	7	38.00				
	2	73	15.90	25	12.00				
	3	73	42.60						
	4	73	44.60	8	41.00	23	63.00		
	5	73	27.30	6	25.50	26	34.00		
	6	73	12.40	9	16.70				
	7	73	8.43	10	9.90				
	8 9	73 73	3.16	3	9.60	31	1.70		
	10	73	3.66 2.26						
3634E1	11	72	3.96	5	15.80				
333761	12	72	3.10	3	7.80				
	1	73	4.98	7	10.50				
	ş	73	4.52	25	3.40				
	3	73	12.19		5440				
	4	73	12.60	8	9.15	23	14.20		
	5	73	7.75	6	5.75	26	7.60		
•	6	73	3.53	9	3.70				
	7	73	2.40	10	2.08				
	8	73	0.90	3	1.62	31	0.30		
	9	73	1.05						
	10	73	0.64						
3634F1	11	72	3.96	5	17.60				
	12	72	3.10	3	8.75				
	1	73	4.98	7	11.70				
	2	73	4.52	25	3.80				
	3	73 73	12.10		0.40	22	14 00		
	4 5	73 73	12.60 7.75	8 6	9.60	23	16.80		
	6	73	3.53	9	5.28 2.80	26	7.56		
	7	73	2.40	10	0.80				
	8	73	0.90	3	1.46	31	0.21		
	9	73	1.05	•	1440	<i>J</i> .	0.21		
	1ó	73	0.64						
3634G1	ii	72	39.40	5	145.00				
	12	72	31.10	3	71.20				
	1	73	49.60	7	96.00				
	2	73	45.10	25	30.20				
	3	73	121.00						
	4	73	126.00	8	122.00	23	218.00		
	5	73	77.10	6	64.00	26	94.00		
	6	73	35.20	9	41.50				
	7	73	23.90	10	21.80				•
	8	73	8.94	3	26.20	31	6.70		
	9	73	10.40						
	10	73	6.46						

TRIBUTARY FLOW INFORMATION FOR NEW YORK

11/26/74

LAKE CODE 3634 SCHROON LAKE

MEAN MONTHLY FLOWS AND DAILY FLOWS

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
3634ZZ	11	72	352.00	5	678.00				
	15	72	278.00	3	336.00				
	1	73	442.00	7	450.00				
	2	73	402.00	25	144.00				
	3	73	1080.00						
	4	73	130.00	8	1140.00	23	2060.00		
	5	73	689.00	6	568.00	26	865.00		
	6	73	315.00	Ğ	364.00				
	7	73	213.00	10	178.00				•
	8	73	80.10	3	244.09	31	79.30		
	9	73	92.70	3	2,1,00	J.	, , , ,		
	10	73	57.40						

APPENDIX B

PHYSICAL and CHEMICAL DATA

363401 43 45 48.0 073 45 48.0 SCHROON LAKE 36 NEW YORK

						11EP	ALES		1202 FEET DEP	тн	
DATE FROM	TIME DEPTH	00010 WATER TEMP	00300 DO	00077 TRANSP SECCHI	00094 CNDUCTVY FIELD	00400 PH	00410 T ALK CACO3	00630 N028N03 N-TOTAL	00610 NH3-N TOTAL	00665 PHOS-TOT	00666 PHOS-DIS
TO	DAY FEET	CENT	MG/L	INCHES	MICROMHO	SU	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/06/01	17 35 0000	18.8	9.0	118	30	6.70	10	0.140	0.010K	0.003	0.002
	17 35 0010	17.9	9.4		50	7.00	11	0.140	0.010	0.003	0.003
	17 35 0037	5.5	10.9		50	6.90	10K	0.200	0.010	0.004	0.002
72/07/25	16 40 0000			140	55	7.50	15	0.090	0.050	0.007	0.004
	16 40 0004	25.0	8.5		55	7.50	14	0.100	0.050	0.007	0.006
	16 40 0015	21.9	8.4		50K	7.30	13	0.120	0.060	0.007	0.006
	16 40 0025	11.2	9.2		50K	7.20	11	0.180	0.060	0.006	0.004
	16 40 0035	7.2			50K	7.20	10	0.220	0.060	0.005	0.004
72/10/10	15 30 0000			180	60	7.10	10	0.080	0.040	0.004	0.004
	15 30 0004	13.8	9.0		60	7.10	11	0.090	0.070	0.004	0.004
	15 30 0015	13.5	9.0		59	7.10	11	0.080	0.040	0.004	0.003
	15 30 0030	11.0	7.6		59	7.10	10K		0.050	0.004	0.003
	15 30 0041	7.8	7.1		57	6.65	19K	0.220	0.030	0.007	0.003

				32217
DATE	TI	1E 1	DEPTH	CHESPHAF
FROM	OF	=		Δ
. 10	DAY	1 (FEET	UGZL
72/06/01	17	35	0000	1.9
72/07/25	16	40	Ú000	1.80
72/10/10	15	30	0000	2.1

K VALUE KNOWN TO BE LESS THAN INDICATED

J VALUE KNOWN TO BE IN ERROR

363402 43 49 54.0 073 47 00.0 SCHROON LAKE 36 NEW YORK

							11EP 3	ALES		1202 FEET DEP	тн	
DATE FROM TO	TIME OF Day	DEPTH FEET	00010 WATER TEMP CENT	00300 DO MG/L	00077 TRANSP SECCHI INCHES	00094 CNDUCTVY FIELD MICROMHO	00400 PH	00410 T ALK CACO3	00630 N026N03 N-TOTAL	00610 NH3-N TOTAL	00665 PH0S-T0T	00666 PHOS-DIS
10	DAT	FEET	CENT	MIT	TMCHES	WICKOWHO	SU	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/06/01	18 2	0 0000	17.7	9.5	132	30	6.60	106	0.160	0.020	0.002	0.002K
	18 2	0 0010	16.0	10.0		30	6.60	10K	0.160	0.030	0.004	0.002K
	18 20	0 0032	5.4	11.2		20	6.90	10	0.180	0.010	0.002	0.002
72/07/25	17 19	5 0000			144	55	7.50	108	0.090	0.060	0.006	0.005
	17 1	5 0004	25.6	8.4		55	7.50	15	0.080	0.030	0.006	0.005
	17 19	5 0014	51.0	8.8		55	7.50	14	0.110	0.060	0.007	0.004
72/10/10	16 0	0 0000			168	59	7.20	10	0.090	0.050	0.006	0.005
	16 0	0 0004	13.8	9.1		58	7.15	11	0.080	0.040	0.003	0.001
	16 0	0 0015	13.7	9.1		58	7.10	10K	0.140	0.040	0.003	0.001
	16 0	0 0024	13.3	9.1		59	7.10	10	0.080	0.050	0.003	0.003

				32217
DATE	TIN	4E	DEPTH	CHLRPHYL
FROM	OF	-		Δ
TO	DAY	1	FEET	UGZL
72/06/01	18	20	0000	1.9J
72/07/25	17	15	0000	2.9J
72/10/10	16	0.0	0000	1.9J

K VALUE KNOWN TO BE LESS THAN INDICATED

J VALUE KNOWN TO BE IN ERROR

APPENDIX C

TRIBUTARY DATA

3634A1 LS3634A1
43 52 30.0 073 44 30.0
SCHROON RIVER
36 15 PARADOX LAKE
I/SCHROON LAKE
ST HWY 73 CULVERT
11EPALES 2111204
4 0000 FEET DEPTH

			00630	00625	00610	00671	00665
DATE	TIME	DEPTH	N028N03	TOT KUEL	NH3-N	PHOS-DIS	PHOS-TOT
FROM	OF		N-TOTAL	N	TOTAL	ORTHO	
TO	ÛAY	FEET	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/11/05	11 49	5	0.030	0.230	0.039	0.005K	0.006
72/12/03			0.033	0.210	0.011	0.005K	0.005K
73/01/07			0.231	0.170	0.026	0.005K	0.005K
73/02/25			0.028	1.320	0.110	0.007	0.020
73/04/08			C.100	2.100	0.200	0.005K	0.005K
73/04/23			0.056	1.200	0.040	0.005K	0.005K
73/05/06			0.031	1.400	0.03P		0.015
73/05/26			0.044	2.600	0.024	0.010	0.015
73/06/09			0.069	0.300	0.018	0.008	0.015
73/07/10			0.010K	0.830	0.033	0.005K	0.005K
73/08/03			0.139	0.810	0.008	0.005K	0.020
73/08/31			0.016	2.400	0.069	0.005K	0.005K
73/09/29			0.021	0.100K	0.036	0.005K	0.010

K VALUE KNOWN TO BE LESS THAN INDITATED

3634A2 LS3634A2
43 43 00.0 073 48 30.0
SCHROON RIVER
36 15 NORTH CREEK
0/SCHROON LAKE
ALONG BANK FROM LGT DUTY RD
11EPALES 2111204
4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 NOZKNO3 N-TOTAL MG/L	00625 TOT KJEL N MG/I	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
72/11/05	10 3	0	0.140	0.600	0.160	0.006	0.039
73/02/25			0.018	0.540	0.026	0.005K	0.005K
73/04/08			0.210	4.000	0.160	0.005K	0.010
73/04/23			0.074	0.910	0.021	0.005K	0.015
73/05/06			0.060	0.920	0.157	0.012	0.030
73/05/26			0.056	1.890	0.068	0.005K	0.055
73/06/09			0.067	0.460	0.031	0.020	0.035
73/07/10			0.018	1.050	0.075	0.00A	0.025
73/08/03			0.063	0.400	0.032	0.005K	0.025
73/08/31			0.294	2.000	0.100		
73/09/29			0.019	0.340	0.058	0.005K	0.015

K VALUE ENGUN TO HE LESS THAN INDICATED

363481 LS363481 43 52 30.0 073 44 00.0 UNNAMED BROOK 36 15 PARADOX LAKE T/SCHROON LAKE ST HWY 73 BRDG 11EPALES 2111204 4 0000 FEET DEPTH

K VALUE A 40 W TO HE LESS THAN IMUTCATED

3634C1 L53634C1 43 51 30.0 073 43 30.0 ALDER CREEK 36 15 PARADOX LAKE T/SCHROON LAKE BRDG S OF OWLS HEAD MTN 11EPALES 2111204 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 NO28NO3 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3+N TOTAL MG/L	00671 PHOS-OIS OPTHO MG/L P	00665 PHOS-TOT MG/L P
72/11/05	11 30	0	0.052	0.310	0.04P	0.005K	0.012
72/12/03			0.048	0.210	0.005K	0.005K	0.005K
73/01/27			0.085	0.190	0.020	0.006	0.006
73/04/08			0.054	2.200	0.176	0.005K	0.005K
73/04/23			0.020	0.660	0.021	0.005K	0.005K
73/05/06			0.030	0.600	0.021	0.005K	0.015
73/05/26			0.048	1.400	0.020	0.005K	0.015
73/06/09			0.056	0.420	0.022	0.006	0.015
73/07/10			0.033	0.560	0.022	0.006	0.017
73/08/03			0.080	0.630	0.080	0.005K	0.030
73/09/31			0.025	0.230	0.038	0.006	0.015
73/09/29			0.021	0.295	0.033	0.005K	0.010

K VALUE KNOWN TO BE LESS THAN IMDICATED

363401 LS3634D1
43 50 00.0 073 45 30.0
ROGERS BROOK
36 15 SCHROON LAKE
T/SCHROON LAKE
FROM BANK IN SCHROON LAKE BELO STP
11EPALES 2111204
4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 NO25NO3 N-TOTAL MG/L	00625 TOT KJEL M MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
72/11/05			0.085	0.360	0.096	0.005K	0.013
72/12/03			0.140	0.290	0.016	0.006	0.006
73/01/07			0.198	0.240	0.025	0.005K	0.010
73/01/27			0.290	0.250	0.035	0.007	0.015
73/02/25			0.010K	0.130	0.018	0.005K	0.007
73/04/08			0.198	1.760	0.147	0.005K	0.010
73/04/23			0.138	1.260	0.029	0.007	0.025
73/05/06			0.066	1.100	0.026	0.006	0.025
73/05/26			0.058	1.890	0.037	0.012	0.050
73/06/09			0.066	0.460	0.030	0.006	0.015
73/07/10			0.130	0.520	0.022	0.006	0.020
73/08/03			0.052	0.500	0.154	0.005K	0.010
73/09/31			0.320	1.320	0.555	0.006	0.055
73/09/29			9.110	0.520	0.053	0.005K	0.010

K VALUE KNOWN TO BE LESS THAN INDICATED

 $(x_{i_1}, \dots, x_{i_{m-1}})^{-1}$

363492 LS3634D2
43 50 00.0 073 46 30.0
ROGERS BROOK
36 15 SCHROON LAKE
T/SCHROON LAKE
BROG .5 MI W OF SCHROON LAKE ABOV STP
11EPALES 2111204
4 0000 FEET DEPTH

DATE FROM	TIME OF	ОЕРТН	00630 NO?&NO3 N-TOTAL	00625 TOT KUEL N	00610 NH3-N TOTAL	00671 PHOS-DIS ORTHO	00665 PHOS-TOT
T 0	DAY	FEET	MG/L	MG/L	MG/L .	MG/L P	MG/L P
72/11/05			0.052	0.390	. 0.044	0.005K	0.011
73/01/07			0.170	0.220	0.028	0.005K	0.010
73/02/25			0.016	0.110	0.021	0.005K	0.005K
73/04/08			0.198	2.900	0.160	0.005K	0.005K
73/04/23			0・13お	1.540	0.034	0.005K	0.005K
73/05/06			0.054	1.320	0.054	0.005K	0.010
73/05/26			0.05?	2.400	0.033	0.010	•
73/06/09			0.066	0.400	0.015	0.005K	0.015
73/07/10			0.110	0.680	0.026	0.005	0.015
73/08/03			0.028	0.600	0.032	0.005K	0.030
73/08/31			0.378	0.580	0.046	0.010	0.050
73/09/29			0.058	0.860	0.067	0.005K	0.010

K VALUE KNOWN TO HE LESS THAN INDICATED

3634E1 LS3634E1
43 50 00.0 073 46 30.0
HORSESHOE POND BROOK
36 15 SCHROON LAKE
T/SCHROON LAKE
US HWY 9 BRDG IN SCHROON LAKE
11EPALES 2111204
4 0000 FEET DEPTH

DATE FROM	TIME OF	DEPTH	00630 NO350N N-TOTAL	00625 TOT KJEL N	00610 NH3-N Total	00671 PHOS-DIS ORTHO	00665 PHOS-TOT
10	DAY	FEET	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/11/05	11 1	5	0.110	0.360	0.065	0.005K	0.017
72/12/03			0.147	0.280	0.013	0.006	0.009
73/01/07			0.094	0.180	0.018	0.005K	0.005K
73/01/27			0.110	0.210	0.025	0.007	0.007
73/02/25			0.017	3.400	0.029	0.009	
73/04/08			0.042	2.400	0.200	0.005K	0.005K
73/04/23			0.034	0.700	0.019	0.005K	0.005K
73/05/06			0.032	0.630	0.037	0.005K	0.010
73/05/26			0.044	0.520	0.035	0.006	
73/06/09			0.064	0.380	0.016	0.007	0.020
73/07/10			0.115	0.570	0.056	0.005K	0.010
73/08/03			0.088	C.750	0.110	0.005K	0.015
73/08/31			0.220	2.600	0.900	0.006	0.035
73/09/29			0.080	0.920	0.790	0.005K	0.005K

K VALUE KNOWN TO HE LESS THAN INDICATED

3634F1 LS3634F1
43 46 30.0 073 47 30.0
UNNAMED BROOK
36 15 SCHROON LAKE
T/SCHROON LAKE
RD BRDG JOINING US HWY 9
11EPALES 2111204
4 0000 FEET DEPTH

			00630	00625	00610	00671	00665
DATE	TIME	DEPTH	N054N03	TOT KJEL	NH3-N	PHOS-DIS	PHOS-TOT
FRUM	OF		N-TOTAL	N	TOTAL	ORTHO	
ΤΟ.	DAY	FEET	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/11/05	11 2	5	0.057	0.240	0.048	0.005K	0.011
72/12/03			0.069	0.250	0.022	0.005K	0.005K
73/01/07			0.077	0.270	0.058	0.005K	0.005K
73/01/27			0.069	1.500	0.150	0.006	0.010
73/02/25			0.015	0.560	0.042	0.005K	0.005K
73/04/08			0.036	1.760	0.105	0.005K	0.010
73/04/23			0.074		0.072	0.005K	0.010
73/05/06	•		0.039	1.200	0.027	0.005K	0.020
73/05/26			0.048	0.260	0.008	0.006	0.010
73/06/09			0.063	0.360	0.018	0.005K	0.015
73/07/10			0.138	0.770	0.046	0.006	0.013
73/08/03			0.078	1.050	0.330	0.005K	0.015
73/08/31			0.198	1.095	0.126	0.005K	0.010
73/09/29			0.043	0.520	0.115	0.005K	0.005K

K VALUE KNOWN TO BE LESS THAN INDICATED

3634G1 LS3634G1 43 46 00.0 073 45 30.0 MILL BROOK 36 15 SCHROON LAKE T/SCHROON LAKE BRDG IN ADIRONDACK 11EPALES 2111204 4 0000 FEET DEPTH

2475	T 1145	DEDIM	00630	00625	00610	00671	00665
DATE		DEPTH		TOT KJEL	NH3-N	PHOS-DIS	PHOS-TOT
FROM	0F		N-TOTAL	N	TOTAL	ORTHO	
10	DAY	FEET	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/11/05	10 49	5	0.060	0.440	0.063	0.005K	0.010
72/12/03			0.048	0.150	0.009	0.005K	0.015
73/01/07			0.084	0.240	0.037	0.005K	0.005K
73/01/27			9.078	0.180	0.028	0.005K	0.010
73/04/08			0.052	4.100	0.252	0.005K	0.015
73/04/23			0.052	3.500	0.132	0.005K	0.005K
73/05/06			0.i15	1.000	0.020	0.005K	0.010
73/05/26			0.052	1.800	0.037	0.006	
73/06/09			0.063	0.430	0.012	0.005K	0.015
73/07/10			0.078	1.050	0.044	0.005K	0.020
73/08/03			0.078	1.700	0.330	0.005K	0.030
73/08/31			0.054	0.260	0.009	0.005K	0.015
73/09/29			0.050	1.100	0.027	0.005K	0.005K

K VALUE KNOWN TO BE LESS THAN INDICATED

3634H1 LS3634H1
43 43 30.0 073 49 30.0
TROUT BROOK
36 15 NORTH CHEEK
T/SCHROON LAKE
US HWY 9 BROG
11EPALES 2111204
4 0000 FEET DEPTH

DATE FROM	TIME OF	DEPTH	00630 NO2&NO3 N-TOTAL	00625 TOT KJEL N	00610 NH3-N TOTAL	00671 PHOS-DIS OPTHO	00665 PHOS-TOT
TO	DAY	FEET	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/11/05	10 3	5	0.211	0.340	0.058	0.005K	0.007
72/12/03			0.190	0.240	0.008	0.005K	0.006
73/01/07			0.189	0.300	0.058	0.005K	0.010
73/04/08			0.260	3.020	0.430	0.005K	0.010
73/04/23			0.210	1.050	0.030	0.005K	0.010
73/05/06			0.126	1.150	0.034	0.005K	0.015
73/05/26			0.052	0.940	0.017	0.005K	0.025
73/06/09			0.0 68	1.470	0.031	0.005K	0.020
73/07/10			0.066	0.800	0.044	0.005K	0.015
73/08/31			0.180	1.260	0.280	0.005K	0.015
73/09/29			0.075	0.460	0.024	0.005K	0.025

K VALUE KNOWN TO BE LESS THAN INDICATED