

**U.S. ENVIRONMENTAL PROTECTION AGENCY
NATIONAL EUTROPHICATION SURVEY**

WORKING PAPER SERIES



REPORT
ON
LAKE HAINES
POLK COUNTY
FLORIDA
EPA REGION IV
WORKING PAPER No. 255

**CORVALLIS ENVIRONMENTAL RESEARCH LABORATORY - CORVALLIS, OREGON
and
ENVIRONMENTAL MONITORING & SUPPORT LABORATORY - LAS VEGAS, NEVADA**

C

REPORT
ON
LAKE HAINES
POLK COUNTY
FLORIDA
EPA REGION IV
WORKING PAPER No. 255

WITH THE COOPERATION OF THE
FLORIDA DEPARTMENT OF ENVIRONMENTAL REGULATION
AND THE
FLORIDA NATIONAL GUARD
DECEMBER, 1977

CONTENTS

	<u>Page</u>
<u>Foreword</u>	ii
List of Florida Lakes	iv
Lake and Drainage Area Map	v
 <u>Sections</u>	
I. Conclusions	1
II. Lake and Drainage Basin Characteristics	3
III. Lake Water Quality Summary	4
IV. Nutrient Loadings	8
V. Literature Reviewed	12
VI. Appendices	13

F O R E W O R D

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nationwide threat of accelerated eutrophication to freshwater lakes and reservoirs.

OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

- a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.
- b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.
- c. With such a transformation, an assessment of the potential for eutrophication control can be made.

LAKE ANALYSIS

In this report, the first stage of evaluation of lake and watershed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning [§303(e)], water quality criteria/standards review [§303(c)], clean lakes [§314(a,b)], and water quality monitoring [§106 and §305(b)] activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

ACKNOWLEDGMENT

The staff of the National Eutrophication Survey (Office of Research & Development, U. S. Environmental Protection Agency) expresses sincere appreciation to the Florida Department of Environmental Regulation for professional involvement and to the Florida National Guard for conducting the tributary sampling phase of the Survey.

Joseph W. Landers, Jr., Secretary of the Department of Environmental Regulation; John A Redmond, former Director of the Division of Planning, Technical Assistance, and Grants; and Dr. Tim S. Stuart, Chief of the Bureau of Water Quality, provided invaluable lake documentation and counsel during the survey, reviewed the preliminary reports, and provided critiques most useful in the preparation of this Working Paper series.


Major General Henry W. McMillan (Retired), then the Adjutant General of Florida, and Project Officer Colonel Hugo F. Windham, who directed the volunteer efforts of the Florida National Guard, are also gratefully acknowledged for their assistance to the Survey.

NATIONAL EUTROPHICATION SURVEY

STUDY LAKES

STATE OF FLORIDA

<u>LAKE NAME</u>	<u>COUNTY</u>
Alligator	Columbia
Apopka	Lake, Orange
Banana	Polk
Crescent	Flagler, Putnam
Doctors	Clay
Dora	Lake
East Tohopekaliga	Osceola
Effie	Polk
Eloise	Polk
George	Putnam, Volusia
Gibson	Polk
Glenada	Highlands
Griffin	Lake
Haines	Polk
Hancock	Polk
Horseshoe	Seminole
Howell	Orange, Seminole
Istokpoga	Highlands
Jessie	Polk
Jessup	Seminole
Kissimmee	Osceola
Lawne	Orange
Lulu	Polk
Marion	Polk
Minnehaha	Orange
Minneola	Lake
Monroe	Seminole, Volusia
Munson	Leon
Okeechobee	Glades, Hendry, Martin, Okeechobee, Palm Beach
Poinsett	Brevard, Orange, Osceola
Reedy	Polk
Seminole	Jackson, FL; Decatur, Seminole, GA
Seminole	Pinellas
South	Brevard
Talquin	Gadsden, Leon
Tarpon	Pinellas
Thonotosassa	Hillsborough
Tohopekaliga	Osceola
Trout	Lake
Weohyakapka	Polk
Yale	Lake



LAKE HAINES
STORET NO. 1215

I. CONCLUSIONS

A. Trophic Condition:

Survey data indicate that Lake Haines is eutrophic. It ranked 15th in overall trophic quality when the 41 Florida lakes sampled in 1973 were compared using a combination of six parameters*. Eleven of the lakes had less and one had the same median total phosphorus and median dissolved phosphorus, eight had less and one had the same median inorganic nitrogen, 17 had less mean chlorophyll a, and 29 had greater mean Secchi disc transparency.

B. Rate-Limiting Nutrient:

The assay results indicate Lake Haines was limited by nitrogen at the time the sample was collected (03/08/73). The lake data indicate nitrogen limitation at all sampling times.

C. Nutrient Controllability:

1. Point sources--It is estimated that the Lake Alfred wastewater treatment plant accounted for 76.5% of the total phosphorus and 4.9% of the total nitrogen inputs to Lake Haines during the sampling year.

The estimated sampling year phosphorus loading of 0.68 g/m² is a little over three times that proposed by Vollenweider (Vollenweider and Dillon, 1974) as a eutrophic loading; and even though

* See Appendix A.

the critical loading for Florida lakes may be somewhat higher than that suggested by Vollenweider (see page 11), the trophic condition of the lake is indicative of excessive nutrient loads. It is calculated that diversion of the Lake Alfred treatment plant effluent would reduce the phosphorus loading to $0.16 \text{ g/m}^2/\text{yr}$ (a mesotrophic loading).

2. Non-point sources--It is estimated that non-point sources, including precipitation, contributed approximately 23% of the total phosphorus and 94% of the total nitrogen inputs to Lake Haines.

The 1970 Winter Haven U.S.G.S. quadrangle map indicates that land use around the lake is mainly agricultural (citrus) and marsh.

II. LAKE AND DRAINAGE BASIN CHARACTERISTICS[†]A. Morphometry^{††}:

1. Surface area: 2.90 kilometers².
2. Mean depth: 2.5 meters.
3. Maximum depth: 4.3 meters.
4. Volume: 7.250×10^6 m³.
5. Mean hydraulic retention time: 2.9 years.

B. Tributary and Outlet:
(See Appendix C for flow data)

1. Tributaries -

<u>Name</u>	<u>Drainage area (km²)^{†††}</u>	<u>Mean flow (m³/sec)</u>
Minor tributaries & immediate drainage -	<u>18.3</u>	<u>0.08</u>
Totals	18.3	0.08*

2. Outlet -

Unnamed Canal A-1	21.2**	0.08
-------------------	--------	------

C. Precipitation^{***}:

1. Year of sampling: 112.8 centimeters.
2. Mean annual: 134.1 centimeters.

† Table of metric conversions--Appendix B.

†† Hulbert, 1973; depths estimated from soundings reported in Appendix D.

††† For limits of accuracy, see Working Paper No. 175, "...Survey Methods, 1973-1976".

* Based on net effect of factors affecting flows in outlet canal during period of October, 1960, through September, 1968 (Anderson, 1974).

** Includes area of lake.

*** See Working Paper No. 175.

III. WATER QUALITY SUMMARY

Lake Haines was sampled three times during 1973 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from two or more depths at one station on the lake (see map, page v). During each visit, a single depth-integrated (near bottom to surface) sample was collected for phytoplankton identification and enumeration, and a similar sample was collected for chlorophyll a analysis. During the first visit, a single 18.9-liter depth-integrated sample was taken for algal assays. The maximum depth sampled was 3.7 meters.

The sampling results are presented in full in Appendix D and are summarized in the following table.

A. SUMMARY OF PHYSICAL AND CHEMICAL CHARACTERISTICS FOR LAKE HAINES
STORET CODE 1215

1ST SAMPLING (3/ 8/73)

2ND SAMPLING (9/ 6/73)

3RD SAMPLING (11/ 7/73)

1 SITES

1 SITES

1 SITES

PARAMETER	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN
TEMP (C)	17.6 - 23.2	21.1	22.4	27.8 - 28.8	28.3	28.3	23.1 - 23.2	23.1	23.1
DISS OXY (MG/L)	4.8 - 8.9	6.8	6.8	4.4 - 6.2	5.3	5.3	7.0 - 7.2	7.1	7.1
CNDCTVY (MCROMU)	205. - 220.	213.	215.	218. - 221.	220.	220.	189. - 190.	190.	190.
PH (STAND UNITS)	8.2 - 9.1	8.6	8.6	6.6 - 7.3	6.9	6.9	6.8 - 7.0	6.9	6.9
TOT ALK (MG/L)	10. - 12.	11.	11.	15. - 17.	16.	16.	14. - 15.	14.	14.
TOT P (MG/L)	0.053 - 0.074	0.060	0.054	0.044 - 0.054	0.049	0.049	0.072 - 0.078	0.074	0.072
ORTHO P (MG/L)	0.009 - 0.030	0.016	0.010	0.008 - 0.014	0.011	0.011	0.014 - 0.015	0.015	0.015
NO2+NO3 (MG/L)	0.050 - 0.140	0.087	0.070	0.060 - 0.060	0.060	0.060	0.030 - 0.040	0.037	0.040
AMMONIA (MG/L)	0.060 - 0.120	0.080	0.060	0.060 - 0.080	0.070	0.070	0.050 - 0.050	0.050	0.050
KJEL N (MG/L)	0.700 - 1.000	0.833	0.800	0.900 - 1.400	1.150	1.150	1.100 - 1.300	1.200	1.200
INORG N (MG/L)	0.110 - 0.260	0.167	0.130	0.120 - 0.140	0.130	0.130	0.080 - 0.090	0.087	0.090
TOTAL N (MG/L)	0.770 - 1.050	0.920	0.940	0.960 - 1.460	1.210	1.210	1.130 - 1.340	1.237	1.240
CHLRPYL A (UG/L)	37.3 - 37.3	37.3	37.3	25.3 - 25.3	25.3	25.3	17.1 - 17.1	17.1	17.1
SECCHI (METERS)	0.8 - 0.8	0.8	0.8	1.2 - 1.2	1.2	1.2	0.9 - 0.9	0.9	0.9

B. Biological characteristics:

1. Phytoplankton -

<u>Sampling Date</u>	<u>Dominant Genera</u>	<u>Algal Units per ml</u>
03/08/73	1. <u>Microcystis sp.</u>	1,566
	2. <u>Anabaena sp.</u>	944
	3. <u>Synedra sp.</u>	693
	4. <u>Kirchneriella sp.</u>	572
	5. <u>Melosira sp.</u>	482
	Other genera	<u>2,881</u>
	Total	7,138
09/06/73	1. <u>Melosira sp.</u>	3,816
	2. <u>Merismopedia sp.</u>	2,589
	3. <u>Lyngbya sp.</u>	1,567
	4. Flagellates	1,431
	5. <u>Dactylococcopsis sp.</u>	1,295
	Other genera	<u>8,450</u>
	Total	19,148
11/07/73	1. <u>Lyngbya sp.</u>	13,689
	2. Centric diatoms	5,513
	3. Flagellates	2,661
	4. <u>Chroococcus sp.</u>	2,567
	5. <u>Dactylococcopsis sp.</u>	2,091
	Other genera	<u>7,895</u>
	Total	34,416

2. Chlorophyll a

<u>Sampling Date</u>	<u>Station Number</u>	<u>Chlorophyll a (μg/l)</u>
03/08/73	1	37.3
09/06/73	1	25.3
11/07/73	1	17.1

C. Limiting Nutrient Study:


1. Autoclaved, filtered, and nutrient spiked -

<u>Spike (mg/l)</u>	<u>Ortho P Conc. (mg/l)</u>	<u>Inorganic N Conc. (mg/l)</u>	<u>Maximum yield (mg/l-dry wt.)</u>
Control	0.028	0.152	4.1
0.050 P	0.078	0.152	4.2
0.050 P + 1.0 N	0.078	1.152	17.4
1.0 N	0.028	1.152	8.2

2. Discussion -

The control yield of the assay alga, Selenastrum capricornutum, indicates that the potential primary productivity of Lake Haines was moderately high at the time the sample was collected (03/08/73). The results also indicate nitrogen limitation. Note that the addition of orthophosphorus alone did not result in an increase in yield, but the addition of nitrogen alone resulted in a yield double that of the control.

The lake data substantiate nitrogen limitation at the time the sample was collected; i.e., the mean inorganic nitrogen/orthophosphorus ratio was 10/1. Mean N/P ratios in September and November were 12/1 and 6/1, respectively.



IV. NUTRIENT LOADINGS (See Appendix E for data)

For the determination of outlet nutrient loadings, the Florida National Guard collected monthly near-surface grab samples from the outlet site indicated on the map (page v). Sampling was begun in March, 1973, and was completed in February, 1974.

Through an interagency agreement, flow estimates for the year of sampling and a "normalized" or average year were provided by the Florida District Office of the U.S. Geological Survey for the outlet canal.

In this report, nutrient loads for the outlet canal were determined by using a modification of a U.S. Geological Survey computer program for calculating stream loadings*. Nutrient loads for unsampled "minor tributaries and immediate drainage" ("ZZ" of U.S.G.S.) were estimated using the mean nutrient loads from citrus groves, in kg/km²/year (Brezonik and Shannon, 1971), and multiplying by the ZZ area in km².

The operator of the Lake Alfred wastewater treatment plant provided monthly effluent samples and corresponding flow data.

* See Working Paper No. 175.

A. Waste Sources:

1. Known municipal -

<u>Name</u>	<u>Pop. Served</u>	<u>Treatment</u>	<u>Mean Flow (m³/d)</u>	<u>Receiving Water</u>
Lake Alfred	670	stabilization pond	656.3	Lake Haines

2. Known industrial - None

B. Annual Total Phosphorus Loading - Average Year:

1. Inputs -

<u>Source</u>	<u>kg P/ yr</u>	<u>% of total</u>
a. Tributaries (non-point load) -		
None	-	-
b. Minor tributaries & immediate drainage (non-point load) -	330*	16.7
c. Known municipal STP's -		
Lake Alfred	1,515	76.5
d. Septic tanks** -	5	0.2
e. Known industrial - None	-	-
f. Direct precipitation* -	<u>130</u>	<u>6.6</u>
Total	1,980	100.0

2. Outputs -

Lake outlet - Unnamed Canal A-1 165

3. Net annual P accumulation - 1,815 kg.

* Brezonik and Shannon, 1971.

** Estimate based on 24 lakeshore dwellings; see Working Paper No. 175.

C. Annual Total Nitrogen Loading - Average Year:

1. Inputs -

<u>Source</u>	<u>kg N/ yr</u>	<u>% of total</u>
a. Tributaries (non-point load) - None		
b. Minor tributaries & immediate drainage (non-point load) -	40,990*	90.8
c. Known municipal STP's - Lake Alfred	2,230	4.9
d. Septic tanks**	255	0.6
e. Known industrial - None	-	-
f. Direct precipitation* -	<u>1,680</u>	<u>3.7</u>
Total	45,155	100.0

2. Outputs -

Lake outlet - Unnamed Canal A-1 4,080

3. Net annual N accumulation - 41,075 kg.

* Brezonik and Shannon, 1971.

** Estimate based on 24 lakeshore dwellings; see Working Paper No. 175.

D. Yearly Loads:

In the following table, the existing phosphorus loadings are compared to those proposed by Vollenweider (Vollenweider and Dillon, 1974). Note, however, that Florida lakes may be able to assimilate phosphorus at a somewhat higher level than that suggested by Vollenweider (Shannon and Brezonik, 1972).

Essentially, Vollenweider's "dangerous" loading is one at which the receiving water would become eutrophic or remain eutrophic; his "permissible" loading is that which would result in the receiving water remaining oligotrophic or becoming oligotrophic if morphometry permitted. A mesotrophic loading would be considered one between "dangerous" and "permissible".

Vollenweider's model may not be applicable to water bodies with short hydraulic retention times.

	<u>Total Phosphorus</u>		<u>Total Nitrogen</u>	
	<u>Total</u>	<u>Accumulated</u>	<u>Total</u>	<u>Accumulated</u>
grams/m ² /yr	0.68	0.63	15.6	14.2

Vollenweider phosphorus loadings
(g/m²/yr) based on estimated mean depth and
hydraulic retention time of Lake Haines:

"Dangerous" (eutrophic loading)	0.20
"Permissible" (oligotrophic loading)	0.10

V. LITERATURE REVIEWED

- Anderson, Warren, 1974. Personal communication (hydrology of Winter Haven chain of lakes). U.S. Geol. Surv., Winter Park.
- Brezonik, Patrick L., and Earl E. Shannon; 1971. Trophic state of lakes in north central Florida. Publ. No. 13, FL Water Resources Res. Ctr., U of FL, Gainesville.
- Hulbert, James, 1973. Personal communication (lake surface area). FL Dept. of Env. Reg., Tallahassee.
- Shannon, Earl E., and Patrick L. Brezonik, 1972. Relationships between lake trophic state and nitrogen and phosphorus loading rates. Env. Sci. & Techn. 6 (8): 719-725.
- Vollenweider, R. A., and P. J. Dillon, 1974. The application of the phosphorus loading concept to eutrophication research. Natl. Res. Council of Canada Publ. No. 13690, Canada Centre for Inland waters, Burlington, Ontario.

VI. APPENDICES

APPENDIX A

LAKE RANKINGS

LAKE DATA TO BE USED IN RANKINGS

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLOR A	15- MIN DO	MEDIAN DISS ORTHO P
1201	ALLIGATOR LAKE	0.620	0.260	474.000	87.733	13.100	0.336
1202	LAKE APOPKA	0.102	0.230	484.176	46.611	8.200	0.019
1203	LAKE BANANA	0.660	0.260	482.667	208.600	3.600	0.293
1206	LAKE CRESCENT	0.065	0.130	473.884	10.211	10.200	0.033
1207	DOCTORS LAKE	0.084	0.120	465.555	27.100	10.600	0.028
1208	LAKE DORA	0.102	0.240	482.889	59.978	7.400	0.022
1209	LAKE EFFIE	1.480	0.410	489.000	261.433	15.000	0.950
1210	LAKE GEORGE	0.129	0.165	469.308	35.000	11.000	0.063
1211	LAKE GIBSON	0.167	0.115	470.000	19.675	10.200	0.069
1212	GLENADA LAKE	0.134	0.165	454.167	27.667	14.700	0.072
1214	LAKE GRIFFIN	0.119	0.260	481.333	66.855	6.600	0.038
1215	LAKE HAINES	0.063	0.115	462.667	26.567	10.600	0.014
1217	LAKE HANCOCK	0.772	0.195	483.500	97.900	5.600	0.158
1219	LAKE HORSESHOE	0.034	0.130	459.000	12.067	11.500	0.023
1220	LAKE HOWELL	1.260	0.285	464.000	54.117	9.000	1.175
1221	LAKE ISTOKPOGA	0.039	0.120	464.222	6.594	8.600	0.010
1223	LAKE JESSUP	0.492	0.290	487.000	76.550	7.600	0.288
1224	LAKE KISSIMEE	0.034	0.145	463.667	24.142	8.800	0.007
1227	LAKE LILU	1.490	1.065	483.000	276.566	14.300	1.030
1228	LAKE MARION	0.044	0.260	468.833	29.967	7.600	0.016
1229	LAKE MINNEHAHA	0.038	0.080	435.000	8.733	7.700	0.012
1230	LAKE MINNEOLA	0.018	0.070	406.333	3.333	7.400	0.009
1231	LAKE MONROE	0.188	0.300	474.555	14.225	10.800	0.128
1232	LAKE OKEECHOBEE	0.063	0.185	472.366	14.524	9.800	0.010
1234	LAKE POINSETT	0.085	0.150	469.000	6.500	10.600	0.051
1236	LAKE PEEDY	0.033	0.330	468.500	34.837	10.600	0.008
1238	LAKE SOUTH	0.074	0.130	464.000	23.167	9.000	0.028
1239	LAKE TALQUIN	0.085	0.290	462.167	9.483	14.400	0.031

LAKE DATA TO BE USED IN RANKINGS

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLORA	15- MIN DO	MEDIAN DISS ORTHO P
1240	LAKE THONOTOSASSA	0.695	0.095	466.167	37.700	10.200	0.565
1241	LAKE TOHOPEKALIGA	0.246	0.200	472.917	30.633	10.500	0.152
1242	TROUT LAKE	1.110	0.650	472.000	76.967	12.900	0.970
1243	LAKE WEOHYAKAPKA	0.047	0.080	458.667	7.767	8.200	0.011
1246	LAKE YALE	0.027	0.160	441.000	25.367	7.600	0.014
1247	LAKE MUNSON	1.475	0.925	486.667	140.317	12.200	0.852
1248	LAKE SEMINOLE	0.234	0.175	473.833	102.000	8.600	0.026
1249	LAKE LAWNE	2.560	1.350	494.667	84.900	10.400	0.117
1250	LAKE TARPON	0.041	0.070	400.889	6.867	9.000	0.027
1252	LAKE ELOISE	0.486	0.170	465.333	70.233	12.200	0.339
1258	LAKE JESSIE	0.051	0.090	452.667	26.300	10.800	0.011
1261	EAST LAKE TOHOPEKALIGA	0.042	0.070	440.833	5.167	9.400	0.007
1264	PAYNE'S PRAIRIE LAKE (NO	1.260	0.140	476.000	98.200	7.400	1.210

PERCENT OF LAKES WITH HIGHER VALUES (NUMBER OF LAKES WITH HIGHER VALUES)

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLORA	15- MIN DO	MEDIAN DISS ORTHO P	INDEX NO
1201	ALLIGATOR LAKE	25 (10)	29 (10)	30 (12)	18 (7)	10 (4)	18 (7)	130
1202	LAKE APOPKA	50 (20)	38 (15)	10 (4)	38 (15)	74 (29)	70 (28)	280
1203	LAKE BANANA	23 (9)	29 (10)	20 (8)	5 (2)	100 (40)	23 (9)	200
1206	LAKE CRESCENT	65 (26)	70 (27)	33 (13)	80 (32)	48 (18)	50 (20)	346
1207	DOCTORS LAKE	60 (24)	76 (30)	60 (24)	55 (22)	34 (12)	56 (22)	341
1208	LAKE DORA	53 (21)	35 (14)	18 (7)	33 (13)	90 (35)	68 (27)	297
1209	LAKE EFFIE	5 (2)	10 (4)	3 (1)	3 (1)	0 (0)	10 (4)	31
1210	LAKE GEORGE	45 (18)	54 (21)	48 (19)	43 (17)	23 (9)	43 (17)	256
1211	LAKE GIBSON	40 (16)	81 (32)	45 (18)	70 (28)	48 (18)	40 (16)	324
1212	GLENADA LAKE	43 (17)	54 (21)	85 (34)	53 (21)	3 (1)	38 (15)	276
1214	LAKE GRIFFIN	48 (19)	29 (10)	23 (9)	30 (12)	95 (38)	48 (19)	273
1215	LAKE HAINES	70 (28)	81 (32)	75 (30)	58 (23)	34 (12)	78 (31)	396
1217	LAKE MANCOCK	18 (7)	43 (17)	13 (5)	13 (5)	98 (39)	28 (11)	213
1219	LAKE HORSESHOE	93 (37)	70 (27)	80 (32)	78 (31)	20 (8)	65 (26)	406
1220	LAKE HOWELL	11 (4)	23 (9)	69 (27)	35 (14)	60 (23)	3 (1)	201
1221	LAKE ISTOKPOGA	85 (34)	76 (30)	65 (26)	93 (37)	69 (27)	89 (35)	477
1223	LAKE JESSUP	28 (11)	18 (7)	5 (2)	25 (10)	83 (32)	25 (10)	184
1224	LAKE KISSIMMEE	90 (36)	63 (25)	73 (29)	65 (26)	65 (26)	99 (39)	455
1227	LAKE LULU	3 (1)	3 (1)	15 (6)	0 (0)	8 (3)	5 (2)	34
1228	LAKE MARION	78 (31)	29 (10)	53 (21)	50 (20)	83 (32)	73 (29)	366
1229	LAKE MINNEHABA	88 (35)	91 (36)	95 (38)	85 (34)	78 (31)	80 (32)	517
1230	LAKE MINNEOLA	100 (40)	98 (38)	98 (39)	100 (40)	90 (35)	93 (37)	579
1231	LAKE MONROE	38 (15)	15 (6)	28 (11)	75 (30)	26 (10)	33 (13)	215
1232	LAKE OKEECHOBEE	68 (27)	45 (18)	40 (16)	73 (29)	53 (21)	89 (35)	368
1234	LAKE POINSETT	58 (23)	60 (24)	50 (20)	95 (38)	34 (12)	45 (18)	342
1236	LAKE REEDY	95 (38)	13 (5)	55 (22)	45 (18)	34 (12)	95 (38)	337
1238	LAKE SOUTH	63 (25)	70 (27)	69 (27)	68 (27)	60 (23)	56 (22)	386
1239	LAKE TALQUIN	55 (22)	20 (8)	78 (31)	83 (33)	5 (2)	53 (21)	294

PERCENT OF LAKES WITH HIGHER VALUES (NUMBER OF LAKES WITH HIGHER VALUES)

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLOR A	15- MIN DO	MEDIAN DISS ORTHO P	INDEX NO
1240	LAKE THONOTOSASSA	20 (8)	85 (34)	58 (23)	40 (16)	48 (18)	15 (6)	265
1241	LAKE TOHOPEKALIGA	33 (13)	40 (16)	38 (15)	48 (19)	40 (16)	30 (12)	229
1242	TROUT LAKE	15 (6)	8 (3)	43 (17)	23 (9)	13 (5)	8 (3)	110
1243	LAKE WEOHYAKAPKA	75 (30)	91 (36)	83 (33)	88 (35)	74 (29)	84 (33)	495
1246	LAKE YALE	98 (39)	58 (23)	90 (36)	63 (25)	83 (32)	75 (30)	467
1247	LAKE MUNSON	8 (3)	5 (2)	8 (3)	8 (3)	16 (6)	13 (5)	58
1248	LAKE SEMINOLE	35 (14)	48 (19)	35 (14)	10 (4)	69 (27)	63 (25)	260
1249	LAKE LAWNE	0 (0)	0 (0)	0 (0)	20 (8)	43 (17)	35 (14)	98
1250	LAKE TARPON	83 (33)	98 (38)	100 (40)	90 (36)	60 (23)	60 (24)	491
1252	LAKE ELOISE	30 (12)	50 (20)	63 (25)	28 (11)	16 (6)	20 (8)	207
1258	LAKE JESSIE	73 (29)	88 (35)	88 (35)	60 (24)	26 (10)	84 (33)	419
1261	EAST LAKE TOHOPEKALIGA	80 (32)	98 (38)	93 (37)	98 (39)	55 (22)	99 (39)	523
1264	PAYNE'S PRAIRIE LAKE (NO	11 (4)	65 (26)	25 (10)	15 (6)	90 (35)	0 (0)	206

LAKES RANKED BY INDEX NOS.

RANK	LAKE CODE	LAKE NAME	INDEX NO
1	1230	LAKE MINNEOLA	579
2	1261	EAST LAKE TOHOPEKALIGA	523
3	1229	LAKE MINNEHAHA	517
4	1243	LAKE WEOHYAKAPKA	495
5	1250	LAKE TARPUN	491
6	1221	LAKE ISTOKPOGA	477
7	1246	LAKE YALE	467
8	1224	LAKE KISSIMMEE	455
9	1258	LAKE JESSIE	419
10	1219	LAKE HORSESHOE	406
11	1215	LAKE HAINES	396
12	1238	LAKE SOUTH	386
13	1232	LAKE OKEECHUBEE	368
14	1228	LAKE MARION	366
15	1206	LAKE CRESCENT	346
16	1234	LAKE POINSETT	342
17	1207	DOCTORS LAKE	341
18	1236	LAKE REEDY	337
19	1211	LAKE GIBSON	324
20	1208	LAKE DORA	297
21	1239	LAKE TALQUIN	294
22	1202	LAKE APOPKA	280
23	1212	GLENADA LAKE	276
24	1214	LAKE GRIFFIN	273
25	1240	LAKE THONOTOSASSA	266
26	1248	LAKE SEMINOLE	260
27	1210	LAKE GEORGE	256
28	1241	LAKE TOHOPEKALIGA	229

LAKES RANKED BY INDEX NOS.

RANK	LAKE CODE	LAKE NAME	INDEX NO
29	1231	LAKE MONROE	215
30	1217	LAKE MANCOCK	213
31	1252	LAKE ELOISE	207
32	1264	PAYNE'S PRAIRIE LAKE (NO	206
33	1220	LAKE HOWELL	201
34	1203	LAKE BANANA	200
35	1223	LAKE JESSUP	184
36	1201	ALLIGATOR LAKE	130
37	1242	TROUT LAKE	110
38	1249	LAKE LAWNE	98
39	1247	LAKE MUNSON	58
40	1227	LAKE LULU	34
41	1209	LAKE EFFIE	31

APPENDIX B

CONVERSION FACTORS

CONVERSION FACTORS

Hectares x 2.471 = acres

Kilometers x 0.6214 = miles

Meters x 3.281 = feet

Cubic meters x 8.107×10^{-4} = acre/feet

Square kilometers x 0.3861 = square miles

Cubic meters/sec x 35.315 = cubic feet/sec

Centimeters x 0.3937 = inches

Kilograms x 2.205 = pounds

Kilograms/square kilometer x 5.711 = lbs/square mile

APPENDIX C

TRIBUTARY FLOW DATA

TRIBUTARY FLOW INFORMATION FOR FLORIDA

8/25/75

LAKE CODE 1215 LAKE HAINES

TOTAL DRAINAGE AREA OF LAKE(SQ KM) 21.2

TRIBUTARY	SUB-DRAINAGE AREA(SQ KM)	NORMALIZED FLOWS(CMS)												
		JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
1215A1	21.2	0.06	0.06	0.11	0.08	0.03	0.06	0.09	0.09	0.16	0.14	0.06	0.06	0.08
1215ZZ	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

SUMMARY

TOTAL DRAINAGE AREA OF LAKE =	21.2	TOTAL FLOW IN =	0.0
SUM OF SUB-DRAINAGE AREAS =	0.0	TOTAL FLOW OUT =	1.01

NOTE *** SEE WRITE UP ON WINTER HAVEN CHAIN OF LAKES

MEAN MONTHLY FLOWS AND DAILY FLOWS(CMS)

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
1215A1	3	73	0.03	18	0.02				
	4	73	0.03	15	0.03				
	5	73	0.02	18	0.01				
	6	73	0.03	17	0.01				
	7	73	0.02	14	0.03				
	8	73	0.02	17	0.02				
	9	73	0.04	16	0.03				
	10	73	0.03	13	0.03				
	11	73	0.01	18	0.01				
	12	73	0.02	16	0.01				
	1	74	0.02	20	0.01				
	2	74	0.01	17	0.01				

APPENDIX D

PHYSICAL and CHEMICAL DATA

STORET RETRIEVAL DATE 75/08/25

121501
28 05 24.0 081 42 31.0
LAKE HAINES
12105 FLORIDA

11EPALES
3

2111202
0014 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00010 WATER TEMP CENT	00300 DO MG/L	00077 TRANSP SECCHI INCHES	00094 CONDUCTVY FIELD MICROMHO	00400 PH SU	00410 T ALK CAC03 MG/L	00610 NH3-N TOTAL MG/L	00625 TOT KJEL N MG/L	00630 NO2&NO3 N-TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P
73/03/08	10 30	0000	23.2		30	215	9.10	12	0.060	1.000	0.050	0.009
	10 30	0004	22.4	8.9		205	8.60	11	0.060	0.700	0.070	0.010
	10 30	0010	17.6	4.8		220	8.20	10	0.120	0.800	0.140	0.030
73/09/06	10 15	0000	28.8	6.2	48	221	7.30	15	0.060	1.400	0.060	0.008
	10 15	0012	27.8	4.4		218	6.60	17	0.080	0.900	0.060	0.014
73/11/07	13 30	0000	23.2		34	190	6.90	14	0.050	1.300	0.040	0.015
	13 30	0005	23.1	7.2		190	6.80	14	0.050	1.100	0.030	0.015
	13 30	0010	23.1	7.0		189	7.00	15	0.050	1.200	0.040	0.014

DATE FROM TO	TIME OF DAY	DEPTH FEET	00665 PHOS-TOT MG/L P	32217 CHLRPHYL A UG/L
73/03/08	10 30	0000	0.054	37.3
	10 30	0004	0.053	
	10 30	0010	0.074	
73/09/06	10 15	0000	0.054	25.3
	10 15	0012	0.044	
73/11/07	13 30	0000	0.072	17.1
	13 30	0005	0.072	
	13 30	0010	0.078	

APPENDIX E

TRIBUTARY and WASTEWATER TREATMENT PLANT DATA

STORET RETRIEVAL DATE 75/08/25

1215A1
 28 05 00.0 081 43 00.0
 UNNAMED STREAM
 12069 7.5 WINTER HAVEN
 T/LAKES HAINES
 BANK OF STREAM BTWN LKS HAINES & ROCHELL
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 NO2&NO3 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
73/03/18	09 43		0.033	0.580	0.048	0.013	0.035
73/04/15	15 30		0.040	2.300	0.138	0.016	0.100
73/05/18	12 10		0.035	4.700	0.032	0.008	0.035
73/06/17	13 55		0.044	2.940	0.200	0.016	0.050
73/07/14	11 25		0.046	0.840	0.058	0.008	0.050
73/08/17	13 45		0.095	0.720	0.032	0.014	0.060
73/09/16	09 40		0.072	0.900	0.039	0.019	0.050
73/10/13	10 50		0.170	0.700	0.071	0.043	0.065
73/11/18	11 10		0.120	0.600	0.032	0.020	0.045
73/12/16	14 36		0.350	0.700	0.036	0.020	0.045
74/01/20	14 35		0.044	0.900	0.028	0.024	0.100
74/02/17	14 45		0.036	0.900	0.015	0.010	0.090

STORET RETRIEVAL DATE 75/08/25

1215DA PD1215DA P001700
28 05 30.0 081 43 00.0
LAKE ALFRED
12069 7.5 WINTERHAVEN
D/LAKE HAINES
LAKE HAINES
11EPALES 2141204
4 0000 FEET DEPTH

DATE	TIME	DEPTH	00630 NO2&NO3 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P	50051 FLOW RATE INST MGD	50053 CONDUIT FLOW-MGD MONTHLY
FROM	OF								
TO	DAY	FEET							
74/02/28	07	30							
CP(T)-			7.400	1.000K	0.230	7.100	7.500	0.175	0.160
74/02/28	16	30							

K VALUE KNOWN TO BE
LESS THAN INDICATED