

**U.S. ENVIRONMENTAL PROTECTION AGENCY
NATIONAL EUTROPHICATION SURVEY
WORKING PAPER SERIES**



REPORT
ON
HERRINGTON LAKE
BOYLE, GARRARD, AND MERCER COUNTIES
KENTUCKY
EPA REGION IV
WORKING PAPER No. 353

**CORVALLIS ENVIRONMENTAL RESEARCH LABORATORY - CORVALLIS, OREGON
and
ENVIRONMENTAL MONITORING & SUPPORT LABORATORY - LAS VEGAS, NEVADA**

REPORT
ON
HERRINGTON LAKE
BOYLE, GARRARD, AND MERCER COUNTIES
KENTUCKY
EPA REGION IV
WORKING PAPER No. 353

WITH THE COOPERATION OF THE
KENTUCKY DEPARTMENT FOR NATURAL RESOURCES AND ENVIRONMENTAL CONTROL
AND THE
KENTUCKY NATIONAL GUARD
JUNE, 1977

CONTENTS

	<u>Page</u>
<u>Foreword</u>	ii
List of Kentucky Reservoirs	iv
Lake and Drainage Area Map	v
 <u>Sections</u>	
I. Conclusions	1
II. Lake and Drainage Basin Characteristics	4
III. Lake Water Quality Summary	5
IV. Nutrient Loadings	9
V. Literature Reviewed	15
VI. Appendices	16

F O R E W O R D

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nation-wide threat of accelerated eutrophication to freshwater lakes and reservoirs.

OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

- a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.
- b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.
- c. With such a transformation, an assessment of the potential for eutrophication control can be made.

LAKE ANALYSIS

In this report, the first stage of evaluation of lake and watershed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning [§303(e)], water quality criteria/standards review [§303(c)], clean lakes [§314(a,b)], and water quality monitoring [§106 and §305(b)] activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

ACKNOWLEDGEMENT

The staff of the National Eutrophication Survey (Office of Research & Development, U. S. Environmental Protection Agency) expresses sincere appreciation to the Kentucky Department for Natural Resources and Environmental Protection for professional involvement, to the Kentucky National Guard for conducting the tributary sampling phase of the Survey, and to those Kentucky wastewater treatment plant operators who voluntarily provided effluent samples.

Harold Snodgrass, former Director, and Mercer M. Peters of the Division of Water Quality provided invaluable lake documentation and counsel during the Survey; and the staff of the Division reviewed the preliminary reports and provided critiques most useful in the preparation of this Working Paper series.

Major General Richard L. Frymire, the Adjutant General of Kentucky, and Project Officers Lt. Colonel Thomas Buyher, Jr., and Lt. Colonel David B. May, who directed the volunteer efforts of the Kentucky National Guardsmen, are also gratefully acknowledged for their assistance to the Survey.

NATIONAL EUTROPHICATION SURVEY

STUDY RESERVOIRS

STATE OF KENTUCKY

NAMECOUNTY

Barkley

Lyon, Trigg, KY;
Cheatham, Montgomery,
Stewart, TN

Barren River

Allen, Barren

Cumberland

Clinton, Pulaski, Russell,
Wayne

Dale Hollow


Clinton, Cumberland, KY;
Clay, Overton, Pickett, TN

Herrington

Boyle, Garrard, Mercer

Kentucky

Calloway, Livingston, Lyon,
Marshall, Trigg, KY;
Benton, Decatur, Henry,
Houston, Humphreys, Perry,
Stewart, TN



HERRINGTON LAKE

STORET NO. 2103

I. CONCLUSIONS

A. Trophic Condition:

Survey data indicate that Herrington Lake is eutrophic; i.e., well-supplied with nutrients and quite productive. Whether nutrient enrichment is beneficial or deleterious depends on the actual or potential effect on the uses of the lake. In this regard, personnel of the Kentucky Department for Natural Resources and Environmental Protection indicate additional study is needed to determine whether the beneficial uses of Herrington Lake are impaired.

This water body ranked last in overall trophic quality when the five Kentucky reservoirs sampled in 1973 were compared using a combination of six parameters*. All of the other reservoirs had less median total phosphorus, median orthophosphorus, median inorganic nitrogen, and mean chlorophyll a. Two of the water bodies had greater mean Secchi disc transparency.

Marked depression or depletion of dissolved oxygen with depth and chlorophyll a levels at all sampling sites in May and August and at station 4 in October indicate lake deterioration.

B. Rate-Limiting Nutrient:

The algal assay results indicate Herrington Lake was limited

* See Appendix A.

by phosphorus at the time the sample was collected (05/26/73). These results are substantiated by the lake data obtained at that time; i.e., the mean inorganic nitrogen to orthophosphorus ratio was 16 to 1. At the other two sampling times (08/20/73 and 10/25/73), the lake data indicate that nitrogen was the limiting nutrient; i.e., the mean inorganic nitrogen to orthophosphorus ratios were 8 to 1 and less.

C. Nutrient Controllability:

1. Point sources--The phosphorus contribution of known point sources amounted to 26.7% of the total phosphorus load reaching Herrington Lake during the sampling year. The wastewater treatment facilities at Danville, Lancaster, Stanford, and Herrington Haven collectively contributed 23.7% of the total. The untreated wastes from Burgin and Junction City together accounted for 3.0% of the load.

The phosphorus loading of 9.24 g/m^2 measured during the sampling year is seven times that proposed by Vollenweider (Vollenweider and Dillon, 1974) as a eutrophic loading (see page 14). Even complete removal of phosphorus at the listed point sources would only reduce the loading to $6.78 \text{ g/m}^2/\text{yr}$ (over five times the eutrophic level), and unless a significant portion of the non-point load can also be reduced, it does not seem likely that point-source removal would result in a significant improvement in the trophic condition of the lake.

2. Non-point sources--Non-point sources accounted for 73.3% of the total phosphorus input to the lake. The Dix River and Clarks Run contributed 29.7% and 28.2% of this total, respectively. The remaining three tributaries contributed a combined load amounting to 10.0% of the total. The ungaged tributaries contributed an estimated 5.1% of the total phosphorus load.

Clarks Run, Mocks Branch, and Spears Creek had export rates which were extremely high (see page 13) as compared to the tributaries of Cumberland Lake*; e.g., Indian Creek ($10 \text{ kg/km}^2/\text{yr}$), Meadow Creek ($16 \text{ kg/km}^2/\text{yr}$), and the Laurel River ($24 \text{ kg/km}^2/\text{yr}$). The high rates of the Herrington Lake tributaries may be due to unknown point sources impacting those streams.

* See Working Paper No. 351.

II. LAKE AND DRAINAGE BASIN CHARACTERISTICS[†]

A. Morphometry^{††}:

1. Surface area: 11.90 kilometers².
2. Mean depth: 23.9 meters.
3. Maximum depth: >61 meters.
4. Volume: 284.410 x 10⁶ m³.
5. Mean hydraulic retention time: 196 days.

B. Tributary and Outlet: (See Appendix C for flow data)

1. Tributaries -

<u>Name</u>	<u>Drainage area (km²)*</u>	<u>Mean flow (m³/sec)*</u>
Dix River	823.6	12.15
Clarks Run	73.0	0.98
Mocks Branch	42.5	0.56
Spears Creek	19.8	0.26
Cane Run	25.3	0.33
Minor tributaries & immediate drainage -	<u>140.9</u>	<u>2.50</u>
Totals	1,125.1	16.78**

2. Outlet -

Dix River	1,137.0***	16.78
-----------	------------	-------

C. Precipitation****:

1. Year of sampling: 126.9 centimeters.
2. Mean annual: 115.8 centimeters.

[†] Table of metric conversions--Appendix B.

^{††} Jett, 1974.

* For limits of accuracy, see Working Paper No. 175, "Survey Methods, 1973-1976".

** Sum of inflows adjusted to equal outflow.

*** Includes area of lake.

**** See Working Paper No. 175.

III. WATER QUALITY SUMMARY

Herrington Lake was sampled three times in 1973 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from a number of depths at four stations on the lake (see map, page v). During each visit, a single depth-integrated (4.6 m to surface) sample was composited from the stations for phytoplankton identification and enumeration; and during the first visit, a single 18.9-liter depth-integrated sample was composited for algal assays. Also each time, a depth-integrated sample was collected from each of the stations for chlorophyll a analysis. The maximum depths sampled were 61.0 meters at station 1, 55.8 meters at station 2, 44.2 meters at station 3, and 9.1 meters at station 4.

The sampling results are presented in full in Appendix D and are summarized in the following table.

A. SUMMARY OF PHYSICAL AND CHEMICAL CHARACTERISTICS FOR HERRINGTON LAKE
STORET CODE 2103

1ST SAMPLING (5/26/73)

2ND SAMPLING (8/20/73)

3RD SAMPLING (10/25/73)

4 SITES

4 SITES

4 SITES

PARAMETER	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN
TEMP (C)	7.0 - 23.9	16.4	17.5	5.4 - 28.1	21.4	24.7	6.8 - 21.4	18.1	20.6
DISS OXY (MG/L)	2.2 - 16.0	7.1	6.5	0.0 - 8.0	2.0	0.2	0.4 - 6.2	2.5	2.8
CNDCTVY (MCROMO)	***** -*****			201. - 318.	242.	236.	205. - 282.	232.	230.
PH (STAND UNITS)	7.4 - 9.2	8.1	7.9	6.4 - 7.9	7.1	7.0	7.0 - 8.0	7.5	7.4
TOT ALK (MG/L)	78. - 117.	103.	106.	75. - 150.	103.	98.	98. - 144.	108.	106.
TOT P (MG/L)	0.027 - 0.231	0.090	0.079	0.028 - 0.412	0.126	0.087	0.046 - 0.445	0.113	0.074
ORTHO P (MG/L)	0.009 - 0.110	0.054	0.054	0.012 - 0.214	0.073	0.056	0.012 - 0.196	0.063	0.042
NO2+NO3 (MG/L)	0.180 - 1.590	0.795	0.815	0.040 - 1.400	0.332	0.120	0.020 - 1.020	0.145	0.040
AMMONIA (MG/L)	0.030 - 0.260	0.094	0.065	0.040 - 1.620	0.280	0.090	0.040 - 0.980	0.144	0.060
KJEL N (MG/L)	0.200 - 1.000	0.480	0.450	0.300 - 2.700	0.863	0.700	0.200 - 1.800	0.587	0.450
INORG N (MG/L)	0.220 - 1.630	0.889	0.950	0.080 - 1.740	0.612	0.560	0.070 - 1.060	0.288	0.100
TOTAL N (MG/L)	0.750 - 1.890	1.275	1.265	0.540 - 2.820	1.195	1.060	0.220 - 1.840	0.732	0.710
CHLRPYL A (UG/L)	6.1 - 26.5	17.6	18.9	8.0 - 11.0	9.2	9.0	1.7 - 64.6	17.8	2.5
SECCHI (METERS)	0.9 - 1.5	1.1	1.0	1.0 - 2.7	1.6	1.3	0.5 - 2.7	1.6	1.7

B. Biological characteristics:

1. Phytoplankton -

<u>Sampling Date</u>	<u>Dominant Genera</u>	<u>Algal Units per ml</u>
05/26/73	1. Flagellates	4,970
	2. <u>Fragilaria sp.</u>	4,970
	3. <u>Scenedesmus sp.</u>	155
	4. <u>Stephanodiscus sp.</u>	133
	5. <u>Asterionella sp.</u>	89
	Other genera	<u>289</u>
	Total	10,606
08/20/73	1. <u>Raphidiopsis sp.</u>	1,711
	2. Flagellates	1,498
	3. <u>Stephanodiscus sp.</u>	642
	4. <u>Oscillatoria sp.</u>	585
	5. <u>Synedra sp.</u>	313
	Other genera	<u>768</u>
	Total	5,517
10/25/73	1. Flagellates	7,501
	2. <u>Melosira sp.</u>	1,162
	3. <u>Stephanodiscus sp.</u>	563
	4. <u>Cyclotella sp.</u>	246
	5. <u>Merismopedia sp.</u>	211
	Other genera	<u>741</u>
	Total	10,424

2. Chlorophyll a -

<u>Sampling Date</u>	<u>Station Number</u>	<u>Chlorophyll <u>a</u> ($\mu\text{g/l}$)</u>
05/26/73	1	21.7
	2	26.5
	3	6.1
	4	16.2
08/20/73	1	9.4
	2	8.6
	3	8.0
	4	11.0
10/25/73	1	2.0
	2	1.7
	3	3.1
	4	64.6

C. Limiting Nutrient Study:

1. Autoclaved, filtered, and nutrient spiked -

<u>Spike (mg/l)</u>	<u>Ortho P Conc. (mg/l)</u>	<u>Inorganic N Conc. (mg/l)</u>	<u>Maximum yield (mg/l-dry wt.)</u>
Control	0.032	0.856	8.1
0.050 P	0.082	0.856	13.5
0.050 P + 1.0 N	0.082	1.856	18.5
1.0 N	0.032	1.856	7.1

2. Discussion -

The control yield of the assay alga, Selenastrum capricornutum, indicates that the potential primary productivity of Herrington Lake was high at the time the sample was collected (05/26/73). Also, the significant increase in yield with the addition of orthophosphorus indicates phosphorus limitation. Note that the addition of nitrogen alone resulted in a yield not significantly different from that of the control.

The lake data also indicate phosphorus limitation at that time; i.e., the mean inorganic nitrogen/orthophosphorus ratio was 16/1. At the other sampling times (08/20/73 and 10/25/73), the lake data indicate that nitrogen was the limiting nutrient; i.e., the mean inorganic nitrogen/orthophosphorus ratios were 8/1 and less.

IV. NUTRIENT LOADINGS (See Appendix E for data)

For the determination of nutrient loadings, the Kentucky National Guard collected monthly near-surface grab samples from each of the tributary sites indicated on the map (page v), except for the month of May when two samples were collected at most of the sites. Sampling was begun in March, 1973, and was completed in April, 1974.

Through an interagency agreement, stream flow estimates for the year of sampling and a "normalized" or average year were provided by the Kentucky District Office of the U.S. Geological Survey for the tributary sites nearest the lake.

In this report, nutrient loads for sampled tributaries were determined by using a modification of a U.S. Geological Survey computer program for calculating stream loadings*. Nutrient loads shown are those measured minus point-source loads, if any.

Nutrient loadings for unsampled "minor tributaries and immediate drainage" ("ZZ" of U.S.G.S.) were estimated using the nutrient exports of the Dix River at station D-3, in kg/km²/year, and multiplying by the ZZ area in km².

The operators of the Lancaster, Stanford, and Danville wastewater treatment plants provided monthly effluent samples and corresponding flow data. The operator of the Herrington Haven State Hospital wastewater treatment plant did not participate in the sampling program, and nutrient loads were estimated at 1.134 kg P and 3.401 kg N/capita/

* See Working Paper No. 175.

year. The untreated nutrient loads from Burgin and Junction City were estimated at 1.587 kg P and 4.263 kg N/capita/year.

A. Waste Sources*:

1. Known municipal -

<u>Name</u>	<u>Pop. Served</u>	<u>Treatment</u>	<u>Mean Flow (m³/d)</u>	<u>Receiving Water</u>
Lancaster	3,230	trickling filter	844.1	White Oak Creek
Stanford	2,474	trickling filter	1,327.6	Logan Creek
Danville	11,542	trickling filter	6,056.6	Clarks Run
Herrington Haven State Hospital**	800	ext. aer.	302.8	Herrington Lake
Junction City**	1,046	none	395.9	Knoblick Creek
Burgin**	1,002	none	379.3	Unnamed Creek/ Herrington Lake

2. Known industrial - None

* Anonymous, 1971; population served is 1970 Census.

** Flow based on 0.3785 m³/capita/day.

B. Annual Total Phosphorus Loading - Average Year:

1. Inputs -

<u>Source</u>	<u>kg P/ yr</u>	<u>% of total</u>
a. Tributaries (non-point load) -		
Dix River	32,685	29.7
Clarks Run	30,980	28.2
Mocks Branch	6,570	6.0
Spears Creek	2,555	2.3
Cane Run	1,835	1.7
b. Minor tributaries & immediate drainage (non-point load) -	5,635	5.1
c. Known municipal STP's -		
Lancaster	5,845	5.3
Stanford	4,960	4.5
Danville	14,350	13.1
Herrington Haven	905	0.8
Junction City	1,660	1.5
Burgin	1,590	1.4
d. Septic tanks* -	155	0.1
e. Known industrial - None	-	
f. Direct precipitation** -	<u>210</u>	<u>0.2</u>
Total	109,935	100.0

2. Outputs -

Lake outlet - Dix River 51,955

3. Net annual P accumulation - 57,980 kg.

* Estimate based on 513 lakeshore dwellings and four camps; see Working Paper No. 175.

** See Working Paper No. 175.

C. Annual Total Nitrogen Loading - Average Year:

1. Inputs -

<u>Source</u>	<u>kg N/ yr</u>	<u>% of total</u>
a. Tributaries (non-point load) -		
Dix River	719,415	61.8
Clarks Run	121,640	10.5
Mocks Branch	70,010	6.0
Spears Creek	23,475	2.0
Cane Run	25,865	2.2
b. Minor tributaries & immediate drainage (non-point load) -	123,145	10.6
c. Known municipal STP's -		
Lancaster	7,880	0.7
Stanford	6,030	0.5
Danville	36,220	3.1
Herrington Haven	2,720	0.2
Junction City	4,460	0.4
Burgin	4,270	0.4
d. Septic tanks* -	5,745	0.5
e. Known industrial - None	-	-
f. Direct precipitation** -	<u>12,845</u>	<u>1.1</u>
Total	1,163,720	100.0

2. Outputs -

Lake outlet - Dix River 1,565,170

3. Net annual N loss - 401,450 kg.

* Estimate based on 513 lakeshore dwellings and four camps; see Working Paper No. 175.

** See Working Paper No. 175.

D. Non-point Nutrient Export by Subdrainage Area:

<u>Tributary</u>	<u>kg P/km²/yr</u>	<u>kg N/km²/yr</u>
Dix River	40	874
Clarks Run	424	1,666
Mocks Branch	155	1,647
Spears Creek	129	1,186
Cane Run	73	1,022

E. Yearly Loadings:

In the following table, the existing phosphorus loadings are compared to those proposed by Vollenweider (Vollenweider and Dillon, 1974). Essentially, his "dangerous" loading is one at which the receiving water would become eutrophic or remain eutrophic; his "permissible" loading is that which would result in the receiving water remaining oligotrophic or becoming oligotrophic if morphometry permitted. A mesotrophic loading would be considered one between "dangerous" and "permissible".

Note that Vollenweider's model may not be applicable to water bodies with short hydraulic retention times.

	<u>Total Phosphorus</u>		<u>Total Nitrogen</u>	
	<u>Total</u>	<u>Accumulated</u>	<u>Total</u>	<u>Accumulated</u>
grams/m ² /yr	9.24	4.87	97.8	loss*

Vollenweider phosphorus loadings
(g/m²/yr) based on mean depth and mean
hydraulic retention time of Herrington Lake:

"Dangerous" (eutrophic loading)	1.28
"Permissible" (oligotrophic loading)	0.64

* There was an apparent loss of nitrogen during the sampling year. This may have been due to nitrogen fixation in the lake, solubilization of previously sedimented nitrogen, recharge with nitrogen-rich ground water, or unsampled point sources discharging directly to the lake. Whatever the cause, a similar nitrogen loss has occurred at Shagawa Lake, Minnesota, which has been intensively studied by EPA's former National Eutrophication and Lake Restoration Branch (Malueg et al., 1975).

V. LITERATURE REVIEWED

Anonymous, 1971. Inventory of municipal waste facilities. EPA Publ. OWP-1, Vol. 4, Wash., D.C.

Jett, Shelby, 1974. Personal communication (lake morphometry). Dept. for Nat. Resources and Env. Protection, Frankfurt.

Malueg, Kenneth W., D. Phillips Larsen, Donald W. Schults, and Howard T. Mercier; 1975. A six-year water, phosphorus, and nitrogen budget for Shagawa Lake, Minnesota. Jour. Environ. Qual., vol. 4, no. 2, pp. 236-242.

Vollenweider, R. A., and P. J. Dillon, 1974. The application of the phosphorus loading concept to eutrophication research. Natl. Res. Council of Canada Publ. No. 13690, Canada Centre for Inland Waters, Burlington, Ontario.

VI. APPENDICES

APPENDIX A

LAKE RANKINGS

LAKE DATA TO BE USED IN RANKINGS

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLORA	15- MIN DO	MEDIAN DISS P
2101	LAKE CUMBERLAND-	0.016	0.330	432.381	3.805	14.900	0.007
2102	DALE HOLLOW RESERVOIR	0.010	0.270	330.000	3.594	14.800	0.005
2103	HERRINGTON LAKE	0.079	0.550	442.667	14.908	14.900	0.047
2104	KENTUCKY LAKE	0.072	0.520	461.972	9.089	12.600	0.030
2105	BARREN RIVER RESERVOIR	0.027	0.460	452.594	8.216	14.900	0.006

PERCENT OF LAKES WITH HIGHER VALUES

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLORA	15- MIN DO	MEDIAN DISS P	INDEX NO
2101	LAKE CUMBERLAND	60 (3)	60 (3)	60 (3)	60 (3)	20 (0)	40 (2)	300
2102	DALE HOLLOW RESERVOIR	80 (4)	80 (4)	80 (4)	80 (4)	60 (3)	80 (4)	460
2103	HERRINGTON LAKE	0 (0)	0 (0)	40 (2)	0 (0)	20 (0)	0 (0)	60
2104	KENTUCKY LAKE	20 (1)	20 (1)	0 (0)	20 (1)	80 (4)	20 (1)	160
2105	BARREN RIVER RESERVOIR	40 (2)	40 (2)	20 (1)	40 (2)	20 (0)	60 (3)	220

LAKES RANKED BY INDEX NOS.

RANK	LAKE CODE	LAKE NAME	INDEX NO
1	2102	DALE HOLLOW RESERVOIR	460
2	2101	LAKE CUMBERLAND	300
3	2105	BARREN RIVER RESERVOIR	220
4	2104	KENTUCKY LAKE	160
5	2103	HERRINGTON LAKE	60

APPENDIX B

CONVERSION FACTORS

CONVERSION FACTORS

Hectares x 2.471 = acres

Kilometers x 0.6214 = miles

Meters x 3.281 = feet

Cubic meters x 8.107×10^{-4} = acre/feet

Square kilometers x 0.3861 = square miles

Cubic meters/sec x 35.315 = cubic feet/sec

Centimeters x 0.3937 = inches

Kilograms x 2.205 = pounds

Kilograms/square kilometer x 5.711 = lbs/square mile

APPENDIX C

TRIBUTARY FLOW DATA

TRIBUTARY FLOW INFORMATION FOR KENTUCKY

2/3/75

LAKE CODE 2103 HERRINGTON LAKE

TOTAL DRAINAGE AREA OF LAKE(SQ KM) 1137.0

TRIBUTARY	SUB-DRAINAGE AREA(SQ KM)	NORMALIZED FLOWS(CMS)												
		JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
2103A1	1137.0	27.24	32.51	37.32	25.17	13.90	10.65	10.56	6.46	5.58	4.33	9.34	19.23	16.78
2103A3	823.6	22.99	29.17	30.21	16.88	9.97	6.68	5.83	2.10	1.30	0.62	6.17	14.92	12.15
2103B1	73.0	1.59	2.18	2.63	1.50	1.02	0.48	0.42	0.21	0.15	0.10	0.48	1.10	0.98
2103C1	42.5	0.88	1.25	1.53	0.88	0.59	0.27	0.25	0.12	0.08	0.06	0.28	0.62	0.56
2103D1	19.8	0.40	0.57	0.71	0.40	0.28	0.12	0.11	0.05	0.04	0.03	0.13	0.28	0.26
2103E1	25.3	0.51	0.74	0.91	0.51	0.37	0.16	0.15	0.07	0.05	0.03	0.17	0.37	0.33
2103Z2	143.0	3.20	4.28	5.18	3.00	1.98	0.99	0.85	0.42	0.31	0.22	0.96	2.15	1.95

SUMMARY

TOTAL DRAINAGE AREA OF LAKE = 1137.0
SUM OF SUB-DRAINAGE AREAS = 1127.2

TOTAL FLOW IN = 196.22
TOTAL FLOW OUT = 202.30

MEAN MONTHLY FLOWS AND DAILY FLOWS(CMS)

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
2103A1	3	73	28.32	31	5.66				
	4	73	37.38						
	5	73	45.87	1	36.81	31	36.81		
	6	73	22.63						
	7	73	19.54	3	36.81				
	8	73	7.59	3	5.66				
	9	73	5.78	6	19.82				
	10	73	4.76	3	5.66				
	11	73	20.05	4	7.08				
	12	73	27.35	1	50.97				
	1	74	82.69	5	39.64	20	59.47		
	2	74	24.55	15	11.89				
2103A3	3	74	37.94	1	17.84	16	24.35		
	4	74	32.85	2	53.80				
	3	73	23.47	31	14.38				
	4	73	30.07						
	5	73	38.99	1	15.01	30	44.17		
	6	73	12.66						
	7	73	12.35	3	9.46				
	8	73	0.97	1	3.77	31	0.25		
	9	73	0.09						
	10	73	0.04	2	0.05	31	0.13		
	11	73	21.04						
	12	73	16.57	31	41.63				
	1	74	71.25	5	41.06				
	2	74	11.33						
	3	74	26.73	1	7.19				
	4	74	18.69						

TRIBUTARY FLOW INFORMATION FOR KENTUCKY

2/3/75

LAKE CODE 2103 HERRINGTON LAKE

MEAN MONTHLY FLOWS AND DAILY FLOWS (CMS)

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
2103B1	3	73	2.07	31	1.27				
	4	73	2.66						
	5	73	3.99	1	1.56	30	4.50		
	6	73	0.91						
	7	73	0.91	3	0.71				
	8	73	0.10	1	0.37	31	0.03		
	9	73	0.01						
	10	73	0.01	2	0.01	31	0.03		
	11	73	1.64						
	12	73	1.22	31	3.06				
	1	74	4.93	31	1.25				
2103C1	2	74	0.85						
	3	74	2.32	1	0.62				
	4	74	1.67	2	6.91				
	3	73	1.19	31	0.74				
	4	73	1.56	29	1.98				
	5	73	2.32	1	0.88	30	2.63		
	6	73	0.51						
	7	73	0.54	3	0.42				
	8	73	0.05	1	0.21	31	0.01		
	9	73	0.01						
	10	73	0.00	2	0.00	31	0.01		
	11	73	0.96						
2103D1	12	73	0.68	31	1.70				
	1	74	2.72	31	0.68				
	2	74	0.48						
	3	74	1.36	1	0.37				
	4	74	0.96	2	3.99				
	3	73	0.57	31	0.34				
	4	73	0.71	29	0.91				
	5	73	1.10	30	1.25				
	6	73	0.23						
	7	73	0.24	3	0.18				
	8	73	0.02	1	0.09	31	0.01		
	9	73	0.00						
	10	73	0.00	2	0.00	31	0.01		
	11	73	0.45						
	12	73	0.31	31	0.79				
	1	74	1.22	31	0.31				
	2	74	0.22						
	3	74	0.62	1	0.17				
	4	74	0.42	2	1.76				

TRIBUTARY FLOW INFORMATION FOR KENTUCKY

2/3/75

LAKE CODE 2103 HERRINGTON LAKE

MEAN MONTHLY FLOWS AND DAILY FLOWS (CMS)

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
2103E1	3	73	0.71	31	0.42				
	4	73	0.91	29	1.16				
	5	73	1.44	31	1.70				
	6	73	0.31						
	7	73	0.31	3	0.24				
	8	73	0.03	3	0.07				
	9	73	0.00	6	0.01				
	10	73	0.00	3	0.01				
	11	73	0.57	4	0.01				
	12	73	0.40						
	1	74	1.59	5	0.91	20	0.79		
	2	74	0.28	15	0.12				
2103ZZ	3	74	0.79	1	0.21	12	1.78		
	4	74	0.57						
	3	73	4.05						
	4	73	5.35						
	5	73	7.76						
	6	73	1.87						
	7	73	1.81						
	8	73	0.20						
	9	73	0.02						
	10	73	0.01						
	11	73	3.28						
	12	73	2.38						
	1	74	9.91						
	2	74	1.67						
	3	74	4.59						
	4	74	3.34						

APPENDIX D

PHYSICAL and CHEMICAL DATA

STORET RETRIEVAL DATE 75/01/27

210301
37 46 53.0 084 42 34.0
HERRINGTON LAKE
21167 KENTUCKY

11EPALES
3

2111202
0184 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00010 WATER TEMP CENT	00300 DO MG/L	00077 TRANSP SECCHI INCHES	00094 CONDUCTVY FIELD MICROMHO	00400 PH SU	00410 T ALK CAC03 MG/L	00610 NH3-N TOTAL MG/L	00625 TOT KJEL N MG/L	00630 NO2&NO3 N-TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P
73/05/26	11 30	0000	22.3		42	270	9.00	94	0.040	1.000	0.320	0.023
	11 30	0004	22.2	13.6		270	8.90	94	0.040	0.800	0.300	0.030
	11 30	0015	19.4	7.2		275	8.20	102	0.120	0.400	0.360	0.014
	11 30	0024	18.0	6.0		285	7.90	110	0.170	0.400	0.530	0.016
	11 30	0045	16.4	6.2		285	7.90	112	0.200	0.400	0.660	0.031
	11 30	0065	14.0	7.0		280	7.90	112	0.180	0.400	0.700	0.048
	11 30	0095	11.9	7.6		275	8.00	110	0.100	0.300	0.860	0.047
	11 30	0130	10.8	6.6		295	7.70	116	0.030	0.200K	1.340	0.076
	11 30	0180	7.0	6.0		285	7.50	117	0.040	0.300	1.530	0.110
73/08/20	12 15	0000	28.0	8.0	40	265	7.90	76	0.070	0.800	0.120	0.019
	12 15	0015	27.5	5.4		274	7.60	78	0.040	0.500	0.050	0.022
	12 15	0037	22.0	0.0		242	7.20	98	0.120	0.500	0.040	0.056
	12 15	0060	18.6			234						
	12 15	0100	16.1			223						
	12 15	0170	10.4	0.1		221	6.90	115	0.300	0.700	0.780	0.123
	12 15	0200	5.4	0.0		208	6.90	127	0.690	1.300	0.150	0.214
73/10/25	10 34	0000	21.2		67	247	8.00	102	0.060	0.700	0.040	0.053
	10 34	0005	21.2	4.8		247	7.90	100	0.050	0.800	0.020	0.042
	10 34	0015	21.4	5.0		244	7.90	100	0.050	0.700	0.020	0.038
	10 34	0030	21.1	4.6		244	7.80	101	0.060	0.700	0.030	0.046
	10 34	0050	20.9	2.8		247	7.70	102	0.080	0.200	0.030	0.052
	10 34	0070	18.1	0.6		214	7.40	102	0.040	0.200K	0.500	0.072
	10 34	0110	15.1	0.6		215	7.40	106	0.040	0.200K	1.020	0.101
	10 34	0150	12.4	0.6		219	7.30	125	0.390	0.600	0.580	0.148
	10 34	0185	6.8	0.6		205	7.30	133	0.810	1.300	0.110	0.196

K VALUE KNOWN TO BE
LESS THAN INDICATED

STORET RETRIEVAL DATE 75/01/27

210301
37 46 53.0 084 42 34.0
HERRINGTON LAKE
21167 KENTUCKY

11EPALES
3

2111202
0184 FEET DEPTH

DATE	TIME	DEPTH	00665	32217
FROM	OF		PHOS-TOT	CHLRPHYL
TO	DAY	FEET	MG/L P	A
				UG/L
73/05/26	11 30	0000	0.099	21.7
	11 30	0004	0.099	
	11 30	0015	0.042	
	11 30	0024	0.030	
	11 30	0045	0.041	
	11 30	0065	0.056	
	11 30	0095	0.053	
	11 30	0130	0.089	
	11 30	0180	0.150	
73/08/20	12 15	0000	0.087	9.4
	12 15	0015	0.043	
	12 15	0037	0.065	
	12 15	0170	0.123	
	12 15	0200	0.235	
73/10/25	10 34	0000	0.079	2.0
	10 34	0005	0.062	
	10 34	0015	0.059	
	10 34	0030	0.068	
	10 34	0050	0.076	
	10 34	0070	0.081	
	10 34	0110	0.106	
	10 34	0150	0.181	
	10 34	0185	0.445	

STORET RETRIEVAL DATE 75/01/27

210302
37 45 20.0 084 42 12.0
HERRINGTON LAKE
21167 KENTUCKY

11EPALES
3

2111202
0185 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00010 WATER TEMP CENT	00300 DO MG/L	00077 TRANSP SECCHI INCHES	00094 CNDUCTVY FIELD MICROMHO	00400 PH SU	00410 T ALK CAC03 MG/L	00610 NH3-N TOTAL MG/L	00625 TOT KJEL N MG/L	00630 NO2&NO3 N-TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P
73/05/26	12 35	0000	22.1		39	220	9.20	91	0.030	0.600	0.200	0.014
	12 35	0005	21.5	16.0		220	9.10	94	0.040	0.700	0.180	0.013
	12 35	0015	19.4	8.8		240	8.60	108	0.040	0.400	0.350	0.009
	12 35	0025	18.0	6.2		260	8.20	112	0.080	0.400	0.510	0.013
	12 35	0050	16.0	6.0		260	7.90	114	0.150	0.400	0.840	0.065
	12 35	0075	13.0	6.6		280	7.90	105	0.070	0.400	0.960	0.060
	12 35	0110	11.6	7.0		280	7.70	108	0.030	0.300	1.260	0.076
	12 35	0150	9.8	6.4		295	7.60	113	0.040	0.300	1.590	0.092
	12 35	0180	7.3	4.4		295	7.40	109	0.040	0.300	1.400	0.096
73/08/20	13 15	0000	28.1	7.6	108	236	7.80	75	0.050	0.900	0.040	0.014
	13 15	0010	27.9			236						
	13 15	0015	26.9	0.8		251	7.20	90	0.040	0.500	0.040	0.022
	13 15	0020	25.2			264						
	13 15	0060	18.5	0.0		222	7.00	101	0.110	0.500	0.490	0.034
	13 15	0100	16.0	0.4		217	6.90	103	0.050	0.300	1.400	0.109
	13 15	0150	13.1			208						
	13 15	0183	7.4	0.2		218	6.70	139	0.850	1.400	0.510	0.214
73/10/25	11 16	0000	21.0		106	230	7.60	98	0.060	0.600	0.040	0.037
	11 16	0010	20.9	4.6		230	7.60	98	0.050	0.300	0.040	0.031
	11 16	0020	20.9	4.6		231	7.60	98	0.040	0.400	0.040	0.030
	11 16	0050	20.9	4.6		230	7.60	99	0.040	0.200	0.040	0.032
	11 16	0070	18.1	0.4		211	7.30	102	0.040	0.300	0.540	0.114
	11 16	0110	15.4	0.6		217	7.20					
	11 16	0150	12.7	0.6		242	7.10	144	0.980	1.200	0.060	0.117
	11 16	0171	7.1	0.6		220	7.20					

STORET RETRIEVAL DATE 75/01/27

210302
37 45 20.0 084 42 12.0
HERRINGTON LAKE
21167 KENTUCKY

11EPALES
3

2111202
0185 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00665 PHOS-TOT MG/L P	32217 CHLRPHYL A UG/L
73/05/26	12 35	0000	0.053	26.5
	12 35	0005	0.062	
	12 35	0015	0.035	
	12 35	0025	0.027	
	12 35	0050	0.079	
	12 35	0075	0.071	
	12 35	0110	0.078	
	12 35	0150	0.097	
	12 35	0180	0.098	
73/08/20	13 15	0000	0.028	8.6
	13 15	0015	0.032	
	13 15	0060	0.035	
	13 15	0100	0.109	
	13 15	0183	0.294	
73/10/25	11 16	0000	0.052	1.7
	11 16	0010	0.046	
	11 16	0020	0.046	
	11 16	0050	0.047	
	11 16	0070	0.136	
	11 16	0150	0.188	

STORET RETRIEVAL DATE 75/01/27

210303
37 42 58.0 084 44 41.0
HERRINGTON LAKE
21167 KENTUCKY

11EPALES
3

2111202
0150 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00010 WATER TEMP CENT	00300 DO MG/L	00077 TRANSP SECCHI INCHES	00094 CNDUCTVY FIELD MICROMHO	00400 PH SU	00410 T ALK CAC03 MG/L	00610 NH3-N TOTAL MG/L	00625 TOT KJEL N MG/L	00630 NO2&NO3 N-TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P
73/05/26	14 40	0000	23.3		60	225	8.90	95	0.060	0.600	0.440	0.026
	14 40	0005	21.7	13.6		225	8.90	94	0.040	0.600	0.420	0.019
	14 40	0015	18.7	7.2		250	8.00	96	0.130	0.500	0.650	0.043
	14 40	0030	17.1	4.4		245	7.70	95	0.170	0.500	0.980	0.103
	14 40	0050	15.9	5.4		265	7.60	106	0.120	0.500	1.040	0.091
	14 40	0070	13.5	5.8		260	7.60	103	0.040	0.500	1.040	0.085
	14 40	0100	11.8	7.0		280	7.60	114	0.030	0.400	1.040	0.072
	14 40	0145	9.7	2.2		280	7.50	117	0.140	0.500	1.090	0.066
73/08/20	14 00	0000	28.0	7.6	66	223	7.80	84	0.060	0.700	0.090	0.012
	14 00	0015	26.8	1.6		251	7.10	97	0.040	0.500	0.050	0.029
	14 00	0021	25.3			274						
	14 00	0030	23.0	0.2		236	6.80	101	0.240	0.800	0.320	0.118
	14 00	0060	18.3	0.2		203	6.70	97	0.040	0.400	0.700	0.093
	14 00	0090	16.6	0.2		201	6.60	98	0.130	0.600	1.180	0.105
	14 00	0129	15.2	0.0		268	6.40	150	1.620	2.700	0.120	0.066
73/10/24	11 54	0000	20.8		65	228	7.40	106	0.070	0.500	0.040	0.049
	11 54	0005	20.8	3.0		229	7.40	106	0.060	0.200	0.020	0.041
	11 54	0015	20.8	3.0		228	7.40	106	0.050	0.200	0.020	0.040
	11 54	0030	20.8	2.8		227	7.40	105	0.060	0.200	0.020	0.040
	11 54	0055	20.6	0.8		231	7.30	106	0.110	0.400	0.020	0.043
	11 54	0070	18.1	0.6		212	7.10	108	0.140	0.400	0.100	0.146
	11 54	0114	15.2	0.4		282	7.00					

STORET RETRIEVAL DATE 75/01/27

210303
37 42 58.0 084 44 41.0
HERRINGTON LAKE
21167 KENTUCKY

11EPALES
3

2111202
0150 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00665 PHOS-TOT MG/L P	32217 CHLRPHYL A UG/L
73/05/26	14 40	0000	0.060	6.1
	14 40	0005	0.064	
	14 40	0015	0.070	
	14 40	0030	0.132	
	14 40	0050	0.108	
	14 40	0070	0.109	
	14 40	0100	0.080	
	14 40	0145	0.098	
73/08/20	14 00	0000	0.028	8.0
	14 00	0015	0.042	
	14 00	0030	0.139	
	14 00	0060	0.105	
	14 00	0090	0.219	
	14 00	0129	0.412	
73/10/24	11 54	0000	0.069	3.1
	11 54	0005	0.060	
	11 54	0015	0.057	
	11 54	0030	0.059	
	11 54	0055	0.072	
	11 54	0070	0.181	

STORET RETRIEVAL DATE 75/01/27

210304
37 40 16.0 684 41 24.0
HERRINGTON LAKE
21079 KENTUCKY

11EPALES
3

2111202
0034 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00010 WATER TEMP CENT	00300 DO MG/L	00077 TRANSP SECCHI INCHES	00094 CONDUCTVY FIELD MICROMHO	00400 PH SU	00410 T ALK CAC03 MG/L	00610 NH3-N TOTAL MG/L	00625 TOT KJEL N MG/L	00630 NO2&NO3 N-TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P
73/05/26	15 40	0000	23.9		37	230	9.10	99	0.050	0.600	0.500	0.028
	15 40	0005	20.0	7.8		230	8.50	99	0.150	0.500	0.790	0.085
	15 40	0015	18.4	5.6		190	7.90	78	0.200	0.600	0.970	0.076
	15 40	0030	18.0	4.2		210	7.70	87	0.260	0.600	0.990	0.095
73/08/20	16 15	0000	27.9	5.8	40	236	7.80	94	0.060	1.000	0.060	0.012
	16 15	0005	27.5			244						
	16 15	0015	26.9	0.4		262	7.00	99	0.090	0.800	0.090	0.032
	16 15	0017	26.6			292						
	16 15	0023	24.7	0.1		318	6.70	134	0.720	1.500	0.080	0.089
73/10/25	12 21	0000	19.4		18	247	7.60	117	0.060	1.800	0.040	0.016
	12 21	0002	19.4	6.2		247	7.60	117	0.050	1.100	0.040	0.014
	12 21	0005	18.9	5.8		250	7.50	118	0.060	0.900	0.060	0.012

DATE FROM TO	TIME OF DAY	DEPTH FEET	00665 PHOS-TOT MG/L P	32217 CHLRPHYL A UG/L
73/05/26	15 40	0000	0.125	16.2
	15 40	0005	0.145	
	15 40	0015	0.231	
	15 40	0030	0.224	
73/08/20	16 15	0000	0.044	11.0
	16 15	0015	0.075	
	16 15	0023	0.284	
73/10/25	12 21	0000	0.246	64.6
	12 21	0002	0.149	
	12 21	0005	0.140	

APPENDIX E

TRIBUTARY and WASTEWATER TREATMENT PLANT DATA

STORET RETRIEVAL DATE 75/01/13

2103A1 LS2103A1
 37 47 00.0 084 42 00.0
 DIX RIVER
 21 7.5 WILMORE
 O/HERRINGTON LAKE
 BANK BELO DAM LIGHT DUTY DAM ACCESS RD
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE	TIME	DEPTH	00630	00625	00610	00671	00665
FROM	OF		NH3-N	TOT KJEL	NH3-N	PHOS-DIS	PHOS-TOT
TO	DAY	FEET	N-TOTAL	N	TOTAL	ORTHO	
			MG/L	MG/L	MG/L	MG/L P	MG/L P
73/03/31	11	10	1.160	0.270	0.008	0.060	0.070
73/05/01	16	30	1.060	0.310	0.014	0.056	0.080
73/05/31	15	00	0.830	0.380	0.125	0.043	0.050
73/07/03	14	00	0.930	0.540	0.020	0.036	0.040
73/08/03	12	15	0.930	3.000	0.084	0.069	0.090
73/09/06	08	00	0.830	4.400	0.147	0.073	0.108
73/10/03	16	00	0.670	2.400	0.092	0.082	0.105
73/11/04	09	00	0.368	1.700	0.064	0.078	0.110
73/12/01	06	30	0.430	1.000	0.028	0.064	0.105
74/01/05	12	40	1.680	2.500	0.115	0.092	0.130
74/02/15	16	30	1.340	2.600	0.050	0.090	0.125
74/03/01	09	00	1.340	1.200	0.035	0.090	0.135
74/03/16	09	30	1.360	1.900	0.040	0.085	0.113
74/04/02	11	00	4.400	1.500	0.065	0.075	

STORET RETRIEVAL DATE 75/01/13

2103A3 LS2103A3
 37 38 30.0 084 39 30.0
 DIX RIVER
 21 7.5 BRYANTSVILLE
 T/HERRINGTON LAKE
 HWY, 52 BRDG NEAR GAGING STATION
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 NO2&NO3 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
73/03/31	09 00		0.790	0.420	0.020	0.042	0.085
73/05/01	12 30		1.100	0.190	0.009	0.037	0.075
73/05/30	14 30		0.960	0.560	0.044	0.050	0.145
73/07/03	08 30		1.100	1.890	0.230	0.070	0.205
73/08/01	13 00		0.980	2.000	0.069	0.078	0.160
73/08/31	11 00		0.020	1.200	0.039	0.079	0.150
73/10/02	09 30		0.037	0.900	0.048	0.052	0.110
73/10/31	10 12		0.132	2.500	0.048	0.052	0.095
73/12/31	10 15		1.760	0.700	0.036	0.060	0.135
74/01/05	09 30		1.520	0.700	0.080	0.065	0.090
74/03/01	12 30		1.200	0.800	0.075	0.080	0.120

STORET RETRIEVAL DATE 75/01/13

210381 LS210381
 37 38 30.0 084 43 30.0
 CLARKS RUN
 21 7.5 BRYANTSVILLE
 T/HARRINGTON LAKE
 FORD ON RD 5 MI N OF ST HWY52
 ILEPALES 2111204
 4 0000 FEET DEPTH

DATE	TIME	DEPTH	00630 NO2&N03 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
FROM TO	OF DAY	FEET					
73/03/31	09 25		1.960	0.860	0.210	0.480	0.640
73/05/01	12 00		2.400	0.480	0.120	0.390	0.550
73/05/30	14 00		1.360	4.300	0.058	0.340	2.400
73/07/03	09 00		2.400	1.400	0.168	0.620	0.840
73/08/01	13 30		1.660	1.630	0.138	0.780	0.940
73/08/31	10 00		2.100	4.350	0.480	3.500	4.100
73/10/02	16 00		2.000	9.000	5.100	4.200	4.700
73/10/31	10 00		2.400	5.100	2.200	3.200	
73/12/31	10 35		3.700	0.800	0.096	0.152	0.300
74/01/31	10 00		3.300	2.100	0.440	0.360	0.520
74/03/01	13 00		2.600	1.500	0.400	0.620	0.880
74/04/02	10 00		2.200	1.500	0.090	0.180	0.530

STORET RETRIEVAL DATE 75/01/20

2103C1 LS2103C1
 37 41 30.0 084 46 00.0
 MOCKS BRANCH
 21 7.5 DANVILLE
 T/HERRINGTON LAKE
 BRDG ON HWY 33 2.5 MI N OF DANVILLE
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 NO2&NO3 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P	50051 FLOW RATE INST MGD	50053 CONDUIT FLOW-MGD MONTHLY
73/03/31	10 30		2.000	0.320	0.013	0.132	0.210		
73/04/29	08 55		0.430	0.130	0.026				
73/05/01	12 30		1.760	0.480	0.011	0.095	0.150		
73/05/30	13 00		2.000	1.050	0.310	0.310	0.610		
73/07/03	09 30		2.100	8.500	0.870	0.200	0.315		
73/08/01	14 30		0.980	1.760	0.058	0.154	0.210		
73/08/31	09 30		0.490	2.100	0.205	0.220	0.345		
73/10/02	11 00		3.700	3.000	0.092	0.560	0.735		
73/10/31	09 00		1.000	1.850	0.184	0.368	0.520		
73/12/31	11 15		4.900	0.600	0.036	0.200	0.270		
74/01/31	11 00		4.200	1.100	0.030	0.165	0.230		
74/03/01	13 30		2.700	0.400	0.040	0.130	0.200		
74/04/02	09 00		2.300	1.700	0.040	0.250	0.830		

STORET RETRIEVAL DATE 75/01/20

210301
 37 40 55.0 084 44 45.0
 SPEARS CREEK
 21 7.5 BRYANTSVILLE
 T/LAKE HERRINGTON
 BANK END OF SPEARS CREEK RD
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 N02&N03 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P	50051 FLOW RATE INST MGD	50053 CONDUIT FLOW-MGD MONTHLY
73/03/31	09 50		1.820	0.150	0.008	0.230	0.250		
73/04/29	11 00		1.780	0.200	0.020	0.210	0.250		
73/05/30	13 30		0.720	1.150	0.058	0.110	0.250		
73/07/03	09 00		0.033	1.050	0.044	0.026	0.095		
73/08/01	14 30		0.010	1.150	0.085	0.028	0.090		
73/08/31	08 30		2.200	1.760	0.100	0.350	0.390		
73/10/02	10 30		2.300	6.000	0.150	0.400	0.575		
73/10/31	09 30		1.120	0.500	0.028	0.336	0.380		
73/12/31	11 00		3.600	0.500	0.036	0.264	0.345		
74/01/31	10 30		2.400	0.500	0.015	0.175	0.220		
74/03/01	13 25		2.500	0.300	0.012	0.230	0.300		
74/04/02	09 30		2.100	1.100	0.030	0.290	0.780		

STORET RETRIEVAL DATE 75/01/13

2103E1
 37 45 15.0 084 44 25.0
 CANE RUN
 21- 7.5 WILMORE
 T/LAKE HERRINGTON
 BANK 3 MI SE OF BURGIN
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 NO2&NO3 N-TOTAL MG/L	00625 TOT KjEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
73/03/31	13 30		2.040	0.440	0.022	0.220	0.290
73/04/29	10 00		2.200	0.100K	0.012	0.210	0.260
73/05/31	16 00		2.400	0.440	0.115	0.231	0.340
73/07/03	14 00		2.700	0.540	0.026	0.273	0.410
73/08/03	14 00		1.540	0.100K	0.026	0.245	0.280
73/09/06	11 00		0.400	0.440	0.073	0.336	0.380
73/10/03	14 00		0.390	0.550	0.075	0.420	0.500
73/11/04	09 20		1.090	2.400	0.216	0.320	0.390
74/01/05	12 00		3.400	0.300	0.015	0.250	0.315
74/02/15	16 00		1.600	2.200	0.050	0.120	0.160
74/03/01	10 00		2.100	4.400	0.090	0.230	0.390

STORET RETRIEVAL DATE 75/02/03

2103AA TF2103AA P003446
37 36 50.0 084 35 10.0
LANCASTER
21 7.5 LANCASTER
T/HERRINGTON LAKE
WHITE OAK CREEK/DIA RIVER
11EPALES 2141204
4 0000 FEET DEPTH

DATE	TIME	DEPTH	00630 NO2&NO3 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P	50051 FLOW RATE INST MGD	50053 CONDUIT FLOW-MGD MONTHLY
FROM	OF								
TO	DAY	FEET							
74/07/02	09	30							
CP(T)-			0.320	23.000	14.000	30.000	30.000	0.250	0.225
74/07/02	11	30							

