

**U.S. ENVIRONMENTAL PROTECTION AGENCY
NATIONAL EUTROPHICATION SURVEY
WORKING PAPER SERIES**



REPORT
ON
BLEWETT FALLS LAKE
ANSON AND RICHMOND COUNTIES
NORTH CAROLINA
EPA REGION IV
WORKING PAPER No. 378

PACIFIC NORTHWEST ENVIRONMENTAL RESEARCH LABORATORY

An Associate Laboratory of the

NATIONAL ENVIRONMENTAL RESEARCH CENTER - CORVALLIS, OREGON

and

NATIONAL ENVIRONMENTAL RESEARCH CENTER - LAS VEGAS, NEVADA

REPORT

ON

BLEWETT FALLS LAKE

ANSON AND RICHMOND COUNTIES

NORTH CAROLINA

EPA REGION IV

WORKING PAPER No. 378

828

WITH THE COOPERATION OF THE

NORTH CAROLINA DEPARTMENT OF NATURAL AND ECONOMIC RESOURCES

AND THE

NORTH CAROLINA NATIONAL GUARD

JUNE, 1975

CONTENTS

	<u>Page</u>
Foreword	ii
List of North Carolina Study Lakes	iv
Lake and Drainage Area Map	v
 <u>Sections</u>	
I. Conclusions	1
II. Lake and Drainage Basin Characteristics	5
III. Lake Water Quality Summary	7
IV. Nutrient Loadings	11
V. Literature Reviewed	17
VI. Appendices	18

F O R E W O R D

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nationwide threat of accelerated eutrophication to fresh water lakes and reservoirs.

OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

- a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.
- b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.
- c. With such a transformation, an assessment of the potential for eutrophication control can be made.

LAKE ANALYSIS

In this report, the first stage of evaluation of lake and watershed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning [§303(e)], water quality criteria/standards review [§303(c)], clean lakes [§314(a,b)], and water quality monitoring [§106 and §305(b)] activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

ACKNOWLEDGMENT

The staff of the National Eutrophication Survey (Office of Research & Development, U. S. Environmental Protection Agency) expresses sincere appreciation to the North Carolina Department of Natural and Economic Resources for professional involvement and to the North Carolina National Guard for conducting the tributary sampling phase of the Survey.

Lewis R. Martin, Director of the Division of Environmental Management; Darwin L. Coburn, Chief of the Water Quality Section; and Julian R. Taylor, Supervisor of the Monitoring Program Unit; provided invaluable lake documentation and counsel during the Survey, reviewed the preliminary reports, and provided critiques most useful in the preparation of this Working Paper series.


Major General William M. Buck, formerly Adjutant General of North Carolina, and Project Officer Colonel Arthur J. Bouchard, who directed the volunteer efforts of the North Carolina National Guardsmen, are also gratefully acknowledged for their assistance to the Survey.

NATIONAL EUTROPHICATION SURVEY

STUDY LAKES

STATE OF NORTH CAROLINA

<u>LAKE NAME</u>	<u>COUNTY</u>
Badin	Montgomery, Stanly
Blewett Falls	Anson, Richmond
Chatuge	Clay, NC; Towns, GA
Fontana	Graham, Swain
Hickory	Alexander, Caldwell, Catawba
High Rock	Davidson, Rowan
Hiwassee	Cherokee
James	Burke, McDowell
John H. Kerr (Nut Bush Creek)	Granville, Vance, Warren, NC; Halifax, Mecklenburg, VA
Junaluska	Haywood
Lookout Shoals	Alexander, Catawba, Iredell
Mountain Island	Gaston, Mecklenburg
Norman	Catawba, Iredell, Lincoln, Mecklenburg
Rhodhiss	Burke, Caldwell
Santeetlah	Graham
Tillery	Montgomery, Stanly
Waccamaw	Columbus
Waterville	Haywood
Wylie	Gaston, Mecklenburg, NC; York, SC



BLEWETT FALLS LAKE

STORET NO. 3702

I. CONCLUSIONS

A. Trophic Condition:

Survey data show that Blewett Falls Lake is eutrophic. However, since the lake has a mean hydraulic retention time of only seven days, it more closely resembles a slow-moving river; and the term "over-enriched" may be a more appropriate description of the condition of this lake.

Blewett Falls Lake ranked thirteenth in overall trophic condition when the 16 North Carolina lakes sampled in 1973 were compared using a combination of six parameters*. Thirteen lakes had less and one had the same median total phosphorus, 14 had less median dissolved phosphorus, 13 had less median inorganic nitrogen, but only four lakes had less mean chlorophyll a. The relatively low concentrations of chlorophyll a and the high nutrient levels indicate primary productivity may have been light-limited. Fourteen of the lakes had greater mean Secchi disc transparency, and Survey limnologists noted that the lake was quite turbid at all sampling times. However, it is likely that primary productivity would not be very high in any case because of the very short hydraulic retention time of the lake.

* See Appendix A.

Depression of dissolved oxygen with depth occurred at sampling station 3 in July, and dissolved oxygen in the shallowest samples collected in September did not exceed 77% of saturation.

Survey limnologists did not observe rooted aquatic vegetation or concentrations of algae during their visits to the lake.

B. Rate-Limiting Nutrient:

Because of a loss of nutrients in the assay sample from the time of collection to the beginning of the assay, the results are not representative of conditions in the lake at the time the sample was taken (03/23/73).

The lake data indicate phosphorus limitation at all three stations in March and July and at station 1 in September also. However, nitrogen limitation is indicated at stations 2 and 3 in September.

C. Nutrient Controllability:

1. Point sources--The phosphorus contributions of the listed point sources amounted to only 2.3% of the total load to Blewett Falls Lake. However, there are at least eight municipal wastewater treatment plants in the Rocky River drainage beyond the 40-kilometer limit of the Survey* (Anonymous,

* See Working Paper No. 175, "...Survey Methods, 1973-1976".

1971). These plants serve a combined 1970 population of 47,281 people.

Also there are major point sources in the Yadkin River drainage upstream from Lake Tillery (Anonymous, 1973); and during the sampling year, some 264,000 kg of total phosphorus were exported from Lake Tillery to Blewett Falls Lake (see Working Paper No. 390, "Report on Lake Tillery"). However, the total phosphorus load measured in the Pee Dee River at the inlet to Blewett Falls Lake, below the confluence of the Rocky River, amounted to more than 900,000 kg during the sampling year. If most of the difference is attributable to the Rocky River phosphorus load, it appears that drainage contributed on the order of 600,000 kg of total phosphorus during the year. Considering the estimated 4,000 km² drainage area of the Rocky River, this phosphorus load seems quite high (150 kg/km²), and a need for a more-detailed study of this drainage and point sources is indicated.

The present Blewett Falls Lake phosphorus loading rate of 91 g/m²/yr is 18 times that proposed by Vollenweider (Vollenweider and Dillon, 1974) as a eutrophic rate (see page 16). However, the mean hydraulic retention time of the lake is a very short seven days, and it is probable that Vollenweider's model is not applicable to this water body.

As noted in the discussion of trophic condition, the primary productivity of Blewett Falls Lake may have been light-limited because of turbidity. If the turbidity observed during Survey sampling is a persistent characteristic of this lake, point-source phosphorus control would not be expected to result in a significant improvement in the trophic condition. However, if the turbidity is not a typical condition, every effort should be made to minimize the phosphorus inputs to the lake.

2. Non-point sources--The phosphorus export rate of the Pee Dee River was nearly six times the mean of the rates of the other Blewett Falls Lake tributaries (see page 16). This high rate probably was due in large part to point-source inputs in the Rocky River and Yadkin River drainages as noted above.

II. LAKE AND DRAINAGE BASIN CHARACTERISTICS[†]

A. Lake Morphometry^{††}:

1. Surface area: 10.36 kilometers².
2. Mean depth: 11.6 meters.
3. Maximum depth: 11.9 meters.
4. Volume: 120×10^6 m³.
5. Mean hydraulic retention time: 7 days.

B. Tributary and Outlet:

(See Appendix C for flow data)

1. Tributaries -

<u>Name</u>	<u>Drainage area (km²)*</u>	<u>Mean flow (m³/sec)*</u>
Pee Dee River	16,397.3	198.0
Cedar Creek	40.4	0.3
Savannah Creek	37.0	0.3
Smith Creek	26.2	0.2
Mountain Creek	194.2	2.6
Little River	911.7	9.6
Dry Creek	20.7	0.2
Cartledge Creek	78.5	1.2
Minor tributaries & immediate drainage -	<u>76.9</u>	<u>1.2</u>
Totals	17,782.9	213.6

2. Outlet -

Pee Dee River	17,793.3**	213.6**
---------------	------------	---------

[†] Table of metric conversions--Appendix B.

^{††} Park, 1974.

* For limits of accuracy, see Working Paper No. 175.

** Includes area of lake; outflow adjusted to equal sum of inflows.

C. Precipitation*:

1. Year of sampling: 140.6 centimeters.
2. Mean annual: 121.8 centimeters.

* See Working Paper No. 175.

III. LAKE WATER QUALITY SUMMARY

Blewett Falls Lake was sampled three times during the open-water season of 1973 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from three stations on the lake and from one or more depths at each station (see map, page v). During each visit, a single depth-integrated (4.6 m or near bottom to surface) sample was composited from the three stations for phytoplankton identification and enumeration; and during the first visit, a single 18.9-liter depth-integrated sample was composited for algal assays. Also each time, a depth-integrated sample was collected from each of the stations for chlorophyll a analysis. The maximum depths sampled were 1.2 meters at station 1, 6.4 meters at station 2, and 9.1 meters at station 3.

The lake sampling results are presented in full in Appendix D and are summarized in the following table.

A. SUMMARY OF PHYSICAL AND CHEMICAL CHARACTERISTICS FOR BLEWETT FALLS LAKE
STORED CODE 3702

PARAMETER	1ST SAMPLING (3/23/73)				2ND SAMPLING (7/ 5/73)				3RD SAMPLING (9/19/73)			
	3 SITES				3 SITES				3 SITES			
	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN
TEMP (C)	11.5 - 12.5	12.5	12.0	25.5 - 28.5	26.4	26.0	26.0 - 26.3	26.2	26.2	26.0 - 26.2	26.0	26.2
DISS OXY (MG/L)	9.8 - 10.2	10.0	10.0	4.2 - 7.9	5.3	4.9	5.2 - 6.2	6.0	6.1	5.2 - 6.2	6.0	6.1
CHLOROBY (MICROMO)	80. - 90.	84.	83.	60. - 90.	78.	80.	62. - 158.	107.	94.	62. - 158.	107.	94.
PH (STAND UNITS)	7.1 - 7.4	7.3	7.2	6.8 - 8.9	7.3	6.9	6.5 - 7.1	6.7	6.7	6.5 - 7.1	6.7	6.7
TOT ALK (MG/L)	16. - 19.	18.	18.	17. - 27.	22.	23.	24. - 36.	30.	28.	24. - 36.	30.	28.
TOT P (MG/L)	0.087 - 0.109	0.100	0.100	0.043 - 0.092	0.065	0.064	0.020 - 0.184	0.078	0.058	0.020 - 0.184	0.078	0.058
ORTHOP (MG/L)	0.032 - 0.045	0.043	0.040	0.006 - 0.028	0.013	0.012	0.015 - 0.150	0.058	0.035	0.015 - 0.150	0.058	0.035
NO2+NO3 (MG/L)	0.500 - 0.570	0.543	0.540	0.290 - 0.640	0.462	0.450	0.230 - 0.380	0.284	0.270	0.230 - 0.380	0.284	0.270
AMMONIA (MG/L)	0.150 - 0.190	0.163	0.160	0.080 - 0.140	0.124	0.130	0.040 - 0.080	0.064	0.060	0.040 - 0.080	0.064	0.060
KJEL N (MG/L)	0.400 - 1.100	0.645	0.600	0.300 - 0.900	0.489	0.400	0.260 - 0.400	0.262	0.200	0.260 - 0.400	0.262	0.200
INHAB N (MG/L)	0.580 - 0.740	0.707	0.700	0.430 - 0.770	0.587	0.570	0.270 - 0.460	0.347	0.340	0.270 - 0.460	0.347	0.340
TOTAL N (MG/L)	0.430 - 1.640	1.188	1.170	0.760 - 1.190	0.951	0.930	0.430 - 0.780	0.546	0.470	0.430 - 0.780	0.546	0.470
CHLORYL A (MG/L)	2.3 - 2.6	2.5	2.5	2.2 - 6.4	4.0	2.9	2.1 - 8.9	6.0	7.1	2.1 - 8.9	6.0	7.1
SECCHI (METERS)	0.3 - 0.3	0.3	0.3	0.3 - 0.5	0.4	0.5	0.3 - 1.5	1.1	1.0	0.3 - 1.5	1.1	1.0

B. Biological characteristics:

1. Phytoplankton -

<u>Sampling Date</u>	<u>Dominant Genera</u>	<u>Algal units per ml</u>
03/23/73	1. <i>Synedra</i> 2. <i>Oscillatoria</i> 3. <i>Cryptomonas</i> 4. <i>Tabellaria</i> 5. <i>Scenedesmus</i> Other genera	30 29 22 7 7 <u>15</u>
	Total	110
07/05/73	1. <i>Melosira</i> 2. <i>Nitzschia</i> 3. <i>Cryptomonas</i> 4. <i>Oscillatoria</i> 5. <i>Stephanodiscus</i> Other genera	202 130 72 58 58 <u>171</u>
	Total	691
09/19/73	1. <i>Raphidiopsis</i> 2. Flagellates 3. <i>Melosira</i> 4. <i>Nitzschia</i> 5. <i>Synedra</i> Other genera	1,173 156 118 98 59 <u>155</u>
	Total	1,759

2. Chlorophyll a -

<u>Sampling Date</u>	<u>Station Number</u>	<u>Chlorophyll a ($\mu\text{g/l}$)</u>
03/23/73	01	2.5
	02	2.3
	03	2.6
07/05/73	01	2.2
	02	2.9
	03	6.9
09/19/73	01	2.1
	02	7.1
	03	8.9

C. Limiting Nutrient Study:

A 30% loss of orthophosphorus and 61% loss of inorganic nitrogen occurred in the assay sample between the time of collection and the beginning of the assay, and the assay results are not representative of conditions in the lake at the time the sample was taken (03/23/73).

The lake data indicate phosphorus limitation at all three sampling stations in March and July and at station 1 in September also (the mean inorganic nitrogen/orthophosphorus ratios were 18/1 or greater). However, the data indicate nitrogen limitation at stations 2 and 3 in September (the mean N/P ratios were 4/1 and 9/1, respectively).

IV. NUTRIENT LOADINGS
(See Appendix E for data)

For the determination of nutrient loadings, the North Carolina National Guard collected monthly near-surface grab samples from each of the tributary sites indicated on the map (page v), except for the high runoff months of January and February when two samples were collected. Sampling was begun in March, 1973, and was completed in March, 1974.

Through an interagency agreement, stream flow estimates for the year of sampling and a "normalized" or average year were provided by the North Carolina District Office of the U.S. Geological Survey for the tributary sites nearest the lake.

Except for Smith Creek, nutrient loads for sampled tributaries were determined by using a modification of a U.S. Geological Survey computer program for calculating stream loadings*. Nutrient loads shown are those measured minus point-source loads, if any. However, the estimated phosphorus load from Lilesville exceeded the load measured in Smith Creek at station D-1, and the non-point nutrient contributions of this stream were estimated.

Nutrient loads for Smith Creek and unsampled "minor tributaries and immediate drainage" ("ZZ" of U.S.G.S.) were estimated using the means of the nutrient exports, in kg/km²/year, at stations B-1, C-1, G-1, and H-1 and multiplying the means by the Smith Creek and ZZ areas in km².

* See Working Paper No. 175.

The operators of the Biscoe and Wadesboro wastewater treatment plants provided monthly effluent samples and corresponding flow data (the Wadesboro plant does not impact Blewett Falls Lake; however, the analytical data are included in Appendix E for the record). The operators of the Norwood, Mount Gilead, Ellerbe, Lilesville, Troy, and Albemarle wastewater treatment plants did not participate in the Survey, and nutrient loads were estimated at 1.134 kg P and 3.401 kg N/capita/year.

Other wastewater treatment plants are located within the direct drainage basin of the lake (see map, page v). Nutrient loads from these plants were not determined since they are well beyond the 40-kilometer limit of the Survey. Also, major point sources are located in the Yadkin River drainage upstream from Lake Tillery which also impact Blewett Falls Lake. The nutrient contributions of all of these sources are included in the Pee Dee River loads measured at station A-2, however.

A. Waste Sources:

1. Known municipal -

<u>Name</u>	<u>Pop. Served</u>	<u>Treatment</u>	<u>Mean Flow (m³/d)</u>	<u>Receiving Water</u>
Norwood	1,896*	Imhoff tank	605.6**	Rocky River
Biscoe	1,200†	act. sludge	567.8	Little River
Mt. Gilead	1,286*	stab. pond	378.5**	Clarks Creek/ Pee Dee River
Ellerbe	913*	stab. pond	345.6 ^{††}	Little Mountain Creek
Lilesville	641**	Imhoff tank	242.6 ^{††}	Smith Creek
Troy	2,429*	stab. pond	794.9**	Denson Creek/ Little River
Albermarle	11,126*	trickling filter	26,495.0**	Long Creek

2. Known industrial - None

* 1970 Census.

** Anonymous, 1971.

† STP questionnaire.

†† Estimate based on 0.3785 m³/capita/day.

B. Annual Total Phosphorus Loading - Average Year:

1. Inputs -

<u>Source</u>	<u>kg P/ yr</u>	<u>% of total</u>
a. Tributaries (non-point load) -		
Pee Dee River	907,845	96.2
Cedar Creek	270	<0.1
Savannah Creek	290	<0.1
Smith Creek	230	<0.1
Mountain Creek	2,410	0.3
Little River	8,705	0.9
Dry Creek	255	<0.1
Cartledge Creek	655	0.1
b. Minor tributaries & immediate drainage (non-point load) -		675
		0.1
c. Known municipal STP's -		
Norwood	2,150	0.2
Biscoe	1,280	0.1
Mount Gilead	1,460	0.2
Ellerbe	1,035	0.1
Lillesville	725	0.1
Troy	2,755	0.3
Albemarle	12,615	1.4
d. Septic tanks - Unknown		-
		-
e. Known industrial - None		-
		-
f. Direct precipitation* -		<u>180</u>
		<u><0.1</u>
Total	943,535	100.0

2. Outputs -

Lake outlet - Pee Dee River 517,685

3. Net annual P accumulation - 425,850 kg.

* See Working Paper No. 175.

C. Annual Total Nitrogen Loading - Average Year:

1. Inputs -

<u>Source</u>	<u>kg N/ yr</u>	<u>% of total</u>
a. Tributaries (non-point load) -		
Pee Dee River	7,711,655	93.1
Cedar Creek	6,280	0.1
Savannah Creek	7,305	0.1
Smith Creek	6,055	0.1
Mountain Creek	95,875	1.2
Little River	333,285	4.0
Dry Creek	4,965	0.1
Cartledge Creek	26,025	0.3
b. Minor tributaries & immediate drainage (non-point load) -	17,770	0.2
c. Known municipal STP's -		
Norwood	6,450	0.1
Biscoe	1,275	<0.1
Mount Gilead	4,375	<0.1
Ellerbe	3,105	<0.1
Lilesville	2,180	<0.1
Troy	8,260	0.1
Albemarle	37,840	0.5
d. Septic tanks - Unknown	-	-
e. Known industrial - None	-	-
f. Direct precipitation* -	<u>11,185</u>	<u>0.1</u>
Total	8,283,885	100.0

2. Outputs -

Lake outlet - Pee Dee River 7,005,540

3. Net annual N accumulation - 1,278,345 kg.

* See Working Paper No. 175.

D. Mean Annual Non-point Nutrient Export by Subdrainage Area:

<u>Tributary</u>	<u>kg P/km²/yr</u>	<u>kg N/km²/yr</u>
Pee Dee River	55	470
Cedar Creek	7	155
Savannah Creek	8	197
Smith Creek	9	231
Mountain Creek	12	494
Little River	10	366
Dry Creek	12	240
Cartledge Creek	8	332

E. Yearly Loading Rates:

In the following table, the existing phosphorus loading rates are compared to those proposed by Vollenweider (Vollenweider and Dillon, 1974). Essentially, his "dangerous" rate is the rate at which the receiving water would become eutrophic or remain eutrophic; his "permissible" rate is that which would result in the receiving water remaining oligotrophic or becoming oligotrophic if morphometry permitted. A mesotrophic rate would be considered one between "dangerous" and "permissible".

Note that Vollenweider's model may not be applicable to water bodies with short hydraulic retention times.

	Total Phosphorus		Total Nitrogen	
	Total	Accumulated	Total	Accumulated
grams/m ² /yr	91.07	41.11	799.6	123.4

Vollenweider loading rates for phosphorus (g/m²/yr) based on mean depth and mean hydraulic retention time of Blewett Falls Lake:

"Dangerous" (eutrophic rate)	5.00
"Permissible" (oligotrophic rate)	2.50

V. LITERATURE REVIEWED

Anonymous, 1971. Inventory of municipal waste facilities. EPA Publ. OWP-1, vol. 4, Washington, DC.

Park, David, 1974. Personal communication (lake morphometry). NC Dept. of Nat. & Econ. Resources, Raleigh.

Vollenweider, R. A., and P. J. Dillon, 1974. The application of the phosphorus loading concept to eutrophication research. Natl. Res. Council of Canada Publ. No. 13690, Canada Centre for Inland Waters, Burlington, Ontario.

Weiss, Charles M., 1972. A proposal to the Water Resources Research Institute of the University of North Carolina. The trophic state of North Carolina lakes, covering the period July 1, 1972 to June 30, 1973. U. of North Carolina, Chapel Hill.

VI. APPENDICES

APPENDIX A

LAKE RANKINGS

LAKES RANKED BY INDEX NOS.

RANK	LAKE CODE	LAKE NAME	INDEX NO
1	3719	LAKE WACCAMAW	534
2	3716	SANTEELAH LAKE	446
3	3711	MOUNTAIN ISLAND LAKE	419
4	3707	HIWASSEE LAKE	414
5	3704	FONTANA LAKE	392
6	3713	LAKE NORMAN	346
7	3708	LAKE JAMES	334
8	3710	LOOKOUT SHOALS	327
9	3715	RHODHMISS LAKE	296
10	3705	LAKE HICKORY	283
11	3717	LAKE TILLERY	246
12	3709	LAKE JUNALUSKA	220
13	3702	BLEWETT FALLS LAKE	200
14	3718	WATERVILLE RESERVOIR	140
15	3701	BADIN LAKE	124
16	3706	HIGH ROCK LAKE	76

PERCENT OF LAKES WITH HIGHER VALUES (NUMBER OF LAKES WITH HIGHER VALUES)

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLORA	15- MIN DO	MEDIAN DISS ORTHO P	INDEX NO
3701	BADIN LAKE	33 (5)	7 (1)	27 (4)	27 (4)	3 (0)	27 (4)	124
3702	BLEWETT FALLS LAKE	7 (1)	13 (2)	7 (1)	73 (11)	93 (14)	7 (1)	200
3704	FONTANA LAKE	100 (15)	33 (5)	93 (14)	100 (15)	3 (0)	63 (9)	392
3705	LAKE HICKORY	27 (4)	60 (9)	53 (8)	13 (2)	80 (12)	50 (7)	283
3706	HIGH ROCK LAKE	13 (2)	20 (3)	0 (0)	0 (0)	23 (2)	20 (3)	76
3707	HIWASSEE LAKE	87 (13)	80 (12)	87 (13)	47 (7)	50 (7)	63 (9)	414
3708	LAKE JAMES	60 (9)	87 (13)	80 (12)	7 (1)	23 (2)	77 (11)	334
3709	LAKE JUNALUSKA	47 (7)	27 (4)	43 (6)	20 (3)	50 (7)	33 (5)	220
3710	LOOKOUT SHOALS	53 (8)	47 (7)	60 (9)	67 (10)	60 (9)	40 (6)	327
3711	MOUNTAIN ISLAND LAKE	73 (11)	73 (11)	43 (6)	53 (8)	87 (13)	90 (13)	419
3713	LAKE NORMAN	67 (10)	53 (8)	73 (11)	40 (6)	23 (2)	90 (13)	346
3715	RHOHMISS LAKE	20 (3)	67 (10)	33 (5)	93 (14)	70 (10)	13 (2)	296
3716	SANTEELAH LAKE	93 (14)	93 (14)	100 (15)	60 (9)	23 (2)	77 (11)	446
3717	LAKE TILLERY	40 (6)	40 (6)	13 (2)	33 (5)	70 (10)	50 (7)	246
3718	WATERVILLE RESERVOIR	0 (0)	0 (0)	20 (3)	80 (12)	40 (6)	0 (0)	140
3719	LAKE WACCAMAW	80 (12)	100 (15)	67 (10)	87 (13)	100 (15)	100 (15)	534

LAKE DATA TO BE USED IN RANKINGS

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLORA	15- MIN DO	MEDIAN DISS ORTHO P
3701	BADIN LAKE	0.042	0.680	466.750	7.190	14.900	0.012
3702	BLEWETT FALLS LAKE	0.090	0.655	476.889	4.167	10.800	0.034
3704	FONTANA LAKE	0.011	0.550	392.650	3.438	14.900	0.007
3705	LAKE HICKORY	0.047	0.320	461.000	7.275	13.400	0.008
3706	HIGH ROCK LAKE	0.090	0.580	477.454	14.283	14.800	0.017
3707	HIWASSEE LAKE	0.015	0.240	420.555	5.678	14.200	0.007
3708	LAKE JAMES	0.020	0.160	428.866	7.660	14.800	0.006
3709	LAKE JUNALUSKA	0.031	0.560	462.000	7.233	14.200	0.009
3710	LOOKOUT SHOALS	0.026	0.370	459.167	4.200	13.800	0.008
3711	MOUNTAIN ISLAND LAKE	0.018	0.270	462.000	5.580	12.800	0.005
3713	LAKE NORMAN	0.019	0.330	446.667	5.807	14.800	0.005
3715	RHODHISS LAKE	0.061	0.305	462.111	3.578	13.600	0.019
3716	SANTEELAH LAKE	0.011	0.160	366.400	5.360	14.800	0.006
3717	LAKE TILLERY	0.040	0.470	468.600	6.827	13.600	0.008
3718	WATERVILLE RESERVOIR	0.103	0.860	468.333	3.817	14.400	0.041
3719	LAKE WACCAMAW	0.018	0.120	455.667	3.583	9.800	0.004

APPENDIX B

CONVERSIONS FACTORS

CONVERSION FACTORS

Hectares x 2.471 = acres

Kilometers x 0.6214 = miles

Meters x 3.281 = feet

Cubic meters x 8.107×10^{-4} = acre/feet

Square kilometers x 0.3861 = square miles

Cubic meters/sec x 35.315 = cubic feet/sec

Centimeters x 0.3937 = inches

Kilograms x 2.205 = pounds

Kilograms/square kilometer x 5.711 = lbs/square mile

APPENDIX C

TRIBUTARY FLOW DATA

TRIBUTARY FLOW INFORMATION FOR NORTH CAROLINA

10/21/75

LAKE CODE 3702 BLEWETT FALLS RESERVOIR

TOTAL DRAINAGE AREA OF LAKE(SQ MI) 6870.00

TRIBUTARY	SUB-DRAINAGE AREA(SQ MI)	NORMALIZED FLOWS(CFS)												MEAN
		JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	
3702A1	6870.00	10140.0	12030.0	12510.0	10550.0	6598.0	5353.0	5341.0	5660.0	5938.0	5795.0	5268.0	7080.0	7662.9
3702A2	6331.00	9400.0	10930.0	11320.0	9600.0	6210.0	5040.0	5020.0	5400.0	5120.0	5040.0	4840.0	6280.0	6994.0
3702B1	15.60	19.00	28.00	27.00	18.00	8.00	5.50	6.50	6.00	5.80	5.00	6.50	11.00	12.10
3702C1	14.30	17.00	24.00	24.00	16.00	6.00	4.80	5.50	5.30	5.00	4.00	4.50	7.00	10.17
3702D1	10.10	14.00	20.00	19.00	12.00	5.50	4.00	4.50	4.30	4.00	3.50	6.00	9.00	8.75
3702E1	75.00	113.00	155.00	150.00	135.00	75.00	57.00	63.00	57.00	55.00	70.00	75.00	90.00	90.84
3702F1	352.00	450.00	650.00	650.00	610.00	240.00	200.00	190.00	145.00	135.00	250.00	230.00	320.00	337.11
3702G1	8.00	7.90	13.00	12.00	11.00	4.30	3.50	3.30	2.60	2.40	4.50	5.00	5.60	6.21
3702H1	30.30	55.00	125.00	95.00	88.00	22.00	16.00	15.00	10.00	9.00	23.00	16.00	33.00	41.68
3702Z1	33.70	60.00	81.00	78.00	62.00	30.00	25.00	29.00	26.00	25.00	27.00	28.00	38.00	42.18

SUMMARY

TOTAL DRAINAGE AREA OF LAKE =	6870.00	TOTAL FLOW IN =	90830.87
SUM OF SUB-DRAINAGE AREAS =	6869.99	TOTAL FLOW OUT =	92263.00

MEAN MONTHLY FLOWS AND DAILY FLOWS(CFS)

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
3702A1	3	73	17800.00	25	12500.00				
	4	73	23700.00	29	20400.00				
	5	73	10400.00						
	6	73	9860.00	3	10700.00				
	7	73	7320.00	8	1170.00				
	8	73	6960.00	9	9180.00				
	9	73	3630.00	9	1580.00				
	10	73	3990.00	14	602.00				
	11	73	4540.00	4	2890.00				
	12	73	8100.00	2	445.00				
	1	74	16500.00	6	12500.00	18	10100.00		
	2	74	17400.00	3	26100.00	14	13600.00		
3702A2	3	74	9400.00	3	11000.00				
	3	73	16300.00	25	10600.00				
	4	73	21500.00	29	18600.00				
	5	73	9770.00						
	6	73	9140.00	3	7310.00				
	7	73	6910.00	8	550.00				
	8	73	6700.00	5	8620.00				
	9	73	3520.00	9	150.00				
	10	73	3850.00	14	160.00				
	11	73	4300.00	4	1730.00				
	12	73	7600.00	2	280.00				
	1	74	15500.00	6	7620.00	18	8560.00		
	2	74	16000.00	3	26400.00	14	9320.00		
	3	74	8500.00	3	11700.00				

TRIBUTARY FLOW INFORMATION FOR NORTH CAROLINA

10/21/75

LAKE CODE 3702 BLEWETT FALLS RESERVOIR

MEAN MONTHLY FLOWS AND DAILY FLOWS(CFS)

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
3702B1	3	73	39.00	25	18.00				
	4	73	55.00	29	23.00				
	5	73	8.10						
	6	73	13.00	3	3.50				
	7	73	17.00	8	4.00				
	8	73	4.00	5	8.00				
	9	73	1.70	9	1.10				
	10	73	1.30	14	0.90				
	11	73	1.20	4	1.10				
	12	73	9.50	2	1.50				
	1	74	30.00	6	14.00	18	5.60		
	2	74	40.00	3	140.00	14	14.00		
3702C1	3	74	12.00	3	9.20				
	4	73	35.00	25	15.00				
	5	73	50.00	29	37.00				
	6	73	12.00						
	7	73	16.00	3	1.50				
	8	73	3.70	8	1.70				
	9	73	1.60	5	2.40				
	10	73	1.60	9	0.30				
	11	73	1.20	14	0.10				
	12	73	1.50	4	0.20				
	1	73	8.70	2	0.10				
	2	74	27.00	6	16.00	18	4.30		
3702D1	3	74	36.00	3	175.00	14	13.00		
	4	74	11.00	3	6.40				
	5	73	25.00	25	11.00				
	6	73	35.00	29	25.00				
	7	73	5.20						
	8	73	8.10	3	0.80				
	9	73	11.00	8	1.20				
	10	73	2.60	5	1.70				
	11	73	1.10	9	0.20				
	12	73	0.90	14	0.10				
	1	73	1.10	4	0.10				
	2	74	6.20	2	0.10				
3702E1	3	74	19.00	6	11.00	18	3.00		
	4	74	26.00	3	125.00	14	9.40		
	5	74	8.00	3	4.50				
	6	73	190.00	25	100.00				
	7	73	270.00	29	105.00				
	8	73	72.00						
	9	73	75.00	3	40.00				
	10	73	49.00	8	40.00				
	11	73	36.00	5	65.00				
	12	73	15.00	9	9.20				
	1	74	5.80	14	5.90				
	2	74	9.80	4	7.80				
	3	74	41.00	2	14.00				
	4	74	125.00	6	56.00	18	34.00		
	5	74	145.00	3	450.00	14	68.00		
	6	74	80.00	3	59.00				

TRIBUTARY FLOW INFORMATION FOR NORTH CAROLINA

10/21/75

LAKE CODE 3702 BLEWETT FALLS RESERVOIR

MEAN MONTHLY FLOWS AND DAILY FLOWS(CFS)

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
3702F1	3	73	900.00	25	470.00				
	4	73	1400.00	29	490.00				
	5	73	340.00						
	6	73	350.00	3	190.00				
	7	73	230.00	8	185.00				
	8	73	170.00	5	310.00				
	9	73	70.00	9	43.00				
	10	73	27.00	14	28.00				
	11	73	45.00	4	37.00				
	12	73	195.00	2	63.00				
	1	74	575.00	6	270.00	18	160.00		
	3	74	380.00	3	280.00				
3702G1	3	73	21.00	25	11.00				
	4	73	27.00	29	11.00				
	5	73	7.50						
	6	73	8.00	3	4.50				
	7	73	5.00	8	4.30				
	8	73	3.80	5	7.20				
	9	73	1.50	9	1.00				
	10	73	0.60	14	0.60				
	11	73	1.00	4	0.80				
	12	73	4.40	2	1.40				
	1	74	13.00	6	6.20	18	3.60		
	2	74	16.00	3	48.00	14	7.20		
3702H1	3	74	8.50	3	6.40				
	4	73	80.00	25	60.00				
	5	73	120.00	29	63.00				
	6	73	36.00						
	7	73	37.00	3	15.00				
	8	73	20.00	8	15.00				
	9	73	13.00	5	32.00				
	10	73	3.50	9	1.60				
	11	73	0.80	14	0.80				
	12	73	1.70	4	1.30				
	1	74	15.00	2	3.00				
	2	74	80.00	6	26.00	18	12.00		
3	74	105.00	3	180.00	14	33.00			
3702Z2	3	74	44.00	3	26.00				
	4	73	83.00						
	5	73	130.00						
	6	73	18.00						
	7	73	27.00						
	8	73	37.00						
	9	73	8.80						
	10	73	3.70						
	11	73	2.60						
	12	73	2.60						
	1	74	20.00						
	2	74	65.00						
3	74	85.00							
		27.00							

APPENDIX D

PHYSICAL and CHEMICAL DATA

STORET RETRIEVAL DATE 75/10/20

370201
35 04 30.0 079 53 46.0
BLEWETT FALLS LAKE
37153 NORTH CAROLINA

DATE FROM TO	TIME OF DAY	DEPTH FEET	WATER TEMP CENT	00010 DO	00300 TRANSP	00077 SECCHI	00094 FIELD MICROMHO	00400 PH	00410 TALK CACO3	00610 NH3-N TOTAL	00625 TOT KJEL	00630 NO2&NO3 N-TOTAL	00671 PHOS-DIS ORTHO MG/L P
				00010 DO	00300 TRANSP	00077 SECCHI	00094 FIELD MICROMHO	00400 PH	00410 TALK CACO3	00610 NH3-N TOTAL	00625 TOT KJEL	00630 NO2&NO3 N-TOTAL	00671 PHOS-DIS ORTHO MG/L P
73/03/23	09 45	0000	11.8		10	81	7.20	16	0.190	1.100	0.540	0.044	
	09 45	0004	11.8	10.2		83	7.10	18	0.150	0.400	0.530	0.032	
73/07/05	16 00	0000	25.7		10	80	6.90	17	0.120	0.600	0.450	0.012	
	16 00	0004	25.7	5.4		60	6.90	18	0.120	0.400	0.450	0.012	
73/09/19	16 05	0000	26.2	6.2	60L	62	7.10	24	0.060	0.200	0.280	0.015	

DATE FROM TO	TIME OF DAY	DEPTH FEET	00665 PHOS-TOT MG/L P	32217 CHLRPHYL A UG/L
			00665 PHOS-TOT MG/L P	32217 CHLRPHYL A UG/L
73/03/23	09 45	0000	0.103	2.5
	09 45	0004	0.087	
73/07/05	16 00	0000	0.080	2.2
	16 00	0004	0.045	
73/09/19	16 05	0000	0.020	2.1

L ACTUAL VALUE IS KNOWN TO BE
GREATER THAN VALUE GIVEN

STORET RETRIEVAL DATE 75/10/20

370202
35 00 17.0 079 53 37.0
BLEWETT FALLS LAKE
37153 NORTH CAROLINA

DATE FROM TO	TIME OF DAY	DEPTH FEET	WATER TEMP CENT	00310 DO MG/L	00300 TRANSP SECCHI	00077 INCHES	00094 CONDUTCTVY FIELD MICROMHO	00400 PH SU	00410 ALK CACO3 MG/L	00610 NH3-N TOTAL MG/L	00625 TOT KJEL N MG/L	00630 NO2&NO3 N-TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	11EPALES 3		2111202 0025 FEET DEPTH	
73/03/23	10 25	0000	12.2			10	80	7.20	16	0.180	0.800	0.500	0.041				
	10 25	0006	11.7	10.0			83	7.20	18	0.160	0.500	0.560	0.040				
	10 25	0013	11.5	10.0			90	7.20	18	0.160	0.600	0.570	0.042				
	10 25	0021	11.5	10.0			88	7.20	19	0.170	0.600	0.570	0.045				
73/07/05	15 15	0000	26.7			18	82	8.10	23	0.120	0.700	0.390	0.009				
	15 15	0006	26.2	4.9			90	7.00	25	0.130	0.300	0.640	0.028				
	15 15	0014	26.0	5.0			80	6.90	27	0.140	0.300	0.540	0.018				
73/09/14	16 20	0000	26.3			34	158	7.00	36	0.080	0.400	0.380	0.150				
	16 20	0005	26.3	6.0			140	6.80	36	0.060	0.400	0.330	0.102				
	16 20	0013	26.1	6.0			149	6.70	35	0.060	0.300	0.310	0.075				

DATE FROM TO	TIME OF DAY	DEPTH FEET	PHOS-TOT MG/L P	00665 CHLOROPHYL A UG/L	32217		
73/03/23	10 25	0000	0.094	2.3			
	10 25	0006	0.165				
	10 25	0013	0.134				
	10 25	0021	0.144				
73/07/05	15 15	0000	0.064	2.4			
	15 15	0006	0.092				
	15 15	0014	0.074				
73/09/14	16 20	0000	0.184	7.1			
	16 20	0005	0.129				
	16 20	0013	0.044				

STORED RETRIEVAL DATE 75/10/20

370203
34 59 07.0 079 52 21.0
BLEWETT FALLS LAKE
37007 NORTH CAROLINA

DATE FROM TO	TIME OF DAY	DEPTH FEET	00010 WATER TEMP CENT	00300 DO MG/L	00077 TRANSP SECCHI INCHES	00094 CONDUTCTVY FIELD MICROMHO	00400 PH SU	00410 ALK CACO3 MG/L	00610 NH3-N TOTAL MG/L	00625 TOT KJEL N MG/L	00630 NO2&NO3 N-TOTAL MG/L	11EPALES 3		2111202 0028 FEET DEPTH	
														00671 PHOS-DIS ORTHO MG/L P	
73/03/23	13 00	0000	12.5			10	80	7.40	18	0.170	0.700	0.530	0.038		
	13 00	0006	12.4	10.0			80	7.40	17	0.150	0.600	0.530	0.036		
	13 00	0012	12.2	10.0			80	7.30	17	0.150	0.700	0.530	0.038		
	13 00	0018	12.0	9.8			85	7.30	18	0.150	0.500	0.550	0.039		
	13 00	0024	12.0	9.8			90	7.30	19	0.160	0.600	0.560	0.040		
73/07/05	14 30	0000	28.3			18	80	8.90	24	0.140	0.900	0.290	0.008		
	14 30	0006	27.4	7.9			80	7.60	23	0.080	0.400	0.360	0.006		
	14 30	0015	25.9	4.3			75	6.90	23	0.130	0.400	0.530	0.015		
	14 30	0030	25.5	4.2			73	6.80	21	0.140	0.400	0.510	0.013		
73/09/19	16 40	0000	26.3			38	77	6.60	26	0.070	0.200	0.240	0.017		
	16 40	0005	26.2	6.2			85	6.60	26	0.040	0.200K	0.230	0.031		
	16 40	0015	26.1	6.2			89	6.50	26	0.060	0.200	0.240	0.031		
	16 40	0025	26.0	5.2			98	6.70	29	0.080	0.200	0.260	0.040		

DATE FROM TO	TIME OF DAY	DEPTH FEET	00665 PHOS-TOF MG/L P	32217 CHLOROPHYL A UG/L	
73/03/23	13 00	0000	0.095	2.6	
	13 00	0006	0.096		
	13 00	0012	0.100		
	13 00	0018	0.100		
	13 00	0024	0.102		
73/07/05	14 30	0000	0.089	5.9	
	14 30	0006	0.043		
	14 30	0015	0.048		
	14 30	0030	0.052		
73/09/19	16 40	0000	0.042	8.9	
	16 40	0005	0.040		
	16 40	0015	0.046		
	16 40	0025	0.070		

K VALUE KNOWN TO BE
LESS THAN INDICATED

APPENDIX E

TRIBUTARY and WASTEWATER TREATMENT PLANT DATA

STORED RETRIEVAL DATE 75/10/20

3702A1 LS3702A1
 34 57 00.0 079 52 30.0
 PEE DEE RIVER
 37167 7.5 ROCKINGHAM
 U/BLEWETT FALLS RESVR
 US HWY 74 HRDG 6.6 MI ESE OF LILESVILLE
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE	TIME	DEPTH	00630 NO2&NO3	00625 TOT KJEL	00610 NH3-N	00671 PHOS-DIS	00665 PHOS-TOT
FROM	OF		N-TOTAL	N	TOTAL	URTHO	
TO	DAY	FEET	MG/L	MG/L	MG/L	MG/L P	MG/L P
73/03/25	12	00	0.460	0.360	0.091	0.029	0.075
73/04/29	09	35	0.399	0.460	0.072	0.042	0.085
73/06/03	08	30	0.390	0.310	0.042	0.019	0.070
73/07/08	08	20	0.460	0.540	0.086	0.025	0.055
73/08/05	08	15	0.270	0.420	0.060	0.007	0.065
73/09/09	08	40	0.180	0.900	0.077	0.019	0.040
73/10/14	08	45	0.357	3.100	0.180	0.029	0.100
73/11/04	08	20	0.290	0.725	0.083	0.046	0.070
73/12/02	08	40	0.310	0.800	0.036	0.032	0.060
74/01/06	08	40	0.580	0.700	0.192	0.032	0.085
74/01/18	10	15	0.672	0.600	0.092	0.032	0.050
74/02/03	08	35	0.560	0.500	0.052	0.036	0.090
74/02/14	11	30	0.570	0.600	0.060	0.035	0.120
74/03/03	08	25	0.528	0.500	0.045	0.040	0.085

STORED RETRIEVAL DATE 75/10/20

3702A2 LS3702A2
 35 05 00.0 030 00 00.0
 PEE DEE RIVER
 37 7.5 MANGUM
 I/BLEWETT FALLS RESVR
 HWY 109 BRDG 2.75 MI SSW OFMANGUM
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 N026N03 H-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
73/03/25	11 25		0.510	0.480	0.063	0.062	0.123
73/04/29	09 15		0.420	0.520	0.096	0.033	0.070
73/06/03	08 55		0.530	0.490	0.061	0.058	0.105
73/07/08	09 20		0.610	0.350	0.044	0.054	0.090
73/08/05	08 30		0.340	0.630	0.040	0.115	0.150
73/09/09	09 45		0.315	0.480	0.070	0.019	0.035
73/10/14	08 50		0.300	2.100	0.074	0.174	0.220
73/11/04	08 45		0.300	0.650	0.126	0.054	0.060
73/12/02	08 30		0.440	0.900	0.104	0.368	0.390
74/01/06	08 40		0.770	0.600	0.108	0.056	0.115
74/01/18	09 30		0.770	0.400	0.120	0.116	0.180
74/02/03	08 45		0.630	0.900	0.140	0.056	0.210
74/02/14	10 20		0.720	0.400	0.045	0.065	0.150
74/03/03	08 35		0.640	0.400	0.030	0.105	0.155

STORED RETRIEVAL DATE 75/10/20

370281 LS370281
 35 04 30.0 079 59 30.0
 CEDAR CREEK
 37 7.5 MANGUM
 T/BLEWETT FALLS RESVR
 RD 1634 BRUG .9 MI E OF NC HWY 109
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 N02&N03 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
73/03/25	11	35	0.072	0.190	0.033	0.012	0.020
73/04/29	08	50	0.034	0.350	0.058	0.012	0.030
73/06/03	09	05	0.079	0.330	0.072	0.010	0.025
73/07/08	09	30	0.054	0.280	0.022	0.008	0.015
73/08/05	09	55	0.048	0.270	0.024	0.018	0.030
73/09/09	08	50	0.015	0.710	0.036	0.011	0.030
73/10/14	08	30	0.014	1.400	0.176	0.020	0.040
73/12/02	08	45	0.028	0.825	0.044	0.020	0.020
74/01/06	09	00	0.076	0.200	0.020	0.012	0.025
74/01/18	09	40	0.024	0.700	0.040	0.008	0.008
74/02/03	09	00	0.105	0.500	0.020	0.015	0.020
	09	25	0.088	0.500	0.028		
74/02/14	10	35	0.176	0.400	0.015	0.015	0.035

STORED RETRIEVAL DATE 76/10/20

3702C1 LS3702C1
 35 04 30.0 079 55 30.0
 SAVANNAH CHURCH
 37 7.5 MANGUM
 T/BLEWETT FALLS RESVR
 RD 1704 BRDG .25 MI S SAVANNAH CHURCH
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 N-26N03	00625 TOT KJEL	00610 NH3-N N	00671 PHOS-DIS ORTHO	00665 PHOS-TOT MG/L P
			MG/L	MG/L	MG/L	MG/L P	
73/03/25	11 50		0.072	0.290	0.046	0.009	0.020
73/04/29	04 40		0.034	0.400	0.054	0.012	0.030
73/06/03	08 55		0.110	0.320	0.054	0.009	0.025
73/07/08	09 45		0.058	0.210	0.038	0.008	0.020
73/08/05	08 45		0.048	0.275	0.026	0.013	0.025
73/09/04	08 45		0.015	0.640	0.052	0.014	0.050
73/10/14	08 35		0.010K	1.850	0.056	0.017	0.055
73/11/04	08 30		0.016	1.800	0.063	0.026	0.060
73/12/02	08 40		0.024	1.700	0.084	0.012	0.040
74/01/06	08 45		0.044	0.500	0.032	0.012	0.025
74/01/15	09 50		0.012	1.400	0.068	0.008	0.015
74/02/03	08 30		0.064	0.100	0.015	0.010	0.015
	09 15		0.040	0.700	0.032		
74/02/14	10 45		0.112	0.400	0.020	0.010	0.035

K VALUE KNOWN TO BE
 LESS THAN INDICATED

STORED RETRIEVAL DATE 75/10/20

370201 LS370201
34 59 30.0 079 55 30.0
SMITH CREEK
37 7.5 LILESVILLE
T/BLEWETT FALLS RESVR
RD 1744 BRDG 3.25 MI NE OF LILESVILLE
11EPALES 2111204
4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 N02&N03 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
73/03/25	12	10	0.058	0.190	0.019	0.012	0.020
73/04/29	08	20	0.075	0.280	0.054	0.013	0.025
73/06/03	08	35	0.082	0.380	0.034	0.006	0.025
73/07/08	10	00	0.044	0.230	0.029	0.008	0.020
73/08/05	08	20	0.010	0.350	0.019	0.011	0.040
73/09/09	08	30	0.018	0.330	0.058	0.013	0.030
73/10/14	08	20	0.040	0.450	0.074	0.013	0.020
73/11/04	08	15	0.038	0.850	0.052	0.014	0.020
73/12/02	08	50	0.044	0.900	0.068	0.005K	0.035
74/01/05	08	30	0.112	0.400	0.056	0.008	0.015
74/01/18	10	10	0.100	0.800	0.056	0.008	0.020
74/02/03	08	15	0.072	0.100K	0.010	0.010	0.015
	09	45	0.068	0.250	0.020	0.012	0.040
74/02/14	11	00	0.120	0.200	0.020	0.010	0.035

K VALUE KNOWN TO BE
LESS THAN INDICATED

STORED RETRIEVAL DATE 75/10/20

3702E1 LS3702E1
 35 03 30.0 079 52 30.0
 MOUNTAIN CREEK
 37 7.5 MANGUM
 T/BLEWETT FALLS RESVR
 RD 1148 BRDG 4.5 MI W OF LENZTON
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 N02&N03 MG/L	00625 TOT KJEL MG/L	00610 NH3-N MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
73/03/25	12 35		0.520	0.300	0.039	0.018	0.025
73/04/24	09 20		0.430	0.370	0.033	0.035	0.060
73/06/03	09 25		0.500	0.500	0.042	0.016	0.045
73/07/08	04 00		0.490	0.330	0.026	0.017	0.035
73/08/05	09 25		0.360	0.750	0.115	0.021	0.050
73/09/09	09 25		0.290	0.720	0.058	0.013	0.055
73/10/14	09 25		0.420	2.300	0.350	0.017	0.040
73/11/04	04 00		0.270	1.450	0.095	0.042	0.075
73/12/02	04 10		0.480	0.900	0.068	0.028	0.038
74/01/05	09 30		0.740	0.600	0.064	0.020	0.030
74/01/18	10 45		0.820	0.400	0.056	0.020	0.040
74/02/03	09 25		0.410	0.700	0.044	0.024	
74/02/14	10 33		0.704	0.300	0.020	0.015	0.040
74/03/03	04 00		0.700	0.200	0.040	0.010	0.020

STORET RETRIEVAL DATE 7/10/20

3702F1 LS3702F1
35 06 30.0 079 54 00.0
LITTLE RIVER
37 7.5 MANGUM
T/BLEWETT FALLS RESVR
RD 1148 HRDG 3.5 MI SW OF COVINGTON
11EPALES 2111204
4 0000 FEET DEPTH

DATE	TIME	DEPTH	00630 N02&N03	00625 TOT KJEL	00610 NH3-N	00671 PHOS-DIS	00665 PHOS-TOT
FROM	OF		N-TOTAL	N	TOTAL	ORTHO	
TO	DAY	FEET	MG/L	MG/L	MG/L	MG/L P	MG/L P
73/03/25	12	50	0.120	0.340	0.036	0.016	0.025
73/04/29	08	30	0.140	0.650	0.072	0.027	0.060
73/06/03	09	30	0.230	0.300	0.040	0.019	0.045
73/07/08	09	15	0.410	1.890	0.096	0.038	0.065
73/08/05	10	15	0.190	0.480	0.072	0.029	0.055
73/09/09	09	20	0.663	0.950	0.061	0.015	0.037
73/10/14	09	40	0.022	4.100	0.176	0.015	0.025
73/11/04	09	15	0.014	1.550	0.058	0.014	0.020
73/12/07	09	25	0.028	0.700	0.028	0.005K	0.015
74/01/06	09	50	0.264	0.700	0.068	0.028	0.070
74/01/18	11	00	0.250	0.400	0.044	0.032	0.055
74/02/03	09	35	0.120	0.700	0.036	0.024	
74/02/14	10	30	0.203	0.300	0.020	0.020	0.045
74/03/03	09	10	0.146	0.500	0.045	0.015	0.035

K VALUE KNOWN TO BE
LESS THAN INDICATED

STORED RETRIEVAL DATE: 7-16-2003

370261 LS370261
 35 06 30.0 080 00 00.0
 DRY CREEK
 37 7.5 MANGUM
 T/BLEWETT FALLS RESVR
 HWY 109 BRDG #4 S INTERSEC WITH RD 1145
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE	TIME	DEPTH	WQSN03	00630	00625	00610	00671	00665
FROM	OF		N-TOTAL	TOT KJEL	N	TOTAL	PHOS-DIS	PHOS-TOT
TO	DAY	FFET	MG/L	MG/L	MG/L	MG/L	MG/L P	MG/L P
73/03/25	11	06		0.143	0.400	0.028	0.013	0.027
73/04/29	09	06		0.150	0.730	0.115	0.016	0.040
73/06/03	04	15		0.231	0.580	0.056	0.022	0.070
73/07/08	04	35		0.350	0.580	0.048	0.029	0.065
73/08/05	08	05		0.210	0.460	0.029	0.024	0.045
73/09/04	09	10		0.613	0.905	0.033	0.015	0.050
73/10/14	04	10		0.013	3.150	0.094	0.012	0.065
73/11/04	04	05		0.010A	1.500	0.034	0.013	0.040
73/12/02	08	50		0.012	1.000	0.028	0.005K	0.040
74/01/05	08	50		0.520	0.400	0.040	0.012	0.030
74/01/12	04	26		0.276	0.300	0.034	0.008	0.025
74/02/03	04	16		0.040	0.400	0.052	0.032	0.085
74/02/14	10	10		0.365	0.300	0.015	0.010	0.032
74/03/03	08	10		0.180	0.300	0.030	0.010	0.020

K VALUE KNOWN TO BE
 LESS THAN INDICATED

STORTEL RETRIEVAL DATE: 7/21/2000

3702H1 LS3702H1
 34 58 30.0 079 51 30.0
 CARTLEDGE CREEK
 37 7.5 RUCKINGHAM
 THLEWETT FALLS RESVR
 RD 1141 HRDG 3.5 SSW OF HOLLY GROVE
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE	TIME	DEPTH	NH2SN03	TOT KJEL	NH3-N	00671	00665
FROM	HHR		N-TOTAL	N	TOTAL	PHOS-DIS	PHOS-TOT
TO	DAY	FILE	MG/L	MG/L	MG/L	MG/L P	MG/L P
73/03/20	12:20		0.147	0.200	0.016	0.014	0.015
73/04/24	04:30		0.100	0.330	0.024	0.007	0.020
73/05/03	04:00		0.092	0.280	0.032	0.006	0.020
73/05/08	04:45		0.100	0.710	0.032	0.011	0.020
73/05/05	08:45		0.056	0.560	0.061	0.011	0.025
73/05/09	04:30		0.028	0.870	0.048	0.011	0.025
73/10/14	04:05		0.011	1.250	0.085	0.009	0.015
73/11/04	08:45		0.010K	0.610	0.066	0.015	0.015
73/12/02	03:50		0.032	0.600	0.040	0.005K	0.015
74/01/05	04:10		0.130	1.300	0.072	0.005K	0.015
74/01/10	10:30		0.100	0.200	0.024	0.005K	0.015
74/02/03	04:00		0.084	0.610	0.032	0.008	
74/02/14	11:00		0.232	0.400	0.040	0.005	0.020
74/03/03	08:40		0.263	0.400	0.025	0.005K	0.005

K VALUE KNOWN TO BE
 LESS THAN INDICATED

STORET RETRIEVAL DATE 75/10/20

3702FA AS3702FA P001200*

35 22 00.0 079 47 30.0

BISCOE NC

37167 MONTGOMERY CO HW

T/BLEWETT FALLS LAKE

LITTLE RIVER

11EPALES 2141204

4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 N026N03	00625 TOT KJEL	00610 NH3-N	00671 PHOS-DIS	00665 PHOS-TOT	50051 FLOW RATE	50053 CONDUIT FLOW-MGD
			MG/L	MG/L	MG/L	MG/L P	MG/L P	INST MGD	MONTHLY
73/05/02	13 00	8.520	4.500	0.910	3.750	4.000	0.179	0.183	
73/06/01	13 30	0.020	3.100	0.120	3.050	3.500	0.125	0.136	
73/07/05	14 00	0.090	4.600	0.050	3.540	4.400	0.133	0.172	
73/08/01	15 00	0.230	4.200	0.230	4.420	9.800	0.121	0.119	
73/09/04	11 30	0.026	3.050	0.610	3.520		0.101	0.103	
73/10/04	11 00	0.180	4.700	0.066	7.100	7.700	0.118	0.113	
73/11/01	15 00	0.720	5.950	0.094	5.400	6.200	0.097	0.095	
73/12/10	08 00	2.600	6.100	0.052	7.700	8.700	0.122	0.099	
74/01/04	11 00	4.300	2.700	0.140	6.700	7.000	0.137	0.172	
74/02/04	09 00	1.400	4.100	0.062	4.700	5.200	0.214	0.204	
74/03/04	10 30	1.200	3.000	0.072	3.300	3.400	0.122	0.267	
74/04/02	13 00	0.520	4.000	0.110	3.000	3.700	0.124	0.144	
74/06/07	11 00	12.000	6.550	1.250	10.000	10.500	0.124	0.140	

STORED RETRIEVAL DATE 75/10/20

3702JA 1F3702JA P004000*
34 57 30.0 080 04 00.0
WADESBORO NC
37 ANSON CO HWY MAP
T/BLEWETT FALLS RES
BRUSHY FURK CREEK
11EPALES 2141204
4 0000 FEET DEPTH