U.S. ENVIRONMENTAL PROTECTION AGENCY NATIONAL EUTROPHICATION SURVEY

WORKING PAPER SERIES



REPORT
ON
LYNN RESERVOIR
BOX ELDER COUNTY
UTAH
EPA REGION VIII
WORKING PAPER No. 843

CORVALLIS ENVIRONMENTAL RESEARCH LABORATORY - CORVALLIS, OREGON and ENVIRONMENTAL MONITORING & SUPPORT LABORATORY - LAS VEGAS, NEVADA

REPORT
ON
LYNN RESERVOIR
BOX ELDER COUNTY
UTAH
EPA REGION VIII
WORKING PAPER No. 843

WITH THE COOPERATION OF THE

UTAH STATE DIVISION OF HEALTH

AND THE

UTAH NATIONAL GUARD

NOVEMBER, 1977

CONTENT

		Page
For	reward	ii
Lis	t of Utah Study Lakes and Reservoirs	iv
Lak	e and Drainage ARea Map	v
Sec	tions	
I.	Introduction	1
II.	Conclusions	1
III.	Reservoir and Drainage Basin Characteristics	2
IV.	Water Quality Summary	3
٧.	Appendices	7

FOREWORD

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nation-wide threat of accelerated eutrophication to freshwater lakes and reservoirs.

OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

- a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.
- b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.
- c. With such a transformation, an assessment of the potential for eutrophication control can be made.

LAKE ANALYSIS

In this report, the first stage of evaluation of lake and water-shed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning [§303(e)], water quality criteria/standards review [§303(c)], clean lakes [§314(a,b)], and water quality monitoring [§106 and §305(b)] activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

ACKNOWLEDGEMENT

The staff of the National Eutrophication Survey (Office of Research and Development, U.S. Environmental Protection Agency) expresses sincere appreciation to the Utah Department of Social Services and the Utah Department of Natural Resources for professional involvement, to the Utah National Guard for conducting the tributary sampling phase of the Survey, and to those Utah wastewater treatment plant operators who voluntarily provided effluent samples and flow data.

The staffs of the Bureau of Water Quality of the Division of Health and the Division of Wildlife Resources provided invaluable lake documentation and counsel during the Survey, reviewed the preliminary reports, and provided critiques most useful in the preparation of this Working Paper series.

Major General Maurice L. Watts, the Adjutant General of Utah, and Project Officer Lt. Colonel T. Ray Kingston, who directed the volunteer efforts of the Utah National Guardsmen, are also gratefully acknowledged for their assistance to the Survey.

NATIONAL EUTROPHICATION SURVEY

STUDY LAKES AND RESERVOIRS

STATE OF UTAH

NAME

Bear Deer Creek Echo

Fish

Flaming Gorge

Huntington Joes Valley Lower Bowns

Lynn

Minersville

Moon Nava.io Newcastle Otter Creek Panquich Pelican Pineview Piute Porcupine Powell.

Pruess Sevier Bridge Starvation Steinaker Tropic Utah

Willard Bay

COUNTY

Rich, UT; Bear Lake, ID

Wasatch Summit Sevier Daggett, UT; Sweetwater, WY


Emery Emery Garfield Box Elder Beaver Duchesne Kane

Iron Piute Garfield Uintah Weber Piute Cache

Garfield, Kane, San Juan, UT; Coconino, AZ

Millard Juab, Sanpete Duchesne Uintah Garfield Utah

Box Elder



LYNN RESERVOIR

STORET NO. 4905

I. INTRODUCTION

Lynn Reservoir was included in the National Eutrophication Survey as a water body of interest to the Utah Bureau of Environmental Health. Tributaries were not sampled, and this report primarily relates to the reservoir sampling data.

II. CONCLUSIONS

A. Trophic Condition:

Survey data indicate Lynn Reservoir is eutrophic. It ranked twenty-fourth in overall trophic quality when the 27 Utah lakes and reservoirs sampled in 1975 were compared using a combination of six parameters*. Twenty-four of the water bodies had less median total phosphorus, 25 had less median dissolved orthophosphorus, 20 had less median inorganic nitrogen, 24 had less mean chlorophyll <u>a</u>, and ll had greater mean Secchi disc transparency. Some depression of dissolved oxygen with depth occurred in August (4.6 mg/l at 5.5 meters).

Survey limnologists observed an algal bloom in the reservoir in September.

B. Rate-Limiting Nutrient:

The algal assay results indicate the reservoir was nitrogen limited in September.

The reservoir data indicate nitrogen limitation all three sampling times.

^{*} See Appendix A.

RESERVOIR AND DRAINAGE BASIN CHARACTERISTICS[†] III.

- A. Morphometry: Unknown.
- B. Precipitation*:
 - 1. Year of sampling: 49.4 centimeters.
 - 2. Mean annual: 44.7 centimeters.

[†] Table of metric equivalents--Appendix B. * See Working Paper No. 175, "... Survey Methods, 1973-1976".

IV. WATER QUALITY SUMMARY

Lynn Reservoir was sampled three times during the open-water season of 1975 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from two or more depths at one station on the reservoir (see map, page v). During each visit, a single depth-integrated (4.6 m or near bottom to surface) sample was collected for phytoplankton identification and enumeration, and a similar sample was taken for chlorophyll <u>a</u> analysis. During the last visit, a single 18.9-liter depth-integrated sample was collected for algal assays. The maximum depth sampled was 5.5 meters.

The sampling results are presented in full in Appendix C and are summarized in the following table.

A. SUMMARY OF PHYSICAL AND CHEMICAL CHARACTERISTICS FOR LYNN RESERVOIR STORET CODE 4905

	1ST SAMPLING (5/15/75) 2ND SAMPLING (8/ 5/75) 1 SITES 1 SITES			2ND SAMP	PLING (8/ 5/75)	3RD SAMPLING (9/17/75)			
				1 SITES					
RAMETER	RANGE	MEAN	MEDIAN	RANGE	MEAN MEDIAN	RANGE	MEAN MEDIAN		
MP (C)	7.5 - 11.5	10.1	11.3	17.2 - 20.9	19.4 20.1	12.7 - 12.8	12.7 12.7		
ISS OXY (MG/L)	8.8 - 8.8	8.5	8.4	4.6 - 7.0	6.2 7.0	7.0 - 8.2	7.6 7.6		
ADCTVY (MCROMO)	193 234.	220.	233.	316 361.	332. 320.	468 470.	469. 469.		
(STAND UNITS)	7.7 - 7.9	7.8	7.7	8.0 - 8.4	8.2 8.3	8.3 - 8.6	8.4 8.4		
T ALK (MG/L)	112 180.	135.	114.	143 181.	156. 144.	98 202.	150. 150.		
T P (MG/L)	0.107 - 0.140	0.127	0.135	0.075 - 0.158	0.107 0.088	0.094 - 0.234	0.164 0.164		
THO P (MG/L)	0.042 - 0.058	0.051	0.052	0.048 - 0.140	0.080 0.052	0.012 - 0.057	0.034 0.034		
)2+N03 (MG/L)	0.340 - 0.400	0.363	0.350	0.020 - 0.050	0.030 0.020	0.020 - 0.030	0.025 0.025		
MONIA (MG/L)	0.050 - 0.080	0.063	0.060	0.030 - 0.150	0.073 0.040	0.0 - 0.040	0.020 0.020		
JEL N (MG/L)	0.600 - 0.600	0.600	0.600	0.400 - 0.400	0.400 0.400	0.400 - 0.400	0.400 0.400		
1026 N (MG/L)	0.390 - 0.460	0.427	0.430	0.050 - 0.200	0.103 0.060	0.070 - 0.070	0.070 0.070		
TAL N (MG/L)	0.940 - 1.000	0.963	0.950	0.420 - 0.450	0.430 0.420	0.420 - 0.430	0.425 0.425		
HERPYL A (UGZL)	3.0 - 3.0	3.0	3.0	2.3 - 2.3	2.3 2.3	113.5 - 113.5	113.5 113.5		
CCHI (METERS)	0.4 - 0.4	0.4	0 • 4	3.4 - 3.4	3.4 3.4	2.5 - 2.5	2.5 2.5		

B. Biological characteristics:

1. Phytoplankton -

Sampling Date	Dom [*]	inant era	Algal Units per ml		
05/15/75	1. 2. 3. 4. 5.	Navicula sp. Phormidium sp. Pandorina sp. Cryptomonas sp. Trachelomonas sp. Other genera	239 159 106 80 27 132		
		Total	743		
08/05/75	1.	Melosira sp. Schroederia sp. Total	386 15 401		
09/17/75	1. 2. 3. 4. 5.	Aphanizomenon sp. Synedra sp. Diatoma sp. Anabaena sp. Asterionella sp. Other genera	7,130 4,045 548 548 480 2,058		
		Total	14,809		

2. Chlorophyll \underline{a} -

Sampling Date	Station <u>Number</u>	Chlorophyll <u>a</u> $(\mu g/1)$
05/15/75	1	3.0
08/05/75	1	2.3
09/17/75	1	113.5

C. Limiting Nutrient Study:

1. Autoclaved, filtered, and nutrient spiked -

Spike (mg/1)	Ortho P Conc. (mg/l)	Inorganic N Conc. (mg/l)	Maximum yield (mg/l-dry wt.)
Control	0.135	0.328	12.4
0.050 P	0.185	0.328	12.0
0.050 P + 1.0 N	0.185	1.328	37.0
1.0 N	0.135	1.328	29.4

2. Discussion -

The control yield of the assay alga, <u>Selenastrum capri-cornutum</u>, indicates that the potential primary productivity of Lynn Reservoir was very high at the time the sample was collected (09/17/76). Also, the significant increase in yield when only nitrogen was added, and the lack of yield response when phosphorus alone was added, indicate the reservoir was nitrogen limited at that time.

The reservoir data indicate nitrogen limitation all three sampling times; i.e., the mean inorganic nitrogen to orthophosphorus ratios were 8 to 1 in May, 1 to 1 in August, and 2 to 1 in September.

V. APPENDICES

APPENDIX A

LAKE RANKINGS

LAKE DATA TO BE USED IN RANKINGS

_							
CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- Mean sec	MEAN Chlora	15- MIN DO	MEDIAN DISS ORTHO P
0408	LAKE POWELL	0.010	0.410	339.830	3.081	13.800	0.007
4901	BEAR LAKE	0.011	0.040	253.167	0.945	9.200	0.003
4902	LOWER BOWN+S RESERVOIR	0.031	0.040	336.000	5.567	9.400	0.006
4903	DEER CREEK RESERVOIR	0.038	0.215	430.333	9.078	14.800	0.006
4904	ECHO RESERVOIR	0.947	0.170	450.333	6.967	14.000	0.012
4905	LYNN RESERVOIR	0.121	0.200	417.667	39.600	10.400	0.052
4906	FISH LAKE	0.023	0.040	152.000	12.483	10.400	0.004
4907	HUNTINGTON NORTH RESERVO	0.013	0.040	392.000	1.900	7.800	0.005
4908	JOE'S VALLEY RESERVOIR	0.012	0.045	400.000	2.483	11.200	0.003
4909	HINERSVILLE RESERVOIR	0.192	0.060	445.000	33.583	8.600	0.107
4910	MOON LAKE	0.008	0.040	381.000	2.700	9.600	0.002
4911	NAVAJO LAKE	0.016	0.040	368.000	2.000	6.000	0.003
4912	NEWCASTLE RESERVOIR	0.051	0.040	428.667	12.467	13.600	0.009
4913	OTTER CREEK RESERVOIR	0.067	0.040	453.667	11.767	10.600	0.033
4914	PANGUITCH LAKE	0.071	0.040	426.500	45.950	14.200	0.010
4915	PELICAN LAKE	0.044	0.050	438.500	6.350	8.400	0.004
4916	PINEVIEW RESERVOIR	0.028	0.300	435.083	5.692	14.600	0.006
4917	PIUTE RESERVOIR	0.047	0.150	482.625	25.329	11.600	0.007
4913	PORCUPINE RESERVOIR	0.025	0.110	440.000	7.860	12.400	0.011
4919	PRUESS RESERVOIR (GARRIS	û∙057	0.140	491.000	4.533	8.800	0.008
4920	SEVIER BRIDGE RESERVOIR	0.026	0.355	449.778	18.222	12.400	0.008
4921	STARVATION RESERVOIR	0.016	0.040	394.583	5.675	13.200	0.004
4922	STEINAKER RESERVOIR	0.011	0.040	316.750	1.844	12.600	0.005
4923	TROPIC RESERVOIR	0.021	0.050	425.000	7.200	8.400	0.006
4924	UTAH LAKE	C.132	0.320	490.583	72.012	11.400	0.012
4923	KILLARD BAY RESERVOIR	0.044	0.060	457.182	7.567	11.000	0.009
5605	FLAMING GORGE RESERVOIR	0.011	0.690	285.636	2.500	10.400	0.003

PERCENT OF LAKES WITH HIGHER VALUES (NUMBER OF LAKES WITH HIGHER VALUES)

CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- Mean Sec	MEAN CHLORA	15- Min do	MEDIAN DISS ORTHO P	INDEX NO
0408	LAKE POWELL	96 (25)	4 (1)	·81 (21)	73 (19)	15 (4)	42 (11)	311
4901	BEAR LAKE	90 (23)	87 (19)	96 (25)	100 (26)	77 (20)	90 (23)	540
4902	LOWER BOWN+S RESERVOIR	46 (12)	87 (19)	85 (22).	65 (17)	73 (19)	50 (13)	406
4903	DEER CREEK RESERVOIR	42 (411)	19 (5)	42 (11),	35 (' 9)	0 (0)	58 (14)	196
4904	ECHO RESERVOIR	31 (8)	27 (7)	19 (5),	50 (13)	12 (3)	13 (3)	152
4905	LYNN RESERVOIR	8 (2)	23 (6)	58 (15)	8 (2)	62 (15)	4 (1)	163
4906	FISH LAKE	62 (16)	65 (16)	100 (26)	23 (6)	62 (15)	79 (20)	391
4937	HUNTINGTON NORTH RESERVO	77 (20)	· 65 (16)	69 (18)	92 (24)	96 (25)	69 (18)	468
4908	JOE'S VALLEY RESERVOIR	81 (21)	58 (15)	62 (16)	85 (22)	46 (12)	96 (25)	428
4909	MINERSVILLE RESERVOIR	0 (0)	44 (11)	27 (7)	. 15 (3)	85 (22)	0 (0)	168
4910	HOON LAKE	100 (26)	87 (19)	73 (19)	77 (20)	69 (18)	100 (26)	506
4911	NAVAJO LAKE	69 (18)	87 (19)	77 (20)	88 (23)	100 (26)	85 (22)	506
4912	NEWCASTLE RESERVOIR	23 (6)	87 (19)	46 (12)	27 (7)	19 (5)	27 (7)	229
4913	OTTER CREEK RESERVOIR	15 (4)	87 (19)	15 (4)	31 (8)	54 (14)	8 (2)	210
4914	PANBUITCH LAKE	12 (3)	65 (16)	50 (13)	4 (1)	8 (2)	23 (6)	162
4915	PELICAN LAKE	37 (9)	54 (14)	35 (9)	54 (14)	90 (23)	73 (19)	343
4916	PINEVIEW RESERVOIR	50 (13)	15 (4)	38 (10)	58 (15)	-4 (1)	58 (14)	223
4917	PIUTE RESERVOIR	27 (7)	31 (8)	8 (2)	15 (4)	38 (10)	46 (12)	165
4918	PORCUPINE RESERVOIR	58 (15)	38 (10)	31 (8)	38 (10)	33 (8)	19 (5)	217
4919	PRUESS RESERVOIR (GARRIS	19 (5)	35 (9)	0 (0),	69 (18)	81 (21)	37 (9)	241
4920	SEVIER BRIDGE RESERVOIR	54 (14)	8 (2)	23 (6)	19 (5)	33 (8)	37 (9)	174
4921	STARVATION RESERVOIR	73 (19)	87 (19)	65 (17)	62 (16)	23 (6)	79 (20)	389
4922	STEINAKER RESERVOIR	85 (22)	87 (19)	88 (23)	96 (25)	27 (7)	65 (17)	448
4923	TROPIC RESERVOIR	65 (17)	50 (13)	54 (14)	46 (12)	90 (23)	58 (14)	363
4924	UTAH LAKE	4 (1)	12 (3)	4 (1)	0 (0)	42 (11)	13 (3)	75
4925	WILLARD BAY RESERVOIR	37 (9)	44 (11)	12 (3)	42 (11)	50 (13)	31 (8)	216
5605	FLAMING GORGE RESERVOIR	90 (23)	0 (0)	92 (24)	81 (21)	62 (15)	90 (23)	415

LAKES RANKED BY INDEX NOS.

	MANUES OF	INDEX NOOT	
RANK	LAKE CODE	LAKE NAME	INDEX NO
1	4901	BEAR LAKE	540
2	4911	NAVAJO LAKE	506
3	4910	MOON LAKE	506
4	4907	HUNTINGTON NORTH RESERVO	468
. 5	4922	STEINAKER RESERVOIR	448
6	4908	JOE'S VALLEY RESERVOIR	428
7	5605	FLAMING GORGE RESERVOIR	415
8	4902	LOWER BOWN'S RESERVOIR	406
9	4906	FISH LAKE	391
10	4921	STARVATION RESERVOIR	389
11	4923	TROPIC RESERVOIR	363
12	4915	PELICAN LAKE	343
13	0408	LAKE POWELL	311
14	4919	PRUESS RESERVOIR (GARRIS	241
15	4912	NEWCASTLE RESERVOIR	229
16	4916	PINEVIEW RESERVOIR	223
17	4918	PORCUPINE RESERVOIR	217
18	4925	WILLARD BAY RESERVOIR	216
19	4913	CTTER CREEK RESERVOIR	210
20	4903	DEER CREEK RESERVOIR	196
21	4920	SEVIER BRIDGE RESERVOIR	174
22	4909	MINERSVILLE RESERVOIR	168
23	4917	PIUTE RESERVOIR	165
24	4905	LYNN RESERVOIR	163
25	4914	PANQUITCH LAKE	162
26	4904	ECHO RESERVOIR	152
27	4924	UTAH LAKE	75

APPENDIX B

CONVERSION FACTORS

CONVERSION FACTORS

Hectares \times 2.471 = acres

Kilometers \times 0.6214 = miles

Meters x 3.281 = feet

Cubic meters $\times 8.107 \times 10^{-4} = acre/feet$

Square kilometers x 0.3861 = square miles

Cubic meters/sec x 35.315 = cubic feet/sec

Centimeters x 0.3937 = inches

Kilograms x 2.205 = pounds

Kilograms/square kilometer x 5.711 = 1bs/square mile

APPENDIX C

PHYSICAL and CHEMICAL DATA

490501 41 50 52.0 113 43 30.0 3 LYNN RESERVOIR 49003 UTAH

150791

00671

PHOS-DIS

MG/L P

0.042

0.052

0.058

0.048 0.052

0.140

0.012 0.057

ORTHO

0.020K

0.020K

2111202 11EPALES 0022 FEET DEPTH CLASS 00

DATE FROM TO	TIME DEPTH OF DAY FEET	00010 WATER TEMP CENT	00300 00 MG/L	00077 TRANSP SECCHI INCHES	00094 CNDUCTVY FIELD MICROMHO	00400 PH SU	00410 T ALK CACO3 MG/L	00610 NH3-N Total MG/L	00625 TOT KJEL N MG/L	00630 NO2&NO3 N-TOTAL MG/L
75/05/15	10 40 0000	11.3	8.2	15	233	7.70	114	0.080	0.600	0.350
	10 40 0005		8.8		234	7.90	112	0.050	0.600	0.340
	10 40 0018		8.4		193	7.70	180	0.060	0.600	0.400
75/08/05			7.0	132	316	8.40	143	0.030	0.400	0.020
	11 10 0005	20.1	7.0		320	8.30	144	0.040	0.400	0.020
	11 10 0018		4.6		361	8.00	181	0.150	0.400	0.050
75/09/17	10 45 0000	12.7	7.0	100	470	8.30	98	0.040	0.400	0.030
	10 45 .0005	12.8	8.2		468	8.60	202	0.000R	0.400	0.020
DATE FROM TO	TIME DEPTH OF DAY FEET	00665 PHOS-TOT	32217 CHLRPHYL A UG/L	00031 INCDT LT REMNING PERCENT						
75/05/15	10 40 0000 10 40 0005 10 40 0018	0.135	3.0							
75/08/05	11 10 0000 11 10 0005	0.075	2.3							

K VALUE KNOWN TO BE LESS THAN INDICATED

0.158

0.094

0.234

113.5

11 10 0018

10 45 0005

75/09/17 10 45 0000