


**U.S. ENVIRONMENTAL PROTECTION AGENCY  
NATIONAL EUTROPHICATION SURVEY  
WORKING PAPER SERIES**



REPORT  
ON  
NEWCASTLE RESERVOIR  
IRON COUNTY  
UTAH  
EPA REGION VIII  
WORKING PAPER No. 849

**CORVALLIS ENVIRONMENTAL RESEARCH LABORATORY - CORVALLIS, OREGON  
and  
ENVIRONMENTAL MONITORING & SUPPORT LABORATORY - LAS VEGAS, NEVADA**

REPORT  
ON  
NEWCASTLE RESERVOIR  
IRON COUNTY  
UTAH  
EPA REGION VIII  
WORKING PAPER No. 849

WITH THE COOPERATION OF THE  
UTAH STATE DIVISION OF HEALTH  
AND THE  
UTAH NATIONAL GUARD  
SEPTEMBER, 1977

## CONTENTS

	<u>Page</u>
Foreward	ii
List of Utah Study Lakes and Reservoirs	iv
Lake and Drainage Area Map	v
 <u>Sections</u>	
I. Introduction	1
II. Conclusions	1
III. Reservoir and Drainage Basin Characteristics	2
IV. Water Quality Summary	3
V. Literature Reviewed	7
VI. Appendices	8

## F O R E W O R D

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nationwide threat of accelerated eutrophication to freshwater lakes and reservoirs.

### OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

### ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

- a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.
- b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.
- c. With such a transformation, an assessment of the potential for eutrophication control can be made.

### LAKE ANALYSIS

In this report, the first stage of evaluation of lake and watershed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning [§303(e)], water quality criteria/standards review [§303(c)], clean lakes [§314(a,b)], and water quality monitoring [§106 and §305(b)] activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

#### ACKNOWLEDGEMENT

The staff of the National Eutrophication Survey (Office of Research and Development, U.S. Environmental Protection Agency) expresses sincere appreciation to the Utah Department of Social Services and the Utah Department of Natural Resources for professional involvement, to the Utah National Guard for conducting the tributary sampling phase of the Survey, and to those Utah wastewater treatment plant operators who voluntarily provided effluent samples and flow data.

The staffs of the Bureau of Water Quality of the Division of Health and the Division of Wildlife Resources provided invaluable lake documentation and counsel during the Survey, reviewed the preliminary reports, and provided critiques most useful in the preparation of this Working Paper series.

Major General Maurice L. Watts, the Adjutant General of Utah, and Project Officer Lt. Colonel T. Ray Kingston, who directed the volunteer efforts of the Utah National Guardsmen, are also gratefully acknowledged for their assistance to the Survey.

NATIONAL EUTROPHICATION SURVEY  
STUDY LAKES AND RESERVOIRS

STATE OF UTAH

NAME

COUNTY

Bear  
Deer Creek  
Echo  
Fish  
Flaming Gorge

Rich, UT; Bear Lake, ID  
Wasatch  
Summit  
Sevier  
Daggett, UT;  
Sweetwater, WY

Huntington  
Joes Valley  
Lower Bowns  
Lynn  
Minersville  
Moon  
Navajo  
Newcastle  
Otter Creek  
Panguich  
Pelican  
Pineview  
Piute  
Porcupine  
Powell


Emery  
Emery  
Garfield  
Box Elder  
Beaver  
Duchesne  
Kane  
Iron  
Piute  
Garfield  
Uintah  
Weber  
Piute  
Cache  
Garfield, Kane, San  
Juan, UT; Coconino, AZ  
Millard  
Juab, Sanpete  
Duchesne  
Uintah  
Garfield  
Utah  
Box Elder

Pruess  
Sevier Bridge  
Starvation  
Steinaker  
Tropic  
Utah  
Willard Bay

# NEWCASTLE RESERVOIR

X Lake Sampling Site

Land Subject to Inundation



Map Location

outlet  
dam

Little  
Pinto Creek

Pinto  
Creek

37°39'

37°38'

113°32'

113°31'

## NEWCASTLE RESERVOIR

STORET NO. 4912

### I. INTRODUCTION

Newcastle Reservoir was included in the National Eutrophication Survey as a water body of interest to the Utah Bureau of Environmental Health. Tributaries and nutrient sources were not sampled, and this report relates only to the lake sampling data.

### II. CONCLUSIONS

#### A. Trophic Condition:

Survey data indicate that Newcastle Reservoir is eutrophic. It ranked fifteenth in overall trophic quality when the 27 Utah lakes and reservoirs sampled in 1975 were compared using a combination of six parameters\*. Twenty of the water bodies had less median total phosphorus, 18 had less and one had the same median orthophosphorus, none had less and ten had the same median inorganic nitrogen, 19 had less mean chlorophyll a, and 14 had greater mean Secchi disc transparency. Significant depression of dissolved oxygen with depth occurred in August (1.4 mg/l at 9.4 meters).

Survey limnologists noted an algal bloom in progress in May.

#### B. Rate-Limiting Nutrient:

Because of nutrient changes in the samples, the algal assay results are not considered representative of conditions in the reservoir at the times the samples were taken. The reservoir data indicate nitrogen limitation in May and August and phosphorus limitation in September.

---

\* See Appendix A.



### III. RESERVOIR AND DRAINAGE BASIN CHARACTERISTICS<sup>†</sup>

#### A. Morphometry<sup>††</sup>:

1. Surface area: 0.66 kilometers<sup>2</sup>.
2. Mean depth: 7.2 meters.
3. Maximum depth: 23.5 meters.
4. Volume:  $4.736 \times 10^6$  m<sup>3</sup>.

#### B. Precipitation\*:

1. Year of sampling: 33.8 centimeters.
2. Mean annual: 26.2 centimeters.

---

<sup>†</sup> Table of metric conversions--Appendix B.

<sup>††</sup> Sudweeks, 1975; maximum depth from Ikner (1975).

\* See Working Paper No. 175, "...Survey Methods, 1973-1976".

#### IV. WATER QUALITY SUMMARY

Newcastle Reservoir was sampled three times during the open-water season of 1975 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from a number of depths at a single station on the reservoir (see map, page v). During each visit, a depth-integrated (4.6 m to surface) sample was collected for phytoplankton identification and enumeration; and a similar sample was taken for chlorophyll a analysis. During the first and last visits, an 18.9-liter depth-integrated sample was collected for algal assays. The maximum depth sampled was 9.4 meters.

The sampling results are presented in full in Appendix C and are summarized in the following table.

A. SUMMARY OF PHYSICAL AND CHEMICAL CHARACTERISTICS FOR NEWCASTLE RESERVOIR  
STORET CODE 4912

PARAMETER	1ST SAMPLING ( 5/ 8/75)				2ND SAMPLING ( 8/13/75)				3RD SAMPLING ( 9/26/75)			
	1 SITES				1 SITES				1 SITES			
	RANGE	MEAN	MEDIAN		RANGE	MEAN	MEDIAN		RANGE	MEAN	MEDIAN	
TEMP (C)	9.8 - 11.3	10.2	9.9		18.0 - 21.3	20.3	21.0		17.8 - 18.5	18.1	18.1	
DISS OXY (MG/L)	9.2 - 9.2	9.2	9.2		1.4 - 8.2	4.8	4.8		7.0 - 8.4	7.5	7.3	
CNDCTVY (MCROMO)	613. - 630.	621.	621.		495. - 596.	559.	573.		528. - 538.	534.	535.	
PH (STAND UNITS)	8.6 - 8.6	8.6	8.6		8.0 - 8.6	8.3	8.3		8.4 - 8.5	8.5	8.5	
TOT ALK (MG/L)	248. - 258.	255.	257.		180. - 186.	183.	182.		192. - 196.	194.	195.	
TOT P (MG/L)	0.071 - 0.079	0.074	0.072		0.034 - 0.064	0.048	0.046		0.033 - 0.045	0.040	0.040	
ORTHO P (MG/L)	0.041 - 0.050	0.044	0.042		0.007 - 0.040	0.016	0.009		0.002 - 0.005	0.003	0.003	
NO2+NO3 (MG/L)	0.020 - 0.020	0.020	0.020		0.020 - 0.020	0.020	0.020		0.020 - 0.020	0.020	0.020	
AMMONIA (MG/L)	0.020 - 0.030	0.022	0.020		0.020 - 0.040	0.030	0.030		0.020 - 0.040	0.025	0.020	
KJEL N (MG/L)	0.300 - 0.800	0.425	0.300		0.600 - 0.700	0.650	0.650		0.400 - 0.500	0.425	0.400	
INORG N (MG/L)	0.040 - 0.050	0.042	0.040		0.040 - 0.060	0.050	0.050		0.040 - 0.060	0.045	0.040	
TOTAL N (MG/L)	0.320 - 0.820	0.445	0.320		0.620 - 0.720	0.670	0.670		0.420 - 0.520	0.445	0.420	
CHLORPYL A (UG/L)	16.1 - 16.1	16.1	16.1		5.7 - 5.7	5.7	5.7		15.6 - 15.6	15.6	15.6	
SECCHI (METERS)	0.9 - 0.9	0.9	0.9		3.0 - 3.0	3.0	3.0		1.5 - 1.5	1.5	1.5	

## B. Biological Characteristics:

## 1. Phytoplankton -

<u>Sampling Date</u>	<u>Dominant Genera</u>	<u>Algal Units per ml</u>
05/08/75	1. <u>Stephanodiscus sp.</u>	6,117
	2. <u>Nitzschia sp.</u>	1,001
	3. <u>Chroomonas sp.</u>	278
	4. <u>Glenodinium sp.</u>	56
	5. <u>Schroederia sp.</u>	56
	Total	7,508
08/13/75	1. <u>Elakatothrix sp.</u>	1,771
	2. <u>Cyclotella sp.</u>	227
	3. <u>Oocystis sp.</u>	91
	4. <u>Endorina sp.</u>	45
	Total	2,134
09/26/75	1. <u>Asterionella sp.</u>	3,798
	2. <u>Oocystis sp.</u>	1,825
	3. <u>Fragilaria sp.</u>	831
	4. <u>Synedra sp.</u>	592
	5. <u>Aphanizomenon sp.</u>	543
	Other genera	1,043
	Total	8,632

2. Chlorophyll a -

<u>Sampling Date</u>	<u>Station Number</u>	<u>Chlorophyll <u>a</u> (<math>\mu</math>g/l)</u>
05/08/75	1	16.1
08/13/75	1	5.7
09/26/75	1	15.6

## C. Limiting Nutrient Study:

Significant nutrient changes occurred in the assay samples during shipment from the field to the laboratory, and the results

are not considered representative of conditions in the reservoir at the times the samples were taken (05/08/75 and 09/26/75).

The reservoir data indicate nitrogen limitation in May and August and phosphorus limitation in September (the mean inorganic nitrogen/dissolved orthophosphorus ratios were 1/1, 3/1, and 15/1, respectively).

## V. LITERATURE REVIEWED

Ikner, James, 1975. Personal communication (reservoir morphometry).  
U.S. Geol. Surv., Salt Lake City.

Sudweeks, Calvin K., 1975. Personal communication (reservoir morphometry). UT Bur. of Env. Health, Salt Lake City.

VI. APPENDICES

APPENDIX A

LAKE RANKINGS

## LAKE DATA TO BE USED IN RANKINGS

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLORA	15- MIN DO	MEDIAN DISS O2/HO P
0408	LAKE POWELL	0.010	0.410	339.830	3.081	13.800	0.007
4901	BEAR LAKE	0.011	0.040	253.167	0.945	9.200	0.003
4902	LOWER BOWN'S RESERVOIR	0.031	0.040	336.000	5.567	9.400	0.006
4903	DEER CREEK RESERVOIR	0.038	0.215	430.333	9.078	14.800	0.006
4904	ECHO RESERVOIR	0.047	0.170	450.333	6.967	14.000	0.012
4905	LYNN RESERVOIR	0.121	0.200	417.667	39.600	10.400	0.052
4906	FISH LAKE	0.023	0.040	152.000	12.483	10.400	0.004
4907	HUNTINGTON NORTH RESERVO	0.013	0.040	392.000	1.900	7.800	0.005
4908	JOE'S VALLEY RESERVOIR	0.012	0.045	400.000	2.483	11.200	0.003
4909	MINERSVILLE RESERVOIR	0.192	0.060	445.000	33.583	8.600	0.107
4910	MOON LAKE	0.008	0.040	381.000	2.700	9.600	0.002
4911	NAVAJO LAKE	0.016	0.040	368.000	2.000	6.000	0.003
4912	NEWCASTLE RESERVOIR	0.051	0.040	428.667	12.467	13.600	0.009
4913	OTTER CREEK RESERVOIR	0.067	0.040	453.667	11.767	10.600	0.033
4914	PANQUITCH LAKE	0.071	0.040	426.500	45.950	14.200	0.010
4915	PELICAN LAKE	0.044	0.050	438.500	6.350	8.400	0.004
4916	PINEVIEW RESERVOIR	0.028	0.300	435.083	5.692	14.600	0.006
4917	PIUTE RESERVOIR	0.047	0.150	482.625	25.329	11.600	0.007
4918	PORCUPINE RESERVOIR	0.025	0.110	440.000	7.860	12.400	0.011
4919	PRUESS RESERVOIR (GARRIS	0.057	0.140	491.000	4.533	8.800	0.008
4920	SEVIER BRIDGE RESERVOIR	0.026	0.355	449.778	18.222	12.400	0.008
4921	STARVATION RESERVOIR	0.016	0.040	394.583	5.675	13.200	0.004
4922	STEINAKER RESERVOIR	0.011	0.040	316.750	1.844	12.600	0.005
4923	TROPIC RESERVOIR	0.021	0.050	425.000	7.200	8.400	0.006
4924	UTAH LAKE	0.132	0.320	490.583	72.012	11.400	0.012
4925	WILLARD BAY RESERVOIR	0.044	0.060	457.182	7.567	11.000	0.009
5605	FLAMING GORGE RESERVOIR	0.011	0.690	285.636	2.500	10.400	0.003



PERCENT OF LAKES WITH HIGHER VALUES (NUMBER OF LAKES WITH HIGHER VALUES)

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLOR A	15- MIN DO	MEDIAN DISS ORT-O P	INDEX NO
0408	LAKE POWELL	96 ( 25)	4 ( 1)	81 ( 21)	73 ( 19)	15 ( 4)	42 ( 11)	311
4901	BEAR LAKE	90 ( 23)	87 ( 19)	96 ( 25)	100 ( 26)	77 ( 20)	90 ( 23)	540
4902	LOWER BOWN'S RESERVOIR	46 ( 12)	87 ( 19)	85 ( 22)	65 ( 17)	73 ( 19)	50 ( 13)	406
4903	DEER CREEK RESERVOIR	42 ( 11)	19 ( 5)	42 ( 11)	35 ( 9)	0 ( 0)	58 ( 14)	196
4904	ECHO RESERVOIR	31 ( 8)	27 ( 7)	19 ( 5)	50 ( 13)	12 ( 3)	13 ( 3)	152
4905	LYNN RESERVOIR	8 ( 2)	23 ( 6)	58 ( 15)	8 ( 2)	62 ( 15)	4 ( 1)	163
4906	FISH LAKE	62 ( 16)	65 ( 16)	100 ( 26)	23 ( 6)	62 ( 15)	79 ( 20)	391
4907	HUNTINGTON NORTH RESERVO	77 ( 20)	65 ( 16)	69 ( 18)	92 ( 24)	96 ( 25)	69 ( 18)	468
4908	JOE'S VALLEY RESERVOIR	81 ( 21)	58 ( 15)	62 ( 16)	85 ( 22)	46 ( 12)	96 ( 25)	428
4909	MINERSVILLE RESERVOIR	0 ( 0)	44 ( 11)	27 ( 7)	12 ( 3)	85 ( 22)	0 ( 0)	168
4910	MOON LAKE	100 ( 26)	87 ( 19)	73 ( 19)	77 ( 20)	69 ( 18)	100 ( 26)	506
4911	NAVAJO LAKE	69 ( 18)	87 ( 19)	77 ( 20)	88 ( 23)	100 ( 26)	85 ( 22)	506
4912	NEWCASTLE RESERVOIR	23 ( 6)	87 ( 19)	46 ( 12)	27 ( 7)	19 ( 5)	27 ( 7)	229
4913	OTTER CREEK RESERVOIR	15 ( 4)	87 ( 19)	15 ( 4)	31 ( 8)	54 ( 14)	8 ( 2)	210
4914	PANQUITCH LAKE	12 ( 3)	65 ( 16)	50 ( 13)	4 ( 1)	8 ( 2)	23 ( 6)	162
4915	PELICAN LAKE	37 ( 9)	54 ( 14)	35 ( 9)	54 ( 14)	90 ( 23)	73 ( 19)	343
4916	PINEVIEW RESERVOIR	50 ( 13)	15 ( 4)	38 ( 10)	58 ( 15)	4 ( 1)	58 ( 14)	223
4917	PIUTE RESERVOIR	27 ( 7)	31 ( 8)	8 ( 2)	15 ( 4)	38 ( 10)	46 ( 12)	165
4918	PORCUPINE RESERVOIR	58 ( 15)	38 ( 10)	31 ( 8)	38 ( 10)	33 ( 8)	19 ( 5)	217
4919	PRUESS RESERVOIR (GARRIS	19 ( 5)	35 ( 9)	0 ( 0)	69 ( 18)	81 ( 21)	37 ( 9)	241
4920	SEVIER BRIDGE RESERVOIR	54 ( 14)	8 ( 2)	23 ( 6)	19 ( 5)	33 ( 8)	37 ( 9)	174
4921	STARVATION RESERVOIR	73 ( 19)	87 ( 19)	65 ( 17)	62 ( 16)	23 ( 6)	79 ( 20)	389
4922	STEINAKER RESERVOIR	85 ( 22)	87 ( 19)	88 ( 23)	96 ( 25)	27 ( 7)	65 ( 17)	448
4923	TROPIC RESERVOIR	65 ( 17)	50 ( 13)	54 ( 14)	46 ( 12)	90 ( 23)	58 ( 14)	363
4924	UTAH LAKE	4 ( 1)	12 ( 3)	4 ( 1)	0 ( 0)	42 ( 11)	13 ( 3)	75
4925	WILLARD BAY RESERVOIR	37 ( 9)	44 ( 11)	12 ( 3)	42 ( 11)	50 ( 13)	31 ( 8)	216
5605	FLAMING GORGE RESERVOIR	90 ( 23)	0 ( 0)	92 ( 24)	81 ( 21)	62 ( 15)	90 ( 23)	415

## LAKES RANKED BY INDEX NOS.

RANK	LAKE CODE	LAKE NAME	INDEX NO
1	4901	BEAR LAKE	540
2	4911	NAVAJO LAKE	506
3	4910	MOON LAKE	506
4	4907	HUNTINGTON NORTH RESERVO	468
5	4922	STEINAKER RESERVOIR	448
6	4908	JOE'S VALLEY RESERVOIR	428
7	5605	FLAMING GORGE RESERVOIR	415
8	4902	LOWER BOWN'S RESERVOIR	406
9	4906	FISH LAKE	391
10	4921	STARVATION RESERVOIR	389
11	4923	TROPIC RESERVOIR	363
12	4915	PELICAN LAKE	343
13	0408	LAKE POWELL	311
14	4919	PRUESS RESERVOIR (GARRIS	241
15	4912	NEWCASTLE RESERVOIR	229
16	4916	PINEVIEW RESERVOIR	223
17	4918	PORCUPINE RESERVOIR	217
18	4925	WILLARD BAY RESERVOIR	216
19	4913	OTTER CREEK RESERVOIR	210
20	4903	DEER CREEK RESERVOIR	196
21	4920	SEVIER BRIDGE RESERVOIR	174
22	4909	MINERSVILLE RESERVOIR	168
23	4917	PIUTE RESERVOIR	165
24	4905	LYNN RESERVOIR	163
25	4914	PANQUITCH LAKE	162
26	4904	ECHO RESERVOIR	152
27	4924	UTAH LAKE	75

## APPENDIX B

### CONVERSION FACTORS

## CONVERSION FACTORS

Hectares x 2.471 = acres

Kilometers x 0.6214 = miles

Meters x 3.281 = feet

Cubic meters x  $8.107 \times 10^{-4}$  = acre/feet

Square kilometers x 0.3861 = square miles

Cubic meters/sec x 35.315 = cubic feet/sec

Centimeters x 0.3937 = inches

Kilograms x 2.205 = pounds

Kilograms/square kilometer x 5.711 = lbs/square mile

## APPENDIX C

### PHYSICAL and CHEMICAL DATA

491201  
37 39 00.0 113 31 00.0 3  
NEWCASTLE RESERVOIR  
49021 UTAH

11EPALES 760114 2111202  
0034 FEET DEPTH CLASS 00

DATE FROM TO	TIME OF DAY	DEPTH FEET	00010 WATER TEMP CENT	00300 DO MG/L	00077 TRANSP SECCHI INCHES	00094 CONDUCTVY FIELD MICROMHO	00400 PH SU	00410 T ALK CAC03 MG/L	00610 NH3-N TOTAL MG/L	00625 TOT KJEL N MG/L	00630 NO2&N03 N-TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P
75/05/08	14 35	0000	11.3	9.2	34	624	8.60	258	0.030	0.800	0.020K	0.050
	14 35	0005	9.9	9.2		630	8.60	258	0.020	0.300	0.020K	0.043
	14 35	0015	9.9	9.2		618	8.60	248	0.020K	0.300	0.020K	0.042
	14 35	0030	9.8	9.2		613	8.60	256	0.020K	0.300	0.020K	0.041
75/08/13	08 45	0000	21.3	7.8	120	574	8.60	182	0.020K	0.700	0.020K	0.007
	08 45	0005	21.3	8.2		572	8.10	186	0.040	0.700	0.020K	0.012
	08 45	0020	20.7	1.8		596	8.60	180	0.020K	0.600	0.020K	0.007
	08 45	0031	18.0	1.4		495	8.00	182	0.040	0.600	0.020K	0.040
75/09/26	13 10	0000	18.5	8.4	60	536	8.50	192	0.020K	0.400	0.020K	0.004
	13 10	0005	18.2	7.6		528	8.50	194	0.020K	0.400	0.020K	0.002
	13 10	0015	18.1	7.0		534	8.45	195	0.020K	0.400	0.020K	0.002
	13 10	0021	17.8	7.0		538	8.50	196	0.040	0.500	0.020K	0.005

DATE FROM TO	TIME OF DAY	DEPTH FEET	00665 PHOS-TOT MG/L P	32217 CHLRPHYL A UG/L	00031 INCDT LT REMNING PERCENT
75/05/08	14 35	0000	0.079	16.1	
	14 35	0005	0.071		
	14 35	0015	0.072		
	14 35	0030	0.073		
75/08/13	08 45	0000	0.035	5.7	
	08 45	0005	0.058		
	08 45	0020	0.034		
	08 45	0031	0.064		
75/09/26	13 10	0000	0.039	15.6	
	13 10	0005	0.042		
	13 10	0015	0.033		
	13 10	0021	0.045		

K VALUE KNOWN TO BE  
LESS THAN INDICATED