

Draft
3/14/86

AMBIENT AQUATIC LIFE WATER QUALITY CRITERIA FOR
PARATHION

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF RESEARCH AND DEVELOPMENT
ENVIRONMENTAL RESEARCH LABORATORIES
DULUTH, MINNESOTA
NARRAGANSETT, RHODE ISLAND

NOTICES

This document has been reviewed by the Criteria and Standards Division, Office of Water Regulations and Standards, U.S. Environmental Protection Agency, and approved for publication.

Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

This document is available to the public through the National Technical Information Service (NTIS), 5285 Port Royal Road, Springfield, VA 22161.

FOREWORD

Section 304(a)(1) of the Clean Water Act of 1977 (P.L. 95-217) requires the Administrator of the Environmental Protection Agency to publish water quality criteria that accurately reflect the latest scientific knowledge on the kind and extent of all identifiable effects on health and welfare that might be expected from the presence of pollutants in any body of water, including ground water. This document is a revision of proposed criteria based upon a consideration of comments received from other Federal agencies, State agencies, special interest groups, and individual scientists. Criteria contained in this document replace any previously published EPA aquatic life criteria for the same pollutant(s).

The term "water quality criteria" is used in two sections of the Clean Water Act, section 304(a)(1) and section 303(c)(2). The term has a different program impact in each section. In section 304, the term represents a non-regulatory, scientific assessment of ecological effects. Criteria presented in this document are such scientific assessments. If water quality criteria associated with specific stream uses are adopted by a State as water quality standards under section 303, they become enforceable maximum acceptable pollutant concentrations in ambient waters within that State. Water quality criteria adopted in State water quality standards could have the same numerical values as criteria developed under section 304. However, in many situations States might want to adjust water quality criteria developed under section 304 to reflect local environmental conditions and human exposure patterns before incorporation into water quality standards. It is not until their adoption as part of State water quality standards that criteria become regulatory.

Guidelines to assist States in the modification of criteria presented in this document, in the development of water quality standards, and in other water-related programs of this Agency, have been developed by EPA.

James M. Conlon
Acting Director
Office of Water Regulations and Standards

ACKNOWLEDGMENTS

Loren J. Larson
(freshwater author)
University of Wisconsin-Superior
Superior, Wisconsin

Jeffrey L. Hyland
Sam R. Petrocelli
(saltwater authors)
Battelle New England Laboratory
Duxbury, Massachusetts

Charles E. Stephan
(document coordinator)
Environmental Research Laboratory
Duluth, Minnesota

David J. Hansen
(saltwater coordinator)
Environmental Research Laboratory
Narragansett, Rhode Island

Clerical Support: Shelley A. Heintz
Terry L. Highland
Diane L. Spehar
Nancy J. Jordan

CONTENTS

	<u>Page</u>
Foreword	iii
Acknowledgments	iv
Tables	vi
Introduction	1
Acute Toxicity to Aquatic Animals	3
Chronic Toxicity to Aquatic Animals	4
Toxicity to Aquatic Plants	5
Bioaccumulation	6
Other Data	6
Unused Data	9
Summary	12
National Criteria	13
References	37

TABLES

	<u>Page</u>
1. Acute Toxicity of Parathion to Aquatic Animals	15
2. Chronic Toxicity of Parathion To Aquatic Animals	22
3. Ranked Genus Mean Acute Values with Species Mean Acute-Chronic Ratios	23
4. Toxicity of Parathion to Aquatic Plants	27
5. Bioaccumulation of Parathion by Aquatic Organisms	28
6. Other Data on Effects of Parathion on Aquatic Organisms	29

Introduction*

Parathion (O,O-diethyl O-4-nitrophenyl phosphorothioate, sometimes called ethyl parathion or parathion-ethyl) is one of several organophosphorus pesticides developed to replace the more persistent organochlorine pesticides. It is now a restricted-use pesticide that is effective against a wide-range of insect pests on many fruit, nut, vegetable, and field crops. It is usually formulated as an emulsifiable concentrate, but is also available in granules, dusts, aerosols, oil sprays, and wettable powders. These formulations often contain large percentages of unspecified ingredients, which are often considered inert. Although no studies have compared the relative toxicities of technical-grade parathion and its various formulations, other organophosphorus insecticides (e.g., chlorpyrifos) have been shown to differ substantially in this regard. Although some data obtained from studies on formulations are discussed, data from such studies are not used in deriving criteria.

The toxicity of parathion is the result of metabolic conversion to its oxygen analogue, parathion-oxon (paraoxon) and its subsequent binding to and inhibition of various enzyme systems (e.g., cholinesterases, carboxylases, acetylcholinesterases and mitochondrial oxidative phosphorylases). Its inhibition of acetylcholinesterase (AChE) is generally accepted to be

* An understanding of the "Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Aquatic Organisms and Their Uses" (Stephan et al. 1985), hereafter referred to as the Guidelines, is necessary in order to understand the following text, tables, and calculations.

its most critical toxic effect. Inhibition of AChE results in accumulation of the neurotransmitter acetylcholine in synapses, disrupting normal neural transmission. Although even substantial reductions in brain AChE activity in fish have not always been fatal, the effect of this condition on normal activities (e.g., feeding, reproduction, predator-prey relationships, etc.) in nature is not known. Parathion has also been demonstrated to produce teratogenic effects in fish embryos (Solomon 1977; Solomon and Weis 1979; Tomita and Matsuda 1961).

Parathion is less persistent than organochlorine pesticides and has such a great affinity for organic material that it is quickly sorbed to sediments and suspended particulate matter. Miller et al. (1967) observed a rapid decrease in concentration after application of parathion to irrigation water and attributed it to degradation. It is more likely that sorption contributed greatly to this decrease. The persistence of parathion in water is dependent on chemical hydrolysis and biodegradation (Ahmed and Casida 1958; Faust 1975; Faust and Gomaa 1972; Gomaa and Faust 1977; Ludemann and Herzel 1973; Mackiewicz et al. 1969; Mulla 1963; Van Middlem 1966; Zuckerman et al. 1970). Graetz et al. (1970) reported that the portion of parathion degradation attributable to abiological in natural lake sediments means was negligible. Movement and persistence of parathion has been described in a natural pond (Mulla et al. 1966; Nicholson et al. 1962), a model stream (Laplanche et al. 1981), and a model ecosystem (Dortland 1980). Several studies report concentrations of parathion in water (Braun and Frank 1980; Dick 1982; Harris and Miles 1975; Greve et al. 1972; Kannan and Job 1979; Sethunathan et al. 1977) and in biota (Chovelon et al. 1984; Haddadin and Alawi 1974; Hesselberg and Johnson

1972; Perry et al. 1983). U.S. EPA (1975) and vom Runkler et al. (1974) reviewed the use, distribution, fate, and effects of parathion.

Unless otherwise noted, all concentrations reported herein are expressed as parathion, not as the material tested. The criteria presented herein supersede previous aquatic life water quality criteria for parathion (U.S. EPA 1976) because these new criteria were derived using improved procedures and additional information. Whenever adequately justified, a national criterion may be replaced by a site-specific criterion (U.S. EPA 1983a), which may include not only site-specific criterion concentrations (U.S. EPA 1983b), but also site-specific durations of averaging periods and site-specific frequencies of allowed excursions (U.S. EPA 1985). The latest literature search for information for this document was conducted in February, 1985; some newer information was also used.

Acute Toxicity to Aquatic Animals

The results of acute tests that were considered useful for deriving water quality criteria for parathion are listed in Table 1. The most striking disparity of values within a species in Table 1 is for the crayfish, Orconectes nais. An early instar was 375 times more sensitive to parathion than adults. The LC50 of 0.04 µg/L for this early instar of Orconectes nais is the lowest available acute value.

Freshwater Species Mean Acute Values (Table 1) were calculated as geometric means of the available acute values, and then Genus Mean Acute Values (Table 3) were calculated as geometric means of the available freshwater Species Mean Acute Values. Of the 31 genera for which acute values are available, the most sensitive genus, Orconectes, is over 130,000 times more sensitive than the most resistant, Tubifex and

Limnodrilus. Although nine of the 31 freshwater genera are fishes, the fifteen most sensitive genera are all invertebrates. However, the two most resistant genera are also invertebrates. Acute values are available for more than one species in each of five genera, and the range of Species Mean Acute Values within four of the genera is less than a factor of 1.9. In the fifth genus, the Gammarus, the acute values for mature individuals of the two species are similar, but acute values are available for younger individuals, which are apparently more sensitive, for only one of the species. The freshwater Final Acute Value for parathion was calculated to be 0.1298 µg/L using the procedure described in the Guidelines and the Genus Mean Acute Values in Table 3. The acute value for the crayfish, Orconectes nais, is about one-third the Final Acute Value.

Data on the acute toxicity of parathion are only available for two saltwater species (Table 1). The 96-hr LC50 for the Korean shrimp, Palaemon macrodactylus, was 11.5 µg/L in a static test and 17.8 µg/L in a flow-through test (Earnest 1970). Korn and Earnest (1974) reported that the 96-hr LC50 for the striped bass, Morone saxatilis, was 17.8 µg/L. Acute values are not available for enough species to allow calculation of a saltwater Final Acute Value.

Chronic Toxicity to Aquatic Animals

Chronic tests on parathion have been conducted with three freshwater invertebrates and three freshwater fishes (Spacie 1976; Spacie et al. 1981). Because of experimental problems, especially high control mortality, only data for Daphnia magna, the fathead minnow, and the bluegill are considered acceptable for use in deriving criteria. Some data are available from these studies on long-term effects on survival (Table 5), but it is not known whether other effects were more or less sensitive. For example, in

an exposure that began with brook trout embryos, an LC50 of 75 µg/L was obtained, but the duration was not given. In addition, 32 µg/L caused developmental abnormalities, whereas 10 µg/L reduced percent hatch, but did not cause abnormalities.

In the life-cycle test with D. magna, the 21-day LC50 was 0.14 µg/L, and the number of young produced was reduced by 0.12 µg/L, but not by 0.0817 µg/L. Chronic exposure of bluegill larvae produced no statistically significant effect on length at 30, 60, and 90 days. There was also no statistically significant effect on number of eggs spawned, percent hatch, or survival of larvae at 7, 14, 21, and 30 days.

Fathead minnows were reported to be significantly affected by chronic exposure to parathion at 9.0 µg/L, but not at 4.4 µg/L. The Acute-Chronic Ratio for fathead minnows is 79.45, whereas that for bluegills is 2,121 (Table 2).

The three Acute-Chronic Ratios available for parathion are 10.10, 79.45, and 2,121. The two highest ratios were obtained with two fish species that are acutely quite resistant to parathion. The ratio of 10.10 was obtained with an invertebrate that is acutely fairly sensitive to parathion. Thus it seems reasonable to use 10.10 as the Final Acute-Chronic Ratio. Division of the freshwater Final Acute Value of 0.1298 µg/L by the Final Acute-Chronic Ratio of 10.10 results in a Final Chronic Value of 0.01285 µg/L.

No data are available on the chronic toxicity of parathion to saltwater animals.

Toxicity to Aquatic Plants

Data are available on the toxicity of parathion to two freshwater algae (Table 4). The blue-green alga, Microcystis aeruginosa, was affected

at 30 µg/L, whereas the green alga, Scenedesmus quadricauda was not affected by concentrations below 390 µg/L.

No data are available concerning the toxicity of parathion to saltwater plants.

Bioaccumulation

Spacie (1976) and Spacie et al. (1981) reported long-term bioconcentration factors (BCFs) for the brook trout, fathead minnow, and bluegill (Table 5), and short-term BCFs are available for the brown trout, brook trout, and bluegill (Table 6). The BCFs determined with brook trout did not show a consistent relationship with either concentration in water or duration of exposure (Tables 5 and 6); the 260-day BCFs ranged from 31 to 232 for muscle tissue. The 260-day BCFs based on whole-body measurements with fathead minnows ranged from 32.9 to 201.4. The short-term BCFs measured with bluegills increased steadily with duration of exposure from 80.5 at 12 hr to 462 at 72 hr, but the BCF at 540 days was only 27.

No data are available on the bioaccumulation of parathion by saltwater species that can be used in the derivation of water quality criteria.

No U.S. FDA action level or other maximum acceptable concentration in tissue is available for parathion, and, therefore, no Final Residue Value can be calculated.

Other Data

Additional data on the effects of parathion on aquatic organisms are given in Table 6. The majority of the data are LC50s for durations other than 96 hours. Ahmed (1977) observed a range in 24-hr LC50s from 1.8 µg/L to 40 µg/L with six freshwater coleopteran species. Because of its wide use as a mosquito larvicide, many data are available on acute toxicity

to mosquito larva. However, standard methods for testing effectiveness of larvicides prescribe a 24-hr test duration. The 24-hr LC50s for seven species of mosquitos in three genera range from 0.47 to 68 µg/L. Gutierrez et al. (1977) reported LC50s from 1.8 to 70 µg/L for larvae of resistant populations of Culex pipiens.

Kynard (1974) observed avoidance of parathion by mosquitofish, and Weiss (1961) found inhibition of AChE in brains of several freshwater fishes. Effects on locomotor behavior of goldfish, bluegills, and largemouth bass were reported by Rand (1977a,b) and Rand et al. (1975). Sun and Taylor (1983) studied effects of parathion on acquisition and retention of a conditioned response in goldfish.

Various studies have examined the effect of a detergent (Solon and Nair 1970; Solon et al. 1969), herbicides (Lichtenstein et al. 1975), and an N-alkyl compound, SKF-525A (Gibson and Ludke 1973) on the toxicity of parathion. Banas and Sprague (1981) reported that prior exposure of rainbow trout did not affect the LC50.

Several studies evaluated the effectiveness of using trout for detecting pollutants including parathion (Jung 1973; Morgan 1975,1976; Van Hoof 1980). Morgan (1977) reported that fishes were able to detect parathion at 15% 48 hr-LC50. Mount and Boyle (1969) examined the use of the concentration of parathion in fish blood to diagnose causes of fish kills.

Ghetti and Gorbi (1985) studied the effects of a simulated parathion spill on a stream. Albright et al. (1983), Gasith and Perry (1980,1983,1985), Gasith et al. (1983a,b), and Grzenda et al. (1962) reported community effects of parathion on a pond. Warnick et al. (1966) found that increases in the concentrations of organochlorine compounds in water correlated with application of parathion to a pond. They postulated that these compounds were released from decomposing tissues of intoxicated organisms.

At a concentration of 1,000 $\mu\text{g/L}$, parathion reduced the rate of growth of natural saltwater plankton communities by 9.9% in 4 hr (Butler 1964). Juvenile pink shrimp, Penaeus duorum, had a 48-hr EC50 of 0.24 $\mu\text{g/L}$, whereas the EC50s for other penaeid and palaemonid shrimp ranged from 1.0 to 5.5 $\mu\text{g/L}$ (Butler 1964; Lowe et al. 1970; U.S. Bureau of Commercial Fisheries 1966,1967). Grass shrimp, Palaemonetes pugio, exposed to 0.1 or 0.5 $\mu\text{g/L}$ were more susceptible to predation by gulf killifish, Fundulus grandis (Farr 1977). Limb regeneration and time to molting of the fiddler crab, Uca pugilator, were apparently unaffected by exposure to parathion for 2 to 3 weeks, but all crabs exposed to 100 $\mu\text{g/L}$ died (Weis and Mantel 1976). The 96-hr EC50 based on shell deposition was 850 $\mu\text{g/L}$ or higher for the eastern oyster, Crassostrea virginica (Butler 1963,1964; Lowe et al. 1970; U.S. Bureau of Commercial Fisheries 1966). Lowe et al. (1971) found that growth of juvenile oysters was not reduced by exposure to 0.8 $\mu\text{g/L}$ for 252 days. Davis and Hidu (1969) reported a 78% reduction in length of oyster larvae after a 12-day exposure to 1,000 $\mu\text{g/L}$.

The sensitivity of saltwater fishes to parathion did not differ greatly. The 48-hr LC50s were 15 $\mu\text{g/L}$ for longnose killifish, Fundulus similis; 18 $\mu\text{g/L}$ for spot, Leiostomus xanthurus; 36 $\mu\text{g/L}$ for sheepshead minnows, Cyprinodon variegatus; and 100 $\mu\text{g/L}$ for striped mullet, Muzil cephalus (Butler 1964; Lowe 1979; and U.S. Bureau of Commercial Fisheries 1966,1967). Fin regenerative ability was reduced in adult mummichogs, Fundulus heteroclitus, exposed to 10 $\mu\text{g/L}$ for 10 weeks (Weis and Weis 1975) and this fish had a 50% incidence of circulatory failure when exposed to 10,000 $\mu\text{g/L}$ for three days (Weis and Weis 1974).

Inhibition of acetylcholinesterase (AChE) in saltwater fishes is a function of degree and duration of acute exposure and appears associated with death. Regardless of concentration and duration of exposure, when 40 to 60% of the sheepshead minnows, pinfish, and spot died, survivors had AChE reductions of \geq 82.3% (Coppage 1972, Coppage and Mathews 1974). Reductions of AChE of up to 82% were observed without death in a 120-hr exposure of sheepshead minnows to 5 μ g/L parathion (Coppage 1972).

White et al. (1979) reported 57 to 90% inhibition of brain cholinesterase activity in dead laughing gulls, Larus atricilla, contaminated with parathion applied to crops. Death of chicks was suspected to be a result of parathion in their food.

Unused Data

Some data on the effects of parathion on aquatic organisms were not used because the studies were conducted with species that are not resident in North America (e.g., Basak and Konar 1976; Bellavere and Gorbi 1984; Bowman et al. 1981; Butler 1964; Dortland 1980; Fleming 1981; Gregory et al. 1969; Gupta et al. 1979; Hashimoto and Nishiuchi 1981; Hudson et al. 1979; Iuhnke and Ludemann (1978); Nishiuchi and Hashimoto 1967; Nishiuchi and Yoshida 1972; Palawski et al. 1983; Panwar et al. 1976; Price 1976, 1978; Rattner 1982; Shah et al. 1983; Siva-Prasada et al. 1983; Weiss 1959) or because the test species was not obtained in North America and was not identified well enough to determine if it is resident in North America (e.g., Lahav and Sarig 1969). Tarpley (1958) conducted tests with brine shrimp, which species are too atypical to be used in deriving national criteria. Data were not used if parathion was a component of a mixture (e.g., Macek 1975) or if the test chamber contained sediment (D'Asaro

1982; Farr 1977,1978). Cole and Plapp (1974) did not verify that the parathion was dissolved off the test tubes by the test solution.

Anderson (1959), Chiou et al. (1977), Henderson et al. (1959), LeBlanc (1984), Ramke (1969), Sato and Kubo (1965), Surber (1948), and Tarzwell (1959a,b) only presented data that had been published elsewhere. Some studies were not used because test procedures or materials were not adequately described (e.g., Gillies et al. 1974; Hart and Womeldorf 1977; Kleerekoper 1974; Konar and Basak 1973; Lahav and Sarig 1969; Lewallen and Wilder 1962; Micks and Rougeau 1977; Moore 1970; Mulla 1980; Wilder and Schaefer 1969; and Zboray and Gutierrez 1979).

Data were not used if the organisms were exposed to parathion by injection or gavage or in food (e.g., Benke et al. 1973,1974; Carlson 1973; Hashimoto and Fukami 1969; King et al. 1984; Loeb and Kelly 1963; and Murphy et al. 1968).

Bradbury (1973a,b), Chambers (1976), Dortland (1978), Dortland et al. (1976), Estenik and Collins (1979), Goldsmith et al. (1976), Hiltibran (1974,1982), Hitchcock and Murphy (1971), Huddart (1978), Ludke et al. (1972), McDonald and Fingerman (1979), Murphy et al. (1968), Nollenberger (1982), Nollenberger et al. (1981), Schoor and Brausch 1980, Weiss (1959), Weiss and Gakstatter (1964,1965), Whitmore and Hodges (1978), and Yahalomi and Perry (1981) only exposed enzymes, excised tissues, or cell cultures or conducted other biochemical or histological studies. Bourquin et al. (1977a,b), Garnas and Crosby (1979), Lewis et al. (1984), Pritchard and Bourquin (1977), and Pritchard et al. (Manuscript a,b) only studied the metabolism of parathion.

Results of some laboratory tests were not used because the tests were conducted in distilled or deionized water without addition of appropriate salts (e.g., Burchfield and Storrs 1954; Goldsmith 1978; Goldsmith and Carlson 1979; Lewallen 1959,1962; Lichtenstein et al. 1966; and Yasuno et al. 1965) or if too few test organisms were exposed (e.g., Carlson 1973; Ludemann and Neumann 1961).

Hughes (1970,1973) did not acclimate the test organisms to the dilution water for a long enough period of time. Laboratory studies using formulations of parathion were not used (e.g., Alexander et al. 1982; Basak and Konar 1976a,b; Chang and Lange 1967; Davey et al. 1976; Gaufin et al. 1961; Hilsenhoff 1959; Labrecque et al. 1956; Mohamed and Gupta 1984; Panwar et al. 1982; Singh and Singh 1981, Sreenivasan and Swaminathan 1967; Srivastava et al. 1977; Verma et al. 1981). Field studies in which the concentration of parathion was not measured were not used (e.g., Ahmed and Washino 1977; Bengel and Fronk 1970; Chang and Lange 1967; Davey and Meisch 1977; Davey et al. 1976; Gahan 1957; Grigarick and Way 1982; Labrecque 1956; Mulla and Isaak 1961; Mulla et al. 1963,1964,1978; Myers et al. 1969; Stewart 1977).

High control mortalities occurred in tests reported by Fleming et al. (1982). High pesticide residues were found in field collected worms by Naqvi (1973), and the concentration of solvent was too high in studies by Poorman (1973).

Microcosm studies were not used (e.g., Dortland 1980; Francis et al. 1980; Miller et al. 1966; Yu and Sanborn 1975).

Results of laboratory bioconcentration tests were not used if the test was not flow-through or renewal (e.g., Verma and Gupta 1976) or the concentration in water was not measured (e.g., Kortus et al. 1971). A

bioconcentration study by Schmidt and Weidaas (1961) was not used because radio-labeled parathion was not adequately identified as the source of residue radioactivity. Reports of concentrations of parathion in wild aquatic organisms (e.g., Badawy and El-Dib 1984; Bradbury 1973a,b; Butler and Schutzmann 1978) were not used to calculate bioaccumulation factors if the number of measurements of the concentration was too small or if the range of the measured concentrations was too great.

Summary

The acute values for thirty-five freshwater species in twenty-nine genera range from 0.04 $\mu\text{g/L}$ for an early instar of a crayfish, Orconectes nais, to 5,230 $\mu\text{g/L}$ for two species of tubifid worms. Chronic toxicity values are available for two freshwater fish species, the bluegill and the fathead minnow, with chronic values of 0.24 $\mu\text{g/L}$ and 6.3 $\mu\text{g/L}$, and acute-chronic ratios of 2,121 and 79.45, respectively. Two freshwater algae were affected by toxaphene concentrations of 30 and 390 $\mu\text{g/L}$, respectively. Bioconcentration factors determined with three fish species ranged from 27 to 573.

The acute values that are available for saltwater species are 11.5 and 17.8 $\mu\text{g/L}$ for the Korean shrimp, Palaemon macrodactylus and 17.8 $\mu\text{g/L}$ for the striped bass, Morone saxitalis. No data are available concerning the chronic toxicity of parathion to saltwater species, toxicity to saltwater plants, or bioaccumulation by saltwater species. Some data indicate that parathion is acutely lethal to commercially important saltwater shrimp at concentrations as low as 0.24 $\mu\text{g/L}$. Measurement of AChE might be useful for diagnosing fish kills caused by parathion.

National Criteria

The procedures described in the "Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Aquatic Organisms and Their Uses" indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms and their uses should not be affected unacceptably if the four-day average concentration of parathion does not exceed 0.013 $\mu\text{g/L}$ more than once every three years on the average and if the one-hour average concentration does not exceed 0.065 $\mu\text{g/L}$ more than once every three years on the average.

The procedures described in the "Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Aquatic Organisms and Their Uses" require the availability of specified data for the derivation of a criterion. Very few of the required data are available concerning effects on parathion on saltwater species.

The allowed average excursion frequency of three years is the Agency's best scientific judgment of the average amount of time it will take an unstressed aquatic ecosystem to recover from a pollution event in which exposure to parathion exceeds the criterion. Stressed systems, for example one in which several outfalls occur in a limited area, would be expected to require more time for recovery. The resiliencies of ecosystems and their abilities to recover differ greatly, however, and site-specific criteria may be established if adequate justification is provided.

Use of criteria for developing water quality-based permit limits and for designing waste treatment facilities requires selection of an appropriate wasteload allocation model. Dynamic models are preferred for the application of these criteria. Limited data or other considerations might make their use impractical, in which case one must rely on a steady-state model. The

Agency recommends interim use of 1Q5 and 1Q10 for the Criterion Maximum Concentration (CMC) design flow and 7Q5 and 7Q10 for the Criterion Continuous Concentration (CCC) design flow in steady-state models for unstressed and stressed systems, respectively. These matters are discussed in more detail in the Technical Support Document for Water Quality-Based Toxics Control (U.S. EPA 1985).

Table 1. Acute Toxicity of Parathion to Aquatic Animals

<u>Species</u>	<u>Method</u> [#]	<u>Chemical</u> [#]	<u>LC50 or EC50 (µg/L)</u> ^{***}	<u>Species Mean Acute Value (µg/L)</u>	<u>Reference</u>
<u>FRESHWATER SPECIES</u>					
<u>Tubificid worm, Limnodrilus sp.</u>	S, U	Analytical (99.6%)	5,230****	5,230	Whitten and Goodnight 1966
<u>Tubificid worm, Tubifex sp.</u>	S, U	Analytical (99.6%)	5,230****	5,230	Whitten and Goodnight 1966
<u>Cladoceran, Daphnia magna</u>	S, U	-	1.8	-	Bringmann and Kuhn 1960
<u>Cladoceran (<24 hr), Daphnia magna</u>	S, M	Reagent (99%)	1.27	-	Spacie 1976; Spacie et al. 1981
<u>Cladoceran (<24 hr), Daphnia magna</u>	S, U	Analytical (99%)	1.3	-	Dortland 1980
<u>Cladoceran (<24 hr), Daphnia magna</u>	F, M	Reagent (99%)	1.0	1.0	Spacie 1976; Spacie et al. 1981
<u>Cladoceran (1st instar), Daphnia pulex</u>	S, U	Technical (98.7%)	0.60	0.60	Johnson and Finley 1980
<u>Cladoceran (1st instar), Simoccephalus serrulatus</u>	S, U	Technical (98.7%)	0.47	0.47	Johnson and Finley 1980
<u>Isopod, Asellus brevicaudus</u>	S, U	Technical (98.7%)	600	-	Sanders 1972
<u>Isopoda (mature), Asellus brevicaudus</u>	S, U	Technical (98.7%)	2,130	1,130	Johnson and Finley 1980
<u>Amphipod (mature), Gammarus fasciatus</u>	S, U	Technical (98.7%)	2.1 [†]	-	Sanders 1972
<u>Amphipod (mature), Gammarus fasciatus</u>	F, U	Technical (98.7%)	4.5 [†]	-	Sanders 1972
<u>Amphipod (mature), Gammarus fasciatus</u>	S, U	Technical (98.7%)	1.3 [†]	-	Johnson and Finley 1980; Sanders 1972

15

Table 1. (continued)

<u>Species</u>	<u>Method</u> ^a	<u>Chemical</u> ^{b,c}	<u>LC50 or EC50 ($\mu\text{g/L}$)^d</u>	<u>Species Mean Acute Value ($\mu\text{g/L}$)</u>	<u>Reference</u>
<u>Amphipod (immature), Gammarus fasciatus</u>	F, M	Reagent (99%)	0.43	-	Spacie 1976; Spacie et al. 1981
<u>Amphipod (immature), Gammarus fasciatus</u>	F, M	Reagent (99%)	0.62	-	Spacie 1976; Spacie et al. 1981
<u>Amphipod (immature), Gammarus fasciatus</u>	F, M	Reagent (99%)	0.26	-	Spacie 1976; Spacie et al. 1981
<u>Amphipod (immature), Gammarus fasciatus</u>	F, M	Reagent (99%)	0.25	0.3628	Spacie 1976; Spacie et al. 1981
<u>Amphipod (mature), Gammarus lacustris</u>	S, U	Technical (98.7%)	3.5	3.5	Johnson and Finley 1980; Sanders 1969
<u>Prawn, Palaemonetes kadiakensis</u>	F, U	Technical (98.7%)	5.0	-	Sanders 1972
<u>Prawn (mature), Palaemonetes kadiakensis</u>	S, U	Technical (98.7%)	1.5	2.739	Johnson and Finley 1980; Sanders 1972
<u>Crayfish (mature), Orconectes nais</u>	S, U	Technical (98.7%)	15 [†]	-	Sanders 1972
<u>Crayfish (early instar), Orconectes nais</u>	S, U	Technical (98.7%)	0.04	0.04	Sanders 1972; Johnson and Finley 1972
<u>Crayfish (mature), Procambarus sp.</u>	S, U	Technical (98.7%)	<250	<250	Johnson and Finley 1980
<u>Phantom midge, Chaoborus sp.</u>	S, U	-	48 ^{††}	-	Collins and Shank 1983
<u>Phantom midge, Chaoborus sp.</u>	S, U	-	0.8 ^{†††}	-	Collins and Shank 1983
<u>Phantom midge, Chaoborus sp.</u>	S, U	-	1.0 ^{†††}	0.8944	Collins and Shank 1983

91

Table 1. (continued)

<u>Species</u>	<u>Method^a</u>	<u>Chemical^{ab}</u>	<u>LC50 or EC50 ($\mu\text{g/L}$)^{abc}</u>	<u>Species Mean Acute Value ($\mu\text{g/L}$)</u>	<u>Reference</u>
<u>Mayfly, Clooson dipterum</u>	S, U	Analytical (99%)	2.5	-	Dortland 1980
<u>Mayfly, Clooson dipterum</u>	S, U	Analytical (99%)	2.6	-	Dortland 1980
<u>Mayfly, Clooson dipterum</u>	R, U	Analytical (99%)	1.7	2.227	Dortland 1980
<u>Mayfly (juvenile), Hexagenia bilineata</u>	S, U	Technical (98.7%)	15	15	Johnson and Finley 1980
<u>Damselfly (juvenile), Ischnura ventralis</u>	S, U	Technical (98.7%)	0.64	0.64	Johnson and Finley 1980
<u>Damselfly, Lestes congener</u>	S, U	Technical (>94%)	3.0	3.0	Federle and Collins 1976
<u>Stonefly, Pteronarcys badia</u>	S, U	Technical (98.7%)	4.2	4.2	Johnson and Finley 1980; Sanders and Cope 1968
<u>Stonefly (naid), Pteronarcys californica</u>	S, U	Technical (95%)	32	-	Jensen and Gauflin 1964
<u>Stonefly (2nd year class), Pteronarcys californica</u>	S, U	Technical (98.7%)	5.4	13.15	Johnson and Finley 1980; Sanders and Cope 1968
<u>Stonefly (naid), Acronuria pacifica</u>	S, U	Technical (95%)	2.9	2.9	Jensen and Gauflin 1964
<u>Stonefly (2nd year class), Claassenia sabulosa</u>	S, U	Technical (98.7%)	1.5	1.5	Johnson and Finley 1980; Sanders and Cope 1968

Table 1. (continued)

<u>Species</u>	<u>Method</u> ^a	<u>Chemical</u> ^{bb}	<u>LC50 or EC50 (µg/L)</u> ^{ccc}	<u>Species Mean Acute Value (µg/L)</u>	<u>Reference</u>
Crawling water beetle (adult), <u>Peltodytes</u> sp.	S, U	Technical (>94%)	7.0	7.0	Federle and Collins 1976
Midge, <u>Chironomus riparius</u>	S, U	-	8.4 ^{††}	-	Collins and Shank 1983
Midge, <u>Chironomus riparius</u>	S, U	-	1.6 ^{†††}	-	Collins and Shank 1983
Midge, <u>Chironomus riparius</u>	S, U	-	1.8 ^{†††}	1.697	Collins and Shank 1983
Chironomid (4th instar), <u>Chironomus tentans</u>	F, M	Reagent (99%)	31.0	31.0	Spacie 1976; Spacie et al. 1981
Cutthroat trout (0.3 g), <u>Salmo clarki</u>	S, U	Technical (98.7%)	1,560	1,560	Johnson and Finley 1980
Rainbow trout (1.0 g), <u>Salmo gairdneri</u>	S, U	Technical (98.7%)	1,430	-	Johnson and Finley 1980
Rainbow trout (embryo, 0 hr), <u>Salmo gairdneri</u>	R, U	(99%)	10,000 [†]	-	Van Leeuwen et al. 1985
Rainbow trout (embryo, 24 hr), <u>Salmo gairdneri</u>	R, U	(99%)	10,000 [†]	-	Van Leeuwen et al. 1985
Rainbow trout (embryo, 14 day), <u>Salmo gairdneri</u>	R, U	(99%)	10,000 [†]	-	Van Leeuwen et al. 1985
Rainbow trout (embryo, 28 day), <u>Salmo gairdneri</u>	R, U	(99%)	10,000 [†]	-	Van Leeuwen et al. 1985
Rainbow trout (fry, 42 day), <u>Salmo gairdneri</u>	R, U	(99%)	10,000 [†]	-	Van Leeuwen et al. 1985

81

Table 1. (continued)

<u>Species</u>	<u>Method</u> ^a	<u>Chemical</u> ^b	<u>LC50 or EC50 ($\mu\text{g/L}$)^c</u>	<u>Species Mean Acute Value ($\mu\text{g/L}$)</u>	<u>Reference</u>
Rainbow trout (fry, 77 day), <u>Salmo gairdneri</u>	R, U	(99%)	1,400	1,415	Van Leeuwen et al. 1985
Brown trout (16-19 cm), <u>Salmo trutta</u>	F, M	Reagent (99%)	1,510	1,510	Spacie 1976; Spacie et al. 1981
Brook trout (juvenile), <u>Salvelinus fontinalis</u>	F, M	Reagent (99%)	1,760	1,760	Spacie 1976; Spacie et al. 1981
Lake trout (0.7 g), <u>Salvelinus namaycush</u>	S, U	Technical (98.7%)	1,920	1,920	Johnson and Finley 1980
Goldfish (juvenile), <u>Carassius auratus</u>	S, U	Technical (99%)	2,700	-	Pickering et al. 1962
Goldfish (0.9 g), <u>Carassius auratus</u>	S, U	Technical (98.7%)	1,830	2,223	Johnson and Finley 1980
Fathead minnow (adult), <u>Pimephales promelas</u>	S, M	Reagent (99%)	1,600	-	Spacie 1976; Spacie et al. 1981
Fathead minnow (1-1.5 g), <u>Pimephales promelas</u>	S, U	Technical (96.5%)	1,400	-	Henderson and Pickering 1958
Fathead minnow (1-1.5 g), <u>Pimephales promelas</u>	S, U	Technical (96.5%)	1,600	-	Henderson and Pickering 1958
Fathead minnow (1-1.5 g), <u>Pimephales promelas</u>	S, U	Technical (96.5%)	2,800	-	Henderson and Pickering 1958
Fathead minnow (1-1.5 g), <u>Pimephales promelas</u>	S, U	Technical (96.5%)	3,700	-	Henderson and Pickering 1958
Fathead minnow (juvenile), <u>Pimephales promelas</u>	S, U	Technical (99%)	1,300	-	Pickering et al. 1962

19

Table 1. (continued)

<u>Species</u>	<u>Method^a</u>	<u>Chemical^{ab}</u>	<u>LC50 or EC50 (μg/L)^{***}</u>	<u>Species Mean Acute Value (μg/L)</u>	<u>Reference</u>
<u>Fathead minnow (0.8 g), Pimephales promelas</u>	S, U	Technical (98.7%)	2,350	-	Johnson and Finley 1980
<u>Fathead minnow (1.8-4.0 cm), Pimephales promelas</u>	F, M	Analytical (98.7%)	1,410	-	Solon et al. 1969; Solon and Mair 1970
<u>Fathead minnow (adult), Pimephales promelas</u>	F, M	Reagent (99%)	500	839.6	Spacie 1976; Spacie et al. 1981
<u>Channel catfish (1.4 g), Ictalurus punctatus</u>	S, U	Technical (98.7%)	2,650	2,650	Johnson and Finley 1980
<u>Mosquitofish (1.1 g), Gambusia affinis</u>	S, U	Technical (98.7%)	320	320	Johnson and Finley 1980
<u>Guppy (6 mo), Poecilia reticulata</u>	S, U	Technical (99%)	56	56	Pickering et al. 1962
<u>Green sunfish (1.1 g), Lepomis cyanellus</u>	S, U	Technical (98.7%)	930	930	Johnson and Finley 1980
<u>Bluegill (juvenile), Lepomis macrochirus</u>	S, U	Technical (99%)	95	-	Pickering et al. 1962
<u>Bluegill (1.5 g), Lepomis macrochirus</u>	S, U	Technical (96.5%)	710	-	Henderson and Pickering 1958
<u>Bluegill (1.0 g), Lepomis macrochirus</u>	S, U	Technical (98.7%)	400	-	Johnson and Finley 1980
<u>Bluegill (juvenile), Lepomis macrochirus</u>	F, M	Reagent (99%)	510	510	Spacie 1976; Spacie et al. 1981
<u>Largemouth bass (0.7 g), Micropterus salmoides</u>	S, U	Technical (98.7%)	620	620	Johnson and Finley 1980
<u>Western chorus frog (1 wk), Pseudacris triseriata</u>	S, U	Technical (98.7%)	1,000	1,000	Sanders 1970

20

Table 1. (continued)

<u>Species</u>	<u>Method</u> [#]	<u>Chemical</u> ^{**}	<u>LC50 or EC50 ($\mu\text{g/L}$)</u> ^{***}	<u>Species Mean Acute Value ($\mu\text{g/L}$)</u>	<u>Reference</u>
<u>SALTWATER SPECIES</u>					
<u>Korean shrimp (adult), Palaeomon macrodactylus</u>	F, U	(99%)	17.8	-	Earnest 1970
<u>Korean shrimp (adult), Palaeomon macrodactylus</u>	S, U	(99%)	11.5	14.31	Earnest 1970
<u>Striped bass (juvenile), Morone saxatilis</u>	F, U	(99%)	17.8	17.8	Korn and Earnest 1974

* S = static; R = renewal; F = flow-through; U = unmeasured; M = measured.

** Percent purity is given in parentheses when available.

*** If the concentrations were not measured and the published results were not reported to be adjusted for purity, the published results were multiplied by the purity if it was reported to be less than 97%.

**** Limnodrilus sp. and Tubifex sp. were tested together, but appeared to be equally resistant.

† Not used in calculation of Species Mean Acute Value because data are available for a more sensitive life stage.

†† 4°C; not used in calculations.

††† 22°C.

21

Table 2. Chronic Toxicity of Parathion to Aquatic Animals

<u>Species</u>	<u>Test*</u>	<u>Chemical**</u>	<u>Limits (µg/L)***</u>	<u>Chronic Value (µg/L)</u>	<u>Reference</u>
<u>FRESHWATER SPECIES</u>					
<u>Cladoceran, Daphnia magna</u>	LC	Reagent (99%)	0.0817-0.12	0.0990	Spacie 1976; Spacie et al. 1981
<u>Fathead minnow, Pimephales promelas</u>	LC	Reagent (99%)	4.4-9.0	6.293	Spacie 1976; Spacie et al. 1981
<u>Bluegill, Lepomis macrochirus</u>	LC	Reagent (99%)	0.17-0.34	0.2404	Spacie 1976; Spacie et al. 1981

* LC = life-cycle or partial life-cycle.

** Percent purity is given in parentheses when available.

*** Results are based on measured concentrations of parathion.

22

<u>Species</u>	<u>Acute-Chronic Ratio</u>		
	<u>Acute Value (µg/L)</u>	<u>Chronic Value (µg/L)</u>	<u>Ratio</u>
<u>Cladoceran, Daphnia magna</u>	1.00	0.0990	10.10
<u>Fathead minnow, Pimephales promelas</u>	500	6.293	79.45
<u>Bluegill, Lepomis macrochirus</u>	510	0.2404	2,121

Table 3. Ranked Genus Mean Acute Values with Species Mean Acute-Chronic Ratios

<u>Rank^a</u>	<u>Genus Mean Acute Value (µg/L)</u>	<u>Species</u>	<u>Species Mean Acute Value (µg/L)^{ab}</u>	<u>Species Mean Acute-Chronic Ratio^{abc}</u>
<u>FRESHWATER SPECIES</u>				
31	5,230	<u>Tubificid worm, Tubifex sp.</u>	5,230	-
30	5,230	<u>Tubificid worm, Limnodrilus sp.</u>	5,230	-
29	2,650	<u>Channel catfish, Ictalurus punctatus</u>	2,650	-
28	2,223	<u>Goldfish, Carassius auratus</u>	2,223	-
27	1,838	<u>Brook trout, Salvelinus fontinalis</u>	1,760	-
		<u>Lake trout, Salvelinus namaycush</u>	1,920	-
26	1,494	<u>Cutthroat trout, Salmo clarki</u>	1,560	-
		<u>Brown trout, Salmo trutta</u>	1,510	-
		<u>Rainbow trout, Salmo gairdneri</u>	1,415	-
25	1,130	<u>Isopod, Asellus brevicaudus</u>	1,130	-
24	1,000	<u>Western chorus frog, Pseudacris triseriata</u>	1,000	-
23	839.6	<u>Fathead minnow, Pimephales promelas</u>	839.6	79.45
22	688.7	<u>Green sunfish, Lepomis cyanellus</u>	930	-
		<u>Bluegill, Lepomis macrochirus</u>	510	2,121

23

Table 3. (continued)

<u>Rank^a</u>	<u>Genus Mean Acute Value ($\mu\text{g/L}$)</u>	<u>Species</u>	<u>Species Mean Acute Value ($\mu\text{g/L}$)^b</u>	<u>Species Mean Acute-Chronic Ratio^c</u>
21	620	Largemouth bass, <u>Micropterus salmoides</u>	620	-
20	320	Mosquitofish, <u>Gambusia affinis</u>	320	-
19	<250	Crayfish, <u>Procambarus sp.</u>	<250	-
18	56	Guppy, <u>Poecilia reticulata</u>	56	-
17	31.0	Midge, <u>Chironomus tentans</u>	31.0	-
16	15	Mayfly, <u>Hexagenia bilineata</u>	15	-
15	13.15	Stonefly, <u>Pteronarcys californica</u>	13.15	-
14	7.0	Beetle, <u>Peltodytes spp.</u>	7.0	-
13	4.2	Stonefly, <u>Pteronarcella badia</u>	4.2	-
12	3.0	Damselfly, <u>Lestes congener</u>	3.0	-
11	2.9	Stonefly, <u>Acronuria pacifica</u>	2.9	-
10	2.739	Prawn, <u>Palaeomonetes kadlakensis</u>	2.739	-
9	2.227	Mayfly, <u>Cloeon dipterum</u>	2.227	-

42

Table 3. (continued)

Rank ^a	Genus Mean Acute Value (µg/L)	Species	Species Mean Acute Value (µg/L) ^{**}	Species Mean Acute-Chronic Ratio ^{***}
8	1.697	Midge, <u>Chironomus riparius</u>	1.697	-
7	1.5	Stonefly, <u>Cliaassenia sabulosa</u>	1.5	-
6	1.127	Amphipod, <u>Gammarus fasciatus</u>	0.3628	-
		Amphipod, <u>Gammarus lacustris</u>	3.5	-
5	0.8944	Phantom midge, <u>Chaoborus sp.</u>	0.8944	-
4	0.7746	Cladoceran, <u>Daphnia magna</u>	1.0	10.10
		Cladoceran, <u>Daphnia pulex</u>	0.60	-
3	0.64	Damselfly, <u>Ischnura ventralis</u>	0.64	-
2	0.47	Cladoceran, <u>Simocephalus serrulatus</u>	0.47	-
1	0.04	Crayfish, <u>Orconectes nais</u>	0.04	-

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value. Inclusion of "greater than" values does not necessarily imply a true ranking, but does allow use of all genera for which data are available so that the Final Acute Value is not unnecessarily lowered.

** From Table 1.

*** From Table 2.

25

Table 3. (continued)

Fresh water

Final Acute Value = 0.1298 $\mu\text{g/L}$

Criterion Maximum Concentration = $(0.1298 \mu\text{g/L}) / 2 = 0.0649 \mu\text{g/L}$

Final Acute-Chronic Ratio = 10.10 (see text)

Final Chronic Value = $(0.1298 \mu\text{g/L}) / 10.10 = 0.01285 \mu\text{g/L}$

Table 4. Toxicity of Parathion to Aquatic Plants

<u>Species</u>	<u>Chemical</u> [*]	<u>Duration</u> <u>(days)</u>	<u>Effect</u>	<u>Result</u> <u>($\mu\text{g/L}$)^{**}</u>	<u>Reference</u>
<u>FRESHWATER SPECIES</u>					
Blue-green alga, <u>Microcystis aeruginosa</u>	-	8	Incipient inhibition	30 (34)	Bringmann and Kuhn 1978a,b
Green alga, <u>Scenedesmus quadricauda</u>	-	8	Incipient inhibition	390	Bringmann and Kuhn 1977; 1978a,b

* Percent purity is given in parentheses when available.

** If the concentrations were not measured and the published results were not reported to be adjusted for purity, the published results were multiplied by the purity if it was reported to be less than 97%.

Table 5. Bioaccumulation of Parathion by Aquatic Organisms

<u>Species</u>	<u>Chemical</u> ^a	<u>Concentration in Water (µg/L)</u> ^{**}	<u>Duration (days)</u>	<u>Tissue</u>	<u>BCF or BAF</u> ^{***}	<u>Reference</u>	
<u>FRESHWATER SPECIES</u>							
Brook trout, <u>Salvelinus fontinalis</u>	Reagent (99%)	0.6	180	Muscle	258	Spacie 1976; Spacie et al. 1981	
		0.6			312		
		1.4			299		
		2.6			439		
		4.0			471		
		6.7			573		
		0.44			260		124
		0.53					86
		1.26					31
		1.45					43
		2.76					99
		2.86					91
		4.24					86
		5.53					88
		8.30					232
8.72	179						
Fathead minnow, <u>Pimephales promelas</u>	Reagent (99%)	0.15	260	Whole body	93.3	Spacie 1976; Spacie et al. 1981	
		4.2			169.4		
		9.0			104.6		
		15.5			32.9		
		21.7			66.8		
		49.0			201.4		
Bluegill, <u>Lepomis macrochirus</u>	Reagent (99%)	4.00	540	Muscle	27	Spacie 1976; Spacie et al. 1981	

* Percent purity is given in parentheses when available.

** Measured concentration of parathion.

*** Bioconcentration factors (BCFs) and bioaccumulation factors (BAFs) are based on measured concentrations of parathion in water and in tissue.

Table 6. Other Data on the Effects of Parathion on Aquatic Organisms

<u>Species</u>	<u>Chemical*</u>	<u>Duration</u>	<u>Effect</u>	<u>Result (µg/L)**</u>	<u>Reference</u>
<u>FRESHWATER SPECIES</u>					
<u>Bacterium, Pseudomonas putida</u>	-	16 hr	Incipient Inhibition	>1	Bringmann and Kuhn 1977
<u>Ciliate, Colpidium campylum</u>	-	43 hr	Change in growth rate	10,000	Dive et al. 1980
<u>Worm, Tubifex tubifex</u>	-	18 hr 18-36 hr	Onset of symptoms Onset of death	10,000 100,000	Ludemann and Neumann 1960b
<u>Cladoceran, Daphnia magna</u>	-	24 hr	LC50	4	Ghetti and Gorbi 1985
<u>Cladoceran (<24 hr old), Daphnia magna</u>	-	26 hr	LC50	0.8	Frear and Boyd 1967
<u>Cladoceran, Daphnia magna</u>	Reagent (99%)	7 days 14 days 21 days	EC50	0.39 0.31 0.16	Spacie 1976; Spacie et al. 1981
<u>Cladoceran (adult), Daphnia pulex</u>	Technical	3 hr	LC50	0.8	Nishiyuchi and Hashimoto 1967, 1969
<u>Cladoceran (adult), Moina macrocopa</u>	Technical	3 hr	LC50	8.1	Nishiyuchi and Hashimoto 1967, 1969
<u>Prawn, Palaemonetes kadiakensis</u>	Technical	24 hr	LC50	7.1 11.8† 7.4† 6.6†	Naqvi and Ferguson 1970
<u>Mayfly, Stenonema femoratum</u>	-	48 hr	EC50 (4 °C) (22 °C)	30.0 1.7	Collins and Shank 1983
<u>Mayfly, Stenonema vicarium</u>	-	48 hr	EC50	29.0	Collins and Shank 1983
<u>Mayfly, (larva), Baetis rhodani</u>	-	65 min	LT50	1,000	Ghetti and Gorbi 1985

29

Table 6. (continued)

<u>Species</u>	<u>Chemical^a</u>	<u>Duration</u>	<u>Effect</u>	<u>Result ($\mu\text{g/L}$)^{ab}</u>	<u>Reference</u>
Stonefly, <u>Allocaonia sp.</u>	-	48 hr	EC50	2.2	Collins and Shank 1983
Beetle (larva), <u>Hydrophilus triangularis</u>	Technical	24 hr	LC50	17	Ahmed 1977
Beetle (adult), <u>Hygrotus sp.</u>	Technical	24 hr	LC50	28	Ahmed 1977
Beetle (adult), <u>Laccophilus decipiens</u>	Technical	24 hr	LC50	12	Ahmed 1977
Beetle (adult), <u>Thermonectus basillaris</u>	Technical	24 hr	LC50	1.8	Ahmed 1977
Beetle (adult), <u>Tropisternus lateralis</u>	Technical	24 hr	LC50	32	Ahmed 1977
Beetle (larva), <u>Tropisternus lateralis</u>	Technical	24 hr	LC50	40	Ahmed 1977
Water bug (adult), <u>Belostoma sp.</u>	Technical	24 hr	LC50	60	Ahmed 1977
Caddisfly, <u>Cheumatopsyche sp.</u>	-	48 hr	EC50 (4°C) (22°C)	21.0 2.5	Collins and Shank 1983
Caddisfly (larva), <u>Hydropsyche pellucidula</u>	-	110 min	LT50	1,000	Ghetti and Gorbi 1985
Caddisfly, <u>Hydropsyche sp.</u>	-	48 hr	EC50 (4°C) (22°C)	36.0 1.3	Collins and Shank 1983
Mosquito (4th instar) <u>Aedes aegypti</u>	32-P labeled	24 hr	LC50	4.8	Schmidt and Weidaas 1961
Mosquito (larva), <u>Aedes nigromaculis</u>	Technical	24 hr	LC50	40 35 3.5	Mulla et al. 1970

Table 6. (continued)

<u>Species</u>	<u>Chemical</u> ^a	<u>Duration</u>	<u>Effect</u>	<u>Result</u> <u>($\mu\text{g/L}$)^{b,c}</u>	<u>Reference</u>
Mosquito (4th instar), <u>Aedes nigromaculis</u>	Technical	24 hr	LC50	27 68	Mulla et al. 1978
Mosquito (4th instar), <u>Aedes taeniorhynchus</u>	32-P labeled	24 hr	LC50	3.6	Schmidt and Woldaas 1961
Mosquito (4th instar), <u>Anopheles freeborni</u>	Technical	24 hr	LC50	2.2-15.0 (24 values)	Womeldorf et al. 1970
Mosquito (larval), <u>Anopheles freeborni</u>	Technical	24 hr	LC50	0.7	Ahmed 1977
Mosquito (4th instar), <u>Anopheles quadrimaculatus</u>	32-P labeled	24 hr	LC50	6.0	Schmidt and Woldaas 1961
Mosquito (4th instar), <u>Culex pipiens</u>	Technical	24 hr	LC50	4.5	Mulla et al. 1962
Mosquito (4th instar), <u>Culex pipiens</u>	Technical	24 hr	LC50	4.5	Mulla et al. 1964
Mosquito (3rd-4th instar), <u>Culex pipiens</u>	Technical	24 hr	LC50	0.45 5.0	Chen et al. 1971
Mosquito (larval), <u>Culex tarsalis</u>	Technical	24 hr	LC50	5.8	Ahmed 1977
Midge (larval), <u>Chironomus plumosus</u>	-	24 hr	LC50	39	Ludemann and Neumann 1960c
Midge (4th instar), <u>Chironomus riparius</u>	Technical	24 hr	LC50	2.5	Estenik and Collins 1979
Midge (2nd and 4th instar), <u>Chironomus tentans</u>	Reagent (99%)	1 day 2 day 5 day 8 day 14 day	LC50	660 135 7.3 2.2 2.6	Spacie 1976; Spacie et al. 1981

31

Table 6. (continued)

<u>Species</u>	<u>Chemical^a</u>	<u>Duration</u>	<u>Effect</u>	<u>Result ($\mu\text{g/L}$)^{ab}</u>	<u>Reference</u>
Brown trout, <u>Salmo trutta</u>	Reagent (99%)	64 hr	BCF	61 77	Spacie 1976; Spacie et al. 1981
Brook trout, <u>Salvelinus fontinalis</u>	Reagent (99%)	-	LC50	75	Spacie 1976; Spacie et al. 1981
Brook trout, <u>Salvelinus fontinalis</u>	Reagent (99%)	-	Reduced percent hatch	10	Spacie 1976; Spacie et al. 1981
Brook trout, <u>Salvelinus fontinalis</u>	Reagent (99%)	8 hr 114 hr 140 hr 144 hr	BCF	88.5 102.5 301.5 192.5	Spacie 1976; Spacie et al. 1981
Goldfish (1.0 g), <u>Cyprinus auratus</u>	Technical	48 hr	LC50	1,700	Nishiyuchi and Hashimoto 1967,1969
Common carp (3.9 g), <u>Cyprinus carpio</u>	-	48 hr	LC50	3,500	Ludemann and Neumann 1960a
Common carp (1.1 g), <u>Cyprinus carpio</u>	Technical	48 hr	LC50	3,200	Nishiyuchi and Hashimoto 1967,1969
Golden shiner, (DDT-susceptible), <u>Notemigonus crysoleucas</u>	Technical	48 hr	LC50	1,895	Minchew and Ferguson 1970
Golden shiner (DDT-resistant), <u>Notemigonus crysoleucas</u>	Technical	48 hr	LC50	2,800	Minchew and Ferguson 1970
Golden shiner, <u>Notemigonus crysoleucas</u>	-	24 hr	LC50	931	Gibson 1971

W
N

Table 6. (continued)

<u>Species</u>	<u>Chemical</u> ^a	<u>Duration</u>	<u>Effect</u>	<u>Result</u> (<u>µg/L</u>) ^{b,c}	<u>Reference</u>
<u>Fathead minnow</u> (DDT-susceptible), <u>Pimephales promelas</u>	Technical	48 hr	LC50	48	Culley and Ferguson 1969
<u>Fathead minnow</u> (DDT-resistant), <u>Pimephales promelas</u>	Technical	48 hr	LC50	199	Culley and Ferguson 1969
<u>Mosquitofish,</u> <u>Gambusia affinis</u>	Technical	24 hr	LC50	140	Ahmed and Washino 1977
<u>Mosquitofish (15-30 mg),</u> <u>Gambusia affinis</u>	Analytical	24 hr	LC50	1,400	Krieger and Lee 1973
<u>Mosquitofish (adult)</u> (DDT-resistant), <u>Gambusia affinis</u>	Analytical (99%)	48 hr	LC50	390 950	Chambers and Yarbough 1974
<u>Mosquitofish (adult)</u> (DDT-susceptible), <u>Gambusia affinis</u>	Analytical (99%)	48 hr	LC50	350 610	Chambers and Yarbough 1974
<u>Guppy,</u> <u>Poecilia reticulata</u>	-	72 hr	LC50	29	Nagasawa et al. 1968
<u>Guppy (7 wk old),</u> <u>Poecilia reticulata</u>	Technical	24 hr	LC50	80 45	Chen et al. 1971
<u>Green sunfish</u> (DDT-susceptible), <u>Lepomis cyanellus</u>	Technical	48 hr	LC50	207	Minchew and Ferguson 1970
<u>Green sunfish</u> (DDT-resistant), <u>Lepomis cyanellus</u>	Technical	48 hr	LC50	275	Minchew and Ferguson 1970
<u>Green sunfish,</u> <u>Lepomis cyanellus</u>	-	24 hr	LC50	155	Gibson 1971
<u>Bluegill,</u> <u>Lepomis macrochirus</u>	-	24 hr	LC50	141	Gibson 1971

33

Table 6. (continued)

<u>Species</u>	<u>Chemical^a</u>	<u>Duration</u>	<u>Effect</u>	<u>Result ($\mu\text{g/L}$)^{b,c}</u>	<u>Reference</u>
<u>Bluegill,</u> <u>Lepomis macrochirus</u>	Reagent (99%)	12 hr	BCF	80.5	Spacie 1976; Spacie et al. 1981
		18 hr		145	
		24 hr		173	
		29 hr		175.3	
		46 hr		253.0	
		70 hr		311	
		72 hr		462	
<u>Largemouth bass,</u> <u>Micropterus salmoides</u>	-	24 hr	Change in opercular rhythm	160	Morgan 1976
<u>Frog (tadpole),</u> <u>Rana catesbeiana</u>	-	96 hr	BCF	50.1	Hall and Kolbe 1980
<u>SALTWATER SPECIES</u>					
<u>Natural phytoplankton</u> <u>communities</u>	(99.6%)	4 hr	9.9% decrease in population growth	1,000	Butler 1964
<u>Eastern oyster (juvenile),</u> <u>Crassostrea virginica</u>	-	96 hr	EC50 (shell deposition)	850	Butler 1963
<u>Eastern oyster (juvenile),</u> <u>Crassostrea virginica</u>	(99.6%)	96 hr	22% reduction in shell deposition	1,000	Butler 1964; Lowe et al 1970; U.S. Bureau of Commercial Fisheries 1966
<u>Eastern oyster (juvenile</u> <u>to adult),</u> <u>Crassostrea virginica</u>	(99.6%)	336 days	No significant effects on growth	0.8	Lowe et al. 1971
<u>Eastern oyster (larva),</u> <u>Crassostrea virginica</u>	-	12 days	78% reduction in average length	1,000	Davis and Hidu 1969
<u>Grass shrimp (juvenile),</u> <u>Palaemonetes pugio</u>	(99.6%)	48 hr	EC50 (mortality and loss of equilibrium)	2.8	U.S. Bureau of Commercial Fisheries 1967
<u>Grass shrimp,</u> <u>Palaemonetes pugio</u>	(99.6%)	24-72 hr	Increased predation by gulf killifish, <u>Fundulus grandis</u>	0.1-0.5	Farr 1977

34

Table 6. (continued)

<u>Species</u>	<u>Chemical</u> ^a	<u>Duration</u>	<u>Effect</u>	<u>Result</u> ($\mu\text{g/L}$) ^{ab}	<u>Reference</u>
Brown shrimp (adult), <u>Penaeus aztecus</u>	(99.6%)	48 hr	EC50 (mortality and loss of equilibrium)	1	Butler 1964; U.S. Bureau of Commercial Fisheries 1966
Pink shrimp (juvenile), <u>Penaeus duorarum</u>	(99.6%)	48 hr	EC50 (mortality and loss of equilibrium)	0.24	Lowe et al. 1970; U.S. Bureau of Commercial Fisheries 1967
Fiddler crab, <u>Uca pugilator</u>	(95%)	2-3 wk	No effect on limb regeneration or time to molt	10	Wels and Mantel 1976
Fiddler crab, <u>Uca pugilator</u>	(95%)	2-3 wk	100% mortality	100	Wels and Mantel 1976
Sheepshead minnow (juvenile), <u>Cyprinodon variegatus</u>	(99.6%)	48 hr	LC50	60	Butler 1964
Sheepshead minnow (juvenile), <u>Cyprinodon variegatus</u>	(99.6%)	48 hr	LC50	36	U.S. Bureau of Commercial Fisheries 1966
Sheepshead minnow (adult), <u>Cyprinodon variegatus</u>	-	2 hr	40-60% mortality; brain AChE activity reduced >82%	5,000	Coppage 1972
Sheepshead minnow (adult), <u>Cyprinodon variegatus</u>	-	24 hr	40-60% mortality; brain AChE activity reduced >82%	2,000	Coppage 1972
Sheepshead minnow (adult), <u>Cyprinodon variegatus</u>	-	48 hr	40-60% mortality; brain AChE activity reduced >82%	100	Coppage 1972
Sheepshead minnow (adult), <u>Cyprinodon variegatus</u>	-	72 hr	40-60% mortality; brain AChE activity reduced >82%	10	Coppage 1972
Sheepshead minnow (adult), <u>Cyprinodon variegatus</u>	-	120 hr	Greatest reduction (78-82%) in normal brain AChE activity obtained without causing death	5	Coppage 1972

35

Table 6. (continued)

<u>Species</u>	<u>Chemical</u> [*]	<u>Duration</u>	<u>Effect</u>	<u>Result</u> (<u>µg/L</u>) ^{**}	<u>Reference</u>
<u>Mummichog (adult), Fundulus heteroclitus</u>	(91%)	2 wk	Significant reduction in fin regeneration	10	Wels and Wels 1975
<u>Mummichog (embryo), Fundulus heteroclitus</u>	(95%)	3 days	50% incidence of circulatory failure	10,000	Wels and Wels 1974
<u>Longnose killifish (juvenile), (99.6%) Fundulus similis</u>		48 hr	LC50	15	Lowe et al. 1970
<u>Pinfish (65-125 mm), Lagodon rhomboides</u>	Technical	24 hr	40-60% mortality; brain AChE activity reduced 90%	10	Coppage and Matthews 1974
<u>Spot (juvenile), Leiostomus xanthurus</u>	(99.6%)	48 hr	LC50	18	U.S. Bureau of Commercial Fisheries 1966
<u>Spot (65-150 mm), Leiostomus xanthurus</u>	Technical	24 hr	40-60% mortality; brain AChE activity reduced 88%	10	Coppage and Matthews 1974
<u>Striped mullet (juvenile) Mugil cephalus</u>	(99.6%)	48 hr	LC50	100	U.S. Bureau of Commercial Fisheries 1967
<u>Laughing gull (chick), Larus atricilla</u>	-	Field collections	75-90% inhibition of brain ChE in dead chicks contaminated with parathion	-	White et al. 1979
<u>Laughing gull (adult), Larus atricilla</u>	-	Field collections	57-89% inhibition of brain ChE in dead adults contaminated with parathion	-	White et al. 1979

* Percent purity is given in parentheses when available.

** If the concentrations were not measured and the published results were not reported to be adjusted for purity, the published results were multiplied by the purity if it was reported to be less than 97%.

† Organisms collected at sites potentially contaminated by pesticides.

REFERENCES

- Ahmed, M.K. and J.E. Casida. 1958. Metabolism of some organophosphorus insecticides by microorganisms. *J. Econ. Entomol.* 51:59-63.
- Ahmed, W. 1977. A laboratory and field study on the toxicity of mosquito larvicides to non-target insects found in California rice fields. Ph.D. thesis, University of California-Davis, Davis, California.
- Ahmed, W. and R.K. Washino. 1977. Toxicity of pesticides used in rice culture in California on Gambusia affinis. Ph.D. thesis, University of California-Davis, Davis, California.
- Albright, L.J., A. Gasith, G.H. Green, Y. Mozel and A.S. Perry. 1983. The influence of organophosphorous insecticides acephate and parathion upon the heterotrophic bacteria of two freshwater ecosystems. In: *Agrochemical-biota interactions in soil and water using nuclear techniques*. Technical Document 283. International Atomic Energy Agency, Vienna. pp. 33-44.
- Alexander, T.C., M.V. Meisch, W.B. Kottkamp and A.L. Anderson. 1982. Effect of notonectids on mosquito larvae and preliminary observations of insecticide toxicity. *Ark. Farm Res.* 31:5.
- Anderson, B.G. 1959. The toxicity of organic insecticides to Daphnia. *Trans. 2nd Sem. Biol. Probl. Water Pollut.* 94-95.
- Banas, W.P. and J.P. Sprague. 1981. Absence of acclimation to parathion by rainbow trout. In: *Proceedings of the eighth annual aquatic toxicity workshop*. Kaushik, N.K. and K.R. Solomon (Eds.). Canadian Technical Report of Fisheries and Aquatic Sciences No. 1151. Department of Fisheries and Oceans, Ottawa, Ontario, Canada. p. 170.

Basak, P.K. and S.K. Konar. 1976a. Toxicity of six insecticides to fish. *Geobios* 3:209-210.

Basak, P.K. and S.K. Konar. 1976b. Pollution of water by pesticides and protection of fishes: Parathion. *Proc. Natl. Acad. Sci., India* 46(B):382-392.

Bellavere, C. and G. Gorbi. 1984. Biological variability and acute toxicity of parathion, dichlorobenil and TPBS to Biompharia glabrata and Brachydanio rerio. *Environ. Technol. Lett.* 5:389-396.

Benge, M.K. and L.E. Fronk. 1970. Effects of pesticides on non-target aquatic organisms on the Ogden Bay Waterfowl Management Area, Utah. *Proc. Annu. Meet. Utah Mosq. Abat. Assoc.* 23:34-40.

Benke, G.M., K.L. Cheever and S.D. Murphy. 1973. Comparative toxicity, anticholinesterase action and metabolism of methyl parathion, parathion and guthion in sunfish and mice. *Proc. 12th Annu. Meet. Soc. Toxicol.* pp. 473-474.

Benke, G.M., K.L. Cheever, F.E. Mirer and S.D. Murphy. 1974. Comparative toxicity, anticholinesterase action and metabolism of methyl parathion and parathion in sunfish and mice. *Toxicol. Appl. Pharmacol.* 28:97-109.

Bourquin, A.W., M.A. Hood and R.L. Garnas. 1977. An artificial microbial ecosystem for determining effects and fate of toxicants in a salt-marsh environment. *Dev. Ind. Microbiol.* 18:185-191.

Bourquin, A.W., R.L. Garnas, P.H. Pritchard, F.G. Wilkes, C.R. Cripe and N.I. Rubinstein. 1979. Interdependent microcosms for the assessment of

pollutants in the marine environment. Intern. J. Environ. Studies.

13:131-140.

Bowman, M.C., W.L. Oller, T. Cairns, A.B. Gosnell and K.H. Oliver. 1981.

Stressed bioassay systems for rapid screening of pesticide residues. Part

I: Evaluation of bioassay systems. Arch. Environ. Contam. Toxicol. 19:9-24.

Bradbury, S.J. 1973a. The effect of parathion on crustacean skeletal

muscle - I. The mechanical threshold and dependence on Ca^{2+} ions. Comp.

Biochem. Physiol. 44A:1021-1032.

Bradbury, S.J. 1973b. The effect of parathion on crustacean skeletal

muscle - II. Disruption of excitation-contraction coupling. Comp. Biochem.

Physiol. 44A:1033-1046.

Braun, H.E. and R. Frank. 1980. Organochlorine and organophosphorus

insecticides: Their use in eleven agricultural watersheds and their loss

to stream waters in southern Ontario, Canada, 1975-1977. Sci. Total.

Environ. 15:169-192.

Bringmann, G. and R. Kuhn. 1960. The water-toxicological detection of

insecticides. Gesund.-Ing. 8:243-244.

Bringmann, G. and R. Kuhn. 1977. Limiting values for the damaging action of

water pollutants to bacteria (Pseudomonas putida) and green algae

(Scenedesmus quadricauda) in the cell multiplication inhibition test. Z.

Wasser Abwasser Forsch. 10:87-98.

Bringmann, G. and R. Kuhn. 1978a. Limiting values for the noxious

effects of water pollutant material to blue algae (Microcystis aeruginosa)

and green algae (Scenedesmus quadricauda) in cell propagation inhibition tests.

Vom Wasser 50:45-60.

Bringmann, G. and R. Kuhn. 1978b. Testing of substances for their toxicity threshold: Model organisms Microcystis (Diplocystis) aeruginosa and Scenedesmus quadricauda. Mitt. Int. Ver. Theor. Angew. Limnol. 21:275-284.

Burchfield, H.P. and E.E. Storrs. 1954. Kinetics of insecticidal action based on the photomigration of larvae of Aedes aegypti. Contrib. Boyce Thompson Inst. 17:439-452.

Butler, P.A. 1963. Commercial fishery investigations. In: Pesticide-wildlife studies. A review of Fish and Wildlife Service investigations during 1961 and 1962. Fish and Wildlife Service Circular 167. U.S. Department of the Interior, Gulf Breeze, FL. pp. 11-25.

Butler, 1964. Commercial fishery investigations. In: Pesticide-wildlife studies, 1963: A review of Fish and Wildlife Service investigations during the calendar year. Fish and Wildlife Service Circular 199. U.S. Department of the Interior, Gulf Breeze, FL. pp. 5-28.

Butler, P.A. and R.L. Schutzmann. 1978. Residues of pesticides and PCBs in estuarine fish, 1972-76. National Pesticide Monitoring Program. Pesti. Monit. J. 12:51-59.

Carlson, G.P. 1973. Comparison of the metabolism of parathion by lobsters and rats. Bull. Environ. Contam. Toxicol. 9:296-300.

Chambers, J.E. 1976. The relationship of esterases to organophosphorus insecticide tolerance to mosquitofish. Pestic. Biochem. Physiol. 6:517-522.

Chambers J.E. and J.D. Yarbrough. 1974. Parathion and methyl parathion toxicity to insecticide-resistant and susceptible mosquitofish (Gambusia affinis). Bull. Environ. Contam. Toxicol. 11:315-320.

- Chang, V.C. and W.H. Lange. 1967. Laboratory and field evaluation of selected pesticides for control of the red crayfish in California rice fields. *J. Econ. Entomol.* 60:473-477.
- Chen, P.S., Y.N. Lin and C.L. Chung. 1971. Laboratory studies on the susceptibility of mosquito-eating fish, Lebistes reticulatus and the larvae of Culex pipiens fatigans to insecticides. *Tai Wan I-Hsueh-Hui Tsa-Chih* 70:28-35.
- Chovelon, A., L. George, C. Gulayets, Y. Hoyano, E. McGuinness, J. Moore, S. Ramamoorthy, P. Singer, K. Smiley and A. Wheatley. 1984. Pesticide and PCB levels in fish from Alberta. *Chemosphere* 13:19-32.
- Cole, D.R. and F.W. Plapp. 1974. Inhibition of growth and photosynthesis in Chlorella pyrenoidosa by a polychlorinated biphenyl and several insecticides. *Environ. Entomol.* 3:217-220.
- Chiou, C.T., V.H. Freed, D.W. Schmedding and R.L. Kohnert. 1977. Partition coefficient and bioaccumulation of selected organic chemicals. *Environ. Sci. Technol.* 11:475-478.
- Collins, W.J. and R. Shank. 1983. Surface water contamination by insecticides. Data from aquatic insect tests that pertain to water quality criteria. PB84-105378. National Technical Information Service, Springfield, VA.
- Coppage, D.L. 1972. Organophosphate pesticides: Specific level of brain AChE inhibition related to death in sheepshead minnow. *Trans. Am. Fish. Soc.* 101:534-536.

Coppage, D.L. and E. Matthews. 1974. Short-term effects of organophosphate pesticides on cholinesterases of estuarine fishes and pink shrimp. Bull. Environ. Contam. Toxicol. 11:483-487.

Cripe, G.M., D.R. Nimmo and T.L. Hamaker. 1981. Effects of two organophosphate pesticides on swimming stamina of the mysid, Mysidopsis bahia. In: Biological monitoring of marine pollutants. Vernberg, F.J., A. Calabrese, F.P. Thurnberg and W.B. Vernberg (Eds.). Academic Press, New York, NY. pp. 21-36.

Culley, D.D., Jr. and D.E. Ferguson. 1969. Patterns of insecticide resistance in the mosquitofish, Gambusia affinis. J. Fish. Res. Board Can. 26:2395-2401.

D'Asaro, C.N. 1982. Cycling of xenobiotics through marine and estuarine sediments. EPA-600/3-82-074. National Technical Information Service, Springfield, VA.

Davey, R.B. and M.V. Meisch. 1977. Dispersal of mosquitofish, Gambusia affinis, in Arkansas rice field. Mosq. News 37:777-778.

Davey, R.B., M.V. Meisch and F.L. Carter. 1976. Toxicity of five rice field pesticides to mosquitofish, Gambusia affinis, and green sunfish, Lepomis cyanellus, under laboratory and field conditions in Arkansas. Environ. Entomol. 5:1053-1056.

Davis, H.C. and H. Hidu. 1969. Effects of pesticides on embryonic development of clams and oysters and on survival and growth of the larvae. Fish. Bull. 67:393-404.

- Dick, M. 1982. Pesticide and PCB concentrations in Texas water, sediment and fish tissue. Report 264. Texas Department of Water Resources, Austin, TX.
- Dive, D., H. Leclerc and G. Perscone. 1980. Pesticide toxicity on the ciliate protozoan Colpidium campylum: Possible consequences of the effect of pesticides in the aquatic environment. *Ecotoxicol. Environ. Safety* 4:129-133.
- Dortland, R.J. 1978. Aliesterase-(ALI-E) activity in Daphnia magna as a parameter for exposure to parathion. *Hydrobiologia* 59:141-144.
- Dortland, R.J. 1980. Toxicological evaluation of parathion and azinphosmethyl in freshwater model ecosystems. *Versl. Landbouwk. Onderz.* 898:1-112.
- Dortland, R.J., H.R. Smitsaert, J. Hamoen and M. Scholte. 1976. In vitro and in vivo inhibition of ali-esterases and cholinesterases by paraoxon and parathion in Daphnia magna. *Meded. Fac. Landbouwet. Rijksuniv. Gent.* 41:1477-1482.
- Earnest, R. 1970. Progress in sport fishery research. Bureau of Sport Fisheries and Wildlife Resource Publication 106. Fish-Pesticide Research Laboratory, Columbia, MO. pp. 10-13.
- Eisler, R. 1969. Acute toxicity of insecticides to marine decapod crustaceans. *Crustaceana* 16:302-310.
- Estenik, J.F. and W.J. Collins. 1979. In vivo and in vitro studies of mixed-function oxidase in an aquatic insect. Chironomus riparius. *Am. Chem. Soc. Symp. Ser.* 99:349-370.
- Farr, J.A. 1977. Impairment of antipredator behavior in Palaemonetes pugio by exposure to sublethal doses of parathion. *Trans. Am. Fish. Soc.* 106:287-290.

Faust, S.D. 1975. Nonbiological degradation and transformations of organic pesticides in aqueous systems. In: Marine chemistry in the coastal environment. Church, T.M. (Ed.). ACS Symposium Series 18. American Chemical Society, Washington, DC. pp. 572-595.

Faust, S.D. and H.M. Goma. 1972. Chemical hydrolysis of some organic phosphorus and carbamate pesticides in aquatic environments. Environ. Lett. 3:171-201.

Federle, P.F. and W.J. Collins. 1976. Insecticide toxicity to three insects from Ohio ponds. Ohio J. Sci. 76:19-24.

Fleming, W.J. 1981. Recovery of cholinesterase activity in mallard ducklings administered organophosphorus pesticides. J. Toxicol. Environ. Health. 8:885-897.

Fleming, W.J., H. deChacin, O.H. Pattee and T.G. Lamont. 1982. Parathion accumulation in cricket frogs and its effect on American kestrels. J. Toxicol. Environ. Health 10:921-927.

Francis, B.M., L.G. Hansen, T.R. Fukuto, P. Lu and R.L. Metcalf. 1980. Ecotoxicology of phenylphosphonothioates. Environ. Health Perspect. 36:187-196.

Frear, D.E. and J.E. Boyd. 1967. Use of Daphnia magna for the microbioassay of pesticides. I. Development of standardized techniques for rearing Daphnia and preparation of dosage-mortality curves for pesticides. J. Econ. Entomol. 60:1228-1236.

Gahan, J.B. 1957. Further studies with water-soluble insecticides for the control of mosquito larvae in irrigation water. Mosq. News 17:198-201.

Garnas, R.L. and D.G. Crosby. 1979. Comparative metabolism of parathion by intertidal invertebrates. In: Marine pollution: Functional responses. Vernberg, W.B., F.P. Thurberg, A. Calabrese and F.J. Vernberg (Eds.). Academic Press, New York, NY. pp. 291-306.

Gasith, A. and A.S. Perry. 1980. Fate of parathion in a fish pond ecosystem and its impact on food chain organisms. Proc. Rep. Agrochem. Residue-Biota interaction in soil and aquatic ecosystems. International Atomic Energy Agency. pp. 125-151.

Gasith, A. and A.S. Perry. 1983. Effect of containment, presence of fish and repetitive exposure to parathion on the composition and dynamics of the natural fish pond zooplankton community. Isr. J. Zool. 32:165-166.

Gasith, A. and A.S. Perry. 1985. Use of limnocorrals for pesticide toxicity studies: Effect on zooplankton composition and dynamics. Verh. Int. Ver. Limnol. 22:2432-2436.

Gasith, A., A.S. Perry and A. Halevy. 1983a. The effect of repeated exposures to sublethal levels of parathion on the survival, growth, reproduction, and biomass production of three species of fish in a eutrophic fish pond ecosystem. In: Environmental Quality and Ecology: Proceedings of the 2nd International Conference on Ecology and Environmental Quality. Shoval, H.I. (Ed.). pp. 399-407.

Gasith, A., J.L. Albright, A.S. Perry and Y. Mozel. 1983b. Effects of parathion on the ecology of a eutrophic aquatic ecosystem: Limnocorral experiment. In: Agrochemical-biota interactions in soil and water using nuclear techniques. Technical Document 283. International Atomic Energy Agency. pp. 19-32.

Gaufin, A.R., L. Jensen and T. Nelson. 1961. Bioassays determine pesticide toxicity to aquatic invertebrates. *Water Sewage Works* 108:355-359.

Gaufin, A.R., L.D. Jensen, A.V. Nebeker, T. Nelson and R.W. Teel. 1965. The toxicity of ten organic insecticides to various aquatic invertebrates. *Water Sewage Works* 12:276-279.

Ghetti, P.F. and G. Gorbi. 1985. Effects of acute parathion pollution on macroinvertebrates in a stream. *Verh. Int. Ver. Limnol.* 22:2426-2431.

Gibson, J.R. 1971. Comparative biochemistry of parathion metabolism in three species of fishes. Ph.D. dissertation. Mississippi State University. Available from: University Microfilms, Ann Arbor, MI. Order No. 71-27,040.

Gibson, J.R. and J.L. Ludke. 1973. Effect of SKF-525A on brain acetylcholinesterase inhibition by parathion in fishes. *Bull. Environ. Contam. Toxicol.* 9:140-142.

Gillies, P.A., D.J. Womeldorf, E.P. Zboray and K.E. White. 1974. Insecticide susceptibility of mosquitoes in California: Status of organophosphorus resistance in larval Aedes nigromaculis and Culex tarsalis through 1973. *Proc. Pap. Annu. Conf. Calif. Mosq. Control Assoc.* 42:107-112.

Goldsmith, L.A. 1978. The toxicity of parathion to Orconectes rusticus and Viviparus malleatus. Ph.D. dissertation. University of Rhode Island, Kingston, RI. Available from: University Microfilms, Ann Arbor, MI. Order No. 7915457.

Goldsmith, L.A. and G.P. Carlson. 1979. Divergent toxicity of parathion in two freshwater invertebrates, Orconectes rusticus and Viviparus malleatus. *J. Environ. Sci. Health.* B14:579-588.

- Goldsmith, L.A., G.P. Carlson and G.C. Fuller. 1976. Toxicity and identification of parathion metabolites in freshwater crayfish (Cambarus) and snail (Viviparus). *Pharmacologist* 18:170.
- Gomaa, H.M. and S.D. Faust. 1971. Chemical hydrolysis and oxidation of parathion and paraoxon in aquatic environments. In: Fate of organic pesticides in the aquatic environment. Faust, S.D. (Ed.). American Chemical Society, Washington, DC. pp. 189-201.
- Graetz, D.A., G. Chesters, T.C. Daniel, L.W. Newland and G.B. Lee. 1970. Parathion degradation in lake sediments. *J. Water Pollut. Control Fed.* 42:R76-R94.
- Gregory, W.W., Jr., J.K. Reed and L.E. Priester, Jr. 1969. Accumulation of parathion and DDT by some algae and protozoa. *J. Protozool.* 16:69-71.
- Greve, P.A., J. Freudenthal and S.L. Wit. 1972. Potentially hazardous substances in surface waters. II. Cholinesterase inhibitors in Dutch surface waters. *Sci. Total. Environ.* 1:253-265.
- Grigarick, A.A. and M.O. Way. 1982. Role of crayfish (Decapoda: Astacidae) as pests of rice in California and their control. *Entomol. Soc. Am.* 75: 633-636.
- Grzenda, A.R., G.J. Lauer and H.P. Nicholson. 1962. Insecticide contamination in a farm pond. Part II. Biological effects. *Trans. Am. Fish. Soc.* 91:213-222.
- Gupta, S.P., S.R. Verma and R.C. Dalela. 1979. Studies on the toxicity of a few pesticides on Colisa fasciatus and Notopterus notopterus with

reference to histopathy and histochemistry. Part 1: Bioassay studies.

Acta Hydrochim. Hydrobiol. 78:223-238.

Gutierrez, M.C., E.P. Zboray and K.E. White. 1977. Insecticide susceptibility of mosquitoes in California: Status of organophosphorus resistance in larval Culex pipiens subsp., Proc. Pap. Annu. Conf. Calif. Mosq. Vector Control Assoc. 45:124-131.

Haddadin, M.J. and A.A. Alawi. 1974. Dead populations of fish in the rivers Jordan and Zarga. Biol. Conserv. 6:215-216.

Hall, R.J. and E. Kolbe. 1980. Bioconcentration of organophosphorus pesticides to hazardous levels by amphibians. J. Toxicol. Environ. Health 6:853-860.

Harris, C.R. and J.R.W. Miles. 1975. Pesticide residues in the Great Lakes region of Canada. Pest. Rev. 57:27-79.

Hart, N.R. and D.J. Womeldorf. 1977. Insecticide susceptibility of mosquitoes in Utah. Status of organophosphorus larvicide resistance through 1976. Proc. Annu. Meet. Utah Mosq. Abatement Assoc. 29:23-28.

Hashimoto, Y. and J. Fukami. 1969. Toxicity of orally and topically applied pesticide ingredients to carp, Cyprinus carpio. Botyukagaku 34:63-66.

Hashimoto, Y. and Y. Nishiuchi. 1981. Establishment of bioassay methods for the evaluation of acute toxicity of pesticides to aquatic organisms. J. Pest. Sci. 6:257-264.

Henderson, C. and Q.H. Pickering. 1958. Toxicity of organic phosphorus insecticides to fish. Trans. Am. Fish. Soc. 87:39-51.

- Henderson, C., Q.H. Pickering and C.M. Tarzell. 1959. The toxicity of organic phosphorus and chlorinated hydrocarbon insecticides to fish. Trans. 2nd Sem. Biol. Probl. Water Pollut., U.S. Public Health Service, Robert A. Taft Sanitary Engineering Center, Cincinnati, OH.
- Hesselberg, R.J. and J.L. Johnson. 1972. Column extraction of pesticides from fish, fish food and mud. Bull. Environ. Contam. Toxicol. 7:115-120.
- Hilsenhoff, W.L. 1959. The evaluation of insecticides for the control of Tendipes plumosus. J. Econ. Entomol. 52:331-332.
- Hiltibran, R.C. 1974. Oxygen and phosphate metabolism of bluegill liver mitochondria in the presence of some insecticides. Trans. Ill. State Acad. Sci. 67:228-237.
- Hiltibran, R.C. 1982. Effects of insecticides on the metal-activated hydrolysis of adenosine triphosphate by bluegill liver mitochondria. Arch. Environ. Contam. Toxicol. 11:709-717.
- Hitchcock, M. and S.D. Murphy. 1971. Activation of parathion and guthion by mammalian, avian, and piscine liver homogenates and cell fractions. Toxicol. Appl. Pharmacol. 19:37-45.
- Huddart, H. 1978. Parathion- and DDT-induced effects on tension and calcium transport in molluscan visceral muscle. Comp. Biochem. Physiol. 61C:1-6.
- Hudson, R.H., M.A. Haegele and R.K. Tucker. 1979. Acute oral and percutaneous toxicity of pesticides to mallards: Correlations with mammalian toxicity data. Toxicol. Appl. Pharmacol. 47:451-460.

- Hughes, J.S. 1970. Tolerance of striped bass, Morone saxatilis, larvae and fingerlings to nine chemicals used in pond culture. Proc. Annu. Conf. Southeast. Assoc. Game Fish Comm. 24:431-438.
- Hughes, J.S. 1973. Acute toxicity of thirty chemicals to striped bass (Morone saxatilis). Louisiana Department of Wildlife and Fisheries, Fish and Game Division, Monroe, LA.
- Jensen, L.D. and A.R. Gaufin. 1964. Effects of ten organic insecticides on two species of stonefly naiads. Trans. Am. Fish. Soc. 93:27-34.
- Johnson, W.W. and M.T. Finley. 1980. Handbook of acute toxicity of chemicals to fish and aquatic invertebrates. Resource Publication 137. U.S. Fish and Wildlife Service, Washington, DC.
- Juhnke, V.I. and D. Ludemann. 1978. Results of the investigation of zoo chemical compounds for acute fish toxicity with the golden orfe test. Wass. Abwass. Forsch. 11:161-164.
- Jung, K.D. 1973. Substances extremely toxic to fish and their importance for a fish test warning system. Gas-Wasserfach, Wasser-Abwasser. 114:232-233.
- Kannan, V. and S.V. Job. 1979. Studies on the residual levels of pesticide pollution in the Sathiar reservoir. J. Radioanal. Chem. 53:247-253.
- Khattat, F.H. and S. Farley. 1976. Acute toxicity of certain pesticides to Acartia tonsa Dana. EPA-600/3-76-033. National Technical Information Service, Springfield, VA.
- King, K.A., D.H. White and C.A. Mitchell. 1984. Nest defense behavior and reproductive success of laughing gulls sublethally dosed with parathion. Bull. Environ. Contam. Toxicol. 33:499-504.

Kleerekoper, H. 1974. Effects of exposure to a sublethal concentration of parathion on the interaction between chemoreception and water flow in fish. In: Pollution and physiology of marine organisms. Vernberg, F.J. and W.B. Vernberg (Eds.). Academic Press, New York, NY. pp. 237-245.

Konar, S.K. and P.K. Basak. 1973. Some effects of mechanical and chemical injuries on catfish. Proc. Indian Sci. Cong. Assoc. 60:551-552.

Korn, S. and R. Earnest. 1974. Acute toxicity of twenty insecticides to striped bass, Morone saxatilis. Calif. Fish Game. 60:128-131.

Kortus, J., P. Macuch, J. Mayer, K. Durcek and V. Krcmery. 1971. Uptake of 32-P parathion and 32-P imidam by Euglena gracilis. J. Hyg. Epidemiol. Microbiol. Immunol. (Prague). 15:101-103.

Krieger, R.I. and P.W. Lee. 1973. Inhibition of in vivo and in vitro epoxidation of aldrin and potentiation of toxicity of various insecticide chemicals by diquat in two species of fish. Arch. Environ. Contam. Toxicol. 1:112-121.

Kynard, B. 1974. Avoidance behavior of insecticide susceptible and resistant populations of mosquitofish to four insecticides. Trans. Am. Fish. Soc. 103:557-561.

Labrecque, G.C., J.R. Noe and J.B. Gahan. 1956. Effectiveness of insecticides on granular clay carriers against mosquito larvae. Mosq. News 16:1-3.

- Lahav, M. and S. Sarig. 1969. Sensitivity of pond fish to cotnion (azinphosmethyl) and parathion. *Bamidgeh* 21:67-74.
- Laplanche, A., M. Bouvet, F. Venien, G. Martin and A. Chabrolles. 1981. Modeling parathion changes in the natural environment laboratory experiment. *Water Res.* 15:599-607.
- LeBlanc, G.A. 1984. Interspecies relationships in acute toxicity of chemicals to aquatic organisms. *Environ. Toxicol. Chem.* 3:47-60.
- Lewallen, L.L. 1959. Toxicity of several organophosphorus insecticides to Gambusia affinis in laboratory tests. *Mosq. News* 19:1-2.
- Lewallen, L.L. 1962. Toxicity of certain insecticides to hydrophilid larvae. *Mosq. News* 22:112-113.
- Lewallen, L.L. and W.H. Wilder. 1962. Toxicity of certain organophosphorus and carbamate insecticides to rainbow trout. *Mosq. News* 22:369-372.
- Lewis, D.L., R.E. Hodson and L.F. Freeman III. 1984. Effects of microbial community interactions on transformation rates of xenobiotic chemicals. *Appl. Environ. Microbiol.* 48:561-565.
- Lichtenstein, E.P., T.T. Liang and B.N. Anderegg. 1975. Synergism of insecticides by herbicides under various environmental conditions. *Environ. Conserv.* 2:148
- Lichtenstein, E.P., K.R. Schulz, R.F. Skrentny and Y. Tsukano. 1966. Toxicity and fate of insecticide residues in water. *Arch. Environ. Health* 12:199-212.

Loeb, H.A. and W.H. Kelly. 1963. Acute oral toxicity of 1,496 chemicals force-fed to carp. U.S. Fisheries and Wildlife Service Special Report No. 471. Washington, DC.

Lowe, J.I., P.D. Wilson and R.B. Davison. 1970. Laboratory bioassays. In: Progress report for fiscal year 1969. U.S. Fish and Wildlife Circular 335. Pesticide Field Station, Gulf Breeze, FL.

Lowe, J.I., P.D. Wilson, A.J. Rick and A.J. Wilson, Jr. 1971. Chronic exposure of oysters to DDT, toxaphene and parathion. Proc. Nat. Shellfish. Assoc. 61:71-79.

Ludemann, D. and F. Herzel. 1973. Toxicity and concentration changes with time of the insecticide parathion in water. Z. Angew. Zool. 60:177-181.

Ludemann, D. and H. Neumann. 1960a. Studies on the acute toxicological action of modern contact insecticides to carp (Cyprinus carpio). Z. Angew. Zool. 47:11-33.

Ludemann, D. and H. Neumann. 1960b. Studies on the acute toxicity of modern contact insecticides in fresh water. Z. Angew. Zool. 47:303-321.

Ludemann, D. and H. Neumann. 1960c. Studies on the acute toxicity of modern contact insecticides in fresh water. III. Chironomid larvae. Z. Angew. Zool. 47:493-505.

Ludemann, D. and H. Neumann. 1961. Study of the acute toxicity of modern contact insecticides for freshwater animals. Z. Angew. Zool. 48:87-96.

Ludke, J.L., J.R. Gibson and C.I. Lusk. 1972. Mixed function oxidase activity in freshwater fishes. Aldrin epoxidation and parathion activation. Toxicol. Appl. Pharmacol. 21:89-97.

- Macek, K.J. 1975. Acute toxicity of pesticide mixtures to bluegills. Bull. Environ. Contam. Toxicol. 14:648-652.
- Mackiewicz, M., K.H. Deubert, H.B. Gunner and B.M. Zuckerman. 1969. Study of parathion biodegradation using gnotobiotic techniques. J. Agric. Food Chem. 17:129-130.
- McDonald, T.O. and M. Fingerman. 1979. Effects of DDT and parathion on goldfish retina/pigment epithelium/choroid concentrations of serotonin, dopamine, and norepinephrine. Toxicol. Appl. Pharmacol. 48:A199.
- Micks, D.W. and D. Rougeau. 1977. Organophosphorus tolerance in Culex quinquefasciatus in Texas. Mosq. News 37:233-239.
- Miller, C.W., B.M. Zuckerman and A.J. Charig. 1966. Water translocation of diazinon-14C and parathion-35S off a model cranberry bog and subsequent occurrence in fish and mussels. Trans. Am. Fish. Soc. 95:345-349.
- Miller, C.W., W.E. Tomlinson and R.L. Norgen. 1967. Persistence and movement of parathion in irrigation waters. Pestic. Monit. J. 1:47-48.
- Minchew, C.D. and D.E. Ferguson. 1970. Toxicities of six insecticides to resistant and susceptible green sunfish and golden shiners in static bioassays. Miss. Acad. Sci. J. 15:29-32.
- Mohamed, M.P. and R.A. Gupta. 1984. Effects of sublethal concentration of ethyl parathion on oxygen consumption and random swimming activity of Cirrhinus mrigala. Indian J. Exp. Biol. 22:42-44.
- Moore, R.B. 1970. Effects of pesticides on growth and survival of Euglena gracilis Z. Bull. Environ. Contam. Toxicol. 5:226-230.

- Morgan, W.S. 1975. Monitoring pesticides by means of changes in electric potential caused by fish opercular rhythms. Prog. Water Technol. 7:33-40.
- Morgan, W.S. 1976. Fishing for toxicity: Biological automonitor for continuous water quality control. Effluent Water Treat. J. 16:471-472, 474-475.
- Morgan, W.S. 1977. Biomonitoring with fish: An aid to industrial effluent and surface water quality control. Prog. Water Technol. 9:703-711.
- Mount, D.I. and H.W. Boyle. 1969. Parathion-use of blood concentration to diagnose mortality of fish. Environ. Sci. Technol. 3:1183-1185.
- Mulla, M.S. 1963. Persistence of mosquito larvicides in water. Mosq. News 23:234-237.
- Mulla, M.S. 1980. New synthetic pyrethroids - effective mosquito larvicides. Proc. Pap. Annu. Conf. Calif. Mosq. Vector Control Assoc. 48:92-93.
- Mulla, M.S. and L.W. Isaak. 1961. Field studies on the toxicity of insecticides to the mosquitofish, Gambusia affinis. J. Econ. Entomol. 54:1237-1242.
- Mulla, M.S., R.L. Metcalf and L.W. Isaak. 1962. Some new and highly effective mosquito larvicides. Mosq. News 22:231-238.
- Mulla, M.S., L.W. Isaak and H. Axelrod. 1963. Field studies on the effects of insecticides on some aquatic wildlife species. J. Econ. Entomol. 56:184-188.
- Mulla, M.S., R.L. Metcalf and G. Kats. 1964. Evaluation of new mosquito larvicides with notes on resistant strains. Mosq. News 24:312-319.

- Mulla, M.S., J.O. Keith and F.A. Gunther. 1966. Persistence and biological effects of parathion residues in waterfowl habitats. J. Econ. Entomol. 59:1085-1090.
- Mulla, M.S., J. St. Amant and L.D. Anderson. 1967. Evaluation of organic pesticides for possible use as fish toxicants. Prog. Fish-Cult. 29:36-42.
- Mulla, M.S., H.A. Darwazeh and P.A. Gillies. 1970. Evaluation of aliphatic amines against larvae and pupae of mosquitoes. J. Econ. Entomol. 63:1472-1475.
- Mulla, M.S., H. Navvab-Gojrati and H.A. Darwazeh. 1978. Biological activity and longevity of new synthetic pyrethroids against mosquitoes and some non-target insects. Mosq. News 38:90-96.
- Murphy, S.D., R.R. Lauwerys and K.L. Cheever. 1968. Comparative anticholinesterase action of organophosphorus insecticides in vertebrates. Toxicol. Appl. Pharmacol. 12:22-35.
- Myers, C.M., P.A. Gillies and R.F. Frohli. 1969. Field test of Abate and parathion granules in catch basins. Proc. Pap. Annu. Conf. Calif. Mosq. Control Assoc. 37:61-63.
- Nagasawa, K., E. Koshimura and H. Fukuda. 1968. LD50 and ED50 of parathion and potassium cyanide and their bioassay using guppies (Lebistes reticulatus). Bull. Natl. Inst. Hyg. Sci. (Tokyo) 86:32-36.
- Naqvi, S.M. 1973. Toxicity of twenty-three insecticides to a tubificid worm Branchiura sowerbyi from the Mississippi delta. J. Econ. Entomol. 66:70-74.
- Naqvi, S.M. and D.E. Ferguson. 1970. Levels of insecticide resistance in freshwater shrimp, Palaemonetes kadiakensis. Trans. Am. Fish. Soc. 99:696-699.

- Nicholson, H.P., H.J. Webb, G.J. Lauer, R.E. O'Brien, A.R. Grzenda and D.W. Shanklin. 1962. Insecticide contamination in a farm pond. Part I. Origin and duration. *Trans. Am. Fish. Soc.* 91:213-222.
- Nimmo, D.R., T.L. Hamaker, E. Matthews and J.C. Moore. 1981. An overview of the acute and chronic effects of first and second generation pesticides on an estuarine mysid. In: *Biological monitoring of marine pollutants*. Vernberg, F.J., A. Calabrese, F.P. Thurberg and W.B. Vernberg (Eds.). Academic Press, New York, NY. pp. 3-19.
- Nishiuchi, Y. and Y. Hashimoto. 1967. Toxicity of pesticide ingredients to some freshwater organisms. *Botyu-Kagaku (Sci. Pest. Control)* 32:5-11.
- Nishiuchi, Y. and Y. Hashimoto. 1969. Toxicity of pesticides to some freshwater organisms. *Rev. Plant Prot. Res.* 2:137-139.
- Nishiuchi, Y. and K. Yoshida. 1972. Toxicities of pesticides to some freshwater snails. *Bull. Agr. Chem. Insp. Stn.* 12:86-92.
- Nollenberger, E.L. 1982. Toxicant-induced changes in brain, gill, liver and kidney of brook trout exposed to carbaryl, atrazine, 2,4-D, and parathion: A cytochemical study. *Diss. Abstr. Int. B.* 42:3922.
- Nollenberger, E.L., W. Neff and A. Anthony. 1981. Cytochemical analysis of brain nucleic acid changes in brook trout (Salvelinus fontinalis) exposed to carbaryl and parathion toxication. *Proc. Pa. Acad. Sci.* 55:97.
- Panwar, R.S., D. Kapoor, H.C. Joshi and R.A. Gupta. 1976. Toxicity of some insecticides to the weed fish, Trichogaster fasciatus. *J. Inl. Fish. Soc. India* 8:129-130.

Panwar, R.S., R.A. Gupta, H.C. Joshi and D. Kapoor. 1982. Toxicity of some chlorinated hydrocarbon and organophosphorus insecticides to gastropod, Viviparus bengalensis. J. Environ. Biol. 3:31-36.

Perry, A.S., A. Gasith and Y. Mozel. 1983. Pesticide residues in fish and aquatic invertebrates. Arch. Toxicol. Suppl. 6:199-204.

Pickering, Q.H., C. Henderson and A.E. Lemke. 1962. The toxicity of organic phosphorus insecticides to different species of warmwater fishes. Trans. Am. Fish. Soc. 91:175-184.

Poorman, A.E. 1973. Effects of pesticides on Euglena gracilis. I. Growth studies. Bull. Environ. Contam. Toxicol. 10:25-28.

Price, N.R. 1976. The effects of two insecticides on the $Ca^{+2} + Mg^{+2}$ activated ATPase of the sarcoplasmic reticulum of the flounder, Platichthys flesus. Comp. Biochem. Physiol. 55C:91-94.

Price, N.R. 1978. Disruption of excitation-contraction coupling by organic insecticides. Mode of action in the muscle of the flounder, Platichthys flesus. Comp. Biochem. Physiol. 59C:127-133.

Ramke, D. 1969. Development of organophosphorus resistant Aedes nigromaculis in the Tulane Mosquito Abatement District. Proc. Pap. Annu. Conf. Calif. Mosq. Control Assoc. 37:63.

Rand, G.M. 1977a. The effect of subacute parathion exposure on the locomotor behavior of the bluegill sunfish and largemouth bass. In: Aquatic toxicology and hazard evaluation. Mayer, F.L. and J.L. Hamelink (Eds.). ASTM STP 634. American Society for Testing and Materials, Philadelphia, PA. pp. 253-268.

Rand, G.M. 1977b. The effect of exposure to a subacute concentration of parathion on the general locomotor behavior of the goldfish. Bull. Environ. Contam. Toxicol. 18:259-266.

Rand, G., H. Kleerekoper and J. Matis. 1975. Interaction of odor and flow perception and the effects of parathion in the locomotor orientation of the goldfish Carassius auratus. J. Fish Biol. 7:497-504.

Rattner, B.A. 1982. Diagnosis of anticholinesterase poisoning in birds: Effects of environmental temperature and underfeeding on cholinesterase activity. Environ. Toxicol. Chem. 1:329-335.

Sanders, H.O. 1969. Toxicity of pesticides to the crustacean Gammarus lacustris. Technical Paper No. 25. U.S. Fish and Wildlife Service, Columbia, MO.

Sanders, H.O. 1970. Pesticide toxicities to tadpoles of the western chorus frog, Pseudis triseriata, and Fowler's toad, Bufo woodhousii fowleri. Copeia 2:246-251.

Sanders, H.O. 1972. Toxicity of some insecticides to four species of malacostracan crustaceans. Technical Paper No. 66. Bureau of Sport Fisheries and Wildlife, U.S. Fish and Wildlife Service, Washington, DC.

Sanders, H.O. and O.B. Cope. 1968. The relative toxicities of several pesticides to naiads of three species of stoneflies. Limnol. Oceanogr. 13:112-117.

Sato, R. and H. Kubo. 1965. The water pollution caused by organophosphorus insecticides in Japan. In: Advances in water pollution research. Jaag, O. (Ed.), Pergamon Press, London. pp. 95-99.

- Schmidt, C.H. and D.E. Weidaas. 1961. The toxicological action of three organophosphorus insecticides with three species of mosquito larvae. J. Econ. Entomol. 54:583-586.
- Sethunathan, N., R. Siddaramappa, K.P. Rajaram, S. Barik and P.A. Wahid. 1977. Parathion: Residues in soil and water. Residue Rev. 68:92-122.
- Shah, P.V., R.J. Monroe and F.E. Guthrie. 1983. Comparative penetration of insecticides in target and non-target species. Drug Chem. Toxicol. 6:155-179.
- Singh, H. and T.P. Singh. 1981. Effect of parathion and aldrin on survival, ovarian 32-P uptake and gonadotrophic potency in a freshwater catfish, Heteropneustes fossilis. Endokrinologie 77:173-178.
- Siva Prasada, K., K.R. Sambasiva and K.V. Ramana. 1983. Effect of parathion on tissue ionic changes in fish, Channa punctatus. Geobios 10:60-62.
- Solomon, H.M. 1977. The teratogenic effects of the insecticides DDT, carbaryl, malathion, and parathion on developing medaka eggs (Oryzias latipes). Diss. Abstr. Int. B 39:2176-2177.
- Solomon, H.M. and J.S. Weis. 1979. Abnormal circulatory development in medaka caused by the insecticides carbaryl, malathion and parathion. Teratology 19:51-61.
- Solon, J.M. and J.H. Nair, III. 1970. The effect of a sublethal concentration of LAS on the acute toxicity of various phosphate pesticides to the fathead minnow (Pimephales promelas Rafinesque). Bull. Environ. Contam. Toxicol. 5:408-413.

- Solon, J.M., J.L. Lincer and J.H. Nair. 1969. The effect of sublethal concentrations of LAS on the acute toxicity of various insecticides to the fathead minnow (Pimephales promelas). Water Res. 3:767-775.
- Spacie, A. 1976. Acute and chronic parathion toxicity to fish and invertebrates. PB257800. National Technical Information Service, Springfield, VA.
- Spacie, A., A.G. Vilkas, G.F. Doebbler, W.J. Kuc and G.R. Iwan. 1981. Acute and chronic parathion toxicity to fish and invertebrates. PB81-245862 or EPA-600/3-81-047. National Technical Information Service, Springfield, VA.
- Sreenivasan, A. and G.K. Swaminathan. 1967. Toxicity of six organophosphorus insecticides to fish. Curr. Sci. 36:397-398.
- Srivastava, G.N., R.A. Gupta, M.P. Mohamed and D. Nath. 1977. Effect of sub-lethal ethyl parathion on the metabolism and activity of Colisa fasciata. Indian J. Environ. Health 19:63-66.
- Stephan, C.E., D.I. Mount, D.J. Hansen, J.H. Gentile, G.A. Chapman and W.A. Brungs. 1985. Guidelines for deriving numerical national water quality criteria for the protection of aquatic organisms and their uses. PB85-227049. National Technical Information Service, Springfield, VA.
- Stewart, J.P. 1977. Synergism of chlorpyrifos by DEF in the control of organophosphorus resistant Culex pipiens quinquefasciatus larvae, with notes on synergism of parathion and fenthion. Proc. Pap. Annu. Conf. Calif. Mosq. Vector Control Assoc. 45:132-133.

Sun, T.J. and D.H. Taylor. 1983. The effects of parathion on acquisition and retention of shuttlebox avoidance - conditioning in the goldfish Carassius auratus. Environ. Pollut. (Series A) 31:119-131.

Surber, E.W. 1948. Chemical control agents and their effects on fish. Prog. Fish-Cult. 10:125-131.

Tarpley, W.A. 1958. Studies on the use of the brine shrimp, Artemia salina, as a test organism for bioassay. J. Econ. Entomol. 51:780-783.

Tarzwel, C.M. 1959a. Some effects of mosquito larviciding and the new pesticides on fishes. Proceedings of the Symposium on Coordination of Mosquito Control and Wildlife Management, Washington, DC. April 1-2, 1959.

Tarzwel, C.M. 1959b. Pollutional effects of organic insecticides. Transactions of the 24th North American Wildlife Conference. Wildlife Management Institute, Washington, DC. pp. 132-142.

Tomita, H. and N. Matsuda. 1961. Deformity of vertebrae induced by lathyrogenic agents and phenyl-thiourea in medaka (Oryzias latipes). Embryologia 5:413-422.

U.S. Bureau of Commercial Fisheries. 1966. Unpublished laboratory data. Gulf Breeze, FL.

U.S. Bureau of Commercial Fisheries. 1967. Unpublished laboratory data. Gulf Breeze, FL.

U.S. EPA. 1975. Substitute chemical program: Initial scientific and minieconomic review of parathion. EPA-540/1-75-001. National Technical Information Service, Springfield, VA.

- U.S. EPA. 1976. Quality criteria for water. EPA-440/9-76-023. National Technical Information Service, Springfield, VA.
- U.S. EPA. 1983a. Water quality standards regulation. Fed. Regist. 48:51400-51413. November 8.
- U.S. EPA. 1983b. Water quality standards handbook. Office of Water Regulations and Standards, Washington, DC.
- U.S. EPA. 1985. Technical support document for water-quality based toxic controls. Office of Water, Washington, DC. September.
- Van Hoof, F. 1980. Evaluation of an automatic system for detection of toxic substances in surface water using trout. Bull. Environ. Contam. Toxicol. 25:221-225.
- Van Leeuwen, C.J., P.S. Griffioen, W.H. Vergouw and J.L. Maas-Diepeveen. 1985. Differences in susceptibility of early life stages of rainbow trout (Salmo gairdneri) to environmental pollutants. Aquat. Toxicol. 7:59-78.
- Van Middelen. C.H. 1966. Fate and persistence of organic pesticides in the environment. In: Organic pesticides in the environment. Rosen, A.A. and H.F. Kraybill (Eds.). American Chemical Society, Washington, DC. pp. 228-242.
- Verma, S.R. and S.P. Gupta. 1976. Pesticides in relation to water pollution. Accumulation of aldrin and ethyl parathion in the tissues of Colisa fasciatus and Notopterus notopterus. Indian J. Environ. Health 18:10-14.
- Verma, S.R., V. Kumar and R.C. Dalela. 1981. Studies on the accumulation and elimination of three pesticides in the gonads of Notopterus notopterus and Colisa fasciatus. Indian J. Environ. Health 23:275-281.

von Rumker, R., E.W. Lawless, A.F. Meiners, K.A. Lawrence, G.L. Kelso and F. Horay. 1974. Production, distribution, use and environmental impact potential of selected pesticides. EPA-540/1-74-001. National Technical Information Service, Springfield, VA. pp. 181-195.

Walsh, G.E. and S.V. Alexander. 1980. A marine algal bioassay method: Results with pesticides and industrial wastes. Water Air Soil Pollut. 13:45-55.

Warnick, S.L., R.F. Gaufin and A.R. Gaufin. 1966. Concentrations and effects of pesticides in aquatic environments. J. Am. Water Works Assoc. 58:601-608.

Weis, J.S. and L.H. Mantel. 1976. DDT as an accelerator of limb regeneration and molting in fiddler crabs. Estuarine Coastal Mar. Sci. 4:461-466.

Weis, J.S. and P. Weis. 1975. Retardation of fin regeneration in Fundulus by several insecticides. Trans. Am. Fish. Soc. 104:135-137.

Weis, P. and J.S. Weis. 1974. Cardiac malformations and other effects due to insecticides in embryos of the killifish, Fundulus heteroclitus. Teratology 10:263-268.

Weiss, C.M. 1959. Stream pollution: Response of fish to sub-lethal exposures of organic phosphorus insecticides. Sewage Ind. Wastes 31:580-593.

Weiss, C.M. 1961. Physiological effect of organic phosphorus insecticides on several species of fish. Trans. Am. Fish. Soc. 90:143-152.

Weiss, C.M. and J.H. Gakstatter. 1964. Detection of pesticides in water by biochemical assay. J. Water Pollut. Control Fed. 36:240-253.

- Weiss, C.M. and J.H. Gakstatter. 1965. The decay of anticholinesterase activity of organic phosphorus insecticides on storage in waters of different pH. In: Advances in water pollution research. Jaag, O. (Ed.), Pergamon Press, London. pp. 83-95.
- White, D.H., K.A. King, C.A. Mitchell, E.F. Hill and T.G. Lamont. 1979. Parathion causes secondary poisoning in a laughing gull breeding colony. Bull. Environ. Contam. Toxicol. 23:281-284.
- Whitmore, D.H. and D.H. Hodges. 1978. In vitro pesticide inhibition of muscle esterases of the mosquitofish, Gambusia affinis. Comp. Biochem. Physiol. 59C:145-149.
- Whitten, B.K. and C.J. Goodnight. 1966. Toxicity of some common insecticides to tubificids. J. Water Pollut. Control Fed. 38:227-235.
- Wilder, W.H. and C.S. Schaefer. 1969. Organophosphorus resistance levels in adults and larvae of the pasture mosquito, Aedes nigromaculis, in the San Joaquin Valley of California. Proc. Pap. Annu. Conf. Calif. Mosq. Control Assoc. 37:64-67.
- Womeldorf, D.J., R.K. Washino, K.E. White and P.A. Gieke. 1970. Insecticide susceptibility of mosquitoes in California: Response of Anopheles freeborni Aitken larvae to organophosphorus compounds. Mosq. News 30:374-382.
- Yahalomi, Z. and A.S. Perry. 1981. Microsomal mixed-function oxidases in fish in relation to environmental contamination by organochlorine insecticides. Comp. Biochem. Physiol. 70C:97-102.

Yasuno, M., S. Hirakoso, M. Sasa and M. Uchida. 1965. Inactivation of some organophosphorus insecticides by bacteria in polluted water. Japan. J. Exp. Med. 35:545-563.

Yu, C.C. and J.R. Sanborn. 1975. The fate of parathion in a model ecosystem. Bull. Environ. Contam. Toxicol. 13:543-550.

Zboray, E.P. and M.C. Gutierrez. 1979. Insecticide susceptibility of mosquitoes in California: Status of organophosphorus resistance in larval Culex tarsalis through 1978, with notes on mitigating the problem. Proc. Pap. Annu. Conf. Calif. Mosq. Vector Control Assoc. 47:26-28.

Zuckerman, B.M., K. Deubert, M. Mackiewicz and H. Gunner. 1970. Studies on the biodegradation of parathion. Plant Soil 33:273-281.