

PB85-173433


Chemical Transformations in Acid Rain
Volume 2. Investigation of Kinetics and
Mechanism of Aqueous-Phase Peroxide Formation

Brookhaven National Lab., Upton, NY

Prepared for

Environmental Protection Agency
Research Triangle Park, NC

Mar 85



EPA/600/3-85/017
March 1985

CHEMICAL TRANSFORMATIONS IN ACID RAIN

Volume II. Investigation of Kinetics and Mechanism
of Aqueous-Phase Peroxide Formation

Yin-Nan Lee

Environmental Chemistry Division
Department of Applied Science
Brookhaven National Laboratory
Upton, New York 11973

Interagency Agreement DW 930256

Project Officer

Marcia C. Dodge

Atmospheric Chemistry and Physics Division
Atmospheric Sciences Research Laboratory
Research Triangle Park, NC 27711

ATMOSPHERIC SCIENCES RESEARCH LABORATORY
OFFICE OF RESEARCH AND DEVELOPMENT
U.S. ENVIRONMENTAL PROTECTION AGENCY
RESEARCH TRIANGLE PARK, NC 27711

TECHNICAL REPORT DATA (Please read Instructions on the reverse before completing)		
1. REPORT NO. EPA/600/3-85/017	2.	3. RECIPIENT'S ACCESSION NO. PDB 5 173433 /AS
4. TITLE AND SUBTITLE CHEMICAL TRANSFORMATIONS IN ACID RAIN Volume II. Investigation of Kinetics and Mechanism of Aqueous-Phase Peroxide Formation		5. REPORT DATE March 1985
		6. PERFORMING ORGANIZATION CODE
7. AUTHOR(S) Yin-Nan Lee		8. PERFORMING ORGANIZATION REPORT NO.
9. PERFORMING ORGANIZATION NAME AND ADDRESS Environmental Chemistry Division Brookhaven National Laboratory Upton, NY 11973		10. PROGRAM ELEMENT NO. CCVNIA/02-3167 (FY-85)
		11. CONTRACT/GRANT NO. Interagency Agreement DW930256
12. SPONSORING AGENCY NAME AND ADDRESS Atmospheric Sciences Research Laboratory-RTP, NC Office of Research and Development U.S. Environmental Protection Agency Research Triangle Park, North Carolina 27711		13. TYPE OF REPORT AND PERIOD COVERED Final (8/83-7/84)
		14. SPONSORING AGENCY CODE EPA/600/09
15. SUPPLEMENTARY NOTES		
16. ABSTRACT The aqueous-phase reactions of O ₃ with a number of species have been studied in an effort to identify pathways leading to the production of hydrogen peroxide in solution. The aqueous-phase systems studied included the decomposition of O ₃ in pure water and the interaction of O ₃ with (1) NO ₂ , (2) PAN, (3) ethylene, (4) formic acid, (5) formaldehyde and (6) formaldehyde in the presence of NO ₂ . Except for the O ₃ -ethylene reaction, peroxide was not found as a reaction product. From the results obtained, it is concluded that the reactions studied in this research effort are not significant with respect to atmospheric peroxide formation.		
17. KEY WORDS AND DOCUMENT ANALYSIS		
a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
18. DISTRIBUTION STATEMENT RELEASE TO PUBLIC	19. SECURITY CLASS (This Report) UNCLASSIFIED	21. NO. OF PAGES 60
	20. SECURITY CLASS (This page) UNCLASSIFIED	22. PRICE

NOTICE

The information in this document has been funded by the United States Environmental Protection Agency under Interagency Agreement DW 930256 to Brookhaven National Laboratory. It has been subject to the Agency's peer and administrative review, and it has been approved for publication as an EPA document. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

ABSTRACT

The aqueous-phase reaction kinetics of dissolved O_3 with a number of atmospheric components have been investigated, with special attention focused on the formation of H_2O_2 or organic peroxide as reaction products. The reagent concentrations employed, rate laws, rate constants, and peroxide yields (y) determined for the specified substrates are:

$$H_2O: -d[C_3]/dt = k[O_3], k = 2.1 \times 10^{-4} s^{-1} (pH \sim 6), y \leq 0.5\%;$$

$$H_2O_2 (\sim 5 \times 10^{-5} M): -d[O_3]/dt = -d[H_2O_2]/dt = k[O_3][H_2O_2],$$

$$k = 2.6 \times 10^3 M^{-1} s^{-1} (pH \sim 6);$$

$$HCO_2H (\sim 1 \times 10^{-5} M): -d[O_3]/dt = k[O_3][HCO_2^-],$$

$$k = 4.3 \times 10^3 M^{-1} s^{-1}, y < 0.5\%;$$

$$H_2CO (\sim 1 \times 10^{-5} M): -d[O_3]/dt = k[O_3]^{1/2} [H_2CO]^{1/2},$$

$$k = 1.2 \times 10^{-3} s^{-1}, y < 2\%;$$

$$C_2H_4 (\sim 15 \text{ ppm}): d[\text{peroxide}]/dt = k[O_3][C_2H_4],$$

$$k = 3.0 \times 10^5 M^{-1} s^{-1};$$

$$PAN (\sim 100 \text{ ppb}): d[\text{peroxide}]/dt = k[PAN][O_3], k \leq 3 \times 10^3 M^{-1} s^{-1};$$

$$NO_2 (40 \text{ ppb}): d[\text{peroxide}]/dt = k {}^HNO_3 PNO_3,$$

$$k {}^HNO_3 \leq 4 \times 10^{-3} M \text{ atm}^{-1} s^{-1};$$

$$NO_2 + HCO_2 (1 \times 10^{-4} M): d[\text{peroxide}]/dt = k {}^HNO_3 [H_2CO] PNO_3,$$

$$k {}^HNO_3 \leq 6 \text{ atm}^{-1} s^{-1}.$$

With the use of these data, the rates of the aqueous-phase peroxide production of these reactions under typical atmospheric conditions are calculated to be $\sim 1 \times 10^{-6} M \text{ hr}^{-1}$ or smaller. It is therefore concluded that the reactions studied in this work contribute insignificantly to the formation of peroxides in atmospheric water.

CONTENTS

ABSTRACT.....	iii
TABLES.....	vi
FIGURES.....	vii
ACKNOWLEDGMENT.....	viii
INTRODUCTION.....	1
EXPERIMENTAL.....	4
Material.....	4
Ozone Generation.....	4
Gas-phase O_3 Concentration.....	5
Preparation of O_3 Solution.....	5
Aqueous-phase O_3 Concentration.....	5
H_2O_2 and Peroxide Concentrations.....	7
Kinetic Measurements.....	7
RESULTS.....	12
O_3 Decomposition in Optical Cell.....	12
O_3 - H_2O_2 Reaction.....	12
O_3 - HCO_2H Reaction.....	15
O_2 - H_2CO Reaction.....	19
O_3 - C_2H_4 Reaction.....	22
Aqueous-phase Reactions of PAN.....	27
O_3 - NO_2 Reaction.....	29
O_3 - NO_2 - H_2CO Reactions.....	30
DISCUSSION.....	31
O_3 Decomposition.....	31
O_3 - H_2O_2 Reaction.....	33
O_3 - HCO_2H Reaction.....	34
O_3 - H_2CO Reaction.....	36
O_3 - C_2H_4 Reaction.....	37
Aqueous-phase PAN Reactions.....	38
O_3 - NO_2 - H_2CO Reactions.....	39
Aqueous-phase H_2O_2 Formation.....	41
SUMMARY.....	45
REFERENCES.....	47

TABLES

<u>Number</u>		<u>Page</u>
1	Second-order Rate Constant of O_3 - HCO_2H Reaction at $25 \pm 2^\circ C$	17
2	Second-order Rate Constant of Aqueous O_3 - C_2H_4 Reaction.....	26
3	Summary of the Reaction Kinetics and Product Analysis of Some Aqueous-phase O_3 Reactions.....	46

FIGURES

<u>Number</u>		<u>Page</u>
1.	Schematic diagram of the experimental setup.....	6
2.	Aqueous-phase O_3 spectra, taken during the course of saturation process.....	8
3.	Time dependence of the O_3 saturation in a 2-liter bubbler. Flow of $O_2 = 70 \text{ cc min}^{-1}$	9
4.	Schematic of the construction of a bubbler-type gas-liquid reaction vessel.....	11
5.	Kinetics of O_3 decomposition in a 5 cm cylindrical quartz optical cell.....	13
6.	Time dependence of O_3 disappearance due to reaction with H_2O_2 . Initial concentrations of H_2O_2 and O_3 were $3.5 \times 10^{-5} \text{ M}$ and $\sim 4 \times 10^{-6} \text{ M}$, respectively.....	14
7.	Plot of a second-order kinetics of $O_3 - HCO_2H$ reaction at neutral pH and $[O_3]_0 \cong [HCO_2H]_0 = 2 \times 10^{-5} \text{ M}$	16
8.	Plot of the pH dependence of the second-order rate constants for $O_3 - HCO_2H$ reaction.....	18
9.	Time dependence of O_3 disappearance due to the reaction with H_2CO . Initial concentrations of H_2CO and O_3 were $1.65 \times 10^{-4} \text{ M}$ and $1.7 \times 10^{-5} \text{ M}$, respectively.....	20
10.	Reaction order determination for H_2CO-O_3 reaction. Plots were made according to eq. 8 and eq. 9. See text for definitions.....	23
11.	Time dependence of the concentration of peroxides as a reaction product in the $O_3 - C_2H_4$ reaction; $pO_3 = 0.5 \text{ ppm}$ and $pC_2H_4 = 28 \text{ ppm}$	25
12.	Contribution of the aqueous-phase HO_2 recombination reaction to the generation of H_2O_2 , calculated for pH = 4.7. Line A represents the aqueous-phase rate of H_2O_2 production and line B the characteristic reaction time for gaseous HO_2 at $L = 10^{-6}$. Line C indicates the contribution to aqueous H_2O_2 from the gas-phase recombination of HO_2 ; H_2O_2 produced is assumed to be rapidly incorporated into the liquid water ($L = 10^{-6}$).....	44

ACKNOWLEDGMENT

The author acknowledges Mr. Paul J. Klotz for his assistance in carrying out part of the laboratory experiments, and Dr. Thomas J. Kelly for his effort in making the H_2O_2 detection technique available to us. The author also wishes to thank Dr. Kelly and Dr. Roger L. Tanner for helpful discussions.

INTRODUCTION

The major acids found in the rain and precipitation, namely, sulfuric acid and nitric acid, are believed to be derived from SO_2 and NO_x emitted mainly from the burning of fossil fuel by power plants, smelters, mobile sources, and space heating. The production of final acids from their precursors in the atmosphere involves chemical transformation in conjunction with transport and deposition. In order to formulate an efficient strategy to control the amount of acid deposition and to minimize its environmental impact, a thorough understanding of each of these atmospheric processes has to be acquired.

It has been recognized that the chemical reactions which produce the strong acids can take place either in the gas phase or in the liquid phase. This notion was established because convincing evidence has been collected to indicate that the atmospheric oxidation of SO_2 is strongly affected by aqueous-phase reactions, particularly by ozone and hydrogen peroxide (Penkett et al., 1979; Martin, 1983; Kelly et al., 1984). In order to assess the importance of these aqueous reactions, the atmospheric concentrations of O_3 and H_2O_2 have to be determined. Although the gas-phase concentration of ozone can be accurately determined by various techniques such as ethylene-chemiluminescence, no viable method is currently available for the measurement of gas-phase concentrations of H_2O_2 . As a result, the major gas-phase routes for peroxide generation, i.e., the recombination of hydroperoxy radicals and the photolysis of formaldehyde, cannot be confidently employed in a numerical model as the sole source for this species. Furthermore, recent attempts to determine the gas-phase concentrations of peroxide using bubbler series have revealed the existence of in-situ

production of artifact peroxide (Heikes, 1984; ten Brink et al., 1984; Heikes et al., 1982). This observation suggests that aqueous pathways for peroxide formation might exist, in addition to gas-phase counterparts. Clearly, these pathways have to be identified and characterized before it is possible to accurately model the atmospheric budget of H_2O_2 and the rate of SO_2 oxidation.

Among the various potential precursors of aqueous peroxide, it has been speculated that O_3 might be a plausible candidate based on the following considerations: (1) O_3 produces peroxides upon reaction with certain organic compounds, e.g., olefins (Gilbert, 1976), (2) aqueous-phase O_3 reactions were found to involve free radicals derived from O_3 decomposition that might serve as H_2O_2 precursors (Hoigne and Bader, 1976; Böhler et al., 1984), and (3) the bubbler series experiments (Heikes et al., 1982) demonstrated that the levels of artifact H_2O_2 do not diminish rapidly along the bubbler train, consistent with the presumption that the precursor species might be present in relatively high concentration and has a low aqueous solubility. O_3 appears to fit the description.

Although the aqueous-phase reactions of ozone have been the subject of numerous studies for the past several decades, major gaps exist in the understanding of the detailed features of these reactions. Additionally, with the emphasis focused on drinking water treatment by O_3 , no major efforts have been directed to the product analysis, particularly H_2O_2 . In this current laboratory research we have examined a series of aqueous-phase reactions of O_3 and the formation of H_2O_2 and organic peroxides as their reaction products in an attempt to identify the direct aqueous sources of peroxides. The aqueous-phase reaction systems examined include: (1) O_3

self-decomposition, (2) O_3 reaction with formaldehyde (with and without the presence of NO_2), formic acid, ethylene, and peroxyacetyl nitrate (PAN). Except for the O_3 -ethylene reaction, peroxide was not found as a reaction product. From this study it is concluded that the reactions studied here are not significant with respect to atmospheric peroxide formation. As a result, knowledge of direct aqueous-phase sources of H_2O_2 remains highly uncertain.

EXPERIMENTAL

Material

N₂ (Ultra-High Purity, Linde) and O₂ (Ultra-Zero, Matheson) which contained less than 0.5 ppm hydrocarbons (as CH₄) were used as the diluent gas and O₃ source, respectively. NO₂ gas was prepared using a NO₂ permeation device (Metronics, Inc., wafer type) enclosed in a constant temperature oven. With a 15 cc/min flow of N₂ carrier gas the concentration of NO₂ from the permeation source was determined to be 6.0 ppm. Samples of gaseous PAN (6-12 ppm) in N₂ were prepared from concentrated PAN stock solution in n-tridecane prepared according to a modified method of Nielsen (Nielsen et al., 1982; Gaffney et al., 1984). Working stock of ethylene (0.140%) in N₂ was prepared from pure ethylene (99.5%, Scott) and UHP N₂. H₂O₂ (30%), H₂CO (37%) and HCO₂H (88%), all of reagent grade from either Mallinckrodt or Baker, were used without purification. EDTA, Trisma Base, and horseradish peroxidase were of the highest purity available from Sigma Chemical Company. Concentrated HCl and NaOH and inorganic salts such as KCl were all of reagent grade and used without further purification. Fresh solutions of standards and reagents were prepared for the same-day use. Distilled water (resistance \geq 16 Mohm at 25°C), which had been further purified by Millipore Milli-Q System, was used for all of the studies.

Ozone Generation

Two ozone generators were used. For higher O₃ output (up to ~200 ppm at 1 l/min flow rate) a 10-inch Pen-Ray UV lamp was employed. This source was used mainly for the preparation of saturated O₃ solutions for batch-type reactions. For lower O₃ output, an AID Ozone Generator (Model 565) equipped

with a continuously adjustable shutter for O_3 output control was employed (up to ~ 1 ppm at 1 l/min flow). This source was used for the continuous-flow reaction system. For both generators, high purity O_2 purchased from Matheson was used in order to minimize possible interferences from NO_x and organic impurities.

Gas-Phase O_3 Concentration

Gas-phase O_3 concentrations were determined either by a Dasibi O_3 monitor (Model 1008-PC) operated based on UV absorption (LOD ~ 2 ppb) or by a Monitor Labs Ozone Analyzer (Model 8410) based on the O_3 -ethylene chemiluminescence, with the former as the primary standard.

Preparation of O_3 Solution

For batch-type experiments, O_3 solutions were prepared in a 2-liter bubbler through which O_3 was continuously bubbled at a total O_2 flow of 70 cc min⁻¹ (Fig. 1). Each Pyrex O_3 bubbler (volume ~ 2 l) was equipped with a coarse-sized frit for the enhancement of mixing and with two ports for liquid transfer. The plumbing was constructed with parts made either of stainless steel or Teflon for purity. Bubbler 1 was needed for the humidifying of the gas-stream and the removal of any soluble substances; bubbler 2 supplied saturated O_3 solutions to be used for the batch studies. For the study of some continuous-flow reactions, valve M-1 was switched so that O_3 would flow through the gas-liquid reaction cell to initiate the reactions.

Aqueous-Phase O_3 Concentration

The concentration of aqueous-phase O_3 was measured by a UV-vis spectrophotometer (Beckman Model DU-7) using either a 10 cm or a 5 cm cylindrical




Figure 1. Schematic diagram of the experimental setup.

optical cell. Spectra of the O_3 solutions in the bubbler were taken during the saturation process (Fig. 2) and the concentrations of O_3 were determined from the absorption peak at 260.0 nm using an extinction coefficient $\epsilon = 2930 \text{ M}^{-1} \text{ cm}^{-1}$ (Kosak-Channing and Helz, 1983). The time dependence of O_3 saturation in the 2-liter bubbler is shown in Figure 3. Using the 10-inch Pen-ray UV ozonator at O_2 flow rate of $70\text{--}90 \text{ cc min}^{-1}$, a typical aqueous-phase O_3 concentration of $1\text{--}2 \times 10^{-5} \text{ M}$ was obtained after 40 min of bubbling. The limit of detection for O_3 with the use of the 10-cm cell is $1 \times 10^{-7} \text{ M}$.

H_2O_2 and Organic Peroxide Concentrations

Concentrations of aqueous-phase H_2O_2 and organic peroxides were determined by the horseradish peroxidase-fluorescence technique (HRPF) (Guilbault et al., 1968) modified by NCAR (Lazrus et al., 1983). In our arrangement, a Perkin-Elmer fluorometer (Model 204S) was employed in conjunction with a liquid flow reaction system equipped with a rotary injection valve (Altex, sample loop size 0.5 ml). The limit of detection of the HRPF technique is $1 \times 10^{-7} \text{ M}$. Since this technique does not distinguish organic peroxides from inorganic H_2O_2 , the determination of the concentration of organic peroxides was achieved by a difference method in which H_2O_2 is preferentially destroyed (or inactivated) by the enzyme catalase (Schonbaum and Chance, 1976).

Kinetic Measurements

Two different types of kinetic methods were used. In the batch-type study, reactions were initiated by mixing the reagents with O_3 solution in an optical cell and the decrease of $[O_3]$ accompanying the reactions were




Figure 2. Aqueous-phase O_3 spectra, taken during the course of saturation process.




Figure 3. Time dependence of the O_3 saturation in a 2-liter bubbler.
Flow of $O_2 = 70 \text{ cc min}^{-1}$.

followed spectrophotometrically at $\lambda = 260$ nm. In the continuous-flow reaction, a reagent gas mixture containing constant concentrations of O_3 and other gaseous reactants was continuously bubbled through a solution contained in a bulk-type gas-liquid reactor. The detailed construction of the gas-liquid reactor is shown schematically in Figure 4. The kinetics of H_2O_2 generation were followed by an aliquot method in which H_2O_2 concentrations were determined by the HRPD technique. Temperature of the reaction vessel was maintained at $22.0 \pm 0.1^\circ C$ for the latter method, but for the spectrophotometric technique the uncertainty was somewhat greater ($\pm 2^\circ C$) due to the lack of temperature control of the sample compartment for a cylindrical cell.




Figure 4. Schematic of the construction of a bubbler-type gas-liquid reaction vessel.

RESULTS

O₃ Decomposition in Optical Cell

In order to study the reaction kinetics of O₃ with other substrates using a spectrophotometric method, it is necessary to determine the stability of O₃ solution in the optical cell employed. Figure 5 shows a typical decay curve of O₃ in a 5 cm cylindrical cell under neutral pH (no acid or base added). The first-order rate constant was found to be $2.1 \times 10^{-4} \text{ s}^{-1}$. When pH of the solution was adjusted to 2 by HCl the rate constant of O₃ decomposition dropped to $1.4 \times 10^{-4} \text{ s}^{-1}$. These values were repeatable within 15% and were found to be slow compared to other O₃ reaction.

The final decomposition mixture of the O₃ solutions (containing initial [O₃] as high as $2.0 \times 10^{-5} \text{ M}$) was analyzed for H₂O₂. The concentration of H₂O₂ was found to be below the limit of detection of $1 \times 10^{-7} \text{ M}$.

O₃ - H₂O₂ Reaction

In order to examine possible H₂O₂ formation in O₃ reactions, the reaction between O₃ and the reaction product, H₂O₂, should be quantified. The kinetics of this reaction was studied under pseudo-first-order condition, i.e., $[\text{H}_2\text{O}_2]_0 \gg [\text{O}_3]_0$. A typical trace of O₃ decomposition under such a condition is shown in Figure 6. The O₃ decay kinetics was found to conform to a pseudo-first-order reaction for ~3 half-lives. The second-order rate constant, obtained by dividing the pseudo-first-order rate constants by $[\text{H}_2\text{O}_2]_0$, which had been varied from $8 \times 10^{-6} \text{ M}$ to $6.4 \times 10^{-5} \text{ M}$, was determined to be $(2.6 \pm 0.4) \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ at neutral pH. The good agreement obtained between the second-order rate constants determined at widely different H₂O₂ concentrations permits the conclusion that the reaction kinetics are also first order with respect to [H₂O₂]. At lower pH the




Figure 5. Kinetics of O_3 decomposition in a 5 cm cylindrical quartz optical cell.




Figure 6. Time dependence of O_3 disappearance due to reaction with H_2O_2 . Initial concentrations of H_2O_2 and O_3 were 3.5×10^{-5} M and $\sim 4 \times 10^{-6}$ M, respectively.

reaction rate decreased rapidly. A second-order rate constant of $38 \pm 8 \text{ M}^{-1} \text{ s}^{-1}$ was determined at $\text{pH} = 4$, where the solutions contained $[\text{HCl}] = 1.00 \times 10^{-4} \text{ M}$.

$\text{O}_3 - \text{HCO}_2\text{H}$ Reaction

Formic acid has been identified as a major organic acid component in rainwater (Keene et al., 1983) and therefore the sources and sinks of this species may be of importance to the understanding of acid rain formation mechanisms. We examined the aqueous-phase reaction of O_3 with HCO_2H to determine the reaction rate and the yield of peroxide as a product.



The kinetic study of reaction (1) was made by monitoring the change of $[\text{O}_3]$ followed spectrophotometrically at $\lambda = 260 \text{ nm}$. Since the rate of this reaction under current conditions is too fast to be studied under pseudo-first-order condition, i.e., high concentrations of HCO_2H , initial concentrations of O_3 and HCO_2H were made approximately equal at ca. $2 \times 10^{-5} \text{ M}$. Treating the reaction with an overall second-order kinetics, i.e.,

$$\text{Rate} = \frac{-d[\text{O}_3]}{dt} = k_1' [\text{O}_3] [\text{HCO}_2\text{H}] \quad (2)$$

the rate constant was determined by using a second-order plot

$$\frac{1}{[\text{O}_3]_t} - \frac{1}{[\text{O}_3]_0} = k_1' t \quad (3)$$

Plots of $[\text{O}_3]_t^{-1}$ vs. time were found to be linear for at least three half-lives. A typical run is shown in Fig. 7. The rate constants obtained are listed in Table 1 as a function of pH.




Figure 7. Plot of a second-order kinetics of $O_3 - HCO_2H$ reaction at neutral pH and $[O_3]_0 \approx [HCO_2H]_0 = 2 \times 10^{-5} M$.

Table 1
Second-Order Rate Constant of O_3 - HCO_2H Reaction at $25 \pm 2^\circ C$

pH	k_1' , $M^{-1} s^{-1}$
3.00 ± 0.02	$(2.5 \pm 0.2) \times 10^2$
4.00 ± 0.05	$(1.8 \pm 0.2) \times 10^3$
4.60 ± 0.2	$(3.3 \pm 0.5) \times 10^3$

The pH dependence of k_1' , increasing with higher pH, can be fitted to a rate law which assumes the rate determining step involves O_3 and the dissociated formate ion, i.e.,

$$\begin{aligned}
 \text{Rate} &= k_4 [O_3][HCO_2^-] \\
 &= k_4 \left(\frac{K_a}{[H^+] + K_a} \right) [O_3][HCO_2H]_T \\
 &= k_1' [O_3][HCO_2H]_T
 \end{aligned} \tag{4}$$

where K_a is the acid dissociation constant of formic acid and $[HCO_2H]_T$ is the total analytical concentration of formic acid. Fitting Eq. (4) to the experimental data allowed the values of k_4 and K_a to be determined; they were found to be $4.3 \times 10^3 M^{-1} s^{-1}$ and $8.9 \times 10^{-5} M^{-1}$, respectively.

Although the agreement between the data and the calculated curve appears to be quite reasonable (Fig. 8), the pK_a value of formic acid determined here, 4.1 ± 0.1 , is nearly 0.3 unit higher than the literature value of 3.75 (Riddick and Bunger, 1970). While this discrepancy may be acceptable in view of the fact that the kinetic approach generally would yield a larger uncertainty for pK_a determination (Sillen and Martell, 1964) than




Figure 8. Plot of the pH dependence of the second-order rate constants for $\text{O}_3 - \text{HCO}_2\text{H}$ reaction

other methods (e.g., conductometric), we nonetheless consider this base-catalyzed mechanism tentative.

Final reaction mixtures of the O_3 - HCO_2H reaction were analyzed for peroxide by the HRP method. The level of peroxide was found to be smaller than the LOD of the instrument, namely, 1×10^{-7} M. Under the present reaction conditions, e.g., $[O_3]_0 = [HCO_2H]_0 = 2 \times 10^{-5}$ M, a value of 0.5% is estimated as the upper limit for the peroxide yield of the O_3 - HCO_2H reaction.

O_3 - H_2CO Reaction

Since formaldehyde is a relatively abundant atmospheric organic constituent (Grosjean, 1982; Tanner and Meng, 1984) which is derived mainly from combustion emission and the atmospheric oxidation of higher hydrocarbons, its chemistry is of considerable interest to the understanding of atmospheric oxidant formation mechanisms. The aqueous-phase reaction of O_3 with H_2CO was studied in a similar fashion to the O_3 - HCO_2H reaction, namely, the extent of the reaction in a 5-cm optical cell was monitored by UV absorption for $[O_3]$ decrease.

The kinetics, under conditions where initial $[H_2CO]$ was in large excess of $[O_3]$, did not conform to a pseudo-first-order reaction. In fact, the effective second-order rate constant appeared to increase as $[O_3]$ was decreasing, indicating a reaction order of less than unity with respect to O_3 . This behavior is illustrated by a typical run shown in Figure 9 where $[H_2CO]_0 = 1.65 \times 10^{-4}$ M and $[O_3]_0 = 1.75 \times 10^{-5}$ M.

Since it has been shown that formic acid is produced as a product in the O_3 - H_2CO reaction (Kuo and Wen, 1977), i.e.,






Figure 9. Time dependence of O_3 disappearance due to the reaction with H_2CO . Initial concentrations of H_2CO and O_3 were 1.65×10^{-4} M and 1.7×10^{-5} M, respectively.

the O_3 decomposition may be enhanced by the subsequent O_3 - HCO_2H reaction, thereby manifesting a non-first-order kinetics. Comparing the initial rates of the O_3 - H_2CO reaction with that for O_3 - HCO_2H reaction indicated that the O_3 decomposition due to the secondary reaction with the intermediate product, HCO_2H , can be significant, if HCO_2H is indeed produced stoichiometrically. However, since the product analysis reported by Kuo and Wen (1977) did not appear to be quantitative and it is not clear to us whether or not formic acid is produced from an elementary reaction step, we analyzed our experimental data without invoking the reaction of HCO_2H , but strictly in terms of O_3 and H_2CO . It should be noted that the general kinetic behavior so deduced may be applied only to reaction conditions similar to that employed in this study.

To determine the reaction order with respect to each of the reagent concentrations, we consider the equation

$$\text{Rate} = \frac{-d[O_3]}{dt} = k[O_3]^n[H_2CO]^m \quad (6)$$

Under conditions that $[H_2CO]_0 \gg [O_3]_0$, Eq. (6) is rearranged to yield

$$(\text{Rate}/[H_2CO]^m) = k [O_3]^n \quad (7)$$

where $[H_2CO]^m$ is a constant throughout the reaction course. Taking logarithm of Eq. (7), we obtain

$$\log (\text{Rate}/[H_2CO]^m) = \log k + n \log [O_3] \quad (8)$$

Fitting Eq. (8) to the data obtained from Figure 9 (from which a set of Rate vs. $[O_3]$ can be determined for any specified time), the value of the slope,

n, was found to be $\sim 1/2$ (Fig. 10). Using the same approach, an analogous equation is obtained as

$$\log (\text{Rate}_0/[\text{O}_3]^{1/2}) = \log k + m \log [\text{H}_2\text{CO}]_0 \quad (9)$$

To determine m, we measured the initial rate for three initial formaldehyde concentrations: $[\text{H}_2\text{CO}]_0 = 1.65 \times 10^{-4}$ M (Fig. 9), 8.2×10^{-5} M, and 4.1×10^{-5} M. Fitting Eq. (9) to the rate data, m was found to have a value also close to $1/2$ (Fig. 10). Eq. (6) is now tentatively identified as

$$\text{Rate} = k [\text{O}_3]^{1/2} [\text{H}_2\text{CO}]^{1/2} \quad (10)$$

with $k = (1.15 \pm 0.7) \times 10^{-3} \text{ s}^{-1}$, at $t = 20 \pm 2^\circ\text{C}$, for the reaction conditions employed.

Product analysis of the final reaction mixtures showed that H_2O_2 was a minor product. With $[\text{H}_2\text{CO}]_0 = 8 \times 10^{-5}$ M and $[\text{O}_3]_0 = 8 \times 10^{-6}$ M the H_2O_2 formed was found to be 1.5×10^{-7} M at the conclusion of the reaction.

$\text{O}_3\text{-C}_2\text{H}_4$ Reaction

Olefins represent an important class of organic constituents of the atmosphere because of their reactivity towards O_3 and OH oxidation (Niki et al., 1983). Since the Criegee intermediates formed in O_3 -olefin reactions may transform into peroxides in protic solvents, we have examined the kinetics of the aqueous O_3 -ethylene reaction in connection to its rate of peroxide production. Due to the low solubility of C_2H_4 , i.e., $H_{\text{C}_2\text{H}_4} = 5.0 \times 10^{-3} \text{ M atm}^{-1}$ at 25°C (Wilhelm et al., 1977), the kinetics of this reaction was studied by the use of the continuous-flow method. A constant flow of N_2 (typically 2.0 l min^{-1}) was first allowed to flow through the gas-liquid




Figure 10. Reaction order determination for $\text{H}_2\text{CO}-\text{O}_3$ reaction. Plots were made according to eq. 8 and eq. 9. See text for definitions.

reaction cell containing a known volume of liquid water to remove the dissolved CO_2 . When the conductivity of the liquid water had stabilized, reagent C_2H_4 was added to the gas stream; the concentration range of C_2H_4 employed was 7-28 ppm. After the reaction was initiated by the addition of O_3 (concentration range: 0.5-1.0 ppm), the reaction mixture was analyzed for peroxide concentration at known time intervals using the HRPF technique. The concentration of peroxide of the reaction mixture was found to increase linearly with time (Fig. 11) and the reaction rate defined as

$$\text{Rate} = \frac{d[\text{peroxide}]}{dt} \quad (11)$$

was determined.

Assuming that the aqueous-phase reaction of O_3 and C_2H_4 is first order with respect to each reagent, eq. (11) is given as

$$\frac{d[\text{peroxide}]}{dt} = k_{12} [\text{O}_3][\text{C}_2\text{H}_4] \quad (12)$$

Additionally, assuming that the liquid water is saturated with the reagent gases according to Henry's Law, i.e.,

$$[X] = H_X P_X \quad (13)$$

which allows eq. (12) to be rewritten as

$$\frac{d[\text{peroxide}]}{dt} = k_{12} H_{\text{O}_3} H_{\text{C}_2\text{H}_4} P_{\text{O}_3} P_{\text{C}_2\text{H}_4} \quad (14)$$

the second-order aqueous-phase rate constant k_{12} can readily be determined. The values of k_{12} determined at various reagent concentrations and pHs are




Figure 11. Time dependence of the concentration of peroxides as a reaction product in the $O_3 - C_2H_4$ reaction; $pO_3 = 0.5$ ppm and $pC_2H_4 = 28$ ppm.

listed in Table 2. It should be noted that the average values were obtained from repeating runs as well as from runs with different reagent concentrations. The fact that the rate constant determined from eq. (14) remained essentially constant while the P_{O_3} and $P_{C_2H_4}$ were varied by a factor of 2 and 4, respectively, lends support to the assumed overall second-order kinetics.

For a gas-liquid reaction taking place in bulk liquid, particular attention has to be given to the mass transfer characteristics of the apparatus in order to identify the extent of mass transfer limitations on the

Table 2
Second-Order Rate Constant of Aqueous O_3 - C_2H_4 Reaction*

pH	$10^{-5} \times k_{12}, M^{-1} s^{-1}$
neutral (6 ~ 7)	2.60 ± 0.26
5.0	3.0 ± 0.25
4.0	3.0 ± 0.23
3.0	3.3 ± 0.41

*[KCl] = 1×10^{-3} M, except for runs at pH 3; $T = 22.0 \pm 0.1^\circ C$.

overall reaction rates (Danckwerts, 1970). A simple approach to this identification is via the comparison of reaction time constant (τ_r) to the mass mass transfer time constant (τ_m): when $\tau_r \geq 10 \tau_m$ the reaction is purely chemical reaction limited; when $\tau_r \leq 10 \tau_m$ mass transfer will become a rate limiting process. For our currently employed gas-liquid reactor, the phase-mixing rate constant, k_m , was determined to be $0.47 s^{-1}$ (at liquid volume = 15 ml, total gas flow rate = $2.0 l \min^{-1}$, cf. Lee and Schwartz, 1981). The time constants for the removal of aqueous O_3 and C_2H_4 by each

other are estimated as $\tau_{O_3} = (3 \times 10^5 [C_2H_4])^{-1} = 24 \text{ s}$ and $\tau_{C_2H_4} = (3 \times 10^5 [O_3])^{-1} = 300 \text{ s}$. These reaction time constants are significantly longer than τ_m ($\approx k_m^{-1} = 2.14 \text{ s}$); it is readily established that the chemical reaction is the rate limiting process and the reaction rate is not significantly affected by mass transfer processes. The assumption of the establishment of gas-liquid equilibrium is therefore validated.




From these results it is also seen that peroxide formation from the aqueous $C_2H_4-O_3$ reaction is independent of the solution pH;; the average rate coefficient $k_{12} = (3.0 \pm 0.3) \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$, is applicable over the pH range 3 - 7. It should be noted that the fraction of organic peroxide formed in the $C_2H_4-O_3$ reaction was determined using the enzyme catalase technique in which the H_2O_2 is preferentially destroyed (Kelly et al., 1984; Schonbaum and Chance, 1976). The results of this test showed that $\approx 60\%$ (average of 3 runs made at pH 3) of the total peroxide was present as organic peroxide which appeared to be reasonably stable. The identity of the organic peroxide was not determined, but from the mechanism advanced for the formation of ozonide and Criegee radical, CH_3O_2H is considered a possible candidate.

Aqueous-phase Reactions of PAN

Peroxyacetyl nitrate (PAN), formed from the reaction of $CH_3C(O)O_2$ and NO_2 , is an important atmospheric species because of its capacity as a reservoir for NO_2 and free radical species in transport processes. PAN was synthesized according to the method described by Nielsen (1982) and was stored in a heavy lipid, i.e., n-tridecane. Gas samples of PAN were prepared by mixing N_2 with the distillate from PAN/n-tridecane solution. With PAN vapor

pressure (over n-tridecane) in the order of 10 torr, 10 ppm gaseous PAN standards can be readily prepared. Because it was found that PAN decomposes on standing with a half-life of approximately 110 hrs (at 25°C, in the absence of NO, Senum and Gaffney, 1984), PAN standards were always freshly prepared for the same day use.

The three reactions of PAN examined include



The typical concentrations used for PAN, O₂, and O₃ were 100 ppb, 20%, and 1.0 ppm, respectively. Due to the low solubilities of O₂ and O₃, as well as of PAN ($H_{\text{PAN}} = 3.6 \text{ M atm}^{-1}$, cf. Lee, 1984), these reactions were again studied by the use of the gas-liquid reaction cell described above. Final reaction mixtures (with total reaction time up to 1 hr) of these reactions were analyzed for peroxide and it was found that the levels of peroxide were all below the detection limit of our HRPD technique. Using a general rate expression

$$\frac{d[\text{peroxide}]}{dt} = k [\text{PAN}] [\text{O}_x] \quad (18)$$

the upper limits for rate constants k_{15} , k_{16} , and k_{17} were estimated to be $2 \times 10^{-4} \text{ s}^{-1}$, $3 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$, and $3 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$, respectively.

O₃ - NO₂ Reaction

In view of the suggestion that NO₃ is potentially important in cloud water chemistry (Heikes and Thompson, 1983), we have conducted some preliminary studies of NO₃ + H₂O. Since the solubility of NO₂ was not expected to be large, the reaction was carried out in the bubbler-type gas-liquid reactor into which the reagent gases were continuously replenished. Study of the background reaction involving H₂O and 1.3 and 43 ppm of O₃ showed that no detectable H₂O₂ was produced after 40 min of reaction. This result is consistent with that observed for the pure aqueous-phase O₃ decomposition study conducted in an optical cell (see above). To conduct the NO₂ experiment a pre-reaction chamber of 1-liter size was placed upstream of the bubbler to allow the production of NO₃ from the O₃- NO₂ gas-phase reaction. Since the second-order gas-phase rate constant for



is $3 \times 10^{-17} \text{ cm}^3 \text{ s}^{-1}$ (Baulch et al., 1982), the 1/e time for NO₃ formation is calculated to be 31 s at pO₃ = 43 ppm, the O₃ concentration employed. With 30 s residence time in the mixing chamber (total gas flow rates = 2 l min⁻¹), nearly 70% of the NO₂ should be converted to NO₃ (pNO₂ employed was 40 ppb). It should be noted that the experiments were carried out in the dark in order to avoid the photolysis of NO₃. At the end of 40 min the reaction mixture in the gas-liquid reactor was analyzed for H₂O₂ content. Concentration of H₂O₂ was found to be $\sim 1 \times 10^{-7} \text{ M}$. This low concentration seems to indicate that NO₃ has a limited role in the direct aqueous production of peroxide, at least in pure water and under neutral pH.

O₃-NO₂-H₂CO Reaction

Reactions of NO₃ with atmospheric constituents have been found to be important in nighttime chemistry (see, for example, Pitts et al., 1984). Since NO₃ can react with aldehydes to initiate free radical chain reactions, we have tested the possibility of the formation of peroxide by the aqueous reaction of NO₃ and H₂CO. In our approach, NO₂ (40 ± 1 ppb) and O₃ (40 ± 1 ppm) in N₂ (with 2% O₂) were allowed to react in a pre-reaction chamber (ca. 1 l size) for ~30 s before entering the gas-liquid bubbler that contained 20 ml of a 1.0 × 10⁻⁴ M H₂CO solution. According to our calculation given in the previous section, nearly 70% of the NO₂ should be converted to NO₃. The reaction extent of the formaldehyde solution was followed by the peroxide concentration determined by an aliquot method. Despite the long reaction time (up to 100 min) no discernible peroxide was detected as a product.

DISCUSSION

A number of selected aqueous-phase reactions of O_3 have been studied in this research project. The main purpose of this work is to identify and characterize those aqueous O_3 reactions that would produce either hydrogen peroxide or organic peroxides as reaction products. Since the effort was mainly directed at the identification of those reactions, the entire project was scoping in nature; rather than spending a great length of time to characterize the detailed kinetics of a reaction which does not produce peroxide, we tried to maximize the number of reactions to be examined for peroxide production. As a consequence, the kinetic information obtained here with limited variations in reaction conditions are necessarily not comprehensive. For instance, reagent concentration variations were generally limited, the solution acidity was normally confined between neutral and pH 3, and the temperature was fixed either at $22.0 \pm 0.1^\circ\text{C}$ or room temperature. Nevertheless, we have attempted to make use of the data obtained in this work to qualitatively assess the atmospheric importance of these aqueous-phase O_3 reactions to the formation of peroxide and to the depletion of the reagent species.

O_3 Decomposition

Despite the fact that the kinetics of O_3 decomposition in aqueous solutions have been studied extensively, no agreement was reached with respect to its reaction order. The reaction order reported in the literature ranged from 1 (for example, Sullivan and Roth, 1979) to 2 (for example, Gurol and Singer, 1982), with some as $3/2$ (for example, Kilpatrick et al., 1956). This discrepancy was suggested to result from a number of possible sources

such as mass transfer limitations, ionic strengths, impurities, as well as effects of buffers with which the solution pH were adjusted. The main features of O_3 decomposition observed in this current study are in agreement with some of the recently published results: (1) O_3 decomposition is insensitive to pH in the acidic region (Gurol and Singer, 1982), (2) decay of O_3 is first order in O_3 concentration (Sullivan and Roth, 1979), and (3) the first order rate constants determined in this work, i.e., $2.1 \times 10^{-4} \text{ s}^{-1}$ at neutral pH and $1.4 \times 10^{-4} \text{ s}^{-1}$ at pH = 2, are within a factor of 2 of those determined by Sullivan and Roth (1979), i.e., $4.1 \times 10^{-4} \text{ s}^{-1}$ and $2.4 \times 10^{-4} \text{ s}^{-1}$ for pH = 4 and 2, respectively. It should be noted that the aqueous-phase concentrations of O_3 used in this study were smaller than those used in previous studies by a factor of at least 10, and, therefore, may more closely resemble the actual atmospheric conditions. Using the aqueous-phase decomposition rate expressed as the gas-phase reaction rate (Lee and Schwartz, 1981), i.e.,

$$-\frac{PO_3}{dt} = k \text{ } H_2O_3PO_3 \text{ LRT} \quad (20)$$

where L is the liquid water content of the atmosphere (volume fraction), we estimated the reaction time constant of O_3 against aqueous-phase decomposition to be $5 \times 10^6 \text{ hr}$ or longer for solution pH of neutral or lower at $L = 10^{-6}$. If wall effects in the laboratory study are partially responsible for the O_3 decomposition, then these estimated time constants would represent lower limits, rendering the atmospheric self-decomposition of O_3 in aqueous droplets even less important.

O₃-H₂O₂ Reaction

O₃ decomposition is known to be initiated by OH⁻ ion under basic conditions (Guroi and Singer, 1982; Sullivan and Roth, 1979; Staehelin and Hoigne, 1982). According to Staehelin and Hoigne, the conjugate base of H₂O₂, i.e., HO₂⁻, also reacts with O₃ to produce free radicals such as OH[•] and O₂^{•-}. From studies made in alkaline solutions, the rate constant for the O₃ - HO₂⁻ reaction, defined as

$$-\frac{d[O_3]}{dt} = k_{21} [O_3] [HO_2^-] \quad (21)$$

was given as $(2.8 \pm .5) \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$. Replacing [HO₂⁻] with the total analytical concentration of H₂O₂, [H₂O₂]_T, we obtain

$$\begin{aligned} -\frac{d[O_3]}{dt} &= k_{21} \frac{K_a}{K_a + [H^+]} [O_3] [H_2O_2]_T \\ &= k_{21}' [O_3] [H_2O_2]_T \end{aligned} \quad (22)$$

where K_a is the acid dissociation constant of H₂O₂. For pK_a = 11.6 and pH = 6, k₂₁' is calculated to be 7.0 M⁻¹ s⁻¹. This value is significantly smaller than that determined here; it is therefore concluded that the second-order rate constant determined in this study may reflect the direct reaction of O₃ and H₂O₂. The lower rate constant determined at pH 4 maintained with [HCl] = 1 × 10⁻⁴ M is qualitatively consistent with a negative [H⁺] dependence, although the possibility that the lower rate constant resulted from Cl⁻ ion inhibition (Taube and Bray, 1940) can not be excluded.

The fact that H_2O_2 can be destroyed by O_3 requires the effect of this O_3 - H_2O_2 reaction on H_2O_2 concentrations be quantitatively assessed. To evaluate the time constant for the removal of H_2O_2 by the O_3 - H_2O_2 reaction, we employ

$$\begin{aligned}\tau_{\text{H}_2\text{O}_2} &= (k_{23} [\text{O}_3])^{-1} \\ &= (k_{23} \text{H}_0\text{O}_3 \text{pO}_3)^{-1}\end{aligned}\tag{23}$$

With $\text{pO}_3 = 1$ ppm, $\tau_{\text{H}_2\text{O}_2}$ is calculated to be 10 hr at neutral pH and 660 hr at pH 4. These time constants are apparently too long to be important for our current laboratory studies.

The time constant for removal of atmospheric gas-phase O_3 by the aqueous O_3 - H_2O_2 reaction is given as

$$\tau_{\text{O}_3} = (k_{23} \text{H}_0\text{O}_3 [\text{H}_2\text{O}_2] \text{LRT})^{-1}\tag{24}$$

For $[\text{H}_2\text{O}_2] = 3 \times 10^{-5}$ M, a representative summer cloudwater H_2O_2 concentration (cf. Kelly et al., 1984), and $L = 10^{-6}$, τ_{O_3} is calculated to be 1.4×10^4 hr and 1.0×10^6 hr at neutral pH and pH 4, respectively. Since the aqueous-phase O_3 concentration in atmospheric water is significantly smaller than that of H_2O_2 , the removal of the dissolved atmospheric H_2O_2 by the aqueous-phase O_3 - H_2O_2 reaction will be extremely slow.

O_3 - HCO_2H Reaction

Kinetics of this aqueous-phase reaction have been reported. Kuo and Wen (1977) found that the rate of O_3 - HCO_2H reaction is first order with respect to $[\text{O}_3]$, and is dependent on $[\text{HCO}_2\text{H}]$ to the 1.2 power. The rate constant defined by

$$-\frac{d[O_3]}{dt} = K_{25} [O_3] [HCO_2H]^{1.2} \quad (25)$$

was determined as 2.6×10^2 and $8.7 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ at pH 2 and pH 7, respectively. These values, along with an additional measurement at pH 11, showed a pH dependence in which the rate constant is approximately proportional to $[OH^-]^{1/2}$. Although the rate constant at pH 7 is in reasonable agreement with that determined in this work, i.e., $4.3 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$, the pH dependence of the reaction rate is quite different from that observed here.

The results reported by Hoigne and Bader (1983a), on the other hand, exhibited a pH dependence similar to that observed in this work, i.e., the rate law is consistent with a mechanism for which the reacting species consists of the dissociated formate ion rather than the undissociated formic acid. The rate constant reported by them for the $O_3\text{-HCO}_2^-$ reaction was $140 \text{ M}^{-1} \text{ s}^{-1}$, which is a factor of 28 smaller than that determined in this work. Their smaller value may very well have resulted from the doping of the reaction mixture with propanol, a free radical scavenging material. According to Hoigne and Bader, the presence of such a substance serves to inhibit the O_3 decomposition via free radical pathways. However, it may be somewhat uncertain whether the scavenger itself may enter the reaction pathways and whether certain scavengers may be more efficient than others.

Using the expression given by Eq. (4), the reaction time constants referred to gas phase for O_3 and HCO_2H against reaction (1) are estimated as

$$\tau_{O_3} = [k_4 \left(\frac{K_a}{[H^+] + K_a} \right) H_{O_3} H_{HCO_2H} H_{HCO_2H} LRT]^{-1} \quad (26)$$

and

$$\tau_{\text{HCO}_2\text{H}} = [k_4 \left(\frac{K_a}{[\text{H}^+] + K_a} \right) \text{H}_2\text{O}_3 \text{H}_2\text{CO}_2\text{H} \text{PO}_3 \text{LRT}]^{-1} \quad (27)$$

Using $\text{H}_2\text{CO}_2\text{H} = 230 \text{ M atm}^{-1}$ (at pH 4, Gaffney and Senum, 1984) and the following conditions: pH = 4, $L = 10^{-6}$, $\text{PO}_3 = 50 \text{ ppb}$, $\text{PHCO}_2\text{H} = 10 \text{ ppb}$, and $T = 25^\circ\text{C}$, O_3 and HCO_2H are calculated to be $2.4 \times 10^5 \text{ hr}$ and $4.8 \times 10^4 \text{ hr}$, respectively. These values are clearly too long to make this reaction significant.

$\text{O}_3\text{-H}_2\text{CO}$ Reaction

Kuo and Wen (1977) studied this reaction and reported an O_3 decomposition rate which is dependent on $[\text{H}_2\text{CO}]$ to the first power and on $[\text{O}_3]$ to the $1/2$ power. The rate constant defined by

$$-\frac{d[\text{O}_3]}{dt} = k_{28}[\text{O}_3]^{1/2}[\text{H}_2\text{CO}] \quad (28)$$

changed from $2.2 \times 10^{-2} \text{ M}^{-1/2} \text{ s}^{-1}$ to $0.53 \text{ M}^{-1/2} \text{ s}^{-1}$ as pH increased from 2.7 to 7.2. On the other hand, Hoigne and Bader (1983b) suggested an overall second-order reaction, first order in each reagent, for which the rate constant was determined to be $0.1 \pm 0.03 \text{ M}^{-1} \text{ s}^{-1}$ at pH 2. In contrast to both of these rate laws, we have suggested a tentative rate law, namely, that given by Eq. (10) where the power dependences of the rate on $[\text{O}_3]$ and on $[\text{H}_2\text{CO}]$ are assigned as one-half. The discrepancies observed in these studies may result from the different reaction conditions employed. For example, Hoigne and Bader used a H_2CO concentration in the range of $6\text{-}70 \times 10^{-2} \text{ M}$, which is higher than the concentrations used in this study by 3 to 4 orders of magnitude. Since it is known that some aliphatic alcohols can serve as free radical scavengers, the hydrolyzed formaldehyde, i.e.,

dihydroxymethane, might efficiently block the free radical pathways of this reaction and diminish the overall observed rate.

The fact that different kinetic behavior is observed for varied reaction conditions suggests that these observed rate laws may at best be applied to those conditions used and should not be extrapolated to conditions which are vastly different. Since the conditions employed in this study involved the lowest concentrations for both O_3 and H_2CO among these studies, the kinetic information obtained in this study may be used to estimate the reaction time constants of atmospheric O_3 and H_2CO against the aqueous O_3 -HCHO reaction. For the following conditions, namely, pH = neutral, $P_{H_2CO} = 10$ ppb, $P_{O_3} = 50$ ppb, and $L = 10^{-6}$, these reaction time constants are found to be 6×10^6 hrs or longer. Again, it may be concluded that the aqueous-phase reaction of O_3 and H_2CO affects very little the atmospheric life times of these species.

O_3 - C_2H_4 Reaction

Due to the low solubility of C_2H_4 , this reaction was studied using a continuous flow technique with the gas-phase reagent concentrations maintained constant throughout the reaction course. The second-order aqueous-phase rate constant determined from the peroxide formation rate, i.e., $3 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$, was found to be of similar magnitude to those determined for some substituted ethylenes such as 1-hexene-4-ol ($2 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$) and styrene ($3 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$) studied as homogeneous aqueous-phase reactions (Hoigne and Bader, 1983b). It is interesting to note that the substituent effect observed for the gas-phase O_3 -olefin reactions seems to be absent in the aqueous phase.

The rate of depletion of O_3 and C_2H_4 from the gas phase due to aqueous O_3 - C_2H_4 reaction may be estimated from

$$\tau = (k_{H_2O_3} C_{C_2H_4} LRT p)^{-1} \quad (29)$$

For $p_{C_2H_4} = 20$ ppb, $p_{O_3} = 50$ ppb, and $L = 10^{-6}$, the values of τ s are found to be in the order of 1×10^7 hr. Evidently, these long reaction time constants would have minor consequences on the atmospheric residence times of either O_3 or C_2H_4 .

The aqueous-phase rate of peroxide production from this reaction is estimated to be $6 \times 10^{-11} \text{ M hr}^{-1}$ for $p_{O_3} = 50$ ppb and $p_{C_2H_4} = 20$ ppb. It should be noted that the O_3 -olefin reaction will not constitute an important source of aqueous-phase peroxide unless the condition $(Hp)_{\text{olefin}} > 1 \times 10^{-4} \text{ M}$ is met. The current knowledge, however, indicates that such a species is not to be found (NSF, 1977).


Aqueous-Phase PAN Reactions

The contribution of aqueous-phase PAN reactions to peroxide production in the atmosphere can be estimated from the upper limits assigned to reactions (15) to (17). For a typical ambient condition where $p_{O_3} = 50$ ppb and $p_{\text{PAN}} = 2$ ppb, the upper limits of the rate of peroxide formation are evaluated as 4×10^{-9} , 1×10^{-6} , and $4 \times 10^{-11} \text{ M hr}^{-1}$ for PAN hydrolysis, O_2 -PAN, and O_3 -PAN reactions, respectively. The atmospheric residence times of PAN against hydrolysis, O_2 -PAN, and O_3 -PAN reactions under the same conditions and $L=10^{-6}$ are found to be 2×10^4 , 1×10^2 , and 7×10^7 hr, respectively. Since these time constants are all significantly longer than the

typical cloud lifetime of ~1 hr (Pruppacher and Klett, 1978) and also longer than that for the gas-phase PAN-NO reaction, it is concluded that the atmospheric lifetime of PAN is not affected by these reactions.

O₃-NO₂-H₂CO Reactions

NO₃ has been examined extensively as an intermediate of the nighttime chemical transformations of nitrogen oxides (Winer et al., 1984). This species is suggested to be not only the source of HNO₃ but also a free radical initiator (Calvert and Stockwell, 1983), even for cloudwater droplets during the daytime (Thompson, 1983). According to the studies made in this work, no peroxide was found to be produced in the NO₂-O₃-H₂O reaction system regardless of whether H₂CO was present. In the case of NO₃ reacting with water the upper limit for the reaction



can be estimated by equating the assumed rate expression

$$\text{Rate} = \frac{d[\text{H}_2\text{O}_2]}{dt} = k_{30}[\text{NO}_3] \quad (31)$$

to the maximum possible rate estimated from the limit of detection of the HRPFF method, i.e.,

$$k_{30}[\text{NO}_3] \lesssim 4 \times 10^{-11} \text{ M s}^{-1} \quad (32)$$

It should be noted that the presence of high concentrations of O₃ (43 ppm) in these experiments might affect the aqueous-phase concentrations of H₂O₂ as it has been shown that their reaction can be rapid. From the rate

constants determined for the aqueous-phase $O_3-H_2O_2$ reaction the characteristic reaction times of H_2O_2 for $[O_3] = 4.7 \times 10^{-7} M$ are 13 min and 930 min at pH = neutral and 2, respectively. Although the $O_3-H_2O_2$ reaction is unimportant at pH 2, it may keep the $[H_2O_2]$ from building up at higher pH. If it is assumed that a steady-state concentration of H_2O_2 is established from the competing actions of $O_3-H_2O_2$ reaction and reaction (30), we may observe the following relationship, i.e.,

$$\frac{k_{30} [NO_3]}{k_{21}' [O_3]} \leq 1 \times 10^{-7} M \quad (33)$$

For the conditions applied in this work, the inequality, $k_{30} H_{NO_3} P_{NO_3} \leq 1.2 \times 10^{-10} M s^{-1}$ is obtained. Using the previously calculated value for P_{NO_3} , the product $k_{30} H_{NO_3}$ is estimated to be $4 \times 10^{-3} M atm^{-1} s^{-1}$ or smaller. Although this value is approximately three times greater than that calculated from Eq. (32), the contribution of reaction (30) to the peroxide production in the cloudwater is still negligible, being only $5 \times 10^{-8} M hr^{-1}$ at $P_{NO_3} = 3$ ppb.

In the examination of the aqueous-phase reaction of NO_3 with H_2CO , the reaction between O_3 and H_2CO may consume the available H_2CO . According to Eq. (10), the rate of H_2CO disappearance due to reaction (5) is calculated to be $1.5 \times 10^{-7} M hr^{-1}$ for $p_{O_3} = 43$ ppm and $[H_2CO] = 1.0 \times 10^{-4} M$. This rate is too small to influence the aqueous-phase concentration of H_2CO .

An upper-limit of the rate of the aqueous-phase reaction



can therefore be estimated from

$$\begin{aligned}
 \frac{d[\text{Peroxide}]}{dt} &\approx k_{34} [\text{NO}_3] [\text{H}_2\text{CO}] \\
 &= k_{34} \text{HNO}_3 \text{PNO}_3 [\text{H}_2\text{CO}] \\
 &\leq 1.7 \times 10^{-11} \text{ M s}^{-1}
 \end{aligned}
 \tag{35}$$


using the limit of detection of HRPFF, i.e., 1×10^{-7} M. The product, $k_{34} \text{HNO}_3$, was calculated to be smaller than $6 \text{ atm}^{-1} \text{ s}^{-1}$. For the condition where $\text{PNO}_3 = 3 \text{ ppb}$ and $\text{PH}_2\text{CO} = 20 \text{ ppb}$, the rate of peroxide formation in the aqueous-phase is estimated to be $8 \times 10^{-9} \text{ M hr}^{-1}$. Again, this rate is found to be unimportant. It should be noted that the NO_3 concentration used here was calculated from the known kinetic information and has not been verified by an analytical method such as a spectrophotometric technique. Although our approach might suffer from interferences such as wall effects or the formation of N_2O_5 , the experimental arrangement nonetheless mimics the bubbler H_2O_2 sampling methodology which manifested the artifact production of peroxide.

Aqueous-Phase H_2O_2 Formation

Although it has been suggested that aqueous-phase reactions of O_3 might be responsible for the in situ production of H_2O_2 in the bubbler experiments, the reactions examined in this work thus far did not provide evidence to support this contention. The maximum rate of peroxide formation of the reactions studied in this work, under typical atmospheric conditions, is found to be smaller than $1 \times 10^{-6} \text{ M hr}^{-1}$. Naturally, the calculations were made for pure water systems and the possible effects of, say, metal ions have not been included. This conclusion may not be at variance with the observations reported by Heikes (1984), although the surface catalyzes O_3

decomposition leading to H_2O_2 production was not observed in our pure systems, where pure gases, instead of ambient air, were used.

One reaction which might produce H_2O_2 in the aqueous phase is the recombination of the dissolved HO_2 and its conjugate base O_2^- , i.e.,



Using the Henry's law coefficient of HO_2 ($2 \times 10^3 \text{ M atm}^{-1}$) and rate constant k_{36} ($1 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$) calculated from a thermochemical cycle (Schwartz, 1984) and pulse-radiolysis studies (Bielski, 1978), respectively, the aqueous-phase rate of H_2O_2 formation can be calculated from

$$\frac{d[\text{H}_2\text{O}_2]}{dt} = \frac{k_{36} K_a}{[\text{H}^+]} [\text{HO}_2] \quad (37)$$

$$= k_{36}' [\text{HO}_2]^2 [\text{PHO}_2]^2$$

where K_a is the acid dissociation constant of HO_2 , being $2.0 \times 10^{-5} \text{ M}$ (Bielski, 1978). Accordingly, the gas-phase reaction time constant of HO_2 against reaction (36) is obtained as

$$\tau_{\text{HO}_2} = (2k_{36}' [\text{HO}_2]^2 [\text{PHO}_2]^2 \text{LRT})^{-1} \quad (38)$$

This quantity, calculated as a function of p_{HO_2} , is shown in Figure 12. These values, calculated under the assumption that the aqueous solution is saturated with the solute HO_2 according to Henry's law, suggest that the aqueous-phase recombination of HO_2 is potentially important as a source for the aqueous-phase H_2O_2 when p_{HO_2} is equal to or greater than 2×10^{-12}

atm. It should be noted that the calculations made here are limited to a single pH, i.e. $\text{pH} = \text{pK}_a$, for simplicity. Additionally, it should be pointed out that a steady-state concentration of HO_2 has been assumed in these calculations. If the consumption rate of HO_2 of reaction (36) is significantly faster than the rate of HO_2 production, then the steady-state assumption may not apply and the rate of H_2O_2 production will be primarily controlled by the source rate of HO_2 .

The aqueous-phase production rate of H_2O_2 from the gas-phase recombination of HO_2 is also shown in Figure 12. Here we used a second-order rate constant of $2.5 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ and have assumed that the incorporation of the H_2O_2 into the liquid phase is instantaneous. The liquid water content is again assumed as 10^{-6} . It is interesting to note that the acquiring of the aqueous-phase H_2O_2 from the reactions of dissolved HO_2 is nearly two orders of magnitude faster than that can be supplied by the concomitant gas-phase recombination reaction of HO_2 . Clearly, laboratory studies have to be carried out to examine this potentially important source of aqueous H_2O_2 .




Figure 12. Contribution of the aqueous-phase H_2O_2 recombination reaction to the generation of H_2O_2 , calculated for $\text{pH} \approx 4.7$. Line A represents the aqueous-phase rate of H_2O_2 production and line B the characteristic reaction time for gaseous H_2O_2 at $L = 10^{-6}$. Line C indicates the contribution to aqueous H_2O_2 from the gas-phase recombination of H_2O_2 ; H_2O_2 produced is assumed to be rapidly incorporated into the liquid water ($L = 10^{-6}$)

SUMMARY

Aqueous-phase reactions of O_3 with a number of important atmospheric components have been examined. Reaction kinetics and product analysis for hydrogen peroxide and organic peroxides were determined for the following reaction systems: (1) $O_3 - H_2O$, (2) $O_3 - H_2O_2$, (3) $O_3 - HCO_2H$, (4) $O_3 - H_2CO$, (5) $O_3 - C_2H_4$, (6) $O_3 - PAN$, (7) $O_3 - NO_2$, and (8) $O_3 - NO_2 - H_2CO$. The results obtained are listed in Table 3. Using these values calculations were made to estimate the atmospheric importance of these reactions to the removal of O_3 and the reagent species, as well as the production of peroxides in the liquid-phase. For typical atmospheric conditions ($p_{O_3} = 50$ ppb, $p_{H_2CO} \approx p_{HCO_2H} \approx p_{C_2H_4} = 20$ ppb, $p_{PAN} < 2$ ppb, $p_{NO_2} < 2$ ppb, $L = 10^{-6}$, and $pH = 2 \sim 6$) it is concluded that none of these aqueous-phase reactions is important as a source of aqueous-phase peroxides or a sink for gaseous O_3 and its corresponding reagents.

Table 3
Summary of the Reaction Kinetics and Product Analysis
of Some Aqueous-Phase O_3 Reactions ($22 \pm 2^\circ C$)

Reaction System	Reaction Condition	$-\frac{d[O_3]}{dt}$	Yield or $\frac{d[\text{Peroxide}]}{dt}$	Rate Constant
$O_3 - H_2O$	$[O_3] \leq 2 \times 10^{-5} M$	$k[O_3]$	$\leq 0.5\%$	$2.1 \times 10^{-4} s^{-1}$, pH ~6 $1.4 \times 10^{-4} s^{-1}$, pH = 2
$O_3 - H_2O_2$	$[O_3] \leq 2 \times 10^{-5} M$ $[H_2O_2] = 8-64 \times 10^{-6} M$	$k [O_3][H_2O_2]$		$2.6 \times 10^3 M^{-1} s^{-1}$, pH ~6 $38 M^{-1} s^{-1}$, pH = 4
$O_3 - HCO_2H$	$[O_3] \approx [HCO_2H]$ $= 1 \times 10^{-5} M$	$k [O_3] [HCO_2^-]$	$\leq 0.5\%$	$4.3 \times 10^3 M^{-1} s^{-1}$ pH = 4.6
$O_3 - H_2CO$	$[O_3] \leq 2 \times 10^{-5} M$ $[H_2CO] = 4-17 \times 10^{-5} M$	$k [O_3]^{1/2} [H_2CO]^{1/2}$	$\leq 2\%$	$1.2 \times 10^{-3} s^{-1}$
$O_3 - C_2H_4$	$P_{O_3} = 0.5 - 1.0 \text{ ppm}$ $P_{C_2H_4} = 7-28 \text{ ppm}$		$k [O_3][C_2H_4]$	$3.0 \times 10^5 M^{-1} s^{-1}$ pH 3-6
$O_3 - PAN$	$P_{PAN} = 100 \text{ ppb}$ $P_{O_3} = 1.0 \text{ ppm}$		$k [PAN] [O_3]$	$\leq 3 \times 10^3 M^{-1} s^{-1}$
$O_3 - NO_2$	$P_{O_3} = 43 \text{ ppm}$ $P_{NO_2} = 40 \text{ ppb}$		$k_{NO_3} P_{NO_3}$	$k_{NO_3} \leq 4 \times 10^{-3} M \text{ atm}^{-1} s^{-1}$
$O_3 - NO_2 - H_2CO$	$[H_2CO] = 1 \times 10^{-4} M$		$k_{HNO_3} [H_2CO] P_{NO_3}$	$k_{HNO_3} \leq 6 \text{ atm}^{-1} s^{-1}$

REFERENCES

- Baulch, D.L., Cox, R.A., Crutzen, P.J., Hampson, Jr., R.F., Kerr, J.A., Troe, J. and Watson, R.T., Evaluated kinetic and photochemical data for atmospheric chemistry. *J. Phys. Chem. Ref. Data* 11, 327-496 (1982).
- Bielski, B.H.J., Reevaluation of the spectra and kinetic properties of HO₂ and O₂⁻ free radicals. *Photochem. Photobiol.* 28, 645 (1978).
- Bühler, R.E., Staehelin, J., and Hoigne, J., Ozone decomposition in water studied by pulse radiolysis. 1. HO₂/O₂⁻ and HO₃/O₃⁻ as intermediates. *J. Phys. Chem.* 88, 2560 (1984).
- Calvert, J.G. and Stockwell, W.R., Acid generation in the troposphere by gas-phase chemistry. *Environ. Sci. Technol.* 17, 428A (1983).
- Danckwerts, P.V., Gas-liquid Reactions, McGraw-Hill Book Company, New York, (1970).
- Gaffney, J.S. and Senum, G.I., Peroxides, peracids, aldehydes and PANs and their links to natural and anthropogenic organic sources. In Proceedings of the "Conference on Gas-Liquid Chemistry of Natural Waters", Brookhaven National Laboratory, Upton, New York, April, 1984.
- Gaffney, J.S., Fajer, R., and Senum, G.I., An improved procedure for high purity gaseous peroxyacyl nitrate production: Use of heavy liquid solvents. *Atmos. Environ.* 18, 215 (1984).
- Gilbert, E., Reaction of ozone with trans-trans muconic acid in aqueous solution. *Water Research* 14, 1637 (1980).
- Grosjean, D., Formaldehyde and other carbonyls in Los Angeles ambient air. *Environ. Sci. Tech.* 16, 254 (1982).
- Guilbault, G.G., Brignac, P.J., Jr., and Juneau, M., New substrates for the fluorometric determination of oxidative enzymes. *Anal. Chem.* 40, 1256 (1968).

- Gurol, M.D. and Singer, P.C., Kinetics of ozone decomposition: a dynamic approach. *Environ. Sci. Technol.* 16, 377 (1982).
- Heikes, B., Aqueous H_2O_2 production from O_3 in glass impingers. *Atmos. Environ.* 18, 1433 (1984).
- Heikes, B.G. and Thompson, A.M., Effects of heterogeneous processes on NO_3 , HONO, and HNO_3 chemistry in the troposphere. *J. Geophys. Res.* 88, 10,883 (1983).
- Heikes, B.G., Lazrus, A.L., Kok, G.L., Kunen, S.M., Gandrud, B.W., Gittlin, S.N., and Sperry, P.D., Evidence for aqueous phase hydrogen peroxide synthesis in the troposphere. *J. Geophys. Res.* 87, 3045 (1982).
- Hoigne, J. and Bader, H. Rate constants of reactions of ozone with organic and inorganic compounds in water - I. Nondissociating organic compounds. *Water Res.* 17, 173 (1983b).
- Hoigne, J. and Bader, H., Rate constants of reactions of ozone with organic and inorganic compounds in water-II. Dissociating organic compounds. *Water Res.*, 17, 185 (1983a).
- Hoigne, J. and Bader, H., The role of hydroxyl radical reactions in ozonation processes in aqueous solutions. *Water Research* 10, 377 (1976).
- Keene, W.C., Galloway, J.N., and Holden, J.D., Jr., Measurement of weak organic acidity in precipitation from remote areas of the world. *J. Geophys. Res.* 88, 5122 (1983).
- Kelly, T.J., Daum, P.H., and Schwartz, S.E., Measurements of peroxides in cloudwater and rain. *J. Geophys. Res.*, submitted (1984).
- Kilpatrick, M.L., Herrick, C.C., Kilpatrick, M., The decomposition of ozone in aqueous solution. *J. Amer. Chem. Soc.* 78, 1784 (1956).

- Kosak-Channing, L.F. and Helz, G.R., Solubility of ozone in aqueous solutions of 0-0.6 M ionic strength at 5-30°C. *Environ. Sci. Technol.* 17, 145 (1983).
- Kuo, C.H., and Wen, C.P., Ozonations of formic acid, formaldehyde, and methanol in aqueous solutions. *AIChE Symposium, Series No. 166*, 73, 272 (1977).
- Lazrus, A.L., Kok, G.L., Lind, J.A., and Sperry, P.D., A fluorometric technique for H₂O₂ in precipitation and in cloudwater. *EOS Trans. Am. Geophys. Union* 64, 670 (1983).
- Lee, Y.-N. and Schwartz, S.E., Evaluation of the rate of uptake of nitrogen dioxide by atmospheric water and surface liquid water, *J. Geophys. Res.* 86, 11971 (1981).
- Lee, Y.-N. and Schwartz, S.E., Reaction kinetics of nitrogen dioxide with liquid water at low partial pressures, *J. Phys. Chem.* 85, 840 (1981).
- Lee, Y.-N., Kinetics of some aqueous-phase reactions of peroxyacetyl nitrate. In *Proceedings of "The Conference on Gas-Liquid Chemistry of Natural Waters,"* Brookhaven National Laboratory, Upton, New York, April 1984.
- Martin, L.R., Measurements of sulfate production in natural clouds. *Atmos. Environ.* 17, 1603 (1983).
- National Academy of Sciences, Vapor-Phase Organic Pollutants, Committee on Medical and Biological Effects of Environmental Pollutants, Washington, D.C., 1976.
- Nielsen, T., Hansen, A.M., and Thomsen, E.L., A convenient method for preparation of pure standards of peroxyacetyl nitrate for atmospheric analyses. *Atmos. Environ.* 16, 2447 (1982).

- Niki, H., Maker, P.O., Savage, C.M., and Breitenbach, L.P., Atmospheric ozone-olefin reactions, *Environ. Sci. Technol.* 17, 312A (1983).
- Penkett, S.A., Jones, M.R., Brice, K.A., and Eggleton, A.E.J., The importance of atmospheric ozone and hydrogen peroxide in oxidizing sulfur dioxide in cloud and rainwater. *Atmos. Environ.* 13, 123 (1979).
- Pitts, J.N., Jr., Biermann, H.W., Atkinson, R., and Winer, A.M., Atmospheric implications of simultaneous nighttime measurements of NO_3 radicals and HONO. *Geophys. Res. Letts.* 11, 557 (1984).
- Pruppacher, H.R. and Klett, J.D., Microphysics of Clouds and Precipitation, D. Reidel Publishing Co., Boston, 1978.
- Riddick, J.A. and Bunger, W.B., Techniques of Chemistry, Vol. II: Organic Solvents. Third ed., Wiley-Interscience, New York, 1970.
- Schonbaum, G.R. and Chance, B., Catalase. In The Enzymes, P.D. Boyer, ed., Vol. XIII, pp. 363-408, Academic Press, New York, 1976.
- Schwartz, S.E., Gas- and aqueous-phase chemistry of HO_2 in liquid-water clouds, *J. Geophys. Res.* 89, 11589 (1984).
- Senum, G.I. and Gaffney, J.S., Private communication (1984).
- Sillen, L.G., and Martell, A.E., Stability Constants of Metal-Ion Complexes. The Chemical Society, London, Special Publication No. 17, 1964.
- Sullivan, D.E. and Roth, J.A., Kinetics of ozone self-decomposition in aqueous solution. *AIChE Symposium Series, Water*, 1979, PP. 142-149.
- Tanner, R.L. and Meng, Z., Seasonal variations in ambient atmospheric levels of formaldehyde and acetaldehyde. *Environ. Sci. Technol.* 18, 723 (1984).

- Taube, H. and Bray, W.C., Chain reactions in aqueous solutions containing ozone, hydrogen peroxide, and acid. J. Amer. Chem. Soc. 62, 3357 (1940).
- ten Brink, H.M., Kelly, T. J., Lee, Y.-N., and Schwartz, S.E., Attempted measurement of gaseous H_2O_2 in the ambient atmosphere. In Proceedings of the Third European Symposium on the Physico-Chemical Behavior of Atmospheric Pollutants, Varese, Italy, April 11-13, 1984.
- Thompson, A.M. Night-time oxidant chemistry as a source of free radicals in cloudwater. Presented at the 5th CACGP Conference: Symposium on Tropospheric Chemistry, Oxford, UK, Aug. 28-Sept. 3, 1983.
- Wilhelm, E., R. Battino, and R.J. Willcock, Low pressure solubility of gases in liquid water. Chem. Rev. 77, 219 (1977).
- Winer, A.M., Atkinson, R., and Pitts, J.M., Jr., Gaseous nitrate radical: possible nighttime atmospheric sink for biogenic organic compounds. Science 224, 156 (1984).

Page Intentionally Blank