


Oxygen-18 Study of SO<sub>2</sub> Oxidation in  
Rainwater by Peroxides

Argonne National Lab., IL

Prepared for

Environmental Protection Agency  
Research Triangle Park, NC

Jul 86



U.S. Department of Commerce  
National Technical Information Service

**NTIS**

| TECHNICAL REPORT DATA<br>(Please read Instructions on the reverse before completing)   |   |  |
|--|---|--|
| 1. REPORT NO.<br>EPA/600/3-86/035  | 2.  | 3. RECIPIENT'S ACCESSION NO.<br>PB86-2167517AS |
| 4. TITLE AND SUBTITLE<br>OXYGEN-18 STUDY OF SO <sub>2</sub> OXIDATION IN<br>RAINWATER BY PEROXIDES   | 5. REPORT DATE<br>July 1986                                   | 6. PERFORMING ORGANIZATION CODE                |
|  | 8. PERFORMING ORGANIZATION REPORT NO.                         |  |
| 7. AUTHOR(S)<br>B. D. Holt and R. Kumar  | 10. PROGRAM ELEMENT NO.<br>CCVNI/A/02-3058 (FY-86)            |  |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br>Chemical Technology Division<br>Argonne National Laboratory<br>Argonne, Illinois 60439  | 11. CONTRACT/GRANT NO.<br>IAG DW 89930060                     |  |
|  | 13. TYPE OF REPORT AND PERIOD COVERED<br>Final (10/82 - 5/86) |  |
| 12. SPONSORING AGENCY NAME AND ADDRESS<br>Atmospheric Sciences Research Laboratory-RTP, NC<br>Office of Research and Development<br>U.S. Environmental Protection Agency<br>Research Triangle Park, North Carolina 27711   | 14. SPONSORING AGENCY CODE<br>EPA/600/09                      |  |
|  | 15. SUPPLEMENTARY NOTES                                       |  |
| 16. ABSTRACT<br><br>A new analytical method was developed for the determination of oxygen isotope ratios in peroxides in rainwater. In the method, rainwater samples were quantitatively degassed of dissolved air by a combined treatment of evacuation, ultrasonic agitation, and helium sparging (VUS), followed by a permanganate oxidation of the dissolved peroxide to O <sub>2</sub> . The O <sub>2</sub> was then quantitatively removed from the rainwater by the VUS treatment and converted to CO <sub>2</sub> for mass spectrometric analysis. Using this method, 14 rainwater samples collected at four sites (Argonne, IL, Research Triangle Park, NC, Whiteface Mountain, NY and Dearborn, MI) were analyzed to determine the fraction of the sulfate in the samples that was produced as a result of the aqueous-phase reaction of dissolved SO <sub>2</sub> with hydrogen peroxide. It was concluded that 40% or more of the sulfate in the samples was formed by peroxide oxidation. |   |  |
| 17. KEY WORDS AND DOCUMENT ANALYSIS  |   |  |
| a. DESCRIPTORS   | b. IDENTIFIERS/OPEN ENDED TERMS                               | c. COSATI Field/Group                          |
|  |   |  |
| 18. DISTRIBUTION STATEMENT<br><br>RELEASE TO PUBLIC  | 19. SECURITY CLASS (This Report)<br>UNCLASSIFIED              | 21. NO. OF PAGES<br>43                         |
|  | 20. SECURITY CLASS (This page)<br>UNCLASSIFIED                | 22. PRICE<br>\$ 9.95                           |

## DISCLAIMER

The information in this document has been funded by the United States Environmental Protection Agency under Interagency Agreement DW 930060-01-0 to Argonne National Laboratory. It has been subject to the Agency's peer and administrative review, and it has been approved for publication as an EPA document. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

## ABSTRACT

Oxygen isotope ratio measurements were used to examine the importance of peroxide oxidation of  $\text{SO}_2$  to sulfate in the atmosphere. A new analytical method was developed for the determination of oxygen isotope ratios in peroxides (concentrations in the ppb range) in rainwater. According to the method, 20-L samples were quantitatively degassed of dissolved air by a combined treatment of evacuation, ultrasonic agitation, and helium sparging (VUS), followed by permanganate oxidation of the dissolved peroxide to  $\text{O}_2$ ; the  $\text{O}_2$  was then quantitatively removed from the rainwater by the VUS treatment and converted to  $\text{CO}_2$  for mass spectrometric analysis. Stock solutions of  $\text{H}_2\text{O}_2$  of various  $^{18}\text{O}$  enrichments were prepared by a high-voltage discharge method; and, by using these solutions to oxidize  $\text{SO}_2$  to  $\text{SO}_4^{2-}$ , the following isotopic relationship was established:

$$\delta^{18}\text{O}_{\text{SO}_4^{2-}} = 0.57 \delta^{18}\text{O}_{\text{SO}_2} + 0.43 \delta^{18}\text{O}_{\text{H}_2\text{O}} + 8.4\text{‰}.$$

This relationship was used to calculate the  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  formed by peroxide oxidation and a similar previously established relationship was used to calculate the  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  formed by metal-catalyzed aqueous oxidation. These sets of calculated values were compared to the measured values for evaluation of the estimated fraction of sulfate in rainwater that was formed by peroxide oxidation, assuming that metal (or carbon) catalysis was the only other major sulfate formation mechanism. It was concluded that  $\sim 40\%$  or more of the sulfate in summer rains in the northeastern U. S. was formed by peroxide oxidation.

## CONTENTS

|  |      |
|--|------|
| ABSTRACT . . . . .   | iii  |
| TABLES . . . . .   | vi   |
| FIGURES . . . . .  | vii  |
| ACKNOWLEDGMENT . . . . .   | viii |
| 1. INTRODUCTION . . . . .  | 1    |
| 2. EXPERIMENTAL . . . . .  | 3    |
| Development of Analytical Method . . . . .   | 3    |
| Conversion of O <sub>2</sub> to CO <sub>2</sub> . . . . .  | 7    |
| Removal of Dissolved O <sub>2</sub> from Water . . . . .   | 9    |
| Oxidation of H <sub>2</sub> O <sub>2</sub> to O <sub>2</sub> in Water . . . . .                  | 13   |
| Synthesis of <sup>18</sup> O-enriched H <sub>2</sub> O <sub>2</sub> . . . . .                    | 13   |
| Collection of Rain Samples . . . . .   | 14   |
| 3. RESULTS . . . . .   | 16   |
| Recovery of Added H <sub>2</sub> O <sub>2</sub> . . . . .  | 16   |
| Test for Isotopic Interference . . . . .   | 16   |
| Effects of Autodecomposition of H <sub>2</sub> O <sub>2</sub> . . . . .                          | 16   |
| Effects of KMnO <sub>4</sub> Decomposition . . . . .   | 18   |
| Procedure Blank . . . . .  | 18   |
| Isotopic Relationships . . . . .   | 18   |
| $\delta^{18}\text{O}$ and Concentrations of H <sub>2</sub> O <sub>2</sub> in Rainwater . . . . . | 22   |
| 4. DISCUSSION . . . . .  | 26   |
| $\delta^{18}\text{O}$ of H <sub>2</sub> O <sub>2</sub> in Rainwater . . . . .                    | 26   |
| $\delta^{18}\text{O}$ of Atmospheric Sulfates: Measured vs. Calculated . . . . .                 | 27   |
| Fraction of Atmospheric Sulfate Formed by Peroxide Oxidation . . . . .                           | 29   |
| 5. SUMMARY . . . . .   | 32   |
| REFERENCES . . . . .   | 33   |

TABLES

| <u>Number</u>   | <u>Page</u> |
|---|-------------|
| 1. Concentration and isotopic results for rainwater collected at Argonne National Laboratory (ANL), Dearborn, MI (DEA), Whiteface Mountain, NY (WFM), and Research Triangle Park, NC (RTP), May-September, 1985 . . . . . | 23          |

## FIGURES

| <u>Number</u>  | <u>Page</u> |
|--|-------------|
| 1. Four-step method for conversion of oxygen in H <sub>2</sub> O <sub>2</sub> (ppb range in water) to CO <sub>2</sub> for mass spectrometric analysis . . . . .        | 4           |
| 2. Analytical train . . . . .  | 5           |
| 3. Reaction chamber . . . . .  | 6           |
| 4. Recovery of O <sub>2</sub> as CO <sub>2</sub> after reaction with charcoal in contact with platinum at various temperatures . . . . .                               | 8           |
| 5. Recovery of CO <sub>2</sub> after exposure to charcoal and platinum at various temperatures . . . . .   | 10          |
| 6. Removal of O <sub>2</sub> from water by various degassing techniques . . . . .  | 12          |
| 7. Isotopic effects of water solvent on the $\delta^{18}\text{O}$ of CO <sub>2</sub> originating from H <sub>2</sub> O <sub>2</sub> . . . . .                          | 17          |
| 8. Dependence of $\delta^{16}\text{O}_{\text{H}_2\text{O}_2}$ on $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ in a high-voltage discharge process of formation . . . . . | 19          |
| 9. Dependence of $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$ on $\delta^{18}\text{O}_{\text{H}_2\text{O}_2}$ in oxidation of SO <sub>2</sub> to sulfate . . . . .         | 21          |
| 10. Measured isotopic data for H <sub>2</sub> O <sub>2</sub> , SO <sub>4</sub> <sup>2-</sup> , and H <sub>2</sub> O in the samples of rainwater . . . . .              | 24          |
| 11. $\delta^{18}\text{O}$ of sulfate in rainwater: calculated compared to measured . . . . .   | 28          |
| 12. Percent of sulfate formed by peroxide oxidation, assuming negligible primary sulfates . . . . .  | 30          |

## ACKNOWLEDGMENT

The authors wish to acknowledge that some of the laboratory studies involving the determination of isotopic relationships using solutions of known  $^{18}\text{O}/^{16}\text{O}$  ratios were conducted under a related program for the U.S. Department of Energy's Office of Energy Research, Basic Energy Sciences, Division of Materials Science. Also, the authors are grateful to Dr. Mark Dubois, Whiteface Mountain Field Station, Atmospheric Sciences Research Center, Wilmington, NY; to Dr. Cary Eaton, Research Triangle Institute, Research Triangle Park, NC; and to Dr. William Pierson and Ms. Wanda Brachaczek, Ford Research Engineering Staff, Dearborn, MI, for collection and shipment of rainwater samples to Argonne for analysis.



## SECTION 1

### INTRODUCTION

Considerable interest has been shown in the possibility of using oxygen isotopy to elucidate the role of  $\text{H}_2\text{O}_2$  in the oxidation of  $\text{SO}_2$  to sulfates in the atmosphere. The potential importance of  $\text{H}_2\text{O}_2$  lies in the view that aqueous-phase oxidation of  $\text{SO}_2$  to  $\text{H}_2\text{SO}_4$  probably accounts for a major fraction of the observed  $\text{SO}_4^{2-}$  in the precipitation occurring in the northeastern United States. The key reactants responsible for this oxidation are not well known, although a large number of possible catalysts and oxidants exist in the atmosphere, including carbon, transition metal ions, hydroxyl and organic free radicals, hydrogen peroxide (including organic peroxides) and ozone (Penkett *et al.*, 1979). It has been postulated that the atmospheric oxidation of  $\text{SO}_2$  is limited by the availability of these oxidants and catalysts, rather than by the availability of  $\text{SO}_2$  itself. This is the so-called phenomenon of non-linearity between the concentration (and, by implication, the emission) of  $\text{SO}_2$  in the atmosphere and the conversion of this  $\text{SO}_2$  to  $\text{H}_2\text{SO}_4$ . If such is indeed the case, it is important to determine if peroxides have a pivotal role in acid formation and deposition from the atmosphere [oxidation by ozone decreases in significance with decreasing pH of the rainwater].

Oxygen isotopic studies are useful in distinguishing between the different oxidation mechanisms effective in the atmosphere. Laboratory simulation of several different atmospheric reaction sequences has shown that the oxygen isotope ratio in the product  $\text{SO}_4^{2-}$  is uniquely related to the reaction pathway followed in its formation (Holt *et al.*, 1982). It may therefore be possible to determine if the atmospheric hydrogen peroxide is responsible for significant oxidation of  $\text{SO}_2$  to sulfate, and if it is this oxidant that limits the aqueous-phase formation of sulfuric acid. The results of these studies could have significant implications for energy technology, particularly if they indicate that, because of the non-linearity in  $\text{SO}_2$  conversion discussed above, it may be more important to reduce the

ambient concentrations of peroxides than of  $\text{SO}_2$ .

In the 1981 JASON Committee Report to the U.S. Department of Energy (Chamberlain *et al.*, 1981), some recommendations for further research were based, at least in part, on our earlier work on  $\text{SO}_2$  oxidation by  $\text{H}_2\text{O}_2$  (Holt *et al.*, 1981). According to our earlier work, the  $\delta^{18}\text{O}$  [deviation in parts per thousand ( $\text{‰}$ ) of the  $^{18}\text{O}/^{16}\text{O}$  ratio of the sample from that of the standard reference material, Standard Mean Ocean Water (SMOW)] of sulfates produced by  $\text{H}_2\text{O}_2$  oxidation were significantly lower than the  $\delta^{18}\text{O}$  of sulfates found in rainwater. However, the  $\delta^{18}\text{O}$  of the reagent-grade  $\text{H}_2\text{O}_2$  used in those experiments was not known. The results suggested the need for isotopic analysis of  $\text{H}_2\text{O}_2$  in dilute solutions, and for a methodology whereby the  $\delta^{18}\text{O}$  values of  $\text{H}_2\text{O}_2$ ,  $\text{H}_2\text{O}$ , and  $\text{SO}_4^{2-}$  in rainwater could be compared, in order to assess the importance of  $\text{H}_2\text{O}_2$  in the formation of sulfate-constituted acid rain.

The plan of this investigation was to develop a method for the determination of the  $\delta^{18}\text{O}$  of  $\text{H}_2\text{O}_2$  in dilute aqueous solutions (simulating rainwater); to prepare solutions of  $\text{H}_2\text{O}_2$  of various  $^{18}\text{O}$  enrichments; to use the freshly prepared solutions of  $\text{H}_2\text{O}_2$  to oxidize  $\text{SO}_2$  to  $\text{SO}_4^{2-}$  for evaluation of the relationship between  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  and  $\delta^{18}\text{O}_{\text{H}_2\text{O}_2}$ ; and to apply this relationship to the measured  $\delta^{18}\text{O}_{\text{H}_2\text{O}_2}$ ,  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$ , and  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  in precipitation water, for assessment of the importance of  $\text{H}_2\text{O}_2$  in the atmospheric transformation of  $\text{SO}_2$  to sulfate.

## SECTION 2

### EXPERIMENTAL

#### Development of Analytical Method

No suitable analytical method was available for the quantitative extraction of the oxygen in  $\text{H}_2\text{O}_2$ , (dissolved in water, ppb range), for isotopic analysis. Figure 1 shows a 4-step method which we developed for this purpose. It consists of the removal of dissolved  $\text{O}_2$  from 20-liter samples of water by a combination of evacuation, ultrasonic agitation, and sparging with helium; oxidation of the dissolved  $\text{H}_2\text{O}_2$  to  $\text{O}_2$  in the water with  $\text{KMnO}_4$ ; removal of the newly formed  $\text{O}_2$  from the water in a carrier-gas stream; and conversion of the  $\text{O}_2$  to  $\text{CO}_2$  by reaction with platinum-catalyzed carbon at  $600^\circ\text{C}$ . The  $\text{CO}_2$  is then mass spectrometrically analyzed to determine its  $\delta^{18}\text{O}$  (identical to that of the oxygen in the original  $\text{H}_2\text{O}_2$ ).

**Analytical Train.** Diagrams of the apparatus are shown in Figs. 2 and 3. Fig. 2 shows the all-glass analytical train. It consists of a bed of molecular sieve at  $-196^\circ\text{C}$  for the removal of traces of  $\text{O}_2$  from helium carrier gas, a gas pipet for the injection of known amounts of  $\text{O}_2$  or  $\text{CO}_2$  into the carrier gas stream during standardization, the water sample chamber detailed in Fig. 3, a 20-mm o.d. cold trap at  $-78^\circ\text{C}$  for the removal of residual water vapor from the gas stream, a cold trap at  $-196^\circ\text{C}$  for the removal of  $\text{CO}_2$  that is scrubbed from the water sample by the helium stream, a bed of activated charcoal (3 g, 8-10 mesh, coconut grade, in a vertically mounted quartz tube, lined with platinum gauze) at  $600^\circ\text{C}$  for the conversion of  $\text{O}_2$  to  $\text{CO}_2$  in the carrier gas stream, another cold trap at  $-196^\circ\text{C}$  for the collection of the newly formed  $\text{CO}_2$  from the gas stream, a capillary open-well mercury manometer for measurement of the  $\text{CO}_2$ , and a gas sample bulb, attached to the train for transfer of the  $\text{CO}_2$  to a mass spectrometer for isotopic analysis. A cold trap at  $-196^\circ\text{C}$ , not shown in the diagram, is in the vacuum manifold to protect the analytical train from vapor contamination from the mechanical vacuum pump.




Figure 1. Four-step method for conversion of oxygen in  $\text{H}_2\text{O}_2$  (ppb range in water) to  $\text{CO}_2$  for mass spectrometric analysis




Figure 2. Analytical train




Figure 3. Reaction chamber

**Reaction Chamber.** The reaction chamber assembly, Fig. 3, consists of a 20-L round bottom flask, a rotatable side-arm tube for the addition of crystalline  $\text{KMnO}_4$ , a cold water condenser to limit the amount of water vapor swept by the helium stream into the  $-78^\circ\text{C}$  cold trap, three stopcocks for manipulation of the helium stream, and a funnel for the addition of solutions of acid and oxidants (used only during the development of the method). The round-bottom flask was supported by a stainless steel rack in an 83-L stainless-steel tank, interior  $40.6 \times 50.8 \times 40.6$  cm, (Model ATH 1620-24), of an ultrasonic cleaning system (Model EMa 50-24) manufactured by Branson Cleaning Equipment Co., Shelton, CT. The water in the ultrasonic tank was maintained at about  $10^\circ\text{C}$  by circulation through a refrigeration unit.

The components of the glass analytical train were connected by No. 18 ball joints, sealed with solidified black wax. The hollow plug of stopcock 22 was modified to provide extra volume in the capillary manometer (Holt, 1955).

Conversion of  $\text{O}_2$  to  $\text{CO}_2$ . The first part of the method to be developed was the conversion of  $\text{O}_2$  to  $\text{CO}_2$  in the carrier gas stream (4th box in the flow sheet, Fig. 1). The procedure is a modification of an established vacuum technique for the conversion of oxygen in air to  $\text{CO}_2$  for mass spectrometric analysis. By the vacuum procedure, the air is circulated at low pressure over graphite and platinum at  $600^\circ\text{C}$  and the resulting  $\text{CO}_2$  is cryogenically removed from the closed system (Horibe *et al.*, 1973; Holt, 1977). By our new method, in which helium is conducted through a bed of 8–10 mesh activated coconut charcoal at  $600^\circ\text{C}$  and in contact with platinum gauze, the yield of  $\text{CO}_2$  was found to be affected by both the bed temperature and the axial temperature gradient. With a suitably long furnace to obtain an essentially isothermal bed, the recovery of  $\text{O}_2$  (as  $\text{CO}_2$ ) as a function of bed temperature is shown in Fig. 4. The data show that the optimum bed temperature for maximum yield of  $\text{CO}_2$  was  $\sim 600^\circ\text{C}$ , the same as that used by Horibe




Figure 4. Recovery of O<sub>2</sub> as CO<sub>2</sub> after reaction with charcoal in contact with platinum at various temperatures



*et al.*, (1973) and Holt (1977) in vacuum-line applications.

The data in Fig. 4 also indicate that the maximum recovery of the added O<sub>2</sub>, measured as CO<sub>2</sub>, was ~97%. Recoveries higher than 97% were apparently unattainable because of competing reactions. At temperatures lower than 600°C, the formation of CO<sub>2</sub> was decreased due to the incomplete reaction,



and at higher temperatures, the yield of CO<sub>2</sub> was decreased by the competing reaction,



The effect of reaction (2) was demonstrated by injecting CO<sub>2</sub> into the carrier gas instead of O<sub>2</sub>. The results, Fig. 5, confirm that about 3% of the CO<sub>2</sub> decomposed at 600°C.

The 97% recovery is adequate for isotopic studies; and, at 600°C, oxygen isotope fractionation within the equilibrated system of CO<sub>2</sub>, CO, unconverted CO<sub>2</sub>, and fixed oxygen on the charcoal is negligible. The demonstrated reliability of the 97% recovery at 600°C allows the applicability of the technique to the quantitative determination of O<sub>2</sub> in gas streams, as well as to isotopic studies.

#### Removal of Dissolved O<sub>2</sub> from Water

After it was experimentally demonstrated that O<sub>2</sub> could be reliably converted to CO<sub>2</sub> in a carrier-gas stream, the technique for the quantitative removal of dissolved oxygen from multi-liter quantities of water was developed (1st and 3rd boxes in Fig. 1). Various combinations of sparging with helium, vacuum pumping, and ultrasonic agitation were tested. The best results were obtained by a combination of all three. (The commonly used degassing method of alternately freezing and thawing the water under vacuum was much too impractical and time-consuming for large samples of water.)




Figure 5. Recovery of CO<sub>2</sub> after exposure to charcoal and platinum at various temperatures

**Helium Sparge.** Attempts were made at first to remove dissolved oxygen from water (saturated with air) by only sparging with helium. Quantitative removal of air oxygen from the water by this technique was intolerably slow. Curve SP-1 in Fig. 6 shows the percent removal of O<sub>2</sub> (estimated to be dissolved in the air-saturated water at room temperature) plotted vs. minutes of sparging. After 50 min of sparging, only ~1/3 of the dissolved O<sub>2</sub> had been removed.

T. G. Holt (1983) cited a comparison of various degassing techniques in Technical Brief No. 101 of Waters Associates, Milford, Massachusetts. The techniques, listed in the order of decreasing effectiveness for the removal of dissolved gases from liquids, were: (1) vacuum pumping combined with ultrasonic agitation, (2) vacuum filtration using a Waters Solvent Clarification Kit, (3) ultrasonic agitation, (4) vacuum pumping, (5) boiling, and (6) sparging.

**Vacuum-Ultrasonic Agitation (VU).** The apparatus was arranged to accommodate treatment of the water sample by a combination of vacuum pumping and ultrasonic agitation. The results of two experiments, VU-1 and VU-2, are plotted in Fig. 6. The rate of removal of dissolved oxygen was found to be greatly enhanced by "rinsing" the water with helium. That is, the evacuated sample chamber was filled with pure helium to atmospheric pressure by allowing the gas to bubble through the water; then it was exhausted through the analytical train. The sequential dissolution and removal of the helium from the water had the effect of "rinsing" the O<sub>2</sub> from the water. Curve VUR-1 in Fig. 6 shows the improved results of a third experiment in which the water sample was rinsed with He at regular intervals throughout the O<sub>2</sub>-removal treatment.

**Vacuum-Ultrasonication-Sparging (VUS).** The improved rate of O<sub>2</sub> removal by frequently rinsing the O<sub>2</sub> from the water with helium suggested that a low-pressure sparge by helium should be combined with the vacuum-ultrasonic agitation treatment. This technique, tested in runs VUS-1, VUS-2, VUS-3, and VUS-4 yielded the best O<sub>2</sub> removal rates.




Figure 6. Removal of  $O_2$  from water by various degassing techniques

The results show removal of ~99% of the dissolved oxygen in 1 h of VUS for 3-L water samples. (20-liter samples were later found to require ~5 h of the treatment for complete air removal.)

#### Oxidation of $H_2O_2$ to $O_2$ in Water

Bromine water was tested as an oxidant of  $H_2O_2$  to  $O_2$  in the degassed water sample. Although it was very effective in converting a known amount of  $H_2O_2$  to  $O_2$ , its use was complicated by its exposure to stopcock grease and by contamination of the analytical train. Curve "VU-1 +  $H_2O_2$  +  $Br_2$ " in Fig. 6 shows the results of one experiment in which a degassed water sample (VU-1) was spiked with  $H_2O_2$  and treated with bromine water, yielding a recovery of about 97% in 50 min of VU treatment.

Potassium permanganate was used in all subsequent experiments. Curve "VUS-2 +  $H_2O_2$  +  $KMnO_4$ " in Fig. 6 shows the results of an experiment made to compare its effectiveness to that of bromine. The recovery of oxygen, added as  $H_2O_2$  to a degassed water sample (see curve VUS-2), was also ~97% after 1 h of vacuum ultrasonic sparging. In earlier experiments with  $KMnO_4$ , it was added as a 0.1N solution through the funnel, Fig. 3; later, the rotatable side arm was made a part of the apparatus to permit the addition of  $KMnO_4$  crystals, with correspondingly lower procedure blanks.

#### Synthesis of $^{18}O$ -enriched $H_2O_2$

Hydrogen peroxides of various  $\delta^{18}O$  were not commercially available. A suitable method of synthesis (Vol'nov *et al.*, 1964) was identified and successfully applied to the laboratory preparations of four stock solutions of  $H_2O_2$  of different  $\delta^{18}O$ . By this method,  $H_2O_2$  is formed by exposure of supplies of water vapor, each differing in  $\delta^{18}O$ , to a high-voltage (~1.4 kV) discharge in ~100 cm of 10-mm o.d. glass tubing between two water-

cooled aluminum electrodes. Some of the HO radicals, formed by the dissociative reaction,



are condensed in a cold trap ( $-196^\circ\text{C}$ ) where they combine to form  $\text{H}_2\text{O}_2$ , leaving the H radicals to combine in formation of  $\text{H}_2$  and be pumped away through the vacuum line.

Other techniques which we experimentally found to give inadequate yields of  $\text{H}_2\text{O}_2$  were conduction of an electric arc across a stream of aerosolized water droplets (Kok, 1982), excitation of water vapor by a radio-frequency silent discharge in a glass chamber (7.5 cm dia  $\times$  20 cm long) in a commercially available plasma cleaner unit, and excitation by a glow-discharge unit (4.8 cm dia  $\times$  70 cm long) that had uncooled aluminum disk electrodes (Jarnagin and Wang, 1958).

#### Collection of Rain Samples

Each rain sampling station consisted of four 1-m<sup>2</sup> plastic skylights, inverted to form funnels and fitted with plastic nipples and hoses to allow the water to flow into two 25-L plastic collection bottles. From each collected sample of  $>20$ -L, a 4-L bottle was filled for subsequent determination of  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  and  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$ , and a 25-L bottle was filled for subsequent determination of  $\delta^{18}\text{O}_{\text{H}_2\text{O}_2}$ . The 4-L sample was treated with 20 mg CuCl to prevent bacterial alteration of the sulfate during storage before analysis; the 25-L sample was treated with 25 ml concentrated  $\text{H}_2\text{SO}_4$  and was refrigerated at  $\sim 5^\circ\text{C}$  to prevent autodecomposition of the  $\text{H}_2\text{O}_2$  during storage before analysis.

During the period June–August 1985, rainwater collection stations, in addition to the one at Argonne, IL, were operated by the Research Engineering Staff, Ford Motor Company, Dearborn, MI; Research Triangle Institute, Research Triangle Park, NC; and Whiteface Mountain Field Station, State University of New York, Albany, NY. Approximately two rain events per month were sampled at each site during the 3-month period.

The rain collectors were installed by GP Engineering, Downers Grove, IL, at each of the three sites. Insulated containers were used for shipment of 25-L bottles of acidified, chilled rainwater by air express to Argonne for analyses.

## SECTION 3

### RESULTS

#### Recovery of added H<sub>2</sub>O<sub>2</sub>

The recovery of reagent grade H<sub>2</sub>O<sub>2</sub>, added to 20 liters of distilled water, was about 96%. This recovery is adequate for determining the oxygen isotope ratio and the concentration of H<sub>2</sub>O<sub>2</sub> in rainwater.

#### Test for Isotopic Interference

Schumb *et al.* (1955) cite references showing that dissolved H<sub>2</sub>O<sub>2</sub> does not exchange oxygen atoms with the solvent water, dissolved molecular oxygen, or oxygenated products. In decomposition of H<sub>2</sub>O<sub>2</sub> by oxidation to molecular oxygen, the O-O bond is not broken and no fractionation occurs. The MnO<sub>4</sub><sup>-</sup> ion does not cause isotopic interference (Cahill and Taube, 1952). However, in decomposition of H<sub>2</sub>O<sub>2</sub> by reduction (not applicable to our experimentation), the bond is severed and fractionation may occur.

To confirm the absence of appreciable isotopic interference in our procedure by oxygen exchange between the H<sub>2</sub>O and either the H<sub>2</sub>O<sub>2</sub> or the O<sub>2</sub>, before, during, or after the oxidation reactions, the oxidation was carried out in the presence of three different water supplies of various δ<sup>18</sup>O. The results in Fig. 7 show that the δ<sup>18</sup>O of the CO<sub>2</sub> product was unaffected by the δ<sup>18</sup>O of the water solvent.

#### Effects of Autodecomposition of H<sub>2</sub>O<sub>2</sub>

Results from two sets of experiments on the autodecomposition of H<sub>2</sub>O<sub>2</sub> (spikes added to 20-L samples of rainwater), over storage times of up to 11 days, showed that if the rainwater is stored unacidified at room temperature, the H<sub>2</sub>O<sub>2</sub> concentration declines rapidly (~30% depletion in 2 days and ~98% in 11 days); and that the δ<sup>18</sup>O of the






Figure 7. Isotopic effects of water solvent on the  $\delta^{18}\text{O}$  of  $\text{CO}_2$  originating from  $\text{H}_2\text{O}_2$

undecomposed  $\text{H}_2\text{O}_2$  undergoes a corresponding increase ( $\sim 5\%$  during the first 5 days). However, if the rainwater is first acidified ( $\sim 20$  ml conc.  $\text{H}_2\text{SO}_4$  in 20-L) and kept cold ( $1-9^\circ\text{C}$ ) during storage, the changes in the concentration and in the  $\delta^{18}\text{O}$  of the  $\text{H}_2\text{O}_2$  are negligible.

#### Effects of $\text{KMnO}_4$ Decomposition

Decomposition of excess  $\text{KMnO}_4$  by organic matter in rainwater was shown to produce only  $\text{CO}_2$ , which is cryogenically removed in the analytical train before the  $\text{H}_2\text{O}_2$ -derived  $\text{O}_2$  is converted to  $\text{CO}_2$ . The reaction between  $\text{KMnO}_4$  and organic matter does not produce  $\text{O}_2$  and is therefore not a source of interference in the method.

#### Procedure Blank

The blank of the analytical procedure was reduced from 6.5 to 3.5  $\mu\text{moles}$  of  $\text{O}_2$  by adding the  $\text{KMnO}_4$  to the 20-L sample as pulverized crystals rather than as a pre-boiled  $\text{KMnO}_4$  solution. (In 20-L of water, 3.5  $\mu\text{moles}$  is  $\sim 3$  ppb.)

#### Isotopic Relationships

High-Voltage Preparation of  $\text{H}_2\text{O}_2$  from  $\text{H}_2\text{O}$ . The dependence of the  $\delta^{18}\text{O}$  of  $\text{H}_2\text{O}_2$  on the  $\delta^{18}\text{O}$  of the  $\text{H}_2\text{O}$  from which it was prepared by the high-voltage discharge method described above is shown in Fig. 8. The equation of the best fit regression curve is

$$\delta^{18}\text{O}_{\text{H}_2\text{O}_2} = 1.03\delta^{18}\text{O}_{\text{H}_2\text{O}} + 29.4\text{‰}. \quad (4)$$

The results indicate that the  $\delta^{18}\text{O}_{\text{H}_2\text{O}_2}$  is controlled directly by the  $\delta^{18}\text{O}$  of the  $\text{H}_2\text{O}$  and that the  $\delta^{18}\text{O}$  of the  $\text{H}_2\text{O}_2$  is substantially higher ( $\sim 29\%$ ) than that of the water from which it was formed. Although the high-voltage process used in these preparations




Figure 8. Dependence of  $\delta^{18}\text{O}_{\text{H}_2\text{O}_2}$  on  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  in a high-voltage discharge process of formation

of  $\text{H}_2\text{O}_2$  may simulate lightning in the atmosphere, the importance of lightning, relative to other sources of  $\text{H}_2\text{O}_2$  in the atmosphere, is open to question (Kok, 1982).

$\text{H}_2\text{O}_2$  Oxidation of  $\text{SO}_2$  to  $\text{SO}_4^{2-}$ . Using the four stock solutions of hydrogen peroxide, each of different  $\delta^{18}\text{O}$ , sulfate solutions of correspondingly different  $\delta^{18}\text{O}$  were prepared by oxidation of  $\text{SO}_2$  which was in isotopic equilibrium with water of constant  $\delta^{18}\text{O} = -7.9\text{‰}$ . In Fig. 9 the  $\delta^{18}\text{O}$  of each resulting sulfate is plotted versus the  $\delta^{18}\text{O}$  of the  $\text{H}_2\text{O}_2$  and the equation of the best-fit regression curve is

$$\delta^{18}\text{O}_{\text{SO}_4^{2-}} = 0.43 \delta^{18}\text{O}_{\text{H}_2\text{O}_2} + 3.5\text{‰}. \quad (5)$$

The regression curve of the previously determined relationship (Holt *et al.*, 1981) between  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  and  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  in aqueous-phase oxidation of  $\text{SO}_2$  by  $\text{H}_2\text{O}_2$  was

$$\delta^{18}\text{O}_{\text{SO}_4^{2-}} = 0.57 \delta^{18}\text{O}_{\text{H}_2\text{O}} - 2.4\text{‰}. \quad (6)$$

Assuming that all significant effects of the  $\delta^{18}\text{O}$  of the  $\text{SO}_2$  on the  $\delta^{18}\text{O}$  of the  $\text{SO}_4^{2-}$  are lost by rapid isotopic exchange between the  $\text{SO}_2$  and the large excess of water, prior to appreciable oxidation (Holt *et al.*, 1981),  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  and  $\delta^{18}\text{O}_{\text{H}_2\text{O}_2}$  remain as the only two complementary variables in the equation for  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$ ; therefore, the comprehensive regression curve for  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  is

$$\delta^{18}\text{O}_{\text{SO}_4^{2-}} = 0.57 \delta^{18}\text{O}_{\text{H}_2\text{O}} + 0.43 \delta^{18}\text{O}_{\text{H}_2\text{O}_2} + C \quad (7)$$

$$= \sim \frac{3}{5} \delta^{18}\text{O}_{\text{H}_2\text{O}} + \sim \frac{2}{5} \delta^{18}\text{O}_{\text{H}_2\text{O}_2} + C. \quad (8)$$

The constant,  $C$ , was evaluated at  $8.4\text{‰}$  from the data given in Fig. 9 by substituting the corresponding measured values for  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  and  $\delta^{18}\text{O}_{\text{H}_2\text{O}_2}$ , and  $-7.9\text{‰}$  for  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ .

The comprehensive equation then becomes

$$\delta^{18}\text{O}_{\text{SO}_4^{2-}} = 0.57 \delta^{18}\text{O}_{\text{H}_2\text{O}} + 0.43 \delta^{18}\text{O}_{\text{H}_2\text{O}_2} + 8.4\text{‰} \quad (9)$$




Figure 9. Dependence of  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  on  $\delta^{18}\text{O}_{\text{H}_2\text{O}_2}$  in oxidation of  $\text{SO}_2$  to  $\text{SO}_4^{2-}$

and can be used to calculate  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  from  $\delta^{18}\text{O}_{\text{H}_2\text{O}_2}$  and  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  of rainwater for comparison with corresponding measured values of  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$ . This comparison may prove to be uniquely useful in the assessment of the importance of  $\text{H}_2\text{O}_2$  in the oxidation of  $\text{SO}_2$  to  $\text{SO}_4^{2-}$  in the atmosphere.

The slope of 0.43 (approximately 2/5) in Eq. (5) of the regression curve through the data of Fig. 9 confirms the evidence of the intermediate species,  $\text{H}_2\text{O}_2\cdot\text{SO}_3^{2-}$ , which was previously proposed (Holt *et al.*, 1981). Apparently, the  $\delta^{18}\text{O}$  of the sulfate product is 2/5-controlled by the two oxygens in the  $\text{H}_2\text{O}_2$  of the adduct, and 3/5-controlled by the  $\text{HSO}_3^-$ , which, in turn, is isotopically controlled by rapid oxygen exchange with the large excess of water with which it is associated.

#### $\delta^{18}\text{O}$ and Concentration of $\text{H}_2\text{O}_2$ in Rainwater

Forty-four samples of rainwater, collected at Argonne from September 1984 through November 1985 and at the other three sites (in Michigan, New York, and North Carolina) during the summer months of 1985, were analyzed by the new method. Results obtained for 14 of these samples are given in Table 1 and in Fig. 10. Isotopic data for  $\text{H}_2\text{O}$ ,  $\text{SO}_4^{2-}$ , and  $\text{H}_2\text{O}_2$  in rainwater from the four sites are plotted vs. time in Fig. 10; in addition, Table 1 gives concentrations of  $\text{H}_2\text{O}_2$  and  $\text{SO}_4^{2-}$  in the rainwater, and the elapsed time between collection and analysis of each sample. Our analytical method does not discriminate  $\text{H}_2\text{O}_2$  from organic peroxides in rainwater; consequently, in the report of field results and in the discussions that follow, " $\text{H}_2\text{O}_2$ " will refer to combined  $\text{H}_2\text{O}_2$  and other active peroxides.

Results for 25 of the samples are not reported because the concentrations of  $\text{H}_2\text{O}_2$  were too low (<15 ppb) to yield sufficient amounts of  $\text{CO}_2$  to give reliable isotopic data by mass spectrometric analysis. Discarding of these samples of low  $\text{H}_2\text{O}_2$  concentration has the effect of estimating a lower than actual proportion of sulfate formed by peroxide oxidation of  $\text{SO}_2$  discussed later in the report. [The cause(s) of very low peroxide concentrations in rainwater samples may be one or more of the following: (1) during early use of the method,

Table I.

Concentration and isotopic results for rainwater collected at Argonne National Laboratory, IL, (ANL); Dearborn, MI, (DEA); Whiteface Mountain, NY, (WFM); and Research Triangle Park, NC, (RTP); May-September, 1985.

| Sample No.  | Storage<br>(days) | Concentration                   |  | $\delta^{18}\text{O}$                    |  |                                      |
|-------------|-------------------|---------------------------------|--|--|--|--------------------------------------|
|             |                   | $\text{H}_2\text{O}_2$<br>(ppb) | $\text{SO}_4^{2-}$<br>( $\text{mg L}^{-1}$ ) | $\text{H}_2\text{O}_2$<br>( $\text{‰}$ ) | $\text{H}_2\text{O}$<br>( $\text{‰}$ ) | $\text{SO}_4^{2-}$<br>( $\text{‰}$ ) |
| ANL-5/27/85 | 2                 | 330                             | 2.4  | 44.8                                     | -3.1                                   | 15.8                                 |
| WFM-6/07/85 | 6                 | 124                             | 1.5  | 50.5                                     | -3.7                                   | 15.0                                 |
| RTP-6/12/85 | 6                 | 147                             | 2.4  | 47.5                                     | -3.5                                   | 14.4                                 |
| RTP-6/18/85 | 9                 | 374                             | 2.1  | 45.0                                     | -2.5                                   | 15.8                                 |
| ANL-7/02/85 | 7                 | 316                             | 2.1  | 41.4                                     | -3.1                                   | 12.6                                 |
| DEA-7/02/85 | 9                 | 23                              | 4.4  | 58.3                                     | -6.6                                   | 12.9                                 |
| ANL-7/09/85 | 7                 | 90                              | 4.4  | 54.2                                     | -1.0                                   | 16.9                                 |
| RTP-7/10/85 | 8                 | 433                             | 3.6  | 53.8                                     | -2.6                                   | 14.3                                 |
| WFM-7/15/85 | 17                | 201                             | 2.0  | 48.4                                     | -3.9                                   | 15.3                                 |
| RTP-7/16/85 | 20                | 224                             | 3.6  | 58.0                                     | -1.5                                   | 16.4                                 |
| WFM-7/31/85 | 64                | 160                             | 1.2  | 41.0                                     | -7.9                                   | 14.1                                 |
| DEA-8/14/85 | 127               | 16                              | 2.9  | 47.9                                     | -1.3                                   | 16.8                                 |
| RTP-8/20/85 | 79                | 184                             | 4.1  | 49.6                                     | -3.5                                   | 12.8                                 |
| ANL-9/06/85 | 76                | 33                              | 1.2  | 44.8                                     | +0.4                                   | 16.1                                 |




Figure 10. Measured isotopic data for  $\text{H}_2\text{O}_2$ ,  $\text{SO}_4^{2-}$ , and  $\text{H}_2\text{O}$  in the samples of rainwater



storage procedures were inadequate for prevention of autodecomposition of peroxides; (2) several samples were collected during winter months of 1984-85 when concentrations of peroxides in precipitation were very low, and (3) high concentrations of SO<sub>2</sub>, relative to peroxides, in the atmosphere depletes the peroxide that might otherwise be present in collected samples of rainwater.] Similarly, three samples were disqualified because of very low sulfate concentrations (<0.3 mg L<sup>-1</sup>), and the data for two other samples were not included because of analytical difficulties (back diffusion of CO<sub>2</sub> from the charcoal furnace into cold trap 2, when the line was inadvertently evacuated between the furnace and the water sample with no helium flow).

## SECTION 4

### DISCUSSION

#### $\delta^{18}\text{O}$ of $\text{H}_2\text{O}_2$ in Rainwater

A unique characteristic of atmospheric peroxides, as shown for the first time by application of our newly developed analytical method, is the very high  $\delta^{18}\text{O}$ , relative to the  $\delta^{18}\text{O}$  of atmospheric water, sulfates, and air. As demonstrated in Fig. 10, the measured  $\delta^{18}\text{O}$  of atmospheric peroxides ranged from 45 to 60‰; sulfates, 13 to 17‰; and rainwater 0 to -6‰. (The  $\delta^{18}\text{O}$  of air oxygen is constant at 23.5‰.) In contrast to the very high  $\delta^{18}\text{O}$  of atmospheric peroxides, the  $\delta^{18}\text{O}$  of reagent-bottle  $\text{H}_2\text{O}_2$  (Fisher Scientific, H325) was found to be -6‰.

The oxygen isotopy of peroxide formation in the atmosphere, resulting in such high  $\delta^{18}\text{O}$ , is intriguing because it is necessarily related to the mechanism(s) by which the peroxide is formed. For the various sources of atmospheric peroxides (Lee, 1985), each possible mechanism of formation is characterized by its own oxygen isotopy. Consequently, our results suggest that oxygen isotopic studies may well be uniquely applicable to investigations of the origin of peroxides in the atmosphere.

As mentioned earlier, the  $\delta^{18}\text{O}$  of peroxides in rainwater was higher than might be expected for  $\text{H}_2\text{O}_2$  that is formed by high-voltage discharge, such as lightning. For example, application of Eq. 4 to the range of  $\delta^{18}\text{O}$  in water (Fig. 10) would lead to an estimated range for  $\text{H}_2\text{O}_2$  of ~24‰ instead of the observed 45-60‰.

A phenomenon that may contribute substantially to the relatively high  $\delta^{18}\text{O}$  of peroxides in rainwater (depending on the conditions of the rainfall, the sample collection, and the storage) is autodecomposition of the peroxide. As the peroxides decompose (under favorable conditions of catalyzing contaminants, temperature and pH), the  $\delta^{18}\text{O}$  by the residual peroxides is expected to increase. Therefore, depending on the extent of autodecomposition between the time of liquid-phase oxidation of  $\text{SO}_2$  in the atmosphere and the

time of collection, acidification, and refrigeration on the ground, the  $\delta^{18}\text{O}$  of the residual  $\text{H}_2\text{O}_2$  in the analyzed sample may be significantly higher than that of the  $\text{H}_2\text{O}_2$  involved in the  $\text{SO}_2$  oxidation. Further experimentation to establish the relationship between the  $\delta^{18}\text{O}$  of peroxide in cloud water and that of concurrently collected rainwater on the ground might provide a correction factor which would improve the significance of the isotope data that is conveniently available from the analysis of water in ground-based collectors.

#### $\delta^{18}\text{O}$ of Atmospheric Sulfates: Measured vs. Calculated

The measured  $\delta^{18}\text{O}$  values for sulfates, Fig. 10, are co-plotted in Fig. 11 with correspondingly calculated values for  $\text{H}_2\text{O}_2$  oxidation, metal-catalyzed aqueous oxidation, and primary sulfates. The measured values are clearly higher than those calculated for metal-catalyzed oxidation and clearly lower than those calculated either for  $\text{H}_2\text{O}_2$  oxidation or for primary sulfates. Calculated values for  $\text{H}_2\text{O}_2$  oxidation were obtained by use of Eq. (9) for metal-catalyzed aqueous oxidation by

$$\delta^{18}\text{O}_{\text{SO}_4^{2-}} = 0.8\delta^{18}\text{O}_{\text{H}_2\text{O}} + 10\text{‰} \quad (\text{Holt } et \text{ al.}, 1982) \quad (10)$$

and for primary sulfates by

$$\delta^{18}\text{O}_{\text{SO}_4^{2-}} = 0.015\delta^{18}\text{O}_{\text{H}_2\text{O}} + 45\text{‰} \quad (\text{Holt } et \text{ al.}, 1984). \quad (11)$$

In 1982, Holt *et al.* proposed that since their measured values for  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  in precipitation water ranged consistently between the calculated values for primary sulfates and secondary sulfates produced by metal-catalyzed oxidations, the fraction of atmospheric sulfates at a given site originating as primary sulfates, could be estimated. At that time, however, the isotopic qualities of atmospheric peroxides and of the sulfates which they might produce were unknown. Since then field experiments were performed near a strong source of primary sulfates, the isotopic results of which indicated that scavengement was  $\sim 300$  times more efficient for sulfates than for  $\text{SO}_2$  (Holt *et al.*, 1983). Consequently, we expect that, during precipitation, scavengement of primary sulfates is essentially complete




Figure 11.  $\delta^{18}\text{O}$  of sulfate in rainwater: calculated compared to measured

within a few kilometers of the source, whereas beyond a few kilometers, the scavenged sulfates are mainly secondary sulfates, formed earlier or within the storm system by one or more mechanisms of SO<sub>2</sub> oxidation.

#### Fraction of Atmospheric Sulfates Formed by Peroxide Oxidation

Assuming that primary sulfates of characteristically high  $\delta^{18}\text{O}$  (Holt *et al.*, 1984) are effectively scavenged by rain within a short distance from their sources, an estimation of the fraction of sulfates in rainwater several kilometers from a strong source can be made from relative deviations of the measured  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  values from the corresponding calculated  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  values for peroxide oxidation and metal-catalyzed O<sub>2</sub> oxidation, respectively.

Using the following relationship for calculation of the percentage of sulfate in rainwater formed by H<sub>2</sub>O<sub>2</sub> oxidation,

$$(\%) = \frac{\delta^{18}\text{O}_{\text{SO}_4^{2-}}(\text{meas.}) - \delta^{18}\text{O}_{\text{SO}_4^{2-}}(\text{calc., O}_2 \text{ oxdn.})}{\delta^{18}\text{O}_{\text{SO}_4^{2-}}(\text{calc., H}_2\text{O}_2 \text{ oxdn.}) - \delta^{18}\text{O}_{\text{SO}_4^{2-}}(\text{calc., O}_2 \text{ oxdn.})} \times 100, \quad (12)$$

the percentages for our four sampling sites during 1985 are plotted vs. time in Fig. 12. The average for all of the values at all of the sites is  $37 \pm 8\%$  sulfate, formed by peroxide oxidation.

Equation (12) assumes that all sulfate measured in the rainwater samples was produced within the raining cloud and that only H<sub>2</sub>O<sub>2</sub> and metal-catalyzed aqueous oxidation contributed to the oxidation of the dissolved SO<sub>2</sub>. Other potential sources of sulfate in the rainwater samples were aqueous-phase ozone oxidation of SO<sub>2</sub>, dissolution after gas-phase reaction of SO<sub>2</sub> with OH radicals, and dissolution of sulfate nuclei previously formed in non-precipitating clouds by any of the above oxidation processes.

In reviewing our work, Dodge (1986) pointed out that the results of other researchers (Scott, 1982; Hegg and Hobbs, 1984; Scire and Venkatram, 1985) variously indicate that 65 to 85% of sulfate in cloudwater is due to *in situ* aqueous phase reactions and the remainder




Figure 12. Percent of sulfate formed by peroxide oxidation, assuming negligible primary sulfates

to cloud scavenging of pre-existing sulfates. However, since the residence time of aerosol sulfate in the atmosphere is rather short, any aerosol sulfate scavenged could be expected to be about the same in isotopic quality as that in the precipitating cloud system. Further, ozone is expected to play a minor role in the oxidation of  $\text{SO}_2$  in cloudwater of pH lower than about 5 (Penkett *et al.*, 1979).

At best, these data establish a lower limit for peroxide oxidation. As discussed above, the  $\delta^{18}\text{O}$  of atmospheric peroxides may be somewhat lower than that which is finally measured from the ground-based rain collector. It follows that the calculated  $\delta^{18}\text{O}_{\text{SO}_4^{2-}}$  would be correspondingly lower, and the % of sulfate formed by peroxide oxidation would be correspondingly higher.

The technological significance of our findings (that ~40% or more of acid sulfate in rainwater originates from peroxide oxidation of  $\text{SO}_2$ ) is that further investigations of the origin(s) and of possible methods of control of atmospheric peroxides are of prime importance.

## SECTION 5

### SUMMARY

A new method was developed for the determination of the  $\delta^{18}\text{O}$  of peroxide (ppb range) in rainwater. Experimental results showed the method to be reliable with respect to recovery of added  $\text{H}_2\text{O}_2$ , blanks, and freedom from isotopic interference. A procedure for collection and treatment of 25-L samples of rainwater with minimum peroxide decomposition between collection and analysis was developed. An isotopic relationship was established between the oxygen in sulfate and the oxygen in the peroxide and in the water involved in the peroxide oxidation of  $\text{SO}_2$  to  $\text{SO}_4^{2-}$ . By using this relationship for peroxide oxidation and a similar relationship previously determined for aqueous metal-catalyzed oxidation, the fraction of sulfates formed by peroxide oxidation in rainwater from four sampling sites in Illinois, North Carolina, New York, and Michigan was estimated to be  $\sim 40\%$ . This fraction could be substantially higher if a substantial amount of peroxide undergoes decomposition between the time of oxidation of  $\text{SO}_2$  in the atmosphere and the time of collection and peroxide-stabilization treatment of the rainwater sample on the ground. Our results show that atmospheric peroxides play a major role in the formation of sulfates in the atmosphere, and, therefore, that it is essential to obtain all possible information on their origin and control.



## REFERENCES

- Cahill, A. E. and Taube, H. The use of heavy oxygen in the study of reactions of hydrogen peroxide. *J. Amer. Chem. Soc.*, **74**, 2312 (1952).
- Chamberlain, J., Foley, H., Hammer, D., MacDonald, G., Rothaus, O., and Ruderman, M. The Physics and Chemistry of Acid Precipitation, Technical Report JSR-81-25, prepared for U.S. Department of Energy, September 1981.
- Dodge, M. C. Private communication, U.S. Environmental Protection Agency, Research Triangle Park, NC, 1986.
- Hegg, D. A., Hobbs, P. V., and Radke, L. F. Measurements of the scavenging of sulfate and nitrate in clouds. *Atmos. Environ.*, **18**, 1939 (1984).
- Holt, B. D. Use of capillary trap in microdetermination of carbon. *Anal. Chem.*, **27**, 1500 (1955).
- Holt, B. D. Preparation of carbon dioxide from sulfates, sulfur dioxide, air, and water for determination of oxygen isotope ratio. *Anal. Chem.*, **49**, 1664 (1977).
- Holt, B. D., Kumar, R., and Cunningham, P. T. Primary sulfates in atmospheric sulfates: estimation by oxygen isotope ratio measurements. *Science*, **217**, 51 (1982).
- Holt, B. D., Kumar, R., and Cunningham, P. T. Oxygen-18 study of the aqueous-phase oxidation of sulfur dioxide. *Atmos. Environ.*, **15**, 557 (1981).
- Holt, B. D. and Kumar, R. Oxygen-18 study of high-temperature air oxidation of SO<sub>2</sub>. *Atmos. Environ.*, **18**, 2089 (1984).
- Holt, B. D., Nielsen, E., and Kumar, R. Oxygen-18 estimation of primary sulfate in total sulfate scavenged by rain from a power plant plume. "Precipitation Scavenging, Dry Deposition, and Resuspension," Vol. 1, "Precipitation Scavenging," H. R. Pruppacher, R. G. Semonin, and W. G. N. Slinn, eds., pp. 357-368, Elsevier, New York, 1983.
- Holt, T. G. Private communication, University of Illinois, Urbana, 1983.

- Horibe, Y., Shigehara, K., and Takakawa, Y. Isotope separation factor of carbon dioxide-water system and isotopic composition of atmospheric oxygen. *J. Geophys. Res.*, **78**, 2625 (1973).
- Jarnagin, R. C. and Wang, J. H. Investigation of the catalytic mechanism of catalase and other ferric compounds with doubly  $^{18}\text{O}$ -labeled hydrogen peroxide. *J. Amer. Chem. Soc.*, **80**, 786 (1958).
- Kok, G. L. The production of hydrogen peroxide by lightning. Paper presented at the American Geophysical Union Meeting, San Francisco, 1982.
- Lee, Y. N. Chemical transformations in acid rain; Vol. II. Investigation of kinetics and mechanism of aqueous-phase peroxide formation. U.S. Environment Protection Agency Technical Report *EPA/600/09* (1985).
- Penkett, S. A., Jones, B. M. R., Brice, K. A., and Eggleston, A. E. J. The importance of atmospheric ozone and hydrogen peroxide in oxidizing sulfur dioxide in cloud and rainwater. *Atmos. Environ.*, **13**, 123 (1979).
- Schumb, W. C., Satterfield, C. N., and Wentworth, R. L. "Hydrogen Peroxide," Reinhold Publishing Corporation, New York, (1955).
- Scire, J. S., and Venkatram, A. The contribution of in-cloud oxidation of  $\text{SO}_2$  to wet scavenging of sulfur in convective clouds. *Atmos. Environ.*, **19**, 637 (1985).
- Scott, B. C. Predictions of in-cloud conversion rates of  $\text{SO}_2$  to  $\text{SO}_4$  based upon a simple chemical and kinematic storm model. *Atmos. Environ.*, **16**, 1735 (1982).
- Vol'nov, I. I., Tsentsiper, A. B., Chamova, V. N., Latysheva, E. I., and Kuznetsova, Z. I. Synthesis of hydrogen [ $^{18}\text{O}$ ] peroxide from the dissociated vapor of [ $^{18}\text{O}$ ] water in a glow discharge. *Russian J. Phys. Chem.*, **38**, 645 (1964).

Page Intentionally Blank