




Adsorption of Energy-Related Organic Pollutants

A Literature Review



RESEARCH REPORTING SERIES

Research reports of the Office of Research and Development, U.S. Environmental Protection Agency, have been grouped into nine series. These nine broad categories were established to facilitate further development and application of environmental technology. Elimination of traditional grouping was consciously planned to foster technology transfer and a maximum interface in related fields. The nine series are:

1. Environmental Health Effects Research
2. Environmental Protection Technology
3. Ecological Research
4. Environmental Monitoring
5. Socioeconomic Environmental Studies
6. Scientific and Technical Assessment Reports (STAR)
7. Interagency Energy-Environment Research and Development
8. "Special" Reports
9. Miscellaneous Reports

This report has been assigned to the ECOLOGICAL RESEARCH series. This series describes research on the effects of pollution on humans, plant and animal species, and materials. Problems are assessed for their long- and short-term influences. Investigations include formation, transport, and pathway studies to determine the fate of pollutants and their effects. This work provides the technical basis for setting standards to minimize undesirable changes in living organisms in the aquatic, terrestrial, and atmospheric environments.

ADSORPTION OF ENERGY-RELATED ORGANIC
POLLUTANTS: A LITERATURE REVIEW

by

K. A. Reinbold, J. J. Hassett,
J. C. Means, and W. L. Banwart
Institute for Environmental Studies
and
Department of Agronomy
University of Illinois at Urbana-Champaign
Urbana, Illinois 61801

Contract No. 68-03-2555

Project Officer

David S. Brown
Environmental Processes Branch
Environmental Research Laboratory
Athens, Georgia 30605

ENVIRONMENTAL RESEARCH LABORATORY
OFFICE OF RESEARCH AND DEVELOPMENT
U.S. ENVIRONMENTAL PROTECTION AGENCY
ATHENS, GEORGIA 30605

DISCLAIMER

This report has been reviewed by the Environmental Research Laboratory, U.S. Environmental Protection Agency, Athens, GA, and approved for publication. Approval does not signify that the contents necessarily reflect the views and policies of the U.S. Environmental Protection Agency, nor does mention of trade names or commercial products constitute endorsement or recommendation for use.

FOREWORD

Environmental protection efforts are increasingly directed towards prevention of adverse health and ecological effects associated with specific compounds of natural or human origin. As part of this Laboratory's research on the occurrence, movement, transformation, impact, and control of environmental contaminants, the Environmental Processes Branch studies the microbiological, chemical, and physico-chemical processes that control the transport, transformation, and impact of pollutants in soil and water.

Efforts to achieve our national goal of energy independence will require increasing use of our country's vast domestic coal reserves. The combustion of coal or its conversion to a gaseous or liquid fuel, however, can release numerous organic compounds that are potentially toxic, carcinogenic, or mutagenic. This report reviews the literature on the adsorption of energy-related organic pollutants and other compounds on sediments and soils. Information on the adsorption of these pollutants onto sediments is needed to predict their movement and fate in aquatic systems so that potential environmental problems can be anticipated.

David W. Duttweiler
Director
Environmental Research Laboratory
Athens, Georgia

ABSTRACT

This report is a literature review which was completed as the first phase of a research project on sorption properties of sediments and energy-related organic pollutants. Adsorption of organic compounds in general is discussed, and analytical methodology in soil thin-layer chromatography and chemical analysis as applicable to measurement of sorption properties is summarized. The literature on the adsorption of energy-related organic pollutants is reviewed. Reported constants for the adsorption of organic compounds on several adsorbents are tabulated, and factors which influence the adsorption are discussed.

This report was submitted in partial fulfillment of Contract No. 68-03-2555 by the University of Illinois Institute for Environmental Studies in cooperation with the Department of Agronomy under the sponsorship of the U.S. Environmental Protection Agency. This report covers the period of July 1977 to April 1978, and work was completed as of June 1979.

CONTENTS

FOREWORD	iii
ABSTRACT	iv
FIGURE	vi
TABLES	vi
ACKNOWLEDGMENTS	vii
1. INTRODUCTION	1
2. SUMMARY AND CONCLUSIONS	8
3. ADSORPTION OF ORGANIC COMPOUNDS	9
The Solid-Solution Interface	9
Adsorptive Forces	9
Adsorption Isotherms	11
4. ANALYTICAL PROCEDURES: SOIL THIN-LAYER CHROMATOGRAPHY	18
5. ANALYTICAL PROCEDURES: CHEMICAL ANALYSIS	22
Recovery of Organic Compounds from Environmental Samples	22
Fractionation, Cleanup, and	
Separation of Organic Extract Components	23
Quantitation of Trace Organic Compounds in Solvent Extracts	24
6. REVIEW AND INTERPRETATION OF ADSORPTION DATA	30
Factors Influencing Adsorption	30
Interpretation of Tabulated Data	34
BIBLIOGRAPHY	114
Adsorption of Organic Compounds	114
Analytical Procedures	134
Compound Characteristics	153
Occurrence and Distribution of Energy-related Organic Compounds	158
APPENDIX: FORMULAS OF ORGANIC COMPOUNDS	161

FIGURE

1. Functional Group Model of Bituminous Coal 2

TABLES

1. Organic Contaminants Present in Coal Tar
from the Synthane Coal Conversion Process 3
2. Contaminants Present in Product Water
from the Gasification of Illinois No. 6 Coal 4
3. Classes of Known or Suspected Carcinogenic or Cocarcinogenic
Compounds Associated with Processing and Utilization of Coal 5
4. Selected Energy-Related Organic Compounds 7
5. K_{ow} , Calculated K_{oc} , Linear K_p and Measured K_{oc} Values for Sorption of
Pyrene, Dibenzothiophene and Acetophenone by Soils and Sediments 33
6. Adsorption Constants for Organic Compounds 36
7. Average Values of Adsorption Constants for Organic Pesticides 89
8. Relationship between Octanol/Water
Partition and R_f Values of Pesticides on a Soil 107
9. Leaching of Pesticides from a Soil 108
10. Sorption Dependence on Sorbate Properties 109

ACKNOWLEDGMENTS

The authors gratefully acknowledge the assistance and coordination provided by Dr. David Brown, Project Officer, of the EPA Environmental Research Laboratory in Athens, Georgia. Co-principal investigators for the project under which this review was done were Dr. John J. Hassett, associate professor of soils, and Dr. Jay C. Means, assistant research chemist.

The literature review presented in this report was conducted under the direction of Dr. Keturah A. Reinbold, associate research biologist, with the assistance of Ms. Dee Condon and Ms. Carol Wells, research assistants, in conducting the literature search. Information on soil thin-layer chromatography was summarized by Dr. Wayne L. Banwart, assistant professor of soils. Appreciation is expressed to Thomas Knecht, publications director of the Institute for Environmental Studies, and Cindy Bohde, student editorial assistant, for the technical editing of this report and to Jean Clarke, Judith Jones, Trace Black, and Sharon Sparling for typing the report.

1 INTRODUCTION

Fossil fuels are a major source of anthropogenic organic compounds in the environment. Because supplies of petroleum and natural gas are dwindling, it will be necessary to utilize increasing amounts of coal to meet our energy needs in the near future. As Figure 1 illustrates, coal is a complex organic chemical. Either the combustion of coal or its conversion to a gaseous or liquid fuel breaks the coal into numerous simpler organic compounds, which then appear in process waste streams and may be released into the environment. These coal fragments include a great variety of polycyclic aromatics, heterocyclic- and carbonyl-polycyclics, and aromatic amines, groups which all contain known or suspected carcinogens, as well as phenolics and long-chain aliphatic hydrocarbons (TRW Systems and Energy, 1976; Sharkey et al., 1976).

Coal conversion processes produce large volumes of gaseous and aqueous effluents (Magee, Bertrand, and Jahnig, 1976). Hundreds of thousands of tons of gases containing volatile organics, particulates, and combustion gases are released each day. These emissions may disperse over hundreds of miles, distributing the effluent materials in both terrestrial and aquatic systems. With such large volumes of effluents, even trace components are distributed in significant amounts--up to hundreds of pounds per year. Some of these materials are potentially toxic, carcinogenic, or mutagenic.

Aqueous effluents may be released at a rate of up to 5 million gallons each day. Also, large volumes of aqueous leachate from stockpiles of coal or solid wastes, such as ash and char, are produced. The wastes from the various steps of the coal conversion process contain a variety of organic materials.

A number of organic products have been detected in the wastes from coal conversion pilot plants. Forney et al. (1974) identified some of the major organic compounds in tars produced by the Synthane coal gasification process (Table 1), and Schmidt, Sharkey, and Friedel (1974) analyzed the process water (Table 2). TRW Systems and Energy (1976) listed several organic compounds which are associated with the processing and utilization of coal and which are known or suspected carcinogenic or cocarcinogenic compounds (Table 3).

Clearly, large quantities of organic compounds are introduced into aquatic systems, either directly in aqueous effluents or indirectly from gaseous effluents. The behavior of these compounds in aquatic systems depends largely upon the extent to which they are adsorbed on suspended or settled sediments.




Figure 1. Functional group model of bituminous coal (Wiser).

Source: Wewerka, Williams, and Vanderborgh, 1976.

TABLE 1. ORGANIC CONTAMINANTS PRESENT IN
COAL TAR FROM THE SYNTHANE COAL CONVERSION PROCESS

Structural Type	Volume (percent)			
	Illinois No. 6	Lignite	Montana Subbituminous	Pittsburg Seam
Benzenes	2.1	4.1	3.9	1.9
Indenes	8.6	1.5	2.6	6.1
Indanes	1.9	3.5	4.9	2.1
Naphthalenes	11.6	19.0	15.3	16.5
Fluorenes	9.6	7.2	9.7	10.7
Acenaphthenes	13.5	12.0	11.1	15.8
3-ring aromatics	13.8	10.5	9.0	14.8
Phenyl naphthalenes	9.8	3.5	6.4	7.6
4-ring aromatics - peri	7.2	3.5	4.9	7.6
4-ring aromatics - cata	4.0	1.4	3.0	4.1
Phenols	2.8	13.7	5.5	3.0
Naphthols	+	9.7	9.6	+
Indanols	0.9	1.7	1.5	0.7
Acenaphthols	-	2.5	4.6	2.0
Phenanthrols	2.7	-	0.9	-
Dibenzofurans	6.3	5.2	5.6	4.7
Dibenzothiophenes	3.5	1.0	1.5	2.4
Benzonaphthothiophenes	1.7	-	-	-
N-heterocycles	10.8	3.8	5.3	8.8

Source: Forney *et al.*, 1974

TABLE 2. CONTAMINANTS PRESENT IN PRODUCT WATER
FROM THE GASIFICATION OF ILLINOIS NO. 6 COAL

Compound	Concentration (ppm)
Phenols	2,660 to 3,400
Cresols	2,610 to 2,840
C ₂ -phenols	560 to 1,170
C ₃ -phenols	70 to 150
Dihydric phenols	60 to 300
Benzofuranols	70 to 120
Indanols	60 to 210
Acetophenones	
Benzoic Acids	40 to 210
Hydroxybenzaldehyde	
Naphthols	110 to 160
Indenols	90
Benzofurans	10 to 30
Dibenzofurans	10
Biphenols	20 to 40
Benzothiophenols	60 to 110
Pyridines	60 to 580
Quinolines	10 to 20
Indoles	20 to 70

Source: Schmidt, Sharkey, and Friedel, 1974.

OBJECTIVES

To examine the adsorption of energy-related organic compounds in the environment, a research project was initiated in the Institute for Environmental Studies at the University of Illinois at Urbana-Champaign. The project is supported by the U.S. Environmental Protection Agency under contract number 68-03-2555. This report presents the results of the first phase of the project, the objective of which was to review published literature on (1) the adsorption of energy-related organic pollutants on sediments, (2) the theory of adsorption, and (3) analytical techniques pertinent to adsorption measurements. Because the literature survey produced only a limited amount

of information on the adsorption of energy-related organic compounds, a variety of organic compounds were included in this review to provide a background of information.

APPROACH

The literature was searched by a combination of computer and manual methods. The 1970-77 volumes of *Chemical Abstracts* were searched using the computerized Bibliographic Retrieval Service System. The key words used were terms such as *adsorption*, *desorption*, and *sediments* accompanied by the names of compound groups or of specific compounds which may be energy-related or organic pollutants, including all those listed in Table 4. Earlier volumes of *Chemical Abstracts* were searched manually, as were pertinent books, series of *Residue Reviews*, current journals, and the 1977 *Weekly Government Abstracts* from the National Technical Information Service.

The computerized search produced more than 900 citations. Copies of approximately one-third of these references were obtained for review. With the addition of those acquired by manual searching, a total of 670 references were obtained for review. A bibliography is included at the end of this report.

TABLE 3. CLASSES OF KNOWN OR SUSPECTED CARCINOGENIC
OR COCARCINOGENIC COMPOUNDS ASSOCIATED WITH
PROCESSING AND UTILIZATION OF COAL













Compound Class	Representative Compound	Structure
<u>Polynuclear Aromatic Hydrocarbons</u>		
Anthracenes	9-,10-dimethylantracene	
Chrysenes	chrysene	
Benzanthracenes	benzo(a)anthracene	
Fluoranthenes	benzo(j)fluoranthene	
Cholanthrenes	3-methylcholanthrene	

TABLE 3, continued

Compound Class	Representative Compound	Structure
Benzopyrenes	benzo(a)pyrene	
Dibenzpyrenes	dibenzo(a,h)pyrene	
<u>Nitrogen-, Sulfur-, and Oxygen-Containing Polycyclic Compounds</u>		
Mono- and dibenzacridines	dibenz(a,h)acridine	
Benzocarbazoles	7H-benzo(c,g)carbazole	
Benzathrones	7H-benz(d,e)anthracen-7-one	
<u>Aromatic Amines</u>		
Aminoazobenzenes	4-dimethylaminoazobenzene	
Naphthylamines	α -naphthylamine	
<u>Cocarcinogens and Promoting Agents</u>		
Phenols/naphthols	α -naphthol	
Long-chain aliphatic hydrocarbons	n-dodecane	$\text{CH}_3 - (\text{CH}_2)_{10} - \text{CH}_3$

Source: TRW Systems and Energy, 1976.

TABLE 4. SELECTED ENERGY-RELATED ORGANIC COMPOUNDS

<u>Polynuclear Aromatics</u>	<u>S-Heterocyclics</u>	<u>Phenolics</u>
Anthracene	2,3-Benzothiophene	1-Naphthol
Phenanthrene	Dibenzothiophene	2-Naphthol
Acenaphthene		Acenaphthol
Fluorene		4-Indanol
Naphthacene	<u>N-Heterocyclics</u>	4-Benzofuranol
Chrysene	Carbazole	4-Hydroxybenzothiophene
Pyrene	Indole	2,3,4-Trimethyl Phenol
Triphenylene	Acridine	
Perylene	Pyridines	
1,2-Benzopyrene	Pyrroles	<u>Miscellaneous</u>
3,4-Benzopyrene	Benzocarbazole	Acetophenone
1,2-Benzanthracene	Dibenzocarbazole	Anthraquinone
1,2,7,8-Dibenzanthracene	Benzoquinoline	Benzidine
1,2,5,6-Dibenzanthracene		Benzophenone
1,2,3,4-Dibenzanthracene		
7,12-Dimethylbenzanthracene		
3-Methylcholanthrene		

REFERENCES

- Forney, A. J., W. P. Haynes, S. J. Gasior, G. E. Johnson, and J. P. Strakey, Jr. 1974. *Analyses of Tars, Chars, Gases, and Water Found in Effluents from the Synthane Process*. Progress Report 76. Bureau of Mines, U.S. Department of the Interior.
- Magee, E. M., R. R. Bertrand, and C. E. Jahnig. 1976. Environmental impact and R and D needs in coal conversion. In *Symposium Proceedings: Environmental Aspects of Fuel Conversion Technology, III*, pp. 395-403. Report No. EPA-600/2-76-149. Washington, D. C.: Office of Research and Development, U. S. Environmental Protection Agency.
- Schmidt, C. E., A. G. Sharkey, Jr., and R. A. Friedel. 1974. *Mass Spectrometric Analysis of Product Water from Coal Gasification*. Technical Progress Report 86. Bureau of Mines, U. S. Department of the Interior.
- Sharkey, A. G., J. L. Schultz, C. White, and R. Lett. 1976. *Analysis of Organic Material in Coal, Coal Ash, Fly Ash, and Other Fuel and Emission Samples*. Report No. EPA-600/2-76-075. Washington, D. C.: Office of Research and Development, U. S. Environmental Protection Agency.
- TRW Systems and Energy. 1976. *Carcinogens Relating to Coal Conversion Processes*. Oak Ridge, Tenn.: U. S. Energy Research and Development Administration.
- Wewerka, E. M., J. M. Williams, and N.E. Vanderborgh. 1976. *Contaminants in Coals and Coal Residues*. Report No. LA-UR 76-2197. Los Alamos, N. M.: Los Alamos Scientific Laboratory.

2 SUMMARY AND CONCLUSIONS

We have reviewed the literature on the adsorption of energy-related organic pollutants and other organic compounds on sediments, soils, and other selected adsorbents. Adsorption constants reported in the literature for organic compounds are compiled in Tables 6 and 7, and factors which influence adsorption are discussed.

Among the many factors which influence adsorption are several molecular properties of compounds. It has been shown that as chain length, molecular volume, molecular weight, and carbon number increase, and as polarity decreases, adsorption of hydrophobic compounds increases. These properties are related to water solubility of the compound. The adsorption of hydrophobic compounds increases with decreasing water solubility, but for more polar compounds adsorption increases with decreasing water solubility only within a family of compounds. Soil organic matter content influences adsorption and has been shown to correlate with the partitioning of nonpolar organic compounds between octanol and water. This relationship makes possible the calculation of K_{oc} and K_p values when K_{ow} is known.

Little of the published information pertains specifically to the adsorption of energy-related organic pollutants onto sediments or soils. Data are available on the adsorption of a few such compounds onto carbon, but those results cannot be directly extrapolated to sediments or soils. Investigation of adsorption of these compounds on sediments is beginning, as in the laboratory phase of this project. Information on the adsorption of these pollutants onto sediments is needed to predict their movement and fate in aquatic systems. To obtain sufficient information, further research is necessary.

3 ADSORPTION OF ORGANIC COMPOUNDS

J. J. Hassett

THE SOLID-SOLUTION INTERFACE

The adsorption or concentration of organic materials at the solid-solution interface is of particular importance in natural waters. Streams, lakes, and rivers contain a variety of colloidal materials both in their bottom sediments and dispersed throughout the aqueous phase. These colloids can actively sorb organic materials, removing them from solution or suspension, and hence can have a marked effect on the chemical and physiological properties of the organic pollutants.

The sediments in a natural body of water consist predominantly of organic colloids (such as clay minerals and metal oxides, hydroxides, and carbonates), of organic colloids, and of living organisms. The metal oxides and hydroxides and the organic colloids may exist as discrete particles or as coatings on other colloids such as the clay minerals. A major source of these inorganic colloids (and, to a certain extent, of the organic colloids) is the erosion of watershed soils. The properties of the sediments can be quite similar to those of the soils from which they are derived, or they may be substantially altered from the original soil, since the sediments may be subject to physical sorting by the water and often to different chemical environments (e.g., lower redox potentials).

Adsorption at the solid-liquid interface results when the forces of attraction between the surface (adsorbent) and the solute (adsorbate) overcome the forces of attraction between the solvent and the solute. Hence, the degree of adsorption depends on the relative strengths of the adsorbate-adsorbent interactions and the solute-solvent interactions. If the adsorbate-adsorbent interactions dominate due to a strong adsorbate-adsorbent interaction or due to a weak solvent-solute interaction, adsorption will take place and the adsorbing species will be primarily associated with the solid phase. The net interaction of the surface and the adsorbate may result from a variety of chemical and electrical interactions (Stumm and Morgan, 1970; Adamson, 1967).

ADSORPTIVE FORCES

1. *Coulombic attraction.* This force of attraction results when a charged surface such as a clay mineral attracts an oppositely charged ion to maintain electrical neutrality. The force of attraction is given by Coulomb's law:

$$F = q_1 q_2 / DX^2 \quad (\text{Eq. 1})$$

where q_1 is the charge density of the surface, q_2 is the charge of the adsorbing species, D is the dielectric constant of the solvent and is a measure of the shielding of q_1 from q_2 by the solvent, and X is the distance between the charges. Sorption of inorganic cations by negatively charged soil colloids in the process of cation exchange is one example of coulombic attraction. The sorption of the organic cations paraquat and diquat by montmorillonite and kaolinite (Weber *et al.*, 1965) is also an example of coulombic attraction. In the case of paraquat and diquat sorption within the interlayer of montmorillonite, other sorptive forces add to the coulombic attraction.

2. *London-van der Waals dispersion forces* result from the oscillating electron cloud of one atom rotating in phase with a nonoverlapping oscillating electron cloud of another atom, producing a dipole-like attraction. The differential heats of adsorption are of the order of 1 to 2 kcal mole⁻¹ for small molecules and atoms. For large molecules the heats of adsorption may be much larger. Corkill *et al.* (1966) gave the heats of adsorption of methane, ethane, pentane and hexane on carbon black as being 3, 4.3, 9.2 and 11.4 kcal mole⁻¹.

3. *Orientation energy* results from the attraction of a permanent dipole for another permanent dipole. The resulting energy of attraction is less than 2 kcal mole⁻¹.

4. *Induction forces* result from the attraction of an induced dipole for the inducing species which can be either a permanent dipole or a charged site or species. This force often adds to the adsorptive forces present in cases of coulombic or orientation energy attraction. The energy of attraction is less than 2 kcal mole⁻¹.

5. *Hydrogen bonding* occurs in compounds such as water where electrons are unequally shared between the more electronegative oxygen and hydrogen. This arrangement results in a slight negative charge on the oxygen atom and a slight positive charge on the hydrogen atoms, producing (in the case of water) a dipole moment of 1.84 Debye units (Douglas and McDaniel, 1965). Hence, attraction is possible between dissimilarly charged sites of the adsorbate and the adsorbent. The energy of attraction ranges from 2 to 10 kcal mole⁻¹.

6. *Chemical forces* result when the adsorbate-adsorbent bond approaches an ordinary chemical bond in strength ($>>10$ kcal mole⁻¹). Chemical forces extend over only very short ranges and often result in the nature of the adsorbate being significantly different in the adsorbed state. Such adsorption is often termed chemisorption to distinguish it from the less specific low-energy physical sorption. It is often difficult to distinguish between chemical and physical sorption because a chemisorbed layer may have physically sorbed layers upon it.

For some solutes the attractive force of the adsorbate for the solid surface can play a subordinate role to the hydrophilic-hydrophobic balance of the solute with the solvent (Hance, 1967; Stumm and Morgan, 1970). This type

of adsorption is of particular interest when considering the adsorption of organic molecules that are concentrated at the solid-solution interface because of the hydrophobic nature of their hydrocarbon parts. For this type of adsorption the effect of the adsorbent on the interfacial tension or surface free energy would appear to be an important consideration in explaining the observed phenomena. Conversely, some ions that show a strong affinity for the solvent (for example, ones that are highly hydrated) may stay in solution even if they are specifically attracted to the adsorbent.

Traube observed a regularity in the lowering of surface tension by members of three homologous series of esters, alcohols, and fatty acids (Kipling, 1965). The rule derived from that study states, in essence, that the tendency to adsorb organic compounds from aqueous solution increases with increasing molecular weight for members of a homologous series. Hence, adsorptive energy increases systematically for each additional CH_2 group.

Other general rules (Stumm and Morgan, 1970) concerning the adsorption of organics state that a polar adsorbent adsorbs the most polar constituent of a nonpolar solution in preference to the least polar constituent(s). In contrast, a nonpolar surface adsorbs the nonpolar component preferentially from a polar solution.

ADSORPTION ISOTHERMS

Several mathematical expressions--some with a theoretical basis and others entirely empirical in nature--have been employed to describe the relationship between the amount adsorbed and the equilibrium solution concentration. Those equations that have a theoretical basis can (if the assumptions each is based upon are met) provide valuable information about bonding or adsorption energies (affinities), adsorption maxima (capacities), and interfacial free energies. The empirical equations provide a framework for predicting the distribution of adsorbate between the solid and aqueous phases.

Langmuir Adsorption Isotherm

The Langmuir equation (1918), originally developed to describe the adsorption of a gas by a clean solid surface, has been used by numerous investigators to describe adsorption at the solid-liquid interface. The equation usually takes the following form (Veith and Sposito, 1977):

$$\frac{x}{m} = \frac{KbC}{(1 + KC)} \quad (\text{Eq. 2})$$

where

- x/m = amount of adsorbate adsorbed per unit mass of adsorbent
- C = the equilibrium concentration of the adsorbate in solution
- K = a constant related to the bonding energy of the adsorbate to the adsorbent
- b = adsorption maximum or capacity factor

Correct use of the equation requires that two assumptions be met:

1. That the adsorbed ions be bound in a monolayer on a homogeneous surface with localized sites.
2. That the energy of adsorption be the same for each molecule of adsorbate regardless of the degree to which the monolayer is completed.

Veith and Sposito (1977) have shown that it is necessary not only to meet the basic assumptions of the Langmuir equation but also to have an independent means of determining that the only process taking place is adsorption, since the Langmuir equation will also fit data obtained in situations where secondary precipitation is taking place. Other investigators (Stumm and Morgan, 1970; Kipling, 1965) have also cautioned that although a set of data may fit the equation, that fact is not of itself sufficient evidence that the assumptions have been met.

The normal application of the Langmuir equation to adsorption data involves a least-squares fitting of the data to a linear form of equation 2. At least three linear forms of the Langmuir equation have been used in adsorption studies:

$$C/(x/m) = 1/Kb = C/b \quad (\text{Eq. 3})$$

$$1/(x/m) = 1/b + 1/KbC \quad (\text{Eq. 4})$$

$$(x/m) = b - (x/m)/KC \quad (\text{Eq. 5})$$

Equation 3 has been used extensively in adsorption studies involving soils. A plot of $C/(x/m)$ against C for this equation should yield a straight line having a slope of $1/b$ and an intercept at $1/Kb$. For equation 4 the plot would be $1/(x/m)$ against $1/C$, yielding a slope of $1/Kb$ and an intercept at $1/b$. This type of plot is very similar to the double-reciprocal or Lineweaver-Burk plot used in enzyme studies employing Michaelis-Menten kinetics. For equation 5 (x/m) is plotted against $(x/m)/C$, yielding a slope equal to $1/K$ and an intercept at b . This plot is of the same form as the Eadie-Hofstee plot (Hofstee, 1952; 1960) also employed in Michaelis-Menten kinetics.

Dowd and Riggs (1965) compared the statistical fit and the predictability of the three linear forms of the Michaelis-Menten equation, which have the same form as the Langmuir adsorption isotherm and its linear equations:

$$v = \frac{V_{\max} C_s}{K + C_s} \quad (\text{Eq. 6})$$

$$C_s/v = K/v_{\max} + C_s/v_{\max} \quad (\text{Eq. 7})$$

$$1/v = 1/v_{\max} + K/v_{\max} C_s \quad (\text{Eq. 8})$$

$$v = V_{\max} - K v/C_s \quad (\text{Eq. 9})$$

where

v = the initial velocity of the reaction

C_s = the concentration of the substrate

V_{\max} = the maximum initial velocity

K_m = the Michaelis constant

They found that equation 7, which has the same form as equation 3, and equation 9, which has the same form as equation 5, gave comparable predictions of the two constants when the error in determining the dependent variable (i.e., v or x/m) was small, although equation 9 would better show deviations from linearity. Equation 9 gave the best results when the error associated with determining the dependent variable was large. These results are supported by Syers *et al.* (1973), who in a soil adsorption study compared the form of the Langmuir equation given in equation 3 with that of equation 5. Dowd and Riggs (1965) concluded that the Lineweaver-Burk or double reciprocal plot--equation 8, which has the same form as equation 4--should not be used even if it fits the data well.

Several nonlinear forms of the Langmuir equation are used in adsorption studies:

$$\theta = KC/(1+KC) \quad (\text{Eq. 10})$$

$$(x/m) = bC/(K'+C) \quad (\text{Eq. 11})$$

Equation 10 is the same as equation 2 except that adsorption is expressed in terms of the percentage of the monolayer that is occupied by adsorbate molecules. This results in b , the monolayer capacity, having a value of 1 unit of adsorption sites and disappearing from the adsorption equation. Equation 11 reduces to equation 2 if K' is replaced with $1/K$; hence, $K' = 1/K$.

Freundlich Adsorption Isotherm

To describe adsorption from dilute solutions, Freundlich (1922) applied the adsorption isotherm:

$$a = \alpha C^{1/n} \quad (\text{Eq. 12})$$

or in its more commonly used form:

$$(x/m) = KC^{1/n} \quad (\text{Eq. 13})$$

where

C = the equilibrium concentration in the solution after adsorption

a or x/m = the adsorption value

α , K , and $1/n$ = constants

In studies of the adsorption of a variety of organic compounds on charcoals Freundlich found the adsorption exponent $1/n$ to vary between 0.1 and 0.5.

The Freundlich adsorption equation is normally considered an empirical equation relating the amount adsorbed to the equilibrium adsorbate concentration. Kipling (1965) cites a derivation of the Freundlich equation by Henry (1922) based on combining an expression for the free energy of a surface with the Gibbs equation. The resulting equation defines the adsorption exponent in terms of the monolayer capacity and surface free energy:

$$(x/m) = KC^{(RT(x/m)m/(\sigma_0 - \sigma_1))} \quad (\text{Eq. 14})$$

hence

$$1/n = (RT(x/m)m/(\sigma_0 - \sigma_1)) \quad (\text{Eq. 15})$$

where

$(x/m)m$ = the monolayer capacity

σ_0 = the surface free energy in contact with pure solvent

σ_1 = the surface free energy of the surface covered with a monolayer of solute

R = the gas constant

T = temperature

This equation is applicable to dilute solutions where Gibb's surface excess equals (x/m) , the amount adsorbed.

When $1/n \approx 1$ then the Freundlich isotherm reduces to a partition equation:

$$K = \frac{(x/m)}{C} = \frac{C_s}{C_w} \quad (\text{Eq. 16})$$

where

C_s = the concentration of adsorbate in the solid phase

C_w = the concentration of adsorbate in equilibrium with the adsorbed phase

Partition equations have been written for both adsorption and desorption (LaFleur, 1976):

$$K_f = \frac{C_s}{C_w} \quad \text{adsorption} \quad (\text{Eq. 17})$$

$$K_b = \frac{C_s}{C_w} \quad \text{desorption} \quad (\text{Eq. 18})$$

Usually $K_b > K_f$; that is, a given surface coverage will be in equilibrium with a greater concentration of the solute for an adsorption process than for the desorption process.

The Freundlich equation is normally statistically fit to adsorption data in its linear form:

$$\log(x/m) = \log K + 1/n \log C \quad (\text{Eq. 19})$$

A plot of $\log(x/m)$ against $\log C$ yields a slope of $1/n$ and an intercept equal to $\log K$.

In many studies the difference in adsorption of an organic material by several soils has been correlated with the organic carbon content of the adsorbate. When the adsorption constants are put on an organic carbon basis, the differences in adsorption are often removed.

$$K_{oc} = \frac{K \times 100}{\%OC} \quad (\text{Eq. 20})$$

Gibbs Adsorption Equation

The Gibbs equation, originally derived for the adsorption at the liquid-liquid interface (Kipling, 1965) has been applied to the adsorption of solutes from dilute solutions by solid surfaces (Stumm and Morgan, 1970):

$$\Gamma_i = \frac{\partial \gamma}{\partial \mu_i} \quad (\text{Eq. 21})$$

$$\mu_i = \mu_i^0 + RT \ln a_i \quad (\text{Eq. 22})$$

hence,

$$\Gamma_i = \frac{-1}{RT} \frac{\partial \gamma}{\partial \ln a_i} \quad (\text{Eq. 23})$$

$T, P, \text{ all } \mu\text{'s except } \mu_i$
and μ_{H_2O} constant

where

Γ_i = the adsorption density of component i

γ = the interfacial tension

μ_i = the chemical potential

a_i = the activity of component i

The Gibbs equation defines adsorption in terms of the effect of a solute in either increasing or decreasing interfacial tension. The equation illustrates that solutes which lower the interfacial tension tend to be concentrated at the interface. Many organic substances tend to lower the interfacial tension and hence are accumulated at the interface.

REFERENCES

- Adamson, A. W. 1967. *Physical Chemistry of Surfaces*. New York: Wiley.
- Corkill, J. M., J. F. Goodman, and J. R. Tate. 1966. Adsorption of non-ionic surface-active agents at the graphon/solution interface. *Trans. Far. Soc.* pp. 535-44.
- Douglas, B. E., and D. H. McDaniel. 1965. *Concepts and Models of Inorganic Chemistry*. Waltham, Mass.: Blaisdell Publishing Co.
- Dowd, J. E., and D. S. Riggs. 1965. A comparison of estimates of Michaelis-Menten kinetic constants from various linear transformations. *J. Biol. Chem.* 240:863-69.
- Freundlich, H. 1922. *Colloid and Capillary Chemistry*. London: Methion & Co.
- Hance, R. J. 1967. Relationship between partition data and the adsorption of some herbicides by soils. *Nature* 214:630-31.
- Henry, D. C. 1922. A kinetic theory of adsorption. *Philos. Mag.* 44:689-705.
- Hofstee, B. H. J. 1952. On the evaluation of the constants V_m and K_m in enzyme reactions. *Science* 116:329-31.
- Hofstee, B. H. J. 1960. Nonlogarithmic linear titration curves. *Science* 131:39.
- Kipling, J. J. 1965. *Adsorption from Solutions of Non-electrolytes*. New York: Academic Press.
- LaFleur, K. S. 1976. Carbaryl desorption and movement in soil columns. *Soil Sci.* 121:212-16.
- Langmuir, I. 1918. The adsorption of gases on plane surfaces of glass, mica and platinum. *J. Amer. Chem. Soc.* 40:1361-1403.
- Stumm, W., and J. J. Morgan. 1970. *Aquatic Chemistry*. New York: Wiley Interscience.
- Syers, J. K., M. G. Browman, G. W. Smillie, and R. B. Corey. 1973. Phosphate sorption by soils evaluated by the Langmuir adsorption equation. *Soil Sci. Soc. Amer. Proc.* 37:358-63.
- Veith, J. A., and G. Sposito. 1977. On the use of the Langmuir equation in the interpretation of "adsorption" phenomena. *Soil Sci. Soc. Amer. J.* 41:697-702.

Weber, T. B., P. W. Perry, and R. P. Upchurch. 1965. The influence of temperature and time on the adsorption of paraquat, diquat, 2,4-D and prometone by clays, charcoal, and an anion-exchange resin. *Soil Sci. Soc. Am., Proc.* 29:678-88.

4 ANALYTICAL METHODS: SOIL THIN-LAYER CHROMATOGRAPHY

W. L. Banwart

The principle of thin-layer chromatography (TLC) and evidence of its widespread use are presented by Stahl (1965). The basis of all chromatographic separations is similar in that a mobile phase passes over a stationary phase and thereby transports different substances with varying speeds in the direction of the flow. Conventional TLC is a form of elution chromatography in which molecules of the test compounds are exchanged between the mobile and stationary phase as a solvent carrying the compounds migrates across a uniform thin layer of stationary phase fixed to a glass plate. The rate or relative rate of movement of the test compounds is dependent on their physical properties and on experimental parameters. The solvent employed may be water, an organic solvent, or a mixture of solvents allowing movement of the test compounds.

The mobility or migration distances of substances on the TLC chromatograms can be expressed in terms of their relative mobility, R_f (Stahl, 1965),

where $R_f = \frac{\text{distance of spot center from starting point}}{\text{distance of solvent from starting point}}$.

The use of relative mobilities provides reproducible data, whereas absolute mobilities, which are defined simply as the distances of the spot centers from the starting point, may vary considerably with experimental conditions.

Using TLC to study the mobility of compounds requires a method for documenting the substances' movement. The final position of the compounds on the chromatogram can be made visible with dye indicators, fluorescence under ultraviolet light (Pullan, Howard, and Perry, 1966), a bioassay (Helling, Kaufman, and Dieter, 1971), a spark-chamber apparatus (Pullan et al, 1966), a beta camera (Snyder, 1970), and autoradiography using X-ray film (Mangold, Kammereck, and Malins, 1962). In the last of these methods, chromatograms are developed using ^{14}C labeled test compounds; X-ray film is then exposed to the chromatogram for several days before development. The developed films make it possible to observe visually the movement of the test compounds and can be used for easy calculations of R_f values. For quantitative measurements, radioactive compounds on TLC plates can be counted directly by commercially available instruments such as strip scanners (Ravenhill and James, 1967) or by zonal analysis where small successive segments of the chromatogram are scraped from the plate and radio-assayed by liquid scintillation spectrometry (Brown and Johnston, 1962). Thin-layer chromatography has been used as a standard method for separating and identifying many synthetic and natural organic compounds (Maier and Mangold, 1964; Stahl and Mangold, 1975), including polynuclear aromatic hydrocarbons (Zoccolillo and Liberti, 1976; Candeli et al., 1975).

Soil thin-layer chromatography was introduced by Helling and Turner (1968). In soil TLC a uniform but relatively thin layer (often less than 1 mm) of a soil-water slurry is spread on TLC plates and allowed to dry. The soil then serves as the stationary phase interacting with compounds carried by the mobile phase. Soil TLC has provided a much simpler and faster way of determining compound mobilities in soil than the traditional leaching columns. Helling and Turner (1968) found that pesticide mobilities determined by soil TLC correlated well with published data. The mobility of substances on soil TLC plates have been reported by Helling and Turner (1968) as frontal R_f values where

$$R_f = \frac{\text{distance of frontal edge of spot or streak from starting point}}{\text{distance of solvent edge from starting point}}$$

Standard or reference compounds can be spotted on each plate to improve reproducibility.

Data obtained by soil TLC provide information of a nature somewhat different from that obtained with sorption isotherms or partition coefficients. A compound on a soil TLC chromatogram may move as a compact band or as a diffuse streak. From the type of movement it is possible to draw conclusions about the homogeneity of the soil material or of the compound itself. It is also possible to obtain a relative measure of the distance compounds might leach in a soil system when finite amounts of water are applied. R_b values have been used (Rhodes, Belasco, and Pease, 1970), where

$$R_b = \frac{\text{distance moved by bottom of spot}}{\text{distance traveled by solvent}}$$

to measure the relative soil depth through which essentially all of a given organic compound (pesticide) applied to a soil has leached. Thus, soil TLC provides some kinds of information that cannot be obtained strictly from adsorption constants or partition coefficients.

Data recalculated from Rhodes, Belasco, and Pease (1970) showed simple correlation coefficients of $r = 0.95$ between Freundlich K and frontal R_f values for four agricultural chemicals applied to two soils. Helling (1971c) reported highly significant negative correlation coefficients for soil adsorption of nonionic herbicides and R_f values of the same or chemically similar herbicides. As determined by leaching experiments, the mobilities of six organophosphorus insecticides (McCarty and King, 1966), three acidic herbicides (Hamaker, Goring and Youngson, 1966) and their five s-triazine herbicides (Harris, 1966) were inversely related to their adsorption by the soil. Data by Hance (1967), using two soils and 29 organic compounds, show correlation coefficients ranging from 0.85 to 0.91 for the amount of compound adsorbed by the soil and the R_f values for TLC using 40% aqueous ethanol as a solvent. Other workers (Martin and Synge, 1941; Stahl, 1965) have also discussed the relationship between K and R_f values. If a correlation between R_f (mobility) and K (adsorption isotherm) values for a given set of organic compounds can be established, it should be possible to predict K values for additional test compounds on a particular soil.

The greatest use of soil TLC to date has been the study of pesticide mobility in soils (Helling 1971a,b,c; Hance, 1967), but applying the technique to other mobile organic compounds should provide useful data.

REFERENCES

- Brown, J. L., and J. M. Johnston. 1962. Radioassay of lipid components separated by thin-layer chromatography. *J. Lipid Res.* 3:480-81.
- Candeli, A., G. Morozzi, A. Paolacci, and L. Zoccolillo. 1975. Analysis using thin-layer and gas-liquid chromatography of polycyclic aromatic hydrocarbons in the exhaust products from a European car running on fuels containing a range of concentrations of these hydrocarbons. *Atmos. Environ.* 9:843-49.
- Hamaker, J. W., C. A. I. Goring, and C. R. Youngson. 1966. Sorption and leaching of 4-amino-3,5,6-trichloropicolinic acid in soils. *Advan. Chem Ser.* 60:23-37.
- Hance, R. J. 1967. Relationship between partition data and the adsorption of some herbicides by soils. *Nature* 214:630-31.
- Harris, C. I. 1966. Adsorption, movement, and phytotoxicity of monuron and s-triazine herbicides in soil. *Weeds* 14:6-10.
- Helling, C. S. 1971a. Pesticide mobility in soils. I: Parameters of soil thin-layer chromatography. *Soil Sci. Soc. Amer. Proc.* 35:732-37.
- Helling, C. S. 1971b. Pesticide mobility in soils. II: Applications of soil thin-layer chromatography. *Soil Sci. Soc. Amer. Proc.* 35:737-43.
- Helling, C. S. 1971c. Pesticide mobility in soils. III: Influence of soil properties. *Soil Sci. Soc. Amer. Proc.* 35:743-48.
- Helling, C. S., and B. C. Turner. 1968. Pesticide mobility: Determination by soil thin-layer chromatography. *Science* 162:562-63.
- Helling, C. S., D. D. Kaufman, and C. T. Dieter. 1971. Algae bioassay detection of pesticide mobility in soils. *Weed Sci.* 19:685-90.
- McCarty, P. L., and P. H. King. 1966. The movement of pesticides in soils. *Purdue Univ. Eng. Bull.*, Ext. Ser. no. 121:156-71.
- Maier, R., and H. K. Mangold. 1964. Thin-layer chromatography. *Advan. Anal. Chem. and Instrum.* 3:369-477.
- Mangold, H. K., R. Kammereck, and D. C. Malins. 1962. Thin-layer chromatography as an analytical and preparative tool in lipid radiochemistry. *Microchemical Journal Symposium Series* 2:697-714.
- Martin, A. J. P., and R. L. M. Synge. 1941. A new form of chromatography employing two liquid phases. I: A theory of chromatography. II: Application to the micro-determination of the higher monoaminoacids in proteins. *Biochem. J.* 35:1358-68.

- Pullan, B. R., R. Howard, and B. J. Perry. 1966. Measuring radionuclide distribution with crossed-wire spark chambers. *Nucleonics* 24:72-75.
- Ravenhill, J. R., and A. T. James. 1967. A simple sensitive radioactive scanner for thin-layer chromatograms. *J. Chromatog.* 26:89-100.
- Rhodes, R. C., I. J. Belasco, and H. L. Pease. 1970. Determination of mobility and adsorption of agrichemicals on soils. *J. Agric. Food Chem.* 18:524-28.
- Snyder, R. 1970. Thin-layer radiochromatography and related procedures. In *Progress in Thin-layer Chromatography and Related Methods*, ed. A. Niederwieser and G. Pataki, Vol. 1, pp. 52-73. Ondon: Ann Arbor-Humphrey Science Publishers.
- Stahl, E., ed. 1965. *Thin-layer Chromatography*. New York: Academic Press.
- Stahl, E., and H. K. Mangold. 1975. Techniques of thin-layer chromatography. In *Chromatography*. 3d ed., ed. E. Heftmann, pp. 164-88. New York: Van Nostrand Reinhold Company.
- Zoccolillo, L., and A. Liberti. 1976. Determination of polycyclic hydrocarbons by channel thin-layer chromatography. *J. Chromatog.* 120:485-88.

5 ANALYTICAL PROCEDURES: CHEMICAL ANALYSIS

J. C. Means

The analysis of trace organic compounds in water or bound to soil and sediment samples is a complicated task because of a number of factors. First, the water and soil or sediment may contain a wide variety of organic compounds of different polarities, molecular weights, and structures. Second, the affinities of different types of organic compounds for different soil-sediment types span a large range. Third, the solubilities of organic compounds in different solvents vary considerably. Fourth, interactions between soil components and many types of organic compounds are poorly understood.

In general, the process of analyzing trace organic components in water and soil or sediment samples may be divided into these tasks:

1. Quantitative recovery of the organic compound of interest
2. Separation of that compound from other types of organic compounds
3. Quantitation of the organic compound using spectroscopy, chromatography, and radioactivity or a combination of techniques.

RECOVERY OF ORGANIC COMPOUNDS FROM ENVIRONMENTAL SAMPLES

To accurately determine the amount of a given organic compound in a water sample or bound to a soil or sediment sample, the compound must be quantitatively recovered from the gross sample. In the case of a soil or sediment sample, recovery is routinely accomplished by extracting the compound from the soil in a Soxhlet apparatus. Depending upon the variety of compounds which require analysis, a single solvent, a series of solvents, or a mixture of solvents may be required. For example, a cyclohexane or benzene extraction (Sawicki, 1964; Hermann, 1974) is typically used to recover aliphatic and aromatic hydrocarbons from sediments, whereas a benzene and methanol mixture may be used to remove a broad spectrum of nonpolar and polar organics (Giger and Blumer, 1974; Blumer and Youngblood, 1976). Acetone has been used to extract polar organics and humic substances.* Regardless of the extraction procedure used, it is always necessary to perform recovery studies using a reference compound to verify that the expected extraction efficiencies are achieved.

Organic compounds may be quantitatively recovered from water samples using one of three techniques. Compounds having low boiling points and low or intermediate polarities may be stripped from the water by passing a pure

*F. J. Stevenson, 1977: personal communication.

inert gas through the sample and collecting it in a trap containing a gas chromatographic column packing (e.g., Tenax GC) (Bellar and Lichtenberg, 1974; Kuo *et al.*, 1977). Soluble organic compounds may be recovered from water by a series of extractions using single or mixed solvents. In some analytical schemes, the pH of the water may be altered (i.e., made basic, neutral, and acidic) to achieve partial separation of compounds having ionizable functional groups. (U.S. EPA, 1977; Acheson *et al.*, 1976; Chang, 1976; Webb *et al.*, 1973). A third technique, one which has been used very successfully, is to collect organic compounds by sorption on purified activated carbon (Keith *et al.*, 1976) or on purified sorbent resins (e.g., XAD-2, 4, and 8) (Malcolm, Thurman, and Aiken, 1977; Junk *et al.*, 1974; Adams, Menzies, and Levins, 1977). These techniques have the advantage that the organics contained in a very large sample volume may be collected on a relatively small amount of resin. However, the recovery efficiencies for different types of organics adsorbed from the water and subsequently desorbed from the resin vary significantly. Therefore, the recovery of each compound of interest should be evaluated. Another advantage of the resin sorption technique is that both polar and non-polar organics may be sorbed to the resin and eluted for analysis.

Once an extract has been prepared from either a solid or a water sample, it is usually necessary to reduce the volume so as to bring the concentrations of the extracted components into a detectable (ppm) analytical range. The best general technique available for this process is the use of a Kuderna-Danish evaporator (Webb *et al.*, 1973; U.S. EPA, 1977). In some cases air evaporation or evaporation under a stream of nitrogen may be sufficient, but several studies have shown that significant losses of extracted organics may occur (Chiba and Mosley, 1968; Goldberg, Delong, and Sinclair, 1973). Similarly, rotary vacuum evaporation may be appropriate for certain compounds, but losses of many extracted components may occur. Extracts are typically concentrated by a factor of 500 to 10,000, depending upon the origin of the extract and the sensitivity of the analytical systems being used. Here again, it is advisable to test the recovery efficiency for the compound of interest using the concentration technique being considered.

FRACTIONATION, CLEANUP, AND SEPARATION OF ORGANIC EXTRACT COMPONENTS

Organic extracts of soil or sediments and of water, particularly those from industrial areas, may contain hundreds of components. These multiple components tend to complicate the accurate quantitation of individual constituents. Therefore, some steps may be needed to fractionate the extract prior to analysis. To some extent, fractionation begins with the selection of an extracting solvent or sorption resin. However, solvent selectivities are rarely sufficient, especially when concentration factors are high. Solvent-solvent partitioning of the organics in an extract may be useful in crude fractionations (e.g., separating polar compounds from nonpolar compounds).

Liquid chromatography and thin-layer chromatography (TLC) provide the best selectivities in fractionating complex mixtures of organics. For relatively large extracts, separation on liquid chromatographic columns containing such adsorbents as silica gel, alumina, or Florisil is the method of choice. Excellent separations of complex mixtures from acidic, basic, and neutral fractions of coal wastes and cigarette-smoke condensates have been achieved

using these three adsorbents (Swain, Cooper, and Stedman, 1969; Bell, Ireland, and Spears, 1969; Severson *et al.*, 1976).

In the last few years, a number of advances in liquid chromatographic column packings have provided investigators with a number of gel-permeation and adsorption materials which can be extremely useful in fractionating and separating complex mixtures into subsamples which can be analyzed and quantitated.

For relatively small samples, thin-layer chromatography can be used successfully to fractionate a complex mixture. The same basic substrates mentioned above for use in column chromatography are used for TLC separations. Careful selection of solvents and the use of two-dimensional development can yield excellent purifications of individual components prior to quantitation. A large number of investigators have employed TLC to purify compounds for study or to fractionate extracts prior to quantitative analysis (Treiber, 1976; Grant and Meiris, 1977; Bender, 1968; White and Howard, 1967; Pierce and Katz, 1975; Stanley, Bender, and Elbert, 1973; additional references are given in the analytical techniques section of the bibliography at the end of this report).

QUANTITATION OF TRACE ORGANIC COMPOUNDS IN SOLVENT EXTRACTS

The quantitation of individual components of a complex mixture after partial cleanup or fractionation may generally be achieved using the techniques of spectroscopy, gas chromatography, liquid chromatography, or, in appropriate cases, liquid scintillation counting or combinations of the above techniques.

Spectroscopic techniques, which are based on some characteristic absorption wavelength (in the UV, visible, or infrared spectrum) of the compound or a derivative, vary in their selectivity and sensitivity. In cases where the extracts contain relatively few components, spectroscopic methods may be successfully used to quantitate a component without additional cleanup or separation. In most cases, however, the spectroscopic techniques must be used in combination with compatible separation systems (e.g., liquid chromatography or thin-layer chromatography). A large body of literature exists on separations and quantitations of organic pollutants using the above techniques. Representative reports are those of Caton, Matthews, and Walters (1976); Kelly (1967); Willis (1973); Freudenthal *et al.* (1975); Jenkins and Baird (1975); Brocco, Cantuti, and Cartoni (1970); and IUPAC Applied Chemistry Division (1974). Additional references are listed in the bibliography.

One spectroscopic technique which offers a relatively high degree of both selectivity and sensitivity is the use of fluorescence spectra of various compounds or their fluorescent derivatives (Woo *et al.*, 1978). These techniques are particularly useful when combined with liquid chromatography for the analysis of many coal-derived substances (Stroupe *et al.*, 1977) and pesticides (Mallet, Belliveau, and Frei, 1975). All of these spectroscopic techniques have the advantage that they are not destructive to the sample being analyzed and that the column packing materials and thin-layer supports used

for separations prior to spectral analysis can cover an almost unlimited range of molecular weights.

Gas chromatography (GC) is perhaps the most widely used technique for the separation and quantitation of organic compounds. Methods are available for almost every class of organic pollutant including many energy-related compounds. Column packing materials are available for the separation of compounds representing a broad spectrum of polarities and functional groups. Although fewer liquid phases are readily available, capillary columns have demonstrated increasing utility for the separation of the highly complex mixtures of organics typically found in environmental extracts.

Gas chromatographic separations in general are better than those offered by liquid chromatography, but the compounds which can be analyzed by GC are limited to those having significant vapor pressures at temperatures below 400°C. In most cases, gas chromatographic analysis is destructive of the sample. A wide variety of GC detector systems are available. The flame ionization detector is the most common universal type. A number of other types of detectors demonstrate selectivities for specific types of compounds (e.g., the electron capture detector for halogens, the thermionic detector for nitrogen and phosphorus, and the flame photometric detector for sulfur and phosphorus). All of these detection systems may be used to quantitate organic compounds with the use of appropriate internal standardization techniques. The literature contains a large amount of information on this general topic. Pertinent references are included in the bibliography.

In the last decade, gas chromatography has been combined with mass spectrometry, providing investigators with a very powerful analytical system which can be used to both identify and quantitate organic compounds (John and Nickless, 1977; Janini *et al.*, 1976; Alford, 1977; Oswald, Albro, and McKinney, 1974; O'Reilly and Murrmann, 1974; Lao, Thomas, and Monkman, 1975; and McGuire, Alford, and Carter, 1973).

In certain types of experiments, radiolabeled organic compounds may be used successfully to follow the movement of trace organics in environmental samples. When labeled compounds are used, many types of samples may be analyzed directly without the need for extraction, concentration, or cleanup procedures. Care must be taken, however, to insure that the labeled compound introduced into the experimental system is not degraded in such a way that the radioactivity is lost or transferred to other compounds. The sample is typically separated by (1) thin-layer chromatography followed by radioautography or liquid scintillation counting or (2) by liquid chromatography followed by liquid scintillation spectrometry. Carbon -14 or tritium -3 labeled compounds can be detected in liquid samples by liquid scintillation spectrometry on aliquots of the samples. If solid samples are to be analyzed, the radiolabeled compound may be extracted and then analyzed as a liquid sample. Solids may also be analyzed directly by pyrolytic combustion of the solid and recovery of the radioactivity as $^{14}\text{CO}_2$ or $^3\text{H}_2\text{O}$.

REFERENCES

- Acheson, M. A., R. M. Harrison, R. Perry, and R. A. Wellings. 1976. Factors affecting the extraction and analysis of polynuclear aromatic hydrocarbons in water. *Water Res.* 10:207-12.
- Adams, J., K. Menzies, and P. Levins. 1977. *Selection and Evaluation of Sorbent Resins for the Collection of Organic Compounds*. Report No. EPA 600/7-77-044. Research Triangle Park, N.C.: Industrial Environmental Research Lab., Office of Research and Development, U.S. Environmental Protection Agency.
- Alford, A. 1977. Environmental applications of mass spectrometry. *Bio-medical Mass Spectrometry* 4:1-22.
- Bell, J. H., S. Ireland, and A. W. Spears. 1969. Identification of aromatic ketones in cigarette smoke condensate. *Anal. Chem.* 41:310-13.
- Bellar, T. A. and J. J. Lichtenberg. 1974. Determining volatile organics at microgram per litre levels by gas chromatography. *J. Am. Water Works Assoc.* 66:739-44.
- Bender, D. F. 1968. Thin-layer chromatographic separation and spectrophotofluorometric identification and estimation of dibenzo[a,e] pyrene. *Environ. Sci. Technol.* 2:204-6.
- Blumer, M., and W. W. Youngblood. 1976. *Polycyclic Aromatic Hydrocarbons in the Environment: Homologous Series in Soils and Recent Marine Sediments*. NTIS No. AD A023637. Office of Naval Research.
- Brocco, D., V. Cantuti, and G. P. Cartoni. 1970. Determination of polynuclear hydrocarbons in atmospheric dust by a combination of thin-layer and gas chromatography.
- Caton, R. D. Jr., J. B. Matthews, and E. A. Walters. 1976. *Development of High Pressure Liquid Chromatographic Techniques*. NTIS No. AD/A039644. Tyndall AFB, Fla.: Air Force Civil Engineering Center, Air Force Systems Command.
- Chang, R. C. 1976. *Concentration and Determination of Trace Organic Pollutants in Water*. Ph.D. dissertation, Ames Laboratory, Iowa State University.
- Chiba, M., and H. V. Mosley. 1968. Studies of losses of pesticides during sample preparation. *J. Assoc. Off. Anal. Chem.* 51:55-62.
- Freudenthal, R. I., A. P. Leber, D. Emmerling, and P. Clarke. 1975. The use of high pressure liquid chromatography to study chemically induced alterations in the pattern of benzo[a]pyrene metabolism. *Chem. - Biol. Interact.* 11:449-58.

- Giger, W., and M. Blumer. 1974. Polycyclic aromatic hydrocarbons in the environment: Isolation and characterization by chromatography, visible ultraviolet, and mass spectrometry. *Anal. Chem.* 46:1663-71.
- Goldberg, M. C., L. Delong, and M. Sinclair. 1973. Extraction and concentration of organic solutes from water. *Anal. Chem.* 45:89-93.
- Grant, D. W., and R. B. Meiris. 1977. Application of thin-layer and high-performance liquid chromatography to the separation of polycyclic aromatic hydrocarbons in bituminous materials. *J. Chromatogr.* 142:339-51.
- Hermann, T. S. 1974. *Development of Sampling Procedures for Polycyclic Organic Matter and Polychlorinated Biphenyls*. Washington, D.C.: Office of Research and Development, U. S. Environmental Protection Agency.
- IUPAC Applied Chemistry Division. 1974. Analytical methods for use in occupational hygiene: Determination of benzo[a]pyrene and benzo[k]fluoranthene in airborne particulates (chromatography and optical fluorescence). *Pure Appl. Chem.* 40:36.1-36.7.
- Janini, G. M., G. M. Muschik, and W. L. Zielinski, Jr. 1976. N,N'-Bis[p-butoxybenzylidene]- α,α' -bi-p-toluidine: thermally stable liquid crystal for unique gas-liquid chromatography separation of polycyclic aromatic hydrocarbons. *Anal. Chem.* 48:809-13.
- Jenkins, R. L., and R. B. Baird. 1975. The determination of benzidine in waste waters. *Bull. Environ. Contam. Toxicol.* 13:436-42.
- John, E. D., and G. Nickless. 1977. Gas chromatographic method for the analysis of major polynuclear aromatics in particulate matter. *J. Chromatogr.* 138:399-412.
- Junk, G. A., J. J. Richard, M. D. Grieser, D. Witiak, J. L. Witiak, M. D. Arguello, R. Vick, H. J. Suec, J. S. Fritz, and G. V. Calder. 1974. Use of macroreticular resins in the analysis of water for trace organic contaminants. *J. Chromatogr.* 99:745-62.
- Keith, L. H., A. W. Garrison, F. R. Allen, M. H. Carter, T. L. Floyd, J. D. Pope, and A. D. Thurston, Jr. 1976. Identification of organic compounds in drinking water from thirteen U. S. cities. In *Identification and Analysis of Organic Pollutants in Water*, ed. L. H. Keith. Ann Arbor, Mich.: Ann Arbor Science Publishers, Inc.
- Kelly, J. A. 1967. *The Determination of Phenolic-type Compounds in Water by High-pressure Liquid Chromatography*. NTIS No. ORD-4254-15. Ph.D. dissertation, Oklahoma State University.
- Kuo, P. P. K., E. S. K. Chian, J. H. Kim, and F. B. DeWalle. 1977. Study of the gas stripping, sorption, and thermal desorption procedures for pre-concentrating volatile polar organics from water samples for analysis by gas chromatography. *Anal. Chem.* 49:1023-29.

- Lao, R. C., R. S. Thomas, and J. L. Monkman. 1975. Computerized gas chromatographic-mass spectrometric analysis of polycyclic aromatic hydrocarbons in environmental samples. *J. Chromatogr.* 112:681-700.
- McGuire, J. M., A. L. Alford, and M. H. Carter. 1973. *Organic Pollutant Identification Utilizing Mass Spectrometry*. Report No. EPA-R2-73-234. Corvallis, Oregon: National Environmental Research Center, Office of Research and Monitoring, U.S. Environmental Protection Agency.
- Malcolm, R. L., E. M. Thurman, and G. R. Aiken. 1977. The concentration and fractionation of trace organic solutes from natural and polluted waters using XAD-8, methylmethacrylate resin. In *Proceedings of the 11th Annual Conference on Trace Substances in Environmental Health*. Columbia, Mo.: University of Missouri. In press.
- Mallet, V. N., P. E. Belliveau, and R. W. Frei. 1975. In situ fluorescence spectroscopy of pesticides and other organic pollutants. *Res. Dev.* 59:51-90.
- O'Reilly, W. F., and R. P. Murrmann. 1974. *Identification of Soil Organics using a Gas Chromatographic/Mass Spectrometric Method*. Washington, D.C.: Directorate of Military Engineering and Topography Office, Chief of Engineers, U. S. Army.
- Oswald, E. O., P. W. Albrow, and J. D. McKinney. 1974. Utilization of gas-liquid chromatography coupled with chemical ionization and electron impact mass spectrometry for the investigation of potentially hazardous environmental agents and their metabolites. *J. Chromatogr.* 98:363-448.
- Pierce, R. C., and M. Katz. 1975. Determination of atmospheric isomeric polycyclic arenes by thin-layer chromatography and fluorescence spectrophotometry. *Anal. Chem.* 47:1743-48.
- Sawicki, E. 1964. The separation and analysis of polynuclear aromatic hydrocarbons present in the human environment. *Chem. Anal.* 53:24-30.
- Severson, R. F., M. E. Snook, H. C. Higman, O. T. Chortyk, and F. J. Akin. 1976. Isolation, identification, and quantitation of the polynuclear aromatic hydrocarbons in tobacco smoke. In *Carcinogenesis--A Comprehensive Survey*, Vol. 1, ed. R. Freundenthal and P. W. Jones, pp. 253-70. New York: Raven Press.
- Stanley, T. W., D. F. Bender, and W. C. Elbert. 1973. Quantitative aspects of thin-layer chromatography in air pollution measurements. In *Quantitative Thin Layer Chromatography*, ed. J. C. Touchstone, pp. 305-22. New York: Wiley - Interscience.
- Stroupe, R. C., P. Tokousbalides, R. B. Dickinson, Jr., E. L. Wehry, and G. Mamantov. 1977. Low temperature fluorescence spectrometric determination of polycyclic aromatic hydrocarbons by matrix isolation. *Anal. Chem.* 49:701-5.

- Swain, A. P., J. E. Cooper, and R. L. Stedman. 1969. Large-scale fractionation of cigarette smoke condensate for chemical and biologic investigations. *Cancer Res.* 29:579.
- Treiber, L. R. 1976. An extension of the programmed multiple development (PMD) technique. *J. Chromatogr.* 124:69-72.
- U. S. Environmental Protection Agency [U. S. EPA]. 1977. *Sampling and Analysis Procedure for Screening of Industrial Effluents for Priority Pollutants*. Cincinnati, Ohio: Environmental Monitoring and Support Laboratory, U. S. EPA.
- Webb, R. G., A. W. Garrison, L. H. Keith, and J. M. McGuire. 1973. *Current Practice in GC-MS Analysis of Organics in Water*. NTIS No. PB-224-947. Southeast Environmental Research Laboratory, Athens, Ga.: U. S. Environmental Protection Agency.
- Woo, Ching S., A. P. D'Silva, V. A. Fassel, and G. J. Oestreich. 1978. Polynuclear aromatic hydrocarbons in coal--identification by their x-ray excited optical luminescence. *Environ. Sci. Technol.* 12:173-74.
- White, R. H., and J. W. Howard. 1967. Thin-layer chromatography of polycyclic aromatic hydrocarbons. *J. Chromatogr.* 29:108-14.
- Willis, R. B. 1973. *High Pressure Liquid Chromatography of Phenols and Metal Ions*. U. S. Atomic Energy Commission.

6 REVIEW AND INTERPRETATION OF ADSORPTION DATA

K. A. Reinbold and J. J. Hassett

A review of the literature revealed relatively few data on the adsorption of energy-related organic pollutants by sediments or soils. The summary presented here, therefore, covers literature on a variety of types of organic compounds, including pesticide data from references not covered by Farmer (1976).

Some of the published reports included constants for the sorption of organic compounds on various adsorbents. These constants are tabulated at the end of this chapter. However, many sources containing sorption data did not report sorption constants. The latter publications are included in the bibliography.

Except in the case of pesticides, most of the adsorption constants reported for organics were derived using adsorbents other than sediments, soils, and clay minerals. The most commonly used adsorbents were activated carbon or ion exchange resins, which are often used for the removal of organics in water treatment processes. Nylon and cellulose triacetate have also been used. The pesticide data, however, do pertain to sorption on sediments, soils, or clay minerals. The adsorption constants were determined in most cases by fitting experimentally derived data to various forms of the Freundlich or Langmuir adsorption equations.

FACTORS INFLUENCING ADSORPTION

Adsorption is a process in which a solution component is concentrated at the solid-solution interface. Adsorption results when the forces of attraction between the solution component and the surface, that is, the adsorbate-adsorbent interaction, overcomes the forces of attraction between the solution component and the solvent, that is, the solute-solvent interaction. There are two general cases where the adsorbate-adsorbent interaction is greater than the solute-solvent interaction and adsorption results.

In the first case there is a strong positive interaction between the surface and the adsorbate, and this interaction is strong enough to overcome even a fairly strong solute-solvent interaction. In this case adsorption is primarily related to the nature of the bonding between the adsorbate and the adsorbent. The adsorption of organic cations such as paraquat by clay minerals or polar organic molecules within the montmorillonitic interlayer are examples of this type of adsorption (Weber *et al.*, 1965). Reviews emphasizing the effect of adsorbate and adsorbent properties on adsorption have been written by Mortland (1970), Adams (1973), Bailey and White (1970), and Weber (1972).

In the second case, adsorption takes place not because of a strong adsorbate-adsorbent interaction, but rather due to a weak solute-solvent interaction. In this case even a small positive adsorbate-adsorbent interaction can overcome the solute-solvent interaction and result in adsorption. The degree of adsorption or partitioning of the adsorbing material between the solid and aqueous phases is primarily determined by the suitability of the aqueous phase as a solvent for the material. The poorer the aqueous phase as a solvent for the adsorbing species, the weaker the solute-solvent interaction and the greater the adsorption.

The adsorption of nonpolar aromatic hydrocarbons of low water solubility by organic surfaces is an example of this type of adsorption (Karickhoff *et al.*, 1979; Means *et al.*, 1979). This type of adsorption has been called "hydrophobic adsorption" because of the emphasis on the weak solute-solvent interaction in determining the degree of adsorption in aqueous systems (Horvath and Melander, 1978).

Hydrophobic or nonpolar adsorption can be considered an example of a nonpolar organic compound (adsorbate) partitioning between a polar aqueous phase and a stationary organic phase (adsorbent). In a soil or sediment system the aqueous phase would be the soil solution or the interstitial water in the sediment, while the organic phase would be the naturally occurring humic materials. Factors which either increase the affinity of the adsorbate for the humic surfaces or decrease the affinity of the adsorbate (solute) for the solvent (water) would result in greater adsorption.

Van der Waals forces have been identified as the main source of adsorbate-adsorbent interactions between nonpolar compounds and nonpolar organic surfaces (Horvath and Melander, 1978). The differential heats of adsorption for van der Waals forces are of the order of 1 to 2 kcal mole⁻¹ for small molecules; these forces may be much greater for larger molecules, especially with an increase in the number of double and triple bonds (Bailey and White, 1970). This is illustrated by the adsorption of methane, ethane, pentane and hexane by carbon black. These hydrocarbons gave differential heats of adsorption of 3, 4.3, 9.2 and 11.4 kcal mole⁻¹, respectively (Corkill *et al.*, 1966). The differential heats of adsorption do not increase indefinitely with increasing molecular size; eventually a point is reached where increasing molecular size does not increase adsorption and may even decrease adsorption due to steric hindrance.

Many factors influence the solute-solvent interaction and hence influence adsorption. Molecular properties such as chain length, molecular volume, molecular weight, carbon number, and polarity have all been shown to influence adsorption (Bailey and White, 1970; Gustafson and Paleos, 1971; Lailach *et al.*, 1968; Cummings *et al.*, 1959; Bartell and Miller, 1924; Kipling, 1965; Hansen and Craig, 1954; Parkash, 1974). It has been demonstrated that as chain length, molecular volume, molecular weight, and carbon number increase, and as polarity decreases, the solute-solvent interaction weakens and hydrophobic adsorption increases. The effect of all of these properties on the solute-solvent interaction is integrated into the water solubility of the compound.

The adsorption of hydrophobic compounds has been shown to increase with decreasing water solubility of the compound (Karickhoff *et al.*, 1979; Means *et al.*, 1979). The adsorption of more polar compounds has also been shown to increase with decreasing water solubility, but only within a family of compounds (Bailey and White, 1970). Aqueous solubilities of organic compounds, while useful in predicting adsorption, are often difficult to determine. Some of the difficulties encountered include the approach of the water solubilities to analytical detection limits, the formation of stable suspensions and the long equilibration times required to establish equilibrium.

Lambert (1968) discussed the similarity between the role of soil organic matter in the sorption of organic compounds and the role of an organic solvent in a liquid-liquid extraction. He observed that the partitioning of a nonpolar organic compound between the soil solution and soil organic matter was highly correlated with the partitioning of the compound between water and an organic solvent. Karickhoff *et al.* (1979) reported, for sorption placed on an organic carbon basis (Koc), a significant correlation between the sorption of several aromatic hydrocarbons and the partitioning of the compounds between octanol and water (Kow).

$$\log K_{oc} = 1.00 \log K_{ow} - 0.21 \quad (\text{Eq. 24})$$

The linear partition coefficients (Kp) for many compounds may be calculated from the following equation:

$$K_p = C_s/C_w \quad (\text{Eq. 25})$$

where

Cs = the concentration of the compound in the solid phase at equilibrium

Cw = equilibrium solution concentration of the compound

When the individual linear partition coefficients for the sorption of a hydrophobic organic compound by a variety of different sediments and soils are divided by the respective sediment or soil organic carbon contents, a unique constant Koc is produced.

$$K_{oc} = \frac{K_p \times 100}{\%OC} \quad (\text{Eq. 26})$$

This constant is independent of soil or sediment (adsorbent) properties and is only dependent on the nature of the adsorbing species (Karickhoff *et al.*, 1979; Means *et al.*, 1979).

The relationship between Koc values and Kow values for hydrophobic compounds has several distinct merits. First, Kow determinations are more reliable than water solubility determinations, particularly for the highly hydrophobic compounds. Second, once a compound's Kow value has been measured or calculated, its Koc can be determined from equation 24. Third, if the organic carbon contents of the individual soils or sediments are known, then their respective Kp values for the adsorption of the compound can be calculated.

Table 5 gives an example of linear Kp values, measured Koc values and Koc values calculated from Kow values for the sorption of three hydrophobic compounds, pyrene, dibenzothiophene and acetophenone by a variety of sediments.

TABLE 5. Kow, CALCULATED Koc, LINEAR Kp AND MEASURED Koc VALUES FOR SORPTION OF PYRENE, DIBENZOTHIOPHENE AND ACETOPHENONE BY SOILS AND SEDIMENTS

Compound	Kow	Koc (calc'd)	Sample No.	%OC	Kp	Koc (meas'd)	Average Koc
Pyrene	124,000	76,400	B2	1.21	774	63,991	63,400
			4	2.07	1098	53,019	
			5	2.28	1191	52,250	
			6	0.72	633	87,847	
			8	0.15	125	83,333	
			9	0.11	79	71,818	
			14	0.48	285	59,271	
			15	0.95	783	82,453	
			18	0.66	509	77,182	
			20	1.30	747	57,469	
			21	1.88	1159	61,628	
			22	1.67	811	48,557	
			23	2.38	1130	47,487	
			26	1.48	1023	69,108	
Dibenzo- thiophene	24,000	14,700	B2	1.21	117.5	9711	11,230
			4	2.07	180.6	8725	
			5	2.28	167.1	7329	
			6	0.72	60.8	8444	
			8	0.15	9.4	6267	
			9	0.11	5.8	5273	
			14	0.48	49.7	10,354	
			15	0.95	179.9	18,937	
			18	0.66	65.1	9864	
			20	1.30	101.4	7800	
			21	1.88	276.0	14,681	
			22	1.67	176.3	10,557	
			23	2.38	388.6	16,328	
			26	1.48	134.5	9088	
Aceto- phenone	38.6	23.8	B2	1.21	0.44	36	38.6
			4	2.07	0.89	43	
			5	2.28	0.56	24	
			6	0.72	0.68	95	
			8	0.15	0.07	48	
			9	0.11	0.09	82	
			14	0.48	0.12	25	
			15	0.95	0.27	28	
			18	0.66	0.30	46	
			20	1.30	0.29	22	
			21	1.88	0.85	45	
			22	1.67	0.53	31	
			23	2.38	0.68	29	
			26	1.48	0.66	45	

The range in Kp values for the adsorption of hydrophobic compounds depends on the compound being adsorbed and the range in organic carbon contents found in the soils or sediments studied. The upper limit for Koc values depends on the compound, but appears to be around 2,000,000 due to present analytical chemistry limitations. The lower limit for the validity of the Koc-Kow relationship has not yet been defined. This limit will be reached when case 1 adsorption, i.e., a specific strong adsorbate-adsorbent interaction, is encountered.

A relationship has been demonstrated for nonlinear Freundlich isotherms when the data are expressed on a molar basis instead of a mass basis (Osgerby, 1970). Molar Kd values may be calculated from mass Kd values by the following equation:

$$K_d(\text{Molar}) = \frac{K_d(\text{Mass}) \times \text{Molecular weight}^{1/n}}{\text{Molecular weight}} \quad (\text{Eq. 27})$$

Where Kd(Mass) and 1/n are Freundlich constants and the molecular weight is that of the adsorbate.

INTERPRETATION OF TABULATED DATA

Various combinations of the influencing factors discussed above are necessary to explain the adsorption data obtained from the literature and tabulated at the end of this chapter. For the nonpesticide organics, the adsorbents used in most of the studies reviewed were montmorillonite clay or carbon rather than natural sediments or soils. Thus, rather than a variety of adsorbents with a range of characteristics, two distinctly different types are represented: an inorganic clay mineral and an organic.

The chemical characteristics of the adsorbate account for much of the variation in adsorption behavior shown in the tabulated data. These effects were best shown in studies of groups of related compounds. Linner and Gortner (1935) studied the adsorption of 31 organic acids on carbon; the results are summarized in Table 6. Traube's rule that adsorption from aqueous solutions increases with molecular weight for a homologous series was demonstrated for fatty acids but not for other acids. The branched chain had little effect on the maximum adsorption of the acids, but the double bond showed a tendency to decrease adsorption. The introduction of polar groups--carboxyl, hydroxyl, or keto--caused decreased adsorption. The decrease was more pronounced as the number of carboxyl groups increased or as a second hydroxyl group was introduced. The decrease in adsorption caused by the keto group was dependent on the length of the chain and the position of the polar group in the chain.

For 52 structurally related N-phenylcarbamates, acetanilides, and anilines from aqueous 2% ethanol solutions, the inverse relationship between solubility and adsorption accounted for 60 percent of the total variation in adsorption, shown by the data in Table 6 (Ward and Upchurch, 1965). All of the compounds were adsorbed on nylon and cellulose triacetate. Other than solubility, the principal variable factor affecting adsorption was the difference in the molecular structures of the compounds. An investigation of

compounds having systematic variations in molecular structure shows which sites in the molecule can be involved in adsorptive processes and how various substituents may influence the extent of adsorption. For those compounds, differences in molecular structure caused differences in adsorption as a result of steric hindrance, tautomerism, chelation, and induction. The results suggest that the preferred adsorption mechanism of the amido compounds from aqueous solution is via the adsorbate's imino hydrogen and the adsorbent's carbonyl oxygen. If neither of these is available, however, alternative binding sites are utilized.

As a means of establishing a quantitative relationship between soil sorption equilibria and chemical structure, Lambert (1967) proposed the following relationship between parachor of uncharged organic chemicals for which no appreciable hydrogen bonding occurs and the soil sorption of those chemicals:

$$K_e = aP^bU \quad (\text{Eq. 25})$$

The relationship is based upon extrathermodynamic linear free energy approximations and uses of parachor as an approximate measure of the molar volume of the chemical. Distribution equilibria between soil and water for a number of chemical homologs of two chemical classes, including anilines (see Table 6) were used to establish the relationship. The relationship emphasizes the importance of using the partition or distribution coefficient, defined with respect to organic matter, as the most representative index of soil sorption equilibria.

Hance (1969) extended this relationship. A factor given by (parachor-45N), where N is the number of sites in a molecule which can participate in the formation of a hydrogen bond, was correlated with the logarithm of the Freundlich K value for the adsorption of 29 aromatic herbicides. Again, the relationship is valid for soils in which organic matter is the dominant adsorbing constituent.

In the case of a few compounds, adsorption constants were determined for more than one adsorbent (Table 6). Where data were obtained on both a clay mineral and a sediment, the adsorption was generally greater on the sediment. Some data on sediments and soils, including those for benz(a)pyrene, pyrene, methoxychlor, and carbaryl, show a correlation between adsorption and organic matter content (Table 6) (Smith, Mabey, Bohonos, Holt, Lee, Chou, Mill, and Bomberger, 1976a; Karickhoff, Brown, and Scott, 1978; La Fleur, 1976a).

In addition to the adsorption constants for specific compounds and adsorbents, additional pertinent data are tabulated here. Adsorption constants for a number of pesticides averaged over several soils are shown in Table 7. Also, Tables 8 and 9 show the inverse relationship between R_f values obtained by soil thin-layer chromatographs and adsorption or adsorbate characteristics.

(Note: the references cited in this chapter are listed beginning on p. 110.)

TABLE 6: ADSORPTION CONSTANTS FOR ORGANIC COMPOUNDS

Compound Name				Ref.
<u>COMPOUNDS OTHER THAN PESTICIDES</u>				
<u>Acids, Aliphatic</u>				
acetic acid	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		Linner and Gortner, 1935
$C_2H_4O_2$	nordite			
<u>Compound properties</u> ^a	(decolorizing carbon)			
m.w. 60.05		$\alpha\beta$	α	β
m.p. 16.604°C		0.462	0.266	1.736
b.p. 117.9°C		Adsorption equation: $a = \alpha C^{1/n}$		
density 1.0492 ₄ ⁰		α	$1/n$	
water sol ∞		2.46	0.351	

Experimental Conditions: 1 gm of adsorbent and
concentrations or reagents varying from 0.01
to 0.25 molar.

^aSee notes at end of table

TABLE 6: Continued

Compound Name					Ref.
adipic acid					Linner and Gortner, 1935
$C_6H_{10}O_4$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$			
	nordite				
	(decolorizing carbon)	$\alpha\beta$	α	β	
<u>Compound properties</u>		2.347	1.886	1.245	
m.w. 146.14 ^a		Adsorption equation: $a = \alpha C^{1/n}$			
m.p. 151°C ^a					
b.p. 265°C at 100mm ^a		α	1/n		
density 1.360 ₄ ^{5a}		1.79	0.163		
water sol 1.5 ^{15b}					
Experimental Conditions: same as for acetic acid.					
butyric acid					Linner and Gortner, 1935
$C_4H_8O_2$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$			
	nordite				
	(decolorizing carbon)	$\alpha\beta$	α	β	
<u>Compound properties</u> ^a		1.689	0.863	1.957	
m.w. 88.12°C		Adsorption equation: $a = \alpha C^{1/n}$			
m.p. -4.26°C					
b.p. 163.53°C		α	1/n		
density 0.9577 ₄ ²⁰		2.46	0.177		
water sol ∞					
Experimental Conditions: same as for acetic acid.					

TABLE 6: Continued

Compound Name		Ref.
butyric acid		Spahn et al., 1974
$C_4H_8O_2$	<u>Adsorbent</u> activated charcoal B10	Adsorption equation: $y = a \cdot x^b$
<u>Compound properties</u> ^a		
m.w. 88.12		x - range I (mmol/dm ³)
m.p. -4.26 °C		a b
b.p. 163.53 °C		0.08 - 7.18 0.878 0.4
density 0.9577 ₄ ²⁰		
water sol =		
n-butyric acid		Gustafson et al., 1968
	<u>Adsorbent</u> XAD-2 (amberlite resin)	Adsorption equation: $\frac{c}{q} = \frac{1}{Kb} + \frac{c}{b}$ (Langmuir)
		K(25°C) b(25°C)
		23.7 liter/mole 1.46X10 ⁻³ mole/g
		Enthalpy of adsorption:
		$\Delta H^\circ = 2.303 \frac{RT_1 T_2}{(T_2 - T_1)} (\log K'_2 - \log K'_1)$
		q -ΔH°*
		7X10 ⁻⁴ $\frac{\text{mole}}{q}$ 4.1 kcal/mole
		9X10 ⁻⁴ $\frac{\text{mole}}{q}$ 2.4 kcal/mole
		1.2X10 ⁻³ $\frac{\text{mole}}{q}$ 2.1 kcal/mole

*ΔH decreases with increasing surface coverage, i.e. energetically preferred sites are utilized first.

TABLE 6: *Continued*

Compound Name		Ref.
caproic acid		Linner and Gortner, 1935
$\text{CH}_3(\text{CH}_2)_4\text{CO}_2\text{H}$	<u>Adsorbent</u> nordite (decolorizing carbon)	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$
<u>Compound properties</u>		
m.w. 116.16 ^a		$\alpha\beta$ α β 8.772 4.636 1.892
m.p. -2 to -1.5°C ^a		Adsorption equation: $a = \alpha C^{1/n}$
b.p. 205°C ^a		
density 0.9274 ₄ ^a		α $1/n$ 3.03 0.175
water sol 0.4 gm/100 ml ^b		

Experimental Conditions: same as for acetic acid.

citraconic acid		Linner and Gortner, 1935
$\text{C}_5\text{H}_6\text{O}_4$	<u>Adsorbent</u> nordite (decolorizing carbon)	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$
<u>Compound properties</u>		
m.w. 130.10 ^a		$\alpha\beta$ α β 1.356 1.014 1.337
m.p. 91°C ^b		Adsorption equation: $a = \alpha C^{1/n}$
density 1.617 ^a		α $1/n$ 1.69 0.167
water sol 238 cold ^b		

Experimental Conditions: same as for acetic acid.

TABLE 6: Continued

Compound Name				Ref.
citric acid				Linner and Gortner, 1935
$C_6H_8O_7$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		
<u>Compound properties</u>	nordite (decolorizing carbon)	$\alpha\beta$	α	β
m.w. 192.14 ^a		1.444	2.757	0.524
m.p. 153°C (anhydrous) ^a		Adsorption equation: $a = \alpha C^{1/n}$		
b.p. decomp ^a		α	$1/n$	
density 1.542 ^{1ab}		0.73	0.203	
water sol 133 cold gm/100 ml ^b				

Experimental Conditions: same as for acetic acid.

dibromosuccinic acid				Linner and Gortner, 1935
$C_4H_4O_2Br_2$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		
<u>Compound properties</u> ^b	nordite (decolorizing carbon)	$\alpha\beta$	α	β
m.w. 275.90		1.397	1.119	1.248
m.p. 151-3°C		Adsorption equation: $a = \alpha C^{1/n}$		
		α	$1/n$	
		2.58	0.320	

Experimental Conditions: same as for acetic acid.

TABLE 6: Continued

TABLE 6: CONTINUED

Compound Name			Ref.						
formic acid			Linner and Gortner, 1935						
HCO_2H	<u>Adsorbent</u> nordite (decolorizing carbon)	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$							
<u>Compound properties</u> ^a		<table><tr><th>$\alpha\beta$</th><th>α</th><th>β</th></tr><tr><td>0.273</td><td>0.159</td><td>1.710</td></tr></table>	$\alpha\beta$	α	β	0.273	0.159	1.710	
$\alpha\beta$	α	β							
0.273	0.159	1.710							
m.w. 46.03									
m.p. 8.4°C									
b.p. 100.7°C		Adsorption equation: $a = \alpha C^{1/n}$							
density 1.220 ₄ ²⁰		<table><tr><th>α</th><th>$1/n$</th></tr><tr><td>2.47</td><td>0.435</td></tr></table>	α	$1/n$	2.47	0.435			
α	$1/n$								
2.47	0.435								
water sol ∞									

Experimental Conditions: same as for acetic acid.

fumaric acid				Linner and Gortner, 1935						
$C_4H_4O_4$	<u>Adsorbent</u> nordite (decolorizing carbon)	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$								
<u>Compound properties</u>		<table><tr><td>$\alpha\beta$</td><td>α</td><td>β</td></tr><tr><td>7.097</td><td>5.798</td><td>1.224</td></tr></table>	$\alpha\beta$	α	β	7.097	5.798	1.224		
$\alpha\beta$	α	β								
7.097	5.798	1.224								
m.w. 116.07 ^b										
m.p. 287°C ^b										
b.p. 200 subl. ^b 165 subl. ^a										
density 1.635 ^{0b}										
water sol 0.70 ²⁵ ; 9.8 ^{100b}										
		Adsorption equation: $a = \alpha C^{1/n}$								
		<table><tr><td>α</td><td>$1/n$</td></tr><tr><td>2.81</td><td>0.248</td></tr></table>	α	$1/n$	2.81	0.248				
α	$1/n$									
2.81	0.248									

Experimental Conditions: same as for acetic acid.

TABLE 6: Continued

Compound Name			Ref.						
glutaric acid			Linner and Gortner,						
$C_5H_8O_4$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$	1935						
<u>Compound properties</u>	nordite (decolorizing carbon)	<table><tr><td>$\alpha\beta$</td><td>α</td><td>β</td></tr><tr><td>3.697</td><td>3.104</td><td>1.192</td></tr></table>	$\alpha\beta$	α	β	3.697	3.104	1.192	
$\alpha\beta$	α	β							
3.697	3.104	1.192							
m.w. 132.11 ^a									
m.p. 99°C ^a									
b.p. 304 decomp. ^b		Adsorption equation: $a = \alpha C^{1/n}$							
density 1.424 ^{25a}		<table><tr><td>α</td><td>1/n</td></tr><tr><td>1.96</td><td>0.201</td></tr></table>	α	1/n	1.96	0.201			
α	1/n								
1.96	0.201								
water sol 64 ^{20a}									
Experimental Conditions: same as for acetic acid.									
glyceric acid			Linner and Gortner,						
$C_3H_6O_4$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$	1935						
<u>Compound properties</u> ^a	nordite (decolorizing carbon)	<table><tr><td>$\alpha\beta$</td><td>α</td><td>β</td></tr><tr><td>0.668</td><td>0.812</td><td>0.823</td></tr></table>	$\alpha\beta$	α	β	0.668	0.812	0.823	
$\alpha\beta$	α	β							
0.668	0.812	0.823							
m.w. 106.08									
b.p. disintegrates		Adsorption equation: $a = \alpha C^{1/n}$							
water sol ∞		<table><tr><td>α</td><td>1/n</td></tr><tr><td>1.29</td><td>0.267</td></tr></table>	α	1/n	1.29	0.267			
α	1/n								
1.29	0.267								
Experimental Conditions: same as for acetic acid.									

TABLE 6: Continued

TABLE 6: Continued

Compound Name			Ref.						
glycolic acid			Linner and Gortner, 1935						
HOCH ₂ CO ₂ H	<u>Adsorbent</u> nordite (decolorizing carbon)	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$							
<u>Compound properties</u> ^a		<table><tr><td>$\alpha\beta$</td><td>α</td><td>β</td></tr><tr><td>0.239</td><td>0.249</td><td>0.958</td></tr></table>	$\alpha\beta$	α	β	0.239	0.249	0.958	
$\alpha\beta$	α	β							
0.239	0.249	0.958							
m.w. 76.05									
m.p. 80°C									
b.p. decomposes		Adsorption equation: $a = \alpha C^{1/n}$							
		<table><tr><td>α</td><td>$1/n$</td></tr><tr><td>1.54</td><td>0.390</td></tr></table>	α	$1/n$	1.54	0.390			
α	$1/n$								
1.54	0.390								
Experimental Conditions: same as for acetic acid.									
glyoxylic acid			Linner and Gortner, 1935						
C ₂ H ₂ O ₃	<u>Adsorbent</u> nordite (decolorizing carbon)	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$							
<u>Compound properties</u>		<table><tr><td>$\alpha\beta$</td><td>α</td><td>β</td></tr><tr><td>0.508</td><td>0.223</td><td>2.275</td></tr></table>	$\alpha\beta$	α	β	0.508	0.223	2.275	
$\alpha\beta$	α	β							
0.508	0.223	2.275							
m.w. 74.04 ^a									
m.p. 70-5°C ^a (+1/2w) 98 (anhydrous)		Adsorption equation: $a = \alpha C^{1/n}$							
water sol very soluble ^b		<table><tr><td>α</td><td>$1/n$</td></tr><tr><td>3.89</td><td>0.455</td></tr></table>	α	$1/n$	3.89	0.455			
α	$1/n$								
3.89	0.455								
Experimental Conditions: same as for acetic acid.									

TABLE 6: Continued

TABLE 6: Continued									
Compound Name			Ref.						
isobutyric acid			Linner and Gortner,						
(CH ₃) ₂ CHCO ₂ H	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$	1935						
<u>Compound properties</u>	nordite (decolorizing carbon)	<table><tr><td>$\alpha\beta$</td><td>α</td><td>β</td></tr><tr><td>0.883</td><td>0.497</td><td>1.776</td></tr></table>	$\alpha\beta$	α	β	0.883	0.497	1.776	
$\alpha\beta$	α	β							
0.883	0.497	1.776							
m.w. 88.11 ^f		Adsorption equation: $a = \alpha C^{1/n}$							
m.p. -46.1°C ^a		<table><tr><td>α</td><td>1/n</td></tr><tr><td>2.36</td><td>0.273</td></tr></table>	α	1/n	2.36	0.273			
α	1/n								
2.36	0.273								
b.p. 153.7°C ^a									
density 0.96815 ²⁰ ₄ ^a ; .949 ²⁰ ₄ ^b									
water sol 20 ²⁰ gm/100 ml ^b									

Experimental Conditions: same as for acetic acid.

isovaleric acid				Linner and Gortner,
$\text{C}_5\text{H}_{10}\text{O}_2$	<u>Adsorbent</u>	Adsorption equation:	$a = \frac{\alpha\beta C}{1+\alpha C}$	1935
<u>Compound properties</u>	nordite (decolorizing carbon)			
m.w. 102.13 ^a		$\alpha\beta$	α	β
m.p. -29.3°C ^a		1.630	0.902	1.807
b.p. 176.7°C ^a				
density 0.9286 ²⁰ ₄ ^a		Adsorption equation:	$a = \alpha C^{1/n}$	
water sol 4.2 ²⁰ ₄ ^b		α	1/n	
		2.51	0.227	

Experimental Conditions: same as for acetic acid.

TABLE 6: *Continued*

Compound Name			Ref.
itaconic acid			Linner and Gortner,
$C_5H_6O_4$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$	1935
<u>Compound properties</u>	nordite (decolorizing carbon)		
m.w. 130.10 ^a		$\alpha\beta$ α β	
m.p. 175°C ^a		1.167 0.904 1.291	
b.p. decomposes ^a			
density 1.632 ^b		Adsorption equation: $a = \alpha C^{1/n}$	
water sol 8.33 ^{20b}		α $1/n$	
		1.54 0.148	

Experimental Conditions: same as for acetic acid.

lactic acid (DL)			Linner and Gortner,
$C_3H_6O_3$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$	1935
<u>Compound properties</u>	nordite (decolorizing carbon)		
m.w. 90.08 ^a		$\alpha\beta$ α β	
m.p. 18°C ^a		0.437 0.415 1.054	
b.p. 122°C ^a			
density 1.249 ^{15b}		Adsorption equation: $a = \alpha C^{1/n}$	
water sol ∞ ^b		α $1/n$	
		1.66 0.335	

Experimental Conditions: same as for acetic acid.

TABLE 6: Continued

Compound Name				Ref
levulinic acid				
$C_5H_8O_3$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		
<u>Compound properties</u>	nordite (decolorizing carbon)	$\alpha\beta$	α	β
m.w. 116.13 ^a		2.990	2.289	1.307
m.p. 37.2 °C ^a				
b.p. 246°C slight decomp. ^a		Adsorption equation: $a = \alpha C^{1/n}$		
density 1.1335 ^{20a} ; 1.1395 ^{20b}		α	1/n	
water sol very sol ^b		1.83	0.183	
Experimental conditions: same as for acetic acid				
maleic acid				
$C_4H_4O_4$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		
<u>Compound properties</u>	nordite (decolorizing carbon)	$\alpha\beta$	α	β
m.w. 116.07 ^a		1.233	0.884	1.395
m.p. 139-140°C ^a				
b.p. 135 decomp. ^b		Adsorption equation: $a = \alpha C^{1/n}$		
density 1.590 ^{20ba}		α	1/n	
water sol 78.8 ²⁵ ; 392.6 ^{97.5b}		1.90	0.203	
Experimental conditions: same as for acetic acid				

Linner
and Gortner,
1935Linner
and Gortner,
1935

TABLE 6: Continued

Compound Name			Ref
malic acid (l)			
$C_4H_6O_5$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$	Linner and Gortner,
<u>Compound properties</u> ^b	nordite		1935
m.w. 134.09	(decolorizing carbon)	$\alpha\beta \quad \alpha \quad \beta$	
m.p. 100°C		0.531 0.574 0.927	
b.p. 140°C decomp.			
density 1.595		Adsorption equation: $a = \alpha C^{1/n}$	
water sol very sol		$\alpha \quad 1/n$	
		1.28 0.252	
Experimental conditions: same as for acetic acid			
malonic acid			
$C_3H_4O_4$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$	Linner and Gortner,
<u>Compound properties</u>	nordite		1935
m.w. 104.06 ^b	(decolorizing carbon)	$\alpha\beta \quad \alpha \quad \beta$	
m.p. 135.6°C subl. ^b		1.897 1.540 1.232	
b.p. decomp. at 140°C ^a			
density 1.631 ₄ ^{15b} ; 1.619 ^a		Adsorption equation: $a = \alpha C^{1/n}$	
water sol 61.1 ₂₀ ⁰ gm/100 ml ^b		$\alpha \quad 1/n$	
73.5 ₂₀ ⁰ gm/100 ml ^b		3.88 0.410	
92.6 ₂₀ ⁰ gm/100 ml ^b			
Experimental conditions: same as for acetic acid			

TABLE 6: Continued

Compound Name				Ref
mesaconic acid				Linner and Gortner, 1935
$C_5H_6O_4$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		
<u>Compound properties</u>	nordite (decolorizing carbon)	$\alpha\beta$	α	β
m.w. 130.10 ^a		2.706	1.886	1.435
m.p. 204.5		Adsorption equation: $a = \alpha C^{1/n}$		
b.p. 250°C decomp. ^b		α	$1/n$	
density 1.466 ²⁰ g/ml ^b		1.80	0.133	
water sol 2.7 ¹⁸ ^b ; 118 ¹⁰⁰ ^b				
Experimental conditions: same as for acetic acid				
methylosuccinic acid				Linner and Gortner, 1935
$C_5H_8O_4$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		
<u>Compound properties</u> ^b	nordite (decolorizing carbon)	$\alpha\beta$	α	β
m.w. 132.11		0.664	0.608	1.092
m.p. 111°C		Adsorption equation: $a = \alpha C^{1/n}$		
b.p. decomp.		α	$1/n$	
density 1.410g/ml		1.30	0.172	
water sol 66.7 ²⁰				
Experimental conditions: same as for acetic acid				

TABLE 6: Continued

Compound Name				Ref
monobromosuccinic acid				
$C_4H_5O_4Br$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		Linner and Gortner,
<u>Compound properties</u>	nordite			1935
m.w. 197.00 ^b	(decolorizing carbon)	$\alpha\beta$ α β		
m.p. 159°C ^b 161°C ^a		0.934 0.643 1.451		
density 2.073 ^a				
water sol 19 ^{15b}		Adsorption equation: $a = \alpha C^{1/n}$		
		α $1/n$		
		1.82 0.195		
	Experimental conditions: same as for acetic acid			
oxalic acid				
$C_2H_2O_4$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		Linner and Gortner,
<u>Compound properties</u>	nordite			1935
m.w. 90.04 ^a	(decolorizing carbon)	$\alpha\beta$ α β		
m.p. 189.5°C ^a		0.440 0.332 1.325		
b.p. 157°C subl. ^a				
density 1.900 ^{17a}		Adsorption equation: $a = \alpha C^{1/n}$		
water sol 19.5 ¹⁵ gm/100ml ^b		α $1/n$		
120 ⁹⁰ gm/100ml ^b (hydrated form)		3.62 0.551		
	Experimental conditions: same as for acetic acid			

TABLE 6: Continued

Compound Name			Ref.
propionic acid			
$C_3H_6O_2$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$	Linner and Gortner,
<u>Compound properties</u> ^a	nordite (decolorizing carbon)		1935
m.w. 74.08		$\alpha\beta$ α β	
m.p. -20.8°C		0.925 0.491 1.885	
b.p. 140.99°C			
density 0.9930 ²⁰		Adsorption equation: $a = \alpha C^{1/n}$	
water sol ∞		α $1/n$	
		2.46 0.236	

Experimental conditions: same as for acetic acid.

propionic acid				Spahn et al.,
$C_3H_6O_2$	<u>Adsorbents</u>	Adsorption equation: $y = a \cdot x^b$		
<u>Compound properties</u>	Activated charcoal B10	Adsorption constants:		1974
m.w. 74.08		x-range I		
m.p. -20.8°C		(mmol/dm ³) a b		
b.p. 140.99°C		0.1 - 8.3 0.497 0.4		
density 0.9930 ²⁰				
water sol ∞				

TABLE 6: Continued

Compound Name				Ref.
pyruvic acid				Linner and Gortner,
$C_3H_4O_3$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		1935
<u>Compound properties</u>	nordite (decolorizing carbon)			
m.w. 88.06 ^a		$\alpha\beta$ α β		
m.p. 13.6°C ^a		0.979 0.585 1.674		
b.p. 165°C slight decomp. ^b				
density 1.2272 ²⁰ ₄ ^a		Adsorption equation: $a = \alpha C^{1/n}$		
water sol ∞ ^b		α $1/n$		
		2.44 0.273		
Experimental Conditions: same as for acetic acid.				
succinic acid				Linner and Gortner,
$C_4H_6O_4$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		1935
<u>Compound properties</u> ^b	nordite (decolorizing carbon)			
m.w. 118.09		$\alpha\beta$ α β		
m.p. 185°C		0.865 0.467 1.854		
b.p. 235°C dissint.				
density 1.564 ¹⁵ ₄ g/ml		Adsorption equation: $a = \alpha C^{1/n}$		
water sol 6.8 at 20°C		α $1/n$		
121 at 100°C		2.44 0.273		
Experimental Conditions: same as for acetic acid.				

TABLE 6: Continued

Compound Name				Ref
tartaric acid				
$C_4H_6O_6$	<u>Adsorbent</u>	Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		Linner and Gortner,
<u>Compound properties</u>	nordite (decolorizing carbon)			1935
m.w. 150.09 ^b		$\alpha\beta$ α β		
m.p. 100°C ^b		0.322 0.468 0.687		
density 1.697 ^b ; 1.788 ^a				
water sol 20.6 ^{20b} ; 9.23 ^{0b} ; 185 ^{100b}		Adsorption equation: $a = \alpha C^{1/n}$		
		α $1/n$		
		0.94 0.275		
	Experimental conditions: same as for acetic acid			
valeric acid				
$CH_3(CH_2)_3CO_2H$	<u>Adsorbent</u>	Adsorption equation: $a = \alpha C^{1/n}$		Linner and Gortner
<u>Compound properties</u>	nordite (decolorizing carbon)			1935
m.w. 102.13 ^a		α $1/n$		
m.p. -33.83°C ^a		2.84 0.182		
b.p. 186.05°C ^a				
density 0.9391 ^{20a} ; 0.9424 ^{0b}		Adsorption equation: $a = \frac{\alpha\beta C}{1+\alpha C}$		
water sol 3.7gm/100ml ^b		$\alpha\beta$ α β		
		1.878 0.872 2.154		
	Experimental conditions: same as for acetic acid			

TABLE 6: Continued

Compound Name	Ref						
<u>Acids, Aromatic</u>							
benzoic acid							Ward and Getzen, 1970
$C_7H_6O_2$	<u>Adsorbents</u>	Adsorption equation: $\frac{x}{m} = \frac{abC}{(1+aC)}$					
<u>Compound properties</u>	Type BL - activated carbon (Pittsburgh Chemical Co.)	Adsorption constants:					
m.w. 122.13 ^a	Surface area 1000-1100 μ^2/g	<u>Compound</u>	<u>pka</u>	<u>pH of soln.</u>	<u>a*</u>	<u>b</u>	<u>x/m</u>
m.p. 122.4°C ^a		benzoic acid	4.20	3 7 11	0.238 0.108 0.081	510 124 75	489 113 67
b.p. 249°C ^b		-2,4-dichloro	2.76	3 7 11	0.259 0.123 0.108	676 159 73	651 147 69
density 1.2659 ^{15b}		-3-amino-2,5- dichloro (amiben)	3.40	3 7 11	0.928 0.283 0.025	515 131 72	510 127 51
water sol 0.18 ^{4b} 0.27 ^{18b} 2.2 ^{75b}		-3-nitro-2,5- dichloro	3.23	3 7 11	0.360 0.118 0.035	505 130 93	491 120 72
		-2-methoxy-3,6- dichloro (dicamba)	1.94	3 7 11	0.181 0.317 0.068	394 154 68	313 149 59
Experimental conditions: 100 ml aqueous acid solution: 10mg carbon							

Experimental conditions: 100 ml aqueous acid solution: 10mg carbon

*Units: a = liters/ μ mole
b = μ moles/g
c = μ moles/liter
 $\frac{x}{m}$ = μ moles/g

TABLE 6: Continued

Compound Name			Ref.
benzoic acid			Spahn et al., 1974
$C_7H_6O_2$	<u>Adsorbent</u> activated charcoal B10	Adsorption equation: $y = a \cdot x^b$	
<u>Compound properties</u>		Adsorption constant:	
m.w. 122.13 ^a		x-range I	
m.p. 122.4°C ^a		(mmol/dm ³)	
b.p. 249°C ^b		a	b
density 1.2659 ¹⁵ ₄ ^b		0.1 - 6	3.06 0.181
water sol 0.18 ⁴ _{0.27} ¹⁸ _{2.2} ⁷⁵			
benzoic acid and substituted benzoic acids	<u>Adsorbent</u> charcoal, activated, 200 mesh	Adsorption equation: $\log x/m = \log k + 1/n \log c$	Hartman, Kern, Bobalek, 1946
$C_7H_6O_2$			
		1/n and log k Values	
		log k	
Acids	1/n	20°	30° 40° 50°
Benzoic	0.3680	0.0613	0.0265 -0.1091 -0.2551
o-Chlorobenzoic	0.4060	0.2208	0.1834 0.1303 0.0936
o-Aminobenzoic	0.3693	0.3472	0.3031 0.2678 0.2218
o-Hydroxybenzoic	0.3840	0.2700	0.2385 0.2091 0.1844
o-Toluic	0.3879	-0.0215	-0.0719 -0.2510 -0.5232
m-Toluic	0.3581	-0.0092	-0.0582 -0.1791 -0.3732
p-Toluic	0.3696	0.1351	0.0762 -0.0324 -0.1886

Adsorption from benzene solutions.

TABLE 6: *Continued*

Compound Name		Ref.							
phenyl acetic acid		Spahn et al., 1974							
$C_6H_5CH_2CO_2H$									
<u>Compound properties^a</u>		<u>Adsorbent</u>	Adsorption equation: $y = a \cdot x^b$						
		Activated charcoal							
m.w. 136.16									
m.p. 77°C									
b.p. 265.5°C		Adsorption constants:							
density 1.091 ⁷⁷ ; 1.228 ⁴		x-range I (mmol/dm ³)		a	b	x-range II (mmol/dm ³)		a	b
water sol slightly sol.		B10	0.1 - 10	2.51	0.16	0.001 - 0.1	2.6	0.32	
		LW	0.05 - 12	1	0.165				
		Lev 634	0.1 - 8	0.84	0.38	0.01 - 0.11	1.76	0.56	
		BD	0.1 - 22	1.14	0.15				
		D1	0.05 - 7	1.93	0.162				
		D2	0.1 - 8	1.57	0.245				
		B2	0.1 - 11	1	0.145				
		LS-supra	0.1 - 7	1.75	0.18	0.001 - 0.11	1.53	0.23	
		Decaka 9	0.1 - 8	1.82	0.03				
		106/427/1	0.1 - 7	1.53	0.11	0.001 - 0.12	2.05	0.33	
		106/427/5	0.1 - 6.5	1.87	0.144	0.001 - 0.11	1.85	0.3	
		106/427/3	0.1 - 8	2.08	0.244	0.001 - 0.12	2.6	0.44	

TABLE 6: *Continued*

Compound Name			Ref.
<u>Carboxylic acids, aromatic</u>			
Phenoxyacetic acid			Ward and Getzen, 1970
$C_8H_8O_3$	<u>Adsorbents</u>	Adsorption equation: $\frac{x}{m} = \frac{abC}{(1+aC)}$	
<u>Compound properties</u>	Type BL - activated carbon (Pittsburgh Chemical Co.)	Adsorption constants:	
m.w. 152.14			
m.p. 99°C	Surface Area 1000-1100 μ^2/g	<u>Compound</u> <u>pka</u> <u>pH of soln.</u> <u>a*</u> <u>b</u> <u>$\frac{x}{m}$</u>	
b.p. 285°C slight decomp.		Phenoxyacetic Acid 3.03 3 0.423 446 436	
water sol 1.2 ¹⁰		7 0.229 105 86	
		11 0.209 64 43	
		-4-chloro- 2.36 3 0.362 575 560	
		7 0.101 198 180	
		11 0.115 145 133	
		-2,4-dichloro- 3.31 3 0.279 629 607	
		7 0.112 223 205	
		11 0.070 154 135	
		-2,4,6- trichloro- 3.35 3 0.672 676 666	
		7 0.095 239 216	
		11 0.073 144 127	
		2,4-dichloro- 3.92 3 0.431 645 630	
		7 0.125 235 218	
		11 0.100 59 54	

Experimental Conditions: 100 ml aqueous acid solution:
10 mg carbon.

Units: a = liters/ μ mole
b = μ moles/g
c = μ moles/liter
 $\frac{x}{m}$ = μ moles/g

TABLE 6: Continued

Compound Name						Ref.	
2,4-dichloro- phenylacetic acid						Ward and Getzen, 1970	
	<u>Adsorbents</u>	Adsorption equation: $\frac{x}{m} = \frac{abC}{(1+aC)}$					
	Type BL - activated carbon (Pittsburgh Chemical Co.)						
	Adsorption constants:						
	Surface area 1000-1100 μ^2/g	<u>pka</u>	<u>pH of soln.</u>	<u>a*</u>	<u>b</u>		<u>x/m</u>
		3.92	3	0.431	645		630
			7	0.125	235		218
		11	0.100	59	54		
Experimental conditions: 100 ml aqueous acid solution: 10 mg carbon							

*Units: a = liters/ μ mole
 b = μ moles/g
 c = μ moles/liter
 $\frac{x}{m}$ = μ moles/g

TABLE 6: Continued

Compound Name					Ref.
<u>Alcohol</u>					
n-butyl alcohol	<u>Adsorbent</u>	Adsorption equation: $\frac{x'}{m} = \frac{x}{m}(1+N_b \bar{v}_b / N_a \bar{v}_a)$			Bartell, Thomas, and Fu, 1951
C ₄ H ₉ OH	Graphite	N = mole fraction			
<u>Compound properties</u>	Blood Char.	\bar{v} = partial molar volume			
m.w. 74.12 ^c		a = solvent			
m.p. -90°C ^c		b = solute			
b.p. 117-118°C ^c ; 117.71°C ^b		<u>t°C</u>	<u>K*</u>	<u>ΔF°*** kcal/mole</u>	<u>ΔH°**** kcal/mole</u>
density 0.810 ₄ ^{0c}		0	63.3	-2.24	-0.16
		25	61.6	-2.43	-1.47
water sol 7.9 gm/100 ml ^b		45	52.7	-2.50	
		0	477	-3.34	-0.94
		25	412	-3.54	-0.52
		45	390	-3.76	

*Equilibrium constant K obtained from the limiting slope of the adsorption isotherms at zero concentration

$$**\Delta F^\circ = -RT \ln K$$

$$***\Delta H^\circ = R \left(\frac{T_2 T_1}{T_2 - T_1} \right) \ln \frac{K_2}{K_1}$$

TABLE 6: Continued

Compound Name		Ref.		
<u>Amides</u>				
acetanilide and derivatives, adsorption from 2% ethanol		Adsorption equation: $\left(\frac{x}{m}\right) = KC^{1/n}$ $1/n = 1$		Ward and Upchurch,
C_8H_9ON		solubility		1965
<u>Compound properties^b</u>	<u>Compound</u>	<u>(x 10⁻⁴M) 2% ethanol</u>	<u>k_N[*]</u>	<u>k_A^{**}</u>
m.w. 135.16	N-Ethylacetanilide	2320	0.05	0.20
m.p. 114°C	4-Hydroxyacetanilide	1580	0.25	0.13
b.p. 305°C	Acetanilide	624	0.21	0.32
density 1.21 ₄	2-Chloroacetanilide	375	0.25	0.47
water sol 0.563 ²⁵ ; 3.5 ⁸⁰	4-Aminoacetanilide	352	0.06	0.05
	N-n-Butylacetanilide	167	0.23	0.89
	2-Nitroacetanilide	138	0.16	0.72
	2-Hydroxyacetanilide	32	0.26	0.15
	N-Phenylacetanilide	31	0.26	1.59
	4-Chloroacetanilide	16	1.19	2.64
	3,4-Dichloroacetanilide	12	6.05	6.83
	4-Nitroacetanilide	11	1.02	2.18
	4-Bromoacetanilide	11	1.73	3.30
	2,5-Dichloroacetanilide	10	0.92	1.59

Method of analysis: UV spectrophotometry with Beckman DU spectrophotometer

* k_N = adsorption on nylon** k_A = adsorption on cellulose triacetate

TABLE 6: *Continued*

Compound Name		Ref.		
aniline and derivatives, adsorption from 2% ethanol		Adsorption equation: $(\frac{x}{m}) = KC^{1/n}$; $1/n = 1$		
$C_6H_5NH_2$		solubility ($\times 10^4 M$) 2% ethanol		
<u>Compound properties</u>		k_N^* k_A^{**}		
m.w. 93.13 ^a	Aniline	4450	0.07	0.27
m.p. -6.3°C ^a	2-Methylaniline	1670	0.10	0.44
b.p. 184.13°C ^a	4-Methylaniline	1240	0.05	0.50
density 1.02173 ₄ ^{0a}	N-Methylaniline	560	0.24	0.92
water sol 3.4 ²⁰ ; 6.4 ^{90b}	4-Chloroaniline	520	0.68	3.31
	2-Chloroaniline	450	0.55	1.94
	N-Ethylaniline	250	0.32	1.10
	2-Nitroaniline	103	0.91	2.84
	N,N-Dimethylaniline	95	0.52	2.12
	2,3-Dichloroaniline	85	3.45	7.20
	3,4-Dichloroaniline	52	4.32	10.20
	2,5-Dichloroaniline	49	3.78	11.20
	4-Nitroaniline	38	1.10	3.40
	N-n-Butylaniline	16	1.42	6.55
	2-Nitro-4-chloroaniline	9	3.70	7.80
	2,4-Dinitroaniline	5	1.94	5.65
	N-Benzylaniline	4	3.46	19.48
	N-Phenylaniline	3	16.50	24.00

Method of analysis: UV spectrophotometry with Beckman DU spectrophotometer.

* k_N = adsorption on nylon

** k_A = adsorption on cellulose triacetate

TABLE 6: *Continued*

Compound Name	TABLE 6. Continued					Ref.
4-(R ₂ sulfonyl)-2,6-dinitro- <i>N,N</i> -(di-R ₁) aniline	Adsorption equation: $\ln K_e = aPAU^*$					Lambert, 1967
	K _e values					
	SD	R ₁	R ₂	Ripperdan soil 1% 0.m	Sacramento soil 5% 0.m	
	11830	CH ₃	CH ₃	125	145	
	12639	C ₂ H ₅	CH ₃	230	193	
	11831	C ₃ H ₇	CH ₃	500	520	
	13207	C ₂ H ₅	C ₂ H ₅	320	269	
	12030	C ₃ H ₇	C ₂ H ₅	750	702	
	12346	C ₃ H ₇	C ₃ H ₇	1170	-	
	12400	CH ₃	iso-C ₃ H ₇	222		

*K_e = equilibrium constant, estimated from adsorption data

$P = \frac{M\gamma^{1/4}}{\rho}$, P = parachor, ρ = density of the liquid, γ = surface tension, M = molecular weight

ΔU = difference in internal pressures of the solvent phases

TABLE 6: *Continued*

Compound Name		Ref.	
crystal violet (gentian violet)	<u>Adsorbent</u>	Adsorption equation: $y = a \cdot x^b$	Spahn et al.
$C_{25}H_{30}ClN_3$	activated charcoal B10		1974
<u>Compound properties^c</u>		Adsorption constants:	
m.w. 408.06		x-range I	
water sol soluble		$(mmol/dm^3)$ <u>a</u> <u>b</u>	
		0.1 - 2 0.84 0.09	
<u>Amino acid</u>			
dl-tryptophan, from aqueous urea solution	<u>Adsorbent</u>	Adsorption equation: $M = \frac{abC}{1+bC}$	Nogami, Nagai, and Uchida,
$C_{11}H_{12}N_2O_2$	carbon black		1968
<u>Compound properties</u>			
m.w. 204.22	Urea Concentration M	$\frac{a}{10^{-3} \text{ liter}} \frac{b}{mole}$	$\frac{\Delta F}{Kcal} \frac{\Delta H}{mole} \frac{\Delta S}{e.u.} \frac{\Delta S_t}{e.u.} \frac{\Delta S_s}{e.u.}$
m.p. 283-5°C	0	1.63	1.70
water sol slightly sol cold sol hot	0.11	1.60	1.70
	1.01	1.54	1.08
	3.05	1.28	0.615
	5.03	1.18	0.472
	7.07	0.92	0.397
			-4.45
			0.22
			15.4
			0
			15.4
			15.4
			14.9
			14.7
			14.6
			14.6
			12.5
			2.08
			14.6

ΔF = free energy change
 ΔS = entropy change
 $\Delta S_s = \Delta S + \Delta S_t$
 ΔS_t = entropy change of the transfer of tryptophan from
aqueous solution to aqueous urea solution
e.u. = entropy units

TABLE 6: Continued

Compound Name		Ref.		
<u>Benzene</u>				
benzene - adsorption from binary mixture		Adsorption equation:		
C_6H_6		$\frac{X_1 X_2}{n^\circ \Delta X/m} = \frac{1}{n^s} \left[X_1 + \frac{1}{K-1} \right]$		
<u>Compound properties</u>		Zettlemoyer and Micale, 1971		
m.w. 78.11 ^d				
density .87901 ^b				
m.p. 5.5°C ^e				
b.p. 80.1°C ^e				
water sol 820 ppm at 22°C ^f				
OCT/water part coeff 130 ^g				
<u>Solvent</u>	<u>Adsorbent</u>	<u>K</u>	<u>Area of adsorbent</u> <u>m²/g</u>	<u>($\Delta H_1^\circ - \Delta H_2^{\circ*}$)</u> <u>cal/g</u>
Ethylene dichloride	Graphon	18.5	119	0.54
	Spheron 6	5	57	-
Cyclohexane	Coconut shell charcoal	19.9	677	12.0
	Decolorizing charcoal	16.4	596	7.1
Carbon tetrachloride	Decolorizing charcoal	16.1	374	4.0
	Coconut shell charcoal	18.2	350	3.8

$$*\Delta H = X_1^s \Delta H_1^\circ + X_2^s \Delta H_2^\circ$$

$$K = \exp \left(- \left(\frac{\Delta H_a}{RT} - \frac{\Delta S_a}{R} \right) \right)$$

TABLE 6: *Continued*

Compound Name		Ref.		
carbamate derivatives, adsorption from 2% ethanol		Adsorption equation: $\frac{x}{m} = KC^{1/n}$ (1/n = 1)		
		Ward and Upchurch, 1965		
Adsorbent	Compound	k_N		
		10°C	26.5°C	50°C
Nylon (Zytel-101) (80 mesh)	Ethyl- <i>N</i> -phenyl	1.03	1.03	1.03
	Isopropyl- <i>N</i> -phenyl	1.16	1.16	1.16
	Ethyl- <i>N</i> -methyl- <i>N</i> -phenyl	0.22	0.22	0.22
	Ethyl- <i>N</i> -ethyl- <i>N</i> -benzyl	0.25	0.25	0.25
	Ethyl- <i>N</i> -benzyl- <i>N</i> -phenyl	1.33	1.85	1.85
	Ethyl- <i>N,N</i> -diphenyl	0.66	1.02	1.02
	Isopropyl- <i>N</i> -(3-chlorophenyl)	9.00	9.00	9.00
	Ethyl- <i>N</i> -(2-nitrophenyl)	0.75	0.90	0.90
	Ethyl- <i>N</i> -(4-nitrophenyl)	3.65	3.70	3.70
	Ethyl- <i>N</i> -(2,3-dichlorophenyl)	3.70	3.70	3.70

Method of analysis: UV spectrophotometry with Beckman DU spectrophotometer.

TABLE 6: Continued

Compound Name		Ref.		
<hr/>				
<u>Carbonyls</u>				
acetophenone				Spahn et al.,
$C_6H_5COCH_3$	<u>Adsorbent</u>	Adsorption equation: $y = a \cdot x^b$		
<u>Compound properties</u>	activated charcoal B 10			1974
m.w. 120.16 ^h		Adsorption constants:		
m.p. 20.5°C ^h		x-range I		
b.p. 202.0°C ^h		(mmol/dm ³)	a	b
density 1.0281 at 20°C/4°C ^h		0.1 - 4	3.46	0.155
water sol 5.6x10 ⁻³ M at 25°C ⁱ				
 <u>N-Heterocyclics</u>				
benzo[f]quinoline				Smith, Mabey, Bohonos, Holt, Lee, Chou, Bomberger, Mill,
$C_{13}H_9N$		Adsorption equation: $S_s = K_p S_w$, n=1		
<u>Compound properties</u>	<u>Adsorbent</u>	<u>TOC*</u>	<u>CEC**</u>	<u>Kp</u>
m.w. 179.22	Coyote Creek sediment	1.4	13.5	1313±170
m.p. 93.5°C				
b.p. 350°C (721 torr)				
202-5°C (8 torr)				
vapor pressure at 20°C (torr)				
water sol (µg ml ⁻¹)				
76.1±2.2				
*TOC - total organic carbon				
**CEC - cation exchange capacity				

TABLE 6: Continued

Compound Name					Ref.
9H-carbazole					Smith, Mabey, Bohonos, Holt, Lee, Chou, Mill, Bomberger, 1977b
C ₆ H ₄ NHC ₆ H ₄					
<u>Compound properties</u>		<u>Adsorbent</u>	<u>K_p</u>		
m.w. 167.21		Ca-montmorillonite	3.20±1.06		
m.p. 247-248°C		Coyote creek sediment	175 ±20.9		
water sol (1.0 µg/ml)					
7H-dibenzocarbazole (DBC)					Smith, Mabey, Bohonos, Holt, Lee, Chou, Mill, Bomberger, 1977a
C ₁₀ H ₆ NHC ₁₀ H ₆					
<u>Compound properties</u>		<u>Adsorbent</u>	<u>TOC*</u>	<u>CEC**</u> <u>K_p ***</u>	
m.w. 267.31		Des Moines River sediment	0.8	10.5 32,600	
m.p. 155-159°C		Coyote Creek sediment	1.9	13.5 18,500	
water sol 2.4X10 ⁻⁷ M		Searsville Lake sediment	5.0	34.5 27,600	

*TOC = total organic carbon

**CEC = cation exchange capacity

***Based on analysis of the supernatant at equilibrium

TABLE 6: Continued

Compound Name		Ref.			
pyridine		Adsorption equation: $y = a \cdot x^b$			
C_5H_5N					
<u>Compound properties^j</u>		Spahn et al., 1974			
m.w. 79.10	<u>Adsorbent</u> activated charcoal B 10	x-range I			
m.p. 42.0°C		<u>(mmol/dm³)</u>			
b.p. 115.3°C		<u>a</u> <u>b</u>			
density .982 at 20°C		0.14 - 7.4 1.22 0.2			
water sol =					
pyridine		Adsorption equation: $X = KCe^n$, $X = \text{mg/g}$; $Ce = \text{mg/l}$			
<u>Adsorbent</u>	<u>Temp.</u>	<u>pH</u>	<u>K</u>	<u>n</u>	Baker and Luh, 1971
Na-kaolinite	1°	2	0.03	1.01	
	24°	2	0.01	1.03	
Na-montmorillonite	1°	2	0.12	1.04*	
	24°	2	0.06	1.02	

Method of analysis: liquid scintillation with Packard Tri-Carb spectrometer

Note: clay: pyridine ratio was varied from 12.15 to 6.25×10^4
 for ratio (12.15) effect of pH was studied:
 max adsorption for Na-kaolinite was found at pH 5.5; pka pyridine = 5.25
 max adsorption for Na-montmorillonite was found at pH 4.0

*discrepancy noted between data given Table 2 p.842 and regression equation p.843, believe this correct value.

TABLE 6: *Continued*

Compound Name					Ref.
quinoline					Smith, Mabey, Bohonos, Holt, Lee, Chou, Mill, Bomberger, 1976b
C_9H_7N					
Adsorption equation:					
$K_p = \frac{\mu g \text{ adsorbed}}{g} / \frac{\mu g \text{ in solution}}{ml}$					
<u>Compound properties</u>					
m.w. 129.15	<u>Adsorbent</u>	<u>TOC</u>	<u>CEC</u>	<u>K_p</u>	
water sol 6.11 μg/ml	Ca-montmorillonite	0.05	69	7.28±0.52	
pK _b = 9.5*	Coyote Creek sediment	1.4	13.5	10.9±0.4	
m.p. -14.5°C					
b.p. 161.9°C					

Experimental conditions: quinoline concentrations 4 and 8 $\mu\text{g/ml}^{-1}$;
sediment concentrations 1000 to 3000 times that of quinoline.

Method of analysis: UV spectrophotometry with Cary Model 11
spectrophotometer.

*Quinoline is calculated to be 97%, 24%, and 0.32% protonated
at pH 3, 5, and 7, respectively.

TABLE 6: Continued

TABLE 6. CONTINUED

Compound Name	Ref.
<u>S-Heterocyclics</u>	
benzo[b]thiophene	
C_8H_6S	Adsorption equation: $S_s = K_p S_w$, $n=1$
<u>Compound properties</u>	<u>Adsorbent</u> <u>TOC</u> <u>CEC</u> <u>K_p</u>
m.p. 31.3°C	Coyote Creek sediment 1.4 13.5 50±5
b.p. 212.9°C	
water sol 127.3±2.5 at 20°C	
Experimental conditions: sediment loadings of 2000:1 and 5000:1 sediment: BT by weight	
dibenzothiophene	
$C_{12}H_8S$	Adsorption equation: $S_s = K_p S_w$, $n=1$
<u>Compound properties</u>	<u>Toc</u> <u>CEC</u> <u>K_p</u>
m.w. 184.27	Coyote Creek sediment 1.4 13.5 1380±130
m.p. 99-100°C	
b.p. 332-333°C	
water sol 1.11±0.09	

Smith, Mabey,
Bohonos, Holt,
Lee, Chou,
Mill, Bomberger,
1976c

Smith, Mabey,
Bohonos, Holt,
Lee, Chou,
Bomberger, Mill,
1977

TABLE 6: Continued

Compound Name		Ref.		
<u>Phenols</u>		Adsorption equation: $y = a \cdot x^b$		
phenol	<u>Adsorbent</u>	Spahn et al., 1974		
C_6H_5OH	activated charcoal B 10	X-range I (mmol/dm ³)	a	b
<u>Compound properties</u>		0.1-8	2.16	0.23
m.w. 94.11 ^d		X-range II		
m.p. 40.9°C ^e		(mmol/dm ³)	a	b
b.p. 180°C/740mm ^k		0.001-0.06	3.6	0.39
density 1.072 g/ml ^b				
water sol 84.12 mg/ml ^k				
<u>Polynuclear aromatics</u>		Adsorption equation: $s_s = K \frac{s_p s_w}{p_w}, n=1$		
benz[a]anthracene		Smith, Mabey, Bohonos, Holt, Lee, Chou, Bomberger, Mill, 1977		
$C_{18}H_{12}$		TOC	CEC	K_p
<u>Compound properties</u>	<u>Adsorbent</u>	1.4	13.5	26,200±1700
m.w. 228.28	Coyote Creek sediment			
m.p. 155-7°C				
b.p. (at 760 torr) 435°C				
water sol at 27°C				

TABLE 6: Continued

Compound Name		Ref		
benzo[a]pyrene		Smith, Mabey, Bohonos, Holt, Lee, Chou, Mill, Bomberger, 1976a		
C ₂₀ H ₁₂		Adsorption equation: $x = K_p C$		
<u>Compound properties</u>		<u>Total organic carbon (%)</u>	<u>CEC meq. 100g⁻¹</u>	<u>K_p (x10⁻⁴)</u>
<u>Adsorbent</u>				
Ca-montmorillonite		0.06	69.0	1.7±0.5
Des Moines River		0.6	10.5	3.5±2.7
Coyote Creek		1.4	13.5	7.6±2.4
Searsville Pond		3.8	34.5	15.0±2.2
m.w. 252.32 ^a				
m.p. 178°C ^l				
b.p. ~500°C ^l				
311°C at 10 torr				
water sol 1.2±0.1 ng/ml				
pyrene		Groszek, 1975		
C ₁₆ H ₁₀		Adsorption equation: $\frac{c}{q} = \frac{1}{Kq^{\circ}} + \frac{c}{q^{\circ}}$		
<u>Compound properties</u>		q = heat evolved when the [cyclohex] is c		
m.w. 202 ^l		q° = total heat evolved when complete		
m.p. 150°C ^l		Equilibrium constants:		
b.p. 393°C ^e				
density 1.1271 ^a		<u>Est. from the Langmuir isotherm*</u>	<u>Est. from heat of ads. data</u>	<u>ΔF**</u>
water sol .135±.005 mg/l ^m				<u>Kj/mole</u>
Oct water part coeff: ^g		<u>Adsorbent</u>		<u>ΔH</u>
KOW=150,000		graphon	2200	1750
		oleophilic graphite	5930	4800
		"polar" graphite	2300	2340

Method of analysis: UV spectrophotometry with Unicam SP 500 spectrophotometer.

*at 21°

**ΔF° = -RTlnK

TABLE 6: Continued

Compound Name				Ref.
pyrene				Karickhoff, Brown, and Scott,
<u>Compound properties</u>	Adsorption equation: $X = K_p C$; $KOC = K_p / OC$			1979
water sol (mole fraction $\times 10^9$) = 12				
K octanol water = 150,000				
<u>Adsorbent</u>	<u>% O.C.</u>	<u>K_p</u>	<u>$KOC (\times 10^{-5})$</u>	
Hickory Hill				
sand	0.13	42	0.32	
coarse silt	3.27	3000	0.92	
medium silt	1.98	2500	1.3	
fine silt	1.34	1500	1.1	
clay	1.20	1400	1.2	
Doe Run				
sand	0.086	9.4	0.11	
coarse silt	2.78	2100	0.76	
medium silt	2.34	3000	1.3	
fine silt	2.89	3600	1.2	
clay	3.29	3800	1.2	
Oconee River				
sand	0.57	68	0.12	
coarse silt	2.92	3200	1.1	
medium silt	1.99	2300	1.2	
fine silt	2.26	2500	1.1	
clay	-	-	-	
Experimental conditions: sorbent concentrations 400 mg/ml of suspension for sand, 20 mg/ml for coarse and medium silt, 10 mg/ml for fine silt and 1 mg/ml for clay.				
Method of analysis: UV spectrophotometry with Perkin Elmer 356 spectrophotometer.				

TABLE 6: Continued

Compound Name		Ref.	
<u>Quinone</u>			
alizarin	adsorbent		
$C_{14}H_8O_4$	activated charcoal B 10	Adsorption equation: $y = a \cdot x^b$	Spahn et al., 1974
<u>Compound properties^a</u>		Adsorption constants:	
m.w. 240.23		x-range I	
m.p. 289-90°C (cor)		(mmol/dm ³)	a b
b.p. 430°C (sub)		0.1 - 8	1.39 0.095
water sol slightly sol			
<u>Sulfonate</u>			
sodium naphthalenesulfonate	Adsorbent	Adsorption equation: $\frac{C}{q} = \frac{1}{Kb} + \frac{C}{b}$	Gustafson et al., 1968
	Amberlite XAD-2		
	t	K	b
	l/mole	mole/g	
	25°C	319	4.58x10 ⁻⁵
		ΔH^{**}	ΔFu^{***} ΔSu^{****}
		Kcal/mole	Kcal Kcal
		-4.4	-5.7 +4.4
Method of analysis: UV spectrophotometry			
$*\Delta H^{\circ} = 2.303 \frac{RT_1T_2}{(T_2-T_1)} (\log K_2' - \log K_1')$			
$**\Delta Fu^{\circ} = RT \ln K; K=55.5K; K=\theta/(1-\theta)C; \theta=\text{fractional surface coverage}$			
$***\Delta Su^{\circ} = (\Delta H^{\circ} - \Delta Fu^{\circ})/T$			

TABLE 6: *Continued*

Compound Name

Ref.

PESTICIDESCarbamates

carbaryl

LaFleur,

N-methyl-1-naphthylcarbamateDesorption Equation: $K_b = P_s/P_w$

1976a

 $C_{12}H_{11}NO_2$ Adsorption Equation: $K_f = P_s/P_w$ Compound properties^pAdsorbent:

pH %o.m. 1/0.5 1/1 1/2 1/4 1/1

m.w. 201

Lakeland s

5.3 0.22 0.44 0.21 0.16 0.13

water sol 350 μ mol/l

Norfolk ls

6.0 0.57 1.4 0.36 0.31 0.25

Norfolk scl

5.4 0.15 0.72 0.12 0.10 0.08

Cecil sl

6.3 1.77 5.2 1.8 1.6 1.3 0.79

Cecil c

5.7 0.53 0.88 0.27 0.22 0.19

Okenee sl

4.65 5.16 7.6 3.7 2.6 2.3

Method of analysis: UV spectrophotometry.

Chlorinated hydrocarbonsAdsorption equation: $x/m = KC^{1/n}$ Huang and
Liao,

DDT

1970

1,1'-(2,2,2-trichloroethylidene)-
bis[4-chlorobenzene]AdsorbentK1/n

illite

 2.72×10^{-3} 3.28

kaolinite

 7.37×10^{-6} 5.08

montmorillonite

 1.10×10^{-5} 5.97Compound properties

m.w. 354.49

Method of analysis: GC - electron capture

TABLE 6: Continued

Compound Name

Ref.

dieldrin

1,2,3,4,10,10-hexachloro-6,7-epoxy-1,4,4a,5,6,7,8,8a-octahydro-*endo*-1,4-*exo*-5,8-dimethanonaphthalene

C₁₂H₈Cl₆O

Compound properties

m.w. 380.91

Adsorbent

illite

kaolinite

montmorillonite

K

9.45X10⁻¹⁶

1.46X10⁻²¹

1.05X10⁻¹⁶

1/n

8.82

11.63

9.24

Adsorption equation: $\frac{x}{m} = KC^{1/n}$

Method of analysis: GC-electron capture.

Huang and Liao, 1970

heptachlor

1,4,5,6,7,8,8-heptachloro-3a,4,7,7a-tetrahydro-4,7-methano-1*H*-indene

C₁₀H₅Cl₇

Adsorbent

illite

kaolinite

montmorillonite

K

1.09X10⁻⁹

5.00X10⁻⁶

1.48X10⁻⁴

1/n

6.07

4.51

3.52

Adsorption equation: $\frac{x}{m} = KC^{1/n}$

Method of analysis: GC-electron capture.

Huang and Liao, 1970

TABLE 6: Continued

Compound Name					Ref.
methoxychlor	Adsorption equation: $X = K_p C$; $KOC = K_p / OC$				Karickhoff, Brown, and Scott,
$C_{16}H_{15}Cl_3O_2$					1979
<u>Compound properties^c</u>	<u>Adsorbent</u>	<u>% O.C.</u>	<u>K_p</u>	<u>KOC (X10⁻⁵)</u>	
m.w. 345.65	Hickory Hill				
m.p. 78-78.2°C or 86-88°C	sand	0.13	53	0.41	
	coarse silt	3.27	2600	0.80	
	medium silt	1.98	1800	0.91	
insol. in water	fine silt	1.34	1400	1.0	
	clay	1.20	1100	0.92	
	Doe Run				
	sand	0.086	8.3	0.097	
	coarse silt	2.78	2200	0.80	
	medium silt	2.34	1700	0.73	
	fine silt	2.89	2300	0.80	
	clay	3.29	2400	0.73	
	Oconee River				
	sand	0.57	95	0.17	
	coarse silt	2.92	2500	0.86	
	medium silt	1.99	2000	1.0	
	fine sand	2.26	2100	0.93	
	clay	--	--	--	

Experimental conditions: sorbent concentrations 400 mg/ml of suspension for sand, 20 mg/ml for coarse and medium silt, 10 mg/ml for fine silt and 1 mg/ml for clay.

Method of analysis: UV spectrophotometry with Perkin Elmer 356 spectrophotometer.

TABLE 6: *Continued*

Compound Name					Ref
mirex					
C ₁₀ Cl ₁₂	Adsorption equation: $S_s = K \frac{S_w}{P}$, n=1				Smith, Mabey, Bohonos, Holt, Lee, Chou, Bomber- ger, Mill,
<u>Compound properties</u>	<u>Adsorbent</u>	<u>TOC</u>	<u>CEC</u>	<u>K_p</u>	1977
m.w. 546.0	Coyote Creek sediment	1.4	13.5	460,000 ± 110,000	
m.p. 485 (decomp)					
vapor pressure at 50° torr 6 x 10 ⁻⁶					
water sol at 22°C (pg/ml) 70±20					
<u>Organophosphates</u>					
aminoparathion					
C ₁₀ H ₁₆ NO ₃ PS	Adsorption equation: $x/m = KC_e^{1/n}$				Bowman and Sans,
O,O-diethyl O-p-aminophenyl phosphorothioate	<u>Adsorbents</u>	<u>1/n</u>	<u>K</u>		1977
<u>Compound properties</u>	Na-montmorillonite*	0.954	46.3		
m.w. 261.3	Ca-montmorillonite*	1.151	43.3		
water sol 390.0 µg/ml	Fe-montmorillonite*	>99.9%	adsorption		
Experimental conditions: 450 mg clay in 30 ml aqueous insecticide solution					
Method of analysis: GC - alkali flame ionization detector					

*Prepared from Wyoming bentonite, < 2 µm fraction

TABLE 6: Continued

Compound Name				Ref.
fenitrothion				Bowman and Sans, 1977
0,0-dimethyl 0-(3-methyl-4-nitrophenyl) phosphorothioate				
C ₉ H ₁₂ NO ₅ PS				
<u>Compound properties</u>	<u>Adsorbent</u>	<u>1/n</u>	<u>K</u>	
m.w. 277.2	Na-montmorillonite*	1.163	71.1	
b.p. 118°C	Ca-montmorillonite*	1.952	64.0	
density 1.3227 ₄ ⁵	Fe-montmorillonite*	1.773	740.1	
water sol 25.4 µg/ml				
Experimental conditions: same as for aminoparathion.				
Method of analysis: GC-alkali flame ionization detector.				
methyl parathion				Bowman and Sans, 1977
0,0-dimethyl 0-p-nitrophenyl phosphorothioate				
C ₈ H ₁₀ NO ₅ PS				
<u>Compound properties</u>	<u>Adsorbent</u>	<u>1/n</u>	<u>K</u>	
m.w. 263.2	Na-Montmorillonite*	1.032	65.4	
m.p. 37-38°C ^c	Ca-Montmorillonite*	1.663	56.3	
density 1.358 ₄ ^{0c}	Fe-Montmorillonite*	1.463	147.3	
water sol 45.0 µg/ml				
50 ppm ^e				
Experimental conditions: same as for aminoparathion.				
Method of analysis: GC-alkali flame ionization detector.				

*Prepared from Wyoming bentonite <2 µm fraction.

TABLE 6: *Continued*

Compound Name				Ref.
methyl parathion <i>O,O</i> -dimethyl <i>O</i> - <i>p</i> - nitrophenyl phosphorothioate $C_8H_{10}NO_5PS$ <u>Compound properties</u> m.w. 263.12	Adsorption equation: $K_p \left[\frac{\mu g \text{ substate in solution}}{ml \text{ solution}} \right]^{1/n}$ $n \approx 1.0$			Smith, Mabey, Bohonos, Holt, Lee, Mill, Bomberger, 1976
	<u>Sediment</u>	<u>K_p</u>	<u>95% confidence limit</u>	
	Ca-montmorillonite	46.1	±2.2	
	Coyote Creek	50.6	±4.5	
		55.0	±6.8	
	Searsville Pond	60.3	±3.5	
	Navarro River	47.6	±3.2	
	Des Moines River	41.6	±2.0	
	Oconee River	23.6	±1.3	

Experimental conditions: methyl parathion concentrations at 4 and 8 µg/ml. Sediment concentrations were 1000 to 2000 times as concentrated as methyl parathion at 8 µg/ml and 1000 to 3000 times as concentrated as MP at 4 µg/ml.

TABLE 6: Continued

Compound Name

Ref

parathion

Adsorption equation: $x/m = KC^{1/n}$ or
 $\log x/m = \log K + 1/n \log C$

Wahid and Sethunathan, 1978

Adsorbent Soil No.*	pH	Soil organic matter, %	CEC	Natural K	Soils 1/n	Oxidized K	Soils 1/n
10	6.20	0.75	18.6	7.67	1.04		
8	6.25	1.62	26.6	12.30	1.05	3.16	1.33
11	6.30	2.88	42.8	38.02	1.11	10.72	1.33
13	5.20	5.52	19.2	125.90	1.05		
15	3.50	8.21	21.2	213.80	1.03		
14	3.30	24.62	28.9	457.10	1.02		

Experimental conditions: 1 g soil in 10 ml aqueous parathion solution

Method of analysis: liquid scintillation

amitrole

3-amino-1,2,4-triazole

$C_2H_4N_4$

pKa = 4.14 (3-ATH⁺=H⁺ +3-AT)

Adsorption equation: $X = X_m bC/(1+bC)$

Nearpass, 1969

	Adsorbent	X_m	b^{**}
Compound properties	H sat. organic matter	63 mM/100g	0.015
water sol 28 g/100 ml (23°C)	Al sat. organic matter	49 mM/100g	0.010

Experimental conditions: 20:1 water to organic matter

Method of analysis: liquid scintillation

*all soils from India

**calculated from figure 2

TABLE 6: *Continued*

Compound Name								Ref.
<u>s-Triazine</u>								
cyanazine								Majka and Lavy, 1977
2-[[4-chloro-6-(ethylamino)- s-triazin-2-yl]amino]-2- methylpropionitrile								
Adsorption equation: $\frac{x}{m} = KC^{1/n}$								
Chromatography thin-layer (soil)								
$C_9H_{13}ClN_6$	<u>Adsorbents</u>	<u>pH</u>	<u>% o.m.</u>	<u>CEC</u>	<u>K</u>	<u>1/n</u>	<u>Rf</u>	
	Monona silty clay loam (Typic Hapludoll)	6.5	2.9	21.2	4.6	0.96	0.39	
	Valentine loamy fine sand (Typic Ustipsamment)	6.6	1.4	10.1	3.4	0.86	0.74	
Experimental conditions: 1 g soil with cyanazine added in 0.2 ml methanol: 10 ml water.								
Method of analysis: liquid scintillation with Packard 3320 spectrometer.								

TABLE 6: Continued

Compound Name										Ref
prometryne										LaFleur,
$C_{10}H_{19}N_5S$										1976b
2,4-bis(isopropylamino)- 6-methylthio-1,3,5- triazine										
	Desorption $K_D = \frac{P_s}{P_w}$									
									ΔG_{sp}^{**} (Kcal/mol)	$K_O(\text{calc})^{***}$
<u>Compound properties</u>	<u>Adsorbent</u>	<u>pH</u>	<u>%O.M</u>	<u>CCC*</u>	<u>1/0.5</u>	<u>1/1</u>	<u>1/2</u>	<u>1/4</u>		
	Lakeland s	5.3	0.22	3.0	1.5	0.86	0.74	0.67	-4.1	1060
m.w. 241	Norfolk ls	6.0	0.57	6.5	2.1	1.5	1.2	1.0	-4.1	990
water sol 200 $\mu\text{M/l}$ at 20°C	Norfolk scl	5.4	0.15	28	5.7	3.4	2.6	2.3	-3.6	420
	Cecil sl	6.3	1.77	15	7.9	4.0	3.2	2.8	-4.1	1110
pKa = 4.05	Cecil c	5.7	0.53	18	3.7	2.3	1.7	1.6	-3.3	280
	Okenee sl	4.65	5.16	19	19	12	9.0	7.0	-4.6	2340
Method of analysis: UV spectrophotometry										

*Cation combining capacity - methylene blue adsorption at pH5.

** $\Delta G_{sp} = -RT \ln K_O$

*** $K_O = P_a/P_w P_a$ is P_s adjusted to soil weight

TABLE 6: *Continued*

Compound Name

Ref.

Urea

diuron

3-(3,4-dichlorophenyl)-1,1-dimethylurea

C₉H₁₀Cl₂N₂O

Compound properties^c

m.w. 233.10

m.p. 158-159°C

water sol 42 ppm at 25°C

Adsorbents

Monona silty clay loam (Typic Hapludoll)

Valentine loamy fine sand (Typic Ustipsamment)

pH

6.5

6.6

% o.m.

2.9

1.4

CEC

21.2

10.1

K

14.3

6.5

1/n

0.77

0.74

R_f

0.18

0.39

Adsorption equation: $\frac{x}{m} = KC^{1/n}$

Chromatography thin-layer (soil)

Majka and Lavy,

1977

Experimental conditions: same as for cyanazine.

Method of analysis: liquid scintillation with Packard 3320 spectrometer.

fluometuron

1,1-dimethyl-3-(α,α,α -trifluoro-*m*-tolyl)urea

C₁₀H₁₁F₃N₂O

Compound properties^c

m.w. 232.21

m.p. 163-164°C

water sol 80 ppm at 25°C

Adsorbent

Bethony silt loam

Hector loam

Eufaula sand

% o.m.

4.4

4.8

0.3

CEC

12.4

9.2

0.6

pH

6.3

6.3

6.7

R_f

0.57

0.65

0.98

Chromatography thin-layer (soil)

Chang and Stritzke,

1977

Experimental conditions: 10 g soil: 10 ml of 0.01 N CaCl₂ solution containing 1,2,4, or 8 ppm fluometuron.

Method of analysis: liquid scintillation.

TABLE 6: *Continued*

Compound Name

Ref.

tebuthiuron

Chang and
Stritzke,*N*-[5-(1,1-dimethyl)ethyl]-1,3,4-
thiadiazol-2-yl-*N,N'*-dimethylurea

1977

 $C_9H_{16}N_4OS$

Chromatography thin-layer (soil)

Water solvent

Compound propertieswater sol 2500 μ g/mlAdsorbent% o.m.CECpHR_f

Bethony silt loam

4.4

12.4

6.3

0.58

Hector loam

4.8

9.2

6.3

0.66

Eufaula sand

0.3

0.6

6.7

0.98

Experimental conditions: 10 gm of soil: 10 ml of 0.01 N
CaCl₂ solutions containing 1,2,4, or 8 ppm tebuthiuron.

Method of analysis: liquid scintillation.

TABLE 6: *Continued*

Compound Name				Ref.
paraoxon				Bowman and Sans, 1977
Adsorption equation: $\frac{x}{m} = KC_e^{1/n}$				
O,O -diethyl O - p -nitrophenyl phosphate				
$C_{10}H_{14}NO_6P$				
<u>Compound properties</u>	<u>Adsorbent</u>	<u>1/n</u>	<u>K</u>	
m.w. 275.2	Na-Montmorillonite*	1.157	53.7	
water sol 3,250.0 μ g/ml	Ca-Montmorillonite*	1.449	42.9	
	Fe-Montmorillonite*	1.376	81.7	
Experimental conditions: same as for aminoparathion.				
Method of analysis: GC-alkali flame ionization detector.				
parathion				Bowman and Sans, 1977
Adsorption equation: $\frac{x}{m} = KC_e^{1/n}$				
O,O -diethyl O - p -nitrophenyl phosphorothioate				
$C_{10}H_{14}NO_5PS$				
<u>Compound properties</u>	<u>Adsorbent</u>	<u>1/n</u>	<u>K</u>	
m.w. 291.3	Na-Montmorillonite*	1.042	119.8	
m.p. 6.1 ^a	Ca-Montmorillonite*	1.513	120.8	
b.p. 375°C ^a	Fe-Montmorillonite*	1.194	605.8	
Experimental conditions: same as for aminoparathion.				
Method of analysis: GC-alkali flame ionization detector.				
density 1.2704 ₂₀ ^a				
water sol 12.9 μ g/ml				

*Prepared from Wyoming bentonite, <2 μ m fraction.

NOTES, TABLE 6

- ^aWeast, R. C., S. M. Selby, J. W. Long, and I. Sunshine, eds. 1974. *CRC Handbook of Chemistry and Physics*. 54th ed. Cleveland, Ohio: Chemical Rubber Company Press.
- ^bHodgman, C. D., R. C. Weast, R. S. Shankland, and S. M. Selby. 1961. *Handbook of Chemistry and Physics*. 43d ed. Cleveland, Ohio: Chemical Rubber Publishing Co.
- ^cWindholz, M., S. Budavari, L. Y. Stroumstos, and M. N. Fertig. 1976. *The Merck Index*. 9th ed. Rahway, N. J.: Merck & Co.
- ^dProtivova, J., and J. Pospisil. 1974. Antioxidants and stabilizers. XLVII: Behavior of amine antioxidants and antiozonants and model compounds in gel permeation chromatography. *J. Chromatogr.* 88:99-107.
- ^eShults, W. D., ed. 1976. *Chemical and Biological Examination of Coal-derived Materials*. Oak Ridge, Tenn.: Oak Ridge National Laboratory.
- ^fChiou, C. T., and V. H. Freed. 1977. Partition coefficient and bioaccumulation of selected organic chemicals. *Environ. Sci. Technol.* 11:475-78.
- ^gKarickhoff, S. W., D. S. Brown, and T. A. Scott. 1979. Sorption of hydrophobic pollutants on natural sediments. *Water Res.* (in press).
- ^hDorigan, J., B. Fuller, and R. Duffy. 1976. Chemistry, production and toxicity of chemicals A-C. Appendix I in *Preliminary Scoring of Selected Organic Air Pollutants*. Report No. EPA-450/3-77-008b. Research Triangle Park, N. C.: Strategies and Air Standards Division, Office of Air Quality Planning and Standards, U.S. Environmental Protection Agency.
- ⁱFendler, J. H., E. J. Fendler, G. A. Infante, P. Shih, and L. K. Patterson. 1974. Adsorption and proton magnetic resonance spectroscopic investigation of the environment of acetophenone and benzophenone in aqueous micellar solutions. *J. Am. Chem. Soc.* 97:89-95.
- ^jDorigan, J., B. Fuller, and R. Duffy. 1976. Chemistry, production and toxicity of chemicals O-Z. Appendix IV in *Preliminary Scoring of Selected Organic Air Pollutants*. Report No. EPA-450/3-77-008b. Research Triangle Park, N. C.: Strategies and Air Standards Division, Office of Air Quality Planning and Standards, U.S. Environmental Protection Agency.
- ^kHansen, R. S., Y. Fu, and F. E. Bartell. 1949. Multimolecular adsorption from binary liquid solutions. *J. Phys. Colloid Chem.* 53:769-85.

- ^lHangerbrauck, R. P., D. J. Von Lehmden, and J. E. Meeker. 1964. Emissions of polynuclear hydrocarbons and other pollutants from heat-generation and incineration processes. *J. Air Pollut. Control Assoc.* 14:267-78.
- ^mMackay, D., and W. Y. Shiu. 1977. Aqueous solubility of polynuclear aromatic hydrocarbons. *J. Chem. Eng. Data* 22:399-402.

TABLE 7: AVERAGE VALUES OF ADSORPTION CONSTANTS FOR ORGANIC PESTICIDES[†]

Compound Name

S-triazines

	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils (No.)	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils (No.)
simazine	0.820±3.48		6	1.93±6.69	135±6.33	174
6-chloro- <i>N,N'</i> -diethyl-1,3,5- triazine-2, 4-diamine				<u>16.3±30</u>	<u>159±5.66</u>	<u>2</u>
C ₇ H ₁₂ ClN ₅ m.w. 201.66 ^a						
propazine	0.95±1.7	2.68±33.0	4	2.37±7.06	160±11.6	54
2-chloro-4,6-bis (isopropylamino)- <i>s</i> -triazine				<u>26.7±9.71</u>	<u>152±3.7</u>	17
C ₉ H ₁₆ ClN ₅ m.w. 229.71 ^a						

[†]Hamaker, J. W., and J. M. Thompson. 1972. Adsorption.
In *Organic Chemicals in the Soil Environment*, Vol. 1,
ed. C. A. I. Goring and J. W. Hamaker, pp. 51-143.
New York: Marcel Dekker, Inc.

$$*K = \frac{C_s}{C_w^{1/n}}$$

$$**K_d = \frac{C_s}{C_w}$$

$$***K_{oc} = \frac{X/m(\mu g/g \text{ of organic carbon})}{C_{eq}} = \frac{K_d(\mu g/g \text{ of soil})}{\% \text{ organic carbon}}$$

Value given is average of K and K_d adjusted for organic carbon

TABLE 7: Continued

Compound Name

	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils (No.)	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils (No.)
atrazine	0.772±8.65		19	2.94±10.8	172±6.42	79
6-chloro- <i>N</i> -ethyl- <i>N'</i> -(1-methylethyl)-1,3,5-triazine-2,4-diamine	0.709±10.7		6	<u>25.5±20.4</u>	<u>102±5.80</u>	<u>24</u>
C ₈ H ₁₄ ClN ₅						
m.w. 215.69 ^a						
prometone	0.81±2.1	3.65±26	4	3.31±16.5	300±19.5	49
2,4-bis(isopropylamino)-6-methoxy- <i>s</i> -triazine				<u>13.1±11.5</u>	<u>73.8±5.1</u>	<u>15</u>
C ₁₀ H ₁₉ N ₅ O						
m.w. 225.30 ^a						
ametryne				6.17±11.1	380±10.0	33
<i>N</i> -ethyl- <i>N'</i> -(1-methylethyl)-6-(methylthio)-1,3,5-triazine-2,4-diamine				<u>167.4</u>	<u>802</u>	<u>1</u>
C ₉ H ₁₇ N ₅ S						
m.w. 227.33 ^a						
prometryne	0.855±1.1	7.83±31	4	7.03±8.78	513±13.0	53
2,4-bis(isopropylamino)-6-(methylthio)- <i>s</i> -triazine	<u>1.76</u>	<u>50.3</u>	<u>1</u>	<u>55.8±8.46</u>	<u>311±5.12</u>	<u>18</u>
C ₁₀ H ₁₉ N ₅ S						
m.w. 241.36 ^a						

TABLE 7: Continued

Compound Name	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils <u>(No.)</u>
ipazine				47.7	2571	1
6-chloro- <i>N,N</i> -diethyl- <i>N'</i> -(1-methylethyl)-1,3,5- triazine-2,4-diamine						
C ₁₀ H ₁₈ ClN ₅						
m.w. 243.74 ^a						
hydroxyatrazine				51.5	888	1
G30026	<u>0.702</u>	<u>37.4</u>			<u>81.8</u>	
Norazine						
C ₇ H ₁₂ ClN ₅						
m.w. 201.66 ^a						
<u>Ureas and Uracils</u>						
Urea	<u>0.717</u>	<u>5.22</u>			<u>14.3</u>	<u>1</u>
CH ₄ N ₂ O						
<u>Compound properties</u> ^b						
density 1.32 ₄ ¹⁸						
m.w. 60.06						
m.p. 132.7°C						

TABLE 7: Continued

Compound Name	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils <u>(No.)</u>
fenuron	0.953±0.95	0.554±20.6	3	0.781±17.0	0.554±20.6	13
<i>N,N</i> -dimethyl- <i>N'</i> -phenylurea; 1,1-dimethyl-3-phenylurea	<u>1.01±3.20</u>	<u>6.22±35.3</u>	<u>3</u>	<u>1.01±3.2</u>	<u>0.622±35.3</u>	<u>1</u>
$C_9H_{12}N_2O$						
m.w. 164.21 ^a						
methylurea	0.604±2.38	1.73±26.3	3		70.2±7.59	
	<u>0.832±4.71</u>	<u>12.5±48.3</u>	<u>3</u>		<u>57.6±8.43</u>	
phenylurea	0.744±1.34	1.80±14.8	3		76.7±8.83	
$C_7H_8N_2O$	<u>0.733±2.90</u>	<u>21.7±52.9</u>	<u>3</u>		<u>98.3±12.9</u>	
<u>Compound properties^b</u>						
m.w. 136.15						
density 1.302						
m.p. 147°C						
b.p. 238°C						
bromacil	0.58, 0.85	0.08, 1.5	2	19, 123		
5-bromo-6-methyl-3-(1-methylpropyl)- uracil						
$C_9H_{13}BrN_2O_2$						
<u>Compound properties^b</u>						
m.w. 261.11						
m.p. 157.5-160°C						

TABLE 7: Continued

Compound Name

	Average* <u>l/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils <u>(No.)</u>
terbacil 5-chloro-3-(1,1-dimethylethyl)- 6-methyl-2,4(1 <i>H</i> ,3 <i>H</i>)- pyrimidinedione C ₉ H ₁₃ ClN ₂ O ₂ m.w. 216.67 ^a	0.50, 0.96	0.15, 1.7	2		37, 140	
monuron N'-(4-chlorophenyl)-N,N- dimethylurea C ₉ H ₁₁ ClN ₂ O <u>Compound properties</u> ^b m.w. 198.65 m.p. 170.5-171.5°C water sol 230 ppm at 25°C	0.74 0.83	5.98 23.6	1 1	2.17±18.9 33.3±22.3	83.1±22.9 163±16.7	31 6
diuron 3-(3,4-dichlorophenyl)- 1,1-dimethylurea C ₉ H ₁₀ Cl ₂ N ₂ O <u>Compound properties</u> ^b m.w. 233.10 water sol 42 ppm at 25°C	0.818±6.45 <u>0.707±5.10</u>		10 <u>3</u>	6.29±9.33 <u>196±39.2</u>	351±8.95 <u>902±31.9</u>	79 <u>4</u>

TABLE 7: *Continued*

Compound Name	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils <u>(No.)</u>
metobromuron					60±20	4
3-(<i>p</i> -bromophenyl)-1-methoxy-1-methylurea						
C ₉ H ₁₁ BrN ₂ O ₂						
m.w. 259.11 ^a						
chlorobromuron					217±20	4
monolinuron	0.810±0.302	9.24±9.75	3	9.24±9.75	195±38.9	7
3-(4-chlorophenyl)-1-methoxy-1-methylurea	<u>0.832±2.33</u>	<u>43.0±48.5</u>	<u>3</u>	<u>43.0±48.5</u>	<u>198±8.67</u>	<u>3</u>
C ₉ H ₁₁ ClN ₂ O ₂						
m.w. 214.65 ^a						
neburon	0.956±2.51	27.9±53.4			787±6.89	3
1-butyl-3-(3,4-dichlorophenyl)-1-methylurea						
C ₁₂ H ₁₆ Cl ₂ N ₂ O						
m.w. 275.18						
linuron	0.75, 0.75	45.2, 67.8	2	18.1±18.7	840±20	26
<i>N</i> '-(3,4-dichlorophenyl)- <i>N</i> -methoxy- <i>N</i> -methylurea				<u>147±77.2</u>	<u>653±13.4</u>	<u>3</u>
C ₉ H ₁₀ Cl ₂ N ₂ O ₂						
<u>Compound properties^b</u>						
m.w. 249.10						
m.p. 93-94°C						
water sol 75 ppm						

TABLE 7: *Continued*

Compound Name

	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils (No.)	Average** <u>K_d±S%</u>	Average*** <u>K_{OC}±S%</u>	Soils (No.)
<u>Halogenated Hydrocarbons</u>						
ethylene dibromide	0.967, 0.970	0.408, 0.803	2	0.537±24.8	32.4±23.7	3
1,2-dibromoethane	<u>1.04</u>	<u>2.16</u>	<u>1</u>	<u>2.13±1.41</u>	<u>14.4±19.4</u>	<u>2</u>
C ₂ H ₄ Br ₂						
m.w. 187.87 ^a						
beta-BHC	0.861	148	1		4254	
1,2,3,4,5,6-hexachlorocyclo- hexane						
C ₆ H ₆ Cl ₆	<u>0.950</u>	<u>456</u>	<u>1</u>		<u>3573</u>	
m.w. 290.85 ^a						
lindane	0.841,080	45.7, 7.91	2	24.7±27.8	1342±15.3	6
1 α ,2 α ,3 β ,4 α ,5 α ,6 β - hexachlorocyclohexane	<u>0.981</u>	<u>331</u>	<u>1</u>	<u>321±6.10</u>	<u>1943±20.4</u>	<u>4</u>
C ₆ H ₆ Cl ₆						
<u>Compound properties^b</u>						
m.w. 290.85						
m.p. 112.5°C						
insol. in water						

TABLE 7: Continued

Compound Name	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{OC}±S%</u>	Soils <u>(No.)</u>
N-serve				6.80±30.1	271±11.5	9
2-chloro-6-(trichloromethyl)- pyridine				<u>44.5</u>	<u>238</u>	<u>1</u>
$C_6H_3Cl_4N$						
<u>Compound properties^b</u>						
m.w. 230.93						
m.p. 62.5-62.9°C						
b.p. 136-137.5°C						
DDT				1.3X10 ⁵ , 1.37X10 ⁴	1.31X10 ⁵ 3.55X10 ⁵	2
				<u>1.063X10⁵</u>	<u>2.29X10⁵</u>	<u>1</u>
<u>Carbamates</u>						
chloropropham	1.00	21	1	9.28±11.8	590±8.35	65
isopropyl-m-chlorocarbanilate	<u>1.00</u>	<u>132</u>	<u>1</u>			
$C_{10}H_{12}ClNO_2$						
m.w. 213.67 ^a						
propham					51	4
isopropyl carbanilate						
$C_{10}H_{13}NO_2$						
m.w. 179.22 ^a						

TABLE 7: *Continued*

Compound Name

	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils <u>(No.)</u>
EPTC				5.96±24.6	283±18.9	3
<u>S</u> -ethyl dipropylcarbamothioate				<u>52.6</u>	<u>109</u>	<u>1</u>
C ₉ H ₁₉ NOS						
m.w. 189.32 ^a						
cycloate				7.32±25	345±17.4	3
<u>S</u> -ethyl cyclohexylethyl- carbamothioate				<u>66.6</u>	<u>222</u>	<u>1</u>
C ₁₁ H ₂₁ NOS						
m.w. 215.36 ^a						
pebulate				14.5±20.1	719±21.4	3
<u>S</u> -propyl butylethylthio- carbamate				<u>109</u>	<u>363</u>	<u>1</u>
C ₁₀ H ₂₁ NOS						
m.w. 203.35 ^a						

TABLE 7: *Continued*

Compound Name

	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils <u>(No.)</u>
<u>Phosphates</u>						
nellite				0.525±72.5	30.89±25.9	8
$C_8H_{13}N_2O_2P$				<u>6.30, 6.02</u>	<u>43.5, 33.2</u>	<u>2</u>
m.w. 200.18 ^a						
crotoxyphos				6.09±15	173±34	3
(E)-1-phenylethyl- 3-[(dimethoxyphosphinyl)oxy]- 2-butenolate						
$C_{14}H_{19}O_6P$						
m.w. 314.28 ^a						
phorate	0.97±2.1	8.76±39	4		3199±25	
O,O-diethyl S-[(ethylthio) methyl] phosphorodithioate						
$C_7H_{17}O_2PS_3$						
m.w. 260.38 ^a						

TABLE 7: *Continued*

Compound Name

	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils <u>(No.)</u>
disulfoton	0.930±2.38	20.1±18.9	10		2132±25	16
<i>O,O</i> -diethyl <i>S</i> -[2-(ethylthio)ethyl]phosphorodithioate $C_8H_{19}O_2PS_3$ <u>Compound properties^b</u> m.w. 274.41 b.p. 108°C density 1.144 ₄ ⁰ insol. in water						
methyl parathion	1.04	18.43±20	4		9799±41	
<i>O,O</i> -dimethyl <i>O</i> -(4-nitrophenyl)phosphorothioate $C_8H_{10}NO_5PS$ <u>Compound properties^b</u> m.w. 263.23 m.p. 37-38°C density 1.358 ₄ ⁰ water sol 50 ppm						
parathion	1.03±1.5	2.19±32	4		10454±38	
<i>O,O</i> -diethyl <i>O</i> -(4-nitrophenyl)phosphorothioate $C_{10}H_{14}NO_5PS$ m.w. 291.26						

TABLE 7: Continued

Compound Name	Average*	Average*	Soils	Average**	Average***	Soils
	<u>l/n±S%</u>	<u>K±S%</u>	<u>(No.)</u>	<u>K_d±S%</u>	<u>K_{oc}±S%</u>	<u>(No.)</u>
ethion	0.945±2.5	28.5±15.9	4		15435±39.3	
$0,0,0',0'$ -tetraethyl S,S' - methylenebisphosphorodithioate $C_9H_{22}O_4P_2S_4$ <u>Compound properties^b</u> m.w. 384.48 m.p. -12 to -13°C density 1.220 ₄ ⁰ slightly sol in water						
carbophenothion	0.940±3.7	74.5±13	4		45368±40.6	
$S[[(4\text{-chlorophenyl})\text{thio}]\text{methyl}]$ $0,0$ -diethyl phosphorodithioate $C_{11}H_{16}ClO_2PS_3$ <u>Compound properties^b</u> m.w. 342.85 b.p. 82°C density 1.271 ₄ ⁵ insol. in water						

TABLE 7: Continued

Compound Name


	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils <u>(No.)</u>
<u>Amino, Nitrophenyl Sulfones</u>						
						
nitralin				50, 26	500, 520	2
(R ₁ = CH ₃ H ₇ ; R ₂ = CH ₃)						
4-(methylsulfonyl)-2,6-dinitro- N,N-dipropylbenzenamine						
C ₁₃ H ₁₉ N ₃ O ₆ S						
m.w. 345.38						
SD11830				12.5, 7.25	125, 145	2
(R ₁ = CH ₃ ; R ₂ = CH ₃)						
SD12639				23, 9.65	230, 193	2
(R ₁ = C ₂ H ₅ ; R ₂ = CH ₃)						
SD13207				32, 13.5	320, 269	2
(R ₁ = C ₂ H ₅ ; R ₂ = C ₂ H ₅)						
SD12030				75, 35.1	750, 702	2
(R ₁ = C ₃ H ₇ ; R ₂ = C ₂ H ₅)						
SD12346				117	1170	2
(R ₁ = C ₃ H ₇ ; R ₂ = C ₃ H ₇)						
SD12400				22.2	222	2
(R ₁ = CH ₃ ; R ₂ = C ₃ H ₇)						

TABLE 7: Continued

Compound Name

	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils <u>(No.)</u>
<u>Carboxylic Acids</u>						
dicamba				<u>0.9</u>	<u>3.8</u>	<u>1</u>
3,6-dichloro-2-methoxybenzoic acid						
$C_8H_6Cl_2O_3$						
m.w. 221.04 ^a						
chloramben NH_4^+ salt				0.320±30.4	15.4±37	9
chloramben: 3-amino-2,5-dichlorobenzoic acid				<u>4.1</u>	<u>17.4</u>	<u>1</u>
$C_7H_5Cl_2NO_2$						
m.w. 206.03 ^a						
chloramben methyl ester				3.5±35	507±24	4
MCPA				0.420±37.4		
(4-chloro-2-methylphenoxy)-acetic acid						
$C_9H_9ClO_3$						
m.w. 200.62						

TABLE 7: *Continued*

Compound Name	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils <u>(No.)</u>
picloram				0.453±32.2	12.7±21.7	14
4-amino-3,5,6-trichloro- 2-pyridinecarboxylic acid						
$C_6H_3O_2Cl_3N_2$						
m.w. 241.46 ^a						
2,4-D				1.59±54.9		7
(2,4-dichlorophenoxy) - acetic acid						
$C_8H_6Cl_2O_3$						
m.w. 221.04 ^a						
2,4,5-T				1.05±42.3		7
(2,4,5-trichlorophenoxy) - acetic acid						
$C_8H_5Cl_3O_3$						
m.w. 255.49 ^a						
dichlobenil		167		4.19±11.4	164±3.57	16

TABLE 7: Continued

Compound Name	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{OC}±S%</u>	Soils <u>(No.)</u>
<u>Acids</u>						
linear alkyl				34.2±57.4	1222±10.2	30 (K _d) 20 (K _{OC})
sulfonates				<u>55.3</u>	<u>170</u>	<u>1</u>
pentachlorophenol				8.96±20.38		10
<chem>C6HCl5O</chem>						
<u>Compound properties</u> ^b						
m.w. 266.35						
m.p. 190-191°C						
b.p. 309-310°C						
density 1.978 ₄ ²²						
water sol. 8 mg/100 ml						
methane arsonate				12.4±38.9	770±19.1	4
<chem>CH5AsO3</chem>						
<u>Compound properties</u> ^b						
m.w. 139.96						
m.p. 161°C						
freely sol. in water						

TABLE 7: *Continued*

Compound Name	Average* <u>1/n±S%</u>	Average* <u>K±S%</u>	Soils <u>(No.)</u>	Average** <u>K_d±S%</u>	Average*** <u>K_{oc}±S%</u>	Soils <u>(No.)</u>
<u>Miscellaneous</u>						
silvex	0.639, 0.987	42.1, 34.2	2		2786, 4682	
2-(2,4,5-trichlorophenoxy)- propionic acid	<u>1.05</u>	<u>162</u>	<u>1</u>		<u>440</u>	
C ₉ H ₇ Cl ₃ O ₃						
<u>Compound properties</u> ^b						
m.w. 269.53						
m.p. 181.6°C						
0.014% sol. in water at 25°C						
chlorthiamid	0.868±1.28	4.72±8.13	5		107.2±6.42	
2,6-dichlorobenzene carbothioamide						
C ₇ H ₅ Cl ₂ NS						
chloroneb	1.3	20	1		1159	
1,4-dichloro-2,5-dimethoxy- benzene						
C ₈ H ₈ Cl ₂ O ₂						
m.w. 207.06 ^a						
paraquat	0.566±19.6	353±97			20152±65	
1,1'-dimethyl-4,4'- bipyridinium	0.360±13.3	5501±16.6				
C ₁₂ H ₁₄ N ₂						
m.w. 186.26 ^a						

NOTES, TABLE 7

^a Wiswesser, W.J., ed. 1976. *Pesticide Index*. 5th ed. College Park, Maryland: The Entomological Society of America.

^b Windholz, M., S. Budavari, L.Y. Stroumtsos, and M.N. Fertig. 1976. *The Merck Index*. 9th ed. Rahway, N.J.: Merck and Co.

TABLE 8: RELATIONSHIP BETWEEN OCTANOL/WATER PARTITION COEFFICIENT AND R_f VALUES OF PESTICIDES
ON A SOIL

<u>Mobility Class</u>	<u>R_f</u>	<u>$\log P$</u> [*]	<u>Q</u> ^{**}	<u>Ref</u>
Immobile	0 - 0.09	>3.78	>398	Briggs, 1973
Low	0.10 - 0.34	3.78-2.39	398-74	
Intermediate	0.35 - 0.64	2.39-1.36	74-29	
Mobile	0.65 - 0.89	1.36-0.08	29-4.5	
Very Mobile	0.90 - 1.00	<0.08	<4.5	

* P = Octanol/water partition coefficient

** Q = Soil organic matter/water partition coefficient

TABLE 9: LEACHING OF PESTICIDES FROM A SOIL

Compound Name			Ref.
Adsorbent = Hagerstown silty clay loam			Hamaker,
$\%0.M = 2.41, \%0.C = \%0.M/1.724 = 1.40$			1974
Pesticide	R_f (soil TLC)	Mobility class	K_{oc}^*
chloramben	0.96	5	12.8
2,4-D	0.69	4	32
propham	0.51	3	51
bromacil	0.69	4	71
monuron	0.48	3	83
simazine	0.45	3	135
propazine	0.41	3	152
dichlobenil	0.22	2	164
atrazine	0.47	3	172
chlorpropham	0.18	2	245
prometone	0.60	3	300
ametryne	0.44	3	380
diuron	0.24	2	485
prometryne	0.25	2	513
chloroxuron	0.09	1	4,986
paraquat	0.00	1	20,000
DDT	0.00	1	243,000

* K_{oc} = adsorption coefficient on the basis of organic carbon

TABLE 10: SORPTION DEPENDENCE ON SORBATE PROPERTIES

Compound Name	Compound Properties			Ref.
Compound	Water solubility (mole fraction $\times 10^9$)	$K_{OC} (X10^{-3})^*$	$K_{OW} (X10^{-3})^{**}$	Karickhoff, Brown, and Scott, 1979
pyrene	12	84	150	
methoxychlor	6.3	80	120	
naphthalene	4460	1.3	2.3	
2-methylnaphthalene	3220	8.5	13	
anthracene	7.57	26	35	
9-methylanthracene	24.4	65	117	
phenanthrene	130	23	37	
tetracene	0.037	650	800	
hexachlorobiphenyl	0.048	1200	2200	
benzene	189,000	0.083	0.13	

* K_{OC} = partition coefficient based on organic carbon

** K_{OW} = octanol/water distribution coefficient

REFERENCES

- Adams, R. S., Jr. 1973. Factors influencing soil adsorption and bioactivity of pesticides. *Residue Rev.* 47:1-53.
- Bailey, G. W., and J. L. White. 1970. Factors influencing the adsorption, desorption and movement of pesticides in soil. *Res. Rev.* 32:29-92.
- Baker, R. A., and M. Luh. 1971. Pyridine sorption from aqueous solution by montmorillonite and kaolinite. *Water Res.* 5:839-48.
- Bartell, F. E., and E. J. Miller. 1924. Adsorption by activated sugar charcoal. III: The mechanics of adsorption. *J. Phys. Chem.* 28:992-1000.
- Bartell, F. E., T. L. Thomas, and Y. Fu. 1951. Thermodynamics of adsorption from solution. *J. Physical Colloid Chem.* 55:1456-62.
- Bowman, B. T., and W. W. Sans. 1977. Adsorption of parathion, fenitrothion, methyl parathion, aminoparathion and paraoxon by Na^+ , Ca^{2+} , and Fe^{3+} montmorillonite suspensions. *Soil Sci. Soc. Am. J.* 41:514-19.
- Briggs, G. G. 1973. A simple relationship between soil adsorption of organic chemicals and their octanol/water partition coefficients. In *Proceedings of the 7th British Insecticide and Fungicide Conference*, pp. 475-78.
- Chang, S. S., and J. F. Stritzke. 1977. Sorption, movement, and dissipation of tebuthiuron in soils. *Weed Sci.* 25:184-87.
- Cork J. M., J. F. Goodman, and J. R. Tate. 1966. Adsorption of non-ionic surface-active agents at the graphon/solution interface. *Trans. Far. Soc.* 62:979-86.
- Cummings, T., H. C. Garven, C. H. Giles, S. M. K. Rahman, J. G. Sneddon, and C. E. Stewart. 1959. Adsorption at inorganic surfaces. Part IV. Mechanism of adsorption of organic solutes by chromatographic alumina. *J. Chem. Soc.* pp. 535-44.
- Farmer, W. J. 1976. Leaching, diffusion, and sorption of benchmark pesticides. In *A Literature Survey of Benchmark Pesticides*, pp. 185-244. Washington, D.C.: George Washington University Medical Center.
- Groszek, A. J. 1975. Adsorption of polycyclic aromatic hydrocarbons onto graphite. *Faraday Discuss. Chem. Soc.* 59:109-16.
- Gustafson, R. L., R. L. Albright, J. Heisler, J. A. Lirio, and O. T. Reid, Jr. 1968. Adsorption of organic species by high surface area styrene-divinyl-benzene copolymers. *Ind. Eng. Chem. Prod. Res. Dev.* 7:107-15.

- Gustafson, R., and J. Paleos. 1971. Interactions responsible for the selective adsorption of organics on organic surfaces. In *Organic Compounds in Aquatic Environments*, ed. S. D. Faust and J. V. Hunter, pp. 213-37. New York: Marcel Dekker, Inc.
- Hamaker, J. W. 1974. The interpretation of soil leaching experiments. In *Environmental Dynamics of Pesticides*, ed. R. Haque and V. H. Freed. New York: Plenum Press.
- Hance, R. J. 1969. Influence of pH, exchangeable cation and the presence of organic matter on the adsorption of some herbicides by montmorillonite. *Can. J. Soil Sci.* 49:357-64.
- Hansen, R. S., and R. P. Craig. 1954. The adsorption of aliphatic alcohols and acids from aqueous solutions by non-porous carbons. *J. Phys. Chem.* 58:211-15.
- Hartman, R. J., R. A. Kern, and E. G. Bobalek. 1946. Adsorption isotherms of some substituted benzoic acids. *J. Colloid Sci.* 1:271-76.
- Horvath, C., and W. Melander. 1978. Reversed-phase chromatography and the hydrophobic effect. *Amer. Lab.*, October 1978, 17-36.
- Huang, J., and C. Liao. 1970. Adsorption of pesticides by clay minerals. *J. San. Eng. Div., Proc. Am. Soc. Civil Eng.* 96:1057-78.
- Karickhoff, S. W., D. S. Brown, and T. A. Scott. 1979. Sorption of hydrophobic pollutants on natural sediments. *Water Res.* (in press).
- Kipling, J. J. 1965. *Adsorption from Solutions of Non-electrolytes*. New York: Academic Press.
- LaFleur, K. S. 1976a. Carbaryl desorption and movement in soil columns. *Soil Sci.* 121:212-16.
- LaFleur, K. S. 1976b. Prometryne desorption and movement in soil columns. *Soil Sci.* 121:9-15.
- Lailach, G. E., T. D. Thompson, and B. W. Brindley. 1968. Adsorption of pyrimidines, purines, and nucleosides by Li-, Na-, Mg-, and Ca-montmorillonite. *Clays and Clay Minerals* 16:285-93.
- Lambert, S. M. 1967. Functional relationship between sorption in soil and chemical structure. *J. Agric. Food Chem.* 15:572-76.
- Lambert, S. M. 1968. Omega (Ω), a useful index of soil sorption equilibria. *J. Agric. Food Chem.* 16:340-43.
- Linner, E. R., and R. A. Gortner. 1935. Interfacial energy and the molecular structure of organic compounds. III: The effect of organic structure on adsorbability. *J. Phys. Chem.* 39:35-67.

- Majka, J. T., and T. L. Lavy. 1977. Adsorption, mobility, and degradation of cyanazine and diuron in soils. *Weed Sci.* 25:401-6.
- Means, J. C., J. J. Hassett, S. G. Wood, and W. L. Banwart. 1979. Sorption properties of energy-related pollutants and sediments. In *Carcinogenesis -- A Comprehensive Survey; Polynuclear Aromatic Hydrocarbons: Analysis Chemistry and Biology*, Vol. 4 (in press).
- Mortland, M. M. 1970. Clay-organic complexes and interactions. *Adv. Agron.* 22:75-117.
- Nearpass, D. C. 1969. Exchange adsorption of 3-amino-1,2,4-triazole by an organic soil. *Soil Sci. Soc. Am. Proc.* 33:524-28.
- Nogami, H., T. Nagai, and H. Uchida. 1968. Physico-chemical approach to biopharmaceutical phenomena. III: Hydrophobic bonding in the adsorption of tryptophan by carbon black from aqueous solution. *Chem. Pharm. Bull.* 16:2263-66.
- Osgerby, J. M. 1970. Sorption of un-ionized pesticides by soils. In *Sorption and Transport Process in Soils*. Monograph 37. London: Society of Chemical Industry.
- Parkash, S. 1974. Adsorption of weak and non-electrolytes by activated carbon. *Carbon* 12:37-43.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, D. C. Bomberger, and T. Mill. 1977. *Environmental Pathways of Selected Chemicals in Freshwater Systems. Part II: Laboratory Studies*. Athens, Ga.: Environmental Research Laboratory, Office of Research and Development, U. S. Environmental Protection Agency.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, T. Mill, and D. C. Bomberger. 1976a. *Laboratory Investigation of Benzo(a)pyrene*. Cincinnati, Ohio: National Environmental Research Center, Office of Research and Development, U. S. Environmental Protection Agency.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, T. Mill, and D. C. Bomberger. 1976b. *Laboratory Investigation of Benzo(b)thiophene*. Cincinnati, Ohio: National Environmental Research Center, Office of Research and Development, U. S. Environmental Protection Agency.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, T. Mill, and D. C. Bomberger. 1976c. *The Fate of Quinoline in Freshwater Aquatic Systems*. Cincinnati, Ohio: National Environmental Research Center, Office of Research and Development, U. S. Environmental Protection Agency.

- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, T. Mill, and D. C. Bomberger. 1977a. Laboratory investigation of 9H-carbazole. In *The Fate of Selected Pollutants in Freshwater Aquatic Systems. Part II: Laboratory Studies*, pp. 145-68. Athens, Ga.: Environmental Research Laboratory, Office of Research and Development, U. S. Environmental Protection Agency.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, T. Mill, and D. C. Bomberger. 1977b. Laboratory investigation of 7H-dibenzo(c,g)carbazole. In *The Fate of Selected Pollutants in Freshwater Aquatic Systems. Part II: Laboratory Studies*, pp. 169-93. Athens, Ga.: Environmental Research Laboratory, Office of Research and Development, U. S. Environmental Protection Agency.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. Mill, and D. C. Bomberger. 1976. *The Fate of Methyl Parathion in Freshwater Aquatic Systems*. Cincinnati, Ohio: National Environmental Research Center, Office of Research and Development, U. S. Environmental Protection Agency.
- Spahn, H., V. Branch, E. U. Schlunder, and H. Sontheimer. 1974. Explanation of activated charcoal filters for water purification. Part I: Investigation of the adsorption on single granules. *Verfahrenstechnik* 8:224-31.
- Wahid, P. A., and N. Sethunathan. 1978. Sorption-desorption of parathion in soils. *J. Agric. Food Chem.* 26:101-5.
- Ward, T. M., and F. W. Getzen. 1970. Influence of pH on the adsorption of aromatic acids on activated carbon. *Environ. Sci. Technol.* 4:64-7.
- Ward, T. M., and R. P. Upchurch. 1965. Role of the amido group in adsorption mechanisms. *J. Agric. Food Chem.* 13:334-40.
- Weber, J. B. 1972. Interaction of organic pesticides with particulate matter in aquatic and soil systems. In *Fate of Organic Pesticides in the Aquatic Environment*, ed. R. F. Gould, pp. 55-120. Advances in Chemistry, Series III. Washington, D.C.: American Chemical Society.
- Weber, T. B., P. W. Perry, and R. P. Upchurch. 1965. The influence of temperature and time on the adsorption of paraquat, diquat, 2,4-D and prometon by clays, charcoal, and an ion-exchange resin. *Soil Sci. Soc. Am., Proc.* 29:678-88.
- Zettlemoyer, A. C., and F. J. Micale. 1971. Solution adsorption thermodynamics for organics on surfaces. In *Organic Compounds in Aquatic Environments*, ed. S. Faust and J. Hunter, pp. 165-85. New York: Marcel Dekker.

BIBLIOGRAPHY

ADSORPTION OF ORGANIC COMPOUNDS

- Adams, R. S., Jr. 1971. Effect of soil organic matter on the movement and activity of pesticides in the environment. In *Proceedings of the University of Missouri's 5th Annual Conference on Trace Substances in Environmental Health*, pp. 81-93. Columbia, Mo.: University of Missouri.
- Adams, R. S., Jr. 1973. Factors influencing soil adsorption and bioactivity of pesticides. *Residue Rev.* 47:1-53.
- Adams, R. S., Jr., and P. Li. 1971. Soil properties influencing sorption and desorption of lindane. *Soil Sci. Soc. Am., Proc.* 35:78-81.
- Adamson, A. W. 1967. *Physical Chemistry of Surfaces*. 2d ed. New York: John Wiley & Sons.
- Allingham, M. M., J. M. Cullen, C. H. Giles, S. K. Jain, and J. J. Woods. 1954. Adsorption at inorganic surfaces. II: Adsorption of dyes and related compounds by silica. *J. Appl. Chem.* 8:108-116.
- Ardizzi, P. G., M. Tschapek, and S. G. de Bussetti. 1973. Thermodynamics of water adsorption by kaolin from benzene. *Kolloid - Z. Z. Polym.* 251:490-93.
- Bader, R. G., and J. B. Smith. 1956. Significance of adsorption isotherms for specific organic materials on sedimentary minerals. *Bull. Geol. Soc. Am.* 70:1564.
- Bailey, G. W. 1971. Adsorption of pesticides by clay minerals. *J. Sanit. Eng. Div. Proc., Am. Soc. Civil Eng.* 97:533-35.
- Bailey, G. W., and J. L. White. 1964. Review of adsorption and desorption of organic pesticides by soil colloids, with implications concerning pesticide bioactivity. *Agricultural and Food Chemistry* 12:324-32.
- Bailey, G. W., and J. L. White. 1970. Factors influencing the adsorption, desorption and movement of pesticides in soil. *Res. Rev.* 32:29-92.
- Baker, R. A., and M. Luh. 1971. Pyridine sorption from aqueous solution by montmorillonite and kaolinite. *Water Res.* 5:839-48.
- Bami, H. L. 1961. Sorption of 75% DDT water-dispersible powder on different mud surfaces. *Bull. World Health Org.* 24:567-75.

- Barlow, F., and A. B. Hadaway. 1955. Studies on aqueous suspensions of insecticides, Part V: The sorption of insecticides by soils. *Bull. Entomological Res.* 46:547-59.
- Barlow, F., and A. B. Hadaway. 1958. Studies on aqueous suspensions of insecticides, Part VI: Further notes on the sorption of insecticides by soils. *Bull. Entomological Res.* 49:315-31.
- Barlow, F., and A. B. Hadaway. 1958. Studies on aqueous suspensions of insecticides, Part VII: The influence of relative humidity upon the sorption of insecticides by soils. *Bull. Entomological Res.* 49:333-54.
- Barres, R. M., and K. E. Kelsey. 1961. Thermodynamics of interlamellar complexes, Part I: Hydrocarbons in methylammonium montmorillonites. *Trans. Faraday Soc.* 57:452-62.
- Barrer, R. M., and K. E. Kelsey. 1961. Thermodynamics of interlamellar complexes, Part 2: Sorption by dimethyldioctadecylammonium bentonite. *Trans. Faraday Soc.* 57:625-40.
- Barrer, R. M., and D. M. MacLeod. 1955. Activation of montmorillonite by ion exchange and sorption complexes of tetra-alkyl ammonium montmorillonites. *Trans. Faraday Soc.* 51:1290-1301.
- Barshad, I. 1952. Factors affecting the interlayer expansion of vermiculite and montmorillonite with organic substances. *Soil Sci. Soc. Am., Proc.* 16:176.
- Bartell, F. E., and Y. Fu. 1929. Adsorption from aqueous solutions by silica. *J. Phys. Chem.* 33:676-87.
- Bartell, F. E., and E. J. Miller. 1924. Adsorption by activated sugar charcoal. III: The mechanics of adsorption. *J. Phys. Chem.* 28:992-1000.
- Bartell, F. E., T. L. Thomas, and Y. Fu. 1951. Thermodynamics of adsorption from solution. *J. Physical Colloid Chem.* 55:1456-62.
- Bertagna, P. 1959. Residual insecticides and the problem of sorption. *Bull. World Health Org.* 20:861-89.
- Blom, B. E., T. F. Jenkins, D. C. Leggett, and R. P. Murrman. 1976. *Effect of Sediment Organic Matter on Migration of Various Chemical Constituents During Disposal of Dredged Material*. NTIS No. AD-A027-394. Vicksburg, Miss.: Environmental Effects Laboratory, U. S. Army Engineer Waterways Experiment Station.
- Blumer, M. 1961. Benzpyrenes in soil. *Science.* 134:474.
- Bowman, B. T., and W. W. Sans. 1977. Adsorption of parathion, fenitrothion, methyl parathion, aminoparathion and paraoxon by Na^+ , Ca^{2+} , and Fe^{3+} montmorillonite suspensions. *Soil Sci. Soc. Am., J.* 41:514-19.

- Bradley, W. F. 1945. Molecular associations between montmorillonite and some polyfunctional organic liquids. *J. Am. Chem. Soc.* 67:975-81.
- Briggs, G. G. 1973. A simple relationship between soil adsorption of organic chemicals and their octanol/water partition coefficients. In *Proceedings of the 7th British Insecticide and Fungicide Conference*, pp. 475-78.
- Brindley, G. W. 1965. Clay-organic studies. X: Complexes of primary amines with montmorillonite and vermiculite. *Clay Minerals* 6:91-96.
- Brindley, G. W., and G. Ertem. 1971. Preparation and solvation properties of some variable charge montmorillonites. *Clays Clay Miner.* 19:399-404.
- Brindley, G. W. and M. Rustom. 1958. Adsorption and retention of an organic material by montmorillonite in the presence of water. *Am. Mineral.* 43:627-40.
- Brindley, G. W., and T. D. Thompson. 1966. Clay-organic studies. XI: Complexes of benzene, pyridine, and piperidine 1,3,-substituted propanes with a synthetic Ca-fluorhectorite. *Clay Minerals* 6:345-50.
- Burns, I. G., and M. H. B. Hayes. 1974. Some physico-chemical principles involved in the adsorption of the organic cation paraquat by soil humic materials. *Residue Rev.* 52:117-46.
- Burns, I. G., M. H. B. Hayes, and M. Stacey. 1973. Some physico-chemical interactions of paraquat with soil organic materials and model compounds. I: Effects of temperature, time and adsorbate degradation on paraquat adsorption. *Weed Res.* 13:67-78.
- Cason, J., and G. A. Gillies. 1955. Adsorption and chromatography of fatty acids on charcoal. *J. Org. Chem.* 20:419-27.
- Cerofolini, G. F. 1975. Extension of the asymptotically-correct approximation to Fowler-Guggenheim adsorption. *Surf. Sci.* 52:195-98.
- Champion, D. F., and S. R. Olsen. 1971. Adsorption of DDT on solid particles. *Soil Sci. Soc. Am., Proc.* 35:887-91.
- Chang, S. S., and J. F. Stritzke. 1977. Sorption, movement, and dissipation of tebuthiuron in soils. *Weed Sci.* 25:184-87.
- Chen, N. Y. 1976. Hydrophobic properties of zeolites. *J. Phys. Chem.* 80:60-64.
- Clark, A. 1970. Localized-adsorption-independent systems. In *The Theory of Adsorption and Catalysis*. pp. 3-16. New York: Academic Press.
- Clark, A. 1970. Thermodynamics of adsorption. In *The Theory of Adsorption and Catalysis*. pp. 17-61. New York: Academic Press.

- Clark-Monks, C., and B. Ellis. 1972. N-butylamine adsorption onto amorphous silica. *Can. J. Chem.* 50:907-11.
- Colbert, F. D., V. V. Volk, and A. P. Appleby. 1975. Sorption of atrazine, terbutryn, and GS-14254 on natural and lime-amended soils. *Weed Sci.* 23:390-94.
- Conway, B. E., H. Angerstein-Kozłowska, and H. P. Dhar. 1974. On selection of standard states in adsorption isotherms. *Electrochim. Acta* 19:455-60.
- Corkill, J. M., J. F. Goodman, and J. R. Tate. 1966. Adsorption of non-ionic surface-active agents at the graphon/solution interface. *Trans. Faraday Soc.* 62:979-86.
- Corkill, J. M., J. F. Goodman, and J. R. Tate. 1967. Adsorption of alkyl-sulphinyllkanols on graphon. *Trans. Faraday Soc.* 63:2264-69.
- Cummings, T., H. C. Garven, C. H. Giles, S. M. K. Rahman, J. G. Sneddon, and C. E. Stewart. 1959. Adsorption at inorganic surfaces. Part IV. Mechanism of adsorption of organic solutes by chromatographic alumina. *J. Chem. Soc.* pp. 535-44.
- Davidson, J. M., C. E. Rieck, and P. W. Santelmann. 1968. Influence of water flux and porous material on the movement of selected herbicides. *Soil Sci. Soc. Am., Proc.* 32:629-33.
- Davies, R. A., H. J. Kaempf, and M. M. Clemens. 1973. Removal of organic material by adsorption on activated carbon. *Chem. Ind. (London)* 1:827-31.
- Defay, R., and I. Prigogine. 1966. Gibbs' adsorption equation. In *Surface Tension and Adsorption*, pp. 85-95. New York: John Wiley & Sons.
- Defay, R., and I. Prigogine. 1966. Model of a surface and the definition of adsorption, and the energy and entropy of surfaces. In *Surface Tension and Adsorption*, pp. 21-33, New York: John Wiley & Sons.
- Defosse, C., P. Canesson, and B. Delmon. 1976. Evidence of superficial reduction of NH_4^+ zeolite silicon upon pyridine adsorption at 150°C. *J. Phys. Chem.* 80:1028-30.
- Dodd, G. C., and S. Ray. 1959. Semiquinone cation adsorption on montmorillonite as a function of surface acidity. *Clays Clay Miner.* 8:237-51.
- Doebler, R. W., and W. A. Young. 1960. Some conditions affecting the adsorption of quinoline by clay minerals in aqueous suspensions. *Clays Clay Miner.* 9:468-83.
- Doherty, P. J., and G. F. Warren. 1969. The adsorption of four herbicides by different types of organic matter and a bentonite clay. *Weed Res.* 9:20-26.

- Dollimore, D. G., R. Heal, and D. R. Martin. 1973. Thermodynamics of adsorption of a series of related organic molecules on graphite and on a carbon black, Part 2: Entropies of adsorption. *J. Chem. Soc., Faraday Trans. 1* 10:1784-96.
- Douglas, B. E., and D. H. McDaniel. 1965. *Concepts and Models of Inorganic Chemistry*. 1st ed. Waltham, Mass.: Blaisdell Publishing Co.
- Dowd, J. E., and D. S. Riggs. 1965. A comparison of estimates of Michaelis-Menten kinetic constants from various linear transformations. *J. Biol. Chem.* 240:863-69.
- Dowdy, B. H., and M. M. Mortland. 1968. Alcohol-water interactions on montmorillonite surfaces. II: Ethylene glycol. *Soil Sci.* 105:36-43.
- Eden, C., and D. Ashboren. 1972. The use of ternary phase diagrams in the study of composite adsorption isotherms. *J. Colloid Interface Sci.* 39:409-12.
- Edwards, C. A. 1964. Factors affecting the persistence of insecticides in soil. *Soils and Fertilizers* 27:451-54.
- Edwards, C. A. 1966. Insecticide residues in soils. *Residue Rev.* 13:83-132.
- Ensminger, L. E., and J. E. Gieseking. 1941. The adsorption of proteins by montmorillonitic clays and its effect on base-exchange capacity. *Soil Sci.* 51:125-32.
- Evcim, N., and M. Barr. 1955. Adsorption of some alkaloids by different clays. *J. Am. Pharm. Assoc.* 44:570-73.
- Everett, D. H. 1975. Adsorption and molecular structuring at the solid/liquid interface. *Isr. J. Chem.* 14:267-77.
- Everett, D. H., and W. I. Whitton. 1955. A thermodynamic study of the adsorption of benzene vapour by active charcoals. *Roy. Soc. London Proc.* 230A:91-110.
- Fadayomi, O., and G. F. Warren. 1971. Adsorption, desorption, and leaching of nitrofen and oxyfluorfen. *Weed Sci.* 25:97-100.
- Farmer, W. J. 1976. Leaching, diffusion, and sorption of benchmark pesticides. In *A Literature Survey of Benchmark Pesticides*, pp. 185-244. Washington, D.C.: George Washington University Medical Center.
- Felmeister, A., D. Tsai, and N. D. Weiner. 1972. Interaction of 3,4-benzpyrene with monomolecular films. *J. Pharm. Sci.* 61:1065-68.
- Fields, D. E. 1976. *CHNSED: Simulation of Sediment and Trace Contaminant Transport with Sediment/Contaminant Interaction*. Oak Ridge, Tenn.: Oak Ridge National Laboratory.

- Forrester, S. D., and C. H. Giles. 1972. From manure heaps to monolayers: One hundred years of solute-solvent adsorption isotherm studies. *Chem. Ind.* (London) 8:318-25.
- Frank, H. S., and M. W. Evans. 1945. Free volume and entropy in condensed systems. III: Entropy in binary liquid mixtures; partial molal entropy in dilute solutions; structure and thermodynamics in aqueous electrolytes. *J. Chem. Phys.* 13:507-32.
- Freundlich, H. 1922. *Colloid and Capillary Chemistry*. London: Methion & Co.
- Frissel, M. J., and G. H. Bolt. 1962. Interaction between certain ionizable organic compounds (herbicides) and clay minerals. *Soil Sci.* 94:284-91.
- Fu, Y., R. S. Hansen, and F. E. Bartell. 1949. Thermodynamics of adsorption from solution. II: Free energy changes and surface pressure-area relationships of adsorbed layers. *J. Phys. Colloid Chem.* 53:141-52.
- Fulk, R., D. Gruber, and R. Wullschleger. 1975. *Laboratory Study of the Release of Pesticide and PCB Materials to the Water Column During Dredging and Disposal Operations*. NTIS No. AD A026685. Vicksburg, Miss.: Environmental Effects Laboratory, U. S. Army Engineer Waterways Experiment Station.
- Furukawa, T., and G. W. Brindley. 1973. Adsorption and oxidation of benzidine and aniline by montmorillonite and hectorite. *Clays Clay Miner.* 21:279-88.
- Gabriel, H., and R. J. Cooky. 1955. Adsorption of a homologous series of aliphatic acids on a nonionic adsorbent. *Ind. Eng. Chem. Fundam.* 47: 1236-39.
- Geissbühler, H., C. Haselbach, H. Aebi, and L. Ebner. 1963. The fate of N'-(4-chlorophenoxy)-phenyl-NN-dimethylurea (C-1983) in soils and plants. *Weed Res.* 3:181-94.
- Gerakis, P. A., and A. G. Sficas. 1974. The presence and cycling of pesticides in the ecosphere. *Residue Rev.* 52:69-87.
- Gerolt, P. 1961. Investigation into the problem of insecticide sorption by soils. *Bull. World Health Org.* 24:577-92.
- Ghosh, A. K., P. K. Bandopadhyay, S. K. Ghosh, and D. P. Rajwar. 1974. Adsorption by flow method and determination of BET surface area by elution of the sorbed component. *Indian J. Technol.* 12:34-37.
- Giesecking, J. E. 1939. The mechanism of cation exchange in the montmorillonite-beidellite-montronite type of clay minerals. *Soil Sci.* 47:1-13.

- Giles, C. H., H. V. Mehta, S. M. K. Rahman, and C. E. Stewart. 1959. Adsorption at inorganic surfaces. V: Adsorption of sulphonated dyes by the anodic film on aluminium. *J. Appl. Chem.* 9:457-66.
- Giles, C. H., H. V. Mehta, C. E. Stewart, and R. V. R. Subramanian. 1954. Adsorption at inorganic surfaces, Part 1: An investigation into the mechanism of adsorption of organic compounds by the anodic film on aluminium. *J. Chem. Soc.* pp. 4360-74.
- Giles, C. H., T. H. MacEwan, S. N. Nakhwa, and D. Smith. 1960. Studies in adsorption, Part XI: A system of classification of solution adsorption isotherms, and its use in diagnosis of adsorption mechanisms and in measurement of specific surface areas of solids. *J. Chem. Soc.* pp. 3973-93.
- Graham, D. 1954. The characterization of physical adsorption systems. II: The effects of attractive interaction between adsorbed molecules. *J. Phys. Chem.* 58:869-72.
- Green, R. E., and J. C. Corey. 1971. Pesticide adsorption measurement by flow equilibration and subsequent displacement. *Soil Sci. Soc. Am., Proc.* 35:561-65.
- Greene-Kelly, R. 1955. An unusual montmorillonite complex. *Clay Miner. Bull.* 2:226-32.
- Greene-Kelly, R. 1955. Sorption of aromatic organic compounds by montmorillonite, Part 1: Orientation studies. *Trans. Faraday Soc.* 51:412-24.
- Greene-Kelly, R. 1955. Sorption of aromatic organic compounds by montmorillonite, Part 2: Packing studies with pyridine. *Trans. Faraday Soc.* 51:425-30.
- Greenland, D. J. 1965. Interaction between clays and organic compounds in soil, Part 1: Mechanism of interaction between clays and defined organic compounds. *Soils and Fertilizers* 28:415-24.
- Greenland, D. J., and E. W. Russell. 1955. Organo-clay derivatives and the origin of the negative charge on clay particles. *Trans. Faraday Soc.* 51:1300.
- Groszek, A. J. 1975. Adsorption of polycyclic aromatic hydrocarbons onto graphite. *Faraday Discuss. Chem. Soc.* 59:109-16.
- Groves, T. E., S. T. Bowden, and W. J. Jones. 1947. Quantity of adsorbent and temperature as factors in adsorption from solution. *Recl. Trav. Chim. Pays-Bas* 66:645-54.
- Gustafson, R. L., R. L. Albright, J. Heisler, J. A. Lirio, and O. T. Reid, Jr. 1968. Adsorption of organic species by high surface area styrene-divinyl-benzene copolymers. *Ind. Eng. Chem., Prod. Res. Dev.* 7:107-15.

- Gustafson, R. L., and J. Paleos. 1971. Interactions responsible for the selective adsorption of organics on organic surfaces. In *Organic Compounds in Aquatic Environments*, ed. S. D. Faust and J. V. Hunter, pp. 213-37. New York: Marcel Dekker.
- Hadaway, A. B., and F. Barlow. 1949. Further studies on the loss of insecticides by adsorption into mud and vegetation. *Bull. of Entomological Res.* 40: 323-43.
- Hadaway, A. B., and F. Barlow. 1951. Sorption of solid insecticides by dried mud. *Nature* 167:854.
- Hadaway, A. B., and F. Barlow. 1952. Studies on aqueous suspensions of insecticides, Part III: Factors affecting the persistence of some synthetic insecticides. *Bull. Entomological Res.* 43:281-311.
- Hadaway, A. B., and F. Barlow. 1964. A note on the sorption of insecticides on tropical soils. *Bull. World Health Org.* 30:146-48.
- Hamaker, J. W. 1966. Mathematical prediction of cumulative levels of pesticides in soils. In *Organic Pesticides in the Environment*, ed. A. A. Rosen and H. F. Kraybill, pp. 122-31. Advances in Chemistry Series No. 60. American Chemical Society.
- Hamaker, J. W. 1974. The interpretation of soil leaching experiments. In *Environmental Dynamics of Pesticides*, ed. R. Haque and V. H. Freed. New York: Plenum Press.
- Hance, R. J. 1967. Relationship between partition data and the adsorption of some herbicides by soils. *Nature* 214:630-31.
- Hance, R. J. 1969. An empirical relationship between chemical structure and the sorption of some herbicides by soils. *J. Agr. Food Chem.* 17:667-68.
- Hance, R. J. 1969. Influence of pH, exchangeable cation and the presence of organic matter on the adsorption of some herbicides by montmorillonite. *Can. J. Soil Sci.* 49:357-64.
- Hance, R. J. 1977. The adsorption of atraton and monuron by soils at different water contents. *Weed Res.* 17:197-201.
- Hansch, C., and F. Helmer. 1968. Extrathermodynamic approach to the study of the adsorption of organic compounds by macromolecules. *J. Polym. Sci., Polym. Chem.* 6:3295-3302.
- Hansch, C., J. E. Quinlan, and G. L. Lawrence. 1968. The linear free-energy relationship between partition coefficients and the aqueous solubility of organic liquids. *J. Org. Chem.* 33:347-50.
- Hansen, R. S., and R. P. Craig. 1954. The adsorption of aliphatic alcohols and acids from aqueous solutions by non-porous carbons. *J. Phys.*

Chem. 58:211-15.

- Hansen, R. S., Y. Fu, and F. E. Bartell. 1949. Multimolecular adsorption from binary liquid solutions. *J. Phys. Colloid Chem.* 53:769-85.
- Haque, R. and V. H. Freed. 1974. Behavior of pesticides in the environment: Environmental chemodynamics. *Residue Rev.* 52:89-116.
- Harris, C. I. 1966. Movement of herbicides in soil. *Weeds* 14:214-16.
- Harris, C. I., D. D. Kaufman, T. J. Sheets, R. G. Nash, and P. C. Kearney. 1968. Behavior and fate of s-triazines in soils. *Adv. Pest Control Res.* 8:1-55.
- Harris, C. I., E. A. Woolson, and B. E. Hummer. 1969. Dissipation of herbicides at three soil depths. *Weed Sci.* 17:27-31.
- Harris, C. R. 1966. Influence of soil type on the activity of insecticides in soil. *J. Econ. Entomol.* 59:1221-25.
- Harris, C. R., and E. P. Lichtenstein. 1961. Factors affecting the volatilization of insecticidal residues from soils. *J. Econ. Entomol.* 54:1038-45.
- Harris, C. R., and W. W. Sans. 1972. Behavior of heptachlor epoxide in soil. *J. Econ. Entomol.* 65:336-40.
- Harter, R. D., and J. L. Ahlrichs. 1969. Effect of acidity on reactions of organic acids and amines with montmorillonite clay surfaces. *Soil Sci. Soc. Am., Proc.* 33:859-63.
- Hartley, G. S. 1964. Herbicide behavior in the soil. I: Physical factors and action through the soil. In *The Physiology and Biochemistry of Herbicides*, ed. L. J. Audus, pp. 111-61. New York: Academic Press.
- Hartman, R. J., R. A. Kern, and E. G. Bobalek. 1946. Adsorption isotherms of some substituted benzoic acids. *J. Colloid Sci.* 1:271-76.
- Hayes, M. H. B., M. E. Pick, and B. A. Toms. 1975. Interactions between clay minerals and bipyridylum herbicides. *Res. Rev.* 57:1-25.
- Healey, T. W. 1971. Selective adsorption of organics on inorganic surfaces. In *Organic Compounds in Aquatic Environments*, ed. S. J. Faust and J. V. Hunter, pp. 187-212. New York: Marcel Dekker.
- Helling, C. S. 1970. Movement of s-triazine herbicides in soils. *Residue Rev.* 32:175-210.
- Helling, C. S., G. Chesters, and R. B. Corey. 1964. Contribution of organic matter and clay to soil cation-exchange capacity as affected by the pH of the saturating solution. *Soil Sci. Soc. Am., Proc.* 28:517-20.

- Hendricks, S. B. 1941. Base exchange of the clay mineral montmorillonite for organic cation and its dependence upon adsorption due to van der Waals forces. *J. Phys. Chem.* 45:65-81.
- Henry, D. C. 1922. A kinetic theory of adsorption. *Philos. Mag.* 44:689-705.
- Hofstee, B. H. J. 1952. On the evaluation of the constants V_m and K_M in enzyme reactions. *Science* 116:329-31.
- Hofstee, B. H. J. 1960. Nonlogarithmic linear titration curves. *Science* 131:39.
- Holford, I. C. R., R. W. M. Wedderburn, and G. E. G. Mattingly. 1974. A Langmuir two-surface equation as a model for phosphate adsorption by soils. *J. Soil Sci.* 25:242-55.
- Hsu, C. C., W. Rudzinski, and B. W. Wojciechowski. 1976. A new isotherm for multilayer adsorption on heterogeneous surfaces. *J. Am. Chem. Soc., Faraday Trans. 1* 72:453.
- Huang, J., and C. Liao. 1970. Adsorption of pesticides by clay minerals. *J. San. Eng. Div. Proc., Am. Soc. Civil Eng.* 96:1057-78.
- Huang, P. M., T. S. C. Wang, M. K. Wang, M. H. Wu, and N. W. Shu. 1977. Retention of phenolic acids by noncrystalline hydroxy-aluminum and iron compounds and clay minerals of soils. *Soil Sci.* 123:213-19.
- Inscoc, M. N. 1964. Photochemical changes in thin layer chromatograms of polycyclic, aromatic hydrocarbons. *Anal. Chem.* 36:2505-6.
- Jaroniec, M. 1975. Adsorption of gas mixtures on heterogeneous surfaces: Analytical solution of integral equation for Jovanović adsorption isotherm. *J. Colloid Interface Sci.* 53:422-28.
- Jaroniec, M. 1975. Adsorption of gas mixtures on homogeneous surfaces: Extension of Jovanović equation on adsorption from gaseous mixtures. *Chem. Zvesti* 29:512-16.
- Jaroniec, M., and W. Rudzinski. 1975. Adsorption of gas mixtures on heterogeneous surfaces: The integral representation for a monolayer total adsorption isotherm. *Surf. Sci.* 52:641-52.
- Jaroniec, M., S. Sokolowski, and W. Rudzinski. 1976. Some remarks on the maximum adsorption energy. *Surf. Sci.* 54:189-93.
- Jellinek, H. H. G., and H. L. Northey. 1954. Adsorption of high polymers from solution on to solids. II: Adsorption of polystyrene on charcoal. *J. Polym. Sci.* 14:583-87.

- John, P. T., and R. K. Aggarwal. 1975. An adsorption isotherm for determining monolayer capacity at any relative pressure and mean pore size. *Indian J. Technol.* 13:556-60.
- John, P. T., and K. K. Datta. 1974. Derivation of an adsorption equation. *Indian J. Technol.* 12:34.
- Jordan, J. W. and F. J. Williams. 1954. Organophilic bentonites. III: Inherent properties. *Kolloid Zeitschrift* 137:40-48.
- Kalb, G. W., and R. B. Curry. 1969. Determination of surface area by surfactant adsorption in aqueous suspension. I: Dodecylamine hydrochloride. *Clays Clay Miner.* 17:47-57.
- Kamprath, E. J., and C. D. Welch. 1962. Retention and cation-exchange properties of organic matter in coastal plain soils. *Soil Sci. Soc. Am., Proc.* 26:263-65.
- Karickhoff, S. W., and G. W. Bailey. 1976. Protonation of organic bases in clay-water systems. *Clays Clay Miner.* 24:170-76.
- Karickhoff, S. W., D. S. Brown, and T. A. Scott. 1979. Sorption of hydrophobic pollutants on natural sediments. *Water Res.* (in press).
- Karpinski, K., and J. Garbacz. 1973. A modified equation of adsorption isotherm in non-localized monolayers on homogeneous surfaces. *Rocz. Chem.* 47:2179-82.
- Kavanagh, B. V., A. M. Posner, and J. P. Quirk. 1976. The adsorption of polyvinyl alcohol on gibbsite and goethite. *J. Soil Sci.* 27:467-77.
- Kawale, G. B., V. D. Joglekar, V. P. Barve, and H. S. Mahal. 1972. Use of mercurous nitrate as a spray reagent for the detection of organic compounds on TLC. *Sci. Cult.* 38:373-76.
- Khan, S. V. 1974. Humic substances reactions involving bipyridylum herbicides in soil and aquatic environments. *Res. Rev.* 52:1-26.
- Kington, G. L., and W. Laing. 1955. The crystal structure of charbazite and its sorptive properties. *Trans. Faraday Soc.* 51:287-98.
- Kipling, J. J. 1965. *Adsorption from Solutions of Non-electrolytes*. New York: Academic Press.
- Knight, B. A. G., and P. J. Denny. 1970. The interaction of paraquat with soil: Adsorption by an expanding lattice clay mineral. *Weed Res.* 10:40-8.
- Knight, B. A. G., and T. E. Tomlinson. 1967. The interaction of paraquat (I:I'-dimethyl 4:4'-dipyridylum dichloride) with mineral soils. *J. Soil Sci.* 18:233-43.

- Kown, B. T., and B. B. Ewing. 1969. Effects of the organic adsorption on clay ion-exchange property. *Soil Sci.* 108:321-25.
- Kratochvic, V., M. Nepras, and K. Obruba. 1974. The basicity of mono-halogeno derivatives of 9,10-anthraquinone and their chromatography on a thin-layer of silica gel. *Collect. Czech. Chem. Commun.* 39:271-74.
- Kressman, T. R. E. 1952. Ion exchange separations based upon ionic size. *J. Phys. Chem.* 56:118-23.
- LaFleur, K. S. 1974. Toxaphene-soil-solvent interactions. *Soil Sci.* 117:205-10.
- LaFleur, K. S. 1973. Fluometuron-soil-solvent interactions. *Soil Sci.* 116:376-82.
- LaFleur, K. S. 1976. Carbaryl desorption and movement in soil columns. *Soil Sci.* 121:212-16.
- LaFleur, K. S. 1976. Prometryne desorption and movement in soil columns. *Soil Sci.* 121:9-15.
- Lahav, N. 1972. Interaction between montmorillonite and benzidine in aqueous solutions. III: The color reaction in the air dry state. *Isr. J. Chem.* 10:925-34.
- Lahav, N. 1973. Montmorillonite-benzidine reactions in the frozen and dry states. *Clays Clay Miner.* 21:137-39.
- Lahav, N., and D. M. Anderson. 1973. Interaction between montmorillonite and benzidine in aqueous solutions. IV: The color reaction in the frozen state. *Isr. J. Chem.* 11:549-55.
- Lahav, N., and S. Raziel. 1971. Interaction between montmorillonite and benzidine in aqueous solutions. I: Adsorption of benzidine on montmorillonite. *Isr. J. Chem.* 9:683-89.
- Lahav, N., and S. Raziel. 1971. Interaction between montmorillonite and benzidine in aqueous solutions. II: A general kinetic study. *Isr. J. Chem.* 9:691-94.
- Lailach, G. E., T. D. Thompson, and G. W. Brindley. 1968. Adsorption of pyrimidines, purines, and nucleosides by Co-, Ni-, Cu-, and Fe (III)-montmorillonite. *Clays Clay Miner.* 16:295-301.
- Lailach, G. E., T. D. Thompson, and G. W. Brindley. 1968. Adsorption of pyrimidines, purines, and nucleosides by Li-, Na-, Mg-, and Ca-montmorillonite. *Clays Clay Miner.* 16:285-93.
- Lambert, S. M. 1966. The influence of soil-moisture content on herbicidal response. *Weeds* 14:273-75.

- Lambert, S. M. 1967. Functional relationship between sorption in soil and chemical structure. *J. Agric. Food Chem.* 15:572-76.
- Lambert, S. M. 1968. Omega (Ω), a useful index of soil sorption equilibria. *J. Agric. Food Chem.* 16:340-43.
- Lambert, S. M., P. E. Porter, and R. H. Schieferstein. 1965. Movement and sorption of chemicals applied to the soil. *Weeds* 13:185-90.
- Langmuir, I. 1918. The adsorption of gases on plane surfaces of glass, mica and platinum. *J. Am. Chem. Soc.* 40:1361-1403.
- Lavy, T. L. 1968. Micromovement mechanisms of s-triazine in soil. *Soil Sci. Soc. Am., Proc.* 32:377-80.
- Lawrence, J., and H. M. Tosine. 1976. Adsorption of polychlorinated biphenyls from aqueous solutions and sewage. *Environ. Sci. Technol.* 10:381-83.
- LeFrancois, M., and G. Malbois. 1971. The nature of the acidic sites on mordenite: Characterization of adsorbed pyridine and water by infra-red study. *J. Catal.* 20:350-58.
- Leistra, M. 1973. Computation models for the transport of pesticides in soil. *Res. Rev.* 32:87-130.
- Leo, A., C. Hansch, and D. Elkins. 1971. Partition coefficients and their uses. *Chem. Rev.* 71:525-616.
- Leo, A., P. Y. C. Jow, C. Silipo, and C. Hansch. 1975. Calculation of hydrophobic constant (log P) from π and f constants. *J. Med. Chem.* 18:865-68.
- Leopold, A. C., P. van Schaik, and M. Neal. 1960. Molecular structure and herbicide adsorption. *Weeds* 8:48-54.
- Lewis, T. E., and F. E. Broadbent. 1961. Soil organic matter-metal complexes. 4: Nature and properties of exchange sites. *Soil Sci.* 91:393-99.
- Lichtenstein, E. P. 1958. Movement of insecticides in soils under leaching and non-leaching conditions. *J. Econ. Entomol.* 51:380-83.
- Lichtenstein, E. P., and J. B. Polivka. 1959. Persistence of some chlorinated hydrocarbon insecticides in turf soils. *J. Econ. Entomol.* 52:289-93.
- Lichtenstein, E. P., and K. R. Schulz. 1959. Persistence of some chlorinated hydrocarbon insecticides as influenced by soil types, rate of application and temperature. *J. Econ. Entomol.* 52:124-31.

- Lindstrom, F. T., R. Haque, V. H. Freed, and L. Boersma. 1967. Theory on the movement of some herbicides in soils linear diffusion and convection of chemicals in soils. *Environ. Sci. Technol.* 1:561-65.
- Linner, E. R., and R. A. Gortner. 1935. Interfacial energy and the molecular structure of organic compounds. III: The effect of organic structure on adsorbability. *J. Phys. Chem.* 39:35-67.
- Lloyd, C. L., and B. L. Harris. 1954. Binary liquid phase adsorption. *J. Phys. Chem.* 58:899-903.
- Luh, M. D., and R. A. Baker. 1970. Vapor phase sorption of phenol on selected clays. *J. Colloid Interface Sci.* 33:539-47.
- Luh, M. D., and R. A. Baker. 1971. Sorption and desorption of pyridine-clay in aqueous solution. *Water Res.* 5:849-59.
- MacEwan, D. M. C. 1948. Complex formation between montmorillonite and halloysite and certain organic liquids. *Trans. Faraday Soc.* 44:349-67.
- McGinnes, P. R., and V. L. Snoeyink. 1974. *Determination of the Fate of Polynuclear Aromatic Hydrocarbons in Natural Water Systems*. Urbana, Ill.: Water Resources Center, University of Illinois at Urbana-Champaign.
- Mackay, D. 1977. Partition coefficients and bioaccumulation of selected organic chemicals. *Environ. Sci. Technol.* 11:1219-20.
- Mackay, D., and W. K. Shiu. 1977. Aqueous solubility of polynuclear aromatic hydrocarbons. *J. Chem. Eng. Data* 22:399-402.
- Mackenzie, R. C. 1948. Complexes of clays with organic compounds, Part II: Investigation of the ethylene glycol-water montmorillonite system using the Karl Fischer reagent. *Trans. Faraday Soc.* 44:368-75.
- Majka, J. T., and T. L. Lavy. 1977. Adsorption, mobility, and degradation of cyanazine and diuron in soils. *Weed Sci.* 25:401-6.
- Masuda, T., and H. Takahashi. 1975. Calorimetric study of the interaction of montmorillonite with amines. *Bull. Inst. Chem. Res., Kyoto Univ.* 53:147-52.
- Miguel, A. H. 1976. I: Pyrene adsorption onto fly ash; II: Thiol oxidation and adsorption by activated carbon. In *Studies of Gas-Solid Reactions of Environmental Significance*, Ph.D. dissertation, University of Illinois at Urbana-Champaign.
- Mustafa, M. A., and Y. Gamar. 1972. Adsorption and desorption of diuron as a function of soil properties. *Soil Sci. Soc. Am., Proc.* 36:561-65.

- Nearpass, D. C. 1969. Exchange adsorption of 3-amino-1,2,4-triazole by an organic soil. *Soil Sci. Soc. Am., Proc.* 33:524-28.
- Nearpass, D. C. 1971. Adsorption interactions in soils between amitrole and s-triazines. *Soil Sci. Soc. Am., Proc.* 35:64-71.
- Nearpass, D. C. 1976. Adsorption of picloram by humic acids and humin. *Soil Sci.* 121:272-77.
- Nogami, H., T. Nagai, and H. Uchida. 1968. Physico-chemical approach to biopharmaceutical phenomena. III: Hydrophobic bonding in the adsorption of tryptophan by carbon black from aqueous solution. *Chem. Pharm. Bull.* 16:2263-66.
- Nogami, H., T. Nagai, and H. Uchida. 1969. Physico-chemical approach to biopharmaceutical phenomena. IV: Adsorption of barbituric acid derivatives by carbon black from aqueous solution. *Chem. Pharm. Bull.* 17:168-75.
- Nomura, N. S., and H. W. Hilton. 1971. The adsorption and degradation of glyphosate in five Hawaiian sugarcane soils. *Weed Res.* 17:113-21.
- O'Connor, G. A., and J. U. Anderson. 1974. Soil factors affecting the adsorption of 2,4,5-T. *Soil Sci. Soc. Am., Proc.* 38:433-36.
- Oddson, J. K., J. Letey, and L. V. Weeks. 1970. Predicted distribution of organic chemicals in solution and adsorbed as a function of position and time for various chemical and soil properties. *Soil Sci. Soc. Am., Proc.* 34:412-17.
- Oehme, C., and F. Martinola. 1973. Removal of organic matter from water by resinous adsorbents. *Chem. Ind. (London)* 1:823-26.
- Parfitt, R. L., and D. J. Greenland. 1970. Adsorption of polysaccharides by montmorillonite. *Soil Sci. Soc. Am., Proc.* 34:862-66.
- Parkash, S. 1974. Adsorption of weak and non-electrolytes by activated carbon. *Carbon* 12:37-43.
- Parochetti, J. V. 1973. Soil organic matter effect on activity of acetanilides, CDAA, and atrazine. *Weed Sci.* 21:157-60.
- Parry, E. P. 1963. An infrared study of pyridine adsorbed on acidic solids: Characterization of surface acidity. *J. Catal.* 2:371-79.
- Patrick, W. A., and D. C. Jones. 1925. Studies in the adsorption from solution from the standpoint of capillarity, I. *J. Phys. Chem.* 29:1-10.
- Pionke, H. B., and G. Chesters. 1973. Pesticide-sediment-water interactions. *J. Environ. Qual.* 2:29-45.

- Polley, M. H., W. D. Schaeffer, and W. R. Smith. 1955. Physical adsorption studies in carbon black technology. *Can. J. Chem.* 33:314-19.
- Pylev, L. N., and G. D. Yankova. 1974. Possible adsorption of benzo(a) pyrene and its level in some Soviet industrial carbon blacks. *Gig. Tr. Prof. Zabol.* pp. 52-53.
- Radke, C. J., J. M. Pravsnitz. 1972. Adsorption of organic solutes from dilute aqueous solution on activated carbon. *Ind. Chem. Eng., Fundam.* 11:445-51.
- Richmond, P. 1977. Note on multilayer adsorption and wetting. *J. Chem. Soc., Faraday Trans. 2* 2:316-20.
- Roberts, A. L., G. B. Street, and D. White. 1964. The mechanism of phenol adsorption by organo-clay derivatives. *J. Appl. Chem.* 14:261-65.
- Rosenberg, J. L., and I. Brinn. 1972. Excited state dissociation rate constants in naphthols. *J. Phys. Chem.* 76:3558-62.
- Russell, J. D., M. J. Cruz, and J. L. White. 1968. The adsorption of 3-aminotriazole by montmorillonite. *J. Agr. Food Chem.* 16:21-24.
- Saltzman, S., and S. Yariv. 1975. Infrared study of the sorption of phenol and p-nitrophenol by montmorillonite. *Soil Sci. Soc. Am., Proc.* 39:474-79.
- Schnitzer, M. 1971. Metal-organic matter interactions in soils and waters. In *Organic Compounds in Aquatic Environments*, ed. S. J. Faust and J. V. Hunter, pp. 237-315. New York: Marcel Dekker.
- Schreiber, H. P., and R. McIntosh. 1954. Some thermodynamic properties of hydrocarbons adsorbed on rutile. *Can. J. Chem.* 32:842-57.
- Scott, D. C., and J. B. Weber. 1967. Herbicide phytotoxicity as influenced by adsorption. *Soil Sci.* 104:151-58.
- Scott, H. D., and J. F. Lutz. 1971. Release of herbicides from clay minerals as a function of water content. I: Kaolinite. *Soil Sci. Soc. Am., Proc.* 35:374-79.
- Shabad, L. M. 1968. On the distribution and the fate of the carcinogenic hydrocarbon benz(a)pyrene (3,4 benzpyrene) in the soil. *Z. Krebsforsch. Klin. Onkol.* 70:204-10.
- Sheets, J. T., A. S. Crafts, and H. R. Drever. 1962. Influence of soil properties on the phytotoxicities of the s-triazine herbicides. *J. Agric. Food Chem.* 10:458-62.
- Simsman, G. V., T. C. Damel, and G. Chesters. 1976. Diquat and endothall: Their fates in the environment. *Res. Rev.* 62:131-74.

- Singhal, J. P., and D. Kumar. 1976. Adsorption of telone on kaolinite, Part 1. *Soil Sci.* 21:156-61.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, D. C. Bomberger, and T. Mill. 1977. *Environmental Pathways of Selected Chemicals in Freshwater Systems. Part II: Laboratory Studies*. Athens, Ga.: Environmental Research Laboratory, Office of Research and Development, U. S. Environmental Protection Agency.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, T. Mill, and D. C. Bomberger. 1976. *The Fate of p-Cresol in Freshwater Aquatic Systems*. Cincinnati, Ohio: National Environmental Research Center, Office of Research and Development, United States Environmental Protection Agency (in draft).
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, T. Mill, and D. C. Bomberger. 1976. *The Fate of Quinoline in Freshwater Aquatic Systems*. Cincinnati, Ohio: National Environmental Research Center, Office of Research and Development, U. S. Environmental Protection Agency.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, T. Mill, and D. C. Bomberger. 1976. *Laboratory Investigation of Benzo(a)pyrene*. Cincinnati, Ohio: National Environmental Research Center, Office of Research and Development, U. S. Environmental Protection Agency.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, T. Mill, and D. C. Bomberger. 1976. *Laboratory Investigation of Benzo(b)thiophene*. Cincinnati, Ohio: National Environmental Research Center, Office of Research and Development, U. S. Environmental Protection Agency.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, T. Mill, and D. C. Bomberger. 1977. Laboratory investigation of 7H-dibenzo(c,g)carbazole. In *The Fate of Selected Pollutants in Freshwater Aquatic Systems, Part II: Laboratory Studies*, pp. 169-93. Athens, Ga.: Environmental Research Laboratory, Office of Research and Development, U. S. Environmental Protection Agency.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. W. Chou, T. Mill, and D. C. Bomberger. 1977. Laboratory investigation of 9H-carbazole. In *The Fate of Selected Pollutants in Freshwater Aquatic Systems, Part II: Laboratory Studies*, pp. 145-68. Athens, Ga.: Environmental Research Laboratory, Office of Research and Development, U. S. Environmental Protection Agency.
- Smith, J. H., W. R. Mabey, N. Bohonos, B. R. Holt, S. S. Lee, T. Mill, and D. C. Bomberger. 1976. *The Fate of Methyl Parathion in Freshwater Aquatic Systems*. Cincinnati, Ohio: National Environmental Research Center, Office of Research and Development, U. S. Environmental Protection Agency.

- Solomon, D. H. 1968. Clay minerals as electron acceptors and/or electron donors in organic reactions. *Clays Clay Miner.* 16:31-39.
- Spahn, H., V. Brauch, E. U. Schlunder, and H. Sontheimer. 1974. Explanation of activated charcoal filters for water purification. Part I: Investigation of the adsorption on single granules. *Verfahrenstechnik* 8:224-31.
- Steele, W. A. 1974. Monolayer adsorption. In *The Interaction of Gases with Solid Surfaces*, pp. 128-219. Oxford: Pergamon Press.
- Steele, W. A. 1974. Multilayer adsorption. In *The Interaction of Gases with Solid Surfaces*, pp. 221-69. Oxford: Pergamon Press.
- Strazhesko, D. N., and L. A. Tarkovskaya. 1973. Ion-exchange and sorptive properties of activated charcoals: Chemical nature of the surface selective ion exchange, and surface complex-formation on oxidized charcoal. In *Adsorption and Adsorbents*, ed. D. N. Strazhesko, pp. 7-24 New York: John Wiley & Sons.
- Street, G. B., and D. White. 1963. The adsorption of phenol by organo-clay derivatives. *J. Appl. Chem.* 13:203-6.
- Stumm, W., and J. J. Morgan. 1970. *Aquatic Chemistry*. New York: Wiley Interscience.
- Suess, M. J. 1964. Retardation of ABS in different aquifers. *J. Am. Water Works Assoc.* 56:89-91.
- Swanson, C. L. W., F. C. Thorp, and R. B. Friend. 1954. Adsorption of lindane by soils. *Soil Sci.* 78:379-88.
- Swoboda, A. R., and G. W. Kunze. 1964. Infrared study of pyridine adsorbed on montmorillonite surfaces. *Clays Clay Miner.* 25:277-88.
- Swoboda, A. R., and G. W. Thomas. 1968. Movement of parathion in soil columns. *J. Agr. Food Chem.* 16:923-27.
- Syers, J. K., M. G. Browman, G. W. Smillie, and R. B. Corey. 1973. Phosphate sorption by soils evaluated by the Langmuir adsorption equation. *Soil Sci. Soc. Am., Proc.* 37:358-63.
- Talibudeen, O. 1955. Complex formation between montmorillonoid clays and amino-acids and proteins. *Trans. Faraday Soc.* 51:581-90.
- Tamamushi, B., and K. Tamaki. 1959. Adsorption of long-chain electrolytes at the solid/liquid interface, Part 2: The adsorption on polar and non-polar adsorbents. *Trans. Faraday Soc.* 55:1007-12.
- Tennakoon, D. T. B., J. M. Thomas, and M. J. Tricker. 1974. Surface and intercalate chemistry of layered silicates, Part 1: General intro-

- duction and the uptake of benzidine and related organic molecules by montmorillonite. *J. Chem. Soc., Dalton Trans.* 20:2207-11.
- Terriere, L. C., and D. W. Ingalsbe. 1953. Translocation and residual action of soil insecticides. *J. Econ. Entomol.* 46:751-53.
- Theng, B. K. G. 1971. Adsorption of molybdate by some crystalline and amorphous soil clays. *New Zealand J. Sci.* 14:1040-56.
- Theng, B. K. G. 1971. Mechanisms of formation of colored clay-organic complexes: A review. *Clays Clay Miner.* 19:383-90.
- Tikhomirova, N. N., I. V. Nikolaeva, E. N. Rosolovskaya, V. V. Demkin, and K. V. Topchieva. 1975. Electron spin resonance study of various cation-exchanged zeolites: Adsorption of various species. *J. Catal.* 40:61-66.
- Towers, G. H. N., and D. C. Mortimer. 1954. Alternation in the adsorption of aliphatic acids on porous carbon. *Nature* 174:1188-89.
- Tucker, B. V., D. E. Pack, and J. N. Ospenson. 1967. Adsorption of bipyridylum herbicides in soil. *J. Agr. Food Chem.* 15:1005-8.
- Turnipseed, S. G., and J. K. Reed. 1963. The rate of leaching of dieldrin from attapulgite clay granules. *J. Econ. Entomol.* 56:410-12.
- Van Middeltem, C. H. 1966. Fate and persistence of organic pesticides in the environment. In *Organic Pesticides in the Environment*, ed. R. F. Gould, pp. 228-49. Washington, D. C.: American Chemical Society.
- Veith, J. A., and G. Sposito. 1977. On the use of the Langmuir equation in the interpretation of "adsorption" phenomena. *Soil Sci. Soc. Am. J.* 41:697-702.
- Wahid, P. A., and N. Sethunathan. 1978. Sorption-desorption of parathion in soils. *J. Agric. Food Chem.* 26:101-5.
- Wang, L. K., R. P. Leonard, M. H. Wang, and D. W. Goupil. 1975. Adsorption of dissolved organics from industrial effluents onto activated carbon. *J. Appl. Chem. Biotechnol.* 25:491-502.
- Ward, T. M., and F. W. Getzen. 1970. Influence of pH on the adsorption of aromatic acids on activated carbon. *Environ. Sci. Technol.* 4:64-67.
- Ward, T. M., and R. P. Upchurch. 1965. Role of the amido group in adsorption mechanisms. *J. Agric. Food Chem.* 13:334-40.
- Watson, J. R., A. M. Posner, and J. P. Quirk. 1973. Adsorption of the herbicide 2,4-D on goethite. *J. Soil Sci.* 24:503-11.

- Weber, J. B. 1970. Adsorption of s-triazines by montmorillonite as a function of pH and molecular structure. *Soil Sci. Soc. Am., Proc.* 34:401-4.
- Weber, J. B. 1972. Interaction of organic pesticides with particulate matter in aquatic and soil systems. In *Fate of Organic Pesticides in the Aquatic Environment*, ed. R. F. Gould, pp. 55-120. Advances in Chemistry, Series 111. Washington, D. C.: American Chemical Society.
- Weber, T. B., P. W. Perry, and R. P. Upchurch. 1965. The influence of temperature and time on the adsorption of paraquat, diquat, 2,4-D and prometone by clays, charcoal, and an anion-exchange resin. *Soil Sci. Soc. Am., Proc.* 29:678-88.
- Weber, J. B., and S. B. Weed. 1968. Adsorption and desorption of diquat, paraquat, and prometone by montmorillonite and kaolinitic clay minerals. *Soil Sci. Soc. Am., Proc.* 32:485-87.
- Weber, J. B., S. B. Weed, and T. J. Sheets. 1972. Pesticides: How they move and react in the soil. *Crops and Soils* 25:14-17.
- Weber, W. J., Jr., and J. P. Gould. 1966. Sorption of organic pesticides from aqueous solution. In *Organic Pesticides in the Environment*, ed. R. F. Gould, pp. 280-304. Washington, D. C.: American Chemical Society.
- Weber, W. J., and J. C. Morris. 1963. Kinetics of adsorption on carbon from solution. *J. San. Eng. Div., Proc. Am. Soc. Civil Eng.* 69:31-59.
- Weed, S. B., and J. B. Weber. 1969. The effect of cation exchange capacity on the retention of diquat²⁺ and paraquat²⁺ by three-layer type clay minerals. I: Adsorption and release. *Soil Sci. Soc. Am., Proc.* 33:379-82.
- Weed, S. B., and J. B. Weber. 1974. Pesticide-organic matter interactions. In *Pesticides in Soil and Water*, ed. W. D. Guenzi, pp. 39-66. Madison, Wis.: Soil Science Society of America.
- Wheatley, G. A. 1965. The assessment and persistence of residues of organochlorine insecticides in soils and their uptake by crops. *Ann. Appl. Bio.* 55:325-29.
- Yariv, S., J. D. Russel, and V. C. Farmer. 1966. Infrared study of the adsorption of benzoic acid and nitrobenzene in montmorillonite. *Isr. J. Chem.* 4:201-13.
- Zettlemoyer, A. C., and F. J. Micale. 1971. Solution adsorption thermodynamics for organics on surfaces. In *Organic Compounds in Aquatic Environments*, ed. S. Faust and J. Hunter, pp. 165-85. New York: Marcel Dekker.

ANALYTICAL PRODEDURES

- Abbott, D. C., A. S. Burridge, J. Thomson, and K. S. Webb. 1967. A thin-layer chromatographic screening test for organophosphorus pesticide residues. *Analyst* (London) 92:170-75.
- Abbott, D. C., and H. Egan. 1967. Determination of residues of organophosphorus pesticides in foods. *Analyst* (London) 92:475-92.
- Abbott, D. C., and J. Thomson. 1965. The application of thin-layer chromatographic techniques to the analysis of pesticide residues. *Residue Rev.* 11:1-59.
- Acheson, M. A., R. M. Harrison, R. Perry, and R. A. Wellings. 1976. Factors affecting the extraction and analysis of polynuclear aromatic hydrocarbons in water. *Water Res.* 10:207-12.
- Adams, J., K. Menzies, and P. Levins. 1977. *Selection and Evaluation of Sorbent Resins for the Collection of Organic Compounds*. Report No. EPA 600/7-77-044. Research Triangle Park, N.C.: Industrial Environmental Research Lab., Office of Research and Development, U.S. Environmental Protection Agency.
- Alexander, M., J. M. Duxbury, A. J. Francis, and J. Adamson. 1972. *Detection of Soil Microorganisms In Situ by Combined Gas Chromatography-Mass Spectrometry*. Report No. NCR 33-010-127. Ithaca, N.Y.: Cornell University.
- Alford, A. 1977. Environmental applications of mass spectrometry. *Bio-medical Mass Spectrometry* 4:1-22.
- Ames, B. 1975. *The Detection and Analysis of Mutagens*. Progress Report for 1 July 1974 to 30 June 1975. NTIS no. UCB-34P 156 X 7. Energy Research and Development Administration.
- Anderson, G. M. 1975. Quantitation of tryptophan metabolites in rat feces by thin-layer chromatography. *J. Chromatogr.* 105:323-28.
- Anwar, M., C. Hanson, and A. N. Patel. 1968. Gas chromatographic separation of pyridine homologues, chloroanilines and toluidines. *J. Chromatogr.* 34:529-30.
- Argauer, R., and J. D. Warthen, Jr. 1975. Separation of 1- and 2-naphthols and determination of trace amounts of 2-naphthyl methylcarbamate in carbaryl formulations by high pressure liquid chromatography with confirmation by spectrofluorometry. *Anal. Chem.* 47:2472-74.
- Arustamova, L. G., V. G. Berezkin, M. J. Rustamov, and N. T. Sultanov. 1977. Surface-layer sorbents for group analysis of aromatic hydrocarbons in petroleum distillates. *J. Chromatogr.* 140:319-21.

- Askew, J., J. H. Rusicka, and B. B. Wheals. 1969. A general method for the determination of organophosphorus pesticide residues in river water and effluents by gas, thin-layer and gel chromatography. *Analyst* (London) 14:275-83.
- Austern, B. M., R. A. Dobbs, and J. M. Cohen. 1975. Gas chromatographic determination of selected organic compounds added to wastewater. *Environ. Sci. Technol.* 9:588-90.
- Bardner, R., K. A. Lord, and S. R. B. Solly. 1963. A cholinesterase inhibition method of determining the distribution of organophosphorus insecticides in soils. *Chem. Ind.* (London) No. 3 (Jan. 19):123-24.
- Bark, L. S., and R. J. T. Graham. 1966. Studies in the relationship between molecular structure and chromatographic behavior. VIII: The reversed phase thin-layer chromatography of some halogenated phenols and some halogeno-; alkyl-substituted phenols. *J. Chromatogr.* 25:357-66.
- Baumgarten, E., F. Weinstrauch, and H. Höffkes. 1977. Adsorption isotherms of hydrocarbons on γ -alumina. *J. Chromatogr.* 138:347-54.
- Bell, J. H., S. Ireland, and A. W. Spears. 1969. Identification of aromatic ketones in cigarette smoke condensate. *Anal. Chem.* 41:310-13.
- Bellamy, L. J. 1955. The infrared spectra of substituted aromatic compounds in relation to chemical reactivities of their substituents. *J. Chem. Soc.*, Part III:2818-21.
- Bellar, T. A., and J. J. Lichtenberg. 1974. Determining volatile organics at microgram per litre levels by gas chromatography. *J. Am. Water Works Assoc.* 66:739-44.
- Bender, D. F. 1968. Thin-layer chromatographic separation and spectrophotofluorometric identification and estimation of dibenzo(*a,e*)pyrene. *Environ. Sci. Technol.* 2:204-6.
- Bergman, R. G., and C. A. Wachtmeister. 1976. Impregnation of silica gel with tetraalkylammonium salts in adsorption chromatography of neutral aromatic compounds. *J. Chromatogr.* 123:231-36.
- Bernofsky, C., and W. J. Gallagher. 1975. Liquid chromatography of pyridine nucleotides and associated compounds and isolation of several analogs of nicotinamide adenine dinucleotide phosphate. *Anal. Biochem.* 67:611-24.
- Bertsch, W., E. Anderson, and G. Holzer. 1976. Characterization of coal-derived fluids by capillary column gas chromatography mass spectrometry. *J. Chromatogr.* 126:213-24.
- Beugeling, T., M. Boduszynski, F. Goudriaan, and J. W. M. Sonnemans. 1971. Gas-liquid chromatographic analysis of products formed by the hydrogenolysis of pyridine. *Anal. Lett.* 4:727-35.

- Bhatia, K. 1976. Gas chromatographic determination of polycyclic aromatic hydrocarbons. *Anal. Chem.* 43:609-10.
- Blumer, M., and W. W. Youngblood. 1976. *Polycyclic Aromatic Hydrocarbons in the Environment: Homologous Series in Soils and Recent Marine Sediments*. NTIS No. AD AO 23637. Office of Naval Research.
- Boehm, P. D., and J. G. Quinn. 1973. *Solubilization of Hydrocarbons by the Dissolved Organic Matter in Sea Water*. Research report. Kingston, R.I.: Graduate School of Oceanography, University of Rhode Island.
- Borneff, J., and H. Kunte. 1969. Carcinogenic substances in water and soil. XXVI: A routine method for the determination of PAH in water. *Arch. Hyg. Bakt.* 153:220-29.
- Bose, K. S., and R. H. Sarma. 1975. Delineation of the intimate details of the backbone conformation of pyridine nucleotide coenzymes in aqueous solution. *Biochem. Biophys. Res. Commun.* 66:1173-79.
- Bosin, T. R., R. Buckpitt, and R. P. Maickel. 1974. Comparative gas-liquid chromatography of biologically important indoles, and their benzo(b)-thiophene and 1-methylindole analogs. *J. Chromatog.* 94:316-20.
- Bowen, B. E. 1976. Determination of aromatic amines by an adsorption technique with flame ionization gas chromatography. *Anal. Chem.* 48:1584-87.
- Brocco, D., V. Cantuti, and G. P. Cartoni. 1970. Determination of polynuclear hydrocarbons in atmospheric dust by a combination of thin-layer and gas chromatography. *J. Chromatogr.* 49:66-69.
- Brown, J. A., ed. 1976. *Proceedings of the 6th Annual Symposium on Trace Analysis and Detection in the Environment*. NTIS No. AD A021948. Aberdeen, Md.: Edgewood Arsenal, Aberdeen Proving Ground.
- Brown, J. L., and J. M. Johnston. 1962. Radioassay of lipid components separated by thin-layer chromatography. *J. Lipid Res.* 3:480-81.
- Burchfield, K. P., E. E. Green, R. J. Wheeler, and S. M. Billedeau. 1974. A direct gas-phase isolation and injection system for the analysis of polynuclear arenes in air particulates by gas-liquid chromatography. *J. Chromatogr.* 99:697-708.
- Byrd, D. J., W. Kochen, D. Idzko, and E. Knorr. 1974. The analysis of indolic tryptophan metabolites in human urine. *J. Chromatogr.* 94:85-106.
- Candeli, A., G. Morozzi, A. Paolacci, and L. Zoccolillo. 1975. Analysis using thin-layer and gas-liquid chromatography of polycyclic aromatic hydrocarbons in the exhaust products from a European car running on fuels containing a range of concentrations of these hydrocarbons. *Atmos. Environ.* 9:843-49.

- Carlson, R. M., R. E. Carlson, and H. L. Kopperman. 1975. Determination of partition coefficients by liquid chromatography. *J. Chromatogr.* 107:219-23.
- Caton, R. D., Jr., J. B. Matthews, and E. A. Walters. 1976. *Development of High Pressure Liquid Chromatographic Techniques*. NTIS No. AD/A039644. Tundall AFB, Fla.: Air Force Civil Engineering Center, Air Force Systems Command.
- Chamberlain, W. J., D. B. Walters, and O. T. Chartyk. 1975. Some pitfalls in high-pressure liquid chromatography. *Anal. Chem. Acta* 76:213-14.
- Chang, R. C. 1976. *Concentration and Determination of Trace Organic Pollutants in Water*. Ph.D dissertation, Ames Laboratory, Iowa State University.
- Chiba, M., and H. V. Mosley. 1968. Studies of losses of pesticides during sample preparation. *J. Assoc. Off. Anal. Chem.* 51:55-62.
- Clementi, S., G. Savelli, and M. Vergoni. 1972. Relative molar response of flame ionization detector to some heteroaromatic compounds. *Chromatographia* 5:413-14.
- Cowan, C. T., and J. M. Hartwell. 1961. An organo-clay complex for the separation of isomeric dichlorobenzenes using gas chromatography. *Nature* 190:712.
- Cronin, D. A., and J. Gilbert. 1974. A technique for the detection of nitrogen-containing compounds in gas chromatographic eluates by means of hydrogenolysis and colour reaction. *J. Chromatogr.* 89:209-14.
- Dhont, J. H., C. Vinkenborg, H. Compaan, F. J. Ritter, R. P. Labadie, A. Verweij, and R. A. De Zeeuw. 1972. Application of R_F correction in thin-layer chromatography by means of two reference R_F values II: Results obtained with a polar multi-component solvent system. *J. Chromatogr.* 71:283-89.
- Doran, T., and N. G. Taggart. 1974. The combined use of high efficiency liquid and capillary gas chromatography for the determination of polycyclic aromatic hydrocarbons in automotive exhaust condensates and other hydrocarbon mixtures. *J. Chromatogr.* 12:715-21.
- Dunn, B. P. 1976. Techniques for determination of benzo(a)pyrene in marine organisms and sediments. *Environ. Sci. Technol.* 10:1018-21.
- Durbin, D. E., and A. Zlatkis. 1970. A gas chromatographic method for the determination of pyridine and quinoline type bases. *J. Chromatogr. Sci.* 8:608-10.
- Eastin, E. F. 1972. Separation of SAN-6706 and some related compounds by thin-layer chromatography. *J. Chromatogr.* 66:386-87.

- Eichelberger, J. W., W. M. Middleton, and W. L. Budde. 1975. *Analytical Quality Assurance for Trace Organics Analysis by Gas Chromatography/Mass Spectrometry*. NTIS No. PB-245-823. Cincinnati, Ohio: Environmental Monitoring and Support Laboratory, Office of Research and Development, U. S. Environmental Protection Agency.
- Eon, C., C. Pommier, and G. Guiochon. 1971. Gas chromatographic study of donor-acceptor complexes. II: Associations between aromatic heterocyclic derivatives and dibutyltetrachlorophthalate. *Chromatographia* 4:241-49.
- Eshel, Y., and G. F. Warren. 1967. A simplified method for determining phytotoxicity, leaching, and adsorption of herbicides in soils. *Weeds* 15:115-18.
- Fishbein, L., and W. Zielinski, Jr. 1965. Gas chromatography of trimethylsilyl derivatives. 1: Pesticidal carbamates and ureas. *J. Chromatogr.* 20:9-14.
- Franken, J. J., and C. Vidal-Madjar. 1971. Gas-solid chromatographic analysis of aromatic amines, pyridine, picolines, and lutidines on cobalt phthalocyanine with porous-layer open-tube columns. *Anal. Chem.* 43:2034-37.
- Frei, R. W., and V. Mallet. 1971. Quantitative thin-layer chromatography of organothiophosphorus pesticides by *in situ* fluorimetry. *Int. J. Environ. Anal. Chem.* 1:99-111.
- Freudenthal, R. I., A. P. Leber, D. Emmerling, and P. Clarke. 1975. The use of high pressure liquid chromatography to study chemically induced alterations in the pattern of benzo(a)pyrene metabolism. *Chem.-Biol. Interact.* 11:449-58.
- Frissel, M. J., P. Poelstra, and R. Reiniger. 1970. Chromatographic transport through soils. III: A simulation model for the evaluation of the apparent diffusion coefficient in undisturbed soils with tritiated water. *Plant Soil* 33:161-76.
- Gasparick J., Z. Deyl, M. Lederer, K. Macek, and J. Janak, eds. 1977. Chromatographic data: Supplement to Chromatographic reviews, 1977. *J. Chromatogr.* 141:D1-D64.
- Getz, M. E. 1971. Instrumentation for quantitative thin layer chromatography. In *Pesticide Chemistry: Proceedings of the International IUPAC Congress Pesticide Chemistry 2d Tel Aviv, Israel February 22-26, 1971*, pp. 43-63.
- Giam, C. S., S. D. Abbott, and W. B. Davis. 1969. Sorbate-sorbent interactions of nitrogen heterocycles in gas chromatography pyridine derivatives. *J. Chromatogr.* 42:457-63.

- Giger, W., and M. Blumer. 1974. Polycyclic aromatic hydrocarbons in the environment: Isolation and characterization by chromatography, visible ultraviolet, and mass spectrometry. *Anal. Chem.* 46:1663-71.
- Gold, A. 1975. Carbon black adsorbates: Separation and identification of a carcinogen and some oxygenated polyaromatics. *Anal. Chem.* 47:1469-72.
- Goldberg, M. C., L. Delong, and M. Sinclair. 1973. Extraction and concentration of organic solutes from water. *Anal. Chem.* 45:89-93.
- Gomez-Taylor, M., D. Kuehl, and P. R. Griffiths. 1976. Vibrational spectrometry of pesticides and related materials on thin-layer chromatography adsorbents. *Appl. Spectrosc.* 30:447-52.
- Grant, D. W., and R. B. Meiris. 1977. Application of thin-layer and high-performance liquid chromatography to the separation of polycyclic aromatic hydrocarbons in bituminous materials. *J. Chromatogr.* 142:339-51.
- Gregg, J. S. 1975. A simple method for comparing the shapes of closely related isotherms. *J. Chem. Soc., Chem. Commun.* 16:699-700.
- Groves, J. K., H. J. Anderson, and H. Nagy. 1971. Pyrrole chemistry, Part XIII: New synthesis of 3-alkylpyrroles. *Can. J. Chem.* 49:2427-32.
- Gutenmann, W. H., and D. J. Lisk. 1965. Gas chromatographic determination of phenolic pesticides and residues. *J. Offic. Agric. Chem.* 48:1173.
- Hamaker, J. W., C. A. I. Goring, and C. R. Youngson. 1966. Sorption and leaching of 4-amino-3,5,6-trichloropicolinic acid in soils. In *Organic Pesticides in the Environment*, ed. R. F. Gould, pp. 23-37. Advances in Chemistry Series 60. Washington, D. C.: American Chemical Society.
- Hance, R. J. 1967. Relationship between partition data and the adsorption of some herbicides by soils. *Nature* 214:630-31.
- Hansch, C. 1969. A quantitative approach to biochemical structure-activity relationships. *Acc. Chem. Res.* 2:232-39.
- Hansen, S. A. 1976. Thin-layer chromatographic method for the identification of organic acids. *J. Chromatogr.* 124:123-26.
- Haque, R., and D. Schmedding. 1975. A method of measuring the water solubility of hydrophobic chemicals: Solubility of five polychlorinated biphenyls. *Bull. Environ. Contam. Toxicol.* 14:13-18.
- Harris, C. I. 1966. Adsorption, movement, and phytotoxicity of monuron and s-triazine herbicides in soil. *Weeds* 14:6-10.

- Hartung, G. K., and D. M. Jewell. 1962. Carbazoles, phenazines and dibenzofuran in petroleum products, methods of isolation, separation and determination. *Anal. Chim. Acta* 26:514-28.
- Hawthorne, A. R., and R. B. Gammage. 1977. *Assessment of Dosimetry Requirements and Techniques for Measuring Polycyclic Aromatic Hydrocarbons*. NTIS No. CONF 770301-4. Oak Ridge, Tenn.: Health Physics Division, Oak Ridge National Laboratory.
- Helling, C. S. 1971. Pesticide mobility in soils. I: Parameters of soil thin-layer chromatography. *Soil Sci. Soc. Am. Proc.* 35:732-37.
- Helling, C. S. 1971. Pesticide mobility in soils. II: Applications of soil thin-layer chromatography. *Soil Sci. Soc. Am. Proc.* 35:737-43.
- Helling, C. S. 1971. Pesticide mobility in soils. III: Influence of soil properties. *Soil Sci. Soc. Am. Proc.* 35:743-48.
- Helling, C. S., D. D. Kaufman, and C. T. Dieter. 1971. Algae bioassay detection of pesticide mobility in soils. *Weed Sci.* 19:685-90.
- Helling, C. S., and B. C. Turner. 1968. Pesticide mobility: Determination by soil thin-layer chromatography. *Science* 162:562-63.
- Hermann, T. S. 1974. *Development of Sampling Procedures for Polycyclic Organic Matter and Polychlorinated Biphenyls*. NTIS No. PB-243 362. Washington, D. C.: Office of Research and Development, U. S. Environmental Protection Agency.
- Holik, M., J. Janak, and M. Ferles. 1967. Studies in the pyridine series. XXIV: Correlation of the structure of several N,C-dimethyl-tetrahydropyridines and their gas chromatographic retention data. *Collect. Czech. Chem. Commun.* 32:3546-52.
- Huibregtse, K. R., and J. H. Moser. 1976. *Handbook for Sampling and Sample Preservation of Water and Wastewater*. NTIS No. PB-259 946. Cincinnati, Ohio: Office of Research and Development, United States Environmental Protection Agency.
- Hurtubise, R. J. 1973. Instrumentation for thin-layer chromatography. *J. Chromatogr. Sci.* 11:476-91.
- Huss, M., and V. M. Adamovic. 1973. Determination of ametrine and atrazine residues in soil by thin-layer chromatography. *J. Chromatogr.* 80:137-39.
- Ingle, P. H. B., H. Y. Koh, and R. H. Perrett. 1973. Gas-liquid chromatographic analysis of aqueous solutions of hydroxypyridines and hydroxyquinolines. *J. Chromatogr.* 81:79-83.
- Iassaq, H. J., and E. W. Barr. 1977. Recent developments in thin-layer chromatography. *Anal. Chem.* 49:83A-84A *passim*.

- IUPAC Applied Chemistry Division. 1974. Analytical methods for use in occupational hygiene: Determination of benzo(a)pyrene and benzo(k)fluranthene in airborne particulates (chromatography and optical fluorescence). *Pure Appl. Chem.* 40:36.1-36.7.
- Janini, G. M., K. Johnston, and W. L. Zielinski, Jr. 1975. Use of a nematic liquid crystal for gas-liquid chromatographic separation of polyaromatic hydrocarbons. *Anal. Chem.* 47:670-74.
- Janini, G. M., G. M., Muschik, J. A. Schroer, and W. L. Zielinski, Jr. 1976. Gas-liquid chromatographic evaluation and gas-chromatography/mass spectrometric application of new high temperature liquid crystal stationary phases for polycyclic aromatic hydrocarbon separation. *Anal. Chem.* 48:1879-83.
- Janini, G. M., G. M. Muschik, and W. L. Zielinski, Jr. 1976. N,N'-bis (p-butoxybenzylidene)- α,α' -bi-p-toluidine: Thermally stable liquid crystal for unique gas-liquid chromatography separation of polycyclic aromatic hydrocarbons. *Anal. Chem.* 48:809-13.
- Janini, G. M., B. Shaikh, and W. L. Zielinski, Jr. 1977. Gas-liquid chromatographic analysis of benzo(a)pyrene in cigarette smoke on a nematic liquid crystal. *J. Chromatogr.* 132:136-39.
- Jenkins, R. L., and R. B. Baird. 1975. The determination of benzidine in wastewaters. *Bull. Environ. Contam. Toxicol.* 13:436-42.
- John, E. D., and G. Nickless. 1977. Gas chromatographic method for the analysis of major polynuclear aromatics in particulate matter. *J. Chromatogr.* 138:399-412.
- John, P. T., and R. K. Aggarwal. 1975. An adsorption isotherm for determining monolayer capacity at any relative pressure and mean pore size. *Indian J. Technol.* 13:556-60.
- Johnson, J. H., E. E. Sturino, and S. Bourne. 1977. *An Automated Gas Chromatographic System for Pesticide Residue Analysis*. NTIS No. 905 14-77-001. Chicago, Ill.: Region V, United States Environmental Protection Agency.
- Jones, G. R. N. 1973. Detection of primary arylamines on thin-layer chromatograms by diazotisation and coupling: Comparison of a new reagent with existing methods. *J. Chromatogr.* 77:357-67.
- Jones, L. A., and R. S. Foote. 1975. Cannabis smoke condensate: Identification of some acids, gases, and phenols. *J. Agric. Food Chem.* 22:1129-31.
- Junk, G. A., J. J. Richard, M. D. Grieser, D. Witiak, J. L. Witiak, M. D. Arguello, R. Vick, H. J. Svec, J. S. Fritz, and G. V. Calder. 1974. Use of macroreticular resins in the analysis of water for trace organic contaminants. *J. Chromatogr.* 99:745-62.

- Jupille, T. H. 1977. Programmed multiple development: High performance thin-layer chromatography. *J. Am. Oil Chem. Soc.* 54:179-82.
- Karger, B. L., M. Martin, J. Loheac, and G. Guiochon. 1973. Separation of polyaromatic hydrocarbons by liquid-solid chromatography using 2,4,7-trinitrofluorenone impregnated corasil 1 columns. *Anal. Chem.* 45:496-99.
- Kasamatsu, K. 1975. Isolation of alkyl naphthalenes from the middle distillate of petroleum by liquid-solid and liquid-liquid chromatography. *Bull. Jpn. Pet. Inst.* 17:21-27.
- Katrolia, S. P., R. K. Mehrotra, and S. Ramaniyam. 1974. Thin layer chromatography of pesticides and their residues. *Def. Sci. J.* 24:113-19.
- Kawale, G. B., and V. D. Joglekar. 1976. Tollen's reagent for the detection of carbamate and organophosphate insecticides. *Curr. Sci.* 45:57-58.
- Keith, L. H., A. W. Garrison, F. R. Allen, M. H. Carter, T. L. Floyd, J. D. Pope, and A. D. Thruston, Jr. 1976. Identification of organic compounds in drinking water from thirteen U. S. cities. In *Identification and Analysis of Organic Pollutants in Water*, ed. L. H. Keith Ann Arbor, Mich.: Ann Arbor Science Publishers, Inc.
- Kelly, J. A. 1967. *The Determination of Phenolic-type Compounds in Water by High-pressure Liquid Chromatography*. NTIS No. ORD-4254-15. Ph.D. dissertation, Oklahoma State University.
- Khan, S. U. 1975. Chemical derivatization of herbicide residues for gas liquid chromatographic analysis. *Res. Rev.* 52:21-50.
- King, P. H., and P. L. McCarty. 1968. A chromatographic model for predicting pesticide migration in soils. *Soil Sci.* 106:248-61.
- Kingston, D. G. I., and B. T. Li. 1975. Preliminary investigation of the use of high-pressure liquid chromatography for the separation of indole alkaloids. *J. Chromatogr.* 104:431-37.
- Kirchner, J. G. 1974. Modern techniques in TLC. *J. Chromatogr. Sci.* 13:558-63.
- Kirchner, J. G. 1974. Thin-layer chromatography--yesterday, today, and tomorrow. *Chem. Technol.* 4:79-82.
- Kovac, J., and M. Henselova. 1977. Detection of triazine herbicides in soil by a Hill-reaction inhibition technique after thin-layer chromatography. *J. Chromatogr.* 133:420-22.

- Kubota, H., W. H. Griest, and M. R. Guerin. 1975. *Determination of Carcinogens in Tobacco Smoke and Coal-derived Samples--Trace Polynuclear Aromatic Hydrocarbons*. NTIS No. CONF-750603-3. Oak Ridge, Tenn.: Analytical Chemistry Division, Oak Ridge National Laboratory.
- Kuehl, D. W. 1977. Identification of trace contaminants in environmental samples by selected ion summation analysis of gas chromatographic-mass spectral data. *Anal. Chem.* 49:521.
- Kuo, P. P. K., E. S. K. Chian, J. H. Kim, and F. B. DeWalle. 1977. Study of the gas stripping, sorption, and thermal desorption procedures for preconcentrating volatile polar organics from water samples for analysis by gas chromatography. *Anal. Chem.* 49:1023-29.
- Kuratsune, M., and T. Hirohata. 1962. Decomposition of polycyclic aromatic hydrocarbons under laboratory illuminations. In *National Cancer Institute Monograph no. 9*, pp. 117-125.
- LaFleur, P. D., ed. 1976. *Accuracy in Trace Analysis: Sampling, Sample Handling, Analysis*, volume I. NTIS No. PB 258 091. Washington, D.C.: National Bureau of Standards, Department of Commerce.
- LaFleur, P. D., ed. 1976. *Accuracy in Trace Analysis: Sampling, Sample Handling, Analysis*, volume II. NTIS No. PB 258 092. Washington, D.C.: National Bureau of Standards, Department of Commerce.
- Lane, D. A., H. K. Moe, and M. Katz. 1973. Analysis of polynuclear aromatic hydrocarbons, some heterocyclics, and aliphatics with a single gas chromatograph column. *Anal. Chem.* 45:1776-78.
- Lao, R. C., R. S. Thomas, and J. L. Monkman. 1975. Computerized gas chromatographic-mass spectrometric analysis of polycyclic aromatic hydrocarbons in environmental samples. *J. Chromatogr.* 112:681-700.
- Lao, R. C., R. S. Thomas, H. Oja, and L. Dubois. 1973. Application of a gas chromatograph-mass spectrometer-data processor combination to the analysis of the polycyclic aromatic hydrocarbon content of airborne pollutants. *Anal. Chem.* 45:908-15.
- Leoni, V., G. Puccetti, and A. Grella. 1975. Preliminary results on the use of Tenax for the extraction of pesticides and polynuclear aromatic hydrocarbons from surface and drinking waters for analytical purposes. *J. Chromatogr.* 106:119-24.
- Lepri, L., P. G. Desideri, and V. Coas. 1974. Chromatographic and electrophoretic behavior of primary aromatic amines on anion-exchange thin layers. *J. Chromatogr.* 90:331-39.
- Lewis, D. L., and D. F. Paris. 1974. Direct determination of carbaryl by gas-liquid chromatography using electron capture detection. *J. Agric. Food Chem.* 22:148-49.

- Li, C. Y., K. C. Lu, J. M. Trappe, and W. B. Bollen. 1970. Separation of phenolic compounds in alkali hydrolysates of a forest soil by thin-layer chromatography. *Can. J. Soil Sci.* 50:458-60.
- Liebman, S. A., D. H. Ahlstron, T. C. Creighton, G. D. Pruder, R. Averitt, and E. J. Levy. 1972. On-line elemental analysis of gas-chromatographic effluents. *Anal. Chem.* 44:1411-15.
- Litman, G. W., R. T. Litman, and C. J. Henry. 1976. Analysis of lipophilic carcinogen-membrane interactions using a model human erythrocyte membrane system. *Cancer Res.* 36:438-44.
- Loft, P. S., and R. J. Hurtubise. 1971. Instrumentation for thin-layer chromatography. *J. Chem. Educ.* 48:A437-A444.
- Lopez, M. C. 1976. *High Pressure Liquid Chromatography and Its Application to the Separation of Polynuclear Aromatic Hydrocarbons Found in Atmospheric Dust and the Residue of Combustion.* NTIS No. CEA R 4678. Grenoble, France: Center for Nuclear Studies.
- Luedecke, E. 1976. *Analysis of Benzo(a)pyrene in Airborne Particulates by Gas Chromatography.* NTIS No. N76-18247. Washington, D. C.: National Aeronautics and Space Administration.
- McCall, J. M. 1975. Liquid-liquid partition coefficients by high-pressure liquid chromatography. *J. Med. Chem.* 18:549.
- McCarty, P. L., and P. H. King. 1966. The movement of pesticides in soils. *Purdue Univ. Eng. Bull.*, Ext. Ser. no. 121:156-71.
- Macek, K., J. Janak, and Z. Deyl. 1977. Thin-layer chromatography: 1. Reviews & books (bibliography). *J. Chromatogr.* 133:B49-B65.
- McFarren, E., F. Raymond, J. Lishka, and H. Parker. 1970. Criterion for judging acceptability of analytical methods. *Anal. Chem.* 42:358-65.
- McGinnes, P. R., and V. L. Snoeyink. 1974. *Determination of the Fate of Polynuclear Aromatic Hydrocarbons in Natural Water Systems.* NTIS No. PB 232 168. Urbana, Ill.: Water Resources Center, University of Illinois at Urbana-Champaign.
- McGuire, J. M., A. L. Alford, and M. H. Carter. 1973. *Organic Pollutant Identification Utilizing Mass Spectrometry.* Report No. EPA-R2-73-234. Corvallis, Oreg.: National Environmental Research Center, Office of Research and Monitoring, U. S. Environmental Protection Agency.
- MacNeil, J. D., and R. W. Frei. 1975. Quantitative thin-layer chromatography of pesticides. *J. Chromatogr. Sci.* 13:279-85.
- Magallona, E. D. 1975. Gas chromatographic determination of residues of insecticidal carbamates. *Res. Rev.* 56:1-77.

- Mahel'ová, H., H. Sackmauerova, A. Szokolay, and J. Kovac. 1974. Determination of BHC isomers in soils by gas-liquid and thin-layer chromatography after extraction with light-petroleum. *J. Chromatogr.* 89:177-83.
- Maier, R., and H. K. Mangold. 1964. Thin-layer chromatography. *Advan. Anal. Chem. and Instrum.* 3:369-477.
- Maier-Bode, H., and M. Riedmann. 1975. Gas chromatographic determination of nitrogen-containing pesticides using the nitrogen flame ionization detector. *Res. Rev.* 54:113-81.
- Maini, P. 1976. Electron-capture gas chromatographic determination of residues of anthraquinone bird repellent. *J. Chromatogr.* 128:174-77.
- Malaspina, L., G. Bardi, and R. Gigli. 1974. Simultaneous determination by Knudsen-effusion microcalorimetric technique of the vapor pressure and enthalpy of vaporization of pyrene and 1,3,5-triphenylbenzene. *J. Chem. Thermodyn.* 6:1053-64.
- Malcolm, R. L., E. M. Thurman, and G. R. Aiken. 1977. The concentration and fractionation of trace organic solutes from natural and polluted waters using XAD-8, methylmethacrylate resin. In *Proceedings of the 11th Annual Conference on Trace Substances in Environmental Health*. Columbia, Mo.: University of Missouri (in press).
- Mallet, V. N., P. E. Belliveau, and R. W. Frei. 1975. *In situ* fluorescence spectroscopy of pesticides and other organic pollutants. *Res. Rev.* 59:51-90.
- Mangold, H. K., R. Kammereck, and D. C. Malins. 1962. Thin-layer chromatography as an analytical and preparative tool in lipid radiochemistry. *Microchem. J. Symposium Series* 2:697-714.
- Martin, A. J. P., and R. L. M. Synge. 1941. A new form of chromatography employing two liquid phases. I: A theory of chromatography. II: Application to the micro-determination of the higher monoaminoacids in proteins. *Biochem. J.* 35:1358-68.
- Mattson, A. M., R. A. Kahrs, and R. T. Murphy. 1970. Quantitative determination of triazine herbicides in soils by chemical analysis. *Res. Rev.* 32:371-99.
- Metcalf, R. L. 1974. Screening compounds for early warnings about environmental pollution. In *Proceedings of the 8th Annual Trace Substances in Environmental Health Conference*, pp. 213-17. Columbia, Mo.: University of Missouri.
- Morales, R., S. M. Rappaport, R. W. Weeks, Jr., E. E. Campbell, and H. J. Ettinger. 1976. *Development of Sampling and Analytical Methods for Carcinogens*. Progress report from 1 January 1975 - 30 June 1975. NTIS No. LA-6160-PR. Los Alamos, N. Mex.: Los Alamos Scientific Laboratory

- Morris, K. M., and R. J. Moon. 1974. Microanalysis of tryptophan metabolites in mice. *Anal. Biochem.* 61:313-27.
- Mosnaim, A. D., M. Wolf, I. Saavedra, A. Rosenkranz, S. Diaz, and D. C. Nonhebel. 1973. Gas-liquid chromatography and thin-layer chromatography of some 3- and 3,x-substituted pyrenes. *J. Chromatogr.* 80:259-62.
- Mukherjee, G., T. V. Mathew, A. K. Mukherjee, and S. N. Mitra. 1971. Identification and separation of chlorinated pesticides by TLC on magnesium hydroxide. *J. Food Sci. Technol.* 8:152-53.
- Neumann, M. G. 1976. The calculation of adsorption isotherms from chromatographic peak shapes. *J. Chem. Educ.* 53:708-10.
- O'Hara, J. R., M. S. Chin, B. Dainius, and J. H. Kilbuck. 1974. Determination of benzo(a)pyrene in smoke condensates by high pressure rapid liquid-liquid chromatography. *J. Food Sci.* 39:38-41.
- O'Reilly, W. F., and R. P. Murrmann. 1974. *Identification of Soil Organics Using a Gas Chromatographic/Mass Spectrometric Method*. Washington, D. C.: Directorate of Military Engineering and Topography, Office, Chief of Engineers, U. S. Army.
- Oscik, J., and G. Chojnacka. 1974. Investigation of the chromatographic process by the correlation between the R_M -values in adsorption and partition chromatography. *Chromatographia* 7:708-12.
- Oscik, J., and G. Chojnacka. 1974. Investigation on the adsorption process in thin-layer chromatography by using two component mobile phases. *J. Chromatogr.* 93:167-76.
- Oswald, E. O., P. W. Albro, and J. D. McKinney. 1974. Utilization of gas-liquid chromatography coupled with chemical ionization and electron impact mass spectrometry for the investigation of potentially hazardous environmental agents and their metabolites. *J. Chromatogr.* 98:363-448.
- Parsons, J. S., and S. Mitzner. 1975. Gas chromatographic method for concentration and analysis of traces of industrial organic pollutants in environmental air and stacks. *Environ. Sci. Technol.* 9:1053-58.
- Pellizzari, E. D. 1975. *Development of Analytical Techniques for Measuring Ambient Atmospheric Carcinogenic Vapors*. NTIS No. PB 250-620. Research Triangle Park, N.C.: Environmental Sciences Research Laboratory, Office of Research and Development, U. S. Environmental Protection Agency.
- Pierce, J. M. 1974. *Mass Spectrometry as Used to Determine Trace Organic Constituents in Aquatic Systems*. Course Syllabus for Chemistry 6050: Analytical Chemistry of Natural Waters. NTIS No. ORO-4254-23.

- Pierce, R. C., and M. Katz. 1975. Determination of atmospheric isomeric polycyclic arenes by thin-layer chromatography and fluorescence spectrophotometry. *Anal. Chem.* 47:1743-48.
- Pionke, H. B., G. Chesters, and D. E. Armstrong. 1968. Extraction of chlorinated hydrocarbon insecticides from soils. *Agron. J.* 60:289-92.
- Popl, M., J. Fährnich, and M. Stejskal. 1976. Adsorption effect in GPC separation of polycyclic aromatic hydrocarbons. *J. Chromatogr. Sci.* 14:537-40.
- Potti, E. E., P. M. Kaimal, and P. G. Nair. 1975. Detection and identification of mixed benzoic dialkyl phosphorodithioic anhydride pesticides by thin layer chromatography. *J. Forensic Sci. Soc.* 15:309-11.
- Poulson, R. E. 1969. Stationary phases for separation of basic and non-basic nitrogen compounds or hydrocarbons by gas-liquid chromatography. *J. Chromatogr. Sci.* 7:152-57.
- Pullan, B. R., R. Howard, and B. J. Perry. 1966. Measuring radionuclide distribution with crossed-wire spark chambers. *Nucleonics* 24:72-75.
- Purkayastha, R. 1971. Simultaneous detection of the residues of atrazine and linuron in water, soil, plant, and animal samples by thin-layer chromatography. *Int. J. Environ. Anal. Chem.* 1:147-58.
- Rai, P. P., and T. D. Turner. 1975. A method for the estimation of anthraquinones using densitometric thin-layer chromatography. *J. Chromatogr.* 104:196-99.
- Rai, P. P., T. D. Turner, and S. A. Matlin. 1975. High pressure liquid chromatography of naturally occurring anthraquinones. *J. Chromatogr.* 110:401-2.
- Ravenhill, J. R., and A. T. James. 1967. A simple sensitive radioactive scanner for thin-layer chromatograms. *J. Chromatogr.* 26:89-100.
- Rhodes, R. C., I. J. Belasco, and H. L. Pease. 1970. Determination of mobility and adsorption of agrichemicals on soils. *J. Agr. Food Chem.* 18:524-28.
- Rummel, W. 1969. On the organic pollution of waters and the methods of their control and purification. *Wiss. Z. Univ. Rostock, Math.-naturwiss. Reihe.* 18:847-58.
- Safar, W., V. Galik, Z. Kafka, and S. Landa. 1973. Chromatographic separation of nitrogen-containing substances in the system gas-liquid, III: Separation of some 1-alkylpiperidines. *Collect. Czech. Chem. Commun.* 38:1655-58.

- Sakodyskii, K. I., L. C. Le, and P. P. Alikhanov. 1973. Separation of isotopically substituted polar compounds derived from hydrocarbons. *J. Chromatogr.* 77:21-27.
- Sawicki, E. 1964. The separation and analysis of polynuclear aromatic hydrocarbons present in the human environment. *Chem. Anal.* 53:24-30.
- Schabron, J. F., R. J. Hurtubise, and H. F. Silver. 1977. Separation of hydroaromatics and polycyclic aromatic hydrocarbons and determination of tetralin and naphthalene in coal derived solvents. *Anal. Chem.* 49:2253-59.
- Schamp, N., and F. van Wassenhove. 1972. Determination of benzo(a)pyrene in bitumen and plants. *J. Chromatogr.* 69:421-25.
- Schulman, S. G., R. M. Threatte, A. C. Capomacchia, and W. L. Paul. 1974. Fluorescence of 6-methoxyquinoline, quinine, and quinidine in aqueous media. *J. Pharm. Sci.* 63:876-80.
- Selkirk, J. K., R. G. Croy, and H. V. Gelboin. 1974. Benzo(a)pyrene metabolites: Efficient and rapid separation by high-pressure liquid chromatography. *Science* 184:169-71.
- Sevcik, J. 1971. Selective detection of sulphur, chlorine, and nitrogen with help of the combination of flame-ionization and flame-photometric detectors. *Chromatographia* 4:195-97.
- Severson, R. F., M. E. Snook, H. C. Higman, O. T. Chortyk, and F. J. Akin. 1976. Isolation, identification, and quantitation of the polynuclear aromatic hydrocarbons in tobacco smoke. In *Carcinogenesis--A Comprehensive Survey* vol. 1, ed. R. Freudenthal and P. W. Jones, pp. 253-70. New York: Raven Press.
- Silhankova, A., M. Holik, and M. Ferles. 1968. Studies in the pyridine series. XXVI: Reduction of some dialkylpyridines and their methiodides with aluminum hydride. *Collect. Czech. Chem. Commun.* 33:2494-2501.
- Singhal, J. P., and R. P. Singh. 1977. Mobility of trace elements in soils by thin layer chromatography, part I. *Colloid Polym. Sci.* 255:488-91.
- Sliwiok, J. and L. Ogierman. 1974. The separation of alkyl aromatic ketones by means of thin-layer chromatography. *J. Chromatogr.* 94:340-41.
- Smart, N. A. 1976. Collaborative studies of methods for pesticides residues analysis. *Res. Rev.* 64:1-16.
- Smith, R. V., J. P. Rosazza, and R. A. Nelson. 1974. Thin-layer chromatographic determination of simple phenols in microbial extracts. *J. Chromatogr.* 95:247-49.

- Snyder, L. R. 1969. Rapid separations by liquid-solid column chromatography. Qualitative and quantitative analysis of hydrogenated quinoline mixture. *J. Chromatogr. Sci.* 7:595-603.
- Snyder, R. 1970. Thin-layer radiochromatography and related procedures. In *Progress in Thin-layer Chromatography and Related Methods*, ed. A. Niederwieser and G. Pataki, vol. 1, pp. 52-73. London: Ann Arbor-Humphrey Science Publishers.
- Soczewinski, E., and J. Kuczynski. 1975. Extraction of organic bases from strongly acidic aqueous solutions. I: Partition systems formed by nonpolar or weakly polar solvent and solutions of sulfuric hydrochloric, hydrobromic or perchloric acid. *Chemia Analityczna* 20:927-41.
- Soczewinski, J., and H. Szumilo. 1973. Investigations on the mechanism and selectivity of chromatography on thin layers of polyamide: Systems of the type cyclohexane + polar solvent-polyamide. *J. Chromatogr.* 81:99-107.
- Soderquist, C. J., and D. G. Crosby. 1972. The gas chromatographic determination of paraquat in water. *Environ. Contam. Toxicol.* 8:363-68.
- Sovocool, G. W., R. G. Lewis, R. L. Harless, N. K. Wilson, and R. D. Zehr. 1977. Analysis of technical chlordane by gas chromatography/mass spectrometry. *Anal. Chem.* 49:734-40.
- Stahl, E., ed. 1965. *Thin-layer Chromatography*. New York: Academic Press.
- Stahl, E., and H. K. Mangold. 1975. Techniques of thin-layer chromatography. In *Chromatography*. 3d ed., ed. E. Heftmann, pp. 164-88. New York: Van Nostrand Reinhold Company.
- Stanford Research Institute. 1976. *NIOSH Analysis Methods for Set F*. NTIS No. NIOSH-SCP-J. Cincinnati, Ohio: National Institute for Occupational Safety and Health.
- Stanford Research Institute. 1976. *NIOSH Analytical Methods for Set L*. NTIS No. 250-159. Cincinnati, Ohio: National Institute for Occupational Safety and Health.
- Stanley, C. W. 1966. Derivatization of pesticide related acids and phenols for gas chromatographic determination. *J. Agric. Food Chem.* 14:321-23.
- Stanley, T. W., D. F. Bender, and W. C. Elbert. 1973. Quantitative aspects of thin-layer chromatography in air pollution measurements. In *Quantitative Thin Layer Chromatography*, ed. J. C. Touchstone, pp. 305-22. New York: Wiley-Interscience.
- Stejskal, J., and P. Kratochvil. 1975. Determination of the coefficients of selective sorption in the system polymer-ternary solvent by the density and refractive index increments methods. *J. Polym. Sci.* 13:715-25.

- Stepanova, M. E., R. E. Elina, and U. K. Schoshnikov. 1972. Determination of polynuclear aromatic hydrocarbons in exhaust from chemical and petrochemical manufacturing. *J. Anal. Chem.* 27:1201-4.
- Stroupe, R. C., P. Tokousbalides, R. B. Dickinson, Jr., E. L. Wehry, and G. Mamantov. 1977. Low temperature fluorescence spectrometric determination of polycyclic aromatic hydrocarbons by matrix isolation. *Anal. Chem.* 49:701-5.
- Suzuki, K., H. Nagayoshi, and T. Kashiwa. 1973. Systematic separation and identification of 13 carbamate pesticides in their mixture. *Agric. Biol. Chem.* 37:2181-84.
- Swain, A. P., J. E. Cooper, and R. L. Stedman. 1969. Large-scale fractionation of cigarette smoke condensate for chemical and biologic investigations. *Cancer Res.* 29:579.
- Szumilo, H., and E. Soczewinski. 1976. Investigations on the mechanism and selectivity of chromatography on thin layers of polyamide. IV: Chromatography of amino acids and complex phenolic substances. *J. Chromatogr.* 124:394-400.
- Takacs, J. M., E. Kocsi, E. Garamvolgyi, E. Eckhart, T. Lombosi, S. Z. Nyiredy, Jr., I. Borbely, and G. Y. Krasznai. 1973. Contribution to the theory of the retention index system. VI: Calculation of the retention indices of compounds containing halogen atoms or hydroxyl groups, amines, ketones, esters, heterocyclic compounds, adamantanes, silanes and steroids on apolar and polar stationary phases in gas-liquid chromatography. *J. Chromatogr.* 81:1-8.
- Terrill, J. B., and E. S. Jacobs. 1970. Application of gas-liquid chromatography to the analysis of anthraquinone dyes and intermediates. *J. Chromatogr. Sci.* 8:604-7.
- Tesarik, K., and S. Ghyczy. 1974. Separation of pyridine bases of coal tar light oil by means of capillary gas chromatography. *J. Chromatogr.* 91:723-31.
- Thielemann, H. 1974. Identification and determination of 1,4-benzoquinone in waste waters from the coal-processing industry. *Z. Wasser-Abwasser-Forschung* 7:91-93. NTIS No. ORNL-tr-2945. Oak Ridge, Tenn.: Oak Ridge National Laboratory.
- Thornburg, W. 1971. Pesticide residues. *Anal. Chem.* 43:145-62.
- Thorstad, O., and K. Undheim. 1974. Mass spectrometry of onium compounds. *Chem. Scr.* 6:222-25.
- Toneby, M. I. 1974. Thin-layer chromatographic fluorimetry of indole derivatives after condensation by a paraform aldehyde spray reagent. *J. Chromatogr.* 97:47-55.

- Treiber, L. R. 1976. An extension of the programmed multiple development (PMD) technique. *J. Chromatogr.* 124:69-72.
- Tsuda, T., H. Yanagihara, and D. Ishii. 1974. Gas-modified solid chromatography using organic vapours as carrier gas. *J. Chromatogr.* 101:95-102.
- U. S. Environmental Protection Agency (U. S. EPA). 1977. *Sampling and Analysis Procedure for Screening of Industrial Effluents for Priority Pollutants*. Cincinnati, Ohio: Environmental Monitoring and Support Laboratory, U. S. EPA.
- Van Der Meeren, A. A. F., and A. L. T. Verhaar. 1968. Gas-liquid chromatographic determination of pyridine bases without tailing effects. *Anal. Chim. Acta* 40:343-46.
- Van Duyne, R. P., and D. A. Aikens. 1968. A chemically selective polarographic detector for gas chromatography. *Anal. Chem.* 40:254-56.
- Voyatzakis, E., G. Vasilikiotis, and H. Alexaki-Tzivanidou. 1972. Separation of some anthracene derivatives by thin layer chromatography. *Anal. Lett.* 5:445-49.
- Waksmudzki, A., M. Jaroniec, S. Sokotowski, and A. Dawidowicz. 1976. Studies of energetic heterogeneity of adsorbents by liquid chromatography. II: Graphical method for investigating mechanism of adsorption from liquids. *Chromatographia* 9:43-47.
- Wasik, S. P., and R. L. Brown. 1976. Analysis of complex aromatic hydrocarbon mixtures with solid silver nitrate columns. *Anal. Chem.* 48:2218-20.
- Webb, R. G., A. W. Garrison, L. H. Keith, and J. M. McGuire. 1973. *Current Practice in GC-MS Analysis of Organics in Water*. NTIS No. PB-224-947. Athens, Ga.: Southeast Environmental Research Laboratory, U. S. Environmental Protection Agency.
- Weeks, R. W., Jr., R. Morales, S. M. Rappaport, H. J. Ettinger, and E. E. Campbell. 1975. *Development of Sampling and Analytical Methods for Carcinogens*. Progress report from 1 July 1974 - 31 December 1974. NTIS No. LA-6042-PR. Energy Research and Development Administration.
- Wessel, J. R. 1968. Collaborative study of three gas chromatographic dual detection systems for analysis of multiple chlorinated and organophosphorus pesticides. *J. Assoc. Off. Anal. Chem.* 51:666.
- Wheatley, G. A., and J. A. Hardman. 1959. The bioassay of residues of insecticides in soil. *Ann. Appl. Biol.* 48:423-27.
- White, D., and C. T. Cowan. 1958. The sorption properties of dimethyldioctadecyl ammonium bentonite using gas chromatography. *Trans. Faraday Soc.* 54:557-61.

- White, R. H., and J. W. Howard. 1967. Thin-layer chromatography of polycyclic aromatic hydrocarbons. *J. Chromatogr.* 29:108-14.
- Willis, R. B. 1973. *High Pressure Liquid Chromatography of Phenols and Metal Ions*. NTIS No. TS-T-601. Ames, Iowa: Iowa State University.
- Woo, C. S., A. P. D'Silva, V. A. Fassel, and G. J. Oestreich. 1978. Polynuclear aromatic hydrocarbons in coal--identification by their x-ray excited optical luminescence. *Environ. Sci. Technol.* 12:173-74
- Yavorovskaya, S. F. 1973. *Gas-chromatography--A Method of Determining Harmful Substances in the Air and in Biological Media*. NTIS No. JPRS-59872. Arlington, Va.: Joint Publications Research Service.
- Young, J. C. 1976. Detection of N-aryl-N-nitrosamines on thin-layer chromatographic plates with 2,4-dinitrophenylhydrazine and phosphomolybdic acid. *J. Chromatogr.* 124:115-19.
- Zielinski, W. L., Jr. 1977. Liquid crystals-anisotropic GLC phases for novel high-temperature separations. *Analabs* Sept. 1977:2-9.
- Zoccolillo, L., and A. Liberti. 1976. Determination of polycyclic hydrocarbons by channel thin-layer chromatography. *J. Chromatogr.* 120:485-88.
- Zoccolillo, L., A. Liberti, and D. Brocco. 1972. Determination of polycyclic hydrocarbons in air by gas chromatography with high efficiency packed columns. In *Atmospheric Environment*, pp. 715-20. New York: Pergamon Press.
- Zweig, G., and J. Sherma. 1976. Paper and thin-layer chromatography. *Anal. Chem.* 48:66R-83R.

COMPOUND CHARACTERISTICS

- Bewick, A. L., and D. Brown. 1977 Effects of ion-pairing and adsorption on the stereochemistry of the cathodic pinacolisation of acetophenone. *J. Chem. Soc., Perkin Trans.* 2:99-102.
- Bisanz, T., and M. Bukowska. 1974. Transmission of substituent effects through the naphthalene system. *Rocz. Chem.* 48:777-86.
- Bloom, T. F., and R. S. Kramkowski. 1974. *Health Hazard Evaluation/Toxicity Determination*. NTIS no. PB-246-473. Rockville, Maryland: National Institute for Occupational Safety and Health.
- Bock, E., R. Wasylishen, and B. E. Gaboury. 1973. Electric dipole moments and conformations of ortho-, meta-, and para-fluoroacetophenones and of ortho-, meta-, and para-trifluoromethylacetophenones. *Can. J. Chem.* 51:1906-9.
- Brian, R. C. 1964. The classification of herbicides and types of toxicity. In *The Physiology and Biochemistry of Herbicides*, ed. L. J. Audus, pp. 1-37. New York: Academic Press.
- Broadbent, A. D., and R. J. Melanson. 1975. The redox behavior of 9,10-anthraquinone-2-sulfonate in acidic aqueous solution. *Can. J. Chem.* 53:3757-60.
- Chiou, C. T., and V. H. Freed. 1977. Partition coefficient and bioaccumulation of selected organic chemicals. *Environ. Sci. Technol.* 11:475-78.
- Christensen, H. E., E. J. Fairchild, B. S. Carroll, and R. J. Lewis, eds. 1976. *Registry of Toxic Effects of Chemical Substances*, 1976 ed. NTIS no. PB-266-295. Rockville, Md.: National Institute for Occupational Safety and Health.
- Conway, B. E., H. P. Ohar, and S. Gottesfeld. 1973. Molecular orientation in adsorption of pyridine and pyrazine at water/mercury and water/air interfaces: electrocapillary and reflectance studies. *J. Colloid Interf. Sci.* 43:303-18.
- Coupek, J., S. Pokorny, and J. Pospisil. 1974. Gel chromatographic behaviour of mononuclear aromatic hydrocarbons and phenols. *J. Chromatogr.* 95:103-12.
- Dewing, J., G. T. Monks, and B. Youll. 1976. Competitive adsorption of pyridine and sterically hindered pyridines on alumina. *J. Catal.* 44:226-35.
- Dorigan, J., B. Fuller, and R. Duffy. 1976. *Preliminary Scoring of Selected Organic Air Pollutants, Appendix I: Chemistry, Production, and Toxicity of Chemicals A through C*. NTIS no. PB-264-443. Research Triangle Park, N.C.: Office of Air Quality Planning and Standards, Strategies and Air Standards Division, U.S. Environmental Protection Agency.

- Dorigan, J., B. Fuller, and R. Duffy. 1976. *Preliminary Scoring of Selected Organic Air Pollutants, Appendix II: Chemistry, Production, and Toxicity of Chemicals D-E*. NTIS no. PB-264-444. Research Triangle Park, N.C.: Office of Air Quality Planning and Standards, Strategies and Air Standards Division, U.S. Environmental Protection Agency.
- Dorigan, J., B. Fuller, and R. Duffy. 1976. *Preliminary Scoring of Selected Organic Air Pollutants, Appendix III: Chemistry, Production, and Toxicity of Chemicals, F-N*. NTIS no. PB-264-445. Research Triangle Park, N.C.: Office of Air Quality Planning and Standards, Strategies and Air Standards Division, U.S. Environmental Protection Agency.
- Dorigan, J., B. Fuller, and R. Duffy. 1976. *Preliminary Scoring of Selected Organic Air Pollutants, Appendix IV: Chemistry, Production, and Toxicity of Chemicals O-Z*. NTIS no. PB-264-446. Research Triangle Park, N.C.: Office of Air Quality Planning and Standards, Strategies and Air Standards Division, U.S. Environmental Protection Agency.
- Fendler, J. H., J. Fendler, G. A. Infante, P. S. Shih, and L. K. Patterson. 1974. Absorption and proton magnetic resonance spectroscopic investigation of the environment of acetophenone and benzophenone in aqueous micellar solutions. *J. Am. Chem. Soc.* 97:89-95.
- Freed, V. H., and R. Haque. 1976. Chemical structure and properties of selected benzene compounds in relation to biological activity. *Environ. Health Perspect.* 13:23-26.
- Fuller, B., J. Hushon, M. Kornreich, R. Ouellette, L. Thomas, and P. Walker. 1976. *Preliminary Scoring of Selected Organic Air Pollutants*. NTIS no. PB-264-442. Research Triangle Park, N.C.: Office of Air Quality Planning and Standards, Strategies and Air Standards Division, U.S. Environmental Protection Agency.
- Gammage, R. B. 1976. *Characterization and Measurement with a View Toward Personnel Protection*. Abstracts of the 1st ORNL workshop on polycyclic aromatic hydrocarbons, NTIS no. ORNL-TM-5598. Oak Ridge, Tenn.: Oak Ridge National Laboratory.
- Ghersetti, S., S. Giorgianni, A. Passerini, and G. Spunta. 1975. Infrared spectra and vibrational features of acetophenones in the range 700-50 cm^{-1} . *Spectrosc. Lett.* 8:391-97.
- Gibson, D. T., V. Mahadevan, D. M. Jerina, H. Yagi, and J. C. Yeh. 1975. Oxidation of the carcinogens benzo(a)pyrene and benzo(a)anthracene to dihydrodiols by a bacterium. *Science* 189:295-97.
- Grant, D. J. W., and T. R. Al-Najjar. 1976. Degradation of quinoline by a soil bacterium. *Microbios.* 15:177-89.
- Grantham, P. H., E. K. Weisburger, and J. H. Weisburger. 1960. Ionization constants of derivatives of fluorene and other polycyclic compounds. *J. Org. Chem.* 26:1008-17.

- Hamann, S. D. and M. Linton. 1974. Influence of pressure on the ionization of substituted phenols. *J. Chem. Soc., Faraday Trans. 1* 12:2239-49.
- Hoshi, T., J. Yoshino, and K. Hayashi. 1973. Electronic spectra of phenolate and naphtholate anions. *Z. Phys. Chem. (Frankfurt am Main)* 83:31-40.
- Keinath, T. M. 1976. *Benzidine: Wastewater Treatment Technology*. NTIS no. PB-254-024. Washington, D.C.: Criteria and Standards Division, Office of Water Planning and Standards, U.S. Environmental Protection Agency.
- Lu, P. Y., and R. L. Metcalf. 1975. Environmental fate and biodegradability of benzene derivatives as studied in a model aquatic ecosystem. *Environ. Health Perspect.* 10:269-84.
- Lumbroso, W., C. Segard, and B. Roques. 1973. The electric dipole moments of various benzene- and thiophene-chromium tricarbonyl compounds. *J. Organomet. Chem.* 61:249-60.
- Malaspina, L., G. Bardi, and R. Gigli. 1974. Simultaneous determination by Knudsen-effusion microcalorimetric technique of the vapor pressure and enthalpy of vaporization of pyrene and 1,3,5-triphenylbenzene. *J. Chem. Thermodyn.* 6:1053-64.
- Marshall, K., and C. H. Rochester. 1975. Infra-red studies of adsorption at the solid liquid interface. *Faraday Discuss. Chem. Soc.* 59:117-26.
- Matkovics, B., Z. S. Fatray, and L. M. Simon. 1975. TLC of substituted pyridines. XII: Hydroxy derivatives. *Microchem. J.* 20:476-82.
- McKenna, E. J., and R. D. Heath. 1976. *Biodegradation of Polynuclear Aromatic Hydrocarbon Pollutants by Soil and Water Microorganisms*. Research Report No. 113. Urbana, Ill.: Water Resources Center, University of Illinois at Urbana-Champaign.
- Moriconi, E. J., B. Rakoczy, and W. F. O'Connor. 1961. Ozonolysis of polycyclic aromatics. VIII: Benzo(a)pyrene. *J. Am. Chem. Soc.* 83:4618-23.
- Muller, W. P., and F. Korte. 1975. Microbial degradation of benzo(a)pyrene, monolinuron, and dieldrin in waste composting. *Chemosphere* 3:195-98.
- Oscik, J., and G. Chojnacka. 1973. Correlation of the R_M -values of some heterocyclic bases, naphthols, naphthylamines, and nitroanilines in adsorption and partition chromatography. *Chromatographia* 6:133-38.
- Parker, V. D. 1976. Energetics of electrode reactions. II: The relationship between redox potentials, ionization potentials, electron affinities, and solvation energies of aromatic hydrocarbons. *J. Am. Chem. Soc.* 98:98-103.

- Radding, S. B., D. H. Liu, H. L. Johnson, and T. Mill. 1977. *Review of the Environmental Fate of Selected Chemicals*. NTIS no. PB-267-121. Washington D.C.: Office of Toxic Substances, U.S. Environmental Protection Agency.
- Radding, S. B., T. Mill, C. W. Gould, D. H. Liu, H. L. Johnson, D. C. Bomberger, and C. V. Fojo. 1976. *The Environmental Fate of Selected Polynuclear Aromatic Hydrocarbons*. NTIS no. PB-250-948. Washington D.C.: Office of Toxic Substances, U.S. Environmental Protection Agency.
- Rao, P. S., and E. Hayon. 1974. Redox potentials of free radicals. II: Pyrimidine bases. *J. Am. Chem. Soc.* 96:1295-1300.
- Roychowdhury, P., and B. S. Basak. 1975. The crystal structure of indole. *Acta Crystallogr.* B:1559-63.
- Sagardia, F., J. J. Rigau, A. Martinez-Lahoz, F. Fuentes, C. Lopez, and W. Flores. 1975. Degradation of benzothiophene and related compounds by a soil pseudomonas in an oil-aqueous environment. *Appl. Microbiol.* 29:722-25.
- Schulman, S. G. 1973. Correspondence of fluorescing states of naphthols and naphtholate anions and its effect on the calculation of pK_a from spectral shifts. *Spectrosc. Lett.* 6:197-202.
- Schwarz, F. P., and S. F. Wasik. 1976. Fluorescence measurements of benzene, naphthalene, anthracene, pyrene, fluoranthene, and benzo(e)pyrene in water. *Anal. Chem.* 48:524-28.
- Seifert, B. 1977. Stability of benzo(a)pyrene on silica gel plates for high-performance thin-layer chromatography. *J. Chromatogr.* 131:417-21.
- Serratosa, J. M. 1965. Infrared analysis of the orientation of pyridine molecules in clay complexes. *Clays Clay Miner.* 26:385-91.
- Shults, W. D., ed. 1976. *Chemical and Biological Examination of Coal-derived Materials*. NTIS no. ORNL/NSF/EATC 18. Oak Ridge, Tenn.: Oak Ridge National Laboratory.
- Slifkin, M. A., and A. O. Al-Chalabi. 1975. The absorption spectrum of the pyrene excimer in different solvents. *Chem. Phys. Lett.* 3:198-200.
- Suess, M. J. 1972. Aqueous solutions of 3,4-benzpyrene. *Water Res.* 6:981-85
- U.S. Environmental Protection Agency. 1976. *Criteria Document: Benzidine*. NTIS no. PB-254-023. EPA report no. 440/9-76-017. Washington, D.C.: Office of Water Planning and Standards, U.S. Environmental Protection Agency.

- U.S. Environmental Protection Agency. 1976. *Identification of Selected Federal Activities Directed to Chemicals of Near-term Concern*. NTIS no. PB-257-494. Washington, D.C.: Office of Toxic Substances, U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. 1976. *Summary Characterizations of Selected Chemicals of Near-term Interest*. NTIS no. PB-255-817. Washington, D.C.: Office of Toxic Substances, U.S. Environmental Protection Agency.

OCCURRENCE AND DISTRIBUTION OF
ENERGY-RELATED ORGANIC COMPOUNDS

- Andelman, J. B., and J. E. Snodgrass. 1974. Incidence and significance of polynuclear aromatic hydrocarbons in the water environment. In *CRC Critical Reviews in Environmental Control*, January 1974, 69-83.
- Andelman, J. B., and M. J. Suess. 1970. Polynuclear aromatic hydrocarbons in the water environment. *Bull. World Health Org.* 43:479-508.
- Ayer, F. A. 1976. *Symposium Proceedings: Environmental Aspects of Fuel Conversion Technology, II*. NTIS No. PB-257 182. Industrial Environmental Research Laboratory, Office of Research and Development, U. S. Environmental Protection Agency.
- Blumer, M., W. Blumer, and T. Reich. 1977. Polycyclic aromatic hydrocarbons in soils of a mountain valley: Correlation with highway traffic and cancer incidence. *Environ. Sci. Technol.* 11:1082-84.
- Blumer, M., and W. W. Youngblood. 1975. Polycyclic aromatic hydrocarbons in soils and recent sediments. *Science* 188:53-55.
- Borneff, J. 1974. *Polycyclic Aromatics in Surface and Ground Water*. NTIS No. PB-237-786-T. Research Triangle Park, N. C.: U. S. Environmental Protection Agency.
- Cavagnaro, D. M., ed. 1977. *Polychlorinated Biphenyls in the Environment: A Bibliography with Abstracts*. NTIS No. PS-770-792. Springfield, Va.: National Technical Information Service.
- Clemo, G. R. 1973. Some aromatic basic constituents of coal soot. *Tetrahedron* 29:3987-90.
- Clugston, D. M., A. E. George, D. S. Montgomery, G. T. Smiley, and H. Sawatzky. 1976. Sulfur compounds in oils from the western Canada tar belt. In *Shale Oil, Tar Sands and Related Fuel Sources*, ed. R. F. Gould, pp. 11-27. Washington, D. C.: American Chemical Society.
- Coalgate, J. L. 1975. *A Study of Coal Associated Wastes Resulting from the Mining, Processing and Utilization of Coal: Literature Survey--Coal Associated Wastes*. NTIS No. FE-1218 T1. Morgantown, W. Va.: Coal Research Bureau, West Virginia University
- Dvorak, A. J., C. D. Brown, E. H. Dettman, R. A. Hinchman, J. D. Jastrow, F. C. Kornegay, C. R. LaFrance, B. G. Lewis, R. T. Lundy, R. D. Olsen, J. I. Parker, E. D. Pentecost, J. L. Saquinsin, and W. S. Vinikour. 1977. *The Environmental Effects of Using Coal for Generating Electricity*. NTIS No. PB-267-237. Division of Site Safety and Environmental Analysis, Office of Nuclear Reactor Regulation, U. S. Nuclear Regulatory Commission.

- Gammage, R. B. 1977. I: Medical Surveillance. II: Industrial experiences, personnel protection and monitoring. In *Proceedings of the 2d ORNL Workshop on Exposure to Polynuclear Aromatic Hydrocarbons in Coal Conversion Processes*. NTIS NO. CONF-770361. Oak Ridge, Tenn.: Oak Ridge National Laboratory.
- Hangerbrauck, R. P., D. J. Von Lehmden, and J. E. Meeker. 1964. Emissions of polynuclear hydrocarbons and other pollutants from heat-generation and incineration processes. *J. Air Pollut. Control Assoc.* 14:267-78.
- Harrison, R. M., R. Perry, and R. A. Wellings. 1975. Polynuclear aromatic hydrocarbons in raw, potable and waste waters. *Water Res.* 9:331-46.
- Hayatsu, R., R. G. Scott, L. P. Moore, and M. H. Studier. 1975. Aromatic units in coal. *Nature* 257:378-80.
- Neely, W. B., D. R. Branson, and G. E. Blau. 1974. Partition coefficient to measure bioconcentration potential of organic chemicals in fish. *Environ. Sci. Technol.* 8:1113-15.
- Olsen, D., and J. L. Haynes. 1969. *Consumer Protection and Environmental Health Series: Air Pollution Aspects of Organic Carcinogens*. NTIS No. PB-188-090. Bethesda, Md.: Litton Systems, Inc.
- Palmer, H. D., K. T. S. Tzou, and A. Swain. (undated). *Transport of Chlorinated Hydrocarbons in Sediments of the Upper Chesapeake Bay*. NTIS No. PB-255 688. Washington, D. C.: Office of Water Resources Research, U. S. Department of the Interior.
- Ray, S. S., and F. G. Parker. 1977. *Characterization of Ash from Coal-fired Power Plants*. NTIS No. PB-265 374. Research Triangle Park, N. C.: Industrial Environmental Research Laboratory, Office of Research and Development, U. S. Environmental Protection Agency.
- Schiller, J. E. 1977. Nitrogen compounds in coal derived liquids. *Anal. Chem.* 49:2292-94.
- Shabad, L. M., Y. L. Cohan, A. P. Ilnitsky, A. Y. Khesina, N. P. Shcherbak, and G. A. Smirnov. 1971. The carcinogenic hydrocarbon benzo(a)pyrene. *J. Natl. Cancer Inst.* 47:1179-91.
- Shackelford, W. M., and L. H. Keith. 1976. *Frequency of Organic Compounds Identified in Water*. NTIS No. PB-267-470. Athens, Ga.: Environmental Research Laboratory, Office of Research and Development, U. S. Environmental Protection Agency.
- Spangler, C., and N. de Nevers. 1975. *Benzo(a)pyrene and Trace Metals in Charleston, S. C.* NTIS No. PB-243-465. Research Triangle Park, N. C.: Office of Air Quality Planning and Standards, Office of Air and Waste Management, U. S. Environmental Protection Agency.


- TRW Energy Systems. 1976. *Carcinogens Relating to Coal Conversion Processes*. NTIS No. FE-2213-1. Oak Ridge, Tenn.: U. S. Energy Research and Development Administration Technical Information Center.
- Van Meter, W. P., and R. E. Erickson. 1975. *Environmental Effects from Leaching of Coal Conversion By-Products*. Interim report for the period June - September 1975. NTIS No. FE-2019-1. Missoula, Mont.: University of Montana.
- Van Meter, W. P., and R. E. Erickson. 1975. *Environmental Effects from Leaching of Coal Conversion By-Products*. Interim report for the period October - December 1975. NTIS No. FE-2019-2. Missoula, Mont.: University of Montana.
- Van Meter, W. P., and R. E. Erickson. 1976. *Environmental Effects from Leaching of Coal Conversion By-Products*. Interim report for January - March 1976. NTIS No. FE-2019-3. Missoula, Mont.: University of Montana.
- Wedgwood, P., and R. L. Cooper. 1954. The detection and determination of traces of polynuclear hydrocarbons in industrial effluents and sewage. *Analyst* 79:163-69.
- Wewerka, E. N., J. M. Williams, and N. E. Vanderborgh. 1976. *Contaminants in Coals and Coal Residues*. Report No. LA-UR-76-2197. Los Alamos, N. M.: Los Alamos Scientific Laboratory.

APPENDIX: FORMULAS OF ORGANIC COUMPOUNDS


COMPOUNDS OTHER THAN PESTICIDES

Acids, Aliphatic


acetic acid




adipic acid




butyric acid




caproic acid




citraconic



citric acid




dibromosuccinic




formic acid




fumaric acid




glutaric acid



glyceric acid




glycolic acid




glyoxylic acid




isobutyric acid




isovaleric acid




itaconic acid




lactic acid




levulinic acid




maleic acid




malic acid




malonic acid




mesaconic acid



methylsuccinic acid




monobromosuccinic




oxalic




propionic acid




pyruvic acid




succinic acid



tartaric acid




valeric acid




Acids, Aromatic


benzoic acid



phenylacetic acid




phenoxyacetic acid




Alcohol

n-butyl alcohol




Amines and derivatives


acetanilide




aniline



4-(R_2 sulfonyl)-2,6-dinitro-N.N.-
(di- R_1) aniline




crystal violet



Amino Acid


tryptophan



Benzene




Carbamate derivatives




Carbonyl compounds

acetophenone




N-Heterocyclics


benzo[f]quinoline



9H carbazole




7-H dibenzocarbazole



pyridine




quinoline




S-Heterocyclics

benzo[b]thiophene




dibenzothiophene




Phenols

phenol




Polynuclear aromatics


benz[a]anthracene




benz[a]pyrene




pyrene



Quinone




alizarin



Sulfonate


sodium naphthalenesulfonate




PESTICIDES

Carbamates


carbaryl




cycloate




EPTC



pebulate




propham




Carboxylic Acids


chloramben




2,4-D




dicamba




dichlobenil




MCPA



picloram




2,4,5-T




Halogenated Hydrocarbons


BHC




DDT




dieldrin




ethylene dibromide




heptachlor




lindane




methoxychlor



mirex




n-serve




Organophosphates


carbophenothion




crotoxyphos




disulfoton




ethion




fenitrothion




methyl parathion




nellite




paraoxon



parathion




phorate




Amino, Nitrophenyl Sulfones

Nitralin




S-Triazines


ametryne




amitrole




atrazine




cyanazine




ipazine




norazine




prometone




prometryne



fluometuron




propazine




linuron



simazine




methy lurea




Ureas & Uracils

bromacil




metrobromuron




monolinuron




diuron




monuron




fenuron




neburon




phenylurea




tebuthiuron



terbacil




urea




Miscellaneous


chloroneb




chlorthiamid



paraquat



silvex



TECHNICAL REPORT DATA

(Please read Instructions on the reverse before completing)

1. REPORT NO. EPA-600/3-79-086		2.	3. RECIPIENT'S ACCESSION NO.	
4. TITLE AND SUBTITLE Adsorption of Energy-Related Organic Pollutants: A Literature Review			5. REPORT DATE August 1979 issuing date	
			6. PERFORMING ORGANIZATION CODE	
7. AUTHOR(S) K.A. Reinbold, J.J. Hassett, J.C. Means, and W.L. Banwart			8. PERFORMING ORGANIZATION REPORT NO.	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Institute for Environmental Studies University of Illinois at Urbana-Champaign Urbana, Illinois 61801			10. PROGRAM ELEMENT NO. 1BB770	
			11. CONTRACT/GRANT NO. 68-03-2555	
12. SPONSORING AGENCY NAME AND ADDRESS Environmental Research Laboratory--Athens, Ga. Office of Research and Development U.S. Environmental Protection Agency Athens, Georgia 30605			13. TYPE OF REPORT AND PERIOD COVERED Final, 7/77 to 4/78	
			14. SPONSORING AGENCY CODE EPA/600/01	
15. SUPPLEMENTARY NOTES				
16. ABSTRACT <p>This report is a literature review on sorption properties of sediments and energy-related organic pollutants. Adsorption of organic compounds in general is discussed, and analytical methodology in soil thin-layer chromatography and chemical analysis as applicable to measurement of sorption properties is summarized. The literature on the adsorption of energy-related organic pollutants is reviewed. Reported constants for the adsorption of organic compounds on several adsorbents are tabulated, and factors that influence adsorption are discussed.</p>				
17. KEY WORDS AND DOCUMENT ANALYSIS				
a. DESCRIPTORS		b. IDENTIFIERS/OPEN ENDED TERMS		c. COSATI Field/Group
Adsorption Chemical analysis Coal Coal gasification Energy Organic compounds Sediments				68C 68D 99A 99D
18. DISTRIBUTION STATEMENT RELEASE TO PUBLIC		19. SECURITY CLASS (This Report) UNCLASSIFIED		21. NO. OF PAGES 178
		20. SECURITY CLASS (This page) UNCLASSIFIED		22. PRICE