


---

Research and Development

---



# Geological and Hydrochemical Sensitivity of the Eastern United States to Acid Precipitation



## **RESEARCH REPORTING SERIES**

Research reports of the Office of Research and Development, U.S. Environmental Protection Agency, have been grouped into nine series. These nine broad categories were established to facilitate further development and application of environmental technology. Elimination of traditional grouping was consciously planned to foster technology transfer and a maximum interface in related fields. The nine series are:

1. Environmental Health Effects Research
2. Environmental Protection Technology
3. Ecological Research
4. Environmental Monitoring
5. Socioeconomic Environmental Studies
6. Scientific and Technical Assessment Reports (STAR)
7. Interagency Energy-Environment Research and Development
8. "Special" Reports
9. Miscellaneous Reports

This report has been assigned to the ECOLOGICAL RESEARCH series. This series describes research on the effects of pollution on humans, plant and animal species, and materials. Problems are assessed for their long- and short-term influences. Investigations include formation, transport, and pathway studies to determine the fate of pollutants and their effects. This work provides the technical basis for setting standards to minimize undesirable changes in living organisms in the aquatic, terrestrial, and atmospheric environments.

GEOLOGICAL AND HYDROCHEMICAL SENSITIVITY OF THE  
EASTERN UNITED STATES TO ACID PRECIPITATION

by

George R. Hendrey  
Brookhaven National Laboratory  
Upton, New York 11973

James N. Galloway  
University of Virginia  
Charlottesville, VA 22903

Stephen A. Norton  
University of Maine  
Orono, ME 04443

Carl L. Schofield  
Cornell University  
Ithaca, NY 14850

Paul W. Shaffer  
University of Virginia  
Charlottesville, VA 22903

Douglas A. Burns  
University of Virginia  
Charlottesville, VA 22903

Project Officer

Charles F. Powers  
Terrestrial Division  
Corvallis Environmental Research Laboratory  
Corvallis, Oregon 97330

This study was conducted by Brookhaven National Laboratory and Associated Universities, Inc. for the U.S. Environmental Protection Agency in cooperation with the U.S. Department of Energy under Interagency Agreement EPA 79-D-X-0672.

CORVALLIS ENVIRONMENTAL RESEARCH LABORATORY  
OFFICE OF RESEARCH AND DEVELOPMENT  
U.S. ENVIRONMENTAL PROTECTION AGENCY  
CORVALLIS, OREGON 97330

## DISCLAIMER

One of the principal reasons for the preparation of this report for the Environmental Protection Agency was to supply scientifically valid information which could be incorporated into the EPA SO<sub>2</sub>-Particulate Matter criteria document, presently in the final stages of preparation. A strict requirement pertaining to that document is that any scientific information used there must be published (or at least in press) by January 1, 1980. Because of this demanding time constraint, it was necessary that the contractor prepare this report in a shorter time than would ordinarily be attempted, and that it be published by EPA without undergoing peer review. We feel that early publication of these results in order to stimulate the broadest scientific discussion prior to completion of the criteria document justified waiving our normally more rigorous prepublication review requirements. Publication, however, does not signify that the contents necessarily reflect the views and policies of EPA, nor does mention of trade names or commercial products constitute endorsement or recommendation for use.

## FOREWORD

Effective regulatory and enforcement actions by the Environmental Protection Agency would be virtually impossible without sound scientific data on pollutants and their impact on environmental stability and human health. Responsibility for building this data base has been assigned to EPA's Office of Research and Development and its 15 major field installations, one of which is the Corvallis Environmental Research Laboratory.

The primary mission of the Corvallis Laboratory is research on the effects of environmental pollutants on terrestrial, freshwater, and marine ecosystems; the behavior, effects and control of pollutants in lakes and streams; and the development of predictive models on the movement of pollutants in the biosphere.

This report assembled detailed maps of the eastern U.S., showing bedrock geology and surface alkalinity, and examined the relationship of the two as indicators of environmental sensitivity to acid precipitation. Biological impacts of acidification on aquatic systems were also summarized.

Thomas A. Murphy, Director  
Corvallis Environmental Research Laboratory

## ABSTRACT

A new analysis of bedrock geology maps of the eastern U.S. constitutes a simple model for predicting areas which might be impacted by acid precipitation and it allows much greater resolution for detecting sensitivity than has previously been available for the region. Map accuracy has been verified by examining current alkalinities and pH's of waters in several test states, including Maine, New Hampshire, New York, Virginia and North Carolina. In regions predicted to be highly sensitive, alkalinities in upstream sites were generally low,  $<200 \mu\text{eq}/\ell$ . Many areas of the eastern U.S. are pinpointed in which some of the surface waters, especially upstream reaches, may be sensitive to acidification. Pre-1970 data were compared to post-1975 data, revealing marked declines in both alkalinity and pH of sensitive waters of two states tested, North Carolina, where pH and alkalinity have decreased in 80% of 38 streams ( $p < 0.001$ ) and New Hampshire, where pH in 90% of 49 streams and lakes has decreased ( $p < 0.001$ ) since 1949. These sites are predicted to be sensitive by the geological map on the basis of their earlier alkalinity values. Thus this mapping of sensitive areas is validated by the observed temporal trends. The map is to be improved by the addition of a soils component.

This study was conducted by Brookhaven National Laboratory and Associated Universities, Inc. for the U.S. Environmental Protection Agency in cooperation with the U.S. Department of Energy under Interagency Agreement EPA 79-D-X-0672.

## CONTENTS

Foreword. . . . .	iii
Abstract. . . . .	iv
Figures . . . . .	vi
Tables. . . . .	ix
Acknowledgement . . . . .	x
I. Introduction . . . . .	1
II. Preparation of Sensitive Areas Map . . . . .	5
III. Verification of the Sensitive Areas Map. . . . .	27
IV Temporal Trends in pH and Alkalinity: Validation of the Sensitive Areas Map. . . . .	58
V Impacts of Acidification on Aquatic Biota. . . . .	74
VI. Regional Assessment of Acidification Impacts on Fish Population . . . . .	83
Endnote . . . . .	91
References. . . . .	93

## FIGURES

<u>Number</u>	<u>Page</u>
II-1a Unmodified tracing of all boundaries of geologic formations shown on a portion of the North Carolina state geologic map. . . . .	8
II-1b Boundaries of rock formations in North Carolina having comparable acid-neutralizing capacities . . . . .	9
II-1c Smoothing of geologic contacts among areas of differing acid-neutralizing capacities shown in Figure II-1b. . . . .	10
II-2a Unmodified tracing of all boundaries of geologic formations shown on a portion of the New York state geologic map. . . . .	11
II-2b Boundaries of rock formations in New York having comparable acid-neutralizing capacities. . . . .	12
II-2c Smoothing of geologic contacts among areas of differing acid-neutralizing capacities in New York State. . . . .	13
II-3a Unmodified tracing of all boundaries of geologic formations shown on a portion of the Tennessee state geologic map in a region of dendritic drainage in flat-lying rocks and trellis drainage in the valley and ridge provinces . . . . .	14
II-3b Boundaries of rock formations in Tennessee, having comparable acid-neutralizing capacities, in the same region as Figure II-3a. . . . .	15
II-3c Smoothing of geologic contacts among areas of differing acid-neutralizing capacities in Tennessee, shown in Figure II-3b. . . . .	16
II-4 Predicted susceptibility of surface waters in six regions of the eastern United States to acidification as determined by bedrock type. . . . .	17-23
II-5 High altitude lakes (elevation > 600 m) in the Adirondack Mountains of New York which have lost their fish populations . . . . .	24
III-1 Location map of Virginia showing the Shenandoah National Park, the original survey area and the smaller study area. . . . .	26
III-2a Key to bedrock geology in the study area in and adjacent to the Shenandoah National Park . . . . .	30



## Figures (cont.)

<u>Number</u>	<u>Page</u>
III-2b Depiction of bedrock geology in the study area in and adjacent to the Shenandoah National Park . . . . .	31
III-3a Key to pH of streams in the Shenandoah survey area. Shaded areas represent watershed areas upstream from sampling locations . . . . .	34
III-3b Depiction of streams in the Shenandoah survey area . . . . .	35
III-4a Key to alkalinity of streams in the Shenandoah survey area. Shaded areas represent watershed areas upstream from sampling locations . .	36
III-4b Depiction of alkalinity of streams in the Shenandoah survey area .	37
III-5 Changes in water chemistry in White Oak Run during the storm of July 16, 1979. . . . .	39
III-6 Geologically-based sensitive areas map of New Hampshire. . . . .	41
III-7 Geologically-based sensitive areas map of North Carolina . . . . .	42
III-8 Location and alkalinity of New Hampshire sites sampled during the summer of 1979 . . . . .	44
III-9 Location and alkalinity of North Carolina sites sampled during the summer of 1979 . . . . .	45
III-10 Location of New Hampshire sites where pH and alkalinity data was available for the period 1976-1979 . . . . .	47
III-11 Location of North Carolina sites where pH and alkalinity data was available for the period 1976-1979 . . . . .	48
III-12 Minimum alkalinity of counties in New Hampshire for the period 1976-1979 . . . . .	49
III-13 Minimum alkalinity of counties in North Carolina for the period 1976-1979. . . . .	50
III-14 Average alkalinity of counties in New Hampshire for the period 1976-1979 . . . . .	51
III-15 Average alkalinity of counties in North Carolina for the period 1976-1979 . . . . .	52
IV-1 Distribution of alkalinity in samples from North Carolina fisheries data, 1960-1963. . . . .	63
IV-2 Distribution of alkalinity in samples from North Carolina field trip, 1979 . . . . .	64

## Figures (cont.)

<u>Number</u>	<u>Page</u>
IV-3    Distribution of pH in samples from North Carolina fisheries data, 1960-1963. . . . .	65
IV-4    Distribution of pH in samples from North Carolina field trip, 1979 .	66
IV-5    Comparative plot of alkalinity for North Carolina streams, 1960- 1964 versus 1979 . . . . .	67
IV-6    Comparative plot of pH for North Carolina streams, 1960-1964 versus 1979 . . . . .	68
IV-7    Distribution of pH in samples from New Hampshire fisheries data 1937-1939. . . . .	69
IV-8    Distribution of pH in samples from New Hampshire field trip 1979	70
IV-9    Comparative plot of pH for New Hampshire streams, 1937-1939 versus 1979 . . . . .	71
IV-10   Calcium versus pH for headwater streams in North Carolina and New Hampshire plotted over Henriksen's line. . . . .	73
VI-1    Acidification of Adirondack lakes in relation to pH measured in 1975 . . . . .	86
VI-2    Acidification of Adirondack lakes in relation to SO <sub>4</sub> measured in 1975 . . . . .	87
VI-3    Nomograph for predicting lake pH and fish population status from lake Ca levels and either regional acidification or precipitation pH . . . . .	89
VI-4    Changes in the H <sup>+</sup> ion concentrations in 36 Adirondack lakes from the 1930's to 1975 in relation to Ca levels measured in 1975 . . .	90

## TABLES

<u>Number</u>		<u>Page</u>
II-1	States and scales of geologic maps used for preparing the maps of sensitive areas. . . . .	7
III-1	pH and alkalinity of streams in major bedrock formations of the Blue Ridge in Central Virginia. . . . .	28
III-2	Comparison of the geologically based map of sensitive areas (Figure III 7) to county-by-county maps of alkalinity (Figures III-13 and III 15), for 49 of 100 North Carolina counties . . . . .	55
III-3	Comparison of the geologically-based sensitive areas map (Figure III-6) to county-by-county maps of alkalinity (Figures III-12 and III 14) for all 10 counties in New Hampshire . . . . .	55
III-4	Comparison of the geologically-based binary number map of sensitive areas (Figure II-4b) to county-by-county maps of alkalinity (Figures III-13 and III 15), for 49 of 100 North Carolina counties. . . . .	56
III-5	Comparison of the geologically-based binary code sensitive areas map (Figure II-4a) to county-by-county maps of alkalinity (Figures III-13 and III-15) for all 10 counties in New Hampshire. . . . .	56
IV-1	A comparison of old and new data for headwater streams in North Carolina . . . . .	60
IV-2	A comparison of old and new data for New Hampshire streams and lakes. . . . .	61-62
V-1	Damages to aquatic biota likely to occur with increasing acidity .	82
V-2	Summary of damages to aquatic organisms with decreasing pH . .	82
VI-1	Lake classification for Adirondack Mountain lakes on the nomograph of Henriksen . . . . .	91
VI-2	Fish population status in 24 Adirondack Mountain lakes, classified according to the nomograph of Figure VI-3 . . . . .	91

#### ACKNOWLEDGMENTS

The authors wish to thank Avis Newell for technical work in the field and laboratory; Fred Liotta, Susan Hall, Lisa Thurlow, Roland Dupuis and Marilyn Morrison for assistance in map preparation; and Ronnie Evans and Charlie Bores for data base management, data reduction, and computer graphics. We appreciate the generous cooperation and assistance by personnel of the Shenandoah National Park, the North Carolina Department of Inland Fisheries, and the New Hampshire Department of Fish and Game.

## I. INTRODUCTION

In recent decades, man's activities have led to significant changes in atmospheric and precipitation chemistry. Among these activities, metal smelting and combustion of fossil fuels have resulted in atmospheric emissions of sulfur and nitrogen oxides. The effect of these emissions has been a dramatic increase in the acidity of rainfall, particularly in areas downwind of major industrial areas. Although the phenomenon of acidic precipitation has been developing on a hemispheric scale, only recently have workers tried to identify those areas in the United States receiving acidic precipitation<sup>16,71</sup> and associated heavy metals.<sup>67</sup> While a reasonably good understanding of the sources and distribution of acid precipitation has been developed,<sup>16,18,19,27,71,72,77</sup> our understanding of the effects of acidic precipitation on terrestrial and aquatic systems is still fragmentary. Reductions of pH and alkalinity of lakes affected by acidic precipitation have been well documented, particularly in the northeastern United States, southeastern Canada, and Scandinavia. In these same areas, declines in fish populations and adverse effects on other aquatic fauna have been widely observed.<sup>1,23,37,38,41,42,49,57,68,86,91,92,103,104</sup>

Studies of the hydrobiological problems caused by acidic precipitation have been primarily qualitative. Biological aspects of the synoptic lake surveys in Sweden,<sup>1</sup> Norway,<sup>49</sup> Canada,<sup>18</sup> and the United States<sup>90</sup> have concentrated on changes in presence or absence of fish and in the kinds and numbers of species. Very little quantitative information is available concerning the effects of acidification on biomass. This is especially true for the bacteria, and other microorganisms. There is strong evidence indicating that processes such as phytoplankton production and microbial decomposition are inhibited, but quantitative data are scarce. Changes in the availability of inorganic nutrients and in nutrient recycling are hypothesized but have not been demonstrated; effects on food chains or webs are not well known, and no quantitative data are available concerning the effects on energy processing in natural aquatic ecosystems.


The overall objective of the sensitive areas project has been to evaluate the eastern United States and determine and map which areas are vulnerable to adverse impacts from acid precipitation. In addition, a review of the biological consequences of freshwater acidification is presented and a regional assessment of effects on fish, which provides a basis for predicting such impacts, is described.

### A. Geologic Controls


Impact to aquatic ecosystems is largely based on the chemical characteristics of the bedrock. Limestone terrains yield "infinite" acid-neutralizing (buffering) capacity to acidic precipitation whereas granites (and

related igneous rocks), their metamorphic equivalent, and noncalcareous sandstones yield minimal buffering. Several hundred common minerals comprise the huge variety of rocks which comprise the crust of the earth, the surficial overburden, and soils. Of interest in terms of susceptibility of a landscape to acidification from precipitation is the total capacity of the minerals, no matter what their configuration, to assimilate protons ( $H^+$ ). Specific considerations are solubility of the phases and the kinetics of solution.


Certain minerals are soluble and rapidly dissolved but yield no buffering capacity, e.g.:



Others are relatively insoluble and yield no buffering capacity, e.g.






Common rock-forming minerals, when placed in water, yield pH's in excess of 7 (commonly in excess of 8) but the kinetics are sufficiently slow so that the total potential  $H^+$  consuming capacity of these minerals is not realized, e.g.:



$$K_{sp} \sim 10 \sim \left[ \frac{[Na^+]^2}{[H^+]^2} \right] \left[ H_4SiO_4 \right]^4$$


Assuming an initial pH of 4.0 and buffering from the  $CO_2 - H_2O$  system ( $HCO_3^- \sim 10^{-5.7}$ ), the expected pH after complete reaction would be approximately 12. This is never achieved.

Additional  $H^+$  sinks available in soils include Al and Fe hydroxides and silicates, e.g.:




These minerals are either developed in sub-soils (B-horizon) in young soils (and thus not necessarily available to overland flow) or dominate in mature/old soils where they may provide extensive acid neutralizing capacity.


The most important mineral for neutralizing acidic waters is calcite (or nearly any other carbonate mineral). The solution at low and at intermediate pH of this mineral is given by:



Further addition of acid to the resulting (7b) solution ( $2\text{Ca}^{++} + 2\text{HCO}_3^{-}$ ) will result in the protonation of the  $\text{HCO}_3^{-}$ :



These reactions (7 and 8) are rapid and for each mole of  $\text{CaCO}_3$  consumed, 2 moles of  $\text{H}^+$  are consumed before the pH is reduced significantly below 5. Reactions 4, 5, and 6 or similar reactions consume large amounts of  $\text{H}^+$  ions at pH's below 5 and may be the primary acid-consuming mechanism for very acidic solutions. They do not contribute  $\text{HCO}_3^{-}$  for buffering except by virtue of the dissociation of  $\text{H}_2\text{CO}_3$  as the pH is raised by the reaction:



The amount of  $\text{HCO}_3^{-}$  produced will be a function of the total  $\text{CO}_2$  in the system. It should be noted that the gain or loss of molecular  $\text{CO}_2$  to the water, by itself, with no addition of cations to balance  $\text{HCO}_3^{-}$  production, will not change alkalinity although pH does change.

Thus we have a spectrum of response of minerals (and rocks) to the changing acidity of atmospheric precipitation. Accordingly, rocks may be classified by the buffering capacity or acid-neutralizing capacity which they render to surface waters.

Additional controls on acidification of aquatic ecosystems include: hydrologic characteristics of the terrain (i.e., overland flow versus ground-water flow, soil porosity/permeability, residence time of water in the soil, distribution of precipitation through time, type of precipitation, thickness of soil), types of soils [residual, glacial (till, ice-contact stratified, etc.), aeolian, lacustrine, alluvial, etc.], mineralogy and age of soil.

Any prediction about the vulnerability of a terrain to acidification from precipitation based solely on bedrock geology must be tempered with consideration of these other factors. For example, Florida is underlain by highly calcareous and phosphatic rocks suggesting that acidification of lakes and streams is highly unlikely. However, many of the soils (particularly in northern Florida) are very mature, highly leached of  $\text{CaCO}_3$ , and as a result,

acidification of some lakes with minimal deep groundwater influx has occurred (P. Brezonik, pers. comm. 1979). Conversely, there are some areas in Maine underlain by granite and receiving precipitation with an average pH of about 4.3<sup>78</sup> where lakes have not been acidified because of lime-bearing till and marine clay in the drainage basins of the lakes.

#### B. Mapping of Areas Sensitive to Acid Precipitation

Other investigations<sup>33,64,71</sup> have defined areas of the United States and Canada as susceptible to impacts from acidic precipitation, from interpretations of large-scale geologic maps. These areas were largely concentrated in Precambrian terrains (the Canadian shield, the Adirondack Massif), the Precambrian core of the Appalachian Mountains, and in younger igneous rocks of the Northwest and Rocky Mountains.

In the United States surface waters acidified from atmospheric deposition have now been identified from New England and New York to Florida, and in the Boundary Waters Canoe Area, Minnesota. Acidic precipitation exists (with pH < 5.6) in all states contiguous to the Mississippi River and east to the Atlantic Ocean<sup>78</sup> as well as in California<sup>72</sup> and in Washington.<sup>6</sup> Because of this widespread aspect of acidic precipitation, it is important to understand the natural characteristics of the landscape which render an area's aquatic ecosystems susceptible to impact. Three major factors can be identified as important. They are: meteorology, pedology, and geology. This report stresses the geological control of sensitivity.

#### C. Water Quality Data Collection

The utility of the geologically-based maps of sensitive areas is limited by the accuracy with which they identify areas in which surface waters are actually sensitive. We define sensitive waters as those with alkalinity values less than 500  $\mu\text{eq}$  (micro equivalents) of alkalinity per liter. The accuracy of the maps was checked against actual measurements of lake and stream water chemistry.

Data from existing records in many states of the eastern U.S., from the STORET system, and from field measurements conducted in New Hampshire and North Carolina in 1979 are combined in the Acidification Chemistry Information Database (ACID) at Brookhaven National Laboratory. Data is registered by state, county, station code, station name, data source code, latitude, longitude, and sample date. Data sets include pH, alkalinity ( $\mu\text{eq/l}$ ), Ca ( $\mu\text{eq/l}$ ), conductivity (microSiemens), color (mgPt) and fish status. Information on methods used for chemical determinations is also stored in ACID.



## II. PREPARATION OF SENSITIVE AREAS MAP

An analysis of the vulnerability of aquatic ecosystems to acidification from precipitation must start with the bedrock geology. The scale of variability of rock types is such that one must literally look at the geology on a drainage basin basis to make predictions about individual lakes and streams. Igneous rocks normally have maximum dimensions less than 10 km. Folded/faulted metasedimentary/sedimentary rocks may have essentially 1-dimensional distribution. Flat lying sedimentary rocks may be widely distributed but topographic relief commonly intersects many rock types over short distances.

Small amounts of limestone in a drainage basin exert an overwhelming influence on water quality in terrains which otherwise would be very vulnerable to acidification. Consequently, in the regional analysis of vulnerability, areas with intimate mixtures of rocks of varying acid neutralizing capacity have been classified according to the more influential rock types. Analysis of geologic terrain has been undertaken on the scale of the most recent state geologic maps (Table II-1). Rock formations have been classified according to their potential buffering capacity. This judgment was based on map explanations, various stratigraphic lexicons<sup>61,62</sup> and S. Norton's personal knowledge of the geology of areas.

Rock classification was as follows:

- Type I - Low to no buffering capacity (labeled 1 on maps)  
(Widespread impact from acidic precipitation expected.)  
Granite/Syenite or metamorphic equivalent  
Granite gneisses  
Quartz sandstones or metamorphic equivalent
- Type II - Medium/Low buffering capacity (labeled 2 on maps)  
(Impact from acidic precipitation restricted to first and second order streams and small lakes. Complete loss of alkalinity unlikely in large lakes.)  
Sandstones, shales, conglomerates or their metamorphic equivalent (no free carbonate phases present).  
High grade metamorphic felsic to intermediate volcanic rocks.  
Intermediate igneous rocks.  
Calc-silicate gneisses with no free carbonate phases.
- Type III - Medium/High buffering capacity (labeled 3 on maps)  
(Impact from acidic precipitation improbable except for overland run-off effects in areas of frozen ground.)  
Slightly calcareous rocks.  
Low grade intermediate to mafic volcanic rocks.  
Ultramafic rocks.  
Glassy volcanic rocks.

Type IV - "Infinite" buffering capacity (labeled 4 on maps)  
(No impact to aquatic ecosystems.)  
Highly fossiliferous sediments or metamorphic equivalents.  
Limestones or dolostones.

The development of vulnerability maps for 3 types of terrains is illustrated in several steps. Figure II-1a represents, unmodified, a tracing of all boundaries of geologic formations shown on part of the North Carolina state geologic map. This area is underlain by Precambrian and possibly lower Paleozoic rocks which are highly folded, faulted and metamorphosed in the core of the Appalachian Mountains. Figure II-1b differs from II-1a in that boundaries between contiguous formations with comparable acid-neutralizing capacities are not shown. Figure II-1c is modified from II-1b by the smoothing of contacts, favoring rocks with higher buffering capacity, and the deletion of small areas or linear outcroppings of low buffer capacity rocks.

Figure II-2a represents, unmodified, a tracing of all boundaries of geologic formations shown on part of the New York state geologic map. This Precambrian terrain is characterized by high grade metamorphic rocks intruded by many igneous rocks, largely granitic. Figures II-2b and II-2c are developed by the same techniques as Figures II-1b and II-1c.

Figures II-3a, b and c represent similar development for a portion of the state of Tennessee in an area of dendritic drainage in flat-lying sedimentary rocks (western half) and trellis drainage in valley and ridge topography in the highly folded Paleozoic section (eastern half).

The state geologic maps already represent a smoothing and generalization of geologic boundaries. Most state maps utilized in this synthesis have been drawn at a scale of 1:250,000 or 1:500,000 from maps at a scale of 1:24,000 or 1:62,500 with the loss of much small detail. At a scale of 1:250,000 the largest geologic unit that is commonly deleted is approximately 0.2 miles in diameter or width.

"Zero"-dimensional rocks of this size (generally intrusions or erosional remnants of flat lying strata) will not normally affect the chemistry of a large number of streams or lakes. However, one-dimensional rock map units (generally the intersection with the land surface of moderately to steeply dipping strata or dikes or metamorphic equivalents) commonly affect large numbers of primary and secondary streams and lakes which lie in topographic lows occupied by certain strata. Thus, thin limestones in the Valley and Ridge topographic province (Figure II-3a) may not be shown on a scale of 1:500,000 (and thus not on these maps) but may dominate the area's water chemistry and vulnerability to acidic precipitation.

The geologically based sensitivity maps may have also been organized on a county basis to make these data compatible with existing data bases for soils, agriculture, forestry, and other land uses. (A new soils map, being prepared by a separate project<sup>74</sup> will also be organized on a county basis so analyses of county susceptibility to acidification of surface waters may be undertaken.) Each county in the study area states (Table II-1) was analyzed by planimetry for percentage of area underlain by rock types I and II. These data are shown for the eastern United States on Figures II-4. Although individual drainage

Table II-1. States and scales of geologic maps used for preparing the maps of sensitive areas.

Alabama	1:500,000	Mississippi	1:500,000
Arkansas	1:500,000	Missouri	1:500,000
California	1:750,000	New Hampshire	1:250,000
Connecticut	1:500,000	New Jersey	1:250,000
Delaware	1:300,000	New York	1:250,000
Florida	1:3,125,000	North Carolina	1:500,000
Georgia	1:500,000	Ohio	1:500,000
Illinois	1:500,000	Oregon	1:500,000
Indiana	1:1,000,000	Pennsylvania	1:250,000
Iowa	1:500,000	Rhode Island	1:250,000
Kentucky	1:500,000	South Carolina	1:250,000
Louisiana	1:1,250,000	Tennessee	1:250,000
Maine	1:500,000	Vermont	1:250,000
Maryland	1:250,000	Virginia	1:500,000
Massachusetts	1:250,000	Washington	1:500,000
Michigan	1:2,500,000	West Virginia	1:250,000
Minnesota	1:1,000,000	Wisconsin	1:1,000,000




Figure II-1a. Unmodified tracing of all boundaries of geologic formations shown on a portion of the North Carolina state geologic map. Numbers refer to buffering classification. Dashed lines are county boundaries.





Figure II-1b. Boundaries of rock formations in North Carolina having comparable acid-neutralizing capacities. Numbers refer to buffering classifications. Dashed lines are county boundaries.






Figure II-2a. Unmodified tracing of all boundaries of geologic formations shown on a portion of the New York state geologic map. Dashed lines are county boundaries.




Figure II-2b. Boundaries of rock formations in New York having comparable acid-neutralizing capacities. Dashed lines are county boundaries.






Figure II-2c. Smoothing of geologic contacts among areas of differing acid-neutralizing capacities in New York State. Areas of the Adirondack Mountains, shown in Figure II-2b, are located in the center of the northeastern third of the state. Numbers refer to buffering classification. Dashed lines are county boundaries.



Figure II-3a. Unmodified tracing of all boundaries of geologic formations shown on a portion of the Tennessee state geologic map in a region of dendritic drainage in flat-lying rocks and trellis drainage in the valley and ridge provinces. Dashed lines are county boundaries.




Figure II-3b. Boundaries of rock formations in Tennessee, having comparable acid-neutralizing capacities, in the same region as Figure II-3a. Numbers refer to buffering classification. Dashed lines are county boundaries.




Figure II-3c. Smoothing of geologic contacts among areas of differing acid-neutralizing capacities in Tennessee, shown in Figure II-3b. Numbers refer to buffering classification. Dashed lines are county boundaries.

Figure II-4. Predicted susceptibility of surface waters in six regions of the eastern United States to acidification as determined by bedrock type. Data are organized on a county-by-county basis. The first character of the 2-digit code represents the decade-percent of Type-I rock (0 = 0-9%, 1 = 10-19%, 2 = 20-29%, etc.) and the second character represents the decade-percent of rock type II. The states included are (a) Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, and New York; (b) Pennsylvania, New Jersey, Delaware, Maryland, West Virginia, Virginia, and North Carolina; (c) Minnesota (excluding western-most counties), Wisconsin, Michigan and Iowa; (d) Missouri, Illinois, Indiana, Ohio, and Kentucky; (e) Arkansas, Mississippi, Alabama, and Tennessee; and (f) South Carolina, Georgia and Florida.




Figure II-4a





Figure II-4b.








Figure II-4d




Figure II-4e




Figure II-4f.




Figure II-5. High altitude lakes (elevation > 600 m) in the Adirondack Mountains of New York which have lost their fish populations (A) are shown superimposed on a map of rock-types I and II. Lakes which still have fish are indicated (P).

systems or lakes cannot be analyzed in this fashion, regions of high vulnerability can be identified. The first digit of the binary number for each county represents the percentage (0-9% = 0, 10-19% = 1, etc.) of rock type I in the county. The second digit represents a similar decade value for rock type II. For example the number 23 indicates that 20-29% and 30-39% of the county is underlain by rock types I and II, respectively. Rock types III and IV are not represented in these maps.

The predictive accuracy of the maps can be established by comparing the geology-based maps against surface water chemical data in areas receiving acidic precipitation as discussed in Section III of this report or by looking at the biological response of aquatic ecosystems to acidic precipitation. A number of lakes in the Adirondack Mountains have been examined in terms of lake pH and fish populations,<sup>92</sup> as discussed in Section VI of this report, and a close correlation between the two was found. Figure II-5 depicts those lakes which have lost their fish populations and the rock type underlying the area. A geographic correlation exists between fish absence and the existence of rock types I and II in this region. Topography and orographic precipitation are factors contributing to the distribution of lakes in which fish are present or absent.

Correlations also exist between acidification of surface waters and bedrock geology (Section III) in a number of states and areas which have thin soils, including New England.<sup>22</sup> This suggests that bedrock geology exerts a very strong influence on the regional extent of acidification of aquatic ecosystems in response to atmospheric loading of acid. Regionally, soils and vegetative types are of secondary importance. However, locally, soils may overwhelm bedrock influences.




Figure III-1. Location map of Virginia showing the Shenandoah National Park, the original survey area (large, lightly stippled area), and the smaller study area (darkly shaded). The small study area is the same area as shown in Figure III-2 to III-4.

### III. VERIFICATION OF THE SENSITIVE AREAS MAP

The usefulness of the geological delineation of sensitive areas is wholly dependent upon the accuracy with which the geological mapping identifies those areas correctly. To verify the accuracy of the maps, three types of studies were conducted covering geographic regions on three scales. First, an intensive investigation of small headwater streams along a seventy-five mile stretch of the ridgeline of the Blue Ridge Mountains in Shenandoah National Park, Virginia was conducted. Streams were sampled over a variety of geological bedrock types. This work was to verify the assumption that stream water chemistry is strongly influenced by bedrock in regions with thin soils. Second, extensive surveys of low alkalinity lakes and streams in New Hampshire and North Carolina mountainous regions were conducted in the period June-August 1979 so that two sets of highly reliable water quality data would be available. Stations were sampled which were located in regions expected to be sensitive and for which we could obtain data taken prior to 1970. These data sets were then to be used to test the map. The third method of verifying the maps was a geographically large-scale comparison between stream water quality derived from Federal, State and local records and the sensitivity predicted by the maps.

#### A. Intensive Study

Within the overall context of the sensitive areas project, this section focuses on the Blue Ridge of central Virginia.<sup>93</sup> Within this area, pH and alkalinity were measured in headwater streams, and bedrock geology and soils data (where available) were determined from the literature. This work represents both an integrated example and a test of the sensitivity model described in other portions of this report. By working on a small scale, local variation in bedrock geology or soil properties can be considered. In addition, by sampling extensively within a small area, one can determine variability of stream chemistry within a single geological formation, while repeated sampling allows one to assess the stability of individual streams and their response to precipitation events.

##### 1. Site Description.

The study area was located in the Blue Ridge of central Virginia (Figure III-1). Sampling sites extended over approximately 75 miles along both sides of the ridgeline, with major effort centered along a 20 mile section at the southern end of the Shenandoah National Park. With the exception of data presented in Table III-1, discussion will be limited to work carried out within the smaller area (hereafter referred to as the study area). Chemistry/bedrock trends observed within the large area were consistent with those occurring in the study area. Within the study area, the Blue Ridge occurs as

Table III-1. pH and alkalinity of streams in major bedrock formations of the Blue Ridge in Central Virginia.

Formation	Major Rock Type <sup>a</sup>	Age <sup>b</sup>	Number of Samples <sup>c</sup>	pH		Alkalinity (µeq/l) <sup>d</sup>	
				mean	range	mean	range
Pedlar	grandodiorite	pC	22	6.9	5.8- 7.7	188	40- 380
Catoctin	meta-basalt	pC	18	7.0	6.5- 7.65	265	115- 655
Lovington	biotite quartz monzonite gneiss	pC	6	6.8	6.25-7.15	227	198- 270
Marshall	biotite quartz monzonite gneiss	pC	3	7.0	6.8 -7.0	254	205- 305
Old Rag	granite	pC	9	6.5	5.7 -6.8	96	39- 195
Antietam	quartzite	C	10	5.7	5.25-6.1	7	-10-17.5
Hampton + Antietam	quartzite, shale	C	19	5.7	5.1 -6.3	11	-5- 35
various	limestone, dolomite	O	5	8.1	7.65-8.3	3600	3360-3950

<sup>a</sup> For a better description, refer to text.

<sup>b</sup> pC = Precambrian, C = Cambrian, O = Ordovician.

<sup>c</sup> Number refers to number of sampling locations. Where samples were taken at one location on more than one date, an average pH and alkalinity for that location are used. Streams passing through more than one bedrock formation are not included in these calculations, except for streams passing through both the Hampton and Antietam formations.

<sup>d</sup> 1 µeq/l = 0.05 ppm CaCO<sub>3</sub>.



a single major ridge line 2500-4000 feet above sea level, with smaller discontinuous ridges to the east and west. The western slopes are drained by numerous small streams which feed the south fork of the Shenandoah River, while the eastern side is drained by three river systems, the north and south forks of the Moorman's River and Doyles River. Oak-hickory forest covers most of the study area, with pine and hemlock occurring in many of the deeper stream valleys. Nearly all of the sampling stations are in or adjacent to the Shenandoah National Park, and much of the area is designated as wilderness; thus human impact on stream chemistry is negligible.

Soils within the study area fall into two groups; western slopes have coarse thin soils formed from weathering of sedimentary bedrock, while kaolin-itic clays are present in large quantities in soils of the eastern slopes. In both cases, soil pH is low (4.5-5.5), and cation exchange capacity, percent base saturation, and organic content are low in all cases.<sup>5,75,87</sup>


Bedrock geology in the study area (Figure III-2a, 2b) and other areas of the Shenandoah National Park is based on the work of Gathright,<sup>36</sup> while bedrock in other parts of the survey area are based on Werner<sup>97</sup> and Bartholomew.<sup>7</sup> Within the study area, five major bedrock formations occur; these are listed below starting with the oldest and easternmost formation:

1. Pedlar Fm -- Massive or layered granodiorite or quartz monzonite. Usually fine to medium-grained, composed largely of plagioclase and plagioclase-feldspars and quartz.
2. Catoctin Fm -- Thick (2000 ft) beds or flow of metamorphosed basalt, composed primarily of feldspar in a matrix of chlorite, actinolite, epidote and pyroxene. Commonly includes cavities containing quartz and calcite.
3. Weverton Fm -- Thin formation (100-500 ft). Pebbly quartzite beds cemented locally by iron oxides. Interlayered with phyllites.
4. Hampton Fm (Harpers) -- Thick beds (1800-2200 ft) of metamorphosed sandstones and shale regularly interbedded, locally iron-cemented.
5. Antietam Fm (Erwin) -- Thick (700-1000 ft) bedded, very resistant quartzite.


In general, the Pedlar and Catoctin formations occur in the eastern slopes of the Blue Ridge, while the three clastic rock formations are found on the western slopes. Exceptions occur at Madison Run and at the southern end of the study area; the significance of these will be discussed later.

In addition to these five formations, several others are found in the initial survey area:

1. Lovingson Formation -- augen gneiss, in gradational contact with the Marshall Formation. Composed primarily of potassic feldspar, quartz, oligoclase-andesine, and biotite.



Antietam (Erwin) Formation. Cambrian.  
Light-gray to white very resistant quartzite



Hampton (Harper's) Formation. Cambrian.  
Metamorphosed shale and siltstone, interbedded with dark, resistant quartzite



Weverton Formation. Cambrian.  
Conglomeritic quartzite and shale



Catoctin Formation. Precambrian.  
Dark green metamorphosed basalt interbedded with purple phyllites, trace calcite



Pedlar Formation. Precambrian.  
Coarse-grained massive to banded granodiorite



Heavy black line denotes east-west watershed divide




Figure III-2a. Key to bedrock geology in the study area in and adjacent to the Shenandoah National Park.




Figure III-2b. Depiction of bedrock geology in the study area in and adjacent to the Shenandoah National Park.

2. Marshall Formation -- gray to green, medium grained gneissic granite, in gradational contact with the Pedlar Formation. Major constituents are potassic feldspar, quartz, oligoclase-andesine, and biotite.
3. Old Rag Granite -- massive plutonic granite, in gradational or fault contact with the Pedlar Formation. Light gray in color, composed mainly of blue quartz and perthitic microcline.
4. Various Limestone and Dolomite Formations -- light gray, mostly fine grained limestone and dolomite. Beds are a few inches to ten feet thick. Occurrence is at the base of the Blue Ridge in the Shenandoah Valley.

## 2. Methods and Materials.

Field samples were collected in 250 ml polyethylene bottles and stored on ice until returned to the laboratory, then refrigerated until analysis. pH was measured potentiometrically in the field; temperature and conductivity were also measured in the field for each sample. For sequential sampling, samples were collected using an automatic sampler (model S-4040, Manning Environmental Corp.). These samples were stored on ice for return to the laboratory, and pH and conductivity were measured in the laboratory.

Alkalinity and acidity were determined using a double endpoint titration for alkalinity (to pH = 4.5 and 4.2) and single endpoint (pH = 8.3) for acidity.<sup>3</sup> Titrants were 0.005N HCl and 0.005 N NaOH for the respective procedures. Major cations were analyzed by atomic absorption spectrophotometry using an Instrumentation Laboratories 751 spectrophotometer. Major anions and silica were analyzed using standard procedures adopted for Technicon II Auto-analyzer.<sup>32</sup>

## 3. Results and Discussion.

Alkalinity and pH data for the large survey area are summarized in Table III-1, while data for the smaller study area around Shenandoah National Park are presented in Figures III-3 and III-4. Soil chemistry and vegetation are generally similar throughout the survey area, and precipitation chemistry does not vary significantly in the survey area; mean annual pH is approximately 4.2.<sup>34</sup> Consequently, bedrock geology is interpreted to be the primary factor determining stream chemistry in Virginia Blue Ridge; this conclusion is carried through the remainder of this discussion.

Data presented in Table III-1 indicate a wide range of pH and alkalinity regimes for the Virginia Blue Ridge. As should be expected, pH and buffer capacity are highest in those streams originating in limestone and dolomite bedrock. These carbonate rocks react readily with both weak and strong acids, resulting in high bicarbonate concentrations in solution and thus the high pH and alkalinity observed. Only a few such streams were included in this survey, their purpose being to put the relatively low buffer capacity of other systems into perspective.

Streams from the four Precambrian, metamorphic formations (Pedlar, Catoctin, Lovington, and Marshall) have similar pH and alkalinity ranges, with near neutral pH and alkalinities of about 200  $\mu\text{eq/l}$  (10 ppm as  $\text{CaCO}_3$ ). While these levels are far below those of the carbonate streams, buffer capacities in this range are sufficient to maintain a stream in good condition except during extreme episodes of acid input. Even in such cases, the changes should be short-lived and relatively minor. Old Rag granite, while quite similar to the Pedlar and Lovington formations, imparts a significantly lower buffer capacity to streams flowing from it. Long term acidification appears unlikely, but buffer capacity is sufficiently low that these streams probably undergo moderate to substantial pH shifts during acid precipitation events such as those observed in Madison Run (discussed below).

Streams in the clastic bedrock formations (Antietam and Hampton and to a lesser extent, Weverton) are clearly the most vulnerable to both short and long term acidification. The pH of these streams is well above that of incident precipitation, indicating that some neutralization occurs in the watershed, but the buffer capacity of streamwater is minuscule, at best.


Figures III-3 and III-4 show trends similar to those discussed above, and clearly show the effects of different bedrock formations on the eastern and western slopes of the Blue Ridge in the study area. These figures also show the consequence of a stream passing through several bedrock formations; the pH and alkalinity of such streams reflect the most reactive bedrock in their watersheds. Madison Run (upper left corner) and several streams at the bottom left side of the figures have much higher pH and alkalinity than other streams on the western slopes; these differences are presumably attributable to the presence of Catoctin bedrock in their watersheds.

Many of the streams in the survey area were sampled several times during the summer to determine whether there were substantial changes in pH and alkalinity as a result of seasonal variability or changes in stream discharge. With the exception of Madison Run (see below) there seemed to be relatively little change in chemistry of the streams involved.


It should be noted that in all streams studied the carbonic acid-bicarbonate system was the dominant buffer system. Alkalinity titrations showed pH behavior indicative of bicarbonate as the principal buffer, and there were no data to suggest the presence of any other buffer.

Following the initial survey work in the Blue Ridge, Madison Run was chosen for more intensive study during the remainder of the study period. This stream has major tributaries flowing out of the Catoctin formation and from clastic bedrock (Weverton and Hampton formations). During initial sampling these tributaries had pH and alkalinity values typical of the respective bedrock formations. Two kinds of sampling were done; regular (at least bi-weekly) sampling at several stations on the main stream and tributaries, and also sequential samples (usually 2 hour intervals) taken at various stations during both dry periods and during and following individual rainstorms. Sequential sampling during storms has allowed us to observe the response of these streams to acid precipitation episodes.


Madison Run is a typical watershed on the western slope of the Blue



pH = 5.0-5.5



pH = 5.5-6.0



pH = 6.0-6.5



pH greater than 6.5  
(sample station = 0)

Scale in miles




Figure III-3a. pH of streams in the survey area. Shaded areas represent watershed areas upstream from sampling locations.







Figure III-3b. Depiction of streams in the Shenandoah survey area.




Alkalinity = 0-20  $\mu\text{eq}/\ell$



Alkalinity = 20-50  $\mu\text{eq}/\ell$



Alkalinity = 50-125  $\mu\text{eq}/\ell$



Alkalinity = 125-250  $\mu\text{eq}/\ell$



Alkalinity greater than 250  $\mu\text{eq}/\ell$   
(sample station = 0)




Figure III-4a. Alkalinity of streams in the survey area. Shaded areas represent watershed areas upstream from sampling locations.






Figure III-4b. Depiction of alkalinity of streams in the Shenandoah survey area.

Ridge, covering about six square miles. Vegetation cover is virtually complete, consisting mostly of oak forest and extensive blueberry coverings. Soils are coarse, thin, and characterized by low pH, cation exchange capacity, and organic matter. The stream system is not gauged, so no budgets can be determined for individual ions.

Total dissolved solids are low in all of the streams of the watershed, typically around 20-30 mg/l. Silica is the major substance in solution, ranging from 7 to 15 mg/l as  $\text{SiO}_2$ . Bicarbonate is the dominant anion under most conditions, with chloride and sulfate present in lower concentrations on an equivalent basis. Magnesium is the major cation in solution with lower concentrations (equivalents per liter) of calcium, sodium, and potassium. Magnesium is a major constituent of the bedrock formations in the watershed, particularly in the Catoctin formation; so its abundance in comparison to other cations is not unexpected.

Unlike other streams in the survey area, pronounced changes occurred in both Madison Run (Catoctin bedrock), and its major tributary, White Oak Run (clastic bedrock) during the study period. From early June until mid-August, pH increased in both streams, from 6.05 to 6.5 in Madison Run, and from 5.7 to 6.45 in White Oak Run. Alkalinity showed corresponding increases, from 70 to 150  $\mu\text{eq/l}$  and 18 to 80  $\mu\text{eq/l}$  for Madison and White Oak Runs, respectively. We are uncertain of the cause of these changes, but changes in stream discharge/groundwater flow and primary production by mosses and algae growing on bedrock probably both play a role.

In the absence of precipitation, no significant diurnal changes occur in any of the measured parameters on either Madison or White Oak Run. Alkalinity and pH may undergo very small diurnal cycles on White Oak Run in response to photosynthesis. During major precipitation events, however, substantial changes occur, particularly on White Oak Run. On July 16, slightly less than one inch of rain fell during a thunderstorm over the watershed, resulting in major but brief changes in water chemistry on White Oak Run (Figure III-5). Both pH and alkalinity fell substantially, with a corresponding increase in acidity. Nitrate and sulfate levels showed a sharp increase, presumably caused by precipitation input and washoff from leaf surfaces during the storm, while silicon levels declined, probably from dilution of streamwater by precipitation rainwater. The concentrations returned to previous levels over about an eight hour period.

A second storm, on August 11 and 12, was observed on Madison Run. About 2.5 inches of rain fell over a 24 hour period during this storm, resulting from passage of a frontal system. Major ion analysis for this storm indicates that its effects were minimal, despite the much larger amount of precipitation (rainwater was not collected for chemical analysis during either storm). Slight increases were observed in nitrate and sulfate concentration, and silicon levels declined slightly during the storm, but no change was observed in pH, alkalinity or acidity. Apparently the higher buffer capacity of the stream, combined with more gradual input of acid (inferred from sulfate and nitrate values) allowed the stream chemistry to remain relatively constant throughout the storm.




Figure III-5. Changes in water chemistry in White Oak Run during the storm of July 16, 1979. Rainfall occurred between 1300 and 1900 hours.

A third storm, the remnant of Hurricane David, passed through the watershed on September 3, resulting in about 6 inches of rain over a twelve to sixteen hour period. Although the pH of rain from this storm was rather high (4.9), stream chemistry changed tremendously. Sequential samples on White Oak Run showed a severe depression of pH to 5.1, alkalinity decreased to zero, and acidity showed a major increase as bicarbonate was converted to carbonic acid. Thirty-six hours after the storm, pH was still well below normal (5.6), and alkalinity was only 12  $\mu\text{eq/l}$ , about 20 percent of the pre-storm buffer capacity. Madison Run was also sampled following the storm and it too showed much lower pH and alkalinity (5.85 and 60  $\mu\text{eq/l}$ ) than normal.

Data on precipitation events is presented here only to indicate the type of response one might observe in stream chemistry. Clearly, the specific response will depend upon both the pre-storm chemistry of the stream and groundwater in the watershed, and on the composition and magnitude of the precipitation events. In retrospect, Madison Run was probably not an ideal watershed to choose for concentrated study; the pH and alkalinity of Madison and White Oak Run seem to show more seasonal variability than occur on other streams. White Oak Run is clearly not typical of streams flowing from clastic bedrock; by late summer its pH and alkalinity were well above those of other streams originating in clastic rocks.

#### 4. Summary of Intensive Study.

Within the Blue Ridge of central Virginia, the pH and buffer capacity of streams varies greatly, with bedrock geology the major factor controlling stream chemistry. With the exception of streams in areas with carbonate-bearing bedrock, buffer capacity is generally low, and virtually nonexistent in streams originating in clastic bedrock. Sensitivity to acidification is presumably greatest in those streams with the lowest buffer capacity, but we have no historical data available to determine whether significant changes in the chemistry of these streams has occurred during recent years. Thus, these data are valuable as baseline data to determine ongoing long-term changes in stream chemistry.

Storm data from the Madison Run watershed clearly demonstrate short-term effects of acid rain on stream chemistry, particularly on White Oak Run. Many streams in the survey area have pH and buffer capacities significantly lower than occur on White Oak Run; episodic changes in the chemistry of such streams are probably more pronounced. The effects of such changes on stream fauna may be substantial,<sup>43</sup> but are beyond the scope of this study.

#### B. Extensive Surveys of Mountainous Areas of New Hampshire and North Carolina

The states of New Hampshire and North Carolina, separated by over 800 km, receive acid precipitation with volume weighted mean annual  $\text{H}^+$  concentrations of approximately 114  $\mu\text{eq/l}$  (pH 3.94) and 50  $\mu\text{eq/l}$  (pH 4.2).<sup>71</sup> Both states have variable topography with the White Mountains of central New Hampshire and the Blue Ridge Mountains of western North Carolina providing terrain with exposed bedrock, grading to regions of thin soil, then to deep soils. The



Figure III-6. Geologically-based sensitive areas map of New Hampshire. Highest sensitivity (Type-I) is indicated by 1 on the map with 2, 3, and 4 indicating progressively less sensitive areas. Dashed lines are county boundaries.




Figure III-7. Geologically-based sensitive areas map of North Carolina. Highest sensitivity (Type I) is indicated by 1 on the map with 2, 3 and 4 indicating progressively less sensitive areas. Dashed lines are county boundaries.

bedrock also is variable with respect to sensitivity, as can be seen in Figures III-6 and III-7 with abundant outcrops of slow-weathering, igneous and metamorphic rock. Thus these states were selected as ideal regions in which to test the accuracy of the sensitive areas map. The test consisted of identifying specific sites which have low alkalinity ( $< 200 \mu\text{eq/l}^{-1}$ ) from historical data, verifying these observations by new sampling, then comparing the locations of these sensitive waters to the map of sensitive areas. The sensitive areas maps were not consulted when selecting sites for sampling, so that the site selection, within the two regions expected to contain sensitive areas, was not coupled to the map-making process.

## 1. Field Methods.

The first step in selecting field sites where samples were to be collected, was to examine existing water quality records of the departments of fisheries in each state so that new data could be compared to old data in an analysis of temporal trends, as described below (Section IV). Stations which had been previously sampled were then identified on county road maps. Whenever possible, samples were collected at about the same time of year and at the same sites used previously. Sites with alkalinity values greater than  $200 \mu\text{eq/l}^{-1}$  or pH greater than 7.0 were excluded. If recent cultural activities such as farming, timber harvest, residential or recreational housing were significant on site or upstream, the site was not included in the sampling. At each location, two samples were taken in 250 ml acid washed polyethylene bottles, rinsed three times in sample water before filling. One sample was acidified for metal analysis to a pH of 2 by the addition of 1 ml of 6N HCl. These samples were not acidified, but because suspended solids were low, and because most calcium occurs in dissolved rather than particulate form,<sup>70</sup> the amount of calcium added in particulate form is believed to be negligible. The other sample was placed in a cooler filled with ice and kept at or below  $4^{\circ}\text{C}$  until pH and alkalinity measurements could be made (usually every two days). Conductivity in microSiemens was measured with either a Beckman or Presto-tek conductivity bridge. For pH and alkalinity determinations, a Beckman pH meter with combination electrode was used. The samples were brought to room temperature before pH was measured. Alkalinity was determined using Gran's plots. Calcium analyses were performed on acidified samples by atomic absorption spectrophotometry.

## 2. Results of Field Measurements.

The observed alkalinity values (Tables IV-1 and IV-2) in the ranges  $< 0 \mu\text{eq l}^{-1}$ ,  $0-50 \mu\text{eq l}^{-1}$ ,  $50-100 \mu\text{eq l}^{-1}$  and  $100-200 \mu\text{eq l}^{-1}$  are shown for each of the stations sampled in Figures III-8 and III-9. These can be compared to Figure III-6 and Figure III-7, the respective state sensitive areas maps. In New Hampshire all of these low alkalinity waters in the regions sampled fall within regions predicted to have Types I and II sensitivity, and in North Carolina all stations fall in Types I and II sensitive areas. Since stations were selected for sampling only if historical data showed them to have low alkalinities, they were expected to lie in regions underlain by sensitive bedrock. The fact that the geologically based sensitive areas map confirms this expectation constitutes a partial verification of this method of determining sensitive areas.




Figure III-8. Location and alkalinity of New Hampshire sites sampled during the summer of 1979.






Figure III-9. Location and alkalinity of North Carolina sites sampled during the summer of 1979.

## C. Large-Scale Comparison

1. Acidification Chemistry Information Database. Data on Water Chemistry were collated from many Federal, state and local agency sources (Appendix II). Information stored in computerized data bases (e.g. STORET), as well as raw data from field reports gathered from these sources, were used to establish the Acidification Chemistry Information Database (ACID) at Brookhaven National Laboratory. Two factors governed the selection of data. Since alkalinity is the dominant factor of water chemistry which determines sensitivity to acidification, all waters with alkalinity values in excess of 500  $\mu\text{eq/l}$  were excluded. This exclusion selected for moderately to highly sensitive waters and greatly reduced the amount of data to be included in ACID. Second, the time scale for this project limited us to including about half of the available and relevant data which we were able to locate.

A generalized data base management system, SYSTEM 2000 (proprietary system of MRI Systems Corp., Austin, Texas) was used for ACID. Analyses were done using a combination of the CDC 6600 and CDC 7600 computers at Brookhaven National Laboratory. Plotting of large scale maps was accomplished with a 36-inch Versatec electrostatic plotter. Some maps and tables were prepared using a FR 80 COM device and many data tables were put into micro-fiche from the FR 80. Plotting software was developed at BNL and is based on DISSPLA (a plotting software proprietary package of Integrated Software Systems Corp., San Diego).

2. County-by-County Maps. As data are transferred from maps with a scale as fine as 1:62,500 to maps at 1:2,500,000, fine resolution is completely lost. A problem arises in trying to preserve the identity of the isolated sensitive areas. Preserving this information can be accomplished by over-representing the size of the sensitive area. Data in ACID are used to map the current alkalinity and pH on a county by county basis. Alkalinity values were divided into 3 categories: a)  $< 100 \mu\text{eq/l}$ ; b)  $100\text{--}200 \mu\text{eq/l}$ ; and c)  $> 200 \mu\text{eq/l}$  and values at each station for the period (1976-1979) were used for this analysis. The stations included are shown in Figures III-10 and III-11.

Two types of maps are presented:

1. An entire county is assigned the alkalinity category determined by the lowest alkalinity value for the county and shaded according to that value (Figures III-12 and III-13). These maps ensure that areas known to be sensitive on the basis of current water chemistry are identified.
2. A county was assigned the average value of alkalinity, observed at all stations in the county (Figures III-14 and III-15). The existence of low alkalinity waters is frequently lost in these maps, but, when compared to the minimum value maps, the true sensitivity of a whole county is more realistically interpreted.

Maps for all states included in the ACID system are shown in the Appendix (see Endnote, page 91).




Figure III-10. Location of New Hampshire sites where pH and alkalinity data was available for the period 1976-1979.





Figure III-11. Location of North Carolina sites where pH and alkalinity data was available for the period 1976-1979.

# NEW HAMPSHIRE - COUNTIES

## MIN ALK (UEQ/L) 1976-79



**bni**  
**oui**

Figure III-12. Minimum alkalinity of counties in New Hampshire for the period 1976-1979.





Figure III-13. Minimum alkalinity of counties in North Carolina for the period 1976-1979.

# NEW HAMPSHIRE - COUNTIES

AVG ALK (UEQ/L) 1976-79



bni  
qui

Figure III-14. Average alkalinity of counties in New Hampshire for the period 1976-1979.




Figure III-15. Average alkalinity of counties in North Carolina for the period 1976-1979.



Some reservations about these maps must be noted. First, it is most unlikely that ACID contains information on all streams of any county; the sample density must be judged by the site maps for each state (Appendix I), recalling that only waters with alkalinities  $< 500 \mu\text{eq/l}$  were included in ACID. Second, for the average pH or alkalinity maps, a single low value may be all that was available for a given county. Again, the site maps (Appendix I) must be consulted. On the other hand, most state water quality data, and especially data in STORET, are biased in favor of downstream locations of water withdrawal for irrigation or municipal use. The water chemistry data from state fisheries agencies are of great value since we suspect that they give a better representation of upstream waters. Inclusion of additional data which we have located from state and local sources into ACID will improve subsequent versions of these maps.

3. Comparing Geologically-Based Maps to Chemistry-Based Maps. The accuracy of the geologically-based sensitive areas maps of North Carolina and New Hampshire (Figures III-6 and III-7) was evaluated by comparing them to the county-by-county maps of current minimum or average alkalinity (Figures III-12 through III-15), which were assumed to be the "correct" maps. (At this stage the county-by-county maps which showed geology as a binary number, Figure 11-4, were not tested. One cannot distinguish 0 percent from 9 percent on those maps. They are less precise than the parent maps such as Figure III-6). The comparison consisted of first, making a prediction of "sensitivity" or "non-sensitivity" for a particular county based on the bedrock geology alone. A county was predicted to be sensitive if either Type I or Type II bedrock was present, and non-sensitive if neither Type I nor Type II was present. The prediction was then compared to the corresponding surface water alkalinity.

A correct prediction of sensitivity was scored if a county contained Type I or Type II geological areas and the county alkalinity value was  $\leq 200 \mu\text{eq/l}$ . A correct prediction of non-sensitivity was scored if the county contained no Type I or Type II geological areas and the county alkalinity value was  $> 200 \mu\text{eq/l}$ . A false prediction of sensitivity was scored if the county had only Type I or Type II geological areas but the alkalinity value was  $> 200 \mu\text{eq/l}$ . A false prediction of non-sensitivity was scored if the county had no Type I or Type II geological areas and the alkalinity value was  $\leq 200 \mu\text{eq/l}$ . Finally, an indeterminate prediction occurred when a county had Type I and/or Type II plus Type III and/or Type IV geological areas and alkalinity values  $> 200 \mu\text{eq/l}$  were in ACID.

Percentages of these predictions were determined by adding the number of each score class (i.e. in North Carolina correct predictions of sensitivity = 38) and dividing by the number of counties with the appropriate alkalinity score (i.e. in North Carolina  $\leq 200 \mu\text{eq/l}$  = 42), since the alkalinity score is presumed to be "correct." Thus, in North Carolina,

$$\text{correct prediction of sensitivity} = \frac{38}{42} = 90\%$$

Several factors confound the reliability of the geologically based map of sensitivity. The false prediction of sensitivity may be due to insufficient

data in ACID; there may be streams of  $< 200 \mu\text{eq/l}$  for which no data have been located in counties where a false prediction of sensitivity was scored. The false prediction of non-sensitivity may be due to lost resolution of the geological map as contact surfaces were merged and small areas deleted in the mapping process (see Section II of this report). Counties with both sensitive (Types I and II) and non-sensitive (Types III and IV) bedrock may be expected to have waters with alkalinities higher or lower than  $200 \mu\text{eq/l}$  so absence of low alkalinity data in this evaluation of the geologically based map of sensitivity results in the indeterminate score.

The degree of agreement is rather good between the maps based on water chemistry and the geologically based map. The results, shown in Table III-2 and III-3 indicate that the geologically based map predicted the possible sensitivity in 90% of 42 counties which did indeed have sensitive waters in North Carolina and all of such counties in New Hampshire. The "correct" prediction of non-sensitivity was 17%, one of six counties in North Carolina. These counties are located in the eastern portion of the state where soil factors are more important than in the central and western parts of the state. The other five of these six counties, where the lowest alkalinity in ACID was  $> 200 \mu\text{eq/l}$ , were identified by the geologically based map as having Type III or IV bedrock (in some cases, in addition to Type I and Type II). The false prediction of non-sensitivity occurred in only 10% of 42 counties.

In New Hampshire, the geologically based map "correctly" predicted the sensitivity of all 8 counties with minimum alkalinities of  $\leq 200 \mu\text{eq/l}$ , and all 5 counties with average alkalinity values  $\leq 200 \mu\text{eq/l}$ . All 10 counties of New Hampshire had Type I and Type II geological areas and 6 had Type III or Type IV. Both counties with minimum alkalinity greater than  $200 \mu\text{eq/l}$  included Type III or IV geology. Thus, two indeterminate predictions and no false sensitivity predictions were scored.

Following this evaluation of the predictive capability of bedrock geology, the utility of the less precise binary number geological map was assessed by using it to also predict the sensitivity or non-sensitivity of the same counties. A correct prediction of sensitivity was scored if the binary number map indicated the presence of at least 10 percent of Type I or Type II rock in a county and the county alkalinity value was  $\leq 200 \mu\text{eq/l}$ . This resulted because, as pointed out above, zero on the binary number map may actually include up to 9 percent of the particular rock type. Predictions based on the binary number map (Tables III-4 and III-5) agreed quite well with the contour maps. The principal difference, because of the range represented by each number, was an increase in the indeterminate category and a decrease in false-sensitive.

#### D. Map Verification Summary

The ability of the geologically based map to accurately define areas which may be sensitive to acidification has been verified in three ways. An intensive study within the Shenandoah National Park found stream water pH and alkalinity along the Blue Ridge to be controlled primarily by bedrock geology. Data obtained during storm events demonstrate that pH and alkalinity in

Table III-2. Comparison of the geologically-based sensitive areas map (Figure III-7) to county-by-county maps of alkalinity (Figures III-13 and III-15), for 49 of 100 North Carolina counties. See text for definitions of predictions.

	Minimum Alkalinity Basis	Average Alkalinity Basis
Correct prediction of sensitivity	90% of 42 counties	93% of 15 counties
Correct prediction of non-sensitivity	17% of 6 counties	15% of 33 counties
False prediction of sensitivity	17% of 6 counties	39% of 33 counties
False prediction of non-sensitivity	10% of 42 counties	0% of 15 counties
Indeterminate	8% of 48 counties	33% of 48 counties

Table III-3. Comparison of the geologically-based sensitive areas map (Figure III-6) to county-by-county maps of alkalinity (Figures III-12 and III-14) for all 10 counties in New Hampshire. See text for definitions of predictions.

	Minimum Alkalinity Basis	Average Alkalinity Basis
• Correct prediction of sensitivity	100% of 8 counties	100% of 5 counties
Correct prediction of non-sensitivity	0% of 2 counties	0% of 5 counties
False prediction of sensitivity	0% of 2 counties	60% of 5 counties
False prediction of non-sensitivity	0% of 8 counties	0% of 5 counties
Indeterminate	20% of 10 counties	20% of 10 counties

Table III-4. Comparison of the geologically-based binary code sensitive areas map (Figure II-4b) to county-by-county maps of alkalinity (Figures III-13 and III-15), for 49 of 100 North Carolina counties.

	Minimum Alkalinity Basis	Average Alkalinity Basis
Correct prediction of sensitivity	86% of 42 counties	100% of 15 counties
Correct prediction of non-sensitivity	17% of 6 counties	21% of 33 counties
False prediction of sensitivity	83% of 6 counties	0% of 33 counties
False prediction of non-sensitivity	0% of 42 counties	0% of 15 counties
Indeterminate	13% of 48 counties	54% of 48 counties

Table III-5. Comparison of the geologically-based binary code sensitive areas map (Figure II-4a) to county-by-county maps of alkalinity (Figures III-13 and III-15) for all 10 counties in New Hampshire.

	Minimum Alkalinity Basis	Average Alkalinity Basis
Correct prediction of sensitivity	100% of 8 counties	100% of 5 counties
Correct prediction of non-sensitivity	0% of 2 counties	0% of 5 counties
False prediction of sensitivity	0% of 2 counties	60% of 5 counties
False prediction of non-sensitivity	0% of 8 counties	0% of 5 counties
Indeterminate	20% of 10 counties	20% of 10 counties

streams lying on sensitive bedrock fall sharply, presumably in response to inputs of acid precipitation.

Extensive surveys of surface waters with alkalinities  $\leq 200$   $\mu\text{eq/l}$  in North Carolina and New Hampshire, conducted in 1979, found 83-100% of these sites were located in Type I and Type II sensitive areas, as defined by the sensitive areas map.

Information obtained from numerous sources of raw data and from computerized data bases was used to establish the Acidification Chemistry Information Database (ACID). This was used to plot surface water alkalinity on a county-by-county basis. This map, which for testing purposes was considered to be the "correct" map, was compared to the geologically-based map definitions of sensitive areas for each county in North Carolina and New Hampshire. The geologically based sensitive areas map was "correct" for 90% of the 42 counties in North Carolina, and for 100% of the 8 counties of New Hampshire, which had alkalinities  $\leq 200$   $\mu\text{eq/l}$ .

#### IV. TEMPORAL TRENDS IN pH AND ALKALINITY: VALIDATION OF THE SENSITIVE AREAS MAP

Historical records have been used to document temporal changes in the chemistry of lakes brought about by acid precipitation.<sup>1,23,91,96,104</sup> The sensitive areas map, when combined with projections of trends into future air pollution and rain chemistry, may become useful in predicting which regions will be acidified as the deposition of anthropogenic atmospheric acids increases. The fact that acidification has taken place in regions which are both heavily loaded with anthropogenic acids, and are predicted to be susceptible to acidification by the sensitive areas map, is in itself a validation of the predictive value of the map.

Temporal trends in water chemistry were determined for low alkalinity ( $< 200 \mu\text{eq/l}$ ) sites in New Hampshire and North Carolina by comparing historical records to data obtained in the 1979 extensive surveys described in Section III of this report. Historical data from a variety of Federal, state and local sources for most of the other states were also collected and pre-1970 data were compared to post-1975 data.

##### A. North Carolina and New Hampshire: Data

All of the old data obtained for sites resampled in the 1979 field trip to North Carolina came from the North Carolina Department of Inland Fisheries river basin surveys of the early 1960s. Table IV-1 lists all of the sampling sites from 1979 with the corresponding earlier data. For the old surveys, alkalinity determinations were made using the methyl orange technique, and pH was determined colorimetrically.

These methods are less precise and less accurate than methods currently used which are potentiometric pH measurement for pH and multiple end-point potentiometric alkalinity titration. The problems associated with comparing data sets obtained by these different methods are discussed by Zimmerman and Harvey.<sup>105</sup> Determination of alkalinity using a fixed end-point titration to pH 4.5 will require 30 to 40  $\mu\text{eq/l}$  of titrant to remove the free  $\text{H}^+$  in solution at this pH, so alkalinity values are overestimated by this amount, compared to the multiple end-point method (Gran's plot). The old alkalinity values were corrected, therefore, by subtracting 40  $\mu\text{eq/l}$ . In some cases this has resulted in obviously erroneous alkalinity values for the old (corrected) data, e.g. Boylston Creek, with pH 6.9 and a corrected alkalinity value of -18  $\mu\text{eq/l}$  (Table IV-1). Use of color-indicator solutions for pH in low-conductivity, poorly buffered waters will alter the water's pH and cause a spurious reading. There is no way to systematically correct these pH errors mathematically.<sup>105</sup>

One might also question how accurately a single sample represents the water quality of a lake or stream since pH and alkalinity of natural waters

fluctuate daily and seasonally. It is assumed that daily and seasonal fluctuation, when sampled during the day and at the same time of year as the previous sampling, are random and normally distributed. Thus, when reasonably large populations of sample pairs obtained in synoptic surveys are compared, the positive and negative errors should cancel each other out and regional, temporal trends may be analyzed.

Distributions of apparent pH and alkalinity changed in the fifteen to nineteen years between samples (Figures IV-1 to IV-6, uncorrected data). Mean hydrogen ion concentration changed from pH 6.77 in the early 1960's to pH 6.51 in 1979, an 82% increase in hydrogen ion concentration. The pH decreased at 79% of the sampling sites. The differences between the two sets of values was found to be significant at the 0.02 probability level. Mean alkalinity values have also experienced a significant decline since the early 60s from 116  $\mu\text{eq/l}$  (corrected) to 80  $\mu\text{eq/l}$  in 1979, with 71% of the 1979 values being lower than the old (corrected) values.

For New Hampshire, extensive sets of old data, collected in the late 1930s, were found. No other data for small streams and lakes were located except for some scattered records from the 1970s done by the New Hampshire Water Pollution Control Commission. Unfortunately, this older survey contained only pH data for small streams; alkalinity titrations were performed only on lakes. It is not known exactly what methods were used to determine pH and alkalinity for these old surveys, but it was almost certainly some form of colorimetric method for both. The old and new data for New Hampshire are listed in Table IV-2.

A comparison of Figures IV-7 to IV-9 reveals that in New Hampshire there has been an apparent decrease in stream pH values since the 1930s. The mean hydrogen ion concentration for the old data is pH 6.66 and in 1979 it is 6.12, a 247% increase in free hydrogen ion. The pH values at 90% of the sampling sites decreased between the 1930s and 1979. Again, the differences between the two sets of data are significant at the 0.02 probability level. Of the five sites where old alkalinity data were available, four of the new samples had lower alkalinity.

Table IV-1. A comparison of old and new data for headwater streams in North Carolina. Old alkalinity values were corrected by subtracting 40  $\mu\text{eq/l}$ .

Stream	Date	pH	Old		New				
			Alkalinity (as reported) ( $\mu\text{eq/l}$ )	Alkalinity (corrected) ( $\mu\text{eq/l}$ )	Date	pH	Alkalinity ( $\mu\text{eq/l}$ )	Specific Conductance ( $\mu\text{Siemens}$ )	Calcium ( $\mu\text{eq/l}$ )
Roaring Fork	08-03-61	7.4	200	160	06-29-79	6.7	96	21	41
Little Elkin Creek	08-24-61	6.9	180	140	06-29-79	7.1	206	35	103
Pike Creek	08-24-60	7.4	180	140	06-29-79	6.8	96	18	33
Gardner Creek	07-25-61	6.7	200	160	06-29-79	6.9	110	19	35
Rich Mountain Creek	08-25-61	6.9	180	140	06-29-79	6.92	128	22	57
Bullhead Creek	07-02-61	7.0	160	120	06-29-79	6.89	89	15	26
Fall Creek	08-14-61	7.2	200	140	06-30-79	6.8	80	15	35
North Harper Creek	07-19-63	6.9	42	2	06-30-79	6.5	46	10	32
Webb Prong	07-31-63	6.9	176	136	06-30-79	6.78	90	14	44
Wilson Creek	07-31-63	6.9	200	160	07-01-79	6.5	23	18	88
Boone Fork	06-06-63	6.9	200	160	07-01-79	6.52	73	16	77
Laurel Fork	06-06-63	6.7	180	140	07-01-79	6.62	116	32	106
Cold Prong	06-10-63	6.9	140	100	07-01-79	6.3	49	14	41
Moody's Mill Creek	06-12-63	6.7	180	140	07-01-79	6.7	86	19	71
Beech Creek	06-01-63	6.9	200	160	07-01-79	6.8	73	41	108
Buckeye Creek	06-20-63	6.9	160	120	07-01-79	6.59	63	26	76
Blevins Creek	06-25-63	6.7	200	160	07-01-79	6.7	107	50	147
Beetree Creek	06-14-62	6.0	36	-4	07-02-79	6.33	68	12	23
Kiessee Creek	06-14-62		40	0	07-02-79		43	10	24
Cathay's Creek	06-12-62	6.8	100	60	07-02-79	6.4	60	10	22
Cove Creek	06-12-62	6.8	180	140	07-02-79	6.5	77	15	44
Davidson River	06-19-62	6.9	120	80	07-02-79	6.4	61	14	38
Looking Glass Creek	06-12-62	7.0	140	100	07-02-79	6.34	61	14	40
Cooper Creek	07-23-61	6.6	40	0	07-03-79	6.2	60	12	26
Deep Creek	06-28-61	6.7	22	-18	07-03-79	6.4	70	16	48
Dick's Creek	06-27-61	6.8	120	80	07-03-79	6.41	77	19	48
Culowhee Creek	07-19-61	7.2	160	120	07-03-79	6.8	192	29	111
Tennessee Creek	07-24-61	6.6	80	40	07-03-79	6.4	64	13	34
Carson Creek	06-19-62	7.6	140	100	07-04-79	6.4	65	15	38
Williamson Creek	06-19-62	6.9	160	120	07-04-79	6.9	109	18	43
Turkey Creek	06-29-62	6.9	140	100	07-04-79	6.7	67	12	29
South Fork Mill River	07-17-62	6.8	140	100	07-04-79	6.5	57	12	27
Boylston Creek	07-09-62	6.9	22	-18	07-04-79	6.63	145	23	74
North Mill River	07-17-62	6.9	180	140	07-04-79	6.6	67	13	31
South Fork Toe River	06-02-64	6.8	180	140	07-04-79	6.2	23	10	24
Middle Creek	06-08-64	6.8	200	160	07-04-79	5.99	16	10	21
Locust Creek	06-09-64	6.8	180	140	07-04-79	6.4	56	13	38
South Fork Toe River	06-02-64	6.8	200	160	07-04-79	6.7	84	17	48



Table IV-2. A comparison of old and new data for New Hampshire streams and lakes. Old alkalinity values were corrected by subtracting 40 µeq/l. Old alkalinity values were also estimated from new Ca concentration using Henricksen's model<sup>51</sup> and the estimated change in alkalinity calculated by subtracting new alkalinity from estimated old alkalinity.

Stream or Lake	Date	pH	Old		Date	pH	New				
			Alkalinity (as reported) (µeq/l)	Alkalinity (corrected) (µeq/l)			Alkalinity (µeq/l)	Specific Conductance (µSiemens)	Calcium (µeq/l)	Estimated old Alk. from Ca	Estimated ΔAlkalinity
Kimball Creek, Coos County	08-01-38	6.6			07-26-79	6.6	196	37	84	87	+109
Millsfield Pond Bk., Coos Co.	07-13-37	7.0			07-26-79	6.9	124	28	155	188	-64
Horne Brook, Coos County	07-08-37	7.0			07-27-79	6.5	151	25	154	187	-36
Upper Ammonoosuc, Coos County	08-03-37	7.0			07-27-79	7.0	189	37	173	214	-25
Jericho Brook, Coos County	08-08-37	7.0			07-27-79	6.4	139	23			
Wild River, Coos County	07-07-37	6.8			07-27-79	6.4	39	9	55	46	-8
Imp Brook, Coos County	07-06-37	6.8			07-27-79	6.5	63	16	20	-4	+67
Peabody River, Coos County	07-06-37	6.8			07-27-79	6.0	8	44	68	65	-57
Nineteen Mile Brook, Coos Co.	07-06-37	7.0			07-27-79	6.4	63	19	76	76	-13
Ellis River, Carroll County	06-28-37	7.2			07-27-79	6.1	85	14	52	42	+43
Wildcat Brook, Carroll County	06-24-37	7.0			07-27-79	6.2	77	20	70	67	+10
Great Brook, Carroll County	06-24-37	7.2			07-27-79	6.1	85	14	73	72	+13
Slippery Brook, Carroll Co.	06-24-37	7.0			07-27-79	6.4	78	13	81	83	-5
Mountain Pond Outlet, Carroll County	07-15-37	6.1			07-27-79	5.6	25	14	61	55	-30
Paugus Brook, Carroll County	06-15-37	7.0			07-29-79	6.2	53	16	100	110	-57
Swift River, Carroll County	06-15-37	7.0			07-29-79	6.8	155	32	157	191	-36
Wonalancet Brook, Carroll Co.	06-15-37	6.8			07-29-79	6.4	94	29	147	177	-83
Whiteface River, Carroll Co.	06-11-37	6.4			07-29-79	6.3	91	25	118	136	-45
Cold River, Carroll County	06-20-37	7.0			07-29-79	6.4	66	21	107	120	-54
Atwood Brook, Carroll County	06-17-37	6.8			07-29-79	6.3	127	26	195	245	-118
Bearcamp Brook, Carroll Co.	06-19-37	6.6			07-29-79	6.3	127	26	197	248	-121
White Lake, Carroll County	06-21-37	6.5	100	60	07-30-79	5.8	4	12	58	50	-46
Slater Brook, Carroll County	06-12-37	6.2			07-30-79	6.3	119	21	129	151	-32
Chocorua River, Carroll Co.	06-15-37	7.0			07-30-79	6.3	89	20	129	151	-62
Cream Brook, Carroll County	06-29-37	7.2			07-30-79	6.3	189	52	188	235	-46
Moat Brook, Carroll County	06-23-37	6.7			07-30-79	6.3	138	26	162	198	-60
Mason Brook, Carroll County	06-21-37	6.4			07-30-79	6.4	138	21	162	194	-60
Weeks Brook, Carroll County	06-26-37	7.0			07-30-79	6.6	79	23	118	136	-57
Upper Kimbal Lake, Carroll Co.	07-23-37	6.0	140	100	07-30-79	6.2	42	16	91	97	-55
Allard Brook, Carroll County	06-29-37	7.2			07-30-79	6.5	105	21	95	106	-1
Hobbs Brook, Carroll County	06-29-37	6.8			07-30-79	6.2	27	20	98	107	-80
Steam Mill Brook, Carroll Co.	06-30-37	6.4			07-30-79	6.0	50	18	79	80	-30
Oliverian Brook, Carroll Co.	06-29-37	6.4			07-30-79	6.1	40	23	95	103	-63
White Brook, Carroll County	06-29-37	6.2			07-30-79	6.0	69	23	109	123	-54
Downes Brook, Carroll County	06-29-37	6.8			07-30-79	5.6	110	16	57	49	-59
Douglass Brook, Carroll Co.	06-29-37	7.0			07-30-79	6.6	134	18	135	160	-26
Razor Brook, Carroll County	06-28-37	7.0			07-31-79	6.1	103	30	143	171	-68
Albany Brook, Carroll County	06-29-37	7.0			07-31-79	6.1	127	29	157	191	-64
Sawyer River, Carroll County	06-25-37	6.8			07-31-79	5.8	32	24	101	111	-79
Nancy Brook, Carroll County	06-25-37	6.4			07-31-79	5.8	18	18	79	80	-62
Gibbs Brook, Coos County	07-26-38	6.8			07-31-79	5.1	15	17	52	42	-57

(continued)

Table IV-2. (continued)

Stream or Lake	Date	pH	Old		Date	pH	New				
			Alkalinity (as reported) ( $\mu\text{eq/l}$ )	Alkalinity (corrected) ( $\mu\text{eq/l}$ )			Alkalinity ( $\mu\text{eq/l}$ )	Specific Conductance ( $\mu\text{Siemens}$ )	Calcium ( $\mu\text{eq/l}$ )	Estimated old Alk. from Ca	Estimated $\Delta$ Alkalinity
Mt. Pleasant Bk., Carroll Co.	07-25-38	6.9			07-31-79	6.0	6	15	79	80	-74
Clay Brook, Coos County	07-25-38	6.9			07-31-79	6.2	29	12	40	25	+4
Jefferson Brook, Coos County	07-25-38	6.9			07-31-79	6.4	18	12	42	28	-10
South Br. Israel R., Coos Co.	07-13-38	7.0			07-31-79	6.3	63	19	103	114	-51
Deception Brook, Coos County	07-26-38	6.8			07-31-79	6.3	150	32	155	188	-38
Zeland Brook, Grafton County	07-26-38	6.8			07-31-79	6.4	64	25	119	137	-73
Little River, Coos County	07-26-38	6.4			07-31-79	6.4	52	16	94	101	-49
Tucker Brook, Grafton County	07-14-38	6.8			07-31-79	6.4	123	222	449	607	-484
Reel Brook, Grafton County	07-14-38	6.8			08-01-79	5.9	52	6	98	107	-55
Ham Branch, Grafton County	07-14-38	6.8			08-01-79	6.2	150	37	164	201	-51
Black Brook, Grafton County	07-12-38	6.8			08-01-79	6.3	111	29	107	120	-9
Lost River, Grafton County	08-27-39	6.5			08-01-79	6.1	43	124	105	117	-74
Walker Brook, Grafton County	08-27-39	6.7			08-01-79	6.4	76	21	75	75	+1
Cascade Brook, Grafton County	08-27-39	6.1			08-01-79	6.5	116	64	150	181	-65
Hancock Branch, Grafton Co.	08-29-39	6.9			08-01-79	6.0	65	36	116	133	-68
Cedar Brook, Grafton County	08-29-39	6.2			08-01-79	5.9	63	28	97	106	-43
E. Branch Pemigewasset, Grafton County	08-28-39	6.9			08-01-79	6.4	51	24	107	120	-69
Baker River, Grafton County	08-25-39	6.9			08-02-79	6.4	33	18	57	49	-16
Merrill Brook, Grafton County	08-26-39	6.8			08-02-79	6.7	173	38	148	178	-5
Still Brook	07-19-38	7.0			08-02-79	6.5	111	25	101	111	0
Stinson Lake, Grafton County	08-06-38	6.8	70	30	08-02-79	6.2	33	20	88	93	-60
Ellsworth Outlet, Grafton Co.	08-10-38	6.8	100	60	08-02-79	6.1	57	19	106	119	-62
Drakes Brook, Grafton County	07-21-39	6.8			08-02-79	6.2	46	24	103	114	-68
Osceola Brook, Grafton County	07-24-39	6.8			08-02-79	6.3	74	38	117	134	-60
Eastman Brook, Grafton County	07-24-38	7.0			08-02-79	6.6	90	25	103	114	-24
Little East Pond Brook, Grafton County	07-24-38	7.0			08-02-79	6.6	90	25	89	94	-4
Russell Pond, Grafton County	07-24-38	6.6	90	50	08-02-79	6.1	49	20	101	111	-62

NORTH CAROLINA FISHERIES DATA (JUNE, JULY  
or AUGUST 1960-1963)




Figure IV-1. Distribution of alkalinity in samples from North Carolina fisheries data, 1960-1963.

# NORTH CAROLINA FIELD TRIP (6-29-79-7-4-79)




Figure IV-2. Distribution of alkalinity in samples from North Carolina field trip, 1979.

# NORTH CAROLINA FISHERIES DATA (JUNE, JULY or AUGUST 1960-1963)




Figure IV-3. Distribution of pH in samples from North Carolina fisheries data, 1960-1963.

# NORTH CAROLINA FIELD TRIP (6-29-79 - 7-4-79)




Figure IV-4. Distributions of pH in samples from North Carolina field trip, 1979.

# ALKALINITY vs. ALKALINITY FOR NORTH CAROLINA STREAMS




Figure IV-5. Comparative plot of alkalinity for North Carolina streams, 1960-1964 versus 1979.




Figure IV-6. Comparative plot of pH for North Carolina streams, 1960-1964 versus 1979.



# NEW HAMPSHIRE FISHERIES DATA (SUMMERS of 1937, 1938, 1939)




Figure IV-7. Distribution of pH in samples from New Hampshire fisheries data 1937-1939.

# NEW HAMPSHIRE FIELD TRIP (7-26-79-8-2-79)




Figure IV-8. Distribution of pH in samples from New Hampshire field trip, 1979.

pH vs. pH FOR NEW HAMPSHIRE  
STREAMS & LAKES




Figure IV-9. Comparative plot of pH for New Hampshire streams, 1937-1939 versus 1979.

## B. North Carolina and New Hampshire: Discussion

A method for estimating the expected alkalinity value from the observed Ca concentration has been derived by Henriksen,<sup>51</sup> based on water chemistry observations obtained in acidified and non-acidified regions underlain by chemically similar bedrock (described further in Section VI of this report). These calculated values, Alkalinity ( $\mu\text{eq/l}$ ) =  $1.42 \text{ Ca } (\mu\text{eq/l}) - 32$  for the New Hampshire data are presented in Table IV-2. Acidification, defined as loss of alkalinity<sup>51</sup> was estimated by subtracting the expected alkalinity from current measured alkalinity. In this data set, 59 stations had alkalinity values in 1979 which were lower, by an average of 57  $\mu\text{eq/l}$ , than expected. Regional acidification was thus calculated as 57  $\mu\text{eq/l}$  for New Hampshire.

This method of calculating acidification depends on an approximately constant relationship between Ca and Mg. We found that it could not be applied to the North Carolina data.

In 1955-56 the Blue Ridge area of North Carolina was receiving precipitation with a weighted annual average hydrogen ion concentration of pH 4.7-5.2.<sup>69</sup> By 1972-73, the same area was receiving precipitation with a weighted annual average of about pH 4.2.<sup>69</sup> This decline in pH appears to have caused a significant decline in the pH and alkalinity of sensitive streams in the Blue Ridge area of North Carolina. If the pH of rainfall continues to decline in the southeast these streams may experience a continued decline in pH and alkalinity.

The 1979 data for New Hampshire and North Carolina are plotted with Henriksen's<sup>52</sup> empirically derived curve (Figure IV-10) of pH versus [Ca], which separates acidified and non-acidified lakes (for a more detailed discussion, see Section VI). Those points lying above the empirically drawn curve are presumed to have been significantly acidified by precipitation and those below presumably have not. None of the samples from North Carolina lie above the curve, while there has been a significant decrease in pH and alkalinity. Henriksen's curve assumes a constant Ca/Mg ratio, which may not be valid for the North Carolina waters, which are underlain by sedimentary and metamorphic rocks. Approximately 54% of the data points for New Hampshire lie on or above the curve, indicating that many of the sensitive streams and lakes have become significantly acidified. The average pH of samples in New Hampshire is lower, and their calcium values are higher, than in North Carolina.

# CALCIUM vs. pH FOR HEADWATER STREAMS IN N.C. & N.H.




Figure IV-10. Calcium versus pH for headwater streams in North Carolina and New Hampshire. The curve is Henriksen's line.

## V. IMPACTS OF ACIDIFICATION ON AQUATIC BIOTA

Acidification of surface waters in sensitive areas causes a variety of changes in the aquatic flora and fauna. The extent of this impact ranges from changing population strengths to complete elimination of many species. The purpose of this section is to provide a review and description of the kinds of biological impacts which have been observed and which may be anticipated in sensitive waters which are altered by acid precipitation. Impacts on organisms other than fish are stressed here. Impacts on fish are discussed in Section VI. This section is a collation of materials previously presented, and includes references to affects of acid mine drainage on biota.

Although the literature concerning effects of acid mine drainage (AMD) on freshwater ecosystems is extensive, it is not directly applicable to the acid precipitation problem. Often the uncontaminated waters of coal mining areas have higher alkalinity and hardness than the very soft waters acidified by acid precipitation. AMD water usually has a heavy load of iron and other heavy metals, and frequently depresses the oxygen concentration of receiving waters. High turbidity and the presence of chemical floc are also common and greatly alter aquatic habitats. These factors make it very difficult to extrapolate observations from AMD situations to those in the Laurentian Shield, for example.

The variety of species of both plants and invertebrate animals occurring in fresh waters is enormous and speciation differs markedly from one locale to another even though water chemistry may be similar. It is at best difficult and probably futile to try to interpret ecosystem damages at lower trophic levels by comparing lists of species. On the other hand, changes in major processes such as primary and secondary production, and decomposition, can be broadly described and compared. Effects of stress on the major functional guilds may be compared from place to place. Finally a few groups of organisms seem to be remarkably insensitive to strong mineral acidity and are common to many acid environments, while some other groups are clearly intolerant of pH levels below 6.0 to 5.5.

### A. Effects on Microbiota

The production of fish and other animal life in a lake is ultimately dependent upon the availability of organic food resources, primarily plant materials. The sources of organic materials may be divided into two major categories: autochthonous, originating by primary production in the lake, and allochthonous, transported into the lake by inflowing water, airborne litter, or dissolved in rain. The relative importance of each of these sources varies greatly from lake to lake. One principal route for both autochthonous and allochthonous organic matter into the trophic system of a lake is via the detritus.

Bacterial consumption and mineralization of detrital organic matter, both particulate (POM) and dissolved (DOM), allows a cycling of carbon which dominates the structure and the functioning of the system and provides what Wetzel<sup>98</sup> has called a fundamental stability to the system. In the deep, open water of the pelagic zone, where phytoplankton production normally provides a substantial portion of the nonrefractory organic matter, bacteria rapidly assimilate dissolved labile organic substances (PDOM) derived from photosynthesis<sup>29,46</sup> and convert it into bacterial biomass. Particulate organic matter (POM) from phytoplankton is assimilated at a somewhat slower rate. Only a small portion of the PDOM is refractory material likely to survive longer than 24 hours.<sup>88</sup> The bacterial mineralization rate appears to be rather slow, a few percent per day,<sup>17,99</sup> so that this new biomass is actually available to other trophic levels. Not only do the bacteria conserve the energy stored in labile PDOM, which otherwise would be lost from the system, but they also convert (at a slower rate) some of the refractory DOM into a usable form. Fungal and bacterial communities render other POC into forms which are useful for detritivores.<sup>13</sup> The significance of these activities to ecosystem energetics can be better appreciated when one considers that on the order of 90% of the organic carbon in the water column is DOM and that detrital POM is many times larger than the total living carbon biomass.

There are other sources of detritus in the pelagic zone. In some lakes, particularly smaller and/or shallow lakes, macrophytes and benthic algae are important sources of autochthonous organic carbon. Material from these plants may contribute significantly to detritus in the pelagic zone. In deciduous forest lakes, leaf litter falling or blown onto the surface annually has been found to be 200 to 500 g dry leaves per meter of wooded shore line.<sup>35,60</sup> This forest litter, plus that which is added by stream inputs, contributes to both DOM (after leaching) and POM.

Water column detritus generated from all of these sources has three possible fates. It can be transformed biologically, it can sink to the sediments where it accumulates and/or is transformed biologically, or it is lost from the system by outflow. In the first two cases, microbial activity plays a key role in removing detritus.

The inhibition of microbial decomposition can have profound effects throughout an aquatic ecosystem. Detritus removal, conservation of energy, nutrient recycling, primary production, detritivore production and thus production at higher trophic levels, can all be affected by changes in microbial activity. Several investigations have indicated that microbial decomposition is greatly inhibited in waters affected by acid precipitation.

An abnormal accumulation of coarse organic detritus has been observed on the bottom of six Swedish lakes where the pH decreased by 1.4 to 1.7 units in the past three to four decades.<sup>37</sup> Bacterial activity apparently decreased, while in some of the lakes the sediment surfaces over large areas were made up of dense felts of fungal hyphae. In one of the lakes, Gårdsjön, 85% of the bottom in the 0 to 2 m depth zone was covered with a thick felt of algae and organic debris, plus fungal hyphae. Lime treatment caused a rapid decomposition of the organic litter as well as great reductions of the felt,<sup>4</sup> indicating that an inhibition of bacterial activities had taken place at low pH. Similar neutralizations of acidified lakes in Canada resulted in a significant

increase in aerobic heterotrophic bacteria in the water column.<sup>89</sup> Results from field and laboratory experiments with litter bags in Norway<sup>49</sup> also indicate reduced weight loss from leaves in acidic waters. Dissolved organic carbon (DOC) in the inflowing water was found to contribute approximately 50% of allochthonous inputs and 8% of all organic carbon in Mirror Lake, while fine particulate organic carbon (FPOC) was negligible.<sup>60</sup> The extent to which this DOC input is converted to bacterial biomass or otherwise enters into the energetics of a lake is not known. Observations of abnormal accumulations of organic debris have also been made in AMD waters in South Africa<sup>44</sup> and West Virginia (J. DeCosta pers. comm.).

In laboratory experiments the decomposition rates of peptone by microbiota decreased with pH and the oxidation of ammonia ceased below pH 5. Bacterial cell counts and the species number of ciliates also decreased.<sup>12</sup> Numerous other studies indicate that the microbial decomposition of organic materials is markedly reduced at pH levels commonly encountered in lakes affected by acid precipitation.<sup>49</sup>

Accumulations of organic debris and extensive mats of fungal hyphae, as observed in the Swedish lakes,<sup>37</sup> both seal off the mineral sediments from interactions with the overlying water and retain organically bound nutrients which would otherwise have become available if normal decomposition had occurred. The reduction in nutrient availability can be expected to have a negative feedback effect on both plants and bacteria, further inhibiting their activities. The reduction of nutrient supplies to the water column from the sediments, because of the physical covering and from reduced mineralization of organic materials in the water itself, will lead to reduced phytoplankton productivity. These ideas have been formulated into the hypothesis of "self-accelerating oligotrophication" by Grahn *et al.*<sup>37</sup> Qualitative observations support this hypothesis but a quantitative evaluation is lacking.

Reduction of microdecomposer activities may have a direct effect upon the invertebrates. Although certain benthic invertebrates appear to feed directly on the allochthonous detritus material, it seems that "conditioned" (colonized by microorganisms) material is preferred, and that the nutritional value of the detritus is highly increased by conditioning.<sup>13</sup> Bacteria may also be a food source to be removed by the filtering apparatus of organisms such as Calanoida. An inhibition of the microbiota or a reduction in microbial decomposition processes would therefore have a direct impact on the lakes' animal communities.

## B. Effects on Benthic Plants

In waters affected by acid precipitation major changes occur within plant communities. Most of the available data are qualitative and descriptive although some experimentation has been done. Intact lake sediment cores which included the rooted macrophyte *Lobelia dortmana* were incubated at three pH levels (4.0, 4.3 to 5.5, 6.0) at Tovdal in southern Norway. The growth and productivity of the plant ( $O_2$  production) were reduced by 75% at pH 4 compared to the control (pH 4.3 to 5.5) and the period of flowering was delayed ten days at the low pH.<sup>66</sup>



In five lakes of the Swedish west coast, a region severely affected by acid precipitation, in the past three to five decades the macrophyte communities dominated by Sphagnum have expanded. In the sheltered and shaded locality of Lake Orvattnet, in the 0 to 2 m depth zone, the bottom area covered by Sphagnum increased from 8 to 63% between 1967 and 1974. In the 4 to 6 m depth zone, the increase was from 4 to 30%. At the same time, pH in Orvattnet decreased 0.8 units to approximately 4.8.<sup>38</sup> Similar growth of Sphagnum occurred in other Swedish lakes, in Norwegian lakes, and in AMD water as well.<sup>42,44,45,49</sup> At the pH of these acid waters, essentially all of the available inorganic carbon is in the form of  $\text{CO}_2$  or  $\text{H}_2\text{CO}_3$ . Conditions are more favorable for Sphagnum, an acidophile which appears to simply outgrow the flowering plants under acid conditions.

In developing their hypothesis on oligotrophication, Grahn et al.<sup>37</sup> have stressed two biologically important consequences of this Sphagnum expansion. First, Sphagnum has an ion-exchange capacity which results in the withdrawal of base ions such as Ca from solution, thus reducing their availability to other organisms. Secondly, dense growths of Sphagnum form a distinct biotope which is unsuitable for many members of the bottom fauna.

Under some acid conditions, unusual accumulations of both epiphytic and epilythic algae may occur. In the Swedish lakes Mougeotia and Batrachospermum become important components of the benthos.<sup>37</sup> In Lake Oggevatn (pH 4.6), a clear-water lake in southern Norway, not only is Sphagnum beginning to choke out Lobelia dortmana and Isoetes lacustris, but these macrophytes have been observed to be festooned with filamentous algae.<sup>49</sup>

A floristic survey, conducted in Lake Colden (elevation 842 m) in 1932, does not mention Sphagnum, but describes a macrophyte community typical of clear oligotrophic lakes of the Adirondack Mountains (Eriocaulon, Lobelia, Myriophyllum, Isoetes, Utricularia, etc.). On 6/11/58 Colden's surface pH was 5.8. In mid-July, 1979, pH was 4.9 and the flora from the shore to 0.5 m depth (about 2-5 m from shore) around most of the lake perimeter was dominated by extremely dense, uniform stands of Sphagnum pylaesii Brid. (390 g/m<sup>2</sup>). In some areas Utricularia alone or with Sphagnum also formed dense stands which were covered over by flocculent growth of algae in which Fragilaria virescence, Tabellaria fenestrata, Eunotia triodon and Mougeotia sp. were abundant. Prominent mats composed primarily of blue-green algae, fungal hyphae and plant detritus were spreading over the dense Utricularia stands (G. H. Hendrey and F. Vertucci, unpublished data). Colden, once famous for its trout fishing, is now too acidic to support fish life. These observations, the first of their kind in North America, parallel findings in some acidified Swedish and Norwegian lakes. The question of why some acidified lakes show these extreme floristic changes, while many others do not, remains a mystery.

Heavy growths of filamentous algae and mosses occur not only in acidified lakes but have also been reported in streams in Norway affected by acidification. In experiments in artificial stream channels using water and the naturally seeded algae from an acidified brook (pH 4.3 to 5.5), an increase in acidity to pH 4.0 by addition of sulfuric acid led to an increased accumulation of algae compared to an unmodified control.<sup>48</sup> The flora was dominated by Binuclearia tatrana, Mougeotia spp., Eunotia lunaris, Tabellaria flocculosa, and Dinobyron spp., each accounting for at least 20% of the flora at one time

or another. The rate of radioactive carbon uptake per unit of chlorophyll in the channels, measured on just two occasions, was found to be lower in the acid channel by about 30%, suggesting greater algal accumulation at low pH despite lower productivity.

Several factors may contribute to these unusual accumulations of certain algae. The intolerance of various species to low pH or to consequent chemical changes<sup>76</sup> will allow just a few algal species to utilize the nutrients available in these predominantly oligotrophic waters. Many species of invertebrates are absent at low pH, and removal of algae by grazing is probably diminished. Microbial decomposition is inhibited, as was previously noted, also reducing the removal of algal mass.

### C. Effects on Phytoplankton

There is no consistency among various investigations as to which taxa are likely to be dominant under conditions of acidification. The Pyrrophyta may be more common (e.g., species of Peridinium and Gymnodinium) than others in lakes near 4.0. With decreasing pH in the range of 6.0 to 4.0, many species of the Chlorophyta are eliminated although a few tolerant forms are found in the acid range. In the survey of 155 Swedish west coast lakes, blue-green algae became less important with decreasing pH,<sup>1</sup> but Kwiatkowski and Roff<sup>65</sup> found the opposite to be true in lakes of the Sudbury, Ontario, region.

There are, however, conspicuous decreases in phytoplankton species number, species diversity, biomass, and production per unit volume ( $\text{mg}/\text{m}^3$ ) with decreasing pH. Lake clarity and the compensation depth increase with lake acidification, so that areal primary production ( $\text{mg}/\text{m}^2$ ), although lower in acid than nonacid lakes, is not as severely depressed as is production per unit volume.<sup>1,58,65</sup> The low phytoplankton biomass ( $< 1 \text{ mg}/\text{l}$ ) has been correlated to the concentration of available phosphorus, which generally decreases with lower lake pH.<sup>1</sup> Low availability of inorganic carbon has also been suggested as a factor limiting primary production in acidic lakes.<sup>58,63</sup>

Kwiatkowski and Roff<sup>65</sup> have carried out a highly quantitative study of phytoplankton in six lakes of the Sudbury region, which are impacted by atmospheric effluents from a huge nickel-copper smelter complex. Curvilinear equations are presented for phytoplankton diversity, secchi transparency depth, chl a, and productivity.

### D. Effects on Invertebrates

Zooplankton analyzed from net samples collected from 84 lakes in Sweden showed that acidification caused the elimination of many species and led to simplification of zooplankton communities.<sup>1</sup> Crustacean zooplankton were sampled in 57 lakes during a Norwegian lake survey in 1974,<sup>47</sup> and the number of species was found to decrease with pH. The distributions and associations of crustacean zooplankton in 47 lakes of a region of Ontario affected by acid precipitation were strongly related to pH and to the number of fish species present in the lakes. However, fish and zooplankton were each correlated with the same limnological variables, especially pH.<sup>94</sup> Zooplankton communities

become less complex with fewer species present as acidity increases. Food supply, feeding habits, and grazing of zooplankton will probably be altered following acidification, as a consequence of decreased biomass and species composition of planktonic algae and bacteria. In streams continuously polluted by AMD, the number of zooplankton species was low compared to the number of individuals in less polluted conditions downstream.<sup>84</sup>

Surveys at many sites receiving acid precipitation in Norway, Sweden, and North America<sup>4,14,18,47</sup> have shown that waters affected by acid precipitation have fewer species of benthic invertebrates than localities which are less acid. In 832 Norwegian lakes, J. Økland<sup>81</sup> found no snails at pH values below 5.2; snails were rare in the pH range 5.2 to 5.8 and occurred less frequently in the pH range 5.8 to 6.6 than in more neutral or alkaline waters. The amphipod Gammarus lacustris, an important element in the diet of trout in Norwegian lakes where it occurs, is not found in lakes with pH less than 6.0.<sup>82</sup> Experimental investigations have shown that the adults of this species cannot tolerate 24 to 48 hours of exposure to pH 5.0.<sup>15</sup>

In the River Duddon in England, pH is the overriding factor which prevents permanent colonization by a number of species of benthic invertebrates, primarily herbivores, of the upper acidified reaches of the river.<sup>95</sup> In the more acid tributaries (pH < 5.7), the fauna consisted of an impoverished Plecopteran community. Ephemeroptera, Trichoptera, Ancylus (Gastropoda), and Gammarus (Amphipoda) were absent. The epiphytic algal flora was reduced (in contrast to increases noted in Norway), and leaf litter decomposition was retarded. The food supply of the herbivores was apparently decreased, and this may have played a role in the simplification of the benthic fauna. Quantitative data concerning the effects of low pH on the benthic fauna are also available for some acid Norwegian lakes,<sup>49</sup> where notably low standing crops have been observed.

Many studies of invertebrate communities in streams receiving AMD have been conducted. Comparisons are usually made between affected and unaffected zones or tributaries, and experimental acidification has been performed.<sup>54</sup> The numbers of species, species diversity, and biomass are usually greatly reduced. Generally, in AMD waters Chironomidae (midges) and Sialis (alderfly) are the most tolerant macroinvertebrates. The order Trichoptera has more tolerant species than does Plecoptera (stone flies), and Plecoptera has more tolerant species than does Ephemeroptera (mayflies).<sup>25,26,44,45,84,102</sup>

This order of tolerance is essentially the same as seen in waters acidified by acid precipitation. However, the Hemiptera, Notonectidae (backswimmer), Corixidae (water boatman), and Gerridae (waterstrider) are often abundant in acidified soft waters at pH as low as 4.0. This may, in part, be due to lack of fish predation.

Benthic plant communities in lakes may be greatly altered as a consequence of lake or stream acidification (as discussed above). Under these conditions, benthic invertebrate populations may be affected by starvation, evacuation, or extinction due to the loss of preferred habitat. Chironomids<sup>83</sup> and other benthic invertebrates,<sup>20</sup> present in many of the poorly buffered northeastern lakes, have diverse feeding habits and habitats. These invertebrates, in many situations, will be affected by altered decomposition cycles and variations in available foods caused by increased acidification.

The tolerance of aquatic invertebrates to low pH varies over their life cycles, and the emergence of adult insects seems to be a period particularly sensitive to low pH levels. Bell<sup>11</sup> and Moss,<sup>76</sup> in similar studies with Trichopter and Ephemeroptera, found emergence patterns to be affected at pH levels which were higher than the 30-day survival limits. Many species of aquatic insects emerge early in the spring, even through cracks in the ice and snow cover. Because of the contamination of spring meltwaters by atmospheric pollutants, including heavy metals,<sup>41,50,55,56</sup> the early emergers must, in many cases, be exposed to the least desirable water conditions.

Damage to invertebrate communities will influence other components of the food chains. Benthic invertebrates assist with the essential function of removing dead organic material. In litter bag experiments, the effects of invertebrates on leaf composition were much more evident at higher pH than at low pH.<sup>49</sup> A reduction of grazing by benthic invertebrates may also contribute to the accumulation of attached algae in acidified lake and streams.

A short reach of Norris Brook, a tributary to Hubbard Brook in New Hampshire, was acidified to pH 4 in the spring-summer of 1977, to evaluate the effects of acidification on a stream ecosystem. Excessive accumulations of algae occurred, bacterial biomass and heterotrophic activity per unit of organic matter were reduced, and both invertebrate diversity and biomass decreased.<sup>43</sup>

In unstressed lake ecosystems there tends to be a continuous emergence of different insect species available to predators from spring to autumn. In acid-stressed ecosystems the variety of prey is reduced and periods may be expected to occur in which the amount of prey available to fish (and waterfowl) is diminished.

#### E. Effects on Vertebrates

Pough<sup>86</sup> has described effects of acid precipitation on spotted salamanders (Ambystoma jeffersonianum and A. maculatum), which breed in temporary rain pools. Below pH 5 and 7, respectively, these species suffered high mortality during hatching in laboratory tests. This mortality was associated with distinctive embryonic malformations. The development of salamander eggs in five ponds near Ithaca, New York, ranging from pH 4.5 to 7.0, were observed. An abrupt transition from low to high mortality occurred below pH 6. Although a synergistic effect of several stresses may have been possible, the studies suggested that pH was the critical variable. Pough cites studies which indicate a decline in British frog populations.<sup>86</sup>

In Tranevatten, a lake acidified by acid precipitation, near Gothenburg, Sweden, the lake pH has declined to 4.0 to 4.5, and all fish have been eliminated. The frog species Rana temporaria is being eliminated as well. Currently, only adults 8 to 10 years old are found. While many egg masses were observed in 1974, few were found in 1977. The few larvae observed in 1977 subsequently died. Another species, Bufo bufo, is also being eliminated from this lake.<sup>42</sup>

Frogs and salamanders are important predators on invertebrates, including mosquitoes and other pests, in lakes and puddles or pools. In turn, they are themselves important prey for higher trophic levels in an ecosystem.

#### F. Biological Effects Summary

Acid precipitation, by causing increased acidity in lakes, streams, pools, and puddles, can cause slight-to-severe alteration in communities of aquatic organisms. The effects are similar to those observed in waters receiving acid mine drainage (AMD), but the toxicology and chemistry are not as complicated by the presence of high concentrations of heavy metals, chemical flocs, turbidity, etc., such as are found in AMD.

The kinds of effects that are likely to result from increasing acidification are shown in Table V-1. In order to provide step-functions for damages, which may be of use in modeling ecosystem acidification, a summary of damages to aquatic organisms as functions of pH is presented in Table V-2.

Table V-1. Damages to aquatic biota likely to occur with inceasing acidity.

1. Bacterial decomposition is reduced and fungi dominate saprotrophic communities. Organic debris accumulates rapidly.
2. The ciliate fauna is greatly inhibited.
3. Nutrient salts are taken up by plants tolerant of low pH (mosses, filamentous algae) and by fungi. Thick mats of these materials may develop which inhibit sediment-to-water nutrient exchange and choke out other aquatic plants.
4. Phytoplankton species diversity, biomass, and production are reduced.
5. Zooplankton and benthic invertebrate species diversity and biomass are reduced. Remaining benthic fauna consists of tubificids and Chironomus (midge) larvae in the sediments. Some tolerant species of stone flies and mayflies persist as does the alderfly. Air-breathing bugs (water-boatman, backswimmer, water strider) may become abundant.
6. Fish populations are reduced or eliminated (see Section VI).

Table V-2. Summary of damages to aquatic organisms with decreasing pH.

pH range	Effects
8.0-6.0	<p>Long-term changes of less than 0.5 pH units likely to alter the biotic composition of freshwaters to some degree. The significance of these slight changes is however, not great.</p> <p>A decrease of 0.5 to 1.0 pH units in the range 8.0 to 6.0 may cause detectable alterations in community composition. Productivity of competing organisms will vary. Some species will be eliminated.</p>
5.5-5.0	<p>Many species will be eliminated, and species numbers and diversity indices will be reduced. Crustacean zooplankton, phytoplankton, molluscs, amphipods, most mayfly species, and some stone fly species will begin to drop out. In contrast, several pH-tolerant invertebrates will become abundant, especially the air-breathing forms (e.g., Gyrinidae, Notonectidae, Corixidae), those with tough cuticles which prevent ion losses (i.e. <u>Sialis lutaria</u>), and some forms which live within the sediments (<u>Oligochaeta</u>, Chironomidae, and Tubificidae). Overall, invertebrate biomass will be greatly reduced.</p>
5.0-4.5	<p>Decomposition of organic detritus will be severely impaired. Autochthonous and allochthonous debris will accumulate rapidly. Most fish species will be eliminated.</p>
< 4.5	<p>All of the above changes will be greatly exacerbated, and all fish will be eliminated.</p>

## VI. REGIONAL ASSESSMENT OF ACIDIFICATION IMPACTS OF FISH POPULATION

### A. Introduction

Acidification of surface waters in geologically sensitive regions by acid precipitation may be viewed as a continuous process of water quality change, analogous to the acidimetric titration of a bicarbonate solution. The "titration curve" can be segmented into three stages, which define both the extent and nature of water quality change and the biological impacts. The initial stage of acidification is characterized by decreased alkalinity, but pH levels remain greater than 6.0 and bicarbonate buffering is maintained. No significant impacts on fish population have been observed at this level of acidification. Loss of  $\text{HCO}_3$  buffering, and resulting severe temporal pH fluctuations occur in the second stage of acidification. Stress, reproductive inhibition, and episodic mortalities may initiate recruitment failure and eventual extinction of fish populations during this stage of acidification. The final stage of acidification is characterized by chronically depressed pH and elevated toxic metal levels. Fish are generally absent from waters at this level of acidification. This general view of acidification impact levels on fish populations is supported by extensive, carefully documented data from sensitive regions in Norway<sup>103</sup> and eastern North America.<sup>9,91</sup>

More recently, Henriksen,<sup>52,53</sup> has developed models of the acidification process which quantitatively relate the stages of acidification described above to precipitation acidity levels in sensitive regions. These models provide a basis for predicting the regional extent and levels of acidification which have occurred in sensitive areas where historical data are lacking. The models also permit evaluation of expected fish population status on a regional level. Considering the paucity of carefully documented fish population data for waters in sensitive areas of the eastern United States and the time constraints for obtaining and interpreting new data, it was decided to utilize the predictive models to obtain a regional perspective of acidification impacts on fish populations. The approach taken was to first calibrate the Norwegian model using North American data and verify the predictions obtained by comparison with known water quality and fish population data from the Adirondack region (New York) and Experimental Lakes Area (Ontario, Canada). The next step will be to apply the models to water quality data from other sensitive regions in the United States to evaluate acidification and population status.

### B. Procedure

1. Data sources. Water quality and fish population data obtained from the following sources were utilized in model calculation and testing.

1. Water chemistry and fish survey data for Adirondack Mountain lakes (New York): Schofield (1976 and file data); New York State Biological Surveys of the Upper Hudson, Raquette, Mohawk-Hudson, and Black River Watersheds (1931-1934, NYS Dept. Conservation).
2. Stream water quality and fish population status in selected areas of West Virginia: Personal Communication and file data from Donald Gasper, West Virginia Dept. of Natural Resources.
3. Water quality data for lakes in the BWCA, N.E. Minnesota: G. E. Glass, EPA Research Laboratory (Duluth), Draft report on Impacts of Air Pollutants on Wilderness Areas of N. Minnesota.
4. Water chemistry and fish population data for Connecticut lakes.<sup>24,79</sup>
5. Water chemistry and fish population data for Rhode Island lakes.<sup>39,40</sup>
6. Water chemistry for Massachusetts lakes.<sup>21</sup>
7. Water chemistry and fish population for ELA lakes.<sup>9</sup>
8. Water chemistry and fish population data for the North Carolina bay lakes.<sup>30,31</sup>
9. Water chemistry for coastal plain streams in South Carolina and Georgia.<sup>10</sup>

#### C. Model Calibration and Verification

1. Acidification. Acidification is defined as the decrease in alkalinity, which results from an equivalent increase in strong acid input. Three independent approaches to estimating regional acidification were applied to the Adirondack lake data base. The first approach developed by Henriksen<sup>51</sup> utilized the empirical relationships between precipitation pH (volume weighted annual mean) and estimated regional acidification of lakes in southern Norway. Regional acidification, a decrease in alkalinity associated with excess sulfate (corrected for seasalt) in precipitation (Ac-S,  $\mu\text{eq/l}$ ), was calculated by regression as:

$$\text{Ac-S} = \frac{(10^{-\text{pH}} \times 10^6) - 17}{0.38} \quad (1)$$

The volume weighted, mean annual hydrogen ion concentration for precipitation collected in the western Adirondack region (Hinckley, New York; USGS network) in 1975 was pH 4.26. Substituting this value in Equation (1) yields an estimated acidification level of 99.9  $\mu\text{eq/l}$  as the average regional acidification of the western Adirondacks.

The second approach to estimating acidification was also as given by Henriksen<sup>51</sup>: Acidification = preacidification alkalinity - present day alka-



linity. In this case, preacidification alkalinities (ALKp) for the Adirondack lakes were estimated from 1975 measurements of calcium concentrations and the empirical relationships between calcium concentration and alkalinity observed in the Experimental Lakes Area (Ontario) waters:

$$\text{ALKp } (\mu\text{eq/l}) = 1.42 (\text{Ca } \mu\text{eq/l}) - 32 \quad (2)$$

The assumptions made in employing this method of estimating preacidification alkalinities are as follows:

- a) Alkalinity in pristine (unacidified) lakes (e.g., the ELA region) is due primarily to  $\text{HCO}_3^-$ , which is accompanied by approximately equivalent amounts of calcium and magnesium, and the Ca/Mg ratio is relatively constant. This relationship (Equation 2) is of a general nature and can be applied to other regions with similar geological settings. This assumption is supported by the similarity of the calcium-alkalinity relationship for pristine lakes in northern Norway and the ELA.<sup>52</sup>
- b) Calcium levels do not increase or decrease in response to acidification. Historical calcium data are not available for the Adirondack region, but estimates of calcium from the 1930s alkalinity data and Equation (2) were not significantly different ( $p = 0.05$ ) from calcium levels measured in 1975 for 36 Adirondack lakes. Additional support for this assumption comes from studies in Sweden<sup>73</sup> which demonstrated that no significant changes in calcium concentration were detected in acidified lakes of south-central Sweden, where significant increases in  $\text{H}^+$  and  $\text{SO}_4^{2-}$  did occur. Also, Wiklander<sup>100</sup> reported that base exchange efficiency is very low in acidic podzols, where the exchange complex is typically dominated by  $\text{H}^+$  and  $\text{Al}^{3+}$  ions, rather than  $\text{Ca}^{2+}$ . Watt (1979) compared the pH and ionic composition of lakes in Nova Scotia sampled in 1955 and 1977. Significant decreases in pH levels were observed but no significant change in  $\text{Ca}^{++}$  was found.<sup>96</sup>

Acidification (Ac-Ca) was then calculated as the difference between preacidification alkalinity (Equation 2) and current alkalinity measurements (ALK\*):

$$\text{Ac-Ca } \mu\text{eq/l} = \text{ALKp} - \text{ALK*} \quad (3)$$

The estimated mean acidification for 215 Adirondack lakes sampled in 1975 was  $\text{Ac-Ca} = 99.6 \mu\text{eq/l}$  ( $\text{SD} = 30.6$ ), which compares well with the first estimate of acidification ( $\text{Ac-S} = 99.9 \mu\text{eq/l}$ ). These estimates of lake acidification are not correlated with current pH levels (Figure VI-1), which is indicative of the regional uniformity of acidification. Estimated acidification levels are significantly correlated with sulfate levels (Figure VI-2), indicating that atmospheric strong acid ( $\text{H}_2\text{SO}_4$ ) inputs are the primary source of acidification.

The third approach to estimating acidification was to directly compare historical (1930s data) and current (1975) alkalinity measurements. Alkalinity measurements obtained in the 1930s for Adirondack lakes employed methyl orange indicator for end point determination in the titration of water samples




Figure VI-1. Acidification of Adirondack lakes in relation to pH measured in 1975 (215 lakes). Mean acidification = 99.6  $\mu\text{eq/l}$ , SD = 30.6, where acidification ( $\mu\text{eq/l}$ ) = 1.42 (Ca  $\mu\text{eq/l}$ ) - 32 (alk  $\mu\text{eq/l}$ ).




Figure VI-2. Acidification of Adirondack lakes in relation to  $\text{SO}_4$  measured in 1975 (215 lakes).  
 Regression: (a) acidification =  $0.676 (\text{SO}_4 \mu\text{eq./l}) + (\text{data})$ ,  $r = 0.62$ ,  $p < 0.01$ ;  
 (b) acidification =  $0.89 (\text{exc } \text{SO}_4 \mu\text{eq./l}) - 12$ .

for total alkalinity measurement. This method yields an apparent end point pH of approximately 4.4, which overestimates the true alkalinity (as measured by Gran plot location of the titration curve inflection for the 1975 data) by the amount of free  $H^+$  ion present in solution at this pH level ( $40 \mu\text{eq/l } H^+$ ). The 1930s alkalinity measurements were corrected to correspond to true alkalinity levels by subtracting  $40 \mu\text{eq/l}$  from each value. Acidification was then estimated as the difference between corrected 1930s alkalinity levels and 1975 alkalinity measurements. The mean acidification for 36 Adirondack lakes, where data were available from both periods, was  $104.7 \mu\text{eq/l}$  (SD 50.9). Estimated acidification for the same lakes, using the calcium-alkalinity model (Equation 2), was  $105.7 \mu\text{eq/l}$  (SD 31.8) which is not significantly different from the previous estimate ( $p = 0.01$ ).

2. Lake classification. Lakes were classified into three categories, which define acidification and fish population status as discussed previously, utilizing regional acidification estimates and lake calcium concentrations. Henriksen<sup>53</sup> constructed a nomogram by grouping data from 719 lakes in southern Norway into discrete pH classes and then determined the least squares regressions of calcium on acidification for each class. The resulting nomograph (Figure VI-3) can then be used to predict lake pH (and fish population status) from lake calcium levels and either regional acidification or precipitation pH. The two regression lines derived from the Norwegian lake data define the boundaries for the three classifications:

$$\text{HCO}_3 \text{ lake boundary} \quad \text{Ca } (\mu\text{eq/l} = 49.9 (0.28 + 0.0192 \text{ Ac-S}) \quad (4)$$

$$\text{Acid lake boundary} \quad \text{Ca } (\mu\text{eq/l} = 49.9 (0.07 + 0.0139 \text{ Ac-S})$$

The characteristics, expected pH values, and boundary calcium levels for this classification, as applied to the Adirondack data, are given in Table VI-1.

The observed pH levels for lakes classified as described below are illustrated in Figure VI-3. For the Adirondack data, 70% of the lakes were correctly classified and 98% of the ELA lakes were correctly designated as  $\text{HCO}_3$  lakes. These results are similar to those obtained by Henriksen<sup>53</sup> for classification of lakes in southern Norway. An additional test of the validity of this model for predicting lake pH changes resulting from acidification is illustrated in Figure VI-4. Increase in  $H^+$  concentration between the 1930s and 1975, as determined from differences in pH, were compared to calcium concentrations measured in 1975 for 36 Adirondack lakes. The break in the plot of  $H^+$  against calcium occurs precisely at the calcium level predicted from Equation (4). The results of these model predictions for Adirondack and ELA lake data appear to be quite adequate for determining acidification status.

Fish population status in the Adirondack lakes is presented in Table VI-2. The frequencies of lakes not supporting fish life in each of the three lake classifications are comparable to those reported by Henriksen<sup>53</sup> for lakes in southern Norway. The lakes of the ELA are still predominantly pristine,  $\text{HCO}_3$  type waters and the fish populations are unaffected.<sup>9</sup>




Figure VI-3. Nomograph for predicting lake pH and fish population status from lake Ca levels and either regional acidification or precipitation pH. Regression lines for the boundary between HCO<sub>3</sub> and transition lakes, and for the acid lakes boundary with transition lakes, are given in the text as Equations (4) and (5) respectively.




Figure VI-4. Changes in the  $H^+$  ion concentrations in 36 Adirondack lakes from the 1930s to 1975, in relation to Ca levels measured in 1975. The arrow designates the critical Ca level for loss of  $HCO_3$  buffering as predicted from Henriksen's model<sup>53</sup> for a region exposed to precipitation with a weighted mean pH of 4.26.

Table VI-1. Lake classification for Adirondack Mountain lakes, based on the nomograph of Henriksen.<sup>53</sup>

Lake Class	Expected pH	Fish Population	Boundary Ca $\mu$ eq/l (Adirondack)
HCO <sub>3</sub>	> 5.3	Good	> 110
Transitional	4.7 - 5.3	Sparse	73 - 100
Acid	< 4.7	Barren	< 73

Table VI-2. Fish population status in 214 Adirondack Mountain lakes, classified according to the nomograph of Figure VI-3.

Lake Type	HCO <sub>3</sub>	Trans	Acid
Expected pH	>5.3	4.7-5.3	<4.7
Number of lakes	69	97	48
Number in pH range	61	65	27
% correctly classified	88.4	67.0	56.3
% without fish	2.9	54.6	91.7

#### ENDNOTE

The 94 pages of Appendix I and II are available separately from National Technical Information Service (NTIS) as EPA-600/3-80-024b under the same title as this volume (EPA-600/3-80-024a). The NTIS address is Springfield, VA 22161.

Appendix I contains four maps for each of 22 states, depicting minimum and average alkalinity of surface waters on a county basis, and indicating the location of pH and alkalinity sampling stations with the measured data range for 1976-1979. The states included are: Connecticut, Delaware, Florida, Georgia, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, North Carolina, Pennsylvania, Rhode Island, South Carolina, Tennessee, Vermont, Virginia, West Virginia, and Wisconsin.

Appendix II is a three page list of additional sources for water chemistry data which have been used in this report.

## REFERENCES

1. Almer, B., W. Dickson, C. Ekström, E. Hörnström, and U. Miller. 1974. Effects of acidification of Swedish Lakes. *Ambio* 3:330-336.
2. Almer, B., W. Dickson, C. Ekström, and E. Hörnström. 1978. In Sulfur in the Environment, Part 2 (ed. Nriagu, J. O.) 272-311 (Wiley, New York).
3. American Public Health Association. 1971. Standard Methods for the Examination of Water and Wastewater. Thirteenth ed. APHA. Washington, D.C.
4. Andersson, I., O. Grahn, H. Hultberg, and L. Landner. 1974. Jamforande undersökning av olika tekniker för alterställande av försurade sjöar. Institute for Water and Air Research, Stockholm, STU Report 73-3651.
5. Anonymous. 1979. Soil Survey of Augusta County, Virginia. USDA Soil Conserv. Ser. and For. Ser.
6. Barnes, R. S., D. E. Spyridakis, and P. B. Birch. 1979. Atmospheric input of trace metals to lake sediments in western Washington State (abs.). Am. Chem. Soc. Ann. Meet., Environ. Div.
7. Bartholomew, M. J. 1978. Geology of the Greenfield and Sherando Quadrangles, Virginia. Virginia Dept. of Mineral Resour. Pub. No. 4.
8. Beamish, R. J. and H. H. Harvey. 1972. *J. Fish Res. B. Can.* 29:1131-1143.
9. Beamish, R. S., L. M. Bloues, and G. A. McFarlane. 1976. Tech. Rep. No. 607, Dept. of Environ., Fish and Marine Serv. Can.
10. Beck, K. C., J. A. Router, and E. M. Perdue. 1974. *Geochem. et. Cosmochem.* 38:344.
11. Bell, H. L. 1971. Effect of low pH on the survival and emergence of aquatic insects. *Water Res.* 5:313-319.
12. Bick, H. and E. F. Drews. 1973. Selbstreinigung und Ciliatenbesiedlung in saurem Milieu (Modellversuche). *Hydrobiologia* 42:393-404.
13. Boling, R. H., Jr, E. D. Goodman, J. A. VanSickel, J. O. Zimmer, K. W. Cummins, R. C. Petersen, and S. R. Reice. 1975. Toward a model of detritus processing in a woodland stream. *Ecology* 56:141-151.



14. Borgström, R., J. Brittain, and A. Lillehammer. 1976. Evertebrater og surt vann: oversikt over innsamlingslokaliteter. SNSF IR 21/76. SNSF-Project, 1432 Aas-NLH, Norway.
15. Borgström, R. and G. R. Hendrey. 1976. pH tolerance of the first larval stages of Lepidurus articus (Palas) and adult Gammarus lacustris G. O. Sars. SNSF IR 22/76. SNSF-Project, 1432 Aas-NLH, Norway.
16. Cogbill, C. V. and G. E. Likens. 1974. Precipitation in the north-eastern United States. Water Resour. Res. 10:1133-1137.
17. Cole, J. J. and G. E. Likens. 1978. Measurement of mineralization of phytoplankton detritus in an oligotrophic lake. (Manuscript in press).
18. Conroy, N., K. Hawley, W. Keller, and C. LaFrance. 1975. Influences of the atmosphere on lakes in the Sudbury area. J. Great Lake Res. 2, Suppl. 1(1976):146-165.
19. Cragin, J. H., M. M. Herron, and C. C. Langway, Jr. 1975. The chemistry of 700 years of precipitation at Dye 3, Greenland. CRREL Res. Report No. 341, 18 pp.
20. Cummins, K. W. 1973. Trophic relations of aquatic insects. Ann. Rev. Entomol. 18:183-206.
21. Daly, L. M. 1965. DJ Completion Report F-20-R-2, Mass. Fish and Game Dept.
22. Davis, R. B., S. A. Norton and D. F. Brakke. 1979. Heavy metal deposition in bottom sediments and acidification of New England lakes from atmospheric inputs, and the effect on lake biota (abs.). Conf. on Great Lakes Res.
23. Davis, R. O., M. O. Smith, J. H. Bailey, and S. A. Norton. 1978. Acidification of Maine lakes by acid precipitation. Verh. Intern. Verein. Limnol., (XX Congress, Copenhagen, 1977). Vol. 20:532-537.
24. Deevey, E. S. and J. S. Bishop. 1942. Bull. No. 63, Conn. State Bd. Fish and Game.
25. Dills, G. and D. T. Rogers. 1974. Macroinvertebrate community structure as an indicator of acid mine pollution. Environ. Pollut. 6:239-262.
26. Dinsmore, B. H. 1968. The aquatic ecology of Toms Run, Clarion County, Pennsylvania, preceding watershed reclamation. Pennsylvania Dept. of Health, Bureau of Sanitary Engineering, Div. of Water Qual. Pub. No. 21.
27. Dochinger, L. S. and T. A. Seliga (eds.). 1976. Proc. of the First Int. Symp. on Acid Precip. and the For. Ecosystem. USDA For. Ser. Gen. Tech. Rep. NE-23.
28. Edgerton, E. Personal communication.

29. Fogg, G. E. 1977. Excretion of organic matter by phytoplankton. *Limnol. Oceanogr.* 22:576-577.
30. Frey, D. G. 1949. *J. Elisha Mitch. Soc.* 65:1.
31. Frey, D. G. 1951. *J. Elisha Mitch. Soc.* 67:1.
32. Galloway, J. N. and G. E. Likens. 1976. Calibration of collection procedures for the determination of precipitation chemistry. *Water, Air, and Soil Pollut.* 6:241.
33. Galloway, J. N. and E. B. Cowling. 1978. The effects of precipitation on aquatic and terrestrial ecosystems: a proposed precipitation chemistry network. *APCA J.* 28:229.
34. Galloway, J. N. 1979. Unpublished data.
35. Gasith, A. and A. D. Hasler. 1976. Airborne litterfall as a source of organic matter in lakes. *Limnol. Oceanogr.* 21:253-258.
36. Gathright, T. M. II. 1976. Geology of the Shenandoah National Park. Virginia Dept. of Mineral Resour. Bull. 86.
37. Grahn, O., H. Hultberg and L. Landner. 1974. Oligotrophication--a self-accelerating process in lakes subjected to excessive supply of acid substances. *Ambio* 3:93-94.
38. Grahn, O. 1975. Macrophyte succession in Swedish lakes caused by deposition of airborne acid substances. In *Proc. of the First Int. Symp. on Acid Precip. and the For. Ecosystem*, Ohio State Univ., 12-15 May 1975. USDA Forest Service Gen. Tech. Report NE-23, 1976, p. 519-530.
39. Guthrie, R. C., J. A. Stolgitis, and W. L. Bridges. 1973. Fish Rep. No. 1, Rhode Island Dept. Nat. Res.
40. Guthrie, R. C. and H. A. Stolgitis. 1977. Fish Rep. No. 3, Rhode Island Dept. Nat. Res.
41. Hagen, A. and A. Langeland. 1973. Polluted snow in southern Norway and the effect of the meltwater on freshwater and aquatic organisms. *Environ. Pollut.* 5:45-57.
42. Hagström, T. 1977. Grodornas Försvinande i en Försurad sjö (The disappearance of frogs in an acidified lake). *Sveriges Natur* 11:367-369.
43. Hall, R. J., G. E. Likens, S. Fiance, and G. Hendrey. 1980. Experimental stream acidification in the Hubbard Brook Experimental Forest. Accepted for publication in *Ecology*.
44. Harrison, A. D. 1958. The effects of sulphuric acid pollution on the biology of streams in the Transvaal, South Africa. *Verh. Internat. Ver. Limnol.* 13:603-610.

45. Harrison, A. D. and J. D. Agnew. 1962. The distribution of invertebrates endemic to acid streams in the western and southern Cape Province. *Ann. Cape Prov. Mus.* 2:273-291.
46. Hellebust, J. A. 1974. Extracellular products. In W. D. Steward (ed.) *Algal physiology and biochemistry*. Univ. Calif.
47. Hendrey, G. R. and R. F. Wright. 1975. Acid precipitation in Norway: Effects on aquatic fauna. In Proc. First Specialty Symp. Atmos. Contrib. Chem. Lake Waters, Int. Assoc. Great Lake Res., Orillia, Ontario, 28 Sept-1 Oct 1975. *J. Great Lakes Res.* 2 Suppl. 1:192-207.
48. Hendrey, G. R. 1976. Effects of low pH on the growth of periphytic algae in artificial stream channels. SNSF-Project, 1432 Aas-NLH, Norway.
49. Hendrey, G. R., K. Baalsrud, T. Traaen, M. Laake and G. Raddum. 1976. Acid precipitation: Some hydrobiological changes. *Ambio* 5:224-227.
50. Henriksen, A. and R. F. Wright. 1976. Concentrations of heavy metals in small Norwegian lakes. SNSF-Project Report. SNSF-Project, 1432 Aas-NLH, Norway.
51. Henriksen, A. 1979. A simple approach for identifying and measuring acidification on freshwater. *Nature* Vol. 278.
52. Henriksen, A. 1979a. *Nature* 278:542.
53. Henriksen, A. Personal communication. (Manuscript in preparation).
54. Herricks, E. E. and J. Cairns, Jr. 1974. Rehabilitation of streams receiving acid mine drainage. Bull. 66, Virginia Water Resour. Res. Center, VPI, Blacksburg, Virginia 24061.
55. Hultberg, H. 1975. Thermally stratified acid water in late winter--a key factor inducing self-accelerating process which increase acidification. In Proc. First Int. Symp. Acid Precip. and the For. Ecosystem, Ohio State Univ., 12-15 May, 1975. USDA For. Ser. Gen. Tech. Rep. NE-23, 1976.
56. Johannessen, M., T. Dale, E. Gjessing, A. Henriksen, and R. F. Wright. 1975. Acid precipitation in Norway: The regional distribution of contaminants in snow and the chemical concentration processes during snowmelt. Proc. Int. Symp. Isotopes and Impurities in Snow and Ice, Int. Assoc. Hydrol. Sci., Grenoble, France, Aug 1975. SNSF FR 4/75. SNSF-Project, 1432 Aas-NLH, Norway.
57. Johnson, A. J. 1979. Acidification of headwater streams in the New Jersey Pine Barrens. *J. of Environ. Qual.*, Vol. 3. (In press).
58. Johnson, M. G., M. F. P. Michalski, and A. E. Christie. 1970. Effects of acid mine waste on phytoplankton communities of two northern Ontario lakes. *J. Fish. Res. Bd. Can.* 27:425-444.

59. Johnson, N. M. 1979. Acid rain: neutralization within the Hubbard Brook ecosystem and regional implications. *Science* 204:497.
60. Jordan, M. and G. E. Likens. 1975. An organic carbon budget for oligotrophic lakes in New Hampshire, USA. *Verh. Int. Verein. Limnol.* 19:994-1003.
61. Keroher, G. C., et al. 1966. Lexicon of geologic names of the United States for 1936-1960. Parts 1, 2, and 3: U.S. Geol. Sur. Bull. 1200.
62. Keroher, G. C., et al. 1970. Lexicon of geologic names of the United States for 1961-1967. U.S. Geol. Sur. Bull. 1350.
63. King, D. L. 1970. The role of carbon in eutrophication. *J. Water Pollut. Control Fed.* 42:2035-2051.
64. Kramer, J. R. 1976. Geochemical and lithological factors in acid precipitation. In Proc. of the First Int. Sym. on Acid Precip. and the For. Ecosystem. U.S. Dept. Agric. For. Ser. Gen. Tech. Rep. NE-23, p. 611-618.
65. Kwiatkowski, R. E. and J. C. Roff. 1976. Effects of acidity on the phytoplankton and primary productivity of selected northern Ontario lakes. *Can. J. Bot.* 54:2546-2561.
66. Laake, M. 1976. Effekter av lav pH p roduksjon, nedbrytning og stoffkretslop i litteralsoner. SNSF-Projekt, 1432 Aas-NLH, Norway.
67. Lazrus, A. L., E. Lorange, and J. P. Lodge, Jr. 1970. Lead and other metal ions in United States precipitation. *Environ. Sci. and Tech.* 4:55-58.
68. Leivestad, H., G. Hendrey, I. P. Muniz, and E. Snekvik. 1976. p. 87-111 In Impact of Acid Precipitation on Forest and Freshwater Ecosystems in Norway (Summary Report, Phase I of SNSF-Projekt).
69. Likens, G. E. 1976. Acid precipitation. *Chem. Eng. News* 54:29-44.
70. Likens, G. E., F. H. Bormann, R. S. Pierce, J. S. Eaton and N. M. Johnson. 1977. Biogeochemistry of a Forested Ecosystem. Springer-Verlag, New York.
71. Likens, G. E., R. F. Wright, J. N. Galloway, and T. J. Butler. 1979. Acid rain. *Sci. Am.* 241:43-51.
72. Liljestrand, H. M. and J. J. Morgan. 1979. Acidic precipitation in southern California (abs.). *Am. Chem. Soc. Ann. Meet., Environ. Div.*
73. Malmer, N. 1975. SNV Res. Rep. No. PM 402E, Univ. Lund, Sweden.
74. McFee, W. W. 1980. Sensitivity of soil regions to long-term acid precipitation. U.S. Environmental Protection Agency Rep. EPA-600/3-80-013.

75. Mills, A. 1979. Dept. of Environ. Sci., Univ. of Vir., Charlottesville, Virginia. Personal communication.
76. Moss, B. 1973. The influence of environmental factors on the distribution of freshwater algae: an experimental study, II. The role of pH and the carbon dioxide bicarbonate system. J. Ecol. 61:157-177.
77. National Academy of Sciences. 1978. Sulfur Oxides.
78. National Atmospheric Deposition Program. 1979. NADP First data report: July 1978 through February 1979. Nat. Resour. Ecol. Lab., Colo. State Univ.
79. Norvell, W. A. and C. R. Frink. 1975. Bull. No. 759, Conn. Agric. Exp. Stn., New Haven.
80. Odén, S. 1976. The acidity problem--an outline of concept. WASP 6:137-166.
81. Økland, J. 1969. Distribution and ecology of the freshwater snails (Gastropoda) of Norway. Malacologia 9:143-151.
82. Økland, K. A. 1969. On the distribution and ecology of Gammarus lacustris G. O. Sars in Norway, with notes on its morphology and biology. Nor. J. Zool. 17:111.
83. Oliver, D. R. 1971. Life history of the Chironomidae. Ann. Rev. Entomol. 16:211-229.
84. Parsons, J. D. 1968. The effects of acid strip-mine effluents on the ecology of a stream. Arch. Hydrobiol. 65:25-50.
85. Parsons, J. D. 1977. Effects of acid mine wastes on aquatic ecosystems. W.A.S.P. 7:333-354.
86. Pough, F. H. 1976. Acid precipitation and embryonic mortality of spotted salamanders, Ambystoma maculatum. Science 192:68-70.
87. Rockingham County, Virginia, Soil Conservation Service. 1979. (Unpublished data).
88. Saunders, G. W. and T. A. Storch. 1971. Coupled oscillatory control mechanism in a planktonic system. Nature 230:58-60.
89. Scheider, W., J. Adamski, and M. Paylor. 1975. Reclamation of acidified lakes near Sudbury, Ontario. Ont. Min. of the Environ., Texdale, Ontario, Canada.
90. Schofield, C. L. 1975. p. 477 In Proc. First Int. Symp. on Acid Precip. and the For. Ecosystem. USDA For. Ser. Gen. Tech. Rep. NE-23.
91. Schofield, C. L. 1976. Acid precipitation: effects on fish. Ambio. 5:228.

92. Schofield, C. L. 1979. Acidification impacts on fish populations (abs.). Am. Chem. Soc. Ann. Meet., Environ. Div.
93. Shaffer and Galloway. Unpublished data, to be presented as an M.S. thesis. Dept. of Environ. Sci., Univ. of Virginia.
94. Sprules, G. W. 1975. Midsummer crustacean zooplankton communities in acid-stressed lakes. J. Fish. Res. Bd. Can. 32:389-395.
95. Sutcliffe, D. W. and T. R. Carrick. 1973. Studies on mountain streams in the English lake district, I. pH, calcium and the distribution of invertebrates in the River Duddon. Freshwater Biol. 3:437-462.
96. Watt, D. W., D. Scott, and S. Ray. 1979. Acidification and other chemical changes in Halifax County lakes after 21 years. Limnol. Oceanogr. 24:1154-1161.
97. Werner, J. J. 1966. Geology of the Vesuvius Quadrangle, Virginia. Vir. Dept. of Mineral Resour. Rep. of Investigations No. 7.
98. Wetzel, R. G. 1975. Limnology. W. B. Saunders Co. 743 pp.
99. Wiebe, W. J. and D. F. Smith. 1977. Direct measurement of dissolved organic carbon release by phytoplankton and incorporation by microheterotrophs. Marine Biol. 42:213-223.
100. Wiklander, L. 1979. EPRI Rep. No. SOA77-403.
101. Winkler, E. M. 1976. Natural dust and acid rain. p. 209-218 In: Proc. First Int. Symp. on Acid Precip. and the For. Ecosystem. USDA For. Ser. Gen. Tech. Rep. NE-23.
102. Wojcik, B. and L. Butler. 1977. Aquatic insects as indicators of stream environmental quality in northern West Virginia. Bull. 653T. West Vir Agric. and For. Exp. Stn., Morgantown, West Virginia.
103. Wright, R. F. and E. Snekvik. 1979. Verh. I. Verein. Limnol. 20:765.
104. Wright, R. F. and E. T. Gjessing. 1976. Changes in the chemical composition of lakes. Ambio 5:219.
105. Zimmerman, A. P. and H. H. Harvey. 1979. Sensitivity to acidification of waters of Ontario and neighboring states. Final rep. for Ontario Hydro. Univ. Toronto, Ontario. 48 pp.

**TECHNICAL REPORT DATA**  
(Please read Instructions on the reverse before completing)

1. REPORT NO. EPA-600/3-80-024		2.		3. RECIPIENT'S ACCESSION NO.	
4. TITLE AND SUBTITLE Geological and Hydrochemical Sensitivity to the Eastern United States to Acid Precipitation				5. REPORT DATE January 1980 issuing date	
				6. PERFORMING ORGANIZATION CODE	
7. AUTHOR(S) G. R. Hendrey, J. N. Galloway, S. A. Norton, and C. L. Schofield				8. PERFORMING ORGANIZATION REPORT NO.	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Brookhaven National Laboratory Upton, New York				10. PROGRAM ELEMENT NO.	
				11. CONTRACT/GRANT NO.  IA 79-D-X0672	
12. SPONSORING AGENCY NAME AND ADDRESS Corvallis Environmental Research Laboratory U. S. Environmental Protection Agency Corvallis, OR 97330				13. TYPE OF REPORT AND PERIOD COVERED Final, May-Oct. 1979	
				14. SPONSORING AGENCY CODE  EPA/600/02	
15. SUPPLEMENTARY NOTES					
16. ABSTRACT State geologic maps were used to produce a detailed map of the bedrock geology of the states east of the Mississippi River, on the premise that impact of acidic precipitation to aquatic ecosystems is largely a function of the buffering capacity of the underlying bedrock. Available alkalinity data for surface waters were simultaneously compiled, on a county-by-county basis in each of the 27 eastern states, and compared with the bedrock to verify the geologic definition of sensitivity. Intensive studies in the Blue Ridge Mountains in Virginia demonstrated the relationship; pH and buffer capacity of streams there varied greatly, with bedrock geology the major factor controlling stream chemistry. In North Carolina and New Hampshire, the geological map was evaluated by comparing it to county-by-county maps of current minimum alkalinity. In North Carolina, predictions were 83 percent correct, and in New Hampshire, 100 percent. Impacts of acidification on aquatic biota are discussed.					
17. KEY WORDS AND DOCUMENT ANALYSIS					
a. DESCRIPTORS		b. IDENTIFIERS/OPEN ENDED TERMS		c. COSATI Field/Group	
Acidification, precipitation, geological maps, geology, water chemistry, Aquatic biology		Acid precipitation Acid rain Atmospheric deposition		08-G 08-H	
18. DISTRIBUTION STATEMENT  Release to public		19. SECURITY CLASS (This Report) unclassified		21. NO. OF PAGES 110	
		20. SECURITY CLASS (This page) unclassified		22. PRICE	