


Research and Development



Effects of Environmental Contaminants on Cell Mediated Immunity



RESEARCH REPORTING SERIES

Research reports of the Office of Research and Development, U.S. Environmental Protection Agency, have been grouped into nine series. These nine broad categories were established to facilitate further development and application of environmental technology. Elimination of traditional grouping was consciously planned to foster technology transfer and a maximum interface in related fields. The nine series are:

1. Environmental Health Effects Research
2. Environmental Protection Technology
3. Ecological Research
4. Environmental Monitoring
5. Socioeconomic Environmental Studies
6. Scientific and Technical Assessment Reports (STAR)
7. Interagency Energy-Environment Research and Development
8. "Special" Reports
9. Miscellaneous Reports

This report has been assigned to the ENVIRONMENTAL HEALTH EFFECTS RESEARCH series. This series describes projects and studies relating to the tolerances of man for unhealthful substances or conditions. This work is generally assessed from a medical viewpoint, including physiological or psychological studies. In addition to toxicology and other medical specialities, study areas include biomedical instrumentation and health research techniques utilizing animals — but always with intended application to human health measures.

EPA-600/1-78-071
December 1978

EFFECTS OF ENVIRONMENTAL
CONTAMINANTS ON CELL MEDIATED
IMMUNITY

by

Loren D. Koller
School of Veterinary Medicine
Oregon State University
Corvallis, Oregon 97331

Grant No. R 804200

Project Officer

Richard J. Bull
Toxicological Assessment Branch
Health Effects Research Laboratory
Cincinnati, Ohio 45268

HEALTH EFFECTS RESEARCH LABORATORY
OFFICE OF RESEARCH AND DEVELOPMENT
U.S. ENVIRONMENTAL PROTECTION AGENCY
CINCINNATI, OHIO 45268

DISCLAIMER

This report has been reviewed by the Health Effects Research Laboratory U.S. Environmental Protection Agency, and approved for publication. Approval does not signify that the contents necessarily reflect the views and policies of the U.S. Environmental Protection Agency, nor does mention of trade names or commercial products constitute endorsement of recommendations for use.

FOREWORD


Man and his environment must be protected from the adverse effects of pesticides, radiation, noise and other forms of pollution, and the unwise management of solid waste. Efforts to protect the environment require a focus that recognizes the interplay between the components of our physical environment--air, water, and land. In Cincinnati, the Environmental Research Center possesses this multidisciplinary focus through programs engaged in

- ° studies on the effects of environmental contaminants on man and the biosphere, and
- ° a search for ways to prevent contamination and to recycle valuable resources.

The Health Effects Research Laboratory conducts studies to identify environmental contaminants singly or in combination, discern their relationships, and to detect, define, and quantify their health and economic effects utilizing appropriate clinical, epidemiological, toxicological, and socio-economic assessment methodologies.

The immune system is one of the body's primary defense systems. If a chemical interferes with the function of this system, an increased frequency of infectious disease may be expected in a population exposed to such chemicals. Newer scientific information also suggests that the immune system plays an important role in preventing tumor growth.

Previous work has shown that both lead and cadmium acted synergistically with certain infectious agents. The present work has established which components of the immune system are compromised by exposure to these metals and suggests an approach that may be useful for detecting similar effects by other environmental chemicals.



R.J. Garner
Director
Health Effects Research Laboratory

ABSTRACT

The effect of lead and cadmium on mouse peritoneal macrophages was investigated. Lead and cadmium given orally to mice for 10 weeks stimulated phagocytosis and increased acid phosphatase levels in peritoneal macrophages. It was concluded that these two environmental contaminants activated macrophages and, therefore, the macrophage does not contribute to the immunosuppressive activity produced by lead and cadmium. The effect of lead and cadmium on macrophage related functions is discussed.

The immune response in aged mice was studied. Mice were exposed to 0, 13, or 1,300 ppm lead in drinking water for 18 months. The immunological assays performed were mitogen (lipopolysaccharide *E. coli*, concanavalin A, and phytohemagglutinin-P) stimulation of lymphocytes; erythrocyte-antibody (EA), erythrocyte-antibody-complement (EAC), phagocytosis of macrophages; and EAC of splenic lymphocytes. As measured by the majority of these assays, the low dosage (13 ppm) of lead tended to stimulate certain immune responses (lymphocyte mitosis, EA, and EAC) while the high dosage (1,300 ppm) did not provoke any appreciable alteration. The results were interpreted by comparing data on aged mice with data on young adult mice. It was apparent from this comparison that the aged mice are naturally immunosuppressed. Therefore, the results obtained from lead-exposed aged mice were unpredictable.

B-lymphocyte response after exposure to lead and cadmium was studied. CBA/J mice were exposed to lead acetate or cadmium chloride in the drinking water for 10 weeks. The ability of bone marrow-derived lymphocytes (B cells) to form rosettes was measured using an erythrocyte-antibody-complement (EAC) assay. Fewer EAC rosettes were formed by splenic B lymphocytes from mice exposed to lead and cadmium than by B cells from control animals. The direct effect of these compounds on B cells could account in part for suppression of the humoral immune response reported in previous studies.

CBA/J mice were exposed to lead and cadmium, and the mitogen stimulation of lymphocytes was studied. Mice were given lead or cadmium in their drinking water for 10 weeks. In addition, some mice were injected with BCG. Con A, LPS, or PPD was cultured with the splenic lymphocytes. The effect of Pb, Cd, and BCG on mitogen stimulation is discussed.

This report was submitted in fulfillment of Grant No. R 8042000 by School of Veterinary Medicine, Oregon State University, under the sponsorship of the U.S. Environmental Protection Agency. This report covers the period February 1, 1976, to January 31, 1978, and work was completed as of February 28, 1978.

CONTENTS

Foreword	iii
Abstract	iv
Tables	vi
1. Introduction	1
2. Conclusions	2
3. Recommendations	5
4. Materials and Methods	6
5. Results	10
References	23
Bibliography	25

TABLES

<u>Number</u>		<u>Page</u>
1	Percent EAC Rosettes Formed by Spleen Cells from Mice Exposed to Lead or Cadmium for 10 Weeks	13
2	Percent Phagocytosis by Peritoneal Macrophages from Mice	13
3	Percent Viability of Peritoneal Macrophages from Mice	14
4	Mean Concentration of Acid Phosphatase in Peritoneal Macrophages from Mice	14
5	Comparison of Immunological Assay Values of Control Young Adult Mice to Aged Mice	15
6	Mean Body Weight of Mice	15
7	Mean Renal Concentrations of Lead and Cadmium	16
8	Lymphocyte Proliferation Response of Mice Exposed to Lead and Cadmium, <u>Macrophages Removed</u>	17
9	Lymphocyte Proliferation Response of Mice Inoculated with BCG and Exposed to Lead and Cadmium, <u>Macrophages Removed</u>	18
10	Lymphocyte Proliferation Response of Mice Inoculated with BCG and Exposed to Lead and Cadmium, <u>Macrophage Not Removed</u>	19
11	Summary of Lymphocyte Proliferation Response of Mice Exposed to Lead and Cadmium	20
12	Lymphocyte Stimulation Induced by LPS, Con A, and PHA from Mice Exposed to Lead for 18 Months	21
13	Percentage of Phagocytosis by Peritoneal Macrophages	21
14	Percentage of EA and EAC Rosettes Formed by Peritoneal Macrophages and Splenic Lymphocytes from Mice	21
15	Kidney, Liver, and Brain Concentration of Lead from Mice	22

SECTION 1

INTRODUCTION

Cadmium (Cook et al., 1975), lead (Cook et al., 1975, Hemphill et al., 1971), methylmercury (Koller, 1975), polychlorinated biphenyls (Friend & Trainer, 1970) and sulfur dioxide (Fairchild et al., 1972) are a few of the environmental contaminants that have been demonstrated to be synergistic to infectious agents in animals.

Prolonged exposure to certain environmental contaminants may suppress the immune system of a host (Koller, 1973; Koller and Thigpen, 1973; Koller et al., 1977; Miller and Zarkower, 1974). Exposure to lead has resulted in reduced circulating antibody titers to infectious disease (Koller, 1973), decreased antibody synthesis of splenic lymphocytes (Koller and Kovacic, 1974), and inhibited formation of erythrocyte-antibody-complement (EAC) (B cell) rosettes (Koller and Brauner, 1977).

Since the macrophage has an important role as an accessory cell by cooperating with T cells in aiding the response of B cells to antigens, the effect of lead and cadmium on peritoneal exudate cells of young adult mice was investigated. The cells were examined for viability, phagocytic properties and acid phosphatase content. In addition, the response of B lymphocytes, T lymphocytes, and macrophages in aged mice chronically exposed to lead was investigated.

Subpopulations of lymphoid cells have been differentiated on the basis of characteristic cell surface antigens and receptors (Gmelig-Meyling et al., 1976; Gormus and Shands, 1975). Splenic B cells were identified by detecting their surface receptors for the activated third component of complement (C'3 receptor) (Nussenzweig, 1974). The procedure used to detect B cells involved the binding of sheep erythrocytes sensitized to antibody and complement (EAC complexes) to the C'3 receptor on B cells to form rosettes (Bianco et al., 1970). In this study, we were concerned with the effect of lead and cadmium on the B cell population in murine spleens.

Mitogens can induce blast transformation in normal lymphocytes. Concanavalin A (Con A) and phytohemagglutinin (PHA) activate thymus derived (T) lymphocytes while lipopolysaccharide E. coli (LPS) stimulates bone marrow derived (B) lymphocytes. Interference of mitogen proliferation induced by Con A and PHA suggests alteration of cell mediated immune responses while LPS indicates humoral involvement. The purpose of this study was to determine if lymphocytes collected from CBA/J mice after exposure to lead or cadmium would respond abnormally to mitogen stimulation.

SECTION 2

CONCLUSIONS

The support from this grant enabled us to examine the effects that lead and cadmium have on the immune system of mice. Several important features were established which revealed that lead and cadmium are generally detrimental to the immune response in animals.

First, in previous studies, we demonstrated that both lead and cadmium inhibited antibody production and thus, diminished circulating antibody titers to antigens. The lowest dosage examined (lead - 13 ppm, cadmium - 3 ppm) resulted in significant suppression.

We next wanted to determine if lead and cadmium directly affected the B lymphocyte which produces antibody. Swiss Webster mice were exposed to lead or cadmium in the drinking water for 10 weeks. The ability of bone marrow-derived lymphocytes (B cells) to form rosettes was measured using an erythrocyte-antibody-complement (EAC) assay. Fewer EAC rosettes were formed by splenic B lymphocytes from mice exposed to lead (130 or 1,300 ppm) and cadmium (30 ppm) than B cells from control animals (Table 1). The direct effect of these compounds on B cells could account in part for suppression of the humoral immune response reported in previous studies.

Macrophages indirectly influence the humoral immune system and were investigated. Lead and cadmium given orally to mice for 10 weeks stimulated phagocytosis (Table 2) and increased acid phosphatase levels (Table 4) in peritoneal macrophages. Since lead and cadmium primarily suppress the secondary immune response, the moderately reduced primary response may be due to direct action of these compounds on B cells. The increased ability of macrophages to phagocytize and contribute to immunogenicity of the antigen may not be great enough to compensate for this inhibition. Also, other factors such as opsonins are involved in *in vivo* studies. Lead and cadmium may inhibit opsonins or other soluble factors (alpha-2-globulin, etc.) that could perhaps oppose the stimulatory effect observed in the *in vitro* experiments. Finally, the results from the B cell and macrophage studies suggest that lead and cadmium may impair the complement receptor but not the Fc receptor of lymphocytes.

Concurrent with the above investigations was a study to determine the immune response in aged mice exposed to lead. Mice were exposed to 0, 13, or 1,300 ppm lead in drinking water for 18 months. The immunological assays performed were mitogen (lipopolysaccharide *E. coli*, concanavalin A, and phytohemagglutinin-P) stimulation of lymphocytes; erythrocyte-antibody (EA), erythrocyte-antibody-complement (EAC), and phagocytosis of macrophages; an

EAC of splenic lymphocytes. As measured by the majority of these assays, the low dosage (13 ppm) of lead tended to stimulate certain immune responses (lymphocyte mitosis, EA and EAC) while the high dosage (1,300 ppm) did not provoke any appreciable alteration. The results were interpreted by comparing data on aged mice with data on young adult mice (Table 5). It was apparent from this comparison that aged mice were naturally immunosuppressed. Therefore, the results obtained from lead-exposed aged mice were unpredictable.

Mitogens can induce blast transformation in normal lymphocytes. Concanavalin A (Con A) and phytohemagglutinin (PHA) activate thymus derived (T) lymphocytes while lipopolysaccharide *E. coli* (LPS) stimulates bone marrow derived (B) lymphocytes. Interference of mitogen proliferation induced by Con A and PHA suggests alteration of cell mediated immune responses while LPS indicates humoral involvement. Few studies concerning lymphocyte blastogenesis after exposure of a host to environmental pollutants have been conducted. The purpose of the following study was to determine if lymphocytes collected from CBA mice exposed to lead or cadmium would respond abnormally to mitogen stimulation.

Mice were exposed to 3, 30, or 300 ppm cadmium as cadmium chloride or 13, 130, 1,300 ppm lead as lead acetate orally in deionized water for 70 days. The controls were given deionized water.

Lead and cadmium had little effect on lymphocyte proliferation by Con A. Counts per minute (CPM) and relative proliferation indices (RPI) were similar for lead, cadmium and control animals from non-BCG treated mice (Table 8) and from BCG treated mice without removing macrophages (Table 10). However, when lymphocytes from BCG treated animals were used after removal of macrophages, 13 ppm lead and 30 ppm cadmium resulted in a slight increase in CPM without a noticeable change in RPI (Table 9). Conversely, the low cadmium dose (3 ppm) impaired lymphocyte stimulation by Con A as seen by a decrease in CPM and RPI.

Lead reduced lymphocyte CPM and RPI by LPS from non-BCG and BCG treated mice at all levels except 13 ppm in BCG mice (Tables 8 and 9). However, when lymphocytes from BCG treated mice with macrophages were stimulated by LPS, 1,300 ppm lead enhanced CPM and RPI (Table 10). Cadmium, on the other hand, inhibited LPS stimulation of lymphocytes in most treatments at 3 ppm but resulted in significantly elevated CPM and RPI at 30 and 300 ppm (Tables 8, 9 and 10).

Response to PPD was similar to LPS. Lead in most cases decreased CPM and RPI and in some instances was as much as one-half that of the controls (Tables 1, 2 and 4). Cadmium at 3 ppm tended to inhibit CPM and RPI stimulation while a slight increase in CPM and RPI often occurred at 30 and 300 ppm cadmium dosages. In the BCG treated animals, the RPI were especially high (Table 9).

Table 11 is a summary of the mitogen studies and indicates changes for each compound and mitogen. The interpretation of proliferative responses to

mitogens has been difficult primarily due to assay variations since these measures are not stable. Furthermore, mitogens may dysfunction and vary between lots in absolute CPM which makes it difficult to combine data from identical experiments for purposes of longitudinal monitoring and analysis. Since assay variation and data analysis is a common problem encountered to measure lymphocyte function with mitogens or alloantigens by tritiated thymidine, alterations in CPM were considered significant when they were 10-15 percent larger or smaller than the control and RPI when they were greater than 1.15 or less than .85.

A summary of the work completed in the two year study revealed that lead (130 and 1,300 ppm) and cadmium (30 ppm) directly affect B lymphocytes by reducing numbers of rosettes in the EAC assay. The complement receptor site on the surface of the B cell was apparently altered by these two compounds. Lead and cadmium, on the other hand, stimulated phagocytosis and digestive functions of mouse peritoneal macrophages in vitro. However, macrophages release several soluble factors which are regulated by T cells. These factors which were not examined in this study often influence the immune response.

Finally, lead, even at low dosages, tended to inhibit mitosis of B lymphocytes while cadmium stimulated proliferation. Therefore, lead should inhibit antibody synthesis which has been demonstrated by other studies while cadmium may amplify the response. However, B lymphocytes are regulated by T cells in amount of antibody produced. T cell proliferation in this study was not altered by lead or cadmium but this technique does not directly measure helper or suppressor activity that is responsible for B cell performances.

An observation discovered during these studies is that strains of mice responded differently to lead in particular. CBA mice seem to be very resistant to lead since 1,300 ppm given orally for 18 months did not result in death. Also, antibody synthesis is not affected in these animals by lead as it is in Swiss Webster mice. We have examined other inbred strains of mice which also have not exhibited an appreciable affect to lead. Furthermore, inclusion bodies in the kidneys cannot be demonstrated by special strains as they are in other animals even though they appear to be present in H & E sections. We are currently investigating other inbred strains of mice to determine if they respond accordingly.

The smallest dose of lead used in these studies was 13.75 ppm lead in the drinking water, so the mice were ingesting about 0.069 mg of lead per day. As approximately 10 percent of ingested lead is absorbed, each mouse was actually receiving about 0.0069 mg of lead into its system each day. A significant decrease in antibody forming cells, particularly 7A, occurred at this dose. The adult human normally ingests 0.3 mg of lead per day and 2 mg per day can produce toxicity.

SECTION 3

RECOMMENDATIONS

I feel these studies have been valuable by providing data which helps to better understand the mechanism by which two environmental contaminants (lead and cadmium) affect the immune system of a host. These studies not only lead us closer to describing the complete mechanism of action but also provide dose response for each compound and assay. It is apparent that subclinical dosages of lead and cadmium alter the immune system of a host. Furthermore, these assays and techniques that were developed will greatly benefit future environmental investigations.

SECTION 4

MATERIAL AND METHODS

ANIMALS AND DIET

Swiss Webster and CBA/J mice 28 days of age were given 3, 30, or 300 ppm cadmium as cadmium chloride or 13, 130, or 1,300 ppm lead as lead acetate orally in deionized water for 70 days. CBA/J mice were also given 13, 130, or 1,300 lead orally as lead acetate in deionized water for 18 months. The controls were given deionized water. Oregon State University Rodent Chow fed to all mice contained less than 1.12 ppm lead by analysis. There were 300 mice in each group. Mice were housed in polycarbonate cages (five per cage) with stainless steel lids and cedar shavings for bedding. These mice were free of apparent infections and ectoparasites during the experiments.

PHAGOCYTOSIS

Five days before termination, mice were inoculated intraperitoneally (ip) with three ml mineral oil. Mice were killed by cervical dislocation and peritoneal exudate cells (PEC) were obtained by injecting seven ml cold Hank's balanced salt solution (HBSS, Flow Laboratories) containing heparin (10 U/ml) ip using a 20 gauge 1.5 in. disposable needle. Leaving the needle in place, the cavity was gently massaged and the fluid containing PEC was withdrawn. The collected PEC were washed three times on cold HBSS and centrifuged at 200 x g for 10 min. to remove excess oil. The washed pellet was transferred to a clean tube and contaminating red blood cells were lysed in 0.83% NH_4CL . The pellet was then diluted to six ml in culture medium composed of medium 199, 15% fetal calf serum (FCS), 100 U/ml penicillin, 100 ug/ml streptomycin, and N-2-hydroxyethyl-piperazine-N'-2-ethanesulfonic acid (HEPES) buffer, pH 7.2. Two-milliliter aliquots of the culture media were transferred to Leighton tubes (Bellco) containing a cover slip (Wheaton no. 1, 9x50mm). The cells were incubated for two hrs. at 37° C and washed once in warm phosphate-buffer saline (PBS) followed by the addition of two ml of 10% opsonized sheep red blood cells (SRBC). The preparation was incubated one hr. at 37° C.

The cover slips were then carefully removed, rinsed in two solutions of PBS, fixed in methanol, stained with Giemsa, mounted onto slides, and the cells counted. A macrophage was considered positive for phagocytosis if two or more SRBC were seen engulfed. A separate count was taken counting cells with three or more SRBC around the macrophage as in the form of a rosette. Two slides of macrophages were prepared from each mouse and 200 cells were counted per slide.

OPSONIN

To obtain optimal phagocytosis, a 10% solution of washed SRBC's was incubated for one hour with an equal amount of diluted opsonin (rabbit anti-SRBC, Cappell Laboratories, Inc.). Preliminary testing showed the optimal opsonin dilution to be 1:1600 hemagglutination units in microtiter plates. The opsonized SRBC's were washed three times in PBS and diluted to 10% in culture medium before addition to PEC monolayers.

ACID PHOSPHATASE

Acid phosphatase concentrations in macrophages were determined using a Coleman U.V. Digital Spectrophotometer. A 0.2 ml solution containing 1×10^6 macrophages was added to 1.0 ml of a buffered substrate (Boehringer Manneheim Corporation, Acid Phosphatase Test Kit, Catalog No. 15988) which contained 50 mM citrate buffer (pH 4.8), 550 u moles p-nitrophenylphosphate and 12.8 M sodium chloride. This solution was kept at 37° C for 30 min. at which time the reaction was terminated by adding 10 ml of 0.02 M sodium hydroxide. Color change was determined with the spectrophotometer at a wavelength of 405 nm by comparing treatment samples to a reagent blank.

SPLENIC LYMPHOCYTES

At termination, mice were etherized, their spleens sterily removed, scissor minced, and smashed lightly through a stainless steel screen. The spleen cells were washed three times in cold HBSS at 200 x g for ten minutes. The spleen cell suspension was then layered on top of a Ficoll-Hypaque density gradient (Ficoll, 400,000 molecular weight, Sigma; Hypaque, 50% Winthrop) with a specific gravity of 1.09 and spun at 400 x g for thirty minutes. The interface band containing mononuclear cells was removed, washed three times in cold HBSS at 200 x g for ten minutes, counted, and checked for viability using the trypan dye exclusion test. The cells were then diluted to 1×10^6 /ml in culture medium (medium 199, Flow; 10% Fetal Calf Serum, Flow; Penicillin, 100 U/ml, Squibb; Streptomycin, 100 ug/ml, Squibb; Gentamicin, 50 ug/ml; L-Glutamine 2 mM/ml, Microbiological Associates; 2 mM HEPES buffer, pH 7.2, Sigma; 2-Mercaptoethanol 5×10^{-5} M, Baker Co.).⁷ Additional mononuclear cells were plated at a cell concentration of 1×10^7 in 75 cm² tissue culture flask (Corning, #25110). The cells were incubated in culture medium at 37° C 5%CO₂ for two hours after which time the flasks were gently shaken to remove the non-adherent mononuclear cells. The non-adherent cells were spun, diluted to a concentration of 1×10^6 /ml and pipetted 0.2 ml per microtiter well (Flow). Percent macrophages before and after removal by adherence was determined by using the non-specific esterase stain procedure of Koski (IN VITRO METHODS IN CELL-MEDIATED AND TUMOR IMMUNITY, ed. Bloom and David).

MITOGEN STIMULATION

Stock mitogens were prepared in aliquots with sterile HBSS and stored at -20° C until used. These solutions were then diluted in culture medium to the concentration required. Optimal concentrations of mitogens were

determined to be: lipopolysaccharide *E. coli* (LPS), 055:B5 (Difco), 160 ug/ml; LPS 0111:B4 (Difco), 40 ug/ml; phytohemagglutinin-P (PHA-P) (Difco), 4 ug/ml; and concanavalin A (Con A) (Sigma, 1V), 2 ug/ml, and purified protein derivative (PPD) (Connought Laboratories, LTD.), 40 ug/ml. All cells, mitogen-stimulated and unstimulated, were cultured in quadruplicate for 72 hr. at 37° C in an atmosphere of 5% CO₂. At 48 hr. 0.5 uCi tritiated thymidine (6.7 Ci/mmol, New England Nuclear) was added to each well; 24 hr. later the cells were harvested using the Analog Digital Automatic Program Systems (Dedham, Massachusetts) automated system and collected on glass fiber filter paper strips (Reeve-Angel). Each filter disk was placed in a minivial (New England Nuclear) with three ml Aquasol (New England Nuclear) scintillation fluid and counted for 10 min. in a Packard Tri-Carb liquid scintillation counter (model 3375). The final values in counts per min. (CPM) were reported as the means of quadruplicate samples. The stimulation index (SI) was calculated by dividing the CPM of spleen cells with mitogen by that of spleen cells with culture medium only. A relative proliferative index (RPI) was calculated by dividing the net CPM of the test groups (mitogen stimulated CPM minus background CPM) by the net CPM of the control group (mitogen stimulated CPM minus background CPM). An RPI value greater than 1.15 was arbitrarily picked to represent an increase from the control, and a value of less than 0.85 to represent a decrease from the control. Values between 0.85 and 1.15 were considered to be of no change from the control.

MACROPHAGE EAC

The PEC's were pooled according to their respective groups and characterized by the EAC rosette assay. Sheep red blood cells were stored at 4° C in Alsever's solution for no longer than seven days. Before use, the cells were washed three times in PBS and resuspended to a final concentration of 5% Rabbit anti-sheep red blood serum (Cappel Laboratories, Inc.) was heat inactivated at 56° C for thirty min. and stored at -20° C. The hemmagglutination titer was 1:400. Autologous mouse complement serum was obtained from blood collected by cardiac puncture and stored at -70° C.

Sheep erythrocytes were sensitized with an equal volume of a 1:800 dilution of rabbit anti-sheep red blood cell serum for thirty min. at 37° C. The EA complex was an equal volume of a 1:10 dilution of mouse complement serum for thirty min. at 37° C. After two washes in VBS, the EAC complex was resuspended in HBSS, eight times the initial volume of SRBC's. A suspension of EA was prepared in a similar manner.

The PEC's (1.5×10^6 in 15 ml HBSS) were incubated with 0.5 ml EAC at 37° C for five min. After centrifugation at 200 x five min., the cells were incubated in ice for one hr. without removing the supernatant (Jondal et al., 1972). After gentle resuspension, an aliquot of cells was mixed with an equal volume of crystal violet solution (1 mg/ml in minimal essential medium with 10% FCS). Two hundred cells were examined microscopically in a hemacytometer. Only stained cells with three or more bound erythrocytes were considered as EAC rosettes. The PEC's were similarly assayed with EA.

LYMPHOCYTE EAC

Spleen lymphocyte cells derived from a Ficoll-Hypaque gradient were similarly characterized by the EAC rosette assay and the direct B-cell fluorescent antibody technique. Fluorescein-conjugated rabbit anti-mouse IgG (FITC-Rxmg) (Cappel Laboratories, Inc.) was filtered through a 0.45 μ m Millipore filter, stored at -70° C. An equal volume of spleen lymphocyte cells (1.5×10^6 in 0.05 ml HBSS with 0.1% NaN_3) was incubated with the FITC-Rxmg for thirty min. on ice. The cells were washed twice by centrifugation at $200 \times g$ at 4° C for six min. on a gradient of 2.5 ml HBSS with 30% FCS and 0.1% NaN_3 . After resuspension in 0.05 ml HBSS with 0.1% NaN_3 , the cells were stored on ice until they were examined by fluorescent microscopy.

BCG

Six weeks before termination 75 of the lead exposed, 75 of the cadmium exposed and 25 control mice were injected subcutaneously with BCG (Bacillus-Calmette Guerin; University of Illinois) diluted to 1 mg with equal amounts of CFA (Complete Freund's Adjuvant; Difco). Backs of mice were shaved and a total of four injection sites, 0.125 ml/site, or a total of 0.5 ml was injected. One week before termination the BCG treated mice received an additional injection of 0.25 mg BCG intraperitoneally.

TISSUE RESIDUES

The brain, liver, and one kidney from each mouse was collected at necropsy and stored at -70° C for lead and cadmium analysis. These tissues were analyzed for content by atomic absorption spectrophotometry using a microcarbon furnace.

PATHOLOGY

Sections of liver, kidney, brain, lungs, heart, urinary bladder, muscle, stomach, intestine, and spleen were collected at necropsy, fixed in 10% buffered formalin, and stained with Harris' hematoxylin and eosin (H&E). In addition, kidney sections were stained by the periodic acid-Schiff (PAS), Ziehl-Neelsen acid-fast, and Masson Trichrome methods.

SECTION 5

RESULTS

The percent of EAC rosette formation by spleen cells from mice exposed to lead and cadmium for 10 weeks was generally less than rosette formation by control animals (Table 1). The number of EAC rosettes was significantly less in animals which received 130 or 1,300 lead or 30 ppm cadmium. The impaired rosetting was not due to toxicity of lead and cadmium since the percent viability was similar for treated and control groups.

The inhibition of EAC rosettes produced by cadmium at 30 ppm but not 300 ppm demonstrates the variable response of animals to different dosages of cadmium which we so often observe in our laboratory. A similar response occurred with acid phosphatase concentrations in macrophages that were collected from cadmium-exposed mice (Koller and Roan, 1977).

Antibody responses to many antigens require cooperation between at least two types of lymphocytes for optimal expression. One cell type is thymus derived (T cells). T cells may amplify, help, or suppress B cells as well as function as cytotoxic cells. T cells do not produce antibody. The other cell type is bone marrow derived (B cell) that mature independently of thymic influence. B cells differentiate into antibody-producing cells and are often influenced by T cells. A third cell type is the macrophage, which is important as an accessory cell in cooperating with T cells and aiding B cells in response to antigens.

Humoral antibody response to sheep red blood cells (SRBC) requires cooperation between B and T cells (Elliott and Haskill, 1974). Recent studies demonstrated suppressed antibody synthesis to SRBC in mice exposed to lead or cadmium (Koller and Kovacic, 1974; Koller et al., 1974). It was postulated that the T helper cell may be the primary target for these two environmental contaminants since the secondary immune response was most severely inhibited. However, in this study, we demonstrated that these two compounds also affected the B cells. B cells from animals exposed to lead and cadmium were inhibited in rosette formation. This could account for the diminished antibody response that was reported to occur during primary immunity after exposure to lead and cadmium (Koller and Kovacic, 1974; Koller et al., 1975). Recently it was reported that lead and cadmium fed to mice stimulated both ingestion and digestive functions of peritoneal macrophages (Koller and Roan, 1977). Studies of the effects of these contaminants on T cells are currently in progress.

Only B cells formed rosettes by the EAC technique. This was confirmed by examining nylon wool-separated T and B cells with fluorescein-conjugated

antiglobulin. All rosetted cells fluoresced with antiglobulin, indicating that they were B cells. T cells do not react with antiglobulin, nor do they form rosettes by the EAC technique. The complement receptor site on the surface of the B cell was apparently altered by lead and cadmium since lymphocyte viability and percentage T and B cells were similar in all groups.

Macrophages from CBA and Swiss Webster mice exposed to lead and cadmium were stimulated to phagocytize SRBC. Phagocytosis increased as the dose increased in the Swiss Webster mice (Table 2) while the greatest response in the CBA mice was at the high dose of both lead and cadmium and the low dose of lead. When macrophages were examined for SRBC rosettes, the response was similar for cells prepared from both treated and control animals.

Viability for macrophages was comparable for treated and control groups (Table 3). Cell viability was considered excellent for most groups and four of the six exposure groups had 96 to 99% viability.

Acid phosphatase levels were increased in macrophages of each cadmium exposure and the two highest lead exposures (Table 4). The greatest concentration of the enzyme occurred in macrophages from the medium dosages of both lead and cadmium.

The body weight of mice that received 300 ppm cadmium were significantly lower than those of animals in the other treated and control groups (Table 6). Renal concentrations of lead and cadmium increased significantly as the metal dosage increased (Table 7).

Lymphocytes from control and lead-exposed mice cultured in the presence of LPS 055:B5, LPS 0111:B4, Con A and PHA-P were classified according to the SI values (Table 12). The SI values for LPS 055:B5, LPS 0111:B4, and Con A were markedly elevated in mice exposed to 13 ppm lead compared to controls or to mice given 1,300 ppm lead. However, the SI of PHA-P was elevated in the mice exposed to 1,300 ppm lead compared to controls or to mice exposed to 13 ppm lead.

The ability of macrophages to phagocytize SRBC's was inhibited in mice exposed to 13 ppm lead (Table 13). However, the EA and EAC values for macrophages (Table 14) and lymphocytes were slightly elevated in the same 13 ppm lead group where as they were suppressed in the 1,300 ppm group compared to controls. Lymphocyte viability before culturing for mitogen stimulation and the EAC test was generally 85-95%.

Tissue residues of lead increased with an increase in dosage (Table 15). The kidneys contained the largest quantities of lead in both exposures; larger quantities of lead were found in the brain than in the liver in the 13 ppm group, while the converse was found in the 1,300 ppm group.

Histopathology disclosed a wide zone of necrosis at the cortical medullary junction in the kidneys of mice exposed to 1,300 ppm lead. The principle lesion was necrosis of tubular epithelial cells. Within these necrotic areas there were frequent granulomas composed of large mononuclear cells, occasional

multinucleated giant cells, and fibrous connective tissue. Many intranuclear inclusion-like bodies were identified in H&E sections, but they did not stain well by the acidfast method.

The lesions in mice exposed to 13 ppm lead were much less severe and were characterized by necrosis of the tubular epithelial cells at the cortical medullary junctions. Granulomas were infrequent in these lesions. The PAS stain demonstrated thickening of the glomerular basement membranes and Bowman's capsule in the lead-exposed animals. Masson's trichrome stain revealed a slight increase of fibrous connective tissue in the necrotic areas of the kidneys of many of the lead-exposed mice. Spontaneous tumors were found in four lead-exposed mice and in one control animal.

Lead and cadmium had little effect on lymphocyte proliferation by Con A. Counts per minute (CPM) and relative proliferation indexes (RPI) were similar for lead, cadmium and control animals when lymphocytes were examined from non-BCG treated mice (Table 8) and from BCG treated mice without removing macrophages (Table 10) in the lymphocyte suspensions. However, when lymphocytes from BCG treated animals were used after removal of macrophages, 13 ppm lead and 30 ppm cadmium resulted in a slight increase in CPM without a noticeable change in RPI (Table 9). Conversely the low cadmium dose (3 ppm) impaired lymphocyte stimulation by Con A reflected by both decreased CPM and RPI.

Lead reduced lymphocyte CPM and RPI by LPS from non-BCG and BCG treated mice at all levels except 13 ppm in BCG mice (Tables 8 and 9). However, when lymphocyte from BCG treated mice with macrophages were stimulated by LPS, 1,300 ppm lead enhanced CPM and RPI (Table 10). Cadmium, on the other hand, inhibited LPS stimulation of lymphocytes in most treatments at 3 ppm but resulted in significantly elevated CPM and RPI at 30 and 300 ppm (Tables 8, 9, and 10).

Response to PPD was similar to LPS. Lead in most cases decreased CPM and RPI and in some instances were as much as one half that of the controls (Tables 8, 9, and 10). Cadmium at 3 ppm tended to inhibit CPM and RPI of PPD stimulation while a slight increase in CPM and RPI often occurred at 30 and 300 ppm cadmium dosages. In the BCG treated animals, the RPI were especially high (Table 9).

Cell viability averaged 87 percent from non-BCG mice and slightly less (84%) for the BCG treated animals. Most of the macrophages were removed by the Ficoll-Hypaque density gradient but for the non-BCG and BCG without macrophate groups, the cells were plated in plastic flasks for two hours. Two to three percent were present after plating. However, in the 300 ppm cadmium group, eight percent macrophages were present before plating and three percent after.

TABLE 1. PERCENT EAC ROSETTES FORMED BY
SPLEEN CELLS FROM MICE EXPOSED
TO LEAD OR CADMIUM FOR 10 WEEKS +

Dose (ppm)		EAC (%)	
Lead	Cadmium	Lead	Cadmium
0	0	53.5	45.1
13	3	46.5	42.9
130	30	41.5* ¹	30.9* ²
1,300	300	41.3* ¹	42.7

+ = 25 mice per group

* = Significant $P < 0.05$ (one way analysis of variance)

¹ = LSD - 9.64 ² = LSD - 13.63

TABLE 2. PERCENT PHAGOCYTOSIS BY PERITONEAL
MACROPHAGES FROM MICE EXPOSED TO
LEAD OR CADMIUM FOR 10 WEEKS

Dose (ppm)		Phagocytosis (%)			
Lead	Cadmium	CBA		SW	
		Lead	Cadmium	Lead	Cadmium
0	0	56	54	43	43
13	3	67	56	49	38
130	30	54	56	57*	45
1,300	300	72*	63*	62**	59*

* = Significant at $P < 0.05$ (t test)

** = Significant at $P < 0.01$ (t test)

TABLE 3. PERCENT VIABILITY OF PERITONEAL MACROPHAGES FROM MICE EXPOSED TO LEAD OR CADMIUM FOR 10 WEEKS

Dose (ppm)		Macrophage viability (%) ^a	
Lead	Cadmium	Lead	Cadmium
0	0	94	93
13	3	96	89
130	30	96	99
1,300	300	89	96

^a = Ten mice in each exposure group; 2 samples per mouse

TABLE 4. MEAN CONCENTRATION OF ACID PHOSPHATASE IN PERITONEAL MACROPHAGES FROM MICE EXPOSED TO LEAD OR CADMIUM FOR 10 WEEKS

Dose (ppm)		Concentration acid phosphatase ^a	
Lead	Cadmium	Lead	Cadmium
0	0	18.37	21.72
13	3	12.81	28.12
130	30	57.63**	34.80*
1,300	300	24.50	28.40

^a = SI units

* = Significant at $P < 0.05$ (t test)

** = Significant at $P < 0.05$ (t test)

TABLE 5. COMPARISON OF IMMUNOLOGICAL ASSAY VALUES OF CONTROL YOUNG ADULT MICE TO AGED MICE^a

Immunological assay	Young adult mice ^b	Aged mice
Phagocytosis by macrophages (%)	56	40
EAC lymphocyte rosettes (%)	54	43
Lymphocyte stimulation index		
LPS 055:B5	27.5	6.9
LPS 0111:B4	17.9	4.6
Con A	32.6	9.3
PHA-P	3.2	1.8

^aErythrocyte-antibody-complement (EAC)
Lipopolysaccharide *E. coli* (LPS)
Concanavalin A (Con A)
Phytohemagglutinin-P (PHA-P)

^bYoung adult mice were 14 weeks old

TABLE 6. MEAN BODY WEIGHTS OF MICE EXPOSED TO LEAD OR CADMIUM FOR 10 WEEKS

Dose (ppm)		Body weight (g) ^a	
Lead	Cadmium	Lead	Cadmium
0	0	26	27
13	3	27	25
130	30	27	26
1,300	300	24	18**

^a = Ten mice each exposure group

** = Significant at P < 0.01

TABLE 7. MEAN RENAL CONCENTRATIONS OF LEAD AND CADMIUM FROM MICE EXPOSED FOR 10 WEEKS

Lead	Dose (ppm)		Renal concentration (ppm) ^{a,b}	
	Lead	Cadmium	Lead	Cadmium
0	0	0	0.07	0.31
13	3	3	0.35	1.07
130	30	30	1.29	8.35
1,300	300	300	8.10	44.80

^a Wet weight

^b Ten mice in each exposure group

TABLE 8. LYMPHOCYTE PROLIFERATION RESPONSE OF MICE EXPOSED TO LEAD AND CADMIUM, MACROPHAGES WERE REMOVED FROM THE LYMPHOCYTE SUSPENSIONS BY PLATING

Dosage	CPM ¹	Con A	RPI ³	CPM ¹	LPS	RPI ³	CPM ¹	PPD	RPI ³
	(CV)	SI ²		(CV)	SI ²		(CV)	(CV)	
Control	108,000 ⁴ (.36)	47 (.58)		38,000 (.49)	15 (.27)		27,500 (.08)	7 (.20)	
Lead									
13	100,000 (.23)	79 (.66)	1.01 (.40)	25,000 (.08)	19 (.53)	0.80 (.56)	13,000 (.24)	12 (.71)	0.52 (.41)
130	105,000 (.17)	64 (.39)	1.08 (.42)	28,000 (.22)	17 (.47)	0.88 (.52)	19,500 (.33)	13 (.62)	0.77 (.49)
1300	119,000 (.25)	52 (.35)	1.18* (.38)	31,000 (.65)	11 (.23)	0.95 (.64)	18,000 (.86)	6 (.13)	0.66 (.93)
Cadmium									
3	102,000 (.23)	52 (.18)	1.04 (.45)	26,000 (.12)	13 (.04)	0.84 (.61)	*14,500 (.05)	7 (.20)	*0.51 (.14)
30	109,000 (.12)	29 (.35)	1.10 (.37)	55,000 (.28)	14 (.19)	1.61 (.34)	32,500 (.11)	6.5 (.33)	1.15 (.06)
300	105,000 (.07)	24 (.19)	1.05 (.38)	*64,000 (.14)	15 (.17)	2.17 (.71)	32,500 (.11)	.9 (.04)	1.22 (.22)
	(.20)	(.39)	(.40)	(.28)	(.27)	(.56)	(.25)	(.32)	(.38)

¹ = mean counts per minute

² = mean stimulation index

³ = mean relative proliferative index

⁴ = coefficient of variance

* = Significantly different from control population at P < 0.05

TABLE 9. LYMPHOCYTE PROLIFERATION RESPONSE OF MICE INOCULATED WITH BCG AND EXPOSED TO LEAD AND CADMIUM. MACROPHAGES WERE REMOVED FROM THE LYMPHOCYTE SUSPENSIONS BY PLATING

Dosage (ppm)	CPM ¹ (CV)	Con A		CPM ¹ (CV)	LPS		CPM ¹ (CV)	PPD	
		SI ² (CV)	RPI ³ (CV)		SI ² (CV)	RPI ³ (CV)			
Control	125,000 ⁴ (.18)	43 (.53)		52,500 (.48)	15 (.14)		9,000 (.56)	2.5 (.40)	
Lead									
13	139,000 (.18)	38 (.33)	1.13 (.22)	54,000 (.36)	14 (.12)	1.12 (.38)	11,000 (.92)	2.4 (.44)	1.09 (.52)
130	133,000 (.26)	52 (.27)	1.09 (.29)	41,000 (.26)	16 (.21)	0.98 (.59)	5,000 (.74)	1.8 (.54)	0.54 (1.42)
1300	126,000 (.15)	56 (.30)	1.03 (.17)	37,000 (.34)	15 (.12)	0.77 (.33)	4,000 (.30)	1.8 (.12)	0.40 (.36)
Cadmium									
3	108,000 (.28)	40 (.23)	0.86 (.17)	49,000 (.29)	19 (.36)	1.16 (.56)	7,000 (.68)	2.7 (.55)	0.93 (.97)
30	139,000 (.16)	33 (.67)	1.07 (.08)	67,000 (.44)	15 (.15)	1.31 (.21)	12,000 (.50)	2.5 (.42)	1.34 (.08)
300	128,000 (.19)	39 (.50)	1.03 (.16)	61,000 (.30)	17 (.24)	1.44 (.52)	9,000 (.29)	2.4 (.35)	1.52 (.96)
	(.20)	(.40)	(.19)	(.35)	(.19)	(.43)	(.57)	(.40)	(.72)

- 1 = mean counts per minute
 2 = mean stimulation index
 3 = mean relative proliferative index
 4 = coefficient of variance

TABLE 10. LYMPHOCYTE PROLIFERATION RESPONSE OF MICE INOCULATED WITH BCG AND EXPOSED TO LEAD AND CADMIUM. MACROPHAGES WERE NOT REMOVED FROM THE LYMPHOCYTE SUSPENSION BY PLATING

Dosage (ppm)	CPM ¹ (CV)	Con A	RPI ³ (CV)	CPM ¹ (CV)	LPS	RPI ³ (CV)	CPM ¹ (CV)	PPD	RPI ³ (CV)
		SI ² (CV)			SI ² (CV)			SI ² (CV)	
Control	169,000 ⁴ (.07)	40 (.29)		58,000 (.35)	13 (.12)		11,000 (.30)	2.6 (.07)	
Lead									
13	170,000 (.11)	41 (.07)	1.01 (.12)	53,000 (.28)	13 (.16)	0.93 (.17)	11,000 (.49)	2.5 (.36)	0.85 (.40)
130	181,000 (.16)	47 (.12)	1.07 (.13)	54,000 (.04)	15 (.25)	1.00 (.31)	5,500 (.63)	2.1 (.03)	0.68 (.47)
1300	166,000 (.07)	36 (.21)	0.98 (.13)	67,000 (.29)	14 (.12)	1.17 (.06)	13,000 (.44)	2.8 (.31)	1.16 (.22)
Cadmium									
3	172,000 (.10)	38 (.32)	1.02 (.11)	53,000 (.33)	11 (.19)	0.90 (.15)	10,000 (.45)	2.1 (.24)	0.77 (.42)
30	178,000 (.12)	30 (.99)	1.03 (.06)	76,000 (.45)	10 (.25)	1.29 (.45)	12,000 (.54)	2.3 (.27)	1.12 (.12)
300	177,000 (.10)	25 (.14)	1.03 (.10)	78,000 (.13)	11 (.14)	1.42 (.33)	17,000 (.50)	2.3 (.50)	1.49 (1.02)
	(.10)	(.31)	(.11)	(.27)	(.18)	(.25)	(.48)	(.25)	(.44)

- 1 = mean counts per minute
 2 = mean stimulation index
 3 = mean relative proliferative index
 4 = coefficient of variance

TABLE 11. SUMMARY OF LYMPHOCYTE PROLIFERATION RESPONSE OF MICE EXPOSED TO LEAD AND CADMIUM. GROUP 2 AND 3 IN ADDITION WERE INOCULATED WITH BCG WHILE GROUP 2 ALSO HAD MACROPHAGES REMOVED FROM THE LYMPHOCYTE SUSPENSION BY PLATING

Lead	Cadmium	Concanavalin A					
		Non-BCG		BCG		BCG w/Macs	
		CPM	RPI	CPM	RPI	CPM	RPI
13		-	-	-	-	-	-
130		-	-	-	-	-	-
1300		-	I	-	-	-	-
	3	-	-	-	-	-	-
	30	-	-	-	-	-	-
	300	-	-	-	-	-	-
Lipopolysaccharide							
13		D	D	-	-	-	-
130		D	-	D	-	-	-
1300		D	-	D	D	I	I
	3	D	D	-	I	-	-
	30	I	I	I	I	I	I
	300	I	I	I	I	I	I
Purified Protein Derivative							
13		D	D	-	-	-	D
130		D	D	D	D	D	D
1300		D	D	D	D	I	I
	3	D	D	D	-	-	D
	30	I	-	I	I	-	-
	300	I	I	-	I	I	I

I = increase
D = decrease
- = no significant change

TABLE 12. LYMPHOCYTE STIMULATION INDUCED BY LPS, CON A, AND PHA FROM MICE EXPOSED TO LEAD FOR 18 MONTHS^a

Lead (ppm)	Stimulation index			
	LPS 055:B5	LPS 0111:B4	Con A	PHA-P
0	6.9	4.6	9.3	1.8
13	11.1	9.0 ^b	12.2	2.0
1,300	7.9	5.7	7.1	2.7

^a Thirty mice per group; LPS, lipopolysacchride *E. coli*; Con A, concanavalin A; PHA-P, phytohemagglutinin-P

^b Significant at $P < 0.05$ (one-way analysis of variance; LSD = 4.27)

TABLE 13. PERCENTAGE OF PHAGOCYTOSIS BY PERITONEAL MACROPHAGES FROM MICE EXPOSED TO LEAD FOR 18 MONTHS

Lead (ppm)	Phagocytosis (%)
0	40
13	32 ^b
1,300	42

^a Thirty mice per group

^b Significant at $P < 0.05$ (one-way analysis of variance; LSD = 5.86)

TABLE 14. PERCENTAGE OF EA AND EAC ROSETTES FORMED BY PERITONEAL MACROPHAGES AND SPLENIC LYMPHOCYTES FROM MICE EXPOSED TO LEAD FOR 18 MONTHS

Lead (ppm)	Macrophages		Lymphocytes
	EA (%)	EAC (%)	EAC (%)
0	53	57	44
13	55	59	45
1300	50	52	39

^a EA, erythrocyte-antibody; EAC, erythrocyte-antibody-complement

TABLE 15. KIDNEY, LIVER, AND BRAIN CONCENTRATIONS OF LEAD FROM MICE EXPOSED FOR 18 MONTHS

Lead (ppm)	Mean lead (ppm)		
	Brain	Liver	Kidney
0	0.10	0.10	0.14
13	0.51	0.25	0.97
1300	5.94	7.32	17.79

^a Wet weight

REFERENCES

- Bianco, C., Patrick, R., and Nussenzweig, V. (1970). A population of lymphocytes bearing a membrane receptor or antigen-antibody complement complexes. *J. Exp. Med.* 132, 702-720.
- Cook, Z.A., Hoffmann, E.O., and DiLuiz, N.R. (1975). Influence of lead and cadmium on the susceptibility of rats to bacterial challenge. *Proc. Soc. Exp. Med.* 150, 741.
- Elliott, B.E., and Haskill, J.S. (1974). Separation of T effector cells in humoral and cellular immunity. *Nature (London)* 252, 607-608.
- Fairchild, G.S., Roan, J., and McCarroll, J. (1972). Atmospheric pollutants and the pathogenesis of viral respiratory infection. Sulfur dioxide and influenza infection in mice. *Arch. Environ. Health* 25, 174.
- Friend, M., and Trainer, D.O. (1970). Polychlorinated biphenyl: interaction with duck hepatitis virus. *Science* 170, 1314.
- Gmelig-Meyling, F., Van Der Ham, F., and Ballieux, R.E. (1976). The use of frozen stored sheep erythrocytes for demonstrating lymphocyte membrane markers. *J. Immunol. Methods* 11, 257-264.
- Gormus, B.J., and Shands, J.W. (1975). Capping of the lymphocyte C3 receptor and temperature-dependent loss of C3 rosettes. *J. Immunol.* 114, 1221-1225.
- Hemphill, F.E., and Kaeberle, M.L. (1971). Lead suppression of mouse resistance to *Salmonella typhimurium*. *Science* 172, 1031.
- Koller, L.D. (1973). Immunosuppression produced by lead, cadmium and mercury. *Am. J. Vet. Res.* 34, 1457.
- Koller, L.D., and Thigpen, J.E. (1973). Reduction of antibody in pseudo-rabies virus on polychlorinated biphenyl-exposed rabbits. *Am. J. Vet. Res.* 34, 1605-1606.
- Koller, L.D., and Kovacic, S. (1974). Decreased antibody formation in mice exposed to lead. *Nature* 250, 148.
- Koller, L.D., Exon, J.H., and Roan, J. (1975). Antibody suppression by cadmium. *Arch. Environ. Health* 30, 598.

- Koller, L.D. (1975). Methylmercury: effect on oncogenic and nononcogenic viruses in mice. *Am. J. Vet. Res.* 36, 1501.
- Koller, L.D., Exon, J.H., and Roan, J. (1977). Effects of lead and cadmium on mouse peritoneal macrophages. *J. Reticuloendothel. Soc.* 21(1), 7-12.
- Koller, L.D., and Brauner, J.A. (1977). Decreased B-lymphocyte response after exposure to lead and cadmium. *Toxicol. Appl. Pharmacol.* 42, 621-624.
- Koller, L.D., Exon, J.H., and Arbogast, B. (1977). Methylmercury: Effect on serum enzymes and humoral antibody. *J. Toxicol. Environ. Health* 2, 1115-1123.
- Miller, S.D., and Zarkower, A. (1974). Alterations of murine immunological responses after silica dust inhalation. *J. Immunol.* 113, 1533-1543.
- Nussenzweig, V. (1974). *Advances in Immunology*, Vol. 19, p. 217. Academic Press, New York and London.

BIBLIOGRAPHY

- Bar-Eli, M., and Gallily, R. (1975). The effect of macrophage hydrolytic enzyme levels on the uptake and degradation of antigen and immune complexes. *J. Reticuloendothel. Soc.* 18, 317.
- Bevan, M.J., and Cohn, M. (1975). Cytotoxic effects of antigens and mitogen induced T cells on various targets. *J. Immunol.* 114, 559-565.
- Blumenstock, F., Saba, T.M., Weber, P., and Cho, E. (1976). Purification and biochemical characterization of a macrophage stimulating alpha-2-globulin opsonic protein. *J. Reticuloendothel. Soc.* 19, 157.
- Boyum, A. (1968). Separation of leukocytes from blood and bone marrow. *Scand. J. Lab. Invest.* 21, 97.
- Cohn, Z.A. (1968). The structure and function of monocytes and macrophages. *Adv. Immunol.* 9, 163.
- Elliott, B.E., and Haskill, J.S. (1974). Separation of T effector cells in humoral and cellular immunity. *Nature* 252, 607.
- Fishel, C.W., Pross, S.H., and Klein, T.W. (1976). The effect of *Bordetella pertussis* on rosette-forming cells present in the peritoneal fluids of mice. *J. Reticuloendothel. Soc.* 19, 97.
- Gatti, P.A., and Good, R.A. (1971). Occurrence of malignancy in immunodeficiency disease. *Cancer* 28, 89-98.
- Gillette, R.W., and Swanson, M.H. (1974). Kinetic studies of macrophages. V. Effect of antigen and adjuvant stimulation. *J. Reticuloendothel. Soc.* 15, 31.
- Herberman, R.B. (1974). Cell-mediated immunity to tumor cells. *Adv. Cancer Res.* 19, 107-263.
- Hirschhorn, R. (1974). Lysosome mechanism in the inflammatory process. In *Inflammatory Process*, (B.W. Zweifach, ed.) 2nd edition, p. 259. Academic Press, New York.
- Jondal, M., Holm, G., and Wigzell, H. (1972). Surface markers on human T and B lymphocytes. 1. A large population of lymphocytes forming non-immune rosettes with sheep red blood cells. *J. Exp. Med.* 136, 207-215.

- Karl, L., Chvapli, M., and Zukoski, C.F. (1973). Effect of zinc on the viability and phagocytic capacity of peritoneal macrophages. *Proc. Soc. Exp. Med.* 142, 1123.
- Koller, L.D. (1973). Immunosuppression produced by lead, cadmium and mercury. *Am. J. Vet. Res.* 34, 1457-1458.
- Koller, L.D., Exon, J.H., and Roan, J.G. (1975). Antibody suppression by cadmium. *Arch. Environ. Health* 30, 598-601.
- Koller, L.D., Exon, J.H., and Roan, J.G. (1976). Humoral antibody response in mice after single dose exposure to lead or cadmium. *Proc. Soc. Exp. Biol. Med.* 151, 339.
- Koller, L.D., and Kovacic, S. (1974). Decreased antibody formation in mice exposed to lead. *Nature (London)* 250, 148-150.
- Koller, L.D., and Roan, J.G. (1977). Effects of lead and cadmium on mouse peritoneal macrophages. *J. Reticuloendothel. Soc.* 21, 7-12.
- Mitchison, N.A. (1969). The immunogenic capacity of antigen taken up by peritoneal exudate cells. *Immunology* 16, 1.
- Page, R.D., Davies, R., and Allison, A.C. (1974). Participation of mononuclear phagocytes in chronic inflammatory diseases. *J. Reticuloendothel. Soc.* 15, 413.
- Segre, D., and Segre, M. (1976). Humoral immunity in aged mice. Increased suppressor T cell activity in immunologically deficient old mice. *J. Immunol.* 116, 735-738.
- Shou, L., Schwartz, S.A., and Good, R.A. (1976). Suppressor cell activity after concanavalin A treatment of lymphocytes from normal donors. *J. Exp. Med.* 143, 1100-1110.
- Soulami, R.L., and Bradfield, J.W.B. (1974). The recovery of hepatic phagocytosis after blockade of Kupffer cells. *J. Reticuloendothel. Soc.* 16, 75.
- Tizard, I.R., Holmes, W.L., and Parappally, N.P. (1974). Phagocytosis of sheep erythrocytes by macrophages: A study of the attachment phase by scanning electron microscopy. *J. Reticuloendothel. Soc.* 15, 225.
- Treagan, L. (1975). Metals and the immune response. *Chem. Pathol. Pharmacol.* 12, 189.
- Trejo, R.A., DiLuizo, N.R., Loose, L.D., and Hoffman, E. (1972). Reticuloendothelial and hepatic functional alterations following lead acetate administration. *Exp. Mol. Pathol.* 17, 145.

Van Ginckel, R.F., and Hoebeke, J. (1975). Carbon clearance enhancing factor in serum from levamisole treated mice. J. Reticuloendothel. Soc. 17, 65.

TECHNICAL REPORT DATA
(Please read Instructions on the reverse before completing)

1. REPORT NO. EPA-600/1-78-071	2.	3. RECIPIENT'S ACCESSION NO.
4. TITLE AND SUBTITLE Effects of Environmental Contaminants on Cell Mediated Immunity	5. REPORT DATE December 1978 issuing date	6. PERFORMING ORGANIZATION CODE
	8. PERFORMING ORGANIZATION REPORT NO.	
7. AUTHOR(S) Loren D. Koller	10. PROGRAM ELEMENT NO. ICC 614	
9. PERFORMING ORGANIZATION NAME AND ADDRESS School of Veterinary Medicine Oregon State University Corvallis, Oregon 97331	11. CONTRACT/GRANT NO. R804200	
	13. TYPE OF REPORT AND PERIOD COVERED Final Report	
12. SPONSORING AGENCY NAME AND ADDRESS Health Effects Research Laboratory - Cinn, OH Office of Research and Development U.S. Environmental Protection Agency Cincinnati, Ohio 45268	14. SPONSORING AGENCY CODE EPA/600/10	

15. SUPPLEMENTARY NOTES

16. ABSTRACT

The effect of lead and cadmium on cell-mediated immunity was studied in peritoneal macrophages, B-, and T-lymphocytes of mice. Lead and cadmium were administered in drinking water for 10 weeks in short-term experiments and up to 18 months to deal with immune responses in aged mice.

Lead and cadmium both tended to stimulate phagocytosis in peritoneal macrophages. Consequently, depressed humoral immune response could not be explained on the basis of an effect on the macrophage. The splenic B-lymphocyte response was depressed by both lead and cadmium treatment. The direct effect of these metals on B cells could account at least in part, for the suppression of the humoral immune response reported in previous studies.

In long-term studies in aged mice low doses of lead (13 mg/l) tended to stimulate certain immune responses. Results obtained with higher doses (up to 1300 mg/l) were complicated by a natural immunosuppression in aged mice. As a consequence, no significant alterations were observed with high doses and the impact of Pb on the immune system in the long term cannot be predicted on the basis of this limited experimentation.

17. KEY WORDS AND DOCUMENT ANALYSIS

a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
Lead (metal) Cadmium Lymphocytes Toxicity Mice	Immune response Peritoneal macrophage	13B

18. DISTRIBUTION STATEMENT Release to public.	19. SECURITY CLASS (This Report) Unclassified	21. NO. OF PAGES 34
	20. SECURITY CLASS (This page) Unclassified	22. PRICE