

EPA-600/4-76-024
May 1976

Environmental Monitoring Series

EFFECT OF TEMPERATURE ON STABILITY OF SULFUR DIOXIDE SAMPLES COLLECTED BY THE FEDERAL REFERENCE METHOD



**Environmental Monitoring and Support Laboratory
Office of Research and Development
U.S. Environmental Protection Agency
Research Triangle Park, North Carolina 27711**

RESEARCH REPORTING SERIES

Research reports of the Office of Research and Development, U.S. Environmental Protection Agency, have been grouped into five series. These five broad categories were established to facilitate further development and application of environmental technology. Elimination of traditional grouping was consciously planned to foster technology transfer and a maximum interface in related fields. The five series are:

1. Environmental Health Effects Research
2. Environmental Protection Technology
3. Ecological Research
4. Environmental Monitoring
5. Socioeconomic Environmental Studies

This report has been assigned to the ENVIRONMENTAL MONITORING series. This series describes research conducted to develop new or improved methods and instrumentation for the identification and quantification of environmental pollutants at the lowest conceivably significant concentrations. It also includes studies to determine the ambient concentrations of pollutants in the environment and/or the variance of pollutants as a function of time or meteorological factors.

EFFECT OF TEMPERATURE
ON
STABILITY OF SULFUR DIOXIDE SAMPLES COLLECTED
BY THE FEDERAL REFERENCE METHOD

by

Robert G. Fuerst, Frank P. Scaringelli, and John H. Margeson
Quality Assurance Branch
Environmental Monitoring and Support Laboratory
Research Triangle Park, North Carolina 27711

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF RESEARCH AND DEVELOPMENT
ENVIRONMENTAL MONITORING AND SUPPORT LABORATORY
RESEARCH TRIANGLE PARK, NORTH CAROLINA 27711

DISCLAIMER

This report has been reviewed by the Environmental Monitoring and Support Laboratory, U.S. Environmental Protection Agency, and approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

ABSTRACT

This report describes an evaluation of the effect of temperature on the stability of collected samples according to the Environmental Protection Agency (EPA) reference procedure for measurement of ambient sulfur dioxide. This evaluation was carried out over the range of 35 to 278 $\mu\text{g SO}_2/\text{m}^3$.

Collected samples decay at a critical temperature-dependent rate. The rate of decay increases five-fold for every 10°C increase in temperature over the range 20 to 40°C. The rate of decay is independent of concentration over the range studied, and the decay reaction follows first-order kinetics.

At 20, 30, 40 and 50°C (68, 86, 104 and 122°F) SO_2 is lost at the rate of 0.9, 5.0, 25, and 74 percent per day, respectively.

A mathematical model was developed that allows sample decay to be calculated if the temperature history of the sample is known.

Temperature specifications, and changes in the procedure necessary to eliminate the decay problem are proposed.

CONTENTS

	<u>Page</u>
Abstract	iii
List of Figures	v
List of Tables	vi
Acknowledgments	vii
I Introduction	1
II Experimental	3
III Results and Discussion	5
IV Conclusions	19
V Recommendations	20
VI References	22
Technical Report Data Sheet	23

LIST OF FIGURES

<u>Number</u>	<u>Page</u>
1. Effect of Temperature on Stability of Dichlorosulfitomercurate Complex, 50°C.	6
2. Effect of Temperature on Stability of Dichlorosulfitomercurate Complex, 40°C	7
3. Effect of Temperature on Stability of Dichlorosulfitomercurate Complex, 30°C	8
4. Effect of Temperature on Stability of Dichlorosulfitomercurate Complex, 20°C	9
5. Determination of Order and Rate of Reaction	13
6. Effect of Temperature on Decay Rate Constant of Dichlorosulfitomercurate Complex	15

LIST OF TABLES

<u>Number</u>	<u>Page</u>
1. Effect of Temperature on Percent Decay Per Day	12

ACKNOWLEDGMENTS

The authors wish to thank Dr. Joseph E. Knoll of the Quality Assurance Branch, Environmental Monitoring and Support Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina for helpful discussions during this project.

SECTION I

INTRODUCTION

On April 30, 1971, the Environmental Protection Agency (EPA) promulgated in the Federal Register the national primary and secondary air quality standards for sulfur dioxide (SO_2). At the same time, the reference method for the determination of ambient concentrations of SO_2 was also described.¹

Previous research² on the SO_2 reference method indicated that a stable complex was formed when SO_2 in the ambient air was collected in a solution of potassium tetrachloromercurate and subsequently analyzed for SO_2 concentration with formaldehyde and pararosaniline. The complex formed, dichlorosulfito-mercurate (SO_2 -TCM), had suspected thermal instability³ (1 percent per day at 22°C), but sample refrigeration at 5°C resulted in no decay for up to 30 days.

With the completion of two collaborative tests^{4,5} of the SO_2 reference method, no indication of any major effect of thermal instability was found. This was due mostly to the experimental design of the tests, which did not address itself to elevated temperatures of sampling, shipment, and storage of SO_2 -TCM samples prior to analysis.

At present, it is a common practice to sample for ambient concentrations of SO_2 in a thermostated sampling box (35°C) and then to transport the collected 24-hour SO_2 samples back to the laboratory for analysis in a non-temperature controlled container. This could involve either direct transportation of samples to the laboratory by staff personnel or transportation by the U.S. Postal Service. Upon receipt at the laboratory, the samples are stored either in the

dark at room temperature, or in a refrigerator. Thus, prior to analysis, the collected samples are exposed to a variety of temperature conditions for various lengths of time. Therefore, it became necessary to define the effect of temperature and time on the stability of SO₂-TCM samples, since it had never before been investigated at elevated temperatures.

SECTION II

EXPERIMENTAL

The specific reagents and procedures used for sample preparation and analysis are those specified in the Federal Register for the determination of SO_2 in the atmosphere (Pararosaniline Method).¹ Other pertinent experimental details not contained in that procedure are described in the following sections:

SAMPLE PREPARATION

Sufficient quantities of five concentrations of SO_2 in TCM (0, 0.2, 0.4, 0.8, and 1.6 $\mu\text{g SO}_2/\text{ml}$) were prepared by dilution of an SO_2 -TCM stock standard prepared from sodium metabisulfite. Four sets of the five different concentrations were prepared by filling (~ 80 ml) each 100-ml polypropylene centrifuge tube with a specific concentration and capping it with a polypropylene tube closure. This is the same tube and closure recommended¹ for use when sampling for SO_2 in the ambient air for 24 hours.

TEMPERATURE CONTROL

Each set of tubes was placed in a constant temperature bath. Each bath was thermostated at either 20, 30, 40 or 50°C. The specific temperature was maintained within $\pm 0.1^\circ\text{C}$. Each tube was submerged below the solution volume line.

SAMPLE ANALYSIS

After selected periods of time, portions of the thermostated samples were taken, and rapidly cooled to 20°C in a constant temperature bath to prevent further decay before analysis. Aliquots of these portions were analyzed immediately for SO_2 concentration by the SO_2 reference method. This temperature (20°C) was also the temperature used for dye formation in the analysis part of

the procedure.

DATA ANALYSIS

Best fit regression analyses and equations were derived using a programmable calculator-plotter.

SECTION III

RESULTS AND DISCUSSION

In sampling and analysis of ambient concentrations of SO_2 by the Federal Register method, there were two areas where temperature and length of exposure to various temperatures needed investigation. These areas were: (1) the effect of temperature on the ability to collect a valid sample and (2) the effect of temperature on the stability of the collected sample. We chose the second area for our investigations, assuming that if the decay is substantial at higher temperatures, we would need to define it statically before any investigation of the dynamic collection system could be attempted. This report then discusses only area two.

To define the effect of temperature on the determination of ambient concentrations of SO_2 collected in 0.04 M TCM, we needed to investigate four areas: (1) the reaction rate at which the SO_2 -TCM complex is decaying with time at a specific ambient temperature, (2) the reaction order indicating how the reaction rate varies with SO_2 concentration, or with another species present, (3) a general outcome (decay) when any concentration of SO_2 in TCM is exposed to elevated ambient temperatures, and (4) a method to estimate the effect of sample exposure to varying temperature.

RATE OF REACTION

The rate of reaction (decay) of the SO_2 -TCM complex was determined by analyzing aliquots of the thermostated samples for SO_2 concentration by the SO_2 reference method, at various periods of time, and then plotting the data to derive a best fit regression equation (Figures 1 to 4). Using this technique the slope of the curve described the rate of decay. The general




Figure 1. Effect of temperature on stability of dichlorosulfitomercurate complex, 50°C .




Figure 2. Effect of temperature on stability of the dichlorosulfitomercurate complex, 40°C.




Figure 3. Effect of temperature on stability of dichlorosulfitomercurate complex, 30°C.




Figure 4. Effect of temperature on stability of dichlorosulfitomercurate complex, 20°C.

equation best fitting these data is described by an exponential curve of the formula

$$C = Ae^{-kt} \quad (1)$$

where

C = concentration measured, $\mu\text{g SO}_2/\text{ml}$

A = a constant, $\mu\text{g SO}_2/\text{ml}$

k = rate of decay, day^{-1}

t = time, day

If we assume no decay has taken place at $t = 0$ ($k = 0$), then $C = A$, which defines A as the initial concentration, C_0 , and the equation

$$C = C_0 e^{-kt} \quad (2)$$

will describe the data.

Comparing the rates of decay at each temperature (20, 30, 40 and 50°C), we found the rate of decay at a given temperature to be independent of concentration in the range of 0.2 to 1.6 $\mu\text{g SO}_2/\text{ml}$. This represents a 24-hour SO_2 sample concentration of 35 to 278 $\mu\text{g SO}_2/\text{m}^3$ collected at a sample flow rate of 0.2 l/min.

The rate of decay of the 0.2 $\mu\text{g SO}_2/\text{ml}$ sample at 50°C differs slightly from the other rates of decay at that temperature. This effect is also evident at the other temperatures, and is probably due to poor precision in the measurements at this low concentration. This is evident when one considers that the minimum sensitivity is approximately 0.14 $\mu\text{g SO}_2/\text{ml}$ and that small net losses ($< 0.14 \mu\text{g/ml}$) are being measured at this concentration.

By determining an average rate of decay (\bar{k}) at each temperature, we can describe decay per day of a collected SO_2 -TCM sample at a specific temperature (Table 1) with the equation

$$P = 100 (1 - e^{-\bar{k}t}) \quad (3)$$

where

P = percent decay/day

100 = conversion factor

\bar{k} = average rate of decay, day^{-1}

t = time of exposure to a specific temperature, day

The data in Table 1 indicate that, with this reaction, the average rate of decay (\bar{k}) increases about 5-fold for each 10°C rise in temperature.

ORDER OF REACTION

Although our data indicated exponential decay and therefore, a first order reaction, we applied the Van't Hoff equation⁶ to the data as an independent means of determining the order of the decay reaction. Van't Hoff stated that the true order of reaction can be determined using a log-log plot of the velocity of the reaction against different initial sample concentrations at a specific temperature, e.g.

$$V = kc^n \quad (4)$$

$$\ln V = \ln k + n \ln c \quad (5)$$

where:

\ln = logarithm, base e

V = velocity of reaction, $\mu\text{g SO}_2/\text{ml-day}$

k = rate of reaction, day^{-1}

n = order of reaction (slope of line)

c = concentration of sample, $\mu\text{g SO}_2/\text{ml}$

Table 1. EFFECT OF TEMPERATURE ON PERCENT DECAY PER DAY

Temperature, °C	Average rate constant of decay (\bar{k})	Percent loss per day
20	0.009 day ⁻¹	0.9
30	0.051 day ⁻¹	5.0
40	0.287 day ⁻¹	25.0
50	1.33 day ⁻¹	73.6




Figure 5. Determination of order and rate of reaction.

If this plot gives a straight line, then the slope of the line is an estimate of the order of the reaction with respect to initial sample concentration and the y-intercept is an estimate of the logarithm of the rate of decay. Our data indicate that the average order of reaction is 0.97 (Figure 5), thus confirming that the decay reaction is first order.

GENERAL OUTCOME (DECAY)

Arrhenius⁷ showed that for certain reactions there is a linear relationship between the logarithm of the reaction rate (decay in this case) and the inverse of the absolute temperature at which the rate was measured. Therefore, by determining rates of decay at a few specific temperatures, we can derive an equation that will describe the rate of decay at any temperature within the range measured.

After determining the average rates of decay at 20, 30, 40 and 50°C, we used the Arrhenius equation (given below) to calculate an expression that would describe the rate of decay at any temperature within this range.

$$k = Ae^{-E/RT} \quad (6)$$

where

- k = rate of reaction (decay), day⁻¹
- A = frequency factor, day⁻¹
- E = energy of activation, calories/mole
- R = gas constant, 1.987 calories/degree-mole
- T = absolute temperature of reaction, °K

Taking logarithms, we have the expression

$$\ln k = \ln A - (E/R)(1/T) \quad (7)$$

which has the form $Y = b + mx$

A plot of this equation, using the above decay data, is shown in Figure 6.




Figure 6. Effect of temperature on decay rate constant of dichlorosulfito-mercurate complex.

An Arrhenius equation will hold only if the activation energy remains independent of temperature. This is the case as indicated by the linear relationship shown in Figure 6 where

$$\ln k = 48.735 - 15661 \left(\frac{1}{T} \right) \quad (8)$$

This equation can now be used to calculate the rate of decay at any temperature within the range 20 to 50°C and to estimate the rate of decay outside this range.

This equation also indicates that the energy of activation of the decay reaction is 31.2 kcal/mole.

Scaringelli³ found that at 22°C, the SO₂-TCM complex decayed at the rate of 1.0 percent per day and could be stored at 5°C for up to 30 days without any noticeable decay. Using equation (8) we find that at 22°C the rate of decay is 1.3 percent per day and at the end of 30 days at 5°C a sample will have decayed a total of only 1.5 percent. Thus, the current study confirms the values previously reported.

EFFECT OF VARYING TEMPERATURE

Because field samples are exposed to different temperatures for various periods of time, a mathematical model was developed to estimate total decay that would be experienced under varying but known temperature conditions.

Using equation (2) and breaking the time of exposure (t) into parts of a day, we have

$$C_m = C_o e^{-[(k_1 t_1)_{T_1} + (k_2 t_2)_{T_2} + (k_3 t_3)_{T_3} + \dots (k_i t_i)_{T_i}]} \quad (9)$$

taking logarithms, we have

$$\ln C_m = \ln C_o - [(k_1 t_1)_{T_1} + (k_2 t_2)_{T_2} + (k_3 t_3)_{T_3} + \dots (k_i t_i)_{T_i}] \quad (10)$$

$$\ln C_m = \ln C_o - [\Sigma(k_i t_i)]_{T_i} \quad (11)$$

where

C_m = concentration measured, $\mu\text{g SO}_2/\text{ml}$

C_o = initial concentration, $\mu\text{g SO}_2/\text{ml}$

k_i = rate of decay at a specific temperature, day^{-1}

t_i = time of exposure to a specific temperature, day

T_i = a specific temperature, $^{\circ}\text{K}$

Thus, by using the Arrhenius equation, we can determine the decay rate (k_i) for any temperature (T_i). By knowing the time of exposure (t_i) to a particular temperature, we can determine the overall effect of varying temperature on sample concentration. As practical examples of the use of this equation, we have estimated the decay of collected SO_2 -TCM samples on exposure to the temperature conditions of a typical spring and summer day at the Research Triangle Park. If the initial concentration C_o is set equal to $1.0 \mu\text{g SO}_2/\text{ml}$ ($174 \mu\text{g SO}_2/\text{m}^3$), C_m will also represent the fraction of SO_2 remaining after decay for any concentration of SO_2 exposed to these conditions. The calculations are:

<u>Temperature Conditions, Spring Day</u>	<u>k_i</u>	<u>$k_i t_i$</u>
16 hours at 22°C (72°F)	0.013	0.009
4 hours at 25°C (77°F)	0.023	0.004
4 hours at 30°C (86°F)	0.054	0.009
$[\Sigma(k_i t_i)]_{T_i} = 0.022$		

$$\ln C_m = 0 - 0.022 = -0.022$$

$$C_m = 0.978 \text{ } \mu\text{g SO}_2/\text{ml}$$

or a sample decay of 2.2 percent per day

<u>Temperature Conditions, Summer Day</u>	<u>k_i</u>	<u>$k_i t_i$</u>
16 hours at 22°C (72°F)	0.013	0.009
4 hours at 30°C (86°F)	0.054	0.009
4 hours at 35°C (95°F)	0.124	0.021

$$[\Sigma(k_i t_i)]_{T_i} = 0.039$$

$$\ln C_m = 0 - 0.039 = -0.039$$

$$C_m = 0.962 \text{ } \mu\text{g SO}_2/\text{ml}$$

or a sample decay of 3.8 percent per day.

These examples are intended to show the weighted effect, of even short-time exposure to temperature of 35°C or above on the overall decay rate experienced by SO₂-TCM samples.

SECTION IV

CONCLUSIONS

The data generated in this study indicated that collected SO_2 -TCM samples decay at a critical temperature-dependent rate. The rate of decay is much greater than had been suspected and follows first order reaction kinetics. If the temperature history of the collected sample is known, the decay can be determined and corrected for by use of a mathematical model.

Since the SO_2 -TCM complex is formed during sampling, one would expect that elevated temperatures would also cause decay during sampling. This has recently been confirmed in our laboratory.

SECTION V

RECOMMENDATIONS

The EPA Reference Method for measurement of SO_2 has been shown to be a reliable procedure when used under more restrictive temperature conditions than those in common field practice today.^{4,5} Therefore, to maintain the integrity of field samples, we recommend that the solution temperature be kept at 25°C or less during sampling and at 5°C throughout shipment and storage prior to analysis. To meet these more restrictive temperature specifications, the following techniques are suggested as possible solutions.

REDESIGN OF GAS SAMPLER AND SHIPPING CONTAINER

This technique will in fact solve the decay problem altogether, if the more restrictive temperature specifications mentioned above are met. Physically separating the pump from the sampler, maintaining the temperature inside the sampler below 25°C year-round and shipping the samples in a refrigerated container will help meet these specifications.

RELOCATION OF SAMPLING SITE

If the sampling site could be relocated into an air conditioned room (22°C) and the sample maintained at room temperature, no significant decay would take place during sampling. After sampling, the samples must still be placed in a 5°C chamber during shipment and storage prior to analysis. This solution would presumably require relocating a large number of sampling sites.

CONTINUOUS MONITORS

The use of continuous monitors would provide no possibility for sample decay and also would give short-term pollutant profiles, but the initial cost per site would be high. The instrument would also require temperatures of approximately 22°C year-round for proper operation.

SAMPLE OF KNOWN CONCENTRATION

Instead of minimizing decay by temperature control, the decay can be measured and corrected by exposure of a sample of known concentration to the same environmental conditions as the field sample. This technique could be used to determine sample decay after collection, but would not provide an accurate measure of decay during sampling, because the concentration of the field sample is unknown and changing during sampling.

SECTION VI

REFERENCES

1. Federal Register 36:8186-8187, April 30, 1971.
2. Scaringelli, F. P., B. E. Saltzman, and S. A. Frey. Anal. Chem. 39:1709-1719, December 1967.
3. Scaringelli, F. P., L. Elfers, D. Norris, and S. Hochheiser. Anal. Chem. 42: (14):1818-1820, December 1970.
4. McKee, H. C., R. E. Childers, and O. Saenz, Jr. Collaborative Study of Reference Method for Determination of Sulfur Dioxide in the Atmosphere (Pararosaniline Method), Prepared by Southwest Research Institute, Houston, Texas for the U.S. Environmental Protection Agency, Research Triangle Park, North Carolina, No. PB-205-891, September 1971.
5. McCoy, R. A., D. E. Camann, and H. C. McKee. Collaborative Study of Reference Method for Determination of Sulfur Dioxide in the Atmosphere (Pararosaniline Method)(24-Hour Sampling). Prepared by Southwest Research Institute, Houston, Texas, for U.S. Environmental Protection Agency, Research Triangle Park, North Carolina, No. EPA-650/4-74-027, December 1973.
6. Laidler, K. J. Chemical Kinetics, 2nd Ed., New York, McGraw-Hill Company, 1965. p. 15-16.
7. Arrhenius, S. Z., Physik Chem. 4:226, 1889.

TECHNICAL REPORT DATA (Please read Instructions on the reverse before completing)		
1. REPORT NO. EPA-600/4-76-024	2.	3. RECIPIENT'S ACCESSION NO.
4. TITLE AND SUBTITLE EFFECT OF TEMPERATURE ON STABILITY OF SULFUR DIOXIDE SAMPLES COLLECTED BY THE FEDERAL REFERENCE METHOD	5. REPORT DATE May 1976	6. PERFORMING ORGANIZATION CODE
7. AUTHOR(S) Robert G. Fuerst, Frank P. Scaringelli, and John H. Margeson	8. PERFORMING ORGANIZATION REPORT NO.	
9. PERFORMING ORGANIZATION NAME AND ADDRESS U.S. Environmental Protection Agency Office of Research and Development Environmental Monitoring and Support Laboratory Research Triangle Park, North Carolina 27711	10. PROGRAM ELEMENT NO. 1HD621	11. CONTRACT/GRANT NO.
12. SPONSORING AGENCY NAME AND ADDRESS Environmental Monitoring and Support Laboratory Office of Research and Development U.S. Environmental Protection Agency Research Triangle Park, North Carolina 27711	13. TYPE OF REPORT AND PERIOD COVERED Final	14. SPONSORING AGENCY CODE EPA-ORD
15. SUPPLEMENTARY NOTES		
16. ABSTRACT This report describes an evaluation of the effect of temperature on the stability of samples collected according to the Environmental Protection Agency procedure for measurement of ambient sulfur dioxide. This evaluation was carried out over the range 35 to 278 micrograms per cubic meter of air sampled. Collected samples were found to decay at a critical temperature-dependent rate. The rate of decay increases five-fold for every 10 degree centigrade increase in temperature over the range 20 to 40 degrees. The rate of decay is independent of concentration over the range studied, and the decay reaction follows first-order kinetics. A mathematical model was developed that allows sample decay to be calculated if the temperature history of the sample is known. Temperature specifications and changes in the procedures necessary to eliminate the decay problem are proposed.		
17. KEY WORDS AND DOCUMENT ANALYSIS		
a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
Air pollution Sampling Mathematical model Temperature	Sulfur dioxide	13B
18. DISTRIBUTION STATEMENT Document is available to the public through National Technical Information Service, Springfield, Virginia 22161	19. SECURITY CLASS (This Report) Unclassified 20. SECURITY CLASS (This page) Unclassified	21. NO. OF PAGES 30 22. PRICE

INSTRUCTIONS

1. **REPORT NUMBER**
Insert the EPA report number as it appears on the cover of the publication.
2. **LEAVE BLANK**
3. **RECIPIENTS ACCESSION NUMBER**
Reserved for use by each report recipient.
4. **TITLE AND SUBTITLE**
Title should indicate clearly and briefly the subject coverage of the report, and be displayed prominently. Set subtitle, if used, in smaller type or otherwise subordinate it to main title. When a report is prepared in more than one volume, repeat the primary title, add volume number and include subtitle for the specific title.
5. **REPORT DATE**
Each report shall carry a date indicating at least month and year. Indicate the basis on which it was selected (*e.g., date of issue, date of approval, date of preparation, etc.*).
6. **PERFORMING ORGANIZATION CODE**
Leave blank.
7. **AUTHOR(S)**
Give name(s) in conventional order (*John R. Doe, J. Robert Doe, etc.*). List author's affiliation if it differs from the performing organization.
8. **PERFORMING ORGANIZATION REPORT NUMBER**
Insert if performing organization wishes to assign this number.
9. **PERFORMING ORGANIZATION NAME AND ADDRESS**
Give name, street, city, state, and ZIP code. List no more than two levels of an organizational hierarchy.
10. **PROGRAM ELEMENT NUMBER**
Use the program element number under which the report was prepared. Subordinate numbers may be included in parentheses.
11. **CONTRACT/GRANT NUMBER**
Insert contract or grant number under which report was prepared.
12. **SPONSORING AGENCY NAME AND ADDRESS**
Include ZIP code.
13. **TYPE OF REPORT AND PERIOD COVERED**
Indicate interim final, etc., and if applicable, dates covered.
14. **SPONSORING AGENCY CODE**
Leave blank.
15. **SUPPLEMENTARY NOTES**
Enter information not included elsewhere but useful, such as: Prepared in cooperation with, Translation of, Presented at conference of, To be published in, Supersedes, Supplements, etc.
16. **ABSTRACT**
Include a brief (*200 words or less*) factual summary of the most significant information contained in the report. If the report contains a significant bibliography or literature survey, mention it here.
17. **KEY WORDS AND DOCUMENT ANALYSIS**
 - (a) **DESCRIPTORS** - Select from the Thesaurus of Engineering and Scientific Terms the proper authorized terms that identify the major concept of the research and are sufficiently specific and precise to be used as index entries for cataloging.
 - (b) **IDENTIFIERS AND OPEN-ENDED TERMS** - Use identifiers for project names, code names, equipment designators, etc. Use open-ended terms written in descriptor form for those subjects for which no descriptor exists.
 - (c) **COSATI FIELD GROUP** - Field and group assignments are to be taken from the 1965 COSATI Subject Category List. Since the majority of documents are multidisciplinary in nature, the Primary Field/Group assignment(s) will be specific discipline, area of human endeavor, or type of physical object. The application(s) will be cross-referenced with secondary Field/Group assignments that will follow the primary posting(s).
18. **DISTRIBUTION STATEMENT**
Denote releasability to the public or limitation for reasons other than security for example "Release Unlimited." Cite any availability to the public, with address and price.
19. & 20. **SECURITY CLASSIFICATION**
DO NOT submit classified reports to the National Technical Information service.
21. **NUMBER OF PAGES**
Insert the total number of pages, including this one and unnumbered pages, but exclude distribution list, if any.
22. **PRICE**
Insert the price set by the National Technical Information Service or the Government Printing Office, if known.