Geographic Information Systems Applications Notebook

United States Environmental Protection Agency Region III - Philadelphia

September, 1991

Produced by
The Geographic Information Center
Information Resources Management Branch

Contents

Introduction

Water Management Division

- 1. Non-point Source Pollution From Agricultural Practices
- 2. Targeting Areas for Wellhead Protection
- 3. Underground Injection Control Site Discovery
- 4. Ground Water Pollution Potential Assessment
- 5. Regional Ground Water Risk Assessment
- 6. Sub-Regional Ground Water Risk Assessment
- 7. Federal Facilities Compliance Initiative

Water Management and Hazardous Waste Management Divisions

- 8. CERCLA Site Discovery
- 9. GIS Automation of Hazard Ranking System, Ground Water Migration Pathway

Environmental Services Division

- Delaware River Water Trends Report
- 11. Assessing Toxicological Impacts

Air, Radiation and Toxics Division

- 12. Monitoring Radon Levels
- 13. Air Quality Monitoring

Hazardous Waste Management Division

- 14. Superfund Applications
- 15. Oil Spill Emergency Response

Regional Initiatives

16. Regional Strategic Planning

Contents (continued)

Regional Projects

Building Regional Geographic Databases

List of GIS Personnel

List of Project Leads

Introduction

Geographic Information Systems (GIS) is a rapidly developing technology that allows users the capability of collecting, managing, displaying, and analyzing large volumes of data from various environmental media. By better understanding the spatial relationship between natural resources, human populations and potential sources of contamination, the Region can more effectively accomplish the Agency's mission.

Region III obtained GIS capabilities in December of 1988. The Geographic Information Center (GIC) was established within the Information Resources Management Branch (IRMB) to provide GIS expertise and support to EPA programs, and to manage GIS related hardware and software. Since that time, GIS projects have been completed in almost all EPA program areas. This document contains brief descriptions and sample GIS output from some of the Region's most successful GIS efforts. Unless stated otherwise, GIS technical and cartographic support was provided by Region III's GIC.


The maps contained in this document are color xerox reproductions of originals produced on a Calcomp 5800 series color electrostatic plotter. The majority of the maps were originally designed to be printed at a 40 inch by 34 inch format, and were reduced prior to xeroxing. As a result, there may be some loss of resolution and readability. We hope this document will provide an insightful overview of GIS activities in Region III.


Non-point Source Pollution From Agricultural Practices


EPA Region III's Ground Water Protection Section is using GIS to assist Jefferson County, West Virginia and the State Department of Agriculture in developing their Pesticide in Ground Water Management Plan. An interagency workgroup of federal, state, and local agencies was established to help in this effort. Each agency has contributed important data layers to the project and a detailed agricultural practices survey of 118 farmers was conducted by EPA and the County Planning Council.

Several different types of analysis are being conducted by EPA staff using GIS. The maps on the following pages were created to identify areas where pesticides are being applied to vulnerable ground water areas. Vulnerability was determined by the State Department of Natural Resources using EPA's "DRASTIC" methodology (see map on the following page). "DRASTIC" is an acronym representing the most important factors controlling ground water pollution potential. GIS will allow EPA to further refine the vulnerability model by considering soil leaching properties, pesticide leaching properties, ground water flow directions, and the locations of sink holes, cavernous zones, and faults. The second map displays pesticide usage in pounds per farm as determined from by the agricultural practices survey. In the third map, the vulnerability and the pesticide usage maps were combined in GIS to identify where large amounts of pesticides are being applied to areas of high ground water vulnerability.

EPA is seeking funds to compare predictions regarding the effects of agricultural practices on ground water quality made using GIS with actual ground water monitoring data. The USGS National Water Quality Assessment of the Potomac River Basin is a potential source of funding for this proposed validation effort.








Targeting Areas for Wellhead Protection

In the interest of assisting Region III states in developing Wellhead Protection Programs, the Ground Water Protection Section (GWPS) is using GIS technology to gather data and produce high quality GIS maps designed to stimulate interest in wellhead protection on the part of state and local officials. As seen in the following example, GWPS and the Geographic Information Center (GIC) produced an intriguing map that targeted areas in Anne Arundel County, Maryland for wellhead protection. The map shows the spatial relationships between potential sources of contamination and public water supply wells. These features were overlaid with a relative ground water vulnerability assessment map. The relative vulnerability was based on the weighted ranking of seven key hydrogeologic factors, referred to as DRASTIC by the National Water Well Association (NWWA).

Maps showing potential for ground water contamination were provided to state and county officials along with data necessary for delineating wellhead protection areas. GIS served as the catalyst in bringing together key state and local officials. They have now formed a workgroup to delineate wellhead protection areas using GIS as one of the primary tools.





Underground Injection Control Site Discovery

The Underground Injection Control Section is responsible for regulating the discharge of waste to the subsurface through injection wells. GIS was used to identify certain industrial facilities for inspection that are not regulated by EPA but that may be contaminating ground water. Sewered and unsewered areas were compared with locations for industrial facilities, gasoline service stations, and automobile related businesses. Since those located in unsewered areas are not discharging to publicly owned treatment facilities, it is possible that some of them may be discharging waste through Class V injection wells into shallow aquifers.


Seven counties in southeastern Pennsylvania that rely heavily on ground water as a drinking water source were selected for this study. For each county, maps of sewered and unsewered areas were obtained from the county planning commissions and digitized. Data from EPA's FINDS database and from a Dun and Bradstreet database were used to identify the locations of certain types of industries, and gasoline service stations.

Six maps were created for each county: industrial facilities, industrial facilities with twenty or more employees, industrial facilities with sales of \$1,000,000 a year or more, gasoline service stations, automobile related industries with twenty or more employees, and automobile related industries with sales of \$1,000,000 or more. These maps were used by inspectors to prioritize the facilities that they inspected. The two maps on the following pages are included as examples. The first identifies gasoline service stations located in unsewered areas. The second identifies industrial facilities for selected SIC codes that are located in unsewered areas.

The results were very favorable and numerous violations were encountered. Prioritizing the facilities for inspection allowed the Underground Injection Control Section to achieve significant cost savings and they are currently repeating the process for additional counties. In addition, they are using GIS to determine the spatial relationship between the identified injection wells and municipal and residential wells.

GASOLINE SERVICE STATIONS IN LANCASTER COUNTY, PENNSYLVANIA Areas Without Severs D Areas That Are Severed Jula: Pasemay dels from 1865 | 1:100,000 bild. Intermibile related dels from Dans & Projection: Produced | 100 per 1800 per 1800

INDUSTRIAL FACILITIES IN LANCASTER COUNTY, PENNSYLVANIA



Ground Water Pollution Potential Assessment

The Ground Water Protection Section is charged with assisting states in developing Wellhead Protection Programs. Limited funding has been appropriated to the states for these activities and the Region needed a technical assistance vehicle. GIS filled this void and was used to gather data and stimulate interest on the part of state and local offices by providing a management tool that would be useful in wellhead protection programs.

The project also sought to demonstrate an application on a countylevel scale. New Castle County, Delaware was selected and areas in the county that are at high risk for ground water contamination were identified and prioritized for further action. Over sixty data layers were converted from New Castle County Water Resource Authority's (NCC WRA) GIS system to Region III's system. Locations of potential sources of contamination were obtained from various EPA databases. These data were used to produce the maps on the following pages. The first map illustrates the location of potential sources of contamination with respect to wellhead protection areas and ground water recharge areas. In the second map the DRASTIC index was used to identify areas of high ground water vulnerability. GIS was used to identify areas of high vulnerability within recharge protection areas or within two miles of a public water supply well. These areas were further prioritized by using GIS to identify and attach weighing factors to potential sources of contamination located in proximity to these vulnerable ground water areas. The third map shows the algorithm used to prioritize the areas and the results of the analysis.


Maps and databases produced for this project were provided to NCC WRA officials. They have since modernized their own GIS capabilities and are using GIS for a wide variety of environmental activities, including the passage of a Wellhead Protection Ordinance. The methodologies developed were later used by the Ground Water Protection Section to conduct a regional ground water risk assessment.

The GIS Project resulted in both tangible and intangible benefits to the Region and the NCC WRA:

 A true federal/local partnership was developed to work towards a common goal - with little money involved.

- EPA provided a technical assistance vehicle for states/locals in a program where resources are limited.
- The development of a management tool to assist managers in determining environmental program priorities.
- The Region's understanding and assessment of the scope and nature of environmental problems was dramatically improved.

LOCATION OF POTENTIAL SOURCES OF CONTAMINATION RELATIVE TO GROUND WATER RECHARGE AREAS AND WELLHEAD PROTECTION AREAS NEW CASTLE COUNTY, DELAWARE



POTENTIAL VULNERABILITY BASED ON DRASTIC INDEX





TARGETING OF HIGH RISK GROUND WATER AREAS


RANKING OF HIGH RISK AREAS


Those portions of the Wellhead and Recharge Protection Areas determined by DRASTIC Index to be highly vulnerable to ground water pollution were selected for analysis. These areas were evaluated relative to each other by the equation shown below. The equation incorporates a range of risk-related factors including potential sources and known incidents of contamination. This map shows the 10 areas at highest risk for groundwater contamination. Their characteristics are summarized in the table below.

RATING • 2(MCL VIOLATIONS) • 2(RCRA RELEASES)

- . 2(USTs 15 YEARS OR OLDER)
- CERCLA 2(RCRA) 3(USTs)

TEN GEOGRAPHIC AREAS AT HIGHEST RISK FOR GROUND WATER CONTAMINATION






Regional Ground Water Risk Assessment

One of the earliest GIS projects in Region III, this study's purpose was to collect various program-related data and to combine them in such a way as to reveal the spatial variation of existing or potential ground water quality problems. The project was accomplished by Temple University Lab for Geographic Information Systems under an FY'89 grant from the Drinking Water/Ground Water Branch. The entire region was examined using counties as the basic study units to identify particular problem areas or "hot spots" needing further attention.


To prioritize those counties in need of the greatest EPA effort and resources, various layers of the USGS 1:2,000,000 Digital Line Graph data were combined with retrievals made from a variety of EPA program databases associated with drinking water quality and potential sources of contamination. For each database, summary statistics were generated by county and related to the digital county map in ARC/INFO. Program data included in the GIS were:

- Public Water Supply (PWS) systems primarily dependent on ground water
 - Population Served by each system
 - Maximum Contaminant Level Violations for each system
- DRASTIC score (to estimate vulnerability to contamination)
- Soluble Pesticides Usage value for each county
- Superfund NPL Sites
- RCRA Treatment, Storage and Disposal Sites
- Class V Underground Injection Wells
- Known ground water contamination incidents

To estimate each county's relative degree of risk to ground water contamination, a weighted ranking scheme was developed to combine all of these factors into a single summary score. The map on the following page shows those counties determined to be at greatest potential risk to ground water contamination.



FY '89 RANKING OF EPA REGION III COUNTIES, BASED UPON GROUND WATER CONTAMINATION RISKS



United States Environmental Protection Agency Drinking Water/Ground Water Protection Branch



Sub-Regional Ground Water Risk Assessment

As in the Regional Ground Water Assessment, a grant from the Drinking Water/Ground Water Protection Branch was issued to the Temple University Lab for Geographic Information Systems. Building on the experience gained in the Regional Assessment, this effort examined the state of Pennsylvania in greater detail. The goal of the project was to synthesize various EPA program data and to identify areas with existing or potential ground water quality problems.


Many layers of data were included in the analysis. Political boundaries and Land Use/Land Cover data layers were built from the USGS 1:250,000 digital data. Program data included in the GIS were:

- Public Water Supply (PWS) systems primarily dependent on ground water
 - Population Served by each system
 - Maximum Contaminant Level Violations for each system
- DRASTIC score (to estimate vulnerability to contamination)
- · Soluble Pesticides Usage value for each county
- Superfund NPL Sites
- RCRA Large Quantity Generators and Treatment, Storage and Disposal Sites
 - · Groundwater and Land Ban Violations
- Sites reporting to the Toxic Release Inventory System SARA Title III)
- Class V Underground Injection Wells
- A select set of businesses from the Dun & Bradstreet Conquest File

Another objective of the project was to try to analyze the data on a site-by-site basis, rather than aggregated by county. Wherever possible, attempts were made to retrieve program data at a more detailed level than was done in Regional Assessment. Although the regional level of analysis was considered better than county aggregates, the quality of the locational data contained within the various program databases proved to be inadequate for the intended analysis. Therefore, zipcode areas were chosen as the basic unit of analysis.

Although a weighted ranking (similar to the Regional Assessment) has not been completed, the data sets and resulting maps created for this project have supported various management-level investigations into existing or potential ground water problems within Pennsylvania. Several different agencies within the state will be receiving the digital data sets to support their own program needs.

The map on the following page shows maximum contaminate level (MCL) violations in drinking water supplies aggregated by zipcode areas and is representative of various maps produced for the project.



Federal Facilities Compliance Initiative


The Permits Enforcement Branch of the Water Management Division created a series of Chesapeake Bay Watershed maps. These maps show the locations of all facilities encompassed by the Chesapeake Bay Watershed. The accompanying maps were used in various executive briefings and public meetings. The first presents status of compliance for federal facilities. The second shows the locations of all NPDES permitted facilities in the Chesapeake Bay Watershed.




EPA REGION III

PERMITS ENFORCEMENT BRANCH

Chesapeake Bay FEDERAL FACILITIES COMPLIANCE INITIATIVE






CERCLA Site Discovery

The Ground Water Protection Section (WMD) and the Site Assessment Section (HWMD) with GIS technical support from the Geographic Information Center (GIC) are conducting a joint effort in Allegheny County, Pennsylvania to assist in the CERCLA Site Discovery process. The methodology uses aerial photography for the Allegheny County area interpreted by the Environmental Photographic Interpretation Center (EPIC) to identify potential hazardous waste disposal sites. EPIC examined aerial photography as far back as the 1930's and 1940's up to the present and interpreted the photography for the signature patterns typically associated with the disposal of hazardous waste. The locations for the waste disposal areas were transferred from the aerial photographs to topographic basemaps and digitized into ARC/INFO. Since EPIC identified over 800 areas of possible hazardous waste disposal a ranking scheme will be applied to prioritize sites for further investigation. This prioritization methodology, based conceptually on the Superfund Hazard Ranking System (HRS), calculates the number of people who rely on ground water for their source of drinking water supply, living within a four mile radius of each of the EPIC-identified sites.

The following map of the Allegheny County area displays locations of EPIC-identified sites and CERCLA sites from the CERCLIS database. The locations will be compared to determine if any EPIC-identified sites are already represented in the CERCLIS database.

The second map covers the Pittsburgh East Quadrangle in Allegheny County. It includes census tract shaded by population and public water supply well locations with associated population served. As mentioned previously, this information will be used to calculate the number of people residing within a four mile radius of each of the EPIC-identified sites who rely on ground water as their drinking water source.

CERCLA LOCATIONS ALLEGHENY COUNTY



LEGEND

Strip Mines


Severed Areas

Polestial Mazardous

. CERCLA Locations


CERCLA Facility Sames





Produced by: Security Produced to the Control of the Produced for Security Webs Produced to Section Produced by: Security Produced by Section Produced by: Section Produced by: Section Produced Produced by: Section Produced Books and Security 1990





PITTSBURGH EAST QUADRANGLE

CENSUS TRACT POPULATION

\$1 - 391 \$11 - 1001 1001 - 2500 2001 - 3000 \$101 - 2000

PUBLIC SUPPLY WELLS

POPULATION: 1991

FOTUTIAL BAZAROOM WASH GAIN KNIMION

IA SILL Area. INDICATE Indicated Millional
III bridge Department Office Control Parket
III bridge Department Office Control Parket
III bridge Department Office Office Parket
III bridge Department Office Office Parket
III bridge Department Office Office


. Righert Potential for

Strip None

Nontial Searcher
Vaste Stee

CERCLA Facility Manua







GIS Automation of Hazard Ranking System Ground Water Migration Pathway

The purpose of this project was to evaluate the feasibility of using GIS as a support tool in the Hazard Ranking System (HRS) scoring process. This project was a joint effort between the Ground Water Protection Section of the Water Management Division and the Site Assessment Section of the Hazardous Waste Management Division. The Temple University Laboratory for Geographic Information Systems was given a grant to perform all GIS-related work over the summer of 1990.

The Hazard Ranking System is a scoring system used to assess the relative threat associated with actual or potential releases of hazardous substances at sites (Federal Register, 40 CFR Part 300, December 14, 1990). To assist a user with little GIS experience in characterizing the potential for contamination from a particular site, a series of programs and graphical menus, written in Arc Macro Language, allow the user to interact directly with the data stored in the GIS. Individual procedures from the Ground Water Migration Pathway scoring process were automated so that the user could access them all in their proper order in the scoring process, or individually, so that the user could examine the effect a particular variable has on the overall score. The controlling procedures keep track of all variables that affect the flow of operations within the scoring process, thus keeping the user from improperly scoring a site.

The Targets section of the scoring process accounts for the populations and resources potentially at risk from contamination at the site being considered. Given that the Targets tend to have the greatest influence on the overall pathway score, the GIS data layers focused on the data needed to characterize populations at risk from a given site. As this was a demonstration project with limited funding, data sets were built for Bucks County, Pennsylvania. Included in the GIS analysis are the following layers:

- Updated locations of sites from the CERCLIS database
- 1980 Census tracts and associated total population values
- Public Water Supply (PWS) Intake locations
 - Populations served by Public Water Supply systems
 - PWS Service Area Boundaries
 - · Associated Wellhead Protection Areas

Waterbodies and roads data layers (at an input scale of 1:24,000) from the Pennsylvania Department of Transportation were incorporated for the purpose of providing the user with some regional context for each site during a scoring session.

The graphic on the following page represents a "snapshot" of the computer display at one point in a typical user scoring session. In this image, four different graphic elements perform specialized functions. The largest window is a map displaying the location of the site and all potential targets within four miles. The bottom window is a dialogue area where the user responds to various prompts throughout the session. The large blue "Text Window" is the Scoresheet which summarizes all sub-scores within the Pathway score. This is accessed by choosing the appropriate option from the forth graphic element, the main "Pulldown Menu".

| ITS NO | GROUND WATER MIGRATION PATHWAY SCORESHEET | | |
|--|---|---|-----------------|
| THE PARTY | Factor Categories and Factors | | |
| + | Likelihood of Release to an Aquifer | Maximum Value | Val ue Assigned |
| JIII X | 1. Observed Release | 550 | 550 |
| | 2. Potential to Release | | |
| 0013.6200 | 2a. Cont ai nment | 10 | 8 |
| 75 TOOTTO-1 | 2b. Net Precipitation | 10 | 8 |
| 300175-0 | 2c. Depth to Aquifer | 5 | 9 |
| Y LY | 2d. Travel Time | 35 | 8 |
| 7 1-1 | Ze. Potential to Release | | |
| 1990年1990年 | (lines $2a \times (2b + 2c + 2d)$) | 588 | 8 |
| 200469-002 | 3. Likelihood of Release (higher of | | |
| | lines 1 or 2e | 550 | 550 |
| 00000-000 | Waste Characteristics ' | | |
| 102284700 | 4. Toxicity/Mobility | * | 4500 |
| T JA A V | 5. Hazardous Waste Quantity | * | 4558 |
| MIDIN | 6. Waste Characteristics | 100 | 56 |
| R | | | |
| 101259-004 101258-083 | Targets | | |
| Reservant | 7. Nearest Well | 50 | 58 |
| X THAN S | 8. Population | | |
| | 8a. Level I Concentrations | ** | 8 |
| - Table 21- | 8b. Level II Concentrations | ** | 8 |
| 77 | 8c. Potential Contamination | ** | 1223.7 |
| | 8d. Population (lines 8a + 8b + 8c) | ** | 1223.7 |
| 36 101241-010 PERKASIE | 9. Resources | 5 | 5 |
| 37 100133-003 NO PENN | 18. Wellhead Protection Area | 28 | 28 |
| 38 101241-008 PERKASIE | 11. Targets (lines 7 + 8d + 9 + 18) | | 1298.7 |
| 39 101243-014 SELLERSV 40 100496-002 GROUSE G | | Sales of the Sales of | |
| the above information co | Ground Mater Migration Score for an A | Ground Water Migration Score for an Aquifer | |

Delaware River Water Trends Report

The Environmental Services Division (ESD) initiated an assessment of water quality, sediment quality and biological integrity of the Delaware River as a pilot to a regional Water Quality Trends Report. GIS was used to display monitoring stations along the Delaware River and the level of their dissolved oxygen, fecal coliform bacteria, and the water quality trends for each monitoring station.

For the purposes of the Water Trends Report, the Delaware River was divided into two sections; the Delaware Bay to Trenton, NJ and Trenton, NJ to Hancock, NY. For each of these sections, Two maps were created and included in the Water Quality Trends Report. One displayed the readings for fecal coliform, the second dissolved oxygen. Both indicate trends in water quality.

TRENDS IN FECAL COLIFORM DENSITY Trenton, NJ to Hancock, NY

1 Increasing levels

Decreasing levels

Fecal Coliform


- Less than 100
- 100 500
- **501 900**
- Greater than 900

Station Percentile*

| 1.8 | 260 |
|---|--|
| 10 | 260 790 1300 260 170 316 1600 |
| 20 | 7 9 0 1 3 0 0 2 6 0 |
| 20 | 1300 |
| 41 | 200 |
| 22 | 170 |
| 23 | 316 |
| 24 | 1600 |
| 20 | 108 |
| 97 | 100 |
| 61 | 80 |
| 20 | 4 9 |
| 29 | 63 |
| 3.0 | 6.8 |
| 31 | 200 |
| 32 | 120 |
| 3 3 | 170 |
| 34 | 200 120 170 458 1008 |
| 35 | 1008 |
| 36 | 5.6 |
| 37 | 8.8 |
| 3.8 | 6.9 |
| 30 | 28 |
| 10 | 9.5 |
| 11 | 9.0 |
| 4 1 | 90 |
| 4.6 | 01 |
| 4.5 | 90 |
| 44 | 44 |
| 4.5 | 54 |
| 4.6 | 484 |
| 47 | 1004 |
| 48 | 164 |
| 18901224567890123456789012345678944456789444444444444444444444444444444444444 | 260 790 1300 2600 1706 31668 108 4968 688 2000 12700 458 1008 688 2000 1270 458 1008 444 4484 1004 484 1004 280 |
| | |

*90% of Values \le






DISSOLVED OXYGEN CONCENTRATION TRENDS Delaware Bay to Trenton, New Jersey

- † Increasing levels
- ↓ Decreasing levels

Dissolved Oxygen
10th Percentile


- Greater than or equal to 5.00
- **4**.0 4.99
- Less than 4.00

Station Percentile*



*10% of Values \le






Assessing Toxicological Impacts

The Emergency Planning Community Right-to-Know Act (EPCRA) requires that certain industries report their emissions and off-site transfers for specified chemicals each year. This data is contained in the Toxics Release Inventory (TRI) database. Region III utilized the information in TRI and the available information regarding the toxicological impacts of these chemicals to provide a "weighted" emission for each of the sites. Based on the results of this approach, the accompanying maps were generated using the Region's GIS capabilities.

The first map indicates the location of facilities in New Castle County, Delaware, that reported to the TRI database for 1987, and their relationship with other EPA programs. The pie charts indicate the relative importance of the emissions for one of the facilities. Similar charts can be created for all of the facilities using a program that was written to interface the TRI database with GIS.

Using 1988 TRI data, the top fifty facilities in terms of weighted emissions for air and land disposal were identified. The weighting factor took into consideration the toxicity of each of the chemicals emitted at the individual facilities. The second map in the series shows the air facilities in Pennsylvania with population density to indicate risk to human health. The legend contains the name of each facility and the reported emission for each chemical. The third map shows the land disposal facilities with ground water vulnerability to indicate potential threat to drinking water supplies.





1988 TRI AIR EMISSIONS IN PENNSYLVANIA FOR CANCER POTENCY

| | PACILITY | CHEMICAL NAME | CHEMICAL IN LBS/YEAR |
|-------|---------------------------|--|-------------------------|
| 3. | AMBRICAN ZINC CORP. | CADMIUM COMPOUNDS | 3,090 |
| 4. | BAST CORP. | 1,3-BUTADIENE | 68,920 |
| ₿. | BETHLEHEM STEEL CORP. | BENZENE | 540,000 |
| 10. | CORNING ASAHI YIDRO PROBU | ARSENIC COMPOUNDS | 500 |
| iā. | GRIFFITH MICRO SCIENCE IN | RTHYLENE OXIDE | 90,384 |
| 20. | | CHLOROPORM | 315,700 |
| | LTV STEEL CO.INC: | BENZENE | 858,508 |
| 2 J . | | ARSENIC | |
| | LUBBRO DIBBL LU. | | 500 |
| 27. | OCCIDENTAL CHEMICAL CORP. | VINYL CHLORIDE | 214,810 |
| 28. | OI-NEG TV PRODUCTS INC. | ARSENIC COMPOUNDS | 750 |
| 36. | USS CLAIRTON VORKS | BENZENE | 1,400,770 |
| | VEST CO. | ETRYLENE OXIDE | 38.800 |
| • ' ' | | ************************************** | 70,000 |

POPULATION DENSITY is persons/sq. mi.

M 8 - 58 M 50 - 500 M 500 - 1,000 M greater than 1,000

Bata frem:


USCS 1:2,000,000 BLC.
Population denoity data
derived from USCS Geonames
database. TRI data from the
1948 Textes Release Inventory.
UTR Some 18
daguet, 1998
EPA Region III - Philadelphia
TRI Stillication Workgroup

Projection: Sate: Produced by:





1988 TRI AIR EMISSIONS IN PENNSYLVANIA FOR CANCER POTENCY



1988 TRI LAND RELEASES IN PENNSYLVANIA FOR CANCER POTENCY

| | FACILITY | | CHEMICAL IN LBS/YR |
|-------------|--|----------------------------------|-----------------------|
| 1. | AMBRICA ZINC CORP. AMBRICAN ZINC CORP. ARMCO ADVANCED MATERIALS BETHLEHEM STEEL CORP. BETHLEHEM STEEL CORP. BETHLEHEM STEEL CORP. CARPENTER TECHNOLOGY CORPEMPIRE STEEL CASTINGS INC GRINNELL CORP. LATROBE STEEL CO. LENOX TORKS — BAR ROD & W NGK METALS CORP. OCCIDENTAL CHEMICAL CORP. PENNSYLVANIA STEEL FOUNDR PENNTECE PAPERS INC. RAYMARK FRICTION CO. RENEER FILMS CORP. ST. GRORGE CRYSTAL LTD. STANDARD STEEL STANDARD STEEL STANDARD STEEL TELEDYNE INC. TELEDYNE VASCO | LEAD COMPOUNDS | 120,000 |
| 2. | AMBRICAN ZINC CORP. | CADMIUM COMPOUNDS | 2,400 |
| 3. | AKNUU ADVANUKU MATKKIALS | NICKEL | 75,000 |
| 5. | RETHLEMEN STEEL CORP. | TRAD COMPOUNDS | 1,300 |
| <u> 5</u> . | RELUTERE DIER CABBI CODD | RICERL COMPOUNDS | 20,000 |
| 7 | STREETS TO THE CONTRACTOR SORT | NICKEL COMPOUNDS | 11,000 |
| 10. | CARLBUIRE INCOMPTON INC | NICERL | 187,700 |
| 12. | CDIANELS CODD | | 750 |
| 11. | TRICTOR CTPS: CO | PURMALUENIUS NICESI COMPONNAC | 310,000 |
| 10. | INNUA UBAGATI | I PAR COMPONING | 9,700 |
| 10. | INTER MARKS - RIP DAN & W | I FAR COMPONING | 10,000 |
| 22 | NCE METALS CORP | REPUBLISH COMPOUNTS | 1,990 |
| 99 | OCCIDENTAL CHEMICAL CORP | ALIAN CALUALUA | 1 4 4 6 |
| 25. | PRNNSYLVANIA STREE POSINDR | NICERA | 1,050 |
| 26 | PRINTECH PAPERS INC. | ĈŔĹŌŖŎPOŖW | 254 |
| 27 | RAYMARE PRICTION CO. | LRAD | 23 891 |
| 28. | RENEER FILMS CORP. | CĂDNIUM COMPOUNDS | 20,010 |
| 29. | ST. GBORGE CRYSTAL LTD. | LEAD COMPOUNDS | 22.320 |
| 30. | STANDARD STREL | NICKBL | 1.431 |
| 31. | STANDARD STREL | NICKEL | 5,068 |
| 32. | TRLEDYNE INC. | NICKEL | 250 |
| 33. | TELEDYNE VASCO | NICKEL | 10.000 |

Potential Yulmerability to Contamination, Bused on BRASTIC

This map indicates the relative valuerability of the teglos, by county, to ground water contamination, as indicated by a "BRASTIC" (ACDRASTIC) more. The secret represent an offregated assessment of each county's valuerability range.


The BRASTIC methodology was developed by RPA and the BVVA to evaluate ground water pollution potential. "BRASTIC" is an acronym representing the most important factors controlling ground water pollution potential. Booth to water, set Becharge, Aquifer Media, Soil media, Topography, Impact of the values zone, and hydraulic Conductivity of the aquifor.

☐ Fo Bata ■ 48 - 100 ■ 101 - 141 ■ Greater than 141

Pais from: USCS 1:2,000,000 BLC.
Population denoity data
derived from USCS Connamer
database. THI data from the
1925 Tester Release Inventory.
Projection: UTH Ione 18
August, 1998
Produced by: EPA Region III - Philadelphia
TRI Utilisation Forkgroup




1988 TRI LAND RELEASES IN PENNSYLVANINA FOR CANCER POTENCY




Monitoring Radon Levels

The Air, Radiation and Toxics Division has been using GIS as the centerpiece of their radon outreach program for the last two years. Radon levels are collected as home owners voluntarily place sampling canisters in their homes. This data has been assembled by several private radon testing firms who have made the information available to EPA aggregated by zipcode area.

The Region III Geographic Information Center has produced regional maps with radon levels by county (see map on the following page), county maps with radon levels by zipcode, and radon levels by congressional districts (see second map). Congressmen were provided with radon level maps for their districts to make them aware of environmental conditions in their districts. The maps have been effective in media presentations and as visual aides in testimony before state legislatures.



PERCENTAGE OF RADON READINGS ABOVE 4 pCi/I IN REGION III




PERCENTAGE OF RADON READINGS ABOVE 4 pCi/l PENNSYLVANIA CONGRESSIONAL DISTRICT 9 REPRESENTATIVE BUD SHUSTER

PERCENTAGE OF RADON READINGS ABOVE 4 pCi/I

- Over 60%
- 40% to 60%
- 20% to 40%
- 0% to 20%
- ☐ Insufficient Data

Data:

Radon data from Key Technology,
Teledyne, AirChek Inc. and The
Radon Project.
UTM Zone 18
Plotted:
Produced By:
EFA Region III - Philadelphia
Information Resources Management
Branch
Produced For:
EFA Region III - Philadelphia
Branch
Produced For:
EFA Region BII - Philadelphia
Air Programs Branch






Air Quality Monitoring

GIS has been used over the last two and a half years to monitor air quality in the region. Data from the AIRS database is periodically downloaded into the GIS and ARC/INFO's contouring capabilities are utilized to create concentration isopleths for various pollutants. The maps on the following two pages were created for ozone concentrations and total suspended particulates (TSP) and were included in the Air Quality Trends Report. The next map in this series identifies counties in the northeastern United States that are designated as non-attainment and do not meet minimum ozone standards.


The New Source Review Section of the Air Enforcement Branch used GIS to help visualize the impact of SO2 from recently permitted or proposed sources in Virginia. The fourth map in this series shows the locations of proposed or permitted sources indicated by a colored circle. The diameter of the circle indicates the amount of proposed SO2 emissions in tons per year.


ARC/INFO was used to display the output from air dispersion models by the Technical Assessment Section of the Air Programs Branch. SO2 emissions from two General Services Administration power plants were modeled under various atmospheric and operational scenarios. The final map in this series shows the dispersion plume for the worse case scenario. This project successfully used GIS running on a minicomputer as a preprocessor and post processor for a PC air dispersion model.

1986 - 1988 OZONE DESIGN VALUES FOR REGION III




SUSPENDED PARTICULATE CONCENTRATIONS IN REGION III 1988 GEOMETRIC MEAN






NEW MAJOR SOURCES RECENTLY PROPOSED OR PERMITTED IN VIRGINIA



HEALTH RELATED S 0 2 STANDARD

MAXIMUM ALLOWABLE EMISSIONS

(ALL ELEVEN BOILERS OPERATING)



Superfund Applications

The Hazardous Waste Management Division (HWMD) has used GIS mapping capabilities for a variety of applications. Seven maps representing these applications are contained on the following pages. Numerous maps have been produced to display data contained in CERCLIS and HWDMS databases. The first map shows all of the NPL sites in the region. In addition to this map the Geographic Information Center (GIC) has produced maps of NPL by state, Regional CERCLA, RCRA TSD, and RCRA large and small quantity generators. The second map was prepared for the Regional Administrator and shows all of the hazardous waste sites in the Philadelphia area.

The Eastern Pennsylvania Section of the Superfund Remedial Branch is using GIS to assist in the RD/DA process. ARC/INFO was used to display ground water contaminate concentration data for five Superfund sites in the North Penn area of Montgomery County, PA. The two accompanying maps display TCE for North Penn Area 2 and PCE concentrations for North Penn Area 1. These maps were used to graphically display some of the data contained in the RD/DA workplan. The maps were use in Headquarters briefings and will be used in future public meetings to convey the nature of the problem to the general public.

GIS was an integral part of the RD/RA process at the Berks Sand Pit Superfund Site, PA. An initial GIS map was created from detailed engineering drawings and contains precise well locations, surface water quality monitoring data and analytical results from residential and monitoring wells: This map was use by EPA staff in the field to locate additional monitoring wells and was updated over the course of the sampling phase of the project to reflect new data. integrating residential, monitoring well, and surface water data on the GIS map, EPA staff were able to base decisions regarding the location of ground water extraction wells. This eliminated the need for an interim report, thereby saving the agency significant resources. The map included in this document contains all of the above data, as well as, the location of the treatment plant (shown in brown) required by the remedial action. The map was also used during a community outreach television broadcast and will continue to be updated as new information becomes available.

The Technical Support Section of the Superfund Programs Branch is conducting an in-house Remedial Investigation at the Berkeley

Products NPL Sites. The GIS system is being utilized to illustrate and aid in the analysis of data collected at the site. Site base maps, derived from the interpretation of aerial photography, as well as, magnetometer and soil gas survey data have been assembled to date. Monitoring well, soil trenching, and ecological characterization data will also be evaluated with the GIS. The last two maps in this series of Superfund applications display magnetometer and soil gas survey data contoured using the GIS.

NPI SITES - REGION III

Site Names by State

Delivare

Waryland

Virginia de la companya de la compan

. Numbers reference site sames to siles or map




Opdared: | fam. 1991


Flotted by: EFA Degine | 111
| fabruarious Secure - Management Search

Projection: Universal Transcense Menagement Lance | 18

Data: Base May Information from 1968 DIG (+ 1,010,690 NE) Sine Location From Calific Data Base CfECLIS Data Detrieves Ats (, 1891



NORTH PENN AREA 2 SUPERFUND SITE TRICHLOROETHENE CONCENTRATIONS



LEGEND

- MALLEGED PRP PACILITIES
- TANKS
- SAMPLED WELL LOCATIONS


CONCENTRATIONS OF TRICHLORORTHENE

- 0.5 5 UG/L
 5 25 UG/L
 25 100 UG/L
 100 500 UG/L
 500 1,000 UG/L
 1,000 5,000 UG/L
 5,000 10,000 UG/L
 Greater than 10,000 UG/L
- Bala from: USGS 1:24,000 BLC data, belt sampling data were analyzed by the North Penn Faiter Authority located in Hontgomery County, Pennsylvania.

 Projection: UTL Zone IS Plotted: Pervary 7, 1990
 Produced by: PR Region 111 Philadelphia Information Resources Hangement Branch For PR Region 111 Philadelphia Information Resources Hangement Branch Eastern Pennsylvania Section



NORTH PENN AREA 1 SUPERFUND SITE TETRACHLOROETHENE CONCENTRATIONS



LEGEND

- MALLEGED PRP FACILITIES
- TANKS
- · SAMPLED WELL LOCATIONS


CONCENTRATIONS OF TETRACHLOROETHENE SAMPLED ON FEBRUARY 28, 1986


- 25 UG/L 50 UG/L 15 UG/L - 25 UG/L 5 UG/L - 15 UG/L 0.5 UG/L - 5 UG/L
- Data from: USCS 1:24,000 bic data. Fell sampling data were analyzed by the North Penn Faler Authority localed in Honlyomery County, Pennsylvania.


 Projection: UTL fore 18
 Platted: February 7, 1990
 Produced by: PA Region 111 Philadelphia Information Resources Hangement Branch Fraduced for: PA Region 111 Philadelphia Section Bastern Pennsylvania Section



BERKS SAND PIT - RI/FS Longswamp Township, Berks Co., PA







Oil Spills in the Chesapeake Bay


The Oil and Title III Section of the Hazardous Waste Management Division is using GIS to evaluate the impact of oil shipping on sensitive environmental areas in the Chesapeake Bay. A basemap for the Chesapeake Bay was created and information from various tabular reports were input in the GIS and the following data layers were created:

- · Shipping ports,
- Refineries,
- Facilities that receive petroleum products by barge or ship,
- Oil spills (vessel related only).

These data layers are being used for planning and presentation purposes and eventually, may be developed as an emergency response tool. The map on the following page shows the locations of oil shipping facilities and oil spills in the Chesapeake Bay.

A second phase of the project will map sensitive environmental areas and analyze their proximity to oil spills, refineries, and shipping facilities.

CHESAPEAKE BAY OIL FACILITIES & SPILLS



SPILLS REPORTED TO THE MRC VESSEL RELATED ONLY 1980 - 1987

- 10,000 99,999 Gallons
- 1,000 9,999 Gallons
- 500 999 Gallons
- 100 499 Gallons
- Less Then 100 Callons

M Oil Pipeline

* = Facilities that recieve bulk oil by barge or ship



Profession: Profession: Profession: Profession:

Farman data from 8363 1:250,000 blc. Facility data from the U.S. Count Dan 878 June 10 Juneary, 1991 EPA Region III - Philadelphia 1547-1516 Farman Farmannat Fran



Regional Strategic Planning Initiative

The overall goal of this project is to support EPA Region III's on-going Strategic Planning efforts by using GIS technology to identify geographic areas representing the greatest potential single-program and cross-program environmental risks. Senior management has charged a multi-disciplinary, multi-program task team with identifying the risks to the environment and to human health. The team is interpreting EPA's program data, to define and rank areas for purposes of program resource targeting and environmental indicator selection.

Data from a variety of sources are being retrieved and integrated to determine the current condition of both the natural resources (air, land and water) and the public health in Region III. Data are being displayed on a geographic basis, by county, zipcode and point locations, depending on the data georeferencing. To date, over twenty data layers have been assembled representing a wide range of environmental conditions and risks.

The RISC project's first phase will identify threatened geographic areas for more concentrated evaluation. Seven data layers (TRI-Water, TRI-Air, 304l, Superfund, RCRA, Radon, Ozone) have been selected for thirteen overlay combinations so as to spatially define areas of risk. Later phases will focus on further analyzing the relationship between sources and receptors of environmental risk, both ecological and human health.

Project results will be used by managers to target: pollution prevention initiatives, intensive monitoring and enforcement, priority-setting within programs, multiple-program initiatives, areas of highest multi-media risk, and risk communication and outreach.

This GIS project is significantly fostering cross-program teamwork in developing holistic environmental solutions, is improving data collection and management, and is helping set regional environmental priorities. GIS is an essential tool for managing enormous amounts of data, in a variety of formats, from a large number of EPA programs. Automated data retrieval and analysis enable the project team to perform sophisticated assessments. This project has also directed attention towards helping implement the Agency's Locational Data Policy, which states that by 1995 locational data will be accurate to within twenty-five meters.

Building Regional Geographic Databases


The Region III Geographic Information Center (GIC) of the Information Resources Management Branch is actively building Region-wide geographic databases. The most time consuming part of most projects is data collection and many potential users become discouraged when they learn that significant time and resources are required to collect the data that they will need. It is also anticipated that as GIS hardware become less expensive and a graphic user interface to ARC/INFO becomes available; the GIS user community will need, and expect, immediate access to commonly needed databases.

The GIC is therefore working to build the following Region-wide databases:

- soils,
- land use,
- · wetlands,
- geology,
- · edgematched 1:100,000 transportation,
- RF3 hydrology,
- population,
- and political boundaries.

The land use map of southeastern Pennsylvania on the following page represents a portion of the data that will be included in the region-wide land use database. It was created for USGS GIRAS land use data.

The GIC is also working on methodologies for periodic updates of ARC/INFO coverages created from EPA programmatic data, such as NPL and RCRA site locations.



LEGEND

URBAN OR BUILT-UP LAND

TRANSPORTATION, COMMUNICATIONS, AND UTILITIES

INDUSTRIAL AND COMMERCIAL COMPLEXES
MIXED URBAN OR BUILT-UP LAND
OTHER URBAN OR BUILT-UP LAND

AGRICULTURAL LAND

CROPLAND AND PASTURE
ORCHARDS, GROVES, VINEYARDS, NURSERIES,
AND ORNAMENTAL HORTICULTURAL AREAS
CONTINED FEEDING OPERATIONS
OTHER AGRICULTURAL LAND

FOREST LAND

DECIDUOUS FOREST LAND EVERGREEN FOREST LAND MIXED FOREST LAND

WATER

WATER BODIES

WETLANDS

FORESTED WETLANDS

BARREN LAND

SANDY AREAS OTHER THAN BEACHES
STRIP MINES, QUARRIES, AND
GRAVEL PITS

TRANSITIONAL AREAS

Data from: U.S.G.S GIRAS 1:250,000 Landuse/Landcover Source Date: 1973

Projection: UTN Zone 18 Platted: April 8, 1989

EPA Region III - Philadelphia Information Resources Management Branch Produced by:

Produced for: EPA Region III - Philadelphia Pennsylvania Remedial Response Section

SCALE 1:150,000



GIS Personnel

All Addresses:

United States Environmental Protection Agency 841 Chestnut Street Philadelphia, PA 19107

Information Resources Management Branch

Edward Kratz, Chief Programming Applications and Support Section Mail Stop 3PM53 FTS 597-9855

Geographic Information Center

David West, GIS Team Leader Environmental Scientist Mail Stop 3PM53 FTS 597-1198

Brian Burch Environmental Scientist Mail Stop 3PM53 FTS 597-1157

Doug Freehafer, CSC Senior Mini-Computer Specialist Mail Stop 3PM53 FTS 597-1198

Nancy Coleman, CSC Technical Information Specialist Mail Stop 3PM53 FTS 597-1595

Don Evans, CSC Mini-Computer Specialist Mail Stop 3PM53 FTS 597-9769

GIS Personnel (continued)

Mike Perpiglia, CSC Mini-Computer Specialist Mail Stop 3PM53 FTS 597-1198

Water Management Division

Peter Weber Environmental Protection Specialist Division GIS Coordinator Mail Stop 3WM42 FTS 597-4283

Sumner Crosby Environmental Scientist Ground Water Protection Section Mail Stop 3WM42 FTS 597-3424 1. Non-point Source Pollution from Agricultural Practices

Cindy Greene
Environmental Scientist
Ground Water Protection Section
Mail Stop 3WM42
FTS 597-8399

2. Targeting Areas for Wellhead Protection

Virginia Thompson, Chief Ground Water Protection Section Mail Stop 3WM42 FTS 597-2786

3. Underground Injection Control - Site Discovery

Ronnie M. Davis
Environmental Engineer
Underground Injection Control Section
Mail Stop 3WM43
FTS 597-2923

4. Ground Water Pollution Potential Assessment

Stuart Kerzner, Deputy Chief Drinking Water/Ground Water Protection Branch Mail Stop 3WM40 FTS 597-8826

5. Regional Ground Water Risk Assessment

Peter Weber Environmental Protection Specialist Ground Water Protection Section Mail Stop 3WM43 FTS 597-4283

Sumner Crosby
Environmental Scientist
Ground Water Protection Section
Mail Stop 3WM42
FTS 597-3424

6. Sub-Regional Ground Water Risk Assessment

Peter Weber Environmental Protection Specialist Ground Water Protection Section Mail Stop 3WM43 FTS 597-4283

Sumner Crosby Environmental Scientist Ground Water Protection Section Mail Stop 3WM42 FTS 597-3424

7. Federal Facilities Compliance Initiative

Koge Suto Environmental Engineer General Enforcement Section Mail Stop 3WM52 FTS 597-4462

8. CERCLA Site Discovery

Fred Sturniolo Hydrogeologist Ground Water Protection Section Mail Stop 3WM42 FTS 597-9388

Lorie Acker Environmental Engineer Site Assessment Section Mail Stop 3HW13 FTS 597-3165

9. GIS Automation of Hazard Ranking System, Ground Water Migration Pathway

Fred Sturniolo Hydrogeologist Ground Water Protection Section Mail Stop 3WM42 FTS 597-9388

Sumner Crosby Environmental Scientist Ground Water Protection Section Mail Stop 3WM42 FTS 597-3424

10. Delaware River Water Trends Report

Maggie Passmore Environmental Scientist Philadelphia Operations Section Mail Stop 3ES11 FTS 597-6149

11. Assessing Toxicological Impacts

Jeffrey Burke Regional Total Quality Coordinator Office of Deputy Regional Administrator Mail Stop 3DA00 FTS 597-7808

Susan McDowell Ecologist Wetlands Section Mail Stop 3ES43 FTS 597-0355

12. Monitoring Radon Levels

Lewis Felleisen, Chief Technical Assessment Section Mail Stop 3AT12 FTS 597-8326

13. Air Quality Monitoring

David L. Arnold, Chief Program Planning Section Mail Stop 3AT13 FTS 597-4556

Virginia SO2

Dennis M. Lohman, Chief New Source Review Section Mail Stop 3AT22 FTS 597-3024

Washington D.C. - SO2 Modelling

Alan Cimorelli Environmental Scientist Technical Assessment Section Mail Stop 3AT12 FTS 597-6563

14. Superfund Applications

Berks Sand Pit Superfund Site

Christopher Corbett Remedial Project Manager Central Pennsylvania Section Mail Stop 3HW27 FTS 597-6906

North Penn Areas

Patrick McManus Environmental Engineer Eastern Pennsylvania Section Mail Stop 3HW22 FTS 597-1265

Berkeley Superfund Site

Bruce Rundell Hydrogeologist Technical Support Section Mail Stop 3HW15 FTS 597-1268

15. Oil Spill Emergency Response

Linda Ziegler Environmental Protection Specialist Oil and Title III Section Mail Stop 3HW34 FTS 597-1395

16. Regional Strategic Planning Initiative

Alvin R. Morris, Director Water Management Division Mail Stop 3WM00 FTS 597-9410

Peter Weber Environmental Protection Specialist Ground Water Protection Section Mail Stop 3WM43 FTS 597-4283