U.S. ENVIRONMENTAL PROTECTION AGENCY NATIONAL EUTROPHICATION SURVEY

WORKING PAPER SERIES



REPORT
ON
LAKE POYGAN
WINNEBAGO AND WAUSHARA COUNTIES
WISCONSIN
EPA REGION V
WORKING PAPER No. 45

PACIFIC NORTHWEST ENVIRONMENTAL RESEARCH LABORATORY

An Associate Laboratory of the
NATIONAL ENVIRONMENTAL RESEARCH CENTER - CORVALLIS, OREGON
and

NATIONAL ENVIRONMENTAL RESEARCH CENTER - LAS VEGAS, NEVADA

REPORT
ON
LAKE POYGAN
WINNEBAGO AND WAUSHARA COUNTIES
WISCONSIN
EPA REGION V
WORKING PAPER No. 45

WITH THE COOPERATION OF THE
WISCONSIN DEPARTMENT OF NATURAL RESOURCES
AND THE
WISCONSIN NATIONAL GUARD
NOVEMBER, 1974

CONTENTS

		<u>Page</u>
For	reword	ii
Lis	t of Wisconsin Study Lakes	iv, v
Lak	e and Drainage Area Map	vi
Sec	tions	
I.	Conclusions	1
II.	Introduction	3
III.	Lake and Drainage Basin Characteristics	4
IV.	Lake Water Quality Summary	5
٧.	Nutrient Loadings	10
VI.	Literature Reviewed	15
VII	Appendices	16

FOREWORD

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nation-wide threat of accelerated eutrophication to fresh water lakes and reservoirs.

OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

- a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.
- b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.
- c. With such a transformation, an assessment of the potential for eutrophication control can be made.

LAKE ANALYSIS

In this report, the first stage of evaluation of lake and water-shed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning [§303(e)], water quality criteria/standards review [§303(c)], clean lakes [§314(a,b)], and water quality monitoring [§106 and §305(b)] activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

ACKNOWLEDGMENT

The staff of the National Eutrophication Survey (Office of Research & Development, U. S. Environmental Protection Agency) expresses sincere appreciation to the Wisconsin Department of Natural Resources for professional involvement and to the Wisconsin National Guard for conduct of the tributary sampling phase of the Survey.

Francis H. Schraufnagel, Acting Assistant Director, and Joseph R. Ball of the Bureau of Water Quality, and Donald R. Winter, Lake Rehabilitation Program, provided invaluable lake documentation and counsel during the Survey. Central Office and District Office personnel of the Department of Natural Resources reviewed the preliminary reports and provided critiques most useful in the preparation of this Working Paper series.

Major General James J. Lison, Jr., the Adjutant General of Wisconsin, and Project Officer CW-4 Donald D. Erickson, who directed the volunteer efforts of the Wisconsin National Guardsmen, are also gratefully acknowledged for their assistance to the Survey.

NATIONAL EUTROPHICATION SURVEY

STUDY LAKES

STATE OF WISCONSIN

LAKE NAME	COUNTY
Altoona Beaver Dam Beaver Dam Big Eau Pleine Browns Butte des Morts Butternut Castle Rock Flowage Como Crystal Delavan Eau Claire	Eau Claire Barron Dodge Marathon Racine Winnebago Price, Ashland Juneau Walworth Vilas Walworth Eau Claire
Geneva Grand Green Kegonsa Koshkonong Lac La Belle Middle Nagawicka Oconomowoc Okauchee Petenwell Flowage Pewaukee Pigeon Pine	Walworth Green Lake Green Lake Dane Jefferson, Rock, Dane Waukesha Walworth Waukesha Waukesha Waukesha Juneau Waukesha Waukesha Juneau Waukesha Waukesha Waukesha
Poygan Rock	Winnebago, Waushara Jefferson
Rome Pond Round Shawano	Jefferson, Waukesha Waupaca Shawano

LAKE NAME

Sinnissippi Swan Tainter Tichigan Townline Trout Wapogassett Wausau Willow Winnebago


Wisconsin Wissota Yellow

COUNTY

Dodge Columbia Dunn Racine Oneida Vilas Polk Marathon Oneida

Winnebago, Fond Du Lac,

Calumet Columbia Chippewa Burnett



LAKE POYGAN

STORET NO. 5538

I. CONCLUSIONS

A. Trophic Condition:

Survey data indicate Lake Poygan is eutrophic. This lake is listed in "Problem Lakes of the United States" (Ketelle and Uttormark, 1971).

B. Rate-Limiting Nutrient:

Algal assay results indicate Lake Poygan was nitrogen limited at the time the sample was collected. However, a significant loss of nitrogen occurred in the sample, and the results are not considered reliable (see discussion, pages 8 and 9).

C. Nutrient Controllability:

1. Point sources--During the sampling year, Lake Poygan received a phosphorus load at a rate well in excess of that proposed by Vollenweider (in press) as "dangerous"; i.e., a eutrophic rate (see page 14). Of that load, it is estimated that the point sources included within the Survey limit* contributed less than 1%. If only these point sources were considered, it would be concluded that point-source phosphorus control would not significantly improve the trophic condition of Lake Poygan.

^{*} See Working Paper No. 1, "Survey Methods".

However, there are many point sources of phosphorus in the drainage beyond the 25-mile limit of the Survey*, and a more detailed study is needed to assess the effectiveness of phosphorus control in the drainage.

2. Non-point sources (see page 14)--Except for the Pine River, the phosphorus exports of the streams tributary to Lake Poygan were quite high and are indicative of the unmeasured point sources noted above (assuming the export of the Pine River is representative of non-point sources in the Wolf River drainage).

Whether effective control of the phosphorus now reaching Lake Poygan can be achieved is questionable in view of the rather high drainage area to lake area ratio of 226 to 1. With zero point-source contributions, the mean non-point phosphorus export of the tributaries to the lake would have to be reduced to about 37 lbs/mi²/yr to achieve a loading rate just equal to Vollenweider's eutrophic rate.

^{*} McKersie, et al., 1971.

II. INTRODUCTION

Lake Poygan is the terminus of the Wolf River drainage in northeastern Wisconsin. The entire drainage is in a glaciated part of the State.

Land use in the drainage is predominately agricultural, and dairy plants and canneries are common in the southern part of the drainage (McKersie, et al., op. cit.).

The primary uses of Lake Poygan are recreational, including swimming, boating, and fishing. Game fish present are muskellunge, northern pike, walleyes, largemouth bass, and panfish (Anonymous, 1972).

Public access is provided, and commercial facilities offer services.

III. LAKE AND DRAINAGE BASIN CHARACTERISTICS

- A. Lake Morphometry[†]:
 - 1. Surface area: 10,992 acres.
 - 2. Mean depth: 7 feet.
 - 3. Maximum depth: 11 feet.
 - 4. Volume: 76,944 acre/ft.
 - 5. Mean hydraulic retention time: 13 days.
- B. Tributary and Outlet: (See Appendix A for flow data)
 - 1. Tributaries -

Name	Drainage area*	
Wolf River Willow Creek Pine River Rat River	3,440.0 mi ² 114.0 mi ² 88.4 mi ² 73.6 mi	2,599.2 cfs 91.9 cfs 72.5 cfs 60.2 cfs
Minor tributaries & immediate drainage -	166.8 mi ²	147.9 cfs
Totals	3,882.8 mi ²	2,971.7 cfs

2. Outlet -

Wolf River

3,900.0 mi²** 2,971.7 cfs**

- C. Precipitation***:
 - 1. Year of sampling: 41.1 inches.
 - 2. Mean annual: 28.4 inches.

[†] Ball, 1973.

^{*} Drainage areas are accurate within $\pm 0.5\%$; mean daily flows are accurate within $\pm 40\%$; mean monthly flows are accurate within $\pm 35\%$; and normalized monthly flows are accurate within $\pm 35\%$.

^{**} Includes area of lake; outflow adjusted to equal sum of inflows.
*** See Working Paper No. 1.

IV. LAKE WATER QUALITY SUMMARY

Lake Poygan was sampled three times during the open-water season of 1972 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from three stations on the lake and generally from two depths at each station (see map, page vi). During each visit, a single depth-integrated (near bottom to surface) sample was composited from the stations for phytoplankton identification and enumeration; and during the last visit, a single five-gallon depth-integrated sample was composited for algal assays. Also each time, a depth-integrated sample was collected from each of the stations for chlorophyll <u>a</u> analysis. The maximum depths sampled were 6 feet at station 1, 4 feet at station 2, and 5 feet at station 3.

The results obtained are presented in full in Appendix B, and the data for the fall sampling period, when the lake was essentially well-mixed, are summarized below. Note, however, the Secchi disc summary is based on all values.

For differences in the various parameters at the other sampling times, refer to Appendix B.

A. Physical and chemical characteristics:

FALL VALUES

(11/08/72)

Parameter	Minimum	Mean	<u>Median</u>	<u>Maximum</u>
Temperature (Cent.) Dissolved oxygen (mg/l) Conductivity (µmhos) pH (units) Alkalinity (mg/l) Total P (mg/l) Dissolved P (mg/l) NO ₂ + NO ₃ (mg/l) Ammonia (mg/l)	6.0 10.4 320 7.7 135 0.051 0.025 0.350 0.060	6.1 10.6 327 7.7 137 0.068 0.035 0.412 0.066	6.2 10.4 325 7.7 137 0.071 0.041 0.450 0.070	6.2 10.9 335 7.7 139 0.080 0.043 0.460 0.070
		ALL VALUE	<u>ES</u>	
Secchi disc (inches)	16	19	18	24

B. Biological characteristics:

1. Phytoplankton -

Sampling Date	Dominant Genera	Number per ml
06/22/72	 Melosira Anabaena Cyclotella Cryptomonas Dinobryon Other genera 	3,333 2,536 2,391 1,268 1,196 1,124
	Total	11,848
08/21/72	 Anabaena Melosira Nitzschia Fragilaria Cyclotella Other genera 	7,826 6,848 1,014 870 399 1,376
	Total	18,333
11/08/72	 Melosira Cyclotella Raphidiopsis Dinobryon Navicula Other genera 	1,181 253 205 193 144 928
	Total	2,904

2. Chlorophyll \underline{a} - (Because of instrumentation problems during the 1972 sampling, the following values may be in error by plus or minus 20 percent.)

Station	Chlorophyll <u>a</u>
<u>Number</u>	(μ g /l)
01	26.1
02	18.3
	14.2 27.9
02	36.3
03	24.9
01	11.9
02	5.6
03	9.7
	Number 01 02 03 01 02 03 01 02 03

C. Limiting Nutrient Study:

1. Autoclaved, filtered, and nutrient spiked -

Spike (mg/l)	Ortho P Conc. (mg/l)	Inorganic N Conc. (mg/l)	Maximum yield (mg/l-dry wt.)
Control	0.035	0.390	8.9
0.006 P	0.041	0.390	9.2
0.012 P	0.047	0.390	9.8
0.024 P	0.059	0.390	9.3
0.060 P	0.095	0.390	9.3
0.060 P + 10.0 N	0.095	10.390	44.9
10.0 N	0.035	10.390	13.2

2. Discussion -

The control yield of the assay alga, <u>Selenastrum capri-cornutum</u>, indicates that the potential primary productivity of Lake Poygan was relatively high at the time the sample was collected. The results also indicate the lake was nitrogen limited. However, there was a significant nitrogen loss

(ca 90 μ g/l) in the sample between the time of collection and the time the assay was begun. Had this loss not occurred, it is quite likely the sample would have been phosphorus limited (the lake data indicate a nitrogen/phosphorus ratio of 14/l; i.e., phosphorus limitation would be expected).

The lake data indicate that Lake Poygan was nitrogen limited in June and August. The mean N/P ratios were 8/1 and 5/1, respectively, at those times.

D. Trophic Condition:

Survey data indicate Lake Poygan is eutrophic. Of the 46 Wisconsin lakes studied, 26 had less mean total phosphorus, 20 had less mean inorganic nitrogen, 43 had greater Secchi disc transparency, and 31 had less mean chlorophyll \underline{a} .

Survey limnologists noted algal blooms in progress in June and in August of 1972.

V. NUTRIENT LOADINGS (See Appendix C for data)

For the determination of nutrient loadings, the Wisconsin National Guard collected a monthly near-surface grab sample from each of the tributary sites indicated on the map (page vi), except for the high runoff months of April and May when two samples were collected. Sampling was begun in September, 1972, and was completed in August, 1973.

Through an interagency agreement, stream flow estimates for the year of sampling and a "normalized" or average year were provided by the Wisconsin District Office of the U.S. Geological Survey for the tributary sites nearest the lake.

In this report, nutrient loads for sampled tributaries were determined by using a modification of a U.S. Geological Survey computer program for calculating stream loadings*. Nutrient loadings for unsampled "minor tributaries and immediate drainage" ("ZZ" of U.S.G.S.) were calculated using the mean concentrations in the Pine River at station C-2 and the mean ZZ flow.

None of the waste sources listed in the following table were sampled. Loads attributed to the communities are estimates (2.5 lbs P and 9.4 lbs N/capita/year*) and are based on reports of failing septic tank systems. Stream loads do not include community loads.

As far as is known, the dairy plants listed do not contribute nutrients to the drainage.

^{*} See Working Paper No. 1.

A. Waste Sources:

Known community* -

Name	Pop. Served**	Treatment	Mean Flow (mgd)	Receiving Water
Poy Sippi SD	450	Septic tanks	?	Pine River
Fremont	330	Septic tanks	?	Wolf River
Winchester	665	Septic tanks	?	Rat River
N. Lake Poygan SD #1	300	Septic tanks	?	Lake Poygan

2. Industrial* -

<u>Name</u>	Product	Treatment	Mean Flow (mgd)	Receiving Water	
Silver- field Cheese Co Fremont	cheese	land disposal	0.0125*	Wolf River	(?)
Daisy Dairy Co. Poy Sippi	cheese ,	land disposal	?	Pine River	(?)

^{*} McKersie, et al., 1971. ** Anonymous, 1971.

B. Annual Total Phosphorus Loading - Average Year:

1. Inputs -

Sou	<u>rce</u>	lbs P/	% of total
a.	Tributaries (non-point load)	-	
	Wolf River Willow Creek Pine River Rat River	484,090 15,110 4,950 22,050	88.9 2.8 0.9 4.1
b.	Minor tributaries & immediate drainage (non-point load) -	e 11,940	2.2
с.	Known community systems -		
	Poy Sippi Fremont Winchester N. Lake Poygan SD #1	1,120 820 1,660 750	0.2 0.1 0.3 0.1
d.	Septic tanks* -	170	<0.1
e.	Known industrial - None	-	-
f.	Direct precipitation** -	1,710	0.3
	Total	544,370	100.0

2. Outputs -

Lake outlet - Wolf River (inlet to L. Butte des Morts) 527,330

3. Net annual P accumulation - 17,040 lbs.

^{*} Estimated 270 dwelling on lakeshore; see Working Paper No. 1. ** See Working Paper No. 1.

C. Annual Total Nitrogen Loading - Average Year:

1. Inputs -

Sou	<u>irce</u>	lbs N/ yr	% of total
a.	Tributaries (non-point load	d) -	
	Wolf River Willow Creek Pine River Rat River	7,619,760 338,350 328,100 245,560	82.5 3.7 3.6 2.7
b.	Minor tributaries & immedia drainage (non-point load)		6.3
с.	Known community systems -		
	Poy Sippi SD Fremont Winchester N. Lake Poygan SD #1	4,230 3,100 6,250 2,820	<0.1 <0.1 <0.1 <0.1
d.	Septic tanks* -	6,340	0.1
e.	Known industrial - None	-	-
f.	Direct precipitation** -	105,900	1.1
	Total	9,241,270	100.0

2. Outputs -

Lake outlet - Wolf River (inlet to L. Butte des Morts) 9,268,300

3. Net annual N loss - 27,030 lbs.

^{*} Estimated 270 dwellings on lakeshore; see Working Paper No. 1. ** See Working Paper No. 1.

D. Mean Annual Non-point Nutrient Export by Subdrainage Area:

Tributary	<u>lbs P/mi²/yr</u>	lbs N/mi ² /yr
Wolf River	141	2,215
Willow Creek	133	2,970
Pine River	56	3,712
Rat River	300	3,336

E. Yearly Loading Rates:

In the following table, the existing phosphorus loading rates are compared to those proposed by Vollenweider (in press). Essentially, his "dangerous" rate is the rate at which the receiving waters would become eutrophic or remain eutrophic; his "permissible" rate is that which would result in the receiving water remaining oligotrophic or becoming oligotrophic if morphometry permitted. A mesotrophic rate would be considered one between "dangerous" and "permissible".

	Tota	1 Phosphorus	Total Nitrogen		
Units	Total	Accumulated	Total	Accumualted	
lbs/acre/yr grams/m²/yr	49.5	1.6	840.1	loss*	
grams/m²/yr	5.55	0.17	94.2	-	

Vollepweider loading rates for phosphorus (g/m²/yr) based on mean depth and mean hydraulic retention time of Lake Poygan:

"Dangerous" (eutrophic rate) 1.48
"Permissible" (oligotrophic rate) 0.74

^{*} There was an apparent loss of nitrogen during the sampling year. This may have been due to nitrogen fixation in the lake, solubilization of previously sedimented nitrogen, recharge with nitrogen-rich ground water, unknown and unsampled point sources discharging directly to the lake, or underestimation of the nitrogen loads from the point sources. Whatever, the cause, a similar loss has occurred at Shagawa Lake, Minnesota, which has been intensively studied by EPA's National Eutrophication Research and Lake Restoration Branch.

VI. LITERATURE REVIEWED

- Anonymous, 1971. Inventory of municipal waste facilities. Publ. OWP-1, Vol. 5, EPA, Washington.
- Anonymous, 1972. Wisconsin lakes. Publ. 218-72, Dept. of Natural Resources, Madison.
- Ball, Joseph R., 1973. Personal communication (morphometry of Lake Poygan). Dept. of Natural Resources, Madison.
- Ketelle, Martha J., and Paul D. Uttormark, 1971. Problem lakes in the United States. EPA Water Poll. Contr. Res. Series, Proj. 16010 EHR.
- Lueschow, Lloyd A., 1972. Biology and control of selected aquatic nuisances in recreational waters. Tech. Bull, #57, Dept. of Natural Resources, Madison.
- McKersie, Jerome R., Robert M. Krill, James L. Lissack, and Ronald K. Krueger, 1971. Wolf River pollution investigation survey. Dept. of Natural Resources, Madison.
- Vollenweider, Richard A., (in press). Input-output models. Schweiz. A. Hydrol.

VII. APPENDICES

APPENDIX A

TRIBUTARY FLOW DATA

LAKE CODE 5538 LAKE POYGAN

TOTAL DRAINAGE AREA OF LAKE 3900.00

•	SUB-DRAINAGI	Ε					NO	HALIZED	FLOWS					
TRIBUTARY	AREA	JAN	FEB	MAR	APR	YAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
5538A1	3900.00	1600.00	1800.00	3200.00	7100.00	4400.00	3700.00	2100.00	1900.00	2100.00	2220.00	2400.00	2000.00	2875.12
5538A3	3440.00	1400.00	1600.00	3000.00	6200.00	4000.00	3200.00	2100.00	1800.00	2100.00	1900.00	2200.00	1700.00	2599.18
553891	114.00	56.00	62.00	180.00	140.00	100.00	90.00	70.00	59.00	67.00	96.00	100.00	81.00	91.91
5538C1	88.40	44.00	48.00	150.00	100.00	78.00	73.00	52.00	46.00	52.00	77.00	83.00	65.00	72.48
553801	73.60	37.00	40.00	120.00	85.00	64.00	60.00	43.00	38.00	43.00	66.00	71.00	54.00	60-20
5538ZZ	184.00	89.00	97.00	270.00	240.00	170.00	160.00	110.00	96.00	110.00	150.00	160.00	120.00	147.87

SUMMARY

TOTAL DRAINAGE AREA OF LAKE = 3900.00 TOTAL FLOW IN = 35662.00 SUM OF SUB-DRAINAGE AREAS = 3900.00 TOTAL FLOW OUT = 34520.00

MEAN MONTHLY FLOWS AND DAILY FLOWS

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
5538A1	9	72	3800.00	23	3000.00				
	10	72	5400.00	15	4900.00				
	11	72	4900.00	12	5000.00				
	12	72	3000.00	10	2800.00				
	1	73	3900.00						
	2	73	3100.00	11	4000.00				
	2 3	73	11000.00	11	14000.00				
	4	73	9700.00	1	9400.00	14	8000.00		
	4 5	73	11000.00	5	12000.00	20	10000.00		
		73	7100.00	26	4400.00				
	6 7	73	2500.00	14	2300.00				
	8	73	1900.00	11	2100.00				
5538A3	9	72	3000.00	23	2100.00				
	10	72	4600.00	14	4400.00				
	11	72	3700.00	12	4300.00				
	12	72	2400.00	10	2500.00				
	1	73	2900.00						
	2	73	2300.00	11	2500.00				
	2 3	73	10000.00	11	15800.00				
	4	73	8600.00	1	7200.00	14	7200.00		
	5	73	9900.00	6	1100.00	20	8200.00		
	6	73	6500.00	26	4000.00				
	6 7	73	2400.00	14	2200.00				
	8	73	1900.00	11	2000.00				

LAKE CODE 5538 LAKE POYGAN

MEAN MONTHLY FLOWS AND DAILY FLOWS

rean .	HOWING	COWS AN							
TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
553881	9	72	96.00	23	68.00				
	10	72	220.00	14	220.00				
	11	72	180.00	12	200.00				
	12	72	110.00	2	110.00				
	1	73	180.00						
	2	73	140.00	4	190.00	•			
	3	73	670.00	4	120.00				
	4	73	200.00	14	160.00	15	160.00	19	210.00
	5 6	73	260.00	6	280.00	27	180.00		
	6	73	190.00	30	110.00				
	7	73	76.00	23	57.00				
	8	73	66.00	4	62.00				
5538C1	9	72	74.00	24	52.00				
	10	72	180.00	14	170.00				
	11	72	150.00	12	160.00				
	12	72	88.00	2	88.00				
	1	73	140.00						
	2	73	120.00	4	160.00				
	2 3 4	73 73	540.00	4	99.00		150.00		150.00
	-	73 73	150.00	15	120.00	19	150.00	28	150.00
	5 6	73	200.00	6	210.00	27	130.00		
	7	73 73	150.00	30	86.00				
	é	73	58.00	23	44.00				
5538D1	9	72	51.00	4 23	48.00				
222001	10	72	62.00 150.00	14	44.00 150.00				
	11	72	120.00	12	140.00				
	iż	72	75.00	10	78.00				
		73	120.00	10	70.00				
	5 J	73	100.00	11	110.00				
	3	73	460.00	ii	660.00				
	4	73	120.00	i	98.00	14	98.00		
	5	73	160.00	ė	170.00	27	110.00		
	6	73	120.00	26	78.00		110000		
	7	73	49.00	14	46.00				
	8	73	43.00	11	45.00	0	0.0		
5538ZZ	9	72	160.00	23	110.00	•			
	10	72	340.00	14	330.00				
	11	72	280.00	12	310.00				
	12	72	170.00	10	170.00				
	1	73	260.00						
	2	73	210.00	4	290.00	11	230.00		
	3	73	990.00	4	180.00	11	1400.00		
	4	73	330.00	1	270.00	14	270.00	15	270.00
	5 6	73	440.00	6	460.00	20	360.00	27	290.00
		73	320.00	26	200.00	30	190.00	28	350.00
	7	73	120.00	14	110.00	23	94.00		
	8	73	110.00	4	100.00	11	110.00		

APPENDIX B

PHYSICAL and CHEMICAL DATA

553801 44 07 12.0 088 53 00.0 POYGAN LAKE 55 WISCONSIN

					11EP	ALES	2111202 0005 FEET DEPTH				
	OF		0 0 300 D0 MG/L	00077 TRANSP SECCHI INCHES	00094 CNDUCTVY FIELD MICROMHO	00400 PH SU	00410 T ALK CACO3 MG/L	00630 NOZ&NO3 N-TOTAL MG/L	00610 NH3-N TOTAL MG/L	00665 PHOS-TOT MG/L P	00666 PH05-DIS MG/L P
72/06/22	07 20 0000	18.5	5.8	24	285	7.90	148	0.180	0.100	0.046	0.026
	07 20 0006	18.6	9.6		280	8.10	145	0.180	0.100	0.110	0.027
	09 30 0000			19	280	8.65	134	0.060	0.040	0.048	0.018
12/11/08	08 00 0000			16	320	7.70	137	0.350	0.070	0.051	0.026
	08 00 0004	6.0	10.9		320	7.70	135	0.350	0.070	0.063	0.025

DATE FROM	TII		DEPTH	32217 CHLRPHYL A
TO	DA'	Y	FEET	UG/L
72/06/22	07	20	0000	26.13
72/08/21	09	30	0000	27.9J
72/11/08	08	00	0000	11.6J

J VALUE KNOWN TO BE IN ERROR

553802 44 09 24.0 088 49 00.0 POYGAN LAKE 55 WISCONSIN

						11EP	ALES		1202 FEET DEP	тн	
FROM	TIME DEPTH OF DAY FEET	00010 Water Temp Cent	00300 D0 MG/L	00077 TRANSP SECCHI INCHES	00094 CNDUCTVY FIELD MICROMHO	00400 PH SU	00410 T ALK CACO3 MG/L	00630 NO26NO3 N-TOTAL	00610 NH3-N TOTAL	00665 PHOS-TOT	00666 PHOS-DIS
-				11101123	HILKOHIO	30	HUIL	MG/L	MG/L	MG/L P	MG/L P
	08 40 0000 09 05 0000	9.4	8.7	18	285	8.30	155	0.220	0.110	0.163	0.050
12,00,21	09 05 0004	23.7	8.4	17	360 350	8.60	129	0.070	0.070	0.065	0.026
72/11/08	07 40 0000			17	358 325	8.60 7.70	130	0.060	0.050	0.075	- 0.025
	07 40 0004	6.2	10.4		325	7.70	139	0.460	0.070	0.080	0.043

DATE FROM TO	Of	-	DEPTH FEET	32217 CHLRPHYL A UG/L
72/06/22 72/08/21				18.3. 36.3.
72/11/08	07	40	0000	5•6J

J VALUE KNOWN TO BE IN ERROR

553803 44 08 18.0 088 45 54.0 POYGAN LAKE 55 WISCONSIN

							11EP/ 3	ALES		1202 FEET DEP	тн	
DATE FROM	TIME OF	DEPTH	00010 Water Temp	0 0300 D0	00077 Transp Secchi	00094 CNDUCTVY FIELD	00400 PH	00410 T ALK CACO3	00630 N-3003 N-TOTAL	00610 NH3-N Total	00665 PHOS-TOT	00666 PHOS-DIS
TO	DAY	FEET	CENT	MG/L	INCHES	MICROMHO	รบ	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/06/22	08 5	0 0000	18.3	8.8		2 7 5	8.40	150	0.080	0.100	0.062	0.024
_	08 5	0 0005	18.3	9.9		270	8.60	149	0.070	0.100	0.088	0.024
72/08/21	08 4	0 0000			18	270	8.50	130	0.050	0.050	0.077	0.023
	08 4	0 0004	23.7	7.4		275	8.50	131	0.060	0.070	0.079	0.026
72/11/08	07 20	0 0000			21	335	7.70	138	0.450	0.060	0.071	0.042
	07 2	0 0004	6.2	10.4		335	7.70	137	0.450	0.060	0.074	0.041

DATE		DEPTH	32217 CHLRPHYL
FROM	OF		A
10	DAY	FEET	UG/L
72/06/22	08 5	0000	14.2J
72/08/21	08 4	0 0000	24.9J
72/11/08			9.7J

J VALUE KNOWN TO BE IN ERROR

APPENDIX C

TRIBUTARY DATA

S538A2 LS5538A2

44 15 30.0 088 52 00.0

WOLF RIVER

55 15 WEYAUWEGA

T/LAKE POYGAN

US 10 PRDG ABOVE FREMONT STP

11EPALES 2111204

4 0000 FEET DEPTH

DATE FROM	TIME OF	DEPTH	00630 006300 0063003 00630	00625 TOT KJEL N	00610 NH3-N TOTAL	00671 PHOS-DIS ORTHO	00665 PH0S-TOT
т О	DAY	FEET	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/09/23	11 39	5	0.440	0.850	0.210	0.082	0.096
72/10/14	14 20)	0.240	0.900	0.098	0.046	0.096
72/11/12	14 10)	0.366	0.750	0.024	0.035	0.061
73/01/13	14 30)	0.870	1.100	J.126	0.048	0.080
73/02/11	14 19	5	0.790	0.880	0.071	0.038	0.070
73/03/11	14 30)	0.630	1.470	0.410	0.089	0.188
73/04/01	09 50)	0.160	0.660	0.008	0.010	0.035
73/04/14	14 20)	0.110	0.780	0.014	0.013	0.030
73/05/06	09 30)	0.092	1.380	0.078	0.036	0.060
73/05/20	14 30)	0.058	1.540	0.042	0.040	0.080
73/05/26	10 19	5	0.280	1.470	0.072	0.069	0.120
73/07/14	14 45	5	0.147	1.150	0.023	0.016	0.095
73/08/11	11 00)	0.330	0.840	0.014	0.023	0.090

5538A3 LS5538A3
44 13 00.0 088 55 00.0
WOLF RIVER
55 15 POY SIPPI
T/LAKE POYGAN
BANK ON HWY HH BELO FREMONTSTP
11FPALES 2111204
4 0000 FEET DEPTH

			00630	00625	00610	00671	00665
DATE	TIME	DEPTH	K0N%20N	TOT KJEL	NH3-N	PHOS-DIS	PHOS-TOT
FROM	OF		N-TOTAL	N	TOTAL	CHTRO	
CT	DAY	FEET	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/09/23			0.190	0.650	0.105	0.015	0.105
72/10/14	14 4	0	0.234	0.950	0.088	0.046	0.096
72/11/12	14 3	0	0.420	1.050	0.022	0.035	0.061
72/12/10	11 3	0	1.080	0.600	0.076	0.038	0.063
73/01/07	14 4	5	0.990	1.050	0.120	0.042	0.070
73/02/11	14 3	0	0.810	0.900	0.080	0.038	0.070
73/03/11	14 1	0	0.660	1.500	0.410	0.095	0.290
73/04/01	10 0	0	0.170	1.760	0.024	0.010	0.040
73/04/14	14 4	0	0.095	0.640	0.016	0.011	0.030
73/05/06	09 4	5	0.077	1.200	0.033	0.023	0.055
73/05/20	14 4	5	0.044	2.000	0.054	9.037	0.075
73/06/26	10 0	0	0.280	0.780	0.039	0.067	0.100
73/07/14	15 0	0	0.189	0.810	0.030	0.019	0.105
73/08/11	11 1	5	0.370	0.690	0.012	0.028	0.095

553881 LS553881
44 06 00.0 088 56 00.0
WILLOW CREEK
55 15 POY SIPPI
T/LAKE POYGAN
CO HWY D BRUG SE OF POY SIPPI
11EPALES 2111204
4 0000 FEET DEPTH

DATE FROM	TIME OF	DEPTH	0630 8002500 1ATOT-N	00625 TOT KUFL N	OOK1O NH3-H TUTAL	00671 PHOS-DIS ORIHO	00665 PH05-TOT
10	DAY	FEET	MG/L	46/L	MG/L	MG/L P	MG/L P
72/09/24	16 00	ס	0.056	1.650	0.115	0.089	0.110
72/10/14	10 30)	0.500	1.550	0.180	0.045	0.092
72/11/13	13 5	5	0.440	1.200	0.120	0.035	0.063
72/12/02	10 1	5	0.820	0.635	0.032	0.014	0.029
73/01/07	09 00)	0.880	1.200	0.160	0.022	0.045
73/02/04	14 00)	0.770	1.150	0.168	0.025	0.060
73/03/04	11 30)	1.260	1.980	0.730	0.075	0.145
73/04/14	08 40)	0.150	1.230	0.100	0.040	0.045
73/04/15	12 30)	0.164	2.600	0.138	0.018	0.045
73/04/19	15 30)	0.035	1.800	0.063	0.018	0.060
73/05/06	08 49	5	0.050	2.100	0.040	0.030	0.075
73/05/27	13 49	5	0.045	1.500	0.054	0 • 0 3 ਦ	0.119
73/06/30	11 30)	0.079	0.730	0.087	0.082	0.120
73/07/23	13 50)	0.037	0.540	0.031	0.033	0.070
73/03/04	03 4	5	0.030	1.800	0.064	0.026	0.065

5538C1 LS553AC1
44 07 30.0 088 57 30.0
PINE RIVER
55 15 POY SIPPI
T/LAKE POYGAN
N-S RD XING SE OF POY SIPPIBELOW STP
11EPALES 2111204
4 0000 FEET DEPTH

DATE	TIME	DEPTH	· · · 	00625 TOT KJEL	00610 NH3-N	00671 PHOS-DIS	00665 PHOS-TOT
FROM	OF		N-TOTAL	N	TOTAL	ORTHO	
ΤO	DAY	FEET	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/09/24	15 16		0.065	1.000	0.140	0.022	0.044
					· · · -		
72/10/14		-	0.740	0.575	0.066	0.017	0.036
72/11/10	13 00)	1.320	1.300	0.065	0.015	0.032
72/12/02	10 30)	1.620	0.430	0.025	0.011	0.025
73/01/07	09 15	5	1.800	0.790	0.072	0.015	0.050
73/02/04	14 30)	1.420	0.800	0.084	0.022	0.050
73/03/04	11 30)	1.440	1.980	0.380	0.084	0.130
73/04/15	11 59	5	0.960	1.600	0.091	0.012	0.035
73/04/19	15 10)	1.000	1.260	0.031	0.024	0.055
73/04/28	08 15	5	1.120	2.520	0.100	0.015	0.035
73/05/06	08 10)	1.040	1.050	0.042	0.019	0.045
73/06/30	11 10)	0.950	0.580	0.040	0.018	0.040
73/07/23	13 29	5	0.950	0.390	0.046	0.016	0.040
73/08/04	08 30)	1.000	1.800	0.160	0.015	0.040

5538C? LS5538C2
44 08 00.0 089 00 00.0
PINE RIVER
55 15 POY SIPPI
T/LAKE POYGAN
ST HWY 49 BRDG ABOV POY SIPPI STP
11EPALES 2111204
4 0000 FEET DEPTH

DATE FROM	TIME DEPTH OF	00630 NO28NO3 N-TOTAL	00625 TOT KJEL N	00610 NH3-N Total	00671 PHOS-DIS ORTHO	00665 PHOS-TOT
10	DAY FEET	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/09/24 72/10/14	14 45 09 45	0.052 0.860	1.050 0.550	0.154 0.073	0.013 0.012	0.037 0.037
72/11/10	12 10	1.340	0.610	0.033	0.011	0.030
72/12/02	10 45	1.640	0.560	0.024	0.009	0.022
73/01/07	09 30	1.900	0.720	0.071	0.011	0.030
73/02/04	15 00	1.460	1.600	0.294	0.015	0.035
73/03/04	11 50	1.440	1.320	0.280	0.066	0.110
73/04/15	11 30	1.000	1.500	0.052	0.009	0.030
-	15 00	1.020	1.400	0.060	0.009	0.040
73/04/28		1.140	0.980	0.023	0.010	0.020
-	08 30	1.100	1.050	0.072	0.016	0.040
73/05/27	13 30	0.810	1.300	0.039	0.022	0.055
73/06/30	10 45	1.000	0.630	0.038	0.015	0.040
73/07/23	13 05	1.000	0.480	0.046	0.012	0.040
73/08/04	08 15	1.060	1.980	0.066	0.014	0.055

5538D1 LS5538D1 44 12 00.0 08A 46 30.0 RAT RIVER 55 15 POY SIPPI T/LAKE POYGAN CO HWY MM BRDG S AND E OF ZITTAN 11EPALES 2111204 4 C000 FEET DEPTH

			00630	00625	00610	00671	00665
DATE	TIME	DEPTH	E ON & SON	TOT KJEL	NH3-N	PHOS-DIS	PHOS-TOT
FROM	0F		N-TOTAL	N	TOTAL	ORTHO	
TO	DAY	FFET	MG/L	MG/L	MG/L	MG/L P	MG/L P
72/09/23	11 19	5	0.210	2-300	0.074	0.126	0.154
72/10/14	14 00	0	0.130	1.750	0.204	0.058	0.088
72/11/12	14 00	0	0.044	1.600	0.060	0.020	0.046
72/12/10	10 25	5	0.200	2.100	0.168	0.021	0.063
73/01/13	14 00	0	0.100	2.600	0.450	0.115	0.300
73/02/11	13 59	5	0.063		0.110	0.075	
73/03/11	13 4	5	0.720	1.440	0.220	0.052	0.095
73/04/01	09 30	0	0.013	1.540	0.034	0.016	0.065
73/04/14	13 59	5	0.061	1.300	0.022	0.012	0.035
73/05/06	09 19	5	0.046	1.500	0.028	0.046	0.085
73/05/27	14 19	5	0.019	1.680	0.029	0.160	0.230
73/06/26	10 30	D	0.022	2.450	0.080	0.340	0.470
73/07/14	14 30	0	0.012	2.600	0.046	0.310	0.440
73/08/11	10 39	5	0.010K	2.200	0.039	0.273	0.370

K VALUE KNOWN TO HE LESS THAN INDICATED