

**U.S. ENVIRONMENTAL PROTECTION AGENCY  
NATIONAL EUTROPHICATION SURVEY**

**WORKING PAPER SERIES**



REPORT  
ON  
BEAVER DAM LAKE (SOUTH BASIN)  
BARRON COUNTY  
WISCONSIN  
EPA REGION V  
WORKING PAPER No. 69

**PACIFIC NORTHWEST ENVIRONMENTAL RESEARCH LABORATORY**

**An Associate Laboratory of the**

**NATIONAL ENVIRONMENTAL RESEARCH CENTER - CORVALLIS, OREGON**

**and**

**NATIONAL ENVIRONMENTAL RESEARCH CENTER - LAS VEGAS, NEVADA**

REPORT  
ON  
BEAVER DAM LAKE (SOUTH BASIN)  
BARRON COUNTY  
WISCONSIN  
EPA REGION V  
WORKING PAPER No. 69

WITH THE COOPERATION OF THE  
WISCONSIN DEPARTMENT OF NATURAL RESOURCES  
AND THE  
WISCONSIN NATIONAL GUARD  
JUNE, 1975

## CONTENTS

	<u>Page</u>
Foreword	ii
List of Wisconsin Study Lakes	iv, v
Lake and Drainage Area Map	vi
 <u>Sections</u>	
I. Introduction	1
II. Conclusions	1
III. Lake Characteristics	3
IV. Lake Water Quality Summary	4
V. Literature Reviewed	8
VI. Appendix	9

## F O R E W O R D

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nation-wide threat of accelerated eutrophication to fresh water lakes and reservoirs.

### OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

### ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.

b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.

c. With such a transformation, an assessment of the potential for eutrophication control can be made.

### LAKE ANALYSIS\*

In this report, the first stage of evaluation of lake and watershed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning [§303(e)], water quality criteria/standards review [§303(c)], clean lakes [§314(a,b)], and water quality monitoring [§106 and §305(b)] activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

---

\* The lake discussed in this report was included in the National Eutrophication Survey as a water body of interest to the Wisconsin Department of Natural Resources. Tributaries and nutrient sources were not sampled, and this report relates only to the data obtained from lake sampling.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

#### ACKNOWLEDGMENT

The staff of the National Eutrophication Survey (Office of Research & Development, U. S. Environmental Protection Agency) expresses sincere appreciation to the Wisconsin Department of Natural Resources for professional involvement and to the Wisconsin National Guard for conducting the tributary sampling phase of the Survey.

Francis H. Schraufnagel, Acting Assistant Director, and Joseph R. Ball of the Bureau of Water Quality, and Donald R. Winter, Lake Rehabilitation Program, provided invaluable lake documentation and counsel during the Survey. Central Office and District Office personnel of the Department of Natural Resources reviewed the preliminary reports and provided critiques most useful in the preparation of this Working Paper series.

Major General James J. Lison, Jr., the Adjutant General of Wisconsin, and Project Officer CW-4 Donald D. Erickson, who directed the volunteer efforts of the Wisconsin National Guardsmen, are also gratefully acknowledged for their assistance to the Survey.

## NATIONAL EUTROPHICATION SURVEY

## STUDY LAKES


STATE OF WISCONSIN

<u>LAKE NAME</u>	<u>COUNTY</u>
Altoona	Eau Claire
Beaver Dam	Barron
Beaver Dam	Dodge
Big Eau Pleine	Marathon
Browns	Racine
Butte des Morts	Winnebago
Butternut	Price, Ashland
Castle Rock Flowage	Juneau
Como	Walworth
Crystal	Vilas
Delavan	Walworth
Eau Claire	Eau Claire
Geneva	Walworth
Grand	Green Lake
Green	Green Lake
Kegonsa	Dane
Koshkonong	Jefferson, Rock, Dane
Lac La Belle	Waukesha
Middle	Walworth
Nagawicka	Waukesha
Oconomowoc	Waukesha
Okauchee	Waukesha
Petenwell Flowage	Juneau
Pewaukee	Waukesha
Pigeon	Waupaca
Pine	Waukesha
Poygan	Winnebago, Waushara
Rock	Jefferson
Rome Pond	Jefferson, Waukesha
Round	Waupaca
Shawano	Shawano
Sinnissippi	Dodge

LAKE NAMECOUNTY

Swan  
Tainter  
Tichigan  
Townline  
Trout  
Wapogasset  
Wausau  
Willow  
Winnebago  
  
Wisconsin  
Wissota  
Yellow

Columbia  
Dunn  
Racine  
Oneida  
Vilas  
Polk  
Marathon  
Oneida  
Winnebago, Fond Du Lac,  
Calumet  
Columbia  
Chippewa  
Burnett





## BEAVER DAM LAKE (SOUTH BASIN)

STORET NO. 5503

### I. INTRODUCTION

The South Basin of Beaver Dam Lake was included in the National Eutrophication Survey as a water body of interest to the Wisconsin Department of Natural Resources. Tributaries and nutrient sources were not sampled, and this report only relates to the lake sampling data.

The wastewater treatment plant serving the City of Cumberland discharges to the South Basin, and the Stokely-Van Camp cannery discharges chlorinated cooling water there also (McKersie, et al., 1971). At this time, the City of Cumberland is considering alternate methods of effluent treatment or disposal; the Stokely-Van Camp waste treatment facilities are considered to be satisfactory (Schraufnagel, 1975).

### II. CONCLUSIONS

#### A. Trophic Condition:

Survey data indicate the South Basin of Beaver Dam Lake is eutrophic. Of the 46 Wisconsin lakes sampled, 43 had less mean total phosphorus, 42 had less mean dissolved phosphorus, ten had less mean inorganic nitrogen, only two had less mean chlorophyll a, and 44 had greater mean Secchi disc transparency.

B. Rate-Limiting Nutrient:

The algal assay results indicate the South Basin was nitrogen limited at the time the sample was taken (11/03/72). The lake data indicate nitrogen limitation at the other sampling times as well.

### III. LAKE CHARACTERISTICS

#### A. Lake Morphometry (entire lake)\*:

1. Surface area: 1,112 acres.
2. Mean depth: 32.1 feet.
3. Maximum depth: 106 feet.
4. Volume: 35,700 acre-feet.

#### B. Precipitation\*\*:

1. Year of sampling: 33.4 inches.
2. Mean annual: 30.1 inches.

---

\* Ball, 1973.

\*\* See Working Paper No. 1, "Survey Methods, 1972".

#### IV. LAKE WATER QUALITY SUMMARY

The South Basin of Beaver Dam Lake was sampled three times during the open-water season of 1972 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from a number of depths at a single station on the lake (see map, page vi). During each visit, a single depth-integrated (near bottom to surface) sample was collected for phytoplankton identification and enumeration, and a similar sample was taken for chlorophyll a analysis. During the last visit, a single five-gallon depth-integrated sample was collected for algal assays. The maximum depth sampled was 7 feet.

The results obtained are presented in full in Appendix A, and the data for the fall sampling period, when the lake essentially was well-mixed, are summarized below. Note, however, the Secchi disc summary is based on all values.

For differences in the various parameters at the other sampling times, refer to Appendix A.

## A. Physical and chemical characteristics:

FALL VALUES

(11/03/72)

<u>Parameter</u>	<u>Surface Sample Only</u>
Temperature (Cent.)	4.2
Dissolved oxygen (mg/l)	13.8
Conductivity ( $\mu$ mhos)	170
pH (units)	9.2
Alkalinity (mg/l)	49
Total P (mg/l)	0.900
Dissolved P (mg/l)	0.680
NO <sub>2</sub> + NO <sub>3</sub> (mg/l)	0.520
Ammonia (mg/l)	0.120

ALL VALUES

	<u>Minimum</u>	<u>Mean</u>	<u>Median</u>	<u>Maximum</u>
Secchi disc (inches)	22	35	36	48

## B. Biological characteristics:

## 1. Phytoplankton -

<u>Sampling Date</u>	<u>Dominant Genera</u>	<u>Number per ml</u>
06/26/72	1. Melosira	2,892
	2. Microcystis	1,988
	3. Fragilaria	1,295
	4. Scenedesmus	452
	5. Anabaena	210
	Other genera	<u>1,325</u>
	Total	8,162
08/26/72	1. Scenedesmus	10,688
	2. Anabaena	6,377
	3. Gloeocapsa	652
	4. Cryptomonas	471
	Other genera	<u>1,087</u>
	Total	19,275
11/03/72	1. Cyclotella	6,244
	2. Franceia	4,072
	3. Scenedesmus	1,629
	4. Synedra	1,584
	5. Microcystis	1,222
	Other genera	<u>4,616</u>
	Total	19,367

2. Chlorophyll a -

(Because of instrumentation problems during the 1972 sampling, the following values may be in error by plus or minus 20 percent.)

<u>Sampling Date</u>	<u>Station Number</u>	<u>Chlorophyll <u>a</u> (<math>\mu\text{g/l}</math>)</u>
06/26/72	01	15.0
08/26/72	01	42.0
11/03/72	01	152.9

## C. Limiting Nutrient Study:

## 1. Autoclaved, filtered, and nutrient spiked -

<u>Spike (mg/l)</u>	<u>Ortho P Conc. (mg/l)</u>	<u>Inorganic N Conc. (mg/l)</u>	<u>Maximum yield (mg/l-dry wt.)</u>
Control	0.770	0.650	23.3
0.006 P	0.776	0.650	22.5
0.012 P	0.782	0.650	22.9
0.024 P	0.794	0.650	24.8
0.060 P	0.830	0.650	23.6
0.060 P + 10.0 N	0.830	10.650	44.8
10.0 N	0.770	10.650	40.9

## 2. Discussion -

The control yield of the assay alga, Selenastrum capricornutum, indicates that the potential primary productivity of South Basin of Beaver Dam Lake was very high at the time the sample was collected (11/03/72). Also, the lack of yield response to increased levels of orthophosphorus and the marked increase in yield when only nitrogen was added show the lake was nitrogen limited at that time.

The lake data indicate nitrogen limitation at the other sampling times as well; the N/P ratio in June was 6/1 and in August was less than 1/1, and nitrogen limitation would be expected.

## V. LITERATURE REVIEWED

Ball, Joseph R., 1973. Personal communication (lake morphometry).  
WI Dept. Nat. Resources, Madison.

McKersie, Jerome R., Robert M. Krill, Charles Kozel, and Danny J.  
Ryan; 1971. Lower Chippewa River pollution investigation  
survey. WI Dept. Nat. Resources, Madison.

Sather, LaVerne M., and C. W. Threinen, 1964. Surface water resources  
of Barron County. WI Cons. Dept., Madison.

Schraufnagel, Francis H., 1975. Personal communication (status of  
waste treatment facilities). WI Dept. Nat. Resources, Madison.



## VI. Appendix

### APPENDIX A

#### PHYSICAL and CHEMICAL DATA

STORET RETRIEVAL DATE 74/11/01

553301  
45 31 48.0 092 01 00.0  
BEAVER DAM LAKE  
55 WISCONSIN

11EPALES  
3

2111202  
0007 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00010 WATER TEMP CENT	00300 DO MG/L	00077 TRANSP SECCHI INCHES	00094 CONDUCTVY FIELD MICROMHO	00400 PH SU	00410 T ALK CAC03 MG/L	00630 NO2&N03 N-TOTAL MG/L	00610 NH3-N TOTAL MG/L	00665 PHOS-TOT MG/L P	00666 PHOS-DIS MG/L P
72/06/26	06 05	0000	19.8		48	140	8.10	50	0.050	0.060	0.079	0.023
	06 05	0007	19.2	7.5		150	7.90	58	0.060	0.120	0.070	0.034
72/08/26	11 40	0000			22	135	8.90	54	0.050	0.080	0.381	0.168
	11 40	0004	18.7	8.6		130	8.90	55	0.050	0.080	0.443	0.239
72/11/03	14 30	0000	4.2	13.8	36	170	9.20	49	0.520	0.120	0.900	0.680

DATE FROM TO	TIME OF DAY	DEPTH FEET	32217 CHLOROPHYL A UG/L
72/06/26	06 05	0000	15.8J
72/08/26	11 40	0000	42.0J
72/11/03	14 30	0000	152.4J

J VALUE KNOWN TO BE IN ERROR