


**U.S. ENVIRONMENTAL PROTECTION AGENCY
NATIONAL EUTROPHICATION SURVEY
WORKING PAPER SERIES**



**CORVALLIS ENVIRONMENTAL RESEARCH LABORATORY - CORVALLIS, OREGON
and
ENVIRONMENTAL MONITORING & SUPPORT LABORATORY - LAS VEGAS, NEVADA**

REPORT
ON
LAKE STOREY
KNOX COUNTY
ILLINOIS
EPA REGION V
WORKING PAPER No. 318

WITH THE COOPERATION OF THE
ILLINOIS ENVIRONMENTAL PROTECTION AGENCY
AND THE
ILLINOIS NATIONAL GUARD
JUNE, 1975

CONTENTS

	<u>Page</u>
Foreword	ii
List of Illinois Study Lakes	iv
Lake and Drainage Area Map	v

Sections

I. Conclusions	1
II. Lake and Drainage Basin Characteristics	3
III. Lake Water Quality Summary	4
IV. Nutrient Loadings	8
V. Literature Reviewed	13
VI. Appendices	14

FOREWORD

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nationwide threat of accelerated eutrophication to fresh water lakes and reservoirs.

OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

- a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.
- b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.
- c. With such a transformation, an assessment of the potential for eutrophication control can be made.

LAKE ANALYSIS

In this report, the first stage of evaluation of lake and watershed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning [§303(e)], water quality criteria/standards review [§303(c)], clean lakes [§314(a,b)], and water quality monitoring [§106 and §305(b)] activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

ACKNOWLEDGMENT

The staff of the National Eutrophication Survey (Office of Research & Development, U. S. Environmental Protection Agency) expresses sincere appreciation to the Illinois Environmental Protection Agency for professional involvement and to the Illinois National Guard for conducting the tributary sampling phase of the Survey.

Dr. Richard H. Briceland, Director of the Illinois Environmental Protection Agency; and Ronald M. Barganz, State Survey Coordinator, and John J. Forneris, Manager of Region III, Field Operations Section of the Division of Water Pollution Control, provided invaluable lake documentation and counsel during the Survey, reviewed the preliminary reports, and provided critiques most useful in the preparation of this Working Paper series.


Major General Harold R. Patton, the Adjutant General of Illinois, and Project Officer Colonel Daniel L. Fane, who directed the volunteer efforts of the Illinois National Guardsmen, are also gratefully acknowledged for their assistance to the Survey.

NATIONAL EUTROPHICATION SURVEY

STUDY LAKES

STATE OF ILLINOIS

<u>LAKE NAME</u>	<u>COUNTY</u>
Baldwin	Randolph
Bloomington	McLean
Carlyle	Bond, Clinton, Fayette
Cedar	Lake
Charleston	Coles
Coffeen	Montgomery
Crab Orchard	Jackson, Williamson
Decatur	Macon
DePue	Bureau
East Loon	Lake
Fox	Lake
Grass	Lake
Highland Silver	Madison
Holiday	LaSalle
Horseshoe	Madison
Long	Lake
Lou Yaeger	Montgomery
Marie	Lake
Old Ben Mine	Franklin
Pistakee	Lake, McHenry
Raccoon	Marion
Rend	Franklin, Jefferson
Sangchris	Christian
Shelbyville	Moultrie, Shelby
Slocum	Lake
Springfield	Sangamon
Storey	Knox
Vandalia.	Fayette
Vermilion	Vermilion
Wee Ma Tuk	Fulton
Wonder	McHenry



LAKE STOREY

STORET NO. 1751

I. CONCLUSIONS

A. Trophic Condition:

Survey data indicate that Lake Storey is eutrophic. It ranked eleventh in overall trophic quality when the 31 Illinois lakes sampled in 1973 were compared using a combination of six lake parameters*. Eight lakes had less median total phosphorus, 11 had less median dissolved phosphorus, 22 had less median inorganic nitrogen, ten had less mean chlorophyll a, and three had greater mean Secchi disc transparency. Marked depression of dissolved oxygen with depth occurred at both sampling stations in August, and depletion occurred at station 1 at 20 feet in depth in October.

Survey limnologists noted a blue-green algal bloom in progress in October and reported much emergent and submerged vegetation in the shoreline shallows.

B. Rate-Limiting Nutrient:

The algal assay results indicate the lake was phosphorus limited at the time the sample was collected (05/12/73). These findings are substantiated by the lake data; i.e., the mean N/P ratios were 32/1 or greater at all sampling times.

* See Appendix A.

C. Nutrient Controllability:

1. Point sources--The phosphorus contribution of known point sources amounted to 54.5% of the total reaching Lake Storey during the sampling year. These point sources were Carl Sandburg Junior College (51.5%) and the Hawthorn Motel (3.0%).

The present phosphorus loading rate of $2.18 \text{ g/m}^2/\text{yr}$ is nearly five times that proposed by Vollenweider (Vollenweider and Dillon, 1974) as a eutrophic rate (see page 12). A 100% reduction in phosphorus loading from the known point sources would only lower the overall loading rate to $0.99 \text{ g/m}^2/\text{yr}$ (about twice the eutrophic rate). However, because Lake Storey is phosphorus limited, all phosphorus inputs to the lake should be minimized to the greatest practicable extent to slow the aging of this water body.

2. Non-point sources--Over 45% of the total phosphorus input to Lake Storey came from non-point sources during the sampling year. The Unnamed Creek A-2 contributed 31.2%, and the unaged tributaries were estimated to have contributed 13.4% of the total phosphorus load.

The phosphorus export rate of the Unnamed Creek A-2 was a relatively low $29 \text{ kg/km}^2/\text{yr}$ (see page 12).

II. LAKE AND DRAINAGE BASIN CHARACTERISTICS[†]

A. Lake Morphometry^{††}:

1. Surface area: 0.53 kilometers².
2. Mean depth: 4.6 meters.
3. Maximum depth: >7.2 meters.
4. Volume: 2.438×10^6 m³.
5. Mean hydraulic retention time: 282 days.

B. Tributary and Outlet:

(See Appendix C for flow data)

1. Tributaries -

<u>Name</u>	<u>Drainage area (km²)*</u>	<u>Mean flow (m³/sec)*</u>
Unnamed Creek (A-2)	12.4	0.1
Minor tributaries & immediate drainage -	<u>5.3</u>	<u><0.1</u>
Totals	17.7	0.1

2. Outlet -

Unnamed Creek (A-1)	18.2**	0.1
---------------------	--------	-----

C. Precipitation***:

1. Year of sampling: 140.7 centimeters.
2. Mean annual: 85.5 centimeters.

[†] Table of metric conversions--Appendix B.

^{††} Forneris, 1973.

* For limits of accuracy, see Working Paper No. 175, "...Survey Methods, 1973-1976".

** Includes area of lake.

*** See Working Paper No. 175.

III. LAKE WATER QUALITY SUMMARY

Lake Storey was sampled three times during the open-water season of 1973 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from two stations on the lake and from a number of depths at each station (see map, page v). During each visit, a single depth-integrated (4.6 m or near bottom to surface) sample was composited from the stations for phytoplankton identification and enumeration; and during the first visit, a single 18.9-liter depth-integrated sample was composited for algal assays. Also each time, a depth-integrated sample was collected from each of the stations for chlorophyll a analysis. The maximum depths sampled were 6.1 meters at station 1 and 3.7 meters at station 2.

The sampling results are presented in full in Appendix D and are summarized in the following table.

A. SUMMARY OF PHYSICAL AND CHEMICAL CHARACTERISTICS FOR LAKE STORY
STORET CODE 1751

PARAMETER	1ST SAMPLING (5/12/73)				2ND SAMPLING (8/9/73)				3RD SAMPLING (10/17/73)			
	> SITES				2 SITES				2 SITES			
	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN	RANGE	MEAN	MEDIAN
TEMP (C)	14.8 - 17.7	16.1	16.5	19.1 - 24.0	25.0	25.5	17.0 - 17.9	17.7	17.8			
DISS OXY (MG/L)	7.3 - 10.7	9.1	9.7	8.2 - 6.8	3.2	2.6	0.0 - 9.0	5.4	5.0			
CNDCTVY (MCHOMO)	460. - 675.	554.	545.	380. - 477.	422.	408.	390. - 485.	408.	399.			
PH (STAND UNITS)	7.4 - 8.4	8.2	8.2	7.3 - 9.1	8.1	8.0	7.3 - 8.5	8.0	7.9			
TOT ALK (MG/L)	175. - 240.	194.	182.	112. - 210.	150.	141.	125. - 248.	173.	167.			
TOT P (MG/L)	0.072 - 0.203	0.097	0.076	0.034 - 0.070	0.052	0.054	0.053 - 0.321	0.125	0.075			
ORTHOP (MG/L)	0.016 - 0.046	0.025	0.019	0.010 - 0.021	0.015	0.013	0.021 - 0.233	0.066	0.031			
NU2+NU3 (MG/L)	2.000 - 5.130	3.430	3.160	0.810 - 3.690	1.597	1.180	0.160 - 2.890	1.224	0.990			
AMMONIA (MG/L)	0.160 - 0.660	0.370	0.280	0.080 - 1.700	0.567	0.355	0.040 - 4.720	0.920	0.430			
KJEL N (MG/L)	0.700 - 1.000	0.833	0.800	1.500 - 2.800	2.050	2.000	0.900 - 6.200	1.900	1.300			
INORG N (MG/L)	2.240 - 5.460	3.800	3.595	1.040 - 4.130	2.163	2.075	1.340 - 4.880	2.144	1.440			
TOTAL N (MG/L)	2.700 - 6.030	4.263	4.000	2.960 - 5.190	3.647	3.440	2.180 - 6.360	3.124	2.360			
CHLRPHYL A (UG/L)	6.6 - 23.0	14.4	14.8	6.1 - 7.8	6.9	6.9	16.2 - 43.8	30.0	30.0			
SECCHI (METERS)	1.4 - 1.5	1.4	1.4	0.7 - 0.9	0.8	0.8	0.8 - 0.9	0.9	0.9			

B. Biological characteristics:

1. Phytoplankton -

<u>Sampling Date</u>	<u>Dominant Genera</u>	<u>Algal units per ml</u>
05/12/73	1. <u>Asterionella sp.</u> 2. <u>Stephanodiscus sp.</u> 3. <u>Melosira sp.</u> 4. <u>Anabaena sp.</u> 5. <u>Synedra sp.</u> Other genera	1,491 983 813 576 68 <u>168</u>
	Total	4,099
08/09/73	1. Coccoid cells 2. <u>Oscillatoria sp.</u> 3. <u>Coelastrum sp.</u> 4. <u>Oocystis sp.</u> 5. <u>Crucigenia sp.</u> Other genera	360 206 129 77 77 <u>256</u>
	Total	1,105
10/17/73	1. <u>Aphanizomenon sp.</u> 2. <u>Coelastrum sp.</u> 3. <u>Anabaena sp.</u> 4. <u>Microcystis sp.</u> 5. <u>Melosira sp.</u> Other genera	2,524 677 431 431 246 <u>801</u>
	Total	5,110

2. Chlorophyll a -

<u>Sampling Date</u>	<u>Station Number</u>	<u>Chlorophyll a ($\mu\text{g/l}$)</u>
05/12/73	01	6.6
	02	23.0
08/09/73	01	6.1
	02	7.8
10/17/73	01	16.2
	02	43.8

C. Limiting Nutrient Study:

1. Autoclaved, filtered, and nutrient spiked -

<u>Spike (mg/l)</u>	<u>Ortho P Conc. (mg/l)</u>	<u>Inorganic N Conc. (mg/l)</u>	<u>Maximum yield (mg/l-dry wt.)</u>
Control	0.015	4.475	3.3
0.050 P	0.065	4.475	30.6
0.050 P + 1.0 N	0.065	5.475	32.2
1.0 N	0.015	5.475	3.3

2. Discussion -

The control yield of the assay alga, Selenastrum capricornutum, indicates that the potential primary productivity of Lake Storey was moderately high at the time the sample was collected (05/12/73). Also, the addition of phosphorus alone produced a significant increase in yield, but no such increase occurred with the addition of only nitrogen. This indicates limitation by phosphorus.

The lake data substantiate phosphorus limitation. At all sampling times, the mean inorganic nitrogen/orthophosphorus ratios were 32/1 or greater.

IV. NUTRIENT LOADINGS
(See Appendix E for data)

For the determination of nutrient loadings, the Illinois National Guard collected monthly near-surface grab samples from each of the tributary sites indicated on the map (page v), except for the high runoff months of February and March when two samples were collected. Sampling was begun in June, 1973, and was completed in May, 1974.

Through an interagency agreement, stream flow estimates for the year of sampling and a "normalized" or average year were provided by the Illinois District Office of the U.S. Geological Survey for the tributary sites nearest the lake.

In this report, nutrient loads for sampled tributaries were determined by using a modification of a U.S. Geological Survey computer program for calculating stream loadings*. Nutrient loads shown are those measured minus point-source loads, if any.

Nutrient loads for unsampled "minor tributaries and immediate drainage" ("ZZ" of U.S.G.S.) were estimated using the nutrient loads, in kg/km²/year, at station A-2 and multiplying by the ZZ area in km².

The wastewater treatment plant operators of Carl Sandburg Junior College and the Hawthorn Motel did not participate in the Survey, and nutrient loads were estimated at 1.134 kg P and 3.401 kg N/capita/year.

* See Working Paper No. 175.

A. Waste Sources:

1. Known municipal* -

<u>Name</u>	<u>Pop. Served</u>	<u>Treatment</u>	<u>Mean Flow (m³/d)</u>	<u>Receiving Water</u>
Carl Sandburg Junior College	525**	sand filter	198.7	Lake Storey
Hawthorn Motel	30	sand filter	11.4	Lake Storey

2. Known industrial - None

* Anonymous, 1972.

** 1,400 students x 0.375 population equivalent.

B. Annual Total Phosphorus Loading - Average Year:

1. Inputs -

<u>Source</u>	<u>kg P/ yr</u>	<u>% of total</u>
a. Tributaries (non-point load) -		
Unnamed Creek (A-2)	360	31.2
b. Minor tributaries & immediate drainage (non-point load) -	155	13.4
c. Known municipal STP's -		
Carl Sandburg Junior College	595	51.5
Hawthorn Motel	35	3.0
d. Septic tanks - Unknown	-	-
e. Known industrial - None	-	-
f. Direct precipitation* -	<u>10</u>	<u>0.9</u>
Total	1,155	100.0

2. Outputs -

Lake outlet - Unnamed Creek (A-1) 355

3. Net annual P accumulation - 800 kg.

* See Working Paper No. 175.

C. Annual Total Nitrogen Loading - Average Year:

1. Inputs -

<u>Source</u>	<u>kg N/ yr</u>	<u>% of total</u>
a. Tributaries (non-point load) -		
Unnamed Creek (A-2)	14,445	62.6
b. Minor tributaries & immediate drainage (non-point load) -	6,175	26.8
c. Known municipal STP's -		
Carl Sandburg Junior College	1,785	7.7
Hawthorn Motel	100	0.4
d. Septic tanks - Unknown	-	-
e. Known industrial - None	-	-
f. Direct precipitation* -	<u>570</u>	<u>2.5</u>
Total	23,075	100.0

2. Outputs -

Lake outlet - Unnamed Creek (A-1) 16,620

3. Net annual N accumulation - 6,455 kg.

* See Working Paper No. 175.

D. Mean Annual Non-point Nutrient Export by Subdrainage Area:

<u>Tributary</u>	<u>kg P/km²/yr</u>	<u>kg N/km²/yr</u>
Unnamed Creek (A-2)	29	1,165

E. Yearly Loading Rates:

In the following table, the existing phosphorus loading rates are compared to those proposed by Vollenweider (Vollenweider and Dillon, 1974). Essentially, his "dangerous" rate is the rate at which the receiving water would become eutrophic or remain eutrophic; his "permissible" rate is that which would result in the receiving water remaining oligotrophic or becoming oligotrophic if morphometry permitted. A mesotrophic rate would be considered one between "dangerous" and "permissible".

Note that Vollenweider's model may not be applicable to water bodies with short hydraulic retention times.

	Total Phosphorus		Total Nitrogen	
	Total	Accumulated	Total	Accumulated
grams/m ² /yr	2.18	1.51	43.5	12.2

Vollenweider loading rates for phosphorus (g/m²/yr) based on mean depth and mean hydraulic retention time of Lake Storey:

"Dangerous" (eutrophic rate)	0.48
"Permissible" (oligotrophic rate)	0.24

V. LITERATURE REVIEWED

Anonymous, 1972. Wastewater treatment works data book. IL Env. Prot. Agency, Springfield.

Forneris, John J., 1973. Personal communication (lake morphometry). IL Env. Prot. Agency, Springfield.

Vollenweider, R. A., and P. J. Dillon, 1974. The application of the phosphorus loading concept to eutrophication research. Natl. Res. Council of Canada Publ. No. 13690, Canada Centre for Inland Waters, Burlington, Ontario.

VI. APPENDICES

APPENDIX A

LAKE RANKINGS

LAKE DATA TO BE USED IN RANKINGS

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500-MEAN SEC	MEAN CHLORA	15-MIN DO	MEDIAN DISS ORTHO P
1703	LAKE BLOOMINGTON	0.050	5.730	464.667	26.200	14.800	0.020
1706	LAKE CARLYLE	0.084	1.270	477.889	17.367	11.000	0.032
1708	LAKE CHARLESTON	0.160	4.680	490.667	12.000	8.400	0.065
1711	COFFEEN LAKE	0.032	0.260	456.222	7.700	14.900	0.012
1712	CRAB ORCHARD LAKE	0.082	0.200	482.222	59.867	13.800	0.013
1714	LAKE DECATUR	0.129	3.750	479.571	43.000	14.500	0.062
1725	LONG LAKE	0.704	1.190	482.667	49.333	8.800	0.398
1726	LAKE LOU YAEGER	0.186	1.600	489.583	10.662	11.400	0.076
1727	LAKE MARIE	0.098	0.370	467.667	39.533	14.700	0.057
1733	PISTAKEE LAKE	0.203	0.370	485.667	75.867	7.000	0.062
1735	REND LAKE	0.071	0.210	471.500	23.533	12.700	0.012
1739	LAKE SHELBYVILLE	0.062	3.290	461.333	17.161	14.800	0.019
1740	SILVER LAKE (HIGHLAND)	0.226	0.970	489.500	5.822	14.800	0.057
1742	LAKE SPRINGFIELD	0.108	3.265	483.385	13.013	10.800	0.059
1748	VERMILION LAKE	0.109	4.695	481.500	31.150	14.200	0.050
1750	WONDER LAKE	0.426	0.890	486.000	98.533	7.800	0.132
1751	LAKE STORY	0.072	2.510	459.333	17.250	14.800	0.021
1752	DEPUE LAKE	0.438	4.050	490.000	58.833	7.600	0.276
1753	LAKE SANGCHRIS	0.050	1.970	475.417	19.292	14.500	0.009
1754	LAKE HOLIDAY	0.167	3.135	485.167	51.217	7.200	0.046
1755	FOX LAKE	0.214	0.375	486.167	63.850	8.800	0.083
1756	GRASS LAKE	0.301	0.820	481.000	83.500	5.900	0.093
1757	EAST LOON LAKE	0.076	0.120	450.000	22.300	14.900	0.018
1758	SLOCUM LAKE	0.865	0.200	487.333	221.100	5.800	0.362
1759	CEDAR LAKE	0.029	0.170	400.333	5.767	12.800	0.013
1761	LAKE WEMATUK	0.069	1.770	466.333	7.967	14.500	0.031
1762	RACCOON LAKE	0.105	0.310	484.333	19.217	13.800	0.020
1763	BALUWIN LAKE	0.044	0.140	461.167	11.333	13.200	0.007

LAKE DATA TO BE USED IN RANKINGS

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLORA	15- MIN DO	MEDIAN DISS ORTHO P
1764	LAKE VANDALIA	0.116	0.480	478.111	11.276	14.800	0.023
1765	OLD BEN MINE RESERVOIR	0.930	0.205	478.333	31.433	11.200	0.575
1766	HORSESHOE LAKE	0.127	0.705	482.833	182.250	6.800	0.018

PERCENT OF LAKES WITH HIGHER VALUES (NUMBER OF LAKES WITH HIGHER VALUES)

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLORA	15- MIN DO	MEDIAN DISS ORTHO P	INDEX NU
1703	LAKE BLOOMINGTON	88 (26)	0 (0)	80 (24)	47 (14)	13 (2)	68 (20)	296
1706	LAKE CARLYLE	63 (19)	40 (12)	63 (19)	63 (19)	63 (19)	53 (16)	345
1708	LAKE CHARLESTON	37 (11)	7 (2)	0 (0)	77 (23)	77 (23)	27 (8)	225
1711	COFFEEN LAKE	97 (29)	77 (23)	93 (28)	93 (28)	2 (0)	92 (27)	454
1712	CRAB ORCHARD LAKE	67 (20)	90 (27)	43 (13)	20 (6)	42 (12)	85 (25)	347
1714	LAKE DECATUR	40 (12)	13 (4)	53 (16)	33 (10)	30 (8)	32 (9)	201
1725	LONG LAKE	7 (2)	43 (13)	40 (12)	30 (9)	72 (21)	3 (1)	195
1726	LAKE LOU YAEGER	30 (9)	37 (11)	7 (2)	87 (26)	57 (17)	23 (7)	241
1727	LAKE MARIE	60 (18)	68 (20)	73 (22)	37 (11)	23 (7)	42 (12)	303
1733	PISTAKEE LAKE	27 (8)	68 (20)	23 (7)	13 (4)	90 (27)	32 (9)	253
1735	REND LAKE	77 (23)	80 (24)	70 (21)	50 (15)	53 (16)	92 (27)	422
1739	LAKE SHELBYVILLE	83 (25)	17 (5)	83 (25)	70 (21)	13 (2)	73 (22)	339
1740	SILVER LAKE (HIGHLAND)	20 (6)	47 (14)	10 (3)	97 (29)	13 (2)	42 (12)	229
1742	LAKE SPRINGFIELD	53 (16)	20 (6)	33 (10)	73 (22)	67 (20)	37 (11)	283
1748	VERMILION LAKE	50 (15)	3 (1)	47 (14)	43 (13)	37 (11)	47 (14)	227
1750	WONDER LAKE	13 (4)	50 (15)	20 (6)	7 (2)	80 (24)	13 (4)	183
1751	LAKE STORY	73 (22)	27 (8)	90 (27)	67 (20)	13 (2)	63 (19)	333
1752	DEPUE LAKE	10 (3)	10 (3)	3 (1)	23 (7)	83 (25)	10 (3)	139
1753	LAKE SANGCHRIS	88 (26)	30 (9)	67 (20)	57 (17)	30 (8)	97 (29)	369
1754	LAKE HOLIDAY	33 (10)	23 (7)	27 (8)	27 (8)	87 (26)	50 (15)	247
1755	FOX LAKE	23 (7)	63 (19)	17 (5)	17 (5)	72 (21)	20 (6)	212
1756	GRASS LAKE	17 (5)	53 (16)	50 (15)	10 (3)	97 (29)	17 (5)	244
1757	EAST LOON LAKE	70 (21)	100 (30)	97 (29)	53 (16)	2 (0)	77 (23)	399
1758	SLOCUM LAKE	3 (1)	87 (26)	13 (4)	0 (0)	100 (30)	7 (2)	210
1759	CEDAR LAKE	100 (30)	93 (28)	100 (30)	100 (30)	50 (15)	85 (25)	528
1761	LAKE WEMATUK	80 (24)	33 (10)	77 (23)	90 (27)	30 (8)	57 (17)	367
1762	RACCOON LAKE	57 (17)	73 (22)	30 (9)	60 (18)	42 (12)	68 (20)	330
1763	BALDWIN LAKE	93 (28)	97 (29)	87 (26)	80 (24)	47 (14)	100 (30)	504

PERCENT OF LAKES WITH HIGHER VALUES (NUMBER OF LAKES WITH HIGHER VALUES)

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500- MEAN SEC	MEAN CHLORA	15- MIN DO	MEDIAN DISS ORTHO P	INDEX NU
1764	LAKE VANDALIA	47 (14)	60 (18)	60 (18)	83 (25)	13 (2)	60 (18)	323
1765	OLD BEN MINE RESERVOIR	0 (0)	83 (25)	57 (17)	40 (12)	60 (18)	0 (0)	240
1766	HORSESHOE LAKE	43 (13)	57 (17)	37 (11)	3 (1)	93 (28)	80 (24)	313

LAKES RANKED BY INDEX NOS.

RANK	LAKE CODE	LAKE NAME	INDEX NO
1	1759	CEDAR LAKE	528
2	1763	BALDWIN LAKE	504
3	1711	COFFEEN LAKE	454
4	1735	REND LAKE	422
5	1757	EAST LOON LAKE	399
6	1753	LAKE SANGCHRIS	369
7	1761	LAKE WEMATUK	367
8	1712	CRAB ORCHARD LAKE	347
9	1706	LAKE CARLYLE	345
10	1739	LAKE SHELBYVILLE	339
11	1751	LAKE STORY	333
12	1762	RACCOON LAKE	330
13	1764	LAKE VANDALIA	323
14	1766	HORSESHOE LAKE	313
15	1727	LAKE MARIE	303
16	1703	LAKE BLOOMINGTON	296
17	1742	LAKE SPRINGFIELD	283
18	1733	PISTAKEE LAKE	253
19	1754	LAKE HOLIDAY	247
20	1756	GRASS LAKE	244
21	1726	LAKE LOU YAEGER	241
22	1765	OLD BEN MINE RESERVOIR	240
23	1740	SILVER LAKE (HIGHLAND)	229
24	1748	VERMILION LAKE	227
25	1708	LAKE CHARLESTON	225
26	1755	FOA LAKE	212
27	1758	SLOCUM LAKE	210
28	1714	LAKE DECATUR	201

LAKES RANKED BY INDEX NOS.

RANK	LAKE CODE	LAKE NAME	INDEX NO
29	1725	LONG LAKE	195
30	1750	WONDER LAKE	183
31	1752	DEPUE LAKE	139

APPENDIX B

CONVERSIONS FACTORS

CONVERSION FACTORS

Hectares x 2.471 = acres

Kilometers x 0.6214 = miles

Meters x 3.281 = feet

Cubic meters x 8.107×10^{-4} = acre/feet

Square kilometers x 0.3861 = square miles

Cubic meters/sec x 35.315 = cubic feet/sec

Centimeters x 0.3937 = inches

Kilograms x 2.205 = pounds

Kilograms/square kilometer x 5.711 = lbs/square mile

APPENDIX C

TRIBUTARY FLOW DATA

TRIBUTARY FLOW INFORMATION FOR ILLINOIS

10/23/75

LAKE CODE 1751 LAKE STOREY

TOTAL DRAINAGE AREA OF LAKE(SQ KM) 18.2

TRIBUTARY	SUB-DRAINAGE AREA(SQ KM)	NORMALIZED FLOWS(CMS)												MEAN
		JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	
1751A1	18.2	0.11	0.18	0.21	0.22	0.17	0.15	0.09	0.04	0.03	0.03	0.05	0.06	0.11
1751A2	12.4	0.07	0.12	0.14	0.15	0.12	0.10	0.06	0.03	0.02	0.02	0.03	0.04	0.07
1751Z2	5.7	0.04	0.06	0.07	0.07	0.05	0.05	0.03	0.01	0.01	0.01	0.02	0.02	0.04

SUMMARY

TOTAL DRAINAGE AREA OF LAKE =	18.2	TOTAL FLOW IN =	1.33
SUM OF SUB-DRAINAGE AREAS =	18.2	TOTAL FLOW OUT =	1.33

MEAN MONTHLY FLOWS AND DAILY FLOWS(CMS)

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY		FLOW	DAY	FLOW	DAY	FLOW
				1	2					
1751A1	6	73	0.40	2	0.48					
	7	73	0.20	8	0.21					
	8	73	0.05	13	0.12					
	9	73	0.03	16	0.01					
	10	73	0.34	14	0.45					
	11	73	0.10	4	0.13					
	12	73	0.34	1	0.14					
	1	74	0.59	5	0.37					
	2	74	0.24	3	0.42	16	0.15			
	3	74	0.31	2	0.20	17	0.37			
	4	74	0.42	6	0.20					
	5	74	0.42	19	0.99					
1751A2	6	73	0.28	2	0.34					
	7	73	0.13	8	0.14					
	8	73	0.03	13	0.08					
	9	73	0.02	16	0.01					
	10	73	0.23	14	0.28					
	11	73	0.07	4	0.09					
	12	73	0.23	1	0.09					
	1	74	0.40	5	0.24					
	2	74	0.16	3	0.28	16	0.10			
	3	74	0.21	2	0.14	17	0.25			
	4	74	0.28	6	0.15					
1751Z2	5	74	0.28	19	0.68					
	6	73	0.12	2	0.14					
	7	73	0.07	8	0.07					
	8	73	0.02	13	0.04					
	9	73	0.01	16	0.01					
	10	73	0.11	14	0.17					
	11	73	0.03	4	0.04					
	12	73	0.10	1	0.04					
	1	74	0.20	5	0.13					
	1	74	0.18	5	0.11					
	2	74	0.07	3	0.14	16	0.05			
	3	74	0.10	2	0.07	17	0.12			
	4	74	0.14	6	0.05					
	5	74	0.14	19	0.31					

APPENDIX D

PHYSICAL and CHEMICAL DATA

STORET RETRIEVAL DATE 75/10/23

175101
40 59 20.0 090 24 30.0
LAKE STORY
17095 ILLINOIS

DATE FROM TO	TIME OF DAY	DEPTH FEET	00010 WATER TEMP CENT	00300 DO MG/L	00077 TRANSP SECCHI INCHES	00094 CNDUCTVY FIELD MICROMHO	11EPALES 3		2111202 0018 FEET DEPTH				00671 PHOS-DIS ORTHO MG/L P
							00400 PH SU	00410 TALK CACO3 MG/L	00610 NH3-N TOTAL MG/L	00625 TOT KJEL N MG/L	00630 N2&N03 N-TOTAL MG/L		
73/05/12	13 35 0000		16.2		60	540	8.10	240	0.660	0.800	2.130	0.032	
	13 35 0004		15.9	8.9		540	8.10	240	0.630	0.700	2.000	0.022	
	13 35 0015		14.8	6.3		675	7.90	176	0.230	1.000	2.010	0.046	
	15 00 0000		25.8	6.4	34	380	9.00	112	0.100	2.000	0.990	0.012	
73/08/09	15 00 0005		25.3			405							
	15 00 0009		24.2	1.0		433	7.80	148	0.440	1.500	3.690	0.010	
	15 00 0015		19.1	0.2		475	7.30	210	1.700	2.800	0.810	0.020	
	11 00 0000		17.9		37	399	7.90	167	0.450	1.300	0.990	0.022	
73/10/17	11 00 0005		17.8	5.0		400	7.80	167	0.430	1.200	0.980	0.031	
	11 00 0015		17.8	4.8		399	7.40	170	0.640	1.400	0.960	0.076	
	11 00 0020		17.0	0.0		485	7.30	248	4.720	6.200	0.160	0.233	

DATE FROM TO	TIME OF DAY	DEPTH FEET	00665 PHOS-TOT MG/L P	32217 CHLRPHYL A UG/L	
73/05/12	13 35 0000		0.079	6.6	
	13 35 0004		0.073		
	13 35 0015		0.203		
	15 00 0000		0.061	6.1	
73/08/09	15 00 0009		0.034		
	15 00 0015		0.041		
	11 00 0000		0.053	16.2	
	11 00 0005		0.075		
73/10/17	11 00 0015		0.106		
	11 00 0020		0.321		

STORET RETRIEVAL DATE 75/10/23

175102
 40 59 20.0 090 23 25.0
 LAKE STORY
 17095 ILLINOIS

11EPALES
 3 2111202
 0015 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	WATER TEMP CENT	00010 DO	00300 TRANSP	00077 SECCHI INCHES	00094 CNDCTVY FIELD MICRUMHO	00400 PH	00410 TALK CACO3	00610 NH3-N TOTAL MG/L	00625 TOT KJEL N MG/L	00630 NO2&NO3 N-TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P
73/05/12	13 55	0000	17.0			54	560	8.30	175	0.210	0.800	4.190	0.016
	13 55	0004	17.0		10.7		550	8.40	185	0.330	0.900	5.130	0.017
	13 55	0011	16.8		10.5		460	8.30	178	0.160	0.800	5.120	0.017
73/08/09	15 25	0000	28.0	6.8		26	393	9.10	113	0.080	2.000	0.960	0.012
	15 25	0005	27.4				404						
	15 25	0009	27.4		4.3		410	8.20	134	0.270	1.900	1.370	0.014
	15 25	0012	22.5		0.4		477	7.40	180	0.810	2.100	1.760	0.021
73/10/17	11 15	0000	17.9			33	390	8.50	167	0.090	1.400	1.250	0.021
	11 15	0005	17.9		8.4		392	8.50	167	0.070	0.900	1.340	0.031
	11 15	0010	17.8		9.0		391	8.50	125	0.040	0.900	2.890	0.047

DATE FROM TO	TIME OF DAY	DEPTH FEET	PHOS-TOT MG/L P	00665 CHLRPHYL A UG/L	32217
73/05/12	13 55	0000	0.074	23.0	
	13 55	0004	0.079		
	13 55	0011	0.072		
73/08/09	15 25	0000	0.060	7.8	
	15 25	0009	0.070		
	15 25	0012	0.049		
73/10/17	11 15	0000	0.070	43.8	
	11 15	0005	0.071		
	11 15	0010	0.178		

APPENDIX E

TRIBUTARY DATA

STORET RETRIEVAL DATE 75/10/23

1751A1
 40 59 20.0 090 24 40.0
 UNNAMED CREEK
 17111 KNOX CO MAP
 0/LAKE STOREY
 SEC RD BRDG JUST BELO DAM
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 N02&N03 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
73/06/02	11 00		3.800	4.400	0.160	0.019	0.065
73/07/08	09 15		2.500	3.230	0.231	0.015	0.030
73/08/13	08 45		0.520	4.400	0.096	0.021	0.065
73/09/16	10 20		0.240	1.540	0.325	0.027	0.115
73/10/14	16 00		2.900	2.650	0.095	0.098	0.250
73/11/04	14 00		1.140	1.750	0.357	0.011	0.065
73/12/01	10 00		1.400	1.450	0.224	0.012	0.021
74/01/05	10 15		2.400	0.600	0.160	0.009	0.030
74/02/03	13 50		3.740	1.700	0.210	0.210	0.315
74/02/16	10 00		2.760	1.400	0.130	0.095	0.145
74/03/02	11 00		4.700	1.200	0.050	0.015	0.075
74/03/17	09 45		9.600	1.000	0.035	0.030	0.057
74/04/06	13 20		4.200	1.500	0.050	0.010	0.035
74/05/19	10 40		3.300	1.000	0.030	0.010	0.045

STORET RETRIEVAL DATE 75/10/23

1751A2
 40 59 20.0 090 22 55.0
 UNNAMED CREEK
 17 KNUX CO MAP
 I/LAKE STOREY
 US HWY 150 BRDG AT E END OF LAKE
 11EPALES 2111204
 4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 NO2&N03 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
73/06/02	10 40		4.500	3.500	0.076	0.013	0.110
73/07/08	08 50		2.500	2.940	0.074	0.022	0.110
73/08/13	08 30		0.990	4.600	0.096	0.030	0.080
73/09/16	10 00		0.280	2.300	0.294	0.054	0.195
73/10/14	15 40		0.560	1.000	0.095	0.011	0.050
73/11/04	13 45		9.200	0.100K	0.075	0.042	0.100
73/12/01	09 45		8.300	0.116	0.116	0.044	0.270
74/01/05	10 30		9.100	0.300	0.048	0.036	
74/02/03	13 40		5.300	1.100	0.230	0.310	0.490
74/02/16	09 30		3.200	0.900	0.110	0.100	0.150
74/03/02	11 00		7.200	0.800	0.050	0.020	0.075
74/03/17	10 05		4.100	1.100	0.025	0.010	0.065
74/04/06	13 40		5.460	0.800	0.015	0.010	0.025
74/05/19	11 10		9.700	0.700	0.075	0.075	0.135

K VALUE KNOWN TO BE
 LESS THAN INDICATED